

## BIXACEAE LIPSTICKTREE FAMILY

Wendy Hodgson  
Desert Botanical Garden  
1201 N. Galvin Parkway  
Phoenix, AZ 85008

Trees, shrubs, or rhizomatous herbs. LEAVES alternate, from simple, entire and palmately veined to palmately lobed or palmately compound, often with a mucilaginous epidermis; stipules well developed. INFLORESCENCES panicles, racemes, or cymes, sometimes appearing before the leaves. FLOWERS perfect, actinomorphic or zygomorphic; sepals and petals 5, distinct; stamens numerous, arranged in 2-5 groups, sometimes borne on a nectariferous disk; ovary superior, carpels 2-5, styles 1 scarcely lobed, the ovules numerous. FRUITS loculicidal capsules. SEEDS with oily, proteinaceous, or starchy endosperm. --3 genera, 12 spp., chiefly of tropical and subtropical regions.

### *Amoreuxia* Moc. & Sessé ex DC.

Herbaceous perennials, the rootstock starchy to woody. LEAVES long-petioled, subentire or deeply parted into 5-9 lobes, orbicular to reniform. INFLORESCENCES cymes, terminal, few-flowered. FLOWERS zygomorphic, 5-8 cm in diameter; sepals linear to oblong-ovate, imbricate; petals showy, flesh-salmon, yellow, to orange-salmon, the lateral petals with 1 conspicuous maroon mark at the base, upper petals with 2 conspicuous maroon marks at the base; stamens unequal, in 2 sets, the filaments and anthers of the lower set slightly longer and tinged with maroon or cream; anthers with 2 apical pores; ovary with 3 carpels. FRUITS globose to ellipsoid, pendulous, the exocarp woody at maturity, endocarp very thin, transparent. SEEDS reniform, subglobose, or globose, the seed coat subglabrous, pilose, or strigose; endosperm oily. --4 spp; sw U.S., n Mex. through C. Amer. to Colombia and Peru, also on Curacao. (for P. J. Amoreux). Sprague, T. 1922. Kew Bull. 1922(3):97-105. Poppendieck, H. 1981. Flora Neotropica Monogr. 27:18-26.

1. Capsules ellipsoid, 4.5-8 cm long; seeds globose; ovary densely silky; arillode loosely fitting and often shed before dehiscence of capsule; lower set of anthers cream-colored or light maroon; leaves 5-7(-9)-lobed ..... *A. gonzalezii*
- 1' Capsule broadly ovoid, 2-4.2 cm long; seeds strongly reniform; ovary densely puberulent-papillose; arillode closely fitting and persistent; lower set of anthers dark maroon; leaves (5-)7-9-lobed ..... *A. palmatifida*

*Amoreuxia gonzalezii* Sprague & Riley (for Gonzalez Ortega). Saiya. --LEAVES cuneate to broadly cuneate, (5-)7(-9)-lobed, 2.6-5.5 cm long (not including 3-10 cm long petiole), 4.7-8.5 cm wide. FLOWERS 7-8 cm long, 6-8 cm wide; sepals long-pilose; petals pale yellow-salmon or deep salmon tinged with maroon, the basal spots pale maroon; upper anthers cream, the lower set of anthers cream or pale maroon; ovary densely silky. CAPSULES ellipsoid, striated longitudinally (less so in ours). 5.7-8 cm long, 2.3-3.9 cm wide, long-acuminate, short-pilose. SEEDS globose, black, dull, 4.7 mm long, 4.3 mm wide; seed coat light brown, easily detached, sparsely short-pilose, the hairs often forked. --Limestone hills and granitic slopes; Pima and Santa Cruz cos., to be expected in Cochise Co.; 1310 m (4300 ft); Jul-Sep; Baja C., Sin., Son., Mex.

*Amoreuxia palmatifida* Moc. & Sessé (palmately divided). Saiya, Tamaquí. --Leaves narrow to broadly cuneate, 7(-9)-lobed, 2.2-5.3(-8) cm long (not including 3-10 cm long petiole), 4-7.7(-13.5) cm wide. FLOWERS 4.3-6 cm long, 4.5-6.4 cm wide; sepals puberulent; petals yellow or salmon-orange to salmon-maroon, with deep maroon basal spots; anthers all dark maroon or only the lower set dark maroon; ovary

densely puberulent-papillose. CAPSULES broadly ovoid, not noticeably striated longitudinally, 2-3.6 cm long, 2-2.6 cm wide, short-acuminate, puberulent. SEEDS reniform, black, shiny, 5.3 mm long, 3.3 mm wide; seed coat light brown, strigillose, closely fitting and persistent. -- Dry rocky slopes and mesas; Cochise, Pima, and Santa Cruz cos.; 800-1500 m (2600-4900 ft); Jul-Sep; Mex. to C. Amer. and Colombia.

The roots, young leaves, flowers, fruits and seeds of both species were a food source to tribes in southwestern U.S. and northern Mex.




Figure 1. p = *A. palmatifida*: pf=fruit(x.8), pr=root(x.5), ph=habit(x.5), ps=seed with overlying seed coat(x1.8), pfs=fruit surface(x2), po=ovary(x2.5); distribution in map insert = circles. g = *A. gonzalezii*: gf=fruit(x.5), gs=seed with overlying seed coat(x1.5), gfs=fruit surface(x2), go=ovary(x2.5); distribution in map insert = diamonds.