



Chemical Constituents of *Cymbocarpum erythraeum* (DC.) Boiss., and Evaluation of Its Anti-*Helicobacter pylori* Activity

Cymbocarpum erythraeum (DC.) Boiss.'in Kimyasal Bileşikleri ve *Helicobacter pylori*'ye Karşı Etkisi

© Samaneh HEIDARI¹, © Azadeh MANAYI², © Soodabeh SAEIDNIA², © Hossein MIGHANI¹, © Hamid Reza MONSEF ESFAHANI³, © Ahmad Reza GOHARI^{2*}, © William N SETZER⁴

¹Golestan University, Faculty of Science, Department of Chemistry, Golestan, Iran

²Tehran University of Medical Sciences, Faculty of Pharmacy, Medicinal Plants Research Center, Tehran, Iran

³Tehran University of Medical Sciences, Faculty of Pharmacy, Department of Pharmacognosy, Tehran, Iran

⁴University of Alabama in Huntsville, Faculty of College of Science, Departments of Chemistry, Huntsville, USA

ABSTRACT

Objective: *Cymbocarpum erythraeum* (Apiaceae) is an endemic species in Iran. Up to now, there have been no phytochemical and biological investigations on this species. Therefore, isolation of the main secondary metabolites of the plant along with its anti-*H. pylori* activity have been considered in this paper.

Materials and Methods: The dried parts of the plant were extracted with different solvents using solvent percolation and the antibacterial activity of the extracts evaluated by the disk diffusion method. Four compounds were isolated using different column chromatography methods.

Results: The compounds were identified by proton nuclear magnetic resonance and carbon-13 nuclear magnetic resonance as isoquercetin (1), rutin (2), β -sitosterol (3) and 2-decenol (4).

Conclusion: Anti-*H. pylori* evaluation of the extracts and isolated compounds against three clinical isolates of *H. pylori* revealed that hexane extract of the plant inhibited all *H. pylori* strains.

Key words: Apiaceae, *C. erythraeum*, chromatography, disk diffusion, *H. pylori*

ÖZ

Amaç: *Cymbocarpum erythraeum* (Apiaceae) İran'a endemik bir türdür. Bu zamana kadar, bu tür üzerinde herhangi bir fitokimyasal ve biyolojik aktivite çalışmasına rastlanmamıştır. Bu nedenle, bu çalışmada, bitkinin ana sekonder metabolitlerinin ve *Helicobacter pylori*'ye karşı etkisinin araştırılması amaçlanmıştır.

Gereç ve Yöntemler: Kurutulmuş bitki kısımları perkolasyon yöntemi ile farklı çözücüler kullanılarak ekstre edilmiş ve ekstrelerin antibakteriyel etkisi disk difüzyon yöntemiyle değerlendirilmiştir. Farklı kolon kromatografisi yöntemleri kullanılarak dört bileşik izole edilmiştir.

Bulgular: Bileşiklerin yapıları, proton nükleer manyetik rezonans ve karbon-13 nükleer manyetik rezonans tekniği ile izokersetin (1), rutin (2), β -sitosterol (3) ve 2-dekenol (4) olarak belirlenmiştir.

Sonuç: Ekstrelerin ve izole edilen bileşiklerin üç *H. pylori* klinik izolatına karşı etkilerinin değerlendirildiği bu çalışmada, bitkinin hekzan ekstresinin tüm *H. pylori* suşlarını ihibe ettiği tespit edilmiştir.

Anahtar kelimeler: Apiaceae, *C. erythraeum*, kromatografi, disk difüzyon, *H. pylori*

*Correspondence: E-mail: a_goharii@tums.ac.ir, Phone: +9821-64121202 ORCID-ID: orcid.org/0000-0001-9071-1663

Received: 16.04.2017, Accepted: 27.07.2017

©Turk J Pharm Sci, Published by Galenos Publishing House.

INTRODUCTION

Cymbocarpum is represented in Iran by three species that occur naturally in the wild, including *Cymbocarpum marginatum*, *Cymbocarpum erythraeum*, and *Cymbocarpum anethoides*.^{1,2} The essential oils had been obtained from these three species of *Cymbocarpum* from Iran and analyzed indicating that the oils of all three plants were rich in aliphatic aldehydes. Furthermore, the oil of *C. erythraeum*, with main constituent (*E*)-2-decenal (52.22%), showed larvicidal activity.³ The main components of the essential oils of the fruits and herbal parts of *Cymbocarpum wiedemannii* were found to be aliphatic aldehydes and aliphatic acids.⁴

A major cause of bacterial gastrointestinal infections is *Helicobacter pylori*. In fact, *H. pylori* has been designated as a class I carcinogen by World Health Organization and its eradication has been reported to be beneficial in preventing gastric disorders specially ulcer and cancer.⁵ Given the extensive treatment with antibiotics for decades, the failure rates due to antimicrobial resistance range from 20% to 40% and the eradication failure rate remains as high as 5-20%.⁶⁻⁸ Regarding the previous study which indicated that methanol extract of the plant showed antibacterial activity⁹ and anti-*H. pylori* activity of some flavonoids¹⁰ the present study was designed to evaluate anti-*H. pylori* property of the plant extracts and its flavonoids.

MATERIALS AND METHODS

Plant material

The flowering aerial parts of *C. erythraeum* were collected from the East Azerbaijan province (June 2010) with voucher No. 214 at the Herbarium of Institute of Medicinal Plants, Iranian Academic Centre for Education, Culture and Research, Karaj, Iran.

General

Silica gel (70-230 mesh, F254 pre-coated plates) (Merck, Germany), reverse phase silica gel 90 (RP-18C; Fluka, Switzerland) and Sephadex LH-20 (Fluka, Switzerland) were used for isolation of compounds. Semi-preparative high-performance liquid chromatography [HPLC (RP-18C; Knauer, Germany)] and medium-pressure liquid chromatography [MPLC (Silica gel, 230-400 mesh; Butchi, Switzerland)] were used for more purification. Nuclear magnetic resonance (NMR) experiments were performed on Bruker (Billerica, USA) DRX 500 instrument [500 MHz for proton NMR (¹H-NMR), 125 MHz for carbon-13 (¹³C-NMR)] with tetramethylsilane as an internal standard. Ultraviolet (UV) spectra were measured on Optizen (Daejeon, Korea) model 2021 UV plus. All solvents were distilled before use.

Isolation procedure

Dried aerial parts of *C. erythraeum* (500 g) were extracted with hexane, methanol and water-methanol (1:1) using the solvent percolation method at room temperature. Extracts were concentrated to obtain 11, 80, 34 g of hexane, methanol and methanol-water (1:1) extracts, respectively. The methanol extract was further partitioned by petroleum ether, butanol and

water. The butanol fraction (8 g) was subjected to Sephadex LH₂₀ column to afford B_A-B_E fractions and those suspected to contain flavonoids under UV light were loaded on another column for more purification. A sub-fractions (B_A: 3980 mg, B_B: 84 mg) were loaded on Sephadex LH₂₀ column with methanol as eluent to obtain compounds 1 (8.4 mg) and 3 (12 mg). The other sub-fraction (B_C: 106.6 mg) was injected to HPLC (RP-18C) eluted with different percentage of methanol-water (2:3, 1:1, and 3:2) to yield compound 2 (27 mg). In addition, the hexane extract was injected to MPLC (normal phase silica gel column) eluted with hexane-chloroform (4:1 and 0:1) to give 12 primary fractions (H_A-H_L). One fraction (H_J: 239 mg) was subjected to Sephadex LH₂₀ column with chloroform-ethyl acetate-methanol (1:1:1) as eluent to provide pure compound 4 (4.1 mg).

Anti-*H. pylori* assay

Clinical *H. pylori* (HP₁, HP₂, and HP₃) strains were used to determine antimicrobial susceptibility following previously published protocols using the disk diffusion method.^{11,12} Serial dilutions of the test samples were made in dimethyl sulfoxide (DMSO). Suspensions of bacteria in normal saline were prepared with the turbidity of McFarland standard No. 2 (6×10⁸ cell/mL). Plates of non-selective blood agar were inoculated with 100 μL of each bacterial suspension and allowed to dry at room temperature. Ten μL of test samples was introduced into a sterile blank disks and deposited on the surface of the inoculated plates. Negative and positive control included blank disks impregnated with 10 μL of DMSO and amoxicillin (1 μg/mL), respectively. Plates were incubated at 37°C under microaerobic conditions and examined after 3-5 days. The mean inhibition zone diameters (IZD) ± standard deviation were recorded.

RESULTS AND DISCUSSION

The isolated compounds from aerial parts of *C. erythraeum* were identified as isoquercetin (1), rutin (2),^{13,14} along with β-sitosterol (3),¹⁵⁻¹⁷ and 2-decenol (4) based on the spectroscopic data (¹H-NMR, ¹³C-NMR) and compared to the pertinent spectroscopic data in previously published literature (Figure 1).⁹

To the best of our knowledge, this is the first report on the isolation and elucidation of secondary metabolites of *C.*




Figure 1. Chemical structures of the isolated compounds from *C. erythraeum*

Table 1. Inhibition zone diameters \pm standard deviation of some active extracts of *C. erythraeum* against clinical isolates of *H. pylori*

Samples	IZD \pm SD (mm)											
	1 Conc.			½ Conc.			¼ Conc.			1/8 Conc.		
	HP1	HP2	HP3	HP1	HP2	HP3	HP1	HP2	HP3	HP1	HP2	HP3
Hexane ¹	15 \pm 0.03	17 \pm 0.04	20 \pm 0.03	14 \pm 0.05	15 \pm 0.03	18 \pm 0.01	10 \pm 0.03	14 \pm 0.05	16 \pm 0.04	9 \pm 0.01	10 \pm 0.02	12 \pm 0.05
Methanol ²	-	-	14 \pm 0.04	-	-	13 \pm 0.04	-	-	10 \pm 0.02	-	-	-
PE ³	-	-	15 \pm 0.02	-	-	12 \pm 0.03	-	-	10 \pm 0.03	-	-	-

Conc.: Concentration, ¹: 440 mg/mL, ²: 464 mg/mL, ³: 226 mg/mL, HP: Strains of *H. pylori*, PE: Petroleum ether, -: Not sensitive, IZD for amoxicillin (1 μ g/mL): 30 \pm 0.04 mm, SD: Standard deviation

erythraeum and its anti-*H. pylori* activity. The susceptibility of the tested bacteria were different related to the extracts and HP₃ was the most susceptible strain. The IZD of the effective test samples against *H. pylori* are summarized in (Table 1). The growth of all bacterial strains were suppressed by amoxicillin (1 μ g/mL) with IZD of 30 \pm 0.04 mm, while DMSO showed no inhibitory activity toward the tested organisms. Methanol (464 mg/mL) and petroleum ether (226 mg/mL) extracts had inhibited one of the clinical isolates of *H. pylori* (HP₃) with IZD of 14 \pm 0.04 and 15 \pm 0.02 mm, respectively, while compounds 1 (16 mg/mL), 2 (14 mg/mL) and the flavonoid fraction (138 mg/mL), along with other extracts were not effective against the isolated strains.

In a previous study, rutin isolated from *Gardenia jasminoides* exhibited antibacterial activity toward *H. pylori*, however this present study showed the compound not to be active against the isolated strains.¹⁸ Additionally, some flavonoids are known to be antibacterial agents.¹⁹ For example, the flavonoid-rich extract of *Glycyrrhiza glabra* inhibited *H. pylori* growth by inhibition of protein synthesis, DNA gyrase and dihydrofolate reductase¹⁰, while the flavonoid fraction of *C. erythraeum* was not active against examined strains, which may be presumably due to the difference in the flavonoid constituents of the plants and/or the susceptibility of bacteria. Whereas in this study, the hexane extract of the plant mainly inhibited the *H. pylori* strains attributed to the presence of non-polar compounds. It was reported that essential oil of *C. erythraeum*, which was rich in (*E*)-2-decenal (52.22%), was biologically active and exhibited larvicidal activity.³ In the present study, the alcoholic compound 2-decenol (**4**), which was isolated from hexane fraction, is likely to be one of the active constituents of this fraction as well as other constituents in this fraction. The medium-chain-length alcohol decanol, from *Coriandrum sativum* oil, showed antibacterial activity especially against Gram-positive strains.^{20,21} As a matter of fact, the relatively polar substances such as flavonoids and other polar fractions are not generally able to effectively prevent the bacterium growth, suggesting that the anti-*H. pylori* activity of this plant can be largely attributed to other non-polar secondary metabolites.

Conflict of Interest: No conflict of interest was declared by the authors.

REFERENCES

- Ajani Y, Mojab F, Tabatabaai S, Ramazan-Pour B. Analysis of the essential oils from 3 species of *Cymbocarpum* (Apiaceae) from Iran. *Res Pharm Sci.* 2012;7:s708.
- Ajany Y. Studied taxonomic genus, *Cymbocarpum*, *Diplotaenia*, *Johrenia*, *Dorema haussknechtia*, *Azilia*, *Opopanax*, *Smyrniopsis* of *Umbelliferus* Iran; Tehran University; 2004.
- Aksakal Ö, Uysal H, Çolak DA, Mete E. Inhibition effects of essential oil of *Cymbocarpum erythraeum* (DC.) Boiss., on percentage of survival from larvae to adult in *Drosophila melanogaster* and its chemical composition. *MAKÜ FEBED.* 2012;3:32-36.
- Baser KHC, Özek T, Vural M. Essential Oil of *Cymbocarpum wiedemannii* Boiss. *J Essent Oil Res.* 1999;11:679-680.
- Zaidi SF, Muhammad JS, Shahryar S, Usmanghani K, Gilani AH, Jafri W, Sugiyama T. Anti-inflammatory and cytoprotective effects of selected Pakistani medicinal plants in *Helicobacter pylori*-infected gastric epithelial cells. *J Ethnopharmacol.* 2012;141:403-410.
- Li Y, Xu C, Zhang Q, Liu JY, Tan RX. *In vitro* anti-*Helicobacter pylori* action of 30 Chinese herbal medicines used to treat ulcer diseases. *J Ethnopharmacol.* 2005;98:329-333.
- Lien HM, Wang CY, Chang HY, Huang CL, Peng MT, Sing YT, Chen CC, Lai CH. Bioevaluation of *Anisomeles indica* extracts and their inhibitory effects on *Helicobacter pylori*-mediated inflammation. *J Ethnopharmacol.* 2013;145:397-401.
- Zhang XQ, Gu HM, Li XZ, Xu ZN, Chen YS, Li Y. Anti-*Helicobacter pylori* compounds from the ethanol extracts of *Geranium wilfordii*. *J Ethnopharmacol.* 2013;147:204-207.
- Ajani Y, Mojab F, Hajiaghahi R, Taghizadeh F, Jamalifar H. Assaying of volatile oil components and antimicrobial effects of methanolic extract of it in *Cymbocarpum erythraeum* (DC.) Boiss and *C anethoides* DC; 2012.
- Asha MK, Debraj D, Prashanth D, Edwin JR, Srikanth HS, Muruganantham N, Dethe SM, Anirban B, Jaya B, Deepak M, Agarwal A. *In vitro* anti-*Helicobacter pylori* activity of a flavonoid rich extract of *Glycyrrhiza glabra* and its probable mechanisms of action. *J Ethnopharmacol.* 2013;145:581-586.
- Manayi A, Khanavi M, Saeidnia S, Azizi E, Mahmoodpour MR, Vafi F, Malmir M, Siavashi F, Hadjiakhoondi A. Biological activity and microscopic characterization of *Lythrum salicaria* L. *Daru.* 2013;21:61.
- Siavoshi F, Saniee P, Latifi-Navid S, Massarrat S, Sheykholeslami A. Increase in resistance rates of *H. pylori* isolates to metronidazole and tetracycline--comparison of three 3-year studies. *Arch Iran Med.* 2010;13:177-187.
- Gohari A, Saeidnia S, Shahverdi A, Yassa N, Malmir M, Mollazade K, Naghinejad AR. Phytochemistry and antimicrobial compounds of *Hymenocrater calycinus*. *Eur Asia J BioSci.* 2009;3:64-68.

14. Jafari S, Saeidnia S, Hajimehdipoor H, Ardekani MR, Faramarzi MA, Hadjiakhoondi A, Khanavi M. Cytotoxic evaluation of *Melia azedarach* in comparison with *Azadirachta indica* and its phytochemical investigation. *Daru*. 2013;21:37.
15. Saeidnia S, Ghamarinia M, Gohari AR, Shakeri A. Terpenes from the root of *Salvia hypoleuca* Benth. *Daru*. 2012;20:66.
16. Saeidnia S, Nourbakhsh MS, Gohari AR, Davood A. Isolation and identification of the main compounds of *Satureja sahendica* Bornm. *Aust J Basic Appl Sci*. 2011;5:1450-1453.
17. Saeidnia S, Gohari AR, Malmir M, Moradi-Afrapoli F, Ajani Y. Tryptophan and sterols from *Salvia limbata*. *J Med Plants*. 2011;10:41-44.
18. Lee JH, Kang KJ, Lee YM, Kim PN, Jeong CS. Effects of *Gardenia jasminoides* Ellis ethanol extract and its constituents on anti-gastritis and anti-gastric cancer cells. *Planta Med*. 2008;74:289.
19. Sahreen S, Khan MR, Khan RA, Shah NA. Estimation of flavonoids, antimicrobial, antitumor and anticancer activity of *Carissa opaca* fruits. *BMC Complement Altern Med*. 2013;13:372.
20. Huhtanen CN. Inhibition of *Clostridium botulinum* by Spice Extracts and Aliphatic Alcohols. *J Food Prot*. 1980;43:195-196.
21. Kubo I, Muroi H, Kubo A. Antibacterial activity of long-chain alcohols against *Streptococcus mutans*. *J Agric Food Chem*. 1993;41:2447-2450.