



Review

Tetraoxygenated naturally occurring xanthones

Valdir Peres^a, Tanus Jorge Nagem^{b,*}, Fernando Faustino de Oliveira^b^aDepartamento de Química, Universidade Federal de Viçosa, Av. PH Rolfs, 36571-000, Viçosa, Minas Gerais, Brazil^bDepartamento de Química da Universidade Federal de Ouro Preto, Universidade Campus do Cruzeiro, 35400-000, Ouro Preto, Minas Gerais, Brazil

Received 4 June 1999; received in revised form 19 June 2000

Abstract

This review, with 350 references, gives information on the chemical study of 234 naturally occurring tetraoxygenated xanthones in 12 families, 53 genus and 182 species of higher plants, and two which are described as fungal and lichen metabolites. The value of these groups of substances in connection with pharmacological activity and therapeutic use of some species is described. The structural formulas of 135 isolated compounds, and their distribution, are also given. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: Xanthone; Tetraoxygenated xanthones; Pharmacological activity; Fungus; Lichen; Biosynthesis

Contents


1. Introduction.....	684
2. Classification.....	684
3. Methods of isolation and structural investigations.....	684
4. Bioactivities of xanthones.....	685
5. Use of plant sources containing tetraoxygenated xanthones	685
6. Tetraoxygenated xanthones	689
7. Synthesis and biosynthesis of xanthones	694
8. Conclusion	705
Acknowledgements.....	705
References	705

* Corresponding author. Tel.: +55-31-9611551; fax: 55-31-5591707.
E-mail address: tjnagem@bhnet.com.br (T.J. Nagem).

1. Introduction

Xanthones are secondary metabolites commonly occurring in a few higher plant families, and in fungi and lichens. Their taxonomic importance in such families and their pharmacological properties has aroused great interest (Cardona et al., 1990).

The symmetrical nature of the xanthone nucleus coupled with its mixed biogenetic origin in higher plants necessitates that the carbons be numbered according to a biosynthetic convention. Carbons 1–4 are assigned to the acetate-derived ring A, and carbons 5–8 to the shikimate-derived ring B (Bennett and Lee, 1989). The numbering system is based on xanthene-9-one as the basic skeleton (Gottlieb, 1968), and in cases where only ring B is oxygenated the lowest numbers are used, except for biosynthetic discussions (Bennett and Lee, 1989).



Xanthone basic skeleton

2. Classification

The xanthones isolated so far may be classified into five major groups: simple oxygenated xanthones, xanthone glycosides, prenylated and related xanthones, xanthonolignoids, and miscellaneous xanthones (Mandal et al., 1992). The simple oxygenated xanthones can again be subdivided into six groups according to the degree of oxygenation, and we have previously reviewed trioxygenated xanthones (Peres and Nagem, 1997). In a continuation of our interest in these compounds, simple tetraoxigenated xanthones, as well as prenylated and related xanthones with the same degree of oxygenation, are listed in this review (Tables 2–6).

3. Methods of isolation and structural investigations

Xanthones are commonly separated by silica gel chromatography using different solvent mixtures (Harborne, 1973). Xanthones are also separated and identified by comparison with authentic samples by TLC (Chawla et al., 1975) and by HPLC (Hostettmann and Guillarmod, 1976a,b; Hostettmann and McNair, 1976). The structures of simple oxygenated xanthones have been established mainly from analysis of the UV, IR, MS and NMR spectroscopic data of these compounds (Roberts, 1961; Arends and Helboe, 1972; Arends et al., 1973; Sultanbawa, 1980).

Xanthones can be detected by their colour in UV light with and without ammonia, or by using a general phenolic spray (Harborne, 1973). The UV spectrum varies in a characteristic manner, depending on the oxygenation pattern, and with the availability of a considerable amount of data, assignments can be readily made. By the use of AlCl_3 shifts for chelated hydroxyls, as well as sodium acetate, sodium hydroxide and boric acid shifts, considerable information on the position of hydroxyl groups can be obtained (Mesquita et al., 1968; Sultanbawa, 1980).

The use of IR spectroscopy in xanthone chemistry is generally limited to the detection of the carbonyl stretching frequency (Roberts, 1961; Mourão et al., 1966). For example, the effect of chelation on the IR carbonyl frequency of hydroxy-xanthones may be a useful feature for analysis of spectra of substituted and extended xanthones (Scheinmann, 1962; Chawla et al., 1973). The use of IR for detecting other functional groups, such as unchelated hydroxyl and methyl groups, has been described elsewhere (Roberts, 1961; Santesson, 1968a,b, 1969).

By introducing lichen samples into a mass spectrometer via a direct inlet system, Santesson (1968a,b) obtained mass spectra of volatile lichen substances. The method proved to be well suited for tentative identification of lichen xanthones (Santesson, 1968a,b, 1969). Apart from discussion of the mass spectra of lichen xanthones, no systematic investigation of the electron-impact-induced fragmentation of xanthones appears to have been made, except by Arends et al. (1973) in their study of electron-impact-induced fragmentation of monohydroxy- and monomethoxyxanthones. Mass spectrometry has not been applied extensively to the study of naturally occurring xanthones, but the MS data available have been very valuable for preliminary examination (Barnes and Occolowitz, 1964).

The data obtained from ^1H NMR spectroscopic analysis are of great value in characterizing and identifying naturally occurring xanthones (Mathis and Goldstein, 1964). Indeed, these have been used to determine the structure of substituents and for locating aromatic protons, through comparison with reference data and analysis of spin–spin coupling. Closer scrutiny of the chemical shifts of aromatic protons allows prediction of the oxygenation pattern (Barraclough et al., 1970), and there are detailed NMR results for this class of compounds (Westerman et al., 1977).

^1H NMR and ^{13}C NMR spectroscopic analyses are the most useful tools in the structure elucidation of xanthones (Chaudhuri et al., 1978). The ^{13}C NMR spectra of a great number of naturally-occurring xanthones are reported, with all chemical shifts assigned (Castelão et al., 1977; Westerman et al., 1977; Frahm and Chaudhuri, 1979). Hambloch and Frahm (1980) introduced a computer program named SEOX 1, which

rapidly identifies unknown xanthones with the help of additivity rules; this lends remarkable facility to structure elucidation.

4. Bioactivities of xanthones

The study of xanthones is interesting not only from a chemosystematic viewpoint, but also from a pharmacological point of view. Xanthones possess antidepressant action and antitubercular activity, while xanthone glycosides exhibit depressant action. The choleric, diuretic, antimicrobial, antiviral and cardiotonic action of some xanthones has also been established (Hostettmann and Wagner, 1977; Suzuki et al., 1980, 1981; Kitanov and Blinova, 1987). The inhibition of type A and type B monoamine oxidases by a number of xanthones has also been observed (Suzuki et al., 1980, 1981). And in spite of their restricted occurrence in the plant kingdom, some xanthones are reported to possess antileukaemic, antitumour, antiulcer, antimicrobial, antihepatotoxic and CNS-depressant activities (Banerji et al., 1994). For example, bellidifolin (1,5,8-trihydroxy-3-methoxy-xanthone) was found to be a selective inhibitor of MAO A, whereas psorospermin, a dihydrofuranoxanthone epoxide, exhibits significant activity against leukaemia and in colon and mammary tumour models (Beerhues and Berger, 1994). In general, xanthones and their derivatives have also been shown to be effective as allergy inhibitors and bronchodilators in the treatment of asthma (Balasubramanian and Rajagopalan, 1988).

A series of isoprenylated xanthones isolated from moraceous plants showed interesting biological activities such as hypotensive effect, anti-rhinoviral activity, inhibition of the formation of some prostanoids and anti-tumour promoting activity (Hano et al., 1990). In a study of structure–activity relationships, it was found that 1,3,5,6-, 1,3,6,7-, 2,3,6,7- and 3,4,6,7-tetraoxxygenated xanthones possess antiplatelet effects, and that the mechanism of action of 1,3,6,7-tetraoxxygenated xanthones is due to both inhibition of thromboxane formation and phosphoinositide breakdown (Lin et al., 1993). Some xanthone dicarboxylic acids have shown a potent inhibition of the binding of leukotriene B₄ to receptors on intact neutrophils. Norathyriol showed anti-inflammatory effects mediated partly through the suppression of mast cell degranulation, and partly through (at least at higher doses) a non-selective blockade of the increase in vascular plasma exudation caused by various mediators (Lin et al., 1996). Bellidifolin, isolated from *Swertia japonica*, was found to be a potent hypoglycemic agent in STZ-induced diabetic rats by both oral and intraperitoneal administration (Basnet et al., 1995).

Recently, various bioactivities of xanthones including cytotoxic and antitumour activities, anti-inflammatory activity, antifungal activity, enhancement of choline

acetyltransferase activity and inhibition of lipid peroxidase have been described (Iinuma et al., 1994). In 1994 Mehta et al., reported the total synthesis of novel xanthone antibiotics, cervinomycins A₁ and A₂, with promising activity against anaerobic bacteria, mycoplasma and some gram positive bacteria (Table 1).

5. Use of plant sources containing tetraoxxygenated xanthones

Anthocleista vogelli (Planch), Loganiaceae, is a small tree which grows in the tropical rain forest areas of west Africa. The stem bark is used in local medicine for curing fever, stomach ache and as purgative (Okorie, 1976).

Among *Calophyllum* species, *C. inophyllum* is the most widespread and has superior timber qualities, particularly in Malaysia (Al-Jeboury and Locksley, 1971). The balsam from the bark of *C. inophyllum* "Alexandrian Laurel" is an oleoresin and used as a cicatrisant, whereas an infusion or decoction of the leaves has been traditionally used as an eye remedy in Asian medicine (Basnet et al., 1995). Xanthones from *C. inophyllum* produce CNS depression in rats and mice (Bennett and Lee, 1989). The tropical American tree *C. brasiliense* produces a strong and durable timber known as "Santa Maria" or "jacareuba" (King et al., 1953).

Canscora decussata Schult (Gentianaceae), an erect annual plant of height 0–6 m, finds use in indigenous medicine for a variety of purposes. The roots are used as a laxative, diuretic, for liver troubles, as a nerve tonic and in tuberculosis and fevers, while the aerial portions are used to treat cases of insanity, epilepsy and nervous debility (Ghosal and Chaudhuri, 1973; Okorie, 1976). The extract of *C. decussata* Schult is used in the treatment of certain mental disorders and tuberculosis (Ghosal and Chaudhuri, 1975).

The botanical source of the Taiwan folk remedy "Hwang-jin-guey" is the root and stem of *Cudrania cochinchinensis* (Lour.) Kudo and Masamune var. *gerontogea* (S. & Z.) Kudo and Masamune (Gottlieb et al., 1975). It was used in the treatment of neuralgia, rheumatism, hepatitis and contusions (Chang et al., 1994). Ethanolic extracts of the roots of this species also showed significant anti-inflammatory and liver protective effects, and the extract as well as isolated xanthone compounds exhibited anti-lipid peroxidative activities both in vivo and in vitro (Chang et al., 1995). *C. tricuspidata* (Carr.) Bur. (Japanese name "Hariguwa", Moraceae) is a deciduous tree whose cortex and root bark have been used as a Chinese crude drug (Hano et al., 1990a,b,c).

Plant extracts obtained from genera *Eustoma* are used to treat various ailments, including constipation, nervous debility, tuberculosis, fever and anorexia (Sullivan et al., 1977).

Table 1

Natural source of tetraoxxygenated xanthone

Family Asteraceae	
Genus Senecio — <i>S. mikanioides</i> [60]	<i>G. subelliptica</i> [103, 162, 163, 168, 169, 247, 248] <i>G. terpnophylla</i> [27] <i>G. thwaitesii</i> [135]
Family Betulaceae	
Genus Alnus — <i>A. glutinosa</i> [197]	Genus Haploclathra — <i>H. leiantha</i> [256, 257, 259] <i>H. paniculata</i> [258, 259]
Family Caryophyllaceae	
Genus Saponaria — <i>S. vaccaria</i> [192, 193]	Genus Hypericum — <i>H. androsaemum</i> [263, 176] <i>H. aucheri</i> [200] <i>H. canariensis</i> [54] <i>H. ericoides</i> [55, 117] <i>H. maculatum</i> [14] <i>H. patulum</i> [175, 176, 178, 179] <i>H. paturum</i> [177] <i>H. reflexum</i> [53] <i>H. roeperanum</i> [290] <i>H. sampsonii</i> [71]
Family Clusiaceae	
Genus Allanblackia — <i>A. floribunda</i> [226, 227]	Genus Kayea — <i>K. stylosa</i> [131]
Genus Archytaea — <i>A. multiflora</i> [206]	Genus Kielmeyera — <i>K. candidissima</i> [98] <i>K. coriacea</i> [59] <i>K. ferruginea</i> [121] <i>K. rupestris</i> [74, 91] <i>K. speciosa</i> [120]
Genus Bonnetia — <i>B. dinizii</i> [270]	Genus Lorostemon — <i>L. coelholi</i> [48] <i>L. negrensis</i> [48]
Genus Calophyllum — <i>C. austroindicum</i> [172]	Genus Mammea — <i>M. africana</i> [56] <i>M. acuminata</i> [330]
<i>C. apetalum</i> [165]	Genus Mesua — <i>M. ferrea</i> [337]
<i>C. bracteatum</i> [312]	Genus Ochrocarpus — <i>O. odoratus</i> [226]
<i>C. brasiliensis</i> [72, 124, 198, 280]	Genus Pentaphalangium — <i>P. solomonse</i> [271]
<i>C. calaba</i> [209, 312]	Genus Poeciloneuron — <i>P. pauciflorum</i> [56]
<i>C. canum</i> [57]	Genus Psorospermum — <i>P. febrifugum</i> [2, 4, 138]
<i>C. cordato-oblongum</i> [134]	Genus Rheedia — <i>R. benthamiana</i> [80]
<i>C. cuneifolium</i> [130]	<i>R. brasiliensis</i> [82] <i>R. gardneriana</i> [83, 84]
<i>C. fragrans</i> [224]	Genus Symphonia — <i>S. globulifera</i> [220, 221, 222, 223]
<i>C. inophyllum</i> [10, 119, 171, 173, 184, 208]	Genus Tovomita — <i>T. brasiliensis</i> [46]
<i>C. neo-ebudicum</i> [303]	<i>T. macrophylla</i> [245, 269] <i>T. mangle</i> [237] <i>T. pyrifolium</i> [47, 245]
<i>C. ramiflorum</i> [39]	Genus Pentadesma — <i>P. butyracea</i> [81]
<i>C. sclerophyllum</i> [182]	Genus Vismia — <i>V. guaramirangae</i> [132] <i>V. guineensis</i> [45]
<i>C. scriblitifolium</i> [183]	
<i>C. thwaitesii</i> [77]	
<i>C. tomentosum</i> [28, 191]	Family Gentianaceae
<i>C. trapezifolium</i> [128]	
<i>C. walkeri</i> [77]	Genus Blackstonia — <i>B. perfoliata</i> [334]
<i>C. zeylanicum</i> [133]	Genus Canscora — <i>C. decussata</i> [66, 107–109] <i>C. cachanlahuen</i> [335]
Genus Caraipa — <i>C. densiflora</i> [59, 214, 215]	Genus Centaurium — <i>C. erythraea</i> [35, 190, 243, 261, 262, 327] <i>C. linearifolium</i> [114, 116, 272–274] <i>C. littorale</i> [35, 333] <i>C. pulchellum</i> [246]
<i>C. grandifolia</i> [104]	
<i>C. psidifolia</i> [104]	Genus Chironia — <i>C. krebsii</i> [342]
<i>C. valloii</i> [104]	Genus Erythraea — <i>E. centaurium</i> [327]
Genus Cratoxylum — <i>C. celebicum</i> [317, 318]	
<i>C. cochinchinense</i> [36, 310]	
<i>C. formosanum</i> [166]	
<i>C. pruniflorum</i> [199]	
Genus Garcinia — <i>G. cowa</i> [202, 211, 280]	
<i>G. densivenia</i> [339]	
<i>G. dioica</i> [179]	
<i>G. dulcis</i> [148, 164, 166, 167]	
<i>G. echinocarpa</i> [27]	
<i>G. eugenifolia</i> [184]	
<i>G. forbesii</i> [147]	
<i>G. mangostana</i> [20, 26, 92, 126, 152, 202, 275, 276, 298, 305, 306, 307, 308, 316, 328]	
<i>G. multiflora</i> [70]	
<i>G. nervosa</i> [12]	
<i>G. opaca</i> [118]	
<i>G. ovalifolia</i> [338]	
<i>G. pedunculata</i> [289]	
<i>G. polyantha</i> [12]	
<i>G. pyrifera</i> [12]	
<i>G. staudtii</i> [340]	

Table 1 (continued)

Genus Eustoma — <i>E. grandiflorum</i> [319]	<i>S. petiolata</i> [195, 207]
Genus Frasera — <i>F. albicaulis</i> [314]	<i>S. punicea</i> [101]
<i>F. albomarginata</i> [90]	<i>S. purpureascens</i> [8, 111]
<i>F. carolinensis</i> [313]	<i>S. racemosa</i> [329]
<i>F. speciosa</i> [90]	<i>S. speciosa</i> [194, 238]
Genus Gentiana — <i>G. algida</i> [51, 326]	<i>S. tetrapetala</i> [7]
<i>G. barbata</i> [285]	
<i>G. bavarica</i> [156, 158]	Genus Veratrilla — <i>V. baillonii</i> [348]
<i>G. bellidifolia</i> [52, 233, 234, 235]	
<i>G. brachyphylla</i> [155, 156]	Family Gesneriaceae
<i>G. campestris</i> [52, 188]	Genus Orphium — <i>O. frutescens</i> [297]
<i>G. cerastioides</i> [187]	
<i>G. ciliata</i> [239]	Family Iridaceae
<i>G. detonsa</i> [187]	Genus Iris — <i>I. nigricans</i> [11]
<i>G. favratii</i> [156]	
<i>G. germanica</i> [52, 186]	Family Loganiaceae
<i>G. karelinii</i> [50, 326]	Genus Anthocleista — <i>A. vogelli</i> [99]
<i>G. kochiana</i> [136, 292]	
<i>G. lactea</i> [302]	Family Lytraceae
<i>G. nivalis</i> [155, 156]	Genus Lawsonia — <i>L. inermis</i> [41, 42, 229]
<i>G. ramosa</i> [52, 186]	
<i>G. rostanii</i> [156]	Family Moraceae
<i>G. schleicherii</i> [156]	Genus Chlorophora — <i>C. tinctoria</i> [122]
<i>G. strictiflora</i> [187]	Genus Cudrania — <i>C. cochinchinensis</i> [62, 64]
<i>G. tenella</i> [52]	<i>C. javanensis</i> [255]
<i>G. utriculosa</i> [156]	<i>C. tricuspidata</i> [140, 141, 142, 264]
<i>G. verna</i> [153, 156]	Genus Maclura — <i>M. aurantica</i> [17]
Genus Gentianopsis — <i>G. paludosa</i> [349]	<i>M. pomifera</i> [75, 76, 86, 343–346]
Genus Halenia — <i>H. asclepidea</i> [315]	Genus Morus — <i>M. insignis</i> [143, 144]
<i>H. campanulata</i> [161]	<i>M. tinctoria</i> [185]
<i>H. corniculata</i> [266, 286, 296]	
<i>H. elliptica</i> [38, 87, 88, 322]	Family Polygalaceae
Genus Hoppea — <i>H. dichotoma</i> [110]	Genus Bredemeyera — <i>B. brevifolia</i> [268]
<i>H. fastigiata</i> [251–254]	Genus Monnina — <i>M. obtusifolia</i> [283]
Genus Schultesia — <i>S. lisianthoides</i> [328]	<i>M. sylvatica</i> [31]
Genus Swertia — <i>S. bimaculata</i> [113]	Genus Polygala — <i>P. arillata</i> [232]
<i>S. calycina</i> [295]	<i>P. nyikensis</i> [236]
<i>S. chirata</i> [61, 79, 112, 231]	<i>P. tenuifolia</i> [100, 174, 181]
<i>S. chirayita</i> [22]	<i>P. virgata</i> [32]
<i>S. cuneata</i> [196]	
<i>S. decussata</i> [78, 292]	Family Polypodiaceae
<i>S. dilatata</i> [329]	Genus Athyrium — <i>A. mesosorum</i> [265, 331, 332]
<i>S. gracilisces</i> [329]	
<i>S. hookeri</i> [106]	Fungus
<i>S. erythrosticta</i> [160]	Genus Aspergillus — <i>A. versicolor</i> [49, 151]
<i>S. iberica</i> [85]	
<i>S. japonica</i> [18, 19, 21, 33, 180, 189, 203]	Lichen
<i>S. lawii</i> [19]	Genus Diploschistes — <i>Diploschistes</i> sp [95]
<i>S. macroperma</i> [350]	
<i>S. mileensis</i> [150, 219]	
<i>S. mussotii</i> [31]	
<i>S. nervosa</i> [329]	
<i>S. paniculata</i> [13]	
<i>S. patens</i> [149]	
<i>S. perennis</i> [154, 292]	
<i>S. perfoliata</i> [145]	

Garcinia (Clusiaceae) is best known in Malaysia as a genus of fruit trees. The fruit of many species are edible and serves as a substitute for tamarinds in curries. Many species produce a yellow resin which is used in the making of varnishes and in treatment of wounds. Some species are used in traditional medicine and as a source of edible oils. Extracts and pure isolates of *Garcinia* species have been shown to exhibit significant antimicrobial and pharmacological activities (Goh et al., 1992).

Garcinia dulcis grows mainly in southeast Asia, and its leaves and seeds have been used in traditional medicine against lymphatitis, parotitis, struma and other disease conditions (Iinuma et al., 1996). The oil obtained from the seeds of *G. echinocarpa* is used for lighting lamps (Bandaranayake et al., 1975). *G. livingstonei* is a small to medium-sized tree producing edible fruits and growing at low altitude. It is found, particularly in South Africa, in riverine fringes and in open woodland. Extracts of the leaves and flowers are reported to exhibit antibiotic properties (Diserens et al., 1992). The fruit hull of *G. mangostana* L., the "Mangosteen" tree, is used in Thai folk medicine for healing skin infections and wounds, and for the relief of diarrhea (Mahabusarakam et al., 1987). It is fairly widespread in India, Sri Lanka and Burma, and is known for the sweet fruits called mangosteen. In the Ayurvedic system of medicine, the fruit hull of this plant finds wide application, mainly as an anti-inflammatory agent and in the treatment of diarrhoea (Balasubramanian and Rajagopalan, 1988).

Garcinia subelliptica is a small shrub 4–5 m in height or a large tree sometimes reaching 15–20 m, and has been extensively cultivated as a windbreak in the Yaeyama islands of Japan. Its bark has been utilized as a source of a yellow coloured dye (Fukuyama et al., 1991). The timber of *G. terpnophylla* is used for building purposes (Bandaranayake et al., 1975).

The roots and rhizomes of the yellow gentian (*Gentiana lutea*) have been much used for the production of a bitter tonic. Any medicinal value possessed by gentian preparations appears to be due to bitter, complex glycosides (Roberts, 1961). *Hoppea dichotoma* Willd. is used in the Ayurvedic system of medicine in the treatment of haemorrhoids, in cardiac disorders and in certain mental disorders (Ghosal et al., 1978).

A number of species of the genus *Hypericum* have been found to possess various biological properties. Among the activities is inhibition of monoamine oxidase (MAO; EC 1.4.3.4). MAO, which exists as two isoenzymes, MAO-A and MAO-B, plays a key role in the regulation of some physiological amines and is the target of inhibitors used as antidepressive drugs. The leaves and young twigs of *H. balearicum* contain lipophilic glandular products (Wollenweber et al., 1994). The xanthones of *H. brasiliense* inhibit MAO activity (Rocha et al., 1994); the stems, leaves and flowers of

H. ericoides are used in Valentian folk medicine (Cardona and Seoane, 1982), and ethanol extracts of the aerial parts of *H. mysorensis* show antifungal action (Nattaya et al., 1996). Extracts of *H. perforatum* are widely used in Europe as a drug for depression treatment. *H. rooperanum*, a shrub or small tree growing in central, eastern and south tropical Africa, is employed, alone or in association with various plants or animal parts, to cure female sterility (Rath et al., 1996). *H. sampsonii* Hance is known as a tumour inhibiting plant in Formosan folk medicine (Chen and Chen, 1985).

Lawsonia inermis L. ("Henna") is a plant used by ancient Egyptians for preserving mummies. It is presently used in India and Africa for dyeing hair and for staining hands and feet. It was also reported that the leaves are used for treating some skin diseases including Tinea of the legs (Mahmoud et al., 1980).

Mesua ferrea is commonly called ironwood in Malaya. The hard and durable trunkwood is used widely in agricultural tools and vehicles in southeast Asian countries (Chow and Quon, 1968). From the expressed oil of its seeds, two crystalline antibiotic principles have been isolated, mesuol and mesuone (Govindachari et al., 1967). In contrast to mangiferin, which is reported to be a CNS stimulant (Bhattacharya et al., 1972), xanthones from *M. ferrea* result in CNS depression in rats and mice (Bennett and Lee, 1989).

Monnieria obtusifolia is used in the folk medicine of Ecuador as an antifungal, antitumoural, antipyrrhea and antiseptic and as a skin cleanser (Pinto et al., 1994).

The roots of *Polygala tenuifolia* are used as an expectorant, tonic and sedative agent under the names "Onji" in Japan and "Yuan zhi" in China (Ito et al., 1977; Ikeya et al., 1991). It is also effective in inhibiting congestive oedema in rats (Fujita et al., 1992).

Psorospermum febrifugum (Clusiaceae) is a woody plant of tropical Africa which has been used as a febrifuge, a leprosy treatment, a poison antidote, and a purgative. An ethanolic extract of *P. febrifugum* showed antileukaemic activity in vivo in the P388 lymphocytic leukaemia in mice and in vitro in the KB cell culture system (Kupchan et al., 1980). Six compounds isolated from the extract were found to exhibit significant in vitro cytotoxic activity against 9PS cells in culture, and another exhibited both in vitro cytotoxic and in vivo antitumour activity (Abou-Shoer et al., 1988, 1989a,b). One of these six xanthones exhibited significant cytotoxicity in the HT-29 human colon adenocarcinoma in vitro cell line (Abou-Shoer et al., 1993); the xanthone, sorospermin, isolated from *P. febrifugum*, exhibited cytotoxic and in vivo antitumour activity in the P388 mouse leukaemia, mammary (CD) and colon (C6) models (Habib et al., 1987).

Saponaria vaccaria is an annual herb growing as a weed in cultivated fields in northern Pakistan, and is widespread in the flora of Russia. It is reputed to possess

sudorific, emetic and laxative properties, and is also used in the indigenous system of medicine for the treatment of jaundice, rheumatism, hepatic eruption and venereal ulcers. It also shows toxicity against worms, and paramecium and other unicellular organisms (Kazmi et al., 1989).

Plants of the genus *Swertia* (Gentianaceae) have been used in traditional medicine for many years. Because these herbs taste extremely bitter and possess the ability to reduce fever, detoxify and act as choleric and liver tonics, they have been mainly used for the treatment of hepatic, choleric and inflammatory diseases, such as hepatitis, cholecystitis, pneumonia, osteomyelitis, dysentery, scabies, spasm, pain and neurasthenia. Both methanolic and dichloromethane extracts of *Swertia calycina* exhibit strong antifungal activity against *Cladosporium cucumerinum* and *Candida albicans* (Rodriguez et al., 1995).

Swertia chirata is widely distributed in India in the temperate Himalayas between 4000 and 10,000 ft, from Kashmir to Bhutan, and in Khasia Hills between 4000 and 5000 ft. It also grows abundantly in Nepal. The plant is well known for its use in the Indian system of medicine for a variety of purposes. The extract of the plant is used as a bitter tonic, stomachic, antimalarial and as a remedy for liver disorders, a febrifuge, an anthelmintic, a remedy for scanty urine, in epilepsy and for certain types of mental disorders (Ghosal et al., 1973). Recently, the hexane extract of the plant has been shown to possess promising hypoglycaemic activity (Chakravarty et al., 1994). *S. chirata* found a place among useful medicinal plants of east and southeast Asia as a bitter tonic, febrifuge, stomachic, laxative and blood purifier (Mukherjee and Mukherjee, 1987).

Swertia chirayita (Roxb. ex. Flem.) Karsh grows abundantly in the temperate regions of the Himalayas. It is used in India as a traditional remedy for chronic fever, anaemia, asthma and liver disorders, and as a bitter tonic. A recent report revealed hypoglycaemic activity in the hexane extract of this plant (Asthana et al., 1991).

The extract of *Swertia hookeri* is used by people for a variety of therapeutic purposes, e.g. in the treatment of microbial infections in man, in hypertension and as a mood enhancer (Ghosal et al., 1980). *S. japonica* Makino, a popular medicinal herb in Japan, usually grows in a cold climate. The whole plant has been widely used as a folk medicine for stomach complaints, and has a characteristic bitter taste. The *Swertia* herb has also been widely used in Ayurveda and Unani medicines as an anthelmintic, febrifuge and bitter tonic. The aqueous ethanolic extract of *S. japonica* and its fractions showed potent hypoglycaemic activity in STZ-induced diabetic rats (Basnet et al., 1994).

By using chemiluminescence assays (CL) Ashida et al. (1994) recently showed that a natural Japanese folk

medicine, named "Senburi" (*Swertia japonica* Makino), contained components with antioxidant activity. Antioxidant activities of six isolated xanthone derivatives are comparable to α -tocopherol and BHT (Ashida et al., 1994). In India, *S. chirata* decoctions are a commonly used household remedy for constipation and weak liver function in children. An infusion of the plant has been used by Hindu physicians as a bitter tonic, stomachic, febrifuge and anthelmintic. Its use has also been reported in Unani medicine as a substitute for official gentiana preparations.

Swertia macrosperma is a medicinal plant used as a febrifuge, an antidote and a stomach tonic by the indigenous population in the southwestern part of China (Zhou et al., 1989). *S. mileensis* and *S. mussotii* are especially efficacious for treating acute viral hepatitis, some preparations have been produced industrially in China.

Swertia petiolata is bitter in taste and is used for its laxative and antimalarial properties in the folk medicine of the high altitude Himalayan region (Khetwal et al., 1990). *S. punicea* grows in moist, mountainous areas in the central district of mainland China, and has been used in the folk medicine as a bitter stomachic (Fukamiya et al., 1990). Extracts of *S. purpureescens* Wall are very commonly used as a tonic and febrifuge in the indigenous system of medicine in India (Ahmad et al., 1973).

Vismia guineensis is a small tree growing in west tropical Africa, whose roots and bark are used as a remedy for skin diseases (Gunasekera et al., 1977).

6. Tetraoxygenated xanthones

The first reported naturally occurring tetraoxygenated xanthone is mangostin, isolated in 1855 from *Garcinia mangostana*; in spite of numerous attempts, its structure was not fully elucidated until more than 100 years later (Roberts, 1961). The first naturally occurring tetraoxygenated xanthone whose structure is described in the literature is 1,5,8-trihydroxy-3-methoxyxanthone, isolated from *S. japonica* in 1942 (Asahina et al., 1942).

The review by Roberts (1961) refers to the natural occurrence of seven tetraoxygenated xanthones, namely decussatin, jacareubin, mangostin, swerchirin, swertianol, swertinin and asperxanthone. Afzal et al. (1979) described 51 tetraoxygenated xanthones isolated from higher plants, and also described fungal and lichen metabolites. The review by Sultanbawa (1980) included 37 tetraoxygenated xanthones from tropical plants, isolated from different mentioned species, with a dramatic increase in the number of known natural tetraoxygenated xanthones.

In a review of *Hypericum* genus, Kitanov and Blinova (1987) listed only five simple tetraoxygenated xanthones; two years later, in a review of xanthones from

Table 2

Simple tetraoxxygenated naturally occurring xanthones

Laxanthone II (1) <i>Lawsonia inermis</i> [42–229]		<i>C. erythraea</i> [190] <i>C. linariifolium</i> [272] <i>C. pulchellum</i> [246] <i>Chironia krebsii</i> [342] <i>Eustoma grandiflorum</i> [319] <i>Gentiana barbata</i> [285] <i>G. bavarica</i> [156–158] <i>G. brachyphylla</i> [155, 156] <i>G. campestris</i> [52] <i>G. ciliata</i> [239] <i>G. detonosa</i> [187] <i>G. favrati</i> [156] <i>G. kochiana</i> [136–292] <i>G. nivalis</i> [155, 156] <i>G. ramosa</i> [52] <i>G. rostanii</i> [156] <i>G. schleicherii</i> [156] <i>G. tenella</i> [52] <i>G. utriculosa</i> [156] <i>G. verna</i> [156] <i>Gentianopsis paludosa</i> [349] <i>Orphium frutescens</i> [297] <i>Swertia bimaculata</i> [113] <i>S. calycina</i> [295] <i>S. chirata</i> [112–161] <i>S. decussata</i> [78] <i>S. dilatata</i> [329] <i>S. gracilescens</i> [329] <i>S. hookeri</i> [106] <i>S. iberica</i> [160] <i>S. lawii</i> [105] <i>S. mileensis</i> [150] <i>S. mussotii</i> [321] <i>S. nervosa</i> [329] <i>S. paniculata</i> [13] <i>S. patens</i> [149] <i>S. perennis</i> [154] <i>S. perfoliata</i> [145] <i>S. punicea</i> [101] <i>S. purpurascens</i> [111] <i>S. racemosa</i> [329] <i>Schultesia lisianthoides</i> [328]
1-Hydroxy-2,3,5-trimethoxyxanthone (3) <i>Frasera albicaulis</i> [314] <i>F. albomarginata</i> [90] <i>F. carolinensis</i> [313] <i>F. speciosa</i> [90] <i>Halenia asclepidea</i> [315] <i>H. campanulata</i> [161] <i>H. corniculata</i> [266, 286] <i>H. elliptica</i> [87] <i>S. millensis</i> [219] <i>S. tetrapetala</i> [7] <i>Veratrilla baillonii</i> [348]		<i>Gentianopsis paludosa</i> [349] <i>Orphium frutescens</i> [297] <i>Swertia bimaculata</i> [113] <i>S. calycina</i> [295] <i>S. chirata</i> [112–161] <i>S. decussata</i> [78] <i>S. dilatata</i> [329] <i>S. gracilescens</i> [329] <i>S. hookeri</i> [106] <i>S. iberica</i> [160] <i>S. lawii</i> [105] <i>S. mileensis</i> [150] <i>S. mussotii</i> [321] <i>S. nervosa</i> [329] <i>S. paniculata</i> [13] <i>S. patens</i> [149] <i>S. perennis</i> [154] <i>S. perfoliata</i> [145] <i>S. punicea</i> [101] <i>S. purpurascens</i> [111] <i>S. racemosa</i> [329] <i>Schultesia lisianthoides</i> [328]
1-Hydroxy-2,3,7-trimethoxyxanthone (4) <i>F. carolinensis</i> [313] <i>Halenia elliptica</i> [38, 87] <i>Veratrilla baillonii</i> [348]		2-Hydroxy-1,3,7-trimethoxyxanthone (12) <i>Frasera albicaulis</i> [314]
1-Hydroxy-3,4,7-trimethoxyxanthone (6) <i>Frasera albicaulis</i> [314] <i>F. albomarginata</i> [90]		2-Hydroxy-5,6,7-trimethoxyxanthone (13) <i>Hypericum ericoides</i> [55–117]
1-Hydroxy-3,5,6-trimethoxyxanthone (7) <i>Canscora decussata</i> [108] <i>Centaurium erythraea</i> [190] <i>Hoppea dichotoma</i> [110]		3-Hydroxy-1,2,4-trimethoxyxanthone (14) <i>Psorospermum febrifugum</i> [138]
1-Hydroxy-3,5,8-trimethoxyxanthone (8) <i>Centaurium littorale</i> [333] <i>Swertia chirata</i> [112–161] <i>S. hookeri</i> [106] <i>S. japonica</i> [189] <i>S. lawii</i> [105] <i>S. paniculata</i> [13] <i>S. perfoliata</i> [145]		3-Hydroxy-1,2,7-trimethoxyxanthone (15) <i>Polygala tenuifolia</i> [100]
1-Hydroxy-3,6,7-trimethoxyxanthone (9) <i>Polygala tenuifolia</i> [174]		3-Hydroxy-1,5,6-trimethoxyxanthone (16) <i>Haploclathra leiantha</i> [256] <i>Kilmeyera rupestris</i> [74, 91]
1-Hydroxy-3,6,8-trimethoxyxanthone (10) (Ellipticol) <i>Halenia elliptica</i> [88]		3-Hydroxy-1,7,8-trimethoxyxanthone (17) <i>Swertia paniculata</i> [13]
1-Hydroxy-3,7,8-trimethoxyxanthone (11) (Decussatin) <i>Anthocleista vogelli</i> [267] <i>Canscora decussata</i> [66] <i>Centaurium cachanlahuen</i> [335]		4-Hydroxy-2,3,6-trimethoxyxanthone (18) <i>Hypericum reflexum</i> [53]
		5-Hydroxy-1,2,3-methoxyxanthone (19) <i>Schultesia lisianthoides</i> [328]

Table 2 (continued)

6-Hydroxy-1,3,5-trimethoxyxanthone (20)	1,5-Dihydroxy-3,8-dimethoxyxanthone (38) (Chiratol)
<i>Calophyllum austroindicum</i> [172]	<i>Swertia chirayita</i> [22]
7-Hydroxy-1,2,3-trimethoxyxanthone (21) (Onjixanthone I)	1,5-Dihydroxy-6,7-dimethoxyxanthone (39)
<i>Polygala tenuifolia</i> [100, 174]	<i>Carapa densiflora</i> [215]
7-Hydroxy-1,3,8-trimethoxyxanthone (22)	<i>C. grandifolia</i> [104]
(Anthaxanthone)	<i>C. psidifolia</i> [104]
<i>Haploclathra leiantha</i> [257]	<i>C. valioi</i> [104]
8-Hydroxy-1,3,5-trimethoxyxanthone (23)	<i>Tovomita brasiliensis</i> [46]
<i>Swertia bimaculata</i> [113]	
<i>S. mussotii</i> [321]	
1,2-Dihydroxy-5,6-dimethoxyxanthone (24)	1,6-Dihydroxy-3,5-dimethoxyxanthone (40)
<i>Garcinia subelliptica</i> [247]	<i>Canscora decussata</i> [108]
1,3-Dihydroxy-2,5-dimethoxyxanthone (25)	<i>Centaurium linearifolium</i> [273, 274]
<i>Bonnetia dinizii</i> [270]	<i>Chironia krebsei</i> [342]
<i>Calophyllum apetalum</i> [165]	<i>Hoppea dichotoma</i> [110]
<i>Centaurium linearifolium</i> [114]	
<i>Monnieria obtusifolia</i> [283]	
1,3-Dihydroxy-2,7-dimethoxyxanthone (26)	1,6-Dihydroxy-3,7-dimethoxyxanthone (41)
<i>Veratrilla baillonii</i> [348]	<i>Iris nigricans</i> [11]
1,3-Dihydroxy-2,8-dimethoxyxanthone (27)	<i>Polygala tenuifolia</i> [100–174]
<i>Kielmeyera candidissima</i> [98]	
1,3-Dihydroxy-4,5-dimethoxyxanthone (28)	1,6-Dihydroxy-5,7-dimethoxyxanthone (42)
<i>Frasera carolinensis</i> [313]	<i>Carapa densiflora</i> [214]
<i>F. speciosa</i> [90]	<i>Hypericum canariensis</i> [54]
<i>Swertia bimaculata</i> [113]	
1,3-Dihydroxy-4,7-dimethoxyxanthone (29)	1,6-Dihydroxy-7,8-dimethoxyxanthone (43)
<i>Frasera albicaulis</i> [314]	<i>Bonnetia dinizii</i> [270]
1,3-Dihydroxy-5,6-dimethoxyxanthone (30) (Leiaxanthone)	<i>Calophyllum trapezifolium</i> [128]
<i>Canscora decussata</i> [107]	<i>Polygala nyikensis</i> [236]
<i>Centaurium linearifolium</i> [116, 272]	
<i>Haploclathra leiantha</i> [256]	
1,3-Dihydroxy-5,8-dimethoxyxanthone (31)	1,6-Dihydroxy-7,8-dimethylenedioxyxanthone (44)
<i>Swertia petiolata</i> [207]	<i>Carapa densiflora</i> [215, 214]
1,3-Dihydroxy-6,7-dimethoxyxanthone (32)	1,7-Dihydroxy-2,3-dimethoxyxanthone (45)
(Laxanthone I)	<i>Frasera speciosa</i> [90]
<i>Lawsonia inermis</i> [42, 229]	<i>Halenia corniculata</i> [266–296]
1,3-Dihydroxy-7,8-dimethoxyxanthone (33) (Swertinin)	<i>H. elliptica</i> [322]
<i>Gentiana kochiana</i> [292]	<i>Polygala tenuifolia</i> [174, 100]
<i>Swertia decussata</i> [78]	
<i>Carapia inophyllum</i> [171]	
1,4-Dihydroxy-3,5-dimethoxyxanthone (34)	1,7-Dihydroxy-3,4-dimethoxyxanthone (46)
<i>Centaurium erythraea</i> [261]	<i>Polygala virgata</i> [32]
1,5-Dihydroxy-2,3-dimethoxyxanthone (35)	1,7-Dihydroxy-3,5-dimethoxyxanthone (47)
<i>Calophyllum thwaitesii</i> [77]	<i>Hoppea fastigiata</i> [252]
<i>Calophyllum walkeri</i> [77]	
<i>Halenia elliptica</i> [322]	1,7-Dihydroxy-3,6-dimethoxyxanthone (48)
<i>H. corniculata</i> [296]	<i>Calophyllum inophyllum</i> [208]
<i>Monnieria sylvatica</i> [31]	
<i>Schultesia lisianthoides</i> [328]	1,7-Dihydroxy-3,8-dimethoxyxanthone (49) (Gentiacaulein)
1,5-Dihydroxy-3,4-dimethoxyxanthone (36) (Tovopyrifolin-B)	<i>Gentiana barbata</i> [285]
<i>Tovomita macrophylla</i> [245]	<i>G. bavarica</i> [156, 158]
<i>T. pyrifolium</i> [245, 47]	<i>G. brachiphylla</i> [155, 156]
1,5-Dihydroxy-3,7-dimethoxyxanthone (37)	<i>G. ciliata</i> [239]
<i>Hoppea fastigiata</i> [254]	<i>G. detonsa</i> [187]
	<i>G. favratii</i> [156]
	<i>G. karelinii</i> [50]
	<i>G. kochiana</i> [292]
	<i>G. nivalis</i> [155, 156]
	<i>G. rostanii</i> [156]
	<i>G. schleicheri</i> [156]
	<i>G. utriculosa</i> [156]
	<i>G. verna</i> [156]
	<i>Gentianopsis pludosa</i> [349]
	<i>Haploclathra leiantha</i> [256]
	<i>H. paniculata</i> [259]
	<i>Swertia dilatata</i> [329]
	<i>S. gracilescens</i> [329]
	<i>S. nervosa</i> [329]
	<i>S. perenis</i> [154]
	<i>S. punicea</i> [101]

Table 2 (continued)

<i>S. racemosa</i> [329]	<i>S. punicea</i> [150]
<i>S. speciosa</i> [238]	<i>S. racemosa</i> [329]
1,8-Dihydroxy-2,6-dimethoxyxanthone (50)^a	
<i>Centaurium erythraea</i> [327]	<i>S. speciosa</i> [238]
<i>C. linearifolium</i> [274]	1,8-dihydroxy-4,6-dimethoxyxanthone (54)^a
1,8-Dihydroxy-3,5-dimethoxyxanthone (51)^a	
(Methylbellidifolin or Swerchirin)	
<i>Blackstonia perfoliata</i> [334]	2,4-dihydroxy-3,6-dimethoxyxanthone (55)
<i>Centaurium cahanlahuen</i> [335]	<i>Hypericum reflexum</i> [53]
<i>C. erythraea</i> [35, 190]	2,5-dihydroxy-1,6-dimethoxyxanthone (56)
<i>C. littorale</i> [35, 333]	<i>Garcinia thwaitesii</i> [135]
<i>C. pulchellum</i> [246]	2,6-dihydroxy-1,5-dimethoxyxanthone (57)
<i>Frasera albicaulis</i> [314]	<i>Garcinia sublliptica</i> [247]
<i>F. albomarginata</i> [90]	2,7-Dihydroxy-1,8-dimethoxyxanthone (58)
<i>F. carolinensis</i> [313]	<i>Cratoxylum formosanum</i> [166]
<i>Gentiana algida</i> [51]	3,8-Dihydroxy-1,2-dimethoxyxanthone(59)
<i>Gentiana bellidifolia</i> [233–252]	<i>Cratoxylum formosanum</i> [166]
<i>G. karelinii</i> [326]	<i>Calophyllum apetalum</i> [165]
<i>G. lactea</i> [302]	3,7-Dihydroxy-1,8-dimethoxyxanthone (60)
<i>Saponaria vaccaria</i> [192]	<i>Haploclathra paniculata</i> [259]
<i>Swertia bimaculata</i> [113]	3,6-Dihydroxy-1,5-dimethoxyxanthone (61)
<i>S. chirata</i> [79, 112]	<i>Calophyllum austroindicum</i> [172]
<i>S. chirayita</i> [22]	3,8-Dihydroxy-1,7-dimethoxyxanthone (62)
<i>S. dilatata</i> [329]	(Isogentiacaulein)
<i>S. gracilescens</i> [329]	<i>Gentiana kochiana</i> [136, 292]
<i>S. japonica</i> [180–189]	<i>Haploclathra paniculata</i> [259]
<i>S. milensis</i> [150]	4,6-Dihydroxy-1,3-dimethoxyxanthone (63)
<i>S. mussotii</i> [321]	<i>Alnus glutinosa</i> [197]
<i>S. nervosa</i> [329]	5,6-Dihydroxy-1,3-dimethoxyxanthone (64) (Ferrxanthone)
<i>S. paniculata</i> [13]	<i>Haploclathra leiantha</i> [256]
<i>S. patens</i> [219]	<i>H. piniculata</i> [259]
<i>S. petiolata</i> [195]	<i>Mesua ferrea</i> [337]
<i>S. racemosa</i> [329]	1,2,3-Trihydroxy-5-methoxyxanthone (65)
<i>S. speciosa</i> [238]	<i>Centaurium erythraea</i> [162–243]
<i>S. tetrapetala</i> [7]	1,2,8-Trihydroxy-3-methoxyxanthone (66)^b
1,8-Dihydroxy-3,6-dimethoxyxanthone (52)	
<i>Diploschistes sp</i> [95]	<i>Archytaea multiflora</i> [206]
1,8-Dihydroxy-3,7-dimethoxyxanthone (53)^a	
(Methylswertianin or Swertiaperenmin)	
<i>Blackstonia perfoliata</i> [334]	1,3,5-Trihydroxy-2-methoxyxanthone (67)
<i>Canscora decussata</i> [66]	(Tovopyrifolin-C)
<i>C. cahanlahuen</i> [335]	<i>Bonnetia dinizii</i> [270]
<i>C. pulchellum</i> [246]	(Haploxanthone)
<i>Gentiana ciliata</i> [239]	<i>Calophyllum apetalum</i> [165]
<i>G. karelinii</i> [50]	<i>Calophyllum bracteatum</i> [312]
<i>G. tenella</i> [52]	<i>C. inophyllum</i> [171]
<i>Schultesia lisianthoides</i> [328]	<i>C. tomentosum</i> [28]
<i>Swertia chirata</i> [112]	<i>C. trapezifolium</i> [128]
<i>S. chirayita</i> [22]	<i>Centaurium linearifolium</i> [114]
<i>S. dilatata</i> [329]	<i>Kayeaa stylosa</i> [131]
<i>S. erythrosticta</i> [160]	<i>Monnieria obtusifolia</i> [283]
<i>S. gracilescens</i> [329]	<i>Pentadesma butyracea</i> [81]
<i>S. iberica</i> [85]	<i>Tovomita pyrifolium</i> [245]
<i>S. japonica</i> [180–189]	1,3,5-Trihydroxy-6-methoxyxanthone (68)
<i>S. lawii</i> [105]	<i>Canscora decussata</i> [66–107]
<i>S. milensis</i> [150]	<i>Haploclathra leiantha</i> [256]
<i>S. mussotii</i> [321]	<i>Hoppea dichotoma</i> [110]
<i>S. nervosa</i> [329]	
<i>S. paniculata</i> [13]	
<i>S. patens</i> [149]	
<i>S. perennis</i> [154, 292]	

Table 2 (continued)

1,3,6-Trihydroxy-5-methoxyxanthone (69)	(Gentiakochianin or Swertianin)
<i>Canscora decussata</i> [107, 108]	<i>Chironia krebsii</i> [342]
<i>Hophea dichotoma</i> [110]	<i>G. lactea</i> [302]
1,3,6-Trihydroxy-7-methoxyxanthone (70)	<i>G. ramosa</i> [186]
<i>Poeciloneuron pauciflorum</i> [330]	<i>G. strictiflora</i> [187]
1,3,7-Trihydroxy-6-methoxyxanthone (71) (Isoathyriol)	<i>Swertia chirata</i> [112–161]
<i>Athyrium mesosorum</i> [331,332]	<i>S. dilatata</i> [329]
1,3,7-Trihydroxy-8-methoxyxanthone (72)^b	<i>S. erythrosticta</i> [160]
<i>Hophea fastigiata</i> [253]	<i>S. gracilescens</i> [329]
<i>Haploclathra leiantha</i> [259]	<i>S. hookeri</i> [106]
<i>H. paniculata</i> [259]	<i>S. japonica</i> [180–188]
1,3,8-Trihydroxy-5-methoxyxanthone (73)	<i>S. nervosa</i> [329]
(Isobellidifolin or Swertianol)	<i>S. paniculata</i> [13]
<i>Gentiana algida</i> [326]	<i>S. perennis</i> [154]
<i>Gentiana bellidifolia</i> [52-234]	<i>S. perfoliata</i> [145]
<i>G. campestris</i> [52]	<i>S. purpurescens</i> [8–111]
<i>G. germanica</i> [52]	<i>S. racemosa</i> [329]
<i>G. karelinii</i> [50]	<i>S. speciosa</i> [90]
<i>G. ramosa</i> [52]	1,6,7-Trihydroxy-3-methoxyxanthone (82)^b (Athyriol)
<i>G. tenella</i> [52]	<i>Athyrium mesosorum</i> [331, 332]
<i>Swertia chirata</i> [112]	1,6,8-Trihydroxy-2-methoxyxanthone (83)
<i>S. erythrosticta</i> [160]	<i>Iris nigricans</i> [11]
<i>S. hookeri</i> [106]	1,7,8-Trihydroxy-3-methoxyxanthone (84)
<i>S. japonica</i> [19]	1,7,8-Trihydroxy-6-methoxyxanthone (85)
<i>S. mussotii</i> [321]	<i>Archytaea multiflora</i> [206]
<i>S. paniculata</i> [13]	2,4,5-Trihydroxy-1-methoxyxanthone (86)
<i>S. purpurascens</i> [111]	<i>Gentiana bavarica</i> [156–158]
<i>Schultesia lisianthoides</i> [328]	<i>G. brachyphylla</i> [155, 156]
1,3,8-Trihydroxy-7-methoxyxanthone (74)	<i>G. ciliata</i> [239]
(Isogentiakochianin)	<i>G. favratii</i> [156]
<i>Calophyllum inophyllum</i> [171]	<i>G. karelinii</i> [50]
<i>Canscora decussata</i> [66]	<i>G. kochiana</i> [136–292]
<i>G. Cerastioides</i> [187]	<i>G. nivalis</i> [155, 156]
<i>G. Germanica</i> [186]	<i>G. rostani</i> [156]
<i>Kielmeyera speciosa</i> [120]	<i>G. verna</i> [153, 156]
<i>Swertia iberica</i> [85]	<i>Orphium frutescens</i> [297]
<i>Schultesia lisianthoides</i> [328]	<i>Swertia calycina</i> [295]
1,4,7-Trihydroxy-3-methoxyxanthone (75)	<i>S. decussata</i> [292]
<i>Garcinia eugenifolia</i> [171]	<i>S. dilatata</i> [78]
1,4,7-Trihydroxy-8-methoxyxanthone (76)	<i>S. erythrosticta</i> [160]
<i>Cratoxylum formosanum</i> [166]	<i>S. gracilescens</i> [329]
1,4,8-Trihydroxy-3-methoxyxanthone (77)^b	<i>S. iberica</i> [85]
<i>Gentiana bellidifolia</i> [235]	<i>S. japonica</i> [21–33]
1,4,8-Trihydroxy-6-methoxyxanthone (78)	<i>S. lawii</i> [105]
<i>Swertia japonica</i> [33]	<i>S. mussotii</i> [321]
1,5,6-Trihydroxy-3-methoxyxanthone (79)	<i>S. nervosa</i> [329]
<i>Canscora decussata</i> [109]	<i>S. racemosa</i> [329]
<i>Hypericum androsaemum</i> [263]	<i>S. speciosa</i> [238]
<i>Swertia chirata</i> [231]	(BR-xanthone-B)
<i>Schultesia lisianthoides</i> [328]	<i>Garcinia mangostana</i> [26]
1,5,7-Trihydroxy-3-methoxyxanthone (80)^b	3,5,6-Trihydroxy-1-methoxyxanthone (87)
<i>Hophea fastigiata</i> [251]	<i>Calophyllum sclerophyllum</i> [182]
1,5,8-Trihydroxy-3-methoxyxanthone (81) (Bellidifolin)	1,3,5,6-Tetrahydroxyxanthone (88)
<i>Gentiana algida</i> [326]	<i>Calophyllum brasiliense</i> [198]
<i>G. bellidifolia</i> [233]	<i>C. sclerophyllum</i> [182]
<i>G. campestris</i> [188]	<i>Canscora decussata</i> [66, 107]
	<i>Cratoxylum cochinchinense</i> [310]
	<i>Hypericum androsaemum</i> [263]
	<i>Hypericum patulum</i> [178]

Table 2 (continued)

<i>Mammea africana</i> [56]	<i>Symphonia globulifera</i> [220, 222]
<i>Morus tinctoria</i> [185]	1,3,7,8-Tetrahydroxyxanthone (92) (Norswertianin)
<i>Ochrocarpus odoratus</i> [227]	<i>Canscora decussata</i> [66]
<i>Psorospermum febrifugum</i> [4]	<i>Chironia krebsei</i> [342]
<i>Sympmania globulifera</i> [220–222]	<i>Gentiana bavarica</i> [158]
1,3,5,7-Tetrahydroxyxanthone (89)	<i>Orphium frutescens</i> [297]
<i>Garcinia pedunculata</i> [289]	<i>Swertia chirata</i> [112]
1,3,5,8-Tetrahydroxyxanthone (90) (Demethylbellidifolin)	<i>S. dilatata</i> [329]
<i>Gentiana bellidifolia</i> [52–233]	<i>S. erythrosticta</i> [160]
<i>G. lactea</i> [302]	<i>S. gracilis</i> [329]
<i>G. ramosa</i> [52–186]	<i>S. hookeri</i> [106]
<i>G. strictiflora</i> [187]	<i>S. iberica</i> [85]
<i>Iris nigricans</i> [11]	<i>S. japonica</i> [21, 189]
<i>Swertia chirata</i> [112]	<i>S. lawii</i> [105]
<i>S. dilatata</i> [329]	<i>S. nervosa</i> [329]
<i>S. erythrosticta</i> [160]	<i>S. perennis</i> [154]
<i>S. gracilis</i> [329]	<i>S. purpurascens</i> [111]
<i>S. hookeri</i> [106]	<i>S. racemosa</i> [329]
<i>S. japonica</i> [19, 21, 189]	1,7-Dimethoxy-2,3-methylenedioxylxanthone (93)
<i>S. lawii</i> [105]	<i>Polygala tenuifolia</i> [100]
<i>S. macroperma</i> [350]	1,8-Dimethoxy-2,3-methylenedioxylxanthone (94)
<i>S. nervosa</i> [329]	<i>Polygala nyikensis</i> [236]
<i>S. purpureascens</i> [111]	1,2,3,7-Tetramethoxyxanthone (95)
<i>S. racemosa</i> [329]	<i>Frasera albicaulis</i> [314]
1,3,6,7-Tetrahydroxyxanthone (91) (Norathyriol)	<i>Polygala tenuifolia</i> [174–181]
<i>Allanblackia floribunda</i> [227]	1,2,6,8-Tetramethoxyxanthone (96)
<i>Athyrium mesosorum</i> [265, 332]	<i>Bredemeyera brevifolia</i> [268]
<i>Canscora decussata</i> [109]	1,3,4,5-Tetramethoxyxanthone (97)
<i>Chlorophora tinctoria</i> [122]	<i>Frasera albicaulis</i> [314]
<i>Cratoxylum pruniflorum</i> [199]	1,3,4,7-Tetramethoxyxanthone (98)
<i>Garcinia echinocarpa</i> [27]	<i>Frasera albicaulis</i> [314]
<i>G. mangostana</i> [152]	1,3,5,8-Tetramethoxyxanthone (99)
<i>G. multiflora</i> [70]	<i>Swertia hookeri</i> [106]
<i>G. pedunculata</i> [289]	1,3,6,7-Tetramethoxyxanthone (100)
<i>G. terpnophylla</i> [27]	<i>Allanblackia floribunda</i> [226]
<i>Hypericum androsaemum</i> [263]	<i>Polygala tenuifolia</i> [100]
<i>H. aucheri</i> [200]	2,3,4,5-Tetramethoxyxanthone (101)
<i>Iris nigricans</i> [11]	<i>Halenia campanulata</i> [161]
<i>Maclura pomifera</i> [343]	
<i>Mammea africana</i> [56]	
<i>Morus tinctoria</i> [185]	
<i>Ochrocarpus odoratus</i> [227]	
<i>Pentaphalangium solomonse</i> [271]	

^a The compounds 51/53, 51/54, 66/84, 72/82, 77/80, have the same structure, but they are cited under different names in the references.

^b The compounds 51/53, 51/54, 66/84, 72/82, 77/80, have the same structure, but they are cited under different names in the references.

Clusiaceae, Bennett and Lee (1989) listed 50 tetraoxygenated xanthones from Clusiaceae species. A review by Mandal et al. (1992) 113 simple tetraoxygenated, prenylated and related xanthones are listed. Between 1942 and mid 1995, approx. 115 tetraoxygenated naturally occurring xanthones were presented in reviews. In the present review, 228 of these compounds found in the literature are listed (Tables 2–6).



7. Synthesis and biosynthesis of xanthones

Some authors have reported the preparation of hydroxyxanthones (Atkinson and Heilbron, 1926; Grover


et al., 1955; Shah and Shah, 1956; Quillinan and Scheinmann, 1972, 1973; Simoneau and Brassard, 1984), but the first xanthone synthesis was proposed by Kostanecki (1892) and the last by Ravi et al. (1994) describing a new route to xanthone synthesis, and Vitale et al. (1994) also describing a novel route for the preparation of xanthones and chromanones.

The preparation of nine 1,2,3,8-tetraoxygenated xanthones was proposed by Gil et al. (1990) and 1,3,5,6-, 3,4,5,6-, 3,4,6,7- and 2,3,6,7-tetrahydroxyxanthone were synthesized from benzophenone precursors (Sundholm, 1978). The total synthesis of lichen xanthones (Patel and Trivedi, 1983), synthesis of furanoxanthones (Pinto and Polonia, 1974; Patel and Trivedi, 1991), synthesis of new


Table 3

Mono C₅ tetraoxxygenated naturally occurring xanthones**1,3,5,6-Tetrahydroxy-2-(3-methylbut-2-enyl)xanthone (102)***Calophyllum brasiliensis* [72]*C. austroindicum* [172]*C. canum* [57]*C. neo-ebudicum* [303]*C. sclerophyllum* [182]*C. scribilitifolium* [183]*C. fragrans* [224]*C. inophyllum* [10, 184](102 R = H) (103 R = CH₃)**1-Hydroxy-3,5,6-trimethoxy-2-(3-methylbut-2-enyl)xanthone (103)***Calophyllum ramiflorum* [39]**Paxanthone (104)***Hypericum patulum* [177]


(104)

Forbexanthone (105)*Garcinia forbesii* [147]


(105)

Toxyloxanthone C (106)*Calophyllum zeylanicum* [133]*Kielmeyera ferruginea* [121]*Maclura pomifera* [86]


(106 R=H; 107 R=Me)

Geronthoxanthone J (107)*Cudrania cochinchinensis* [64]**2-Deprenylrheediaxanthone B (108)***Hypericum roeperanum* [290]


(108 R=H; 109 R=Me)

5-Methoxy-2-deprenylrheediaxanthone B (109)*Hypericum roeperanum* [290]**5-O-Methylisojacareubin (110)**


(110)

Lorostemin A (111)*Lorostemon coelholi* [48]*L. negrensis* [48]


(111)

1,3,5-Trihydroxy-6',6'-dimethylpyrano(2',3':6,7)xanthone (112)*Rheedia brasiliensis* [82]


(112)

Morusignin D (113)*Morus insignis* [144]*Hypericum patulum* [179]


(113)

Toxyloxanthone-B (114)*Hypericum androsaemum* [263]*H. paturum* [177]*H. sampaonii* [71]*Maclura pomifera* [76–86]*Morus insignis* [144]


(114)

1,4,6-Trihydroxy-5-methoxy-7-(3-methylbut-2-enyl)xanthone (115)*Garcinia dulcis* [148]

(115)

Cudraniaxanthone (116)*Cudrania cochinchinensis* [62]*Cudrania javanensis* [255]

(116)

Jacareubin (117)*Calophyllum austroindicum* [172]*C. bracteatum* [312]*C. brasiliense* [72, 280]*C. calaba* [209, 312]*C. canum* [57]*C. cordato-oblongum* [134]*C. cuneifolium* [130]*C. fragrans* [224]*C. inophyllum* [10–184, 208]*C. neo-ebudicum* [303]*C. ramiflorum* [39]*C. sclerophyllum* [182]*C. scribilitifolium* [183]*C. thwaitesii* [77]*C. tomentosum* [191]*C. trapezifolium* [128]*C. walkerii* [77]*C. zeylanicum* [133]*Kielmeyera ferruginea* [121]*Pentadesma butyracea* [81]

(117)

Table 3 (continued)














Isojacareubin (118)*Hypericum roeperanum* [290]**1,5,8-Trihydroxy-3-methoxy-2-(3-methylbut-2-enyl)xanthone (119)***Garcinia mangostana* [275, 276]**Morusignin C (120)***Morus insignis* [144]**1,6,7-Trihydroxy-4,4,5-trimethyldihydrofuran-(2,3;3,4) xanthone (121)***Allanblackia floribunda* [226]**Celebixanthone (122)***Cratoxylum celebicum* [317, 318]**Symphoxanthone (123)***Garcinia subelliptica* [248]*Symphoxanthone (123)* [*Garcinia subelliptica*] [223]**1-Methylsymphoxanthone (124)***Garcinia subelliptica* [248]**1,3,5,6-Tetrahydroxy-2-(3-hydroxy-3-methylbutyl)xanthone (125)***Calophyllum inophyllum* [119]**Ugaxanthone (126)***Symphoxanthone (126)* [*Symphoxanthone*] [221]**Morusignin B (127)***Morus insignis* [144]**Morusignin A (128)***Morus insignis* [144]**1,3,6,7-Tetrahydroxy-4-(1,1-dimethylallyl)xanthone (129)***Allanblackia floribunda* [226]**1,3,6,7-Tetrahydroxy-8-(3-methylbut-2-enyl)xanthone (130)***Hypericum androsaemum* [176–263]**Subelliptenone F (131)***Garcinia subelliptica* [168]*Garcinia dulcis* [166]

Table 4
Di-C₅ tetraoxxygenated naturally occurring xanthones
















Tovopyrifolin D (132) <i>Tovomita pyrifolium</i> [47]		10-O-methylmacluraxanthone (140) <i>Kayea stylosa</i> [131]	
Kayeaxanthone (133) <i>Kayeaxanthone</i> [131]		Gerontoxanthone E (141) <i>Cudrania cochinchinensis</i> [62]	
Pyranojacareubin (134) <i>Garcinia densivenia</i> [339] <i>G. forbesii</i> [147] <i>Rheedia brasiliensis</i> [82] <i>R. gardneriana</i> [83, 84]		Caloxanthone B (142) <i>Carapia inophyllum</i> [173]	
Padiaxanthone (135) <i>Hypericum patulum</i> [175]		Tovopyrifolin A (143) <i>Tovomita pyrifolium</i> [47, 245]	
Rheediaxanthone A (136) <i>Garcinia staudtii</i> [340] <i>Rheedia benthamiana</i> [80] <i>R. brasiliensis</i> [82] <i>R. gardneriana</i> [83]		3-Isomangostin (144) <i>Garcinia mangostana</i> [228, 305]	
Gerontoxanthone A (137) <i>Cudrania cochinchinensis</i> [64] <i>Rheedia brasiliensis</i> [82]		3-Isomangostin hydrate (145) <i>Garcinia mangostana</i> [228]	
1,5-Dihydroxy-6,6-dimethylpyrano(2,3:6,7)-6'',6''-dimethyldihydropyrano(2'',3'':3,4)xanthone (138) <i>Rheedia brasiliensis</i> [82]		β-Mangostin (146) <i>Cratoxylum cochinchinense</i> [36] <i>Garcinia mangostana</i> [20, 126, 228, 298]	
1,5-Dihydroxy-6,6-dimethylpyrano(2,3:6,7)-6'',6''-dimethyldihydropyrano(2'',3'':2,3)xanthone (139) <i>Rheedia brasiliensis</i> [82]			

Table 4 (continued)


















BR-xanthone A (147)*Garcinia mangostana* [26]**Manglexanthone (148)***Tovomita mangle* [237]**Caloxanthone D (149)***Calophyllum inophyllum* [171]**3,5-Dihydroxy-6,6-dimethylpyrano(2,3:6,7)-6'',6''-dimethyldihydroxanthone (150)***Rheedia brasiliensis* [82]**1-Isomangostin (151)***Garcinia mangostana* [228]**11-Hydroxy-1-isomangostin (152)***Cratoxylum cochinchinense* [310]**1-Isomangostin hydrate (153)***Garcinia mangostana* [228]**Rheediaxanthone C (154)***Rheedia brasiliensis* [82]*R. gardneriana* [83]*R. benthamiana* [80]**Subelliptenone H (155)***Garcinia subelliptica* [169]**Gerontoxanthone B (156)***Cudrania cochinchinensis* [64]**1,3,5-Trihydroxy-6,6-dimethylpyrano(2,3:6,7)-4-(1,1-dimethylallyl)xanthone (157)***Rheedia brasiliensis* [82]*Garcinia opaca* [118]**Cudraxanthone F (158)***Cudrania tricuspidata* [142]**Cudraxanthone E (159)***Cudrania tricuspidata* [142]**Garcinone D (160)***Cratoxylum cochinchinense* [36]*Garcinia mangostana* [307]**Cratoxylene (161)***Cratoxylum cochinchinense* [36]**Cudraxanthone C (162)***Cudrania tricuspidata* [141]

Table 4 (continued)


1,3,6-Trihydroxy-7-methoxy-2,5-bis(3-methyl-but-2-enyl)xanthone (163)
Garcinia cowa [280]



(163)

Mangostin (164)

Cratoxylum cochinchinense [36]
Garcinia echinocarpa [27]
G. mangostana
[20, 92, 126–228, 298, 316]
G. terpnophylla [27]




(164 R = Me; 165 R = H)

 γ -Mangostin or Normangostin (165)

Garcinia mangostana
[20, 126, 228, 298]
Hypericum patulum [177]
Garcinia mangostana [260]

Mangostanol (166)


Garcinia mangostana [260]



(166)

Garcinone B (167)


Garcinia mangostana [298, 306, 308]
Hypericum patulu [176]
Morus insignis [144]



(167)

Cudraxanthone K (168)


Cudrania tricuspidata [141]



(168)

Cudraxanthone B (169)


Cudrania tricuspidata [264]



(169)

Paxanthone B (170)


Hypericum patulum [176]



(170)

Subelliptenone I (171)


Garcinia subelliptica [169]



(171)

Macluraxanthone (172)


Calophyllum inophyllum [173]
Garcinia opaca [118]
G. ovalifolia [338]
Maclura aurantica [17]
M. pomifera [344, 345]
Rheedia benthamiana [80]
R. brasiliensis [82]
R. gardneriana [83]



(172)

Xanthone V₁ (173)


Vismia guineensis [45]



(173)

7-Prenyljacareubin (174)


Rheedia gardneriana [83]



(174)

Rheediaxanthone B (175)


Garcinia polyantha [12]
Rheedia benthamiana [80]
R. brasiliensis [82]
R. gardneriana [83]



(175)

Isorheediaxanthone B (176)


Garcinia polyantha [12]
Rheedia brasiliensis [82]
R. gardneriana [83]



(176)

Gerontoxanthone C (177)


Cudrania cochinchinensis [64]



(177)

Gerontoxanthone G (178)


Cudrania cochinchinensis [62]




(178)

Table 4 (continued)


Morusignin E (179)
Morus insignis [143]




Morusignin F (180)
Morus insignis [143]




Cudraxanthone M (181)
Cudrania tricuspidata [140]




Caloxanthone A (182)
Carapia inophyllum [173]




Subelliptenone B (183)
Garcinia subelliptica [163]




Gerontoxanthone I (184)
Cudrania cochinchinensis [62]




Xanthone V_{1a} (185)
Vismia guineensis [45]




1,3,5,6-Tetrahydroxy-4-(1,1-dimethylallyl)-7-(3-methylbut-2-enyl)xanthone (186)
Rheedia brasiliensis [82]




Toxyloxanthone-D (187)
Machura pomifera [86]




Alvaxanthone (188)
Maclura aurantica [17]
M. pomifera [344, 346]



Gartanin (189)
Garcinia mangostana
[20, 126, 228, 260, 275, 298]
Morus insignis [144]



Cudraxanthone L (190)
Cudrania tricuspidata [140]



Garcinone C (191)
Garcinia mangostana [306]






Table 5

Tri C₅ tetraoxxygenated naturally occurring xanthones


1,5-Dihydroxy-6,6-dimethylpyrano(2,3:6,7)-4'',4'',5''-trimethylfurano(2'',3':3,4)-2-(3-methylbut-2-enyl)xanthone (192)
Garcinia opaca [118]




Tovophyllin B (193)
Tovomita macrophylla [269]
T. pyrifolium [47, 245]




1,3,5-Trihydroxy-6,6-dimethylpyrano(2,3:6,7)-2-(3-methylbut-2-enyl)-4-(1,1-dimethylallyl)xanthone (194)
Garcinia opaca [118]




Tovophyllin A (195)
Cratoxylum cochinchinense [36]
Tovomita macrophylla [269]
T. pyrifolium [47, 245]




Pentadesmaxanthone (196)
Pentadesma butyracea [81]




Subelliptenone E (197)
Garcinia subelliptica [168]




Nervosaxanthone (198)
Garcinia nervosa [12]




Garcinone E (199)
Garcinia mangostana
[20, 92, 202, 260, 298]




7-O-methylgarcinone (200)
Garcinia cowa [202]



Subelliptenone A (201)
Garcinia subelliptica [163, 248]



Dulciol B (202)
Garcinia dulcis [166]



Dulciol C (203)
Garcinia dulcis [166]

Table 6

More complex tetraoxxygenated naturally occurring xanthones













Cadensin A (204)*Carapa densiflora* [59]*C. grandifolia* [104]*C. valloii* [104]*Kielmeyera coriacea* [59]*Vismia guaramirangae* [132]**Cadensin G (210)***Psorospermum febrifugum* [2]**Cadensin-B (205)***Carapa densiflora* [59]*Kielmeyera coriacea* [59]*Vismia guaramirangae* [132]**Sapxanthone (211)***Saponaria vaccaria* [193]**Cadensin C (206)***Vismia guaramirangae* [132]**Garciniaxanthone E (212)***Garcinia subelliptica* [248]**5-Methoxysterigmatocystin (207)***Aspergillus versicolor* [151]**6-Methoxysterigmatocystin (208)***Aspergillus versicolor* [49]**Roeperanone (214)***Hypericum roeperanum* [290]**5'-Demethoxycadensin G (209)***Cratoxylum cochinchinense* [310]**5-O-Demethylpaxanthonin (215)***Hypericum roeperanum* [290]*H. patulum* [175]

Table 6 (continued)














Vaccaxanthone (216)*Saponaria vaccaria* [192]**Isocowanin (222)***Garcinia pyrifera* [12]**1,2,5-Trihydroxy-4-(1,1-dimethylallyl)-furano(2,3:6,7)xanthone (217)***Garcinia subelliptica* [162]**Dulciol A (223)***Garcinia dulcis* [166]**Cowaxanthone (218)***Garcinia cowa* [280]**Isocowanol (224)***Garcinia pyrifera* [12]**Rubraxanthone (219)***Garcinia cowa* [211]*G. nervosa* [12]*G. pyrifera* [12]**Cowanin (220)***Garcinia cowa* [280]**Subelliptenone C (225)***Garcinia subelliptica* [162]*G. dulcis* [166]**Cowanol (221)***Garcinia cowa* [280]**Maculatoxanthone (226)***Hypericum maculatum* [14]

Table 6 (continued)


Norcowanin (227)
Garcinia cowa [280]




6-Methoxykielcorin (228)
Hypericum reflexum [53]




Isocadensin D (229)
Psorospermum febrifugum [2]




Cadensin F (230)
Psorospermum febrifugum [2]



Isocadensin D — monoacetate (231)
Psorospermum febrifugum [2]




Paxanthonin (232)
Hypericum patulum [176, 179]




1,2-Dihydro-3,6,8-trihydroxy-1,1-bis(3-methylbut-2-enyl)xanthen-2,9-dione (233)

Hypericum patulum [178]



Muraxanthone (234)
Seneico mikanioides [60]



xanthones (Patolia and Trivedi, 1983), a new one-step method (Atkinson and Lewis, 1969) and a general study for synthesis of dioxygenated xanthones are in a recent report by Lin et al. (1992).

The biosynthetic pathways to xanthones have been discussed in recent years. Initially, these attempted to interrelate the observed oxygen pattern of natural xanthones and correlate them with recognized oxygenation patterns. In general it seems that ring A and the attached CO group are provided by the shikimic acid pathway, whereas ring B arises via the acetate–malonate polyketide route (Locksley and Murray, 1970; Afzal and Al-Hassan, 1980; Sultanbawa, 1980). Therefore Locksley et al. (1967) reported the significance of maclurin in xanthone biosynthesis and the biogenetic-type synthesis of xanthones from their benzophenone precursors (Locksley and Murray, 1970). Gottlieb (Gottlieb and Magalhães, 1966; Gottlieb, 1968; Gottlieb

et al., 1968) offered biogenetic proposals regarding xanthones, and Bhanu et al. (1972) related the biogenetic implications in the conversion of 4-phenylcoumarins into xanthones. More recently; biosynthetic studies on tajixanthone and shamixanthone (Ahmed et al., 1992) were reported.

Some xanthones in lower plants have been proven to be totally acetate-derived, from seven acetate units (McMaster et al., 1960; Birch et al., 1976). However, the oxygenation patterns of all xanthones in higher plants suggest that these are formed by a mixed shikimate–acetate pathway. This involves the condensation of shikimate and acetate-derived moieties to form benzophenones or benzophenone-like intermediates, which then react intramolecularly to form xanthones. Mechanisms for this intramolecular reaction have been postulated involving either direct phenol oxidative coupling (Lewis, 1963), quinone addition (Ellis et al., 1967), dehydration

between hydroxyl groups on the acetate and shikimate-derived rings (Markham, 1965), or spirodienone formation and subsequent rearrangement to form the xanthone (Gottlieb, 1968; Carpenter et al., 1969).

Regarding lichen and fungi biosynthesis, it is important to say that many secondary metabolites found in lichen-forming fungi play a dominant role in the systematics of these organisms due to extensive morphological parallels and their clear ecological significance. Despite their common occurrence in a number of important genera, lichen xanthones have not featured prominently in the repertoire of lichen taxonomists. Nevertheless, with the availability of more sensitive methods of detection and synthetic materials for comparison, these compounds have been effectively employed in recent systematic studies of lichens (Leuckert et al., 1990; Elix and Crook, 1992; Elix et al., 1992).

Elix and Crook (1992) reported the unambiguous total synthesis of 17 chlorine-containing derivatives of norlichexanthone, achieved using the condensation of an appropriately substituted methyl or ethyl orsellinate and phloroglucinol or 2-chlorophloroglucinol in the key step. Previously, Elix et al. (1992) reported the synthesis of several trioxigenated lichen xanthones (Elix et al., 1984).

8. Conclusion

The majority of xanthones that have been isolated so far are of the tetraoxigenated-type. The main natural source of these xanthones are plants from the Gentianaceae (14 genus, 72 species), Clusiaceae (23 genus, 84 species), Moraceae (4 genus, 8 species) and Polygalaceae (3 genus, 7 species) families. Other families of higher plants that produce tetraoxigenated xanthones are Betulaceae, Caryophyllaceae, Gesneriaceae, Iridaceae, Loganiaceae, Lytracaeae and Polypodiaceae. Only three tetraoxigenated xanthones have been identified as fungal and lichen metabolites: two in *Aspergillus versicolor* and one in *Diploschistes* sp. Therefore, tetraoxigenated xanthones have no application in chemosystematics studies of these organisms.

Acknowledgements

Part of the PhD thesis of Valdir Peres, submitted at Universidade Federal de Minas Gerais, Belo Horizonte, 31270-901, Minas Gerais, Brasil.

References

- [1] Abou-Shoer, M., Boettner, F.E., Chang, C.J., Cassady, J.M., 1988. Phytochemistry 27, 2795.
- [2] Abou-Shoer, M., Habib, A.A., Chang, C.J., Cassady, J.M., 1989. Phytochemistry 28, 2483.
- [3] Abou-Shoer, M., Suwanborirux, K., Chang, C.J., Cassady, J.M., 1989. Tetrahedron Letters 30, 3385.
- [4] Abou-Shoer, M., Suwanborirux, K., Habib, A.A.M., Chang, C.J., Cassady, J.M., 1993. Phytochemistry 34, 1413.
- [5] Afzal, M., Al-Hassan, J.M., 1980. Heterocycles 14, 1173.
- [6] Afzal, M., Al-Hassan, J.M., Al-Masad, F.N., 1979. Heterocycles 12, 269.
- [7] Agata, I., Sekizaki, H., Sakushima, A., Nishibe, S., Hisada, S., Kimura, K., 1981. Yakugaku Zasshi 101, 1067.
- [8] Ahmad, S., Ikram, M., Khan, I., Galbraith, M.N., 1973. Phytochemistry 12, 2542.
- [9] Ahmed, S.A., Bardshiri, E., McIntyre, C.R., Simpson, T.J., 1992. Australian Journal of Chemistry 45, 249.
- [10] Al-Jeboury, F.S., Locksley, H.D., 1971. Phytochemistry 10, 603.
- [11] Al-Khalil, S., Tosa, H., Iinuma, M., 1995. Phytochemistry 38, 729.
- [12] Ampofo, S.A., Waterman, P.G., 1986. Phytochemistry 25, 2351.
- [13] Anand, S.M., Basumatary, P.C., Ghosal, S., Handa, S.S., 1982. Planta Medica 45, 61.
- [14] Arends, P., 1969. Tetrahedron Letters 55, 4893.
- [15] Arends, P., Helboe, P., 1972. Acta Chemica Scandinavica 26, 4180.
- [16] Arends, P., Helboe, P., Moller, J., 1973. Organic Mass Spectrometry 7, 667.
- [17] Artamonova, N.A., Nikonov, G.K., Kiseleva, V.V., 1989. Chemical Natural Compounds 3, 373.
- [18] Asahina, Y., Asano, J., Uyeno, Y., 1942. Bulletin of the Chemical Society Japan 17, 104.
- [19] Asahina, Y., Asano, J., Uyeno, Y., 1942. Journal of the Pharmaceutical Society Japan 62, 22.
- [20] Asai, F., Tosa, H., Tanaka, T., Iinuma, M., 1995. Phytochemistry 39, 943.
- [21] Ashida, S., Noguchi, S.F., Suzuki, T., 1994. Journal of the America Oil Chemical Society 71, 1095.
- [22] Asthana, R.K., Sharma, N.K., Kulshreshtha, D.K., Chatterjee, S.K., 1991. Phytochemistry 30, 1037.
- [23] Atkinson, H., Heilbron, I.M., 1926. Journal of the Chemical Society, 2688.
- [24] Atkinson, J.E., Lewis, J.R., 1969. Journal of the Chemical Society, 281.
- [25] Balachandran, S., Vishwakarma, R.A., Popli, S.P., Kapil, R.S., 1988. Indian Journal of Chemistry 27B, 385.
- [26] Balasubramanian, K., Rajagopalan, K., 1988. Phytochemistry 27, 1552.
- [27] Bandaranayake, W.M., Selliah, S.S., Sultanbawa, M.U.S., Ollis, W.D., 1975. Phytochemistry 14, 1878.
- [28] Banerji, A., Deshpande, A.D., Prabhu, B.R., Pradhan, P., 1994. Journal of Natural Products 57, 396.
- [29] Barnes, C.S., Ocelowitz, J.L., 1964. Australian Journal of Chemistry 17, 975.
- [30] Barraclough, D., Locksley, H.D., Scheinmann, F., Magalhães, M.T., Gottlieb, O.R., 1970. Journal of the Chemical Society B, 603.
- [31] Bashir, A., Hamburger, M., Gupta, M.P., Solis, P., Hostettmann, K., 1992. Phytochemistry 31, 3203.
- [32] Bashir, A., Hamburger, M., Msomthi, J.D., Hostettmann, K., 1992. Phytochemistry 31, 309.
- [33] Basnet, P., Kadota, S., Namba, T., 1994. Planta Medica 60, 507.
- [34] Basnet, P., Kadota, S., Shimizu, M., Takata, Y., Kobayashi, M., Namba, T., 1995. Planta Medica 61, 402.
- [35] Beerhues, L., Berger, U., 1994. Phytochemistry 35, 1227.
- [36] Bennett, G.J., Harrison, L.J., Sai, G.L., Sim, K.Y., 1993. Phytochemistry 32, 1245.
- [37] Bennett, G.J., Lee, H.H., 1989. Phytochemistry 28, 967.
- [38] Bennett, G.J., Lee, H.H., 1991. Phytochemistry 30, 1347.
- [39] Bhanu, S., Scheinmann, F., Jefferson, A., 1975. Phytochemistry 14, 298.

- [40] Bhanu, S., Saroja, T., Seshadri, T.R., Mukerjee, S.K., 1972. Indian Journal of Chemistry 10, 577.
- [41] Bhardwaj, D.K., Jain, R.K., Jain, B.C., Mehta, C.K., 1978. Phytochemistry 17, 1440.
- [42] Bhardwaj, D.K., Seshadri, T.R., Singh, R., 1977. Phytochemistry 16, 1616.
- [43] Bhattacharya, S.K., Ghosal, S., Chaudhuri, R.K., Sanyal, A.K., 1972. Journal of Pharmaceutical Science 61, 1838.
- [44] Birch, A.J., Baldas, J., Hlubucek, J.R., Simpson, T.J., Westerman, P.W., 1976. Journal of the Chemical Society Perkin Transactions I, 898.
- [45] Botta, B., Delle-Monache, G., Delle-Monache, F., Bettolo, G.B.M., Menichini, F., 1986. Phytochemistry 25, 1217.
- [46] Braz Filho, R., Miranda, C.A.S., Gottlieb, O.R., Magalhães, M.T., 1982. Acta Amazonica 12, 801.
- [47] Braz Filho, R., Costa, S.B.S., Rocha, A.I., 1983. Arquivo da Universidade Federal Rural do Rio de Janeiro 6, 31.
- [48] Braz Filho, R., Lemos, M.J.C., Gottlieb, O.R., 1973. Phytochemistry 12, 947.
- [49] Bullock, E., Kirkaldy, D., Roberts, J.C., Underwood, J.G., 1963. Journal of the Chemical Society, 829.
- [50] Butayarov, A.V., Batirov, E.K., Tadzhibaev, M.M., Malikov, V.M., 1993. Journal Natural Compounds 3, 469.
- [51] Butayarov, A.V., Batirov, E.K., Tadzhibaev, M.M., Ibragimov, E.E., Malikov, V.M., 1993. Khimiya Prirodnykh Soedinenii 6, 901.
- [52] Carbonnier, J., Massias, M., Molho, D., 1977. Bulletin of the Museum of National History Natural Paris 13, 23.
- [53] Cardona, M.L., Fernandez, M.I., Pedro, J.R., Serrano, A., 1990. Phytochemistry 29, 3003.
- [54] Cardona, M.L., Fernandez, M.I., Pedro, J.R., Seoane, E., Vidal, R., 1986. Journal of Natural Products 49, 95.
- [55] Cardona, M.L., Seoane, E., 1982. Journal of Natural Products 45, 134.
- [56] Carpenter, I., Locksley, H.D., Scheinmann, F., 1969. Journal of the Chemical Society C, 2421.
- [57] Carpenter, I., Locksley, H.D., Scheinmann, F., 1969. Journal of the Chemical Society C, 486.
- [58] Carpenter, I., Locksley, H.D., Scheinmann, F., 1969. Phytochemistry 8, 2013.
- [59] Castelão, J.F. Jr., Gottlieb, O.R., Lima, R.A., Mesquita, A.A.L., Gottlieb, H.E., Wenkert, E., 1977. Phytochemistry 16, 735.
- [60] Catalano, S., Luschi, S., Flamini, G., Luigi, P., Neri, E.M., Morelli, I., 1996. Phytochemistry 42 (6), 1605.
- [61] Chakravarty, A.K., Mukhopadhyay, S., Moitra, S.K., Das, B., 1994. Indian Journal of Chemistry 33B, 405.
- [62] Chang, C.H., Lin, C.C., Kawata, Y., Hattori, M., Namba, T., 1989. Phytochemistry 28, 823.
- [63] Chang, C.H., Lin, C.C., Hattori, M., Namba, T., 1989. Phytochemistry 28, 595.
- [64] Chang, C.H., Lin, C.C., Hattori, M., Namba, T., 1994. Journal of Ethnopharmacology 44, 79.
- [65] Chang, C.H., Lin, C.C., Kadota, S., Hattori, M., Namba, T., 1995. Phytochemistry 40, 945.
- [66] Chaudhuri, R.K., Ghosal, S., 1971. Phytochemistry 10, 2425.
- [67] Chaudhuri, R.K., Zymalkowski, F., Frahm, A.W., 1978. Tetrahedron 34, 1837.
- [68] Chawla, H.M., Chibber, S.S., Khera, U., 1975. Journal of Chromatography 111, 246.
- [69] Chawla, H.M., Chibber, S.S., Seshadri, T.R., 1973. Proceedings Indian Academic Science, 141
- [70] Chen, F.C., Lin, Y.M., Hung, J.C., 1975. Phytochemistry 14, 300.
- [71] Chen, M.T., Chen, C.M., 1985. Heterocycles 23, 2543.
- [72] Chilpa, R.R., Estrada, M.J., Muñiz, E.E., 1997. Journal of Chemical Ecology 23, 1901.
- [73] Chow, Y.L., Quon, H.H., 1968. Phytochemistry 7, 1871.
- [74] Correa, D.B., Silva, L.G.F., Gottlieb, O.R., Gonçalves, S.J., 1970. Phytochemistry 9, 447.
- [75] Cotterill, P.J., Scheinmann, F., 1975. Journal of the Chemical Society Chemical Communications, 664.
- [76] Cotterill, P.J., Scheinmann, F., 1980. Journal of the Chemical Society Perkin Transactions I, 2353.
- [77] Dahanayake, M., Kitagawa, I., Somanathan, R., Sultanbawa, M.U.S., 1974. Journal of the Chemical Society, Perkin Transactions I, 2510.
- [78] Dalal, M.S.R., Sethna, S., Shah, R.C., 1953. Journal of the Indian Chemical Society 30, 457.
- [79] Dalal, S.R., Shah, R.C., 1956. Chemistry and Industry 7, 664.
- [80] Delle-Monache, F., Botta, B., Nicoletti, M., Coelho, J.S.B., Lyra, F.D.A., 1981. Journal of the Chemical Society Perkin Transactions I, 484.
- [81] Delle-Monache, F., Mac-Quhae, M.M., Delle-Monache, G., Bettolo, G.B.M., Lima, R.A., 1983. Phytochemistry 22, 227.
- [82] Delle-Monache, G., Botta, B., Mello, J.F., Coelho, J.S.B., Menichini, F., 1984. Journal of Natural Products 47, 620.
- [83] Delle-Monache, G., Delle-Monache, F., Bettolo, G.B.M., Lima, R.A., 1983. Journal of Natural Products 46, 655.
- [84] Delle-Monache, G., Delle-Monache, F., Waterman, P.G., Crichton, E.G., Lima, R.A., 1984. Phytochemistry 23, 1757.
- [85] Denisova, O.A., Glyzin, V.I., Patudin, A.V., Fesenko, D.A., 1980. Chemical and Natural Compounds 2, 145.
- [86] Deshpande, V.H., Rao, A.V.R., Varadarajan, M., Venkataraman, K., 1973. Indian Journal of Chemistry 11, 518.
- [87] Dhasmana, H., Garg, H.S., 1989. Phytochemistry 28, 2819.
- [88] Dhasmana, H., Garg, H.S., 1990. Phytochemistry 29, 961.
- [89] Diserens, I.S., Rogers, C., Sordat, B., Hostettmann, K., 1992. Phytochemistry 31, 313.
- [90] Dreyer, D.L., Bourell, J.H., 1981. Phytochemistry 20, 493.
- [91] Duarte, A.P., Correa, D.B., Silva, L.G.F., Janot, S., Gottlieb, O.R., 1968. Anais da Academia Brasileira de Ciências 40, 307.
- [92] Dutta, P.K., Sem, A.K., Sarkar, K.K., Banerji, N., 1987. Indian Journal of Chemistry 26B, 281.
- [93] Elix, J.A., Crook, C.E., Jiang, H., Zhi-Ning, Z., 1992. Australian Journal of Chemistry 45, 845.
- [94] Elix, J.A., Crook, C.E., 1992. The Bryologist 95, 52.
- [95] Elix, J.A., Gaul, K.L., Lumbsch, H.T., 1987. Australian Journal of Chemistry 40, 1031.
- [96] Elix, J.A., Whitton, A.A., Sargent, M.V., 1984. Chemical Organic Naturstoffe 45, 103.
- [97] Ellis, R.C., Whalley, W.B., Ball, K., 1967. Chemical Communications, 803.
- [98] Ferreira, G.A.L., Gottlieb, O.R., Mesquita, A.A.L., 1972. Phytochemistry 11, 1512.
- [99] Frahm, A.W., Chaudhuri, R.K., 1979. Tetrahedron 35, 2035.
- [100] Fujita, T., Liu, D.Y., Ueda, S., Takeda, Y., 1992. Phytochemistry 31, 3997.
- [101] Fukamiya, N., Okano, M., Kondo, K., Tagahara, K., 1990. Journal of Natural Products 53, 1543.
- [102] Fukuyama, Y., Kamiyama, A., Mima, Y., Kodama, M., 1991. Phytochemistry 30, 3433.
- [103] Fukuyama, Y., Mitsunami, H., Yoshizawa, T., Taki, J., Nakagawa, K., 1996. Heisei July, 7.
- [104] Gabriel, S.J., Gottlieb, O.R., Lima, R.A., Mesquita, A.A.L., 1977. Acta Amazonica 7, 289.
- [105] Ghosal, S., Sharma, P.V., Chaudhuri, R.K., 1975. Phytochemistry 14, 1393.
- [106] Ghosal, S., Biswas, K., Jaiswal, D., 1980. Phytochemistry 19, 123.
- [107] Ghosal, S., Chaudhuri, R.K., 1975. Journal of Pharmaceutical Science 64, 888.
- [108] Ghosal, S., Chaudhuri, R.K., Nath, A., 1973. Journal of Pharmaceutical Science 62, 137.

- [109] Ghosal, S., Chaudhuri, R.K., 1973. *Phytochemistry* 12, 2035.
- [110] Ghosal, S., Jaiswal, D.K., Biswas, K., 1978. *Phytochemistry* 17, 2119.
- [111] Ghosal, S., Sharma, P.V., Chaudhuri, R.K., Bhattacharya, S.K., 1975. *Journal of Pharmaceutical Science* 64, 80.
- [112] Ghosal, S., Sharma, P.V., Chaudhuri, R.K., Bhattacharya, S.K., 1973. *Journal of Pharmaceutical Science* 62, 926.
- [113] Ghosal, S., Sharma, P.V., Chaudhuri, R.K., 1975. *Phytochemistry* 14, 2671.
- [114] Gil, S., Palanca, P., Sanz, V., Tortajada, A., 1991. *Journal of Natural Products* 54, 127.
- [115] Gil, S., Palanca, P., Sanz, V., Tortajada, A., 1990. *Journal of Natural Products* 53, 1198.
- [116] Gil, S., Parra, M., Sanz, V., Tortajada, A., 1988. *Journal of Natural Products* 51, 339.
- [117] Gil, S., Sanz, V., Tortajada, V., 1987. *Journal of Natural Products* 50, 301.
- [118] Goh, S.H., Jantan, I., Gray, A.I., Waterman, P.G., 1992. *Phytochemistry* 31, 1383.
- [119] Goh, O.S.H., Jantan, O.I., 1991. *Phytochemistry* 30, 366.
- [120] Gottlieb, O.R., Mesquita, A.A.L., Oliveira, G.G., Melo, M.T., 1970. *Phytochemistry* 9, 2537.
- [121] Gottlieb, O.R., Mesquita, A.A.L., Silva, E.M., Melo, M.T., 1969. *Phytochemistry* 8, 665.
- [122] Gottlieb, O.R., Lima, R.A., Mendes, P.H., Magalhães, M.T., 1975. *Phytochemistry* 14, 1674.
- [123] Gottlieb, O.R., Magalhães, M.T., 1966. *Anais da Academia Brasileira de Ciências* 38, 439.
- [124] Gottlieb, O.R., Magalhães, M.T., Pereira, M.O.S., Mesquita, A.A.L., Correa, D.B., Oliveira, G.G., 1968. *Tetrahedron* 24, 1601.
- [125] Gottlieb, O.R., 1968. *Phytochemistry* 7, 411.
- [126] Govindachari, T.R., Kalyanaraman, P.S., Muthukumaraswamy, N., Pai, B.R., 1971. *Tetrahedron* 27, 3919.
- [127] Govindachari, T.R., Pai, B.R., Subramaniam, P.S., Rao, U.R., Muthukumaraswamy, N., 1967. *Tetrahedron* 23, 243.
- [128] Govindachari, T.R., Subramaniam, P.S., Pai, B.R., Kalyanaraman, P.S., Rao, U.R., 1971. *Indian Journal of Chemistry* 9, 772.
- [129] Grover, P.K., Shah, G.D., Shah, R.C., 1955. *Journal of the Chemical Society*, 3982.
- [130] Gunasekera, S.P., Jayatilake, G.S., Selliah, S.S., Sultanbawa, M.U.S., 1977. *Journal of the Chemical Society, Perkin Transactions I*, 1505.
- [131] Gunasekera, S.P., Selliah, S.S., Sultanbawa, M.U.S., 1975. *Journal of the Chemical Society, Perkin Transactions I*, 1539.
- [132] Gunasekera, S.P., Sivapalan, K., Sultanbawa, M.U.S., Ollis, W.D., 1977. *Journal of the Chemical Society, Perkin Transactions I*, 11.
- [133] Gunasekera, S.P., Sotheeswaran, S., Sultanbawa, M.U.S., 1981. *Journal of the Chemical Society, Perkin Transactions I*, 1831.
- [134] Gunasekera, S.P., Sultanbawa, M.U.S., 1975. *Journal of the Chemical Society, Perkin Transactions I*, 2215.
- [135] Gunatilaka, A.A.L., Sriyani, H.T.B., Sotheeswaran, S., Waight, E.S., 1983. *Phytochemistry* 22, 233.
- [136] Guyot, M., Massicot, J., Rivaille, P., Normant, H.C.R., 1968. *Academic Science Paris* 267, 423.
- [137] Habib, A.M., Ho, D.K., Masuda, S., McCloud, T., Reddy, K.S., Abou-Shoer, M., McKenzie, A., Byrn, S.R., Chang, C.J., Cassady, J.M., 1987. *The Journal of Organic Chemistry* 52, 412.
- [138] Habib, A.M., Reddy, K.S., McCloud, T.G., Chang, C.J., Cassady, J.M., 1987. *Journal of Natural Products* 50, 141.
- [139] Hamblon, H., Frahm, A.W., 1980. *Tetrahedron* 36, 3273.
- [140] Hano, Y., Matsumoto, Y., Shinohara, K., Sun, J.Y., Nomura, T., 1991. *Planta Medica* 57, 172.
- [141] Hano, Y., Matsumoto, Y., Sun, J.Y., Nomura, T., 1990. *Planta Medica* 56, 478.
- [142] Hano, Y., Matsumoto, Y., Sun, J.Y., Nomura, T., 1990. *Planta Medica* 56, 399.
- [143] Hano, Y., Okamoto, T., Nomura, T., 1991. *Heterocycles* 32, 1357.
- [144] Hano, Y., Okamoto, T., Nomura, T., Momose, Y., 1990e. *Heterocycles* 31, 1345.
- [145] Haqqani, M.H., 1981. *Fitoterapia* 52, 5.
- [146] Harborne, J.B., 1973. *Phytochemical Methods*. Chapman & Hall, London, pp. 278.
- [147] Harrison, L.J., Leong, L.S., Sai, G.L., Sim, K.Y., Tan, H.T.W., 1993. *Phytochemistry* 33, 727.
- [148] Harrison, L.J., Leong, L.S., Leong, Y.W., Lee, S.G., Sai, K.Y., Sim, K.Y. et al., 1994. *Natural Product Letters* 5, 11.
- [149] He, R., Feng, S., Nie, R., 1984. *Yunnan Zhiwu Yanjin* 6, 341.
- [150] He, R., Shuji, F., Nie, R., 1982. *Yunnan Zhiwu Yanjin* 4, 68.
- [151] Holker, J.S.E., Kagal, S.A., 1968. *Chemical Communications*, 1574.
- [152] Holloway, D.M., Scheinmann, F., 1975. *Phytochemistry* 14, 2517.
- [153] Hostettmann, K., Guillarmod, A.J., 1974. *Helvetica Chimica Acta* 57, 1155.
- [154] Hostettmann, K., Guillarmod, A.J., 1976. *Helvetica Chimica Acta* 59, 1584.
- [155] Hostettmann, K., Guillarmod, A.J., 1976. *Journal of Chromatography* 124, 381.
- [156] Hostettmann, K., Guillarmod, A.J., 1977. *Phytochemistry* 16, 481.
- [157] Hostettmann, K., McNair, H.M., 1976. *Journal of Chromatography* 116, 201.
- [158] Hostettmann, K., Tabacchi, R., Guillarmod, A.J., 1974. *Helvetica Chimica Acta* 57, 294.
- [159] Hostettmann, K., Wagner, H., 1977. *Phytochemistry* 16, 821.
- [160] Hu, B., Hongfa, S., Shufen, F., Jingye, D., 1992. *Zhiwu Xuebao* 34, 886.
- [161] Iglesias, M.C.R., Marston, A., Hostettmann, K., 1992. *Phytochemistry* 31, 1387.
- [162] Inuma, M., Tosa, H., Tanaka, T., Asai, F., Shimano, R., 1995. *Heterocycles* 40, 279.
- [163] Inuma, M., Tosa, H., Tanaka, T., Shimano, R., Asai, F., Yonemori, S., 1994. *Phytochemistry* 35, 1355.
- [164] Inuma, M., Ito, T., Tosa, H., Tanaka, T., 1996. *Journal of Natural Products* 59, 472.
- [165] Inuma, M., Ito, T., Tosa, H., Tanaka, T., Miyake, R., Chelladurai, V., 1997. *Phytochemistry* 46, 1423.
- [166] Inuma, M., Tosa, H., Ito, T., Tanaka, T., Madulid, D.A., 1996. *Phytochemistry* 42, 1195.
- [167] Inuma, M., Tosa, H., Ito, T., Tanaka, T., Riswan, S., 1996. *Chemical and Pharmaceutical Bulletin* 4 (9), 1744.
- [168] Inuma, M., Tosa, H., Tanaka, T., Asai, F., Shimano, R., 1995. *Phytochemistry* 38, 247.
- [169] Inuma, M., Tosa, H., Tanaka, T., Asai, F., Shimano, R., 1995. *Phytochemistry* 39, 945.
- [170] Inuma, M., Tosa, H., Tanaka, T., Riswan, S., 1996. *Chemical and Pharmaceutical Bulletin* 44 (1), 232.
- [171] Inuma, M., Tosa, H., Tanaka, T., Yonemori, S., 1995. *Phytochemistry* 38, 725.
- [172] Inuma, M., Tosa, H., Toriyama, N., Tanaka, T., Ito, T., Chelladurai, V., 1996. *Phytochemistry* 43, 681.
- [173] Inuma, M., Tosa, H., Tanaka, T., Yonemori, S., 1994. *Phytochemistry* 35, 527.
- [174] Ikeya, Y., Sugama, K., Okada, M., Mitsuhashi, H., 1991. *Phytochemistry* 30, 2061.
- [175] Ishiguro, K., Fukumoto, H., Suitani, A., Nakajima, M., Itoi, K., 1996. *Phytochemistry* 42, 435.
- [176] Ishiguro, K., Nakajima, M., Fukumoto, H., Itoi, K., 1995. *Phytochemistry* 38, 867.

- [177] Ishiguro, K., Fukumoto, H., Nakajima, M., Isoi, K., 1993. *Phytochemistry* 33, 839.
- [178] Ishiguro, K., Nagareya, N., Suitani, A., Fukumoto, H., 1997. *Phytochemistry* 44, 1065.
- [179] Ishiguro, K., Nakajima, M., Fukumoto, H., Isoi, K., 1995. *Phytochemistry* 39, 903.
- [180] Ishimaru, K., Sudo, H., Satake, M., Matsunaga, Y., Hasegawa, Y., Takemoto, S. et al., 1990. *Phytochemistry* 29, 1563.
- [181] Ito, H., Taniguchi, H., Kita, T., Matsuki, Y., Tachikawa, E., Fujita, T., 1977. *Phytochemistry* 16, 1614.
- [182] Jackson, B., Locksley, H.D., Scheinmann, F., 1966. *Journal of the Chemical Society C*, 178.
- [183] Jackson, B., Locksley, H.D., Scheinmann, F., 1967. *Journal of the Chemical Society C*, 2500.
- [184] Jackson, B., Locksley, H.D., Scheinmann, F., 1969. *Phytochemistry* 8, 927.
- [185] Jefferson, A., Scheinmann, F., 1965. *Nature* 207, 1193.
- [186] Kaldas, M.H., Guillarmod, A.J., 1978. *Phytochemistry* 17, 2083.
- [187] Kaldas, M.H., Hostettmann, K., Sticher, O., 1981. *Phytochemistry* 20, 443.
- [188] Kaldas, M., Hostettmann, K., Guillarmod, A.J., 1974. *Helvetica Chimica Acta* 57, 2557.
- [189] Kanamori, H., Sakamoto, I., Mizuta, M., Hashimoto, K., Tanaka, O., 1984. *Chemical and Pharmaceutical Bulletin* 32, 2290.
- [190] Kaouadji, M., Vaillant, I., Mariotte, A.M., 1986. *Journal of Natural Products* 49, 359.
- [191] Karunanayake, S., Sotheeswaran, S., Sultanbawa, M.U.S., Balasubramaniam, S., 1981. *Phytochemistry* 20, 1303.
- [192] Kazmi, S.N.U.H., Ahmed, Z., Malik, A., 1989. *Heterocycles* 29, 1923.
- [193] Kazmi, S.N.U.H., Ahmed, Z., Malik, A., 1989. *Phytochemistry* 28, 3572.
- [194] Khetwal, K.S., Bisht, R.S., 1988. *Phytochemistry* 27, 1910.
- [195] Khetwal, K.S., Joshi, B., Bisht, R.S., 1990. *Phytochemistry* 29, 1265.
- [196] Khetwal, K.S., Pande, S., 1997. *Indian Journal of Chemistry* 36B, 833.
- [197] Khorost, O.P., Serbin, A.G., Komissarenko, N.F., Bublik, N.P., 1987. *Chemical Natural Compounds* 4, 506.
- [198] King, F.E., King, T.J., Manning, L.C., 1953. *Journal of the Chemical Society*, 3932.
- [199] Kitanov, G.M., Assenov, I., Van, A.T., 1988. *Pharmazie* 43, 879.
- [200] Kitanov, G.M., Blinova, K.F., 1980. *Khimiya Prirodnnykh Soedinenii* 2, 256.
- [201] Kitanov, G.M., Blinova, K.F., 1987. *Chemical Natural Compounds* 2, 151.
- [202] Kittisak, L., Thatree, P., Chulabhorn, M., Somsak, R., 1997. *Phytochemistry* 45 (6), 1299.
- [203] Komatsu, M., Tomimori, T., Mikuriya, N., 1969. *Chemical and Pharmaceutical Bulletin* 17, 155.
- [204] Kostanecki, V., 1892. *Ber. D. Chemistry* 25, 1640.
- [205] Kostanecki, V., Rutishauser, R., 1892. *Ber. D. Chemistry* 25, 1648.
- [206] Kubitzki, K., Mesquita, A.A.L., Gottlieb, O.R., 1978. *Biochemical Systematics Ecology* 6, 185.
- [207] Kulanthaivel, P., Pelletier, S.W., Khetwal, O.K.S., Verna, D.L., 1988. *Journal of Natural Products* 51, 379.
- [208] Kumar, V., Ramachandran, S., Sultanbawa, M.U.S., 1976. *Phytochemistry* 15, 2016.
- [209] Kumar, V., Sotheeswaran, S., Surendrakumar, S., Balasubramaniam, S., 1982. *Phytochemistry* 21, 807.
- [210] Kupchan, S.M., Streelman, D.R., Sneden, A.T., 1980. *Journal of Natural Products* 43, 296.
- [211] Lee, H.H., Chan, H.K., 1977. *Phytochemistry* 16, 2038.
- [212] Leuckert, C., Ahmadjian, V., Culberson, C.F., Johnson, A., 1990. *Mycologia* 82, 370.
- [213] Lewis, J.R., 1963. *Proceedings of the Chemistry Society* 363, 373.
- [214] Lima, R.A., Gottlieb, O.R., Mesquita, A.A.L., 1970. *Anais da Academia Brasileira de Ciências* 42, 133.
- [215] Lima, R.A., Gottlieb, O.R., Mesquita, A.A.L., 1972. *Phytochemistry* 11, 2307.
- [216] Lin, C.N., Chung, Mi, Liou, S.J., Lee, T.H., Wang, J.P., 1996. *Journal of Pharmaceutical Pharmacology* 48, 532.
- [217] Lin, C.N., Liou, S.S., Ko, F.N., Teng, C.M., 1993. *Journal of Pharmaceutical Science* 82, 11.
- [218] Lin, C.N., Liou, S.S., Ko, F.N., Teng, C.M., 1992. *Journal of Pharmaceutical Science* 81, 1109.
- [219] Liu, J., Huang, M., 1982. *Zhongcaoyao* 13, 433.
- [220] Locksley, H.D., Moore, I., Scheinmann, F., 1966. *Journal of the Chemical Society C*, 430.
- [221] Locksley, H.D., Moore, I., Scheinmann, F., 1966. *Journal of the Chemical Society C*, 2265.
- [222] Locksley, H.D., Moore, I., Scheinmann, F., 1967. *Tetrahedron* 23, 2229.
- [223] Locksley, H.D., Moore, I., Scheinmann, F., 1966. *Journal of the Chemical Society C*, 2186.
- [224] Locksley, H.D., Murray, I.G., 1969. *Journal of the Chemical Society C*, 1567.
- [225] Locksley, H.D., Murray, I.G., 1970. *Journal of the Chemical Society C*, 392.
- [226] Locksley, H.D., Murray, I.G., 1971. *Journal of the Chemical Society C*, 1332.
- [227] Locksley, H.D., Murray, I.G., 1971. *Phytochemistry* 10, 3179.
- [228] Mahabusarakam, W., Wiriyachitra, P., Taylor, W., 1987. *Journal of Natural Products* 50, 474.
- [229] Mahmoud, Z.F., Salam, N.A.A., Khafagy, S.M., 1980. *Fitoterapia* 51, 153.
- [230] Mandal, S., Das, P.C., Joshi, P.C., 1992. *Journal of the Indian Chemical Society* 69, 611.
- [231] Mandal, S., Chatterjee, A., 1987. *Tetrahedron Letters* 28, 1309.
- [232] Mao, S., Liao, S., Wu, J., Liang, H., Chen, H., Zhang, C., 1996. *Yaoxue Xuebao* 31 (2), 118.
- [233] Markham, K.R., 1964. *Tetrahedron* 20, 991.
- [234] Markham, K.R., 1965. *Tetrahedron* 21, 1449.
- [235] Markham, K.R., 1965. *Tetrahedron* 21, 3687.
- [236] Marston, A., Hamburger, M., Diserens, I.S., Msonthi, J.D., Hostettmann, K., 1993. *Phytochemistry* 33, 809.
- [237] Marta, M., Delle-Monache, F., Bettolo, G.B.M., Lyra, D.A., Mello, J.F., Lima, O.G., 1976. *Gazzetta Chimica Italiana* 106, 651.
- [238] Massias, M., Carbonnier, J., Molho, D., 1977. *Bulletin of the Museum of National History Natural Paris* 13, 55.
- [239] Massias, M., Carbonnier, J., Molho, D., 1976. *Bulletin of the Museum of National History Natural Paris* 10, 45.
- [240] Mathis, C.T., Goldstein, J.H., 1964. *Spectrochimica Acta* 20, 871.
- [241] McMaster, W.J., Scott, A.I., Trippett, S., 1960. *Journal of the Chemical Society*, 4628.
- [242] Mehta, G., Shah, S.R., Venkateswarlu, Y., 1994. *Tetrahedron* 50, 11729.
- [243] Meravy, L., 1987. *Biology of Plants* 29, 81.
- [244] Mesquita, A.A.L., Correa, D.B., Gottlieb, O.R., Magalhães, M.T., 1968. *Anais Chimica Acta* 42, 311.
- [245] Mesquita, A.A.L., Oliveira, W.G., Neiva, R.M.T., Gottlieb, O.R., 1975. *Phytochemistry* 14, 803.
- [246] Miana, G.A., Al-Hazimi, H.M.G., 1984. *Phytochemistry* 23, 1637.
- [247] Minami, H., Kinoshita, M., Fukuyama, Y., Kodama, M., Yoshizawa, T., Sugiura, M., Nakagawa, K., Tago, H., 1994. *Phytochemistry* 36, 501.

- [248] Minami, H., Takahashi, E., Kodama, M., Fukuyama, Y., 1996. *Phytochemistry* 41, 629.
- [249] Mourão, J.C., Gottlieb, O.R., Magalhães, M.T., 1966. *Anais da Academia Brasileira Ciências* 38, 435.
- [250] Mukherjee, B., Mukherjee, S.K., 1987. *International Journal of Crude Drug Research* 25, 97.
- [251] Mukherjee, K.S., Chakraborty, C.K., Chatterjee, T.P., Bhattacharjee, D., Laha, S., 1991. *Phytochemistry* 30, 1036.
- [252] Mukherjee, K.S., Chakraborty, C.K., Laha, S., Bhattacharya, D., Chatterjee, T.P., 1990. *Journal Indian Chemical Society* 67, 1003.
- [253] Mukherjee, K.S., Laha, S., Manna, T.K., Roy, S.C., 1995. *Journal of the Indian Chemical Society* 72, 63.
- [254] Mukherjee, K.S., Manna, T.K., Laha, S., Chakravorty, C.K., 1994. *International Journal Pharmacology* 32, 201.
- [255] Murti, V.V.S., Seshadri, T.R., Sivakumaran, S., 1972. *Phytochemistry* 11, 2089.
- [256] Nagem, T.J., Silveira, J.C., 1986. *Phytochemistry* 25, 503.
- [257] Nagem, T.J., Silveira, J.C., 1988. *Phytochemistry* 27, 646.
- [258] Nagem, T.J., Silveira, J.C., 1986. *Phytochemistry* 25, 2681.
- [259] Nagem, T.J., Silveira, J.C., 1989. *Phytochemistry* 28, 2211.
- [260] Nattaya, C., Kazuya, T., Yasushi, O., Shigeo, N., Tomohisa, O., 1996. *Phytochemistry* 43 (5), 1099.
- [261] Neshta, N.M., Glyzin, V.I., Patudin, A.V., 1984. *Chemical Natural Compounds* 1, 108.
- [262] Neshta, N.M., Glyzin, V.I., Savina, A.A., Patudin, A.V., 1983. *Chemical Natural Compounds* 6, 750.
- [263] Nielsen, H., Arends, P., 1979. *Journal of Natural Products* 42, 301.
- [264] Nomura, T., Hano, Y., Fujimoto, T., 1983. *Heterocycles* 20, 213.
- [265] Noro, T., Ueno, A., Mizutani, M., Hashimoto, T., Miyase, T., Kuroyanage, M., Fukushima, S., 1984. *Chemical and Pharmaceutical Bulletin* 32, 4455.
- [266] Odontuya, G., Purev, O., Sambuu, D.G., Nasreen, A., Atta-Ur, R., 1995. *Natural Products Letters* 5, 269.
- [267] Okorie, D.A., 1976. *Phytochemistry* 15, 1799.
- [268] Oliveira, M.C.F., Silveira, E.R., 1995. In: 18th Reunião Anual da Sociedade Brasileira de Química, PN 029.
- [269] Oliveira, W.G., Gottlieb, O.R., Mesquita, A.A.L., 1972. *Phytochemistry* 11, 3323.
- [270] Oliveira, W.G., Mesquita, A.A.L., Kubitzki, K., Gottlieb, O.R., 1990. *Phytochemistry* 29, 1893.
- [271] Owen, P.J., Scheinmann, F., 1974. *Journal of the Chemical Society, Perkin Transactions I*, 1018.
- [272] Parra, M., Picher, M.T., Seoane, E., Tortajada, A., 1985. *Journal of Natural Products* 48, 998.
- [273] Parra, M., Picher, M.T., Seoane, E., Tortajada, A., 1984. *Journal of Natural Products* 47, 123.
- [274] Parra, M., Seoane, E., Tortajada, A., 1984. *Journal of Natural Products* 47, 868.
- [275] Parveen, M., Khan, N.U.D., 1988. *Phytochemistry* 27, 3694.
- [276] Parveen, M., Khan, N.U.D., 1987. *Chemistry and Industry* 12, 418.
- [277] Patel, G.N., Trivedi, K.N., 1983. *Indian Journal of Chemistry* 22B, 755.
- [278] Patel, G.N., Trivedi, K.N., 1991. *Indian Journal of Chemistry* 30B, 437.
- [279] Patolia, R.J., Trivedi, K.N., 1983. *Indian Journal of Chemistry* 22B, 444.
- [280] Pattalung, P., Thongtheeraparp, W., Wiriyachitra, P., Taylor, W.C., 1994. *Planta Medica* 60, 365.
- [281] Pereira, M.O.S., Gottlieb, O.R., Magalhães, M.T., 1966. *Anais da Academia Brasileira de Ciências* 38, 425.
- [282] Peres, V., Nagem, T.J., 1997. *Phytochemistry* 44, 191.
- [283] Pinto, D.C.G., Fuzzati, N., Pazmino, X.C., Hostettmann, K., 1994. *Phytochemistry* 7, 875.
- [284] Pinto, M.M., Polonia, J., 1974. *Helvetica Chimica Acta* 57, 286.
- [285] Pureb, O., Vansan, Y.Z., Oyuun, K., 1991. *Chemical Natural Compounds* 2, 245.
- [286] Purev, O., Odontuya, G., Oyun, H., Maksimovna, T.L., Nasreen, A., Atta-Ur, R., 1995. *Natural Products Letters* 5, 261.
- [287] Quillinan, A.J., Scheinmann, F., 1972. *Journal of the Chemical Society, Perkin Transactions I*, 1382.
- [288] Quillinan, A.J., Scheinmann, F., 1973. *Journal of the Chemical Society, Perkin Transactions I*, 1329.
- [289] Rao, A.V.R., Sarma, M.R., Venkataraman, K., Yemul, S.S., 1974. *Phytochemistry* 13, 1241.
- [290] Rath, G., Potterat, O., Mavi, S., Hostettmann, K., 1996. *Phytochemistry* 43, 513.
- [291] Ravi, P., Vathani, P., Reddy, G.C., 1994. *Indian Journal of Heterocycles Chemistry* 3, 209.
- [292] Rivaille, P., Massicot, J., Guyot, M., Plouvier, V., Massias, M., 1969. *Phytochemistry* 8, 1533.
- [293] Roberts, J.C., 1961. *Chemical Reviews* 61, 591.
- [294] Rocha, L., Marston, A., Kaplan, M.A.C., Evans, H.S., Thull, U., Testa, B., Hostettman, K., 1994. *Phytochemistry* 36, 1381.
- [295] Rodriguez, S., Wolfender, J.L., Hakizamungu, E., Hostettmann, K., 1995. *Planta Medica* 61, 362.
- [296] Rodriguez, S., Wolfender, J.L., Odontuya, G., Purev, O., Hostettmann, K., 1995. *Phytochemistry* 40, 1265.
- [297] Roitman, J.N., Wollenweber, E., Giner, F.J.A., 1992. *Journal Plant Physiology* 139, 632.
- [298] Sakai, S.I., Katsura, M., Takayama, H., Aimi, N., Chokethaworn, N., Suttajit, M.E., 1993. *Chemical and Pharmaceutical Bulletin* 41, 958.
- [299] Santesson, J., 1968a. *Arkiv Kemi* 30, 363.
- [300] Santesson, J., 1968b. *Acta Chemica Scandinavica* 22, 2393.
- [301] Santesson, J., 1969. *Arkiv Kemi* 30, 479.
- [302] Schaufelberger, D., Hostettmann, K., 1988. *Planta Medica* 54, 219.
- [303] Scheinmann, F., Sripong, N.A., 1971. *Phytochemistry* 10, 1331.
- [304] Scheinmann, F., 1962. *Tetrahedron* 18, 853.
- [305] Sem, A.K., Sarkar, K.K., Majumder, P.C., Banerji, N., Uusvuori, R., Hase, T.A., 1980. *Phytochemistry* 19, 2223.
- [306] Sen, A.K., Sarkar, K.K., Majumder, P.C., Banerji, N., 1980. *Indian Journal of Chemistry* 19B, 1008.
- [307] Sen, A.K., Sarkar, K.K., Majumder, P.C., Banerji, N., 1986. *Indian Journal of Chemistry* 25B, 1157.
- [308] Sen, A.K., Sarkar, K.K., Majumder, P.C., Banerji, N., Uusvuori, R., Hase, T.A., 1982. *Phytochemistry* 21, 1747.
- [309] Shah, G.D., Shah, R.C., 1956. *Journal of Science Industry Research* 15B, 630.
- [310] Sia, G.L., Bennett, G.J., Harrison, L.J., Sim, K.Y., 1995. *Phytochemistry* 38, 1521.
- [311] Simoneau, B., Brassard, P.A., 1984. *Journal of the Chemical Society, Perkin Transactions I*, 1507.
- [312] Somanathan, R., Sultanbawa, M.U.S., 1992. *Journal of the Chemical Society, Perkin Transactions I*, 1935.
- [313] Stout, G.H., Balkenhol, W.J., 1969. *Tetrahedron* 25, 1947.
- [314] Stout, G.H., Christensen, E.N., Balkenhol, W.J., Stevens, K.L., 1969. *Tetrahedron* 25, 1961.
- [315] Stout, G.H., Fries, J.L., 1970. *Phytochemistry* 9, 235.
- [316] Stout, G.H., Krahn, M.M., Yates, P., Bhat, H.B., 1968. *Chemical Communications*, 211.
- [317] Stout, G.H., Stout, V.F., Welsh, M.J., Jensen, L.H., 1962. *Tetrahedron Letters* 13, 541.
- [318] Stout, G.H., Stout, V.F., Welsh, M.J., 1963. *Tetrahedron* 19, 667.
- [319] Sullivan, G., Stiles, F.D., Rosler, K.H.A., 1977. *Journal of Pharmaceutical Science* 66, 828.
- [320] Sultanbawa, M.U.S., 1980. *Tetrahedron* 36, 1465.
- [321] Sun, H., Jingye, D., 1981. *Zhiwu Xuebao* 23, 464.
- [322] Sun, H., Hu, B., Fan, S., Ding, J., 1983. *Zhiwu Xuebao* 25, 460.

- [323] Sundholm, E.G., 1978. *Tetrahedron* 34, 577.
- [324] Suzuki, O., Katsumata, Y., Oya, M., Chari, V.M., Klapfenberger, R., Wagner, H., Hostettman, K., 1980. *Planta Medica* 39, 19.
- [325] Suzuki, O., Katsumata, Y., Oya, M., Chari, V.M., Vermes, B., Wagner, H., Hostettman, K., 1981. *Planta Medica* 42, 17.
- [326] Tadzhibaev, M.M., Butayarov, A.V., Batirov, E.K., Malikov, V.M., 1992. *Chemical Natural Compounds* 2, 244.
- [327] Takagi, S., Yamari, M., 1982. *Yakugaku Zasshi* 102, 546.
- [328] Terreaux, C., Maillard, M., Gupt, M.P., Hostettmann, K., 1995. *Phytochemistry* 40, 1791.
- [329] Tomimori, T., Yoshizaki, M., Namba, T., 1974. *Yakugaku Zasshi* 94, 647.
- [330] Tosa, H., Linuma, M., Murakami, K., Ito, T., Tanaka, T., Chelladura, V. et al., 1997. *Phytochemistry* 45, 133.
- [331] Ueno, A., 1962. *Yakugaku Zasshi* 82, 1486.
- [332] Ueno, A., 1962. *Yakugaku Zasshi* 82, 1482.
- [333] Van Der Sluis, W.G., Labadie, R.P., 1985. *Phytochemistry* 24, 2601.
- [334] Van Der Sluis, W.G., 1985. *Plant System Evolution* 149, 253.
- [335] Versluis, C., Cortés, M., López, J.T., Sierra, J.R., Razmilié, I., 1982. *Experientia* 38, 771.
- [336] Vitale, A.A., Romanelli, G.P., Autino, J.C., Pomilio, A.B., 1994. *Journal of Chemical Research S*, 82.
- [337] Walia, S., Mukerjee, S.K., 1984. *Phytochemistry* 23, 1816.
- [338] Waterman, P.G., Crichton, E.G., 1980. *Planta Medica* 40, 351.
- [339] Waterman, P.G., Crichton, E.G., 1980. *Phytochemistry* 19, 2723.
- [340] Waterman, P.G., Hussain, R.A., 1982. *Phytochemistry* 21, 2099.
- [341] Westerman, P.W., Gunasekera, S.P., Sultanbawa, M.U.S., Kazlauskas, R., 1977. *Organic Magnetic Resonance* 9, 631.
- [342] Wolfender, J.L., Hamburger, M., Msonthi, J.D., Hostettmann, K., 1991. *Phytochemistry* 30, 3625.
- [343] Wolfrom, M.L., Bhat, H.B., 1965. *Phytochemistry* 4, 765.
- [344] Wolfrom, M.L., Dickey, E.E., McWain, P., Thompson, A., Looker, J.H., Windrath, O.M. et al., 1964. *The Journal of Organic Chemistry* 29, 689.
- [345] Wolfrom, M.L., Komitsky, F. Jr., Fraenkel, G., Looker, J.H., Dickey, E.E., McWain, P. et al., 1963. *Tetrahedron Letters* 12, 749.
- [346] Wolfrom, M.L., Komitsky, F. Jr., Mundell, P.M., 1965. *Journal of Organic Chemistry* 30, 1088.
- [347] Wollenweber, E., Dorr, M., Roitman, J.N., Giner, F.J.A., 1994. *Zhurnal of Naturforsch* 49c, 393.
- [348] Yan-Bin, Y., Zhou, J., 1980. *Yun-nan, Chih Wu Yen Chiu* 2, 468.
- [349] Zang, B.C., Cheng, J.T., Hu, P.L., Ho, S.H., 1980. *Chun Ts'ao Yao* 11, 149.
- [350] Zhou, H.M., Liu, Y.L., Blasko, G., Cordell, G.A., 1989. *Phytochemistry* 28, 3569.