



On *Plesionika persica* (Kemp, 1925) and *P. reflexa* Chace, 1985 (Crustacea: Decapoda: Pandalidae) from India

TIN-YAM CHAN^{1,2}, REKHA DEVI CHAKRABORTY³, P. PURUSHOTHAMAN³,
G. KUBERAN³ & CHIEN-HUI YANG^{1,2,4}

¹Institute of Marine Biology, National Taiwan Ocean University, 2 Pei-Ning Road, Keelung 20224, Taiwan, R.O.C.

²Center of Excellence for the Oceans, National Taiwan Ocean University, 2 Pei-Ning Road, Keelung 20224, Taiwan, R.O.C.

³Crustacean Fisheries Division, Central Marine Fisheries Research Institute, Ernakulam North P.O., P.B. No. 1603, Cochin-682018, Kerala, India

⁴Corresponding author. E-mail: chyang.03@ntou.edu.tw

Abstract

The availability of Indian specimens of *Plesionika persica* (Kemp, 1925) and *P. reflexa* Chace, 1985 provided more information on the taxonomy around these two species. Moreover, it is the first record of *P. persica* to India. Although *P. taiwanica* Chan and Yu, 2000 is superficially rather similar to *P. persica*, there are many differences between them and probably it is inappropriate to establish a species group for these two species. It is likely that all previous records of *P. ensis* (A. Milne-Edwards, 1881) from India actually represent *P. reflexa* Chace, 1985. Nevertheless, the present Indian specimens of *P. reflexa* have more than 10% COI sequence divergence from the topotypic materials of both *P. ensis* and *P. reflexa*, and the epipods at the pereopods III and IV reduced or absent. This data further highlights the confusing taxonomy in the “*P. ensis*” group.

Key words: *Plesionika*, deep-sea, shrimps, taxonomy, India, first record

Introduction

Deep-sea shrimps of the genus *Plesionika* Bate, 1888 are very diverse, with 93 species known to date (Chan 2016). The taxonomy in many species of this genus is still rather confusing (Chan & Crosnier 1997; Li & Chan 2013). Recently specimens of *P. persica* (Kemp, 1925) and *P. reflexa* Chace, 1985 from India are available for study, with the former species being a new record of India. This work reports the findings after examining the Indian material. *Plesionika persica* differs from most species of the genus in having overlong thread-like posterior pereopods but that is also a distinctive characteristic of the Taiwanese species *P. taiwanica* Chan & Yu, 2000. A comparison between *P. persica* and *P. taiwanica* is provided. *Plesionika reflexa* belongs to the “*P. ensis* (A. Milne-Edwards, 1881)” group including only two species but with very confusing taxonomy (Chan & Crosnier 1997; Fransen 2006). The characteristics of the Indian *P. reflexa* material are discussed.

The specimens examined are deposited at the Central Marine Fisheries Research Institute, Cochin (CMFRI), at the National Taiwan Ocean University (NTOU) and at the Muséum national d’Histoire naturelle, Paris (MNHN). The measurement given is carapace length (cl) measured dorsally from the postorbital margin to the posterior margin of the carapace. The synonymy provided is restricted to important taxonomic works and Indian reports on the species. For comparing genetic relatedness, partial sequences of mitochondrial cytochrome c oxidase I (COI, 657 bp) gene barcoding data (Hebert *et al.* 2003; Hajibabaei *et al.* 2007) were generated by following the methods given in Yang *et al.* (2010).

Systematic account

Plesionika persica (Kemp, 1925)

(Fig. 1)

Parapandalus persicus Kemp, 1925: 273, fig. 8. (type locality: Gulf of Oman)

Parapandalus filipes Calman, 1939: 202, fig. 2. (type locality: Gulf of Oman, see Remarks)

Plesionika filipes—Chace 1985: 46.

Plesionika persica—Chace, 1985: 46; Tiefenbacher 1992: 120; Fransen 2006: 73, fig. 21; De Grave & Fransen 2011: 450.

Material examined. India, Kalamuku, Cochin, 250–300 m, April 2017, 1 male cl 11.8 mm, 1 specimen sex undetermined cl 10.5 mm (NTOU); Kerala, Cochin, 250–300 m, April 2017, 4 males cl 10.0–12.0 mm, 5 females cl 10.0–12.0 mm (CMFRI ED.2.4.3.5).

Diagnosis. Rostrum slightly shorter than carapace, curving downwards at basal region, slightly recurved upwards and nearly straight after passing antennular peduncle; with low rostral crest and bearing 6–8 dorsal teeth (including 3 teeth posterior to orbital margin and posteriormost tooth sometimes with incomplete basal suture) restricted to proximal half of rostrum and none with barbed tip; ventral margin armed with 5–7, rarely 3 teeth. Postrostral ridge blunt but strong, extending to about middle of carapace and not highly elevated. Carapace with faint hepatic and branchial grooves; antennal and pterygostomial spines present and distinct, similar sized; orbital margin with upper part straight and dorsally inclined forwards, lower part slightly convex. Eyes without distinct ocellus. Stylocerite tapered anteriorly and not markedly folded upwards laterally, extending to about middle of second segment of antennular peduncle. Scaphocerite slender, 4.6–4.8 times as long as maximum width, distolateral tooth just exceeding distal margin of lamina, basicerite spine moderately long and just exceeding proximal end of lateral margin of scaphocerite. Maxilliped III and pereopods all lacking epipods. Pereiopod I overreaching scaphocerite by dactylus and half propodus, propodus about 0.41–0.43 times as long as carpus. Pereiopod II subequal, extending to tip of scaphocerite and bearing 14–20 carpal articles. Posterior 3 pereopods very long, with pereopods IV and V even thread like (see Remarks). Abdomen with tergite III without any spine, dorsal ridge or groove; pleuron IV posteroventrally rounded, while pleuron V with posteroventral angle somewhat pointed; somite VI 2.2–2.5 times longer than maximum height; telson as long as somite VI and bearing 3 pairs of dorsolateral spines (excluding pair adjacent to posterior margin of telson).

Distribution. Arabian Sea and Red Sea, mesopelagic, at depths of 0–2500 m (see Tiefenbacher 1992; Fransen 2006).

Remarks. *Plesionika persica* is characterized in having extremely long thread-like pereopods IV and V. Only three described species in genus *Plesionika* have such overlong posterior pereopods: *P. persica* (Kemp, 1925), *P. filipes* (Calman, 1939) and *P. taiwanica* Chan & Yu, 2000 (Chan & Yu 2000). *Plesionika persica* and *P. filipes* are very similar (Chace 1985) and have their type localities nearby. Although Calman (1939) mentioned that the holotype of *P. filipes* was collected from the “South Arabian Coast”, the exact position of the station (*i.e.*, St. 76B) for the holotype is 24°13'54"–24°16'42"N/59°16'42"E/59°03'30"E–59°06'30"E (see Tiefenbacher 1992), which is actually at the entrance of Gulf of Oman and very near to the type locality of *P. filipes* (23°44'30"N, 58°52'15"E; Kemp 1925). Therefore, these two species are treated as synonyms by Tiefenbacher (1992) and followed in recent works (e.g., Fransen 2006; De Grave & Fransen 2011). *Plesionika taiwanica* is endemic to Taiwan and superficially rather similar to *P. persica* except for having a much longer rostrum that bearing numerous ventral rostral teeth. Nevertheless, detailed comparisons revealed many more differences between these two species. There are strap-like epipods on the maxilliped III and anterior four pereopods in *P. taiwanica* but all these appendages lack epipods in *P. persica*. In *P. taiwanica*, there is no rostral crest (versus a low rostral crest in *P. persica*) and the upper part of the orbital margin is slightly convex (versus straight in *P. persica*). The stylocerite is obtuse and distinctly folded upwards laterally in *P. taiwanica* but tapered anteriorly and not particularly folded upwards laterally in *P. persica*. Moreover, the lateral carapace is smooth in *P. taiwanica* but bearing faint hepatic and branchial grooves in *P. persica*. Therefore, more studies are necessary to determine if these two species with overlong pereopods are really closely related and can be grouped together. For example, the presence or absence of epipods on the maxilliped III and anterior pereopods have been considered by many workers as of generic importance (see Chace 1985; Holthuis 1993).




FIGURE 1. *Plesionika persica* (Kemp, 1925), Kalamuku, 250–300 m, sex undetermined cl 10.5 mm (NTOU). A, lateral carapace; B, anterior carapace, lateral; C, right antennular peduncle and stylocerite, lateral; D, abdominal somites II to VI and tail fan, lateral. Scales = 2 mm.

The “Diagnosis” provided above is based on the Indian material except for the posterior three pereopods that are all missing or very incomplete in the specimens examined. Nevertheless, the present material can still be satisfactorily identified as *P. persica* by the remaining characters especially by the shape of the rostrum (short and with low rostral crest, bearing few teeth on both dorsal and ventral borders and with dorsal teeth restricted to proximal half of rostrum), smooth abdominal tergite III and telson having 3 pairs of dorsolateral spines (see Kemp 1925; Calman 1939; Tiefenbacher 1992; Fransen 2006). Moreover, the male specimen deposited at NTOU has fewer ventral rostral teeth (only three) than other specimens reported for the species.

Plesionika persica is recorded for the first time from India (off the Arabian coast), though this species appears to be rather common in the other parts of the Arabian Sea and the Red Sea (Calman 1939; Tiefenbacher 1992; Fransen 2006). Nevertheless, the coloration of this species is still unknown and with all known material preserved in formalin we are unable to work for genetic sequencing.

***Plesionika reflexa* Chace, 1985**

(Figs. 2, 3)

Pandalus ? *ensis*—Alcock & Anderson, 1899: 284. (non A. Milne-Edwards, 1881).

Pandalus (Plesionika) ensis—Alcock 1901: 96. (non A. Milne-Edwards, 1881).

Plesionika ensis—Suseelan & Mohamed 1969: 88, fig. 1. (non A. Milne-Edwards, 1881).

Plesionika reflexa Chace, 1985: 108, fig. 49. (type locality: Philippines); Hayashi 1986: 137, fig. 88; Hanamura & Takeda 1987: 116; Kensley *et al.* 1987: 318; Takeda & Hanamura 1994: 23, fig. 10; Chan & Crosnier 1997: 194, figs. 3, 24; Fransen 2006: 77, fig. 23; Li 2006: 1290; De Grave & Fransen 2011: 450.

Material examined. Sakthikulangara fishing port, Kollam district, Kerala, 200–300 m 23 Nov 2015, 1 ovig. female cl 15.0 mm (CMFRI); 250 m, 27 April 2017, 1 male cl 14.3 mm (CMFRI ED.2.4.3.8), 1 male cl 14.6 mm, 1 ovig. female cl 18.9 mm, 1 specimen sex undetermined cl 12.8 mm (NTOU M02081).

Diagnosis. Rostrum long, about 2 times longer than carapace length and far overreaching scaphocerite, slightly curved downwards at basal part but slightly recurved upwards and nearly straight after passing middle of antennular peduncle; dorsal margin armed with 6 teeth at basal part as well as 1 subapical tooth, none with barbed tip, basal teeth restricted behind distal end of antennular peduncle and not forming crest, with 2–3 of them posterior to orbital margin and posterior 2–4 teeth bearing incomplete basal suture; ventral margin with 40–43 teeth. Postrostral ridge distinct but blunt, extending to about middle of carapace and not highly elevated. Carapace only with short, faint hepatic grooves; antennal and pterygostomial spines similar and moderately large; orbital margin with upper part nearly vertically convex, lower part also convex. Eye with distinct ocellus. Stylocerite obtuse but distally pointed, laterally slightly folded upwards, just overreaching basal segment of antennular peduncle. Scaphocerite slender, 4.6–5.0 times as long as maximum width, with distolateral tooth just exceeding distal margin of lamina, basicerite spine long and markedly overreaching proximal end of lateral margin of scaphocerite. Maxilliped III bearing well developed epipods, penultimate segment slightly shorter than terminal segment. Only anterior 3 pereopods with epipods, those at pereopods I and II well developed, epipod at pereopod III small and delicate. Pereiopod II subequal and bearing 15–19 carpal articles. Pereiopod III exceeding scaphocerite by length of dactylus and a short portion of propodus, dactylus 0.41–0.47 times as long as propodus, accessory distal spine minute and little separated from base of terminal spine. Abdomen with tergite III bearing well-developed posteromesial spine, which more or less recurved upwards; pleuron IV posteroventrally rounded while pleuron V bearing distinct posteroventral denticle; somite VI 2.5–2.9 times longer than maximum height; telson 0.8 times as long as somite VI and bearing 3 pairs of dorsolateral spines (excluding pair adjacent to posterior margin of telson).

Distribution. Widely distributed in the Indo-West Pacific from Gulf of Aden to Japan and French Polynesia, at depths of 191–910 m.

Remarks. The “Diagnosis” given above was based on the present four Indian specimens; with only one of them (male cl 14.8 mm) having intact rostrum and another one (male cl 14.3 mm) without any complete pereopod III. *Plesionika reflexa* belongs to the “*P. ensis*” group, which currently includes only two recognized species (Chan & Crosnier 1997). Species of the “*P. ensis*” group are very characteristic in having a distinct posteromesial spine on the abdominal tergite III, rostrum long but with only a few dorsal teeth restricted to base and bearing numerous




FIGURE 2. *Plesionika reflexa* Chace, 1985, Sakthikuylangara fishing port, 250 m, A–D, ovig. female cl 18.9 mm, E, male cl 14.6 mm (NTOU M02081). A, lateral carapace; B, anterior carapace, lateral; C, propodus and dactylus of left pereopod III, lateral; D, abdominal somites III to VI and tail fan, lateral; E, abdominal somites III to V, lateral. Scales: A, B, D, E = 5 mm; C = 2 mm.




FIGURE 3. A, *Plesionika ensis* (A. Milne-Edwards, 1881), Guadeloupe, KARUBENTHOS 2015 stn DW 4609, MNHN IU-2013-19054; B, *P. reflexa* Chace, 1985, the Philippines, AURORA stn CP2695, NTOU M02080.

ventral teeth, posterior pereopods rather robust and not particularly long. The two species in this group are assumed to have discrete distribution, with *P. ensis* occurring in the Atlantic while *P. reflexa* is distributed in the Indo-West Pacific. The two species differ only in *P. ensis* having the posteromesial spine on the abdominal somite III straight and the pereopod III with shorter dactylus. On the other hand, *P. reflexa* has the abdominal somite III

posteromesial spine more or less recurved upwards and longer dactylus at the pereopod III (see Chace 1985). However, as noticed by many workers (see Hanamura & Takeda 1987; Chan & Crosnier 1997; Fransen 2006), Indo-West Pacific material of this group has large variations in the curvature of the abdominal posteromesial spine from very straight to distinctly recurved though this spine is always straight in Atlantic specimens. The proportional length of the dactylus of the pereopod III is also highly variable (from 0.09 to 0.46 as long as propodus) at various localities in both the Atlantic and Indo-West Pacific (see Chace 1985; Chan & Crosnier 1997; Fransen 2006). Nevertheless, the pereopod III dactylus is shorter (0.11–0.25 as long as propodus) in the topotypic material of *P. ensis* in the western Atlantic (type locality: near Barbados) but longer (0.30–0.46 as long as propodus) in the topotypic material of *P. reflexa* in the Philippines. The present Indian specimens have the abdominal posteromesial spine from nearly straight (Fig. 2E) to distinctly recurved (Fig. 2D), and even slightly longer pereopod III dactylus (Fig. 2C, 0.41–0.47 as long as propodus).

As discussed in Chan & Crosnier (1997), the large variations in the curvature of the abdominal posteromesial spine, length of pereopod III dactylus and rostrum in the material of *P. ensis* and *P. reflexa* may indicate that there are actually many more species in the “*P. ensis*” group similar to the situation in the “*P. martia* (A. Milne-Edwards, 1883)” and “*P. narval* (Fabricius, 1787)” groups (Chace 1985; Chan & Crosnier 1991). The present Indian material further differs in the epipods at the pereopods III and IV more reduced or even absent. The present specimens have the epipods of the pereopods III half the size of those at preceding pereopods and rather delicate. Moreover, the epipod is completely absent at the pereopod IV in all specimens. The presence or absence of epipods at the pereopods has often been considered as of specific value in *Plesionika* (Chace 1985; Li & Komai 2003). Fransen (2006) also reported that *P. reflexa* material in the western Arabian Sea and Burma from northern Indian Ocean differs from the typical form in having the epipods at the pereopods III and IV reduced though a minute epipod is still present at the pereopod IV. However, Suseelan & Mohamed (1968) mentioned that their SW Indian specimens carry well developed epipods at the anterior four pereopods, and their fig. 1a showed that the dactylus of pereopod III is about one fifth as long as the propodus. Genetic comparisons of the present Indian specimens (2 males, 1 ovig. female, CMFRI ED.2.4.3.8, NTOU M02081; GenBank nos. MG729438–729440) with topotypic material of *P. ensis* (Fig. 3A from the Lesser Antilles: Guadeloupe, KARUBENTHOS 2015 stn DW 4609, MNHN IU-2013-19054, GenBank no. MG729442, Poupin & Corbari 2016: fig. 6e, versus type locality Barbados) and *P. reflexa* (Fig. 3B from the Philippines: eastern Luzon, AURORA stn CP2695, NTOU M02080, GenBank no. MG729441, versus type locality S.E. Luzon) showed that there are high divergence in the barcoding gene COI amongst the specimens from different localities (9.3%, 10.3–10.7%, 14.2–14.5% divergences between material from Lesser Antilles/Philippines, Philippines/India and Lesser Antilles/India, respectively). However, there are only 0.0–0.3% genetic divergence amongst the Indian material including those with the abdominal posteromesial spine variably recurved (Fig. 2D, E). The high genetic difference of the Indian form from topotypic material of both *P. ensis* and *P. reflexa*, and the reduction or absence of epipods at the pereopod III and IV may urge the separation of the Indian form as another species. As pointed out by Chakraborty *et al.* (2015), however, species delimitation using COI data alone is still rather controversial for *Plesionika*. For example, the COI sequence divergences amongst seven *Plesionika* species from Northeast Atlantic and Mediterranean ranged from 2.5% to 18.3% (Matzen da Saliva *et al.* 2013) while that of *P. quasigrandis* Chace, 1985 materials from India and the Philippines can be as high as 5.8–8.4% (Chakraborty *et al.* 2015). In views that *P. ensis/reflexa* have a very wide geographical range with highly variable morphological characters (Chan & Crosnier 1997; Fransen 2006) but no coloration difference (Fig. 3A, B as compared to Kuberan *et al.* in press: fig. 1 and the color description of the Indian material provided by Suseelan & Mohamed 1968); and the recent finding of the development of the pereopodal epipods can be rather variable in certain species of *Plesionika* (see Chan 2016), it may be more appropriate to erect further new species in the “*P. ensis*” group after a full revision on the group. For example, the recent review by Ahamed *et al.* (2017: tables 1, 2 and key) on Asian species of *Plesionika* included both *P. ensis* and *P. reflexa*. However, *P. ensis* was omitted in their systemic account. It is likely that all previous reports of *P. ensis* from India (e.g. Alcock & Anderson, 1899; Alcock, 1901; George 1966; Suseelan & Mohamed 1968; Suseelan 1974; Kurup *et al.* 2008; Rajool Shanis *et al.* 2012; Radhakrishnan *et al.* 2012; Samuel *et al.* 2016) represented the present form. Examination of more specimens from India, as well as the other parts of the Indian Ocean, will be necessary to fully understand the development of pereopodal epipods and the variations in the length of the dactyli of the posterior pereopods in the Indian Ocean material.

Acknowledgements

We sincerely thanks L. Corbari and P. Martin-Lefevre of the Muséum national d'Histoire naturelle, Paris (MNHN) for sending the tissues of a Guadeloupe *P. ensis* specimen for sequencing work, J. Poupin of Institut de Recherche de l'École Navale, Brest, for allowing us to use the photograph of the same Guadeloupe specimen, C.C. Lin of National Taiwan Ocean University (NTOU) for preparing the line-drawings. The AURORA 2007 expedition, onboard M/V DA-BFAR, was a collaboration between the MNHN (Principal investigator, Philippe Bouchet), the Philippines National Museum (Principal investigator, Marivene Manuel-Santos), the Philippines Bureau of Fisheries and Aquatic Resources (BRAR), the Smithsonian Institution, the Aurora State College of Technology, National University of Singapore, the University of San Carlos (USC) and NTOU. The cruise was affiliated with the Census of Margins (CoMarges) component of the Census of Marine Life, and we acknowledge funding from the Lounsbery Foundation and the Total (Philippines) Corporation. Malcolm Sarmiento, former director of BFAR, kindly made M/V DA-BFAR available, and Noel Saguil (USC Associate) organized the logistics. We also thank the Department of Science and Technology, India for a financial grant towards the Fast Track Scheme for Young Scientists (SR/FT/LS-73/2012, SERB), and the Director of CMFRI for the facilities provided and encouragements. This work was supported by grants from the Ministry of Science and Technology, Taiwan, R.O.C.

References

- Ahamed, F., Cardoso, I.A., Ahmed, Z.F., Hossain, M.Y. & Ohtomi, J. (2017) An overview of the genus *Plesionika* Bate, 1888 (Decapoda, Caridea, Pandalidae) in Asian waters. *Zootaxa*, 4221 (5), 575–593.
<https://doi.org/10.11646/zootaxa.4221.5.6>
- Alcock, A. (1901) *A descriptive catalogue of the Indian deep-sea Crustacea Decapoda Macrura and Anomala, in the Indian museum. Being a revised account of the deep-sea species collected by the Royal Indian marine survey ship "Investigator"*. Indian Museum, Calcutta, 286 pp., 3 pls.
- Alcock, A. & Anderson, A.R. (1899) Natural history notes from H.M. royal Indian marine survey ship "Investigator," Commander T.H. Heming, R.N., Commanding. Series 3. Number 2. An account of the deep-sea Crustacea dredged during the surveying season of 1897–98. *Annals and Magazine of Natural History*, 3, 1–27, 278–292.
<https://doi.org/10.1080/00222939908678071>
- Bate, C.S. (1888) Report of the Crustacea Macrura collected by the *Challenger* during the years 1873–1876. *Report on the Scientific Results of the Voyage of H. M. S. Challenger during the years 1873–1876*, 24, 1–942, pls. 1–157.
- Calman, W.T. (1939) Crustacea: Caridea. In: *The John Murray Expedition 1933–34 Scientific Reports*, 6 (4), 183–224.
- Chace, F.A. Jr. (1985) The Caridean shrimps (Crustacea: Decapoda) of the *Albatross* Philippine Expedition, 1907–1910, Part 3: Families Thalassocarididae and Pandalidae. *Smithsonian Contributions to Zoology*, 411, 1–143.
<https://doi.org/10.5479/si.00810282.411>
- Chakraborty, R.D., Chan, T.Y., Maheswarudu, G., Kuberan, G., Purushothaman, P., Chang, S.C. & Jomon, S. (2015) On *Plesionika quasigrandis* Chace, 1985 (Decapoda, Caridea, Pandalidae) from southwestern India. *Crustaceana*, 88 (7–8), 923–930.
<https://doi.org/10.1163/15685403-00003451>
- Chan, T.Y. (2016) New records of the rare pandalid shrimp *Plesionika exigua* (Rathbun, 1906) (Crustacea: Decapoda: Caridea) in the western Pacific. *Zootaxa*, 4205 (1), 97–100.
<https://doi.org/10.11646/zootaxa.4205.1.11>
- Chan, T.Y. & Crosnier, A. (1991) Crustacea Decapoda: Studies of the *Plesionika narval* (Fabricius, 1787) group (Pandalidae), with descriptions of six new species. In: Crosnier, A. (Ed.), Résultats des Campagnes MUSORSTOM, vol. 9. *Mémoires du Muséum national Histoire naturelle*, 152, 413–461.
- Chan, T.Y. & Crosnier, A. (1997) Crustacea Decapoda: Deep sea shrimps of the genus *Plesionika* Bate, 1888 (Pandalidae) from French Polynesia, with descriptions of five new species. In: Crosnier, A. (Ed.), Résultats des Campagnes MUSORSTOM. Vol. 18. *Mémoires du Muséum National d'Histoire naturelle*, 176, 187–234.
- Chan, T.Y. & Yu, H.P. (2000) A new deep-sea shrimp of the genus *Plesionika* Bate, 1888 (Crustacea: Decapoda: Pandalidae) from Taiwan. *National Taiwan Museum Special Publication Series*, 10, 119–127.
- De Grave, S. & Franssen, C.H.J.M. (2011) Carideorum catalogus: The recent species of the Dendrobranchiate, Stenopodidean Procarididean and Caridean shrimps (Crustacea: Decapoda). *Zoologische Mededelingen*, 85 (9), 195–589.
- Fabricius, J.C. (1787) *Mantissa Insectorum sistens eorum species nuper detectas adjectis characteribus genericis, differentiis specificis, emendationibus, observationibus. Vol. 1*. Christ. Gottl. Prof., Hafniae (t), xx + 348 pp.
<https://doi.org/10.5962/bhl.title.36471>
- Franssen, C.H.J.M. (2006) Pandalidae (Crustacea: Decapoda) of the SONNE, VALDIVIA and METEOR expeditions 1977–1987 to the Red Sea and the Gulf of Aden. *Senckenbergiana maritima*, 36 (1), 51–82.
<https://doi.org/10.1007/BF03043702>
- George, M.J. (1969) Prawn fisheries of India, Part II: Systematics-Taxonomic consideration and general distribution. *Central*

- Hanamura, Y. & Takeda, M. (1987) Family Pandalidae (Crustacea, Decapoda, Caridea) collected by the RV "SOELA" from the Northwest Australian Shelf. *Bulletin of the National Science Museum, Tokyo*, (A), 13 (3), 103–121.
- Hajibabaei, M., Singer, G.A.C., Hebert, P.D.N. & Hickey, D.A. (2007) DNA barcoding: how it complements taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics*, 23 (4), 167–172.
<https://doi.org/10.1016/j.tig.2007.02.001>
- Hayashi, K.I. (1986) Penaeoidea and Caridea. In: Baba, K., Hayashi, K.I. & Toriyama, M. (Eds.), *Decapod Crustaceans from continental shelf and slope around Japan*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 38–149, 232–279.
- Hebert, P.D.N., Cywinska, A., Ball, S.L. & de Waard, J.R. (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B*, 270, 313–321.
<https://doi.org/10.1098/rspb.2002.2218>
- Holthuis, L.B. (1993) *The Recent genera of the caridean and stenopodidean shrimps (Crustacea, Decapoda), with an appendix on the order Amphionidacea*. Nationaal Natuurhistorisch Museum, Leiden, 328 pp.
- Kemp, S. (1925) Notes on Crustacea Decapoda in the Indian museum, XVII: On various Caridea. *Records of the Indian Museum*, 27 (4), 249–343.
- Kuberan, G., Chakraborty, R.D., Purushothaman, P. & Maheswarudu, G. (2018) *Plesionika reflexa* Chace, 1985 (Decapoda: Pandalidae) a new record of deep-sea caridean shrimp from the south-west coast of India. *Marine Fisheries Information Service, Technical and Extension Series*. [in press]
- Kurup, B.M., Rajasree, R. & Venu, S. (2008) Distribution of deep sea prawns off Kerala. *Journal of the Marine Biological Association of India*, 50 (2), 122–126.
- Li, X.Z. (2006) Report on some pandalid shrimps from the East China Sea (Decapoda Caridea). *Crustaceana*, 79 (11), 1281–1296.
<https://doi.org/10.1163/156854006779277303>
- Li, X. & Chan, T.Y. (2013) Pandalid shrimps (Crustacea, Decapoda, Caridea) collected from the Philippines PANGLAO 2005 deep-sea expedition. In: Ahyong, S.T., Chan, T.Y., Corbari, L. & Ng, P.K.L. (Eds.), *Tropical Deep-sea Benthos*, vol. 27. *Mémoires du Muséum national d'Histoire naturelle*, 204, pp. 129–154.
- Li, X. & Komai, T. (2003) Pandaloid shrimps from the northern South China Sea, with description of a new species of *Plesionika* (Crustacea: Decapoda: Caridea). *Raffles Bulletin of Zoology*, 51, 257–275.
- Matzen de Silva, J., Dos Santos, A., Cunha, M.R., Coata, F.O., Creer, S. & Carvalho, G.R. (2013) Investigating the molecular systematic relationships amongst selected *Plesionika* (Decapoda: Pandalidae) from the Northeast Atlantic and Mediterranean Sea. *Marine Ecology*, 34, 157–170.
<https://doi.org/10.1111/j.1439-0485.2012.00530.x>
- Milne-Edwards, A. (1881) Description de quelques Crustacés Macroures provenant des grandes profondeurs de la mer des Antilles. *Annales des Sciences Naturelles, Zoologie, Series 6*, 11 (4), 1–16.
- Milne-Edwards, A. (1883) *Recueil de figures de crustacés nouveaux oupeuconnus*. Published by the Author, Paris, 3 pp., 44 pls.
- Poupin, J. & Corbari, L. (2016) A preliminary assessment of the deep-sea Decapoda collected during the KARUBENTHOS 2015 expedition to Guadeloupe Island. *Zootaxa*, 4190 (1), 1–107.
<https://doi.org/10.11646/zootaxa.4190.1.1>
- Radhakrishnan, E.V., Deshmukh, V.D., Maheswarudu, G., Josileen, J., Dineshababu, A.P., Philipose, K.K., Sarada, P.T., Pillai, S.L., Saleela, K.N., Chakraborty, R., Dash, G., Sajeev, C.K., Thirumilu, P., Sridhara, B., Muniyappa, Y., Sawant, A.D., Vaidya, N.G., Johny, R.D., Verma, J.B., Baby, P.K., Unnikrishnan, C., Ramachandran, N.P., Vairamani, A., Palanichamy, A., Radhakrishnan, M. & Raju, B. (2012) Prawn fauna (Crustacea: Decapoda) of India – An annotated checklist of the penaeoid, segestoid, stenopodid and caridean prawns. *Journal of Marine Biological Association of India*, 54 (1), 50–72.
- Rajool Shanis, C.P., Akhilesh, K.V., Manjebraayakath, H., Ganga, U. & Pillai, N.G.K. (2012) Shrimps of the family Pandalidae (Caridea) from Indian waters, with new distributional record of *Plesionika adensameri* (Balss, 1914). *Journal of the Marine Biological Association of India*, 54 (1), 45–49.
- Samuel, V.K.D., Sreeraj, C.R., Krishnan, P., Parthiban, C., Sekar, V., Chamundeeswari, K., Immanuel, T., Shesdev, P., Purvaja, R. & Ramesh, R. (2016) An updated checklist of shrimps on the Indian coast. *Journal of Threatened Taxa*, 8 (7), 8977–8988.
<https://doi.org/10.11609/jott.2628.8.7.8977-8988>
- Suseelan, C. (1974) Observations on the deep-sea prawn fishery off the South-West coast of India with special references to pandalids. *Journal of the Marine Biological Association of India*, 16 (2), 491–511.
- Suseelan, C. & Mohamed, K.H. (1969) On the occurrence of *Plesionika ensis* (A. Milne Edwards) (Pandalidae, Crustacea) in the Arabian Sea with notes on its biology and fishery potentialities. *Journal of the Marine Biological Association of India*, 10 (1), 88–94.
- Takeda, M. & Hanamura, Y. (1994) Deep-sea shrimps and lobsters from the Flores Sea collected by the RV HAKUHO-MARU during KH-85-1 cruise. *Bulletin of the National Science Museum, Tokyo*, (A), 20 (1), 1–37.
- Tiefenbacher, L. (1992) Beiträge zur Kenntnis der Natantia des Arabischen Meeres und zu ihrer horizontalen und vertikalen Verbreitung unter Berücksichtigung der Sauerstoffminimum-Schicht. *Spixiana*, 15 (2), 113–136.
- Yang, C.H., Chan, T.Y. & Chu, K.H. (2010) Two new species of the “*Heterocarpus gibbosus* Bate, 1888” species group (Crustacea: Decapoda: Pandalidae) from the western Pacific and north-western Australia. *Zootaxa*, 2372, 206–220.