

New species, reports, observations and taxonomical changes of southern African rust fungi (Uredinales)

Reinhard Berndt · Elisabeth Uhlmann

Received: 10 February 2006 / Revised: 29 June 2006 / Accepted: 30 June 2006 / Published online: 15 August 2006
© German Mycological Society and Springer 2006

Abstract This work presents research on the diversity of the southern African rust mycobiota (Uredinales). It describes new species, lists new reports and adds new information on several rust fungi. *Puccinia cornurediata*, *Puccinia dioscoreae-mundtii*, *Puccinia horti-kirstenboschi*, *Puccinia othonnoides*, *Puccinia rapipes*, *Puccinia subindumentana*, *Uredo otholobii* and *Uromyces lotononidicola* are described as new; *Puccinia verwoerdiana* is assigned to *Puccinia lycii* as a synonym, and *Uredo lotononi* to *U. lotononidicola*. Comprehensive accounts and keys are presented for *Puccinia* species on *Lycium* (Solanaceae), *Helichrysum* and *Othonna* (Asteraceae). *Puccinia butleri* and *Uromyces bidenticola* are new reports for South Africa, and *Puccinia spinulosa* is new for Namibia. So far, the latter species has only been known from Madagascar, and *P. butleri* from the Indian subcontinent. Taxonomical novelties are *P. cornurediata* R. Berndt; *P. dioscoreae-*

mundtii R. Berndt, A.R. Wood & E. Uhlmann; *P. horti-kirstenboschi* R. Berndt & E. Uhlmann; *P. othonnoides* R. Berndt, A.R. Wood & E. Uhlmann; *P. rapipes* R. Berndt & E. Uhlmann; *P. subindumentana* R. Berndt; *U. otholobii* R. Berndt, A.R. Wood & E. Uhlmann and *U. lotononidicola* R. Berndt

Introduction

Our knowledge of the taxonomy and diversity of South African rust fungi (Basidiomycota, Uredinales) is still mainly based on a series of papers (compare Doidge 1950) published by the eminent South African mycologist Ethel M. Doidge (1887–1965). Since her time, new species and observations have been reported only erratically, although a commendable checklist of South African phytopathogenic fungi was compiled (Crous et al. 2000 onwards). Additionally, important contributions were made to the understanding of the biology of certain rust fungi to use them as biocontrol agents against plants indigenous to South Africa but which have become invasive weeds elsewhere (e.g. Kleinjan et al. 2004; Morris 1982; Wood 2002; Wood et al. 2004). Recently, a research project was initiated to investigate the composition of the rust mycobiota of southwestern Africa (Berndt et al. 2002). So far, this has led to the publication of findings of several new species and observations on South African rust fungi (e.g. Mennicken et al. 2003, 2005). With the present paper we aim to contribute to the knowledge of the southern African rust mycobiota in several respects: (1) By describing new species and reporting new findings from South Africa, we emphasise the importance and necessity of continuing to survey the area mycologically. (2) We present “micro-monographs” of groups of rust species on selected host taxa

R. Berndt (✉)
Herbarium turicense, Institute of Integrative Biology (IBZ),
ETH Zurich, CHN,
Universitätsstr. 16,
CH-8092 Zurich, Switzerland
e-mail: reinhard.berndt@env.ethz.ch

E. Uhlmann
Systematic Botany and Mycology, University of Tübingen,
Botanical Institute,
Auf der Morgenstelle 1,
D-72076 Tübingen, Germany
e-mail: elisabeth.uhlmann@uni-tuebingen.de
e-mail: elisabeth.uhlmann@ufz.de

Present address:
E. Uhlmann
Department Soil Ecology (BOOEK),
UFZ Centre for Environmental Research Leipzig-Halle,
Theodor-Lieser-Strasse 4,
D-06120 Halle, Germany

with detailed species descriptions, keys and illustrations. These treatments are intended to contribute to the future project of a rust flora of the region.

Materials and methods

Spores and hand sections of herbarium material were mounted in lactophenol and gently heated to boiling. The preparations were examined with a C. Zeiss "Axiphof" light microscope and photographs were taken with a C. Zeiss MC-80 camera on Kodak Ektachrome 64 Professional slide film. All micrographs were taken using differential interference contrast optics. At least 30 spores were measured for each spore stage; exceptions are mentioned in the descriptions. The arithmetic means are given after the ranges of measurements (in brackets). Names of herbaria are abbreviated by their acronyms according to *Index herbariorum* (Stafleu et al. 1981). The rust species are listed under their respective host families, which are ordered alphabetically.

Results

On Acanthaceae

Puccinia species with loculate telia on Acanthaceae

Several *Puccinia* spp. on members of Acanthaceae are characterised by loculate telia: *Puccinia mackenensis* Cumm. (including *Puccinia boerhaviaefoliae* Thirum.), *Puccinia multiloculata* Cumm., *Puccinia namibiana* Mennicken et al., *Puccinia semiloculata* Laundon and *Puccinia thunbergiae* Cooke. These species are very similar in the telial stage but reveal characters in the aecial stage that may help to distinguish them.

The aecidia of *P. multiloculata* (holotype and PUR 16117 et 16116) and *P. thunbergiae* (IMI 61368a) were virtually identical. The aecidiospores revealed numerous "light refractile bodies" (or "pore plugs") which were approximately muffin-shaped and detached easily from the spore surface (Fig. 1a). Both species are also very similar in the telial stage and may not be specifically different. As we could not examine type material of *P. thunbergiae*, we prefer to keep them separate, however. Light refractile bodies were not present in *P. namibiana* or *P. mackenensis*. In *Puccinia blepharidis* P. Henn., a species on *Blepharis* with non-loculate telia, we found aeciospores with isolated coarse granules which did not assume the size of the described globules and only occasionally separated from the spore surface. Light refractile bodies have been described in unrelated groups




Fig. 1 **a** *Puccinia multiloculata* (type), aeciospores with numerous, more or less muffin-shaped light refractile bodies. Scale bar=10 µm. **b** *Puccinia horti-kirstenboschi* (type), aeciospores. Scale bar=10 µm. **c** *Puccinia horti-kirstenboschi* (type), peridial cells; note the finely granular surface. Scale bar=10 µm

of *Puccinia/Uromyces*, and their value for systematic purposes is probably restricted (Berndt 2004; Holm 1966; Sato and Sato 1982).

Puccinia namibiana was described for the reason that the peridial cells of the aecia had very thick (ca. 9–13 µm) outer walls (Mennicken et al. 2005). *Aecidium acanthopsidis* Syd. & P. Syd. was listed as a synonym. We measured only 5.5–8 µm for the thickness of the external periclinal walls of the peridial cells in the type specimen. Additionally, we found them to be verruculose rather than finely striate as described. These peridial characters overlap or coincide with those of other *Puccinia* species on Acanthaceae (Laundon 1963) and do not seem to suffice to delimit *P. namibiana* as a separate species. One should note here that the thickness of the walls of the aecial peridial cells can be variable, as Mayus (1904) observed in the field and Iwanoff (1907) demonstrated experimentally that the thickness of peridial walls varied considerably within in a single rust species according to the light exposition of the infected plants.

We found that *P. namibiana* differs from the other species with loculate telia by certain teliospore characters (compare key). The differences are subtle, however, and it appears that the species are closely related. We did not investigate whether *Ae. acanthopsidis* really is a synonym of *P. namibiana*.

Key to the *Puccinia* species with loculate telia on Acanthaceae:

- 1 Telia distinctly loculate (teliospores produced in “chambers” bounded by densely aggregated, slenderly cylindrical paraphyses), more or less compact, spores on average broader than 15 µm
- 2 Aeciospores with conspicuous, “muffin-shaped” globules
- 3 Teliospore pedicels generally darker than the spore wall—*P. multiloculata*
- 3* Teliospore pedicels coloured like spores or paler—*P. thunbergiae*
- 2* Aeciospores without such globules
- 4 Teliospore wall light chestnut brown, 1.5–2 µm thick at sides—*P. makenensis*
- 4* Teliospores paler, 1–1.5 µm thick at sides—*P. namibiana*
- 1* Telia loculate, but not distinctly so, and thus appearing “velvety”, teliospores 10–15 µm broad, with a pale spore wall—*P. semiloculata*

Material examined: *Puccinia blepharidis*: Africa, Angola (?), am Knebe bei Manonge, on *Blepharis buchneri*, leg. H. Baum (no. 835 et 855), 21/23 April 1900 (type, Z+ZT). *Puccinia makenensis*: Africa, Sierra Leone, Makene, on *Blepharis maderaspatensis* Heyne ex Roth, leg F.C. Deighton (no. 1741), 28 January 1939 (holotype, PUR 9558). *Puccinia multiloculata*: Africa, Sierra Leone, Segbwama, on *Thunbergia cynanchifolia* Benth., leg. F.C. Deighton (no. 1460), 11 December 1937 (holotype, PUR

9568). Gold Coast (=Ghana), Kumasi, on *Justicia? insularis*, leg. L. Piening (no. 2216 et 2319), 17 January 1956 et 4 April 1956 (PUR 16117 et 16116 ex IMI 62148a et 63544a). *Puccinia namibiana*: Africa, Namibia, between Okahandja and Wilhelmstal, on *Blepharis obmitrata* C.B. Clarke, leg. M. Mennicken, 6 April 2002 (type, to be deposited in PREM). *Puccinia thunbergiae*: Africa, Gold Coast (=Ghana), on *Justicia* sp., leg. L. Piening, 1955 (IMI 61368a). Uganda, Kiboga, on *Asystasia schimperi* T. Anders., leg. G. Hakiza, 19 August 1988 (IMI 327948).

On Asteraceae

Puccinia species on *Helichrysum* (*Gnaphalieae*) in Africa

Recent collections of *Puccinia* rust on *Helichrysum* spp. from South Africa could not be assigned to known species readily. A study of the relevant rusts led to the recognition of three new species, *Puccinia horti-kirstenboschi*, *Puccinia subindumentana* and *P. cornurediata*.




Fig. 2 *Puccinia horti-kirstenboschi* (type), teliospores. Scale bar=20 µm

Puccinia horti-kirstenboschi R. Berndt & E. Uhlmann, sp. nov. (Figs. 1b,c and 2).

Etymology: Named after the collection site close to Kirstenbosch Botanical Garden.

Spermogonia typi 4 adsunt. Aecia aecidiomorpha, sparsa vel laxe aggregata in maculis atro-brunneis, non vel paulum hypertrophis paginae abaxialis foliorum; peridio albo, anguste cylindrico vel inaperte subconico, ca. 0.5–1 mm longo et 0.2–0.3 mm lato, apicaliter aperenti et laceranti, interdum longitudinaliter inciso; cellulis peridii hyalinis, intus verrucis subtilissimis verruculosis vel fere subreticulatis, ca. 4–5 µm crassis, extus verrucis inconspicuis, humilibus praeditis et ca. 3–5 µm crassis; aeciosporae subangulariter subglobosae, globosae vel late ellipsoideae, 22–26.5×19.5–22.5 µm (medium 24.1×21.0 µm), pariete hyalino, ca. 1.5 µm crasso (including ornamentum), verrucis moderate grossis, ca. 1 µm diam. dense obsito; poris germinationis non visis. Telia abaxialia in foliis, sub tomento foliorum occulta, minuta, rotundata, pulvinata, laxe fibroso-carnosa; teliosporae ellipsoideae, oblongae vel, rariore, late ellipsoideae vel subclavatae, secundum septum leniter vel moderate constrictae, 40–52 (55)×(17.5) 20–23 µm (medium 45.4×21.0 µm), pariete levi, dilute ochraceo ad stramineo, ca. 0.5–1 µm crasso, in apice usque ad 1.5 µm, poris germinationis apicaliter in cellula distali, septum juxta in cellula proximali, apapillatis; teliosporae post maturitatem basidiis germinantes, pedicellis brevibus, hyalinis, tenui tunicatis praeditae. Mesosporae interdum adsunt.

In foliis *Helichrysi* sp. (Asteraceae).

Spermogonia of type 4 present. Aecia *Aecidium*-like, abaxially on leaves, singly or in loose groups on blackish-brown spots, causing no or only slight hypertrophy of leaf tissue; peridium white, slenderly cylindrical or subconical when still closed, ca. 0.5–1 mm long and 0.2–0.3 mm wide, opening and lacerating irregularly at the apex, sometimes longitudinally incised; peridial cells delicately verruculose to subreticulate on the inner surface and finely verrucose on the outer surface by low, inconspicuous warts, inner cell wall ca. 4–5, outer wall ca. 3–5 µm thick; aeciospores subangularly subglobose, globose or broadly ellipsoidal, 22–26.5×19.5–22.5 µm (mean 24.1×21.0 µm), spore wall hyaline, ca. 1.5 µm thick (including ornament), densely covered by moderately coarse, flat-topped warts with an irregular, subpolygonal outline and about 1 µm in diameter, germ pores not seen. Telia present abaxially on leaves, hidden under the wooly indument, scattered, tiny, rounded and pulvinate, ferruginous, soft and somewhat sticky; teliospores ellipsoidal, oblong, more rarely broadly ellipsoidal or subclavate, slightly to moderately constricted at the septum, 40–52 (55)×(17.5) 20–23 µm (mean 45.4×21.0 µm), spore wall smooth, light ochraceous or straw-coloured, ca. 0.5–1 µm thick, to 1.5 µm at the apex, germ pores apical in distal cells and close to septum in

proximal cells, without papillae, spores germinating with basidia upon maturity, pedicels short, reaching up to spore length, hyaline, thin-walled. One-celled mesospores occurred occasionally.

On leaves of *Helichrysum* sp. (Asteraceae).

Holotype (PREM): South Africa, Western Cape Province, Cape Peninsula, at Klaasen Road adjacent to Kirstenbosch Botanic Garden, on *Helichrysum* sp. (Asteraceae), leg. R. Berndt and E. Uhlmann, 2 November 2004 (isotype Z+ZT).

Puccinia horti-kirstenboschi differs from *P. cornurediata* R. Berndt, *Puccinia kalchbrenneri* De Toni vars., *Puccinia macowani* Winter and *Puccinia rocherpaniana* Mennicken & Oberw. and *P. subindumentana* R. Berndt in teliospores (Fig. 2), which are not apically thickened and from *P. pienarii* Pole Evans in the smooth and thin teliospore wall. Uredinia were not observed in the present rust and are probably lacking. The species was collected at several other sites in the Western Cape Province and may be quite common.





Fig. 3 *Puccinia subindumentana* (type), teliospores. Scale bar=20 µm

Fig. 4 **a** *Puccinia subindumentana* (type), aeciospores. Scale bar=10 µm. **b** *Puccinia subindumentana* (type), peridial cells; note the finely granular surface. Scale bar=10 µm. **c** *Puccinia cornurediata* (type), aeciospores. Scale bar=10 µm. **d** *Puccinia cornurediata* (type), distal part of uredinial peridium with enclosed urediniospores. Scale bar=10 µm



Puccinia subindumentana R. Berndt, sp. nov. (Figs. 3 and 4a,b).

Etymology: Indicating that the sori are hidden under the indumentum of the leaves.

Aecia aecidioidea, in maculis violaceis ad brunneis paginae abaxialis foliorum aliquot aggregata vel singulatim sparsa,

peridio cylindrico, eburneo ad ochraceo, longitudinaliter inciso, copia sporarum ochroleuco; aeciosporae subglobosae vel globosae, subangulares, 27–33×25–30 µm (medium 29.1×27.4 µm), pariete hyalino, ca. 2–3 µm crasso, verrucis irregulariter rotundatis vel subpolygonalibus, ca. 0.8–1.5 µm latis et ca. 0.8–1.5 µm altis dense vel densissime obsito;

cellulae peridii extus subtilissime subreticulatae ad subfoveolatae verrucis delicatissimis, basaliter inter se coniunctis, intus ornamento simili vel leniter grossiori ornatae. Telia in pagina abaxiali foliorum aliquot aggregata vel singulatim sparsa, rotundata et pulvinata, 0.3–0.5 mm diam., primum armeniaca, deinde ferruginea et post germinationem teliosporae basidiis pruinosa, sub indumento foliorum subocculta; teliosporae fusiformes vel ellipsoideae, secundum septum non vel leniter constrictae, basaliter vel leniter oblique pedicellatae pedicellis hyalinis, delicatis, tenui tunicatis, usque ad 55 µm longis, (36) 41–68 (71)×16–22.5 µm (medium 53.7×18.7 µm), pariete levi, stramineo ad dilute aureo, ca. 1 µm crasso, incrassato secundum septum, apicaliter (sub)lateraliter usque ad 10 µm incrassato, poris germinationis (sub)apicalibus et septum juxta.

In foliis *Helichrysi chrysophori* (Asteraceae).

Aecia *Aecidium*-like, scattered on violet to light brown spots of abaxial leaf surface, in small groups or singly on thickened areas, with a cylindrical, whitish to ochraceous, longitudinally incised peridium, spore mass cream-coloured; aeciospores subglobose or globose, subangular by mutual pressure, 27–33×25–30 µm (mean 29.1×27.4 µm), spore wall hyaline, ca. 2–3 µm thick including the irregularly rounded or subpolygonal, densely situated warts which are ca. 0.8–1.5 µm high and 0.8–1.5 µm in diameter, warts sometimes confluent and then forming almost smooth areas; outer surface of peridial cells very finely subreticulate to almost foveolate by basally coalescing very fine warts, inner surface with similar, but slightly coarser, ornament. Telia scattered or in small groups on the abaxial leaf surface, first apricot to orange, later ferruginous and pruinose after germination of teliospores, pulvinate, rounded, 0.3–0.5 mm in diameter, subcompact, almost hidden under the indumentum of the leaves; teliospores fusiform to ellipsoidal, not or slightly constricted at the septum, basally stalked by a delicate, thin-walled, hyaline, collapsing pedicel up to 55 µm long, sometimes the pedicel is slightly offset, (36) 41–68 (71)×16–22.5 µm (mean 53.7×18.7 µm), spore wall smooth, straw-coloured to light golden, ca. 1 µm thick, thickened around septum and to 10 µm in the most often laterally situated thickening of the spore apex, germ pores (sub)apical and at septum.

On leaves of *Helichrysum chrysophorum* (Asteraceae).

Holotype: Africa, Nyasaland (=Malawi), Mlanje Mt., on *H. chrysophorum* S. Moore, leg. P.O. Wiehe (as “Niehe”), 17 November 1949 (IMI 45352, sub *Aecidium helichrysi*).

The present species is similar to *P. horti-kirstenboschi* by the pulvinate, apricot to ferruginous telia and the pallid teliospores. The teliospores are longer and apically thickened (Fig. 3), however, and the larger aeciospores (Fig. 4a) have thicker walls and coarser warts. The aecial stage is also different from *Ae. helichrysi* by larger aeciospores with coarser warts.




Fig. 5 *Puccinia cornurediata* (type), teliospores. Scale bar=20 µm

Puccinia cornurediata R. Berndt, sp. nov. (Figs. 4c,d and 5).

Etymology: Named after the corniculate peridium of the uredinia.

Spermogonia typi 4 adsunt, adaxialia, aeciis opposita. Aecia aecidioidea, maculo pallescenti insidentia, peridio albo, cylindrico, laceranti praedita; cellulae peridii extus labyrinthoide delicate striatae, intus verrucis moderate grossis, coalescentibus et superficiem irregulariter labyrinthicam ad foveolatam efficientibus; aeciosporae irregulariter subglobosae, vel late ellipsoideae, cum ornamento 29–36 (41)×23–31 µm (medium 32.3×27.9 µm), pariete hyalino, cum ornamento ca. 2–2.5 µm crasso, verrucis cylindricis, ca. 0.5–1 µm altis, usque ad 1.5 µm latis moderate dense obsito. Uredinia in maculis pallescentibus paginae abaxialis foliorum singulatim laxeque sparsa, ampullacea, peridio tenui, aurantio-flavo, ca. 0.7–1 mm longo et ca. 0.08–0.12 mm lato, hyphis arctissime coalitis, linearibus vel fusiformibus, longitudinalibus, aureis, crassule tunicatis, in apice subacutis composito; urediniosporae pedicellatae, late ellipsoideae, late obovoideae ad subglo-

bosae, 28–34.5×24.5–29.5 µm (medium 31.2×27.3 µm), pariete hyalino, ca. 2 µm crasso, aequaliter et moderate sparse echinulatae spinis ca. 1 µm longis et 2–4 µm inter se distantibus, poris germinationis non visis. Telia in gregibus parvis maculis pallescentibus paginae abaxialis foliorum insidentia, nitide plumbea quamdiu epidermide tecta, postea ferruginea, mollia ad pulverulenta; teliosporae ellipsoideae vel subclavatae, saepe leniter incurvatae vel deformes, cellulis distalibus apicaliter subacutis vel rotundatis, cellulis proximalibus basim versus attenuatis vel plus minusve rotundatis, secundum septum leniter, rariore paullulum vel distincte constrictae, (48) 50–69 (80)×20.5–25.5 (27) µm (medium 58.3×23.5 µm), pariete levi, aureo, vel dilute aureo in cellula proximali et poro germinationis apicali, ca. 1 µm crasso, usque ad 4–8 µm incrassato in apice, poris germinationis apicalibus et juxta septum.

In foliis *Helichrysi petiolati* (Asteraceae).

Spermogonia of type 4 present, adaxial, opposite aecia. Aecia *Aecidium*-like, in small groups (always?) on round, bleached leaf spot, peridiate with a white, cylindrical, lacerating peridium, peridial cells with a delicate, labyrinthine pattern of striae on outside and moderately coarse warts on the inner side which generally coalesce to form an irregular labyrinthine or almost foveolate surface; aeciospores irregularly subglobose or broadly ellipsoidal, including ornament 29–36 (41)×23–31 µm (mean 32.3×27.9 µm), spore wall hyaline, ca. 2–2.5 µm thick (including ornament), rather densely or moderately densely covered by cylindrical warts ca. 0.5–1 µm high and up to 1.5 µm in diameter. Uredinia singly and sparsely scattered on small bleached spots on abaxial side of leaves, flask-shaped, with a slender, tubular and slightly tapering, orange-yellow peridium ca. 0.7–1 mm long and ca. 0.08–0.12 mm wide, composed of firmly adherent linear or spindle-shaped, longitudinal, slightly thick- and ochraceous-walled hyphae ending in subacute apices; urediniospores pedicellate, broadly ellipsoidal, broadly obovoidal to subglobose, 28–34.5×24.5–29.5 µm (mean 31.2×27.3 µm), spore wall hyaline, about 2 µm thick, evenly and moderately sparsely echinulate, spines ca. 1 µm long and ca. 2–4 µm apart, germ pores not seen. Telia found on bleached leaf spots abaxially on leaves, pulvinate, ferruginous, or with plumbaceous lustre as long as they are covered by epidermis, soft to pulverulent; teliospores ellipsoidal or subclavate, often slightly bent or deformed by mutual pressure within sori, distal cells apically subacute to rounded, lower cells tapering into the pedicel or more or less rounded, slightly constricted at septum, sometimes hardly or more deeply constricted, (48) 50–69 (80)×20.5–25.5 (27) µm (mean 58.3×23.5 µm), spore wall smooth, golden brown or lighter pigmented in proximal cells and at spore apex, ca. 1 µm thick in proximal cells, thickening to 4–8 µm towards the apex in distal cells, germ pores apical and at the septum.

On leaves of *Helichrysum petiolatum* (Asteraceae).

Holotype: Africa, Nyasaland (=Malawi), Zomba Mt., on *H. petiolatum* D. Don. (as *Helichrysum petrolatum*), leg. P. O. Wiehe, 17 September 1950 (IMI 44327, sub *P. kalchbrenneri*).

This is a remarkable member of the genus *Puccinia*, belonging to a group of species sometimes classified in the separate genus *Miyagia* and characterised by peridiate uredinia and telia with or without peridia. It is well distinguished from the other members of *Miyagia*, *Miyagia anaphalis* Syd. & P. Syd. and *Miyagia macrospora* Hirats. f., by its very long and slender, corniculate uredinal peridium (Fig. 4d). *Miyagia anaphalis* and *M. macrospora* occur in Asia on *Anaphalis*, which belongs to tribe Gnaphalieae, as does *Helichrysum*. *Miyagia pseudosphaeria* (Mont.) Jørstad differs morphologically and grows on *Sonchus* (Lactuceae). In some specimens of *M. pseudosphaeria*, one-celled teliospores predominate (Wilson and Henderson 1966). *Corbulopsora* is a related genus differing from *Miyagia* by consistently one-celled teliospores. It is interesting to note that *Corbulopsora cumminsii* Thirum. occurs on *Lactuca*, like *Sonchus*, a member of Lactuceae, and is morphologically quite similar to *M. pseudosphaeria* (Jørstad 1956b).


Because of similar morphological traits and host relationships, we speculate that the species on *Helichrysum* and *Anaphalis* are more closely related and that *M. pseudosphaeria* and *C. cumminsii* on Lactuceae may be linked. A closer affinity with the remaining two *Corbulopsora* species on *Olearia* (tribe Astereae) is uncertain. As *Miyagia* and *Corbulopsora* may not circumscribe natural relationships, we prefer to retain the present rust in *Puccinia*.

Assuming a closer relationship of the *Miyagia*-like rust species on Gnaphalieae and not knowing additional locations between Malawi and Asia, these rusts may represent an astonishing southern African–Asian disjunction.

Puccinia kalchbrenneri De Toni 1888. Sacc. Syll. Fung. vol. VII:645 (Figs. 6a–d and 7a)

The aecial stage is probably represented by *Ae. helichrysi* Doidge (compare below). Uredinia predominantly abaxial on leaves, semi-immersed in host tissue, opening irregularly at apex and liberating the pulverulent, pallid cinnamon spore mass; urediniospores obovoidal, subglobose or broadly ellipsoidal, 24–32 (38)×20–27 µm [mean 25.7×23.2 µm in PREM 29862, 28.8×22.8 µm in PREM 26021 ($n=19$), 30.6×23.5 µm in PREM 23440], spore wall ca. 1–1.5 µm thick, subhyaline to straw-coloured, finely echinulate with slender, sharp spines about 2–3 µm apart, germ pores obscure, scattered. Telia are abaxial on leaves, loosely scattered on bleached areas, blackish brown, most often tiny and crust-like (almost phakopsoroid), surrounded by a layer of tangled, thick-walled, light brown hyphae,


Fig. 6 **a** *Puccinia kalchbrenneri* (type), teliospores. Scale bar=20 µm. **b** *Puccinia* cf. *kalchbrenneri* (Cape peninsula specimen), teliospores; note the knotty wall thickenings. Scale bar=20 µm. **c** *Puccinia kalchbrenneri* var. *valida* (PREM 12823), teliospores. Scale bar=20 µm. **d** *Puccinia kalchbrenneri* var. *valida* (PREM 23440), teliospores. Scale bar=20 µm



permanently (?) covered by the epidermis, more rarely bullate, with a compact texture and naked; teliospores 37–65 (76)×19–27 µm (mean 43.6×22.8 µm in PREM 29862, 50.6×23.7 µm in PREM 26021, 50.9×23.1 µm in Z+ZT, 58.3×24.2 in PREM 23440), ellipsoidal, broadly ellipsoidal or subclavate but often bent or deformed by mutual pressure within the densely packed sori, slightly to strongly constricted at septum, distal cells rounded to subapiculate,

proximal cells tapering towards the hilum, pedicels short, thin-walled, subhyaline to ochraceous, collapsing, spore wall light brown to light chestnut (generally more lightly pigmented at the apex and where the spore wall is thinner), smooth, 1–1.5 µm thick laterally, thickening to 8 µm in the apex, germ pores apical and close to the septum, usually with a shallow, conical pit. Mesospores rare or scattered.

Fig. 7 **a** *Puccinia kalchbrenneri* (PREM 32752), aeciospores. Scale bar=10 µm. **b** *Aecidium helichrysi* (PREM 10128), aeciospores. Scale bar=10 µm. **c** *Puccinia macowanii* (type), aeciospores. Scale bar=10 µm. **d** *Puccinia macowanii* (type), peridial cells of aecial peridium. Scale bar=10 µm



Material examined: South Africa, Transvaal, Zoutpansberg Distr., Piesanghoek, on *Helichrysum nudifolium* var. *quinquenerve* (Thunb.) Moes. (=*Helichrysum quinquenerve* Less.), leg. P. Watson, 1 June 1929 (PREM 29862). South Africa, Transvaal, Barberton Distr., on *H. quinquenerve*, leg. L.C.C. Liebenberg, June 1931 (PREM 26021). South Africa, Transvaal, Haenertsburg, on *H. nudifolium* var.

quinquenerve, leg. K. Putterill, 13 November 1938 (PREM 32752, only aecidia present, det. E.M. Doidge). South Africa, Natal, Port Shepstone Distr., Oribi Gorge, on *H. nudifolium* var. *quinquenerve*, leg. H. Schuepp, 16 May 1959 (Z+ZT). South Africa, Pretoria Distr., Silverton Ridge, on *H. nudifolium* var. *leiopodium* (DC.) Moes., leg. E.M. Doidge, 12 June (?) 1928 (PREM 23440). South

Africa, Transvaal, Olifantsfontein, on *Helichrysum coriaceum*, leg. Pienaar, 21 February 1920 (?) (PREM 12823, var. *valida*). Uganda, Kigezi, Mpalo, on *H. nudifolium* Less. (as *H. nudiflorum*), leg. C.G. Hansford, August 1937 (IMI 55385).

Puccinia kalchbrenneri may be a variable species. Doidge (1927) distinguished a variety *valida* for specimens with longer teliospores and a more strongly thickened teliospore apex. In a specimen assigned to the latter variety (PREM 12823), a few telia were discovered among many uredinia. The teliospores measured $48\text{--}63 \times 22\text{--}27$ (29) μm (mean $57.2 \times 24.7 \mu\text{m}$) and were apically thickened to 7 μm (Fig. 6c). This is within the observed range of var. *kalchbrenneri*. It is possible that the telium morphology (crustose or erumpent) could influence the shape and the size of the teliospores. If this is true, the separation of vars. *valida* and *kalchbrenneri* would be artificial. No type specimen was designated by Doidge (1927) in the protologue of *P. kalchbrenneri* var. *valida*, and a lectotype needs to be selected. This is not done here as none of the syntypes could be studied. A uredinial specimen from Uganda (IMI 55385) assigned to *P. kalchbrenneri* differed from the other specimens by considerably smaller urediniospores with thinner walls [$22\text{--}28$ (30.5) $\times 17\text{--}22.5 \mu\text{m}$ (mean $25.0 \times 19.3 \mu\text{m}$), spore wall ca. 1 μm thick]. It is doubtful whether it belongs to the present species.

We collected two specimens of *Puccinia* on *Helichrysum* spp. on the Cape Peninsula. One was telial, the other revealed uredinia and telia, as well as some aecia almost hidden in the dense tomentum of the lower leaf surface. The teliospores were virtually indistinguishable, and both specimens most probably belong to the same species. They were similar to the studied *P. kalchbrenneri* but showed some differences in all present spore stages: telia occurred on leaves or stems and were chestnut brown to blackish brown, pulvinate and subcompact. Teliospores measured $52\text{--}65 \times 20\text{--}27 \mu\text{m}$ (mean $59.0 \times 24.0 \mu\text{m}$) and (44) $51\text{--}73$ (76) $\times 21\text{--}27 \mu\text{m}$ (mean $61.1 \times 23.9 \mu\text{m}$). They were distinguished by a prominent apical thickening (8–13 μm) of the cell wall and additional knotty thickenings (Fig. 6b). The urediniospores were very similar to those of *P. kalchbrenneri* [$28\text{--}31$ (33) $\times 23\text{--}27 \mu\text{m}$, mean $29.6 \times 25.7 \mu\text{m}$] but were extremely fine and rather densely echinulate (ca. 1–2 μm between spines). The aeciospores were much more delicately verruculose than spores of *Ae. helichrysi* and had thinner spore walls [1.5–2 μm instead of 2–2.5 (3) μm]. Doidge (1927) suggested that *Ae. helichrysi* (Fig. 7b) belonged to the life cycle of *P. kalchbrenneri*; however, this remains unproven. The only evidence is a specimen of *Ae. helichrysi*, which occurred together with the uredium stage of *P. kalchbrenneri* on the same leaves. Despite the differences observed in the *Puccinia* from Cape Peninsula, we assign it to




Fig. 8 *Puccinia macowani* (type), teliospores. Scale bar=20 μm

P. kalchbrenneri. More material needs to be studied to evaluate the constancy or variability of the relevant characters.

Puccinia kalchbrenneri has also been reported from India based on a uredinal collection on *Helichrysum buddleoides* DC. (Ragunathan and Ramakrishnan 1972). If the determination is correct, this would be an example for a South African–Indian pattern of rust distribution.

Puccinia macowani Winter, *Puccinia pienaarii* Pole Evans and *P. rocherpaniana* Mennicken & Oberw. are the other *Puccinia* species occurring on *Helichrysum* in South Africa. *Puccinia pienaarii* is unmistakable in the telial stage by very irregularly thickened, bulging teliospore walls and more or less equatorial germ pores (Fig. 9). *Puccinia rocherpaniana* has thicker urediniospore walls than *P. kalchbrenneri* and broader teliospores. The teliospores of *P. macowani* are quite similar to those of *P. kalchbrenneri* while the aeciospores are entirely different (Fig. 7a,c). The teliospore apex is thickened and does not show a pit before germination but a subglobose to broadly ellipsoidal thickening (Fig. 8). The aeciospores are characteristically




Fig. 9 *Puccinia pienarii* (type), teliospores. Scale bar=20 µm

ornamented by flat and broad, button-like warts (Fig. 7c). Aeciospores of *P. horti-kirstenboschi* have similar warts that are smaller, however. The peridial cells of the aecial peridium are very finely warty to granular in *P. macowani* (Fig. 7d) and resemble those of *P. subindumentana*.

Key to the *Puccinia* species on *Helichrysum*:

- 1 Teliospores with very irregularly thickened, knotty or bulged cell wall, germ pores more or less equatorial; uredinia unknown—*P. pienarii*
- 1* Teliospores with other characters, germ pores apical and close to septum
- 2 Teliospores uniformly thin-walled, straw-coloured to ochraceous, aeciospores with flat, button-like warts, uredinia unknown—*P. horti-kirstenboschi*
- 2* Teliospores with apically thickened walls, straw-coloured to light brown (to chestnut)
- 3 Uredinia with a slenderly cylindrical to subconical peridium up to 1 mm long, composed of linear, golden, thick-walled hyphae—*P. cornurediata*
- 3* Uredinia not peridiate or unknown
- 4 Telia ferruginous, soft to subcompact; teliospores ellipsoidal to fusiform, with a subglobose or broadly

ellipsoidal thickening at germ pores; aeciospores with broad, flat, button-like warts, uredinia unknown—*P. macowani*

- 4* Teliospores apically thickened, with or without conical pits at the germ pores but not biconvexly thickened; aeciospores not with button-like warts, verruculose to verrucose or aecial stage absent
- 5 Telia apricot to ferruginous, erumpent; teliospores fusiform to ellipsoidal with straw-coloured wall and an oblique thickening at the apex, often conspicuously thickened in the angles of the septum—*P. subindumentana*
- 5* Telia dark brown to blackish, erumpent or crustose; teliospore wall darker coloured, apically thickened but normally not obliquely
- 6 Urediniospore wall 2–3 µm thick, yellow-brown, teliospores relatively broad (21–37 µm), aecia unknown—*P. rocherpaniana*
- 6* Urediniospore wall 1–2 µm thick, straw-coloured, teliospores narrower on average, aecial stage probably present (*Ae. helichrysi*)—*P. kalchbrenneri* (vars.)

Rust on *Othonna* (Senecioneae)

The genus *Othonna* comprises ca. 120 species and is centred in South Africa. It is host to *Uredo othonnae* Jørstad and *Puccinia othonnae* Doidge of which only the telial stage is known. We collected several *Puccinia* specimens on *Othonna* spp. in the Western Cape Province that revealed both uredinia and telia. To elucidate the identity of these specimens and to check a possible connection with *U. othonnae*, we made a comparison of the rusts. This resulted in the recognition of a new species, *Puccinia othonnoidea*, and a detailed evaluation of morphological characters particular to these rusts.

Puccinia othonnae Doidge 1927. Bothalia 2:203 (Fig. 10a–c)

Puccinia othonnae is known from *Othonna natalensis* Sch. Bip. in Transvaal (=Gauteng Province, NE South Africa). We studied the type and the paratype and present a description to supplement the diagnosis published by Doidge (1927).

Telia amphigenous on leaves, though predominant on the abaxial side, dark brown, rounded or irregularly shaped by confluence, ca. 0.3–2.4 mm in diameter, slightly pulvinate, pulverulent but with a somewhat fibrous texture as the pedicels appear to stabilise the spore mass; teliospores broadly ellipsoidal, rounded at both ends, not or very slightly constricted at the septum, 53–65 (68)×38–46 µm (mean 59.8×42.2 µm), pedicels stout, thick-walled, hyaline, up to 120 µm long, inserted basally but generally




Fig. 10 **a** *Puccinia othonnae* (type), teliospores. Scale bar=10 µm. **b** *Puccinia othonnae* (type), teliospores; note the delicately verrucose spore surface. Scale bar=10 µm. **c** *Puccinia othonnae* (type), teliospores; note the fine ripple mark pattern underlying the verrucose ornament. Scale bar=10 µm. **d** *Puccinia othonnoides* (type), telio-

spores. Scale bar=10 µm. **e** *Puccinia othonnoides* (type), urediniospores; note the ripple mark pattern of the spore wall. Scale bar=10 µm. **f** *Puccinia othonnoides* (type), urediniospores; note the striate pattern of the spore wall in the optical section. Scale bar=10 µm

shifted sideways or, more rarely, spores stalked laterally, spore wall 7–9 µm thick, at germ pores up to 12 µm thick, laminate, with an outer, ochraceous layer ca. 4–5 µm thick and an inner, light chestnut brown, ca. 3–4-µm-thick layer (layers are not sharply delimited but tend to blend), outer wall layer more or less evenly ornamented by flat, broad warts (ca. 1 µm in diameter) about 2.5–3.5 µm apart and by an inconspicuous ripple mark-like pattern that seems to

underly the surface, germ pores apical in distal cells and subequatorial to equatorial in the proximal cell, without papillae but with a thickening of the outer wall layer and a pit in the inner layer. In the telia, very few, mostly old, urediniospores were found with a bilaminate, echinulate wall and numerous, scattered, papillate germ pores.

Material examined: *Puccinia othonnae*. South Africa, Transvaal, Olifantsfontein, on *O. natalensis* Sch. Bip., leg.

Pienaar (no. 260), 14 April 1920 (holotype, PREM 13052). South Africa, Transvaal, Roodepoort, on *O. natalensis*, leg. I.B. Pole Evans, 4 June 1919 (paratype, PREM 14184).

Puccinia othonnoides R. Berndt, A.R. Wood & E. Uhlmann, sp. nov. (Fig. 10d–f).

Etymology: The epithet designates morphological similarity to *P. othonnae*.

Spermogonia et aecia ignota. Uredinia in foliis amphigena, subepidermalia, erumpentia, rotundata, ca. 0.5–1 mm diam., ferruginea, pulverulenta, nonnunquam telios evolventia; urediniosporae subglobosae ad late ellipsoideae, 27–45×25–36 µm, pariete 2.5–3.5 µm crasso, in hilo et poris germinationis crassiori (interdum usque ad 8 µm), bilaminato, lamina exteriori tenui, straminea vel subhyalina, sate dense moderate delicate echinulata, lamina inferiori crassiori, aurea ad dilute castanea, striis finissimis, impressioni digitalis similibus crebre praedita, poris germinationis sparsis, 9–13, infra incrassatis, extra papillis latis humilibusque. Telia urediniis similia, atro-brunnea ad nitide atra, pulvinata, textura subfibrosa, 0.5–1.2 mm diam.; teliosporae late ellipsoideae, late obovoideae vel - rariter - subglobosae, secundum septum non vel leniter constrictae, (39) 42–64×32–48 µm, pariete (5) 6–8 (10) µm crasso, usque ad 12 µm in poris germinationis, indistincte bilaminato lamina exteriori tenui, usque ad 2 µm crasso, ochracea ad aurantio-brunnea, levi vel inconspicue verrucosa verrucis humilibus, lamina interiori crassiori, aurantio-brunnea ad castanea, poris germinationis apicalibus in cellula distali, aequatorialibus vel subaequatorialibus in cellula proximali, papillatis; pedicello basaliter inserto vel obliquo, rariter laterali, usque ad 150 µm longo, hyalino vel subhyalino, crasse tunicato et persistenti. Mesosporae absunt, vel rarae ad sparsae adsunt.

In foliis *Othonnae* specierum (Asteraceae).

Uredinia amphigenous on leaves, originally subepidermal, erumpent, rounded, ca. 0.5–1 mm in diameter, ferruginous, pulverulent, sometimes giving rise to telia; urediniospores subglobose to broadly ellipsoidal, 27–45×25–36 µm, spore wall 2.5–3.5 µm thick, thicker at germ pores and generally at hilum (there occasionally up to 8 µm), two-layered with a thin, straw-coloured or subhyaline outer layer and a thicker, golden to light chestnut brown inner layer, germ pores numerous, scattered, mostly 9–13, with an internal thickening and a broad flat papilla, outer wall layer evenly moderately fine and rather densely echinulate, inner layer with an inconspicuous to conspicuous ripple mark or fingerprint-like pattern around and between the germ pores. Telia similar to uredinia, blackish brown to shiny black, pulvinate with subfibrous texture, ca. 0.5–1.2 mm in diameter; teliospores broadly ellipsoidal, broadly ovoidal or rarely subglobose, not or very slightly constricted at the septum, (39) 42–64×32–48 µm, spore

wall (5) 6–8 (10) µm thick, at germ pores to 12 µm, two-layered, with an indistinctly delimited thin, ochraceous, golden or orange brown outer layer up to 2 µm thick and a much thicker, orange brown to chestnut brown inner layer, entirely smooth to (very) inconspicuously verrucose by flat warts, sometimes only visible on young spores, germ pores apical in the distal cell and equatorial to subequatorial in the proximal cell, with a pit in the inner wall layer and a thickening of the outer one, pedicels inserted basally or slightly shifted sideways, rarely almost lateral, persistent and up to 150 µm long, hyaline to subhyaline, thick-walled. One-celled mesospores absent, rare or scattered.

On leaves of *Othonna* species (Asteraceae)

Holotype (PREM): South Africa, Western Cape Province, at the side of the road to Little Boy Kraal ca. 500 m from road between Citrusdal and Algeria, on *Othonna* cf. *coronopifolia* L., leg. E. Uhlmann and R. Berndt (no. RSA 26), 21 October 2004 (isotype Z+ZT).

Paratypes: South Africa, Western Cape Province, Cederberg Mts., at Wolfberg, on *Othonna* sp., leg. R. Berndt (no. RSA 7), 13 October 2004 (PREM, Z+ZT). South Africa, Western Cape Province, road from Citrusdal to Ceres (R 303), towards Middelberg Pass, on *Othonna* cf. *parviflora* L., leg. E. Uhlmann and R. Berndt (no. RSA 38), 22 October 2004 (PREM, Z+ZT). South Africa, Western Cape Province, on side road to Touws Rivier, ca. 1 km after turn-off from Ceres to Calvinia road, on *Othonna* sp., leg. E. Uhlmann and R. Berndt (no. RSA 46), 23 October 2004 (PREM, Z+ZT). South Africa, Western Cape Province, on road from Barrydale to Heidelberg, towards Tradouw Pass, on *O. cf. parviflora*, leg. E. Uhlmann and R. Berndt (no. RSA 56), 24 October 2004 (PREM, Z+ZT). South Africa, Western Cape Province, Cape Peninsula, Miller's Point S Simon's Town, on *Othonna arborescens* L., leg. A.R. Wood, 18 November 2000 and 28 September 2002 (no. 253 and 398).

Puccinia othonnoides appears to be a variable species. The investigated collections differ mainly with regard to teliospore size and ornament and to urediniospore size and shape. The differences are listed in Table 1. All collections share essential characters, however, and we consider it best to regard them as pertaining to a single species as the characters intergrade and sizes overlap. It is unclear at the moment whether the observed differences correlate with different *Othonna* hosts or just represent small sections from a morphological continuum.

Comparing the characters listed in Table 1, one can see that *P. othonnoides* is very similar to *P. othonnae* and *U. othonnae*: the urediniospores have many scattered germ pores and a thick, inconspicuously to distinctly bilaminated wall whose inner layer generally shows a fine striate or ripple mark pattern (Fig. 10e,f). The teliospores are thick-walled and two-layered with germ pores apical and more or less equatorial (Fig. 10d). *Puccinia othonnae* differs from

Table 1 Characters of telio- and urediniospores of rust fungi on *Othonna*

Specimen	Teliospores	Urediniospores
<i>P. othonnoides</i> , (Wood no. 398, [II] and no. 253 [III]), on <i>O. arborescens</i>	(39) 44–58 (65)×33–44 µm (mean 51.1×37.3), wall 6–7 µm thick, at germ pores 8–12 µm, bilaminar, smooth	32–36×28–36 µm (mean 34.7×31.6), wall indistinctly bilaminar, 3–3.5 µm thick, inner layer with ripple mark pattern, germ pores 9–12 (14?)
<i>P. othonnoides</i> (RSA 7), on <i>Othonna</i> sp.	42–52×34–40 µm (mean 45.9×36.9), wall about 6 µm thick, at germ pores ca. 8 µm, bilaminar, smooth (but indistinct warts visible in young spores)	32–39.5×29–33 µm (mean 36.2×30.8), wall bilaminar, 3–4 µm thick, with ripple mark pattern, germ pores (10) 11–13
<i>P. othonnoides</i> , (RSA 26, holotype), on <i>O. cf. coronopifolia</i>	45–57×31.5–38 (41) µm (mean 51.2×35.3), wall 5–7 µm thick, apically to 12 µm, indistinctly bilaminar, smooth	27–33×26–29 µm (mean 29.6×27.9), wall 2.5–3 µm thick, thicker at hilum, uniform, with ripple mark pattern, germ pores (8) 9–11
<i>P. othonnoides</i> , (RSA 38), on <i>O. cf. parviflora</i>	44–58 (60)×36–42.5 µm (mean 49.1×39.7), wall 7–10 µm thick, apically to 12 µm, appearing smooth to very inconspicuously verrucose with flat warts	31–37.5×27–33.5 µm (mean 34.3×29.8), wall 2.5–3 µm thick, bilaminar, inner layer with ripple mark pattern, germ pores 9–12
<i>P. othonnoides</i> , (RSA 46), on <i>Othonna</i> sp.	53.5–64×40–48.5 µm (mean 59.1×45.0), wall 7–8 µm thick, apically to 12 µm, indistinctly bilaminar, indistinctly verrucose with small flat warts	33.5–45×25.5–31 µm (mean 36.9×28.1), wall about 2.5 µm thick, much thicker at hilum and slightly thicker apically, indistinctly bilaminar, inner layer with ripple mark pattern, germ pores 12–14
<i>P. othonnoides</i> , (RSA 56), on <i>Othonna</i> sp.	42.5–48×33.5–38 (40) µm (mean 45.0×36.1), wall 5.5–7 µm thick, apically to 9 µm, indistinctly bilaminar, almost smooth but with inconspicuous flat warts	28–35×28–32 µm (mean 31.2×29.6), wall about 2.5 µm thick, much thicker at hilum, very indistinctly bilaminar, wall with ripple mark pattern, germ pores mostly 11–13
<i>P. othonnae</i>	53–65 (68)×38–46 µm (mean 59.8×42.2), wall 7–9 µm thick, at germ pores to 12 µm, distinctly bilaminar with thick outer layer, verrucose and with underlying “ripple mark-pattern” (Dodge: 43–50×33–40 µm, wall 6.5–8 µm thick, not thickened at pores or apically)	Very scarce: with numerous, scattered, papillate germ pores and a two-layered, echinulate wall, ripple mark pattern not observed
<i>Uredo othonnae</i> (holotype)	Not present	(31) 33–39×(29) 31–35 µm (mean 35.9×32.7), wall ca. 3–3.5 µm thick, bilaminar with ripple mark pattern of inner layer, germ pores 6–10. (Jørstad: 30–42×25–32 µm, wall 3–5 µm thick, bilaminar, germ pores ca. 8, scattered)

P. othonnoides, however, by larger teliospores with a thick, distinctly verrucose outer wall layer and an underlying ripple mark pattern (Fig. 10c). The latter may not always be visible. *Uredo othonnae* [South Africa, Western Cape Province, Koude Bokkeveld, Wagedrift, altitude of 5,500 ft, on *Othonna multicaulis* Harv. var., leg. Schlechter, 21 January 1897 (type, S F32636)] could not be assigned to either species with confidence. Its urediniospores (Fig. 11) have fewer germ pores (6–10, mostly 8) than those of *P. othonnoides*, otherwise it is very similar. Our study showed that *P. othonnae* has urediniospores, too, but they were too scanty to make a proper comparison.

Puccinia butleri Syd. & P. Syd. 1906. Annales Mycologici 4:431. New for South Africa (Fig. 12).

Rust collected on *Launaea dregeana* DC. (Eastern Cape Province, Kayser's Beach S of East London, 20 August 2002, leg. A.R. Wood) was determined as *P. butleri*, known so far from India and Pakistan. The South African specimen (Fig. 12) differs in certain aspects from the type of *P. butleri* (Z+ZT): it has teliospores with thinner walls (ca. 1.5–2 µm versus ca. 2.5 µm) with a more prominently verrucose

ornament and urediniospores that are larger [22.5–28 (30)×18.5–22.5 µm (mean 24.9×20.9 µm) versus 18–21.5×17.5–21 µm (mean 20.7×19.4 µm)]. These differences may lie within the variability of *P. butleri*, but it will be interesting to collect more material from southern Africa to evaluate whether they occur constantly.

As the original description of *P. butleri* is short, we present here additional observations obtained from the type (Z+ZT): Urediniospores obovoidal to subglobose, 18–21.5×17.5–21 µm (mean of 20 spores 20.7×19.4 µm), spore wall ochraceous to light brown, ca. 1.5 µm thick, moderately fine and closely echinulate, spines mostly 1.5–2.5 µm apart, germ pores inconspicuous, about five, scattered. Teliospores broadly ellipsoidal to elongate broadly ellipsoidal, rounded at both ends and slightly constricted at the septum, 36–42.5×24–29 µm (mean 39.2×25.6 µm), pedicels subhyaline, thin-walled, up to as long as the spores but most often breaking shorter, spore wall light chestnut brown, about 2.5 µm thick, not thickened at the apex, rather densely verrucose by flat warts, germ pores apical or subapical in the distal cell, equatorial to subequatorial in the proximal cell, with flat, rather broad, ochraceous papillae.




Fig. 11 *Uredo othonnae* (type), urediniospores; a small section of the ripple mark pattern observable within the spore wall has been delineated (not to scale). Scale bar=20 µm

Uromyces bidenticola Arthur 1917. *Mycologia* 9:71. New for South Africa, formerly reported as *Uromyces bidentis* Lagh.

Doidge (1927) reported *U. bidentis* from Natal and the Cape Province. She stated that only the urediniospores had been found so far in South Africa. *Uromyces bidentis* is a micro-cyclic species, however, producing only teliospores (Arthur 1917). Doidge's reports refer to *U. bidenticola*, therefore, and *U. bidentis* must be deleted from the South African rust list.

Material examined: South Africa, Western Cape Province, Cape Peninsula, Kirstenbosch Botanical Garden, on weed *Bidens* cf. *pilosa* L., 4 November 2004, leg. E. Uhlmann and R. Berndt.

On Dioscoreaceae

Puccinia dioscoreae-mundtii R. Berndt, A.R. Wood, & E. Uhlmann sp. nov. (Figs. 13 and 14).

Etymology: Named after the host species, *Dioscorea mundtii*, a climber in forest and thicket vegetation in South Africa.




Fig. 12 *Puccinia butleri* (Z+ZT), teliospores. Scale bar=20 µm

Uredinia in pagina abaxiali foliorum dense sparsa, subepidermalia, minuta, 0.2–0.4 mm diam. vel maiores si coalescent, primo pallide ochracea, deinde aurantio-brunnea, pulverulenta; urediniosporae obovoideae vel pyriformes, 23.5–36×18–20.5 µm (mean 28.3×19.2 µm), pariete subhyalino ad pallide ochraceo, ca. 1 µm crasso, moderate dense et aliquantum delicate echinulato, spinis 1.5–2 µm inter se distantibus, aegre echinulato hilum versus, poris germinationis obscuris, 4–5 sparsis vel cum duobus poris basalibus et duobus subapicalibus. Telia non visa; teliosporae inter urediniosporas, ellipsoideae, clavatae, rarer late ellipsoideae vel oblonge ellipsoideae, paulum constrictae secundum septum, cellula distali ellipsoidea, late ellipsoidea vel subglobosa, quam cellula proximali, cuneato, breviori est, 39–65×14.5–20 µm (mean 49.4×17.9 µm), pariete levi, ca. 0.5–1 µm crasso, apicaliter 1–2 µm, subhyalino usque ad pallide ochraceo, poris germinationis obscuris, pedicellis tenui tunicatis, plus minusve hyalino, brevibus; sporae tricellulares rarer adsunt.




Fig. 13 *Puccinia dioscoreae-mundtii* (type), teliospores. Scale bar=20 μm




Fig. 14 *Puccinia dioscoreae-mundtii* (type), urediniospores. Scale bar=20 μm

In foliis *Dioscorea mundtii* Baker (Dioscoreaceae)

Uredinia densely scattered on abaxial side of leaves that are marbled by blurred discoloured to ochraceous or pallid green areas, subepidermal, tiny, 0.2–0.4 mm in diameter, sometimes larger when confluent, pallid ochraceous when young to orange brown when older, pulverulent; urediniospores obovoidal or pyriform, 23.5–36×18–20.5 μm (mean 28.3×19.2 μm), spore wall subhyaline to light ochraceous, ca. 1 μm thick, moderately dense and rather finely echinulate (spines ca. 1.5–2 μm apart), echinulation vanishing towards the hilum; thus, more or less bald around hilum, germ pores difficult to discern, mostly by a thickening of the wall, four to five, scattered or, quite often, two pores basal and two subapical. Telia not seen, uredinia in which teliospores were found appeared slightly wax-like; teliospores ellipsoidal, clavate, rarely broadly ellipsoidal or oblong ellipsoidal, slightly constricted at septum, the distal cell ellipsoidal, broadly ellipsoidal or subglobose most often shorter than the proximal, cuneate one, 39–65×14.5–20 μm (mean 49.4×17.9 μm), wall smooth, ca. 0.5–1 μm thick, at the apex 1–2 μm , subhyaline to light ochraceous, germ pores not seen, pedicels thin-walled, more or less hyaline and breaking shortly from the hilum; three-celled spores occur rarely.

On *Dioscorea mundtii* Baker (Dioscoreaceae)

Holotype (PREM): South Africa, Western Cape Province, at road from Knysna to George shortly after Homtini pass, in the afromontane forest of a river valley, on *Dioscorea mundtii* Baker (Dioscoreaceae), 27 October 2004, leg. E. Uhlmann and R. Berndt. Isotype in Z+ZT.

Additional material studied: South Africa, Western Cape Province, Victoria Bay E of George, on *D. mundtii*, 25 July 2001, leg. A.R. Wood (no. 337, only II present).


The present species differs from known *Puccinia* spp. on *Dioscorea*, namely *Puccinia dioscoreae* Kom. and *Puccinia valida* Arthur, by uniformly thin-walled teliospores (Fig. 13). Its uredinal stage appears to be different from that of the described *Uredo* spp. on *Dioscorea*, though it is quite similar to *Uredo dioscoreae-filiformis*, which differs, however, by deep-seated uredinia, which liberate thicker-walled urediniospores through a central aperture (Ono 1982).

On Fabaceae

Uredo otholobii R. Berndt, A.R. Wood & E. Uhlmann, sp. nov. (Fig. 15a,b).

Etymology: Named after the host genus *Otholobium* of Fabaceae

Fig. 15 **a** *Uredo otholobii* (type), urediniospores; optical section. Scale bar=10 µm. **b** *Uredo otholobii* (type), urediniospores; echinulate spore surface. Scale bar=10 µm



Uredinia foliicola, amphigena, parva, pulverulenta, castanea; urediniosporae obovoideae, late ellipsoideae vel (rariter) subglobosae, 21–27×18–22 µm (medium 23.4×15.0 µm), pariete brunneo, ca. 1.5–2 µm crasso (in poris germinationis parum crassiore), delicate moderate denseque echinulato, spinis inter se 2–3 µm distantibus, poris germinationis plerumque 2–3, aequatorialibus et aequidistantibus, papillis subhyalinis, humilibus vel moderate altis et tumore debili interno praeditae.

In foliis *Otholobium* cf. *candidans* (Fabaceae, Psoraleae)

Uredinia amphigenous on leaves, small, chestnut brown, pulverulent; urediniospores obovoidal, broadly ellipsoidal or rarely subglobose, 21–27×18–22 µm (mean 23.4×15 µm), spore wall brown, ca. 1.5–2 µm thick, slightly thicker at the germ pores, evenly fine and moderately dense echinulate, spines mostly 2–3 µm apart, two to three germ pores, in most cases approximately equatorial and equidistant with a flat to moderately high subhyaline papilla and a slight internal swelling.

On leaves of *Otholobium* cf. *candidans* (Fabaceae, Psoraleae)

Holotype (PREM): South Africa, Western Cape Province, Barrydale, at the southern border of the village, on *Otholobium* cf. *candidans* (Eckl. & Zeyh.) C.H. Stir., 23 October 2004, leg. E. Uhlmann and R. Berndt (isotype Z+ZT).

Additional material investigated: South Africa, Western Cape Province, Stellenbosch, Jan Marais Nature Reserve, on *Otholobium hirtum* (L.) C.H. Stirton, 16 September 2002, leg. A.R. Wood (no. 394). South Africa, Western Cape Province, E Bredasdorp, De Hoop Nature Reserve, on

Otholobium fruticans (L.) C.H. Stirton, 24 June 1998, leg. A.R. Wood (no. 53).

Other rust fungi on *Otholobium* are *Phakopsora meibomiae* (Arthur) Arthur with paraphysate uredinia and *Uromyces psoraleae* Peck, a demicyclic rust. *Uredo otholobii* is also different from *U. psoraleae-polystictae* Doidge and *Uromyces abbreviatus* Arthur, which occur on other members of the tribe Psoraleae.

Rust on *Lotononis* (Fabaceae, Crotalarieae)

Rust collected on *Lotononis falcata* (E. Mey.) Benth. revealed the telial stage of *Uromyces* sp. together with the uredinial stage. According to the authors' knowledge, the only known rust on *Lotononis* spp. is *Uredo lotononi* Doidge. A comparison of both rust fungi did not reveal significant differences. Teliospores were found in the uredinia of the type specimen of *U. lotononi*, and therefore, a new species, *Uromyces lotononidicola*, is proposed for the holomorph:

Uromyces lotononidicola R. Berndt sp. nov. (Figs. 16 and 17)

Syn. *U. lotononi* Doidge 1927. Bothalia 2:213.

Uredinia in foliis amphigena, sparsa, minuta (0.3–0.5 mm diam.), subepidermalia, ferruginea ad dilute badia, pulverulenta; urediniosporae obovoideae, late ellipsoideae vel, rariter, subglobosae, 25–33×20–24.5 µm (medium 28.1×22.5 µm), pariete dilute brunneo, ca. 2 µm crasso, delicatissime echinulato, 1.5–2.5 µm inter spinas, poris




Fig. 16 *Uromyces lotononidicola* (type), teliospores. Scale bar=15 µm

germinationis 3–5 (praecipue 3–4), aequatorialibus vel sparsis cum papillis latis, hyalinis et echinulatis. Telia non visa; teliosporae urediniosporis immixtae, deciduae, late ellipsoideae, obovoideae (ad subglobosae), (23) 25–32×20–23 µm (medium 27.7×21.7 µm), pariete dilute brunneo, ca. 2–3 µm crasso, grosse denseque verrucoso, cum verrucis ca. 1.5–2.5 µm latis, conicis vel subacutis, poro germinationis apicali, papilla lata, ca. 1.5–2.5 µm alta, subhyalina, rugosa vel rugulosa praedita; pedicello hyalino fragili.

In foliis *Lotononis cytisoides* Benth. (Fabaceae)

Uredinia amphigenous on leaves, scattered, tiny (0.3–0.5 mm in diameter), subepidermal, ferruginous to light chestnut brown, pulverulent; urediniospores obovoidal, broadly ellipsoidal or, more rarely, subglobose, 25–33×20–24.5 µm (mean 28.1×22.5 µm), wall light brown, ca. 2 µm thick, very finely echinulate, ca. 1.5–2.5 µm between spines, germ pores 3–5 (predominantly 3–4), almost equatorial to scattered, with a moderate internal thickening of the wall and a broad, hyaline, echinulate papilla. Telia not seen; teliospores among the urediniospores, broadly ellipsoidal, obovoidal (to subglobose), (23) 25–32×20–23 µm (mean 27.7×21.7 µm), spore wall light brown, ca. 2–3 µm thick, coarsely verrucose, warts ca. 1.5–




Fig. 17 *Uromyces lotononidicola* (type), urediniospores. Scale bar=15 µm

2.5 µm broad, conical to subacute, densely situated, germ pores apical with a broad, subhyaline rugose to rugulose papilla ca. 1.5–2.5 µm high, spores deciduous with fragile, hyaline and delicate pedicels that normally break off shortly below the hilum.

On leaves of *Lotononis cytisoides* Benth. (Fabaceae)

Holotype: South Africa, Natal, Mont-aux-Sources, 20 April 1919, leg. Mogg (holotype of *U. lotononi*, PREM 12955).

Additional material studied: Paratype (PREM): South Africa, Northern Cape Province, at road no. 7 north of Gariep, on *Lotononis falcata* Benth., 14 October 2004, leg. E. Uhlmann and R. Berndt (isoparatypus in Z+ZT).

Uredo lotononi: South Africa, Pretoria Distr., Irene, on *Lotononis hirsuta* Schinz, 14 March 1917, leg. Pole Evans (paratype of *U. lotononi*, PREM 10986).

The urediniospores of the specimen from the Northern Cape Province were very similar to those of the type but slightly smaller: Urediniospores broadly ellipsoidal, obovoidal or subglobose, 22–30 (31.5)×18–22 µm (mean 24.7×20.3 µm), spore wall ochraceous to light brown, about 1.5 (2) µm thick, evenly rather sparsely to moderately dense and finely echinulate, three to four germ pores, mostly three, more or less equatorial and equidistant, sometimes almost scattered, with broad, flat, subhyaline papillae. Telia amphigenous on leaves, more rarely on pedicels, subepidermal, first singly then confluent, rounded or elongated, about 0.4–2 mm in diameter, up to 3 mm long, pulvinate, dark chestnut brown, pulverulent; teliospores broadly ellipsoidal, subglobose or subpyriform,

sometimes slightly irregularly deformed, with short remnants of the fragile, thin-walled, hyaline pedicels, $25\text{--}32 \times 20.5\text{--}25 \mu\text{m}$ (mean $27.6 \times 22.8 \mu\text{m}$), spore wall $2.5\text{--}3 \mu\text{m}$ thick, orange brown to light chestnut brown, densely covered by rather coarse, broadly conical or hemispherical warts, germ pore apical, without or with inconspicuous to conspicuous broad, ochraceous and rugose papilla.

In a slide prepared from PREM 10986 two coarsely verrucose teliospores with apical papillae were present. The urediniospores measured $21.5\text{--}29 \times 18\text{--}22.5 \mu\text{m}$ (mean $24.2 \times 19.8 \mu\text{m}$) and the specimen is assigned to *U. lotononidicola* as well. Other *Uromyces* spp. on members of Crotalarieae, namely *Uromyces africanus* (Gjærum) Ono, *Uromyces bolusii* Massee, *Uromyces crotalariae* (Arth.) Baxter, *Uromyces decoratus* Syd. & P. Syd., *Uromyces harmsianus* (Henn.) Doidge, and *Uromyces occidentalis* Dietel, are clearly different from the present species.




Fig. 18 *Puccinia aurea* (type of *P. satyrii*), teliospores. Scale bar= $20 \mu\text{m}$

On Orchidaceae

Puccinia aurea Winter 1884. Flora 67:260

Syn.: *Puccinia satyrii* P. Syd. & Syd. 1903. Monogr. Uredin. Vol. I:594 (Fig. 18).

Puccinia rust collected on a sterile orchid in Fynbos vegetation at Bain's Kloof above Paarl resembled *P. aurea* and *P. satyrii*. A comparison was made, therefore, with original material.

Material examined: *Puccinia* aff. *aurea*: South Africa, Western Cape Province, Paarl, Witte Rivier at Bain's Kloof, on sterile orchid (cf. *Ceratandra*), 11 December 2004, leg. E. Uhlmann and R. Berndt. *Puccinia satyrii*: South Africa, Western Cape Province, Cape flats near False Bay, on *Satyrium carneum* R. Br., September 1884, leg. P. Mac Owan (holotype, Berlin) and Rabh.-Winter Fungi Europ. no. 3614 (isotype, Berlin, sub *P. aurea*).

Puccinia satyrii is very similar to *P. aurea* and, according to Sydow and Sydow (1904), differs essentially by slightly smaller teliospores and the presence of urediniospores. Jørstad (1956a) considered *P. satyrii* to be conspecific with *P. aurea* and listed six different host genera. We studied the type of *P. satyrii* and another specimen issued under *P. aurea* in Rabh.-Winter Fungi Europ. no. 3614. Our measurements (Table 2) show that the teliospores of *P. satyrii* are not smaller than those of *P. aurea*, and we follow the opinion of Jørstad (1956a) that both species are the same.

The specimen recently collected at Bain's Kloof is distinct from *P. aurea* by smaller teliospores (Fig. 19), which are less thickened at the apex and have a lighter-coloured spore wall. Despite these differences, we prefer to assign this specimen to *P. aurea* at the moment as it shows the same set of characters in principal (leptosporic teliospores with a much thickened apex and tiny, punctiform, subcompact telia). *Puccinia aurea* may therefore be regarded as a variable species able to infect a rather broad spectrum of orchids.

On Solanaceae

Puccinia on *Lycium* species in Africa and the Near and Middle East

Five species of *Puccinia* have been described on *Lycium* spp. in Africa, the Near and Middle East and southern Europe: *Puccinia afra* Winter (South Africa, Spain), *Puccinia lycii* Kalchbr. [South Africa, Yemen (Island of Abd-al-Kuri)], *Puccinia spinulosa* Jørstad (Madagascar), *Puccinia turgida* P. Syd. & Syd. (Egypt, Israel, Pakistan) and *Puccinia verwoerdiana* Van der Byl (South Africa). Specimens of *Puccinia* collected recently on *Lycium* in Namibia and in South Africa could not be determined by

Table 2 Spore measurements of *Puccinia aurea* (including *P. satyrii*)

Taxon	Teliospores	Average	Urediniospores	Average
<i>P. aurea</i> [after Sydow and Sydow (1904)]	35–65×13–20 µm, apex thickened to 16 µm	–	Not described	–
<i>P. satyrii</i> , holotype	36–56×15–20 µm, apex thickened to 14 µm [after Sydow and Sydow (1904); 32–52×13–19 µm, apex thickened to 13–19 µm]	45.7×17.6 µm	Not seen in available fragment of type [after Sydow and Sydow (1904); globose or subglobose, 16–24 µm in diameter]	–
<i>P. satyrii</i> (sub <i>P. aurea</i>), isotype issued in Rabh. Fg. Europ. no. 3614	(37.5) 44–61 (64)×16–20 µm, apex thickened to 8–17 µm [after Sydow and Sydow (1904); 35–65×13–20 µm]	50.9×17.9 µm	Only a few collapsed spores present in specimen from B [after Jørstad (1956a); 19–26×15–19 µm, germ pores obscure]	–
<i>Puccinia</i> aff. <i>aurea</i> , (Bain's Kloof)	30–47×13.5–18 µm, apex thickened to 6–11 µm	38.6×15.8 µm	18–23.5×14.5–18 µm, germ pores not seen	20.8×16.4 µm

comparison with available descriptions. To evaluate their status, they were compared to original material of the listed *Puccinia* species.

The specimen from South Africa was different from the known species and is described as new:

Puccinia rapipes R. Berndt & E. Uhlmann, sp. nov.
(Figs. 20a,b, and 21a)

Etymology: Named after the shape of the swollen teliospore pedicel.




Fig. 19 *Puccinia* aff. *aurea* (specimen from Bain's Kloof), teliospores. Scale bar=20 µm

Spermogonia et aecia ignota. Uredinia in foliis amphigena, sparsa, parva, subepidermalia, ferruginea, pulverulentia, 0.3–1 mm diam.; urediniosporae fusiformes ad ellipsoideae, apicaliter subacute, basim versus attenuatae vel truncatae, 38–60×16.5–22 µm (medium 50.8×18.5 µm), pariete ochraceo ad dilute brunneo, ca. 1.5 µm crasso, apicaliter leniter incrassato usque ad 2–3 µm, delicate et moderate dense echinulato spinis brevibus, inter se ca. 2.5–3.5 µm distantibus, hilum versus decrescentibus, deinde levi, poris germinationis 3–4, praecipue 4, plusminusve subaequatorialibus et aequidistantibus, papillis parvis, humilibus, subhyalinis praeditis. Telia urediniis similia, atrobrunnea vel atra; teliosporae ellipsoideae, late ellipsoideae vel subfusiformes, non vel paululum constrictae ad septum, apicaliter saepe apiculo usque ad 6 µm alto praeditae, 41.5–61×21.5–27 µm (medium 48.8×24.2 µm), pariete castaneo, 1.5–2.5 µm crasso, crebre verruculoso verrucis parvis, humilibus, non-nihil longitudinaliter dispositis, areas fere leves includentibus, poro germinationis cellulae distalis subapicali ad aequatoriali, plusminusve aequatoriali in cellula proximali, cum vel sine papillis verruculosis et inconspicuis, pedicello basaliter inserto, levi, subhyalino, dilute brunneo hilum versus, crasse tunicato, ca. 25–30 µm ab hilo vesiculoso inflato vel obovoideo-caudato.

In foliis *Lycii* cf. *ferocissimi* Miers (Solanaceae)

Spermogonia and aecia not present. Uredinia amphigenous on leaves, scattered, subepidermal, small, more or less rounded, 0.3–1 mm in diameter, ferruginous, pulverulent, often developing to telia later; urediniospores fusiform to ellipsoidal, apically subacute, tapering towards the hilum or truncate basally, 38–60×16.5–22 µm (mean 50.8×18.5 µm), spore wall ochraceous to light brown, ca. 1.5 µm thick, apically slightly thickened to 2–3 µm, rather finely and moderately densely echinulate with short spines, spines spaced about 2.5–3.5 µm, or slightly closer apically, becoming smaller towards the hilum and fading, smooth

around the hilum, three to four germ pores, predominantly four, more or less subequatorial and equidistant, with small and flat, subhyaline papillae. Telia like uredinia, blackish-brown to black; teliospores ellipsoidal, broadly ellipsoidal or subfusiform, not or hardly constricted at the septum, apically often with an inconspicuous to conspicuous yellow-brown apiculus up to 6 μm high, 41.5–61 \times 21.5–27 μm (mean 48.8 \times 24.2 μm), spore wall chestnut brown, 1.5–2.5 μm thick, densely verrucose with small, flat,

irregular warts that tend to be arranged in broad longitudinal stripes including almost smooth areas, germ pores subapical to equatorial in the distal cell and more or less equatorial in the proximal cell, without or with inconspicuous verrucose papillae, pedicel broadly (8–10 μm) attached at base of spore, smooth, light brown below the hilum, then subhyaline, slightly thick-walled, swelling conspicuously ca. 25–30 μm from the hilum and then ovoid-caudate to almost globose.




Fig. 20 **a** *Puccinia rapipes* (type), teliospores; in one spore, the fine warty spore surface has been delineated. Scale bar=20 μm . **b** *Puccinia rapipes* (type), urediniospores. Scale bar=20 μm . **c** *Puccinia lycii* (type), teliospores; in one spore, a section of the surface has been delineated with the verrucose ornament. Scale bar=20 μm . **d** *Puccinia lycii* (type), urediniospores. Scale bar=20 μm




Fig. 21 **a** *Puccinia rapipes* (type), teliospores. Scale bar=10 µm. **b** *Puccinia lycii* (type of *P. verwoerdiana*, PREM 46255), teliospores. Scale bar=10 µm. **c** *Puccinia spinulosa* (Namibian specimen), teliospores. Scale bar=10 µm

On leaves of *Lycium* cf. *ferocissimum* Miers (Solanaceae)

Holotype (PREM): South Africa, Western Cape Province, Simon's Town, on the shore above the penguin sanctuary at Boulders, on *Lycium* cf. *ferocissimum*, leg. E. Uhlmann and R. Berndt, 1 November 2004 (isotype Z+ZT).

The present species is morphologically intermediate between *P. lycii* and *P. afra*. It differs from *P. afra* by more slender and thinner-walled teliospores that are finely

verrucose and by smaller, thinner-walled urediniospores that are more finely echinulate and have inconspicuously papillate germ pores. It is very similar to *P. lycii* but has more slender and longer, thinner-walled, often apiculate teliospores and smaller urediniospores, which are more finely echinulate and hardly thickened apically.

Puccinia lycii Kalchbr. 1882. Grevillea 11:21 (Figs. 20c,d and 21b).

Syn.: *P. verwoerdiana* Van der Byl 1927. S. Afr. J. Science 24:226.

The diagnosis of *P. verwoerdiana* is very short and does not allow positioning the species among the other Puccinias on *Solanum*. A study of the type (PREM 46255, Fig. 21b) did not reveal significant differences to *P. lycii* (type, Z+ZT, Fig. 20c,d). *Puccinia verwoerdiana* is therefore reduced to a synonym of the latter.

Puccinia spinulosa Jørstad 1957. Arkiv för Botanik 3 (17):592. New report for Namibia (Fig. 21c).

Syn.: *P. turgida* auct., non P. Syd. & Syd.: Mennicken, Maier and Oberw. 2005. Mycol. Progr. 4:69.

Rust collected in Israel on *Lycium europaeum* L. by Bornmüller was first assigned to *P. lycii* by Magnus (1898), but was later regarded as a separate species, *P. turgida* (Sydow and Sydow 1904). It has been known hitherto only from the Near and Middle East but was reported recently from Namibia by Mennicken et al. (2005). We studied the Namibian material and found that the specimens are different from *P. turgida* but not distinguishable from *P. spinulosa* described by Jørstad (1956a) from Madagascar. This is the first report of the latter rust from outside Madagascar and a new report for Namibia.

A specimen of *P. lycii* reported from the Island of Abd-al-Kuri (Yemen) by Gjærum (1987) may belong to *P. turgida* as the location is close to the reported area of this rust.

The Puccinias on *Lycium* considered here share a set of characteristic morphological traits: the proximal or central parts of their teliospore pedicels swell conspicuously in aqueous fluids, their teliospores are verrucose and have about equatorial germ pores and the urediniospores are slender with more or less equatorial pores and an even echinulation that becomes very fine or fades towards the hilum. This similarity clearly indicates that these rusts are closely related and belong to a natural group. It may also suggest that speciation within this group took place relatively recently. Fukuda et al. (2001) studied the phylogeny and biogeographical aspects of *Lycium* with a cpDNA sequence analysis. Their results indicate that *Lycium* originated in the New World from where it dispersed to South Africa, and that the Eurasian *Lycium* spp. probably derive from South African ancestors. With

the exception of *P. turgida*, all Old World species of *Puccinia* on *Lycium* occur in southern Africa, and one can assume that this group came into being there. It is interesting to note that the New World species, *Puccinia globosipes* Peck and *Puccinia paradoxopoda* Speg. (=*Puccinia tumidipes* Peck) on members of *Lycium* are characterised by swelling pedicels as well. In these species, however, the entire pedicel swells to an ellipsoid body, and they differ by several other characters from the Old World *Lycium* rusts.

Key to the *Puccinia* species on Old World *Lycium*:

- 1 Teliospores predominantly or often apiculate or subapiculate
- 2 Teliospore wall 3–4 µm thick, grossly verrucose—*P. afra*
- 2* Teliospore wall 1.5–2.5 µm thick, finely verrucose—*P. rapipes*
- 1* Teliospores rounded at apex, only occasionally subapiculate or apiculate
- 3 Teliospores densely and finely verrucose or rather densely verrucose with dome-shaped or knob-like warts
- 4 Teliospores verrucose by irregular, dome-shaped, knob-like (to shortly cylindrical) warts, walls ca. 4–5 µm thick, urediniospores ellipsoid and rather short (mean ca. 44×21 µm)—*P. turgida*
- 4* Teliospores finely verrucose, walls ≤3 µm thick, urediniospores ± fusiform and considerably longer on average—*P. lycii*
- 3* Teliospores sparingly verrucose or spinulose by slenderly cylindrical, conical or obconical warts—*P. spinulosa*

Discussion

At present, we know roughly 480 taxa of plant rust occurring in South Africa (Crous et al. 2000 onwards; personal data). This number is low if related to the very rich and diverse phanerogam flora of the country comprising more than 21,000 species (Cowling and Hilton-Taylor 1997). In our work, we contribute eight new rust species from the Western and Northern Cape Provinces, and it is certain that new rust species will continue to be discovered (e.g. Wood and Scholler 2005). In recent years, attempts were made to estimate the species richness of plant parasitic fungi extrapolating parasite–host ratios calculated for restricted areas of the world (compare Rossman 1994). Referring to such ratios, it has been assumed that up to 5,000 rust species may occur in South Africa (Mennicken and Oberwinkler 2004). We think that it is problematic to apply parasite–host ratios assessed for a specific country or area to other regions of the world. The flora of South Africa's cape floristic region (CFR), for example, is very

species-rich with almost 9,000 species of seed plants (Linder 2003), a high rate of endemism and an astonishing predominance of a few very large taxa. It is interesting to note that the latter are generally hosts to very few rust fungi or are even not known to support any rusts. The following list was prepared after Linder (2003) and comprises 12 of the most species-rich clades of seed plants of the CFR with their species numbers (first number in the parentheses). For each clade, the number of known rust fungi in the CFR is given (second number in the parentheses):

Ericaceae, *Erica* (658/none); Iridoideae, Iridoideae and Nivenioideae (ca. 516/ca. 20); Restionaceae (340/none); Fabaceae, *Aspalathus* and *Rafnia* (291/3); Rutaceae, Diosmeae (268/1); Proteaceae (264/none); Geraniaceae, *Pelargonium* (148/2); Rhamnaceae, Phytoliceae (134/1); Asteraceae, Relhaniinae (131/2); Fabaceae, Podalyriaceae and Liparieae (120/none); Polygalaceae, *Muraltia* and *Nylandia* (108/none); Scrophulariaceae, *Selago* (101/none).

Together, the listed plant taxa comprise ca. 3,079 species and represent ca. 34% of the flora of the CFR. They are hosts to ca. 29 known rust species. Omitting Iridoideae from the list, there remain 2,563 plant species (ca. 28% of flora of CFR) with only nine known rust fungi. Though the numbers of rust species on the listed clades may be underestimated, they are, nonetheless, exceedingly low. Hennen and McCain (1993) calculated a rust fungus–host ratio of between 1:4 and 1:20. Even only applying the low-limit ratio of one rust species per 20 plant species assumed by the latter authors, one should expect 154 rusts instead of 29 for the listed plant taxa in the CFR (or 128 instead of 9 when omitting Iridoideae from the list).

In a relatively well-studied area like the Cape region, this discrepancy between real and calculated parasite–host ratios can hardly be explained by insufficient collecting alone. It is very likely that other factors, such as the composition of a given flora as well as the “relative susceptibility” of its components to rust fungi or even abiotic conditions, are responsible for or contribute to the observed paucity. It will not suffice, therefore, just to extrapolate assumed or known parasite–host ratios to an area of interest, but it will be necessary to consider the mentioned factors to obtain more reliable estimates of the true species-richness.

The CFR offers a unique possibility to study the rust mycobiota of an extra-tropical area with a very species-rich and well distinguished flora and to test assumptions on parasite–host ratios. Floristic and taxonomic data as contributed by the present study are an integral part of such studies.

Acknowledgements This work was carried out as a part of the “BIOTA-Southern Africa” research project (subproject S03b) within the framework of BIOLOG-Germany. We thank all nature conservation authorities for permissions and support: in South Africa, the

Northern Cape Department of Nature and Environmental Conservation and the Western Cape Nature Conservation Board; in Namibia, the Ministry of Environment and Tourism, as well as the Ministry of Agriculture, Water and Rural Development. Many thanks go to all landowners for the permission to work on their land, to the German Ministry of Education and Research for financial support and to the herbaria PREM, PUR, TUB and S for the loan of specimens. We also thank A. Ritschel (Tübingen, Germany) for corrections and suggestions on the manuscript and A.R. Wood (Stellenbosch, South Africa) for rust specimens and discussions.

References

- Arthur JC (1917) Uredinales of Porto Rico based on collections by H. H. Whetzel and E. W. Olive. *Mycologia* 9:55–104
- Berndt R (2004) A checklist of Costa Rican rust fungi. In: Agerer R, Piepenbring M, Blanz P (eds) *Frontiers of Basidiomycote Mycology*. IHW, Eching, Germany, pp 185–236
- Berndt R, Görke C, Mennicken M, Uhlmann E, Oberwinkler F (2002) The BIOTA southern Africa project: concepts and first results. Abstract. 7th international mycological congress, Oslo, August 2002
- Cowling RM, Hilton-Taylor C (1997) Phyogeography, flora and endemism. In: Cowling RM, Richardson DM, Pierce SM (eds) *Vegetation of southern Africa*. Cambridge University Press, Cambridge, UK, pp 43–61
- Crous PW, Phillips AJL, Baxter AP (2000) Phytopathogenic fungi from South Africa. <http://nt.ars-grin.gov/fungaldatabases/southafrica/Index.cfm> (online version, designed and hosted by Systematic Botany and Mycology Laboratory, United States Department of Agriculture)
- Doidge EM (1927) A preliminary study of the South African rust fungi. *Bothalia* 2:1–228
- Doidge EM (1950) The South African fungi and lichens to the end of 1945. *Bothalia* 5:1–1094
- Fukuda T, Yokoyama J, Ohashi H (2001) Phylogeny and biogeography of the genus *Lycium* (Solanaceae): inferences from chloroplast DNA sequences. *Mol Phylogen Evol* 19:246–258
- Gjærum HB (1987) Rust fungi from Socotra and Abd-al-Kuri. Notes R Bot Gard Edinb 44:411–413
- Hennen JF, McCain JW (1993) New species and records of Uredinales from the Neotropics. *Mycologia* 85:970–986
- Holm L (1966) Études Urédinologiques. 4. Sur les *Puccinia* caricoles et leurs alliés. *Sven Bot Tidskr* 60:23–32
- Iwanoff B (1907) Untersuchungen über den Einfluss des Standortes auf den Entwicklungsgang und den Peridienbau der Uredineen. Centbl Bakteriol Parasitenkd Infektkrankh 2 Abt 18:1–5
- Jørstad I (1956a) Reliquiae Lagerheimiana. African Uredinales. *Ark Bot* 3(17):563–598
- Jørstad I (1956b) On the *Sonchus* rust *Peristemma pseudosphaeria* (Mont.) n. comb. (syn. *Puccinia sonchi* Rob.) *Friesia* 5:278–283
- Kleinjan CA, Morin L, Edwards PB, Wood AR (2004) Distribution, host range and phenology of the rust fungus *Puccinia myrsiphylli* in South Africa. *Australas Plant Pathol* 33: 263–271
- Laundon GF (1963) Rust fungi I: on Acanthaceae. *Mycol Pap* 89:1–89
- Linder HP (2003) The radiation of the Cape flora, southern Africa. *Biol Rev* 78:597–638
- Magnus P (1898) Ein kleiner Beitrag zur Kenntnis der *Puccinia lycii* Kalchbr. *Hedwigia* 1898:91–93 (Beibl.)
- Mayus O (1904) Die Peridienzellen der Uredineen in ihrer Abhängigkeit von Standortsverhältnissen. Dissertation, University of Bern, Bern, Switzerland
- Mennicken M, Oberwinkler F (2004) A contribution to the rust flora (Uredinales) of southern Africa, with an emphasis on South Africa. *Mycotaxon* 90:1–28
- Mennicken M, Berndt R, Oberwinkler F (2003) A new rust fungus (Uredinales) on Penaeaceae: *Uredo sarcocollae* on *Saltera sarcocolla*. *Mycotaxon* 85:147–151
- Mennicken M, Maier W, Oberwinkler F (2005) A contribution to the rust flora (Uredinales) of southern Africa, with an emphasis on Namibia. *Mycol Progress* 4:55–75
- Morris MJ (1982) A systemic rust fungus infecting *Chrysanthemoides monilifera* subsp. *monilifera* in South Africa. *Phytophylactica* 14:31–34
- Ono Y (1982) Rusts of yams in Southeast Asia and the South Pacific. *Trans Br Mycol Soc* 79:423–429
- Ragunathan AN, Ramakrishnan K (1972) Rust fungi of Madras State. IV. *Puccinia*. *Mysore J Agric Sci* 6:450–460
- Rossman AY (1994) A strategy for an all-taxa inventory of fungal biodiversity. In: Peng C-I, Chou CH (eds) *Biodiversity and terrestrial ecosystems*. Institute of Botany, Academia Sinica Monograph series no. 14, Taipei, Taiwan, pp 169–194
- Sato T, Sato S (1982) Aeciospore surface structure of the Uredinales. *Trans Mycol Soc Jpn* 23:51–63
- Stafleu FA, Holmgren PK, Keuken W, Schofield EK (1981) *Index herbariorum*, 7th edn. Bohn, Scheltema and Holkema, Utrecht/W. Jonk B.V., The Hague, Netherlands
- Sydow P, Sydow H (1904) *Monographia Uredinearum*, vol 1. *Puccinia*. Gebr. Bornträger, Leipzig, Germany
- Wilson M, Henderson DM (1966) *British rust fungi*. Cambridge University Press, Cambridge, UK
- Wood AR (2002) Infection of *Chrysanthemoides monilifera* ssp. *monilifera* by the rust fungus *Endophyllum osteospermi* is associated with a reduction in vegetative growth and reproduction. *Australas Plant Pathol* 31:409–415
- Wood AR, Scholler M (2005) *Uromyces euryopsidicola* sp. nov., a rust species that forms witches' brooms on *Euryops* (Asteraceae) in South Africa. *Sydowia* 57:137–143
- Wood AR, Crous PW, Lennox CL (2004) Predicting the distribution of *Endophyllum osteospermi* (Uredinales, Pucciniaceae) in Australia based on its climatic requirements and distribution in South Africa. *Australas Plant Pathol* 33:549–558