



## Kobe University Repository : Kernel

|                         |   |
|-------------------------|---|
| タイトル<br>Title           | Chromosome number determinations in the Australian Astereae (Asteraceae)  |
| 著者<br>Author(s)         | Watanabe, Kuniaki / Short, P.S. / Denda, Tetsuo / Suzuki, Y / Ito, Motomi / Yahara, Tetsukazu / Kosuge, Keiko       |
| 掲載誌・巻号・ページ<br>Citation  | Muelleria,9:197-228   |
| 刊行日<br>Issue date       | 1996  |
| 資源タイプ<br>Resource Type  | Journal Article / 学術雑誌論文  |
| 版区分<br>Resource Version | publisher   |
| 権利<br>Rights            |   |
| DOI                     |   |
| URL                     | <a href="http://www.lib.kobe-u.ac.jp/handle_kernel/90001252">http://www.lib.kobe-u.ac.jp/handle_kernel/90001252</a> |

Create Date: 2017-12-18



## Chromosome number determinations in the Australian Astereae (Asteraceae)

K. Watanabe<sup>1</sup>, P.S. Short<sup>2</sup>, T. Denda<sup>1</sup>, Y. Suzuki<sup>3</sup>, M. Ito<sup>3</sup>, T. Yahara<sup>4</sup> & K. Kosuge<sup>1</sup>

<sup>1</sup>Department of Biology, Faculty of Science, Kobe University, Kobe 657, Japan.

<sup>2</sup>National Herbarium of Victoria, Birdwood Avenue, South Yarra, 3141, Victoria, Australia.

<sup>3</sup>Department of Biology, Faculty of Science, Chiba University, Chiba, Japan.

<sup>4</sup>Department of Biology, Faculty of Science, Kyushu University, Hukuoka 812, Japan.

### ABSTRACT

Chromosome number determinations from 200 populations attributed to 99 species or infraspecific taxa of ten genera are presented in Table I. They include the first substantiated reports for *Brachyscome glandulosa* ( $2n = 36$ ), *B. muellerooides* ( $n = 3$ ), *B. obovata* ( $2n = 18$ ), *B. petrophila* ( $2n = 18$ ), *B. tatei* ( $n = 9$ ), *Erodiocephalum elderi* ( $2n = 16$ ), *Minuria gardneri* ( $n = 9$ ), *M. integrerrima* ( $n = 18$ ), *Olearia astrotricha* ( $n = 9$ ), *O. ferresii* ( $n = 9$ ), *O. imbricata* ( $n = 9$ ), *O. plucheacea* ( $n = 9$ ), *O. ramulosa* ( $n = 9$ ), *O. xerophila* ( $n = 9$ ), *Vittadinia cuneata* ( $n = 9$ ), *V. gracilis* ( $n = 9$ ), *V. muelleri* ( $n = 9$ ), *V. pterochaeta* ( $n = 9$ ), *V. pustulata* ( $n = 9$ ) and *V. sulcata* ( $n = 9$ ). Polyploidy is reported for the first time in *Brachyscome dissectifolia*, *B. cardiocarpa*, *Calotis anthemoides* and in the *Olearia phlogopappa* complex. Remaining counts support previously reported determinations and add to the knowledge of chromosome number distribution within taxa. Some taxonomic problems are briefly discussed. It is concluded that chromosomal variation in Australian Astereae is similar to that found in North America.

### Introduction

In a recent census of Australian vascular plants 831 native species of Asteraceae were listed under c. 130 genera (Hnatiuk 1990). About 330 formally recognised species belong to the tribe Astereae, and most are currently distributed amongst four genera, i.e. *Olearia* Moench (c. 130), *Brachyscome* Cass. (c. 70), *Vittadinia* A.Rich. (29) and *Calotis* R.Br. (c. 28). The remaining genera here considered to belong to the Astereae are *Achnophora* F.Muell. (1), *Camptacra* N.T.Burb (2), *Celmisia* Cass. (c. 10), *Centipeda* Lour. (5), *Ceratogyne* Turcz. (1), *Dichrocephala* DC. (1), *Dichromochlamys* Dunlop (1), *Dimorphocoma* F.Muell. & Tate (1), *Elachanthus* F.Muell. (2), *Erigeron* L. (c. 10), *Erodiocephalum* F.Muell. (2), *Eurybiopsis* DC. (1), *Isoetopsis* Turcz. (1), *Ixiochlamys* F.Muell. & Sond. (4), *Kippistia* F.Muell. (1), *Minuria* DC. (10 or 11), *Lagenifera* Cass. (4) and *Solenogyne* Cass. (3).

Chromosome number surveys have been primarily restricted to just two genera, i.e. *Brachyscome* (Smith-White *et al.* 1970, Carter 1978, Watanabe & Short 1992) and *Calotis* (Stace 1978, 1982). In this paper we report further chromosome number determinations for both of these genera as well as new determinations for native species of *Erodiocephalum*, *Lagenifera*, *Minuria*, *Olearia*, *Solenogyne* and *Vittadinia*, briefly examine previous reports, and comment on some associated taxonomic problems.

A number of species of Astereae belonging to *Aster* L., *Bellis* L., *Conyza* Less., *Erigeron* L. and *Solidago* L. are naturalised in Australia but with the exception of *Erigeron karvinskianus* they have not been examined.

### Materials and methods

Chromosome counts were obtained from either floral bud material fixed in the field, or from root tips obtained from seedlings grown from fruit of known provenance. For the

TABLE 1. CONTINUED

| Species & locality   | n     | 2n |
|--|-------|----|
| 1.2 km NW of MacKenzie Falls, Vict.,<br>27 Sept. 1993, <i>Watanabe 97</i>  | 9II   |    |
| <b>B. nivalis</b> F.Muell.<br>Club Lake, Mt Kosciusko N.P.,<br>N.S.W., 5 Feb. 1993, <i>Short 4005</i>                | 11III | 22 |
| <b>B. nodosa</b> P.S.Short & K.Watan.<br>Bracket Creek, Qld, 29 Sept. 1992,<br><i>Watanabe 19</i>                    | 3II   |    |
| 15.5 km E of Inglewood, Qld,<br>29 Sept. 1992, <i>Watanabe 20</i>  | 3II   |    |
| 2 km E of Inglewood, Qld,<br>29 Sept. 1992, <i>Watanabe 24</i>   | 3II   |    |
| 69 km N of Coonabarabran, N.S.W.,<br>25 Sept. 1992, <i>Watanabe 8</i>  | 3II   |    |
| <b>B. nova-anglica</b> G.L.R.Davis<br>Girraween N.P., Qld,<br>29 Jan. 1993, <i>Watanabe 79</i>                       | 7II   | 14 |
| 26.5 km SE of Wongwibinda, N.S.W.,<br>31 Jan. 1993, <i>Short 3978</i>  | 7II   | 14 |
| <b>B. aff. nova-anglica (A)</b><br>3 km S of Glen Aplin, Qld,<br>30 Jan. 1993, <i>Short 3969</i>                     | 5II   | 10 |
| <b>B. aff. nova-anglica (B)</b><br>Mt Kaputar N.P., N.S.W.,<br>25 Sept. 1992, <i>Watanabe 9</i>                      | 6II   | 12 |
| <b>B. obovata</b> G.L.R.Davis<br>Daners Gap, Mt Kosciusko N.P.<br>N.S.W., 4 Feb. 1993, <i>Short 3993</i>             |       | 18 |
| Blue Lake to Charlotte Pass<br>Mt Kosciusko N.P., N.S.W.,<br>5 Feb. 1993, <i>Short 3997</i>                          |       | 18 |
| Mustering Flat, Baw Baw Plateau,<br>Vict., 26 Feb. 1991, <i>Walsh 3051</i>   |       | 18 |
| <b>B. oncocarpa</b> Diels<br>100 km N of Galena Bridge on NW<br>Coastal Hwy, W.A., 6 Oct. 1993,<br><i>Short 4084</i> |       | 18 |

TABLE 1. CONTINUED

| Species & locality   | n                  | 2n |
|--|--------------------|----|
| <i>B. papillosa</i> G.L.R.Davis<br>28.7 km S of Ivanhoe, N.S.W.,<br>6 Oct. 1992, <i>Watanabe 73</i>  | 4II or<br>4II + 1B |    |
| Muggabah Creek, 14 km N of Booligal,<br>N.S.W., 6 Oct. 1992, <i>Watanabe 75</i>  |                    | 8  |
| <i>B. perpusilla</i> (Steetz) J.M.Black<br>Totadgin Rock, W.A., 3 Oct. 1993,<br><i>Watanabe 115</i>  |                    | 18 |
| Bonnie Rock, W.A., 11 Oct. 1993,<br><i>Watanabe 160</i>  |                    | 18 |
| <i>B. petrophila</i> G.L.R.Davis<br>Angler's Rest, Vict.<br>Unvouchered colln, Feb. 1993   |                    | 18 |
| <i>B. procumbens</i> G.L.R.Davis<br>Mt Kaputar N.P., N.S.W.,<br>28 Jan. 1993, <i>Short 3951</i>  |                    | 18 |
| <i>B. ptychocarpa</i> F.Muell.<br>Mt Mittamatite Regional Park, Vict.,<br>3 Dec. 1993, <i>Short 4151</i>   | 6II                | 12 |
| <i>B. pusilla</i> Steetz<br>W edge of Lake King, W.A.,<br>1 Nov. 1995, <i>Short 4533</i>   | 9II                |    |
| <i>B. radicans</i> Steetz ex Lehm.<br>Mother of Ducks Nature Reserve, Guyra,<br>N.S.W., 31 Jan. 1993, <i>Short 3976</i>  | 13II               | 26 |
| <i>B. readeri</i> G.L.R.Davis<br>Ulupna Island, Vict.,<br>23 Sept. 1992, <i>Watanabe 4</i>   | 5II                | 10 |
| <i>B. rigidula</i> (DC.) G.L.R.Davis<br>Lake Omeo, c. 0.4 km S of<br>intersection of Blowhard road<br>and Lake Omeo road, Vict.,<br>9 Feb. 1993, <i>Short 4016</i> | 9II                | 18 |
| <i>B. scapigera</i> (Sieber ex Sprengel) DC.<br>3.5 km NW of Backwater, N.S.W.,<br>30 Jan. 1993, <i>Short 3972</i>   | 9II                | 18 |
| Buckety Plains, Bogong N.P., Vict.,<br>9 Feb. 1993, <i>Short 4023</i>  | 9II                |    |
| <i>B. smithwhitei</i> P.S.Short & K.Watan.<br>28.7 km S of Ivanhoe, N.S.W.,<br>6 Oct. 1992, <i>Watanabe 72</i>   | 6II or<br>3IV      |    |

TABLE 1. CONTINUED

| Species & locality   | n                           | 2n |
|--|-----------------------------|----|
| 1.4 km W of Menindee, N.S.W.,<br>22 Oct. 1993, <i>Watanabe 210</i>   | 6II or<br>3IV               |    |
| <i>B. spathulata</i> Gaudich.<br>Sawyers Hill, Kosciusko N.P., N.S.W.,<br>3 Feb. 1993, <i>Short 3985</i>   | 27II                        |    |
| Daners Gap, Kosciusko N.P., N.S.W.,<br>4 Feb. 1993, <i>Short 3994</i>  | 27II                        |    |
| Club Lake, Kosciusko N.P., N.S.W.,<br>5 Feb. 1993, <i>Short 4003</i>   | 9II                         | 18 |
| Cathcart to Rocky Hall, N.S.W.,<br>6 Feb. 1993, <i>Short 4011</i>  | 27II                        |    |
| <i>B. stolonifera</i> G.L.R.Davis<br>Club Lake, Kosciusko N.P., N.S.W.,<br>5 Feb. 1993, <i>Short 4006</i>  | 15II                        | 30 |
| <i>B. stuartii</i> Benth.<br>Girraween N.P., Qld, 29 Jan. 1993,<br><i>Watanabe 81</i>  | 6II                         | 12 |
| 11 km from Deepwater towards<br>Tenterfield, N.S.W., 29 Jan. 1993,<br><i>Short 3964</i>  | 6II                         |    |
| <i>B. tatei</i> J.M.Black<br>42.5 km E of Border Village, S.A.,<br>18 Oct. 1993, <i>Watanabe 184</i>   | 9II                         | 18 |
| <i>B. tenuiscapa</i> Hook.f.<br>var. <i>pubescens</i> (Benth.) G.L.R.Davis<br>3.5 km NW of Backwater, N.S.W.,<br>30 Jan. 1993, <i>Short 3975</i> | 9II                         |    |
| aff. var. <i>tenuiscapa</i><br>3.5 km NE of Mt Reynard, Snowy<br>Plains, Vict., <i>Walsh s.n.</i> ,<br>Cultivated RBG, Accession No. 905484      |                             | 28 |
| <i>B. trachycarpa</i> F.Muell.<br>Ceduna, S.A., 12 Sept. 1990,<br><i>Short 3781</i>  | 18 + 0-2Bs                  |    |
| <i>B. sp. aff. trachycarpa</i> F.Muell.<br>25 km W of Dalby, N.S.W.,<br>30 Sept. 1992, <i>Watanabe 31</i>  | 27I or 1III<br>+ 0-9II + Is |    |
| <i>Brachyscome</i> sp.<br>8 km NW of Glendambo, S.A.,<br>26 Aug. 1989, <i>Short 3673</i>   | 9II                         |    |

TABLE 1. CONTINUED

| Species & locality  | n          | 2n |
|---|------------|----|
| 119 km N of Glendambo, S.A.,<br>26 Aug. 1989, <i>Short 3682</i>   | 9II        |    |
| <b>C. calotis</b> R.Br.   |            |    |
| <i>C. anthemoides</i> F.Muell.<br>Ulupna Is., Vict.<br>26 Sept. 1993, <i>Watanabe 92</i>  | 14II       |    |
| <i>C. cuneata</i> (F.Muell. ex Benth.) G.L.R.Davis<br>var. <i>cuneata</i><br>25 km W of Dalby, N.S.W.,<br>30 Sept. 1992, <i>Watanabe 32</i> | 16II       |    |
| <i>C. cuneifolia</i> R.Br.<br>17 km SW of Dubbo, N.S.W.<br>24 Sept. 1992, <i>Watanabe 5</i>   | 8II        | 16 |
| 15.5 km E of Inglewood, Qld,<br>29 Sept. 1992, <i>Watanabe 23</i>   | 8II        |    |
| 20 km E of Elmore, Vict.,<br>7 Oct. 1992, <i>Watanabe 78</i>  | 8II        |    |
| 25 km W of Dalby, N.S.W.,<br>30 Sept. 1992, <i>Watanabe 34</i>  | 16II       |    |
| <i>C. cymbacantha</i> F.Muell.<br>Tilpa, N.S.W., 2 Oct. 1992,<br><i>Watanabe 48</i>   | 14II       |    |
| <i>C. dentex</i> R.Br.<br>3 km S of Glen Alpin, Qld,<br>30 Jan. 1993, <i>Short 3967</i>   | 8II        | 16 |
| <i>C. erinacea</i> Steetz<br>15 km SW of Kimba, S.A.,<br>21 Sept. 1982, <i>Short 1766</i>   | 12II + 1IV |    |
| <i>C. lappulacea</i> Benth.<br>15.5 km E of Inglewood, Qld,<br>29 Sept. 1992, <i>Watanabe 21</i>  | 14II       |    |
| <i>C. aff. lappulacea</i><br>15.5 km E of Inglewood, Qld,<br>29 Sept. 1992, <i>Watanabe 22</i>  | 8II        |    |
| <i>C. multicaulis</i> (Turcz.) Druce<br>38 km N of Pt Augusta, S.A.,<br>5 Oct. 1992, <i>Watanabe 62</i>                                     | 4II        |    |
| <i>C. plumulifera</i> F.Muell.<br>24 km NE of Wirrealpa, S.A.,<br>4 Oct. 1992, <i>Watanabe 56</i>   | 5II + BII  |    |

TABLE 1. CONTINUED

| Species & locality  | n    | 2n        |
|---|------|-----------|
| 72 km S of Charleville, Qld,<br>1 Oct. 1992, <i>Watanabe</i> 38   |      | 5II + BII |
| 45 km SE of Broken Hill, N.S.W.,<br>6 Oct. 1992, <i>Watanabe</i> 66   |      | 5II + 2BI |
| 80 km SW of Broken Hill, N.S.W.,<br>6 Oct. 1992, <i>Watanabe</i> 69   |      | 5II + 2BI |
| <i>C. scabiosifolia</i> Sond. & F.Muell<br>var. <i>scabiosifolia</i><br>15 km SW of Louth, N.S.W.,<br>2 Oct. 1992, <i>Watanabe</i> 45 | 8II  |           |
| 28.7 km S of Ivanhoe, N.S.W.,<br>23 Oct. 1993, <i>Watanabe</i> 213  | 8II  |           |
| Sawpit Creek, Kosciusko N.P.,<br>N.S.W., 4 Feb. 1993, <i>Short</i> 3995   | 16II |           |
| <i>C. scapigera</i> Hook.<br>Louth, N.S.W., 2 Oct. 1992,<br><i>Watanabe</i> 42  | 8II  |           |
| 6 km NE of Jerilderie, N.S.W.,<br>26 Jan. 1993, <i>Short</i> 3933   | 16II |           |
| <i>Erigeron</i> L.  |      |           |
| <i>E. karvinskianus</i> DC.<br>4 km N of Mt Slide, Vict.,<br>March 1991, <i>H.Manson</i> s.n.,<br>(MEL 1592608)                       |      | 36        |
| <i>Erodiophyllum</i> F.Muell.   |      |           |
| <i>E. elderi</i> F.Muell.<br>41 Km ENE of Iron Knob,<br>S.A., 12 Oct. 1990, <i>Short</i> 3779   |      | 16        |
| <i>Kippistia</i> F.Muell.   |      |           |
| <i>K. suaedifolia</i> F.Muell.<br>Lake Austin, W.A., 25 Aug. 1995,<br><i>Short</i> 4221   |      | 9II       |
| <i>Lagenifera</i> Cass.   |      |           |
| <i>L. huegellii</i> Benth.<br>4 km N of Zumstein, Grampians,<br>Vict., 26 Oct. 1993, <i>Watanabe</i> 225                              |      | 18        |
| <i>L. stipitata</i> (Labill.) Druce<br>Mt Kaputar N.P., N.S.W. 28 Jan. 1993<br><i>Short</i> 3952                                      | 9II  | 18        |
| Barington Tops State Forests,<br>N.S.W., 1 Feb. 1993, <i>Short</i> 3980   |      | 18        |

TABLE 1. CONTINUED

| Species & locality   | n    | 2n         |
|--|------|------------|
| <b>Minuria DC.</b>   |      |            |
| <i>M. cunninghamii</i> (DC.) Benth.<br>Lake Austin, W.A., 25 Aug. 1995<br><i>Short 4223</i>  | 9II  |            |
| Tilpa, N.S.W., 2 Oct. 1992<br><i>Watanabe 49</i>   | 9II  |            |
| <i>M. gardneri</i> Lander & R.Barry<br>Lake Austin, W.A., 25 Aug. 1995,<br><i>Short 4222</i>                                       | 9II  |            |
| <i>M. integrerrima</i> (DC.) Benth.<br>Dalby, N.S.W., 30 Sept. 1992,<br><i>Watanabe 30</i>   | 18II |            |
| 28 km W of Condamine, N.S.W.,<br>30 Sept. 1992, <i>Watanabe 35</i>   | 18II |            |
| 0.9 km SW of Louth, N.S.W.<br>2 Oct. 1992, <i>Watanabe 44</i>  | 18II |            |
| <i>M. leptophylla</i> DC.<br>Wudinna Hill, S.A., 19 Oct. 1993,<br><i>Watanabe 186</i>  | 9II  |            |
| 28.5 km N of Orroroo, S.A.,<br>21 Oct. 1993, <i>Watanabe 194</i>   | 9II  | 18 + 0-2Bs |
| <b>Olearia Moench</b>  |      |            |
| <i>O. astrotricha</i> (F.Muell.) F.Muell. ex Benth.<br>Victoria Range road, Grampians,<br>Vict., 2 Dec. 1986, <i>Corrick 10103</i> | 9II  |            |
| <i>O. ciliata</i> (Benth.) F.Muell. ex Benth.<br>10 km SE of Ravensthorpe, W.A.,<br>3 Sept. 1986, <i>Short 2663</i>                | 9II  |            |
| Between Karkoo and Mount Hill, S.A.,<br>29 Sept. 1993, <i>Watanabe 100</i>   | 9II  |            |
| <i>O. ferresii</i> (F.Muell.) F.Muell. ex Benth.<br>S. boundary of Ormiston Gorge N.P.,<br>N.T., 12 Aug. 1988, <i>Short 3148</i>   | 9II  |            |
| <i>O. humilis</i> Lander<br>Sandstone to Yuinmery, W.A.,<br>14 Oct. 1986, <i>Short 2563</i>  | 9II  |            |
| Yuinmery, W.A., 14 Oct. 1986,<br><i>Short 2565</i>   | 9II  |            |

TABLE 1. CONTINUED

| Species & locality   | n   | 2n       |
|--|-----|----------|
| <i>O. imbricata</i> (Turcz.) Benth.<br>10 km SE of Ravensthorpe, W.A.,<br>3 Sept. 1986, <i>Short 2662</i>          | 9II |          |
| <i>O. phlogopappa</i> (Labill.) DC. complex<br>Falls Creek, Vict., 2 March 1987<br><i>Short 3035</i>               |     | 18II     |
| <i>O. pimelioides</i> (DC.) Benth.<br>53.5 km E of 'Nallan', W.A.,<br>1986, <i>Lander 1391</i>                     | 9II |          |
| Sandstone-Paynes Find road, W.A.,<br>24 Aug. 1986, <i>Short 2564</i>   | 9II |          |
| Parachilna to Blinman road, S.A.,<br>20 Oct. 1993, <i>Watanabe 193</i>   |     | 18 + 2Bs |
| <i>O. plucheacea</i> Lander<br>Kennedy Range, W.A.,<br>20 Aug. 1986, <i>Short 2535</i>                             | 9II |          |
| <i>O. ramulosa</i> (Labill.) Benth. complex<br>Golton Gorge, Grampians,<br>Vict. 11 Sept. 1986, <i>RBG 86/1735</i> | 9II |          |
| <i>O. rufis</i> (Benth.) Benth.<br>19 km SW of Three Springs, W.A.,<br>9 Sept. 1986, <i>Short 2795</i>             | 9II |          |
| 10 km S of Mt Hope, N.S.W.,<br>16 Sept. 1987, <i>Short 3078</i>  | 9II |          |
| 17 km N of Patchewollock, Vict.,<br>25 Aug. 1988, <i>Short 3174</i>  | 9II |          |
| <i>O. stuartii</i> (F.Muell.) Benth.<br>Kennedy Range, W.A., 20 Aug. 1986<br><i>Short 2538</i>                     | 9II |          |
| <i>O. xerophila</i> (F.Muell.) Benth.<br>Dales Gorge, W.A., 30 Aug. 1995,<br><i>Short 4275</i>                     | 9II |          |
| <i>Olearia sp.</i><br>23.6 km N of Maroubra Rd/Scotsman Rd<br>intersection, W.A., 1986, <i>Lander 1408</i>         | 9II |          |
| <i>Solenogyne</i> Cass.<br><i>S. dominii</i> L.G.Adams<br>Ulupna Island, Vict.,<br>25 Jan. 1993, <i>Short 3923</i> |     | 18       |

TABLE I. CONTINUED

| Species & locality   | n   | 2n |
|--|-----|----|
| <b>Vittadinia A.Rich.</b>  |     |    |
| <i>V. cuneata</i> DC.  |     |    |
| var. <i>hirsuta</i> N.T.Burb.<br>4 km NE of Milmerran, Qld,<br>29 Sept. 1992, Watanabe 26                        | 9II | 18 |
| <i>V. gracilis</i> (Hook.f.) N.T.Burb.<br>2 km W of Hines Hill, W.A.,<br>23 Oct. 1995, Short 4450                | 9II |    |
| Ulupna Island, Vict.,<br>25 Jan. 1993, Short 3925  | 9II |    |
| <i>V. muelleri</i> N.T.Burb.<br>44 km NE of Narrabri, N.S.W.,<br>26 Sept. 1992, Watanabe 14                      | 9II |    |
| <i>V. pterochaeta</i> (F.Muell. ex Benth.) J.M.Black<br>24 km NE of Wirrealpa, S.A.,<br>3 Oct. 1992, Watanabe 57 | 9II |    |
| <i>V. pustulata</i> N.T.Burb.<br>28 km W of Condamine, N.S.W.,<br>30 Sept. 1992, Watanabe 36                     | 9II |    |
| <i>V. sulcata</i> N.T.Burb.<br>c. 10 km S. of Wild Dog Glen, S.A.,<br>20 Oct. 1993, Watanabe 190                 | 9II |    |

## BRACHYSCOME

As in Watanabe & Short (1992) our references to species and species complexes in Tables I & II are largely the concepts of Davis (1948, 1949, 1955, 1959) and Smith-White *et al.* (1970) although the concept of 'superspecies' is not used. As we are primarily concerned here with the presentation of new chromosome number determinations and with reviewing chromosome numbers in the Australian Astereae, further comments on species or generic delimitation are here kept to a minimum. Such problems will be the subject of future papers and will follow after the completion of detailed anatomical, morphological, cytological and macromolecular studies.

As currently recognised *Brachyscome* is a genus of more than 80 species and occurs in Australia, New Guinea and New Zealand. Of these, over 70 species are confined to Australia. Davis (1948) recognised two subgenera within *Brachyscome*, i.e. '*Eubrachyscome*' and '*Metabrachyscome*'. They are illegitimate names but are frequently adopted in cytological papers (e.g. Smith-White *et al.* 1970).

The majority of species were referred by Davis to 'subgenus *Eubrachyscome*' and this group was the subject of cytological investigations by Smith-White *et al.* (1970). They found an array of chromosome numbers in the group and suggested that it represents a reducing series, i.e.  $x = 9$  to  $x = 2$ . It was also noted that most species with  $n = 9$  (including polyploids) are mesic, perennial species. In contrast, species of arid regions frequently display an annual habit and this is correlated with lower chromosome numbers. Chromosome number determinations by Watanabe & Short (1992), and in this paper for species not examined by Smith-White *et al.*, are consistent with these observations. Thus *B. goniocarpa* ( $n = 4$ ), *B. gracilis* ( $n = 4$ ), *B. muelleri* ( $n = 3$ ) and *B. muelleroides* ( $n = 3$ ) are all annual species which occur in arid, or seasonally dry areas.

As noted by Carter (1978a), species referred by Davis to 'subgenus *Metabrachyscome*' are chromosomally conservative compared to members of '*Ebrachyscome*'. All species have  $x = 9$ , with polyploidy mainly confined to the *B. ciliaris* complex. Many of the species are also annuals confined to arid regions.

The *B. iberidifolia* complex referred to in both Tables is mainly confined to Western Australia and includes *B. bellidioides* and *B. pusilla*. The complex belongs to 'subgenus *Metabrachyscome*'. There is considerable diversity in collections referred by us to this complex and additional taxa undoubtedly should be recognised.

The chromosome complements of *B. nivalis* ( $2n = 22$ ), *B. radicans* ( $2n = 26$ ), *B. tenuiscapa* aff. var. *tenuiscapa* ( $2n = 28$ ) and *B. stolonifera* ( $2n = 30$ ) are markedly different from those found in most other species, their karyotypes displaying two different sizes of chromosomes. This is in marked contrast to the unimodal karyotypes found in species with  $2n = 18$ . However, species with chromosome numbers less than  $2n = 18$  frequently have bimodal karyotypes, a feature explained by descending aneuploidy caused by reciprocal translocation and loss of centromeric fragments. Species with chromosome numbers greater than  $2n = 18$  are perhaps amphidiploids originating as a result of hybridization between taxa with chromosome numbers less than  $n = 9$ .

Both the array of chromosome numbers and the discrepancies in numbers reported by us with those published by Smith-White *et al.* (1970) for *B. diversifolia*, *B. microcarpa*, *B. multifida* and *B. nova-anglica* may in part be the result of misidentification. However, they primarily reflect the morphological complexity of the taxa concerned. More work is required to clarify the delimitation of the various entities in these complexes and therefore discussion of the significance of chromosome numbers in these taxa is premature.

#### CALOTIS

A genus containing 28 formally recognised species *Calotis* mainly occurs in semi-arid and arid regions of Australia.

Stace (1978, 1982) has reported on the cytoevolution of the genus and noted two significant trends, i.e. a reduction from a base of  $x = 8$  down to  $x = 4$  in the mainly arid zone annuals, and a high degree of polyploidy. Two-thirds of the species were found to be polyploids and this condition was found to be as frequent in low chromosome number annuals as in high chromosome number perennials.

Our data presented in Table I support the observations made by Stace.

#### CELMISIA

*Celmisia* contains about 70 species confined to Australia and New Zealand. They are mainly found in alpine regions and most are endemic to New Zealand, only about ten species occurring in Australia. Celmisas are perennial herbs or subshrubs of diverse habit and a cytological survey of 65 species (Hair 1980, Given & Gray 1986) indicates that most are at least 12-ploid, with  $n = 54$ . Haploid numbers of  $n = 54$  and  $n = 108$  have been found in Australian species.

#### CERATOGYNE

*Ceratogyne obionoides*, the only member of this genus, is an annual species with  $n = 6$  and is found throughout much of arid Australia.

#### ERODIOPHYLLUM

A ditypic genus, *Erodiophyllum* occurs in semi-arid and arid mainland Australia. Both species are perennial herbs with  $n = 8$ .

TABLE II. SUMMARY OF CHROMOSOME NUMBER DETERMINATIONS IN NATIVE AUSTRALIAN ASTEREAE

|  | n               | 2n           |   |
|--|-----------------|--------------|---|
| <b>Brachyscome Cass.</b>   |                 |              |   |
| <i>B. aculeata</i> (Labill.) Cass. ex Lessing                    | 9,18<br>0-6Bs   | 18           | Stace 1981;<br>Watanabe <i>et al.</i> 1996  |
| <i>B. aff. aculeata</i><br>(Mt Gingera)                          | 9 + 1B          |              | Stace 1981  |
| <i>B. aff. aculeata</i><br>(Halls Gap)                           | 27              |              | Stace 1981  |
| <i>B. angustifolia</i> A.Cunn.ex DC.<br>var. <i>angustifolia</i> | 9               |              | Smith-White <i>et al.</i><br>1970   |
| var. <i>heterophylla</i> (Benth.)<br>G.L.R.Davis                 | 9               |              | Smith-White <i>et al.</i><br>1970   |
| <i>B. sp. aff. angustifolia</i>                                  | 5               | 10           | Watanabe <i>et al.</i> 1996   |
| <i>B. basaltica</i> F.Muell.<br>var. <i>basaltica</i>            | 8               | 16           | Smith-White <i>et al.</i> 1970;<br>Watanabe & Short 1992;<br>Watanabe <i>et al.</i> 1996                                    |
| var. <i>gracilis</i> Benth.                                      | 6               | 12           | Smith-White <i>et al.</i> 1970;<br>Watanabe & Short 1992;<br>Watanabe <i>et al.</i> 1996                                    |
| <i>B. breviscapis</i> C.R.Carter                                 | 4               | 8            | De Jong 1963; Smith-<br>White <i>et al.</i> 1970; Carter<br>1978c; Watanabe &<br>Short 1992                                 |
| <i>B. campylocarpa</i> J.M.Black                                 | 5               | 10           | Smith-White <i>et al.</i> 1970,<br>as 'B. campylocarpa sp.<br>B', Watanabe & Short<br>1992                                  |
| <i>B. cardiocarpa</i> F.Muell. ex Benth.                         | 9               | 36           | Smith-White <i>et al.</i> 1970;<br>Watanabe <i>et al.</i> 1996  |
| <i>B. cheilocarpa</i> F.Muell.                                   | 9               |              | Carter 1978a; Watanabe<br><i>et al.</i> 1996  |
| <i>B. aff. cheilocarpa</i> (A)                                   |                 | 18           | Watanabe <i>et al.</i> 1996   |
| <i>B. aff. cheilocarpa</i> (B)                                   | 9               |              | Watanabe <i>et al.</i> 1996   |
| <i>B. chrysoglossa</i> F.Muell.                                  | 4               | 8 +<br>0-3Bs | Smith-White <i>et al.</i> 1970;<br>Watanabe & Short 1992  |
| <i>B. ciliaris</i> (Labill.) Less. complex                       | 9,18<br>27, 36I | 27,36<br>81  | De Jong 1963; Smith-<br>White <i>et al.</i> 1970; Carter<br>1978a; Watanabe &<br>Short 1992; Watanabe<br><i>et al.</i> 1996 |




Fig. 1. Mitotic and meiotic chromosomes in Australian Astereae. A - *Brachyscome muellerioides*  $n = 3II$  (Watanabe 3). B - *B. readeri*  $n = 5II$  (Watanabe 4). C - *B. sp. aff. nova-anglica*  $n = 5II$  (Short 3969). D - *B. sp. aff. angustifolia*  $2n = 10$  (Short 3979). E - *B. tatei*  $n = 9II$  (Watanabe 184). F - *Minuria leptophylla*  $2n = 18$  (Watanabe 194). G - *Vittadinia gracilis*  $n = 9II$  (Short 3925). H - *Erodiocephalum elderi*  $2n = 16$  (Short 3779). Scale: 10  $\mu\text{m}$ .

TABLE II. CONTINUED

|  | <b>n</b>                   | <b>2n</b>                  |   |
|--|----------------------------|----------------------------|---|
| <i>B. ciliocarpa</i> W.Fitzg.  | 9                          | 18 +<br>0-2Bs              | Carter 1978a; Watanabe & Short 1992; Watanabe et al. 1996   |
| <i>B. sp. aff. ciliocarpa</i>  | 9                          |                            | Watanabe et al. 1996  |
| <i>B. cuneifolia</i> Tate  | 9                          | 18                         | Stace 1981  |
| <i>B. curvicarpa</i> G.L.R.Davis   | 4                          | 8 +<br>0-5Bs               | Smith-White et al. 1970;<br>Watanabe & Short 1992;<br>Watanabe et al. 1996  |
| <i>B. debilis</i> Sond.  | 3                          | 6                          | Smith-White et al. 1970;<br>Watanabe & Short 1992;<br>Watanabe et al. 1996  |
| <i>B. decipiens</i> Hook.f.  | 9,27                       | 18,54                      | Solbrig et al. 1964;<br>Smith-White et al. 1970;<br>Watanabe et al. 1996  |
| <i>B. dentata</i> Gaudich.   | 4,8,12<br>+ 0-4Bs          | 8,16<br>24 +<br>0-4Bs      | Smith-White et al. 1970;<br>Watanabe & Short 1992; Watanabe et al. 1996   |
| <i>B. dichromosomatica</i> C.R.Carter                                      | 2 +<br>0-3Bs +<br>micro Bs | 4 +<br>0-3Bs<br>+ micro Bs | Smith-White 1968,<br>as <i>B. lineariloba</i> ;<br>Smith-White et al. 1970, as 'B.<br><i>lineariloba</i> sp. A';<br>Smith-White & Carter 1970, as 'B.<br><i>lineariloba</i> sp. A';<br>Watanabe et al. 1975, as 'B.<br><i>lineariloba</i> race A';<br>Carter 1978b; Carter 1978c; Smith-White & Carter 1981;<br>Nagl & Pfeifer 1988;<br>Watanabe et al. 1991; Watanabe & Short 1992 |
| <i>B. dissectifolia</i> G.L.R.Davis  | 6,12                       | 12,24                      | Smith-White et al. 1970; Watanabe et al. 1996   |
| <i>B. diversifolia</i> (Hook.) Fischer & C. Meyer var. <i>diversifolia</i> | 12,16<br>18,c.20           | 24,<br>36 +<br>0-1B        | Smith-White et al. 1970; Watanabe & Short 1992;<br>Watanabe et al. 1996   |

TABLE II. CONTINUED

|  | n           | 2n |  |
|--|-------------|----|--|
| <i>B. eriogona</i> (J.M.Black) G.L.R.Davis | 4           | 8  | Smith-White <i>et al.</i><br>1970, as ' <i>B. campylocarpa</i> sp. A';<br>Watanabe & Short<br>1992   |
| <i>B. exilis</i> Sond.                     | 9           | 18 | Carter 1978a;<br>Watanabe <i>et al.</i> 1996   |
| <i>B. formosa</i> P.S.Short                | 9           |    | Short 1988;<br>Watanabe <i>et al.</i> 1996   |
| <i>B. glandulosa</i> (Steetz) Benth.       |             | 36 | Watanabe <i>et al.</i> 1996  |
| <i>B. goniocarpa</i> Sond. & F.Muell.      | 4           | 8  | Watanabe <i>et al.</i><br>1991; Watanabe &<br>Short 1992   |
| <i>B. gracilis</i> G.L.R.Davis             | 4           | 8  | Smith-White <i>et al.</i><br>1970, as <i>B. diversifolia</i> var.<br><i>dissecta</i> G.L.R.<br>Davis; Watanabe &<br>Short 1992;<br>Watanabe <i>et al.</i> 1996   |
| <i>B. graminea</i> (Labill.) F.Muell.      | 9           | 18 | Smith-White <i>et al.</i><br>1970; Watanabe <i>et<br/>al.</i> 1996   |
| <i>B. halophila</i> P.S.Short              | 9           | 18 | Short 1988;<br>Watanabe & Short<br>1992; Watanabe <i>et<br/>al.</i> 1996   |
| <i>B. iberidifolia</i> Benth.              | 9 +<br>0-1B | 18 | De Jong 1963;<br>Chouksanova <i>et al.</i><br>1968; Gupta 1969;<br>Turner 1970,<br>including specimens<br>referred to <i>B. bellidioides</i> ;<br>Carter 1978a;<br>Keighery 1978;<br>Gupta & Gill 1983,<br>1989; Watanabe <i>et<br/>al.</i> 1996 |
| <i>B. latisquamea</i> F.Muell.             | 9           | 18 | Carter 1978a;<br>Watanabe <i>et al.</i> 1996   |
| <i>B. leptocarpa</i> F.Muell.              | 3           | 6  | Smith-White <i>et al.</i><br>1970, probably  |

TABLE II. CONTINUED

|  | n                  | 2n            |  |
|--|--------------------|---------------|--|
|  |                    |               | conspecific with <i>B. debilis</i>   |
| <i>B. lineariloba</i> (DC.) Druce      | 6,8<br>4II + 2I    | 10,12<br>16   | Smith-White 1968;<br>Smith-White <i>et al.</i><br>1970; Carter &<br>Smith-White 1972;<br>Kyhos <i>et al.</i> 1977;<br>Watanabe <i>et al.</i><br>1985; Watanabe &<br>Smith-White 1987;<br>Watanabe & Short<br>1992; Watanabe <i>et al.</i><br>1996; Watanabe <i>et al.</i> 1996 |
| <i>B. lyrifolia</i> J.M.Black          | 9                  | 18            | Smith-White <i>et al.</i><br>1970 and Carter<br>1978, as to Parachila<br>collections of <i>B. ciliaris</i> ; Watanabe &<br>Short 1992  |
| <i>B. melanocarpa</i> Sond. & F.Muell. | 6,12               | 12,30         | Smith-White <i>et al.</i><br>1970; Watanabe &<br>Short 1992; Watanabe <i>et al.</i><br>1996  |
| <i>B. microcarpa</i> F.Muell.          | 6, 10<br>10II + 4I | 12            | Smith-White <i>et al.</i><br>1970; Watanabe <i>et al.</i> 1996   |
| <i>B. muelleri</i> Sond.               | 3                  | 6             | Watanabe & Short<br>1992   |
| <i>B. muelleroides</i> G.L.R.Davis     | 3                  | 6             | Watanabe <i>et al.</i> 1996  |
| <i>B. multicaulis</i> F.Muell.         | 9                  | 18            | Watanabe & Short<br>1992, as <i>B. rigidula</i>  |
| <i>B. aff. multicaulis</i>             | 9                  | 18 +<br>0-2Bs | Watanabe <i>et al.</i> 1996  |
| <i>B. multifida</i> DC. complex        | 7,9                | 14,18         | Smith-White <i>et al.</i><br>1970, including var.<br><i>dilatata</i> Benth. and<br>var. <i>multifida</i> ;<br>Watanabe & Short<br>1992; Watanabe <i>et al.</i><br>1996   |
| <i>B. nivalis</i> F.Muell.             | 11                 | 22            | Smith-White <i>et al.</i><br>1970, excluding<br>record of n = 9,<br>re Stace 1981;<br>Watanabe <i>et al.</i> 1996  |

TABLE II. CONTINUED

|   | n         | 2n    |  |
|---|-----------|-------|--|
| <i>B. nodosa</i> P.S.Short & K.Watan.   | 3         | 6     | Smith-White <i>et al.</i> 1970, as ' <i>B. goniocarpa</i> '; Watanabe & Short 1992, as ' <i>B. sp. aff. goniocarpa</i> '; Short & Watanabe 1993; Watanabe <i>et al.</i> 1996 |
| <i>B. nova-anglica</i> G.L.R.Davis      | 7         | 14    | Smith-White <i>et al.</i> 1970, a count of 3n = 18 is probably erroneous; Watanabe <i>et al.</i> 1996  |
| <i>B. aff. nova-anglica (A)</i>         | 5         | 10    | Watanabe <i>et al.</i> 1996  |
| <i>B. aff. nova-anglica (B)</i>         | 6         | 12    | Smith-White <i>et al.</i> 1970, as <i>B. nova-anglica</i> ; Watanabe <i>et al.</i> 1996  |
| <i>B. obovata</i> G.L.R.Davis           |           | 18    | Watanabe <i>et al.</i> 1996  |
| <i>B. oncocarpa</i> Diels               | 9         | 18    | Carter 1978a; Watanabe <i>et al.</i> 1996  |
| <i>B. papillosa</i> G.L.R. Davis        | 4 + 0-2Bs | 8     | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996  |
| <i>B. parvula</i> Hook.f.               | 9         |       | Smith-White <i>et al.</i> 1970; Carter 1978a   |
| <i>B. perpusilla</i> (Steetz) J.M.Black | 9         | 18,36 | Smith-White <i>et al.</i> 1970; Carter 1978a; Watanabe & Short 1992; Watanabe <i>et al.</i> 1996   |
| <i>B. petrophila</i> G.L.R. Davis       |           | 18    | Watanabe <i>et al.</i> 1996  |
| <i>B. procumbens</i> G.L.R.Davis        | 9         | 18    | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996  |
| <i>B. ptychocarpa</i> F.Muell.          | 6         | 12    | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996  |
| <i>B. pusilla</i> Steetz                | 9         | 18    | Carter 1978a; Watanabe <i>et al.</i> 1996  |

TABLE II. CONTINUED

|  | <b>n</b>         | <b>2n</b>       |   |
|--|------------------|-----------------|---|
| <i>B. radicans</i> Steetz  | 13               | 26              | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996   |
| <i>B. rara</i> G.L.R. Davis  | 6                | 12              | Watanabe & Short 1992   |
| <i>B. readeri</i> G.L.R. Davis   | 5                | 10              | Watanabe & Short 1992; Watanabe <i>et al.</i> 1996  |
| <i>B. rigidula</i> (DC.) G.L.R. Davis  | 9                | 18              | ?Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996  |
| <i>B. riparia</i> G.L.R. Davis   | 9                |                 | Watanabe & Short 1992   |
| <i>B. scapigera</i> (Sprengel) DC.   | 9                | 18              | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996   |
| <i>B. sieberi</i> DC.<br>var. <i>gunnii</i> DC.                                | 9 + 0-1B         |                 | Stace 1981  |
| <i>B. smithwhitei</i> P.S. Short & K. Watan.                                   | 3,6              | 6,12 +<br>0-2Bs | Smith-White <i>et al.</i> 1970, as ' <i>B. campylocarpa</i> sp. C'; Watanabe & Short 1992, as ' <i>B. sp. aff. campylocarpa</i> '; Short & Watanabe 1993; Watanabe <i>et al.</i> 1996 |
| <i>B. spathulata</i> Gaudich.<br>subsp. <i>glabra</i> (DC.) Stace              | 9                | 18              | Stace 1981  |
| subsp. <i>spathulata</i>   | 9,18,27<br>36,45 |                 | Stace 1981;<br>Watanabe <i>et al.</i> 1996  |
| <i>B. stolonifera</i> G.L.R. Davis   | 15               | 30              | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996   |
| <i>B. stuartii</i> Benth.  | 6                | 12              | Smith-White <i>et al.</i> 1970; Watanabe <i>et al.</i> 1996   |
| <i>B. tatei</i> J.M. Black   | 9                | 18              | Watanabe <i>et al.</i> 1996   |
| <i>B. tenuiscapa</i> Hook.f.<br>var. <i>pubescens</i> (Benth.)<br>G.L.R. Davis | 9                | 18              | Smith-White <i>et al.</i>   |

TABLE II. CONTINUED

|   | <b>n</b>                       | <b>2n</b>         |  |
|---|--------------------------------|-------------------|--|
|   |                                |                   | 1970; Watanabe <i>et al.</i> 1996  |
| <i>aff. var. tenuiscapa</i>                           |                                | 28                | Watanabe <i>et al.</i> 1996  |
| <i>B. tesquorum</i> J.M.Black                         | 9                              | 18                | Smith-White <i>et al.</i> 1970   |
| <i>B. tetrapterocarpa</i> G.L.R. Davis                | 4 + 0-1B                       | 8                 | Watanabe & Short 1992  |
| <i>B. trachycarpa</i> F.Muell.                        | 27                             | 18 + 0-2Bs,<br>36 | De Jong 1963; Carter 1978a; Watanabe <i>et al.</i> 1996                        |
| <i>B. sp. aff. trachycarpa</i>                        | 27I or<br>1III +<br>0-9II + Is |                   | Watanabe <i>et al.</i> 1996  |
| <i>B. uliginosa</i> G.L.R.Davis                       | 9                              |                   | Smith-White <i>et al.</i> 1970   |
| <i>B. whitei</i> G.L.R.Davis                          | 5                              | 10                | Smith-White <i>et al.</i> 1970; Watanabe & Short 1992                          |
| <i>Brachyscome</i> sp.                                | 9                              |                   | Watanabe <i>et al.</i> 1996, referred to <i>B. iberidifolia</i> , Fl. S. Aust. |
| <i>Calotis</i> R.Br.                                  |                                |                   |  |
| <i>C. cancyrocarpa</i> J.M.Black                      | 4                              | 8                 | Stace 1978   |
| <i>C. anthemoides</i> F.Muell.                        | 7,14                           | 14                | Solbrig <i>et al.</i> 1964;<br>Stace 1978;<br>Watanabe <i>et al.</i> 1996      |
| <i>C. cuneata</i> (F.Muell. ex Benth.)<br>G.L.R.Davis | 16                             | 32 + 0-10Bs       | Stace 1978;<br>Watanabe <i>et al.</i> 1996                                     |
| <i>C. cuneifolia</i> R.Br.                            | 8,16                           | 16,32             | Solbrig <i>et al.</i> 1964;<br>Stace 1978;<br>Watanabe <i>et al.</i> 1996      |
| <i>C. cymbacantha</i> F.Muell.                        | 7,14                           | 28                | Stace 1978;<br>Watanabe <i>et al.</i> 1996                                     |
| <i>C. dentex</i> R.Br.                                | 8                              | 16                | Stace 1978;<br>Watanabe <i>et al.</i> 1996                                     |

TABLE II. CONTINUED

|   | n             | 2n          |  |
|---|---------------|-------------|--|
| <i>C. erinacea</i> Steetz   | 7,14<br>21,28 | 14,28<br>56 | Turner 1970; Stace<br>1978; Watanabe <i>et al.</i> 1996  |
| <i>C. glandulosa</i> F.Muell.   | 8             | 16          | Stace 1978   |
| <i>C. hispidula</i> (F.Muell.) F. Muell.  |               | 16          | Stace 1978   |
| <i>C. inermis</i> Maiden & Betche   | 8             | 16          | Stace 1978   |
| <i>C. lappulacea</i> Benth.   | 14            |             | Stace 1978;<br>Watanabe <i>et al.</i> 1996   |
| <i>C. ? aff. lappulacea</i>   | 8             |             | Watanabe <i>et al.</i> 1996  |
| <i>C. latiuscula</i> F.Muell. & Tate  | 7,14<br>28    | 14,21       | Stace 1978, 1982   |
| <i>C. multicaulis</i> (Turcz.) Druce  | 4,8           | 8,16        | Turner 1970; Stace<br>1978, as <i>C.<br/>multicaulis</i> sp. B;<br>Watanabe <i>et al.</i> 1996 |
| <i>C. plumulifera</i> F.Muell.  | 5 +<br>02Bs   | 10,20       | Stace 1978, as<br><i>C. multicaulis</i> sp. B;<br>Watanabe <i>et al.</i> 1996                  |
| <i>C. porphyroglossa</i> F. Muell.  | 10            | 20          | Stace 1978   |
| <i>C. scabiosifolia</i> Sond. & F.Muell.<br>var. <i>integrifolia</i> F.Muell. ex Benth. | 8,16          | 16,32       | Stace 1978   |
| var. <i>scabiosifolia</i>   | 8,16          | 16,32       | Stace 1978;<br>Watanabe <i>et al.</i> 1996   |
| <i>C. scapigera</i> Hook.   | 8,16          |             | Stace 1982;<br>Watanabe <i>et al.</i> 1996   |
| <i>C. squamigera</i> C.T.White  |               | 16          | Stace 1978   |
| <i>C. xanthosoidea</i> Domin  | 8             | 16          | Stace 1978   |
| <i>Celmisia</i> Cass.<br><i>C. asteliifolia</i> Hook.f. complex                         | 54<br>108     | 108<br>216  | Hair 1980  |
| <i>C. longifolia</i> Cass. complex  | 54<br>108     | 108         | Hair 1980  |
| <i>C. sericophylla</i> J.H.Willis   |               | 108         | Hair 1980  |
| <i>Ceratogyne</i> Turcz.<br><i>C. obionoides</i> Turcz.                                 |               | 6           | Turner 1970  |

TABLE II. CONTINUED

|  | n          | 2n            |  |
|--|------------|---------------|--|
| <b>Erodiophyllum</b> F.Muell.                          |            |               |  |
| <i>E. acanthocephalum</i> Stapf                        | 8          |               | Solbrig <i>et al.</i> 1964   |
| <b>E. elderi</b> F.Muell.                              |            | 16            | Watanabe <i>et al.</i> 1996  |
| <b>Isoetopsis</b> Turcz.                               |            |               |  |
| <i>I. graminifolia</i> Turcz.                          | 17         |               | Turner 1970  |
| <b>Kippistia</b> F.Muell.                              |            |               |  |
| <i>K. suaedifolia</i> F.Muell.                         | 9          |               | Short 1986;<br>Watanabe <i>et al.</i> 1996                         |
| <b>Lagenifera</b> Cass.                                |            |               |  |
| <i>L. huegelii</i> Benth.                              | 9          | 18            | Turner 1970;<br>Watanabe <i>et al.</i> 1996                        |
| <i>L. stipitata</i> (Labill.) Druce                    | 9          | 18            | Smith-White <i>et al.</i><br>1970; Watanabe <i>et<br/>al.</i> 1996 |
| <b>Minuria</b> DC.                                     |            |               |  |
| <i>M. cunninghamii</i> (DC.) Benth.                    | 9          | 18            | Turner 1970;<br>Watanabe <i>et al.</i> 1996                        |
| <i>M. gardneri</i> Lander & R.Barry                    | 9          |               | Watanabe <i>et al.</i> 1996  |
| <i>M. integerrima</i> (DC.) Benth.                     | 18         |               | Watanabe <i>et al.</i> 1996  |
| <i>M. leptophylla</i> DC                               | 9<br>0-2Bs | 18 +<br>1986; | Turner 1970; Short<br>Watanabe <i>et<br/>al.</i> 1996              |
| <b>Olearia</b> Moench                                  |            |               |  |
| <i>O. adenolasia</i> F.Muell.                          | 9          |               | Turner 1970  |
| <i>O. algida</i> N.A.Wakef.                            |            | 18            | Beuzenberg & Hair<br>1984  |
| <i>O. argophylla</i> F.Muell.                          | 54         |               | Solbrig <i>et al.</i> 1964   |
| <i>O. astroloba</i> Lander & N.G.Walsh                 | 9          |               | Short in Lander &<br>Walsh 1989                                    |
| <i>O. astrotricha</i> (F.Muell.)<br>F.Muell. ex Benth. | 9          |               | Watanabe <i>et al.</i> 1996  |
| <i>O. axillaris</i> (DC.) F.Muell. ex Benth.           | 9          |               | Solbrig <i>et al.</i> 1964   |
| <i>O. ciliata</i> (Benth.) F.Muell. ex Benth.          | 9          |               | Turner 1970;<br>Watanabe <i>et al.</i> 1996                        |
| <i>O. ferresii</i> (F.Muell.) F.Muell.<br>ex Benth.    | 9          |               | Watanabe <i>et al.</i> 1996  |

TABLE II. CONTINUED

|  | n    | 2n            |  |
|--|------|---------------|--|
| <i>O. floribunda</i> (Hook.f.) Benth.              | 9    |               | Pai 1964   |
| <i>O. frostii</i> (F.Muell.) J.H.Willis            |      | 18            | Beuzenberg & Hair<br>1984  |
| <i>O. humilis</i> Lander                           | 9    |               | Turner 1970, as<br><i>Olearia</i> sp.; Short in<br>Lander 1989;<br>Watanabe <i>et al.</i> 1996 |
| <i>O. imbricata</i> (Turcz.) Benth.                | 9    |               | Watanabe <i>et al.</i> 1996  |
| <i>O. muelleri</i> (Sond.) Benth.                  | 9    |               | Turner 1970  |
| <i>O. pannosa</i> Hook.                            |      | 90            | Beuzenberg & Hair<br>1984  |
| <i>O. phlogopappa</i> (Labill.) DC. complex        | 9,18 |               | Solbrig <i>et al.</i> 1964;<br>Watanabe <i>et al.</i> 1996                                     |
| <i>O. pimelioides</i> (DC.) Benth.                 | 9    | 18 +<br>0-2Bs | Turner 1970, and as<br><i>O. propinqua</i> ;<br>Watanabe <i>et al.</i> 1996                    |
| <i>O. plucheacea</i> Lander                        | 9    |               | Short in Lander<br>1990; Watanabe <i>et<br/>al.</i> 1996                                       |
| <i>O. ramulosa</i> (Labill.) Benth. complex        | 9    |               | Watanabe <i>et al.</i> 1996  |
| <i>O. rудis</i> (Benth.) Benth.                    | 9    |               | Turner 1970;<br>Watanabe <i>et al.</i> 1996  |
| <i>O. stuartii</i> (F.Muell.) F.Muell.<br>ex Benth | 9    |               | Short in Lander<br>1989; Watanabe <i>et<br/>al.</i> 1996.                                      |
| <i>O. xerophila</i> (F.Muell.) Benth.              | 9    |               | Watanabe <i>et al.</i> 1996  |
| <i>Olearia</i> sp.                                 | 9    |               | Watanabe <i>et al.</i><br>1996, syn. <i>Eurybia</i><br><i>dampieri</i> DC.                     |
| <i>Solenogyne</i> Cass.                            |      |               |  |
| <i>S. bellidioides</i> Cass.                       | 9    |               | Smith-White <i>et al.</i><br>1970; Adams 1979  |
| <i>S. dominii</i> L.G.Adams                        | 9    | 18            | Adams 1979;<br>Watanabe <i>et al.</i> 1996   |
| <i>S. gunnii</i> (Hook.f.) Cabrera                 | 9    |               | Adams 1979   |
| <i>Vittadinia</i> A.Rich.                          |      |               |  |
| <i>V. cuneata</i> DC.                              |      |               |  |

TABLE II. CONTINUED

|  | n | 2n |   |
|--|---|----|---|
| <i>var. hirsuta</i> N.T.Burb.  | 9 | 18 | Watanabe <i>et al.</i> 1996                               |
| <i>V. dissecta</i> (Benth.) N.T.Burb.<br><i>var. hirta</i> N.T.Burb. | 9 |    | Turner 1970, as <i>V. triloba</i> & <i>Vittadinia</i> sp. |
| <i>V. gracilis</i> (Hook.f.) N.T.Burb.                               | 9 |    | Watanabe <i>et al.</i> 1996                               |
| <i>V. muelleri</i> N.T.Burb.   | 9 |    | Watanabe <i>et al.</i> 1996                               |
| <i>V. pterochaeta</i> (F.Muell.<br>ex benth.) J.M.Black              | 9 |    | Watanabe <i>et al.</i> 1996                               |
| <i>V. pustulata</i> N.T.Burb.  | 9 |    | Watanabe <i>et al.</i> 1996                               |
| <i>V. sulcata</i> N.T.Burb.  | 9 |    | Watanabe <i>et al.</i> 1996                               |

## ERIGERON

Given (1973) and Given & Gray (1986) noted that Australian species of *Erigeron* should be excluded from that genus. Nesom (1994a,b) has subsequently referred most Australian species to three genera, i.e. *Iotasperma* Nesom, *Lagenithrix* Nesom and *Pappochroma* Labill. (syn. *Lagenopappus* Nesom), but has left the placement of *E. conyzoides* F.Muell. unresolved. Cladistic studies (by PSS) in Australian Astereae do not wholly support Nesom's treatment and will be the subject of a future paper.

The alpine *E. pappochroma* complex (*Lagenithrix* & *Pappochroma* sensu Nesom) seemingly has affinities with the *Olearia-Celmisia* complex (Given & Gray 1986) and could be expected to have a base of  $x = 9$ .

The count of  $2n = 36$  for the introduced *E. karvinskianus* is consistent with those obtained by Montgomery & Yang (1960, as *E. mucronatus* DC.).

## LAGENIFERA &amp; SOLENOGYNE

Whether or not *Solenogyne* is deserving of generic rank or should be relegated to synonymy under *Lagenifera* has been the subject of debate for some years (Drury 1974, Adams 1979). They are here maintained as separate genera pending further work. All species are perennials and tend to be found in mesic conditions. Our chromosome number determinations, i.e.  $n = 9$  and  $2n = 18$ , are consistent with the observations of previous workers (Smith-White *et al.* 1970, Turner 1970, Adams 1979).

## KIPPSTIA &amp; MINURIA

Eleven species are currently placed in this genus (Lander & Barry 1980b, Lander 1987b, Short 1991) although one, *M. macrorhiza*, should possibly be reinstated as *Eurybiopsis* (see below). All but one, *M. annua* (Tate) J.M.Black, are apparently perennial herbs or small shrubs although several, including *M. multiseta* P.S.Short may be short lived. We have confirmed earlier reports of  $n = 9$  for *M. cunninghamii* and *M. leptophylla* (Turner 1970, Short 1986) and here record the tetraploid condition for *M. integrerrima*. The presence of B chromosomes is here reported for the first time in this genus.

*Kippstia*, a monotypic genus, at one time relegated to synonymy under *Minuria* but reinstated by Lander & Barry (1980a), is seemingly very closely related to the latter genus. *K. suaedifolia* is a small shrub with  $n = 9$ .

cultivation of specimens and the preparation of material for chromosome number determinations the procedures of Smith-White *et al.* (1970) and Watanabe *et al.* (1975) were followed. Herbarium vouchers have been gathered for all but two species (*Brachyscome petrophila* and *Olearia ramulosa*) and a complete set of vouchers is housed at MEL. An incomplete set will be deposited in TI and voucher specimens of non-Victorian populations will be deposited in the major government herbarium (AD, BRI, NSW, PERTH) of the State from which collections were gathered.

### Results

Chromosome number determinations from 200 populations attributed to 99 species or infraspecific taxa of ten genera are presented in Table I. They include the first substantiated reports for *Brachyscome glandulosa* ( $2n = 36$ ), *B. muelleroides* ( $n = 3$ ), *B. obovata* ( $2n = 18$ ), *B. petrophila* ( $2n = 18$ ), *B. tatei* ( $n = 9$ ), *Erodiocephalum elderi* ( $2n = 16$ ), *Minuria gardneri* ( $n = 9$ ), *M. integriflora* ( $n = 18$ ), *Olearia astrotricha* ( $n = 9$ ), *O. ferresii* ( $n = 9$ ), *O. imbricata* ( $n = 9$ ), *O. plucheacea* ( $n = 9$ ), *O. ramulosa* ( $n = 9$ ), *O. xerophila* ( $n = 9$ ), *Vittadinia cuneata* ( $n = 9$ ), *V. gracilis* ( $n = 9$ ), *V. muelleri* ( $n = 9$ ), *V. pterochaeta* ( $n = 9$ ), *V. pustulata* ( $n = 9$ ) and *V. sulcata* ( $n = 9$ ). Some reports for apparently distinct, but undescribed, taxa such as *Calotis* aff. *lappulacea* ( $n = 8$ ) and *B. tenuiscapa* aff. var. *tenuiscapa* ( $2n = 28$ ) are also presented, as are new determinations for some previously examined taxa, e.g. *Brachyscome microcarpa* ( $n = 6$ ,  $2n = 12$ ; cf.  $n = 10$  reported by Smith-White *et al.* 1970). Polyploidy is reported for the first time in *Brachyscome dissectifolia*, *B. cardiocarpa*, *Calotis anthemoides* and in the *Olearia phlogopappa* complex and B chromosomes are newly reported in *Brachyscome ciliocarpa*, *B. diversifolia*, *B. iberidifolia*, *B. trachycarpa*, *Minuria leptophylla* and *Olearia pimelioides*.

The meiotic or mitotic chromosomes of some species are displayed in Fig. 1.

The new data is incorporated (by reference to Watanabe *et al.* 1996) in Table II which summarises known chromosome numbers for taxa of Australian Astereae.

### Discussion

As previously pointed out (Watanabe & Short 1992) there are many taxonomic problems within *Brachyscome*. There are unresolved species complexes, e.g. *B. nova-anglica* and its allies and the *B. ciliaris* complex, and generic delimitation is also a problem. The same is true for many of the other genera and the placement of *Isoetopsis* in the Astereae is debatable and perhaps not tenable (Bremer & Anderberg in Bremer 1987, Bruhl & Quinn 1990, Bremer & Humphries 1993). Although there are taxonomic problems it is none the less possible to generally review the state of knowledge of chromosome numbers in the Australian Astereae. An absence of chromosome numbers for some genera, i.e. *Achnophora*, *Camptacra*, *Dichrocephala*, *Dichromochlamys*, *Dimorphocoma* and *Elachanthus* is unfortunate, although between them these genera contain only 12 Australian species.

TABLE 1. NEW CHROMOSOME NUMBER DETERMINATIONS FOR AUSTRALIAN ASTEREAE

| Species & locality   | n | 2n  |
|--|---|-----|
| <i>Brachyscome</i> Cass.<br><i>B. aculeata</i> (Labill.) Less.<br>5.5 km SE of Sawyers Hill,<br>Kosciusko N.P., N.S.W.,<br>3 Feb. 1993, Short 3982 |   | 9II |
| Sawpit Creek, Kosciusko N.P.,<br>N.S.W., 4 Feb. 1993, Short 3996   |   | 9II |

### OLEARIA

*Olearia* consists of about 180 species and occurs in Australia, New Guinea and New Zealand. About 130 species occur in Australia (Lander 1992) and the genus is currently under revision. All species are shrubs and the genus is widespread in alpine, temperate and arid regions of Australia. Few species have been cytologically examined but our data are consistent with the finding for New Zealand species that the base number is  $x = 9$ . However, most Australian species are diploids, a contrast to the situation in New Zealand where only 12-, 14-, 32- and 48-ploid levels have been recorded (Beuzenberg & Hair 1984). For the first time B chromosomes are also reported for this genus, having been observed by us in *O. pimelioides*.

### VITTADINIA, CAMPTACRA & EURYBIOPSIS

Burbridge (1982) revised *Vittadinia* and recognised 29 species, most of which are confined to Australia. She segregated two further genera, describing ditypic *Camptacra* (Lander 1987b) and reinstating monotypic *Eurybiopsis*. Lander (1987a) has subsequently reduced *Eurybiopsis* to synonymy under *Minuria* although more recent work by Wiggins (1990) suggests that the former genus should be maintained.

Chromosome counts are lacking for *Camptacra* and *Eurybiopsis* and until now for all but one species of *Vittadinia*. Chromosome numbers are now known for seven species (Table II). Of these, three are, or possibly are, annuals. All seven species have  $n = 9$ . One of these is more or less confined to temperate areas, the others extend from temperate to arid areas.

### Base numbers and comparisons

With the exception of *Isoetopsis*, which possibly should be placed in the Gnaphalieae (Bremer & Humphries 1993), ditypic *Erodiophyllum*, monotypic *Ceratogyne*, and the essentially arid-zone genus *Calotis*, all Australian genera for which data are available have one or more species with  $n = 9$  or a base number that is a multiple of 9. Therefore, it seems reasonable to conclude that  $x = 9$  is the base number for the Australian Astereae, a conclusion that is in keeping with previous findings for the tribe Astereae (Raven *et al.* 1960, Solbrig *et al.* 1964, 1969). Following a survey of mainly Northern Hemisphere genera (38 out of 53) and some Southern Hemisphere genera (22 out of 65) Solbrig *et al.* (1964) also noted that many species with low chromosome numbers ( $n = 4$  & 5) belong to mainly annual genera that are concentrated in southwestern North America and that low numbers were probably correlated with dry habitats. As noted above, the same correlations are apparent in annual species of *Brachyscome* 'subgenus *Ebrachyscome*', *Calotis* and *Ceratogyne*. Chromosomal variation in Australian Astereae is clearly similar to that found in North America. In fact, although fewer chromosome number determinations were available to him, Turner (1970) previously suggested this to be the case, not just for the tribe Astereae, but the entire family.

### Acknowledgement

Research was supported by grants (nos. 04041071 and 07041140) in Aid for Scientific Research and Field Research from the Ministry of Education, Science and Culture, Japan.

### References

- Adams, L.G. (1979). A review of the genus *Solenogyne* (Asteraceae) in Australia and New Zealand. *Brunonia* 2: 43-65.
- Beuzenberg, E.J. & Hair, J.B. (1984). Contributions to a chromosome atlas of the New Zealand flora - 27 Compositae. *New Zealand Journal of Botany* 22: 353-356.
- Bremer, K. (1987). Tribal interrelationships of the Asteraceae. *Cladistics* 3: 210-253.
- Bremer, K. & Humphries, C.J. (1993). Generic monograph of the Asteraceae-Anthemideae. *Bulletin of the Natural History Museum London (Botany)* 23: 71-177.
- Bruhl, J.J. & Quinn, C.J. (1990). Cypsela anatomy in the 'Cotuleae' (Asteraceae-Anthemideae). *Botanical Journal of the Linnean Society* 102: 37-59.

- Burbidge, N.T. (1982). A revision of *Vittadinia* A. Rich. (Compositae) together with reinstatement of *Eurybiopsis* DC. and description of a new genus, *Camptacra*. *Brunonia* 5: 1-72.
- Carter, C.R. (1978a). The cytology of *Brachycome*. II. The subgenus *Metabrachycome*: a general survey. *Australian Journal of Botany* 26: 699-706.
- Carter, C.R. (1978b). The cytology of *Brachycome*. 8. The inheritance, frequency and distribution of B chromosomes in *B. dichromosomatica* ( $n = 2$ ), formerly included in *B. lineariloba*. *Chromosoma (Berlin)* 67: 109-121.
- Carter, C.R. (1978c). Taxonomy of the *Brachycome lineariloba* complex (Asteraceae). *Telopea* 1: 387-393.
- Carter, C.R. & Smith-White, S. (1972). The cytology of *Brachycome lineariloba*. 3. Accessory chromosomes. *Chromosoma (Berlin)* 39: 361-379.
- Chouksanova, N.A., Sveshnikova, L.I. & Alrandrova, T.U. (1968). Data on karyology of the family Compositae Gisecke. *Cytologija* 10: 198-206.
- Davis, G.L. (1948). Revision of the genus *Brachycome* Cass. Part. I. Australian species. *Proceedings of the Linnean Society of New South Wales* 73: 142-241.
- Davis, G.L. (1949). Revision of the genus *Brachycome* Cass. III. Description of three new Australian species, and some new locality records. *Proceedings of the Linnean Society of New South Wales* 74: 145-152.
- Davis, G.L. (1955). Supplementary notes on the genus *Brachycome* Cass. *Proceedings of the Linnean Society of New South Wales* 79: 203-210.
- Davis, G.L. (1959). Two new Australian species of *Brachycome* Cass. *Muelleria* 1: 111-113.
- De Jong, D.C.D. (1963). Documented chromosome numbers in plants. *Madrono* 17: 116-117.
- Drury, D.G. (1974). A broadly based taxonomy of *Lagenifera* sect. *Lagenifera* and *Solenogyne* (Compositae-Astereae) with an account of their species in New Zealand. *New Zealand Journal of Botany* 12: 365-395.
- Given, D.R. (1973). *Damnamenia* gen. nov. A new subantarctic genus allied to *Celmisia* Cass. (Astereae-Compositae). *New Zealand Journal of Botany* 11: 785-796.
- Given, D.R. & Gray, M. (1986). *Celmisia* (Compositae-Astereae) in Australia and New Zealand. In Barlow, B.A. (ed.), *Flora and fauna of alpine Australia*. (CSIRO: Melbourne), pp. 451-470.
- Gupta, P.K. (1969). Cytological investigations in some Indian Compositae. *Cytologia* 34: 429-438.
- Gupta, R.C. & Gill, B.S. (1989). Cytopalynology of north and central Indian Compositae. *Journal of Cytology and Genetics* 24: 96-105.
- Hair, J.B. (1980). Contributions to a chromosome atlas of the New Zealand flora - 20 *Celmisia* and *Damnamenia* (Compositae). *New Zealand Journal of Botany* 18: 553-558.
- Hnatiuk, R.J. (1990). *Census of Australian vascular plants*. (AGPS: Canberra).
- Keighery, G.J. (1978). Chromosome numbers in Western Australian plants 1. *Journal of the Royal Society of Western Australia* 60: 105-106.
- Kyhos, D.W., Carter, C.R. and Smith-White, S. (1977). The cytology of *Brachycome lineariloba*. 7. Meiosis in natural hybrids and race relationships. *Chromosoma (Berlin)* 65: 81-101.
- Lander, N.S. (1987a). New combinations in *Camptacra* N.Burb. (Asteraceae, Astereae). *Nuytsia* 6: 61.
- Lander, N.S. (1987b). New combinations in *Minuria* DC. (Asteraceae, Astereae). *Nuytsia* 6: 63-66.
- Lander, N.S. (1989). Taxonomy of *Olearia stuartii* (Asteraceae: Astereae) and allied species. *Nuytsia* 7: 25-36.
- Lander, N.S. (1990). New species of *Olearia* (Asteraceae: Astereae) endemic to Western Australia. *Nuytsia* 7: 141-159.
- Lander, N.S. (1992). *Olearia*. In Harden, G.J. (ed.), *Flora of New South Wales*. (NSW University Press: Kensington.), Vol. 3, pp.185-197.
- Lander, N.S. & Barry, R. (1980a). Reinstatement of the genus *Kippistia* F.Muell. (Asteraceae, Astereae). *Nuytsia* 3: 215-219.
- Lander, N.S. & Barry, R. (1980b). A review of the genus *Minuria* DC. (Asteraceae, Astereae). *Nuytsia* 3: 221-237.
- Lander, N.S. & Walsh, N.G. (1989). *Olearia astroloba* (Asteraceae: Astereae), a new species endemic to Victoria. *Muelleria* 7: 123-125.
- Montgomery, F.H. & Yang, S. (1960). Cytological studies in the genus *Erigeron*. *Canadian Journal of Botany* 38: 381-386.
- Nagl, W. & Pfeifer, M. (1988). Karyological stability and microevolution of a cell suspension culture of *Brachycome dichromosomatica* (Asteraceae). *Plant Systematics and Evolution* 158: 133-139.
- Nesom, G.L. (1994a). Taxonomic dispersal of Australian *Erigeron* (Asteraceae: Astereae). *Phytologia* 76: 143-159.
- Nesom, G.L. (1994b). *Pappochroma* Rafin. is the correct generic name for *Erigeron pappochroma* Labill. *Phytologia* 76: 426
- Pai, O.J. (1964). Some chromosome numbers of cultivated plants. *Journal of Heredity* 55: 186-192.
- Raven, P.H., Solbrig, O.T., Kyhos, D.W. & Snow, R. (1960). Chromosome numbers in Compositae 1. Astereae. *American Journal of Botany* 47: 124-132.
- Short, P.S. (1986). Chromosome number reports in Australian Asteraceae. *Taxon* 35: 610.
- Short, P.S. (1988). Two new species of *Brachyscome* Cass. (Compositae: Astereae), with a note on the orthography of the generic name. *Muelleria* 6: 389-398.
- Short, P.S. (1991). A new species of *Minuria* DC. (Asteraceae: Astereae). *Muelleria* 7: 361-367.
- Short, P.S. & Watanabe, K. (1993). Two new species of *Brachyscome* Cass. (Asteraceae: Astereae) from eastern Australia. *Australian Systematic Botany* 6: 335-342.
- Smith-White, S. (1968). *Brachycome lineariloba*: a species for experimental cytogenetics. *Chromosoma (Berlin)* 23: 359-364.
- Smith-White, S. & Carter, C.R. (1970). The cytology of *Brachycome lineariloba*. 2. The chromosome species and their relationships. *Chromosoma (Berlin)* 30: 129-153.

- Smith-White, S. & Carter, C.R. (1981). The maintenance of B chromosomes in *Brachycome dichromosomatica*. In Atchley, W.R. & Woodruff, D. (eds), *Evolution and Speciation. Essays in honour of M. J. D. White*. (Cambridge University Press: Cambridge). pp. 335-355.
- Smith-White, S., Carter, C.R. and Stace, H.M. (1970). The cytology of *Brachycome*. I. The subgenus *Eubrachycome*: a general survey. *Australian Journal of Botany* 18: 99-125.
- Solbrig, O.T., Anderson, L.C., Kyhos, D.W., Raven, P.H. & Rudenberg, L. (1964). Chromosome number in the Compositae. Asteraceae. *American Journal of Botany* 51: 513-519.
- Solbrig, O.T., Anderson, L.C., Kyhos, D.W. & Raven, P.H. (1969). Chromosome number in the Compositae VII. *American Journal of Botany* 56: 348-353.
- Stace, H.M. (1981). Biosystematics of the *Brachyscome aculeata* (Labill.) Less. sensu G. L. Davis species complex (Compositae: Astereae). *Australian Journal of Botany* 29: 425-440.
- Stace, H.M. (1978). Cytoevolution in the genus *Calotis* R.Br. (Compositae: Astereae). *Australian Journal of Botany* 26: 287-307.
- Stace, H.M. (1982). *Calotis* (Compositae), a Pliocene arid zone genus? In Barker, W. R. & Greenslade P. J. M. (eds), *Evolution of the flora and fauna of arid Australia*. (Peacock Publins: Frewville). pp. 357-367.
- Turner, B.L. (1970). Chromosome numbers in the Compositae. xii. Australian species. *American Journal of Botany* 57: 382-389.
- Watanabe, K., Carter, C.R. and Smith-White, S. (1975). The cytology of *Brachycome lineariloba*. 5. Chromosome relationships and phylogeny of the race A cytodesmes ( $n = 2$ ). *Chromosoma (Berlin)* 52: 383-397.
- Watanabe, K., Carter, C.R. and Smith-White, S. (1976). The cytology of *Brachycome lineariloba*. 6. Asynchronous chromosome condensation and meiotic behaviour in *B. lineariloba* A ( $n = 2$ ) x *B. campylocarpa* A ( $n = 4$ ). *Chromosoma (Berlin)* 57: 319-331.
- Watanabe, K., Carter, C.R. and Smith-White, S. (1985). The cytology of *Brachycome lineariloba*. 9. Chromosomal heterogeneity in natural populations of cytodeme C ( $2n = 16$ ). *Canadian Journal of Genetics and Cytology* 27: 410-420.
- Watanabe, K. & Short, P.S. (1992). Chromosome number determinations in *Brachyscome* Cass. (Asteraceae: Astereae) with comments on species delimitation, relationships and cytogeography. *Muelleria* 7: 457-471.
- Watanabe, K., Short, P.S., Kosuge, K. & Smith-White, S. (1991). The cytology of *Brachyscome* Cass. (Asteraceae: Astereae). 11. Hybridization between *B. goniocarpa* Sond. & F. Muell. ex Sond. ( $n = 4$ ) and *Brachyscome dichromosomatica* C. R. Carter ( $n = 2$ ). *Australian Journal Botany* 39: 475-485.
- Watanabe, K. and Smith-White, S. (1987). Phylogenetic and evolutionary relationships of *Brachycome lineariloba* 6 (Compositae). *Plant Systematics and Evolution*. 157: 121-141.
- Wiggins, D.J. (1990). A cladistic analysis of the generic concepts of *Vittadinia* A.Richard (Asteraceae: Astereae). Unpublished B.Sc.(Hons.) thesis, University of New South Wales.

Revised paper received 20 April 1995; last update 24 November 1995.

TABLE 1. CONTINUED

| Species & locality  | n          | 2n |
|---|------------|----|
| 6 km from Cathcart toward Rocky Hall, 9 Feb. 1993, N.S.W., <i>Short 4010</i>  | 9II        |    |
| Bundara R. crossing on Omeo to Tallangatta road, Vict., 9 Feb. 1993, <i>Short 4018</i>                                    | 18II + 2Bs |    |
| <i>B. sp. aff. angustifolia</i> A. Cunn. ex DC.<br>Mt Kaputar N.P., N.S.W., 28 Jan. 1993,<br><i>Short 3944</i>            | 5II        | 10 |
| Stewart Forest, Barrington Tops State Forest, N.S.W., 1 Feb. 1993, <i>Short 3979</i>                                      | 5II        | 10 |
| Polblue Picnic Site, Barrington Tops State Forest, N.S.W., 1 Feb. 1993, <i>Short 3981</i>                                 |            | 10 |
| <i>B. basaltica</i> F.Muell.<br>var. <i>basaltica</i><br>4 km NE of Millmerran, Qld,<br>29 Sept. 1992, <i>Watanabe 25</i> | 8II        | 16 |
| var. <i>gracilis</i> Benth.<br>14 km N of Booligal, N.S.W.,<br>6 Sept. 1992, <i>Watanabe 76</i>                           |            | 12 |
| Barmah State Park, Vict.,<br>23 Sept. 1992, <i>Watanabe 1</i>   | 6II        |    |
| 20 km N of Apsley, Vict.,<br>25 Oct. 1993, <i>Watanabe 222</i>  | 6II        |    |
| <i>B. cardiocarpa</i> F.Muell. ex Benth.<br>Digby to Strathdownie road, Vict.,<br>3 Sept. 1990, <i>Short 3919</i>         |            | 36 |
| 19 km NW of Apsley, Vict.,<br>12 Oct. 1995, <i>Watanabe 306</i>   |            | 36 |
| <i>B. cheilocarpa</i> F.Muell.<br>94 km N of Galena Bridge on NW Coastal Hwy, W.A., 6 Oct. 1993,<br><i>Short 4083</i>     |            | 18 |
| 1 km W of NW Coastal Hwy along road to Blowholes, W.A., 7 Oct. 1993,<br><i>Short 4097</i>                                 |            | 18 |
| <i>B. aff. cheilocarpa</i> (A)<br>Pt Quobba, W.A., 7 Oct. 1993,<br><i>Short 4091</i>                                      |            | 18 |

TABLE 1. CONTINUED

| Species & locality  | n   | 2n |
|---|-----|----|
| <b>B. aff. cheilocarpa (B)</b>  |     |    |
| Cleary to Paynes Find road, W.A.,<br>17 Sept. 1990, <i>Short 3823</i>                                   | 9II |    |
| Yalgo, W.A., 10 Oct. 1993,<br><i>Short 4108</i>   | 9II |    |
| <b>B. ciliaris</b> (Labill.) Less. complex  |     |    |
| Cape Riche, W.A., 14 Oct. 1993,<br><i>Short 4126</i>  | 9II |    |
| Tooreburrup Scenic Lookout, W.A.,<br>14 Oct. 1993, <i>Watanabe 173</i>                                  | 9II |    |
| 35 km E of Jerramungup, Fitzgerald<br>River Crossing, W.A., 15 Oct. 1993,<br><i>Watanabe 174</i>        | 9II |    |
| 6 km W of Coolgardie, W.A.,<br>2 Oct. 1993, <i>Watanabe 107</i>   | 27  |    |
| 4 km NE of Millmerran, Qld,<br>29 Sept. 1992, <i>Watanabe 27</i>  | 27  |    |
| 52 km SW of Louth, N.S.W.,<br>2 Oct. 1992, <i>Watanabe 47</i>   | 27  |    |
| 13 km N of Carnamah, W.A.,<br>5 Oct. 1993, <i>Short 4075</i>  | 36  |    |
| 33 km from Blowholes along road<br>to Carnarvon, W.A., 7 Oct. 1993,<br><i>Short 4094</i>                | 36  |    |
| 5 km E of Pindar, W.A., 9 Oct. 1993,<br><i>Short 4105</i>   | 36  |    |
| 27 km from Great Northern Hwy along<br>road to Warriedar Hmsd, W.A.,<br>10 Oct. 1993, <i>Short 4116</i> | 36  |    |
| 63 km W of Balladonia, W.A.,<br>1 Oct. 1993, <i>Watanabe 103</i>  | 36  |    |
| 6 km W of Coolgardie, W.A.,<br>2 Oct. 1993, <i>Watanabe 106</i>   | 36  |    |
| 12.5 km W of Coolgardie, W.A.,<br>2 Oct. 1993, <i>Watanabe 110</i>                                      | 36  |    |
| Bonnie Rock, W.A., 11 Oct. 1993,<br><i>Watanabe 158</i>   | 36  |    |
| E edge of Lake Wallambin, W.A.,<br>11 Oct. 1993, <i>Watanabe 163</i>                                    | 36  |    |

TABLE 1. CONTINUED

| Species & locality   | n                  | 2n       |
|--|--------------------|----------|
| 25 km NW of Pt Augusta, S.A.,<br>29 Sept. 1993, <i>Watanabe 101</i>  | 36                 |          |
| 61 km N of Hawker, S.A.,<br>20 Oct. 1993, <i>Watanabe 191</i>  | 36                 |          |
| 10 km NNE of Ashford, N.S.W.,<br>26 Sept. 1992, <i>Watanabe 15</i>   | 36                 |          |
| <i>B. ciliocarpa</i> W. Fitzg.<br>'The Loop', Murchison River,<br>Kalbarri N.P., W.A., 6 Oct. 1993,<br><i>Short 4080</i> |                    | 18 + 2Bs |
| 40 km from Yalgoo, W.A.,<br>9 Oct. 1993, <i>Short 4107</i>   | 9II                |          |
| 30 km from Paynes Find along road<br>to Cleary, W.A., 10 Oct. 1993,<br><i>Short 4119</i>                                 |                    | 18       |
| <i>B. sp. aff. ciliocarpa</i><br>41 km E of Quilpie, Qld,<br>17 Aug. 1989, <i>Short 3607</i>                             | 9II                |          |
| <i>B. curvicarpa</i> G.L.R.Davis<br>59.5 km SW of Bourke, N.S.W.,<br>2 Oct. 1992, <i>Watanabe 41</i>                     | 4II + BI           |          |
| 0.9 km SW of Louth, N.S.W.,<br>2 Oct. 1992, <i>Watanabe 43</i>   | 4II + BI           |          |
| <i>B. debilis</i> Sond.<br>Mt Arapiles, Vict., 25 Oct. 1993,<br><i>Watanabe 217</i>                                      | 3II                | 6        |
| <i>B. decipiens</i> Hook.f.<br>Sawyers Hill, Kosciusko N.P.<br>N.S.W., 3 Feb. 1993, <i>Short 3986</i>                    | 9II                | 18       |
| <i>B. dentata</i> Gaudich.<br>Dalby, N.S.W., 30 Sept. 1992,<br><i>Watanabe 29</i>  | 4II or<br>4II + BI |          |
| 44 km SW of Louth, N.S.W.,<br>2 Oct. 1992, <i>Watanabe 46</i>  | 4II                |          |
| 24 km NE of Wirrealpa, N.S.W.,<br>3 Oct. 1992, <i>Watanabe 55</i>  | 4II                |          |
| 80 km SE of Broken Hill, N.S.W.,<br>6 Oct. 1992, <i>Watanabe 70</i>  | 4II + BI           |          |
| 48 km SE of Martins Well, N.S.W., 8II<br>3 Oct. 1992, <i>Watanabe 53</i>   |                    |          |

TABLE 1. CONTINUED

| Species & locality  | n    | 2n        |
|---|------|-----------|
| 27 km S of Tilpa, N.S.W.,<br>2 Oct. 1992, Watanabe 52   | 12II |           |
| c. 45 km SE of Broken Hill, N.S.W.,<br>6 Oct. 1992, Watanabe 67   | 12II |           |
| 33 km SW of Broken Hill, N.S.W.,<br>22 Oct. 1993, Watanabe 201  | 12II |           |
| c. 43 km SE of Broken Hill, N.S.W.,<br>22 Oct. 1993, Watanabe 203   | 12II |           |
| 20 km E of Elmore, Vict.,<br>7 Oct. 1992, Watanabe 77   | 12II |           |
| <b>B. dissectifolia</b> G.L.R.Davis<br>3.5 km NW of Backwater, N.S.W.,<br>30 Jan. 1993, Short 3973                                | 12II | 24        |
| <b>B. diversifolia</b> (Hook.) Fischer & Meyer<br>17 km SE of Halls Gap, Vict.,<br>27 Sept. 1993, Watanabe 95                     | 12II |           |
| Polblue Creek, Barrington Tops<br>State Forests, N.S.W., 1 Feb.<br>1993, Short 3981   | 18II | 36 + 0-1B |
| <b>B. exilis</b> Sond.<br>Mullett Lake Nature Reserve, near<br>Esperance, W.A., 16 Oct. 1993,<br>Watanabe 178                     | 9II  | 18        |
| <b>B. formosa</b> P.S.Short<br>1.6 km SE of Coonabarabran, N.S.W.,<br>24 Sept. 1992, Watanabe 6                                   | 9II  |           |
| <b>B. glandulosa</b> (Steetz) Benth.<br>Boyagin Rock, W.A., 12 Oct. 1993,<br>Watanabe 165   |      | 36        |
| <b>B. gracilis</b> G.L.R. Davis<br>7.5 km towards Ashford from Bonshaw<br>to Glen Innes road, N.S.W., 29 Jan.<br>1993, Short 3962 |      | 18        |
| <b>B. graminea</b> (Labill.) F.Muell.<br>Sawpit Creek, Kosciusko N.P.<br>N.S.W., 6 Feb. 1993, Short 4008                          | 9II  |           |
| Green Cape Lighthouse, N.S.W.<br>7 Feb. 1993, Short 4015  | 9II  | 18        |
| <b>B. halophila</b> P.S.Short<br>13 km N of Carnamah, W.A.,<br>5 Oct. 1993, Short 4072  | 9II  |           |

TABLE 1. CONTINUED

| Species & locality  | n                  | 2n  |
|---|--------------------|-----|
| 11 km W of Pindar, W.A.,<br>9 Oct. 1993, <i>Short 4101</i>  | 9II                |     |
| <i>B. iberidifolia</i> Benth. complex<br>Moora, W.A., 4 Oct. 1993,<br><i>Short 4065</i>                       | 9II                |     |
| 13 km N of Carnamah, W.A.,<br>5 Oct. 1993, <i>Short 4074</i>  |                    | 18  |
| 2.5 km S of Binnu, W.A.,<br>5 Oct. 1993, <i>Short 4077</i>  | 9II                |     |
| 21 km from Carnarvon along road<br>to Gascoyne Junction, W.A.,<br>7 Oct. 1993, <i>Short 4087</i>              |                    | 18  |
| 29 km N of Galena Bridge, W.A.,<br>8 Oct. 1993, <i>Short 4100</i>   | 9II                |     |
| Cosy Corner, W.A., 13 Oct. 1993,<br><i>Short 4124</i>   | 9II                |     |
| Mt Chudalup, W.A., 13 Oct. 1993,<br><i>Short 4125</i>   | 9II or<br>9II + BI |     |
| 9 km N of Nallan Homestead<br>turn-off along Great Northern Highway,<br>W.A., 26 Aug. 1995, <i>Short 4227</i> | 9II                |     |
| Cape Leeuwin, W.A., 3 Nov. 1995,<br><i>Short 4550</i>   | 9II                |     |
| 74 km WSW of Coolgardie, W.A.,<br>2 Oct. 1993, <i>Watanabe 111</i>  | 9II                |     |
| Yellowdine, W.A., 2 Oct. 1993,<br><i>Watanabe 113</i>   |                    | 9II |
| c. 5 km NW of Bonnie Rock, W.A.,<br>11 Oct. 1993, <i>Watanabe 161</i>   | 9II                |     |
| Lake Wallanbin, W.A., 11 Oct. 1993,<br><i>Watanabe 164</i>  | 9II                |     |
| 35 km E of Jerramungup, Fitzgerald<br>River crossing, W.A., 15 Oct. 1993,<br><i>Watanabe 175</i>              | 9II                |     |
| <i>B. latisquamea</i> F. Muell.<br>Pt Quobba, W.A., 7 Oct. 1993,<br><i>Short 4090</i>                         |                    | 18  |

TABLE 1. CONTINUED

| Species & locality  | n    | 2n         |
|---|------|------------|
| 33 km from Blowholes along road<br>to Carnarvon, W.A., 7 Oct. 1993,<br><i>Short 4093</i>                            |      | 18         |
| <i>B. lineariloba</i> (DC.) Druce<br>S of Border Village on S.A./W.A.<br>border, 29 Sept. 1993, <i>Watanabe 102</i> |      | 12         |
| 12.5 km W of Kimba, S.A.,<br>19 Oct. 1993, <i>Watanabe 187</i>  |      | 12         |
| 6 km W of Coolgardie, W.A.,<br>2 Oct. 1993, <i>Watanabe 105</i>   |      | 16         |
| 7 km S of Bimbijy Hmsd turn-off<br>along Paynes Find to Cleary Road,<br>W.A., 10 Oct. 1993, <i>Short 4122</i>       |      | 16         |
| 33 km from Broken Hill on Menindee<br>road, N.S.W., 22 Oct. 1993,<br><i>Watanabe 200</i>                            |      | 16         |
| <i>B. melanocarpa</i> Sond. & F.Muell<br>1.4 km W of Menindee, N.S.W.,<br>22 Oct. 1993, <i>Watanabe 209</i>         | 12II |            |
| <i>B. microcarpa</i> F.Muell.<br>Girraween N.P., Qld,<br>29 Jan. 1992, <i>Watanabe 80</i>                           | 6II  | 12         |
| 6 km S of Boonoo Boonoo, N.S.W.,<br>30 Jan. 1993, <i>Short 3970</i>   | 6II  |            |
| <i>B. muelleroides</i> G.L.R.Davis<br>Ulupna Island, Vict.,<br>23 Sept. 1992, <i>Watanabe 3</i>                     | 3II  |            |
| <i>B. aff. multicaulis</i> F.Muell.<br>8 km W of Kiandra, N.S.W.,<br>3 Feb. 1993, <i>Short 3988</i>                 | 9II  | 18 + 0-2Bs |
| <i>B. multifida</i> DC. complex<br>3 km S of Glen Alpin, Qld,<br>30 Jan. 1993, <i>Short 3965</i>                    | 7II  |            |
| Warrumbungles N.P., N.S.W.,<br>25 Sept. 1992, <i>Watanabe 7</i>   | 7II  |            |
| Mt Kaputar N.P., N.S.W.,<br>25 Sept. 1992, <i>Watanabe 11</i>   | 7II  | 14         |
| Nr Clover Flat along Big River<br>Fire Track, Vict., 9 Feb. 1993,<br><i>Short 4021</i>                              | 9II  | 18         |