

Ifz
klosterneuburg

MITTEILUNGEN KLOSTERNEUBURG

64(2014) 1

www.mitt-klosterneuburg.com

Bimonthly ISSN: 0007-5922

Short communication

First record of *Alosa fallax* (Lacépède, 1803) (Teleostei: Clupeiformes:Clupeidae) in the Azores Archipelago, Portugal (NE Atlantic).

João P. Barreiros ⁽¹⁾*, Rui P. Vieira ⁽²⁾ and Leonardo Machado ⁽³⁾

- (1) Azorean Biodiversity Group (CITA-A) and Platform for Enhancing Ecological Research & Sustainability (PEERS), Universidade dos Açores, Dep. Ciências Agrárias, 9700-042 Angra do Heroísmo, Portugal.
- (2) Departamento de Biologia & CESAM, Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal. r.vieira87@gmail.com
- (3) Universidade Federal do Espírito Santo, Dep. Ciências Agrárias e Biológicas-DCAB/CEUNES. Rodovia BR 101 Norte, Km 60, Bairro Litorâneo. 29932-540, São Mateus, ES, BRASIL. dusky.grouper@gmail.com

* Corresponding author. E-mail: joao.pedro@uac.pt

Key words: Clupeidae; Azorean ichthyofauna; anadromous species; vulnerable.

ABSTRACT

This note reports the first record of *Alosa fallax* (Lacépède, 1803) in the Azores Archipelago (NE Atlantic, Portugal). Being an anadromous species, this occurrence is quite unexpected since there are no suitable breeding habitats for this species in these volcanic islands, isolated and river less. Although *A. fallax* is known to migrate offshore, it has never been previously reported in oceanic environments.

We discuss this occurrence and theorize on the possibility of the species' being found in Madeira, another Macaronesian Archipelago, albeit much older and closer to European coasts and thus more suitable to be crossed by *A. fallax* during their migration routes in pelagic environments.

With this record, the number of Clupeid fishes for Azorean waters increases to two, the other being the well known and resident *Sardina pilchardus*.

INTRODUCTION

The twaite shad, *Alosa fallax* (Lacépède, 1803), is a well known Clupeidae widely distributed in the Northeast Atlantic, from the southern coasts of Scandinavia to off Morocco, including the Mediterranean and Black Seas (Whitehead 1985; Froese & Pauly 2012).

A. fallax populations are being negatively affected by several human activities, such as river and coastal pollution, dam constructions and overfishing (Collares-Pereira et al. 1999). Consequently it is a matter for conservation concerns and classified on the Portuguese Vertebrate Red List as “Vulnerable” (Costa et al. 2001; Cabral et al. 2006), although globally considered of “Least Concern” status for IUCN (Freyhof & Kottelat, 2008). It is listed in Appendix III of the Bern Convention (2002) and in Annexes II and V of the EU Habitats Directive (2007).

Alosa fallax is a marine pelagic anadromous species performing short distance migrations up to rivers for spawning between May and June (Pina et al. 2003; Alexandrino et al. 2006). Successively, larvae migrate downstream and the 1 year-olds spend several months in estuaries before the journeying to the sea (Jolly et al. 2012).

A. fallax feeds on small fish, euphausiids, mysids, isopods, copepods and insects (Whitehead 1984, 1985; Assis et al. 1992; Oesmann & Thiel 2001).

The occurrence of *A. fallax* has never been reported for the Azores archipelago (Santos et al. 1997; Porteiro et al. 2010).

MATERIAL AND METHODS

This specimen (Fig. 1) was caught at a depth of 45 m, 2,3 nm off the SE coast of Terceira Island (Azores), the 8th November, 2010.



Fig. 1. First specimen of *Alosa fallax* (Lacépède, 1803) found in Azorean waters (Catalogue number ALOFAL-UAç-DCA-1/10). Photo by F. Chauvin, editing by R.P.V.

The specimen shows a characteristic dark spot posterior to the gill opening, followed by eight horizontal spots along the flank just anterior and below the dorsal fin base. Its main descriptive measurements, morphometrics and meristics (see below) are in agreement with diagnostic species' data from Whitehead (1984), Rochard & Elie (1994) and Kottelat & Freyhof (2007):

Total length: 38.6cm

Fork length: 32.1cm (83% of TL)

Standard length: 30.0cm (78% of TL)

Gillrakers on the first arc (total): 78

The possibility of a misidentification with other clupeids was carefully checked using a photo database and the adequate meristics and biometric data for clupeid fish (see Froese & Pauly, 2013). After this process, and with all data available cross checked, every possible misidentification was discarded. Another clupeid besides *Sardina pilchardus* (Walbaum, 1792), in this case *Sardinella maderensis* (Lowe, 1838) was given as a possible occurrence in these waters (Santos et al. 1997). However, the above mentioned cross check definitively separated these two species and confirmed *A. fallax* as the correct identified species.

The specimen is preserved frozen in the Laboratory of Veterinary Anatomy of the Department of Agricultural Sciences (University of the Azores) under the provisional catalogue number, ALOFAL-UAç-DCA-1/10.

RESULTS AND DISCUSSION

The number of gillrakers is a distinctive feature between the recognized subspecies (Quignard & Douchement, 1991). Although, the number of gillrakers counted in our specimen does not fit with those from Whitehead's (1984) diagnosis and the author gives no references to subspecies with more than 45 gillrakers (except for *Alosa a. lacustris* (76) - now considered a synonym *A. agone* (Scopoli, 1836)), Kottelat & Freyhof 2007 indicate 30-80 gill rakers in the first arc as a diagnostic characteristic of *A. fallax*. Kottelat (1997) strongly recommends the reclassification as species for some of the presently valid subspecies and one must take note that hybrids may well interbreed (Quignard & Douchement 1991; Alexandrino et al. 1996). In fact, the whole genus *Alosa* Linck, 1790, represents a continuous challenge for many taxonomists and ichthyologists (Jolly et al. 2012).

The occurrence of *A. fallax* in the Azores was totally unexpected even when one considers it's strongly migratory habits (Whitehead, 1985; Stankus, 2009). This occurrence is even more peculiar since there are simply no estuaries in the Azores and, apparently, the specimen was isolated. According to de Groot (2002) and Astthorsson & Palsson (2006) these types of erratic/unexpected occurrences could be related with changes in the marine environment.

Moreover, this new record increases to two the number of Clupeiformes' species known to occur in Azorean waters, the other being *Sardina pilchardus* (Santos et al.

1997; Porteiro et al. 2010). It is also plausible that *A. fallax* will soon be detected in Madeira Archipelago due to its proximity to the species' southern limit records.

Acknowledgements

The authors wish to thank the crew of F/V "Pérola da Praia" and the Maritime Police of Praia da Vitória which, respectively, caught and kept this specimen.

REFERENCES

- Alexandrino, P.J., Ferrand, N., & Rocha, J. (1996). Genetic polymorphism of a haemoglobin chain and adenosine deaminase in European shads: evidence for the existence of two distinct genetic entities with natural hybridization. *Journal of Fish Biology*, 48, 447-456.
- Alexandrino, P., Faria, R., Linhares, D., Castro, F., LeCorre, M., Sabatier, R., Baglinie, J.L., & Weiss, S. (2006). Interspecific differentiation and intraspecific substructure in two closely related clupeids with extensive hybridization, *Alosa alosa* and *Alosa fallax*. *Journal of Fish Biology* 69(Supplement B), 242–259.
- Assis, C.A., Almeida, P.R., Moreira, F., Costa, J.L., & Costa, M.J. (1992). Diet of the twaite shad *Alosa fallax* (Lacépède) (Clupeidae) in the River Tagus Estuary, Portugal. *Journal of Fish Biology*, 41(6), 1049–1050.
- Astthorsson, O.S., & Palsson, J. (2006). New fish records and records of rare southern species in Icelandic water in the warm period 1996-2005. *ICES CM*, 2006/C:20.
- Cabral, M.J., Queiroz, A.I., Palmeirim, J., Almeida, J., Rogado, L., Santos-Reis, M., Oliveira, M.E., Ferrand Almeida, N., Almeida, P.R., & Dellinger, T. (2006). *Livro Vermelho dos Vertebrados de Portugal*. Instituto de Conservação da Natureza, Lisboa.
- Collares-Pereira, M.J., Cowx, I.G., Luís, T.S., Pedrosa, N., & Santos-Reis, M. (1999). Observations on the ecology of a landlocked population of allis shad in Aguiaria Reservoir, Portugal. *Journal of Fish Biology*, 55, 658–664. doi: 10.1111/j.1095-8649.1999.tb00706.x
- Costa, M.J., Almeida, P.R., Domingos, I.M., Costa, J.L., Correia, M.J., Chaves, M.L., & Teixeira, C.M. (2001). Present status of the main shads' populations in Portugal. *Bulletin Français de la Pêche et de la Pisciculture*, 362/363, 1109–1116.
- de Groot, S.J. (2002). A review of the past and present status of anadromous fish species in the Netherlands: is restocking the Rhine feasible? *Hydrobiologia*, 478, 205-218.



- Freyhof, J., & Kottelat, M. (2008). *Alosa fallax*. IUCN 2010. *IUCN Red List of Threatened Species*. Version 2010.4. <www.iucnredlist.org>.
- Froese, R., & Pauly, D. (Eds.) (2013). FishBase. World Wide Web electronic publication. www.fishbase.org, version (12/2013).
- Jolly, M. T., Aprahamian, M.W., Hawkins, S.J., Henderson, P.A., Hillman, R., OMaoileidigh, N., Maitland, P.S., Piper, R., & Genner, M.J. (2012). Population genetic structure of protected allis shad (*Alosa alosa*) and twaite shad (*Alosa fallax*). *Marine Biology*, 159, 675 - 687.
- Kottelat, M. (1997). European freshwater fishes. *Biologia* 52, Suppl. 5, 1-271.
- Kottelat, M., & Freyhof, J. (2007). *Handbook of European Freshwater Fishes*. Publications Kottelat, Cornol, Switzerland. 646 pp.
- Oesmann, S., & Thiel, R. (2001). Feeding of juvenile twaite shad (*Alosa fallax* Lacépède, 1803) in the Elbe Estuary. *Bulletin Français de la Pêche et de la Pisciculture*, 362/363, 785-800.
- Pina, T., Esteves, E., & Andrade, J.P. (2003). Gross and histological observations of ovarian development in twaite shad, *Alosa fallax fallax*, from the Rivers Mira and Guadiana (Portugal). *Scientia Marina*, 67(3), 313-322.
- Porteiro, F.M., Menezes, G.M., Afonso, P., Monteiro, J.G., & Santos, R.S. (2010). *Marine Fish (Chondrichthyes, Actinopterygii)*. Pp. 326-345 In: Borges, P.A.V., Costa, A., Cunha, R., Gabriel, R., Gonçalves, V., Martins, A.F., Melo, I., Parente, M., Raposeiro, P., Rodrigues, P., Santos, R.S., Silva, L., Vieira, P., & Vieira, V. (Eds.). *A List of the Terrestrial and Marine Biota from the Azores*. Principia, Cascais.
- Quignard, J.-P., & Doucement, C. (1991). *Alosa fallax fallax* (Lacépède 1803). Pp. 225-253 In: Hoestlandt, H. (Ed.). *The Freshwater Fishes of Europe*. Vol. 2. Clupeidae, Anguillidae. AULA-Verlag, Wiesbaden.
- Rochard, E., & Elie, P. (1994). *La Macrofaune Aquatique de l'estuaire de la Gironde. Contribution au Livre Blanc de l'Agence de l'Eau Adour Garonne*. Pp. 1-56 in: Mauvais, J.-L., & Guillaud, J.-F. (Eds.). *État des connaissances sur l'estuaire de la Gironde. Agence de l'Eau Adour-Garonne*. Éditions Bergeret, Bordeaux.
- Santos, R.S., Porteiro, F.M., & Barreiros, J.P. (1997). *Marine fishes of the Azores: An annotated checklist and bibliography*. Arquipélago. Life and Marine Sciences Supplement 1: xxiii + 242pp. Ponta Delgada.
- Stankus, S. (2009). Spawning Migration and Population Condition of Twaite Shad (*Alosa fallax*, Lacépède 1803) in Lithuania. *Environmental Research, Engineering and Management*, 4(50), 20-29.



Whitehead, P.J.P. (1984). *Clupeidae*. Pp. 268-281 In: Whitehead, P.J.P., Bauchot, M.-L., Hureau, J.-C., Nielsen, J., & Tortonese, E. (Eds.). *Fishes of North-eastern Atlantic and the Mediterranean*. Vol. I. UNESCO, Paris.

Whitehead, P.J.P. (1985). *Clupeoid fishes of the world (suborder Clupeoidei)*. FAO Species Catalogue. Vol. 7 in: *An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, shads, anchovies and wolf-herrings. Part 1 - Chirocentridae, Clupeidae and Pristigasteridae*. FAO Fisheries Synopsis 125(7/1), 1-303.