

Kuroshio Science 8-1, 109-112, 2014

Parasitic Crustaceans and Marine Invasions: Two Case Studies from Kuroshio Region (extended abstract of the 7th Kuroshio Symposium)

Gyo Itani^{1,2*}, Chiharu Yamada¹, Hodaka Asama¹, Yumi Henmi², Hiroshi Kume³ and John W. Chapman⁴

¹ Graduate School of Kuroshio Science, Kochi University, 2-5-1 Akebono, Kochi, Kochi 780-8520, Japan.

² Graduate School of Education, Kochi University, 2-5-1 Akebono, Kochi, Kochi 780-8520, Japan.

³ Fisheries Research Center, Ehime Prefecture, 1611 Ikeda, Tambara, Saijo, Ehime 791-0508, Japan.

⁴ Hatfield Marine Science Center, Oregon State University, Newport, Oregon 97365-5296, USA.

ABSTRACT

Parasitic crustaceans are ubiquitous in marine environment. They have negative effects on host individuals and populations, and even on the ecosystem where the host species plays a great role. In this short review, we show two case studies of marine invasions related with parasitic crustaceans from Kuroshio Region. One is the native pinnotherid crab parasitizing introduced mussels and the other is the bopyrid isopod introduced from Asia to US Pacific Coast. The green mussel *Perna viridis*, native to the tropical Indo-Pacific region, has recently been introduced worldwide. Although the native mussel populations in India suffer from parasitic effects of the pinnotherid crab *Arcotheres placunae*, mussels are relatively free from parasitic crabs in invaded regions (parasite escape). We investigated pea crab infections in *P. viridis* in Uranouchi Inlet, Kochi Prefecture, Japan. Results showed that the native generalist pea crab *A. sinensis* utilized the non-indigenous mussel *P. viridis*, though the prevalence is much lower. The introduced Asian parasitic bopyrid isopod, *Orthione griffenis*, was first discovered on the Pacific coast of North America ca. 25 years ago. High prevalence of *Orthione* infestations decreased many local populations of the host mud shrimp *Upogebia pugettensis* as well as the diverse symbiotic fauna in the host burrows. We investigated the distribution and host specificity of the bopyrid in western Japan. Results showed that *O. griffenis* was a rare species in Japan. Further ecological studies on parasitic crustaceans are needed in both native (source) and introduced (recipient) habitats.

Keywords : crustacean parasite, invasion, pinnotherid crab, *Perna* mussel, bopyrid isopod, *Upogebia* shrimp

Introduction

Parasitic crustaceans are ubiquitous in marine environment (Rohde, 2005). They have negative effects on host individuals and populations, and even on the ecosystem where the host species plays a great role. Parasites present several aspects to marine invasions.

The enemy release hypothesis (ERH) or parasite escape can sometimes explain the success of introduced species (Torchin *et al.*, 2002; Torchin and Lafferty, 2009; Blakeslee *et al.*, 2013). That is, many introduced species lack natural enemies and parasites in non-native (recipient) habitats, thus giving them an advantage over

native species. However, some native parasites may utilize the introduced species over time. Reports of marine parasites invading marine environments are few (Torchin *et al.*, 2002; Rohde, 2005). When parasite invasion is successful, the effect on the new hosts in recipient habitats is often deleterious (Griffen, 2009).

In this short review, we show two case studies of marine invasions related with parasitic crustaceans from Kuroshio Region. One is the native Japanese pinnotherid crab parasitizing introduced tropical mussels and the other is the bopyrid isopod introduced from Asia to US Pacific Coast.

*Corresponding author: e-mail itani@kochi-u.ac.jp

Case study 1: Utilization of the non-indigenous green mussel by the native pinnotherid crab in Japan

The green mussel *Perna viridis* is native to the tropical Indo-Pacific region, but has recently been introduced worldwide (Baker *et al.*, 2007). Pea crabs of the family Pinnotheridae are often found living together with mussels worldwide, and in some cases parasitic effects of the crabs on the host have been confirmed (Haines *et al.*, 1994). In the case of *P. viridis*, native populations suffer from pea crabs (Rajagopal *et al.*, 2006). Jose and Deepthi (2005) reported that about 6% of the *P. viridis* population in Kerala, India, was parasitized by the pea crab *Arcotheres placunae* (= *Pinnotheres placunae*) and infested mussels showed significant reductions in shell length and live weight.

Yamada *et al.* (2009) showed utilization of *P. viridis* by the native pinnotherid crab *Arcotheres sinensis* (Shen, 1932) (= *Pinnotheres sinensis*) in Uranouchi Inlet, Kochi



Figure 1. Invaded green mussel, *Perna viridis*, in natural rocky habitats at Uranouchi Inlet, Kochi, Japan.



Figure 2. A native pinnotherid crab, *Arcotheres sinensis*, parasitizing *Perna viridis* collected at Uranouchi Inlet, Kochi, Japan.

Prefecture, southwestern Japan (Yamada *et al.*, 2009). In Uranouchi Inlet, *P. viridis* is densely distributed in both artificial and natural habitats (Figure 1), where many individuals can overwinter (Yamada *et al.*, 2010; Ueda *et al.*, 2013). About 30 *P. viridis* individuals with shell lengths from 70 to 95 mm were collected at random monthly from February 2008 to January 2009 from the undersurfaces of raft floats. Consequently, 1.8% of 340 green mussels were parasitized by *A. sinensis* (Figure 2). Infested mussels had lower Condition Index values, indicating parasitic effects by the crabs.

Yamada *et al.*, (2009) revealed that the newly invaded mussel *P. viridis* in Japan is not parasitized by the original parasite *A. placunae*, but parasitized by a native generalist parasite *A. sinensis* in the recipient habitats. This is a similar situation to that recorded from New Zealand, where the invasive mussel *Musculista senhousia* was parasitized by the native generalist pea crab *Pinnotheres novaezelandiae* (Miller *et al.*, 2008). The relatively low prevalence of *A. sinensis* in *P. viridis* in Yamada *et al.* (2009) indicates that the parasitic effect of pea crabs on the green mussel population in Japan is low. However, the reduced condition index means that the parasite can cause negative effects to the host individual. Future studies must investigate whether *A. sinensis* comes to utilize *P. viridis* more often over time, reducing the impacts of mussel invasion to the native environment, as discussed by Mouristen and Poulin (2002). In addition, ecological studies of parasites in *P. viridis* in native (source) habitats are still limiting. Further ecological studies on parasitic crustaceans are needed in both native (source) and introduced (recipient) habitats.

Case study 2: An introduced Asian bopyrid isopod threatens northeastern Pacific estuarine ecosystems

Bopyrid isopods are parasites that castrate host decapod crustaceans (Williams and Boyko, 2012). The introduced Asian parasitic bopyrid isopod, *Orthione griffenis* (Figure 3), was first discovered on the Pacific coast of North America ca. 25 years ago (Markham, 2004; Chapman *et al.*, 2012). High prevalence of *Orthione* infestations decreased many local populations of the host mud shrimp *Upogebia pugettensis* as well as the diverse symbiotic fauna in the host burrows (Griffen, 2009; Dumbauld *et al.*, 2011; Chapman *et al.*, 2012). Chapman *et al.* (2012) revealed using six criteria that *Orthione* is introduced to North America: “its conspecificity with disjunct Asian populations, its earliest collec-



Figure 3. Invaded bopyrid isopod, *Orthione griffenis*, collected at Yaquina Bay, Oregon, USA. A female (larger) and a male (on the female abdomen).

tions in Asia, its late discovery among symbiotic species associated with *Upogebia*, its historical absence, and its appearance in North America coincident with extensive new ballast water traffic from Asia”.

Itani (2004) investigated the distribution and host specificity of the bopyrid isopods infesting upogebiid shrimps in western Japan. As a result, nine species (*Gyge ovalis*, *Progebiophilus villosus*, *Progebiophilus* sp. 1, *Progebiophilus* sp. 2, Pseudioninae sp. 1, Pseudioninae sp. 2, *Procepon insolitum*, *Upogebione* sp., and *Phylloporus* sp.) were collected (Itani, 2004), although Pseudioninae sp. 1 was later identified with *Orthione griffenis* (Chapman *et al.*, 2012). *Orthione griffenis* was collected from various localities, parasitizing *Upogebia major*, *U. issaeffi*, and *Austinogebia narutensis* (Itani, 2004). Our unpublished researches (by GI, YH, and HK) suggested that prevalence of *O. griffenis* was extremely low compared to the commonest species *Gyge ovalis* in Japan. Future studies must explain why *O. griffenis* was successfully invaded to North Eastern Pacific, instead of *G. ovalis*, abundant in Japanese waters. Further ecological studies on *O. griffenis* are needed in both native (source) and introduced (recipient) habitats, as well as on many other bopyrid isopods.

We must take the greatest care not to let bopyrid isopods invade, considering the deleterious effects of this parasite on the host decapod crustaceans (Griffen, 2009; Dumbauld *et al.*, 2011; Chapman *et al.*, 2012). Bopyrid isopods of the subfamily Orbioninae are parasites of penaeid prawns (Figure 4) that are not distributed



Figure 4. A penaeid prawn with Orbioninae parasite in the left branchial chamber, bought in a supermarket at Kochi, Japan.

in American continents (Markham, 1986). When one of the Orbioninae species is introduced to America, it will damage penaeid shrimp populations and shrimp industries in American continents. Ecology and life history of bopyrids including Orbioninae species should also be investigated.

Acknowledgment

This work was partly supported by JSPS KAKENHI Grant Number 24510328.

References

- Baker, P., Fajans, J. S., Arnold, W. S., Ingrao, D. A., Marelli, D. C., and Baker, S. M., 2007. Range and dispersal of a tropical marine invader, the Asian green mussel, *Perna viridis*, in subtropical waters of the Southeastern United States. *Journal of Shellfish Research*, 26: 345-355.
- Blakeslee, A. M., Fowler, A. E., and Keogh, C. L., 2013. Marine invasions and parasite escape: updates and new perspectives. *Advances in Marine Biology*, 66: 87-169.
- Chapman, J. W., Dumbauld, B. R., Itani, G., and Markham, J. C., 2012. An introduced Asian parasite threatens northeastern Pacific estuarine ecosystems. *Biological Invasions*, 14: 1221-1236.
- Dumbauld, B.R., Chapman, J. W., Torchin, M. E., and Kuris, A. M., 2011. Is the collapse of mud shrimp (*Upogebia pugettensis*) populations along the Pacific coast of North America caused by outbreaks of a previously unknown bopyrid isopod parasite (*Orthione griffenis*)? *Estuaries and Coasts* 34: 336–350.
- Griffen, B. D., 2009. Effects of a newly invasive parasite on the burrowing mud shrimp, a widespread ecosystem engineer. *Marine Ecology Progress Series* 391: 73-83.

- Haines, C. M. C., Edmunds, M., and Pewsey, A. R., 1994. The pea crab, *Pinnotheres pisum* (Linnaeus, 1767), and its association with the common mussel, *Mytilus edulis* (Linnaeus, 1758), in the Solent (UK). *Journal of Shellfish Research*, 13: 5-10.
- Itani, G., 2004. Host specialization in symbiotic animals associated with thalassinidean shrimps in Japan. In: A. Tamaki, (ed), Proceedings of the symposium on "Ecology of large bioturbators in tidal flats and shallow sublittoral sediments -from individual behavior to their role as ecosystem engineers." Nagasaki University, pp. 33-43.
- Jose, B. and Deepthi, T. R., 2005. Green mussel *Perna viridis*, a new host for the pea crab *Pinnotheres placunae* along the Malabar coast, Kerala. *Current Science*, 89: 1090-1091.
- Markham, J. C., 1986. Evolution and zoogeography of the Isopoda Bopyridae, parasites of Crustacea Decapoda. In: R. H. Gore and K. L. Heck (eds). *Crustacean Issues 4 Crustacean Biogeography*. Rotterdam: A.A. Balkema. pp 143-164.
- Markham, J. C., 2004. New species and records of Bopyridae (Crustacea : Isopoda) infesting species of the genus *Upogebia* (Crustacea : Decapoda : Upogebiidae): the genera *Orthione* Markham, 1988, and *Gyge* Cornalia & Panceri, 1861. *Proceedings of the biological society of Washington*, 117: 186-198.
- Miller, A., Inglis, G. J., and Poulin, R., 2008. Use of the introduced bivalve, *Musculista senhousia*, by generalist parasites of native New Zealand bivalves. *New Zealand Journal of Marine and Freshwater Research*, 42: 143-151.
- Mouritsen, K. M. and Poulin, R., 2002. Parasitism, community structure and biodiversity in intertidal ecosystems. *Parasitology*, 124: S101-S117.
- Poulin, R. and Mouillot, D., 2003. Host introductions and the geography of parasite taxonomic diversity. *Journal of Biogeography*, 30: 837-845.
- Rajagopal, S., Venugopalan, V. P., Van Der Velde, G., and Jenner, H. A., 2006. Greening of the coasts: a review of the *Perna viridis* success story. *Aquatic Ecology*, 40: 273-297.
- Rohde K. (ed.), 2005. *Marine parasitology*. CSIRO Publishing. Collingwood, Australia. 565 pp.
- Torchin, M. E. and Lafferty, K. D., 2009. Escape from parasites. In : G. Rilov and J. A. Crooks, (eds.), *Biological invasions in marine ecosystems*, Springer-Verlag, Berlin Heidelberg, *Ecological studies*, 204: 203-214.
- Torchin, M. E., Lafferty, K. D., and Kuris, A. M., 2002. Parasites and marine invasions. *Parasitology* 124: S137-151.
- Ueda, I., Sakaguchi, I., Ogita, T., Yamada, C., and Itani, G., 2013. Thermal conditions of overwintering for tropical green mussel *Perna viridis* in Uranouchi Inlet, Tosa Bay, Shikoku, Japan in 2010. *Sessile organisms*, 30: 29-36.
- Williams, J. D. and Boyko, C. B., 2012. The global diversity of parasitic isopods associated with crustacean hosts (Isopoda: Bopyroidea and Cryptoniscoidea). *PLoS One* 7: e35350.
- Yamada, C., Itani, G., and Asama, H., 2009. Utilization of the non-indigenous green mussel, *Perna viridis*, by the native pinnotherid crab *Arcotheres sinensis* in Uranouchi Inlet, Kochi, Japan. *Crustacean Research*, 38: 70-76.
- Yamada, C., Itani, G., and Ueda, H., 2010. Habitat use and horizontal distribution of the green mussel, *Perna viridis*, in Uranouchi Inlet, Kochi Prefecture. *Sessile organisms*, 27: 41-50.