

HOSTED BY



ELSEVIER

Contents lists available at [ScienceDirect](http://www.elsevier.com/locate/japb)

Journal of Asia-Pacific Biodiversity

journal homepage: <http://www.elsevier.com/locate/japb>Journal of
Asia-Pacific
Biodiversity

Original Article

Two new records of the orthotyline plant bug (Hemiptera: Heteroptera: Miridae) from the Korean Peninsula

Junggon Kim, Sunghoon Jung*

Laboratory of Systematic Entomology, Department of Applied Biology, College of Agriculture and Life Sciences, Chungnam National University, Daejeon, South Korea

ARTICLE INFO

Article history:

Received 27 July 2016

Received in revised form

23 August 2016

Accepted 13 September 2016

Available online xxx

Keywords:

Miridae

new record

Orthotylus bilineatus

the Korean Peninsula

Zanichius tarasovi

ABSTRACT

Two new records of the subfamily Orthotylinae (Heteroptera: Miridae: Orthotylinae), *Zanichius tarasovi* Kerzhner, 1988 and *Orthotylus bilineatus* (Fallén, 1807), are reported for the first time to the Korean fauna. The genus *Zanichius* is also first recorded from the Korean Peninsula. Redescriptions of genitalia, diagnoses of each species and genus, and biological notes are presented with the photographs and illustrations.

Copyright © 2016, National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA). Production and hosting by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The subfamily Orthotylinae is the third largest group in the family Miridae (Insecta: Hemiptera: Heteroptera) in the world (Schuh and Slater 1995), and is one of the remarkable groups economically because it contains insect pest groups and predator groups (Schaefer and Panizzi 2000). Among them, some species of the predatory groups have been developed and used as biological control agents, such as *Cyrtorhinus fulvus* (Wheeler 2000).

In this paper, two orthotyline species are newly reported from the Korean Peninsula, *Zanichius tarasovi* Kerzhner, 1988 and *Orthotylus bilineatus* (Fallén, 1807). Regarding *Z. tarasovi*, its higher taxa, genus *Zanichius*, is also reported for the first time. Morphological information including diagnoses and descriptions of male genital structures, and biological notes are provided along with photographs and illustrations.

Materials and methods

Photographs of dorsal habitus and genitalia were taken by a Leica DMC 2900 camera combined with a Leica M165C microscope.

* Corresponding author. Tel.: +82 42 821 5767; fax: +82 42 823 8679.

E-mail address: jung@cnu.ac.kr (S. Jung).

Peer review under responsibility of National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA).

<http://dx.doi.org/10.1016/j.japb.2016.09.004>

pISSN2287-884X eISSN2287-9544/Copyright © 2016, National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA). Production and hosting by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Measurements were taken using a software program equipped with the same microscope. All measurements are given in millimeters. To observe male genital structures, the genital segment was detached and then boiled in 10% KOH solution at 70°C for 3–5 minutes until it was transparent. After placing in distilled water, it was dissected to examine endosoma and parameres. Terminology mainly follows Yasunaga (1999). The depository of examined specimens is Laboratory of Systematic Entomology, Chungnam National University (CNU), Daejeon, Korea. Host with an asterisk means it is a new record.

Taxonomic accounts

Genus *Zanichius* Distant, 1904 연초록들장님노린재속 (신칭)
Zanichius Distant, 1904: 477. Type species: *Zanichius annulatus* Distant, 1904.

Zonodorus Distant, 1909: 522 (synonymized by Carvalho, 1952: 79).
Type species: *Zonodorus gubernator* Distant, 1909.

Uzeliella Poppius, 1911: 31 (junior homonym of *Uzeliella* Bagnal, 1908, Thysanoptera) (synonymized by Carvalho, 1952: 79). Type species: *Uzeliella flavovirens* Poppius, 1911.

Poppiella Bergroth, 1911: 188. New name for *Uzeliella* Poppius, 1911.
Habrocoris Wagner, 1951: 153 (synonymized by Linnavuori, 1964: 329). Type species: *Habrocoris breviceps* Wagner, 1951.

Diagnosis. Recognized by body generally green coloration and elongated oval; frons anteriorly projected; compound eye not

touching anterior margin of pronotum; antennae with reddish or dark markings; lateral margin of hemelytron rounded; legs without patterns (Figures 1A–C); endosoma membranous with sclerites (Figure 2D).

Zanchius tarasovi Kerzhner, 1988 연초록들장님노린재 (신칭) (Figures 1A, 1C, 2, 3A–C)

Zanchius tarasovi Kerzhner, 1988: 49; Yasunaga, 1999: 167; Lin, 2005: 191.

Diagnosis. Recognized by head with reddish marking behind compound eyes; first antennal segment with reddish stripe laterally (Figure 1C); scutellum usually with reddish marking in middle; apex of clavus with reddish marking; inner part of corium usually with reddish marking, reddish marking sometimes individual variation in size (Figure 1A); sclerite I broad with approximately eight teeth (Figures 2D and 2E).

Description. See Kerzhner (1988) and Yasunaga (1999) for original and detailed description, respectively.

Male genitalia. Left paramere scythe shaped with sharp hypophysis and some sparse setae (Figure 2A); right paramere slightly curved, with distinct project medially (Figures 2B and 2C); endosoma membranous with three sclerites, sclerite I broad with approximately eight teeth, sclerite II somewhat shin and tapered at apex, sclerite III thicker than sclerite II and curved (Figures 2D and 2E).

Measurements (in mm). Male ($n = 2$)/female ($n = 1$) body length, tylus–apex of membrane: 4.48–4.59/5.01; head length, excluding collar: 0.30–0.32/0.33; head width, including compound eyes: 0.77–0.79/0.79; vertex width: 0.32–0.33/0.38; first antennal segment length: 0.42–0.45/0.43; second antennal segment length: 1.90–1.93/2.10; third antennal segment length: 1.08–1.12/missing; fourth antennal segment length: 0.68–0.74/missing; total antennal length: 4.08–4.24/2.53; first rostral segment length: 0.30–0.32/0.36; second rostral segment length: 0.34–0.36/0.36; third rostral segment length: 0.39–0.43/0.40; fourth rostral segment length: 0.35–0.37/0.41; total rostral length: 1.38–1.48/1.53; mesal pronotal length: 0.47–0.49/0.62; basal pronotal maximal width (straight): 1.16–1.17/1.24; anterior scutellumal width: 0.75–0.78/0.78; mesal scutellumal length: 0.92–0.94/0.93; outer embolial margin length (straight): 2.39–2.47/2.60; outer cuneal margin length (straight): 0.72–0.76/0.84; maximal width across hemelytron: 0.79–0.85/0.93; foreleg (femur:tibia:tarsus): 0.96–1.00:1.20–1.32:0.33–0.35/1.22:1.48:0.35; midleg (femur:tibia:tarsus): 1.10–1.14:1.31–

1.42:0.33–0.37/1.29:1.61:0.37; hindleg (femur:tibia:tarsus): 1.67–1.73:2.47–2.51:0.41–0.42/1.78:2.85:0.48.

Specimen examined. [CNU] 1♂, Gung-dong, Yuseong-gu, Daejeon, Korea, 16 ix 2014, by Light trap (JG Kim); 1♂, 1♀, Saeun-ri, Chilseong-myeon, Goesan-gun, Chungcheongbuk-do, Korea, 6 viii 2015, by Light trap (JG Kim); [CNU] 2♂♂, 1♀, Pocheon-si, Gyeonggi-do, Korea, 9 vii 2016, on *Quercus aliena* (JH Park).

Host. *Juglans ailantifolia* (Juglandaceae), *Tilia japonica* (Tiliaceae), *Quercus crispula* (Fagaceae), *Alnus* spp. (Betulaceae), *Salix* spp. (Salicaceae) (Yasunaga 1999), *Q. aliena* (Fagaceae)*.

Distribution. Korea (new record), China, Japan, Russia, Taiwan.

Biology. This species is reported as a predator for many leafhoppers, especially typhlocybinid leafhopper such as *Limassolla* species on persimmon tree in China and Japan (Zheng and Liang 1991; Zheng and Liu 1993; Yasunaga 1999; Wheeler 2001). This species has been attracted to light trap.

Remarks. Last instar of this species is mostly pale greenish in color. First and second antennal segments are pale brown, with slight reddish tinge at the base and apex (Figure 3C), while those of adult are reddish (Figures 3A and 3B).

Genus *Orthotylus* Fieber, 1858 들장님노린재속

Orthotylus Fieber, 1858: 315. Type species: *Cimex nassatus* Fabricius sensu Fieber, 1858 (= *Orthotylus marginalis* Reuter, 1883).

Litosoma Douglas & Scott, 1865: 334 (synonymized by Putton, 1869: 26). Type species: *Litosoma nassata* sensu Douglas & Scott, 1865 (= *Orthotylus marginalis* Reuter, 1883).

Diommatus Uhler, 1887: 32 (synonymized by Horvath, 1908: 9).

Type species: *Diommatus congrex* Uhler, 1887 (= *Lygus dorsalis* Provancher, 1872).

Diagnosis. Recognized by body generally greenish, elongated or elongated oval; compound eye reaching or almost reaching to anterior margin of pronotum; lateral margin of hemelytron straight or slightly rounded; pronotum and hemelytra covered with brown to dark setae, or entirely covered with pale setae; male genitalia with several branch-shaped sclerites.

Orthotylus bilineatus (Fallén, 1807) 검은줄들장님노린재 (신칭) (Figures 1B, 3D, 4)

Capsus bilineatus Fallén, 1807: 102.

Capsus kirschbaumii Flor, 1860: 465, 614 (synonymized by Douglas & Scott, 1866: 246).

Pseudorthotylus sordidus Poppius, 1914: 66 (synonymized by Linnavuori, 1994: 31).




Figure 1. Dorsal habitus and diagnostic character of *Zanchius tarasovi* and *Orthotylus bilineatus*: A, habitus of *Z. tarasovi*; B, ditto, *O. bilineatus*; C, reddish markings of *Z. tarasovi* in lateral view. <scale bar: 1 mm>.




Figure 2. Male genitalia of *Zanchius tarasovi*: A, left paramere; B, right paramere in dorsal view; C, ditto, in ventral view; D, endosoma; E, sclerite I in detail view. sc = sclerite. <scale bar: 0.1 mm>.




Figure 3. *Zanchius tarasovi* and *Orthotylus bilineatus*: A–C, *Z. tarasovi* (A and B, adult; C, final nymph); D, *O. bilineatus* (adult).

Orthotylus (Neomecomma) bilineatus, Southwood, 1953: 433; Wagner & Weber, 1964: 313; Miyamoto, 1969: 78; Wagner, 1974: 182.

Orthotylus (Pseudorthotylus) bilineatus Linnavuori, 1994: 31.

Diagnosis. Recognized by body generally greenish pale brown; head pale brown with longitudinal dark stripe, antennae entirely dark; pronotal collar dark brown; anterior and posterolateral part of pronotum dark; mesal part of scutellum dark; inner part of clavus and corium dark; legs mostly greenish brown; cuneus greenish (Figure 1B); endosoma with long and sharp spicule with triangular process subapically (Figures 4A–C).

Description. See Wagner and Weber (1964) for detailed description.

Male genitalia. Sensory lobe of left paramere with blunt process sub-basally, hypophysis broad at apex (Figure 2A), right paramere curved, hypophysis blunt laterally (Figures 4A and 4B); endosoma

with long and sharp spicule, with triangular process subapically; basal part thicker (Figure 4C).

Measurements (in mm). Male ($n = 2$)/female ($n = 1$) body length, tylus–apex of membrane: 4.82–4.87/4.89; head length, excluding collar: 0.45–0.47/0.47; head width, including compound eyes: 0.96–0.97/0.98; vertex width: 0.28–0.29/0.30; first antennal segment length: 0.42–0.44/0.43; second antennal segment length: 1.80–1.83/1.83; third antennal segment length: 1.30–1.33/1.34; fourth antennal segment length: 0.28–0.29/0.29; total antennal length: 3.80–3.89/3.89; first rostral segment length: 0.34–0.35/0.35; second rostral segment length: 0.32–0.33/0.33; third rostral segment length: 0.26–0.28/0.27; fourth rostral segment length: 0.31–0.32/0.32; total rostral length: 1.23–1.28/1.27; mesal pronotal length: 0.66–0.68/0.69; basal pronotal maximal width (straight): 1.17–1.18/1.19; anterior scutellum width: 0.55–0.56/0.56; mesal scutellum length: 0.80–0.82/0.83; outer embolial margin length (straight): 2.11–2.13/2.14; outer cuneal margin length (straight):




Figure 4. Male genitalia of *Orthotylus bilineatus*: A, left paramere; B, right paramere; C, endosoma. <scale bar: 0.1 mm>.

0.84–0.86/0.88; maximal width across hemelytron: 0.70–0.72/0.73; foreleg (femur:tibia:tarsus): 1.09–1.11:1.13–1.14:0.41–0.43/1.11:1.13:0.43; midleg (femur:tibia:tarsus): 1.12–1.14:1.51–1.53:0.48–0.49/1.13:1.54:0.49; hindleg (femur:tibia:tarsus): 1.56–1.58:2.28–2.31:0.50–0.52/1.58:2.32:0.54.

Specimen examined. [CNU] 3♂, 1♀, Hoenggye-ri, Daegwanryeong-myeon, Pyeongchang-gun, Gangwon-do, Korea, 29–30 vi 2016, by Light trap (JG Kim).

Host. *Populus tremula*, *Salix* sp. (Salicaceae) (Kerzhner 1964; Putshkov 1971; Wagner 1974; Yasunaga 1999; Kondorosy and Szeoke 2004).

Distribution. Korea (new record), China, Japan, Russia, Taiwan.

Remarks. This species has been attracted to light trap near *Salix* habitat at an altitude of approximately 900 m.

Acknowledgments

We would like to thank Mr WG Kim and Mr JH Park for providing pictures and valuable samples for this study. This research was supported by a grant from the National Institute of Biological Resources, funded by the Ministry of Environment of the Republic of Korea, and was supported by the Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Science, ICT & Future Planning (NRF-2014R1A1A1005338).

References

- Bergroth E. 1911. On some recently described Hemiptera chiefly from India. *Annales de la Societe Entomologique de Belgique* 55:184–189.
- Distant WL. 1904. *The fauna of British India, including Ceylon and Burma. Rhynchota*. London: Taylor & Francis.
- Distant WL. 1909. Descriptions of Oriental Capsidae. *Annals and Magazine of Natural History* 4:509–523.
- Douglas JW, Scott J. 1865. *The British-Hemiptera*, In *Hemiptera-Heteroptera*, Vol. 1. London: The Ray Society.
- Fallén CF. 1807. *Monographia Cimicum Sveciae*. Hafniae. C.G. Proft.
- Fieber FX. 1858. Kriterien zur generischen Theilung der Phytocoriden (Capsini auct.). *Wiener Entomologische Monatschrift* 2. 289–327, 329–347, 388, 1 pl.
- Flor G. 1860. *Die Rhynchoten Livlands in systematischer Folge beschrieben*, Vol. 1; 1860. Dorpat: Schulz.
- Kerzhner IM. 1964. Family Isometopidae. Family Miridae (Capsidae). In: , Bienko GY, editor. *Opredeelite'nasekomykh evropeiskoichasti SSSR [Keys to the*

Insects of the European part of the USSR], Vol. 1. Moskova and Leningrad: Nauka [In Russian; English translation: 1967, Jerusalem: Israel Program for Scientific Translations].

Kerzhner IM. 1988. *Novye i maloizvestnye poluzhestkiokrylye nasekomye (Heteroptera) Dal'nego Vostoka SSSR [New and little known heteropteran insects (Heteroptera) from the Far East of the USSR]*. Vladivostok: Academy of Science USSR [In Russian].

Kondorosy E, Szeoke K. 2004. Two mirid bugs (Heteroptera: Miridae) from the Koszeg Mountains, new for the Hungarian fauna. *Folia Entomologica Hungarica* 50:242–244 [In Hungarian].

Lin CS. 2005. The genus *Zanchius* (Hemiptera: Miridae) of Taiwan. *Formosan Entomologist* 25:185–194.

Linnavuori RE. 1994. Orthotylineae of west, central and north-east Africa (Heteroptera, Miridae). *Acta Zoologica Fennica* 193:1–84.

Miyamoto S. 1969. Some Orthotylineae from Japan (Hemiptera-Heteroptera: Miridae). *Sieboldia* 4:75–83.

Poppius B. 1911. Beiträge zur Kenntnis der Miriden-Fauna von Ceylon. *Öfversigt af Finska Vetenskaps societeters Förhandlingar* 53:1–36.

Poppius B. 1914. Die Miriden der Äthiopischen Region II—Macrolophinae, Heterotominae, Phylinae. *Acta Societatis Scientiarum Fennicae* 44:1–136.

Putshkov VG. 1971. On the ecology of some little known Heteroptera from the European regions of the USSR. Communication IV. Miridae. *Vestnik Zoologii* 1971:30–35 [In Russian].

Schaefer CW, Panizzi AR. 2000. *Heteroptera of economic importance*. Boca Raton: CRC Press.

Schuh RT, Slater JA. 1995. *True bugs of the world (Hemiptera: Heteroptera). Classification and natural history*. Ithaca, NY: Cornell University Press.

Southwood TRE. 1953. The morphology and taxonomy of the genus *Orthotylus* Fieber (Hem., Miridae), with special reference to the British species. *Transactions of the Royal entomological Society of London* 104:415–449.

Uhler PR. 1887. Observations on some Capsidae with descriptions of a few new species. (No. 2). *Entomologica Americana* 3:29–35.

Wagner E. 1951. Einige neue Miridenarten aus Ägypten. *Bulletin de la Societe Fouad Ter d'entomologie* 35:153–161.

Wagner E. 1974. Die Miridae Hahn, 1831, des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 2. *Entomologische Abhandlungen Staatlichen Museum für Tierkunde Dresden* 39:1–421.

Wagner E, Weber HH. 1964. *Hétéroptères, Miridae. Faune de France* 67. Paris: Fédération Française des Sociétés de Sciences Naturelles.

Wheeler AG. 2000. Predacious plant bugs (Miridae). In: Schaefer CW, Panizzi AR, editors. *Heteroptera of economic importance*. Ithaca, NY: Cornell University Press.

Wheeler AG. 2001. *Biology of the plant bugs: pests, predators, opportunists*. Ithaca, NY: Cornell University Press.

Yasunaga T. 1999. The plant bug tribe Orthotylini in Japan (Heteroptera: Miridae: Orthotylineae). *Tijdschrift voor Entomologie* 142:143–183.

Zheng LY, Liang LJ. 1991. Mirid bugs preying on Persimmon Leafhoppers from China. *Acta Scientiarum Naturalium Universitatis Nankaiensis* 1991:84–87.

Zheng LY, Liu GQ. 1993. Two new species of genus *Zanchius* Dist. (Insecta: Hemiptera: Heteroptera: Miridae). *Reichenbachia* 30:17–20.