



ELSEVIER

Topology and its Applications 103 (2000) 203–219

**TOPOLOGY  
AND ITS  
APPLICATIONS**

[www.elsevier.com/locate/topol](http://www.elsevier.com/locate/topol)

## Darboux-like functions within the classes of Baire one, Baire two, and additive functions <sup>☆</sup>

Krzysztof Ciesielski <sup>a,\*</sup>, Jan Jastrzębski <sup>b,1</sup>

<sup>a</sup> *Department of Mathematics, West Virginia University, Morgantown, WV 26506-6310, USA*

<sup>b</sup> *Department of Mathematics, Gdańsk University, Wita Stwosza 57, 80-952 Gdańsk, Poland*

Received 21 July 1998; received in revised form 3 November 1998

---

### Abstract

In the paper we present an exhaustive discussion of the relations between Darboux-like functions within the classes of Baire one, Baire two, Borel, and additive functions from  $\mathbb{R}^n$  into  $\mathbb{R}$ . In particular we construct an additive extendable discontinuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$ , answering a question of Gibson and Natkaniec (1996–97, p. 499), and show that there is no similar function from  $\mathbb{R}^2$  into  $\mathbb{R}$ . We also describe a Baire class two almost continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  which is not extendable. This gives a negative answer to a problem of Brown, Humke, and Laczkovich (1988, Problem 1). (See also Problem 3.21 of Gibson and Natkaniec (1996–97).) © 2000 Elsevier Science B.V. All rights reserved.

*Keywords:* Extendable; Almost continuous; Connectivity; Additive functions

*AMS classification:* Primary 26A15; 54C30, Secondary 26A21; 54C08

---

### 1. Introduction

The study of different generalizations of continuity of functions from  $\mathbb{R}^n$  into  $\mathbb{R}$  has a long history. In this paper we will be interested in the functions with some of these generalized continuities that are known under the common name of Darboux-like functions. The readers unfamiliar with their definitions can find them in the next section.

The basic relations between these classes, for the functions from  $\mathbb{R}$  to  $\mathbb{R}$ , are given in the following chart, in which arrows  $\rightarrow$  denote strict inclusions. Moreover, all other possible

---

<sup>☆</sup> This work was partially supported by NSF Cooperative Research Grant INT-9600548 with its Polish part being financed by Polish Academy of Science PAN. The results were obtained during a visit of the second author at West Virginia University.

\* Corresponding author. Email: [KCies@wvnmvms.wvnet.edu](mailto:KCies@wvnmvms.wvnet.edu).

<sup>1</sup> Email: [jjas@ksinet.univ.gda.pl](mailto:jjas@ksinet.univ.gda.pl).

“natural intersection” inclusions (in a form of  $AC \cap CIVP \subseteq Conn \cap CIVP$ ) obtained from different classes of this chart remain strict.




Chart 1.

The inclusions  $C \subset Ext$ ,  $Conn \subset D \subset PC$ ,  $SCIVP \subset CIVP \subset WCIVP$ , and  $PR \subset PC$  are obvious from the definitions. The inclusions  $Ext \subset AC \subset Conn$  were proved by Stallings [27]. The inclusion  $CIVP \subset PR$  was stated without the proof in [11]. The proof can be found in [10, Theorem 3.8]. The inclusion  $Ext \subset SCIVP$  was proved by Rosen, Gibson, and Roush in [26]. An excellent discussion of this chart can be found in a recent survey by Gibson and Natkaniec [10, Section 3]. The examples concerning the properness of all intersection inclusions can be also found, in stronger versions, in Theorems 1.1, 1.2, and 1.5 stated below. Also, function  $F$  from Corollary 3.3 is the first simple ZFC example of almost continuous SCIVP function which is not extendable.

For the functions from  $\mathbb{R}^n$  into  $\mathbb{R}$  with  $n > 1$  the classes from the lower part of Chart 1 are not defined. The relations between the classes in the upper part of the chart change to the following.




Chart 2.

The inclusions  $C(\mathbb{R}^n) \subset Ext(\mathbb{R}^n) \subset Conn(\mathbb{R}^n)$  are obvious from the definitions. The inclusion  $Conn(\mathbb{R}^n) \subset Ext(\mathbb{R}^n)$  was recently proved by Ciesielski, Natkaniec, and Wojciechowski [6]. The containment  $Conn(\mathbb{R}^n) \subset PC(\mathbb{R}^n)$  was proved by Hamilton [15] and Stallings [27], and the inclusion  $PC(\mathbb{R}^n) \subset Conn(\mathbb{R}^n)$  by Hagan [14]. (See also Whyburn [28] and [10, Theorem 8.1].) The relation  $Conn(\mathbb{R}^n) \subset AC(\mathbb{R}^n) \cap D(\mathbb{R}^n)$  was proved by Stallings [27]. The examples concerning the properness of the inclusions can be found, in the Baire class one, in Theorem 1.3.

The main goal of this paper is to discuss these two charts when we restrict the function in all these classes to the following four classes of functions: Baire one  $B_1$ , Baire two  $B_2$ , Borel  $\mathcal{B}or$ , and additive functions  $Add$ . Notice that an intersection of any two of these classes is trivial, since  $B_1 \subset B_2 \subset \mathcal{B}or$  and  $Add \cap \mathcal{B}or \subset C$ .


**Theorem 1.1.** *For the Baire one functions  $B_1$  from  $\mathbb{R}$  to  $\mathbb{R}$  the following holds.*

$$C \subsetneq Ext = AC = Conn = D = PC = SCIVP = CIVP = PR \subsetneq WCIVP.$$

**Proof.** The proof of the equation  $\text{Ext} \cap B_1 = \text{PC} \cap B_1$  can be found in [3]. This equation and Chart 1 imply all the other equations. The properness of the inclusions is justified as follows.

- $B_1 \cap D \setminus C \neq \emptyset$ . It is witnessed by the function  $f_0: \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f_0(x) = \sin(1/x)$  for  $x \neq 0$  and  $f_0(0) = 0$ . (See [22, Example 1.1].)
- $B_1 \cap \text{WCIVP} \setminus \text{PC} \neq \emptyset$ . It is witnessed by the function  $g: \mathbb{R} \rightarrow \mathbb{R}$  defined by  $g(x) = x^2$  for  $x \neq 0$  and  $g(0) = -1$ .  $\square$

**Theorem 1.2.** *The classes from Chart 1 restricted to either Baire two functions  $B_2$  or Borel functions from  $\mathbb{R}$  to  $\mathbb{R}$  leads to the following chart. Moreover, all possible “natural intersection” inclusions obtained from different classes of this chart remain strict.*



**Proof.** To see that the inclusion  $D \subseteq \text{SCIVP}$  holds in the class of Borel functions let  $x < y$  and  $K$  be a perfect set between  $f(x)$  and  $f(y)$ . Since  $f$  is Darboux,  $f^{-1}(K) \cap (x, y)$  is an uncountable Borel set. Thus, it contains a perfect subset  $C_0$ . Moreover, we can find a perfect set  $C \subset C_0$  for which  $f \upharpoonright C$  is continuous. Similarly we can argue that the inclusion  $\text{CIVP} \subseteq \text{SCIVP}$  holds in the class of Borel functions.

The properness of the inclusions is justified as follows.

- $B_2 \cap \text{Ext} \setminus C \neq \emptyset$ . See Theorem 1.1.
- $B_2 \cap \text{AC} \setminus \text{Ext} \neq \emptyset$ . See Corollary 3.3.
- $B_2 \cap \text{Conn} \setminus \text{AC} \neq \emptyset$ . See Brown [2] or Jastrzębski [17].
- $B_2 \cap D \setminus \text{Conn} \neq \emptyset$ . See Brown [2].
- $B_2 \cap \text{SCIVP} \setminus D \neq \emptyset$ . See Example 3.5.
- $B_2 \cap \text{WCIVP} \cap \text{PR} \setminus \text{CIVP} \neq \emptyset$ . Let  $\{F_q: q \in \mathbb{Q}\}$  be a family of pairwise disjoint  $c$ -dense  $F_\sigma$  sets. Then  $f = \sum_{q \in \mathbb{Q} \cap (0,1)} q \chi_{F_q}$  is as desired.
- $B_2 \cap \text{WCIVP} \cap \text{PC} \setminus \text{PR} \neq \emptyset$ . It is witnessed by  $g = f + 2\chi_D$ , where  $f$  is as above and  $D$  is a countable dense subset of  $F_2$ .
- $B_2 \cap \text{WCIVP} \setminus \text{PC} \neq \emptyset$ . See Theorem 1.1.  $\square$

Restricting functions from Chart 2 to Baire one functions has a lot simpler solution.

**Theorem 1.3.** *For the Baire one functions  $B_1(\mathbb{R}^n)$  from  $\mathbb{R}^n$  to  $\mathbb{R}$ ,  $n > 1$ , Chart 2 remains unchanged.*


**Proof.** The properness of all the inclusions, as well as of their other possible combinations, is justified by the following facts.

- $B_1(\mathbb{R}^n) \cap \text{Conn}(\mathbb{R}^n) \setminus C(\mathbb{R}^n) \neq \emptyset$ . It is witnessed by a function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  given by  $f(x, y) = \sin(1/(x^2 + y^2))$  for  $\langle x, y \rangle \neq \langle 0, 0 \rangle$  and  $f(0, 0) = 0$ . It is in  $\text{PC}(\mathbb{R}^2)$  straight from the definition.

- $B_1(\mathbb{R}^n) \cap AC(\mathbb{R}^n) \cap D(\mathbb{R}^n) \setminus \text{Conn}(\mathbb{R}^n) \neq \emptyset$ . Rosen, Gibson and Roush [26, Example 1] proved that a function  $f : [-1, 1] \times [0, 1] \rightarrow [-1, 1]$  given by  $f(x, y) = \sin(1/y)$  for  $y > 0$  and  $f(x, 0) = x$  is Baire one, almost continuous, Darboux, but not connectivity. It is easy to extend it to a finite support function on  $\mathbb{R}^2$  with the same properties.
- $B_1(\mathbb{R}^n) \cap D(\mathbb{R}^n) \setminus AC(\mathbb{R}^n) \neq \emptyset$ . It is justified by the function  $F(x, y) = f_0(x)$ , where  $f_0$  is a function from Theorem 1.1. (See Natkaniec [22, Example 1.7] or [23, Example 1.1.9].)
- $B_1(\mathbb{R}^n) \cap AC(\mathbb{R}^n) \setminus D(\mathbb{R}^n) \neq \emptyset$ . Let  $f(x) = \sin(1/x)$  for  $x \neq 0$  and  $f(0) = 1$ , and let  $F : [-1, 1]^2 \rightarrow [-1, 1]$  be given by the formula  $F(x, y) = yf(x)$ . It was proved by Natkaniec [23, Example 1.1.10] that  $F$  is Baire one, almost continuous, and not Darboux. It is easy to extend  $F$  to a function  $\bar{F} : \mathbb{R}^2 \rightarrow [0, 1]$  with compact support while preserving these properties.  $\square$

The study of classes of additive functions from Charts 1 and 2 were initiated by Banaszewski [1]. (See also [10, Section 5].) In this direction we have the following results.

**Theorem 1.4.** *For the additive functions  $\text{Add}(\mathbb{R}^n)$  from  $\mathbb{R}^n$  to  $\mathbb{R}$ ,  $n > 1$ , Chart 2 changes as follows:*



**Proof.** The inclusion  $\text{Add}(\mathbb{R}^n) \cap AC(\mathbb{R}^n) \cap D(\mathbb{R}^n) \subset C(\mathbb{R}^n)$  is proved in Theorem 4.8. The properness of the inclusions is justified by the following facts.

- $\text{Add}(\mathbb{R}^n) \cap AC(\mathbb{R}^n) \setminus D(\mathbb{R}^n) \neq \emptyset$ . See Example 4.9.
- $\text{Add}(\mathbb{R}^n) \cap D(\mathbb{R}^n) \setminus AC(\mathbb{R}^n) \neq \emptyset$ . See Example 4.10.  $\square$

**Theorem 1.5.** *For the additive functions  $\text{Add}$  from  $\mathbb{R}$  to  $\mathbb{R}$  we have the equation  $\text{PR} = \text{WCIVP}$ . The other inclusions of Chart 1 remain unchanged, except possibly for the inclusion  $AC \subset \text{Conn}$ . Thus, we have*



Moreover all possible “natural intersection” inclusions obtained from the different classes of this chart and not involving  $AC \subset \text{Conn}$  remain strict. The inclusion  $\text{Add} \cap \text{CIVP} \cap AC \subset \text{Conn}$  is strict if union of less than continuum many meager subsets of  $\mathbb{R}$  is meager in  $\mathbb{R}$ .

**Proof.** The properness of all the inclusions is justified by the following facts.

- $\text{Add} \cap \text{Ext} \setminus C \neq \emptyset$ . See Corollary 4.4.

- $\text{Add} \cap \text{SCIVP} \cap \text{AC} \setminus \text{Ext} \neq \emptyset$ . See Ciesielski and Rosłanowski [8]. Compare also Ciesielski [5, Theorem 3.1].
- $\text{Add} \cap \text{SCIVP} \cap \text{D} \setminus \text{Conn} \neq \emptyset$ . See Example 5.3. (An example of a function from  $\text{Add} \cap \text{D} \setminus \text{Conn}$  was earlier given in [1].)
- $\text{Add} \cap \text{PC} \setminus \text{D} \neq \emptyset$ . See Example 5.2. (An example of a function from  $\text{Add} \cap \text{CIVP} \setminus \text{D}$  was earlier given in [1].)
- $\text{Add} \cap \text{AC} \cap \text{CIVP} \setminus \text{SCIVP} \neq \emptyset$ . See Ciesielski [5, Theorem 4.1].
- $\text{Add} \cap \text{AC} \cap \text{PR} \setminus \text{CIVP} \neq \emptyset$ . See Example 5.1.
- $\text{Add} \cap \text{AC} \setminus \text{PR} \neq \emptyset$ . See Banaszewski [1].
- $\text{Add} \cap \text{CIVP} \cap \text{Conn} \setminus \text{AC}$ . Such a function, under the assumption that union of less than continuum many meager subsets of  $\mathbb{R}$  is meager, has been constructed by Ciesielski and Rosłanowski [8]. (Example for  $\text{Add} \cap \text{Conn} \setminus \text{AC}$  requires only that union of less than continuum many meager subsets of  $\mathbb{R}$  does not cover  $\mathbb{R}$ .)  $\square$

The following questions, which are variants of Banaszewski's question [10, Question 5.5], remain open.

**Problem 1.1.**

- (1) Does there exist a ZFC example of an additive connectivity function  $f: \mathbb{R} \rightarrow \mathbb{R}$  which is not almost continuous?
- (2) Does there exist an additive connectivity function  $f: \mathbb{R} \rightarrow \mathbb{R}$  with SCIVP property which is not almost continuous?

**2. Definitions and notation**

Our terminology is standard and follows [4]. We consider only real-valued functions of one or more real variables. No distinction is made between a function and its graph. A restriction of a function  $f: X \rightarrow Y$  to a set  $A \subset X$  is denoted by  $f \upharpoonright A$ . Symbol  $\chi_A$  will be used for a characteristic function of a subset  $A$  of a fixed space  $X$ . By  $\mathbb{R}$  and  $\mathbb{Q}$  we denote the set of all real and rational numbers, respectively. We will consider  $\mathbb{R}^n$  as linear spaces over  $\mathbb{Q}$ . In particular, for  $X \subset \mathbb{R}^n$  we will use the symbol  $\text{LIN}_{\mathbb{Q}}(X)$  to denote the smallest linear subspace of  $\mathbb{R}^n$  over  $\mathbb{Q}$  that contains  $X$ . Recall also that if  $D \subset \mathbb{R}^n$  is linearly independent over  $\mathbb{Q}$  and  $f: D \rightarrow \mathbb{R}$  then

$$F = \text{LIN}_{\mathbb{Q}}(f) \subset \mathbb{R}^{n+1}$$

is an additive function (see definition below) from  $\text{LIN}_{\mathbb{Q}}(D)$  into  $\mathbb{R}$  which extends  $f$ . Any linear basis of  $\mathbb{R}$  over  $\mathbb{Q}$  will be referred as a *Hamel basis*.

The ordinal numbers will be identified with the sets of all their predecessors and cardinals with the initial ordinals. In particular  $2 = \{0, 1\}$  and the first infinite ordinal  $\omega$  number is equal to the set of all natural numbers  $\{0, 1, 2, \dots\}$ . The family of all functions from a set  $X$  into  $Y$  is denoted by  $Y^X$ . In particular  $2^n$  will stand for the set of all sequences  $s: \{0, 1, 2, \dots, n-1\} \rightarrow \{0, 1\}$ , while  $2^{<\omega} = \bigcup_{n < \omega} 2^n$  is the set of all finite sequences into 2. The symbol  $|X|$  stands for the cardinality of a set  $X$ . The cardinality of  $\mathbb{R}$  is denoted by

$c$  and referred as *continuum*. A set  $S \subset \mathbb{R}$  is said to be *c-dense* if  $|S \cap (a, b)| = c$  for every  $a < b$ . The closure of a set  $A \subset \mathbb{R}^n$  is denoted by  $\text{cl}(A)$ , its boundary by  $\text{bd}(A)$ , and its diameter by  $\text{diam}(A)$ .

We will use also the following terminology [10]. For  $X \subseteq \mathbb{R}^n$  a function  $f : X \rightarrow \mathbb{R}$  is:

- *additive* if  $X$  is closed under the addition and  $f(x + y) = f(x) + f(y)$  for every  $x, y \in X$ ;
- *Darboux* if  $f[K]$  is a connected subset of  $\mathbb{R}$  (i.e., an interval) for every connected subset  $K$  of  $X$ ;
- *almost continuous* (in sense of Stallings) if each open subset of  $X \times \mathbb{R}$  containing the graph of  $f$  contains also a continuous function from  $X$  to  $\mathbb{R}$  [27];
- *connectivity* function if the graph of  $f \upharpoonright Z$  is connected in  $Z \times \mathbb{R}$  for any connected subset  $Z$  of  $X$ ;
- *extendability* function provided there exists a connectivity function  $F : X \times [0, 1] \rightarrow \mathbb{R}$  such that  $f(x) = F(x, 0)$  for every  $x \in X$ ;
- *peripherally continuous* if for every  $x \in X$  and for all pairs of open sets  $U$  and  $V$  containing  $x$  and  $f(x)$ , respectively, there exists an open subset  $W$  of  $U$  such that  $x \in W$  and  $f[\text{bd}(W)] \subset V$ .

The classes of these functions are denoted by  $\text{Add}(X)$ ,  $\text{D}(X)$ ,  $\text{AC}(X)$ ,  $\text{Conn}(X)$ ,  $\text{Ext}(X)$ , and  $\text{PC}(X)$ , respectively. The class of continuous functions from  $X$  into  $\mathbb{R}$  is denoted by  $\text{C}(X)$ . We will drop the index  $X$  if  $X = \mathbb{R}$ .

Recall also that if the graph of  $f : \mathbb{R} \rightarrow \mathbb{R}$  intersects every closed subset  $B$  of  $\mathbb{R}^2$  which projection  $\text{pr}(B)$  onto the  $x$ -axis has nonempty interior then  $f$  is almost continuous. (See, e.g., [22].) Similarly, if the graph of  $f : \mathbb{R} \rightarrow \mathbb{R}$  intersects every compact connected subset  $B$  of  $\mathbb{R}^2$  with  $|\text{pr}(B)| > 1$  then  $f$  is connectivity. This follows from the following well-known fact.

**Fact 2.1.** *If  $S \subset \mathbb{R}^2$  disconnects  $\mathbb{R}^2$  then it contains a nontrivial compact connected subset.*

A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has:

- *Cantor intermediate value property* if for every  $x, y \in \mathbb{R}$  and for each perfect set  $K$  between  $f(x)$  and  $f(y)$  there is a perfect set  $C$  between  $x$  and  $y$  such that  $f[C] \subset K$ ;
- *strong Cantor intermediate value property* if for every  $x, y \in \mathbb{R}$  and for each perfect set  $K$  between  $f(x)$  and  $f(y)$  there is a perfect set  $C$  between  $x$  and  $y$  such that  $f[C] \subset K$  and  $f \upharpoonright C$  is continuous;
- *weak Cantor intermediate value property* if for every  $x, y \in \mathbb{R}$  with  $f(x) < f(y)$  there exists a perfect set  $C$  between  $x$  and  $y$  such that  $f[C] \subset (f(x), f(y))$ ;
- *perfect road* if for every  $x \in \mathbb{R}$  there exists a perfect set  $P \subset \mathbb{R}$  having  $x$  as a bilateral (i.e., two-sided) limit point for which  $f \upharpoonright P$  is continuous at  $x$ .

The above classes of these functions are denoted by  $\text{CIVP}$ ,  $\text{SCIVP}$ ,  $\text{WCIVP}$ , and  $\text{PR}$ , respectively.

### 3. Almost continuous Baire two class function which is not extendable

The main example described in this section answers Problem 2 and the main part of Problem 1 from [3], as well as Problem 3.21 from [10].

Let  $C \subset [0, 1]$  be the ternary Cantor set and let  $\mathcal{J}$  be the family of all component intervals of  $[0, 1] \setminus C$ . We put

$$\mathcal{J}_0 = \{J \in \mathcal{J} : \text{the length of } J \text{ is } 3^{-n} \text{ with } n < \omega \text{ even}\}$$

and

$$\mathcal{J}_1 = \{J \in \mathcal{J} : \text{the length of } J \text{ is } 3^{-n} \text{ with } n < \omega \text{ odd}\}.$$

Let  $\{(a_n, b_n) : n < \omega\}$  and  $\{(c_n, d_n) : n < \omega\}$  be the enumerations of  $\mathcal{J}_0$  and  $\mathcal{J}_1$ , respectively. Define function  $f : [0, 1] \rightarrow [0, 1]$  in the following way:

- for every  $n < \omega$  we put  $f(a_n) = f(d_n) = 0$ ,  $f(b_n) = f(c_n) = 1$ , and extend it linearly on  $[a_n, b_n]$  and  $[c_n, d_n]$ ;
- for all other  $x$ 's we put  $f(x) = 0$ .

**Theorem 3.1.** *The function  $f$  is almost continuous, Baire class two, but not extendable.*

**Proof.** We will start with showing that  $f$  is not extendable. By way of contradiction, assume that  $f$  can be extended, that is, that there is a connectivity function  $F : [0, 1]^2 \rightarrow [0, 1]$  with  $F(x, 0) = f(x)$  for all  $x \in [0, 1]$ . Thus  $F$  is peripherally continuous. We will deduce from this that there exists a perfect set  $P \subset C \times \{0\}$  on which  $F$  is constantly equal to 1, which evidently contradicts our definition of  $f$ .

We will define the following families:  $\langle p_s \in C : s \in 2^{<\omega} \rangle$ ,  $\langle B_s \subset [0, 1]^2 : s \in 2^{<\omega} \rangle$ , and  $\langle U_s \subset [0, 1]^2 : s \in 2^{<\omega} \rangle$  such that the following conditions hold for every  $s \in 2^n$  and different  $t, t' \in 2^{n+1}$  extending  $s$ .

- (i)  $\langle p_s, 0 \rangle \in U_s$ ,  $f(p_s) = 1$ , and  $U_s$  is open with a diameter at most  $2^{-n}$ .
- (ii)  $B_s$  is closed, connected, and  $F[B_s] \subset [1 - 2^{-n}, 1]$ .
- (iii)  $\text{cl}(U_t) \cup B_t \subset U_s$ ,  $\langle p_s, 0 \rangle \notin U_t$ , and  $B_t \cap B_s \neq \emptyset$ .
- (iv)  $\text{cl}(U_t) \cap \text{cl}(U_{t'}) = \emptyset$ .

For  $s = \emptyset \in 2^0$  we put  $U_s = B_s = [0, 1]^2$  and choose an arbitrary  $p_s$  with  $f(p_s) = 1$ . Farther, the construction goes by the induction on the length  $n$  of  $s \in 2^{<\omega}$ . Thus, assume that for some  $s \in 2^n$  the sets  $B_s$ ,  $U_s$ , and the point  $p_s$  are already chosen. Let  $t$  and  $t'$  be different sequences from  $2^{n+1}$  extending  $s$ . To choose  $B_t$ ,  $p_t$ , and  $U_t$  we proceed as follows.

Note that  $p_s$  is an endpoint of some  $J \in \mathcal{J}$  since  $f(p_s) = 1$ . If  $p_s$  is a left endpoint of  $J$  we put  $I = [0, p_s)$ . Otherwise we put  $I = (p_s, 1]$ . Choose an  $\varepsilon > 0$  less than the diameters of  $I$  and  $B_s$  and such that

$$\begin{aligned} &\text{for every } J \in \mathcal{J} \text{ with } J \subset I \text{ if the distance from } J \text{ to } p_s \text{ is less than } \varepsilon \\ &\text{then } J \times \{0\} \subset U_s. \end{aligned} \tag{*}$$

Let  $W_s \subset [0, 1]^2$  be an open neighborhood of  $\langle p_s, 0 \rangle$  with diameter less than  $\varepsilon$  and such that  $F[\text{bd}(W_s)] \subset [1 - 2^{-n}, 1]$ . It exists since  $F$  is peripherally continuous at  $\langle p_s, 0 \rangle$  and

$F(p_s, 0) = f(p_s) = 1$ . Without loss of generality we can assume that  $\text{bd}(W_s)$  is connected. (Replacing  $W_s$  by its component, if necessary, we can assume that  $W_s$  is connected. Then we can increase  $W_s$  to  $[0, 1]^2 \setminus V_s$ , where  $V_s$  is an “unbounded” component of  $[0, 1]^2 \setminus W_s$ , that is the one which contains the boundary of  $[0, 1]^2$ . This decreases  $\text{bd}(W_s)$  and makes it connected.) Note that the choice of  $\varepsilon$  guarantees that  $\text{bd}(W_s) \cap B_s \neq \emptyset \neq \text{bd}(W_s) \cap (I \times \{0\})$  since  $B_s$  and  $I \times \{0\}$  are connected and  $\text{bd}(W_s)$  disconnects  $[0, 1]^2$ . Let  $z \in I$  be such that  $\langle z, 0 \rangle \in \text{bd}(W_s) \cap (I \times \{0\})$ . Since  $F(z, 0) \in F[\text{bd}(W_s)] \subset [1 - 2^{-n}, 1] \subset (0, 1]$  there exists a  $J \in \mathcal{J}$  such that  $z \in \text{cl}(J)$ . Let  $p_t$  be the endpoint of  $\text{cl}(J)$  for which  $f(p_t) = 1$ . By (\*) we have  $\langle p_t, 0 \rangle \in U_s$ . The set  $B_t$  is defined as a union of  $\text{bd}(W_s)$  and a closed segment joining  $\langle z, 0 \rangle$  and  $\langle p_t, 0 \rangle$ . The open neighborhood  $U_t$  of  $p_t$  is chosen such that its diameter is at most  $2^{-(n+1)}$  and that  $p_s \notin \text{cl}(U_t) \subset U_s$ . It is easy that conditions (i)–(iii) are satisfied.

To choose  $B_{t'}$ ,  $p_{t'}$ , and  $U_{t'}$  we replace  $U_s$  with  $U'_s = U_s \setminus \text{cl}(U_t)$  and repeat the process described above. This finishes the inductive construction.

Now, to finish the argument take an arbitrary  $s \in 2^\omega$  and note that by (i) and (iii) the limit  $\lim_{n \rightarrow \infty} \langle p_s \upharpoonright n, 0 \rangle$  exists and is equal to a point  $p_s$  which belongs to  $C \times \{0\}$ . Also, by (ii), the set

$$B_s = \bigcup_{0 < n < \omega} B_s \upharpoonright n$$

is connected and  $p_s$  is its only accumulation point. Since  $P = \{p_s : s \in 2^\omega\}$  is evidently equal to a perfect set  $\bigcap_{n < \omega} \bigcup \{\text{cl}(U_t) : t \in 2^n\}$  it is enough to prove that  $F(p_s) = 1$  for every  $s \in 2^\omega$ . But if  $F(p_s) \neq 1$  then  $F(p_s) = 0$ , since  $p_s \in C \times \{0\}$ . Take  $\varepsilon > 0$  is less than the diameter of  $B_s$  and let  $U$  be an open neighborhood of  $p_s$  of diameter less than  $\varepsilon$  and such that  $F[\text{bd}(U)] \subset [0, 1/2)$ . Then for a point  $w \in B_s \cap \text{bd}(U)$  we have  $F(w) \in [0, 1/2) \cap [1/2, 1]$ , a contradiction. This finishes the proof that  $f$  is not extendable.

Next we will show that  $f$  is almost continuous. Let  $G$  be an open set contained in  $[0, 1]^2$  containing the graph of  $f$ . For every  $x \in [0, 1]$  there exists an interval  $(a_x, b_x)$  such that

- $x \in (a_x, b_x)$ ,
- $f(a_x) = f(b_x) = 0$ , and
- there is a continuous function  $g_x : [0, 1] \rightarrow \mathbb{R}$  with  $g_x \upharpoonright [a_x, b_x] \subset G$  and such that  $g_x(t) = 0$  for  $t \notin (a_x, b_x)$ .

Indeed, if  $f(x) = 0$  then it is easy to find  $(a_x, b_x)$  for which  $g_x \equiv 0$  works. If  $f(x) \neq 0$  then  $x \in J$  for some  $J \in \mathcal{J}$ , say  $J = (A, B)$ . Assume that  $f$  is increasing on  $J$ , the other case is similar. Then  $f(B) = 1$ . Find an interval  $J' = (C, D) \in \mathcal{J}$  on which function  $f$  is decreasing and such that  $[B, C] \times \{1\} \subset G$ . Put  $a_x = A$ ,  $b_x = D$ , define  $g_x(B) = g_x(C) = 1$ ,  $g_x(0) = g_x(a_x) = g_x(b_x) = g_x(1) = 0$ , and extend  $g_x$  in a linear way on each interval with these endpoints. Thus  $g_x$  has a hat shape.

Now choose a finite subcover  $\{(a_{x_i}, b_{x_i}) : i < n\}$ , with  $n < \omega$ , of the cover  $\{(a_x, b_x) : x \in [0, 1]\}$  of the interval  $[0, 1]$ . Then the function

$$g(x) = \max\{g_{x_i}(x) : i < n\}$$

is continuous and  $g \subset G$ . This ends the proof that  $f$  is almost continuous.



To see that  $f$  is of Baire class two it is enough to notice that the preimage of every open (and closed) set  $U$  is a countable union of  $F_\sigma$  (closed) sets  $f^{-1}(U) \cap \text{cl}(J)$  with  $J \in \mathcal{J}$  and, possibly, of a  $G_\delta$  set  $C \setminus \bigcup \{\text{cl}(J) : J \in \mathcal{J}\}$ .  $\square$

In [3, Problem 1] the authors asked also whether the function as above can be in the class  $J_1$  of all functions (from  $\mathbb{R}$  or  $[0, 1]$  into  $\mathbb{R}$ ) that are pointwise limits of functions which have only discontinuities of the first class, that is, these functions for which both one sided limits exist at each point. Clearly  $B_1 \subset J_1 \subset B_2$ . Our function gives also an answer to this question.

**Theorem 3.2.** *There exists a  $J_1$  function  $f : [0, 1] \rightarrow [0, 1]$  which is almost continuous but not extendable.*

**Proof.** Let  $f$  be a function from Theorem 3.1 and let  $\chi_D$  stand for a characteristic function of  $D \subset [0, 1]$ . If  $\{J_n : n < \omega\}$  is an enumeration of  $\{\text{cl}(J) : J \in \mathcal{J}\}$  and  $D_n = \bigcup_{i < n} J_i$  then  $f$  is a pointwise limit of functions  $f \chi_{D_n}$ . Thus,  $f$  is in  $J_1$ .  $\square$

**Corollary 3.3.** *There exists a  $J_1$  function  $F : \mathbb{R} \rightarrow \mathbb{R}$  which is almost continuous but not extendable.*

**Proof.** Extend the function  $f$  from Theorem 3.2 to  $F$  by putting  $F(x) = 0$  for all  $x \in \mathbb{R} \setminus [0, 1]$ .  $\square$

The main core of the proof that the function  $f$  from Theorem 3.1 is not extendable is that the set  $f^{-1}(1)$  is countable. The next proposition shows that it is essentially the only obstacle for  $f$  to be extendable, in a sense that we can redefine  $f$  on a subset of  $C$  to get an extendable function.

**Proposition 3.4.** *If  $f$  is from Theorem 3.1 then there exists a meager  $F_\sigma$  subset  $B$  of  $C_0 = C \setminus \bigcup \{\text{cl}(J) : J \in \mathcal{J}\}$  such that  $f_0 = f + \chi_B$  is extendable.*

**Proof.** Gibson and Roush [12] proved that a function  $g : [0, 1] \rightarrow [0, 1]$  is extendable if and only if there exists a sequence  $\langle \langle I_n, J_n \rangle : n < \omega \rangle$  of pairs of open intervals, called a *PI family* for  $g$ , such that

- (a)  $\lim_{n \rightarrow \infty} \text{diam}(I_n) = 0$ ,
- (b)  $g[\text{bd}(I_n)] \subset J_n$  for every  $n < \omega$ ,
- (c) for every  $x \in [0, 1]$  and  $\varepsilon > 0$  there exists an  $n < \omega$  such that  $x \in I_n$ ,  $g(x) \in J_n$ , and  $\max\{\text{diam}(I_n), \text{diam}(J_n)\} < \varepsilon$ ,
- (d) for every  $n < m < \omega$  if the sets  $I_n \cap I_m$ ,  $I_n \setminus I_m$ , and  $I_m \setminus I_n$  are nonempty then  $J_n \cap J_m \neq \emptyset$ .

It was also noticed in [6] that in (c) it is enough to consider only discontinuity points  $x$  of  $g$ . In what follows we will construct a PI family for our future  $f_0$ .

Let  $T$  be the set of all endpoints of all intervals  $J \in \mathcal{J}$  and let  $\{t_n: n < \omega\}$  be an enumeration of  $T$ . We construct by induction on  $n < \omega$  the finite families  $F_n$  of triples  $\langle t, I, J \rangle$  such that

- (1)  $I$  and  $J$  are open intervals in  $[0, 1]$ ,  $t \in T \cap I$ , and  $f(t) \in J$ ;
- (2)  $f[\text{bd}(I)] \subset (0, 1) \cap J$  and  $\text{diam}(J) \leq 2^{-n}$ ;
- (3)  $I \cap I' = \emptyset$  for different  $\langle t, I, J \rangle, \langle t', I', J' \rangle \in F_n$ ;
- (4)  $C \subset \bigcup \{I: \langle t, I, J \rangle \in F_n\}$  and  $\{t_i: i < n\} \subset \{t: \langle t, I, J \rangle \in F_n\}$ ;
- (5) if  $n > 0$  then for every  $\langle t, I, J \rangle \in F_n$  there exists  $\langle t_0, I_0, J_0 \rangle \in F_{n-1}$  such that  $I \subset I_0$ .

The induction can be started with  $F_0 = \{\langle t_0, (0, 1), (0, 1) \rangle\}$ , and can be easily carried through since  $C$  is compact zero-dimensional and the sets  $f^{-1}(1)$  and  $T \cap f^{-1}(0)$  are dense in  $C$ .

Now, let  $\{\langle t_n, I_n, J_n \rangle: n < \omega\}$  be an enumeration of  $\bigcup_{n < \omega} F_n$ . We claim that the sequence  $\langle \langle I_n, J_n \rangle: n < \omega \rangle$  is a PI family for  $f_0 = f + \chi_B$  for an appropriately chosen set  $B$ .

Clearly (3) and (5) imply that condition (d) is satisfied in void. Condition (a) can be deduced from the density of  $T$  in  $C$  and (1)–(3). (b) For  $g = f_0$  is implied by (2) and the fact that  $f_0(x) = f(x)$  for  $x \in [0, 1] \setminus C$ . Similarly (c) for the points  $x \in T$  is implied by (4) and  $f_0 \upharpoonright T = f \upharpoonright T$ . To finish the proof it is enough to show that (c) holds for points  $x \in C_0$  for an appropriate choice of  $B$ . But every  $x \in C_0$  the set  $S_x = \{k < \omega: x \in I_k\}$  is infinite. Let  $B$  be the set of all those points  $x \in C_0$  for which the set  $\{k \in S_x: f(t_k) = 0\}$  is finite. Then for every  $x \in C_0$  the set  $\{k \in S_x: f_0(x) \in J_k\}$  is infinite proving (b).

The fact that  $B$  is a meager  $F_\sigma$  subset of  $C_0$  is left as an exercise.  $\square$

**Example 3.5.** There exists an  $f \in B_2 \cap \text{SCIVP} \setminus D$ .

**Proof.** Define  $g$  from  $[-1, 1]$  onto  $(0, 1]$  by  $g(x) = (x^2 - 1) \sin^2(1/x) + 1$  for  $x \neq 0$  and  $f(0) = 1$ . Note that  $g(-1) = g(1) = 1$  and that  $g$  is SCIVP. For each component  $J$  of  $[0, 1] \setminus C$  of length  $1/3^{-n}$  define  $f \upharpoonright \text{cl}(J)$  as  $(-1)^n g \circ h_J$ , where  $h_J$  is an increasing linear function with  $h_J[\text{cl}(J)] = [-1, 1]$ . For all other points  $x$  we put  $f(x) = 1$ . Note that  $f[\mathbb{R}] = [-1, 0) \cup (0, 1]$  and that preimage of every open set is a union of a  $G_\delta$  and an  $F_\sigma$  set. So  $f$  is Baire two and not Darboux. To see that it is SCIVP take  $x < y$  and a perfect set  $K$  between  $f(x)$  and  $f(y)$ . We have to find a perfect set  $P \subset (x, y)$  on which  $f$  is continuous and with  $f[P] \subset K$ . If both  $x$  and  $y$  belong to the closure of the same component of  $C$  then the existence of  $K$  follows from SCIVP property of  $g$ . Otherwise there exist two components  $J_0$  and  $J_1$  of  $C$  between  $x$  and  $y$  with  $f[J_0 \cup J_1] = [-1, 0) \cup (0, 1]$ . This we can choose appropriate  $K \subset J_0 \cup J_1$ .  $\square$

#### 4. Additive extendable discontinuous function

Let  $h \in \text{Ext}(\mathbb{R})$ . We say that a set  $G \subset \mathbb{R}$  is  $h$ -negligible for a function  $h$  provided  $f \in \text{Ext}$  for every function  $f: \mathbb{R} \rightarrow \mathbb{R}$  for which  $f = h$  on a set  $\mathbb{R} \setminus G$ .

**Lemma 4.1** (Rosen [25], Ciesielski and Reclaw [7]). *There exists an extendable function  $h : \mathbb{R} \rightarrow \mathbb{R}$  such that some dense  $G_\delta$ -set  $G \subset \mathbb{R}$  is  $h$ -negligible.*

**Lemma 4.2.** *If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is homeomorphism,  $h \in \text{Ext}$ , and  $G$  is  $h$ -negligible then  $h \circ g^{-1} \in \text{Ext}$  and  $g[G]$  is  $(h \circ g^{-1})$ -negligible.*

**Proof.** This is a simple corollary from [18, Lemma 2.2]. (See also [24].)  $\square$

**Proposition 4.3.** *For every  $\mathfrak{c}$ -dense meager  $F_\sigma$ -set  $F \subset \mathbb{R}$  there exists an extendable function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $\mathbb{R} \setminus F$  is  $f$ -negligible.*

**Proof.** Let  $h$  and  $G$  be as in Lemma 4.1. By Lemma 4 of [13] there exists a homeomorphism  $g : \mathbb{R} \rightarrow \mathbb{R}$  such that  $g[\mathbb{R} \setminus G] \subset F$ . Then  $\mathbb{R} \setminus F \subset g[G]$  so, by Lemma 4.2, function  $f = h \circ g^{-1}$  is extendable and the set  $\mathbb{R} \setminus F$  is  $f$ -negligible.  $\square$

**Corollary 4.4.** *There exists an additive extendable function  $g : \mathbb{R} \rightarrow \mathbb{R}$  with a dense graph. In particular  $g \in \text{Add} \cap \text{Ext} \setminus \mathcal{C}$ .*

**Proof.** Let  $F$  be  $\mathfrak{c}$ -dense meager  $F_\sigma$ -set which is linearly independent over  $\mathbb{Q}$ . Such a set can be easily constructed from a linearly independent perfect set, which description can be found in [20, Theorem 2, Chapter XI, Section 7]. By Proposition 4.3 there exists an  $f \in \text{Ext}$  such that  $\mathbb{R} \setminus F$  is  $f$ -negligible. In particular  $f \upharpoonright F$  must be discontinuous. Let  $g$  be a linear extension of  $f \upharpoonright F$ . Then  $g \in \text{Ext}$  since  $\mathbb{R} \setminus F$  is  $f$ -negligible. Clearly  $g$  is additive and discontinuous, so it has a dense graph.  $\square$

Next we prove that  $\text{Add}(\mathbb{R}^n) \cap \text{AC}(\mathbb{R}^n) \cap \text{D}(\mathbb{R}^n) \subset \mathcal{C}(\mathbb{R}^n)$  for  $n > 1$ . Its proof will be based on the following two propositions, the first of which was proved by Lipiński [19]. (See also Maliszewski and Natkaniec [21].) This fact was noticed independently by the authors of this paper and our proof is enclosed below.

**Proposition 4.5.** *If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is discontinuous then  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  given by  $F(x, y) = f(x)$  is not almost continuous.*

**Proof.** Assume that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is discontinuous at some point  $x$ . Taking a translation of a graph of  $f$ , if necessary, we can assume that  $x = 0$  and  $f(0) = 0$ . So, there exists a sequence  $\{x_n\}_{n < \omega}$  converging to 0 such that  $\lim_{n \rightarrow \infty} f(x_n) = L_0 \neq 0$ . Multiplying  $f$  by  $3/L_0$ , if necessary, we can assume that  $L_0 = 3$ . We will also assume that  $f(x_n) > 2$  for all  $n < \omega$ .

Consider the closed set

$$A = \{ \langle x, y, z \rangle : x = x_n \text{ for some } n < \omega, y \in \mathbb{R}, \text{ and } z \leq 2 \} \cup (\{0\} \times \mathbb{R}^2)$$

and let  $G = \mathbb{R}^3 \setminus A$ . Then  $G$  is an open set containing all the graph of  $F$  except of a line  $L = \{0\} \times \mathbb{R} \times \{0\}$ . Let

$$H = \{ \langle x, y, z \rangle \in \mathbb{R}^3 : x^2 + z^2 < e^{-y^2} \}.$$

Then  $H$  is an open and contains  $L$ , so  $U = G \cup H$  contains  $F$ . We will show that  $U$  does not contain a graph of a continuous function.

By way of contradiction assume that there exists a continuous  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  with  $g \subset U$ . Then  $\langle 0, 0, g(0, 0) \rangle \in H$ . In particular,  $g(0, 0) \in (-1, 1)$ . So, by the continuity of  $g$ , there exists an  $n < \omega$  such that  $g(x_n, 0) \in (-1, 1)$ . We claim that  $g \upharpoonright (\{x_n\} \times \mathbb{R})$  is discontinuous. Indeed, notice that

$$(\{x_n\} \times \mathbb{R}^2) \cap U \subset \{x_n\} \times (G_0 \cup H_0),$$

where  $G_0 = \mathbb{R} \times (2, \infty)$ ,  $H_0 = (-b, b) \times (-\infty, 1)$ , and  $b > 0$  is such that  $e^{-(|x_n|+b)^2} = 1$ . Moreover,  $\langle x_n, 0, g(x_n, 0) \rangle \in \{x_n\} \times H_0$ . But clearly there is no continuous function on  $\mathbb{R}$  whose graph is contained in  $G_0 \cup H_0$  and intersects  $H_0$ . This contradiction finishes the proof.  $\square$

Notice that Proposition 4.5 stay in contrast with the following fact.

**Fact 4.6** (Natkaniec [22, Corollary 4.2(1)]). *If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is almost continuous and  $Y$  is a compact topological space then  $F : \mathbb{R} \times Y \rightarrow \mathbb{R}$  given by  $F(x, y) = f(x)$  is almost continuous.*

**Proposition 4.7.** *If  $n > 1$  and  $F \in \text{Add}(\mathbb{R}^n) \cap \text{D}(\mathbb{R}^n)$  then  $F^{-1}(0)$  contains a straight line.*

**Proof.** If  $F$  is constantly equal to 0 then there is nothing to prove. So, assume that this is not the case. Then  $f[\mathbb{R}^2] = \mathbb{R}$ . In particular  $V = F^{-1}(0)$  disconnects  $\mathbb{R}^2$ . (Otherwise its complement  $\mathbb{R}^2 \setminus V$  would be connected, while  $F[\mathbb{R}^2 \setminus V] = \mathbb{R} \setminus \{0\}$  would not, contradicting Darboux property.) Thus, by Fact 2.1,  $V$  contains a nontrivial compact connected subset  $K$ . Pick  $a, b \in K$  with  $\text{diam}(\{a, b\}) = \text{diam}(K)$ . Since every rotation  $r$  is a linear homeomorphism, replacing  $F$  with  $F \circ r$  for an appropriate  $r$  if necessary, we can assume that  $a$  and  $b$  are on the same vertical line. If  $\text{pr}(K)$  is a singleton, then  $K$  contains a straight line segment connecting  $a$  with  $b$ . This, and the fact that  $V$  is linear over  $\mathbb{Q}$ , easily imply that  $V$  contains a straight line. So, assume that  $\text{pr}(K) = [x_0, x_1]$  for some  $x_0 < x_1$ . We claim that this implies that

$$\text{there exists a bounded open set } U \subset \mathbb{R}^2 \text{ with } \text{bd}(U) \subset V. \quad (*)$$

To see it take  $c, d \in K$  with  $\text{pr}(c) = x_0$  and  $\text{pr}(d) = x_1$ , and let  $[y_0, y_1]$  be a projection of  $K$  to the second coordinate. Let  $P = [x_0, x_1] \times [y_0, y_1]$ . Notice that  $K \subset P$  and that  $a$  and  $b$  lie on the opposite horizontal sides of  $P$ , and  $c$  and  $d$  on opposite vertical sides of it. Note also that vectors  $v = b - a$  and  $w = d - c$  belong to  $V$ . Now consider the parallelogram-like set

$$B = \bigcup_{m=0}^4 [mv + (K \cup (4w + K)) \cup mw + (K \cup (4v + K))] \subset V.$$

(The “sides” are formed from translated “roads” from  $a$  to  $b$  and from  $c$  to  $d$ .) Note that the interior  $U_0$  of  $2v + w + P$  is disjoint with  $B$  and that  $B$  separates it from the infinity. Then the component  $U$  of  $\mathbb{R}^2 \setminus B$  containing  $U_0$  satisfies (\*).

Now, as in [9, Lemma 5.4] we can find a nonempty bounded connected open set  $W \subset \mathbb{R}^2$  with connected boundary  $\text{bd}(W) \subset B \subset V$ . Take an  $\varepsilon > 0$  such that some open disk of radius  $\varepsilon$  is contained in  $B$ . Then, if  $B(\varepsilon) = \{v \in \mathbb{R}^2: \text{length of } v \text{ is less than } \varepsilon\}$  then for every  $v \in B(\varepsilon)$  we have  $\text{bd}(W) \cap (v + \text{bd}(W)) \neq \emptyset$ . So,  $v \in \text{bd}(W) - \text{bd}(W) \subset V$ . Therefore,  $B(\varepsilon) \subset V$ , and so  $V = \mathbb{R}^2$  contradicting our assumption that  $F$  is non-zero.  $\square$

**Theorem 4.8.**  $\text{Add}(\mathbb{R}^n) \cap \text{AC}(\mathbb{R}^n) \cap \text{D}(\mathbb{R}^n) \subset \text{C}(\mathbb{R}^n)$  for  $n > 1$ .

**Proof.** Let  $F \in \text{Add}(\mathbb{R}^n) \cap \text{AC}(\mathbb{R}^n) \cap \text{D}(\mathbb{R}^n)$ . We will show that it is continuous.

Since it is enough to prove continuity of any restriction of  $F$  to a plane containing the origin, we can assume that  $n = 2$ . Now, by Proposition 4.7,  $f^{-1}(0)$  contains a line  $L_0$ . Since  $f^{-1}(0)$  is close under addition, it contains also a parallel line  $L$  which contains the origin. Since all the classes under consideration are closed under inner composition with a rotation, we can assume that  $L$  is a vertical line. But this means that there exists additive  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $F(x, y) = f(x)$  for every  $x, y \in \mathbb{R}$ . So, by Proposition 4.5, function  $f$  is continuous. Thus  $F \in \text{C}(\mathbb{R}^2)$ .  $\square$

**Example 4.9.** There exists an  $f \in \text{Add}(\mathbb{R}^2) \cap \text{AC}(\mathbb{R}^2) \setminus \text{D}(\mathbb{R}^2)$ .

**Proof.** Let  $\mathcal{B}$  be a family of all closed subsets  $B$  of  $\mathbb{R}^2 \times \mathbb{R}$  for which the projection  $\text{pr}(B)$  onto first coordinate  $\mathbb{R}^2$  has cardinality  $\mathfrak{c}$ . (Thus, it contains a perfect set.) It is known that if  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  intersects every element of  $\mathcal{B}$  then  $f \in \text{AC}(\mathbb{R}^2)$ . (See [22, Proposition 1.2].) Let  $\{F_\xi: \xi < \mathfrak{c}\}$  be an enumeration of  $\mathcal{B}$ . By induction on  $\xi < \mathfrak{c}$  define a sequence  $\langle \langle a_\xi, b_\xi, y_\xi \rangle: \xi < \mathfrak{c} \rangle$  such that for every  $\xi < \mathfrak{c}$  the following conditions hold:

- (i)  $a_\xi \in \text{pr}(B_\xi) \setminus \text{LIN}_{\mathbb{Q}}(\{a_\zeta: \zeta < \xi\} \cup \{b_\zeta: \zeta < \xi\})$ ,
- (ii)  $b_\xi \in \text{pr}(B_\xi) \setminus \text{LIN}_{\mathbb{Q}}(\{a_\zeta: \zeta \leq \xi\} \cup \{b_\zeta: \zeta < \xi\})$ ,
- (iii)  $\langle a_\xi, y_\xi \rangle \in B_\xi$ .

Let  $A = \{a_\xi: \xi < \mathfrak{c}\}$ ,  $B = \{b_\xi: \xi < \mathfrak{c}\}$ , and  $H$  be a Hamel base containing  $A \cup B$ . Define  $f_0: H \rightarrow \mathbb{R}$  by putting  $f_0(a_\xi) = y_\xi$  and  $f_0(x) = 1$  for  $x \in H \setminus A$ . Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be a linear extension of  $f_0$ . Then  $f$  is additive and almost continuous, since it intersects every  $F \in \mathcal{F}$ . However, it is not Darboux, since the set  $D = B \cup \{0\}$  is connected, as it intersects every perfect subset of  $\mathbb{R}^2$ , while  $f[D] = \{0, 1\}$  is not connected.  $\square$

**Example 4.10.** There exists an  $f \in \text{Add}(\mathbb{R}^2) \cap \text{D}(\mathbb{R}^2) \setminus \text{AC}(\mathbb{R}^2)$ .

**Proof.** Take an additive Darboux function  $g: \mathbb{R} \rightarrow \mathbb{R}$  with the property that

$$g[(a, b)] = \mathbb{R} \quad \text{for every } a < b. \tag{*}$$

A function  $f$  from Example 5.3 has the property. Define  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  by  $f(x, y) = g(x)$ . Clearly  $f$  is additive and discontinuous. To see that  $f$  is Darboux take a nonempty connected set  $D \subset \mathbb{R}^2$ . If  $J = \text{pr}(D)$  is a singleton then  $f[D] = g[\text{pr}(D)]$  is a singleton, so it is connected. Otherwise  $\text{pr}(D)$  contains an interval  $(a, b) \neq \emptyset$  and, by (\*),  $f[D] = g[\text{pr}(D)] \supset g[(a, b)] = \mathbb{R}$ . Now, by Theorem 4.8,  $f$  is not almost continuous.  $\square$

As noted above, function  $f$  from Example 4.10 cannot be almost continuous. It is interesting however, that if function  $g$  used to define  $f$  is almost continuous, which can be easily constructed, then  $f \upharpoonright \mathbb{R} \times [-k, k]$  is almost continuous for every  $k > 0$ . This follows from Fact 4.6.

## 5. Some missing examples of additive Darboux-like functions on $\mathbb{R}$

**Example 5.1.** There exists an  $f \in \text{Add} \cap \text{AC} \cap \text{PR} \setminus \text{CIVP}$ . Moreover,  $f[K]$  is not nowhere dense for every perfect set  $K \subset \mathbb{R}$ .

**Proof.** Let  $\mathcal{P}$  be a family of pairwise disjoint perfect sets such that the set  $\bigcup \mathcal{P}$  is linearly independent and  $|\{P \in \mathcal{P}: P \subset (a, b)\}| = \mathfrak{c}$  for every  $a < b$ . Such a family can be easily constructed from a linearly independent perfect set. (See, e.g., [20, Theorem 2, Chapter XI, Section 7].) Let  $\mathcal{J}$  be a family of all nonempty open intervals and let  $\{\langle I_\xi, J_\xi \rangle: \xi < \mathfrak{c}\}$  be an enumeration of  $\mathcal{J} \times \mathcal{J}$ . By an easy induction we can find a one-to-one sequence  $\{P_\xi \in \mathcal{P}: \xi < \mathfrak{c}\}$  such that  $P_\xi \subset I_\xi$  for every  $\xi < \mathfrak{c}$ . Let  $H \subset \mathbb{R}$  be a Hamel basis containing  $\bigcup \mathcal{P}$  and for each  $h \in H$  let  $J_h = J_\xi$  if  $h \in P_\xi$ , and  $J_h = (0, 1)$  otherwise. Our  $f$  will be a linear extension of some function  $f_0: H \rightarrow \mathbb{R}$  such that

$$f_0(h) \in J_h \quad \text{for every } h \in H. \quad (*)$$

It is easy to see that any such an  $f$  will have a perfect road. To make sure that  $f$  has the additional property, which make it not CIVP, we will need to make some more work.

Let  $\mathcal{F}$  be the family of all perfect nowhere dense subsets of  $\mathbb{R}$  and let  $\{\langle K_\xi, S_\xi \rangle: \xi < \mathfrak{c}\}$  be an enumeration of  $\mathcal{F} \times \mathcal{F}$ . We will make sure for every  $\xi < \mathfrak{c}$  there exists an  $x_\xi \in K_\xi$  such that  $f(x_\xi) \notin S_\xi$ . Clearly such a function will have all the desired property. For this we will construct a sequences  $\langle H_\xi: \xi < \mathfrak{c} \rangle$  of pairwise disjoint finite subsets of  $H$  and  $\langle g_\xi: \xi < \mathfrak{c} \rangle$  of functions from  $H_\xi$  into  $\mathbb{R}$  such that for every  $\xi < \mathfrak{c}$

- (i)  $g_\xi(h) \in J_h$  for every  $h \in H_\xi$ ,
- (ii) there exists an  $x_\xi \in K_\xi \cap \text{LIN}_{\mathbb{Q}}(\bigcup \{H_\zeta: \zeta \leq \xi\})$  such that  $G_\xi(x_\xi) \notin S_\xi$ , where  $G_\xi = \text{LIN}_{\mathbb{Q}}(\bigcup \{g_\zeta: \zeta \leq \xi\})$ .

Now, the inductive choice of  $H_\xi$  and  $g_\xi$  is quite simple. We choose an  $x_\xi \in K_\xi \setminus \text{LIN}_{\mathbb{Q}}(\bigcup \{H_\zeta: \zeta < \xi\})$  and represent  $x_\xi$  as  $z + q_1 h_1 + \dots + q_n h_n$  where  $z \in \text{LIN}_{\mathbb{Q}}(\bigcup \{H_\zeta: \zeta < \xi\})$ ,  $h_1, \dots, h_n \in H \setminus \bigcup \{H_\zeta: \zeta < \xi\}$ , and  $q_1, \dots, q_n \in \mathbb{Q}$ . We put  $H_\xi = \{h_1, \dots, h_n\}$  and define  $g_\xi$  such that (i) and (ii) are satisfied. This can be done, since we have an open interval of possible choices for each value of  $g_\xi(h_i)$ , while we have to omit only a nowhere dense set  $S_\xi$  for the value of  $G_\xi(x_\xi) = G_\xi(z) + q_1 g_\xi(h_1) + \dots + q_n g_\xi(h_n)$ . Now, if  $f_0: H \rightarrow \mathbb{R}$  is any extension of  $\bigcup_{\xi < \mathfrak{c}} g_\xi$  which satisfies (\*) then a linear extension  $f$  of  $f_0$  has all the desired properties.  $\square$

**Example 5.2.** There exists an  $f \in \text{Add} \cap \text{SCIVP} \cap \text{PC} \setminus \text{D}$ .

**Proof.** Let  $H_0$  be a Hamel basis such that  $|H_0 \cap K| = \mathfrak{c}$  for every perfect set  $K \subset \mathbb{R}$ . (See [4, Corollary 7.3.7].) Let  $h_0 \in H_0$  and  $V = \text{LIN}_{\mathbb{Q}}(H_0 \setminus \{h_0\})$ . Then  $V$  is a proper linear subspace of  $\mathbb{R}$  with  $|V \cap K| = \mathfrak{c}$  for every perfect  $K \subset \mathbb{R}$ .

As in the example above, let  $\mathcal{P}$  be a family of pairwise disjoint perfect sets such that the set  $\bigcup \mathcal{P}$  is linearly independent and  $|\{P \in \mathcal{P}: P \subset (a, b)\}| = \mathfrak{c}$  for every  $a < b$ . Let  $\mathcal{J}$  be a family of all nonempty open intervals and let  $\{(I_\xi, v_\xi): \xi < \mathfrak{c}\}$  be an enumeration of  $\mathcal{J} \times V$ . Find a one-to-one sequence  $\{P_\xi \in \mathcal{P}: \xi < \mathfrak{c}\}$  such that  $P_\xi \subset I_\xi$  for every  $\xi < \mathfrak{c}$  and let  $H \subset \mathbb{R}$  be a Hamel basis containing  $\bigcup \mathcal{P}$ .

Define  $f_0: H \rightarrow V$  by putting  $f_0(h) = v_\xi$  for  $h \in P_\xi$  and choose an arbitrary  $f_0(h) \in V$  for all other  $h \in H$ . Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a linear extension of  $f_0$ . Then  $f$  is additive and non-zero, so  $f$  has a dense graph. Thus  $f \in \text{PC}$ . Also  $f[\mathbb{R}] = V$  implying  $f \notin \text{D}$ . It is SCIVP since for every perfect set  $K \subset \mathbb{R}$  and  $a < b$  there exist  $v \in V \cap K$  and  $\xi < \mathfrak{c}$  such that  $\langle I_\xi, v_\xi \rangle = \langle (a, b), v \rangle$ . So  $f[P_\xi] = \{v_\xi\} = \{v\} \subset K$  and  $f \upharpoonright P_\xi$  is continuous.  $\square$

**Example 5.3.** There exists an  $f \in \text{Add} \cap \text{SCIVP} \cap \text{D} \setminus \text{Conn}$ .

**Proof.** The construction is very similar to that for function  $g$  from Example 4.10. Let  $L = \{(x, x + 1): x \in \mathbb{R}\}$ . As above choose a family  $\mathcal{P}$  of pairwise disjoint perfect sets such that the set  $\bigcup \mathcal{P}$  is linearly independent and  $|\{P \in \mathcal{P}: P \subset (a, b)\}| = \mathfrak{c}$  for every  $a < b$ . Let  $\{A, B\}$  be a partition of continuum  $\mathfrak{c}$  with  $|A| = |B| = \mathfrak{c}$ . Put  $\mathcal{J} = \{(a, b): a < b\}$  and let  $\{(I_\xi, r_\xi): \xi \in A\}$  be an enumeration of  $\mathcal{J} \times \mathbb{R}$ . Also, find a one-to-one sequence  $\{P_\xi \in \mathcal{P}: \xi < \mathfrak{c}\}$  such that  $P_\xi \subset I_\xi$  for every  $\xi < \mathfrak{c}$ . Finally, let  $H \subset \mathbb{R}$  be a Hamel basis containing  $\bigcup \mathcal{P}$  and let  $H = \{h_\xi: x \in B\}$ .

By induction on  $\xi < \mathfrak{c}$  construct a sequence  $\langle (K_\xi, y_\xi): \xi < \mathfrak{c} \rangle$  such that for every  $\xi < \mathfrak{c}$  the following conditions hold.

- (i)  $K_\xi \cap \text{LIN}_{\mathbb{Q}}(\bigcup \{K_\zeta: \zeta < \xi\}) = \emptyset$ .
- (ii)  $L \cap \text{LIN}_{\mathbb{Q}}(\bigcup \{K_\zeta \times \{y_\zeta\}: \zeta \leq \xi\}) = \emptyset$ .
- (iii) If  $\xi \in A$  then  $K_\xi$  is a perfect subset of  $P_\xi$  and  $y_\xi = r_\xi$ .
- (iv) If  $\xi \in B$  then  $|K_\xi| \leq 1$  and  $h_\xi \in \text{LIN}_{\mathbb{Q}}(\bigcup \{K_\zeta: \zeta \leq \xi\})$ .

To find such a sequence assume that a sequence  $\langle (a_\zeta, y_\zeta): \zeta < \xi \rangle$  is already constructed for some  $\xi < \mathfrak{c}$ . If  $\xi \in A$  put  $y_\xi = r_\xi$  and look at the set

$$Z = L \cap \text{LIN}_{\mathbb{Q}}\left(\bigcup \{K_\zeta \times \{y_\zeta\}: \zeta \leq \xi\}\right) \cup (P_\xi \times \{y_\xi\}).$$

Notice that  $Z$  has cardinality less than  $\mathfrak{c}$  since  $Z \subset \mathbb{R} \times \text{LIN}_{\mathbb{Q}}(\bigcup \{y_\zeta: \zeta \leq \xi\})$  and  $L$  intersects every horizontal line at exactly one point. Let  $T = \text{pr}(Z)$  and notice that by the inductive assumption of (ii) we have  $T \cap \text{LIN}_{\mathbb{Q}}(\bigcup \{K_\zeta: \zeta < \xi\}) = \emptyset$ . For every  $t \in T$  choose an  $h_t \in H \setminus \bigcup \{K_\zeta: \zeta < \xi\}$  with  $t \notin \text{LIN}_{\mathbb{Q}}(H \setminus \{h_t\})$ . Then the set

$$S = \{h_t: t \in T\} \cup \bigcup \{K_\zeta: \zeta < \xi \text{ and } \zeta \in B\}$$

has cardinality less than  $\mathfrak{c}$ . Choose a perfect set  $K_\xi \subset P_\xi \setminus S$ . Then  $K_\xi$  satisfies (i)–(iii).

Next assume that  $\xi \in B$  and let  $V = \text{LIN}_{\mathbb{Q}}(\bigcup\{K_{\zeta} : \zeta < \xi\})$ . If  $h_{\xi} \in V$  we put  $K_{\xi} = \emptyset$  and choose  $y_{\xi}$  arbitrarily. So assume that  $h_{\xi} \notin V$ . We put  $K_{\xi} = \{h_{\xi}\}$ . This guarantees (i) and (iv). To get (ii) we have to find  $y_{\xi}$  such that the set

$$\text{LIN}_{\mathbb{Q}}(V \cup \{h_{\xi}, y_{\xi}\}) = \{\langle x, y \rangle + q\langle h_{\xi}, y_{\xi} \rangle : \langle x, y \rangle \in V \text{ and } q \in \mathbb{Q}\}$$

is disjoint with  $L$ . Thus, we must have  $y + qy_{\xi} \neq x + qh_{\xi} + 1$ , that is,  $y_{\xi} \neq px + h_{\xi} - py + p$  for every  $\langle x, y \rangle \in V$  and  $q = p^{-1} \in \mathbb{Q} \setminus \{0\}$ . Therefore it is enough to choose

$$y_{\xi} \notin \text{LIN}_{\mathbb{Q}}\left(\bigcup\{K_{\zeta} : \zeta \leq \xi\} \cup \{y_{\zeta} : \zeta < \xi\} \cup \{1\}\right)$$

which can be done, since  $\text{LIN}_{\mathbb{Q}}(\bigcup\{K_{\zeta} : \zeta \leq \xi\})$  has co-dimension  $c$ . This finishes the inductive construction.

Now put  $f = \text{LIN}_{\mathbb{Q}}(\bigcup\{K_{\xi} \times \{y_{\xi}\} : \xi < c\})$ . Then  $f : \mathbb{R} \rightarrow \mathbb{R}$  is additive and misses  $L$ , so it is not connectivity. It is Darboux and SCIVP since for every  $a, b, y \in \mathbb{R}$ ,  $a < b$ , there exists  $\xi < c$  such that  $\langle I_{\xi}, y_{\xi} \rangle = \langle (a, b), y \rangle$ . So  $K_{\xi} \subset P_{\xi} \subset I_{\xi} = (a, b)$ ,  $f[K_{\xi}] = \{y_{\xi}\}$ , and  $f \upharpoonright K_{\xi}$  is continuous.  $\square$

## References

- [1] D. Banaszkowski, On some subclasses of additive functions, PhD Thesis, Łódź University, 1997 (in Polish).
- [2] J.B. Brown, Almost continuous Darboux functions and Reed's pointwise convergence criteria, *Fund. Math.* 86 (1974) 1–7.
- [3] J. Brown, P. Humke, M. Laczko, Measurable Darboux functions, *Proc. Amer. Math. Soc.* 102 (1988) 603–610.
- [4] K. Ciesielski, *Set Theory for the Working Mathematician*, London Math. Soc. Student Texts 39, Cambridge Univ. Press, Cambridge, 1997.
- [5] K. Ciesielski, Some additive Darboux-like functions, *J. Appl. Anal.* 4 (1) (1998) 43–51. (Preprint\* available. <sup>2</sup>)
- [6] K. Ciesielski, T. Natkaniec, J. Wojciechowski, Extending connectivity functions on  $\mathbb{R}^n$ , *Topology Appl.*, to appear. (Preprint\*.)
- [7] K. Ciesielski, I. Reclaw, Cardinal invariants concerning extendable and peripherally continuous functions, *Real Anal. Exchange* 21 (1995–96) 459–472. (Preprint\* available.)
- [8] K. Ciesielski, A. Rosłanowski, Two examples concerning almost continuous functions, *Topology Appl.* 103 (2000) 187–202 (this issue). (Preprint\* available.)
- [9] K. Ciesielski, J. Wojciechowski, Sums of connectivity functions on  $\mathbb{R}^n$ , *Proc. London Math. Soc.* 76 (2) (1998) 406–426. (Preprint\* available.)
- [10] R.G. Gibson, T. Natkaniec, Darboux like functions, *Real Anal. Exchange* 22 (2) (1996–97) 492–533. (Preprint\* available.)
- [11] R.G. Gibson, F. Roush, Connectivity functions with a perfect road, *Real Anal. Exchange* 11 (1985–86) 260–264.
- [12] R.G. Gibson, F. Roush, A characterization of extendable connectivity functions, *Real Anal. Exchange* 13 (1987–88) 214–222.
- [13] W.J. Gorman III, The homeomorphic transformation of  $c$ -sets into  $d$ -sets, *Proc. Amer. Math. Soc.* 17 (1966) 826–830.

<sup>2</sup> Preprints marked by \* are available in electronic form accessible from *Set Theoretic Analysis Web page*: <http://www.math.wvu.edu/homepages/kcies/STA/STA.html>



- [14] M.R. Hagan, Equivalence of connectivity maps and peripherally continuous transformations, *Proc. Amer. Math. Soc.* 17 (1966) 175–177.
- [15] O.H. Hamilton, Fixed points for certain noncontinuous transformations, *Proc. Amer. Math. Soc.* 8 (1957) 750–756.
- [16] W. Hurewicz, H. Wallman, *Dimension Theory*, Princeton Univ. Press, Princeton, NJ, 1948.
- [17] J. Jastrzębski, An answer to a question of R.G. Gibson and F. Roush, *Real Anal. Exchange* 15 (1989–90) 340–341.
- [18] J. Jastrzębski, T. Natkaniec, On sums and products of extendable functions, *Real Anal. Exchange*, to appear.
- [19] J. Lipiński, On some extension of almost continuous functions, *Tatra Mt. Math. Publ.* 2 (1993) 15–18.
- [20] M. Kuczma, *An Introduction to the Theory of Functional Equations and Inequalities*, Polish Scientific Publishers, PWN, Warsaw, 1985.
- [21] A. Maliszewski, T. Natkaniec, Almost continuous functions of two variables, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales* 87 (1993) 396–404.
- [22] T. Natkaniec, Almost continuity, *Real Anal. Exchange* 17 (1991–92) 462–520.
- [23] T. Natkaniec, *Almost Continuity*, WSP, Bydgoszcz, 1992.
- [24] T. Natkaniec, Extendability and almost continuity, *Real Anal. Exchange* 21 (1995–96) 349–355.
- [25] H. Rosen, Every real function is the sum of two extendable connectivity functions, *Real Anal. Exchange* 21 (1995–96) 299–303.
- [26] H. Rosen, R.G. Gibson, F. Roush, Extendable functions and almost continuous functions with a perfect road, *Real Anal. Exchange* 17 (1991–92) 248–257.
- [27] J. Stallings, Fixed point theorems for connectivity maps, *Fund. Math.* 47 (1959) 249–263.
- [28] G.T. Whyburn, *Analytic Topology*, Amer. Math. Soc. Colloquium Publication 28, Amer. Math. Soc., Providence, RI, 1942.