

1996

Phylogeny of Polemoniaceae Based on Nuclear Ribosomal Internal Transcribed Spacer DNA Sequences

J. Mark Porter

Rancho Santa Ana Botanic Garden

Follow this and additional works at: <http://scholarship.claremont.edu/aliso>



Part of the [Botany Commons](#)

Recommended Citation

Porter, J. Mark (1996) "Phylogeny of Polemoniaceae Based on Nuclear Ribosomal Internal Transcribed Spacer DNA Sequences," *Aliso: A Journal of Systematic and Evolutionary Botany*: Vol. 15: Iss. 1, Article 6.
Available at: <http://scholarship.claremont.edu/aliso/vol15/iss1/6>

PHYLOGENY OF POLEMONIACEAE BASED ON NUCLEAR RIBOSOMAL INTERNAL TRANSCRIBED SPACER DNA SEQUENCES

J. MARK PORTER

Rancho Santa Ana Botanic Garden
1500 North College Avenue
Claremont California 91711-3157
e-mail: porterj@cgs.edu

ABSTRACT

Nuclear ribosomal internal transcribed spacer (ITS) DNA sequences are used to estimate the phylogeny of 53 members of Polemoniaceae, representing all but two genera of the family. Fitch parsimony analysis of equal-weighted nucleotide sites result in 1080 minimal-length trees. However, when alignment-ambiguous positions are removed and an 11:10 transition to transversion weighting is imposed only eight trees are found. These data are used to address two issues: 1) patterns of diversification in Polemoniaceae, and 2) the circumscription and monophyly of the genus *Gilia*. Although the monophyly of Polemoniaceae is well supported, relationships inferred among the earliest diverging lineages are altered by character weighting, treatment of indels, and taxon inclusion. In spite of the lack of reliable resolution at the basal nodes, ITS data provide evidence that *Gilia*, as currently interpreted, is polyphyletic and comprises at least five independent lineages.

Key words: DNA sequences, *Gilia*, Polemoniaceae, phylogeny.

INTRODUCTION

The Polemoniaceae are a small, mostly New World, group of between 300 and 340 species (Grant 1959; Willis 1966; Cronquist 1981; Thorne 1992; Patterson 1993). Evolutionary relationships and classification within this family have long been controversial (Bentham 1845; Gray 1870; Kuntze 1891; Brand 1907; Rydberg 1913; Dawson 1936; Wherry 1940; Mason 1945; Mason and Grant 1948; Grant 1959; Cronquist 1984; Day and Moran 1986). Although generally considered a "natural" group by most authors, generic and species relationships within the family have remained obscure: the circumscription of species, genera and higher taxonomic groups differing radically from author to author. The most recent family-wide classification of Polemoniaceae is that of Grant (1959), who recognizes 18 genera, placed into five tribes (Table 1). Over the past 30 years, a broad spectrum of data has accumulated bearing on relationships within the family. These include family- and generic-level surveys of palynological (Stuchlik 1967; Loeblich 1964; Taylor and Levin 1975; Chuang et al. 1978; Timbrook 1986; Lord and Eckard 1986; Day and Moran 1986), histological (Carlquist et al. 1984), chemosystematic (Smith et al. 1977; Dean et al. 1978; Harborne and Smith 1978; Harborne 1980; Smith et al. 1982; Wilken et al. 1982; Kiredjiad and Smith 1985), and reproductive data (Grant and Grant 1965; Plitmann and Levin 1983; 1990). Although neither phenetic nor phylogenetic analyses have been conducted on the family using this wide array of morphological and chemical data (but

see Wilken and Hartman 1991), recent phylogenetic analyses of nucleotide sequences of the chloroplast gene *matK* (Steele 1991, 1992; Steele and Vilgalys 1994; Johnson and Soltis 1995; Johnson et al. 1996) suggest these tribes are not monophyletic. In light of these data, realignment of Grant's (1959) tribes appears inevitable. This study provides additional insight into classification and evolutionary history of Polemoniaceae from an independent source of evidence. Presented is a molecular phylogenetic study using sequences from the nuclear genes coding for ribosomal RNA, the internal transcribed spacers (ITS1 and ITS2).

Taxonomic Background

The contrast between various classifications of Polemoniaceae has been reviewed by Grant (1959: 4). There is little consensus among historical classifications. The few general similarities (particularly for classifications since 1900) include: 1) the tendency to place the tropical genera *Cantua*, *Cobaea* and *Bonplandia* in separate tribes apart from the remainder of the so-called "temperate genera" (Brand 1907; 1908; Grant 1959); and 2) the inclusion of *Linanthus* and *Leptodactylon* in the same tribe with *Gilia* (Brand 1907; 1908; Wherry 1940; 1945; Grant 1959) while placing *Phlox* in a different tribe (Wherry 1940; 1945; Grant 1959). A particularly striking disparity among historical classifications in the family is the circumscription of the genus *Gilia*. These differences range from broad circumscriptions that include nearly all of

the "temperate genera" except *Polemonium*, *Phlox*, and *Collomia* (e.g., Bentham and Hooker 1876) to the present treatment by Grant (1959) that segregates *Allophyllum*, *Ipomopsis*, *Eriastrum*, *Langloisia*, *Loeseliastrum*, and *Navarretia* from *Gilia*.

Grant's currently accepted tribal classification has not found support from recent molecular studies. Steele and Vilgalys (1994) provided evidence from cpDNA *matK* sequences that some genera included in tribe Polemoniaceae share more recent common ancestry with genera placed in tribe Gilieae. For example *Collomia* and *Allophyllum* (tribe Polemoniaceae) are more closely related to *Gilia* and *Navarretia* (tribe Gilieae) than to either *Phlox* or *Polemonium* (tribe Polemoniaceae). Indeed, Steele and Vilgalys suggest that there is no support for two temperate tribes, rather Polemoniaceae possess a single temperate lineage. These results were not only corroborated by Johnson and Soltis (1995) and Johnson et al. (1996) but due to the increased taxon sampling, evidence was presented that the genus *Gilia*, as currently circumscribed, is a polyphyletic assemblage. Some members of *Gilia* (e.g., *G. filiformis*) are more closely related to genera never associated with tribe Gilieae (e.g., *Phlox*) than to other members of *Gilia*.

This controversy over relationships within Polemoniaceae is problematic because Polemoniaceae have been used as a model system for the study of many aspects of character evolution and reproductive biology. The family as well as individual species have been used as primary examples for the pattern and modes of evolution of pollination mechanisms (Grant and Grant 1965; Galen and Kevan 1983; Paige and Whitham 1985, 1987a; Galen and Stanton 1989), evolution of plant sexual systems (Grant 1959; Schoen 1982a, b, c, 1983; Plitmann and Levin 1983, 1990), hybridization and its evolutionary consequences (Grant 1950, 1954a, b, 1957, 1979; Grant and Grant 1960; Day 1965; Levin 1979; Coyne 1974; Grant and Wilken 1987), natural selection on floral traits (Ellstrand 1983; Paige and Whitham 1985; Wolf et al. 1986; Ellstrand and Mitchell 1988; Campbell 1989; 1991), and evolution life history (Paige and Whitham 1987a, b). An accurate phylogenetic backdrop is necessary to interpret evolutionary changes in these traits (Donoghue 1989). The chloroplast phylogenies provide an important inroad into understanding the phylogeny of Polemoniaceae. However, a single gene phylogeny may not correspond with an organismal phylogeny (Dwyer et al. 1991; Doyle 1992; Neigel and Avise 1986; Sanderson and Doyle 1992). A phylogeny based on an independent source of data is both desirable and necessary to either corroborate or refute phylogenetic inferences from the chloroplast genome.

In this paper I provide a preliminary assessment of phylogenetic relationships of Polemoniaceae using

DNA sequences of the internal transcribed spacers (ITS1 and ITS2) of the nuclear ribosomal cistron. Because these spacer sequences are encoded in the nuclear genome, they represent an independent source of phylogenetic information from chloroplast sequences.

METHODS

Sampled Taxa

An attempt was made to include representatives of all of the 19 currently recognized genera of Polemoniaceae (Table 1). Because of the controversy over the limits of the genus *Gilia* and its relation to *Ipomopsis*, 13 species of *Gilia* (members of sections *Arachnion*, *Gilia*, *Giliandra*, *Giliastrum*, *Gilmania*, and *Saltugilia*) and five species of *Ipomopsis* (members of sections *Ipomopsis*, *Microgilia*, *Phloganthea* [including the *Giliopsis* group]). In addition, a largely unknown woody member of Polemoniaceae, *Gilia scabra* (treated by Grant 1959 as a synonym of *Linanthus nuttallii*), is also included. Only two genera are not represented. *Gymnosteris*, with two species, is not included because the DNAs did not sequence well enough for reasonable interpretation (nearly all of the nucleotide sites appeared polymorphic). In addition, the DNA sequence from *Huthia coerulea* is not included because it is unalignable with other Polemoniaceae sequences.

Selection of an outgroup for Polemoniaceae is problematical. In spite of most present classifications (Cronquist 1981; Takhtajan 1980; Thorne 1968, 1992), molecular data from the chloroplast gene *rbcL* (Olmstead et al. 1992; Chase et al. 1993) provide evidence that Polemoniaceae are more closely related to Ericalean groups, usually placed in the Dilleniidae, than to the Asteridae families with which they are traditionally placed: i.e., Convolvulaceae, Hydrophyllaceae, and Solanaceae (but see Anderberg 1992; Martin and Dowd 1991). In particular, chloroplast sequence data from *rbcL* (Olmstead et al. 1992) and *matK* (Johnson et al. 1996) place Fouquieriaceae as the sister group to Polemoniaceae. By contrast, the Chase et al. study of *rbcL* supports a close relationship between Polemoniaceae and Diapensiaceae. Close relationships between Polemoniaceae and both Diapensiaceae (Linnaeus 1753; Don 1822) and Fouquieriaceae (Humbolt et al. 1823; Nash 1903; Engler and Gilg 1924; Abrams 1951; Thorne 1977) have been suggested based on morphological evidence. Based on this evidence, I have selected Diapensiaceae and Fouquieriaceae as outgroups. Attempts were made to also include sequences from Solanales, e.g., *Nicotiana rustica* L. (Venkateswarlu and Nazar 1991); however, these sequences are unalignable.

Collection and DNA Extraction

Material collected in the field was placed on ice for transport to storage at -80°C . Total genomic DNA extracts from leaf material, powdered under liquid nitrogen, used a 2X CTAB protocol, modified from Doyle and Doyle (1987). In some cases a 4X CTAB (with 0.1% PVP-40) extraction buffer was used for species with high polysaccharide content. The DNAs were further purified using cesium chloride/ethidium bromide gradients (Maniatis et al. 1982). The DNAs of *Acanthogilia gloriosa* and *Gilia scabra* were extracted from leaf material desiccated using silica gel (Liston and Rieseberg 1990; Chase and Hillis 1991). The DNAs of six taxa were obtained from leaf fragments taken from herbarium collections (Table 1). Subsequently, all DNA samples obtained from herbarium specimens have been verified using freshly collected material, and in no case did the DNA sequences of ITS differ at more than a single, uninformative nucleotide site (Porter unpubl.).

PCR Amplification and Sequencing

Sequencing was performed on 39 samples using the method of Sanger et al. (1977). Single-stranded DNAs for both strands of the two internal transcribed spacers of the nuclear ribosomal cistron, ITS1 and ITS2, were amplified directly by asymmetric polymerase chain reaction (PCR), using a 1:20 ratio of the primers "ITS5" (5'-GGA AGT AAA AGT CGT AAC AAG G-3') and "ITS2" (5'-GCT GCG TTC TTC ATC GAT GC-3') for the ITS1 spacer, and the primers "ITS3" (5'-GCA TCG ATG AAG AAC GCA GC-3') and "ITS4" (5'-TCC TCC GCT TAT TGA TAT GC-3') for the ITS2 spacer (White et al. 1990). PCR amplifications, PCR product purification and direct single-stranded DNA sequencing followed the procedures described by Baldwin (1992), except for the following modifications. The amplification of DNA from herbarium material employed a modified 10 \times *Taq* polymerase reaction buffer consisting of 67mM Tris-HCl (pH 8.8), 2 mM MgCl_2 , and 2 mg/ml bovine serum albumin (Pääbo et al. 1989; Pääbo 1990). The PCR products were electrophoresed using a 1.5% agarose gel (pre-stained with ethidium bromide) in a 1 \times TBE (pH 8.3) buffer, to confirm the single-stranded product and purified using differential filtration in Millipore Ultra-Free-MC microfuge tubes (Millipore UFC-3 THK00). The purified DNAs were sequenced using the dideoxy chain termination reaction, following the procedures outlined in the TAQuence[®] kit (U. S. Biochemical Co.). Sequencing products were labeled using α - ^{35}S dATP. To reduce base compressions, 7-deazadGTP was substituted for dGTP in all of the sequencing reactions. The gels were fixed in 5% acetic acid/5% methanol for 45 min and vacuum dried at

80°C for 1 hr. The autoradiographs were exposed for a minimum of 24 hr. All sequences were checked by sequencing the opposite strand.

Sequence data for 14 species were obtained using an Applied Biosystems Model 373A Automated DNA Sequencing System (Perkin Elmer). Template DNAs were prepared by symmetric PCR of the entire ITS region using a 1:1 ratio of primers ITS5 and ITS4. Direct cycle-sequencing of the ITS region followed manufacturers specifications, using the PRISM[™] DyeDeoxy[™] Terminator Kit (Perkin Elmer) and employed primers ITS2, ITS3, "ITS5I" (5'-AGG TG ACC TGC GGA AGG ATC ATT-3') and "ITS4I" (5'-GGT AGT CCC GCC TGA CCT GG-3'). Primers ITSP5I and ITSP4I are internal to and flanking ITS5 and ITS4, respectively. The four primers provide sequences for overlapping fragments that collectively cover both strands of the entire ITS region (ITS1, 5.8S and ITS2).

Sequence Alignment

The 56 sequences were truncated to include only ITS1 and ITS2. Identification of the terminal ends of ITS1 and ITS2 was based on comparisons with published sequences for *Daucus carota* L., *Nicotiana rustica*, *Vicia faba* L., species of *Astragalus*, and representatives of *Madiinae* (Baldwin 1992, 1993; Venkateswarlu and Nazar 1991; Wojciechowski et al. 1993; Yokota et al. 1989). The two spacer regions were aligned separately, using PILEUP (Feng and Doolittle 1987; Higgins and Sharp 1987; see also Needleman and Wunch 1987), a multiple sequence alignment program of the University of Wisconsin Genetics Computer Group (UWGCG) software package. The alignment parameters were varied from GapWeight= 1.0 and GapLengthWeight= 0.1 to GapWeight= 5.0 and GapLengthWeight= 3.0. With higher gap and length penalty values, fairly large blocks of aligned sequences are produced, but single taxa are sometimes grossly misaligned. Low gap and length penalties result in alignments lacking grossly misaligned taxa, but with a greater frequency of gaps. Evenso, alternative alignments were evident. The resulting alignments were both analyzed directly and manually realigned and reanalyzed. The PILEUP aligned sequences were manually realigned to maximize the number of invariant nucleotide positions. The results presented refer to sequences which were realigned manually; however, the conclusions apply to both alignments.

Phylogenetic Analysis

Estimations of phylogenetic relationships, based on combined ITS1 and ITS2 sequences, were obtained using Fitch parsimony as implemented in PAUP (version 3.0; Swofford 1991). A variety of heuristic approaches

Table 1. Classification of Polemoniceae, sensu Grant (1959) and collections utilized in the phylogenetic analysis of nrDNA internal transcribed spacer sequences. The cross (†) indicates those taxa for which herbarium collections were used for DNA extractions. The asterisk (*) indicates those species for which cycle sequencing and automated sequencing procedures were used. Vouchers are cited and housed as indicated. The taxa included to serve as the outgroup are designated as such parenthetically.

Tribe Cantuaeeae (2: ca. 9)		
<i>Acanthogilia</i> (1 species)	<i>n</i> = 9	<i>A. gloriosa</i> (Brandeg.) Day & Moran. 2.4 km west of Punta Prieta, Baja California, Mexico. [J. M. Porter & K. D. Heil 7987; SJNM]
<i>Cantua</i> (ca. 12 species)	<i>n</i> = 27	<i>C. quercifolia</i> Cav. Cult. San Francisco State University, San Francisco, California. [R. Patterson sn; RSA]
<i>Huthia</i> (2 species)	<i>n</i> = ?	†* <i>H. coerulea</i> Brand. Arequipa, Peru. [O. Tovar 3543; MO]
Tribe Bonplandieae (2: ca. 17)		
<i>Bonplandia</i> (2 species)	<i>n</i> = 15	<i>B. geminiflora</i> Cav. Cult. San Francisco State University, San Francisco, California. [R. Patterson sn; RSA]
<i>Loeselia</i> (ca. 15 species)	<i>n</i> = 9	<i>L. glandulosa</i> (Cav.) Don. Patagonia Mountains, Santa Cruz County, Arizona. [J. M. Porter & C. Campbell 9231; AZ, SJNM]
Tribe Cobaeaeae (1: 19)		
<i>Cobaea</i> (19 species)	<i>n</i> = 27	<i>C. scandens</i> Cav. Cult. San Francisco State University, San Francisco, California. [R. Patterson sn; RSA]
	<i>n</i> = ?	† <i>C. baiurita</i> Standl. Puerto Vienda, Chiapas, Mexico. 20 Aug. 1967. [O. Clarke 293; AZ]
Tribe Polemonieae (5: ca. 102)		
<i>Allophyllum</i> (5 species)	<i>n</i> = 9	* <i>A. divaricatum</i> (Nutt.) A. Grant & V. Grant. Mt. Konocti, Lake County, CA. [J. M. Porter & L. Machen 10819; RSA]
	<i>n</i> = 9	<i>A. gilioides</i> (Benth.) Grant & Grant. 5 miles south of Oracle, Santa Catalina Mtns., Pinal County, Arizona. [J. M. Porter & L. Machen 8751; AZ, SJNM]
<i>Collomia</i> (15 species)	<i>n</i> = 8	* <i>C. grandiflora</i> Lindl. Libre Mtns., Los Angeles County, California. [T. Ross & S. Boyd 8142; RSA]
	<i>n</i> = 8	<i>C. linearis</i> Nutt. 10 miles northeast of Pogosa Springs, Archuleta County, Colorado. [J. M. Porter 8565; AZ]
<i>Gymnosteris</i> (2 species)	<i>n</i> = 6	—
<i>Phlox</i> (ca. 60 species)	<i>n</i> = 7	<i>P. stanburyi</i> (Torr.) Heller. 4 miles south of the junction of Nevada Highway 160 and US Highway 95, Nye County Nevada. [J. M. Porter 8841; SJNM]
	<i>n</i> = 7	* <i>P. (Microsteris) gracilis</i> E. Greene. Mt. Emma, Los Angeles County, California [J. M. Porter 10566; RSA]
<i>Polemonium</i> (ca. 20 species)	<i>n</i> = 9	<i>P. foliosissimum</i> Gray. Lizard Head Pass, along Colorado Highway 145, Dolores County, Colorado. [J. M. Porter 7526; SJNM]
Tribe Gilieae (8: ca. 200)		
<i>Eriastrum</i> (14 species)	<i>n</i> = 7	<i>E. diffusum</i> (Gray) Mason. Tucson Mountains, 4 miles west of Tucson, Pima County, Arizona. [J. M. Porter 8754; AZ]
<i>Gilia</i> (ca. 70 species)	<i>n</i> = 8	<i>G. caespitosa</i> Gray. Capitol Reef National Park, Wayne County, Utah [J. M. Porter & K. Heil 7352; SJNM]
	<i>n</i> = 9	†* <i>G. campanulata</i> Gray. 5 mi W Independence, Inyo County, CA [Kerr 638; RSA]
	<i>n</i> = 9	† <i>G. capitata</i> Sims. E of Washougal, Skamania County, Washington. 14 July 1984. [R. S. Halse 2900; AZ]
	<i>n</i> = 72	†* <i>G. crassifolia</i> Benth. Rio Molles, Endesa, Coquimbo, Chile. 5 Nov. 1970. [J. Simon 161.; RSA]
	<i>n</i> = 9	* <i>G. filiformis</i> Gray. 12 mi E. Independence, Inyo County, CA [J. M. Porter & L. Machen 10849; RSA]
	<i>n</i> = 18	<i>G. flavocincta</i> A. Nels subsp. <i>australis</i> A. & V. Grant. Ca. 6 miles east of Roosevelt Reservoir, roadside, Pinal County, Arizona. [J. M. Porter 7060; SJNM]
	<i>n</i> = 9	† <i>G. foetida</i> Gil. ex Benth. Las Heras camino a Yalguaraz, Provincia de Mendoza, Argentina. 27 Dec. 1947. [R. Leal, 11,113; AZ]
	<i>n</i> = 9	<i>G. hutchinsifolia</i> Rydb. 5 mi S of Lathrop Wells, Nye County, Nevada. [J. M. Porter 8209; AZ, SJNM]
	<i>n</i> = ?	* <i>G. inyoensis</i> I. M. Johnston. Anchorite Pass, Mineral County, NV. [J. M. Porter & L. Machen 10842; RSA]
	<i>n</i> = 18	<i>G. latifolia</i> Wats. Cataract Canyon, Canyonlands National Park, San Juan County, Utah. [J. M. Porter & M. Heil 8804; AZ, SJNM]
	<i>n</i> = 9	* <i>G. maculata</i> Parish. Whitewater Canyon, Riverside County, CA. [J. M. Porter & L. Machen 10560; RSA]
	<i>n</i> = 8	<i>G. mcvickerae</i> M. E. Jones. Ca. 4 mi S of Panguitch on UT89, Garfield County, Utah. [J. M. Porter and L. Machen 7184; SJNM]

Table 1. Continued.

	<i>n</i> = 8	<i>G. nyensis</i> J. Reveal. Red001 burned area, Red Rock Canyon, Nevada Nuclear Test Site, Nye County, Nevada. [<i>J. M. Porter</i> 8202; AZ]
	<i>n</i> = 9	<i>G. ochroleuca</i> M. E. Jones. Ca. 4 mi. S of Big Pine, Inyo County, California. [<i>J. M. Porter</i> 8860; AZ, SJNM]
	<i>n</i> = 8	<i>G. pentstemonoides</i> M. E. Jones. N side of the Gunnison River, Cimarron, Montrose County, Colorado. [<i>J. M. Porter</i> 7192; AZ, SJNM]
	<i>n</i> = 9	<i>G. rigidula</i> Benth. 7 miles southeast of Sonoita, Santa Cruz County, Arizona. [<i>J. M. Porter</i> 8723; RSA, SJNM]
	<i>n</i> = 9	<i>G. scabra</i> Brandegee. Cerro Prietto, near San Hiralito, Viscaino Desert, Baja California Sur, Mexico. [<i>J. M. Porter and K. Heil</i> 7991; SJNM]
	<i>n</i> = 9	<i>G. scopulorum</i> M. E. Jones. Hualapai Indian Reservation, Mohave County, Arizona. 21 Apr. 1973. [<i>P. S. Martin, sn</i> ; AZ]
	<i>n</i> = 9	* <i>G. splendens</i> Dougl. ex Lindl. Pacific Crest Trail, San Bernardino County, CA. [<i>J. M. Porter</i> 10142; RSA]
	<i>n</i> = 9	<i>G. stellata</i> Heller. Roadside along AZ85, 9 mi N of Why, Pima County, AZ. [<i>J. M. Porter</i> 9492; SJNM]
	<i>n</i> = 8	<i>G. subnuda</i> Gray. W of Steamboat Canyon, Apache County, AZ. [<i>J. M. Porter & K. Heil</i> 7355; SJNM]
	<i>n</i> = 18	† <i>G. tenerrima</i> A. Gray. 70 mi NW of Elko, Elko County, Nevada. 26 June 1937. [<i>N. Nichols & L. Lund</i> 206; AZ]
	<i>n</i> = 9	<i>G. tricolor</i> Benth. Cult. University of Arizona, seed from retail seed source. [<i>J. M. Porter sn</i> ; AZ]
<i>Ipomopsis</i> (30 species)	<i>n</i> = 7	<i>I. gunnisonii</i> (Torr. and Gray) V. Grant. Navajo Mine, 12 miles south of Waterflow, San Juan County, New Mexico. [<i>J. M. Porter</i> 9295; AZ, SJNM]
	<i>n</i> = 7	<i>I. multiflora</i> (Nutt.) V. Grant. Molino Basin, Santa Catalina Mountains, Pima County, Arizona. [<i>J. M. Porter</i> 8052; AZ, SJNM]
	<i>n</i> = 7	<i>I. longiflora</i> (Torr.) V. Grant. 8 mi south of Safford, Greenlee County, Arizona. [<i>J. M. Porter</i> 7142; SJNM]
	<i>n</i> = ?	†* <i>I. sonora</i> (Rose) A. Grant. Sonora, Mexico. [<i>T. R. VanDevender</i> 93-20; AZ, RSA]
	<i>n</i> = 7	† <i>I. tenuifolia</i> (Gray) V. Grant. 11 May 1980. Sierra San Pedro Martir, Mexico. [<i>S. P. McLaughlin</i> 2465; AZ]
<i>Langloisia</i> (1 species)	<i>n</i> = 7	<i>L. setosissima</i> (Torr. & Gray) Greene subsp. <i>setosissima</i> . 4 miles south of the junction of Nevada Highway 160 and US Highway 95, Nye County Nevada. [<i>J. M. Porter</i> 8561; SJNM]
	<i>n</i> = 7	<i>L. setosissima</i> (Torr. & Gray) Greene subsp. <i>punctata</i> (Cov.) Timbr. 4 miles south of the junction of Nevada Highway 160 and US Highway 95, Nye County Nevada. [<i>J. M. Porter</i> 8840; SJNM]
<i>Leptodactylon</i> (7 species)	<i>n</i> = 9	* <i>L. pungens</i> (Torr.) Rydb. 3 mi S Natural Bridges Nat. Mon., Kane County, Utah. [<i>J. M. Porter</i> 8561; SJNM]
	<i>n</i> = 9	<i>L. watsonii</i> (Gray) Rydb. Waterpocket Fold, Capitol Reef National Park, Wayne County, Utah. [<i>J. M. Porter</i> 8571; SJNM]
<i>Linanthus</i> (41 species)	<i>n</i> = 9	<i>L. aureus</i> (Nutt.) Greene. 2 miles north of Searchlight, along US Highway 95, Clark County, Nevada. [<i>J. M. Porter</i> 8822; AZ, SJNM]
	<i>n</i> = 9	* <i>L. bicolor</i> (Nutt.) Greene. Hayfork, Trinity County, CA. [<i>J. M. Porter & L. Machen</i> 10821; RSA]
	<i>n</i> = 9	<i>L. nuttallii</i> (Gray) Greene ex Mkn. 8 miles north of Granville, near Grey Peak, White Mountains, Greenlee County, Arizona. [<i>J. M. Porter & L. Machen</i> 9004; RSA, SJNM]
	<i>n</i> = 9	* <i>L. parryae</i> (Gray) Greene. Valyermo, Los Angeles County, CA. [<i>J. M. Porter</i> 10553 RSA]
<i>Loeseliastrum</i> (2 species)	<i>n</i> = 7	<i>L. matthewsii</i> (Gray) Timbr. San Bernardino County, California. [<i>J. M. Porter</i> 8869; SJNM]
	<i>n</i> = 7	<i>L. schottii</i> (Torr.) Timbr. 14 mi S of Parker, along AZ95, Yuma County, Arizona. [<i>J. M. Porter</i> 8856; RSA]
<i>Navarretia</i> (32 species)	<i>n</i> = 9	<i>N. breweri</i> (Gray) Greene, Columbine Pass, Uncompahgre Plateau, Montrose County, Colorado. [<i>J. M. Porter</i> 8599; RSA]
Diapensiaceae (Outgroup)	<i>n</i> = 6	*† <i>D. laponica</i> L. Mt. McKinley, Alaska. [<i>M. Fay</i> 325; ARIZ]
Fouquieriaceae (Outgroup)	<i>n</i> = ?	<i>F. columnaris</i> (Kell.) Hendrickson. Cult., The University Mall, University of Arizona Campus, Tucson, Arizona. [<i>J. M. Porter sn</i> ; SJNM]
	<i>n</i> = 6	<i>F. splendens</i> Engelm. Cult., Old Main, University of Arizona Campus, Tucson, Arizona. [<i>J. M. Porter sn</i> ; SJNM]

was employed to help assure that all of the most parsimonious solutions were obtained. Initially, CLOSEST addition and TBR (tree bisection-reconnection) swapping was performed. This was followed by 100 replicates of RANDOM addition and TBR branch swapping. Finally, 500 replicates of RANDOM addition were carried out with no swapping, followed by TBR swapping on the resulting set of trees (Maddison 1991). This should insure that all minimal length trees are found, even if multiple "islands" of equally parsimonious trees exist. The data matrix was analyzed with equal weights at all nucleotide positions and using an 1.1 to 1 transition/transversion bias (based on estimates from Polemoniaceae sequences). Insertions and/or deletions (indels) were coded initially as missing data. Subsequent analyses treated indels as a "fifth nucleotide" (GAPMODE=FIFTHSTATE option in PAUP) and as additional binary characters.

Robustness and Sampling

An assessment of the relative support for clades was performed using the decay index (Bremer 1988; Donoghue et al. 1992) and the bootstrap (Felsenstein 1985). Decay values were calculated for one, two and three steps longer than the shortest trees. My purpose is to identify only the weakly supported nodes. Bootstrap values were calculated from 100 replicate Fitch parsimony analyses, as implemented by PAUP 3.0, using heuristic searches based on CLOSEST addition and TBR branch swapping.

The distribution of tree-lengths, based on 10,000 randomly generated trees was evaluated for skewness (g_1). A strongly left-skewed tree length distribution has been suggested to indicate that a data set is highly structured and likely contains strong phylogenetic signal (Goodman et al. 1979; Fitch 1984; Werman 1986; Hillis and de Sa 1988; Hillis and Dixon 1989; Hillis and Huelsenbeck 1992; Hillis 1991; Huelsenbeck 1991). However, this test actually assesses whether the g_1 of a given data set deviates from the expected g_1 of random data (see also Kallersjo et al. 1992).

To investigate the sensitivity of these data to taxon sampling, an analysis based on resampling terminal taxa was conducted. Lanyon's jackknife is a heuristic tool to explore the effect of taxon sampling on internal branch support and tree structure (Lanyon 1985a, b; Felsenstein 1988; Siddall 1995). Lanyon (1985a) points out that trees generated during this jackknife procedure simulate possible outcomes had the investigator failed to sample a taxon or if extinction had occurred. Clearly, taxon sampling (inclusion) can have profound consequences on assessment of monophyly (Doyle and Donoghue 1987; Gauthier et al. 1988; Donoghue et al. 1989; Novacek 1991). The manually realigned data matrix was analyzed by sequentially re-

moving each member of the ingroup, one at a time, and performing a parsimony analysis (CLOSEST addition, HOLD= 2, TBR branch swapping) on the resulting data sets (pseudoreplicates). In all cases, only the strict-consensus tree from analyses of each pseudoreplicate was retained. The trees were compared and pseudoreplicate-consensus trees were generated. This was done by first returning the excluded taxon from each pseudoreplicate run back into the pseudoreplicate strict-consensus tree. The excluded taxon was placed in the same position it is found in the strict-consensus tree from the analysis of the entire data set. This is based on the assumption that the analysis of the full data set provides evidence for the taxon's placement. After the excluded taxon is returned to the tree, a consensus of the entire set of trees was obtained, using strict and majority rule consensus procedures.

RESULTS

ITS Sequence Variation

The ITS1 regions of all representatives of Polemoniaceae, as well as the outgroups, were consistently larger than the ITS2 regions. Within Polemoniaceae the ITS1 region is generally 249–253 bases. Extremes beyond this range include the sequences of *Gilia rigidula* at 242 base pairs (bp) and *Gilia scabra* at 262 bp. The outgroups possessed the largest ITS1 sequence of this analysis, 267 bp. In Polemoniaceae, percent G+C content of ITS1 ranged from 42.2% in *Cobaea baiurita* to 65.0% in *Acanthogilia gloriosa*. ITS2 sequences were mostly between 188 and 193 bases, but ranged from 187 bp in *Phlox standsburyi* and *Linanthus nuttallii* to 195 in *Gilia foetida*. The ITS2 region of the outgroups was slightly larger, 196 bp. Percent G+C content of ITS2 in Polemoniaceae varied from 48.3% in *Cobaea baiurita* to 61.8% in *Acanthogilia gloriosa*.

Alignment

Both ITS1 and ITS2 possess several highly conserved regions, flanked by more variable regions (Baldwin et al. 1995). The conserved regions are easily and nearly unambiguously aligned; however, the variable regions become increasingly difficult to align as pairwise distance increases. As a result, alignment of some taxa (i.e., outgroup taxa) should be viewed with skepticism. The variable regions of ITS in Polemoniaceae are frequently associated with short indels of one to a few nucleotides. Sequences of ITS1 and ITS2, aligned using variable weights in PILEUP produce similar alignments; however, the more variable regions seemed better aligned (i.e., there is a higher frequency of invariant sites) using GapWeight= 1.0 and GapLengthWeight= 0.1 (the aligned sequences us-

ing PILEUP are not shown). The manually realigned sequences are used in all of the following comparisons. Even so, the manually aligned sequence data set is not without alternative alignments and alignment in some regions remains ambiguous (Appendix I).

The aligned, combined sequences of ITS1 and ITS2 required, on average, gaps at 14.39% of the sites of any given sequence. The inclusion of the outgroup was responsible for gaps at 4.53% of these nucleotide positions. The combined data matrix, possesses 529 characters, of which 120 are invariant and 346 (65.28%) are potentially informative. The pair-wise levels of divergence range from 0.18% (between *Langloisia setosissima* subsp. *setosissima* and *Langloisia setosissima* subsp. *punctata*) to 42.4% (between *Diapensia lapponica* and *Cantua quercifolia*).

Phylogenetic Analysis

All of the parsimony analyses of the full, equal-weighted data set, with indels treated as missing, resulted in recovery of the same 1080 minimum-length trees. For the data set including only potentially informative sites, the most parsimonious trees were of 1552 steps (tree length including all sites is 1646), with a consistency index (C.I.) of 0.423, retention index (R.I.) of 0.622, and rescaled consistency index (R.C.) of 0.263. The strict consensus tree (Fig. 1) shows that the primary areas of disagreement in the set of minimal length trees involves 1) relationships involving *Ipomopsis sonora* and *Loeseliastrum matthewsii* in the Loeseliae Clade, 2) relationship of *Gilia campanulata* within the Linanthieae Clade, and 3) relationships between three clades of *Gilia* species as well as *Gilia splendens*, within the Gilieae Clade. Aside from the basal nodes, the phylogenetic estimate is not highly sensitive to coding of the indel regions or transition-transversion weighting. Figure 2 shows the strict consensus trees from analyses that 1) treat each indel position as a fifth character state while all other sites are equally weighted, and 2) remove all indels and alignment-ambiguous positions while imposing a 1.0:1.1 transition to transversion weighting. In both analyses many fewer trees are recovered, 12 and 8 respectively. The topologies differ primarily in the inferred relationships of *Cobaea*, *Cantua*, *Bonplandia* and *Acanthogilia* to the remainder of Polemoniaceae. Both of these trees are very similar in structure to the full, equal-weighted analysis.

Tree Support

Bootstrap.—The bootstrap 50% majority rule tree (not shown) is structurally identical to the strict consensus insofar as the majority rule tree is resolved. Many of the clades are supported with high bootstrap percentage values (see Fig. 1). Ten clades are supported by

values of 95% or greater. Hillis and Bull (1993), suggest that 70% bootstrap percentage values or greater might be a better estimate of a “95% confidence interval” for estimating the correct phylogeny. If this can be generalized to all data (but see Felsenstein and Kashino 1993), another 14 clades fall within the confidence limit.

Some portions of the trees, show little bootstrap support. In general the highest values are associated with the more terminal portions of the phylogeny. The values tend to decrease toward the more basal nodes. The only notable exception is the ancestral node of Polemoniaceae, that possesses a significant value of 99%.

Decay index.—The decay index is here employed to identify those clades that are weakly supported or may be prone to errors in phylogenetic estimation. Decay values shown in Fig. 1 range from one to three. In general, the most weakly supported nodes are the more basal nodes within Polemoniaceae. Again, the only exception to this trend is the ancestral node of Polemoniaceae.

Lanyon's jackknife.—The jackknife procedure required 53 analyses, the consensus of these analyses is shown in Fig. 3. The regions of the phylogeny that are affected by the inclusion of particular taxa are represented as unresolved polytomies. The lack of resolution indicates that a major portion of the phylogenetic estimate can change based solely on which terminal taxa are included in the analysis. However, it is the removal of *Acanthogilia* and the placement of *Cantua* that are responsible for much of the lack of resolution (i.e., the Loeseliae Clade). Resolved portions of the consensus tree represent relationships not effected by the current sampling. This includes the Gilieae Clade, the *Giliandra* clade, the *Gilia foetida*-*G. rigidula* clade, the Linanthieae Clade, the *Ipomopsis longiflora*-*I. multiflora*-*I. gunnisonii* clade, and the *Cobaea scanned*-*C. biaurita* clade.

Skewness of the distribution of random trees.—One of the primary limitations to evaluation of the skewness of the distribution of sets of random trees for a data set of this size is that no 95% confidence limit for skewness values (g_1) has been determined for data sets in excess of 25 terminal taxa (Hillis 1991; Hillis and Huelsenbeck 1992; Huelsenbeck 1991). However, as a conservative test, I have used the 95% confidence limit of the 25-taxon tree for those trees with 25 or more taxa (Hillis and Huelsenbeck 1992).

The distribution of tree length for a set of 10,000 trees based on all 56 sequences is significantly skewed, with a g_1 statistic of -0.6622 . Following Hillis (1991), I removed well-supported clades (as measured by bootstrap values of 95% or greater), except for one





Fig. 1. The strict-consensus of 1080 trees of 1646 steps (CI = 0.423, RC = 0.263), resulting from the PAUP analysis of the manually realigned PILEUP alignment of 53 members of Polemoniaceae and three outgroup members. The number below each internode is the decay index for that branch, ranging from d1 to d3. The frequency at which clades occur in 100 bootstrap replications is shown as a percentage, above each internode. Tribal classification of Grant (1959) is indicated by the symbol following each ingroup taxon name (+ = Cobaeae; ◆ = Cantuaeae; ▲ = Bonplandieae; □ = Polemonieae; ● = Gilieae). Intrageneric classification of *Gilia* is indicated by the numbered brackets (1 = sect. *Gilia*; 2 = sect. *Arachnion*; 3 = sect. *Saltugilia*; 4 = sect. *Giliastrum*; 5 = sect. *Giliaandra*). *Gilia scabra* is labeled “?” because Grant erroneously considered it synonymous with *Linanthus nuttallii*.

randomly selected representative, leaving 46 taxa. Another set of 10,000 random trees was generated. Again, the distribution is significantly skewed: $g1 = -0.4441$. This was followed by removal of clades with bootstrap values of 90% or greater, save one representative, leaving 31 taxa, and skewness was determined. Similarly, the distribution remains significantly skewed ($g1 =$

-0.4881). This process continued, using 21, 15, 12, 11, 10, nine, and eight taxa, each time removing members of the most well supported clades. In all cases, the $g1$ value significantly differs from the expected value of random data. Similar results (Porter unpublished) are obtained from permutation tail probability tests (Faith and Cranston 1991).



DISCUSSION

Sequence data from the two intergenic ITS spacers provide a compelling insight into some aspects of phylogeny. However, the phylogenetic inferences from ITS sequences contains both well-supported conclusions and highly questionable ones. In general, the weakly supported portions of the phylogeny are nodes associated with the earliest diverging lineages within the family. The lack of well-supported relationships at the base of the tree could be due to biological reasons or artifact. Biological reasons could include rapid diversification, low nucleotide substitution rate, or spurious long-branch attraction. Artifacts also could result from the difficulty in aligning the more divergent lineages and the treatment of the large blocks of question marks in the data matrix that result from indels. Fortunately, even with the ambiguity of relationships associated with the earliest diverging lineages, the robust portions of the phylogeny provides an understanding of relationships in Polemoniaceae that directly conflicts with current classification. This insight includes the pattern of diversification of the phlox family as well as issues of monophyly of the genera *Gilia*, *Ipomopsis* and *Linanthus*.

Monophyly of Polemoniaceae

The only well-supported node (based on bootstrap, decay and Lanyon's jackknife) in the basal region of the ITS phylogeny is the Polemoniaceae clade. Regardless of differential weighting of indels and/or transitions versus transversions, ITS sequences support the monophyly of Polemoniaceae. There is no support for the treatment of *Cobaea* as a separate family, Cobaeaceae (Don 1824). However, this study provides at best a weak test of the monophyly of Polemoniaceae. If the phylogeny is treated as unrooted, it can be recognized that a monophyletic Polemoniaceae is the result of the two longest branches (*Diapensia* and the *Fouqueria* clade) being united. These two lineages are characterized by nucleotide changes at nearly 30% of all potentially informative sites and between 37% and 50% of potentially informative sites not involving indels, raising the potential for long-branch attraction. This caution is tempered by the concordant support for a monophyletic Polemoniaceae from both chloroplast genes *matK* (Steele and Vilgalys 1994; Johnson et al. 1996) and *ndhF* (A. Prather, pers. comm.) and nuclear 18S sequences (as cited in Johnson et al. 1996).

Diversification of Polemoniaceae

The evolution and diversification of Polemoniaceae have been characterized by recent authors (e.g., Grant 1959) as a lineage of tropical origin that has radiated into the temperate regions. Grant postulates the inde-

Fig. 2. Lanyon's jackknife strict-consensus tree, identifying those regions of the phylogeny affected by taxon inclusion. Resolved portions of the tree indicate those nodes not influenced single taxon removal. Unresolved portions (compare with Fig 1) change based on the removal of a single terminal taxon.




Fig. 3. Comparison of two parsimony analyses differing in the weighting of gaps and transitions vs. transversions. Tree A is the strict consensus of 12 trees, resulting from the analysis in which no transition/transversion bias is imposed but all gap positions were treated as a fifth nucleotide state (GAPMODE = FIFTHSTATE option in PAUP). The analysis for tree B utilized a 1.1 to 1.0 transition to transversion bias but all gap and alignment-ambiguous positions were removed. Tree B is the strict consensus of the resulting eight trees.

pendent origin of two temperate lineages, tribes Polemoniaceae and Gilieae, from different tropical groups (tribes Cobaceae and Bonplandiaeae, respectively). This tropical-temperate distinction has persisted. The phylogenetic inference based in *matK*, presented by Steele and Vilgalys (1994), is suggested to support a tropical origin although the strict consensus tree shows ambiguous relationships between the included genera of tropical distribution. Steele and Vilgalys also propose a single "temperate" lineage, rather than two. Johnson et al. (1996) present a more complete *matK* phylogeny in which *Cantua*, *Cobaea*, and *Bonplandia* form a "tropical clade." Aside from *Acanthogilia*, the remaining genera are included in a large "temperate clade." Given that *Acanthogilia* is not tropical in distribution, the ancestral condition, if reconstructed using parsimony, is dependent on the distribution of the outgroups. Because Diapensiaceae are temperate in distribution and Fouquieriaceae are subtropical to tropical the ancestral character state assignment is ambiguous.

The phylogeny based on nucleotide sequences from ITS 1 and 2 show *Acanthogilia*, *Cantua*, *Cobaea*, and *Bonplandia* as a paraphyletic group, positioned as the earliest diverging lineages of the family. This could be interpreted as support for a tropical origin of Polemoniaceae. However, the relationships between these four genera, as inferred by ITS, have been shown to change based on taxon inclusion, character weighting (indels and transition-transversion) and how alignment-ambiguous sites are treated. It is difficult to argue that the ITS phylogeny is error free in this region. Given the uncertainty of relationships at the basal nodes the ancestral character state assignment is also uncertain. The question of a tropical origin for the family remains unanswered. Such determination again requires a more rigorous assessment of sister and outgroup relationships for Polemoniaceae than is presented here or in previous studies (e.g., Chase et al. 1993; Steele and Vilgalys 1994; Johnson et al. 1996).

The ITS spacer sequences support the remaining genera of Polemoniaceae (i.e., *Gilia*, *Ipomopsis*, *Eriastrum*, *Langloisia*, *Loeseliastrum*, *Loeselia*, *Allophyllum*, *Collomia*, *Navarretia*, *Phlox*, *Linanthus*, *Lepidactylon* and *Polemonium*) as a monophyletic group. This lineage has been referred to as the "Temperate Clade" (Steele and Vilgalys 1994; Johnson et al. 1996). While it is true that the majority of species in this clade occur in temperate regions, such a characterization may obscure the geographic patterns associated with diversification and dispersal within the lineage. This lineage includes several clades that are either wholly or partly tropical in distribution. For example, *Loeselia* ranges from southernmost Arizona, USA, through Mexico and Central America to Columbia, but all species occur in the tropics of Mexico.

Similarly, *Gilia* section *Giliastrum* (exclusive of *Gilia tenerrima*, *G. filiformis*, *G. campanulata*, *G. inyoensis*, and *G. maculata*) occurs not only in the temperate southwestern United States, but also the subtropical and tropical latitudes of both Mexico and South America (Peru, Bolivia, and Argentina). These examples serve to point out this clade includes tropical, subtropical, temperate, and even boreal lineages. It is not evident where the early diversification took place. Indeed, Grant's (1959) hypothesis of several independent lineages dispersing to and radiating within the temperate regions cannot be ruled out. For convenience I will refer to this clade as the Polemoniaceae Clade (Fig. 1).

The Polemoniaceae Clade is comprised of members from three of Grant's (1959) tribes (tribes Gilieae, Polemoniaceae, and Bonplandiaeae). None of these three tribes is supported as monophyletic (Fig. 1). Rather, three primary lineages, in addition to the genus *Polemonium*, make up this clade in all of the most parsimonious trees. The support for relationships between these clades is weak, but the membership of the three lineages is consistent regardless of weighting of transitions vs. transversions or indels and it is corroboration from other molecular phylogenetic studies (Steele and Vilgalys 1994; Prather 1995; Johnson et al. 1996). I refer to these clades as the Loeseliaeae, Gilieae, and Linanthiaeae Clades (Fig. 1). Note that the endings are used to distinguish the clades from generic names and are not intended to imply a particular hierarchic rank.

The Loeseliaeae Clade, which includes *Gilia* section *Giliandra*, *Gilia* section *Giliastrum* (exclusive of *Gilia campanulata*, *G. inyoensis*, *G. maculata* and *G. filiformis*), *Gilia scabra*, *Loeselia*, *Ipomopsis*, *Langloisia*, *Loeseliastrum*, and *Eriastrum*, is the sister group to the remaining members of the Polemoniaceae Clade. The members of this clade have traditionally been included within tribe Gilieae, with the exception of *Loeselia*, which has been placed in Bonplandiaeae (Grant 1959). The potential relationship between *Loeselia* and certain members of tribe Gilieae was recognized by Grant (1959), given his hypothesis that tribe Gilieae has its origin within or near *Loeselia*. Support, as measured by bootstrap, jackknife and decay, is low for many of the nodes within the Loeseliaeae Clade. Even so, the phylogenetic inference is highly consistent with that of *matK* (Johnson et al. 1996). Several noteworthy points can be made concerning the phylogenetic inferences based on ITS sequences. First, the two subspecies of *Langloisia setosissima* are supported as the sister group of *Eriastrum*. The sister group of this (the *Eriastrum*-*Langloisia* Clade) is *Loeseliastrum schottii*, supporting Timbrook's (1986) assertion that *Langloisia s.s.* and *Loeseliastrum* do not represent a monophyletic group. This represents a direct conflict between ITS and phylogenetic inferences from chloro-

plast *matK* (Johnson et al. 1996) and *trnL* intron (Porter, unpubl.) sequences, that support *Langloisia* and *Loeseliastrum* as sharing most recent common ancestry. Second, note that *Ipomopsis* is not monophyletic. The placement of *Ipomopsis tenuifolia* as sister to *Ipomopsis*, *Eriastrum*, *Langloisia*, and *Loeseliastrum* is not surprising, given that this member of the *Giliopsis* group was formerly included in *Loeselia* (Gray 1876; 1886). *Ipomopsis sonora* is also not unambiguously more closely related to other *Ipomopsis* species than to other genera. There may be three lineages, representing a grade, included in the current circumscription of *Ipomopsis*. The paraphyly of *Ipomopsis* is corroborated by chloroplast DNA sequence data (Johnson et al. 1996; Porter, unpubl.) and a more detailed ITS analysis of the Loeseliae Clade (Porter, in prep.). Third, the members of *Gilia* within this clade are neither monophyletic (although, *Gilia* section *Giliandra* is monophyletic) nor closely related to those *Gilia* associated with the type (i.e., *Gilia* section *Gilia*, type = *G. laciniata* R. & P.). In fact, *Gilia* section *Giliastrum*, as circumscribed by Grant (1959) is polyphyletic. In addition, *Gilia scabra*, considered by Grant (1959: 140) synonymous with *Linanthus nuttallii*, is suggested to share common ancestry with *Loeselia*.

The Gilieae Clade includes *Allophyllum*, *Collomia*, *Navarretia*, *Gilia* sections *Saltugilia*, *Arachnion*, *Gilia*, and one member of section *Giliastrum*. Historically, *Allophyllum* has been variously treated within *Collomia* (Bentham 1845; Gray 1870), close to *Collomia* in tribe Polemonieae (Grant and Grant 1955; Grant 1959), or included within *Gilia* (Nuttall 1848; Brand 1907; Mason and Grant 1948; Cronquist 1984), in tribe Gilieae. *Navarretia* has been of equally uncertain affiliation: included within *Gilia* by Gray (1870); allied with *Collomia*, *Cantua* and *Gymnosteris* in the *Collomia* tribe (Wherry 1945); or within Tribe *Gilieae*, near *Langloisia* and *Leptodactylon* (Grant 1959). It is perhaps not surprising that ITS sequences support a close relationship between *Navarretia*, *Collomia*, *Allophyllum* and a portion of what is now considered *Gilia*. Even so, this is a striking noncorrespondance with traditional taxonomy because it not only segregates only a portion of *Gilia* and allies it with *Navarretia* (both members of Grant's tribe Gilieae), but also strongly supports the relationships between this group and two members of tribe Polemonieae, *Allophyllum* and *Collomia*. The monophyly of the Gilieae Clade, supported here by ITS sequences, is both well supported in terms of bootstrap values and Lanyon's jack-knife analysis and also corroborated by sequence data from the chloroplast gene *matK* (Steele 1992; Steele and Vilgalys 1994; Johnson and Soltis 1995). The correspondence between ITS and *matK* gene phylogenies and the relative robustness of this portion of the ITS

phylogeny strongly supports the monophyly of this group.

Perhaps the most intriguing of the hypothesized relationships based on ITS sequence data involves the Linanthieae Clade. This clade is comprised of the genera *Linanthus*, *Leptodactylon* and *Phlox*, as well as four species of *Gilia* currently included in section *Giliastrum*. The Linanthieae Clade is well supported but very inconsistent with current classification. Although there is a long history associating *Linanthus* and *Leptodactylon* together (Gray 1870; Brand 1907; Wherry 1940; 1945; Grant, 1959) and they are currently both included in tribe Gilieae, *Phlox* (currently in tribe Polemonieae) has not been considered closely related. While floral morphology of *Leptodactylon* has been compared with that of *Phlox* (Gray 1870), no direct relationship was espoused. It is noteworthy that *Linanthus*, *Phlox* and *Leptodactylon* all possess pantoporate pollen grains (Marticorena 1961; Stuchlik 1967; Taylor and Levin 1975), very similar leaf flavonoid and phenolic chemistry (Smith et al. 1977; Dean et al. 1978; Smith et al. 1982), and possess opposite leaves, an unusual trait in Polemoniaceae.

The Linanthieae Clade is also intriguing because four species of *Gilia* (*G. campanulata*, *G. inyoensis*, *G. filiformis*, and *G. maculata*) are also members. These species have historically been closely aligned (see for example Brand 1907); however, most treatments prior to 1989 maintained *G. maculata* in *Linanthus*, while the remaining species were retained in *Gilia* sect. *Giliastrum*. Patterson (1989) has cautiously argued that, in spite of the unique traits possessed by *G. maculata*, there is sufficient morphological similarity between *Gilia campanulata*, *G. inyoensis*, *G. filiformis* and *G. maculata* to include them in the same genus, i.e., *Gilia*. ITS sequence data, presented here, supports in essence Patterson's proposition; however, this group of species is related not to members of *Gilia*, but to *Linanthus*, *Leptodactylon* and *Phlox*. It is also important to note that ITS sequences do not support the monophyly of these species.

Gilia is not monophyletic

The genus *Gilia* has remained, over the last 100 years, a recurring taxonomic problem, the circumscription of which has changed radically. Mason and Grant (1948) correctly point out that all of the herbaceous genera of the Polemoniaceae, with the exception of *Polemonium* and *Phlox*, have historically been placed in *Gilia*. One of the most broad interpretations of the genus was that of Asa Gray (1886). While recognizing that *Gilia* was "certainly a polymorphous . . . genus" (Gray 1870: 262), he included the currently recognized genera *Langloisia*, *Loeseliastrum*, *Gymnosteris*, *Leptodactylon*, *Linanthus*, *Navarretia*, *Ipomopsis* and *Er-*

iastrum within *Gilia*. The most recent classification of the family (Grant 1959) has maintained all these segregate genera. However, as noted above, Cronquist (Cronquist 1959; Cronquist 1984) has included *Ipomopsis* and *Allophyllum* within *Gilia*.

ITS sequence data provide insight into this conflict. The monophyly of representatives of *Gilia* sects. *Gilia* (here represented by *G. capitata* and *G. tricolor*), *Arachnion* (represented by *G. crassifolia*, *G. ochroleuca* and *G. flavocincta*), *Saltugilia* (represented by *G. stellata* and *G. scopulorum*), and *Giliastrum*, sensu Grant 1959 (represented by *G. tenerrima*) is supported (Fig. 2). Conspicuously absent from this clade are *Gilia scabra* and members of *Gilia* sects. *Giliastrum* (represented by *G. foetida*, *G. rigidula*, *G. latifolia*, *G. campanulata*, *G. inyoensis*, *G. filiformis*, and *G. maculata*), and *Giliandra* (represented by *G. caespitosa*, *G. subnuda*, *G. pentstemonoides*, *G. mcvickeriae*, *G. hutchinsifolia* and *G. nyensis*). All of these samples, except *G. campanulata*, *G. inyoensis*, *G. filiformis*, and *G. maculata* (in the Linanthieae Clade), are within the Loeseliae Clade. The most parsimonious placement of any one of these species into the Gilieae Clade requires between 27 and 51 steps (ACCTRANS reconstruction). Moreover, if representatives of *Gilia* sensu Grant (1959) are constrained to be monophyletic, three minimal-length trees are found of tree length 1693, 47 steps longer than the unconstrained analysis. The inclusion of representatives of *Ipomopsis* into *Gilia* (as proposed by Cronquist) does not ameliorate this situation. If *Ipomopsis* is constrained to be included in *Gilia*, sensu Cronquist, the eight trees found are 59 steps longer than the unconstrained analyses. Including additional taxa to make the representatives of *Gilia* monophyletic—if one considers the minimal-length ITS trees—will result in a group as morphologically diverse as Gray's circumscription of *Gilia*. If this is considered an unacceptable treatment (i.e., considering the Polemoniaceae Clade as a single genus, *Polemonium*), then the genus *Gilia* must be recognized in the more strict sense (i.e., *G. tenerrima* and its the sister group), it will be necessary to remove those clades that do not share common ancestry.

Conclusions

Sequences of the two internal transcribed spacers, ITS1 and ITS2, have been demonstrated to provide reliable phylogenetic estimates for portions of Polemoniaceae. The bootstrap, decay index, Lanyon's jackknife, and distribution of the tree length of random trees provide support for much of the phylogenetic inference. These include the *Giliandra* clade, *Gilia foetida*-*G. rigidula* clade, the *Allophyllum*-*Gilia* clade, the *Linanthus*-*Phlox*-*Leptodactylon* clade, and the *Ipomopsis gunnisonii*-*I. longiflora*-*I. multiflora* clade. Un-

fortunately, resolution of relationships among most of the major lineages, representing the basal radiation of the family, is suspect. Even given this limitation, ITS data unambiguously fail to support the current classification of Polemoniaceae that recognizes two temperate tribes, Polemoniaceae and Gilieae (Table 1). Rather, all of the genera, exclusive of *Bonplandia*, *Cantua* (and presumably *Huthia*), *Cobaea*, and *Acanthogilia*, are within a single clade, with members of Grant's tribes, Polemoniaceae and Gilieae variously related. Further, it has been shown that ITS sequence data does not support *Gilia*, as currently circumscribed, as a monophyletic group. The taxa now treated as *Gilia* represent at least five independent lineages, and potentially more (Fig. 1).

ACKNOWLEDGMENTS

Thanks to Bruce Baldwin, Chris Campbell, Mike Sanderson, Marty Wojciechowski for assistance with field collections; thanks to Mike Donoghue, Bruce Baldwin, Geeta Bharathan, Mike Sanderson, Carol von Dolen, Rob Robichaux, Kelly Steele, Dieter Wilken, and Alva Day for their helpful discussions and comments during the various stages of this project. The manuscript was greatly improved by the comments and suggestions provided by Mike Donoghue, Lucinda McDade, Leigh Johnson, Doug Soltis, Robert Thorne and one anonymous reviewer. Special thanks to RSAGB Molecular Lab Coordinator, Mary Debacon, for technical assistance.

A portion of this research was carried out in partial fulfillment of a Ph.D. degree at the Department of Ecology and Evolutionary Biology, University of Arizona, and a portion under NSF research grant, DEB-9509121.

LITERATURE CITED

- ABRAMS, L. 1951. Polemoniaceae, pp. 396–474. *In*: Illustrated flora of the Pacific States Vol. 3. Stanford University Press, Stanford, California.
- ANDERBERG, A. A. 1992. The circumscription of the Ericales, and their cladistic relationships to other families of "higher" Dicotyledons. *Syst. Bot.* 17: 660–675.
- BALDWIN, B. G. 1992. Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: an example from the Compositae. *Molec. Phylogen. Evol.* 1: 1–16.
- . 1993. Molecular phylogenetics of *Calycadenia* (Compositae) based on ITS sequences of nuclear ribosomal DNA: Chromosomal and morphological evolution reexamined. *Amer. J. Bot.* 80: 222–238.
- , M. J. SANDERSON, J. M. PORTER, M. F. WOJCIECHOWSKI, C. S. CAMPBELL, AND M. J. DONOGHUE. 1995. The ITS region of nuclear ribosomal DNA: a valuable source of evidence on angiosperm phylogeny. *Ann. Missouri Bot. Gard.* 82: 247–277.
- BENTHAM, G. 1845. Polemoniaceae, pp. 302–322. *In*: A. de Candolle, *Prodromus systematis naturalis regni vegetabilis*, Vol. 9. Paris, France.
- , AND J. D. HOOKER. 1876. Polemoniaceae, pp. 820–824. *In*: *Genera Plantarum*, Vol. 2. London, England.

- BRAND, A. 1907. Polemoniaceae, pp. 1–203. In: A. Engler, Das Pflanzenreich. 4 (250).
- . 1908. Polemoniaceae peruviana, pp. 174–175. In: I. Urban [ed.] *Plantae novae andinae imprimis Weberbauerianae* IV. Engler's Bot. Jhrb. 42.
- BREMER, K. 1988. The limits of amino acid sequence data in angiosperm reconstruction. *Evolution* 42: 795–803.
- CAMPBELL, D. R. 1989. Measures of selection in a hermaphroditic plant: Variation in male and female pollination success. *Evolution* 43: 318–334.
- . 1991. Effects of floral traits on sequential components of fitness in *Ipomopsis aggregata*. *Amer. Naturalist* 6: 713–737.
- CARLQUIST, S., V. M. ECKHART, AND D. C. MICHENER. 1984. Wood anatomy of Polemoniaceae. *Aliso* 10: 547–572.
- CHASE, M. W., AND H. H. HILLIS. 1991. Silica gel: an ideal material for field preservation of leaf samples for DNA studies. *Taxon* 40: 215–220.
- CHASE, M. W., D. E. SOLTIS, R. G. OLMSTEAD, D. MORGAN, D. H. LES, B. D. MISHLER, M. R. DUVAL, R. A. PRICE, H. G. HILLS, Y.-L. QIU, K. A. KRON, J. H. RETTIG, E. CONTI, J. D. PALMER, J. R. MANHART, K. J. SYTSMAN, H. G. MICHAELS, W. J. KRESS, K. G. KAROL, W. D. CLARK, M. HEDREN, B. S. GAUT, R. K. JANSEN, K.-J. KIM, C. F. WIMPEE, J. F. SMITH, G. R. FURNIER, S. H. STRAUSS, Q.-Y. XIANG, G. M. PLUNKETT, P. S. SOLTIS, S. M. SWENSEN, S. E. WILLIAMS, P. A. GADEK, C. J. QUINN, L. E. EGUIARTE, E. GOLDENBERG, G. H. LEARN, JR., S. W. GRAHAM, S. C. H. BARRETT, S. DAYANANDAN, AND V. A. ALBERT. 1993. Phylogenetics of seed plants: an analysis of nucleotide sequences from the plastid gene *rbcL*. *Ann. Missouri Bot. Gard.* 80: 528–580.
- CHUANG, T.-I., W. C. HSIEH, AND D. H. WILKEN. 1978. Contribution of pollen morphology to systematics of *Collomia* (Polemoniaceae). *Amer. J. Bot.* 65: 450–458.
- COYNE, J. A. 1974. The evolutionary origin of hybrid inviability. *Evolution* 28: 505–506.
- CRONQUIST, A. 1959. Polemoniaceae, pp. 1–510. In: C. L. Hitchcock, A. Cronquist, M. Ownbey and J. W. Thompson [eds.] *Vascular plants of the Pacific Northwest. Part 4: Ericaceae through Campanulaceae*. University of Washington Publications in Botany Vol. 17, Pt. 4. Seattle, Washington.
- . 1981. An integrated system of classification of flowering plants. Columbia University Press. New York, New York. 1262 p.
- . 1984. Polemoniaceae, pp. 86–155. In: Cronquist, A., A. Holmgren, N. Holmgren, J. Reveal, and P. Holmgren [eds.], *Intermountain Flora: Vascular Plants of the Intermountain West, USA. Vol. 4. Subclass Asteridae (except Asteraceae)*. New York Botanical Garden. Bronx, New York.
- DAWSON, M. L. 1936. The floral morphology of the Polemoniaceae. *Amer. J. Bot.* 23: 501–511.
- DAY, A. 1965. The evolution of a pair of sibling allotetraploid species of Cobwebby Gilias (Polemoniaceae). *Aliso* 6(1): 25–75.
- , AND R. MORAN. 1986. *Acanthogilia*, a new genus of Polemoniaceae from Baja California, Mexico. *Proc. Calif. Acad. Sci.* 44: 111–126.
- DEAN, F. M., A. M. COSTA, J. B. HARBORNE, AND D. M. SMITH. 1978. Leptodactylone, a yellow coumarin from *Leptodactylon* and *Linanthus* species. *Phytochemistry* 17: 505–509.
- DON, D. 1822. Observations on the natural family of plants called Polemoniaceae; with descriptions of the genera belonging to it, and of a genus improperly referred to that order by botanists. *Edinburgh Philos. J.* 7: 283–291.
- . 1824. Observations on a new natural family of plants to be called Cobeeaceae. *Edinburgh Philos. J.* 10: 109–112.
- DONOGHUE, M. J. 1989. Phylogenies and the analysis of evolutionary sequences, with examples from seed plants. *Evolution* 43: 1137–1156.
- , J. A. DOYLE, J. GAUTHIER, A. KLUGE, AND T. ROWE. 1989. The importance of fossils in phylogeny reconstruction. *Annual Rev. Ecol. Syst.* 20: 431–460.
- , R. G. OLMSTEAD, J. F. SMITH, AND J. D. PALMER. 1992. Phylogenetic relationships of Dipsacales based on *rbcL* sequences. *Ann. Missouri Bot. Gard.* 79: 333–245.
- DOYLE, J. A., AND M. J. DONOGHUE. 1987. The importance of fossils in elucidating seed plant phylogeny and macroevolution. *Rev. Palaeobot. Palynol.* 50: 63–95.
- DOYLE, J. J. 1992. Gene trees and species trees: Molecular systematics as one character taxonomy. *Syst. Bot.* 17: 144–163.
- , AND J. L. DOYLE. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19: 11–15.
- DWYER, K. G., M. A. BALENET, J. B. NASRALLAH, AND M. E. NASRALLAH. 1991. DNA sequences of self-incompatibility genes from *Brassica campestris* and *B. oleracea*: polymorphism predating speciation. *Pl. Molec. Biol.* 16: 481–486.
- ELLSTRAND, N. C. 1983. Floral formula inconsistency within and among plants and populations of *Ipomopsis aggregata* (Polemoniaceae). *Bot. Gaz. (Crawfordsville)* 144: 119–123.
- , AND R. J. MITCHELL. 1988. Spatial and temporal patterns of floral inconsistency in plants and populations of *Ipomopsis aggregata* (Polemoniaceae). *Bot. Gaz. (Crawfordsville)* 149: 209–212.
- ENGLER, A., AND E. GILG. 1924. *Syllabus der Pflanzenfamilien*. 10th Ed. Berlin, Germany.
- FAITH, D. P., AND P. S. CRANSTON. 1991. Could a cladogram this short have arisen by chance alone?: On permutation tests for cladistic structure. *Cladistics* 7: 1–28.
- FELSENSTEIN, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- . 1988. Phylogenies and quantitative characters. *Annual Rev. Ecol. Syst.* 19: 445–471.
- , AND H. KASHINO. 1993. Is there something wrong with the bootstrap on phylogenies? A reply to Hillis and Bull. *Syst. Biol.* 42: 193–199.
- FENG, D. F., AND R. F. DOOLITTLE. 1987. Progressive sequence alignment as a prerequisite to correct phylogenetic trees. *J. Molec. Evol.* 25: 351–360.
- FITCH, W. M. 1984. Cladistic and other methods: problems, pitfalls, and potentials, pp. 221–252. In: T. Duncan and T. Stuessy [eds.] *Cladistics: Perspectives on the reconstruction of evolutionary history*. Columbia University Press. New York, New York.
- GALLEN, C., AND P. G. KEVAN. 1983. Bumblebee foraging and floral scent dimorphism: *Bombus kirbyellus* Curtis (Hymenoptera: Apidae) and *Polemonium viscosum* Nutt. (Polemoniaceae). *Canad. J. Zool.* 61: 1207–1213.
- , AND M. L. STANTON. 1989. Bumblebee pollination and floral morphology: factors influencing pollen dispersal in the alpine sky pilot, *Polemonium viscosum* (Polemoniaceae). *Amer. J. Bot.* 76: 419–426.
- GAUTHIER, J., A. G. KLUGE, AND T. ROWE. 1988. Amniote phylogeny and the importance of fossils. *Cladistics* 4: 105–209.
- GOODMAN, M., J. CZELUSNIAK, AND G. W. MOORE. 1979. Further remarks on the parameter of gene duplication and expression events in parsimony reconstructions. *Syst. Zool.* 28: 379–385.
- GRANT, A., AND V. GRANT. 1955. The genus *Allophyllum* (Polemoniaceae). *El Aliso* 3: 93–110.
- GRANT, V. 1950. Genetic and taxonomic studies in *Gilia*. I. *Gilia capitata*. *El Aliso* 2: 239–316.
- . 1954a. Genetic and taxonomic studies in *Gilia*. IV. *Gilia achilleaefolia*. *El Aliso* 3: 1–18.
- . 1954b. Genetic and taxonomic studies in *Gilia*. V. Interspecific relationships in the leafy-stemmed Gilias. *El Aliso* 3: 35–49.
- . 1957. The plant species in theory and practice. pp. 39–80.

- In: E. Mayr [ed.], The species problem. American Association for the Advancement of Science, Washington, D.C.
- . 1959. Natural history of the phlox family. Martinus Nijhoff. The Hague, Netherlands.
- . 1979. Character coherence in natural hybrid populations of plants. *Bot. Gaz. (Crawfordsville)* **140**: 443–448.
- , AND A. GRANT. 1960. Genetic and taxonomic studies in *Gilia*. XI. Fertility relationships of the diploid Cobwebby *Gilias*. *El Aliso* **4**: 435–481.
- , AND D. H. WILKEN. 1987. Secondary integration between *Ipomopsis aggregata candida* and *I. a. collina* (Polemoniaceae) in Colorado. *Bot. Gaz. (Crawfordsville)* **148**: 372–378.
- , AND K. GRANT. 1965. Flower pollination in the phlox family. Columbia University Press. New York, New York. 180 p.
- GRAY, A. 1870. Revision of North American Polemoniaceae. *Proc. Amer. Acad. Arts* **8**: 247–282.
- . 1876. Miscellaneous botanical contributions. *Proc. Amer. Acad. Arts* **11**: 71–92.
- . 1886. Polemoniaceae, pp. 128–151 In: Synoptical flora of North America, Ed 2, Vol. 1 and Supplement: 406–412. Smithsonian Institution. Washington, D.C.
- HARBORNE, J. B. 1980. New experimental approaches to plant chemosystematics, pp. 39–70. In: F. A. Bisby, J. G. Vaughn and C. A. Wright [eds.], Chemosystematics: principles and practice. Academic Press, New York, New York.
- , AND D. M. SMITH. 1978. Correlations between anthocyanin chemistry and pollination ecology in Polemoniaceae. *Biochem. Syst. Ecol.* **6**: 127–130.
- HIGGINS, D. G., AND P. M. SHARP. 1987. Fast and sensitive multiple sequence alignments on a microcomputer. *Computer Appl. Biol. Sci.* **5**: 151–153.
- HILLIS, D. M. 1991. Discriminating between signal and random noise in DNA sequences, pp. 278–294. In: M. M. Miyamoto and J. Cracraft [eds.], Phylogenetic analysis of DNA sequences. Oxford University Press, Inc. New York, New York.
- , AND J. J. BULL. 1993. An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analyses. *Syst. Biol.* **42**: 182–192.
- , AND R. DE SA. 1988. Phylogeny and taxonomy of the *Rana palmipes* group (Salientia: Ranidae). *Herpetological Monog.* **2**: 1–26.
- , AND M. T. DIXON. 1989. Vertebrate phylogeny: evidence from 28S ribosomal DNA sequences, pp. 355–367. In: B. Fernholm, K. Bremer, and H. Jornvall [eds.], The hierarchy of life. Molecules and morphology in phylogenetic analysis. Elsevier Science Publishers B. V. Amsterdam, Netherlands.
- , AND J. P. HUELSENBECK. 1992. Signal, noise, and reliability in molecular phylogenetic analyses. *J. Heredity* **83**: 189–195.
- HUELSENBECK, J. P. 1991. Tree-length distribution skewness: an indicator of phylogenetic information. *Syst. Zool.* **40**: 257–270.
- HUMBOLT, A., A. BONLAND, AND C. S. KUNTH. 1823. Nova genera et species plantarum. Vol. 6, pp. 83–85. Paris.
- JOHNSON, L. A., AND D. E. SOLTIS. 1995. Phylogenetic inference in Saxifragaceae sensu strictu and *Gilia* (Polemoniaceae) using *matK* sequences. *Ann. Missouri Bot. Gard.* **82**: 149–175.
- , J. L. SCHULTZ, D. E. SOLTIS, AND P. S. SOLTIS. 1996. Monophyly and generic relationships of Polemoniaceae based on *matK* sequences. *Amer. J. Bot.* **83**: 1207–1224.
- KALLERSJO, M., J. S. FARRIS, A. G. KLUGE, AND C. BULT. 1992. Skewness and permutation. *Cladistics* **8**: 275–287.
- KIREDJAD, M., AND D. M. SMITH. 1985. A chemosystematic investigation on *Ipomopsis*. *Biochem. Syst. Ecol.* **13**: 141–143.
- KUNTZE, O. 1891. Polemoniaceae, pp. 432–434. In: *Revisio generum plantarum* Vol. 2. Arthur Felix, Leipzig, Germany.
- LANYON, S. M. 1985a. Detecting internal inconsistencies in distance data. *Syst. Zool.* **43**: 397–403.
- . 1985b. Molecular perspective on higher-level relations in the Tyrannoidea (Aves). *Syst. Zool.* **43**: 404–418.
- LEVIN, D. A. [ed.]. 1979. Benchmark Papers in Genetics, Vol. 11. Hybridization: an Evolutionary Perspective. Dowden Hutchinson and Ross, Inc. Stroudsburg, Pennsylvania. 320 p.
- LINNAEUS, C. 1753. Species Plantarum, Vol. 2. Stockholm, Sweden.
- LISTON, A., AND L. H. RIESEBERG. 1990. A method for collecting dried plant specimens for DNA and isozyme analysis, and the results of a field test in Xinjiang, China. *Ann. Missouri Bot. Gard.* **77**: 859–863.
- LOEBLICH, A. R. 1964. The pollen grain morphology of *Collomia* as a taxonomic tool. *Madroño* **17**: 205–216.
- LORD, E. M., AND K. J. ECKARD. 1986. Ultrastructure of the dimorphic pollen and stigmas of the cleistogamous species *Collomia grandiflora* (Polemoniaceae). *Protoplasma* **132**: 12–22.
- MADDISON, D. R. 1991. The discovery and importance of multiple islands of most-parsimonious trees. *Syst. Zool.* **40**: 315–328.
- MANIATIS, T., E. F. FRITSCH, AND J. SAMBROOK. 1982. Molecular cloning: A laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, New York.
- MARTICORENA, C. 1961. Morfología de los granos de polen de las Polemoniaceae Chilenas. *Gayana* **2**: 5–12.
- MARTIN, P. G., AND J. M. DOWD. 1991. Studies of angiosperm phylogeny using protein sequences. *Ann. Missouri Bot. Gard.* **78**: 296–337.
- MASON, H. L. 1945. The genus *Eriastrum* and the influence of Bentham and Gray upon the problem of generic confusion in Polemoniaceae. *Madroño* **8**: 33–64.
- , AND A. GRANT. 1948. Some problems in the genus *Gilia*. *Madroño* **9**: 201–220.
- NASH, G. V. 1903. A revision of the family Fouquieriaceae. *Bull. Torrey Bot. Club* **30**: 449–459.
- NEEDLEMAN, S. B., AND C. D. WUNCH. 1987. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Molec. Biol.* **48**: 443–453.
- NEIGEL J. E., AND J. C. AVISE. 1986. Phylogenetic relationships of mitochondrial DNA under various demographic models of speciation, pp. 515–534. In: S. Karlin and E. Nevo [eds.], Evolutionary processes and theory. Academic Press, New York, New York.
- NOVACEK, M. J. 1991. “All tree histograms” and the evaluation of cladistic evidence: Some ambiguities. *Cladistics* **7**: 345–349.
- NUTTALL, T. 1848. Polemoniaceae, pp. 185–187. In: The genera of North American plants Vol. 1. D. Heart, Philadelphia, Pennsylvania.
- OLMSTEAD, R. G., H. J. MICHAELS, K. M. SCOTT, AND J. D. PALMER. 1992. Monophyly of the Asteridae and identification of their major lineages inferred from DNA sequences of *rbcL*. *Ann. Missouri Bot. Gard.* **79**: 294–265.
- PÄÄBO, S. 1990. Amplifying ancient DNA, pp. 159–166. In: PCR protocols: A guide to methods and applications. M. Innis, D. Gelfand, J. Sninsky, and T. White [eds.], Academic Press, San Diego, California.
- , R. G. HIGUCHI, AND A. C. WILSON. 1989. Ancient DNA and the polymerase chain reaction. *J. Biol. Chem.* **264**: 9709–9712.
- PAIGE, K. N., AND T. G. WHITHAM. 1985. Individual and population shifts in flower color by scarlet *gilia* (*Ipomopsis aggregata*) a mechanism for pollinator tracking. *Science* **227**: 315–317.
- , AND ———. 1987a. Flexible life history traits: Shifts by scarlet *gilia* in response to pollinator abundance. *Ecology* **68**: 1691–1695.
- , AND ———. 1987b. Overcompensation in response to mammalian herbivory: the advantage of being eaten. *Amer. Naturalist* **129**: 407–416.
- PATTERSON, R. 1989. Taxonomic relationships of *Gilia maculata* (Polemoniaceae). *Madroño* **36**: 15–27.
- . 1993. Polemoniaceae, pp. 824–852. In: J. C. Hickman

- In: E. Mayr [ed.], The species problem. American Association for the Advancement of Science, Washington, D.C.
- . 1959. Natural history of the phlox family. Martinus Nijhoff. The Hague, Netherlands.
- . 1979. Character coherence in natural hybrid populations of plants. *Bot. Gaz. (Crawfordsville)* **140**: 443–448.
- , AND A. GRANT. 1960. Genetic and taxonomic studies in *Gilia*. XI. Fertility relationships of the diploid Cobwebby *Gilias*. *El Aliso* **4**: 435–481.
- , AND D. H. WILKEN. 1987. Secondary integration between *Ipomopsis aggregata candida* and *I. a. collina* (Polemoniaceae) in Colorado. *Bot. Gaz. (Crawfordsville)* **148**: 372–378.
- , AND K. GRANT. 1965. Flower pollination in the phlox family. Columbia University Press. New York, New York. 180 p.
- GRAY, A. 1870. Revision of North American Polemoniaceae. *Proc. Amer. Acad. Arts* **8**: 247–282.
- . 1876. Miscellaneous botanical contributions. *Proc. Amer. Acad. Arts* **11**: 71–92.
- . 1886. Polemoniaceae, pp. 128–151 In: Synoptical flora of North America, Ed 2, Vol. 1 and Supplement: 406–412. Smithsonian Institution. Washington, D.C.
- HARBORNE, J. B. 1980. New experimental approaches to plant chemosystematics, pp. 39–70. In: F. A. Bisby, J. G. Vaughn and C. A. Wright [eds.], Chemosystematics: principles and practice. Academic Press, New York, New York.
- , AND D. M. SMITH. 1978. Correlations between anthocyanin chemistry and pollination ecology in Polemoniaceae. *Biochem. Syst. Ecol.* **6**: 127–130.
- HIGGINS, D. G., AND P. M. SHARP. 1987. Fast and sensitive multiple sequence alignments on a microcomputer. *Computer Appl. Biol. Sci.* **5**: 151–153.
- HILLIS, D. M. 1991. Discriminating between signal and random noise in DNA sequences, pp. 278–294. In: M. M. Miyamoto and J. Cracraft [eds.], Phylogenetic analysis of DNA sequences. Oxford University Press, Inc. New York, New York.
- , AND J. J. BULL. 1993. An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analyses. *Syst. Biol.* **42**: 182–192.
- , AND R. DE SA. 1988. Phylogeny and taxonomy of the *Rana palmipes* group (Salientia: Ranidae). *Herpetological Monog.* **2**: 1–26.
- , AND M. T. DIXON. 1989. Vertebrate phylogeny: evidence from 28S ribosomal DNA sequences, pp. 355–367. In: B. Fernholm, K. Bremer, and H. Jornvall [eds.], The hierarchy of life. Molecules and morphology in phylogenetic analysis. Elsevier Science Publishers B. V. Amsterdam, Netherlands.
- , AND J. P. HUELSENBECK. 1992. Signal, noise, and reliability in molecular phylogenetic analyses. *J. Heredity* **83**: 189–195.
- HUELSENBECK, J. P. 1991. Tree-length distribution skewness: an indicator of phylogenetic information. *Syst. Zool.* **40**: 257–270.
- HUMBOLT, A., A. BONLAND, AND C. S. KUNTH. 1823. Nova genera et species plantarum. Vol. 6, pp. 83–85. Paris.
- JOHNSON, L. A., AND D. E. SOLTIS. 1995. Phylogenetic inference in Saxifragaceae sensu strictu and *Gilia* (Polemoniaceae) using *matK* sequences. *Ann. Missouri Bot. Gard.* **82**: 149–175.
- , J. L. SCHULTZ, D. E. SOLTIS, AND P. S. SOLTIS. 1996. Monophyly and generic relationships of Polemoniaceae based on *matK* sequences. *Amer. J. Bot.* **83**: 1207–1224.
- KALLERSJO, M., J. S. FARRIS, A. G. KLUGE, AND C. BULT. 1992. Skewness and permutation. *Cladistics* **8**: 275–287.
- KIREDJAD, M., AND D. M. SMITH. 1985. A chemosystematic investigation on *Ipomopsis*. *Biochem. Syst. Ecol.* **13**: 141–143.
- KUNTZE, O. 1891. Polemoniaceae, pp. 432–434. In: Revisio generum plantarum Vol. 2. Arthur Felix, Leipzig, Germany.
- LANYON, S. M. 1985a. Detecting internal inconsistencies in distance data. *Syst. Zool.* **43**: 397–403.
- . 1985b. Molecular perspective on higher-level relations in the Tyrannoidea (Aves). *Syst. Zool.* **43**: 404–418.
- LEVIN, D. A. [ed.]. 1979. Benchmark Papers in Genetics, Vol. 11. Hybridization: an Evolutionary Perspective. Dowden Hutchinson and Ross, Inc. Stroudsburg, Pennsylvania. 320 p.
- LINNAEUS, C. 1753. Species Plantarum, Vol. 2. Stockholm, Sweden.
- LISTON, A., AND L. H. RIESEBERG. 1990. A method for collecting dried plant specimens for DNA and isozyme analysis, and the results of a field test in Xinjiang, China. *Ann. Missouri Bot. Gard.* **77**: 859–863.
- LOEBLICH, A. R. 1964. The pollen grain morphology of *Collomia* as a taxonomic tool. *Madroño* **17**: 205–216.
- LORD, E. M., AND K. J. ECKARD. 1986. Ultrastructure of the dimorphic pollen and stigmas of the cleistogamous species *Collomia grandiflora* (Polemoniaceae). *Protoplasma* **132**: 12–22.
- MADDISON, D. R. 1991. The discovery and importance of multiple islands of most-parsimonious trees. *Syst. Zool.* **40**: 315–328.
- MANIATIS, T., E. F. FRITSCH, AND J. SAMBROOK. 1982. Molecular cloning: A laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, New York.
- MARTICORENA, C. 1961. Morfología de los granos de polen de las Polemoniaceae Chilenas. *Gayana* **2**: 5–12.
- MARTIN, P. G., AND J. M. DOWD. 1991. Studies of angiosperm phylogeny using protein sequences. *Ann. Missouri Bot. Gard.* **78**: 296–337.
- MASON, H. L. 1945. The genus *Eriastrum* and the influence of Bentham and Gray upon the problem of generic confusion in Polemoniaceae. *Madroño* **8**: 33–64.
- , AND A. GRANT. 1948. Some problems in the genus *Gilia*. *Madroño* **9**: 201–220.
- NASH, G. V. 1903. A revision of the family Fouquieriaceae. *Bull. Torrey Bot. Club* **30**: 449–459.
- NEEDLEMAN, S. B., AND C. D. WUNCH. 1987. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Molec. Biol.* **48**: 443–453.
- NEIGEL J. E., AND J. C. AVISE. 1986. Phylogenetic relationships of mitochondrial DNA under various demographic models of speciation, pp. 515–534. In: S. Karlin and E. Nevo [eds.], Evolutionary processes and theory. Academic Press, New York, New York.
- NOVACEK, M. J. 1991. “All tree histograms” and the evaluation of cladistic evidence: Some ambiguities. *Cladistics* **7**: 345–349.
- NUTTALL, T. 1848. Polemoniaceae, pp. 185–187. In: The genera of North American plants Vol. 1. D. Heart, Philadelphia, Pennsylvania.
- OLMSTEAD, R. G., H. J. MICHAELS, K. M. SCOTT, AND J. D. PALMER. 1992. Monophyly of the Asteridae and identification of their major lineages inferred from DNA sequences of *rbcL*. *Ann. Missouri Bot. Gard.* **79**: 294–265.
- PÄÄBO, S. 1990. Amplifying ancient DNA, pp. 159–166. In: PCR protocols: A guide to methods and applications. M. Innis, D. Gelfand, J. Sninsky, and T. White [eds.], Academic Press, San Diego, California.
- , R. G. HIGUCHI, AND A. C. WILSON. 1989. Ancient DNA and the polymerase chain reaction. *J. Biol. Chem.* **264**: 9709–9712.
- PAIGE, K. N., AND T. G. WHITHAM. 1985. Individual and population shifts in flower color by scarlet *gilia* (*Ipomopsis aggregata*) a mechanism for pollinator tracking. *Science* **227**: 315–317.
- , AND ———. 1987a. Flexible life history traits: Shifts by scarlet *gilia* in response to pollinator abundance. *Ecology* **68**: 1691–1695.
- , AND ———. 1987b. Overcompensation in response to mammalian herbivory: the advantage of being eaten. *Amer. Naturalist* **129**: 407–416.
- PATTERSON, R. 1989. Taxonomic relationships of *Gilia maculata* (Polemoniaceae). *Madroño* **36**: 15–27.
- . 1993. Polemoniaceae, pp. 824–852. In: J. C. Hickman

Appendix I

	10	20	30	40	50	60	70	80	90	100	110
<i>Fouquieria columnaris</i>	TCGATACCTGCCTAGCAGAACGACCTGTGAACTCGT	CTCCAAAATGAGGTGAT	TCGC	GGGTGGGC	CTCC	GGTCCCCCGTGGGGAA	CCTTCTCGGGAAAGCG				
<i>Fouquieria splendens</i>	TCGATACCTGCCTAGCAGAACGACCTGTGAACTCGT	CTCCAAAATGAGGTGAT	TCAY	GGGTGGGC	CTCT	GGTCCCTCCCGTGCAGAA	CCTTATCGGGAAACG				
<i>Diapensia lapponica</i>	TCGATATCTGCACAGCAGAACGACCCGCGAACTAGTATTTCACTTTG	GGAGAT	GCATTTGGGCGGTC	TAAT	CCCCCT	CCTTTGTGCT	CTCCAC	GGC			
<i>Acanthogilia gloriosa</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTGTATTTCAAACTCGGTGGC	ACG	GGCTTTGGC	CTC	GGTCCCT	CGCCCGTTCC	CCTT	CCGGA			
<i>Cantua quercifolia</i>	TCGAAA	CTGTCCCGAAACACGACCCGTGAACTTGTATTCTCAYATCGGGT	GC	ATG	GTCGATCGG	CTTC	GGTCCCT	CGGCCATTGCTACCCCGTTGGTGGT	GT		
<i>Bonplandia geminiflora</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTCGT	TAAACATATCGGG	GC	AC	TCGGC	CTTC	GGA	CT	CGGTCTCG	GTGCTTT	GCCCTCGT
<i>Cobaea biaurita</i>	TCGAAACCTGCATRGCAGAAAGACCCYGTGAACTTAT	TTTTAAACTCAG	TGGC	ATG	ATTGGGGA	CTYC	GGTCCCT	CCTTTGTTGY	CCCTTT	RCTTGTG	
<i>Cobaea scandens</i>	TCGAAACCTGCATGGCAGAAAGACCTGTGAACTTAT	TTTTAAACTCAG	TGGC	ATG	ATTGGGGA	CTTC	GGTCCCT	CATTTGTTGC	CCCTTT	GCT	GTTG
<i>Gilia caespitosa</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	AGCATGGTC	CTTT	GGTCCA	AACTCGTTGC	CCCCA	GCTGGGT		
<i>Gilia subnuda</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	AGCATGGTC	CTTT	GGTCCA	AACTCGTTGC	CCCCA	GCTGGGT		
<i>Gilia pentstemonoides</i>	TCGAAACCTGCCTAGCAGAMCGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GGCATGSS	CTTT	GGTCCCT	AGCCCGTTGC	CCCCA	GCTGGGT		
<i>Gilia mcvickerae</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GRCATGGTC	CTTT	GGTCCCT	AGTCTGTTGC	CCCCA	GCTGGGT		
<i>Gilia nyensis</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	AGCATGGT	CTTC	GGTCCCT	AGTCTGTTGC	CCCCA	GCTGGGT	T	
<i>Gilia hutchinsifolia</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TTGGGTGGC	ACG	AGCATGGTC	CTTT	GGTCCCT	ACCTCGTTGC	CCCCA	GCTGGGC		
<i>Gilia latifolia</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GGCGTGTGTA	CTCC	GGTCCCT	CGTCTGTTGC	CCCTT	GCTGGGTGT		
<i>Gilia foetida</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGA	GC	ACG	GTCGTGGAC	CTTT	GGTCCCT	TGCTCG	TGC	CCC	A
<i>Gilia rigidula</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GTCGTGGAC	CTTC	GGTCCA	CGATCGATGC	CCC	A	GCTGGGA	
<i>Gilia scabra</i>	TCGAAACCTGTCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GACGTGGAC	CTCC	GGTCCCT	TGGCCGTTGC	ACCCA	GCCGGAT		
<i>Loeselia glandulosa</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	GCG	GGCGTGGC	CTTTT	GGTCCCT	CGTCTGTTGC	CCCCA	GCCGACA		
<i>Ipomopsis longiflora</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GGTGTGGTC	CTAT	GGTCCCT	CCCCCGTTGC	CCCTA	GCTGACA		
<i>Ipomopsis gunnisonii</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GGTGTGGTC	CTAT	GGTCCCT	CACCCATTGC	CCCTA	GCTGACG		
<i>Ipomopsis multiflora</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GGTGTGGTC	CTAT	GGTCCCT	CCCCCGTTGC	CCCTA	GCTGCCA		
<i>Ipomopsis tenuifolia</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	AC	GGCTGGTC	CTTTT	GGTCCCT	CGCCCGTTGC	CCCTA	GCTGACT		
<i>Ipomopsis sonorae</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGT	GCTGGC	GTGGGGTC	CTTC	GGTCCCT	CGTCTGTTGC	CCCTA	GCCGACA		
<i>Eriastrum diffusum</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TTGGGCGGC	ACG	GGTGTGGC	CTCTAT	GGTCCCT	CGTCTGTTGC	CCCTA	GCCGACA		
<i>Loeseliastrum matthewsii</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	GGTGTGGTC	CTWT	GGTCCCT	CGTCTGTTGC	CCCTA	GCTGACA		
<i>Loeseliastrum schottii</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TTGGGTGGC	ACG	GGTGTGGTC	CTTT	GGTCCCT	CGTCTGTTGC	CCCTA	GCTGACA		
<i>Langloisia setosissima</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TTGGGTGGC	ACG	GGCTGGTC	CTCT	GGTCCCT	CGTCTGTTGC	CCCTA	GCCGGCA		
<i>Langloisia set. punctata</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TTGGGTGGC	ACG	GGCTGGTC	CTCC	GGTCCCT	CGTCTGTTGC	CCCTA	GCCGGCA		
<i>Gilia stellata</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	GCCSTTGTATCTC		GGTACG	CGCTCGTTCC	CCCAAC	GCTAGGA		
<i>Gilia crassifolia</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	GCG	GGCTGTATCTC		GGTACT	CGCTCGTTCC	CCCAAC	GCTAGGA		
<i>Gilia scopulorum</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	GCG	GGCTGTATCTC		GGTACT	CGCTCGTTCC	CCC	AC	GCTAGGA	
<i>Gilia ochroleuca</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	GCG	GGTGTGTATCTC		GGTACT	CGCTCGTTCC	CCCCAC	GCTAGGA		
<i>Gilia flavocincta</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	GCG	GGTGTGTATCTC		GGTACT	CGCTCGTTCC	CCCCAC	GCTAGGA		
<i>Gilia tenerrima</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	GCG	GGCTGTATCTC		GGTACT	TGCHCGTTCC	CCCCAC	GCTAGGA		
<i>Gilia capitata</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	GCG	GGCTGTATCTC		GGTACT	CGCCCGTTCC	CACCTC	GCTAGGA		
<i>Gilia tricolor</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	RCG	GGTGTGTATCTC		GGTACT	CGCTCGTTCC	CCCCAC	GCTAGGA		
<i>Allophyllum gilioides</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	ACG	GGCTGTATCTC		GGTACT	CGCTCGTTCC	CCCCAC	GCTAGGA		
<i>Allophyllum divaricatum</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCAGGTGGT	ACG	GGCTGTATCTC		GGTACT	CGCTCGTTCC	CCCCAC	GCTAGGA		
<i>Gilia splendens</i>	TCGAGACCTGTCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	TCG	GGCGTGTGTA	CTTC	GGTACT	CGCCGTTGC	CCCCAC	GCTAGGA		
<i>Collomia linearis</i>	TCGACAC	TGCATAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	GGTGTGTATCTC	C	GGTACT	CGCTCGTTAC	CCCCAC	GCTAGGA	
<i>Collomia grandiflora</i>	TCGACACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	GGTGTGTATCTC	C	GGTACT	CGCTCGTTAC	CCCCAC	GCTAGGA		
<i>Navarretia breweri</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	GGCTGTATCATC		GGTACT	CGCTCGTTAC	CCCCAC	GCTAGGA		
<i>Polemonium foliosissima</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACG	ARGCTAGTA	CTTC	GGTACT	CGCCCGTYGC	CCCTC	GCTGTGA		
<i>Linanthus aureus</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	AC	TTGTG	CTTC	GGTACT	CACGTTGC	CCCTC	GCTGGTG		
<i>Linanthus parryae</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	ACA	TGTG	CTTC	GGTACT	CGCATGTTGC	CCCTTTGCTGGTG			
<i>Gilia maculata</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	GAC	ATGTG	CTTC	GGTACT	TGTCCGTTGC	CCCTT	GCTGGTG	
<i>Gilia campanulata</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	ATG	GCG	TTGTG	CTTC	GGTACT	CGCCGTTGC	CCCTT	GCTGGTG	
<i>Gilia inyoensis</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGG	GCF	GCG	GGC	TTGTG	CTTC	GGTACT	CGCCCGTTGC	CCCTC	GCTGGTG
<i>Gilia filiformis</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	GCG	GGC	TTGTG	CTTC	GGTACT	CGTCCGTTGC	CCCTC	GCTGGTG	
<i>Phlox gracilis</i>	TCGACACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	TGC	TTGCG	CTTC	GGCCGT	TGCTCG	TGCATTG	CCCTC	GCTGGTG
<i>Phlox standsburyi</i>	TCGAAACCTGCCTAGCATAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GCG	TGC	TTCT	CTTC	GGCCGT	TGCTCG	TGT	AACCWC	GCTGGTG
<i>Leptodactylon watsonii</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGY	GCA	GGC	TTGTG	CTTC	GGTACT	CGCTGTTGC	CCCTTTGCTGGTG		
<i>Leptodactylon pungens</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGC	GCA	TGC	TTGTG	CTTC	GGTACT	CGCTGTTGC	CCCTTTGCTGGTG		
<i>Linanthus bicolor</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GAT	GAC	TTGTG	CTTT	GGTACT	TGTATGTTGC	CCCTT	GCTGGTG	
<i>Linanthus nuttallii</i>	TCGAAACCTGCCTAGCAGAACGACCCGCGAACTTGTAAACTAAA	TCGGGTGGT	GAC	GAC	TTGTG	CTTC	GGTACT	TGCACGTTGC	CCCTT	GCTGGTG	

Appendix 1. Aligned nrDNA ITS1 and ITS2 nucleotide sequences from 53 representatives of Polemoniaceae and three outgroup members. Nucleotide sites are numbered 5' to 3', beginning with the first base of ITS1. Position 278 is the final base of ITS1 and position 279 is the first base of ITS2 (the 5.8S region has been removed). Nucleotide coding follows standard IUB-IUPAC codes. Gaps at nucleotide positions are denoted by dots.

	120	130	140	150	160	170	180	190	200	210	220
<i>Fouquieria columnaris</i>											
<i>Fouquieria splendens</i>											
<i>Diapensia lapponica</i>											
<i>Acanthogilia gloriosa</i>											
<i>Cantua quercifolia</i>											
<i>Bonplandia geminiflora</i>											
<i>Cobaea baurita</i>											
<i>Cobaea scandens</i>											
<i>Gilia caespitosa</i>											
<i>Gilia subnuda</i>											
<i>Gilia pentstemonoides</i>											
<i>Gilia mcvickerae</i>											
<i>Gilia nyensis</i>											
<i>Gilia hutchinsifolia</i>											
<i>Gilia latifolia</i>											
<i>Gilia foetida</i>											
<i>Gilia rigidula</i>											
<i>Gilia scabra</i>											
<i>Loeselia glandulosa</i>											
<i>Ipomopsis longiflora</i>											
<i>Ipomopsis gunnisonii</i>											
<i>Ipomopsis multiflora</i>											
<i>Ipomopsis tenuifolia</i>											
<i>Ipomopsis sonorae</i>											
<i>Eriastrum diffusum</i>											
<i>Loeseliastrum matthewsii</i>											
<i>Loeseliastrum schottii</i>											
<i>Langloisia setosissima</i>											
<i>Langloisia set. punctata</i>											
<i>Gilia stellata</i>											
<i>Gilia crassifolia</i>											
<i>Gilia scopulorum</i>											
<i>Gilia ochroleuca</i>											
<i>Gilia flavocincta</i>											
<i>Gilia tenerima</i>											
<i>Gilia capitata</i>											
<i>Gilia tricolor</i>											
<i>Allophyllum gilioides</i>											
<i>Allophyllum divaricatum</i>											
<i>Gilia splendens</i>											
<i>Collomia linearis</i>											
<i>Collomia grandiflora</i>											
<i>Navarretia breweri</i>											
<i>Polemonium foliosissima</i>											
<i>Linanthus aureus</i>											
<i>Linanthus parryae</i>											
<i>Gilia maculata</i>											
<i>Gilia campanulata</i>											
<i>Gilia inyoensis</i>											
<i>Gilia filiformis</i>											
<i>Phlox gracilis</i>											
<i>Phlox standsburyi</i>											
<i>Leptodactylon watsonii</i>											
<i>Leptodactylon pungens</i>											
<i>Linanthus bicolor</i>											
<i>Linanthus nuttallii</i>											

	230	240	250	260	270	280	290	300	310	320	330
				. ITS1 .	.	<----> [.---->	. ITS2 .				
<i>Fouquieria columnaris</i>	CCCCGCGCTCC.GTTCGCGGATGCACG..GGGACGGTTGCATCTTTTC.ACAGAACATCGCGTGCCTCCCAACCC.CTCCCGCCGTTCG.GGGCGCGGGTGTTCGAGG										
<i>Fouquieria splendens</i>	CCCCGACGCTCC.GTTTGGCGGATGCACG..GGGACGGTTGCATCTTTTC.ACAGAACATCGCGTGCCTCCCAACCC.CTCCCGCCCTTGT.GGGCGCGGGTGTTCGAGG										
<i>Diapensia lapponica</i>	TTCCCTCTTGCCCGTTTACTCGGGTGTACGGGGGAAGTTGG.TGTCTCTGAAAACATCGCGTGCCTCCCAACCCCTCTCCCTTTTGTGGTGTAGTGGAGTGTGGGG										
<i>Acanthogilia gloriosa</i>	TC.CGGAGGCCCC.GTTCGCGGGTGA.CGAAGGGCGG.TGCCATCTGA.C.ACAAAACATCGCGTGCCTCCCAACCC..TCTGCCCTCAG.GGGGACGGGCAATG.GGA										
<i>Cantua quercifolia</i>	TCCGTCCCTCGCCCGTTACGGGGTGT.CGAAGGGGCG.CGCGATCD.A.C.GCAATGATTGCGTGCCTCCCAACCC..TGTCCCTCGG.GGAATCRG..ATTG.GS.										
<i>Bonplandia geminiflora</i>	CC.CGGAGGCCCG.CTTTCG..GGT.C.CGAAGGGCAA.TGCCATCTTA.C.TCAATACATCGCGTGCCTCCCAACCC..TACCCTTCG.GGCGTGACGGGGTTC.GGA										
<i>Cobaea baurita</i>	TC.CTCTTTCCCATTTTATGG..GT.CAAAGGGCAA.TGTCRTCTTA.C.ACAATACATCTCGTGCCTCCCAACCC..TATTTCCACAA.GAAAKYRTGTGTGT.CGA										
<i>Cobaea scandens</i>	TC.CTCTTTCCCATTTTATGGGTGT.CAGAGGGCAA.TGTCGTCCTA.CAACAATACATCTCGTGCCTCCCAACCC..TATTTCTACAA.GAAATGTGTGTGT.CGA										
<i>Gilia caespitosa</i>	TT.TGGAAGCCCC.GTTCGCGGGTGC.CGAAGTCAA.TGCCATCTKA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GGATTTGAGGTATTG.GGA										
<i>Gilia subnuda</i>	TT.TGGAAGCCCC.GTTCGCGGGTGC.CGAAGG.CAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTTCCCTTTG.GGATTTGATGTATTG.GGA										
<i>Gilia pentstemonoides</i>	TT.TGGAAGTCCC.GTTCGCGGGTGC.CGAAGGACAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GGATTTAAGGTATTG.GGA										
<i>Gilia mcvickeriae</i>	TT.TGGAAGCCCC.GTTCGCGGGTGC.CGAA.GACAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GGATTTAAGGTATTG.GGA										
<i>Gilia nyensis</i>	TT.CGAAGGCCCC.GTTCGCGGGTGC.CGAAGAACAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GGATTTAAGGTATTG.GGA										
<i>Gilia hutchinsifolia</i>	TT.CGGAGGCCCC.GTTCGCGG.GTGC.TGAAGGGCAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GGATTTMAGGTCTTG.GGA										
<i>Gilia latifolia</i>	TC.TCAAAGCCCC.GTTCGC.GCTGC.CGAAGG.CAA.TG.CATCTTA.C.ACGAACATCGCGTGCCTCCCAACCC..TATTTCCCTCTG.GGATTTACGGTTTTT.GGA										
<i>Gilia foetida</i>	TC.CGGATGCCTT.GTTCGCGGGTGC.GGAAGGGCAA.TGCCATCTTA.C.TCAAACATCGCGTGCCTCCCAACCC..CTTCCCTATG.GGAATATGGGTATTG.GGA										
<i>Gilia rigidula</i>	...AGAAGCCTT.GTTCGCGGAGTGC..AAGGGTAT.TGCCATCTTA.C.WMAAACATCGCGTGCCTCCCAACCC..TCTTCCCATTG.GGA.GATGGTTTTG.GGA										
<i>Gilia scabra</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.GGAAGGGCAA.CGCGATCTTG.C.ACAAACATCGCGTGCCTCCCAACCC..CTTCCCTCTG.GGATTTGCGGGCTTG.GGA										
<i>Loeselia glandulosa</i>	CC.TGGAAGCCCC.GTTCGCGGGTGT.GGAAGGGCAGATGCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTCG.GGAATATGGGTATTG.GGA										
<i>Ipomopsis longiflora</i>	TTGCGGAA.CCCC.GTTCAGGGTGC.GGAAGTCAA.TGCAATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GAATTTGTTGTTTT.GGA										
<i>Ipomopsis gunnisonii</i>	CT.CGAAAGCCCC.GTTCGCGRRWCAAGAAGGGCAA.TGCAATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTAG.GGAATACGGTGTG.GGA										
<i>Ipomopsis multiflora</i>	TT.CRGAAGCCCC.GTTCGCGGGTGC.GGAAGTCAA.TGCAATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TATTTCCCTTTG.GGATTTAAGGTATTG.GGA										
<i>Ipomopsis tenuifolia</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.GGAAG.GCAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTTCCCTTCG.GGAATGTGGTCTTG.GGA										
<i>Ipomopsis sonorae</i>	TT.CGGAAGCCCC.GTTCGCGGGTGC.AGAAG.GCAATTGCATCTTA.C.AAAAA.CTAAACGTCGCGTGCCTCCCAACCC..TCTTCCCTCCG.GGAAGCACGGTCTTG.GGA										
<i>Eriastrum diffusum</i>	TC.CGGAAG.CCC.GTTCGC.GGTTCGCGGAAGGGCAA.TG.CATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTT.CCTCT..GGAATACGGTCTTG.GGA										
<i>Loeseliastrum matthewsii</i>	TT.CGGAAGCCCC.GTTCGCGGGTGC.CGAAGKCAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTT.CCTCT..GGAATACGGTCTT.GGA										
<i>Loeseliastrum schottii</i>	TT.CGGAAG.CCC.GTTCGCGGGTGC.CGAAGG.CAA.TG.CATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTTCCCTCTG.GGATTTACGGTCTTG.GGA										
<i>Langloisia setosissima</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.GGAAGGGCAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTTCCCTCTG.GGAATACGGTCTTG.GGA										
<i>Langloisia set. punctata</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.GGAAGGGCAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCTT.CCTCT..GGAATACGGTCTTG.GGA										
<i>Gilia stellata</i>	CT.TGGAAGCCCC.GTCCGCGGGTGC.TGAAGGGCAG.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTACG.GGAATCATGG.TTCG.GGG										
<i>Gilia crassifolia</i>	TT.TCGATGCCCC.GTCCGCGGGTGC.TGAAGGGCAA.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTAAG.GGAATCATGG.TTCG..GG										
<i>Gilia scopulorum</i>	CT.TGGAAGCCCC.GTCCGCGGGTGC.TGAAGGGCAA.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTATG.GGAATCATGG.GTCG.GGG										
<i>Gilia ochroleuca</i>	CT.TAGATGCCCC.GTTT.CGCGTGC.TGAAGGGCAA.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTACG.GGAATCATGG.TTCG.GGG										
<i>Gilia flavocincta</i>	CT.TGGATGCCCC.GTTCGCGGGTGC.TGAAGGGCAA.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTAAG.GGAATCATGGTTTTCG.GGA										
<i>Gilia tenerrima</i>	CC.TGGATGCCCC.GTTCGCGGGTGC.WGAAGGGTAGTGACATC.TA.CAAAAAACATCGCGTGCCTCCCAACCC..TCTTCCCTATG.GGAATCATGG.TTCG.GGA										
<i>Gilia capitata</i>	CT.TGGATGCCCC.GTTCGCGGGTGC.TGAAGGGCAA.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTACG.GGAATCATGG.TTCG.GGG										
<i>Gilia tricolor</i>	CT.TGGATGCCCC.GTTCGCGGGTGC.TGAAGGGCAA.GCCATCTTA.C.AAAAAATATCGCGTGCCTCCCAACCC..TCTTCCCTATG.GGAATCATGG.TTCG.GGG										
<i>Allophyllum giloides</i>	TT.CGGAAGCCCC.GTTCGCGGGTGC.TGAAGGGCAA.TGCCATCTTA.C.AAATACATCGCGTGCCTCCCAACCC..TCTTCCCTATG.GGAATCATGK.TCTG.GGA										
<i>Allophyllum divaricatum</i>	TT.CGGAAGCCCC.GTTCGCGGGTGC.TGAAGGGCAA.TGCCATCTTA.C.AAATATATACGCGTGCCTCCCAACCC..TCTTCCCTATG.GGAATCATGG.TTCG.GGA										
<i>Gilia splendens</i>	CT.TGGATGCCCC.GTCCGCGGGTGC.TGAAGGGCAA.TGCCATCTTA.C.AAAAA.CATCGCGTGCCTCCCAACCC..TCTTCCCTATG.GGAATCATGG.TGTG.GGA										
<i>Collomia linearis</i>	CT.TGGATGCCCC.GTCCGCGGGTGC.TGAAGTCAA.TGCCATCTTG.C.AAAAAACATCGCGTGCCTCCCAACCC..TCACTCCCTATG.GGAATCTTGG.TTCG.GGA										
<i>Collomia grandiflora</i>	CT.TGGATGCCCC.GTTTCGCGGGTGC.TGAAGAACAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCACTCCCTATG.GGAAGCATGG.TTCG.GGA										
<i>Navarretia breweri</i>	CT.TGGAAGCCCCGTTTCGCGGGTGC.TGAAGTCAA.TGCCATCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCACTCCCTATG.GGAATCATGG.TTCG.GGA										
<i>Polemonium foliosissimum</i>	C..CAGAAGCCC..GTTTCGCGGGTGC..GAAAGTCAA..GCG.TCTTA.C.AAAAAACATCGCGTGCCTCCCAACCC..TCATTTCCCTTTG.GGAATCAGGGTCTTG.GGA										
<i>Linanthus aureus</i>	TT.TGGAAGCCCC.GTTCGCGGGTGC.CAAAGGGCAG.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..ATTCCCTTTG.GAAACACGGG.TTCG.GGA										
<i>Linanthus parryae</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.GGAAGGGCAT.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..TTTCCCTCTG.GGAATCATGGGCTTG.GGA										
<i>Gilia maculata</i>	TC.CAGAAGCCCC.GTTTTCGCGGGTGC.CGAAGGGCAT.TG.CATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..TTTCCCTTTG.GGAATCAGGGGATTG.GGA										
<i>Gilia campanulata</i>	TC.TGGAAGCCCC.GTTCGCGGGTGC.CGAAGGGCAT.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..TTTCCCTTTG.GGAATCAGGGGTTTG.GGA										
<i>Gilia inyoensis</i>	TC.TCGAAGCCCC.GTTCGCGGGTGC.TGAAGGGCAT.TGCCATCTTA.C.AAAAACTTCGCGTGCCTCCCAACCC..CTTTCCTTTG.GGAATCAGGGGCTTG.GGA										
<i>Gilia filiformis</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.CGAAGGGCAT.TGCCATCTTA.C.ACAAACTTCGCGTGCCTCCCAACCC..TTTCCCTTTG.GGAATCAGGGGCTTG.GGA										
<i>Phlox gracilis</i>	TC.TGGAAGCCCC.GTTCGCGGGTGC.CGAAGGGCAG.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..ATTCCCTTCG.GGAATCAGGGGTTTG.GGA										
<i>Phlox standsburyi</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.CGAAGGGCAT.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..ATTTCCTTCG.GGAATCAACGGTATTG.GGA										
<i>Leptodactylon watsonii</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.GGAAGGGCAT.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..TTTTCCTTTT.GGAATCAGGGGCTTG.GGA										
<i>Leptodactylon pungens</i>	TC.CGGAAGCCCC.GTTCGCGGGTGC.CAAAGGGCAT.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAACCC..TTTCCCTTTT.GGAATCAGGGGCTTG.GGA										
<i>Linanthus bicolor</i>	TC.TGGAAGCCCC.GTTCGCGGGTGC.AAAAGGGCAA.TGCCATCTTA.C.ACAAACTACGCGTGCCTCCCAACCA..ATT.CCTTTT.GGAATCAGGGTCTTG.GGA										
<i>Linanthus nuttallii</i>	CC.TGGAAGTCCC.GTTCGCGGGTGC.CAAAGGGCAT.TGCCATCTTA.C.ACAAAACATCGCGTGCCTCCCAATCC..ATT.CCTTTT.GGAATCATGGTTTTG.GGA										

Appendix 1. Continued.

	340	350	360	370	380	390	400	410	420	430	440	
<i>Fouquieria columnaris</i>	GCGGAA	.ATGG	.CCCCCGTGC	CGTCA	GTGCGG	TGCGCCT	AAAAAGAG	AGTCCC	.CGGCGACT	GACG	.TCACGA	.CAAGTGGTGGT
<i>Fouquieria splendens</i>	GCGGAA	.ATGG	.CCCCCGTGC	CACATC	AGTGC	CGCGGT	TGCGCCT	AAAAAGCG	AGTCCC	.CGACGACT	GACG	.TCACGA
<i>Diapensia lapponica</i>	GCGGAA	.TTGG	.CCCCCGTGC	AGACAG	TCTG	CGCGG	TGCGC	AAAAATA	.CAGTCCA	TCGACG	AAAWTAT	GTGCACGA
<i>Acanthogilia gloriosa</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CAC	.GCTG	CGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAG
<i>Cantua quercifolia</i>	GCRGAGA	.TTGG	.CCTCCCGTGC	CACATC	GTG	CGCGGT	TGCGC	AAAAATA	.AGTGA	.CGTGGATG	GGGCG	.TCACGA
<i>Bonplandia geminiflora</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CGCTAG	CCG	TGCGGT	TGCGC	AAAAATA	.CGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Cobaea biaurita</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CACAT	TTG	TGTGGT	TGCGC	AAAAATA	.TGAGTSCA	.CGTGGATG	AGAGT	.TCGCAA
<i>Cobaea scandens</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CACAT	TTG	TGTGGT	TGCGC	AAAAATA	.TGAGTSCA	.CGTGGATG	AGAGT	.TCGCAA
<i>Gilia caespitosa</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CTTT	.TCTG	TGCGGT	TGCGC	AAAAATA	.CGAGTCGA	.CGTGGCGAG	CG	.TCACAA
<i>Gilia subnuda</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ATTT	.TCTG	TGCGGT	TGCGC	AAAAATA	.CAAGTCGA	.CGTGGCGAG	CG	.TCACAA
<i>Gilia pentstemonoides</i>	GCRGAAA	.TTGG	.TCTCCCGTGC	CTCT	.WTTG	TGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia mcvickeriae</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CTCT	.TCTG	TGCGGT	TGCGC	AAAAATA	.AGAGTMGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia nyensis</i>	GCGGAA	.TTGG	.TCTCCCGTGC	CTCT	.TCTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGAMG	AGCG	.TCACAA
<i>Gilia hutchinsifolia</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ATTT	.TCTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia latifolia</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ATTT	.TCTG	TGCGGT	TGCGC	AAAAATA	.TGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia foetida</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAG	.CGG	.CGTGGCGAG	CG
<i>Gilia rigidula</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTAT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGG	.CGTGGATG	AGCG	.TCACAA
<i>Gilia scabra</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	AGCGGT	TGCGC	AAAAATA	.CGAGTCGA	.CGTGAACG	AGCG	.TCACAA
<i>Loeselia glandulosa</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTC	AGATG	CGCGGT	TGCGC	AAAAATA	.AGAGTCGG	.CGTGAACG	AGCG	.TCACAA
<i>Ipomopsis longiflora</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Ipomopsis gunnisonii</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGATG	AGCG	.TCACAA
<i>Ipomopsis multiflora</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTTTG	.CTKCGG	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA	
<i>Ipomopsis tenuifolia</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Ipomopsis sonorae</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Eriastrum diffusum</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGKAG	AGCG	.TCACAA
<i>Loeseliastrum matthewsii</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGCGAG	CG	.TCACAA
<i>Loeseliastrum schottii</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	CGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGCGAG	CG	.TCACAA
<i>Langloisia setosissima</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGCGAG	CG	.TCACAA
<i>Langloisia set. punctata</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGCGAG	CG	.TCACAA
<i>Gilia stellata</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACACCG	ATG	CGCGGT	TGCGC	AAAAATA	.TGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia crassifolia</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACACCG	ATG	CGCGGT	TGCGC	AAAAATA	.TGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia scopulorum</i>	GCGGAA	.TTGG	.CCTCCCGTGC	ACAC	ATG	CGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia ochroleuca</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACACCG	ATG	CGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia flavocincta</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACATC	ATG	CGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia tenerrima</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACCG	TTG	YCGCGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia capitata</i>	..GAGA	.TTGG	.TCTCCCGTGC	ACTCCG	ATG	CGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia tricolor</i>	ACGGAA	.TTGG	.TCTCCCGTGC	ACTCCG	ATG	CGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Allophyllum gilioides</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTCT	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGAACG	AGCG	.TCATAA
<i>Allophyllum divaricatum</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTACG	TTG	TGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGAACG	AGCG	.TCATAA
<i>Gilia splendens</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACAATC	CTG	CGCGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TTACAA
<i>Collomia linearis</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACATC	GTG	CGTGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGAACG	AGCG	.GTCACAA
<i>Collomia grandiflora</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTACG	TTG	CGTGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGAACG	AGCG	.TCACAA
<i>Navarretia breweri</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACAT	TTG	CGTGGT	TGCGC	AAAAATA	.AGAGTGA	.CGTGGACG	AGCG	.TCACAA
<i>Polemonium foliosissima</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACGCT	TTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGAACG	AGCG	.TCACAA
<i>Linanthus aureus</i>	.CGGAGA	.TTGG	.TCTCCCGTGC	ACAT	.GTTG	TGTGGT	TGCGC	AAAAATA	.AGAGTGG	.CGTGTACG	AGATG	.CSACAA
<i>Linanthus parryae</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTT	.GTTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGTACG	AGATG	.CSACAA
<i>Gilia maculata</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTT	.GTTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia campanulata</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACAACG	CTG	CGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Gilia inyoensis</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTTCG	CTG	CGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGAGG	AGCG	.TCACAA
<i>Gilia filiformis</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTT	.GTTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGGACG	AGCG	.TCACAA
<i>Phlox gracilis</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTTCG	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGT	.GA	.CGCGGACT	AGCG
<i>Phlox standsburyi</i>	GCGGAA	.TTGG	.TCTCCCGTGC	ACTTCG	GTG	TGCGGT	TGCGC	AAAAATA	.AGAGT	.GA	.CGCGGACT	AGCG
<i>Leptodactylon watsonii</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTT	.GTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGTACG	AGCG	.TCACAA
<i>Leptodactylon pungens</i>	GCGGAGA	.TTGG	.TCTCCCGTGC	ACTT	.GTTG	TGCGGT	TGCGC	AAAAATA	.AGAGTCGA	.CGTGTACG	AGCG	.TCACAA
<i>Linanthus bicolor</i>	GCGGAGA	.TTGG	.TCTCCCATAC	ACAT	.GTTG	TGTGGT	TGCGC	AAAAATA	.AGAGTCGG	.CATGAGT	AGATG	.TCACAA
<i>Linanthus nuttallii</i>	GCGGAGA	.TTGG	.TCTCCCGTAC	ACAT	.GCTG	TGTGGT	TGCGC	AAAAATA	.AGAGTCGG	.CGTGAATG	AGATG	.CCACAG

	450	460	470	480	490	500	510	520
<i>Fouquieria columnaris</i>	TTTCCTGTCG	TGCGTCGCA	TCGTCGCAACGGGCTGG	...CTCAC	...GTGACCC	TGAAG	TG	.CGGTAACAG
<i>Fouquieria splendens</i>	TTTCCTGTCG	TGCGTCGCA	TCGTCGCAACGGGCTGG	...CTCAT	...GTGACCC	TGAAG	CG	.CGGTAACAGCTGTCGCTCGACC
<i>Diapensia lapponica</i>	CCT	.TGTCG	TGA	TC	...TTCTCGCCGATGT?GCA	.CTTTA	...	TGACCC
<i>Acanthogilia gloriosa</i>	CTGT	TGTTG	TGCGT	GCT	TCGTCCTCC	TCGT	AA	...CTCCA
<i>Cantua quercifolia</i>	TTGCC	GTTG	TGCGT	.CG	TCGTCCTCCGG	...	AA	.GCKCCA
<i>Bonplandia geminiflora</i>	TT	CCTGTTGTTG	GT	.CG	TTCTCCCGG	.TT	AG	.CGTCCG
<i>Cobaea biaurita</i>	TTAT	TGTTG	TCTGA	.CC	TCGTCCTC	...	TACAA	...CWCCA
<i>Cobaea scandens</i>	TTAT	TGTTG	TCTGA	.CC	TCGTCCTC	...	AAAA	...CTCCA
<i>Gilia caespitosa</i>	TTGC	TGTTG	TGCGT	.TT	TCATCCTCT	TCGTCGA	...	CCCCA
<i>Gilia subnuda</i>	TTGC	TGATG	TGCGT	.TT	TCATCCTCT	TCGTCGA	...	CCCCA
<i>Gilia pentstemonoides</i>	TTGC	TGTTG	TGCGT	.TT	TCATCCTCA	TCGTCGA	...	CTACA
<i>Gilia mcvickerae</i>	TTGC	TGTTG	TGCGT	.TT	TCATCCTCT	TCGTCGA	...	CTCCA
<i>Gilia nyensis</i>	TTGC	TGTTG	TACGT	.TT	TCATCCTCT	TCGTCGA	...	CTCCA
<i>Gilia hutchinsifolia</i>	TTGC	TGTTG	TACGT	.TT	TCATCCTTT	TCGTCGA	...	CTTCA
<i>Gilia latifolia</i>	TTGC	TGTTG	TGCGT	.TA	CCGTCTCT	TCGCCAA	...	CTCCA
<i>Gilia foetida</i>	TTGT	TGTTG	TGCGT	.TA	AGTCGTGCTY	CGTCCAGCCTTACATAGTTGACCC	...	TGTTG
<i>Gilia rigidula</i>	TTGT	TGTTG	TGCGT	.TA	A	TCCT	CT	TCGTCAA
<i>Gilia scabra</i>	TGGT	TGTTG	TGCGT	.CA	TCGTACTCA	CCGTTGG	...	TTCCG
<i>Loeselia glandulosa</i>	TTGT	TGTTG	TAGCGT	.CAGCCGCTCTGT	ACGCSAA	...	CTCAA	...
<i>Ipomopsis longiflora</i>	TTGC	TGTTG	TGCGT	.CA	CCGTCTC	ACGTCAA	...	CTCCA
<i>Ipomopsis gunnisonii</i>	ATTC	TGTTG	WGCCT	.AA	KCATCTCT	ATGTMGA	...	CTCCA
<i>Ipomopsis multiflora</i>	TTGC	TGTTG	TGCGT	.CA	TGCTCTCT	CCGTCAA	...	CTCCA
<i>Ipomopsis tenuifolia</i>	TTGC	TGTTG	TGCGT	.CA	CCGTCTCT	CCGTCAA	...	CTCAA
<i>Ipomopsis sonora</i>	TTGT	TGTTG	TGCGT	.CG	TCGTCGCT	CAGTCGA	...	CTCCA
<i>Eriastrum diffusum</i>	TTGCGT	TGTTG	TGCGT	.GCGTCTCTCT	CCGTCAA	...	CTCCA	...
<i>Loeseliastrum matthewsii</i>	TTGC	TGTTG	AGCGT	.CA	CCGTCTCT	CCGTCAA	...	CTCCA
<i>Loeseliastrum schottii</i>	TTGM	TGTTG	TGCGT	.CA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Langloisia setosissima</i>	TTGC	TGTTG	TGCGT	.CA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Langloisia set. punctata</i>	TTGC	TGTTG	TGCGT	.CA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Gilia stellata</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Gilia crassifolia</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Gilia scopulorum</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Gilia ochroleuca</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTMT	CCGTCAA	...	CTCCA
<i>Gilia flavocincta</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTCT	CCGTCAA	...	CTCCA
<i>Gilia tenerrima</i>	TTGT	TGTTG	TGCGT	.TT	TCGTCTCT	WCCTCAA	...	CTCCA
<i>Gilia capitata</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTCT	CGTACAG	...	CTCCA
<i>Gilia tricolor</i>	TTGT	TGTTG	TGCGT	.TA	TCGTCTCT	CGTACAT	...	CTCCA
<i>Allophyllum gilioides</i>	TTAT	TGTTG	TAGCGT	.TAGTTGTTCTCT	TCGTTAA	...	CTCCA	...
<i>Allophyllum divaricatum</i>	TTAT	TGTTG	GCGT	.TA	TTGTTCTCT	TCGTTAA	...	CTCCA
<i>Gilia splendens</i>	TTGC	TGTTG	TGCGT	.CT	TCGCCCTCT	TCGTTAA	...	CTCCA
<i>Collomia linearis</i>	TTGT	TGTTG	TGCGT	.TA	TTGTTCTCT	CCGTCAA	...	CTCCA
<i>Collomia grandiflora</i>	TTGT	TGTTG	TGCGT	.TG	TTGTTCTCT	CCGTCAA	...	CTCCA
<i>Navarretia breweri</i>	TTGT	TGTTG	WGCCT	.TA	TTGTTCTCT	TCGTTAA	...	CTCCA
<i>Polemonium foliosissimum</i>	TTGC	TGTTG	WGCCT	.TA	SCGTCTCT	TCGTTMG	...	CTCCA
<i>Linanthus aureus</i>	TTACAT	GTTG	TG	...	TGCTTTTCCG	...	MTGT	...
<i>Linanthus parryae</i>	TTGC	TGTTG	TGCGT	.TG	TCGTCTCG	TC	...	CGA
<i>Gilia maculata</i>	TTAC	TGTTG	TGCGT	.TA	ACGTCTCT	TCGTCGA	...	CTCCA
<i>Gilia campanulata</i>	TTAC	TGTTG	TGCGT	.TG	TCGTCTCT	TCGTCGA	...	CTCCA
<i>Gilia inyoensis</i>	TTAC	TGTTG	TGCGT	.CG	TTGCTCTCT	TCGTCGA	...	CTCCA
<i>Gilia filiformis</i>	TTAC	TGTTG	TGCGT	.TG	TCGTCTCT	TCGTCGA	...	CTCCA
<i>Phlox gracilis</i>	TTAC	TGTTG	TTCGT	.CTTTGCTCTCG	...	TCTA	...	CTCCA
<i>Phlox standsburyi</i>	TTAC	TGTTG	TGCGT	.CTTTGCTCTCG	...	TCTA	...	CTCCA
<i>Leptodactylon watsonii</i>	TTGC	TGTTG	TGCGT	.TG	TCGTCTCT	TCGTCGA	...	CTCCA
<i>Leptodactylon pungens</i>	TTGC	TGTTG	TGCGT	.TG	TCGTCTCT	TCGTCGA	...	CTCCA
<i>Linanthus bicolor</i>	T	.C	TGTTG	TGCG	...	GCTGT	...	TTCCA
<i>Linanthus nuttallii</i>	TTACAT	GTTG	TG	...	CTGTCAT	...	GTTAT	...

Appendix 1. Continued.