

Adapting agriculture to climate change:
collecting, protecting and preparing crop wild relatives

Uganda



Seed Collecting Guide

Please cite this guide as:
RBG Kew (2015) Uganda Seed Collecting Guide

NOTE: This guide has been abridged owing to copyright restrictions for selected images and/or datasets. Full, unabridged guides are only available from the Kew Millennium Seed Bank Library Archive. To request access to a full, unabridged guide, please contact the Seed Bank Librarian at msbplib@kew.org.

The content of this collecting guide is intended only as a general reference for future collecting missions; the contents and data within are not guaranteed to be complete, correct, timely, current or up-to-date at the time of publishing. For general information and resources on collecting crop wild relatives, visit cwrdiversity.org.

Cover photos

TOP LEFT: Sorghum, CREDIT: Neil Palmer/CIAT;

TOP RIGHT: Black beans, CREDIT: Neil Palmer/CIAT;

BOTTOM LEFT: Bean breeding in Uganda, CREDIT: Neil Palmer/CIAT;

BOTTOM RIGHT: Rice CREDIT: Neil Palmer/CIAT

This work was undertaken as part of the initiative “Adapting Agriculture to Climate Change” which is supported by the Government of Norway. The project is managed by the Global Crop Diversity Trust with the Millennium Seed Bank of the Royal Botanic Gardens, Kew, in partnership with national and international genebanks and plant breeding institutes around the world. It is implemented in accordance with the International Treaty on Plant Genetic Resources for Food and Agriculture. For further information see the project website: www.cwrdiversity.org/

Many individual scientists, herbaria, genebanks and specialist institutes are contributing advice and information to the Project and these guides. The Project aims to collect the wild relatives of 29 key crops, conserve them in genebanks, and prepare them for use in plant improvement programs to breed new crop varieties adapted to future climates.



MILLENNIUM
SEED BANK
PARTNERSHIP



The boundaries and names shown on the maps included in this guide do not imply official endorsement or acceptance by the Adapting Agriculture to Climate Change Project. Data source: GADM, Version 1.0 via diva-gis.org

This work is licenced by RBG Kew (2016) under the Creative Commons CC BY-NC-ND 4.0 Licence. To view a copy of the licence visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>. If use in any further context is required please contact RBG Kew for permission. Copyright restrictions for the datasets and images used in the compilation remain as stated at their primary sources.

The Harlan and de Wet Crop Wild Relatives Checklist was developed by Holly Vincent and Nigel Maxted at the University of Birmingham.

UNIVERSITY OF
BIRMINGHAM



International Center for Tropical Agriculture
Since 1967 *Science to cultivate change*

The Gap Analysis work which informed the list of species included in this guide, and all the map files, were produced by the Gap Analysis team at CIAT: Andy Jarvis, Nora Castañeda, Colin Khoury and Julian Ramirez-Villegas.

RBG Kew is involved in the research and collection phases of the project. This collecting guide was developed based on the work of the Millennium Seed Bank Enhancement Project Species Targeting Team.

Royal Botanic Gardens
Kew



CROP
TRUST

The Crop Wild Relatives Project is led by the Global Crop Diversity Trust. This work was undertaken as part of the initiative.

Specimen data was kindly provided to this project by many individuals and organisations who are listed on the website: <http://www.cwrdiversity.org/home/data-sources>

This data set will be made available for download. Please refer to the website for more information on this dataset.

This collecting guide has been compiled by:

Ruth Harker

Collecting Guide Compiler
Conservation Science Department
Herbarium, Library Art & Archives
Royal Botanic Gardens, Kew

Dr Ruth Eastwood

Crop Wild Relatives Project Co-ordinator
Millennium Seed Bank Partnership
Conservation Science Department
Royal Botanic Gardens, Kew

This collecting guide consists of species profiles and information sheets contained within this folder, alongside a CD which contains localities of the taxa in an excel file.


The species included in this guide are a selection of the wild relatives of the 29 key crops which this project covers (African Rice, Alfalfa, Apple, Aubergine, Bambara groundnut, Banana, Barley, Bread Wheat, Butter Bean, Carrot, Chickpea, Common Bean, Cowpea, Faba bean, Finger millet, Grasspea, Lentil, Oat, Pea, Pearl millet, Pigeon pea, Plantain, Potato, Rice, Rye, Sorghum, Sunflower, Sweet potato, Vetch). It is not a definitive guide to the Crop Wild Relatives in this country.

The guides are designed to be used both in the planning of a collecting trip, and also in the field.

At the front of this guide there is a phenology table showing the flowering and fruiting times of all the taxa to indicate which species may be found at a certain time of year, or when to collect target species.

Synonyms for each species are listed in the Appendix at the end of this guide.


On each species profile, there is a collection of images to help identify the target species, accompanied by a series of symbols :



Conservation Assessments

Conservation Status:

Assessments are completed using 2001 IUCN Red List Categories and Criteria version 3.1 with the following categories:



Where a full conservation assessment has not been completed, a preliminary conservation rating may be indicated. Preliminary assessments are produced using specimen locality data and GIS, which calculates two parameters accepted by IUCN as suitable measures of range: namely extent of occurrence (EOO) and area of occupancy (AOO). These values derived for each species are then compared with thresholds set out by IUCN under Criterion B.

Where a preliminary conservation assessment has been calculated this is indicated by the word PRELIM:



Maps

Two maps are provided for each target species. The first map shows a point distribution of all the known localities of this species based on herbarium specimen records and existing data-sets. The area shaded on this map shows the predicted distribution based on Maxent.



The second map shows the potential gaps in gene bank collections, where seed collections should be targeted.



Useful resources

The following resources are available online.

Kew technical information sheets

- Assessing a potential seed collection:
<http://brahmsonline.kew.org/Content/Projects/msbp/resources/Training/02-Assessing-population.pdf>
- Post-harvest handling of seed collections:
<http://brahmsonline.kew.org/Content/Projects/msbp/resources/Training/04-Post-harvest-handling.pdf>

Other sheets covering the following topics are available from

<http://brahmsonline.kew.org/msbp/Training/Resources>

- Protocol for comparative seed longevity testing
- Measuring seed moisture status using a hygrometer
- Selecting containers for long-term seed storage
- Low-cost monitors of seed moisture status
- Small-scale seed drying methods
- Equilibrating seeds to specific moisture levels
- Identifying desiccation-sensitive seeds
- Seed bank design: seed drying rooms
- Seed bank design: cold rooms for seed storage
- Cleaning seed collections for long-term conservation

ENSCONET seed collecting manual for wild species

http://ensconet.maich.gr/PDF/Collecting_protocol_English.pdf

Seed conservation: turning science into practice

<https://academic.oup.com/aob/article/95/5/888/201951>

Collecting plant genetic diversity: Technical guidelines (Bioversity)

http://cropgenebank.sgrp.cgiar.org/index.php?option=com_content&view=article&id=390&Itemid=557

FAO – Commission on Genetic Resources for Food and Agriculture

<http://www.fao.org/nr/cgrfa/en/>

IUCN Red List Categories and Criteria (Version 3.1)

<https://iucn-csg.org/red-list-categories/>

Plants of the World Online

<http://plantsoftheworldonline.org/>

For more information about the Crop Wild Relatives Project and to access the Harlan and de Wet Crop Wild Relatives checklist, please visit the website:

www.cwrdiversity.org

Identification Keys

Interactive identification keys can be accessed using the links below.

Kew Grassbase interactive identification key

<http://www.kew.org/data/grasses-db/ident.htm>

African Vigna: an interactive key

http://keys.lucidcentral.org/keys/African_Vigna/default.htm

Clayton, W.D., Vorontsova, M.S., Harman, K.T. and Williamson, H. (2006 onwards). GrassBase - The Online World Grass Flora. <http://www.kew.org/data/grasses-db.html>. [accessed 15 March 2012; 14:30 GMT]

Maxted, N., Mabuza-Dlamini, P., Moss, H., Padulosi, S., Jarvis, A. & Guarino, L., (2004). An Ecogeographic Survey: African Vigna. *Systematic and Ecogeographic Studies of Crop Genepools* 10. pp. 1-468. IPGRI, Rome.

Al-Atawneh, N., Shehadeh, A., Amri, A. & Maxted, N., (2009). Conservation Field Guide to Medics of the Mediterranean Basin. Pp. 1-214. ICARDA, Syria.

Seed Collecting Techniques

Michael Way and Kate Gold, Seed Conservation Department

Seed collecting from wild plants requires care, resourcefulness and determination. There are many different collecting techniques. The most appropriate technique will depend on the species, particularly the type of dispersal unit (fleshy fruit, dry fruit, individual seeds etc). This information sheet outlines the manual techniques most commonly used to make seed collections of adequate quality and quantity, for long term conservation.

Hand picking of whole fruits

The most basic and flexible of techniques, hand picking or plucking, has many benefits. Consider though, if you can use a more efficient technique.




Plucking is particularly suitable when:

- target fruits can easily be selected by eye (e.g. due to colour or texture change of fruit coat, or swelling of fruit);
- non-target (e.g. immature or damaged) fruit cannot be excluded from the collection by more efficient techniques;
- fruits are easily accessible and collectors can tie buckets or similar containers around the waist, releasing both hands for collecting;
- collecting many-seeded fleshy or dry indehiscent fruits; and
- making small seed collections.

Pruning clusters of fruit

This technique is typically used to collect tree seeds. Cut groups or clusters of fruits using secateurs or tree pruners. Assess for ripeness and damage before adding seeds to the collection.




This is a very effective technique when:

- seed is clustered at the distal (terminal) parts of branches;
- the species is abundant and a small associated loss of branch and foliage is acceptable;
- seed is beyond reach of the collectors and has to be obtained using tree pruners.

Shaking branches


Careful shaking of branches will sometimes dislodge the best available seed, which can be collected in buckets or on a tarpaulin held or spread out beneath the plant. Start with gentle taps, and carefully check each sample of seed dislodged. Light shaking will often dislodge fully ripe fruits and seeds, leaving immature, poorly developed and damaged seeds to be retained on the parent plant. Too-heavy beating of branches may cause damage to the tree, and may also dislodge other plant material and associated insects, necessitating additional cleaning of the collection.



Shaking branches may be useful when collecting:

- dehiscent fruits with medium large seeds;
- seeds with irritant plumes (e.g. *Cercocarpus* of the Rosaceae);
- spiny trees such as *Prosopis* (Fabaceae);
- on level, open terrain suitable for tarpaulin use.


This technique may not be suitable for light, plumed seed from *Bombacaceae* and *Asclepiadaceae*, which may be carried away by air currents.



ABOVE: Stripping seed heads may be appropriate for grasses
Credit: Global Crop Diversity Trust/Britta Skagerfalt

Stripping entire seed-heads

This is a popular technique for collecting seed from grasses and may be suitable for other species with erect infructescences (seedheads). Grasp the seed-heads at the base with a gloved hand and slide the hand upwards, dislodging many or all of the seeds. This technique may introduce a proportion of immature seeds into the collection. Such seeds might need further postharvest ripening which can be time consuming and is best avoided.




The stripping technique is most suitable for:

- dense, mono-specific stands of target species with no weed or other species present; and
- infructescences which are completely and consistently at the natural dispersal stage.


Bagging seed-heads

If there is frequent access to the collecting site, and if seeds would otherwise be lost, fix a well-tied mesh bag loosely over pre-dispersal seed heads. Seeds are captured as soon as they are shed, and can be periodically removed. This has been successfully used on a small scale, e.g. for collecting *Fouquieria* sp.



Collecting from the ground

You will frequently find seeds on the ground below trees or shrubs, but they will often be damaged by pests or pathogens. The seeds may have been on the ground for several months, and could even date from the previous year. Such seed will have aged and life-span in storage will be reduced. Inspect the seed carefully, noting any variation in the fruit, seed coat and internal tissues.



In general, only collect from the ground when:

- the parent tree(s) can be determined without doubt;
- you are certain that you are collecting recently dispersed seeds;
- seeds have not suffered significant damage from pests or pathogens; and
- other techniques or collecting options are unsuitable.

Collecting fleshy fruits

- Collect fleshy fruits directly into strong plastic bags or tubs with as much air as possible.
- Pack the bags in a rigid plastic container to ensure that the fruits are not squashed and help prevent them getting too hot and fermenting during transit.
- You may need to remove the seeds from fleshy fruits either during or immediately after the field trip.



ABOVE Collecting small seeds into paper bags
Credit: Ruth Harker/ RBG Kew

Containers

Collect into buckets, cloth or paper bags, and check each person's sample carefully before combining into a single population collection.

Using buckets has the advantage of allowing you to monitor the quality of the collection whilst associated insects disperse freely.


Place collections of dry, ripe seed into cloth or paper bags for transit. Store any awned seed or hooked fruit, that would damage or get stuck in cotton bags, in cardboard boxes or strong paper bags. Never collect or store seeds in plastic bags.

Label all seed containers inside and out with a unique collection number, and seal them securely. It is best to prepare sufficient labels before filling the containers.

Each target species in this guide is a wild relative of a crop. On each species profile it is indicated how closely related the target species is to the crop using either the Gene Pool concept or the Taxon Group concept. Species more closely related to the crop are higher priorities for collecting.

Gene Pool Concept

Harlan and de Wet, 1971



Taxon Group Concept

Maxted et al. 2006

Taxon Group 1 – cultivated/wild form of the crop


Taxon Group 2 – species in same series/section as crop

Taxon Group 3 – species in same subgenus as crop


Harlan, J. and J. de Wet (1971). Towards a rational classification of cultivated plants. *Taxon* 20: 509-517.

Maxted, N., B.V. Ford-Lloyd, S.L. Jury, S.P. Kell and M.A. Scholten (2006). Towards a definition of a crop wild relative. *Biodiversity and Conservation* 14: 1-13.


Occurrences of all taxa in this guide, as a point distribution



Species richness



Collecting Gaps



Species in this guide - High priority for collecting

Family	Taxon	Genepool	Sheet	Conservation Status
LEGUMINOSAE	<i>Vigna unguiculata</i> ssp. <i>burundensis</i> Walp.	Gene Pool 1B relative of <i>Vigna unguiculata</i>	1	VU
POACEAE	<i>Sorghum bicolor</i> ssp. <i>verticilliflorum</i> (L.) Moench	Gene Pool 1B of Sorghum	2	LC (prelim)
POACEAE	<i>Eleusine africana</i> Kenn.-O'Byrne	Genepool 1B of Finger millet	3	LC (prelim)
POACEAE	<i>Pennisetum purpureum</i> Schumach.	Genepool 2 of Pearl millet	4	LC (prelim)
POACEAE	<i>Sorghum purpureosericeum</i> (Hochst. ex A. Rich.) Asch. & Schweinf.	Genepool 3 of Sorghum	5	LC (prelim)
SOLANACEAE	<i>Solanum campylacanthum</i> Hochst.	Genepool 2 of Eggplant	6	LC (prelim)
SOLANACEAE	<i>Solanum dasyphyllum</i> Schumach.	Genepool 3 of Eggplant	7	LC (prelim)
SOLANACEAE	<i>Solanum hastifolium</i> Hochst. ex Dunal	Genepool 3 of Eggplant	8	LC (prelim)
SOLANACEAE	<i>Solanum anguivi</i> Lam.	Genepool 2 of Eggplant	9	LC (prelim)
SOLANACEAE	<i>Solanum cyaneopurpureum</i> De Wild.	Genepool 3 of Eggplant	10	DD

Species in this guide - Lower priority for collecting

Family	Taxon	Genepool	Sheet	Conservation Status
CONVOLVULACEAE	<i>Ipomoea ochracea</i> (Lindl.) G. Don	Wild relative of sweet potato	11	LC (prelim)
LEGUMINOSAE	<i>Lens ervoides</i> Grande	Genepool 2 of lentil	12	LC (prelim)
LEGUMINOSAE	<i>Vicia sativa</i> ssp. <i>nigra</i> (L.) Ehrh.	Genepool 1B relative of <i>Vicia sativa</i>	13	LC (prelim)
LEGUMINOSAE	<i>Vigna unguiculata</i> ssp. <i>pubescens</i> Walp.	Genepool 1B of Cowpea	14	LC (prelim)
POACEAE	<i>Eleusine kigeziensis</i> S.M.Phillips	Provisional Genepool 1B of finger millet	15	NT (prelim)
POACEAE	<i>Oryza eichingeri</i> Peter	Genepool 2 relative of rice	16	LC (prelim)
POACEAE	<i>Oryza longistaminata</i> A.Chev. & Roehrich	Genepool 1B of Rice	17	LC (prelim)
POACEAE	<i>Oryza punctata</i> Kotschy ex Steud.	Genepool 2 of rice	18	LC (prelim)
SOLANACEAE	<i>Solanum aculeatissimum</i> Jacq.	Genepool 2 of eggplant	19	LC (prelim)

Taxon	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
<i>Oryza longistaminata</i>	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower
<i>Oryza punctata</i>	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit	Species in fruit
<i>Solanum aculeatissimum</i>	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower	Species in flower

KEY

Species in flower



Species in fruit



data gathered from literature and herbarium specimens

Appendix - Synonyms

Taxon		Synonyms
Vigna unguiculata ssp. burundiensis Walp	1	
Sorghum bicolor ssp. verticilliflorum (L.) Moench	2	Sorghum verticilliflorum (Steud.) Stapf; Sorghum brevicarinatum Snowden; Andropogon sorghum (L.) Brot. var. aethiopicus Hack.; Andropogon sorghum (L.) Brot. subsp. vogelianus Piper; Sorghum vogelianum (Piper) Stapf; Sorghum usambarense Snowden; Sorghum macrochaeta Snowden; Sorghum bicolor (L.) Moench subsp. arundinaceum (Desv.) de Wet & J. R. Harlan ex Davidse; Rhaphis arundinacea Desv.; Sorghum virgatum (Hack.) Stapf; Sorghum stapfii (Hook. f.) C. E. C. Fisch.; Holcus sorghum L. var. effusus Hitchc.; Andropogon arundinaceus Willd.; Andropogon sorghum (L.) Brot. var. virgatus Hack.; Andropogon sorghum (L.) Brot. var. effusus Hack.; Andropogon verticilliflorus Steud.; Sorghum pugionifolium Snowden; Holcus sorghum L. var. verticilliflorus (Steud.) Hitchc.; Sorghum arundinaceum (Desv.) Stapf; Sorghum lanceolatum Stapf; Sorghum aethiopicum (Hack.) Rupr. ex Stapf; Andropogon stapfii Hook. f.
Eleusine africana Kenn.-O'Byrne	3	Eleusine coracana subsp. africana (Kenn.-O'Byrne) Hilu & de Wet; Eleusine indica subsp. africana (Kenn.-O'Byrne) S.M.Phillips
Pennisetum purpureum Schumach.	4	Pennisetum benthamii Steud.; Pennisetum purpureum subsp. benthamii (Steud.) Maire & Weiller; Pennisetum purpureum subsp. flexispica (K.Schum.) Maire & Weiller
Sorghum purpureosericeum (Hochst. ex A. Rich.) Asch. & Schweinf.	5	Andropogon pappii Gand.; Andropogon purpureosericeus Hochst. ex A.Rich.; Andropogon purpureosericeus var. calomelas Hack.; Andropogon purpureosericeus var. pallidior Hack.; Sarga purpureosericea (Hochst. ex A.Rich.) Spangler; Sorghum deccanense Stapf ex Raizada; Sorghum dimidiatum Stapf; Sorghum purpureosericeum subsp. deccanense Garber; Sorghum purpureosericeum subsp. dimidiatum (Stapf) Garber
Solanum campylacanthum Hochst.	6	Solanum suaveolens Bojer; Solanum bojeri var. sinuatorepandum Dunal, in DC.; Solanum bojeri Dunal, in DC.; Solanum benguelense Peyr.; Solanum antidotum Dammer; Solanum fischeri Dammer; Solanum urbanianum Dammer; Solanum psilostylum Dammer; Solanum volkensii Dammer; Solanum astrochlaenoides Dammer; Solanum magdalenae Dammer; Solanum neumannii Dammer; Solanum merkeri Dammer; Solanum deckenii Damme; Solanum bussei Dammer; Solanum delpierrei De Wild.; Solanum endlichii Dammer; Solanum himatacanthum Dammer; Solanum iodes Dammer; Solanum macrosepalum Dammer; Solanum secedens Dammer; Solanum repandifrons Bitter; Solanum verbascifrons Bitter; Solanum volkensii var. himatiacanthum (Dammer) Bitter; Solanum ukerewense Bitter; Solanum malacochlamys var. transgrediens Bitter; Solanum malacochlamys Bitter; Solanum maranguense Bitter; Solanum mesomorphum Bitter; Solanum neumannii var. schoense Bitter; Solanum pembae Bitter; Solanum merkeri var. tobleri Bitter; Solanum merkeri var. endastrophorum Bitter; Solanum merkeri var. intermontanum Bitter; Solanum merkeri var. mediidominans Bitter; Solanum merkeri var. ruandense Bitter; Solanum merkeri subsp. militans Bitter; Solanum delagoense var. astrochlaenoides (Dammer) Bitter; Solanum delagoense var. fischeri (Dammer) Bitter; Solanum delagoense var. benguelense (Peyr.) Bitter; Solanum bojeri var. deckenii (Dammer) Bitter; Solanum bojeri var. houyanum Bitter; Solanum bojeri var. integrum Bitter; Solanum sennii Chiov.; Solanum cufodontii Lanza; Solanum melongenifolium Lanza; Solanum goniocalyx Lanza

Appendix - Synonyms

b

Solanum dasyphyllum Schumach	7	Solanum duplosinuatatum Klotzsch
Solanum hastifolium Hochst. ex Dunal	8	Solanum longestamineum Dammer; Solanum hastifolium subsp. velutinellum Bitter; Solanum cynanchoides Chiov.
Solanum anguivi Lam.	9	Solanum indicum L.; Solanum indicum var. lividum (Link) Bitter; Solanum indicum var. maroanum Bitter; Solanum lividum Link; Solanum sodomeum L.Solanum scalare C. H. Wright
Solanum cyaneopurpureum De Wild.	10	Solanum kitivuense subsp. ukerewense Bitter; Solanum tanganikense Bitter
Ipomoea ochracea (Lindl.) G. Don	11	Ipomoea curtisii House; Ipomoea ochracea var. curtissii (House) Stearn
Lens ervoides Grande	12	Cicer ervoides Brign.; Ervum ervoides (Brign.) Hayek; Ervum hohenakeri Fisch. & C.A. Mey.; Ervum lenticula Schreb. ex Sturm; Lens lenticula; Lens nigricans (M. Bieb.) Godr. subsp. ervoides (Brign.)Ladiz.; Vicia ervoides (Brign.) Fiori; Vicia lenticula (Schreb.) Janka
Vicia sativa ssp. nigra (L.) Ehrh.	13	Vicia angustifolia L.; Vicia angustifolia subsp. angustifolia L.; Vicia angustifolia subsp. pusilla Boiss.; Vicia angustifolia var. segetalis (Thuill.) Arcang.; Vicia angustifolia subsp. segetalis (Thuill.) Arcang.; Vicia angustifolia var. uncinata (Desv.) Rouy; Vicia bobartii E.Forst.; Vicia bobartii Koch; Vicia canadensis Zuccagni; Vicia cuneata Guss.; Vicia debilis Perez Lara; Vicia heterophylla C.Presl; Vicia lanciformis Lange; Vicia maculata C.Presl; Vicia pilosa M.Bieb.; Vicia sativa var. angustifolia (L.) Wahlb.; Vicia sativa var. angustifolia L.; Vicia sativa subsp. angustifolia (L.) Gaudin; Vicia sativa subsp. angustifolia (L.) Batt.; Vicia sativa subsp. consobrina (Pomel) Quezel & Santa; Vicia sativa subsp. cordata (Hoppe) Batt.; Vicia sativa subsp. cuneata (Guss.) Maire; Vicia sativa subsp. heterophylla (C.Presl) J.Duvign.; Vicia sativa var. minor (Bertol.) Ohwi; Vicia sativa var. nigra L.; Vicia segatalis Thuill.
Vigna unguiculata ssp. pubescens Walp.	14	Vigna pubescens R. Wilczek; Vigna unguiculata subsp. protracta; Vigna unguiculata var. protracta; Vigna unguiculata var. pubescens (R. Wilczek) Maréchal, Mascherpa & Stainier.
Eleusine kigeziensis S.M.Phillips	15	
Oryza eichingeri Peter	16	Oryza collina (Trimen) S.D.Sharma & Shastry; Oryza glauca Robyns; Oryza latifolia var. collina (Trimen) Hook.f.; Oryza rhizomatis D.A.Vaughan ; Oryza sativa var. collina Trimen ; Oryza ubanghensis A.Chev.
Oryza longistaminata A.Chev. & Roehrich	17	Oryza barthii auct.
Oryza punctata Kotschy ex Steud.	18	Oryza eichingeri var. longiaristata Peter; Oryza sativa var. punctata (Kotschy ex Steud.) Kotschy; Oryza schweinfurthiana Prodoehl
Solanum aculeatissimum Jacq.	19	Solanum horridum Salisb.; Solanum reflexum Schrank; Solanum aculeatissimum var. hispidissimum Dunal, in DC.; Solanum khasianum C.B.Clarke; Solanum cavaleriei H.Lév. & Vaniot; Solanum angustispinosum De Wild.