

Switzerland - Alpine Flowers of the Upper Engadine

Naturetrek Tour Report

5 - 12 July 2015



Androsace alpina



Cirsium spinosissimum



Eritrichium nanum



Primula integrifolia

Report and images by David Tattersfield



Naturetrek

Mingledown Barn

Wolf's Lane

Chawton

Alton

Hampshire

GU34 3HJ

UK

T: +44 (0)1962 733051

E: info@naturetrek.co.uk

W: www.naturetrek.co.uk

Tour participants: David Tattersfield (leader) with eight Naturetrek clients

Day 1

Sunday 5th July

The temperature had reached 39°C, when the group assembled at Zurich airport. We made our way down to the railway station and caught the train to Zurich main station. Once on the intercity express we settled down, in comfort, to enjoy the scenic journey to the Alps. We passed Lake Zurich and the Walensee, meeting the Rhine as it flows into Liechtenstein, and then changed to the regional railway at Chur. Dramatic scenery and many loops, tunnels and bridges followed, as we made our way through the mountains to Sameden. After a final short train ride to Pontresina, we were met by the hotel minibus and we were soon enjoying a delicious meal. After a brief chat about our plans for the week, we retired to bed.

Day 2

Monday 6th July

We caught the 9.06am train at Surovas and travelled south along Val Bernina, crossing the watershed beside the milky waters of Lago Bianco and alighting at Alp Grum. As it was our first day, there were many flowers to be discovered and we made slow progress up the hillside, to the Belvedere, for coffee. There were superb specimens of St Bruno's Lily *Paradisea liliastrum*, Giant Cat's-ear *Hypochaeris uniflora*, *Arnica montana* and a number of orchids included Heath Spotted-orchid *Dactylorhiza maculata*, Vanilla Orchid *Gymnadenia rbellicani* and Small-white Orchid *Pseudorchis albida*.

In the forest, below the station, there were Martagon Lilies, Yellow Melancholy Thistle *Cirsium eristhales* and a spectacular large area of Alpine Blue-sowthistle *Cicerbita alpina*. It was hot and sunny and we were glad of some shade while we ate our lunch. We continued down through the forest, seeing *Clematis alpina*, Large Yellow Foxglove *Digitalis grandiflora* and showy patches of Carthusian Pink *Dianthus carthusianorum* subsp. *vaginatus*. Butterflies included Swallowtail, Clouded Apollo and the bright-orange Purple-edged Copper. We emerged into colourful meadows, around the delightful village of Cavaglia and visited the Gletscher Garten, where we explored an amazing series of potholes, left behind after the last glacial retreat. The circular walk finished at Cavaglia station where we boarded our return train to Surovas

Day 3

Tuesday 7th July

We took the train to Bernina Lagalb, where we crossed the road into Val Minor. For the first few hundred metres the flora was very rich and colourful, influenced by the limestone screes and boulders tumbling down from Piz Alv. Lots of Red Vanilla Orchids *Nigritella rubra* were scattered in the turf, with *Aster alpinus*, Striped Daphne *Daphne striata* and Jacob's-ladder *Polemonium caeruleum* around the boulders. Large-flowered Common Rockrose and the smaller-flowered Alpine Rockrose grew together with Alpine Gypsophila *Gypsophila repens*, making a stunning display on one area of scree and in one gully, there were large numbers of the deep purple Dragon-mouth *Horminum pyrenaicum*. Something about the soil conditions here suits its requirements and it appears to grow nowhere else in the vicinity. Beyond here, we were back on acidic soils and followed an easy path through pastures, grazed by a placid herd of cattle. In the upper part of the valley, there were many Alpine Pasqueflower *Pusatilla alpina* subsp. *apifolia* and deep-purple Broad-leaved Primrose *Primula latifolia*. We had lunch by the lakes, where there were countless Dwarf Snowbell *Soldanella pusilla* and Entire-leaved Primrose *Primula integrifolia* in the soggy, peaty turf.

After a short steep climb, the path crossed an outcrop of metamorphosed limestone with a striking flora, dominated by dwarf shrubs; Net-veined Willow *Salix reticulata*, Retuse-leaved Willow *Salix retusa* and Mountain Avens *Dryas octopetala*, with blue-grey cushions of *Saxifraga caesia*, Purple Saxifrage *Saxifraga oppositifolia* and Yellow Whitlowgrass *Draba aizoides*. Round the corner we reached our highest point at Fuorcla Minor, offering us splendid views of the Poschiavo Valley, to the south, and the glaciers of Piz Cambrena, ahead of us. From here, it was a gentle walk down to the road, just below the Bernina Pass and then down to Ospizio Bernina, where we caught the train and were back in Surovas for 6pm.

Day 4

Wednesday 8th July

We had light rain in the morning until 10.30am. After taking the train to Morteratsch, we set off along the broad track to the glacier. At intervals, the various positions of the retreating glacier-front since the late nineteenth century are marked, allowing us to fully appreciate the timescale and the processes of colonisation. A colourful flora on the moraines included the brick-red flowers of *Sempervivum arachnoideum*, the deep blue bells of Fairy's-thimble *Campanula cochleariifolia* and the attractive pink willowherb *Epilobium fleischeri*. We approached quite close to the dirty slope of wasting ice, and the torrents of meltwater flowing from beneath it. We returned, in sunshine, to the station, just before another shower and had lunch under the shelter of the platform buildings.

Four of the group decided to walk back to the hotel, while the rest of us took the next train to Ospizio Bernina and set off along the shore of the lake. It remained sunny apart from one heavy shower. There was plenty to interest us on the way, including fine colonies of the delicate yellow Rhaetic Poppy *Papaver aurantiacum*, the robust Spotted Gentian *Gentiana punctata* and the deep-red beaked flowers of Kerner's Lousewort *Pedicularis kernerii*. A final section, through the forest, brought us back to the Belvedere, where we had refreshments, before catching the train at Alp Grum.

Day 5

Thursday 9th July

Clear skies brought perfect weather for our last three days. We took the cable-car to the dizzy heights of the Diavolezza restaurant, from where we looked out over a stunning panorama of peaks from Piz Palu to Piz Morteratsch and the ice-fields and glaciers draining from them. We followed the ridge, leading towards Munt Pers, where huge white mounds of Glacier Crowfoot *Ranunculus glacialis* decorated the screes. Further on was Creeping Avens, *Geum reptans* with glowing yellow flowers and red runners, and nestling in the rocks were bright blue patches of King-of-the-Alps *Eritrichium nanum* and pale pink splashes of Alpine Rock-jasmine *Androsace alpina*.

We returned on the cable-car to the valley, crossed the road, and had our lunch at the foot of Val da Fain. Nearby, a series of springs and wet flushes provided a fascinating range of plants, including the rare Flecked Marsh-orchid *Dactylorhiza cruenta*, Marsh Lousewort *Pedicularis palustris* and the easily-overlooked Marsh Arrowgrass *Triglochin palustris*. Below the limestone cliffs of Piz Alv, we compared the huge pleated leaves of Great Yellow Gentian *Gentiana lutea* and White False-helleborine *Veratrum album* and found the rare hybrid between Greater Yellow Gentian and Spotted Gentian *Gentiana x charpentieri*. Martagon Lily, Alpine Clematis *Clematis alpina* and Alpine Columbine *Aquilegia alpina* were flowering around the shrub-clad boulders that dotted the hillside. Some of us continued to the limestone screes, running down from the mountain summit, where there was an interesting mixture of dwarf shrubs, including *Dryas octopetala*, dwarf willows, Alpine Bearberry

Arctostaphylos alpinus and cushions of Blue Saxifrage *Saxifraga caesia*. Hidden among these, we found the tiny Dwarf Orchid *Chamorchis alpina*. We, eventually, had to tear ourselves away from this botanical wonderland and followed the farm track to the railway at Bernina Suot, where we caught our train.

Day 6

Friday 10th July

For a change, this morning, we caught the bus, in Pontresina, and travelled through St Moritz and past the lakes to Surlej. We took the cable-cars all the way to the top station, at 3,300 metres, just below Piz Murtel, where we enjoyed a magnificent 360° panorama of the surrounding peaks and valleys. Returning to Murtel, on the cable-car, we followed a broad track to Fuorcla Surlej. Wet hillsides had lots of *Soldanella pusilla* and purple *Primula integrifolia*, where the snow had not long receded, and on the rocks were *Primula latifolia*, occasional hybrids between it and *Primula integrifolia* (*Primula* x *muretiana*) and the yellow *Saxifraga seguieri*. On our final ascent to the col, there were more exquisite blue cushions of King-of-the-Alps *Eritrichium nanum* on the cliffs. We had our lunch overlooking the peaks and glaciers to the south and the ice-field at the head of Val Roseg.

It was then a long descent to the valley floor at Hotel Roseg, with the flora gradually changing from that of high-mountain to forest. Here was the option of a horse-drawn carriage or a much more interesting seven kilometre walk. The open forest of Arolla Pine provides a magical environment for birds, such as Great Spotted Woodpecker, Spotted Nutcracker and Eurasian Nuthatch, and an interesting ground flora, with occasional One-flowered Wintergreen *Moneses uniflora* and carpets of Twinflower *Linnaea borealis*, covering mossy banks and boulders. We finally emerged from the forest at Surovas Station and the familiar walk back to the hotel.

Day 7

Saturday 11th July

Our final full day had arrived, all too soon. We travelled by bus to St. Moritz, boarded the two-stage funicular railway to Corviglia and finally the cable-car to Piz Nair. The views from the restaurant were amazing and we enjoyed these over coffee, before venturing out and climbing the short distance to the summit, where cushions of King-of-the-Alps, White Musky Saxifrage *Saxifraga exarata* subsp. *exarata* and dark-blue *Gentiana brachyphylla* vied for our attention. A rocky knoll, a little way down the mountain, was home to a similar community, with the addition of Alpine Toadflax *Linaria alpina*, the tiny yellow Glacier Cinquefoil *Potentilla frigida*, Alpine Rock-jasmine *Androsace alpina* and the aromatic, silver-leaved Black Genipi *Artemisia genipi*.

We returned to Corviglia and enjoyed a leisurely lunch, in the sunshine, before continuing down the mountain on foot. The path threaded its way through a fascinating area of sugar limestones, with many deep solution-hollows creating an almost lunar landscape. An astonishing diversity of flowers included lots of Edelweiss *Leontopodium alpinum*, many intensely-fragrant Short-spurred Fragrant Orchids *Gymnadenia odoratissima* and Lesser Butterfly Orchid *Platanthera bifolia* and we were delighted to find more Dwarf Orchid *Chamorchis alpina*, in flower. Further on, wet pastures were thick with Broad-leaved Marsh-orchid *Dactylorhiza majalis* and a thorough search yielded some nice specimens of Round-headed Orchid *Traunsteinera globosa*. The walk finished at Chantarella, where we caught the funicular back to St. Moritz and relaxed over drinks and ice-creams. It had been an amazing day, with some of the best flora that the region has to offer and the perfect way to end our week in the Upper Engadine.

Day 8

Sunday 12th July

After breakfast and packing, we had time for a short excursion to see the rare Northern Dragon-head *Dracocephalum ruyschiana* and the delicate fern, Oblong Woodsia *Woodsia ilvensis*. We thanked the hotel staff for their tireless efforts to make our stay such a pleasant one, before being driven down to the station. The scenic journey back to Zurich went smoothly, leaving plenty of time to check-in for our return flight to London.

Receive our e-newsletter

Join the Naturetrek e-mailing list and be the first to hear about new tours, additional departures and new dates, tour reports and special offers. Visit www.naturetrek.co.uk to sign up.

Naturetrek Facebook

We are delighted to launch the Naturetrek Facebook page so that participants of Naturetrek tours can remain in touch after the holiday and share photos, comments and future travel plans.

Setting up a personal profile at www.facebook.com is quick, free and easy. The [Naturetrek Facebook page](#) is now live; do please pay us a visit!

Species Lists

Plants (‡ = cultivated or naturalised)

| Latin Name | Common Name | Notes |
|---------------------------------|-----------------------|-----------------------------|
| LYCOPODIOPSIDA | | |
| Lycopodiaceae | | |
| <i>Diphasiastrum alpinum</i> | Alpine Clubmoss | Occasional. Mountain turf |
| <i>Huperzia selago</i> | Fir Clubmoss | Frequent. Acidic pasture |
| <i>Lycopodium annotinum</i> | Interrupted Clubmoss | Rare. Forest. Alp Grum |
| <i>Lycopodium clavatum</i> | Stag's-horn Clubmoss | Mountain turf |
| Selaginellaceae | | |
| <i>Selaginella selaginoides</i> | Lesser Clubmoss | Frequent in turf |
| EQUISETOPSIDA | | |
| Equisetaceae | | |
| <i>Equisetum arvense</i> | Field Horsetail | Meadows and tracksides |
| <i>Equisetum hyemale</i> | Rough Horsetail | Limestone turf. Val da Fain |
| <i>Equisetum palustre</i> | Marsh Horsetail | Wet pasture |
| <i>Equisetum sylvaticum</i> | Wood Horsetail | Woods and wet pasture |
| <i>Equisetum variegatum</i> | Variiegated Horsetail | Wet flushes |
| PTEROPSIDA | | |
| Aspleniaceae | | |
| <i>Asplenium ruta-muraria</i> | Wall-rue | Walls |
| <i>Asplenium septentrionale</i> | Forked Spleenwort | Rare on rocks |
| <i>Asplenium trichomanes</i> | Maidenhair Spleenwort | Walls and cliffs |
| <i>Asplenium viride</i> | Green Spleenwort | Damp mountain rocks |
| Dennstaedtiaceae | | |
| <i>Pteridium aquilinum</i> | Bracken | Valley woodland |
| Dryopteridaceae | | |
| <i>Dryopteris carthusiana</i> | Narrow Buckler-fern | Damp scrub |
| <i>Dryopteris dilatata</i> | Broad Buckler-fern | Frequent |
| <i>Dryopteris filix-mas</i> | Male-fern | Widespread |
| <i>Gymnocarpium dryopteris</i> | Oak Fern | Woods. Local |
| <i>Gymnocarpium robertianum</i> | Limestone Fern | Calcareous scree |
| <i>Polystichum lonchitis</i> | Holly-fern | Mountain rocks |
| Ophioglossaceae | | |
| <i>Botrychium lunaria</i> | Adder's-tongue | Common in short turf |
| Polypodiaceae | | |
| <i>Polypodium vulgare</i> | Common Polypody | Occasional on rocks |
| Pteridaceae | | |
| <i>Cryptogramma crispa</i> | Maidenhair Fern | Acidic rocks and screes |

| Latin Name | Common Name | Notes |
|---|--|--|
| Thelypteridaceae <i>Phegopteris connectilis</i> | Marsh Fern Family Beech Fern | Woods. Occasional |
| Woodsiaceae <i>Athyrium distentifolium</i> <i>Athyrium filix-femina</i> <i>Cystopteris fragilis</i> <i>Woodsia alpina</i> <i>Woodsia ilvensis</i> | Lady-fern Family Alpine Lady-fern Lady-fern Brittle Bladder-fern Alpine Woodsia Oblong Woodsia | Mountain rocks Occasional Frequent on rocks Rare on cliffs Rare on scree |
| PINOPSIDA | CONIFERS | |
| Cupressaceae <i>Juniperus communis</i> subsp. <i>nana</i> | Juniper Family Common Juniper | Common |
| Pinaceae <i>Abies alba</i> <i>Larix decidua</i> <i>Picea abies</i> <i>Pinus cembra</i> <i>Pinus mugo</i> <i>Pinus sylvestris</i> | Pine Family European Silver-fir Larch Norway Spruce Arolla Pine Dwarf Mountain-pine Scots Pine | Occasional in forests Frequent Common in northern valleys Forests Forests at the treeline Common in northern valleys |
| MAGNOLIOPSIDA | FLOWERING PLANTS | |
| MAGNOLIIDAE | DICOTYLEDONS | |
| Apiaceae <i>Aegopodium podagraria</i> <i>Angelica sylvestris</i> <i>Astrantia minor</i> <i>Bupleurum stellatum</i> <i>Chaerophyllum hirsutum</i> <i>Chaerophyllum villarsii</i> <i>Heracleum sphondylium</i> <i>Imperatoria ostruthium</i> <i>Laserpitium gaudinii</i> <i>Laserpitium halleri</i> <i>Laserpitium latifolium</i> <i>Ligusticum mutellina</i> <i>Ligusticum mutellinoides</i> <i>Pimpinella major</i> <i>Pimpinella saxifraga</i> | Carrot Family Ground Elder Wild Angelica Lesser Astrantia Starry Hare's-ear Hairy Chervil Villars' Chervil Hogweed Masterwort Gaudin's Sermountain Haller's Sermountain Broad-leaved Sermountain Alpine Lovage Unbranched Lovage Greater Burnet-saxifrage Burnet-saxifrage | Waste places Occasional. Lower valleys Common Lago Bianco, Val Minor Woods Woods Common Common Rocky places Rocky places Common Mountain pasture High mountain pasture Meadows Meadows |
| Araliaceae <i>Hedera helix</i> | Ivy Family Ivy | Occasional in woods |
| Asteraceae <i>Adenostyles alliariae</i> <i>Adenostyles leucophylla</i> <i>Achillea atrata</i> <i>Achillea erba-rotta</i> subsp. <i>moschata</i> <i>Achillea macrophylla</i> | Daisy Family Adenostyles White-leaved Adenostyles Dark-stemmed Sneezewort Musk Milfoil Large-leaved Sneezewort | Val Roseg River gravels. Morteratsch Rocky places on limestone Rocky places on acidic soils Occasional |

| Latin Name | Common Name | Notes |
|---|---------------------------|--------------------------------|
| <i>Achillea millefolium</i> | Yarrow | Frequent |
| <i>Achillea nana</i> | Dwarf Milfoil | Screes |
| <i>Antennaria carpatica</i> | Carpathian Cat's-foot | Mountain turf |
| <i>Antennaria dioica</i> | Cat's-foot | Common |
| <i>Arnica montana</i> | Arnica | Common |
| <i>Artemisia genipi</i> | Black Genipi | Screes |
| <i>Artemisia umbelliformis</i> | Yellow Genipi | Screes and cliffs |
| <i>Aster alpinus</i> | Alpine Aster | Limestone rocks and turf |
| <i>Aster bellidiastrum</i> | False Aster | Frequent |
| <i>Bellis perennis</i> | Daisy | Frequent |
| <i>Carduus defloratus</i> | Alpine Thistle | Common |
| <i>Carduus personata</i> | Great Marsh Thistle | Pontresina |
| <i>Carlina acaulis</i> | Stemless Carlina Thistle | Mountain turf |
| <i>Carlina vulgaris</i> | Common Carlina Thistle | Occasional. Lower valleys |
| <i>Centaurea scabiosa</i> | Greater Knapweed | Meadows |
| <i>Cicerbita alpina</i> | Alpine Blue-sowthistle | Forest below Alp Grum |
| <i>Cirsium acaule</i> | Dwarf Thistle | Occasional in pasture |
| <i>Cirsium eriophorum</i> | Woolly Thistle | Morteratsch |
| <i>Cirsium erisithales</i> | Yellow Melancholy Thistle | Alp Grum |
| <i>Cirsium heterophyllum</i> | Melancholy Thistle | Meadows |
| <i>Cirsium palustre</i> | Marsh Thistle | Damp places |
| <i>Cirsium spinosissimum</i> | Spiniest Thistle | Mountain turf |
| <i>Crepis aurea</i> | Golden Hawk's-beard | Common |
| <i>Crepis bocconeii</i> | Boccone's Hawk's-beard | Damp pasture |
| <i>Crepis paludosa</i> | Marsh Hawk's-beard | Wet meadows |
| <i>Crepis uniflora</i> | One-flowered Hawk's-beard | Rocky pasture |
| <i>Doronicum clusii</i> | Clusius' Leopard's-bane | Screes |
| <i>Erigeron acris</i> subsp. <i>politus</i> | Blue Fleabane | Moraines |
| <i>Erigeron alpinus</i> | Alpine Fleabane | Local |
| <i>Erigeron neglectus</i> | Neglected Fleabane | Local |
| <i>Erigeron uniflorus</i> | One-flowered Fleabane | Local |
| <i>Eupatorium cannabinum</i> | Hemp Agrimony | Lowland |
| <i>Gnaphthodium norvegicum</i> | Norwegian Cudweed | Occasional in mountain pasture |
| <i>Gnaphthodium supinum</i> | Dwarf Cudweed | Snowbeds |
| <i>Hieracium alpinum</i> | Alpine Hawkweed | Frequent |
| <i>Hieracium angustifolium</i> | Glacier Hawkweed | Morteratsch |
| <i>Hieracium aurantiacum</i> † | Orange Hawkweed | Pontresina |
| <i>Hieracium hoppeanum</i> | Hoppe's Hawkweed | Stony places |
| <i>Hieracium intybaceum</i> | Whitish Hawkweed | Stony places |
| <i>Hieracium lactucella</i> | | Common |
| <i>Hieracium murorum</i> | | Occasional |
| <i>Hieracium piliferum</i> | | Stony places |
| <i>Hieracium pilosella</i> | Mouse-ear Hawkweed | Common |
| <i>Hieracium staticifolium</i> | Stalice-leaved Hawkweed | Morteratsch |
| <i>Hieracium villosum</i> | Hairy Hawkweed | Stony pasture |
| <i>Homogyne alpina</i> | Alpine Colt's-foot | Common |
| <i>Hypochaeris uniflora</i> | Giant Cat's-ear | Alp Grum |
| <i>Jacobaea abrotanifolia</i> | Pinnate-leaved Ragwort | Occasional. Local |
| <i>Jacobaea incana</i> subsp. <i>carniolica</i> | Carniolian Ragwort | Locally common |
| <i>Lapsana communis</i> | Nipplewort | Waste places |
| <i>Leontodon helveticus</i> | Swiss Hawkbit | Mountain turf |
| <i>Leontodon hispidus</i> | Alpine Hawkbit | Acidic pasture |
| <i>Leontopodium alpinum</i> | Edelweiss | Limestone turf |

| Latin Name | Common Name | Notes |
|--|--------------------------------|---------------------------|
| <i>Leucanthemopsis alpina</i> | Alpine Moon Daisy | Common. Mountain rocks |
| <i>Leucanthemum atratum</i> agg. | Saw-leaved Moon Daisy | Mountain turf |
| <i>Leucanthemum vulgare</i> | Ox-eye Daisy | Meadows |
| <i>Matricaria discoidea</i> | Pineappleweed | Ruderal |
| <i>Mycelis muralis</i> | Wall Lettuce | Walls |
| <i>Prenanthes purpurea</i> | Purple Lettuce | Forests |
| <i>Saussurea alpina</i> | Alpine Saw-wort | Mountain turf |
| <i>Scorzoneroides autumnalis</i> | Autumn Hawkbit | Stony pasture |
| <i>Senecio doricum</i> | Chamois Ragwort | Locally common |
| <i>Senecio hercynicus</i> | Shrub Ragwort | Forest below Alp Grum |
| <i>Senecio ovatus</i> | Wood Ragwort | Forests |
| <i>Senecio rupestris</i> | Rock Groundsel | Rocky places |
| <i>Senecio vulgaris</i> | Groundsel | Ruderal |
| <i>Solidago virgaurea</i> subsp. <i>minuta</i> | Goldenrod | Mountain rocks |
| <i>Stemnacantha rhapsantica</i> | Giant Stemnacantha | Occasional in meadows |
| <i>Taraxacum alpinum</i> | Alpine Dandelion | Mountain rocks |
| <i>Taraxacum officinale</i> | Dandelion | Widespread |
| <i>Tragopogon pratensis</i> | Goat's-beard | Meadows |
| <i>Tussilago farfara</i> | Colt's-foot | Common. Damp rocky places |
| Betulaceae | Birch Family | |
| <i>Alnus glutinosa</i> | Common Alder | Valleys, by rivers |
| <i>Alnus incana</i> | Grey Alder | Valleys, by rivers |
| <i>Alnus viridis</i> | Green Alder | Mountain scrub |
| <i>Betula pendula</i> | Silver Birch | Widespread |
| <i>Betula pubescens</i> | Downy Birch | Frequent |
| Boraginaceae | Borage Family | |
| <i>Echium vulgare</i> | Viper's Bugloss | Waste places |
| <i>Eritrichium nanum</i> | King-of-the-Alps | Acidic mountain rocks |
| <i>Myosotis alpestris</i> | Alpine Forget-me-not | Common |
| <i>Myosotis sylvatica</i> | Wood Forget-me-not | Woods |
| <i>Pulmonaria australis</i> | Southern Lungwort | Above Pontresina |
| Brassicaceae | Cabbage Family | |
| <i>Arabis alpina</i> subsp. <i>alpina</i> | Alpine Rock-cress | Mountain rocks |
| <i>Arabis soyeri</i> subsp. <i>subcoriacea</i> | Soyer's Rock-cress | Rocky places |
| <i>Barbarea vulgaris</i> | Winter-cress | Pontresina |
| <i>Biscutella laevigata</i> | Buckler Mustard | Rocky places |
| <i>Capsella bursa-pastoris</i> | Shepherd's-purse | Widespread |
| <i>Cardamine alpina</i> | Alpine Bitter-cress | Acidic rocks and scree |
| <i>Cardamine amara</i> | Large Bitter-cress | Streamsides |
| <i>Cardamine resedifolia</i> | Mignonette-leaved Bitter-cress | Mountain rocks |
| <i>Draba aizoides</i> | Yellow Whitlowgrass | Rocky places on limestone |
| <i>Draba dubia</i> | Austrian Whitlowgrass | Acidic rocks. Corvatsch |
| <i>Draba fladnizensis</i> | Bald White Whitlowgrass | Alpine rocks |
| <i>Draba tomentosa</i> | Tomentose Whitlowgrass | Limestone rocks |
| <i>Kernera saxatilis</i> | Rock Kernera | Rocks |
| <i>Pritzelago alpina</i> | Chamois Cress | Mountain rocks |
| <i>Sisymbrium austriacum</i> | Austrian Rocket | Waste places |
| <i>Thlaspi perfoliatum</i> | Perfoliate Penny-cress | Near habitation |

| Latin Name | Common Name | Notes |
|--|---------------------------|---------------------------|
| Campanulaceae | Bellflower Family | |
| <i>Campanula barbata</i> | Bearded Bellflower | Mountain turf |
| <i>Campanula cochlearifolia</i> | Fairy's-thimble | Mountain rocks |
| <i>Campanula glomerata</i> | Clustered Bellflower | Occasional. Meadows |
| <i>Campanula scheuchzeri</i> | Scheuchzer's Bellflower | Mountain turf |
| <i>Campanula thyrsoides</i> | Yellow Bellflower | Alp Nova |
| <i>Campanula trachelium</i> | Nettle-leaved Bellflower | Woods |
| <i>Phyteuma betonicifolium</i> | Betony-leaved Rampion | Common |
| <i>Phyteuma globulariifolium</i> | Globularia-leaved Rampion | Alpine rocks |
| <i>Phyteuma hedraianthifolium</i> | Rhaetian Rampion | Rocks. Lago Bianco |
| <i>Phyteuma hemisphaericum</i> | Globe-headed Rampion | Acidic rocks |
| <i>Phyteuma orbiculare</i> | Round-headed Rampion | Meadows. Frequent |
| <i>Phyteuma scheuchzeri</i> | Scheuchzer's Rampion | Occasional |
| Caprifoliaceae | Honeysuckle Family | |
| <i>Linnaea borealis</i> | Twinflower | Locally common in forests |
| <i>Lonicera alpigena</i> | Alpine Honeysuckle | Forests |
| <i>Lonicera caerulea</i> | Blue-berried Honeysuckle | Rocky places |
| <i>Lonicera nigra</i> | Black-berried Honeysuckle | Woods |
| <i>Lonicera xylosteum</i> | Fly Honeysuckle | Woods |
| <i>Sambucus nigra</i> | Common Elder | Woods |
| <i>Sambucus racemosa</i> | Alpine Elder | Frequent |
| Caryophyllaceae | Pink Family | |
| <i>Arenaria biflora</i> | Two-flowered Sandwort | Scree |
| <i>Arenaria ciliata</i> | Ciliate-leaved Sandwort | Scree |
| <i>Cerastium alpinum</i> | Alpine Mouse-ear | Mountain turf |
| <i>Cerastium arvense</i> subsp. <i>strictum</i> | Field Mouse-ear | Frequent |
| <i>Cerastium cerastoides</i> | Starwort Mouse-ear | Rocks and pasture |
| <i>Cerastium fontanum</i> | Common Mouse-ear | Common |
| <i>Cerastium latifolium</i> | Broad-leaved Mouse-ear | Moraines and scree |
| <i>Dianthus carthusianorum</i> subsp. <i>vaginatus</i> | Carthusian Pink | Local. Open woodland |
| <i>Dianthus superbus</i> | Large Pink | Damp turf |
| <i>Dianthus sylvestris</i> | Wood Pink | Rocky places |
| <i>Gypsophila repens</i> | Alpine Gypsophila | Frequent |
| <i>Herniaria glabra</i> | Smooth Rupturewort | Railway ballast |
| <i>Minuartia biflora</i> | Northern Sandwort | Wet mountain turf |
| <i>Minuartia sedoides</i> | Mossy Cyphel | Alpine rocks and turf |
| <i>Minuartia rupestris</i> | Rock Sandwort | Rocks and scree |
| <i>Minuartia verna</i> | Vernal Sandwort | Stony places |
| <i>Moehringia ciliata</i> | Creeping Sandwort | Scree |
| <i>Sagina procumbens</i> | Procumbent Pearlwort | Common |
| <i>Sagina saginoides</i> | Alpine Pearlwort | Alpine turf and scree |
| <i>Saponaria ocymoides</i> | Rock Soapwort | Chantarella |
| <i>Scleranthus annuus</i> | Annual Knawel | Occasional |
| <i>Silene acaulis</i> | Moss Campion | Mountain turf |
| <i>Silene dioica</i> | Red Campion | Meadows |
| <i>Silene exscapa</i> | Stemless Moss Campion | Mountain turf |
| <i>Silene flos-cuculi</i> | Ragged-Robin | Wet meadows. Celerina |
| <i>Silene nutans</i> | Nottingham Catchfly | Rocks and meadows |
| <i>Silene rupestris</i> | Rock Campion | Rocks |
| <i>Silene vulgaris</i> | Bladder Campion | Meadows |
| <i>Stellaria holostea</i> | Greater Stichwort | Occasional |

| Latin Name | Common Name | Notes |
|--|--------------------------------|---------------------------------|
| <i>Stellaria graminea</i> | Lesser Stitchwort | Woods and meadows |
| <i>Stellaria media</i> | Common Chickweed | Occasional |
| <i>Stellaria nemorum</i> | Wood Stitchwort | Woods |
| Chenopodiaceae | Goosefoot Family | |
| <i>Chenopodium album</i> | Fat-hen | Occasional weed |
| <i>Chenopodium bonus-henricus</i> | Good-King-Henry | Frequent weed |
| Cistaceae | Rock-rose Family | |
| <i>Helianthemum nummularium</i> subsp. <i>grandiflorum</i> | Large-flowered Common Rockrose | Rocky places |
| <i>Helianthemum oelandicum</i> subsp. <i>alpestre</i> | Alpine Rockrose | Calcareous turf |
| Crassulaceae | Stonecrop Family | |
| <i>Sedum acre</i> | Biting Stonecrop | Frequent |
| <i>Sedum alpestre</i> | Alpine Stonecrop | Mountain rocks |
| <i>Sedum annuum</i> | Annual Stonecrop | Moraines. Morteratsch |
| <i>Sedum atratum</i> | Dark Stonecrop | Mountain rocks |
| <i>Sedum villosum</i> | Pink Stonecrop | Acidic flushes. Rare. Val Roseg |
| <i>Sempervivum arachnoideum</i> | Cobweb Houseleek | Common. Stony places |
| <i>Sempervivum montanum</i> | Mountain Houseleek | Common. Stony places |
| <i>Sempervivum tectorum</i> | Common Houseleek | Occasional. Rocky turf |
| <i>Sempervivum wulfenii</i> | Wulfen's Houseleek | Occasional. Rocky turf |
| Dipsacaceae | Teasel Family | |
| <i>Knautia dipsacifolia</i> | Wood Scabious | Shady places |
| <i>Scabiosa columbaria</i> | Small Scabious | Occasional |
| Empetraceae | Crowberry Family | |
| <i>Empetrum nigrum</i> subsp. <i>hermaphroditum</i> | Hermaphrodite Crowberry | Acidic heaths |
| Ericaceae | Heath Family | |
| <i>Arctostaphylos alpinus</i> | Alpine Bearberry | Limestone turf and screes |
| <i>Arctostaphylos uva-ursi</i> | Bearberry | Rocky pasture |
| <i>Calluna vulgaris</i> | Ling | Frequent |
| <i>Kalmia procumbens</i> | Trailing Azalea | Acidic heaths |
| <i>Rhododendron ferrugineum</i> | Alpenrose | Common |
| <i>Rhododendron hirsutum</i> | Hairy Alpenrose | Limestone rocks |
| <i>Vaccinium myrtillus</i> | Bilberry | Common |
| <i>Vaccinium uliginosum</i> | Bog Whortleberry | Common |
| <i>Vaccinium vitis-idaea</i> | Cowberry | Common |
| Euphorbiaceae | Spurge Family | |
| <i>Euphorbia cyparissias</i> | Cypress Spurge | Frequent on limestone |
| Fabaceae | Pea Family | |
| <i>Anthyllis vulneraria</i> subsp. <i>alpestris</i> | Alpine Kidney Vetch | Mountain pasture |
| <i>Astragalus alpinus</i> | Alpine Milk-vetch | Mountain turf |
| <i>Astragalus australis</i> | Southern Milk-vetch | Mountain turf |
| <i>Astragalus frigidus</i> | Pallid Milk-vetch | Limestone pasture |
| <i>Astragalus penduliflorus</i> | Mountain Lentil | Pasture and stony places. Local |
| <i>Hedysarum hedysaroides</i> | Alpine Sainfoin | Mountain pasture |
| <i>Hippocrepis comosa</i> | Horseshoe Vetch | Common on limestone |

| Latin Name | Common Name | Notes |
|---|------------------------------|---------------------------|
| <i>Lathyrus pratensis</i> | Meadow Vetchling | Meadows |
| <i>Lotus alpinus</i> | Alpine Bird's-foot Trefoil | Mountain turf |
| <i>Lotus corniculatus</i> | Bird's-foot Trefoil | Meadows |
| <i>Medicago lupulina</i> | Black Medick | Common |
| <i>Medicago sativa</i> † | Lucerne | Occasional |
| <i>Melilotus albus</i> | White Melilot | Waste places |
| <i>Melilotus officinalis</i> | Ribbed Melilot | Waste places |
| <i>Onobrychis arenaria</i> | Small Sainfoin | Occasional. Pontresina |
| <i>Onobrychis montana</i> | Mountain Sainfoin | Mountain pasture |
| <i>Oxytropis campestris</i> | Yellow Milk-vetch | Common in pasture |
| <i>Oxytropis jacquinii</i> | Mountain Milk-vetch | Stony places on limestone |
| <i>Oxytropis lapponica</i> | Northern Milk-vetch | Stony places |
| <i>Trifolium alpinum</i> | Alpine Clover | Common |
| <i>Trifolium badium</i> | Brown Clover | Mountain meadows |
| <i>Trifolium montanum</i> | Mountain Clover | Frequent |
| <i>Trifolium pallescens</i> | Pale Clover | Pasture and rocky places |
| <i>Trifolium pratense</i> | Red Clover | Meadows |
| <i>Trifolium repens</i> | White Clover | Common |
| <i>Trifolium thalii</i> | Thal's Clover | Mountain turf |
| <i>Vicia cracca</i> | Tufted Vetch | Frequent |
| <i>Vicia sepium</i> | Bush Vetch | Occasional |
| Gentianaceae | Gentian Family | |
| <i>Gentiana acaulis</i> | Trumpet Gentian | Mountain pasture |
| <i>Gentiana bavarica</i> | Bavarian Gentian | Marshes |
| <i>Gentiana brachyphylla</i> | Short-leaved Gentian | Alpine turf |
| <i>Gentiana x charpentieri</i> (<i>G. lutea</i> x <i>G. punctata</i>) | Charpentier's Gentian | With both parents |
| <i>Gentiana clusii</i> | Clusius' Gentian | Limestone turf |
| <i>Gentiana lutea</i> | Great Yellow Gentian | Limestone pasture |
| <i>Gentiana nivalis</i> | Snow Gentian | Mountain turf |
| <i>Gentiana orbicularis</i> | Orbiculate-leaved Gentian | Occasional |
| <i>Gentiana punctata</i> | Spotted Gentian | Mountain pasture |
| <i>Gentiana verna</i> | Spring Gentian | Common |
| <i>Gentianella campestris</i> | Field Gentian | Pasture |
| Geraniaceae | Crane's-bill Family | |
| <i>Geranium robertianum</i> | Herb Robert | Common |
| <i>Geranium sylvaticum</i> | Wood Crane's-bill | Common |
| Globulariaceae | Globularia Family | |
| <i>Globularia cordifolia</i> | Matted Globularia | Frequent on limestone |
| <i>Globularia nudicaulis</i> | Leafless-stemmed Globularia | Frequent on limestone |
| Hypericaceae | St John's-wort Family | |
| <i>Hypericum maculatum</i> | Imperforate St John's-wort | Frequent |
| Lamiaceae | Dead-nettle Family | |
| <i>Ajuga pyramidalis</i> | Pyramidal Bugle | Locally common |
| <i>Ajuga reptans</i> | Common Bugle | Common |
| <i>Clinopodium alpinum</i> | Alpine Calamint | Rocky turf on limestone |
| <i>Dracocephalum ruyschiana</i> | Northern Dragon-head | Open forest. Rare |
| <i>Galeopsis tetrahit</i> | Common Hemp-nettle | Occasional |
| <i>Horminum pyrenaicum</i> | Dragon-mouth | Val Minor |

| Latin Name | Common Name | Notes |
|---|----------------------------------|------------------------------|
| <i>Lamiasstrum galeobdolon</i> | Yellow Archangel | Woods. Alp Grum |
| <i>Lamium album</i> | White Dead-nettle | Frequent |
| <i>Prunella vulgaris</i> | Self-heal | Common |
| <i>Salvia pratensis</i> | Meadow Clary | Roadside banks. Pontresina |
| <i>Teucrium montanum</i> | Mountain Germander | Limestone rocks. Chantarella |
| <i>Thymus polytrichus</i> | Wild Thyme | Common |
| Lentibulariaceae | Butterwort Family | |
| <i>Pinguicula alpina</i> | Alpine Butterwort | Mountain turf and rocks |
| <i>Pinguicula leptoceras</i> | Southern Butterwort | Wet flushes |
| <i>Pinguicula vulgaris</i> | Common Butterwort | Wet flushes |
| Linaceae | Flax Family | |
| <i>Linum catharticum</i> | Fairy Flax | Common |
| Onagraceae | Willowherb Family | |
| <i>Chamerion angustifolium</i> | Rosebay Willowherb | Occasional |
| <i>Epilobium alpestre</i> | Whorled-leaved Willowherb | Frequent |
| <i>Epilobium alsinifolium</i> | Chickweed Willowherb | Frequent |
| <i>Epilobium anagallidifolium</i> | Alpine Willowherb | Frequent |
| <i>Epilobium fleischeri</i> | Fleischer's Willowherb | River gravels and moraines |
| <i>Epilobium montanum</i> | Broad-leaved Willowherb | Occasional |
| Oxalidaceae | Wood-sorrel Family | |
| <i>Oxalis acetosella</i> | Wood-sorrel | Shaded rocks |
| Papaveraceae | Poppy Family | |
| <i>Papaver aurantiacum</i> | Rhaetic Poppy | Scree. Local |
| <i>Papaver croceum</i> † | Islandic Poppy | River gravels. |
| Parnassiaceae | Grass-of-Parnassus Family | |
| <i>Parnassia palustris</i> | Grass-of-Parnassus | Damp flushes |
| Plantaginaceae | Plantain Family | |
| <i>Plantago alpina</i> | Alpine Plantain | Mountain turf |
| <i>Plantago atrata</i> | Dark Plantain | Mountain turf |
| <i>Plantago maritima</i> subsp. <i>serpentina</i> | Fleshy Plantain | Frequent |
| <i>Plantago media</i> | Hoary Plantain | Occasional |
| Polemoniaceae | Jacob's-ladder Family | |
| <i>Polemonium caeruleum</i> | Jacob's-ladder | Limestone rocks |
| Polygalaceae | Milkwort Family | |
| <i>Polygala alpestris</i> | Mountain Milkwort | Common |
| <i>Polygala alpina</i> | Alpine Milkwort | Mountain turf |
| <i>Polygala chamaebuxus</i> | Shrubby Milkwort | Frequent |
| Polygonaceae | Dock Family | |
| <i>Oxyria digyna</i> | Mountain Sorrel | Rocks and screes |
| <i>Persicaria bistorta</i> | Common Bistort | Meadows |
| <i>Persicaria vivipara</i> | Alpine Bistort | Common |
| <i>Rumex acetosa</i> | Common Sorrel | Widespread |
| <i>Rumex acetosella</i> | Sheep's Sorrel | Acidic pasture |

| Latin Name | Common Name | Notes |
|---|----------------------------|-------------------------|
| <i>Rumex alpestris</i> | Alpine Sorrel | Widespread |
| <i>Rumex alpinus</i> | Monk's-rhubarb | Disturbed areas |
| <i>Rumex obtusifolius</i> | Broad-leaved Dock | Occasional |
| <i>Rumex scutatus</i> | French Sorrel | Screes and rocky places |
| Portulacaceae | Purslane Family | |
| <i>Montia fontana</i> | Blinks | Springs |
| Primulaceae | Primrose Family | |
| <i>Androsace alpina</i> | Alpine Rock-jasmine | Alpine scree |
| <i>Androsace chamaejasme</i> | Ciliate Rock-jasmine | Frequent |
| <i>Androsace obtusifolia</i> | Blunt-leaved Rock Jasmine | Limestone turf |
| <i>Primula x berninae</i> (<i>P. latifolia</i> x <i>P. hirsuta</i>) | A hybrid Primrose | Val Minor |
| <i>Primula elatior</i> | Oxlip | Snowbeds |
| <i>Primula farinosa</i> | Bird's-eye Primrose | Mountain turf |
| <i>Primula halleri</i> | Long-flowered Primrose | Rare. Below Alp Grum |
| <i>Primula x heerii</i> (<i>P. hirsuta</i> x <i>P. integrifolia</i>) | A hybrid Primrose | Val Minor |
| <i>Primula hirsuta</i> | Hairy Primrose | Mountain rocks |
| <i>Primula integrifolia</i> | Entire-leaved Primrose | Common. Wet turf |
| <i>Primula latifolia</i> | Broad-leaved Primrose | Locally common |
| <i>Primula x muretiana</i> (<i>P. integrifolia</i> x <i>P. latifolia</i>) | A hybrid Primrose | Fuorcla Surlej |
| <i>Soldanella alpina</i> | Alpine Snowbell | Mountain turf |
| <i>Soldanella pusilla</i> | Dwarf Snowbell | Wet turf |
| <i>Trientalis europaea</i> | Chickweed Wintergreen | Local in forest |
| Pyrolaceae | Wintergreen Family | |
| <i>Moneses uniflora</i> | One-flowered Wintergreen | Forest |
| <i>Orthilia secunda</i> | Serrated Wintergreen | Forest |
| <i>Pyrola minor</i> | Lesser Wintergreen | Forest |
| <i>Pyrola rotundifolia</i> | Round-leaved Wintergreen | Alp Nova |
| Ranunculaceae | Buttercup Family | |
| <i>Aconitum lycotonum</i> subsp. <i>vulparia</i> | Wolf's-bane | Frequent. Forests |
| <i>Aconitum napellus</i> agg. | Monk's-hood | Damp pasture |
| <i>Aconitum napellus</i> subsp. <i>vulgare</i> | Compact Monk's-hood | Damp pasture |
| <i>Aconitum variegatum</i> subsp. <i>paniculatum</i> | Variiegated Monk's-hood | Damp pasture |
| <i>Anemone narcissiflora</i> | Narcissus-flowered Anemone | Mountain pasture |
| <i>Aquilegia alpina</i> | Alpine Columbine | Limestone pasture |
| <i>Caltha palustris</i> | Marsh Marigold | Flushes and streamsides |
| <i>Clematis alpina</i> | Alpine Clematis | Forest and pasture |
| <i>Pulsatilla alpina</i> subsp. <i>apiifolia</i> | Alpine Pasqueflower | Mountain pasture |
| <i>Pulsatilla vernalis</i> | Spring Pasqueflower | Mountain pasture |
| <i>Ranunculus aconitifolius</i> | Aconite-leaved Buttercup | Forest |
| <i>Ranunculus acris</i> | Meadow Buttercup | Common |
| <i>Ranunculus alpestris</i> | Alpine Buttercup | Common |
| <i>Ranunculus glacialis</i> | Glacier Crowfoot | Alpine scree |
| <i>Ranunculus kuepferi</i> | Kupfer's Buttercup | Mountain turf |
| <i>Ranunculus montanus</i> | Mountain Buttercup | Mountain turf |
| <i>Ranunculus platanifolius</i> | Maple-leaved Buttercup | Rare in open forest |
| <i>Ranunculus repens</i> | Creeping Buttercup | Meadows |
| <i>Thalictrum aquilegifolium</i> | Great Meadow-rue | Meadows |
| <i>Thalictrum foetidum</i> | Stinking Meadow-rue | Pasture |
| <i>Thalictrum minus</i> | Lesser Meadow-rue | Pasture |

| Latin Name | Common Name | Notes |
|---------------------------------|---------------------------|---------------------------|
| <i>Trollius europaeus</i> | Globeflower | Damp pasture |
| Rhamnaceae | Buckthorn Family | |
| <i>Rhamnus pumilus</i> | Dwarf Buckthorn | Mountain Rocks |
| Rosaceae | Rose Family | |
| <i>Alchemilla alpina</i> | Alpine Lady's Mantle | Common |
| <i>Alchemilla conjuncta</i> | Silver Lady's Mantle | Frequent |
| <i>Alchemilla glabra</i> | Smooth Lady's Mantle | Frequent |
| <i>Alchemilla hybrida</i> agg. | Hybrid Lady's Mantle | Occasional |
| <i>Alchemilla pentaphyllea</i> | Cut-leaved Lady's Mantle | Damp mountain turf |
| <i>Alchemilla vulgaris</i> agg. | Lady's-mantle | Common |
| <i>Alchemilla xanthochlora</i> | Pale Lady's-mantle | Frequent |
| <i>Cotoneaster integerrima</i> | Wild Cotoneaster | Rocks |
| <i>Dryas octopetala</i> | Mountain Avens | Mountain rocks and scree |
| <i>Filipendula ulmaria</i> | Meadowsweet | Meadows |
| <i>Fragaria vesca</i> | Wild Strawberry | Woods and pasture |
| <i>Geum montanum</i> | Alpine Avens | Mountain turf |
| <i>Geum reptans</i> | Creeping Avens | Alpine rocks and scree |
| <i>Geum rivale</i> | Water Avens | Occasional. Wet places |
| <i>Potentilla anserina</i> | Silverweed | Waste places |
| <i>Potentilla argentea</i> | Hoary Cinquefoil | Rocky places, open woods |
| <i>Potentilla aurea</i> | Golden Cinquefoil | Occasional. Mountain turf |
| <i>Potentilla crantzii</i> | Alpine Cinquefoil | Common. Mountain turf |
| <i>Potentilla erecta</i> | Tormentil | Acidic turf |
| <i>Potentilla frigida</i> | Glacier Cinquefoil | Acidic alpine rocks |
| <i>Potentilla grandiflora</i> | Large-flowered Cinquefoil | Acidic pasture |
| <i>Potentilla rupestris</i> | Rock Cinquefoil | Local |
| <i>Poterium sanguisorba</i> | Salad Burnet | Widespread |
| <i>Prunus padus</i> | Bird Cherry | Near Hotel Palu |
| <i>Rosa canina</i> | Dog-rose | Wooded slopes |
| <i>Rosa pendulina</i> | Alpine Rose | Frequent. Mountains |
| <i>Rubus idaeus</i> | Raspberry | Woods |
| <i>Rubus saxatilis</i> | Stone Bramble | Occasional. Rocky places |
| <i>Sanguisorba officinalis</i> | Great Burnet | Meadows |
| <i>Sibbaldia procumbens</i> | Sibbaldia | Stony places in mountains |
| <i>Sorbus aria</i> | Whitebeam | Occasional in forest |
| <i>Sorbus aucuparia</i> | Rowan | Woods and scrub |
| <i>Sorbus chamaemespilus</i> | False-medlar | Mountain scrub |
| Rubiaceae | Madder Family | |
| <i>Asperula cynanchica</i> | Squinancywort | Limestone rocks |
| <i>Cruciata laevipes</i> | Crosswort | Meadows |
| <i>Galium anisophyllum</i> | Dwarf Bedstraw | Frequent |
| <i>Galium boreale</i> | Northern Bedstraw | Occasional. Grassy places |
| <i>Galium mollugo</i> | Hedge Bedstraw | Occasional |
| <i>Galium odoratum</i> | Woodruff | Woods |
| <i>Galium pumilum</i> | Slender Bedstraw | Frequent |
| <i>Galium verum</i> | Lady's Bedstraw | Meadows |
| Salicaceae | Willow Family | |
| <i>Populus tremula</i> | Aspen | Occasional |
| <i>Salix aurita</i> | Eared Willow | Frequent |

| Latin Name | Common Name | Notes |
|---|-----------------------------------|----------------------------|
| <i>Salix breviserrata</i> | Finely-toothed Willow | Occasional |
| <i>Salix caprea</i> | Goat Willow | Lowland woods |
| <i>Salix eleagnos</i> | Hoary Willow | Occasional. Streamsides |
| <i>Salix foetida</i> | Stinking Willow | Val da Fain |
| <i>Salix glaucosericea</i> | Silky Willow | Occasional |
| <i>Salix hastata</i> | Large-stipuled Willow | Occasional |
| <i>Salix hegetschweileri</i> | Alpine Willow | Mountain turf |
| <i>Salix helvetica</i> | Swiss Willow | Rare. Val Muragl |
| <i>Salix herbacea</i> | Least willow | Mountain turf |
| <i>Salix laggeri</i> | Lagger's Willow | Occasional |
| <i>Salix myrsinites</i> | Whortle-leaved Willow | Mountain turf |
| <i>Salix reticulata</i> | Net-leaved Willow | Mountain rocks and turf |
| <i>Salix retusa</i> | Retuse-leaved Willow | Mountain rocks and turf |
| <i>Salix serpyllifolia</i> | Thyme-leaved Willow | Mountain rocks and turf |
| Santalaceae | Bastard-toadflax Family | |
| <i>Thesium alpinum</i> | Alpine Bastard-toadflax | Occasional. Pasture |
| <i>Thesium pyrenaicum</i> | Pyrenaean Bastard-toadflax | Rocky pasture |
| Saxifragaceae | Saxifrage Family | |
| <i>Chrysosplenium alternifolium</i> | Alternate-leaved Golden-saxifrage | Occasional. Wet places |
| <i>Saxifraga aizoides</i> | Yellow Mountain Saxifrage | Wet turf and screes |
| <i>Saxifraga androsacea</i> | Scree Saxifrage | Damp screes and rocks |
| <i>Saxifraga aspera</i> | Rough Saxifrage | Rocky places |
| <i>Saxifraga bryoides</i> | Mossy Saxifrage | Rocky places |
| <i>Saxifraga caesia</i> | Blue Saxifrage | Limestone scree and turf |
| <i>Saxifraga exarata</i> subsp. <i>exarata</i> | White Musky Saxifrage | Rocky places |
| <i>Saxifraga exarata</i> subsp. <i>moschata</i> | Musky Saxifrage | Rocky places |
| <i>Saxifraga oppositifolia</i> | Purple Saxifrage | Moraines and screes |
| <i>Saxifraga paniculata</i> | Livelong Saxifrage | Rocks and cliffs |
| <i>Saxifraga seguieri</i> | Seguier's Saxifrage | Damp alpine turf and rocks |
| <i>Saxifraga stellaris</i> | Starry Saxifrage | Wet acidic flushes |
| Scrophulariaceae | Figwort Family | |
| <i>Bartsia alpina</i> | Alpine Bartsia | Alpine turf |
| <i>Chaenorhinum minus</i> | Small Toadflax | Railway tracks |
| <i>Digitalis grandiflora</i> | Large Yellow Foxglove | Local. Woods |
| <i>Euphrasia alpina</i> | Alpine Eyebright | Mountain turf |
| <i>Euphrasia minima</i> | Dwarf Eyebright | Mountain rocks |
| <i>Linaria alpina</i> | Alpine Toadflax | Screes and rocky places |
| <i>Linaria vulgaris</i> | Common Toadflax | Below Alp Grum |
| <i>Melampyrum pratense</i> | Common Cow-wheat | Forest |
| <i>Melampyrum sylvaticum</i> | Small Cow-wheat | Forest |
| <i>Pedicularis kernerii</i> | Kerner's Lousewort | Alpine turf |
| <i>Pedicularis oederi</i> | Crimson-tipped Lousewort | Alpine turf |
| <i>Pedicularis palustris</i> | Marsh Lousewort | Marshes |
| <i>Pedicularis recutita</i> | Beakless Red Lousewort | Wet pasture. Val Muragl |
| <i>Pedicularis tuberosa</i> | Long-beaked Yellow Lousewort | Acidic mountain turf |
| <i>Pedicularis verticillata</i> | Whorled Lousewort | Mountain turf |
| <i>Rhinanthus alectorolophus</i> | Greater Yellow-rattle | Meadows |
| <i>Rhinanthus minor</i> | Yellow-rattle | Frequent |

| Latin Name | Common Name | Notes |
|--|------------------------------|-----------------------------|
| <i>Verbascum nigrum</i> | Dark Mullein | Lowland |
| <i>Veronica alpina</i> | Alpine Speedwell | Stony pasture |
| <i>Veronica bellidioides</i> | Violet Speedwell | Acidic pasture |
| <i>Veronica chamaedrys</i> | Germander Speedwell | Occasional |
| <i>Veronica fruticans</i> | Rock Speedwell | Mountain rocks |
| <i>Veronica fruticoso</i> | A Rock Speedwell | Rocky moraines. Morteratsch |
| <i>Veronica officinalis</i> | Heath Speedwell | Woods and banks |
| <i>Veronica serpyllifolia</i> | Thyme-leaved Speedwell | Common |
| <i>Veronica urticifolia</i> | Nettle-leaved Speedwell | Woods |
| Tamaricaceae | Tamarisk Family | |
| <i>Myricaria germanica</i> | German Tamarisk | River gravels. Morteratsch |
| Thymelaeaceae | Mezereon Family | |
| <i>Daphne mezereum</i> | Mezereon | Limestone rocks. Local |
| <i>Daphne striata</i> | Striped Daphne | Stony pasture |
| Urticaceae | Nettle Family | |
| <i>Urtica dioica</i> | Common Nettle | Widespread |
| Valerianaceae | Valerian Family | |
| <i>Valeriana montana</i> | Mountain Valerian | Limestone rocks |
| <i>Valeriana officinalis</i> | Common Valerian | Damp meadows |
| <i>Valeriana tripteris</i> | Three-leaved Valerian | Woods |
| Violaceae | Violet Family | |
| <i>Viola biflora</i> | Yellow Wood Violet | Common |
| <i>Viola calcarata</i> | Long-spurred Pansy | Locally common |
| <i>Viola palustris</i> | Marsh Violet | Common |
| <i>Viola riviniana</i> | Common Dog-violet | Common |
| <i>Viola rupestris</i> | Teesdale Violet | Limestone meadows |
| <i>Viola tricolor</i> | Wild Pansy | Waste places |
| LILIIDAE | MONOCOTYLEDONS | |
| Amaryllidaceae | Daffodil Family | |
| <i>Allium schoenoprasum</i> | Chives | Occasional |
| <i>Allium victorialis</i> | Alpine Leek | Local. Alp Grum |
| Asparagaceae | Asparagus Family | |
| <i>Convallaria majalis</i> | Lily-of-the-Valley | Occasional |
| <i>Maianthemum bifolium</i> | May Lily | Forests |
| <i>Paradisea liliastrum</i> | St Bruno's Lily | Local. Pasture |
| <i>Polygonatum odoratum</i> | Angular Solomon's-seal | Woods |
| <i>Polygonatum verticillatum</i> | Whorled Solomon's Seal | Woods and Juniper scrub |
| Colchicaceae | Meadow Saffron Family | |
| <i>Colchicum autumnale</i> | Meadow Saffron | Occasional. Meadows |
| Cyperaceae | Sedge Family | |
| <i>Carex atrata</i> subsp. <i>aterrima</i> | Black Alpine Sedge | Wet pasture. Frequent |
| <i>Carex atrata</i> subsp. <i>atrata</i> | Black Alpine Sedge | Frequent |
| <i>Carex bicolor</i> | Two-coloured Sedge | Glacial alluvium. Val Roseg |

| Latin Name | Common Name | Notes |
|---------------------------------|--------------------------|--------------------------------|
| <i>Carex canescens</i> | White Sedge | Wet heaths |
| <i>Carex capillaris</i> | Hair Sedge | Rocky places |
| <i>Carex curvula</i> | An alpine Sedge | Locally dominant |
| <i>Carex davalliana</i> | Davall's Sedge | Frequent |
| <i>Carex dioica</i> | Dioecious Sedge | Calcareous flushes. Val Minor |
| <i>Carex echinata</i> | Star Sedge | Common |
| <i>Carex ferruginea</i> | Rusty Sedge | Locally dominant on limestone |
| <i>Carex flacca</i> | Glaucous Sedge | Common |
| <i>Carex flava</i> | Large Yellow Sedge | Fen. Corvaglia |
| <i>Carex frigida</i> | Ice Sedge | Moraines. Morteratsch |
| <i>Carex lachenalii</i> | Hare's-foot Sedge | Wet acid rocks |
| <i>Carex leporina</i> | Oval Sedge | Moraines. Morteratsch |
| <i>Carex maritima</i> | Curved Sedge | Alluvium. Val Roseg |
| <i>Carex microglochin</i> | Bristle Sedge | Base-rich flushes |
| <i>Carex montana</i> | Soft-leaved Sedge | Limestone grassland. Val Minor |
| <i>Carex nigra</i> | Black Sedge | Common |
| <i>Carex ornithopoda</i> | Bird's-foot Sedge | Frequent |
| <i>Carex pallescens</i> | Pale Sedge | Frequent |
| <i>Carex panicea</i> | Carnation Sedge | Common |
| <i>Carex parviflora</i> | Small-flowered Sedge | Frequent |
| <i>Carex pauciflora</i> | Few-flowered Sedge | Oligotrophic flushes |
| <i>Carex rostrata</i> | Bottle Sedge | Swamps and lake margins |
| <i>Carex rupestris</i> | Rock Sedge | Limestone scree |
| <i>Carex sempervirens</i> | Evergreen Sedge | Common |
| <i>Eriophorum angustifolium</i> | Common Cottongrass | Common |
| <i>Eriophorum latifolium</i> | Broad-leaved Cottongrass | Wet places |
| <i>Eriophorum scheuchzeri</i> | Scheuchzer's Cottongrass | Lakes and pools |
| <i>Eriophorum vaginatum</i> | Hare's-tail Cottongrass | Occasional |
| <i>Kobresia myosuroides</i> | Naked Sedge | Alpine turf |
| <i>Kobresia simpliciuscula</i> | False-sedge | Base-rich alpine flushes |
| <i>Trichophorum cespitosum</i> | Northern Deergrass | Common |
| Juncaceae | | |
| <i>Juncus alpinoarticulatus</i> | Alpine Rush | Wet mountain pastures |
| <i>Juncus arcticus</i> | Arctic Rush | Alluvium. Rare. Val Roseg |
| <i>Juncus filiformis</i> | Thread Rush | Occasional |
| <i>Juncus jacquinii</i> | Jacquin's Rush | Frequent |
| <i>Juncus trifidus</i> | Three-leaved Rush | Acidic turf |
| <i>Juncus triglumis</i> | Three-flowered Rush | Wet flushes. Val da Fain |
| <i>Luzula alpinopilosa</i> | Alpine Wood-rush | Mountain pasture |
| <i>Luzula lutea</i> | Yellow wood-rush | Mountain pasture |
| <i>Luzula luzuloides</i> | White Wood-rush | Woods |
| <i>Luzula multiflora</i> | Heath Wood-rush | Mountain pasture |
| <i>Luzula nivea</i> | Snowy Wood-rush | Woods |
| <i>Luzula spicata</i> | Spiked Wood-rush | Mountain pasture |
| <i>Luzula sudetica</i> | Sudetic Wood-rush | Occasional |
| <i>Luzula sylvatica</i> | Great Wood-rush | Woods |
| Juncaginaceae | | |
| <i>Triglochin palustris</i> | Marsh Arrowgrass | Local. Wet flushes |
| Liliaceae | | |
| <i>Gagea serotina</i> | Snowdon Lily | Mountain turf |

| Latin Name | Common Name | Notes |
|---|-------------------------------|--------------------------------|
| <i>Lilium bulbiferum</i> subsp. <i>croceum</i> | Orange Lily | Woods and cliffs. Local |
| <i>Lilium martagon</i> | Martagon Lily | Woods |
| <i>Streptopus amplexifolius</i> | Streptopus | Woods. Cavaglia |
| Melanthiaceae | Herb-Paris Family | |
| <i>Veratrum album</i> | White False-helleborine | Mountain pasture |
| Orchidaceae | Orchid Family | |
| <i>Chamorchis alpina</i> | Dwarf Orchid | Scree and mountain turf. Local |
| <i>Coeloglossum viride</i> | Frog Orchid | Frequent |
| <i>Corallorhiza trifida</i> | Coralroot Orchid | Forests |
| <i>Dactylorhiza fuchsii</i> | Common Spotted-orchid | Common |
| <i>Dactylorhiza incarnata</i> subsp. <i>cruenta</i> | Flecked Marsh-orchid | Calcareous flushes. Local |
| <i>Dactylorhiza maculata</i> | Heath Spotted-orchid | Common |
| <i>Dactylorhiza majalis</i> | Broad-leaved Marsh-orchid | Wet pasture. Common |
| <i>Gymnadenia conopsea</i> | Fragrant-orchid | Common |
| <i>Gymnadenia odoratissima</i> | Short-spurred Fragrant-orchid | Locally common |
| <i>Gymnadenia rhellicani</i> | Vanilla Orchid | Locally common |
| <i>Gymnadenia rubra</i> | Red Vanilla Orchid | Locally common |
| <i>Neottia ovata</i> | Twayblade | Occasional. Meadows |
| <i>Orchis mascula</i> | Early-purple Orchid | Locally common |
| <i>Platanthera bifolia</i> | Lesser Butterfly-orchid | Local. Grassland |
| <i>Pseudorchis albida</i> | Small-white Orchid | Common |
| <i>Traunsteinera globosa</i> | Round-headed Orchid | Local. Chantarella |
| Poaceae | Grass Family | |
| <i>Agropyron repens</i> | Common Couch | Waste places |
| <i>Agrostis capillaris</i> | Common Bent | Common |
| <i>Alopecurus pratensis</i> | Meadow Foxtail | Common in lower meadows |
| <i>Anthoxanthum alpinum</i> | Alpine Vernal-grass | Common |
| <i>Anthoxanthum odoratum</i> | Sweet Vernal-grass | Common |
| <i>Arrhenatherum elatius</i> | False Oat-grass | Occasional |
| <i>Brachypodium sylvaticum</i> | False Brome | Forests. Common |
| <i>Briza media</i> | Quaking-grass | Common |
| <i>Bromus hordeaceus</i> | Soft Brome | Lowland |
| <i>Cynosurus cristatus</i> | Crested Dog's-tail | Meadows |
| <i>Dactylis glomerata</i> | Cock's-foot | Waste places |
| <i>Deschampsia cespitosa</i> | Tufted Hair-grass | Wet turf. Common |
| <i>Deschampsia flexuosa</i> | Wavy Hair-grass | Common |
| <i>Festuca ovina</i> | Sheep's-fescue | Common |
| <i>Festuca rubra</i> | Red Fescue | Common |
| <i>Festuca violacea</i> | Violet Fescue | Mountains. Common |
| <i>Holcus lanatus</i> | Yorkshire-fog | Frequent |
| <i>Holcus mollis</i> | Creeping Soft-grass | Frequent |
| <i>Koeleria macrantha</i> | Crested Hair-grass | Frequent |
| <i>Lolium perenne</i> | Perennial Rye-grass | Waste places |
| <i>Melica uniflora</i> | Wood Melick | Frequent |
| <i>Molinia caerulea</i> | Purple Moor-grass | Oligotrophic flushes |
| <i>Nardus stricta</i> | Mat-grass | Common |
| <i>Oreochloa disticha</i> | Two-lined Moor-grass | Alpine Rocks |
| <i>Phleum alpinum</i> | Alpine Timothy | Common |
| <i>Phleum pratense</i> | Timothy | Meadows |
| <i>Poa alpina</i> | Alpine Meadow-grass | Common |

| Latin Name | Common Name | Notes |
|------------------------------|---------------------------------|---------------------|
| <i>Poa annua</i> | Annual Meadow-grass | Common on trackways |
| <i>Poa nemoralis</i> | Wood Meadow-grass | Forests |
| <i>Poa pratensis</i> | Smooth Meadow-grass | Meadows |
| <i>Poa supina</i> | Creeping Meadow-grass | Meadows |
| <i>Poa trivialis</i> | Rough Meadow-grass | Meadows |
| <i>Schedonorus pratensis</i> | Meadow Fescue | Common |
| <i>Sesleria caerulea</i> | Blue Moor-grass | Limestone |
| Tofieldiaceae | Scottish Asphodel Family | |
| <i>Tofieldia calyculata</i> | Tofield's Asphodel | Common |
| <i>Tofieldia pusilla</i> | Scottish Asphodel | Rare. Wet flushes |

Birds (✓=recorded but not counted; H = heard only)

| | Common name | Latin name | July | | | | | | | | |
|----|--------------------------|-----------------------------------|------|---|---|---|---|----|----|----|---|
| | | | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | |
| 1 | Mute Swan | <i>Cygnus olor</i> | ✓ | | | | | | | | |
| 2 | Little Grebe | <i>Tachybaptus ruficollis</i> | ✓ | | | | | | | | |
| 3 | Great Crested Grebe | <i>Podiceps cristatus</i> | ✓ | | | | | | | | |
| 4 | Common Buzzard | <i>Buteo buteo</i> | ✓ | | | | | | | | |
| 5 | Golden Eagle | <i>Aquila chrysaetos</i> | | | ✓ | | | | | | |
| 6 | Common Kestrel | <i>Falco tinnunculus</i> | | | ✓ | | ✓ | | | | |
| 7 | Eurasian Coot | <i>Fulica atra</i> | ✓ | | | | | | ✓ | | |
| 8 | Black-headed Gull | <i>Chroicocephalus ridibundus</i> | ✓ | | | | | | | | |
| 9 | Yellow-legged Gull | <i>Larus michahellis</i> | ✓ | | | | | | | | |
| 10 | Common Swift | <i>Apus apus</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 11 | Great Spotted Woodpecker | <i>Dendrocopos major</i> | | | | | | H | | | |
| 12 | Eurasian Jay | <i>Garrulus glandarius</i> | ✓ | | | | | | | | |
| 13 | Spotted Nutcracker | <i>Nucifraga caryocatactes</i> | ✓ | H | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 14 | Alpine Chough | <i>Pyrrhocorax graculus</i> | | | ✓ | ✓ | ✓ | ✓ | | | |
| 15 | Carrion Crow | <i>Corvus corone</i> | ✓ | ✓ | | ✓ | | | ✓ | ✓ | |
| 16 | Northern Raven | <i>Corvus corax</i> | | | | | | | ✓ | | |
| 17 | Coal Tit | <i>Parus ater</i> | | ✓ | | ✓ | ✓ | ✓ | | | |
| 18 | Eurasian Crag Martin | <i>Ptyonoprogne rupestris</i> | | | | | | ✓ | | | |
| 19 | Barn Swallow | <i>Hirundo rustica</i> | ✓ | | | | | | | | |
| 20 | Common House Martin | <i>Delichon urbica</i> | ✓ | ✓ | | | | | | | |
| 21 | Common Chiffchaff | <i>Phylloscopus collybita</i> | | H | | ✓ | | | ✓ | | |
| 22 | Eurasian Blackcap | <i>Sylvia atricapilla</i> | | H | | | | | | | |
| 23 | Common Firecrest | <i>Regulus ignicapilla</i> | | H | | | | H | ✓ | | |
| 24 | Eurasian Nuthatch | <i>Sitta europaea</i> | | | | ✓ | | | ✓ | | |
| 25 | Common Starling | <i>Sturnus vulgaris</i> | ✓ | | | | | | | | |
| 26 | Common Blackbird | <i>Turdus merula</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 27 | Fieldfare | <i>Turdus pilaris</i> | ✓ | ✓ | ✓ | | | ✓ | ✓ | | |
| 28 | Mistle Thrush | <i>Turdus viscivorus</i> | | ✓ | | | | | ✓ | | |
| 29 | Black Redstart | <i>Phoenicurus ochruros</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 30 | European Stonechat | <i>Saxicola rubicola</i> | | | ✓ | | | | | | |
| 31 | Northern Wheatear | <i>Oenanthe oenanthe</i> | | ✓ | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 32 | White-throated Dipper | <i>Cinclus cinclus</i> | | | | ✓ | | | | | |
| 33 | House Sparrow | <i>Passer domesticus</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 34 | White-winged Snowfinch | <i>Montifringilla nivalis</i> | | | | ✓ | | | ✓ | | |
| 35 | White Wagtail | <i>Motacilla alba</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 36 | Meadow Pipit | <i>Anthus pratensis</i> | | | | | | ✓ | | | |

| | Common name | Latin name | July | | | | | | | |
|----|------------------|-----------------------------|------|---|---|---|---|----|----|----|
| | | | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| 37 | Water Pipit | <i>Anthus spinoletta</i> | | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 38 | Common Chaffinch | <i>Fringilla coelebs</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 39 | European Serin | <i>Serinus serinus</i> | | ✓ | ✓ | ✓ | | ✓ | ✓ | |
| 40 | Citril Finch | <i>Carduelis citrinella</i> | | | | ✓ | | | | |
| 41 | Common Redpoll | <i>Acanthis flammea</i> | | | | ✓ | ✓ | | | |
| 42 | Common Linnet | <i>Linaria cannabina</i> | | ✓ | | | | | | |

Mammals

| | | | | | | | | | | |
|---|---------------|----------------------------|--|--|---|---|---|---|---|--|
| 1 | Red Squirrel | <i>Sciurus vulgaris</i> | | | | ✓ | | | | |
| 2 | Alpine Marmot | <i>Marmota marmota</i> | | | | ✓ | ✓ | | ✓ | |
| 3 | Common Vole | <i>Microtus arvalis</i> | | | | | | | ✓ | |
| 4 | Snow Vole | <i>Microtus nivalis</i> | | | | | | ✓ | | |
| 5 | Roe Deer | <i>Capreolus capreolus</i> | | | ✓ | | | | | |

Amphibians

| | | | | | | | | | | |
|---|-------------|------------------------|--|---|---|---|---|---|--|--|
| 1 | Common Frog | <i>Rana temporaria</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | | |
|---|-------------|------------------------|--|---|---|---|---|---|--|--|

Butterflies

| | | | | | | | | | | |
|----|---------------------------------|-------------------------------------|--|---|---|---|---|---|---|--|
| 1 | Swallowtail | <i>Papilio machaon</i> | | ✓ | | | | | | |
| 2 | Apollo | <i>Parnassius apollo</i> | | | ✓ | | | | | |
| 3 | Clouded Apollo | <i>Parnassius mnemosyne</i> | | ✓ | ✓ | | | | | |
| 4 | Large White | <i>Pieris brassicae</i> | | ✓ | | | | | | |
| 5 | Green-veined White | <i>Artogeia napi</i> | | | | | | ✓ | | |
| 6 | Purple-edged Copper | <i>Palaeochrysophanus hippothoe</i> | | ✓ | | ✓ | | | | |
| 7 | Little Blue | <i>Cupido minimus</i> | | ✓ | ✓ | | ✓ | ✓ | ✓ | |
| 8 | Mazarine Blue | <i>Cyaniris semiargus</i> | | ✓ | | | ✓ | ✓ | ✓ | |
| 9 | Common Blue | <i>Polyommatus icarus</i> | | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 10 | Painted Lady | <i>Vanessa cardui</i> | | | | ✓ | | ✓ | ✓ | |
| 11 | Small Tortoiseshell | <i>Aglais urticae</i> | | ✓ | | | | | | |
| 12 | Small Pearl-bordered Fritillary | <i>Boloria selene</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 13 | Alpine Grayling | <i>Oeneis glacialis</i> | | | ✓ | ✓ | | | | |
| 14 | De Pruner's Ringlet | <i>Erebia triaria</i> | | | | ✓ | | | | |
| 15 | Piedmont Ringlet | <i>Erebia meolans</i> | | ✓ | ✓ | | ✓ | ✓ | | |
| 16 | Ringlet | <i>Aphantopus hyperantus</i> | | ✓ | | | | | | |
| 17 | Grizzled Skipper | <i>Pyrgus malvae</i> | | ✓ | | | | | | |
| 18 | Oberthur's Grizzled Skipper | <i>Pyrgus armoricanus</i> | | | | | | ✓ | | |
| 19 | Dingy Skipper | <i>Erynnis tages</i> | | ✓ | | | | | | |

Moths

| | | | | | | | | | | |
|---|-----------------|-----------------------------|--|---|---|---|---|---|---|--|
| 1 | Six-spot Burnet | <i>Zygaena filipendulae</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 2 | Chimney Sweeper | <i>Odezia atrata</i> | | ✓ | | | | | | |