

To Professor Lindley from the Author

NARCISSINEÄRUM MONOGRAPHIA.

AUTORE

A. H. HAWORTH,

Soc. Linn. Lond.—Soc. Horticult. Lond.—Soc. Caesar. Nat. Curios. Mosc.—necnon
Soc. Reg. Horticult. Belgic. Socius. &c. &c.

AUTHOR OF LEPIDOPTERA BRITANNICA; SYNOPSIS PLANTARUM SUCCULENTA-
RUM; AND MANY OTHER WORKS.

Multum adhuc restat, multumque restabit.

THE SECOND EDITION,
WITH ADDITIONS AND IMPROVEMENTS.

LONDON :
JAMES RIDGWAY, PICCADILLY.

M. DCCC. XXXI.

" ... Sulphur Phoenix (or Crown) was omitted casually. I was not aware of it.

The paper was printed in a periodical in a great hurry and with many typographic errors. I had 25 copies for the manuscript (now all given to British and foreign Authors) save one for myself (and 2 proofs full of corrections). Before these said 25 were worked, I revised the press and advised Ridgway to cancel the others and strike off new from the revised types. But I know not whether he did so.

- From a letter by Haworth to
the Rev. H.T. Ellacombe, now
in the Rowley Library.

P R E F A C E.

THE present MONOGRAPH on the beautiful family of NARCISSINEAN Plants, the finest early group in our Gardens, was compiled from the author's manuscripts, at a very short notice, to be ready for publication at the end of the last volume of his friend Sweet's *British Flower Garden*; and the consequence of this haste was, typographical errors, a few other inaccuracies, and insufficient descriptions; which have been corrected in the present edition.

Notwithstanding the short time which has elapsed since the first publication of the MONOGRAPH, many material new varieties (four of which may prove to be species) of these charming plants, have been added to their showy and fragrant affinities; each hereunder inserted in its most appropriate place.

An idea may, perhaps, be entertained by some, who are insufficiently acquainted with AMARILLIDEOUS plants, of which NARCISSINEÆ form the fifth *sub-ordo*, that many of the articles hereunder enumerated as *species*, by the author, are not primitive and originally distinct species, but seminal variations only, originating in the Flowery Gardens of the Dutch. But, both the author, and his acute friend, Mr. Sweet, (whose knowledge of living plants is, perhaps, unrivalled,) have found good reason to determine otherwise; not only on account of the characteristics given, but because the *species* themselves were actually fostered in the gardens of our forefathers, one or two hundred years ago; and they were managed too, with a degree of skill and success that has never since been equalled, particularly in Britain.

And this very early period was, before the Gardens of Holland had either raised, or acquired from other countries, many seedling varieties of NARCISSINEAN plants; as we satisfactorily learn from Miller, who says, in the last edition of the Gardeners Dictionary, printed in his time, in the year 1768, under the article *Narcissus Tazetta*:—"The printed catalogues, by Dutch Florists, contain more than *thirty varieties*;" and he enumerates a great many of them. But of Miller more hereunder.

Moreover, the various countries from whence the plants were brought from their native wilds, are likewise carefully stated by some or other of the elder Botanists, (however, since neglected,) as Clusius, Tournefort, and Barrellier; Rudbeck, Besler, and Swertius; by Lobel, Chabréus, Dodoneus, and Garidel; and others among foreigners; and in our own country, by Gerrard, Johnson, and the faithful Parkinson; by the great Ray; and Morison; and by Trew, in his fine and beautiful works. By the celebrated Madam Merian, whose journey to, and long sojourn at Surinam, in quest of natural knowledge, cannot be forgotten; and by the ingenious author of *Theatrum Floræ*, abounding in *Narcissinean* Plants: and last, but not least, by a name nearer to our

own days, and which the Botanists of Chelsea rejoice in repeating that of *Phillip Miller*, whom foreigners have emphatically styled "the Prince of Gardeners;"

"Hortulanorum Princeps;"

and of whose Dictionary the great Linnæus said,

"Non erit Lexicon Hortulanorum, sed Botanicorum."

For the culture and management of these plants, I refer the reader to that famous Dictionary; but, as the book is both high priced and not in every ones possession, it may likewise be desirable to detail a few particulars on this head, in the present place.

NARCISSINÆ have all annually quiescent, perennial, bulbous roots, of the tunicated kind, of an ovato-pyramidal form, covered externally with brownish thin membranous scariose coats, and ~~are~~ from the size of a large Spanish nut or filbert, in the Genera *Corbularia* and *Ganymedes*, to that of the clenched fist of a man, in *Hermione cupularis*. A few new inner tunics are annually formed in the centre of the bulb, one by the base of every leaf of the year, which is completed on the annual quiescence of the bulb, when the leaf decays down to its base; and a few of the extreme and living tunics are annually absorbed and exsiccated by the regular growth of the plant, into exterior membranous ones. These last, though dead, endure about the bulb for a long time, and help to protect it from frosts; therefore I never willfully destroy those coats, but re-plant and close them round it.

The roots require transplantation once in three or four years, to take off their offsets, and to encrease them; but if they are transplanted annually, as in Holland, they form the largest, roundest, and finest roots. This may be done as soon as their leaves decay in summer, and not long after, lest they should form new fibres; after which, transplanting would be very injurious; and if they are moved in winter or spring, they are rarely able to flower for two or more years afterwards. After taking them up, the flowering sized roots may be kept out of the ground dry, like Tulips, a few weeks, or more, if desirable; but they are best planted late in September, except the very early species, which should be planted late in October, as by so retarding them, they more easily escape from frost, pushing later out of the ground. All offsets, and weak bulbs, ought never be kept out at all, because, drying shrivels and greatly weakens them.

The soil these plants thrive best in, is a rich or loamy one, a foot and a half beneath their bulbs, which should be six inches apart, filling up with the same, or any soil that is light, three or four inches over the tops of the larger roots, and two or three over small ones; but there should be no stones or gravel, which would cut the swelling bulbs, and prevent their due and equal enlargement. All barren sandy soils

are to be avoided ; the flowers in such, especially double white ones, are never fine, and rarely perfect. The Genera *Ganymedes*, and its nearest allies, and *Corbularia*, should be planted in pure loam, and covered over their tips three inches, to protect them from cold.

All *NARCISSINÆ*, (except those from cold native situations,) require to be planted in beds, raised two or three inches above the walks, in an unshaded situation, or near a South or West aspected wall, and to be covered, in frosty weather, with a single mat, or a light protection of dry straw or litter of any kind, but not near so thick as is used for early radishes ; for that would blanch their early leaves ; and it should be removed occasionally, in open weather, for the same reason. Common empty garden pots placed over single plants, or patches of plants, equally answer the purposes against frost, and should be removed in the same manner, and for the same reasons, stopping up the holes at the bottom of the pots ; or single large roots, covered with earth over their tips, four or five inches, often escape very well in dry situations, without any other protection.

These plants are nearly all natives of Europe, or Asia Minor, a few excepted, as stated by authors, under the account of the species respectively, which are found near Gibraltar, Algiers, Morocco, and the most northern coast of Africa ; and one solitary specimen is extant in the Herbarium of the late Sir J. E. Smith, brought from Nipaul by the late Dr. Hamilton : but my excellent friend Dr. Wallich, (the best of all judges of Indian plants,) joins with me in supposing it not to be an Indian plant, but to have been sent there as a lovely flagrant flower, through the way of Persia. This plant is very closely allied to, if not the same as *Hermione tazetta*, of Linn.

All the *NARCISSINÆ* hereunder enumerated, are at this present time actually growing in the noble Gardens of the Horticultural Society at Chiswick ; or those of Mr. Sweet, and his friends ; except such only as are marked with an * in the following pages ; and any so marked, the writer of this *Monograph* would be glad to receive bulbs of, from any one, addressed to him at the Publisher's, Mr. RIDGWAY, Bookseller, Piccadilly, London : and for such communications the author would cheerfully and thankfully return any of the species or varieties he has described, which are duplicate with him, and which may be requested.

Before closing this Preface, it may not be unuseful to observe that vast numbers of chiefly Dutch, imported bulbs, of the Genera *Hermione* and *Jonquilla*, with a few of *Phylloigne*, *Corbularia*, and *Ajax*, are annually and easily forced into early and beautiful bloom, from the end of January, to the end of April, by Nurserymen, Gardeners, and others, for the purpose of sale, both at home, and in the vegetable Markets of London, and elsewhere ; where they are in great demand, and sell readily, and very profitably, before their periods of flowering in the open ground.

The process of forcing these plants, is effected in the following manner :—

The largest and roundest, and, consequently, the heaviest and hardest bulbs, of their respective sorts, are carefully selected for this purpose ; and one, two, or three of them are planted in rich loamy earth, in rather large pots, from the size of that called 32 by the Gardeners, to any size you please ; but the larger the pot, the stronger the plant. The bulbs should be only just covered over with earth, that their fibres may have more room. They are planted at various times, or at short intervals, to secure a regular succession ; and those which are planted the earliest, of course, bloom the first, if of the same sort.

As soon as planted, they should be plunged in the earth, or old tan of some pit, at the front of a hot or greenhouse, or in garden frames ; or may be placed on shelves in such houses themselves ; but they should always be near the glass. They also succeed pretty well in the sunny windows of a dwelling house ; and may be even flowered, but not vigorously, in common bulb-glasses, filled with water, occasionally changed in the usual way ; the darker the glass the better ; as all exposure to light is injurious and unnatural to the white fibres of the roots, and prevents their flowering again for at least two seasons of the best subsequent care.

After plunging and covering the pots as abovementioned, the plants will particularly require to have air and water duly and regularly given to them, at all proper and favourable opportunities ; and the more close they are kept, the weaker will be their beautiful blooming, which will occur from the end of January, in the early kinds, to the middle or end of April, in the late ones ; and there is one fine white-flowered species of *Hermione*, which blooms, in its wild state, near Naples, in November and December, which is figured by Tenore, in his *Flora Neapolitana*, and named *Narcissus unicolor*. But this plant, although it has been, is not, at present, in the British gardens ; or, at least, the writer has never seen it.

In March or April, in the neighbourhood of London, roots that have been forced, and whose flowers have faded, are usually turned carefully out of their pots, into some cool border of rich soil, near a wall or fence, and covered with earth 3 or 4 inches over their bulbs, and duly watered until they recover ; which (unless forced in water) they will sometimes do, so as to flower again the following season.

It must not be forgotten, that it is as necessary to defend these forced plants from weeds, and from worms—from slugs, and from snails ; as it is requisite to protect them from cold, by straw, or mats, &c. during every frost, or loss and disappointment will ensue ; for blighted will be the precocious beauties, and in vain the gardener's every care :

“ And DAFFODILLIES fill their cups with tears.”

Chelsea, November, 1831.

NARCISSINEÆ.

Nat. Ordo. AMARYLLIDÆ. *Brown prodr.* 1. 296. *Subordo* 5.
Hexandria Monogynea Linn. *Spatha monophylla* 1, usque ad 20 flora.
Corolla tubulosa limbo sexpartito hexapetaloida coronata. *Ovarium*
subturbanatum seu oblongum vel cylindricum triloculare, trivalve; valvis
biseptiferis. Semina numerosa, in 2-4-ordinibus inserta.

- * Genitalia adscendentia.
- I. CORBULARIA. *Narc.* *Revis.* 111. *Corollæ laciniis* exiguis coronâ truncatâ sèpissimè brevioribus. *Ovarium* subturbanatum cylindricum, semina triangularia numerosa bifariam inserta. *Folia* filiformia canaliculata viridia.
- † Albifloræ.
- * 1 *cantabrica.* C. (the lesser white) candida: corona crenata, stylo inclusu. *Clus. Hist.* 166. f. 2.
- * 2 *albicans.* C. (the larger white) corona sulphureo-albicans integerrima, stylo exerto. *Park. Par.* 106. f. 4.
- † Flavifloræ.
- 3 *Bulbocodium.* C. (the small entire-crowned) parva: corona integra, stylo inclusu. *Linn. sp. pl.* 3. 417. atque ejus *Herbarii*.
- 4 *tenuifolia.* C. (slender leaved) parva: corona 6-lobata, stylo exerto. *Narc. Revis.* p. 121. *Swt. Br. Fl. Gard.* t. 114. Martio floret.
- * 5 *aurea.* C. (the shorter crowned) corona repando-crenata laciniis breviore stylo exerto. *Schult. syst.* 7. 985. *Moris.* s. 4. t. 9. f. 15. *H. Eyst. ord.* 3. fol. 13. f. 4.
- 6 *obesa.* C. (the inflated) corona inflata integrâ, stylo exerto, foliis semper humifusis. *Narc. Revis.* p. 122. *exclus.* synon. Martio floret.
- 7 *lobulata.* C. (the lesser hoop-petticoat) corona undulato-lobata stylo fere semper inclusu. *Narc. Revis.* 122. *Park. Par.* 107. f. 7. *Clus. Hist.* 166. f. 1. Apr. floret.
- 8 *conspicua.* C. (the showy) magna: indora: corona lèvissimè repandulata, stylo fere semper exerto, foliis sub anthesin erectis. *Park. Par.* 107. f. 6. Floret f. Ap.
- 9 *serotina.* C. (the large, late) magna: corona integra, stylo inclusu. Pseud. *Narc. serotinus*, &c. *Park. Par.* 106. f. 8. N. *turgidus. Salisb. Prod.* p. 222. N. *Bulbocodium. Bot. Mag.* nec *Linn.* neque ejus *Herbarii*, med. Maii floret.
- * 10. *Gigas* C. (the giant) corollæ laciniis lanceolato-linearibus coronam maximam campanularem plusquam 2-unciam aequantibus, stylo alte porrecto. *Pseudo Narcissus pyrenæus* flore sublato amplo calyce juncifolius. *Theatr. flor.* t. 21. Fig. inferior media. Cæteris 2-3-plo major.
- ** Genitalia recta.
- II. AJAX. *Narc. Revis.* 111. *Corona lobata*, crenata, serrata, fere vel plus longitudine laciniarum sèpissimè imbricatarum corollæ. *Ovula* in valvis quadrifariam inserta. *Folia* ensiformia vel lorata obtusa sèpe infernè carinata, et supra, ab inflexis lateribus latè canaliculata.
- * *Minores*, flaviflores, tubo longiusculo. Corolla laciniis semiexpansis.
- 1 *minimus.* A. (the least) corollæ laciniis imbricatis, corona patula 6-lobata valde crispa, ante anthesin ipsum humum nutantim tangente. *Obs.* foliis perglaucis trientalibus humi subjacentibus. *Pseudo-Narcissus minimus. Park. Par.* 107. 3. *Malu.* *Narcissus minor. Bot. Mag.* 6. spatha supra germen ex errore posita!
- 2 *minor.* A. (the lesser) corollæ laciniis imbricatis, corona patula 6-lobata valde crispa, scapo erectiore. *Obs.* foliis patulo adscendentibus perglaucis 6 uncialibus.
- A. minor. ♂. *Narc. Revis.* 112. N. minor. *Linn. sp. pl.* p. 415. et ejus *Herbarii*. *Ps. Narcis. minus. Park. Par.* 107. f. 2.
- ♂. foliis longioribus erectioribus quam a. *Pseudo-Narcissus mediuss. Park. Par.* 105.
- γ. *angustus*, foliis erectis longè angustioribus quam in prioribus. Forte propria species: Omnes var. med. Martii florent excepta forte hæc, quæ non vidi florens.
- 3 *pumilus.* A. (the small) corollæ laciniis obcuneatis distinctis, corona precedentium. *Ajax cuneifolius. Narc. Revis.* 113. *Narcis. pumilus. Salisb. Prod.* p. 220.

- † Reflexiflores, corollæ laciniis reflexis.
- * 4 cyclamineus. A. (small reflexing-flowered). N. hispanicus minor luteus amplio calyce foliis reflexis. *Theat. Flor.* t. 220.
- ** Pallidiflores : corollæ mediocris seu majuscule, tubo abbreviato, laciniis quam prioribus magis expansis albicantibus seu albis, corona alte lutea.
- 5 nanus. A. (the dwarf sulphur) corollæ laciniis subsemierectis ovatis sulphureis tubo sesquifloribus, corona per lutea subrecta, ore lobatim crenato plicatulo criso. N. minor δ. conspicuus. *Narc. Revis.* p. 112.
Obs. *Folia* glauccissima semunciam lata. *Flos* sesquifloralis fere inodorus aperiens f. Feb. seu i. Mart.
 β . Corolla laciniis horizontalibus, foliis duplo angustioribus minus glaucis. Amicus Dom. Penny communicavit. Floret cum α . γ. *Jl. pl.* apud amicum Rev. Dom. Ellacom. Non vidi.
- 6 breviflos. A. (short whitish petalled yellow crown) corollæ laciniis albescensibus, tube coronæ luteæ cylindricæ rectissimæ crenato-incisæ duplo breviore, foliis glaucis. Ajax loriifolius, β . *Narc. Revis.* 119.
- 7 loriifolius. A. (whitish petalled yellow crown) corona lutea, corollæ laciniis albis s. albicantibus, scapo teretiusculo ancipiiti, foliis vix glaucescentibus. A. loriifolius α . *Narc. Revis.* 119. N. bicolor, *Bot. mag.* 1187, nec Linn., neque ejus Herbarii. *Obs.* Elatiior quam priore foliis magis viridis corolla longiore.
 β anceps. Scapo ancipiiti, magis compresso. A. loriifolius γ. anceps. *Narc. Revis. l.c.*
- 8 bicolor. A. (great white petalled yellow crown) corollæ tubo longitudine pericarpii, laciniis niveis incurvo horizontalibus subtortis tubo triplo longioribus ovato-lanceolatis corona lutea basi ventricosa brevissime 6-lobata repandocrenata, vix plicata. *Narc. Revis.* 119. *Narcissus bicolor.* Linn. *sp. pl.* 1. 415. N. tubæflorus. Salisb. *Prod.* p. 221.
- *** Albitflores, corollæ mediocribus seu majusculis, primo sulphureis demum albis, corona longissima.
- 9 albicans. A. (the greatest Spanish white) corollæ laciniis ovato-lanceolatis planiusculis albidis : corona crenata sulphurascente. Pseudo-Narcissus hispanicus maximus albidus. *Park. Par.* p. 109. t. 101. f. 1. Vidi florentem, apud Dom. Young's Epsomæ. Apr. 1831.
- 10 tortuosus. A. (the great tortuous white) corollæ laciniis prætortuosis sulphureo-albicantibus, corona citrina crenata demum alba longe brevioribus. N. tortuosus. *Nob. in Misc. Nut.* p. 179. N. moschatus. *Bot. Mag.* t. 924, nec Linn., neque ejus Herbarii. N. longiflorus. *Salisb. in Hort. Tr.* 1. p. 349.
- 11 cernuus. A. (the drooping creamy white) corollæ cernuæ denique albæ laciniis prætortuosis sulphureis, corona citrina ore sexlobate plicatulato, lobis rotundatis recurvantibus margine aliquantulum repandulo. N. cernuus. *Roth. Cat. Bot. fasc.* 1. p. 43. *Sweet br. fl. gar.* v. 2. s. 2. t. 101. Pseudo-N. hispanicus flore albo minor. *Park. Par.* 100. β . flore elegantissime pleno. *Nob. in Phil. Mag.* Jun. 1830. p. 134. *Sweet br. fl. gar.* v. 2. s. 2. t. 101. Vidi in horto Whitley, Bræmes, & Milne, apud Fulham, Aprili florentem.
- 12 moschatus. A. (the lesser Spanish white) corollæ laciniis tortis primo sulphurascentibus demum albis, coronæ longitudine. N. moschatus. Linn. *sp. pl.* 1. p. 415. et ejus Herbarii. N. moschatus δ. *Bot. Mag.* t. 1800. N. candidissimus. *Redout. Lil.* t. 188. Ajax patulus. *Salisb. in Hort. Tr.* 1. p. 348. Pseudo N. hispanicus fl. albo minor. *Park. Par.* p. 100. t. 101. f. 4.
- **** Lutei, floribus plerumque luteis, saepè sulphureis, rarissimè stramineo-albicantibus, corona plus minusve saepè saturatiore.
- † *Serricornæ*, statura mediocri, coronæ minus profundè serrata seu crenata quam in sequentibus.
- 13 Pseudo Narcissus. A. (the garland) scapo ancipiote teretiusculo recto striato ; corollæ laciniis pallidè sulphureis, exterioribus lanceolato-ellipticis tubum vix superantibus : corona lutea ore recto crenatum serrato longitudine laciniarum. Linn. *spec. I.* p. 414. Ajax festalis. *Salisb. in Hort. Tr.* 5. 243. *Narc. Pseudo-Narcissus. Eng. Bot.* t. 17.
- α . pallidus, corollæ laciniæ pallide luteæ.
 β . albus. Corollæ laciniis albis.
 γ . luteus. Corollæ laciniis pallidè luteis.
 ζ . plenus, laciniis pallidis. *Park. Par.* t. 101. f. 8. *Parthenocallis* (?)
 ε . plenissimus. John Tradescant's great Rose Daffodill. *Park. Par.* t. 101. f. 6.
 ζ . ryticarpus. Capsulis rugosis. An propria species ? Habitat in anglia.

14 *serratus*. A. (the serrate) scapo striato subcompresso; corollæ laciniis planis sulphureis, exterioribus ovato-acuminatis: coronam luteam rectam plicatam alte serratam non attingentibus.
N. serratus. Nob. in *Misc. Nat.* p. 179.
 $\beta?$ *suavis*.

γ . *radians*. Corona perlutea intus crebre rugulosa, ore patulo plicato subregulariter valde radianter serrato, laciniis (corollæ) semi-expansis ovato-oblongis sulphureis, tubo duplo longioribus.

From the Epsom Nursery, March, 1831.

δ . *præcox*. Corollæ laciniis saturate sulphureis planis plusquam semiexpansis ovato-acuminatis tubo 5 lineas superantibus, coronæ patulæ luteæ grande serrata subæquantibus.

Forte omnium præcocius floret in Hort. nobili, Hort. Londini Soc.

• 15 *nobilis*. A. (the noble serrate) scapo teretum ancipiæ alte striato, laciniis corollæ patentissimis tortis ellipticis luteis, nectario perluteo ore patulo profundi sime serrato brevioribus. *Nob. in Narc. Rev.* p. 115.

Narcissus pseudo Narcissus. *Red. Lill.* t. 158.

†† **LOBATO-CORONÆ**, mediocres lobis integroribus.

16 *lobularis*. A. (Tenby 6-lobed Daffodill) corollæ laciniis luteis tubo obconico exactè duplo longioribus: corona perlutea patula sexlobata (lobis integris) laciniis tres lineas superante. *Nob. in Phil. Mag. May*, 1830, p. 131.

This grows wild near Tenby in Pembrokeshire, which by mistake was written Truby in Derbyshire, in the place above cited.

$\beta?$ *amplicorona*.

Obs. In Chelsea Garden a wild bulb from Devonshire, with double flowers, produced offsets bearing for the three last years permanently single ones. Was its source genuine, it would form a good species with the following essential characters.

Corollæ laciniis ovato-acutis valde imbricantibus luteis, tubo sesquiongioribus; corona amplissima perlutea, basi ad apicem per dilatata sub-20 lineas expansa crenatim incisa subplicata crispa, intus rugosiuscula, laciniis superante.

γ . *plenus*. (The common deep yellow double Daffodill.) β . *Gawler's* *

δ . *scoticus*. Scapo compresso ancipiæ torta alte striato, corollæ laciniis lanceolato-ellipticis sulphureis tubum longe superantibus, corona lutea, ore magno expanso in seiso-crenato ruguloso.

Forte propria species. Exemplarium unum solum examinavi.

17 *rugilobus*. A. (the early rough lobed) corollæ laciniis ovato-acuminatis luteis, tubo tertia parte fere longioribus; corona perlutea altè sexlobata, lobis patulis plicatim criso-rugosis.

Amicus Dom. Johannes Denson, Junior; communicavit. Floret i, Martio prope Londinum.

18 *cambricus*. A. (the recurring-lobed) corollæ laciniis luteis ovato-acuminatis tubo tertia parte longioribus, corona perlutea alte sexlobata, lobis patulis vix rugosiusculis plicato-crenatis recurvantibus.

Communicavit amicus Dom. Penny, fine Martii, 1831. *Habitat* in Cambria. Nova species, vel fortasse precedentis varietas. Florem unum marcescentem solum vidi.

19 *obcallaris*. A. (short-tubed spread crown) corollæ laciniis tubo $\frac{1}{2}$ longioribus rectis ovatis, interioribus multo angustioribus valde imbricatis; corona infundibuliformi basi cylindracea 6-fida, repando-dentata superne plicata.

A. obvarialis. *Narc. Reris.* 120.

N. obvarialis. *Salsib. Prod.* p. 221.

N. major. γ . *Gawler in Bot. mag.* t. 1381, fig. ima sine foliis.

†† **INSCISIOB. I.** Majores, et maxi-
mi, *coronæ* lobis profundis propriis sex,
semper plus minus irregulariter laci-
niato-serratis, s. crenatis.

* 20 *spurius*. A. (deep yellow) scapo lœvi compresso, ore coronæ perluteæ alte 6-fido patulo, laciniis corollæ semie-rectis lanceolatis luteis. *Narcis. Reris.* p. 115. Floret Aprili. *Narc. major* luteus calyx prælongo alter. *Rudb. Elys.* 729.

21 *Telamonius*. A. (the long-flowered) corona ore 6-fido expanso alte irregulariter crenato. *Narciss. Reris.* p. 115. Floret Aprili. In solo fertili, flos est fere trientalis longus.

β . *grandifolius*. (the great double.)

Park. Par. t. 101. f. 7. *Telamonius plenus*
 γ . *pleno pallidus acuminatus* (the acu-
ter pointed paler double.) In the Fulham Nursery.

22 *propinquus*. A. (the great gash-crowned) corollæ perluteæ laciniis semierectis tortis incurvo-patulis, corona laciniarum longitudine, ore subpatente pro-
fundissime irregulariter-dentato-cre-
nato; foliis planis perglaucis.

N. propinquus. *Salsib. Prod.* p. 221.

Ajax laciniularis ejusd. in *Hort. trans.* 1. p. 344. *Ajax propinquus*. β . *Narc. Reris.* p. 116. *N. major*. β . *Gawler in Bot. mag.* t. 1301, figura superior.

23 *maximus*. A. (the intermediate large yellow) corollæ perluteæ laciniis patulis

* in view of *Telamonius* *Nervosus* being listed separately (21).

- tortulis, corona ampla, 6-lobata lobis patentibus magnicrenatis.
- Ajax propinquus. a. *Narc. Retis*. p. 116.
- N. maximus. *Hortorum Batavorum*.
- N. major totus lutes calyce prælongo.
- Rudb. *Elys.* t. 71. f. 19.
- * 24 major. A. (the great) corollæ perluteæ, laciniis patulis, corona patentissima amplissima, lobis omnium maximis recurvis subundato-crenatis; foliis glaucissimis spiralibus.
- Narcissus major*. *Bot. Mag.* t. 51.
- III. OILEUS. (*Poetarum Ajax minor*.) *Spatha uniflora*. *Corolla* luteæ laciniis patentibus. *Corona* longior, vel longitudine laciniarum integra seu integerima truncata. *Stylo* inclusio. *Genitalia* non vidi. *Statura* et *folia* *Ajacium minorum*, discrepante corona abque fimbria serratur.
- * 1 *abscissus*. O. (the greater clipt-trunk) *Pseudo-Narcissus* tubo quasi absciso. *Park. Par.* t. 107. f. 1. *Narc. flavus* tubo rotundo majore. Rudb. *Elys.* t. 69. f. 4. *Suertius Florileg.* t. 22. f. 4. β flore pleno. Soleil Brillante. *Trew flor.* *imag.* t. 124. *Narcissus IV*.
- * 2 *hexangularis*. O. (the 6-angled clipt-trunk) coronae integræ hexangularis, angulis in totidem mucronis obsoletis fimbriis.
- Narcissus subflavus* tubo sexangulari. Rudb. *Elys.* 60. f. 3. *Suertius Florileg.* t. 22. f. 1. *Pseudo-Narcissus* tubo sexangulari. *Park. Par.* t. 101. f. 5.
- * 3 *minor*. O. (the lesser clipt-trunk) *Narcissus flavus* tubo rotundo minor. Rudb. *Elys.* t. 68. f. 4.
- Obs.* Penultima simillima, at triplo minor.
- * 4 *albus*. O. (the white) *Narcissus albus*, calyce seu tubo non fimbriato. Rudb. *Elys.* t. 70. f. 7.—*Obs.* Facies *Ajacis minoris*, stylo exerto.
- * 5 *pumilus*. O. (the small clipt-trunk) *Narcissus hispanicus* tubo non fimbriato. Rudb. *Elys.* t. 72. f. 15. *Suertius Florileg.* t. 21. f. 9.
- Obs.* *Ajacis minimi* supra, valde similis, at corona, truncata.
- Obs.* I have not seen any of these species, and insert them with a view to excite inquiry after them. Q. capax. Salisb. may be among them.
- IV. ASSARACUS. (*Ganymedis frater*.) *Spatha biflora*. *Corolla* ochroleuca nutantis laciniis subsemireclexis. *Corona* subundata, poculiformis, laciniis æquans potius superans. *Stamina* clusa, *tria* ore tubi, alia *tria*, fere orem coronæ attingentes, ut in figuris infra citatis. *Folia* lorata plana et statura ut in *Ajacium minorum*, sed *Ganymedis facie*.
- * 1 *Capax*. A.—*Narcissus Calathinus*. *Re-*
- dout.* *Lill.* t. 177. nec *Linn.* *Queltia capax*. *Salisb.* in *Hort. Tr.* 1. 353. N. capax. *Schult. syst. reg.* 7. 950.
- β . *plenus*. *Nob.* in *Phil. Mag.* Jun. 1830. p. 134. *Park. Parad.* t. 107. f. 4. *Flores* flavi dicuntur.
- * 2 *reflexus*. A. *Corolla* pallide flava.
- N. *reflexus*. *Schult. syst.* 7. 952. *Narcissus calathinus*. *Red. Lill.* t. 410. nec *Linn.* *Linnei* N. *calathinus* vide infra sub *Phlogyne*.
- V. ILLUS. (*Ganymedis Frater*) *Spatha* 2-4-flora. *Corolla* absolute cernua seu pendula laciniis cyclaminum more reflexis, usque ad pedunculos, alba sive ochroleuca. *Corona* poculiformis integræ laciniis subduplicata brevior. *Stamina* inclusa. *Genitalia* recta gracilia valde inæqualia. *Filamenta tria* tubo breviora adnata fere ad *antheras* erecta: alia *tria* e tubo progradientia usque ad 3 lineas, libera ultra tubum at corona breviora. *Stylus* exsertus 2-3 lineas vel plus. *Folia* juncoidea valde canaliculata.
- * 1 *cernuus*. I. (the pale yellow) corollæ ochroleucae laciniis planis oris albis, corona sesquilonigioribus: scapo teretiæculo. *Narcissus cernuus*. *Salisb. prodr.* p. 223. *Ganymedes cernuus*. *Salisb. hort. trans. v. 1. p.* 352. *Narcissus triandrus*. *Bot. Mag.* 48. non *L. Ganymedes* ochroleucus. *Revis. pl. succ.* 206. N. *coronatus*. *Schult. syst. reg.* 7. p. 986. *Suert. Flor.* t. 65. f. 7.
- 2 *triandrus*. I. (the white twisting flowered) corollæ albæ laciniis tortis corona duplo longioribus, scapo compresso. *Ganymedes albus*. *Revis. pl. succ.* 208. *Narcissus triandrus*. *Linn. sp. pl.* 2. p. 416. N. *Coorniei*. *Schult. syst. reg.* 7. 986. N. *coronatus*. *ibid.* *Suert. Flor.* t. 29. f. 4.
- VI. GANYMEDES. *Corolla* laciniis semireflexis, nutantibus, vix cernuis, *corona* cyathiformi, laciniis 3-4-plo breviore, *stylus* subinde inclusus. *Semina* ovata placentis duabus longitudinalibus linearibus affixa, duobus ordinibus in singula placenta, cetera ut in genere precedente.
- 1 *pulchellus*. G. (the pretty white cupped) 3-7-flora: corollæ reflexæ laciniis luteis, corona alba, stylo inclusio semper. G. *pulchellus*. *Salisb.* in *Hort. Trans.* 1. 354. *Sweet. Br. Fl. Gard. Series* 2. t. 99.
- N. *triandrus* v. *luteus*. *Bot. Mag.* 1262.
- 2 *concolor*. G. (the self-coloured) corolla totâ lutea nutante, stylo semper alte exerto. *Narc. Revis.* 134. 4. *Sweet's Br. Fl. Gard. Series* 2. t. 113.
- 3 *striatulus*. G. (the faint striped) subuniflorus: laciniis corollæ luteis tortuosis

- non imbricantibus, corona sulphureo albidente striatula valde repanda triplo longioribus, stylo excluso. *Narciss. Revis.* 132.
- β *parvicorona*: humilior, biflorus; omni parte temiore, stigmate coronam minorem aequante. From the Horticultural Society's Garden.
- 4 *nutans*. G. (the yellow cupped) sub3-florus: corollæ laciniis pallide luteis corona saturiore, stylo valde exerto. *Narciss. Revis.* 131.
- Narcissus trilobus*, Linnæi. *Gawler in Bot. Mag.* 945.
- * 5 *albus*. G. (the white) corollæ albæ laciniis semireflexis corona integra subcupulari subduplo longioribus.
Narcissus totus albus reflexus. Moris. s. 4. t. 9. f. 13.
- N. juncifolius albus*. Park. *Parad.* t. 93. f. 1.
- β *major*. N. *juncifol. alb.* magno calyce. Park. *Parad.* t. 92.
- VII. DIOMEDES. *Corollæ tubus clavatum cylindricus validus, corona poculiformi magna laciniis plusquam semiæquante. Genitalia recta. Filamenta subæqualia tubo semiplusve deorsum connata*. Nob. in *Phil. Mag.* Nov. 1823.
- 1 *minor*. D. (the small white petalled) filamentis tribus apice solum liberis, stylo corona brevirore. *Floret Aprili*. N. Macleayi. *Bot. Mag.* 761.
- 2 *major*. D. (the large white petalled) corollæ laciniis oris reflexis filamentis plusquam semiliberis, stylo coronam aequante. Nob. in *Phil. Mag.* l. c.—N. Sabini. *Bot. Reg.* 762.
- N. *albus tubo luteo*. Rubd. *Elys.* t. 69. f. 5.
- * 3 *Parkinsoni*. D. (shorter cupped) corollæ laciniis, corona plus duplo longioribus. *Narcissus albus oblongo calyce*. Park. *Parad.* t. 71. f. 7. fig. 13.
- VIII. TROS. (Ganymedius Pater.) *Spatha* 1-2 flora. *Corolla e sulphureo albidente, ante aperientiam ad scapum tortim inflexa plusque, postquam valde cernua, et tortula, laciniis corona recta, majuscula poculiformi delicatori eleganter plicata, ore plicatim-crenulato, plus duplo longioribus. Filamenta æqualia tria plus dimidiatis, cæteris altius tubo adnata et libera plus quam antherarum longitudine. Anthera superne curvatim semirectæ non versatiles. Cætera ut in Ajace, ut longissimè graciliores*.
- 1 *poculiformis*. T. (the great plaited cup) corollæ pendulae mox nutantis laciniis rectis semiexpansis oblongiusculis cum acumine concavo inflexis corona ore crenulatum erosulo. Queltia poculiformis. Salisb. in *Hort. trans.* 1. 352.
- Narc. montanus*. *Bot. Reg.* 2. 123.
- Obs.* Individuus, forte varietas, per tres annos remanet omni parte triplo minor.
- 2 *Galanthifolius*. T. (snow drop-leaved) corollæ laciniæ lacteæ nutantis subtortis substellatis ovato-lanceolatis lateribus infernè deflexis, corona poculiformi plicatissima. *Synops. succ. app.* p. 328. *Obs.* variat. Stylus longitudine coronæ vel plus, et 10 dies serius floret. Et stylo semper inclusu 10 dies præcocior.—Presented by Mr. Milne of Fulham, from his private Garden.
- IX. QUELTIA. Omnia fere ut in genere precedentibus, at omni parte grossiore. *Spatha* 1-flora. *Corolla* *nutans*, laciniis *corona* duplo longioribus, *corona* late campanulata ore valde dilatata et sexlobulatum crispo-undulatum. *Filamenta* paulo magis tubo adnata.
- * *Sulphureæ*, corona lutea.
- 1 *aurantia*. Q. (the orange rimmed) corollæ laciniis pallide luteis corona calyciformi perlutea, ore patulo 6-lobato, lobulis crispatis undulatis plicatis subimbricantibus. *Narcis. Revis.* 124. *Obs.* Semel vidi duos flores in eadem spatha.
- β *plena*: luteo-aurantia. Vulgo Butter and Eggs.
- 2 *Gouani*. Q. (the lobed orange rim) laciniis luteis per oras demum recurvas, distincti pallidè luteis, corona perlutea, supernè subaurantia sexlobata plicata crista undulata.
- Narcissus* -*Gouani*. *Redout. Lill. tab.* 158.
- Obs.* *Folia* florendi tempore pedalia glaucescentia valde nervoso lineata. *Scapus* *folia* æquans anceps striatulus.
- Q. aurantiæ proxima, sed minor, *coronæ* lobis certè longè distinctioribus profundioribus. Paulo serius floret.
- 3 *incomparabilis*. Q. (the great incomparable) corollæ laciniis sulphureis, corona late campanulari lutea, ore valde patulo, lobis magnis crispo-undulatim quasi imbricantibus.
- Narcissus incomparabilis*. *Bot. mag.* t. 121.
- * 4 *concolor*. Q. (the self-coloured incomparable) corollæ laciniis coronaque late campanulata lobata crispa uniformiter flavis concoloribus.
- Narc. odorus*. ð. Linn. *amæn. Acad.* t. 4. p. 311.
- Narc.* omnium maximus flore et calyce flavo. Park. *Par.* p. 68.
- β *plena-sulphurea*. The Sulphur Crown, and the Sulphur Phoenix; Hortulanorum. Varietas omnium plenarum forsitan sumptuosissima.
- 5 *semipartita*. Q. (the cloven-cupped) corollæ laciniis pallide sulphurascensibus, corona patentii-campanulata semisexpartita lutea, lobis parabolicis

- late concavo canaliculatis distinctis obsolete rugosulis. *Narcis. Revis.* p. 126.
- ** *Albiflores*, corolla laciñiis albis primo sulphurascensibus, vel omnino albis sive niveis; corona lutea.
- 6 *alba*. Q. (the great white) corolla laciñiis primo sulphurascensibus mox albis imbricantibus, corona saturate citrina, lobis maximè crispis. *Narcis. Revis.* p. 127. Narc. (odorus) latifolius maximus albo flore calyce brevi tubo. *Linn. amæn. acad.* 4. 311. β? plena niveo-aurantia. Vulgo, the orange Phoenix.
- 7 *nivea*. Q. (the lesser white) corolla stellaris laciñiis tortuosis, basi (per oras reflexas) distinctis niveis corona lutea angustè campanulata plicata crista undulata. Communicavit florentem amicus Johannes Denison, Junior, Mense Aprili 1831. Obs. corona minus patula, *Scapus* anceps pedalis subviridis ut *folia* striata, vix glaucescentia, ut in priore, in qua etiam omni parte major.
- X. SCHISANTHES. *Spatha* 3-4-flora. *Corolla* sulphureæ stellatim-patentes. *Corona* perflava semitripartito-patula, recta, laciñiis circiter duplo brevior. *Antheræ* trilineares. *Scapus* valde compressus fistulosus gracilis laevis substratalis. *Folia* lorata plana erecta viridia, ad latera arcè erecto-incurva. *orientalis*. S. (the triple-gashed.) *Narc. Revis.* p. 129.
- N. orientalis. *Linn. Aiton. Willd.* N. incomparabilis β. *Bot. Mag.* 948. Hermione ambigena. *Salisb. Hort. Tr. v. 1. p. 361.*
- XI. PHILOGYNE. *Spatha* 2-4-flora. Corolla concolor lutea. *Filamenta* gracilia tubo adnata, tria libera circiter antherarum longitudine, tria plus. *Antheræ* contiguae erectæ linearies altitudine tubi. *Stylus* subgracilis antheras longe superans corona brevior. *Corona* varians saepe heminæformis laciñiis subduplo brevior rarissime fere eadem longitudine. *Ovarium* oblongum: *ovula* oblonga subcapitata basi attenuata bifaria inserta.
- * *Distinctilobæ*. *Corona* sexlobata, lobis distinctissimis planis, laciñiis corolla distinctis.
- 1 *odora*. P. (the great 6-lobed) corolla stellaris laciñiis distinctis, corona laevi distinctissimè sexlobata patula campanulata, scapis 3-4 floris folio altioribus. *Narciss. Revis.* p. 134. N. odorus. *Linn. sp. pl. ed. 3. p. 416. et ejus Herbarii.* N. calathinus. *Bot. Mag.* 934, nec aliorum.
- Obs. Omnia major. *Scapus* 2-4 florus.
- 2 *Campernelli*. P. (the late great 6-lobed) scapis numerosis, 2-3-floris folio humilioribus, corolla laciñiis flexuosis distinctis corona 6-lobata patula duplo longioribus.
- N. Campernelli. *Hortulanorum*. Priori simillima, et forte varietas senior subpallidior corolla speciosior, corona lobis paululum minoribus, majis patulis seu recurvantibus, foliis altioribus florendi tempore Fl. Aprili in Maiam. Forte formosissima varietas solum.
- ** *Subsexlobata*: *Corona* 6-lobata, lobulis obsoletioribus seu minus distincte definitis crispis undulatis; corolla laciñiis valde imbricatis, prænæ qualibus.
- 3 *rugulosa*. P. (the great wrinkled cup) corona plicata interne rugulosa, lobis profundi semicircularibus basi crispatis imbricantibus, laciñiis corolla vix duplo breviore. *Narc. Revis.* p. 135.
- 4 *interjecta*. P. (the great-curved cup) sub-2-flora: corona obsoletæ crispatis duplice trilobulata seu irregulariter undulata laevi, externè obtuse angulata, laciñiis corolla imbricata non duplo breviore: scapo sesquipedali. *Narc. Revis.* p. 135.
- Obs. *scapus* solo fertili sesquipedalis, 2-3 florus.
- 5 *Curtisi*. P. (Curtis's) corolla laciñiis latè ovatis obtusis luteis, corona recta campanulari ore crispatis lobulato subsaturatione, subsesquilonioribus.
- N. odorus. *Curtis Bot. mag. t. 78. non Linnæi*, neque ejus Herbarii.
- N. laetus. *Hort. Kew. ed. 2. v. 2. p. 216. non Salisb. qui est Philog. triloba, infra.*
- * 6 *calathina*. P. (the greatest-cupped) spatha multiflora, nectario campanulato subcrenato æquante petala, foliis planis. *Linn. sp. pl. 1. 415.*
- Narcissus angustifolius* flavus magno calyce. *Rudb. Elys.* 2. p. 60. f. 5.
- Species propria, corona maxima quæ non vidi vivam neque sicciam, sed dubio sine bona; et praecedenti affinis; a qua nihilominus differt quandoquidem magna corona. Forte affinior *Phil. heminali* infra.
- N. calathinus. *Redout. Pl. Lill.* 417. est *Assaraci* species supra descripta.
- 7 *triloba*. P. (the lesser 3-lobed) spatha sub-2-3-flora: nectario campanulato subtrifido integerrimo dimidio breviore petalis.
- N. triloba. *Linn. sp. pl. 1. 147.* Philogyne triloba, *Narc. Revis.* p. 135. exclusis synonymis. *Bot. mag.* 78. et calathina, *Salisb.* et *Park. Par.* et *Observationibus finientibus*.
- Obs. *Corolla* laciñiæ inæquales subplanæ valde imbricata, *corona* sub-

- campanulari crispatim lobata duplo longiores, late ovatae submucronatae, interiores parum tortae ovato-ellipticæ. Corona recta ore obsolete 6-lobato plicatim undulato-crenato.
- * 8 minor. P. (the lesser) spatha biflora corona crispatim undulato-lobata: scapo dodrantali. *Narc. Revis. p. 137.*
N. rupicola. Schult. syst. v.7. p.958.
N. juncifolius. Schult. syst. p. 962. sed N. Assoanus. l.c. potius ad N. tenuirem. Bot. Mag. 379. pertinet.
N. Jonquilloides. Schult. syst. 7. p.988?
 $\beta.$ fl. pl. minor. Queen Ann's Jonquil.
 $\gamma.$ fl. pl. major.
 $\delta.$ fl. pl. maxima.
- All in the Epsom collection.
- *** Undato-coronæ. Corolla laciniis substellantibus seu angustioribus vix imbricantibus magis aquilibus, quam in precedentibus, corona heminiformi angustiora integriora sesqui et subsesquiongioribus.
- 9 heminalis. P. (the narrow cupped) spatha 2-3 flora, corollæ laciniis subæqualibus subimbricatis tortuosis, corona heminiformi subintegra sesquiongioribus. *Narc. Revis. p. 436.*
N. Juncifolius luteus magno calyce. Park. Par. t. 93. f. 4.
- Obs. Folia mediorius hujus generis, Juncoidea atro-viridia.
- β. pluriflora. P. duplo major; scapi duplo numerosiores 1-3 flores longè altiores, corona saepe nec semper longiore: foliis duplo majoribus pallidioribus. Forte propria species; communicavit amicus Dom. Peuny. *Narcissus Juncifolius maximus amplio calyce. Park. Par. t. 89. f. 5?*
- XII. JONQUILLA. Nomen antiquum genericum, sed non Linnæi more constructum.
- Spatha* 2-6 flora. Corollaæ concoloris flavæ laciniis in stellam quasi patentibus, corona parva ore eroso-crenato acetabuliformi seu patellari 4-5-plo longioribus. Scapi lèves obtusè subcompressi. Folia Juncina semiteretia, supra concavo-canaliculata erecta, sèpius atro-viridia. Flores delicatissimi suaveolentes. Semina incipientia subreniformia. Cætera ut in *Hermione*.
- 1 major. J. (the great Jonquil) 3-6 flora: sessipedalis atroviridis: corolla laciniis anguste-obovatis, tubo respectu horizontalibus, corona sub4-plo longioribus. *N. Juncifolius luteus vulgaris major. Park. Par. t. 94. 7. tab. 93. f. 5.*
- 2 media. J. (the middle Jonquil) pedalis, 3-4 flora, atroviridis subtriflora, corollaæ flavæ, laciniis late obovatis sub-
- spatuliformibus semireflexis, corona plus 3-plo longioribus.
- N. Juncifolius luteus mediocris. *Park. Par. p. 94. n. 8.*
- 3 minor. J. (the lesser Jonquil) 2-flora: 6-9-entalis, viridis, corollaæ paulo pallidioris, laciniis cuneato-ovatis, corona majore magisque expansa, 4-plo longioribus.
- N. Juncifolius luteus minor. *Park. Par. p. 93. No. 9. Hermione similis. Salisb. in Hort. Tr. 1. p. 359.*
- 4 parvicorona. J. (the little-crowned) pedalis: atroviridis: gracilis: biflora: corolla laciniis stellaribus flavis, corona saturatio paulo, omnium minore.
- XIII. CHLORASTER. *Spatha* 1-3-flora. Corollaæ viridis laciniæ in gracillimam stellam patentes. Corona minima integra, vel 6-partita, partibus incurvatum cochleatis incrassatis. Filamenta omnino tubo adnata, 3 tubo breviora, 3 ejus longitudine.
- * 1 fissus. C. (the cloven cupped) florendi tempore aphyllus, corona hexapetaloidæ. *Phil. Mag. Mar. 1824.*
- Narcissus viridiflorus. Bot. Mag. f. 1687.*
- * 2 integer. C. (the intire cupped) spatha 3 flora, corona integra ante foliatione. *Narcissus Juncifolius autumnalis flore viridi. Park. Par. 94. 11. tab. 93. f. 6.*
- XIV. HERMIONE. *Spatha* 3-20 flora. Corollaæ laciniis in stellam patentes, corona parva calycoformi seu cyathiformi vel cupulari 2-5-plo-longiores. Tubus gracilis angulato-cylindricus: corolla laciniis longior viridis. Filamenta fere toto tubo adnata apice solum semilineam libera: tria tubo longe breviora, tria ejus longitudine. Antheræ parvulae trigono-ovatae erectæ, tres tubo inclusæ, tres id semisuperantes. Stylus rectus inclusus. Stigma plus minus trilobato-trifidum, lobis rotundatis.
- * Flavocoloratus; levigati pervirides, scapis teretiusculis vel parum compressis, foliis plus minus Juncaginosis, erectis, seu altè, concavo-canaliculatis, subulato-acutis. Genus proprium? Flavae.
- * 1 intermedia. H. (the intermediate) foliis canaliculatis viridibus basi subsemicylindracis, scapo subcylindrico 1-3 flora, corona cyathiformi ore dilatato criso subintegro, segmentis quadruplo breviore.
- N. intermedius. *Red. Lill. t. 427.*
- Narcissus major Juncifolius. Hort. Eyst. tet. tertius Ord. fol. 4. f. 1. N. intermedius γ. Schult. syst. 7. 963. sed var. β. l. c. est Philogyne triloba Nob. Var. α Schult. nescio, nisi est H. compressa infra.*

- * 2 *bifrons*. H. (the long flowered) 1-5 flora, pedunculis longis effusis gracilibus, corollæ laciniis stellaribus, corona calyciformi sexlobata ter longioribus.
N. bifrons. Bot. mag. 1186. N. longiflorus. Willd. enum. v. 1. p. 351.
N. polyanthus flore minore stellato toto luteo. Rudb. Elys. t. 60. f. 7.
Folia semic. lata.
- 3 *compressa*. H. (the Jasmine Jonquil) 2-6 flora: pedunculus suberectis parvum ad germen paululum flexis, corollæ laciniis imbricatis, corona erecta eroso-crenata, subtrilobata longioribus.
Obs. folia singulariter lata angustaque eodem bulbo.
N. compressus. Nob. in Lin. Tr. v. 5. p. 245. H. intermedium a. Schult. syst. v. 7. p. 965? quæ non vidi? (nisi est speciem presentem) neque figuram.
N. radiatus. Red. Lill. t. 459.
N. tripartitus. Schult. syst. 7. p. 965.
- 4 *primulina*. H. (the cowslip cupped) subquadriflora: scapo obtusissimè comppresso, corolla laciniis perluteis ovatis imbricatis, corona subaurantia patula sexlobata 3-4-plo longioribus. *Narc.* Revis. 140. *Narc. bifrons* β. Bot. Mag. 1299.
- 5 *biscrenata*. H. (large cowslip cupped) scapo lœvissimo 6-9-floro vix compresso, corollæ laciniis perluteis latissimè ovatis, corona sub 6-lobata, lobis plicato-crispulis bilobulatis, plus duplo longioribus. H. primulina β. *Narcis.* Revis. p. 140.
Obs. A priore precipue differt, statura majore, floribus duplo numerosioribus, lobis normalibus sex, itidem rebulatibus seu bifidis; inde corona sub 12 lobulata-crenata est.
- †† *Alba*.
- * 6 *alba*. H. (the rush-leaved white) subsex-flora: corollæ-laciniis ovatis acutis imbricantibus albis, corona concinna crenata fere duplo longioribus: foliis juncagineis obtusulis.
Narcissus juncifolius polyanthus albus. Hort. Eystt. Ord. 3. fol. 17. f. 1.
 Non vidi, sed l. c. solum descripsi.
 Species valde singularis flore albo.
H. biscrenata, supra dubio sine proxima: a qua precipue alba corolla discrepat.
- ** *Rotuliflora*, ochroleucæ seu lutescentes cupulari corona.
- 7 *tereticaulis*. H. (the green-leaved cream coloured) viridis: sub6-flora: corollæ laciniis rotundo-ovatis imbricatis flos lacticoloribus, corona citrina subpatula subintegra lobulatae sesquiduplo longioribus, scapo inferne teretiusculo. *Narc.* Revis. p. 140. *N. orientalis*. Bot. Mag. 1298. *N. ochroleucus*. Loi-
- sel. not.* p. 163. *Narc.* p. 169. *Flor.* gall. ed. 2. 1. p. 236. *Schult. syst.* 7. p. 963.
- Obs.* Corona subindè definitè sexlobata. *Folia* angustiora, quam in sequentibus, sed saepè sex lineas lata, extus magis convexa, supra magis inflexo-concava. *Flores* elegantissimè fastigiatim capitulati.
- 8 *rotularis*. H. (the glaucous leaved) sub8-flora, glaucescens: corolla rotularis ochroleucæ laciniis imbricatissimis, corona lutea duplo longioribus, foliis secundum latis planis obtusissimis. *Narc.* III. *Trew Flor. Imagin.* t. 105. *Barrel. icon.* 918.
- Obs.* Forte omnium obtusissima *folia*, scapus sesquipedalis teretiusculus. *Flores* demum lactei eleganter capiti, sed in juventute incipientes, seu ante aperientiam insigniter abbreviati crassi, et quasi plenos flores simulantes. *Corolla* laciniis amplissimè rotundato-ovatis variè flexis mucronatis et quasi plicatulatum rugosulis. *Corona* subcupulari paululum ore patulo, plicatim quandoque irregulari integra. *Præcocius* floret quam priore.
- 9 *flos-lactis*. H. (rich creamy white) sub7-flora: *Corolla* rotularis laciniis flos lacticoloribus apice rotundatis maximè imbricatis (tubo respectu) horizontalibus, tubo tertia parte brevioribus, corona cupulari integra perlutea duplo longioribus.
- Obs.* Nova species. *Folia* lorata plana viridia unciam basi lata altitudine scapi, sub anthesin fistulosi, subcompressi grossi subancipitis.
- 10 *multiflora*. H. (the yellowish) sub14 flora, corolla rotularis laciniis pallide luteis, corona flava saturatiore subcupulari ore truncato-repandulato sub duplo longioribus: foliis lorato-ensiformibus. Herm. multiflora. β. *Narcis.* Revis. p. 146. *N. orientalis* δ. Bot. Mag. 1026.
- β. floribus pallidioribus.
- *** *Flaviflora*. Corolla laciniis longioribus seu acutioribus quam in sectione priore, plus minus luteis flavisve, corona sæpiùs aurantia.
- † Corona plus minus cupulari.
- 11 *sublatea*. H. (the pale yellow) sub9-flora, corolla subrotularis laciniis pallide luteis ovatis, corona saturatissime lutea subcupulari ore patulo integro fere triplo longioribus.
Obs. Communicavit florentem amicus Dom. Penny. f. Ap. 1831, cum sequenti.
- 12 *perlutea*. H. (the full yellow) sub4-flora, corolla laciniis (tubo respectu) horizontalibus perluteis, corona cupulari

aurantio-lutea suberenata duplo plusve longioribus.

Obs. Similis parum, H. cupulari. *Bot. Mag. t. 925.* Sed differt coronā longē pallidiore integriore, &c., et senior in f. Aprilis floret. Affinior precedentem est, et varietates bellicas vidi ambas valde simulantes: et inde fortè solum prioris varietas.

13 *flaveola*. H. (the lesser pale yellow) sub-flora; corollæ laciñis ovatis imbricatis flavescentibus, corona cupulari, subintegra lètè aurantia duplo longioribus.

Obs. Inter minores est. Florendi tempore subdodrantalis. *Folia* viridia obtusa vix glaucescentia 5-6-lineas lata altitudine *scapi* florantis obtuse compressi lèvis. *Corolla* laciñis tubo fere duplo brevioribus. *Stylus* et tres humiliores *antheræ* cum *polline* auroeo, inclusæ. Communicavit cum sequente, ex Caldario, amicus Dom. Sweet, Martio, 1831.

14 *deflexicaulis*. H. (the self-coloured yellow) floribus subseptenis luteo-subcoloribus secundis, laciñis latè ovatis imbricatis, corona cupulari, scapo declinante, spatha abbreviata acuta.

Narc. bizantinus luteus. *Park. Parad. p. 78. No. 3.*

Obs. *Folia* lata lorata erecta plana glaucescentia obtusa *scapus* ad anthesin subpedalis teretiusculus s. subcompressus substriatulus firmus sed inanis, apicem versus precipue, et foliis altior, et concolor. *Flores* inter mediocres; subhorizontales vel nuntantes. *Pedunculi* $\frac{1}{2}$ -ad 3 uncias longi erecti triangulares virides altitudine *spathæ*. Germen trigono-ovale perviride. *Corollæ* *tubus* lutescens trigonus, *laciñis* *exterioribus* lato-ovatis acuminate horizontalibus vel demum parum reflexis planiusculis, tubi longitudine vel brevioribus, *interioribus* minoribus ut in plurimis. *Corona* cupularis laciñis in ætate paulo saturator ore præplicatum subcrenulato, *laciñis* plusquam duplo brevior. *Antheræ* *corona* concolores. Cetera ordinaria.

B. plena. Narcis. cyprius flore pleno luteo polyanthos. *Park. Parad. p. 86. N. 6.*

15 *trifida*. H. (trifid crowned yellow) sub-3-flora: corollæ laciñis pallide luteis, corona saturatice subcampanulata trifida (lobis bifidis) subtriplo longioribus: foliis 9-lineas latis loratis planis obtusis glaucis.

Obs. In charactere, foliis exceptis, fere cum N. *tripartito*, Schult. syst. t. 7. p. 965, concordat, sed, in fo-

liis glaucis latis obtusis, (non ullo modo Juncagincis) abundè discrepat.

Obs. In my rambles after Narcissi in the Nursery Gardens near London, I detected this species going out of bloom in the open ground of the Nursery of Messrs. Allen and Co., in the Five Fields, King's Road; who, when I expressed a desire to publish it as a new species, immediately presented it to me, with a view of aiding me in my long-tried and well-known endeavours to elucidate these charming and fragrant plants; and ever friendly to the science of Botany, they have often so assisted me before.

† *Corona acetabuliformi*.

16 *aperticorona*. H. (spreading orange cupped) sub6-flora: corollæ laciñis semireflexis late ovalibus preimbricatis perluteis, tube brevioribus, corona aurantia patula plicatim subrepandula sesquiduplo longioribus.

Obs. *Folia* lorata plana glaucescentia obtusa, 10 lineas lata. *Scapus* medioris compressus striatus. *Floret* medio Aprilis.

**** RUFO-AURANTIE, Corollæ laciñis flavissimis, corona rufo-aurantia, seu saturatissimè aurantia.

17 *cupularis*. H. (the deepest orange cup) sub10-flora, corollæ laciñis saturate flavis, corona cupulari rufo-aurantia truncata subrepandula subtriplo longioribus.

H. cupularis. *Sulisb. in Hort. Trans. l. p. 361. N. flavus. Lagasc. nov. gen. et spec. p. 13. Narcissus Tazzetta, Bot. Mag. 925. et Redout. Lill. t. 17. Nec Herbarii, Linn. N. africanus aureus major. Park. Parad. t. 81.f. 1. Soleil d'or Hortulanorum.*

18 *solaris*. H. (the starry orange) corollæ stellaris, laciñis subovatis distinctis ad oras reflexis flavissimis, corona rufo-aurantia cupulari subintegra, sesquiduplo longioribus.

Obs. A precedentem cui simillima, pulcherrimè radiantibus laciñis valde distinctis et non imbricatis ut in ea, discrepat.

Fine Aprilis, A.D. 1831, florentem, Dom. Allen and Co.: mihi donabant.

19 *latifolia*. H. (broad-leaved orange cup) subtriflora: corollæ laciñis oblongo-rotundis valde imbricatis flavis, corona rufo-aurantia recta integra fissave, 2-3-plo longioribus, styllo tubo incluso. *Narciss. Revis.* 145. *Floret* f. Apr. 1. Maii.

Obs. H. cupulari, humilior, *corolla* minus rotularis, *corona* subinde rupratim 6-fissa, seu 6-partita, et tunc

- plus minus patula. *Folia* eusiformia lorata obtusa basi uncian lata. Seminalem exemplum habeo, cum floribus septem, et corona integra, sed vix distinctum.
- ***** *GRANDIFLORÆ*, spatha 3-8-flora ; corollæ laciniis omnium hujus generis longioribus et speciosioribus imbricantibus albis rarius sulphureis luteis corona conspicua patula acetabuliformi lutea subesquiduplo longioribus. Semina incipientia subcylindrico-clavata. Genus proprium ?
- 20 *Trewiana*. H. (great broad petalled) sub8-flora : corollæ amplissimæ rotularis laciniis albis planis, corona patula seu plicatim crenulata toto lutea.
H. Trewianus. Sweet. Br. Fl. Gard. Ser. 2. t. 118.
N. Trewianus. Kerr. in Bot. Mag. fol. 1298. N. grandiflorus. *Synops. succ. appendix*. 330. N. orientalis. a. Gauver in Bot. Mag. 940. Bazleman major. *Trew. flor. Imagin. Narcissus*, I. t. 23.
- * β *fissicorona*. (the cloven cropped) subtriflora : corona majore patente trilobatum fisco, lobis magnis 1-4 relobulatis truncatis.
Forsan propria species.
- 21 *flexiflora*. H. (flexuose large flowered) sub3-flora : corollæ laciniis (tubo respectu) horizontalibus obovatis variè deflexis et flexulis niveis, corona lutea plicatim subundulata patula sesquiduplo longioribus.
Obs. Corollæ laciniæ exteriore precepit semper plus minus ad oras deflexæ, et variè in aliis locis oblique tortuæ, apice mucronulatæ, ipsa basi luteæ. *Stylus* (in Caldario solum vidi) brevissimus fere ad basin tubi.
Obs. Pedalis (florend. temp.) foliis loratis glaucescentibus.
- 22 *subcrenata*. H. (great yellow cupped) 3-5-flora : corollæ laciniis subreflectentibus albis, corona patula plicatulum crenata toto lutea 2-3-plo longioribus, foliis loratis glaucis.
Cum priore floret. Hortulanorum Bazleman medius.
- 23 *crenulata*. H. (great saffron rimmed) sub3-flora : corollæ laciniis subreflexis albis, corona patula lutea, croceo marginata plicatulum crenulata 2-3-plo longioribus, foliis loratis angustioribus. *N. crenulatus*. Nob. in Lin. trans. v. 5. p. 245. N. Mussarti. Park. Parad. t. 81. f. 5. Bazleman minor. *Trew. flor. imagin. Narcissus*, II. t. 62.
- 24 *sulphurea*. H. (great sulphury) sub5-flora : corollæ laciniis varie tortulis ad oras reflexo-stellantibus sulphureis; corona crocea oris pallidis fimbriatis.
- Narciss. polyanthemos præcox sulphureus maximus calyce croceo oris pallidis et fimbriatis. *Barrel* ic. 936.
- 25 *lutea*. H. (great yellow) lutea : sub3-flora : corollæ laciniis magnis longis et acutè oblongis flexulisstellantibus luteis. *Narcissus luteus polyanthos lusitanicus*. *Rudb. Elys. tab. 69. f. 2.* Bazleman luteo, Hortulanorum.
Obs. quatuor precedentibus similis sed corollæ laciniis longioribus in figura supra citata et magis distinctis stellantibusque. Non vidi, neque precedentem.
- Obs.* The fine large flowered plants of this section are known to Gardeners by the several names of "Bazleman major, medius, minor, sulphureus et luteus." The last two, which appear to be distinct species, I have never yet possessed, but taken up on the authorities abovementioned, with a special view towards inducing all lovers of these plants to search for them. They are nearest allied to *H. flexiflora* above, and should perhaps have preceded it.
- ***** *SUBLONGIFLORÆ* : spatha 4-20-flora : corollæ laciniis longiusculis albis seu lacteis acutis corona lutea seu aurantiaca 2-3-plo longioribus.
- 26 *Tazzetta*. H. (the white, yellow cup) sub5-6-flora : corollæ laciniis albis oblongo-ovatis acutis imbricatis, corona recta subintegra lutea, sæpe triplo longioribus in eadem spatha.
H. Tazzetta. *Narc. Revis.* p. 442.
N. Tazzetta. Lin. sp. pl. ed. 3.
N. Narbonensis medio luteus. *Park. Parad.* t. 81. f. 3.
- 27 *Cypræ*. H. (the obovate cyprian white) scapo gracili, sub4-floro : corollæ laciniis obovato mucronatis subimbricantibus semireflexis et varie subflexulis, corona cupulari, lutea truncata sesquiduplo longioribus.
H. Cypræ. Nob. in Phil. Mag. Mart. 1831. Sweet Brit. Flower Garden, t. 92. series 2.
 β *semiplena*. *Park. Parad.* 85. f. 22.
 γ *plena*. *N. orientalis* var. *fl. plen.* *Bot. mag.* t. 1011.
Obs. I have seen plants this season at Messrs. Allen and Rogers' Nursery, in the King's-Road, Five Fields, as finely grown, and with very nearly as many flowers as the fine figure above cited, of this most favourite and fragrant plant.
- 28 *lacticolor*. H. (milk-coloured Grecian) sub7-flora : corollæ laciniis lacteis (tubo respectu) horizontalibus planis ovatis valde imbricatis, corona recta perlutea cupulari lobulatum crenata subplicata subrugulosa sesquiduplo longioribus ; foliis glaucis planis.

N. Tazzetta. *Fl. Græc.* t. 308.

A specimen of this was communicated by my very intelligent friend Mr. Penny, foreman to the fine nursery of Messrs. Young, at Epsom, the middle of last April.

***** CURTIFLORÆ, scapi multiflori teretiusculi saepè plus minus fistulosi. *Folia* lata plana lorata plus minus glauca. *Corolla* laciniis sèpius æqualibus et latioribus quam cæteris, et inde brevioribus, seu quasi curtis; semper albis, *corona* lutea, rarius subaurantia sesquivel rarius duplo plusve longioribus. *Medio-luteæ*, *Hortulanorum*.

* 29 *cerina*. H. (large waxen cuffed) sub4-flora; corona crateriformi præampla subincrassata truncata integra luteo-*cerina*, laciniis (*corolla*), albis ovatis basi imbricatis duplo breviore. *Narc.* *Revis.* p. 329.
Floret Maio.

30 *fistulosa*. H. (the pipe stalked) sub9-flora: corolla laciniis albis subreflexo-incurvis ovatis æqualibus valde imbricatis, corona recta inflata subintegra lutea, fere duplo longioribus. *Narc.* *Revis.* p. 142.

31 *patula*. H. (lesser, patulous) foliis glaucescentibus subcanaliculatis patulis scapo subcylindrico 2-6-floro, corona cyathiformi subintegerrima aurea petalis niveis alterna latioribus dimidio breviore. *Schult. syst. v. 7. p. 977.* N. *patulus*. *Loisel. Journ. d. 1809. 2. p. 276.* *Narc.* p. 625. *Not. p. 52. et p. 159.* *Flor. gall. ed. 2. 1. p. 235.*
Vidi, sed non examinavi. H. *tazzetta* supra duplo minor, et precedentis affinior.

Obs. I think there are many varieties of this plant in gardens, and that one of them, whose flowers have the (perhaps unique) quality of dyeing off with tints of dull or dirty purple colour, have been found apparently wild in two different counties of England; and I have seen plants from both, growing in the rich Nursery at Epsom. If distinct, it may be called *rubescens*.

* 32 *breviflora*. H. (the short round flowered) sub4-flora: corolla laciniis æqualibus albis orbicularibus imbricatissimis; corona lutea integra subdupo longioribus.

Obs. Dubio sine, nova species, *corolla* laciniis omnium rotundioribus; foliis loratis et in nostro exemplari, solum 6-lineas latis. *Descriptio* e sicco hortense specimeine solum.

33 *auranticrona*. H. (the orange-crowned white) sub9-flora: corolla laciniis niveis ovatis variè flexis imbricatis,

corona cupulari acetabuliformi subrepandula lätè aurantia plus duplo longioribus.

Cum priore et sequente fortè accedit ad *H. fistulosam*, sed distincta *corollæ* majore, etiamque *coronæ* colore.

Communicavit eum sequente; fine Aprilis A.D. 1831. Amicus Dom. Penny.

34 *crispicrona*. H. (the curly-crown) sub11-flora: Corolla rotularis, laciniis albis rotundo-ovatis subrugoso-flexili imbricatissimis; corona aurantiolutea cupulari campanulata plicata, (ore præcrispo) laciniis fere duplo breviore.

Obs. *Corolla*, laciniis, tubo respectu, horizontalibus variè incurvulis et quasi ruguloso-flexilis. *Corona* ore indefinitè 6-lobulato criso, lobulis *Philogynum* modo imbricantibus. *Folia* ensiformi-lorata plana glauca obtusa scapum florendi tempore æquantia.

Obs. Forsan hujus sectionis tres ultimæ species, melius associent inter se. *Rotulifloræ*, supra.

***** CITROCORONÆ, 5-20-floræ. Corolla laciniis albis minus abbreviatis quam in sectione priore, coronis sub-cupularibus primè citrinis, denique albanticibus albisis.

35 *floribunda*. H. (the Nosegay) sub16-flora: corolla magna, laciniis rotundo-ovalibus subreflexo-incurvulis albis, corona ampla citrina subrecta integra 3-4-plo longioribus. *Narc. Revis.* p. 141. The Grand Monarque of Gardeners, of which I have seen many minor Dutch varieties.

36 *citrina*. H. (the common Citronniere) sub10-flora: corolla laciniis albis rotundo-ovatis, corona citrina subpatula irregulariter saepè fissa subtriplo longioribus. *Narcis. Rev.* p. 141. *Narcissus orientalis* γ. *Gawler in Bot. Mag.* 946.

Obs. Priori affinis, at differt structura scapi striati teretiusculi, foliorumque.

37 *decora*. H. (the neat lemon cuffed) corolla laciniis obovatis niveis, corona pallidè citrina cupulari (ore, repandulo) duplo longioribus.

Obs. Folia ensiformi-lorata mediocria subglaucescentia. *Scapus* anceps; striis elevatis. *Corolla* concinnæ, mucronulatae, ut in plurimis.

Obs. Flowered finely in the splendid Nursery of Mr. Colvill, in the King's-Road, Chelsea, in the end of April last, in the open ground, along with the next but one; and to him I am greatly obliged for a root in bloom.

38 *polyantha*. H. (the many-flowered pale cup) foliis planiusculis virescentibus, scapo subcylindrico, ancipti, 8-20 floro, nectario cyathiformi, subintegerrimo, petalis ovatis, alternis latioribus, subtriplo breviore; floribus albis. *Narcissus polyanthus*. *Loisel. Schult. syst.* 7. p. 973. N. orient. totus albus major. *Hort Eyst. ord.* 3. fol. 11. f. 1. N. orientalis major *polyanthus* totus albus. *Moris. sect.* 4. tab. 8. f. 2.

Obs. Folia viridia. *Nectarium* pallidissime flavum serius omnino album. *Loisel.* l. c. Vidi 16-flores solum, in una spatha, laciniis ovatis acutis.

β. sub-3-8-flora, petalis angustioribus, corona citrina, foliis latissimis subensiformi-loratis viridibus. *Floret f. Martii. Habitat in Gallia?*

39 *sulcicaulis*. H. (many pale flowered, furrowed) 8-16 flora : corollæ laciniis ellipticis mucronato-acutis ad oras recurvis, subimbricatis niveis, corona cupulari tenuiter crenulata citrina 3-plo longioribus.

Obs. Folia mediocria ensiformi-lorata subglaucouscentia. *Scapus* foliis concolor compressus, utraque longitudinaliter obtusè sed profundè sulcatus, angulis obtusissimis. Communicat amicus Dom. Sweet.

40 *Luna*. H. (the pale few-flowered) 3-5 flora corollæ laciniis niveis imbricatis, corona primo pallidissime citrina, mox alba, sub3-plo longioribus : foliis ensiformi-loratis viridibus planis.

Obs. Folia seriora sæpe 8-lineas lata plana, sæpe superne debiliter flexuoso-recurva.

***** ANCIPITES: Italicae: *Scapis* omnium præcocioribus, valde compressis gracilibus, acutè ancipitibus. Genus proprium?

† *Sulphureæ*: 3-9-floræ, corollæ laciniis primo sulphureis seu rarius flavescentibus demum plus minus albicanibus, sæpius ovato-lanceolato-acute-mati, fere elongatis; vel simpli-citer lanceolatis.

41 *chrysanthæ*. H. (the golden) subtriflora : corollæ laciniis saturatissimè sulphureis lanceolatis distinctis, (tubo respectu) flexuose horizontalibus, coronæ cupulari obsoletè crenato-repandulæ aurantiæ, 4-plo plusve longioribus.

N. *chrysanthus*. *Redout. Lill. vol. t. Schult. syst.* v. 7. p. 971.

Habitat, ex structurâ, in Italia. *Floret* i. Apr. sub dio.

H. *tenuifloræ*. Nob. *affinis*. Folia lorata plana glauca ordinaria, uti planta, inter minores. *Scapus* com-

pressus gracilis acutè anceps. *Pendunculi* triangulares fere ut in omnibus. *Flores* horizontaliter subnantes. *Corollæ* laciniæ *tubo* breviore sub 9-lineas longæ. *Corona* rectissima saturatissimè lutea potius subaurantia.

Obs. I have seen in Messrs. Young's fine Nursery, at Epsom, 2 semi-double (plenicoronæ) varieties of a plant, extremely near to this, but rather paler; and two more as large again, but similar. Few points amongst monocotyledonous plants are more difficult, than to ascertain the true parents of double varieties by study only. They may become single, and may again become double. They are still varieties from seed in every stage.

Obs. 2. Since the printing of the former edition, the author has obtained, and has at this time actually growing in his garden, no less than eight different *Herminies* with double crowns; (*Plenicoronæ*.)

These he hopes to flower in the ensuing spring, and to be able to determine and refer to the proper species from which they sprung; for they are merely seminal variations; but they are very beautiful, and extremely fragrant. November, 1831.

42 *italica*. H. (the large early sulphur) sub10-flora : corollæ laciniis elliptico-lanceolatis subflexiosis basi imbricatis primo sulphureis, corona citrina subquadripli longioribus : *stylo* coronam crenato-erosam patulo-cupularem aquanti.

N. *italicus*. *Bot. Mag.* 1188.

Obs. Spatha inter alias magna viudi-membranacea, fere foliacea.

* 43 *præcox*. N. (the middle early sulphur) corollæ laciniis lanceolato-acuminatis tortuosu-undatis subdistinctis primo sulphureis, corona recta citrina subcampanulata plicata 6-partita quadripli longioribus.

N. *præcox*. *Tenore Fl. Neap.* p. 146. t. 27.

44 *straminea*. H. (the lesser sulphur petaled) corollæ laciniis lanceolatis cum mucronulo sulphureis mox stramineo-albicantibus sub-distinctis, tubo sæpe brevioribus, corona cupulari citrina plicatim erosio-crenata, sub4-plo longioribus.

Obs. Priori simillima, sed in latioribus laciniis discrepat.

Ex Caldario cum sequente, communicavit amicus Dom. R. Sweet.

45 *subalbida*. H. (the straw-coloured) sub4-flora : corollæ laciniis anguste lanceolatis lacteis basi subimbricantibus, corona cyathiformi straminea plicatim

- eroso-crenulata, sub4-plo longioribus.
N. subalbidus. *Loisel. Schult. syst. v. 7.*
p. 967.
- Obs.* Plusquam pedalis omni parte
gracilis. *Folia* 7 lineas lata viridia,
uti *scapus* apicem versus altè stri-
atus.
- * 46 *tenuiflora*. H. (the slender-petaled sul-
phur) sub4-flora : corollæ laciniis sor-
dide albis lanceolatis stellatis distinc-
tis, corona minuta lutea sublacera in-
curvo-erecta 5-plo longioribus. *Philos.*
Mag. for 1830. p. 133. *Floret* sub dio-
i. Martii. An precedentis varietas?
†† *Niveafloræ*, Floribus toto niveis gra-
ciliere elegantissimis suaveolentibus.
- 47 *papyratia*. H. (the great paper white)
sub11-flora : corolle niveæ laciniis
subovo-lanceolatis imbricatis, coronâ
cupulata subplicatim erosula sub4-plo
longioribus.
N. *papyratius*. *Bot. Mag.* 947. N.
niveus. *Loisel. Schult. syst. veg.*
v. 7. p. 975.
- 48 *dubia*. H. (the lesser white) sub9-flora :
petalis ovatis nectario campanulato
integerrimo triplo breviore.
N. *dubius*. *Gouan. Illustr.* p. 22.
Willd. sp. pl. 2. p. 40. N. totus
alb. medium polyanthus, &c. *Red.*
Lill. t. 428. Hort. Eyst. ord. 3. fol.
2. f. 2. *Tazzettae* duplo humilio.
Petala ovata uniformia, nectario
campanulato integerrimo albo, triplo
longiora. *Willd. l. c.*
Obs. Dodrantalis plusve, *scapo* anci-
piti, *flores* toto albi. Init. Maio
florebat.
An vero hujus sectionis?
- * 49 *unicolor*. H. (the winter flowering) mul-
tiflora: petalis oblongis, nectario cam-
panulato plicato lacero petalis sextu-
plo breviore, scapo foliis planis ligulatis
longiore. *Tenore* (sub *Narciso*) *Flora*
Neap. v. 1. p. 144. t. 26. *Schult. syst.*
v. 7. p. 976.
Floret in loco natali, Novembro in
Decembrem.
- 50 *Jasminea*. H. (the Jasmine-like) corollæ
elegantissimæ, laciniis lanceolatis dis-
tinctis, corona erosula 5-plo longiori-
bus. *Phil. Mag. for 1830,* p. 133.
- †† *Serotinae*. Folia synanthia.
- * 51 *elegans*. H. (the slender Jasmine) 1-7 flo-
ra: pedunculis gracillimis elongatis, co-
rollæ albae gracilis elegantissimæ, laciniis
lineari-lanceolatis acuminate albis, co-
rona minuta integra multoties longiori-
bus.
Narcissus serotinus. *Desfontain. Flor.*
Atlant. v. 1. p. 283. tab. 82.
Habitat in Algeriæ arvis, secundum
Desf. l. c.
- Obs.* Haec planta omnium forte ele-
gantissima, sed non est, N. serotin-
nus Linnæi, neque ejus Herbarii,
quandoquidem *lavis* non trinodosus
scapus.
- * 52 *obsoleta*. H. (the white leafy autumnal)
2-flora : corolla laciniis albis ovato-
oblongis acutis imbricantibus, corona
obsoleta lutea multoties longioribus.
Narcissus autumnalis, medio obsoletus,
the white autumn Daffodill with a sul-
len crown. *Park. Par. tab. 89. fig. 4.*
p. 90. No. 1. *Swert. Flor. t. 30. f. 2.*
Obs. *Folia* 2-3 valde angustata et fere
Juncina, sed basi paulo latiora, et
apice magis acuta. Non vidi, sed
Parkinsoni figura etc. solum de-
scripsi.
- †† *Nudicaules*: *scapo*, florendi tempore
aphyllo nodoso vel articulato. Genus
proprium?
- * 53 *obliterata*. H. (the obliterated cup) sub8-
flora : nectario oblitterato, foliis linear-
lanceolatis.
N. oblitteratus. *Schult. syst. v. 7. p. 981.*
Habitat in Magadore.
- Obs.* *Scapus* pedalis. *Folia* scapo bre-
viora $\frac{1}{2}$ poll. circiter lata obtusius-
cula. *Flores* subumbellati erecti pe-
dicellati. *Spatha* floribus parum bre-
vior. *Schult. l. c. De Schlectend. Fil.*
Obs. An hujus Generis? Ab ultima,
penultima quoque satis differt foli-
orum latitudine.
Forte H. *Jasminea* supra magis con-
cordat; vel H. elegans. Non vidi.
An N. *Broussonetii*. *Lagasca, Nov.*
Gen. et spec. p. 13? "ob folia linear-
lanceolata."
- * 54 *serotina*. H. (the knottystalked autumnal)
uniflora : scapo digitali nodoso, corol-
lae albae laciniis patent-stellatis oblon-
go-ovatis acutis, corona lutea sexden-
tata 7-plo longioribus. *Swert. Fl. t. 30.*
f. 4.—*Obs.* foliis post anthesin pro-
gredientibus fere ut in *Scilla* autumnali.
N. *serotinus*. *Linn. sp. pl. ed. 3. p. 417,*
et ejus Herbarii. N. *autumnalis* mi-
nor albus. *Park. Par. tab. 89. f. 3.*
An verè hujus generis, ob scapum ano-
malum nodosum.
- XV. HELENA. (Hermionis formosissima
mater.)
Spatha 1-3-flora. *Corollæ* flavæ, ochro-
leucæ s. albæ, laciniis in stellam ex-
pansis, tubo multo brevioribus. *Tubus*
grossus valde compressus, angulis obtusis,
fere ut in *Narciso*, at longior,
magisque compressus. *Filamenta* omnia
fere omnino toto tubo adnata. *An-*
thera demum subspiraliter tortæ, tres
superne parte tubi inclusæ; tres tubum
semisuperantes. *Stylus* tubum æquans,
stigmatibus tribus rotularibus. *Ovalæ*

- numerosæ, bifariam insertæ, obovato-spathulatae. Cætera ut in *Narciso*. Character H. gracili sumptus.
- Habitus* in omnibus gracilior multo quam in ultimo, vel sequenti genere.
- Folia* linearia fere plana, sublineam lata, et viridia seu glauca.
- * *Flavissimæ*. Foliis viridibus.
- 1 *gracilis*. H. (the slender yellow) 1-3-flora: scapo gracillimo compresso supernè ancipti lœvi, spatha pedunculo brevior infernè integra: corolla flavissimæ, laciniis ovato-acutis valde imbricatis horizontalibus, corona saturatiore patula acetabuliformi ore plicatim eroso-crenulato marcescente, 5-6-plo longioribus.
- N. gracilis. *Bot. reg. t. 816. Swt. br. fl. gar. ser. 2. v. 2. ined.*
- N. lœtus. *Red. Lill. 428. nec 427. vel Salisb.*
- Obs.* Maio usque ad medium vel fere in finem floret.
- ** *Floribus ochroleucis, foliis angustissimis, sed planis nec Juncinis, vel infernè parum convexis, supernè parum concavis, viridibus.*
- 2 *tenuior*. H. (the slender straw coloured) 1-2-flora: prægracilis: corolla laciniis ochroleucis mox albicantibus, corona lutea patellari 3-4-plo longioribus.
- Hermione tenuior. *Narc. Rev. p. 145. N. tenuior. Curtis in Bot. Mag. 379.*
- 3 *planicorona*. H. (the slender, flat-crowned) corollar laciniis ochroleucis, corona lutea deplanata, foliis lineam lati. *Obs.* Omnim planifoliarum angustissima folia.
- Narcis. minimus Juncifolii flore. *Park. Par. 88. No. 4. tab. 89. f. 2.*
- Amicus Rev. Dom. Ellincomb, fere post florescentiam, Narcissorum eximius cultor, Aprili mense, communicavit.
- Obs.* Priore plus duplo gracilior flore simili, at in corona plana discoidea sufficienter discrepat.
- *** *Dodrantales: floribus albis solitariis corona parva lutea, alio colore, marginata.*
- *4 *purpureo-cincta*. H. (the dwarf purple rimmed) corolla diminuta alba, corona parva lutea, purpureo tenuissime cincta. *Narcissus minimus medio purpureus. Park. Par. 87. No. 3.*
- Non vidi, ex l. c. solum enumeravi.
- * 5 *crocco-cincta*. H. (the dwarf saffron rimmed) dodrantalis: foliis 4-5 spathulatis, corolla parva solitaria alba, corona lutea, margine crocco, radice parva rotunda sub-oblonga tunicis exterioribus nigricantibus.
- Narcissus medio croceus tenuifolius. Park. Par. p. 87. No. 2.*
- N. medio croceus graminifolius. *Rudb. Elys. t. 61. 5.*
- Obs.* non vidi, et l. c. solum enumeraui. Haec et ultima (secund. Park. l. c.) in habitu, simulant N. poeticum, in fere omnibus, at multoties minoribus.
- **** *Floribus omnino albis, foliis glaucis.*
- * 6 *pumila*. H. (the dwarf white) uniflora: foliis linearibus (valde glaucis): corollæ laciniis ovato-subrotundis (albis), nectario concolore crasse crenato, triplo longioribus, antheris 3, tubo inclusis, 3-exsertis. *Redout. Lill. 409. Ganymedis species?*

XVI. NARCISSUS.

Spatha 1-3-flora. *Corona* parva patellari, *corolla* sæpius nivea laciniis multoties brevior toto crocea seu flava et croceo fere semper cincta. *Filamenta* tubo adnata 3-inferiora cum antheris, parum inclusa; et 3, etiam inclusa cum antheris semisexsertis.

* Præcociores.

1 *poetarum*. N. (saffron-capped) corollæ laciniis amplissimis cuneato-ovobatis planis valde imbricatis niveis, corona sub-patellari (mox subdiscoidea) primo toto crocea, ore plicatissimè criso saturatiore.

Narcissus maximus medio purpureus.

Merian. florileg. t. 144. fig. infer.

Narcissus poetarum.

—croceum pro corpore florem Inveniunt, foliis medium cingentibus albis. *Ovid.*

Obs. *Corolla* nivea trientalis. *Corona* primo omnino pulcherrimè crocea, mox a basi seorsum intènse lutea, denique albicante periente. *Folia* glauca infernè carinata 7 lineas lata apice flaccide recurva.

Florebat i. Apr. 1831, in Nobili Horto, Hort. Soc. Londini, copiosè et pulcherrimè omnium.

Obs. Non est Linnaei *Narcissus poeticus*, nec Salshuri, neque aliorum Botanicorum.

2 *ornatus*. N. (the flat crowned saffron rim) corollæ laciniis (præcipue exterioribus) subreflexentibus, basi imbricatis; corona mature omnino complanatim expansa: foliis erectis perglancis angustis.

N. poeticus. Salisb. in Hort. Trans.

1. 363, sine ullo charaktere. N. medio purpureo præcox. *Park. Par. p. 76. tab. 75. f. 3. N. tripodalis. Salisb. MSS.* ex laciniis 3 sepe semireflexis (tripodam quasi simulantibus, et inde ex errore N. tripodalis. *Schult. syst. v. 7. p. 987.*)

Floret initio Aprilis cum sequente.

3 *angustifolius*. N. (narrow leaved saffron-rim) corollæ laciniis horizontalibus spatulato obovatis distinctis, corona

- acetabuliformi ore scarioso crenulatissimo internè coccineo, antheris 3 inclusis.
- N. angustifolius.* *Bot. Mag.* 193. N. radiiflorus. *Salisb. Prod.* p. 225. et in *Hort. Trans.* v. 1. p. 365, excluso synon. *Parkinsoni*, quod est *N. stellaris* *infra*. *Narciss. Revis.* p. 140. *Habitat* Helvetia pratis humidis subalpinis. *Floret* i. Aprilis. *Variat.* a. ut supra, foliis angustè loratis glaucis. β. foliis angustis viridibus. Vidi apud Messrs. Young Epsomæ. γ. foliis glaucis floreque majoribus. Communicavit amicus Johannes Denson, Junior. ** Medio tempore florescentes.
- 4 *spathulatus*. N. (the lesser saffron rim) corollæ minoris, laciniis obtusis seu spathulatis, corona lutea, margine plicato criso croceo. In nobili Horto, Hort. Soc. vidi florēt, medio seu fine Aprilis. An varietas minor certè præcocior *N. patellaris* *infra*?
- 5 *albus*. N. (the slightly saffron rimmed) corollæ mediocris niveæ, laciniis exterioribus obovatis, interioribus subovatis valde imbricatis, corona patula lutea, margine plicato-crispo subcroceo, mox marcescente albido, multo longioribus. N. albus. *Mill. Dic. ed. 8. No. 5. Schult. syst. v. 7. p. 985.* Florebat in Nobili Hort. Hortic. Soc. Londini, med. Aprilis, 1831.
- 6 *dianthus*. N. (the two-flowered saffron rim) 2-florus : corollæ laciniis subovatis niveis flexuosis, (per latera recurva) basi distinctis substellatis, corona acetabuliformi aurantia valde plicatum lobata crispa, lobis valde imbricatis plicato repandulis. Obs. *N. biflorus* *infra*, similis sed discrepat *caule* sesquipedali, corollæ laciniis angustioribus niveis, coronā aurantia et foliis subglaucis carinatis basi, oris inflexis et inde cavo-triangularibus, 8-lineas latis. Florebat in nobili Horto, Hort. Soc. Lond. medio Aprilis, 1831.
- 7 *triflorus*. N. (three flowered) 2-3, rarissimè 4-florus : corollæ laciniis niveis ovatis planis, mox semireflexis imbricatis, corona patellari perlutea plicatula. Florebat cum priore. Obs. *Foliū* fere precedentis sed viridia lorata carinata 5-6 lineas lata acuta. Differt etiam corollæ laciniis non flexuosestellatis sed planioribus et imbricatis, corona lutea nec aurantia.
- 8 *biflorus* N. (the cream-coloured) 1-3 florus : corollæ ochroleucæ mox albidae, la-
- ciniis amplissimis præinæqualibus late ovatis, corona patellari lutea, margine marcescente pallido, plicatulatim crenulato.
- N. biflorus. Narciss. Revis.* 151. *Bot. Mag.* 197. Obs. This plant is reputed to be a Mule, because in gardens it bears not even the embryos of seeds : but as it spreads over whole fields in a wild state in Devonshire, &c. may it not, like the Hautboy Strawberry, have both fertile and barren individuals proper to its species ; those now in gardens having arisen from the barren ones ?
- *** *Seriōres florendi.*
- 9 *recurvus*. N. (the drooping leaved saffron rim) uniflorus : foliis semunciam latis perglauca supernè debilitè recurvo-dependentibus scapo humilioribus, corollæ niveæ, laciniis imbricatis late ovatis retusis cum mucrone, lateribus inflexis : corona præplicata, ore crenulato croceo. *Narciss. Revis.* p. 151. *Floret* i. Maio, post medium.
- Obs. Flores 1 rarissimè 2 in eadem spatha. Coronæ margo sublatè croceus demum interne parum albidus, sed minus quam in sequente.
- β *gracilior*, foliis scapoque fere duplo angustioribus glaucescentibus minus planis, paucis solum recurvis sed erectis inde scapi sèpius altitudine : coronæ croceo margine sèpe tenuiori.
- 10 *poeticus*. Lin. N. (middle-sized May) foliis erecto-subincutantibus glaucis, 4 lineas latis, corollæ laciniis imbricatis lateribus deflexis, corona patellari lutea, minutissime plicatula margine subscariose circulo albo-gracili, externè coccineo, crenulato ; stigmate longitudine staminum interiorum. N. poeticus. Linn. sp. pl. ed. 3. p. et ejus *Herbarii*. N. majalis. *Curtis Bot. Mag.* 193 at the end. *Narciss. Revis.* 150. β flor. plen. albo.
- γ omnibus antheris exsertis. N. poeticus. Red. *Lill. t. 260.*
- 11 *patellaris* N. (the large broad May) corollæ amplissimæ laciniis orbiculato-lobovatis imbricatissimis niveis, oris deflexis, corona patellari lutea supernè alba ore denticulato-crispo croceo ; foliis 8 lineas latis, glaucis carinatis. N. poeticus. Eng. *Bot. tab.* 275. *Narcis. purpureus maximus. Park. Par.* 75. f. 2.
- β flor. pleno albo cum croceo.
- Obs. Corollæ laciniæ 3, exteriores subsemireflexæ, 3-interiores tubo respectu horizontales.

12 *stellaris*. N. (the long petalled saffron rim)
corollæ niveæ elongatæ stellaris, la-
ciniis obovato cuneatis mucronatis tor-
tis oris undatis reflexis distinctis, co-
rona perlutea patellari, margine pli-
cato crenulato croceo mox intus albido
multoties longioribus, antheris omni-
bus sub-exsertis.
Narcissus medio purpureus serotinus.
Park. Parad. tab. 76. f. 2.

Communicavit amicus Dom. Ander-
son, ex Hort. Chels. Initio Maii flo-
ribusque post medium.

Obs. corollæ laciniæ tubum 3-lineas
longiores. *Scapus* gracilis anceps.
Folia lorata viridia sive aliquantum
glaucescentia. *Antheræ* tres paulo
exsertæ, tres semiexsertæ sed cæ-
teris certe humiliores.

POSTSCRIPT.

For the liberal use of books, and for the presentation both of many Roots and Specimens, without which the foregoing Enumeration of Narcissinæan Plants could not have been completed, I beg leave very much to thank the following Royal and Public Gardens, Gentlemen, and Nurserymen.

The Royal Gardens of Kew.

The Royal Gardens of Berlin.

The Horticultural Garden at Chiswick.

The Botanic Garden at Bury St. Edmund's.

Chelsea Physic Garden.

The Rev. W. T. Bree, Rector of Allesley, near Coventry.

The Rev. H. T. Ellicomb, Bitton, near Bath.

Aylmer Bourke Lambert, Esq. for the use of his Library.

Messrs. David & George Don, for shewing me the Linnean Herbarium.

Messrs. Loddiges, of Hackney.

Messrs. Whitley, Brames, and Milne, Fulham.

Messrs. Young, Epsom, Surry.

Messrs. Allen, and Co., King's Road, Chelsea.

Mr. Colvill, King's Road, Chelsea.

Mr. Knight, King's Road, Chelsea.

Mr. Tate, Sloane Street, Chelsea.

Mr. R. Sweet, Cook's Ground, Chelsea, for roots, and valuable remarks.

Mr. John Denson, junior, late Curator of the Bury Botanic Garden.

Mr. William Pamplin, Lavender Hill Nursery, Wandsworth Road.

In the course of the above Enumeration, will be found all the Species and Varieties contained in the just published Seventh Volume of Schulte's celebrated edition of Linnæus's *Systema Vegetabilium*, and all, as far as practicable, duly appreciated; nearly all the Species and Varieties of the faithful Parkinson, are likewise given; a few from the less accurate *Hortus Eystettensis*; the rare *Campi Elysii* of Rudbeck; Barrellier's *Icones*; the *Florilegium of Swertiae*; Trew's *Floræ Imagines*; the *Theatrum Floræ*; Madam Merian's *Icones*; and many other works.

I have carefully preserved Specimens of almost the whole; and they will be hereafter my Vouchers for the truth.

The Genera and Species of this group of Plants here, and heretofore proposed by me, are very natural; and though many years have elapsed without their having been adopted by others.* So much the better it is, for my future reputation; for adopted they will be; and then must be proved, my having seen *just so far, before* my competitors.

On the other hand, half a century's experience constantly amongst the living Plants, might be expected to have effected *more*. But, "non omnia, possumus omnes."

The characters I have given throughout, are *short*, but they are the result of *experience*, founded on the solid laws of botanical proportions; and I know their *validity*.

" His L.B. utere mecum; dum alii plura melioraque sistant."

A. H. HAWORTH.

Chelsea, 1831.

* Since the first edition was written, my acute friend, Mr. Sweet, has adopted them all, and published some in his beautiful British Flower Garden.