

# The MUSORSTOM-TDSB deep-sea benthos exploration programme (1976-2012): An overview of crustacean discoveries and new perspectives on deep-sea zoology and biogeography

Bertrand Richer de Forges<sup>(1)</sup>, Tin-Yam Chan<sup>(2)</sup>, Laure Corbari<sup>(3)</sup>, Rafael Lemaitre<sup>(4)</sup>,  
Enrique Macpherson<sup>(5)</sup>, Shane T. Ahyong<sup>(6)</sup> & Peter K. L. Ng<sup>(7)</sup>

<sup>(1)</sup> Kiwa Consulting - 5, rue Félix Frachette - 98800 Nouméa - New Caledonia  
b.richerdeforges@gmail.com

<sup>(2)</sup> Institute of Marine Biology and Center of Excellence for the Oceans  
National Taiwan Ocean University, Keelung 20224 - Taiwan, R.O.C  
tychan@mail.ntou.edu.tw

<sup>(3)</sup> Muséum national d'Histoire naturelle - Département Systématique & Évolution - CP 26 - 43, rue Cuvier - 75005 Paris, France  
corbari@mnhn.fr

<sup>(4)</sup> Smithsonian Institution - National Museum of Natural History - Department of Invertebrate Zoology  
MRC 4210 Silver Hill Road, Suitland, MD 20746, USA  
lemaitr@si.edu

<sup>(5)</sup> Centro de Estudios Avanzados de Blanes (CEAB-CSIC) - C. Acces Cala Sant Francesc 14 - 17300 Blanes, Girona, Spain  
macpherson@ceab.csic.es

<sup>(6)</sup> Australian Museum - 6 College St. - Sydney, NSW 2010, Australia  
Shane.Ahyong@austmus.gov.au

<sup>(7)</sup> Tropical Marine Science Institute and Department of Biological Sciences - National University of Singapore  
Kent Ridge - Singapore 119260, Republic of Singapore  
peterng@nus.edu.sg


## INTRODUCTION

In 1976, the French research vessel “Vauban” was scheduled to travel from France to New Caledonia. Through the efforts of Alain Crosnier (Director of the Oceanography, ORSTOM), it was decided that the craft take a detour through the Philippines —this became the first MUSORSTOM cruise. The principal objective of this cruise was to find new

RICHER DE FORGES B., CHAN T.-Y., CORBARI L., LEMAITRE R., MACPHERSON E., AHYONG S. T. & NG P.K.L. — The Musorstom-TDSB deep-sea benthos exploration program (1976-2012): An overview of crustacean discoveries and new perspectives on deep-sea zoology and biogeography, in AHYONG S. T., CHAN T.-Y., CORBARI L. & NG P.K.L. (eds), *Tropical Deep-Sea Benthos 27*. Muséum national d'Histoire naturelle, Paris: 13-66 (Mémoires du Muséum national d'Histoire naturelle; 204). ISBN: 978-2-85653-692-6.

specimens of the living fossil glypheid *Neoglyphea inopinata* described one year earlier by Jacques Forest and Michèle de Saint Laurent from a single specimen collected by the U.S. Fish Commission Steamer research vessel "Albatross" in the beginning of the 20th century (Forest 1981, 2006; Richer de Forges & Justine 2006; Bouchet *et al.* 2008). This first French deep-sea cruise in the Indo-West Pacific was a great success for two reasons. Firstly, nine fresh specimens of *Neoglyphea inopinata* were collected near Lubang island in a very small area (Figure 1A). The detailed description of this only extant species of glypheid contributed significantly to our knowledge on the evolution of decapod crustaceans (Schram & Ahyong 2002). Secondly, the trawls in the Philippine archipelago also collected a very rich benthic fauna including many rare and new species, demonstrating that the exploration of the deep-sea fauna was far from complete.

In the last quarter of the 20th century, our knowledge of the deep-sea fauna was mainly based on the results of the so-called "Great Expeditions" of preceding decades. These expeditions, which took place in the era between the CHALLENGER in 1873 and the GALATHEA in 1952, sampled the oceans on a world-wide scale. They studied and mapped the geography and bathymetry of the sea, and brought up a totally new fauna, resulting in the re-writing of marine zoology (Thomson 1878; Bruun 1953). However, despite their many significant discoveries, these expeditions only sampled about 1000 deep-sea stations across the planet. Considering the size of Earth's oceans, huge areas remained totally unexplored.



**FIGURE 1**

The species of extant glypheids were considered as "living fossils" supposedly extinct 50 million years ago. **A**, *Neoglyphea inopinata*, male, cl 120 mm, Philippines (photograph J. Forest). **B**, *Laurentaeglyphaea neocaledonica*, female, cl 60.0 mm, Chesterfield Island (photograph J. Lai).





FIGURE 2

Updated map of the MUSORSTOM-TDSB cruises.


This first MUSORSTOM cruise in 1976 was to be the starting point of a long series of deep-sea cruises in the Indo-Pacific (Figure 2). Under the leadership of Alain Crosnier, a modern pioneer in deep-sea biodiversity research, these cruises rewrote the annals of deep-sea biodiversity. The saga of the MUSORSTOM cruises is also intricately linked to the late Jacques Forest and Michèle de Saint Laurent. Their discovery of, and search for, *Neoglyphea* was the catalyst that started the whole exploration process. To appreciate the significance of these cruises and explorations, one must remember that they took place 10 years before the invention of the word “biodiversity” and 30 years before the “CENSUS OF MARINE LIFE” (CoML) began. With the publication of the 27th volume of scientific results of the MUSORSTOM-TDSB cruises in 2013, after 37 years of exploration across the Indo-West Pacific, it is timely for a preliminary assessment of its impact. As noted above, the cruises first started in 1976. As these were guided and inspired by carcinologists like Crosnier, Forest and de Saint Laurent, it is not surprising that a “carcinological slant” continued over the years. Of all the animals collected by these cruises, the best studied and best reported on were the crustaceans. Some 33% of all papers resulting from these cruises were on these crustaceans (Figure 3). It is thus appropriate to summarize how our increased knowledge of crustaceans resulting from the MUSORSTOM-TDSB program contributed to knowledge of deep-sea biology.

#### CONTRIBUTIONS OF THE TDSB PROGRAMME TO ZOOLOGY

The programme of Tropical Deep-Sea Benthos (TDSB), formerly MUSORSTOM, is ongoing and very ambitious: to describe the deep-sea fauna of the tropical Indo-Pacific. The first three cruises in the Philippines (1976, 1980, 1985) mainly devoted to the catch of *Neoglyphea* showed that even the deep-sea fauna from the area sampled by the U.S. Fish Commission Steamer research vessel “Albatross”, was still poorly known. The question then was: what about the other archipelagoes never sampled because they stayed outside of the cruise track of the “Great Expeditions”? What started in 1976 with the research vessel “Vauban” was soon expanded to other parts of the western Pacific (Richer de Forges 1990). Between 1984 to 1994, 12 cruises totally devoted to the zoological exploration (after 1988 using the R.V. “Alis”), 7 cruises targeting chemical products in the deep-sea animals and 3 cruises focused on resources were realized in the New Caledonia EEZ (Exclusive Economic Zone). These cruises captured a huge amount of deep-sea material. From this

**FIGURE 3**

Publications from the MUSORSTOM-TDSB programme between 1977 and 2012: total number 1411 of which 468 (33 %) concern Crustacea (grey bars).

**FIGURE 4**

Histogram from CoML showing the species discovered during the 10 years of the program. Crustaceans dominate (after Ausubel et al. 2010: fig. 12).


material, deposited primarily in Muséum national d'Histoire naturelle, actively managed by Alain Crosnier for many years, and later by Philippe Bouchet, more than 1400 scientific articles were published, totally renewing knowledge of the Tropical Deep-Sea Benthos (Figure 3). For New Caledonia alone, of the 1619 species reported in 2000, 60.7 % were new to science (Richer de Forges *et al.* 2000, 2005). Since 1994 other areas have been extensively sampled: Wallis & Futuna, Fiji, Tonga, Marquesas, Australes, Vanuatu, Solomons, Indonesia, Papua New Guinea, Taiwan, Philippines, Mozambique and Madagascar. The collections have covered the Central Pacific, the Melanesian Arc in the Southern Hemisphere, part of the Southeast Asia in the Northern Hemisphere targeting the "Coral Triangle" and in the Indian Ocean. It has made possible broad geographical taxonomic and systematic revisions, and for the first time, resulted in well grounded hypothesis about the origin and evolution of the deep-sea fauna.

Among the deep-sea fauna, crustaceans are dominant across all depths, sometimes with high diversity. The TDSB programme overlapped and contributed significantly to the CoML initiative. Ausubel *et al.* (2010) observed that crustaceans dominated the number of new species found during the decade of the CoML, with 452 species named (Figure 4) — some 20% of all discovered species. Molluscs and fish came out as second and third in diversity. Much of this crustacean diversity was reported by the carcinologists working on the TDSB collections. Below, we provide a summary of some of the accomplishments of the TDSB programme.

## DISCOVERY SUMMARY AND HIGHLIGHTS OF KEY CRUSTACEAN GROUPS

**TRUE CRABS (BRACHYURA)** — The brachyuran crabs are the most diverse of all decapods, with half the known taxa (Ng *et al.* 2008; De Grave *et al.* 2009) and over 7000 species. The MUSORSTOM-TDSB collections have been invaluable in the discovery of hundreds of species, not just in obtaining specimens of new species but also providing valuable comparative material to resolve the taxonomy of many others. The collections from the first few expeditions were first reported on by Serène & Vadon (1981) but the few new species described belies this paper's significance as it was just a preliminary study. A series of papers by Guinot and others demonstrated that the deep-sea brachyuran fauna was not just rich, but extremely rich (e.g., see Guinot & Richer de Forges 1981a, b). This was not restricted to the Philippines but in waters off New Caledonia (see Richer de Forges & Ng 2006) and even Taiwan (e.g., see Ho *et al.* 2004). The expedition material has been the basis of a large number of studies in the following families: Atelecyclidae (Guinot 1989b); Cancridae (Davie 1991), Calappidae and Leucosiidae (Chen 1989, 1990; Tan & Ng 1993, 1995; Galil 1997, 2003; Galil & Ng 2007, 2009, 2010; Ng & Richer de Forges 2007), Dorippidae (Chen 1986, 1993), Dromiidae (McLay 1993, 2001; McLay & Ng 2005, 2007), Ethusidae (Chen 1997, 2000; Ng & Ho 2003; Castro 2005; Naruse *et al.* 2009), Euryplacidae (Ng & Castro 2007; Castro & Ng 2010a), Gonoplacidae (Guinot 1989a, 1990; Castro 2007, 2009, 2013; Castro & Ng 2010b), Latreilliidae (Castro *et al.* 2003), various families in the Majoidea (Guinot & Richer de Forges, 1982a, b, 1986; Tavares 1991a; Poupin 1995; Ahyong & Ng 2007; Richer de Forges & Ng 2007b, 2008a, 2009a, b, 2012, 2013; Richer de Forges 2010; Richer de Forges & Corbari 2012; Ng & Richer de Forges 2013), Mathildellidae (Crosnier & Ng 2004), Parthenopidae (Tan & Ng 2007a, b; McLay & Tan 2009), Pilumnidae (Chia & Ng 1995, 2000; Komai & Motoh 2012), Plagusiidae (Crosnier 2001), Pinnotheridae (Ahyong & Ng 2008), Portunidae (Moosa 1996; Crosnier 2002; Davie & Crosnier 2006; Nguyen & Ng 2010), Progeryonidae (Castro & Ng 2008), Trapeziidae (Castro *et al.* 2004), Trichopeltarionidae (Tavares & Cleva 2010), and Xanthidae (Davie 1993, 1997; Ng & Chia 1994; Ng & Naruse 2007b; Manuel-Santos & Ng 2007; Mendoza & Ng 2008a, b, 2010; Mendoza 2012; Mendoza & Manuel-Santos 2012).

Among the Brachyura, members of the family Homolidae are interesting, with some deep-sea taxa even displaying what seems like "a human face" on the dorsal shell (e.g., *Lamoha personata*, Figure 5A). Almost all are deep-water genera, with only the genus *Latreillopsis* having some representatives in waters shallower than 100 m. It took five years just to gather the few specimens collected by the "Great Expeditions" before they could be compared with the MUSORSTOM-TDSB collections. Some new taxa from the study were written up (Guinot & Richer de Forges 1981c) but a world-wide revision of the family was eventually published in the MUSORSTOM series (Guinot & Richer de Forges 1995). Before 1995, the family contained 14 genera and some 40 species. Their 1995 revision added four new genera and 17 new species! This group shows a very special shape of



the 5th ambulatory legs, forming a forceps-like structure to carry other animals like sponge for camouflage. The morphological diversity of this supplementary structure was commented in relation with the behavior of these animals (Guinot & Richer de Forges 1995). The mysterious pigmented spot situated on the chelae of some species of *Lamoha* (formerly *Hypsophrys*, see Ng 1998a) and interpreted as a luminescent organ were also documented. Since this key revision, many more species were described from new cruises in the Philippines, Salomon and Vanuatu archipelago. The family now has 68 species (Ng 1998b; Ng & Chen 1999; Ng & Wang 2002; Richer de Forges & Ng 2007a, 2008b; Naruse & Richer de Forges 2010; Ng & Eldredge 2012).

Associated with the homolids were several essentially deep-water families that many carcinologists regard as "primitive crabs". The Homolodromiidae was revised by Guinot (1993, 1995), and although no new genera were established, the affinities of this enigmatic family were substantially clarified. Even so, eight new species were described. In subsequent years, expeditions have obtained two more species from TDSB cruises (Ng & McLay 2005; Ng & Naruse 2007a). The small family Dynomenidae was revised by McLay (1999), and in the process, he named 13 species in five genera, two of which were newly described. Using new collections from the subsequent expeditions, seven more species have since been added by McLay & Ng (2004, 2005) and Ng & McLay (2010). Recently, in a major reappraisal of the systematics of the family, Guinot (2008) recognised three new subfamilies in the Dynomenidae. Perhaps one of the most exciting discoveries to French Polynesia was a new family, Poupiniidae, described by Guinot (1991). A revision of Cyclodorippidae and Cymonomidae from the Indo-West Pacific conducted by Tavares (1991b, 1992, 1993), which had previously only nine genera and 48 species, resulted in eight new genera and 21 new species described! In a follow-up study (Tavares 1998), several genera were removed into their own family, the Phyllotymolinidae Tavares, 1998. Since then, using these studies as the basis, seven other species have been described from the Indo-West Pacific as a result of new MUSORSTOM-TDSB collections (Ahyong & Ng 2009, 2011; Naruse 2013)!

From the crabs collected in the Philippines, a new superfamily was described by de Saint Laurent (1989): Retroplumoidea. Retroplumid crabs live in very muddy habitats of the deep-sea. They have a very flat carapace and with the fifth ambulatory legs greatly modified and sometimes feather-like. There are only two genera, *Retropluma* and *Bathypluma*, in the Indo-West Pacific. Description on the newly collected MUSORSTOM-TDSB material allowed detailed comparisons with the crab fossils of this group. Based on morphological characters, de Saint Laurent (1989) proposed a phylogeny of this group of crabs as separated from the Eubrachyura since the upper Cretaceous. Using a larger collection of Retroplumidae, McLay (2006) was able to provide full descriptions on these animals and added more paleogeographical comments. There are now seven species of *Retropluma* and three species of *Bathypluma*. Another poorly known group are the palicids. Studying the large Indo-Pacific collection, Castro (2000) completed a revision of the family Palicidae. There are now two subfamilies, 10 genera and 43 species, with four genera and 17 species newly described. More species were treated by Castro (2010) from additional collections. A strange crab, presumably associated with deep-water wood and debris, collected from recent expeditions to the Philippines, *Vultocinus anfractus*, proved to belong to a new family, Vultocinidae (Ng & Manuel-Santo 2007). Surprisingly, it was later also discovered in New Caledonia from among the older MUSORSTOM collections (Ng & Richer de Forges 2009).

**SHRIMPS AND LOBSTERS** — The MUSORSTOM-TDSB collection has contributed significant knowledge that has often led to a world-wide or Indo-West Pacific revision of many shrimp and lobster groups (Figures 1, 5E, 6). These include Solenoceridae (Crosnier 1978, 1985a, 1988a, 1989, 1994c; Crosnier & Dall 2004; Chan 2012), Sicyoniidae (Crosnier 2003), *Parapenaeus* (Crosnier 1985b, 2005), *Metapenaeopsis* (Crosnier 1987, 1991, 1994a, b), *Trachypenaeopsis* (Crosnier 2007), Stenopodidae (de Saint Laurent & Cleva 1981; Saito *et al.* 2009), Spongicolidae (de Saint Laurent & Cleva 1981; Komai & Saito 2006; Saito & Komai 2008), Stylodactylidae (Cleva 1990, 1997, 2004), Processidae (Noël 1985), Pasiphaeidae (Crosnier

#### FIGURE 5

Remarkable crustaceans collected by the MUSORSTOM-TDSB programme. **A**, *Lamoha personata*, New Caledonia (photograph B. Richer de Forges, carapace size about 40 x 50 mm). **B**, *Lithodes richeri*, Vanuatu (photograph B. Richer de Forges, carapace size about 70 x 100 mm). **C**, Giant ostracod *Azygocypridina brynmawria*, measuring 17 mm across, Coral Sea (photograph B. Richer de Forges). **D**, The living fossil barnacle *Waikalasma boucheti*, New Caledonia (photograph B. Richer de Forges, size about 20 mm). **E**, *Glyphocrangon panglao*, Philippines (photograph T.-Y. Chan, cl 15.0 mm). **F**, *Sympagurus brevipes*, Philippines (photograph T.-Y. Chan, size about 10 mm). **G**, *Babamunida callista*, New Caledonia (photograph B. Richer de Forges, cl 12.0 mm). **H**, *Eurysquilloides sibogae*, Philippines (photograph T.-Y. Chan, size about 50 mm).



FIGURE 6

*Dinochelus ausubeli*, male, cl 31.1 mm, Philippines (photograph T.-Y. Chan). An icon of the Census of Marine Life.

1988b; Hayashi 1999, 2004, 2006), Bathypalaemonellidae (Cleva 2001, 2004), deep-sea Pontoniinae (Bruce 1981, 1990, 1991, 1996; Li *et al.* 2008; Mitsuhashi & Chan 2006, 2009; Mitsuhashi *et al.* 2010, 2012), *Heterocarpus* (Crosnier 1988c, 1999b; Cleva & Crosnier 2006; Li 2006; Yang *et al.* 2010; Li & Chan 2013), *Plesionika* (Crosnier 1986; Chan & Crosnier 1991, 1997; Chan & Yu 1991c; Chan 2004; Komai & Chan 2010; Li & Chan 2013), *Eugonatonotus* (Chan & Yu 1991b), Nematocarcinidae (Burukovsky 2013), *Glyphocrangon* (Komai 2004, 2006b, 2007, 2011; Komai & Chan 2008, 2013), many Crangonidae genera (*Aegaeon*, *Pontocaris*, *Parapontocaris* (Chan 1996), *Parapontophilus* (Komai 2008), *Prionocrangon* (Kim & Chan 2005), *Lissosabinea* (Komai 2006a), *Metacrangon* (Komai 1997, 2012)), Glypheidae (Forest & de Saint Laurent 1981, 1989; Richer de Forges 2006; Forest 2006a, b; Boisselier-Dubayle *et al.* 2010), *Enoplometopus* (de Saint Laurent 1988; Chan & Yu 1998; Chan & Ng 2008), *Nephropsis* (Macpherson 1990b; Chan 1997), *Metanephrops* (Macpherson 1990b; Chan & Yu 1991a; Chan 1997), Thaumastochelidae (Chan & de Saint Laurent 1999; Ahyong *et al.* 2010), *Justitia* (Poupin 1994), *Palinnustus* (Chan & Yu 1995), *Puerulus* (Chan *et al.* 2013), Scyllarinae (Holthuis 2002; Yang *et al.* 2008, 2011; Yang & Chan 2010, 2012) and Polychelidae (Galil 2000; Ahyong & Galil 2006; Ahyong & Chan 2008). Altogether, 29 genera and 263 species were described from these works, and many of these reports have become the essential taxonomic references for deep-sea shrimps and lobsters. The high diversity for several classical deep-sea caridean groups such as Pasiphaeidae, *Glyphocrangon* and *Plesionika* has certainly exceeded all expectations.

Of the numerous crustacean species collected during the MUSORSTOM-TDSB programme, the most famous are certainly the glypheids. The discovery of extant species by Forest of this group, previously known only as fossils and supposedly extinct at the end of the Cenozoic era, was a major event in the taxonomy of crustaceans. The glypheids, well known from the Mezozoic, was considered to be extinct in the Eocene, 50 million years ago. They were originally placed under the infraorder Palinura and considered as probable ancestors of part of the remaining Decapoda Reptantia. However, detailed studies on the first extant species *N. inopinata* showed that the glypheids are closer to the Astacidae (Forest 2006). In 2005, the cruise EBISCO discovered a second species of glypheid (Figure 1B) living in a very different habitat in the Coral Sea (Richer de Forges 2006). This new species was so different from *Neoglyphea inopinata* that Forest decided to establish a new genus *Laurentaeglyphea* for it. This lobster-like animal, *Laurentaeglyphea neocalledonica* soon became famous under the nickname “Jurassic shrimp” (the first species discovered was called the “Phoenix lobster”, Trew Crist *et al.* 2009). The second species, *L. neocalledonica*, is even closer to those glypheids known from the Mezozoic and Eocene (Forest 2006).

Another bizarre lobster *Dinochelus ausubeli* (Figure 6), recently collected from the Philippines AURORA expedition (Ahyong *et al.* 2010) shows all the ‘monster-like’ features associated with deep-sea animals. This species was chosen as an icon of CoML and has been used by popular media to promote deep-sea biodiversity and conservation (*e.g.*, Trew Crist *et al.* 2009; Snelgrove 2010).

**ANOMURANS** — Perhaps no other group of decapods has been as substantially impacted by the MUSORSTOM-TDSB expeditions than the Paguroidea (*sensu* McLaughlin *et al.* 2007). Prior to 1976 when collecting efforts began on cruises of the VAUBAN, the extent of deep-water paguroid specimens from the Indo-West and Central Pacific available in museums, was minimal and insufficient for scientists to decipher the morphological boundaries and distributions of many species. Taxonomic research was confined mostly to studying type material, with only scant additional preserved or fresh specimens accessible. The first major work on paguroids that was produced with MUSORSTOM-TDSB material was that by Forest (1987), who expanded exponentially our knowledge of the intriguing and mysterious “symmetrical” hermit crabs of the family Pylochelidae, a presumably ancient group considered key to understanding hermit crab evolution (Tsang *et al.* 2011). Before Forest’s work, only 19 species were known, based on 60 specimens obtained at 30 stations from around the world. He added 24 new species and subspecies using 400 specimens from about 200 sites, primarily from MUSORSTOM-TDSB cruises, and complemented by specimens from earlier deep-sea expeditions on the ALBATROSS. Subsequent MUSORSTOM-TDSB and other French cruises obtained considerably more material which was used to further increase our knowledge of the diversity of pylochelids, resulting in revisions of their classification and phylogeny (McLaughlin & Lemaitre 2008; Lemaitre *et al.* 2009; McLaughlin *et al.* 2009).

Among the hermit crabs of the family Diogenidae, the genus *Trizopagurus* benefitted greatly from a major revision by Forest (1995) of their taxonomy and morphology, largely based on specimens obtained during MUSORSTOM-TDSB. This genus contained only 10 species before the study by Forest, who expanded it to contain (in addition to *Trizopagurus*) two new genera (*Ciliopagurus* Forest, 1995, *Strigopagurus* Forest, 1995) and 24 species, 14 of them new. Particularly fascinating are the stridulatory structures in these three genera, the morphology of which Forest (1995) described for the first time in great detail. He also discussed their anatomy in an evolutionary context. From the extraordinary collections in Polynesia, another new *Ciliopagurus* was described by Poupin (2001), as well as new species and colour data of *Calcinus* species (Poupin & McLaughlin 1998; Poupin & Lemaitre 2003). The taxonomy, diversity and morphology of what is probably the most speciose and complex diogenid (indeed paguroid) genus, *Paguristes*, and other taxonomically difficult diogenid genera, were significantly improved with a number of important taxonomic and morphological studies by Rahayu (2005, 2006, 2007, 2008, 2010) and Rahayu & Forest (2009). Even a new species of burrowing barnacle of the genus *Tomlinsonia* that is a parasite on three species of *Calcinus* was documented by Williams & Boyko (2006).

A seminal paper on the Paguridae from Indonesian waters was published by McLaughlin (1997). That paper was based on specimens obtained during the KARUBAR expedition, and revealed an unexpected wealth of hermit crabs of this family in the Indo-West Pacific. Seven new genera and 26 new species were discovered. Previous to McLaughlin’s paper, most hermit crabs inhabiting the tropical environs of the Indo-West Pacific were thought to be species of Diogenidae. In fact, McLaughlin’s paper showed that the Paguridae are as rich or even richer in this region of the world oceans than the Diogenidae, and opened the door to a complete re-evaluation of the Paguroidea in general from the Indian and Pacific Oceans or indeed the world over. Whereas brachyurans are certainly the most diverse in number of species among the Decapoda, the anomurans, and in particular the Paguroidea, are the most diverse in body shape and morphology. Additional discoveries based on MUSORSTOM-TDSB include: a new species of *Xylopagurus* A. Milne-Edwards, 1880, a genus previously known only from the western Atlantic and eastern Pacific (Forest 1997); several more new genera and new species (Lemaitre & McLaughlin 1995; McLaughlin & Forest 1997); new species of *Porcellanopagurus* and *Solitariopagurus*, two genera that include remarkably carcinized forms (McLaughlin 2000); a remarkable new genus and species with reduced corneas (Lemaitre 2003); two astounding and minute new species of a new genus, *Pteropagurus* McLaughlin & Rahayu, 2006, found living in empty planktonic pteropod shells accumulated on the seafloor in a very unusual hermit crab-mollusc association; new species or records, and taxonomic clarifications of species of *Cestopagurus*,

*Pagurixus*, *Pseudopagurus* (Rahayu & Komai 2013) and *Pagurus* (Komai & Chan 2006; Komai 2010; Komai & Poupin 2012); and a new monotypic genus with a dwarf new species, *Pumilopagurus tuberculomanus* McLaughlin & Rahayu, 2008b. In addition, a major taxonomic and morphological review of *Nematopagurus*, a genus with 26 species distributed primarily in the Indo-West Pacific, was published by McLaughlin (2004b).

Within the paguroids, no other family has been impacted as much by the rich collections from the MUSORSTOM-TDSB projects, than the deep-water hermit crabs of the family Parapaguridae. In the last 40 years, primarily because of the wealth of deep-water material collected by French expeditions, the number of species has increased from 31 to 86, and the number of genera from 4 to 10, an increase of 277% and 250%, respectively (Figure 5F). Numerous parapagurid specimens were collected from many deep-water habitats and regions that had never been sampled before. Before 1986, when French expeditions using the S.M.C.B. "Marara" began to collect in French Polynesia including the Marquesas, there were no specimens or records of parapagurids from that immense region of the central South Pacific. Works on those collections resulted in the description of several remarkable new species, and more importantly a better picture of the anomurans from that region emerged. The Polynesian parapagurids as well as other paguroids and decapods, were summarized in a number of catalogues or atlases, including fisheries data and many color photographs (Poupin *et al.* 1990; Poupin, 1996a, b). Furthermore, a database of the species of decapods (shrimps, lobsters, crabs, hermit crabs) and stomatopods (mantis shrimps), recorded from the French Polynesian Islands (Marquesas, Society, Tuamotu, Austral, and Gambier), Pitcairn, Easter Island and Clipperton, is available through a free database in the internet (<http://decapoda.free.fr/>). The numerous studies on the taxonomy and systematics of parapagurids based on specimens from all these French-sponsored faunistic studies, greatly expanded and clarified the world-wide picture of diversity and biogeography of the parapagurids, an evolutionary key group of hermit crabs. Taxonomic and systematic information of this family can be found in a number of convenient, broad generic reviews or revisions (Lemaitre 1999, 2004a, b, 2013), as well as in a number of papers reporting on parapagurids from particular regions and describing new species, or discussing morphological and biological aspects (Lemaitre 1993, 1994, 1997, 1998, 2004a, b, 2006; Lemaitre & Poupin 2003).

The specialized burrowing crabs of the superfamily Hippoidea from the tropical Indo-West Pacific region had been the subject of few systematic studies until many specimens of this poorly known group of crabs became available from collections obtained by MUSORSTOM-TDSB. As a result, three landmark and detailed studies on the systematics of the families Albuneidae and Hippidae, including descriptions of several new species, have been published (Boyko & Harvey 1999; Boyko 2000, 2002). In those papers, the diagnostic morphology of the group was radically redefined and deciphered to the benefit of future studies on their biology and evolution.

Squat lobsters are a major component of the deep-sea crustacean fauna (Baba *et al.* 2008). They are divided into two main groups. The Galatheoidea (with four families Galatheidae, Munididae, Munidopsidae and Porcellanidae) has a large number of free-living species living on the ocean bottom, and generally have numerous eggs with planktonic larvae. The Chirostyloidea (with three families Chirostylidae, Eumunididae and Kiwaidae) are often associated with other benthic invertebrates like sponges, antipatharians or gorgonians, and produce few eggs with advanced larvae and limited dispersion. The abundant galatheoid material obtained in the MUSORSTOM-TDSB collection has resulted in intensive taxonomic studies on these animals that have revolutionised our understanding of their diversity. After the TDSB studies initiated in 1976, these families have been enriched by 13 new genera (Baba 1991, 2004; Baba & de Saint Laurent 1996; Macpherson 1998, 2006; Macpherson & Machordom 2000; Cabezas *et al.* 2008). The total number of known species has increased nearly 2.5 fold since 1976, with 937 taxa today (Baba *et al.* 2008; Appeltans *et al.* 2012). Most of the newly described species have been based on the MUSORSTOM-TDSB material. Some of them living in the upper part of the bathyal zone (200–600 m) can be surprisingly colourful, although some live colours and patterns have to be seen to be believed (Figure 5G).

The first contribution towards galatheoid systematics using these collections was published by Baba (1990) on specimens collected by Alain Crosnier on board of the R.V. "Vauban", in Madagascar. Subsequently, Baba (1991a, b), using material from some expeditions carried out around New Caledonia, described new genera and species of chirostylids and galatheids of the genera *Alainius*, *Leiogalathea*, *Phylladiorhynchus*, *Gastroptychus* and *Chirostylus*. These papers,

together with the review of the genus *Eumunida* (de Saint Laurent & Macpherson 1990a) along the Indian and Pacific Oceans, were the starting point of the many subsequent studies of the group. The genus *Eumunida* was revised again by de Saint Laurent & Poupin (1996), adding numerous specimens from the MUSORSTOM expeditions, and describing six new species from the Indian and Pacific Oceans. Baba & de Saint Laurent (1996) described some new genera of small galatheids (*Anoplонida*, *Neonida*, *Plesionida*, among others), closely related to the genus *Bathymunida*. Furthermore, they divide the galatheids in two large groups (*i.e.* having one or two pairs of male gonopods) and reorganizing the taxonomy of the family. The phylogenetic value of this division was later validated by molecular tools (Machordom & Macpherson 2004).

Baba (2004) erected the genus *Uroptychodes* from *Uroptychus*, using material collected in New Caledonia, Indonesia and adjacent waters. An additional study on the genus *Uroptychus* from Taiwan was published by Baba & Lin (2008), Cabezas *et al.* (2012), and a large and complete revision of the genus including more than 100 species will be published soon (Baba in preparation). This major taxonomic revision of the Chirostyliidae (Baba in preparation) will also form the basis for new hypotheses on seamount endemism.

Macpherson (1993a, b) and Macpherson & Baba (1993) studied the material collected in the early MUSORSTOM-TDSB cruises carried out in the Philippines and Indonesia, adding numerous new species belonging to the genera *Agononida*, *Munida* and *Paramunida*. In addition, they solved the confusing status of *Munida japonica* and their relatives. In 1994, a study on the collections obtained mostly in New Caledonia, Chesterfield, Matthew and Hunter Islands, demonstrated the existence of 56 new species of *Agononida* and *Munida* (Macpherson 1994). The existence of a very rich fauna in the western Pacific was also supported by the results obtained in MUSORSTOM-TDSB expeditions from Wallis and Futuna (Macpherson 1996), Kei Islands (Indonesia) (Macpherson 1997), Vanuatu (Macpherson 1999a), Marquesas Islands (Macpherson 2000), Fiji and Tonga (Macpherson 2004), Austral Islands (Macpherson 2006a), Solomon Islands (Cabezas *et al.* 2009) and Taiwan (Lin *et al.* 2004, 2005; Cabezas *et al.* 2011b). Other studies using specimens from recent expeditions are completing our knowledge of the group along the continental shelf and slope of the western and central Pacific, and eastern Indian Ocean (Macpherson 1999b, 2006b, 2009; Macpherson & Baba 2006).

The deep-sea fauna, mostly belonging to the genera *Munidopsis* and *Galacantha*, was also studied using numerous specimens collected in all MUSORSTOM-TDSB cruises, from Madagascar to French Polynesia (Baba & de Saint Laurent 1992, Macpherson 2007, 2012), adding more than 25 new species. Furthermore, numerous new species have been collected in recent expeditions along the coasts of Taiwan and the Philippines (Lin *et al.* 2007, 2011; Osawa *et al.* 2006a, b, 2007, 2008, 2013).

The shallow-water genera (*e.g.*, *Allogalathea*, *Galathea*, *Lauriea*, *Sadayoshia*) have also been enriched by the MUSORSTOM-TDSB expeditions confirming the existence of a highly diverse fauna with similar patterns than in the continental shelf, slope and abyssal basins (Macpherson & Baba 2010; Macpherson & Cleva 2010; Cabezas *et al.* 2011), and a major revision of the diverse genus *Galathea* is in preparation (Macpherson in preparation).


The results obtained from MUSORSTOM-TDSB programme have demonstrated that squat lobsters are abundant, speciose and widely distributed. They occur in all marine habitats at depths ranging from the surface of the sea to more than 5000 m (Baba *et al.* 2008). Squat lobsters are most diverse in the western Pacific area and, at present, the most speciose genera are *Munida* and *Munidopsis*. The highest numbers of squat lobsters are found in the Coral Sea (Solomon-Vanuatu-New Caledonia), and Indo-Malay-Philippines Archipelago (Macpherson *et al.* 2010; Osawa *et al.* 2013). Most species of *Munida* are from shelf and slope depths, whereas those of *Munidopsis* and *Galacantha* are found mainly on continental slopes and the continental rise zones. The distribution of endemism along the Pacific Ocean indicates that there are several major centers of diversity, for example in the Coral Sea, Indo-Malay-Philippines Archipelago and French Polynesia.

The squat lobsters are becoming one of the best-known groups of deep-sea crustaceans. As a summary of the work that started in 1976, the different authors have produced many important papers that have formed the basis for other major, synthetic works. For instance, studies on the fauna of some basins, *e.g.*, Taiwan (Baba *et al.* 2009), the checklist of squat lobsters of the world (Baba *et al.* 2008), and a book on the biology of squat lobsters (Poore *et al.* 2011).

The king crabs (family Lithodidae) are also well represented in MUSORSTOM-TDSB collections (Figure 5B). The first contribution using specimens collected by Alain Crosnier in Madagascar was published by Macpherson (1988). Other studies covering the area of New Caledonia and French Polynesia were published by Macpherson (1990) and (1991), respectively. The hydrothermal vents also provide some interesting new species (de Saint Laurent & Macpherson 1997), and the most recent MUSORSTOM-TDSB expeditions collected additional material, including interesting new occurrences and species from New Caledonia, Vanuatu, Solomon, Fiji and Tonga areas (Macpherson 2001, 2003). More recently, some interesting species have been described in expeditions carried out in Taiwan (Macpherson & Chan 2008; Ahyong & Chan 2010).

**OTHER REMARKABLE CRUSTACEANS** — Not all the work done on crustaceans during the MUSORSTOM-TDSB exercise was on decapods. From the crustaceans collected in the deep-sea some were remarkable by their giant size such as the isopods *Bathynomus* and ostracod *Azygocypridina*. Other than “living fossils” like the glypheids *Neoglyphea inopinata* and *Laurentaeglyphea neocalaledonica*, the ancient cirriped *Waikalasma boucheti* also merits mention. These have since been used as “icons” for deep-sea biodiversity research.

*Bathynomus* is an isopod genus containing the largest known isopod species, its members living between 100 and 1000 m deep. Based mainly on the MUSORSTOM-TDSB collection, Lowry & Dempsey (2006) published a revision of this group of giant scavengers from the Indo-West Pacific. Six species are redescribed and six new species were added. The authors separated the genus into two groups, the giants and the super-giants having an adult size between 150 and 500 mm long (Figure 7).



**FIGURE 7**

A supergiant of the isopod genus *Bathynomus* — *B. kensleyi*, Philippines (photograph P. K. L. Ng). A saga of the MUSORSTOM-TDSB deep-sea expeditions.

Another giant described from the MUSORSTOM-TDSB material was the ostracod *Myodocopa* in the family Cyprinidae, *Azygocypridina brynmawria*. It was collected in the Coral Sea at about 400 m deep (Diamond, Richer de Forges & Kornicker 2008). The majority of ostracods have adult sizes of only a few millimetres, but this species has a diameter of 17 mm (Figure 5C). *Azygocypridina* ostracods possess iridescence on the first antennae. The description of this phenomenon by Andrew Parker on a species of the same genus was the starting point of the fascinating speculations on the color vision in fossil crustaceans and its role in the so called "Cambrian Explosion" of life (Parker 1998, 2003; Nielsen & Parker 2010).

Cirripeds of the genus *Waikalasma* was known only from fossils in the Miocene of New Zealand. However, during the cruise MUSORSTOM 8 in Vanuatu a living species was collected: *Waikalasma boucheti*. In 2011, the cruise EXBODI discovered another specimen of this species from New Caledonia. This genus is characterised by an extra layer of plates around the body (Figure 5D).

Although a much smaller group than the decapods, the mantis shrimps (Order Stomatopoda) are generally best known from shallow waters. Knowledge of their deep-water diversity, however, has been significantly increased through MUSORSTOM-TDSB sampling programmes, with discovery of many new species, genera and even a new family, Alainosquillidae, from New Caledonia (Moosa 1991). Further discoveries are ongoing with many more new species under study. Many uncommon species were often collected by the MUSORSTOM-TDSB cruises, such as *Bathysquilla microps*, *B. crassispinosa* and *Gonodactyloideus cracens*, and others that were until that time thought particularly rare, such as the enigmatic, *Eurysquilloides sibogae* (Figure 5H), which is now known to be common in the Philippines. Studies dealing with Stomatopoda include major studies from the Philippines (Moosa 1986; Ahyong 2010, 2013), New Caledonia (Moosa 1991; Ahyong 2007), Indonesia and the Marquesas (Ahyong 2002a, b).

## OTHER CONTRIBUTIONS OF THE MUSORSTOM-TDSB PROGRAMME

### BIOGEOGRAPHY

The science of biogeography synthesises knowledge from taxonomy, phylogeny and genetics, and helps us understand how the marine environment has evolved, the impacts of sea level variations and plate tectonics on biodiversity. All these different aspects of science contribute to reconstructing the spatio-temporal scenario explaining the origin and current distribution of species (Heads 2005, 2008; Hoeksema 2007). The sparseness of deep-sea samplings often impairs a meaningful interpretation of the deep-sea biogeography. Hypotheses of deep-sea biogeography, inferred primarily from shallow-water patterns are far from accurate. One exception is among the squat lobsters (Galatheoidea and Chirostyloidea) because they have been extensively sampled by the MUSORSTOM-TDSB programme from a wide range of depths (littoral to 1500 m deep), having a large number of species supported by solid taxonomy.

Using the taxonomic results of squat lobsters coming largely from the MUSORSTOM-TDSB collections, two papers were published in 2010 marking the end of the first phase of the Census of Marine Life operations (Macpherson *et al.* 2010; Rowden *et al.* 2010). The first paper was trying to define through statistical analysis the characteristics of each biogeographical province in the Pacific Ocean. The data set composed of 54 deep-sea cruises and 3200 stations with 402 species. The multivariate analyses clearly showed a gradient of decreasing specific richness from Western to Central Pacific. This fundamental concept, well known for the littoral marine fauna, was demonstrated for the first time in the bathyal zone. The richest areas are the Coral triangle (Indo-Malayan-Philippines) and the Coral Sea.

The second work focused on the seamounts of the southwest Pacific and was trying to determine if the faunistic composition is different between seamounts and the slopes at comparable depths. They used a data set from a total of 502 species (Rowden *et al.* 2010) and concluded that the different assemblages of squat lobsters between seamounts and non-seamounts are mainly a consequence of hydrological conditions and especially the availability of food.

Both studies have clearly demonstrated that good taxonomic work at species level is a prerequisite for any meaningful analysis on ecology or biogeography.

### SPECIAL ECOSYSTEMS

During the MUSORSTOM-TDSB explorations two unusual habitats were discovered, representing rich oasis in the deep-sea desert: the summit of seamounts and sunken woods. Both have a special assemblage of life with remarkable crustaceans.

**SUNKEN WOOD FAUNA** — The presence of animals on plant remains at the deep-sea floor was first documented by the CHALLENGER expedition (1872-1876). The GALATHEA expedition (1950-1952) also recorded the abundant presence of plant remains on the deep-sea floor (Bruun 1959). The importance of plant material in the deep sea was first emphasised by Wolf (1979). Associated with these remains is a unique fauna, which has mainly been examined by the taxonomists. The ecological aspects are less well studied until recently. Apart from sporadic taxonomic considerations, plant-associated organisms were seen as zoological and/or ecological curiosities and thus only anecdotally studied. Sunken wood and plant material has long been considered a mere curiosity, but this habitat was recently emphasized as deep-sea environment of major evolutionary importance. Many of the species living on sunken wood are phylogenetically related to those living on hydrothermal vents, cold seeps and other organic substrates (e.g., whale falls).

Distel *et al.* (2000) underlined the evolutionary affinities of the giant hydrothermal vent mussels with those found on sunken wood or whale bones (so-called whale falls). This result was then confirmed by several studies (Samadi *et al.* 2007; Lorion *et al.* 2009; Fujita *et al.* 2009; Kyuno *et al.* 2009). The zoological affinities of organisms associated with sunken woods and whale falls with those from hydrothermal vent or cold seep ecosystems suggests that these ecosystems, developing on organic substrata in the deep-sea, could be an evolutionary step toward the colonization of hydrothermal vents (the so-called “wooden steps” hypothesis of Distel *et al.* (2000)). However, few data are available for sunken wood organisms. Recent results suggest that the evolutionary history of deep-sea mussels associated with organic falls is completely mixed up with that of vent and seep mussels, with evolutionary scenarios probably more complex than the “wooden steps” hypothesis (Lorion *et al.* 2010; Thubaut 2012). Moreover, geography, rather than habitat specificity or chemosymbiotic requirements, seems to significantly structure diversification patterns in sunken wood-associated mytilids (Lorion *et al.* 2009, 2010).


In 2004, the TDSB programme started specific series of cruises particularly devoted to the exploration of sunken-wood environments (*Biodiversité des Organismes Associés aux bois coulés — BOA*). Since the first MUSORSTOM cruises in the Philippines (see Forest 1981) and Indonesia (see Moosa 1984), it became clear that the slopes of oceanic islands within these regions accumulate large amounts of decomposing vegetation. Subsequent cruises in the Solomon Islands, Vanuatu Archipelago, and Papua New Guinea specifically targeted sunken vegetation to further characterize its fauna. Several cruises provided specimens associated with sunken wood in the depth range between 100-1500 m from different locations corresponding to the most important bathyal area where sunken woods accumulate in the Indo-Pacific region: (i) Philippines (PANGLAO, 2005; AURORA, 2007; LUMIWAN, 2008), (ii) Solomon Islands (SALOMON 2, 2004; SALOMONBOA 3, 2007), (iii) Vanuatu Archipelago (BOA 0, 2004; BOA 1, 2005 and SANTOBOA, 2006) and Papua New Guinea (BIOPAPUA, 2010). Additional samples from other cruises, far from these basins, also provided specimens associated with organic remains. Indeed, for example, on seamounts, small pieces of wood and seeds (e.g., coconuts) are variously pulled up by the trawls or the dredges and these erratic substrates, are usually also colonized by the organisms typically found on pieces of wood in accumulation areas.

This sampling is, however, focussed on substrates that are easy to locate on the deep-sea floor (such as large pieces of wood off the mouth of the rivers). Therefore, to more accurately catch the diversity of the organisms associated with a larger diversity of organic substrates that can be found on the deep-sea floor, pots covered by netting with mesh of 3 mm were deployed (Samadi *et al.* 2010). The “bait” placed in the pots were seeds and pieces of wood of various tree species, sugar cane stems, whale bones, green turtle shell, stag horn, feathers, cephalopod beaks and cuttlefish-bones. These lines of pots were deployed in 2003 off the barrier of New Caledonia and off Santo island in Vanuatu and left for periods ranging from 10 to 30 months.

These studies demonstrated that such organic substrates were associated with a rich and unique fauna (Samadi *et al.* 2007) and its confirmation catalyzed new research interests for the TDSB team (reviewed in Samadi *et al.* 2010). With the large scale sampling conducted by TDSB cruises, new recent taxonomic and phylogenetic studies have been

launched, mainly focused on different key-groups like molluscs such as Mytilidae (Lorion *et al.* 2010, 2011; Thubaut 2012) and polyplacophoran chitons (Duperron *et al.* 2012). Several studies have also been done on the feeding behaviour of echinoderms (Becker *et al.* 2009) and molluscs (Zbinden *et al.* 2010), botanical identification of the plant-remains (Pailleret *et al.* 2007), prokaryotic diversity (Palacios *et al.* 2009), fungal diversity (Dupont *et al.* 2009) and the association of mytilids with chemosynthetic bacteria (reviewed in Duperron *et al.* 2009).

For crustaceans, isopods, amphipods and decapods as well as barnacles have been found among plants debris in the trawls or dredges. It is, however, difficult to know to what extent they depend on the decaying plant substrates, in particular for the vagrant fauna. Among the decapods, the main groups repeatedly caught with such substrata include squat lobsters (e.g., *Munidopsis andamanica*, *M. nitida*, *M. bispinosculata*, *M. similior*), two different families of symmetrical hermit crabs (all the species of the genus *Xylopagurus* and several species of *Pylocheles* inhabit centimetre-sized pieces of wood), ghost shrimp (e.g., *Rayllianassa amboinensis*), some Axiidae and a number of snapping shrimps (Alpheidae) (Figure 8A-E).



**FIGURE 8**

**A-D**, Wood-associated crustaceans. **A**, *Galacantha* sp. (photograph B. Richer de Forges). **B**, *Parapylocheles scorpio* (photograph T.-Y. Chan). **C**, *Bathyceradocus* sp. **D**, *Onesimoides* sp. **E-H**, *Bathyceradocus* sp. **E**, General view, total length 18 mm. **F**, SEM picture (magnification x 40) of the mouthparts illustrating the dense bacterial colonisation. **G, H**, SEM pictures (respectively x 800 and x 6000) of the diversity of bacterial morphotypes (rod-shaped and filamentous bacteria) fixed on the mouthpart setae (photographs C-H, L. Corbari).

With regard to peracarid crustaceans, the most recurrent specimens collected in association with wood were: Isopoda (Sphaeromatidae, *Cassidias*; Limnoriidae, *Limnoria* sp.; Gnathiidae, *Elaphognathia* sp.), Amphipoda (Lysianassidae, *Onesimoides* sp.; Maeridae, *Bathyceradocus* sp.). The same organisms were also collected in the pots that were submerged off New Caledonia or Vanuatu.

The main challenge is now to understand the interactions between crustaceans and organic substrates (e.g., habitat, feeding relationship or even dependence in relation to reduced compounds produced by the wood-degradation via microorganisms such as fungi). Preliminary studies on the diet of wood-associated decapods have been conducted on squat lobsters, ghost shrimp and pylochelid hermit crabs (TDSB samples from Vanuatu, New Caledonia, Solomon Archipelago, Philippines). A multidisciplinary approach based on electron microscopy (SEM, TEM); fluorescence in situ hybridization (FISH) and stable isotopic analyses revealed their affiliation to different diets and emphasizes their trophic relationships by the presence (or absence) of bacterial symbioses (Hoyoux 2010). The results revealed that two main diet-groups can be identified for the studied wood-associated decapods: (i) predators such as *Xylocheles macrops*, *Bathycheles incisus*, *B. integer* and *Xylopagurus caledonicus*; and (ii) detritivorous species such as *Munidopsis bispinoculata*, *M. pilosa*, *M. tida*, *M. andamanica* and the thalassinid *Raylianassa amboinensis*. The most striking results deal with the detritivorous groups, especially *M. andamanica* and *M. nitida*. Both species have detritivorous and xylophagous diets and harbour symbiotic digestive microflora, indicating their close trophic interactions with wood and organic substrates. Indeed, a careful examination of the feeding appendages, gut contents and gut lining of *M. andamanica* caught with wood falls revealed this species to be a truly detritivorous species which uses wood and the biofilm-covering as its two main food sources (Hoyoux *et al.* 2009). *Munidopsis andamanica* also possesses a resident gut microflora consisting of bacteria and fungi possibly involved in the digestion of the wood fragments. This species can therefore be regarded as a xylophagous specialist. In this work (Hoyoux *et al.* 2012), authors also examined the feeding appendages, gut contents and gut lining of *M. nitida*, *M. bispinoculata* and *M. pilosa* from natural sunken woods and compared them with specimens of the same species having survived and grown on different hard to-digest substrates (*i.e.*, woods, turtle shells and whale bones) experimentally submerged in the deep South Pacific. In both cases, all three species directly ingest large wood fragments that have been substantially degraded by micro-organisms. However, *M. nitida* also feeds on experimentally submerged whale bone and turtle shell fragments. *Munidopsis nitida* is also the only species known to host a resident gut microflora, although the bacterial morphotypes vary according to the ingested substrate. The results suggest that the three species are most probably opportunistic, bacterivorous detritivores and *M. nitida* could be at the beginning of an evolutionary process leading towards xylophagy within the genus *Munidopsis*.

New investigations are in progress on wood-associated amphipods. Taxonomic and phylogenetic studies have been conducted on several groups. Identified as a key-group is the genus *Bathyceradocus* (family Maeridae; Figure 8E-H) that revealed an unsuspected diversity. *Bathyceradocus* has only three species: *B. iberiensis*, described from Mediterranean Sea; *B. stephensi* described from three specimens collected during cruises of the SIBOGA in the Celebes sea (Pirlot 1934) and *B. wuzzae* recently described by Larsen & Krapp-Schickel (2007) from the vicinity of hydrothermal vents (Juan de Fuca). In all these descriptions, the specimens were collected from dropped wood or plant remains but without more ecological information. About 80 specimens of *Bathyceradocus* sp. have been collected during MUSORSTOM-TDSB cruises from Madagascar to Fiji islands through Philippines and Papua New Guinea. The first morphological hypothesis suggests that these specimens are representative of *B. stephensi*. Phylogenetic analyses (COI and 18S) have revealed that this group is much more diversified than expected: 12 species (based on morphological and molecular data) have been differentiated, revealing the cryptic diversity in this genus (L. Corbari unpublished data). Moreover, microscopic and ultra-structural investigations on the digestive tract have revealed the presence of wood-fragments in the digestive content and the occurrence of micro-organisms (bacteria and fungi) in the posterior part of the intestine. These results confirm that the diet of the *Bathyceradocus* group as strictly wood-feeders (L. Corbari unpublished data). More interesting is the recurrent presence of bacterial ectosymbiosis on the mouthparts of these specimens that suggests an original pathway of nutrition that could involve bacteria in wood degradation (L. Corbari unpublished data). The same approach

of integrative taxonomy (combining morphological and molecular characters) is also being conducted for amphipods in the *Onesimoides* group.

**SEAMOUNTS** — Seamounts are mountains under the sea level and elevated above the sea bottoms more than 1000 m. Seamounts are major features of the oceans. Recent mapping estimated that their number is at least to 14,000 but less than 400 of them had been biologically sampled (Richer de Forges *et al.* 2000; Samadi *et al.* 2007; McClain 2007; Koslow 2007; Richer de Forges & Tessier 2012). They form a benthic habitat of exceptional richness, both in diversity and abundance, and they became a theme of a special section in the CoML called CenSeam, Census of Seamounts (2005-2010). Several MUSORSTOM-TDSB cruises were devoted to this rich habitat in the Pacific (CHALCAL 2, AZTEQUE, NORFOLK 1 & 2, EBISCO, TERRASSE, TARASOC). A preliminary hypothesis on the endemism of the seamount fauna was proposed after the first Norfolk cruises (Richer de Forges *et al.* 2000). More recent studies of population genetics on several animal groups such as molluscs and crustaceans have shown that seamounts are functioning as oases in the deep-sea (Samadi *et al.* 2006; Castelin *et al.* 2012). On the other hand, some species generally considered to be very “rare” were also caught on the Norfolk Ridge seamounts like *Palinustus unicornutus*; *Homola ranunculus*; *Sphaerodromia lamellata*, and *Takedromia longispina*. But the notion of rarity in ecology is a complex issue because sampling in the sea is generally not representative and with some habitats largely neglected, particularly at steep slopes of islands and seamounts (see Mendoza *et al.* 2009).

**EVOLUTION** — Traditional phylogenetic studies based on morphology have been complicated by the identification of homologies, which is often difficult. Many modern techniques have been recently developed to overcome these difficulties. However, all these modern techniques require fresh specimens or specimens preserved in a particularly way when fresh. Most museum specimens are or have been preserved in formalin and unsuitable for these modern techniques. Taxa from the deep-sea are generally not easy to collect because of the access of ship-time and very high running cost of research vessels. Unfortunately, many species and even the whole genus or family in various animal groups are entirely from the deep-sea. Thus, phylogenetic studies using modern techniques often encounter difficulties in sourcing enough taxa to be included in the analysis.

The abundant and diverse specimens collected during the MUSORSTOM-TDSB programme from a very broad geographical area are almost all preserved in ethanol or with the specimens specially treated immediately after collection for later molecular or histological work. For example, a fruitful collaboration was established amongst Bertrand Richer de Forges of the IRD, Danièle Guinot of the Muséum national d’Histoire naturelle and Barrie Jamieson of the University of Queensland on the ultrastructure of spermatozoa in decapod crustaceans. Morphology of reproductive cells is very conservative and can be used to reconstruct phylogeny. Before the MUSORSTOM programme very little was known on the morphology of sperm in decapod crustaceans. Having large access to fresh material from the deep-sea new data were rapidly obtained for the following families: Brachyura (Homolidae, Homolodromiidae, Dromiidae, Dynomeniidae, Latreilliidae, Cyclodorippidae, Dorippidae, Calappidae, Atelecyclidae, Raninidae, Palicidae, Retroplumidae, Leucosiidae, Geryonidae, Gonoplacidae, Cancridae, Grapsidae, Xanthidae, Portunidae, Majidae, Hymenosomatidae), Anomura (Paguridae, Pylochelidae, Parapaguridae, Chirostyliidae, Galatheidae, Lithodidae, Hippidae, Porcellanidae), and Macrura (Palinuridae, Polychelidae, Nephropidae). A large part of these results has been published (Guinot *et al.* 1994; Richer de Forges *et al.* 1997; Tudge *et al.* 2012). Using the sperm characters a new phylogeny of Brachyura was proposed (Jamieson *et al.* 1995). The phylogeny trees obtained are coherent with the classification of Guinot (1977) and distinguishing the three groups Podotremata, Heterotremata and Thoracotremata.

During the last decade, the fast development of molecular biology has provided supplementary information that can strengthen the outcomes of traditional morphology-based studies. The vast and diverse samples collected by the MUSORSTOM-TDSB programme are now a major focus of the international BarCode of Life programme on crustaceans. On the other hand, the availability of many deep-sea taxa that can be used for molecular sequencing has allowed comprehensive studies on the higher classification and evolution of decapod crustaceans (e.g., Tong *et al.* 2000;

Ahyong *et al.* 2007; Chan *et al.* 2008, 2009, 2010; Tsang, Lin *et al.* 2008; Tsang, Ma *et al.* 2008; Ahyong *et al.* 2009; ; Chu *et al.* 2009; Lai *et al.* 2009; Ma *et al.* 2009; Sin *et al.* 2009; Tshudy *et al.* 2009; Tsang *et al.* 2009, 2011; De Grave *et al.* 2010; Boisselier-Dubayle *et al.* 2010; Lai *et al.* 2011; Li *et al.* 2011; Schnabel *et al.* 2011; Tsoi *et al.* 2011; Yang *et al.* 2012). Some studies have shown the importance of the southwest Pacific area, specifically the island arc of Fiji, Tonga, Vanuatu, Wallis and Futuna, for diversification of squat lobsters, probably associated with a global warming, high tectonic activity, and changes in oceanic currents that took place in this region during the Oligocene-Miocene period (Cabezas *et al.* 2012).

The results of these molecular studies have revolutionized the classification scheme of decapod crustaceans; such as the hermit crabs, squat lobsters, the mud shrimps Thalassinidea, the subfamilies in Nephropidae and many caridean superfamilies are not likely to be monophyletic groups, while the family or generic status of Sicyoniidae, Thaumastochelidae, and Synaxidae may not be valid. On the other hand, the status of Enoplometopidae is settled and considered as a superfamily in Astacidea. Moreover, the molecular data provide insights into the evolution within some of the infraorders, such as parallel carcinization within the Anomura (Tsang *et al.* 2011).

Phylogenetics and molecular studies of paguroids in particular, and anomurans or decapods in general, have taken advantage of the new availability of specimens as well as the greatly expanded morphological and distributional knowledge generated during nearly 40 years of publications based on sampling programs in the deep-seas of the Indo-West Pacific. Discoveries of remarkable new forms of what may be considered primitive or advanced taxa, have provided improved insight into the evolution of paguroids. Cladistic and genetic studies have incorporated the data or molecular information extracted from the vast MUSORSTOM-TDSB collections. Significantly improved evolutionary scenarios have been produced, and although theories are still controversial and hotly debated, we are much closer than before to the understanding of the true relationships and origins of paguroids in particular, and anomurans in general (e.g., McLaughlin *et al.* 2007; Tsang *et al.* 2008, 2011; Ahyong *et al.* 2009; Lemaitre & McLaughlin 2009; Lemaitre *et al.* 2009).

The molecular techniques have been providing new findings on the taxonomy, phylogeny and classification of squat lobsters, e.g., Machordom & Macpherson (2001, 2004), Chu *et al.* (2009), Ahyong *et al.* (2009), Cabezas *et al.* (2008, 2010), Puillandre *et al.* (2011). The most recent and complete studies have proposed a new classification of the squat lobsters (Ahyong *et al.* 2010; Schnabel & Ahyong 2010), with two superfamilies (Galatheoidea and Chirostyloidea). The material obtained along the Indian and Pacific Oceans has allowed some studies on the phylogeny and origin of this rich fauna. These studies suggest that the galatheid diversification in the southwestern Pacific may have begun some 7-14 million years ago, during the Middle or Late Miocene (Machordom & Macpherson 2004), and the Coral Sea area seems to be the origin of several genera of squat lobsters like *Paramunida* and *Plesionida* (Cabezas *et al.* 2009, 2012).

All in all, the MUSORSTOM-TDSB collection now forms an important part in specimen sourcing for the molecular component in the international Assembling the Tree of Life project on decapod crustaceans.

### THE ALAIN CROSNIER EFFECT

All crustacean researchers who have worked on the MUSORSTOM-TDSB collections were invariably influenced by the “Alain Crosnier effect”. His energy, passion and dedication were legendary. In the years between the first cruise (1976) to 2008 (when he decided to fully retire), he was instrumental in arranging expeditions, having the material centralized in the Muséum national d’Histoire naturelle where he sorted it out, then invited specialists from around the world to work on them (or occasionally sending it to them for study) and then assiduously arranging to have them published. The series of MUSORSTOM volumes for which he personally edited and readied for press exceeds 10,000 printed pages, and with hundreds of new genera and species described within.

More significantly, many of the papers published as a result of Crosnier’s efforts were not just straightforward new genera and new species descriptions. A good number were major revisions or reviews of genera, subfamilies and families that are extremely important and useful for biologists. Crosnier used the extensive MUSORSTOM-TDSB collections as a nexus and a catalyst for such revisions that brought together types and material from all over the world. Crosnier was directly or indirectly responsible for more such major revisions than any other period in carcinological science.

Overall, the efforts he put in to secure the collections and the funds to have the experts work in Paris, and then to have them published were nothing short of phenomenal. And he was a tough taskmaster, driving the researchers hard to have the job done and done well. He was a “one-man army”, almost single-handedly pushing carcinological and deep-sea research to heights it has never seen before. It was a Herculean effort that has never been exceeded and may never be equaled.

Through all these years, Alain also created a special community of carcinologists devoted to the MUSORSTOM enterprise and the advancement of crustacean systematic knowledge (see Macpherson *et al.* 1998). This included the many researchers he invited to the museum to work on his material (the first was the late Chen Hui-Lian from Qingdao, China, who worked on Leucosiidae and Dorippidae; the last being Tomoyuki Komai from Chiba, Japan, a specialist of crangonids and glyptocrangonids) as well as those he generously sent material to. Affectionately known as “Crosnier’s cronies”, this community continues their works to this day across almost all continents and over a dozen countries. He was even honoured by his peers and “cronies” in a special volume of *Zoosystema* in 1998 (Volume 20 number 2). Alain was an influential carcinologist himself, specialising in penaeoid shrimps, but also publishing on a wide variety of other decapods. Even today, several years after his retirement from science, many researchers of this “fellowship of the MUSORSTOM” still correspond with him. For us, he is a classic demonstration of the French proverb “la foi soulève des montagnes” (faith that moves mountains). And he did move “many large mountains” in advancing carcinology and deep-sea research. The Alain Crosnier effect is now deeply entrenched in carcinological history.

### THE INDISPENSABLE ROLE OF TAXONOMY

The endeavours of the MUSORSTOM-TDSB rammed home the importance of taxonomy as a science. Nowadays, with the rapid development of powerful new molecular tools, there is a tendency for many people, including respected colleagues and scientific leaders, to decry morphological taxonomy as “old fashioned” and obsolete, dated, unimportant and irrelevant. They think new techniques like DNA barcoding and fingerprinting will replace morphologists. They are wrong.

The MUSORSTOM-TDSB programme is a clear demonstration that taxonomy is at the heart of good biological science. Exploration and surveys can collect extensive material for study, but for this material to be useful for science, the right questions must be asked and hypotheses tested. To have a solid knowledge of zoology, the species in this material have to be formally described and named. Detractors who argue that a COI gene sequence alone is enough to characterise a species are fooling themselves. A genetic barcode is not unlike a person’s identification number — it indicates your position in a list, but not who you are (see also Kwong *et al.* 2012; Taylor & Harris 2012).

It is often forgotten (or ignored) that taxonomic classifications and species names, represent hypotheses of biological relationship. The population to which a species name has been applied has unique biological properties, shared within the group, and different from the next species. The information embodied in the species’ classification tells us about its diagnostic and distinctive features, and provides indications as to what the animals may do in life.

There can be no denying that molecular methods are powerful aids to the work of taxonomists, but they cannot be used in isolation. Moreover, genetic tools are also limited in their ability to distinguish species, without reference to morphology, as recently demonstrated for deep-sea corals (Baco & Cairns 2012). Ultimately, knowledge of the morphology is integral to the human understanding of what animals are and do. Far from being “old-fashioned”, morphological taxonomy remains a timeless and very essential discipline.

The strong taxonomic foundation deriving from the MUSORSTOM-TDSB programme has led to numerous advances in our knowledge of deep-sea biology. The many named taxa (families, genera and species) have been used for dozens of important phylogenetic, biogeographic and ecological questions, many of which have been reviewed above.

### NEW PARADIGMS

In the middle of the 19th century, our deep-sea knowledge was very limited and the dominant idea at this time was that the bathyal world is an extremely harsh environment that was primarily azoic. In the second half of the 19th century,

especially after the brilliant success of the deep-sea dredgings by Charles Wyville Thomson, the scientific community began to realise that there is a high diversity of animals in the deep-sea. England started the race to discover and sample the world oceans with the CHALLENGER expedition (1872-1876) funded by the Royal Society. Other countries quickly followed suit: The Netherlands (SIBOGA 1899-1900), Germany (VALDIVIA 1898-1899), British India (INVESTIGATOR 1884-1897), United States (ALBATROSS 1891-1905) and Denmark (GALATHEA 1950-1952). As a result of these surveys, by the middle of the 20th century, the revised thinking was that there is life in the deep-sea at all depths, but the diversity is relatively poor with the animals widely distributed.

From 1970 onwards, ORSTOM, under the leadership of Alain Crosnier, started sampling the deep-sea fauna in Madagascar, originally aimed at exploring fish and crustacean resources. But these explorations were boosted by the discovery of *Neoglyphea* in 1975 and its rediscovery in 1976 in the Philippines; the same year the American submersible "Alvin" discovered the astonishing fauna living in and around hydrothermal vents. This extraordinary discovery completely changed our concept of life in the deep-sea. Several decades of deep-sea studies mainly using "high technology" submersibles have since focused on this remarkable fauna that is dependent on chemosynthetic ecosystems along the oceanic ridges (Desbruyères & Segonzac 1997; Laubier 2008; [www.noc.soton.ac.uk/chess](http://www.noc.soton.ac.uk/chess)).

At the same time, the MUSORSTOM-TDSB programme used "low technology" efforts to sample the bathyal zone around tropical Islands and started to document the "more ordinary fauna". These studies have opened several new paradigms in our understanding of the deep-sea environment: (i) the deep-sea benthos in tropical areas is very diverse in the upper part of the bathyal zone; (ii) some ecosystems like steep hard cliffs are neglected and very hard to sample but nevertheless very species rich; (iii) even if some species have a wide distribution, each archipelago generally has its own unique faunal assemblage; (iv) for the deep-sea epi-benthos in the Pacific, there is a decreasing gradient of diversity from West to East; with a similar pattern from the West Pacific to West Indian Ocean; (v) organic substrates such as sunken wood provides a key food resource in the deep-sea, and acts as stepping stones for the dispersion of chemosynthetic species; and (vi) the thousands of seamounts in the world oceans are generally oases that are concentrate with deep-sea species and play an important role in the global dispersion of species.

## CONCLUSIONS

With the rapid changes of environment, the huge MUSORSTOM-TDSB collection of deep-sea fauna that spans the last 40 years constitutes a very valuable archive to understand deep-sea evolution on our planet. Climate change is a hot topic nowadays. With the Arctic ice disappearing rapidly, sea level rise may increase faster than expected. The increasing erosion of the coast and discharge of anthropogenically impacted rivers bringing huge amounts of sediments and organic material into the oceans. The changing hydrological conditions on the surface modify the plankton productivity and may also modify oceanic currents and deep-sea habitats. All this will ultimately influence the deep-sea fauna significantly.

In only 160 years, our scientific knowledge of the deep-sea has transitioned from a trickle to a flood, giving scientists an almost "panoramic vision" of the great diversity of life existing on the sea floor of our planet. Not surprisingly, this diversity is mostly localized at ridges, seamounts and continental slopes. The richest deep-sea areas appear to be situated in the western tropical Pacific Ocean. However, we have to remain very humble about our knowledge of the deep-sea, remembering that a large part of it remains unexplored (Observatory Biodiversity Information System: [www.iobis.org](http://www.iobis.org)). For example, less than 3% of large seamounts have been sampled, and a large part of the zoological collections already obtained from seamounts and deep-water systems are still unstudied because of an acute lack of taxonomic specialists. Even for studied species very little is known about the reproduction, growth and behavior of these deep-sea animals. Deep-sea explorations now mainly focus on the role of chemosynthesis, methane seeps, hydrothermal vents and the origin of chemosynthetic fauna. To understand, conserve and manage the deep-sea environment, deep-sea biodiversity survey programs such as TDSB are more important than ever.

## ACKNOWLEDGEMENTS

A great number of people have contributed to the MUSORSTOM saga in discovering and describing the deep-sea tropical fauna. This short assessment on crustaceans is an opportunity to thank them all: the crews of the oceanographic and fishing boats involved in the hundreds of cruises; the technical teams associate with specimen collection, preservation and curation; the artists for making photographs and drawings; the editors and publishers of the MUSORSTOM-TDSB volumes; the administrations supporting the researches; the sponsors helping to fund the boats expenses; the network of international researchers spending time to study the fauna. A special mention for Alain Crosnier who was a pioneer in many ways animated the research groups (not limited to Crustacea), from the beginning of obtaining funding for the cruises till the end of publications of the results.

## REFERENCES

- AHYONG S. T. 2002a — Stomatopod Crustacea from the Marquesas Islands: results of MUSORSTOM 9. *Zoosystema* 24: 347-372.
- AHYONG S. T. 2002b — Stomatopod Crustacea of the Karubar Expedition in Indonesia. *Zoosystema* 24: 373-383.
- AHYONG S. T. 2007 — Shallow water Stomatopoda of New Caledonia (0-100 m), in PAYRI C. E. & RICHER DE FORGES, B. (ed.) *Compendium of Marine Species from New Caledonia. Documents Scientifiques et Techniques II*: 333-335. IRD Noumea, Noumea.
- AHYONG S. T. 2013 — Stomatopoda collected primarily by the Philippine AURORA 2007 expedition (Crustacea: Stomatopoda: Squilloidea), in AHYONG S. T., CHANT T.-Y., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 85-106.
- AHYONG S. T., BABA K., MACPHERSON E. & POORE G. C. B. 2010 — A new classification of the Galatheoidea (Crustacea: Decapoda: Anomura). *Zootaxa* 2676: 57-68.
- AHYONG S. T. & CHAN T.-Y. 2008 — Polychelidae from the Bohol and Sulu Seas collected by PANGLAO 2005 (Crustacean: Decapoda: Polychelidae). *Raffles Bulletin of Zoology*, suppl. 19: 63-70.
- AHYONG S. T. & CHANT T.-Y. 2010 — *Lithodes formosae*, a new species of king crab from Taiwan (Crustacea: Decapoda: Lithodidae). *Zootaxa* 2332: 61-68.
- AHYONG S. T., CHANT T.-Y. & BOUCHET P. 2010 — Mighty claws: a new genus and species of lobster from the Philippines deep-sea (Crustacea, Decapoda, Nephropidae). *Zoosystema* 32 (3): 525-535.
- AHYONG S. T. & GALIL B. S. 2006 — Polychelidae from the southern and western Pacific (Decapoda, Polychelida). *Zoosystema* 28 (3): 757-767.
- AHYONG S. T., LAI J. C. Y., SHARKEY D., COLGAN D. J. & NG P. K. L. 2007 — Phylogenetics of the brachyuran crabs (Crustacea: Decapoda): The status of Podotremata based on small subunit nuclear ribosomal RNA. *Molecular Phylogenetics and Evolution* 45: 576-586.
- AHYONG S. T. & NG P. K. L. 2007 — *Pleistacantha ori*, a new species of deep-water spider crab (Crustacea: Decapoda: Brachyura: Majidae) from the western Indian Ocean. *Raffles Bulletin of Zoology*, suppl. 16: 67-74.
- AHYONG S. T. & NG P. K. L. 2008 — *Alain raymondi*, a new species of deepwater pinnotherid crab (Crustacea: Decapoda: Brachyura) from the Philippines, commensal with holothurians. *Zootaxa* 1919: 61-68.
- AHYONG S. T. & NG P. K. L. 2011 — Cyclodorippoid crabs from the Philippines collected by the PANGLAO 2004-2005 and AURORA 2006 expeditions. *Zoologische Anzeiger* 250: 479-487.
- AHYONG S. T. & NG P. K. L. 2009 — The Cymonomidae of the Philippines (Crustacea: Decapoda: Brachyura), with descriptions of four new species. *Raffles Bulletin of Zoology*, suppl. 20: 233-246.
- AHYONG S. T., SCHNABEL K. E. & MAAS E. 2009 — Anomuran phylogeny: new insights from molecular data, in MARTIN J. W., CRANDALL K. A. & FELDER D. L. (eds), *Decapod crustacean phylogenetics. Crustacean Issues* 18: 399-414.
- APPELTANS W., AHYONG S. T., ANDERSON G., ANGEL M. V., ARTOIST T., BAILLY N., BAMBER R., BARBER A., BARTSCH I., BERTA A., BLAZEWICZ-PASZKOWYCZ M., BOCK P., BOXSHALL G., BOYKO C. B., BRANDÃO S. N., BRAY R. A., BRUCE N. L., CAIRNS S. D., CHAN T.-Y., CHAN L., COLLINS A. G., CRIBB T., CURINI-GALLETI M., DAHDOUH-GUEBAS F., DAVIE P. J. F., DAWSON M. N., DE CLERCK O., DECOCK W., DE GRAVE S., VOOGD N. J. DE, DOMMING D. P., EMIG C. C., ERSÉUS C., ESCHMEYER W., FAUCHALD K., FAUTIN D. G., FEIST S. W., FRANSEN C. H. J. M., FURUYA H., GARCIA-ALVAREZ O., GERKEN S., GIBSON D., GITTEMBERGER A., GOFAS S., GÓMEZ-DAGLIO L., GORDON D. P., GUIRY M. D., HERNANDEZ F., HOEKSEMA B. W., HOPCROFT R., JAUME D., KIRK P., KOEDAM N., KOENEMANN S., KOLB J. B., KRISTENSEN R. M., KROH A., LAMBERT G., LAZARUS D. B., LEMAÎTRE R., LONGSHAW M., LOWRY J., MACPHERSON E., MADIN L. P., MAH C., MAPSTONE G., MCLAUGHLIN P., MEES J., MELAND K., MESSING C. G., MILLS C. E., MOLODTSOVAT T. N., MOOI R., NEUHAUS B., NG P. K. L., NIELSEN C., NORENBURG J., OPRESKO D. M., OSAWA M., PAULAY G., PERRIN W., PILGER J. F., GARY C. B. P., PUGH P., READ G. B., REIMER J. D., RIUS M., ROCHA R. M., SAIZ-SALINAS J. I., SCARABINO V., SCHIERWATER B., SCHMIDT-RHAESA A., SCHNABEL K. E., SCHOTTE M., SCHUCHERT P., SCHWABE E., SEGERS H., SELF-SULLIVAN C., SHENKAR N., SIEGEL V., STERRER W., STÖHR S., SWALLA B., TASKER M. L., THUESEN E. V., TIMM T., TODARO A., TURON X., TYLER S., UETZ P., VAN DER LAND J., VANHOORNE B., VAN OFWEGEN L. P., VAN SOEST R. W. M., VANAVERBEKE J., WALKER-SMITH G., WALTER T. C., WARREN A., WILLIAMS G., WILSON S. P. & COSTELLO M. J. 2012 — The magnitude of global marine species diversity. *Current Biology* 22: 2189-2202.
- AUSUBEL J. H., TREW CRIST D. & WAGGONER P. E. 2010 — *First Census of Marine Life 2010. Highlights of a Decade of Discovery*. Census of Marine Life, 64p.
- BABA K. 1990 — Chirostylid and galatheid crustaceans of Madagascar (Decapoda, Anomura). *Bulletin du Muséum national d'Histoire naturelle*, Paris (4) (A) 11: 921-975.
- BABA K. 1991a — Crustacea Decapoda: *Chirostylus* Ortmann, 1892, and *Gastroptychus* Caullery, 1896 (Chirostyliidae) from New Caledonia, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM Vol. 9. Mémoires du Muséum national d'Histoire naturelle* 152: 463-477.

- BABA K. 1991b — Crustacea Decapoda: *Alainius* gen. nov., *Leiogalathea* Baba, 1969, and *Phylladiorhynchus* Baba, 1969 (Galatheidae) from New Caledonia, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 479-491.
- BABA K. 2004 — *Uroptychodes*, new genus of Chirostylidae (Crustacea: Decapoda: Anomura), with description of three new species. *Scientia Marina* 68: 97-116.
- BABA K. & LIN C. W. 2008 — Five new species of chirostyliid crustaceans (Crustacea: Decapoda: Anomura: Chirostylidae) from Taiwan. *Zootaxa* 1919: 1-24.
- BABA K., MACPHERSON E., POORE G. C. B., AHYONG S. T., BERMUDEZ A., CABEZAS P., LIN C. W., NIZINSKI M., RODRIGUES C. & SCHNABEL K. E. 2008 — Catalogue of squat lobsters of the world (Crustacea: Decapoda: Anomura—families Chirostylidae, Galatheidae and Kiwaidae). *Zootaxa* 1905: 1-220.
- BABA K., MACPHERSON E., LIN C. W. & CHAN T.-Y. 2009 — Crustacean Fauna of Taiwan. Squat lobsters (Chirostylidae and Galatheidae). National Taiwan Ocean University, Keelung: ix + 312 p.
- BABA K. & SAINT LAURENT DE M. 1992 — Chirostyliid and galatheid crustaceans (Decapoda: Anomura) from active thermal vent areas in the southwest Pacific. *Scientia Marina* 56: 321-332.
- BABA K. & SAINT LAURENT DE M. 1996 — Crustacea Decapoda: Revision of the genus *Bathymunida* Balss, 1914, and description of six new related genera (Galatheidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 433-502.
- BACO A. R. & CAIRNS S. D. 2012 — Comparing molecular variation to morphological species designations in the deep-sea coral *Narella* reveals new insights into seamount coral ranges. *PLoS One* 7 (9): e45555.
- BECKER P., SAMADI S., ZBINDEN M., HOYOUX C., COMPÈRE P. & DE RIDDER C. 2009 — First insights into the gut microflora associated with an echinoid from wood falls environments. *Cahiers de Biologie Marine* 50: 343-352.
- BOISSELIER-DUBAYLE M. C., BONILLO C., CRUAUD C., COULOUX A., RICHER DE FORGES B. & VIDAL N. 2010 — The phylogenetic position of the "living fossils" *Neoglyphea* and *Laurentaeglyphea* (Decapoda: Glypheidea). *Comptes Rendus de l'Académie des Sciences; Biologies* 333 (10): 755-759.
- BOUCHET P., HEROS V., LOZOUET P. & MAESTRATI P. 2008 — A quarter-century of deep-sea malacological exploration in the South and West Pacific: Where do we stand? How far to go? in HEROS V., COWIE R. H. & BOUCHET P. (eds), Tropical Deep-Sea Benthos Vol. 25. *Mémoires du Muséum national d'Histoire naturelle* 196: 9-40.
- BOUCHET P., NG P. K. L., LARGO D. & TAN S. H. 2009 — PANGLAO 2004 — Investigations of the marine species richness in the Philippines. *Raffles Bulletin of Zoology*, suppl. 20: 1-19.
- BOURDON R. & BOYKO C. B. 2005 — Redescription of *Bopyrophrynxus brachiabdominalis* Codreanu, 1965 (Crustacea: Isopoda: Bopyridae) with a reappraisal of the subfamily Bopyrophryxinae Codreanu, 1965. *Proceedings of the Biological Society of Washington* 118 (1): 108-116.
- BOYKO C. B. 2003 — A new genus and species of bopyrid isopod (Crustacea, Isopoda, Bopyridae, Orbioninae) parasitic on *Sicyonia* (Crustacea, Decapoda, Penaeoidea) from New Caledonia. *Zoosystema* 25 (4): 593-600.
- BOYKO C. B. 2004 — The Bopyridae (Crustacea, Isopoda) parasites of the Styloactylidae (Crustacea, Decapoda, Caridea). *Zoosystema* 26 (2): 199-210.
- BOYKO C. B. 2004 — The Bopyridae (Crustacea: Isopoda) of Taiwan. *Zoological Studies* 43 (4): 677-703.
- BRUCE A. J. 1981 — Decapod Crustacea: Pontoniinae, in DORST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 1. *Mémoires ORSTOM* 91: 189-215.
- BRUCE A. J. 1990 — Crustacea Decapoda: Deep-sea Palaemonoid shrimps from New Caledonian waters, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 149-215.
- BRUCE A. J. 1991 — Crustacea Decapoda: Further deep-sea Palaemonoid shrimps from New Caledonian waters, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 299-411.
- BRUCE A. J. 1996 — Crustacea Decapoda: Palaemonoid shrimps from the Indo-West Pacific region mainly from New Caledonia, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 197-267.
- BRUUN A. F. 1956 — Animal life of the deep sea bottom, in BRUUN A. F., GREVE S. V., MIELCHE H. & SPÄRK R. (eds), *The Galathea Deep Sea Expedition, 1950-1952. Described by members of the expedition*. New York, The MacMillan company: 149-195.
- BRUUN A. F. 1959 — General introduction to the reports and list of deep-sea stations. *Galathea Report* 1: 7-28.
- BURUKOVSKY R. N. 2013 — Shrimps of the family Nematocarcinidae Smith, 1884 (Crustacea, Decapoda, Caridea) from Taiwan and the Philippines collected by the TAIWAN, PANGLAO 2005 and AURORA expeditions in the western Pacific, in AHYONG S. T., CHAN T.-Y., CORBARI L. & NG P. K. L. (eds), *Tropical Deep-Sea Benthos Vol. 27. Mémoires du Muséum national d'Histoire naturelle* 204: 155-190.
- CABEZAS P., LIN C.-W. & CHAN, T.-Y. 2011 — Two new species of the deep-sea squat lobster genus *Munida* Leach, 1820 (Crustacea: Decapoda: Munididae) from Taiwan: morphological and molecular evidence. *Zootaxa* 3036: 26-38.
- CABEZAS P., LIN C.-W. & CHAN T.-Y. 2012 — A new species of *Uroptychus* (Decapoda: Anomura: Chirostylidae) from Taiwan. *Bulletin of Marine Science* 88: 113-118.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2008 — A new genus of squat lobster (Decapoda: Anomura: Galatheidae) from the South West Pacific and Indian Ocean inferred from morphological and molecular evidence. *Journal of Crustacean Biology* 28: 68-75.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2009 — Morphological and molecular description of new species of squat lobster (Crustacea: Decapoda: Galatheidae) from the Solomon and Fiji Islands (South-West Pacific). *Zoological Journal of the Linnean Society* 156: 465-493.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2010 — Taxonomic revision of the genus *Paramunida* Baba, 1988 (Crustacea: Decapoda: Galatheidae): a morphological and molecular approach. *Zootaxa* 2712: 1-60.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2011 — *Allogalathea* (Decapoda: Galatheidae): a monospecific genus of squat lobsters? *Zoological Journal of the Linnean Society* 156: 465-493.
- CABEZAS P., SANMARTÍN I., PAULAY G., MACPHERSON E. & MACHORDOM A. 2012 — Deep under the sea: Unraveling the evolutionary history of the deep-sea squat lobster *Paramunida* (Decapoda, Munididae). *Evolution* 66: 1878-1896.
- CASTELIN M., PUILLANDRE N., LOZOUET P., SYSOEV A., RICHER DE FORGES B. & SAMADI S. 2012 — Molluskan species richness and endemism on New Caledonian seamounts: Are they enhanced compared to adjacent slopes. *Deep-sea Research I*, 58: 637-646.

- CASTRO P. 2000 — Crustacea Decapoda: A revision of the Indo-west Pacific species of palicid crabs (Brachyura Palicidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 437-610.
- CASTRO P. 2005 — Crabs of the subfamily Ethusinae Guinot, 1977 (Crustacea, Decapoda, Brachyura, Dorippidae) of the Indo-West Pacific region. *Zoosystema* 27 (3): 499-600.
- CASTRO P. 2007 — A reappraisal of the family Gonoplacidae MacLeay, 1838 (Crustacea, Decapoda, Brachyura) and revision of the subfamily Gonoplacinae, with the description of ten new genera and eighteen new species. *Zoosystema* 29 (4): 609-773.
- CASTRO P. 2009 — Two new species of *Carcinoplax* H. Milne Edwards, 1852, and *Pycnoplax* Castro, 2007, from the western Pacific, and a description of the female of *Thyraplax truncata* Castro, 2007 (Crustacea, Decapoda, Brachyura, Gonoplacidae). *Zoosystema* 31 (4): 949-957.
- CASTRO P. 2010 — A new species and new records of palicoid crabs (Crustacea, Decapoda, Brachyura, Palicoidea, Palicidae, Crossotonotidae) from the Indo-West Pacific region. *Zoosystema* 32 (1): 73-86.
- CASTRO P. 2013 — Brachyuran crabs (Crustacea, Brachyura: Crossotonotidae, Ethusidae, Euryplacidae, Gonoplacidae, Latreilliidae, Palicidae, Tetraliidae, Trapeziidae) of the MAINBAZA, MIRIKY, and ATIMO VATAE expeditions to the Mozambique Channel and Madagascar, in AHYONG S.T., CHANT T.Y., CORBARI L. & NG P.K.L. (eds), Tropical Deep-Sea Benthos Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 437-466.
- CASTRO P. & NG P.K.L. 2008 — *Rhadinoplax*, a new genus of Progeryonidae Števićić, 2005, for *Carcinoplax microphthalmus* Guinot & Richer de Forges, 1981, and a redescription of *Paragalene longicrura* (Nardo, 1868) (Crustacea: Decapoda: Brachyura: Gonoplaocoidea). *Zootaxa* 1777: 53-68.
- CASTRO P. & NG P.K.L. 2010a — Revision of the family Euryplacidae Stimpson, 1871 (Crustacea: Decapoda: Brachyura: Gonoplaocoidea). *Zootaxa* 2375: 1-130.
- CASTRO P. & NG P.K.L. 2010b — A new genus and species of gonoplacid crab (Decapoda, Brachyura, Gonoplacidae) from the western Pacific, in CASTRO P., DAVIE P.J.F., NG P.K.L. & RICHER DE FORGES B. (eds), Studies on Brachyura: a homage to Danièle Guinot. *Crustaceana Monographs* 11: 51-60.
- CASTRO P., NG P.K.L. & AHYONG S.T. 2004 — Phylogeny and systematics of the Trapeziidae Miers, 1886 (Crustacea: Brachyura), with the description of a new family. *Zootaxa* 643: 1-70.
- CASTRO P., WILLIAMS A. B. & COOPER L. L. 2003 — Revision of the family Latreilliidae Stimpson, 1858 (Crustacea, Decapoda, Brachyura). *Zoosystema* 25 (4): 601-634.
- CHAN T.-Y. 1996 — Crustacea Decapoda Crangonidae: Revision of the three closely related genera *Aegaeon* Agassiz, 1846, *Pontocaris* Bate, 1888 and *Parapontocaris* Alcock, 1901, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 269-336.
- CHAN T.-Y. 1997 — Crustacea Decapoda: Palinuridae, Scyllaridae and Nephropidae collected in Indonesia by the KARUBAR cruise, with an identification key for the species of *Metanephrops*, in CROSNIER A. & BOUCHET P. (eds), Résultats des Campagnes MUSORSTOM Vol. 16. *Mémoires du Muséum national d'Histoire naturelle* 172: 409-431.
- CHANT T.-Y. 2004 — The "Plesionika rostricrescentis" (Bate, 1888)" and "P. lophotes Chace, 1985" species groups of *Plesionika* Bate, 1888, with descriptions of five new species (Crustacea: Decapoda: Pandalidae), in MARSHALL B. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos Vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 293-318.
- CHAN T.-Y. 2012 — A new genus of deep-sea solenocerid shrimp (Decapoda: Penaeoidea) from Papua New Guinea. *Journal of Crustacean Biology* 32 (3): 489-495.
- CHANT T.-Y. & CROSNIER A. 1991 — Crustacea Decapoda: Studies of the *Plesionika* *nerval* (Fabricius, 1787) group (Pandalidae), with descriptions of six new species, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 413-461.
- CHANT T.-Y. & CROSNIER A. 1997 — Crustacea Decapoda: Deep sea shrimps of the genus *Plesionika* Bate, 1888 (Pandalidae) from French Polynesia, with descriptions of five new species, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 187-234.
- CHANT T.-Y., HO K.C., LI C.P. & CHU K.H. 2009 — Origin and diversification of the clawed lobster genus *Metanephrops* (Crustacea: Decapoda: Nephropidae). *Molecular Phylogenetics and Evolution* 50: 411-422.
- CHANT T.-Y., LEI H.C., LI C.P. & CHU K.H. 2010 — Phylogenetic analysis using rDNA reveals polyphyly of Ophelophoridae (Decapoda: Caridae). *Invertebrate Systematics* 24: 172-181.
- CHANT T.-Y., MA K.Y. & CHU K.H. 2013 — The deep-sea spiny lobster genus *Puerulus* Ortmann, 1897 (Crustacea, Decapoda, Palinuridae), with descriptions of five new species, in AHYONG S.T., CHAN T.-Y., CORBARI L. & NG P.K.L. (eds), Tropical Deep-Sea Benthos Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 191-230.
- CHAN T.-Y. & NG P.K.L. 2008 — *Enoplometopus* A. Milne-Edwards, 1862 (Crustacea: Nephropidea) from the Philippines, with description of one new species and a revised key to the genus. *Bulletin of Marine Science* 83 (2): 347-365.
- CHAN T.-Y. & SAINT LAURENT DE M. 1999 — The rare lobster genus *Thaumastocheles* (Decapoda: Thaumastocheilidae) from the Indo-Pacific, with description of a new species. *Journal of Crustacean Biology* 19 (4): 891-901.
- CHAN T.-Y., TONG J., TAM Y.K., CHU K.H. 2008 — Phylogenetic relationships among the genera of the Penaeidae (Crustacea: Decapoda) revealed by mitochondrial 16S rRNA gene sequences. *Zootaxa* 1694: 38-50.
- CHAN T.-Y. & YU H.P. 1991a — Studies on the *Metanephrops japonicus* group (Decapoda, Nephropidae), with descriptions of two new species. *Crustaceana* 60 (1): 18-51.
- CHAN T.-Y. & YU H.P. 1991b — *Eugonatonotus chacei* sp. nov., second species of the genus (Crustacea, Decapoda, Eugonatonotidae). *Bulletin du Muséum national d'Histoire naturelle* (4) 13 (A) (1-2): 143-152.
- CHAN T.-Y. & YU H.P. 1991c — Two similar species: *Plesionika edwardsii* (Brandt, 1851) and *Plesionika crosnieri*, new species (Crustacea: Decapoda: Pandalidae). *Proceedings of the Biological Society of Washington* 104 (3): 545-555.
- CHAN T.-Y. & YU H.P. 1995 — On the rare lobster genus *Palinustus* A. Milne Edwards, 1880 (Decapoda: Palinuridae), with description of a new species. *Journal of Crustacean Biology* 15 (2): 376-394.
- CHAN T.-Y. & YU H.P. 1998 — A new reef lobster of the genus *Enoplometopus* A. Milne Edwards, 1862 (Decapoda: Nephropoidea) from the western and southern Pacific. *Zoosystema* 20 (2): 183-192.
- CHEN H.L. 1986 — Decapod Crustacea: Dorippidae, in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 2. *Mémoires du Muséum national d'Histoire naturelle* 133: 179-204, pls 1, 2.
- CHEN H.L. 1989 — Leucosiidae (Crustacea, Brachyura), in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 5. *Mémoires du Muséum national d'Histoire naturelle* 144: 181-264, pls 1-6.

- CHEN H. 1990 — Leucosiidae (Crustacea Decapoda: Brachyura) collected in Indonesia during the Corindon II Expedition. *Indo-Malayan Zoology* (1989) 6: 173-184.
- CHEN H. L. 1993 — Crustacea Decapoda: Dorippidae of New Caledonia, Indonesia and the Philippines, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 315-345.
- CHEN H. L. 1997 — Crustacea Decapoda: Ethusinae (Dorippidae) mainly from the KARUBAR cruise, in CROSNIER A. & BOUCHET P. (eds), Résultats des Campagnes MUSORSTOM Vol. 16. *Mémoires du Muséum national d'Histoire naturelle* 172: 613-625.
- CHEN H. L. 2000 — Crustacea Decapoda: New species and new records of Ethusinae (Dorippidae) from Vanuatu, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 425-435.
- CHIA D. G. B. & NG P. K. L. 1995 — A revision of the genus *Rhabdonotus* A. Milne-Edwards, 1879, with descriptions of two new species and the first zoeal stage of *R. pictus* A. Milne-Edwards, 1879, (Brachyura: Eumelonidae). *Crustacean Research* 24: 104-127.
- CHIA D. G. B. & NG P. K. L. 2000 — A revision of *Eumedonous* H. Milne Edwards, 1834 and *Gonatonotus* White, 1847 (Crustacea: Decapoda: Brachyura: Eumelonidae), two genera of crabs symbiotic with sea urchins. *Journal of Natural History* 34: 15-56.
- CHU K. H., TSANG L. M., MA K. Y., CHAN T.-Y. & NG P. K. L. 2009 — Decapod phylogeny: What can protein-coding genes tell us? in MARTIN J. W., CRANDALL K. A. & FELDER D. L. (eds), *Decapod crustacean phylogenetics*. *Crustacean Issues* 18: 89-99.
- CLEVA R. 1990 — Crustacea Decapoda: Les genres et les espèces indo-ouest pacifiques de Stylodactylidae, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 71-136.
- CLEVA R. 1997 — Crustacea Decapoda: Stylodactylidae récoltés en Indonésie, aux îles Wallis et Futuna et au Vanuatu (campagnes KARUBAR, MUSORSTOM 7 et 8). Données complémentaires sur les Stylodactylidae de Nouvelle-Calédonie, in CROSNIER A. & BOUCHET P. (eds), Résultats des Campagnes MUSORSTOM Vol. 16. *Mémoires du Muséum national d'Histoire naturelle* 172: 385-407.
- CLEVA R. 2001 — Les Bathypalaemonellidae de Saint-Laurent, 1985 (Crustacea, Decapoda, Caridea) avec description d'une espèce nouvelle et définition d'un genre nouveau. *Zoosystema* 23: 757-782.
- CLEVA R. 2004 — Stylodactylidae and Bathypalaemonellidae from Taiwan (Crustacea, Decapoda, Caridea). *Raffles Bulletin of Zoology* 52: 497-511.
- CLEVA R. & CROSNIER A. 2006 — *Heterocarpus tenuidentatus*, a new species of shrimp from the Solomon Islands (Crustacea, Decapoda, Caridea, Pandalidae). *Zootaxa* 1200: 61-68.
- CROSNIER A. 1978 — Crustacés Décapodes pénéides Aristeidae (Benthescyminae, Aristeinae, Solenocerinae). *Faune de Madagascar* 46: 1-197.
- CROSNIER A. 1985a — Penaeoid shrimps (Benthescymidae, Aristeidae, Solenoceridae, Sicyoniidae) collected in Indonesia during the Corindon II and IV Expeditions. *Marine Research in Indonesia* 24: 19-47.
- CROSNIER A. 1985b — Crustacés Décapodes: Penaeidae. Les espèces indo-ouest-pacifiques du genre *Parapenaeus*, in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 2. *Mémoires du Muséum national d'Histoire naturelle* 133: 303-353.
- CROSNIER A. 1986 — Crevettes de la famille des Pandalidae récoltées durant ces dernières années en Polynésie française. Description de *Plesionika chacei* et *P. carsini* spp. nov. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 8: 361-377.
- CROSNIER A. 1987 — Les espèces indo-ouest-pacifiques d'eau profonde du genre *Metapenaeopsis* (Crustacea, Decapoda Penaeoidea). *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 9: 409-453.
- CROSNIER A. 1988a — Contribution à l'étude des genres *Haliporus* Bate, 1881 et *Gordonella* Tirmizi, 1960 (Crustacea Decapoda Penaeoidea). Description de deux espèces nouvelles. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 10: 563-601.
- CROSNIER A. 1988b — Les *Éupasiphæ* (Crustacea Decapoda Pasiphæidae) du sud-ouest de l'océan Indien. Description d'*E. paucidentata* sp. nov. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 10: 785-797.
- CROSNIER A. 1988c — Sur les *Heterocarpus* (Crustacea, Decapoda, Pandalidae) du sud-ouest de l'océan Indien. Remarques sur d'autres espèces ouest-pacifiques du genre et description de quatre taxa nouveaux. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 10: 57-103.
- CROSNIER A. 1989 — Benthescymidae, Aristeidae, Solenoceridae (Crustacea Penaeoidea), in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 5. *Mémoires du Muséum national d'Histoire naturelle* 144: 37-67.
- CROSNIER A. 1991 — Crustacea Decapoda: Les *Metapenaeopsis* indo-ouest pacifiques sans appareil stridulant (Penaeidae). Deuxième partie, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 9. *Mémoires du Muséum national d'Histoire naturelle*: 155-297.
- CROSNIER A. 1994a — Crustacea Decapoda: Les *Metapenaeopsis* indo-ouest-pacifiques avec un appareil stridulant (Penaeidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 12. *Mémoires du Muséum national d'Histoire naturelle* 161: 255-337.
- CROSNIER A. 1994b — Crustacea Decapoda: Observations complémentaires sur les *Metapenaeopsis* indo-ouest pacifiques sans appareil stridulant (Penaeidae). Description de deux espèces nouvelles, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 12. *Mémoires du Muséum national d'Histoire naturelle* 161: 339-349.
- CROSNIER A. 1994c — *Sphaerodromia lamellata*, espèce nouvelle de Nouvelle-Calédonie (Decapoda, Brachyura, Dromiidae). *Crustaceana* 67 (3): 341-347, pl. 1.
- CROSNIER A. 1994c — Crustacea Decapoda: Penaeoidea récoltés lors de la campagne KARUBAR en Indonésie, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 12. *Mémoires du Muséum national d'Histoire naturelle* 161: 351-365.
- CROSNIER A. 1999 — Un *Heterocarpus* nouveau (Crustacea, Decapoda, Pandalidae) du Pacifique Sud-Ouest. *Zoosystema* 21: 345-351.
- CROSNIER A. 2001 — Grapsidae (Crustacea, Decapoda, Brachyura) d'eau profonde du Pacifique sud-ouest. *Zoosystema* 23 (4): 783-796.
- CROSNIER A. 2002 — Révision du genre *Parathranites* Miers, 1886 (Crustacea, Brachyura, Portunidae). *Zoosystema* 24 (4): 799-825.
- CROSNIER A. 2003 — *Sicyonia* (Crustacea, Decapoda, Penaeoidea, Sicyoniidae) de l'Indo-ouest Pacifique. *Zoosystema* 25: 197-350.
- CROSNIER A. 2005 — Deux *Parapenaeus* nouveaux (Crustacea, Decapoda, Penaeidae) du sud-ouest Pacifique. *Zoosystema* 27: 257-266.
- CROSNIER A. & DALL W. 2004 — Redescription of *Hymenopenaeus obliquirostris* (Crustacea, Decapoda, Penaeoidea, Solenoceridae) and description of two new species of *Hymenopenaeus* from the Indo-West Pacific. *Zootaxa* 600: 1-26.

- CROSNIER A., MACHORDOM A. & BOISSELIER-DUBAYLE M. C. 2007 — Les espèces du genre *Trachypenaopsis* (Crustacea, Decapoda, Penaeidae). Approches morphologiques et moléculaires. *Zoosystema* 29 (3): 471-489.
- CROSNIER A. & NG P. K. L. 2004 — Remarques sur le genre *Intesius* (Crustacea, Decapoda, Brachyura, Goneplacidae) et description de deux espèces nouvelles. *Zoosystema* 26 (2): 263-277.
- DAVIE P. J. F. 1991 — Crustacea Decapoda: the genus *Platepistoma* Rathbun, 1906 (Cancridae) with the description of three new species, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 493-514.
- DAVIE P. J. F. 1993 — Deepwater xanthid crabs from French Polynesia (Crustacea, Decapoda, Xanthoidea). *Bulletin du Muséum national d'Histoire naturelle* (4) 14 (A2): 501-561, pls 1-13.
- DAVIE P. J. F. 1997 — Crustacea Decapoda: Deep water Xanthoidea from the South-Western Pacific and the Western Indian Ocean, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 337-387.
- DAVIE P. J. F. & CROSNIER A. 2006 — *Echinolatus* n. gen. (Crustacea, Decapoda, Portunidae) with description of two new species from the South-West Pacific, in RICHER DE FORGES B. & JUSTINE J. L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 393-410.
- DE GRAVE S., CHU K. H. & CHAN T.-Y. 2010 — On the systematic position of *Galatheacaris abyssalis* (Decapoda: Galatheacaridoidea). *Journal of Crustacean Biology* 30 (3): 521-527.
- DE GRAVE S., PENTCHEFF N. D., AHYONG S., CHAN T.-Y., CRANDALL K. A., DWORSCHAK P., FELDER D. L., FELDMANN R. M., FRANSEN C. H. J. M., GOULDING L. Y. D., LEMAÎTRE R., LOW M. L., MARTIN J. W., NG P. K. L., SCHWEITZER C. E., TANS H., TSHUDY D. & WETZER R. 2009 — A classification of Recent and fossil genera of decapod crustaceans. *Raffles Bulletin of Zoology* suppl 21: 1-109, colour figures 1-7.
- DESBRUYERES D. & SEGONZAC M. 2007 — *Handbook of deep-sea hydrothermal vent fauna*. Editions IFREMER, 279 p.
- DIAMOND E. A., RICHER DE FORGES B. & KORNICKER L. S. 2008 — *Azygocypriolina brynmawria* a new myodocopid ostracod off the southwest coast of New Caledonia (Crustacea; Ostracoda; Myodocopa; Cypridinidae). *Proceedings of the Biological Society of Washington* 121 (3): 354-364.
- DISTEL D. L., BACO A. R., CHUANG E., MORRILL W., CAVANAUGH C. & SMITH C. R. 2000 — Do mussels take wooden steps to deep-sea vents? *Nature* 403: 725-726.
- DUPERRON S., POTTIER M. A., LEGER N., PUILLANDRE N., LE PRIEUR S., SIGWART J., RAVAUX J. & ZBINDEN M. 2012 — A tale of two chitons: is habitat specialisation linked to distinct associated bacterial communities? *FEMS Microbiology Ecology* 83 (3): 552-567.
- DUPERRON S., LORION J., SAMADI S., GROS O. & GAILL F. 2009 — Symbioses between deep-sea mussels (Mytilidae: Bathymodiolinae) and chemosynthetic bacteria: diversity, function and evolution. *Comptes Rendus de l'Académie des Sciences* 332: 298-310.
- DUPONT J., MAGNIN S., ROUSSEAU F., ZBINDEN M., FREBOURG G., SAMADI S., RICHER DE FORGES B. & JONES E. B. G. 2009 — Molecular and ultrastructural characterization of two ascomycetes found on sunken wood off Vanuatu Islands in the deep Pacific Ocean. *Mycological Research* 113: 1351-1364.
- FOREST J. 1981 — Compte rendu et remarques générales, in DORST J. (ed.), Résultats des campagnes MUSORSTOM Vol. 1. *Mémoires ORSTOM* 91: 9-50.
- FOREST J. 1987 — Les Pylochelidae ou "Pagures symétriques" (Crustacea Coenobitoidea), in DORST J. (ed.), Résultats des campagnes MUSORSTOM Vol. 1. *Mémoires du Muséum national d'Histoire naturelle* 137: 1-254.
- FOREST J. 1995 — Crustacea Decapoda Anomura: Révision du genre *Trizopagurus* Forest, 1952 (Diogenidae), avec l'établissement de deux genres nouveaux, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 13. *Mémoires du Muséum national d'Histoire naturelle* 163: 9-149.
- FOREST J. 1997 — Présence du genre *Xylopagurus* A. Milne-Edwards, 1880 (Crustacea, Decapoda, Paguridae) dans l'Indo-Ouest Pacifique, avec la description de deux espèces nouvelles. *Zoosystema* 19: 421-435.
- FOREST J. 2006a. - *Laurentaglypheia*, un nouveau genre pour la seconde espèce actuelle de Glypheide récemment découverte (Crustacea Décapoda Glypheidae). *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*, Paris, 329: 841-846.
- FOREST J. 2006b — The recent glypheids and their relationship with their fossil relatives (Decapoda, Reptantia). *Crustaceana* 79 (7): 795-820.
- FOREST J. & SAINT LAURENT DE M. 1981 — La morphologie externe de *Neoglypheia inopinata*, espèce actuelle de Crustacé Décapode Glypheide, in DORST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 1. *Mémoires ORSTOM* 91: 51-84.
- FOREST J. & SAINT LAURENT DE M. 1989 — Nouvelle contribution à la connaissance de *Neoglypheia inopinata* Forest & de Saint Laurent, à propos de la description de la femelle adulte, in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 5. *Mémoires du Muséum national d'Histoire naturelle* 144: 75-92.
- FUJITA T., STAMPANATO S. & JANGOUX M. 1994 — *Belyaevostella hyugaensis*, a New Species of Deep-sea Asteroid (Asteroidea, Caymanostellidae) found on a Sunken Wood from off Southern Japan. *Bulletin of the National Science Museum (A) Zoology* 20: 183-188.
- GALIL B. S. 1997 — Crustacea Decapoda: A revision of the Indo-Pacific species of genus *Calappa* Weber, 1795 (Calappidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 271-335.
- GALIL B. S. 2000 — Crustacea Decapoda: A revision of the family Polychelidae, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 285-387.
- GALIL B. S. 2003 — Four new genera of leucosiid crabs (Crustacea: Brachyura: Leucosiidae) for three new species and nine species previously in the genus *Randallia* Stimpson, 1857, with a redescription of the type species, *R. ornata* (Randall, 1939). *Proceedings of the Biological Society of Washington* 116 (2): 395-422.
- GALIL B. S. & NG P. K. L. 2009 — Calappoidea and Leucosioidea (Crustacea: Decapoda: Brachyura) from Luzon, Philippines, with descriptions of two new species of *Mursia*. *Zootaxa* 2085: 45-60.
- GALIL B. S. & NG P. K. L. 2010 — On a collection of calappoid and leucosiod crabs (Decapoda, Brachyura) from Vanuatu, with description of a new species of Leucosiidae, in CASTRO P., DAVIE P. J. F., NG P. K. L. & RICHER DE FORGES B. (eds), Studies on Brachyura: a homage to Danièle Guinot. *Crustacean Monographs* 11: 139-152.
- GUINOT D. 1977 — Propositions pour une nouvelle classification des Crustacés Décapodes Brachyures. *Comptes rendus hebdomadaires des Séances de l'Académie des Sciences*, Paris 285: 1049-1052.
- GUINOT D. 1989a — Le genre *Carcinoplax* H. Milne-Edwards, 1852 (Crustacea, Brachyura: Goneplacidae), in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 5. *Mémoires du Muséum national d'Histoire naturelle* 144: 265-345.

- GUINOT D. 1989b — Les genres *Trachycarinus* Faxon et *Trichopeltarion* A. Milne-Edwards (Crustacea, Brachyura: Atelecyclidae), in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 5. *Mémoires du Muséum national d'Histoire naturelle* 144: 347-385, pls 1-5.
- GUINOT D. 1990 — Crustacea Decapoda: Le genre *Psopheticus* Wood-Mason, 1892 (Goneplacidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 145: 331-367.
- GUINOT D. 1991 — Établissement de la famille des Poupinidae pour *Poupinia hirsuta* gen. nov., sp. nov. de Polynésie (Crustacea Decapoda Brachyura Homoloidea). *Bulletin du Muséum national d'Histoire naturelle* (A) 12 (3-4): 577-605, pls 1-3.
- GUINOT D. 1995 — Crustacea Decapoda Brachyura: Révision de la famille des Homolodromiidae Alcock, 1900, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 13. *Mémoires du Muséum national d'Histoire naturelle* 163: 155-282.
- GUINOT D. 1993 — Données nouvelles sur les Crabes primitifs (Crustacea Decapoda Brachyura Podotremata). *Comptes rendus de l'Académie des Sciences* (III), 316 (10): 1225-1232, figs. 1-8.
- GUINOT D. 2008 — A re-evaluation of the Dynomenidae Ortmann, 1892 (Crustacea, Decapoda, Brachyura, Podotremata), with the recognition of four subfamilies. *Zootaxa* 1850: 1-26.
- GUINOT D., JAMIESON B. G. M & RICHER DE FORGES B. 1994 — Relationship of Homolidae and Dromiidae: Evidence from Spermatozoal Ultrastructure (Crustacea, Decapoda). *Acta Zoologica* 75 (3): 255-267.
- GUINOT D. & RICHER DE FORGES B. 1981a — Crabes de profondeur, nouveaux ou rares, de l'Indo-Pacifique (Crustacea, Decapoda, Brachyura) (Première partie). *Bulletin du Muséum national d'Histoire naturelle* (4) 2 (1980) section A (4): 1113-1153.
- GUINOT D. & RICHER DE FORGES B. 1981b — Crabes de profondeur, nouveaux ou rares, de l'Indo-Pacifique (Crustacea, Decapoda, Brachyura) (Deuxième partie). *Bulletin du Muséum national d'Histoire naturelle* 3 (1981) (A) 11: 227-260, pls 1-7.
- GUINOT D. & RICHER DE FORGES B. 1981c — Homolidae, rares ou nouveaux, de l'Indo-Pacifique (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle* (A) 8 (2): 523-581, pls 1-8.
- GUINOT D. & RICHER DE FORGES B. 1982a — Nouvelles récoltes des genres *Cyrtomaia* Miers et *Pleistacantha* Miers (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 3 (4): 1087-1125, pls I-IV.
- GUINOT D. & RICHER DE FORGES B. 1982b — Révision du genre Indo-Pacifique *Cyrtomaia* Miers, 1886: Campagnes océanographiques du Challenger, de l'Albatross, du Siboga et du Vauban (Crustacea Decapoda Brachyura). *Annales de l'Institut Océanographique New Series* 58 (1): 5-87.
- GUINOT D. & RICHER DE FORGES B. 1986 — Crustacés Décapodes: Majidae (genres *Platymaia*, *Cyrtomaia*, *Pleistacantha*, *Sphenocarcinus* et *Naxioides*), in Résultats des Campagnes MUSORSTOM I et II -Philippines (1976, 1980). Tome 2. *Mémoires du Muséum National d'Histoire Naturelle* 133: 83-179, pls 1-9.
- GUINOT D. & RICHER DE FORGES B. 1995 — Crustacea Decapoda Brachyura: Révision de la famille des Homolidae de Haan, 1839, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 13. *Mémoires du Muséum national d'Histoire naturelle* 163: 283-517.
- HAYASHI K. I. 1999 — Crustacea Decapoda: Revision of *Pasiphaea sivado* (Risso, 1816) and related species, with descriptions of one new genus and five new species (Pasiphaeidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 20. *Mémoires du Muséum national d'Histoire naturelle* 180: 267-302.
- HAYASHI K. I. 2004 — Revision of the *Pasiphaea cristata* Bate, 1888 species group of *Pasiphaea* Savigny, 1816, with descriptions of four new species, and referral of *P. australis* Hanamura, 1989 to *Alainopasiphaea* Hayashi, 1999 (Crustacea, Decapoda, Pasiphaeidae), in MARSHALL B. A. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos Vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 319-373.
- HAYASHI K. I. 2006 — Revision of the *Pasiphaea alcocki* species group (Crustacea, Decapoda, Pasiphaeidae), in RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 193-241.
- HEADS M. 2005 — Towards a panbiogeography of the seas. *Biological Journal of the Linnean Society* 84: 675-723.
- HEADS M. 2008 — Panbiogeography of New Caledonia, south-west Pacific: Basal angiosperms on basement terranes, ultramafic endemics inherited from volcanic island arcs and old taxa endemic to young islands. *Journal of Biogeography* 35: 2153-2175.
- HO P.H., NG P.K. L., CHAN T.-Y. & LEE D. A. 2004 — New records of 31 species of brachyuran crabs from the joint Taiwan-France Expeditions, "TAIWAN 2000" and "TAIWAN 2001", off deep waters in Taiwan. *Crustaceana* 77 (6): 641-668.
- HOEKSEMA B. W. 2007 — Delineation of the Indo-Malayan Centre of Maximum Marine Biodiversity: The Coral Triangle, in RENEMA W. (ed.), *Biogeography, Time, and Place: Distributions, Barriers, and Islands*. Springer: 117-178.
- HOLTHUIS L. B. 2002 — The Indo-Pacific scyllarine lobsters (Crustacea: Decapoda: Scyllaridae). *Zoosystema* 24 (3): 499-683.
- HOYOUX C. 2010 — Crustacés décapodes de bois coulés en océan profond: régimes alimentaires et symbioses microbiennes. *PhD thesis, University of Liege*, Belgium. 347 pp.
- HOYOUX C., ZBINDEN M., SAMADI S., GAILL & COMPÈRE P. 2009 — Wood-based diet and gut microflora of a galatheid crab associated with Pacific deep-sea wood falls. *Marine Biology* 156: 2421-2439.
- HOYOUX C., ZBINDEN M., SAMADI S., GAILL & COMPÈRE P. 2012 — Diet and gut microorganisms of *Munidopsis* squat lobsters associated with natural woods and mesh-enclosed substrates in the deep South Pacific. *Marine Biology Research* 8: 28-47.
- JAMIESON B. G. M., GUINOT D. & RICHER DE FORGES B. 1995 — Phylogeny of the brachyura: Evidence from spermatozoal ultrastructure (Crustacea, Decapoda), in JAMIESON B. G. M., AUSIO J. & JUSTINE J.-L. (eds), Advances in spermatozoal Phylogeny and taxonomy. *Mémoires du Muséum national d'Histoire naturelle* 166: 265-283.
- KIM J. N & CHANT T.-Y. 2005 — A revision of the genus *Prionocrangon* (Crustacea: Decapoda: Caridea: Crangonidae). *Journal of Natural History* 39 (19): 1597-1625.
- KOMAIT. 1997 — A review of the *Metacrangon jacqueti* group, with descriptions of two new species (Decapoda, Caridea, Crangonidae). *Zoosystema* 19: 651-681.
- KOMAI T. 2004 — A review of the Indo-West Pacific species of the genus *Glyphocrangon* A. Milne-Edwards, 1881 (excluding the *G. caeca* species group) (Crustacea, Decapoda, Caridea, Glyphocrangonidae), in MARSHALL B. A. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos Vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 375-610.
- KOMAI T. 2006a — A review of the crangonid genus *Lissosabinea* Christoffersen, 1988 (Crustacea, Decapoda, Caridea), with descriptions of three new species from the western Pacific. *Zoosystema* 28: 31-59.
- KOMAI T. 2006b — Revision of the *Glyphocrangon caeca* species group (Crustacea, Decapoda, Glyphocrangonidae), in RICHER DE FORGES B.

- & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 243-264.
- KOMAI T. 2007 — A new species of *Glyphocrangon* (Crustacea, Decapoda, Caridea, Glyphocrangonidae) from the Austral Islands, French Polynesia. *Zoosystema* 29: 565-573.
- KOMAI T. 2008 — A world-wide revision of species of the deep-water crangonid genus *Parapontophilus* Christoffersen, 1988 (Crustacea, Decapoda, Caridea), with descriptions of ten new species. *Zoosystema* 30: 261-332.
- KOMAI T. 2010 — New species and new records of the hermit crab genus *Pagurixus* Melin, 1939 (Crustacea: Decapoda: Anomura: Paguridae) from the Indo-West Pacific. *Journal of Natural History* 44: 21: 1269-1342.
- KOMAI T. 2011 — Further records of deep-sea shrimps of the genus *Glyphocrangon* (Crustacea: Decapoda: Caridea: Glyphocrangonidae) from the Southwestern Pacific, with descriptions of two new species. *Species Diversity* 16: 113-135.
- KOMAI T. 2012 — A review of the western Pacific species of the crangonid genus *Metacrangon* Zarenkov, 1965 (Decapoda: Caridea), with descriptions of seven new species. *Zootaxa* 3468: 1-77.
- KOMAI T. & CHANT T.-Y. 2006 — A new species of the hermit crab genus *Pagurus* (Crustacea: Decapoda: Anomura: Paguridae) from deep waters off Taiwan. *Proceedings of the Biological Society of Washington* 119 (2): 239-250.
- KOMAI T. & CHAN T.-Y. 2008 — Further records of deep-sea shrimps of the genus *Glyphocrangon* A. Milne-Edwards, 1881 (Crustacea: Decapoda: Caridea) from the Philippines, with descriptions of three new species. *Raffles Bulletin of Zoology*, Supplement 19: 39-62.
- KOMAI T. & CHANT T.-Y. 2010 — Two new pandalid shrimps and the discovery of the second specimen of the rare hippolytid shrimp *Leontocaris bulga* Taylor & Poore, 1998 (Crustacea, Decapoda) from the Mozambique MAINBAZA cruise. *Zoosystema* 32 (4): 625-641.
- KOMAI T. & CHAN T.-Y. 2013 — New records of *Glyphocrangon* A. Milne-Edwards, 1881 (Crustacea, Decapoda, Caridea, Glyphocrangonidae) from recent French expeditions off the Mozambique Channel and Papua New Guinea, with description of one new species, in AHYONG S.T., CHANT T.-Y., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 107-128.
- KOMAI T. & MOTOH H. 2012 — The identity of *Pilumnus doylei* Bals, 1933 (Crustacea: Decapoda: Brachyura: Pilumnidae), with descriptions of three new species from the Western Pacific. *Zootaxa* 3305: 1-27.
- KOMAI T. & POUPIN J. 2012 — Two new species of shallow-water hermit crabs (Crustacea: Decapoda: Paguridae) from Mayotte, Comoro Islands, southwestern Indian Ocean. *Zootaxa* 3277: 56-68.
- KOMAI T. & SAITO T. 2006 — A new genus and two new species of Spongicolidae (Crustacea: Decapoda: Stenopodidea) from the South-West-Pacific, in RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 265-284.
- KOSLOW T. 2007 — *The Silent Deep. The Discovery, Ecology and Conservation of the Deep Sea*. UNSW Press, Sydney: 270 pp.
- KWONG S., SRIVATHSAN A. & MEIER R. 2012 — An update on DNA barcoding: low species coverage and numerous unidentified sequences. *Cladistics* (DOI: 10.1111/j.1096-0031.2012.00408.x).
- LAI J. C. Y., AHYONG S. T., JENG M. S. & NG P. K. L. 2009 — Are coral-dwelling crabs monophyletic? A phylogeny of the Trapezioidae (Crustacea: Decapoda: Brachyura). *Invertebrate Systematics* 23: 402-408.
- LAI J. C. Y., MENDOZA J. C. E., GUINOT D., CLARK P. F. & NG P. K. L. 2011 — Xanthidae Macleay, 1838 (Decapoda: Brachyura: Xanthoidea) systematics: a multi-gene approach with support from adult and zoeal morphology. *Zoologischer Anzeiger* 250: 407-448.
- LARSEN K. & KRAPP-SCHICKEL T. 2007 — Amphipoda (Crustacea: Peracarida) from chemically reduced habitats: the hydrothermal vent system of the north-east Pacific. Part II. Melitidae and Eusiridae. *Journal of the Marine Biological Association of the UK* 87: 1207-1217.
- LAUBIER L. 2008 — *Ténèbres océaniques. Le triomphe de la vie dans les abysses*. Buchet & Chastel, Ecologie, 298 p.
- LEMAITRE R. 1993 — A new genus of Parapaguridae (Decapoda: Anomura). *Crustacean Research* 22: 11-20.
- LEMAITRE R. 1994 — Crustacea Decapoda: Deep-water hermit crabs (Parapaguridae) from French Polynesia with descriptions of four new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM Vol. 12. Mémoires du Muséum national d'Histoire naturelle* 161: 375-419.
- LEMAITRE R. 1997 — Crustacea Decapoda: Parapaguridae from the KARUBAR cruise in Indonesia, with descriptions of two new species, in CROSNIER A. & BOUCHET P. (eds), *Résultats des Campagnes MUSORSTOM Vol. 16. Mémoires du Muséum national d'Histoire naturelle* 172: 573-596.
- LEMAITRE R. 1998 — A new species of hermit crab of the family Parapaguridae (Decapoda, Anomura) from French Polynesia. *Zoosystema* 20 (1): 101-107.
- LEMAITRE R. 1999 — Crustacea Decapoda: A review of the species of the genus *Parapagurus* Smith, 1879 (Parapaguridae) from the Pacific and Indian Oceans, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM Vol. 20. Mémoires du Muséum national d'Histoire naturelle* 180: 303-378.
- LEMAITRE R. 2003 — A new genus and species of hermit crab (Crustacea: Anomura: Paguridae) from Taiwan. *Memoirs of Museum Victoria* 60 (1): 105-110.
- LEMAITRE R. 2004a — A review of *Strobopagurus* Lemaître, 1989 (Crustacea: Decapoda: Paguroidea: Parapaguridae), with description of a new species. *Scientia Marina* 68 (3): 355-372.
- LEMAITRE R. 2004b — Crustacea Decapoda: A worldwide review of the hermit crab species of the genus *Sympagurus* Smith, 1883, in MARSHALL B. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos Vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 85-149.
- LEMAITRE R. 2004c — Discovery of the first hermit crab (Crustacea: Decapoda: Parapaguridae) associated with hydrothermal vents. *Cahiers de Biologie marine*, 45: 325-334.
- LEMAITRE R. 2006 — Two new species of Parapaguridae (Crustacea, Anomura, Paguroidea) with subconical corneas, and new data on biology of some rare species. *Zoosystema* 28 (2): 517-532.
- LEMAITRE R. 2013 — The genus *Paragiopagurus* Lemaître, 1996 (Crustacea, Decapoda, Anomura, Paguroidea, Parapaguridae): A worldwide review of the species, with descriptions of five new species, in AHYONG S. T., CHANT T.-Y., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 311-422.
- LEMAITRE R. & POUPIN J. 2003 — A strikingly coloured new species of *Paragiopagurus* Lemaître, 1996 (Crustacea: Decapoda: Anomura: Parapaguridae) from French Polynesia. *Zootaxa* 386: 1-11.
- LEMAITRE R. & MCLAUGHLIN P. A. 1995 — *Alainopagurus crosnieri*, gen. et sp. nov. (Decapoda: Anomura: Paguridae) from the Western Pacific. *Bulletin du Muséum national d'Histoire naturelle* (4) 17A (3-4): 273-282.
- LEMAITRE R. & MCLAUGHLIN P. A. 2009 — Recent advances and conflicts in concepts of anomuran phylogeny (Crustacea: Malacostraca). *Arthropod Systematics & Phylogeny* 67: 119-135.

- LEMAITRE R., MCLAUGHLIN P. A. & SORHANNUS U. 2009 — Phylogenetic relationships within the Pylochelidae (Decapoda: Anomura: Paguroidea): a cladistic analysis based on morphological characters. *Zootaxa* 2022: 1-14.
- LI C. P., DE GRAVE S., CHAN T.-Y., LEI H. C. & CHU K. H. 2011 — Molecular systematics of caridean shrimps based on five nuclear genes: implications for superfamily classification. *Zoologischer Anzeiger* 250: 270-279.
- LI X. & CHAN T.-Y. 2013 — Pandalid shrimps (Crustacea, Decapoda, Caridea) collected from the Philippines PANGLAO 2005 deep-sea expedition, *in* AHYONG S. T., CHAN T.-Y., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos. Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 129-154.
- LI X. 2006 — Additional pandaloid shrimps from the South China Sea (Crustacea: Decapoda: Caridea), with description of one new species. *Raffles Bulletin of Zoology* 54: 361-372.
- LI X., MITSUHASHI M. & CHANT-T.Y. 2008 — Deep-sea pontoniines (Decapoda: Palaemonidae) from the Philippine "PANGLAO 2005" expedition, with descriptions of four new species. *Journal of Crustacean Biology* 28 (2): 385-411.
- LIN C.-W. & CHANT-T.Y. 2005 — A new squat lobster, *Munida rupicola* (Crustacea: Decapoda: Galatheidae), from Taiwan. *Proceedings of the Biological Society of Washington* 118: 237-243.
- LIN C.-W., CHANT-T.Y. & CHU K. H. 2004 — A new squat lobster of the genus *Raymunida* (Decapoda: Galatheidae) from Taiwan. *Journal of Crustacean Biology* 24: 149-156.
- LIN C.-W., OSAWA M. & CHAN T.-Y. 2007 — A new *Munidopsis* (Crustacea: Decapoda: Galatheidae) associated with gorgonian corals from the deep waters off Taiwan. *Proceedings of the Biological Society of Washington* 120: 167-174.
- LIN C.-W. & CHAN T.-Y. 2011 — Two new deep-sea squat lobsters of the genus *Munidopsis* Whiteaves, 1874 (Crustacea: Decapoda: Munidopsidae) from Taiwan. *Zootaxa* 2754: 51-59.
- LORION J., DUPERRON S. B., GROS O., CRUAUD C. & SAMADI S. 2009 — Several deep-sea mussels and their associated symbionts are able to live both on wood and on whale falls. *Proceedings of the Royal Society B: Biological Sciences* 276: 177-185.
- LORION J., BUGE B., CRUAUD C. & SAMADI S. 2010 — New insights into diversity and evolution of deep-sea Mytilidae (Mollusca: Bivalvia). *Molecular Phylogenetics and Evolution* 57: 71-83.
- LOWRY J. K. & DEMPSEY K. 2006 — The giant deep-sea scavenger genus *Bathynomus* (Crustacea, Isopoda, Cirolanidae) in the Indo-West Pacific, *in* RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 163-192.
- MA K. Y., CHAN T.-Y. & CHU K. H. 2009 — Phylogeny of penaeoid shrimps (Decapoda: Penaeoidea) inferred from nuclear protein-coding genes. *Molecular Phylogenetics and Evolution* 53: 45-55.
- MACHORDOM A. & MACPHERSON E. 2004 — Rapid radiation and cryptic speciation in galatheid crabs of the genus *Munida* and related genera in the South West Pacific: molecular and morphological evidence. *Molecular Phylogenetics and Evolution* 33: 259-279.
- MACPHERSON E. 1988 — Lithodid crabs (Crustacea, Decapoda, Lithodidae) from Madagascar and La Reunion (SW Indian Ocean). *Bulletin du Muséum national d'Histoire naturelle* (4) (10) (A) 1: 117-133.
- MACPHERSON E. 1990a — Crustacea Decapoda: On some species of Lithodidae from the Western Pacific, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 217-226.
- MACPHERSON E. 1990b — Crustacea Decapoda: On a collection of Nephropidae from the Indian Ocean and Western Pacific, *in* CROSNIER, A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 289-329.
- MACPHERSON E. 1991 — A new species of the genus *Lithodes* (Crustacea, Decapoda, Lithodidae) from the French Polynesia. *Bulletin du Muséum national d'Histoire naturelle* (4) (13), A (1-2): 153-158.
- MACPHERSON E. 1993a — Crustacea Decapoda: species of the genus *Munida* Leach, 1820 (Galatheidae) collected during the MUSORSTOM and CORINDON cruises in the Philippines and Indonesia, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 421-442.
- MACPHERSON E. 1993b — Crustacea Decapoda: species of the genus *Paramunida* Baba, 1988 (Galatheidae) from the Philippines, Indonesia and New Caledonia, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 443-473.
- MACPHERSON E. 1994 — Crustacea Decapoda: Studies on the genus *Munida* Leach, 1820 (Galatheidae) in New Caledonia and adjacent waters with descriptions of 56 new species, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 12. *Mémoires du Muséum national d'Histoire naturelle* 161: 421-569.
- MACPHERSON E. 1996a — Crustacea Decapoda: species of the genera *Munida* Leach, 1820 and *Paramunida* Baba, 1988 (Galatheidae) from the seas around the Wallis and Futuna Islands, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 387-421.
- MACPHERSON E. 1996b — Crustacea Decapoda: new records of species of the genera *Munida* Leach, 1820 and *Paramunida* Baba, 1988 (Galatheidae) from New Caledonia, with the descriptions of three new species, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 423-431.
- MACPHERSON E. 1997 — Crustacea Decapoda: species of the genera *Agononida* Baba & de Saint Laurent, 1996 and *Munida* Leach, 1820 (Galatheidae) from the KARUBAR cruise, *in* CROSNIER A. & BOUCHET P. (eds), Résultats des Campagnes MUSORSTOM Vol. 16. *Mémoires du Muséum national d'Histoire naturelle* 172: 597-612.
- MACPHERSON E. 1998 — A new genus of Galatheidae (Crustacea, Anomura) from the western Pacific Ocean. *Zoosystema* 20: 351-355.
- MACPHERSON E. 1999a — Crustacea Decapoda: Species of the genera *Agononida* Baba & de Saint Laurent, 1996 and *Munida* Leach, 1820 (Galatheidae) collected during the MUSORSTOM 8 cruise in Vanuatu, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 20. *Mémoires du Muséum national d'Histoire naturelle* 180: 407-426.
- MACPHERSON E. 1999b — Three new species of the genus *Munida* Leach, 1820 (Decapoda, Galatheidae) from the Seychelles Islands (Indian Ocean). *Zoosystema* 21: 473-482.
- MACPHERSON E. 2000 — Crustacea Decapoda: species of the genera *Crosnierita* Macpherson, 1998, *Munida* Leach, 1820, and *Paramunida* Baba, 1988 (Galatheidae) collected during the MUSORSTOM 9 cruise to the Marquesas Islands, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 415-423.
- MACPHERSON E. 2001 — New species and new records of lithodid crabs (Crustacea, Decapoda, Lithodidae) from the Southwestern and Central Pacific Ocean. *Zoosystema* 23: 797-805.

- MACPHERSON E. 2003 — Some lithodid crabs (Crustacea, Decapoda, Lithodidae) from the Salomon Islands (SW Pacific Ocean) with the description of a new species. *Scientia Marina* 67: 413-418.
- MACPHERSON E. 2004 — Species of the genus *Munida* Leach, 1820 and related genera from Fiji and Tonga (Crustacea: Decapoda: Galatheidae), in MARSHALL B. A. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos Vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 231-292.
- MACPHERSON E. 2006a — Galatheidae (Crustacea, Decapoda) from the Austral Islands, Central Pacific, in RICHER DE FORGES B. & JUSTINE J. L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 285-333.
- MACPHERSON E. 2006b — New species and new occurrence of Galatheoidea (Crustacea, Decapoda) from New Caledonia. *Zoosystema* 28: 669-681.
- MACPHERSON E. 2007 — Species of the genus *Munidopsis* Whiteaves, 1784 from the Indian and Pacific Oceans and reestablishment of the genus *Galacantha* A. Milne-Edwards, 1880 (Crustacea, Decapoda, Galatheidae). *Zootaxa* 1417: 1-135.
- MACPHERSON E. 2009 — New species of squat lobsters of the genera *Munida* and *Raymunda* (Decapoda, Galatheidae) from Vanuatu and New Caledonia. *Zoosystema* 31: 431-451.
- MACPHERSON E. 2012 — New deep-sea squat lobsters of the genus *Galathea* Fabricius, 1793 (Decapoda, Galatheidae) from Vanuatu and New Caledonia. *Zoosystema* 34: 409-427.
- MACPHERSON E. & BABA K. 1993 — Crustacea Decapoda: *Munida japonica* Stimpson, 1858, and related species (Galatheidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 381-420.
- MACPHERSON E. & BABA K. 2006a — New species and records of small galatheids (Crustacea, Decapoda, Galatheidae) from the southwest and central Pacific Ocean. *Zoosystema* 28: 443-456.
- MACPHERSON E. & BABA K. 2009 — New species of squat lobsters of the genera *Agononida* and *Paramunida* (Crustacea: Decapoda: Anomura: Galatheidae) from the western Pacific. *Zootaxa* 2024: 56-68.
- MACPHERSON E. & BABA K. 2010 — Revision of the genus *Sadyoshia* (Crustacea: Anomura: Galatheidae) with description of three new species, in FRANSEN C. H. J. M., DE GRAVE S. & NG P. K. L. (eds), Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume. *Crustaceana Monographs* 14: 415-452.
- MACPHERSON E. & CHANT-Y. 2008 — Some lithodid crabs (Crustacea: Decapoda: Lithodidae) from Taiwan and adjacent waters, with the description of one new species from Guam. *Zootaxa* 1924: 43-52.
- MACPHERSON E. & CLEVA R. 2010 — Shallow-water squat lobsters (Crustacea, Decapoda, Galatheidae) from Mayotte (Comoros Island), La Réunion and Madagascar, with the description of a new genus and two new species. *Zootaxa* 2612: 57-68.
- MACPHERSON E., LEMAITRE R., RICHER DE FORGES B. & MANNING R. B. (eds) 1998 — Les "Crosnier's cronies": par la fenêtre du grenier. Crosnier's cronies and their view from the attic window. *Zoosystema* 20 (2): 139-141.
- MACPHERSON E. & MACHORDOM A. 2000 — *Raymunda*, new genus (Decapoda: Anomura: Galatheidae) from the Indian and Pacific Oceans. *Journal of Crustacean Biology* 20: 253-258.
- MACPHERSON E. & MACHORDOM A. 2001 — Phylogenetic relationships of species of *Raymunda* (Decapoda: Galatheidae) based on morphology and mitochondrial cytochrome oxidase sequences, with the recognition of four new species. *Journal of Crustacean Biology* 21: 696-714.
- MACPHERSON E. & MACHORDOM A. 2005 — Use of morphological and molecular data to identify three new sibling species of the genus *Munida* Leach, 1820 (Crustacea, Decapoda, Galatheidae) from New Caledonia. *Journal of Natural History* 39: 819-834.
- MACPHERSON E., RICHER DE FORGES B., SCHNABEL K. & SAMADI S. 2010 — Biogeography of the Pacific Galatheids. *Deep-Sea Research I*. 57: 228-238.
- MANUEL-SANTOS M. R. & NG P. K. L. 2007 — On the genus *Ladomedaeus* Števčić, 2005, from the Philippines and Japan, and the status of the Ladomedaeidae Števčić, 2005 (Decapoda: Brachyura: Xanthoidea). *Raffles Bulletin of Zoology* suppl. 16: 177-185.
- MCCAIN C. R. 2007 — Seamounts: identity crisis or split personality? *Journal of Biogeography* 34 (12): 2001-2184.
- MCLAUGHLIN P. A. 1997 — Crustacea Decapoda: Hermit crabs of the family Paguridae from the KARUBAR cruise in Indonesia, in CROSNIER A. & BOUCHET P. (eds), Résultats des Campagnes MUSORSTOM Vol. 16. *Mémoires du Muséum national d'Histoire naturelle* 172: 433-572.
- MCLAUGHLIN P. A. 1998 — Hermit crabs of the genus *Nematopagurus* (Crustacea, Decapoda, Paguridae) from south-eastern South Africa and Madagascar: new records and new species. *Zoosystema* 20 (2): 315-338.
- MCLAUGHLIN P. A. 2000 — Crustacea: Decapoda: *Porcellanopagurus* Filhol and *Solitariopagurus* Türky (Paguridae), from the New Caledonia area, Vanuatu, and the Marquesas: new records, new species, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 389-414.
- MCLAUGHLIN P. A. 2003 — Reassignment and redescription of "*Eupagurus*" *microps* Balss, 1911 (Crustacea, Decapoda, Paguridae), with notes on *Bathypaguropsis kuroshioensis* (Miyake, 1978). *Zoosystema* 25: 635-642.
- MCLAUGHLIN P. A. 2004a — Redescription of *Tomopaguroides valdiviae* (Balss, 1911) (Crustacea, Decapoda, Anomura, Paguroidea, Paguridae) with notes on variation and female morphology. *Zoosystema* 26: 469-481.
- MCLAUGHLIN P. A. 2004b — Crustacea Decapoda: A review of the hermit crab genus *Nematopagurus* A. Milne-Edwards & Bouvier, 1892 and the descriptions of five new species (Crustacea: Decapoda: Paguridae), in MARSHALL B. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos Vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 151-229.
- MCLAUGHLIN P. A. 2007a — A new species of *Pteropagurus* McLaughlin & Rahayu, 2006 and a new genus and species of scaphopod dweller (Crustacea, Decapoda, Anomura: Paguroidea, Paguridae). *Zoosystema* 29 (3): 503-513.
- MCLAUGHLIN P. A. 2007b — New records and a new species in the genus *Turleania* McLaughlin, 1997. (Crustacea, Decapoda, Anomura, Paguridae) from MUSORSTOM cruises, with a key to species. *Zoosystema* 29 (3): 583-593.
- MCLAUGHLIN P. A. & FOREST J. 1997 — Crustacea Decapoda: *Diacanthurus* gen. nov., a new genus of hermit crabs (Paguridae) with both Recent and fossil representation, and the descriptions of two new species, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 236-259.
- MCLAUGHLIN P. A. & LEMAITRE R. 2004 — The discovery of two new species of *Lithopagurus* Provenzano, 1968 (Crustacea, Decapoda, Anomura, Paguroidea, Paguridae) and the first records of the genus in the western Pacific. *Zoosystema* 26 (3): 483-494.
- MCLAUGHLIN P. A. & LEMAITRE R. 2008 — Larvae of two species of *Trizochelus* (Decapoda: Anomura: Paguroidea: Pylochelidae: Trizochelinae), description of the adult of one, and preliminary implications of development on pylochelid phylogeny. *Zootaxa* 1911: 52-68.

- MCLAUGHLIN P. A. & RAHAYU D. L. 2006 — A new genus with two new species of hermit crabs (Crustacea, Decapoda, Paguroidea, Paguridae) from an unique habitat. *Zootaxa* 1116: 55-68.
- MCLAUGHLIN P. A. & RAHAYU D. L. 2008a — *Pteropagurus* and *Catapagurus* (Decapoda, Anomura, Paguridae): resource sharing or “any port in a storm?” *Zoosystema* 30 (4): 899-916.
- MCLAUGHLIN P. A. & RAHAYU D. L. 2008b — A new genus and species of hermit crab of the family Paguridae (Crustacea: Anomura: Paguroidea) from the Vanuatu Archipelago. *Proceedings of the Biological Society of Washington* 121: 365-373.
- MCLAUGHLIN P. A., LEMAITRE R. & SORHANNUS U. 2007 — Hermit crab phylogeny: A reappraisal and its “fall out”. *Journal of Crustacean Biology* 27: 97-115.
- MCLAY C. L. 1999 — Crustacea Decapoda: Revision of the Family Dynomenidae, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM Vol. 20. Mémoires du Muséum national d’Histoire naturelle* 180: 427-569.
- MCLAY C. L. 2001 — Dynomenidae and Dromiidae crabs (Decapoda: Brachyura) from Guam, Philippines Islands, Tonga and Samoa. *Zoosystema* 23 (4): 807-856.
- MCLAY C. L. 2006 — Retroplumidae (Crustacea, Decapoda) from the Indo-Malayan archipelago (Indonesia, Philippine) and the Melanesian arc islands (Solomon Islands, Fiji and New Caledonia), and paleogeographical comments in: RICHER DE FORGES B. & JUSTINE J. L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d’Histoire naturelle* 193: 375-391.
- MCLAY C. L. & NG P. K. L. 2004 — A taxonomic revision of the genus *Paradynomene* Sakai, 1963 (Crustacea: Decapoda: Brachyura: Dynomenidae). *Zootaxa* 657: 1-24.
- MCLAY C. L. & NG P. K. L. 2005 — On a collection of Dromiidae and Dynomenidae from the Philippines, with description of a new species of *Hirsutodynamene* McLay, 1999 (Crustacea: Decapoda: Brachyura). *Zootaxa* 1029: 1-30.
- MCLAY C. L. & NG P. K. L. 2007 — Revision of the Indo-West Pacific sponge crabs of the genus *Petalomera* Stimpson, 1858 (Decapoda: Brachyura: Dromiidae). *Raffles Bulletin of Zoology* 55 (1): 107-120.
- MCLAY C. L. & TAN S. H. 2009 — Revision of the crab genus *Garthambrus* Ng, 1996, with the description of two new genera and discussion of the status of *Tutankhamen* Rathbun, 1925 (Crustacea: Brachyura: Parthenopidae). *Zootaxa* 2122: 1-50.
- MENDOZA J. C. E. 2010 — A new species of *Lipkemera* Davie, 2010 (Decapoda, Brachyura, Xanthidae, Liomerinae) from the Bohol Sea, Philippines, in FRANSEN C. H. J. M., DE GRAVE S. & NG P. K. L. (eds), Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume. *Crustaceana Monographs* 14: 505-513.
- MENDOZA J. C. E. & MANUEL-SANTOS M. R. 2012 — Revision of *Garthiella* Titgen, 1986 (Crustacea: Decapoda: Brachyura: Xanthidae), with description of a new subfamily and a new species from the central Philippines. *Zootaxa* 3446: 32-48.
- MENDOZA J. C., NARUSE T., TAN S. H., CHAN T.-Y., NG P. K. L. & RICHER DE FORGES B. 2010 — Rarity in marine invertebrates? Case studies based on decapod crustaceans collected from deep, steep slopes in the Philippines. *Biological Conservation* 19: 575-586.
- MENDOZA J. C. E. & NG P. K. L. 2008a — A new species of *Alainodaeus* Davie, 1993 (Crustacea: Decapoda: Brachyura: Xanthidae) from Balicasag Island, Philippines, with a key to the genus. *Zootaxa* 1897: 53-63.
- MENDOZA J. C. E. & NG P. K. L. 2008b — New genera and species of euxanthine crabs (Crustacea: Decapoda: Brachyura: Xanthidae) from the Bohol Sea, the Philippines. *Raffles Bulletin of Zoology* 56 (2): 385-404.
- MENDOZA J. C. E. & NG P. K. L. 2010 — Euxanthine crabs (Crustacea: Brachyura: Xanthidae) of the Philippines. *Raffles Bulletin of Zoology* 58 (1): 57-74.
- MITSUHASHI M. & CHAN T.-Y. 2006 — A new genus and species of deep-water pontoniine shrimp (Decapoda: Caridea: Palaemonidae) from Taiwan. *Zoosystema* 28 (2): 389-398.
- MITSUHASHI M. & CHAN T.-Y. 2009 — A new deep-sea pontoniine shrimp (Decapoda: Palaemonidae) of the “*Periclimenes foresti* Bruce, 1981” species group from Taiwan. *Crustaceana* 82 (7): 919-929.
- MITSUHASHI M., LI X. & CHAN T.-Y. 2010 — *Neoclimenes holthuisi* n. gen., n. sp., a new deep-sea pontoniine shrimp from the South China Sea (Decapoda: Palaemonidae), in FRANSEN C. H. J. M., DE GRAVE S. & NG P. K. L. (eds), Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume. *Crustaceana Monographs* 14: 515-527.
- MITSUHASHI M., LI X. & CHAN T.-Y. 2012 — Additional deep-sea pontoniine shrimps (Decapoda: Palaemonidae) from Taiwan, with description of one new species, in KOMATSU H., OKUNO J. & FUKUOKA K. (eds), Studies on Eumalacostraca: a homage to Takeda Masatsune. *Crustaceana Monographs* 17: 211-223.
- MOOSA M. K. 1984 — Report on the CORINDON cruises. *Marine Research in Indonesia* 24: 1-6.
- MOOSA M. K. 1986 — Stomatopod Crustacea, in FOREST J. (ed.), *Résultats des Campagnes MUSORSTOM Vol. 2. Mémoires du Muséum national d’Histoire naturelle*, 133: 367-414, pl. I.
- MOOSA M. K. 1991 — The Stomatopoda of New Caledonia and Chesterfield Islands, in RICHER DE FORGES B. (ed.) *Le benthos de fonds meubles des lagons de Nouvelle-Calédonie* 1: 149-219.
- MOOSA M. K. 1996 — Crustacea Decapoda: Deep-water swimming crabs from the South-West Pacific, particularly New Caledonia (Brachyura, Portunidae), in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM Vol. 15. Mémoires du Muséum national d’Histoire naturelle* 168: 503-530.
- NARUSE T. 2013 — Species of *Corycodus* A. Milne-Edwards, 1880 (Crustacea, Brachyura, Cyclodorippidae) collected from the Mozambique MAINBAZA and Madagascar MIRIKY expeditions, with description of a new species, in AHYONG S. T., CHANT-T., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos Vol. 27. *Mémoires du Muséum national d’Histoire naturelle* 204: 485-494.
- NARUSE T., CASTRO P. & NG P. K. L. 2009 — A new genus and new species of *Ethusidae* (Crustacea: Decapoda: Brachyura) from Vanuatu, western Pacific. *Crustaceana* 82 (7): 931-938.
- NARUSE, T. & RICHER DE FORGES B. 2010 — A new species of *Ihlopsis* Guinot & Richer de Forges, 1995 (Crustacea: Decapoda: Brachyura: Homolidae) from Taiwan. In: CASTRO P., DAVIE P. J. F., NG P. K. L. & RICHER DE FORGES B. (eds). Studies on Brachyura: Homage to Danièle Guinot. *Crustaceana Monographs* 11: 221-228.
- NG P. K. L. 1998a — *Lamoha*, a replacement name for *Hypsophrys* Wood Mason & Alcock, 1891 (Brachyura, Homolidae), a junior synonym of *Hypsophrys* Agassiz, 1859 (Pisces, Teleostei, Cichlidae). *Crustaceana* 71 (1): 121-125.
- NG P. K. L. 1998b — *Lamoha hystrix*, a new species of deep-water porter crab (Crustacea: Decapoda: Brachyura: Homolidae) from the central Pacific. *Proceedings of the Biological Society of Washington* 111 (4): 893-898.
- NG P. K. L. & CASTRO P. 2007 — On a new genus and species of euryplacid crab (Crustacea: Decapoda: Brachyura: Goneplacoidea) from the Philippines. *Zootaxa* 1549: 43-53.

- NG P. K. L. & CHEN H. L. 1999 — On the identities of two Pacific species of deep-water porter crabs, *Hypsophris longirostris* Chen, 1986, and *Homologenus donghaiensis* Chen, 1986 (Crustacea: Decapoda: Brachyura: Homolidae). *Proceedings of the Biological Society of Washington* 112 (4): 759-767.
- NG P. K. L. & CHIA D. G. B. 1994 — The genus *Glyptocarcinus* Takeda, 1973, with descriptions of a new subfamily, two new genera and two new species from New Caledonia (Crustacea: Decapoda: Brachyura: Xanthidae). *Raffles Bulletin of Zoology* 42 (3): 701-730.
- NG P. K. L. & ELDREDGE L. G. 2012 — A new species of *Yaldwynopsis* from O'ahu, Hawai'i (Crustacea: Decapoda: Brachyura: Homolidae), in EVENHUIS N. L. & ELDREDGE L. G. (eds), Records of the Hawaii Biological Survey for 2011. *Bishop Museum Occasional Papers*, Honolulu 112: 29-38.
- NG P. K. L., GUINOT D. & DAVIE P. J. F. 2008 — Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *Raffles Bulletin of Zoology* suppl 17: 1-286.
- NG P. K. L. & HO P.-H. 2003 — On the deep-water dorippid crabs of the genus *Ethusina* Smith, 1884 (Crustacea: Decapoda: Brachyura) from Taiwan. *Raffles Bulletin of Zoology* 51 (1): 71-85.
- NG P. K. L. & MANUEL-SANTOS M. R. 2007 — Establishment of the Vultocinidae, a new family for an unusual new genus and new species of Indo-West Pacific crab (Crustacea: Decapoda: Brachyura: Gonoplacoidea), with comments on the taxonomy of the Gonoplacidae. *Zootaxa* 1558: 39-68.
- NG P. K. L. & MCLAY C. L. 2005 — *Dicranodromia danielae*, a new species of homolodromiid crab from the Philippines (Crustacea: Decapoda: Brachyura). *Zootaxa* 1029: 39-46.
- NG P. K. L. & MCLAY C. L. 2010 — *Metadynomene tuamotu*, a new species of dynomenid crab from French Polynesia (Crustacea: Decapoda: Brachyura). *Zootaxa* 2405: 48-54.
- NG P. K. L. & NARUSE T. 2007a — On two species of deep sea homoloid crabs of the genus *Dicranodromia* (Crustacea: Decapoda: Brachyura: Homolodromiidae) from the Philippines, with description of a new species. *Raffles Bulletin of Zoology* suppl. 16: 47-53.
- NG P. K. L. & NARUSE T. 2007b — *Lia Gore pulchella*, a new species of xanthid crab (Crustacea: Decapoda: Brachyura) from Vanuatu. *Zootaxa* 1665: 53-60.
- NG P. K. L. & RICHER DE FORGES B. 2007 — A new genus and new species of leucosiid crab from New Caledonia, with a note on the validity of *Tanaoa serenei* (Richer de Forges, 1983) (Crustacea: Decapoda: Brachyura). *Zootaxa* 1662: 15-24.
- NG P. K. L. & RICHER DE FORGES B. 2009 — *Vultocinus anfractus* Ng & Manuel-Santos, 2007 (Decapoda, Brachyura, Vultocinidae): a new record for New Caledonia, with notes on female characters. *Crustaceana* 82 (5): 627-634.
- NG P. K. L. & RICHER DE FORGES B. 2012 — *Pleisticanthoides* Yokoya, 1933, a valid genus of inachid spider crab (Crustacea: Decapoda: Brachyura: Majooidea), with descriptions of two new species from the Philippines, Papua-New Guinea and Vanuatu. *Zootaxa* 3351: 65-81.
- NG P. K. L. & WANG C. H. 2002 — Notes on two species of deep-water porter crabs of the genus *Lamoha* (Crustacea: Decapoda: Brachyura: Homolidae) from the Pacific. *Journal of the National Taiwan Museum* (2001) 54 (2): 13-24.
- NGUYEN T. S. & NG P. K. L. 2010 — A new genus of the family Portunidae (Crustacea: Decapoda: Brachyura) and the identity of *Portunus (Cycloachelous) yoronensis* Sakai, 1974. *Zootaxa* 2677: 38-48.
- NIELSEN C. & PARKER A. 2010 — Morphological novelties detonated the Ediacaran-Cambrian "explosion". *Evolution & Development* 12: 345-346.
- NOËL P. 1985 — Crustacés Décapodes: Processidae de l'Indo-Ouest-Pacifique, in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 2. Mémoires du Muséum national d'Histoire naturelle 133: 261-301.
- OSAWA M., LIN C.-W. & CHANT T.-Y. 2006a — *Munidopsis* (Decapoda: Anomura: Galatheidae) from the abyssal depths of Taiwan, with description of one new species. *Journal of Crustacean Biology* 26: 420-428.
- OSAWA M., LIN C.-W. & CHAN T.-Y. 2006b — A new species of *Munidopsis* (Crustacea: Decapoda: Galatheidae) from deep waters off Taiwan. *Proceedings of the Biological Society of Washington* 119: 251-258.
- OSAWA M., LIN C.-W. & CHAN T.-Y. 2007 — A new deep-sea squat lobster of the genus *Munidopsis* Whiteaves, 1874 (Crustacea: Decapoda: Anomura: Galatheidae) collected by the Panglao 2005 expedition to the Philippines. *Raffles Bulletin of Zoology* suppl. 16: 15-20.
- OSAWA M., LIN C.-W. & CHANT T.-Y. 2008 — Species of *Galacantha* and *Munidopsis* (Crustacea: Decapoda: Anomura: Galatheidae) from the deep-waters off Taiwan, with the description of two new species. *Scientia Marina* 72: 35-57.
- OSAWA M., C.-W. LIN & CHAN T.-Y. 2013 — Munidopsidae Ortmann, 1898 (Crustacea, Decapoda, Anomura) collected by the PANGLAO 2005 and AURORA expeditions to the Philippines, with descriptions of four new species from the Philippines and one new species from Taiwan, in AHYONG S.T., CHANT-T.Y., CORBARI L. & NG P.K. L. (eds), Tropical Deep-Sea Benthos Vol. 27. Mémoires du Muséum national d'Histoire naturelle 204: 231-286.
- PAILLERET M., HAGAT, PETIT P., PRIVÉ-GILL C., SAEDLOU N., GAILL F. & ZBINDEN M. 2007 — Sunken wood from the Vanuatu Islands: identification of wood substrates and preliminary description of associated fauna. *Marine Ecology* 28: 233-241.
- PALACIOS C., ZBINDEN M., PAILLERET M., GAILL F. & LEBARON P. 2009 — High similarity in the microbial community structure of sunken woods at shallow marine waters and deep-sea sites across the oceans. *Microbial Ecology* 58 (4): 737-752.
- PARKER A. R. 1998 — Colour in Burgess Shale animals and the effect of light on evolution in the Cambrian. *Proceedings of the Royal Society of London* 265: 967-972.
- PARKER A. R. 2003 — *In the blink of an eye: the cause of the most dramatic event in the history of life*. The Free Press, Simon & Scuster Inc., London: 316 p.
- PIRLOT J. M. 1934 — Les amphipodes de l'Expédition du Siboga. Deuxième partie. Les amphipodes gammarides. II. Les amphipodes de la Mer Profonde. 2. *Siboga Expeditions* 33: 167-235.
- POORE G. C. B., AHYONG S. T. & TAYLOR J. (eds) 2011 — *The biology of squat lobsters*. CSIRO Publishing: Melbourne and CRC Press: Boca Raton, 363 pp.
- POUPIN J. 1994 — The genus *Justitia* Holthuis, 1946, with description of *J. chani* and *J. vericeli* spp. nov. (Crustacea: Decapoda: Palinuridae). *Journal of Taiwan Museum* 47 (1): 37-56.
- POUPIN J. 1995 — Étude des *Naxioides* du groupe *robilliardi* (Miers, 1882) (Brachyura: Majidae: Pisinae). *Journal of Natural History* 29 (1): 85-109.
- POUPIN J. 1996a — *Atlas des Crustacés marins profonds de Polynésie Française. Récoltes du navire MARARA (1986/1996)*. Service mixte de Surveillance radiologique et biologique (SMSRB), Montlhéry, 59 pp.
- POUPIN J. 1996b — Crustacea Decapoda of French Polynesia (Astacidea, Palinuridea, Anomura, Brachyura). *Atoll Research Bulletin* 442: i-iv, 1-114.
- POUPIN J. 2001 — New collections of *Ciliopagurus* from French Polynesia, with the description of a new species from the Marquesas Islands (Crustacea: Decapoda: Anomura: Diogenidae). *Raffles Bulletin of Zoology* 49 (2): 291-300.

- POUPIN J. & LEMAIRE R. 2003 — Hermit crabs of the genus *Calcinus* Dana, 1851 (Decapoda: Anomura: Diogenidae) from the Austral Islands, French Polynesia, with description of a new species. *Zootaxa* 391: 1-20.
- POUPIN J. & MCLAUGHLIN P. 1998 — Additional *Calcinus* (Decapoda, Anomura, Diogenidae) from French Polynesia with three new species and a key to Indo-West Pacific species. *Crustacean Research* 27: 9-27.
- POUPIN J., TAMARII T. & VANDENBOOMGAERDE A. 1990 — (N/O MARARA - 1986/1989). *Notes et Documents ORSTOM* Papeete, sér. *Océanographie* (42): 1-97.
- PUILLANDRE N., MACPHERSON E., LAMBOURDIERE J., CRUAUD C., BOISSELIER-DUBAYLE M.C. & SAMADI S. 2011 — Barcoding type-specimens helps to reliably link species names to molecular taxonomic units in *Eumunida* Smith, 1883 (Decapoda, Eumunididae). *Invertebrate Systematics* 25: 322-333.
- RAHAYU D. L. 2005 — Additions to the Indonesian fauna of the hermit crab genus *Pseudopaguristes* McLaughlin and a further division of the genus *Paguristes* Dana (Crustacea: Decapoda: Paguroidea: Diogenidae). *Zootaxa* 831: 1-42.
- RAHAYU D. L. 2006 — The genus *Paguristes* (Crustacea, Decapoda, Diogenidae) from Indonesia, in RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 349-374.
- RAHAYU D. L. 2007 — The hermit crabs *Paguristes* Dana, 1851 sensu lato (Crustacea, Decapoda, Anomura, Diogenidae) from the western Indian Ocean. *Zoosystema* 29 (3): 515-534.
- RAHAYU D. L. & KOMAI T. 2013 — Two new species of *Pseudopagurodes* McLaughlin, 1997 (Crustacea, Decapoda, Anomura, Paguridae) from the Philippines, in AHYONG S. T., CHAN T.-Y., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos, Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 423-436.
- RICHER DE FORGES B. 1990 — Les campagnes d'exploration de la faune bathyale dans la zone économique de la Nouvelle-Calédonie. Explorations for bathyal fauna in the New Caledonian economic zone, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 9-54.
- RICHER DE FORGES B. 1994 — A new genus of deep-sea majid crab: *Griffinia* gen. nov. (Crustacea, Decapoda, Brachyura). *The Beagle, Records of the Museum and Art Gallery of the Northern Territory* 11: 65-72.
- RICHER DE FORGES B. 2006 — Découverte en mer du Corail d'une deuxième espèce de glypéide (Crustacea, Decapoda, Glypheoidea). *Zoosystema* 28 (1): 17-29.
- RICHER DE FORGES B. 2010 — Majoid crabs from the Mozambique Channel with the description of a new species of *Oxyleurodon* Miers, 1886 (Decapoda, Brachyura), in FRANSEN C. H. J. M., DE GRAVE S. & NG P. K. L. (eds), Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume. *Crustaceana Monographs* 14: 645-653.
- RICHER DE FORGES B. & CORBARI L. 2012 — A new species of *Oxyleurodon* Miers, 1886 (Crustacea, Brachyura, Majoidea) from the Bismarck Sea, Papua New Guinea. *Zootaxa* 3320: 56-60.
- RICHER DE FORGES B., HOFFSCHIR C., CHAUVIN C. & BERTHAULT C. 2005 — Inventaire des espèces de profondeur de Nouvelle-Calédonie/Census of deep-sea species of New Caledonia. *Documents scientifiques et techniques* /l6 volume spécial, IRD/Nouméa; 113 pp.
- RICHER DE FORGES B. & JUSTINE J.-L. 2006 — Introduction, in: RICHER DE FORGES B. & JUSTINE J. L. (eds), Tropical Deep-Sea Benthos, Vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 15-32.
- RICHER DE FORGES B., KOSLOW J. A. & POORE G. C. B. 2000 — Diversity and endemism of the benthic seamount fauna in the south-west Pacific. *Nature* 405: 944-947.
- RICHER DE FORGES B., NEWELL P., SCHLACHER-HOENLINGER M., SCHLACHER T., NATING D., CESAR F. & BOUCHET P. 2000 — La campagne MUSORSTOM 10 dans l'archipel des îles Fidji. Compte rendu et liste des stations, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 9-23.
- RICHER DE FORGES B. & NG P. K. L. 2006 — The Brachyura of New Caledonia, in: *Compendium of marine species from New Caledonia*. Documents Scientifiques et Techniques, Institut de recherches pour le développement, PAYRI C. E. & RICHER DE FORGES B. (eds), special volume 117: 273-289.
- RICHER DE FORGES B. & NG P. K. L. 2007a — New records and new species of Homolidae De Haan, 1839, from the Philippines and French Polynesia (Crustacea: Decapoda: Brachyura). *Raffles Bulletin of Zoology suppl.* 16: 29-45.
- RICHER DE FORGES B. & NG P. K. L. 2007b — Notes on deep-sea spider crabs of the genus *Cyrtomaia* Miers, 1886, from the Philippines (Crustacea: Decapoda: Brachyura: Majidae), with description of a new species. *Raffles Bulletin of Zoology suppl.* 16: 55-65.
- RICHER DE FORGES B. & NG P. K. L. 2008a — New records of deep-sea spider crabs of the genus *Cyrtomaia* Miers, 1886, from the Pacific Ocean, with description of a new species (Crustacea: Decapoda: Brachyura: Majidae). *Zootaxa* 1861: 17-28.
- RICHER DE FORGES B. & NG P. K. L. 2008b — New Western Pacific records of Homolidae De Haan, 1839, with descriptions of new species of *Homolochunia* Doflein, 1904, and *Latreillopsis* Henderson, 1888 (Crustacea: Decapoda: Brachyura). *Zootaxa* 1967: 1-35.
- RICHER DE FORGES B. & NG P. K. L. 2009a — On the majoid genera *Oxyleurodon* Miers, 1886, and *Sphenocarcinus* A. Milne-Edwards, 1875 (Crustacea: Brachyura: Epialtidae), with descriptions of two new genera and five new species. *Raffles Bulletin of Zoology suppl.* 20: 247-266.
- RICHER DE FORGES B. & NG P. K. L. 2009b — New genera, new species and new records of Indo-West Pacific spider crabs (Crustacea: Brachyura: Epialtidae: Majoidea). *Zootaxa* 2025: 1-20.
- RICHER DE FORGES B. & NG P. K. L. 2012 — *Griffinia takedai*, a new species of deep sea majoid crab (Decapoda, Brachyura, Epialtidae) from the Philippines, in KOMATSU H., OKUNO J. & FUKUOKA K. (eds), Studies on Eumalacostraca: a homage to Takeda Masatsune. *Crustaceana Monographs* 17: 275-284.
- RICHER DE FORGES B. & NG P. K. L. 2013 — On a collection of spider crabs of the genera *Rochinia* A. Milne-Edwards, 1875 and *Naxioides* A. Milne-Edwards, 1875 (Crustacea, Brachyura, Majoidea: Epialtidae) from Mozambique Channel, Solomon, Vanuatu and Philippines Islands, with description of a new species of *Rochinia*, in AHYONG S. T., CHAN T.-Y., CORBARI L. & NG P. K. L. (eds), Tropical Deep-Sea Benthos, Vol. 27. *Mémoires du Muséum national d'Histoire naturelle* 204: 467-484.
- RICHER DE FORGES B. & NG P. K. L. & AHYONG S. 2013 — *Parapleisticantha* Yokoya, 1933, a valid genus of deep-sea inachid spider crab from Japan and Philippines (Crustacea: Decapoda: Brachyura: Majoidea). *Zootaxa* 3635: 15-26.
- RICHER DE FORGES B., TAN S. H., BOUCHET P., NG P. K. L. & SAGUIL N. 2009 — PANGLAO 2005 — Survey of the deep-water benthic fauna of the Bohol Sea and adjacent waters. *Raffles Bulletin of Zoology suppl.* 20: 19-36.
- RICHER DE FORGES B. & TESSIER E. 2012 — Les monts sous-marins: éléments morphologiques essentiels des planchers océaniques. *Le Courrier de la Nature* mai-juin 2012 (288): 1-8.

- ROWDEN A. A., SCHNABEL K. E., SCHLACHERT. A., MACPHERSON E., AHYONG S. T. & RICHER DE FORGES B. 2010 — Squat lobster assemblages on seamounts differ from some, but not all, deep-sea habitats of comparable depth. *Marine Ecology* 31 (supl. 1): 63-83.
- SAINT LAURENT DE M. 1988 — Enoplotetrapoda, nouvelle superfamille de Crustacés décapodes Astacidea. *Comptes Rendus hebdomadaires de l'Académie des Sciences*, Paris (3) 307: 59-62.
- SAINT LAURENT DE M. 1989 — La nouvelle superfamille des Retroplomoidea Gill, 1894 (Decapoda, Brachyura): systématique, affinités et évolution, in FOREST, J. (ed.), Résultats des campagnes MUSORSTOM Vol. 5. *Mémoires du Muséum national d'Histoire naturelle* 144: 103-179.
- SAINT LAURENT DE M. & CLEVA R. 1981 — Crustacés Décapodes: Stenopodidea, in DORST J. (ed.), Résultats des Campagnes MUSORSTOM Vol. 1. *Mémoires ORSTOM* 91: 151-188.
- SAINT LAURENT DE M. & MACPHERSON E. 1990 — Crustacea Decapoda: Le genre *Eumunida* Smith, 1883 (Chirostyliidae) dans les eaux néo-calédoniennes, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 227-288.
- SAINT LAURENT DE M. & MACPHERSON E. 1997 — Une nouvelle espèce du genre *Paralomis* White, 1856, des sources hydrothermales du Sud-ouest Pacifique (Crustacea, Decapoda, Lithodidae). *Zoosystema* 19: 721-727.
- SAINT LAURENT DE M. & POUPIN J. 1996 — Crustacea, Anomura: Les espèces indo-ouest pacifiques du genre *Eumunida* Smith, 1880 (Chirostyliidae) Description de six espèces nouvelles, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 337-385.
- SAITOH T. & KOMAI T. 2008 — A review of species of the genera *Spongicola* de Haan, 1844 and *Paraspongicola* de Saint Laurent & Cleva, 1981 (Crustacea, Decapoda, Stenopodidea, Spongicolidae). *Zoosystema* 30: 87-147.
- SAITOH T., OKUNO J. & CHANT.-Y. 2009 — A new species of *Stenopus* (Crustacea: Decapoda: Stenopodiidae) from the Indo-West Pacific, with a redefinition of the genus. *Raffles Bulletin of Zoology* suppl. 20: 109-120.
- SAMADI S., BOTTAN L., MACPHERSON E., RICHER DE FORGES B., BOISSELIER M.-C. 2006 — Seamount endemism questioned by the geographic distribution and population genetic structure of marine invertebrates. *Marine Biology* 149: 1463-1475.
- SAMADI S., CORBARI L., LORION J., HOURDEZ S., HAGAT, DUPONT J., BOISSELIER M.-C. & RICHER DE FORGES B. 2010 — Biodiversity of deep-sea organisms associated with sunken-wood or other organic remains sampled in the tropical Indo-Pacific. *Cahier de Biologie Marine* 51: 459-466.
- SAMADI S., QUEMERE E., LORION J., TILLIER A., VON COSEL R., LOPEZ P., CRUAUD C., COULOUX A. & BOISSELIER M.-C. 2007 — Molecular phylogeny in mytilids supports the wooden steps to deep-sea vents hypothesis. *Comptus Rendus Biologie* 330: 1995.
- SAMADI S., SCHLACHERT. & RICHER DE FORGES B. 2007 — Seamount Benthos, in: PITCHER T. J., MORATO T., HART P. J. B., CLARK M. R., HAGGAN N. & SANTOS R. S. (eds). *Seamounts. Ecology, Fisheries, Conservation*. Blackwell, Fisheries and Aquatic Resources Series: 119-140.
- SCHNABEL K.E., AHYONG S.T. & MAAS, E.W. 2011 — Galatheoidea are not monophyletic – molecular and morphological phylogeny of the squat lobsters (Decapoda: Anomura) with recognition of a new superfamily. *Molecular Phylogenetics and Evolution* 58: 157-168.
- SERENE R. & VADON C. 1981 — Crustacés Décapodes: Brachyoures. Liste préliminaire, descriptions de formes nouvelles et remarques taxonomiques. Résultats des Campagnes MUSORSTOM Vol. 1. *Collection Mémoires ORSTOM* 91: 117-140, pls 1-4.
- SCHNABEL K. E. & AHYONG S.T. 2010 — A new classification of the Chirostyloidea (Crustacea: Decapoda: Anomura). *Zootaxa* 2687: 56-64.
- SCHRAM F. R. & AHYONG S. T. 2002 — The higher affinities of *Neoglypeha inopinata* Forest & de Saint Laurent in particular and the Glypoidea (Reptantia: Decapoda) in général. *Crustaceana* 75: 629-635.
- SIN Y. W., LAI J. C. Y., NG P. K. L. & CHU K. H. 2009 — Phylogeny of Dorippoidea (Crustacea: Decapoda: Brachyura) inferred from three mitochondrial genes. *Invertebrate Systematics* 23: 223-230.
- SNELGROVE P. V. R. 2010 — Discoveries of the Census of Marine Life. Making Ocean Life Count. Cambridge University Press, London: 1-286.
- TAN C. G. S. & NG P. K. L. 1993 — On two new species of *Oreotlos* Ihle, 1918 (Crustacea Decapoda, Brachyura, Leucosiidae) from French Polynesia. *Bulletin du Muséum nationale d'Histoire naturelle* (4) 14A (3-4): 797-804.
- TAN C. G. S. & NG P. K. L. 1995 — A revision of the Indo-Pacific genus *Oreophorus* Rüppell, 1830 (Crustacea: Decapoda: Brachyura: Leucosiidae), in RICHER DE FORGES B. (ed.), *Les fonds meubles des lagons de Nouvelle-Calédonie* (Sédimentologie, Benthos) Vol. 2. *Études & Thèses ORSTOM*, Paris: 101-205, pls 1-16.
- TAN S. H. & NG P. K. L. 2007a — Descriptions of new genera from the subfamily Parthenopinae (Crustacea: Decapoda: Brachyura: Parthenopidae). *Raffles Bulletin of Zoology* suppl. 16: 95-119.
- TAN S. H. & NG P. K. L. 2007b — Review of the Daldorfiinae Ng & Rodríguez, 1986 (Crustacea: Decapoda: Brachyura: Parthenopidae). *Raffles Bulletin of Zoology* suppl. 16: 121-167.
- TAVARES M. S. 1991a — Redéfinition des genres *Rochinia* A. Milne Edwards, *Sphenocarcinus* A. Milne Edwards et *Oxypleurodon* Miers, et établissement du genre *Nasutocarcinus* gen. nov. (Crustacea, Brachyura, Majidae). *Bulletin du Muséum national d'Histoire naturelle* A 13(1-2): 159-179.
- TAVARES M. 1991b — Révision préliminaire du genre *Tymolus* Stimpson, avec la description de *Tymolus brucei* sp. nov. d'Australie occidentale (Crustacea, Brachyura, Cyclodorippioidea). *Bulletin du Muséum national d'Histoire naturelle*, (4) A (13): 439-456.
- TAVARES M. S. 1992 — Tendances évolutives chez les Crabes primitifs, avec la description d'un nouveau type de chambre incubatrice (Crustacea, Decapoda: Cyclodorippinae Ortmann, 1892, et Xeinostomatinae subfam. nov.). *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* (3) 312: 509-514.
- TAVARES M. 1993 — Crustacea Decapoda: Les Cyclodorippidae et Cymonomidae de l'Indo-Ouest-Pacifique à l'exclusion du genre *Cymonomus*, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM Vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 253-313.
- TAVARES M. 1998 — Phyllotymolinidae, nouvelle famille de Brachyoures Podotremata (Crustacea, Decapoda). *Zoosystema* 20 (1): 109-122.
- TAVARES M. & CLEVA R. 2010 — Trichopeltariidae (Crustacea, Decapoda, Brachyura), a new family and superfamily of eubrachyuran crabs with description of one new genus and five new species. *Papéis Avulsos de Zoologia*, Museu de Zoologia da Universidade de São Paulo 50 (9): 97-157.
- TAYLOR H. R. & HARRIS W. E. 2012 — An emergent science on the brink of irrelevance: a review of the past 8 years of DNA barcoding. *Molecular Ecology Resources* 12: 377-388.
- THUBAUT J. 2012 — Histoire évolutive et biologie des populations des mytilidés associés aux substrats organiques coulés. Approche comparative au sein des Bathymodiolinae des milieux réducteurs profonds. *Thèse du Muséum national d'Histoire naturelle*, Paris, pp. 226.

- THOMSON C. W. 1878 — *The Atlantic. A preliminary account of the general results of The Exploring voyage of H.M.S. "Challenger" during the year 1873 and the early part of the year 1876.* In two volumes. New York, Harper & Brothers, Publishers, Franklin Square: 391 pp.
- TONG J. G., CHANT-T.-Y. & CHU K. H. 2000 — A preliminary phylogenetic analysis of *Metapenaeopsis* (Decapoda: Penaeidae) based on mitochondrial DNA sequences of selected species from the Indo-West Pacific. *Journal of Crustacean Biology* 20 (3): 541-549.
- TREW CRIST, D., SCOCROFT G. & HARDING J. M. JR. 2009 — *World Ocean Census. A global survey of Marine Life.* CoML, Firefly Books; 256 pp.
- TSANG L. M., LIN F.-J., CHU K. H. & CHANT-T.-Y. 2008 — Phylogeny of Thalassinidea (Crustacea, Decapoda) inferred from three rDNA sequences: implications for morphological evolution and superfamily classification. *Journal of Zoological Systematics and Evolutionary Research* 46 (3): 216-223.
- TSANG, L. M., MA K. Y., AHYONG S. T., CHANT-T.-Y. & CHU K. H. 2008 — Phylogeny of Decapoda using two nuclear protein-coding genes: Origin and evolution of the Reptantia. *Molecular Phylogenetics and Evolution* 28: 359-368.
- TSANG L. M., CHANT-T.-Y., CHEUNG M. K. & CHU K. H. 2009 — Molecular evidence for the Southern Hemisphere origin and deep sea diversification of spiny lobsters (Crustacea: Decapoda: Palinuridae). *Molecular Phylogenetics and Evolution* 51: 304-311.
- TSANG L. M., CHAN T.-Y., AHYONG S. T. & CHU K. H. 2011 — Hermit to kind, or hermit to all: Multiple transitions to crab-like forms from hermit crab ancestors. *Systematic Biology* 60 (5): 616-629.
- TSHUDY D., ROBLES R., CHANT-T.-Y., HO K. C., CHU K. H., AHYONG S. T. & FELDER D. L. 2009 — Phylogeny of marine clawed lobster families Nephropidae Dana 1852 and Thaumastochelidae Bate 1888 based on mitochondrial genes, in MARTIN J. W., CRANDALL K. A. & FELDER D. L. (eds), *Decapod crustacean phylogenetics. Crustacean Issues* 18: 357-367.
- TSOI K. H., CHAN T.-Y. & CHU K. H. 2011 — Phylogenetic and biogeographic analysis of the spear lobsters *Linuparus* (Decapoda, Palinuridae), with the description of a new species. *Zoologischer Anzeiger* 250: 302-315.
- TUDGE C. C., SCHELTINGA D. M., JAMIESON B. G. M., GUINOT D. & RICHER DE FORGES B. 2012 — Comparative ultrastructure of the spermatozoa of the Majoidea (Crustacea, Decapoda, Brachyura) with new data on six species in five genera. *Acta Zoologica* (Stockholm), doi: 10.1111/azo.12005: 1-20.
- WILLIAMS J. D. & BOYKO C. B. 2006 — A new species of *Tomlinsonia Turquier, 1885* (Crustacea, Cirripedia, Trypetesidae) in hermit crab shells from the Philippines, and a new parasite species of *Hemioniscus Buchholz, 1866* (Crustacea, Isopoda, Hemioniscidae). *Zoosystema* 28 (2): 285-305.
- YANG C.-H., BRACKEN-GRISOM H., KIM D., CRANDALL K. A. & CHAN T.-Y. 2012 — Phylogenetic relationships, character evolution, and taxonomic implications within the slipper lobsters (Crustacea: Decapoda: Scyllaridae). *Molecular Phylogenetics and Evolution* 62: 237-250.
- YANG C.-H. & CHANT-T.-Y. 2010 — A new slipper lobster of the genus *Galearctus* Holthuis, 2002 (Decapoda: Scyllaridae) from Taiwan and Japan, in FRANSEN C. H. J. M., DE GRAVE S. & NG P. K. L. (eds), *Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume. Crustaceana Monographs* 14: 735-745.
- YANG C.-H. & CHAN T.-Y. 2012 — On the taxonomy of the slipper lobster *Chelarctus cultrifer* (Ortmann, 1897) (Crustacea: Decapoda: Scyllaridae), with description of a new species. *Raffles Bulletin of Zoology* 60 (2): 449-460.
- YANG C.-H., CHANT-T.-Y. & CHU K. H. 2010 — Two new species of the "*Heterocarpus gibbosus* Bate, 1888" species group (Crustacea: Decapoda: Pandalidae) from the western Pacific and north-western Australia. *Zootaxa* 2372: 206-220.
- YANG C.-H., CHEN I.-S. & CHANT-T.-Y. 2008 — A new slipper lobster of the genus *Petrarctus* (Crustacea: Decapoda: Scyllaridae) from the West Pacific. *Raffles Bulletin of Zoology* suppl. 19: 71-81.
- YANG C.-H., CHEN I.-S. & CHANT-T.-Y. 2011 — A new slipper lobster of the genus *Galearctus* Holthuis, 2002 (Crustacea: Decapoda: Scyllaridae) from the New Caledonia. *Zoosystema* 32 (2): 207-217.
- ZBINDEN M., PAILLERET M., RAVAUX J., GAUDRON S. M., HOYOUX C., LAMBOURDIERE J., WAREN A., LORION J., HALARY S. & DUPERRON S. 2010 — Bacterial communities associated with the wood feeding gastropod *Pectinodonta* sp. (Patellogastropoda, Mollusca). *FEMS Microbiology and Ecology* 74: 450-463.

## APPENDIX 1

Cruises in the *Tropical Deep-Sea Benthos* programme and satellite programmes that have yielded deep-sea zoological material, and relevant published cruise narratives or station lists (updated list since Bouchet *et al.* (2008).

CRUISE NAME OR ACRONYM	YEAR	SHIP	AREA	NO. OF STATIONS	STATION NUMBERS (CONSECUTIVE NUMBERING)	CRUISE REPORT IF ANY
ATIMO VATAE	2010	<i>Nosy b�� 11</i>	South Madagascar	119	3508-3627	
AURORA 2007	2007	DA-BFAR	Philippines	117	2653-2764	
AZTEQUE	1990	<i>Alis</i>	New Caledonia	11		Grandperrin <i>et al.</i> 1990
BATHUS 1	1993	<i>Alis</i>	New Caledonia	75	639-713	Richer de Forges & Chevillon 1996
BATHUS 2	1993	<i>Alis</i>	New Caledonia	58	714-771	Richer de Forges & Chevillon 1996
BATHUS 3	1993	<i>Alis</i>	New Caledonia	78	773-849	Richer de Forges & Chevillon 1996
BATHUS 4	1994	<i>Alis</i>	New Caledonia	74	882-955	Richer de Forges & Chevillon 1996
BENTHAUS	2002	<i>Alis</i>	Austral Islands	161	1861-2021	
BERYX 11	1992	<i>Alis</i>	New Caledonia	60		Lehodey <i>et al.</i> 1992
BERYX 2	1991	<i>Alis</i>	New Caledonia	19		Grandperrin & Lehodey 1992
BIOCAL	1985	Jean-Charcot	New Caledonia	68		L��vi, 2006
BIOGEOCAL	1987	<i>Coriolis</i>	New Caledonia	152		Cotillon P. & Monniot C. 1987
BIOPAPUA	2010	<i>Alis</i>	Papua New-Guinea	156	3628-3783	
BOAO	2004	<i>Alis</i>	Vanuatu	27	2304-2330	
BOA1	2005	<i>Alis</i>	Vanuatu	71	2411-2480	
BORDAU 1	1999	<i>Alis</i>	Fiji	117	1391-1507	Richer de Forges 2000
BORDAU 2	2000	<i>Alis</i>	Tonga	138	1508-1645	
CALSUB	1989	<i>Cyana</i>	New Caledonia	22		Grandperrin & Richer de Forges 1989; Roux 1994
CHALCAL 1	1984	<i>Coriolis</i>	Coral Sea	27		Richer de Forges & Pianet 1984
CHALCAL 2	1986	<i>Coriolis</i>	New Caledonia	38		Richer de Forges <i>et al.</i> 1987
CONCALIS	2008	<i>Alis</i>	New Caledonia	102	2929-3030	
CORAIL 2	1988	<i>Coriolis</i>	Coral Sea	19		Richer de Forges <i>et al.</i> 1988
CORINDON 2	1980	<i>Coriolis</i>	Indonesia	70		Moosa 1985
EBISCO	2005	<i>Alis</i>	Coral Sea	187	1-14 and 2492-2652	
EXBODI	2011	<i>Alis</i>	New Caledonia	162	3784-3945	
GEMINI	1989	<i>Alis</i>	New Hebrides Arc	15		Bargibant <i>et al.</i> 1989
HALICAL 1	1994	<i>Alis</i>	New Caledonia	4		Grandperrin <i>et al.</i> 1995a
HALIPRO 1	1994	<i>Alis</i>	New Caledonia	32	850-881	Grandperrin <i>et al.</i> , 1995b
HALIPRO 2	1996	<i>Tangaroa</i>	New Caledonia	106		Grandperrin <i>et al.</i> 1997
KARUBAR	1991	<i>Baruna Jaya 1</i>	Indonesia	91		Crosnier <i>et al.</i> 1997
LAGON	1984-89	Vauban, then <i>Alis</i>	New Caledonia	62		Richer de Forges 1991
LITHIST	1999	<i>Alis</i>	New Caledonia	18		
MADANG	2012	<i>Alis</i>	Papua New-Guinea	137	3946-4082	
MAINBAZA	2009	<i>Vizconde de Eza</i>	Mozambique Channel	46	3130-3175	
MIRIKY	2009	<i>Miriky</i>	Madagascar	119	3176-3294	
MUSORSTOM 1	1976	<i>Vauban</i>	Philippines	75	1-73	Forest 1981
MUSORSTOM 2	1980	<i>Coriolis</i>	Philippines	82	1-83	Forest 1986
MUSORSTOM 3	1985	<i>Coriolis</i>	Philippines	59	86-145	Forest 1989
MUSORSTOM 4	1985	<i>Vauban</i>	New Caledonia	148	146-249	Richer de Forges 1986
MUSORSTOM 5	1986	<i>Coriolis</i>	Coral Sea	137	250-390	Richer de Forges <i>et al.</i> 1986
MUSORSTOM 6	1989	<i>Alis</i>	New Caledonia	103	391-493	Richer de Forges & Laboute 1989
MUSORSTOM 7	1992	<i>Alis</i>	Wallis & Futuna	147	494-638	Richer de Forges & Menou 1993
MUSORSTOM 8	1994	<i>Alis</i>	Vanuatu	186	956-1141	Richer de Forges <i>et al.</i> 1996
MUSORSTOM 9	1997	<i>Alis</i>	Marquesas	166	1142-1307	Richer de Forges <i>et al.</i> 1999
MUSORSTOM 10	1998	<i>Alis</i>	Fiji	83	1308-1390	Richer de Forges <i>et al.</i> 2000
NORFOLK 1	2001	<i>Alis</i>	New Caledonia	88	1651-1739	
NORFOLK 2	2003	<i>Alis</i>	New Caledonia	142	2022-2161	

CRUISE NAME OR ACRONYM	YEAR	SHIP	AREA	NO. OF STATIONS	STATION NUMBERS (CONSECUTIVE NUMBERING)	CRUISE REPORT IF ANY
PALEO-SURPRISE	1999	<i>Alis</i>	New Caledonia	29		Garrigue <i>et al.</i> 2000
PANGLAO 2005	2005	<i>DA-BFAR</i>	Philippines	80	2331-2409	Richer de Forges <i>et al.</i> 2009
SALOMON 1	2001	<i>Alis</i>	Solomon Islands	121	1740-1860	
SALOMON 2	2004	<i>Alis</i>	Solomon Islands	142	2164-2303	
SALOMONBOA 3	2007	<i>Alis</i>	Solomon Islands	97	2765-2860	
SANTO 2006	2006	<i>Alis</i>	Vanuatu	112		
SMCB	1990-91	<i>Marara</i>	French Polynesia	80		Poupin 1991
SMIB 1	1986	<i>Vauban</i>	New Caledonia	15		Richer de Forges 1990
SMIB 2	1986	<i>Vauban</i>	New Caledonia	31		Richer de Forges 1990
SMIB 3	1987	<i>Vauban</i>	New Caledonia	32		Richer de Forges 1990
SMIB 4	1989	<i>Alis</i>	New Caledonia	36		Richer de Forges 1990
SMIB 5	1989	<i>Alis</i>	New Caledonia	36		Richer de Forges 1993
SMIB 6	1990	<i>Alis</i>	New Caledonia	32		Richer de Forges 1993
SMIB 8	1993	<i>Alis</i>	New Caledonia	55		Richer de Forges & Chevillon 1996
SMIB 9	1993	<i>Alis</i>	New Caledonia	1		
SMIB 10	1995	<i>Alis</i>	New Caledonia	17		Richer de Forges 1990
TAIWAN 2000	2000	<i>Fisheries Researcher 1</i>	Taiwan	62		
TAIWAN 2001	2001	<i>Chung Tung Long 26</i>	Taiwan	80		
TAIWAN 2002	2002	<i>Cheng-Ming Fa and Ocean Researcher 1</i>	Taiwan	35		
TAIWAN 2004	2004	<i>Rih-Jheng 101 and Ocean Researcher 2</i>	Taiwan	23		
TARASOC	2009	<i>Alis</i>	French Polynesia	213	3295-3507	
TERRASSES	2008	<i>Alis</i>	New Caledonia	99	3031-3129	
Vauban 1978-79	1978-79	<i>Vauban</i>	New Caledonia	47		
VOLSMAR	1989	<i>Alis</i>	New Hebrides Arc	25		Laboute <i>et al.</i> 1989

**APPENDIX 2**

Check-list of deep-sea decapod crustaceans (recorded below 100 m depth) from New Caledonia. An asterisk \* indicates that the species was originally described from New Caledonia. References to the original descriptions are listed in Appendix 4. The classification and the succession of the families follow De Grave *et al.* (2009).

**PENAEOIDEA****ARISTEIDAE**

- Aristaeomorpha foliacea* (Risso, 1827)  
*Aristaeopsis edwardsiana* (Johnson, 1868)  
*Aristeus mabahissae* Ramadan, 1938  
*Aristeus semidentatus* Bate, 1881  
*Aristeus virilis* (Bate, 1881)  
*Austropenaeus nitidus* (Barnard, 1947)  
*Hemipenaeus carpenteri* Wood-Mason & Alcock, 1891  
*Plesiopenaeus armatus* (Bate, 1881)

**BENTHESICYMIDAE**

- Benthesicymus howensis* Dall, 2001  
*Benthesicymus investigatoris* Alcock & Anderson, 1899  
*Benthesicymus urinator* Burkenroad, 1936  
*Gennadas bouvieri* Kemp, 1909  
*Gennadas capensis* Calman, 1925  
*Gennadas clavigarpus* de Man, 1907  
*Gennadas crassus* Tirmizi, 1960  
*Gennadas gilchristi* Calman, 1925  
*Gennadas incertus* (Balss, 1927)

**PENAEIDAE**

- Funchalia villosa* (Bouvier, 1905)  
*Metapenaeopsis difficilis* Crosnier, 1991  
*Metapenaeopsis evermanni* (Rathbun, 1906)  
*Metapenaeopsis provocatoria* Racek & Dall, 1965  
*Metapenaeopsis sibogae* (De Man, 1907)  
*Metapenaeopsis velutina* (Dana, 1852)  
*Parapenaeus cayrei* Crosnier, 2005  
*Parapenaeus fissurus* (Bate, 1881)  
*Parapenaeus investigatoris* Alcock & Anderson, 1899  
*Parapenaeus kensleyi* Crosnier, 2005 \*  
*Parapenaeus murrayi* Ramadan, 1938  
*Parapenaeus perezfarfantei* Crosnier, 1986  
*Parapenaeus sextuberculatus* Kubo, 1949  
*Penaeopsis balsii* Ivanov & Hassan, 1976  
*Penaeopsis challengerii* De Man, 1911  
*Penaeopsis eduardoi* Pérez Farfante, 1977  
*Penaeopsis jerryi* Pérez Farfante, 1979  
*Penaeopsis mclaughlini* Crosnier, 2006 \*  
*Penaeopsis rectacuta* (Bate, 1881)

**SICYONIIDAE**

- Sicyonia adunca* Crosnier, 2003 \*  
*Sicyonia altirostrum* Crosnier, 2003 \*  
*Sicyonia australiensis* Hanamura & Wedley, 1998  
*Sicyonia benthophila* De Man, 1907  
*Sicyonia curvirostris* Balss, 1913  
*Sicyonia dejouannei* Crosnier, 2003 \*  
*Sicyonia fallax* De Man, 1907  
*Sicyonia inflexa* (Kubo, 1949)  
*Sicyonia laevis* Bate, 1881  
*Sicyonia lancifer* (Olivier, 1811)

**SICYONIIDAE**

- Sicyonia rotunda* Crosnier, 2003 \*  
*Sicyonia trispinosa* De Man, 1907  
*Sicyonia truncata* (Kubo, 1949)  
*Sicyonia vitulans* (Kubo, 1949)

**SOLENOCERIDAE**

- Cryptopenaeus elevai* Crosnier, 1985  
*Cryptopenaeus crozieri* Pérez Farfante & Kensi, 1985  
*Gordonella kensleyi* Crosnier, 1988a \*  
*Gordonella paravillosa* Crosnier, 1988a \*  
*Hadropenaeus lucasi* (Bate, 1881)

**HADROPOENAEIDAE**

- Hadropenaeus spinicaudatus* Liu & Zhong, 1983  
*Haliporoides cristatus* Kensi, Tranter & Griffin, 1987  
*Haliporoides sibogae* sibogae (De Man, 1907)  
*Haliporus thetis* Faxon, 1893  
*Hymenopenaeus debillis* Smith, 1882  
*Hymenopenaeus equalis* (Bate, 1888)  
*Hymenopenaeus halli* Bruce, 1966  
*Hymenopenaeus methalli* Crosnier & Dall, 2004 \*  
*Hymenopenaeus neptunus* (Bate, 1881)  
*Hymenopenaeus obliquirostris* (Bate, 1881)  
*Hymenopenaeus propinquus* (De Man, 1907)  
*Mesopenaeus brucei* Crosnier, 1986  
*Solenocera comata* Stebbing, 1915

**SERGESTOIDEA**

- SERGESTIDAE**  
*Sergia talismani* (Barnard, 1947)

**STENOPODIODEA**

- SPONGICOLIDAE**  
*Globospongicola spinulatus* Komai & Saito, 2006  
*Paraspóngicola inflatus* (de Saint Laurent & Cleva, 1981)  
*Spongicola andamanicus* Alcock, 1901  
*Spongicola depressus* Saito & Komai, 2008 \*  
*Spongicola goyi* Saito & Komai, 2008 \*  
*Spongicola levigatus* Hayashi & Ogawa, 1987

**CARIDEA**

- ACANTHEPHYRIDAE**  
*Acanthephyra armata* A. Milne-Edwards, 1881  
*Acanthephyra curtirostris* Wood-Mason & Alcock, 1891  
*Acanthephyra eximia* Smith, 1884  
*Acanthephyra quadrispinosa* Kemp, 1939  
*Acanthephyra sanguinea* Wood-Mason [in Wood-Mason & Alcock, 1892]  
*Acanthephyra smithi* Kemp, 1939  
*Acanthephyra tenuipes* (Bate, 1888)

- ALPHEIDAE**  
*Heterogenys microphthalmus* (Smith, 1885)  
*Meningodora vesca* (Smith, 1886)

**ANCHISTIOIDIIDAE**

- Anchistioides willeyi* (Borradaile, 1900)

**BATHYPALEMONELLIDAE**

- Bathypalaemonella hayashii* Komai, 1995  
*Bathypalaemonella serratipalma* Pequegnat, 1970  
*Bathypalaemonetes brevirostris* (Bruce, 1986)  
*Bathypalaemonetes pilosipes* (Bruce, 1986)

**BRESILIIDAE**

- Bresilia antipodarum* Bruce, 1990

**CRANGONIDAE**

- Aegaeon lacazei* (Gourret, 1887)  
*Aegaeon rathbuni* De Man, 1918  
*Lissosabinea armata* Komai, 2006 \*  
*Lissosabinea indica* (De Man, 1918)  
*Lissosabinea unispinosa* Komai, 2006 \*  
*Parapontocaris aspera* Chace, 1984  
*Parapontophilus caledonicus* Komai, 2008 \*  
*Parapontophilus cyton* Komai, 2008 \*  
*Parapontophilus demani* (Chace, 1984)  
*Parapontophilus difficilis* Komai, 2008  
*Parapontophilus junceus* (Bate, 1888)  
*Parapontophilus longirostris* Komai, 2008  
*Parapontocaris levigata* Chace, 1984  
*Pontocaris propensalata* Bate, 1888  
*Pontocaris sibogae* (De Man, 1918)  
*Pseudopontophilus serratus* Komai, 2004a \*

**EUGONATONOTIDAE**

- Eugonatonotus chacei* Chan & Yu, 1991 \*

**GLYPHOCRANGONIDAE**

- Glyphocrangon armata* Komai, 2004b \*  
*Glyphocrangon conodactylus* Komai, 2004b \*  
*Glyphocrangon cornuta* Komai, 2004b \*  
*Glyphocrangon dimorpha* Komai, 2004b \*  
*Glyphocrangon musorstromia* Komai, 2006  
*Glyphocrangon parviocellus* Komai, 2006 \*  
*Glyphocrangon punctata* Komai, 2004 \*  
*Glyphocrangon regalis* Bate, 1888  
*Glyphocrangon richeri* Komai, 2004b  
*Glyphocrangon similior* Komai, 2004  
*Glyphocrangon speciosa* Komai, 2004b \*

**HIPPOLYTIDAE**

- Gelastreutes crosnieri* Bruce, 1990b \*

**NEMATOCARCINIDAE**

- Nematocarcinus combensis* Burukovsky, 2000  
*Nematocarcinus crosnieri* Burukovsky, 2000 \*  
*Nematocarcinus gracilis* Bate, 1888  
*Nematocarcinus parvus* Burukovsky, 2000

- NEMATOCARCINIDAE**
- Nematocarcinus paucidentatus* Bate, 1888
  - Nematocarcinus productus* Bate, 1888
  - Nematocarcinus richeri* Burukovsky, 2000
  - Nematocarcinus tenuipes* Bate, 1888
  - Nematocarcinus tenuirostris* Bate, 1888
  - Nigmatullinus acanthitelonis* (Pequegnat, 1970)
- OPLOPHORIDAE**
- Jancicella spinicauda* (A. Milne-Edwards, 1883)
  - Oplophorus gracilirostris* A. Milne-Edwards, 1881
  - Oplophorus spinosus* (Brullé, 1839)
  - Systellaspis debilis* (A.Milne- Edwards, 1881)
  - Systellaspis lanceoaustrata* Bate, 1888
  - Systellaspis pellucida* (Filhol, 1884)
- PALAEOMONIDAE**
- Altopontonia disparostris* Bruce, 1990a \*
  - Amphipontonia kanak* Bruce, 1991 \*
  - Anchistus pectinis* Kemp, 1925
  - Ancylomenes tenuirostris* (Bruce, 1991) \*
  - Cupates tenuipes* (Borradaile, 1898)
  - Dactylymon monnioti* (Bruce, 1990a) \*
  - Mesopontonia gorgoniophila* Bruce, 1967
  - Mesopontonia gracilicarpus* Bruce, 1990a \*
  - Mesopontonia monodactylus* Bruce, 1991 \*
  - Palaemonella dolichodactylus* Bruce, 1991 \*
  - Palaemonella rotumana* (Borradaile, 1898)
  - Paraclimenes franklini* (Bruce, 1990a) \*
  - Paraclimenes setirostris* (Bruce, 1990a) \*
  - Periclimenes jearcharoti* Bruce, 1991 \*
  - Periclimenes acanthimerus* Bruce, 2006 \*
  - Periclimenes albatrossae* Chace & Bruce, 1993
  - Periclimenes alcocki* Kemp, 1922
  - Periclimenes aleator* Bruce, 1991 \*
  - Periclimenes brevirostris* Bruce, 1991 \*
  - Periclimenes forcipulatus* Bruce, 1991 \*
  - Periclimenes forgesi* Li & Bruce, 2006 \*
  - Periclimenes foveolatus* Bruce, 1981
  - Periclimenes fujinoi* Bruce, 1990a \*
  - Periclimenes hertwigi* Bals, 1913
  - Periclimenes laccadivensis* (Alcock & Anderson, 1894)
  - Periclimenes latipollex* Kemp, 1922
  - Periclimenes leptodactylus* Bruce, 1991 \*
  - Periclimenes loyautensis* Li & Bruce, 2006 \*
  - Periclimenes ordinarius* Bruce, 1991 \*
  - Periclimenes paraleator* Li & Bruce, 2006 \*
  - Periclimenes parispinatus* Bruce, 1990a \*
  - Periclimenes pectinipes* Bruce, 1991 \*
  - Periclimenes platyrhynchus* Bruce, 1991 \*
  - Periclimenes rectirostris* Bruce, 1981
  - Periclimenes richeri* Bruce, 1990a \*
  - Periclimenes uniunguiculatus* Bruce, 1990a \*
  - Periclimenes vaubani* Bruce, 1990a \*
  - Urocaridella urocaridella* (Holthuis, 1950)
- PANDALIDAE**
- Bitias brevis* (Rathbun, 1906)
  - Calipandalus elachys* Komai & Chan, 2003
  - Chelonika macrochela* Fransen, 1997 \*
  - Heterocarpus amacula* Crosnier, 1988b
- HETEROCARPUS**
- Heterocarpus dorsalis* Bate, 1888
  - Heterocarpus ensifer* A. Milne-Edwards, 1881
  - Heterocarpus hayashii* Crosnier, 1988b \*
  - Heterocarpus intermedius* Crosnier, 1999 \*
  - Heterocarpus laevigatus* Bate, 1888
  - Heterocarpus lepidus* De Man, 1917
  - Heterocarpus longirostris* MacGilchrist, 1905
  - Heterocarpus parvispina* De Man, 1917
  - Heterocarpus sibogae* De Man, 1917
  - Heterocarpus tricarinatus* Alcock & Anderson, 1894
  - Heterocarpus woodmasoni* Alcock, 1901
  - Pandalus (Parapandalus) longirostris* Borradaile, 1900
  - Plesionika alcocki* (Anderson, 1896)
  - Plesionika bimaculata* Chan, 2004
  - Plesionika echinocola* Chan & Crosnier, 1991 \*
  - Plesionika ensis* (A. Milne-Edwards, 1881)
  - Plesionika erythrocyclus* Chan & Crosnier, 1997
  - Plesionika flavicauda* Chan & Crosnier, 1991
  - Plesionika hsuehyui* Chan, 2004
  - Plesionika laurentae* Chan & Crosnier, 1991 \*
  - Plesionika macropoda* Chace, 1939
  - Plesionika reflexa* Chace, 1985
  - Plesionika ruformaculata* Chan, 2004 \*
  - Plesionika scopifera* Chan, 2004 \*
  - Plesionika semilaevis* Bate, 1888
  - Plesionika serratifrons* (Borradaile, 1900)
  - Plesionika spinipes* Bate, 1888
  - Plesionika suffusa* Chan, 2004 \*
  - Pseudopandalus curvirostris* Crosnier, 1997a \*
- PASIPHAEIDAE**
- Leptochela (Leptochela) japonica* Hayashi & Miyake, 1969
  - Pasiphaea fragilis* Hayashi, 1999 \*
  - Pasiphaea gracilis* Hayashi, 1999 \*
  - Pasiphaea kapala* Kensley, Tranter & Griffin, 1987
  - Pasiphaea major* Hayashi, 2006 \*
  - Pasiphaea mclaughlinae* Hayashi, 2006 \*
  - Pasiphaea planidorsalis* Hayashi, 2004 \*
  - Pasiphaea sirenkoi* Burukovsky, 1987
  - Pasiphaea telacantha* Hayashi, 2004 \*
- PSALIDOPODIDAE**
- Psalidopus huxleyi* Wood-Mason [in Wood-Mason & Alcock, 1892]
- STYLODACTYLIDAE**
- Neostyloactylus affinis* Hayashi & Miyake, 1968
  - Parastyloactylus bimacillaris* Bate, 1888
  - Parastyloactylus richeri* Cleva, 1990 \*
  - Parastyloactylus semblata* Cleva, 1990 \*
  - Parastyloactylus tranterae* Cleva, 1990 \*
  - Stylocyloides crozieri* Cleva, 1990 \*
  - Stylocyclus discissipes* Bate, 1888
  - Stylocyclus laurentae* Cleva, 1990 \*
  - Stylocyclus libratus* Chace, 1983
  - Stylocyclus licinus* Chace, 1983
  - Stylocyclus macropus* Chace, 1983
  - Stylocyclus multidentatus* multidentatus Kubo, 1942
  - Stylocyclus profundus* Cleva, 1990 \*
  - Stylocyclus tokarensis* Zarenkov, 1968
- ASTACIDEA**
- ENOPLOMETOPIDAE**
- Enoplometopus gracilipes* (Saint Laurent, 1988)
- NEPHROPIDAE**
- Acanthacaris tenuimana* Bate, 1888
  - Nephropsis acanthura* Macpherson, 1990
  - Nephropsis suhmi* Bate, 1888
  - Nephropsis sulcata* Macpherson, 1990
  - Taumastochelus japonicus* Calman, 1913
- GLYPHEIDEA**
- GLYPHEIDAE**
- Laurentae glyphea neocaldonica* (Richer de Forges, 2006) \*
- AXIIDAE**
- AXIIDAE**
- Acanthaxius miyazakiensis* (Yokoya, 1933)
  - Ambiaxioides alcocki* (McArdle, 1900)
  - Bouvieraxius rufidus* (Rathbun, 1906)
  - Oxyrhynchaxius tricarinatus* Lin, 2006 \*
  - Spongianus holthuisi* Poore & Collins, 2010 \*
- CALLIANASSIDAE**
- Callianassa jaculatoria* De Man, 1905
  - Cheramus propinquus* (De Man, 1905)
- MICHELEIDAE**
- Meticonaxius dentatus* Lin, 2006 \*
  - Meticonaxius noumea* Poore, 1997 \*
  - Tethisea indica* Poore, 1994
- STRAHLAXIIDAE**
- Neaxius trondlei* Ngoc-Ho, 2005
- ACHELATA**
- PALINURIDAE**
- Linuparus meridionalis* Tsoi, Chan & Chu, 2011 \*
  - Linuparus sordidus* Bruce, 1965
  - Nupalirus chanii* (Poupin, 1994) \*
  - Nupalirus japonicus* Kubo, 1955
  - Palinustus unicornutus* Berry, 1979
  - Puerulus mesodontus*, Chan, Ma & Chu, this volume
  - Puerulus quadridentis*, Chan, Ma & Chu, this volume \*
  - Puerulus richeri*, Chan, Ma & Chu, this volume
  - Puerulus sericus*, Chan, Ma & Chu this volume \*
- SCYLLARIDAE**
- Bathyarctus chani* Holthuis, 2002 \*
  - Bathyarctus rubens* (Alcock & Anderson, 1894)
  - Eduarctus martensi* (Pfeffer, 1881)
  - Galearctus aurora* (Holthuis, 1982)
  - Galearctus avulsus* Yang, Chen & Chan, 2011 \*
  - Galearctus timidus* (Holthuis, 1960)
  - Ibacus brevipes* Bate, 1888
  - Ibacus brucei* Holthuis, 1977
  - Petrarctus brevicornis* (Holthuis, 1946)
  - Remiarctus bertholdii* (Paulson, 1875)
  - Scammarctus batei* (Holthuis, 1946)

**POLYCHELIDA****POLYCHELIDAE**

- Pentacheles laevis* Bate, 1878  
*Pentacheles obscurus* Bate, 1878  
*Pentacheles validus* A. Milne-Edwards, 1880  
*Polycheles baccatus* Bate, 1878  
*Polycheles coccifer* Galil, 2000  
*Polycheles enthrrix* (Bate, 1878)  
*Polycheles typhlops* Heller, 1862  
*Stereomastis aculeata* (Galil, 2000) \*  
*Stereomastis auriculata* (Bate, 1878)  
*Stereomastis helleri* (Bate, 1878)  
*Stereomastis nana* (Smith, 1884)  
*Stereomastis phosphorus* (Alcock, 1894)  
*Stereomastis surda* (Galil, 2000)

**ANOMURA****ALBUNEIDAE**

- Austrolepidopa caledonia* Boyko & Harvey, 1999 \*  
*Paralbunea dayriti* (Serène & Umalı, 1965)  
*Zygopa nortoni* Serène & Umalı, 1965

**CHIROSTYLIDAE**

- Chirostylus novaecaledoniae* Baba, 1991a \*  
*Gastropychus brevipropodus* Baba, 1991a \*  
*Gastropychus hendersoni* (Alcock & Anderson, 1899)  
*Gastropychus paucispina* Baba, 1991a \*  
*Gastropychus sternoornatus* (Van Dam, 1933)  
*Uroptychodes epigaster* Baba, 2004 \*  
*Uroptychodes musorstomi* Baba, 2004 \*  
*Uroptychodes spinimarginatus* (Henderson, 1885)  
*Uroptychus cavirostris* Alcock & Anderson, 1899  
*Uroptychus tridentatus* (Henderson, 1885)

**DIOGENIDAE**

- Areopaguristes micheleae* (Rahayu, 2005)  
*Areopaguristes ngochoae* (Rahayu, 2005)  
*Ciliopagurus alcocki* Forest, 1995 \*  
*Ciliopagurus kremppi* (Forest, 1952)  
*Ciliopagurus pacificus* Forest, 1995  
*Paguristes aciculus* Grant, 1905  
*Paguristes alcocki* McLaughlin & Rahayu, 2005  
*Paguristes ortmanni* Miyake, 1978  
*Paguristes palythophilus* Ortmann, 1892  
*Paguristes puniceus* Henderson, 1896  
*Paguristes versus* Komai, 2001  
*Pseudopaguristes laurentae* (Morgan & Forest, 1991)  
*Pseudopaguristes monoporus* (Morgan, 1987)  
*Strigopagurus boreonotus* Forest, 1995 \*

**EUMUNIDAE**

- Eumunida annulosa* de Saint Laurent & Macpherson, 1990 \*  
*Eumunida capillata* de Saint Laurent & Macpherson, 1990 \*  
*Eumunida karubar* de Saint Laurent & Poupin, 1996  
*Eumunida keijii* de Saint Laurent & Macpherson, 1990 \*  
*Eumunida marginata* de Saint Laurent & Macpherson, 1990 \*  
*Eumunida minor* de Saint Laurent & Macpherson, 1990 \*  
*Eumunida pacifica* Gordon, 1930  
*Eumunida parva* de Saint Laurent & Macpherson, 1990 \*  
*Eumunida spinosa* Macpherson, 2006b \*

- Eumunida sternomaculata* de Saint Laurent & Macpherson, 1990 \*  
*Pseudomunida fragilis* Haig, 1979

**GALATHEIDAE**

- Alainius crozieri* Baba, 1991b \*  
*Allogalathea babai* Cabezas, Macpherson & Machordom, 2011 \*  
*Allogalathea elegans* (Adams & White, 1848)  
*Allogalathea inermis* Cabezas, Macpherson & Machordom, 2011 \*  
*Galathea barbellata* Macpherson, 2012 \*  
*Galathea echinata* Macpherson, 2012 \*  
*Galathea profunda* Macpherson, 2012  
*Galathaea sanctae* Macpherson, 2012  
*Leiogalathea laevirostris* (Balss, 1913)  
*Phylladiohynchus ikedai* (Miyake & Baba, 1965)  
*Phylladiohynchus pusillus* (Henderson, 1885)

**LITHODIDAE**

- Lithodes richeri* Macpherson, 1990 \*  
*Neolithodes bronwynae* Ahyong, 2010  
*Paralomis dawsoni* Macpherson, 2001 \*  
*Paralomis haigae* Eldredge, 1976

**MUNIDIDAE**

- Agononida andrewi* (Macpherson, 1994) \*  
*Agononida callirhoe* (Macpherson, 1994) \*  
*Agononida eminens* (Baba, 1988)  
*Agononida incerta* (Henderson, 1888)  
*Agononida laurentae* (Macpherson, 1994) \*  
*Agononida marini* (Macpherson, 1994) \*  
*Agononida norfocerta* Poore & Andreakis, 2012  
*Agononida normani* (Henderson, 1885)  
*Agononida ocyrhoe* (Macpherson, 1994) \*  
*Agononida rubrizonata* Macpherson & Baba, 2009 \*  
*Agononida sabatesae* (Macpherson, 1994) \*  
*Agononida soelae* (Baba, 1986)  
*Agononida sphecia* (Macpherson, 1994) \*  
*Agononida squamosa* (Henderson, 1885)  
*Anoplionda inermis* (Baba, 1994)  
*Babamunida callista* (Macpherson, 1994) \*  
*Babamunida javieri* (Macpherson, 1994) \*  
*Bathymunida balssi* Van Dam, 1938  
*Bathymunida eurybregma* Baba & de Saint Laurent, 1996 \*  
*Bathymunida nebulosa* Baba & de Saint Laurent, 1996 \*  
*Bathymunida ocularis* Baba & de Saint Laurent, 1996 \*  
*Bathymunida rufa* Baba & de Saint Laurent, 1996 \*  
*Bathymunida sibogae* Van Dam, 1938  
*Crosnierita dicata* Macpherson, 1998 \*  
*Crosnierita urizae* (Macpherson, 1994) \*  
*Crosnierita yante* (Macpherson, 1994)  
*Enriquea leviantennata* (Baba, 1988)  
*Heteronida aspinirostris* (Khodkina, 1981)  
*Munida acantha* Macpherson, 1994 \*  
*Munida alonsoi* Macpherson, 1994 \*  
*Munida amblytes* Macpherson, 1994 \*  
*Munida angusta* Macpherson, 2004  
*Munida armilla* Macpherson, 1994 \*  
*Munida asprosoma* Ahyong & Poore, 2004  
*Munida aulakodes* Macpherson, 2006b \*  
*Munida barangae* Macpherson, 1994 \*
- Munida brachytes* Macpherson, 1994 \*  
*Munida clinata* Macpherson, 1994 \*  
*Munida congesta* Macpherson, 2000  
*Munida delicata* Macpherson, 2004 \*  
*Munida devestiva* Macpherson, 2006b \*  
*Munida distiza* Macpherson, 1994 \*  
*Munida eclepsis* Macpherson, 1994 \*  
*Munida elachia* Macpherson, 1994 \*  
*Munida erato* Macpherson, 1994 \*  
*Munida gillii* Macpherson, 1993  
*Munida gordoe* Macpherson, 1994 \*  
*Munida guttata* Macpherson, 1994 \*  
*Munida hyalina* Macpherson, 1994 \*  
*Munida idyia* Macpherson, 1994 \*  
*Munida inornata* Henderson, 1885  
*Munida japonica* Stimpson, 1858  
*Munida leagora* Macpherson, 1994 \*  
*Munida leptitis* Macpherson, 1994 \*  
*Munida lineola* Macpherson, 1994 \*  
*Munida magniantennulata* Baba & Türkay, 1992  
*Munida masi* Macpherson, 1994 \*  
*Munida microps* Alcock, 1894  
*Munida moliae* Macpherson, 1994 \*  
*Munida notata* Macpherson, 1994 \*  
*Munida ornata* Macpherson, 2004 \*  
*Munida oritea* Macpherson & Baba, 1993  
*Munida pagesi* Macpherson, 1994 \*  
*Munida parca* Macpherson, 1996a \*  
*Munida parile* MacPherson & Machordom, 2005 \*  
*Munida pectinata* MacPherson & Machordom, 2005 \*  
*Munida pontoporeia* Macpherson, 1994 \*  
*Munida proto* Macpherson, 1994 \*  
*Munida psamathe* Macpherson, 1994 \*  
*Munida pseliophora* Macpherson, 1994 \*  
*Munida psylla* Macpherson, 1994 \*  
*Munida pygmaea* Macpherson, 1996a \*  
*Munida rhodonia* Macpherson, 1994 \*  
*Munida rogeri* Macpherson, 1994 \*  
*Munida rosula* Macpherson, 1994 \*  
*Munida rubridigitata* Baba, 1994  
*Munida rufiantennulata* Baba, 1969  
*Munida runcinata* Macpherson, 1994 \*  
*Munida sacksi* Macpherson, 1993  
*Munida sao* Macpherson, 1994 \*  
*Munida semoni* Ortmann, 1894  
*Munida simulatrix* MacPherson & Machordom, 2005 \*  
*Munida spilota* Macpherson, 1994 \*  
*Munida stia* Macpherson, 1994 \*  
*Munida stigmatica* Macpherson, 1994 \*  
*Munida striola* Macpherson & Baba, 1993  
*Munida taenia* Macpherson, 1994 \*  
*Munida theo* Macpherson, 1994 \*  
*Munida tiresias* Macpherson, 1994 \*  
*Munida tuberculata* Henderson, 1885  
*Munida tyche* Macpherson, 1994 \*  
*Munida typhle* Macpherson, 1994 \*  
*Munida zebra* Macpherson, 1994 \*  
*Onconida alaini* Baba & de Saint Laurent, 1996 \*  
*Onconida tropis* Baba & de Saint Laurent, 1996  
*Paramunida amphitrita* Macpherson, 1996b

- Paramunida antares* Cabezas, Macpherson & Machordom, 2010 \*
- Paramunida belone* Macpherson, 1993 \*
- Paramunida granulata* (Henderson, 1885)
- Paramunida labis* Macpherson, 1996b
- Paramunida longior* Baba, 1988
- Paramunida microrhina* Cabezas, Macpherson & Machordom, 2010 \*
- Paramunida parvispina* Cabezas, Macpherson & Machordom, 2010 \*
- Paramunida pictura* Macpherson, 1993 \*
- Paramunida pronoe* Macpherson, 1993 \*
- Paramunida setigera* Baba, 1988
- Paramunida stictas* Macpherson, 1993 \*
- Paramunida tenera* Cabezas, Macpherson & Machordom, 2010 \*
- Paramunida thalie* Macpherson, 1993 \*
- Plesionida aliena* (Macpherson, 1996a) \*
- Plesionida psila* Baba & de Saint Laurent, 1996 \*
- Raymunida bellior* (Miyake & Baba, 1967)
- Raymunida confundens* Macpherson & Machordom, 2001 \*
- Raymunida dextralis* Macpherson & Machordom, 2001 \*
- Torbenella calvata* (Macpherson, 2006b) \*
- Torbenella insolita* (Macpherson, 2004) \*
- Torbenella orbis* (Baba, 2005)
- MUNIDOPSISAE**
- Galacantha bellis* Henderson, 1885
- Galacantha quiquei* Macpherson, 2007
- Galacantha rostrata* A. Milne Edwards, 1880
- Leiogalathea laevirostris* (Balss, 1913)
- Munidopsis analoga* Macpherson, 2007 \*
- Munidopsis andamanica* MacGilchrist, 1905
- Munidopsis arenula* Macpherson, 2007 \*
- Munidopsis carinimarginata* Baba, 1988
- Munidopsis centrina* Alcock & Anderson, 1894
- Munidopsis ceres* Macpherson, 2007 \*
- Munidopsis cornuata* Macpherson, 2007 \*
- Munidopsis cylindrophthalma* (Alcock, 1894)
- Munidopsis debilis* (Henderson, 1885)
- Munidopsis kensleyi* Ahyong & Poore, 2004
- Munidopsis latimanata* Miyake & Baba, 1966
- Munidopsis nitida* (A. Milne Edwards, 1880)
- Munidopsis orcina* McArdle, 1901
- Munidopsis sinclairi* McArdle, 1901
- Munidopsis tafrii* Osawa, Lin & Chan, 2006
- Munidopsis ternaria* Macpherson, 2007 \*
- Munidopsis treis* Ahyong & Poore, 2004
- Munidopsis trifida* Henderson, 1885
- PAGURIDAE**
- Alainopaguroides megalophthalmus* McLaughlin, 2006 \*
- Alainopagurus crozieri* Lemaitre & McLaughlin, 1995 \*
- Anapagurus bonnieri* Nobili, 1905
- Catapagurus danida* McLaughlin, 2002
- Catapagurus franklinae* McLaughlin, 2004a \*
- Catapagurus imperialis* (Asakura, 2001)
- Catapagurus spinicarpus* de Saint Laurent & McLaughlin, 2000
- Catapagurus tanimbarensis* McLaughlin, 1997
- Diacanthurus ecpyma* McLaughlin & Forest, 1997 \*
- Diacanthurus richeri* McLaughlin & Forest, 1997 \*
- Icelopagurus crozieri* McLaughlin, 1997
- Icelopagurus undulatus* McLaughlin, 2006 \*
- Michelopagurus limatulus* (Henderson, 1888)
- Micropagurus Polynesiensis* (Nobili, 1906)
- Micropagurus spinimanus* Asakura, 2005 \*
- Nematopagurus alcocki* McLaughlin, 1997
- Nematopagurus australis* (Henderson, 1888)
- Nematopagurus diadema* Lewinsohn, 1969
- Nematopagurus gardineri* Alcock, 1905
- Nematopagurus indicus* Alcock, 1905
- Nematopagurus kosiensis* McLaughlin, 1998
- Nematopagurus lepidochirus* (Doflein, 1902)
- Nematopagurus lewinsohni* Türkay, 1986
- Nematopagurus meiringae* McLaughlin, 1998
- Nematopagurus ricei* McLaughlin, 2004b \*
- Nematopagurus richeri* McLaughlin, 2004b \*
- Nematopagurus scutelliformis* McLaughlin, 1997
- Nematopagurus spinulosensis* McLaughlin & Brock, 1974
- Nematopagurus spongiorparticeps* McLaughlin, 2004b \*
- Nematopagurus vallatus* (Melin, 1939)
- Pagurojacquesia polymorpha* (de Saint Laurent & McLaughlin, 1999)
- Porcellanopagurus chiltoni* de Saint Laurent & McLaughlin, 2000
- Porcellanopagurus filholi* de Saint Laurent & McLaughlin, 2000
- Porcellanopagurus haptodactylus* McLaughlin, 2000 \*
- Porcellanopagurus tridentatus* Whitelegge, 1900
- Propagurus haigae* (McLaughlin, 1997)
- Pteropagurus inermis* McLaughlin & Rahayu, 2006 \*
- Pteropagurus spina* McLaughlin & Rahayu, 2006 \*
- Pteropagurus spinulocarpus* McLaughlin, 2007a \*
- Solitariopagurus triporbolus* Poupin & McLaughlin, 1996
- Solitariopagurus trullirostris* McLaughlin, 2000 \*
- Solitariopagurus tuerkayi* McLaughlin, 1997
- Tomopaguroides valdiviae* (Bals, 1911)
- Turleania boucheti* McLaughlin, 2007b
- Turleania multispina* McLaughlin, 1997
- Turleania senticosia* (McLaughlin & Haig, 1996)
- Xylopagurus caledonicus* Forest, 1997 \*
- PARAPAGURIDAE**
- Oncopagurus conicus* Lemaitre, 2006 \*
- Paragiopagurus acutus* (de Saint Laurent, 1972)
- Paragiopagurus bicarinatus* (de Saint Laurent, 1972)
- Paragiopagurus boletifer* (de Saint Laurent, 1972)
- Paragiopagurus diogenes* (Whitelegge, 1900)
- Paragiopagurus hirsutus* (de Saint Laurent, 1972)
- Paragiopagurus orthotenes*, Lemaitre, this volume \*
- Paragiopagurus oxychelos*, Lemaitre, this volume \*
- Paragiopagurus ruticcheles* (A. Milne-Edwards, 1891)
- Paragiopagurus schnauzeri* Lemaitre, 2006 \*
- Paragiopagurus trilineatus*, Lemaitre, this volume \*
- Paragiopagurus umbonatus*, Lemaitre, this volume \*
- Parapagurus furici* Lemaitre, 1999 \*
- Parapagurus latimanus* Henderson, 1888
- Parapagurus richeri* Lemaitre, 1999 \*
- Strobopagurus breviacus* Lemaitre, 2004a \*
- Strobopagurus gracilipes* (A. Milne-Edwards, 1891)
- Strobopagurus sibogae* (de Saint Laurent, 1972)
- Sympagurus acinops* Lemaitre, 1989 \*
- Sympagurus affinis* (Henderson, 1888)
- Sympagurus aurantium* Lemaitre, 2004b \*
- Sympagurus brevipes* (de Saint Laurent, 1972)
- Sympagurus burkenroadi* Thompson, 1943
- Sympagurus dofleini* (Balss, 1912)
- Sympagurus planimanus* (de Saint Laurent, 1972)
- Sympagurus pouponi* Lemaitre, 1994
- Sympagurus soela* Lemaitre, 1996 \*
- Sympagurus symmetricus* Lemaitre, 2004b \*
- Sympagurus trispinosus* (Balss, 1911)
- Sympagurus villosus* Lemaitre, 1996 \*
- Tylaspis anomala* Henderson, 1885
- PYLOCHELIDAE**
- Cheiroplata pumicicola* Forest, 1987
- Trizocheles caledonicus* Forest, 1987 \*
- Trizocheles piligrimi* Forest & McLaughlin, 2000
- Trizocheles pulcher* Forest, 1987 \*
- Trizocheles spinosus* (Henderson, 1888)
- Trizocheles vaubanae* McLaughlin & Lemaitre, 2008 \*
- BRACHYURA**
- AETHRIDAE**
- Aethra scruposa* (Linnaeus, 1764)
- CALAPPIDAE**
- Mursia armata* De Haan, 1837
- Mursia australiensis* Campbell, 1971
- Mursia longispina* Crosnier, 1997b \*
- Mursia microspina* Davie & Short, 1989
- Mursia musorstromia* Galil, 1993 \*
- Mursia trispinosa* Parisi, 1914
- CANCRIDAE**
- Platepistoma nanum* Davie, 1991 \*
- CRYPTOCHIRIDAE**
- Cecidocarcinus zibrowii* Manning, 1991 \*
- CYCLODORIPPIDAE**
- Ketamia depressa* (Ihle, 1916)
- Krangalangia orstom* Tavares, 1993 \*
- Krangalangia spinosa* (Zarenkov, 1970)
- Tymolus daviei* Tavares, 1997
- Xeinostoma richeri* Tavares, 1993 \*
- CYMONOMIDAE**
- Elassopodus stellatus* Tavares, 1993 \*
- DROMIIDAE**
- Cryptodromia coronata* Stimpson, 1858
- Cryptodromia hilgendorfi* de Man, 1888
- Eodromia denticulata* McLay, 1993 \*
- Epigodromia areolata* (Ihle, 1913)
- Epigodromia ebalioides* (Alcock, 1899)
- Epigodromia nodosa* Sakai, 1936
- Epigodromia rotunda* McLay, 1993 \*
- Frodromia atypica* (Sakai, 1936)
- Lewindromia unidentata* (Rüppell, 1830)
- Metadromia wilsoni* (Fulton & Grant, 1902)
- Petalomera pulchra* Miers, 1884
- Sphaerodromia lamellata* Crosnier, 1994 \*

*Stirometra foresti* (McClay, 1993) \*  
*Takidromia cristatipes* (Sakai, 1969)  
*Takidromia longispina* McClay, 1993 \*

**DYNOMENIDAE**

*Dynomene hispida* (Latreille, in Milbert, 1812)  
*Dynomene pilumnoides* Alcock, 1900  
*Metadynomene tanensis* (Yokoya, 1933)  
*Paradynomene demon* McClay & Ng, 2004 \*  
*Paradynomene diablo* McClay & Ng, 2004 \*  
*Paradynomene quasimodo* McClay & Ng, 2004  
*Paradynomene tuberculata* Sakai, 1963

**EPIALTIDAE**

*Guinotinia cordis* Richer de Forges & Ng, 2009a \*  
*Oxypleurodon alaini* Richer de Forges & Ng, 2009b \*  
*Oxypleurodon mammatum* (Guinot & Richer de Forges, 1986b) \*  
*Oxypleurodon orbiculatum* (Guinot & Richer de Forges, 1986a) \*  
*Oxypleurodon stuckiae* (Guinot & Richer de Forges, 1986a) \*  
*Stegopleurodon planirostrum* Richer de Forges & Ng, 2009b \*

**ETHUSIDAE**

*Ethusa abbreviata* Castro, 2005a  
*Ethusa crosnieri* Chen, 1993 \*  
*Ethusa curvipes* Chen, 1993 \*  
*Ethusa furca* Chen, 1993 \*  
*Ethusa granulosa* Ihle, 1916  
*Ethusa indica* Alcock, 1894  
*Ethusa izuensis* Sakai, 1937  
*Ethusa magnipalmata* Chen, 1993 \*  
*Ethusa minuta* Sakai, 1937  
*Ethusa obliquedens* Chen, 1993 \*  
*Ethusa orientalis* Miers, 1886  
*Ethusa parapygmaea* Chen, 1993 \*  
*Ethusa pygmaea* Alcock, 1894  
*Ethusa quadrata* Sakai, 1937  
*Ethusina brevidentata* Chen, 1993 \*  
*Ethusina ciliacirrata* Castro, 2005a  
*Ethusina coronata* Castro, 2005a  
*Ethusina dilobatus* Chen, 1993 \*  
*Ethusina microspina* Chen, 2000  
*Ethusina paralongipes* Chen, 1993 \*  
*Ethusina pubescens* Chen, 1993 \*  
*Ethusina robusta* (Miers, 1886)  
*Ethusina stenommata* Castro, 2005a  
*Ethusina vanuatensis* Chen, 2000

**EURYPLACIDAE**

*Eucrate laevis* (Borradaile, 1903)

**GERYONIDAE**

*Chaceon bicolor* Manning & Holthuis, 1989 \*

**GONEPLACIDAE**

*Carcinoplax cooki* Rathbun 1906  
*Carcinoplax cracens* Castro, 2007 \*  
*Carcinoplax ischurodous* (Stebbing, 1923)  
*Carcinoplax specularis* Rathbun, 1914  
*Carcinoplax tenuidentata* Castro, 2007  
*Carcinoplax tuberosa* Castro, 2007 \*

*Carcinoplax uncinata* Castro, 2009 \*  
*Goneplacoides marivenae* (Komatsu & Takeda, 2004)  
*Menoplax longispinosa* (Chen, 1984)  
*Microgoneplax elegans* (Chen, 1998)  
*Microgoneplax prius* Castro, 2007 \*  
*Neogoneplax renoculis* (Rathbun, 1914)  
*Neogoneplax serratipes* Castro, 2007 \*

*Paragoneplax serenei* (Zarenkov, 1972)  
*Psopheticus stridulans* Wood-Mason, 1892  
*Psopheticus vocans* Guinot, 1985  
*Pycnoplax aspera* Castro, 2009 \*

*Pycnoplax suruguensis* (Rathbun, 1932)  
*Thyraplax cooki* (Rathbun, 1906)  
*Thyraplax crosnieri* (Guinot & Richer de Forges, 1981a) \*  
*Thyraplax truncata* Castro, 2007

**HEXAPODIDAE**

*Hexaplex megalops* Doflein, 1904

**HOMOLIDAE**

*Dagnaudus petterdi* (Grant, 1905)  
*Homola coriolisi* Guinot & Richer de Forges, 1995 \*  
*Homola mienensis* Sakai, 1979  
*Homola orientalis* Henderson, 1888  
*Homola ranunculus* Guinot & Richer de Forges, 1995 \*  
*Homolochunia kular* Griffin & Brown, 1976  
*Homolochunia valdiviae* Doflein, 1904  
*Homologenus levii* Guinot & Richer de Forges, 1995 \*  
*Homolomanria sibogae* Ihle, 1912  
*Ihlopsis tirardi* Guinot & Richer de Forges, 1995 \*  
*Lamoha inflata* (Guinot & Richer de Forges, 1981b) \*  
*Latreillopsis antennata* Guinot & Richer de Forges, 1995 \*  
*Latreillopsis bispinosa* Henderson, 1888  
*Latreillopsis gracilipes* Guinot & Richer de Forges, 1981b \*  
*Moloha majora* (Kubo, 1936)  
*Paromola bathyalis* Guinot & Richer de Forges, 1995 \*  
*Paromolopsis boosi* Wood-Mason & Alcock, 1891

**HOMOLODROMIIDAE**

*Dicranodromia foersteri* Guinot, 1993 \*  
*Dicranodromia spinulata* Guinot, 1995 \*  
*Homolodromia kai* Guinot, 1993

**HYPOTHALASSIIDAE**

*Hypothalassia armata* (De Haan, 1835)

**INACHIDAE**

*Cyrtomaia coriolisi* Richer de Forges & Guinot, 1988 \*  
*Cyrtomaia cornuta* Richer de Forges & Guinot, 1988 \*  
*Cyrtomaia ericina* Guinot & Richer de Forges, 1982 \*  
*Cyrtomaia furici* Richer de Forges & Guinot, 1988 \*  
*Cyrtomaia griffini* Richer de Forges & Guinot, 1990  
*Cyrtomaia horrida* Rathbun, 1916  
*Cyrtomaia ihlei* Guinot & Richer de Forges, 1982 \*  
*Cyrtomaia platypes* Yokoya, 1933  
*Grypachaeus hyalinus* Alcock & Anderson, 1894  
*Platymaia fimbriata* Rathbun, 1916  
*Platymaia rebierei* Guinot & Richer de Forges, 1986a  
*Platymaia wyvillethomsoni* Miers, 1886  
*Pleistacantha cervicornis* Ihle & Ihle-Landenberg, 1931

*Pleistacantha exophthalmus* Guinot & Richer de Forges, 1982 \*  
*Pleistacantha cf japonica* (Yokoya, 1933)

**LATREILLIIDAE**

*Eplumula australiensis* (Henderson, 1888)  
*Latreillia metanesa* Williams, 1982  
*Latreillia pennifera* Alcock, 1900  
*Latreillia valida* De Haan, 1839

**LEUCOSIIDAE**

*Alox ornatum* (Ihle, 1918)  
*Ancylodactyla elata* (Zarenkov, 1994)  
*Ancylodactyla elongata* (Zarenkov, 1969)  
*Ancylodactyla nana* (Zarenkov, 1990)  
*Arcania elongata* Yokoya, 1933  
*Arcania tuberculata* Bell, 1855  
*Galilia narusei* Ng & Richer de Forges, 2007 \*  
*Myra curtimana* Galil, 2001a \*  
*Myra eudactylus* (Bell, 1855)  
*Myra fugax* (Fabricius, 1798)  
*Tanaoa serenei* (Richer de Forges, 1983) \*  
*Toru septimus* Galil, 2003 \*  
*Urnalana granulimera* Galil, 2005

**MATHILDELLIDAE**

*Intesius pilosus* Guinot & Richer de Forges, 1981a \*  
*Intesius richeri* Crosnier & Ng, 2004 \*  
*Mathildella maxima* Guinot & Richer de Forges, 1981a \*  
*Platypilumnus jamiesoni* Richer de Forges, 1996 \*

**OZIIDAE**

*Eupilumnus laciniatus* (Sakai, 1980)

**PALICIDAE**

*Miropalicus vietnamensis* (Zarenkov, 1968)  
*Paliculus kyusyuensis* (Yokoya, 1933)  
*Parapalicus ambonensis* Moosa & Serène, 1981  
*Parapalicus clinodentatus* Castro, 2000  
*Parapalicus denticulatus* Castro, 2000 \*  
*Parapalicus inanis* Castro, 2000 \*  
*Parapalicus pируensis* Moosa & Serène, 1981  
*Parapalicus trispialis* Castro, 2000  
*Pseudopalicus acanthodactylus* Castro, 2000 \*  
*Pseudopalicus amadaibai* (Sakai, 1963)  
*Pseudopalicus declivis* Castro, 2000 \*  
*Pseudopalicus glaber* Castro, 2000 \*  
*Pseudopalicus investigatoris* Alcock, 1900  
*Pseudopalicus oahuensis* (Rathbun, 1906)  
*Rectopalicus amphiceros* Castro, 2000 \*  
*Rectopalicus ampullatus* Castro, 2000 \*  
*Rectopalicus woodmasoni* (Alcock, 1900)

**PARTHENOPIDAE**

*Garthambus pouponi* (Garth, 1993)  
*Pseudolambrus guinotae* Tan, 2010 \*

**PHYLLOTYMOLINIDAE**

*Genka kejii* Tavares, 1993 \*  
*Phyllotymlinum crosnieri* Tavares, 1993 \*

**PILUMNIDAE**

*Eumedonus brevihynchus* Chia & Ng, 2000 \*

**PLAGUSIIDAE**

*Euchirograpsus timorensis* Türkay, 1975

*Miersiograpsus australiensis* Türkay, 1978

**POLYBIIDAE**

*Benthochascon herningi* Alcock & Anderson, 1899

*Brusinia elongata* (Sakai, 1969)

*Brusinia profunda* Moosa, 1996 \*

*Ovalipes iridescentis* (Miers, 1886)

*Parathranites granosus* Crosnier, 2002b

*Parathranites intermedius* Crosnier, 2002b \*

*Parathranites orientalis* (Miers, 1886)

**PONTUNIDAE**

*Charybdis (Charybdis) rufodactylus* Stephenson & Rees, 1968

*Cycloachelous orbitosinus* (Rathburn, 1911)

*Echinolatus caledonicus* (Moosa, 1996) \*

*Liocarcinus vernalis* (Risso, 1816)

*Lissocarcinus polybioides* Adams & White, 1849

*Lupocyclus philippinensis* Semper, 1880

*Lupocyclus tugelae* Barnard, 1950

*Nectocarcinus pubescens* Moosa, 1996 \*

*Portunus (Monomia) gladiator* Fabricius, 1798

**PORTUNIDAE**

*Portunus (Monomia) lecromi* Moosa, 1996 \*

*Portunus (Xiphonectes) hastatooides* Fabricius, 1798

*Portunus (Xiphonectes) stephensi* Moosa, 1981 \*

*Thalamita spinifera* Borradale, 1902

**PROGERYONIDAE**

*Progeryon vaubani* Guinot & Richer de Forges, 1981a \*

*Rhadinoplax micropthalmus* (Guinot & Richer de Forges, 1981a) \*

**RANINIDAE**

*Cosmonotus mclaughlinae* Tavares, 2006

*Lyreidus brevifrons* Sakai, 1937

*Ranina ranina* (Linnaeus, 1758)

**RETROPLUMIDAE**

*Retropluma serenei* Saint Laurent, 1989

*Retropluma solomonensis* McLay, 2006

**TRAPEZIIDAE**

*Calocarcinus africanus* Calman, 1909

*Calocarcinus crosnieri* Galil & Clark, 1990 \*

*Quadrella maculosa* Alcock, 1898

**TRICHOPELTARIIDAE**

*Trichopeltarion vanuatuensis* Tavares & Cleva, 2010

**XANTHIDAE**

*Alainodaeus alis* Davie, 1997 \*

*Alainodaeus rimatara* Davie, 1993

*Antrocarcinus petrosus* Ng & Chia, 1994 \*

*Demania cultripes* (Alcock, 1898)

*Demania garthi* Guinot & Richer de Forges, 1981a \*

*Demania intermedia* Guinot, 1969

*Demania mortensenii* (Odhner, 1925)

*Demania wardi* Garth & Ng, 1985

*Euryxanthops latifrons* Davie, 1997 \*

*Gaillardiellus bathus* Davie, 1997 \*

*Glyptocarcinus politus* Ng & Chia, 1994 \*

*Liomera nigrimanus* Davie, 1997 \*

*Lophozomyrus bertoni* Guinot & Richer de Forges, 1981a \*

*Medaeops gemini* Davie, 1997 \*

*Medaeops merodontos* Davie, 1997 \*

*Medaeus aztec* Davie, 1997 \*

*Meractaea multidentata* Davie, 1997 \*

*Miersiella haswelli* (Miers, 1886)

*Palatigum trichostoma* Davie, 1997 \*

*Paramedaeus globosus* Serène & Vadon, 1981 \*

*Paraxanthodes cumatodes* (MacGilchrist, 1905)

*Rata chalcal* Davie, 1997 \*

*Xanthias teres* Davie, 1997 \*

**APPENDIX 3**

New species of decapod crustaceans (recorded below 100 m depth) described from South Pacific island groups (other than New Caledonia) based on the *Tropical Deep-Sea Benthos* expeditions material. For references, see Appendix 4.

PENAEOIDEA	
PENAEIDAE	
<i>Parapenaeus cayrei</i> Crosnier, 2005	Tonga
SOLENOCERIDAE	
<i>Maximiliaeus odoceros</i> Chan, 2012	PNG
STENOPODIDEA	
SPONGICOLIDAE	
<i>Globospionicola spinulatus</i> Komai & Saito, 2006	Vanuatu
CARIDEA	
CRANGONIDAE	
<i>Metacrangon clevai</i> Komai, 2012	Solomon
<i>Parapontophilus cornutus</i> Komai, 2008	French Polynesia
<i>Parapontophilus difficilis</i> Komai, 2008	Vanuatu
<i>Parapontophilus longirostris</i> Komai, 2008	French Polynesia
<i>Parapontophilus stenorhinus</i> Komai, 2008	Tonga
GLYPHOCRANGONIDAE	
<i>Glyphocrangon arduus</i> Komai, 2007	French Polynesia
<i>Glyphocrangon boa</i> Komai, 2011	Vanuatu
<i>Glyphocrangon musorstomia</i> Komai, 2006	Wallis & Futuna
<i>Glyphocrangon prostrata</i> Komai, 2011	Solomon
<i>Glyphocrangon richeri</i> Komai, 2004b	Wallis & Futuna
<i>Glyphocrangon rubricinctuta</i> Komai, 2004b	Wallis & Futuna
<i>Glyphocrangon similior</i> Komai, 2004b	Vanuatu
HYPOPLYTIDAE	
<i>Eumanninigia pliarthon</i> Crosnier, 2000	Wallis & Futuna
NEMATOCARCINIDAE	
<i>Nematocarcinus machaerophorus</i> Burukovsky, 2003	French Polynesia
PALAEOMONIDAE	
<i>Izucaris crosnieri</i> Li, 2008	French Polynesia
<i>Palaemonella komaii</i> Li & Bruce, 2006	Tonga
<i>Periclimenes paralcocki</i> Li & Bruce, 2006	Wallis & Futuna
<i>Periclimenes pouponi</i> Bruce, 1989	French Polynesia
PANDALIDAE	
<i>Heterocarpus amacula</i> Crosnier, 1988b	French Polynesia
<i>Heterocarpus corona</i> Yang, Chan & Chu, 2010	Vanuatu
<i>Heterocarpus tenuidentatus</i> Cleva & Crosnier, 2006	Solomon
<i>Plesionika bimaculata</i> Chan, 2004	Vanuatu
<i>Plesionika carsini</i> Crosnier, 1986a	French Polynesia
<i>Plesionika curvata</i> Chan & Crosnier, 1991	French Polynesia
<i>Plesionika erythrocyclus</i> Chan & Crosnier, 1997	French Polynesia
<i>Plesionika fennieri</i> Crosnier, 1986b	French Polynesia
PENAEOIDEA	
<i>Plesionika flavicauda</i> Chan & Crosnier, 1991	French Polynesia
<i>Plesionika payeni</i> Chan & Crosnier, 1997	French Polynesia
<i>Plesionika picta</i> Chan & Crosnier, 1997	French Polynesia
<i>Plesionika pouponi</i> Chan & Crosnier, 1997	French Polynesia
<i>Plesionika protati</i> Chan & Crosnier, 1997	French Polynesia
<i>Plesionika rubrior</i> Chan & Crosnier, 1991	French Polynesia
STYLODACTYLIDAE	
<i>Stylocactus brucei</i> Cleva, 1994	Wallis & Futuna
ASTACIDEA	
ENOPLOMETOPIDAE	
<i>Enoplometopus gracilipes</i> (de Saint Laurent, 1988)	French Polynesia
AXIIDAE	
AXIIDAE	
<i>Acanthaxius clevai</i> Ngoc-Ho, 2006	Solomon
STRAHLAXIIDAE	
<i>Neaxius trondlei</i> Ngoc-Ho, 2005	French Polynesia
THOMASSINIIDAE	
<i>Crosniera dayrati</i> Ngoc-Ho, 2005	French Polynesia
ACHELATA	
PALINURIDAE	
<i>Nupalirus vericeli</i> (Poupin, 1994)	French Polynesia
<i>Puerulus richeri</i> Chan, Ma & Chu, this volume	French Polynesia
SCYLLARIDAE	
<i>Educarctus marginatus</i> Holthuis, 2002	Fiji
<i>Galearctus rapanus</i> (Holthuis, 1993)	French Polynesia
<i>Chelarctus crosnieri</i> Holthuis, 2002	Tonga
POLYCHELIDA	
POLYCHELIDAE	
<i>Stereomastis alis</i> (Ahyong & Galil, 2006)	French Polynesia
ANOMURA	
ALBUNEIDAE	
<i>Albunea marquisiana</i> Boyko, 2000	French Polynesia

<b>DIOGENIDAE</b>		
<i>Calcinus albengai</i> Poupin & Lemaitre, 2003 .....	French Polynesia	
<i>Calcinus anani</i> Poupin & McLaughlin, 1998 .....	French Polynesia	
<i>Calcinus gouti</i> Poupin, 1997 .....	French Polynesia	
<i>Ciliopagurus major</i> Forest, 1995 .....	French Polynesia	
<i>Ciliopagurus pacificus</i> Forest, 1995 .....	French Polynesia	
<i>Ciliopagurus plessisi</i> Forest, 1995 .....	French Polynesia	
<i>Strigopagurus poupinii</i> Forest, 1995 .....	French Polynesia	
<b>EUMUNIDAE</b>		
<i>Eumunida treguieri</i> de Saint Laurent & Poupin, 1996 .....	French Polynesia	
<b>GALATHEIDAE</b>		
<i>Galathea profunda</i> Macpherson, 2012 .....	Vanuatu	
<i>Galathea raventosae</i> Macpherson, 2012 .....	Vanuatu	
<i>Galathea sanctae</i> Macpherson, 2012 .....	Vanuatu	
<b>LITHODIDAE</b>		
<i>Lithodes megacantha</i> Macpherson, 1991 .....	French Polynesia	
<i>Paralomis aerae</i> Macpherson, 2001 .....	Fiji	
<i>Paralomis mendagnai</i> Macpherson, 2003 .....	Solomon	
<b>MUNIDIDAE</b>		
<i>Agononida aequabilis</i> Macpherson, 2006a .....	French Polynesia	
<i>Agononida alisiae</i> Macpherson, 1999 .....	Vanuatu	
<i>Agononida garciai</i> Macpherson, 2004 .....	Fiji	
<i>Agononida imitata</i> Macpherson, 2006a .....	French Polynesia	
<i>Agononida isabelensis</i> Cabezas, Macpherson & Machordom, 2009 .... Solomon		
<i>Agononida similima</i> Macpherson, 2006a .....	French Polynesia	
<i>Anoplоніда patae</i> MacPherson & Baba, 2006 .....	Tonga	
<i>Babamunida corniculata</i> Macpherson, this volume .....	French Polynesia	
<i>Babamunida hystrix</i> (Macpherson & de Saint Laurent, 1991) ... French Polynesia		
<i>Babamunida plexaura</i> (Macpherson & de Saint Laurent, 1991) ... French Polynesia		
<i>Bathymunida avatea</i> MacPherson & Baba, 2006 .....	French Polynesia	
<i>Bathymunida dissimilis</i> Baba & de Saint Laurent, 1996 .....	Wallis & Futuna	
<i>Bathymunida recta</i> Baba & de Saint Laurent, 1996 .....	Wallis & Futuna	
<i>Crozieria tucanae</i> Macpherson, 2004 .....	Fiji	
<i>Crozieria yante</i> (Macpherson, 1994) .....	French Polynesia	
<i>Heteronida clivicola</i> MacPherson & Baba, 2006 .....	French Polynesia	
<i>Munida abelloi</i> Macpherson, 1994 .....	Wallis & Futuna	
<i>Munida amathea</i> Macpherson & de Saint Laurent, 1991 .....	French Polynesia	
<i>Munida angusta</i> Macpherson, 2004 .....	Tonga	
<i>Munida antliae</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida apeleas</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida apodis</i> Macpherson, 2004 .....	Tonga	
<i>Munida aerae</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida atarapa</i> Macpherson, this volume .....	French Polynesia	
<i>Munida caeli</i> Cabezas, Macpherson & Machordom, 2009 .....	Solomon	
<i>Munida clevai</i> Macpherson, 2009 .....	Vanuatu	
<i>Munida columbae</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida congesta</i> Macpherson, 1999 .....	Vanuatu	
<i>Munida cornuta</i> Macpherson, 1994 .....	Fiji	
<i>Munida descensa</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida erugata</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida fasciata</i> Macpherson, 2006a .....	French Polynesia	
<i>Munida fornacis</i> Macpherson, 2006a .....	French Polynesia	
<b>MUNIDOPSISIDAE</b>		
<i>Galacantha quiquei</i> Macpherson, 2007 .....	Wallis & Futuna	
<i>Munidopsis austellus</i> Macpherson, 2007 .....	French Polynesia	
<i>Munidopsis bruta</i> Macpherson, 2007 .....	Solomon	

<i>Munidopsis concava</i> Macpherson, 2007	Fiji	
<i>Munidopsis demeter</i> Macpherson, 2007	Solomon	
<i>Munidopsis denudata</i> Macpherson, 2007	Solomon	
<i>Munidopsis keiji</i> Macpherson, 2007	Solomon	
<i>Munidopsis pericalla</i> Macpherson, 2007	Solomon	
<i>Munidopsis strigula</i> Macpherson, 2007	Solomon	
<i>Munidopsis zarazagai</i> Macpherson, 2007	Vanuatu	
<b>PAGURIDAE</b>		
<i>Lithopagurus boucheti</i> McLaughlin & Lemaitre, 2004	Fiji	
<i>Micropagurus bijdeleyi</i> Lemaitre, 2010	French Polynesia	
<i>Pagurojacquesia polymorpha</i> (de Saint Laurent & McLaughlin, 1999)	Vanuatu	
<i>Solitariopagurus triprobolus</i> Poupin & McLaughlin, 1996	French Polynesia	
<i>Turleania boucheti</i> McLaughlin, 2007b	Wallis&Futuna	
<b>PARAPAGURIDAE</b>		
<i>Oncopagurus oimos</i> Lemaitre, 1998	French Polynesia	
<i>Oncopagurus tuamotu</i> Lemaitre, 1994	French Polynesia	
<i>Paragiopagurus bougainvillei</i> (Lemaitre, 1994)	French Polynesia	
<i>Paragiopagurus fasciatus</i> Lemaitre & Poupin, 2003	French Polynesia	
<i>Paragiopagurus trilineatus</i> Lemaitre, this volume	Tonga	
<i>Paragiopagurus wallisi</i> (Lemaitre, 1994)	French Polynesia	
<i>Sympagurus poupini</i> Lemaitre, 1994	French Polynesia	
<b>PYLOCHELIDAE</b>		
<i>Bathycheles phenax</i> McLaughlin & Lemaitre, 2009	Fiji	
<i>Trizocheles mendanai</i> McLaughlin & Lemaitre, 2009	Solomon	
<b>BRACHYURA</b>		
<b>CALAPPIDAE</b>		
<i>Calappa sebastieni</i> Galil, 1997	French Polynesia	
<i>Mursia diwata</i> Galil & Takeda, 2004	Vanuatu	
<i>Mursia poupini</i> Galil, 2001	French Polynesia	
<b>CYCLODORIPPIDAE</b>		
<i>Ketamia limatula</i> Tavares, 1993	Vanuatu	
<i>Tymolus daviei</i> Tavares, 1997	Vanuatu	
<b>DYNOMENIDAE</b>		
<i>Metadynomene tuamotu</i> Ng & McLay, 2010	French Polynesia	
<b>DROMIIDAE</b>		
<i>Cryptodromia erioxylon</i> McLay, 2001	French Polynesia	
<i>Cryptodromia marquesas</i> McLay, 2001	French Polynesia	
<i>Sphaerodromia ducoussoi</i> McLay, 1991	French Polynesia	
<b>EPIALTIDAE</b>		
<i>Garthinia disica</i> Richer de Forges & Ng, 2009a	Solomon	
<i>Guinotinia lehouarnoi</i> Richer de Forges & Ng, 2009a	Fiji	
<i>Naxioides teatui</i> Poupin, 1995	French Polynesia	
<i>Naxioides vaitahu</i> Poupin, 1995	French Polynesia	
<i>Oxyleurodon barazeri</i> Richer de Forges & Ng, 2009b	Solomon	
<i>Oxyleurodon christiani</i> Richer de Forges & Corbari, 2012	PNG	
<i>Oxyleurodon parallelum</i> Richer de Forges & Ng, 2009b	Solomon	
<i>Oxyleurodon tavaresi</i> Richer de Forges, 1995	Wallis & Futuna	
<i>Rochinia boucheti</i> Richer de Forges & Ng, this volume	Solomon	
<b>ETHUSIDAE</b>		
<i>Ethusa abbreviata</i> Castro, 2005a	Vanuatu	
<i>Ethusa barbata</i> Castro, 2005a	Solomon	
<i>Ethusa crassipodia</i> Castro, 2005a	Solomon	
<i>Ethusina ciliacirrata</i> Castro, 2005a	Vanuatu	
<i>Ethusina coronata</i> Castro, 2005a	Solomon	
<i>Ethusina crenulata</i> Castro, 2005a	French Polynesia	
<i>Ethusina exophthalma</i> Castro, 2005a	Fiji	
<i>Ethusina huiliana</i> Castro, 2005a	French Polynesia	
<i>Ethusina microspina</i> Chen, 2000	Vanuatu	
<i>Ethusina ocellata</i> Castro, 2005a	Solomon	
<i>Ethusina stenommata</i> Castro, 2005a	Wallis & Futuna	
<i>Ethusina vanuatuensis</i> Chen, 2000	Vanuatu	
<i>Parethusa hylophora</i> Castro, 2005a	Solomon	
<i>Serpenthusa brucei</i> Naruse, Castro & Ng, 2009	Vanuatu	
<b>GERYONIDAE</b>		
<i>Chaceon australis</i> Manning, 1993	French Polynesia	
<b>GONEPLACIDAE</b>		
<i>Carcinoplax tenuidentata</i> Castro, 2007	Fiji	
<i>Carcinoplax velutina</i> Castro, 2007	French Polynesia	
<i>Guinoplax aphelis</i> Castro & Ng, 2010	Fiji	
<i>Microgoneplax cope</i> Castro, 2007	Wallis & Futuna	
<i>Thyraplax cristata</i> Castro, 2007	French Polynesia	
<i>Thyraplax digitudentata</i> Castro, 2007	French Polynesia	
<i>Thyraplax truncata</i> Castro, 2007	Fiji	
<b>HOMOLIDAE</b>		
<i>Homola poupini</i> Richer de Forges & Ng, 2007	French Polynesia	
<i>Homolochunia menezi</i> Richer de Forges & Ng, 2008b	Solomon	
<i>Homologenus broussei</i> Guinot & Richer de Forges, 1981b	French Polynesia	
<i>Homologenus wallis</i> Guinot & Richer de Forges, 1995	Wallis & Futuna	
<i>Lamoha longirostris</i> (Chen, 1986)	Wallis & Futuna	
<i>Lamoha personata</i> (Guinot & Richer de Forges, 1981b)	Vanuatu	
<i>Yaldwynopsis guinotae</i> Richer de Forges & Ng, 2007	French Polynesia	
<b>INACHIDAE</b>		
<i>Cyrtomaia polynesica</i> Richer de Forges & Ng, 2008a	French Polynesia	
<i>Platymaia rebierei</i> Guinot & Richer de Forges, 1986a	Vanuatu	
<i>Pleisticanthoides piccardorum</i> Ng & Richer de Forges, 2012	PNG	
<b>LEUCOSIIDAE</b>		
<i>Euclosiaria guinotae</i> Galil & Ng, 2010	Vanuatu	
<i>Euclosiaria vella</i> (Galil, 2007)	Solomon	
<i>Tanaoa nanus</i> Galil, 2003	Vanuatu	
<i>Tokoyo cirrata</i> Galil, 2003	Vanuatu	
<i>Urnalana granulimera</i> Galil, 2005	Fiji	
<b>MATHIELLIDAE</b>		
<i>Beuroisia manquenei</i> Guinot & Richer de Forges, 1981a	French Polynesia	
<i>Intesius crosnieri</i> Davie, 1998	French Polynesia	
<b>PALICIDAE</b>		
<i>Neopalicus simulus</i> Castro, 2010	French Polynesia	
<i>Parapalicus armatus</i> Castro, 2000	Vanuatu	
<i>Parapalicus clinodentatus</i> Castro, 2000	Vanuatu	
<i>Parapalicus inermis</i> Castro, 2000	Fiji	

<i>Pseudopalicus pictus</i> Castro, 2000 .....	Vanuatu
<b>PLAGUSIIDAE</b>	
<i>Euchirograpsus tuerkayi</i> Crosnier, 2001 .....	French Polynesia
<b>POLYBIIDAE</b>	
<i>Parathranites parahexagonus</i> Crosnier, 2002b .....	French Polynesia
<i>Parathranites tuberosus</i> Crosnier, 2002b .....	French Polynesia
<b>PORTUNIDAE</b>	
<i>Echinolatus proximus</i> Davie & Crosnier, 2006 .....	French Polynesia
<i>Laleonectes stridens</i> Crosnier & Moosa, 2002 .....	French Polynesia
<i>Thalamita difficilis</i> Crosnier, 2002a .....	French Polynesia
<b>PROGERYONIDAE</b>	
<i>Progeryon mararae</i> Guinot & Richer de Forges, 1981a .....	French Polynesia
<b>PSEUDOZIIDAE</b>	
<i>Euryozius danielae</i> Davie, 1993 .....	French Polynesia
<b>RETROPLUMIDAE</b>	
<i>Retropluma solomonensis</i> McLay, 2006 .....	Solomon
<b>TRAPEZIIDAE</b>	
<i>Hexagonalia unidentata</i> Castro, 2005b .....	Solomon
<b>TRICHOPELTARIIDAE</b>	
<i>Trichopeltarion dejouanneti</i> Tavares & Cleva, 2010 .....	Solomon
<i>Trichopeltarion parvum</i> Tavares & Cleva, 2010 .....	Solomon
<i>Trichopeltarion vanuatuensis</i> Tavares & Cleva, 2010 .....	Vanuatu
<i>Sphaeropeltarion edentatum</i> Tavares & Cleva, 2010 .....	Wallis & Futuna
<b>XANTHIDAE</b>	
<i>Alainodaeus akiaki</i> Davie, 1993 .....	French Polynesia
<i>Alainodaeus nuku</i> Davie, 1997 .....	French Polynesia
<i>Alainodaeus rimatara</i> Davie, 1993 .....	French Polynesia
<i>Banareia fatuhiva</i> Davie, 1993 .....	French Polynesia
<i>Epistocavea mururoa</i> Davie, 1993 .....	French Polynesia
<i>Lipkemera rufomaculata</i> (Davie, 1993) .....	French Polynesia
<i>Medaeus grandis</i> Davie, 1993 .....	French Polynesia
<i>Meractaea tafai</i> Davie, 1993 .....	French Polynesia
<i>Paramedaeus megagomphios</i> Davie, 1997 .....	Wallis & Futuna
<i>Paraxanthodes polynesiensis</i> Davie, 1993 .....	French Polynesia
<i>Rata tuamotense</i> Davie, 1993 .....	French Polynesia

**APPENDIX 4**

References to taxonomical publications containing original descriptions (listed in Appendix 2 & 3) of decapod crustaceans collected by the *Tropical Deep-Sea Benthos* programme and satellite programmes.

- AHYONG S.T. 2010 — The marine fauna of New Zealand: King crabs of New Zealand, Australia and the Ross Sea (Crustacea: Decapoda: Lithodidae) NIWA *Biodiversity Memoir* 123, Wellington, 196 p.
- AHYONG S.T. & GALIL B.S. 2006 — Polychelidae from the southern and western Pacific (Decapoda, Polychelida). *Zoosystema* 28 (3): 757-767.
- ASAKURA A. 2005 — A review of the genus *Micropagurus* (Crustacea Decapoda Anomura Paguridae). *Zootaxa* 1090: 1-34.
- BABA K. 1991a — Crustacea Decapoda: *Chirostylus* Ortmann, 1892, and *Gastroptychus* Caullery, 1896 (Chirostyliidae) from New Caledonia, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 463-477.
- BABA K. 1991b — Crustacea Decapoda: *Alainius* gen. nov., *Leiogalathea* Baba, 1969, and *Phylladiorhynchus* Baba, 1969 (Galatheidae) from New Caledonia, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 479-491.
- BABA K. 2004 — *Uroptychodes*, new genus of Chirostyliidae (Crustacea: Decapoda: Anomura), with description of three new species. *Scientia Marina* 68: 97-116.
- BABA K. & SAINT LAURENT DE M. 1996 — Crustacea Decapoda: Revision of the genus *Bathymunida* Balss, 1914, and description of six new related genera (Galatheidae), in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 433-502.
- BOYKO C.B. & HARVEY A.W. 1999 — Crustacea Decapoda: Albuneidae and Hippidae of the tropical Indo-West Pacific region, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 20. *Mémoires du Muséum national d'Histoire naturelle* 180: 379-406.
- BOYKO C.B. 2000 — The Hippoidea (Decapoda, Anomura) of the Marquesas Islands, with description of a new species of Albunea. *Zoosystema* 22: 107-116.
- BRUCE A.J. 1989 — *Periclimenes pouponi* sp. nov., a new anemone-associated shrimp from deep-water traps (Crustacea, Decapoda, Palaemonidae). *Bulletin du Muséum national d'Histoire naturelle, Section A, Zoologie, Biologie et Ecologie Animales, Paris*, 4e série 11: 851-863.
- BRUCE A.J. 1990a — Crustacea Decapoda: Deep-sea Palaemonoid shrimps from New Caledonian waters, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 149-215.
- BRUCE A.J. 1990b — Crustacea Decapoda: *Gelastreutes crosnieri* gen. nov., sp. nov. (Hippolytidae) from New Caledonia, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 137-148.
- BRUCE A.J. 1991 — Crustacea Decapoda: Further deep-sea Palaemonoid shrimps from New Caledonian waters, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 299-411.
- BRUCE A.J. 2006 — An unusual new *Periclimenes* (Crustacea, Decapoda, Palaemonidae) from New Caledonia. *Zoosystema* 28 (3): 703-712.
- BURUKOVSKY R.N. 2000 — Taxonomy of shrimps from the genus *Nematocarcinus* (Crustacea, Decapoda, Nematocarcinidae). Description of species from Tenuirostris group (In russian). *Zoologicheskii Zhurnal* 79 (8): 898-906.
- BURUKOVSKY R.N. 2003 — *Shrimps of the family Nematocarcinidae*. Kaliningrad Izdatelistovo KGGU. Kaliningrad State Technical University, Kaliningrad, 250 p.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2009 — Morphological and molecular description of new species of squat lobster (Crustacea: Decapoda: Galatheidae) from the Solomon and Fiji Islands (South-West Pacific). *Zoological Journal of the Linnean Society* 156: 465-493.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2010 — Taxonomic revision of the genus *Paramunida* Baba, 1988 (Crustacea: Decapoda: Galatheidae): a morphological and molecular approach. *Zootaxa* 2712: 1-60.
- CABEZAS P., MACPHERSON E. & MACHORDOM A. 2011 — *Allogalathea* (Decapoda: Galatheidae): a monospecific genus of squat lobsters? *Zoological Journal of the Linnean Society* 156: 465-493.
- CASTRO P. 2000 — Crustacea Decapoda: A révision of the Indo-west Pacific species of palicid crabs (Brachyura Palicidae), in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 437-610.
- CASTRO P. 2005a — Crabs of the subfamily *Ethusinae* Guinot, 1977 (Crustacea, Decapoda, Brachyura, Dorippidae) of the Indo-West Pacific region. *Zoosystema* 27 (3): 499-600.
- CASTRO P. 2005b — A new species of *Hexagonalia* (Crustacea: Brachyura: Trapeziidae) from the Solomon Islands. *Proceedings of the Biological Society of Washington* 118: 539-542.
- CASTRO P. 2007 — A reappraisal of the family *Goneplacidae* MacLeay, 1838 (Crustacea, Decapoda, Brachyura) and revision of the subfamily *Goneplacinae*, with the description of ten new genera and eighteen new species. *Zoosystema* 29 (4): 609-773.
- CASTRO P. 2009 — Two new species of *Carcinoplax* H. Milne Edwards, 1852, and *Pycnoplax* Castro, 2007, from the western Pacific, and a description of the female of *Thyraplax truncata* Castro, 2007 (Crustacea, Decapoda, Brachyura, Goneplacidae). *Zoosystema* 31 (4): 949-957.
- CASTRO P. 2010 — A new species and new records of palicoid crabs (Crustacea, Decapoda, Brachyura, Palicoidea, Palicidae, Crossotonotidae) from the Indo-West Pacific region. *Zoosystema* 32 (1): 73-86.
- CASTRO P. & NG P.K.L. 2010 — A new genus and species of goneplacid crab (Decapoda, Brachyura, Goneplacidae) from the western Pacific, in CASTRO P., DAVIE P.J.F., NG P.K.L. & RICHER DE FORGES B. (eds), *Studies on Brachyura: a homage to Danièle Guinot*. Brill, Leiden; Boston: 51-60 (Crustaceana Monographs; 11).

- CHAN T.-Y. 2004 — The "Plesionika rostricrescentis" (Bate, 1888)" and "P. lophotes Chace, 1985" species groups of *Plesionika* Bate, 1888, with descriptions of five new species (Crustacea: Decapoda: Pandalidae), in MARSHALL B. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos, vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 293-318.
- CHANT T.-Y. 2012 — A new genus of deep-sea solenocerid shrimp (Decapoda: Penaeoidea) from Papua New Guinea. *Journal of Crustacean Biology* 32 (3): 489-495.
- CHAN T.-Y. & CROSNIER A. 1991 — Crustacea Decapoda: Studies of the *Plesionika narval* (Fabricius, 1787) group (Pandalidae), with descriptions of six new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 413-461.
- CHAN T.-Y. & CROSNIER A. 1997 — Crustacea Decapoda: Deep sea shrimps of the genus *Plesionika* Bate, 1888 (Pandalidae) from French Polynesia, with descriptions of five new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 187-234.
- CHAN T.-Y. & YU H.-P. 1991 — *Eugonatonotus chacei* sp. nov., second species of the genus (Crustacea, Decapoda, Eugonatonotidae). *Bulletin du Muséum national d'Histoire naturelle* (4) 13 (A) 1-2): 143-152.
- CHEN H.L. 1986 — Decapod Crustacea: Dorippidae, in FOREST J. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 2. *Mémoires du Muséum national d'Histoire naturelle* 133: 179-204.
- CHEN H.L. 1993 — Crustacea Decapoda: Dorippidae of New Caledonia, Indonesia and the Philippines, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 315-345.
- CHEN H.L. 2000 — Crustacea Decapoda: New species and new records of Ethusinae (Dorippidae) from Vanuatu, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 425-435.
- CHIA D.G.B. & NG P.K.L. 2000 — A revision of *Eumedonous* H. Milne Edwards, 1834 and *Gonatonotus* White, 1847 (Crustacea: Decapoda: Brachyura: Eumelonidae), two genera of crabs symbiotic with sea urchins. *Journal of Natural History* 34: 15-56.
- CLEVA R. 1990 — Crustacea Decapoda: Les genres et les espèces indo-ouest pacifiques de Stylopactylidae, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 71-136.
- CLEVA R. 1994 — Some Australian Stylopactylidae (Crustacea: Decapoda) with descriptions of two new species. *The Beagle, Records of the Museums and Art Galleries of the Northern Territory* 11: 53-64.
- CLEVA R. & CROSNIER A. 2006 — *Heterocarpus tenuidentatus*, a new species of shrimp from the Solomon Islands (Crustacea, Decapoda, Caridea, Pandalidae). *Zootaxa* 1200: 61-68.
- CROSNIER A. 1986a — Crevettes de la famille des Pandalidae récoltées durant ces dernières années en Polynésie française. Description de *Plesionika chacei* et *P. carsini* spp. nov. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 8: 361-377.
- CROSNIER A. 1986b — *Plesionika fenneri*, nouveau nom pour *Plesionika chacei* Crosnier, 1986. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 8: 691.
- CROSNIER A. 1988a — Contribution à l'étude des genres *Haliporus* Bate, 1881 et *Gordonella* Tirmizi, 1960 (Crustacea Decapoda Penaeoidea). Description de deux espèces nouvelles. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 10: 563-601.
- CROSNIER A. 1988b — Sur les *Heterocarpus* (Crustacea, Decapoda, Pandalidae) du sud-ouest de l'océan Indien. Remarques sur d'autres espèces ouest-pacifiques du genre et description de quatre taxa nouveaux. *Bulletin du Muséum national d'Histoire naturelle* (4) (A) 10: 57-103.
- CROSNIER A. 1994 — *Sphaerodromia lamellata*, espèce nouvelle de Nouvelle-Calédonie (Decapoda, Brachyura, Dromiidae). *Crustaceana* 67 (3): 341-347.
- CROSNIER A. 1997a — Crustacea Decapoda: *Pseudopandalus curvirostris*, genre et espèce nouveaux (Pandalidae) de Nouvelle-Calédonie, in CROSNIER A. (ed.), *Résultats des campagnes MUSORSTOM*, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 169-176.
- CROSNIER A. 1997b — Une nouvelle espèce de *Mursia* de Nouvelle-Calédonie (Crustacea, Decapoda, Brachyura, Calappidae). *Zoosystema* 19: 151-158.
- CROSNIER A. 1999 — Un *Heterocarpus* nouveau (Crustacea, Decapoda, Pandalidae) du Pacifique Sud-Ouest. *Zoosystema* 21: 345-351.
- CROSNIER A. 2000 — A new genus and species of hippolytid shrimp (Decapoda, Caridea) from Wallis Island, south-west Pacific. *Journal of Crustacean Biology* 20 (spec. n° 2): 109-115.
- CROSNIER A. 2001 — Grapsidae (Crustacea, Decapoda, Brachyura) d'eau profonde du Pacifique sud-ouest. *Zoosystema* 23 (4): 783-796.
- CROSNIER A. 2002a — Portunidae (Crustacea, Decapoda, Brachyura) de Polynésie française, principalement des îles Marquises. *Zoosystema* 24: 401-449.
- CROSNIER A. 2002b — Révision du genre *Parathranites* Miers, 1886 (Crustacea, Brachyura, Portunidae). *Zoosystema* 24 (4): 799-825.
- CROSNIER A. 2003 — *Sicyonia* (Crustacea, Decapoda, Penaeoidea, Sicyoniidae) de l'indo-ouest Pacifique. *Zoosystema* 25: 197-350.
- CROSNIER A. 2005 — Deux *Parapenaeus* nouveaux (Crustacea, Decapoda, Penaeidae) du sud-ouest Pacifique. *Zoosystema* 27: 257-266.
- CROSNIER A. 2006 — *Penaeopsis* Bate 1881 (Crustacea, Decapoda, Penaeidae) récoltées dans le Pacifique sud-ouest par les campagnes françaises depuis 1976. Description d'une espèce nouvelle. *Zoosystema* 28: 331-340.
- CROSNIER A. & DALL W. 2004 — Redescription of *Hymenopenaeus obliquirostris* (Crustacea, Decapoda, Penaeoidea, Solenoceridae) and description of two new species of *Hymenopenaeus* from the Indo-West Pacific. *Zootaxa* 600: 1-26.
- CROSNIER A. & MOOSA M.K. 2002 — Trois Portunidae (Crustacea, Decapoda, Brachyura) nouveaux de Polynésie française. *Zoosystema* 24: 385-399.
- CROSNIER A. & NG P.K.L. 2004 — Remarques sur le genre *Intesius* (Crustacea, Decapoda, Brachyura, Gonoplacidae) et description de deux espèces nouvelles. *Zoosystema* 26 (2): 263-277.

- DAVIE P.J.F. 1991 — Crustacea Decapoda: the genus *Platepistoma* Rathbun, 1906 (Cancridae) with the description of three new species, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 493-514.
- DAVIE P.J.F. 1993 — Deepwater xanthid crabs from French Polynesia (Crustacea, Decapoda, Xanthoidea). *Bulletin du Muséum national d'Histoire naturelle* (4) 14 (A2): 501-561.
- DAVIE P.J.F. 1997 — Crustacea Decapoda: Deep water Xanthoidea from the South-Western Pacific and the Western Indian Ocean, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 337-387.
- DAVIE P.J.F. 1998 — A new species of *Intesius* (Crustacea, Decapoda, Gonoplacidae) from the deep water of French Polynesia. *Zoosystema* 20 (2): 221-227.
- DAVIE P.J.F. & CROSNIER A. 2006 — *Echinolatus* n. gen. (Crustacea, Decapoda, Portunidae) with description of two new species from the South-West Pacific, *in RICHER DE FORGES B. & JUSTINE J. L. (eds)*, Tropical Deep-Sea Benthos, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 393-410.
- DE GRAVE S. 2004 — A new species of *Batella* (Crustacea, Decapoda, Alpheidae) from New Caledonia. *Zoosystema* 26: 503-509.
- DE GRAVE S., PENTCHEFF N.D., AHYONG S., CHANT T.-Y., CRANDALL K.A., DWORSCHAK P., FELDER D.L., FELDMANN R.M., FRANSEN C.H.J.M., GOULDING L.Y.D., LEMAITRE R., LOW M.L., MARTIN J.W., NG P.K.L., SCHWEITZER C.E., TAN S.H., TSHUDY D. & WETZER R. 2009 — A classification of Recent and fossil genera of decapod crustaceans. *Raffles Bulletin of Zoology* suppl 21: 1-109.
- FOREST J. 1987 — Les Pylochelidae ou "Pagures symétriques" (Crustacea Coenobitoidea), *in Résultats des campagnes MUSORSTOM, vol. 3. Mémoires du Muséum national d'Histoire naturelle* 137: 1-254.
- FOREST J. 1995 — Crustacea Decapoda Anomura: Révision du genre *Trizopagurus* Forest, 1952 (Diogenidae), avec l'établissement de deux genres nouveaux, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 13. *Mémoires du Muséum national d'Histoire naturelle* 163: 9-149.
- FOREST J. 1997 — Présence du genre *Xylopagurus* A. Milne-Edwards, 1880 (Crustacea, Decapoda, Paguridae) dans l'Indo-Ouest Pacifique, avec la description de deux espèces nouvelles. *Zoosystema* 19: 421-435.
- FOREST J. & MC LAUGHLIN P.A. 2000 — Superfamily Coenobitoidea, *in FOREST J., SAINT LAURENT M. DE, MC LAUGHLIN P.A. & LEMAITRE R. (eds)*, The marine fauna of New Zealand: Paguridea (Decapoda: Anomura) exclusive of the Lithodidae. *NIWA Biodiversity Memoir* 114: 31-103.
- FRANSEN C.H. 1997 — Crustacea Decapoda: *Chelonila macrochela*, a new genus and new species of pandalid shrimp (Caridea) from New Caledonian waters, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 177-185.
- GALIL B.S. 1993 — Crustacea Decapoda: A revision of the genus *Mursia* Desmarest, 1823 (Calappidae), *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 347-379.
- GALIL B.S. 1997 — Crustacea Decapoda: A revision of the Indo-Pacific species of genus *Calappa* Weber, 1795 (Calappidae), *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 271-335.
- GALIL B.S. 2000 — Crustacea Decapoda: A revision of the family Polychelidae, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 285-387.
- GALIL B.S. 2001a — A new genus and species of leucosiid crab (Crustacea, Decapoda, Brachyura) from the Indo-Pacific Ocean. *Zoosystema* 23: 65-75.
- GALIL B.S. 2001b — The Calappidae of the Marquesas Islands with a description of a new species of *Mursia* (Crustacea, Decapoda, Brachyura). *Zoosystema* 23: 499-505.
- GALIL B.S. 2003 — Four new genera of leucosiid crabs (Crustacea: Brachyura: Leucosiidae) for three new species and nine species previously in the genus *Randallia* Stimpson, 1857, with a redescription of the type species, *R. ornata* (Randall, 1939). *Proceedings of the Biological Society of Washington* 116 (2): 395-422.
- GALIL B.S. 2005 — Contributions to the knowledge of Leucosiidae III. *Urnalana* gen. nov. (Crustacea: Brachyura). *Zoologische Mededelingen* 79: 9-40.
- GALIL B.S. 2007 — The deep-water Calappidae, Matutidae and Leucosiidae of the Solomon Islands with a description of a new species of *Eucloria* Galil 2003 (Crustacea, Decapoda, Brachyura). *Zoosystema* 29: 555-563.
- GALIL B.S. & CLARK P.F. 1990 — Crustacea Decapoda: Notes on some species of Trapeziidae from New Caledonia including the descriptions of two new ones, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 369-388.
- GALIL B.S. & NG P.K.L. 2010 — On a collection of calappoid and leucosioid crabs (Decapoda, Brachyura) from Vanuatu, with description of a new species of Leucosiidae, *in CASTRO P., DAVIE P.J.F., NG P.K.L. & RICHER DE FORGES B. (eds)*, *Studies on Brachyura: a homage to Danièle Guinot*. Brill, Leiden; Boston: 139-152 (Crustaceana Monographs; 11).
- GALIL B.S. & TAKEDA M. 2004 — On a Collection of *Mursia* (Crustacea, Decapoda, Brachyura, Calappidae) from Balicasag Island, Philippines. *Bulletin of the National Science Museum Series A* 30 : 23-35.
- GUINOT D. 1993 — Données nouvelles sur les Crabes primitifs (Crustacea Decapoda Brachyura Podotremata). *Comptes rendus de l'Académie des Sciences* (III), 316 (10): 1225-1232.
- GUINOT D. 1995 — Crustacea Decapoda Brachyura: Révision de la famille des Homolodromiidae Alcock, 1900, *in CROSNIER A.* (ed.), Résultats des Campagnes MUSORSTOM, vol. 13. *Mémoires du Muséum national d'Histoire naturelle* 163: 155-282.
- GUINOT D. & RICHER DE FORGES B. 1981a — Crabes de profondeur, nouveaux ou rares, de l'Indo-Pacifique (Crustacea, Decapoda, Brachyura) (Première partie). *Bulletin du Muséum national d'Histoire naturelle* (4) 2 (A) (4): 1113-1153.
- GUINOT D. & RICHER DE FORGES B. 1981b — Homolidae, rares ou nouveaux, de l'Indo-Pacifique (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle* (A) 8 (2): 523-581.

- GUINOT D. & RICHER DE FORGES B. 1982 — Nouvelles récoltes des genres *Cyrtomaia* Miers et *Pleistacantha* Miers (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle* (4) 3 (A) (4): 1087-1125.
- GUINOT D. & RICHER DE FORGES B. 1986a — Crustacés Décapodes: Majidae (genres *Platymaia*, *Cyrtomaia*, *Pleistacantha*, *Sphenocarcinus* et *Naxioides*), in FOREST J. (ed.), Résultats des Campagnes MUSORSTOM, vol. 2. *Mémoires du Muséum National d'Histoire Naturelle* 133: 83-179.
- GUINOT D. & RICHER DE FORGES B. 1986b — Découverte d'une nouvelle espèce de *Sphenocarcinus* en Nouvelle-Calédonie, *S. mammatus* sp. nov. (Crustacea, Decapoda, Brachyura). *Indo-Malayan Zoology* 3: 27-37.
- GUINOT D. & RICHER DE FORGES B. 1995 — Crustacea Decapoda Brachyura: Révision de la famille des Homolidae de Haan, 1839, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 13. *Mémoires du Muséum national d'Histoire naturelle* 163: 283-517.
- HAYASHI K.I. 1999 — Crustacea Decapoda: Revision of *Pasiphaea sivado* (Risso, 1816) and related species, with descriptions of one new genus and five new species (Pasiphaeidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 20. *Mémoires du Muséum national d'Histoire naturelle* 180: 267-302.
- HAYASHI K.I. 2004 — Revision of the *Pasiphaea cristata* Bate, 1888 species group of *Pasiphaea* Savigny, 1816, with descriptions of four new species, and referral of *P. australis* Hanamura, 1989 to *Alainopasiphaea* Hayashi, 1999 (Crustacea, Decapoda, Pasiphaeidae), in MARSHALL B.A. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos, vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 319-373.
- HAYASHI K.I. 2006 — Revision of the *Pasiphaea alcocki* species group (Crustacea, Decapoda, Pasiphaeidae), in RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 193-241.
- HOLTHUIS L.B. 2002 — The Indo-Pacific scyllarid lobsters (Crustacea: Decapoda: Scyllaridae). *Zoosystema* 24 (3): 499-683.
- KOMAI T. 2004a — A new genus and new species of Crangonidae (Crustacea, Decapoda, Caridea) from the southwestern Pacific. *Zoosystema* 26: 73-85.
- KOMAI T. 2004b — A review of the Indo-West Pacific species of the genus *Glyphocrangon* A. Milne-Edwards, 1881 (excluding the *G. caeca* species group) (Crustacea, Decapoda, Caridea, Glyphocrangonidae), in MARSHALL B.A. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos, vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 375-610.
- KOMAI T. 2006b — Revision of the *Glyphocrangon caeca* species group (Crustacea, Decapoda, Glyphocrangonidae), in RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 243-264.
- KOMAI T. 2007 — A new species of *Glyphocrangon* (Crustacea, Decapoda, Caridea, Glyphocrangonidae) from the Austral Islands, French Polynesia. *Zoosystema* 29: 565-573.
- KOMAI T. 2008 — A world-wide revision of species of the deep-water crangonid genus *Parapontophilus* Christoffersen, 1988 (Crustacea, Decapoda, Caridea), with descriptions of ten new species. *Zoosystema* 30: 261-332.
- KOMAI T. 2011 — Further records of deep-sea shrimps of the genus *Glyphocrangon* (Crustacea: Decapoda: Caridea: Glyphocrangonidae) from the Southwestern Pacific, with descriptions of two new species. *Species Diversity* 16: 113-135.
- KOMAI T. 2012 — A review of the western Pacific species of the crangonid genus *Metacrangon* Zarenkov, 1965 (Decapoda: Caridea), with descriptions of seven new species. *Zootaxa* 3468: 1-77.
- KOMAI T. & SAITO T. 2006 — A new genus and two new species of Spongicolidae (Crustacea: Decapoda: Stenopodidea) from the South-West-Pacific, in RICHER DE FORGES B. & JUSTINE J.-L. (eds), Tropical Deep-Sea Benthos, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 265-284.
- LEMAITRE R. 1994 — Crustacea Decapoda: Deep-water hermit crabs (Parapaguridae) from French Polynesia with descriptions of four new species, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 12. *Mémoires du Muséum national d'Histoire naturelle* 161: 375-419.
- LEMAITRE R. 1997 — Crustacea Decapoda: Parapaguridae from the KARUBAR cruise in Indonesia, with descriptions of two new species, in CROSNIER A. & BOUCHET P. (eds), Résultats des Campagnes MUSORSTOM, vol. 16. *Mémoires du Muséum national d'Histoire naturelle* 172: 573-596.
- LEMAITRE R. 1989 — Revision of the genus *Parapagurus* (Anomura: Paguroidea: Parapaguridae), including redescriptions of the western Atlantic species. *Zoologische Verhandelingen* 253: 1-106.
- LEMAITRE R. 1998 — A new species of hermit crab of the family Parapaguridae (Decapoda, Anomura) from French Polynesia. *Zoosystema* 20 (1): 101-107.
- LEMAITRE R. 1999 — Crustacea Decapoda: A review of the species of the genus *Parapagurus* Smith, 1879 (Parapaguridae) from the Pacific and Indian Oceans, in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 20. *Mémoires du Muséum national d'Histoire naturelle* 180: 303-378.
- LEMAITRE R. 2004a — A review of *Strobopagurus* Lemaitre, 1989 (Crustacea: Decapoda: Paguroidea: Parapaguridae), with description of a new species. *Scientia Marina* 68 (3): 355-372.
- LEMAITRE R. 2004b — Crustacea Decapoda: A worldwide review of the hermit crab species of the genus *Sympagurus* Smith, 1883, in MARSHALL B. & RICHER DE FORGES B. (eds), Tropical Deep-Sea Benthos, vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 85-149.
- LEMAITRE R. 2004c — Discovery of the first hermit crab (Crustacea: Decapoda: Parapaguridae) associated with hydrothermal vents. *Cahiers de Biologie marine* 45: 325-334.
- LEMAITRE R. 2006 — Two new species of Parapaguridae (Crustacea, Anomura, Paguroidea) with subconical corneas, and new data on biology of some rare species. *Zoosystema* 28 (2): 517-532.
- LEMAITRE R. 2010 — A new species of *Micropagurus* McLaughlin, 1986 (Decapoda, Anomura, Paguridae) from French Polynesia. *Crustaceana Monographs* 14: 379-388.

- LEMAITRE R. & POUPIN J. 2003 — A strikingly coloured new species of *Paragiopagurus* Lemaître, 1996 (Crustacea: Decapoda: Anomura: Parapaguridae) from French Polynesia. *Zootaxa* 386: 1-11.
- LEMAITRE R. & MCLAUGHLIN P.A. 1995 — *Alainopagurus crosnieri*, gen. et sp. nov. (Decapoda: Anomura: Paguridae) from the Western Pacific. *Bulletin du Muséum national d'Histoire naturelle* (4) 17 (A) (3-4): 273-282.
- LI X & BRUCE A.J. 2006 — Further Indo-West Pacific palaemonoid shrimps (Crustacea: Decapoda: Palaemonoidea), principally from the New Caledonian region. *Journal of Natural History* 40: 611-738.
- LI X. 2008 — Report on some species of Palaemonidae (Crustacea, Decapoda) from French Polynesia. *Zoosystema* 30: 203-252.
- LIN F.-J. 2006 — Two new Axioids (Decapoda : Thalassinidea) from New Caledonia. *Journal of Crustacean Biology* 26: 234-241.
- MACPHERSON E. 1990 — Crustacea Decapoda: On some species of Lithodidae from the Western Pacific, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 217-226.
- MACPHERSON E. 1991 — A new species of the genus *Lithodes* (Crustacea, Decapoda, Lithodidae) from the French Polynesia. *Bulletin du Muséum national d'Histoire naturelle* (4) 13 (A) (1-2): 153-158.
- MACPHERSON E. 1993 — Crustacea Decapoda: species of the genus *Paramunida* Baba, 1988 (Galatheidae) from the Philippines, Indonesia and New Caledonia, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 443-473.
- MACPHERSON E. 1994 — Crustacea Decapoda: Studies on the genus *Munida* Leach, 1820 (Galatheidae) in New Caledonia and adjacent waters with descriptions of 56 new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 12. *Mémoires du Muséum national d'Histoire naturelle* 161: 421-569.
- MACPHERSON E. 1996a — Crustacea Decapoda: new records of species of the genera *Munida* Leach, 1820 and *Paramunida* Baba, 1988 (Galatheidae) from New Caledonia, with the descriptions of three new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 423-431.
- MACPHERSON E. 1996b — Crustacea Decapoda: species of the genera *Munida* Leach, 1820 and *Paramunida* Baba, 1988 (Galatheidae) from the seas around the Wallis and Futuna Islands, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 387-421.
- MACPHERSON E. 1998 — A new genus of Galatheidae (Crustacea, Anomura) from the western Pacific Ocean. *Zoosystema* 20: 351-355.
- MACPHERSON E. 1999 — Crustacea Decapoda: Species of the genera *Agononida* Baba & de Saint Laurent, 1996 and *Munida* Leach, 1820 (Galatheidae) collected during the MUSORSTOM 8 cruise in Vanuatu, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 20. *Mémoires du Muséum national d'Histoire naturelle* 180: 407-426.
- MACPHERSON E. 2000 — Crustacea Decapoda: species of the genera *Crozierita* Macpherson, 1998, *Munida* Leach, 1820, and *Paramunida* Baba, 1988 (Galatheidae) collected during the MUSORSTOM 9 cruise to the Marquesas Islands, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 415-423.
- MACPHERSON E. 2001 — New species and new records of Lithodid crabs (Crustacea, Decapoda, Lithodidae) from the Southwestern and Central Pacific Ocean. *Zoosystema* 23: 797-805.
- MACPHERSON E. 2003 — Some lithodid crabs (Crustacea, Decapoda, Lithodidae) from the Salomon Islands (SW Pacific Ocean) with the description of a new species. *Scientia Marina* 67: 413-418.
- MACPHERSON E. 2004 — Species of the genus *Munida* Leach, 1820 and related genera from Fiji and Tonga (Crustacea: Decapoda: Galatheidae), in MARSHALL B.A. & RICHER DE FORGES B. (eds), *Tropical Deep-Sea Benthos*, vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 231-292.
- MACPHERSON E. 2006a — Galatheidae (Crustacea, Decapoda) from the Austral Islands, Central Pacific, in RICHER DE FORGES B. & JUSTINE J.-L. (eds), *Tropical Deep-Sea Benthos*, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 285-333.
- MACPHERSON E. 2006b — New species and new occurrence of Galatheoidea (Crustacea, Decapoda) from New Caledonia. *Zoosystema* 28: 669-681.
- MACPHERSON E. 2007 — Species of the genus *Munidopsis* Whiteaves, 1784 from the Indian and Pacific Oceans and reestablishment of the genus *Galacantha* A. Milne-Edwards, 1880 (Crustacea, Decapoda, Galatheidae). *Zootaxa* 1417: 1-135.
- MACPHERSON E. 2009 — New species of squat lobsters of the genera *Munida* and *Raymunida* (Decapoda, Galatheidae) from Vanuatu and New Caledonia. *Zoosystema* 31: 431-451.
- MACPHERSON E. 2012 — New deep-sea squat lobsters of the genus *Galathea* Fabricius, 1793 (Decapoda, Galatheidae) from Vanuatu and New Caledonia. *Zoosystema* 34: 409-427.
- MACPHERSON E. & BABA K. 2006 — New species and records of small galatheids (Crustacea, Decapoda, Galatheidae) from the southwest and central Pacific Ocean. *Zoosystema* 28: 443-456.
- MACPHERSON E. & BABA K. 2009 — New species of squat lobsters of the genera *Agononida* and *Paramunida* (Crustacea: Decapoda: Anomura: Galatheidae) from the western Pacific. *Zootaxa* 2024: 56-68.
- MACPHERSON E. & MACHORDOM A. 2000 — *Raymunida*, new genus (Decapoda: Anomura: Galatheidae) from the Indian and Pacific Oceans. *Journal of Crustacean Biology* 20: 253-258.
- MACPHERSON E. & MACHORDOM A. 2001 — Phylogenetic relationships of species of *Raymunida* (Decapoda: Galatheidae) based on morphology and mitochondrial cytochrome oxidase sequences, with the recognition of four new species. *Journal of Crustacean Biology* 21: 696-714.

- MACPHERSON E. & MACHORDOM A. 2005 — Use of morphological and molecular data to identify three new sibling species of the genus *Munida* Leach, 1820 (Crustacea, Decapoda, Galatheidae) from New Caledonia. *Journal of Natural History* 39: 819-834.
- MACPHERSON E. & SAINT LAURENT DE M. 1991 — Galatheid crustaceans of the genus *Munida* Leach, 1818, from French Polynesia. *Bulletin du Muséum national d'Histoire naturelle* (4) 13 (A) (3-4): 373-422.
- MANNING R. B. 1991 — Crustacea Brachyura: *Cecidocarcinus zibrowii*, a new deep-waler gall crab (Cryptochiridae) from New Caledonia, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 9. *Mémoires du Muséum national d'Histoire naturelle* 152: 515-520.
- MANNING R. B. 1993 — A new deep-sea crab, genus *Chaceon*, from the Austral Islands, southwestern Pacific Ocean (Decapoda: Geryonidae). *Crustacean Research* 22: 7-10.
- MANNING R. B. & HOLTHIUS L. B. 1989 — Two new genera and nine new species of geryonid crabs (Crustacea, Decapoda, Geryonidae). *Proceedings of the Biological Society of Washington* 102 (1): 50-77.
- MCLAUGHLIN P. A. 2000 — Crustacea: Decapoda: *Porcellanopagurus* Filhol and *Solitariopagurus* Türkay (Paguridae), from the New Caledonia area, Vanuatu, and the Marquesas: new records, new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 21. *Mémoires du Muséum national d'Histoire naturelle* 184: 389-414.
- MCLAUGHLIN P. A. 2003 — Reassignment and redescription of "Eupagurus" *microps* Balss, 1911 (Crustacea, Decapoda, Paguridae), with notes on *Bathypaguropsis kuroshioensis* (Miyake, 1978). *Zoosystema* 25: 635-642.
- MCLAUGHLIN P. A. 2004a — A reappraisal of the hermit crab genera *Catapagurus* A. Milne-Edwards and *Hemipagurus* Smith (Crustacea: Decapoda: Anomura: Paguridae), with the description of a new species. *Zootaxa* 433: 1-16.
- MCLAUGHLIN P. A. 2004b — Crustacea Decapoda: A review of the hermit crab genus *Nematopagurus* A. Milne-Edwards & Bouvier, 1892 and the descriptions of five new species (Crustacea: Decapoda: Paguridae), in MARSHALL B. & RICHER DE FORGES B. (eds), *Tropical Deep-Sea Benthos*, vol. 23. *Mémoires du Muséum national d'Histoire naturelle* 191: 151-229.
- MCLAUGHLIN P. A. 2006 — Two new hermit crabs from New Caledonia and environs, in RICHER DE FORGES B. & JUSTINE J.-L. (eds), *Tropical Deep-Sea Benthos*, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 335-347.
- MCLAUGHLIN P. A. 2007a — A new species of *Pteropagurus* McLaughlin & Rahayu, 2006 and a new genus and species of scaphopod dweller (Crustacea, Decapoda, Anomura: Paguroidea, Paguridae). *Zoosystema* 29 (3): 503-513.
- MCLAUGHLIN P. A. 2007b — New records and a new species in the genus *Turleania* McLaughlin, 1997. (Crustacea, Decapoda, Anomura, Paguridae) from MUSORSTOM cruises, with a key to species. *Zoosystema* 29 (3): 583-593.
- MCLAUGHLIN P. A. & FOREST J. 1997 — Crustacea Decapoda: *Diacanthurus* gen. nov., a new genus of hermit crabs (Paguridae) with both Recent and fossil representation, and the descriptions of two new species, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 236-259.
- MCLAUGHLIN P. A. & LEMAITRE R. 2004 — The discovery of two new species of *Lithopagurus* Provenzano, 1968 (Crustacea, Decapoda, Anomura, Paguroidea, Paguridae) and the first records of the genus in the western Pacific. *Zoosystema* 26 (3): 483-494.
- MCLAUGHLIN P. A. & LEMAITRE R. 2008 — Larvae of two species of *Trizocheles* (Decapoda: Anomura: Paguroidea: Pylochelidae: Trizochelinae), description of the adult of one, and preliminary implications of development on pylochelid phylogeny. *Zootaxa* 1911: 52-68.
- MCLAUGHLIN P. A. & LEMAITRE R. 2009 — A new classification of the Pylochelidae (Decapoda: Anomura: Paguroidea) and descriptions of new taxa. *Raffles Bulletin of Zoology*. Supp. 20: 159-231.
- MCLAUGHLIN P. A. & RAHAYU D. L. 2006 — A new genus with two new species of hermit crabs (Crustacea, Decapoda, Paguroidea, Paguridae) from an unique habitat. *Zootaxa* 1116: 55-68.
- MCLAUGHLIN P. A. & RAHAYU D. L. 2008 — A new genus and species of hermit crab of the family Paguridae (Crustacea: Anomura: Paguroidea) from the Vanuatu Archipelago. *Proceedings of the Biological Society of Washington* 121: 365-373.
- MCLAY C. L. 1991 — A small collection of deep water sponge crabs (Brachyura, Dromiidae) from French Polynesia, including a new species of *Sphaerodromia* Alcock, 1899. *Bulletin du Muséum national d'Histoire naturelle* (4) 13 (A) (3-4): 457-481.
- MCLAY C. L. 1993 — Crustacea Decapoda : The Sponge Crabs (Dromiidae) of New Caledonia and the Philippines with a review of the genera, in CROSNIER A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 111-251.
- MCLAY C. L. 2001 — The Dromiidae of French Polynesia and a new collection of crabs (Crustacea, Decapoda, Brachyura) from the Marquesas Islands. *Zoosystema* 23: 77-100.
- MCLAY C. L. 2006 — Retroplumidae (Crustacea, Decapoda) from the Indo-Malayan archipelago (Indonesia, Philippine) and the Melanesian arc islands (Solomon Islands, Fiji and New Caledonia), and paleogeographical comments, in RICHER DE FORGES B. & JUSTINE J. L. (eds), *Tropical Deep-Sea Benthos*, vol. 24. *Mémoires du Muséum national d'Histoire naturelle* 193: 375-391.
- MCLAY C. L. & NG P. K. L. 2004 — A taxonomic revision of the genus *Paradynomene* Sakai, 1963 (Crustacea: Decapoda: Brachyura: Dynomenidae). *Zootaxa* 657: 1-24.
- MOOSA M. K. 1981 — Crustacés Décapodes: Portunidae, in DORST (J.) (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 1. Éditions de l'Office de la Recherche Scientifique et Technique d'Outre-Mer, Paris: 141-150 (Mémoires ORSTOM; 91).

- MOOSA M. K. 1996 — Crustacea Decapoda: Deep-water swimming crabs from the South-West Pacific, particularly New Caledonia (Brachyura, Portunidae), in CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 503-530.
- NARUSE T., CASTRO P. & NG P. K. L. 2009 — A new genus and new species of Ethusidae (Crustacea: Decapoda: Brachyura) from Vanuatu, western Pacific. *Zoetaceana* 82 (7): 931-938.
- NG P. K. L. & CHIA D. G. B. 1994 — The genus *Glyptocarcinus* Takeda, 1973, with descriptions of a new subfamily, two new genera and two new species from New Caledonia (Crustacea: Decapoda: Brachyura: Xanthidae). *Raffles Bulletin of Zoology* 42 (3): 701-730.
- NG P. K. L. & MCLAY C. L. 2010 — *Metadynomene tuamotu*, a new species of dynomenid crab from French Polynesia (Crustacea: Decapoda: Brachyura). *Zootaxa* 2405: 48-54.
- NG P. K. L. & RICHER DE FORGES B. 2007 — A new genus and new species of leucosiid crab from New Caledonia, with a note on the validity of *Tanaoa serenei* (Richer de Forges, 1983) (Crustacea: Decapoda: Brachyura). *Zootaxa* 1662: 15-24.
- NG P. K. L. & RICHER DE FORGES B. 2012 — *Pleisticanthoides* Yokoya, 1933, a valid genus of deep-sea inachid spider crabs (Crustacea: Decapoda: Brachyura: Majoidea), with descriptions of two new species from the Philippines, Papua New Guinea and Vanuatu. *Zootaxa* 3551: 65-81.
- NGOC-HO N. 2005 — Thalassinidea (Crustacea, Decapoda) from French Polynesia. *Zoosystema* 27: 47-83.
- NGOC-HO N. 2006 — Three species of *Acanthaxius* Sakai & de Saint Laurent, 1989, including two new to science, from the Solomon Islands and New Caledonia (Crustacea, Thalassinidea, Axiidae). *Zootaxa* 1240: 57-68.
- POORE G. 1997 — A review of the thalassinidean families Callianideidae Kossmann, Micheleidae Sakai, and Thomassiniidae de Saint Laurent (Crustacea, Decapoda) with descriptions of fifteen new species. *Zoosystema* 19: 345-420.
- POORE G. C. B. & COLLINS D. J. 2010 — *Parascytopleptus* and *Spongjaxius* redefined with two new species from the southwestern Pacific (Decapoda, Axiidea, Axiidae), in FRANSEN C. H. J. M., DE GRAVE S., NG P. K. L. (eds), *Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume*. Brill, Leiden: 611-625.
- POUPIN J. 1994 — The genus *Justitia* Holthuis, 1946, with description of *J. chani* and *J. vericeli* spp. nov. (Crustacea: Decapoda: Palinuridae). *Journal of Taiwan Museum* 47 (1): 37-56.
- POUPIN J. 1995 — Étude des *Naxioides* du groupe *robilliardi* (Miers, 1882) (Brachyura: Majidae: Pisinae). *Journal of Natural History* 29 (1): 85-109.
- POUPIN J. 1997 — Les pagures du genre *Calcinus* en Polynésie française, avec le description de trois nouvelles espèces (Crustacea, Decapoda, Diogenidae). *Zoosystema* 19 (4): 683-719.
- POUPIN J. & LEMAIRE R. 2003 — Hermit crabs of the genus *Calcinus* Dana, 1851 (Decapoda: Anomura: Diogenidae) from the Austral Islands, French Polynesia, with description of a new species. *Zootaxa* 391: 1-20.
- POUPIN J. & MCLAUGHLIN P. 1996 — A new species of *Solitariopagurus* Türkay (Decapoda, Anomura, Paguridae) from French Polynesia. *Bulletin du Muséum national d'Histoire naturelle* (4) 18 (A) (1-2): 211-224.
- POUPIN J. & MCLAUGHLIN P. 1998 — Additional *Calcinus* (Decapoda, Anomura, Diogenidae) from French Polynesia with three new species and a key to Indo-West Pacific species. *Crustacean Research* 27: 9-27.
- RICHER DE FORGES B. 1983 — Description d'une espèce nouvelle de Leucosiidae du Pacifique, *Randallia serenei* sp. nov. (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle* (4) 5 (A) (2): 633-640.
- RICHER DE FORGES B. 1995 — Nouvelles récoltes et nouvelles espèces de Majidae de profondeur du genre *Oxyleurodon* Miers, 1886. *Crustaceana* 68: 43-60.
- RICHER DE FORGES B. 1996 — The Genus *Platypilumnus* Alcock and description of *P. jamiesoni* n. sp. from New Caledonia (Crustacea, Decapoda, Brachyura). *Records Australian Museum* 48: 1-6.
- RICHER DE FORGES B. 2006 — Découverte en mer du Corail d'une deuxième espèce de glypheide (Crustacea, Decapoda, Glypheoidea). *Zoosystema* 28 (1): 17-29.
- RICHER DE FORGES B. & CORBARI L. 2012 — A new species of *Oxyleurodon* Miers, 1886 (Crustacea, Brachyura, Majoidea) from the Bismarck Sea, Papua New Guinea. *Zootaxa* 3320: 56-60.
- RICHER DE FORGES B. & GUINOT D. 1988 — Description de trois espèces de *Cyrtomaia* Miers, 1886, de Nouvelle Calédonie et des îles Chesterfield (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle* (4) 10 (A) (1): 39-55.
- RICHER DE FORGES & NG P. K. L. 2007 — New records and new species of Homolidae De Haan, 1839, from the Philippines and French Polynesia (Crustacea: Decapoda: Brachyura). *Raffles Bulletin of Zoology supl.* 16: 29-45.
- RICHER DE FORGES B. & NG P. K. L. 2008a — New records of deep-sea spider crabs of the genus *Cyrtomaia* Miers, 1886, from the Pacific Ocean, with description of a new species (Crustacea: Decapoda: Brachyura: Majidae). *Zootaxa* 1861: 17-28.
- RICHER DE FORGES & NG P. K. L. 2008b — New Western Pacific records of Homolidae De Haan, 1839, with descriptions of new species of *Homolochunia* Doflein, 1904, and *Latreillopsis* Henderson, 1888 (Crustacea: Decapoda: Brachyura). *Zootaxa* 1967: 1-35.
- RICHER DE FORGES B. & NG P. K. L. 2009a — On the majoid genera *Oxyleurodon* Miers, 1886, and *Sphenocarcinus* A. Milne-Edwards, 1875 (Crustacea: Brachyura: Epialtidae), with descriptions of two new genera and five new species. *Raffles Bulletin of Zoology supl.* 20: 247-266.
- RICHER DE FORGES B. & NG P. K. L. 2009b — New genera, new species and new records of Indo-West Pacific spider crabs (Crustacea: Brachyura: Epialtidae: Majoidea). *Zootaxa* 2025: 1-20.
- SAINT LAURENT DE M. 1988 — Enoplometopoidea, nouvelle superfamille de Crustacés décapodes Astacidea. *Comptes Rendus hebdomadaires de l'Académie des Sciences, Paris* (3) 307: 59-62.

- SAINT LAURENT DE M. & MACPHERSON E. 1990 — Crustacea Decapoda: Le genre *Eumunida* Smith, 1883 (Chirostyliidae) dans les eaux néo-caledoniennes, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 6. *Mémoires du Muséum national d'Histoire naturelle* 145: 227-288.
- SAINT LAURENT DE M. & MCLAUGHLIN P.A. 1999 — A new genus and species of hermit crabs (Decapoda: Anomura: Paguridae) from the western Pacific. *Zoosystema* 21(1): 77-92.
- SAINT LAURENT DE M. & POUPIN J. 1996 — Crustacea, Anomura: Les espèces indo-ouest pacifiques du genre *Eumunida* Smith, 1880 (Chirostyliidae) Description de six espèces nouvelles, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 15. *Mémoires du Muséum national d'Histoire naturelle* 168: 337-385.
- SAITO T. & KOMAI T. 2008 — A review of species of the genera *Spongicola* de Haan, 1844 and *Paraspóngicola* de Saint Laurent & Cleva, 1981 (Crustacea, Decapoda, Stenopodidea, Spongicolidae). *Zoosystema* 30: 87-147.
- SERENE R. & VADON C. 1981 — Crustacés Décapodes: Brachyures. Liste préliminaire, descriptions de formes nouvelles et remarques taxonomiques, *in* DORST (J.) (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 1. Éditions de l'Office de la Recherche Scientifique et Technique d'Outre-Mer, Paris: 117-140. (Collection Mémoires ORSTOM; 91).
- TAN S. H. & NG P.K. L. 2010 — A new species of *Pseudolambrus* (Brachyura, Parthenopidae) from New Caledonia, *in* CASTRO P., DAVIE P.J.F., NG P.K.L. & RICHER DE FORGES B. (eds), *Studies on Brachyura: a homage to Danièle Guinot*. Brill, Leiden; Boston: 329-333 (Crustaceana Monographs; 11).
- TAVARES M. 1993 — Crustacea Decapoda: Les Cyclodorippidae et Cyromonomidae de l'Indo-Ouest-Pacifique à l'exclusion du genre Cyromonomus, *in* CROSNIER A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 10. *Mémoires du Muséum national d'Histoire naturelle* 156: 253-313.
- TAVARES M. 1997 — Crustacea Decapoda: Cyclodorippidae récoltés dans l'archipel de Vanuatu (Brachyura), *in* CROSNIER A. (eds), Résultats des campagnes MUSORSTOM, vol. 18. *Mémoires du Muséum national d'Histoire naturelle* 176: 261-271.
- TAVARES M. 1998 — Phyllotymolinidae, nouvelle famille de Brachyures Podotremata (Crustacea, Decapoda). *Zoosystema* 20 (1): 109-122.
- TAVARES M. & CLEVA R. 2010 — Trichopeltaridae (Crustacea, Decapoda, Brachyura), a new family and superfamily of eubrachyuran crabs with description of one new genus and five new species. *Papéis Avulsos de Zoologia, Museu de Zoologia da Universidade de São Paulo* 50 (9): 97-157.
- TSOI K.H., CHANT-Y. & CHU K.H. 2011 — Phylogenetic and biogeographic analysis of the spear lobsters *Linuparus* (Decapoda, Palinuridae), with the description of a new species. *Zoologischer Anzeiger* 250: 302-315.
- YANG C.-H. CHAN T.-Y. & CHU K.H. 2010 — Two new species of the "Heterocarpus gibbosus" Bate, 1888 species group (Crustacea: Decapoda: Pandalidae) from the western Pacific and north-western Australia. *Zootaxa* 2372: 206-220.
- YANG C.-H., CHEN I.-S. & CHAN T.-Y. 2011 — A new slipper lobster of the genus *Galearctus* Holthuis, 2002 (Crustacea, Decapoda, Scyllaridae) from New Caledonia. *Zoosystema* 33 (2): 207-217.