



Centaurea Sect. Cyanus: Nuclear Phylogeny, Biogeography, and Life-Form Evolution

Author(s): Igor Boršić, Alfonso Susanna, Svetlana Bancheva, Núria Garcia-Jacas

Source: *International Journal of Plant Sciences*, Vol. 172, No. 2 (February 2011), pp. 238-249

Published by: [The University of Chicago Press](http://www.uchicago.edu)

Stable URL: <http://www.jstor.org/stable/10.1086/657645>

Accessed: 17/02/2011 02:51

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ucpress>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



The University of Chicago Press is collaborating with JSTOR to digitize, preserve and extend access to *International Journal of Plant Sciences*.

CENTAUREA SECT. CYANUS: NUCLEAR PHYLOGENY, BIOGEOGRAPHY, AND LIFE-FORM EVOLUTION

Igor Boršić,*† Alfonso Susanna,¹† Svetlana Bancheva,‡ and Núria Garcia-Jacas†

*State Institute for Nature Protection, Trg Mažuranića 5, 10000 Zagreb, Croatia; †Botanic Institute of Barcelona (Consejo Superior de Investigaciones Científicas–Institut de Cultura de Barcelona), Passeig del Migdia s/n, Parc de Montjuïc, 08038 Barcelona, Spain; and ‡Institute of Botany, Bulgarian Academy of Sciences, Academician Georgi Bonchev Street, block 23, 1113 Sofia, Bulgaria

Section *Cyanus* of *Centaurea* is a group that is very well defined morphologically and, thus, is a good representative of many radiations of eastern groups of the genus in the Mediterranean region. To confirm the existence of the two natural groups, subsect. *Cyanus* (annual species) and subsect. *Perennes* (perennial taxa), typically defined within this section, and to confirm their radiation patterns, a molecular phylogenetic analysis was carried out using the highly variable nuclear-ribosomal spacers ITS (internal transcribed spacer) and 3'ETS (external transcribed spacer). Our results confirm the eastern origin of the group, which probably arose from a Caucasian and North Iranian stock. Both subsections are monophyletic, and annuals (subsect. *Cyanus*) arose from perennials in Anatolia. The radiations of the two subsections follow very different patterns. Inconsistencies between present classifications and molecular results strongly suggest that the present delineation of some species (*Centaurea triumfetti* being the best example) is incorrect, and a deep taxonomic revision is necessary.

Keywords: biogeography, *Centaurea*, *Cyanus*, ITS, ETS, phylogeny.

Introduction

The *Cyanus* group (Compositae, Cardueae-Centaureinae) was first mentioned by Miller (1754) as a genus, *Cyanus* Mill. The first to consider it a section within the genus *Centaurea* was de Candolle (1838), and this was widely accepted by most subsequent synantherologists (Bentham 1873; Boissier 1875; Hoffmann 1894; Stefanov and Gheorghiev 1931; Dittrich 1968; Wagenitz 1975). Today it is usually recognized as a subgenus (Dostál 1976) or as an informal group within the genus *Centaurea* (Wagenitz and Hellwig 1996; Garcia-Jacas et al. 2001). Some authors, however, reassign it to a generic status (Bancheva and Raimondo 2003; Greuter 2003, 2008; Bancheva and Greilhuber 2006). Molecular studies of Centaureinae by Garcia-Jacas et al. (2001) and the tribe Cardueae by Susanna et al. (2006) have shown that the *Cyanus* and *Jacea* groups constitute one natural group, with the *Cyanus* group being sister to the *Jacea* group. In light of the molecular evidence, the latest compilations of the Cardueae (Susanna and Garcia-Jacas 2007, 2009) suggest that *Cyanus* should be left within *Centaurea*.

In contrast with the somewhat controversial taxonomical status of the *Cyanus* group, its delimitation has never posed a major problem. The group itself has many well-defined morphological characteristics, among which the color of the florets was the first to be observed (Miller 1754). The florets are blue or purplish blue (with only a few exceptions of cream- or

pale pink-flowered taxa), which are extremely unusual colors for the subtribe *Centaureinae*. Dittrich (1968) noticed that the achenes are conspicuously barbate at the margins of the insertion areole. Nevertheless, the structure of the phyllaries is the most unique characteristic. They have pectinate-ciliate, spineless appendages that are decurrent nearly to the base of the phyllary (Wagenitz and Hellwig 1996). Finally, there are features in this group that are shared with the *Jacea* and *Acrocentron* groups. These include the marginal sterile florets without staminodes, the smooth pollen type associated with a reaction pollen presentation mechanism, and the lateral hilum of the seed (Garcia-Jacas et al. 2001).

In addition to the morphological features, the *Cyanus* group is also characterized by its pollen types. According to Wagenitz (1955), two out of the eight pollen types present in the genus *Centaurea* s.l. are confined exclusively to this group, within which two subgroups are defined. One is the subgroup of annual species that have the *Cyanus* pollen type, and the other group consists of perennials with the *Montana* pollen type. The accepted names for the two groups are *Centaurea* sect. *Cyanus* (Mill.) DC. subsect. *Cyanus* and *Centaurea* sect. *Cyanus* subsect. *Perennes* Boiss.

These two subgroups are also corroborated by karyological data. Annual species show a complex dysploid chromosome series with base chromosome numbers $x=8, 9, 10$ and 12 . Perennials, on the other hand, are far more uniform in their base chromosome numbers, with $x=10$ or 11 (Wagenitz and Hellwig 1996; Garcia-Jacas et al. 2001).

The *Cyanus* group is distributed throughout central and southern Europe, North Africa, Asia Minor, and the Caucasus, with some species appearing as far east as Iran and Afghanistan (Wagenitz 1975; Dostál 1976; Czerepanov 2001). The

¹ Author for correspondence; e-mail: asusanna@ibb.csic.es.

subject. *Perennes* mainly lies within the mountain areas of southeast Europe, Turkey, and the Caucasus (Meusel and Jäger 1992; Hellwig 2004). The area with the highest number of described species is the Balkans (Dostál 1976; Bancheva and Raimondo 2003), which is where most of the taxonomical issues are concentrated. A synthetic treatment by Dostál (1976) gave most of the small endemics described in the area the subspecies rank. However, this classification is disputed and remains very complicated, especially in the microspecies of the *Centaurea triumfetti* complex (Olšavská et al. 2009).

The annual species of the group have dual distributions, with two widely dispersed weeds and three narrow endemics. One species, *Centaurea cyanus*, has dispersed synanthropically as a crop weed associated with cereal cultivation since prehistorical times (Rösch 1998). *Centaurea depressa* is also widely distributed (southwest and central Asia, Spain, Balkans, and Crimea) and is associated with agriculture. Three other species (*Centaurea cyanoides*, *Centaurea pinardii*, and *Centaurea tchihatcheffii*) are distributed within a much narrower area centered in southwest Asia. This subsection does not present any systematic problems, as the only doubtful designation is associated with one of the Anatolian endemics, *Centaurea tchihatcheffii*. Because of some of its unusual and unique characteristics, such as its funnel-shaped marginal flowers with crenate margins and anther tubes with glands at the tip of appendages, this species was combined with the genus *Melanoloma* Cass. (now a section of *Centaurea*) as *Melanoloma tchihatcheffii* by Boissier (1875). It was included in the *Cyanus* group by Wagenitz (1975) despite its differences.

To date, there has been no extensive molecular investigation of the *Cyanus* group. However, some representative species were included in wider and more general investigations on the subtribe *Centaureinae* (Garcia-Jacas et al. 2001) and the tribe *Cardueae* (Susanna et al. 2006). Considering the copious nuclear-ribosomal DNA sequences available for the *Centaureinae*, we carried out a molecular survey of the group, using the internal transcribed spacer (ITS) region. We also added the same external transcribed spacer (ETS) that has been used in studies on closely related species of *Compositae* (Baldwin and Markos 1998; Clevinger and Panero 2000; Urbatsch et al. 2003; Suárez-Santiago et al. 2007) with the following aims: (1) to verify the existence of the two subgroups (subsects. *Cyanus* and *Perennes*) as defined by life forms, karyology, pollen types, and biogeography; (2) to explore the relationships between both groups and examine molecular data, which may shed light on the geographic origin of the group; and (3) to verify whether molecular data can improve the systematics of the Balkan group of the taxa.

Material and Methods

Plants

Sampling of the plant material focused on providing the most comprehensive representation of the *Cyanus* group. Therefore, representatives of all five annual species (subsect. *Cyanus*) and 32 taxa (species and subspecies) from the subsect. *Perennes* were included. Several widely distributed species from both subsections were represented with more than one population from different parts of their distribution areas.

The entire distribution range of the group was covered with special emphasis on southeast Europe and Turkey. The nomenclature of the taxa was mainly adjusted according to *Flora Europaea* (Dostál 1976) and *Flora of Turkey* (Wagenitz 1975), with some exceptions (*Centaurea lingulata*, according to Susanna and Garcia-Jacas 1996; *Centaurea pseudoaxillaris* and *Centaurea tuberosa*, according to Bancheva 1999).

Published sequences and new sequences were used in the analysis of the ITS sequences. All of the 3'ETS sequences analyzed were new. The sources for the published sequences, voucher data, and GenBank sequence accession numbers for the newly studied taxa are given in the appendix.

Centaurea scabiosa L. and *Centaurea ochrocephala* Wagenitz from sect. *Acrocentron* were chosen as the outgroup species following Garcia-Jacas et al. (2001). Two plastid noncoding regions, the intergenic spacer *ycf3-trnS* and the intron *trnG*, were also sequenced in several representative taxa; however, the very low number of informative characters led us to abandon this work.

DNA Extraction

Total genomic DNA was extracted from herbarium material following the procedure described in Doyle and Doyle (1987) and Cullings (1992). The DNA of older herbarium material was extracted using the DNeasy Plant Kit (Qiagen, Valencia, CA) following the manufacturer's protocol. In some cases, the fresh leaves of plants cultivated in the Botanic Institute of Barcelona or silica gel-dried leaves collected in the field were used (appendix).

nrDNA ITS and ETS Regions Amplification Strategies

Double-stranded DNA of the entire ITS region (ITS1, 5.8S gene, ITS2) was amplified using 1406F (Nickrent et al. 1994) as the forward primer and ITS4 (White et al. 1990) as the reverse primer. Alternatively, if it was not possible to obtain the whole ITS region, the primers ITS1 (White et al. 1990) and 5.8I1 (Sun et al. 1994) were used to amplify the ITS1 region, and primers 5.8I2 (Sun et al. 1994) and ITS4 were used for the amplification of the ITS2 region.

PCR conditions included a hot start at 95°C for 4 min and 30 cycles of amplification that were carried out under the following conditions: 94°C for 1 min 30 s, 55°C for 2 min, and 72°C for 3 min, with an additional extension step of 15 min at 72°C. The 3' portion of the ETS region was amplified using ETS-Car-1 (Kelch and Baldwin 2003) as the forward primer and 18S-ETS (Baldwin and Markos 1998) as the reverse primer. In *Centaurea depressa*, this combination of primers failed to amplify any products, so ETS1F (Linder et al. 2000) was used as the forward primer and 18S2L (Linder et al. 2000) as the reverse primer instead. PCR conditions for this amplification included a hot start at 95°C for 5 min and 30 cycles that were carried out under the following conditions: 94°C for 45 s, 48°C for 45 s, and 72°C for 40 s, with an additional extension step of 7 min at 72°C.

In some taxa, more than one band was produced by the amplification, and some others could not be sequenced directly. PCR products of those taxa were cloned using the TOPO TA Cloning Kit (Invitrogen, Carlsbad, CA) following the manu-

facturer's instructions, except that only one-half of the reagents were used. When possible, eight positive colonies from each reaction were screened with direct PCR using the T7 and M13R universal primers following the protocol described by Vilatersana et al. (2007). Five to eight PCR products were selected for sequencing in both directions using the same primers.

nrDNA Sequencing Strategies

PCR products were purified with the QIAquick PCR Purification Kit (Qiagen) and sequenced with the same primers used in the amplification. Direct sequencing of the amplified DNA segments was performed using the BigDye Terminator Cycle Sequencing v3.1 (Applied Biosystems, Foster City, CA), according to the protocol recommended by the manufacturer. The nucleotide sequencing was performed at the Serveis Científic-Tècnics at the University of Barcelona on an ABI PRISM 3700 DNA Analyzer (Applied Biosystems).

Phylogenetic Analysis

Nucleotide sequences were edited using BioEdit 7.0.1 (Hall 1999) and aligned visually by sequential pairwise comparison (Swofford and Olsen 1990). For the clones, the sequences with unique substitutions in single clones were excluded. Consensus sequences were generated for each of the cloned species, condensing the single base pair differences among clones. This reduced the size of the matrices and the impact of any potential PCR artifacts (chimeric sequences and Taq errors; Cline et al. 1996; Popp and Oxelman 2001). Matrices of obtained data are available on request from A. Susanna.

Two different phylogenetic analyses were performed: maximum parsimony, as implemented in PAUP 4.0b10 (Swofford 2002), and Bayesian inference (BI), using MrBayes 3.1.2 (Huelsenbeck and Ronquist 2001; Ronquist and Huelsenbeck 2003). Separate analyses were conducted on the ITS and the 3'ETS sequence data.

The maximum parsimony analysis involved heuristic searches using tree bisection-reconnection (TBR) branch swapping with character states specified as unordered and unweighted. The in-dels were treated as missing data. All most-parsimonious trees (MPTs) were saved. To locate islands of MPTs (Maddison 1991), 1000 replications were performed with random taxon addition and TBR branch swapping. Bootstrap support (BS) was estimated (Felsenstein 1985) with 100 replicates, simple taxon addition, and TBR branch swapping.

To determine the best-fit model of molecular evolution for Bayesian inference, the ITS and the 3'ETS matrices were analyzed using hierarchical likelihood ratio tests and Akaike Information Criteria, as implemented in MrModeltest 2.2b (Nylander 2004), which considers only those nucleotide substitution models that are currently implemented in PAUP and MrBayes 3.1.2 (Huelsenbeck and Ronquist 2001; Ronquist and Huelsenbeck 2003). In both cases, each criterion selected different evolutionary models under which BI analyses were then performed. They were initiated with random starting trees and run for 1×10^6 generations. Four Metropolis-coupled Monte Carlo Markov chains were sampled every 100 genera-

tions, resulting in 10,000 sample trees. A critical aspect of the Bayesian analysis is to ensure that the Markov chain has reached stationarity. Because all sample points before stationarity are essentially random and usually do not contain useful parameter estimates, 1000 sample trees were discarded as "burn-ins." Internodes with posterior probabilities of at least 95% were considered to be statistically significant. Trees obtained using different evolutionary models were very similar in their topology, but the general time reversible model (Rodríguez et al. 1990) model, which was common to both analyses, showed slightly higher values of clade support.

The nrDNA ITS and 3'ETS data sets were tested for congruence using the incongruence length difference (ILD) test (Farris et al. 1995a, 1995b), as implemented in WinClada 1.00.08 (Nixon 2002). It was conducted with 10,000 replicates and 10 random addition sequences per replicate, holding two trees at each step and saving two trees per replicate.

Results

The numerical results of the analyses of the ITS and 3'ETS data sets are summarized in table 1. The ITS alignment of the 74 taxa consisted of 478 positions and contained 125 phylogenetically informative substitutions. Mean pairwise distances (as calculated by PAUP) within the ingroup varied from 0% (between 30 pairs of taxa of the *Cyanus* group of *Centaurea*) to 9.82% (between *Centaurea tchihatcheffii* and *Centaurea triumfetti* subsp. *stricta*). Pairwise distances between the ingroup and outgroup varied from 5.12% (between *Centaurea scabiosa* and *Centaurea epirota* population 2) to 10.26% (between *Centaurea ochrocephala* and *Centaurea cf. triumfetti* subsp. *adscendens*).

The 3'ETS alignment of the 47 taxa consisted of 668 positions and contained 144 phylogenetically informative substitutions. Mean pairwise distances (as calculated by PAUP) within the ingroup varied from 0% (between eight pairs of taxa of the *Cyanus* group of *Centaurea*) to 23.88% (between *Centaurea depressa* population 3 and *Centaurea pinardii*). Pairwise

Table 1
Comparison of Results Obtained from the Internal Transcribed Spacer (ITS) and 3' External Transcribed Spacer (ETS) Data Sets

Data set	ITS	3'ETS
No. taxa	74	47
Total characters	478	668
Informative characters	125	144
No. MPTs	201,196	532,214
No. steps	264	301
Islands	2	4
Consistency index	0.5451	0.5475
Retention index	0.8056	0.7430
Homoplasy index	0.4549	0.4525
Range of divergence, ingroup (%)	0–9.82	0–23.88
Range of divergence, ingroup-outgroup (%)	5.12–10.26	9.48–24.66

Note. The consistency and homoplasy indices are calculated excluding uninformative characters.

distances between the ingroup and outgroup varied from 9.48% (between *C. ochrocephala* and *C. epirota* population 1) to 24.66% (between *C. scabiosa* and *C. pinardii*).

Only two species were cloned, one for the ITS and another for the ETS region. The first was *Centaurea pichleri* population 1, in which we found four ITS clones. The second cloned species was *Centaurea cf. pindicola*, in which we found three ETS clones.

The topology of the trees obtained by parsimony and Bayesian approaches was consistent, although a slightly better resolution was obtained with the Bayesian trees. Therefore, only Bayesian majority-rule consensus trees are shown, with Bayesian posterior probability values (PP) and parsimony BS percentages added onto the branches (figs. 1, 2).

Some noticeable discrepancies were observed in the topologies of the trees obtained from the ITS and 3'ETS regions. The first incongruence was the changing positions of *Centaurea triumphetti* subsp. *lugdunensis* and *C. triumphetti* subsp. *aligera*. The two species were placed together in the ITS and 3'ETS analyses, but both the composition of their clade and their positions within the *Perennes* clade (figs. 1, 2) were different. Another inconsistency was the position of *C. nigrofimbria*, which was placed within the Caucasian and Iranian clade in the ITS analysis (fig. 1) but formed a clade with the Turkish *C. triumphetti* population 1 in the 3'ETS analysis. Finally, the clade formed by the Iranian and Caucasian representatives was sister to the rest of the *Perennes* clade in the ITS analysis (fig. 1), a position that was not retained in the 3'ETS tree (fig. 2). Regarding ILD, two different tests were carried out, the first with all the taxa and the second omitting the three species of incongruous position. The data sets were incompatible ($P = 0.001$ in both analyses). Considering the differences between the topologies obtained by both data sets and the negative result of the ILD, combining them into a single matrix was not possible.

Trees obtained in both analyses (ITS region, fig. 1; 3'ETS region, fig. 2) showed absolute support for the monophyly of the *Cyanus* group (PP = 1.00, BS = 100%). In the 3'ETS analysis, the *Cyanus* group was divided into two well-supported clades. The first clade included all of the annual species with strong Bayesian support in the 3'ETS (PP = 1.00, BS = 72%; fig. 2), but it was not supported by the Bayesian analysis of the ITS data set (PP = 0.88; fig. 1). The second clade encompassed the perennial taxa, within which two clades were formed in the ITS analysis: the first one with the Caucasian and Iranian endemic species and a robust second clade with the rest of the taxa. In both the ITS and 3'ETS analyses, this second clade was a polytomy with several more or less well-supported groups.

The ITS clones 2 and 4 of *C. pichleri* population 1 were grouped with *C. fuscomarginata* in a strongly supported clade, but clones 1 and 3 were placed as successive sisters to this clade without support (BS < 50%, PP = 0.63 and 0.58, respectively; fig. 1). As for the ETS clones of *C. cf. pindicola* (fig. 2), one of the clones was grouped with population 2 of *C. epirota* with high support (PP = 1.00, BS = 82%), and the other two clones were placed in a polytomy with populations 2, 5, and 6 of *Centaurea lingulata*; *C. fuscomarginata*; populations 1, 3, and 4 of *C. pichleri*; and *Centaurea reuterana* var. *phrygia* with high Bayesian support (PP = 1.00, BS = 73%).

Discussion

Taxonomic Implications

The results of the analyses of the ITS and 3'ETS regions confirm the monophyly of sect. *Cyanus*. As shown in morphological analyses by Wagenitz and Hellwig (1996) and previous molecular analyses by Garcia-Jacas et al. (2001), two subgroups are defined within sect. *Cyanus*: annuals and perennials.

Our results point to new evolutionary inferences in the clade of annual species. First, the populations of widespread *Centaurea cyanus* and *Centaurea depressa*, associated with agriculture, each form strongly supported clades. In the ITS analysis, the clade formed by the populations of *C. cyanus* is the sister to the strongly supported clade formed by the rest of annuals. The surprising position of *C. cyanus* is probably caused by a different mutation rate in the ITS region, which is usually related to differences in the biological history of the species (Smith and Donoghue 2008). In fact, the branches of the annuals are much longer than those of the perennials in both phylograms (figs. 1, 2) due to the accumulation of changes, as suggested by Ainouche and Bayer (1999) and Andreassen and Baldwin (2001). However, this general rule is not free of exceptions, depending on the group (Andreassen 2005; Soria-Hernanz et al. 2008). In contrast, the 3'ETS analysis places *C. cyanus* well nested in the subsect. *Cyanus* (fig. 2). Thus, deviating behavior in annuals, when included in molecular analyses, seems to mainly affect the ITS region, as observed in other groups from the subtribe *Centaureinae* (A. Susanna, M. Galbany, K. Romaschenko, L. Barres, and N. Garcia-Jacas, unpublished manuscript). Finally, all of the analyses have confirmed that *Centaurea tchihatcheffii* is more closely related to the *Cyanus* than to the *Melanoloma* group, which is consistent with the placement by Wagenitz (1975). The ITS results place it as most closely related to the Anatolian group of taxa, including *Centaurea pinardii* and *C. depressa* (fig. 1).

It is very difficult to draw taxonomical conclusions from our results about the clade *Perennes*, other than confirming the need for its taxonomic revision in view of the problems in defining some species. The first of the conflicting species is *Centaurea triumphetti*, which traditionally has been considered to be a difficult taxon (Wagenitz 1975; Gamal-Eldin and Wagenitz 1991). Populations of alleged *C. triumphetti* are placed in different clades (figs. 1, 2). Both the ITS and 3'ETS analyses place *C. triumphetti* subsp. *dominii*, *C. triumphetti* subsp. *axillaris*, and *C. triumphetti* subsp. *pirinensis* into one clade, whereas *C. triumphetti* subsp. *triumfetti* is positioned in a different clade. These results agree with the morphologically based conclusions of Olšovská et al. (2009). Another example is *C. triumphetti* subsp. *lugdunensis*, a subspecies from western Europe that is morphologically very similar to *C. triumphetti* and is alternately placed in both clades of *C. triumphetti* (figs. 1, 2). The last important issue regarding the taxonomy of *C. triumphetti* s.l. is the case of *Centaurea lingulata* from the Iberian Peninsula and North Africa, subordinated to *C. triumphetti* as a subspecies (Dostál 1976). Our results show that the Iberian and North African populations of *C. lingulata* are more closely related to the eastern Mediterranean stock represented by *Centaurea fuscomarginata*, *Centaurea pichleri*, and *Centaurea reuterana* var. *phrygia* (fig.




Fig. 1 Majority-rule consensus tree based on Bayesian Monte Carlo Markov chain analysis of internal transcribed spacer region. Numbers above branches are Bayesian posterior probability values; numbers below branches are bootstrap percentages.




Fig. 2 Majority-rule consensus tree based on Bayesian Monte Carlo Markov chain analysis of 3' external transcribed spacer region. Numbers above branches are Bayesian posterior probability values; numbers below branches are bootstrap percentages.

2). This result supports the status of *C. lingulata* as an independent species. A thorough revision of these taxa is highly necessary, and either the delineation or the independence of some species should be questioned.

Another species of controversial delineation is *C. pichleri* from Greece, Bulgaria, Turkey, and Lebanon. The populations of this taxon included in our study appear to be diversely associated with other eastern taxa and collectively do not form a supported clade in any of our analyses (figs. 1, 2). Thus, this is another taxon in need of taxonomic revision.

Revision of species boundaries in subsect. *Perennes* is a very difficult task because of the scarcity of unifying morphological characteristics present (Wagenitz 1975). Most of the current taxonomy of the group is based on vegetative characters such as the presence of sterile rosettes, the presence of taproots or tubers, and the shape and indument of the leaves (Wagenitz 1975; Dostál 1976; Czerepanov 2001). One of these characters, the presence of napiform tubers, is confirmed by our analyses to be a good indicator of phylogenetic affinity. *Centaurea napulifera*, *Centaurea orbelica*, *Centaurea tuberosa*, and *Centaurea velenovskyi*, which all possess tubers, form a well-supported clade in the ITS analysis (fig. 1). However, other tuberose species like *Centaurea diospolitana* (Bancheva & S. Stoyanov) Bancheva, **comb. nov.** [Basionym *Cyanus diospolitanus* Bancheva & S. Stoyanov, Novon (2009) 421], *C. napulifera* subsp. *thirkei*, *Centaurea nyssana*, and *Centaurea pseudoaxillaris*, which has rudimentary tubers, are not included in this clade. The latter is grouped to another population of *C. napulifera* that is also placed outside the “tuberose” clade. Finally, it was also suggested by Wagenitz (1980) that *Centaurea cheiranthifolia* is related to *Centaurea elbrusensis*, which was corroborated by our findings, suggesting that both of these species belong in the same group.

Introgression and Reticulation

The difficulties in species delimitation may be caused by a still-incomplete evolutionary differentiation, although they can also be due to hybridization and introgression (Hellwig 2004). The hypothesis of introgression was suggested (Font et al. 2002) and recently confirmed (Garcia-Jacas et al. 2009) for *Centaurea* sect. *Acrocentron*. Löser and Hellwig (2007) reported polymorphisms among plastid DNA haplotypes in most populations of perennial species of *Cyanus* that could also indicate hybridization.

Introgression may also be indicated in the results obtained from *Centaurea pindicola*. The 3'ETS analysis includes two cloned sequences of this species. One is grouped with *Centaurea epirota* and the other with *C. pichleri* (fig. 2). The presence of different copies with different affinities and a lack of congruence between the results of the two data sets strongly suggest introgression, as already noted in *C. triumfetti* by Wagenitz (1975) and Gamal-Eldin and Wagenitz (1991). The alternative hypothesis for explaining multiple copies would be the presence of pseudogenes or incomplete lineage sorting of ancestral polymorphisms. Pseudogenes and polymorphisms can be ruled out in our case because cloning resulted in well-preserved sequences in both the ITS and ETS regions (Yamaji et al. 2007; Grimm and Denk 2008).

Biogeographic Implications

Biogeographic implications are also very much of interest. The analysis of the ITS data set defines as the sister to subsect. *Perennes* the Caucasian and North Iranian representatives of the subsection: *C. cheiranthifolia* and *Centaurea woronowii* from Georgia, *C. elbrusensis* from Iran, *Centaurea fischeri* from Armenia, and *Centaurea nigrofimbria* from southeast Russia (fig. 1). Dittrich (1968) reported that *C. cheiranthifolia* diverged earlier than the other perennial species on the basis of achene characteristics. This group corresponds to the so-called subsect. *Fischerianae* Czer. (Czerepanov 2001), with the exception of *C. nigrofimbria*, which was placed into another subsection.

Significantly, a genus from subtribe Centaureinae that has been hypothesized to be related to *Centaurea* sect. *Cyanus* on a morphological basis, *Psephellus* Cass., has a very similar geographic distribution limited to the Caucasus, Anatolia, and northern Iran (Susanna and Garcia-Jacas 2009). The morphological similarities of *Psephellus* and sect. *Cyanus* are evident and include an exine ultrastructure, unarmed appendages on the bracts, large and showy sterile peripheral florets, purple or pink corollas, and seeds with lateral hilum (Dittrich 1968; Wagenitz and Hellwig 2000; Susanna and Garcia-Jacas 2009).

The Iberian and North African *C. lingulata* is more closely related to the eastern Mediterranean stock, as represented by *C. fuscomarginata*, *C. pichleri*, and *C. reuterana* (fig. 2), noted above. In contrast, one species from northern Iberia, *C. triumfetti* subsp. *semidecurrens*, is more related to the eastern European stock (figs. 1, 2). These relationships parallel the case of *Centaurea* sect. *Acrocentron* (Font et al. 2009), in which two migrations were suggested, one from the south connecting Iberia and North Africa and a second one from central Europe.

Evolution of Life Forms

Löser et al. (2009) maintained that the perennial taxa are derived from the annuals on the basis of molecular data, morphology, and karyology. Garcia-Jacas et al. (2001) pointed out that the divergence in the annual *Cyanus* was higher than that of *Perennes*, which could indicate that *Perennes* is more recent. We shall review the basis of the hypothesis regarding the evolutionary direction taken by the group.

Morphology alone cannot be used to decide which subsection is ancestral, as reversals and parallel evolution are a constant in the *Centaureineae* (Garcia-Jacas et al. 2001). Karyology could be a stronger argument, since the annual *C. cyanus* has the highest basic chromosome number of the *Cyanus* group ($x=12$). However, the difference between the *Cyanus* and *Perennes* basic chromosome number ($x=12$ and $x=11$) is minimal, and subsect. *Cyanus* also has the lowest number of the group ($x=8$). Finally, our molecular results cannot provide clarification as to which group is older because *Cyanus* and *Perennes* are sister clades. Branch lengths in the annuals are longer than in perennials, but whether the differences are due to the varying life histories and habits of these two groups is unknown. Comparisons between groups with different life cycles are risky.

There are arguments to support the hypothesis that the annual species of *Cyanus* are derived from perennials. First, we can argue against the biogeographic evidence. Perennial taxa

are, without exception, mesophilous plants growing in the circum-Mediterranean mountains, a conservative habitat where most of the old preglacial flora survived the glaciations (Médail and Diadema 2009). In the Mediterranean region, aridity is a novel condition that arose only in the late Miocene era (Suc 1984). Annual species would have been derived in the east as an adaptation to the increasing aridity, which is consistent with the latest datation of the group that assigned an age of 6–7 million years to *Cyanus* and *Perennes* (Barres, personal communication). The geographic distribution of annuals is very suggestive, with their main diversification center in Anatolia. Our results suggest that the perennial species of *Cyanus* originated in the Caucasus and the adjacent mountains in the extreme north of the Irano-Turanian region (fig. 1), which is consistent with the hypothesis of *Psephellus* being the most plausible ancestor of the *Cyanus* group (Susanna and Garcia-Jacas 2009). Annuals probably appeared very soon after in the more arid foothills.

Another argument supporting the idea that annuals are derived from perennials involves pollen type. There is a clear evolutionary line that connects the pollen types *Psephellus*, *Montana*, and *Cyanus*, as reported by Wagenitz (1955), even though this is unsupported by molecular analyses (Susanna and Garcia-Jacas 2009). These three types share identical exine structure (well-conserved endexine in two layers of columellae and smooth ectexine), and differences between them are due only to shape: spherical in *Psephellus*, prolate in *Montana* and prolate-laterally compressed in *Cyanus* (“stadium form”; Wagenitz 1955). In a recent survey of the *Centaureinae*, Susanna and Garcia-Jacas (2009) correlated the evolution of pollen shape from spherical to prolate with the adaptation to the great oscillations in humidity and temperature typical in arid climates. The laterally compressed *Cyanus* type would be the ultimate adaptation to the extreme oscillations of temperatures in a Mediterranean climate. From our point of view, the most par-

simonious hypothesis would be that the *Cyanus* pollen type (annual species) evolved from the *Montana* type of perennials. And, finally, on the basis of the anatomical characteristics of achenes, Dittrich (1968) concluded that perennials were phylogenetically older. We favor the hypothesis that the group evolved from mesophilous-montane perennials (*Cyanus* subsect. *Perennes*) to annual species more adapted to the Mediterranean aridity (*Cyanus* subsect. *Cyanus*). This would have been through two radical changes: first, the adoption of an annual habit; second, the adaptation of the pollen type to a model more suited for the extreme changes in hygrometry and temperature that are typical of a Mediterranean habitat. More studies within the group are needed to confirm any of the contradicting hypothesis, but difficulties in comparisons between annuals and perennials make molecular methods a less reliable tool.

Acknowledgments

We thank the curators of the herbaria BC, FI, G, K, LE, SO, SOM, UPA, Z, ZA, ZAHO, and ZT for specimen loan and Olja Vasić for providing *C. nyssana* material. Igor Boršić thanks all of our friends from the Botanic Institute of Barcelona who helped in many ways during our laboratory work, data analyses, and the preparation of the manuscript. Kostyantyn Romaschenko kindly prepared the GenBank submissions. Financial support from the Spanish Ministry of Education and Science (project CGL2007–60781/BOS) and the Generalitat de Catalunya (Ajuts a Grups de Recerca Consolidats 2009-SGR 439) is gratefully acknowledged. Igor Boršić benefited from a grant from the Ministry of Foreign Affairs and Cooperation and the Spanish Agency of International Cooperation (MAEC-AECI) for the academic year 2006–2007. Suggestions by two anonymous referees and the editor greatly improved the quality of the manuscript.

Appendix

Voucher Information

The voucher information includes names of taxa, geographic origin, and collection details of the materials; herbaria where the vouchers are deposited; and GenBank accession numbers (new sequences are underlined). A dash indicates a region that was not sequenced.

Centaurea achtarovii Urum., Bulgaria, Blagoevgrad province: Pirin Mountains, under Vihren Peak, *Draževa*, 10.8.1973 (SO), HQ147721, HQ147613, –. *Centaurea angustifolia* Mill., France: Alpes de Savoie, Brezon’s pastures, *Bourgeau 150*, July 1848 (LE), HQ147722, HQ147614, –. *Centaurea cheiranthifolia* Willd., Georgia, South Ossetia: Ermani, left bank, middle Ermani gorge, 2300 m, *Illarionova*, 16.7.1958 (LE), HQ147723, HQ147615, HQ147676. *Centaurea cyanoides* Wahlenb., population 1, *Martins and Hellwig* 2005, AJ867998, AJ867999, –. *Centaurea cyanoides* Wahlenb., population 2, Lebanon, Mount Lebanon Governorate: to Ain Aanoub, 715 m, among young wheat, *Maitland 391*, early May 1959 (LE), HQ147724, HQ147616, HQ147677. *Centaurea cyanus* L., population 1, USA: Washington, Whitman County, Pullman, Hilltop Stables, *Susanna 1450 & Garcia-Jacas*, 5.1993 (WS), HQ147725, HQ147617, HQ147678. *Centaurea cyanus* L., population 2, France, Lozère: Causse de Sauveterre, cultivated fields between le Sec and l’Aumède, near Chanac, *Carretero & Vilatersana 51*, 5.7.1997 (BC), HQ147726, HQ147618, –. *Centaurea cyanus* L., population 3, Susanna et al. 2006, AY826254, HQ147679. *Centaurea depressa* M. Bieb., population 1, Iran: Mazandaran, Chalus Road, Azad-bar Mountains between Gachsar and Valiabad, fields, *Susanna 1619 et al.*, 1.8.1996 (BC), HQ147728, HQ147620, –. *Centaurea depressa* M. Bieb., population 2, Garcia-Jacas et al. 2001, AY012280, AY012316, –. *Centaurea depressa* M. Bieb., population 3, Armenia: Kamo, near Lake Sevan in Cape Noraduz, 2000 m, margins of cultivated fields, *Susanna 1513 et al.*, 16.8.1995 (BC), HQ147727, HQ147619, HQ147680. *Centaurea depressa* M. Bieb., population 4, Spain, Toledo: El Emperador (Urda), *Garcia-Jacas & Susanna 2640*, 5.6.2006 (BC), HQ147729, HQ147621, –. *Centaurea diospolitana* (Bancheva & S. Stoyanov) Bancheva, Bulgaria, Yambol Province: Toundzha hilly country, Lesovo village, ~350 m, *Bancheva 0621*, 25.05.2006 (SOM), HQ147730, HQ147622, HQ147681. *Centaurea elbrusensis* Boiss. & Buhse, Iran: East Azerbaijan (Azarbayjan-e-sharghi), heights of the Sabalan coming

from Movil, *Susanna 1666 et al.*, 5.8.1996 (BC), [HQ147731](#), [HQ147623](#), [HQ147682](#). *Centaurea epirota* Halácsy, population 1, Greece: Ioanninon, Pogoniou, Epirus, Mount Dhouskon (Nëmerçka) 3–4 km N of the village Kakolakkos, 1900–2200 m, *Hartvig & Christiansen 7925*, det. *Landström*, 15.7.1979 (UPA), [HQ147732](#), [HQ147624](#), [HQ147683](#). *Centaurea epirota* Halácsy, population 2, Republic of Macedonia: Prilep District, on stony pasturages on Drenska Mountain, leg. *Nikoloff*, det. *Stojanoff*, 23.5.1918 (SO), [HQ147733](#), [HQ147625](#), [HQ147684](#). *Centaurea fischeri* Willd., Armenia: Idjevan, north side of Sevan Pass, 1900 m, damp margins of pastures, *Susanna 1520 et al.*, 17.8.1995 (BC), [HQ147734](#), [HQ147626](#), -. *Centaurea fuscomarginata* (K. Koch) Juz., Ukraine, Crimea: Demerdzhi plateau, *Hort & Burdeynaya*, 17.6.1974 (LE), [HQ147735](#), [HQ147627](#), [HQ147685](#). *Centaurea lingulata* Lag., population 1, Susanna et al. 2006, AY826258, -. *Centaurea lingulata* Lag., population 2, Spain, Jaén: Mancha Real, Sierra de Mágina, by the relay station on the Almadén Peak, ~1700 m, bush community on limestone, *Susanna 1607 et al.*, 23.6.1996 (BC), [HQ147739](#), [HQ147631](#), [HQ147687](#). *Centaurea lingulata* Lag., population 3, Spain, Teruel: Sierra del Pobo, stoneware upper part of TVE relay, lawn, 1700 m, *Litzler 72/702 E*, 5.7.1972 (ZT), [HQ147737](#), [HQ147629](#), -. *Centaurea lingulata* Lag., population 4, Spain, La Rioja: Soto en Cameros, Luezas, slopes in regeneration, limestone, 1180 m, UTM 30TWM 4282, *Alejandro 1006/88*, 5.6.1988. (Z), [HQ147736](#), [HQ147628](#), -. *Centaurea lingulata* Lag., population 5, Morocco, Taza-Al Hoceima-Taounate: Aknoul, Tizi-Ouzli, ascent to Jbel Azrou Akchar, from the mountain pass, 1932 m, *Hidalgo & Romo 13075*, 12.6.2005 (BC), [HQ147738](#), [HQ147630](#), [HQ147686](#). *Centaurea lingulata* Lag., population 6, Spain, Madrid: in pastures and grasslands near Cercedilla in Sierra de Guadarrama, granite soil, 1300–1400 m, *Vicioso* (LE), [HQ147740](#), [HQ147632](#), [HQ147688](#). *Centaurea maramarosiensis* (Jáv.) Czerep., population 1, Ukraine, Zakarpattia Province: Rakhiv region, Breboia, Yalynoviy forest, *Rocug*, 28.7.1946 (LE), [HQ147742](#), [HQ147634](#), -. *Centaurea maramarosiensis* (Jáv.) Czerep., population 2, Ukraine: Maramureş, in the upper part of Black Tisza's valley, *Margittai*, 14.7.1937 (ZT), [HQ147741](#), [HQ147633](#), -. [HQ147743](#), [HQ147635](#), -. *Centaurea mollis* Waldst. & Kit., Hungary: Bükk Mountains, above Felađháamor, *Hulják*, 1.7.1929 (LE), [HQ147743](#), [HQ147635](#), -. *Centaurea montana* L., population 1, Susanna et al. 2006, L35887, -. *Centaurea montana* L., population 2, France: Hautes-Alpes, path of Crevasses, mountain pass of Lautaret, *Hidalgo 204*, 5.7.2007 (BC), [HQ147744](#), [HQ147636](#), [HQ147689](#). *Centaurea napulifera* Rochel, population 1, Bulgaria, Plovdiv Province: Central Balkan National Park, Central Stara Planina Mountain, rocky meadow near hut "Rai", Kalofersko, 1950 m, *Achtarov*, 3.8.1941 (LE), [HQ147745](#), [HQ147637](#), -. *Centaurea napulifera* Rochel, population 2, Bulgaria: Central Stara Planina Mountain, ~1700 m, leg. *Stoyanov*, det. *Bancheva 03126*, 10.07.2003 (SOM), [HQ147746](#), [HQ147638](#), [HQ147690](#). *Centaurea cf. napulifera* Rochel, Greece: Dramas, Rhodopi Mountains, N of Zagradena forest station, area called Trigono near the Bulgarian border, 1800–1850 m, meadow in opening of *Picea abies* forest, crystalline schist, *Strid 19499 et al.*, 25.7.1981 (G), [HQ147747](#), [HQ147639](#), -. *Centaurea napulifera* Rochel subsp. *thirkei* (Sch. Bip.) Dostál, Garcia-Jacas et al. 2006, DQ319136, -. *Centaurea nigrofimbria* Sosn., Russia: Krasnodarskiy kray region, Krasnaya polyana, Kordon Pslukh, 1600 m, *Mishenkova 163*, 9.9.1971 (LE), [HQ147748](#), [HQ147640](#), [HQ147691](#). *Centaurea nyssana* Petrović, Serbia: Suva Mountain, Divna gorica Peak, 1400 m, limestone, *Niketić*, 13.08.2006 (BEO), [HQ147749](#), [HQ147641](#), [HQ147692](#). *Centaurea ochrocephala* Wagenitz, Font et al. 2002, FJ459678, FJ459622. *Centaurea orbelica* Velen., population 1, Republic of Macedonia: Prilep District, at the foot of Pletvar Mountain, leg. *Nikoloff*, det. *Stojanoff*, 17.4.1918 (SO), [HQ147751](#), [HQ147643](#), -. *Centaurea orbelica* Velen., population 2, Bulgaria, Kyustendil Province: Rila Mountain, Suhoto Lake, ~2400 m, *Bancheva 0626*, 12.06.2006 (SOM), [HQ147750](#), [HQ147642](#), [HQ147693](#). *Centaurea pichleri* Boiss. subsp. *pichleri*, population 1, Turkey: Bursa, Uludağ height, 1950 m, steppe, *Nydegger 14091*, 7.7.1979 (G), [HQ147754](#), [HQ147646](#), [HQ147695](#). *Centaurea pichleri* Boiss. subsp. *pichleri*, population 2, Turkey: Kütahya, Kosik Sögüt at Murat Dağ, 1700 m, clearcut with streamlet, *Nydegger 44058*, 4.6.1989 (G), [HQ147755](#), [HQ147647](#), -. *Centaurea pichleri* Boiss., population 1, Turkey: Ilgaz Dağı Milli Parkı, 40°04', 208'N, 32°43', 850'E, forest of *Abies nordmanniana*, *Romo 14740 et al.*, 19.6.2007 (BC), clone 1, [HQ147758](#), [HQ147650](#), [HQ147696](#); clone 2, [HQ147759](#), [HQ147649](#), [HQ147696](#); clone 3, [HQ147756](#), [HQ147651](#), [HQ147696](#); clone 4, [HQ147757](#), [HQ147648](#), [HQ147696](#). *Centaurea pichleri* Boiss., population 2, Bulgaria, Smolyan Province: Eastern Rhodope Mountains, between Mandritza village and Likan Cheshma locality, ~700 m, *Bancheva 0302*, 25.04.2003 (SOM), [HQ147752](#), [HQ147644](#), -. *Centaurea pichleri* Boiss., population 3, Lebanon, North Governorate: northern Mount Lebanon above Bsharri in cedar forest, 1925 m, *Bornmüller 12009 & Bornmüller*, 3.–5.7.1910 (LE), [HQ147753](#), [HQ147645](#), [HQ147694](#). *Centaurea pichleri* Boiss. population 4, Turkey: between Pinarbasi and Sariz, 38°33', 458'N, 36°26', 705'E, 1781 m, stepic grassland with *Juniperus foetidissima*, *Romo 14864 et al.*, 23.6.2007 (BC), [HQ147760](#), [HQ147652](#), [HQ147697](#). *Centaurea pinardii* Boiss., Garcia-Jacas et al. 2006, DQ319146, [HQ147698](#). *Centaurea cf. pindicola* Griseb., Greece: E-exposed, rocky steep hillside at hut A, east side of Mount Olympus, ~25km SSW from Katerini, 1990–2200 m, leg. *Baltisberger & Meili 11381*, det. *Wagenitz*, 23.6.1987 (ZT), clone 1, [HQ147761](#), [HQ147653](#), [HQ147701](#); clone 2, [HQ147761](#), [HQ147653](#), [HQ147699](#); clone 3, [HQ147761](#), [HQ147653](#), [HQ147700](#). *Centaurea pseudoaxillaris* Stef. & Georg., Bulgaria, Sofia Province: in fields at Nova Mahala, leg. *Štríbrný*, det. *Czerepanov*, May 1900 (LE), [HQ147762](#), [HQ147654](#), [HQ147702](#). *Centaurea reuteriana* Boiss., Turkey: Muğla, Sandras Dağ, W side of the summit area, 37°04'N, 28°50'E, 2100–2200 m, snowbed meadows and rocky slopes, serpentine, leg. *Strid 23345 et al.*, det. *Wagenitz*, 7.7.1984 (G), [HQ147763](#), [HQ147655](#), -. *Centaurea reuteriana* Boiss. var. *phrygia* Bornm., Turkey: Ala Dağ, track above Demir Kazk, *Susanna 2296 et al.*, 3.8.2002 (BC), [HQ147764](#), [HQ147656](#), [HQ147703](#). *Centaurea scabiosa* L., Font et al. 2009, FJ459692, FJ459636. *Centaurea tchibatcheffii* Fisch. & C. A. Mey., Garcia-Jacas et al. 2006, DQ319170, [HQ147704](#). *Centaurea triumfetti* All., population 1, Turkey: Balçesme, at the entrance of the village, *Susanna 2398 et al.*, 7.8.2002 (BC), [HQ147779](#), [HQ147671](#), [HQ147717](#). *Centaurea triumfetti* All., population 2, Slovenia: Primorsko, in meadows near village Senadole at the foot of Vremščica Mountain, ~450 m, *Palma & Trpin*, 18.6.1980 (LE), [HQ147772](#), [HQ147664](#), [HQ147710](#). *Centaurea triumfetti* All. subsp. *adscendens* Bartl., Croatia, Karlovac county: Klek Mountain, on the upper side opposite to Ogulin, ~1100 m, *Kárpáti*, 16.6.1935 (ZT), [HQ147778](#),

HQ147670, HQ147716. *Centaurea* cf. *triumfetti* All. subsp. *adscendens* (Bartl.) Dostál, Croatia, Lika-Senj county: on grassy hills at Ljubovo, on Velebit above Halan, *Farkaš-Vukotinović*, June–July 1856 (ZA), [HQ147769](#), [HQ147661](#), –. *Centaurea triumfetti* All. subsp. *aligera* (Gugler) Dostál, Italy, Lombardy: in shrubs at Mount Lavino, 500 m, *Schröter*, 27.5.1886 (ZT), [HQ147775](#), [HQ147667](#), [HQ147713](#). *Centaurea triumfetti* All. subsp. *axillaris* (Willd.) Stef. & T.Georgiev, population 1, Bulgaria, Sofia City: Lozenska Mountain, in oak bushes above village of Pancharevo, *Ganchev*, 21.5.1953 (SO), [HQ147770](#), [HQ147662](#), –. *Centaurea triumfetti* All. subsp. *axillaris* (Willd.) Stef. & T.Georgiev, population 2, Bulgaria, Sofia Province: Lozenska Mountain, above Dolni Lozen village, ~1000 m, leg. *Stoyanov*, det. *Bancheva* 0715, 24.06.2007 (SOM), [HQ147765](#), [HQ147657](#), [HQ147705](#). *Centaurea triumfetti* All. subsp. *dominii* Dostál, Bulgaria, Plovdiv Province: Trojanska Mountain, Kozja stena, 42°47'18"N, 24°33'30"E, ~1650 m, rocky pasture, lime, *Röthlisberger*, 17.7.2001 (Z), [HQ147774](#), [HQ147666](#), [HQ147712](#). *Centaurea* cf. *triumfetti* All. subsp. *dominii* Dostál, Slovenia: Steinbrück, *Faler*, 28.5.1908 (ZA), [HQ147768](#), [HQ147660](#), –. *Centaurea triumfetti* All. subsp. *lugdunensis* (Jord.) Dostál, France: Department Ain, pass between Bellegarde and Plateau de Retort, ~1130 m west of the pass, in Narcissus meadows and especially on dry edges, *Landolt*, 15.6.1991 (ZT), [HQ147771](#), [HQ147663](#), [HQ147709](#). *Centaurea* cf. *triumfetti* All. subsp. *pirinensis* (Degen, Urum. & J. Wagner) Dostál, Bulgaria, Blagoevgrad Province: Pirin Mountain, stony slopes at the elevated parts of the “Bayuvi Dupli” Natural Reserve, leg. *Evstatieva*, det. *Vichodcevski*, 10.7.1968 (SO), [HQ147767](#), [HQ147659](#), [HQ147708](#). *Centaurea triumfetti* All. subsp. *semidecurrans* (Jord.) Dostál, Spain, Barcelona: Coll de Montnegre de Ponent (St. Iscle de Vallalta), 720 m, leg. *Pannon*, det. *Vilatersana*, June 2000 (BC), [HQ147773](#), [HQ147665](#), [HQ147711](#). *Centaurea triumfetti* All. subsp. *stricta* (Waldst. & Kit.) Dostál, Garcia-Jacas et al. 2006 as *C. mollis*, DQ319133, [HQ147706](#). *Centaurea triumfetti* All. subsp. *tanaitica* (Klokov) Dostál, Ukraine, Luhansk Province: Mt. Bryanka near Zamkovka, *Deripova*, 1.6.1980 (LE), [HQ147766](#), [HQ147658](#), [HQ147707](#). *Centaurea triumfetti* All. subsp. *triumfetti*, Croatia, Split-Dalmatia county: Mosor Mountain, Kotlenice, in the surroundings of village Vladovići, dry grassland and abandoned cultivated areas, *Vladović*, 19.5.1990 (ZA), [HQ147776](#), [HQ147668](#), [HQ147714](#). *Centaurea triumfetti* All. cf. subsp. *triumfetti*, Hungary: Budapest, Sashegy, *Wágner*, 5.6.1944 (LE), [HQ147777](#), [HQ147669](#), [HQ147715](#). *Centaurea tuberosa* Vis., population 2, Greece: Dramas, Mount Falakro, along road from Volakas to the ski center, 41°18'N, 24°01'E, 1000–1100 m, limestone, opening in *Fagus* forest, leg. *Strid* 30110 et al., det. *Landström*, 4.6.1989 (UPA), [HQ147780](#), [HQ147672](#), [HQ147718](#). *Centaurea tuberosa* Vis., population 1, Croatia, Split-Dalmatia county: Mosor Mountain, Kotlenice, in the surroundings of village Vladovići, abandoned cultivated areas, *Vladović*, 24.4.1988 (ZA), [HQ147781](#), [HQ147673](#), [HQ147719](#). *Centaurea velenovskyi* Adamović, Republic of Macedonia: north foothills of Mount Korab, south side of W-foothills of Kula e Ziberit Peak, ~6 km NNE of Mali i Korabit Peak, S-exposed lawns, 2000–2100 m, *Baltisberger* 12450 & *Baltisberger*, 26.7.1991 (ZT), [HQ147782](#), [HQ147674](#), –. *Centaurea woronowii* Borrm. ex Sosn., Georgia, Adjara: Zakavkazie, Batum region, Artvish distr., *Woronow* 6186, 12.7.1911 (LE), [HQ147783](#), [HQ147675](#), [HQ147720](#).

Literature Cited

- Ainouche A-K, RJ Bayer 1999 Phylogenetic relationships in *Lupinus* (Fabaceae: Papilionoideae) based on internal transcribed spacer sequences (ITS) of nuclear ribosomal DNA. *Am J Bot* 86:590–607.
- Andreasen K 2005 Implications of molecular systematic analyses on the conservation of rare and threatened taxa: contrasting examples from Malvaceae. *Conserv Genet* 6:399–412.
- Andreasen K, BG Baldwin 2001 Unequal evolutionary rates between annual and perennial lineages of checker mallows (Sidalcea, Malvaceae): evidence from 18S-26S rDNA internal and external transcribed spacers. *Mol Biol Evol* 18:936–944.
- Baldwin BG, S Markos 1998 Phylogenetic utility of the external transcribed spacer (ETS) of 18S-26S rDNA: congruence of ETS and ITS trees of *Calycadenia* (Compositae). *Mol Phylogenet Evol* 10:449–463.
- Bancheva S 1999 Biosystematic study of genus *Centaurea* L. (sect. *Cyanus* and *Lepteranthus*) in Bulgaria. PhD thesis. Department of Botany, Sofia University.
- Bancheva S, J Greilhuber 2006 Genome size in Bulgarian *Centaurea* s.l. (Asteraceae). *Plant Syst Evol* 257:95–117.
- Bancheva S, FM Raimondo 2003 Biosystematic studies of seven Balkan species from genus *Cyanus* (Compositae). *Bocconea* 16:507–527.
- Bentham G 1873 Compositae. Pages 162–533 in G Bentham, JD Hooker, eds. *Genera Plantarum*. Vol 2(1). Reeve, London.
- Boissier E 1875 Flora orientalis, sive enumeratio plantarum in Oriente a Graecia et Aegyptio ad Indiae fines hucusque observatarum. Vol 3. Georg, Genève.
- Clevinger JA, JL Panero 2000 Phylogenetic analysis of *Silphium* and subtribe Engelmanniinae (Asteraceae: Heliantheae) based on ITS and ETS sequence data *Am J Bot* 87:565–572.
- Cline J, JC Braman, HH Hogrefe 1996 PCR fidelity of pfu DNA polymerase and other thermostable DNA polymerases. *Nucleic Acids Res* 24:3546–3551.
- Cullings KW 1992 Design and testing of a plant-specific PCR primer for ecological and evolutionary studies. *Mol Ecol* 1:233–240.
- Czerepanov SK 2001 Subgenus *Cyanus*. Pages 385–415 in EG Bobrov, SK Czerepanov, eds. *Flora of the USSR*. Vol 28. Compositae, tribes Cynareae and Mutisieae. Smithsonian Institution, Washington, DC.
- de Candolle AP 1838 *Prodromus systematis naturalis regni vegetabilis*. Vol 6. Treuttel & Würtz, Paris.
- Dittrich M 1968 Morphologische Untersuchungen an den Früchten der Subtribus Cardueae-Centaureinae (Compositae). *Willdenowia* 5:67–107.
- Dostál J 1976 *Centaurea* L. Pages 254–301 in TG Tutin, VH Heywood, NA Burges, DM Moore, DH Valentine, SM Walters, DA Webb, eds. *Flora Europaea*. Vol 4. Cambridge University Press, London.
- Doyle JJ, JL Doyle 1987 A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* 19:11–15.
- Farris JS, M Källersjö, AG Kluge, C Bult 1995a Constructing a significance test for incongruence. *Syst Biol* 44:570–572.
- 1995b Testing significance of incongruence. *Cladistics* 10: 315–319.
- Felsenstein J 1985 Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783–791.
- Font M, N Garcia-Jacas, R Vilatersana, C Roquet, A Susanna 2009 Evolution and biogeography of *Centaurea* section *Acrocentron* inferred from nuclear and plastid DNA sequence analyses. *Ann Bot* 103:985–997.

- Font M, T Garnatje, N Garcia-Jacas, A Susanna 2002 Delineation and phylogeny of *Centaurea* sect. *Acrocentron* based on DNA sequences: a restoration of the genus *Crocodylium* and indirect evidence of introgression. *Plant Syst Evol* 234:15–26.
- Gamal-Eldin E, G Wagenitz 1991 *Centaurea* L. Pages 488–525 in A Strid, K Tan, eds. *Mountain flora of Greece*. Vol 2. Edinburgh University Press, Edinburgh.
- Garcia-Jacas N, PS Soltis, M Font, DE Soltis, R Vilatersana, A Susanna 2009 The polyploid series of *Centaurea toletana*: glacial migrations and introgression revealed by nrDNA and cpDNA sequence analyses. *Mol Phylogenet Evol* 52:377–394.
- Garcia-Jacas N, A Susanna, T Garnatje, R Vilatersana 2001 Generic delimitation and phylogeny of the subtribe Centaureinae (Asteraceae): a combined nuclear and chloroplast DNA analysis. *Ann Bot* 87:503–515.
- Greuter W 2003 The Euro+Med treatment of Cardueae (Compositae): generic concepts and required new names. *Willdenowia* 33:49–61.
- 2008 Med-checklist: a critical inventory of vascular plants of the circum-Mediterranean countries. Vol 2. Dicotyledones (Compositae). Conservatoire Botanique de la Ville de Genève, Geneva.
- Grimm GW, T Denk 2008 ITS evolution in *Platanus* (Platanaceae): homoeologues, pseudogenes and ancient hybridization. *Ann Bot* 101:403–419.
- Hall TA 1999 BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98.
- Hellwig FH 2004 Centaureinae (Asteraceae) in the Mediterranean: history of ecogeographical radiation. *Plant Syst Evol* 246:137–162.
- Hoffmann O 1894 Compositae. Pages 324–333 in A Engler, K Prantl, eds. *Die natürlichen Pflanzenfamilien*. Engelmann, Leipzig.
- Huelsenbeck JP, F Ronquist 2001 MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755.
- Kelch DG, BG Baldwin 2003 Phylogeny and ecological radiation of new world thistles (*Cirsium*, Cardueae-Compositae) based on ITS and ETS rDNA sequence data. *Mol Ecol* 12:141–151.
- Linder CR, LR Goertzen, B Vanden Heuvel, J Francisco-Ortega, RK Jansen 2000 The external transcribed spacer of the rDNA repeat: a new nuclear region for low-level taxonomic analysis of the Asteraceae and closely allied families. *Mol Phylogenet Evol* 14:285–303.
- Löser C, G Akaydin, FH Hellwig 2009 Incomplete lineage sorting among annuals and perennials in the genus *Cyanus*. Paper presented at Systematics 2009, 7th Biennial Conference of the Systematics Association, Leiden, August 10–14.
- Löser C, FH Hellwig 2007 Chloroplast DNA haplotype diversity in *Centaurea* sect. *Cyanus*. Paper presented at Botanical Congress 2007, Hamburg, September 3–7. University of Hamburg, Faculty for Mathematics, Informatics and Natural Sciences—Department of Biology.
- Maddison DR 1991 The discovery and importance of multiple islands of most-parsimonious trees. *Syst Zool* 40:315–328.
- Martins L, FH Hellwig 2005 Systematic position of the genera *Serratula* and *Klasea* within Centaureinae (Cardueae, Asteraceae) inferred from ETS and ITS sequence data and new combinations in *Klasea*. *Taxon* 54:632–638.
- Médail F, K Diadema 2009 Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J Biogeogr* 36:1333–1345.
- Meusel H, EJ Jäger 1992 Vergleichende Chorologie der zentral-europäischen Flora. Vol 3. Fischer, Jena.
- Miller P 1754 *The gardeners dictionary*. Abridged ed. 4. Privately published, London.
- Nickrent DL, KP Schuette, EM Starr 1994 A molecular phylogeny of *Arceuthobium* (Viscaceae) based on nuclear ribosomal DNA internal transcribed spacer sequences. *Am J Bot* 81:1149–1160.
- Nixon KC 2002 WinClada, version 1.00.08. Published by the author, Cornell University. <http://www.cladistics.com/>.
- Nylander JAA 2004 MrModeltest, version 2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University. <http://www.abc.se/~nylander/>.
- Olšovská K, M Perný, P Mártonfi, I Hodálova 2009 *Cyanus triumfetti* subsp. *triumfetti* (Compositae) does not occur in the western Carpathians and adjacent parts of Pannonia: karyological and morphological evidence. *Nordic J Bot* 27:21–36.
- Popp M, B Oxelman 2001 Inferring the history of the polyploid *Silene aegaea* (Caryophyllaceae) using plastic and homoeologous nuclear DNA sequences. *Mol Phylogenet Evol* 20:474–481.
- Rodríguez F, JL Oliver, A Marin, JR Medina 1990 The general stochastic model of nucleotide substitution. *J Theor Biol* 142:485–501.
- Ronquist F, JP Huelsenbeck 2003 MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574.
- Rösch M 1998 The history of crops and crop weeds in southwestern Germany from the Neolithic period to modern times, as shown by archaeobotanical evidence. *Veget Hist Archaeobot* 7:109–125.
- Smith SA, MJ Donoghue 2008 Rates of molecular evolution are linked to life history in flowering plants. *Science* 322:86–89.
- Soria-Hernanz DF, O Fiz-Palacios, JM Braverman, MB Hamilton 2008 Reconsidering the generation time hypothesis based on nuclear ribosomal ITS sequence comparisons in annual and perennial angiosperms. *BMC Evol Biol* 8:344.
- Stefanov B, T Gheorghiev 1931 Contribution to distinction of the species of genus *Centaurea* (L.)-sect. *Cyanus* (DC). *Rev Acad Bulg Sci* 44:133–193. (In Bulgarian.)
- Suárez-Santiago VN, MJ Salinas, N Garcia-Jacas, PS Soltis, DE Soltis, G Blanca 2007 Evolution by reticulation of the *Acrolophus* subgroup (*Centaurea* L., Compositae) in the occidental Mediterranean: origin and diversification of the section *Willkommia* Blanca. *Mol Phylogenet Evol* 43:156–172.
- Suc JP 1984 Origin and evolution of the Mediterranean vegetation and climate in Europe. *Nature* 307:429–432.
- Sun Y, DZ Skinner, GH Liang, SH Hulbert 1994 Phylogenetic analysis of *Sorghum* and related taxa using internal transcribed spacers of nuclear ribosomal DNA. *Theor Appl Genet* 89:26–32.
- Susanna A, N Garcia-Jacas 1996 El zurrón de las Centaureíneas. *Fontqueria* 44:25–36.
- 2009 The tribe Cardueae. Pages 293–313 in VA Funk, A Susanna, TF Stuessy, RJ Bayer, eds. *Systematics, evolution, and biogeography of the Compositae*. IAPT, Vienna.
- 2007 The tribe Cardueae. Pages 123–146 in J Kadereit, C Jeffrey, eds. *Flowering plants: eudicots: Asterales*. Vol VIII in K. Kubitzki, ed. *The families and genera of vascular plants*. Springer, Berlin.
- Susanna A, N Garcia-Jacas, O Hidalgo, R. Vilatersana, T Garnatje 2006 The Cardueae (Compositae) revisited: insights from ITS, trnL-trnF, and matK nuclear and chloroplast DNA analysis. *Ann Mo Bot Gard* 93:150–171.
- Swofford DL 2002 PAUP*. Phylogenetic analysis using parsimony (*and other methods), ver. 4.0b10. Sinauer, Sunderland, MA.
- Swofford DL, GJ Olsen 1990 Phylogeny reconstruction. Pages 411–501 in D Hillis, C Moritz, eds. *Molecular systematics*. Sinauer, Sunderland, MA.
- Urbatsch LE, RP Roberts, V Karaman 2003 Phylogenetic evaluation of *Xylothamia*, *Gundlachia*, and related genera (Asteraceae, Astereae) based on ETS and ITS nrDNA sequence data. *Am J Bot* 90:634–649.
- Vilatersana R, AK Brysting, C Brochmann 2007 Molecular evidence for hybrid origins of the invasive polyploids *Carthamus creticus* and *C. turkestanicus* (Cardueae, Asteraceae). *Mol Phylogenet Evol* 44:610–621.

- Wagenitz G 1955 Pollenmorphologie und Systematik in der Gattung *Centaurea* L. s.l. *Flora* 142:213–279.
- 1975 *Centaurea* L. Pages 465–585 in PH Davis, ed. *Flora of Turkey*. Vol 5. Edinburgh University Press, Edinburgh.
- 1980 *Centaurea*. Pages 313–419 in KH Rechinger, ed. *Flora Iranica*. Akademische, Graz.
- Wagenitz G, FH Hellwig 1996 Evolution of characters and phylogeny of the Centaureinae. Pages 491–510 in DJN Hind, HJ Beentje, eds. *Compositae: systematics*. Proceedings of the International Compositae Conference, Kew, 1994. Vol 1. *Compositae: systematics*. Royal Botanic Gardens, Kew.
- 2000 The genus *Psephellus* Cass. (*Compositae*, *Cardueae*) revisited with a broadened concept. *Willdenowia* 30:29–44.
- White TJ, TD Bruns, S Lee, JW Taylor 1990 Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Pages 315–322 in MA Innis, DH Gelfand, JJ Sninsky, TJ White, eds. *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, CA.
- Yamaji H, T Fukuda, J Yokoyama, J-H Pak, C-Z Zhou, C-S Yang, K Kondo, et al 2007 Reticulate evolution and phylogeography in *Asarum* sect. *Asiasarum* (Aristolochiaceae) documented in internal transcribed spacer sequences (ITS) of nuclear ribosomal DNA. *Mol Phylogenet Evol* 44:863–884.