

DRAFT

RECOVERY PLAN FOR THE BLUE WHALE

(Balaenoptera musculus)

REVISION



Office of Protected Resources
National Marine Fisheries Service
National Oceanic and Atmospheric Administration
October 2018



DRAFT

RECOVERY PLAN FOR THE BLUE WHALE

(Balaenoptera musculus)

REVISION

Prepared by:

Office of Protected Resources and West Coast Region
National Marine Fisheries Service

Approved: _____

Chris W. Oliver
Assistant Administrator for Fisheries
National Oceanic and Atmospheric Administration

Date: _____

PREFACE

Congress passed the Endangered Species Act of 1973 (16 U.S.C. 1531 *et. seq.*) (ESA) to provide a means to conserve the ecosystems upon which endangered and threatened species depend, to provide a program for the conservation of such endangered and threatened species, and to take such steps as may be appropriate to achieve the purposes of the treaties and conventions that conserve such species. The National Marine Fisheries Service (NMFS) and the U.S. Fish and Wildlife Service share responsibility for the administration of the Act. NMFS is responsible for most marine mammals including the blue whale (*Balaenoptera musculus*). This Recovery Plan was prepared at the request of the Assistant Administrator for Fisheries to promote the conservation of blue whales.

The goals and objectives of the Plan can be achieved only if a long-term commitment is made to support the actions recommended herein. Achievement of these goals and objectives will require the continued cooperation of the governments of the United States and other nations. Within the United States, the shared resources and cooperative involvement of federal, state, and local governments, industry, academia, nongovernmental organizations, and individuals will be required throughout the recovery period.

ACKNOWLEDGMENTS

This Revised Recovery Plan is based on the 1998 Recovery Plan for the Blue Whale, prepared largely by Randall Reeves, Phillip Clapham, Robert Brownell, Jr., and Gregory Silber. Updates and revisions contained in this Plan were provided by Monica DeAngelis, Nancy Young, and Amanda Keledjian.

For their technical assistance, editing, and review, we are grateful to Gregory Silber, Robert Brownell, Jr., Abigail Machernis, Allison Rosner, Therese Conant, Shannon Bettridge, Larissa Plants, Holly Wheeler, and Lynne Barre.

DISCLAIMER

Recovery plans delineate such reasonable actions as may be necessary, based upon the best scientific and commercial data available, for the conservation and survival of listed species. Plans are published by NMFS, sometimes prepared with the assistance of recovery teams, contractors, State agencies, and others. Recovery plans do not necessarily represent the views, official positions, or approval of any individuals or agencies involved in the plan formulation, other than NMFS. They represent the official position of NMFS only after they have been signed by the Assistant Administrator. Recovery plans are guidance and planning documents only; identification of an action to be implemented by any public or private party does not create a legal obligation beyond existing legal requirements. Nothing in this plan should be construed as a commitment or requirement that any Federal agency obligate or pay funds in any one fiscal year in excess of appropriations made by Congress for that fiscal year in contravention of the Anti-Deficiency Act, 31 U.S.C. 1341, or any other law or regulation. Approved recovery plans are subject to modification as dictated by new findings, changes in species status, or the completion of recovery actions.

Recommended Citation:

National Marine Fisheries Service. 2018. Draft Recovery Plan for the Blue Whale (*Balaenoptera musculus*) - Revision. National Marine Fisheries Service, Office of Protected Resources, Silver Spring, MD.

Recovery plans can be downloaded from the NMFS website:

<https://www.fisheries.noaa.gov/resources/documents>

Cover photo: Blue whale. Credit - Peter Duley, Northeast Fisheries Science Center, NOAA Fisheries, under permit #17355.

TABLE OF CONTENTS

PREFACE.....	i
ACKNOWLEDGMENTS	i
DISCLAIMER	ii
LIST OF TERMS AND ACRONYMS	v
EXECUTIVE SUMMARY	vi
I. BACKGROUND	1
A. Brief Overview	1
B. Species Description, Vocalizations, Taxonomy, Population Structure, and Natural History	1
B.1 Species Description.....	1
B.2 Hearing and Vocalizations	2
B.3 Taxonomy	3
B.4 Population Structure.....	4
B.5 Natural History.....	5
C. Zoogeography.....	7
D. Natural History – Northern Blue Whales (<i>B. m. musculus</i>) in the North Atlantic Ocean.....	8
D.1 Population Structure.....	8
D.2 Distribution and Habitat Use	8
D.3 Feeding and Prey Selection.....	10
D.4 Competition.....	11
D.5 Reproduction.....	11
D.6 Natural Mortality	11
D.7 Abundance and Trends	12
E. Natural History – Northern Blue Whales (<i>B. m. musculus</i>) in the North Pacific Ocean.....	13
E.1 Population Structure.....	13
E.2 Distribution and Habitat Use	14
E.3 Feeding and Prey Selection.....	16
E.4 Competition.....	17
E.5 Reproduction.....	17
E.6 Natural Mortality	17
E.7 Abundance and Trends	18
F. Natural History – Antarctic Blue Whales (<i>B. m. intermedia</i>)	19
F.1 Population Structure.....	19
F.2 Distribution and Habitat Use	20
F.3 Feeding and Prey Selection.....	21
F.4 Competition.....	21
F.5 Reproduction.....	22
F.6 Natural Mortality	22
F.7 Abundance and Trends	22
G. Natural History – Pygmy-type blue whales (<i>B. m. breviceauda</i> , <i>B. m.</i> <i>indica</i> , and <i>B. m. unnamed</i> subsp.).....	23
G.1 Population Structure.....	23

G.2	Distribution and Habitat Use	23
G.3	Feeding and Prey Selection.....	26
G.4	Competition.....	26
G.5	Reproduction.....	26
G.6	Natural Mortality	26
G.7	Abundance and Trends	27
H.	Potential Threats and Other Stressors	28
H.1	Potential Threats	31
H.2	Stressors Evaluated but Determined Not to be Threats	41
I.	Conservation Measures	46
II.	RECOVERY STRATEGY	48
III.	RECOVERY GOALS, OBJECTIVES, AND CRITERIA	48
A.	Goals.....	48
B.	Management Units	48
C.	Objectives and Criteria.....	49
C.1	Downlisting Objectives and Criteria.....	50
C.2	Delisting Objectives and Criteria.....	52
IV.	RECOVERY PROGRAM	55
A.	Recovery Action Outline.....	55
B.	Recovery Action Narrative.....	57
V.	IMPLEMENTATION SCHEDULE.....	67
VI.	LITERATURE CITED	75

LIST OF TERMS AND ACRONYMS

The following is a list of acronyms, abbreviations, and terms used throughout the Revised Recovery Plan.

BOEM	Bureau of Ocean Energy and Management
CI	confidence interval
CITES	Convention on International Trade in Endangered Species of Wild Fauna and Flora
CMS	Convention on the Conservation of Migratory Species of Wild Animals
CV	coefficient of variance
dB	decibels
DDT	dichlorodiphenyl trichloroethane
Delisting	removal from the list of Endangered and Threatened Wildlife and Plants
DNA	deoxyribonucleic acid
Downlisting	considered for reclassification from endangered to threatened under the ESA
EEZ	Exclusive Economic Zone
ENSO	El Niño Southern Oscillation
ESA	Endangered Species Act
ETP	Eastern tropical Pacific
FR	Federal Register
FY	fiscal year
Hz	hertz
ICRW	International Convention for the Regulation of Whaling
IUCN	International Union for Conservation of Nature
IWC	International Whaling Commission
kHz	kilohertz
LFA	low-frequency active (for sonar)
m	meters
MARU	marine acoustic recording unit
MFA	mid-frequency active (for sonar)
MMPA	Marine Mammal Protection Act
NMFS	National Marine Fisheries Service
NOAA	National Oceanic and Atmospheric Administration
PCB	polychlorinated biphenyl
PDO	Pacific (inter) Decadal Oscillation
POP	persistent organic pollutant
PTS	permanent threshold shift
SOWER	IWC's Southern Ocean Whale Ecosystem Research program
SURTASS	Surveillance Towed Array Sensor System
TTS	temporary threshold shift
UAS	unmanned aircraft system
USC	United States Code
USCG	United States Coast Guard

EXECUTIVE SUMMARY

Current Species Status: The blue whale (*Balaenoptera musculus*) was listed as “endangered” throughout its range under the precursor to the Endangered Species Act (ESA), the Endangered Species Conservation Act of 1969 (35 FR 8491; June 2, 1970), and remained on the list of threatened and endangered species after the passage of the ESA in 1973. Blue whales were subject to intensive commercial whaling, with over 380,000 blue whales taken in 1868-1978, mostly from Antarctic waters (Branch *et al.* 2008). The global population abundance is estimated to be 10,000-25,000 blue whales, or between 3-11% of the 1911 population size (Reilly *et al.* 2008). Although still depleted compared to historical abundance, blue whale populations around the world show signs of growth.

Blue whales are global in distribution. The subspecific taxonomy has not been fully resolved, but there are five currently recognized subspecies. *B. m. musculus* (Linnaeus, 1758) is the northern blue whale (North Atlantic and North Pacific Oceans); *B. m. intermedia* (Burmeister 1871) is the Antarctic blue whale; *B. m. brevicauda* (Ichihara, 1966) is the pygmy blue whale (Ichihara 1966, Rice 1977); and *B. m. indica* (Blyth 1859) is the northern Indian Ocean blue whale. There is some uncertainty in the characteristics that distinguish *B. m. indica* from *B. m. brevicauda* and additional research is needed to clarify the separation, but *B. m. indica* is currently considered a distinct subspecies (Perrin *et al.* 2009, Anderson *et al.* 2012, Thomas *et al.* 2016). Additionally, an unnamed subspecies off Chile in the southeastern Pacific Ocean was recently recognized by the Society for Marine Mammalogy’s Committee on Taxonomy (Branch *et al.* 2007a, Committee on Taxonomy 2016).

An improved understanding of subspecific taxonomy and population structure would facilitate management of the species.

Habitat Requirements and Limiting Factors: Although blue whales are listed as endangered under the ESA, it is not known whether and to what extent current threats are putting the globally listed species at risk of extinction. Nonetheless, we identify numerous *potential* threats. A potential threat, as used here, means a stressor that a) contributed to the species’ extinction risk, such as commercial whaling, and has the potential to do so again unless certain measures are taken or remain in place; or b) is known to be affecting one or more subspecies or populations, but more research is needed to understand the extent to which the stressor occurs or affects the globally listed entity. These include directed hunting, ship strikes, entanglement in marine debris and fishing gear, anthropogenic noise, and loss of prey base due to climate and ecosystem change. Other stressors were identified, but it was determined that there is currently no evidence that the effects (which may even include the loss of individual blue whales) have population-level consequences or are significant enough to contribute to the species’ extinction risk.

Recovery Strategy: Commercial whaling was the main cause of blue whales’ historical decline, and is not a current operative threat only because an international moratorium remains in place. Therefore, a primary strategy of this Revised Recovery Plan is to maintain the international ban on commercial hunting that was instituted in 1986. Additionally, this Plan provides a strategy to improve our understanding of how potential threats may be limiting blue whale recovery. Finally, this Plan provides a research strategy to obtain data necessary to determine blue whale taxonomy, population structure, distribution, and habitat, which can then inform estimation of

population abundance and trends. After the populations and their threats are more fully understood, this Plan will be modified to more specifically include actions to minimize any threats that are determined to be limiting recovery.

Because blue whales move freely across international borders, it would be ineffective to confine recovery efforts to U.S. waters, and this Plan stresses the importance of a multinational approach to management. As demonstrated by work on humpback whales involving a number of researchers from different countries (SPLASH (Calambokidis *et al.* 2008a); Year of the North Atlantic Humpback (Palsbøll *et al.* 1997, Smith *et al.* 1999)), considerably more information is gathered for management of whale populations when research is conducted on the basis of biological, rather than political divisions and through multilateral cooperation.

Management Units: For purposes of this Revised Recovery Plan, we define nine blue whale management units:

1. Northern subspecies (*B. m. musculus*) - North Atlantic population
2. Northern subspecies (*B. m. musculus*) - Eastern North Pacific population
3. Northern subspecies (*B. m. musculus*) - Western/Central North Pacific population
4. Northern Indian Ocean subspecies (*B. m. indica*)
5. Pygmy subspecies (*B. m. breviceauda*) - Madagascar population
6. Pygmy subspecies (*B. m. breviceauda*) - Western Australia/Indonesia population
7. Pygmy subspecies (*B. m. breviceauda*) - Eastern Australia/New Zealand population
8. Chilean subspecies (*B. m. unnamed subsp.*)
9. Antarctic subspecies (*B. m. intermedia*)

Blue whale subspecific taxonomy and population structure has not been fully resolved and is an area of active research. We identify at least one management unit for each of the five subspecies currently recognized by the Society for Marine Mammalogy. We also identify management units within two subspecies. In the Northern subspecies, *B. m. musculus*, we identify one North Atlantic and two North Pacific management units. In general, individuals from different ocean basins are unlikely to interbreed when mature, and if a basin population were extirpated, the area would likely not be recolonized in a time period that is meaningful for management purposes (Angliss *et al.* 2002). It is unclear whether blue whales in the eastern and western portions of the North Atlantic Ocean belong to the same population, but until separation is more strongly supported, for this Revised Recovery Plan we consider blue whales in the North Atlantic to comprise one management unit, based on the IWC blue whale stock definition and the current understanding that there is only one blue whale song type in the North Atlantic. The IWC also considers blue whales in the North Pacific to be one stock, but we define two management units there based on multiple lines of evidence (song types, length-frequency data, and movement data from satellite tags and photo-identification) indicating there are at least eastern and western/central populations. In the pygmy blue whale subspecies, *B. m. breviceauda*, the three management units correspond with “acoustic populations,” following recommendations of the IWC (IWC 2016b).

Despite the uncertainties, the delineation of these nine units reflects our current understanding of blue whale taxonomy and population structure and we consider them to be the appropriate units to conserve. We consider recovery of all nine units to be important for achieving geographic and

ecological representation of blue whales in the world's oceans, and to ensure conservation of the breadth of blue whale' genetic variability. If, based on additional research, blue whale subspecific taxonomy is revised, our understanding of population structure changes, or we learn more about whether recovery of all units is necessary for the long-term viability of the species, these management units and their associated species-level recovery criteria should be changed in a future revision of the plan.

Recovery Goals and Criteria: The goal of this Revised Recovery Plan is to promote the recovery of blue whales to the point at which they can be removed from the List of Endangered and Threatened Wildlife and Plants under the provisions of the ESA. The intermediate goal is to reach a sufficient recovery status to reclassify the species from endangered to threatened.

The two main objectives for blue whales are to 1) increase blue whale resiliency and ensure geographic and ecological representation by achieving sufficient and viable populations in all ocean basins and in each recognized subspecies, and 2) increase blue whale resiliency by managing or eliminating significant anthropogenic threats. The recovery criteria take two forms: 1) those that reflect the status of the species itself and 2) those that indicate effective management or elimination of threats. These criteria would apply to the species throughout its range, but the criteria also include specific targets for each of the nine management units to support the objectives of species' viability (e.g., resiliency, redundancy, and representation).

Population viability analyses (PVAs) or other quantitative assessments for predicting a species' or population's future status can be useful in evaluating extinction risk. PVAs have been used in some recent marine mammal status reviews (e.g., (Krahn *et al.* 2004, Oleson *et al.* 2010), but not in others (e.g., (Boveng *et al.* 2009, Boveng *et al.* 2013, Bettridge *et al.* 2015), and are a component of several other large whale recovery plans (e.g., (NMFS 2010a, NMFS 2010b, NMFS 2011, NMFS 2013). Although not required to meet the criteria below, should sufficient data become available, a quantitative PVA demonstrating a low probability of extinction over a reasonable timeframe (e.g., scaled to blue whale generation time, (Taylor *et al.* 2007) could be used to further support downlisting or delisting decisions.

Downlisting Criteria:

Blue whales may be considered for reclassifying from endangered to threatened when all of the following criteria are met.

In each of nine blue whale management units:

1. The minimum abundance is:
 - a. North Atlantic: 2,000 whales
 - b. Eastern North Pacific: 2,000 whales
 - c. Western/Central North Pacific: 2,000 whales
 - d. Northern Indian Ocean: 500 whales
 - e. Madagascar: 2,000 whales
 - f. Western Australia/Indonesia: 2,000 whales
 - g. New Zealand: 500 whales
 - h. Chilean: 2,000 whales

- i. Antarctic: 2,000 whales
2. The trend in abundance is stable or increasing over the most recent 30-year period assessed.
3. Anthropogenic threats have been identified and demonstrably minimized. Specifically, the factors in 4(a)(1) of the ESA described below have been addressed:

Factor A: The present or threatened destruction, modification, or curtailment of a species' habitat or range.

- Effects of anthropogenic noise, entanglement in marine debris and fishing gear, and reduced prey abundance due to climate change have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

Factor B: Overutilization for commercial, recreational, or educational purposes.

- The international ban on commercial hunting has been maintained.
- Any subsistence or scientific hunting that has the potential to overutilize the species is restricted to levels that are sustainable, precautionary, and in accordance with the advice of the IWC's Scientific Committee.

Factor C: Disease or Predation.

- There is no information at this time indicating disease or predation is a threat to blue whale recovery.

Factor D: The inadequacy of existing regulatory mechanisms.

- (Hunting is addressed under Factor B.)

Factor E: Other natural or manmade factors affecting its continued existence.

- Ship strikes have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.
- Entanglement with fishing gear continue to be investigated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

Delisting Criteria:

Blue whales may be considered for removal from the list of Endangered and Threatened Wildlife and Plants under provisions of the ESA when all of the following criteria are met.

In each of nine blue whale management units:

1. The minimum abundance is:
 - a. North Atlantic: 2,500 whales
 - b. Eastern North Pacific: 2,500 whales
 - c. Western/Central North Pacific: 2,500 whales
 - d. Northern Indian Ocean: 1,000 whales
 - e. Madagascar: 2,500 whales
 - f. Western Australia/Indonesia: 2,500 whales
 - g. New Zealand: 1,000 whales

- h. Chilean: 2,500 whales
 - i. Antarctic: 2,500 whales
2. The trend in abundance is stable or increasing over the most recent 30-year period assessed.
 3. Anthropogenic threats have been identified and demonstrably minimized or eliminated, and there is information indicating they are not contributing to the species being in danger of extinction within the foreseeable future throughout all or a significant portion of its range. Specifically, the factors in 4(a)(1) of the ESA described below have been addressed:

Factor A: The present or threatened destruction, modification, or curtailment of a species' habitat or range.

- Effects of anthropogenic noise, entanglement in marine debris and fishing gear, and reduced prey abundance due to climate change have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

Factor B: Overutilization for commercial, recreational, or educational purposes.

- The international ban on commercial hunting has been maintained.
- Any subsistence or scientific hunting that has the potential to overutilize the species is restricted to levels that are sustainable, precautionary, and in accordance with the advice of the IWC's Scientific Committee.

Factor C: Disease or Predation.

- There is no information at this time indicating disease or predation is a threat to blue whale recovery.

Factor D: The inadequacy of existing regulatory mechanisms.

- (Hunting is addressed under Factor B.)

Factor E: Other natural or manmade factors affecting its continued existence.

- Ship strikes have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.
- Entanglement with fishing gear has been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

Estimated Cost of Recovery Actions for the First Five Fiscal Years (FY) (estimates are in thousands of dollars):

Action	FY1	FY2	FY3	FY4	FY5	Total¹
Coordinate State, Federal, and international measures to maintain international regulation of whaling for blue whales	a	a	a	a	a	a
Determine blue whale taxonomy, population structure, occurrence, distribution, and range	400	400	400	400	400	2,000
Estimate population size and monitor trends in abundance	530	530	530	530	530	2,650
Identify, characterize, protect, and monitor habitat important to blue whale populations	400	400	400	400	400	2,000
Investigate human-caused threats and, should they be determined to be impeding blue whale recovery, take steps to minimize their occurrence and severity	730 ^b	730 ^b	700 ^b	400 ^b	400 ^b	2,960^b
Maximize efforts to acquire scientific information from dead, stranded, and entangled or entrapped blue whales	30	30	30	30	30	150^b
Total	2,090_b	2,090_b	2,060_b	1,760_b	1,760_b	9,760_b

^a No cost associated, NMFS staff time only.

^b Given uncertainty in potential threats and actions required to address them, there are additional costs that cannot be determined.

Estimated Cost of Recovery Actions (First 5 Fiscal Years): \$9.76 million plus additional costs that cannot be determined. A total estimate to recovery beyond the first five fiscal years is not practicable given the uncertainty in the potential threats, any actions that might be required to address the potential threats, and the length of time beyond the first five fiscal years that will be necessary to conduct research to fill critical information gaps and evaluate effectiveness of threat reduction.

Anticipated Date of Recovery: It is not possible to predict the time and cost to recovery with the current information because of the uncertainty of potential threats. Thus, an estimate of the time required and the cost to carry out those actions needed to achieve the plan’s goal and to achieve intermediate steps (beyond 5 years) is not practicable. Conducting research necessary to support conclusions regarding the

¹ The totals in this table differ slightly from those in Table 2 in Section V Implementation Schedule. This table provides the estimated cost of recovery actions for the first five fiscal years, while Table 2 additionally includes minimum estimates of the cost for discrete or ongoing recovery actions that are likely to extend into FY6 and beyond. See the footnotes to Table 2 for more information.

impact of the potential threats to blue whale populations, and developing, implementing, and evaluating the effectiveness of recovery actions to reduce threats or potential threats (*i.e.*, to meet recovery criterion 3) may take decades. The minimum data needed to satisfy criteria 1 and 2 for downlisting or delisting are population structure studies and abundance surveys, which will also take decades, given the species' global distribution and the need to evaluate the abundance trend across a minimum of 30 years. If the necessary research is undertaken and demonstrates that the abundance and trend criteria have been met, and potential threats are evaluated and, as necessary, minimized or eliminated, it might be feasible to downlist or delist blue whales in 30 years. In the future, as more information is obtained, it may be possible to develop estimates for the full time to recovery and its expense.

I. BACKGROUND

A. Brief Overview

The blue whale, *Balaenoptera musculus* (Linnaeus 1758), was listed as endangered throughout its range under the precursor to the Endangered Species Act (ESA), the Endangered Species Conservation Act of 1969, and remained on the list of threatened and endangered species after the passage of the ESA in 1973 (35 FR 8491; June 2, 1970). Blue whales are the largest living animal on earth. The species experienced intensive whaling throughout the 20th century, with over 380,000 blue whales taken in 1868-1978, mostly from Antarctic waters (Branch *et al.* 2008). The global population abundance is estimated to be 10,000-25,000 blue whales, or between 3-11% of the 1911 population size (Reilly *et al.* 2008).

Although the species is globally listed, five blue whale subspecies are currently recognized (Committee on Taxonomy 2016). Blue whales generally make seasonal migrations between feeding and breeding locations, with occurrence often linked to prey aggregations. Underwater recordings of blue whale vocalizations have been used to show changing distribution and trends in local abundance at feeding locations around the world. Little is known about interspecific competition and sources of natural mortality. Though still depleted compared to historical abundance, blue whale populations around the world show signs of growth.

Below is a general description of the listed species with information on vocalizations, taxonomy, and population structure for blue whales worldwide. In addition there is general information on several natural history elements which applies to all blue whales. Where there is specific information on natural history for blue whale by subspecies or in particular regions, it is included in the more detailed sections below. Sections I.D-I.G include detailed regional information including distribution, abundance, and trends where available for blue whales in the North Atlantic, North Pacific, and Antarctic, as well as for the pygmy-type blue whales.

B. Species Description, Vocalizations, Taxonomy, Population Structure, and Natural History

B.1 Species Description

The blue whale is a cosmopolitan species of baleen whale (Gambell 1979, Yochem & Leatherwood 1985, Mead & Brownell 1993) found in all oceans except the Arctic (although they are regularly sighted near the ice edge in the North Atlantic), but largely absent from some regional areas such as the Mediterranean, Okhotsk, and Bering Seas north into the Arctic Ocean (few sightings have been observed in the Bering Sea/Sea of Okhotsk). The species is the largest animal ever to have lived: adults can reach 32.6 meters (m) in length and weigh more than 150,000 kg. Blue whales in the Northern Hemisphere are generally smaller than those in the Southern Ocean, with maximum body length in the North Atlantic and North Pacific Oceans being approximately 27 m (True 1904, Reeves *et al.* 1985). Pygmy blue whales, a subspecies of blue whales, may only reach 21 m in length when mature. As is true of other baleen whales, female blue whales are slightly larger than males (Ralls 1976). At birth, blue whales can be 8 m long and weigh 3 tons.

Blue whales are long-bodied and slender, with proportionally smaller dorsal fins compared to

other baleen whales. Their gray-blue skin color appears light blue when viewed through the water and has a unique mottling pattern that has been used to identify individuals. Viewed from above, the blue whale has a broad, flat rostrum, with a single ridge that runs from the splashguard of the blowholes to the tip of the rostrum. The blue whale may have as many as 400 relatively short (0.5 m) plates of typically black baleen on each side of the upper jaw. When a blue whale is feeding, its pleated throat expands to accommodate the enormous intake of seawater and food. As the water is expelled and the filtered prey is swallowed, the body returns to its characteristically slender shape.

Blue whale diet is composed almost exclusively of krill (euphausiids) (Yochem & Leatherwood 1985, Reilly *et al.* 2008). They feed both at the surface and at depths of more than 100 m, following their prey's diel vertical migration through the water column (Sears & Perrin 2009).

B.2 Hearing and Vocalizations

Marine mammal hearing has been reviewed by several authors, notably Popper (1980a, 1980b), Schusterman (1981), Ridgeway (1983), Watkins and Wartzok (1985), Moore and Schusterman (1987), Au (1993), Richardson (1995), and Wartzok and Ketten (1999). Marine mammal hearing has been studied through behavioral or electrophysiological tests and predictions based on inner ear morphology. No direct measurements of the hearing sensitivity of baleen whales (mysticetes) are available. Thus, hearing predictions for mysticetes are based on other methods including anatomical studies and modeling (Houser *et al.* 2001, Parks 2007, Ketten & Mountain 2012, Ketten & Mountain 2014, Cranford & Krysl 2015); vocalizations (Richardson *et al.* 1995, Wartzok & Ketten 1999, Au & Hastings 2008); and behavioral responses to sound (Dahlheim & Ljungblad 1990, Reichmuth 2007). Mysticetes hear at low frequencies and differ from other cetaceans and pinnipeds that hear higher frequency sounds (Southall *et al.* 2007). While little information exists specific to blue whales, recent studies of hearing in other baleen whales may be applicable.

Baleen whales' sensitivity to low-frequency sound has been inferred from observed vocalization frequencies, observed reactions to playback of sounds, and anatomical analyses of the auditory system. The anatomy of the baleen whale inner ear appears well adapted for detecting a wide range of low-frequency sounds (Ketten 1992b, Ketten 1992a, Ketten 1994, Yamato *et al.* 2008). Baleen whale calls are predominantly at low frequencies and their hearing is presumably best at detecting frequency ranges analogous to those in which they vocalize (Richardson *et al.* 1995). Based on these data, Southall *et al.* (2007) estimated the low-frequency hearing range of cetaceans to extend from approximately 7 Hz to 22 kHz. Watkins (1986) reported a variety of mysticete species responding to sounds up to 28 kHz; Au *et al.* (2006) reported humpback whales songs having harmonics that extend beyond 24 kHz; and Frankel (2005) Lucifredi and Stein (2007) reported gray whales potentially responding to sounds above 22 kHz. Thus, the auditory system of baleen whales almost certainly lacks the acuity in high frequency ranges that are attributed to the smaller toothed whales. However, auditory sensitivity in at least some large whale species presumably extends into frequencies higher than the upper frequency ranges of their calls.

Blue whale calls are among the loudest and lowest frequency sounds made by any animal and consist of long-duration, low-frequency pulses (known as type A) and tonal calls (type B),

upswept tones that precede type B calls (type C), moderate-duration downswept tones (type D), and variable amplitude-modulated and frequency-modulated sounds (Thompson *et al.* 1996, Aroyan *et al.* 2000, Thode *et al.* 2000, McDonald *et al.* 2001). Stafford *et al.* (1998) reported that the fundamental frequency for blue whale vocalizations ranged from 8 to 25 Hz. Blue whale vocalizations might vary by region, season, behavior, and time of day. Similar to other baleen whales, blue whales can produce long patterned sets of sounds referred to as songs. Ten song types have been described for blue whales around the world that have remained largely stable in the number of units and phrasing for decades, and might be used to distinguish populations in the absence of definitive genetic data (McDonald *et al.* 2006b, McDonald *et al.* 2009).

In studying the eastern North Pacific blue whales off southern California, researchers found that only males produced type AB calls, with songs (stereotypic, repeated AB call sequences) produced by lone males and singular AB calls generally produced by whales in pairs, which suggests their importance in reproductive activities (Oleson *et al.* 2007a, Calambokidis *et al.* 2010). Type D calls were produced by both males and females during shallow, daytime dives, and have been associated with social interactions between deeper foraging dives (Oleson *et al.* 2007a, Oleson *et al.* 2007b).

Seven of the ten blue whale song types identified worldwide have shifted linearly downward in tonal frequency, though at different rates (Nieukirk *et al.* 2005, McDonald *et al.* 2009). In one of the best documented blue whale song types, from the eastern North Pacific Ocean, the frequency of the song's tonal section is 31% lower than it was in the early 1960s (Nieukirk *et al.* 2005, McDonald *et al.* 2009). McDonald *et al.* (2009) discussed several hypotheses for the observed pattern, and suggested it might be related to increasing population density as blue whales recover from whaling, and related effects on sexual selection pressure. Specifically, females selecting mates may prefer males producing lower frequency songs because it indicates larger body size; the greater number of singing males competing for mates may increase this selection pressure. However, there is a tradeoff: a lower frequency corresponds with a lower amplitude (quieter) song, which has a shorter propagation distance. At higher population densities, though, there is less need to communicate with other blue whales over long distances, so males can afford to sing at a lower frequency despite the decreased detection distance (McDonald *et al.* 2009).

B.3 Taxonomy

The subspecific taxonomy of blue whales is an area of continued research. Studies of intraspecific variability and life history characteristics have led to the recognition of five blue whale subspecies to date. *B. m. musculus* (Linnaeus, 1758) is the northern blue whale (North Atlantic and North Pacific Oceans); *B. m. intermedia* (Burmeister, 1871) is the Antarctic blue whale, sometimes referred to as the “true” blue whale; *B. m. breviceuda* (Ichihara 1966) is the pygmy blue whale, generally occurring in the sub-Antarctic southern Indian Ocean and the southwestern Pacific Ocean (Ichihara 1966, Rice 1977); *B. m. indica* (Blyth, 1859) is the northern Indian Ocean blue whale; and there is a recently recognized, unnamed subspecies that generally occurs off Chile in the southeastern Pacific Ocean (Branch *et al.* 2007a, Committee on Taxonomy 2016).

There is some uncertainty in the characteristics that distinguish *B. m. indica* from *B. m. breviceuda* and additional research is needed to clarify the separation, but *B. m. indica* is

currently considered a distinct subspecies (Perrin *et al.* 2009, Anderson *et al.* 2012, Committee on Taxonomy 2016). *B. m. indica* has a breeding season that is asynchronous with Southern Hemisphere blue whales (Mikhalev 2000), a distinct call type (Alling *et al.* 1991, McDonald *et al.* 2006b), and slightly smaller total length at maturity compared to *B. m. breviceauda* (Branch & Mikhalev 2008).

The unnamed subspecies in the southeastern Pacific, recently recognized by the Society for Marine Mammalogy's Committee on Taxonomy, is considered distinct from the other subspecies in the Southern Hemisphere based on geographic separation (latitudinal from Antarctic blue whales and longitudinal from pygmy blue whales), a difference in the mean length of mature females, a unique call type, and significant genetic differentiation (Branch *et al.* 2007a). LeDuc *et al.* (2016) reported significant mitochondrial and nuclear DNA differentiation between eastern South Pacific and eastern North Pacific blue whales, but considered them to be modest and less than the differences between eastern South Pacific versus Indian Ocean or Antarctic blue whales.

B.4 Population Structure

In the United States blue whales are managed under three constructs: the Marine Mammal Protection Act (MMPA), the ESA, and the International Convention for the Regulation of Whaling (ICRW), all with different objectives and, therefore, different terminology for population structure. The goal of the MMPA is to protect marine mammal species by maintaining marine mammal population “stocks” as functioning elements of their ecosystem. The International Whaling Commission (IWC), established under the terms of the ICRW, manages whales with a goal of maintaining healthy stocks while authorizing hunts to meet aboriginal subsistence needs (and potentially commercial catches), and scientific research and related purposes. The ESA governs the taking of species that have an elevated risk of extinction to ensure that effects of threats are restricted to levels that would allow recovery of the species to the point where it is no longer threatened or endangered (Eagle *et al.* 2008).

In this document, we use the term “stocks” in the context of MMPA or IWC stocks, and use the more generic term “populations” when referring to species subunits. NMFS guidelines define “stock” under the MMPA as a “demographically independent biological population” where internal dynamics (births and deaths) have a larger impact on population dynamics than external dynamics (immigration and emigration) (NMFS 2016). The IWC has adopted two definitions of stock: biological stocks (based on genetic separation) and management stocks (based on functional population units) (Donovan 1991).

The IWC recognizes three blue whale stocks corresponding to three major ocean basins (North Atlantic Ocean, North Pacific Ocean, and Southern Hemisphere), and six management areas in the Southern Hemisphere (Donovan 1991). The IWC's Southern Hemisphere baleen whale management Areas, each encompassing 50° to 70° longitude from the South Pole to the equator, were officially delineated ahead of the 1974/75 season, and are still used today (Donovan 1991). The IWC is preparing for a comprehensive assessment of Southern Hemisphere pygmy blue whales and evaluating the feasibility of conducting assessment of North Pacific blue whales (IWC 2017); the outcomes of these assessments may inform the recognition of blue whale stocks and/or populations. For example, after reviewing relevant scientific literature, in 2016 the IWC Scientific Committee identified six “acoustic populations” in the Southern Hemisphere and

northern Indian Ocean (Antarctic, Chilean, Northern Indian Ocean, New Zealand, Indonesia/western Australia, and southwest Indian Ocean) (IWC 2016a). In the Northern Hemisphere, NMFS recognizes three stocks in United States waters: the western North Atlantic Ocean and the central and eastern North Pacific Ocean stocks (Eagle *et al.* 2008, Waring *et al.* 2010, Carretta *et al.* 2015).

The sub-specific taxonomy of blue whales (see Section I.B.3 above) is the subject of continuing work. Researchers are also working to define and distinguish regional populations or stocks by evaluating differences in genetics, acoustics, morphology, and movements. For example:

- *Genetics.* Conway (2005) conducted a global comparison of nuclear DNA and found a significant genetic separation between Antarctic and southern Indian Ocean samples and moderate substructure suggesting separation between blue whales in the western North Atlantic, eastern North Pacific, southern Indian, and Southern Oceans. LeDuc *et al.* (2007) evaluated nuclear and mitochondrial DNA of blue whale subspecies and populations in the Southern Hemisphere, and found, among other things, significant differentiation between populations of pygmy blue whales. Attard *et al.* (2016) found evidence of three sympatric populations of blue whales on Antarctic feeding grounds which likely have separate (but unknown) breeding grounds, although the IWC Scientific Committee found the evidence for these populations inconclusive (IWC 2016a). There is also evidence of hybridization between fin and blue whales (Berube & Aguilar 1998, NAMMCO 2014) and between pygmy and Antarctic blue whales (Attard *et al.* 2012).
- *Acoustics.* McDonald *et al.* (2006b) used temporally stable blue whale songs to identify regional populations around the world, and suggested that any differences not reflected in genetic analysis represent more recent patterns in movement and distribution. Stafford (2003) and Stafford *et al.* (2001) found that blue whales in the North Pacific Ocean produce two distinct, stereotypic calls that have been termed the northwestern and northeastern call types, and it has been proposed that these represent two distinct populations with some degree of geographic overlap.
- *Morphological differences.* Gilpatrick and Perryman (2008) used whaling records and aerial photographs to examine morphological differences between regional populations to inform subspecies and stock distinctions. They found that blue whales from the eastern North Pacific Ocean are on average 2 meters shorter in length at maturity than those in the central and western areas of the North Pacific Ocean (Gilpatrick & Perryman 2008).
- *Movements.* Satellite tagging and photo-identification research demonstrated that the eastern tropical Pacific Ocean is a migratory destination for blue whales that were tagged and/or photo-identified off California (Mate *et al.* 1999, Calambokidis *et al.* 2009a, Irvine *et al.* 2014).

More information about population structure is described in the region-specific sections of this document (Sections I.D.1, I.E.1, I.F.1, and I.G.1 below).

B.5 Natural History

B.5.1 Feeding and Prey Selection

Blue whales often forage in small groups at depths of approximately 100 m, although some

whales forage around 250-300 m (Calambokidis *et al.* 2008b). They are considered lunge-feeders and filter large quantities of seawater (as much as 70% of their body weight with every feeding lunge) (Pivorunas 1979) to capture krill in their baleen. Sustaining swim speeds with the increased drag required by this feeding behavior is energetically expensive and may explain typical dive times (approximately 10 minutes) being much shorter than would otherwise be predicted based on body size (Croll *et al.* 2001b). In support of this hypothesis, blue whales spend more time recovering from foraging dives compared to shallower non-foraging dives (Acevedo-Gutierrez *et al.* 2002). However, Goldbogen *et al.* (2011) suggested that a large body size enables relatively high feeding efficiency because blue whales have a slow metabolism and consume large quantities of rapidly filtered prey with each feeding event. Blue whales are able to consume and assimilate as much as 90 times the amount of energy expended in obtaining their prey (Goldbogen *et al.* 2011, Knight 2011).

B.5.2 Reproduction

Most blue whale reproductive activity, including birthing and mating, takes place in winter and is marked by annual migratory cycles, but the breeding grounds for some populations are still unknown. Blue whales typically reach sexual maturity beginning at an average length of 23 m, which corresponds to between five and ten years of age in both males and females, though these estimates vary across regions and subspecies (Mizroch *et al.* 1984, Yochem & Leatherwood 1985, Branch & Mikhalev 2008). Recent testosterone profiles suggest that male blue whales reach sexual maturity at approximately 10 years of age (Trumble *et al.* 2013). Little is known about the blue whale mating system, but anatomical characteristics (Brownell & Ralls 1986) and behavioral observations during the breeding season (Sears *et al.* 2013) suggest a polygynous, antagonistic male-male competition strategy. Blue whales are also known occasionally to hybridize with other large whale species, such as fin and humpback whales (Berube & Aguilar 1998, Reeves *et al.* 2002). Calf production is thought to be determined by a combination of age and food availability, with a gestation period lasting approximately ten to twelve months (Christensen *et al.* 1992). Pregnant females have been observed to consume 4% of their body weight daily (Sergeant 1969), amounting to 60% of their overall body weight throughout summer foraging periods (Lockyer 1984). There are no observations of the birth of a calf (Sears *et al.* 2013). Weaning likely occurs six to seven months after birth en route to, or after whales arrive at, summer feeding areas (Reeves *et al.* 1998). The average calving interval for most females is two to three years and depends on the cow's ability to replenish food stores depleted during the lactation period (Lockyer 1984).

B.5.3 Competition

Nearly all baleen whale species that are sympatric with the blue whale eat euphausiids to some extent and are, therefore, potential competitors (Nemoto 1973). However, there is currently little or no direct evidence for interspecific competition involving blue whales (Clapham & Brownell 1996). In some locations, researchers have found evidence of resource partitioning among sympatric baleen whales, including separation in space and/or time and selection of prey species (Doniol-Valcroze 2008, Fossette *et al.* 2017). It seems unlikely that resource competition would be an important factor in preventing blue whale recovery. In addition, blue whales are capable of traveling great distances, which should enable them to take advantage of prey concentrations over very large areas and exploit prey even as oceanographic conditions vary.

B.5.4 Natural Mortality

At maturity, blue whales have little vulnerability to natural predators; however, young or ailing blue whales are likely vulnerable to shark and killer whale predation. Calves are most vulnerable to these attacks and most encounters are likely fatal, based on observations of killer whale attacks with other baleen whale calves. In some locations, ice entrapment is another source of mortality.

C. Zoogeography

As noted above, blue whales are a cosmopolitan species inhabiting nearly all major oceans of the world (Figure 1). Although populations of blue whales were severely depleted by whaling, this exploitation has not resulted in a major change in their distribution, although some areas with historical catches show an apparent lack of recovery, including parts of the eastern North Atlantic Ocean (Christensen *et al.* 1992), western North Pacific Ocean (Forney & Brownell 1996), and eastern South Atlantic Ocean (Branch *et al.* 2007b). In general, blue whale distribution is driven largely by food requirements. Blue whales avoid the oligotrophic central gyres in the Indian, Pacific, and Atlantic Oceans, but are common where phytoplankton densities are high, and where there are dynamic oceanographic processes such as upwelling and oceanic fronts (Branch *et al.* 2007b). Blue whales generally migrate seasonally toward the polar regions in spring to areas with abundant summer zooplankton, and toward the subtropics in the fall to avoid ice entrapment in some areas and engage in reproductive activities in warmer waters. However, blue whales have been acoustically detected year-round in some locations, suggesting resident populations or portions of populations that do not migrate each year (Stafford *et al.* 1999a, Širović *et al.* 2004, Širović *et al.* 2009, Samaran *et al.* 2010, Stafford *et al.* 2011, Samaran *et al.* 2013).



Figure 1. Distribution of blue whales spanning all oceans except the Arctic Ocean and Mediterranean Sea (adapted from <http://maps.iucnredlist.org/map.html?id=2477>). The yellow area represents the species' approximate range.

D. Natural History – Northern Blue Whales (*B. m. musculus*) in the North Atlantic Ocean

D.1 Population Structure

Blue whales in the North Atlantic Ocean may exist as eastern and western populations, with regional feeding subgroups (Christensen 1955, Jonsgard 1955, Sigurjónsson & Gunnlaugsson 1990). Blue whales in the western North Atlantic Ocean generally extend from the Arctic to at least mid-latitude waters, but are most frequently sighted in the waters off eastern Canada, with the majority of recent records in the Gulf of St. Lawrence (Waring *et al.* 2010). Photo identification research has shown that whales inhabiting the Gulf of St. Lawrence and waters off eastern Canada, New England, and Greenland belong to one population, with no known matches to the eastern population. Blue whales identified off West Greenland, the Scotian Shelf, and the Gulf of Maine were observed in subsequent years in the Gulf of St. Lawrence (Wenzel *et al.* 1988, Sears 1990). Based on whaling records, Jonsgard (1955) argued that the blue whales hunted along the coasts of Newfoundland and Labrador belonged to the same population as those hunted in Davis Strait as far north as Disko Island, Greenland. Blue whales in the eastern North Atlantic Ocean are centered around Iceland and have been observed passing through the Azores to northwest Africa (CETAP 1982, Wenzel *et al.* 1988, Sears & Calambokidis 2002, Sears & Larsen 2002, Sears *et al.* 2005, Visser *et al.* 2011).

Clark (1994) found that blue whale songs do not vary significantly in different regions of the North Atlantic Ocean, suggesting that these blue whales may range over the entire ocean basin and comprise a single panmictic population. Other researchers noted subtle differences in the cadence and composition of song phrases between blue whales in the Gulf of St. Lawrence and the rest of the North Atlantic Ocean (Berchok *et al.* 2006).

In some sections below, we discuss information specific to blue whales in the western and eastern North Atlantic Ocean, but recognize that there is uncertainty in whether they represent two separate units. Genetic analyses are needed to further elucidate population structure in the North Atlantic Ocean.

D.2 Distribution and Habitat Use

Blue whale distribution in the North Atlantic Ocean extends from largely temperate areas north to Baffin Bay and the Greenland Sea (Jonsgard 1955, Yochem & Leatherwood 1985, Sears & Larsen 2002), with whales primarily sighted in the northwestern portion of the Gulf of St. Lawrence (Lesage *et al.* 2007, Comtois *et al.* 2010). A stranding at Ocean City, Maryland, in 1891 (True 1904) is the southernmost confirmed record along the U.S. east coast. There are only two records from the Gulf of Mexico, a 1924 stranding near Sabine Pass, Louisiana and a 1940 stranding between Freeport and San Luis Pass, Texas. However, the species identification for both strandings is considered to be questionable, so blue whale occurrence in the Gulf of Mexico remains unconfirmed (Davis & Schmidly 1994 (rev)).

In general, blue whales migrate seasonally between feeding and calving areas. The seasonal and interannual distribution of blue whales is strongly associated with both the static and dynamic oceanographic features that result in krill aggregations (Gendron 1990, Reilly & Thayer 1990,

Del-Angel-Rodriguez 1997, Croll *et al.* 1998, Fiedler *et al.* 1998, Benson *et al.* 2002, Hamazaki 2002, Croll *et al.* 2005, Matteson *et al.* 2010). Blue whales depend on a critical density of prey to feed efficiently, and thus seek out highly productive areas near upwelling zones and thermal fronts where krill abundance is greatest (Doniol-Valcroze *et al.* 2007). Since krill, the primary prey for blue whales, has relatively limited capacity for horizontal displacement, its distribution is strongly linked to areas of phytoplankton blooms and enhanced biomass retention.

Much of the recent research on blue whale distribution has focused on correlating seasonal survey observations with whaling data and whale vocalizations to elucidate potential feeding and breeding areas. This work often corroborates modeling predictions based on oceanographic features such as those noted above. For example, Clark and Gagnon (2004) found that blue whale singers strongly prefer shelf breaks, seamounts, and other highly productive areas throughout the year.

Western North Atlantic

In the 1960s, whalers regularly observed blue whales on the Scotian Shelf throughout summer and fall (Sutcliffe & Brodie 1977), with summer catches made off the northern Newfoundland and southern Labrador coasts, including the Strait of Belle Isle (Sergeant 1953, Jonsgard 1955, Sergeant 1966). In the Gulf of St. Lawrence, blue whales have been described as nomadic with varying degrees of site fidelity (Sears 1990, Comtois *et al.* 2010). Four individuals were documented to have traveled more than 400 km in two weeks (Sears 1990), while others remain in the same area for a month or more (Ramp & Sears 2013). The largest concentrations of blue whales are found in the lower St. Lawrence Estuary, around the eastern tip of the Gaspé Peninsula, along the north shore of the Jacques-Cartier Passage, and in the waters adjacent to Sept-Îles. However, the low numbers of re-sightings within years suggest that blue whales in the Gulf of St. Lawrence are highly mobile. Compared to other baleen whales in the region, blue whales have exhibited the most specific habitat preferences, using areas with high salinity and slow currents (Doniol-Valcroze 2008). Although blue whales come mainly to the Gulf of St. Lawrence to feed in summer, they can occur year-round. The number of sightings tends to increase throughout summer, peaking in late August and early September; however, regional differences are apparent. The first peak in sightings occurs in June/July when blue whales are observed off the Gaspé Peninsula, while sightings in the lower St. Lawrence Estuary tend to drive the main peak in August/September. Seasonally, most blue whales leave the Gulf of St. Lawrence by early winter when ice cover becomes a factor. Blue whales were observed an average of only two days per season (occurrence), with an average occupancy of 22 days (Ramp & Sears 2013).

Outside the Gulf of St. Lawrence in Canadian waters, a few sightings have been made on the Scotian Shelf and two blue whales were sighted in the lower Bay of Fundy in the summer of 1995 (Sutcliffe & Brodie 1977, CETAP 1982). Farther south, blue whales are occasionally seen in eastern U.S. shelf waters, with a few sightings off Cape Cod, MA, in summer and fall (Wenzel *et al.* 1988). This region may represent the current southern limit of the blue whales' feeding range. However, acoustic detections and tracking using the U.S. Navy's Sound Surveillance System indicate blue whales travel can long distances throughout the western North Atlantic, including to waters north of the West Indies and deep waters east of the U.S. Exclusive Economic Zone (Clark 1995). During periods of peak whaling, blue whales were known to

frequent subtropical waters in the North Atlantic Ocean throughout the fall and winter (Reeves *et al.* 2004).

Eastern North Atlantic

Blue whales were regularly hunted from land stations in West Greenland, Iceland, Norway, Ireland, the Shetland Islands, the Hebrides, and the Faroes Islands (True 1904, Thompson 1928, Sergeant 1953, Jonsgard 1955, Sergeant 1966, Jonsgard 1977, Kapel 1979, Sigurjónsson & Gunnlaugsson 1990). The paucity of sightings during recent surveys along the coasts of Finnmark and on the banks west of Bear Island and Svalbard, where blue whales were common in the late 1800s and early 1900s, may indicate that either the historic distribution and migratory pattern have changed or that the population was depleted by whaling (Christensen *et al.* 1992). More recently, however, sighting of blue whales in Svalbard have increased noticeably in the last several years, and there were “many sightings” during the 2015 North Atlantic Sightings Surveys in the Norwegian Sea and part of the area around Jan Mayen (NAMMCO 2014).

Migration patterns for blue whales in the eastern North Atlantic Ocean are poorly understood. However, blue whales have been documented in winter months off Mauritania in northwest Africa (Baines & Reichelt 2014); in the Azores, where their arrival is linked to secondary production generated by the North Atlantic spring phytoplankton bloom (Visser *et al.* 2011); and traveling through deep-water areas near the shelf break west of the British Isles (Charif & Clark 2009). Blue whale calls have been detected in winter on hydrophones along the mid-Atlantic ridge south of the Azores (Nieukirk *et al.* 2004). Sears *et al.* (2005) reported a photo-identification match between a blue whale sighted in Mauritania and Iceland, representing an estimated 5,200 km migration.

D.3 Feeding and Prey Selection

Stomach content analysis provides information on blue whale diet, but it cannot determine whether this reflects preferred prey, or the relative abundance of prey species. Blue whales in the North Atlantic feed on relatively large euphausiid crustaceans or krill species (Jonsgard 1955, Sergeant 1966, Christensen *et al.* 1992). The species *Thysanoëssa raschii* and *Meganctiphanes norvegica* represent important food sources of blue whales in the Gulf of St. Lawrence, based on observations of feeding whales and water sampling (Sears 1987).

Although little is known of their winter distribution and migratory routes (Reeves *et al.* 2004), North Atlantic blue whales are found in summer on a range of high latitude feeding grounds from eastern Canada to Greenland (Sears & Calambokidis 2002). Doniol-Valcroze *et al.* (2012) identified sites of intense feeding activity in the northwestern Atlantic Ocean. The dominant topographic feature of this area is the Laurentian Channel that extends from the Scotian Shelf, outside the Gulf of St. Lawrence, to Tadoussac in the St. Lawrence Estuary (Doniol-Valcroze *et al.* 2012). The 200-300 m deep channel is characterized by steep slopes on both sides, and comes within 100 m of the north shore in some places. Tidal mixing and interaction with bathymetric features result in rich krill aggregations (Simard *et al.* 1986). Sears (1987) suggested that the whales’ preference for the 100 m depth during daylight hours along the north shore of the Gulf of St. Lawrence is linked to krill concentrations that occur regularly at depths of 90–120 m (Doniol-Valcroze *et al.* 2012). In general, blue whales do not typically frequent foraging areas if prey concentration does not reach a minimum threshold (Goldbogen *et al.* 2011)

as is evident in the Gulf of St. Lawrence and in other blue whale foraging areas.

Less is known about blue whale diet composition in the eastern North Atlantic Ocean, but *Thysanoëssa inermis* and *Meganyctiphanes norvegica* are likely important prey in this region (Hjort & Ruud 1929, Christensen *et al.* 1992).

D.4 Competition

Blue whales may have little competition with other baleen whales in the North Atlantic Ocean due to resource partitioning. Doniol-Valcroze (2008) found that all baleen whales in the Gulf of St. Lawrence exhibit specialized niche preferences such that they use habitats and feed on prey that differs from sympatric species. Isotope ratios in whale tissue indicate blue, fin, minke, and humpback whales in the Gulf of St. Lawrence consume different proportions of shared prey species and have different trophic niches (Gavrilchuk *et al.* 2014). In the western North Atlantic Ocean, Clark and Gagnon (2004) found that blue and fin whales overlap in high latitude summer feeding areas while humpback whales were not detected in large numbers in the area. Similarly, in the eastern North Atlantic Ocean, Charif and Clark (2009) found that detection levels of fin, humpback, and blue whales peaked at different times throughout the year, suggesting little competition for food or space.

Blue whale distribution patterns are strongly correlated with the location of dense euphausiid patches (Reilly & Thayer 1990, Croll *et al.* 1998, Fiedler *et al.* 1998, Croll *et al.* 2005).

Therefore, any downward trend in the abundance of blue whales in important feeding areas could be an indication of significant biological changes occurring within the ecosystem. For example, Comtois *et al.* (2010) attributed a decline in numbers of blue whales in the Gulf of St. Lawrence beginning in 1994 to ecosystem changes or diminished prey availability following the groundfish stocks collapsed (Hanson & Chouinard 2002, Savenkoff *et al.* 2007, Harvey & Devine 2008, Comtois *et al.* 2009). The collapse of the groundfish stocks and resulting increase in the small-bodied forage species, such as northern sand lance, capelin, and Atlantic herring that largely feed on krill, might impact blue whales because of the whales' relatively narrow trophic niche, compared to other baleen whales with generalist foraging strategies.

D.5 Reproduction

Only nine blue whale calves were observed from 1970 – 1990 along the north shore of the Gulf of St. Lawrence, suggesting either that lactating females reside in unsurveyed areas, that weaning occurs before reaching the nearshore estuaries, or that few calves were born during that time (Reeves *et al.* 1998). While it has been suggested that reproductive parameters of Antarctic blue whales, such as interbirth interval, have been changing over time due to recovery from whaling pressure (Gambell 1973, Lockyer 1979), it is unknown whether these changes are occurring in the North Atlantic Ocean populations.

D.6 Natural Mortality

Long-term sightings and photo identification data have enabled researchers to understand certain aspects of natural mortality rates among western North Atlantic Ocean blue whales. Ice entrapment is known to injure and kill some blue whales, particularly along the southwest coast of Newfoundland during late winter and early spring (Beamish 1979, Sergeant 1982). Scarring

on the dorsal surface of some whales in the Gulf of St. Lawrence is thought to be from contact with ice and or killer whales, but no direct evidence of predation has been reported (Sears 1987, Sears 1990). Based on sightings records from 1979 to 2002, Ramp *et al.* (2006) provided an estimated annual adult survival rate of 0.975 (95% confidence interval (CI) 0.960-0.985) for whales occurring in the Gulf of St. Lawrence region. Though little is known about survival rates in the eastern North Atlantic Ocean, a similar estimate may apply.

D.7 Abundance and Trends

Few abundance estimates exist for blue whales occurring in the North Atlantic Ocean due to the wide distribution and highly migratory nature of these animals. Historically, as many as 15,000 blue whales may have inhabited the North Atlantic before whaling began (Sergeant 1966, Allen 1970, Rørvik & Jonsgård 1981). Although uncertainties and caveats surround these historical estimates, researchers agree that populations were largely depleted by modern whaling throughout the 20th century and only now may be showing signs of modest recovery. The IWC annual catch database indicates over 10,000 blue whales were killed in the North Atlantic Ocean between 1868 and 1978, although only about 1,600 blue whales were taken after 1914 (Branch *et al.* 2008). Based on compiled survey and photo identification data, it is likely that the number of blue whales throughout the entire the North Atlantic Ocean now ranges from 600 to 1,500 animals (Sears & Calambokidis 2002). There are insufficient data to determine population trends for North Atlantic blue whales. Data gaps highlight the importance of long-term monitoring to understand the trends in abundance.

Western North Atlantic

Based on cumulative catches from 1898 to 1915, researchers estimate that between 1,100 and 1,500 blue whales frequented the western North Atlantic before modern whaling began (Sergeant 1966, Allen 1970), and researchers agree that blue whales were severely depleted by 1955 when IWC prohibitions against killing blue whales in the North Atlantic Ocean came into force (Jonsgard 1955, Best 1993).

Almost all recent information about the abundance of blue whales in the western North Atlantic comes from the Gulf of St. Lawrence, Canada. Within the Gulf of St. Lawrence, a long-term monitoring project provided a rare opportunity to observe changes in baleen whale distribution and ecosystem processes (Ramp & Sears 2013). From 1979 to 2009, about 440 blue whales were identified within and around the estuary area, with between 20 and 105 blue whales identified each year, with varying degrees of site fidelity among and between years (Ramp & Sears 2013); see Section I.D.2). Because the Gulf of St. Lawrence is a small portion of the population's range and it is difficult to determine the percentage of whales within the population that use the area in any given year or at any given time, the known number of individuals in those discrete areas cannot be extrapolated for an overall abundance estimate.

Ramp and Sears (2013) provided a summary of sightings from opportunistic observations from the published literature and other records of individual blue whales outside of the Gulf of St. Lawrence area. While the information was not a complete overview of historic sightings of blue whales outside of the Gulf of St. Lawrence area, it provides insight on the importance of these additional areas and further exhibits the challenge in determining an overall abundance.

Given the small number of blue whales encountered and photographed and the limited survey data, an estimate of abundance of blue whales in the Northwest Atlantic Ocean is not possible with a minimum degree of certainty (Hammond *et al.* 1990, Sears *et al.* 1990, Sears & Calambokidis 2002, Beauchamp *et al.* 2009, Waring *et al.* 2010). Some researchers speculate that there may be between 400 and 600 blue whales there (Mitchell 1974, Waring *et al.* 2010).

Without accurate current abundance estimates, no trend information is available.

Eastern North Atlantic

Estimates of pre-exploitation abundance are likely unreliable. Prior to commercial whaling, there were estimated to be a “few thousand” to 10,000 blue whales in the Denmark Strait (Rørvik & Jonsgård 1981) and 2,500 from northern Norway (Rørvik & Jonsgård 1981).

Based on the results of a summer shipboard survey in 1987, Sigurjónsson and Gunnlaugsson (1990) estimated a maximum of 442 blue whales in Icelandic waters. Sighting data from whaling vessels operating off the west coast of Iceland have demonstrated an increasing trend of approximately 5% per year since the late 1960s (Sigurjónsson & Gunnlaugsson 1990). This observed trend is considered to apply only to the population in these waters and has been corroborated by Pike *et al.* (2009). An estimate of approximately 1,000 individuals in Icelandic waters would suggest that this population is the largest in the North Atlantic (Pike *et al.* 2009). Based on the North Atlantic Sightings Surveys in 1995 and 2001, Pike *et al.* (2004) estimate there are between 1,000 and 2,000 blue whales in the surveyed areas off the east coast of Greenland, Denmark Strait, Iceland, Jan Mayen, Faroe Islands, west coast of Ireland, and north of the United Kingdom. While these figures are uncertain, historical abundance estimates indicate that this population is diminished compared with pre-exploitation estimates.

E. Natural History – Northern Blue Whales (*B. m. musculus*) in the North Pacific Ocean

E.1 Population Structure

The population structure of blue whales in the North Pacific Ocean has been evaluated through the analysis of whaling data, length frequencies, vocal repertoire, and movement data from tagging studies. There may be as many as five populations with an unknown degree of mixing among them: (a) southern Japan; (b) northern Japan/Kuril Islands/Kamchatka; (c) Aleutian Islands (which may winter in deep water north of Hawaii); (d) eastern Gulf of Alaska; and (e) the eastern North Pacific (Reeves *et al.* 1998). Blue whales were heavily exploited in waters off southern Japan (Kasahara 1950), but are no longer sighted in the area, leading researchers to believe that the population may have been extirpated. Blue whales have also been observed ranging from northern Japan in winter to the Kamchatka Peninsula in summer (Nishiwaki 1966, Ohsumi & Wada 1974, Fujise *et al.* 1995). Forney and Brownell (1996) concluded that these whales were a separate group from those exploited around the Aleutian Islands based on whaling catch locations and body size.

More recent data, described below, suggest that blue whales in the North Pacific Ocean consist of at least two existing populations, the eastern and western/central populations (Stafford *et al.* 2001, Stafford 2003, McDonald *et al.* 2006b, Monnahan *et al.* 2014, Calambokidis *et al.* 2015).

NMFS recognizes Eastern and Central North Pacific stocks under the MMPA.

Blue whales in the North Pacific produce two distinct, stereotypic calls that have been termed the northwestern and northeastern North Pacific Ocean call types, and it has been proposed that these are indicative of two distinct populations with some degree of geographic overlap (Stafford *et al.* 2001, Stafford 2003, Monnahan *et al.* 2014). The northeastern Pacific Ocean call predominates in the Gulf of Alaska, the U.S. West Coast, and the eastern tropical Pacific Ocean, while the northwestern Pacific Ocean call is heard among calling blue whales from along the Aleutian Islands to the Kamchatka Peninsula off Russia, though both call types have been recorded concurrently in the Gulf of Alaska (Stafford *et al.* 2001, Stafford 2003). Both call types were recorded in lower latitudes in the central North Pacific Ocean, although with much lower occurrence (Stafford *et al.* 2001).

Length frequency data analysis by Gilpatrick and Perryman (2008) supports separation of North Pacific Ocean blue whales into eastern and western/central populations. They found that blue whales from waters off California to Central America (the eastern North Pacific stock) are on average, two meters shorter than blue whales measured from historic whaling records in the central and western North Pacific Ocean. Additionally, satellite tagging data and photo-identification studies have shown that whales from the eastern population move between the Gulf of Alaska and the Costa Rica Dome off central America (Mate *et al.* 1999, Calambokidis *et al.* 2009a, Bailey *et al.* 2010).

E.2 Distribution and Habitat Use

In general, blue whales migrate seasonally between feeding and calving areas (Sears & Calambokidis 2002, Burtenshaw *et al.* 2004). Blue whales range throughout much of the North Pacific Ocean, from Kamchatka to southern Japan in the west, and from the Gulf of Alaska and waters off California to areas off Central America. Small numbers have been observed in Hawaiian waters and as far north as the Chukchi Sea (Berzin & Rovnin 1966, Northrop *et al.* 1970, Thompson & Friedl 1982, Yochem & Leatherwood 1985, Barlow 1997). Whales in the eastern and central North Pacific Ocean stocks range along the U.S. west and eastern Russian coasts, respectively.

As in the North Atlantic Ocean, blue whales inhabit both coastal and pelagic environments and are frequently found on the continental shelf (Calambokidis *et al.* 1990, Fiedler *et al.* 1998) and also far offshore in deep water (Wade & Friedrichsen 1978). Seasonal and interannual distribution of blue whales is strongly associated with both the static and dynamic oceanographic features that result in krill aggregations (Gendron 1990, Reilly & Thayer 1990, Del-Angel-Rodriguez 1997, Croll *et al.* 1998, Fiedler *et al.* 1998, Benson *et al.* 2002, Hamazaki 2002, Croll *et al.* 2005). Blue whales depend on a critical density of prey to feed efficiently, and thus seek out productive areas near upwelling zones and underwater thermal fronts where krill abundance is highest (Doniol-Valcroze *et al.* 2007). In the North Pacific Ocean, Munger *et al.* (2009) found that distribution was most strongly linked to cool sea surface temperatures, particularly in summer. Irvine *et al.* (2014) found that blue whale “home ranges” and “core areas” are typically near highly productive, strong upwelling centers. Additionally, Stafford *et al.* (2009) found that vocalization occurrence and rates could be predicted by sea surface temperatures, with peaks in calling occurring one to two months after cold spells.

Eastern North Pacific

Blue whales inhabit a broad area throughout the eastern North Pacific coincident with shifting feeding areas (Calambokidis *et al.* 2009a, Calambokidis *et al.* 2015). Since the 1970s, large concentrations of blue whales have been documented feeding off California each summer and fall (Calambokidis *et al.* 1990), and increasingly have been found feeding to the north and south of these summer/fall feeding areas (Calambokidis & Barlow 2004, Barlow & Forney 2007, Calambokidis *et al.* 2009a). Nine “biologically important areas” for blue whale feeding were identified off the California coast by Calambokidis *et al.* (2015), including six in southern California and three in central California. In fall, blue whales migrate northward along the North American coast to secondary feeding areas off Oregon and Washington (Burtenshaw *et al.* 2004), the Alaska Gyre, and Aleutian Islands (Stafford *et al.* 2001, Stafford *et al.* 2007). This migration was first documented in 1997 and may represent the return to a historical migratory route that was prevalent before whaling activities exploited the population (Calambokidis *et al.* 2009a). These whales also range north into the Gulf of Alaska where they seasonally overlap with the central North Pacific stock (Stafford *et al.* 2001, Stafford 2003). It is possible that this northward migration constitutes only a small portion of the California feeding population, though the sex and age composition of this group remains unknown. Bailey *et al.* (2010) monitored migratory routes of tagged whales from 1993 to 1997 and noted interannual variability in the occurrence of the whales undergoing this northward movement, which they attribute to environmental changes affecting the whales’ prey.

In winter and spring, Eastern North Pacific blue whales move to feed and breed in low latitudes, especially in the Gulf of California and the Costa Rica Dome (Reilly & Thayer 1990, Mate *et al.* 1999). Many whales have been documented off the coast of Baja California, Mexico and the relatively deep waters of the southern Gulf of California (Gendron & Hernandez 1995, Mate *et al.* 1999, Calambokidis *et al.* 2009a), but blue whale abundance in this area is relatively low (approximately 5% of the California feeding population) and peaks in March-April (Tershy *et al.* 1990). The Baja California Frontal System, a dynamic region at the convergence of the California and Davidson Currents (Etnoyer *et al.* 2004), appears to be used as a migratory stopover for whales wintering off Central America and those returning north to California waters for the summer (Rice 1974). Whales may also be feeding in this transitory area (Etnoyer *et al.* 2006). Matteson *et al.* (2010) found that blue whale distribution around the Costa Rica Dome during winter was most significantly associated with chlorophyll concentration and subsurface temperature, indicating that feeding is linked to zooplankton aggregations. Blue whales have been observed feeding year-round near the Costa Rica Dome, though Southern Hemisphere blue whales may also frequent the area (Reilly & Thayer 1990).

Blue whale occurrence and distribution may change with prey abundance and oceanographic conditions. The California Current is a dynamic ecosystem governed by large-scale oceanographic patterns, most notably the El Niño Southern Oscillation (ENSO) characterized by unusually warm temperatures. La Niña conditions can feature unusually cool temperatures in the equatorial Pacific. The Pacific (inter) Decadal Oscillation (PDO) is an El Niño-like oscillatory pattern of climate variability centered over the Pacific Ocean and North America. The PDO has considerable influence on climate sensitive living marine resources in the Pacific and over North America, including major marine ecosystems from coastal California to the Gulf of Alaska and Bering Sea (Mantua 2002). Pardo *et al.* (2015) found that inter-annual variation in blue whale

distribution during El Niño anomalies suggests that the local and regional densities of the species decreases in the Equatorial Cold Tongue and the Costa Rica Dome, and occurrence retracts to areas that remain productive (*i.e.*, the California Current, northern Gulf of California, the North Equatorial Countercurrent thermocline ridge, and the southern portion of the Humboldt Current System). During the 1997/1998 ENSO event, blue whale distribution off the U.S. West Coast was significantly different from others years as the whales shifted northward to areas that remained productive (Burtenshaw *et al.* 2004).

Western/Central North Pacific

Little is known about the distribution, migration, and habitat use of blue whales in the central and western North Pacific Ocean. Blue whales appear to feed in summer southwest of Kamchatka, south of the Aleutian Islands, the Gulf of Alaska (Gambell 1973, Watkins *et al.* 2000, Stafford 2003), and waters off Vancouver Island, Canada (Omura & Ohsumi 1964, Ivashin & Rovnin 1967, Ohsumi & Masaki 1975, Branch & Mikhalev 2008). Whaling records indicate whales were commonly taken off the southern portion of the main Japanese islands in winter, off Hokkaido in spring, and off Kamchatka, the Kurils, and the western Aleutians in summer (Kasahara 1950), suggesting a seasonal northward migration.

In winter, blue whales migrate to low latitudes in the western and central Pacific, including Hawaiian waters (Berzin & Rovnin 1966, Northrop *et al.* 1970, Thompson & Friedl 1982, Stafford *et al.* 2001, Bradford *et al.* 2017). It is possible that some individuals from this stock undertake a seasonal migration while others do not. One study, based on vocalizations, documented individual blue whales remaining off the coast of Kamchatka year-round, with peak calling in fall (Watkins *et al.* 2000). Additionally, Stafford (2003) recorded the northwestern Pacific blue whale call type (identified from recordings in Hawaii, Midway Island, the Aleutian Islands, and the northwest Pacific; (Northrop *et al.* 1970, Thompson & Friedl 1982, Stafford *et al.* 2001) in the Gulf of Alaska during fall and early winter, which may indicate some individuals do not undertake a winter migration south.

E.3 Feeding and Prey Selection

North Pacific Ocean blue whales feed primarily on *Euphausia pacifica* and *Thysanoëssa spinifera* (Rice 1986, Schoenherr 1991, Kieckhefer *et al.* 1995, Fiedler *et al.* 1998). In the Gulf of California, Mexico, blue whales feed primarily on *Nyctiphanes simplex* (Gendron 1990, Gendron 1992, Del-Angel-Rodriguez 1997), which often aggregates in dense swarms near the surface except during periods of increased surface water temperatures related to the El Niño Southern Oscillation (Gendron 1990, Sears 1990, Gendron 1992, Gendron & Sears 1993). Other known prey species include *T. inermis*, *T. longipes*, *T. raschii*, and *Nematoscelis megalops* (Kawamura 1980, Yochem & Leatherwood 1985).

One exception to their near-total dependence on euphausiid prey is that blue whales have been observed feeding on pelagic red crabs off Baja California (Rice 1974, Rice 1986), although these observations have not been confirmed by subsequent observations or other analyses (*e.g.*, fecal samples). The discovery of copepods and amphipods in the stomach contents of some whales taken during whaling activities or reports of foraging on schooling fish (Mizue 1951, Sleptsov 1955) are thought to be the result of opportunistic or accidental consumption rather than evidence of a mixed diet (Nemoto 1957, Nemoto & Kawamura 1977).

E.4 Competition

In the California Current Ecosystem, other baleen whales are present throughout the year, but the most overlap with blue whales occurs during periods of high primary productivity. It would seem unlikely that resource competition would occur with abundant prey present. In summer, blue and fin whales off the coast of central California in summer target ephemeral krill patches. Despite the overall similarities in blue and fin whales' dive profiles, Friedlaender *et al.* (2015) found differences in feeding performance that may represent unique predatory strategies that minimize competition yet maximize energy gain. Blue whale competition with another sympatric baleen whale species in the California Current Ecosystem, the humpback whale, may be minimized by the whales' partitioning of prey resources in space, time, and trophic level (Fossette *et al.* 2017).

E.5 Reproduction

No differences in the reproductive biology between blue whales in the North Pacific and North Atlantic Oceans are known or suspected. In the eastern North Pacific stock, blue whales accompanied by young calves have been observed often in the Gulf of California, Mexico from December through March, suggesting that this area is likely an important calving and nursing area (Sears 1990). However, only a portion of females observed off the U.S. West Coast were observed in the Gulf of California, indicating that there is likely another, unknown calving or nursing ground for this population (Sears *et al.* 2013). Observations of females re-sighted in alternating years with calves supports the estimated two-year calving interval noted by Lockyer (1984); Sears *et al.* (2013) estimate a mean calving interval of 2.57 years and an age of first parturition of more than 10 years. In coastal waters off California, observations of females with young whales in late summer support the idea that weaning occurs at approximately six months of age (Reeves *et al.* 1998). No information on reproductive parameters exists specific to blue whales in the central North Pacific stock.

Oleson *et al.* (2007a) assessed eastern North Pacific Ocean blue whale vocalizations and found patterns in behavior, sex, and group size for certain call types. Only males produced song, indicating song is likely involved in reproductive activities, despite being produced year-round and on both feeding and presumed breeding grounds (Stafford *et al.* 2001, Oleson *et al.* 2007a). The purpose of producing song during the feeding season is not known, but suggests that mate selection can occur outside of the breeding season (Oleson *et al.* 2007a).

E.6 Natural Mortality

A high proportion of the blue whales in the Gulf of California show signs of injuries or rake-like scars that are the result of encounters with killer whales (Sears 1990), although the rate of fatal attacks by killer whales is unknown. A well-documented observation of killer whales attacking a blue whale off Baja California Sur, Mexico indicates that blue whales are at least occasionally vulnerable to these predators in this area (Tarpy 1979). Additionally, killer whales were observed feeding on a blue whale calf off the coast of Nicaragua, but the frequency of these events remains unknown (Pitman *et al.* 2007). Unlike in the western North Atlantic Ocean, injury or suffocation from ice entrapment is not known to be a factor in the natural mortality of blue whales in the North Pacific. Based on earplug aging data collected during the early 1960s,

the survival rate for North Pacific blue whales was estimated to range from 0.93 – 0.95, which is lower than that of blue whales in the North Atlantic and Indian Oceans (Ohsumi & Wada 1974, Branch 2008b).

E.7 Abundance and Trends

In the North Pacific Ocean, commercial whaling operations killed 9,773 blue whales from 1905 to 1971 (Monnahan *et al.* 2014). Approximately 65% of these catches are estimated to have been from the western North Pacific population, which suggests that the pre-whaling abundance of the western population was much larger than the eastern population (Monnahan *et al.* 2014).

Omura and Ohsumi (1974) estimated an “initial stock size” of 4,900 for the entire North Pacific. However, this estimate and those by Wada (1975) were actually indices of abundance based on sightings from Japanese whaling catcher boats. The data were not collected or analyzed in the same ways as those from more recent sighting cruises, nor did the area of coverage include several well-known centers of blue whale abundance in the North Pacific (for example, Wada (1975), Figures 1-2, compared with Reilly and Thayer (1990), Figures 1, 3). It is therefore not possible to evaluate trends by comparing the Japanese “indices” from the 1960s and 1970s with the more recent abundance estimates from sighting cruises and photo-identification studies.

Eastern North Pacific

Based on a crude analysis of catch statistics and whaling effort, Rice (1974) estimated an initial (1924) population size of about 6,000 blue whales in the eastern North Pacific (Baja California to Alaska) and a decline to fewer than 2,000 by the early 1970s. In contrast, Monnahan *et al.* (2015) used a population dynamics model to estimate pre-whaling abundance of the eastern North Pacific population as 2,210 (95% Bayesian credible interval 1,823-3,721). The assessment by Monnahan *et al.* (2014) has been accepted by the IWC Scientific Committee (IWC 2016b).

The eastern North Pacific stock is one of the most well-studied population of blue whales, with directed surveys occurring almost every year off the California coast. The size of the feeding stock of blue whales off the U.S. West Coast has been estimated by both line-transect and mark-recapture methods, but because some fraction of the population is always outside the survey area, the line-transect and mark recapture estimation methods provide different measures of abundance for this stock. From ship-based line-transect survey data from 1986-1990, Wade and Gerrodette (1993) estimated 1,400 blue whales in the eastern tropical Pacific. Line-transect abundance estimates from summer/autumn 2001 to 2008 research cruises in the California Current ranged between 400 and 800 individuals (Barlow & Forney 2007, Forney 2007, Barlow 2010). These estimates are considerably lower than previous line-transect estimates of approximately 1,900 whales between 1991 and 1996 (Barlow 2010). Part of the reason for the discrepancy is likely due to the northward shift in blue whale distribution related to oceanographic conditions and prey abundance, and not a population decline (Barlow & Forney 2007, Calambokidis *et al.* 2009a). New abundance estimates based on photographic mark-recapture data for the period 2005 to 2011 range from approximately 1,000 to 2,300 whales (Calambokidis & Barlow 2013). An estimate of 1,647 (CV=0.07) whales is regarded as the best estimate of blue whale abundance for the period 2008-2011, based on Calambokidis and Barlow’s (2013) Chao 4-year model (Carretta *et al.* 2015).

Mark-recapture estimates provide the best indicator of population trends for this stock, because recent northward shifts in blue whale distribution might result in a negative bias in estimates based on line transect surveys due to the locations of those surveys. Based on mark-recapture estimates from the U.S. West Coast and Baja California, Mexico coast, Calambokidis *et al.* (2009b) estimated an abundance increase of just under 3% per year, but it is not known if that corresponds to the maximum growth rate of this stock.

Calambokidis *et al.* (2015) argued that the eastern Pacific blue whale population has not shown signs of recovery from commercial whaling over the last twenty years, particularly compared to other baleen whale populations in the region. In contrast, Monnahan *et al.* (2015) used a population dynamics model to estimate that the eastern Pacific blue whale population was at 97% of carrying capacity (95% Bayesian credible interval 62%–99%) in 2013 and suggested that an observed lack of a population size increase since the early 1990s can be attributed to density dependence. The authors estimated that the minimum population size of the eastern North Pacific Ocean population was at least 460 whales during the last century, despite extensive commercial whaling exploitation. In 2015, the IWC Scientific Committee endorsed the conclusions of Monnahan *et al.* (2015) regarding the population approaching carrying capacity, but recommended refinement of the analysis to address uncertainties when additional data become available (IWC 2016b).

Western/Central North Pacific

No blue whales were observed during a summer/fall 2002 survey of the entire U.S. EEZ around the Hawaiian Islands (Barlow 2006), but some blue whales were observed in these waters during a summer/fall 2010 survey, resulting in an abundance estimate of 133 (CV = 1.09) blue whales (Bradford *et al.* 2017). This is currently the best available abundance estimate for this stock within the U.S. EEZ around the Hawaiian Islands, but the majority of blue whales would be expected to be at higher latitude feeding grounds during the time of year the survey was conducted. Developing abundance estimates for this stock is difficult due to the uncertainty surrounding their seasonal distribution and overlap with the eastern stock in the central North Pacific Ocean. Thus, the status of blue whales in Hawaiian waters is unknown and there are insufficient data to evaluate trends in abundance.

An aerial survey of the former Akutan whaling grounds around the eastern Aleutians in 1984 produced no sightings of blue whales, suggesting that the population remained severely depleted (Stewart *et al.* 1987). No blue whales were sighted during a marine mammal survey south of the Aleutian Islands in 1994 (Forney & Brownell 1996). However, as described in Sections I.E.1 and I.E.2, northwestern blue whale calls have been recorded in summer southwest of Kamchatka, south of the Aleutian Islands, in the Gulf of Alaska, and in lower latitudes of the western and central Pacific, including Hawaii, in winter (Watkins *et al.* 2000, Stafford *et al.* 2001, Stafford 2003, Carretta *et al.* 2015).

F. Natural History – Antarctic Blue Whales (*B. m. intermedia*)

F.1 Population Structure

Antarctic blue whale movement information from Discovery tags (i.e., uniquely numbered metal tubes that were shot into whales' muscles during scientific cruises and recovered when the

animals were killed during whaling), photo-identification, individual genotype identification, and satellite tagging show both small-scale movements (e.g., within IWC Areas, both within and between seasons) and large-scale movements (e.g., up to 180° longitude). Only one blue whale call type has been recorded in the Southern Ocean (Donovan 1991, Rankin *et al.* 2005, McDonald *et al.* 2006b, Širović *et al.* 2009), suggesting a single population, or alternatively, that any discrete populations are not acoustically distinct.

Antarctic blue whales have the greatest haplotypic diversity of any subspecies, likely due to a large historical abundance (LeDuc *et al.* 2007). Sremba *et al.* (2012) found significant genetic differences at fixation indices between some IWC Areas. Attard *et al.* (2016) analyzed a larger set of genetic samples that was not bound by *a priori* population boundaries, and found three genetically differentiated populations that occur sympatrically off Antarctica during the austral summer feeding season. However, in 2016 the IWC Scientific Committee found the evidence for three populations to be inconclusive and recommended alternate methods with greater power to discriminate population structure should be considered (IWC 2016a).

F.2 Distribution and Habitat Use

Blue whales in the Antarctic are almost exclusively from the Antarctic subspecies (Branch *et al.* 2007b). Due to the rarity of sightings, most of the information on Antarctic blue whale distribution and habitat use has been obtained from monitoring of vocalizations and historical whaling data. Acoustic recordings indicate that blue whales are distributed around Antarctica, generally remaining south of the Antarctic Convergence Zone (55° S) during summer, and likely moving into mid- and low-latitude waters in fall and winter (Stafford *et al.* 1999b, Clark & Fowler 2001, Širović *et al.* 2004, Stafford *et al.* 2004, McKay *et al.* 2005, Rankin *et al.* 2005, Stafford *et al.* 2005, McCauley *et al.* 2006, McDonald *et al.* 2006b, Ensor *et al.* 2009, Širović & Hildebrand 2011). However, as described later in this section, not all Antarctic blue whales migrate each year.

The seasonal and inter-annual distribution of Antarctic blue whales is strongly associated with both the static and dynamic oceanographic features that aggregate krill (Gendron 1990, Reilly & Thayer 1990, Del-Angel-Rodriguez 1997, Fiedler *et al.* 1998, Benson *et al.* 2002, Hamazaki 2002, Croll *et al.* 2005). Blue whales depend on a critical density of prey to feed efficiently, and thus occur in highly productive areas near upwelling zones and thermal fronts where krill are most abundant (Doniol-Valcroze *et al.* 2007). In the Antarctic, krill abundance is governed by the combination of a number of oceanographic features, but mainly by the presence of phytoplankton near ice edges and continental shelves (Murase *et al.* 2002, Atkinson *et al.* 2004, Siegel 2005). Many of the sub-Antarctic islands and coastal areas where Antarctic blue whales have been detected are highly productive in winter and sustain abundant zooplankton (Moore & Abbott 2000).

Most Antarctic blue whales are believed to make annual migrations between summer feeding and winter breeding/calving grounds. This theory is supported by detection of Antarctic call types in winter in low latitude areas, including the eastern tropical Pacific, the central Indian Ocean, and off southwestern Australia and northern New Zealand (McCauley *et al.* 2001, Stafford *et al.* 2004, McDonald 2006, Samaran *et al.* 2013). Antarctic blue whales were known to migrate into South African and Namibian waters in a period prior to the onset of intensive

whaling; the region off southwest Africa could be a breeding location based on seasonality of Antarctic blue whale historical catches there (Branch *et al.* 2007a, Branch *et al.* 2007b), but since the cessation of whaling, there have been only two recorded blue whale sightings there (Branch *et al.* 2007b).

Not all Antarctic blue whales migrate each year. Year-round acoustic detections of Antarctic blue whale calls near the West Antarctic Peninsula (Širović *et al.* 2004, Širović *et al.* 2009), the Weddell Sea and along the Greenwich meridian (Thomisch *et al.* 2016), eastern Antarctica at 67° S, 70° E (McKay *et al.* 2005, Širović *et al.* 2009), and South Georgia Island (Hinton 1915, Risting 1928) suggest that only a portion of the population migrates north each year, or that at least one population is resident in Antarctica throughout the year (Attard *et al.* 2016). However, the frequency of calling in these Antarctic areas generally decreases during winter as many of the whales migrate northward (Širović *et al.* 2009). Antarctic blue whales have also been detected in mid-latitude waters year-round. Samaran *et al.* (2010) documented the year-round presence of Antarctic blue whales around the Crozet Islands in the southern Indian Ocean.

F.3 Feeding and Prey Selection

Cotte *et al.* (2006) found that historical abundance of blue whales in the Antarctic, based on whale catches, was highly correlated with the extent of the seasonal ice zone. The timing and extent of sea ice formation varies each year, but data indicate that, in contrast to patterns in the Arctic Ocean, the Southern Ocean around Antarctica has experienced a slight but statistically significant increase in sea ice extent since the 1970s, although regional trends differ (Parkinson & Cavalieri 2012). However, after decades of increase, including record high daily extents in 2015 and a record high winter maximum in 2014, the sea ice minimum extent reached a record low in March 2017 (National Snow & Ice Data Center 2017). Given the variability in these conditions and uncertainty in how Antarctic sea ice is responding to climate change, impacts on baleen whales, including blue whales, are unclear. See Section I.H.1.4 below for more information about potential impacts to blue whale prey due to climate and ecosystem changes.

F.4 Competition

In the Southern Ocean around the West Antarctic Peninsula, fin whales are found farther from shore than other baleen whales foraging near the ice edge, making them less likely to compete with blue whales for resources (Murase *et al.* 2002, Murase *et al.* 2006, Širović 2006), but there is significant spatial overlap between blue, minke, and killer whales throughout known Antarctic feeding areas (Kasamatsu *et al.* 2000). Mori and Butterworth (2004) suggested that blue whales have a competitive advantage over minke whales because their large body size enables better tolerance of low krill density compared to their smaller relatives.

Baleen whales from different species have been shown to preferentially feed on Antarctic krill of specific sizes, supporting the hypothesis that spatial separation and unique feeding habits result in less interspecific competition (Santora *et al.* 2010). It is not clear how climate change may influence the density, distribution, or availability of krill and whether these changes may impact interspecific competition. However, it is unlikely that interspecific competition is limiting recovery of Antarctic blue whales.

F.5 Reproduction

The reproductive biology of Antarctic blue whales is likely very similar to that of blue whales in the North Pacific and North Atlantic Oceans. Little is known about the reproductive behavior of Antarctic blue whales. Branch (2008b) reviewed available information on Antarctic blue whale life history parameters, including age at sexual maturity and inter-calf interval. Based on analysis of a very small sample of earplug layers, Antarctic blue whales are thought to reach sexual maturity between 8-12 years old, which is later than blue whales in the Northern Hemisphere (Ichihara 1966, Ohsumi & Wada 1974, Branch 2008a, Branch 2008b). However, Lockyer (1984) argues that they may reach sexual maturity as young as five years old, which is more consistent with other baleen whales. Antarctic blue whales are thought to give birth every 2.5 years (Ohsumi & Wada 1974, Branch 2008b).

The historically intensive exploitation pressure on this population may have affected reproductive output. A lower post-exploitation density of whales (compared with pygmy blue and other baleen whales; (Branch *et al.* 2007a) that has access to relatively plentiful prey resources may have led individuals to reach sexual maturity more quickly, give birth more often, or live longer (Branch 2008b). However, this has not been confirmed.

F.6 Natural Mortality

Little information exists concerning the natural mortality of Antarctic blue whales. It is possible that calves fall victim to killer whale predation based on similar observations in the north and central Pacific, noted above. No estimates of mortality rates among adults or calves exist for this population, though survival rates are likely comparable to other blue whale populations ranging from 0.93-0.98 (Mizroch *et al.* 1984, Ramp *et al.* 2006, Branch 2008a).

F.7 Abundance and Trends

Some 360,000 blue whales were whaled from the Southern Hemisphere last century (Clapham & Baker 2002). Branch (2008c) estimated that catches of Antarctic blue whales (345,775 whales) reduced the population from 256,000 (95% credibility interval 235,000-307,000) in the 1920s to a low of 395 (235-804) animals in the early 1970s, which is just 0.15% (0.10-0.28%) of pre-exploitation levels. Blue whale sightings remain rare in the Antarctic (0.17-0.52 groups sighted per 1,000 km surveyed) (Branch *et al.* 2007b), although sighting rates increased over three circumpolar surveys corresponding to mid-years of 1981, 1988, and 1998 (Branch 2007). The distribution of recent sightings is narrowly concentrated along the edge of the pack ice and continental shelves, compared to the more broadly distributed historical catches (Branch *et al.* 2007b).

Currently, Antarctic blue whales are estimated to number 2,280 individuals (CV = 0.36) based on the IWC International Decade of Cetacean Research and the Southern Ocean Whale Ecosystem Research (SOWER) annual summer surveys from 1991/92 through 2003/04, which covered 99.7% of the area between the pack ice and 60° S (Branch 2007). The population is estimated to be increasing at a rate of 6.4% per year (95% credibility interval 2.4-8.4%), but remains depleted at less than 1% of pre-exploitation abundance (Branch *et al.* 2004, Branch 2008c).

G. Natural History – Pygmy-type blue whales (*B. m. brevicauda*, *B. m. indica*, and *B. m.* unnamed subsp.)

G.1 Population Structure

The population structure of pygmy-type blue whales was reviewed at the 2016 IWC Scientific Committee meeting, in preparation for population assessments. The IWC noted certain shortcomings in the currently available genetic data. Specifically, most of the pygmy blue whale samples collected in the Southern Hemisphere are from feeding areas or along migratory pathways, many of which are also used by Antarctic blue whales. The geographic stratification of samples by area may be confounded by this mixing, such that any genetic differences found between areas may not represent different breeding populations (IWC 2016a). Therefore, the IWC recommended using “acoustic populations” to delineate pygmy blue whale populations for purposes of assessment, until additional genetic samples are obtained (IWC 2016a).

As discussed in previous sections, stereotypical call types can be an indication of population structure. After reviewing relevant scientific literature, the IWC Scientific committee identified six acoustic populations in the Southern Hemisphere and northern Indian Ocean, including Antarctic blue whales (discussed in Section I.F), Chilean blue whales, and pygmy-type blue whales from the Northern Indian Ocean, New Zealand, the Indonesia/western Australia region, and the southwest Indian Ocean (IWC 2016a). Chilean and Northern Indian Ocean blue whales are considered separate subspecies. The three remaining non-Antarctic acoustic populations belong to *B. m. brevicauda*.

Australia and New Zealand blue whales are not genetically separated (Sremba *et al.* 2015) and are morphologically similar, but they have distinct song types and there are no photo-identification matches between the populations (Galletti Vernazzani *et al.* 2016). Balcazar *et al.* (2015) identified the acoustic boundary between Australian and New Zealand whales as the junction of the Indian and Pacific Oceans.

G.2 Distribution and Habitat Use

G.2.1 *B. m. brevicauda*

Pygmy blue whales undergo seasonal migrations to breeding and feeding locations, and are generally tied to highly productive areas with dense aggregations of krill. Pygmy blue whales mainly remain north of the Antarctic Circumpolar Current (55° S) and are most abundant in waters off Australia, Madagascar, and New Zealand (Reilly *et al.* 2008).

Australia

Australian pygmy blue whales likely spend winter in waters off Indonesia before traveling south along western Australia to feed in summer (Gill 2002, Branch *et al.* 2007b, Rennie *et al.* 2009, McCauley & Jenner 2010, Double *et al.* 2014). Acoustic data indicate that these whales are also distributed in the sub-Antarctic waters of the southern Indian Ocean in summer and fall, including near the Crozet Islands and Amsterdam Island, likely following productive waters of the Subantarctic and Subtropical Fronts (Samaran *et al.* 2010, Samaran *et al.* 2013).

Two main feeding aggregations around Australia have been identified: Perth Canyon off Western Australia (Rennie *et al.* 2009) and between the Great Australian Bight and Bass Strait in the Bonney Upwelling off South Australia and Victoria (Gill 2002, Gill *et al.* 2011). Genetic evidence suggests that the whales comprise one mating group (Attard *et al.* 2010).

A small number of Australian pygmy blue whales, as well as hybrids between Australian pygmy and Antarctic blue whales, have been identified off Antarctica, which may indicate pygmy blue whales are moving into Antarctic waters in response to less competition with the depleted population of Antarctic blue whales for food, or may be a result of environmental conditions related to climate change (Attard *et al.* 2012).

Madagascar

Ljungblad *et al.* (1998) first described the unique call types of whales feeding in sub-Antarctic waters near the Madagascar Plateau and into subtropical areas. These whales are thought to migrate in spring and summer south from the Seychelles and Amirante Islands, through the Mozambique Channel to the Crozet Islands and Prince Edward Islands to feed before returning north in fall (Zemsky & Sazhinov 1982). In summer, pygmy blue whales have a nearly continuous distribution in sub-Antarctic waters between Africa and Australia (Branch *et al.* 2007b).

The Madagascar call type has been recorded throughout this region, including Diego Garcia in May-July (Stafford *et al.* 2011); the Mozambique Channel in November-December (Cerchio *et al.* 2016); the Madagascar Basin south of La Reunion Island from March-June, peaking in April-May (Samaran *et al.* 2013); southwest of Amsterdam Island from December-May, peaking in March (Samaran *et al.* 2013); and off Crozet Island in December-June, peaking in April (Samaran *et al.* 2010). The Madagascar call type has also been recorded at approximately 61.5° S, indicating they may, at least occasionally, travel as far south as the Antarctic continental shelf (Gedamke & Robinson 2010).

Ocean currents running into the steep bathymetric features of the Madagascar Plateau likely stimulate upwelling and zooplankton production that attract whales to the area (Branch *et al.* 2007b). The sighting rates on the Madagascar Plateau (Best *et al.* 2003) are an order of magnitude greater than Antarctic sightings (Branch *et al.* 2007b).

New Zealand

Based on sightings, strandings, and acoustic detections, pygmy-type blue whales are present in New Zealand waters nearly year-round (McDonald *et al.* 2006b, Miller *et al.* 2014, Olson *et al.* 2015). New Zealand blue whale calls have also been detected in winter (June-August) in the Tasman Sea and the Lau Basin near Tonga (Balcazar *et al.* 2015), and as far south as 52° S in summer (January-March) (Miller *et al.* 2014). Whales appear to be concentrated on foraging grounds in the South Taranaki Bight between the North and South Islands, and along the east coast of Northland (North Island), which may be a migratory corridor (Torres 2013). Feeding behavior has been observed off the east and west coasts of the South Island and the Hauraki Gulf (Torres 2013, Olson *et al.* 2015).

G.2.2 *B. m. indica*

Blue whales are considered to be resident within the northwestern Indian Ocean, given sightings and strandings year-round (Yochem & Leatherwood 1985, Branch *et al.* 2007b, Anderson *et al.* 2012, Ilangakoon & Sathasivam 2012), as well as distributional gaps to the south and east (Branch *et al.* 2007b). However, blue whales undertake migrations within the region. Based on catch, sighting, stranding, and acoustic detection data, Anderson *et al.* (2012) proposed that blue whale distribution in the Northern Indian Ocean is driven by oceanographic changes associated with the monsoons. Specifically, most blue whale feed in productive upwelling areas off Somalia and southern Arabia during the southwest monsoon (approximately May-October), while some feed off the southwest coast of India and the west coast of Sri Lanka. The whales then disperse during the northeast monsoon (approximately December-March) to areas such as the east coast of Sri Lanka, west of the Maldives, the Indus Canyon, and parts of the southern Indian Ocean (Anderson *et al.* 2012). A lack of recent records from the northeast Arabia Sea and north of the Seychelles, however, suggests that whales there may have been extirpated by Soviet whalers in the 1960s (Anderson *et al.* 2012).

Acoustic evidence suggests that some of these whales may travel as far south as the sub-Antarctic waters around Crozet Islands in late summer and early fall, though calling was much less frequent compared with the other blue whale populations simultaneously using the area (Samaran *et al.* 2010).

G.2.3 *B. m. unnamed subspecies*

In summer and fall, Chilean blue whales feed along the west coast of South America, particularly in the Chiloense Ecoregion including the Corcovado Gulf, Pacific and northwest coasts of Chiloe Island, and inner sea of Chiloe Island (Hucke-Gaete *et al.* 2004, Cabrera *et al.* 2005, Abramson & Gibbons 2010, Galletti Vernazzani *et al.* 2012). The whales then migrate to lower latitude areas including the Galapagos Islands and the Eastern Tropical Pacific (ETP) (Hucke-Gaete *et al.* 2004, Torres-Florez *et al.* 2015, Hucke-Gaete *et al.* 2016).

The Southeast Pacific blue whale call type has been recorded in the ETP year-round, although the occurrence is an order of magnitude lower than in the Chiloense Ecoregion (Buchan *et al.* 2015). The calls generally peak in June (Buchan *et al.* 2015), and are detected less frequently from September to March (Stafford *et al.* 1999a).

Chilean blue whales may mix with whales from the northeast Pacific in the ETP. Based on genetic analysis, it appears that northeast Pacific blue whales primarily use northern ETP waters (e.g., the Costa Rica Dome off Central America) and Chilean blue whales primarily use southern ETP waters (e.g., off Peru and Ecuador), but they do not do so exclusively. There may gene flow between hemispheres, and indeed, Chilean blue whales' genetic differentiation from northeast Pacific blue whales is less than their differentiation from Antarctic or Indian Ocean blue whales (LeDuc *et al.* 2016). LeDuc *et al.* (2016) suggested that, at least seasonally, the Chilean blue whale range should perhaps include the Costa Rica Dome and mid-latitudes of the eastern North Pacific.

There is also sympatry with Antarctic blue whales in the southern ETP breeding area, and likely on migratory pathways along the western coast of South America. Antarctic blue whale calls acoustically detected in the ETP in May-September, peaking in July (Stafford *et al.* 2004). However, Chilean blue whales are genetically differentiated from Antarctic blue whales (LeDuc *et al.* 2016), so they are unlikely to be interbreeding.

G.3 Feeding and Prey Selection

Similar to other blue whales, pygmy-type blue whales feed on abundant and accessible euphausiids and other crustaceous zooplankton (Yochem & Leatherwood 1985). An analysis of stomach contents revealed a range of krill and other zooplankton within the pygmy whale diet, with a large proportion of *Euphausia frigida*, *Euphausia vallentini*, and *Myctophum punctatum* (Pervushin 1968). The Australian population feeds primarily on *Euphausia recurva* and *Nyctiphanes australis* that are abundant in nearshore upwelling areas off the southern coast (Gill 2002, Rennie *et al.* 2009). In New Zealand's South Taranaki Bight, pygmy blue whales feed predominantly on *Nyctiphanes australis* (Torres *et al.* 2014).

The Chiloense Ecoregion feeding ground is a complex system that receives inputs of both oceanic subantarctic and continental freshwater (Palma & Silva 2004) leading to an array of frontal features that influence primary production. *Euphausia vallentini* dominates the mesoplankton in this Ecoregion (Buchan & Quiñones 2016). It doubles in abundance between winter and spring (González *et al.* 2010) and likely peaks in late summer (Croll *et al.* 2005), which coincides with the seasonal peak in acoustic detections made by Buchan *et al.* (2015).

G.4 Competition

Little or nothing is known about possible competition between pygmy-type blue whales and sympatric species. The high mobility of these whales enables them to take advantage of transitory concentrations of prey over a very large area and adapt to local prey availability with variable oceanographic conditions.

G.5 Reproduction

The reproductive biology of pygmy-type blue whales is likely very similar to that of blue whales in the North Pacific and North Atlantic Oceans. In addition, considerable data on the size and age of pygmy blue whales at sexual maturity has been derived from translated and reanalyzed Japanese and Soviet whaling catch data from the Indian Ocean. Earplug layers, size descriptions, and ovarian corpora reveal that female pygmy-type blue whales in the Indian Ocean are approximately 19 m and 9.9 years old when sexually mature (Branch *et al.* 2007a, Branch & Mikhalev 2008). Similar to other blue whales, pygmy-type blue whales are thought to give birth every 2.6 years (95% CI 2.2 – 3.0) (Branch 2008b), with an average 43% of females pregnant among the population in a given year (Mizroch *et al.* 1984, Mikhalev 2000, Branch *et al.* 2004).

G.6 Natural Mortality

Predation and other sources of natural mortality have not been documented for pygmy-type blue whales. Soviet and Japanese whaling records have been analyzed to reveal the number of pygmy blue whales in various age classes. From this information, researchers estimated that the

mortality rate among this subspecies is approximately 0.06 (95% CI 0.05 – 0.07) (Branch 2008b). Calf mortality rates have not been calculated for blue whales, but it is likely higher than adult mortality, as it is in other baleen whales (Branch *et al.* 2004).

In the Northern Indian Ocean, shark bite scars were observed less frequently in blue whales caught by Soviet whaling fleets in the Arabian Sea compared with pygmy blue whales in the southern Indian Ocean (Mikhalev 2000).

G.7 Abundance and Trends

G.7.1 *B. m. breviceauda*

The status and abundance of pygmy blue whales is less well known compared to the Antarctic subspecies due to their large range and uncertain population structure. Similar to the Antarctic subspecies, pygmy blue whale populations were exploited throughout the 20th century, though they likely had a lower pre-exploitation abundance (Branch *et al.* 2008, Reilly *et al.* 2008, Stafford *et al.* 2011). Some abundance estimates exist, though little work has been done to determine overall pygmy blue whale abundance. Best *et al.* (2003) suggest a minimum abundance of 424 (CV = 0.42) pygmy blue whales on the Madagascar Plateau, or 472 (CV = 0.48) when an unidentified “like blue” whale sighting was included in the analysis. Kato *et al.* (2007) estimated 671 (279–1613) pygmy blue whales from a line-transect survey of a small area off the southern coast of Australia. Acoustic monitoring of Australian pygmy whales during feeding migrations in the southeastern Indian Ocean has yielded an estimate that this population likely ranges between 660-1,750 whales (Jenner *et al.* 2008, McCauley & Jenner 2010, Gill *et al.* 2011). Combining the Australian and Madagascar populations would yield a maximum of 2,950 pygmy blue whales in the southern Indian Ocean. The small size of the New Zealand photo-identification catalog precludes development of a mark-recapture abundance estimate at this time (IWC 2016a).

While little population trend information is available for pygmy blue whales, researchers have estimated pre-exploitation abundance for some of the feeding populations. Zemsky and Sazhinov (1982) estimate a pre-exploitation (1959/1960) abundance of 7,598 pygmy blue whales in the sub-Antarctic region north of 54° S between 0° and 80° E, corresponding with the Madagascar population, and 2,900 pygmy blue whales in the Australian region. Catch records from illegal Soviet whaling throughout the 1960s indicate that more than 12,500 pygmy blue whales were caught in the southern Indian Ocean, indicating a minimum population size for that time period (Branch *et al.* 2004). It is possible that the current combined estimate of nearly 3,000 whales (see above) represents less than 25% of the historical population size.

These figures indicate that pygmy blue whales remain depleted throughout their range, though perhaps to a lesser extent than Antarctic blue whales (Branch *et al.* 2007b, Branch *et al.* 2008).

G.7.2 *B. m. indica*

Nearly 1,300 blue whales were caught illegally in the Arabian Sea by Soviet whalers from 1963-1966 (Mikhalev 2000). There is no current abundance estimate for northern Indian Ocean blue whales.

G.7.3 *B. m.* unnamed subspecies

From 1908-1971, an estimated 5,782 blue whales were caught by whalers in the Southeast Pacific, including waters offshore of Chile, Peru, and Ecuador (Williams *et al.* 2011), with an estimated 4,288 from Chile alone (Branch *et al.* 2007c). Jackson (2016) estimated the median pre-exploitation abundance to be 2,100-3,600 blue whales, while Williams *et al.* (2011) estimate approximately 2,000-6,200. A minimum abundance estimate for the Chiloé Island feeding ground in southern Chile in mid-2008 is 450 animals (CV 0.17) for an open population or 576 (CV-0.16) for a closed population, based on a model that allows for a mixture of resident and transient whales (IWC 2016a). Since this represents only a portion of the population's feeding range, this is considered a minimum abundance for Chilean blue whales.

H. Potential Threats and Other Stressors

In this Revised Recovery Plan, we considered eleven natural and human-related stressors to determine whether they might present a threat to blue whale recovery. As used in this plan, a threat is any factor, natural or human-related, that impedes recovery or contributes to blue whale extinction risk. Our assessment of these stressors is summarized in Table 1, including the relevant ESA listing factor(s)², potential effect pathways for the stressor to act on individual animals, the subspecies and/or populations known to be affected (or if unknown, whether one or more is particularly vulnerable), and the severity of the effects. More specific information is discussed in Sections I.H.1-I.H.2.

While the species is listed as endangered, it is not known whether and to what extent current threats are putting the globally listed species at risk of extinction, although we identify numerous *potential* threats. We use the term potential threat to mean a stressor that a) contributed to the species' extinction risk, such as commercial whaling, and has the potential to do so again unless certain measures are taken or remain in place; or b) is known to be affecting one or more subspecies or population, but more research is needed to understand the extent to which the stressor occurs or affects the globally listed entity. We also describe other stressors where there is sufficient information to indicate that they do not currently pose a threat to the listed global species. For these stressors, there is currently no evidence that the effects (which may even include the loss of individual blue whales) have population-level consequences or are significant enough to contribute to the species' extinction risk.

² Section 4(a)(1) of the ESA and the listing regulations (50 CFR part 424) set forth considerations for listing species. We list a species if it is endangered or threatened because of any one or a combination of the following: (A) the present or threatened destruction, modification, or curtailment of its habitat or range; (B) overutilization for commercial, recreational, scientific, or educational purposes; (C) disease or predation; (D) inadequacy of existing regulatory mechanisms; or (E) other natural or human-made factors affecting its continued existence.

Table 1. Stressors that may be affecting blue whale recovery.

Stressor	ESA Listing Factors	Potential Effect Pathways	Subspecies/Populations Affected	Severity of Impact
Directed Hunting	<ul style="list-style-type: none"> • Overutilization 	<ul style="list-style-type: none"> • Injury and/or mortality 	<ul style="list-style-type: none"> • All 	Historically severe but presently controlled. Requires continued action to ensure it does not return as an operative threat.
Ship Strikes	<ul style="list-style-type: none"> • Other natural or manmade factors 	<ul style="list-style-type: none"> • Injury and/or mortality 	<ul style="list-style-type: none"> • Northern Indian Ocean • Eastern North Pacific • Chilean • Unknown whether or to what extent other populations are affected, but those with ranges overlapping areas of high shipping traffic are vulnerable. 	Unknown; likely varies by population (<i>e.g.</i> , potentially high for Northern Indian Ocean, potentially low/medium for Eastern North Pacific and Chilean). Relatively few strikes have been reported but many likely go unreported or undetected.
Entanglement in Marine Debris and Fishing Gear	<ul style="list-style-type: none"> • Habitat • Other natural or manmade factors 	<ul style="list-style-type: none"> • Injury and/or mortality • Impairment of swimming, feeding, breeding • Physiological response and negative health effects of long-term entanglement 	<ul style="list-style-type: none"> • Northern Indian Ocean • Eastern North Pacific • Unknown but potentially all, particularly those with ranges that include areas of high fishing effort 	Unknown but potentially low; likely varies by population. Very few entanglements have been reported but many likely go unreported or undetected.
Anthropogenic Noise	<ul style="list-style-type: none"> • Habitat 	<ul style="list-style-type: none"> • Behavioral response • Physiological response • Injury (hearing damage or impairment) 	<ul style="list-style-type: none"> • Unknown but potentially all, particularly those with ranges that include areas of high shipping traffic, oil and gas exploration and development, and military activity 	May range from no effect to potentially significant effects on whales' fitness and their habitat; likely varies by population. Research needed to determine degree of impact.
Loss of Prey Based Due to Climate and Ecosystem Change	<ul style="list-style-type: none"> • Habitat • Other natural or manmade factors 	<ul style="list-style-type: none"> • Changes in prey abundance and/or distribution 	<ul style="list-style-type: none"> • Unknown but potentially all, particularly those that feed at high latitudes or are geographically restricted (<i>e.g.</i>, Northern Indian Ocean population) 	Unknown, but impacts to krill under certain future climate scenarios may be severe. Likely varies by population.

Stressor	ESA Listing Factors	Potential Effect Pathways	Subspecies/Populations Affected	Severity of Impact
Environmental Contaminants and Pollutants	<ul style="list-style-type: none"> • Habitat 	<ul style="list-style-type: none"> • Sublethal health effects • Mortality 	<ul style="list-style-type: none"> • Unknown but potentially all 	Low; no evidence of population- or species-level effects; broad distribution and wide-ranging movements expected to lessen impact of localized events (e.g., oil spills).
Disease	<ul style="list-style-type: none"> • Disease or predation 	<ul style="list-style-type: none"> • Sublethal health effects • Mortality 	<ul style="list-style-type: none"> • Chilean • Unknown but potentially all 	Low; no evidence linking toxins or disease to chronic health problems or deaths of blue whales.
Behavioral Disturbance from Vessel and UAS Interactions	<ul style="list-style-type: none"> • Habitat • Overutilization • Inadequacy of existing regulations 	<ul style="list-style-type: none"> • Behavioral response • Physiological response 	<ul style="list-style-type: none"> • Unknown but potentially all, particularly those with ranges that include nearshore areas accessible to whale watching activities 	Low; no evidence linking short-term effects of disturbance to population- or species level impacts.
Research	<ul style="list-style-type: none"> • Other natural or manmade factors 	<ul style="list-style-type: none"> • Behavioral response • Physiological response 	<ul style="list-style-type: none"> • All 	Low; no evidence linking short-term effects of disturbance to population- or species level impacts.
Predation and Natural Mortality	<ul style="list-style-type: none"> • Disease or predation 	<ul style="list-style-type: none"> • Injury and/or mortality 	<ul style="list-style-type: none"> • All 	Low; no evidence this is a threat to recovery of any population or the species.
Competition for Resources	<ul style="list-style-type: none"> • Habitat • Other natural or manmade factors 	<ul style="list-style-type: none"> • Changes in prey abundance and/or distribution 	<ul style="list-style-type: none"> • All 	Low; no evidence that competition with sympatric species is a threat; management limits impacts of krill fisheries on the whales' prey resource.

H.1 Potential Threats

H.1.1 Directed Hunting

Direct hunts were the main cause of initial depletion of blue whales and other large whales. Because of their size and speed, blue whales were safe from early whalers, who could not pursue them in open boats with hand harpoons. However, a Norwegian, Sven Foyn, revolutionized the whaling industry with the invention of the exploding harpoon gun and by using steam and diesel powered factory ships and catcher boats (Schmitt *et al.* 1980, Reeves & Barto 1985). Foyn also perfected the technique of inflating dead whales with air so they would not sink after being harpooned. The eventual introduction of deck-mounted harpoon cannons made it possible to kill and secure blue, fin, and sei whales on an industrial scale (Tønnessen & Johnsen 1982). (Tønnessen & Johnsen 1982). Over 380,000 blue whales were taken from 1868-1978, mostly from Antarctic waters (Branch *et al.* 2008). The directed hunt peaked in 1931 when over 29,000 blue whales were killed in one season. After that, blue whales became so scarce that the whalers turned to other species. The International Whaling Commission banned all hunting of blue whales in 1966 and gave them worldwide protection, although illegal Soviet whaling continued through the 1970s (Yablokov 1994, Mikhalev 2000). The IWC's moratorium on the commercial hunting of all whale species has been in effect since 1986, and it has almost certainly had a positive effect on the species' recovery, although the slow rate of recovery of some populations is likely a direct result of the extent of past hunting. There is currently no commercial, aboriginal subsistence, or scientific whaling for blue whales by IWC member nations party to the moratorium. There is no evidence of illegal whaling of blue whales in recent years.

Directed hunting is presently controlled, but it requires continued U.S. involvement and collaboration through the IWC to ensure it does not return as an operative threat.

H.1.2 Ship Strikes

Blue whales are vulnerable to ship strikes, due at least in part to the seasonally coastal distribution of some populations and the overlap with shipping routes, such as off southern Sri Lanka and southern California (Berman-Kowalewski *et al.* 2010, deVos *et al.* 2013, Priyadarshana *et al.* 2015). Blue whales are found year-round off southern Sri Lanka, an area of very high shipping traffic (Randage *et al.* 2014). Eleven blue whales were known to be killed by ship strikes between 2010 and 2012 around Sri Lanka (Priyadarshana *et al.* 2015)2016 #855}(Priyadarshana *et al.* 2015), and at least two were known to be killed by ship strike in 2014 (Randage *et al.* 2014). A risk assessment found that a 15 nautical mile shift in shipping lanes associated with the Dondra Head traffic separation scheme (TSS) off the southern coast of Sri Lanka could reduce the risk of ship collisions with blue whale by approximately 95% (Priyadarshana *et al.* 2016). To date, Sri Lanka has not brought forward a proposal to the International Maritime Organization to modify the TSS.

Shipping lanes off San Francisco Bay, the Santa Barbara Channel, and Los Angeles/Long Beach in California overlap with important blue whale feeding areas, including identified biologically important areas (Irvine *et al.* 2014, Calambokidis *et al.* 2015). Nine blue whales were known to be killed and one seriously injured by ship strikes between 2007 and 2010 in California waters

(Carretta *et al.* 2013), although no blue whales were reported struck in the most recent five-year period available (2011-2015) (Carretta *et al.* 2017b). Model estimates of blue whale ship strike mortality off the U.S. West Coast range from 18 to 40 in a single year, significantly exceeding the Eastern North Pacific stock's (Rockwood *et al.* 2017) Potential Biological Removal (PBR) level (Rockwood *et al.* 2017), which is the number of individuals that could be taken as a result of human activities while still allowing a stock to recover to or remain within its optimum sustainable population range. In 2013, IMO-approved changes to the shipping lanes off California became effective (although they have yet to be codified in USCG regulations). The modifications to the TSSs were devised in collaboration between the USCG and NOAA staff at the Cordell Bank, Gulf of the Farallones, Monterey Bay, and Channel Islands National Marine Sanctuaries, and NMFS, and were designed, in part, to reduce the co-occurrence of whales and ships in the area of the TSS (USCG 2011).

Chilean blue whales are also known to be affected by ship strikes. Two blue whales were killed by ship strikes in recent years off southern Chile, an important feeding area for this population (Brownell *et al.* 2014, IWC 2017). In addition to co-occurrence between ships and whales, the whales' behavior may also contribute to their vulnerability to ship strikes. Some whales may be unaware of approaching vessels or may be involved in a vital activity (*i.e.*, mating or feeding) which may reduce the likelihood of an avoidance response (Nowacek *et al.* 2004, Silber *et al.* 2010). In one study, blue whales demonstrated little to no lateral movement in response to close approaches by large commercial ships; they sometimes undertook shallow "response dives," mainly when they were already at the surface, but these were considered likely to be ineffective for avoiding collisions (McKenna *et al.* 2015).

The size of a ship and its speed affect the likelihood and severity of the collision. Reviews of stranding and collision records indicate that larger ships (262.5 ft [80 m] or larger) and ships traveling at or above 14 knots (26 km/hour) have a much higher instance of fatal collisions with whales (Laist *et al.* 2001). Vanderlaan and Taggart (2007) demonstrated the relationship between ship speed and likelihood of fatal strikes; and Conn and Silber (2013) found that vessel speed restrictions diminished the risk of a fatal strike by 80-90%. Maritime industries, resource managers, and government agencies have proposed seeking ways to reduce the magnitude of the threat through technological solutions. However, there are no easy technological "fixes" and no technology exists, or is expected to be developed in the near future, that will completely ameliorate, or reduce to zero the chances of, ship strikes of large whales; and no single technology will fit all situations (Silber *et al.* 2009). Vessel speed restrictions have proved successful in reducing fatal whale strikes in some locations (Laist *et al.* 2014); however, reducing the co-occurrence of whales and vessels is likely the only sure means of reducing ship strikes, but it is not possible in many locations.

Increased ship traffic may increase the risk of blue whale ship strikes. The number of trips taken by ships globally has steadily increased in recent decades. Further, changes in the extent of polar sea ice could influence the number of vessels transiting blue whale habitat. For example, the opening of the Northwest Passage and Northern Sea Route may bring an increase in the volume of vessel traffic through polar waters and corresponding lower latitude waters including the Bering Sea, North Pacific, and North Atlantic Ocean blue whale habitat. If this occurs, the potential for negative impacts on blue whales and other cetaceans will increase.

As noted above, some blue whale populations are likely more vulnerable than others, based on differences in distribution relative to shipping traffic. Collisions with blue whales appear to be less common than for other large whale species (Laist *et al.* 2001, Jensen & Silber 2004), but many ship strikes go unreported or undetected, and estimates of mortality and serious injury should be considered minimums. Monnahan *et al.* (2015) evaluated the effects of ship strikes on Eastern North Pacific blue whales and found that they are not threatening the status of the population, although the authors suggest continued monitoring, given that the levels exceed the stock's PBR level. However, Rockwood *et al.* (2017) suggest caution in interpreting these results, since the conclusions of Monnahan *et al.* (2015) are based on lower numbers of annual ship strikes than were estimated by Rockwood *et al.* (2017), as well as other assumptions regarding environmental conditions and threats that may not be accurate. The impact to other populations is largely unknown. Additional information is needed to determine whether and to what extent ship strikes may be impeding the recovery of blue whales. Therefore, we consider ship strikes to be a potential threat to the species.

H.1.3 Entanglement in Marine Debris and Fishing Gear

Harmful marine debris consists of plastic garbage and other materials washed or blown from land into the sea, derelict fishing gear lost or abandoned by recreational and commercial fishers, and solid non-biodegradable floating materials (such as plastics) disposed of by ships at sea and from other sources. Whales may become entangled in debris or may ingest it. Plastics and other debris may be consumed incidental to normal feeding, and some marine species may actually confuse plastic bags, rubber, or balloons with prey and ingest them. The debris may cause a physical blockage in the digestive system, leading to internal injuries or other types of significant complications. Stomach obstruction caused by marine debris has not been documented in blue whales, but there are documented cases of ingestion of marine debris in both odontocete and mysticete species including, but not limited to, sperm, pygmy sperm (*Kogia breviceps*), and minke whales (*B. acutorostrata*) (Viale *et al.* 1991, Tarpley & Marwitz 1993).

Very few confirmed cases of blue whale entanglements in fishing gear (derelict or actively fished) have been documented. The first documented blue whale entanglement off the U.S. West Coast occurred off southern California in 2015. Although it was not possible to confirm the entangling gear to a specific fishery, based on the characteristics of the gear, it was considered to be from some type of deep-water trap/pot fishery (Carretta *et al.* 2017b). In 2016, four blue whales were reported entangled in fishing gear, two of which were confirmed to be entangled with Dungeness crab commercial trap fishing gear (NMFS 2017). Scarring has been observed on some blue whales, indicating past interactions with entangling gear (J. Calambokidis, pers. comm.), but no other information on these interactions is available. There have been no observed entanglements of blue whales in the California swordfish drift gillnet fishery in 26 years of observing the fishery (1990-2015) (Carretta *et al.* 2017a). There have been no observed hookings or entanglements of blue whales in the Hawaii-based deep-set or shallow-set longline fisheries since observer coverage began in 1994 (NMFS 2012, NMFS 2014). There have been no observed fishery mortalities or serious injuries of Western North Atlantic blue whales in U.S. fisheries (Waring *et al.* 2010).

In January 2013, an underwater photographer in Sri Lanka documented a whale with a net wrapped through its mouth, along the sides of its body, and wound around its tail (de Vos 2015).

This individual was noticeably thin, and unable to dive. Scarring patterns around its caudal peduncle indicated it had been entangled in this gear for an extended period. While attempts to approach the animal for rescue were futile (and dangerous for untrained personnel), the fate of this animal remains unknown, but given the extent of the entanglement, it is likely that this resulted in the death of the whale (de Vos 2015).

The small number of documented interactions likely represents only a fraction of interactions with fishing gear and more information is needed to determine if this is a significant cause of mortality. Data on entanglement and entrapment can be largely anecdotal and may not be reported systematically because many fisheries have low or no observer coverage. Like other large whale species, blue whales may break through or carry away fishing gear. Whales carrying gear may die later, become debilitated or seriously injured, or have normal functions impaired, but with no evidence of the incident recorded. Most whales killed by offshore fishing gear likely do not drift far enough to strand on beaches or to be detected floating in the nearshore corridor where most whale watching and other types of boat traffic occur (Heyning & Lewis 1990).

There are also non-lethal effects from entanglements that could affect recovery. Entanglement-related stress may decrease an individual's reproductive success or reduce its life span, which may in turn depress population growth. Additionally, injuries and entanglements that are not initially lethal may result in a gradual weakening of entangled individuals, making them more vulnerable to other causes of mortality (Kenney & Kraus 1993, Moore & Van Der Hoop 2012).

While the number of observed and reported entanglements is currently low, additional information is needed to determine whether entanglement in marine debris and fishing gear is impeding recovery of blue whales, given the uncertainty in the rate/frequency of entanglements and how this may vary across populations. Therefore, we consider this to be a potential threat to the species.

H.1.4 Anthropogenic Noise

Increasing human activity is leading to rising levels of anthropogenic underwater noise, which is changing the acoustic environment and impacting marine species and their habitat through acute, chronic, and cumulative effects (Radford *et al.* 2014). Potential impacts include altering important behavioral patterns, physiological effects such as hearing impairment or stress, and masking critical acoustic cues, and the results of these range from no effect to potentially significant effects on the fitness of marine mammals and their habitat, depending on the context and scale of the noise exposures (Southall *et al.* 2007, NOAA 2016). Below we briefly describe these types of effects, and then discuss several sources of anthropogenic noise: commercial vessels, oil and gas exploration and development, and military sonar and explosives.

Behavioral Response

Exposure to anthropogenic sound can result in a multitude of behavioral effects, ranging from no or minor effects (such as minor or brief avoidance or changes in vocalizations), to those being more potentially severe or sustained (*e.g.*, abandonment of higher quality habitat), and even, in certain circumstances, those that can combine with physiological effects or result in secondary responses that lead to stranding and death. Behavioral reactions to noise can vary not only across species and individuals but also for a given individual, depending on previous experience

with a sound source, hearing sensitivity, sex, age, reproductive status, geographic location, season, health, or its current activities or reproductive status (Richardson *et al.* 1995, Ellison *et al.* 2012, Costa *et al.* 2016). Responses might also vary depending on the sound source itself (*e.g.*, its frequency, whether it is moving or stationary), its proximity to the individual, exposure levels, propagation patterns in a particular area, or other factors (Richardson *et al.* 1995). Exposure to noise might disrupt communication, navigation, foraging, and social behavior, and animals may be displaced from habitat for short or long periods. Sensitization (increased behavioral or physiological responsiveness over time) to noise could also exacerbate other effects, while conversely habituation (decreased behavioral responsiveness over time) to chronic noise could result in animals remaining close to noise sources which could result in an increase in physical injuries or other effects (Richardson *et al.* 1995).

Most observations of marine mammal responses to anthropogenic sounds have been limited to short periods, and included the cessation of feeding, resting, or social interactions. Given the many variables involved and complex interaction with sources and animals (*i.e.*, overlap that varies over time, space, and frequency), it has been difficult to link specific behavioral responses to specific sound sources (Southall *et al.* 2007), although more recent controlled exposure studies have illustrated these connections and discerned the importance of more nuanced contextual factors such as the distance of the sound source or the behavioral state of the animal (Southall *et al.* 2016, Dunlop *et al.* 2017). In addition, it is difficult to quantify disturbance or overall, long-term impacts of exposure to noise (National Research Council 2005). (National Research Council 2003, National Research Council 2005), Lusseau and Bejder (2007) present data from three long-term studies illustrating the connections between disturbance from whale-watching boats and population-level effects in cetaceans, ranging from a 15% decrease in abundance (Bejder *et al.* 2006), to reduced reproductive success and increased stillbirths (Lusseau 2004), to bioenergetically modeled decreased energy intake and increased energy output (Williams *et al.* 2006). Following on the 2005 recommendations of the National Research Council, New (2014) outline a conceptual model of the relationships linking disturbance to changes in behavior and physiology, health, vital rates, and population dynamics. Further, scientists have developed state-space energetic models for several species (southern elephant seal, North Atlantic right whale, beaked whale, and bottlenose dolphin), that illustrate how specific information about anticipated behavioral changes or reduced resource availability can potentially be used to effectively forecast longer-term, population-level impacts (New *et al.* 2013a, New *et al.* 2013b, Schick *et al.* 2013, New 2014) when enough data are available. However, more work and data are needed before these sorts of models can be broadly applied for management use.

Masking

Masking, or “auditory interference,” occurs when sounds interfere with an animal’s ability to detect, recognize, or discriminate between acoustic signals of interest. Marine mammals use acoustic signals for a variety of purposes, which differ among species, but include intraspecific communications, navigation, prey detection, predator detection, social interactions, and acquisition of information about their environment (Aroyan *et al.* 2000, Erbe & Farmer 2000). Masking generally occurs when the interfering noise is louder than or comparable to, and of a similar frequency to, the sound that is being detected. Masking of important acoustic cues may affect social signaling and other sound-mediated functions (Erbe & Farmer 2000, Southall *et al.* 2007, McWilliams & Hawkins 2013).

The size of the “zone of masking” for a marine mammal is highly variable and depends on many factors that affect the received levels of the background noise and the sound signal (Richardson *et al.* 1995, Foote *et al.* 2004). Masking is influenced by the amount of time that the noise is present, as well as the spectral characteristics of the noise source. There are still many uncertainties regarding how masking affects marine mammals. For example, it is not known how loud acoustic signals must be for animals to recognize or respond to another animal’s vocalizations (National Research Council 2003). Richardson *et al.* (1995) argued that the maximum radius of influence of an industrial noise (including broadband low frequency sound transmission) on a marine mammal is the distance from the source to the point at which the noise can barely be heard. This range is determined by the hearing sensitivity of the animal and/or the background noise level present.

Animals may alter their behavior in response to masking. These behavior changes may include producing more calls, longer calls, or shifting the frequency of the calls. For example, studies indicate that North Atlantic right whales (Parks *et al.* 2009), blue whales (Di Iorio & Clark 2009), and killer whales (Holt *et al.* 2009) alter their vocalizations (call parameters or timing of calls) in response to background noise levels. Clark *et al.* (2009) developed a model to quantify changes in an animal’s acoustic communication space that result from changes in the characteristics of background noise. More recently, Redfern *et al.* (2017) assessed the impacts of chronic noise from commercial shipping in southern California on blue, fin, and humpback whales. However, the potential effects that masking may have on energetic costs or behavior are difficult to quantify and remain poorly understood.

Hearing Damage or Impairment

Exposure to anthropogenic noise, in some cases, may impact whales by damaging body tissue or the inner ear and hearing. Noise-induced threshold shifts are defined as increases in the threshold of audibility (*i.e.*, the sound has to be louder to be detected) of the ear at a certain frequency or range of frequencies (ANSI (American National Standards Institute) 1995, Yost 2000), *i.e.*, a loss in hearing sensitivity, and can be temporary or permanent. As mentioned previously, there are no direct measurements of the hearing abilities of baleen whales, but baleen whales’ sensitivity to low-frequency sound has been inferred using a variety of methods, including observed vocalization frequencies, observed reactions to playback of sounds, and anatomical analyses of the auditory system (NOAA 2016). Direct changes in hearing from noise exposure have only been measured in a laboratory on a limited number of species (odontocete and pinniped species only) and for only a handful of individuals within those species (Southall *et al.* 2007, Finneran 2015).

H.1.4.1 Commercial Vessel Noise

Sound emitted from large commercial vessels is a major source of chronic noise in the ocean today (Andrew *et al.* 2002, McKenna *et al.* 2012). Ship propulsion systems, generators, engine gearing, compressors, bilge and ballast pumps, as well as hydrodynamic flow surrounding a ship’s hull contribute to the introduction of noise into the water column by a large vessel. Propeller-driven vessels also generate noise through cavitation, which accounts for approximately 85% or more of the noise emitted by a large vessel (Richardson *et al.* 1995). Large vessels tend to generate sounds that are louder and at lower frequencies than small vessels (Polefka 2004).

Surface shipping is the most widespread source of anthropogenic, low frequency (0 to 1,000 Hz) noise in the oceans (Simmonds & Hutchinson 1996); (Simmonds & Hutchinson 1996, Southall *et al.* 2017). Ross (1976) estimated that between 1950 and 1975, shipping caused a rise in ambient noise levels of 10 decibels (dB) in the areas where shipping dominates, based on information about the total number of ships at sea, increases in average ship speed, propulsion power, and propeller tip speed. He predicted that this would increase by another 5 dB by the beginning of the 21st century. The National Research Council (2003) estimated that the background ocean noise level at 100 Hz has been increasing by about 1.5 dB per decade since the advent of propeller-driven ships, while others have estimated that the increase in background ocean noise is as much as 3 dB per decade in the Pacific Ocean (McDonald *et al.* 2006a). This rate may have slowed in certain periods in the last decade as patterns in global shipping have changed (Silber *et al.* 2015). Tyack (2008) noted that the increase in ambient noise from shipping likely reduced the detectable range of low frequency whale calls from many hundreds of kilometers in the pre-propeller ocean to tens of kilometers in many settings today. Clark *et al.* (2009) provided information on the effects of sound masking on mysticetes (*i.e.*, fin, North Atlantic right, and humpback whales) exposed to noise from ships and reported that, among other things, projected that whale call rates diminished in the presence of passing vessels. More recently, Redfern *et al.* (2017) examined the co-occurrence of blue, fin, and humpback whales with sound from commercial shipping off southern California and identified several regions of overlap where the acoustic habitat of these species was degraded by noise.

Although ship noise may result in negative behavioral, physiological, or auditory effects to blue whales, it is uncertain whether there are consequences at either individual or population levels, though more serious effects are more likely in areas where focused blue whale use or important habitat overlaps with areas of heavy ship traffic, such as shipping lanes. Additional research is needed to determine the degree to which anthropogenic noise from ships is a threat to blue whales.

H.1.4.2 Oil and Gas Exploration and Development

Oil and gas exploration and development activities occur in areas where blue whales are known to feed, breed, or migrate through. A number of activities associated with offshore energy development result in the introduction of underwater sound. Loud sounds from seismic acoustic surveys to locate undersea oil reserves may adversely affect marine mammals. Oil and gas exploration, including seismic surveys (e.g., towed airgun arrays), typically operate with marine mammal observers as part of required mitigation measures as required in incidental take permits issued for the activity (NOAA 2011). Baleen whales are known to detect the low-frequency sound pulses emitted by airguns and have been observed, in some cases, reacting to seismic vessels (Stone 2003, Di Iorio & Clark 2009, Blackwell *et al.* 2015). All seismic systems are towed behind ships (multiple ships may be involved), which themselves may impact whales. In addition, a variety of non-tactical sonar or sonar-like devices and technologies are used for purposes of geophysical research and studies of bathymetry that introduce sound energy into the water. Monitoring of fin whale calls during seismic surveys indicated that fin whales changed their vocalizations and may be displaced in response to these types of activities (Castellote *et al.* 2012).

Supply vessels, low-flying aircraft, construction work, and dredging introduce underwater noise

during various oil and gas exploration-related activities (Gales 1982, Greene 1987). Drilling for oil and gas generally produces low-frequency sounds with strong tonal components—frequency ranges in which large baleen whales communicate. However, recorded noise from one study of one drilling platform and three combined drilling production platforms found that noise was low volume and almost undetectable alongside the platform at Beaufort scale sea states of three or above. The strongest tones were at low frequencies, near 5 Hz (Richardson *et al.* 1995).

The intense pulses produced by seismic airgun acoustic surveys have the potential to cause direct harm at close distances, but subtler impacts can occur at larger distances. Cerchio *et al.* (2014) noted that the breeding display of humpback whales was disrupted by seismic survey activity in waters off northern Angola when seismic activities were conducted in relatively close proximity. Castellote *et al.* (2012) demonstrated that fin whale singing activity and call features were affected by the presence of seismic survey airgun operations in the western Mediterranean Sea. Blue whales in the St. Lawrence Estuary region called more frequently when seismic surveys using sparkers were being conducted than when they were not, likely indicating a response to the survey work (Di Iorio & Clark 2009). McDonald *et al.* (1995) documented that a blue whale ceased to call when it came within 10 km of a seismic survey. Gedamke and Robinson (2010) suggested a reasonable likelihood that whales a kilometer or more from seismic surveys could be potentially susceptible to a temporary impact to hearing, but also noted the importance of uncertainty and variability in risk assessments.

Although underwater noise generated by oil and gas exploration and development activities may have a detrimental effect on some blue whale social or acoustic behavior, none of the studies discussed above point to definitive consequences at either individual or population levels. Additional research is needed to determine the degree to which anthropogenic noise from oil and gas exploration and development is a threat to blue whales.

H.1.4.3 Military Sonar and Explosives

Military training activities by the U.S. Navy and navies of other countries regularly occur in the Atlantic Ocean, Gulf of Mexico, Mediterranean Sea, Indian Ocean, and Pacific Ocean. These activities include anti-submarine warfare, surface warfare, anti-surface warfare, mine warfare, missile, ship scuttling, and aerial combat exercises. In addition to these training activities, navies conduct ship shock trials, which involve detonations of high explosive charges to test combat readiness of a ship and its various on-board systems, and other activities involving the use of underwater explosives.

As part of its suite of training activities, the U.S. Navy employs low-, mid-, and high-frequency active sonar systems. The primary low-frequency active sonar system, of which only four are operated worldwide, is the Surveillance Towed Array Sensor System Low Frequency Active (SURTASS LFA) sonar system, which produces relatively loud sounds between 100–500 Hz, and has operated in the western and central Pacific Ocean. The U.S. Navy employs fleets across the globe utilizing several mid-frequency sonar systems that range from large systems mounted on the hulls of ships (*e.g.*, sonar devices referred to as AN/SQS-53 and -56), to smaller systems that are deployed from helicopters and fixed-wing aircraft, sonobuoys, and torpedoes. These sonar systems produce relatively loud, 1-10 kHz (or greater) frequency sounds (NMFS & U.S. Department of the Navy 2001, U.S. Department of the Navy 2008). The navies of other

countries also employ tactical active sonars across a range of frequencies.

Studies undertaken in 1997–98 pursuant to the U.S. Navy’s Low-Frequency Sound Scientific Research Program found only short-term responses to low frequency sound by mysticetes (fin, blue, and humpback whales), including changes in vocal activity and avoidance of the source vessel (Clark & Fristrup 2001, Croll *et al.* 2001a, Fristrup *et al.* 2003, Nowacek *et al.* 2007). The effect of sonar on blue whales remains uncertain; however, sonar associated with naval training activities might adversely affect blue whales in ways previously described (*i.e.*, hearing damage/impairment, behavioral response, and masking). Melcon *et al.* (2012) observed a decrease in blue whale calls in the Southern California Bight in the presence of mid-frequency active (MFA) sonar. (U.S. Department of the Navy 2008) More recently, Goldbogen *et al.* (2013) and DeRuiter *et al.* (2017) found that exposure to simulated MFA sonar interrupted blue whale feeding, especially when animals were in the deep feeding mode. Responses in this study were heavily dependent on the animal’s behavioral state and are illustrative of the complexity of understanding the impacts of sound on marine mammals.

Underwater detonations associated with military training activities include ship scuttling/sinking; missile, gunnery, and mine-neutralization exercises; and the disposal of unexploded ordnance and grenades. Underwater transmission of explosions includes an initial shock pulse followed by a succession of oscillating bubble pulses and sound fields of varying size. Whales very close to a large detonation might be killed or seriously injured; more distant whales might suffer lesser but nonetheless nearly debilitating injury (*e.g.*, tympanic membrane rupture, slight to extensive lung injury), while distant whales might experience physiological stress responses or behavioral disturbance, with severity of the reaction dependent on their distance from the detonation.

Various measures have been developed to reduce marine mammal exposure to sonar transmissions or underwater detonations during testing and training exercises. For example, exercises involving the SURTASS LFA sonar observe closures to reduce impacts in important areas and times, and also employ visual monitoring for marine mammals during daylight hours, passive sonar to listen for vocalizing marine mammals, and a high-frequency sonar that allows the U.S. Navy to detect large and most small cetaceans. If marine mammals are detected, the U.S. Navy is required to cease use of LFA sonar transmissions until whales have moved away from the area. Prior to and during MFA sonar operations and when explosives may be used, the Navy uses trained watch-standers to look for whales in the vicinity.

As noted above for other anthropogenic sound sources, military sonar and explosives may have a detrimental effect on some blue whale social or acoustic behavior, but there is currently no evidence of definitive consequences at either individual or population levels. Additional research is needed to determine to what degree anthropogenic noise from military sonar and explosives is a threat to blue whales.

H.1.5 Loss of Prey Base Due to Climate and Ecosystem Change

Climate change has received considerable attention in recent years, with growing concerns about warming ocean temperatures and influences on natural climatic oscillations, such as the Pacific Decadal Oscillation or El Niño and La Niña conditions. Evidence suggests that productivity in the North Pacific (Quinn II & Niebauer 1995, Mackas *et al.* 1998) and other ocean areas could

be affected by changes in the environment. Increases in global temperatures are expected to have profound impacts on arctic and subarctic ecosystems, and the ecosystem changes in these regions are projected to accelerate during this century (Aguilar *et al.* 2002, Anisimov *et al.* 2007). Climate and oceanographic change may affect habitat and food availability of blue whales. Whale migration, feeding, and breeding locations may be influenced by changing ocean currents and water temperatures, although the extent of potential change is not known. For example, decadal scale climatic regime shifts have been related to changes in zooplankton in the North Pacific (Brodeur & Ware 1992, Francis *et al.* 1998), and long-term trends of warming sea surface temperatures in the California Current Ecosystem have been linked to major changes in zooplankton abundance (Roemmich & McGowan 1995). Such changes could affect blue whales if prey resources undergo changes in occurrence or densities.

In the Southern Ocean, krill abundance fluctuates with oceanographic conditions, most notably variations in winter sea ice, and is susceptible to environmental change (Braithwaite *et al.* 2015). However, we cannot yet predict how inter-regional variability in the effects of climate change on factors that affect Antarctic krill will affect productivity, and while models indicate that krill biomass will decline with continued surface warming, there is low confidence in the predictions (Larsen *et al.* 2014).

In the North Atlantic, the distribution of copepods (a prey item of some North Atlantic krill species) has shown signs of shifting due to climate change (Hays *et al.* 2005). Doniol-Valcroze *et al.* (2012) noted that blue whale feeding depth and behavior varied across suitable habitats in the Gulf of St. Lawrence. These habitats were used preferentially at different times of the tidal cycle and appeared linked to known prey aggregation mechanisms. Oceanographic conditions might become less favorable to species such as Arctic krill (the preferred prey of blue whales) in higher latitudes like the Gulf of St. Lawrence (Walther *et al.* 2002, Hays *et al.* 2005). How this will influence prey availability and distribution and abundance of blue whales remains uncertain and should be closely monitored (Gavrilchuk *et al.* 2014).

In addition to changes in ocean temperatures, ocean acidification may adversely affect blue whale prey. As increasing amounts of carbon dioxide are released into the atmosphere, more carbon dioxide is absorbed by the oceans, which reduces its pH. Between 1750 and 1994, ocean surface pH decreased by 0.1 (with highest decreases at high latitudes), which corresponds with a 26% increase in acidity (Bindoff *et al.* 2007). Krill embryonic development (Kawaguchi *et al.* 2011), hatch rates (Kawaguchi *et al.* 2013), and post-larval metabolic physiology (Saba *et al.* 2012) are likely to be affected by increasingly acidic conditions. Projections of future acidification indicate large portions of Antarctic krill's present range will be at damaging levels, with potential collapse of the krill population there by 2300 (Kawaguchi *et al.* 2013).

The feeding range of blue whales is broadly distributed geographically and, consequently, it is likely that the blue whale may be more resilient to climate change, should it affect prey, than a species with a narrower range (Silber *et al.* 2016). A potential exception to this is the northern Indian Ocean population, which is restricted geographically and cannot move north as ocean temperatures increase (Thomas *et al.* 2016). However, blue whales have a relatively specialized diet (eating mainly krill), which might make them more vulnerable to climate change impacts on their prey, compared to species with generalist foraging strategies that might adapt to changing conditions by prey switching. The effects of climate-induced shifts in productivity, biomass, and

species composition of prey on the foraging success of blue whales have received little attention and more research is needed to understand possible impacts and the extent to which these impacts might impede blue whale recovery. Therefore, we consider loss of prey base due to climate and ecosystem change to be a potential threat to the species.

H.2 Stressors Evaluated but Determined Not to be Threats

H.2.1 Environmental Contaminants and Pollutants

The manner in which pollutants might negatively impact animals is not well understood, particularly in animals for which many of the key variables and physiological pathways are unknown (Aguilar 1987, O'Shea & Brownell 1994) such as blue whales. Organic chemical contaminants are generally considered less of a concern for mysticetes than odontocetes (Reijnders *et al.* 1999) because baleen whales generally feed at lower trophic levels with less opportunity for bioaccumulation, and contaminants are not considered to be primary factors in slowing the recovery of any populations of large whale species (O'Shea & Brownell 1994). O'Shea and Brownell (1994) indicated that concentrations of organochlorine and metal contaminants in tissues of baleen whales were low, and lower than other marine mammal taxa, in both local and global comparisons. They further stated that there was no firm evidence that levels of organochlorines, organotins, or heavy metals in baleen whales generally were high enough to cause toxic or other damaging effects. However, individual marine mammals with higher contaminant levels in tissues show increased susceptibility to infections, lesions, impairments, and even reproductive failure (De Guise *et al.* 1995, Moore *et al.* 1998, Aguilar *et al.* 2002, Jenssen *et al.* 2003).

Other studies confirm low levels of contaminants in some marine mammal populations. In a review of organochlorine and metal pollutants in southern Pacific marine mammals (Franciscana dolphins, *Pontoporia blainvillei*, from Argentina and pantropical spotted dolphins, *Stenella attenuata*, from the eastern tropical Pacific), Borrell and Aguilar (1999) noted that organochlorine levels suggested low exposure compared to other regions of the world. Although information is extremely scarce, concentrations of organochlorines in marine mammals occurring in the tropical and equatorial fringe of the Northern Hemisphere and throughout the Southern Hemisphere appear to be low. Aguilar *et al.* (2002) found that organochlorine concentrations in marine mammals off South America, South Africa and Australia were invariably low, with the lowest organochlorine concentrations in cetaceans found in the polar regions of both hemispheres. However, due to the systematic long-term transfer of airborne pollutants toward higher latitudes, it is expected that concentrations of organochlorines will increase in the Arctic and Antarctic in the future, warranting long-term monitoring of polar regions (Aguilar *et al.* 2002).

More specific to baleen whales, in a study of organochlorine exposure and bioaccumulation in the North Atlantic right whale, Weisbrod *et al.* (2000) noted that biopsy concentrations of organochlorines were an order of magnitude lower than concentrations in the blubber of seals and odontocetes. They concluded that there was no evidence to indicate that right whales bioaccumulate hazardous concentrations of organochlorines, and further noted that this was consistent with similar studies of baleen whales (Weisbrod *et al.* 2000).

Recently, researchers used the earwax of a male blue whale killed by a collision with a ship off the coast of California to investigate contaminants and hormone profiles. Earwax is continuously deposited throughout a blue whale's lifetime, but forms alternating light and dark layers at approximately 6-month intervals (Trumble *et al.* 2013). The light corresponds to periods in the blue whale's lifecycle when it is feeding, while the dark represents times of fasting and migration. Using the technique of analyzing blue whale earwax, Trumble *et al.* (2013) found markers of the stress hormone cortisol, growth-inducing testosterone, and contaminants such as pesticides, flame retardants, mercury. The reconstructed persistent organic pollutant (POP) profiles of the ship-struck male blue whale demonstrated that a substantial maternal transfer occurred during gestation and/or lactation, approximately 20% of its total lifetime burden (Trumble *et al.* 2013). A review by Wagemann and Muir (1984) highlighted similar maternal transfer of contaminants in a large number of marine mammals throughout the Northern Hemisphere. In blue whales in the Gulf of St. Lawrence, Canada, concentrations of PCBs, dichlorodiphenyl trichloroethane (DDT), and metabolites and several other organochlorine compounds were present at higher concentrations in the blubber of males relative to females; reflecting maternal transfer of these persistent contaminants from females into young (Metcalf *et al.* 2004). The impact from the chronic and acute POP exposure on baleen whales is largely unknown, but may potentially be positively correlated with cortisol.

Anthropogenic mercury is common in the environment and has received much attention among ecologists, environmental chemists, and toxicologists because of its ability to bioaccumulate and impair neurological development. Well-documented research involving humans reveals maternal transfer of mercury in utero and then to the neonate during lactation (Vieira *et al.* 2013). Mercury profiles in the ship-struck male blue whale do not mirror maternal transfer to the same degree as the POPs. Because this blue whale appeared to routinely traverse the coast of California (ship struck near Santa Barbara, CA), Trumble *et al.* (2013) speculated that pulse events of mercury accumulation may be associated with regional environmental and/or anthropogenic increases of mercury.

As noted above, marine mammals with higher contaminant levels may experience sublethal health effects, but there is currently no evidence that the effects are manifesting at a population or species level relevant to recovery and management decisions for blue whales.

Oil Spills and Spill Responses

Oil spills can injure or kill marine mammals. Dispersants used in oil spill response may also cause injury. Actual impacts would depend on factors such as the extent and duration of contact and the characteristics (*e.g.*, the type and degree of weathering) of the oil, but dermal exposure to oil and other chemicals could cause irritation, lesions, or burns, which may increase susceptibility to infection (Deepwater Horizon Natural Resource Damage Assessment Trustees 2016). Animals may also inhale, ingest, or absorb petroleum compounds or dispersants, which can injure their respiratory or gastrointestinal tracts or affect liver or kidney functions (Geraci 1990, Schwacke *et al.* 2014). Oil could also foul baleen, which would reduce its filtration efficiency during feeding (Lambertsen *et al.* 2005). Sub-lethal effects may include impaired health and reproduction and increased susceptibility to other diseases (Harvey & Dahlheim 1994). There are likely to be continuing effects even after clean-up has ended, such as exposure

to compounds in oil that persist in the environment and effects on habitat and prey resources (Deepwater Horizon Natural Resource Damage Assessment Trustees 2016).

We do not know the consequences of oil spills that have occurred in the blue whale's range, nor do we know the extent to which these or any future spills may affect blue whales. However, oil spill impacts to another baleen whale population suggest that, in some circumstances, impacts could be severe. Damage assessments following the April 2010 Deepwater Horizon major oil spill and subsequent response efforts estimated that 17% (95% confidence interval (CI) 7%–24%) of the small, resident Gulf of Mexico population of Bryde's whales was killed. Further, the spill was estimated to have caused 22% (95% CI 10%–31%) of the population's reproductive females to experience reproductive failure and 18% (95% CI 7%–28%) of the population to suffer adverse health effects (Deepwater Horizon Natural Resource Damage Assessment Trustees 2016).

Individual blue whales may experience potentially severe health effects from exposure to oil and other chemicals involved in spill response, but, unlike the small, resident population of Bryde's whales affected by the Deepwater Horizon oil spill, blue whales have a broad distribution and wide-ranging movements, which would be expected to lessen the population-, sub-species-, or species-level impact of such spills. For this reason, oil spills are not considered to be impeding the recovery of blue whales.

H.2.2 Disease

Harmful algal blooms are increasing in frequency and global distribution (Van Dolah 2000), potentially due to human activities' effects on the ecosystem such as nutrient loading from runoff or sewage (Sellner *et al.* 2003) or climate change (Moore *et al.* 2008). Biotoxins produced in the blooms have caused an increasing number of mass mortalities of cetaceans, pinnipeds, and mustelids (Gulland & Hall 2007, Landsberg *et al.* 2014). For example, in 1987 and 1988 fourteen humpback whales in the western North Atlantic died after eating mackerel that contained saxitoxin, a neurotoxin that can cause paralytic shellfish poisoning in humans (Geraci *et al.* 1989). Saxitoxin and another neurotoxin, domoic acid, were also detected in two and three humpback whales, respectively, that were associated with a 2003 unusual mortality event in Maine that killed 16 humpback whales, one fin whale, one minke whale, one pilot whale, and two individuals of unknown species (Gulland & Hall 2007). Domoic acid can cause amnesic shellfish poisoning in humans and has been linked to marine mammal and seabird mass-mortality events, particularly off the U.S. West Coast (reviewed in (Landsberg *et al.* 2014). Domoic acid is likely passed on to krill (Bargu *et al.* 2002), a common food of many baleen whales, including blue whales. Lefebvre *et al.* (2002) found that humpback and blue whales were exposed to the toxin and consumed domoic acid contaminated prey. However, it is unknown how the pervasiveness of biotoxins such as domoic acid affects baleen whales like the blue whale since data is limited. There is currently no evidence linking these toxins to acute impacts (including death) or chronic health problems in blue whales.

It is not known whether blue whales suffer from stress-induced bacterial infections similar to those observed in captive cetaceans (Buck *et al.* 1987). Brownell *et al.* (2007) described three types of skin lesions in blue whales in Chile: 1) resulting from cookie-cutter shark, *Isistius brasiliensis*, bites; 2) vesicular or blister lesions; and 3) a tattoo-like skin disease. The authors

note that the presence of blister lesions in 2006 and 2007 may indicate that this “disease” will be present in the population for a long time. It is unknown if these lesions contribute to mortality of blue whales frequenting Chilean waters, but the tattoo-like skin lesions, if shown to be a pox virus, could cause neonatal and calf mortality (Brownell *et al.* 2007). There is currently no evidence to suggest that the lesions pose a conservation risk to blue whales in general or the Chilean subspecies in particular. There is limited data on the status of morbillivirus or *Brucella* sp. infections in blue whales. Both of these diseases have caused significant impacts in other cetacean species (Guzmán-Verri *et al.* 2012, Van Bresseem *et al.* 2014) and should be monitored in stranded blue whales to provide baseline data for future assessments.

Based on the above information, disease is not currently considered to be limiting blue whale recovery.

H.2.3 Behavioral Disturbance from Vessel and Unmanned Aircraft System (UAS) Interactions

Blue whales are increasingly a target species for educational or recreational activities, including whale watching operations. As a result, individual and groups of blue whales may be closely accompanied by vessels on a regular basis. In addition to generating noise (discussed in Section I.H.1.4.1 above) and presenting a risk of vessel strike (discussed in Section I.H.1.2 above), vessel-based whale watching activities and other types of close approach may disturb the whales’ behavior.

Short-term behavioral changes made in response to whale watching activities have been documented in a number of cetacean species (Parsons 2012), including blue whales (Lesage *et al.* 2017). These include changes in surfacing and diving behavior, other active behavior, swim speed or direction, amount of time spent foraging or resting, and vocalization patterns (Parsons 2012). Animals’ reactions likely depend on a number of factors such as their behavioral context (e.g., feeding versus traveling), the number of vessels approaching, vessel approach distance, and duration of the encounter. It is unclear whether or how short-term changes translate to long-term effects on individuals or populations, such as changes in distribution (including abandonment of important feeding or breeding habitat, use of higher risk areas) or reduced reproductive success. There is currently no evidence indicating that these effects are detrimental to blue whales at the population level, but long-term monitoring would be important to understand any impacts of disturbance.

Unmanned aircraft systems (UAS), also known as model aircraft or drones, are a new way to obtain unique views of wildlife and natural landscapes. In recent years, there has been an increase in the recreational use of UAS, particularly to view marine mammals. Several federal statutes require scientists to obtain research permits to closely approach protected species of wildlife, such as marine mammals, but the lack of available information on the effects of UAS operations on these species has made it difficult to evaluate and reduce possible impacts (Smith *et al.* 2016). UAS can be disruptive to both people and animals if not used safely, appropriately, or responsibly. In the United States, the Federal Aviation Administration provides hobbyists

with some basic guidance for operating UAS to address safety and privacy concerns³. The U.S. National Park System has also recently prohibited the use of UAS in United States National Parks⁴ (some of which provide habitat to marine mammals).

Scientists and wildlife managers are concerned that acute or chronic disturbances of wildlife can impact the animals' health and fitness by disrupting migratory patterns, breeding, feeding, and sheltering. However, there is currently no evidence to suggest adverse effects to blue whale recovery from the use of UAS.

H.2.4 Research

Scientific research can involve close interactions with blue whales. In many countries, directed research activities typically require permits and are closely monitored to ensure any potential negative impacts are minimized. The potential for disturbance or harassment from observing or approaching whales for behavioral studies, photography, tagging, and data collection (including biopsy samples collected for health and genetic analysis) is likely minimal and is far outweighed by the benefits of gaining information that could prove critical in helping manage and recover the species.

The use of UAS offers a new method for scientific researchers and emergency responders to obtain important information about marine mammals that can further support the conservation of these protected species. In the United States, NMFS continues to evaluate scientific research permit applications to ensure that the potential hazards of UAS use do not outweigh the benefits, and other types of public activity that have the potential to negatively impact protected marine species⁵.

We address the effects of research activities that do not involve the direct study of blue whales in other subsections of the Plan's threats analysis section, such as vessel interactions, anthropogenic noise, contaminants and pollutants, oil and gas exploration, and military sonar and explosives.

H.2.5 Predation and Natural Mortality

While there are records of killer whale attacks on blue whales and some shark species likely take individual blue whales, there is no evidence that this is a significant threat to the species.

H.2.6 Competition for Resources

Many sympatric baleen whale species target similar resources and may interact ecologically. Sympatric baleen whales may develop species-specific foraging behaviors to reduce competition; for example, Friedlaender *et al.* (2009) found that humpback and minke whales may partition resources vertically. However, off central California, sympatric blue and fin whales were not

³ <https://www.faa.gov/uas/>

⁴ <https://www.nps.gov/orgs/aviationprogram/upload/unmanned-aircraft-in-national-parks.pdf>

⁵ <https://www.fisheries.noaa.gov/national/marine-life-viewing-guidelines/permitting-scientific-research-using-small-unmanned>

found to feed at significantly different depths when targeting the same prey patch (Friedlaender *et al.* 2015).

Prey distribution and abundance is a primary driver of baleen whale distribution and feeding behavior (Croll *et al.* 2005, Friedlaender *et al.* 2006). While competitive interactions are possible, there is no basis to assume that competition for food among baleen whales is affecting the blue whale's population trend and abundance. Rather, the presence of food may influence the distribution or occurrence of blue whales in certain areas. For example, when large predatory fish declined in the Northeast Atlantic, their prey (herring, capelin, shrimp, and snow crab) eventually increased in abundance (Lilly 1991, Berenboim *et al.* 2000, Garrison & Link 2000, Koeller 2000, Lilly *et al.* 2000, Bundy 2005), and as a consequence, prey needs increased, resulting in changes in the trophic structure in the Gulf of St. Lawrence. During this time, blue whale sightings in the northwestern Gulf of St. Lawrence declined. Gavrilchuk *et al.* (2014) indicated that the effects on non-target species following the collapse of groundfish fishery in the early 1990s might be less favorable for the blue whale, which have a relatively species-specific diet, compared to baleen whales with generalist foraging strategies. Thus, the enhanced competition for krill between blue whale and species at various trophic levels following changes in the structure and dynamic of the northern Gulf of St. Lawrence ecosystem appeared sufficient to account for the decline in food availability for blue whales and their departure from the area (Comtois *et al.* 2009, Comtois *et al.* 2010).

Krill fisheries have the potential to reduce blue whale prey. In the Northern Hemisphere, krill fisheries occur around Japan and off British Columbia, while krill harvest off the U.S. West Coast and Alaska are prohibited. Nicol *et al.* (2012) consider the Antarctic likely to be the main source of krill harvest in the future. In the Southern Ocean, biomass of *Euphausia superba* is currently estimated to be 379 million tons (Atkinson *et al.* 2009). The Commission for the Conservation of Antarctic Marine Living Resources sets precautionary harvest limits to ensure a sustainable fishery while minimizing impacts on the ecosystem. From a catch limit of 620,000 tons, an estimated 260,151 tons of *E. superba* were caught in the Southern Ocean in 2016 (CCAMLR 2017). Overfishing of krill could impact the availability of blue whale prey, but this is considered unlikely given management measures in place.

In summary, there is limited information on blue whale competition for prey with sympatric species, but there is also no evidence that competition with sympatric species is a threat for any large whale species. Indeed, Clapham and Brownell (1996) reviewed evidence for interspecific competition in baleen whales and concluded that it was not possible to establish inter-specific competition as an important factor in the population dynamics of large whales. In addition, changes in trophic structure may affect the prey availability for blue whales, but there is no evidence that this is a threat to blue whales. Krill fisheries, mainly in the Antarctic, compete with blue and other whales for prey, but given the precautionary management and harvest limits, are not considered to be limiting blue whale prey. Therefore, competition for resources is not considered to be a threat to blue whale recovery.

I. Conservation Measures

The blue whale is protected under both the ESA and the MMPA. The blue whale was listed as “endangered” throughout its range under the precursor to the ESA, the Endangered Species

Conservation Act of 1969 (35 FR 8491; June 2, 1970), and remained on the list of threatened and endangered species after the passage of the ESA in 1973. The blue whale is automatically designated as depleted under the MMPA due to its status as endangered under the ESA. The Secretary of Commerce (as delegated to NMFS) administers the ESA for most endangered marine species, including blue whales. NMFS has lead responsibility for developing and implementing a recovery program for this species.

Blue whales are also listed in Appendix I of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and the Convention on the Conservation of Migratory Species of Wild Animals (CMS). The CITES classification is intended to ensure that commercial trade in the products of blue whales does not occur across international borders and are not introduced from the sea. The CMS, also known as the Bonn Convention, aims to conserve terrestrial, aquatic and avian migratory species throughout their range.

Through amendments to the Schedule to the ICRW, blue whale received complete protection from whaling throughout the North Atlantic Ocean in 1955, the North Pacific Ocean in 1966, and the entire Southern Hemisphere in 1968 (Best 1993). In 1982, the IWC set catch limits for all commercial whaling to zero. These catch limits went into effect in 1986, beginning a commercial whaling moratorium that remains in effect today. However, Norway and Iceland continue to commercially take whales under formal objections or reservations to the moratorium. The Russian Federation has also objected to the moratorium but does not exercise it. There have been no recorded commercial catches of blue whales since 1978 (Branch *et al.* 2008). Japan conducts special permit whaling (also called scientific whaling) under Article VIII of the ICRW but has no recorded catches of blue whales.

II. RECOVERY STRATEGY

Commercial whaling was the main cause of blue whales' historical decline, and is not a current operative threat only because an international moratorium remains in place. Therefore, a primary strategy of this Revised Recovery Plan is to maintain the international ban on commercial hunting that was instituted in 1986. Additionally, this Plan provides a strategy to improve our understanding of how potential threats may be limiting blue whale recovery. Finally, this Plan provides a research strategy to obtain data necessary to determine blue whale taxonomy, population structure, distribution, and habitat, which can then inform estimation of population abundance and trends. After the populations and their threats are more fully understood, this Plan will be modified to more specifically include actions to minimize any threats that are determined to be limiting recovery.

Because blue whales move freely across international borders, it would be ineffective to confine recovery efforts to U.S. waters, and this Plan stresses the importance of a multinational approach to management. As demonstrated by work on humpback whales involving a number of researchers from different countries (SPLASH, (Calambokidis *et al.* 2008a) Year of the North Atlantic Humpback, (Palsbøll *et al.* 1997, Smith *et al.* 1999), considerably more information is gathered for management of whale populations when research is conducted on the basis of biological, rather than political divisions and through multilateral cooperation.

III. RECOVERY GOALS, OBJECTIVES, AND CRITERIA

A. Goals

The goal of this Revised Recovery Plan is to promote recovery of blue whales to a level at which it becomes appropriate to “downlist” the species from endangered to threatened status, and ultimately to “de-list”, or remove the species from the list of Endangered and Threatened Wildlife and Plants, under the provisions of the ESA. The Act defines an “endangered species” as “any species which is in danger of extinction throughout all or a significant portion of its range.” A “threatened species” is defined as “any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.”

B. Management Units

For purposes of this Revised Recovery Plan, we identify nine blue whale management units:

1. Northern subspecies (*B. m. musculus*) - North Atlantic population
2. Northern subspecies (*B. m. musculus*) - Eastern North Pacific population
3. Northern subspecies (*B. m. musculus*) - Western/Central North Pacific population
4. Northern Indian Ocean subspecies (*B. m. indica*)
5. Pygmy subspecies (*B. m. breviceauda*) - Madagascar population
6. Pygmy subspecies (*B. m. breviceauda*) - Western Australia/Indonesia population
7. Pygmy subspecies (*B. m. breviceauda*) - Eastern Australia/New Zealand population
8. Chilean subspecies (*B. m. unnamed subsp.*)
9. Antarctic subspecies (*B. m. intermedia*)

As described in Section I.B.3, blue whale subspecific taxonomy has not been fully resolved and is an area of active research. The Society of Marine Mammalogy's (SMM) Taxonomy Committee currently recognizes five subspecies: *B. m. musculus*, *B. m. intermedia*, *B. m. indica*, *B. m. breviceauda*, and an unnamed subspecies off Chile (Committee on Taxonomy 2016). However, there are continuing discussions in the scientific literature regarding whether *B. m. indica* is a subspecies separate from *B. m. breviceauda*, and whether the blue whales off Chile are a separate sub-species. We identify at least one management unit (and associated recovery criteria) for each of the five subspecies recognized by SMM.

We also identify management units within two subspecies. In the Northern subspecies, *B. m. musculus*, we identify one North Atlantic and two North Pacific management units. In general, individuals from different ocean basins are unlikely to interbreed when mature, and if a basin population were extirpated, the area would likely not be recolonized in a time period that is meaningful for management purposes (Angliss et al. 2002). As described in Section I.D.1, it is unclear whether blue whales in the eastern and western portions of the North Atlantic Ocean belong to the same population. Until separation is more strongly supported, for this Revised Recovery Plan we consider blue whales in the North Atlantic to comprise one management unit, based on the IWC blue whale stock definition and the current understanding that there is only one blue whale song type in the North Atlantic. The IWC also considers blue whales in the North Pacific to be one stock, but we define two management units there based on multiple lines of evidence (song types, length-frequency data, and movement data from satellite tags and photo-identification), summarized in Section I.E.1, indicating there are at least eastern and western/central populations. In the pygmy blue whale subspecies, *B. m. breviceauda*, we identify three management units corresponding with "acoustic populations," following recommendations of the IWC (IWC 2016a).

Despite the uncertainties, the delineation of these nine units reflects our current understanding of blue whale taxonomy and population structure and we consider them to be the appropriate units to conserve. We consider recovery of all nine units to be important for achieving geographic and ecological representation of blue whales in the world's oceans, and to ensure conservation of the breadth of blue whale' genetic variability. If, based on additional research, blue whale subspecific taxonomy is revised, our understanding of population structure changes, or we learn more about whether recovery of all units is necessary for the long-term viability of the species, these management units and their associated species-level recovery criteria should be revised.

C. Objectives and Criteria

The two main objectives for blue whales are to 1) increase blue whale resiliency and ensure geographic and ecological representation by achieving sufficient and viable populations in all ocean basins and in each recognized subspecies, and 2) increase blue whale resiliency by managing or eliminating significant anthropogenic threats. A prerequisite to achieving these objectives is obtaining sufficient data to determine whether they have been met, so many of the recovery actions in Section IV below focus on data collection.

The recovery criteria take two forms: 1) those that reflect the status of the species itself and 2) those that indicate effective management or elimination of threats. The criteria include specific targets for each of nine management units to support the objectives of species' viability (e.g.,

resiliency, redundancy, and representation).

Population viability analyses (PVAs) or other quantitative assessments for predicting a species' or population's future status can be useful in evaluating extinction risk. PVAs have been used in some recent marine mammal status reviews (e.g., (Krahn *et al.* 2004, Oleson *et al.* 2010), but not in others (e.g., (Boveng *et al.* 2009, Boveng *et al.* 2013, Bettridge *et al.* 2015), and are a component of several other large whale recovery plans (e.g., (NMFS 2010a, NMFS 2010b, NMFS 2011, NMFS 2013). Although not required to meet the criteria below, should sufficient data become available, a quantitative PVA demonstrating a low probability of extinction over a reasonable timeframe (e.g., scaled to blue whale generation time, (Taylor *et al.* 2007) could be used to further support downlisting or delisting decisions.

C.1 Downlisting Objectives and Criteria

Blue whales may be considered for reclassifying to threatened when all of the following have been met.

Objective 1: Increase blue whale resiliency and ensure geographic and ecological representation by achieving sufficient and viable populations in all ocean basins and in each recognized subspecies.

Criteria: In each of the nine management units:

1. The minimum abundance is:
 - a. North Atlantic: 2,000 whales
 - b. Eastern North Pacific: 2,000 whales
 - c. Western/Central North Pacific: 2,000 whales
 - d. Northern Indian Ocean: 500 whales
 - e. Madagascar: 2,000 whales
 - f. Western Australia/Indonesia: 2,000 whales
 - g. New Zealand: 500 whales
 - h. Chilean: 2,000 whales
 - i. Antarctic: 2,000 whales

The 2015 status review for humpback whales (Bettridge *et al.* 2015) reviews scientific literature on the relationship between population size and extinction risk. Bettridge *et al.* (2015) describe widespread scientific agreement that total population sizes greater than approximately 2,000-2,500 are sufficiently large to maintain genetic diversity and protect against the effects of demographic and environmental stochasticity and natural catastrophes. For management units that are known or believed to have had at least 2,500 individuals prior to the onset of commercial whaling, we adopt the lower end of this range (2,000) as a downlisting criterion, and the upper end (2,500) as a delisting criterion (see Section III.C.2 below). For management units that are known or believed to have had fewer than 2,500 individuals prior to the onset of commercial whaling, we specify lower minimum abundance criteria: 500 individuals as a downlisting criterion and 1,000 individuals as a delisting criterion. As noted in the humpback whale status review (Bettridge *et al.* 2015), an abundance of more than 500 individuals provides protection against the genetic risks of inbreeding and against moderate levels of environmental

variance. For both sets of minimum abundance criteria, approximately half of the total population of each management unit is assumed to be mature individuals capable of reproduction (Taylor *et al.* 2007).

Based on information summarized in the Background sections of this Revised Recovery Plan, the pre-exploitation abundances of blue whales in the (a) North Atlantic, (c) Western/Central North Pacific, (e) Madagascar, (f) Western Australia/Indonesia, and (i) Antarctic management units very likely exceeded 2,500, while pre-exploitation abundance in the (d) Northern Indian Ocean management unit was very likely below 2,500. Pre-exploitation abundance estimates for the (b) Eastern North Pacific (2,210 individuals, 95% Bayesian credible interval 1,823-3,721; Monnahan *et al.* 2015) and (h) Chilean (2,100-3,600 individuals, Jackson 2016; or 2,000-6,200 individuals, Williams *et al.* 2011) management units are not clearly above the 2,500 threshold, but catch information supports applying this higher threshold. In the Eastern North Pacific, an estimated 3,411 whales (95% range 2,593-4,113, (Monnahan *et al.* 2015) were caught from 1905-1971, while 4,288 were caught off Chile from 1908-1967 (Williams *et al.* 2011). There are no estimates of pre-exploitation abundance for blue whales around (g) New Zealand, but Brownell *et al.* (2015) suggest that Soviet catches might provide a minimum estimate, since most of the catches were made during a relatively short period of time. Catches appear to be less than 2,500 individuals, based on data presented in Branch *et al.* (2007b).

Although we use pre-exploitation abundances to assign management units to one of two sets of minimum abundance down- and de-listing criteria, the minimum abundance thresholds themselves are not linked to historical abundance. Because of this, a management unit meeting its minimum abundance criterion might remain a fraction of its pre-exploitation numbers. This is particularly true for Antarctic blue whales, which numbered in the hundreds of thousands prior to commercial whaling. However, management units meeting this criterion are not likely to be at risk of extinctions due to low abundance alone. Continued management (beyond the ESA down- and de-listing goals considered here) under the MMPA and the IWC are expected to encourage continued population growth and ultimately achievement of optimum sustainable population levels (or greater), which would further strengthen the likelihood that blue whales would fully reclaim their previous ecosystem function.

2. The trend in abundance is stable or increasing over the most recent 30-year period assessed.

Trends in abundance are an important measure of a population's viability. This criterion could be met for a particular subspecies or population if a peer-reviewed analysis demonstrates that population is stable or increasing over 30 years, which is the estimated blue whale generation length under pre-disturbance conditions with an assumed stable population (Taylor *et al.* 2007). The specific 30-year time period may differ by management unit, depending on when abundance surveys or analyses have been conducted.

Objective 2: Increase blue whale resiliency by minimizing anthropogenic threats.

Criteria: In each of the nine management units, anthropogenic threats have been identified and demonstrably minimized. Specifically, the factors in 4(a)(1) of the ESA described below have been addressed:

Factor A: The present or threatened destruction, modification, or curtailment of a species' habitat or range.

- Effects of anthropogenic noise, entanglement in marine debris and fishing gear, and reduced prey abundance due to climate change have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

Factor B: Overutilization for commercial, recreational, or educational purposes.

- The international ban on commercial hunting has been maintained.
- Any subsistence or scientific hunting that has the potential to overutilize the species is restricted to levels that are sustainable, precautionary, and in accordance with the advice of the IWC's Scientific Committee.

Factor C: Disease or Predation.

- There is no information at this time indicating disease or predation is a threat to blue whale recovery.

Factor D: The inadequacy of existing regulatory mechanisms.

- (Hunting is addressed under Factor B.)

Factor E: Other natural or manmade factors affecting its continued existence.

- Ship strikes have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.
- Entanglement with fishing gear has been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

As noted in Section I.H above, there are no currently identified threats to the species, but there are numerous potential threats. This recovery criterion, therefore, first requires that additional research be conducted to understand whether and to what extent any of the potential threats are impacting the recovery of each blue whale management unit. For any potential threat that is found to be impeding blue whale recovery, measures must be taken to minimize or eliminate its impact.

C.2 Delisting Objectives and Criteria

Blue whales will be considered for delisting when all of the following criteria are met.

Objective 1: Increase blue whale resiliency and ensure geographic and ecological representation by achieving sufficient and viable populations in all ocean basins and in each recognized subspecies.

Criteria: In each of the nine management units:

1. The minimum abundance is:

- a. North Atlantic: 2,500 whales
- b. Eastern North Pacific: 2,500 whales
- c. Western/Central North Pacific: 2,500 whales
- d. Northern Indian Ocean: 1,000 whales
- e. Madagascar: 2,500 whales
- f. Western Australia/Indonesia: 2,500 whales
- g. New Zealand: 1,000 whales
- h. Chilean: 2,500 whales
- i. Antarctic: 2,500 whales

See Section III.C.1 above for a justification for the selection of a minimum abundance criterion for each management unit. Based on the minimum abundance delisting thresholds described here, management units meeting this criterion are not likely to be at risk of extinction within the foreseeable future due to low abundance alone.

2. The trend in abundance is stable or increasing over the most recent 30-year period assessed.

Objective 2: Increase blue whale resiliency by ensuring anthropogenic activities are not contributing to the species being in danger of extinction within the foreseeable future throughout all or a significant portion of its range.

Criteria: In each of the nine management units, anthropogenic threats have been identified and demonstrably minimized or eliminated, and there is information indicating they are not contributing to the species being in danger of extinction within the foreseeable future throughout all or a significant portion of its range. Specifically, the factors in 4(a)(1) of the ESA described below have been addressed:

Factor A: The present or threatened destruction, modification, or curtailment of a species' habitat or range.

- Effects of anthropogenic noise, entanglement in marine debris and fishing gear, and reduced prey abundance due to climate change have been evaluated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

Factor B: Overutilization for commercial, recreational, or educational purposes.

- The international ban on commercial hunting has been maintained.
- Any subsistence or scientific hunting that has the potential to overutilize the species is restricted to levels that are sustainable, precautionary, and in accordance with the advice of the IWC's Scientific Committee.

Factor C: Disease or Predation.

- There is no information at this time indicating disease or predation is a threat to blue whale recovery.

Factor D: The inadequacy of existing regulatory mechanisms.

- (Hunting is addressed under Factor B.)

Factor E: Other natural or manmade factors affecting its continued existence.

- Ship strikes have been evaluated, and, if determined to be impeding blue whale

- recovery, measures have been taken to minimize effects.
- Entanglement with fishing gear has been investigated, and, if determined to be impeding blue whale recovery, measures have been taken to minimize effects.

As noted in Section I.H above and in the description of downlisting criteria above, there are no currently identified threats to the species, but there are numerous potential threats. This recovery criterion, therefore, first requires that additional research be conducted to understand whether and to what extent any of the potential threats are impacting the recovery of each blue whale management unit. For any potential threat that is found to be impeding blue whale recovery, measures must be taken to minimize or eliminate its impact, and there must be information indicating it is no longer limiting blue whale recovery.

IV. RECOVERY PROGRAM

A. Recovery Action Outline

Recovery actions in this outline are not in order of priority. Priorities are identified in the Implementation Schedule (Section V below). More detail about each recovery action appears in the Recovery Action Narrative in Section IV.B below. Unless otherwise indicated, the relevant “site” for each recovery action is throughout all nine management units.

1.0 Coordinate State, Federal, and International Measures to Maintain International Regulation of Whaling for Blue Whales

2.0 Determine Blue Whale Taxonomy, Population Structure, Occurrence, Distribution, and Range

- 2.1 *Conduct studies to investigate population discreteness and population structure of blue whales.*
- 2.2 *Conduct studies to assess blue whale occurrence, daily and seasonal movements, and inter-area exchange.*
- 2.3 *Support the development of models to yield robust predictions of blue whale distribution.*

3.0 Estimate Population Size and Monitor Trends in Abundance

- 3.1 *Establish collaborative agreements with relevant national governmental bodies and scientific institutions to develop plans for estimating abundance and monitoring trends in abundance.*
- 3.2 *Conduct surveys to estimate blue whale abundance and monitor trends in abundance worldwide.*
- 3.3 *To the extent possible, work with appropriate government agencies in other countries to develop and maintain blue whale photo-identification programs, and educate the public about contributing information about dead, entangled, or ship-struck whales, to continue to support or establish international databases.*

4.0 Identify, Characterize, Protect, and Monitor Habitat Important to Blue Whale Populations

- 4.1 *Characterize blue whale habitat.*
- 4.2 *Monitor important habitat features and blue whale use patterns to assess potentially detrimental shifts in habitat features that might reflect disturbance or degradation of habitat.*
- 4.3 *Promote measures to identify and protect important habitat throughout the species’ range.*

5.0 Investigate Potential Human-Caused Threats and, Should They Be Determined to Be Limiting Blue Whale Recovery, Take Steps to Minimize Their Occurrence and Severity

- 5.1 *Anthropogenic noise*

- 5.1.1 Conduct studies to determine whether anthropogenic noise is adversely affecting blue whale distribution, behavior, or vital rates.
- 5.1.2 Take steps to minimize anthropogenic noises that are found to be detrimental to blue whales.
- 5.2 *Vessel collisions*
 - 5.2.1 Maintain database of known ship strikes of blue whales.
 - 5.2.2 Review photographic databases for evidence of injuries to blue whales from ship strikes to better characterize these events.
 - 5.2.3 Conduct studies to identify areas of high risk for blue whale ship strikes.
 - 5.2.4 Conduct studies to determine whether collisions with ships are adversely affecting blue whale abundance and recovery.
 - 5.2.5 If risk of ship strike is determined to be significant, work with mariners, the shipping industry, appropriate state, federal, international agencies to develop and implement measures to reduce the threat of ship strikes, and report strikes, should they occur. Assess the effectiveness of ship strike measures and adjust, as appropriate.
- 5.3 *Marine debris*
 - 5.3.1 Determine if areas exist in which blue whale habitat and significant deposits of marine debris coincide.
 - 5.3.2 Develop and implement means to reduce fishing gear loss, and improve the reporting of lost gear.
 - 5.3.3 Take steps to reduce or eliminate sources of marine debris if it is found to be detrimental to blue whales.
- 5.4 *Fishing gear entanglement*
 - 5.4.1 Review data on blue whale entanglement in fishing gear, including reports from fishermen and fishery observers.
 - 5.4.2 Review photographic databases for evidence of injury to blue whales caused by fishing gear entanglement.
 - 5.4.3 If found to be detrimental to blue whales, develop means to reduce entanglements, and enhance the effectiveness of disentanglement of blue whales in fishing gear. Determine whether measures to reduce entanglements are effective.
- 5.5 *Climate change*
 - 5.5.1 Conduct research and perform analyses to understand whether climate change is adversely affecting blue whale distribution, abundance, or fecundity.
 - 5.5.2 Take steps to minimize climate change impacts if found to be detrimental to blue whales.

6.0 Maximize Efforts to Acquire Scientific Information from Dead, Stranded, and Entangled or Entrapped Blue Whales

- 6.1 *Make every effort to necropsy stranded blue whales to determine cause of death.*
- 6.2 *Review, analyze, and summarize data on dead blue whales and causes of death.*
- 6.3 *Establish reliable sources of funding for necropsy (including sites to conduct the necropsy), tissue collection, and analysis efforts.*

B. Recovery Action Narrative

Recovery actions in this outline are not in order of priority. Priorities are identified in the Implementation Schedule (Section V below). Unless otherwise indicated, the relevant “site” for each recovery action is throughout all nine management units.

1.0 Coordinate State, Federal, and International Measures to Maintain International Regulation of Whaling for Blue Whales

Work through the IWC and other relevant international bodies or agreements to ensure that the moratorium on blue whale commercial hunting is maintained. The international regulation of whaling is vital to the recovery of whale populations. A full assessment of present status has not been conducted and there is a lack of sufficient information on blue whale population size, trends, and structure to justify the resumption of exploitation.

For meaningful estimates, it will be necessary for the United States to promote and participate in cooperative surveys with other countries. A primary goal should be to foster an international collaboration and cooperation in the study and protection of blue whales.

This action is expected to involve staff time (see Table 2); costs associated with travel to IWC meetings, for example, is included in other large whale recovery plans (NMFS 2010a, NMFS 2010b).

2.0 Determine Blue Whale Taxonomy, Population Structure, Occurrence, Distribution, and Range

Existing knowledge of the population structure of blue whales is insufficient, and a more comprehensive understanding is essential for developing strategies to promote recovery and for classifying the populations according to their recovery status. Subspecies taxonomy remains unresolved, but it is almost certain that the global listing inadequately captures the current population structure.

Analyses should be directed at examining trends over time, and attempts should be made to correlate observed changes in whale populations with physical, biological, or human-induced changes in the environment. As much as possible, data should be presented in peer-reviewed journals and other open publications to ensure that research programs benefit from regular peer scrutiny.

2.1 Conduct studies to investigate population discreteness and population structure of blue whales.

This is among the highest priority actions in this plan because it would improve

understanding and management of the species; however, it cannot be given a Priority 1 ranking because basic studies of population structure would not contribute to preventing extinction.

Researchers equipped to sample other whale species (*e.g.*, fin, right, and humpback whales), particularly in more remote areas where blue whale samples have not previously been obtained, should be encouraged to take advantage of opportunities to obtain samples from blue whales, on an opportunistic basis. Collaborative efforts with agencies, organizations, and researchers in other nations will be necessary to obtain sufficient samples over sufficiently broad areas for conclusions regarding population structure. Standardized sampling protocols and analytical procedures should be developed and used. Genetics work should be complemented by a thorough review of existing data from whaling and other sources. This might include investigation of geographical variation in morphology and acoustics of blue whales. New methods examining stable isotopes and fatty acids have also proven effective auxiliary data in cases where there is population mixing (*i.e.*, genetically distinct groupings mixing spatially on feeding grounds). Any such methods that can assist in resolving population structure should be encouraged.

Costs in the implementation schedule (Table 2) include equipment used and analysis of genetic and other data, as well as costs for opportunistic data collection during ship-based surveys (see 3.2 for additional costs associated with dedicated ship-based surveys).

2.2 Conduct studies to assess blue whale occurrence, daily and seasonal movements, and inter-area exchange.

Telemetry studies using satellite-linked and radio tags can be useful in investigating patterns and ranges of daily, seasonal, and longer-term movements of individual blue whales. Tagging studies are also useful for determining habitat use and identifying locations of any unknown breeding and feeding grounds. Exchange rates between populations might also be addressed to some degree by telemetry studies. Long-term efforts at photo-identification should also be encouraged to continue and opportunistic efforts to photo-document sightings could contribute to knowledge of movements by individuals and residency times. A central repository for blue whale photographs, and a system for curating and analyzing them, should be established. Photographs should be supplemented whenever possible by tissue samples (whether sloughed skin or biopsies) for DNA fingerprinting (Amos & Hoelzel 1992).

Like other baleen whales, blue whales make low-frequency vocalizations that are audible over long distances. Detection of underwater calls of a number of cetacean species has been used in studies of cetacean occurrence and distribution. Large whale species tend to vocalize rather often, depending on behavioral, social, and other contexts, and the technique has proven effective in determining presence (but not absence) of large whales. With arrays of three or more detection elements (to enable “triangulation”), information on distribution, and in some cases abundance, and habitat use can be determined. Bottom-mounted recording systems have been used to detect baleen whales with considerable success. Particularly when considered relative to vessel- or aircraft-based observer surveys, passive acoustic techniques are highly cost-effective, are less limited by poor weather conditions and thus are able to make observations more consistently, and

pose fewer risks to human observers. Therefore, and particularly with regard to their cost-effectiveness, passive acoustic studies should be used, if feasible, in attempts to determine blue whale occurrence, distribution, abundance, trends in abundance, and possibly response to some threats (*e.g.*, anthropogenic underwater noise).

The distribution and habitat use of marine mammals, including blue whales, can be studied along predetermined transect lines using autonomous gliders equipped with instrumentation to: 1) record low and mid-frequency marine mammal vocalizations; 2) detect, classify, and remotely report vocalizations of interest; and 3) measure high-frequency acoustic backscatter, chlorophyll fluorescence, and oceanographic conditions. Therefore, passive acoustic data can be used to document the distribution of acoustically active marine mammals, including blue whales, and accompanying environmental and oceanographic measurements. Autonomous gliders allow researchers to collect data in areas they are not able to access or in seasons they cannot survey using other means. Onboard detectors for baleen whales currently exist for gliders.

Costs in the implementation schedule (Table 2) include equipment deployed and data analysis.

2.3 Support the development of models to yield robust predictions of blue whale distribution.

A number of species distribution models have been developed, using various methods and data sets. The modelling community has recognized the value of using predictions from a set (“ensemble”) of models rather than those from a single model because multi-model weighted averages often yield more robust predictions (Pérez-Jorge *et al.*, Wintle *et al.* 2003, Johnson & Omland 2004, Araujo & New 2007, Forney *et al.* 2015, Jones & Cheung 2015). For blue whales in the Eastern North Pacific and the California Current, the recent creation of species distribution models makes this a feasible candidate for generating robust predictions using ensemble averaging. Once this has been developed for the Eastern Pacific blue whale stock, the applicability of this type of work should be considered for blue whales that inhabit other areas.

3.0 Estimate Population Size and Monitor Trends in Abundance

Along with determining population structure, assessing population abundance and trends in abundance are among the highest priority actions in this plan. Recovery of blue whale populations can only be assessed if reliable estimates of abundance are available and trends in abundance can be determined. Although abundance estimates are available for some management units, estimates are not available for others, or may be imprecise or limited to geographic areas rather than population units. Ongoing work to assess blue whales in the U.S. and in other parts of the world should also be encouraged and supported.

3.1 Establish collaborative agreements with relevant national governmental bodies and scientific institutions to develop plans for estimating abundance and monitoring trends in abundance.

For meaningful estimates, it will be necessary for U.S. agencies and agency scientists to promote and participate in cooperative surveys with scientists from other countries. A primary goal should be to foster an international collaboration and cooperation in the study and protection of the worldwide population of blue whales.

3.2 Conduct surveys to estimate blue whale abundance and monitor trends in abundance

worldwide.

Systematic surveys should be conducted to assess abundance in areas known, primarily from historic whaling data and large-scale sighting surveys, to have been inhabited regularly by blue whales in the past. The planning of such surveys would be important to capture and understand the occurrence and locations of these whales' annual migrations. Findings from population structure studies identified in action 2.0, above, will be useful in interpreting survey results, because abundance and growth rates may differ between populations. Long-term studies should be anticipated to fully capture population growth rates and possible inter-annual variation in regional occurrence. Potential cost savings might be achieved by combining these studies with other large ship-based research projects and other objectives listed in this Revised Recovery Plan.

3.3 To the extent possible, work with appropriate government agencies in other countries to develop and maintain blue whale photo-identification programs, and educate the public about contributing information about dead, entangled, or ship-struck whales, to continue to support or establish international databases.

In addition to systematic survey data, photo-identification and other data are useful in assessing abundance and monitoring trends. Any existing photo-identification catalogs for blue whales (*e.g.*, Cascadia Research Collective, Baja CA Catalog) should be maintained. Photo-identification data may exist or may be gathered for other locations and habitats and populations. Therefore, efforts should be made to work with appropriate agencies in other countries to solicit the gathering, housing, and maintaining of blue whale photo-identification data and to make these data available to researchers engaged in related studies. Efforts should also be made to educate and to solicit information from the public regarding dead or injured whales. The scientific importance of such catalogs and data bases has been demonstrated with numerous species, and the possibilities for obtaining information about managing the species will increase as more information is obtained.

4.0 Identify, Characterize, Protect, and Monitor Habitat Important to Blue Whale Populations

Identifying important habitat and reducing direct and indirect threats to blue whale habitat is integral to recovery. Important habitat may or may not qualify as critical habitat under the ESA. Information is needed on environmental factors that influence blue whale distribution. In addition, adequate protective measures are needed to reduce or eliminate human-related impacts to blue whale habitat.

4.1 Characterize blue whale habitat.

Areas where blue whales are consistently seen and heard are assumed to be important to their survival. Areas used infrequently or for short periods may also be linked to population fitness. Some areas are known to be important habitat while others may be discovered during survey work discussed in actions 2.0 and 3.0, above. More research is needed to define rigorously and specifically, the environmental features that make an area important to blue whales. Relevant physical, chemical, biological, fishery, and other data should be collected or compiled to characterize features of important habitats and potential sources of human-caused degradation of what are determined to be important

areas for blue whales. Habitat characterization also involves, among other things, descriptions of prey types, densities, and abundances, and of associated oceanographic and hydrographic features. It may also involve characterization of the underwater soundscape, including natural and anthropogenic inputs. Inter-annual variability in habitat features, and in blue whale habitat use, is an important component of habitat characterization. A predictive framework for identifying blue whale habitat is already being used as a management tool in the California Current (Hazen *et al.* 2016). Only with information on the ecological needs of the species will managers be able to provide necessary protections.

Research and analyses to characterize blue whale habitat may also inform evaluation of the effects of climate change, described in 5.5.1 below.

Costs in the implementation schedule (Table 2) include collection of data (*e.g.*, during ship-based surveys, see 3.2), collection of data environmental parameters, and analysis.

4.2 Monitor important habitat features and blue whale use patterns to assess potentially detrimental shifts in habitat features that might reflect disturbance or degradation of habitat.

After baseline data are obtained and analyzed, ongoing studies should be conducted to determine if shifts or changes are occurring in essential habitat components, which might explain or predict changes in blue whale distribution or abundance. Blue whale habitat should be assessed periodically. Shifts in distribution or habitat use should be assessed. Similarly to action 4.1, research and analyses to assess shifts in habitat features may inform evaluation of the effects of climate change, described in 5.5.1 below.

Costs associated with this action (Table 2) are listed as TBD and occurring in FY6 or beyond because baseline data and modeling results from action 4.1 are first needed to determine which aspects of blue whale habitat are most influential in resulting in its use and to inform development of habitat monitoring studies.

4.3 Promote measures to identify and protect important habitat throughout the species' range.

Support efforts to collect and compile data on habitat use patterns, and promote measures to protect and reduce threats to important blue whale habitat. Blue whale ranges include international waters, therefore, collaboration with foreign governments should be encouraged including the use of multi-lateral agreements to protect blue whale habitat in multiple EEZs. International efforts to collect and compile data on habitat use patterns for the blue whale population should be supported. Human activities adversely affecting blue whales should be reduced or mitigated, and the United States should support and endorse such efforts. Validation of those areas where blue whales are thought to occur and protection of those areas that are determined as important areas warranting habitat protection should be supported. Finding means to reduce pollution, protect prey resources, and ensure blue whale habitat integrity on large geographic scales and involving multiple nations may be key to the long-term conservation of blue whale populations.

5.0 Investigate Potential Human-Caused Threats and, Should They Be Determined to Be Limiting Blue Whale Recovery, Take Steps to Minimize Their Occurrence and Severity

5.1 Anthropogenic noise

Blue whales may be disturbed by loud or unfamiliar noises. The scale of their movements may expose them to an array of human activities, including shipping or other commercial vessel traffic, mineral exploration and exploitation (*e.g.* seismic surveys, drilling activities), military operations, and research using acoustic sampling. It is therefore important to understand and reduce the effects of anthropogenic noise on these animals.

- 5.1.1 Conduct studies to determine whether anthropogenic noise is adversely affecting blue whale distribution, behavior, or vital rates.

It is important that the effects of underwater noise on baleen whales become better understood. Studies, such as playback experiments, are needed to assess potential adverse effects of underwater noise (including ship noise) on blue whales, including, but not limited to, disturbance of intraspecific communication, disruption of vital functions, distributional shifts, and stress from chronic or frequent exposure to loud noise. Noise sources studied should include, but are not limited to, industrial and shipping activities, oceanographic experiments, and military-related activities.

Costs in the implementation schedule (Table 2) include support for research such as a behavioral response studies (*e.g.*, (Southall *et al.* 2012) and data analysis.

- 5.1.2 Take steps to minimize anthropogenic noises that are found to be detrimental to blue whales.

If particular sources or types of underwater noise are found to have adverse effects on blue whales, develop and implement appropriate management measures to reduce the threat. In countries, including the U.S., that issue permits and/or otherwise review activities conducted by their citizens, continue existing permitting and consultation processes to review project applications for acoustic impacts to blue whales, and require appropriate monitoring and mitigation measures.

Costs in the implementation schedule (Table 2) are listed as TBD because the step(s) taken to minimize noise found to be detrimental to blue whales (beyond those associated with ongoing permitting and consultation processes) will depend on the findings from 5.1.1 and a determination of whether anthropogenic noise is impeding blue whale recovery.

5.2 Vessel collisions

- 5.2.1 Maintain database of known ship strikes of blue whales.

Consolidate records and maintain a database of known ships strikes to support analyses described below.

- 5.2.2 Review photographic databases for evidence of injuries to blue whales from ship strikes to better characterize these events.

Existing databases, especially those with extensive photographic records of blue whale observations, should be searched for evidence of ship strikes. Although the records may not be sufficient to quantify blue whale injury or mortality rates from this source, such a review might help characterize types of vessels and locations that are prone to ship strike interactions. This, in turn, will help in developing threat-reduction measures.

5.2.3 Conduct studies to identify areas of high risk for blue whale ship strikes.

Areas where high vessel traffic and whale densities overlap should be identified. Areas where ship strikes of blue whales are commonplace should also be identified. This information will help determine the severity of ship strike risk and facilitate the developing and implementing measures to reduce ship strikes.

5.2.4 Conduct studies to determine whether collisions with ships are adversely affecting blue whale abundance and recovery.

Evaluate the magnitude of the risk of ship strikes on blue whale populations to determine the relative impact to recovery. Studies similar to that done by Monnahan *et al.* (2015) could be conducted for blue whale populations in various locations.

5.2.5 If risk of ship strike is determined to be significant, work with mariners, the shipping industry, appropriate state, federal, international agencies to develop and implement measures to reduce the threat of ship strikes, and report strikes, should they occur. Assess the effectiveness of ship strike measures and adjust, as appropriate.

The practicality and effectiveness of options to reduce ship strikes should be assessed. Methods and measures developed for other endangered whales (*e.g.*, North Atlantic right whales) should be considered for their possible application to blue whale populations, as should any current options to reduce ship strikes that have been applied to blue whales in various locations.

Costs in the implementation schedule (Table 2) are listed as TBD because the step(s) taken to address risk of ship strikes will depend on the findings from the other actions in 5.2 and a determination of whether ship strikes are impeding blue whale recovery.

5.3 Marine debris

Harmful marine debris can include plastics and other materials washed or blown from land into the sea, abandoned or lost fishing gear, and solid non-biodegradable materials disposed of by ships at sea. Ingested plastics and other materials may cause a physical blockage in the digestive system, leading to internal injuries or other significant complications, including death. Entanglement or entrapment in “ghost” fishing gear can cause injury or death. Research and management measures are recommended to better understand and address the risk posed by marine debris.

5.3.1 Determine if areas exist in which blue whale habitat and significant deposits of marine debris coincide.

- 5.3.2 Develop and implement means to reduce fishing gear loss, and improve the reporting of lost gear.
- 5.3.3 Take steps to reduce or eliminate sources of marine debris if it is found to be detrimental to blue whales.

Costs in the implementation schedule (Table 2) are listed as TBD because the step(s) taken to reduce or eliminate sources of marine debris will depend on the findings from 5.3.1 and a determination of whether marine debris is impeding blue whales blue whale recovery.

5.4 Fishing gear entanglement

- 5.4.1 Review data on blue whale entanglement in fishing gear, including reports from fishermen and fishery observers.

Evaluate existing data on entanglements in fishing gear, and examine potential overlaps in distribution between fishing operations and blue whales to understand the types of fisheries and fishing gear that might pose the greatest risk to blue whales.

- 5.4.2 Review photographic databases for evidence of injury to blue whales caused by fishing gear entanglement.

Information in these databases may supplement data obtained in action 5.4.1 and provide indications about the nature and extent of fishing gear entanglements.

- 5.4.3 If found to be detrimental to blue whales, develop means to reduce entanglements, and enhance the effectiveness of disentanglement of blue whales in fishing gear. Determine whether measures to reduce entanglements are effective.

Research on possible modifications to fishing gear that may allow an entangled whale to free itself should be fostered or continued if underway. These studies might include assessing the potential use of biodegradable lines, studying ways to reduce the number and length of vertical lines in the water column, designing breakaway lines for heavy gear, and researching acoustic deterrents.

Costs in the implementation schedule (Table 2) are listed as TBD because the step(s) taken to address risk of fishing gear entanglement will depend on the findings from 5.4.1 and 5.4.2, and a determination of whether fishing gear entanglement is impeding blue whale recovery.

5.5 Climate change

- 5.5.1 Conduct research and perform analyses to understand whether climate change is adversely affecting blue whale distribution, abundance, or fecundity.

Improved knowledge of the effects of climate change on blue whale feeding ecology and habitat use would be informative for evaluating or predicting shifts in prey abundance or distribution caused by climate change. Similarly, information on the environmental drivers influencing distribution of blue

whales and their prey, and the long-term changes in these factors, will aid in understanding the overall impact, if any, to various blue whale populations resulting from climate change. Although the natural absorption of carbon dioxide (CO₂) by the world's oceans helps mitigate the climatic effects of anthropogenic emissions of CO₂, it is believed that the resulting decrease in pH will have negative consequences. While the full ecological consequences of these changes are not known, organisms such as blue whales may suffer adverse effects, either directly as reproductive or physiological effects or indirectly through negative impacts on their food resources.

Research and analyses on blue whale abundance, distribution, and habitat characteristics conducted as part of other identified recovery actions (2.0, 3.0, and 4.0) may be informative for understanding effects of climate change. Any efforts to combine data collection or analysis would reduce costs.

- 5.5.2 Take steps to minimize climate change impacts if found to be detrimental to blue whales.

Strategies developed through international efforts to minimize the effects of climate change on blue whales and the ecosystems they inhabit should be pursued. However, costs in the implementation schedule (Table 2) are listed as TBD because the step(s) taken to address climate change impacts will depend on the findings from 5.5.1 and a determination of whether climate change is impeding blue whale recovery.

6.0 Maximize Efforts to Acquire Scientific Information from Dead, Stranded, and Entangled or Entrapped Blue Whales

Assessments of the causes and frequency of mortality (either natural or human-caused) is important to understanding population dynamics and threats that may impede the recovery of blue whale populations. Therefore, cause of death determinations are an important part of recovery and threat reduction actions. Discovery of a carcass under circumstances allowing response and, if dead, necropsy in a timely and rigorous manner is a relatively rare event. Accordingly, efforts to detect and investigate blue whale deaths, strandings, and other responses should be as efficient as possible.

6.1 Make every effort to necropsy stranded blue whales to determine cause of death.

The detection and reporting of blue whales, whether stranded or floating at sea, need to be encouraged. Marine Mammal Stranding Network coordinators should continue working with representatives of local, state, and federal agencies, private organizations, academic institutions, and regional and local stranding networks, to ensure use of accepted reporting procedures, implement standardized necropsy protocol, and facilitate information exchange. In areas where protocols do not exist, they should be developed. As part of this effort, responsibilities of all relevant agencies, organizations, and individuals should be clearly defined.

Blue whales may die at sea, but may not be detected or reported. Mariners, Navy, and Coast Guard personnel, commercial and recreational boaters, and fishermen might observe carcasses at sea, but may not recognize the importance of their observation. Mariners should be educated about the importance of reporting carcasses so worthwhile

information might be collected from them.

Each blue whale carcass represents an opportunity for scientific investigation of the cause of death, as well as addressing other questions related to the biology of the species. Delays in attempts to secure or examine a carcass can result in the loss of valuable data, or even of the carcass itself. Stranding Network coordinators should work with appropriate agencies, organizations, and individuals to ensure that, when a blue whale carcass is reported and secured: (i) a necropsy is performed as rapidly and as thoroughly as possible by qualified individuals to gather information regarding the cause of death; (ii) samples are taken and properly preserved for studies of genetics, toxicology, and pathology; and (iii) funding is available to notify and transport appropriate experts to planned necropsy sites rapidly and to distribute tissue samples to appropriate locations for analysis or storage. In addition, coordinators should work with stranding networks and the scientific community, to develop and maintain lists of tissue samples requested by qualified individuals and agencies, and ensure that these samples are routinely collected from each carcass and stored in designated locations or distributed to appropriate researchers.

Strandings of blue whales are relatively rare. Costs in the implementation schedule (Table 2) include necropsy and stranding response for blue whales in the U.S. EEZ. Costs for necropsy and response throughout the blue whales' range may be significantly greater.

6.2 Review, analyze, and summarize data on dead blue whales and causes of death.

Current and complete data from stranding events is essential to developing protective measures. These data are likely to be useful in determining the magnitude of potential threats such as vessel strikes, entanglements, disease, and impacts of climate change.

6.3 Establish reliable sources of funding for necropsy (including sites to conduct the necropsy), tissue collection, and analysis efforts.

Collection of information from blue whale carcasses is essential to recovery efforts. Therefore, identifying and committing predictable sources of funding for completing necropsy, tissue collection, and analysis efforts is also critical.

V. IMPLEMENTATION SCHEDULE

The implementation schedule that follows is used to estimate costs to direct and monitor implementation and completion of recovery actions set forth in this Revised Recovery Plan. It is a guide for meeting recovery goals outlined in this Revised Recovery Plan. The implementation schedule indicates the action numbers, action descriptions, action priorities, duration of the action, the parties potentially involved in the actions, and estimated costs. While parties with authority, responsibility, or expressed interest to implement a specific recovery action are identified in the implementation schedule, this does not require any identified party to implement the action(s) or to secure funding for implementing the action(s).

Priorities in column 3 of the implementation schedule are assigned as follows⁶:

Priority 1 – An action that must be taken to prevent extinction or to identify those actions necessary to prevent extinction.

Priority 2 – An action that must be taken to prevent a significant decline in population numbers or habitat quality, or to prevent other significant negative trends short of extinction.

Priority 3 – All other actions necessary to provide for full recovery of the species.

This implementation schedule accords priorities to individual actions to specify their importance in the recovery effort. It should be noted that even the highest-priority actions within a plan are not given a Priority 1 ranking unless they are actions necessary to prevent extinction or to identify those actions necessary to prevent extinction.

As discussed in Section I.H above, while there are no known current threats to the species, there are numerous potential threats. These include potential threats that are known to be affecting one or more subspecies or population, but more research is needed to understand the extent to which the stressor occurs or affects the globally listed entity. As such, many of the recovery actions addressing potential threats to blue whale are listed as Priority 1 if research and/or other monitoring efforts identify that the potential threats are indeed operative threats to blue whale recovery. We cannot know the outcome of the research or monitoring efforts before they occur. If research does not support identification of these potential threats as operative threats, then the recovery actions addressing them should be considered lower priority. In addition, the research actions will also contribute to monitoring of stressors to confirm that they remain at insignificant levels that do not limit recovery.

For each of the six major recovery actions, Table 2 provides cost sub-totals in ***bold italics***. Recovery actions are linked, in that the ability to complete one may depend on outcomes from a previous study. However, the costs of recovery actions are estimated individually. Additionally,

⁶ NMFS proposed revisions to the Recovery Plan Preparation and Implementation Priority guidelines (82 FR 24944, May 31, 2017), which included changes to the scheme for prioritizing recovery actions contained within a recovery plan. The revised guidelines, once finalized, will be used in the final Revised Recovery Plan.

many of the recovery actions are similar to those identified in other NMFS large whale recovery plans (see <https://www.fisheries.noaa.gov/resources/documents>) and might be expected to provide information on or benefit other large whales in the area. However, the costs of recovery actions are estimated as if efforts are specific to blue whales. Therefore, in some cases the identified costs of actions that benefit multiple species may be overestimates.

Costs are estimated in accordance with the number of years necessary to complete the action once implementation has begun. Some costs are listed over discrete periods of time (*e.g.*, 5 years), while others are listed as ongoing. “Ongoing” means that the action should be conducted regularly or opportunistically for the foreseeable future, until the knowledge gap has been filled, or until the threat has been reduced to the point where it is no longer a concern.

It is not possible to predict the time and cost to recovery with the current information because of the uncertainty of potential threats, as explained previously. Thus, an estimate of the time required and the cost to carry out those actions needed to achieve the plan’s goal and to achieve intermediate steps (beyond 5 years) is not practicable. Conducting research necessary to support conclusions regarding the impact of the potential threats to blue whale populations, and developing, implementing, and evaluating the effectiveness of management measures to reduce threats or potential threats may take decades. The minimum data needed to satisfy criterion 1 for downlisting or delisting are population structure studies and abundance surveys, which will also take decades, given the species’ global distribution. In the future, as more information is obtained, it may be possible to develop estimates for the full time to recovery and its expense.

Table 2. Implementation Schedule by Fiscal Year

Action Number	Action Description	Priority	Action Duration (years)	Agencies/ Organizations Potentially Involved	Cost Estimates by FY (thousands of dollars)						
					FY1	FY2	FY3	FY4	FY5	FY6+ ⁷	Total ⁸
1.0	<i>Coordinate State, Federal, and International Measures to Maintain International Regulation of Whaling for Blue Whales</i>	1	Ongoing	NMFS, IWC, International Partners, Department of State	*	*	*	*	*	*	*
TOTAL 1					*	*	*	*	*	*	*
2.0	<i>Determine Blue Whale Taxonomy, Population Structure, Occurrence, Distribution, and Range.</i>										
2.1	<i>Conduct studies to investigate population discreteness and population structure of blue whales.</i>	2	Ongoing (15+)	NMFS, IWC, International Partners	250	250	250	250	250	250+	1,500+
2.2	<i>Conduct studies to assess blue whale occurrence, daily and seasonal movements, and inter-area exchange.</i>	2	Ongoing (15+)	NMFS, IWC, Navy, International Partners, Non-Governmental Partners	150	150	150	150	150	150+	900+
2.3	<i>Support the development of models to yield robust predictions of blue whale distribution.</i>	2	4-5	NMFS, IWC, Non-Governmental Partners	*	*	*	*	*		*
TOTAL 2					400	400	400	400	400	400+	2,400+

⁷ For actions with a discrete duration exceeding five fiscal years, the FY6+ column includes total costs anticipated after FY1-5. For ongoing actions (see definition of “ongoing” above), costs beyond FY5 are listed as a single year’s cost with the symbol “+” to indicate additional, ongoing annual costs.

⁸ For actions with a discrete duration, the total is the sum of anticipated costs across the action’s duration. For ongoing actions, the total is the sum of the first six fiscal years (FY1-6) with the symbol “+” to indicate additional, ongoing annual costs.

Action Number	Action Description	Priority	Action Duration (years)	Agencies/ Organizations Potentially Involved	Cost Estimates by FY (thousands of dollars)							
					FY1	FY2	FY3	FY4	FY5	FY6+ ⁷	Total ⁸	
3.0	<i>Estimate Population Size and Monitor Trends in Abundance</i>											
3.1	<i>Establish collaborative agreements with relevant national governmental bodies and scientific institutions to develop plans for estimating abundance and monitoring trends in abundance.</i>	2	3-5	NMFS, IWC, International Partners, Department of State	*	*	*	*	*	*	*	*
3.2	<i>Conduct surveys to estimate blue whale abundance and monitor trends in abundance worldwide.</i>	2	Ongoing (15+)	NMFS, International Partners, Non-Governmental Partners	450	450	450	450	450	450+	2,700+	
3.3	<i>To the extent possible, work with appropriate government agencies in other countries to develop and maintain blue whale photo-identification programs, and educate the public about contributing information about dead, entangled, or ship-struck whales, to continue to support or establish international databases.</i>	2-3	Ongoing	NMFS, International Partners, Non-Governmental Partners	80	80	80	80	80	80+	480+	
TOTAL 3					530	530	530	530	530	530+	3,180+	
4.0	<i>Identify, Characterize, Protect, and Monitor Habitat Important to Blue Whale Populations</i>											
4.1	<i>Characterize blue whale habitat.</i>	2	5	NMFS, International Partners, Non-Governmental Partners	400	400	400	400	400		2,000	
4.2	<i>Monitor important habitat features and blue whale use patterns to assess potentially detrimental shifts in habitat features that might reflect disturbance or degradation of habitat.</i>	2	Ongoing	NMFS, International Partners, Non-Governmental Partners						TBD	TBD	

Action Number	Action Description	Priority	Action Duration (years)	Agencies/ Organizations Potentially Involved	Cost Estimates by FY (thousands of dollars)						
					FY1	FY2	FY3	FY4	FY5	FY6+ ⁷	Total ⁸
4.3	<i>Promote measures to identify and protect important habitat throughout the species' range.</i>	2	Ongoing	NMFS, International Partners	*	*	*	*	*	*	*
TOTAL 4					400	400	400	400	400	TBD	2,000+
5.0	<i>Investigate Potential Human-Caused Threats and, Should They Be Determined to Be Limiting Blue Whale Recovery, Take Steps to Minimize Their Occurrence and Severity</i>										
5.1	<i>Anthropogenic noise</i>										
5.1.1	Conduct studies to determine whether anthropogenic noise is adversely affecting blue whale distribution, behavior, or vital rates.	2	5-10 years	NMFS, U.S. Navy, Bureau of Ocean Energy and Management (BOEM); International Partners, Non-Governmental Partners	300	300	300	*	*		900
5.1.2	Take steps to minimize anthropogenic noises that are found to be detrimental to blue whales.	1	Ongoing	NMFS, U.S. Navy, BOEM						TBD	TBD
5.2	<i>Vessel collisions</i>										
5.2.1	Maintain database of known ship strikes of blue whales.	2	Ongoing	NMFS, IWC	*	*	*	*	*	*	*
5.2.2	Review photographic databases for evidence of injuries to blue whales from ship strikes to better characterize these events.	2	Ongoing	NMFS	*	*	*	*	*	*	*
5.2.3	Conduct studies to identify areas of high risk for blue whale ship strikes.	2	2	NMFS, USCG	30	30					60
5.2.4	Conduct studies to determine whether collisions with ships are adversely affecting blue whale abundance and recovery.	2	Ongoing	NMFS	*	*	*	*	*	*	*

Action Number	Action Description	Priority	Action Duration (years)	Agencies/ Organizations Potentially Involved	Cost Estimates by FY (thousands of dollars)						
					FY1	FY2	FY3	FY4	FY5	FY6+ ⁷	Total ⁸
5.2.5	If risk of ship strike is determined to be significant, work with mariners, the shipping industry, appropriate state, federal, international agencies to develop and implement measures to reduce the threat of ship strikes, and report strikes, should they occur. Assess the effectiveness of ship strike measures and adjust, as appropriate.	1	Ongoing	NMFS, USCG, International Maritime Organization, international partners						TBD	TBD
5.3	<i>Marine debris</i>										
5.3.1	Determine if areas exist in which blue whale habitat and significant deposits of marine debris coincide.	2	Ongoing	NMFS, IWC, International Partners	100	100	100	100	100	100+	600+
5.3.2	Develop and implement means to reduce fishing gear loss, and improve the reporting of lost gear.	2	Ongoing	NMFS, International Partners	150	150	150	150	150	150+	900+
5.3.3	Take steps to reduce or eliminate sources of marine debris if it is found to be detrimental to blue whales.	1	TBD	NMFS, International Partners						TBD	TBD
5.4	<i>Fishing gear entanglement</i>										
5.4.1	Review data on blue whale entanglement in fishing gear, including reports from fishermen and fishery observers.	2	Ongoing	NMFS	*	*	*	*	*	*	*
5.4.2	Review photographic databases for evidence of injury to blue whales caused by fishing gear entanglement.	2	Ongoing	NMFS	*	*	*	*	*	*	*
5.4.3	If found to be detrimental to blue whales, develop means to reduce entanglements, and enhance the effectiveness of disentanglement of blue whales in fishing gear. Determine whether measures to reduce entanglements are effective.	1	TBD	NMFS, International Partners, Non-Governmental Partners	TBD	TBD	TBD	TBD	TBD	TBD	TBD

Action Number	Action Description	Priority	Action Duration (years)	Agencies/ Organizations Potentially Involved	Cost Estimates by FY (thousands of dollars)							
					FY1	FY2	FY3	FY4	FY5	FY6+ ⁷	Total ⁸	
5.5	<i>Climate change</i>											
5.5.1	Conduct research and perform analyses to understand whether climate change is adversely affecting blue whale distribution, abundance, or fecundity.	2	Ongoing	NMFS, IWC, NOAA's National Ocean Service (NOS), NOAA Office of Oceanic and Atmospheric Research	150	150	150	150	150	150+	900+	
5.5.2	Take steps to minimize climate change impacts if found to be detrimental to blue whales.	1	TBD	TBD	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
TOTAL 5					730 and TBD	730 and TBD	700 and TBD	400 and TBD	400 and TBD	400+ and TBD	2,960+ and TBD	
6.0	Maximize Efforts to Acquire Scientific Information from Dead, Stranded, and Entangled or Entrapped Blue Whales											
6.1	<i>Make every effort to necropsy stranded blue whales to determine cause of death.</i>	3	Ongoing	NMFS, USCG, States, Stranding networks, other partners	30	30	30	30	30	30+	180+	
6.2	<i>Review, analyze, and summarize data on dead blue whales and causes of death.</i>	3	Ongoing	NMFS, States, NOS	*	*	*	*	*	*	*	
6.3	<i>Establish reliable sources of funding for necropsy (including sites to conduct the necropsy), tissue collection, and analysis efforts.</i>	3	Ongoing	NMFS	*	*	*	*	*	*	*	
TOTAL 6					30	30	30	30	30	30+	180+	

*No cost associated, NMFS staff time

This Page Intentionally Left Blank

VI. LITERATURE CITED

- Abramson, J.Z. and J. Gibbons. 2010. New records of blue whales *Balaenoptera musculus* (Linnaeus, 1758) in winter season in the inlet waters of Chile continental-Chile. *Anales Instituto Patagonia (Chile)* 38:107-109.
- Acevedo-Gutierrez, A., D.A. Croll and B.R. Tershy. 2002. High feeding costs limit dive time in the largest whales. *Journal of Experimental Biology* 205:1747-1753.
- Aguilar, A. 1987. Using organochlorine pollutants to discriminate marine mammal populations - a review and critique of the methods. *Marine Mammal Science* 3:242-262.
- Aguilar, A., A. Borrell and P.J.H. Reijnders. 2002. Geographical and temporal variation in levels of organochlorine contaminants in marine mammals. *Marine Environmental Research* 53:425-452.
- Allen, K.R. 1970. A note on baleen whale stocks of the northwest Atlantic. Report of the International Whaling Commission 20:112-113.
- Alling, A., E.M. Dorsey and J.C.D. Gordon. 1991. Blue whales (*Balaenoptera musculus*) off the northeast coast of Sri Lanka: Distribution, feeding and individual identification. Pages 247-258 in S. Leatherwood and G.P. Donovan eds. *Cetaceans and Cetacean Research in the Indian Ocean Sanctuary*. United Nations Environment Programme.
- Amos, B. and R. Hoelzel. 1992. Applications of molecular genetic techniques to the conservation of small populations. *Biological Conservation* 61:133-144.
- Anderson, R.C., T.A. Branch, A. Alagiyawadu, R. Baldwin and F. Marsac. 2012. Seasonal distribution, movements and taxonomic status of blue whales (*Balaenoptera musculus*) in the northern Indian Ocean. *Journal of Cetacean Research And Management* 12:203-218.
- Andrew, R.K., B.M. Howe, J.A. Mercer and M.A. Dzieciuch. 2002. Ocean ambient sound: Comparing the 1960s with the 1990s for a receiver off the California coast. *Acoustics Research Letters Online-ARLO* 3:65-70.
- Anisimov, O.A., D.G. Vaughan, T.V. Callaghan, C. Furgal, H. Marchant, T.D. Prowse, H. Vilhjálmsson and J.E. Walsh. 2007. Polar regions (Arctic and Antarctic). Pages 653-685 *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK.
- ANSI (American National Standards Institute). 1995. American National Standard: Bioacoustical Terminology (ANSI S3.20-1995). Acoustical Society of America, Woodbury, NY.
- Araujo, M.B. and M. New. 2007. Ensemble forecasting of species distributions. *Trends in Ecology and Evolution* 22:42-47.
- Aroyan, J.L., M.A. McDonald, S.C. Webb, J.A. Hildebrand and C. Clark. 2000. Acoustic models of sound production and propagation. Pages 409-469 in W.W.L. Au, A.N. Popper and R.R. Fay eds. *Hearing by Whales and Dolphins*. Springer-Verlag, New York.
- Atkinson, A., V. Siegel, E. Pakhomov and P. Rothery. 2004. Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature* 432:100-103.
- Atkinson, A., V. Siegel, E.A. Pakhomov, M.J. Jessopp and V. Loeb. 2009. A re-appraisal of the total biomass and annual production of Antarctic krill. *Deep Sea Research I* 56:727-740.
- Attard, C.R.M., L.B. Beheregaray, C. Jenner, P. Gill, M. Jenner, M. Morrice, J. Bannister, R. Leduc and L. Moller. 2010. Genetic diversity and structure of blue whales (*Balaenoptera musculus*) in Australian feeding aggregations. *Conservation Genetics* 11:2437-2441.
- Attard, C.R.M., L.B. Beheregaray, K.C.S. Jenner, P.C. Gill, M.-N. Jenner, M.G. Morrice, K.M.

- Robertson and L.M. Moller. 2012. Hybridization of Southern Hemisphere blue whale subspecies and a sympatric area off Antarctica: Impacts of whaling or climate change? *Molecular Ecology* 21:5715-5727.
- Attard, C.R.M., L.u.B. Beheregaray and L.M. Moller. 2016. Towards population-level conservation in the critically endangered Antarctic blue whale: the number and distribution of their populations. *Sci Rep* 6.
- Au, W.W.L. 1993. *The Sonar of Dolphins*. Springer-Verlag, New York, New York.
- Au, W.W.L. and M.C. Hastings. 2008. *Principles of Marine Bioacoustics*. Springer-Verlag New York.
- Au, W.W.L., A.A. Pack, M.O. Lammers, L.M. Herman, M.H. Deakos and K. Andrews. 2006. Acoustic properties of humpback whale songs. *Journal of the Acoustical Society of America* 120:1103.
- Bailey, H., B.R. Mate, D.M. Palacios, L. Irvine, S.J. Bograd and D.P. Costa. 2010. Behavioural estimation of blue whale movements in the northeast Pacific from state-space model analysis of satellite tracks. *Endangered Species Research* 10:93-106.
- Baines, M.E. and M. Reichelt. 2014. Upwellings, canyons and whales: An important winter habitat for balaenopterid whales off Mauritania, northwest Africa. *Journal of Cetacean Research And Management* 14:57-67.
- Balcazar, N.E., J.S. Tripovich, H. Klinck, S.L. Nieukirk, D.K. Mellinger, R.P. Dziak and T.L. Rogers. 2015. Calls reveal population structure of blue whales across the southeast Indian Ocean and the southwest Pacific Ocean. *Journal of Mammalogy* 96:1184-1193.
- Bargu, S., C.L. Powell, S.L. Coale, M. Busman, G.J. Doucette and M.W. Silver. 2002. Krill: a potential vector for domoic acid in marine food webs. *Marine Ecology Progress Series* 237:209-216.
- Barlow, J. 1997. Preliminary estimates of cetacean abundance off California, Oregon, and Washington based on a 1996 ship survey and comparisons of passing and closing modes. National Marine Fisheries Service, Southwest Fisheries Science Center Administrative Report. 25 pp.
- Barlow, J. 2006. Cetacean abundance in Hawaiian waters estimated from a summer/fall survey in 2002. *Marine Mammal Science* 22:446-464.
- Barlow, J. 2010. Cetacean abundance in the California Current estimated from a 2008 ship-based line-transect survey. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-456. pp.
- Barlow, J. and K.A. Forney. 2007. Abundance and population density of cetaceans in the California Current ecosystem. *Fishery Bulletin* 105:509-526.
- Beamish, P.C. 1979. Behaviour and significance of entrapped baleen whales. Pages 291-309 in H.E. Winn and B.L. Olla eds. *Behavior of Marine Animals: Current Perspectives in Research Vol. 3: Cetaceans*. Plenum Press, N.Y.
- Beauchamp, J., H. Bouchard, P. de Margerie, N. Otis and J.-Y. Savaria. 2009. Recovery Strategy for the blue whale (*Balaenoptera musculus*), Northwest Atlantic population, in Canada [FINAL]. Fisheries and Oceans Canada. *Species at Risk Act Recovery Strategy Series* 62 pp.
- Bejder, L., A. Samuels, H. Whitehead and N. Gales. 2006. Interpreting short-term behavioural responses to disturbance within a longitudinal perspective. *Animal Behaviour* 72:1149-1158.
- Benson, S.R., D.A. Croll, B.B. Marinovic, F.P. Chavez and J.T. Harvey. 2002. Changes in the cetacean assemblage of a coastal upwelling ecosystem during El Niño 1997-98 and La Niña 1999. *Progress in Oceanography* 54:279-291.

- Berchok, C.L., D.L. Bradley and T.B. Gabrielson. 2006. St. Lawrence blue whale vocalizations revisited: Characterization of calls detected from 1998 to 2001. *Journal of the Acoustical Society of America* 120:2340-2354.
- Berenboim, B.I., A.V. Dolgov, V.A. Korzhev and N.A. Yaragina. 2000. The Impact of Cod on the Dynamics of Barents Sea Shrimp (*Pandalus borealis*) as Determined by Multispecies Models. *J. Northw. Atl. Fish. Sci.* 27:69-75.
- Berman-Kowalewski, M., F.M.D. Gulland, S. Wilkin, J. Calambokidis, B. Mate, J. Cordaro, D. Rotstein, Judy St. Leger, Paul Collins, Krista Fahy and S. Dover. 2010. Association Between Blue Whale (*Balaenoptera musculus*) Mortality and Ship Strikes Along the California Coast. *Aquatic Mammals* 36:59-66.
- Berube, M. and A. Aguilar. 1998. A new hybrid between a blue whale, *Balaenoptera musculus*, and a fin whale, *B. physalus*: Frequency and implications of hybridization. *Marine Mammal Science* 14:82-98.
- Berzin, A.A. and A.A. Rovnin. 1966. Distribution, and migration of whales in the northeastern part of the Pacific Ocean, Bering, and Chukchi seas. *Izvestiya Tinro* 58:179-207.
- Best, P.B. 1993. Increase rates in severely depleted stocks of baleen whales. *ICES Journal of Marine Science* 50:169-186.
- Best, P.B., R.A. Rademeyer, C. Burton, D. Ljungblad, K. Sekiguchi, H. Shimada, D. Thiele, D. Reeb and D.S. Butterworth. 2003. The abundance of blue whales on the Madagascar Plateau, December 1996. *Journal of Cetacean Research And Management* 5:253-260.
- Bettridge, S., C.S. Baker, J. Barlow, P.J. Clapham, M. Ford, D. Gouveia, D.K. Mattila, R.M. Pace, III, P.E. Rosel, G.K. Silber and P.R. Wade. 2015. Status Review of the Humpback Whale (*Megaptera novaeangliae*) under the Endangered Species Act. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-540. 156 + Appendices pp.
- Bindoff, N.L., J. Willebrand, V. Artale, A. Cazenave, J. Gregory, S. Gulev, K. Hanawa, C.L. Quéré, S. Levitus, Y. Nojiri, C.K. Shum, L.D. Talley and A. Unnikrishnan. 2007. Observations: Oceanic Climate Change and Sea Level. Pages 385-432 in S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller eds. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Blackwell, S.B., C.S. Nations, T.L. McDonald, A.M. Thode, D. Mathias, K.H. Kim, C.R. Greene, Jr. and A.M. Macrander. 2015. Effects of airgun sounds on bowhead whale calling rates: evidence for two behavioral thresholds. *PLoS one* 10:e0125720.
- Borrell, A. and A. Aguilar. 1999. A review of organochlorine and metal pollutants in marine mammals from Central and South America. *Journal of Cetacean Research And Management Special Issue* 1:195-207.
- Boveng, P.L., J.L. Bengtson, T.W. Buckley, M.F. Cameron, S.P. Dahle, B.P. Kelly, B.A. Megrey, J.E. Overland and N.J. Williamson. 2009. Status Review of the Spotted Seal (*Phoca largha*) U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-AFSC-200. 153 pp.
- Boveng, P.L., J.L. Bengtson, M.F. Cameron, S.P. Dahle, E.A. Logerwell, J.M. London, J.E. Overland, J.T. Sterling, D.E. Stevenson, B.L. Taylor and H.L. Ziel. 2013. Status Review of the Ribbon Seal (*Histiophoca fasciata*) U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-AFSC-255. 174 pp.
- Bradford, A.L., K.A. Forney, E.M. Oleson and J. Barlow. 2017. Abundance estimates of

- cetaceans from a line-transect survey within the U.S. Hawaiian Islands Exclusive Economic Zone. *Fishery Bulletin* 115:129-142.
- Braithwaite, J.E., J.J. Meeuwig, T.B. Letessier, K.C.S. Jenner and A.S. Brierley. 2015. From sea ice to blubber: Linking whale condition to krill abundance using historical whaling records. *Polar Biology* 38:1195-1202.
- Branch, T.A. 2007. Abundance of Antarctic blue whales south of 60° from three complete circumpolar sets of surveys. *Journal of Cetacean Research And Management* 9:253-262.
- Branch, T.A. 2008a. Biological parameters for pygmy blue whales. International Whaling Commission Scientific Committee. 13 pp.
- Branch, T.A. 2008b. Biologically plausible rates of increase for Antarctic blue whales. International Whaling Commission Scientific Committee. 7 pp.
- Branch, T.A. 2008c. Current status of Antarctic blue whales based on Bayesian modelling. International Whaling Commission Scientific Committee. 10 pp.
- Branch, T.A., E.M.N. Abubaker, S. Mkango and D.S. Butterworth. 2007a. Separating southern blue whale subspecies based on length frequencies of sexually mature females. *Marine Mammal Science* 23:803-833.
- Branch, T.A., C. Allison, Y.A. Mikhalev, D. Tormosov and R.L. Brownell, Jr. 2008. Historical catch series for Antarctic and pygmy blue whales. International Whaling Commission Scientific Committee. 11 pp.
- Branch, T.A., K. Matsuoka and T. Miyashita. 2004. Evidence for increases in Antarctic blue whales based on Bayesian modelling. *Marine Mammal Science* 20:726-754.
- Branch, T.A. and Y.A. Mikhalev. 2008. Regional differences in length at sexual maturity for female blue whales based on recovered Soviet whaling data. *Marine Mammal Science* 24:690-703.
- Branch, T.A., K.M. Stafford, D.M. Palacios, C. Allison, J.L. Bannister, C.L.K. Burton, E. Cabrera, C.A. Carlson, B.G. Vernazzani, P.C. Gill, R. Hucke-Gaete, K.C.S. Jenner and M.-N.M. Jenner. 2007b. Past and present distribution, densities and movements of blue whales *Balaenoptera musculus* in the Southern Hemisphere and northern Indian Ocean. *Mammal Review* 37:116-175.
- Branch, T.A., A.N. Zerbini and K. Findlay. 2007c. Abundance of blue whales off Chile from the 1997/98 SOWER survey. International Whaling Commission Scientific Committee. 9 pp.
- Brodeur, R.D. and D.M. Ware. 1992. Long-term variability in zooplankton biomass in the subArctic Pacific Ocean. *Fisheries Oceanography* 1:32-38.
- Brownell, R.L., Jr., E. Cabrera and B. Galletti Vernazzani. 2014. Dead Blue Whale in Puerto Montt, Chile: Another Case of Ship Collision Mortality. International Whaling Commission Scientific Committee. 9 pp.
- Brownell, R.L., Jr., C.A. Carlson, B. Galletti Vernazzani and E. Cabrera. 2007. Skin lesions on blue whales off southern Chile: Possible conservation implications? International Whaling Commission Scientific Committee. 6 pp.
- Brownell, R.L., Jr., B. Galletti Vernazzani, A. deVos, P.A. Olson, K. Findlay, J. Bannister and A.R. Lang. 2015. Assessment of pygmy type blue whales in the Southern Hemisphere. International Whaling Commission Scientific Committee. 25 pp.
- Brownell, R.L., Jr. and K. Ralls. 1986. Potential for sperm competition in baleen whales. *Special Issue* 8:97-112.
- Buchan, S.J. and R.A. Quiñones. 2016. First insights into the oceanographic characteristics of a blue whale feeding ground in northern Patagonia, Chile. *Marine Ecology Progress Series*

554:183-199.

- Buchan, S.J., K.M. Stafford and R. Hucke-Gaete. 2015. Seasonal occurrence of southeast Pacific blue whale songs in southern Chile and the eastern tropical Pacific. *Marine Mammal Science* 31:440-458.
- Buck, J.D., L.L. Shepard and S. Spotte. 1987. *Clostridium perfringens* as the cause of death of a captive Atlantic bottlenosed dolphin (*Tursiops truncatus*). *Journal of Wildlife Diseases* 23:488-491.
- Bundy, A. 2005. Structure and functioning of the eastern Scotian Shelf ecosystem before and after the collapse of groundfish stocks in the early 1990s. *Canadian Journal of Fisheries and Aquatic Sciences* 62:1453-1473.
- Burmeister, H. 1871. Del Año 1871. *Boletín Del Museo Público de Buenos Aires*, 1871:11-20.
- Burtenshaw, J.C., E.M. Oleson, J.A. Hildebrand, M.A. McDonald, R.K. Andrew, B.M. Howe and J.A. Mercer. 2004. Acoustic and satellite remote sensing of blue whale seasonality and habitat in the Northeast Pacific. *Deep Sea Research Part II: Topical Studies in Oceanography* 51:967-986.
- Cabrera, E., C.A. Carlson and B. Galletti Vernazzani. 2005. Presence of blue whale (*Balaenoptera musculus*) in the northwestern coast of Chiloe Island, southern Chile. *Latin American Journal of Aquatic Mammals* 4:73-74.
- Calambokidis, J. and J. Barlow. 2004. Abundance of blue and humpback whales in the eastern North Pacific estimated by capture-recapture and line-transect methods. *Marine Mammal Science* 20:63-85.
- Calambokidis, J. and J. Barlow. 2013. Updated abundance estimates of blue and humpback whales off the US west coast incorporating photo-identifications from 2010 and 2011. Final Report for contract AB133F-10-RP-0106. Document PSRG-2013-13 presented to the Pacific Scientific Review Group, April 2013. 7 pp.
- Calambokidis, J., J. Barlow, J.K.B. Ford, T.E. Chandler and A.B. Douglas. 2009a. Insights into the population structure of blue whales in the eastern North Pacific from recent sightings and photographic identification. *Marine Mammal Science* 25:816-832.
- Calambokidis, J., E. Falcone, A. A. Douglas, L. Schlender and J. Huggins. 2009b. Photographic identification of humpback and blue whales off the U.S. West Coast: results and updated abundance estimates from 2008 field season. Cascadia Research. Final Report for Contract AB133F08SE2786 from Southwest Fisheries Science Center. 18 pp.
- Calambokidis, J., E.A. Falcone, T.J. Quinn, A.M. Burdin, P.J. Clapham, J.K.B. Ford, C.M. Gabriele, R. LeDuc, D. Mattila, L. Rojas-Bracho, J.M. Straley, B.L. Taylor, J. Urbán R., D. Weller, B.H. Witteveen, M. Yamaguchi, A. Bendlin, D. Camacho, K. Flynn, A. Havron, J. Huggins and N. Maloney. 2008a. SPLASH: Structure of Populations, Levels of Abundance and Status of Humpback Whales in the North Pacific. Cascadia Research. Final report for Contract AB133F-03-RP-00078 for U.S. Department of Commerce, Western Administrative Center, Seattle, WA. . 57 pp.
- Calambokidis, J., E. Oleson, J. Goldbogen and M. McKenna. 2010. Use of suction cup attached tags on whales to examine underwater behavior, feeding, vocalizations, and interactions with ships. 2010 Ocean Sciences Meeting. American Geophysical Union, Portland, OR.
- Calambokidis, J., G.S. Schorr, G.H. Steiger, J. Francis, M. Bakhtiari, G. Marshal, E.M. Oleson, D. Gendron and K. Robertson. 2008b. Insights into the underwater diving, feeding, and calling behavior of blue whales from a suction-cup-attached video-imaging tag (CRITTERCAM). *Marine Technology Society Journal* 41:19-29.

- Calambokidis, J., G.H. Steiger, J.C. Cubbage, K.C. Balcomb, C. Ewald, S. Kruse, R. Wells and R. Sears. 1990. Sightings and movements of blue whales off central California 1986-88 from photo-identification of individuals. Report of the International Whaling Commission Special Issue 12:343-348.
- Calambokidis, J., G.H. Steiger, C. Curtice, J. Harrison, M.C. Ferguson, E. Becker, M. DeAngelis and S.M. Van Parijs. 2015. Biologically important areas for selected cetaceans within U.S. waters – West Coast region. *Aquatic Mammals* 41:39-53.
- Carretta, J.V., J.E. Moore and K.A. Forney. 2017a. Regression tree and ratio estimates of marine mammal, sea turtle, and seabird bycatch in the California drift gillnet fishery: 1990-2015. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-568. 83 pp.
- Carretta, J.V., M.M. Muto, J. Greenman, K. Wilkinson, D. Lawson, J. Viezbicke and J. Jannot. 2017b. Sources of human-related injury and mortality for U.S. Pacific West Coast marine mammal stock assessments, 2011-2015. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-579. 126 pp.
- Carretta, J.V., E.M. Oleson, D.W. Weller, A.R. Lang, K.A. Forney, J. Baker, M.M. Muto, B. Hanson, A.J. Orr, H. Huber, M.S. Lowry, J. Barlow, J.E. Moore, D. Lynch, L. Carwsell and R.L. Brownell, Jr. 2015. U.S. Pacific Marine Mammal Stock Assessments: 2014. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-549. 414 pp.
- Carretta, J.V., S.M. Wilkin, M.M. Muto and K. Wilkinson. 2013. Sources of human-related injury and mortality for U.S. Pacific West Coast marine mammal stock assessments, 2007-2011. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-514. 92 pp.
- Castellote, M., C. Clark and M.O. Lammers. 2012. Acoustic and behavioral changes by fin whales (*Balaenoptera physalus*) in response to shipping and airgun noise. *Biological Conservation* 147:115-122.
- CCAMLR. 2017. Catch History of *Euphausia superba*. in C.T.-K.-A. Areas ed., <https://www.ccamlr.org/en/fisheries/krill-fisheries>.
- Cerchio, S., B. Andrianantenaina, T. Guillemain DEchon and A. Guillemain dEchon. 2016. Evidence for the Presence of Pygmy Blue Whales (*Balaenoptera musculus breviceuda*) in the Mozambique Channel off the Northwest Coast of Madagascar. International Whaling Commission Scientific Committee. 6 pp.
- Cerchio, S., S. Strindberg, T. Collins, C. Bennett and H. Rosenbaum. 2014. Seismic surveys negatively affect humpback whale singing activity off northern Angola. *PLoS one* 9:e86464.
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north-Atlantic areas of the U.S. Outer Continental Shelf. Cetacean and Turtle Assessment Program, Bureau of Land Management, BLM/YL/TR-82/03. 538 pp.
- Charif, R.A. and C.W. Clark. 2009. Acoustic monitoring of large whales in deep waters north and west of the British Isles: 1996 - 2005. Preliminary report. Cornell Lab of Ornithology, Bioacoustics Research Program. 40 pp.
- Christensen, G. 1955. The stocks of blue whales in the northern Atlantic. *Norsk Hvalfangst-tid* 44:640-642.
- Christensen, I., T. Haug and N. Øien. 1992. Seasonal distribution, exploitation, and present abundance of stocks of large baleen whales (Mysticeti) and sperm whales (*Physeter macrocephalus*) in Norwegian and adjacent waters. *ICES Journal of Marine Science* 49:341-355.
- Clapham, P.J. and C.S. Baker. 2002. Modern Whaling. Pages 1328-1332 in W.F. Perrin, B.

- Wursig and J.G.M. Thewissen eds. *Encyclopedia of Marine Mammals*. Academic Press, New York.
- Clapham, P.J. and R.L. Brownell, Jr. 1996. The potential for interspecific competition in baleen whales. Report of the International Whaling Commission 46:361-367.
- Clark, C. and M. Fowler. 2001. Status of archival and analysis effort of acoustic recordings during SOWER and IWC cruises 1996-2000. Report of the International Whaling Commission SC/53/IA28:9.
- Clark, C.W. 1994. Blue deep voices: Insights from Navy's Whales '93 Program. *Whalewatcher* 28:6-11.
- Clark, C.W. 1995. Application of U.S. Navy Underwater Hydrophone Arrays for Scientific Research on Whales. Report of the International Whaling Commission 45:210-212.
- Clark, C.W., W.T. Ellison, B.L. Southall, L. Hatch, S.M. Van Parijs, A. Frankel and D. Ponirakis. 2009. Acoustic masking in marine ecosystems: Intuitions, analysis, and implication. *Marine Ecology Progress Series* 395:201-222.
- Clark, C.W. and K.M. Fristrup. 2001. Baleen whale responses to low-frequency human-made underwater sounds. *Journal of the Acoustical Society of America* 110:2751.
- Clark, C.W. and G.J. Gagnon. 2004. Low-frequency vocal behaviors of baleen whales in the North Atlantic: Insights from Integrated Undersea Surveillance System detections, locations, and tracking from 1992 to 1996. *Journal of Underwater Acoustics* 52:48.
- Committee on Taxonomy. 2016. List of marine mammal species and subspecies. Society for Marine Mammalogy, <https://www.marinemammalscience.org>.
- Comtois, S., S. Claude, M. Bourassa, J. Brethes and R. Sears. 2010. Regional distribution and abundance of blue and humpback whales in the Gulf of St. Lawrence. Canadian Technical Report of Fisheries and Aquatic Sciences 2877. vii + 38 pp.
- Comtois, S., C. Savenkoff, M.-N. Bourassa, J.-C. Brethes and R. Sears. 2009. Is the change in distribution and abundance of blue whales related to the groundfish collapse in the northern Gulf of St. Lawrence? Pages 59 Eighteenth Biennial Conference on the Biology of Marine Mammals. Quebec City, Canada.
- Conn, P.B. and G.K. Silber. 2013. Vessel speed restrictions reduce risk of collision-related mortality for North Atlantic right whales. *Ecosphere* 4:Article 43.
- Conway, C.A. 2005. Global population structure of blue whales, *Balaenoptera musculus ssp.*, based on nuclear genetic variation. Doctor of Philosophy Dissertation, University of California at Davis, 115 pp.
- Costa, D.P., L.A. Hückstädt, L.K. Schwarz, A.S. Friedlaender, B.R. Mate, A.N. Zerbini, A. Kennedy and N.J. Gales. 2016. Assessing the exposure of animals to acoustic disturbance: Towards an understanding of the population consequences of disturbance. *Proceedings of Meetings in Acoustics* 27: 010027.
- Cotte, C., L. Dubroca and C. Guinet. 2006. Importance of seasonal ice zone for blue and fin whales in the Southern Ocean feeding ground. *International Whaling Commission Scientific Committee*. 7 pp.
- Cranford, T.W. and P. Krysl. 2015. Fin whale sound reception mechanisms: Skull vibration enables low-frequency hearing. *PLoS one* 10:e116222.
- Croll, D., C.W. Clark, J. Calambokidis, W.T. Ellison and B. Tershy. 2001a. Effect of anthropogenic low-frequency noise on the foraging ecology of *Balaenoptera* whales. *Animal Conservation* 4:13-27.
- Croll, D.A., A. Acevedo-Gutierrez, B.R. Tershy and J. Urban-Ramirez. 2001b. The diving

- behavior of blue and fin whales: Is dive duration shorter than expected based on oxygen stores? *Comparative Biochemistry and Physiology A Molecular and Integrative Physiology* 129:797-809.
- Croll, D.A., B. Marinovic, S. Benson, F.P. Chavez, N. Black, R. Ternullo and B.R. Tershy. 2005. From wind to whales: Trophic links in a coastal upwelling system. *Marine Ecology Progress Series* 289:117-130.
- Croll, D.A., B.R. Tershy, R.P. Hewitt, D.A. Demer, P.C. Fiedler, S.E. Smith, W. Armstrong, J.M. Popp, T. Kiekhefer, V.R. Lopez, J. Urban and D. Gendron. 1998. An integrated approach to the foraging ecology of marine birds and mammals. *Deep Sea Research Part II: Topical Studies in Oceanography* 45:1353-1371.
- Dahlheim, M.E. and D.K. Ljungblad. 1990. Preliminary hearing study on gray whales (*Eschrichtius robustus*) in the field. Pages 335-346 *Sensory Abilities of Cetaceans: Laboratory and Field Evidence*. NATO ASI Series, Series A: Life Sciences. Springer US.
- Davis, W.B. and D.J. Schmidly. 1994 (rev). *The Mammals of Texas--Online Edition*. <http://www.nsr.ttu.edu/tmot1/Default.htm>.
- De Guise, S., D. Martineau, P. Beland and M. Fournier. 1995. Possible mechanisms of action of environmental contaminants on St. Lawrence beluga whales (*Delphinapterus leucas*). *Environmental Health Perspectives* 103:73-77.
- de Vos, A. 2015. Marine life on the line. in D. Braun ed. *Ocean Views*. National Geographic.
- Deepwater Horizon Natural Resource Damage Assessment Trustees. 2016. Deepwater Horizon oil spill: Final Programmatic Damage Assessment and Restoration Plan and Final Programmatic Environmental Impact Statement. Chapter 4: Injury to Natural Resources. Retrieved from <http://www.gulfspillrestoration.noaa.gov/restoration-planning/gulf-plan>. 685 pp.
- Del-Angel-Rodriguez, J.A. 1997. Hábitos alimentarios y distribución espacio-temporal de los rorcuales común (*Balaenoptera physalus*) y azul (*Balaenoptera musculus*) en la Bahía de la Paz, B.C.S., México [Foraging habits and spatial-temporal distribution of fin whales (*Balaenoptera physalus*) and blue whales (*Balaenoptera musculus*) in La Paz Bay, B.S.C., Mexico]. M.Sc. thesis, Centro Interdisciplinario de Ciencias Marinas, Instituto Politécnico Nacional, La Paz, BCS, México, 68 p., 21 pp.
- DeRuiter, S.L., R. Langrock, T. Skirbutas, J.A. Goldbogen, J. Calambokidis, A.S. Friedlaender and B.L. Southall. 2017. A multivariate mixed hidden Markov model for blue whale behaviour and responses to sound exposure. *The Annals of Applied Statistics* 11:362-392.
- deVos, A., T. Wu and R.L. Brownell, Jr. 2013. Recent Blue Whale Deaths Due to Ship Strikes around Sri Lanka. *International Whaling Commission Scientific Committee*. 8 pp.
- Di Iorio, L. and C.W. Clark. 2009. Exposure to seismic survey alters blue whale communication. *Biol Lett* 6:51-54.
- Doniol-Valcroze, T. 2008. Habitat selection and niche characteristics of rorqual whales in the northern Gulf of St. Lawrence (Canada). Ph.D., McGill University, 146 pp.
- Doniol-Valcroze, T., D. Berteaux, P. Larouche and R. Sears. 2007. Influence of thermal fronts on habitat selection by four rorqual whale species in the Gulf of St. Lawrence. *Marine Ecology Progress Series* 335:207-216.
- Doniol-Valcroze, T., V. Lesage, J. Giard and R. Michaud. 2012. Challenges in marine mammal habitat modelling: Evidence of multiple foraging habitats from the identification of feeding events in blue whales. *Endangered Species Research* 17:255-268.
- Donovan, G.P. 1991. A review of IWC stock boundaries. *Special Issue* 13:39-68.

- Double, M.C., V. Andrews-Goff, K.C.S. Jenner, M.-N. Jenner, S.M. Laverick, T.A. Branch and N.J. Gales. 2014. Migratory movements of pygmy blue whales (*Balaenoptera musculus brevicauda*) between Australia and Indonesia as revealed by satellite telemetry. PLoS one 9:e93578.
- Dunlop, R.A., M.J. Noad, R.D. McCauley, L. Scott-Hayward, E. Kniest, R. Slade, D. Paton and D.H. Cato. 2017. Determining the behavioral dose-response relationship of marine mammals to air gun noise and source proximity. Journal of Experimental Biology 220:2878-2886.
- Eagle, T.C., S.X. Cadrin, M.E. Caldwell, R.D. Methot and M.F. Nammack. 2008. Conservation units of managed fish, threatened or endangered species, and marine mammals. U.S. Department of Commerce, NOAA Technical Memorandum. pp.
- Ellison, W.T., B.L. Southall, C.W. Clark and A.S. Frankel. 2012. A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. Conservation Biology 26:21-28.
- Ensor, P., H. Komiya, S. Kumagai, S. Kuningas, P. Olson and Y. Tsunda. 2009. 2008-2009 International Whaling Commission-Southern Ocean whale and ecosystem research (IWC-SOWER) cruise. International Whaling Commission Scientific Committee. 60 pp.
- Erbe, C. and D.M. Farmer. 2000. A software model to estimate zones of impact on marine mammals around anthropogenic noise. Journal of the Acoustical Society of America 108:1327-1331.
- Etnoyer, P., D. Canny, B. Mate and L. Morgan. 2004. Persistent Pelagic Habitats in the Baja California to Bering Sea (B2B) Ecoregion. Oceanography 17:90-101.
- Etnoyer, P., D. Canny, B.R. Mate, L.E. Morgan, J.G. Ortega-Ortiz and W.J. Nichols. 2006. Sea-surface temperature gradients across blue whale and sea turtle foraging trajectories off the Baja California Peninsula, Mexico. Deep Sea Research II 53.
- Fiedler, P., S. Reilly, R. Hewitt, D. Demer, V. Philbrick, S. Smith, W. Armstrong, D. Croll, B. Tershy and M. B. 1998. Blue whale habitat and prey in the Channel Islands. Deep Sea Research Part II-Topical Studies in Oceanography 45:1781-1801.
- Finneran, J.J. 2015. Noise-induced hearing loss in marine mammals: A review of temporary threshold shift studies from 1996 to 2015. Journal of Acoustical Society of America 138:1702-1726.
- Foote, A.D., R.W. Osborne and A.R. Hoelzel. 2004. Whale-call response to masking boat noise. Nature 428:910.
- Forney, K.A. 2007. Preliminary estimates of cetacean abundance along the U.S. West Coast and within four national marine sanctuaries during 2005. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-406. 36 pp.
- Forney, K.A., E.A. Becker, D.G. Foley, J. Barlow and E.M. Oleson. 2015. Habitat-based models of cetacean density and distribution in the central North Pacific. Endangered Species Research 27:1-20.
- Forney, K.A. and R.L. Brownell, Jr. 1996. Preliminary report of the 1994 Aleutian Island marine mammal survey. International Whaling Commission Scientific Committee, Paper SC/48/011. pp.
- Fossette, S., B. Abrahms, E.L. Hazen, S.J. Bograd, K.M. Zilliacus, J. Calambokidis, J.A. Burrows, J.A. Goldbogen, J.T. Harvey, B. Marinovic, B. Tershy and D.A. Croll. 2017. Resource partitioning facilitates coexistence in sympatric cetaceans in the California Current. Ecol Evol 7:9085-9097.
- Francis, R.C., S.R. Hare, A.B. Hollowed and W.S. Wooster. 1998. Effects of interdecadal

- climate variability on the oceanic ecosystems of the NE Pacific. *Fisheries Oceanography* 7:1-21.
- Frankel, A.S. 2005. Gray whales hear and respond to a 21-25 kHz high-frequency whale-finding sonar. Pages 97 Sixteenth Biennial Conference on the Biology of Marine Mammals. San Diego, California.
- Friedlaender, A.S., J.A. Goldbogen, E.L. Hazen, J. Calambokidis and B.L. Southall. 2015. Feeding performance by sympatric blue and fin whales exploiting a common prey resource. *Marine Mammal Science* 31:345-354.
- Friedlaender, A.S., P.N. Halpin, S.S. Qian, G.L. Lawson, P.H. Wiebe, D. Thiele and A.J. Read. 2006. Whale distribution in relation to prey abundance and oceanographic processes in shelf waters of the Western Antarctic Peninsula. *Marine Ecology Progress Series* 317:297-310.
- Friedlaender, A.S., G.L. Lawson and P.N. Halpin. 2009. Evidence of resource partitioning between humpback and minke whales around the western Antarctic Peninsula. *Marine Mammal Science* 25:402-415.
- Fristrup, K.M., L.T. Hatch and C.W. Clark. 2003. Variation in humpback whale (*Megaptera novaeangliae*) song length in relation to low-frequency sound broadcasts. *Journal of the Acoustical Society of America* 113:3411-3424.
- Fujise, Y., T. Kishiro, R. Zenitani, K. Matsuoka, M. Kawasaki and K. Shimamoto. 1995. Cruise report of the Japanese whale research program under a Special Permit for North Pacific minke whales in 1994. International Whaling Commission Scientific Committee. 29 pp.
- Gales, R.S. 1982. Effects of noise of offshore oil and gas operations on marine mammals - an introductory assessment. U.S. Bureau of Land Management and Naval Ocean Systems Center. 79 + appendices pp.
- Galletti Vernazzani, B., C. Burton, M. Double, P. Gill, C. Jenner, M. Jenner, P. Olson and C. Salgado-Kent. 2016. Comparisons among Southern Hemisphere Blue Whale Catalogue off Australia and New Zealand. International Whaling Commission Scientific Committee. 6 pp.
- Galletti Vernazzani, B., C.A. Carlson, E. Cabrera and R.L. Brownell, Jr. 2012. Chilean blue whales off Isla Grande de Chiloe, 2004-2010: distribution, site-fidelity and behaviour. *Journal of Cetacean Research And Management* 12:353-360.
- Gambell, R. 1973. Some effects of exploitation on reproduction in whales. *Journal of Reproductive Fertility Supplement* 19:533-553.
- Gambell, R. 1979. The blue whale. *Biologist* 26:209-215.
- Garrison, L.P. and J.S. Link. 2000. Fishing effects on spatial distribution and trophic guild structure of the fish community in the Georges Bank region. *ICES Journal of Marine Science* 57:723-730.
- Gavrilchuk, K., V. Lesage, C. Ramp, R. Sears, M. Berube, S. Bearhop and G. Beauplet. 2014. Trophic niche partitioning among sympatric baleen whale species following the collapse of groundfish stocks in the Northwest Atlantic. *Marine Ecology Progress Series* 497:285-301.
- Gedamke, J. and S.M. Robinson. 2010. Acoustic survey for marine mammal occurrence and distribution off East Antarctica (30-80°E) in January-February 2006. *Deep Sea Research Part II: Topical Studies in Oceanography* 57:968-981.
- Gendron, D. 1990. Relacion entre la abundancia de eufausidos y de ballenas azules (*Balaenoptera musculus*) en el Golfo de California. M.Sc. thesis, Centro Interdisciplinario de Ciencias Marinas, Instituto Politécnico Nacional, La Paz, BCS, México . 64p.
- Gendron, D. 1992. Population structure of daytime surface swarms of *Nyctiphanes simplex* (*Euphausiacea*) in the Gulf of California, Mexico. *Marine Ecology Progress Series* 87:1-6.

- Gendron, D. and V.Z. Hernandez. 1995. Blue whales of Baja California: A summary of their distribution and preliminary reproductive data based on photoidentification. Pages 43 Eleventh Biennial Conference on the Biology of Marine Mammals. Orlando, Florida.
- Gendron, D. and R. Sears. 1993. Blue whales and *Nyctiphanes simplex* surface swarms: A close relationship in the southwest Gulf of California, Mexico. Pages 52 Tenth Biennial Conference on the Biology of Marine Mammals. Galveston, Texas.
- Geraci, J.R. 1990. Physiologic and toxic effects on cetaceans. Pages 167-197 in J.R. Geraci and D.J.S. Aubin eds. *Sea Mammals and Oil: Confronting the Risks*. Academic Press, San Diego.
- Geraci, J.R., D.M. Anderson, R.J. Timperi, D.J. St. Aubin, G.A. Early, J.H. Prescott and C.A. Mayo. 1989. Humpback whales (*Megaptera novaeangliae*) fatally poisoned by dinoflagellate toxin. *Canadian Journal of Fisheries and Aquatic Sciences* 46.
- Gill, P.C. 2002. A blue whale (*Balaenoptera musculus*) feeding ground in a southern Australian coastal upwelling zone. *Journal of Cetacean Research And Management* 4:179-184.
- Gill, P.C., M.G. Morrice, B. Page, R. Pirzl, A.H. Levings and M. Coyne. 2011. Blue whale habitat selection and within-season distribution in a regional upwelling system off southern Australia. *Marine Ecology Progress Series* 421:243-263.
- Gilpatrick, J.W., Jr. and W.L. Perryman. 2008. Geographic variation in external morphology of North Pacific and Southern Hemisphere blue whales (*Balaenoptera musculus*). *Journal of Cetacean Research And Management* 10:9-21.
- Goldbogen, J.A., J. Calambokidis, E. Oleson, J. Potvin, N.D. Pyenson, G. Schorr and R.E. Shadwick. 2011. Mechanics, hydrodynamics and energetics of blue whale lunge feeding: Efficiency dependence on krill density. *Journal of Experimental Biology* 214:131-146.
- Goldbogen, J.A., B.L. Southall, S.L. DeRuiter, J. Calambokidis, A.S. Friedlaender, E.L. Hazen, E.A. Falcone, G.S. Schorr, A. Douglas, D.J. Moretti, C. Kyburg, M.F. McKenna and P.L. Tyack. 2013. Blue whales respond to simulated mid-frequency military sonar. *Proceedings of the Royal Society B* 280:20130657.
- González, H.E., M.J. Calderón, L. Castro, A. Clement, L.A. Cuevas, G. Daneri, J.L. Iriarte, L. Lizárraga, R. Martínez, E. Menschel, N. Silva, C. Carrasco, C. Valenzuela, C.A. Vargas and C.M. . 2010. Primary production and plankton dynamics in the Reloncaví Fjord and the Interior Sea of Chiloé, Northern Patagonia, Chile. *Marine Ecology Progress Series* 402:13-30.
- Greene, C.R., Jr. 1987. Characteristics of oil industry dredge and drilling sounds in the Beaufort Sea *Journal of Acoustical Society of America* 82:1315-1324.
- Gulland, F.M.D. and A.J. Hall. 2007. Is marine mammal health deteriorating? Trends in the global reporting of marine mammal disease. *Ecohealth* 4:135-150.
- Guzmán-Verri, C., R. González-Barrientos, G. Hernández-Mora, J.-A. Morales, E. Baquero-Calvo, E. Chaves-Olarte and E. Moreno. 2012. *Brucella ceti* and brucellosis in cetaceans. *Frontiers in Cellular and Molecular Microbiology* 2:22.
- Hamazaki, T. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, North Carolina, USA to Nova Scotia, Canada). *Marine Mammal Science* 18:920-939.
- Hammond, P., R. Sears and M. Berube. 1990. A note on problems in estimating the number of blue whales in the Gulf of St Lawrence from photo-identification data. Report of the International Whaling Commission Special Issue 12:141-142.
- Hanson, J. and G. Chouinard. 2002. Diet of Atlantic cod in the southern Gulf of St. Lawrence as an index of ecosystem change, 1959-2000. *Journal of Fish Biology* 60:902-922.
- Harvey, J. and M. Dahlheim. 1994. Cetaceans in oil. Pages 257-264 in T. Loughlin ed. *Impacts*

- of the Exxon Valdez Oil Spill on Marine Mammals*. Academic Press, San Diego.
- Harvey, M. and L. Devine. 2008. Oceanographic conditions in the Estuary and the Gulf of St. Lawrence during 2007: zooplankton. Fisheries and Oceans Canada. pp.
- Hays, G., A. Richardson and C. Robinson. 2005. Climate change and marine plankton. Trends in Ecology and Evolution 20:337-344.
- Hazen, E.L., D.M. Palacios, K.A. Forney, E.A. Howell, E. Becker, A.L. Hoover, L. Irvine, M. DeAngelis, S.J. Bograd, B.R. Mate and H. Bailey. 2016. WhaleWatch: a dynamic management tool for predicting blue whale density in the California Current. Journal of Applied Ecology 10.1111/1365-2664.12820:14.
- Heyning, J.E. and T.D. Lewis. 1990. Entanglements of baleen whales in fishing gear off southern California. Report of the International Whaling Commission 40:427-431.
- Hinton, M.A.C. 1915. Report on the papers left by the late Major Barrett-Hamilton, relating to the whales of South Georgia. Appendix VII. Inter-departmental committee on whaling and the protection of whales. Crown Agents for the Colonies, London. Colonial Office Miscellaneous Number 298:69-193.
- Hjort, J. and T. Ruud. 1929. Whaling and fishing in the North Atlantic. Rapports Et Proces-Verbaux Des Reunions Conseil International Pour L'Exploration de la Mer 56:1-123.
- Holt, M.M., D.P. Noren, V. Veirs, C.K. Emmons and S. Veirs. 2009. Speaking up: Killer whales (*Orcinus orca*) increase their call amplitude in response to vessel noise. Journal of the Acoustical Society of America 125:E127-E132.
- Houser, D., R. Howard and S. Ridgway. 2001. Can diving behavior increase the chance of acoustically driven bubble growth in marine mammals? Pages 103 Fourteenth Biennial Conference on the Biology of Marine Mammals. Vancouver, Canada.
- Hucke-Gaete, R., L.P. Osman, C.A. Moreno, K.P. Findlay and D.K. Ljungblad. 2004. Discovery of a blue whale feeding and nursing ground in southern Chile. Proceedings of the Royal Society of London Series B Biological Sciences 271:S170-S173.
- Hucke-Gaete, R., F.A. Vidali, L. Bedrianana-Romano and A. Zerbini. 2016. Satellite tracking of blue whales (*Balaenoptera musculus*) from Patagonia (Chile) to Galapagos (Ecuador): novel insights into migratory pathways along the Eastern South Pacific. International Whaling Commission Scientific Committee. 11 pp.
- Ichihara, T. 1966. The pygmy blue whale, *Balaenoptera musculus brevicauda*, a new subspecies from the Antarctic. Pages 79-113 *Whales, Dolphins and Porpoises*. K. S. Norris (ed.). University of California Press, Berkeley, CA.
- Ilangakoon, A.D. and K. Sathasivam. 2012. The need for taxonomic investigations on Northern Indian Ocean blue whales (*Balaenoptera musculus*): implications of year-round occurrence off Sri Lanka and India. Journal of Cetacean Research And Management 12:195-202.
- Irvine, L.M., B.R. Mate, M.H. Winsor, D.M. Palacios, S.J. Bograd, D.P. Costa and H. Bailey. 2014. Spatial and temporal occurrence of blue whales off the U. S. West Coast, with implications for management. PLoS one 9:e102959.
- Ivashin, M.V. and A.A. Rovnin. 1967. Some results of the Soviet whale marking in the waters of the North Pacific. Norsk Hvalfangst-Tidende 56:123-135.
- IWC. 2016a. Report of the Scientific Committee. International Whaling Commission. 138 pp.
- IWC. 2016b. Report of the Scientific Committee. Journal of Cetacean Research And Management, 17 (Supplement). 92 pp.
- IWC. 2017. Report of the Scientific Committee. International Whaling Commission. 136 pp.
- Jackson, J.A. 2016. A preliminary population assessment of the recovery status of Chilean blue

- whales. International Whaling Commission Scientific Committee. 22 pp.
- Jenner, C., M. Jenner, C. Burton, V. Sturrock, C.S. Kent, M. Morrice, C. Attard, L. Moller and M.C. Double. 2008. Mark recapture analysis of pygmy blue whales from the Perth Canyon, Western Australia 2000-2005. International Whaling Commission Scientific Committee. 9 pp.
- Jensen, A.S. and G.K. Silber. 2004. Large Whale Ship Strike Database. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-OPR-25. 37 pp.
- Jenssen, B.M., O. Haugen, E.G. Sormo and J.U. Skaare. 2003. Negative relationship between PCBs and plasma retinol in low-contaminated free-ranging gray seal pups (*Halichoerus grypus*). *Environmental Research* 93:79-87.
- Johnson, J.B. and K.S. Omland. 2004. Model selection in ecology and evolution. *Trends in Ecology and Evolution* 19:101-108.
- Jones, M.C. and W.W.L. Cheung. 2015. Multi-model ensemble projections of climate change effects on global marine biodiversity. *ICES Journal of Marine Science* 72:741-752.
- Jonsgard, A. 1955. The stocks of blue whales (*Balaenoptera musculus*) in the northern Atlantic Ocean and adjacent Arctic waters. *Norsk Hvalfangst-Tidende* 44:505-519.
- Jonsgard, A. 1977. Tables showing the catch of small whales (including minke whales) caught by Norwegians in the period 1938-75, and large whales caught in different North Atlantic waters in the period 1868-1975. Report of the International Whaling Commission 27:413-426.
- Kapel, F.O. 1979. Exploitation of large whales in West Greenland in the twentieth century. Report of the International Whaling Commission 29:197-214.
- Kasahara, A. 1950. Whaling and its resources in the Japanese coastal waters. Bulletin of the Research Institute of the Nippon Suisan Company Limited. 4. 103 pp + 195 figures pp.
- Kasamatsu, F., K. Matsuoka and T. Hakamada. 2000. Interspecific relationships in density among the whale community in the Antarctic. *Polar Biology* 23:466-473.
- Kato, H., K. Matsuoka, S. Nishiwaki and J.L. Bannister. 2007. Distributions and abundances of pygmy blue whales and southern right whales in waters off the southern coast of Australia, based on data from the Japan/IWC Blue Whale Cruise 1995-96. International Whaling Commission Scientific Committee. 14 pp.
- Kawaguchi, S., A. Ishida, R. King, B. Raymond, N. Waller, A. Constable, S. Nicol, M. Wakita and A. Ishimatsu. 2013. Risk maps for Antarctic krill under projected Southern Ocean acidification. *Nature Climate Change* 3:843-847.
- Kawaguchi, S., H. Kurihara, R. King, L. Hale, T. Berli, J.P. Robinson and A. Ishimatsu. 2011. Will krill fare well under Southern Ocean acidification? *Biol Lett* 7:288-291.
- Kawamura, A. 1980. A review of food of balaenopterid whales. *Scientific Reports of the Whales Research Institute* 32:155-197.
- Kenney, R.D. and S.D. Kraus. 1993. Right whale mortality - a correction and an update. *Marine Mammal Science* 9:445-446.
- Ketten, D. and D.C. Mountain. 2012. Modeling Minke Whale Hearing. United Kingdom: Joint Industry Programme.
- Ketten, D.R. 1992a. The cetacean ear: Form, frequency, and evolution. Pages 53-75 in J.A. Thomas, R.A. Kastelein and A.Y. Supin eds. *Marine Mammal Sensory Systems*. Plenum Press, New York.
- Ketten, D.R. 1992b. The marine mammal ear: Specializations for aquatic audition and echolocation. Pages 717-754 in D. Webster, R. Fay and A. Popper eds. *The Biology of Hearing*. Springer-Verlag.
- Ketten, D.R. 1994. Functional analyses of whale ears: Adaptations for underwater hearing. Pages

264-270 Oceans 1994.

- Ketten, D.R. and D.C. Mountain. 2014. Inner ear frequency maps: First stage audiograms of low to infrasonic hearing in mysticetes. Pages 41 Fifth International Meeting on the Effects of Sounds in the Ocean on Marine Mammals (ESOMM - 2014). Amsterdam, The Netherlands.
- Kieckhefer, T.R., J. Calambokidis, G.H. Steiger and N.A. Black. 1995. Prey of humpback and blue whales off California based on identification of hard parts in feces. Pages 62 Eleventh Biennial Conference on the Biology of Marine Mammals. Orlando, Florida.
- Knight, K. 2011. Blue whale-sized mouthfuls make foraging super efficient. *Journal of Experimental Biology* 214.
- Koeller, P.A. 2000. Relative importance of abiotic and biotic factors to the management of the Northern shrimp (*Pandalus borealis*) fishery on the Scotian Shelf. *Journal of Northwest Atlantic Fishery Science* 27:21-33.
- Krahn, M., M.J. Ford, W.F. Perrin, P.R. Wade, R.P. Angliss, M.B. Hanson, B.L. Taylor, G.M. Ylitalo, M.E. Dahlheim, J.E. Stein and R.S. Waples. 2004. 2004 Status review of Southern Resident killer whales (*Orcinus orca*) under the Endangered Species Act. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-NWFSC-62 73 pp.
- Laist, D.W., A.R. Knowlton, J.G. Mead, A.S. Collet and M. Podesta. 2001. Collisions between ships and whales. *Marine Mammal Science* 17:35-75.
- Laist, D.W., A.R. Knowlton and D. Pendleton. 2014. Effectiveness of mandatory vessel speed limits for protecting North Atlantic right whales. *Endangered Species Research* 23:133-147.
- Lambertsen, R.H., K.J. Rasmussen, W.C. Lancaster and R.J. Hintz. 2005. Functional morphology of the mouth of the bowhead whale and its implications for conservation. *Journal of Mammalogy* 86:342-352.
- Landsberg, J.H., K.A. Lefebvre and L.J. Flewelling. 2014. Effects of Toxic Microalgae on Marine Organisms. Pages 379-449 in G.P. Rossini ed. *Toxins and Biologically Active Compounds from Microalgae*. Biological Effects and Risk Management. CRC Press.
- Larsen, J.N., O.A. Anisimov, A. Constable, A.B. Hollowed, N. Maynard, P. Prestrud, T.D. Prowse and J.M.R. Stone. 2014. Polar Regions. Pages 1567-1612 in V.R. Barros, C.B. Field, D.J. Dokken, M.D. Mastrandrea, K.J. Mach, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. Maccracken, P.R. Mastrandrea and L.L. White eds. *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- LeDuc, R.G., E.I. Archer, A.R. Lang, K.K. Martien, B. Hancock-Hanser, J.P. Torres-Florez, R. Huckle-Gaete, H.R. Rosenbaum, K. Van Waerebeek, R.L. Brownell, Jr. and B.L. Taylor. 2016. Genetic variation in blue whales in the eastern Pacific: implication for taxonomy and use of common wintering grounds. *Molecular Ecology* 10.1111/mec.13940.
- LeDuc, R.G., A.E. Dizon, M. Goto, L.A. Pastene, H. Kato, S. Nishiwaki, C.A. LeDuc and R.L. Brownell. 2007. Patterns of genetic variation in Southern Hemisphere blue whales, and the use of assignment test to detect mixing on the feeding grounds. *Journal of Cetacean Research And Management* 9:73-80.
- Lefebvre, K.A., S. Bargu, T. Kieckhefer and M.W. Silver. 2002. From sanddabs to blue whales: The pervasiveness of domoic acid. *Toxicon* 40:971-977.
- Lesage, V., J.-F. Gosselin, M. Hammill, M.C.S. Kingsley and J. Lawson. 2007. Ecologically and Biologically Significant Areas (EBSAs) in the estuary and Gulf of St. Lawrence – A marine

- mammal perspective. Fisheries and Oceans Canada. 96 pp.
- Lilly, G.R. 1991. Interannual variability in predation by cod (*Gadus morhua*) on capelin (*Mallotus villosus*) and other prey off Southern Labrador and Northeastern Newfoundland. ICES Marine Symposia 1991 193:133-146.
- Lilly, G.R., D.G. Parsons and D.W. Kulka. 2000. Was the Increase in Shrimp Biomass on the Northeast Newfoundland Shelf a Consequence of a Release in Predation Pressure from Cod? J. Northw. Atl. Fish. Sci. 27:45-61.
- Ljungblad, D.K., C.W. Clark and H. Shimada. 1998. A comparison of sounds attributed to pygmy blue whales (*Balaenoptera musculus breviceauda*) recorded south of the Madagascar Plateau and those attributed to 'true' blue whales (*Balaenoptera musculus*) recorded off Antarctica. Report of the International Whaling Commission 48:439-442.
- Lockyer, C. 1979. Changes in a growth parameter associated with exploitation of southern fin and sei whales. Report of the International Whaling Commission 29:191-196.
- Lockyer, C. 1984. Review of baleen whale (Mysticeti) reproduction and implications for management. Report of the International Whaling Commission Special Issue 6:27-50.
- Lucifredi, I. and P.J. Stein. 2007. Gray whale target strength measurements and the analysis of the backscattered response. Journal of the Acoustical Society of America 121:1383-1391.
- Lusseau, D. 2004. The hidden cost of tourism: Detecting long-term effects of tourism using behavioral information. Ecology and Society 9:2.
- Lusseau, D. and L. Bejder. 2007. The long-term consequences of short-term responses to disturbance experiences from whale watching impact assessment. International Journal of Comparative Psychology 20:228-236.
- Mackas, D.L., R. Goldblatt and A.G. Lewis. 1998. Interdecadal variation in developmental timing of *Neocalanus plumchrus* populations at Ocean Station P in the subArctic North Pacific. Canadian Journal of Fisheries and Aquatic Sciences 55:1878-1893.
- Mantua, N.J. 2002. Impacts of Pacific Decadal Variability on Marine Ecosystems. American Geophysical Union, Spring Meeting 2002. Washington, DC.
- Mate, B.R., B.A. Lagerquist and J. Calambokidis. 1999. Movements of North Pacific blue whales during the feeding season off southern California and their southern fall migration. Marine Mammal Science 15:12.
- Matteson, R.S., J.K. Benoit-Bird, B.R. Mate and J. Calambokidis. 2010. Aggregation characteristics of prey determine blue whale distribution at the Costa Rica Dome. 2010 Ocean Sciences Meeting. American Geophysical Union, Washington DC Portland, OR.
- McCauley, R.D. and C. Jenner. 2010. Migratory patterns and estimated population size of pygmy blue whales (*Balaenoptera musculus breviceauda*) traversing the Western Australian coast based on passive acoustics. International Whaling Commission Scientific Committee. 9 pp.
- McCauley, R.D., C. Jenner, J. Bannister, C. Burton, D. Cato and A. Duncan. 2001. Blue whale calling in the Rottneest Trench - 2000, western Australia. Report R2001-6, Center for Marine Science and Technology, Curtin University of Technology, Perth, Western Australia:56pp.
- McCauley, R.D., C.P.S. Kent, C.L.K. Burton and C. Jenner. 2006. Blue whale calling in Australian waters. Journal of the Acoustical Society of America 120:3266.
- McDonald, M.A. 2006. An acoustic survey of baleen whales off Great Barrier Island, New Zealand. New Zealand Journal of Marine and Freshwater Research 40:519-529.
- McDonald, M.A., J. Calambokidis, A.M. Teranishi and J.A. Hildebrand. 2001. The acoustic calls of blue whales off California with gender data. Journal of the Acoustical Society of America 109:1728-1735.

- McDonald, M.A., J.A. Hildebrand and S. Mesnick. 2009. Worldwide decline in tonal frequencies of blue whale songs. *Endangered Species Research* 9:13-21.
- McDonald, M.A., J.A. Hildebrand and S.C. Webb. 1995. Blue and fin whales observed on a seafloor array in the northeast Pacific. *Journal of the Acoustical Society of America* 98:712-721.
- McDonald, M.A., J.A. Hildebrand and S.M. Wiggins. 2006a. Increases in deep ocean ambient noise in the Northeast Pacific west of San Nicolas Island, California. *Journal of Acoustical Society of America* 120:711-718.
- McDonald, M.A., S.L. Mesnick and J.A. Hildebrand. 2006b. Biogeographic characterization of blue whale song worldwide: Using song to identify populations. *Journal of Cetacean Research And Management* 8:55-66.
- McKay, S., A. Sirovic and D. Thiele. 2005. Preliminary results of blue whale (*Balaenoptera musculus*) call detections in eastern Antarctica. *International Whaling Commission Scientific Committee*. 4 pp.
- McKenna, M.F., J. Calambokidis, E.M. Oleson, D.W. Laist and J.A. Goldbogen. 2015. Simultaneous tracking of blue whales and large ships demonstrates limited behavioral responses for avoiding collision. *Endangered Species Research* 27:219-232.
- McKenna, M.F., D. Ross, S.M. Wiggins and J.A. Hildebrand. 2012. Underwater radiated noise from modern commercial ships. *Journal of the Acoustical Society of America* 131:92-103.
- McWilliams, J.N. and A.D. Hawkins. 2013. A comparison of inshore marine soundscapes. *Journal of Experimental Marine Biology and Ecology* 446:166-176.
- Mead, J.G. and R.L. Brownell. 1993. Order Cetacea. Pages 349-364 in D.E. Wilson and D.M. Reeder eds. *Mammal Species of the World: A Taxonomic and Geographic Reference*. Smithsonian Institution Press, Washington.
- Melcon, M.L., A.J. Cummins, S.M. Kerosky, L.K. Roche, S.M. Wiggins and J.A. Hildebrand. 2012. Blue whales respond to anthropogenic noise. *PLoS one* 7:e32681.
- Metcalfe, C., B. Koenig, T. Metcalfe, G. Paterson and R. Sears. 2004. Intra- and inter-species differences in persistent organic contaminants in the blubber of blue whales and humpback whales from the Gulf of St. Lawrence, Canada. *Marine Environmental Research* 57:245-260.
- Mikhalev, Y.A. 2000. Whaling in the Arabian Sea by the whaling fleets slava and Sovetskaya Ukraina. Pages 141-181 *Soviet Whaling Data (1949-1979)*. Center for Russian Environmental Policy Marine Mammal Council, Moscow.
- Miller, B.S., K. Collins, J. Barlow, S. Calderan, R. Leaper, M. McDonald, P. Ensor, P.A. Olson, C. Olavarria and M.C. Double. 2014. Blue whale vocalizations recorded around New Zealand: 1964-2013. *Journal of the Acoustical Society of America* 135:1616-1623.
- Mitchell, E. 1974. Present status of northwest Atlantic fin and other whale stocks. Pages 108-169 in W.E. Schevill ed. *The Whale Problem*. Harvard University Press, Cambridge, Massachusetts.
- Mizroch, S.A., D.W. Rice and J.M. Breiwick. 1984. The blue whale, *Balaenoptera musculus*. *Marine Fisheries Review* 46:15-19.
- Mizue, K. 1951. Food of whales (in the adjacent waters of Japan). *Scientific Reports of the Whales Research Institute Tokyo* 5:81-90.
- Monnahan, C.C., T.A. Branch and A.E. Punt. 2015. Do ship strikes threaten the recovery of endangered eastern North Pacific blue whales? *Marine Mammal Science* 31:279-297.
- Monnahan, C.C., T.A. Branch, K.M. Stafford, Y.V. Ivashchenko and E.M. Oleson. 2014. Estimating historical eastern North Pacific blue whale catches using spatial calling patterns.

PLoS one 9:e98974.

- Moore, J. and M. Abbott. 2000. Phytoplankton chlorophyll distributions and primary production in the Southern Ocean. *Journal of Geophysical Research* 105:28709-28722.
- Moore, M.J., C.A. Miller, A.V. Weisbrod, D. Shea, P.K. Hamilton, S.D. Kraus, V.J. Rowntree, N. Patenaude and J.J. Stegeman. 1998. Cytochrome P450 1A and chemical contaminants in dermal biopsies of northern and southern right whales. International Whaling Commission Scientific Committee. *Comprehensive Assessment of Right Whales: A Worldwide Comparison*. 18 pp.
- Moore, M.J. and J.M. Van Der Hoop. 2012. The painful side of trap and fixed net fisheries: Chronic entanglement of large whales. *Journal of Marine Biology*:230653.
- Moore, P.W.B. and R.J. Schusterman. 1987. Audiometric assessment of northern fur seals, *Callorhinus ursinus*. *Marine Mammal Science* 3:31-53.
- Moore, S.K., V.L. Trainer, N.J. Mantua, M.S. Parker, E.A. Laws, L.C. Backer and L.E. Fleming. 2008. Impacts of climate variability and future climate change on harmful algal blooms and human health. *Environmental Health* 7:S4.
- Mori, M. and D.S. Butterworth. 2004. Consideration of multispecies interactions in the Antarctic: A preliminary model of the minke whale – blue whale – krill interaction. *African Journal of Marine Science* 26:245-259.
- Munger, L.M., D. Camacho, A. Havron, G. Campbell, J. Calambokidis, A. Douglas and J. Hildebrand. 2009. Baleen whale distribution relative to surface temperature and zooplankton abundance off southern California, 2004-2008. *California Cooperative Oceanic Fisheries Investigations Reports* 50:155-168.
- Murase, H., K. Matsuoka, T. Ichii and S. Nishiwaki. 2002. Relationship between the distribution of euphausiids and baleen whales in the Antarctic (35°E - 145°W). *Polar Biology* 25:135-145.
- Murase, H., T. Tamura, K. Matsuoka and T. Hakamada. 2006. First attempt of estimation of feeding impact on krill standing stock by three baleen whale species (Antarctic minke, humpback and fin whales) in Areas IV and V using JARPA data. Paper SC/D06/J22 presented to the JARPA Review Meeting, December 2006.:7.
- NAMMCO. 2014. NAMMCO Annual Report 2014. North Atlantic Marine Mammal Commission. 247 pp.
- National Research Council. 2003. *Ocean Noise and Marine Mammals*. National Academies Press, Washington, D.C.
- National Research Council. 2005. *Marine Mammal Populations and Ocean Noise*. National Academies Press, Washington, D.C.
- National Snow & Ice Data Center. 2017. Arctic sea ice maximum at record low for third straight year. *Arctic Sea Ice News & Analysis*. <http://nsidc.org/arcticseaicenews/2017/03/arctic-sea-ice-maximum-at-record-low/>.
- Nemoto, T. 1957. Foods of baleen whales in the northern Pacific. *Scientific Reports of the Whales Research Institute* 12:33-89.
- Nemoto, T. 1973. Feeding pattern of baleen whales in the oceans. Pages 241-252 in J.H. Steele ed. *Marine food chains*. University of California Press, Berkeley, California.
- Nemoto, T. and A. Kawamura. 1977. Characteristics of food habits and distribution of baleen whales with special reference to the abundance of North Pacific sei and Bryde's whales. *Report of the International Whaling Commission Special Issue* 1:80-87.
- New, L.F., Clark, J.S., Costa, D.P., Fleishman, E., Hindell, M.A., Klanjš, T., Lusseau, D., Kraus, S., McMahon, C.R., Robinson, P.W., Schick, R.S., Schwarz, L.K., Simmons, S.E., Thomas,

- L., Tyack, P., Harwood, J. . 2014. Using short-term measures of behavior to estimate long-term fitness of southern elephant seals. *Marine Ecology Progress Series* 496:99-108.
- New, L.F., J. Harwood, L. Thomas, C. Donovan, J.S. Clark, G. Hastie, P.M. Thompson, B. Cheney, L. Scott-Hayward and D. Lusseau. 2013a. Modelling the biological significance of behavioural change in coastal bottlenose dolphins in response to disturbance. *Functional Ecology* 27:314-322.
- New, L.F., D.J. Moretti, S.K. Hooker, D.P. Costa and S.E. Simmons. 2013b. Using Energetic Models to Investigate the Survival and Reproduction of Beaked Whales (family Ziphiidae). *PLoS one* 8:e68725.
- Nicol, S., J. Foster and S. Kawaguchi. 2012. The fishery for Antarctic krill – recent developments. *Fish and Fisheries* 13:30-40.
- Nieukirk, S.L., D.K. Mellinger, J.A. Hildebrand, M.A. McDonald and R.P. Dziak. 2005. Downward shift in the frequency of blue whale vocalizations. Pages 205 Sixteenth Biennial Conference on the Biology of Marine Mammals. San Diego, California.
- Nieukirk, S.L., K.M. Stafford, D.K. Mellinger, R.P. Dziak and C.G. Fox. 2004. Low-frequency whale and seismic airgun sounds recorded in the mid-Atlantic Ocean. *Journal of the Acoustical Society of America* 115:1832-1843.
- Nishiwaki, M. 1966. Distribution and migration of the larger cetaceans in the North Pacific as shown by Japanese whaling results. Pages 171-191 *Whales, Dolphins and Porpoises*. University of California Press, Berkeley.
- NMFS. 2010a. Final Recovery Plan for the Fin Whale (*Balaenoptera physalus*). National Marine Fisheries Service, Office of Protected Resources. 121 pp.
- NMFS. 2010b. Final Recovery Plan for the Sperm Whale (*Physeter macrocephalus*). National Marine Fisheries Service, Office of Protected Resources. 165 pp.
- NMFS. 2011. Final Recovery Plan for the Sei Whale (*Balaenoptera borealis*). National Marine Fisheries Service, Office of Protected Resources. 108 pp.
- NMFS. 2012. Endangered Species Act Section 7 Biological Opinion on the Continued Operation of the Hawaii-based Shallow-set Longline Swordfish Fishery under Amendment 18 to the Fishery Management Plan for Pelagic Fisheries of the Western Pacific Region. NMFS Pacific Islands Regional Office. 162 + Technical Correction pp.
- NMFS. 2013. Final Recovery Plan for the North Pacific Right Whale (*Eubalaena japonica*). National Marine Fisheries Service, Office of Protected Resources. 84 pp.
- NMFS. 2014. Endangered Species Act Section 7 Biological Opinion on the Continued Operation of the Hawaii-based Deep-set Pelagic Longline Fishery. NMFS Pacific Islands Regional Office. 216 pp.
- NMFS. 2016. Guidelines for Preparing Stock Assessment Reports Pursuant to the 1994 Amendments to the MMPA NMFS Instruction 02-204-01.
- NMFS. 2017. 2016 West Coast Entanglement Summary. in N.W.C. Region ed., [http://www.westcoast.fisheries.noaa.gov/mediacenter/WCR%202016%20Whale%20Entanglements 3-26-17 Final.pdf](http://www.westcoast.fisheries.noaa.gov/mediacenter/WCR%202016%20Whale%20Entanglements%203-26-17%20Final.pdf).
- NMFS and U.S. Department of the Navy. 2001. Joint interim report Bahamas marine mammal stranding event 15 - 16 March 2000. U.S. Department of the Navy and U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service. pp.
- NOAA. 2011. Effects of oil and gas activities in the Arctic Ocean: Draft environmental impact statement.

- NOAA. 2016. Ocean Noise Strategy Roadmap. 90 + Appendices pp.
- Northrop, J., W.C. Cummings and M.F. Morrison. 1970. Underwater 20-Hz signals recorded near Midway Island. *Journal of the Acoustical Society of America* 49:1909-1910.
- Nowacek, D.P., M.P. Johnson and P.L. Tyack. 2004. North Atlantic right whales (*Eubalaena glacialis*) ignore ships but respond to alerting stimuli. *Proceedings of the Royal Society of London Series B Biological Sciences* 271:227-231.
- Nowacek, D.P., L.H. Thorne, D.W. Johnston and P.L. Tyack. 2007. Responses of cetaceans to anthropogenic noise. *Mammal Review* 37:81-115.
- O'Shea, T. and R.L. Brownell, Jr. . 1994. Organochlorine and metal contaminants in baleen whales: A review and evaluation of conservation implications. *Science of the Total Environment* 154:179-200.
- Ohsumi, S. and Y. Masaki. 1975. Japanese whale marking in the North Pacific, 1963-1972. *Bulletin of the Far Seas Fisheries Research Laboratory* 12:171-219.
- Ohsumi, S. and S. Wada. 1974. Status of whale stocks in the North Pacific, 1972. *Report of the International Whaling Commission* 24:114-126.
- Oleson, E.M., C.H. Boggs, K.A. Forney, M.B. Hanson, D.R. Kobayashi, B.L. Taylor, P.R. Wade and G.M. Ylitalo. 2010. Status Review of Hawaiian Insular False Killer Whales (*Pseudorca crassidens*) under the Endangered Species Act U.S. Department of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-22. 140 + Appendices pp.
- Oleson, E.M., J. Calambokidis, W.C. Burgess, M.A. McDonald, C.A. Leduc and J.A. Hildebrand. 2007a. Behavioral context of call production by eastern North Pacific blue whales. *Marine Ecology Progress Series* 330:269-284.
- Oleson, E.M., S.M. Wiggins and J.A. Hildebrand. 2007b. Temporal separation of blue whale call types on a southern California feeding ground. *Animal Behaviour* 74:881-894.
- Olson, P.A., P. Ensor, C. Olavarria, N. Bott, R. Constantine, J. Weir, S. Childerhouse, M. van der Linde, N. Schmitt, B.S. Miller and M.C. Double. 2015. New Zealand Blue Whales: Residency, Morphology, and Feeding Behavior of a Little - Known Population. *Pacific Science* 69:477-485.
- Omura, H. and S. Ohsumi. 1964. A review of Japanese whale marking in the North Pacific to the end of 1962, with some information on marking in the Antarctic. *Norsk Hvalfangst-Tidende* 53:90-112.
- Omura, H. and S. Ohsumi. 1974. Research on whale biology of Japan with special reference to the North Pacific stocks. Pages 196-208 in W.E. Schevill ed. *The Whale Problem*. Harvard Univ. Press, Cambridge, Massachusetts.
- Palma, S. and N. Silva. 2004. Distribution of siphonophores, chaetognaths, euphausiids and oceanographic conditions in the fjords and channels of southern Chile. *Deep-Sea Research II* 51:513-535.
- Palsbøll, P.J., J. Allen, M. Bérubé, P.J. Clapham, T.P. Feddersen, R.R. Hudson, H. Jørgensen, S. Katona, A.H. Larsen, F. Larsen, J. Lien, D.K. Mattila, J. Sigurjónsson, R. Sears, T. Smith, R. Sponer, P. Stevick and N. Øien. 1997. Genetic tagging of humpback whales. *Nature* 388:767-769.
- Pardo, M.A., T. Gerrodette, E. Beier, D. Gendron, K.A. Forney, S.J. Chivers, J. Barlow and D.M. Palacios. 2015. Inferring cetacean population densities from the absolute dynamic topography of the ocean in a hierarchical bayesian framework. *PLoS one* 10:e0120727.
- Parkinson, C.L. and D.J. Cavalieri. 2012. Antarctic sea ice variability and trends, 1979–2010. *The Cryosphere* 6:871-880.

- Parks, S., D.R. Ketten, J.T. O'Malley, and J. Arruda. 2007. Anatomical Predictions of Hearing in the North Atlantic Right Whale. *The Anatomical Record* 290:734-744.
- Parks, S.E., I. Urazghildiiev and C.W. Clark. 2009. Variability in ambient noise levels and call parameters of North Atlantic right whales in three habitat areas. *Journal of the Acoustical Society of America* 125:1230-1239.
- Pérez-Jorge, S., T. Pereira, C. Corne, Z. Wijtten, M. Omar, J. Katello, M. Kinyua, D. Oro and M. Louzao. Can Static Habitat Protection Encompass Critical Areas for Highly Mobile Marine Top Predators? Insights from Coastal East Africa. *PLoS one* 10:e0133265.
- Perrin, W.F., J.G. Mead and R.L. Brownell, Jr. 2009. Review of the evidence used in the description of currently recognized cetacean subspecies. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-SWFSC-450. 41 pp.
- Pervushin, A.S. 1968. Observations of behavior and feeding of whale-bone whales in the area of the Crozet Islands. *Oceanology* 8:110-115.
- Pike, D.G., G.A. Vikingsson and T. Gunnlaugsson. 2004. Abundance estimates for blue whales (*Balaenoptera musculus*) in Icelandic and adjacent waters. International Whaling Commission Scientific Committee. 10 pp.
- Pike, D.G., G.A. Vikingsson, T. Gunnlaugsson and N. Oien. 2009. A note on the distribution and abundance of blue whales (*Balaenoptera musculus*) in the central and northeast North Atlantic. Pages 19-29 in C. Lockyer and D. Pike eds. *North Atlantic Sightings Surveys: Counting Whales in the North Atlantic, 1987-2006*.
- Pitman, R.L., H. Fearnbach, R. Leduc, J.W. Gilpatrick, Jr., J.K.B. Ford and L.T. Ballance. 2007. Killer whales preying on a blue whale calf on the Costa Rica Dome: Genetics, morphometrics, vocalisations and composition of the group. *Journal of Cetacean Research And Management* 9:151-157.
- Pivorunas, A. 1979. The feeding mechanisms of baleen whales. *American Scientist* 67:432-440.
- Polefka, S. 2004. Anthropogenic noise and the Channel Islands National Marine Sanctuary: How noise affects sanctuary resources, and what we can do about it. Environmental Defense Center. 53 pp.
- Popper, A.N. 1980a. Behavioral measures of odontocete hearing. Pages 469-481 in R.-G. Busnel and J.F. Fish eds. *Animal Sonar Systems*. Plenum Press, New York.
- Popper, A.N. 1980b. Sound emission and detection by delphinids. Pages 1-52 in L.M. Herman ed. *Cetacean Behavior: Mechanisms and Functions*. John Wiley and Sons, New York.
- Priyadarshana, T., S.M. Randage, A. Alling, S. Calderan, J. Gordon, T. Gordon, R. Leaper, T. Lewis, L. Porter and M.V. Thillo. 2015. An update on work related to ship strike risk to blue whales off southern Sri Lanka. International Whaling Commission Scientific Committee. 15 pp.
- Priyadarshana, T., S.M. Randage, A. Alling, S. Calderan, J. Gordon, R. Leaper and L. Porter. 2016. Distribution patterns of blue whale (*Balaenoptera musculus*) and shipping off southern Sri Lanka. *Regional Studies in Marine Science* 3:181-188.
- Quinn II, T.J. and H.J. Niebauer. 1995. Relation of eastern Bering Sea walleye Pollock (*Theragra chalcogramma*) recruitment to environmental and oceanographic variables. Pages 739 in R.J. Beamish ed. *Climate change and northern fish populations*. Canadian Special Publication of Fisheries and Aquatic Sciences.
- Radford, A.N., E. Kerridge and S.D. Simpson. 2014. Acoustic communication in a noisy world: Can fish compete with anthropogenic noise? *Behavioral Ecology* 25:1022-1030.
- Ralls, K. 1976. Mammals in which females are larger than males. *Quarterly Review of Biology*

51:245-276.

- Ramp, C., M. Berube, W. Hagen and R. Sears. 2006. Survival of adult blue whales *Balaenoptera musculus* in the Gulf of St. Lawrence, Canada. *Marine Ecology Progress Series* 319:287-295.
- Ramp, C. and R. Sears. 2013. Distribution, densities, and annual occurrence of individual blue whales (*Balaenoptera musculus*) in the Gulf of St. Lawrence, Canada from 1980-2008. Fisheries and Oceans Canada. 44 pp.
- Randage, S.M., A. Alling, K. Currier and E. Heywood. 2014. Review of the Sri Lanka blue whale (*Balaenoptera musculus*) with observations on its distribution in the shipping lane. *Journal of Cetacean Research And Management* 14:43-49.
- Rankin, S., D. Ljungblad, C. Clark and H. Kato. 2005. Vocalisations of Antarctic blue whales, *Balaenoptera musculus intermedia*, recorded during the 2001/2002 and 2002/2003 IWC/SOWER circumpolar cruises, Area V, Antarctica. *Journal of Cetacean Research And Management* 7:13-20.
- Redfern, J.V., L.T. Hatch, C. Caldw, M.L. DeAngelis, J. Gedamke, S. Hastings, L. Henderson, M.F. McKenna, T.J. Moore and M.B. Porter. 2017. Assessing the risk of chronic shipping noise to baleen whales off Southern California, USA *Endangered Species Research* 32:153-167.
- Reeves, R.R. and M.F. Barto. 1985. Whaling in the Bay of Fundy. *Whalewatcher* 19:14-18.
- Reeves, R.R., P.J. Clapham, R.L. Brownell, Jr. and G.K. Silber. 1998. Recovery Plan for the blue whale (*Balaenoptera musculus*). National Marine Fisheries Service, Office of Protected Resources. 42 pp.
- Reeves, R.R., S. Leatherwood, S.A. Karl and E.R. Yohe. 1985. Whaling results at Akutan (1912-39) and Port Hobron (1926-37), Alaska. Report of the International Whaling Commission 35:441-457.
- Reeves, R.R., T.D. Smith, E.A. Josephson, P.J. Clapham and G. Woolmer. 2004. Historical observations of humpback and blue whales in the North Atlantic Ocean: Clues to migratory routes and possibly additional feeding grounds. *Marine Mammal Science* 20:774-786.
- Reeves, R.R., B.S. Stewart, P.J. Clapham, J.A. Powell and P.F. (illustrator). 2002. Guide to marine mammals of the world. A. A. Knopf, New York.
- Reichmuth, C. 2007. Assessing the hearing capabilities of mysticete whales: A proposed research strategy for the Joint Industry Programme on Sound and Marine Life. E&P Sound & Marine Life Programme. 35 pp.
- Reijnders, P.J.H., G.P. Donovan, A. Aguilar and A. Bjorge. 1999. Report of the workshop on chemical pollution and cetaceans. *Journal of Cetacean Research And Management Special Issue* 1:1-42.
- Reilly, S.B., J.L. Bannister, P.B. Best, M. Brown, R.L. Brownell, Jr., D.S. Butterworth, P.J. Clapham, J. Cooke, G.P. Donovan, J. Urban and A.N. Zerbini. 2008. *Balaenoptera musculus*. IUCN Red List of Threatened Species. International Union for the Conservation of Nature.
- Reilly, S.B. and V.G. Thayer. 1990. Blue whale (*Balaenoptera musculus*) distribution in the eastern tropical Pacific. *Marine Mammal Science* 6:265-277.
- Rennie, S., C.E. Hanson, R.D. McCauley, C. Pattiaratchi, C. Burton, J. Bannister, C. Jenner and M.-N. Jenner. 2009. Physical properties and processes in the Perth Canyon, Western Australia: Links to water column production and seasonal pygmy blue whale abundance. *Journal of Marine Systems* 173:21-44.
- Rice, D.W. 1974. Whales and whale research in the eastern North Pacific. Pages 170-195 in W.E. Schevill ed. *The Whale Problem*. Harvard University Press, Cambridge, Massachusetts.

- Rice, D.W. 1977. A list of the marine mammals of the world. U.S. Department of Commerce, NOAA Technical Report, NMFS SSRF-711. 15 pp.
- Rice, D.W. 1986. Blue whale. *in* D. Haley ed. *Marine mammals of eastern North Pacific and Arctic waters*. Pacific Search Press.
- Richardson, W.J., C.R. Greene, Jr., C.I. Malme and D.H. Thomson. 1995. *Marine Mammals and Noise*. Academic Press, San Diego, California.
- Ridgway, S.H. 1983. Dolphin hearing and sound production in health and illness. Pages 247-296 *in* R.R. Fay and G. Gourevitch eds. *Hearing and Other Senses: Presentations in Honor of E. G. Wever* Amphora Press, Groton, CT.
- Risting, S. 1928. Whales and whale foetuses: statistics of catch and measurements collected from the Norwegian Whalers' Association 1922-1925. *Rapports et Proces-Verbaux Des Reunions* 50:1-122.
- Rockwood, R.C., J. Calambokidis and J. Jahncke. 2017. High mortality of blue, humpback and fin whales from modeling of vessel collisions on the U.S. West Coast suggests population impacts and insufficient protection. *PLoS one* 12:e0183052.
- Roemmich, D. and J. McGowan. 1995. Climate warming and the decline of zooplankton in the California Current. *Science* 267:1324-1326.
- Rørvik, C.J. and Å. Jonsgård. 1981. Review of balaenopterids in the North Atlantic Ocean. Pages 269-286 *Mammals in the Seas. Volume III. General Papers and Large Cetaceans. FAO Fisheries Series No. 5*. Food and Agriculture Organization of the United Nations, Rome.
- Ross, D. 1976. *Mechanics of Underwater Noise*. Pergamon Press, New York.
- Saba, G.K., O. Schofield, J.J. Torres, E.H. Ombres and D.K. Steinberg. 2012. Increased Feeding and Nutrient Excretion of Adult Antarctic Krill, *Euphausia superba*, Exposed to Enhanced Carbon Dioxide (CO₂). *PLoS one* 7:e52224.
- Samaran, F., O. Adam and C. Guinet. 2010. Discovery of a mid-latitude sympatric area for two Southern Hemisphere blue whale subspecies. *Endangered Species Research* 12:157-165.
- Samaran, F., K.M. Stafford, T.A. Branch, J. Gedamke, J.-Y. Royer, R.P. Dziak and C. Guinet. 2013. Seasonal and geographic variation of southern blue whale subspecies in the Indian Ocean. *PLoS one* 8:e71561.
- Santora, J.A., C.S. Reiss, V.J. Loeb and R.R. Veit. 2010. Spatial association between hotspots of baleen whales and demographic patterns of Antarctic krill *Euphausia superba* suggests size-dependent predation. *Marine Ecology Progress Series* 405:255-269.
- Savenkoff, C., M. Castonguay, D. Chabot, M.O. Hammill, H. Bourdages and L. Morissette. 2007. Changes in the northern Gulf of St. Lawrence ecosystem estimated by inverse modelling: Evidence of a fishery-induced regime shift? *Estuarine, Coastal and Shelf Science* 73:711-724.
- Schick, R.S., S. Kraus, R.M. Rolland, A.R. Knowlton, P.K. Hamilton, H.M. Pettis, R.D. Kenney and J.S. Clark. 2013. Using Hierarchical Bayes to Understand Movement, Health, and Survival in the Endangered North Atlantic Right Whale. *PLoS one* 8:e64166.
- Schmitt, F.P., C. de Jong and F.H. Winter. 1980. *Thomas Welcome Roys: America's Pioneer of Modern Whaling*. University of Virginia Press, Newport News, Virginia.
- Schoenherr, J.R. 1991. Blue whales feeding on high concentrations of euphausiids around Monterey Submarine Canyon. *Canadian Journal of Zoology* 69:583-594.
- Schusterman, R.J. 1981. Behavioral capabilities of seals and sea lions: A review of their hearing, visual, learning, and diving skills. *Psychological Record* 31:125-143.
- Schwacke, L.H., C.R. Smith, F.I. Townsend, R.S. Wells, L.B. Hart, B.C. Balmer, Tracy K.

- Collier, S.D. Guise, M.M. Fry, J. Louis J. Guillette, S.V. Lamb, S.M. Lane, W.E. McFee, N.J. Place, M.C. Tumlin, G.M. Ylitalo, E.S. Zolman and T.K. Rowles. 2014. Health of Common Bottlenose Dolphins (*Tursiops truncatus*) in Barataria Bay, Louisiana, Following the Deepwater Horizon Oil Spill. *Environmental Science & Technology* 48:93-103.
- Sears, R. 1990. The Cortez blues. *Whalewatcher* 24:12-15.
- Sears, R., C.L.K. Buton and G. Vikingson. 2005. Review of blue whale (*Balaenoptera musculus*) photoidentification distribution data in the North Atlantic, including the first long-range match between Iceland and Mauritania. Pages 254 Sixteenth Biennial Conference on the Biology of Marine Mammals. San Diego, California.
- Sears, R. and J. Calambokidis. 2002. Update COSEWIC status report on the blue whale *Balaenoptera musculus* (Atlantic population, Pacific population) in Canada. 1-32.
- Sears, R., F.W. Wenzel, and J.M. Williamson. 1987. The Blue Whale: A Catalogue of Individuals from the Western North Atlantic (Gulf of St. Lawrence). Mingan Island Cetacean Study, St. Lambert, Quebec. 27 pp.
- Sears, R. and F. Larsen. 2002. Long range movements of a blue whale (*Balaenoptera musculus*) between the Gulf of St. Lawrence and West Greenland. *Marine Mammal Science* 18:281-285.
- Sears, R. and W.F. Perrin. 2009. Blue whale, *Balaenoptera musculus*. Pages 120-124 in W.F. Perrin, B. Wursig and J.G.M. Thewissen eds. *Encyclopedia of Marine Mammals, Second Edition*. Academic Press, San Diego, California.
- Sears, R., C. Ramp, A.B. Douglas and J. Calambokidis. 2013. Reproductive parameters of eastern North Pacific blue whales *Balaenoptera musculus*. *Endangered Species Research* 22:23-31.
- Sears, R., J.M. Williamson, F.W. Wenzel, M. Berube, D. Gendron and P. Jones. 1990. Photographic identification of the blue whale (*Balaenoptera musculus*) in the Gulf of St. Lawrence, Canada. Report of the International Whaling Commission Special Issue 12:335-342.
- Sellner, K.G., G.J. Doucette and G.J. Kirkpatrick. 2003. Harmful algal blooms: causes, impacts and detection. *Journal of Industrial Microbiology and Biotechnology* 30:383-406.
- Sergeant, D.E. 1953. Whaling in Newfoundland and Labrador waters. *Norsk Hvalfangst-Tidende* 42:687-695.-P.439-447 in Some Vol.).
- Sergeant, D.E. 1966. Populations of large whale species in the western North Atlantic with special reference to the fin whale. Fisheries Research Board of Canada, Arctic Biological Station Circular 9.
- Sergeant, D.E. 1969. Feeding rates of Cetacea. *Fiskeridir. Skr. Havundersok* 15.
- Sergeant, D.E. 1982. Some biological correlates of environmental conditions around Newfoundland during 1970-79: Harp seals, blue whales and fulmar petrels. *Nafo Scientific Council Studies* No.5:107-110.
- Siegel, V. 2005. Distribution and population dynamics of *Euphausia superba*: summary of recent findings. *Polar Biology* 29:1-22.
- Sigurjónsson, J. and T. Gunnlaugsson. 1990. Recent trends in abundance of blue (*Balaenoptera musculus*) and humpback whales (*Megaptera novaeangliae*) off West and Southwest Iceland, with a note on occurrence of other cetacean species. Report of the International Whaling Commission 40:537-551.
- Silber, G., J. Slutsky and S. Bettridge. 2010. Hydrodynamics of a ship/whale collision. *Journal of Experimental Marine Biology and Ecology* 391:10-19.
- Silber, G.K., J.D. Adams, M.J. Asaro, T. Cole, K.S. Moore, L.I. Ward and B.J. Zoodsm. 2015.

- The right whale Mandatory Ship Reporting System: a retrospective. PeerJ 3:e866.
- Silber, G.K., S. Bettridge and D. Cottingham. 2009. Report of a workshop to identify and assess technologies to reduce ship strikes of large whales; Providence, Rhode Island 8-10 July 2008. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-OPR-42. 66 pp.
- Silber, G.K., M. Lettrich and P.O. Thomas. 2016. Report of a Workshop on Best Approaches and Needs for Projecting Marine Mammal Distributions in a Changing Climate. 12-14 January 2016, Santa Cruz, California, USA. U.S. Department of Commerce, NOAA Technical Memorandum, NMFS-OPR-54. 50 pp.
- Simard, Y., R. de Ladurantaye and J.-C. Therriault. 1986. Aggregation of euphasiids along a coastal shelf in an upwelling environment Marine Ecology Progress Series 32:203-215.
- Simmonds, M.P. and J.D. Hutchinson. 1996. The Conservation of Whales and Dolphins - Science and Practice. John Wiley & Sons.
- Širović, A. 2006. Blue and fin whale acoustics and ecology off Antarctic Peninsula. Ph.D., University of California, San Diego, California 163 pp.
- Širović, A. and J.A. Hildebrand. 2011. Using passive acoustics to model blue whale habitat off the Western Antarctic Peninsula. Deep Sea Research Part II: Topical Studies in Oceanography 58:1719-1728.
- Širović, A., J.A. Hildebrand, S.M. Wiggins, M.A. McDonald, S.E. Moore and D. Thiele. 2004. Seasonality of blue and fin whale calls and the influence of sea ice in the Western Antarctic Peninsula. Deep Sea Research Part II: Topical Studies in Oceanography 51:2327-2344.
- Širović, A., J.A. Hildebrand, S.M. Wiggins and D. Thiele. 2009. Blue and fin whale acoustic presence around Antarctica during 2003 and 2004. Marine Mammal Science 25:125-136.
- Sleptsov, M.M. 1955. Biology of whales and the whaling fishery in Far Eastern seas. 'Pishch. Prom.', Moscow. [In Russian.] (Transl. with comments and conclusions only by Fish. Res. Board Can., Transl. Ser. 118, 6 pp.).
- Smith, C.E., S.T. Sykora-Bodie, B. Bloodworth, S.M. Pack, T.R. Spradlin and N.R. LeBoeuf. 2016. Assessment of known impacts of unmanned aerial systems (UAS) on marine mammals: data gaps and recommendations for researchers in the United States. Journal Unmanned Vehicle Systems 4:1-14.
- Smith, T.D., J. Allen, P.J. Clapham, P.S. Hammond, S. Katona, F. Larsen, J. Lien, D. Mattila, P.J. Palsböll, J. Sigurjónsson, P.T. Stevick and N. Øien. 1999. An ocean-basin-wide mark-recapture study of the North Atlantic humpback whale (*Megaptera novaeangliae*). Marine Mammal Science 15:11689.
- Southall, B.L., A.E. Bowles, W.T. Ellison, J.J. Finneran, R.L. Gentry, C.R. Greene, D. Kastak, D.R. Ketten, J.H. Miller, P.E. Nachtigall, W.J. Richardson, J.A. Thomas and P.L. Tyack. 2007. Marine mammal noise and exposure criteria: initial scientific recommendations. Aquatic Mammals 33:411-521.
- Southall, B.L., D. Moretti, B. Abraham, J. Calambokidis, S.L. DeRuiter and P.L. Tyack. 2012. Marine Mammal Behavioral Response Studies in Southern California: Advances in Technology and Experimental Methods. Marine Technology Society Journal 46:46-59.
- Southall, B.L., D.P. Nowacek, P.J.O. Miller and P.L. Tyack. 2016. Experimental field studies to measure behavioral responses of cetaceans to sonar. Endangered Species Research 31:293-315.
- Southall, B.L., A.R. Scholik-Schlomer, L. Hatch, T. Bergmann, M. Jasny, K. Metcalf, L. Weilgart and A.J. Wright. 2017. Underwater Noise from Large Commercial Ships - International Collaboration for Noise Reduction. Encyclopedia of Maritime and Offshore

- Engineering 10.1002/9781118476406.emoe056:1-9.
- Sremba, A., D. Steel, L. Torres, R. Constantine, N. Bott and C.S. Baker. 2015. Genetic identity of blue whales in the surrounding waters of New Zealand. International Whaling Commission Scientific Committee. 9 pp.
- Sremba, A.L., B. Hancock-Hanser, T.A. Branch, R.L. Leduc and C.S. Baker. 2012. Circumpolar diversity and geographic differentiation of mtDNA in the critically endangered Antarctic blue whale (*Balaenoptera musculus intermedia*). PLoS one 7:e32579.
- Stafford, K.M. 2003. Two types of blue whale calls recorded in the Gulf of Alaska. Marine Mammal Science 19:12.
- Stafford, K.M., D.R. Bohnenstiehl, M. Tolstoy, E. Chapp, D.K. Mellinger and S.E. Moore. 2004. Antarctic-type blue whale calls recorded at low latitudes in the Indian and eastern Pacific Oceans. Deep Sea Research Part I: Oceanographic Research Papers 51:1337-1346.
- Stafford, K.M., E. Chapp, D.R. Bohnenstiel and M. Tolstoy. 2011. Seasonal detection of three types of pygmy blue whale calls in the Indian Ocean. Marine Mammal Science 27:828-840.
- Stafford, K.M., J.J. Citta, S.E. Moore, M.A. Daher and J.E. George. 2009. Environmental correlates of blue and fin whale call detections in the North Pacific Ocean from 1997 to 2002. Marine Ecology Progress Series 395:37-53.
- Stafford, K.M., C.G. Fox and D.S. Clark. 1998. Long-range acoustic detection and localization of blue whale calls in the northeast Pacific Ocean. Journal of the Acoustical Society of America 104:3616-3625.
- Stafford, K.M., D.K. Mellinger, S.E. Moore and C.G. Fox. 2007. Seasonal variability and detection range modeling of baleen whale calls in the Gulf of Alaska, 1999-2002. Journal of the Acoustical Society of America 122:3378-3390.
- Stafford, K.M., S.E. Moore and C.G. Fox. 2005. Diel variation in blue whale calls recorded in the eastern tropical Pacific. Animal Behaviour 69:951-958.
- Stafford, K.M., S.L. Nieukirk and C.G. Fox. 1999a. An acoustic link between blue whales in the eastern tropical Pacific and the northeast Pacific. Marine Mammal Science 15:1258-1268.
- Stafford, K.M., S.L. Nieukirk and C.G. Fox. 1999b. Low-frequency whale sounds recorded on hydrophones moored in the eastern tropical Pacific. Journal of the Acoustical Society of America 106:3687-3698.
- Stafford, K.M., S.L. Nieukirk and C.G. Fox. 2001. Geographic and seasonal variation of blue whale calls in the North Pacific. Journal of Cetacean Research And Management 3:65-76.
- Stewart, B.S., S.A. Karl, P.K. Yochem, S. Leatherwood and J.L. Laake. 1987. Aerial surveys for cetaceans in the former Akutan, Alaska, whaling grounds. Arctic 40:33-42.
- Stone, C.J. 2003. The effects of seismic activity on marine mammals in UK waters, 1998-2000. Joint Nature Conservation Committee, JNCC Report No. 323. 78 pp.
- Sutcliffe, W.H., Jr., and P.F. Brodie. 1977. Whale distributions in Nova Scotia waters. Fisheries and Marine Service (Canada) Technical Report. 722. 1-89 pp.
- Tarpley, R.J. and S. Marwitz. 1993. Plastic debris ingestion by cetaceans along the Texas coast: Two case reports. Aquatic Mammals 19:93-98.
- Tarpy, C. 1979. Killer whale attack! National Geographic Magazine 155:542-545.
- Taylor, B.L., S.J. Chivers, J. Larese and W.F. Perrin. 2007. Generation length and percent mature estimates for IUCN assessments of cetaceans. National Marine Fisheries Service, Southwest Fisheries Science Center Administrative Report, LJ-07-01. 24 pp.
- Tershy, B.R., D. Breese and C.S. Strong. 1990. Abundance, seasonal distribution and population composition of balaenopterid whales in the Canal de Ballenas, Gulf of California, Mexico.

- Report of the International Whaling Commission Special Issue 12:369-375.
- Thode, A.M., G.L. D'Spain and W.A. Kuperman. 2000. Matched-field processing, geoacoustic inversion, and source signature recovery of blue whale vocalizations. *Journal of the Acoustical Society of America* 107:1286-1300.
- Thomas, P.O., R.R. Reeves and R.L. Brownell, Jr. 2016. Status of the world's baleen whales. *Marine Mammal Science* 32:682-734.
- Thomisch, K., O. Boebel, C.W. Clark, W. Hagen, S. Spiesecke, D.P. PZitterbart and I. Van Opzeeland. 2016. Spatio-temporal patterns in acoustic presence and distribution of Antarctic blue whales *Balaenoptera musculus intermedia* in the Weddell Sea. *Endangered Species Research* 30:239-253.
- Thompson, D.W. 1928. On whales landed at the Scottish whaling stations during the years 1908-1914 and 1920-1927. Fishery Board for Scotland, Scientific Investigations 1928, No. III. pp.
- Thompson, P.O., L.T. Findley, O. Vidal and W.C. Cummings. 1996. Underwater sounds of blue whales, *Balaenoptera musculus*, in the Gulf of California, Mexico. *Marine Mammal Science* 12:288-293.
- Thompson, P.O. and W.A. Friedl. 1982. A long term study of low frequency sounds from several species of whales off Oahu, Hawaii. *Cetology* 45:1-19.
- Tønnessen, J.N. and A.O. Johnsen. 1982. *The History of Modern Whaling*. University of California Press.
- Torres-Florez, J.P., P.A. Olson, L. Bedrinana-Romano, H. Rosenbaum, J. Ruiz, R. Leduc and R. Huck-Gaete. 2015. First documented migratory destination for eastern South Pacific blue whales. *Marine Mammal Science* 31:1580-1586.
- Torres, L., P. Gill, R. Hamner and D. Glasgow. 2014. Documentation of a blue whale foraging ground in the South Taranaki Bight, New Zealand. *International Whaling Commission Scientific Committee*. 10 pp.
- Torres, L.G. 2013. Evidence for an unrecognised blue whale foraging ground in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 47:235-248.
- True, F.W. 1904. The whalebone whales of the western North Atlantic compared with those occurring in European waters; with some observations on the species of the North Pacific. *Smithsonian Contributions to Knowledge* 33:1-318.
- Trumble, S.J., E.M. Robinson, M. Berman-Kowalewski, C.W. Potter and S. Usenko. 2013. Blue whale earplug reveals lifetime contaminant exposure and hormone profiles. *Proceedings of the National Academy of Sciences* 110:16922-16926.
- Tyack, P.L. 2008. Implications for marine mammals of large-scale changes in the marine acoustic environment. *Journal of Mammalogy* 89:549-558.
- U.S. Department of the Navy. 2008. Draft Environmental Impact Statement/Overseas Environmental Impact Statement: Southern California Range Complex. Vols. 1 and 2.
- USCG. 2011. Port Access Route Study: Approaches to Los Angeles-Long Beach and in the Santa Barbara Channel. 33 pp.
- Van Bresseem, M.-F., P.J. Duignan, A. Banyard, M. Barbieri, K.M. Colegrove, S.D. Guise, G.D. Guardo, A. Dobson, M. Domingo, D. Fauquier, A. Fernandez, T. Goldstein, B. Grenfell, K.R. Groch, F. Gulland, B.A. Jensen, P.D. Jepson, A. Hall, T. Kuiken, S. Mazzariol, S.E. Morris, O. Nielsen, J.A. Raga, T.K. Rowles, J. Saliki, E. Sierra, N. Stephens, B. Stone, I. Tomo, J. Wang, T. Waltzek and J.F. Wellehan. 2014. Cetacean Morbillivirus: Current Knowledge and Future Directions. *Viruses* 6:5145-5181.
- Van Dolah, F.M. 2000. *Marine Algal Toxins: Origins, Health Effects, and Their Increased*

- Occurrence. *Environmental Health Perspectives* 108:133-141.
- Vanderlaan, A.S.M. and C.T. Taggart. 2007. Vessel collisions with whales: the probability of lethal injury based on vessel speed. *Marine Mammal Science* 23:144-156.
- Viale, D., N. Verneau and Y. Tison. 1991. Occlusion gastrique jutale chez un cachalot echoue sur les Iles Lavezzi: Macropollution en Mediterranee. *Journal de Recherche Oceanographique* 16:100-102.
- Vieira, S.M., R. de Almeida, I.B.B. Holanda, M.H. Mussu, R.C.F. Galvão, P.T.B. Crispim, J.G. Dórea and W.R. Bastos. 2013. Total and methyl-mercury in hair and milk of mothers living in the city of Porto Velho and in villages along the Rio Madeira, Amazon, Brazil. *International Journal of Hygiene and Environmental Health* 216:682-689.
- Visser, F., K.L. Hartman, G.J. Pierce, V.D. Valavanis and J. Huisman. 2011. Timing of migratory baleen whales at the Azores in relation to the North Atlantic spring bloom. *Marine Ecology Progress Series* 440:267-279.
- Wada, S. 1975. Indices of abundance of large-sized whales in the North Pacific in 1973 whaling season. Pages 129-165 Report of the International Whaling Commission. London.
- Wade, L.S. and G.L. Friedrichsen. 1978. Recent sightings of the blue whale, *Balaenoptera musculus*, in the northeastern tropical Pacific. *Fishery Bulletin* 76:915-919.
- Wade, P.R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. *Reports of the International Whaling Commission* 43:477-493.
- Wagemann, R. and D.C.G. Muir. 1984. Concentrations of heavy metals and organochlorines in marine mammals of northern waters: Overview and evaluation. *Canadian Technical Report of Fisheries and Aquatic Sciences* 1279:vi + 97.
- Walther, G.-R., E. Post, P. Convey, A. Menzel, C. Parmesan, T.J.C. Beebee, J.-M. Fromentin, O. Hoegh-Guldberg and F. Bairlein. 2002. Ecological responses to recent climate change. *Nature* 416:389-395.
- Waring, G.T., E. Josephson, K. Maze-Foley and P.E. Rosel. 2010. US Atlantic and Gulf of Mexico marine mammal stock assessments-2010. U.S. Department of Commerce, NOAA Technical Memorandum. 606 pp.
- Wartzok, D. and D.R. Ketten. 1999. Marine mammal sensory systems. Pages 117-175 in J.E.R. Iii and S.A. Rommel eds. *Biology of Marine Mammals*. Smithsonian Institution Press, Washington.
- Watkins, W.A. 1986. Whale reactions to human activities in Cape Cod waters. *Marine Mammal Science* 2:251-262.
- Watkins, W.A., J.E. George, M.A. Daher, K. Mullin, D.L. Martin, S.H. Haga and N.A. Dimarzio. 2000. Whale call data for the North Pacific November 1995 through July 1999 occurrence of calling whales and source locations from SOSUS and other acoustic systems. Woods Hole Oceanographic Institution. Woods Hole Oceanographic Institution Technical Report. 161 pp.
- Watkins, W.A. and D. Wartzok. 1985. Sensory biophysics of marine mammals. *Marine Mammal Science* 1:219-260.
- Weisbrod, A.V., D. Shea, M.J. Moore and J.J. Stegeman. 2000. Organochlorine exposure and bioaccumulation in the endangered northwest Atlantic right whale (*Eubalaena glacialis*) population. *Environmental Toxicology and Chemistry* 19:654.
- Wenzel, F.W., D.K. Mattila and P.J. Clapham. 1988. *Balaenoptera musculus* in the Gulf of Maine. *Marine Mammal Science* 4:172-175.
- Williams, R., S.L. Hedley, T.A. Branch, M.V. Bravington, A.N. Zerbini and K.P. Findlay. 2011.

- Chilean blue whales as a case study to illustrate methods to estimate abundance and evaluate conservation status of rare species. *Conservation Biology* 25:526-535.
- Williams, R., D. Lusseau and P.S. Hammond. 2006. Estimating relative energetic costs of human disturbance to killer whales (*Orcinus orca*). *Biological Conservation* 133:301-311.
- Wintle, B.A., M.A. McCarthy, C.T. Volinsky and R.P. Kavanagh. 2003. The Use of Bayesian Model Averaging to Better Represent Uncertainty in Ecological Models. *Conservation Biology* 17:1579-1590.
- Yablokov, A.V. 1994. Validity of whaling data. *Nature* 367.
- Yamato, M., D.R. Ketten, J. Arruda and S. Cramer. 2008. Biomechanical and structural modeling of hearing in baleen whales. *Bioacoustics* 17-Jan:100-102. Special Issue on the International Conference on the Effects of Noise on Aquatic Life. Edited By A. Hawkins, A. N. Popper & M. Wahlberg.
- Yochem, P.K. and S. Leatherwood. 1985. Blue whale *Balaenoptera musculus* (Linnaeus, 1758). Pages 193-240 in S.H. Ridgway and R. Harrison eds. *Handbook of Marine Mammals*. Academic Press, London.
- Yost, W.A. 2000. *Fundamentals of Hearing: An Introduction*. Academic Press, New York.
- Zemsky, V.A. and E.G. Sazhinov. 1982. Distribution and abundance of the pygmy blue whale. Pages 20 in M.A. Donahue and R.L. Brownell, Jr. eds. *National Marine Fisheries Service, Southwest Fisheries Science Center Administrative Report*.