



[Plants of South Florida](#) · [Plants by Conservation Area](#) · [Plants by County](#) · [Plants by Habitat](#)
[Submit Data](#) · [Quick Search](#) · [Advanced Search](#)

Trichomanes punctatum Poir. subsp. *floridanum*
W. Boer
Florida bristle fern

Family: **Hymenophyllaceae**

Group: **Pteridophyte**

Native Status: **Native**

Cultivated Status: **Not Cultivated**

Perennation: **Perennial**

Substrate: **Lithophyte**

Habit: **Herb**

IRC Status: **Critically Imperiled**

State Status: **Endangered**

FNAI State Status: **Critically Imperiled**

FNAI Global Status: **G5T1**

Native Range: **Endemic to peninsular Florida.**



Copyright by: Gil Nelson

Other data on *Trichomanes punctatum* subsp. *floridanum* available from :



[IRC Species Account](#)



Trichomanes punctatum subsp. *floridanum* has been found in the following 8 conservation areas :

	Occurrence	Native Status
Castellow Hammock Park	Present	Native
Deering Estate at Cutler	Present	Native
Everglades National Park	Presumed Extirpated	Presumed Extirpated
Fuchs Hammock Preserve	Present	Native
Matheson Hammock Park	Present	Native
Meissner Hammock	Present	Native
R. Hardy Matheson Preserve	Presumed Extirpated	Presumed Extirpated
Silver Palm Hammock	Present	Native

Trichomanes punctatum subsp. *floridanum* has been found in the following county :

	Occurrence	Native Status
Miami-Dade County		Native

Trichomanes punctatum subsp. *floridanum* has been found in the following 2 habitats :

[Rockland Hammock](#)
[Sinkhole](#)

Gann, G.D., K.A. Bradley and S.W. Woodmansee. 2001-2008.
The Floristic Inventory of South Florida Database Online.
The Institute for Regional Conservation, Miami.