



Species Diversity of Lepidoptera in Western Mindanao State University – Experimental Forest Area, Zamboanga City, Philippines

Clarice Margrethe D. Sebu^a*, Olga M. Nuñez^a

Department of Biological Sciences, Mindanao State University – Iligan Institute of Technology, Andres Bonifacio Avenue, Tibanga, Iligan City, Philippines.

ABSTRACT

Moths and butterflies belonging to Lepidoptera are considered biological indicators of human disturbances in tropical forests. This study aimed to determine the species diversity of Lepidoptera in Western Mindanao State University – Experimental Forest Area, Zamboanga City. Six sites were sampled for 126 person-hours. An opportunistic sampling method using sweep nets was employed. PAST software version 3.0 was used to determine biodiversity indices. Thirty-nine species consisting of 23 species of butterflies and 16 species of moth from eight families were recorded. The family Nymphalidae was dominant, most abundant, and had the highest species richness mainly due to its polyphagous nature. *Eurema hecabe tamiathis* was the most distributed and most abundant species (13.57%) which means that it can thrive in different types of habitats. *Idea electra* was the only Lepidoptera species categorized as vulnerable. Sampling site 4, a secondary dipterocarp forest, was the most diverse ($H'=2.993$), most abundant (30.00%), and with the highest species richness ($S=23$). The results showed that WMSU-EFA had a relatively moderate diversity ($H'=2.2625$) attributed to its diverse vegetation. There was no dominant species since the distribution in all sampling sites was even. Threat observed that can affect the Lepidoptera diversity of the sampling area was severe anthropogenic clearing.

Keywords: *Butterflies, Dipterocarp, Indicators, Moth, Nymphalidae.*

HOW TO CITE THIS ARTICLE: Clarice Margrethe D. Sebu, Olga M. Nuñez; Species Diversity of Lepidoptera in Western Mindanao State University – Experimental Forest Area, Zamboanga City, Philippines, Entomol Appl Sci Lett, 2020, 7 (1): 33-43.

Corresponding author: Clarice Margrethe D. Sebu

E-mail ✉ claricemargrethe.sebu@g.msuiit.edu.ph

Received: 28/09/2019

Accepted: 06/03/2020

INTRODUCTION

Species belonging to Lepidoptera live in interaction with many plant [1] and animal species [2]. Lepidoptera, which consists of moths and butterflies [3, 4], is a plant-feeder during its larval stage and nectar-feeder or fruit-feeder in its adult stage [3]. Both stages are almost highly dependent on vascular plants. This makes it an important herbivore and pollinator in the ecosystem [5]. Larval instars of butterflies and moths are with chewing type mouthparts [6, 7]. Also, Lepidoptera is considered a better biological indicator because of its sensitivity to environmental changes [8, 9]. It has over 157,424 known species worldwide [10] and with this number, it actually is the second largest order of Insecta, surpassed only by the beetles [11].

Nöske et al. [12] reported 282 species and 829 species of arctiid and geometrid moth species, respectively, in southern Ecuador. One hundred twenty-seven species in Kaya Muhaka, 56 species in Kaya Kinondo, and 77 butterfly species in Kaya Diani, all in Kaya forests in southern Kenya, were discovered by Lehmann and Kioko [13]. In the Atlantic Forest of Southeastern Brazil, Uehara-Prado et al. [14] recorded 70 butterfly species from six subfamilies of Nymphalidae (fruit-feeders) and concluded that forest fragmentation can affect fruit-feeding butterflies but not as severe that it can cause an obvious change on species diversity. In Asia, particularly Sabah, Borneo, a similar study, but on selective logging instead of forest fragmentation, was conducted by Dumbrell and Hill [15]. They sampled 58 species, also from Nymphalidae and concluded, based on ground-level data, that there is no

significant difference in species diversity between primary and selectively logged forests even though the primary forest is more diverse than the selectively logged one. In Poring Hill Dipterocarp Forest, also in Sabah, 1,169 macromoth species were documented by Abang and Karim [16] and found that the species diversity was very high due to a higher habitat diversity of their sampling area. Majumder et al. [17] recorded 59 butterfly species in Trishna Wildlife Sanctuary in northeast India and stated that butterflies under Nymphalidae are the dominant species in the said area. This is because they are active fliers and have polyphagous nature, enabling them to survive in different types of habitats. Kudavidanage et al. [18] reported 120 species of butterflies in Sri Lanka. As expected, tropical rainforests have very high species diversity [16]. However, research studies on Lepidoptera diversity in the tropical regions are still very few [19] regardless of all the related studies mentioned above. The Philippines, a tropical country, is also home to a diverse group of Lepidoptera. However, there are only a few biological records on Lepidoptera in Mindanao, which is the second largest island in the country [20]. Among these studies in Mindanao are those conducted in Mt.

Hamiguitan, Davao Oriental [21], Bega Watershed, Agusan del Sur [20], Tandag, Surigao del Sur [22], and Mt. Timpoong and Mt. Hibokhibok, Camiguin Island [23]. Despite these records, no lepidopteran study was conducted in Western Mindanao State University - Experimental Forest Area, the present study area.

METHODOLOGY

Study Area

Sampling was conducted in the experimental forest area of Western Mindanao State University located in Upper La Paz, Zamboanga City (Fig. 1). This is located in the southwestern part of Mindanao Island and is 26 kilometers away from the city proper. The land area of the forest is a total of 1277 hectares. The lowest point is about 600 meters above sea level (masl), located in the southwest, and the highest point is in the northern part at 1200 masl. The area is covered with diverse vegetation. The dominant tree species found are predominantly dipterocarps. There are also abundant non-tree species like rattan, vines, orchids, ground grasses, ferns, etc.




Figure 1. Map of the Philippines showing the location of Upper La Paz, Zamboanga City [24].

Sampling Sites

Site 1 at 7°02'46.0"N, 122°01'05.1"E is an agroecosystem area with an elevation of 875

masl and a moderately rugged slope. The emergent tree is white lauan (*Shorea contorta*), standing 25m with a DBH of 40cm. A small

sandy stream is present. Rattan of family Rhamnaceae and “nito” (*Lygodium circinnatum*) as well as mosses and canopy vines like “palo verde” (*Parkinsonia aculeata*), were observed. The grasses were highly dense which were composed of carabao grass (*Bouteloua dactyloides*), starflower (*Hypoxis hemerocallidea*), tiger grass (*Thysanolaena latifolia*), guinea grass (*Megathyrsus maximus*), and some ferns. Vegetation along the stream consisted of giant fern (*Angiopteris evecta*), Bengal arum (*Typhonium trilobatum*), “gabi-gabi” (*Jussiaea inclinata*), and “dilang-aso” (*Pseudelephantopus spicatus*). The ground was covered with approximately 10cm of leaf litter and 5cm of porous humus. The area had a clay type of soil which indicates soil erosion. Fallen logs and branches and even sedimentary and metamorphic exposed rocks were observed. Cultivation of pineapple (*Ananas comosus*), durian (*Durio zibethinus*), banana (*Musa sp.*), and lime (*Citrofortunella microcarpa*) was common in this site.

Site 2 at 7°01'47.3"N, 122°00'19.4"E is also an agroecosystem area with an elevation of 645 masl with a rolling slope facing east. Coconut (*Cocos nucifera*) was abundant in the site. There were no canopy epiphytes and vines. Understory plants are composed of palo verde, “gabi” (*Colocasia esculenta*), and cacao (*Gliricidia sepium*). Vegetation is composed of “dilang-aso”, water primrose (*Ludwigia adscendens*), “makahiya” (*Mimosa pudica*), “hagonoy” (*Chromolaena odorata*), giant fern, garlic (*Allium sativum*), carabao grass and some ferns. Aside from coconuts, there were also papaya (*Carica papaya*), jackfruit (*Artocarpus heterophyllus*), and bananas. The area has a clay loam type of soil. Moss, fallen logs, exposed rocks, leaf litter, and humus were absent. An intermittent stream is present 100m away from the site.

Site 3, 7°02'46.7"N, 122°00'58.0"E, is a secondary dipterocarp forest with an elevation of 990 masl. It has a rugged slope and a flowing stream. Canopy trees have lichens and canopy vines like the love vines (*Cassytha filiformis*) on their trunks. The understory plants composed of silver fern (*Pityrogramma calomelanos*), “dilang-aso”, water primrose, and fishtail palm (*Caryota mitis*). There was also pandan (*Pandanus sp.*), which was common, and rarely bananas were observed. The ground was covered with few carabao grass and mosses. There were also sed-

imentary rocks. The area has a porous and moist loamy type soil covered with approximately 10cm of humus and 20cm of leaf litter.

Site 4 at 7°01'48.0"N, 122°00'13.5"E is a secondary dipterocarp forest at an elevation of 645 masl, with rugged sloping terrain, and a flowing intermittent stream. The emergent tree was lanceleaf buttonwood (*Conocarpus lancifolius*) standing 18m with a DBH of 40cm. Canopy trees were dominated by “nibung” palm (*Oncosperma tigillarum*) and fishtail palm with a height of 8m. Orchids and canopy vines like Chinese skullcap (*Scutellaria baicalensis*) were present on the trunks of the canopy trees. Understory plants were composed of ferns like “nito”, taro (*Alocasia sp.*), “dilang-aso,” “bamban” (*Donax caniniformis*), oakleaf fern (*Drynaria quercifolia*), rattan vines, bamboo, “hagithit”, and “palmay”. Bananas were also present. No grass was present and mosses were rarely observed. Exposed sedimentary rocks and fallen branches can also be seen. The soil was of a porous loamy type covered with approximately 10cm of leaf litter and thick humus. Tree nursery can be found in the site and anthropogenic clearing was present 25m away.

Site 5 at 7°03'20.0"N 122°00'04.0"E is a forest fragment and a highly disturbed dipterocarp forest due to logging for road construction. It has an elevation of 1019m, a rolling slope and a clear, flowing stream with a small waterfall. The emergent tree was almon tree (*Shorea almon*) with a height of 20m and a DBH of 10cm. The vegetation composed of moderately to highly dense dipterocarps. The underbrush composed of dipterocarp wildlings and carabao grass. The understory also included palo verde (*Parkinsonia aculeata*), guinea grass and “camariang gubat” (*Kibatalia macgregorii*). Mosses, a few weeping figs (*Ficus sp.*), and rattan vines were also present, as well as fallen branches and twigs. The site had a porous, sandy, and loamy type of soil covered with approximately 10cm of leaf litter and humus. Anthropogenic clearing can be observed approximately 10m from the site.

Site 6 at 7°02'48.9"N 122°00'52.9"E is also a forest fragment and a highly disturbed dipterocarp forest due to a human settlement nearby. It is generally exposed to sunlight with an elevation of 842m, flat to rugged slope, and a lateral stagnant clear creek. The canopy was the tan-

bark oak (*Lithocarpus* sp.) with a height of 30m and a DBH of 15cm. Lichens and mosses were present but vines were absent. Carabao grass, some ferns including giant fern, "lipang-aso" (*Urtica dioica*), and *Colocasia esculenta* were observed. The site had a clay loamy type of soil covered with approximately 10cm of leaf litter and humus, as well as a few exposed metamorphic rocks.

Collection, Identification, and Processing of Samples

The opportunistic sampling method was employed for seven field days and 126 person-hours in six sampling sites. Samples were captured using sweep netting and were put in a glassine paper. Only two to three voucher specimens were taken, the rest were released back to the habitat. Glassine papers containing the voucher specimens were enclosed in a plastic container with mothballs to prevent other insects from penetrating the specimens. For the moths, only the diurnal ones were captured. Photo documentation in the field was done. Initial identification was done using the Philippine Lepidoptera website and verified by Dr. Jade Aster T. Badon of Silliman University and a member of Philippine Lepidoptera Inc. while moth identification was verified by Dr. Leana Lahom Cristobal, founder of Philippine Lepidoptera Inc. and a member of Asian Lepidoptera Conservation.

Statistical Analysis

Paleontological Statistics Software Package

(PAST) version 3.0 was used in calculating the biodiversity indices.

RESULTS AND DISCUSSION

Thirty-nine species of Lepidoptera were recorded in six sampling sites in WMSU-EFA, Zamboanga City, Philippines. This record is higher than the ones recorded in Central Kalimantan, Indonesia [25] and Bega Watershed, Agusan del Sur [20]. The high species richness was observed to be due to the rich plant composition of the sampling area, which is mostly dipterocarp forest. However, the results were lower than the Lepidoptera composition on Dinagat Island [26], Mt. Hamiguitan [21], Mt. Kitanglad, Mt. Apo, Mt. Musuan, and Mt. Timpoong [27]. This was observed to be due to the difference in the sampling techniques employed. The studies mentioned used insect traps which can capture more individuals and more species but only sweep netting was employed in the present study.

Out of the 39 species, 23 are butterflies belonging to four families and 20 genera (Table 1). Fifteen species are from family Nymphalidae, three from Lycaenidae, four from Pieridae, and one from Papilionidae. Nymphalidae was the dominant family, most abundant, and had the highest species richness. The same results were obtained by Marchiori & Romanowski [28]. Nymphalids are always dominant in tropical regions because of their polyphagous nature which enables them to survive in different habitats [29]. Family Nymphalidae also has many active flying butterflies that search for food in large areas [30].

Table 1. Species richness and relative abundance (in parenthesis) of butterflies in WMSU-EFA.

Species Name	Agroecosystem		Secondary Forest		Forest Fragment		Total
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1 019 masl)	Site 6 (842 masl)	
Family Lycaenidae							
<i>Celarchus archagathos archagathos</i> (Fruhstorfer, 1910)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	1 (0.72)	2 (1.44)
<i>Jamides</i> sp.	1 (0.72)	0 (0)	1 (0.72)	0 (0)	0 (0)	0 (0)	2 (1.44)
<i>Prosotas</i> sp.	4 (2.90)	0 (0)	0 (0)	1 (0.72)	0 (0)	2 (1.44)	7 (5.07)
Family Nymphalidae							
<i>Cethosia luzonica magindanaica</i> (Semper, 1888)	3 (2.17)	1 (0.72)	0 (0)	1 (0.72)	0 (0)	0 (0)	5 (3.62)
<i>Euploea mulciber mindanensis</i> (Staudinger, 1885)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)

<i>Faunis phaon leucis</i> (Felder & Felder, 1861)	3 (2.17)	0 (0)	3 (2.17)	2 (1.44)	4 (2.90)	4 (2.90)	16 (11.59)
<i>Idea electra electra</i> (Semper, 1878)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	0 (0)	1 (0.72)
<i>Ideopsis gaura glaphyra</i> (Moore, 1883)	1 (0.72)	0 (0)	2 (1.44)	0 (0)	0 (0)	0 (0)	3 (2.17)

Table 1. Species richness and relative abundance (in parenthesis) of butterflies in WMSU-EFA. (cont.)

Species Name	Agroecosystem		Secondary Forest		Forest Fragment		Total
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1 019 masl)	Site 6 (842 masl)	
<i>Junonia hedonia ida</i> (Cramer, 1775)	1 (0.72)	0 (0)	0 (0)	3 (2.17)	0 (0)	5 (3.62)	9 (6.52)
<i>Mycalesis ita imeldae</i> (Aoki & Uemura, 1982)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)
<i>Mycalesis</i> sp.	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.72)	1 (0.72)
<i>Parantica luzonensis luzonensis</i> (C. & R. Felder, 1863)	2 (1.44)	0 (0)	0 (0)	2 (1.44)	0 (0)	0 (0)	4 (2.90)
<i>Ragadia melindena melindena</i> (C. & R. Felder, 1863)	1 (0.72)	0 (0)	0 (0)	2 (1.44)	0 (0)	0 (0)	3 (2.17)
<i>Symbrenthia lilaea semperi</i> (Moore, 1899)	0 (0)	1 (0.72)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.72)
<i>Tanaecia leucotaenia leucotaenia</i> (Semper, 1878)	0 (0)	0 (0)	1 (0.72)	3 (2.17)	0 (0)	0 (0)	4 (2.90)
<i>Ypthima sempera chaboras</i> (Fruhstorfer, 1911)	3 (2.17)	3 (2.17)	0 (0)	2 (1.44)	0 (0)	0 (0)	8 (5.80)
<i>Ypthima stelleri stelleri</i> (Eschscholtz, 1821)	0 (0)	3 (2.17)	0 (0)	4 (2.90)	1 (0.72)	0 (0)	8 (5.80)
Family Papilionidae							
<i>Papilio antonio antonio</i> (Hewitson, 1875)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)
Family Pieridae							
<i>Delias diaphana diaphana</i> (Semper, 1878)	1 (0.72)	2 (1.44)	0 (0)	0 (0)	2 (1.44)	0 (0)	5 (3.62)
<i>Eurema hecabe tamiathis</i> (Fruhstorfer, 1910)	1 (0.72)	6 (4.35)	2 (1.44)	5 (3.62)	3 (2.17)	2 (1.44)	19 (13.77)
<i>Eurema sarilata sarilata</i> (Semper, 1891)	0 (0)	2 (1.44)	0 (0)	0 (0)	0 (0)	0 (0)	2 (1.44)
<i>Pareronia boebera boebera</i> (Eschscholtz, 1821)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)
Total Number of Individuals	21	18	10	30	15	15	104
Total Number of Species	11	7	6	15	5	6	23

For the moth species, 16 species belong to four families and six subfamilies (Table 2). Nine species are from family Erebidae, five from Crambidae, and one species each for families Zygaenidae and Geometridae.

Of the 39 species of Lepidoptera, the most dominant was the butterfly *Eurema hecabe tamiathis* with 19 individuals. The study of Bora et al. [29] also had *Eurema hecabe* as the most dominant species. *Eurema hecabe* is a generalist species

which means that it can thrive in many kinds of habitats [31].

The results of this study showed that butterfly and moth species are highly abundant in sampling site 4, which is a secondary dipterocarp forest. The same results can be observed in the study of Ramirez and Mohagan [22] in Tandag, Surigao del Sur. Jew et al. [32] reported that species richness and abundance highly depend on the heterogeneity of a habitat. The abundance in site 4 was observed to be due to the

higher vegetation diversity present in the site. The site with the least abundance was sampling site 6, a forest fragment and a highly disturbed dipterocarp forest. The same results were recorded in the study of Nuñez et al. [20] in Bega Watershed, Agusan del Sur. According to Leksono et al. [33], overall abundance and species richness decline with increasing site disturbance.

Sampling site 3, which is a secondary dipterocarp forest, had the most number of species. The study of Ramirez and Mohagan [22] in Tandag, Surigao del Sur also had the highest species richness in the dipterocarp forest. This site was moderately disturbed and according to Vu & Vu [34], slightly disturbed areas give rise to more diverse plants. This is a very positive effect be-

cause when there is higher plant diversity in the area, more Lepidoptera species will thrive [35]. The second species-rich site is sampling site 1, an agroecosystem site, similar to the study of Toledo & Mohagan [23] in Mt. Timpoong, Camiguin Island. This site is near a secondary forest or a less disturbed area, which is usually a site with higher species richness [36]. This difference in species composition indicates the presence of an edge effect. A study by Chacoff and Aizen [37] in Argentina showed that forest edges have higher morphospecies number than in plantations or agroecosystems. The proximity of the forests to the agroecosystem site can cause the mixing of species pool [38]. Therefore, it can influence the species composition of the agroecosystem site [39].

Table 2. Species richness and relative abundance (in parenthesis) of moth species in WMSU-EFA.

Species Name	Agroecosystem		Secondary Forest		Forest Fragment		Total
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1019 masl)	Site 6 (842 masl)	
Family Crambidae							
<i>Cnaphalocrocis</i> sp.	0 (0)	2 (1.44)	0 (0)	0 (0)	0 (0)	0 (0)	2 (1.44)
<i>Conogethes</i> sp.	2 (1.44)	2 (1.44)	0 (0)	0 (0)	0 (0)	1 (0.72)	5 (3.62)
<i>Nevrina procopia</i> (Stoll, 1781)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	0 (0)	1 (0.72)
Unidentified Pyraustinae	0 (0)	1 (0.72)	0 (0)	1 (0.72)	0 (0)	0 (0)	2 (1.44)
Unidentified Spilomelinae	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)
Family Erebidae							
<i>Chalciope mygdon</i> (Cramer, 1777)	1 (0.72)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.72)
<i>Mocis frugalis</i> (Fabricius, 1775)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	0 (0)	1 (0.72)
<i>Mocis undata</i> (Fabricius, 1775)	1 (0.72)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	2 (1.44)
<i>Nyctemera adversata</i> (Schaller, 1788)	0 (0)	0 (0)	0 (0)	2 (1.44)	0 (0)	0 (0)	2 (1.44)
<i>Nyctemera coleta</i> (Stoll, 1781)	0 (0)	1 (0.72)	0 (0)	2 (1.44)	0 (0)	0 (0)	3 (2.17)
<i>Nyctemera contrasta contrasta</i>	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)
Unidentified Arctiinae	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	0 (0)	1 (0.72)
Unidentified Arctiinae	2 (1.44)	4 (2.90)	0 (0)	0 (0)	0 (0)	1 (0.72)	7 (5.07)
Unidentified Lisothiini	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.72)	0 (0)	1 (0.72)
Family Geometridae							
<i>Eumelea</i> sp.	0 (0)	0 (0)	0 (0)	2 (1.44)	0 (0)	1 (0.72)	3 (2.17)

Table 2. Species richness and relative abundance (in parenthesis) of moth species in WMSU-EFA. (cont.)

Species Name	Agroecosystem		Secondary Forest		Forest Fragment		Total
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1019 masl)	Site 6 (842 masl)	
Family Zygaenidae							
<i>Eucorma mindanaoensis</i> (Kishida, 1996)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.72)	1 (0.72)
Total Number of Individuals	6	10	2	10	2	4	34
Total Number of Species	4	5	2	7	2	4	16

Table 3 shows the presence and absence of Lepidoptera species in six sampling sites. The most species-rich site was sampling site 4, the secondary dipterocarp forest, where *Eurema hecabe tamiathis*, *Faunis phaon leucis*, and *Tanaecia leucotaenia leucotaenia* were found. Species found in forest fragments, sites 5 and 6, are *Faunis phaon leucis* and *Eurema hecabe tamiathis*, which means that these species can survive in highly disturbed areas even though the vegetation is slowly changing. The species that are found in agroecosystems sites (site 1 and site 2) are *Cethosia luzonica magindanaica*, *Ypthima sempera chaboras*, *Delias diaphana diaphana*, *Eurema hecabe tamiathis*, and an unidentified Arctiinae. The distribution of these species highly depends on food availability, food variation, and light preferences [20, 40]. Of all the species recorded in this study, only *Eurema hecabe tamiathis* can be found in all sites. Aside

from this species' generalist nature, it also prefers places near roads and most of the sampling sites in this study are near roads [31]. There were nine Philippine endemic species recorded in this study, namely: *Celarchus archagathos archagathos*, *Delias diaphana diaphana*, *Eurema sarilata sarilata*, *Idea electra electra*, *Nyctemera contrast contrasta*, *Pareronia boebers boebers*, *Ragadia melindena melindena*, *Ypthima sempera chaboras*, and *Ypthima stelleri stelleri*. Four Mindanao endemic species were recorded, namely *Cethosia luzonica magindanaica*, *Ideopsis gaura glaphyra*, and *Mycalesis ita imeldae*. Overall endemism is 33%.

The butterfly *Idea electra* is a threatened species and red-listed as vulnerable by the IUCN [41]. It was found only in sampling site 3, a secondary forest. The rest of the species have not yet been evaluated by the IUCN.

Table 3. Presence and absence of butterflies and moths in WMSU-EFA.

Species Name	Agroecosystem		Secondary Forest		Forest Fragment	
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1019 masl)	Site 6 (842 masl)
BUTTERFLIES						
Family Lycaenidae						
<i>Celarchus archagathos archagathos*</i>						
<i>Jamides</i> sp.						
<i>Prosotas</i> sp.						
Family Nymphalidae						
<i>Cethosia luzonica magindanaica**</i>						

Table 3. Presence and absence of butterflies and moths in WMSU-EFA. (cont.)

Species Name	Agroecosystem		Secondary Forest		Forest Fragment	
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1019 masl)	Site 6 (842 masl)
<i>Euploea mulciber mindanensis</i>						
<i>Faunis phaon leucis</i>						
<i>Idea electra electra*</i>						
<i>Ideopsis gaura glaphyra**</i>						
<i>Junonia hedonia ida</i>						
<i>Mycalesis ita imeldae**</i>						
<i>Mycalesis</i> sp.						
<i>Parantica luzonensis luzonensis</i>						
<i>Ragadia melindena melindena*</i>						
<i>Symbrenthia lilaea semperi</i>						
<i>Tanaecia leucotaenia leucotaenia</i>						

<i>Ypthima sempera chaboras*</i>						
<i>Ypthima stelleri stelleri*</i>						
Family Papilionidae						
<i>Papilio antonio antonio</i>						
Family Pieridae						
<i>Delias diaphana diaphana*</i>						
<i>Eurema hecabe tamiathis</i>						
<i>Eurema sarilata sarilata*</i>						
<i>Pareronia boebers boebers*</i>						
MOTHS						
Family Crambidae						
<i>Cnaphalocrocis sp.</i>						
<i>Conogethes sp.</i>						
<i>Nevrina procopia</i>						
Unidentified Pyraustinae						
Unidentified Spilomelinae						
Family Erebidae						
<i>Chalciope mygdon</i>						
<i>Mocis frugalis</i>						
<i>Mocis undata</i>						
<i>Nyctemera adversata</i>						
<i>Nyctemera coleta</i>						
<i>Nyctemera contrasta contrasta*</i>						
Unidentified Arctiinae						

Table 3. Presence and absence of butterflies and moths in WMSU-EFA. (cont.)

Species Name	Agroecosystem		Secondary Forest		Forest Fragment	
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1019 masl)	Site 6 (842 masl)
Unidentified Arctiinae						
Unidentified Lisothiini						
Family Geometridae						
<i>Eumelea sp.</i>						
Family Zygaenidae						
<i>Eucorma mindanaoensis</i>						

■ - Presence, □ - Absence, *Philippine endemic, **Mindanao endemic,

Table 4 shows the biodiversity indices of Lepidoptera species in this study. The average Shannon-Wiener Diversity Index of all six sampling sites was 2.2625, which means that WMSU-EFA has moderate diversity [42]. Sampling site 3, a secondary dipterocarp forest, was the most diverse site (H'=2.993). This is because of its diverse vegetation, presence of water, and slight light exposure [35]. This site was near a primary

forest and since secondary forests can also support species from primary forests, some species shown in site 3 of this study may have also come from the primary forest [43]. The second most diverse site was sampling site 1, an agroecosystem site (H'=2.57). According to Tscharrntke et al. [39], the proximity of forests and agroecosystem areas has a strong influence on the diversity of the agroecosystem areas. The site with the lowest species diversity was site 5, a forest

fragment and a highly disturbed dipterocarp forest ($H'=1.633$). It was near an anthropogenic clearing. According to Irwin et al. [44], disturbances, especially severe ones like deforestation,

typically decrease species diversity, including native or endemic ones. Distribution was even in all sampling sites indicating that there was no dominant species.

Table 4. Biodiversity indices of lepidoptera in WMSU-EFA.

Species Name	Agroecosystem		Secondary Forest		Forest Fragment		Average
	Site 1 (875 masl)	Site 2 (645 masl)	Site 3 (990 masl)	Site 4 (645 masl)	Site 5 (1019 masl)	Site 6 (842 masl)	
Taxa	15	12	8	23	6	10	12.3
Individual	27	28	12	42	12	19	23.3
Shannon	2.57	2.317	1.979	2.993	1.633	2.083	2.2625
Evenness	0.8712	0.8452	0.9046	0.8672	0.8529	0.803	0.8574

CONCLUSION

This study has recorded 39 Lepidoptera species consisting of 23 butterflies and 16 moth species. Endemism is at 33% with nine Philippine endemic and three Mindanao endemic species. *Idea electra*, a Philippine endemic butterfly species, was the only recorded threatened species and red-listed as vulnerable by the IUCN. The study area, WMSU-EFA, has moderate diversity ($H'=2.2625$) which is attributed to a variety of food sources and the diverse vegetation of the sampling area. Family Nymphalidae was the dominant, most abundant, and most species-rich family. This is highly due to its polyphagous and active flying nature. *Eurema hecabe tamiathis* was the most distributed species. Sampling site 4, a secondary forest, was the most diverse site ($H'=2.993$), most abundant (30.00%), and with the highest species richness ($S=23$) mainly because it was composed of diverse vegetation and enough light exposure. The severe anthropogenic clearing was the observed threat that is most likely to affect the diversity of Lepidoptera in the area.

ACKNOWLEDGMENT

We acknowledge Prof. Noel B. Salatan of Western Mindanao State University (WMSU) for logistical support and the people of WMSU-Experimental Forest Area for assistance in sampling.

Conflict of Interest

The authors declare no conflict of interest.

REFERENCES

1. Cabrido C, Demayo C G. Antimicrobial and cellular metabolic effects of the ethanolic extract of the dallas red variety of lantana camara. *Pharmacophores*. 2018; 9(1): 10-18.
2. Zeghti S, Bouras A, Kherbouche Y, Sekour M. Assessment of the Diversity of Lepidoptera and the Importance of Useful and Harmful Species in Palm Groves (Algeria). *World Journal of Environmental Biosciences*. 2019; 8(1): 11-20.
3. Regier JC, Zwick A, Cummings MP, Kawahara AY, Cho S, Weller S, Roe A, Baixeras J, Brown JW, Parr C, Davis DR. Toward reconstructing the evolution of advanced moths and butterflies (Lepidoptera: Ditrysia): an initial molecular study. *BMC Evolutionary Biology*. 2009 Dec 1;9(1):280.
4. Saad A, Bounaceur F. Diversity of Lepidoptera (Rhopalocera) in natural and modified habitats of Bousaâda, Algeria. *World Journal of Environmental Biosciences*. 2018; 7(1):79-83.
5. Krenn HW. Feeding mechanisms of adult Lepidoptera: structure, function, and evolution of the mouthparts. *Annual review of entomology*. 2010 Jan 7;55:307-27.
6. Khyade V B. The Importance of Nature (Length) of Proboscis in Hesperidae Butterflies. *World Journal of Environmental Biosciences*. 2019; 8(2): 1-10.
7. Choudhary N K N, Mallya R. Phytochemical investigation and antibacterial activity of a medicinal plant. *Int. J. Pharm. Phytopharm. Res*. 2019; 9(4): 53-58.
8. Thomas JA. Monitoring change in the abundance and distribution of insects using butterflies and other indicator groups. *Philosophical Transactions of the Royal So-*

- ciety B: Biological Sciences. 2005 Feb 28;360(1454):339-57.
9. Bonebrake TC, Ponisio LC, Boggs CL, Ehrlich PR. More than just indicators: a review of tropical butterfly ecology and conservation. *Biological conservation*. 2010 Aug 1;143(8):1831-41.
 10. van Nieuwerkerken EJ, Kaila L, Kitching IJ, Kristensen NP, Lees DC, Minet J, Mitter C, Mutanen M, Regier JC, Simonsen TJ, Wahlberg N. Order Lepidoptera Linnaeus, 1758. In: Zhang, Z.-Q.(Ed.) *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*. 2011 Dec 23;3148(1):212-21.
 11. Wahlberg N, Wheat CW, Peña C. Timing and patterns in the taxonomic diversification of Lepidoptera (butterflies and moths). *PLOS one*. 2013;8(11). <https://www.ncbi.nlm.nih.gov/pmc/article/PMC3839996/>
 12. Nöske NM, Hilt N, Werner FA, Brehm G, Fiedler K, Sipman HJ, Gradstein SR. Disturbance effects on diversity of epiphytes and moths in a montane forest in Ecuador. *Basic and Applied Ecology*. 2008 Jan 2;9(1):4-12.
 13. Lehmann I, Kioko E. Lepidoptera diversity, floristic composition and structure of three Kaya forests on the south coast of Kenya. *Journal of East African Natural History*. 2005 Jan;94(1):121-63.
 14. Uehara-Prado M, Brown Jr KS, Freitas AV. Species richness, composition and abundance of fruit-feeding butterflies in the Brazilian Atlantic Forest: comparison between a fragmented and a continuous landscape. *Global Ecology and Biogeography*. 2007 Jan;16(1):43-54.
 15. Dumbrell AJ, Hill JK. Impacts of selective logging on canopy and ground assemblages of tropical forest butterflies: implications for sampling. *Biological conservation*. 2005 Sep 1;125(1):123-31.
 16. Abang F, Karim C. Diversity of macromoths (Lepidoptera: Heterocera) in the Poring Hill dipterocarp forest, Sabah, Borneo. *Journal of Asia-Pacific Entomology*. 2005 Feb 1;8(1):69-79.
 17. Majumder J, Lodh R, Agarwala BK. Butterfly species richness and diversity in the Trishna Wild-life Sanctuary in South Asia. *Journal of Insect Science*. 2013 Jan 1;13(1):79. <https://doi.org/10.1673/031.013.7901>
 18. Kudavidanage EP, Wanger TC, De Alwis C, Sanjeewa S, Kotagama SW. Amphibian and butterfly diversity across a tropical land-use gradient in Srilanka; implications for conservation decision making. *Animal Conservation*. 2012 Jun;15(3):253-65.
 19. Van Lien V, Yuan D. The differences of butterfly (Lepidoptera, Papilionoidea) communities in habitats with various degrees of disturbance and altitudes in tropical forests of Vietnam. *Biodiversity & Conservation*. 2003 Jun 1;12(6):1099-111.
 20. Nuñez KM, Nuñez OM, Dupo AL. B. Species Richness of Lepidoptera in Bega Watershed, Prosperidad, Agusan del Sur, Philippines. *Bull. Env. Pharmacol. Life Sci*. 2016 Jul 8;5(8):83-90.
 21. Mohagan AB, Treadaway CG. Diversity and status of butterflies across vegetation types of Mt. Hamiguitan, Davao Oriental, Philippines. *Asian Journal of Biodiversity*. 2010 Dec 2;1(1):1-24.
 22. Ramirez RK, Mohagan AB. Diversity and Status of Butterflies in Maitum Village, Tandag, Surigao del Sur, Philippines. *Asian Journal of Biodiversity*. 2012 Jan 1;3(1):74-112.
 23. Toledo JM, Mohagan AB. Diversity and Status of Butterflies in Mt. Timpoong and Mt. Hibok-hibok, Camiguin Island, Philippines. *JPAIR Multidisciplinary Journal*. 2011;6: 103-116.
 24. Google Maps. Philippines. [Image on internet]. 2019 [updated 2019; cited April 9, 2019]. Available from <https://www.google.com/maps/@12.0609728,121.8550578,1660577m/data=!3m1!1e3>
 25. Houlihan PR, Harrison ME, Cheyne SM. Impacts of forest gaps on butterfly diversity in a Bornean peat-swamp forest. *Journal of Asia-Pacific Entomology*. 2013 Mar 1;16(1):67-73.
 26. Mohagan AB, Mohagan DP, Libor R. Diversity and Status of Butterflies in Dinagat Island, Philippines. *SDSSU Multidisciplinary Research Journal*. 2013;1(1):34-40.
 27. Mohagan DP, Tambuli AE, Mohagan AB. Diversity of butterflies in the selected key biodiversity areas of Mindanao, Philippines. *Asian Journal Of Biodiversity*. 2013 Dec 3;2(1):121-148.
 28. Marchiori MO, Romanowski HP. Species composition and diel variation of a butterfly taxocene (Lepidoptera, Papilionoidea and Hesperioidea) in a restinga forest at Itapuã State Park, Rio Grande do Sul, Brazil. *Revista Brasileira de Zoologia*. 2006 Jun;23(2):443-54.
 29. Bora A, Meitei LR, Deb M. Butterfly species richness and diversity in experimental botanic garden, botanical survey of India, ERC, Umiam, Meghalaya, India. *Journal of*

- Entomology and Zoology Studies. 2014;2(5):212-7.
30. Eswaran R, Pramod P. Structure of butterfly community of Anaikatty hills, Western Ghats. Zoos' Print Journal. 2005;20(8):1939-42.
 31. Cleary DF, Boyle TJ, Setyawati T, Menken SB. The impact of logging on the abundance, species richness and community composition of butterfly guilds in Borneo. Journal of applied Entomology. 2005 Feb;129(1):52-9.
 32. Jew EK, Loos J, Dougill AJ, Sallu SM, Benton TG. Butterfly communities in miombo woodland: Biodiversity declines with increasing woodland utilisation. Biological Conservation. 2015 Dec 1;192:436-44.
 33. Leksono AS, Nakagoshi N, Isagi Y. The effects of forest disturbances on flying insect assemblages in Trawas, east Java. Tropics. 2005;14(4):335-43.
 34. Vu LV, Quang Vu C. Diversity pattern of butterfly communities (Lepidoptera, Papilionoidea) in different habitat types in a tropical rain forest of Southern Vietnam. ISRN Zoology. 2011 Apr 7;2011: 1-8.
 35. Hawes J, da Silva Motta C, Overal WL, Barlow J, Gardner TA, Peres CA. Diversity and composition of Amazonian moths in primary, secondary and plantation forests. Journal of Tropical Ecology. 2009 May;25(3):281-300.
 36. Tylianakis JM, Klein AM, Tscharrntke T. Spatiotemporal variation in the diversity of Hymenoptera across a tropical habitat gradient. Ecology. 2005 Dec;86(12):3296-302.
 37. Chacoff NP, Aizen MA. Edge effects on flower-visiting insects in grapefruit plantations bordering premontane subtropical forest. Journal of Applied Ecology. 2006 Feb 1;43(1):18-27.
 38. Axmacher JC, Tünte H, Schrupf M, Müller-Hohenstein K, Lyaruu HV, Fiedler K. Diverging diversity patterns of vascular plants and geometrid moths during forest regeneration on Mt Kilimanjaro, Tanzania. Journal of Biogeography. 2004 Jun;31(6):895-904.
 39. Tscharrntke T, Sekercioglu CH, Dietsch TV, Sodhi NS, Hoehn P, Tylianakis JM. Landscape constraints on functional diversity of birds and insects in tropical agroecosystems. Ecology. 2008 Apr;89(4):944-51.
 40. Kumar V, Reddy PV, Anal AK, Nath V. Outbreak of the looper, *Perixera illepidaria* (Lepidoptera: Geometridae) on litchi, *Litchi chinensis* (Sapindales: Sapindaceae)-a new pest record from India. Florida Entomologist. 2014 Mar;97(1):22-9.
 41. Lepidoptera Specialist Group 1996. *Idea electra*. The IUCN Red List of Threatened Species 1996: e.T10781A3215545. [cited 2019 April 22]. Retrieved from: <https://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T10781A3215545.en>.
 42. Odum EP. *Fundamentals of Ecology*. Third ed. Philadelphia, USA: W.D. Saunders Corporation; 1971. 574 pp.
 43. Dent DH, Wright SJ. The future of tropical species in secondary forests: a quantitative review. Biological conservation. 2009 Dec 1;142(12):2833-43.
 44. Irwin MT, Wright PC, Birkinshaw C, Fisher BL, Gardner CJ, Glos J, Goodman SM, Loiselle P, Rabeson P, Raharison JL, Raherilalao MJ. Patterns of species change in anthropogenically disturbed forests of Madagascar. Biological Conservation. 2010 Oct 1;143(10):2351-62.