Dry grasslands on sandy soils in the forest and forest-steppe zones of the plains region of Ukraine: present state of syntaxonomy

- Anna Kuzemko -

Abstract

This paper compiles and synthesizes the present knowledge of dry grassland vegetation on sandy soils in northern Ukraine, based on published and unpublished sources. In total, 391 relevés from 29 association-level syntaxa of the class Koelerio-Corynephoretea (incl. Sedo-Scleranthetea and Festucetea vaginatae), as well as of the alliance Agrostion vinealis from the class Molinio-Arrhenatheretea were used for the analysis. Based on cluster analysis and comparison of synoptic tables, a compact classification scheme of the sandy dry grassland vegetation in the forest and forest-steppe zones of the plains region of Ukraine has been developed. It comprises 10 associations from the Koelerio-Corynephoretea and one from the Agrostion vinealis (Molinio-Arrhenatheretea).

The Koelerio-Corynephoretea include four orders in Ukraine: Alysso-Sedetalia, Trifolio arvensis-Festucetalia ovinae, Corynephoretalia canescentis, and Sedo acris-Festucetalia. The Alysso-Sedetalia are represented by two associations, the Aurinio saxatilis-Allietum podolici and the Minuartio auctae-Festucetum pallentis from West Podillya. The associations Stipetum pulcherrimae, Astragalo-Stipetum, and Poetum versicoloris, which also had been included in the Alysso-Sedetalia by some authors, are assigned to the class Festuco-Brometea based on their species composition. All Ukrainian stands of the Trifolio arvensis-Festucetalia ovinae are assigned to the association Thymo pulegioides-Festucetum ovinae. They develop on granite outcrops within Ukrainian crystalline shield. Communities of the Corynephoretalia canescentis are described from the Polissya region, where fluvioglacial sediments dominate. These relevés are provisionally assigned to the associations Corniculario aculeatae-Corynephoretum canescentis and Corynephoro-Silenetum tataricae. Sedo acris-Festucetalia communities have mainly been recorded in the Dnieper Valley, and some relevés come from Zhytomyr Polissya. This order includes two alliances. The Koelerion glaucae is represented by two associations, namely Veronico dillenii-Secalietum sylvestris and Diantho borbasii-Agrostietum syreistschikovii. The Festucion beckeri is represented by three associations (Centaureo borysthenicae-Festucetum beckeri, Chamaecytiso ruthenici-Festucetum beckeri and perhaps Poo bulbosae-Caricetum colchicae), which are distributed in the Dnieper Valley. The Agrostion vinealis alliance from the Galietalia veri order has been assigned to the Molinio-Arrhenatheretea. The present analysis revealed a significant similarity among the previously described associations of this alliance, which allowed the reduction of them to one association, the Koelerio-Agrostietum vinealis. It occurs mostly in the Dnieper Valley and its left bank tributaries. Among the right bank tributaries, it occurs only along lower stretches of rivers and around their confluences with the Dnieper. Most of the dry grassland communities on sandy soils within the study area require further detailed investigation.

Zusammenfassung: Sandtrockenrasen in der Wald- und Waldsteppenzone des ukrainischen Flachlandes: aktueller Stand der Syntaxonomie

Dieser Artikel synthetisiert das gegenwärtige Wissen zur Vegetation der Sandtrockenrasen in der nördlichen Ukraine basierend auf publizierten und unpublizierten Quellen. Insgesamt wurden 391 Aufnahmen von 29 Syntaxa auf Assoziationsebene aus der Klasse Koelerio-Corynephoretea (incl. Sedo-Scleranthetea und Festucetea vaginatae) sowie aus dem Verband Agrostion vinealis (Klasse Molinio-Arrhenatheretea) für die Analyse verwendet. Basierend auf einer Clusteranalyse und der erstellten Stetigkeitstabelle wurde eine konsistente Klassifikation der Sandtrockenrasenvegetation in der Waldund Waldsteppenzone des ukrainischen Flachlandes entwickelt. Sie umfasst 10 Assoziationen der Klasse Koelerio-Corynephoretea and eine aus dem Agrostion vinealis (Molinio-Arrhenatheretea).

Zu den Koelerio-Corynephoretea gehören im Gebiet vier Ordnungen: Alysso-Sedetalia, Trifolio arvensis-Festucetalia ovinae, Corynephoretalia canescentis und Sedo acris-Festucetalia. Die Alysso-Sedetalia sind durch zwei Assoziationen vertreten, das Aurinio saxatilis-Allietum podolici und das Minuartio auctae-Festucetum pallentis West-Podoliens. Die Assoziationen Stipetum pulcherrimae, Astragalo-Stipetum und Poetum versicoloris, die von manchen Autoren ebenfalls in den Alysso-Sedetalia

eingeschlossen wurden, werden hier aufgrund ihrer Artenkombination zur Klasse Festuco-Brometea gestellt. Alle ukrainischen Bestände der Ordnung Trifolio arvensis-Festucetalia ovinae werden in eine einzige Assoziation gestellt, das Thymo pulegioides-Festucetum ovinae. Dieses tritt im Bereich von Granitfelsen im Ukrainischen Tafelland auf. Gesellschaften der Ordnung Corynephoretalia canescentis werden aus Polesien beschrieben, wo fluvioglaziale Sedimente vorherrschen. Derartige Aufnahmen werden provisorisch den Assoziationen Corniculario aculeatae-Corynephoretum canescentis und Corynephoro-Silenetum tataricae zugeordnet. Gesellschaften der Ordnung Sedo acris-Festucetalia wurden vor allem im Dnepr-Tal nachgewiesen, wobei einige Aufnahmen auch aus der Oblast Schytomyr stammen. Diese Ordnung ist im Gebiet durch zwei Verbände vertreten. Zum Koelerion glaucae gehören das Veronico dillenii-Secalietum sylvestris und das Diantho borbasii-Agrostietum syreistschikovii. Das Festucion beckeri ist durch drei Assoziationen vertreten (Centaureo borysthenicae-Festucetum beckeri, Chamaecytiso ruthenici-Festucetum beckeri und möglicherweise Poo bulbosae-Caricetum colchicae), die alle im Dnepr-Tal vorkommen. Der Verband Agrostion vinealis aus der Ordnung Galietalia veri wird zur Klasse Molinio-Arrhenatheretea gestellt. Es zeigte sich, dass die verschiedenen zuvor in diesem Verband unterschiedenen Gesellschaften alle untereinander so ähnlich sind, dass sie in einer Assoziation zusammengefasst werden sollten, dem Koelerio-Agrostietum vinealis. Dieses kommt in den Tälern des Dnepr und seiner linksseitigen Zuflüsse vor. Dagegen tritt es an den rechtsseitigen Zuflüssen nur im Unterlauf auf. Zusammenfassend lässt sich feststellen, dass für die Mehrzahl der behandelten Gesellschaften weitere Untersuchungen erforderlich sind.

Keywords: Agrostion vinealis, Festuco-Brometea, Galietalia veri, Koelerio-Corynephoretea, Molinio-Arrhenatheretea, Dnieper Valley, Podillya, Polissya.

1. Introduction

The vegetation of Ukraine has been studied using the Braun-Blanquet approach for about 30 years. The results of the preliminary inventory of coenotic-level phytodiversity were generalized in the first "Prodrome of Ukrainian vegetation" (SOLOMAKHA 1996). During the initial stages of developing a syntaxonomic classification of the country, Ukrainian phytosociologists faced the problem of difficulty of access of important literature on European vegetation. Moreover, there was a lack of vegetation data from the more eastern regions in most cases. This causes difficulty in aligning syntaxa described from the territory of Ukraine with those described from other regions. There were no opportunities to analyze syntaxa within their complete distribution range. Thus, some syntaxa were misapplied in the "Prodrome of Ukrainian vegetation", their distribution actually being limited to more western areas of Europe. Additionally, it turned out that some syntaxa that have been described as new for science from the territory of Ukraine actually had previously been described elsewhere. In particular, some syntaxa have repeatedly been described under different names from the territories of Ukraine, Belarus and the Russian Federation.

During the last 10 years, the preparation of the multi-volume "Vegetation of Ukraine" has commenced, with the first three volumes already published (MALINOVSKIY & KRICH-FALUSHIY 2000, DUBYNA 2006, DUBYNA et al. 2007). Further, substantial information has been accumulated in recent publications, e.g. in series of monographs "Nature Protected Territories of Ukraine. Plant World" (ORLOV & YAKUSHENKO 2005, GAL'CHENKO 2006), in surveys of the vegetation of some regions of Ukraine (GONCHARENKO 2002, DUBYNA et al. 2004, GRIGORA et al. 2005) and in several dissertations dealing with Ukrainian vegetation. As a result, a considerable amount of phytosociological data has been accumulated that requires systematization, generalization and comparison with vegetation of adjacent regions. It is necessary also to clarify which vegetation classes and which regions of Ukraine need further detailed investigation and for which of them the level of knowledge is satisfactory.

Therefore, the aim of this work was to generalize all data on dry sandy grassland vegetation available in literature and archive sources for the forest and forest-steppe zones of the plains region of Ukraine; to compare the syntaxa, and to outline the direction and the tasks of further research.

2. Study area

The study area includes the forest and forest-steppe zones of the plains region of Ukraine. According to the latest version of the physiographic zonation of Ukraine (MARINICH & SHYSHCHENKO 2005, RUDENKO 2007), the forest zone was divided into the zone of mixed forests and the zone of deciduous forests (Fig. 1). The Dnieper River divides Ukraine into two parts, referred to as Right-Bank Ukraine and Left-Bank Ukraine in this paper.

The zone of mixed forests of the East European Plain occupies the northern part of Ukraine. This zone is characterised as having lowland relief with prevalence of sandy and clay-sandy deposits, temperate continental climate with positive moisture balance and a high-density river network with wide river valleys that regularly flood. The mixed forest zone is the most humid region of the plains region of Ukraine. The geomorphological structure of the zone is heterogeneous. In the west it occupies a part of the Volyno-Podilska plate and the Halytsko-Volynska depression, the central portion occupies the northern portion of the Ukrainian crystalline shield, and the eastern portion occupies the Dniprovsko-Donetska depression and the slopes of the Voronezh crystalline massif. The zone occupies polygenic lowland plains, formed mainly from alluvial, glacial, fluvioglacial, and atmogenic sediments (MARINICH & SHYSHCHENKO 2005). Sod-podzol soils, dominate, particularly weakly-developed podzol soils which are distributed on sandy terraces and moraine plains on the tops and gentle slopes of the moraine hills and ridges. On the elevated convex or flat areas of terraces with dry pine forests sod-sandy soils with clearly sandy microstructure have developed. In isolated sites, particularly in Zhytomyr Polissya, sod-skeletal soils occur. They develop on crystalline rock outcrops, and consist of sand and broken stones. In shallow watershed depressions with weak discharge, on river terraces above floodplains, and in the middle and high-level floodplains, sod-gley and meadow soils are formed. Their texture varies between light sandy and loamy (VERNANDER et al. 1986). Zonal vegetation is represented by coniferous and mixed forests with predominance of Pinus sylvestris and Quercus robur.



Fig. 1: Physiographic zonation of Ukraine (adapted from RUDENKO 2007). The study deals with the three highlighted vegetation zones.

Abb. 1: Physiografische Gliederung der Ukraine (nach RUDENKO 2007). Der Artikel beschäftigt sich mit den drei hervorgehobenen Vegetationszonen.

The zone of deciduous forests occupies the western part of the Ukrainian territory between the Carpathians, the mixed forest zone, and the forest-steppe zone. This zone extends over the West-Ukrainian physiographic region, and is partitioned by numerous rivers within the Dnister, Pivdenny Bug, Visla and Prip'yat' basins. The valleys of left tributaries of the Dnister are canyons, cutting through the section of paleozoic and mesoneozoic sediments. The cliffy, partitioned limestone hills of Tovtry are prominent in relief. They are the remainders of barrier reefs and atolls of the Neogenic Sea. The Tovtry rises about 50–60 m above the surrounding area, and its peak altitude is 350–400 m a.s.l. (MARINICH & SHYSHCHENKO 2005). The distribution of sandy soils is connected with crystalline rock exposures, where weathering of some elements takes place. As a result, sandy and skeletal soils of different thickness are formed. The exception is Male Polissya, where the natural and territorial complexes of the forested lowland prevails in the landscape: areas with outwashed, moraine-outwashed and outwash-alluvial plains with sod-weakly developed podzol soils (MARINICH & SHYSHCHENKO 2005). Zonal vegetation includes deciduous forests dominated by *Quercus robur* and *Carpinus betulus*.

The forest-steppe zone stretches from the zone of deciduous forests in the west to the west spurs of the Middle-Russian Highland in the east. Its northern border coincides with that of continuous distribution of north-forest-steppe landscapes. Their indicators are grey forest soils and podzolic chernozems on the loess stratum. The southern border is traceable on the line of the deep middle-humus chernozem. Formation and development of the foreststeppe landscapes are dependent on the optimal balance of heat and moisture, and during the growing season, evaporation almost equals the amount of precipitation. The soil formation in the Dnieper Right-Bank part of the zone is influenced by crystalline outcrops and weathering processes. Sandy soils occupy greater areas in the Dnieper Valley, especially in the Left-Bank region. This is caused by erosion and accumulative processes in the Dnieper Valley and its tributaries. Their floodplains are formed by alluvial sands with the strata of loamy sand, loam and buried soils. The sandy areas with pine forests are raised up to 10-12 m above the floodplain with the peak altitude mark of 140-150 m a.s.l. They are composed of ancient alluvial anisomerous sands, which lie over the fluvioglacial stratum. Their surface topography is diversified by sandy ridges, dunes, and blow-outs. Alluvial sediments are widespread in valleys of the Left-Bank tributaries on palaeogenic sediments (Kharkiv sands, Kiev marl). Sandy areas covered mainly with pine forests are distributed in the valleys of Sula, Psel, and Khorol and other rivers. They rise up to 10–12 m above the floodplains with a peak altitude of 120-128 m a.s.l. Typical of these sandy areas are atmogenic ridge-hilly relief and sod-podzol sandy and sabulous soils. The sandy terrace of the Vorskla River is 2-7 km wide. The sandy hills with a peak altitude of 80-100 m a. s.l. are distributed at its outside edge. In the eastern part of Ukraine, sandy terraces are located in river valleys of the Kharkiv hillside-elevated region, where floodplain sandy terraces form a belt 0.4-4.0 km wide (MARINICH & SHYSHCHENKO 2005). Zonal vegetation is represented by deciduous forests (dominated by Quercus robur and Carpinus betulus, Acer platanoides and Tilia cordata) and meadow steppes (dominated e.g. by Festuca valesiaca, Stipa capillata and Botriochloa ischemum).

3. Material and methods

3.1. Plant nomenclature

In general, the nomenclature for vascular plants follows Cherepanov (1995), that of mosses Ignatov & Afonina (1992), and that of lichens Kondratyuk et al. (1998). As an exception, the nomenclature of some vascular plant taxa follows Mosyakin & Fedoronchuk (1999) because some taxonomic interpretations by Cherepanov (1995) caused difficulties.

In the following cases, microspecies were included in aggregates and were not considered as separate taxa in the present paper: *Medicago falcata* agg. (including *M. procumbens* Besser – absent in Cherepanov's list and *M. romanica* Prodán – present in Cherepanov's list as only species), *Thymus pulegioides* agg. (including *T. clandestinus* Schur. and *T. ucrainicus* (Klokov & Des.-Shost.) Klokov –

absent in Cherepanov's list), Hylotelephium maximum agg. (including H. polonicum (Błocki) Holub – absent in Cherepanov's list, H. stepposum (Boriss.) Tzvelev – present in Cherepanov's list as only species), Hylotelephium telephium agg. (including H. argutum (Haw.) Holub – absent in Cherepanov's list and H. triphyllum (Haw.) Holub – present in Cherepanov's list as only species). Seseli libanotis (L.) W. D. J. Koch subsp. intermedium is considered as a subspecies although all cited authors applied the old name of this taxon Seseli intermedium (Rupr.) V. Vodopianova (but in Cherepanov's list it is given as Seseli libanotis). In the nomenclature of several species absent in Cherepanov's checklist, I followed Mosyakin & Fedoronchuk (1999): Asparagus polyphyllus Steven, Pilosella caespitosa (Dumort.) P. D. Sell & C. West, Seseli campestre Besser, and Stipa borysthenica Klokov ex Prokudin.

3.2. Relevés

For the survey of the Koelerio-Corynephoretea, phytosociological materials published in Ukrainian literature were used. Relevés of the alliance Agrostion vinealis (order Galietalia veri, class Molinio-Arrhenatheretea) were taken from archive data and historical relevés from "phytocoenotheka" (archives of paper-copy relevés of the Department of Geobotany at the M. G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine). Additionally, unpublished data of the author were used. All available phytosciological data (single relevés and synoptic tables) originally assigned either to the Koelerio-Corynephoretea s.l. (incl. Sedo-Scleranthetea, Festucetea vaginatae) or the alliance Agrostion vinealis were used for the analysis. Regarding the Molinio-Arrhenatheretea, only those syntaxa occurring on dry sandy soils were included in the study. Appendix A gives an overview of relevés included in the analysis.

In the analysis, I tried to use the greatest possible number of relevés of each syntaxon. Unfortunately, for some syntaxa newly described from the territory of Ukraine only very few relevés exist. Thus, the range of relevé numbers varies from 3 to 61 in different syntaxa, which may have affected the accuracy of results to a certain extent. The information about plot sizes was absent in literature sources in most cases. In those cases with available information, the size varies noticeably from 2 m² to 100 m². Another problem is the absence of information on bryophytes and lichens in some syntaxa (this is indicated in Table 1 in the Supplement). I fully realize that such variation of raw data may have an essential influence on the reliability of the results but I decided to use all data available from sandy dry grasslands in the forest and forest-steppe zone of the plains region of Ukraine because their small quantity did not allow for further reduction based on quality criteria.

3.3. Phytosociological classification and cluster analysis

A comparative analysis of the 29 "syntaxa" included was carried out to evaluate the diagnostic value of species delimited as diagnostic in original publications/information sources and to assess how well-founded their assignment was to higher syntaxa.

All the data were summarized into a synoptic table where the constancy of each species is expressed in percent (Table 1 in the Supplement). If in sources only constancy classes were given, I transformed these back to percentage constancies approximately (V: 90%; IV: 70%; III: 50%; II: 30%; I: 10%). The relevés were grouped according to the original association names assigned by the authors, with two exceptions. The two localities of the *Veronico dillenii-Secalietum sylvestris* differed considerably in their floristic composition and were therefore analyzed separately. The same was true for the *Centaureo borysthenicae-Festucetum beckeri*.

Species with constancy at least two times as high than in other syntaxa were considered as differential (DENGLER 2003). Delimitation of character species of higher syntaxa was based on literature sources (MUCINA 1997, BERG et al. 2001, MATUSZKIEWICZ 2001, DENGLER 2003, RUSINA 2005, RUPRECHT et al. 2009). Similarity among syntaxa was determined by cluster analysis with the program STATISTICA 7.0 (STATSOFT, INC. 2005). The agglomerative hierarchical method known as Ward's method was applied. As a resemblance measure, Euclidean distance was employed.

A nomenclatural revision of the involved syntaxa was not a focus of this paper. However, misspelled names were corrected according to the regulations of the International Code of Phytosociological Nomenclature (further as ICPN; Weber et al. 2000). Full author citations of all mentioned syntaxa are given in the syntaxonomic overview (section 5.4).

4. Results

4.1. Comparison of syntaxa in synoptic table (Table 1 in the Supplement)

Syntaxa SS1, SS2 and SS3 are characterized by high constancy and abundance of the *Festuco-Brometea* character species and low constancy of the *Koelerio-Corynephoretea* character species with the exception of species of the alliance *Alysso alyssoidis-Sedion*. The high constancy of the later probably was the reason why ABDULOEVA (2002) in the original publication assigned them to the *Sedo-Scleranthetea*.

Diagnostic species of syntaxa SS4 and SS5 are *Thymus* ×oblongifolius, Cephalaria uralensis and Minuartia aucta, which do not occur in any other syntaxon. Vegetation is composed mainly of diagnostic species of the Alysso-Sedion and the Koelerio-Corynephoretea.

Syntaxa SS6, SS7, SS8, SS9 and SS10 contain a number of species of the *Hyperico perfo- rati-Scleranthion perennis*, allowing them to be included into this alliance. Considerable differences among these syntaxa were not observed in the synoptic table, and there appears to be no reason to delimit them as separate associations.

Syntaxa KC1, KC3 and FV8 do not possess good diagnostic species at the association level. Most probably they belong to the Corynephorion canescentis alliance although the group of character species of the named alliance (only Corynephorus canescens, Jasione montana and Thymus serpyllum) and also of the Koelerio-Corynephoretea class (only 14 species with low constancy) is represented weakly. By contrast, syntaxon KC4 has good association-level diagnostic species. This syntaxon is probably the transition from Corynephorion canescentis to Koelerion glaucae alliances as species of both alliances are present.

Syntaxon FV7 has good association-level diagnostic species, and also includes character species of the class *Koelerio-Corynephoretea* and the alliance *Koelerion glaucae*.

Syntaxa SS11, FV5, FV1b and FV2b show a pronounced similarity in floristic composition, and can be assigned to the alliance *Koelerion glaucae* (order *Sedo acris-Festucetalia*, class *Koelerio-Corynephoretea*).

Syntaxa FV3, FV4 and FV6 contain the same set of diagnostic species and approximately equal numbers of character species of the *Koelerion glaucae* and *Festucion beckeri* alliances, indicating the transitional nature of these syntaxa.

Syntaxa FV2a and FV1a are well differentiated by such diagnostic species as Agropyron pectinatum, Centaurea pseudomaculosa, Dianthus platyodon, Erysimum canescens, Kochia prostrata Linaria dulcis and Tragopogon borysthenicus. Character species of both the alliance Festucion beckeri and the order Sedo acris-Festucetalia are well represented.

Syntaxon SS12 has a clear diagnostic species group, and some character species of the Sedo acris-Festucetalia and the Koelerio-Corynephoretea have a high constancy.

Syntaxa AV1, AV2 and AV3 demonstrate considerable similarity in floristic composition, suggesting their inclusion into a single association. This is distinguished by a high number of the *Molinio-Arrhenatheretea* character species as well as character species of the *Koelerio-Corynephoretea*. These syntaxa represent transitional vegetation between both phytosociological classes.

Syntaxon KC2 has no diagnostic species. The *Koelerio-Corynephoretea* character species are nearly absent but the *Molinio-Arrhenatheretea* class is represented quite well with some species having high constancy.

4.2 Cluster analysis

The dendrogram of the cluster analysis demonstrates clear division of the processed syntaxa into two large groups (Fig. 2). The first of them includes syntaxa of communities which occur on stony outcrops while the second contains syntaxa developing on sandy deposits. The extreme branches of the dendrogram include groups questionably positioned syntaxonomically within the *Koelerio-Corynephoretea*. Syntaxa SS1–SS3 are assigned to the *Festuco-Brometea* class in conventional classification schemes, and syntaxa AV1–AV3 traditionally are assigned to the *Molinio-Arrhenatheretea* class.




Fig. 2: Dendrogram of the cluster analysis of 29 sandy dry grassland syntaxa. The code numbers refer to Appendix A.

Abb. 2: Dendrogramm der Clusteranalyse von 29 Sandtrockenrasen-Syntaxa. Die Codes sind in Anhang A aufgeschlüsselt.


In total, six groups of clusters were obtained. The first group is formed by syntaxa SS1, SS2, SS3, SS4, and SS5. They include communities on limestone outcrops in West Podillya. The second group includes syntaxa SS6, SS7, SS8, SS9 and SS10, communities of which are described from granite outcrops of forest and forest-steppe zones of the Right-Bank Ukraine. This group is floristically quite homogeneous.


The third group is more heterogeneous. One branch is formed by syntaxa SS11, FV5, FV1b, and FV2b, and it includes phytocenoses occurring on sands of the Dnieper floodplain. The second branch contains syntaxa FV1a and FV2a with communities that were described on sands of the Left-Bank forest-steppe.

The fourth group is the most heterogeneous. One branch combines syntaxa KC1, FV8, KC3, and KC4, which include communities of sandy deposits, mainly fluvioglacial sediments of forest zone. The other branch is formed by syntaxa FV3, FV4, and FV6, which are represented by the Dnieper sandy terrace communities.

Syntaxa of the fifth group are SS12 and FV7. They demonstrate slight similarity with each other and with other syntaxa. These syntaxa include communities from the Dnieper Valley: FV7 communities are described at the most northern frontier, while SS12 occurs in the south part of the study area.

The sixth group includes syntaxa KC2, AV1, AV2 and AV3. They include communities of high sections of floodplains, and they are the most dissimilar to all other examined syntaxa. Syntaxa AV1, AV2 and AV3 are characterized by greater density of herb vegetation, but syntaxon KC2 is clearly separated from them.








Fig. 3: Distribution of sandy dry grassland communities in the forest and forest-steppe zones of the plains region of Ukraine: a – orders Alysso-Sedetalia and Trifolio arvensis-Festucetalia ovinae (Koelerio-Corynephoretea), b – orders Corynephoretalia canescentis and Sedo acris-Festucetalia (Koelerio-Corynephoretea), c – alliance Agrostion vinealis (Molinio-Arrhenatheretea).

Abb. 3: Verbreitung der Sandtrockenrasengesellschaften in der Wald- und Waldsteppenzone des ukrainischen Flachlandes. a – Ordnungen Alysso-Sedetalia und Trifolio arvensis-Festucetalia ovinae (Koelerio-Corynephoretea), b – Ordnungen Corynephoretalia canescentis und Sedo acris-Festucetalia (Koelerio-Corynephoretea), c – Verband Agrostion vinealis (Molinio-Arrhenatheretea).

5. Discussion

5.1. Festuco-Brometea

The associations Stipetum pulcherrimae (SS1), Astragalo-Stipetum (SS2), and Poetum versicoloris (SS3) were originally assigned to the alliance Alysso-Sedion within the Sedo-Scleranthetea by ABDULOEVA (2002). The current analysis of their floristic composition (Table 1 in the Supplement) revealed a considerable representation of Festuco-Brometea character species, while the Koelerio-Corynephoretea species were less frequent. Based on the results of cluster analysis (Fig. 2), I propose the inclusion of these associations into the Festuco-Brometea class, namely into the alliance Festucion valesiacae in the order Festucetalia valesiacae. Therefore these syntaxa must be considered as steppe communities and not sandy grassland communities. Further studies should be carried out to define their syntaxonomical position more precisely.

5.2. Koelerio-Corynephoretea

5.2.1. Alysso-Sedetalia

The associations Aurinio saxatilis-Allietum podolici (SS4) and Minuartio auctae-Festuce-tum pallentis (SS5), described by ONISHCHENKO (2001) on the territory of Nature Reserve "Medobory", showed considerable distinctiveness from other syntaxa. This can be explained by their unique floristic composition with such endemic species as Allium podolicum and Minuartia aucta. Floristically, the associations correspond well to the Alysso-Sedion alliance

(Koelerio-Corynephoretea; see Table 1 in the Supplement). Some common features with the three associations from the Festuco-Brometea class (Fig. 2) are indicative of the transitional nature of these syntaxa.

Thus, communities of *Alysso-Sedetalia* order (Fig. 3a, Fig. 4) are represented only by two associations in the plains region of Ukraine. Apparently, these two associations are endemic to West Podillya. Further inspection of limestone outcrops of this region, especially Tovtry Ridge, valleys of Dnister and its tributaries probably will make it possible to discover new localities of these communities. Taking into account the specific environmental characteristics of the territory may make possible the discovery of additional plant communities of this alliance new for science.



Fig. 4: Stand of the Aurinio saxatilis-Allietum podolici (order Alysso-Sedetalia) with Allium podolicum, Aurinia saxatilis, and Gypsophila paniculata in the National Nature Park "Podils'ki Tovtry" (A. Kuzemko, 08/2008).

Abb. 4: Bestand des Aurinio saxatilis-Allietum podolici (Ordnung Alysso-Sedetalia) mit Allium podolicum, Aurinia saxatilis und Gypsophila paniculata im Nationalen Naturpark "Podils'ki Tovtry" (A. Kuzemko, 08/2008).

5.2.2. Trifolio arvensis-Festucetalia ovinae

Communities on granite outcrops of forest and forest-steppe zones were assigned by DIDUKH & KONTAR (1998) and KONTAR (2000) to two orders, two newly described alliances and five new associations. Nevertheless, the floristic composition of all five associations is very similar (Table 1 in the Supplement), as confirmed by cluster analysis (Fig. 2). Thus, these syntaxa should be included in one alliance, *Hyperico perforati-Scleranthion perennis* from the order *Trifolio arvensis-Festucetalia ovinae*, and probably must be merged into one association *Thymo pulegioides-Festucetum ovinae*.

At present, few localities of this community are known, namely on granite outcrops within the Ukrainian crystalline shield (Fig. 3a and 5). Further research is needed to reveal any new localities of this association and to assess its overall distribution in Ukraine.

5.2.3. Corynephoretalia canescentis

Those communities described by VICHEREK (1972) in the Dnieper Valley under the name Veronico dillenii-Corynephoretum (FV8) can be included into the Corynephorion canescentis alliance without doubt. Regarding the communities from Polissky Nature Reserve (KC1) which were included into the Helichryso-Jasionetum association by VOROBYOV et al. (1998), I would surmise that their floristic composition is similar to Corniculario aculeatae-Corynephoretum canescentis communities. The comparison of these communities with phytocenoses of Helichryso-Jasionetum from Latvia (RŪSIŅA 2007) and Germany (BERG et al. 2001) make it clear that diagnostic species of this association (e.g. Cladonia rangiformis, Cetraria aculeata, Hypnum cupressiforme, Carex arenaria, and others) are absent in relevés from Polissky Nature Reserve. Probably, the distribution area of Helichryso-Jasionetum lies outside Ukraine.

The associations Artemisio campestris-Dianthetum borbasi (KC3) and Corynephoro-Silenetum tataricae (KC4) from Zhytomyr Polissya were included within the same cluster with the previous two syntaxa by the cluster analysis (Fig. 2). Thus, I assign these syntaxa to the above mentioned alliance, the first one as a synonym of Corniculario-Corynephoretum and a second one as a single association. However, their floristic composition indicates the transitional nature of these associations.

Stands of the *Corniculario-Corynephoretum* have been described from Dnieper Valley and some plots within Zhytomyr Polissya (Fig. 3b and 6). In my opinion, the association has wider distribution in the forest zone. There is need for further research to clarify the distributional range of the association *Corniculario-Corynephoretum*. In addition, it is possible that communities of other associations of this order occur on fluvioglacial sediments of Polissya.

5.2.4. Sedo acris-Festucetalia

The communities of Artemisio dniproicae-Sedetum sexangularis (SS11) were described by SCHEVCHYK et al. (1996) in Dnieper islands at the border of Kaniv Nature Reserve. He assigned them to the Corynephorion canescentis alliance. However, these communities differ considerably from other syntaxa of this alliance (Table 1 in the Supplement). Moreover, this syntaxon was merged together with the Festuco psammophilae-Koelerietum glaucae (FV5) from the Dnieper islands in the Kremenchig Reservoir (GAL'CHENKO 2006), the Veronico dillenii-Secalietum sylvestris (FV1b), and the Centaureo borysthenicae-Festucetum beckeri (FV2b) relevés from islands of the Kaniv Nature Reserve in the cluster analysis (Fig. 2). Based on the results of cluster analysis, all mentioned syntaxa were merged into one association within the Koelerion glaucae alliance with the name Veronico dillenii-Secalietum sylvestris as the oldest available name.

The other associations that were assigned to the class *Festucetea vaginatae* according to Ukrainian phytosociological tradition have to be included into the *Festucion beckeri* alliance. As can be seen from the cluster analysis, the communities from the Left-Bank forest-steppe that were delimited by DIDUKH & KOROTCHENKO (1996) as two different



Fig. 5: Stand of the *Thymo pulegioides-Festucetum ovinae* (order *Trifolio arvensis-Festucetalia ovinae*) with *Festuca ovina* and *Sedum sexangulare* in the Pivdenny Bug Valley (A. Kuzemko, 05/2006).

Abb. 5: Bestand des *Thymo pulegioides-Festucetum ovinae* (Ordnung *Trifolio arvensis-Festucetalia*

ovinae) mit Festuca ovina und Sedum sexangulare im Tal des Pivdenny Bug (A. Kuzemko, 05/2006).



Fig. 6: Stand of the Corniculario aculeatae-Corynephoretum (order Corynephoretalia canescentis) with Corynephorus canescens and Cladonia sp. in the Polissky Nature Reserve (A.Kuzemko, 06/2008).

Abb. 6: Bestand des Corniculario aculeatae-Corynephoretum (order Corynephoretalia canescentis) mit Corynephorus canescens and Cladonia sp. im Naturschutzgebiet Polissky (A.Kuzemko, 06/2008).

associations, Veronico dillenii-Secalietum sylvestris (FV1a) and Centaureo borysthenicae-Festucetum beckeri (FV2a), possess substantial resemblance (Fig. 2). The high level of similarity becomes apparent also in Table 1. Therefore, they should be considered as single association: Centaureo borysthenicae-Festucetum beckeri.

Floristic composition is very similar also between the associations *Chamaecytiso* ruthenicae-Festucetum beckeri (FV3) and Thymo pallasiani-Centauretum sumensis (FV4), which were described by SHEVCHYK & SOLOMAKHA (1996) in Kaniv Nature Reserve. In my opinion, they must be considered as synonyms, with Chamaecytiso ruthenicae-Festucetum beckeri being the oldest available name. They should be placed within the Festucion beckeri and not the Koelerion glaucae as was proposed in the original publication.

The syntaxonomical position of the *Cladonietum* from the *Koelerion glaucae* alliance, which was provisionally described as new by SHEVCHYK & POLISHKO (2000) from a sandy terrace of the Dnieper River, remains unclear. The name of the association is not valid according to the Art. 3g, ICPN. The *Cladonietum* was included in the same cluster as the two previous associations but floristical comparison of these syntaxa is troublesome as non-vascular plants were treated in FV6 but not in syntaxa FV3 and FV4. It is necessary to conduct further investigation to clarify the correct placement of this syntaxon.

The same applies to the *Poetum bulbosi* (SS12), which was described by the same authors and classified in a newly described alliance *Trifolio arvensis-Sedion* within the *Sedo-Scleranthetea* class. However, the floristic composition of this association is very similar to the earlier described association *Poo bulbosae-Caricetum colchicae* from Biruchy Island spit in the Azov Sea (Dubyna et al. 1994), though it differs from the latter in regional peculiarities caused by its distribution in the different zone. The syntaxonomic position of this association remains unclear. In its original diagnosis, it was assigned to the *Festucion beckeri*, but the character species of this alliance are absent in the available relevés of the association (see Table 1). Unfortunately, only three relevés were available, which is insufficient for comparative analysis and final decision on syntaxonomy of these communities.

The Poo bulbosae-Caricetum colchicae created a single cluster together with the Diantho borbasii-Agrostietum syreistschikovii (FV7). The later was described by VICHEREK (1972) from the Dnieper Valley and included by him into the Koelerion glaucae alliance within the class Festucetea vaginatae. The floristic composition of both syntaxa differs substantially (Table 1). Thus, they can be interpreted as separate associations. They are located separately from other syntaxa of both alliances (Koelerion glaucae and Festucion beckeri) in the cluster dendrogram (Fig. 2). Nonetheless, the Diantho borbasii-Agrostietum syreistschikovii should be included into the first alliance, and the Poo bulbosae-Caricetum colchicae into the second alliance based on their floristic composition (Table 1).

Thus, the results of the analysis showed considerable similarity of communities that were included in either of the two classes Festucetea vaginatae and the Koelerio-Corynephoretea by different authors in Ukraine. It is not surprising that the correlation of these syntaxa has repeatedly been discussed in literature. For instance, VICHEREK (1972) included the Koelerion glaucae alliance into the Festucetea vaginatae. Also CHYTRÝ & TICHÝ (2003) recognize the independence of the Festucetea vaginatae class. The same point-of-view is traditional in Ukrainian phytosociology and in some other European countries. However, in some large-scale overviews on the vegetation of Europe, the authors have presented opposing opinions. MUCINA (1997) considers Festucetea vaginatae class as syntaxonomic synonym of the Festuco-Brometea. A similar opinion is shared by RODWELL et al. (2002). DENGLER (2001, 2003) considers the Festucetalia vaginatae order as synonym of the Sedo acris-Festucetalia. I agree with the last point-of-view because this assignment is confirmed by the present analysis of floristic composition of syntaxa and the results of cluster analysis. Concerning the syntaxonomic position of the alliance Koelerion glaucae, I believe that it is more appropriate to assign this alliance to the Sedo acris-Festucetalia order instead of Corynephoretalia canescentis or Festuco-Astragaletalia arenarii.



Fig. 7: Stand of the Veronico dillenii-Secalietum sylvestris (order Sedo acris-Festucetalia) with Anthemis ruthenica and Sedum acre in the Psel River floodplain (A. Kuzemko, 05/2008).

Abb. 7: Bestand des Veronico dillenii-Secalietum sylvestris (Ordnung Sedo acris-Festucetalia) mit Anthemis ruthenica und Sedum acre in der Psel-Aue (A. Kuzemko, 05/2008).



Fig. 8: Communities of the alliance Festucion beckeri (order Sedo acris-Festucetalia, class Koelerio-Corynephoretea) with Secale sylvestre, Rumex acetosella, and Potentilla argentea in the Kolomak River floodplain (A. Kuzemko, 05/2008).

Abb. 8: Gesellschaften des Verbandes Festucion beckeri (Ordnung Sedo acris-Festucetalia, Klasse Koelerio-Corynephoretea) mit Secale sylvestre, Rumex acetosella und Potentilla argentea in der Kolomak-Aue (A. Kuzemko, 05/2008).

Communities of the *Sedo acris-Festucetalia* order are investigated mainly in the Dnieper Valley and in some plots in Zhytomyr Polissya (Fig. 3b, 7 and 8) but predominantly in protected areas. They need further detailed investigation and comparison with other syntaxa of this order that have been described from the steppe zone of Ukraine, where they show a considerable variability (more than 40 associations have been described, but probably a portion of them needs revision).

5.3. Molinio-Arrhenatheretea

The Galietalia veri order is probably one of the most disputable syntaxa in Europe. The alliance Agrostion vinealis was first described from the territory of Ukraine (SIPAYLOVA et al. 1985) and was subsequently included into the newly described order Poo-Agrostietalia vinealis (SHELYAG et al. 1985). Almost simultaneously MIRKIN & NAUMOVA (1986) published the name Galietalia veri. These names must be considered as synonyms (KUZEMKO & DZYUBA 2002). Galietalia veri is the correct name because the name Poo-Agrostietalia vinealis was not validly published according to the rules of the last edition of ICPN. Both orders appeared mainly in publications from the former USSR (STSEPANOVICH 2000, BULOKHOV 2001, YAMALOV et al. 2003; 2004, MARTYNENKO et al. 2005 and others). The majority of European phytosociologists do not agree with the existence of such syntaxa, but see RODWELL et al. (2002), who lists the order Poo-Agrostietalia vinealis with the two alliances Agrostio-Avenulion schellianae Royer 1991 and Agrostion vinealis within the Festuco-Brometea. I disagree with that a point of view because in the present investigation a considerable number of Molinio-Arrhenatheretea species were present in Agrostion vinealis stands (Table 1 in the Supplement) and it is impossible to ignore this fact in classification, whereas the *Festuco-Brometea* species group was noticeably weaker.



Fig. 9: Communities of the alliance Agrostion vinealis (order Galietalia veri, class Molinio-Arrhen-atheretea) with Galium verum, Rumex thyrsiflorus, and Achillea millefolium in the Dnieper River floodplain (T. Vinichenko, 05/2005).

Abb. 9: Gesellschaften des Verbandes Agrostion vinealis (Ordnung Galietalia veri, Klasse Molinio-Arrhenatheretea) mit Galium verum, Rumex thyrsiflorus und Achillea millefolium in der Dnepr-Aue (T. Vinichenko, 05/2005).

Finally, examination of synoptic table and cluster analysis results revealed considerable similarity of the Koelerio-Agrostietum vinealis, the Festuco valesiacae-Agrostietum vinealis and the Agrostio vinealis-Calamagrostietum epigeii associations, which had been considered as separate units for more than 20 years. I propose to consider these syntaxa as one association under the oldest available name Koelerio delavignei-Agrostietum vinealis.

Corynephoretea syntaxa is evidence of their transitional character.

The Euphorbio cyparissiae-Festucetum ovinae, which was originally described from the Desna River floodplain within the Agrostion vinealis (SHELYAG-SOSONKO et al. 1987), but later was transferred to the Corynephorion canescentis (SIPAYLOVA & SHELYAG-SOSONKO 1996), shows high similarity to the Molinio-Arrhenatheretea syntaxa (Fig. 2) as it contains many species of Molinio-Arrhenatheretea, particularly Deschampsia caespitosa, Festuca rubra, Prunella vulgaris, and Trifolium pratense with high constancy (Table 1). Therefore, I consider its original inclusion in the Koelerio-Corynephoretea as a mistake and suggest to place these stands in the Molinio-Arrhenatheretea. However, I refrain from further classifying these stands as they probably comprise a complex of several associations from different alliances and even orders.

The communities of *Agrostion vinealis* are known mostly from the Dnieper Valley and its left tributaries, at Right-Bank they occur only in lower and mouth parts of the Dnieper tributaries (Fig. 3c and 9). At the present stage of knowledge, these communities have been inventoried and investigated rather carefully.

5.4. Classification scheme

As a consequence of the results and the preceding considerations, I suggest the following new compact classification scheme of the dry grassland vegetation on sandy soils in the forest and forest-steppe zones of the plains region of Ukraine. It can be integrated into a modern classification system of European vegetation. Synonyms are indicated with "="; some narrow syntaxa, that were assigned to the broader syntaxa in the present paper are indicated with "incl.". Associations whose syntaxonomic position within the next higher syntaxon is doubtful are marked with "?".

FB Festuco-Brometea Br.-Bl. & Tx. ex Klika & Hada 1944

FB1 Festucetalia valesiacae Br.-Bl. & Tx. ex Br.-Bl. 1950

FB1.1 Festucion valesiacae Klika 1931

FB1.1.1 (?) Stipetum pulcherrimae Soó 1942

FB1.1.2 (?) Astragalo-Stipetum R. Knapp 1944 nom. inval.

FB1.1.3 (?) Poetum versicoloris Kukovitsa et al. 1992

KC Koelerio-Corynephoretea Klika in Klika & Novák 1941 (incl. Sedo-Scleranthetea Br.-Bl. 1955, Festucetea vaginatae Soó ex Vicherek 1972)

KC1 Alysso-Sedetalia Moravec 1967

KC1.1 Alysso-Sedion Oberdorfer & Müller in Müller 1961

KC1.1.1 Aurinio saxatilis-Allietum podolici Onishchenko 2001

KC1.1.2 Minuartio auctae-Festucetum pallentis Onishchenko 2001

KC2 Trifolio arvensis-Festucetalia ovinae Moravec 1967

KC2.1 Hyperico perforati-Scleranthion perennis Moravec 1967 (incl. Thymo pulegioidis-Sedion sexangularis Didukh & Kontar 1998, Poo compressae-Rumicion acetosellae Didukh & Kontar 1998)

KC2.1.1 Thymo pulegioides-Festucetum ovinae Oberdorfer 1957 (incl. Thymo pulegioidis-Sedetum sexangularis Didukh & Kontar 1998, Artemisio austriacae-Teucrietum chamaedrys Didukh & Kontar 1998, Sempervivo ruthenicae-Sedetum ruprechti Didukh & Kontar 1998, Vincetoxico hirundinariae-Rumicetum acetum sessellae Didukh & Kontar 1998,

Melico transsylvanicae-Sedetum ruprechti Kontar 1998)

KC3 Corynephoretalia canescentis Klika 1934

KC3.1 Corynephorion canescentis Klika 1931

KC3.1.1 Corniculario aculeatae-Corynephoretum canescentis Steffen 1931 nom. invers. propos. (incl. Veronico dillenii-Corynephoretum Passarge 1960, Helichryso-Jasionetum sensu Vorobyov et al. 1998 non Libbert 1940, Artemisio campestris-Dianthetum borbasi Yakushenko 2004) KC3.1.2 Corynephoro-Silenetum tataricae Libbert 1931

KC4 Sedo acris-Festucetalia Tx. 1951 nom. invers. propos. (= Festucetalia vaginatae Soó ex Vicherek 1972)

KC4.1 Koelerion glaucae Volk 1931

KC4.1.1 Diantho borbasii-Agrostietum syreistschikovii Vicherek 1972

KC4.1.2 Veronico dillenii-Secalietum sylvestris Shevchyk & Solomakha 1996 (incl. Artemisio dniproicae-Sedetum sexangularis Shevchyk & Solomakha in Shevchyk et al. 1996, Festuco psammophilae-Koelerietum glaucae sensu Gal'chenko 2006, non Klika 1931, Centaureo borysthenicae-Festucetum beckeri sensu Shevchyk & Solomakha 1996, non Vicherek 1972)

KC4.2 Festucion beckeri Vicherek 1972 (incl. Trifolio arvensis-Sedion Shevchyk & Polishko 2000)
KC4.2.1 Chamaecytiso ruthenicae-Festucetum beckeri Shevchyk & Solomakha 1996
(incl. Thymo pallasiani-Centauretum sumensis Shevchyk & Solomakha 1996, Cladonietum Shevchyk et Polishko 2000 nom. inval.)

KC4.2.2 Centaureo borysthenicae-Festucetum beckeri Vicherek 1972 (incl. Veronico dillenii-Secalietum sylvestri sensu Didukh & Korotchenko 1996, non Shevchyk & Solomakha 1996) KC4.2.3 (?) Poo bulbosae-Caricetum colchicae Dubyna et al. 1994 (incl. Poetum bulbosae Shevchyk & Polishko 2000)

MA Molinio-Arrhenatheretea R.Tx. 1937

MA1 Galietalia veri Mirkin & Naumova 1986 (= Poo-Agrostietalia vinealis Shelyag et al. 1985 nom. inval.)

MA1.1 Agrostion vinealis Sipaylova et al. 1985

MA1.1.1 Koelerio-Agrostietum vinealis (Sipaylova et al. 1985) Shelyag et al. 1987 (incl. Festuco valesiacae-Agrostietum vinealis Shelyag et al. in Shelyag et al. 1985, Agrostio vinealis-Calamagrostietum epigeii (Shelyag et al. 1981) Shelyag et al. 1985) MA? (?) Euphorbio cyparissiae-Festucetum ovinae Shelyag-Sosonko et al. 1987 (probably a mixture of several associations)

6. Conclusions

Dry grasslands on sandy soils have a considerable distribution in the forest and steppeforest zones of the plains region of Ukraine. While some of them (communities of the *Trifolio* arvensis-Festucetalia ovinae and Corynephoretalia canescentis) are similar to analogous syntaxa from central and western Europe, the others are characterized by the originality of their floristic composition. A significant part of their diagnostic species belongs to the Podolian (communities of Alysso-Sedion) and Pontic (communities of Koelerion glaucae and Festucion beckeri) centers of endemism. Communities of the Agrostion vinealis probably have no analogs in the vegetation of countries west of Ukraine. Their development seems to be conditioned by hydrological and geomorphological peculiarities of the floodplains of the Dnieper River and its tributaries as well as the regional climate.

Data on sandy dry grassland communities in Ukraine are still very scarce, despite their high diversity and the large size of the study area. Additionally, usage of varying plot sizes, inconsistent vascular plant nomenclature, and ignoring of non-vascular plants in some cases considerably impair the quality of available data. Finally, the absence of a common phytosociological database seriously hinders large-scale syntaxonomic studies in Ukraine, not only of dry grasslands. Presently, phytosociological data are scattered in various archives, theses, monographs, reports, and so on. These methodological and organizational problems need to be solved before the preliminary results of the present study can be replaced by a more complete inventory and consistent analysis of the sandy dry grasslands of Ukraine.

Acknowledgements

The author expresses sincere thanks to Dr. Jürgen Dengler for his idea and encouragement to prepare the article, to Dr. Nicholai Ermakov for his helpful advice in the course of preparing the publication and to Dr. Christian Dolnik for the possibility to present the research at the "Dry Grassland Meeting 2008". I am grateful to Dr. Tatiana Dzyuba for making available several literature sources with limited access and to Mrs. Tatiana Sukhomeilo for her assistance in editing the English version of the manuscript. Special gratitude to Dr. Solvita Rusina for scientific and linguistic editing of the manuscript and her valuable comments. Finally, I thank Curtis Björk for his kind improvement of the English usage.

Appendix A: Descriptions of associations used in this study and their sources

Syntaxon names are corrected orthographically if necessary.

Appendix A: Assoziationsbeschreibungen, die in dieser Arbeit Verwendung fanden, und deren Quellen. Syntaxonnamen wurden erforderlichenfalls orthografisch korrigiert.

Code number	Name of syntaxon in original publication	Syntaxonomic position according to original publication (alliance, order, class)	Source	Number of relevés	Region	Localities of relevés	Ecotope	Plot size (m²)
SS1	Stipetum pulcherrimae	Alysso-Sedion, Alysso- Sedetalia, Sedo- Scleranthetea	ABDULOEVA (2002)	13	Western forest-steppe	Pokuttia (Ivano- Frankovsk region)	chalky outcrops	no information given
SS2	Astragalo-Stipetum	Alysso-Sedion, Alysso- Sedetalia, Sedo- Scleranthetea	ABDULOEVA (2002)	19	Western forest-steppe	National Natural Park "Podilski Tovtry" (Khmelnitsk region)	stony outcrops	no information given
SS3	Poetum versicoloris	Alysso-Sedion, Alysso- Sedetalia, Sedo- Scleranthetea	ABDULOEVA (2002)	6	Western forest-steppe	National Natural Park "Podilski Tovtry" (Khmelnitsk region)	shaly clay	no information given
SS4	Aurinio saxatilis- Allietum podolici	Alysso-Sedion, Alysso- Sedetalia, Sedo- Scleranthetea	ONISHCHENKO (2001)	12	Western forest-steppe	Natural reserve Medobory	limestone outcrops	2-33
SS5	Minuartio auctae- Festucetum pallentis	Alysso-Sedion, Alysso- Sedetalia, Sedo- Scleranthetea	ONISHCHENKO (2001)	6	Western forest-steppe	Kremenetski hills	carbonate rocks	3–20
SS6	Artemisio austriacae- Teucrietum chamaedrys	Thymo pulegioidis-Sedion sexangularis, Festuco- Sedetalia, Sedo- Scleranthetea	DIDUKH & KONTAR (1998)	19	Right-Bank forest zone	Pivdenny Bug Valley (Vinnitsa region)	granite outcrops	no information given
SS7	Thymo pulegioidis- Sedetum sexangularis	Thymo pulegioidis-Sedion sexangularis, Festuco- Sedetalia, Sedo- Scleranthetea	DIDUKH & KONTAR (1998)	10	Right-Bank forest zone	Sluch Valley (Rivno region), Pivdenny Bug Valley (Vinnitsa region), Teteriv Valley (Zhytomyr region)	granite outcrops	no informatior given
SS8	Vincetoxico hirundinariae- Rumicetum acetosellae	Poo compressae- Rumicion acetocellae, Sedo-Sclerathetalia, Sedo-Scleranthetea	DIDUKH &KONTAR (1998)	21	Right-Bank forest zone	Hnylopiat' Valley (Zhytomyr region), Sluch Valley (Rivno region)	granite outcrops	no information given
SS9	Sempervivo ruthenicae-Sedetum ruprechti	Poo compressae- Rumicion acetocellae, Sedo-Sclerathetalia, Sedo-Scleranthetea	DIDUKH & KONTAR (1998)	15	Right-Bank forest zone	Sluch Valley (Rivno region), Pivdenny Bug Valley (Vinnitsa region), Teteriv Valley (Zhytomyr region)	granite outcrops	no information given
SS10	Melico transsylvanicae- Sedetum ruprechti	Thymo pulegioidis-Sedion sexangularis, Festuco- Sedetalia, Sedo- Scleranthetea	KONTAR (2000)	19	Right-Bank forest-steppe	Hnylij Tikych Valley, Ros' Valley (Cherkasy region), Pivdenny Bug (Vinnitsa region)	granite outcrops	no informatior given
SS11	Artemisio dniproicae- Sedetum sexangulare	Corynephorion canescentis, Corynephoretalia canescentis, Sedo- Scleranthetea	SHEVCHYK et al. (1996), GAL'CHENKO (2006), own data of author 1999	11	Middle Dnieper Valley	Kaniv Nature Reserve, Ros' river mouth (Cherkassy region), Kremenchuk Regional Landscape Park, (Poltava region)	alluvial deposits	25–?
SS12	Poetum bulbosi	Trifolio arvensis-Sedion, Trifolio arvensis- Festucetalia ovinae, Sedo-Scleranthetea	SHEVCHYK & POLISHKO (2000)	3	Left-Bank Dnieper Valley	Liplava forestry (Cherkassy region)	pinery terrace	no information given
KCI	Helichryso- Jasionetum	Corynephorion canescentis, Corynephoretalia canescentis, Sedo- Scleranthetea	VOROBYOV et al. (1998)	15	Polissky Nature Reserve	Polissky Nature Reserve (Zhytomyr region)	sandy hills with granite outcrops	no informatior given
KC2	Euphorbio cyparissiae- Festucetum ovinae	Corynephorion canescentis, Corynephoretalia canescentis, Sedo- Scleranthetea	SIPAYLOVA & SHELYAG-SOSONKO (1996)	11	Left-Bank forest zone	Desna floodplain (Chernigiv region)	high plots of the central and near-riverbed parts pf floodplain	no information given
кс3	Artemisio campestris- Dianthetum borbasi	Koelerion glaucae, Corynephoretalia canescentis, Koelerio- Corynephorenea	YAKUSHENKO (2004)	11	Riht-Bank forest zone	Zytomyr Polyssia (Zytomyr region)	disturbed sandy plots	no informatior given
KC4	Corynephoro- Silenetum tataricae	Koelerion glaucae, Corynephoretalia canescentis, Koelerio- Corynephorenea	Orlov & Yakushenko (2005)	3	Riht-Bank forest zone	Projected Korostyshiv National Nature Park (Zytomyr region)	slopes of sandy terrace	no information given
FV1a	Veronico dillenii- Secalietum sylvestris	Festucion beckeri, Festucetalia vaginatae, Festucetea vaginatae	DIDUKH & KOROTCHENKO (1996)	10	Left-Bank forest-steppe	Vorskla river Valley, islands of Dneprodzerzhinsk and Pechenizhsky reservoirs (Poltava region)	sandy terraces and alluvial deposits	100

Code number	Name of syntaxon in original publication	Syntaxonomic position according to original publication (alliance, order, class)	Source	Number of relevés	Region	Localities of relevés	Ecotope	Plot size (m²)
FV1b	Veronico dillenii- Secalietum sylvestris	Festucion beckeri, Festucetalia vaginatae, Festucetea vaginatae	Sнеvснук et al. (1996)	9	Middle Dnieper Valley	Kaniv Nature Reserve, islands (Cherkassy region)	sandy deposits on the tops of hills and downs	no information given
FV2a	Centaureo borysthenicae- Festucetum beckeri	Festucion beckeri, Festucetalia vaginatae, Festucetea vaginatae	DIDUKH & KOROTCHENKO (1996)	5	Left-Bank forest-steppe	Vorskla river mouth, islands of Dneprodzerzhinsk and Pechenizhsky reservoirs (Poltava region)	alluvial deposits	100
FV2b	Centaureo borysthenicae- Festucetum beckeri	Festucion beckeri, Festucetalia vaginatae, Festucetea vaginatae	SHEVCHYK et al. (1996)	5	Middle Dnieper Valley	Kaniv Nature Reserve, islands (Cherkassy region)	plain sections of the inter- ridge depressions in floodplaine	no information given
FV3	Chamaecytiso ruthenicae- Festucetum beckeri	Koelerion glaucae, Festuco-Astragaletalia arenarii, Festucetea vaginatae	SHEVCHYK et al. (1996)	4	Middle Dnieper Valley	Kaniv Nature Reserve, islands (Cherkassy region)	sandy terrace	no information given
FV4	Thymo pallasiani- Centauretum sumensis	Koelerion glaucae, Festuco-Astragaletalia arenarii, Festucetea vaginatae	SHEVCHYK et al. (1996), SHEVCHYK & POLISHKO (2000), POLISHKO (2005)	13	Middle Dnieper Valley	Kaniv Nature Reserve, islands (Cherkassy region)	sandy terrace	no information given
FV5	Festuco psammophilae- Koelerietum glaucae	Koelerion glaucae, Festuco-Astragaletalia arenarii, Festucetea vaginatae	GAL'CHENKO (2006)	8	Middle Dnieper Valley	Kremenchuk Regional Landscape park (Poltava region)	high plots of floodplain	16–100
FV6	Cladonietum	Koelerion glaucae, Festuco-Astragaletalia arenarii, Festucetea vaginatae	SHEVCHYK & POLISHKO (2000), POLISHKO (2005)	7	Middle Dnieper Valley	Chihyryn forestry (Cherkassy region)	pinery teracces	no information given
FV7	Diantho borbasii- Agrostietum syreistschikovii	Koelerion glaucae, Festuco-Astragaletalia arenarii, Festucetea vaginatae	VICHEREK (1972)	10	Lower and Middle Dnieper Valley	Truchaniv Island, Kyiv	alluvial deposits	16
FV8	Veronico dillenii- Corynephoretum	Corynephorion canescentis, Corynephoretalia canescentis, Sedo- Scleranthetea	VICHEREK (1972)	15	Lower and Middle Dnieper valley	Dnieper Valley (Kyiv region)		16
AV1	Koelerio-Agrostietum vinealis	Agrostion vinealis, Galietalia veri, Molinio- Arrhenatheretea	Phytocenotheka (Afanasiev 1949– 1972, Sipaylova 1974, 1976, Sitenko 1976), own data of author (1999, 2005)	61	forest and forest-steppe zones	Dnieper, Desna, Seim, Ros' & Teteriv floodplains	high sections of floodplain	16–100
AV2	Festuco valesiacae- Agrostietum vinealis	Agrostion vinealis, Galietalia veri, Molinio- Arrhenatheretea	Phytocenotheka (Afanasiev 1972, Sipaylova 1980, 1982, Iutina 1998, Fitsailo 1997, 1998), own data of author (2003, 2005)	22	forest and forest-steppe zones	Dnieper, Desna, Seim, Ros', Psel & Pivdenny Bug floodplains	high sections of floodplain	16–100
AV3	Agrostio vinealis- Calamagrostietum epigeii	Agrostion vinealis, Galietalia veri, Molinio- Arrhenatheretea	Phytocoenotheka (Afanasiev 1949– 1968, Sipaylova 1974–1982, Panchenko 1996– 1998), own data of author (1999, 2003, 2005)	28	forest and forest-steppe zones	Dnieper, Desna, Sudost', Ros', Oril', Vilshanka floodplains	high sections of floodplain	16–100

References

- ABDULOEVA, O. S. (2002): Do syntaxonomii kserophitnoi trav'yanystoi roslynnosti Zakhidnogo Lisostepu (Prydnistrovs'ke Podillya, Tovtrovyy kryazh, Kremenetsli hory) (On the syntaxonomy of xerophytic herbaceous vegetation of the Western forest-steppe (Near-Dnister Podillya, Tovtry reef, Kremenetski hills)) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 1(18): 124–144. Kyiv.
- & DIDUKH, Y. P. (1999): Luchno-stepova roslynnist' erodovalykh skhyliv Prydnistrov'ya (natsional'nyy pryrodnyy park "Podilski Tovtry") v aspekti ii okhorony (Meadow-steppe vegetation of the eroded slopes in Prydnistrovia (National Natural Park "Podilski Tovtry") in the aspect of its protection) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 3(14): 10–36. Kyiv.
- Berg, C., Dengler, J. & Abdank, A. (2001) [Eds.]: Die Pflanzengesellschaften Mecklenburg-Vorpommerns und ihre Gefährdung Tabellenband. Weissdorn, Jena: 341 pp.
- -, -, & ISERMANN, M. (2004) [Eds.]: Die Pflanzengesellschaften Mecklenburg-Vorpommerns und ihre Gefährdung – Textband. – Weissdorn, Jena: 606 pp.
- BULOKHOV, A. D. (2001): Travyanaya rastitel'nost' Yugo-Zapadnogo Nechernozem'ya Rossii (Herbal vegetation of South-West Non-Chernozem area of Russia) [in Russian, with English summary]. Izd-vo BGU, Bryansk: 296 pp.

- CHEREPANOV, S. K. (1995): Sosudistye rasteniya Rossii i sopredel'nykh gosudarstv (Vascular plants of Russia and neighbour countries) [in Russian]. Mir i Semya, St. Petersburg: 992 pp.
- CHYTRÝ, M. (2007) [Ed.]: Vegetace České republiky. 1. Travinná a keřičková vegetace (Vegetation of the Czech Republic. 1. Grassland and Heathland Vegetation) [in Czech, with English summary]. Academia, Praha: 526 pp.
- & TICHY, L. (2003): Diagnostic, constant and dominant species of vegetation classes and alliances of the Czech Republic: a statistical revision.
 - Folia Fac. Sci. Nat. Univ. Masaryk. Brun., Biol., 108: 1–231. Praha.
- DENGLER, J. (2001): Erstellung und Interpretation synchorologischer Karten am Beispiel der Klasse Koelerio-Corynephoretea. Ber. R.-Tüxen-Ges. 13: 223–228. Hannover.
- (2003): Entwicklung und Bewertung neuer Ansätze in der Pflanzensoziologie unter besonderer Berücksichtigung der Vegetationsklassifikation. – Arch. Naturwiss. Diss. 14: 297 pp. Galunder, Nümbrecht.
- DIDUKH, Y. P. & KONTAR, I. S. (1998): Syntaksonomiya roslynnosti vidslonen' krystalichnykh porid lisovoi zony Ukrainy. I. Klasy *Asplenietea trichomanes* ta *Sedo-Scleranthetea* (Syntaxonomy of granite outcrops vegetation in forest zone of Ukraine. I. *Asplenietea trichomanes* and *Sedo-Scleranthetea* classes) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A, 2(11): 62–90. Kyiv.
- & KOROTCHENKO, I. A. (1996): Stepova roslynnist' pivdennoi chastyny Livoberezhnogo Lisostepu Ukrainy. I. Klasy Festucetea vaginatae ta Helianthemo-Thymetea (steppe vegetation of the south part of Left-Bank forest-steppe of Ukraine. I. Festucetea vaginatae and Helianthemo-Thymetea classes) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A, 2: 56–63. Kyiv.
- DUBYNA, D. V. (2006): Vyshcha vodna roslynnist'. Roslynnist' Ukrainy (High aquatic vegetation. Vegetation of Ukraine) [in Ukrainian, with English summary]. Phytosociocentre, Kyiv: 534 pp.
- NEUHÄUSLOVA, Z., & SHELYAG-SOSONKO, Y. R. (1994): Coastal vegetation of the "Birjucij Island"
 Spit in the Azov Sea, Ukraine. Preslia 66: 193–216. Průhonice near Praha.
- -, -, DZYUBA, T. P. & SHELYAG-SOSONKO, Y. R. (2004): Klasyfikatsiya ta prodromus roslynnosti vodoym, perezvolozhenykh terytoriy ta aren Pivnichnogo Prychornomor'ya (Classification and prodrome of vegetation of reservoirs, floodlands and arenasof the Northern Black Sea Region) [in Ukrainian, with English summary]. Phytosociocentre, Kyiv: 200 pp.
- –, DZYUBA, Т. Р., NEUHÄUSLOVÁ, Z., SOLOMAKHA, V. А., TYSHCHENKO, O. V. & SHALYAG-SOSONKO, Y. R. (2007): Halofitna roslynnist'. Roslynnist' Ukrainy (Halophytic vegetation. Vegetation of Ukraine) [in Ukrainian, with English summary]. Phytosociocentre, Kyiv: 315 pp.
- GAL'CHENKO, N. P. (2006): Regional'nyy landshaftnyy park "Kremenchutski plavni" (Regional landscape park "Kremenchuk plavni") [in Ukrainian, with English summary]. In: Shelyag-Sosonko, Y. R. [Ed.]: Plant World. Natural Protected territories of Ukraine 5: 176 pp. Phytosociocentre, Kyiv.
- GONCHARENKO, I. V. (2003): Analiz roslynnogo pokryvu pivnichno-skhidnogo Lisostepu Ukrainy (The analysis of vegetation cover of the northeast forest-steppe of Ukraine). Ukr. Phytosociol. Col. Ser. A 1(19): 203 pp. Kyiv.
- GRYGORA, I. M., VOROBYOV, Y. O. & SOLOMAKHA, V. A. (2005): Lisovi bolota Ukrains'kogo Polissya (pokhodzhennya, dynamika, klasyfikatsiya) (Woody bogs of the Ukrainian Polissya (origin, dynamic, classification of vegetation)) [in Ukrainian]. Phytosociocentre, Kyiv: 515 pp.
- IGNATOV, M. S. & AFONINA, O. M. (1992) [Eds.]: Check-list of mosses of the former USSR. Arctoa 1: 1–85. Moscow.
- KONDRATYUK, S. Y., KHODOSOVTSEV, A. Y. & ZELENKO S. D. (1998): The second checklist of lichenforming, lichenicolous and allied fungi of Ukraine. Phytosociocentre, Kyiv: 180 pp.
- KONTAR, I. S. (2000): Syntaksonomiya roslynnosti vidslonen' krystalichnykh porid Lisostepu Ukrainy. II. Klasy *Festuco-Brometea* ta *Sedo-Scleranthetea* (Syntaxonomy of vegetation of granite outcrops forest-steppe of Ukraine. II. Classes *Festuco-Brometea* and *Sedo-Scleranthetea*) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 1(16): 16–28. Kyiv.
- KUKOVITSA, H. S., MOVCHAN, Y. I., SOLOMAKHA, V. A. & SHELYAG-SOSONKO, Y. R. (1992): Novyy syntakson Poetum versicoloris ass.nova stepiv Zakhidnogo Podillya (New syntaxon Poetum versicoloris ass. nova of the steppes of Western Podillya) [in Ukrainian, with English summary]. Ukr. Bot. J. 49(1): 27–30. Kyiv.
- -, -, & (1994): Syntaksonomiya luchnykh stepiv Zakhidnogo Podillya (The meadow-steppe syntaxonomy of the Western Podillya) [in Ukrainian, with English summary]. - Ukr. Bot. J. 51(2-3): 35-48. Kyiv.
- KUZEMKO, A. A. & DZYUBA, T. P. (2002): Syntaksonomichna struktura klasu *Molinio-Arrhenatheretea* R. Tx. 1937 rivnynnoi chastyny Ukrainy (Syntaxonomic structure of the *Molinio-Arrhenatheretea*

- R. Tx. 1937 class in the plain part of Ukraine) [in Ukrainian, with English summary]. In: DIDUKH, Y. P. [Ed.]: Y. D. Kleopov and modern botanic science. Materials of the readings dedicated to the centenary of Yu. D. Kleopov: 238–245. Phytosociocentre, Kyiv.
- MALYNOVS'KYY, K. A. & KRICHFALUSHIY, V. V. (2000): Vysokohirna roslynnist' (High mountain vegetation) [in Ukrainian, with English summary]. In: Vegetation of the Ukraine. Vol. 1 Phytosociocentre, Kyiv: 230 pp.
- MARINICH, A. M., PASHCHENKO, V. M. & SHYSHCHENKO, P. G. (1985): Priroda Ukrainskoy SSR. Landshafty i fiziko-geograficheskoe rayonirovanie (Nature of Ukrainian SSR. Landscapes and physiographic zoning) [in Russian]. Naukova Dumka, Kiev: 224 pp.
- & SHYSHCHENKO, P. G. (2005): Fizychna geografiya Ukrainy. Pidruchnyk (Physiography of Ukraine. Manual) [in Ukrainian]. - Znannia, Kyiv: 511 pp.
- MARTYNENKO, V. B., YAMALOV, S. M., ZHIGUNOV, O. Y. & FILINOV, A. A. (2005): Rastitel'nost' gosudarstvennogo prirodnogo zapovednika "Shul'gan-Tash" (The vegetation of the State Natural Reserve "Shul'gan-Tash") [in Russian]. Gilem, Ufa: 272 pp.
- MATUSZKIEWICZ, W. (2001): Przewodnik do oznaczania zbiorowisk roślinnych Polski (Guidebook for definition of plant communities of Poland) [in Polish]. Wydawnictwo Naukowe PWN, Warszawa: 537 pp.
- MIRKIN, B. M. & NAUMOVA, L. G. (1986): O vysshykh edinitsakh sintaksonomii ravninnykh glikofitnykh lugov evropeyskoy chasti SSSR (About high units of syntaxonomy of the plain glycophyte meadows of European Part of USSR) [in Russian, with English summary]. Bul. MOIP Biol. Dept. 91(2): 93–101. Moscow.
- MOSYAKIN, S. L. & FEDORONCHUK, M. M. (1999): Vascular plants of Ukraine. A nomenclatural checklist. – National Academy of Sciences of Ukraine, Kiev: 346 pp.
- MUCINA, L. (1997): Conspectus of Classes of European Vegetation. Folia Geobot. Phytotaxon. 32: 117–172. Průhonice near Praha.
- ONISHCHENKO, V. A. (2001): Roslynnist' karbonatnykh vidslonen' pryrodnogo zapovidnyka "Medobory" (Calcareous rock vegetation of the Medobory Natural Reserve) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 1(17): 86–104. Kyiv.
- ORLOV, O. O. & YAKUSHENKO, D. M. (2005): Roslynnyy pokryv proektovanogo Korostyshivs'kogo natsional'nogo pryrodnogo parku (Plant cover of projected Korostyshiv Nature National Park) [in Ukrainian, with English summary]. In: SHELYAG-SOSONKO, Y. R. [Ed.]: Plant World. Natural Protected territories of Ukraine 3: 180 pp. Phytosociocentre, Kyiv.
- Polishko, O. D. (2005): Syntaksonomiya roslynnosti dilyanky borovoi terasy Dnipra (Chyhyryns'ke lisnytstvo Cherkass'koi oblasti) (Syntaxonomy of vegetation of the pinery terrace area of the Dnieper (Chyhyryn forestry, Chercassy region)) [in Ukrainian, with English summary]. In: DIDUKH, Y. P. [Ed.]: Actual Problems of Botany and Ecology. Coll. Sci. Works 1: 163–176. Kyiv.
- RODWELL, J. S., SCHAMINÉE, J. H. J., MUCINA, L., PIGNATTI, S., DRING, J. & MOSS, D. (2002): The Diversity of European Vegetation An overview of phytosociological alliances and their relationships to EUNIS habitats. Rapp. EC-LNV 2002/054: 168 pp. National Reference Centre for Agriculture, Nature and Fisheries, Wageningen.
- RUDENKO, L. G. (2007) [Ed.]: Natsional'nyy atlas Ukrainy (National Atlas of Ukraine) [in Ukrainian]. DNVP "Kartografiya", Kyiv: 440 pp.
- RUPRECHT, E., SZABÓ, A., ÉNYEDI, M. Z. & DENGLER, J. (2009): Steppe-like grasslands in Transylvania (Romania): characterisation and influence of management on species diversity and composition. Tuexenia 29: 353–368. Göttingen.
- RÜSIŅA, S. (2005): Diagnostic species of mesophylous and xerophylous grassland plant communities in Latvia. Acta Univ. Latv., Earth Environ. Sci. 685: 69–95. Riga.
- (2007): Latvijas mezofito un kserofito zalaju daudzveidiba un kontaktsabiedribas (Diversity and Contact Communities of Mesophytic and Xerophytic Grasslands in Latvia) [in Latvian, with English summary].
 Latv. Veg. 12: 241 pp. Riga.
- SCHEVCHYK, V. L. & POLISHKO, O. D. (2000): Syntaksonomiya roslynnosti dilyanky borovoi terasy (Liplyavs'ke lisnytstvo Cheras'koi oblasti) (Syntaxonomy of pinery terrace vegetation (Liplavo forestry of Cherkassy region)) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 1(16): 67–89. Kiev.
- SHELYAG-SOSONKO, Y. R., SOLOMAKHA, V. A. & SIPAYLOVA, L. M. (1985): Novye sintaksony poymennykh lugov ravninnoy chasti Ukrainy (New syntaxa of floodplan meadow of the plain part of Ukraine) [in Russian]. Manuscript, deposited at Ukrainian Institute for Scientific and Technical Information (No. N 6525–B85), Kiev: 40 pp.

- SHEVCHYK, V. L. & SOLOMAKHA, V. A. (1996): Syntaksonomiya roslynnosti ostroviv Kruglyk i Shelestiv Kanivs'kogo pryrodnogo zapovidnyka (The syntaxonomy of vegetation Kruglyk and Shelestiv Islands of Kaniv Natural Reserve) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. B 1: 12–27. Kyiv.
- -, & VOYTYUK, Y. O. (1996): Syntaksonomiya roslynnosti ta spysok flory Kanivs'kogo pryrodnogo zapovidnyka (The syntaxonomy of vegetation and list of the flora of Kaniv Natural Reserve) [in Ukrainian, with English summary]. - Ukr. Phytosociol. Col. Ser. B 1: 120 pp. Kiev.
- SIPAYLOVA, L. M. & SHELYAG-SOSONKO, Y. R. (1996): Luchna roslynnist' zaplav richok rivnynnoi chastyny Ukrainy (The meadow vegetation of the flood-plain rivers of the plain part of Ukraine) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 1: 28–40. Kyiv.
- -, MIRKIN, B. M., SHELYAG-SOSONKO, Y. R. & SOLOMAKHA, V. A. (1985): Novi soyuzy Agrostion vinealis ta Festucion pratensis luchnoi roslynnosti (New alliances Agrostion vinealis and Festucion pratensis of meadow vegetation) [in Ukrainian, with English summary]. Ukr. Bot. J. 42(4): 13–18. Kyiv.
- SIRENKO, I. P. (1996): Creation a Databases for Floristic and Phytocenologic Researches. Ukr. Phytosociol. Col. Ser. A(1): 9–11, Kyiv.
- SOLOMAKHA, V. A. (1996): Syntaksonomiya roslynnosti Ukrainy (The Syntaxonomy of Vegetation of the Ukraine) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. A 4(5): 120 pp. Kyiv.
- STATSOFT, INC. (2005): STATISTICA for Windows. Version 7.0. URL: http://www.statsoft.com.
- STSEPANOVICH, Y. M. (2000): Ekolaga-flarystychny dyyahnaz sintaksonau pryrodnay travyanistay raslinnastsi Belarusi (Ecological and floristic diagnosis of syntaxa of natural herbaceous vegetation of Belarusa) [in Belorussian]. Kamtat, Minsk: 140 pp.
- VERNANDER, N. B., GOGOLEV, I. N., KOVALISHIN, D. I., NOVAKOVSKY, L. Y., SIRENKO, N. A. & TIUTIUNNIK, D. A. (1986): Priroda Ukrainskoy SSR. Pochvy (Nature of Ukrainian SSR. Soils) [in Russian]. Naukova Dumka, Kiev: 216 pp.
- VOROBYOV, Y. O., BALASHOV, L. S. & SOLOMAKHA, V. A. (1997): Syntaksonomiya roslynnosti Polis'kogo pryrodnogo zapovidnyka (The syntaxonomy of vegetation of the Polesie Natural Reserve) [in Ukrainian, with English summary]. Ukr. Phytosociol. Col. Ser. B 1(8): 128 pp. Kyiv.
- VICHEREK, J. (1972): Die Sandpflanzengesselschaften des unteren und mittleren Dnieprstromgebietes (die Ukraine). Folia Geobot. Phytotaxon. 7: 9–46. Praha.
- Weber, H. E., Moravec, J. & Theurillat, J.-P. (2000): International Code of Phytosociological Nomenclature. 3rd edition. J. Veg. Sci. 11: 739–768. Uppsala.
- YAKUSHENKO, D. (2004): Nova asotsiatsiya psamofil'noi roslynnosti zi skhodu Zhytomyrs'kogo Polissya (New assotiation of psammophilous vegetation from Zhytomyr Polissia Eastern Part) [in Ukrainian, with English summary]. Visnyk L'viv Univ. Biol. Ser. 35: 95–101. L'viv.
- Yamalov, S. M., Filinov, A. A. & Solomeshch, A. I. (2003): Ostepnennye luga poryadka *Galietalia* veri Mirkin et Naumova 1986 na Yuzhnom Urale (The steppe-meadows of the *Galietalia veri* Mirkin et Naumova 1986 order in the South Ural) [in Russian, with English summary]. In: Vegetation of Russia 5: 62–80. St. Petersburg.
- –, MARTYNENKO, V. B., GOLUB, V. B. & BAISHEVA, E. Z. (2004): Prodromus rastitel'nykh soobshchestv respubliki Bashkortostan (The prodrome of plant communities of Bashkortostan Republic) [in Russian].
 – Gilem, Ufa: 64 pp.

Anna Kuzemko
National dendrological park "Sofiivka"
National Academy of Sciences of Ukraine
12a Kyivska str.
Uman' 20300
UKRAINE
anya meadow@mail.ru

Managing editor: Jürgen Dengler Manuscript received 07.11.2008, accepted 21.02.2009.

Differential species of associations (D; including potential character species) are shaded in dark grey and character species (Ch) of higher syntaxa in light grey. Taxa within differential species blocks are arranged by their constancy in the differentiated syntaxon, while taxa within character species blocks are arranged by their total constancy. Non-diagnostic "Other species" are sorted alphabetically. "?" denotes unknown

Tab. 1 (Teil 1): Stetigkeitstabelle von Sandtrockenrasen in der Wald- und Waldsteppenzone des ukrainischen Flachlandes. Differenzialarten von Assoziationen (D; einschließlich möglicher Assoziationskennarten) sind dunkelgrau hinterlegt, Charakterarten höherer Syntaxa hellgrau. Innerhalb der Differenzialartenblöcke sind die Arten nach fallender Stetigkeit im jeweiligen Block, ansonsten nach fallender Gesamtstetigkeit angeordnet. Nicht diagnostische sonstige Arten sind alphabetisch angeordnet. Das "?" steht für fehlende Daten.

Class		estue				_						Koel	erio	-Coi	rynep	phor	etea			_			_				Arrhe there		
Final designation of association (see classification scheme in the Section 5.4)	FB1.1.1	FB1.1.2	FB1.1.3	KC1.1.1	KC1.1.2			KC2.1.1				KC3.1.1		KC3.1.2	KC4.1.1		VC413	7.1.*			KC4.2.1		20,000	KC4.2.2	KC4.2.3		MA1.1.1		MA?
Original designation of association (see Table 1)	SSI	SS2	SS3	SS4	SSS	9SS	SS7	SS8	6SS	SS10	FV8	KCI	KC3	KC4	FV7	SS11	FV5	FV1b	FV2b	FV4	FV3	FV6	FV2a	FV1a	SS12	AVI	AV2	AV3	KC2
Number of relevés Plot sizes [m²]	13	19	6	2-	3-	19	10 ?	21 ?	15 ?	19 ?	15 16	15 ?	11 ?	3			8 16– 100	9	5 ?	13 ?	4	7	5 100	10 100		16-	22 16– 100	16–	
Cumulative species richness of vascular plants	83	59	55			79	60	102	77	60	14	71	78	28	32			38	27	83	30	48	62	68	25		149		32
Mean species richness of vascular plants	?	?	?	19	13	16	10	13	14	8	4	15	10	20	21	16	14	12	12	22	10	10	23	17	17	18	25	23	?
Treatment of non-vascular plants $(Y = yes; N = no)$	N	N	N	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	N	Y	N	N	N	N	N	N	N
Cumulative species richness of non-vascular plants	?	?	?	18	11	19	10	16	16	15	8	13	2	1	7	?	?	?	?	3	?	10	?	?	?	?	?	?	?
Mean species richness of non-vascular plants	?	?	?	3	2	4	3	3	3	3	4	3	1	1	4	?	?	?	?	0	?	6	?	?	?	?	?	?	?
D Stipetum pulcherrimae																													
Stipa pulcherrima Jurinea arachnoidea	90 50	•	•		•	•		8.08	•	0.0	٠				9	33	27	7	19			39	*			•	*	•	٠
Securigera varia	50		10	17				5	7	•				8	8	8	8); (4						*		14		
Lembotropis nigricans	30	١.					9.5				e*:								12										
D Astragalo-Stipetum																					8				92				
Chamaecytisus austriacus		90	10											6	93	35	82	27	83	12	32	32	37	10		2			
Stipa pennata Teucrium montanum		90		100	17					0.0	•	20% 0.00			38 85	9	99 01		100		8	13 82	33		8		21	12 0	
Alyssum rostratum		70	i											9	8	5	9	9	16				÷				8		
Astragalus monspessulanus		70															,						×						
Picris hieracioides		70	10	0.0						5					33	35			6			94	0			3			
D Poetum versicoloris	20		00																										
Poa versicolor Iris hungarica	30 10		90 50				•	5	13	11	9.5		•		5.5 50	23 00	8 2	3.5	27	8	27	84	2		32		*		*
Stipa capillata			50	8	17									is ja		0					100	92			100				
Ch Festuco-Brometea	0.53		_			29700	0.400	ustrali		1000	0.000			425	308	.856	201	A15	.est.	107	750		455	95	500	67)	301	600	
Salvia verticillata	70	90	70	8		5			7							9	,												
Teucrium chamaedrys	90		70			79	10		7		٠				•	3				8	25	à	*	*	•	٠		٠	٠
Euphorbia cyparissias	50				33		50			16				8	20	27	85	11	i.	38	75	14	15	85	2.		14	•	71
Elytrigia intermedia Bothriochloa ischaemum	50		90	8	17	32 21	10	14	33				9	15.	33	3		3.	3	8	4						5		
Inula ensifolia	30		10		17									1		10							10						
Festuca valesiaca			50		17	5				26						18	100			14	a			10		5	91	11	*
Medicago falcata agg.	10		30					5	13				9	67													14		
Koeleria cristata	30		50	50							٠			33		-	8	•			4	12				7	5	٠	•
Stachys recta Campanula sibirica	50	50 30	*	33	17 33	37	20	10	13	5 . 5	S* (5			1	12	27	97	*		15	87	1	20	**	35	12	*	*	*
Eryngium campestre		50				5	10	5			•					9							40	20			27		
Allium sphaerocephalum	17.5	70				53	10									16			1	8	ž.							÷	
Hieracium cymosum	30	10		25									×			15						į.	40						
Thalictrum minus		10		25	17	5	٠		13		٠	٠		•							•			•				•	
Agrimonia eupatoria Filipendula vulgaris	10	10	10.60		٠	11		5	13	5	•					3	:	59		•			13	٠	٠	5	14 23	7	51
Aurinia saxatilis	10	5.5	10	67			3.5			74					6.	98	8	35 34		10	88	10	35	1.5	88				
Anthyllis macrocephala	10			8	•									13		2				,		ex.			9	2	5		*
Hieracium echioides		30				32																14		10					
Festuca rupicola	10	10								5						3	3						${}^{\circ}$	100	٠	٠			٠
Fragaria viridis Astragalus onobrychis	10		30	8	17	•	:		•		1			:	83	85	35		11	1.5	82	87		10	*		9	•	10
Carex humilis	10			8	17									29	114	-													
Achillea collina	1763		50		,											7							1						
Bupleurum falcatum	10	30													1														
Helictotrichon desertorum	10				17	•	•				•					2	•										•	•	
Aster amellus	10	50	10		17											- 1		्							٠		•	٠	٠
Hypericum elegans Centaurea scabiosa	10	30					•	3.5	50.0					•	10	88	82	82	3.5	12	3	85	85		85	15	25	*	1.5
D Aurinio saxatilis-Allietum podolici	10	50	•			•	•				•													100					
Allium podolicum				58					16	37				1.0	24				9.4	18		9.				×			
Cardaminopsis arenosa				50												9		11											
Thymus ×oblongifolius				50							٠	•				10				39	4	1	100			12		٠	
Cephalaria uralensis Sacali libaratis subsp. intermedium	10		•	25 25			•	2.40						12	100	12	*	8	<u></u>	10	10	10	95		8		*	÷	*
Seseli libanotis subsp. intermedium D Minuartio auctae-Festucetum pallentis	10			23											99			*	•	*		100			**			•	٠
Minuartia aucta					100	١																							
Festuca pallens					83									i.	10	32	: :	14		10		:: ::	10	100					į.
Alyssum turkestanicum					50	5				•												×		10					
Asplenium ruta-muraria				25	50	١.										1			•							٠			٠
Ch Alysso alyssoidis-Sedion				reco																									
Acinos arvensis	20		10			16		10	16		٠		18	-		-	3						19					٠	*
Asperula cynanchica Allium montanum	30 10		10	42			20	19		•			*	8	62	22	87	0 .	27	12	.:t	(đ	10	100	**	5.5 1.5	85		*
Ch Hyperico perforati-Scleranthion perennis	10		50	12	50			1,														•						•	•
Hylotelephium maximum agg.	10	1000		33		53	20	52	73	74		13			10.	27		22	20	8									
Centaurea rhenana		20	10			74								100								ī.					·	Ŷ.	
Festuca ovina								57				53			3.	Þ						÷					9		91
Poa compressa			٠	50	٠			38					64		,	2									•	2		7	
Steris viscaria Sempervivum ruthenicum				33	17		10 10	48	60 67	21		7	9	•	53	35		8	10	8	25	87	G.				14		
Thymus pulegioides agg.				33	1/	74		19		5	•			:	83 82	88 62	SE .		85	31)/* 82	83 82	33	15	88 82	8	18		
Allium oleraceum								19								10			•	10			9	8			9		
Melica transsilvanica	10			50												9													
Phleum phleoides	10			•			19.05	5	33			7		•	6	9				31		97		٠	٠		•	٠	
D Corynephoro-Silenetum tataricae														-										•					
Scabiosa ochroleuca	10	30	10	17													7			15			40		ं	*		20	
Silene tatarica Ch Correspondentian agreements														100		18	ः	11			10	•	40		٠	•		29	
Ch Corynephorion canescentis Jasione montana								14				80	45		10		99	-		46		57	20			2	,		
Corynephorus canescens											100			:		8	: :	:: :						10	1	3	*	7	100
Thymus serpyllum					17									67							,							٠	

Table 1 (part 2)

ass	Bi	ome		-		_						noe.	erto		rynep	nor	0000000										Arrh there	
nal designation of association (see classification heme in the Section 5.4)	FB1.1.1	FB1.1.2	FB1.1.3	KC1.1.1	KC1.1.2			KC2.1.1				KC3.1.1		KC3.1.2	KC4.1.1		VCA13				KC4.2.1		VCATT	KC4.2.2	KC4.2.3		MA1.1.1	
riginal designation of association (see Table 1)	SSI	SS2	SS3	SS4	SSS	9SS	SS7	SS8	6SS	SS10	FV8	KCI	KC3	KC4	FV7	SS11	FV5	FV1b	FV2b	FV4	FV3	FV6	FV2a	FV1a	SS12	AVI	AV2	AV3
Diantho borbasii-Agrostietum syreistschikovii									_						_	_								_		_		
ammophiliella muralis						•			i						80					•			20			25	9	7
erniaria polygama		19	12			-	•		18						70	×								10	33	5	9	18
ochia laniflora			0.7										*	•	70		•	•	•		•		20	Ċ	•			•
econax subconica		112	89	33								٠	•		60			٠	٠	٠		٠			٠	٠	٠	
h Koelerion glaucae												7	21		50		20		20	15		1.4		10			-	
lene borysthenica enothera rubricaulis	2.5	85	32	2.5	98	185	85	1.5	3.5	95	100	7	21 9	33	50	36	38		20	15		14	8 1	10	*	11	9	*
antago arenaria												7	9				13		20							3		
Chamaecytiso ruthenici-Festucetum beckeri	105	9.5	200	10.5	335	300	365		8.5	100	515.	1/1		100						(65)	0.	330	100		150		6.0	270
hamaecytisus ruthenicus		-		8	17	12		10	0			7	27	33				22	. 1	85	100	86	÷				14	
rucedanum oreoselinum	27	334				333			7				18					33		54			20					
entaurea sumensis			27	2.7	10.5	25	et.		8.5	85						12				54	25							
arex ericetorum	8.					132	34	10	82	19		7								31	25		Ÿ				2	
rinea pseudocyanoides		12					9.													23		43				*		
dsatilla pratensis	1	\mathbb{S}_{k}				5			Ü	5			7	7		ů.			.	38	4	72				÷		
Centaureo borysthenicae-Festucetum beckeri																					7			ÿ.				
ianthus platyodon		9				36			į.							(8)			9	×			80	50		٠		
nchusa gmelinii				3					ì					•			•		•	•			60	8888			٠	
entaurea pseudomaculosa		9	9	94	•	9	16		į,			ů,				×		٠	٠	٠		÷	40				14	
chillea micrantha	8.7	127	D.	B.	82	i d	•	82	135			1.5	•	8.	101		•	Ċ	12.		•		40		*	*	*	•
naria dulcis	74	83	34	8		14				15	15	92				12							60			ï		
ochia prostrata	10	0	()			1		10	×	100	2			٠						٠	•	÷	20	1000		*	٠	٠
agopogon borysthenicus			4		•		•					•	•	•				٠		٠	•	•	20					
ysimum canescens		19	19	3			٠				3	٠	٠			×	٠	•	٠	٠			40	10000	٠	٠	٠	
seli tortuosum			O.		*		•						•	ं	•			•		Ċ	Ċ	ė.	20	20	•	*	•	•
rinea charcoviensis	8.3	302	33					•	9.2		•	٠		•					٠	٠	٠		60	*	٠		٠	
olygonum arenarium Symus ×tschernjajevii	5.7	8.5	8.5	8.5	9.5		85	10.5	82	83	3	0	3.5	100			13	*	12			14	60 40		6		5	4
gropyron dasyanthum		-			50.		212	•		12													40					•
renia montana		33.	10.5	3.t		39	8.	58	12	18	8.5	3.5		1.0	*	3.5		3.5	8	38	•	•	+0	40	*	*	*	*
alium ruthenicum											•							7	•	•	•		20	40	•	•	•	•
perula graveolens						124			57														20					
anthus eugeniae	150	150		50	500	560	516	65	20	173	10		100	7.0	66	50	10	220	48	500	100			20	55	85	107	100
berna behen																								20				
h Festucion beckeri	17	0.0		515	0.5					- 12	100			150.					15)		(2)	200		(Internal		170		-
estuca beckeri																18		66	100	92	100	29	60	20		11		14
symus pallasianus																							40					
necio borysthenicus	100	114	3	12	0.5	100	100	10								÷	÷			15			40		÷			
Poetum bulbosi	- 65			3/2		95	100	8	133	100	-	8	34		S	200						•		10	Ö	90	ř.	3
nisantha tectorum						5		10		16								22		8				10	100			
renaria uralensis																9									100			
h Festuco-Sedetalia acris																												
elichrysum arenarium	10	30	10	352		02	32	19	17		90	67	55		40	36	63	** 1	40	31	·	14	100	100	2	10	9	11
peleria glauca												7	18		100									10				7
naria genistifolia	30	10	30		100	80	32	-		100		47	9										20		10	÷		
paragus officinalis		10		8		10	8	10		- 85	100		9	3	40		13					14				20	3	29
eracium umbellatum						32		14	13				9							23		43	20			28		4
ianthus borbasii		117	35	117			35		1				100		100					46	1574	1				52		39
cale sylvestre																	50				50		60	100				
arex colchica		5.	8.			11			12							36	75	44	40	8				30		8		21
agopogon ucrainicus		104				-				94	13	7			50		38		100	15			20	20				18
entaurea borysthenica		13		39													13	11	60	54	25	43	20	30				
uphorbia seguieriana	- 3	23	29	3.					99	34					40	18	50	11		23			100	70			٠	
hondrilla juncea		100	15	10			8.5		2.5	10		45	9			9	25			46			40	20	33		*	7
pidium densiflorum		94	94	:		10		8	92	114					·.			66				Ų.	v		100	(4)		
n Koelerio-Corynephoretea				_																								
mex acetosella		98	×			42	20	67	27		47	67			80					31		29	80				41	57
dum acre	10	97	10	100	67	37	60	14	27	53	•														100		27	39
temisia campestris		17			67	84	10	57	40				100	100	100	100	13	44	100	23	٠		100	90	٠	8	18	11
ifolium arvense				1		63	60	14	13			7	18			63		•	٠	8		14	20	10	100		18	4
dum sexangulare	- 5	17	15			63	100	24		32					50	100			20	٠	٠		¥	10		7	5	4
a bulbosa	9.	97	21	**		•		10		21	7		9		100		*	33	*	8	*	14		40	100		14	
rniaria glabra		95	37	34			40	5	59	4		7	9	1			٠		4				20	10	2.	7		7
ronica dillenii	1.7		27	100					0.5		100	7	14					88	40		25	*				•	•	•
themis ruthenica					•					1	•					18		55	•		•		40	20	100	3	•	7
deranthus annuus		13					20	10	7			27			70				*		*			*		5	5	
tentilla arenaria	30		30	75	50	10	1	V.	12	15		1.5	3				15	0		69		43	3		2	2	•	
osotis micrantha	1.5	10	33	-				83			47		9		80	9	٠		20		•	٠		٠		2		٠
rastium semidecandrum	5.5	3.5	82	88	135	100	83	8.5	82	8	3.5	*	1.5		40	10		*	3	1.5	18	1			100		14	7
ntaurea diffusa		34	8,				10	10							40		38	•			•		20	٠		5	5	11
leranthus perennis	12	9	2	100	16	*	10	10				2	27	1 ·	40			•		*			*		•	. 2		٠
rastium arvense					1	•				3.5					•	9	13		•		•	•		•		3	9	
		14	7.				10		3.		٠	*	٠	٠	*			•			*	*	**	*		8	5	
		95	1	100	**	15	8	1.5	10	:		**	1	100	100	•	*	•	ř	*	1	*	ं		67	5	5	7
arysaspis campestris				100		100			34		47			٠	100			•			•				67	٠		
ianthus deltoides trysaspis campestris rophila verna		-				16		-	1.0																			
arysaspis campestris rophila verna rabidopsis thaliana	54 54	84		837		16	100	5	13	3.5		3		*											100	*		
arysaspis campestris rophila verna rabidopsis thaliana vronica verna	54 54 57					16			13	10	*	3.		*		9	*			*		14			100			
arysaspis campestris rophila verna rabidopsis thaliana pronica verna dum reflexum								5	13		*		9			9	*	*		•		14			100			
arysaspis campestris rophila verna rabidopsis thaliana vronica verna						16 5	10 10	5	13		*	•	9	•		9						14	•		100			

continued on separate sheet

Zu Kuzemko: Dry grasslands on sandy soils in the forest and forest-steppe zones of the plains part of Ukraine: present state of syntaxonomy Table 1 (part 3): Synoptic table of the dry grassland vegetation of sandy soils in the forest and forest-steppe zones of the plains region of Ukraine. Differential species of associations (D; including potential character species) are shaded in dark grey and character species (Ch) of higher syntaxa in light grey. Taxa within differential species blocks are arranged by their constancy in the differentiated syntaxon, while taxa within character species blocks are arranged by their total constancy. Non-diagnostic "Other species" are sorted alphabetically. "?" denotes unknown values.

Tab. 1 (Teil 3): Stetigkeitstabelle von Sandtrockenrasen in der Wald- und Waldsteppenzone des ukrainischen Flachlandes. Differenzialarten von Assoziationen (D; einschließlich möglicher Assoziationskennarten) sind dunkelgrau hinterlegt, Charakterarten höherer Syntaxa hellgrau. Innerhalb der Differenzialartenblöcke sind die Arten nach fallender Stetigkeit im jeweiligen Block, ansonsten nach fallender Gesamtstetigkeit angeordnet. Nicht diagnostische sonstige Arten sind alphabetisch angeordnet. Das "?" steht für fehlende Daten.

104200000°	Br	estue ome	tea		_					-	-	Koei	erio		ryne _l	phor	etea										Arrh ther	
Final designation of association (see classification scheme in the Section 5.4)	FB1.1.1	FB1.1.2	FB1.1.3	KC1.1.1	KC1.1.2			KC2.1.1				KC3.1.1		KC3.1.2	KC4.1.1		61134	NC4:1:2			KC4.2.1			KC4.2.2	KC4.2.3		MA1.1.1	
Original designation of association (see Table 1)	SS1	SS2	SS3	SS4	SS5	9SS	SS7	SS8	6SS	SS10	FV8	KCI	KC3	KC4	FV7	SS11	FV5	FV1b	FV2b	FV4	FV3	FV6	FV2a	FV1a	SS12	+	AV2	AV3
Ch Agrostion vinealis																												
Agrostis vinealis	10		¥		Ε,		28		**	28	28	27			100	9		33		23		14		10				100 9
Carex praecox Koeleria delavignei	- 5	10	10		*	5					59 12		9				•			8					:: ::	70 67	14 27	75 21
Rumex thyrsiflorus					*6	**	*3	•					-			18										34	18	43
Euphorbia virgultosa Amoria montana		20					•	5	10		•	•	14	67	•	18		11	370	31	•			٠		10	9 23	39
Ch Molinio-Arrhenatheretea							*	3	•	40	•					•		•	•	•			•			13	43	11
Achillea millefolium	40	40		33				57		16	10	2	55	100		18		55		15	50	14	20	10				46 9
Plantago lanceolata Agrostis tenuis	91	70	12	1	\$	21	30 10	33 29	7	200	20		9	67	60	27						43	•	30	33	54 11	55 9	54 5
Taraxacum officinale		- 6	10	17			10	5		- 10						9						29				8	27	7 9
Rumex acetosa	81	ΑS	•		*	11	×		٠	•	•	7		33		٠			٠	15	25	٠				8	23	7
Carex hirta Poa pratensis	10 21	10			13	58 VS	10	14 19	8	11	18 18	7	55		1.50	•					25		•		1	3 20		21 1 39 3
Festuca pratensis	-	•		8	17	6	61	•	•	5	•		45		(*)			٠								11		21
Festuca rubra Lotus corniculatus		*	7	20		21		5		5		13	2					65.55			820					33	32 55	14 9
Stellaria graminea										-		7	-			9											36	11
Medicago lupulina		70	Ŷ.	è	20		*	5	6	5	40	40	40		(4)											7	27	7
Amoria repens Arrhenatherum elatius	6	50 70	*	17	5 E		55	10	*8	10 10	10 12	13 13	53 23	1.5				•	•						8	16	14	32 9 11
Dactylis glomerata	*		*	•	40	9	23	*:	•	5	•9	45	49		•			٠			•		•			7	9	14
Alopecurus pratensis Deschampsia cespitosa				•		٠		•	٠	•		•			•	9	٠	•		٠	•		•	•		11 2	9	32
Leucanthemum vulgare		-											e e						2.5						,	2		21 1
Prunella vulgaris		43				·		17	43	i.			23												1	11		25 7
Trifolium pratense Veronica chamaedrys	1	- 62 VI	12			5		8	10	10	10	7	50 20								25					8	23	11 3
Potentilla anserina		*						•	•	•	•	• 5	41														9	11 3
Stellaria hippoctona Allium angulosum					•	٠	•	•	٠	•	•	•			٠				•	٠						2	5	18 21
Allium angulosum Centaurea jacea	41	*5		*	**	*		**	•	•	•	•	63			•			•		•	•	•			23 18	9 14	21 14
Cerastium holosteoides		V			es Vi		13	*	20		8	10	25													10	14	11
Coccyganthe flos-cuculi Leontodon autumnalis	43	55	15	*5	*5	80	•	*	13	10	*0	55	1	٠	3.*3 0.52	(3 * ()	(1.00) (100)			•		•	0.0			8	9	7 14
Ranunculus acris		*	**							-			63													18	36	18
Phleum pratense		٠	٠			٠		٠	٠	٠						•	•									7	5	11
Plantago major Vicia cracca		- 20		*	**			•		- 60			80						•				20			7	5	7 21
Rumex crispus		- 33			27	10	ò	ä		43			-														5	14
Anthoxanthum odoratum Briza media	10	- 63 - 33	*	٠	:5 :0	50	*3	*	*3	53	•35 55	•	9	•	650 660	93 . 93					99 . 9	٠	(*) (*)	•	•	8.5	14	
Ranunculus polyanthemos										-	6	1	1													2	27	
Agrostis gigantea		•	•	•	2	*					•	•				٠	•	٠	٠		٠	٠				2		14
Carex lachenalii Galium boreale	- 41					10			44	6	•		#E		(30)		•				360	•			54	3		11
Gratiola officinalis					2	2	22		;;	8		23	27													10		18
Potentilla erecta	-	10	35	10	10	*		*	•	50	13	10	10		٠	•	200									2		4
Ranunculus repens Inula britannica		- 6				-							8					•	•						3	3 15		4
Juncus atratus		10	-		10		**		*	101	*3		10													2		11
Poa palustris Amoria hybrida	¥	77	77	•	20	20	83			21	•	•3	41		•		•		•				•			5 2		4 18
Mentha arvensis	, a	9	20	9	2	34		20	99	10	20		29		Tes			:*: :::	•						1	2		4
Widespread species of dry grassland and dry ruder															22										٠.	5.2	-	
Galium verum Poa angustifolia			70 10		**	74 27	10	29 5	7	21 11	**	7	9		80	18 18		11 22	•	46 31			20	10	33			21 46 5
Potentilla argentea		500 40	30		20			57			53. 45		91				25			23			40					46
Verbascum lychnitis	30		10	17 8	- 5			10 38			10	13 4	18			27 9		•			100			20		5	18	7
				0		3		30	21				64				13 13				25	/1		30				
						80			7	•			73		10	05		22		54	75	86		40		43	27 18	100
Hieracium pilosella Calamagrostis epigeios Echium vulgare	10	50	30					10	7 27	16	•3 •3	27	18	33			13					86		40		43 3	18 23	
Calamagrostis epigeios Echium vulgare Artemisia absinthium	10	50	30			5	10	24	7 27 53	16 16		27 ·	18 27	33 33			13 13						40		33	43 3	18 23 14	100
Calamagrostis epigeios Echium vulgare	10	50				5		24 14	7 27	16 16 53	•3 •3	27	18 27	33		18	13		•						33	43 3	18 23 14	
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria	. 10	50				5 84	10	24 14 10	7 27 53 60 13 27	16 16 53 5		27	18 27 45	33 33 100		18 36	13 13 13 13	22		23			40 60	40		43 3 15 11	18 23 14 23 36	100 18 7
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis	. 10	** ** **		25		5 84	10	24 14 10 57	7 27 53 60 13 27 7	16 16 53 5		27	18 27 45	33 33 100	•	18 36	13 13 13 13	22 44	40	23	•		40 60			43 3 15 11	18 23 14 23	100 18 7
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma	10	** ** **		25	17	5 84	10 40	24 14 10 57	7 27 53 60 13 27	16 16 53 5		27	18 27 45	33 33 100		18 36	13 13 13 13	22 44 44	40	23 15			40 60	40		43 3 15 11 9 8 7	18 23 14 23 36 5	100 18 7
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum	. 10	** ** **		25	17	5 84 21 16 26	10 40	24 14 10 57 10 14 29	7 27 53 60 13 27 7	16 16 53 5		27	18 27 45	33 33 100		18 36	13 13 13 	22 44 44 22	40 40 80	23 15 46 8	25	14 57	40 60	40		43 3 15 11 0 8 7 11 5	18 23 14 23 36 5 23 27	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis	. 10	** ** **	30	25	17	5 84 21 16 26	10 40 20	24 14 10 57 10 14	7 27 53 60 13 27 7	16 16 53 5		27	18 27 45	33 33 100		18 36	13 13 13 	22 44 44 22	40 40 80	23 15 46	25	14 57	40 60	40		43 3 15 11 0 8 7 11 5	18 23 14 23 36 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum		** ** **	30	25	17	5 84 21	10 40 20	24 14 10 57 10 14 29	7 27 53 60 13 27 7 27 13 7	16 16 53 5		27 	18 27 45 14 45 18 18 9	33 33 100 			13 13 13 		40 40 80		25	14 57	40 60	40		43 3 	18 23 14 23 36 5 23 27 32 14	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria	. 10		30	25	33	5 84 21	10 40 20 20 30 10	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7	16 16 53 5		27 	18 27 45	33 33 100 			13 13 13 		40 40 80		25	14 57	40 60	40		43 3 15 11	18 23 14 23 36 5 23 27 32 14	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens	. 10		30	25	17	5 84 21	10 40	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7	16 16 53 5		27 	18 27 45 14 45 18 18 9	33 33 100 			13 13 13 		40 40 80		25	14 57	40 60	40		43 3 	18 23 14 23 36 5 23 27 32 14	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species			30	25	33	5 84 21	10 40 20 20 30 10	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40	40	. 1000	43 3 15 11	18 23 14 23 36 5 23 27 32 14	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana			30	25	33	5 84 21	10 40 20 20 30 10	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7	16 16 53 5		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40 60	40	. 1000	43 3 15 11	18 23 14 23 36 5 23 27 32 14	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis			30	25	33	5 84 21	10 40 20 20 30 10	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40	. 40	. 1000	43 3 15 11	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum			30	25	33	5 84 21	10 40 20 20	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40	. 40	. 1000	43 3 15 11	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis			30	25	33	5 84 21	10 40 20 20 30 10	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40	. 40	. 1000	43 3 15 11	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria			30	25	33	5 84 21	10 40 20 20	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40 60 40 		. 1000	43 3 15 11 9 8 7 111 5 20 3 7 2	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria			30	25	33	5 84 21	10 40 20	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40 		. 1000	43 3 15 11 9 8 7 111 5 20 3 7 2	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria			30	25	33	5 84 21	10 40 20	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13		40 40 80		25		40 60 40 		. 1000	43 3 15 11 9 8 7 111 5 20 3 7 2	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biemis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris			30	25	33	5 84 21	10 40 20	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 27 13 7 7	16 53 5 		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13 	·			25		40 		. 1000	43 3 15 11 0 8 7 11 5 20 3 7 2	18 23 14 23 36 5 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca			30	25	33	5 84 21	10 40 20	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 27 13 7 7	16 16 53 5		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13 13 	·	40 40 80		25		40 		. 1000	43 3 15 11 9 8 7 111 5 20 3 7 2	18 23 14 23 36 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Aselepias syriaca Asplenium septentrionale Asplenium trichomanes			30	25	33	5 84 21 · · · · · · · · · · · · · · · · · ·	10 40 20	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 	16 16 53 5		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13 13 13 	·			25		40 60 40 20 20 		. 1000	43 3 15 11 0 8 7 11 5 20 3 7 2	18 23 14 23 36 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Arthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium trichomanes Astragalus sulcatus			30	25	33	5 84 21 · · · · · · · · · · · · · · · · · ·	10 40 	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 	16 16 53 5		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13 13 	·			25		40 		. 1000	43 3 15 11 0 8 7 111 5 20 3 7 2	18 23 14 23 36 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Arthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium trichomanes			30	25	33	5 84 21 · · · · · · · · · · · · · · · · · ·	10 40 	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 	16 16 53 5		27 	18 27 45 14 45 18 18 9 55	33 33 100 		18 36	13 13 13 13 13 	·			25		40 60 40 20 20 		. 1000	43 3 15 11 0 8 7 11 5 20 3 7 2	18 23 14 23 36 23 27 32 14 18 5	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Astragalus sulcatus Beckmannia eruciformis Betula pendula Bromopsis inermis			30	25	33	5 84 21	10 40 	24 14 10 57 10 14 29 10 14	7 27 53 60 13 27 7 	16 16 53 5 		27 	18 27 45	33 33 100 		18 36	13 13 13 13 13 	·			25		40 60 40 20 20 		. 1000	43 3 15 11 9 8 7 111 5 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Aljuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium trichomanes Astragalus sulcatus Beckmannia eruciformis Betula pendula Bromopsis inermis Bromus japonicus			30	25	33	5 84 21 · · · · · · · · · · · · · · · · · ·	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 	16 16 53 5 		27 	18 27 45	33 33 100 			13 13 13 13 13 	·			25		40 60 40 20 20 		. 1000	43 3 15 11 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium septentrionale Asplenium septentrionale Bromopsis inermis Bromus japonicus Bromus mollis Bromus secalinus			30	25	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 	16 16 53 5 		27 	18 27 45	33 33 100 			13 13 13 13 13 	·			25		40 60 40 20 20 		333	43 3 15 11 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Arctium nemorosum Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium septentrionale Asplenium septentrionale Asplenium septentrionale Bromopsis inermis Bromus japonicus Bromus secalinus Bromus sequarrosus Bromus squarrosus Bromus squarrosus Bromus squarrosus			30	25	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 		27 	18 27 45	33 33 100 			13 13 13 13 				25		40 60 40 20 20 		. 1000	43 3 15 11 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Arctium nemorosum Artemisis vulgaris Asclepias syriaca Asplenium septentrionale Asplenium septentrionale Asplenium septentrionale Asplenium septentrionale Bromopsis inermis Bromus japonicus Bromus mollis Bromus squarrosus Campanula bononiensis			30	25	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 	16 16 53 5 		27 	18 27 45	33 33 100 			13 13 13 13 				25		40 60 40 20 20 		333	43 3 15 11 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Astragalus sulcatus Beckmannia eruciformis Betula pendula Bromopsis inermis Bromus japonicus Bromus secalinus Bromus secalinus Bromus squarrosus Campanula persicifolia Campanula rapunculoides			30	25 	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 13 13 13 13 13 13 13 13 13 1			23 15 46 8	25		40 60 40 20 20 		333	43 3 15 11 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea nobilis Achillea retacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Astragalus sulcatus Beckmannia eruciformis Beckmannia eruciformis Beromus japonicus Bromus squarrosus Campanula bononiensis Campanula persicifolia Campanula rapunculoides Carduus acanthoides			30	25	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 			23 15 46 8	25		40 60 40 20 20 		333	43 3 15 11 20 3 7 2	18 23 14 23 36	100 ·
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium richomanes Astragalus sulcatus Beckmannia eruciformis Beromopsis inermis Bromus japonicus Bromus squarrosus Campanula bononiensis Campanula persicifolia Campanula rapunculoides Carduus acanthoides Carduus acanthoides Carac contigua			30	25 	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 13 13 13 13 13 13 13 13 13 1			23 15 46 8	25		40 60 40 20 20 		333	43 3 15 11 20 3 7 2	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea nobilis Achillea nobilis Anthemis subtinctoria Alyaga genevensis Aljuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium trichomanes Beschmannia eruciformis Beromopsis inermis Beromopsis inermis Bromus secalinus Bromus secalinus Bromus seculinus Bromus squarrosus Campanula bononiensis Campanula persicifolia Campanula rapunculoides Carduus acanthoides Carduus acanthoides Carex contigua Chenopodium album			30	25 	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 	16 16 53 5		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 			23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 20 3 	18 23 14 23 36	100 ·
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea nobilis Achillea nobilis Anthemis subtinctoria Alyaga genevensis Aljuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium richomanes Astragalus sulcatus Beckmannia eruciformis Betula pendula Bromus japonicus Bromus secalinus Bromus secalinus Bromus squarrosus Campanula bononiensis Campanula persicifolia Campanula rapunculoides Carex contigua Chenopodium album Chrysaspis aurea	500	10	30	25 	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 	16 16 53 5 · · · · · · · · · · · · · · · · ·		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 			23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 20 3 	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea nobilis Achillea nobilis Anthemis subtinctoria Alyaga genevensis Aljuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Astragalus sulcatus Beckmannia eruciformis Betula pendula Bromopsis inermis Bromus secalinus Bromus secalinus Bromus sequarrosus Campanula bononiensis Campanula persicifolia Campanula prapinculoides Carduus acanthoides Carex contigua Chelidonium majus Chenopodium album Chrysaspis aurea Cichorium intybus			30	25 	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7 	16 16 53 5		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 			23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 20 3 	18 23 14 23 36	100 ·
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Oother species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Asstragalus sulcatus Beckmannia eruciformis Betula pendula Bromopsis inermis Bromus secalinus Bromus squarrosus Campanula bononiensis Campanula persicifolia Campanula persicifolia Campanula prayunculoides Carex contigua Chelidonium majus Chenopodium album Chrysaspis aurea Cichorium intybus Clematis recta Clinopodium vulgare	500	10	30	25 ·	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 · · · · · · · · · · · · · · · · ·		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 	22 · 44 44 22 · · · · · · · · · · · · ·		23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 200 3 	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Asstragalus sulcatus Beckmannia eruciformis Betula pendula Bromus sponicus Bromus squarrosus Campanula bononiensis Campanula persicifolia Chelidonium majus Chenopodium album Chrysaspis aurea Cichorium intybus Clematis recta Clinopodium vulgare Convallaria majalis	500	10	30	25 ·	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 · · · · · · · · · · · · · · · · ·		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 	22 · 44 44 22 · · · · · · · · · · · · ·		23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 20 3 	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium trichomanes Asstragalus sulcatus Beckmannia eruciformis Betula pendula Bromus sponicus Bromus squarrosus Campanula bononiensis Campanula persicifolia Campanula persicifolia Campanula persicifolia Campanula persicifolia Campanula persicifolia Campanula persicifolia Carduus acanthoides Carex contigua Chelidonium majus Chenopodium album Chrysaspis aurea Cichorium intybus Clematis recta Clinopodium vulgare	500	10	30	25 ·	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 · · · · · · · · · · · · · · · · ·		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 	22 · 44 44 22 · · · · · · · · · · · · ·		23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 200 3 	18 23 14 23 36	100
Calamagrostis epigeios Echium vulgare Artemisia absinthium Berteroa incana Vicia tetrasperma Vincetoxicum hirundinaria Conyza canadensis Oenothera biennis Veronica spicata Hypericum perforatum Elytrigia repens Artemisia austriaca Phalacroloma annuum Genista tinctoria Anthericum ramosum Artemisia marschalliana Other species Achillea nobilis Achillea setacea Agropyron pectinatum Ajuga genevensis Ajuga orientalis Allium waldsteinii Anthemis subtinctoria Apera spica-venti Arctium nemorosum Aristolochia clematitis Artemisia vulgaris Asclepias syriaca Asplenium septentrionale Asplenium trichomanes Astragalus sulcatus Beckmannia eruciformis Betula pendula Bromopsis inermis Bromus japonicus Bromus squarrosus Campanula bononiensis Campanula persicifolia Chelidonium majus Chenopodium album Chrysaspis aurea Cichorium intybus Clematis recta Clinopodium vulgare Convallaria majalis Convolvulus arvensis	500	10	30	25 ·	33	5 84 21	10 40 	24 14 10 57 10 14 29 10	7 27 53 60 13 27 7	16 16 53 5 · · · · 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · 5 5 · · · · · 5 5 · · · · · 5 5 · · · · · 5 5 · · · · · · 5 5 · · · · · · 5 5 · · · · · · · 5 5 · · · · · · · · 5 5 ·		27 · · · · · · · · · · · · · · · · · · ·	18 27 45	33 33 100 			13 13 13 13 	22 · 44 44 22 · · · · · · · · · · · · ·		23 15 46 8	25		40 60 40 20 20 		333	43 3 . 15 11 . 0 8 7 111 5 200 3 	18 23 14 23 36	100

continued on the back side

Class		estue			_	_						Koel	erio	-Coi	ynep	ohor	etea										Arri	linio hena retea	1 -
Final designation of association (see classification scheme in the Section 5.4)	FB1.1.1	FB1.1.2	FB1.1.3	KC1.1.1	KC1.1.2			KC2.1.1				KC3.1.1		KC3.1.2	KC4.1.1		KC413	7.1.7			KC4.2.1		20100	KC4.2.2	KC4.2.3		MA1.1.1		
Original designation of association (see Table 1)	SSI	SS2	SS3	SS4 I	SSS I	9SS	SS7	SS8 I	6SS	SS10	FV8	KC1	KC3	KC4	FV7 I	SS11	FV5	FV1b	FV2b	FV4	FV3 I	FV6	FV2a	FV1a	SS12 I	-	AV2 N	T	2000
Dianthus pseudosquarrosus					17				<u>.</u>				9										9						
ryopteris carthusiana chinops sphaerocephalus	10	10		•	•						:: ::	•	•				•	•	•	•				e e	•	2	•		
quisetum arvense quisetum pratense					•		•						9)4 33											20	23 9	21 4	
rigeron acris ryngium planum							i.		115			2	9		i.	27	. 25		8		e e		20			2 10	14 14	11	
uonymus europaea uphorbia virgata				8				5	7										•							. 2			
alcaria vulgaris	10	10	10			5		es es		33	87		:		:: ::				•	•							:	•	
allopia convolvulus estuca pseudovina				8		5					•	13	: ::					•	•		•			10				4	
ragaria vesca rangula alnus					17			10	7	S4 S4	(i)			i v	82 84							14 14		7		2		7	
ialium aparine ialium campanulatum		٠	٠	67			•	ě			8			9		٠		•		•	•		٠			2	٠	4	
Galium exoletum	10			8			2	. 5	:: ::		8					,		•											
ilechoma hederacea iypsophila paniculata					17				2		2			13	94 18				•	8	3		20			5	14		
Iieracium vaillantii Iierochloe odorata					•		•								8	9		11	•	8	50					3	•		
lumulus lupulus lylotelephium telephium agg.							•			5	i.	•			60					•			•	•		25		7 22	
uncus bufonius amium album										8.	8.	. 7			lit es			2			31						9	4	
amium purpureum					•				7			7			9					•		•				•	•		
inaria vulgaris uzula pallidula				8			10				:		9							8	25	14		10		2	5		
Aelandrium album Aelilotus albus								er 15	7			:: ::	9	33	: :	9		: :		•	0	14	20	10		2	5		
Aelilotus officinalis Ononis arvensis	30	10	880	8	17	5				0.5	8.		2* 52	13 10	15	*	3.8 10	8.6 (12	## 100	0.	8		*	8. 32	16	*	5	*	
Orchis coriophora			,		٠										a	8	ě			ě	×	×		8	·	3		4	
Priganum vulgare Persicaria maculata	•		•	•	•		20		16			*	•	•				•	:	•	•	•		:		·	5	7	
etasites spurius Hlomoides tuberosa					•	•		3. 3.	3. 2	5	31 31	•		3	2	9	25		20			3		9			14		
impinella saxifraga inus sylvestris (juv.)					•	•	•	5				7 67	45	67			1			23		86				•	9 5	3	
lantago media oa nemoralis	٠	٠	٠	17	è			. 14	27	8	÷	•						9	•	•	•	•	9			10	32	4	
olygala vulgaris												3		17												3	23		
olygonatum odoratum olygonum aviculare	10			8			: :			9			9						•	8		14			67		•	18	
olypodium vulgare otentilla impolita					•		•	5	7	5	•					9		•			•	14		•	67	5	. 23	7	
otentilla reptans runus spinosa	. 10				82.0			0.7	7	e e	i d				11	10	4						20				5	4	
teridium aquilinum								5				13						3									•		
yrus communis Quercus robur (juv.)	10				•	5	2	5	7	9		7 7	18	28		10 14						14					5		
anunculus pedatus eseda lutea	10	10			•		10	5												•	•		•	•					
hamnus cathartica hinanthus aestivalis	810		•	17		2	8	5	87	10	11		8	85	ill lo	2	et.		•	10 12	(† 10	ė.	e e			7			
hinanthus serotinus hinanthus vernalis							•				٠						1.5									5	5 27	٠	
osa canina									13	5					,													14	
umex confertus alix acutifolia (juv.)			•	•							•	•				9			20		•		•	•		3	9	7 7	
alvia pratensis aponaria officinalis	10							8	11		8	12					30				25 25			10			14 5		
enecio erucifolius enecio jacobaea									2			7		. 33		9	25				4		40						
etaria viridis								5			17	13	9			8													
ideritis comosa ieglingia decumbens	30	10				•	:		3				18			:			•	•	•		•	•		2			
ilene eugeniae ilene nutans	10			25	17				22	9		•	9	2	31 32					8								8	
olidago virgaurea pergularia rubra						5		5				47 7	18						•	15				•			:	. 4	
piraea media anacetum vulgare	•	•				5		19	16	95				3.5	e e	45			*	*			20					. 32	
hymus marschallianus	10						2		2	11		ii ii								•									
hymus moldavicus ragopogon major		10 10	30 10																	15	•							4	
rifolium alpestre rifolium medium					•	5		5							•	•	•	•			•			•		•	5		
erbascum densiflorum		10 10	5000	8				43	8	9	8	::			12		13										٠	ě	
erbascum nigrum erbascum thapsus			10									•	9										40	30			•		
'eronica incana 'eronica officinalis			50	58		5	•			•	•	7	9						•	•	•			•					
/iola arvensis /iola matutina				8		11	10	5	7	82		47	9	33	6		8			8	25				:	: Q	14	4	
Non-vascular plants:																													
Ch Alysso-Sedetalia bietinella abietina	?	?	?	25	17	11	12		7	21						?	?	?	?		?		?	?	. ?	2	?	?	
Fortella tortuosa	?	?	?	50	17											?	?	?	?		?		?	?	?	?	?	?	
Camptothecium lutescens Phytidium rugosum	?	?	?	17 17	17 17		19 12	34 35	:: :::		e e	94	es es		8	?	?	?	?		?		?	?	?	?	?	?	
ncalypta streptocarpa incalypta vulgaris	?	?	?	8	17 17		114		73. 13.					14	: :	?	?	?	?	*	?		?	?	?	?	?	?	
seudoleskeella catenulata Ditrichum flexicaule	?	?	?	33	. 33				33		1	8	12	89		?	?	?	?		?		?	?	?	?	?	?	
Ch Trifolio arvensis-Festucetalia ovinae	1		-		33			•	3.						9.5	e e	-	f	f	đ	1		1		•	1	•		
lypnum cupressiforme irimmia laevigata	?	?	?	25	17			19 29				•				?	?	?	?		?	•	?	?	?	?	?	?	
leofusceliaa pulla 'anthoparmelia somloensis	?	?	?			37	10	14 38	27		:			3.5		?	?	?	?		?	÷	?	?	?	?	?	?	
ledwigia ciliata	?	?	?			16	30	33	40	•						?	?	?	?		?		?	?	?	?	?	?	
ryum argenteum armelia sulcata	?	?	?	8		11	30		7 13							?	?	?	?	e e	?		?	?	?	?	?	?	
ladonia coniocraea rimmia pulvinata	?	?	?			32		10	16	37						?	?	?	?	•	?		?	?	?	?	?	?	
h Koelerio-Corynephoretea	9	9	9			-	20	40	27	16	67	72	26	67	100	9	9	9	9	0	9	1.6	9	9	9		9	9	
olytrichum piliferum Teratodon purpureus	?	?	?		100	47	60	48 38	60	47	67	7	45		100	?	?	?	?	8	?		?	?	?	?	?	?	
ortula ruralis Eladonia pyxidata	?	?	?	58	17	32 11		5	7		67				50 10	?	?	?	?	*	?	57	?	?	?	?	?	?	
ladonia foliacea ladonia arbuscula	?	?	?	(*)	34)3	74	9		7						90 50	?	?	?	?	8	?	14	?	?	?	?	?	?	
ladonia rangiformis	?	?	?		120	12 . (4)	33 33	14	2%		67				40	?	?	?	?		?	14	?	?	?	?	?	?	
ladonia furcata leurozium schreberi	?	?	?	50% 80%	302 761	5	3	5 10	7		: :	7				?	?	?	?	*	?	86 86	?	?	?	?	?	?	
ladonia mitis ladonia phyllophora	?	?	?		59.0 95.0	(a)	3	10	39 33		9. 1	2				?	?	?	?	. 8	?	100 86	?	?	?	?	?	?	
ladonia fimbriata	?	?	?	100	330		8		**	8.	34		8.	18	(i)	?	?	?	?	*	?	86 71	?	?	?	?	?	?	
olytrichum commune acomitrium canescens	?	?	?	986 9867	92 () 848	8	12 14	3: 3:	%: %:	15	33		*			?	?	?	?	*	?		?	?	?	?	?	?	
eltigera praetextata t her species	?	?	?		300	25			10			17				?	?	?	?		?		?	?	?	?	?	?	
ladonia coccifera	?	?	?					6						ij.		?	?	?	?	Ÿ	?		?	?	?	?	?	?	
ladonia rangiferina icranum scoparium	?	?	?					10								?	?	?	?		?		?	?	?	?	?	?	
Grimmia ovalis	?	?	?	740			20		84	11			38	10	5	?	?	?	?		?	1	?	?	?	?	?	?	

Grimmia ovalis Homalothecium sericeum Polytrichum juniperinum

(SS10/5), Trinia multicaulis (SS4/8).

Spesies which present only in one column with low constancy (Designation of original syntaxa/percentage constancy).

Vascular plants: Acer tataricum (SS10/11), Adonis vernalis (SS1/10), Allium paczoskianum (FV1a/10), Allium scorodoprasum (FV1a/10), Alnus glutinosa (j.) (KC1/7), Alyssum calycinum (SS4/8), Alyssum gmelinii (SS1/10), Amorpha fruticosa (j) (AV3/7), Anchusa officinalis (AV2/5), Anemonastrum narcissiflorum (SS2/10), Anemone sylvestris (SS1/10), calycinum (SS4/8), Alyssum gmelinu (SS1/10), Amorpha fruticosa (1) (AV3/1), Anchusao officinalis (AV2/5), Anemonastrum narcissiflorum (SS2/10), Anemone sylvestris (SS1/10), Angelica sylvestris (AV3/4), Antennia dioica (KC1/7), Anthemis arvensis (SS1/19), Anthyllis scoiwereckii (SS5/17), Arenaria serpyllifolia (SS6/5), Artemisia abrotanum (AV1/8), Artemisia scoparia (KC1/7), Aruncus vulgaris (KC1/7), Asparagus polyphyllus (FV1a/10), Asparagus tenuifolius (FV5/25), Astragalus austriacus (SS1/10), Astragalus cicer (AV2/5), Athyrium filix-femina (SS8/5), Artiplex sagittata (SS4/8), Bidens tripartita (AV3/11), Bromopsis riparia (SS1/10), Bromus arvensis (FV1a/20), Calluna vulgaris (KC1/2), Campanula cervicaria (AV2/5), Campanula glomerata (KC1/7), Campanula patula (SS8/5), Capsella bursa-pastoris (AV3/4), Cardus crispus (SS4/8), Carex buxbaumii (AV1/2), Carex melanostachya (SS9/7), Carex michelii (AV3/4), Carex muricata (AV2/18), Carex panicea (AV3/7), Carex vulpina (AV3/4), Carlina biebersteinii (SS1/10), Centaurium erythraea (AV2/5), Cerastium glomeratum (AV2/14), Cerasus fruticosa (SS1/30), Chamaecytisus blockianus (SS1/10), Chamaecytisus bodolicus (SS1/10), Chrysaspis dubia (SS1/19), Cirsium arvense (AV3/11), Cirsium palustre (AV3/4), Carex melanostachya (SS2/10), Chamaecytisus podolicus (SS1/10), Centaurium erythraea (SS2/10), Chrysaspis dubia (SS1/19), Cirsium arvense (AV3/11), Cirsium palustre (AV3/4), Contaurium (AV3/4), Carex materia (SS2/10), Caretaurium erythraea vulgare (AV2/5), Clematis integrifolia (SS4/8), Kadenia dubia (AV1/5), Cornus mas (SS3/10), Crataegus leiomonogyna (SS4/8), Cynoglossum officinale (SS9/7), Dactylorhiza incarnata (AV1/2), Dactylorhiza majalis (AV2/5), Dianthus armeria (AV3/4), Dianthus campestris (AV3/4), Dianthus carthusianorum (SS8/19), Dianthus membranaceus incarnata (AV12), Dactylorhiza majatis (AV25), Dianthus armeria (AV34), Dianthus campestris (AV34), Dianthus carthustanorum (SS819), Dianthus membranaceus (SS611), Digitalis grandiflora (SS616), Digitaria sanguinalis (FV614), Dryothus, mas (FV6129), Epilobium montanum (AV25), Eragrostis minor (SS123), Eremogone saxatilis (FV48), Erysimum hieracifolium (SS119), Erysimum pannonicum (SS417), Euonymus verrucosa (SS97), Euphrasia stricta (AV25), Festuca polesica (KC113), Galium album (SS230), Galium palustre (AV344), Galium rubioides (AV112), Geranium pratense (AV344), Geranium purpureum (SS97), Geranium sanguineum (FV48), Glechoma hirsuta (AV34), Filaginella uliginosa (AV34), Heracleum sibiricum (AV344), Hierochloe repens (FV5/13), Pilosella cespitosa (AV344), Holcus lanatus (KC17), Hordeum murinum (SS105), Hypericum montanum (FV48), Inula salicina (AV311), Juncus compressus (AV344), Juniperus communis (SS417), Jurinea thyrsiflora (FV1a/10), Lactuca chaixii (FV6/57), Lactuca serriola (SS4/8), Lappula squarrosa (SS1/10), Lavatera thuringiaca (SS3/10), Leontodon danubialis (SS810), Leonurus quinquelobatus (SS4/8), Lepidium campestre (KC3/9), Linum catharticum (SS4/17), Linum flavum (SS1/10), Linuspermum officinale (SS65), Lolium perenne (AV2/5), Lyvium horgetina (MS2/10), Linusperina (AV2/5), Luzula campestris (SS7/10), Luzula multiflora (AV2/9), Luzula pilosa (KC3/9), Lycium barbatum (SS10/5), Lysimachia nummularia (AV1/5), Lysimachia vulgaris (AV3/4), Lythrum virgatum (AV3/11), Malva sylvestris (KC2/71), Medicago minima (SS3/10), Melampyrum arvense (SS4/17), Melampyrum nemorosum (KC3/18), Melampyrum polonicum (FV4/8), Melampyrum pratense (KC17), Melica nutans (FV4/8), Milium effusum (AV3/7), Mycelis muralis (FV6/57), Myosotis arvensis (AV1/2), Myosotis cespitosa (AV1/2), Nepeta cataria (SS8/5), Onopordum acanthium (FV5/13), Oxytropis pilosa (SS4/8), Petasites hybridus (AV3/11), Phalaroides arundinacea (AV3/4), Polygala podolica (AV2/5), Persicaria minor (AV3/4), Populus tremula (KC1/7), Potentilla heptaphylla (AV3/4), Potentilla obscura (SS6/11), Ptarmica cartilaginea (AV3/14), Pulsatilla patens (FV4/8), Ranunculus auricomus (AV1/3), Ranunculus flammula (AV3/7), Rhinanthus minor (AV1/2), Rorippa austriaca (AV2/9), Rorippa sylvestris (AV3/4), Rosa porrectidens (SS4/8), Rubus caesius (AV3/4), Sagina procumbens (KC1/7), Salix triandra (j) (AV3/4), Salvia nutans (SS2/10), Sambucus ebulus (AV2/5), Saxifraga tridactylites (SS5/17), Scirpoides holoschoenus (SS1/9), Schivereckia podolica (SS4/17), Scolochloa festucacea (AV3/4), Scrophularia nodosa (AV1/2), Senecio vulgaris (FV6/29), Seseli campestre Scirpoides holoschoenus (SS119), Schivereckia podolica (SS4/17), Scolochloa festucacea (AV3/4), Scrophularia nodosa (AV1/2), Senecio vulgaris (FV6/29), Seseli campestre (FV5/25), Sesleria heufleriana (SS1/10), Silene chlorantha (SS3/10), Silene lithuanica (SS9/13), Silene viscosa (SS4/8), Solanum dulcamara (FV6/14), Sorbus aucuparia (FV6/14), Spergula morisonii (FV8/7), Stachys officinalis (AV1/3), Stellaria media (AV2/5), Stipa borysthenica (FV4/23), Succisa pratensis (SS10/5), Swida sanguinea (SS1/10), Taniocpetalum arenarium (FV4/15), Teucrium pannonicum (SS1/10), Thalictrum flavum (AV2/5), Thalictrum lucidum (AV1/2), Thalictrum simplex (AV1/2), Thesium arenarium (FV4/15), Thesium arenarium (SS1/10), Tragopogon orientalis (KC1/7), Tribulus terrestris (FV1a/10), Trisetum sibiricum (AV3/4), Turritis glabra (FV4/15), Ulmus minor (SS10/5), Urtica dioica (SS4/25), Valerianella coronata (FV5/13), Verbascum phlomoides (SS12/67), Veronica arvensis (KC1/7), Veronica fuccium (AV2/5), Veronica prostrata (SS4/8), Veronica scutellata (AV3/7), Veronica teucrium (AV2/5), Veronica triphyllos (SS8/14), Vicia villosa (FV2a/20), Viola reichenbachiana (KC1/7), Viola rupestris (SS4/8), Xanthium strumarium (AV1/2), Non-vascular plants: Bryum subelegans (SS4/8), Candonia polydactyla (KC1/7), Cladonia alpina (KC1/4), Cladonia degenerans (FV8/27), Cladonia polydactyla (KC1/2), Climacium dendroides (KC1/7), Diploschistes scruposus (SS6/11), Encalypta rhaptocarpa (SS4/8), Funaria hygrometrica (KC1/7), Hypnum vaucheri (SS5/17), Leucodon sciuroides (SS4/8), Orthotrichum anomalum (SS4/8), Peltigera canina (SS6/5), Pogonatum urnigerum (SS6/5), Porella platyphylla (SS4/17), Pseudoleskeella tectorum (SS5/17), Ramalina pollinaria (SS10/5), Schistidium trichodon (SS4/8), Tortula subulata (SS10/5), Trinia multicaulis (SS4/8).