



The impact of agriculture on tropical mountain soils in the western Peruvian Andes: a pedo-geoarchaeological study of terrace agricultural systems in the Laramate region (14.5°S)

5 Fernando Leceta^{1,2}, Christoph Binder³, Christian Mader⁴, Bertil Mächtle¹, Erik Marsh⁵, Laura Dietrich⁶,
Markus Reindel⁷, Bernhard Eitel¹, Julia Meister³

¹Geomorphology and Soil Geography, Geographical Institute, Heidelberg University, Heidelberg, D-69120, Germany

²Historical Mapping Laboratory, Espacio Interdisciplinario, Universidad de la República Uruguay (UdelaR), Montevideo, C.P. 11200, Uruguay

10 ³Geoarchaeology and Quaternary Science, Institute of Geography and Geology, University of Würzburg, Würzburg, D-97074, Germany

⁴ArchDepth Research Group, Bonn Center for Dependency and Slavery Studies, University of Bonn, Bonn, D-53113, Germany

⁵National Scientific and Technical Research Council (CONICET), Universidad Nacional de Cuyo, Mendoza, CP. M5502JMA, Argentina

15 ⁶Prehistoric Archaeology, Martin Luther University Halle-Wittenberg, Halle, D-06108, Germany

⁷Commission for Archaeology of Non-European Cultures, German Archaeological Institute, Bonn, D-53173, Germany

Correspondence to: Fernando Leceta (leceta@gmail.com); Julia Meister (julia.meister@uni-wuerzburg.de)

20 **Keywords.** Andes, prehispanic agricultural terraces, agricultural terrace soils, soil science, paleo-pedology, land use dynamics, phytolith analysis

Abstract. This integrated pedo-geoarchaeological study focuses on three abandoned prehispanic terrace agricultural systems near Laramate in the southern Andes of Peru, aiming to unravel the pedological and land-use history of the region. The investigation involved contextualizing the former agricultural management system within its paleoecological framework and assessing the impact of agricultural practices on soil development and quality. The Laramate terrace complex, with its
25 diverse terrace systems and varied geomorphological settings, provided an ideal setting for the investigation. Comparative analyses between non-irrigated agricultural terrace soils and undisturbed reference sequences were conducted, employing a range of methodologies, including surveys, soil analysis, GIS and remote sensing, palaeobotany, and radiocarbon dating.

The study identifies three WRB Reference Soil Groups in the Laramate region: *Phaeozems*, *Andosols*, and *Anthrosols*. Unique characteristics of *Phaeozems* challenge typical descriptions, influenced by factors such as climatic seasonality,
30 vegetation, fauna, lithology, and aeolian inputs. Despite long-term use, terrace soils (*Anthrosols*) show no severe degradation, maintaining balanced acidity and nutrient availability for successful crop cultivation. Tillage horizons of all terrace soils exhibit elevated organic matter content, highlighting the sustainability of prehispanic agricultural practices through a consistent application of organic manure. Phytolith concentrations indicate extensive agricultural activities, particularly



maize cultivation, with varying patterns suggesting changes in cultivation or fertilization practices over time. Starch grain
35 identification aligns with phytolith analyses, reinforcing maize's significance in the region. Notably, the abandonment of the
prehispanic cultivation system was not linked to soil exhaustion or terrace instability.

The prehispanic history of terraced agriculture in the Laramate region extends over four development phases, reflecting
dynamic interactions between environmental, cultural, and agricultural factors. The initial phase, from the Formative Paracas
40 period to the Early Nasca period (800 BCE–200 CE), witnessed the establishment of agricultural terraces with simple terrace
architecture, while the Middle Horizon (600–1000 CE) saw systematic areal expansion influenced by the Wari culture.
Adaptations to drier conditions included terraced agriculture on volcanic soils. The Late Intermediate Period (1000–1450
CE) witnessed hydrological variability and further terrace expansion to lower altitudes and less agriculturally suitable
locations. The final phase, marked by the onset of the Hispanic colonial period in 1532 CE, saw the gradual abandonment of
45 terraced agricultural systems due to demographic shifts and reorganization of production systems. Despite this, the historical
trajectory underscores the adaptability and resilience of prehispanic communities in the Laramate region, showcasing
innovative terrace agriculture as a means of coping with changing environmental conditions across diverse landscape units.

1 Introduction

Terrace farming encompasses a set of specialized techniques crafted to facilitate agriculture in mountainous regions, where
50 steep slopes typically pose challenges for cultivation. This method is globally practiced across diverse climatic conditions
and entails substantial alterations to the natural landscape for agricultural purposes. This involves creating artificial terraces
and forming an artificial tillage 'Ap' horizon as a new topsoil, as can be observed in the tropical Andes or in central
Amazonia (Zavaleta García, 1992; Sandor and Eash, 1995; Glaser and Birk, 2012; Varotto et al., 2019).


55 Similar to other semi-arid and mountainous regions worldwide, terracing in the Peruvian Andes serves the primary goal of
enhancing agricultural productivity by addressing environmental challenges. These challenges include the erratic availability
of soil moisture during dry and rainy seasons, the constrained expanse of cultivable land, and the inherent risk of soil
erosion. In the Peruvian Andes, the availability of fertile and arable land is a limited resource, constituting only 19% of the
national territory (Zavaleta García, 1992; Pulgar Vidal, 1996). The Western Cordillera of the southern Peruvian Andes
60 boasts a longstanding tradition of agricultural terracing, a practice with roots dating back to prehispanic times. These
agricultural terraces typically align with a fertile strip characterized by rich natural soils, notable accumulations of organic
matter in the mineral topsoil, and elevated levels of base saturation. This fertile zone extends along the slopes at elevations
between 2500 and 3700 m asl. Locally referred to as "Serranía Esteparia" (Brack Egg and Mendiola Vargas, 2010), "Region
Quechua" (Pulgar Vidal, 1996) or "Pajonales" (Weberbauer, 1944), this semi-arid and cool ecosystem is predominantly
65 marked by seasonal and sparse shrub and grassland vegetation. It shares agro-ecological characteristics with highly fertile



steppe landscapes in mid-latitudes. While steppe ecosystems and soils have been extensively studied in the Eurasian steppe belt, the Canadian prairies, and eastern Patagonia (Zech et al., 2014; Blume et al., 2010; IUSS Working Group, 2015), the agricultural soils in the Peruvian Andes, which share similar characteristics, have only captured the attention of the scientific community in recent decades (Sandor and Eash, 1995; Kemp et al., 2006; Goodman-Elgar, 2008; Aguirre-Morales, 2009; Kendall and Rodríguez, 2009; Branch et al., 2007; Nanavati et al., 2016; Handley et al., 2023).

The region surrounding the small village of Laramate (14°17'11" S, 74°50'34" W), situated on the upper slopes of the Rio Grande drainage in the Peruvian southern Andes, features numerous terrace agricultural systems (Fig. 1). This terrace complex serves as a compelling illustration of the sustainable and long-term utilization of land and soil in the region during prehispanic times. Recent archaeological findings (Reindel and Isla, 2013a; Soßna, 2015) underscore the region's significance as an agricultural hub, contributing to the development of the local Paracas (800–200 BCE) and Nasca (200 BCE–600 CE) cultures. Additionally, it played a pivotal role in the expansion of the Wari (600–1000 CE) and Inka (1450–1535 CE) cultures in the Andes.

The agricultural terrace systems in the Laramate region offer a valuable opportunity to explore the enduring effects of agricultural practices on soils within the mountain shrubland ecosystem. To create an accurate and up-to-date soil map of the study area at a 1:50,000 scale, comprehensive soil surveys were conducted. The exploration of prehispanic terrace agricultural systems involved pedestrian surveys and detailed mapping, incorporating satellite images. In order to assess the morphological and geochemical soil characteristics resulting from prolonged agricultural use, we compared five non-irrigated agricultural terrace soils from three terrace agricultural systems with two undisturbed reference profiles across various soil groups in diverse lithological and topographical settings (Fig. 1). This evaluation specifically focused on (i) soil hydrological functions, (ii) nutrient availability and organic matter content, and (iii) cultivated plants, employing a pedo-geoarchaeological approach. To analyse these aspects, selected samples underwent standard procedures for bulk chemistry, texture, nutrient budget, bulk density, phytoliths, and starch. Additionally, the chronostratigraphy of the sequences was established through radiocarbon dating.



95

Figure 1: a: Location of the Laramate region within Peru. b: Schematic cross section of the Rio Grande drainage, including major landscape elements and the Laramate terrace complex. c: Soil map of the study area according to the FAO WRB classification and location of major terrace systems, sampling sites and settlements. d-f: Satellite images of the Ayllapampa, Santa María and Sihuilca terrace complexes and location of the soil profiles studied. Sources: figure (a): soil data own sampling and survey;



lithology, river & village data basis INGEMMET 2001; elevation model ASTER GDEM (e) map (b): NaturalEarth; images (c), (d) © DigitalGlobe; image (e): © Google Maps 2023; figure (b): modified from Mächtle 2007 and INGEMMET 2001, some drawing-vectors from Freepik.

100 2 Study site

2.1 Location

The study area is located near the rural village of Laramate (14°17'11"S, 74°50'34"W, 3045 m asl) and covers a total area of 380 km² in the mountain shrubland landscape of the western Cordillera Occidental in Peru (Fig. 1). In terms of hydrology, this area corresponds to the upper and middle drainage region of the Rio Grande de Nasca. The valley follows a south-
105 westerly course, descending from the continental divide to the Pacific Ocean. The altitude of the mountainous region ranges from 2000 m to 4200 m asl.

2.2 Geology

The geology of the Laramate region is complex and can be divided into three main groups. The consolidated sedimentary rocks of the Yura Group, of Jurassic-Lower Cretaceous age, form a thick sedimentary sequence of sandstone, quartzite, shale
110 and limestone, exposed in the valley bottoms and adjacent slopes of the Palpa, Viscas and Ingenio rivers. The location of the Santa María archaeological site corresponds to the Hualhuani Formation subgroup, within which quartzite and quartz rich sandstone beds are arranged in thick beds with some intercalation of laminar stratified silt and claystones (Castillo et al., 1993).

115 Tonalite and granodiorite are the most widespread lithological group. These acidic intrusive rocks belong to the plutonic bodies of the Peruvian Coastal Batholith, which have intruded since the Lower and Upper Cretaceous through older sedimentary rocks and younger volcanic bodies. While tonalites are mainly exposed in the valley bottoms, granodiorite outcrops are restricted to the summits and slopes. Spheroidal weathering is typical of these rocks, best seen in the sector between the villages of Laramate and Ocaña, where intense weathering disintegrates them into boulders or coarse sand
120 masses. The archaeological site of Sihuilca and the control soil profile Pe09-02 are located on this sub-group (Castillo et al., 1993).

A thick volcanic-clastic sedimentary sequence of Paleogene-Neogene age dominates the north-eastern half of the study area, occupying the upper positions of the Cordillera Occidental as it enters the Altiplano. It gathers a series of conglomerates,
125 breccias, tuffs, lava flows and reworked sandstones. The Ayllapampa archaeological site is located in the Castrovirreyna Formation, an Eocene-Oligocene subunit. At this site, breccias of andesitic composition dominate (Castillo et al., 1993). The Pliocene deposits of the Nasca Group sub-unit are exposed at higher elevations within the Rio Grande watershed. The sequence is composed of a package of lapilli tuffs at the base and is completed in the upper levels by volcanic conglomerates



130 containing clasts of rhyolite, dacite and trachyandesite composition within a sandy tuffaceous matrix (Castillo et al., 1993).
The sample site Pe10-32, near Pucará, was developed from the volcanic ejecta of the basal levels.

2.3 Climate

135 The study area covers a transitional zone between the semi-arid western slopes of the Cordillera Occidental and the sub-humid western ranges of the Altiplano. Seasonality and semi-aridity are the main features that characterize the climatology of the Laramate region. There are two reasons for the aridity of the Pacific coast and the western slopes of the Andes. First, the orography of the central Andes prevents the transport of moist air from the Amazon basin and the Atlantic. Secondly, the Southeast Pacific Anticyclone generates very dry and stable conditions (Garreaud et al., 2003), inhibiting the development of local convective activity. The moist air masses that reach the Altiplano and the western flanks of the Cordillera Occidental originate as tropical convective rainfall over the Amazon basin and are subsequently transported to the western Andes by the tropical easterly current during the austral summer months (December to March) (Garreaud et al., 2003).

140

The climate of the Laramate region is classified as alpine semi-arid and temperate, with a pronounced rainy season from December to April in which about 90% of the total annual rainfall of 250–300 mm/year is concentrated. This is followed by a long dry season (May to November) characterized by a lack of rainfall, high insolation and greater temperature amplitude due to the lack of cloud cover and stronger easterly winds. Laramate has an average annual temperature of 10°C, and frost can occur in the winter months. Thermal amplitudes increase proportionally with altitude (ONERN, 1971).

145



Figure 2: Panorama photos of the Ayllapampa (a), Santa María (b, c) and Sihuilca terrace complexes (d).

150 2.4 Vegetation

The mountain shrubland (2500–3800 m asl) is a high-altitude, inner-tropical, low-rainfall vegetation formation that extends across the steep and highly dissected topography of the Cordillera Occidental. Floristically, it is characterized by the gradual substitution of succulent (*Cereus candelaris* and *Opuntia subulate*) and xerophytic shrub communities (*Viguiera sp.*, *Chinopappus benthamin*, *Mutisia acuminata* and *Dodonea viscosa*), which predominate in the semi-arid lower levels. The sub-humid upper levels are dominated by deciduous and frost-tolerant woody shrubs (*Schinus molle*, *Escallonia resinosa*, *Agave americana*, *Dodonea viscosa*, *Tecoma sambucifolia*, *Spartium junceum* and *Caesalpinea spinosa*) and perennial grasses (*Festuca*, *Stipa*, *Deyeuxia* and *Poa*) (Brack Egg and Mendiola Vargas, 2010; MINAM, 2015).



160 An ecosystem of economic importance since prehispanic times, the mountain shrubland of the Peruvian Andes has suffered
more degradation from deforestation, burning and overgrazing than any other vegetation type in the Cordillera Occidental.
Non-intensive agriculture, intended for subsistence, is confined to the wet, narrow banks of the major river valleys, as well
as seasonally wet valley flanks and small episodic streams. Barley (*Hordeum vulgare*), potatoes (*Solanum tuberosum*), wheat
(*Triticum spp.*), maize (*Zea mays*), avocado (*Persea americana*), ulluco (*Ullucus tuberosus*), broad beans (*Vicia faba*), pea
(*Pisum sativum*), oca (*Oxalis tuberosa*) and mashua (*Tropaeolum tuberosum*) are today among the most frequently cultivated
165 species (ONERN, 1971).

2.5 Soils

The study area is characterized by two particularly fertile soil groups, *Phaeozems* and *Andosols*, which form on different
parent rocks and provide the undisturbed pedological context for the agricultural terrace soils, which are classified as
Anthrosols.

170 *Phaeozems* are the dry grassland soils of the semi-arid mountain shrubland of the Peruvian Andes. They typically form on
the middle and upper slopes of the Cordillera Occidental at altitudes between 2500 and 3800 m asl. Due to their excellent
agro-ecological quality, the *Phaeozems* of the mountain shrubland have been used since prehispanic times and are
considered the most productive soils in Peru (Zamora Jimeno and Bao Enríquez, 1972; Brack Egg and Mendiola Vargas,
175 2010; Instituto de Estudios Histórico-Marítimos del Perú., 1996). Although *Phaeozems* are not described in detail for the
study area, the genetically related soil types *Luvic Kastanozems* and *Calcic Kastanozems* are described. While *Luvic
Kastanozems* are characterized by increasing clay content and clay coatings with depth due to clay illuviation, *Calcic
Kastanozems* are alkaline soils with typical soil saturation of more than 50% in the upper 50 cm of the topsoil (Zamora
Jimeno and Bao Enríquez, 1972; Instituto de Estudios Histórico-Marítimos del Perú., 1996).

180 *Andosols* are dark soils typical of volcanic landscapes, developed from glass-rich volcanic ejecta in almost any climate. In
the western Andes, they occur in the undulating landscape of the upper slopes of the Cordillera Occidental and the adjacent
Altiplano above 3600 m asl, where semi-aridity prevents soil nutrient leaching and infrequent frosts do not severely restrict
root growth (Zamora Jimeno and Bao Enríquez, 1972; Instituto de Estudios Histórico-Marítimos del Perú., 1996; Brack Egg
and Mendiola Vargas, 2010). Their development is strongly conditioned by the widespread grassland vegetation of the
185 mountain shrubland of the Peruvian Andes, which provides organic matter (Zamora Jimeno and Bao Enríquez, 1972).
Mountain shrubland *Andosols* are moderately acidic and generally shallow; organic carbon levels within the top 50 cm are
typically above 1% (Zamora Jimeno and Bao Enríquez, 1972). Although not as fertile as *Phaeozems*, *Andosols* have dark
and humus-rich surface horizons, less base-rich than typical mid-latitude steppe soils (*Chernozems*, *Kastanozems*,
190 *Phaeozems*) but potentially valuable for agriculture if managed appropriately (IUSS Working Group, 2015). Whereas *andic*



properties are related to an advanced state in the weathering sequence of pyroclastic deposits (tephric soil material > vitric properties > andic properties), layers with *vitric* properties are distinguished by a lower degree of weathering (IUSS Working Group, 2015).

195 *Anthrosols* are the group of soils that have been heavily modified by human activities such as harvesting, irrigation, fertilization or ploughing (IUSS Working Group, 2015; Zech et al., 2014). In agricultural terraces, therefore, changes in the natural soil sequence and properties caused by intensive agricultural use usually lead to the formation of *Anthrosols*.

2.6 Holocene environmental and prehispanic settlement history of the Laramate region

The history of human occupation and settlement is well documented in the Laramate region, from the Archaic period (8000–
200 3500 BCE) to Inka sites from the early sixteenth century (Reindel, 2009; Unkel et al., 2012; Reindel and Isla, 2013a; Sořna, 2015; Beresford-Jones et al., 2023). The Cerro Llamocca sacred mountain complex (4487 m asl) includes the oldest archaeological site hitherto recorded in this study area: a rock shelter with evidence of human occupation in the Early Archaic period ~ 8000 BCE (Mader et al., 2023b). Settlement and land use activities increased significantly during the Formative period from 1700 to 200 BCE, especially toward the end of the Paracas period (800–200 BCE), coinciding with a
205 more humid climate following a pronounced dry period from 2600 to 2200 BCE. Large-scale archaeological excavations at Collanco (1630 m asl) and Cutamalla (3300 m asl) have provided rich data on the chronology, layout, and use of Paracas settlements in this highland region (Reindel et al., 2015; Reindel and Isla, 2017, 2018; Mader, 2019; Mader et al., 2023a, 2024). These settlement activities were associated with cultivation in the extensive terrace farming systems that surrounded the sites. Cutamalla was an economic and agricultural centre that was integrated into a considerable interregional exchange
210 network of various goods (e.g., obsidian, camelid wool, maize, and other agricultural goods) with llama caravans (Reindel and Isla, 2013b; Mader et al., 2018; Mader, 2019; Mader et al., 2021, 2023a, 2024).

The transitional Initial Nasca period (120 BCE–90 CE) was characterized by intense demographic mobility (Unkel et al., 2012). The florescence of the Nasca archaeological culture during the Early Intermediate Period (200 BCE–600 CE) – a
215 time of high cultural development – coincides with a humid and relatively stable interval. The decline of the Nasca culture at the end of the Early Intermediate Period and a strong presence of the Wari culture during the Middle Horizon (600–1000 CE) were largely contemporaneous with an abrupt turn to a sustained dry period between 600–800 CE. The expansive Wari state, which originated in the highlands, occupied ancient Nasca settlements and extended its influence into coastal regions (Reindel and Isla, 2013a). A representative Nasca and Wari site in the Laramate region is Huayuncalla (3100 m asl), which
220 was also excavated on a large scale to obtain reliable including settlement data (Isla and Reindel, 2014). Dry conditions prevailed until 1250 CE.



More humid but hydrologically highly variable conditions prevailed again after 1250 CE, which apparently had positive effects on settlement activities and an increase in population during the Late Intermediate period (1000–1450 CE), another era of cultural and economic flourishing in the region (Sořna, 2015). These developments were supported by advances in agricultural techniques and a massive expansion of sophisticated and partly irrigated farming terraces, leading to an intensification of agricultural production. The Inka state of the Late Horizon (1450–1535 CE) had a minor presence in the Laramate region; Santa María is one of the few sites with surviving Inka architecture and it was connected to the imperial road network. European contact, beginning in 1532 CE, had severe consequences for Andean communities and agricultural regimes. Introduced diseases led to a demographic decline and thus the available workforce. Moreover, aridization in the eighteenth and nineteenth century led to a profound environmental turnover, marked by dry conditions impacting the Altiplano ranges (Schittek et al., 2018). The current population density in the Laramate region is low and merely a small portion of the terrace agricultural systems are in use today, usually located along rivers.

In sum, important cultural changes seem to have coincided with significant changes in paleoclimate (Eitel and Mächtle, 2009; Mächtle and Eitel, 2013; Reindel and Isla, 2013a; Schittek et al., 2015). The mobility of people, llamas, and commodities was an important component of cultural and economic development (Mader et al., 2021; Beresford-Jones et al., 2023). Archaeological and geoscientific data indicate that a main function of prehispanic highland settlements and their related extensive terrace systems was agricultural production (Mader et al., 2024).

3 Material and Methods

Soil surveys were conducted to generate an accurate and current soil map of the study area at a scale of 1:50,000, adhering to international standards. The investigation into prehispanic agricultural terrace systems involved both pedestrian surveys and aerial mapping, utilizing satellite images. To discern morphological and geochemical features arising from prolonged agricultural activity, detailed pedological examinations and subsequent laboratory analyses were carried out on five key terrace profiles. These profiles were strategically chosen from a cluster of three terrace agricultural systems in the Laramate area, directly associated with the archaeological sites of Ayllapampa (profiles Pe11-07 and Pe10-30), Santa María (profiles Pe10-26 and Pe11-06), and Sihuilca (profile Pe10-31), which exhibit evidence of repeated prehispanic use. Additionally, two reference-sampling sites (Pe10-32, Pe09-02) were selected within the *Phaeozems* and *Andosols* edaphic soil provinces to represent undisturbed pedological contexts (Fig. 1, 2).

3.1 GIS-based mapping of agricultural terraces on satellite images

For precision in mapping the chosen agricultural terrace systems, a thorough analysis of satellite imagery from Google Earth Pro was conducted, focusing on discernible patterns, such as parallel lines indicative of terrace presence. Subsequently, individual terrace walls and boundaries were manually identified and digitized, distinguishing between abandoned and



actively used terraces. Nevertheless, challenges arose in the identification process, especially in regions obscured by
255 vegetation. Distinguishing terrace walls from animal tracks and geological formations proved challenging in certain
instances. Additionally, there were areas where satellite imagery suggested terracing, yet the terrace walls lacked the
necessary detail for accurate mapping.

3.2 Field methods

Fieldwork was conducted in three consecutive campaigns spanning the years 2009 to 2011, occurring during the dry season
260 in the austral winter months. In consideration of local relief conditions, soil profiles were systematically arranged in a
catenary manner along hillslopes. The variations observed in genetically related soils were found to be influenced by
landscape location, anthropogenic impact, drainage characteristics, and the downstream transport of soil material (Blume et
al., 2010; Zavaleta García, 1992). The catena under investigation stretches across a 13 km transect running parallel to the
Laramate-Viscas and Palmareda-Palpa rivers, situated between the villages of San Antonio de Locchas (14°19'01"S,
265 74°55'56"W, 2760 m asl) and Pucará (14°13'32"S, 74°53'36"W, 3520 m asl). Soil field properties were described in
accordance with the FAO Guidelines for soil description (Jahn et al., 2006), and soil classifications were carried out
following the IUSS Working Group on the WRB (IUSS Working Group, 2015).

3.3 Laboratory methods

3.3.1 Soil parameters

270 Twenty-eight samples were collected from the seven soil profiles examined in the field and laboratory (Supplementary Table
S1). Physico-chemical soil analyses were conducted at the Laboratory for Geomorphology and Geocology of the Institute
of Geography at Heidelberg University.

The sample preparation for laboratory analysis was following standard procedures of ISRIC (van Reeuwijk, 2002); the fine-
275 earth fraction (≤ 2.0 mm) of air-dried samples was removed by dry sieving prior to the analyses. All analytical results refer to
the fine earth fraction. Grain size analyses were carried out by a combination of the Köhn-pipette method (silt and clay
fractions) and wet sieving (sand fractions) (Schlichting et al., 1995). The measurement of the field-estimated bulk density
followed the WRB's recommendations (Jahn et al., 2006). A cylindrical coring instrument was used to obtain the samples
volume without disturbing the natural soil structure. The dry mass was determined after 48 hours of drying at 105°C. Bulk
280 density's calculation expresses the quotient of the substrate's dry weight and the core's volume. Soil colour was classified
under laboratory conditions on dry soil samples according to the Munsell Soil Color Charts (Munsell Color, 2000). The pH
value was measured in a 1:2.5 suspension in 1 M KCL solution using a pH-meter (van Reeuwijk, 2002). The cation
exchange capacity (CEC_{pot}) and exchangeable bases (BS) were detected using the Ammonium Acetated Method (van
Reeuwijk, 2002). The concentrations of plant-available cations Magnesium (Mg^{2+}), Kalium (K^{+}), Calcium (Ca^{2+}), Natrium



285 (Na⁺) were determined through Flame Atomic Absorption Spectrometry (Schimadzu AA-6300). Additionally, the
concentration of exchangeable cation Aluminium (Al³⁺) was determined using Graphite Furnace Atomic Absorption
Spectrometry (Analytik Jena-AAS Zenit 60). H⁺ ion's concentrations were obtained from the determination of Exchange
Acidity (H-value) in 1 Mol KCL solution. The soil organic carbon (C_{org}) contents were determined by applying the Wet
Combustion Method following the DIN procedures (DIN 19684-2, 1977). Total carbon, nitrogen, and sulphur contents were
290 measured on milled samples using the CNS analyzer vario MAX (Fa. Elementar). The carbon/nitrogen (C/N) ratios were
calculated to evaluate the mineralization rates in the soil. Total phosphorus contents were photometrical detected in the form
of a molybdenum-blue complex at 880 nm (Analytik Jena Specord 200 Plus) using the Ammonium molybdate spectrometric
method according to Olsen (1954) described by van Reeuwijk (2002). The plant available phosphorus fraction was extracted
with a 1% citric acid solution; phosphate in the extract was determined colorimetrically with the blue ammonium molybdate
295 method with ascorbic acid as reducing agent (van Reeuwijk, 2002).

3.3.2 Radiocarbon dating

To place the terraces systems in time and to compare chronologies to nearby residential contexts, six charcoal samples and
one bulk sediment sample from the terraces at Sihuilca, Santa María and Ayllapampa were radiocarbon dated at the Curt-
Engelhorn-Centre of Archaeometry in Mannheim, Germany (Table 1). Samples were pretreated using the acid-alkali-acid
300 method to remove contamination by carbonates and humic acids. The results were calibrated in OxCal 4.4 (Bronk Ramsey,
2009), using a site-specific mixed curve of SHCal20 (Hogg et al., 2020) and IntCal20 (Reimer et al., 2020) that reflects
modern mixtures of air from the Northern and Southern Hemispheres at Cutamalla (Mader et al., 2024), following recently
published methods (Ancapichún et al., 2022; Marsh et al., 2023).



305

Table 1: Sample information and results of radiocarbon dating in the Laramate terrace complex. $\delta^{13}\text{C}$ values are from AMS stable isotope measurements and are not comparable to $\delta^{13}\text{C}$ values measured with IRMS.

Lab code MAMS	Profile and sample ID	^{14}C age	\pm	$\delta^{13}\text{C}$ AMS (‰)	Median	95% range (rounded by 10 years)	Soil horizon	Horizon depth (cm)	Sample material
13597	Pe10/30-4	1291	22	-25.2	730 CE	670–860 CE	AC	65–100	charcoal
13598	Pe10/30-3	1183	22	-22.6	930 CE	770–990 CE	Ah	35–65	charcoal
13599	Pe10/31-3	1169	13	-24.1	940 CE	880–990 CE	Ah	20–50	charcoal
20089	Pe10/26-4	212	22	-26.8	1760 CE	1650–1810 CE	C1	60–100	charcoal
20090	Pe11/06-3	611	19	-27.8	1350 CE	1310–1410 CE	AC	30–45	charcoal
20091	Pe11/07-2	2920	21	-27.3	1080 BCE	1210–1000 BCE	Ah	25–50	charcoal
18749	Pe10/32-3	3754	19	-28.9	2120 BCE	2210–2030 BCE	2Ahb	20–30	bulk sediment (≤ 2 mm)

3.3.3 Phytolith analyses

310 Phytolith analyses were conducted on twenty-five samples derived from five terrace profiles (Pe10-31, Pe10-26, Pe11-06, Pe10-30, Pe11-07), along with one reference profile (Pe10-32) at the University of Würzburg, following established extraction protocols (Albert et al., 1999). Approximately 1 g of air-dried sample (< 2 mm) was treated with 3 N HCl, 3 N HNO₃, and H₂O₂ (30 %) to remove carbonates, phosphates, and organic material. The insoluble mineral components were then fractionated into three density-based fractions using a ~ 2.4 g/ml sodium polytungstate solution [Na₆ (H₂W₁₂O₄₀) ×

315 H₂O]. The third fraction, anticipated to contain the majority of phytoliths due to its lower density, was used for analysis. Approximately 1 mg of this fraction was weighed, mounted on a microscope slide with Entellan New (Merck), and examined under a KERN OBE-114 microscope at 400x magnification. Image capture of selected phytoliths was achieved using a KERN ODC 825 microscope camera. To establish a suitable basis for statistical evaluation, a minimum of 200 identifiable phytoliths per sample were counted (Albert et al., 1999; Piperno, 2006; Meister et al., 2017).

320

Morphological identification of phytoliths relied on subject-specific literature (Bremond et al., 2008; Pearsall et al., 2003; Piperno, 1988, 2006; Twiss, 1992), with adherence to the International Code for Phytolith Nomenclature where applicable (Madella et al., 2005; International Committee for Phytolith Taxonomy (ICPT), 2019). Phytoliths lacking a clear description were categorized as weathered morphotypes. Phytolith concentrations were calculated as described by Albert et al. (1999).



325 3.3.4 Starch analysis

330 Starch analyses were conducted at the University of Halle on five samples derived from two terrace profiles (Pe10-31, Pe10-26), along with one reference profile (Pe10-32; Table 2). Microfossil extraction followed the procedure developed by Liu et al. (2018). For each sample, 40 mg of sediment was meticulously ground in a mortar and transferred to individual tubes (1.5 ml). To these tubes, 50 µl of EDTA was added, followed by 450 µl of a 2.4 g/ml sodium polytungstate solution [Na₆(H₂W₁₂O₄₀) × H₂O]. The tubes were vortexed to disperse the sediment and then subjected to an ultrasonic bath for 10 minutes. Afterward, the tubes underwent centrifugation at 5000 rpm for 10 minutes to concentrate the starch, and the resulting supernatant was transferred to a new tube and vortexed. Subsequently, 50 µl of the supernatant from each sample was mounted on glass slides and analysed for starch using polarisation filters at 400x magnification with a Bresser polarisation microscope. Photographs were captured using a 12 MP Bresser Microcam.

335

Table 2: Sample information and results of starch analyses in the Laramate terrace complex.

Profile and sample ID	Soil horizon	Horizon depth (cm)	Terrace system	Starch (<i>Zea Mays</i>)
Pe10/31-2	Ap2	10-20	Sihuilca	2 grains
Pe10/31-4	Bwt	50-70	Sihuilca	1 grain
Pe10/26-1	Ap	0-15	Santa María	1 grain
Pe10/26-4	C1	60-100	Santa María	-
Pe10/32-3	2Ahb	20-30	Soil control profile	-



4 Results

340 4.1 The Laramate terrace complex

The Laramate terrace complex encompasses a collection of predominantly abandoned prehispanic agricultural terrace systems near the village of Laramate (14°17'11" S, 74°50'34" W, 3045 m asl; Fig. 1). These terrace systems extend in a scattered pattern across both flanks of the Laramate-Viscas and Palmadera-Palpa rivers, situated between the semi-arid and sub-humid margins of the mountain shrubland ecosystem. Almost each terrace system is closely associated with, and frequently named after, an adjacent archaeological settlement of varying prehispanic age. Notable examples include Sihuilca, Locchas, Santa María, Pocopca, Chaupicancha, Cutamalla, Carhuachuco, Ayllapampa, Lindero, Mauka Llaqta, Plaza Pampa, and Colca Chica (Soßna, 2015).

While the primary purpose of the terrace systems in the Laramate area was for the cultivation of agricultural products, discoveries indicate that these terraces served as more than just cultivation sites (Reindel and Isla, 2013a). They also functioned as processing and storage units for specific products, such as maize, wood, or wool, intended for later commercial exchange. Furthermore, these terraces were occasionally utilized for purposes such as animal husbandry, residential use, erosion control, and even ceremonial practices, as exemplified by the *Challupas* building in Santa María (Soßna, 2015).

The dimensions of each individual terrace system vary, influenced by both soil moisture conditions and the area of potentially cultivable land. Additionally, a discernible connection between the size of the system and its geomorphologic setting exists. The most expansive terrace systems, spanning several square kilometres, are typically situated on lower and gently sloping terrains, such as valley bottoms, alluvial fans, and debris cones. In contrast, those confined to narrow and steep valleys and slopes tend to be significantly smaller in size, exemplified by the terraces surrounding the Chaupicancha site (Fig. 1).

The absence of significant hydraulic engineering features in the Laramate area, such as water reservoirs, aqueducts, or irrigation canals described in other regions like the Rio Grande's floodplain and lower drainage area, or the notable *Puquios* filtration galleries and *Khadin*-like water harvesting systems found in the Nasca-Palpa region of the Andean footzone (Hesse, 2008; Mächtle et al., 2009; Soßna, 2015), suggests that irrigation played a minor role in these agricultural terrace systems. In prehispanic times, the functionality of the Laramate terrace complex essentially relied on a rain-fed system, where moisture management centred on efficient drainage and the storage of rainwater within each terrace.

Only a relatively limited network of irrigation canals supplies certain terraces, primarily concentrated on the adjacent slopes of rivers in the upper catchments, such as the Laramate-Viscas river, which ensures a consistent water supply throughout the year. Currently, the terraces located between Laramate and Atocata stand out as the sole group with a constant water supply



utilized year-round. Additionally, smaller irrigation systems have been identified in the lower slopes and on alluvial terraces, notably in the northern sector of the Santa María settlement. Overall, pinpointing irrigation infrastructures in the Laramate area and determining their age poses challenges due to their often poor preservation. However, in the vicinity of Santa María, an irrigation facility is likely associated with the Middle Horizon Cultural Period, given its connection to a nearby small Wari compound (Soßna, 2015).

Currently, the active terraces are utilized for agriculture only during a brief period determined by the rainy season. Permanent water sources are scarce, occurring mainly in favourable locations where they enable an extension of the regular cropping season and, subsequently, facilitate extensive sheep and goat grazing. The Chaupicancha terrace system stand out as an exceptional case in this regard. Previous research suggests that the construction of terraces and their intensive use in the Laramate area in prehispanic periods can be associated with more favourable climatic conditions than those observed today. This correlation is particularly noted during the Early Horizon (800–200 BCE), the Early Intermediate Period (200 BCE–600 CE), and the Late Intermediate Period (1150–1438 CE) in the upper and middle Rio Grande basin (Eitel et al., 2005; Mächtle, 2007; Mächtle and Eitel, 2013; Schitteck et al., 2015).


4.2 Studied terrace systems and pedo-sedimentary characteristics

4.2.1 Sihuilca terrace system: overview and soil profile Pe10-31

The archaeological site Sihuilca (PAP-888 and PAP-897) was recorded in 2008 by surveys of the Nasca-Palpa Archaeological Project. According to surface ceramics and architectural remnants, PAP-888 is a domestic settlement dating to the Nasca Early Intermediate period (200 BCE–600 CE) and Wari Middle Horizon (600–1000 CE), while PAP-897 probably dates to the Middle Horizon (600–1000 CE) (Reindel and Isla, 2013a; Soßna, 2015).

The Sihuilca terrace system encompasses an area of approximately 650 ha (6.50 km²), with around 318 ha (3.18 km²) clearly identified as terraced agricultural land (Fig. 3). Mapping efforts have identified a total of 2739 terrace walls, amounting to a collective length of 107.49 km. These identified terraces predominantly face west and exhibit a generally heterogeneous pattern. Larger terraces are observed in flatter areas, while smaller terraces are found on steeper slopes. The width of the terraces varies considerably, ranging from 3m to 30m.

Soil profile Pe10-31 (14°19'30.22"S, 74°56'5.01"W, 2750 m asl) is situated approximately 1 km southwest of the village of Locchas within the catchment area (21.8 km²) of a tributary of the Huichulca-Viscas river system. The profile is positioned in the centre of a 650 m long slope that faces west, featuring an inclination of 25°.



405 **Figure 3: Detailed map of the Sihuilca terrace system.**



The sample site exhibits no signs of recent soil erosion. Isolated stones (6–20 cm) and boulders (20–60 cm) likely originate from collapsed terrace walls. The topsoil can be divided into two sections based on its structure. The first section (0–20 cm) is stratified and moderately cohesive, while the second section (20–50 cm) is characterized by coarse, blocky, sub-angular peds and displays moderate to high cohesion. The upper 50 cm of the soil matrix is enriched with roots ranging from 2 to 5 mm in diameter (Fig. 4).

The soil type is a *Terric Anthrosol (Loamic, Escalric)* with two edaphic cycles and six soil horizons (Ap1-Ap2-Ah-Bwt-C1-C2). **Ap1** (0–10 cm) is a slightly compacted, poorly developed, dark brownish-black (10 YR 3/3) tillage horizon. The underlying **Ap2** (10–20 cm) tillage horizon is slightly more compact, less developed and dark brownish-black (10 YR 3/3), containing human artefacts such as obsidian point fragments. The following **Ah** (20–50 cm) is brownish-black (10 YR 2/3), with slightly higher humus content and structural development. It represents the former soil surface before terracing. The horizon contains particles of burnt charcoal. Its lower boundary is gradual. **Bwt** (50–70 cm) is a transitional horizon with a brownish-black colour (10 YR 2/3), higher skeletal content and an increased clay content. It marks the lower limit of human influence on the soil. **C1** (70–85 cm) is a brownish-black (10 YR 2/3) horizon. **C2** (> 85 cm) represents the parent material of the soil, slightly weathered granodiorite; the horizon contains a few boulders, systematically situated at the lower terrace base. A 14C-dated charcoal taken from the **Ah** horizon (20–50 cm) revealed an age of 880–990 CE (Table 1, Fig. 5).





Figure 4: Photographs of the terrace soils Pe10-31, Pe11-06 & Pe11-07.



While the highest CEC_{pot} values are found in the surface horizons Ap1 and Ap2 (16.3 and 16.9 cmol/kg), the underlying Ah has the lowest values (13.9 cmol/kg; Fig. 5). Base saturation reaches very high values in all sampled horizons, starting from 99.4%. **Soil pH** ranges from 6.1 in Ap1 to 5.4 in the Ah, where soil acidity is moderate to weak. **Organic matter** tends to decrease from 3.5% in Ap1 to 1.6% in C1. The lowest value (1.5%) was found in the buried Ah. The **C/N ratio** decreases gradually towards the parent material and varies between 7.5 in the uppermost Ap1 and 6.4 in the C1. The **soil texture** of the uppermost Ap1 is loamy. An increase in the relative proportion of clay fractions in all underlying horizons results in a change in texture class to clay loam. **The skeleton fraction** of the topsoil horizons differs from the subsoil horizons by a higher content of coarse fractions. While the proportions in the upper 60 cm range between 24.7 and 25.4 %, this proportion decreases to 19.3–14.2 % in the lower part of the profile. The **plant available P fraction** remains consistently high (0.4 mg/g) at the cultivable Ap horizon, and its concentration decreases drastically (0.1–0.0 mg/g) in the Ah and with increasing depth.




440 **Figure 5: Profile description and laboratory results for terrace soil Pe10-31: macro-morphology, pedo-chemical analytical data, AMS 14C dating and total phytolith amounts.**

In the context of **phytolith analysis**, a total of 46 phytolith morphotypes were identified across all sites (Supplementary Tables S2 & S3, Fig. 6). For simplicity, the following descriptions are based on the 20 most common morphotypes (Fig. 7). Ten out of the 20 morphotypes can be attributed to the monocotyledonous class (Poaceae), while the remaining morphotypes may also originate from dicotyledons. Four of the monocotyledonous (Poaceae) morphotypes can serve as diagnostic indicators for *Zea mays*. The frequency of these morphotypes exhibits variation across sites.



In soil profile Pe10-31, **phytolith concentrations** varied from 2,343,877 to 950,881 phytoliths per 1 g of sediment (Fig. 5). The highest phytolith concentrations were found in Ap1 (2,343,877 phytoliths per gram of sediment). The lowest concentrations were found in Bwt (950,881 phytoliths per gram of sediment). *Elongated SUM* (mean = 22.6%, $\sigma = 2.9$, $n = 5$) are the most abundant **phytolith morphotypes**, followed by *Rondel* (mean = 19.2%, $\sigma = 0.6$, $n = 5$) and *Rondel elongated/wavy top* morphotypes (mean = 14.1%, $\sigma = 2.7$, $n = 5$). Another *Zea mays* diagnostic morphotype, *Rondel tall* (mean = 1.9%, $\sigma = 1.1$, $n = 5$), shows a significant occurrence, as well as *Polyhedral irregular SUM* (mean = 3.9%, $\sigma = 2.3$, $n = 5$) and *Perforated platelets* (mean = 0.4%, $\sigma = 0.2$, $n = 5$). *Bulliform cells* (mean = 1.1 %, $\sigma = 1.5$, $n = 5$) occur significantly in the Ah horizon while *Prickles* (mean = 3.6 %, $\sigma = 2.5$, $n = 5$) show a significant increase in Bwt. In the tillage horizon Ap2, two **starch grains** were identified, while the Bwt horizon revealed the presence of one starch grain, likely corresponding to *Zea mays* starch grains (Fig. 6f; (Gismondi et al., 2019).



460 **Figure 6: Photomicrographs of selected phytolith morphotypes identified in the samples. The photographs were taken at 400× magnification. (a) Rondel elongated; (b) Rondel wavy top; (c) Cross; (d) Rondel tall; (e) Spheroid echinate; (f) Starch grain from *Zea mays*.**

4.2.2 Santa María terrace system: overview and soil profiles Pe10-26 & Pe11-06

The archaeological site of Santa María (PAP-788) was catalogued in 2006 through surveys carried out by the Nasca-Palpa Archaeological Project. Based on surface ceramics and architecture, PAP-788 is a Middle Horizon (600–1000 CE) and Late Intermediate period (1000–1450 CE) settlement. The site also has an Inka Late Horizon (1450–1535 CE) component (Reindel and Isla, 2013a; Soßna, 2015).


The Santa María terrace system spans an area of approximately 42 ha (0.42 km²), with around 39 ha (0.39 km²) clearly identified as terraced agricultural land (Fig. 8). Mapping efforts have revealed a total of 692 terraces, collectively measuring 33.80 km in length. These identified terraces predominantly face west and exhibit a generally heterogeneous pattern. Larger terraces are observed in flatter areas, while smaller terraces are found on steeper slopes. The width of the terraces varies from about 2m to over 10m.

475 Soil profiles Pe10-26 (14°18'43.83"S, 74°51'46.35"W, 2750 m asl) and Pe11-06 (14°18'42.69"S, 74°51'49.66"W, 2739 m asl) are positioned approximately 3.5 km southwest of the village of Laramate, situated along a northwest-exposed debris cone (0.37 km², 2.55 km long, average slope: 6°) adjacent to the Laramate-Viscas River (Fig. 8). These profiles are located at upper and middle slope positions along a 100 m long, northwest-facing slope with an inclination of 19°.



480 The sample sites exhibit no signs of erosion since the abandonment of the terraces. Individual stones (6–20 cm) and boulders (20–60 cm) likely originate from collapsed terrace walls. The soil structure in both profiles is uniform in the upper 30–40 cm, characterized by coarse and blocky sub-angular peds with moderate to strong cohesion. Abundant roots with a diameter of 2–5 mm are present in the upper 40 cm of the profile.

485 In both cases, the soil type is classified as a *Terric Anthrosol (Loamic, Escalic)*. Within a recurring profile sequence (Ap-Ah-AC-C), up to two edaphic cycles could be identified. Minor differences in morphology between the two profiles primarily pertain to horizon thickness and the formation of horizon boundaries (Fig. 4, 9).



490 **Figure 7: Distribution of the most abundant phytolith morphotypes within the studied soil profiles in the study area.**





Figure 8: Detailed map of the Santa María terrace system.



Profile Pe10-26 (Supplementary Fig. 1) has a slightly compact, poorly developed and brownish-black 10 YR 2/3 **Ap** (0–15 cm) tillage horizon on its surface that contains human artefacts (Fig. 9). The underlying **Ah** (15–40 cm) is brownish-black in colour (10 YR 3/2) and has a higher humus content and a more developed soil structure. The horizon contains fragments of non-diagnostic pottery. It represents the former soil surface before terracing. The lower horizon boundary is very diffuse. The **AC** (40–60 cm) is a transitional horizon to the parent material, with a dark brownish-black (10 YR 2/3) colour and a very diffuse lower boundary. The horizon contains fragments of non-diagnostic pottery and charcoal particles and marks the lower limit of human influence on the soil. **C1** and **C2** (> 60 cm) consist of the parent material of the soil, which is slightly weathered quartzite. A 14C-dated charcoal taken from the **C1** horizon (60–100 cm) gave an age of 1650–1810 CE (Table 1, Fig. 9).



505 **Figure 9: Profile description and laboratory results for terrace soil Pe10-26: macro-morphology, pedo-chemical analytical data, AMS 14C dating and total phytolith amounts.**

Profile Pe11-06 has a slightly more compacted, less developed and brownish-black (10 YR 2/3) **Ap** (0–15 cm) tillage horizon on its surface, which contains human artefacts (Fig. 4, 10). The underlying **Ah** (15–30 cm) is brownish-black in colour (10 YR 3/2) and has a higher humus content and a more developed soil structure. It represents the former soil surface before terracing. The horizon contains large quantities of domestic waste including charcoal particles, bone remains in a poor state of preservation and undiagnostic pottery fragments. **AC** (30–45 cm) is a transitional horizon to the parent material. It is brownish-black brown (10 YR 3/4) in colour and has a very diffuse lower boundary. It contains a layer of charcoal particles at 40 cm depth and non-diagnostic ceramic fragments. The horizon marks the lower limit of human influence on the soil. **C** (> 45 cm) consists of the parent material of the soil, which is composed of slightly weathered quartzite. A 14C-dated charcoal taken from the **AC** horizon (30–45 cm) gave an age of 1310–1410 CE (Table 1, Fig. 10).



The **CECpot** values differ between sampling sites, both in terms of absolute amounts and behaviour in the profiles (Fig. 9, 10). While the values in Pe10-26 are lowest in Ap and Ah (8.8 and 9.6 cmol/kg) and increase slightly towards the bottom (up to 11.9 cmol/kg), the CECpot values in profile Pe11-06 are much higher and remain constant between 17.1 and 23.3 cmol/kg throughout the profile. Base saturation values are very high in all horizons at both sites, with an absolute minimum of 99.5%. **Soil pH** values are very similar in both profiles and increase with depth. Values range from moderately acidic at the surface to slightly alkaline in the soil parent material (pH 5.5 to 7.3 in Pe10-26; pH 5.9 to 7.4 in Pe11-06). **Organic matter** decreases gradually in both profiles, with values ranging from 1.9% to 0.5% in profile Pe10-26 and from 2.3% to 0.8% in profile Pe11-06. **C/N ratios** vary between the two sampling sites. The ratios in profile Pe10-26 are lower, with values ranging from 4.9 (Ah) to 7.3 (Ap). The ratios in profile Pe11-06 are significantly higher, with values between 12.8 (Ah) and 11.0 (AC). The Ap surface horizons of both profiles are significantly more compact than the underlying Ah horizons, as indicated by the **bulk densities** (Ap: 1.53 and 1.53 g/cm³ at Pe10-26 and Pe11-06 respectively; Ah: 1.35 and 1.31 g/cm³ at Pe10-26 and Pe11-06 respectively). The **packing density** coefficients of the topsoil horizons reach a value of 1.6 (Ap, Ah) in profile Pe10-26, while in profile Pe11-06 they show values of 1.7 (Ap) and 1.5 (Ah). The **soil texture** of both profiles is sandy loam with slightly varying clay and sand contents. In profile Pe-11 the clay content is generally slightly higher and the sand content slightly lower than in profile Pe10-26. The **skeleton fraction** marks a clear boundary between the Ap and the underlying Ah in both profiles. The percentage of the coarse fraction (> 2 mm) increases from 13.9 % (Ap) to 20.2 % (Ah) in Pe10-26 and from 13.3 % (Ap) to 14.7 % (Ah) in Pe11-06. In soil profile Pe11-06, the **plant available P** content remains consistently high at 1.7 mg/g in the Ap and Ah horizons and decreases to 0.9 mg/g in the AC.




Figure 10: Profile description and laboratory results for terrace soil Pe11-06: macro-morphology, pedo-chemical analytical data, AMS 14C dating and total phytolith amounts.



540

Phytolith concentrations in Pe10-26 vary considerably (Fig. 9). The Ap shows an amount of 4,119,018 phytoliths per 1 g of sediment while the lowest concentration is found in Ah (532,325 phytoliths per 1 g of sediment). AC (1,106,241 phytoliths per 1 g of sediment) and C1 (1,175,349 phytoliths per 1 g of sediment) show a similar concentration. With respect to **phytolith morphotypes**, *Rondels* (mean = 19.0 %, σ = 5.1, n = 4) are most abundant, followed by *Elongate SUM* (mean = 18.1 %, σ = 0.6, n = 4) and *Zea mays* diagnostic *Rondel elongated/wavy top* (mean = 17.3 %, σ = 2.6, n = 4) morphotypes. Other diagnostic *Zea mays* morphotypes such as *Rondel tall* (mean = 0.9 %, σ = 0.7, n = 4), *Bilobate long* (mean = 0.4 %, σ = 0.4, n = 4) and *Cross* (mean = 1.1 %, σ = 0.7, n = 4), show a detectable signal. *Spheroid echinates* (mean = 1.9 %, σ = 1.0, n = 4), *Acicular hair cells* (mean = 2.2 %, σ = 0.5, n = 4), *Prickle* (mean = 4.8 %, σ = 0.9, n = 4) and *Parallelepiped blocky SUM* (mean = 5.5 %, σ = 1.3, n = 4) also occur in significant numbers. In the tillage horizon Ap, a single **starch grain** was detected, likely corresponding to *Zea mays* (Gismondi et al., 2019). Conversely, no starch was observed in horizon C1.

Profile Pe11-06 shows the highest phytolith concentration in Ah (10,277,195 phytoliths per 1 g of sediment) and the lowest in AC (3,017,753 phytoliths per 1 g of sediment; Fig. 10). In the Ap horizon (0–15 cm depth) an abundance of 7,326,396 phytoliths per 1 g of sediment was observed. Phytolith assemblages from profile Pe11-06 show similar trends to profile Pe10/26. *Elongate SUM* (mean = 20.4 %, σ = 0.4, n = 3) are the most common morphotypes, followed by *Rondel* (mean = 17.5 %, σ = 3.9, n = 3) and *Rondel elongated/wavy top* (mean = 16.0 %, σ = 1.5, n = 3). In addition, the *Zea mays* diagnostic *Cross* (mean = 1.7 %, σ = 1.1, n = 3) and *Rondel tall* (mean = 0.5 %, σ = 0.2, n = 3) morphotypes occur significantly. *Spheroid echinates* (mean = 1.7 %, σ = 0.8, n = 3) are observed with similar frequency. *Acicular hair cells* (mean = 2.2 %, σ = 0.4, n = 3) and *Prickles* (mean = 4.8 %, σ = 1.2, n = 3) occur much more frequently than in profile Pe10/26.

560

4.2.3 Ayllapampa terrace system: overview and soil profiles Pe10-30 & Pe11-07

The archaeological site of Ayllapampa (PAP-967) was recorded in 2009 during surveys carried out by the Nasca-Palpa Archaeological Project. Based on surface ceramics and architectural remains, PAP-967 primarily dates to the Middle Horizon (600–1000 CE), but also has a Late Intermediate period (1000–1450 CE) component. PAP-967 shows the typical spatial layout of a Middle Horizon domestic settlement (SoBna, 2015).

The Ayllapampa terrace system encompasses an area of approximately 373 ha (3.73 km²), with around 191 ha (1.91 km²) clearly identified as terraced agricultural land (Fig. 11). A total of 2504 terrace walls have been revealed, collectively measuring 97.66 km in length. These identified terraces predominantly face west and exhibit a generally heterogeneous pattern. Larger terraces are observed in flatter areas, while smaller terraces are found on steeper slopes. The width of the terraces varies widely, ranging from about 2m to over 10m.

570



Figure 11: Detailed map of the Ayllapampa terrace system.



575

Profiles Pe10-30 (14°12'45.75"S, 74°53'12.35"W, 3410 m asl) and Pe11-07 (14°12'38.62"S, 74°53'10.79"W, 3400 m asl) are situated approximately 1.7 km northeast of the village of Pucará, on the central section of a west-facing debris cone (1.0 km², 4.5 km long, average slope: 17°) adjacent to the Palmadera-Palpa River (Fig. 11). Sampling was conducted at a mid-slope position on a 200 m long slope that faces northwest with an average slope of 18°.

580

The sample sites exhibit no signs of erosion since the terraces were abandoned. Isolated stones (6–20 cm) and boulders (20–60 cm) likely originate from collapsed terrace walls. The soil structure of profile Pe10-30 is uniform in the upper 65 cm, displaying a coherent massive structure with moderate cohesion. In contrast, the soil structure of Pe11-07 exhibits a coherent massive structure in the upper 25 cm and blocky subangular peds with moderate to strong cohesion at a depth of 25–50 cm.

585

Roots with a diameter of 2–5 mm are present in the upper 70 cm of the Pe10-30 profile, whereas slightly more roots are found in the upper 50 cm of profile Pe11-07.


590

In both cases, the soil type is classified as *Terric Anthrosol (Loamic, Escalric)*. However, despite their proximity and the shared parent material, Pe10-30 and Pe11-07 exhibit notable morphological differences in terms of profile thickness, horizon boundaries, and the extent of intermixed soil material (Fig. 4, 12, 13).

595

The Pe10-30 (Supplementary Fig. 1) soil sequence consists of a 100 cm thick brownish-black topsoil (10 YR 2/3) composed of four horizons (Ap1-Ap2-Ah-AC) formed on the parent rock (Fig. 12). The fine-grained soil matrix is very homogeneous and mixed at least to a depth of 65 cm. The two tillage horizons **Ap1** (0–8 cm) and **Ap2** (8–35 cm) are distinguished by a slightly higher skeletal content in Ap1. Ap2 also contains non-diagnostic pottery fragments. Its lower boundary is very diffuse. **Ah** (35–65 cm) contains large amounts of domestic debris: charcoal particles, bone remains in a poor state of preservation, obsidian point fragments and undiagnostic pottery fragments. It probably represents the pre-terraced ground surface, which also served as a settlement area. Its lower boundary is very diffuse. **AC** (65–100 cm) is a transitional horizon to the parent material, which has a higher content of coarse fragments, but at the base of which terrace building material has also been deposited. It marks the lower limit of human influence on the soil. **C** (> 100 cm) consists of the parent material of the soil, which is volcanic tuff. Two 14C-dated charcoals taken from the **Ah** (35–65 cm) and **AC** (65–100 cm) horizons gave ages of 770–990 CE and 670–860 CE respectively (Table 1, Fig. 12).

600



605 **Figure 12: Profile description and laboratory results for terrace soil profile Pe10-30: macro-morphology, pedo-chemical analytical data, AMS 14C dating's and total phytolith amounts.**

The soil sequence of Pe11-07 begins with a fine-grained, slightly compacted and brownish-black (10 YR 3/1) tillage horizon **Ap** (0–25 cm). The underlying buried **Ah** (25–50 cm) is more developed and black (10 YR 2/1). Among other human artefacts (e.g., obsidian point fragments and non-diagnostic pottery) it contains a layer of burnt charcoal particles at a depth of 35–40 cm. The horizon represents the former soil surface before terracing. **AC** (50–60 cm) is a brownish-black (10 YR 2/2) transitional horizon to the parent material. It contains fragments of obsidian points and marks the lower limit of human influence in the soil. **C** (> 60 cm) consists of the soil parent material, volcanic tuff. Unlike profile Pe10-30, the terrace has no construction or fill material at its base. The colour of the substrate is brownish-black (7.5 YR 3/2). A 14C-dated charcoal taken from the **Ah** horizon (25–50 cm) gave an age of 1210–1000 BCE (Table 1, Fig. 13).




Figure 13: Profile description and laboratory results for terrace soil profile Pe11-07: macro-morphology, pedo-chemical analytical data, AMS 14C dating and total phytolith amounts.

620

The **CECpot** values differ at both sampling sites (Fig. 12, 13). While the values in profile Pe10-30 are slightly higher and the variations are small (from 18.1 in Ap1 to 18.8 cmol/kg in AC, but 17.2 cmol/kg in Ap2 and Ah), the CECpot values in profile Pe11-07 show a clear peak in the Ah-horizon (15.2 cmol/kg); the values are lower both in the parent material (12.3 cmol/kg in C) and at the surface (13.6 cmol/kg in Ap). Base saturation reaches very high values in all horizons at both sites, with an absolute minimum of 98.9% (Ap) in profile Pe11-07. The **soil pH** is moderately acidic in both profiles. It varies between 5.1 and 5.7. **Organic matter** decreases significantly with depth in both profiles, from 6.5% (Ap1) to 2.2% (AC) in profile Pe10-30 and from 3.2% (Ap) to 1.0% (C) in profile Pe11-01. The **C/N ratios** are different at the two sampling locations. While the highest value in profile Pe11-07 is found in the Ah (12.3), the Ah in the Pe10-30 profile has one of the lowest values (8.9) within the profile. In general, the ratios in Pe11-07 are slightly higher than in the corresponding horizons in Pe10-30. In both profiles there is a tendency towards lower values in the Ap horizons compared to the underlying Ah horizons (from 8.1 to 8.9 in Pe10-30 and from 11.0 to 12.3 in Pe11-07). Soil compaction tends to increase slightly with depth in both profiles. Profile Pe10-30 has relatively constant **bulk densities** between 1.22 g/cm³ (Ap1) and 1.25 g/cm³ (AC). Profile Pe11-07, on the other hand, shows a much more compact Ap (1.32 g/cm³), while the underlying Ah horizon is less compact (1.22 g/cm³). The highest density (1.40g/cm³) was recorded in the C horizon. **Packing density coefficients** vary between 1.3 and 1.5 in both profiles. The **soil texture** is loam in all Pe10-30 horizons, with relative proportions of sand ranging from 45.5 % to 48.9 %. The soil texture in profile Pe11-07 varies between silty loam in the Ap, Ah and AC horizons and sandy loam with up to 60% sand in the underlying C horizon. The **skeleton fraction** of the soil marks a clear boundary between the tillage horizons and the underlying horizons in profile Pe10-30. While the skeletal fraction (> 2 mm) varies between 23.5 and 25.7 % in Ap1 and Ap2, it increases significantly to 38.9 and 35.4 % in Ah and AC. Although not as

630

635



640 pronounced as in profile Pe10-30, this pattern was also observed in profile Pe11-07, where the proportion of the coarse fraction increases from 22.0 % in the Ap to 26.9 % in the Ah horizon. The **carbonate content** is 0% in both profiles. The plant available **P** fraction remains low (0.1 mg/g) from Ap to AC and does not have measurable concentrations in the C horizon.

645 **Phytolith concentrations** vary considerably across the Ayllapampa site (Fig. 12, 13). In Pe10-30, the highest amount of phytoliths was found in Ap1 (2,446,706 phytoliths per 1 g of sediment), followed by the lowest amount in Ap2 (1,560,504 phytoliths per 1 g of sediment), while Ah (1,778,108 phytoliths per 1 g of sediment) and AC (1,791,331 phytoliths per 1 g of sediment) show similar concentrations. In terms of **phytolith morphotypes**, *Elongate SUM* (mean = 24.3 %, σ = 2.5, n = 4) are the most common morphotypes, followed by *Rondel* (mean = 19.7 %, σ = 1.1, n = 4) and the *Zea mays* diagnostic morphotype *Rondel elongated/wavy top* (mean = 13.4 %, σ = 3.2, n = 4). Another *Zea mays* diagnostic morphotype, *Cross* (mean = 0.7 %, σ = 1.1, n = 4), also shows a notable occurrence in Ap1 (Fig. 7).

Phytolith concentrations in profile Pe11-07 decrease slightly exponentially with depth (Fig. 13). The highest number of phytoliths was reached in Ap (2,357,810 phytoliths per 1 g of sediment), while the C horizon had the lowest concentrations (168,678 phytoliths per 1 g of sediment). In profile Pe11-07, *Rondel elongated/wavy top* (mean = 2.1 %, σ = 2.1, n = 4) morphotypes are rare (Fig. 7). *Rondels* (mean = 34.4 %, σ = 3.4, n = 4) are the most common, followed by *Elongate SUM* (mean = 11.5 %, σ = 3.1, n = 4). Compared to profile Pe10-30, other Poaceae short cells, such as *Trapezoid* (mean = 4.4 %, σ = 0.6, n = 4), *Polylobate* (mean = 9.1 %, σ = 4.0, n = 4) and *Trapezoid sinuate* (mean = 9.4 %, σ = 1.8, n = 4) can be observed more frequently, while *Cross* (mean = 0.0 %, σ = 0.1, n = 4) can be found in very low numbers in Ap. In AC, a significant occurrence of *Bulliform cells* (mean = 1.8 %, σ = 2.6, n = 4) and *Polyhedral irregular SUM* (mean = 3.2 %, σ = 3.8, n = 4) can be observed. *Weathered morphotypes* (mean = 16.2 %, σ = 6.4, n = 4) show an increasing concentration with depth.

4.2.4 Soil reference profiles Pe09-02 on plutonic acid & Pe10-32 on volcanic tuff

665 Profile Pe09-02 (14°16'22.98"S, 74°51'6.96"W, 3184 m asl) is located approximately 1.5 km northwest of the village of Laramate, situated within a modest south-facing catchment (1.8 km² and 6.2 km long) of a tributary to the Laramate-Viscas river system. Positioned in the middle of an 85 m long west-facing slope, the profile location features an inclination of 30°.

No signs of natural or anthropogenic soil erosion were observed at the profile location. Stones (6–20 cm) and cobbles (20–60 cm) are present in limited quantities on the surface. The soil structure is uniform in the upper 50 cm of the topsoil, but surface dry cracks (1–2 cm wide and 0.2–0.5 m apart) are commonly observed. Additionally, the topsoil exhibits coarse and prismatic peds with moderate to strong cohesion. The prevalence of roots with diameters ranging from 2 to 5 mm in the




upper 60 cm of the soil indicates recent biological activity. Bioturbation is evident through krotovinas, insect nests and burrows, as well as the presence of worms and larger animals.



Figure 14: Photographs of soil reference profiles Pe09-2 (Laramate) and Pe10-32 (Pucará).

680 The soil type is a *Someric Phaeozem (Clayic, Pachic)*. Four horizons (Ah1-Ah2-AC-Cw) can be distinguished (Fig 14, 15). **Ah1** (0–15 cm) is the first topsoil horizon, rich in humus and clay. Its brownish-black colour (7.5 YR 2.5/2) is due to fine-grained organic material. The lower boundary of the horizon is gradual and diffuse. The underlying **Ah2** (15–50 cm) differs slightly in its lighter brown colour (7.5 YR 3/3). Its lower boundary is diffuse and gradually merges into the transitional horizon **AC** (50–60 cm). The **C** (> 60 cm) represents the parent material of the soil, a granodiorite in a highly weathered state. The abrupt change in colour (10 YR 5/1) reflects the vertical boundary of pedogenesis.



685

Figure 15: Profile description and laboratory results for the soil control profiles Pe09-02 and Pe10-32: macro-morphology, pedo-chemical analytical data, AMS 14C dating and total phytolith amounts.

The **CEC_{pot}** values are elevated throughout the soil profile (32.8–21.6 cmol/kg), with the highest values being found in the topsoil horizons Ah1 and Ah2 (Fig. 15). Soil saturation reaches a maximum of 100% in all horizons. The **pH values** are in the moderately acidic range and vary minimally between 5.3 (Ah) and 5.1 (C). The **organic matter content** in the soil profile decreases from 2.1% in Ah1 to 1.6% in Ah2. The **C/N ratio** shows values between 9.7 and 10.7 in the topsoil horizons Ah1 and Ah2. As the **bulk density** increases from 1.37 to 1.56 g/cm³, horizon Ah1 becomes slightly looser than the underlying horizon Ah2. The compaction in Ah2 shows an incipient restriction of plant root growth. The **soil texture** is silty clay in both the uppermost Ah1 and Ah2 horizons, with clay contents of 42.8% and 42.1% respectively. The proportions of sand and silt fractions alternate as their total contents reach 32.1% and 25.1% in Ah1 and 28.3% and 29.6% in Ah2, respectively. The **skeleton content** of the topsoil remains constant with values of 7.6 % in Ah1 and 8.8 % in Ah2. The profile has a **carbonate content** of 0%. The plant available **P** fraction increases slightly from 0.1 mg/g in Ah1 and Ah2 to 0.2 mg/g in C.

700

Profile Pe10-32 (14°12'58.61"S, 74°52'17.49"W, 3790 m asl) is situated approximately 2.8 km northeast of the village of Pucará in an elongated and west-facing catchment (4.9 km² and 9.2 km long) of a tributary to the Palmareda-Palpa river system. Sampling was conducted in the lower part of an isolated, northeast-facing 5 m slope with an inclination of 3°.

No evidence of natural or man-made soil erosion was observed at the profile location. Stones (6–20 cm) and cobbles (20–60 cm) are present in limited quantities on the surface. The topsoil (0–20 cm) exhibits a single grain structure with very weak cohesion. The underlying horizons (20–50 cm) are characterized by granular and blocky subangular clods with moderate



cohesion. Roots with a diameter of 2–5 mm are abundant in the upper 30 cm, while a limited number of insect nests indicate low animal activity.

710

The soil type is a *Vitric Andosol (loamic)* with a sequence of five horizons (Ah1-Ah2-2Ahb-2Bwt-2C) containing a buried paleosol (Fig 14, 15). **Ah1** (0–5 cm) has a loose structure, a brownish black brown colour (10YR 3/4) and a high content of undecomposed plant material. **Ah2** (5–20 cm) is a fine-grained, loose and poorly developed topsoil, brown in colour (7.5 YR 4/4). A well-defined lower boundary with colour change marks a clear transition to the horizon below. The **2Ahb** horizon (20–30 cm) is a well preserved paleosol. It differs significantly from the topsoil horizons in its more developed and coherent soil structure, higher humus content and very dark brown colour (7.5 YR 2/3). The following subsoil horizon **2Bwt** (30–50 cm) shows signs of alteration as indicated by its colour (7.5 YR 5/6). Its lower boundary is gradual and diffuse. **2C1** and **2C2** (> 50 cm) consists of the soil parent material, slightly weathered volcanic tuff which marks the lower limit of pedogenesis. A ¹⁴C-dated bulk sediment sample taken from the **2Ahb** horizon (20–30 cm) gave an age of 2210–2030 BCE (Table 1, Fig. 15).

720

CEC_{pot} values in surface horizons Ah1 (7.7 cmol/kg) and Ah2 (6.1 cmol/kg) are higher than in the underlying 2Ahb (2.8 cmol/kg, Fig. 15). The values increase with depth to 11.7 cmol/kg in 2C. **Base saturation** is elevated throughout the soil profile (96.8–99.7 %) and shows a similar behaviour to the CEC_{pot} values. The **pH values** are in the strong to very strong acidic range, decreasing from 4.1 in the Ah1 and 2Ahb horizons to 3.3 in 2C. **Organic matter** decreases with depth from 7.5 % (Ah1) to 0.3 % (2C1), with the highest values found both in the topsoil (Ah1 and Ah2) and in the buried horizon (2Ahb). The **C/N ratio** decreases with depth, ranging from 9.5 (Ah1) at the soil surface to 3.4 (2C1) in the parent material. The surface horizons Ah1 and Ah2, with **bulk densities** of 1.06 and 1.01 g/cm³, are significantly looser than the underlying horizons 2Ahb and 2Bwt, with densities of 1.30 and 1.29 g/cm³. **Packing density coefficients** vary between 1.3 and 1.4. The **soil texture** is loam and silty clay in Ah 1 and Ah 2, with high silt contents between 39.1 % and 44.5 %. A marked decrease in silt content and a simultaneous increase in sand content to 69.6–60.2 % in the underlying 2Ahb, 2Bwt and 2C1 horizons characterize their texture as sandy loam. The **skeleton fraction** of the soil marks a clear boundary between topsoil and subsoil, as shown by the low percentage of only 1.9 % coarse fraction in Ah2, followed by an abrupt increase to 19.0 % in 2Ahb and 40.4% in 2C1. The profile has a **carbonate content** of 0%. Plant available **P** concentrations show very low peaks in Ah1 and 2Ahb and are not measurable in the rest of the profile.

735

Phytolith concentrations in profile Pe10-32 varied significantly from 11,925,773 to 18,690 phytoliths per 1 g of sediment, following an exponential decrease with depth (Fig. 15). The highest numbers of phytoliths were found in the upper Ah1 horizon (11,925,773 phytoliths per 1 g of sediment), while the lowest concentrations were found in the 2Bwt horizon (18,690 phytoliths per 1 g of sediment). In terms of **phytolith morphotypes**, *Rondels* (mean = 28.0 %, $\sigma = 5.9$, $n = 5$) are the most abundant morphotype and show a slight increase with depth, followed by *Elongate SUM* (mean = 23.2 %, $\sigma = 3.2$, $n = 5$),

740



745 which also decrease slightly with depth. In general, Poaceae short cells such as *Trapezoid* (mean = 3.6 %, σ = 3.6, n = 5), *Polylobate* (mean = 5.0 %, σ = 3.6, n = 5) or *Trapezoid sinuate* (mean = 10.5 %, σ = 2.7, n = 5) are common, whereas diagnostic short cells of *Zea mays*, such as *Rondel elongated/wavy top* (mean = 5.0 %, σ = 1.5, n = 5), *Rondel tall* (mean = 0.9 %, σ = 0.3, n = 5) or *Cross* (mean = 0.3 %, σ = 0.5, n = 5), are less frequent. *Weathered morphotypes* (mean = 10.9 %, σ = 4.4, n = 5) increase with depth. In horizon 2A_{hb}, no **starch grain** was detected.

5 Discussion

5.1 Reference Soil Groups in the Laramate region

750 In the following discussion, our focus will be on the WRB Reference Soil Groups, namely *Phaeozems*, *Andosols*, and *Anthrosols*. These soil groups are characteristic of the Laramate area and align with the soil profiles under examination.

5.1.1 Phaeozems

755 Mid-latitude *Phaeozems* are typically described as soils found in subhumid steppe margins within regions featuring vegetation ecosystems ranging from long grass steppe to forested steppe (Zech et al., 2014; IUSS Working Group, 2015). In the Laramate region, *Phaeozems* exhibit a distinctive Ah-(AC)-C profile sequence, marked by a dark and humus-rich surface horizon, significant organic matter accumulation in the mineral topsoil, and a notably high base saturation in the upper meter. While Laramate *Phaeozems* share similarities with well-described mid-latitude grassland steppe soils in their exceptional fertility, they deviate from the typical descriptions by not developing under subhumid conditions. Laramate *Phaeozems* are entirely devoid of carbonates throughout the profile, displaying no leaching features or accumulation of secondary carbonates. These soils originate from intensely weathered granodiorites and tonalites, as well as various
760 unconsolidated parent materials such as reworked volcanic conglomerates and alluvial deposits within the mountain shrubland of the Peruvian Andes. The absence of secondary carbonates precludes categorization as *Chernozems* or *Kastanozems*, related steppe soils. The diagnostic presence of subsoil secondary carbonates (*protocalcic* properties or *calcic* horizons) characterizes *Chernozems*, while *Kastanozems* are distinguished by greater amounts of secondary carbonates (IUSS Working Group, 2015).

765

A characteristic shared by all fertile steppe soils is the development of a *mollic* horizon in the advanced stages of pedogenesis (Eitel, 1999; Blume et al., 2010; Zech et al., 2014; IUSS Working Group, 2015). In profile Pe09-02, the presence of a *mollic* horizon serves as the diagnostic feature for classifying it as a *Phaeozem*. This near-surface mineral horizon is defined by a coherent and homogeneous substrate structure, an organic carbon content ranging from 0.8 to 1.0%, a
770 dark-coloured substrate with Munsell colour and chroma values of 3 or less, a total horizon thickness of 50 cm, and a base saturation of $\geq 50\%$. In the tropical Andes, the development of *mollic* horizons is influenced by seasonality, which dictates the alternation between two phases responsible for organic matter accumulation in the topsoil within an annual cycle. The



775 first phase involves the production of organic matter and the accumulation of undecomposed litter. The second phase is
linked to bioturbation and the intensive enrichment of humified organic matter (Eitel, 1999; Zech et al., 2014). In the
Laramate Region, the first phase coincides with a peak in soil moisture related to monsoonal rainfall activity during the
summer months. Humid conditions lead to rapid vegetation growth, including short and medium grasses, deciduous, and
xeromorphic shrubs, promoting the production of substantial biomass. The accumulation of undecomposed organic material
in the topsoil and upper subsoil layers predominates, although partial decomposition of the litter is stimulated by increased
microbial activity during the warm and humid summer periods (Eitel, 1999; Blume et al., 2010; Zech et al., 2014). The onset
780 of the dry season in April and May initiates significant changes in the physical, chemical, and biological dynamics of the soil
substrate. The colder temperatures during early winter months inhibit microbial activity and decomposition. Conversely,
grazing mammals, rodents, and invertebrates intensify their burrowing activity as a means to escape extreme environmental
conditions, including increased solar radiation in the absence of cloud cover, rapid nocturnal temperature drops, and frequent
frosts. This activity results in the displacement and mixing of undecomposed organic material from the topsoil into the
785 subsoil up to a depth of 1 m, where humification and some remineralization processes occur (Eitel, 1999). The presence of
krotovinas, insect nests, worm remains, and burrows of larger animals in the topsoil of the soils analysed in the Laramate
region attests to the intensive soil mixing by the local fauna. Facilitated by the loose soil structure, Andean mice (*Phyllotis*
spp.), Viscacha (*Lagidium peruanum*), Zari güeya (*Common opossum*), Andean fox (*Pseudalopex culpaeus*) and taruca
(*Hippocamelus antisensis*) are particularly effective in this process (Brack Egg and Mendiola Vargas, 2010). (Eash and
790 Sandor, 1995) identified analogous high-altitude soils in the Colca Valley, classified as *Mollisols* according to the US
classification of soil taxonomy. These soils exhibit similar characteristics of *mollic* horizons as observed in the Laramate
area. The authors also propose that slower decomposition of organic matter at lower temperatures at high altitudes and the
production of grass biomass play pivotal roles in pedogenesis and the development of high organic matter levels.

795 In the context of the prevailing aridity in the Laramate region, processes such as leaching and upward movement of water
play a minimal role in pedogenesis. Additionally, further profile differentiation of Laramate *Phaeozems* is infrequent due to
the heightened water-holding capacity of the topsoil horizons, facilitated by humic substances, which restricts percolation
and nutrient leaching. Furthermore, the presence of soil fauna contributes to the upcycling of non-weathered parent material
into the upper soil layers (Eitel, 1999).

800

While $\delta^{13}\text{C}$ (AMS) numeric values lack standardization, rendering them incomparable with other published data, their
relative variations appear to reflect hydrological changes during prehispanic times. The drought during the Middle Horizon
is suggested by less $\delta^{13}\text{C}$ depleted values (Table 1), while the lowest $\delta^{13}\text{C}$ values correspond to more humid periods,
possibly due to the adapted metabolism of CAM plants like *Prosopis juliflora*, which could have served as wooden fuel
805 sourced from local vegetation. However, this hypothesis warrants further investigation through IRMS measurements in
future studies.



5.1.2 Andosols

Different types of *Andosols* can be found in the southern Peruvian Andes, characterized by the volcanic source material (Zamora Jimeno and Bao Enríquez, 1972). In the Laramate region, *Andosols* typically develop on weathered volcanic material such as ashes, scoria, and vitric pyroclastics. Younger volcanic soils, identified as *Vitric Leptosols* or *Vitric Andosols*, are dominated by volcanic glass, glassy aggregates, and other glass-coated primary minerals. The presence of secondary minerals with higher concentrations of allophane clays and/or ferrihydrite indicates advanced stages of volcanic soil maturity, classifying them as *Andic Andosols* (Zech et al., 2014; Zamora Jimeno and Bao Enríquez, 1972). In the Laramate region, the occurrence of *Andic Andosols* is less frequent due to prevailing semi-arid conditions. In this area, the most common and extensive soil association comprises initial *Leptosols* over volcanic material, along with more developed *Vitric Andosols*. The Pe10-32 soil profile in this region can be categorized as a *Vitric Andosol*, as confirmed by $Al_{ox} + \frac{1}{2}Fe_{ox}$ values exceeding the minimum threshold of 0.4% and phosphate retention values reaching the minimum of 25%.

The *Andosol* in profile Pe10-32, situated on a gently sloping plain, exhibits a well-preserved soil-paleosol sequence associated with two edaphic cycles. A fine-grained and relatively poorly developed 'Ah' topsoil overlays a well-preserved '2Ahb-2Bwt-2C1-2C2' paleosol, offering insights into the paleoenvironmental evolution of the Laramate region. Drawing on earlier investigations and the examination of geoarchives in the Palpa region (Mächtle, 2007), peat sediments from Cerro Llamocca (Schittek et al., 2015), and paleosols in the Laramate area, it is postulated that there were approximately three pedogenesis phases in the study region during the Holocene. These phases were punctuated by two distinct dry periods between 2650–2250 BCE and 750–1200 CE (Fig. 16). The initial soil formation phase, characterized by heightened environmental humidity and geomorphodynamic stability, spanned from approximately 9050 to 2650 BCE and is deemed the most significant in terms of both duration and extent. It was during this phase that many of the *Phaeozems* and *Andosols* observed in the Laramate region formed. Subsequent soil formation phases were of shorter duration and lower intensity, making them difficult to detect.

The ^{14}C age of the fossil topsoil horizon '2Ahb' in Pe10-32, dated to 2210–2030 BCE, likely represents the final manifestation of in-situ pedogenesis in autochthonous material before the horizon was buried by finer-grained sediments. However, the exact initiation time of pedogenesis remains uncertain. Studies of paleosols in the Colca Valley suggest that pedogenesis commenced in the middle Pleistocene, coinciding with the stabilization of land surfaces and glaciers not extending below 4000 m asl (Eash and Sandor, 1995). (Kappl, 2012) investigated soil development along a Pleistocene ground moraine sequence in the nearby Puquio region, approximately 150 km east of Laramate. The author postulates that the formation of soils with *andic* characteristics in the study area spanned approximately 40,000 years with the youngest *Vitric Andosols* developing over surfaces that became stable around 30 and 23.7 ka BP.



840 In the case of profile Pe10-32, sedimentological data reveals a distinct boundary between the uppermost Ah horizons (Ah1, Ah2) and the underlying paleosol sequence. The topsoil horizons exhibit a notably loose structure, low bulk density values, minimal skeleton fraction proportions, and elevated levels of silt and clay. A comparison of the observed grain size composition with aeolian sedimentary archives from the Palpa region indicates similar granulometric properties. This suggests that the fine-grained sediments in Ah1 and Ah2 primarily consist of allochthonous material transported aeolian, likely originating from the loess archives of the western Andean foothills referred to as 'desert margin loess' (Eitel et al., 2005; Mächtle, 2007). It appears plausible that during pronounced dry periods in the Late Holocene, desert loess at the base of the Andes was carried by the wind and redeposited in the highlands. However, mineralogical analyses of sediment origins in the Rio Grande basin have indicated that aeolian sediments in the Laramate region could also originate from the Eastern Cordillera of Peru (Brenner, 2011).

850 5.1.3 Anthrosols

In the terrace profiles, the presence of tillage horizons overlaying buried paleosols resulted in specific profile sequences: 'Ap-Ah-Bwt-C1-C2' in profile Pe10-31 and 'Ap-Ah-AC-C' in profiles Pe10-26, Pe11-06, Pe10-30, and Pe11-07, classifying them as *Anthrosols*. Prolonged cultivation or the addition of materials can alter the properties of the topsoil horizons, but typically only to a certain depth. Consequently, the horizon differentiation of a buried soil may remain intact (IUSS Working Group, 2015). Morphological descriptions and analytical data indicate that all the investigated terrace soils are *Terric Anthrosols*, characterized by *terric* horizons with a total thickness of 50 cm. These *terric* horizons exhibit no clear stratification, possess a colour resembling the parent material, and demonstrate a base saturation of $\geq 50\%$. No other diagnostic horizons for *Anthrosols*, such as *irragric*, *anthraquic*, *hydragric* or *hortic pretic* were identified.

5.2 Characteristics and effects of terrace soil management

860 5.2.1 Soil texture, skeleton fraction and bulk densities

The terrace soils in the Ayllapampa and Santa María systems consistently exhibit a lower proportion of the soil skeleton fraction (up to 8% less) in the arable Ap horizon compared to the underlying buried paleosol. This reduction in the soil fraction is likely attributed to the deliberate removal of gravel and stones during terrace construction. This practice can offer economic advantages by enhancing the terrace's water holding capacity and reducing tillage resistance. This strategy aligns with the findings of (Sandor and Eash, 1995) in the Colca Valley, where farmers placing gravel masses at the forefront of agricultural terraces, possibly to promote water drainage.

The soil texture of the investigated terrace soils is defined by various loam types, creating optimal conditions for soil moisture retention (Blume et al., 2010; Zavaleta García, 1992; USDA-NRCS, 1998b), as documented in various regional studies (Sandor and Eash, 1995; Branch et al., 2007; Goodman-Elgar, 2008; Nanavati et al., 2016). Inherent site conditions,



notably the parent rock, contribute to minor distinctions between terraces. Although the water storage capacity of the Santa María terrace soils are comparatively lower, the Sihuilca and Ayllapampa terrace systems exhibit a slightly higher water storage capacity owing to elevated silt contents in the terrace fill deposits.

875 Concerning bulk densities, the terrace profiles show a distinct boundary between the tillage horizons (Ap) and the underlying topsoil horizons (Ah) of the buried soils, with higher bulk density values in the tillage horizons, as expected for long-term cultivated soils (Zavaleta García, 1992; Blume et al., 2010; USDA-NRCS, 1996b). Among all the studied terraces, the Santa María terraces display the most compact soils. The bulk densities of the topsoil horizons in profile Pe11-06 are even marginally higher than those in profile Pe10-26, potentially adversely affecting root growth. In contrast, the terrace soils in
880 Ayllapampa show minimal compaction, creating favourable conditions for root growth (USDA-NRCS, 1996b). Overall, the slight variations in bulk density do not suggest that the individual terraces served purposes beyond agriculture.

Drawing from various ethnohistorical sources and their own research findings, Goodman-Elgar (2008) and Nanavati et al. (2016) underscore the advantages of non-mechanized tillage tools, such as the *chaquitaclla*, which have been employed in
885 the Peruvian Andes since ancient times. These traditional tools, including the *ranccana* or *rucana*, facilitate soil cultivation with lesser destructiveness compared to industrial ploughing. It is reasonable to infer their utilization for the Laramete terraces.

5.2.2 Soil acidity, nutrient availability and soil quality

The studied terrace soils in the vicinity of Laramate respond in different ways to historical agricultural practices. Overall, the
890 region provides a conducive environment for agriculture. The soil's acidity and nutrient availability support the development of microorganisms, the establishment of robust roots, and the successful germination of seeds. This favourable environment facilitates the cultivation of staple Andean crops (Zavaleta García, 1992), including those identified in the region and associated with prehispanic agricultural systems, such as maize, potatoes, oats, alfalfa, and barley (Sandor and Eash, 1995; Goodman-Elgar, 2008; Kendall and Rodríguez, 2009; Nanavati et al., 2016; Handley et al., 2023).

895 The Sihuilca system stands out for its notably high soil quality. Notably, the pH of the tillage horizon (Ap) in profile Pe10-31 is markedly higher than the underlying Ah of the former soil surface and the pH values observed in the soil control profile Pe09-02. Additionally, the tillage horizon shows a substantial increase in available nutrients (Ca²⁺, K⁺, and Mg²⁺), a contrast that is significant compared to the underlying Ah horizon. When combined with other soil parameters such as
900 elevated total phosphorus and organic matter contents, these findings suggest that the Sihuilca terrace underwent intensive cultivation, directly meliorating the soil properties of the terrace.



Terrace profiles Pe10-30 and Pe11-07 in the Ayllapampa system differ significantly from their undisturbed reference profile Pe10-32. The observed variations in soil properties appear primarily attributed to local differences in the parent substrate on which the terrace soils and the reference soil developed (andesitic breccias and volcanic tuffs), rather than substantial alterations resulting from agricultural use. This assessment is grounded in the similarity between the buried paleosol sequences (Ah-AC-C) of the terrace soils and the characteristics of the cultivated tillage horizons (Ap). The soil quality of the tillage horizons is only marginally superior to that of the paleosols. In profile Pe10-30, the soil properties support agricultural use, as the tillage horizon (Ap) is less acidic and concurrently offers a higher quantity of nutrients compared to the underlying former soil surface (Ah). The significant increase in the base cation K^+ is particularly noteworthy. In contrast, the tillage horizon (Ap) of profile Pe11-07 demonstrates incipient soil acidification and reduced nutrient availability, marked by a clear loss of basic cations Ca^{2+} and Mg^{2+} .

The tillage horizons (Ap) within the Santa María terraces exhibit signs of incipient soil acidification and nutrient leaching, notably marked by a significant loss of alkaline Ca^+ cations. Soil acidification and nutrient leaching are common in various soil types under prolonged increases in soil moisture, such as those resulting from irrigation or more humid conditions (Zavaleta García, 1992). Lower pH values in the terraced topsoil horizons are also common in the Colca Valley and are probably related to higher levels of organic acids, carbonic acid or greater nitrification (Sandor and Eash, 1995). Given the prevailing arid conditions in the Laramate area, where soil acidification and nutrient losses are infrequently observed, the deficiencies noted in Santa María likely stem from the cultivation of the terrace system. It is plausible that livestock production in the region increased following the final abandonment of the agricultural terrace system, influencing soil pH conditions. Presently, the Santa María area is extensively utilized for livestock farming, with its gently sloping riverbanks and proximity to permanent water sources making it particularly appealing for grazing, especially during the dry season. Additionally, Handley et al. (2023) discovered, in the Chicha Soras terrace system in the neighbouring Apurimac region, that the nutrient content of the terraces was likely regularly sustained in prehispanic times through the addition of camelid dung, as indicated by the phosphate content of the terrace soils (Kemp et al., 2006). However, the Laramate area appears to have seen rare additions of organic nitrogen sources, such as urea or compost, as evidenced by the almost negligible total nitrogen contents.

5.2.3 C/N ratios, organic matter and organic phosphorus contents

C/N ratios in the Ap tillage horizons across all terraces reveal rapid mineralization rates, which differ from those observed in the underlying buried Ah horizons and their undisturbed soil context. These elevated mineralization levels are comparable to C/N levels found in terrace soils in the region (Sandor and Eash, 1995; Nanavati et al., 2016) and can be attributed to fertilization and selective cultivation.

In the Sihuilca terrace soils, both total carbon and total nitrogen content surpass the values of the *Phaeozem* reference profile, accompanied by relatively lower C/N ratios (7.5). The Santa María terraces, situated on crystalline parent material,



exhibit similarly low ratios (7.3-11.7) in the Ap horizons, yet possess higher total carbon and total nitrogen contents than the undisturbed *Phaeozem* reference profile. The increase in organic carbon and the greater amounts of nitrate in cultivated horizons can be attributed to long-term management practices involving the addition of organic matter and the cultivation of nitrogen-fixing crops (Sandor and Eash, 1995). Terraces located on volcanic soils within the Ayllapampa system display slightly higher C/N ratios (8.1-11.0) compared to the *Andosol* reference sequence and the terrace soils of the other systems. However, they have comparatively higher total carbon and total nitrogen contents. This difference may arise from the organic matter in these terraces not having fully decomposed, potentially associated with the spread of natural vegetation on less intensively cultivated land (Sandor and Eash, 1995).

The organic matter contents (1.9-3.5%) in the tillage horizons of all terraces are notably elevated, surpassing levels observed in both the undisturbed soil contexts and the underlying Ah horizons of the buried paleosols. Coupled with the absence of acidification and nutrient leaching, as well as the moderate to high C/N ratios observed in the studied terrace systems, these findings indicate generally favourable soil quality. They suggest that activities aimed at promoting the accumulation and maintenance of high organic matter content in the terrace topsoil were integral to prehispanic terrace management practices in the Laramate region. This aligns with findings from other studies on prehispanic agricultural terraces in neighbouring regions, such as the Colca Valley in Arequipa (Sandor and Eash, 1995) the Chicha Soras Valley in Apurimac (Branch et al., 2007; Handley et al., 2023) and the Viejo Sangayaico area in the upper Ica drainage (Nanavati et al., 2016). These studies propose that intensive agricultural use was accompanied by the systematic implementation of diverse fertilizing practices, including organic manuring, seasonal field burning, or manuring through grazing animals. While the terrace soils of the Colca Valley have undergone significant changes due to long-term fertilization since the Middle Horizon, leading to thickening of the soil and modification of its pedo-chemical properties, the terrace soils of Chicha Soras and Viejo Sangayaico seem to have been managed without heavy fertilization but rather through repeated additions of organic material. A study on Formative prehispanic agricultural soils in the Tafi Valley in the Argentinean region of Tucumán provides insights into the effects of long-term tillage and the use of fertilizers to compensate for soil deficits (Vattuone et al., 2011). Organic fertilization appears to have been a common prehispanic practice in the semi-arid Andes. This practice aimed to counteract the potential decline in organic matter and consequent deterioration in soil quality resulting from continuous cultivation and harvesting. The increased aeration of soils through tillage, coupled with the disruption of aggregates, exposes organic matter to aerobic conditions after tillage, stimulating microbial activity and accelerating its decomposition (Stevenson, 1982). Over time, tillage and the depletion of organic matter result in nutrient loss, increased compaction, diminished water infiltration capacity of the soil, and further erosion of soil cover (USDA-NRCS, 1996a).

The decomposition of organic matter in the soil releases nitrogen, carbon and organic phosphorus (P_{org}). Under optimal moisture conditions, carbon and nitrogen become rapidly available for plant uptake, while P_{org} remains stable in an unavailable fraction (Buckman and Brady, 1977). The accumulation of P_{org} in the soil substrate may thus indicate practices



970 involving the continuous and long-term incorporation of biomass (organic fertilization) into the soil to sustain consistently
high levels of organic matter content (Vattuone et al., 2011; USDA-NRCS, 1996a, 1998a). The substantial buildup of P_{org} in
the Santa María and Ayllapampa terraces suggests prolonged agricultural use with ongoing application of organic manure.
This interpretation is reinforced by increased bulk densities in the tillage horizons (A_p) of the terraces, particularly those
with high P_{org} contents (e.g., terrace soils Pe11-06 and Pe11-07). The observation that the younger terrace system of Santa
975 María, dating to the Late Intermediate period, exhibits a stronger P_{org} signal than the older terrace system of Sihuilca, dating
to the Early Horizon, could have multiple causes. On one hand, the Sihuilca system is situated in an area where terrace soils
formed on *Phaeozems* with excellent agroecological characteristics, potentially requiring little artificial improvement of soil
properties through fertilization. On the other hand, the Santa María system likely saw more intensive terrace cultivation
during a cultural period when such practices were more prevalent.

980

Moreover, the replenishment of soil nutrients through seasonal burning of fields is highlighted as a prehispanic agricultural
practice in the region (Nanavati et al., 2016; Handley et al., 2023). However, the limited presence of charcoal particles
within the Laramate terraces and the absence of burnt soil layers hinder the reliable attribution of charred material to a
specific soil horizon or cultural period. Consequently, there is no evidence supporting the notion that nutrient levels were
985 sustained through seasonal burning in the Laramate region.

5.2.4 Phytolith concentrations and assemblages

The phytolith concentrations indicate clear soil mixing in profiles Pe10-26 and Pe11-06 from Santa María, Pe10-30 from
Ayllapampa, and Pe10-31 from Sihuilca. This is evident as the concentrations are more uniformly distributed within the
profile and are notably high at depth. Consequently, it can be inferred that the sampled terraces have been utilized for
990 agricultural activities over an extended period, as the highest phytolith concentrations in natural soils typically occur in the
upper humic horizon and decrease with depth (Qader et al., 2023; Strömberg et al., 2018). Interestingly, the terrace profiles
from Santa María exhibit distinct differences. The concentrations in profile Pe11-06 are markedly higher than those in Pe10-
26, despite both profiles being in close proximity to each other and at a similar distance from a settlement. The
concentrations in both profiles suggest intensive terrace use, hinting at longer use for Pe11-06. However, it is worth noting
995 that the C1 horizon (60-100 cm depth) of profile Pe10-26 appears to be approximately 400 years younger than the AC
horizon (30-45 cm depth) of Pe11-6, introducing a potential contradiction to this theory (Table 1).

The situation differs in terrace profile Pe11-07 from Ayllapampa, where concentrations are high in the A_p horizon but
sharply decrease with depth (Fig. 13). This mirrors the soil control profile Pe10-32 (Fig. 15), devoid of terracing and likely
1000 agriculture, suggesting that the Pe11-07 terrace might not have been used extensively for agriculture over an extended period
(Strömberg et al., 2018; Qader et al., 2023). In contrast, the phytolith concentrations of terrace profile Pe10-30 exhibit clear
signs of soil mixing, indicating agricultural use (Fig. 12). This interpretation finds support in the higher abundance of C3



Pooideae diagnostic Grass Silica Short-Cell Phytoliths (GSSCP) such as *Rondel*, *Trapezoid*, *Polylobate* and *Trapezoid Sinuate* in profiles Pe10-32 and Pe11-07 (Fig. 7). Compared to other sites, the abundance of C3 Pooideae GSSCP is significantly higher. This grassland community is typical in environments of high elevations, increased moisture, and lower temperatures (Twiss, 1992; Bremond et al., 2008; Aleman et al., 2014). Given the proximity of Pe10-32 to the Ayllapampa system, the similarity of phytolith assemblages could be explained, among other factors, by their close location. Assuming that the terrace of profile Pe11-07 has not been used extensively for a long time, the natural vegetation of the surrounding area might have reestablished itself on the site. This interpretation is supported by the radiocarbon dating of the Ah horizons of both terrace soils. The Ah horizon of profile Pe11-07, situated beneath the tillage horizon, dates back to the Initial Period (1210–1000 BCE). Given that the subsequent Paracas period signifies the inception of the first agricultural terrace infrastructure, it is reasonable to assume that the terrace was constructed and utilized during this period, a hypothesis substantiated by the terrace architecture. In contrast, the Ah horizon of profile Pe10-30 is dated to the Middle Horizon (770–990 CE), the period of systematic expansion of terrace agriculture. The dating of terrace soil Pe10-31 from Sihuilca also falls within this period (Fig. 16). The similarity between these two profiles is evident not only in the phytolith concentrations but also in the phytolith assemblages, suggesting a similar use of the two terraces, e.g., maize cultivation. In contrast, the terrace of profile Pe11-07 may have been unused since its potential initial construction and use in the Paracas period. It is plausible that, for various reasons, some terraces were no longer used for agricultural purposes, while others continued to be cultivated or were recultivated.

1020

A notably higher occurrence of the diagnostic morphotype *Rondel elongated/wavy top* associated with *Zea mays* is evident at the other sites (Fig. 7). This could be linked to maize cultivation and the agricultural use of the terraces (Bozarth, 1998; Pearsall et al., 2003; Piperno, 2006; Logan et al., 2012; Handley et al., 2023). Generally, there is a greater presence of phytoliths from maize cobs (*Rondel elongated/wavy top*) compared to those from leaves and husks (*Crosses*) in the profiles.

1025

Interestingly, both Santa María profiles (Pe10-26 and Pe11-06) show a distinct signal of *Cross*-shaped phytoliths, possibly originating from maize (Handley et al., 2023; Piperno, 2006). Andean maize typically produces medium-sized *Crosses* and *Long Bilobates* (Logan et al., 2012), both of which are present in profile Pe10-26 from Santa María. The radiocarbon dates from Santa María suggest a prolonged period of terrace use or recultivation, with potential changes in cultivation or fertilization practices over time. In contrast to Sihuilca or Ayllapampa, where leaves and glumes seem to have been mostly removed, there is a clear presence of *Cross* morphotypes indicative of maize leaves or glumes in the Santa María terrace profiles (Fig. 7). Elevated phosphorus values in Pe10-26 and Pe11-06 also stand out compared to other profiles, potentially indicating fertilization with organic material like maize leaves or corncobs. Another possibility is fertilization with camelid dung or the grazing of camelids on the terraces. Since maize could have been a component of the camelid diet, the presence of leaf phytoliths, especially *Cross*-shaped morphotypes, might suggest such a practice (Handley et al., 2023; Finucane et al., 2006; Cadwallader et al., 2012). However, it's essential to note that the *Cross*-shaped morphotype alone cannot definitively

1035



1040 identify maize, as other grasses of the Panicoideae family also produce such phytoliths (Logan et al., 2012). Therefore, a more precise method, such as discriminant function analysis, would be needed to distinguish *Cross*-shaped phytoliths from *Zea mays* and determine their specific origin (Piperno, 2006). *Cross*-shaped phytoliths from other Panicoideae could also have been part of the camelid diet or used as fertilizer.

1045 Surprisingly, *Spheroid echinate* phytoliths, considered diagnostic for Arecaceae (palms) (Huisman et al., 2018; International Committee for Phytolith Taxonomy (ICPT), 2019), were discovered in both profiles from Santa María (Pe10-26, Pe11-06), while they were present in very low abundance in the soil control profile Pe10-32 and terrace profile Pe10-30 from Ayllapampa (Fig. 7). Palms typically thrive in lower, warmer, and moister forests east of the Andes in Peru (Henderson et al., 2019; Huisman et al., 2018). However, (Handley et al., 2023), working in the nearby Chicha-Soras Valley, also found *Spheroid echinate* phytoliths in similar concentrations in their samples. They suggest that these palm remnants may result from human cultivation or the transport of plant parts such as fruit, leaves, or wooden tools. In comparison to the Arecaceae phytoliths described by (Huisman et al., 2018), the *Spheroid echinate* phytoliths in our study (Fig. 6e) resemble the forms of 1050 *Ceroxylon alpinum* (seed and wood). Nonetheless, there is the potential for confusion with other plant families producing *Spheroid echinate* morphotypes, such as Bromeliaceae. Common genera such as Tillandsias belong to the Bromeliaceae. They are found in the nearby Lomas ecosystem but are also seasonally and locally established in the adjacent Cactus Belt (e.g. supported by various cacti species on the slopes or by *Orthopterygium huaucui* in lower valleys) (Mächtle, 2007; Brack Egg and Mendiola Vargas, 2010; MINAM, 2015; Beresford-Jones et al., 2015). Due to a possible overlap in size, *Spheroid* 1055 *echinates* cannot be conclusively assigned to either Arecaceae or Bromeliaceae, warranting further research (International Committee for Phytolith Taxonomy (ICPT), 2019; Piperno, 1988).

5.2.5 Starch analyses

1060 The identification of *Zea maize* starch grains (Gismondi et al., 2019) in the tillage horizons of terrace profiles Pe10-31 in Sihuilca and Pe10-26 in Santa María aligns with the findings from phytolith analyses. Consequently, it suggests that maize held significant agricultural importance in the region during prehispanic times. The occurrence of starch in the Bwt horizon of profiles Pe10-26 is probably attributed to soil mixing linked with terrace utilization. Conversely, the lack of starch in the C1 horizon of terrace profile Pe10-31 indicates that the soil mixing process did not extend to the parent material. The relatively low starch content in the terrace soils is likely a consequence of unfavourable conservation conditions stemming from starch degradation and damage inflicted by soil bacteria (Hutschenreuther et al., 2017). The lack of maize starch in the 1065 control profile Pe10-32 corroborates the presumption that maize cultivation did not take place in this location.



5.3 Prehispanic history of terraced agriculture in the Laramate region

Drawing on previous research and the findings of our study, the prehispanic history of terraced agriculture in the Laramate region can be broadly delineated into four phases, encompassing the initial establishment, systematic expansion, and eventual abandonment of the agricultural infrastructure at the beginning of the Hispanic colonial period (Fig. 16).

1070




Figure 16: Chronological model for the emplacement and use of the Laramate terrace cultivation system and its paleo-environmental context. Sources: relative chronology modified from Reindel and Isla, 2013a; Sořna, 2015; paleo-environmental and paleo-pedological records modified from Eitel et al., 2005, Mächtle 2007, Schittek et al. 2015 and own sampling; terrace records modified from Sořna, 2015, Mader et al. 2024 and soil records from the current paper.

1075

(1) Initial phase of agricultural terracing in the Formative Late Paracas and transitional Early Nasca periods

1080

Archaeological evidence from the Laramate region indicates that the initial construction and use of agricultural terraces took place during the Formative Late Paracas period (Reindel and Isla, 2013a; Sořna, 2015). Recent geoarchaeological



investigations at the archaeological site of Cutamalla and the surrounding extensive agricultural terrace systems, spanning approximately 220 hectares, revealed a contemporaneous occupation of Cutamalla alongside the utilization of the terrace systems. This agricultural terrace-settlement system was intensively employed for a relatively brief period of about 200 years (~250–40 BCE) during the Formative Late Paracas and transitional Early Nasca periods, with no evidence of reoccupation or
1085 subsequent reuse of the agricultural systems. The absence of identified irrigation canals suggests rainfed agriculture. The architecture of the agricultural terraces in Cutamalla is notably similar and simple: the terrace walls with medium height were constructed using fairly large stones and boulders (up to 0.5 m in diameter), representing a typical architectural pattern of Formative agricultural terraces in the region (Mader et al., 2024).

1090 Most of the terraces near the archaeological site of Sihuilca share a similar architectural style with those of Cutamalla. This resemblance implies their initial construction during the Late Paracas and/or transitional Early Nasca periods, a premise bolstered by the dating of the architectural remains of the PAP-888 settlement to the Nasca Early Intermediate period. However, unlike Cutamalla, modifications involving the use of smaller stones on some terrace walls were noted at Sihuilca, particularly in the upper slope areas. This observation suggests that parts of the terrace system were subject to reuse and
1095 modification during later settlement periods, a hypothesis supported by the ¹⁴C dating of terrace soil Pe10-31, which yielded an age between 880–990 CE, corresponding to the Middle Horizon.

Typical sites for these early terrace systems are usually those that maintained constant and high soil moisture during construction and use, particularly those exhibiting exceptional agro-ecological soil quality, as found in the soil province of
1100 *Phaeozems*. The Late Paracas and Initial Nasca periods were not only highly dynamic with significant population growth but also experienced more humid environmental conditions (Mächtle, 2007; Schitteck et al., 2015), facilitating the widespread adoption of these agricultural practices in the region (Mader et al., 2024). Unfortunately, the conservation status of these old terraces is often poor due to extended periods of disuse.

1105 (2) *Systematic expansion of terrace agricultural systems in the Middle Horizon*

The widespread and systematic expansion of terrace agriculture in the southern Peruvian Andes during subsequent prehispanic periods, especially in the Middle Horizon (600–1000 CE) and influenced by the Wari culture, has been extensively documented. Parallel developments have been observed in various regions, such as the Colca Valley in Arequipa
1110 (Sandor and Eash, 1995), the Chicha Soras Valley in Apurimac (Branch et al., 2007; Handley et al., 2023), and the Viejo-Sangayaico area in the upper Ica catchment (Nanavati et al., 2016).

In the upper Rio Grande basin, the expansion of agricultural systems linked to the Wari culture has been documented, focusing on the nearby Andamarca terraces (Aguirre-Morales, 2009). This trend is also noticeable in the Laramate area,



1115 especially around the Ayllapampa site. With the transition to drier environmental conditions and the gradual shift of the
desert margin to the east, the importance of soil moisture for the construction and utilisation of terrace systems became
crucial. Preferred sites, including some under irrigation, included the adjacent banks of major watercourses and areas with
reduced solar radiation. Rather than solely depending on areas with *Phaeozems* (nutrient-rich soils), terraced agriculture on
volcanic soils (*Andosols*) at higher elevations became more widespread. Radiocarbon dating findings from the Ayllapampa
1120 Pe10-30 terrace profile, indicating ages between 670–860 CE and 770–990 CE, provide support for this hypothesis. The
construction pattern of Ayllapampa terraces is distinct, featuring large stones at the base that, through subsequent
modifications with smaller stones (as seen in Pe10-30), can reach medium to high wall heights of up to 1.3 meters. This
suggests a later utilization of the terraces, likely extending until the Late Intermediate Period.

1125 (3) *Terrace expansion to lower altitudes and less agriculturally suitable locations during the Late Intermediate Period*

The Late Intermediate period (1000–1450 CE) was characterized hydrologically highly variable conditions, featuring phases
of both dry and humid conditions (Eitel et al., 2005; Mächtle, 2007; Schitteck et al., 2015). Particularly noteworthy were the
positive effects of the humid conditions after 1250 CE on settlement activities and population growth in the Laramate region
1130 (Soßna, 2015). This period witnessed advancements in agrarian techniques and a substantial expansion of sophisticated
farming terraces, as observed also in neighbouring regions (Sandor and Eash, 1995; Branch et al., 2007; Handley et al.,
2023; Nanavati et al., 2016). As the desert fringe receded westward across the Cordillera Occidental (Eitel et al., 2005;
Mächtle, 2007; Schitteck et al., 2015), the adoption of terracing extended to lower elevations. Additionally, the heightened
humidity facilitated the utilization of areas that were previously deemed less suitable for agriculture.

1135 A striking illustration of the effective expansion of cropland in regions with less favourable agro-ecological conditions
through terracing is the Chaupicancha system (Fig. 1). Presently, the system relies on seasonal water supply from two
springs activated during the rainy season, though the total number of active springs was likely higher during the Late
Intermediate period. The consistent availability of soil moisture compensated for deficiencies in soil quality (shallow topsoil,
1140 lower nutrient availability, and organic matter, low water retention capacity) and the somewhat unfavourable geomorphology
of the site, marked by steep slopes and high solar radiation (Supplementary Fig. 2; Supplementary Table S1). The distinctive
architecture of the terraces during this period, characterized by small terrace wall stones, and the absence of buried tillage
horizons, indicate that the Chaupicancha terraces were initially built during this period and do not represent a re-use of older
terraces. The Colca Chica terrace group, situated at the westernmost edge of the study area at lower altitudes (Fig. 1),
1145 exhibits similar characteristics. In this case as well, an area with unfavourable agro-ecological conditions was initially built
and utilized during the Late Intermediate period.



Furthermore, some older agricultural terrace systems saw intensive agricultural activity during the Late Intermediate period. The Santa María terrace system, for example, displays indications of repeated use and abandonment spanning multiple
1150 cultural periods. This is substantiated by observed modifications to the terrace structure, the archaeological settlement remains representing various ages, and the radiocarbon ages of the studied terrace soils, ranging from 1310–1410 CE in terrace profile Pe11-06 to 1650–1810 CE in profile Pe10-26. The Inka state of the Late Horizon (1450–1535 CE) had a minor presence in the Laramate region, however, Santa María is one of the few sites with remaining Inka architecture (Soßna, 2015).

1155

(4) *Final abandonment of the terrace agricultural systems at the onset of the colonial period*

Terraces gradually fell into disuse and lacked systematic maintenance after the Late Intermediate period contemporaneous to a rapid decrease in rainfall (Schittek et al., 2015). With the advent of the Hispanic colonial period in 1532 CE, the local
1160 population was relocated, and the production system underwent reorganization (Aguirre-Morales, 2009; Nanavati et al., 2016). Moreover, introduced diseases led to a demographic decline. This marked the beginning of the ultimate abandonment of the Laramate terrace complex. During the Spanish colonial administration, the terraced farming system lost its prominence, resulting in the terraces no longer being systematically used for agriculture.

5 Conclusion

1165 In conclusion, our integrated pedo-geoarchaeological study has centered on three abandoned terrace agricultural systems and their paleo-pedological context in the vicinity of Laramate, situated in the southern Andes of Peru (14.5°S). The principal aim of this investigation was to untangle the pedological and land-use history of the Laramate area, which served as a significant agricultural hub during prehispanic times. This encompassed two key objectives: 1) contextualizing the former agricultural management system within its paleoecological framework and 2) assessing the impact of agricultural practices
1170 on soil development and quality. The Laramate terrace complex emerged as an optimal setting for our inquiry, given its diverse array of terrace installations and exploitation ages, coupled with varied geomorphological and geological settings. To evaluate the enduring effects on mountain soils resulting from ancient agricultural practices, a comparative analysis was conducted by comparing non-irrigated agricultural terrace soils with their undisturbed paleo-pedological counterparts. This comprehensive examination integrated a range of methodologies, including surveys, soil science techniques, GIS and remote
1175 sensing applications, paleobotany, and radiocarbon dating.

The Laramate region is characterized by three WRB Reference Soil Groups, specifically *Phaeozems*, *Andosols*, and *Anthrosols*. Notably, Phaeozems in this region exhibit a deviation from typical descriptions as they do not develop under subhumid conditions. The formation of their *mollic* horizon is strongly influenced by various factors, including climatic



1180 seasonality, dominant herbaceous vegetation, burrowing fauna, local lithology, and inputs of allochthonous aeolian material. In areas with volcanic material, the prevalent soil association consists of *Leptosols*, along with more developed *Vitric Andosols*. The fossil topsoil in Pe10-32, dating back to 2210–2030 BCE, represents the final stage of in-situ pedogenesis. The overlying fine-grained sediments consist of allochthonous material that was deposited after aeolian transport as a loess cover.

1185 Detailed descriptions and analytical data suggest that all the terrace soils investigated in the area can be classified as *Terric Anthrosols*. Examining soil quality indicators of terrace soils reveals no significant signs of severe degradation, even with long-term use. The soil's balanced acidity and nutrient availability contribute to the successful cultivation of staple Andean crops. The final abandonment of the cultivation system is not attributed to soil exhaustion or terrace structural instability, 1190 despite facing phases of landscape instability during dry periods.

The Sihuilca terrace system stands out for its notably high soil quality, evident in improvements such as increased water storage capacity and optimal soil acidity levels. The soil quality of the Ayllapampa terrace system is relatively neutral, with tillage horizons exhibiting a marginal superiority over underlying paleosols, as demonstrated by minimal compaction, lower 1195 acidification, and a higher quantity of available nutrients. In contrast, terrace soils at Santa María show signs of incipient deterioration; slight acidification, nutrient leaching, reduced water storage capacity, and compaction of cultivable horizons are evident in this area.

The tillage horizons of all terraces exhibit significantly elevated organic matter contents, surpassing both undisturbed soil 1200 contexts and buried paleosols. This suggests that the sustainability of prehispanic agricultural practices in the Laramate region was closely tied to maintaining high organic matter content in the terrace topsoil. The substantial accumulation of P_{org} indicates prolonged agricultural use with a consistent application of organic manure. Notably, there is no evidence supporting the idea that nutrient levels were maintained through seasonal burning in the Laramate region. Additionally, irrigation played a minor role in these agricultural terrace systems.

1205 Phytolith concentrations in all terrace profiles strongly indicate soil mixing, signifying intensive agricultural activities over an extended period on the terraces. The presence of the diagnostic morphotype *Rondel elongated/wavy top*, associated with *Zea mays*, is more pronounced in terrace systems with presumed long-term agricultural use, emphasizing a connection to maize cultivation. Notably, there is a higher abundance of phytoliths from maize cobs (*Rondel elongated/wavy top*) compared to those from leaves and husks (*Crosses*). The latter, prominently found in both Santa María profiles, suggests 1210 prolonged terrace use or recultivation, hinting at potential changes in cultivation or fertilization practices over time. The identification of *Zea maize* starch grains in the tillage horizons at Sihuilca and Santa María aligns with phytolith analyses, reinforcing the significance of maize in the region during prehispanic times.



In conclusion, the prehispanic history of terraced agriculture in the Laramate region unfolds across four distinctive phases, each reflecting the dynamic interactions between environmental, cultural, and agricultural factors. The initial phase, during the Formative Late Paracas and transitional Early Nasca periods, witnessed the establishment of agricultural terraces exemplified by sites like Cutamalla and Sihuilca. This humid period of intensive agricultural use is characterized by rainfed agriculture and simple terrace architecture. The systematic expansion of terrace agricultural systems marked the Middle Horizon (600–1000 CE), influenced by the Wari culture. The transition to drier environmental conditions prompted adaptations, such as terraced agriculture on volcanic soils as observed in the Ayllapampa terrace system. During the Late Intermediate Period (1000–1450 CE), characterized by hydrological variability, terraced agriculture experienced further expansion to lower altitudes and less agriculturally suitable locations. The Chaupicancha system serves as a notable example of effective cropland expansion in challenging conditions, reflecting more humid conditions and advancements in agrarian techniques. The final phase marked the gradual abandonment of terraced agricultural systems after the Late Intermediate period, particularly with the onset of the Hispanic colonial period in 1532 CE. Demographic shifts and reorganization of production systems led to the loss of prominence for the terraced farming system in the Laramate region, ultimately resulting in the cessation of systematic agricultural activities on the terraces. This historical trajectory underscores the adaptability and resilience of prehispanic communities in the Laramate region, who innovatively employed terraced agriculture to navigate changing environmental conditions and sustain agricultural practices across diverse landscape units.

1230 **Data availability statement**

Data used in the study are available in the article and Supplement. Inquiries can be directed to the corresponding authors.

Author contributions

FL: conceptualization, fieldwork, methodology, data acquisition, data curation, formal analysis, data interpretation, visualization, funding, writing and editing – original draft, review and editing; CB: formal analysis, data interpretation, visualization, writing and editing – original draft; CM: methodology, funding, writing and editing – original draft; BM: conceptualization, fieldwork, methodology, data interpretation, writing – review and editing; EM: methodology, formal analysis, statistical analysis, writing – review and editing; LD: methodology, formal analysis, data interpretation; MR: funding, resources; BE: conceptualization, fieldwork, methodology, data interpretation, funding, resources, writing – review and editing; JM: conceptualisation, methodology, data interpretation, funding, resources, writing and editing – original draft, review and editing.



Competing interests

The contact author has declared that none of the authors has any competing interests.

Disclaimer

1245 Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Financial support

1250 Fernando Leceta was financially supported by the German Academic Exchange Service (DAAD; Ref. Number A/08/98226) and Heidelberg University. Moreover, this research was funded by the Groundcheck Program of the German Archaeological Institute (DAI). This publication was supported by the Open Access Publication Fund of the University of Würzburg.

Acknowledgments

1255 Fernando Leceta would like to thank the German Academic Exchange Service (DAAD) for financial support (DAAD Ref. Number A/08/98226), the soil laboratory staff at the University of Heidelberg, led by Gerd Schukraft (T), Nicola Manke and Adnan Al Karghuli, as well as Dr. Volker Soßna for the exchange of ideas and support during the field campaigns. We would also like to thank Santiago Ancapichún for calculating the air mixture at Cutamalla.

References

- Aguirre-Morales, M.: Excavaciones en los andenes de Andamarca, cuenca del río Negromayo, Lucanas, Ayacucho, *Arqueología y sociedad*, 223–268, 2009.
- 1260 Albert, R. M., Lavi, O., Estroff, L., Weiner, S., Tsatskin, A., Ronen, A., and Lev-Yadun, S.: Mode of occupation of Tabun Cave, Mt Carmel, Israel during the Mousterian Period: a study of the sediments and phytoliths, *Journal of Archaeological Science*, 26, 1249–1260, <https://doi.org/10.1006/jasc.1999.0355>, 1999.
- Aleman, J. C., Canal-Subitani, S., Favier, C., and Bremond, L.: Influence of the local environment on lacustrine sedimentary phytolith records, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 414, 273–283, <https://doi.org/10.1016/j.palaeo.2014.08.030>, 2014.
- 1265 Ancapichún, S., Pawlyta, J., Rakowski, A. Z., and Siczekowska, D.: Influence of Air Parcels from Northern and Southern Hemispheres on Radiocarbon-Based INCA Chronology, *Radiocarbon*, 64, 1431–1446, 2022.
- Beresford-Jones, D., Pullen, A. G., Whaley, O. Q., Moat, J., Chauca, G., Cadwallader, L., Arce, S., Orellana, A., Alarcón, C., Gorriti, M., Maita, P. K., Sturt, F., Dupeyron, A., Huaman, O., Lane, K. J., and French, C.: Re-evaluating the resource potential of lomas fog oasis environments for Pre-ceramic hunter-gatherers under past ENSO modes on the south coast of Peru, *Quaternary Science Reviews*, 129, 196–215, <https://doi.org/10.1016/j.quascirev.2015.10.025>, 2015.
- 1270 Beresford-Jones, D. G., Mader, C., Lane, K. J., Cadwallader, L., Gräffingholt, B., Chauca, G., Grant, J., Hölzl, S., Coll, L. V. J., Lang, M., Isla, J., French, C., and Reindel, M.: Beyond Inca roads: archaeological mobilities from the high Andes to the



- Pacific in southern Peru, *Antiquity*, 97, 194–212, <https://doi.org/10.15184/aqy.2022.168>, 2023.
- 1275 Blume, H.-P., Brümmer, G. W., Horn, R., Kandeler, E., Kögel-Knabner, I., Kretschmar, R., Stahr, K., Thiele-Bruhn, S., Welp, G., and Wilke, B.-M.: Lehrbuch der Bodenkunde, 16. Auflage., edited by: Scheffer, F. and Schachtschabel, P., Spektrum Akademischer Verlag, Heidelberg, 2010.
- Bozarth, S.: Maize (*Zea mays*) Cob Phytoliths From A Central Kansas Great Bend Aspect Archaeological Site, *Plains Anthropologist*, 43, 279–286, <https://doi.org/10.1080/2052546.1998.11931908>, 1998.
- 1280 Brack Egg, A. and Mendiola Vargas, C.: *Ecología del Perú*, 3a ed., Lima : Bruño : PNUD, 2010. 3a ed., 2010.
- Branch, N. P., Kemp, R. A., Silva, B., Meddens, F. M., Williams, A., Kendall, A., and Pomacanchari, C. V.: Testing the sustainability and sensitivity to climatic change of terrace agricultural systems in the Peruvian Andes: a pilot study, *Journal of Archaeological Science*, 34, 1–9, <http://dx.doi.org/10.1016/j.jas.2006.03.011>, 2007.
- 1285 Bremond, L., Alexandre, A., Wooller, M. J., Hély, C., Williamson, D., Schäfer, P. A., Majule, A., and Guiot, J.: Phytolith indices as proxies of grass subfamilies on East African tropical mountains, *Global and Planetary Change*, 61, 209–224, <https://doi.org/10.1016/j.gloplacha.2007.08.016>, 2008.
- Brenner, E. M.: Sedimentologisch-bodenkundliche und mineralogische Untersuchungen im Llamoca-Moor (Peru, 14°S) – ein Beitrag zur Rekonstruktion der Paläoumweltbedingungen, Unpublished Master's thesis, Ruprecht-Karls-Universität Heidelberg, 122, 2011.
- 1290 Bronk Ramsey, C.: Bayesian Analysis of Radiocarbon Dates, *Radiocarbon*, 51, 337–360, <https://doi.org/10.1017/S0033822200033865>, 2009.
- Buckman, H. O. and Brady, N. C.: *Naturaleza y propiedades de los suelos: texto de edafología para enseñanza*, 1977.
- Cadwallader, L., Beresford-Jones, D. G., Whaley, O. Q., and O'Connell, T. C.: The Signs of Maize? A Reconsideration of What $\delta^{13}\text{C}$ Values Say about Palaeodiet in the Andean Region, *Human Ecology*, 40, 487–509, <https://doi.org/10.1007/s10745-012-9509-0>, 2012.
- 1295 Castillo, J., Barreda Ampuero, J., Vela Velásquez, C., and others: *Geología de los cuadrángulos de Laramate y Santa Ana. Hojas: 29-n, 29-ñ-[Boletín A 45]*, 1993.
- DIN 19684-2: *Bodenuntersuchungsverfahren im Landwirtschaftlichen Wasserbauchemische Laboruntersuchungen; Bestimmung des Humusgehaltes*, 1977.
- 1300 Eash, N. S. and Sandor, J. A.: Soil chronosequence and geomorphology in a semi-arid valley in the Andes of southern Peru, *Geoderma*, 65, 59–79, [https://doi.org/10.1016/0016-7061\(94\)00025-6](https://doi.org/10.1016/0016-7061(94)00025-6), 1995.
- Eitel, B.: *Bodengeographie, Das geographische Seminar*, 1999.
- Eitel, B. and Mächtle, B.: Man and Environment in the Eastern Atacama Desert (Southern Peru): Holocene Climate Changes and Their Impact on Pre-Columbian Cultures, in: *New Technologies for Archaeology: Multidisciplinary Investigations in Palpa and Nasca, Peru*, edited by: Reindel, M. and Wagner, G. A., Springer Berlin Heidelberg, Berlin, Heidelberg, 17–23, https://doi.org/10.1007/978-3-540-87438-6_2, 2009.
- 1305 Eitel, B., Hecht, S., Mächtle, B., Schukraft, G., Kadereit, A., and Wagner, G. A.: Geoarchaeological evidence from desert loess in the Nazca-Palpa region, southern Peru : Palaeoenvironmental changes and their impact on Pre-Columbian cultures, *Archaeometry*, 47, 137–185, 2005.
- 1310 Finucane, B., Agurto, P. M., and Isbell, W. H.: Human and animal diet at Conchopata, Peru: stable isotope evidence for maize agriculture and animal management practices during the Middle Horizon, *Journal of Archaeological Science*, 33, 1766–1776, <https://doi.org/10.1016/j.jas.2006.03.012>, 2006.
- Garreaud, R., Vuille, M., and Clement, A. C.: The climate of the Altiplano: observed current conditions and mechanisms of past changes, *Palaeogeography, palaeoclimatology, palaeoecology*, 194, 5–22, 2003.
- 1315 Gismondi, A., D'Agostino, A., Canuti, L., Di Marco, G., Basoli, F., and Canini, A.: Starch granules: a data collection of 40 food species, *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology*, 153, 273–279, <https://doi.org/10.1080/11263504.2018.1473523>, 2019.
- Glaser, B. and Birk, J. J.: State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Índio), *Geochimica et Cosmochimica Acta*, 82, 39–51, <https://doi.org/10.1016/j.gca.2010.11.029>, 2012.
- 1320 Goodman-Elgar, M.: Evaluating soil resilience in long-term cultivation: a study of pre-Columbian terraces from the Paca Valley, Peru, *Journal of Archaeological Science*, 35, 3072–3086, <https://doi.org/10.1016/j.jas.2008.06.003>, 2008.
- Handley, J., Branch, N., Meddens, F. M., Simmonds, M., and Iriarte, J.: Pre-Hispanic terrace agricultural practices and long-



- distance transfer of plant taxa in the southern-central Peruvian Andes revealed by phytolith and pollen analysis, *Vegetation History and Archaeobotany*, <https://doi.org/10.1007/s00334-023-00946-w>, 2023.
- 1325 Henderson, A., Galeano, G., and Bernal, R.: Field guide to the palms of the Americas, Princeton University Press, 2019.
- Hesse, R.: Fluvial dynamics and cultural landscape evolution in the Rio Grande de Nazca drainage basin, southern Peru, British Archaeological Reports Limited, 2008.
- 1330 Hogg, A. G., Heaton, T. J., Hua, Q., Palmer, J. G., Turney, C. S., Southon, J., Bayliss, A., Blackwell, P. G., Boswijk, G., Ramsey, C. B., Pearson, C., Petchey, F., Reimer, P., Reimer, R., and Wacker, L.: SHCal20 Southern Hemisphere Calibration, 0–55,000 Years cal BP, *Radiocarbon*, 62, 759–778, <https://doi.org/10.1017/RDC.2020.59>, 2020.
- Huisman, S. N., Raczka, M. F., and McMichael, C. N. H.: Palm Phytoliths of Mid-Elevation Andean Forests, *Frontiers in Ecology and Evolution*, 6, 2018.
- 1335 Hutschenreuther, A., Watzke, J., Schmidt, S., Büdel, T., and Henry, A. G.: Archaeological implications of the digestion of starches by soil bacteria: Interaction among starches leads to differential preservation, *Journal of Archaeological Science: Reports*, 15, 95–108, <https://doi.org/10.1016/j.jasrep.2017.07.006>, 2017.
- INGEMMET: Mapa Carta Geológica Nacional 1:50,000 Perú. Cuadrantes Laramate 29-n 1, 29-n 2, 29-n 3 & 29-n 4, Lima : INGENMET, 2001, Lima, Peru, 2001.
- 1340 Instituto de Estudios Histórico-Marítimos del Perú.: El Perú y sus recursos : atlas geográfico y económico., edited by: 2a ed., Auge, 1996.
- International Committee for Phytolith Taxonomy (ICPT): International Code for Phytolith Nomenclature (ICPN) 2.0, *Annals of Botany*, 124, 189–199, <https://doi.org/10.1093/aob/mcz064>, 2019.
- Isla, J. and Reindel, M.: La ocupación Wari en los valles de Palpa, costa sur del Perú, *Arqueología y Sociedad*, 193–226, 2014.
- 1345 IUSS Working Group, W.-F.: World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports N°. 106, FAO Rome, 2015.
- Jahn, R., Blume, H. P., Asio, V. B., Spaargaren, O., and Schad, P.: Guidelines for soil description, FAO, 2006.
- Kappl, B.: Bodenentwicklung einer pleistozänen Grundmoränensequenz in den südlichen Anden von Peru, 14°S, Unpublished Bachelor's thesis, Ruprecht-Karls-Universität Heidelberg, 44, 2012.
- 1350 Kemp, R., Branch, N., Silva, B., Meddens, F., Williams, A., Kendall, A., and Vivanco, C.: Pedosedimentary, cultural and environmental significance of paleosols within pre-hispanic agricultural terraces in the southern Peruvian Andes, *Quaternary International*, 158, 13–22, <http://dx.doi.org/10.1016/j.quaint.2006.05.013>, 2006.
- Kendall, A. and Rodríguez, A.: Desarrollo y perspectivas de los sistemas de andenería de los Andes centrales del Perú, Institut Français d'Études Andines, Lima, 312 pp., <https://doi.org/10.4000/books.ifea.6110>, 2009.
- 1355 Liu, L., Wang, J., Rosenberg, D., Zhao, H., Lengyel, G., and Nadel, D.: Fermented beverage and food storage in 13,000 y-old stone mortars at Raqefet Cave, Israel: Investigating Natufian ritual feasting, *Journal of Archaeological Science: Reports*, 21, 783–793, <https://doi.org/10.1016/j.jasrep.2018.08.008>, 2018.
- Logan, A. L., Hastorf, C. A., and Pearsall, D. M.: “Let's Drink Together”: Early Ceremonial Use of Maize in the Titicaca Basin, *Latin American Antiquity*, 23, 235–258, 2012.
- 1360 Mächtle, B.: Geomorphologisch-bodenkundliche Untersuchungen zur Rekonstruktion der holozänen Umweltgeschichte in der nördlichen Atacama im Raum Palpa, Südperu, Heidelberg, Univ., Diss., 2007, XX, 226 S. pp., 2007.
- Mächtle, B. and Eitel, B.: Fragile landscapes, fragile civilizations — How climate determined societies in the pre-Columbian south Peruvian Andes, *CATENA*, 103, 62–73, <https://doi.org/10.1016/j.catena.2012.01.012>, 2013.
- Mächtle, B., Eitel, B., Schukraft, G., and Ross, K.: Built on Sand: Climatic Oscillation and Water Harvesting During the Late Intermediate Period, in: *New Technologies for Archaeology: Multidisciplinary Investigations in Palpa and Nasca, Peru*, edited by: Reindel, M. and Wagner, G. A., Springer Berlin Heidelberg, Berlin, Heidelberg, 39–46, https://doi.org/10.1007/978-3-540-87438-6_3, 2009.
- 1365 Madella, M., Alexandre, A., and Ball, T.: International code for phytolith nomenclature 1.0., *Ann Bot*, 96, 253–260, <https://doi.org/10.1093/aob/mci172>, 2005.
- 1370 Mader, C.: Sea shells in the mountains and llamas on the coast: the economy of the Paracas Culture (800 to 200 BC) in Southern Peru, Harrassowitz Verlag, 2019.
- Mader, C., Hölzl, S., Heck, K., Reindel, M., and Isla, J.: The llama's share: Highland origins of camelids during the Late Paracas period (370 to 200 BCE) in south Peru demonstrated by strontium isotope analysis, *Journal of Archaeological*



- Science: Reports, 20, 257–270, <https://doi.org/10.1016/j.jasrep.2018.04.032>, 2018.
- 1375 Mader, C., Reindel, M., and Isla, J.: Camelids as cargo animals by the Paracas culture (800–200 BC) in the Palpa valleys of southern Peru, in: *Caravans in Socio-Cultural Perspective*, Routledge, 174–192, 2021.
- Mader, C., Reindel, M., and Isla, J.: Economic Directness in the Western Andes: A New Model of Socioeconomic Organization for the Paracas Culture in the First Millennium BC, *Latin American Antiquity*, 34, 385–403, <https://doi.org/10.1017/laq.2022.40>, 2023a.
- 1380 Mader, C., Reindel, M., Isla, J., Behl, M., Meister, J., and Hölzl, S.: In the land of the apu: Cerro Llamocca as a sacred mountain and central place in the pre-Columbian Andes of southern Peru, *Journal of Archaeological Science: Reports*, 49, 104045, <https://doi.org/10.1016/j.jasrep.2023.104045>, 2023b.
- Mader, C., Godde, P., Behl, M., Binder, C., Hägele, E., Isla, J., Leceta, F., Lyons, M., Marsh, E., Odenthal, R., Fernengel, E., Stryjski, P., Weber, A.-K., Reindel, M., and Meister, J.: An integrative approach to ancient agricultural terraces and forms of dependency: the case of Cutamalla in the prehispanic Andes, *Frontiers in Environmental Archaeology*, 3, <https://doi.org/10.3389/fearc.2024.1328315>, 2024.
- Marsh, E. J., Vranich, A., Blom, D., Bruno, M., Davis, K., Augustine, J., Couture, N. C., Ancapichún, S., Knudson, K. J., Popović, D., and Cunietti, G.: The center cannot hold: A Bayesian chronology for the collapse of Tiwanaku, *PLOS ONE*, 18, e0288798, <https://doi.org/10.1371/journal.pone.0288798>, 2023.
- 1390 Meister, J., Krause, J., Müller-Neuhof, B., Portillo, M., Reimann, T., and Schütt, B.: Desert agricultural systems at EBA Jawa (Jordan): Integrating archaeological and paleoenvironmental records, *Quaternary International*, 434, Part B, 33–50, <https://doi.org/10.1016/j.quaint.2015.12.086>, 2017.
- MINAM, M. del A. del P.: *Mapa nacional de cobertura vegetal: memoria descriptiva*, 2015.
- Munsell Color, L.: *Munsell soil color charts, Year 2000 rev. washable ed.*, Munsell Color, GretagMacbeth, New Windsor, NY, 2000.
- 1395 Nanavati, W. P., French, C., Lane, K., Oros, O. H., and Beresford-Jones, D.: Testing soil fertility of Prehispanic terraces at Viejo Sangayaico in the upper Ica catchment of south-central highland Peru, *Catena*, 142, 139–152, 2016.
- Olsen, S. R.: *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*, US Department of Agriculture, 1954.
- 1400 ONERN, Peru. O. N. de E. de R. N.: *Inventario, evaluación y uso racional de los recursos naturales de la costa : cuenca del Río Grande (Nazca)*, ONERN, 1971.
- Pearsall, D. M., Chandler-Ezell, K., and Chandler-Ezell, A.: Identifying maize in neotropical sediments and soils using cob phytoliths, *Journal of Archaeological Science*, 30, 611–627, [https://doi.org/10.1016/S0305-4403\(02\)00237-6](https://doi.org/10.1016/S0305-4403(02)00237-6), 2003.
- Piperno, D. R.: *Phytolith Analysis: An Archaeological and Geological Perspective*, Elsevier Science, 1988.
- 1405 Piperno, D. R.: *Phytoliths: A Comprehensive Guide for Archaeologists and Paleoecologists*, AltaMira Press, 2006.
- Pulgar Vidal, J.: *Geografía del Perú: Las Ocho Regiones Naturales, la Regionalización Transversal, la Sabiduría Ecológica Tradicional*, 10a ed., Peisa, Lima, Peru, 302 pp., 1996.
- Qader, W., Mir, S. H., Meister, J., Dar, R. A., Madella, M., and Rashid, I.: Sedimentological perspective on phytolith analysis in palaeoecological reconstruction, *Earth-Science Reviews*, 244, 104549, <https://doi.org/10.1016/j.earscirev.2023.104549>, 2023.
- 1410 van Reeuwijk, L. P.: *Procedures for soil analysis, SIXTH EDITION.*, ISRIC, Wageningen, 2002.
- Reimer, P. J., Austin, W., Bard, E., Bayliss, A., Blackwell, P. G., Ramsey, C. B., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP), *Radiocarbon*, 62, 725–757, <https://doi.org/10.1017/RDC.2020.41>, 2020.
- 1420 Reindel, M.: Life at the Edge of the Desert – Archaeological Reconstruction of the Settlement History in the Valleys of Palpa, Peru, in: *New Technologies for Archaeology: Multidisciplinary Investigations in Palpa and Nasca, Peru*, edited by: Reindel, M. and Wagner, G. A., Springer Berlin Heidelberg, Berlin, Heidelberg, 439–461, https://doi.org/10.1007/978-3-540-87438-6_25, 2009.
- Reindel, M. and Isla, J.: Cambio climático y patrones de asentamiento en la vertiente occidental de los Andes del sur del



- Perú, *Diálogo andino*, 83–99, 2013a.
- 1425 Reindel, M. and Isla, J.: Jauranga: una aproximación a la ocupación Paracas en los valles de Palpa, *Boletín de Arqueología PUCP*, 231–262, 2013b.
- Reindel, M. and Isla, J.: Nuevo patrón arquitectónico paracas en Lucanas, sierra sur del Perú, *Boletín de Arqueología PUCP*, 227–254, 2017.
- 1430 Reindel, M. and Isla, J.: De Paracas a Nasca: nuevas evidencias desde la vertiente occidental de la sierra de Lucanas, Ayacucho, *Boletín de Arqueología PUCP*, 229–254, 2018.
- Reindel, M., Isla, J., Gorbahn, H., and Otten, H.: Paracas en Palpa: los fundamentos del poder de la cultura Nasca, *Peruvian Archaeology*, 2, 37–64, 2015.
- Sandor, J. A. and Eash, N. S.: Ancient Agricultural Soils in the Andes of Southern Peru, *Soil Science Society of America Journal*, 59, 170–179, <https://doi.org/10.2136/sssaj1995.03615995005900010026x>, 1995.
- 1435 Schitteck, K., Forbriger, M., Mächtle, B., Schäbitz, F., Wennrich, V., Reindel, M., and Eitel, B.: Holocene environmental changes in the highlands of the southern Peruvian Andes (14° S) and their impact on pre-Columbian cultures, *Climate of the Past*, 11, 27–44, <https://doi.org/10.5194/cp-11-27-2015>, 2015.
- Schitteck, K., Forbriger, M., Berg, D., Hense, J., Schäbitz, F., and Eitel, B.: Last millennial environmental dynamics in the western Peruvian Andes inferred from the development of a cushion-plant peat hillock, *Perspectives in Plant Ecology, Evolution and Systematics*, 30, 115–124, <https://doi.org/10.1016/j.ppees.2017.09.002>, 2018.
- 1440 Schlichting, E., Blume, H.-P., and Stahr, K.: *Bodenkundliches Praktikum: eine Einführung in pedologisches Arbeiten für Ökologen, insbesondere Land- und Forstwirte und für Geowissenschaftler; mit 60 Tabellen*, 2., neubearb. Aufl., Blackwell-Wiss.-Verl., Berlin [u.a.], 1995.
- Sořna, V.: Climate and settlement in southern Peru: the northern Río Grande de Nasca drainage between 1500 BCE and 1532 CE, Dr Ludwig Reichert, 2015.
- 1445 Stevenson, F. J.: *HUMus Chemistry Genesis, Composition, Reactions*, 1982.
- Strömberg, C. A. E., Dunn, R. E., Crifò, C., and Harris, E. B.: Phytoliths in Paleocology: Analytical Considerations, Current Use, and Future Directions, in: *Methods in Paleocology: Reconstructing Cenozoic Terrestrial Environments and Ecological Communities*, edited by: Croft, D. A., Su, D. F., and Simpson, S. W., Springer International Publishing, Cham, 235–287, https://doi.org/10.1007/978-3-319-94265-0_12, 2018.
- 1450 Twiss, P. C.: Predicted world distribution of C3 and C4 grass phytoliths, in: *Phytolith Systematics: Emerging Issues*, edited by: Mulholland, S. C. and Rapp Jr., G., Springer US, 113–128, 1992.
- Unkel, I., Reindel, M., Gorbahn, H., Isla Cuadrado, J., Kromer, B., and Sossna, V.: A comprehensive numerical chronology for the pre-Columbian cultures of the Palpa valleys, south coast of Peru, *Journal of Archaeological Science*, 39, 2294–2303, <https://doi.org/10.1016/j.jas.2012.02.021>, 2012.
- 1455 USDA-NRCS: Soil Quality Indicators: Organic Matter, Soil quality information sheet, 1996a.
- USDA-NRCS: Soil quality resource concerns: Compaction, Soil quality information sheet, 1996b.
- USDA-NRCS: Soil Quality Indicators: Aggregate Stability, Soil quality information sheet, 1998a.
- USDA-NRCS: Soil Quality Indicators: Infiltration, Soil quality information sheet, 1998b.
- 1460 Varotto, M., Bonardi, L., and Tarolli, P. (Eds.): *World Terraced Landscapes: History, Environment, Quality of Life*, Springer International Publishing, Cham, <https://doi.org/10.1007/978-3-319-96815-5>, 2019.
- Vattuone, M. M. S., Roldán, J., Neder, L., Maldonado, M. G., and Vattuone, M. A.: Formative pre-Hispanic agricultural soils in northwest Argentina, *Quaternary Research*, 75, 36–44, 2011.
- Weberbauer, A.: *El mundo vegetal de los Andes peruanos: estudios fitogeográficos*, Lima: Ministerio de Agricultura, 1945., 1944.
- 1465 Zamora Jimeno, Carlos. and Bao Enríquez, Raúl.: *Regiones edáficas del Perú*, ONERN, 1972.
- Zavaleta García, Amaro.: *Edafología: el suelo en relación con la producción*, CONCYTEC, 223 pp., 1992.
- Zech, W., Schad, P., and Hintermaier-Erhard, G.: *Böden der Welt: Ein Bildatlas*, 2. Aufl. 2014., Springer Spektrum, Berlin, Heidelberg, 2014.
- 1470