

**Table 1. Database search results for significant flora known to occur within a 5 km radius of the subject site.**

Species	DBC Status	EPBC Act Status	Likelihood of Occurrence
<i>Banksia nivea subsp. uliginosa</i>	T	Endangered	Unlikely. Sandy clay, gravel.
<i>Banksia squarrosa subsp. argillacea</i>	T	Vulnerable	Unlikely. Prefers sandy or gravel soils and winter wet flats
<i>Brachyscias verecundus</i>	-	Critically Endangered	Unlikely. In a moss sward. On a granite outcrop.
<i>Caladenia busselliana</i>	-	Endangered	Unlikely. Sandy loam. Winter-wet swamps.
<i>Caladenia huegii</i>	-	Endangered	Unlikely. Grey or brown sand, clay loam.
<i>Caladenia procera</i>	-	Critically Endangered	Unlikely. Rich clay loam. Alluvial loamy flats
<i>Daviesia elongata</i>	T	Vulnerable	Unlikely. Prefers sandy soils
<i>Diuris micrantha</i>	T	Vulnerable	Unlikely. Brown loamy clay. Winter-wet swamps.
<i>Drakaea elastica</i>	T	Endangered	Unlikely. winter-wet swamps.
<i>Drakaea micrantha</i>	T	Vulnerable	Unlikely. Prefers white-grey sandy soils
<i>Eucalyptus x phylacis</i>	T	Endangered	Unlikely. Laterite, loam over granite. Coastal areas.
<i>Gastrolobium papilio</i>	T	Endangered	Unlikely. Sandy clay over ironstone and laterite. Flat plains.
<i>Grevillea brachystylis subsp. grandis</i>	T	Critically Endangered	Unlikely. Brown lateritic clay loam soils.
<i>Lambertia echinata subsp. occidentalis</i>	T	Endangered	Unlikely. Red clay or sand over laterite. Seasonally wet flats.
<i>Petrophile latericola</i>	T	Endangered	Unlikely. Red lateritic clay. Winter-wet flats.
<i>Verticordia densiflora var. pedunculata</i>	T	Endangered	Unlikely. Grey/yellow sand, sandy loam. Winter-wet low-lying areas.
<i>Verticordia plumosa var. ananeotes</i>	T	Endangered	Unlikely. Sandy loam. Seasonally inundated plains.
<i>Verticordia plumosa var. vassensis</i>	T	Endangered	Unlikely. White/grey sand. Winter-wet flats.

**Table 2. Significant fauna potentially occurring within the subject site as identified by State and Commonwealth database searches.**

Species	DBCA Status	EPBC Act Status	Likelihood of Occurrence
<i>Botaurus poiciloptilus</i> (Australasian Bittern)	T	Endangered	Unlikely, no potential habitat
<i>Calidris canutus</i> (Red Knot)	T	Endangered	Unlikely, no potential habitat
<i>Calidris ferruginea</i> (Curlew Sandpiper)	T	Critically Endangered	Unlikely, no potential habitat
<i>Calyptorhynchus banksii naso</i> (Forest Red Tailed Black Cockatoo)	T	Vulnerable	Possible, presence of foraging habitat
<i>Calyptorhynchus baudinii</i> (Baudin's Cockatoo)	T	Endangered	Possible, presence of foraging habitat
<i>Calyptorhynchus latirostris</i> (Carnaby's Cockatoo)	T	Endangered	Possible, presence of foraging habitat
<i>Dasyurus geoffroii</i> (Chuditch)	T	Vulnerable	Unlikely, no potential habitat
<i>Engaewa pseudoreducta</i> (Margaret River Burrowing Crayfish)	T	-	Unlikely, no potential habitat
<i>Engaewa reducta</i> (Dunsborough Burrowing Crayfish)	T	Critically Endangered	Unlikely, no potential habitat
<i>Falco hypoleucos</i> (Grey Falcon)	T	Vulnerable	Unlikely, no potential habitat
<i>Numenius madagascariensis</i> (Eastern Curlew)	T	Critically Endangered	Unlikely, no potential habitat
<i>Pseudocheirus occidentalis</i> (Western Ringtail Possum)	T	Critically Endangered	Unlikely, unsuitable habitat
<i>Sternula nereis nereis</i> (Australian Fairy Tern)	T	Vulnerable	Unlikely, no potential habitat
<i>Westralunio carteri</i> (Carter's Freshwater Mussel)	T	Vulnerable	Unlikely, no potential habitat

Migratory bird species have been omitted from this assessment as while they may infrequently visit the subject site, they are unlikely to rely on it for their survival given its degraded condition.