



Title	Biostratigraphy and Correlation of the Permian of Japan
Author(s)	Minato, Masao; Kato, Makoto; Nakamura, Koji; Hasegawa, Yoshiyuki; Choi, Dong Ryong; Tazawa, Jun-ichi
Citation	北海道大学理学部紀要, 18(1-2), 11-47
Issue Date	1978-02
Doc URL	http://hdl.handle.net/2115/34822
Type	bulletin (article)
File Information	18_1-2_p11-47.pdf



[Instructions for use](#)

BIOSTRATIGRAPHY AND CORRELATION OF THE PERMIAN OF JAPAN

by

Masao Minato, Makoto Kato, Koji Nakamura,
Yoshiyuki Hasegawa*, Dong Ryong Choi**
and Jun-ichi Tazawa***

(with 2 text-figures and 3 tables)

(Contribution from the Department of Geology and Mineralogy,
Faculty of Science, Hokkaido University, No. 1526)

Abstract

Permian biostratigraphy is described for seven selected districts in Japan. A new biostratigraphic scheme is presented for the Permian in the Kitakami mountains, based on a number of assemblage zones. New names, Maehama and Nabekoshi Stages are proposed in the Toyoma Series of the Kitakami mountains. An attempt is made to correlate the Japanese Permian in the studied districts.

Intruduction

International Subcommittee on Permian Stratigraphy (President- Prof. D.L. Stepanov) of the IUGS is planning to correlate eventually the World Permian. Dr. E.Ya. Leven of the USSR and Minato have been asked to prepare a correlation table of the Permian in the Tethys province. At the request of Dr. Leven Minato in cooperation with Kato made a draft of correlation chart of the Permian of Japan, and sent it to Dr. Leven sometime ago. Then the present authors jointly prepared the second, modified version of the first draft, with explanatory text. We sent this manuscript also to Dr. Leven in 1976.

In 1977 Dr. Leven circulated his view on a kind of Permian standard in the Tethyan realm, based mainly on his data from Central Asia, but combined with data from the other regions. He asked research workers on Permian for comments on his scheme to be sent to Drs. Leven, Minato and S.V. Meyen (Secretary of the Commission). Thus Minato received several letters from some research workers somehow criticizing Leven's scheme.

However, we feel that the first step we should take is to accumulate as many data as possible in connection with local biostratigraphy in many regions,

* Department of Geology, Faculty of General Education, Niigata University ** Kokusai Aeronautical Co., Ltd. *** Institute of Geology and Palaeontology, Faculty of Science, Tohoku University

before we come to any concrete idea of correlation. We also feel it useful and advisable to publish, if possible, such data as available sources of information on the concerned problem.

It is the aim of the present paper to fulfil such a need. However we are fully aware of that the paper is inevitably a sketch or an outline of the Japanese Permian, since Minato, to his regret, has not been able to really organize a working group for such a task in correlating the Permian deposits of entire Japan.

Marine Permian deposits are widely distributed throughout Japan. They are known in Hokkaido, Honshu, Shikoku, Kyushu and Okinawa.




Fig. 1 Surface distribution of Palaeozoic sediments and their metamorphosed counterparts in Japan, indicating the locations of important districts for Permian deposits.

For biostratigraphy and correlation, however, seven districts are here purposefully chosen for consideration; namely the Kitakami mountains in N.E. Japan, and the Omi, Akasaka, Atetsu, Akiyoshi, Oita and Kuma mountain districts in S.W. Japan (Fig. 1).

In Japan, the southern Kitakami mountains represent one of the best geologically investigated districts where Palaeozoic deposits are most typically distributed. And the Permian deposits are characterized by detrital as well as carbonate facies. Geology of this district is, however, much complicated owing to the repeated crustal movements. Therefore stratigraphic succession is very difficult to establish there. Also fossils are more or less deformed and often obliterated though they are locally abundant.

Minato presents here a new scheme of biostratigraphy for the Permian of the Kitakami mountains (Fig. 2).

In the other districts, the Permian is represented mostly by carbonate facies except for the Kuma mountain district where detrital facies is more predominant. Fossils are better preserved in these districts than those from the Kitakami mountains. But stratigraphic succession in these districts is by no means complete for the entire Permian, and only fusulinid foraminifers have been extensively searched for. Brief description is made on the Permian of each district.

Apart from the above mentioned two types of sedimentary environments, namely the carbonate facies and the mixed facies of carbonate and detritus, another type of deposits is actually distributed in central and the outer zone of Southwest Japan. They are dominantly volcanic in lithologic character at least with regard to the Lower to Middle Permian.

Thus three different types are recognizable amongst Permian deposits of Japan (Minato & Kato, in Minato, Gorai & Hunahashi (eds.), 1965). But as to the last mentioned type, biostratigraphy is by no means thoroughly investigated as yet, and it is not described here.

Detailed and thorough consideration and comprehension of the Japanese Permian are to be presented in the other occasion when we organize a working group on the problems of Permian correlation in Japan in near future.


As a matter of fact Minato has long studied the Palaeozoic deposits distributed in a rather narrow strip called the Setamai-Sakari district in the southern Kitakami mountains in cooperation with a number of his students. Without their effort of detailed mapping, we could hardly establish the lithologic sequence and many bio- and assemblage zones of Palaeozoic there.

The Permian deposits are especially well developed in the Setamai-Sakari district. Accordingly description will be based mainly on the data from this district concerning to the Kitakami mountains. Various information sources

may be found from the accompanied list of references.

Although not mentioned in the text, Dr. Y. Fujiwara made a palaeomagnetic study on the Lower Permian sediments, and Dr. N. Minoura has been studying the sedimentary rocks including limestones in the Kitakami mountains. Prof. M. Hunahashi kindly taught us on the lithologic nature of the volcanic rocks and feldspathic sandstones intercalated in the Permian. These studies are actually of great help to establish a sequence of the Permian

Fig. 2 Correlation chart of the Permian of Japan.
 A – slate; B – sandstone; C – limestone; D – conglomerate; E – alternation;
 G – unconformity; H – fault relation.



formations of the Kitakami mountains.

Then the Permian developed in Southwest Japan will be stated briefly, based on the researches hitherto carried out by a number of workers including ourselves.

We acknowledge Prof. F. Kahler of Klagenfurt, Austria, Dr. H. Kozur of German Democratic Republic, Dr. H. Taraz of Iranian Geological Survey, Prof. J.B. Waterhouse of the University of Queensland, Australia, for their kind comments on the major divisions on Permian, communicated to Minato.

Before stepping into description we would also like to acknowledge with many thanks Mr. S. Kumano for drawing figures, and Miss E. Mima for typing the present manuscript.

KUMA MOUNTAINS <i>Kanmera, 1958, 1963</i>	OITA PREFECTURE <i>Kanmera in Working Group, 1975</i>	AKIYOSHI PLATEAU <i>Hasegawa, 1963</i>	ATETSU PLATEAU <i>Nogami, 1963</i>	AKASAKA LIMESTONE <i>Honjo, 1959</i>	OMI LIMESTONE <i>Hasegawa et al, 1969</i>
	Fukami Formation 60m <i>Staffella Nankinella Palaeofusulina Reichelina Codonofusicella</i>				
Kuma Formation 900m+ <i>Codonofusicella sp.</i> <i>Lepidolina kumaensis</i> <i>Lepidolina multiseptata</i>		<i>Yabeina-Lepidolina Zone</i> 20m+	<i>Yabeina shiraiwensis</i> Z. 100m	<i>Yabeina Zone</i> 200m	<i>Yabeina-Lepidolina Zone</i>
		<i>Neoschwagerina Lepidolina Z.</i> 100m <i>Gifuella douvillei Verbeekina verbeeki Neoschwagerina haydeni Verbeekina heimi Neoschwagerina craticulifera</i>	<i>Neoschwagerina douvillei-margaritae</i> S.Z. 60m <i>Neoschwagerina craticulifera</i> S.Z. ?	<i>Gifuella douvillei</i> S.Z. 55mm <i>Yabeina otawai</i> S.Z. 90m <i>Neoschwagerina craticulifera</i> S.Z. 62m	<i>Neoschwagerina-Gifuella Zone</i>
		<i>Parafusulina Zone</i> 40m	<i>Parafusulina kaermitzensis</i> S.Z. 10m	<i>Pseudodolotina otawai</i> S.Z. 40m <i>Minoella nipponica</i> S.Z. 25m	<i>Parafusulina Zone</i> 50m
		<i>Pseudofusulina Zone</i> <i>Pseudofusulina krafftii magna</i> S.Z. 110m	<i>Pseudofusulina krafftii magna</i> S.Z. 40-60m		<i>Pseudofusulina krafftii</i> S.Z.
		<i>Pseudoschwagerina Z.</i> <i>Pseudofusulina vulgaris</i> S.Z. 150m <i>Pseudoschwagerina murauchensis</i> S.Z. 100m <i>Triticites simplex</i> S.Z. 100m	<i>Pseudofusulina vulgaris</i> Z. 80m <i>Pseudoschwagerina subspheerica</i> S.Z. 40m <i>Quasifusulina longissima ultima</i> - "P." nakazawai S.Z. 50-70m		<i>Pseudofusulina Z. 200m</i> <i>Pseudofusulina vulgaris</i> S.Z. <i>Pseudoschwagerina sp.</i> S.Z. <i>Triticites simplex</i> S.Z.
Yayamadake Lst <i>Pseudoschwagerina minatoi</i> Z. 60m <i>Pseudoschwagerina morkawai</i> Z. 60m					

Kitakami Mountains

A. The boundary between the Carboniferous and Permian formations

The Permian deposits unconformably lie on the Carboniferous formations with the basal conglomerate of the Sakamotozawa Series (Minato, 1942; Nagao and Minato, 1943; Yamada, 1959; Kanmera and Mikami, 1965; Saito, 1966, 1968). The basal conglomerate rests directly on the limestone formation (Nagaiwa formation) with *Profusulinella* fauna, entirely lacking in the formations represented by *Fusulinella* – *Fusulina* –, and *Triticites* fauna. In many places, the basal conglomerate is also observed directly resting on the formations correlatable with the lower Tournaisian.

B. The Sakamotozawa Series

The Permian system in the Kitakami mountains is divisible into three series: the Sakamotozawa, the Kanokura and the Toyoma Series in ascending order. Of them the Sakamotozawa Series is lithologically and palaeontologically divided into two units: Kawaguchi Stage and Kabayama Stage.

1. The Kawaguchi Stage

a. *Lophocarinophyllum suetomii* Zone, Pla

The lower half of the stage is composed of conglomerate at its base. Sandstone and slate are predominant upwards. At the type area it is less than 50 m in thickness, but is locally very thick, reaching almost 300 m in thickness. It is lithologically composed of thick feldspathic sandstone, acidic and basic tuffs, coaly slate, etc. besides thick slate and calcareous slate in alternation. Marine fossils are not rare in the last mentioned part of the stage but have not been studied in detail.

The lower half of the Kawaguchi stage may be designated as *Lophocarinophyllum suetomii* Zone, Pla, since this solitary coral has been found rather widely from the calcareous slate, at the middle and lower part of this formation. Fragmental plant remains are also found in the lower part.

Fossils described from this zone may be listed below. All of them indicate the Permian.

Cordaites principalis (Germer), *Wentzelella* (*Wentzelella*) sp. (Minato and Kato, 1965a), *Lophocarinophyllum suetomii* (Minato), *Waagenoconcha asiatica* Zavodowsky, *Derbyia* sp. (Nakamura, 1972), cfr. *Orthotetes callytharrensensis* (Thomas), *Magniderbyia* sp. (Nakamura, 1972), *Streptorhynchus* sp. (Naka-

mura, 1972), and small sized spiriferid resembling *Munella* (Minato MS).

b. *Zellia nunosei-Ferganites langsonensis* Zone, PIb

The upper half of the Kawaguchi stage is mainly composed of limestone, about 120 to 400 m in thickness. This is named as *Zellia nunosei-Ferganites langsonensis* Zone, PIb. This zone is also an assemblage zone; described fossils from this zone will be listed below.

Fusulinids:

Zellia nunosei Hanzawa, *Quasifusulina tenuissima* (Schellwien), *Paraschwagerina (Acervoschwagerina)* sp. (Kanmera and Mikami, 1965), *Paraschwagerina (Paraschwagerina)* sp. (Kanmera and Mikami, 1965), *Nipponitella explicata* Hanzawa, *Ferganites langsonensis* (Saurin), *Rugosofusulina alpina* (Schellwien), *Rugosofusulina rossica* (Schellwien), *Triticites* sp. A (Choi, 1973), *Triticites* sp. B (Choi, 1973), *Triticites* cf. *simplex* (Schellwien), *Nagatoella ikenoensis* Morikawa et Isomi, *Schwagerina* cf. *krotowi* (Schellwien et Dyhrenfurth), *Pseudofusulina ? jenkinsi* (Thorsteinsson), *Pseudofusulina* sp. A (Choi, 1973), *Pseudofusulina pseudoanderssoni* Shyomina, *Misellina claudiae* (Deprat), *Minojaponella elongata* Fujimoto et Kanuma, *Schubertella irumensis* Fujimoto, *Nagatoella minatoi* Kanmera et Mikami, *Paraschwagerina (Acervoschwagerina)* sp. (Choi, 1973), *Pseudoschwagerina schellwieni* Hanzawa, *Chalaroschwagerina vulgaris* (Schellwien et Dyhrenfurth), *Pseudofusulina* aff. *japonica* (Gümbel), *Pseudofusulina krafftii* (Schellwien et Dyhrenfurth), *Parafusulina motoyoshiensis* (Morikawa), *Toriyamaia laxiseptata* Kanmera.

Corals:

Pavastehphyllum (Sakamotosawanella) sakamotosawanum Minato et Kato, *Duplophyllum (Duplophyllum)* cf. *tenueseptatum* Schoupe et Stacul, *Iranophyllum (Laophyllum) nakamurai* Minato et Kato, *Durhamina kitakamiensis* Minato et Kato.

Calcareous Algae:

Anthracoportella magnipora Endo, *Epimastopra japonica* Endo, *Anchicodium magnum* Endo, *Hikorocodium elegantae* Endo, *Macroporella* sp. (Endo, 1951), *Osagia* sp. (Endo, 1951), *Hikorocodium elegantae* Endo, *Gyroporella* (?) *longipora* Endo, *Diplopora phanerospora* Pia, *Macroporella ?* sp. (Endo, 1952).

Ostracods:

Kirkbya cfr. *subnipponica* Ishizaki, *Kirkbya* sp., *Aurikirkbya* ? *brevis* Ishizaki, *Aurikirkbya*? *hinomataensis* Ishizaki, *Aurikirkbya* ? *lata* Ishizaki, *Aurikirkbya* ? *tenuis* Ishizaki, *Coronakirkbya hataii* Ishizaki, *Knightina hinomataensis* Ishizaki, *Amphissites kitakamiensis* Ishizaki, *Kelletina* ? *japonica* Ishizaki, *Kindlella kitanipponica* Ishizaki, *Hataiella ohazamensis* Ishizaki, *Hataiella longa* Ishizaki, *Hataiella minima* Ishizaki, *Roundyella dorsopapillosa* Sohn, *Bairdia iwaizakiensis* Ishizaki, *Bairdia* cfr. *dissimilis* Cooper, *Bairdia* sp. (Ishizaki, 1967).

Gastropod:

Phymatifer pugiloides Hayasaka

Amongst fossils above listed, *Pseudoschwagerina schellwieni* Hanzawa appears slightly earlier than the rest of fossils, *Chalartoschwagerina vulgaris* for instance.

Durhamina kitakamiensis Minato et Kato also is found from the relatively lower part of this zone. *Kindlella kitakamiensis* Ishizaki is said to be very resembled *Ulrichia minuta* described by Harris and Lalicker (1932) from the Wolfcampian, Kansas (Ishizaki, 1967). All the ostracods listed above are actually described by Ishizaki (1967) not from the Setamai-Sakari district but from the northern area, the Tassobe, as *Wentzelella* sp. listed from Pla in the foregoing paragraph.

2. The Kabayama Stage

The Kabayama stage is lithologically divided into two units: the lower half is chiefly composed of limestone, while the upper half is mainly consisting in slate and sandstone in alternation; although the upper half may be locally replaced by carbonate rocks.

The base of the Kabayama stage may be defined as the thin layer of conglomerate in the limestones as diagrammatically shown in the correlation chart (Fig. 2). At the type area of the Sakamotozawa series, presence of unconformity is proved below the limestone conglomerate (Mikami, 1965) although in other areas, stratigraphically equivalent conglomerate is observed to be intraformational.

The limestone beds of the Kabayama stage may be divided into two assemblage zones (PIc and PId), and the uppermost part of this stage may be distinguished from both as PIe.

a. *Schwagerina* cf. *compacta* Zone, Plc

Fusulinids:

Pseudofusulina aff. *pseudosimplex* (Chen), *Schwagerina* cf. *compacta* (White), *Rugosofusulina* aff. *serrate* Rauser-Chernousova, *Schubertella* sp. (Choi, 1973), *Toriyamaia laxiseptata* Kanmera, *Nankinella* sp. (Choi, 1973), *Staffella* sp. (Choi, 1973), *Nagatoella minatoi* Kanmera et Mikami, *Paraschwagerina* (*Acervoschwagerina*) sp. (Choi, 1973), *Chalaroschwagerina vulgaris* (Schellwien et Dyhrenfurth), *Pseudofusulina krafftii* (Schellwien et Dyhrenfurth), *Parafusulina* aff. *japonica* (Gümbel), *Parafusulina motoyoshiensis* (Morikawa), *Misellina claudiae* (Deprat).

Corals:

Waagenophyllum (*Waagenophyllum*) *polyseptata* Minato, *Yatsengia kabayamensis* Minato, *Michelinia* (*Protomichelinia*) *multitabulata* (Yabe et Hayasaka), *Wentzelophyllum hayasakai* Minato et Kato, *Wentzelophyllum kitakamiense* Minato et Rowett, *Polythecalis* sp.

Calcareous Algae:

Teutloporella cfr. *triasina* Schaueroth, *Mizzia velebitana* Schubert, *Atractyliopsis* ? sp. (Endo, 1951), *Macroporella* sp. (Endo, 1951), *Gyroporella longipora* Endo, *Anthracoporella magnipora* Endo, *Macroporella maxima* Endo, *Atractyliopsis* (?) sp. (Endo, 1952).

Gastropod:

Phymatifer aff. *nodocarinatus* (Wanner)

In the southwestern part of the Kitakami mountains (Maiya area), there is a plant bearing formation, composed of slate with carbonaceous matter, which has been called the Rodai Formation (Mabuti, 1932; Onuki et al, 1960; Ueda, 1963). This formation rests on the limestones with *Pseudoschwagerina schellwieni*, *Zellia nunosei* and *Triticites* cf. *simplex*. Hence, the stratigraphical level of the plant bearing formation may be correlatable with Plc.

The following species is recorded from the Rodai formation (Asama, 1956):

Sphenophyllum cf. *oblongifolium* (Germ. et Kaulf.), *Sphenophyllum Thonii* var. *minor* Sterzel, *Pecopteris* sp., *Odontopteris subcrenulata* Zeiller, *Gigantopteris Whitei* Halle, *Taeniopteris latecostata* Halle, *Taeniopteris* cfr. *Schenckii* Sterzel, *Taeniopteris Nystroemii* Halle, *Cordaites principalis* Germ., *Cordaites* (*Noeggerathiopsis*) *Arakawae* Asama, *Zamiopteris glossopteriodes* (Schmalh.) f. *minor* Kawasaki, *Zamiopteris uedae* Asama, *Psymphyllum flabellatum*

(Lindley et Hutton).

Presence of *Gigantopteris Whitei* is especially worthy of note, since this species is one of characteristic elements of the Shansi formation of Taiyuan, Shansi province, North China (Halle, 1927).

The Rodai formation is overlain by the formation characterized by frequent occurrence of *Monodioxodina matsubaishi* (Fujimoto) probably correspondent to the Lower part of the Kanokura series to be later on described.

b. *Pseudofusulina fusiformis*-*Pseudofusulina ambigua* Zone, PId

Although the lithological boundary between the preceding assemblage zone and the present zone (PId) is hardly discernible in the field, certain types of fusulinids such as *Pseudofusulina fusiformis*, *Pseudofusulina ambigua*, *Parafusulina* aff. *gigantea*, etc. newly appear in this zone. Especially *Pseudofusulina ambigua* occupies the highest level of this zone.

It may be also noted that *Pseudoschwagerina schellwieni* and *Chalartoschwagerina vulgaris* are still found in the lower part of this zone.

Fusulinids:

Monodioxodina kattaensis (Schwager), *Paraschwagerina* (*Paraschwagerina*) sp. (Choi, 1973), *Paraschwagerina* (*Acervoschwagerina*) cf. *endoi* Hanzawa, *Chalartoschwagerina setamaiensis* Choi, *Pseudofusulina tchernyschewi* (Schellwien), *Pseudofusulina fusiformis* (Schellwien et Dyhrenfurth), *Pseudofusulina ambigua* (Deprat), *Parafusulina* aff. *gigantea* (Deprat), *Parafusulina* cfr. *multi-septata* (Schellwien), *Nagatoella minatoi* Kanmera et Mikami, *Pseudoschwagerina schellwieni* Hanzawa, *Chalartoschwagerina vulgaris* (Schellwien et Dyhrenfurth), *Pseudofusulina* aff. *japonica* (Gümbel), *Pseudofusulina krafftii* (Schellwien et Dyhrenfurth), *Minojaponella elongata* Fujimoto et Kanuma, *Parafusulina motoyoshiensis* (Morikawa), *Chusenella pseudocrassa* Kanmera, *Nankinella* sp. C (Choi, 1973), *Eoverbeekina* sp. (Choi, 1973).

Corals:

Michelinia (*Protomichelinia*) *multitabulata* Yabe et Hayasaka, *Yatsengia kabayamensis* Minato, *Yatsengia ibukiensis* Minato, *Waagenophyllum* (*Waagenophyllum*) *polyseptata* Minato.

Calcareous Algae:

Mizzia velebitana Schubert, *Teutloporella* cf. *triasina* Schaubroth.

c. *Mesolobus mesolobus* Zone, Pie

The overlying formation on the limestones represented by *Pseudofusulina fusiformis-ambigua* zone is chiefly composed of slate, and alternation of sandstone and slate, 200 m in total thickness. It is locally intercalated by seams of magnetite sandstone especially in the eastern part of the Setamai-Sakari district. Fossils are rare. The described species from this formation are as follows:

Avonia echidniformis Grabau et Chao, *Juresania juresanensis* (Tschernyschew), *Mesolobus mesolobus* (Norwood et Pratten), *Orthotetina* sp. (Nakamura, 1972), *Meekella mexicana* Girty, *Phricodothyris* sp. (Nakamura, 1959), *Schuchertella chaoi* Nakamura, *Derbyia* sp. (Nakamura, 1972), "*Productus*" *yohi* Chao, *Dictyoclostus* aff. *gruenewaldti* (Krotow), *Dictyoclostus taiyuanfuensis* var. *loczyi* Nakamura, *Dictyoclostus* cf. *inflatiformis* var. *expansus* (Grabau), *Dictyoclostus grabaui* Nakamura, *Dictyoclostus* sp. A (Nakamura, 1960), *Tainoceras kitakamiense* Hayasaka, *Bellerophon* (*Bellerophon*) *kitakamiensis* Murata, "*Murchisonia*" sp. (Murata, 1971), *Towapteria nipponica* Nakazawa et Newell, *Waagenoperna* (*Permoperna*) *hayamii* Nakazawa et Newell, *Aviculopecten* sp. (Murata, 1971), *Pseudoactinodontophora yabei* Murata.

C. The Kanokura Series

The Kanokura Series rests on the Sakamotozawa Series with basal conglomerate. The conglomerate seems to directly cover the limestones of either Pic or PId and sandstones characterized by *Mesolobus* fauna. Presence of unconformity between the Sakamotozawa and the Kanokura Series is almost doubtless, although the unconformity has not been evidently proved yet in the field.

The lower half of the Kanokura Series has been distinguished as the Kattisawa Stage from the upper half, the Iwaizaki Stage. It may be worth while mentioned that a few fusulinid species show a long geological range extending from the Kabayama Stage until the Kattisawa Stage: e.g. *Parafusulina motoyoshiensis*, *Chusenella pseudocrassa* and *Monodiexodina matsubaishi*.

1. The Kattisawa Stage

The stage is mainly composed of sandstone and alternation of sandstone and slate, besides conglomerate at its basal part. There are at least two thin layers of limestone in both upper and lower part of this formation. Fusulinids found from the upper limestone are mostly ranging upwards until the lower

part of the Iwaizaki Stage, while the fusulinids of the lower limestone seem to show somewhat older aspect.

Thus the Kattisawa Stage may be divisible into two assemblage zones.

a. *Monodiexodina kofuganensis* Zone, PIIa

Rauserella alveolaris Choi, *Yangchienia kwangsiensis* Chen, *Chusenella choshiensis* Chisaka, *Monodiexodina kofuganensis* Choi, *Cancellia* sp. (Choi, 1973), *Parafusulina iwaizakiensis* (Morikawa), *Monodiexodina matsubaishi* Fujimoto, *Codonofusiella explicata* Kawano, *Chusenella pseudocrassa* Kanmera.

b. *Leptodus nobilis* Zone, PIIb

From the carbonate rocks stratigraphically positioned in the higher level, the following fusulinids, corals, calcareous algae and bryozoa have been described.

Fusulinids:

Rauserella alveolaris Choi, *Monodiexodina matsubaishi* Fujimoto, *Pseudodoliolina gravitesta* Kanmera, *Kahlerina pachythea* K. Devide et Ramovs, "*Colania*" *kozuboensis* Choi, *Parafusulina* aff. *mccloudensis* Skinner et Wilde, *Chusenella* sp. (Choi, 1973), *Pseudodoliolina elongata* Choi, *Codonofusiella explicata* Kawano, *Verbeekina verbeeki* Geinitz.

Corals:

Waagenophyllum (Waagenophyllum) indicum (Waagen et Wentzel), *Parawentzelella (Parawentzelella) iwaizakiensis* (Yabe et Minato), *Parawentzelella (Miyagiella) miyagiensis* Minato et Kato, *Parawentzelella (Miyagiella) motoyoshiensis* Minato et Kato, *Iranophyllum (Iranophyllum) tunicatum* Igo.

Calcareous Algae:

Mizzia velebitana Schubert.

Brachiopods, bryozoa and pelecypods are not seldom found from the lower part of the Kattisawa Stage, but they are more predominant in the upper part. Fossils mainly described from the upper part may be listed below.

Bryozoas:

Fistulipora kesenumensis Sakagami, *F. megastoma* Sakagami, *Meekopora*

delicata Sakagami, *Batostomella* sp. (Sakagami, 1961), *Leioclema globosa* Crockford, *Fenestella* cf. *retiformis* (Scholtheim), *Polypora fujimotoi* Sakagami, *P. hataii* Sakagami, *P. endoi* Sakagami, *P. sugiyamae* Sakagami, *Tabulipora* ? sp. (Sakagami, 1961), *Septopora kawamatae* Sakagami, *Thamniscus* cf. *dubius* (Scholtheim), *Penniretepora kamiyatsenensis* Sakagami, *Hayasakapora* ? sp. (Sakagami, 1961), *Sulcoretepora nipponica* Sakagami.

From Iwaizaki, south of Kesennuma City, Sakagami (1961) recorded the following bryozoans.

Fistulipora cf. *timorensis* Bassler, *Fistulipora* sp., *Meekopora delicata* Sakagami, *Batostomella sponigera* Bassler, *Fenestella rhomboidea* Nikiforova, *Hayasakapora matsudae* Sakagami, *Saffordotaxis morikawae* Sakagami, *Strebla scopora delicatula* Sakagami, *Sulcoretepora nipponica* Sakagami.

Brachiopoda:

Isogramma paotchowensis (Grabau et Chao), *Isogramma heritschi* Nakamura, *Permudaria asiatica* Nakamura, Kato et Choi, *P. tenuistriata* Tazawa, *Waagenoconcha humboldti* (d'Orbigny), *W. imperfecta* Prendergast, *Spinomarginifera kueichowensis* Huang, *S. huangi* Nakamura, *Dictyoclostus* aff. *spiralis* (Waagen), *D. sino-indicus* (Frech), *Orthotetina hayasakai* Nakamura, *O. elongata* Nakamura, *O. kayseri* (Jäkel), *O. polita* Fliegel, *O. transversa* Nakamura, *O. sp. A* (Nakamura, 1972), *Meekella* cf. *striatocostata* (Cox), *M. cf. eximia* (Eichwald), *M. garnieri* Bayan, *M. nodosa* Nakamura, *M. minato* Nakamura, *M. sp.* (Nakamura, 1972), *Schuchertella frechi* Huang, *S. cf. beyrichi* (Rothpletz), *Streptorhynchus pelargonatus* Schlotheim, *Derbyia nipponica* Nakamura, *D. cf. acutiplicata* Metz, *D. sp. A* (Nakamura, 1972), *Leptodus richthofeni* Kayser em. Hayasaka, "*Productus*" *flemingi* De Koninck, "*Productus*" *horridus* Sowerby, "*Productus*" *vishnu* Waagen, *Dielasma* cf. *biplex* Waagen, "*Spiriferina*" *crinata* Schlotheim, *Leptodus nobilis* Waagen, *Chonetes blanfordi* Reed, *C. sinuosa* Schellwien, *Linoproductus cora* (d'Orbigny), *Rhynchonella (Uncinulus) jabiensis* Waagen, *Spiriferella* cf. *saranae* mut. *lita* (Fredericks), *Orbiculoidea* sp. (Hayasaka, 1963), *Punctospirifer mesocostatus* Hayasaka, *Martiniopsis cathaysiensis* Grabau, *Geyerella arakii* Hayasaka, *G. koizumii* Hayasaka, *Camarophoria humbletonensis* Howse, *Canocrinella villiersi kozlowskiana* Fredericks, *C. cancriniformis spinosa* Hayasaka et Minato, *Martinia semiplana* Waagen, *M. semiplana* var. *lata* Grabau, ? "*Spirifer*" *lyra* Kutorga, *Neospirifer fasciger* (Keyserling), *Orthotetina kitakamiensis* (Hayasaka), *Orthotetes rugosa* Fredericks, *Kiangsiella* cf. *condoni* (Thomas), *Productella patula* Girty, *Martinia rostrata* Girty, *Rhipidomella* sp., *Derbyia hemisphaerica* Waagen, *Waagenites deplanata* (Waagen), *W. barusiensis*

(Davidson), "*Productus*" *gratiosus* Waagen, "*P.*" *carinatus* Chi-Thuan, *Anidanthus* sp. (Tazawa, 1976), *Echinoconchus fasciatus* Kutorga, *E. taimyrensis* Einor, *Cançrinella cançriniformis* (Tschernyschew), *Nantanella* sp. (Tazawa, 1976), *Spiriferella keilhavii* (von Buch), *Callispirina ornata* (Waagen).

Pelecypoda:

Parallelodon cf. *longus* Maslennikov, *P.* cf. *tenuistriatus* (Meek et Worthen), *Aviculopinna rectangularis* (Hayasaka), *Aviculopecten hataii* Murata, *A.* cf. *hataii* Murata, *A.* sp. A (Nakazawa et Newell, 1968), *Aviculopecten (Deltopecten) hiemalis* Salter, *Hayasakapecten sasakii* (Murata), *Etheripecten ? hayasakai* (Murata), *Acanthopecten spinosus* Hayasaka, *A. onukii* Murata, *A.* cf. *coloradoensis* (Newberry), *Streblopteria* sp. A (Nakazawa and Newell, 1968), *S.* ? sp. B (Nakazawa and Newell, 1968), *Cyrtorostra* cf. *lunwalensis* (Reed), *Euchondria ? kesennensis* (Hayasaka), *Annuliconcha kitakamiensis* Murata, *Schizodus tobai* (Hayasaka), *Astartella ?* sp. (Nakazawa and Newell, 1968), *Sanguinolites* sp. (Nakazawa and Newell, 1968), *S. kamiyassensis* Nakazawa et Newell, *Wilkingia* sp. (Nakazawa and Newell, 1968), *Waagenoperna (Permoperna) hayamii* Nakazawa et Newell, *Edmondia* sp. (Hayasaka, 1967), *Edmondia* sp. (Nakazawa and Newell, 1968), *Promytilus maiyaensis* Nakazawa et Newell, *Leptodesma (Leiopteria)* sp., *Towapteria nipponica* Nakazawa et Newell, *Enisipteria onukii* Nakazawa et Newell, *Hayasakapecten shimizui* Nakazawa et Newell, *Pernopecten* sp. (Nakazawa and Newell, 1968), *Lopha ? teranosawensis* Nakazawa et Newell, *Neoschizodus (Neoschizodus) kitakamiensis* Nakazawa et Newell, *Netshajewia* sp. (Nakazawa and Newell, 1968), *Stutchburia ?* sp. (Nakazawa and Newell, 1968), *Myoconcha* sp. (Hayasaka, 1967), *Allorisma* sp. (Hayasaka, 1967).

Cephalopoda:

Cibolites cf. *uddeni* Plummer et Scott, *Tainoceras abukumense* Hayasaka, *Tainoceras kitakamiensis* Hayasaka, *Foordiceras akiyamai* Hayasaka, *Paracel-tites* aff. *elegans* Girty, *Tyronautilus permicus* Hayasaka, *Foordiceras whynei-formis* Hayasaka et Ozaki, *Agathiceras* aff. *suessi* Gemm., *Propinacoceras* aff. *galilaei* Gemm., *Stacheoceras iwaizakiense* Mabuti, *Hanieloceras intermedium* (Wanner), *Medlicottia* sp. (Hayasaka, 1954).

Conulariida:

Neoconularia rectangularis (Hayasaka), *Neoconularia* sp. (Hayasaka, 1963), *Calloconularia ? kitakamiensis* Murata.

Trilobita:

Pseudophillipsia obtusicauda (Kayser).

Monodioxodina matsubaishi is especially dominant throughout the Kattisawa Stage; *Rauserella alveolaris* also ranges from the lower to the upper division of the stage. Presence of *Cancellina* in the *Monodioxodina kofuganensis* zone is remarkable. While, *Colania kozuboensis* Choi found from the upper part of the Kattisawa Stage very resembles "*Neoschwagerina*" *douvillei* Ozawa, although *Colania kozuboensis* shows to range further upwards until the lower horizon of the Iwaizaki Stage.

2. The Iwaizaki Stage

The lower part of the Iwaizaki Stage is lithologically changeable from place to place; usually limestone or limestone and slate in alternation. But it is also represented by thick conglomerate with limestone lenses. Therefore, the lowest level of the Iwaizaki Stage has been defined at the first appearance of the *Lepidolina* fauna.

Lepidolina multiseptata-*L. minatoi* Zone, PIIc

Described fossils from this zone may be listed below:

Fusulinids:

Lepidolina multiseptata (Deprat), *Nankinella* sp. A (Choi, 1973), *Pseudodoliolina gravitesta* Kanmera, *Kahlerina pachythea* K. Devide et Ramovs, *Lepidolina minatoi* Choi, *Lepidolina kumaensis* Kanmera, "*Colania*" *kozuboensis* Choi, *Schwagerina acris* Thompson et Wheeler, *Parafusulina* aff. *mccloudensis* Skinner et Wilde, *Chusenella* sp. (Choi, 1973), *Pseudodoliolina elongata* Choi, *Pseudodoliolina kanokuraensis* Choi, *Dunbarula kitakamiensis* Choi, *Codonofusiella inuboensis* (Chisaka), *Pseudofusulina chihsiensis* (Lee), *Rauserella pachythea* Choi, *Verbeekina verbeeki* (Geinitz).

Brachiopoda:

Aulosteges sp. (Hayasaka, 1964), *Spinomarginifera huangi* Nakamura, *Gratiosina gratiosus* (Waagen), *Waagenoconcha humboldti* (d'Orbigny), *Waagenoconcha purdoni* (Davidson), *Martinia semiplana* Waagen, *Scacchinella* cf. *gigantea* Schellwien, *Richthofenia* sp., *Leptodus richthofeni* (Kayser), *Spirifer* cf. *saranae* Vern. mut. *lita* Fredericks, *Productus* sp. (Mabuti, 1935), *Spiriferina cristata* Schlothheim, *Spirifer* sp. (Mabuti, 1935), *Camarophoria* sp.,

Meekella gigantea Hayasaka, *M. sp.* (Koizumi, 1964), *Orthotetina sp.* (Koizumi, 1964), *Geyerella sp.* (Koizumi, 1964).

Ammonoids:

Propinacoceras aff. galilaei Gemmellaro, *Tainoceras abukumaensis* Hayasaka, *Foordiceras akiyamae* Hayasaka.

Corals:

Waagenophyllum (Waagenophyllum) indicum (Waagen et Wentzel), *Wentzelloides maiyaensis* Yabe et Minato, *Parawentzelella (Parawentzelella) canalicifera sisophonensis* Fontaine, *Wentzelella (Szechuanophyllum) kitakamiensis* Yabe et Minato, *Yatsengia kiangsuensis* var. *mabutii* Minato.

Calcareous Algae:

Mizzia velebitana Schubert

Ostracods:

Holinella elliptica Ishizaki, *Aurikirkbya formula* Ishizaki, *A. subkelllettae* Ishizaki, *Kirkbya atolla* Ishizaki, *K. centrotumida* Ishizaki, *K. magniforma* Ishizaki, *K. multicresta* Ishizaki, *K. subnipponica* Ishizaki, *K. subquadriforma* Ishizaki, *Amphissites centronotus* (Ulrich et Bassler), *Ectodemites globosa* Ishizaki, *Polytylites kitanipponica* Ishizaki, *Roundyella neopapillosa* Ishizaki, *Bairdia eucurvia* Ishizaki, *B. nagaiwensis* Ishizaki, *B. cfr. oklahomaensis* Harlton, *B. trianguliformis* Chen, *Bairdiacypris deloi* Bradfield, *Ceratobairdia ? ambigua* Ishizaki, *Spinobairdia sp.*, *Cavellina ? nipponica* Ishizaki, *Tube-libairdia venusta* (Chen).

Bryozoa:

Fistulipora iwaizakiensis Sakagami, *F. cf. timorensis* Bassler, *F. sp.* (Sakagami, 1961), *Meekopora densa* Sakagami, *Ramipora ambigua* Sakagami, *Goniocladia intricata* Sakagami, *Coscinotrypa ? sp.* (Sakagami, 1961), *Batostomella igoi* Sakagami, *B. microstoma* Sakagami, *Dyscritella cylindrica* (Crockford) ?, *Coeloclemis minima* Sakagami, *Fenestella spp.* (Sakagami, 1961), *Polypora spp.* (Sakagami, 1961), *Penniretepora iwaizakiensis* Sakagami, *P. tenuis* Sakagami, *P. akiyamae* Sakagami, *Hayasakapora erectoradiata* Sakagami, *Streblascopora delicatula* Sakagami, *Clansotrypa exillis* Sakagami.

Many fusulinid species characterizing the *Leptodus nobilis* Zone disappear in the earlier half of the present zone. Rugose corals belonging to Waagenophyllidae seem to have been almost extinct in this zone.

Lepidolina minatoi appears to be more dominant in the upper part. *Schwagerina acris* occurs at the highest level with *Lepidolina multiseptata* and *Condonofusiella explicata*.

D. The Toyoma Series

The thick Toyoma Series is lithologically divisible into two major units: the Maehama Stage and Nabekoshi Stage in ascending order. The former of them is mainly composed of slate, with conglomerate at its base, while the latter is chiefly composed of sandstone with intercalations of thin limestone and conglomerate lense.

Characteristic and described fossils from this series may be listed below.

1. The Maehama Stage

Araxoceras cf. *kiangsiense* Zone, PIIIa

Astartella toyomensis Nakazawa et Newell, *Palaeoneilo ogachiensis* Hayasaka, *Bellerophon* (*Bellerophon*) *yabei* Murata, *Euphemitopsis kitakamiensis* Murata, *Paraconularia siitai* (Sugiyama), *Araxoceras* cf. *kiangsiense* Chao, *Protocycloceras* cfr. *cyclophorum* (Waagen), *Stacheoceras* cf. *trimuvii* Diener, *Eumedlicottia* sp. (Bando, 1975), *Artinskia* sp. (Bando, 1975).

Among them, *Araxoceras* cf. *kiangsiense* occupies rather higher level than the other species.

2. The Nabekoshi Stage

Colaniella parva Zone, PIIIb

Colaniella parva (Colani), *Paracolaniella leei* Wang, *Lantschichites* sp. (Tazawa, 1975), *Orthothrix* cf. *excavata* (Geinitz), *Tschernyschewia typica* Stoyanow, *Megousia nakamurai* Tazawa, *Megousia auriculata* Muir-Wood et Cooper, *Paramarginifera japonica* Tazawa, *Eolyttonia* cf. *nakazawai* Shimizu, *Palaeofusulina* sp.

From the Toyoma Series the following land plants were recorded by Endo (1950) and Konno (1973):

Neocalamites superpermicus Konno, *Paracalamites iwatensis* Konno, *P. takahashii* (Endo).

Exact horizon for them is, however, difficult to mention at present.

Kuma Mountain District

The Kuma mountain district in central Kyushu is a terrain where Permian clastic sediments are predominantly developed, which is intercalated by large and small limestone lenses at various horizons. Biostratigraphy in the Kuma mountain region was worked out in detail by Kanmera 1954, 1958, 1963. But Permian sediments are so much folded and faulted that a continuous stratigraphic succession is difficult to establish there.

The Yayamadake limestone furnishes a good section for *Pseudoschwagerina* limestone, which is said to unconformably cover the *Triticites* limestone (Kanmera, 1952).

According to Kanmera (1958) the upper part of the Yayamadake limestone is defined as *Pseudoschwagerina* Zone which is further divisible into two fossil Zones.

The lower fossil Zone, *Pseudoschwagerina morikawai* Zone, yields *Pseudoschwagerina morikawai* Igo, *Triticites montiparus* (Möller), *T. ozawai* Toriyama, *T. aff. haydeni* (Ozawa), *T. sp.* (Kanmera, 1958), *T. yayamadakensis evectus* Kanmera, *Rugosofusulina prisca* [Ehrenberg (Möller)], *Schwagerina cf. alpina* (Schellwien), *Quasifusulina longissima ultima* Kanmera.

The Upper fossil Zone is called *Pseudoschwagerina minatoi* Zone which yields *Pseudoschwagerina minatoi* Kanmera, *Paraschwagerina shimodakensis* Kanmera, *Triticites samaricus* Rauser-Chernoussova, *T. fornicus* Kanmera. *T. aff. pusillum* (Schellwien), *T. sp.* (Kanmera, 1958), *Rugosofusulina pristina* Kanmera, *Pseudofusulina regularis* (Schellwien), *Pseudofusulina sp.* (Kanmera, 1958), *Psf. sokensis* Rauser-Chernoussova, *Psf. horrida* Kanmera, *Psf. kumasoana* Kanmera, *Schwagerina stabilis* (Rauser-Chernoussova), *Schw. krotowi* (Schellwien), *Schw. grandensis* Thompson, *Nankinella kotakiensis* (Fujimoto et Kawada), *N. kawadai* (Igo).

The upper part of the Yayamadake limestone is superceded by pyroclastic deposits which produce no fossils.

To the south to the area of the Yayamadake limestone there develops the Kozaki formation consisting of slate and conglomerate in alternation with some limestone lenses which yield fossils. The formation is bounded by faults and is lithologically divided into the lower and upper subformations. The lower part is further faunistically contains two fossil horizons (Kanmera, 1961, 1963).

From the lower part the following fusulinids were described:

Misellina claudiae (Deprat), *Nankinella kozakiensis* Kanmera, *Sphaerulina crassispira japonica* Kanmera, *Minojapanella sp.*, *Toriyamaia laxiseptata* Kanmera, *Nagatoella sp.*, *Parafusulina (Skinnerella) gruperaensis* Thompson et Miller, *P. (S.) figueroai* Thompson et Miller, *P. (S.) nakamigawai* Morikawa

et Horiguchi, *Monodiexodina kumensis* Kanmera.

From the upper part of the lower subformation the following fusulinids were described:

Neoschwagerina simplex (Ozawa), *Cancellina tenuitesta* Kanmera, *Verbeekina sphaera* Ozawa, *Yangchienia compressa* Ozawa, *Parafusulina* (*Parafusulina*) *kaerimizensis* (Ozawa), *P.* (*Skinnerella*) cf. *sapperi* (Staff), *Pseudodoliolina pseudolepida* (Deprat), *Verbeekina* sp.

Also from the topmost part of the Kozaki formation Kanmera (1963) recorded *Yabeina globosa* (Yabe), *Neoschwagerina minoensis* Deprat, *N. margaritae* Deprat, *Schwagerina* sp.

In between the area of the Yayamadake limestone and the area of the Kozaki formation there is distributed the Upper Permian Kuma formation, which is composed of sandstone, slate and conglomerate in alternation with several small lenses of limestone. This formation is again in fault contact with the other formations of different ages.

Fossils occur at three different horizons (Kanmera, 1954; Kanmera & Nakazawa, 1973).

From the lowest horizon *Lepidolina multiseptata shiraiwensis* (Ozawa), *Codonofusiella cuniculata* Kanmera and *Schwagerina* aff. *acris* Thompson et Wheeler were obtained.

The middle horizon is characterized by the association of *Lepidolina kumaensis* Kanmera, *L. multiseptata* (Deprat), *Metadoliolina gravitesta* (Kanmera), *Chusenella pseudocrassa* (Kanmera), *Codonofusiella cuniculata* Kanmera, *Dunbarula* sp., *Rauserella* sp., *Reichelina* sp., *Parareichelina subangusta* Sosnina, *Sichotenella maichensis* Sosnina, *Staffella* sp., *Nankinella* sp., *Colaniella* sp., *C.* (*Wanganella*) sp., and *Nodosaria grandecamerata* Sosnina.

The upper horizon yields *Waagenophyllum indicum* (Waagen et Wentzel), *Wentzelella* ? sp., *Codonofusiella* sp., *Pachyphloia* sp., *Nodosaria mirabilis* Lipina, *N. linae* M-Maklay, *Geinitzina* sp., *Cribrogenerina* sp., *Climacamma* sp., *Robuloides* cf. *lens* Reichel, *Hemigordius* sp., *Agathammina* sp., and *Archaeodiscus* sp.

Here it became clear that *Lepidolina kumaensis* occurs later than *Lepidolina multiseptata*, and that it is followed in turn by the uppermost horizon with *Codonofusiella* fauna.

Oita Prefecture

Quite recently the presence of *Codonofusiella* – *Palaeofusulina* fauna was ascertained by Kanmera both in the Oita and Miyazaki Prefectures (Kanmera &

Nakazawa, 1973; Working Group, 1975).

Uppermost part of the Tsukumi limestone formation in the Oita Prefecture is largely dolomitized in lithologic character. The thickness is about 60 m and the limestone is faunistically divisible into two zones.

Staffella sp., *Nankinella* sp., *Danbarula* ? sp., and *Codonofusiella* sp. occur in the upper, while *Staffella* sp., *Nankinella* sp., *Reichelina changhsingensis* Sheng et Chang, *Palaeofusulina* sp., *Codonofusiella kueichowensis* Sheng occur in the lower.

The mentioned part of limestone of the Tsukumi formation is in fault contact with *Yabeina* – *Neoschwagerina* limestone and is considered to represent the uppermost Permian there.

A very similar faunal sequence is also known in the Mitai formation in the Miyazaki Prefecture, where *Palaeofusulina* cf. *simplicata* Scheng was obtained.

Akiyoshi Limestone Plateau

In the Akiyoshi limestone plateau Permian limestone is widely developed. Permian there is supposed to be unconformably overlying the Carboniferous limestone. Also an unconformity is suspected between *Parafusulina* limestone and *Pseudofusulina* limestone. Development of carbonate facies is however terminated by the top of *Yabeina shiraiwensis* limestone which is laterally changed into slate facies. Fossils denoting the uppermost Permian has not been recorded from Akiyoshi.

Since Ozawa (1925) there are a number of important stratigraphical as well as palaeontological contributions on the Permian of Akiyoshi (e.g. Toriyama (1954–1958); Hasegawa (1958, 1963); Ota et al. (1973)).

According to Hasegawa (1963) Permian fossils from the Akiyoshi limestone are as listed below in descending order:

- Limestone of *Yabeina-Lepidolina* zone about 20 m
 Fossils: *Lepidolina shiraiwensis* (Ozawa), *L. yasubaensis* (Toriyama), *Schwagerina* spp., *Akagophyllum hasegawai* Minato et Kato, *Waagenophyllum* (s.s.) *akasakense* (Yabe), *Yatsengia* cf. *ibukiense* Minato, *Ipciphyllum subtimoricum* (Huang), *Pavastehphyllum* sp. and *Lophophyllidium* ? sp.
- Limestone of the *Neoschwagerina-Verbeekina* zone;
Gifuella douvillei subzone 25 m
 Fossils: *Gifuella douvillei* (Ozawa), *Neoschwagerina* sp. and “*Lonsdaleia-straea*” *ozawai* Minato

- Verbeekina verbeeki* subzone 35 m
Fossils: *Verbeekina verbeeki* (Geinitz) and *Schwagerina* spp.
- Neoschwagerina haydeni* subzone 7 m
Fossils: *Neoschwagerina haydeni* Dutkevich et Khabakov, *N. craticulifera* (Schwag.), *N. cf. colaniae* Ozawa, *N. sp.*, *Gifuella gifuensis* Honjo, *G. sp.*, *Pseudodoliolina pseudolepida* (Deprat), *P. ozawai* Yabe et Hanzawa, *P. sp. A.* and *Pseudofusulina gigantea* (Deprat)
- Verbeekina heimi* subzone 18 m
Fossils: *Verbeekina heimi* Thompson, *V. verbeeki* (Geinitz), *Pseudofusulina edoensis* (Ozawa), *P. gigantea* (Deprat), *Afghanella schencki* Thompson, *A. ozawai* Hanzawa, *Pseudodoliolina cf. pseudolepida* (Deprat) and *P. ozawai* Yabe et Hanzawa
- Neoschwagerina craticulifera* subzone 20 m
Fossils: *Neoschwagerina craticulifera* (Schwager), *N. irregularis* Honjo, *N. simplex* Ozawa, *N. sp.*, *Parafusulina kaerimizensis* Ozawa, *Yatsengia* sp. and *Waagenophyllum* sp.
- Limestone of the *Parafusulina* zone 40 m
Fossils: *Parafusulina kaerimizensis* Ozawa, *P. lutugini* (Schellwien), *P. spp.*, *Pseudofusulina* sp., *Afghanella schencki* Thompson, *A. ozawai* Hanzawa and *Ipciphyllum subtimoricum* (Huang)
- Limestone of the *Pseudofusulina* zone:
- Pseudofusulina krafftii magna* subzone 110 m
Fossils: *Pseudofusulina krafftii magna* Toriyama, *P. yobarensis* (Ozawa), *P. lepida* (Deprat), *P. cf. edoensis* (Ozawa), *P. cf. vulgaris globosa* (Schellwien), *Schwagerina krotowi* (Schellwien), *S. cf. regularis* (Schellwien), *S. cf. kueichihensis* (Chen), *S. sp. A.*, *Misellina claudiae* (Deprat), *Nagatoella kobayashii* Thompson, *Schubertella kingi* Dunbar et Skinner, *Staffella yobarensis* Ozawa, *S. sp.*, *Akagophyllum yabei* Minato et Kato, *A. akagoense* (Ozawa) and *Yokoyamaella yokoyamai* (Ozawa)
- Pseudofusulina vulgaris* subzone 150 m
Fossils: *Pseudofusulina vulgaris* (Schellwien), *P. vulgaris megaspherica* Toriyama, *P. globosa exilis* Toriyama, *P. watanabei* (Ozawa), *P. yobarensis* (Ozawa), *P. cf. ambigua* (Deprat), *Schwagerina satoi* (Ozawa), *S. etoi* Toriyama, *S. cf. krotowi* (Schellwien), *Dunbarinella cervicalis* (Chen), *D. cf. densa* Toriyama, *Triticites ellipsoidalis* Toriyama, *T. spp.*, *Paraschwagerina akiyoshiensis* Toriyama, *P. sp.*, and *Schubertella kingi* (Dunbar et Skinner)
- Limestone of the *Pseudoschwagerina* zone:
- Pseudoschwagerina muongthensis* subzone 100 m
Fossils: *Pseudoschwagerina muongthensis* (Deprat), *P. sp. A.*, *Triticites simplex* (Schellwien), *T. ozawai* Toriyama, *T. montipara* (Ehrenberg)

Möller), *T. pseudosimplex* Chen, *T. biconica* Toriyama, *T. cullomensis* Dunbar et Condra, *T. suzukii* (Ozawa), *T. haydeni* (Ozawa), *T. noinskyi paula* Toriyama, *T. tantula* Toriyama, *T. arctica* (Schellwien), *T. cf. ellipsoidalis* Toriyama, *T. spp.*, *Dunbarinella* ? sp., *Schwagerina* cf. *satoi* (Ozawa), *S. regularis* (Schellwien), *Paraschwagerina* cf. *akiyoshiensis* Toriyama and *Schubertella kingi* Dunbar et Skinner

Triticites simplex subzone 100 m

Fossils: *Triticites simplex* (Schellwien), *T. ozawai* Toriyama, *T. montipara* (Ehrenberg Möller), *T. pseudosimplex* Chen, *T. biconica* Toriyama, *T. cullomensis* Dunbar et Condra, *T. suzukii* (Ozawa), *T. noinskyi paula* Toriyama, *T. arctica* (Schellwien), *T. spp.*, *Quasifusulina longissima* (Möller), *Schubertella kingi* Dunbar et Skinner

Apart from fusulinids and corals listed above some bryozoans and algae have been added to the faunal as well as floral contents of the Akiyoshi limestone. But other group of fossils, being scarce except for abundant crinoid stem joints, have been little investigated.

Atetsu Limestone Plateau

On the Permian part of the Atetsu limestone there are two major biostratigraphical contributions; namely by Nogami 1961a,b, 1962 and by Sada, 1965.

Here we follow Nogami for the division and faunal contents of the Permian of Atetsu.

Nogami's scheme is tabulated in the accompanied chart, and his zones and subzones with fossil contents are listed below in ascending order.

Pseudoschwagerina subsphaerica – *Quasifusulina longissima ultima* Zone

Quasifusulina longissima ultima – “*Pseudoschwagerina*” *nakazawai* subzone
Oolitic limestone of about 50–70 m thick.

Fusulinids are *Triticites ozawai* Toriyama, *T. montiparus* [Ehrenberg (Möller)], *T. gravitesta* Nogami, *T. obai* Toriyama, *T. haydeni* (Ozawa), *T. pseudoarcticus* Rauser-Chernousova, *T. aff. contractus* (Schellwien), *T. spp.*, “*Schwagerina*” aff. *alpina* (Schellwien), *Rugosofusulina* aff. *serrata* Rauser-Chernousova and “*Pseudoschwagerina*” *nakazawai* Nogami.

Pseudoschwagerina subsphaerica subzone

Muddy limestone of 40 m thick.

Fusulinids are *Pseudoschwagerina subsphaerica* Nogami, *Paraschwagerina kanmerai* Nogami, *Pseudofusulina atetsuensis* Nogami, *Schwagerina*

primigena Nogami, *Triticites obai* Toriyama, *T. pseudoarcticus* Rauser-Chernoussova, *T. sp.*, and “*Schwagerina*” aff. *alpina* (Schellwien).

Pseudofusulina vulgaris Zone

About 80 thick conglomeratic limestone.

Fossils are *Pseudofusulina vulgaris* (Schellwien), *Psf. vulgaris globosa* (Schellwien), *Psf. vulgaris exigua* (Schellwien), *Schwagerina* spp., *Psf. krafftii magna* Toriyama, *Psf. krafftii* form A, *Psf. aff. fusiformis* (Schellwien), *Nagatoella kobayashii* Thompson, and “*Triticites*” *pseudosimplex* Chen.

Parafusulina kaerimizensis – *Pseudofusulina krafftii magna* Zone

Pseudofusulina krafftii magna subzone

Massive limestone and conglomeratic limestone.

Thickness is about 40–60 m.

Fossils are *Pseudofusulina krafftii magna* Toriyama, *Psf. krafftii* form A, *Psf. globosa exilis* Toriyama, *Psf. aff. fusiformis* (Schellwien), *Nagatoella kobayashii* Thompson, *Schwagerina* sp., *Psf. vulgaris* (Schellwien), *Psf. vulgaris globosa* (Schellwien), *Psf. aff. edoensis* (Ozawa), *Schwagerina semilucera* Nogami, *Sch. semilucera grande* Nogami, *Rauserella* sp. and *Misellina* sp.

Parafusulina kaerimizensis subzone

Massive limestone of about 10 m thick.

Fossils are *Parafusulina kaerimizensis* (Ozawa), *Pseudodoliolina ozawai* Yabe et Hanzawa, and *Pseudofusulina* aff. *edoensis* (Ozawa).

Neoschwagerina douvillei – *N. craticulifera* Zone

Neoschwagerina craticulifera subzone

Conglomeratic limestone. Thickness unknown.

Fossils are *Neoschwagerina craticulifera* (Schwager), *Afghanella* sp., *Pseudodoliolina ozawai* Yabe et Hanzawa, *Pseudofusulina* aff. *edoensis* (Ozawa), *Psf. kusamensis* Nogami, *Parafusulina kaerimizensis* (Ozawa), and *Rauserella* sp.

Neoschwagerina douvillei – *N. margaritae* subzone

Mostly conglomeratic limestone of about 60 m thick.

Fusulinids are *Neoschwagerina douvillei* Ozawa, *N. margaritae* Deprat, *N. cheni* Sheng, *Sumatrana annae* Volz, *Pseudodoliolina pseudolepida* (Deprat), *Pseudofusulina kusamensis* Nogami, *Verbeekina verbeeki* (Geinitz), *Schwagerina globularis* (Gubler), *Neoschwagerina* sp., *Pseudodoliolina* aff. *lepida* (Schwager), *Rauserella* sp., and *Nankinella* sp.

Yabeina shiraiwensis Zone

Yabeina shiraiwensis – *Y. sp. A* subzone

Lithology is conglomeratic limestone in the lower part and is slate with

limestone lenses in the upper part. Thickness is not described.

Yabeina shiraiwensis Ozawa, *Y. sp. A*, *Sumatrina longissima* Deprat, *Pseudodoliolina pseudolepida* (Deprat), *Schwagerina otai* Nogami, *Verbeekina verbeeki* (Geinitz), *Schwagerina sp.*, *Nankinella sp.*, *Schubertella ? sp.*, *Neoschwagerina margaritae* Deprat, *N. douvillei* Ozawa and *Sumatrina annae* Volz.

Yabeina shiraiwensis subzone

Mostly slate with a maximum thickness of about 100 m.

Fossils are *Yabeina shiraiwensis* Ozawa, *Neoschwagerina sp.* and *Sumatrina sp.*

Akasaka Limestone

The Akasaka limestone occupies a small hill, which is bounded to the west by a fault with non calcareous slate presumably of Permian in age. The limestone is covered by Quaternary sediments to the east. Therefore known succession represents "middle" part of Permian only, although fossils are abundant there.

Since the time of Gumbel (1878), Schwager (1883), Gottsche (1884) and Yabe (1902), Akasaka drew the attention of many palaeontologists.

Biostratigraphical studies are by Deprat (1914), Ozawa (1927), Akasaka Research Group (1956), Honjo (1959).

From the Akasaka limestone Honjo (1959) listed in descending order the following fossils:

<i>Yabeina</i> Zone	100 m
<i>Yabeina globosa</i> (Yabe), <i>Y. katoi</i> Ozawa, <i>Neoschwagerina minoensis</i> Deprat.	
<i>Neoschwagerina</i> Zone	
<i>Gifuella douvillei</i> subzone	55 m
<i>Gifuella cf. douvillei</i> (Deprat)	
<i>Yabeina ozawai</i> subzone	90 m
<i>Verbeekina sphaera</i> Ozawa, <i>Yabeina ozawai</i> Honjo, <i>Pseudofusulina spp.</i> , <i>Gifuella amricula</i> Honjo, <i>G. gifuensis</i> Honjo, <i>Neoschwagerina colaniae</i> Ozawa, <i>N. haydeni</i> (D. et K.), <i>Metaschwagerina ovalis</i> Minato et Honjo, <i>Bellerophon sp.</i>	
<i>Neoschwagerina craticulifera</i> subzone	62 m
<i>Neoschwagerina craticulifera</i> (Schwager), <i>N. irregularis</i> Honjo, <i>N. haydeni</i> (D. et K.)	
<i>Pseudodoliolina ozawai</i> subzone	40 m

- Pseudodoliolina ozawai* Yabe et Hanzawa, *Pseudofusulina* spp., *Verbeekina verbeeki* (Geinitz), *Verbeekina* cf. *heimi* Thompson et Foster
Minoella nipponica subzone 25 m
Neoschwagerina (Minoella) nipponica Ozawa, *N. (M.) eonipponica* Honjo,
N. simplex Ozawa, *N. sphaerica* M. Maclay, *Verbeekina minatoi* Honjo
(MS), *Pseudofusulina* spp., *Pseudodoliolina ozawai* Yabe et Hanzawa

Many fossils other than fusulinids have been described from the Akasaka limestone. They may be listed below.

Yabeina globosa Zone

Pelecypoda:

Etheripecten ? *hayasakai* (Murata) = *Aviculopecten interstitialis* of Hayasaka, 1925, *Hayasakapecten minoensis* (Hayasaka), *H. reticularis* (Hayasaka), *Waagenoperna hayamii* Nakazawa et Newell, = *Liebea sinensis* of Hayasaka, 1925, *Schizodus japonicus* (Hayasaka), *Solenomorpha elegantissima* Hayasaka.

Scaphopoda:

Dentalium akasakaensis Hayasaka, *D. neornatum* Hayasaka, *D. (Laevidentalium)* cfr. *priscum* Münster, *D. (Plagioglypta) herculeum* deKoninck.

Gastropoda:

Bellerophon jonesianus deKoninck, *B. jonesianus hiulciformis* Hayasaka, *Pleurotomaria* aff. *multicarinata* Mansuy, *P. yokoyamai* Hayasaka, *Murchisonia yabei* Hayasaka, *Solenospira multicostata* Hayasaka, *Naticopsis wakimizui* Hayasaka, *N. minoensis* Hayasaka, *N. fasciata* Hayasaka, *N.* cfr. *praealta* Wanner, *Naticella japonica* Hayasaka, *T. magna* Hayasaka, *Spiromphalus yabei* Hayasaka, *Porcellia puzoides* Hayasaka, *Euconospira nipponica* Hayasaka.

Cephalopods:

Koninckoceras sp. = *Stearoceras* sp., *Coelogastrioceras giganteum* Nakazawa, *Stacheoceras* sp.

Corals:

Pseudofavosites sp., *Pseudoromingeria kotoi* (Yabe et Hayasaka), *Waagenophyllum (Waagenophyllum) polyseptatum* Minato, *W. (W.) indicum* (Waagen et Wentzel), *W. (W.) akasakense* (Yabe), *W. (W.) compactum* Minato et Kato, *W. (W.) pulchrum* Hamada.

Trilobita:

Pseudophillipsia obtusicauda Kayser.

Calcareous Algae:

Mizzia velebitana Schubert, *M. yabei* Karpinksky, *Solenopora yabei* Endo, *Epimastopora kanumai* Endo, *Ortonella akasakaensis* Endo. (All these algal species are said to occur also from the *Neoschwagerina* limestone in Akasaka).

Neoschwagerina Zone

Hydrozoa ?:

Stromatopora (Paralleropora) minoensis Yabe et Sugiyama.

Corals:

Praewentzelella honjoi Minato et Kato.

Brachiopods:

Squamularia cfr. *inequilateralis* Gemmelaro, *S. asiatica* Chao, *S. waageni* Frech, *Terebratuloida* sp., *Leptodus richthofeni* Kayser, *Scacchinella* cfr. *gigantea* Gemmelaro, *Enteletes akasakensis* Ozawa, *E. minoensis* Hayasaka, *E. suessi* Schellwien, *Geyerella* sp.

Echinoid:

Miocidaris spinulifera Nishiyama.

Incertae sedis:

Shikamaia akasakaensis Ozaki.

Omi Limestone

Permian fusulinids at Omi were first described by Hayasaka (1924), which is followed by a number of biostratigraphical works (Kawada, 1954; Fujita, 1958; Hasegawa et al, 1969, etc.).

To date, however, a detailed description on its fauna is still not available, although overall faunal characteristics in the Omi limestone are quite like those of the Atetsu and the Akiyoshi.

Based on the work of Hasegawa et al (1969) the sequence of fusulinids in Omi may be tabulated in descending order as below:

Yabeina-Lepidolina Zone

“*Yabeina*” *hayasakai* Ozawa, *Sumatrina annae* Volz, *Reichelina* sp.,
Dunbarula sp., *Kahlerina* sp., *Chusenella* sp.

Neoschwagerina-“*Gifuella*” Zone

Neoschwagerina craticulifera (Schwager), *Gifuella* sp., *Gublerina* sp.,
Pseudodoliolina sp., *Neoschwagerina* sp.

Parafusulina Zone

Parafusulina kaerimuzensis (Ozawa), etc.

Pseudofusulina Zone

Pseudofusulina krafftii subzone

Pseudofusulina krafftii (Schellwien), *Schwagerina krotowi* (Schellwien),
Nagatoella kobayashii Thompson, *Yangchientia* sp., *Misellina claudiae*
(Deprat), *Staffella* sp.

Pseudofusulina vulgaris subzone

Pseudofusulina vulgaris (Schellwien), *Paraschwagerina akiyoshiensis*
Toriyama, *Schwagerina satoi* (Ozawa), *Schwagerina krotowi* (Schellwien),
Eoparafusulina ellipsoidalis (Toriyama), *Triticites* spp., *Schubertella* sp.

Pseudoschwagerina Zone

Pseudoschwagerina sp. subzone

Pseudoschwagerina sp., etc.

Triticites simplex subzone

Triticites simplex (Schellwien), *Triticites* spp.

Correlation

As described above, ammonoid cephalopods are extremely scarce within the Japanese Permian (Table 1).

Table 1 Sequence of Permian ammonites in the Kitakami mountains
(after Bando, 1975)

Age	Series	Ammonoid Zones
Late Permian	Toyoma Series	<i>Araxoceras</i> cf. <i>kiangsiensis</i> Zone <i>Eumedlicottia</i> Zone
Early Permian	Kanokura Series	<i>Timorites intermedium</i> Zone <i>Stacheoceras iwaizakiense</i> Zone
	Sakamotozawa Series	(Unknown)

They are by no means sufficient for the correlation. The occurrence of corals is quite sporadic in the Japanese Permian. And Bryozoans and algae are both relatively long ranging. Conodont studies began only quite recently. Only fusulinids have been studied from various districts of Japan, so far. They occur abundantly, and therefore we have to rely heavily upon these fusulinids for the sake of Permian correlation (Toriyama, 1967).

Table 2 Succession of Japanese Permian Fusulinids

<i>Palaeofusulina</i> + <i>Colaniella parva</i> (<i>Araxoceras</i>)	
<i>Lepidolina kumaensis</i> + <i>L. minatoi</i>	
<i>Yabeina globosa</i> + <i>Lepidolina multiseptata</i>	
" <i>Neoschwagerina</i> " <i>douvillei</i>	
<i>Neoschwagerina margaritae</i>	
<i>Neoschwagerina craticulifera</i>	
<i>Neoschwagerina simplex</i> + <i>Parafusulina kaerimizensis</i>	
<i>Cancellina</i> , <i>Minoella</i> , <i>Maklaya</i>	*
<i>Misellina</i> , <i>Toriyamaia</i> , <i>Nagatoella</i>	*
<i>Pseudofusulina krafftii</i> + <i>P. ambigua</i>	*
" <i>Pseudofusulina</i> " <i>vulgaris</i> + <i>Paraschwagerina</i>	
<i>Pseudoschwagerina</i> + <i>Triticites</i>	
<i>Triticites</i> + primitive <i>Pseudoschwagerina</i> + <i>Quasifusulina</i>	

* distinction between these three assemblages has not been firmly established as yet.

Table 2 is a scheme of fusulinid succession in Japan, which is applicable to carbonate facies, and to such mixed environment that we have in the Kitakami mountains, to some extent.

Figure 2 is an attempt to correlate the Japanese Permian in the districts described above. As may be seen from the figure, three fold division of the Permian System may be tenable in Japan, especially in the Kitakami mountains.

International correlation posed difficulty since the world standard regions for Permian, e.g. Uralian and North American basin, are all beyond the Tethyan faunal realm.

With South China, however, the Japanese Permian correlates quite well.

Lower part of the Kawaguchi Stage of the Sakamotozawa Series, with the entry of *Pseudoschwagerina* is correlatable with the Chuanshanian and

Mapingian in China.

The Chihisian in South China now comprises both *Misellina* subzone and *Cancellina* subzone, the latter of which yields *Pseudodoliolina* (Sheng and Lee, 1974). Thus it is correlatable with the upper part of the Kawaguchi Stage and the Kabayama Stage, both of the Sakamotozawa Series, and the lower part of the Kattisawa Stage of the Kanokura Series, from which *Cancellina* sp. was obtained (Choi, 1970). The rest of the Kanokura Series is correlatable with the Maokou limestone which comprises both *Neoschwagerina* and *Yabeina* zones.

In Japan, *Lepidolina kumaensis* and *L. minatoi* occur a little later than the first occurrence of *L. multiseptata*. It is certain that the top part of the Maokou limestone with *Yabeina* fauna corresponds to the Japanese *Yabeina globosa-Lepidolina multiseptata* Zone. But whether the part does include deposits corresponding to the Japanese *L. kumaensis* – *L. minatoi* zone or not is still an open question.

The newly proposed Maehama Stage corresponds to the Wuchiaping limestone based on the common occurrence of *Araxoceras* both in China and Japan. Also the Chinese Changhsing limestone is equivalent to the Nabekoshi Stage in the Kitakami mountains with *Colaniella* – *Palaeofusulina* fauna. Thus almost the whole Lopingian is correlatable with the Toyoma Series (Table 3).

Table 3 Correlation of the Permian of Japan, South China and the USSR

Kitakami Mts., Japan (Minato et al, this paper)		South China (Sheng & Lee, 1974)			U. S. S. R. (Correlation after Kahler, 1974)
Toyoma Series	Nabekoshi Stage	Loping Series	Changhsing Lst.	<i>Palaeofusulina</i>	Tatarian
	Maehama Stage		Wuchiaping Lst.	<i>Codonofusiella</i>	
Kanokura Series	Iwaizaki Stage	Yangsing Series	Maokou Lst.	<i>Yabeina</i>	Kazanian Ufimian
	Kattisawa Stage			<i>Neoschwagerina</i>	
Sakamotozawa Series	Kabayama Stage		Chihisia Lst.	<i>Cancellina</i>	Kungurian
	Kawaguchi Stage	Chuanshan Lst.	<i>Misellina</i>	Artinskian	
			<i>Pseudoschwagerina</i>	Sakmarian Asselian	

For further international correlation we have no new data of our own. But such an attempt based on fusulinids was presented by Kahler (1974), and we wish to follow him.

Thus the lower part of the Kawaguchi Stage, *Lophocarinophyllum suetomii* Zone, is the Asselian, being situated below the bed with *Pseudoschwagerina schellwieni* and *Zellia* of Sakmarian. The upper part of the Kawaguchi Stage, especially the earlier half of it is correlatable with the Sakmarian with the entry of *Pseudoschwagerina schellwieni*. The Artinskian starts with the appearance of *Pseudofusulina vulgaris* and may be equivalent to the rest of the Sakamotozawa Series. The Kungurian is designated by Kahler as *Cancellina* zone, which in the Kitakami mountains may be equivalent to the lower part of the Kattisawa Stage. The Kazanian and the Ufimian are both thought by Kahler as possible equivalent of *Neoschwagerina* zone, which is correlated with the later half of the Kattisawa Stage of the Kanokura Series. Tatarian starts from the *Yabeina* – *Lepidolina* zone. Therefore the Iwaizaki Stage of the Kanokura Series and the Toyoma Series may be the Tatarian.

References

- Akasaka research group, 1956. Geological studies of the Akasaka limestone. *Earth Science (Chikyu Kagaku)*, 26-27: 10-18.
- Asama, K., 1956. Permian plants from Maiya in Northern Honshu, Japan. *Proc. Japan Acad.*, 32(7): 469-471.
- Bando, Y., 1975. On Some Permian Medicottidae from the Toyoma Formation in the Kitakami Massif. *Mem. Fac. Education, Kagawa Univ.*, II, 25(2): 67-81.
- Chao, K., 1965. The Permian ammonoid bearing formations of South China. *Scientia Sinica*, 14: 1813-1825.
- Chisaka, T., 1953. On the Permian System of the southwestern part of the Kitakami Mountainsland (Maiya District). *Studies Geol. Min. Inst. Tokyo Univ. Education*, 2: 1-9.
- Choi, D.R., 1970. On some Permian fusulinids from Iwaizaki, N.E. Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 14(3): 313-325.
- Choi, D.R., 1970. Permian fusulinids from Imo, Southern Kitakami mountains, N.E. Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 14(3): 327-354.
- Choi, D.R., 1972. *Colania douvillei* (Ozawa), a fusulinid foraminifera from the northern Kitakami mountains, N.E. Japan. *Trans. Proc. Palaeont. Soc. Japan*, N.E., 86: 369-374.
- Choi, D.R., 1972. Classification and phylogeny of *Misellina* with description of *Misellina* from the Lower Permian in the Southern Kitakami mountains, N.E. Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 15(3-4): 625-648.
- Choi, D.R., 1972. Discovery of Uralian fusulinids from the Upper Permian conglomerate in the southern Kitakami mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 15(3-4): 479-492.
- Choi, D.R., 1973. Permian fusulinids from the Setamai-Yahagi district, Southern Kitakami mountains, N.E. Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 14(1): 1-13.

- Choi, D.R., 1976. Distribution of the Upper Permian Fusulinids with Relation to Limestone Lithofacies in the Southern Kitakami Mountains, N.E. Japan. *Jour. Geol. Soc. Japan*, 82(2): 113-125.
- Deprat, J., 1914. Etude des Fusulinides du Japon, de Chine et d'Indochine et classification des calcaires a fusulines (III^e Memoir). Etude comparative des Fusulinides d'Akasaka (Japon) et des Fusulinides de Chine et d'Indochine. *Mém. Serv. Geol. l'Indochine*, 3(1): 1-45.
- Endo, R., 1951. Several new species from the Sakamotozawa section, Hikoroichi-mura, Kesen-gun, in the Kitakami mountainous land, Later Palaeozoic calcareous algae in Japan, 1. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 4: 121-129.
- Endo, R., 1952. Several previously described species from the Sakamotozawa section, Hikoroichi-mura, Kesen-gun, in the Kitakami mountainous land, Later Palaeozoic calcareous algae in Japan 2. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 5: 139-144.
- Endo, R., 1952. A new species from the Maiya section, Maiya-machi, Tome-gun, Miyagi-ken, Later Palaeozoic calcareous algae in Japan 3. *Sci. Rep. Saitama Univ.*, 1(1): 23-28.
- Endo, R., 1953. Several species from the Iwaizaki limestone, Motoyoshi-gun, in the Kitakami mountainous land, Later Palaeozoic calcareous algae in Japan, 5. *Japan. Jour. Geol. & Geogr.*, 23: 117-126.
- Endo, R., 1953. Stratigraphical and Palaeontological studies of the later Palaeozoic calcareous algae in Japan, 6. *Sci. Rep. Saitama Univ.*, Ser.B, 1(2):
- Endo, R., 1954. Stratigraphical and Palaeontological studies of the later Palaeozoic calcareous algae in Japan, 8. *Sci. Rep. Saitama Univ.*, Ser.B, 1(3):
- Endo, R., 1961. Stratigraphical and Palaeontological studies of the later Palaeozoic calcareous algae in Japan, 17, Fossil algae from the Akiyoshi Limestone Group. *Sci. Rep. Saitama Univ.*, Ser.B, 3(3): 119-142.
- Endo, R. and Matsumoto, E., 1962. Permo-Carboniferous Trilobites from Japan. *Sci. Rep. Saitama Univ.*, ser.B, 4(2): 149-172.
- Fujita, H., 1958. Geology of Omi Limestone (Permo-Carboniferous), Niigata Prefecture. *Contr. Inst. Geol. & Palaeont., Tohoku Univ.*, 48: 1-60.
- Gottsche, C., 1884. Ueber japanisches Karbon. *Zeit. deutsch. geol. Gesell.*, 36: 653.
- Gümbel, C.W., 1878. Anteilungen zu wissenschaftlichen Beobachtungen auf Alpen-reisen. *Beil. Zeit. deutsch. u. österr. Alpenvereins*, 1878, 1-192.
- Hasegawa, Y., 1958. Note on the geologic structure of Akiyoshi plateau SW Honshu. *Earth Science (Chikyu Kagaku)*, 38: 15-18.
- Hasegawa, Y., 1963. New Find of Fossils in the Reddish Tuffaceous Shale in the Akiyoshi Province. *Earth Science (Chikyu Kagaku)*, 64: 32-37.
- Hasegawa, Y., 1964. On phrenotheca in *Pseudofusulina vulgaris globosa*. *Earth Science (Chikyu Kagaku)*, 74: 13-16.
- Hasegawa, Y., 1966. On the proloculus size of *Lepidolina shiraiwensis* (Ozawa). *Sci. Rep. Dept. Geol. Min., Niigata Univ.*, 1: 21-29.
- Hasegawa, Y., 1973. "Chronological and geographical change of fusulinid assemblages in the Japanese island". *Jour. Fac. General Education, Niigata Univ.*, 3: 51-59.
- Hasegawa, Y. et al., 1969. Palaeozoic of the Omi district, Niigata Prefecture. Geological field excursion guide book. *Geol. Soc. Japan*, 1-23.
- Hanzawa, S., 1941. The stratigraphical relation between the Carboniferous and Permian formation in the Manchuria, Korea and Japan Proper. *Japan. Jour. Geol. & Geogr.*, 17(3): 97-108.
- Hanzawa, S., 1944. Stratigraphical distribution of fusulinid foraminifera found in south Manchuria and Japan. *Japan. Jour. Geol. & Geogr.*, 19(1-4): 1-10.
- Hanzawa, S. and Murata, M., 1963. The paleontologic and stratigraphic considerations on the Neoschwagerininae and Verbeekininae, with the descriptions of Fusulinid

- Foraminifera from the Kitakami Massif, Japan. *Sci. Rep., Tohoku Univ.*, 2nd Ser., 35(1): 1-31.
- Hayasaka, I., 1924. On the fauna of the anthracolithic limestone of Omi-mura in the western part of Echigo. *Sci. Rep. Tohoku Imp. Univ.*, 2nd Ser., 8(1): 1-82.
- Hayasaka, I., 1924. Fossil in the roofing slate of Ogachi Prov. Rikuzen. *Japan. Jour. Geol. Geogr.*, 3: 45-53.
- Hayasaka, I., 1925. On some Palaeozoic molluscs of Japan, 1. *Sci. Rep. Tohoku Imp. Univ.*, 2nd Ser., 8(2):
- Hayasaka, I., 1925. On some brachiopods from the *Lyttonia* horizon of the Kitakami mountains. *Japan. Jour. Geol. & Geogr.*, 4:
- Hayasaka, I., 1932. On three brachiopod species of the subfamily Orthotetinae in Fusulina-Limestone of Kinsyozan, Akasaka-machi, Prov. Mino, Japan. *Mem. Fac. Sci., Agr. Taihoku Imp. Univ.*, 6(1):
- Hayasaka, I., 1938. Two species of *Trachydomia* from Japan. *Mem. Sci. Agri. Taihoku Imp. Univ.*, 22(1): 1-8.
- Hayasaka, I., 1939. *Spiromphalus*, a new gastropod genus from the Permian of Japan. *Mem. Fac. Sci. Agri. Taihoku Imp. Univ.*, 22, (2): 19-23.
- Hayasaka, I., 1940. On two Permian Ammonooids from the Kitakami mountains, North Japan. *Jour. Geol. Soc. Japan*, 46(565):
- Hayasaka, I., 1943. On some Permian gastropods from Kinshozan, Akasaka-machi, Gifu Prefecture. Part I. *Mem. Fac. Sci., Taihoku Imp. Univ.*, Ser.III, 1(2): 23-48.
- Hayasaka, I., 1953. *Hamletella*, a new Permian genus of brachiopoda, and a new species from the Kitakami mountains. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 12:
- Hayasaka, I., 1954. Younger Palaeozoic Cephalopods from the Kitakami mountains. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 8(4):
- Hayasaka, I., 1954. *Euconospira* with color markings from the Permian of Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 8(4): 349-360.
- Hayasaka, I., 1954. An occurrence of *Koninckoceras* from the Japanese Permian. *Japan. Jour. Geol. & Geogr.*, 25(1-2): 57-59.
- Hayasaka, I., 1955. A new Permian species of *Porcellia* from Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 9(1): 21-24.
- Hayasaka, I., 1960. A new Permian species of *Foordiceras* from the Kitakami mountains. *Sci. Rep. National Sci. Museum*, 47:
- Hayasaka, I., 1960. On the occurrence of *Neospirifer fasciger* (Keyerling) in Japan, and a note on some associate Permian brachiopods from around Kesen-numa City, Northeast Japan. *Collection of essays in commemoration of the Tenth Anniversary (1959) of Shimane University (Natural science)*.
- Hayasaka, I., 1963. Some Permian fossils from the Southern Kitakami, I. *Proc. Japan Acad.*, 39(7): 474-478.
- Hayasaka, I., 1963. Some Permian fossils from Southern Kitakami, II, two brachiopod species. *Proc. Japan Acad.*, 39(7): 479-483.
- Hayasaka, I., 1963. Some Permian fossils from Southern Kitakami, III, Ammonoidea. *Proc. Japan Acad.*, 39(8): 594-599.
- Hayasaka, I., 1963. Some Permian fossils of Southern Kitakami, IV, Brachiopod Superfamily Orthotetacea Williams. *Proc. Japan Acad.*, 39(10):
- Hayasaka, I., 1964. Some Permian fossils from Southern Kitakami, V, *Aulosteges* sp. indet. *Proc. Japan Acad.*, 40(7): 528-532.
- Hayasaka, I., 1966. Some Permian fossils from Southern Kitakami, VI, Three brachiopods. *Proc. Japan Acad.*, 42(10): 1223-1228.
- Hayasaka, I., 1967. Some Permian fossils from Southern Kitakami, VII, Two species of *Phymatifer*. *Proc. Japan Acad.*, 43: 143-147.

- Hayasaka, I., 1967. Some Permian fossils from Southern Kitakami, VIII. *Proc. Japan Acad.*, 43(5): 378-383.
- Hayasaka, I., 1967. Some Permian fossils from Southern Kitakami, IX. Two pelecypod genera, *Allorisma* and *Myoconcha*. *Proc. Japan Acad.*, 43(6): 517-521.
- Hayasaka, I. and Minato, M., 1956. Some brachiopods from the Lower Kanokura series of the Kitakami mountains, Japan. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 21:
- Honjo, S., 1959. Neoschwagerinids from the Akasaka Limestone (A Palaeontological Study of the Akasaka Limestone, 1st Report). *Jour. Fac. Sci., Hokkaido Univ.*, Ser.4, 10(1): 11-161.
- Honjo, S., 1960. A study of some primitive *Neoschwagerina* by a new serial section technique. *Jour. Fac. Sci., Hokkaido Univ.*, Ser.4, 10(3): 457-470.
- Ishii, K., Okimura, Y. and Nakazawa, K., 1975. On the Genus *Colaniella* and its Biostratigraphic significance. *Jour. Geosci., Osaka City University*, 19(6): 107-138.
- Ishizaki, K., 1964. Middle Permian ostracods from the Iwaizaki limestone, northeast Japan. *Tohoku Univ., Sci., Rep.* 2nd ser., 36(1): 139-160.
- Ishizaki, K., 1967. Ostracods from the Lower Permian Tassobe formation, Northeast Japan. *Saito Ho-on kai Museum Bull.*, 36: 49-66.
- Kahler, F., 1974. Fusuliniden aus T'ien-schan und Tibet mit Gedanken zur Geschichte der Fusuliniden-Meer im Perm. *The Sino-Swedish Expedition*, 4: 1-148.
- Kanmera, K., 1952. The Upper Carboniferous and the Lower Permian of the Hikawa Valley, Kumamoto Pref., Kyushu, Japan. *Jour. Geol. Soc. Japan*, 58(676): 17-32.
- Kanmera, K., 1954. Fusulinids from the Upper Permian Kuma Formation, Southern Kyushu, Japan- With Special Reference to the Fusulinid Zone in the Upper Permian of Japan. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 4(1): 1-38.
- Kanmera, K., 1958. Fusulinids from the Yayamadake Limestone of the Hikawa Valley, Kumamoto Prefecture, Kyushu, Japan. Part III. -Fusulinids of the Lower Permian. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 6(3): 153-215.
- Kanmera, K., 1963. Fusulines of the Middle Permian Kozaki Formation of Southern Kyushu. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 14(2): 79-141.
- Kanmera, K. and Mikami, T., 1965. Succession and sedimentary features of the Lower Permian Sakamotozawa formation. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 16(3): 265-274.
- Kanmera, K. and Mikami, T., 1965. Fusuline Zonation of the Lower Permian Sakamotozawa Series. *Mem. Fac. Sci., Kyushu Univ.*, ser.D, 16(3): 275-320.
- Kanmera, K. and Nakazawa, K., 1973. Permian-Triassic relationships and faunal change in the Eastern Tethys, The Permian and Triassic systems and their mutual boundary. *Canada Soc. Petrol. Geol.*, 2: 100-119.
- Kanmera, K. and Toriyama, R., 1976. The Evolution and Extinction Patterns of Permian Fusulinaceans. *Geol. & Palaeont. Southeast Asia*, 17: 129-154.
- Karpinsky, A., 1908. Einige problematischen Fossilien aus Japan. *Verhandl. d. Russische-Kaiserlichen Mineral Gesell., St. Petersburg*, 2nd Ser., 46: 255-272.
- Kato, M., Kumano, S. and Takahashi, T., 1970. A new occurrence of *Cordaites* from the Lower Permian of southe Kitakami mountains. *Earth Science (Chikyū Kagaku)*, 24(1): 40.
- Kawada, S., 1954. Stratigraphical and Palaeontological studies of the Omi Limestone in the Itagamine district, Niigata Prefecture. *Bull. Tokyo Univ. Education, Geol. & Min. Inst.*, 32: 15-27.
- Kobayashi, T., 1945. *Notaculites toyomensis*, a new trail found in the Upper Permian Toyoma series in Nippon. *Japan. Jour. Geol. Geogr.*, 20: 13-18.
- Koizumi, H., 1975. *Paleozoic Cephalopods of Japan*. Kanaeishi Bunko, Chiba
- Kon'no, E., 1973. New Species of *Pleuromeia* and *Neocalamites* from the Upper Scythian

- Beds in the Kitakami Massif, Japan, with a brief note on some Equisetacean Plants from the Upper Permian Beds in the Kitakami Massif. *Sci. Rep. Tohoku Univ.*, Second ser., 43(2): 99-115.
- Mabuchi, S., 1935. Stratigraphy of the Iwaizaki Limestone. *Saito Ho-on Kai Jiho*, 101: 1-18.
- Minato, M., 1942. Unconformity of the Pre-Sakamotozawa stage (Pre-Sakmarian) in the Kitakami mountainland, Northeast Japan. *Jour. Geol. Soc. Japan*, 49(581): 47-72.
- Minato, M., 1955. *Isogramma paotechowensis* (Grabau and Chao) from the Permian of Japan. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 18: 29-30.
- Minato, M., 1955. Japanese Carboniferous and Permian Corals. *Jour. Fac. Sci., Hokkaido Univ.*, Ser.IV, 9(2): 1-202.
- Minato, M. (Chief Editor), 1965. *The geologic development of the Japanese Island*. Tsukiji Shokan, Tokyo, 1-142.
- Minato, M., 1966. Palaeozoic stratigraphy in the Kitakami mountains and the Abean movements in Japan. *Prof. S. Matsushita Mem. Vol.*, 143-156.
- Minato, M., 1975. Japanese Palaeozoic Corals. *Jour. Geol. Soc. Japan*, 81(2): 103-126.
- Minato, M., Hashimoto, S., Suyama, K., Takeda, Y., Suzuki, Y., Kakimi, T., Yamada, K., Kimura, S., Ishikawa, T., and Suetomi, H., 1954. Biostratigraphie der permischen Formation des Setamai-Geländes in Sud-Kitakami Gebirge. (Stratigraphische und tektonische Untersuchungen des japanischen Palaeozoikums, Teil 9). *Geol. Soc. Japan. Jour.*, 60(708): 378-387.
- Minato, M. & Honjo, S., 1958. Shell structure of *Metaschwagerina* n.g. from Akasaka limestone. *Earth Science (Chikyu Kagaku)*, 38: frontispiece.
- Minato, M. & Honjo, S., 1959. The axial septula of some Japanese Neoschwagerininae with special remarks of the phylogeny of the subfamily Neoschwagerininae Dunbar and Condra, 1928. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 10(2): 305-336.
- Minato, M. and Honjo, S., 1965. On *Colania kwangsiana* Lee. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 13(7): 1-5.
- Minato, M. and Kato, M., 1965. Durhaminidae (Tetracoral). *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 13(1): 13-86.
- Minato, M. and Kato, M., 1965. Waagenophyllidae. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 12(3-4): 1-241.
- Minato, M., Kato, M. & Hasegawa, Y., 1964. A note on the boundary between *Pseudofusulina* and *Parafusulina* zones in Japan. *Proc. Japan. Acad.*, 40(10): 827-831.
- Minato M, and Nakamura K, (1956): On four brachiopod species of the Subfamily Orthotetinae from the Kanokura series of the Kitakami mountains, Japan. *Trans. Palaeont. Soc. Japan*, N.S., no.21, pp.149-155.
- Minato, M. and Rowett, C.L., 1967. On a new species of *Wentzellophyllum* Hudson from the Kitakami mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 13(4): 343-348.
- Morikawa, R., 1958. Fusulinids from the Akasaka limestone (Part 1). *Sci. Rep. Saitama Univ.*, ser.B, 31(1): 93-130.
- Morikawa, R., Sato, T., Shibasaki, T., Shinada, Y., Okubo, M., Nakazawa, K., Horiguchi, M., Murata, M., Kikuchi, Y., Taguchi, M. and Takahashi, K., 1958. Stratigraphy and biostratigraphy of the "Iwaizaki Limestone" in the Southern Kitakami mountainland. *Jubilee Publ. Commen. Prof. H. Fujimoto Sixtieth Birthday*. 81-90.
- Morikawa, R. & Suzuki, Y., 1961. Fusulinids from the Akasaka Limestone (Part 2). *Sci. Rep. Saitama Univ.*, ser.B, 4(1): 43-74.
- Murata, M., 1964. Geological age of the Kanokura Formation in the southern part of the Kitakami Massif, northeast Japan. *Saito Ho-on Kai Mus., Res. Bull.*, 33: 17-29.
- Murata, M., 1967. Some Permian Conulariidae from the Kitakami Massif, Northeast Japan. *Saito Ho-on Kai Museum Res. Bull.*, 36: 9-17.
- Murata, M., 1969. Molluscan fauna of the Toyoma formation (Late Permian). *Saito Ho-on*

- Kai Museum Research Bull.*, 38: 1-22.
- Murata, M., 1971. Fusulinid Biostratigraphy and Molluscan Fauna from the Uppermost part of the Sakamotozawa Formation, and the Pre-Kanokura Unconformity, in the southern part of the Kitakami Massif, Northeast Japan. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 82: 93-116.
- Murata, M. and Bando, Y., 1975. Discovery of Late Permian *Araxoceras* from the Toyoma formation in the Kitakami Massif, Northeast Japan. *Trans. Proc. Palaeont. Soc. Japan*, N. S., 97: 23-31.
- Nagao, T. und Minato, M., 1943. Über eine bedeutende Diskordanz in jüngeren Palaeozoikum des Kitakami-Gebirges im nördlichen Honshu, Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 7(1): 29-48.
- Nakamura, K., 1959. *Spinomarginifera* found in Japanese Permian. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 35: 143-146.
- Nakamura, K., 1960. *Dictyoclostus* derived from the Middle Permian Kanokura series and the Lower Permian Sakamotozawa series of the Kitakami mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 10(3): 495-511.
- Nakamura, K., 1970. *Isogramma* from the Permian Kanokura series of the Kitakami mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 14(3): 301-311.
- Nakamura, K., 1972. Permian Davidsoniacea from the Southern Kitakami mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 15(3-4) 361-425.
- Nakamura, K., 1972. *Anidanthus* and *Megousia* (Brachiopoda) from the Permian of Japan and Cambodia. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 15(3-4): 427-445.
- Nakamura, K., Kato, M. and Choi, D.R., 1970. On *Permudaria*, a new genus of the brachiopod family Linoproductidae. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4, 14(3): 293-299.
- Nakazawa, K., 1960. Two Permian nautiloids from Japan. *Japanese Jour. Geol. & Geogr.*, 31(3-4): 121-127.
- Nakazawa, K. and Newell, N.D., 1960. Permian Bivalves of Japan. *Mem. Fac. Sci., Kyoto Univ.*, ser. Geol. & Mineral., 35(1): 1-108.
- Nakazawa, K. et al., 1975. The Upper Permian and the Lower Triassic in Kashmir, India. *Mem. Fac. Sci., Kyoto Univ.*, Ser. Geol. & Mineral., 42(1): 1-106.
- Nishiyama, S., 1966. The echinoid fauna from Japan and adjacent regions. Part I. *Palaeont. Soc. Japan, Special papers*, 11:
- Nishiyama, S., 1968. The echinoid fauna from Japan and adjacent regions. Part II. *Palaeont. Soc. Japan, Special papers*, 13:
- Nogami, Y., 1961. Permische Fusuliniden aus dem Atetsu-Plateau Südwest japans. *Mem. Coll. Sci., Univ., Kyoto*, ser.B, 27(3): 159-249, 28(1): 159-243.
- Nogami, Y., 1962. Jungpaläozoikum im Atetsu-Plateau südwest japans. *Mem. Coll. Sci., Univ. Kyoto*, ser.B, 29(2): 161-176.
- Okimura, Y., Ishii, K., and Nakazawa, K., 1975. *Abadehella*, A New Genus of Tetrataxid Foraminifera from the Late Permian. *Mém. Fac. Sci. Kyoto Univ.*, Ser. Geol. & Mineral., 41(1): 35-48.
- Onuki, Y., 1956. *Geology of the Kitakami Massif*; One of the explanatory texts of the geological map (1/100,000) of Iwate Prefecture. Iwate Prefecture, 1-189.
- Onuki, Y., Murata, M., Bando, Y. and Mito, A., 1960. On the Permian System of the Maiya district in the Southern Kitakami Massif, Japan. *Geol. Soc. Japan. Jour.*, 66(782): 717-732.
- Ota, M., Toriyama, R., Sugimura, A. and Haikawa, T., 1973. Restudy on the geologic structure of the Akiyoshi Limestone Group, Southwest Japan. *Jour. Geogr.*, 82(3): 115-135.
- Ozaki, K., 1968. Problematical fossils from the Permian limestone of Akasaka, Gifu

- Prefecture. *Sci. Rep. Yokohama National Univ.*, ser.II, 14:
- Ozawa, T., 1970. Notes on the phylogeny and classification of the Superfamily Verbeekinoidea. *Mem. Fac. Sci., Kyushu Univ.*, ser.D, 20(1): 17-58.
- Ozawa, T., 1975. Evolution of *Lepidolina multiseptata* (Permian Foraminifer) in East Asia. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 23(2): 117-164.
- Ozawa, Y., 1925. Palaeontological and stratigraphical studies on the Permo- Carboniferous limestone of Nagato. Pt.2, Palaeontology. *Jour. Coll. Sci., Imp. Univ. Tokyo*, 45(6): 1-90.
- Ozawa, Y., 1927. Stratigraphical Studies of the Fusulina Limestone of Akasaka, Province of Mino. *Jour. Fac. Sci., Tokyo Imp. Univ.*, Sec.II, 2(3): 121-162.
- Rausser-Chernoussova, D.M., 1965. Foraminifers in the stratotypical section of th Sakmarian Stage (Sakmara river, Southern Urals). *Trans. Geol. Inst. Academy of Sciences, USSR*, 135: 1-78.
- Sada, K., 1965. Carboniferous and Permian Stratigraphy of the Atetsu Limestone in West Japan. *Jour. Sci., Hiroshima Univ.*, Ser.C, 5(1): 21-80.
- Sakagami, S., 1961. Japanese Permian Bryozoa. *Palaeont. Society of Japan, Special Papers*, 7: 1-58.
- Sakagami, S., 1964. Bryozoa of Akiyoshi. part 1. Permian bryozoa from the Shigeyasu quarry. *Bull. Akiyoshidai Sci., Mus.*, 3: 1-24.
- Saito, Y., 1966. Geology of the Setamai District. Southern Kitakami Massif, Northeast Japan. *Contr. Inst. Geol. & Palaeont., Tohoku University*, 62: 55-67.
- Saito, Y., 1968. Geology of the Younger Palaeozoic System of the southern Kitakami Massif, Iwate Prefecture, Japan. *Sci. Rep. Tohoku Univ.*, 2nd ser., 40: 79-139.
- Sato, K., 1969. Stratigraphy of the Permian System in the area of Toyoma Town, Southern Kitakami Massif, Japan. *Jour. Geol. Soc. Japan*, 75(11): 555-570.
- Schwager, C., 1883. Karbonische Foraminiferen aus China und Japan, in Richthofen's *China*. 4: 106-159.
- Sheng, J.C. and Lee, H.H., 1974. Recent Advances in the Permian Biostratigraphy of China. *Mem. Nanking Inst. Geol. & Palaeont.*, 5: 117-122.
- Shiida, I., 1940. On the geology of the vicinity of Kesennumacho, Miyagi Prefecture. *Contr. Inst. Geol. Palaeont., Tohoku Univ.*, 33: 1-72.
- Sugimura, A., 1972. Three new species of the genus *Hayasakapora* from the Akiyoshi limeston group. *Bull. Akiyoshidai Sci. Mus.*, 8: 1-6.
- Tazawa, J., 1973. Geology of the Kamiyasse area, southern Kitakami mountains. *Jour. Geol. Soc. Japan*, 79(10): 677-686.
- Tazawa, J., 1974. Two species of *Permundaria* from the Kitakami mountains, Northeast Japan. *Trans. Proc. Palaeont. Soc. Japan*, N.S., 94: 313-318.
- Tazawa, J., 1974. *Waagenoconcha* (Brachiopoda) from the Permian of the southern Kitakami mountains, Northeast Japan. *Jour. Fac. Sci., Hokkaido Univ.*, ser.4 16(2-3): 121-143.
- Tazawa, J., 1975. Uppermost Permian fossils from the Southern Kitakami mountains, Northeast Japan. *Jour. Geol. Soc. Japan*, 61(10) 629-640.
- Tazawa, J., 1976. The Permian of Kesennuma, Kitakami Mountain: A Preliminary Report. *Earth Science (Chikyu Kagaku)*, 30(3): 175-185.
- Toriyama, R., 1954. Geology of Akiyoshi, part 1. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 4(1): 29-97.
- Toriyama, R., 1954. Geology of Akiyoshi, part 2. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 5(1): 1-46.
- Toriyama, R., 1958. Geology of Akiyoshi, part 3. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D, 7: 1-264.
- Toriyama, R., 1967. The Fusulinacean Zones of Japan. *Mem. Fac. Sci., Kyushu Univ.*, Ser.D,

- 18(1): 35-260.
- Ueda, F., 1963. The geological structure of the Permian and the Triassic systems in the Toyoma and Maiya districts, southern Kitakami Massif, Northeast Japan. *Jour. Toyo Univ., General Education*, 4: Natural Science, 1-78.
- Wakimizu, T., 1902. On the limestone of Kinsyozan, Akasaka, Mino Province. *Jour. Geol. Soc. Tokyo*, 9(102):(104):(105):(108):
- Wilde, G.L., 1975. Fusulinid-Defined Permian Stages in Permian Exploration, Boundaries, and Stratigraphy. West Texas Geol. Soc. and Permian Basin Section, SEPM Publ., 75-76: 67-83.
- Working group on the Permian-Triassic System, 1975. Stratigraphy near the Permian-Triassic boundary in Japan and its correlation. *Jour. Geol. Soc. Japan*. 81(3): 165-184.
- Yabe, H., 1902: Materials for a knowledge of the Anthracolithic fauna. *Jour. Geol. Soc. Tokyo*, 9(110): 1.
- Yabe, H. & Hayasaka, I., 1915-16. Palaeozoic Corals from Japan, Korea and China. *Jour. Geol. Soc. Tokyo*, 22: 55-70, 79-92, 93-109, 127-142, 1915: 23: 57-75, 1916.
- Yabe, H. & Minato, M., 1944. *Wentzelloides maiyaensis* Yabe et Minato Gen. et sp. nov. aus dem Perm des Süd-Kitakami-Gebirges. *Japan. Jour. Geol. Geogr.*, 19(1-4): 141-142.
- Yabe, H. & Minato, M., 1944. Eine neue Art von *Wentzelella* aus dem japanische Perm. *Japan. Jour. Geol. Geogr.* 19(1-4): 139-140.
- Yabe, H. & Minato, M. (1944): Eine *Verbeekiella* Art aus dem Kitakami-Gebirges, Nordöstlichen Honsyu, Japan. *Proc. Imp. Acad. Tokyo*, 20: 159.
- Yabe, H. & Minato, M., 1945. A New Species of *Wentzelella* from the Permian Limestone near Iwaizaki, Kitakami District, Northeast Japan. *Proc. Japan Acad.*, 21(3-10): 469-472.
- Yabe, H. & Sugiyama, T., 1941. *Pseudoromingeria*, a new genus of Auloporid from Japan. *Proc. Imp. Acad. Tokyo*, 17: 379.
- Yamada, Y., 1959. On the unconformity of the Post-Nagaiwa Pre-Sakamotozawa epoch in the Hikoroichi district. *Jour. Geol. Soc. Japan*, 55(771): 713-724.

(Received on Sept. 22, 1977)