

1 ***Artemisia* pollen dataset for exploring the potential ecological
2 indicators in deep time**

3 Li-Li Lu^{a,d†}, Bo-Han Jiao^{a,d†}, Feng Qin^{b†}, Gan Xie^{a†}, Kai-Qing Lu^{a,d}, Jin-Feng Li^a, Bin Sun^a, Min Li^a,
4 David K. Ferguson^c, Tian-Gang Gao^{a,d *}, Yi-Feng Yao^{a,d*}, Yu-Fei Wang^{a,d*}

5 ^a*State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of
6 Sciences, 20 Nanxincun Xiangshan, Beijing 100093, China*

7 ^b*Key Laboratory of Land Surface Pattern and Simulation, Institute of Geographic Sciences and Natural
8 Resources Research, Chinese Academy of Sciences, Beijing 100101, China*

9 ^c*Department of Paleontology, University of Vienna, Althanstrasse 14, Vienna A-1090, Austria*

10 ^d*University of Chinese Academy of Sciences, Beijing 100049, China*

11 [†]These authors contributed equally to this work.

12 *Corresponding authors. Tel: +86 (10) 62836439.

13 Email addresses: gaotg@ibcas.ac.cn (T. G. GAO); yaoyf@ibcas.ac.cn (Y.
14 F. WANG).

15 **Abstract.** *Artemisia*, along with Chenopodiaceae is the dominant component growing in the desert and dry
16 grassland of the Northern Hemisphere. *Artemisia* pollen with its high productivity, wide distribution, and easy
17 identification, is usually regarded as an eco-indicator for assessing aridity and distinguishing grassland from
18 desert vegetation in terms of the pollen relative abundance ratio of Chenopodiaceae/*Artemisia* (C/A).
19 Nevertheless, divergent opinions on the degree of aridity evaluated by *Artemisia* pollen have been circulating
20 in the palynological community for a long time. To solve the confusion, we first selected 36 species from 9
21 clades and 3 outgroups of *Artemisia* based on the phylogenetic framework, which attempts to cover the
22 maximum range of pollen morphological variation. Then, sampling, experiments, photography, and
23 measurements were taken using standard methods. Here, we present pollen datasets containing 4018 original
24 pollen photographs, 9360 statistical pollen morphological traits, information on 30858 source plant
25 occurrences, and corresponding environmental factors. Hierarchical cluster analysis on pollen morphological
26 traits was carried out to subdivide *Artemisia* pollen into three types. When plotting the three pollen types of
27 *Artemisia* onto the global terrestrial biomes, different pollen types of *Artemisia* were found to have different
28 habitat ranges. These findings change the traditional concept of *Artemisia* being restricted to arid and
29 semi-arid environments. The data framework that we designed is open and expandable for new pollen data of
30 *Artemisia* worldwide. In the future, linking pollen morphology with habitat via these pollen datasets will
31 create additional knowledge that will increase the resolution of the ecological environment in the geological
32 past. The *Artemisia* pollen datasets are freely available at Zenodo (<https://doi.org/10.5281/zenodo.6791891>;
33 Lu et al., 2022).

34 **1 Introduction**

35 The concept of global change can be considered as any consistent trend in the environment - past, present, or
36 projected - that affects a substantial part of the globe. Consequently past climates shed light on our future
37 (Tierney et al., 2020). When attempting to reconstruct past global change prior to meteorological records, we
38 need some appropriate biological or abiotic proxies based on long-term, consistently collected data, e.g. leaf
39 wax biomarkers (Bhattacharya et al., 2018), tree-ring data (Moberg et al., 2005), leaf form (Yang et al., 2015),
40 pollen data (Mosbrugger et al., 2005; Guiot and Cramer, 2016; Marsicek et al., 2018), atmospheric carbon
41 dioxide (Zachos et al., 2008; Beerling and Royer, 2011), and isotope records (Zachos et al., 2001;
42 Sánchez-Murillo et al., 2019). Determining a suitable proxy to reconstruct palaeoclimate and
43 palaeoenvironment is a great scientific challenge (Tierney et al., 2020; McClelland et al., 2021).

44 The pollen of *Artemisia* (A), together with that of Chenopodiaceae (C) in arid and semi-arid areas, in the
45 form of the ratio of C/A pollen abundance, was applied to distinguish grassland and desert vegetation types
46 and assess the degree of drought in the geological past (El-Moslimany, 1990; Sun et al., 1994; Davies and Fall,
47 2001; Herzschuh et al., 2004; Xu et al., 2007; Zhao et al., 2009; Zhang et al., 2010; Zhao et al., 2012; Li et al.,
48 2017; Ma et al., 2017; Koutsodendris et al., 2019; Wang et al., 2020), because both Chenopodiaceae and
49 *Artemisia* are dominant elements of desert vegetation (China Vegetation Editorial Committee, 1980; Vrba,
50 1980; Tarasov et al., 1998; Herzschuh et al., 2004; Li et al., 2010; Zhao et al., 2021), and the sum of their
51 pollen relative abundances in the surface soil is usually more than 50% in arid and semi-arid areas (Sun et al.,
52 1994; Lu et al., 2020).

53 Among them, the pollen of *Artemisia*, with its high productivity, wide spatial and temporal distribution,
54 easy identification, and morphological uniformity under the light microscope (LM), is an essential component
55 and useful bio-indicator in pollen-based past vegetation reconstructions and environmental assessments. Some
56 researchers regarded *Artemisia* as an aridity indicator (El-Moslimany, 1990; Yi et al., 2003a; Yi et al., 2003b;
57 Liu et al., 2006; Cai et al., 2019; Cui et al., 2019; Chen et al., 2020; Wu et al., 2020; Cao et al., 2021), while
58 others suggested that the correlation between the relative abundance of *Artemisia* pollen and humidity was
59 insignificant (Weng et al., 1993; Sun et al., 1996; Koutsodendris et al., 2019; Lu et al., 2020; Zhao et al.,
60 2021). Consequently, there is an urgent need to evaluate whether different pollen types of *Artemisia* represent
61 distinct habitats.

62 In the past, *Artemisia* pollen was regarded as very uniform under LM (Wodehouse, 1926; Sing and Joshi,
63 1969; Ling, 1982; Wang et al., 1995). For instance, following the description and statistics of pollen
64 morphology of 27 species of *Artemisia* in Eurasia under LM, Sing and Joshi (1969) stated that the pollen
65 grains of *Artemisia* are consistent and continuous in morphology. Later, some authors recognized a series of
66 pollen types (Chen, 1987; Jiang et al., 2005; Ghahraman et al., 2007; Shan et al., 2007; Hayat et al., 2009;
67 Hayat et al., 2010; Hussain et al., 2019), based on a detailed survey of the pollen micromorphology of
68 different taxa under the scanning electron microscope (SEM).

69 For example, Chen (1987) described the pollen morphology of 77 *Artemisia* species from China under
70 LM and SEM and divided these pollen grains into six types by using pollen characters, such as the shape and
71 size of the spinules as well as the density of spinules and granules. Type I (sparse spinules with granules
72 among them), type II (dense spinules, no or few granules), type III (sparse spinules, no granules), type IV
73 (dense spinules, well-developed granules), type V (small and sparse spinules, smooth tectum) and type VI
74 (dissimilar spinules with granules among them).

75 Shan et al. (2007) investigated the pollen morphology of 32 *Artemisia* species from the Loess Plateau of
76 China under LM and SEM and divided these pollen grains into five types according to exine sculpture: type I
77 (dense spinules with swollen bases, small granules), type II (dense spinules, swollen bases almost united),
78 type III (dense spinules with swollen bases and smooth tectum), type IV (sparse small spinules and smooth
79 tectum) and type V (sparse spinules, small granules).

80 Jiang et al. (2005) observed the pollen morphology of 57 representative plants in 7 groups of *Artemisia*
81 under LM and SEM. This pollen can be divided into two types based on exine sculpture: type I (spinules
82 multi-ruminated with flared bases, connecting the mostly densely arranged spinules) and type II (densely or
83 loosely arranged spinules without flared bases, interspace glandular or smooth) with subtypes II-1, II-2, II-3,
84 and II-4 based on the distribution of the spinules.

85 Ghahraman et al. (2007) studied the pollen morphology of 26 species of the 33 *Artemisia* species in Iran
86 under LM and SEM. Based on exine ornamentation observed under SEM, two types of pollen grains were
87 recognized: type I, exine surface covered with dense acute spinules, Type II, exine surface with few spinules.

88 Hayat et al. (2009, 2010) carried out a palynological study of 22 *Artemisia* species from Pakistan under
89 LM and SEM. Earlier work demonstrated the phylogenetic associations within *Artemisia* based on a
90 phylogenetic analysis of 9 characters (pollen type, pollen shape, spinule arrangement, exine sculpture, spinule
91 base, the length of polar axis, the length of equatorial axis, exine thickness, and colpus width) of pollen grains

92 of *Artemisia*. In the latter work, eight micromorphological characters were identified and pooled by cluster
93 analysis, leading to the recognition of 5 groups.

94 Hussain et al. (2019) studied the pollen morphology of 15 *Artemisia* species in the Gilgit-Baltistan region
95 of Pakistan utilizing SEM and divided these species into four groups based on cluster analysis of seven
96 micromorphological characters (pollen type, pollen shape, spinule arrangement, exine sculpture, spinule base,
97 polar length, and equatorial width).

98 Almost all of the above-mentioned *Artemisia* pollen classifications were designed to solve taxonomic or
99 phylogenetic problems, and only a few were concerned with linking diverse habitats to the different pollen
100 types in *Artemisia*.

101 Here we attempt to 1) present abundant pollen photographs of 36 species from 9 branches and 3
102 outgroups of the genus (ca. 400 species worldwide, see Ling, 1982; Bremer and Humphries, 1993),
103 constrained by the phylogenetic framework of *Artemisia* (Sanz et al., 2008; Malik et al., 2017); 2) describe
104 and measure the morphological traits of these pollen grains; 3) provide a new classification of pollen types
105 and their distribution worldwide, with a key to pollen types in *Artemisia*; 4) explore the diverse ecological
106 niches of *Artemisia* represented by different pollen types in order to evaluate palaeovegetation and reconstruct
107 palaeoenvironments.

108 **2 Materials and methods**

109 **2.1 Sampling strategy**

110 The 36 pollen samples studied were selected from voucher sheets in the PE herbarium at the Institute of
111 Botany, Chinese Academy of Sciences (Fig. 1, Table B1), covering 9 main clades, i.e., Subg. *Tridentata*, Subg.
112 *Artemisia* (contains Sect. *Artemisia*, Sect. *Abrotanum* I, Sect. *Abrotanum* II and Sect. *Abrotanum* III), Subg.
113 *Pacifica*, Subg. *Seriphidium*, Subg. *Absinthium*, and Subg. *Dracunculus*, constrained by the phylogenetic
114 framework of *Artemisia* (Malik et al., 2017) and 3 outgroups (Sanz et al., 2008), reflecting the maximum
115 diversity or morphological variation under LM and SEM.





Figure 1. Phylogenetic tree of *Artemisia* (modified from Malik et al., 2017). The styles of the strokes that were used to draw the branches indicate the traditional subgeneric classification of *Artemisia*, and the yellow spots indicate sampled taxa.

120 2.2 Data acquisition

121 Pollen samples were acetolyzed by the standard method (Erdtman, 1960) and fixed in glycerine jelly. Standard
122 procedures were followed for LM and SEM (Chen, 1987; Wang et al., 1995). The pollen grains were
123 photographed under LM (Leica DM 4000) at a magnification of $\times 1000$ and SEM (Hitachi S-4800) at an
124 accelerating voltage of 30 kV. The pollen terminology followed the descriptions of Hesse et al. (2009) and

125 Halbritter et al. (2018). The statistical pollen morphological traits under LM (Figs. 2a-b, P: Polar length; E:
 126 Equatorial width; P/E; T: Exine thickness; L: Pollen length; T/L) of each species were measured using 20
 127 pollen grains. We chose five pollen grains under SEM for each exine ornamentation trait in each species (Figs.
 128 2c-f, D: Diameter of spinule base; H: Spinule height; D/H; Gs: Granule spacing; Ss: Spinule spacing; Gs/Ss;
 129 Ps: Perforation spacing), and on average, randomly selected four regions of each pollen grain for measuring,
 130 yielding a total of 20 measurements. The mean value (M) and standard deviation (SD) of the pollen grains of
 131 each species were measured and calculated in both polar and equatorial views (Appendix A, Table 1).



133 **Figure 2.** Graphical illustration of measured pollen morphological traits in *Artemisia* (a-b: *A. annua*; c-d: *A.
134 vulgaris*) and outgroups (e: *Kaschagaria brachanthemoides*; f: *Ajania pallasiana*). Scale bar in LM and SEM
135 overview 10 µm, in SEM close-up 1 µm.

136 The scientific names of selected taxa were standardized according to Plants of the World Online
137 (<https://powo.science.kew.org/>). The specimen sampling coordinates of the corresponding taxa were obtained
138 from the Global Biodiversity Information Facility (GBIF, <https://www.gbif.org/>). Only preserved specimens
139 were filtered for GBIF data given their well-documented geographical information and the availability of
140 specimens as definitive vouchers. The distribution data on observations and cultivated collections provided by
141 GBIF were excluded because they may contain incorrect identification or incorrect geo-referencing (Brummitt
142 et al., 2020). Next, the distribution data was standardized cleaned using R package "CoordinateCleaner"
143 (Zizka et al., 2019); no outliers were found.

144 The corresponding environmental factors including altitude and 19 climate parameters of these
145 coordinates were obtained from WorldClim (<https://www.worldclim.org/>) with a spatial resolution of 30
146 seconds (~1 km²) in 1970-2000 by Extract MultiValues To Points using ArcGIS 10.2 software in bilinear
147 interpolation.


148 **2.3 Data processing**

149 OriginPro 2021 software was used for hierarchical cluster analysis on *Artemisia* and its outgroup pollen data.
150 The Euclidean distance was calculated after the normalization of the original data, and the Ward method was
151 used for clustering. Five groups were established, and the center point of each group was calculated according
152 to the sum of distances. Pollen morphological traits for the principal component analysis (PCA) of *Artemisia*
153 and its outgroups and grouped according to the five groups of the cluster analysis. OriginPro 2021 software
154 was used to draw group violin diagrams and run an ANOVA to test for an overall difference between the
155 pollen characters of 3 pollen types, followed by post hoc tests (Tukey). OriginPro 2021 software was also
156 used to run correlation coefficients analysed by the Pearson correlation between pollen morphological traits
157 and environmental factors as well as draw group violin diagrams and run a KWANOVA to test for overall
158 differences between the environmental factors of the 3 pollen types. The images of habitats reproduced in the
159 text are from the websites listed in Table B1.

160 The global distribution data of the 36 representative species and 3 pollen types were plotted on the map
161 of terrestrial ecological regions (Olson et al., 2001) using ArcGIS 10.2 software (Figs. 16, 20).

162 **3 Data description**163 **3.1 *Artemisia* pollen grains and their source plant habitats**


164 Here we provide detailed data on pollen morphological traits, covering 36 species from 9 main clades of
 165 *Artemisia* and 3 outgroups constrained by the phylogenetic framework (Fig. 1, Sanz et al., 2008; Malik et al.,
 166 2017) under LM and SEM, the habitats of their source plants (Figs. 3-14).



167
 168 **Figure 3.** Pollen grains and the habitats of their source plants.
 169 a. *Artemisia cana*; b. *Artemisia tridentata*; c. *Artemisia californica*.

170 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
171 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
172 <https://www.inaturalist.org/photos/54492753> by © Jason Headley, b7 cited from
173 <https://www.inaturalist.org/photos/117436654> by © Matt Berger, c7 cited from
174 <https://www.inaturalist.org/photos/108921528> by © Don Rideout).

175 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




176 **Figure 4.** Pollen grains and the habitats of their source plants.


177 a. *Artemisia indica*; b. *Artemisia argyi*; c. *Artemisia mongolica*.

178 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
179 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
180 <https://www.inaturalist.org/photos/66336449> by © yangting, b7 cited from
181 <https://www.inaturalist.org/photos/66336449> by © yangting, c7 cited from

182 <https://www.inaturalist.org/photos/95820686> by © sergeyprokopenko, c7 cited from
183 <https://www.inaturalist.org/photos/163584035> by © Nikolay V Dorofeev).
184 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.



185
186 **Figure 5.** Pollen grains and the habitats of their source plants.
187 a. *Artemisia vulgaris*; b. *Artemisia selengensis*; c. *Artemisia ludoviciana*.
188 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
189 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
190 <https://www.inaturalist.org/photos/120600448> by © Sara Rall, b7 cited from
191 <https://www.inaturalist.org/photos/46352423> by © Gularjanz Grigoryi Mihajlovich, c7 cited from
192 <https://www.inaturalist.org/photos/77690333> by © Ethan Rose).
193 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




194

195 **Figure 6.** Pollen grains and the habitats of their source plants.

196 a. *Artemisia roxburghiana*; b. *Artemisia rutifolia*; c. *Artemisia chinensis*.

197 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
 198 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 provided
 199 by © Bo-Han Jiao, b7 cited from <https://www.inaturalist.org/photos/62207191> by © Daba, c7 provided by ©
 200 Jia-Hao Shen).


201 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.



202

203 **Figure 7.** Pollen grains and the habitats of their source plants.204 a. *Artemisia kurramensis*; b. *Artemisia compactum*; c. *Artemisia maritima*.205 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
206 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
207 <https://www.inaturalist.org/photos/133758174> by © Andrey Vlasenko, b7 provided by © Chen Chen, c7 cited from
208 <https://www.inaturalist.org/photos/86515371> by © torkild).

209 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




210

211 **Figure 8.** Pollen grains and the habitats of their source plants.

212 a. *Artemisia aralensis*; b. *Artemisia annua*; c. *Artemisia freyniana*.

213 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
 214 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
 215 <https://www.plantarium.ru/lang/en/page/image/id/73063.html> by © Полынь аральская, b7 provided by ©
 216 Chen Chen, c7 cited from <https://www.inaturalist.org/photos/154390279> by © Шильников Дмитрий
 217 Сергеевич).

218 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




219

220 **Figure 9.** Pollen grains and the habitats of their source plants.

221 a. *Artemisia stechmanniana*; b. *Artemisia pontica*; c. *Artemisia frigida*.

222 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
 223 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 provided
 224 by © Bo-Han Jiao, b7 cited from <https://www.inaturalist.org/photos/93438780> by © Martin Pražák, c7 cited
 225 from <https://www.inaturalist.org/photos/125022240> by © Suzanne Dingwell).


226 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.



227

228 **Figure 10.** Pollen grains and the habitats of their source plants.229 a. *Artemisia rupestris*; b. *Artemisia sericea*; c. *Artemisia absinthium*.230 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
231 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 provided
232 by © Bo-Han Jiao, b7 cited from <https://www.inaturalist.org/photos/48033353> by © svetlana_katana, c7 cited
233 from <https://www.inaturalist.org/photos/123569286> by © Станислав Лебедев).

234 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




235

236 **Figure 11.** Pollen grains and the habitats of their source plants.

237 a. *Artemisia abrotanum*; b. *Artemisia blepharolepis*; c. *Artemisia norvegica*.

238 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
 239 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
 240 <https://www.inaturalist.org/photos/116106722> by © Андрей Москвичев, b7 provided by © Ji-Ye Zheng, c7
 241 cited from <https://www.inaturalist.org/photos/161393521> by © Erin Springinotic).

242 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




243

244 **Figure 12.** Pollen grains and the habitats of their source plants.

245 a. *Artemisia tanacetifolia*; b. *Artemisia tournefortiana*; c. *Artemisia dracunculus*.

246 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
247 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
248 <https://www.inaturalist.org/photos/78902853> by © Alexander Dubynin, b7 provided by © Chen Chen, c7 cited
249 from <https://www.inaturalist.org/photos/76312868> by © anatolymikhaltsov).

250 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.




251

252 **Figure 13.** Pollen grains and the habitats of their source plants.253 a. *Artemisia japonica*; b. *Artemisia capillaris*; c. *Artemisia campestris*.

254 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
 255 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 cited from
 256 <https://www.inaturalist.org/photos/44507659> by © 陳 達 智 , b7 cited from
 257 <https://www.inaturalist.org/photos/60639286> by © Cheng-Tao Lin, c7 cited from
 258 <https://www.inaturalist.org/photos/113822257> by © pedrosanz-anapri).

259 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.



260

261 **Figure 14.** Pollen grains and the habitats of their source plants.

262 a. *Kaschgaria brachanthemoides*; b. *Ajania pallasiana*; c. *Chrysanthemum indicum*.

263 Pollen grains in equatorial view under LM (a1, b1, c1) and SEM (a3, a5, b3, b5, c3, c5), in polar view under
264 LM (a2, b2, c2) and SEM (a4, a6, b4, b6, c4, c6), along with the habitats of their source plants (a7 provided
265 by © Chen Chen, b7 cited from <https://www.inaturalist.org/photos/162408714> by © Игорь Пospelов, c7
266 provided by © Bo-Han Jiao).

267 Scale bar in LM and SEM overview 10 µm, in SEM close-up 1 µm.


268 **3.2 Statistical pollen morphological trait data of 36 sampled taxa**

269 The mean values of 10 pollen morphological traits of 36 sampled species are listed in Table 1, and these data
 270 distribution patterns are shown in boxplots (Fig. 15) in the form of variation (25%-75%), and further
 271 described in the form of mean value \pm standard deviation ($M \pm SD$, Appendix A).

272 **Table 1.** Pollen morphological traits of 36 selected species (P: Polar length; E: Equatorial width; T: Exine
 273 thickness; L: Pollen length; D: Diameter of spinule base; H: Spinule height; Gs: Granule spacing; Ss: Spinule
 274 spacing; Ps: Perforation spacing).

No.	Species	P (μm)	E (μm)	P/E	T (μm)	L (μm)	T/L	D (μm)	H (μm)	D/H	Gs (μm)	Ss (μm)	Gs/Ss	Ps (μm)
1	<i>Artemisia cana</i>	23.46	24.5	0.96	3.91	24.58	0.16	0.58	0.46	1.28	0.33	1.60	0.21	0
2	<i>Artemisia tridentata</i>	21.36	20.69	1.04	3.55	22.35	0.16	0.76	0.60	1.30	0.24	1.12	0.22	0
3	<i>Artemisia californica</i>	18.94	19.13	0.99	2.70	18.85	0.14	0.75	0.71	1.08	0.24	1.45	0.17	0
4	<i>Artemisia indica</i>	23.47	23.81	0.99	3.50	23.31	0.15	0.76	0.39	2.04	0.28	1.21	0.24	0
5	<i>Artemisia argyi</i>	21.8	21.67	1.01	3.55	22.24	0.16	0.64	0.38	1.71	0.22	0.90	0.26	0
6	<i>Artemisia mongolica</i>	21.05	20.42	1.03	3.29	19.78	0.17	0.62	0.41	1.54	0.19	0.91	0.22	0
7	<i>Artemisia vulgaris</i>	19.72	19.29	1.03	2.92	18.94	0.16	0.69	0.34	2.13	0.29	1.55	0.20	0
8	<i>Artemisia selengensis</i>	20.67	19.68	1.06	3.72	20.8	0.18	0.67	0.38	1.76	0.22	1.05	0.22	0
9	<i>Artemisia ludoviciana</i>	21.65	20.82	1.04	3.71	20.94	0.18	0.70	0.37	1.94	0.2	1.23	0.16	0
10	<i>Artemisia roxburghiana</i>	23.88	23.69	1.01	3.78	21.81	0.17	0.76	0.39	1.96	0.28	0.79	0.36	0
11	<i>Artemisia rutifolia</i>	22.22	22.7	0.98	3.53	24.93	0.14	0.31	0.26	1.2	0.21	1.27	0.17	0
12	<i>Artemisia chinensis</i>	21.53	22.75	0.95	2.97	23.71	0.13	0.70	0.55	1.29	0.27	0.91	0.31	0
13	<i>Artemisia kurramensis</i>	19.71	19.35	1.02	3.30	19.44	0.17	0.38	0.27	1.41	0.23	1.25	0.19	0
14	<i>Artemisia compactum</i>	22.33	21.97	1.02	2.97	21.67	0.14	0.41	0.28	1.50	0.51	0.92	0.56	0
15	<i>Artemisia maritima</i>	26.24	23.09	1.14	3.54	24.42	0.14	0.28	0.23	1.30	0.53	1.08	0.50	0
16	<i>Artemisia aralensis</i>	22.32	21.91	1.02	3.16	22.76	0.14	0.25	0.22	1.16	0.50	1.09	0.46	0

17	<i>Artemisia annua</i>	19.71	19.45	1.02	3.45	19.2	0.18	0.45	0.39	1.18	0.27	1.29	0.21	0
18	<i>Artemisia freyniana</i>	23.39	21.3	1.10	3.17	21.29	0.15	0.56	0.40	1.40	0.2	1.15	0.18	0
19	<i>Artemisia stechmanniana</i>	26.31	25.16	1.05	3.97	23.45	0.17	0.37	0.35	1.07	0.19	1.40	0.14	0
20	<i>Artemisia pontica</i>	20.64	19.62	1.05	3.01	19.75	0.15	0.6	0.37	1.63	0.17	1.32	0.13	0
21	<i>Artemisia frigida</i>	25.11	24.9	1.01	4.61	24.83	0.19	0.46	0.32	1.44	0.31	1.3	0.24	0
22	<i>Artemisia rupestris</i>	24.45	22.92	1.07	3.18	21.96	0.14	0.55	0.33	1.68	0.25	0.91	0.28	0
23	<i>Artemisia sericea</i>	26.31	27.9	0.94	3.75	26.89	0.14	0.89	0.54	1.71	0.28	1.74	0.16	0
24	<i>Artemisia absinthium</i>	22.79	20.84	1.09	3.39	19.92	0.17	0.59	0.40	1.52	0.18	1.11	0.16	0
25	<i>Artemisia abrotanum</i>	24.47	23.73	1.03	3.15	18.82	0.17	0.72	0.51	1.44	0.22	1.41	0.16	0
26	<i>Artemisia blepharolepis</i>	18.96	19.26	0.99	3.15	18.82	0.17	0.69	0.44	1.64	0.37	1.68	0.23	0
27	<i>Artemisia norvegica</i>	24.51	22.11	1.11	3.48	22.61	0.15	0.67	0.43	1.66	0.19	1.56	0.12	0
28	<i>Artemisia tanacetifolia</i>	28.38	27.75	1.03	3.46	27.63	0.13	0.71	0.32	2.23	0.30	1.08	0.29	0
29	<i>Artemisia tournefortiana</i>	20.76	20.43	1.02	3.33	20.03	0.17	0.73	0.42	1.81	0.26	1.25	0.22	0
30	<i>Artemisia dracunculus</i>	22.89	22.87	1.00	2.82	21.91	0.13	0.68	0.45	1.56	0.31	0.92	0.34	0
31	<i>Artemisia japonica</i>	20.18	21.23	0.95	4.24	21.02	0.2	0.57	0.32	1.8	0.26	1.26	0.21	0
32	<i>Artemisia capillaris</i>	19.53	19.64	1.00	3.54	19.18	0.18	0.51	0.36	1.44	0.26	1.27	0.21	0
33	<i>Artemisia campestris</i>	21.69	21.26	1.02	3.68	21.21	0.17	0.57	0.38	1.53	0.41	1.23	0.34	0
34	<i>Kaschagaria brachanthemoides</i>	23.26	22.09	1.06	3.93	21.01	0.19	0.55	0.44	1.25	0	1.75	0	0.47
35	<i>Ajania pallasiana</i>	35.16	35.92	0.98	10.23	38.31	0.27	4.41	3.47	1.29	0	7.84	0	0.39
36	<i>Chrysanthemum indicum</i>	33.54	34.42	0.98	8.65	34.82	0.25	2.94	3.59	0.82	0	7.11	0	0.37



275

276


Figure 15. Boxplot of 36 sampled taxa, showing the variation in pollen morphological traits.

277 1. *Artemisia cana*; 2. *Artemisia tridentata*; 3. *Artemisia californica*; 4. *Artemisia indica*; 5. *Artemisia argyi*; 6.
 278 *Artemisia mongolica*; 7. *Artemisia vulgaris*; 8. *Artemisia selengensis*; 9. *Artemisia ludoviciana*; 10. *Artemisia*
 279 *roxburghiana*; 11. *Artemisia rutifolia*; 12. *Artemisia chinensis*; 13. *Artemisia kurramensis*; 14. *Artemisia*
 280 *compactum*; 15. *Artemisia maritima*; 16. *Artemisia aralensis*; 17. *Artemisia annua*; 18. *Artemisia freyniana*;
 281 19. *Artemisia stachmanniana*; 20. *Artemisia pontica*; 21. *Artemisia frigida*; 22. *Artemisia rupestris*; 23.
 282 *Artemisia sericea*; 24. *Artemisia absinthium*; 25. *Artemisia abrotanum*; 26. *Artemisia blepharolepis*; 27.
 283 *Artemisia norvegica*; 28. *Artemisia tanacetifolia*; 29. *Artemisia tournefortiana*; 30. *Artemisia dracunculus*;
 284 31. *Artemisia japonica*; 32. *Artemisia capillaris*; 33. *Artemisia campestris*; 34. *Kaschagaria brachanthemoides*;
 285 35. *Ajania pallasiana*; 36. *Chrysanthemum indicum*.

286

3.3 The source plant occurrences

287 The source plant distributions in global terrestrial biomes of 36 sampled species are shown in Fig. 16. In
 288 *Artemisia*, some species have worldwide distributions, such as *A. vulgaris* (Fig. 16-7), *A. absinthium* (Fig.
 289 16-24), and *A. campestris* (Fig. 16-33); a few taxa are limited to East Asia, such as *A. roxburghiana* (Fig.
 290 16-10) and *A. blepharolepis* (Fig. 16-26), while others have narrow and isolated distributions in deserts and
 291 xeric shrublands of Central Asia, e.g. *A. kurramensis* (Fig. 16-13) and *A. aralensis* (Fig. 16-16). In outgroups
 292 of *Artemisia*, *Kaschagaria brachanthemoides* is also confined to deserts and xeric shrublands of Central Asia
 293 (Fig. 16-34), while *Ajania pallasiana* lives in forests of East Asia (Fig. 16-35).



294

295 **Figure 16.** The global distribution maps of 36 sampled taxa in terrestrial biomes (modified from Olson et al.,
296 2001).

- 297 1. *Artemisia cana*; 2. *Artemisia tridentata*; 3. *Artemisia californica*; 4. *Artemisia indica*; 5. *Artemisia argyi*; 6.
298 *Artemisia mongolica*; 7. *Artemisia vulgaris*; 8. *Artemisia selengensis*; 9. *Artemisia ludoviciana*; 10. *Artemisia
299 roxburghiana*; 11. *Artemisia rutifolia*; 12. *Artemisia chinensis*; 13. *Artemisia kurramensis*; 14. *Artemisia
300 compactum*; 15. *Artemisia maritima*; 16. *Artemisia aralensis*; 17. *Artemisia annua*; 18. *Artemisia freyniana*;
301 19. *Artemisia stachmanniana*; 20. *Artemisia pontica*; 21. *Artemisia frigida*; 22. *Artemisia rupestris*; 23.
302 *Artemisia sericea*; 24. *Artemisia absinthium*; 25. *Artemisia abrotanum*; 26. *Artemisia blepharolepis*; 27.
303 *Artemisia norvegica*; 28. *Artemisia tanacetifolia*; 29. *Artemisia tournefortiana*; 30. *Artemisia dracunculus*; 31.
304 *Artemisia japonica*; 32. *Artemisia capillaris*; 33. *Artemisia campestris*; 34. *Kaschagaria brachanthemoides*;
305 35. *Ajania pallasiana*; 36. *Chrysanthemum indicum*.

306 **4 Potential use of the *Artemisia* pollen datasets**

307 **4.1 The pollen classification of *Artemisia***

308 The pollen grains of Anthemideae and Asteraceae under LM could be simply divided into *Artemisia* pollen
309 type (Figs. 3-13, 14a, Appendix A) with indistinct and short spinules and *Anthemis* pollen type such as
310 *Chrysanthemum indicum* and *Ajania pallasiana* (Figs. 14b-c, Appendix A) with distinct and long spines on
311 pollen exine ornamentation (Wodehouse, 1926; Stix, 1960; Chen, 1987; Chen and Zhang, 1991; Martín et al.,
312 2001; Martín et al., 2003; Sanz et al., 2008; Blackmore et al., 2009; Vallès et al., 2011). *Artemisia* pollen
313 grains are difficult to separate from those of other related genera with *Artemisia* pollen type such as
314 *Kaschgaria brachanthemoides* (Figs. 14a1-2, Appendix A), *Elachanthemum*, *Ajanopsis*, *Filifolium*, and
315 *Neopallasia* (Chen and Zhang, 1991) under LM due to their great similarity in pollen exine ornamentation and
316 colporate patterns (Chen, 1987; Martín et al., 2001; Martín et al., 2003; Vallès et al., 2011). Furthermore, Sing
317 and Joshi (1969) questioned the feasibility of recognizing pollen types under LM in the highly uniform pollen
318 of *Artemisia*. Later, SEM made it possible to subdivide the pollen of *Artemisia* and those of other related
319 genera within the *Artemisia* pollen type using pollen exine ultrastructure characters (Chen, 1987; Chen and
320 Zhang, 1991; Sun and Xu, 1997; Jiang et al., 2005; Ghahraman et al., 2007; Shan et al., 2007; Hayat et al.,
321 2009; Hayat et al., 2010; Hussain et al., 2019).

322 Hierarchical cluster analysis (Fig. 17a) revealed that the pollen morphological traits (P/E, H, D, D/H, Ss,
323 Gs, Gs/Ss, and Ps) of *Artemisia* and its outgroups were divided into Clade A with perforations and without
324 granules (Figs. 13a5-6, b5-6, c5-6) and Clade B with granules and without perforations (Figs. 3-13a5-6, b5-6,
325 c5-6) on the pollen exine under SEM.





Figure 17. Hierarchical cluster analysis, showing the dendrogram for pollen types from *Artemisia* and outgroups (a) and the habitat ranges of 36 representative species (b, Tutin et al., 1976; Zhang, 2007; Ling et al., 2011).

In addition, Clade A, as the outgroup of *Artemisia*, includes *Anthemis* type (*Chrysanthemum indicum* and *Ajania pallasiana*) with prominent spines on pollen exine under LM, and *Kaschgaria* type (*Kaschgaria brachanthemoides*) with spinules on pollen exine (Figs. 14a, 17a). Clade B comprises three pollen types from three branches of *Artemisia* (Fig. 17a), i.e., SG type (short and wide spinule pollen type, Clade B1), LNS type (long and narrow spinule pollen type, Clade B2-1), and SG type (sparse granule pollen type, Clade B2-2).


Eight pollen morphological traits (P/E, H, D, D/H, Ss, Gs, Gs/Ss, and Ps) were selected for the principal component analysis (PCA) of 36 taxa of *Artemisia* and its outgroups (Fig. 18) and grouped according to the five clades of the cluster analysis, i.e. the five pollen types (Fig. 17a). The results reveal that *Artemisia* pollen morphology differs significantly from that of the outgroups, and that three *Artemisia* pollen types could be distinguished.



340
341 Figure 18. Principal component analysis of 36 taxa of *Artemisia* and its outgroups.

342 Nine characteristics of *Artemisia* pollen could partially explain the differences between these 3 pollen
343 types (Fig. 19). P/E (the length of polar axis/the length of equatorial axis) in LNS types (0.93-1.06) are
344 significantly different (ANOVA $P < 0.001$) from both SWS (0.97-1.12) and SG (0.98-1.14), so could be used
345 to identify the LNS type. D/H (diameter of spinule base/spinule height) in the SWS type differ significantly
346 (ANOVA $P < 0.001$) from both LNS and SG types. The variation range of D/H is 1.38-2.23 in the SWS type,
347 1.07-1.75 in the LNS type, and 0.98-1.66 in the SG type, indicating that the SWS pollen type is distinguished
348 by short and wide spinules. Gs/Ss (granule spacing/spinule spacing) in the SG type was higher than those of
349 the SWS and LNS types (ANOVA $P < 0.001$), which distinguished the SG type from the other two types.
350 Moreover, the SG type is characterized by sparse granules with the variation range of Gs/Ss spanning
351 0.37-0.64, while the SWS and LNS types show much denser granules whose Gs/Ss are mainly below 0.35.

352 Within the new *Artemisia* pollen classification (Fig. 17a, Key), the SWS type represents a type of pollen
353 with short and wide spinules ($D/H > 1.81$) and dense granules (Figs. 17a, 19). The LNS type represents a type
354 of pollen with long and narrow spinules ($D/H < 1.38$) and dense granules (Figs. 17a, 19). The SG type is
355 characterized by sparse granules ($Gs/Ss > 0.37$) and small, long, and narrow spinules (Figs. 17a, 19).



356
357
358
359
360

Figure 19. Violin diagrams of three pollen types from *Artemisia*, showing the variations ($M \pm SD$) in nine pollen characters (P: length of polar axis; E: length of equatorial axis; D: diameter of spinule base; H: spinule height; Gs: granule spacing; Ss: spinule spacing; Ps: perforation spacing). Asterisks indicate statistically significant differences ($p < 0.001$).

361
362
363
364
365
366
367
368
369

Key to 3 pollen types of *Artemisia* and 3 outgroups

1. Pollen exine with perforations and without granules under SEM 2
1. Pollen exine with granules and without perforations under SEM 3
2. Distinct and long spines on pollen exine, with $H > 3 \mu\text{m}$ *Anthemis* type
2. Indistinct and short spinules on pollen exine, with $H < 1 \mu\text{m}$ *Kaschgaria* type
3. Pollen exine with sparse granules and $Gs/Ss \geq 0.37$ under SEM SG type
3. Pollen exine with dense granules and $Gs/Ss \leq 0.31$ under SEM 4
4. Pollen exine with $D/H < 1.38$ under SEM LNS type
4. Pollen exine with $D/H \geq 1.38$ under SEM SWS type


370

4.2 The ecological implications of *Artemisia* pollen types

371
372


Plotting the distribution data of 33 species from 9 main branches of *Artemisia* constrained by the phylogenetic framework (Fig. 1) onto the global terrestrial biomes (Fig. 20), we noticed that the genus is widely distributed

373 from forest to grassland, desert, and saline habitats (Figs. 16, 17b, 20). Furthermore, different species of
 374 *Artemisia* with SWS pollen type (Fig. 20a) and LNS type (Fig. 20b) have a rather wide distribution with
 375 severely overlapping ranges while those with SG type (Fig. 20c) have narrow and isolated distributions.



376
 377 **Figure 20.** The global distribution pattern of 3 *Artemisia* pollen types in terrestrial biomes (modified from
 378 Olson et al., 2001). a. SG type; b. LNS type; c. SWS type.
 379 14 terrestrial biomes: 01. Tropical and Subtropical Moist Broadleaf Forests; 02. Tropical and Subtropical Dry
 380 Broadleaf Forests; 03. Tropical and Subtropical Coniferous Forests; 04. Temperate Broadleaf and Mixed
 381 Forests; 05. Temperate Coniferous Forests; 06. Boreal Forests/Taiga; 07. Flooded Grasslands and Savannas;
 382 08. Montane Grasslands and Shrublands; 09. Tundra; 10: Mediterranean Forests, Woodlands, and Shrub;
 383 11. Tropical and Subtropical Grasslands, Savannas, and Shrublands; 12. Temperate Grasslands, Savannas, and
 384 Shrublands; 13. Deserts and Xeric Shrublands; 14. Mangroves; 15. Lakes; 16. Rock and Ice.

385 The ecological implications of *Artemisia* pollen types mentioned above fall into four categories. (i)
 386 *Artemisia* with the SG pollen type all belong to the subg. *Seriphidium*, which generally grows in dry habitats
 387 ranging from grassland desert to desert and coastal saline-alkaline environments, with their distribution
 388 largely limited to Eurasia and growing at low altitude (Figs. 17b, 20c, 21). (ii) The habitats of *Artemisia* with
 389 LNS pollen type have a global distribution and occur in forest, grassland and desert, and even coastal areas
 390 (Figs. 17b, 20b, 21), with the highest mean annual temperature (MAT). Hence, the LNS pollen type is a
 391 generalist. (iii) *Artemisia* with SWS pollen type include Sect. *Artemisia* and its habitats range from forest to
 392 desert, although most of the taxa are confined to humid environments from forest to grassland with a global
 393 distribution and the highest mean annual precipitation (MAP, Figs. 17b, 20c, 21). (iv) If the SWS pollen type
 394 and the SG pollen type appear together, the range of vegetation types could be reduced to grassland desert and
 395 desert through niche coexistence (Fig. 17b).



396
 397 **Figure 21.** Violin diagrams of three pollen types from *Artemisia*, showing the variations (25%-75%) in MAT,
 398 MAP, and altitude. Asterisks indicate statistically significant differences ($p < 0.001$).

399 In addition, we noticed that *Kaschgaria brachanthemoides* as an outgroup of *Artemisia* lives in dry
 400 mountain valleys or dry riverbeds of Northwest China (Toksun) and Kazakhstan, with highly characteristic
 401 pollen (Fig. 14a), narrow habitats (Fig. 17b), and regional distribution (Fig. 16-34) and has the potential to
 402 indicate some specific habitats.

403 **5 Data availability**

404 Pollen datasets (Table 2) including pollen photographs under LM and SEM, statistical data of pollen
 405 morphological traits, and their source plant distribution for each species are available at Zenodo
 406 (<https://doi.org/10.5281/zenodo.6791891>; Lu et al., 2022).

407 **Table 2.** *Artemisia* pollen datasets in this study.

Data type	Data format	Data acquisition	Data accessibility
The phylogenetic framework of <i>Artemisia</i> pollen sampling.	.png	Literature survey (modified from Malik et al., 2017).	
A voucher specimen list of 36 representative species.	.doc	Pollen samples were obtained from PE herbarium at the Institute of Botany, Chinese Academy of Sciences.	This article
12 illustrations of pollen grains and the habitats of their source plants.	.png	Habitat photos from online sources (Appendix Table A).	
4018 original pollen photographs (3205 under LM, 813 under SEM).	.jpg	Pollen samples were acetolyzed by the standard method and fixed in glycerine jelly. The pollen grains were photographed under LM and SEM using standard procedures.	
9360 statistical pollen morphological traits.	.xlsx	Statistical data of pollen morphological traits were measured by standard methods.	Zenodo (https://doi.org/10.5281/zenodo.6791891 ;
30858 source plant occurrence information, and corresponding environmental factors including altitude and 19 climate parameters.	.xlsx	Their source plant distribution coordinates were obtained from GBIF (https://doi.org/10.15468/dl.596xd9). The corresponding environmental factors of these coordinates were obtained from WorldClim (https://www.worldclim.org/) with a spatial resolution of 30 seconds between 1970-2000.	Lu et al., 2022)

408

6 Summary

409

To cover the maximum range of *Artemisia* pollen morphological variation, we provide a pollen dataset of 36 species from 9 clades and 3 outgroups of *Artemisia* constrained by the phylogenetic framework, containing high-quality pollen photographs under LM and SEM, statistical data of pollen morphological traits, together with their source plant distribution, and corresponding environmental factors. Here, we attempt to decipher the underlying causes of the long-standing disagreement in the palynological community on the correlation between *Artemisia* pollen and aridity by recognizing the different ecological implications of *Artemisia* pollen types.

416

This dataset should work well for identifying and classifying *Artemisia* pollen from Neogene and

417 Quaternary sediments. While *Artemisia* pollen grains are uniform in morphology under LM, different types
418 can be recognized under SEM. So, the single-grain technique for picking out fossil pollen grains and
419 photographing the same grains under LM and SEM should provide valuable insights in the diversity of fossil
420 *Artemisia* (Ferguson et al., 2007; Grímsson et al., 2011; Grímsson et al., 2012; Halbritter et al., 2018).
421 Furthermore, those *Artemisia* pollen grains could then be compared with the rich photographs from this
422 dataset, and together with the key provided here, possibly attributed to one of the three *Artemisia* pollen types,
423 which in turn may provide a link to the different habitat ranges.

424 However, the application of this dataset probably may not work well for the Palaeogene, as 1) *Artemisia*
425 might have originated in the Palaeocene, although there is no evidence for a specific location or time interval
426 of its origin (e.g. Ling 1982; Wang 2004; Miao 2011); 2) both the lack of macrofossils of *Artemisia* and the
427 strong pollen similarity between *Artemisia* and its closely related taxa under LM might lead to confusion and
428 more uncertainty in tracing the origin of *Artemisia*. On the other hand, the present dataset provides a potential
429 morphological tool to distinguish *Artemisia* pollen grains from those of its related taxa at the SEM level and
430 may shed light on the origin of this genus in the Palaeogene.

431 Moreover, these pollen photographs also have potential and the possibility to be used for deep learning
432 research. We are attempting to automatically identify pollen images using pollen assemblages from the eastern
433 Central Asian desert as an example with deep convolutional neural network (DCNN) of artificial intelligence.
434 Pollen images of the many species of *Artemisia* provided here, and the increasing number of intraspecific
435 replications in the future, will all serve for projected image identification research.

436 Finally and most importantly, the *Artemisia* pollen dataset as designed is open and expandable for new
437 pollen data from *Artemisia* worldwide in order to better serve the global environment assessment and refined
438 reconstruction of vegetation in the geological past as a basis or blueprint for other overarching statistical
439 analyses on pollen morphology.

440 **Appendix A**441 **Text A1**442 Pollen morphological descriptions of 36 representative species from 9 clades of *Artemisia* and 3 outgroups.443 Pollen morphology of *Artemisia*: pollen grains oblate, spherical, or ellipsoidal; apertures tricolporate; almost
444 circular in equatorial view and trilobate circular in polar view; the exine near the colpi gradually thinned; the
445 exine has an obvious double structure of inner and outer layers where the outer is thicker than the inner under
446 LM; the exine ornamentation is psilate (LM), spinulate and granule (SEM).447 **1. *Artemisia cana* (Table 1, Figs. 3a, 15)**448 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
449 Apertures tricolporate. The exine near the colpi gradually thinned. Polar length (P) = $23.46 \pm 1.76 \mu\text{m}$ (M ±
450 SD), equatorial width (E) = $24.50 \pm 2.13 \mu\text{m}$ (M ± SD), P/E = 0.96 ± 0.04 (M ± SD), Exine thickness (T) =
451 $3.91 \pm 0.36 \mu\text{m}$ (M ± SD), Pollen length (L) = $24.58 \pm 1.24 \mu\text{m}$ (M ± SD), T/L = 0.16 ± 0.02 . The exine
452 ornamentation is psilate (LM), spinulate (SEM). Under SEM, diameter of spinule base (D) = $0.58 \pm 0.13 \mu\text{m}$
453 (M ± SD), spinule height (H) = $0.46 \pm 0.08 \mu\text{m}$ (M ± SD), D/H = 1.28 ± 0.38 (M ± SD), granule spacing (Gs)
454 = $0.33 \pm 0.08 \mu\text{m}$ (M ± SD), spinule spacing (Ss) = $1.60 \pm 0.22 \mu\text{m}$ (M ± SD), Gs/Ss = 0.21 ± 0.06 (M ± SD).455 Habitat: grasslands, gravel soils, mountain meadows, stream banks; Wet mountain meadows, stream banks,
456 rocky areas with late-lying snows.457 **2. *Artemisia tridentata* (Table 1, Figs. 3b, 15)**458 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
459 Apertures tricolporate. The exine near the colpi gradually thinned. P = $21.36 \pm 1.54 \mu\text{m}$, E = $20.69 \pm 1.85 \mu\text{m}$,
460 P/E = 1.04 ± 0.07 , T = $3.55 \pm 0.41 \mu\text{m}$, L = $22.35 \pm 1.90 \mu\text{m}$, T/L = 0.16 ± 0.02 . The exine ornamentation is
461 psilate (LM), spinulate (SEM). Under SEM, D = $0.76 \pm 0.08 \mu\text{m}$, H = $0.60 \pm 0.08 \mu\text{m}$, D/H = 1.30 ± 0.23 , Gs
462 = $0.24 \pm 0.06 \mu\text{m}$, Ss = $1.12 \pm 0.22 \mu\text{m}$, Gs/Ss = 0.22 ± 0.08 .463 Habitat: mountains, grasslands, and meadows of western North America. Arid and semi-arid, desert, or
464 semi-desert areas of the growing shrub or semi-shrub environment.465 **3. *Artemisia californica* (Table 1, Figs. 3c, 15)**466 Pollen grains prolate or spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar
467 view. Apertures tricolporate. The exine near the colpi gradually thinned. P = $18.94 \pm 1.30 \mu\text{m}$, E = $19.13 \pm$
468 $1.43 \mu\text{m}$, P/E = 0.99 ± 0.08 , T = $2.70 \pm 0.16 \mu\text{m}$, L = $18.85 \pm 1.12 \mu\text{m}$, T/L = 0.14 ± 0.01 . The exine
469 ornamentation is psilate (LM), spinulate (SEM). Under SEM, D = $0.75 \pm 0.11 \mu\text{m}$, H = $0.71 \pm 0.10 \mu\text{m}$, D/H =
470 1.08 ± 0.20 , Gs = $0.24 \pm 0.05 \mu\text{m}$, Ss = $1.45 \pm 0.23 \mu\text{m}$, Gs/Ss = 0.17 ± 0.05 .

471 Habitat: coastal scrub, dry foothills.

472 **4. *Artemisia indica* (Table 1, Figs. 4a, 15)**

473 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
474 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 23.47 \pm 1.39 \mu\text{m}$, $E = 23.81 \pm 0.86 \mu\text{m}$,
475 $P/E = 0.99 \pm 0.06$, $T = 3.50 \pm 0.27 \mu\text{m}$, $L = 23.31 \pm 0.61 \mu\text{m}$, $T/L = 0.15 \pm 0.01$. The exine ornamentation is
476 psilate (LM), spinulate (SEM). Under SEM, $D = 0.76 \pm 0.10 \mu\text{m}$, $H = 0.39 \pm 0.06 \mu\text{m}$, $D/H = 2.04 \pm 0.53$, Gs
477 $= 0.28 \pm 0.07 \mu\text{m}$, $Ss = 1.21 \pm 0.24 \mu\text{m}$, $Gs/Ss = 0.24 \pm 0.07$.

478 Habitat: roadsides, forest margins, slopes, shrublands; low elevations to 2000 m.

479 **5. *Artemisia argyi* (Table 1, Figs. 4b, 15)**

480 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
481 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 21.80 \pm 1.00 \mu\text{m}$, $E = 21.67 \pm 1.27 \mu\text{m}$,
482 $P/E = 1.01 \pm 0.08$, $T = 3.55 \pm 0.40 \mu\text{m}$, $L = 22.24 \pm 1.13 \mu\text{m}$, $T/L = 0.16 \pm 0.01$. The exine ornamentation is
483 psilate (LM), spinulate (SEM). Under SEM, $D = 0.64 \pm 0.07 \mu\text{m}$, $H = 0.38 \pm 0.04 \mu\text{m}$, $D/H = 1.71 \pm 0.23$, Gs
484 $= 0.22 \pm 0.06 \mu\text{m}$, $Ss = 0.90 \pm 0.17 \mu\text{m}$, $Gs/Ss = 0.26 \pm 0.09$.

485 Habitat: waste places, roadsides, slopes, hills, steppes, forest steppes; low elevations to 1500 m.

486 **6. *Artemisia mongolica* (Table 1, Figs. 4c, 15)**

487 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
488 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 21.05 \pm 0.82 \mu\text{m}$, $E = 20.42 \pm 1.01 \mu\text{m}$,
489 $P/E = 1.03 \pm 0.05$, $T = 3.29 \pm 0.19 \mu\text{m}$, $L = 19.78 \pm 0.99 \mu\text{m}$, $T/L = 0.17 \pm 0.01$. The exine ornamentation is
490 psilate (LM), spinulate (SEM). Under SEM, $D = 0.62 \pm 0.08 \mu\text{m}$, $H = 0.41 \pm 0.05 \mu\text{m}$, $D/H = 1.54 \pm 0.25$, Gs
491 $= 0.19 \pm 0.06 \mu\text{m}$, $Ss = 0.91 \pm 0.14 \mu\text{m}$, $Gs/Ss = 0.22 \pm 0.08$.

492 Habitat: slopes, shrublands, riverbanks, lakeshores, roadsides, steppes, forest steppes, dry valleys; low
493 elevations to 2000 m.

494 **7. *Artemisia vulgaris* (Table 1, Figs. 5a, 15)**

495 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
496 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 19.72 \pm 1.25 \mu\text{m}$, $E = 19.29 \pm 1.82 \mu\text{m}$,
497 $P/E = 1.03 \pm 0.08$, $T = 2.92 \pm 0.23 \mu\text{m}$, $L = 18.94 \pm 1.09 \mu\text{m}$, $T/L = 0.16 \pm 0.02$. The exine ornamentation is
498 psilate (LM), spinulate (SEM). Under SEM, $D = 0.69 \pm 0.07 \mu\text{m}$, $H = 0.34 \pm 0.07 \mu\text{m}$, $D/H = 2.13 \pm 0.52$, Gs
499 $= 0.29 \pm 0.07 \mu\text{m}$, $Ss = 1.55 \pm 0.32 \mu\text{m}$, $Gs/Ss = 0.20 \pm 0.07$.

500 Habitat: roadsides, slopes, canyons, forest margins, forest steppes, subalpine steppes; 1500-3800 m.

501 **8. *Artemisia selengensis* (Table 1, Figs. 5b, 15)**

502 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
503 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 20.67 \pm 1.57 \mu\text{m}$, $E = 19.68 \pm 1.94 \mu\text{m}$,
504 $P/E = 1.06 \pm 0.09$, $T = 3.72 \pm 0.72 \mu\text{m}$, $L = 20.80 \pm 2.21 \mu\text{m}$, $T/L = 0.18 \pm 0.03$. The exine ornamentation is
505 psilate (LM), spinulate (SEM). Under SEM, $D = 0.67 \pm 0.08 \mu\text{m}$, $H = 0.38 \pm 0.05 \mu\text{m}$, $D/H = 1.76 \pm 0.27$, Gs
506 $= 0.22 \pm 0.06 \mu\text{m}$, $Ss = 1.05 \pm 0.15 \mu\text{m}$, $Gs/Ss = 0.22 \pm 0.07$.

507 Habitat: riverbanks, lakeshores, humid areas, meadows, slopes, roadsides.

508 **9. *Artemisia ludoviciana* (Table 1, Figs. 5c, 15)**

509 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
510 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 21.65 \pm 1.02 \mu\text{m}$, $E = 20.82 \pm 1.10 \mu\text{m}$,
511 $P/E = 1.04 \pm 0.08$, $T = 3.71 \pm 0.28 \mu\text{m}$, $L = 20.94 \pm 1.13 \mu\text{m}$, $T/L = 0.18 \pm 0.01$. The exine ornamentation is
512 psilate (LM), spinulate (SEM). Under SEM, $D = 0.70 \pm 0.08 \mu\text{m}$, $H = 0.37 \pm 0.04 \mu\text{m}$, $D/H = 1.94 \pm 0.31$, Gs
513 $= 0.20 \pm 0.05 \mu\text{m}$, $Ss = 1.23 \pm 0.13 \mu\text{m}$, $Gs/Ss = 0.16 \pm 0.04$.

514 Habitat: disturbed roadsides, open meadows, rocky slopes.

515 **10. *Artemisia roxburghiana* (Table 1, Figs. 6a, 15)**

516 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
517 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 23.88 \pm 2.04 \mu\text{m}$, $E = 23.69 \pm 2.00 \mu\text{m}$,
518 $P/E = 1.01 \pm 0.06$, $T = 3.78 \pm 0.39 \mu\text{m}$, $L = 21.81 \pm 1.05 \mu\text{m}$, $T/L = 0.17 \pm 0.02$. The exine ornamentation is
519 psilate (LM), spinulate (SEM). Under SEM, $D = 0.76 \pm 0.07 \mu\text{m}$, $H = 0.39 \pm 0.06 \mu\text{m}$, $D/H = 1.96 \pm 0.37$, Gs
520 $= 0.28 \pm 0.11 \mu\text{m}$, $Ss = 0.79 \pm 0.11 \mu\text{m}$, $Gs/Ss = 0.36 \pm 0.14$.

521 Habitat: roadsides, slopes, dry canyons, grasslands, waste areas, terraces; 700-3900 m.

522 **11. *Artemisia rutifolia* (Table 1, Figs. 6b, 15)**

523 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
524 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 22.22 \pm 1.10 \mu\text{m}$, $E = 22.70 \pm 1.37 \mu\text{m}$,
525 $P/E = 0.98 \pm 0.05$, $T = 3.53 \pm 0.37 \mu\text{m}$, $L = 24.93 \pm 1.05 \mu\text{m}$, $T/L = 0.14 \pm 0.01$. The exine ornamentation is
526 psilate (LM), spinulate (SEM). Under SEM, $D = 0.31 \pm 0.04 \mu\text{m}$, $H = 0.26 \pm 0.04 \mu\text{m}$, $D/H = 1.20 \pm 0.18$, Gs
527 $= 0.21 \pm 0.05 \mu\text{m}$, $Ss = 1.27 \pm 0.19 \mu\text{m}$, $Gs/Ss = 0.17 \pm 0.04$.

528 Habitat: hills, dry river valleys, basins, steppes, semideserts, stony desert; 1300-5000 m.

529 **12. *Artemisia chinensis* (Table 1, Figs. 6c, 15)**

530 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
531 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 21.53 \pm 1.95 \mu\text{m}$, $E = 22.75 \pm 2.00 \mu\text{m}$,
532 $P/E = 0.95 \pm 0.05$, $T = 2.97 \pm 0.40 \mu\text{m}$, $L = 23.71 \pm 2.30 \mu\text{m}$, $T/L = 0.13 \pm 0.01$. The exine ornamentation is

533 psilate (LM), spinulate (SEM). Under SEM, $D = 0.70 \pm 0.05 \mu\text{m}$, $H = 0.55 \pm 0.07 \mu\text{m}$, $D/H = 1.29 \pm 0.19$, Gs
534 $= 0.27 \pm 0.07 \mu\text{m}$, $Ss = 0.91 \pm 0.17 \mu\text{m}$, $Gs/Ss = 0.31 \pm 0.09$.

535 Habitat: littoral plants found on raised coral outcrops.

536 **13. *Artemisia kurramensis* (Table 1, Figs. 7a, 15)**

537 Pollen grains spheroidal. Almost circular in equatorial view and trilobate circular in polar view. Apertures
538 tricolporate. The exine near the colpi gradually thinned. $P = 19.71 \pm 1.28 \mu\text{m}$, $E = 19.35 \pm 1.02 \mu\text{m}$, $P/E = 1.02$
539 ± 0.05 , $T = 3.30 \pm 0.38 \mu\text{m}$, $L = 19.44 \pm 0.92 \mu\text{m}$, $T/L = 0.17 \pm 0.02$. The exine ornamentation is psilate (LM),
540 spinulate (SEM). Under SEM, $D = 0.38 \pm 0.04 \mu\text{m}$, $H = 0.27 \pm 0.03 \mu\text{m}$, $D/H = 1.41 \pm 0.21$, $Gs = 0.23 \pm 0.07$
541 μm , $Ss = 1.25 \pm 0.21 \mu\text{m}$, $Gs/Ss = 0.19 \pm 0.06$.

542 Habitat: foothills, mountain slopes, dry graveyards, field borders with sparse vegetation on gravelly, fine to
543 coarse sandy-clay soils.

544 **14. *Artemisia compactum* (Table 1, Figs. 7b, 15)**

545 Pollen grains spheroidal. Almost circular in equatorial view and trilobate circular in polar view. Apertures
546 tricolporate. The exine near the colpi gradually thinned. $P = 22.33 \pm 1.81 \mu\text{m}$, $E = 21.97 \pm 1.23 \mu\text{m}$, $P/E = 1.02$
547 ± 0.06 , $T = 2.97 \pm 0.43 \mu\text{m}$, $L = 21.67 \pm 0.87 \mu\text{m}$, $T/L = 0.14 \pm 0.02$. The exine ornamentation is psilate (LM),
548 spinulate (SEM). Under SEM, $D = 0.41 \pm 0.07 \mu\text{m}$, $H = 0.28 \pm 0.03 \mu\text{m}$, $D/H = 1.50 \pm 0.33$, $Gs = 0.51 \pm 0.12$
549 μm , $Ss = 0.92 \pm 0.12 \mu\text{m}$, $Gs/Ss = 0.56 \pm 0.12$.

550 Habitat: rocky slopes, semi-deserts, from low elevations to sub-alpine areas.

551 **15. *Artemisia maritima* (Table 1, Figs. 7c, 15)**

552 Pollen grains prolate. Almost circular in equatorial view and trilobate circular in polar view. Apertures
553 tricolporate. The exine near the colpi gradually thinned. $P = 26.24 \pm 1.61 \mu\text{m}$, $E = 23.09 \pm 1.43 \mu\text{m}$, $P/E = 1.14$
554 ± 0.06 , $T = 3.54 \pm 0.44 \mu\text{m}$, $L = 24.42 \pm 1.51 \mu\text{m}$, $T/L = 0.14 \pm 0.02$. The exine ornamentation is psilate (LM),
555 spinulate (SEM). Under SEM, $D = 0.28 \pm 0.04 \mu\text{m}$, $H = 0.23 \pm 0.06 \mu\text{m}$, $D/H = 1.30 \pm 0.34$, $Gs = 0.53 \pm 0.12$
556 μm , $Ss = 1.08 \pm 0.12 \mu\text{m}$, $Gs/Ss = 0.50 \pm 0.13$.

557 Habitat: saltmarsh, dry and calcareous hillsides, seashores, and dry saline or alkaline soils.

558 **16. *Artemisia aralensis* (Table 1, Figs. 8a, 15)**

559 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
560 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 22.32 \pm 1.72 \mu\text{m}$, $E = 21.91 \pm 1.63 \mu\text{m}$,
561 $P/E = 1.02 \pm 0.06$, $T = 3.16 \pm 0.36 \mu\text{m}$, $L = 22.76 \pm 1.45 \mu\text{m}$, $T/L = 0.14 \pm 0.01$. The exine ornamentation is
562 psilate (LM), spinulate (SEM). Under SEM, $D = 0.25 \pm 0.04 \mu\text{m}$, $H = 0.22 \pm 0.04 \mu\text{m}$, $D/H = 1.16 \pm 0.28$, Gs
563 $= 0.50 \pm 0.13 \mu\text{m}$, $Ss = 1.09 \pm 0.18 \mu\text{m}$, $Gs/Ss = 0.46 \pm 0.14$.

564 Habitat: clayey, sandy loam, solonetzic soils.

565 **17. *Artemisia annua* (Table 1, Figs. 8b, 15)**

566 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
567 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 19.71 \pm 0.84 \mu\text{m}$, $E = 19.45 \pm 1.32 \mu\text{m}$,
568 $P/E = 1.02 \pm 0.07$, $T = 3.45 \pm 0.25 \mu\text{m}$, $L = 19.20 \pm 0.92 \mu\text{m}$, $T/L = 0.18 \pm 0.01$. The exine ornamentation is
569 psilate (LM), spinulate (SEM). Under SEM, $D = 0.45 \pm 0.06 \mu\text{m}$, $H = 0.39 \pm 0.05 \mu\text{m}$, $D/H = 1.18 \pm 0.25$, Gs
570 $= 0.27 \pm 0.08 \mu\text{m}$, $Ss = 1.29 \pm 0.16 \mu\text{m}$, $Gs/Ss = 0.21 \pm 0.08$.

571 Habitat: hills, waysides, wastelands, outer forest margins, steppes, forest steppes, dry flood lands, terraces,
572 semidesert steppes, rocky slopes, roadsides, saline soils; 2000-3700 m.

573 **18. *Artemisia freyniana* (Table 1, Figs. 8c, 15)**

574 Pollen grains prolate. Almost circular in equatorial view and trilobate circular in polar view. Apertures
575 tricolporate. The exine near the colpi gradually thinned. $P = 23.39 \pm 1.21 \mu\text{m}$, $E = 21.30 \pm 1.07 \mu\text{m}$, $P/E = 1.10$
576 ± 0.04 , $T = 3.17 \pm 0.26 \mu\text{m}$, $L = 21.29 \pm 0.95 \mu\text{m}$, $T/L = 0.15 \pm 0.01$. The exine ornamentation is psilate (LM),
577 spinulate (SEM). Under SEM, $D = 0.56 \pm 0.05 \mu\text{m}$, $H = 0.40 \pm 0.06 \mu\text{m}$, $D/H = 1.40 \pm 0.15$, $Gs = 0.20 \pm 0.05$
578 μm , $Ss = 1.15 \pm 0.15 \mu\text{m}$, $Gs/Ss = 0.18 \pm 0.05$.

579 Habitat: steppes, slopes, dry river valleys, riverbanks, outer forest margins.

580 **19. *Artemisia stachmanniana* (Table 1, Figs. 9a, 15)**

581 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
582 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 26.31 \pm 1.48 \mu\text{m}$, $E = 25.16 \pm 1.22 \mu\text{m}$,
583 $P/E = 1.05 \pm 0.07$, $T = 3.97 \pm 0.60 \mu\text{m}$, $L = 23.45 \pm 1.38 \mu\text{m}$, $T/L = 0.17 \pm 0.02$. The exine ornamentation is
584 psilate (LM), spinulate (SEM). Under SEM, $D = 0.37 \pm 0.05 \mu\text{m}$, $H = 0.35 \pm 0.05 \mu\text{m}$, $D/H = 1.07 \pm 0.25$, Gs
585 $= 0.19 \pm 0.04 \mu\text{m}$, $Ss = 1.40 \pm 0.24 \mu\text{m}$, $Gs/Ss = 0.14 \pm 0.04$.

586 Habitat: hillsides, roadsides, shrubland, and forest-steppe areas, and often becoming the dominant species or
587 main associated species of plant communities in some areas of mountainous sunny slopes.

588 **20. *Artemisia pontica* (Table 1, Figs. 9b, 15)**

589 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
590 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 20.64 \pm 1.54 \mu\text{m}$, $E = 19.62 \pm 1.59 \mu\text{m}$,
591 $P/E = 1.05 \pm 0.07$, $T = 3.01 \pm 0.39 \mu\text{m}$, $L = 19.75 \pm 0.84 \mu\text{m}$, $T/L = 0.15 \pm 0.02$. The exine ornamentation is
592 psilate (LM), spinulate (SEM). Under SEM, $D = 0.60 \pm 0.11 \mu\text{m}$, $H = 0.37 \pm 0.06 \mu\text{m}$, $D/H = 1.63 \pm 0.37$, Gs
593 $= 0.17 \pm 0.04 \mu\text{m}$, $Ss = 1.32 \pm 0.27 \mu\text{m}$, $Gs/Ss = 0.13 \pm 0.04$.

594 Habitat: rocky slopes, dry valleys, steppes, hills; low to middle elevations.

595 **21. *Artemisia frigida* (Table 1, Figs. 9c, 15)**

596 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
597 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 25.11 \pm 1.75 \mu\text{m}$, $E = 24.90 \pm 1.48 \mu\text{m}$,
598 $P/E = 1.01 \pm 0.07$, $T = 4.61 \pm 0.74 \mu\text{m}$, $L = 24.83 \pm 1.27 \mu\text{m}$, $T/L = 0.19 \pm 0.02$. The exine ornamentation is
599 psilate (LM), spinulate (SEM). Under SEM, $D = 0.46 \pm 0.08 \mu\text{m}$, $H = 0.32 \pm 0.04 \mu\text{m}$, $D/H = 1.44 \pm 0.26$, Gs
600 $= 0.31 \pm 0.08 \mu\text{m}$, $Ss = 1.30 \pm 0.18 \mu\text{m}$, $Gs/Ss = 0.24 \pm 0.06$.

601 Habitat: steppes, sub-alpine meadows, dry hillsides, stable dunes, dry waste areas; 1000-4000 m.

602 **22. *Artemisia rupestris* (Table 1, Figs. 10a, 15)**

603 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
604 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 24.45 \pm 1.41 \mu\text{m}$, $E = 22.92 \pm 1.40 \mu\text{m}$,
605 $P/E = 1.07 \pm 0.08$, $T = 3.18 \pm 0.40 \mu\text{m}$, $L = 21.96 \pm 1.15 \mu\text{m}$, $T/L = 0.14 \pm 0.02$. The exine ornamentation is
606 psilate (LM), spinulate (SEM). Under SEM, $D = 0.55 \pm 0.05 \mu\text{m}$, $H = 0.33 \pm 0.04 \mu\text{m}$, $D/H = 1.68 \pm 0.28$, Gs
607 $= 0.25 \pm 0.07 \mu\text{m}$, $Ss = 0.91 \pm 0.11 \mu\text{m}$, $Gs/Ss = 0.28 \pm 0.09$.

608 Habitat: dry hills, desert or semidesert steppes, grassy marshlands, dry river valleys, riverbeds, scrub, forest
609 margins.

610 **23. *Artemisia sericea* (Table 1, Figs. 10b, 15)**

611 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
612 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 26.31 \pm 1.31 \mu\text{m}$, $E = 27.90 \pm 1.67 \mu\text{m}$,
613 $P/E = 0.94 \pm 0.03$, $T = 3.75 \pm 0.32 \mu\text{m}$, $L = 26.89 \pm 2.12 \mu\text{m}$, $T/L = 0.14 \pm 0.01$. The exine ornamentation is
614 psilate (LM), spinulate (SEM). Under SEM, $D = 0.89 \pm 0.09 \mu\text{m}$, $H = 0.54 \pm 0.10 \mu\text{m}$, $D/H = 1.71 \pm 0.36$, Gs
615 $= 0.28 \pm 0.07 \mu\text{m}$, $Ss = 1.74 \pm 0.31 \mu\text{m}$, $Gs/Ss = 0.16 \pm 0.05$.

616 Habitat: Forest margins, hills, steppes, canyons, waste areas.

617 **24. *Artemisia absinthium* (Table 1, Figs. 10c, 15)**

618 Pollen grains prolate. Almost circular in equatorial view and trilobate circular in polar view. Apertures
619 tricolporate. The exine near the colpi gradually thinned. $P = 22.79 \pm 1.22 \mu\text{m}$, $E = 20.84 \pm 1.11 \mu\text{m}$, $P/E = 1.09$
620 ± 0.05 , $T = 3.39 \pm 0.31 \mu\text{m}$, $L = 19.92 \pm 1.74 \mu\text{m}$, $T/L = 0.17 \pm 0.01$. The exine ornamentation is psilate (LM),
621 spinulate (SEM). Under SEM, $D = 0.59 \pm 0.05 \mu\text{m}$, $H = 0.40 \pm 0.06 \mu\text{m}$, $D/H = 1.52 \pm 0.25$, $Gs = 0.18 \pm 0.04$
622 μm , $Ss = 1.11 \pm 0.15 \mu\text{m}$, $Gs/Ss = 0.16 \pm 0.04$.

623 Habitat: hillsides, steppes, scrub, forest margins, often in locally moist situations; 1100-1500 m.

624 **25. *Artemisia abrotanum* (Table 1, Figs. 11a, 15)**

625 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
626 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 24.47 \pm 1.56 \mu\text{m}$, $E = 23.73 \pm 1.65 \mu\text{m}$,
627 $P/E = 1.03 \pm 0.07$, $T = 3.15 \pm 0.28 \mu\text{m}$, $L = 18.82 \pm 0.81 \mu\text{m}$, $T/L = 0.17 \pm 0.01$. The exine ornamentation is
628 psilate (LM), spinulate (SEM). Under SEM, $D = 0.72 \pm 0.10 \mu\text{m}$, $H = 0.51 \pm 0.05 \mu\text{m}$, $D/H = 1.44 \pm 0.25$, Gs
629 $= 0.22 \pm 0.04 \mu\text{m}$, $Ss = 1.41 \pm 0.19 \mu\text{m}$, $Gs/Ss = 0.16 \pm 0.04$.

630 Habitat: the wasteland of western, southern, central, and southern Europe.

631 **26. *Artemisia blepharolepis* (Table 1, Figs. 11b, 15)**

632 Pollen grains spheroidal. Almost circular in equatorial view and trilobate circular in polar view. Apertures
633 tricolporate. The exine near the colpi gradually thinned. $P = 18.96 \pm 0.98 \mu\text{m}$, $E = 19.26 \pm 0.99 \mu\text{m}$, $P/E = 0.99$
634 ± 0.05 , $T = 3.15 \pm 0.28 \mu\text{m}$, $L = 18.82 \pm 0.81 \mu\text{m}$, $T/L = 0.17 \pm 0.01$. The exine ornamentation is psilate (LM),
635 spinulate (SEM). Under SEM, $D = 0.69 \pm 0.09 \mu\text{m}$, $H = 0.44 \pm 0.07 \mu\text{m}$, $D/H = 1.64 \pm 0.44$, $Gs = 0.37 \pm 0.18$
636 μm , $Ss = 1.68 \pm 0.20 \mu\text{m}$, $Gs/Ss = 0.23 \pm 0.14$.

637 Habitat: low-altitude areas of dry slopes, grasslands, steppes, waste areas, roadsides, dunes near riverbanks.

638 **27. *Artemisia norvegica* (Table 1, Figs. 11c, 15)**

639 Pollen grains prolate. Almost circular in equatorial view and trilobate circular in polar view. Apertures
640 tricolporate. The exine near the colpi gradually thinned. $P = 24.51 \pm 1.40 \mu\text{m}$, $E = 22.11 \pm 1.05 \mu\text{m}$, $P/E = 1.11$
641 ± 0.06 , $T = 3.48 \pm 0.39 \mu\text{m}$, $L = 22.61 \pm 1.31 \mu\text{m}$, $T/L = 0.15 \pm 0.01$. The exine ornamentation is psilate (LM),
642 spinulate (SEM). Under SEM, $D = 0.67 \pm 0.08 \mu\text{m}$, $H = 0.43 \pm 0.11 \mu\text{m}$, $D/H = 1.66 \pm 0.51$, $Gs = 0.19 \pm 0.03$
643 μm , $Ss = 1.56 \pm 0.24 \mu\text{m}$, $Gs/Ss = 0.12 \pm 0.03$.

644 Habitat: bare stony ground, Racomitrium heath, bouldery crests of solifluction terraces, and sometimes
645 hollows between rocks.

646 **28. *Artemisia tanacetifolia* (Table 1, Figs. 12a, 15)**

647 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
648 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 28.38 \pm 0.90 \mu\text{m}$, $E = 27.75 \pm 1.70 \mu\text{m}$,
649 $P/E = 1.03 \pm 0.06$, $T = 3.46 \pm 0.47 \mu\text{m}$, $L = 27.63 \pm 1.06 \mu\text{m}$, $T/L = 0.13 \pm 0.02$. The exine ornamentation is
650 psilate (LM), spinulate (SEM). Under SEM, $D = 0.71 \pm 0.06 \mu\text{m}$, $H = 0.32 \pm 0.04 \mu\text{m}$, $D/H = 2.23 \pm 0.40$, Gs
651 $= 0.30 \pm 0.07 \mu\text{m}$, $Ss = 1.08 \pm 0.16 \mu\text{m}$, $Gs/Ss = 0.29 \pm 0.07$.

652 Habitat: middle and low-altitude areas of forest grasslands, grasslands, meadows, forest edges, open forests,
653 salty grasslands, grass slopes, and brushwood.

654 **29. *Artemisia tournefortiana* (Table 1, Figs. 12b, 15)**

655 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
656 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 20.76 \pm 0.98 \mu\text{m}$, $E = 20.43 \pm 0.83 \mu\text{m}$,

657 P/E = 1.02 ± 0.06 , T = 3.33 ± 0.19 μm , L = 20.03 ± 0.79 μm , T/L = 0.17 ± 0.01 . The exine ornamentation is
658 psilate (LM), spinulate (SEM). Under SEM, D = 0.73 ± 0.06 μm , H = 0.42 ± 0.07 μm , D/H = 1.81 ± 0.33 , Gs
659 = 0.26 ± 0.07 μm , Ss = 1.25 ± 0.20 μm , Gs/Ss = 0.22 ± 0.08 .

660 Habitat: widely distributed on hills, terraces, dry flood lands, waste fields, steppes, open forests,
661 semi-marshlands.

662 **30. *Artemisia dracunculus* (Table 1, Figs. 12c, 15)**

663 Pollen grains spheroidal. Almost circular in equatorial view and trilobate circular in polar view. Apertures
664 tricolporate. The exine near the colpi gradually thinned. P = 22.89 ± 1.24 μm , E = 22.87 ± 1.32 μm , P/E = 1.00
665 ± 0.05 , T = 2.82 ± 0.52 μm , L = 21.91 ± 1.35 μm , T/L = 0.13 ± 0.03 . The exine ornamentation is psilate (LM),
666 spinulate (SEM). Under SEM, D = 0.68 ± 0.05 μm , H = 0.45 ± 0.07 μm , D/H = 1.56 ± 0.31 , Gs = 0.31 ± 0.10
667 μm , Ss = 0.92 ± 0.15 μm , Gs/Ss = 0.34 ± 0.11 .

668 Habitat: dry slopes, steppes, semidesert steppes, forest steppes, forest margins, waste areas, roadsides, terraces,
669 subalpine meadows, meadow steppes, dry river valleys, rocky slopes, saline-alkaline soils; 500-3800 m.

670 **31. *Artemisia japonica* (Table 1, Figs. 13a, 15)**

671 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
672 Apertures tricolporate. The exine near the colpi gradually thinned. P = 20.18 ± 1.28 μm , E = 21.23 ± 1.26 μm ,
673 P/E = 0.95 ± 0.05 , T = 4.24 ± 0.49 μm , L = 21.02 ± 1.14 μm , T/L = 0.20 ± 0.02 . The exine ornamentation is
674 psilate (LM), spinulate (SEM). Under SEM, D = 0.57 ± 0.05 μm , H = 0.32 ± 0.05 μm , D/H = 1.80 ± 0.24 , Gs
675 = 0.26 ± 0.05 μm , Ss = 1.26 ± 0.16 μm , Gs/Ss = 0.21 ± 0.04 .

676 Habitat: forest margins, waste areas, shrublands, hills, slopes, roadsides. Low elevations to 3300 m.

677 **32. *Artemisia capillaris* (Table 1, Figs. 13b, 15)**

678 Pollen grains spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar view.
679 Apertures tricolporate. The exine near the colpi gradually thinned. P = 19.53 ± 1.09 μm , E = 19.64 ± 1.62 μm ,
680 P/E = 1.00 ± 0.08 , T = 3.54 ± 0.34 μm , L = 19.18 ± 0.97 μm , T/L = 0.18 ± 0.01 . The exine ornamentation is
681 psilate (LM), spinulate (SEM). Under SEM, D = 0.51 ± 0.06 μm , H = 0.36 ± 0.04 μm , D/H = 1.44 ± 0.30 , Gs
682 = 0.26 ± 0.04 μm , Ss = 1.27 ± 0.16 μm , Gs/Ss = 0.21 ± 0.05 .

683 Habitat: humid slopes, hills, terraces, roadsides, riverbanks; 100-2700 m.

684 **33. *Artemisia campestris* (Table 1, Figs. 13c, 15)**

685 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
686 Apertures tricolporate. The exine near the colpi gradually thinned. P = 21.69 ± 0.85 μm , E = 21.26 ± 0.89 μm ,
687 P/E = 1.02 ± 0.07 , T = 3.68 ± 0.33 μm , L = 21.21 ± 0.89 μm , T/L = 0.17 ± 0.02 . The exine ornamentation is

688 psilate (LM), spinulate (SEM). Under SEM, $D = 0.57 \pm 0.09 \mu\text{m}$, $H = 0.38 \pm 0.05 \mu\text{m}$, $D/H = 1.53 \pm 0.23$, Gs
689 $= 0.41 \pm 0.09 \mu\text{m}$, $Ss = 1.23 \pm 0.19 \mu\text{m}$, $Gs/Ss = 0.34 \pm 0.08$.

690 Habitat: steppes, waste areas, rocky slopes, dune margins; 300-3100 m.

691 **34. *Kaschgaria brachanthemoides* (Table 1, Figs. 14a, 15)**

692 Pollen grains prolate or spheroidal. Almost circular in equatorial view and trilobate circular in polar view.
693 Apertures tricolporate. The exine near the colpi gradually thinned. $P = 23.26 \pm 1.44 \mu\text{m}$, $E = 22.09 \pm 1.18 \mu\text{m}$,
694 $P/E = 1.06 \pm 0.08$, $T = 3.93 \pm 0.44 \mu\text{m}$, $L = 21.01 \pm 1.28 \mu\text{m}$, $T/L = 0.19 \pm 0.02$. The exine ornamentation is
695 psilate (LM), spinulate (SEM). Under SEM, $D = 0.55 \pm 0.07 \mu\text{m}$, $H = 0.44 \pm 0.05 \mu\text{m}$, $D/H = 1.25 \pm 0.20$, Gs
696 $= 0 \mu\text{m}$, $Ss = 1.75 \pm 0.20 \mu\text{m}$, $Gs/Ss = 0$, Pertorations spacing (Ps) $= 0.47 \pm 0.14 \mu\text{m}$.

697 Habitat: dry mountain valleys, old dry riverbeds; 1000-1500 m.

698 **35. *Ajania pallasiana* (Table 1, Figs. 14b, 15)**

699 Pollen grains spheroidal. Almost circular in equatorial view and trilobate circular in polar view. Apertures
700 tricolporate. The exine near the colpi gradually thinned. $P = 35.16 \pm 2.68 \mu\text{m}$, $E = 35.92 \pm 3.31 \mu\text{m}$, $P/E = 0.98$
701 ± 0.03 , $T = 10.23 \pm 0.85 \mu\text{m}$, $L = 38.31 \pm 2.06 \mu\text{m}$, $T/L = 0.27 \pm 0.03 \mu\text{m}$. The exine ornamentation spinose.
702 Under SEM, $D = 4.41 \pm 0.35 \mu\text{m}$, $H = 3.47 \pm 0.38 \mu\text{m}$, $D/H = 1.29 \pm 0.21$, $Gs = 0 \mu\text{m}$, $Ss = 7.84 \pm 1.25 \mu\text{m}$,
703 $Gs/Ss = 0$, $Ps = 0.39 \pm 0.12 \mu\text{m}$.

704 Habitat: thickets, mountain slopes, 200-2900 m.

705 **36. *Chrysanthemum indicum* (Table 1, Figs. 14c, 15)**

706 Pollen grains prolate or spheroidal or oblate. Almost circular in equatorial view and trilobate circular in polar
707 view. Apertures tricolporate. The exine near the colpi gradually thinned. $P = 33.54 \pm 1.71 \mu\text{m}$, $E = 34.42 \pm$
708 $2.46 \mu\text{m}$, $P/E = 0.98 \pm 0.08$, $T = 8.65 \pm 0.89 \mu\text{m}$, $L = 34.82 \pm 1.65 \mu\text{m}$, $T/L = 0.25 \pm 0.02$. The exine
709 ornamentation spinose. Under SEM, $D = 2.94 \pm 0.33 \mu\text{m}$, $H = 3.59 \pm 0.29 \mu\text{m}$, $D/H = 0.82 \pm 0.10$, $Gs = 0 \mu\text{m}$,
710 $Ss = 7.11 \pm 0.76 \mu\text{m}$, $Gs/Ss = 0$, $Ps = 0.37 \pm 0.13 \mu\text{m}$.

711 Habitat: grasslands on mountain slopes, thickets, wet places by rivers, fields, roadsides, saline places by
712 seashores, under shrubs, 100-2900 m.

713 **Appendix B**714 **Table B1.** List of the voucher specimen in PE Herbarium, Institute of Botany, Chinese Academy of Sciences

Subgenus	Species	Specimen barcodes	Coll. No.	Habitat photograph sources
Subg. Tridentata	<i>Artemisia cana</i>	PE 01668975	H.Mozingo 79-97	© Jason Headley https://www.inaturalist.org/photos/54492753
	<i>Artemisia tridentata</i>	PE 01917565	Debreczy-Racz- Biro s.n.	© Matt Berger https://www.inaturalist.org/photos/17436654
	<i>Artemisia californica</i>	PE 01668942	Lewis S.Rose 69107	© Don Rideout https://www.inaturalist.org/photos/108921528
Subg. Artemisia, Sect. Artemisia	<i>Artemisia indica</i>	PE 00444597	Tian-Lun Dai 104336	© yangting https://www.inaturalist.org/photos/66336449
	<i>Artemisia argyi</i>	PE 00420930	K.M.Liou 9276	© sergeyprokopenko https://www.inaturalist.org/photos/95820686
	<i>Artemisia mongolica</i>	PE 00445665	Cheng-Yuan Yang & Zu-Gui Li 36466a	© Nikolay V Dorofeev https://www.inaturalist.org/photos/163584035
	<i>Artemisia vulgaris</i>	PE 01669703	P.Frost-Olsen 1833	© Sara Rall https://www.inaturalist.org/photos/120600448
	<i>Artemisia selengensis</i>	PE 00479106	Ming-Gang Li et al. 486	© Gularjanz Grigoryi Mihajlovich https://www.inaturalist.org/photos/46352423
	<i>Artemisia ludoviciana</i>	PE 01669278	W.Hess 2405	© Ethan Rose https://www.inaturalist.org/photos/77690333
	<i>Artemisia roxburghiana</i>	PE 00478222	Xingan collection team 70	© Bo-Han Jiao © Daba
	<i>Artemisia rutifolia</i>	PE 00478427	Ke Guo 12528	https://www.inaturalist.org/photos/62207191
	<i>Artemisia chinensis</i>	PE 01565620	Y.Tateishi J.Murata.Y.Endo et al. 15202	© Jia-Hao Shen

	<i>Artemisia kurramensis</i>	PE 01669178	M.Togasi 1672	© Andrey Vlasenko https://www.inaturalist.org/photos/133758174
	<i>Artemisia compactum</i>	PE 00457459	Hexi team 313	© Chen Chen
				© torkild
Subg. <i>Seriphidium</i>	<i>Artemisia maritima</i>	No. 1338063	s.n.	https://www.inaturalist.org/photos/86515371
				© Полынь аральская
	<i>Artemisia aralensis</i>	No. 202006	s.n.	https://www.plantarium.ru/lang/en/page/image/id/73063.html
				Wen-Hong
	<i>Artemisia annua</i>	PE 01197344	Jin-Tian, Kai-Yong Lang, Ge Yang 328	© Chen Chen
				© Шильников Дмитрий
	<i>Artemisia freyniana</i>	PE 01669030	S.Kharkevich 753	Сергеевич https://www.inaturalist.org/photos/154390279
Subg. Artemisia, Sect. Abrotanum I	<i>Artemisia stachmanniana</i>	PE 00478480	Shen-E Liu, Pei-Yun Fu et al. 4715	© Bo-Han Jiao
				© Martin Pražák
	<i>Artemisia pontica</i>	PE 01589110	Gy.Szollat & K.Dobolyi s.n.	https://www.inaturalist.org/photos/93438780
				© Suzanne Dingwell
	<i>Artemisia frigida</i>	PE 00444197	Ren-Chang Qin 0913	https://www.inaturalist.org/photos/125022240
Subg. <i>Absinthium</i>	<i>Artemisia rupestris</i>	PE 00478380	Anonymous 948	© Bo-Han Jiao
				© svetlana_katana
	<i>Artemisia sericea</i>	PE 01669585	N.Maltzev 3175	https://www.inaturalist.org/photos/48033353
				© Станислав Лебедев
	<i>Artemisia absinthium</i>	PE 01668816	G.Bujorean s.n.	https://www.inaturalist.org/photos/123569286
Subg. Artemisia, Sect. Abrotanum II	<i>Artemisia abrotanum</i>	PE 01668792	T.Leonova s.n.	© Андрей Москвичев https://www.inaturalist.org/photos/116106722
				© Ji-Ye Zheng
Subg. Artemisia, Sect. Abrotanum III	<i>Artemisia blepharolepis</i>	PE 00421006	Kun-Jun Fu 7252	© Erin Springinotic https://www.inaturalist.org/photos/161393521

	<i>Artemisia tanacetifolia</i>	PE 00479744	T.P.Wang W.3379	© Alexander Dubynin https://www.inaturalist.org/photos/78902853
	<i>Artemisia tournefortiana</i>	PE 00479786	Ren-Chang Qin 2266	© Chen Chen
	<i>Artemisia dracunculus</i>	PE 00421462	Shen-E Liu et al. 8084	© anatolymikhaltsov https://www.inaturalist.org/photos/76312868
Subg.	<i>Artemisia japonica</i>	PE 00444874	Qianbei team 2850	© 陳達智 https://www.inaturalist.org/photos/44507659
Dracunculus	<i>Artemisia capillaris</i>	PE 00421156	Han-Chen Wang 4078	© Cheng-Tao Lin https://www.inaturalist.org/photos/60639286
	<i>Artemisia campestris</i>	PE 00421097	T.N.Liou L.1008	© pedrosanz-anapri https://www.inaturalist.org/photos/113822257
	<i>Kaschagaria brachanthemoides</i>	PE 01577564	Yun-Wen Tian 22158	© Chen Chen
Outgroups	<i>Ajania pallasiana</i>	PE 00420032	Guang-Zheng Wang 497	© Игорь Пospelов https://www.inaturalist.org/photos/162408714
	<i>Chrysanthemum indicum</i>	PE 01258852	Anonymous 221	© Bo-Han Jiao

715 Note: In the absence of habitat photographs of two species, habitat photographs of species with which they have close
 716 phylogenetic relationships and similar habitats were used in this study instead, i.e. the habitat photograph of *Kaschagaria*
 717 *komarovii* was used instead of *Kaschagaria brachanthemoides*, the habitat photograph of *Artemisia taurica* for *Artemisia*
 718 *kurramensis*.

719 **Author contributions.** YFW, YFY, TGG conceived the ideas, LLL, BHJ, KQL, and BS collected the
720 literature, LLL extracted and compiled the data, LLL, FQ, and BHJ made the statistical analysis, GX and ML
721 collected pictures, LLL, KQL, and BS drew the figures and tables, LLL, YFW, YFY, LJF, FQ, and GX wrote
722 the first draft of this manuscript, DKF corrected the various versions of the manuscript, while all authors
723 contributed substantially to revisions.

724 **Competing interests.** The authors declare that they have no conflict of interest.

725 **Acknowledgments.** We thank Dr. Jian Yang, Institute of Botany, Chinese Academy of Sciences, for his kind
726 help in drafting graphics. We appreciate Chen Chen from Institute of Botany, Chinese Academy of Sciences,
727 Ji-Ye Zheng from No. 1 Middle School of Jiyang Shandong and Jia-Hao Shen from Institute of Botany,
728 Jiangsu Province and Chinese Academy of Sciences, for their enthusiastic assistance in providing habitat
729 photographs.

730 **Financial support.** This research was supported by the Strategic Priority Research Program of the Chinese
731 Academy of Sciences (No. XDB26000000), National Natural Science Foundation of China (Nos. 31970223,
732 32070240, and 42077416), and the Chinese Academy of Sciences President's International Fellowship
733 Initiative (No. 2018VBA0016).

734 **References**

- 735 Beerling, D. J. and Royer, D. L.: Convergent Cenozoic CO₂ history, *Nat. Geosci.*, 4, 418-420, <https://doi.org/10.1038/ngeo1186>,
736 2011.
- 737 Bhattacharya, T., Tierney, J. E., Addison, J. A., and Murray, J. W.: Ice-sheet modulation of deglacial North American
738 monsoon intensification, *Nat. Geosci.*, 11, 848-852, <https://doi.org/10.1038/s41561-018-0220-7>, 2018.
- 739 Blackmore, S., Wortley, A. H., Skvarla, J. J., and Robinson, H., V. A. Funk, A. Susanna, Stuessy, T. F., and Bayer, R. J.
740 (Eds.): Evolution of pollen in Compositae. In *Systematics, Evolution and Biogeography of the Compositae*,
741 International Association of Plant Taxonomy, Vienna, 2009.
- 742 Bremer, K. and Humphries, C. J.: Generic monograph of the Asteraceae-Anthemideae, *Bull. Nat. Hist. Mus.*, 23, 71-177,
743 <https://www.biodiversitylibrary.org/item/19562>, 1993.
- 744 Brummitt, N., Araujo, A. C., and Harris, T.: Areas of plant diversity-What do we know?, *Plants people planet*, 3, 33-44,
745 <https://doi.org/10.1002/ppp3.10110>, 2021.
- 746 Cai, M., Ye, P., Yang, X., and Li, C.: Vegetation and climate change in the Hetao Basin (Northern China) during the last
747 interglacial-glacial cycle, *J. Asian Earth Sci.*, 171, 1-8, <https://doi.org/10.1016/j.jseas.2018.11.024>, 2019.
- 748 Cao, X. Y., Tian, F., Li, K., Ni, J., Yu, X. S., Liu, L. N., and Wang, N. N.: Lake surface sediment pollen dataset for the
749 alpine meadow vegetation type from the eastern Tibetan Plateau and its potential in past climate reconstructions,
750 *Earth Syst. Sci. Data*, 13, 3525-3537, <https://doi.org/10.5194/essd-13-3525-2021>, 2021.
- 751 Chen, J. X., Shi, X. F., Liu, Y. G., Qiao, S. Q., Yang, S. X., Yan, S. J., Lv, H. H., Li, J. Y., Li, X. Y., and Li, C. X.:
752 Holocene vegetation dynamics in response to climate change and hydrological processes in the Bohai region, *Clim.
753 Past.*, 16, 2509-2531, <https://doi.org/10.5194/cp-16-2509-2020>, 2020.
- 754 Chen, S. B.: Pollen Morphology of *Artemisia* L. from China: A Discussion on the Relationship between Pollen
755 Morphology of *Artemisia* L. and Allies, 1987 (in Chinese).
- 756 Chen, S. B. and Zhang, J. T.: A Study on Pollen Morphology of Some Chinese Genera in Tribe Anthemideae, *Acta
757 Phytotax. Sin.*, 29, 246-251, 1991 (in Chinese).
- 758 China Vegetation Editorial Committee, Wu, Z. Y. (Ed.): Chinese Vegetation Science Press, Beijing, 1980 (in Chinese).
- 759 Cui, Q. Y., Zhao, Y., Qin, F., Liang, C., Li, Q., and Geng, R. W.: Characteristics of the modern pollen assemblages from
760 different vegetation zones in Northeast China: Implications for pollen-based climate reconstruction, *Sci.
761 China-Earth Sci.*, 62, 1564-1577, <https://doi.org/10.1007/s11430-018-9386-9>, 2019.
- 762 Davies, C. P. and Fall, P. L.: Modern pollen precipitation from an elevational transect in central Jordan and its
763 relationship to vegetation, *J. Biogeogr.*, 28, 1195-1210, <https://doi.org/10.1046/j.1365-2699.2001.00630.x>, 2001.
- 764 El-Moslimany, A. P.: Ecological significance of common nonarboreal pollen : examples from drylands of the Middle East,
765 *Rev. Palaeobot. Palynol.*, 64, 343-350, [https://doi.org/10.1016/0034-6667\(90\)90150-h](https://doi.org/10.1016/0034-6667(90)90150-h), 1990.
- 766 Erdtman, G.: The acetolysis method, a revised descriptions, *Svensk Botanisk Tidskrift*, 54, 561-564, 1960.
- 767 Ferguson, D. K., Zetter, R., and Paudyal, K. N.: The need for the SEM in palaeopalynology, *C. R. Palevol*, 6, 423-430,
768 <http://doi.org/10.1016/j.crpv.2007.09.018>, 2007.

- 769 Ghahraman, A., Nourbakhsh, N., Mehdi, G. K., and Atar, F.: Pollen Morphology of *Artemisia* L. (Asteraceae) in Iran,
770 Iran. Journ. Bot., 13, 21-29, 2007.
- 771 GBIF.org GBIF Occurrence Download: <https://doi.org/10.15468/dl.596xd9>, last access: 09 November 2021.
- 772 Grímsson, F., Zetter, R., and Hofmann, C.: *Lythrum* and *Peplis* from the Late Cretaceous and Cenozoic of North America
773 and Eurasia: new evidence suggesting early diversification within the Lythraceae, Am. J. Bot., 98, 1801-1815,
774 <https://doi.org/10.3732/ajb.1100204>, 2011.
- 775 Grímsson, F., Zetter, R., and Leng, Q.: Diverse fossil Onagraceae pollen from a Miocene palynoflora of north-east China:
776 early steps in resolving the phytogeographic history of the family, Plant Syst. Evol., 298, 671-687,
777 <https://doi.org/10.1007/s00606-011-0578-0>, 2012.
- 778 Guiot, J. and Cramer, W.: Climate change: The 2015 Paris Agreement thresholds and Mediterranean basin ecosystems,
779 Science, 354, 465-468, <https://doi.org/10.1126/science.aah5015>, 2016.
- 780 Halbritter, H., Silvia, U., Grímsson, F., Weber, M., Zetter, R., Hesse, M., Buchner, R., Svojtka, M., and Frosch-Radivo, A.:
781 Illustrated Pollen Terminology, Springer Open, 2018.
- 782 Hayat, M. Q., Ashraf, M., Khan, M. A., Yasmin, G., and Jabeen, S.: Palynological study of the genus *Artemisia*
783 (Asteraceae) and its systematic implications, Pak. J. Bot., 42, 751-763, <https://doi.org/10.1094/MPMI-23-4-0522>,
784 2010.
- 785 Hayat, M. Q., Ashraf, M., Khan, M. A., Yasmin, G., Shaheen, N., and Jabeen, S.: Phylogenetic analysis of *Artemisia* L.
786 (Asteraceae) based on micromorphological traits of pollen grains, Afr. J. Biotechnol., 8, 6561-6568,
787 <https://doi.org/10.1556/AMicr.56.2009.4.11>, 2009.
- 788 Herzschuh, U., Tarasov, P., Wünnemann, B., and Kai, H.: Holocene vegetation and climate of the Alashan Plateau, NW
789 China, reconstructed from pollen data, Paleogeogr. Paleoclimatol. Paleoecol., 211, 1-17,
790 <https://doi.org/10.1016/j.palaeo.2004.04.001>, 2004.
- 791 Hesse, M., Buchner, R., Froschradivo, A., Halbritter, H., Ulrich, S., Weber, M., and Zetter, R.: Pollen Terminology : An
792 illustrated handbook, Springer, NewYork, 2009.
- 793 Hussain, A., Potter, D., Hayat, M. Q., Sahreen, S., and Bokhari, S. A. I.: Pollen morphology and its systematic
794 implication on some species of *Artemisia* L. from Gilgit-Baltistan Pakistan, Bangladesh J. Plant Taxon., 26, 157-168,
795 <https://doi.org/10.3329/bjpt.v26i2.44576>, 2019.
- 796 Jiang, L., Q., W., Ye, L. Z., and R., L. Y.: Pollen Morphology of *Artemisia* L. and Its Systematic Significance, Wuhan
797 Univ. J. Nat. Sci., 10, 448-454, <https://doi.org/10.1007/BF02830685>, 2005.
- 798 Koutsodendris, A., Allstadt, F. J., Kern, O. A., Kousis, I., Schwarz, F., Vannacci, M., Woutersen, A., Appel, E., Berke, M.
799 A., Fang, X. M., Friedrich, O., Hoorn, C., Salzmann, U., and Pross, J.: Late Pliocene vegetation turnover on the NE
800 Tibetan Plateau (Central Asia) triggered by early Northern Hemisphere glaciation, Glob. Planet. Change, 180,
801 117-125, <https://doi.org/10.1016/j.gloplacha.2019.06.001>, 2019.
- 802 Li, F., Sun, J., Zhao, Y., Guo, X., Zhao, W., and Zhang, K.: Ecological significance of common pollen ratios: A review,
803 Front. Earth Sci. China, 4, 253-258, <https://doi.org/10.1007/s11707-010-0112-7>, 2010.

- 804 Li, X. L., Hao, Q. Z., Wei, M. J., Andreev, A. A., Wang, J. P., Tian, Y. Y., Li, X. L., Cai, M. T., Hu, J. M., and Shi, W.:
805 Phased uplift of the northeastern Tibetan Plateau inferred from a pollen record from Yinchuan Basin, northwestern
806 China, *Sci. Rep.*, 7, 10, <https://doi.org/10.1038/s41598-017-16915-z>, 2017.
- 807 Ling, Y. R.: On the system of the genus *Artemisia* Linn. and the relationship with allies, *Bulletin of Botanical Research*, 2,
808 1-60, 1982 (in Chinese).
- 809 Ling, Y. R., Humphries, C. J., and Gilbert, M. G.: *Flora of China, The Genus Artemisia L.*, Science Press, Beijing, 2011.
- 810 Liu, H. Y., Wang, Y., Tian, Y. H., Zhu, J. L., and Wang, H. Y.: Climatic and anthropogenic control of surface pollen
811 assemblages in East Asian steppes, *Rev. Palaeobot. Palynol.*, 138, 281-289,
812 <https://doi.org/10.1016/j.revpalbo.2006.01.008>, 2006.
- 813 Lu, K. Q., Qin, F., Li, Y., Xie, G., Li, J. F., Cui, Y. M., Ferguson, D. K., Yao, Y. F., Wang, G. H., and Wang, Y. F.: A new
814 approach to interpret vegetation and ecosystem changes through time by establishing a correlation between surface
815 pollen and vegetation types in the eastern central Asian desert, *Paleogeogr. Paleoclimatol. Paleoecol.*, 551, 12,
816 <https://doi.org/10.1016/j.palaeo.2020.109762>, 2020.
- 817 Lu, L. L., Jiao, B. H., Qin, F., Xie, G., Lu, K. Q., Li, J. F., Sun, B., Li, M., Ferguson, D. K., Gao, T. G., Yao, Y. F., and
818 Wang, Y. F.: *Artemisia* pollen dataset for exploring the potential ecological indicators in deep time, Zenodo [data
819 set], <https://doi.org/zenodo.6791891>, 2022.
- 820 Ma, Q. F., Zhu, L. P., Wang, J. B., Ju, J. T., Lu, X. M., Wang, Y., Guo, Y., Yang, R. M., Kasper, T., Haberzettl, T., and
821 Tang, L. Y.: *Artemisia/Chenopodiaceae* ratio from surface lake sediments on the central and western Tibetan Plateau
822 and its application, *Paleogeogr. Paleoclimatol. Paleoecol.*, 479, 138-145,
823 <https://doi.org/10.1016/j.palaeo.2017.05.002>, 2017.
- 824 Malik, S., Vitales, D., Hayat, M. Q., Korobkov, A. A., Garnatje, T., and Valles, J.: Phylogeny and biogeography of
825 *Artemisia* subgenus *Seriphidium* (Asteraceae: Anthemideae), *Taxon*, 66, 934-952, <https://doi.org/10.12705/664.8>,
826 2017.
- 827 Marsicck, J., Shuman, B. N., Bartlein, P. J., Shafer, S. L., and Brewer, S.: Reconciling divergent trends and millennial
828 variations in Holocene temperatures, *Nature*, 554, 92-96, <https://doi.org/10.1038/nature25464>, 2018.
- 829 Martín, J., Torrell, M., and Valles, J.: Palynological features as a systematic marker in *Artemisia* L. and related genera
830 (Asteraceae, Anthemideae), *Plant Biol.*, 3, 372-378, <https://doi.org/10.1055/s-2001-16462>, 2001.
- 831 Martín, J., Torrell, M., Korobkov, A. A., and Valles, J.: Palynological features as a systematic marker in *Artemisia* L. and
832 related genera (Asteraceae, Anthemideae) - II: Implications for subtribe Artemisiinae delimitation, *Plant Biol.*, 5,
833 85-93, <https://doi.org/10.1055/s-2001-16462>, 2003.
- 834 McClelland, H. L. O., Halevy, I., Wolf-Gladrow, D. A., Evans, D., and Bradley, A. S.: Statistical Uncertainty in
835 Paleoclimate Proxy Reconstructions, *Geophys. Res. Lett.*, 48, e2021GL092773,
836 <https://doi.org/10.1029/2021GL092773>, 2021.
- 837 Miao, Y. F., Meng, Q. Q., Fang, X. M., Yan, X. L., Wu, F. L., and Song, C. H.: Origin and development of *Artemisia*
838 (Asteraceae) in Asia and its implications for the uplift history of the Tibetan Plateau: A review, *Quatern. Int.*, 236,
839 3-12, <https://doi.org/10.1016/j.quaint.2010.08.014>, 2011.

- 840 Moberg, A., Sonechkin, D. M., Holmgren, K., Datsenko, N. M., and Karlen, W.: Highly variable Northern Hemisphere
841 temperatures reconstructed from low- and high-resolution proxy data, *Nature*, 433, 613-617,
842 <https://doi.org/10.1038/nature03265>, 2005.
- 843 Mosbrugger, V., Utescher, T., and L. D. D.: Cenozoic continental climatic evolution of Central Europe, *Proc. Natl. Acad.
844 Sci. U. S. A.*, 102, 14964-14969, <https://doi.org/10.1073/pnas.0505267102>, 2005.
- 845 Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'Amico, J. A.,
846 Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F.,
847 Wettenberg, W. W., Hedao, P., and Kassem, K. R.: Terrestrial ecoregions of the worlds: A new map of life on Earth,
848 *Bioscience*, 51, 933-938, [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:teotwa\]2.0.co;2](https://doi.org/10.1641/0006-3568(2001)051[0933:teotwa]2.0.co;2), 2001.
- 849 Sánchez-Murillo, R., Durán-Quesada, A. M., Esquivel-Hernández, G., Rojas-Cantillano, D., and Cobb, K. M.:
850 Deciphering key processes controlling rainfall isotopic variability during extreme tropical cyclones, *Nat. Commun.*,
851 10, 4321, <https://doi.org/10.1038/s41467-019-12062-3>, 2019.
- 852 Sanz, M., Vilatersana, R., Hidalgo, O., Garcia-Jacas, N., Susanna, A., Schneeweiss, G. M., and Vallès, J.: Molecular
853 phylogeny and evolution of floral characters of *Artemisia* and allies (Anthemideae, Asteraceae): Evidence from
854 nrDNA ETS and ITS sequences, *Taxon*, 57, 66-78, <https://doi.org/10.2307/25065949>, 2008.
- 855 Shan, B. Q., He, X. L., and Chen, Y. S.: Pollen Morphology of *Artemisia* in the Loess Plateau, *Acta Botanica
856 Boreali-Occidentalia Sinica*, 27, 1373-1379, 2007 (in Chinese).
- 857 Sing, G. and Joshi, R. D.: Pollen Morphology of Some Eurasian Species of *Artemisia*, *Grana Palynologica*, 9, 50-62,
858 <https://doi.org/10.1080/00173136909436424>, 1969.
- 859 Stix, E.: Pollenmorphologische Untersuchungen an Compositen, *Grana*, 2, 41-104,
860 <https://doi.org/10.1080/00173136009429443>, 1960.
- 861 Sun, J. T. and Xu, Y. T.: Pollen morphology and its taxonomic significance of *Artemisia* Linn. from Shandong, *Journal of
862 Shandong Normal University*, 12, 186-190, 1997 (in Chinese).
- 863 Sun, X. J., Du, N. Q., Weng, C. Y., Lin, R. F., and Wei, K. Q.: Paleovegetation and paleoenvironment of Manasi Lake,
864 Xinjiang, N. W. China during the last 14000 years, *Quaternary Sciences*, 14, 239-248, 1994 (in Chinese).
- 865 Sun, X. J., Wang, F. Y., and Song, C. Q.: Pollen-climate response surfaces of selected taxa from northern China, *Sci.
866 China Ser. D-Earth Sci.*, 39, 486-493, 1996.
- 867 Tarasov, P. E., Cheddadi, R., Guiot, J., Bottema, S., Peyron, O., Belmonte, J., Ruiz-Sanchez, V., And, F. S., and Brewer,
868 S.: A method to determine warm and cool steppe biomes from pollen data; application to the Mediterranean and
869 Kazakhstan regions, *J. Quat. Sci.*, 13, 335-344,
870 [https://doi.org/10.1002/\(SICI\)1099-1417\(199807/08\)13:4<335::AID-JQS375>3.0.CO;2](https://doi.org/10.1002/(SICI)1099-1417(199807/08)13:4<335::AID-JQS375>3.0.CO;2), 1998.
- 871 Tierney, J. E., Poulsen, C. J., Montanez, I. P., Bhattacharya, T., Feng, R., Ford, H. L., Honisch, B., Inglis, G. N., Petersen,
872 S. V., Sagoo, N., Tabor, C. R., Thirumalai, K., Zhu, J., Burls, N. J., Foster, G. L., Godderis, Y., Huber, B. T., Ivany, L.
873 C., Turner, S. K., Lunt, D. J., McElwain, J. C., Mills, B. J. W., Otto-Bliesner, B. L., Ridgwell, A., and Zhang, Y. G.:
874 Past climates inform our future, *Science*, 370, eaay3701, <https://doi.org/10.1126/science.aay3701>, 2020.
- 875 Tutin, T. G., Persson, K., and Gutermann, W.: *Artemisia*, *Flora Europaea* 4, Cambridge University Press, Cambridge,
876 178-186, 1976.

- 877 Vallès, J., Garcia, S., Hidalgo, O., Martín, J., and Garnatje, T.: Biology, Genome Evolution, Biotechnological Issues and
878 Research Including Applied Perspectives in *Artemisia* (Asteraceae), *Adv. Bot. Res.*, 60, 349-419, 2011.
- 879 Vrba, E. S.: Evolution, species and fossils-how does life evolve?, *S. Afr. J. Sci.*, 76, 61-84, 1980.
- 880 Wang, F. X., Qian, N. F., Zhang, Y. L., and Yang, H. Q.: Pollen Morphology of Chinese Plants (2nd edition), Science
881 Press, Beijing, 1995 (in Chinese).
- 882 Wang, W. M.: On the origin and development of *Artemisia* (Asteraceae) in the geological past, *Bot. J. Linnean Soc.*, 145,
883 331-336, <https://doi.org/10.1111/j.1095-8339.2004.00287.x>, 2004.
- 884 Wang, Y., Wang, W., Liu, L. N., Jiang, Y. J., Niu, Z. M., Ma, Y. Z., He, J., and Mensing, S. A.: Reliability of the
885 *Artemisia*/Chenopodiaceae pollen ratio in differentiating vegetation and reflecting moisture in arid and semi-arid
886 China, *Holocene*, 30, 858-864, <https://doi.org/10.1177/0959683620902219>, 2020.
- 887 Weng, C. Y., Sun, X. J., and Chen, Y. S.: Numerical characteristics of pollen assemblages of surface samples from the
888 West Kunlun mountains, *Acta Botanica Sinica*, 35, 69-79, 1993 (in Chinese).
- 889 Wodehouse, R. P.: Pollen Grain Morphology in the Classification of the Anthemideae, *Bull. Torrey Bot. Club*, 53,
890 479-485, <https://doi.org/10.2307/2480028>, 1926.
- 891 Wu, F. L., Fang, X. M., and Miao, Y. F.: Aridification history of the West Kunlun Mountains since the mid-Pleistocene
892 based on sporopollen and microcharcoal records, *Paleogeogr. Paleoclimatol. Paleoecol.*, 547, 109680,
893 <https://doi.org/10.1016/j.palaeo.2020.109680>, 2020.
- 894 Xu, Q. H., Li, Y. C., Yang, X. L., and Zheng, Z. H.: Quantitative relationship between pollen and vegetation in northern
895 China, *Sci. China Ser. D-Earth Sci.*, 50, 582-599, <https://doi.org/10.1007/s11430-007-2044-y>, 2007.
- 896 Yang, J., Spicer, R. A., Spicer, T. E. V., Arens, N. C., Jacques, F. M. B., Su, T., Kennedy, E. M., Herman, A. B., Steart, D.
897 C., Srivastava, G., Mehrotra, R. C., Valdes, P. J., Mehrotra, N. C., Zhou, Z. K., and Lai, J. S.: Leaf form-climate
898 relationships on the global stage: an ensemble of characters, *Glob. Ecol. Biogeogr.*, 24, 1113-1125,
899 <https://doi.org/10.1111/geb.12334>, 2015.
- 900 Yi, S., Saito, Y., Zhao, Q. H., and Wang, P. X.: Vegetation and climate changes in the Changjiang (Yangtze River) Delta,
901 China, during the past 13,000 years inferred from pollen records, *Quat. Sci. Rev.*, 22, 1501-1519,
902 [https://doi.org/10.1016/s0277-3791\(03\)00080-5](https://doi.org/10.1016/s0277-3791(03)00080-5), 2003a.
- 903 Yi, S., Saito, Y., Oshima, H., Zhou, Y. Q., and Wei, H. L.: Holocene environmental history inferred from pollen
904 assemblages in the Huanghe (Yellow River) delta, China: climatic change and human impact, *Quat. Sci. Rev.*, 22,
905 609-628, [https://doi.org/10.1016/s0277-3791\(02\)00086-0](https://doi.org/10.1016/s0277-3791(02)00086-0), 2003b.
- 906 Zachos, J., Pagani, M., Sloan, L., Thomas, E., and Billups, K.: Trends, Rhythms, and Aberrations in Global Climate 65
907 Ma to Present, *Science*, 292, 686-693, <https://doi.org/10.1126/science.1059412>, 2001.
- 908 Zachos, J. C., Dickens, G. R., and Zeebe, R. E.: An early Cenozoic perspective on greenhouse warming and carbon-cycle
909 dynamics, *Nature*, 451, 279-283, <https://doi.org/10.1038/nature06588>, 2008.
- 910 Zhang, X. S.: Vegetation map of China and its geographic pattern: Illustration of the vegetation map of the People's
911 Republic of China (1:1 000 000), Geological Press, Beijing, 2007 (in Chinese).

- 912 Zhang, Y., Kong, Z. C., Wang, G. H., and Ni, J.: Anthropogenic and climatic impacts on surface pollen assemblages
913 along a precipitation gradient in north-eastern China, *Glob. Ecol. Biogeogr.*, 19, 621-631,
914 <https://doi.org/10.1111/j.1466-8238.2010.00534.x>, 2010.
- 915 Zhao, Y., Xu, Q. H., Huang, X. Z., Guo, X. L., and Tao, S. C.: Differences of modern pollen assemblages from lake
916 sediments and surface soils in arid and semi-arid China and their significance for pollen-based quantitative climate
917 reconstruction, *Rev. Palaeobot. Palynol.*, 156, 519-524, <https://doi.org/10.1016/j.revpalbo.2009.05.001>, 2009.
- 918 Zhao, Y., Liu, H. Y., Li, F. R., Huang, X. Z., Sun, J. H., Zhao, W. W., Herzschuh, U., and Tang, Y.: Application and
919 limitations of the *Artemisia*/Chenopodiaceae pollen ratio in arid and semi-arid China, *Holocene*, 22, 1385-1392,
920 <https://doi.org/10.1177/0959683612449762>, 2012.
- 921 Zhao, Y. T., Miao, Y. F., Fang, Y. M., Li, Y., Lei, Y., Chen, X. M., Dong, W. M., and An, C. B.: Investigation of factors
922 affecting surface pollen assemblages in the Balikun Basin, central Asia: Implications for palaeoenvironmental
923 reconstructions, *Ecol. Indic.*, 123, <https://doi.org/10.1016/j.ecolind.2020.107332>, 2021.
- 924 Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Ritter, C. D., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn,
925 R., Svantesson, S., Wengstrom, N., Zizka, V., and Antonelli, A.: CoordinateCleaner: Standardized cleaning of
926 occurrence records from biological collection databases, *Methods Ecol. Evol.*, 10, 744-751,
927 <https://doi.org/10.1111/2041-210X.13152>, 2019.