

Lalramliana

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3346968/publications.pdf>

Version: 2024-02-01

33
papers

193
citations

1039880

9
h-index

1281743

11
g-index

33
all docs

33
docs citations

33
times ranked

144
citing authors

#	ARTICLE	IF	CITATIONS
1	Integrative taxonomy peeps a new torrent catfish species of genus <i>Amblyceps</i> (Siluriformes): Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	1.0	2
2	<p>A new country record of Raorchestes cangyuanensis Wu, Suwannapoom, Xu, Murphy & Che 2019 and additional record of Kurixalus yangi Yu, Hui, Rao & Yang 2018 (Anura: Rhacophoridae: Rhacophorinae) from India</p>. Zootaxa, 2021, 4974, 383-390.	0.2	1
3	<i>Badis kaladanensis</i> , a new fish species (Teleostei: Badidae) from Mizoram, northeast India. PLoS ONE, 2021, 16, e0246466.	1.1	2
4	Susceptibility of the fall armyworm, <i>Spodoptera frugiperda</i> (J.E. Smith, 1797) (Lepidoptera: Noctuidae), to four species of entomopathogenic nematodes (Steinernematidae and Heterorhabditidae) from Mizoram, North-Eastern India. Egyptian Journal of Biological Pest Control, 2021, 31, .	0.8	6
5	Ichthyofauna of Dampa Tiger Reserve Rivers, Mizoram, North-Eastern India. Journal of Environmental Biology, 2020, 41, 884-895.	0.2	1
6	Studies on the distribution and diversity of helminth infection in <i>Xenentodon cancila</i> (Hamilton, 1822) in Mizoram, Northeast India. Journal of Environmental Biology, 2020, 41, 832-839.	0.2	0
7	Characterization and screening of antifungal activity of bacteria associated with entomopathogenic nematodes from Mizoram, North-Eastern India. Journal of Environmental Biology, 2020, 41, 942-950.	0.2	4
8	Evaluation of pathogenicity of indigenous entomopathogenic nematodes (Steinernematidae and) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	0.2	1
9	DNA barcoding revealed a new species of <i>Neolissochilus</i> Rainboth, 1985 from the Kaladan River of Mizoram, North East India. Mitochondrial DNA Part A: DNA Mapping, Sequencing, and Analysis, 2019, 30, 52-59.	0.7	9
10	Morphological and molecular characterization of entomopathogenic nematode, <i>Heterorhabditis baujardi</i> (Rhabditida, Heterorhabditidae) from Mizoram, northeastern India. Journal of Parasitic Diseases, 2018, 42, 341-349.	0.4	11
11	<i>Laubuka parafasciata</i> , a new cyprinid fish species (Teleostei: Cyprinidae) from Mizoram northeastern India. Zootaxa, 2017, 4244, 269-276.	0.2	3
12	Characterization of a new isolate of entomopathogenic nematode, <i>Steinernema sangi</i> (Rhabditida,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Mizoram, northeastern India. Journal of Parasitic Diseases, 2017, 41, 1123-1131.	0.4	13
13	A new diminutive sisorid catfish (Actinopterygii: Siluriformes) from northeastern India. Zootaxa, 2016, 4105, 546-56.	0.2	5
14	<i>Physoschistura chhimtuipuiensis</i> , a new species of loach (Teleostei: Nemacheilidae) from Mizoram, north-eastern India. Zootaxa, 2016, 4173, 192.	0.2	2
15	<i>Channa aurantipectoralis</i> , a new species of snakehead from Mizoram, north-eastern India (Teleostei): Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	0.2	9
16	New records of Indian Nematomorpha, with the description of a new species from the genus <i>Chordodes</i> . Zootaxa, 2016, 4158, 272.	0.2	4
17	Effects of storage temperature on survival and infectivity of three indigenous entomopathogenic nematodes strains (Steinernematidae and Heterorhabditidae) from Meghalaya, India. Journal of Parasitic Diseases, 2016, 40, 1150-1154.	0.4	11
18	<p class="HeadingRunIn">Psilorhynchus kaladanensis</p> a new species (Teleostei: Psilorhynchidae) from Mizoram, northeastern India</p>. Zootaxa, 2015, 3962, 171.	0.2	6

#	ARTICLE	IF	CITATIONS
19	<i>Psilorhynchus khopai</i> , a new fish species (Teleostei: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	0.2	6
20	<i>Pethia rutila</i> (Teleostei: Cyprinidae), a new species from Mizoram, Northeast India. Zootaxa, 2014, 3827, 366.	0.2	10
21	<i>Olyra saginata</i> , a new species of bagrid catfish (Actinopterygii: Tj ETQq1 1 0.784314 rgBT /Over	0.2	4
22	<i>Schistura andrewi</i> , a new species of loach (Teleostei: Nemacheilidae) from Mizoram, northeastern India. Zootaxa, 2014, 3860, 253-60.	0.2	3
23	<i>Eutropiichthys cetosus</i> , a new riverine catfish (Teleostei: Schilbeidae) from northeastern India. Journal of Threatened Taxa, 2014, 6, 6073-6081.	0.1	3
24	<i>Schistura maculosa</i> , a new species of loach (Teleostei: Tj ETQq0 0 0 rgBT /Overlock 10 Tf	0.2	11
25	<i>Pseudolaguvia nubila</i> , a new sisorid catfish (Teleostei: Sisoridae) from northeastern India. Zootaxa, 2013, 3647, .	0.2	4
26	<i>Glyptothorax radiolus</i> , a new species of sisorid catfish (Osteichthyes: Siluriformes) from northeastern India, with a redescription of <i>G. striatus</i> McClelland 1842. Zootaxa, 2013, 3682, 501-12.	0.2	6
27	<i>Garra dampansensis</i> , a new ray-finned fish species (Cypriniformes: Cyprinidae) from Mizoram, northeastern India. Journal of Threatened Taxa, 2013, 5, 4368-4377.	0.1	3
28	<i>Pseudolaguvia nubila</i> , a new sisorid catfish (Teleostei: Sisoridae) from northeastern India. Zootaxa, 2013, 3647, 518-26.	0.2	0
29	Efficacy of indigenous entomopathogenic nematodes from Meghalaya, India against the larvae of taro leaf beetle, <i>Aplosomyx chalybaeus</i> (Hope). Journal of Parasitic Diseases, 2012, 36, 149-154.	0.4	15
30	Evaluation of the efficacy of three indigenous strains of entomopathogenic nematodes from Meghalaya, India against mustard sawfly, <i>Athalia lugens proxima</i> Klug (Hymenoptera: Tenthredinidae). Journal of Parasitic Diseases, 2012, 36, 175-180.	0.4	6
31	Soil moisture effects on the activity of three entomopathogenic nematodes (Steinernematidae and) Tj ETQq1 1 0.784314 rgBT /Over	0.4	14
32	<i>Chordodes mizoramensis</i> (Nematomorpha, Gordiida), a new species of horsehair worm from Mizoram, North-East India. ZooKeys, 2011, 75, 1-8.	0.5	11
33	<i>Pseudolaguvia virgulata</i> , a new sisorid catfish (Teleostei: Sisoridae) from Mizoram, northeastern India. Zootaxa, 2010, 2518, .	0.2	7