

Forgotten Books

— www.forgottenbooks.com —

Copyright © 2016 FB &c Ltd.


All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the publisher, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law.

PUBLICATIONS OF THE ARNOLD ARBORETUM, No. 9

A
MONOGRAPH OF AZALEAS

RHODODENDRON SUBGENUS
ANTHODENDRON

BY
ERNEST HENRY WILSON
AND
ALFRED REHDER



ISSUED APRIL 15, 1921

PRINTED FROM THE INCOME OF THE WILLIAM L. BRADLEY FUND

BY
THE UNIVERSITY PRESS
CAMBRIDGE
1921

LANDSCAPE
ARCHITECTURE

Add'l

Farrand Gift

PREFACE

THE value of Azaleas as garden plants in the eastern United States, where few of the Rhododendrons with persistent leaves can be successfully cultivated, has led to the critical studies of these plants which appear on the following pages.

Mr. Wilson's long journeys in eastern Asia has made it possible for him to study all the species of the Japanese Empire and China as wild plants, and to become familiar with the Azaleas growing in Japanese gardens. For many years the Arboretum has been engaged in a field study of the American species, and has been able to place at Mr. Rehder's disposal the large amount of material which is preserved in this herbarium, and which has been supplemented by that contained in other American collections.

The Azaleas of northern Japan and Korea, and of the northern United States, are already growing in the Arboretum, but the gardens of the southern states and of California have still much to gain by the introduction of the species of the southern states and those of southern Japan, China, and Formosa, and this publication will not have accomplished its purpose if it fails to induce the more general cultivation of these plants in the United States and Europe.

The art of the hybridizer has produced many interesting and beautiful Azaleas. Imperfect records have made the study of many of these hybrids difficult and uncertain, and this difficulty is increased by the fact that many of them are no longer cultivated unless, having escaped the change of fashion in plants, they are still growing in English gardens, planted about the middle of the last century, and I venture to suggest to my associates in the English Rhododendron Society that the study of Azaleas in European gardens might add much to the knowledge of the origin and value of many of these plants.

C. S. SARGENT, *Director.*

ARNOLD ARBORETUM
MARCH, 1921.

TABLE OF CONTENTS

	PAGE
PREFACE	v
THE AZALEAS OF THE OLD WORLD. By <i>Ernest Henry Wilson</i> .	1
THE AZALEAS OF NORTH AMERICA. By <i>Alfred Rehder</i>	107
DOUBTFUL NAMES	197
ILLUSTRATIONS	203
ADDITIONS	206
INDEX	207

AZALEAS OF THE OLD WORLD

BY

ERNEST HENRY WILSON

THE AZALEAS OF THE OLD WORLD

INTRODUCTION

BOTANISTS may hold different views on the classification of the subdivisions of the large genus *Rhododendron*, but gardeners and lovers of plants in general have no difficulty in recognizing Azaleas as distinct from other groups of the genus. It is in the popular rather than in the strictly technical sense that the title of "Azaleas of the Old World" is applied here. These plants with few exceptions are sun-loving; and their wide popularity among the peoples of the Orient, of America, of Europe and Australia is due largely to the brilliancy of their flowers, their floriferousness and the ease with which many of them can be cultivated. In the Orient they have been favorite garden flowers from very early times and in the Occident for more than a century some have been familiar and valued greenhouse plants, and in recent years it has been found that several are more hardy than they were generally supposed to be. In the Arnold Arboretum such hardy kinds as *R. obtusum* var. *Kaempferi* and *R. japonicum* are among the most satisfactory and most beautiful of Asiatic plants. Farther south and in California the old *Azalea amoena* and *A. ledifolia* and the newer "Hinodegiri" are much grown in gardens. Others are less known, but I believe that in the near future both in increased variety and in quantity Azaleas will have a much more important place in gardens. In certain groups the hybridists in Europe have wrought wonderful results, and the field is still full of promise. In Japan, in the city of Kurume, selection and raising from seed has been in progress among a single group for a century, and these plants are now beginning to find their way into the Occident. With these facts in view the need for a critical survey of the species and forms seems necessary. Through confusion with other species and for lack of accurate names many good plants have become lost from our gardens, and others for the same reason have not been introduced. During my travels in China I greatly admired

the red-flowered Azalea of the country so abundant on the hills from the coast to the extreme west, from near sea-level to 1300 m. altitude. So common is it that in May much of the countryside is a blaze of red. For years I unhesitatingly accepted this plant as the *Azalea indica* of Linnaeus, considering the difference in the number of stamens an unimportant triviality. In 1912-13, when working up the Rhododendron material for Sargent's *Plantae Wilsonianae* in conjunction with my colleague, Alfred Rehder, I became conscious that all was not so simple as I had assumed. The question as to what *Azalea indica* really was began to arouse my interest. On the occasion of the Arnold Arboretum Expedition to Japan in 1914 I devoted much time to the problem, and in Yaku-shima in February collected an Azalea which later I discovered to be the true *Azalea indica* of Linnaeus. I questioned many botanists in Japan, but none at that time had any definite views on the subject. Nevertheless, by deduction and comparative study I arrived at the conclusion that the group of Azaleas cultivated in Japan under the name "Satsuki" all belonged to Linnaeus' species and that the material collected in Yaku-shima was its wild parent. On my return to the Arnold Arboretum early in 1915 I critically compared the material I had collected with Linnaeus' description and with the old figures on which his species was based, and the proof of this theory was absolute.

Japan is so rich in Azaleas, both wild on the mountains and cultivated in gardens, parks and temple grounds, that at the end of 1914 I felt that beyond establishing the identity of *Azalea indica* I had acquired no more than a moderate working knowledge of the group. My interest, however, was keener than ever. Being favored with another opportunity to visit the Orient I determined to devote as much time as possible to the study of these Azaleas. During 1917 and 1918 I visited many places famous in Japan for Azaleas and gathered specimens and information from all sides. On my visits to Korea, Liukiu and Formosa I collected much new material. The result is that on my various visits to the Chinese and Japanese Empires I have seen growing wild, with few exceptions, every species known from those regions and almost every known variety and garden form. In Japan in 1914 and again in 1917-18 I received inestimable assistance from Mr. H. Suzuki, President of the

Yokohama Nursery Company and one of the foremost horticulturists in Japan. Together we visited Mt. Kirishima and the famous Azalea district of Kurume in Kyushu, also those near Osaka and Tokyo. He introduced me to all the leading specialists, gave of his own store of knowledge freely, and without his help it would have been quite impossible to have delved so deeply into the cultivated Azaleas of Japan. It is with keen pleasure that I acknowledge my indebtedness to this genial and scholarly gentleman. I know not how many hundreds of specimens of my own collecting have been available for this work, and in addition, through the courtesy of the Director of Kew and the Keeper of the Gray Herbarium of Harvard University, I have had on loan all their material, so rich in old types of these Azaleas. In Tokyo, through the courtesy of Professor J. Matsu-mura, I have examined the material preserved in the Herbarium of the Imperial Botanic Garden. Dr. Merrill has kindly loaned me the material preserved in the Herbarium of the Bureau of Science, Manila. To Dr. T. Nakai I am indebted for much useful information relative to certain critical Azaleas to which S. Komatsu has recently given names with brief descriptive notes in Japanese.

The study of long-cultivated garden plants is exceedingly complex, and to fathom some of the problems is almost beyond human skill. This statement is as trite as it is true, and I freely confess that I have found the problems presented by many of these Azaleas more intricate than those of the Japanese Cherries. Notwithstanding my unusual opportunities in the field and in the gardens of the Orient and the abundant herbarium material at my command I should have been ill equipped for the task but for the collection of living plants cultivated for nearly three-quarters of a century on the Sargent estate in Brookline, Mass. This collection has been zealously maintained and is rich in old garden types introduced through Fortune and others, some of which have apparently been lost in Europe. Professor Sargent's knowledge of these Azaleas is profound, and it is under his sympathetic guidance and help that this little treatise has developed from chaos to order, and to him most largely is due what merit it contains.

HISTORY

THE cultivation of plants and the development of gardens is ever a sign of a nation's advance toward culture and refinement. How early this art began in the Orient we do not know, though history tells us that plants like the Yulan and Moutan Paeony were highly esteemed garden flowers among the Chinese during the Tang Dynasty (618–907 A. D.). It is much to be regretted that we are so ignorant of the early history of China. It is known that during the Han Dynasty (B. C. 206–A. D. 25) the Chinese had intercourse with India and Zanzibar. If proof were needed of their visiting these countries at an early period it is supplied by the presence in south China and Formosa of such typical African plants as Cactus-like Tree Euphorbias. Such species as *E. triangularis* Desf., *E. nerifolia* L. and *E. tirucalli* L. are common hedge-plants and are naturalised in certain places like Hongkong, Kowloon and elsewhere in China, and near Takao in Formosa, as they also are in parts of India. In point of fact *E. tirucalli* was named by Linnaeus from material which came from India. Did we know more about the early travels of the Chinese it is possible that we should find that they were acquainted with America long before its discovery by Columbus. In south China, Formosa and the Liukiu Islands *Agave fourcroydes* Lem. is naturalised, but this may have been introduced by the Portuguese, and so too may have been Maize and Tobacco. Much more difficult is it to account for *Opuntia Dillenii* Haw. naturalised and abundant not only on the Yunnan plateaux but in the valley of the Tung River in a remote part of extreme western China. It is significant also that Boym in his *Flora Sinensis* published in 1656 figures such characteristic American plants as the Pineapple (*Ananas sativa* Schult. f.), the Sweetsop or Custard Apple (*Anona squamosa* L.) and the Guava (*Psidium guajava* L.), which were evidently common plants in China when he was there (1642–53). In what era the love for flowers began to manifest itself in Japan is uncertain, but Buddhism in its Chinese form was introduced from Korea about 552 A. D. and from then until the 8th century Korean and Chinese monks and nuns visited Japan for purposes of proselytism. From the 8th century onward it became more usual for the Japanese

monks to go to China in order to study the doctrines of the best accredited teachers at the fountain-head. These monks introduced into Japan a number of favorite trees and shrubs like the Ginkgo, Yulan, Moutan Paeony, Chinese Quince, the Tea-plant, Buddha's Tree (*Tilia Miqueliana*), *Sophora japonica*, *Prunus tomentosa*, *P. japonica*, *Paulownia tomentosa*, such fruit trees as the Peach, Apricot, Persimmon, Pear, Apple, Plum, Cherry, Loquat, the Orange and its relatives and, in all probability, the Chrysanthemum also. Though they probably did not originate the love for gardens these monks must have greatly stimulated it. The mutual love of flowers among the peoples of the Orient induced and fostered the introduction of plants from one country to another in the Far East even as it did in the Levant and, in later days, as it has in the Occident. And so we find the history of garden plants intimately associated with the social life of the peoples. We know nothing of early intercourse between the Japanese and Chinese prior to the introduction of Buddhism to Japan. We do know that Chinese monks introduced their favorite flowers to Korea, that Chinese and Korean monks carried them to Japan, and it is safe to assume that these same monks and their Japanese disciples carried to Korea and to China plants of Japan which appealed to their aesthetic sense. In fact the presence in temple-grounds and other sacred places in China of the *Cryptomeria*, which is endemic in Japan, and of many varieties of Japanese Camellias and Azaleas in Chinese gardens, is positive proof that Japanese plants were long ago introduced into China.

The illustrious Venetian, Marco Polo, who lived many years in China during the latter part of the 12th century and to whom we owe the first authentic account of that mighty empire, notices from his own observation many of the vegetable productions of China used for economic purposes. After his memorable travels China was for a long time closed to European access and had been altogether forgotten in Europe when in 1516 the Portuguese first arrived in China. The Portuguese claim the honor of having first introduced the Sweet-Orange from China to Portugal some time between 1545 and 1548. About the end of 1542 the Portuguese adventurer, Mendez Pinto, discovered Japan, first touching at the island now known as Tanagashima south of Kagoshima in Kyushu. The Portuguese made their way north as far as Oita in northeast Kyushu where Mendez Pinto in 1543 received a friendly reception from the local Daimyo. The arquebus Pinto carried astounded the Japanese, who had never before seen any explosive weapon. In 1549 St. Francis Xavier reached Japan

and introduced Christianity; he visited Oita, and going north to Hondo established a mission at Yamaguchi, a few miles north of Shimonoseki. This newly introduced religion soon claimed thousands of adherents, much to the alarm of the Japanese Daimyos and military men backed by Buddhist priests, and a period of persecution commenced. The native Christians migrated to Nagasaki, which soon became one of the chief marts of Portuguese trade.

In 1596 the Dutch first visited Java and other islands of the East Indies and in 1602 the Dutch East India Company was established. The war which then ensued between the Dutch, Spaniard and Portuguese for possession of the spice islands lasted until 1610 when the Dutch remained master of these seas. The seat of the Dutch government was first established on the island of Amboyna, but in 1619 it was transferred to the newly founded city of Batavia in Java. In 1600 a Dutch ship visited Japan, and nine years later the Dutch East India Company sent several vessels to Firando (Hirado), northwest of Nagasaki, where they were well received by the Japanese. In 1611 a formal edict in favor of their trade was obtained. A Dutch factory and also an English factory were established at Hirado the same year.

In 1624 Christianity was proscribed in Japan and the land was closed to foreigners. The Portuguese, English and Spaniards, who also had a factory, were expelled and only the Dutch and the Chinese were permitted to carry on trade, and that under galling restrictions. The Dutch factory was removed in 1639 to Deshima, then an islet at the head of Nagasaki harbor but now absorbed into the foreshore. And so Japan secluded herself for over two centuries from contact with the outer world until the advent of the American Expedition in 1853-54 under command of Commodore Perry. All this may seem to have little to do with plants and with Azaleas in particular and yet they are inextricably bound up with the subject. It is to the merchant adventurer that we owe all our first plant introductions from Japan and our early knowledge of the flora. From 1690-92 Engelbert Kaempfer, in the service of the Dutch East India Company, lived in Japan, and in his *Amoenitates Exoticae*, published in 1712, he gives an admirable account of Japanese plants. He gives good figures of many of these under the vernacular names and among them an Azalea now known as *R. obtusum* var. *Kaempferi* Wils. In all Kaempfer mentions twenty-one Azaleas, and it is interesting to note that many of the vernacular names he gives are in common use to-day. C. P. Thunberg in 1775 visited Japan in the capacity of physician to the Dutch Com-

pany and in 1784 published his *Flora Japonica*, the first post-Linnaean work on the flora of the Orient. Thunberg mentions the Azaleas under their vernacular names and relegates them all to *Azalea indica*. According to Juel (*Pl. Thunb.* 391 (1918)) they represent *R. obtusum* Planch., *R. japonicum* Suring. and *R. mucronatum* G. Don.

Lastly in this connection mention must be made of Philipp von Siebold, who joined the service of the Dutch East India Company and lived at Deshima from 1823 to 1829. To him we owe the magnificent *Flora Japonica*; also he introduced about 1830 such ornamental plants as *Lilium elegans* and *L. speciosum* from Japanese gardens, and, twenty years later, several others including *Malus floribunda*, *M. Sieboldii* and certain flowering Cherries.

About 1677 the English East India Company established a factory at Amoy and in 1684, after a conflict with the Portuguese, one at Canton. Through this company many Chinese plants were introduced to India during the 18th century and early in the 19th century to England; they include several Azaleas. However, in China the Jesuit priests were the first to inform us about the flora of the land. In 1790 Joannis de Loureiro, a Portuguese, published his *Flora Cochinchinensis* and mentions one species of Azalea (*A. punctata*), but this remains an obscure plant and may not belong to the genus.

During the last quarter of the 17th century Jakob Breyne, a merchant of Danzig and a distinguished botanist, visited Holland and saw growing there several famous garden plants of the Orient which he duly records in his *Prodromus Plantarum*. On page 24, pt. I, published in 1680, we read "Chamaerhododendron exoticum amplissimus floribus liliaceis. Frutex spectabilis elegans. In horto Beveringiano." This is the plant on which Linnaeus (*Spec.* 1753) bases his *Azalea indica*. Breyne does not say from what country this shrub had been brought, but P. Hermann, in his *Academici Horti lugduno-batavi catalogus*, p. 152 (1687), describes the same plant under the name of "Cistus Indicus Ledi alpini foliis, floribus amplis," figures it on page 153, and reports that it was introduced from Jaccatra, which is Batavia, in Java. No species of Azalea grows in India, and it has been generally supposed that this Azalea had been brought to Batavia by the Chinese, but I am of the opinion that it was taken there from Japan by Dutch traders. In the gardens of Nagasaki and elsewhere, then as now, it was a common garden plant and, moreover, it is indigenous in the warm southern parts of Japan. That at the early date mentioned plants from Japan were growing in gardens in Holland is certain,

and the Tea-plant was introduced from Japan about 1689. Breyne (*Prodromus*, I. 4) states that he saw in the garden of Hieronymus van Beverningk in 1678 a fine specimen of the Camphor-tree which had been introduced from Japan. Again, in his *Prodromus*, II. 66 (1689) Breyne tells of six varieties of Chrysanthemum from Japan which he saw growing in Holland gardens. All these early introductions seem to have been lost. I have carefully looked through many old books of the 18th century and I find no references to any species of Azaleas being in European gardens, and they appear to have been unknown in Europe at the close of the 18th century. The one oriental species was known to Linnaeus through the books. In Batavia, however, several were grown, as we know from the writings of Burmann and Blume.

In Aiton's *Hortus Kewensis* (1810) *Azalea indica* is stated to be in cultivation in Kew, having been introduced from China in 1808 by the Court of Directors of the East India Company in the ship "Cuffnells" Captain Wellbank. This plant was *Rhododendron Simsii* Planch., and here began a confusion between Linnaeus' species and the Chinese Azalea which still survives. Through the activities of officers of the East India Company at the close of the 18th and the beginning of the 19th centuries English gardens were enriched by many Chinese plants. No officer was more active in this work than John Reeves, who as Chief Inspector of Tea in the Company's establishment at Canton resided in Macao and Canton from 1812 to 1831. He was either the immediate or indirect source from which our gardens derived Camellias, Moutan Paeonies, Chrysanthemums, Roses, *Wistaria sinensis* and many Chinese Azaleas including *R. Farrerae* Tate, a pretty Hong-kong species, and also *R. phoeniceum* G. Don, destined to be of so much value as a stock on which to graft "Indian Azaleas," so called.

In 1818 or 1819 Samuel Brooks, a nurseryman at Ball's Pond, Newington Green, sent out to China Joseph Poole, a gardener, and through him secured among other plants *Azalea indica alba* (*R. mucronatum* G. Don) and *Azalea indica purpurea plena* (*R. mucronatum* f. *plenum* Wils.). In 1823 *Azalea sinensis* (*R. molle* G. Don) was received from China by Loddiges of Hackney and by William Wells, nurseryman, of Redleaf. In 1832 Joseph Knight, nurseryman, of Chelsea, London, secured *Azalea indica variegata* (*R. indicum* f. *variegatum* DC.) and *A. indica lateritia* (*R. indicum* Sweet) brought home from China by Mr. M'Killigan. In 1821 *Rhododendron Farrerae* was reintroduced and in 1832 *R. indicum* var. *ignescens* Sweet, both through Captain Farrer of the East Indiaman "Orwell," who gave

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

seeds of Azaleas and by this means introduced into Petrograd *R. reticulatum* G. Don and *R. linearifolium* var. *macrosepalum* Mak. Maximowicz also introduced plants of *R. linearifolium* S. & Z., its variety *macrosepalum* and *R. scabrum* G. Don. In 1861 a Hollander, J. B. Groenwegen, claims to have received direct from Japan seeds of *R. japonicum* Suring., one of the most valuable of all hardy Azaleas. Crossed with the Chinese *R. molle* G. Don this has given rise to the lovely but short-lived Kosterianum hybrids; all these, again mixed with the so-called Ghent hybrids (*R. luteum* × American species), have produced the race of Azaleas named × *R. mixtum* Wils.

Under each species and variety the story of its introduction is recorded, so in this historical chapter I need not repeat it. Curiously enough the most common and one of the most conspicuously beautiful Azaleas of Japan, with flame-red flowers and figured by Kaempfer, was ignored until 1892, when Professor Sargent sent seeds to the Arnold Arboretum where it has proved not only perfectly hardy but one of the most valuable exotic plants. This and *R. japonicum*, raised by Professor Sargent from seeds sent from the Jardin des Plantes in 1876, were the first oriental Azaleas to be raised from seeds in America.

Turning to Korea there is little to say, for prior to the beginning of this century the Occident had received no living plants or seeds direct from that country. A common Azalea of Korea, the lovely *R. Schlippenbachii* Maxim., found its way to Japan, and from there was sent to England by James H. Veitch in 1893. The first seeds of an Azalea sent from Korea was in 1905 when Mr. J. G. Jack sent to the Arnold Arboretum seeds of *R. yedoense* var. *poukhanense* Nakai, a hardy species of compact habit with rose-purple, fragrant flowers.

From the Liukiu Islands no Azaleas, either plants or seeds, have been sent direct to the Occident, but the large-flowered *R. scabrum* G. Don, better known as *R. sublancheolatum* Miq., endemic there, was long ago brought to Japan, and in Kyushu in particular it is a common garden plant. This Azalea was apparently in cultivation in England when G. Don wrote his *General System* in 1834, but seems to have been lost soon afterward. Maximowicz in 1864 sent it to Petrograd, but it appears to have been unknown in England when reintroduced by Mr. Notcutt in 1911. The only Formosan Azalea introduced into Europe prior to my visit in 1918 was *R. Oldhamii* Maxim. which was raised by Messers Veitch from seeds sent in 1878 by Charles Maries. The Philippine Island species has not yet been introduced.

DISTRIBUTION

OF the thirty-four species of Azaleas enumerated here thirty-three are widely scattered over the countries of the Orient having their northern limit in central Hokkaido, their southern on Luzon in the Philippines; from east to west they are found from the shores of the Pacific Ocean in Japan to the extreme west of China proper. They belong to four well-marked sections of the genus *Rhododendron*, two of which (*Tsutsutsi* and *Sciadorhodion*) are peculiar to the Orient; two (*Rhodora* and *Pentanthera*) are represented also in America. The thirty-fourth species is *R. luteum* Sweet which belongs to the last-named section. It grows in the Caucasus, the Pontus region of the Black Sea and in Lithuania, Poland, and southwestern Russia, and is the only species of the groups here discussed known to grow west of China proper.

Geographically the east Asiatic species are found over 45 degrees of longitude and 30 degrees of latitude and altitudinally from sea-level to about 3300 m. They are social plants and where climate and soil conditions are favorable, as in most parts of Japan, of Korea and of China from about the latitude of Shanghai south, these Azaleas are perhaps the most common shrubs. The Chinese *R. Simsii* and *R. Mariesii* grow also in *Formosa*, and a variety of the first-named is endemic on the Kawanabe Islands off the northwest coast of the Liukiu Archipelago. The Japanese *R. serpyllifolium* is said to grow on Okinawa, the main island of the Liukius. At least three (*R. Schlippenbachii*, *R. Tschonoskii* and *R. Weyrichii*) of the four Korean species grow also in Japan. One (*R. phoeniceum*) is unknown in a wild state and is of doubtful standing. The others are endemic in the different countries of the Orient.

It is remarkable that in western China only the section *Tsutsutsi* with three species is represented, and on the Himalayas no member of the four sections is known to grow, yet in these two regions the subgenus *Eurhododendron* has its headquarters and is represented by hundreds of species. In Malaysia, too, where the subgenus *Vireya* is so richly represented, no species of our subgenus is known. The great concentration of species is in Japan. *Formosa*, considering its small

size, is also rich in Azaleas. In China these Azaleas are poorly represented by eight species of which only three are really common and widespread. This is singular in view of the remarkable richness of the Chinese flora in general. In Korea four species grow, but at most only one (*R. yedoense* var. *poukhanense*) is endemic and specimens of an Azalea much like this have been collected in Bitchu province, western Japan. In Japan Azaleas reach their northern limits in central Hokkaido, where *R. Tschonoskii*, *R. Albrechtii*, *R. obtusum* var. *Kaempferi* and *R. reticulatum* are found. The marked absence of limestone in Japan is favorable to the family Ericaceae which is richly represented there, and on the volcanic mountains Azaleas are astonishingly abundant. The small-flowered *R. obtusum*, the wild type of which is the form *japonicum*, is local, being confined to Kirishima and to a few other mountains in Kyushu, but the large-flowered variety *Kaempferi* grows in plenty from the extreme south, northward to central Hokkaido. The same is true of the magenta-flowered *R. reticulatum*. Billions of plants of these two species are scattered on the grass and shrub-clad slopes, in thickets, in thin woods and on the margins of forests. In May from Kyushu to central Hondo travelling by train or along the roads one is never out of sight of the blossoms of these Azaleas; in June the same is true for the north until their limit of distribution is reached. Many districts are famous for their Azaleas, and of these I may mention the slopes of Mt. Fuji, the Hakone mountains and the Nikko region. In Yamato province, and also round Fuji and Nikko *R. japonicum* with its clusters of large flame-colored flowers grows in millions. Indeed it is to her Azaleas that Japan owes much for the color which decks her countryside. And what joy it is to ramble through this flower-clad land in spring and early summer. In autumn the leaves of such species as *R. japonicum*, *R. Tschonoskii*, *R. reticulatum*, *R. quinquefolium*, *R. pentaphyllum* and *R. linearifolium* change to rich shades of yellow, orange, crimson and vinous purple. These autumn tints, like the wealth of brilliantly colored flowers in spring and early summer, arrest attention on every side, and small wonder that the Japanese have taken Azaleas into their gardens, parks and temple grounds and cultivated them from time immemorial. The only truly woodland species are *R. quinquefolium* and *R. pentaphyllum*, but so adaptable to different conditions are *R. reticulatum* and *R. obtusum* var. *Kaempferi* that they too may be considered woodland plants. In dark ravines, by the sides of torrents and in the forests of the Nikko region *R. quinquefolium* is plentiful, lighting up the dark and

solemn depths with its snow-white, almost bell-shaped flowers. Crowning a knoll among mixed trees round the Hashimoto Tea House at Chuzenji and high up on the steep, wooded slope behind the Lakeside Hotel near the same place *R. pentaphyllum* is common, and I shall never forget the day in June, 1914, when I first beheld these bushes covered with masses of bright rose-pink blossoms. These two may be classed as small trees and among the Azaleas of eastern Asia are only approached in size by the pink-flowered *R. Schlippenbachii*.

R. Tschonoskii has the smallest flowers of any species of Azalea and, indeed, in this respect is only approached by the Chinese *R. Seniavinii*. But in size of plant the real dwarf species is *R. serpyllifolium*, which is common in the Idzu province, and is also known from Mt. Unzen near Nagasaki, and from Okinawa in the Liukiu Archipelago. Yet individual plants of *R. obtusum* f. *japonicum* at alpine limits of vegetation are equally dwarf.

By the sides of rock-strewn torrents high up on mountains on Yaku-shima Island *R. indicum*, the first-known of all Asiatic Azaleas, forms dense thickets more than a metre high, but in the gardens of Japan it is a dwarf and hugs the ground closely. The lovely *R. mucronatum* with its pure white, fragrant flowers has not been found wild though its pale purple-flowered variety *ripense* is common by the sides of certain rivers in the island of Shikoku. The pink to rose-colored *R. linearifolium* var. *macrosepalum* with sticky leaves and flower-buds and an extraordinarily variable and glandular calyx is abundant in thin Pine-woods and among scrub round Futagawa, Yoshino and near Osaka. Very remarkable is the distribution of the red-flowered *R. Weyrichii*, which is known only from parts of Shikoku in the east, from the Goto Islands in the Japan Sea, and from the island of Quelpaert off the south of Korea. The small-leaved *R. tosaense* is confined to parts of the island of Shikoku, and *R. Schlippenbachii*, which is really a Korean species, grows sparingly on Chokai-san and one or two other mountains in northern Hondo. The slender-branched *R. Albrechtii* grows in thickets and on the margins of forests from the Nikko region northward to central Hokkaido, but is nowhere really common. On the foot-hills of Adzuma-yama the remarkable *R. nipponicum*, with its campanulate white corolla, shreddy, cinnamon-brown bark and fragile fruit, is common in open grass and scrub-clad country, but is not known elsewhere. On Mt. Kirishima in Kyushu *R. obtusum* f. *japonicum* grows under alpine conditions, but the most truly alpine species of east Asiatic Azaleas is *R. Tschonoskii*, widespread in Japan

on high mountains from Shikoku and central Hondo north to central Hokkaido, and in Korea found on the summit of the Chiri-san. All in all the most widely spread, most variable, most spectacularly beautiful and most useful of the red-flowered Japanese Azaleas is *R. obtusum* and its variety *Kaempferi*.

In Korea, from Chiri-san in the extreme south of the peninsula to fifty miles north of Gensan, grows *R. Schlippenbachii*. It is partial to rocky places and thickets and often in thin woods it is the dominant undergrowth. On the famous Diamond Mountains in northeastern Korea it is particularly plentiful in mixed, rather open woods and on cliffs; in early summer its myriads of blossoms literally form sheets of pure pink for miles and miles. On exposed cliffs the habit is stiff and sturdy, but in the shade of trees the branches are slender. On Chiri-san it is usually a tree-like bush, four metres tall, and it is on this mountain range that this Azalea reaches its maximum size. On bare, grass-clad and shrub-clad slopes and in Pine-woods from the latitude of Seoul southward to the island of Quelpaert *R. yedoense* var. *poukhanense* abounds. In full exposure it often forms low mats several yards wide, but among shrubs it is occasionally 1.5 m. tall. This and *R. Weyrichii* are the only species that grow on Quelpaert, and the last-named is unknown on the mainland of Korea.

In China the red-flowered *R. Simsii* is wild from the coast to the extreme west, the valley of the Yangtze River being the northern limit of its range. In many places it is extraordinarily abundant, and in May whole hillsides are red with its flowers. This species is wild in the extreme south of Formosa, and a well-marked variety (*eriocarpum*) is endemic on the Kawanabe Islands off northern Liukiu. This is the only Chinese species known to grow wild outside of China proper, but phylogenetically it and *R. obtusum* may be considered of common origin, and to these China and Japan owe the wealth of brilliant red color, so dominant a feature in their spring landscapes. The lovely yellow-flowered *R. molle*, better known under Sweet's name of *R. sinense*, is found on open hillsides from the coast near Ningpo to Ichang in central China. It is a social plant though local in distribution. *R. Mariesii* grows in thickets and thin woods from Fokien to western Hupeh and also in Formosa, but its near relative *R. Farrerae* is confined to Hongkong and a few places in the neighboring province of Kwantung. On open mountains in western Yunnan the small-flowered *R. microphyton* is abundant; this is the most alpine of the Chinese species and, with the little known *R. atrovirens*, is the most western of

the section Tsutsutsi. The other two Chinese species are *R. Seniavini* and *R. Mariae*, and these are apparently rare and confined to the east-central and south-central provinces.

In Liukiu the gaudy-flowered *R. scabrum* is endemic, but was long ago taken to the gardens of southern Japan where it is a common plant to-day. This species has the largest flowers of all Azaleas, and in a wild state and in gardens where it flowers freely is strikingly conspicuous. Unfortunately in the Occident its tendency is to flower sparingly. I strongly suspect that *R. phoeniceum*, unknown as a wild plant though introduced from Canton about 1824, and the popular stock for grafting the "Indian Azaleas" of western gardens, is specifically not really distinct from *R. scabrum*.

Although eight species grow wild in Formosa, Azaleas cannot be called a prominent feature of the vegetation. The red-flowered *R. Oldhamii* is widespread, and in districts in the north like Mt. Daiton and round Lake Candidius in the centre of the island it is fairly abundant. On grass-clad mountain slopes east of Ari-san the pretty, pink-flowered *R. rubropilosum* is plentiful. The others, so far as my personal knowledge goes, are local if not actually rare plants. Lastly, on Luzon in the Philippine Islands grows *R. subsessile*, the most southern representative of the group.

The distribution and the number of the species in the four groups are shown in the following table:—

	Tsutsutsi	Sciadorhodion	Rhodora	Pentanthera
Japan	8	4	3	1
Korea	2	2	—	
Liukiu	3	—		
Formosa	7	1		—
China	5	2		1
Philippines	1	—		—
Pontus region, Caucasus, Lithuania, Poland and southwestern Russia	—	—		

CLASSIFICATION

MANY of our favorite garden plants of oriental origin came to us from the gardens of China and Japan and in some instances their wild prototypes have not yet been discovered. Often when the wild plant has been discovered it has been found to differ greatly from that already in cultivation, and in consequence has to be designated as a variety or form of the described garden type. Phylogeny and taxonomy are often opposed, which is regrettable but unalterable. Students of the floras of the Orient, particularly those of China and Japan, cannot too strongly bear in mind the fact that a great many of the earliest described species are based on cultivated plants. Let me illustrate this by reference to Japan. The Japanese, like the Chinese and Koreans, are lovers of the odd and curious in flowers and trees for which they have an innate love. Nearly every house, from the poorest hovel to the castle of the feudal lord or his descendants, boasts a few flowers or a garden of some sort. The Japanese love their mountains, many of which are sacred and on most of them shrines have been raised. Pilgrimages to these mountains and shrines are a passion with the people and as mementoes of their visits it is a common practice, and has been for centuries, to bring back some strange or curious plant for their gardens. The result is that Japanese gardens are full of odd and curious plants — dwarf plants, weeping plants, fastigate plants, plants with variegated and curiously incised foliage, plants with abnormal flowers. This love of flowers has been fostered by Buddhist priests, and this has led to mutual exchange of plants among the peoples of China, Korea, Japan and Liukiu. Now if it be properly realised that it was from such gardens — really museums of oddities — that our early plants were taken and names applied to them the difficulty of identifying them with their wild parents will be better understood and appreciated.

Until after 1860 all the Azaleas introduced from the Orient were from gardens, chiefly those of Batavia, Canton and Shanghai. As the plants flowered they were given specific and varietal names, but no attempt to discover their origin was made. Some were placed under the genus *Azalea* and others under that of *Rhododendron*. Salisbury

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

in *Flore des Serres*, IX. (1854), emends Don's name to *Tsutsia*, includes under it *R. molle* and *R. Farrerae*, and adds *R. Championae* Hook. In the last-named species the flowers develop from clustered, axillary buds, and it belongs to the section *Choniastrum* Franchet. Maximowicz emends the section; calls it *Tsutsia*, but keeps under it *R. Championae*. He also emends Planchon's subgenus *Azalea* and employs it as a section to include *R. Farrerae*, *R. molle* and others; he also places in it *R. macrosepalum*, which obviously belongs to the section *Tsutsutsi*.

Of the new section *Sciadorhodium*, *R. Schlippenbachii* may be taken as the type, and under it we place *R. reticulatum* and *R. Farrerae*, referred by G. Don to his section *Tsutsutsi*, also *R. Weyrichii* which Maximowicz includes in his section *Azalea*. Together these form a very homomorphic group. The type of G. Don's section *Pentanthera* is the Eurasian *R. luteum* Sweet, but as most of the species are American I leave the discussion of it and also of the section *Rhodora* G. Don, the type of which is American though a majority of the species are Japanese, to my colleague Alfred Rehder.

KEY TO THE ASIATIC SECTIONS OF THE SUBGENUS

ANTHODENDRON ENDL.

Flowers and leaves from the same bud.

Shoots with flattened, appressed, bristle-like hairs; leaves persistent or sometimes deciduous, usually dimorphic, scattered on the branches, elliptic to lanceolate or oblanceolate I. *Tsutsutsi*.

Shoots glabrous or villose, without bristle-like hairs; leaves deciduous, not dimorphic, in whorls at the end of the branchlets, scattered only on vigorous shoots, usually rhombic to obovate, rarely ovate . . . II. *Sciadorhodion*.

Flowers from the terminal bud, leaves from lateral buds below.

Corolla rotate-campanulate; stamens 8 to 10 III. *Rhodora*.

Corolla funnel-form; stamens 5. IV. *Pentanthera*.

KEY TO THE ASIATIC SPECIES

SECT. I. TSUTSUTSI G. DON

Bud-scales not viscid; shoots densely clothed with flattened, appressed hairs.

Corolla funnel-form-campanulate.

Stamens 5 (5 to 8 in *R. tosaense*).

Style villose at base *R. breviperulatum*, p. 28.

Style glabrous at base.

Leaves linear-lanceolate to lanceolate or oblanceolate.

Leaves crenate-dentate; flowers red to rose-red, more than 2.5 cm. across. *R. indicum*, p. 22.

Leaves entire; flowers lilac-purple, less than 2 cm. across

R. tosaense, p. 52.

Leaves oval or obovate to lanceolate.

Leaves more than 1 cm. long, the midrib beneath covered with hairs and usually with scattered hairs on the whole under surface

R. obtusum, p. 29.

Leaves 1 cm. to 1 cm. or less long, glabrous beneath, except a few brown hairs on the midrib *R. serpyllifolium*, p. 53.

Stamens 7 to 10.

Style villose at base.

Leaves narrow-lanceolate to oblanceolate, crenate-dentate; stamens 10 *R. Kanehirai*, p. 28.

Leaves oval to obovate or lanceolate, entire; stamens 7 to 10.

Flowers lilac to magenta; leaves generally elliptic-oblong, acute

R. sessile, p. 51.

Flowers pink.

Leaves conspicuously dimorphic, obovate to obovate-oblong, abruptly mucronate, 2 to 4.5 cm. long *R. lasiostylum*, p. 50.

Leaves not conspicuously dimorphic, often widest below the middle, elliptic-ovate to ovate-oblong or elliptic to elliptic-oblong, acute *R. rubropilosum*, p. 52.

Style glabrous.

Flowers more than 2.5 cm. across; leaves elliptic to elliptic-oblong or oblong-obovate, from 2 to 5 cm. long *R. Simsii*, p. 45.

Flowers less than 2.5 cm. across.

Leaves lanceolate, acuminate *R. atrovirens*, p. 50.

Leaves oval to oblanceolate, 1 cm. or less long . *R. Nakaharai*, p. 54.

Corolla with short, rarely long, cylindric tube and spreading lobes; stamens 5, rarely 4.

Style villose at base; flowers white to rose-color *R. Seniavinii*, p. 57.

Style glabrous.

Leaves persistent; flowers 5-partite.

Corolla rose-purple, leaves not conspicuously dimorphic, oval to lanceolate *R. microphytum*, p. 57.

Corolla lilac-colored, with a long tube; leaves conspicuously dimorphic, lanceolate or elliptic to oblong-obovate *R. Mariae*, p. 58.

Leaves deciduous; flowers white, 4 to 5 partite; leaves narrow-lanceolate to ovate-lanceolate or oblong-obovate, often prominently 3-nerved

R. Tschonoskii, p. 55.

Bud-scales viscid on inner surface; corolla wide funnel-form; calyx-lobes green, ample; style glabrous.

Shoots densely clothed with flattened, appressed hairs; leaves conspicuously dimorphic; stamens 10.

Leaves persistent, calyx-lobes glandular-ciliate.

Corolla scarlet to rose-red; calyx-lobes oval to lanceolate; leaves subcrenulate, coriaceous *R. scabrum*, p. 59.

Corolla rose-purple to magenta; calyx-lobes lanceolate; leaves entire, subcoriaceous *R. phoeniceum*, p. 61.

Leaves deciduous or semipersistent, obscurely crenate-serrate, veins impressed above; corolla rose to rose-purple, fragrant, often double; calyx-lobes lanceolate, not glandular. *R. yedoense*, p. 64.

Shoots with few appressed, flattened hairs and many spreading, pilose often glandular hairs.

Stamens 8 to 10; leaves persistent or semipersistent, not conspicuously dimorphic.

Corolla red, not viscid; stamens 10; ovary glandular; leaves not viscid beneath *R. Oldhamii*, p. 66.

Corolla white or pale-purple, viscid outside; stamens 8 to 10; ovary glandless; leaves viscid beneath *R. mucronatum*, p. 68.

Stamens 5; leaves deciduous or semipersistent, lanceolate to oblanceolate-ovate or linear-lanceolate; corolla pink to pale rosy purple, viscid without, funnel-form or divided into linear segments; ovary glandular

R. linearifolium, p. 74.

SECT. II. SCIADORHODION REHD. & WILS.

Leaves broadest below the middle; stamens nearly equal.

Leaves ovate; corolla pale lilac-purple; stamens 8 to 10; fruit conic-ovoid on a curved pedicel *R. Farreræ*, p. 79.

Leaves ovate to ovate-lanceolate; corolla rose-color; stamens 10; fruit short-cylindric, erect *R. Mariesii*, p. 80.

Leaves broadest about or above the middle.

Leaves rhombic; fruit cylindric.

Stamens subequal; corolla red, rotate-funnel-form; style villose

R. Weyrichii, p. 82.

- Stamens unequal, sometimes declinate; corolla rotate-campanulate.
 Corolla rose-purple to magenta; style glabrous, villose or lepidote or both;
 leaves with long scattered hairs when young, glabrescent at maturity;
 fruit cylindrical, furrowed *R. reticulatum*, p. 83.
 Corolla white; style glabrous; leaves densely ciliate, short-villose, at least
 on midrib at maturity; fruit cylindrical, terete *R. quinquefolium*, p. 87.
 Leaves broadly obovate; corolla pink; style glandular; fruit oblong-ovoid,
 verruculose *R. Schlippenbachii*, p. 88.

SECT. III. RHODORA G. DON

- Corolla rotate-campanulate.
 Leaves obovate to oblong-ob lanceolate, pubescent below; flowers 3 to 6, ma-
 genta; fruit conic-ovoid, bristly glandular *R. Albrechtii*, p. 91.
 Leaves elliptic to narrow-elliptic, glabrescent; petiole bearded; flowers 1 or 2,
 bright rose-pink; fruit spindle-shape, glabrous, verrucose
R. pentaphyllum, p. 92.
 Corolla campanulate, white; leaves obovate to panduriform, subsessile with scat-
 tered, bristly hairs; fruit oblong-ovoid, verruculose, with thin valves, wavy
 at the margin when open *R. nipponicum*, p. 93.

SECT. IV. PENTANTHERA G. DON

- Corolla wide-funnel-form, non-glandular.
 Stamens as long as the yellow corolla; leaves softly pubescent, often canescent;
 winter-buds velutinous *R. molle*, p. 95.
 Stamens shorter than the orange- or flame-red, rarely yellow corolla; leaves
 glabrous except on the veins beneath; winter-buds glabrous
R. japonicum, p. 99.
 Corolla tubular with spreading lobes, glandular *R. luteum*, p. 103.

ENUMERATION OF THE SPECIES

SECT. I. TSUTSUTSI G. DON

Rhododendron section **Tsutsutsi** G. Don, *Gen. Syst.* III. 845 (1834), excluding Nos. 30, 31, 36, 39.

Azalea Linnaeus, *Spec.* 150 (1753), as to species No. 1.

Tsutsutsi Adanson, *Fam. Pl.* II. 164 (1763).

Loiseleria Reichenbach, *Fl. Germ. Exc.* I. 416 (1830).

Beverinckia Salisbury ined. ex De Candolle, *Prodr.* VII. pt. 2, 726 (1839).

Rhododendron section *Toutousi* Duchartre in Orbigney, *Dict. Univ. Hist. Nat.* XI. 120 (1849).

Rhododendron subgenus *Tsutsia* Planchon in *Fl. des Serr.* IX. 75 (1854); in *Rev. Hort.* 1854, 45.

Rhododendron section *Tsutsugi* Miquel, *Fl. Ind. Bat.* II. 1059 (1857).

Rhododendron section *Tsusia* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 32 (*Rhodod. As. Or.*) (1870), excluding No. 17.

Azalea subgenus *Tsutsutsi* K. Koch, *Dendr.* II. pt. 1, 171 (1872), excluding species No. 6.

Shrubs with dense twiggy branches; shoots densely or sparsely clothed with appressed, flattened, red-brown or rufous hairs, occasionally also with spreading soft hairs. Flowers from terminal bud, with leafy shoots from axils of lower scales of same bud. Leaves evergreen or half evergreen, rarely deciduous, more or less villose, rarely glandular. Corolla funnel-form, campanulate-funnel-form, rarely cylindric, with short spreading lobes; stamens 5 to 10, equal or unequal in length, never declinate, puberulous, or pubescent; ovary strigose, sometimes glandular. Fruit conic or conic-ovoid, glabrescent or strigose-villose, 5-celled; seeds wingless.

This section is confined to eastern Asia, where it is represented by 22 species.

Rhododendron indicum Sweet, *Brit. Flow. Gard.* ser. 2, II. sub. t. 128 (1833). — G. Don, *Gen. Syst.* III. 845 (1834), excluding varieties. — De Candolle, *Prodr.* VII. pt. 2, 726 (1839), in part. — Siebold & Zuccarini in *Abh. Akad. Münch.* IV. pt. III. 131 (*Fl. Jap. Fam. Nat.* 7) (1846), in part. — Miquel, *Fl. Ind. Bat.* II. 1059 (1857). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 37 (*Rhodod. As. Or.*) (1870), in part. — Franchet & Savatier, *Enum. Pl. Jap.* I. 291 (1875), in part. — Schneider, *Ill. Handb. Laubholz.* II. 506, figs. 331 1-m, 332 a-d (1911). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916). — Komatsu in *Tokyo Bot. Mag.* XXXII. [18] (1918).

Azalea indica Linnaeus, *Spec.* 150 (1753). — Burmann, *Fl. Ind.* 42 (1768). — Willdenow, *Spec.* I. pt. 2, 831 (1798). — Persoon, *Syn.* 212 (1805). — Aiton,

- Hort. Kew.* ed. 2, I. 318 (1810), as to description and synonyms. — Roemer & Schultes, *Syst. Veg.* IV. 374 (1819), excluding β . — Blume, *Cat. Gewass. Buitenz.* 44 (1823); *Bijdr. Fl. Ned. Ind.* 852 (1825). — Sprengel, *Syst. Veg.* I. 628 (1825).
- Azalea indica* var. b. *spatulata* Blume, *Bijdr. Fl. Ned. Ind.* 852 (1825).
- Azalea indica* var. d. *angustifolia* Blume *Bijdr. Fl. Ned. Ind.* 852 (1825.)
- Azalea indica* var. e. *floribunda* Blume, *Bijdr. Fl. Ned. Ind.* 852 (1825).
- Azalea macrantha* Bunge in *Mém. Soc. Étr. Sci. St. Pétersbourg*, II. 115 (*Enum. Pl. Chin. Bor.* 41) (1833).
- Azalea indica lateritia* Lindley in *Bot. Reg.* XX. t. 1700 (1834).
- Rhododendron macranthum* G. Don, *Gen. Syst.* III. 846 (1834). — Sweet, *Brit. Flow. Gard.* ser. 2, III. t. 261 (1834).
- Rhododendron decumbens* D. Don apud G. Don, *Gen. Syst.* III. 846 (1834).
- Azalea Danielsiana* Paxton, *Mag. Bot.* I. 129, t. (1834).
- Rhododendron indicum* δ . ? *lateritium* De Candolle, *Prodr.* VII. pt. 2, 726 (1839).
- Azalea* ? *macrantha* De Candolle, *Prodr.* VII. pt. 2, 718 (1839).
- Azalea decumbens* De Candolle, *Prodr.* VII. pt. 2, 718 (1839).
- Rhododendron Breynii* Planchon in *Fl. des Serr.* IX. 77 (1854).
- Rhododendron Danielsianum* Planchon in *Fl. des Serr.* IX. 79 (1854).
- Rhododendron lateritium* Planchon in *Fl. des Serr.* IX. 80 (1854).
- Rhododendron Sieboldii* β . *serrulatum* Miquel in *Ann. Mus. Ludg.-Bat.* I. 33 (1863).
- Azalea Breynii* André in *Belg. Hort.* XV. 183 (1865).
- Azalea lateritia* Hort. ex André in *Belg. Hort.* XV. 185 (1865).
- Rhododendron indicum* γ . *macranthum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 39 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* II, 292 (1875). — Makino in *Tokyo Bot. Mag.* XXIV. 77 (1910). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 461 (1912).
- Rhododendron indicum* γ . *macranthum* a. *macranthum genuinum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 39 (*Rhodod. As. Or.*) (1870). — Makino in *Tokyo Bot. Mag.* XXIV. 77 (1910).
- Rhododendron indicum* γ . *macranthum* b. *lateritium* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 39 (*Rhodod. As. Or.*) (1870). — Makino in *Tokyo Bot. Mag.* XXIV. 77 (1910).
- Rhododendron lateritium* var. *brachytrichum* Nakai in *Tokyo Bot. Mag.* XXIX [261] (1915).
- Rhododendron hangnoense* Nakai in *Tokyo Bot. Mag.* XXIX. [261] (1915).
- Rhododendron indicum* f. *lateritium* Rehder in *Mitt. Deutsch. Dendr. Ges.* XXIV. 225 (1916); in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916).
- Rhododendron indicum* var. *brachytrichum* Komatsu in *Tokyo Bot. Mag.* XXXI. [18] (1918).

Japan: Kyushu, prov. Osumi, Yaku-shima, side of torrents, alt. 300–800 m. February 19, 1914, *E. H. Wilson* (No. 6003); same place, without collector's name (Herb. Bur. Sci. Manila); prov. Hizen, Nagasaki, 1873, *Capt. Blomfield* (Herb. Kew); same locality, ex Herb. Lugd.-Bat. 1863, as *R. Sieboldii* var. *minor*, in part (Herb. Kew and Herb. Gray); same locality, cultivated, 1863, *C. Maximowicz* (Herb. Kew, co-type of *R. indicum* γ . *macranthum* b. *lateritium* Maximowicz). Hondo, prov. Shinano, 1864, *Tschonoski* (Herb. Kew); prov. Sagami, Hakone, cultivated, June 21, 1914, *E. H. Wilson* (No. 6888); near Uruga, 1862,

C. Maximowicz (Herb. Kew and Herb. Gray, co-type of *R. indicum* γ. *macranthum* subvar. *a. genuinum* Maximowicz); prov. Suruga, foot of Mt. Fuji, cultivated, June 13, 1914, *E. H. Wilson* (No. 6932); prov. Musashi, Yokohama, cultivated, 1862, *C. Maximowicz* (Herb. Kew and Herb. Gray); Tokyo, cultivated, July 2, 1882, *K. Miyabe*; same locality (Herb. Bot. Gard. Tokyo, type of *R. lateritium* var. *brachytrichum* Nakai); prov. Shimotsuke, Nikko, alt. 600–1000 m. cultivated and sub-spontaneous, June 20 flowers, October 20 fruit, 1914, *E. H. Wilson* (Nos. 6816, 6818, 7709).

Cultivated: Java ex Herb. Ludg.-Bat. 1863, as *R. indicum* Sweet (Herb. Kew); Hongkong Bot. Gard. May 12, 14, 31, 1919, *W. J. Tutcher*.

The only place where I have seen this species truly wild is Yaku-shima, an island some ninety miles south of Kagoshima and interesting as being the southern limit of the range of a great many Japanese plants. On that island *R. indicum* Sweet is a common fluviatile shrub growing from a half to two meters high and forming dense masses in open country on the banks of rock-strewn streams. Elsewhere in Japan I found it cultivated only, although in old burial and temple grounds it is often more or less naturalized. Makino says it grows spontaneously in the provinces of Kii and Yamato in Hondo and not so very far from the old capital of Kyoto. In an old burial ground at Nikko the plants were prostrate or nearly so and with their scattered, large red flowers strongly suggested the Rock Cistus, and one saw how very applicable was the name the ancients gave this plant. The plants though often decumbent are naturally upright and very densely branched; the branches are slender but rigid and are clothed with flattened, appressed, strigose, chestnut-brown hairs which in the second and third years lose their color and disappear. The leaves are sub-coriaceous and crowded, short-petioled, narrow-lanceolate, lanceolate to oblanceolate, mucronulate, dark green, rather shining above, pallid or sub-glaucous beneath, ciliolate, usually remotely crenate-dentate, with scattered, closely appressed, red-brown setose hairs on both surfaces; in the autumn the leaves change to crimson and vinous purple. The flowers are terminal, solitary or in pairs; the corolla is broad-funnel shape, opening at the same time or immediately before annual growth commences, bright red to scarlet, sometimes rose-red; there are 5 stamens shorter than the style but equalling or exceeding the corolla in length.

In Japanese gardens this Azalea is a prime favorite and I was told that something like two hundred named forms are recognized. It is planted on or among rocks, singly or in groups, and is used as a garden edging. It bears trimming well, and its naturally low, dense habit makes it particularly useful in the gardens of Japan. Its native name is "Satsuki-tsutsuji," that is *Fifth-month Azalea*, from the fact that it blossoms in June, which is the fifth month of the year reckoning by the old Chinese calendar.

The first mention of this plant in a European book is in Breyne's *Prodromus*, I. 23 (1680), which I have not seen, but in the edition of 1739 it is mentioned in pt. I. 16 as "*Chamaerhododendron exoticum, amplissimis floribus liliaceis.*" Breyne was a merchant of Danzig and a distinguished botanist who visited Holland during the last quarter of the 17th century and in his *Prodromus* described the remarkable plants he saw there. His "*Chamaerhododendron*" was growing in the garden of a Mr. Beveringk, but no mention is made of its country of origin.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Rhododendron indicum f. *Hakatajiro* Komatsu in *Tokyo Bot. Mag.* XXXII. [45] (1918).

This is often seen in old Japanese gardens and has been introduced into England.

A form with salmon-red flowers whitish at the base is: —

Rhododendron indicum f. *tanimanoyuki* Millais, *Rhodod.* 193 (1917). — Komatsu in *Tokyo Bot. Mag.* XXII. [45] (1918).

The best known to-day of all the forms of *R. indicum* is: —

Rhododendron indicum var. *balsaminaeflorum* Nicholson in *Kew Hand-list*, pt. 2, 53 (1896); ed. 2, 499 (1902). — Bean, *Trees and Shrubs Brit. Isl.* II. 362 (1914). — Millais, *Rhodod.* 193 (1917).

Azalea Rollissoni T. Moore in *Florist and Pomol.* 1878, 35 fig. — *Garden* XVIII. 254, t. 249 (1880).

Azalea rosaeflora R. Dean in *Fl. Mag.* XIX. t. 418 (1880).

Azalea balsaminaeflora Carrière in *Rev. Hort.* 1882, 432 t. — Nicholson, *Dict. Gard.* I. 149, fig. 196 (1887).

Azalea indica var. *rosiflora* Rehder in *Bailey, Cycl. Am. Hort.* I. 122, fig. 176 (1900).

Rhododendron indicum var. *rosiflorum* Rehder in *Bailey, Stand. Cycl. Hort.* V. 2944 (1916); in *Mitt. Deutsch. Dendr. Ges.* XXIV. 226 (1916).

Rhododendron indicum *Rollisoni* Millais, *Rhodod.* 193 (1917).

This interesting little plant is common in the gardens of Japan, and it is there, as in the West, of very slow growth. It is uncertain whether it was first introduced into England or France. In the *Garden*, XVIII. p. 254 (1880), it is said to have been "introduced about three years ago by Messrs. Rollisson." Carrière says it was introduced into France from Japan by M. Viésener in 1877. On Long Island, New York, it is hardy in sheltered, shady places. Bean says "it has lived out-of-doors for many years at Kew but prefers a milder climate," Millais states that it is "a great favorite for rock gardens, where it does not always flourish." The Japanese name for the Azalea is "Komanyo-satsuki."

Similar but rather less double is: —

Rhododendron indicum f. *kinnozai* Millais, *Rhodod.* 1913 (1917).

An old form is: —

Rhododendron indicum f. *variegatum* De Candolle, *Prodr.* VII. pt. 2, 726 (1839).

Azalea indica var. *a. variegata* Blume, *Bijdr. Fl. Ned. Ind.* 852 (1825). — Lindley in *Bot. Reg.* XX. t. 1716 (1834). — Hovey, *Mag. Hort.* V. 25 (1839).

Rhododendron indicum f. *Matsushima* Millais, *Rhodod.* 193 (1917).

This has white and red striped and spotted flowers and blooms irregularly the year round. In the nursery district round Osaka it is known as "Shiki-takane-satsuki." In Blume's time it was cultivated in Java, and according to Lindley it was introduced into England by Mr. M'Killigan in 1833, to Knight's nursery in Chelsea, London. From this Azalea Knight raised several seedling forms which received names. Such plants as *Azalea indica exquisita* Lemaire (in *Fl. des Serr.* III. t. 239 (1847), syn. *R. indicum equisitum* Lemaire) and *Azalea indica Iveryana albo-*

cincta Van Houtte (in *Fl. des Serr.* XI. 211, t. 1180 (1856)), are probably seedlings from *Azalea indica variegata*. So too are Azaleas "Gledstanesii," "Iveryana," "Charles Encke" and "Reine Louise." In fact this var. *variegatum* was the parent of most of the best "Indian Azaleas" cultivated up to about 1850. According to Hovey the var. *variegatum* was introduced into Boston, Mass., in 1838, and for many years was a familiar and much admired plant at exhibitions. It is still cultivated in the Holm Lea collection, so also are the above mentioned named forms. These Azaleas were famous in their day and have been grown in this country for more than seventy years. "Gledstanesii" was raised by Messers Rollison of Tooting and exhibited for the first time in London on May 15, 1841. It was imported into this country by Hovey & Co., Boston, Mass., who exhibited it before the Massachusetts Horticultural Society on May 2, 1846. In England the var. *variegatum* appears to have been lost to cultivation until quite recently, when it was reintroduced under the name used by Millais.

A curious and abnormal form is: —

Rhododendron indicum f. polypetalum Wilson, n. comb.

Rhododendron indicum γ . *macranthum* ζ . *polypetalum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 40 (*Rhodod. As. Or.*) (1870).

Cultivated: Hongkong Bot. Gard. May 20, 1919, *W. J. Tutcher*.

I have seen the Herb. Kew co-type of this curious plant with its petaloid, deeply lacinate calyx and no corolla. In Japan its vernacular name is "Kin-no-zai." I have not seen this plant nor the following, which has a proper calyx and a deeply lacinate corolla.

Rhododendron indicum f. laciniatum Wilson, n. comb.

Rhododendron indicum γ . *macranthum* subvar. b. *lateritium* f. *laciniatum* Makino in *Tokyo Bot. Mag.* XXVII. 110 (1910).

Makino gives "Shide-satsuki" as the Japanese name of this curious form. Komatsu enumerates several other forms, but the only other it is necessary to mention here is the very distinct: —

Rhododendron indicum var. crispiflorum Schneider, *Ill. Handb. Laubholz.* II. 506 (1911). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916).

Azalea crispiflora Hooker in *Bot. Mag.* LXXIX. t. 4726 (1853). — Lemaire, *Jard. Fl.* IV. t. 430 (1854). — Hovey, *Mag. Hort.* XXIII. 70 (1857); XXIV. 141 (1858).

Azalea indica crispiflora Van Houtte in *Fl. des Serr.* IX. t. ad p. 79 (1854).

Rhododendron crispiflorum Planchon in *Fl. des Serr.* IX. 83 (1854); in *Rev. Hort.* 1854, 67.

This interesting variety has thicker leaves than the type and the wavy margins of its corolla give it a distinct appearance. The *Azalea indica* var. *grandiflora* Blume, of which there is a specimen in Herb. Kew collected by Thomas Lobb, is similar, but the margin of the corolla is not wavy. *R. indicum* var. *crispiflorum*, was introduced into England from China by Robert Fortune to the nursery of Messers Standish & Noble in 1850 or 1851. It was imported from England to Boston, Mass., in 1855 by C. M. Hovey; soon afterwards a plant was acquired by Ignatius Sargent, Esq., father of Professor C. S. Sargent, and descendants of the

original plant are in the Holm Lea collection to this day. In England it appears to have been lost to cultivation until plants were sent to Kew and elsewhere from Holm Lea by Professor Sargent.

Rhododendron Kanehirai Wilson, n. sp.

Frutex ramosissimus, 1-2.5 m. altus; rami graciles, ramulosi, juveniles pilis compressis castaneis adpressis vestiti. *Folia* petiolata, dimorpha; vernalia sparsa, lineari-lanceolata vel oblongo-lanceolata, 2-5 cm. longa, 0.5-1.5 cm. lata, acuta, glanduloso-mucronulata, basi attenuata, obscure crenato-serrata, supra atrovi-ridia, subtus pallida, utrinque pilis lucide brunneis strigosis adpressis vestita, venis supra impressis; folia aestivalia in apice ramulorum aggregata, lineari-oblan-ceolata vel anguste obovata, 1.5-3 cm. longa, 0.2-0.6 cm. lata, apice acuta vel rotundata, ceterum ut vernalia; petioli 1-5 mm. longi, dense pilis compressis castaneis adpressis vestiti. *Flores* terminales, bini vel raro solitarii, carminei vel scarlatini; pedicelli robusti, erecti, 5-8 mm. longi, dense pilis castaneis strigosis vestiti; calyx plerumque bene evolutus; lobi virides, erecti, rotundati vel ova-les, 1-5 mm. longi, 1-2 mm. lati, pilis adpressis lucide castaneis ciliati; corolla anguste infundibuliformis, 2.5-4 cm. longa et 2-2.5 cm. lata, glabra; stamina 10, inaequalia, corolla breviora, filamentis gracilibus infra medium papillose pubescentibus, antheris 1-1.5 mm. longis; ovarium dense pilis cinereis vel castaneis strigosis ob-tectum; stylus gracilis, 3-4 cm. longus, ad basin pilis paucis compressis strigosis instructus, stigmatate parvo capitato. *Fructus* non visus.

Formosa: prov. Taihoku, Urai, in a garden at the police station, April 1, 1918, E. H. Wilson (No. 10,276).

This new species is closely related to *R. indicum* Sweet, which is distinguished chiefly by its smaller leaves, rather larger, usually solitary flowers, its 5 stamens with anthers from 2 to 3 mm. long and by its stouter and glabrous style. The chief morphological difference is the number of stamens, and it may eventually prove that this new species is a decandrous variety of the old Linnean plant. I saw it in the garden mentioned above and was told that the plants had been brought from the higher slopes of the surrounding mountains. It is distinct from anything else I have seen and apparently has nothing to do with any of the species described by Hayata. The flowers vary in color from carmine-red to scarlet.

Rhododendron breviperulatum Hayata, *Icon. Pl. Formos.* III. 129 (1913). — Kanehira, *Formos. Trees*, 318, fig. 16 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [14] (1918).

Formosa: prov. Nanto, Horisha, planted in Hotel garden, December 6, 1918, E. H. Wilson (No. 11,206).

This red-flowered Azalea is well characterized by its 5 stamens and its pilose style; also by its leaves, which are oblong to oblanceolate or more rarely oblong-ovate, with no conspicuous difference between those of the spring and summer. The calyx-lobes are membranous, reddish, from 3 to 4 mm. long, ciliate and pilose without, glabrous within. The corolla is from 2.5 to 3 cm. long and broad; the stamens are nearly as long as the corolla and have rather large purple anthers. The young shoots and leaves are clothed with red-brown to gray appressed hairs. The plants my specimens came from were from 1 to 1.5 m. tall and had rather slender branches. I was told that they had been brought from the neighboring mountains.

I cannot be absolutely sure of the correct identification of this plant. It agrees very well with Hayata's description, though the calyx and corolla are larger than he describes them, and the apiculate base of the anthers is by no means very apparent.

Rhododendron obtusum Planchon in *Fl. des Serr.* IX. 80 (1854); in *Rev. Hort.* 1854, 64. — Schneider, *Ill. Handb. Laubholz.* II. 506, fig. 332e (1911). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916). — *Jour. Jap. Bot.* I. [173] fig. (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [39] 1918. — Miyazawa in *Tokyo Bot. Mag.* XXXII. [318] (1918).

Azalea indica Thunberg, *Fl. Jap.* 84 (1784), in part, not Linnaeus. — Juel, *Pl. Thunb.* 391 (1918).

Azalea obtusa Lindley in *Jour. Hort. Soc. Lond.* I. 152 (1846); in *Bot. Reg.* XXXII. t. 37 (1846).

Rhododendron Thunbergii Planchon in *Fl. des Serr.* IX. 78 (1854).

Rhododendron Sieboldii minor Miquel in *Ann. Mus. Ludg.-Bat.* I. 33 (1863); II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67).

Azalea Thunbergii André in *Belg. Hort.* XV. 184 (1865).

Rhododendron indicum δ . *obtusum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 40 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 292 (1875). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 461 (1912). — Bean, *Trees and Shrubs Brit. Isles*, II. 362 (1914). — Millais, *Rhodod.* 193 (1917).

Azalea indica var. *obtusum* Rehder in Bailey, *Cycl. Am. Hort.* I. 122 (1900).

Rhododendron indicum Kirishima Millais, *Rhodod.* 193 (1917).

Rhododendron obtusum f. *a. Honkirishima* Komatsu in *Tokyo Bot. Mag.* XXXII. [39] (1918).

Japan: Kyushu, prov. Osumi, Kirishima-jinza in a garden, said to have come from Nishi-Kirishima, May 5, 1918, *E. H. Wilson* (No. 10,328); prov. Hizen, Nagasaki, *G. H. Langsdorff* (Herb. Kew); same locality, cultivated, 1863, *C. Maximowicz* (Herb. Kew, Herb. Gray); probably same locality, *von Siebold* (Herb. Kew, Herb. Gray); Hondo, prov. Kii, May 3, 1886, *H. Mayr*; prov. Musashi, Tokyo, cultivated, May 11, 1883 (Herb. Kew); same locality, May 12, 1904, *K. Sakurai*; cultivated, Imperial Botanic Garden, Tokyo, March 29, 1914, *E. H. Wilson* (No. 6338); Yokohama, cultivated, Yokohama Nursery Company's grounds, April 30, 1918, *E. H. Wilson*; Tokyo, *Hilgendorf*, ex Mus. Bot. Berol. (Herb. Bur. Sci. Manila); prov. Rikuzen, Sendai, cultivated, May 26, 1888, *U. Faurie* (No. 2296, Herb. Kew); prov. Ugo, Akita, cultivated, May 20, 1888, *U. Faurie* (No. 2132, Herb. Kew); Japan, without locality, *L. Savatier* (Herb. Kew); without locality or collector's name, ex Herb. Lugd.-Bat. (Herb. Kew, Herb. Gray as *R. Sieboldii* var. *minor* Miquel).

It is always unfortunate when a selected or garden form has to do duty as the type of a species. Such is the case here and the result is that the phylogenetic order is inverted. Under the name of "Kirishima-tsutsuji" this plant has for centuries been grown in Japanese gardens. By this name it is mentioned by Kaempfer (*Amoen. Exot. fasc. V. 849* [1712]), and his description is perfectly accurate. It is a densely branched, twiggy plant, seldom more than a metre high and usually less, and has small, scarlet, slightly scented flowers. Like all its relatives it is exceedingly floriferous, and in April and early May, as Kaempfer says, the whole plant appears blood red. In the autumn the leaves are lustrous and assume fine crimson tints. The typical form has rather pointed corolla-lobes, as shown in Lindley's figure, and this is the one so abundantly grown around Tokyo and Yokohama. At Kurume in Kyushu this typical form is called "Hiryu." In the nursery district round Osaka a form with rounded corolla-lobes is grown and sold as the Kirishima Azalea, and it is this form that is grown in many western gardens as the real *R. obtusum* Planch. At Kurume what appears to be this form is called "Hino-tsukasa." A hose-in-hose form of the type is grown around Tokyo as "Yayegiri" and at Kurume as "Yayehiryu." In addition a number of other slight forms are grown, some of which are mentioned by Komatsu.

Like a number of other Japanese Azaleas this plant was first sent to western gardens from China. Robert Fortune found it in the Pou-shan Azalea gardens near Shanghai, and sent it to the Horticultural Society of London in 1844. Fortune saw in the same Chinese garden the hose-in-hose (semi-double) form. Fortune's introduction was named and figured by Lindley. The figure shows the obovate, rounded, obtuse, mucronate summer leaves of the previous year and two newly developed elliptic-lanceolate, acute spring leaves. Too little attention has been paid to the fact that the red-flowered species of this section of *Rhododendron* all have dimorphic leaves. Those which are formed immediately after the flowers open are from lanceolate to ovate or elliptic in shape, acute, light green in color and membranous in texture. The "spring leaves" are followed by "summer leaves," which are smaller, more or less obovate in shape, obtuse or rounded at the apex, dark green and fairly coriaceous. These spring leaves are normally deciduous and the summer leaves persistent, but climate exercises a strong influence,—the colder it is the more deciduous the leaves, and vice versa. In the early summer when the larger spring leaves predominate, the plant presents a very different appearance to what it does in late autumn or early spring when the small obovate summer leaves only are present. The overlooking or inappreciation of this simple fact has resulted in much confusion in the classification of the species and forms.

Now as to the origin of *R. obtusum*. Its name "Kirishima Azalea" would appear to suggest its original habitat. Working on this clue I visited Mt. Kirishima and am convinced that the so-called "Kirishima Azalea" is simply a color form of the Azalea so extraordinarily abundant on this mountain and hereunder referred to as *f. japonicum*. Long ago some pilgrim to this sacred mountain took back a plant or plants as a souvenir, planted it in his garden and from it (or them) has sprung all the plants of this particular Azalea in gardens to-day. The Japanese have always been fond of making pilgrimages to their sacred shrines and mountains, and their houses and gardens are often museums filled with souvenirs of such visits. The Azalea known as *amoena* is unquestionably of like origin, and so too is "Hinodegiri," a more recent comer but an established favorite in our gardens. Further, the race of Azaleas known as "Kurume Azaleas," with which we are only just now becoming acquainted, is a garden product derived by long cultivation from the same common stock — the vari-colored Azalea of Mt. Kirishima, *f. japonicum* Wils.

A form with white flowers is: —

Rhododendron obtusum f. album Schneider, *Ill. Handb. Laubholz.* II. 506 (1911). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916).

Azalea ramentacea Lindley in *Jour. Hort. Soc. Lond.* IV. 291, fig. (1849).

Rhododendron ramentaceum Planchon in *Fl. des Serr.* IX. 81 (1854); in *Rev. Hort.* 1854, 65.

Azalea obtusa alba Mottet, *Dict. Hort. Jard.* I. 302 (1892–93).

Rhododendron (Azalea) Indicum obtusum album, *Garden and Forest*, IX. 394, fig. 52 (1896). — Bean, *Trees and Shrubs Brit. Isl.* II. 362 (1914). — Millais, *Rhodod.* 193 (1917).

Azalea indica var. *obtusum f. alba* Rehder in Bailey, *Cycl. Am. Hort.* I. 122 (1900).

Azalea amoena alba Möller's *Deutsch. Gärt. Zeit.* XVIII. 476, fig. (1903).

Cultivated: Hort. Holm Lea, Brookline, Mass.; Hort. Berckmans, Augusta, Georgia, March 18, 1909, April 6, 1914, C. S. Sargent.

This is another of Fortune's introductions from China to the garden of the Horticultural Society of London. It was received in May, 1846, from Hongkong, and certainly from a garden (although this is not expressly stated) and doubtless of Japanese origin. It has the dense, twiggy habit characteristic of its class, pure white flowers with a normal calyx and rather large leaves. In every essential respect it agrees with *f. japonicum* Wils., of which it is nothing but a selected, cultivated form, as are *A. obtusum*, *A. amoena*, "Hinodogiri" and the "Kurume Azaleas." Among the latter are several white-flowered forms with relatively large leaves. Under *f. japonicum* the origin of this "obtusum" group of Azaleas is discussed, but I may add here that variants of *f. japonicum* having white flowers have been found wild on Nishi-Kirishima. Such a plant is growing in a garden at the foot of the mountain and from it I gathered specimens. I have been unable to find out just when this plant was introduced into America but in all probability it was very soon after its appearance in England. For more than sixty years it has been cultivated in the Holm Lea collection and from there was sent by Professor Sargent to England, where it appears that it had become lost, and to the Berckmans and to other gardens.

A form with a small corolla and long exserted stamens is: —

Rhododendron obtusum f. macrostemon Wilson, n. comb.

Rhododendron macrostemon Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 41, t. 3, figs. 15–20 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 292 (1875). — Ōkubo in *Tokyo Bot. Mag.* IX. 40 (1895). — Schneider, *Ill. Handb. Laubholz.* II. 507, fig. 332 h–k (1911). — Millais, *Rhodod.* 205 (1917).

Rhododendron indicum var. *macrostemon* Ōkubo & Makino in *Tokyo Bot. Mag.* XVI. 178 (1902). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 461 (1912).

Rhododendron Kaempferi var. *macrostemon* Makino in *Jour. Jap. Bot.* I. 18 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [43] (1918).

Japan: Hondo, prov. Musashi, Tokyo, cultivated, 1862, C. Maximowicz (Herb. Kew, co-type).

This garden plant is characterised by its small flowers, minute calyx, and long exserted stamens, which are three times as long as the corolla. It is, however,

more a condition of than a distinct form. In all the forms of *R. obtusum*, including the large flowered var. *Kaempferi*, the stamens are variable in length, and are inconstant in this from year to year. In the wild f. *japonicum* and the cultivated "Kurume Azaleas" the stamens are conspicuously variable in length. The same is true of the f. *multicolor*. Besides the co-type cited above there is another Japanese specimen in the Kew Herbarium from Maximowicz, collected by Tschonoski in Shinano province, central Hondo, in 1864, and labelled, apparently in Maximowicz's handwriting, "*Rhod. indicum* a. *Kaempferi* forma *subalpina*." This differs only in having larger flowers than those of his *R. macrostemon*.

A well-known form with magenta-colored flowers is: —

***Rhododendron obtusum* f. *amoenum* Wilson, n. comb.**

Azalea amoena Lindley in Paxton, *Flow. Gard.* III. 81, t. 89 (1852). — Hooker in *Bot. Mag.* LXXIX. t. 4728 (1853). — *Rev. Hort.* 1854, 241, t. — Van Houtte in *Fl. des Serr.* IX. 75, t. 885 (1854). — Hovey, *Mag. Hort.* XXII. 9 (1856); XXIII. 70 (1857); XXIV. 141 (1858).

Rhododendron amoenum Planchon in *Fl. des Serr.* IX. 80 (1854); in *Rev. Hort.* 1854, 64. — Bean, *Trees and Shrubs Brit. Isl.* II. 341 (1914).

Rhododendron Buergeri Miquel in *Ann. Mus. Lugd.-Bat.* I. 34 (1863); in II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67).

Rhododendron indicum ε. *amoenum* d. *genuinum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 41 (*Rhodod. As. Or.*) (1870). — Franchet and Savatier, *Enum. Pl. Jap.* I. 292 (1875). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 461 (1912). — Millais, *Rhodod.* 193 (1917).

Rhododendron indicum ε. *amoenum* b. *normale* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 41 (*Rhodod. As. Or.*) (1870).

Rhododendron indicum ε. *amoenum* c. *Buergeri* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 41 (*Rhodod. As. Or.*) (1870).

Azalea amoena Caldwelli Caldwell apud Van Geert, *Icon. Azal. Ind.* 45, t. 17 (1882).

Azalea indica var. *amoena* Rehder in Bailey, *Cycl. Am. Hort.* I. 122 (1900).

Rhododendron Kaempferi var. *amoenum* Rehder in Sargent, *Trees and Shrubs*, II. 30 (1907). — Schneider, *Ill. Handb. Laubholz.* II. 505, fig. 332, f-g (1911).

Rhododendron obtusum var. *amoenum* Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916). — Komatsu in *Tokyo Bot. Mag.* XXXII. 41 (1918).

Azalea amoena Hatsu-giri Millais, *Rhodod.* 113 (1917).

Japan: Kyushu, prov. Hizen, cultivated, Nagasaki, 1863, C. Maximowicz (Herb. Kew, Herb. Gray). Hokkaido, prov. Oshima, Hakodate, cultivated, 1861, C. Maximowicz (Herb. Kew as f. *normale* Maxim.). Without locality, probably Nagasaki, ex Herb. Lugd.-Bat. (Herb. Kew as *R. Buergeri* Miquel).

Cultivated: Hort. Standish (Herb. Kew, type of *Bot. Mag.* t. 4728). Arnold Arboretum, May 29, 1884, May 24, 1902 (Nos. 862-1, 862-2).

This Azalea, with its small, rich magenta-colored flowers, its small leaves and dense, twiggy habit, is too well known to need further description. It was first sent to England in 1850 or 1851 from Shanghai by Robert Fortune and was soon recognized as the most hardy of its class. It is generally assumed to be of Chinese

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

E. H. Wilson (Nos. 10,327, flowers white). Hondo, prov. Settsu, cultivated round Osaka, May, 1918, *E. H. Wilson* (form "Hinodegiri"); prov. Musashi, cultivated, Hatagaya, Hort. Oishi, May, 1914, *E. H. Wilson* (80 named forms of "Kurume Azaleas").

Cultivated: Hort. J. S. Ames, 1920 (10 forms of "Kurume Azaleas"); Hort. Kew, May 17, 1909; Hort. Holm Lea, March, 1920 (50 forms "Kurume Azaleas").

This is the phylogenetic type of the race of small flowered Azaleas so plentiful in Japan and of which *amoena*, *obtusa*, and "Hinodegiri" are perhaps the best known in western gardens. It is especially abundant on the wind-swept slopes of Nishi-Kirishima, an active volcano in south Kyushu. It is found above the tree-level, growing among lava and in volcanic ashes with coarse grasses and miscellaneous low shrubs. At its lowest altitudinal range the leaves are persistent, but higher up they are deciduous. The plants are seldom a metre tall, more usually less than half of this, and quite commonly they are prostrate or hug boulders closely. The habit is normally dense and twiggy, but when sheltered a few strong shoots develop and the plant becomes relatively tall and sparsely branched. The leaves vary greatly in size according to altitude and exposure and are smallest on the plants which annually shed their foliage. On an average plant the spring leaves are oval to elliptic-ovate, from 1 to 2 cm. long and from 0.5 to 1 cm. wide, but they may be smaller or larger. The upper surface is bright green and the under usually pallid. The summer leaves are darker, more coriaceous and from oblanceolate to obovate in shape, with the mucronate apex rounded or acute. Normally the leaves are tufted at the ends of the twigs, but on free-growing shoots they are scattered; both shoots and leaves are clad with appressed, pale gray to yellowish, straight hairs. The flowers vary a little in size, in degree of fragrance, and greatly in color; the most common shades are rosy mauve to magenta, but salmon and salmon-red are plentiful; pink, scarlet and crimson are rather rare. Plants bearing white flowers are occasionally found. It is very floriferous; each twig terminates in a flower-bud, often in a cluster of from three to five, and from each one to three flowers develop in the spring. The bud-scales are dry, nowhere glandular, and are ciliated with pale gray hairs. The calyx is usually well developed with green, oval to nearly orbicular, ciliate lobes; sometimes it is quite small. The corolla lobes may be rounded or pointed and the stamens may be shorter or longer than the corolla; the anthers vary in color from pale yellow-brown to deep magenta, this being correlated with the color of the flowers. The capsule is variable in size and the calyx-lobes are persistent.

From the above facts is it apparent that this plant is responsive to varying ecological conditions and that its extreme forms present a very different appearance. This is indeed true, nevertheless this form is well marked and, except that there is no definite demarkation between it and the variety *Kaempferi*, presents no difficulty unless an attempt be made to subdivide it. This I find impossible to do, and fortunately it is not necessary, since all the color forms have received names in gardens and may be classed under the general title of "Kurume Azaleas," from a town in Kyushu where much attention has been paid to their cultivation and development.

The type of this form came from Mt. Onsen, a volcano not far from Nagasaki, but I do not know whether it is plentiful there. It also grows on other Kyushu mountains. On Kirishima it is abundant and when in flower forms sheets of color on the bare or grass-clad, bleak slopes and is strikingly attractive. Mt. Kirishima, of

which there are two principal peaks, Nishi and Higashi, is a sacred mountain celebrated in Japanese mythology. It was on Nishi-Kirishima that the god Ninigi, grandson of the sun-goddess Amaterasu, alighted when he came down from heaven to pave the way for the conquest of Japan. A bronze sword, fixed in the ground, hilt upward, crowns the peak and is considered to be a relic of this divinity. To a place so sacred pilgrimages have been made by the Japanese from time immemorial. With their profound love for flowers some of the pilgrims would certainly take back as a souvenir living plants of this charming Azalea. Naturally it was named for the mountain, and in the course of time was distributed widely in the gardens of Japan. It is easily understood that a plant bearing flowers of an unusual color would be that selected as a souvenir by the average pilgrim. It is such forms that reached gardens first, and so we find the red *obtusum* the magenta *amoena*, the white *alba* to be the earliest known. In the autumn of 1918 I collected a quantity of seeds on Nishi-Kirishima for the Arnold Arboretum. These have been widely distributed, and it will be interesting to watch the range of variation in the habit foliage and flowers of resultant plants. I fear, however, that they will not prove so hardy as the var. *Kaempferi*, but that they will find a welcome in gardens I am quite sure.

On visiting Nishi-Kirishima, where between 650 and 1500 m. altitude this form *japonicum* grows in great abundance, and after seeing the wide range of color among the flowers, I am absolutely convinced that this Azalea is the wild parent of the race known as "Kurume Azaleas." The plant (*R. obtusum* Planch.) which, following International Rules, has to do duty as type of this species, also belongs to this race, and of it too there is a hose-in-hose form known as "Yayehiryu" in Kurume. Another of these Azaleas which in recent years has become widely known is the brilliant red-flowered "Hinodegiri." Curiously this form so abundant in Tokyo and Osaka gardens is to-day not grown at Kurume. But good as the forms named above are, they are surpassed by many hardly known in western gardens, and in fact scarcely known in Japan outside of Kurume and a narrow circle of enthusiasts. Millais in his *Rhododendrons* makes the first reference to the race in a European publication. I first became acquainted with "Kurume Azaleas" in 1914, when at the invitation of Mr. H. Suzuki of the Yokohama Nursery Company and a foremost Japanese horticulturist, I accompanied him on a visit to the nursery district at Hatagaya, a few miles north of Tokyo. There, in the garden of Mr. Oishi, I saw thousands of tiny plants bearing white and colored flowers of nearly every hue. I secured a set of fragments and dried them for our herbarium, and later, in 1916, at my suggestion Mr. John S. Ames, North Easton, Mass., purchased a selection of living plants from Mr. Oishi's collection. These arrived in the spring of 1917 and were the first "Kurume Azaleas" introduced into eastern North America. In May, 1917, I saw in the Yokohama Nursery a few large plants of "Kurume Azaleas," trained into umbrella-shape, in full flower and this decided me to visit the place of their origin. This became possible in the spring of 1918 and I was fortunate to have as a companion my friend Mr. H. Suzuki. Kurume is some 800 miles south by west from Tokyo and is quite an important city, but the fame of its Azaleas is likely to make it better known in the near future. The Azaleas are grown in a number of gardens and nurseries, but the oldest and best collection is that of Mr. Kijiro Akashi. This fine old gardener has for more than forty years devoted himself to the development of this race of Azaleas and has raised from seeds and perpetuated by cuttings nearly all the forms in cultivation. Mr. Akashi gave me the history of the race, and it seems that it originated about one hundred years ago with a gentleman named Motozo Sakamoto, who lived at Kurume. The parents came from sacred Mt. Kirishima, but whether brought from there by Sakamoto himself or given to him by some pilgrim is uncertain. At any rate,

Sakamoto cultivated several kinds and raised and selected seedlings, including one he named "Azuma-kagami," from which it is claimed all the pink-flowered forms have descended. After the death of Mr. Sakamoto his collection passed into the possession of Akashi with the result above stated. The original plant of "Azuma-kagami" is still healthy, and I tried but without success to induce Mr. Akashi to part with it. Japanese experts recognize by name more than 250 kinds of Kurume Azaleas, but the differences are often infinitesimal. I made a selection of fifty kinds in duplicate which the following year (1919) safely reached the Arnold Arboretum. The two leading experts, Messers Akashi and Kuwano, at my suggestion selected the following six as the pick of them all: — "Takasago" (pale pink, hose-in-hose), "Azuma Kagami" (deep pink, hose-in-hose), "Kirin" (deep rose shading to silvery rose, hose-in-hose), "Kumo No Uye" (pure salmon), "Kurai No Himo" (carmine, hose-in-hose), "Kureno Yuki" (white, hose-in-hose). My companion and I concurred, but had the number been twenty the task would have been much easier.

To my mind the "Kurume Azaleas" are the loveliest of all Azaleas. The colors are so pure and exquisite and of every hue and shade from white, pink and salmon to scarlet, crimson and the richest magenta. Many of them are delicately fragrant. The hose-in-hose forms have none of the ugly features of double flowers, but on the contrary they are singularly attractive and the flowers last longer. The stamens, always five, are straight, exerted or included and the anthers vary in color from pale yellow-brown to blackish crimson; the color of the corolla and that of the anthers being correlated. The plants are often trained as low standards with a compact umbrella-shaped crown; less commonly they are dense and globose, or open and irregular in form. They are extremely floriferous, and in season the blossoms often completely hide the leaves. The leaves are of course dimorphic, but apart from this they vary considerably in size, in their shades of green, in their autumn-coloring and in their degree of persistence. In a great measure these variations are correlated with the color of the flowers, and experts in Japan can with ease distinguish each variety by its foliage and general appearance. It is strange that a race so rich in forms and of such decorative value should have remained unknown to us until now. Yet the explanation is simple. Interior Kyushu is little known to the western world, and even to those Japanese whose home is on the other islands. The feudal system of government which until comparatively recently obtained in Japan created and preserved this aloofness. Further, Kurume is remote from the horticultural centres of Osaka and Yokohama, from whence we have drawn the bulk of our garden plants and where a business is made of growing for export. Nagasaki is much nearer, but in the days of Thunberg and Siebold, Veitch, Maximowicz and Fortune intercommunication was difficult, and for foreigners impossible. And so it has resulted that the product of Sakamoto's hobby richly developed by Akashi has remained hidden from the outside world until now. During the last decade the fame of the Kurume Azaleas has reached Osaka, Tokyo and other places, and growers have obtained stocks and are propagating them apace. Unfortunately every grower and enthusiast names the plants according to his fancy and the result in a few years will be chaos. And this is helped by the fact that every slight sport or variant is kept and named and no attempt at selection made. I do not see how it is possible to improve upon the strain grown in Kurume unless yellow could be injected. What is needed is rigorous selection and the reduction of the named forms to fifty or less. In the past seedling selection and preservation of sports by vegetative propagation have been the sole means employed in the evolution of the race of "Kurume Azaleas," but now attempts at hybridising them with the large flowered "Indian" and "ledifolia" types are in progress. This may result in a new race, but whether it will be as lovely and

fascinating as the present one is doubtful. In the matter of their origin Mr. Akashi was careful to inform me that for purposes of policy he had not before set forth the facts so clearly and that more or less distorted versions had appeared in print in Japan. To test the story for ourselves Mr. Suzuki and I on leaving Kurume visited and ascended Nishi-Kirishima and were convinced.

In the garden of Mr. Akashi I selected and obtained a double set of the under-mentioned fifty varieties of these Azaleas which I consider the best, and these safely reached the Arnold Arboretum on April 24, 1919. The names are those of the originator, Akashi, and therefore authentic. The plants will be propagated and distributed under these names and, owing to the reprehensible habit of changing names by Japanese nurserymen, will probably remain the only set with standard names in existence.

No. 1.	SEIKAI	flowers white, hose-in-hose
2.	KURENO YUKI	flowers white, hose-in-hose
3.	SHIN SEIKAI	flowers creamy white, hose-in-hose
4.	YOROZUYO	flowers white
5.	NANI WAGATA	flowers white, suffused with salmon-pink
6.	TANCHO	flowers flesh-color, hose-in-hose
7.	HACHIKA TSUGI	flowers white, suffused with lavender
8.	IROHAYAMA	flowers white, margined with pale lavender
9.	HOŌ	flowers white, tinged with pink
10.	SUIYŌHI	flowers flesh-color
11.	TAKASAGO	flowers pale pink, the color of apple-blossoms
12.	KASUMI GASEKI	flowers pale pink
13.	BIJINSUI	flowers pale pink
14.	ASAGASUMI	flowers rose-pink, hose-in-hose
15.	KIMIGAYO	flowers pink
16.	AZUMA KAGAMI	flowers deep pink, hose-in-hose
17.	OSARAKU	flowers white, suffused and margined with lavender
18.	OTOME	flowers blush pink
19.	AYA KAMMURI	flowers rose-color
20.	SHINTOKI NO HAGASANE	flowers rose, shading to pink, hose-in-hose
21.	SAOTOME	flowers rose-color
22.	KIRIN	flowers deep rose, shading to silvery rose
23.	TAMAFUYO	flowers white, striped peach-color
24.	KIRITSUBO	flowers rosy mauve
25.	OMOINE	flowers pale lavender
26.	OINŌ MEZAME	flowers deep rose-color
27.	KATSURA NO HANA	flowers rose-color
28.	SHIN UTENA	flowers pale salmon
29.	KUMO NO UYE	flowers pure salmon
30.	BENIFUDE	flowers salmon
31.	SUGA NO ITO	flowers pure pink
32.	KASANE KAGARIBI	flowers dull salmon-red
33.	TSUTA MOMIJI	flowers bright red
34.	SUETSUMU	flowers crimson
35.	FUDESUTE YAMA	flowers light red
36.	IMA SHŌJŌ	flowers bright red, hose-in-hose
37.	RASHŌ MON	flowers scarlet
38.	WAKA KAYEDE	flowers red
39.	YAYEHIRYU	flowers bright scarlet, hose-in-hose

40. KURAI NO HIMO flowers carmine, hose-in-hose
 41. AGEMAKI flowers carmine
 42. HINODEGIRI flowers bright crimson
 43. AIOI flowers color of almond-blossoms, hose-in-hose
 44. SAKURA TSUKASA flowers rosy mauve
 45. TAMA NO UTENA flowers pale salmon
 46. GOSHO ZAKURA flowers white, striped peach-color
 47. UKAMUSE flowers vermilion, hose-in-hose
 48. HINODE NO TAKA flowers crimson
 49. OSARAKU SEEDLING flowers white, suffused and margined with lavender
 50. HANA ASOBI flowers red

Rhododendron obtusum var. **Kaempferi** Wilson, n. comb.

Azalea indica Thunberg, *Fl. Jap.* 84 (1784), in part, not Linnaeus.

Rhododendron Kaempferi Planchon in *Fl. des Serr.* IX. 77 (1854); in *Rev. Hort.* 1854. — Rehder in Sargent, *Trees and Shrubs*, II. 29, t. 113 (1907), flowers here represented with more than 5 stamens; in Bailey, *Stand. Cycl. Hort.* V. 2944, fig. 3393 (1916). — Schneider, *Ill. Handb. Laubholz.* II. 505, fig. 331 g-k (1911). — Bean, *Trees and Shrubs Brit. Isl.* II. 363 (1914). — Millais, *Rhodod.* 197, fig. (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [16] (1918).

Rhododendron Sieboldii Miquel in *Ann. Mus. Lugd.-Bat.* I. 33 (1863), excluding varieties; in II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67).

Azalea Kaempferi André in *Belg. Hort.* XV. 184 (1865).

Rhododendron indicum a. *Kaempferi* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 38 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 291 (1875). — Matsumura, *Pl. Nikko*, 71 (1894); *Ind. Pl. Jap.* II. pt. 2, 460 (1912). — Shirasawa, *Icon. Ess. For. Jap.* II. t. 61, figs. 1-10 (1908).

Azalea Indica var. *Kaempferi* Rehder in Bailey, *Cycl. Am. Hort.* I. 122 (1900); in Möller's *Deutsch. Gärtner-Zeit.* XVII. 417, figs. (1902).

Rhododendron scabrum var. *Kaempferi* Nakai in *Tokyo Bot. Mag.* XXXIII. 208 (1919).

Rhododendron scabrum var. *Kaempferi* f. 2. *latifolium* Nakai in *Tokyo Bot. Mag.* XXXIII. 209 (1919).

Rhododendron scabrum var. *Kaempferi* f. 3. *tubiflorum* Nakai in *Tokyo Bot. Mag.* XXXIII. 209 (1919). Komatsu in *Icon. Koisikav.* IV. 7, t. 216 (1918).

Rhododendron scabrum var. *Kaempferi* f. 4. *angustifolium* Nakai in *Tokyo Bot. Mag.* XXXIII. 209 (1919). Komatsu in *Icon. Koisikav.* IV. 31, t. 228 (1918).

Japan: Kyushu, prov. Osumi, Yaku-shima February 25, 1914, *E. H. Wilson* (No. 6082); Mt. Kirishima, western slopes, alt. 1000-1500 m., March 8, 1914, *E. H. Wilson* (No. 6228); lower slopes of Nishi-Kirishima, alt. 600 m., May 5, 1918, *E. H. Wilson* (No. 10,334); foothills round base of Nishi-Kirishima, sea-level to alt. 500 m., May 4, 1917, *E. H. Wilson* (Nos. 10,326, 10,330, 10,334); same general locality, April 29, 1917, *Z. Tashiro*; prov. Satsuma, Kagoshima, March 15, 1917, *E. H. Wilson*; same locality, May 6, 1918, *E. H.*

Wilson (No. 10,343); Togo, March 14, 1914, *E. H. Wilson* (No. 6278); prov. Hizen, Nagasaki, 1862, *R. Oldham* (Nos. 505, 507, 508, 607, Herb. Kew, Herb. Gray); same locality, *A. C. Maingay* (Nos. 872, 882, Herb. Kew); prov. Buzen, Kokura, April 20, 1903, *U. Faurie*. Shikoku, prov. Tosa, Osaki, June 5, 1889, *K. Watanabe* (Herb. Gray). Hondo, prov. Yamaguchi, near Onoda, June 17, 1918, *E. H. Wilson* (Nos. 10,379, 10,380, 10,381); between Habu and Asa, May 7, 1918, *E. H. Wilson* (No. 10,368A); round Ajisa, May 17, 1917, *E. H. Wilson* (No. 8424); prov. Yamato, Yoshino, April 22, 1914, *E. H. Wilson* (No. 6575); prov. Shinano, above Narai, alt. 1100 m., September 3, 1905, *J. G. Jack*; without precise locality, 1864, *Tschonoski* (Herb. Kew); Togakushi-yama, July 9, 1884 (Herb. Kew); prov. Idzu, Oshima, March 30, 1917, July, 1918, *E. H. Wilson* (No. 8200); prov. Sagami, Hakone, May 12, 1918, *E. H. Wilson* (No. 10,365); Miyanoshita, August 24, 1892, *C. S. Sargent*; Kamakura, May 2, 1914, *E. H. Wilson* (No. 6610); prov. Musashi, Yokosuka, May, 1875, *H. N. Moseley* (Herb. Kew); Yokohama, 1862, *C. Maximowicz* (Herb. Kew); Mt. Takao, *H. Takeda* (No. 89, Herb. Kew); Mt. Mitake, May 19, 1907, *K. Sakurai*; prov. Shimotsuke, between Nikko and Lake Chuzenji, September 3, 1892, *C. S. Sargent*; in a garden at Umagaeshi, near Nikko, alt. 1000 m., June 4, 1914, *E. H. Wilson* (No. 6807); shores of Lake Chuzenji, November 3, 1892, *C. S. Sargent*; Nikko region, alt. 100–1500 m., May 16, 1914, *E. H. Wilson* (No. 6679); prov. Ugo, Akita, May 17, 1888, *U. Faurie* (No. 2128 Herb. Kew); Lake Towada, June 25, 1894, *U. Faurie* (No. 13,272); prov. Mutsu, Shichi-nobe, June 15, 1886, *U. Faurie* (No. 661); prov. unknown, Mt. Nagi, Mimasaka, May 8, 1903, *S. Arimoto* (Herb. Gray); without locality, 1866–74, *L. Savatier* (Herb. Kew, Herb. Bur. Sci. Manila). Hokkaido, prov. Oshima, Hakodate, *C. Wright* (Herb. Kew, Herb. Gray); same locality, July, 1859, *C. Wilford* (No. 1017, Herb. Kew); same locality, 1861, *M. Albrecht* (Herb. Kew, Herb. Gray); same locality, July 29, 1888, *T. Tokubuchi*; prov. Iburi, Nobori-betsu, June 15, 1893, *U. Faurie* (No. 10,020, Herb. Kew); Chitose, June 12, 1889, *T. Tokubuchi*; prov. Hidaka, *T. Tokubuchi*; Samani-sando, June 19, 1884, *K. Miyabe* (Herb. Gray); prov. Ishikari, Sapporo, July 29, 1888, May 8, 1895, *T. Tokubuchi*; same locality, October, 1891, *K. Miyabe*; same locality, May 24, 1903, *S. Arimoto* (Herb. Gray); ex Agricultural College, Sapporo, June 7, 1885.

Cultivated: Arnold Arboretum (Nos. 3690, 3690–1, 3690–3, 3690–5, 3697).

This is the Mountain or Hill Azalea (Yama-tsutsuji) of Japan, where it is abundant from Yaku-shima in the extreme south to central Hokkaido in the north. Its altitudinal range is from sea-level to about 1600 m., though north of the Nikko region it is much less. In May and June by the wayside and on the hills and mountain-slopes it is most conspicuous with its wealth of brilliant red flowers. It grows among grasses and in thickets, also on the edge of forests and among the undergrowth in thin woods, but it is a sun-loving plant and is seen to best advantage in open thickets on mountain-slopes. The height is from 1 to 3 m., but averages from 1 to 1.5 m., and according to situation it may be densely or laxly branched and form either a broad or a narrow, twiggy bush. In the warmer parts of Japan all the summer leaves and often some of the large spring leaves are persistent, but in the colder parts they are both deciduous. The spring leaves vary in shape from lanceolate or lanceolate-ovate to elliptic, and in size from 1.5 to 6 cm. in length and from 0.8 to 3 cm. in width, and they may be either acute or obtuse; the summer leaves are elliptic to obovate in shape, as is usual in the group to which this plant belongs. The pubescence on the shoot, leaf, calyx, ovary and fruit varies in color from pale gray to rufous. Typically the color of the flowers is red, usually bright red, but it varies considerably and, excluding the forms to be mentioned later, may be any shade from flesh-color to bright red. In size the flowers vary somewhat but average from 2.5 to 4 cm. across. The corolla-lobes are rounded or pointed. The calyx is normally well-developed and the lobes are green, ciliate and usually rounded, and from suborbicular to elliptic or ovate in shape, and from 2 to 8 mm. long; usually they are from 2 to 4 mm. long and occasionally they are scarcely noticeable. The stamens may be included or exerted and this considerably affects the appearance of the flowers; the anthers vary in color from pale yellow-brown to blackish purple. The fruit is woody, erect, ovoid and from 0.6 to 1.5 cm. high, and when open the valves are recurved for about one-third their length.

In its typical form this variety, with its relatively large clustered flowers each with five stamens, cannot be confused with anything else, but there are intermediate forms which inextricably link it with *R. obtusum*. The extremes look widely different, yet on analysis size alone remains, and in this there is every gradation both among the leaves and flowers. The degree of persistence of the leaves, which is dependent upon winter temperature and exposure, strongly affects the appearance of the plants. The leaves, flowers, genital organs and fruit vary in size, and the flowers, anthers and pubescence vary in color. Round the base of Nishi-Kirishima the typical red-flowered var. *Kaempferi* is plentiful; on the middle and upper slopes of the mountain the typical f. *japonicum* abounds. In my ascent of this mountain and after exhausting every means I could think of I failed completely to discover where one variety ended and the other began. The transition was both gradual and complete. At my request my colleague, Alfred Rehder, has critically examined the mass of material in our possession and is in complete agreement as to the specific identity here set forth. I confess that I should much prefer to keep Kaempfer's Azalea as a species distinct from *R. obtusum*, but the facts as I interpret them will not permit of this. Whether var. *Kaempferi* or the f. *japonicum* is best entitled to be considered the phylogenetic type of the species is difficult to determine, but the first named is much more widely distributed, and north of Hamamatsu, in prov. Totomi, central Hondo, is the only representative of the species.

Kaempfer's Azalea was first introduced to the Occident by Professor Sargent, who sent seeds of it from the Nikko region and elsewhere in Japan to the Arnold Arboretum in the autumn of 1892, where plants flowered in May, 1897. It has proved both hardy and amenable to cultivation here, and in late May and early June of each year flowers abundantly, producing most vivid and spectacular floral

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Japan: Kyushu, prov. Chikugo, Koradai, near Kurume, April, 1918, *K. Akashi*; without locality, May, 1918, *B. Miyazawa*.

In this plant the calyx is normal but the stamens and pistil are petaloid. *Akashi's* specimen has rather small flowers salmon-red in color and is from a spontaneous plant. *Miyazawa's* is in size and color typical *Kaempferi* with double flowers. He told me this was also from a wild plant.

A form with petaloid calyx forming hose-in-hose flowers and which also has been found wild is: —

***Rhododendron obtusum* var. *Kaempferi* f. *Komatsui* Wilson, n. comb.**

Rhododendron Kaempferi var. *Komatsui* Nakai in *Tokyo Bot. Mag.* XXIX. [261] (1915).

A form with monstrous flowers may be designated: —

***Rhododendron obtusum* var. *Kaempferi* f. *monstrosum* Wilson, n. f.**

Korea: prov. Keiki, cultivated, said to have come from Japan, Agricultural Experimental Station, Suigen, May 24, 1917, *E. H. Wilson* (No. 8482).

In this curious plant the corolla is white and has a petaloid calyx, the pistil is abnormal and in most flowers is changed to a staminode, the stigma being changed to a swollen, abortive anther which is often conspicuously aristate. The flowers though hose-in-hose are as large as those of ordinary *Kaempferi*. Possibly this is the same as Nakai's f. *album*.

A form with abortive or nearly abortive corolla is: —

***Rhododendron obtusum* var. *Kaempferi* f. *cryptopetalum* Wilson, n. comb.**

Rhododendron ledifolium δ . *cryptopetalum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 36 (*Rhodod. As. Or.*) (1870). — *Matsumura, Ind. Pl. Jap.* II. pt. 2, 462 (1912).

Rhododendron Kaempferi f. ζ . *Kinshibe* Komatsu in *Tokyo Bot. Mag.* XXXII. [43] (1918).

Japan: Hondo, prov. Kawachi, Yamomoto, cultivated, May 8, 1918, *E. H. Wilson*; prov. Musashi, Hatagaya, cultivated, April 29, 1914, *E. H. Wilson* (No. 6593).

This is a monstrous form long cultivated in Japanese gardens under the name of "Kinshibe-tsutsuji." The corolla may be entirely suppressed or represented by from 1 to 5 deeply lacinate, strap-shaped lobes. The stamens vary from 5 to 10 (usually 7 to 10); the filaments are variable in length, often curled, bright crimson in color, and may be either normal in shape or slightly flattened and petaloid; the anthers are yellow. It has no garden value, being simply a curiosity. As to its specific identity there can be no question, and I am at a loss to understand why Maximowicz referred it to *R. ledifolium* (*R. mucronatum* G. Don).

A form with purple flowers and from 5 to 10 stamens is: —

Rhododendron obtusum var. **Kaempferi** f. **mikawanum** Wilson, n. comb.

Rhododendron indicum var. *mikawanum* Makino in *Tokyo Bot. Mag.* XXIII. 251 (1909).

Rhododendron Kaempferi var. *mikawanum* Makino in *Jour. Jap. Bot.* I. 18 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [37] (1918).

Rhododendron purpureum Komatsu in *Tokyo Bot. Mag.* XXXII. [16] (1918).

Rhododendron poukhanense Komatsu l. c. [12] (1918), as to the Japanese plant, not Léveillé.

Rhododendron poukhanense f. *obtusifolium* Komatsu l. c. [37] (1918).

Rhododendron poukhanense f. *acutifolium* Komatsu l. c. [38] (1918).

Rhododendron scabrum Nakai in *Tokyo Bot. Mag.* XXXIII. 207 (1919), not G. Don.

Rhododendron scabrum var. *Kaempferi*, f. 1. *purpureum* Nakai l. c. 208.

Japan: Hondo, prov. Suruga, near Ashitake-yama, lower slopes of Mt. Fuji, June 20, 1918, *H. Suzuki*; prov. Musashi, Ome, near Tokyo, May 31, 1918, *E. H. Wilson*.

This is a form of the common Mountain Azalea of Japan having purple flowers and from 5 to 10 stamens. It is found here and there mixed with the type and is occasionally cultivated, being known as the "Murasaki Yama-tsutsuji" (Purple Hill Azalea). It is in cultivation in the Arnold Arboretum from plants I brought from Japan in the spring of 1919. The increase in the number of stamens is interesting, and is another instance of the extraordinary range of variation which obtains in *R. obtusum*. In Tokyo I saw specimens from Hangno which Komatsu refers to *R. poukhanense*, but it is not that species and I could find no character by which to separate them from Makino's variety. The material was collected on the Chichibu Mountains in the western part of Musashi province. It has flowers in pairs, red-purple in color, with dark purple anthers; the leaves are rather long, lanceolate and glabrescent, and the pedicels and calyx are clothed with gray pubescence. Komatsu's *R. purpureum* is based on a specimen collected in Yumato province. It has the foliage of typical var. *Kaempferi*, pale rose-purple colored flowers each with from 7 to 10 stamens; the calyx-lobes are lanceolate, acuminate and erose.

Nakai in the *Tokyo Bot. Mag.* (XXXIII. 207 [1919]) identified this plant with *R. scabrum* G. Don. I do not understand how he arrived at this conclusion unless he was guided solely by the color of the flowers. This f. *mikawanum* never has flowers 3 inches in diameter, which is the size given by G. Don, neither are the leaves coriaceous. It seems to me that the plant described by G. Don as *R. scabrum* is that which Miquel later described as *R. sublancheolatum*; from the description I do not see how it can have anything to do with *R. obtusum* var. *Kaempferi* or any of its forms.

A hybrid race with richly colored flowers of small to medium size is: —

× **Rhododendron Sanderi** Wilson, n. hyb.

Rhododendron Simsii "Garnet" × *R. obtusum* Planchon.

This is an interesting and valuable race that has been raised and developed at Holm Lea, Brookline, the home of Professor C. S. Sargent. About thirty-five years

ago the gardener, Charles Sander, collected seeds from the Holm Lea collection of "Indian Azaleas" and from them raised many plants. Among these when they flowered were two of dwarf, compact habit which produced relatively small flowers (3 cm. diam.) carmine-red or crimson-lake in color. In the Holm Lea collection was the well-known "Indian Azalea Decora" of European origin and introduced into the garden of Marshall P. Wilder, Dorchester, Mass., about 1847. It is highly probable that this was the parent of the two Azaleas above mentioned. The best of these two Sander named "Garnet." This he crossed with an unknown Azalea having small red flowers obtained from M. Atkinson, gardener for Mr. John L. Gardner, Brookline, Mass., which to-day Sander thinks was *R. obtusum* Planch. After a close study of the race I think this conclusion is correct. The best of the hybrids from the above parents were crossed with *R. obtusum* f. *album* Schneid. and gave rise to forms with rose-pink flowers. In more recent years the well-known "Hinodegiri" has been used with excellent results. The outcome of some three decades of hybridising, selecting and recrossing is a race of beautiful Azaleas of compact twiggy habit with flowers from 2.5 to 4.5 cm. in diameter in varying shades of rose-pink, salmon, red, scarlet and fiery crimson. Among the scarlet and crimsons are deeper and richer shades than I have seen among any of the Azaleas of China and Japan. The influence of *R. Simsii* forms is seen in the number of stamens, which varies from 5 to 10 and is inconstant on the same individual; also in the size of the flowers and in the ease with which the plants may be forced into blossom during the winter and early spring months. Some of the forms have flowers no larger than those of *R. obtusum*, though in the majority the flowers exceed in size those of that plant. The race is singularly attractive and highly ornamental, and it needs no expert knowledge to predict for \times *R. Sanderi* a great future. Among the best forms are "Suzuki," scarlet maroon, "Hebe," white with rose-red stripe, "Hermione," deep salmon-red, "Ruby" crimson maroon, "Uncas," dark scarlet, "Helena," rose-pink, "Hilda Hedlund," deep pink, "Holm Lea," crimson-maroon, "Vulcan," salmon-red, "Natalie," salmon, "Muriel," rose-red, "Alice Sargent," bright salmon-red, "Rose Queen," deep rose, "Havemeyer," rose-red, "Brookline," carmine, "Venus," cerise, "Jupiter," fiery-red, "Mars," intense scarlet, "Vivid," scarlet. To this group must be referred the highly appreciated and well-known Azalea "Hexe"¹ or "Firefly" of American gardens, which is a hybrid between *R. obtusum* f. *amoenum* and *R. Simsii* "Duc de Nassau." This favorite Azalea with rich red, hose-in-hose flowers was raised by Otto Forster, Lebenhof, near Scheibbs, Lower Austria, about 1885, and was first extensively cultivated round Dresden. At first it was not particularly esteemed and was used as a stock on which to graft the ordinary "Indian Azaleas" of commerce; later it became one of the most popular varieties, and before the Great War was imported in hundreds of thousands into this country for decorative purposes. It roots freely from cuttings and flowers profusely when quite small. One of its parents, Azalea "Duc de Nassau," is an old variety, having been exhibited in London in June, 1862, by Charles Turner of Slough. There are several other hose-in-hose Azaleas of much the same parentage as "Hexe." One of these, "Vuylestekiana," has been crossed at Holm Lea with "Flambeau," a French variety, with crimson-maroon flowers of medium size. The result is a race with relatively large (4 to 6 cm. diam.), hose-in-hose flowers varying in color from intense scarlet to crimson-maroon, and perhaps the most highly colored of all Azaleas.

¹ *Azalea indica* (Hexe) Ad. Vanden Heede in *Rev. Hort. Belg.* XXXI. 49, t. (1905).

Azalea Hexe Hort. ex W. Watson, *Rhodod. and Azaleas*, 67 (1911).

Rhododendron indicum Hexe Millais, *Rhodod.* 193 (1917).

Rhododendron Simsii Planchon in *Fl. des Serr.* IX. 78 (1854); in *Rev. Hort.* 1854, 47. — Schneider, *Ill. Handb. Laubholz.* II. 506, figs. 331 l-m, 332 a-d (1911). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916).

Azalea indica Aiton, *Hort. Kew*, ed. 2, 318 (1810), as to the plant, not Linnaeus. — Sims in *Bot. Mag.* XXXVI. t. 1480 (1812). — Bentham, *Fl. Hongk.* 201 (1861), as to the Hongkong plant.

Azalea indica a. punicea Sweet, *Hort. Brit.* 264 (1827).

Rhododendron indicum var. *ignescens* Sweet, *Brit. Flow. Gard.* ser. 2, II. t. 128 (1832). — Rehder & Wilson in Sargent, *Pl. Wilson.* II. 547 (1913).

Rhododendron indicum a. puniceum Sweet in *Brit. Flow. Gard.* ser. 2, II. sub. t. 128 (1832).

Azalea indica ignescens Hovey, *Mag. Hort.* III. 72 (1837).

Rhododendron Calleryi Planchon in *Fl. des Serr.* IX. 81 (1854); in *Rev. Hort.* 1854, 66.

Rhododendron indicum β . *Simsii* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 38 (*Rhodod. As. Or.*) (1870). — Franchet in *Nouv. Arch. Mus. Paris*, ser. 2, VI. 77 (1883); *Pl. David* I. 97 (1884); in *Bull. Soc. Bot. France*, XXXIII. 235 (1886).

Rhododendron indicum Hemsley in *Jour. Linn. Soc.* XXVI. 25 (1889), as to the Chinese plant, not Sweet. — Diels in *Bot. Jahrb.* XXIX. 514 (1900); in *Notes Bot. Gard. Edinburgh*, VII. 392 (*Pl. Chin. Forrest.*) (1913). — Pampañini in *Nuov. Giorn. Bot. Ital.* n. ser. XVIII. 131 (1911). — W. Watson, *Rhodod. & Azaleas*, 79 (1911), in part. — Dunn & Tutcher in *Kew Bull. Misc. Inform.* add. ser. X. 155 (*Fl. Kwangtung and Hongk.*) (1914). — Millais, *Rhodod.* 189 (1917), in part.

Azalea indica var. *Simsii* Rehder in Bailey, *Cycl. Am. Hort.* I. 122 (1900).

Rhododendron indicum var. *Matsumura* in *Tokyo Bot. Mag.* XIV. 69 (1900).

Rhododendron indicum var. *Tamurai* Makino in *Tokyo Bot. Mag.* XVIII. 102, fig. (1904).

Rhododendron indicum γ . *macranthum* subvar. *genuinum* forma *Tamurai* Makino in *Tokyo Bot. Mag.* XXIV. 77 (1910).

Rhododendron indicum var. *formosanum* Hayata, *Icon. Pl. Formos.* III. 134 (1913). — Kanehira, *Formos. Trees*, 321, fig. 9 (1917).

China: Hongkong, Happy Valley, November 5, 1903, C. S. Sargent; without locality, J. G. Champion (Nos. 122, 123, Herb. Kew, Herb. Gray); without locality, C. Wright (No. 195, Herb. Gray, Herb. Kew); April 28, 1914, Mary Strong Clemens (No. 4261, Herb. Bur. Sci. Manila). Kwantung, Canton, March 10, 1835, S. W. Williams (Herb. Gray); same locality, October 27-30, 1916, March 20, 1917, C. O. Levine (No. 438, 599, Herb. Bur. Sci. Manila); Shiuchow, April, 1919, To Kang P'eng (No. 2761, Herb. Bur. Sci. Manila). Chekiang, Pootoo Island, E. Faber; Ningpo, Ning-kan-jou, April 15, 1877, W. Hancock (No. 32, Herb. Kew); Moka-shan, alt. 500 m. August 3, 1915, F. N. Meyer (No. 1607); without locality, 1845, R. Fortune (Nos. 72, 152, 153, Herb. Kew, Herb. Gray); hills near Taihu Lake, April, 1881, W. R. Carles. Kiangsu, near Shanghai,

1915, *D. Macgregor*. Kiangsi, Kuling, alt. 1600 m., July 28, 1907, *E. H. Wilson* (No. 1682); same locality, 1873, *G. Shearer* (Herb. Kew). Hunan, Yoku-shan, alt. 70–300 m., February 21, 1918, *Dr. Handel-Mazzetti* (No. 480). Hupeh, north and south of Ichang, alt. 30–1300 m., May and November, 1907, *E. H. Wilson* (No. 3474); same locality, *A. Henry* (Nos. 782, 1160, Herb. Kew, Herb. Gray); Changyang Hsien, thickets, alt. 1300 m., May and November, 1907, *E. H. Wilson* (No. 3472); Changlo Hsien, cliffs, alt. 600–1300 m., May, 1907, *E. H. Wilson* (No. 3472^a); Patung Hsien, *A. Henry* (No. 1416, Herb. Kew); Nanto and mountains to northward, *A. Henry* (No. 3194, Herb. Kew); Hsing-shan Hsien, alt. 1000–1800 m., May 14, December, 1907, *E. H. Wilson* (No. 569); “Kao-hien-scian,” alt. 800 m., May, 1907, *C. Silvestri*. Szech’uan, Chungking, 1885, *F. S. A. Bourne* (Herb. Kew); Kiating Fu, alt. 300–800 m., May, 1908, *E. H. Wilson* (No. 3475); Mt. Omei, May, 1904, *E. H. Wilson* (Veitch Exped. No. 5143); without locality, *A. von Rosthorn* (No. 2148); Yunnan or Kueichou, 1911, *A. Hosie* (Herb. Kew). Yunnan, Menztsze, alt. 1500–2000 m., *A. Henry* (Nos. 9900^b, 9900^c, 9900^e, Herb. Kew, Herb. Bur. Sci. Manila); same locality, alt. 1600–2300 m., April, 1893, *W. Hancock* (No. 155, Herb. Kew); Milê, *A. Henry* (No. 9900^d); “Momyen,” June 3, 1868, *D. J. Anderson* (Herb. Kew); near Tali Fu, Tsang-chan, alt. 2500 m., *J. M. Delavay* (No. 1122), Herb. Kew); Tali Fu, alt. 2200–2600 m., cultivated, *G. Forrest* (No. 4173); “Ho-yeh-shui,” north end of Lichiang Valley, alt. 2900 m., cultivated by Mossoo natives, May, 1906, *G. Forrest* (No. 2282), Herb. Kew); without locality, *G. Forrest* (Nos. 7832, 11,824); 1910, *R. P. Maire* (Nos. 3713, 3775, Herb. Bur. Sci. Manila). *Jongynak*

Formosa: prov. Koshun, South Cape, *Schmuser* ex *A. Henry* (No. 588 in part, Herb. Kew).

Cultivated: “Jardin du Luxembourg, Orangerie, 14 avril, 1822, 27 avril, 1823 ” ex Herb. *J. Gay* (Herb. Kew).

This species grows in all the temperate parts of China and in south Formosa, and a variety (*eriocarpum* Wils.) on the Kawanabe Islands off Liukiu Oshima, but it has not been found in Japan or Korea. It is particularly abundant in the area of the Yangtze Valley from near Ningpo to Mt. Omei in the far west. Hancock, on a specimen, states that “the Ningpo hills are absolutely crimson in places with this shrub”; the same is true of the Lushan hills round Kiukiang and of the region north and south of Ichang. It delights in rocky places, preferably cliffs, thin dry woods and thickets, blossoms profusely, and its wealth of red flowers makes it one of the most conspicuous of all shrubs. The habit is twiggy and much-branched with a maximum height of three metres, but averaging about half this height. The leaves are variable in size and with the shoots are clothed with appressed,

strigose, gray to shining brown hairs. At low altitudes the leaves are comparatively large, from lanceolate to ovate-elliptic, and are all persistent; at its upper altitudinal limit the leaves are much reduced in size, and nearly all fall off in the late autumn. The leaves immediately below the winterbuds are small, relatively thick, obovate to oblanceolate and more persistent than the other form of leaves. The flowers are clustered, from two to six together, at the end of the shoots and all I have seen are fairly uniform in color. The corolla is broad-funnel shape, varying in color from rose-red through bright to dark red. The calyx varies from an inconspicuous rim to a length of 5 mm.; the stamens are normally ten, rarely eight, never fewer, as long or nearly as long as the corolla but shorter than the style. In central China it is generally known as the "Yin-shan-hung." Millais says that Forrest states under Nos. 4173, 4173^a, from plants cultivated by Chinese in the Tali Valley, that the color of the flowers is "rose to white." In China I have never seen wild an albino of this species; Forrest's No. 4173 in Herb. Kew is described as having rose-colored flowers. The Formosan specimen cited from A. Henry in Herb. Kew represents Hayata's var. *formosanum*, the type of which I have seen in Tokyo. It is indistinguishable from typical *R. Simsii* so abundant in China, but in Formosa known only from the extreme southern end of the island.

For nearly a century and a quarter this plant has usurped the name *indicum*, and in books is almost hopelessly confused with the true *R. indicum* Sweet, with *R. obtusum* Planchon and its varieties, and with *R. scabrum* G. Don (*R. sublanceolatum* Miquel). The first mention of the Chinese plant I can find is in Macartney's *Embassy to China*, II. 524 (1797), where *Azalea indica* is recorded as collected in the provinces of Kiangsi and Canton (Kwangtung) in 1793. In all probability it is the Chinese species that was introduced into Kew "by the Hon. Court of Directors of the East India Company" in the ship "Cuffnells," Captain Wellbank (Aiton, *Hort. Kew*, ed. 2. 318 [1810]), though Aiton's description is of the true *R. indicum* Sweet. Sims was the first to figure the Chinese species (*Bot. Mag.* t. 1480 [1812]) under the erroneous name of *Azalea indica*, and states that it was a rare plant in England and that the only one that had flowered was in the collection of James Veré, Esq. In the Kew Herbarium there are two specimens from the Jardin du Luxembourg collected in 1822 and 1823. These are the oldest cultivated specimens I have seen. Sweet in 1832 gave it a varietal name, and says that the plant was imported in the East India Company's ship "Orwell" from China by Mr. Tate. This Mr. Tate was a nurseryman in Sloane Square, London, and the master of the ship "Orwell" was Captain Farrer, who brought a number of Chinese plants including other Azaleas to England and handed them over to Tate. In Braam's *Icones plantarum*, issued in 1818, there is a figure (t. 27) of a red-flowered Azalea. The drawing is very crude and, while it probably is meant for *R. Simsii*, it might represent any Azalea with red flowers.

The Chinese Azalea appears to have remained rare in the Occident for several decades. In 1837, under the name of *Azalea indica ignescens*, it was cultivated in America by Marshall P. Wilder, Dorchester, Mass. Fortune, from 1843 to 1856, introduced into England from Chinese gardens many Azaleas of different kinds, and these gave great impetus to the cultivation of these ornamental plants. The so-called "Indian Azaleas" of western gardens have been originated almost entirely from *R. Simsii* since about 1850, chiefly in Belgium, but some in France and others in Germany.

There is a very considerable literature on "Indian Azaleas" and at least one illustrated book (*Iconographie Azalées de L'Inde* by Auguste van Geert, published in 1882). Planchon (in *Fl. des Serr.* IX. 77 [1854]) gives a brief history of them, and enumerates varieties in the collections of M. Margottin, M. Delessert, and M. Michel. Some of them were forms of the true *R. indicum*, others of *R. phoe-*

41
44

niceum, and a few of *R. Simsii*; several are recorded as presumed hybrids between *R. phoeniceum* and *R. ledifolium* (*R. mucronatum* G. Don) and between *R. phoeniceum* and *R. Danielsianum* (*R. indicum* Sweet) but such hybrids are unknown to-day and probably never existed. In *La Belgique Horticole* for 1865 (XV. pp. 174–192), there is an excellent article by M. Édouard André with a list of varieties and the names of the originators. In the various volumes of *Flore des Serres* are many colored figures and notes chiefly by Louis van Houtte. In fact in all garden periodicals from about 1850 on, there are many references and figures of these plants. However, scientific exactness is wanting and much of their early beginnings is obscure. As far as I can discover the introduction into England of the striped-flowered *Azalea vittata* by Fortune in 1850 or 1851 led to the raising of seedlings. Further impetus was given by the advent the same years of *A. vittata Bealii* with red and white striped flowers. Many of the best varieties are branch sports perpetuated by grafting. From the crossing of red and white flowered varieties many variegated forms have originated. One of the most successful breeders of "Indian Azaleas" of commerce was Mr. Joseph Vervaene of Ghent, whose work is fittingly commemorated by the lovely Azaleas "Vervaeniana," and "Vervaeniana alba." Other names immutably connected with the raising and perfecting of the "Indian Azaleas" of commerce are Messers Knight & Perry, Ivery and Rollisson in England; L. Eeckhaute, Haerens, Van Houtte, Van Geert, A. Verschaffelt, J. de Kneep and Van der Cruyssen in Belgium; Lesebe, Truffaut, H. de May and Mabire in France; Messers Schulz, E. Liebig, Seidel and Rose in Germany.

The varieties are legion, and each year adds materially to the number. All the modern ones have been raised in Belgium or Germany, where they are cultivated by the millions and exported to almost all parts of the world. Prior to 1914 more than two and a half million plants were annually exported from Belgium alone. In Germany a *Eurhododendron* (Cunningham's White) is much used as a stock for these "Indian Azaleas," but not in Belgium, where *R. phoeniceum* and its f. *concinnum* are chiefly so employed. According to J. Breck (in *The Horticulturist*, I. 512 [1847]), Marshall P. Wilder imported from Germany by way of England and cultivated in his garden in Dorchester, Mass., in 1847, Azalea "Optima" with dark scarlet flowers, Azalea "Prince Albert" with scarlet flowers, and Azalea "Alba Insignis" with large white flowers. The first two were undoubtedly derivatives of *R. Simsii* and representatives of the "Indian Azaleas" of to-day, and the first introduced into America so far as I can discover. The third was probably a form of *R. mucronatum*. Hovey (*Mag. Hort.* XIV. 284 [1848]) records that on May 13, 1848, Marshall P. Wilder exhibited before the Massachusetts Horticultural Society Azalea "Decora," another Indian Azalea. A plant of this was acquired by Ignatius Sargent, and its descendants are still in the Holm Lea collection.

Among the typical large-flowered forms of "Indian Azaleas" of to-day I can detect no influence of any species other than *R. Simsii*, though without question *R. phoeniceum* and its forms have been used. Those with smaller flowers, especially the hose-in-hose kinds like "Hexe" and "Vuylstekeana," are without doubt hybrids between forms of *R. Simsii* and *R. obtusum* f. *amoenum*. It is probable also that typical *R. obtusum* has been employed. *R. mucronatum* and numerous named forms of it are loosely designated *Azalea indica*, but in the evolution of the "Indian Azalea" of present-day gardens neither this species nor the true *A. indica* of Linnaeus have had any part.

Rhododendron Simsii var. **vittatum** Wilson, n. comb.

Azalea vittata Fortune, *Tea Districts of China*, 330 (1852).

Azalea indica vittata Van Houtte in *Fl. des Serr.* IX. t. 886 (1854). — Millais, *Rhodod.* 259 (1917).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Japan: Kawanabe Islands, Suwanose-shima, May, 1917, *H. Ushiwo* (No. 1^c); Takara-shima, May 4, 1917, *H. Ushiwo* (No. 1^d, 1^e); Taira-shima, May, 1917, *H. Ushiwo* (No. 1^f).

These specimens were collected for me by Forestry-officer Ushiwo by courtesy of Dr. Naito, chief of the Kagoshima forestry bureau, and are excellent, though unfortunately unaccompanied by field notes. The flowers of the Takara-shima specimens are white, those of that from Taira-shima pink or rose-colored. The color of the flowers and of the pubescence more readily distinguish this variety from the type than does the shape of the leaves pointed out by Hayata. I am sorry that I did not see this interesting plant in a wild state and am inclined to the view that ultimately it may prove to be a distinct species. The Kawanabe Islands are a number of small rocky eminences which jut out of the sea south of Yaku-shima toward the Liukiu Islands and have a flora like that of Liukiu (see *Jour. Arnold Arb.* I. 183-186 [1920]).

Rhododendron atrovirens Franchet in *Bull. Soc. Bot. France*, XXXIII. 235 (1886). — Hemsley in *Jour. Linn. Soc.* XXVI. 19 (1889). — Millais, *Rhodod.* 124 (1917).

The species from western Yunnan is unknown to me, except from Franchet's description. It would appear to be intermediate between *R. Simsi* Planch. and *R. microphyton* Franch. It is said to be a large bush or small tree with lanceolate, long-acuminate leaves narrowed to the base and glaucescent on the underside. The flowers are described as red, clustered 3 or 4 together, and each 2.5 cm. or less in diameter. It was discovered by Père Delavay at "Tchen-fang-chan" near Takouan in May, 1882, but apparently has not yet been introduced to cultivation.

Rhododendron lasiostylum Hayata, *Icon. Pl. Formos.* III. 135 (1913). — Kanehira, *Formos. Trees*, 321 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII [14] (1918).

Formosa: prov. Nanto, between Shushu and Horisha, March 12, 1918, *E. H. Wilson* (No. 10,020).

This is a pretty pink-flowered Azalea of which I have seen only one plant. It was growing on top of a rock by the trolley line at about 400 m. altitude, and was about 1 m. high and less broad. The branches are slender and twiggy and when young clothed with the usual appressed, flattened chestnut-brown hairs. The young leaves, petioles, pedicels, calyx and ovary are all densely clothed with chestnut-brown pubescence changing to gray. The leaves are dimorphic; the spring leaves are lanceolate to ovate-lanceolate and from 2 to 4.5 cm. long and 1 to 1.5 cm. wide, acute and deciduous; the summer leaves are oblong-obovate to oblanceolate from 2 to 4.5 cm. long and 1 to 2 cm. wide, rounded and mucronate at apex and cuneate at base; both surfaces are clothed with appressed gray to rufous-brown hairs. The flowers are terminal, three or four together at the end of the branchlet. The corolla is funnel-shape, pink, about 2 cm. long with spreading lobes and about 2.5 cm. across, the tube being rather less than 1 cm. long. The calyx is variable in size being either nearly obsolete or with distinct, rounded lobes each a millimetre or more long. The stamens are of unequal length, the longest equalling the corolla-lobes and overtopped by the style. It is with considerable hesitation that I identify my specimens with this species. There are many discrepancies but of trivial character the more important being the difference in

size of the calyx-lobes and of the leaves. I have seen the type specimens in Tokyo but the material is so poor that I could arrive at no definite conclusion in the absence of my own material for comparison.

Rhododendron subsessile Rendle in *Jour. Bot.* XXXIV. 357 (1896).—Merrill in *Govt. Lab. Publ.* (Philipp.) XXIX. 40 (1905).—Millais *Rhodod.* 249 (1917).

Philippine Islands: Luzon, Prov. Benquet, Baguio, March 1907, 1913, A. D. E. Elmer (Nos. 8595, 14,298); same locality, May–June, 1916, J. K. Santos (No. 38); same province, Mt. St. Tomas, February 18, 1916, E. C. Leano (Forestry Bureau No. 25,125); same locality, March, 1904, A. D. E. Elmer (No. 5799); same locality, July 1, November 29, 1904 R. S. Williams (Nos. 1223, 2001); same locality, May 10, 1904, P. T. Barnes (No. 922); same locality, April, 1908, H. N. Whitford (Forestry Bureau No. 11,090); Mt. Tonglon alt. 2200 m., November 12, 1905, May, 1911, E. D. Merrill (Nos. 4815, 736); same place, August, 1906, H. M. Curran (No. 5032); same place, December, 1906, E. A. Mearns; same place, December, 1908, M. L. Merritt (No. 14,148); Bucao, alt. 2300 m. January 4, 1909, R. J. Alvarez (No. 18,362); Mt. Pulog, January 3 & 8, 1909, T. C. Zschokke and M. L. Merritt (Nos. 18,035, 18,172); Pauai, July, 1907, E. A. Mearns (No. 4275); same place, alt. 2100 m., June, 1909, R. C. McGregor (No. 8420); same place, April 17, 1918, J. K. Santos (No. 31,994); District of Lepanto, January 26, 1909, F. W. Darling (No. 16,573); same district, Mt. Data, November 4, 1905, E. D. Merrill (No. 4606); prov. Abra, February 8, 1909, M. Ramos (Nos. 7249, 7105); subprov. Bontoc, alt. 1350 m. April 7, 1910, Père M. Vanoverber (Nos. 351, 364) — all these numbers in Herb. Bureau Science Manila.

This Philippine species in its pubescence and foliage somewhat resembles, as Dr. Rendle points out, the Japanese *R. mucronatum* G. Don, but it is non-glandular, the flowers are not more than one-third the size and have a minute calyx and a more tubular corolla. Its nearest relative is probably the Formosan *R. rubropilosum* Hayata. It is evidently a twiggy plant and the young branches are clothed with the usual flattened, appressed bristles. The oval to lanceolate persistent leaves are from 1.5 to 3.5 cm. long and from 0.7 to 1.2 cm. wide, acute and mucronate, with slightly recurved margins, the midrib impressed above, prominent below. On adult leaves the upper surface is clothed with appressed white hairs but when young both surfaces are densely covered with rufous-gray, silky hairs. The flowers are in terminal two to several flowered clusters. The corolla is campanulate-funnel-form, with the tube less than 1 cm. long and short spreading lobes, and is said to be from lilac to violet-purple in color; the stamens vary from 6 to 10, are slightly exserted and are overtopped by the graceful, curved style which is villose at the base. The pedicels, calyx and ovary are densely clad with ferruginous villose and more or less flattened, straight hairs. The

fruit is conic-ovoid and is densely clothed with rufous, appressed and villose pubescence.

It is widely distributed and very common on the highlands of northern Luzon. It is evidently a very pretty species and ought to be introduced into gardens. The flowers vary considerably in color and No. 364 from Bontoc is said to have white flowers. No. 351, also from Bontoc, has very slender branches and small, elliptic, acute leaves and is very different in appearance from the other specimens. I can find no morphological differences, and it appears to be simply a condition; probably the plant from which the specimen was taken was growing in the crevice of a boulder or of a cliff.

Rhododendron rubropilosum Hayata in *Jour. Coll. Sci. Tokyo*, XXX. art. 1, 173 (*Mat. Fl. Formos.*) (1911). — Kanehira, *Formos. Trees*, 326, fig. 11 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [16] (1918).

Rhododendron caryophyllum Hayata, *Icon. Pl. Formos.* III. 130, t. 23 (1913). — Kanehira, *Formos. Trees*, 318, fig. 17 (1917).

Formosa: prov. Kagi, east of Ari-san, alt. 2500–3300 m. February 2 and 4, October 26, 1919, (Nos. 9728, 9779, 10,939); same locality, June, 1914, *U. Faurie* (Herb. Bur. Sci. Manila).

On the grass-clad mountain-slopes east of Ari-san this species is common between altitudes of 2300 and 3300 m. but I did not see it elsewhere. It is a narrow shrub from 1 to 3 m. tall, with twiggy, rigid, ascending branches, which when young are densely covered with appressed, flattened, gray to red-brown hairs. The leaves are persistent, scattered on the free-growing shoots and crowded at the tip of the branchlets; in shape they vary from oblong-lanceolate to elliptic-lanceolate and are from 1 to 4 cm. long and from 0.5 to 1.7 cm. wide, acute, tipped with a glandular mucro and narrowed at base; the upper surface is dark green, with scattered, appressed, pale gray hairs and the underside is pale green and clothed with flattened, appressed, gray to brown strigose hairs, which are especially prominent on the midrib. The petiole is densely clad with appressed, flattened hairs similar to those on the branches. The bud-scales of the flowers are pubescent but not glandular. The flowers are of medium size, pink spotted with rose, short-stalked and borne several together at the end of the branchlet; the corolla is funnel-shape, with spreading lobes, and from 1.5 to 2.5 cm. across. Hayata gives the number of stamens as 10 but on my specimens I find 7 or 8 only; they are of unequal length but shorter than the corolla. The pistil overtops the stamens and has a relatively stout, slightly curved style, villose at the base, and the ovary is densely clothed with pale gray soft hairs. The fruit is ovoid-conic, from 0.5 to 0.8 cm. long and densely clothed with shaggy, gray-brown hairs. The type specimen of *R. caryophyllum* which I have seen, has smaller leaves and flowers than is usual in the species which is due to the fact that the specimens came from a wind-swept slope near the altitudinal limits of the species. My No. 9779 is similar. In late October I found many plants in full bloom and apparently it is precocious in its habit of flowering. I collected seeds, and plants of this species are now growing in the Arnold Arboretum and in several gardens in England. They will not be hardy except in favored localities like Cornwall and California.

Rhododendron tosaense Makino in *Tokyo Bot. Mag.* VI. 53 (1892); in XVIII. 101 (1904). — Schneider, *Ill. Handb. Laubholz.* II, 506

(1911). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 465 (1912). — Komatsu in *Icon. Pl. Koisikav.* II. 91, t. 130 (1915); in *Tokyo Bot. Mag.* XXXII. [17] (1918).

Japan: Shikoku, prov. Tosa, round Kochi, sea-level to 1000 m. altitude, November 20, 1914, *E. H. Wilson* (No. 7801).

A twiggy, eglandular shrub growing from 0.3 to 2.5 m. but usually from 1 to 1.5 m. tall, with slender branches clothed with appressed, straight gray to gray-brown hairs. The leaves are scattered on the shoots and free-growing branches and crowded at the end of the branchlets. They are more or less deciduous and change in the autumn to purplish crimson; those formed in the spring are oblanceolate or lanceolate from 1.2 to 3.5 cm. long and from 0.5 to 1 cm. broad and are acute and clothed with appressed, straight, gray hairs; the summer leaves are linear to oblanceolate from 0.5 to 1 cm. long and from 1 to 5 mm. wide. The flowers are produced in clusters of from 2 to 6, but occasionally are solitary. The corolla is lilac-purple, about 3 cm. across. The pedicels are short and the calyx is very small and clothed with long, straight white hairs. The stamens vary in number from 5 to 8, are of unequal length, included and overtopped by the style which is exserted. The fruit is ovoid, about 9 mm. long, clothed with appressed, straight hairs and subtended by the persistent calyx-lobes.

The species is abundant on the lower mountain-slopes of Tosa province, Shikoku, from sea-level to altitudes of 1000 m. On exposed slopes it is dwarf and dense in habit, but in thickets it is tall and loosely branched. The summer leaves are minute and with the slender twiggy branchlets easily distinguish this species from others. I have seen a few flowers, but the color is not attractive though doubtless in spring when covered with blossoms, the plant would have a charm of its own. I saw plenty of it in November, 1914, and collected seeds for the Arnold Arboretum. These were distributed but the plants raised here have not proved hardy.

Rhododendron serpyllifolium Miquel in *Ann. Mus. Lugd.-Bat.* II. 165 (1865–66); *Prol. Fl. Jap.* 97 (1866–67). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér 7, XVI. 42, t. 4, figs. 18–25 (*Rhodod. As. Or.*) (1870). Franchet & Savatier, *Enum. Pl. Jap.* I. 292 (1875). — Hooker f. in *Bot. Mag.* CXXII. t. 7503 (1896). — Boissier in *Bull. Herb. Boiss.* V. 919 (1897). — Matsumura in *Bot. Mag. Tokyo* XIV. 69 (1900); *Ind. Pl. Jap.* II. pt. 2, 464 (1912). — Schneider, *Ill. Handb. Laubholz.* II. 507, fig. 333 a–d (1911). — Bean, *Trees & Shrubs Brit. Isl.* II. 378 (1914). — Millais, *Rhodod.* 240 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [15] (1918).

Azalea? *serpyllifolia* A. Gray in Perry, *Jap. Exped.* II. 315 (1857); in *Mém. Am. Acad.* n. ser. VI. 399 (1858–59). — Masters in *Gard. Chron.* n. ser. XVII. 429 (1882).

Japan: Kyushu, prov. Osumi, Mt. Takakuma, March, 1912, *K. Toyohira*; prov. Hizen, Mt. Unzen, April 14, 1914, *K. Sakurai*; Nagasaki, cultivated, 1863, *C. Maximowicz* (Herb. Kew). Hondo, prov. Settsu, near Kobe, April, 1875, *H. N. Moseley* (Herb. Kew); prov.

Idzu, May 19, 1853, *C. Wright* (Herb. Gray); same place, cultivated, April, 1854, *Messers Williams & Morrow* (Herb. Gray, type); Mt. Amagi, April 27, 1913, *K. Sakurai*; prov. Sagami, Hakone, April and May, 1875, *H. N. Moseley* (Herb. Kew).

Cultivated: Arnold Arboretum, flowered in greenhouse, March 14, 1907; Royal Gardens, Kew, April 14, 1896 (Herb. Kew, type of *Bot. Mag.* t. 7503).

During my travels in Japan I did not see this pretty little species in a wild state. On the volcanic mountains of the Hakone region and on those of Idzu province, especially Amagi-san, it is said to be common. This seems to be the northern limit of its range. Matsumura gives two or three other localities in Hondo but I have seen no specimens except one from near Kobe which probably came from a cultivated plant. In the south island of Kyushu it grows on the volcanic Mt. Unzen near Nagasaki, and on Mt. Takakuma in the southern province of Osumi. Matsumura records it from Okinawa Island in Liukiu but I have seen no material from there. In Japan it appears to be confined to purely volcanic soils. It is also frequently cultivated there, especially in the gardens of the peasantry.

According to Messers Williams and Morrow its Japanese name signifies "Rice Azalea" and has reference to the leaves which are minute and smaller than those of any other Japanese species. It was introduced into cultivation in England by Messers Veitch, presumably through their collector Charles Maries, and flowered for the first time in England in 1882. Later it was introduced to America but is still very rare. In the Arnold Arboretum it has not proved hardy. The figure in the *Botanical Magazine* gives a very good idea of this species.

Differing from the type in color of the corolla only is: —

Rhododendron serpyllifolium* var. *albiflorum Makino in *Tokyo Bot. Mag.* XXII. 57 (1908). — Schneider, *Ill. Handb. Laubholz.* II. 507 (1911). — Komatsu in *Tokyo Bot. Mag.* XXXII. [15] (1918).

Of this I have seen only Makino's type which he says is from the mountains of western Japan. Makino names it in the vernacular the "Shirobana-unzen-tsutsuji."

Rhododendron Nakaharai Hayata in *Jour. Coll. Sci. Tokyo* XXV. art. 19, 153 (*Fl. Mont. Formos.*) (1908). — Millais, *Rhodod.* 216 (1917). — Kanehira, *Formos. Trees*, 325, fig. 15 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [14] (1918).

Rhododendron serpyllifolium Hayata in *Tokyo Bot. Mag.* XX. 72 (1906), not Miquel.

Formosa: prov. Taihoku, Mt. Shichisei ex Hayata.

I did not see this plant growing wild but in Tokyo examined the type specimen and consider it to be a good species. As the author points out it is nearest to *R. serpyllifolium* Miq. from which it is at once distinguished by its larger flowers, each with 10 stamens.

Rhododendron Tschonoskii Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. 42, t. 3, figs. 8b–14 (*Rhodod. As. Or.*) (1870); in *Bull. Acad. Sci. St. Pétersbourg*, sér. 3, XV. 230 (1871).—Franchet & Savatier, *Enum. Pl. Jap.* I. 295 (1875). — Boissier in *Bull. Herb. Boiss.* V. 919 (1897). — Matsumura, *Pl. Nikko*, 71 (1894); *Ind. Pl. Jap.* II. pt. 2, 465 (1912). — Schneider, *Ill. Handb. Laubholz.* II. 507, fig. 332 l–o (1911). — Bean, *Trees and Shrubs Brit. Isl.* II. 383 (1914). — Millais, *Rhodod.* 256 (1917). — Nakai in *Tokyo Bot. Mag.* XXXI. 244 (1917); *Fl. Syl. Kor.* VIII. 46, t. 17 (1919). — Komatsu in *Tokyo Bot. Mag.* XXXII. [15] (1918).

Azalea Tschonoskii O. Kuntze *Rev. Gen.* II. 387 (1891).

Rhododendron trinerve Franchet apud Boissier in *Bull. Herb. Boiss.* V. 92 (1897).

Rhododendron Tschonoskii a. typicum Makino in *Tokyo Bot. Mag.* XVIII. 66 (1904).

Rhododendron Tschonoskii a. typicum a. pentamerum Makino *l. c.* — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 465 (1912).

Rhododendron Tschonoskii a. typicum b. tetramerum Makino *l. c.* — Matsumura *l. c.*

Rhododendron Tschonoskii β. trinerve Makino in *Tokyo Bot. Mag.* XVIII. 66 (1904). — Schneider, *Ill. Handb. Laubholz.* II. 507 (1911).

Rhododendron Tschonoskii β. trinerve a. pentamerum Makino *l. c.* — Matsumura *l. c.*

Rhododendron Tschonoskii β. trinerve b. tetramerum Makino *l. c.* — Matsumura *l. c.*

Rhododendron Tschonoskii var. tetramerum Komatsu in *Tokyo Bot. Mag.* XXXII. [15] (1918).

Japan: Shikoku, prov. Tosa, Nakatsu-san, June 20, 1889, *K. Watanabe* (Herb. Gray). Hondo, prov. Ettchu, Mt. Tateyama, July 23, 1884, (Herb. Kew); prov. Shinano, Tsubakura-dake, alt. 1300–1600 m., not common, September 15, 1914, *E. H. Wilson* (No. 7455); Hattasan, July 13, 1894, *U. Faurie* (No. 13,356); Mt. Asama, July, 1904, *U. Faurie* (No. 5831); prov. Shimotsuke, Nikko region, Lake Yumoto, June 22, October 18, 1914, *E. H. Wilson* (No. 7657); same locality, November 6, 1892, *C. S. Sargent*; Lake Chuzenji, August 12, 1905, *J. G. Jack*; Mt. Nyoho, August 3, 1909, *K. Sakurai*; prov. Uzen, Adzuma-san, alt. 800–2000 m. July 20, 1914, *E. H. Wilson* (Nos. 7233, 7234, 7235); mountains round Yamagata, July 12, 1889, *U. Faurie* (No. 4444); prov. Rikuchu, slopes of Hayachine-san, alt. 1300–2000 m. September 28, 1914, *E. H. Wilson* (No. 7583); Mt. Iwate July 16, 1903, *S. Arimoto* (Herb. Gray); high mountains, 1865, *Tschonoskii ex Maximowicz* (Herb. Kew, Herb. Gray, co-type); prov. Mutsu, Nambu, Shichinoki, June 23, 1889, *U. Faurie* (Herb. Kew);

Lake Towada, June 25, 1894, *U. Faurie* (No. 13,279, Herb. Kew); Hakkoda-san, alt. 1600 m. July 5, 1914, *E. H. Wilson* (No. 7122); same locality, *U. Faurie* (No. 875, Herb. Kew). Hokkaido, prov. Oshima, Esashi, August 4, 1890, *K. Miyabe & Y. Tokubuchi* (Herb. Gray); prov. Ishikari, near Sapporo, September 17, 1892, *C. S. Sargent*; Masuzan, July 30, 1897, *U. Faurie* (No. 636).

Korea: prov. South Keisho, Chiri-san, peak Ten-no-bo, alt. 1845 m. November 16, 1917, *E. H. Wilson* (No. 9595); same place, July 30, 1913, *T. Nakai*.

Cultivated: Arnold Arboretum (No. 3691); Royal Gardens, Kew, July 1, 1910; Gardens, Tokyo, May 31, 1882, *K. Miyabe*.

This is an alpine or subalpine species that grows on most of the high mountains in Hondo from Shinano northward; in Hokkaido it grows sparingly as far north as the mountains round Sapporo. From Shikoku I have seen but one specimen and it is not reported from Kyushu. Out of Japan this species has been found only on the Chiri-san range in south Korea. This is an interesting discovery for the plant does not grow on Quelpaert Island where so many Japanese species are common, neither has it been found elsewhere in Korea. In the Nikko region *R. Tschonoskii* is quite common, especially round Lake Yumoto, where it grows on shady rocks. On Adzuma-san it is abundant in places fully exposed to the sun and this locality is about the centre of its distribution. On the wind-swept upper-slopes of Hayachine-san it is common but north of this mountain I found it to be rare.

The plant forms a broad, densely branched twiggy shrub from 0.3 to 2.5 m. high but is usually from 0.5 to 1.5 m. being taller in the forest shade than on exposed rocks. It prefers rocky ground, but seems to be equally happy in the moist woods of Nikko and on the exposed cliffs and slopes of Adzuma and Hayachine mountains. Sometimes it forms a broad low mat but is more usually an upright shrub. It is always very twiggy and the young shoots are densely covered with closely appressed, flattened rufous hairs. The short stalked leaves are deciduous, crowded at the end of the branches, and vary in shape from narrow-lanceolate to ovate-lanceolate or broad-oblongate, they are acute and mucronulate and may have one or three prominent nerves; the undersurface is pale and both surfaces have long, appressed, soft, red-brown hairs which are especially noticeable on the young leaves. The flowers are small, with a white corolla, and are four or five partite; they have exserted stamens, and are borne in terminal clusters of from 4 to 6 flowers in each. Except that the leaves vary in size and may have either one or three prominent nerves, and that the flowers are tetramerous or pentamerous the species is very constant. From one and the same bush on Adzuma-san I gathered specimens which exhibited the whole range of variation on which Makino founded his numerous forms. With its small white flowers *R. Tschonoskii* is very distinct from all other Japanese species but as a garden plant it has little to recommend it though in autumn the leaves change to rich shades of orange-red and crimson.

It was discovered in 1865 by Maximowicz's Japanese collector Tschonoski on the mountains of Rikuchu province, northern Hondo, and is figured and described by Maximowicz. It was introduced into England by Charles Maries who sent seeds to Messers Veitch in 1878; to this country it was introduced by Professor Sargent who sent seeds from the Nikko region to the Arnold Arboretum in 1892.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

China: prov. Hunan, Yün-shan, near town of Wukang, alt. 1350–1400 m. June 9, 1918, *Dr. Handel-Mazzetti* (No. 738).

This rare and very distinct species is unknown to me in a living state. *Dr. Handel-Mazzetti* says it is a shrub some 2 m. tall, and that it grows on the margins of forests of broad-leaved trees and has white flowers with the tube of the corolla suffused with rose-color and the upper-lobe spotted with purple. The gray and rufous-gray appressed hairs which clothe the shoots, both surfaces of the young leaves, the under-side of adult leaves, pedicels, calyx and ovary well characterise the plant. It is evidently a much-branched twiggy shrub, and the shoots are clad for a season with the usual flattened, appressed bristles of the group. The leaves are persistent or semi-persistent, crowded at the end of the branches, dark green, glabrescent above with impressed reticulate veins. In shape the leaves vary from oval to lanceolate, being from 1.5 to 6.5 cm. long and from 1 to 2.5 cm. wide, acute or obtuse and mucronate at apex. The flowers are small, clustered from 3 to 10 together at the end of the shoots; the corolla is about 1.5 cm. across, with a nearly cylindric tube hairy on the outside, and spreading lobes. The calyx is small and hidden beneath the gray or rufous straight hairs. The five stamens are exserted, and are overtopped by the style, which has a capitate, lobed stigma. In *Maximowicz's* figure the petioles vary from 0.5 to 1 cm. in length whereas on the specimen cited none exceed 0.5 cm. but this is the only difference I can find and were more material available this would probably disappear.

The discoverer of this species is unknown but from what *Bretschneider* says it seems probable that the original specimen came from the borders of western Fokien, a region still very little known to us. *Dr. Handel-Mazzetti's* discovery in south-western Hunan is an interesting extension of the range of the species, which has for so long remained obscure. It is not in cultivation.

Rhododendron Mariae Hance in *Jour. Bot.* XX. 230 (1882). — *Hemsley* in *Jour. Linn. Soc.* XXIX. 27 (1889). — *Dunn & Tutcher* in *Kew Bull. Misc. Inform.* add. ser. X. 155 (*Fl. Hongk. and Kwangtung*) (1912). — *Hutchinson* in *Millais, Rhodod.* 207 (1917).

China: prov. Kwangtung, Ting-woo Mt., May 26, 1918, *C. O. Levine*, Canton Christian College Herb. No. 2015 (Herb. Bur. Sci. Manila); Loh-fau-shan, alt. 1000 m. August 10, 1917, *C. O. Levine* (No. 1563, Herb. Bur. Sci. Manila); same place, August 13, 1917, *E. D. Merrill* (No. 11,103, Herb. Bur. Sci. Manila); Tiu-kaen-shan, March, 1918, *To-Kang-P'eng*, Canton Christian College Herb. No. 2370 (Herb. Bur. Sci. Manila).

Known to me only from *Hance's* description with which the above-cited specimens agree exactly. It is evidently a twiggy shrub and the branchlets are clothed with appressed, reddish gray bristles which, changing to gray persist more or less during a couple of seasons. The leaves are dimorphic, clustered at the end of the branches, persistent and for a plant of its group very coriaceous; they are shining dark green and nearly glabrous above, with impressed, reticulate veins, pale below, with raised veins and scattered appressed rufous bristles. The spring leaves are elliptic-lanceolate from 3 to 9 cm. long and from 2 to 3 cm. wide, acute and mucronulate; the summer leaves are elliptic to obovate from 1 to 3 cm. long and from 0.8 to 1.5 cm. wide, obtuse or rounded at the mucronulate apex. The petioles are

strigosely pubescent. The flowers are small, crowded in terminal clusters, and fragrant. The corolla is lilac-colored, with a slender tube about 1 cm. long, and small spreading lobes; the 5 exerted stamens are overtopped by the slender style with its capitate stigma. The pedicels are straight from 1 to 1.5 cm. long and with the annular calyx and the ovary are densely clothed with shining, rufous-brown strigose hairs.

This species is known only from one or two localities near Canton in south China and is evidently closely related to *R. Seniavinii* Maxim. So far as I know it has never been introduced into cultivation, although discovered by the Reverend B. C. Henry as long ago as April, 1882.

Rhododendron scabrum G. Don, *Gen. Syst.* III. 846 (1834). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 49 (*Rhodod. As. Or.*) (1870).

Rhododendron sublaceolatum Miquel in *Ann. Mus. Lugd.-Bat.* II. 163 (1865–66); *Prol. Fl. Jap.* 95 (1866–67). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 35 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 290 (1875). — Hemsley in *Jour. Linn. Soc.* XXVI. 31 (1889). — *Gard. Chron.* XLIX. 342, fig. 157 (1911). — *Jour. Hort. Soc. Lond.* XXXVII. p. cxxxi. fig. 115 (1911). — Schneider, *Ill. Handb. Laubholz.* II. 504 (1911). — Craib in *Bot. Mag.* CXXXIX. t. 8478 (1913). — Bean, *Trees and Shrubs Brit. Isl.* II. 381 (1914). — Millais, *Rhodod.* 249 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [12] (1918).

Rhododendron indicum Hooker and Arnott, *Bot. Voy. Beechey* 266 (1841), not Sweet. — Matsumura in *Tokyo Bot. Mag.* XIV. 69 (1900).

Rhododendron Jodogawa Hort. apud Hasskarl in *Cat. Hort. Bog.* alt. 162 (1844).

Rhododendron indicum β . *sinensis* Miquel in *Ann. Mus. Lugd.-Bat.* I. 33 (1863–64).

Rhododendron ledifolium β . *purpureum* Maximowicz in *Mém. Acad. St. Pétersbourg*, sér. 7, XVI. No. 9, 36 (*Rhodod. As. Or.*) (1870), in part.

Azalea sublaceolata O. Kuntze, *Rev. Gen.* II. 387 (1891).

Rhododendron indicum var. *sublaceolatum* Makino in *Tokyo Bot. Mag.* XVIII. 100 (1904). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 461 (1912).

Rhododendron sublateritium Komatsu in *Tokyo Bot. Mag.* XXXII. [12] (1918).

Rhododendron liukiense Komatsu in *Tokyo Bot. Mag.* XXXII. [12] (1918).

Liukiu: Okinawa Island, near Katena, sea-level to 100 m. March 8, 1917, *E. H. Wilson* (No. 8046); without locality, *C. Wright* (No. 172, Herb. Gray, Herb. Kew): Takuno-shima, hillsides, April 1, 1853, *C. Wright* (Herb. Gray).

Japan: Kyushu, prov. Satsuma, Kagoshima, cultivated, May 6, 1918, *E. H. Wilson* (No. 10,344, 10,345); Kajiki, cultivated, May, 1917, *Z. Tashiro*; prov. Hizen, Nagasaki, 1862, *R. Oldham* (No. 504, Herb. Gray, Herb. Kew); same locality, *G. H. Langsdorff* (Herb. Kew); same locality, 1863, *C. Maximowicz* (Herb. Kew); same locality, cultivated, April 25, 1903, *U. Faurie* (No. 7873); prov. Chikugo, Kurume, cultivated, May 3, 1918, *E. H. Wilson*. Hondo, prov.

Musashi, Yokohama, cultivated, December 22, 1914, *E. H. Wilson* (No. 7873); same locality, May 15, 1918, *E. H. Wilson*.

Cultivated: Hort. Holm Lea, Brookline, Mass., April 15, 1915; Hort. Kew, 1912, type of *Bot. Mag.* t. 8478.

This is a laxly branched shrub from 1 to 2 m. high and has relatively stout branches which are clothed in their first year with appressed, flattened, strigose, gray-brown hairs which gradually disappear. The leaves are shining, coriaceous, nearly glabrous except on the veins at maturity, persistent; those formed in the spring are lanceolate to elliptic-lanceolate, from 4 to 10 cm. long and from 2 to 3.5 cm. broad, acute or obtuse and mucronate at the apex, the base broadly cuneate, the margin slightly recurved ciliate and sub-crenulate, the principal veins impressed above and prominently raised on the under surface; the summer leaves are oblanceolate, smaller and more coriaceous than the spring leaves and subacute or rounded at the apex, which is mucronulate; when young the leaves have appressed, straight gray-brown hairs more especially on the nerves and an occasional sessile gland on the under-side; at maturity they are very sparsely pubescent except on primary and secondary nerves; the petiole is stout, flattened, somewhat appressed to the shoot, and is clothed with pubescence similar to that on the shoots and leaf veins. The winter-buds are ovoid, acute, clothed with rufous, appressed straight hairs; the inner bud-scales cinnamon-brown, ciliate, viscid on inner surface and remain on the base of the pedicels of the expanded flowers. The flowers are borne in terminal clusters from 2 to 6 in each. The calyx is green, ample, the lobes usually oval and rounded, about 6 mm. long and 5 mm. broad, sometimes one and occasionally all the calyx-lobes lance-shape and 1.2 cm. or more long, acuminate, ciliate and glandular, and covered with appressed, straight, gray hairs. The corolla is rose-red to vivid scarlet, broad-funnel-shape about 6 cm. wide, lobed nearly to the middle; stamens 10, included, overtopped by style. The fruit is erect, ovoid, from 1 to 1.2 cm. long, sparsely hairy, and is subtended by persistent calyx-lobes.

This species is endemic in the Liukiu Islands, where it is confined to the middle, or Okinawa, group of islands and to Takuno-shima, the most southern of the northern, or Oshima, group. It grows among shrubs, coarse grasses and *Pinus luchuensis* Mayr, and when in bloom is very conspicuous. It is a common plant in gardens round Kagoshima in south Japan, having been long ago introduced from Liukiu. From Kagoshima it has been taken to Nagasaki and north as far as Tokyo, but it is only in the warmer parts of south Japan that it can be described as a common garden plant. The leaves, flowers and fruit are larger than those of other species of the group; the leaves are also more persistent and more coriaceous. The flowers are normally of an intense scarlet but they vary to rose-red. The calyx-lobes are often very unequal in size on the same flower or flower-cluster and they vary too in degree of pubescence. In habit it is more vigorous but less compact than that of other species of the group. Specimens in Herb. Kew and Herb. Gray labelled *R. Sieboldii* var. *ellipticum* ex Herb. Lugd.-Bat., 1863, would appear to belong here though the flowers are abnormally small for the species. The leaves, bud-scales, calyx and number of stamens agree and it may be that the specimens are from a plant flowering out of season or under abnormal conditions.

Maximowicz, Miquel and Makino refer to it as the "Chinese Azalea" but there does not appear to be any authority for the statement since the plant itself is unknown in China. To-day in Japan generally it is known as the Liukiu (Riukiu) Azalea, but round Kurume in Kyushu it is called "Hiwoge-yodogawa." Maximowicz does not appear to have properly known this species and has confused Wright's Tokuno-shima specimen and others with his *R. ledifolium* β . *purpureum* yet he

states that it was introduced into Petrograd in 1864. In England this Liukiu species does not appear to have been properly known before 1911, when Mr. R. C. Notcutt of Woodbridge exhibited it at the Royal Horticultural Society's Temple Show in London. In 1915 I brought plants from Japan to the Arnold Arboretum. Of all the red-flowered species of eastern Asia except the Formosan *R. Oldhamii* Maxim. it is the least hardy. In Massachusetts it is a greenhouse plant and blossoms sparsely. Miquel did not know the color of the flowers of his *R. sublanceolatum* citing it as *alba?* which it certainly is not for no white form is known even to-day. The color of the flowers varies from intense scarlet to rich magenta. Don gives deep rose-color for the type and the scarlet-flowered form, which is the usual form found growing wild, may be distinguished as f. *coccineum* Wilson, n. name. It is unfortunate that we have to adopt Don's name for this species yet it is undoubtedly the oldest since his description fits this plant and no other. He says that the leaves are "ovate" a term not often applicable to the leaves of any Azalea, they are, however, broader in this species than in any other. Don cites as a synonym *R. maximum* Thunb., which is *R. Metternichii* S. & Z., but no word of his description has reference to this Eurhododendron. This Liukiu species has been cultivated round Nagasaki and other towns in southern Kyushu I know not how long, but Don is wrong in saying that it is native of Japan, though the mistake is natural when the period he wrote in is considered. Komatsu bases his two species on slight differences in the shape of the leaves, in the color of the corolla and on the length of the corolla-tube. Such trivial differences are unimportant in such variable plants as the Azaleas of the Tsutsutsi section.

Rhododendron phoeniceum G. Don, *Gen. Syst.* III. 846 (1834).

Azalea indica var. γ . Sims in *Bot. Mag.* LIII. t. 2667 (1826).

Azalea punicea Sweet apud Poiteau in *Annal. de From.* I. 102, 104 (1829), not *Rhododendron indicum a. puniceum* Sweet.

Rhododendron indicum γ . *phoeniceum* Sweet, *Brit. Flow. Gard.* ser. 2, II. sub. t. 128 (1832).

Rhododendron indicum var. *speciosum* D. Don in Sweet, *Brit. Flow. Gard.* ser. 2, III. t. 284 (1835), excluding synonymy.

Azalea indica phoenicea Hovey, *Am. Gard. Mag.* I. 224 (1835).

Azalea indica Rawsonii Loudon, *Gard. Mag.* n. ser. II. 421 (1836).

Azalea Rawsonii Paxton, *Mag. Bot.* III. 123, t. (1837).

Rhododendron indicum γ . *purpureum* De Candolle, *Prodr.* VII. pt. 2, 726 (1839), in part, not Sweet.

Rhododendron ledifolium β . *phoeniceum* De Candolle, *Prodr.* VII. pt. 2, 727 (1839), in part. — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916).

Rhododendron puniceum Planchon in *Fl. des Serr.* IX. 79 (1854), not Roxburgh; in *Rev. Hort.* 1854, 46, 61. — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 35 (*Rhodod. As. Or.*) (1870). — Schneider, *Ill. Handb. Laubholz.* II. 504 (1911).

Azalea phoenicea Hort. Paris apud Planchon in *Fl. des Serr.* IX. 79 (1854), as a synonym; in *Rev. Hort.* 1854, 61.

Japan: ex Siebold comm. Zuccarini ex Herb. Zucc. 1842 (Herb. Gray, doubtful specimen).

Cultivated: Hort. Holm Lea, March 30, 1917, A. Rehder.

Although introduced into European gardens as long ago as 1824 and used in such vast quantities as a stock on which to graft the varieties of "Indian Azaleas"

so-called, the origin of this plant is obscure. It appears to have been sent from Canton, China, to the Horticultural Society's garden at Chiswick, probably by John Reeves but the facts are not clearly known. One of these plants was figured by Sims in the *Botanical Magazine* (t. 2667), and according to Poiteau it was known in the gardens round Paris in 1829 but was rare. No wild species to which it can be referred unhesitatingly has been discovered. In Japan I saw nothing just like it but it is evident that the "Omurasaki-tsutsuji" is a glorified form of it. In its vigorous habit, size of flower and shyness of blossoming *R. phoeniceum* G. Don resembles *R. scabrum* G. Don, whilst in the calyx and character of pubescence it suggests *R. mucronatum* G. Don; its viscid inner bud-scales agree with both species. Possibly it is a hybrid between these two species though I incline to the belief that it is nothing but an extreme form of *R. scabrum* G. Don. I have not been able to discover just when it reached America but it was cultivated by Hovey & Co., Boston, Mass., in 1835. In England attempts to hybridise this plant with all sorts of species of *Rhododendron* were made and the books record a number of presumed hybrids of remarkable parentage. For example Paxton's *Azalea Rawsonii* is said to be a cross between *A. phoenicea* and *R. dauricum atrovirens* but an unbiased examination of the plate reflects nothing but a good colored form of *R. phoeniceum*. Other presumed hybrids are dealt with hereunder and it is evident that enthusiasm out-did the facts. It is probable that this plant has had something to do in the evolution of the present-day race of "Indian Azaleas." In Belgium *R. phoeniceum* and its variety *concinnum* are the principal stocks used in grafting the "Indian Azaleas" of commerce.

A form of this is: —

***Rhododendron phoeniceum* f. *semiduplex* Wilson, n. f.**

Azalea ledifolia var. β . *phoenicea* Hooker in *Bot. Mag.* LX. t. 3239 (1833).

Rhododendron ledifolium β . *phoeniceum* De Candolle, *Prodr.* VII. pt. 2, 727 (1839), in part.

G. Don and others have merged this into the type but its semi-double flowers make it distinct as a garden plant. The color is poor. I have seen the specimen preserved in Herb. Kew from which the *Botanical Magazine* figure was prepared but no living material, and do not know if the plant is now in cultivation. In the typical *R. phoeniceum* flowers are often seen in which the stamens show a tendency toward becoming petaloid.

Another form is: —

***Rhododendron phoeniceum* f. *Smithii* Wilson, n. comb.**

Rhododendron indicum γ . *Smithii* Sweet, *Hort. Brit.* ed. 2, 343 (1830).

Rhododendron pulchrum Sweet, *Brit. Flow. Gard.* ser. 2, II. t. 117 (1832). —

De Candolle, *Prodr.* VII. pt. 2, 726 (1839). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 36 (*Rhodod. As. Or.*) (1870).

Rhododendron indicum pulchrum G. Don, *Gen. Syst.* III. 845 (1834).

Azalea indica hybrida Loudon, *Gard. Mag.* n. ser. I. 326 (1835). — Hovey, *Am. Gard. Mag.* II. 157 (1836).

This plant is stated variously to be "the produce of *R. ledifolium* Hook. impregnated by the pollen of the Old Red *R. indicum*," and "a hybrid between *R. phoeniceum* and *R. ledifolium*" but to me it appears to be nothing else than a form of *R. phoeniceum*. Under Loudon's name it was introduced into America about 1835 and was exhibited for the first time before the Massachusetts Horticultural Society on March 5, 1836, by Hovey & Co. For many years it was a favorite

plant at exhibitions. It is still cultivated in the Magnolia Gardens, near Charleston, South Carolina.

An old and obscure plant is: —

Rhododendron phoeniceum f. splendens Wilson, n. comb.

Rhododendron phoeniceum var. *splendens* D. Don in Sweet, *Brit. Flow. Gard.* ser. 2, IV. t. 385 (1837).

This is said to be a seedling raised from *R. phoeniceum* G. Don pollinated by *R. catawbiense* Mich. but the figure shows no influence of the latter plant. That it is a seedling may be accepted, but it is certainly nothing more than a good colored form of *R. phoeniceum* with only from 5 to 8 stamens.

A common garden plant in Japan is: —

Rhododendron phoeniceum var. calycinum Wilson, n. comb.

Azalea indica calycina Lindley in Paxton, *Flow. Gard.* II. 169, t. 70 (1852).

Rhododendron calycinum Planchon in *Fl. des Serr.* IX. 81 (1854); in *Rev. Hort.* 1854, 65.

Rhododendron ledifolium β . *purpureum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 36 (*Rhodod. As. Or.*) (1870), in part. — Franchet & Savatier, *Enum. Pl. Jap.* I. 291 (1875). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 462 (1912). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2944 (1916).

Azalea rosmarinifolia var. *purpurea* Rehder in Bailey, *Cycl. Am. Hort.* I. 123 (1900).

Rhododendron rosmarinifolium var. *purpureum* Schneider, *Ill. Handb. Laubholz.* II. 504 (1911). — Komatsu in *Tokyo Bot. Mag.* XXXII. [35] (1918).

Rhododendron rosmarinifolium var. *speciosum* Makino in *Tokyo Bot. Mag.* XXVII. 110 (1913).

Rhododendron Oomurasaki Makino in *Jour. Jap. Bot.* I. 18 (1917).

Rhododendron Osakazuki Komatsu in *Tokyo Bot. Mag.* XXXII. [46] (1918); in *Icon. Koisikav.* IV. 9, t. 217 (1918).

Japan: Kyushu, prov. Hizen, Nagasaki, cultivated, 1863, *C. Maximowicz* (Herb. Kew, Herb. Gray, co-types of *R. ledifolium* β . *purpureum* Maxim.); same locality, 1862, *R. Oldham* (No. 509, Herb. Kew, Herb. Gray; No. 607, Herb. Kew); prov. Chikugo, Kurume, cultivated, May 3, 1918, *E. H. Wilson*. Hondo, prov. Choshu, Shimono-seki, Sanyo Hotel garden, cultivated, May 17, 1917, *E. H. Wilson* (No. 8425); prov. Musashi, Yokohama, cultivated, Hort. Suzuki, May 15, 1918, *E. H. Wilson*; Tokyo, cultivated, May 11, 1902, *K. Sakurai*; same locality, cultivated, May 12, 1910.

As pointed out by Lindley this variety is much superior to the type for it has larger, more richly colored flowers. The corolla is rose-purple with crimson spots. The calyx-lobes are larger and more acuminate but they vary. In Paxton's figure the calyx is glabrous but this is a mistake. It is ciliate and sparsely or densely clothed with long, straight, appressed hairs. Apart from the calyx Paxton's figure is an excellent representation of the Omurasaki-tsutsuji (large purple Azalea) so much grown in the gardens of Yokohama, Tokyo and elsewhere in Japan. At

Kurume in Kyushu it is known as the Kumagai-yodogawa. It is a vigorous free-flowering plant of compact habit of growth, and its large, magenta-red flowers are very handsome. Nothing definite appears to be known in Japan of the origin of this Azalea but it has been cultivated for a long period. Maximowicz's specimen undoubtedly belongs here, but those of Langsdorff, Wright and some of Oldham's included by Maximowicz under his var. *purpureum* belong to *R. scabrum* G. Don. Recent botanists seem to have overlooked Lindley's name and excellent figure. According to Planchon this variety was introduced from China by Robert Fortune to the gardens of the Horticultural Society at Chiswick where it flowered for the first time in 1851.

A very hardy form with carmine-red flowers much cultivated in Europe and, to a less degree, in this country may be distinguished as: —

Rhododendron phoeniceum var. **calycinum** f. **Maxwellii** Wilson, n. comb.

Azalea Maxwellii Hort. Wezelenburg, *Cat.* 6, fig. (1915).

Rhododendron Maxwellii Millais, *Rhodod.* 208, fig. (1917).

Japan: Kyushu, prov. Chikugo, Kurume, cultivated, May 3, 1918, E. H. Wilson.

Cultivated: Hort. Bobbink & Atkins, Rutherford, N. J., April, 1920.

At Kurume this bright-colored form is known as the "Aka-yodogawa."

To this variable species must be added: —

Rhododendron phoeniceum var. **tebotan** Wilson, n. comb.

Rhododendron Tebotan (*Rh. Kaempferi* × *Rh. rosmarinifolium*) Komatsu in *Tokyo Bot. Mag.* XXXII. [46] (1918).

This variety has double flowers with small green leaves showing the centre, and is remarkable for its color which is the same as that of *Bougainvillea glabra* Choisy, and quite unique among Azaleas. It is an old plant in Japanese gardens but is now rare and Japanese nurserymen substitute for it *Azalea* "Fuji-manyo" (*R. mucronatum* f. *plenum* Wils.). In 1919 I secured in Japan a living plant of the true var. *tebotan* for the Holm Lea collection where it has flowered.

Rhododendron yedoense Maximowicz apud Regel in *Gartenfl.* XXXV. 565, t. 1233 a-b (1886). — Millais, *Rhodod.* 264 (1917).

Azalea Yodogawa Grignan in *Rev. Hort.* 1908, 425 t. fig. 2.

Rhododendron phoeniceum Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463 (1912), not G. Don.

Rhododendron Yodogawa Kunert in *Gartenwelt* XVI. 163, t. (1912). — Millais, *Rhodod.* 264 (1917).

Rhododendron poukhanense var. *yodogawa* Rehder in *Mitt. Deutsch. Dendr. Ges.* XXIV. 225 (1916); in Bailey, *Stand. Cycl. Hort.* V. 2943 (1916).

Rhododendron Poukhanense var. *plenum* Nakai in *Tokyo Bot. Mag.* XXXI. 245 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [12] (1918).

Azalea Tado Guno Hort. apud Millais, *Rhodod.* 264 (1917), as a synonym.

Rhododendron Matsumurai Komatsu in *Tokyo Bot. Mag.* XXXII. [13] (1918).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

This Korean Azalea is usually a compact densely branched shrub from a few inches to a metre high, but in shaded places it may be as much as two metres tall and rather loosely branched; on wind-swept slopes and on the tops of rocks it hugs the ground and forms broad mats. The branches are twiggy and clothed with appressed gray hairs which disappear the second year. No part of the plant is glandular; the winter-buds are ovoid, more or less pointed and covered with appressed, gray-brown hairs. The leaves are quite or partially deciduous according to climate and in the autumn are tinted from orange to crimson. The spring leaves are membranous, oblanceolate to lanceolate or ovate-lanceolate, from 3 to 8 cm. long and from 1 to 2 cm. broad, acute and mucronate at apex, with a narrow wedge-shaped base; they are dark green and have impressed veins on the upper surface, and are pallid below where the principal veins are prominent, ascending and arching inward but not extending to the margins of the leaf; both surfaces are clothed with straight appressed gray to shining brown hairs. The summer leaves are thicker, narrowly oblong-lanceolate, from 0.4 to 1 cm. wide, attenuate at the base; the margins are often obscurely crenate-serrate, and the upper surface is glabrous at maturity except for a few, short appressed, white bristles. The flowers are in clusters of two to several, and are remarkably fragrant, with a corolla rose to rosy purple. The calyx is ample, green and clothed with pubescence similar to that of the leaves, the lobes are lanceolate to ovate-lanceolate, from 5 to 10 mm. long, acute, obtuse or rounded. The pistil is exserted but the stamens are included, 10 in number, with purple anthers. The fruit is erect, ovoid, from 7 to 10 mm. long, clothed with strigose hairs and subtended by the persistent calyx-lobes. This is the common Azalea of Korea from about the latitude of Seoul (Keijyo), the capital city southward. It is partial to open country and on grassy mountain-slopes and in thin Pine-woods it forms dense matlike masses from a few inches to a yard high. On boulders in water-courses the growth is low and dense but in thickets the plants are more loosely branched and often two metres high. It grows from sea-level up to about 1600 m. altitude according to latitude and in some places, like the foothills of Chiri-san, it is very abundant. On Quelpaert it grows from sea-level to near the summit of Hallai-san and is common on boulders in the water-courses and on the moorlands. On Poukhan-san behind Keijyo from which it derives its specific name it is rare though there are places not far from there where it is common.

This Azalea was introduced to the Arnold Arboretum by Mr. J. G. Jack who sent seeds from Poukhan-san in the autumn of 1905. Plants flowered for the first time in May, 1914. It has proved perfectly hardy here, grows freely and flowers profusely. The flowers are distinctly and pleasantly fragrant and the plant is a decided acquisition to the list of hardy species of its section.

Japanese botanists have recorded this plant from several places in Hondo, Japan, and Komatsu (in *Tokyo Bot. Mag.* XXXII. [38] (1918)) enumerates fourteen garden forms. I have seen the Japanese material from wild plants referred to this species in the herbarium of the Imperial Botanic Garden, Tokyo, but it is too fragmentary to base a definite opinion upon. That from the Tokyo region has yellow anthers and belongs to *R. obtusum* var. *Kaempferi* f. *mikawanum* Wils. It certainly has nothing to do with *R. yêdoënsë* var. *poukhanense*. That from Bitchu province collected by Z. Yoshino has purple anthers and the veins on the upper surface are indistinctly impressed. It may represent Nakai's variety but the material is too incomplete to judge by.

Rhododendron Oldhamii Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 34 (*Rhodod. As. Or.*) (1870). — Masters in *Gard. Chron.* n. ser. XVII. 524 (1882). — Hemsley in *Jour. Linn. Soc.*

XXVI. 28 (1889). — Henry in *Trans. Asiat. Soc. Jap.* XXIV. suppl. 57 (1896). — J. H. Veitch, *Hortus Veitchii*, 286 (1906). — Matsumura & Hayata in *Jour. Coll. Sci. Tokyo*, XXII. 218 (*Enum. Pl. Formos.*) (1906). — Hayata in *Jour. Coll. Sci. Tokyo*, XXV. art. 19, 153 (*Fl. Mont. Formos.*) (1908). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463 (1912). — Komatsu in *Icon. Pl. Koisikav.* II. 69, t. 119 (1914); in *Tokyo Bot. Mag.* XXXII. [11] (1918). — Kanehira, *Formos. Trees*, 325, fig. 10 (1917). — Millais, *Rhodod.* 220 (1917).

Azalea Oldhamii Hort. Veitch. apud Masters in *Gard. Chron.* n. ser. XVII. 524 (1882).

Rhododendron Oldhamii var. *glandulosum* Hayata in *Jour. Coll. Sci. Tokyo* XXV. art. 19, 153 (*Fl. Mont. Formos.*) (1908).

Rhododendron longiperulatum Hayata, *Icon. Pl. Formos.* III. 138 (1913). — Kanehira, *Formos. Trees*, 323, fig. 13 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [14] (1918).

Formosa: prov. Taihoku, Jyukirin, sea-level to 1000 m., April 8, 1918, *E. H. Wilson* (No. 10,316); Tamsui, 1864, *R. Oldham* (No. 212, Herb. Kew, Herb. Gray); Sekitei, alt. 100–500 m. March 17, 1918, *E. H. Wilson* (No. 10,165); prov. Nanto, round Lake Candidius, March 11, December 7, 1918, *E. H. Wilson* (Nos. 9980, 11,175); beyond Musha up to 2900 m. March 3, 1918, *E. H. Wilson* (No. 10,029); prov. Kagi, beyond Ari-san, alt. 2800–2900 m. February 4, 1918, *E. H. Wilson* (No. 9778); prov. Koshun, South Cape, *Schmüser* ex *A. Henry* (Nos. 588, in part, 588A. Herb. Kew); prov. unknown, “Okasaki,” March, 1914, *U. Faurie* (No. 164, Herb. Bur. Sci. Manila); without locality ex Herb. Imp. Bot. Gard. Tokyo (Herb. Bur. Sci. Manila).

This is the common red-flowered *Azalea* of *Formosa*, where it is endemic and widely spread from sea-level up to 2800 m. altitude but is more abundant in the north than elsewhere. On volcanic Mt. Daiton, not far from the capital city of Taihoku, it is a feature on the wind-swept upper slopes. On sandstone cliffs east of Taihoku it is common, and in thickets round Lake Candidius in central *Formosa* it is a conspicuous shrub. On the grass-clad highlands beyond Musha and beyond Ari-san it is plentiful. I did not see it in the extreme south, which is of coral formation, and suspect that *Schmüser*'s specimen cited really came from the savage territory more to the north. The species is very glandular and very hairy, the leaves shoots, pedicels and calyx being covered with a soft pubescence much of which is long, villose and ferruginous in color but on wind-swept mountains in winter the pubescence may be bleached pale gray and the plant shaggy in appearance. It is a much-branched shrub from 1 to 3 m. tall, bushy in habit with twiggy branchlets densely clothed with glandular, spreading, red-brown hairs among which are scattered flattened, paleaceous hairs spreading and less appressed than is the rule in other red-flowered *Azaleas*. The leaves are persistent, and vary in shape from elliptic or elliptic-oblong to elliptic-ovate or very rarely to lanceolate; they are from 2 to 8 cm. long and from 1 to 4 cm. wide, rounded or subacute at the

apex which is furnished with a long gland-tipped mucro; the base is narrowed. Both surfaces and the petiole are clothed with pubescence similar to that on the shoots; on mature leaves the upper surface is scaberulous from the presence of the tumid persistent base of deciduous hairs; the veins are impressed above, raised and prominently reticulate below. The flowers are borne at the end of the shoots, and at low altitudes appear every month in the year. The corolla is red, funnel-shape, with spreading, rounded lobes, and from 3 to 5 cm. in diameter. The calyx is green and variable in size, and the lobes may be rounded and less than 1 mm. long or triangular-acuminate and as much as 8 mm. long. The ten stamens are of unequal length, the longest almost equalling the corolla but overtopped by the slender, glabrous, exserted style. The fruit is more or less erect, ovoid, from 6 to 10 mm. long, densely glandular-setose, on pedicels from 1 to 1.5 cm. long.

This species varies considerably in size of the leaves and calyx-lobes and somewhat in degree of hairiness and glandulosity but is well marked and easily recognized among all the species of its section. At its altitudinal limits and on bleak mountain-slopes the leaves, which are scattered on the free shoots and clustered at the end of the branchlets, are tinted blackish purple. It was discovered by Richard Oldham round Tamsui in 1864, and introduced to gardens by Charles Maries who sent seeds to Messers Veitch in 1878. Plants raised from these seeds were exhibited in London in the spring of 1882. Whether plants of this origin are still in cultivation I do not know. In 1918 I sent seeds to the Arnold Arboretum which were distributed in America and Europe. As a greenhouse plant Oldham's Azalea is worth cultivating.

In Tokyo I examined the type specimen of *R. longiperulatum* Hayata and could find no characters by which to separate it from *R. Oldhamii*. On Mt. Daiton, the type locality given by Hayata for his species, the variable *R. Oldhamii* is abundant but I could find no other species of the group growing there. Mr. Shimada who accompanied me told me it was all *R. longiperulatum* Hayata so-named from his original specimen collected on this mountain.

Rhododendron mucronatum G. Don, *Gen. Syst.* III. 846 (1834).

Azalea rosmarinifolia Burmann, *Fl. Ind.* 43, t. 3, fig. 3 (1768). — Rehder in Bailey, *Cycl. Am. Hort.* I. 123 (1900).

Azalea mucronata Blume, *Cat. Gewass. Buitenz.* 44 (1823); *Bijdr. Fl. Nederl.* 853 (1825).

Azalea indica alba Lindley, *Bot. Reg.* X. t. 811 (1824). — Loddiges, *Bot. Cab.* XIII. t. 1253 (1827).

Azalea ledifolia Hooker in *Bot. Mag.* LVI. t. 2901 (1829). — Hovey, *Mag. Hort.* IV. 135 (1838).

Azalea liliiflora Poiteau in *Annal. de From.* 104 (1829). — Loiseleur, *Herb. Amat.* sér. 2, I. t. 40 (1839). — Carrière in *Rev. Hort.* 1869, 299.

Azalea alba Sweet apud Poiteau in *Annal. de From.* 104 (1829), as a synonym.

Rhododendron leucanthum Bunge in *Mém. Soc. Étr. Acad. Sci. St. Pétersbourg*, II. 115 (*Enum. Pl. Chin. Bor.* 41) (1833).

Rhododendron ledifolium G. Don, *Gen. Syst.* 846 (1834). — Siebold and Zuccarini in *Abh. Akad. Münch.* IV. pt. III. 131 (*Fl. Jap. Fam. Nat.* II. 7) (1846). — Planchon in *Rev. Hort.* 1854, 48. — Boissier, *Herb. Boiss.* V. 918 (1897). — Bean, *Trees and Shrubs Brit. Isl.* II. 365 (1914). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2943 (1916). — Millais, *Rhodod.* 201, t. (1917).

Rhododendron Burmanni G. Don, *Gen. Syst.* III. 846 (1834).

Rhododendron macranthum var. *albiflorum Souchetianum* Lemaire in Loiseleur, *Herb. Amat.* sér. 2, III. t. 15 (1843).

Rhododendron liliiflorum grandiflorum souchetianum Lemaire in Loiseleur, *Herb. Amat.* sér. 2, III. sub. t. 15 (1843), as a synonym.

Rhododendron ledifolium a. *leucanthum* De Candolle, *Prodr.* VII. pt. 2, 727 (1839). — Miquel in *Ann. Mus. Lugd.-Bat.* I. 34 (1863); II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 36 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 291 (1875). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 462 (1912).

Rhododendron rosmarinifolium Dippel, *Handb. Laubholz.* I. 421 (1889), not Vidal. — Voigtländer in *Gartenwelt* XV. 650, fig. (1911). — Schneider *Ill. Handb. Laubholz.* II. 503, figs. 330 f., 331 c-e (1911). — Komatsu in *Tokyo Bot. Mag.* XXXII. [13] (1918).

Azalea rosmarinifolia var. *alba* Rehder in Bailey, *Cycl. Am. Hort.* I. 123 (1900).

Rhododendron rosmarinifolium var. *album* Schneider, *Ill. Handb. Laubholz.* II. 504 (1911).

Rhododendron ledifolium var. *album* Rehder in *Mitt. Deutsch Dendr. Ges.* XXIV. 225 (1916).

Japan: Kyushu, prov. Hizen, Nagasaki, cultivated, April 25, 1903, *U. Faurie* (No. 5434); prov. Chikugo, Kurume, cultivated, May 3, 1918, *E. H. Wilson*. Hondo, prov. Choshu, Shimonoseki, cultivated, garden Sanyo Hotel, May 17, 1917, *E. H. Wilson* (No. 8426); prov. Settsu, near Osaka, cultivated, May 7, 1918, *E. H. Wilson*; prov. Kishu, May 15, 1911, *K. Sakurai*; prov. Musashi, Yokohama, cultivated, 1862, *C. Maximowicz* (*Herb. Kew*); Tokyo, cultivated, March 29, 1914, *E. H. Wilson* (No. 6339); same locality, cultivated, May 15, 1882, *K. Miyabe*; prov. Ugo, Akita, cultivated, May 20, 1888, *U. Faurie* (No. 2133); prov. Mutsu, Hirosaki, May 27, 1905, *U. Faurie* (No. 6786). Hokkaido, prov. Oshima, Hakodate, cultivated, 1855, *C. Wright* (*Herb. Gray, Herb. Kew*); same locality, 1862, *M. Albrecht* (*Herb. Kew*).

Japan: without locality, 1862-63, *R. Oldham* (No. 506); ex *Herb. Lugd.-Bat.*, 1863 (*Herb. Gray, Herb. Kew*).

China: prov. Hupeh, Ichang, cultivated, *A. Henry* (No. 3503, *Herb. Gray*).

Cultivated: ex *Herb. Bot. Bog.* (*Herb. Bur. Sci. Manila*); Forest Hills Cemetery, Boston, Mass., June 9, 1906, *J. G. Jack*; Butler Plantation, West Feliciana Parish, La., March 28, 1910, *C. S. Sargent*; Hort. Berckmans, Augusta, Georgia, March 18, 1909, *C. S. Sargent*; Hort. Kew, May 28, 1901, *A. Rehder*.

This well-known plant is a much-branched compact, wide-spreading shrub usually from 1.5 to 2 m. tall but sometimes over 3 m. high and more in diameter. The shoots in their first year, as well as the petioles and pedicels, are densely clothed with gray to gray-brown, spreading, soft hairs and intermixed with them, especially on the shoots, are few or many appressed and spreading, flattened, bristle-like hairs. On both shoots and pedicels occasional glandular hairs are present; the inner

scales of the flower-buds are also glandular. The spring leaves are light green, deciduous and rather thin in texture, they are lanceolate to ovate-lanceolate, from 3 to 6 cm. long and from 1 to 2.5 cm. wide, acute or obtuse, mucronulate, clothed on both surfaces with soft, straight, appressed, gray to rufous hairs; the summer leaves are dark green, numerous, persistent and sub-coriaceous, oblong-lanceolate to oblanceolate-oblong, from 1 to 4 cm. long and from 0.6 to 1.2 cm. wide, obtuse or rounded, mucronate, somewhat rugulose above, prominently veined on the lower surface, their margins more or less recurved; both surfaces are clothed with soft, straight, appressed, gray to rufous hairs, and often mixed with these are spreading, glandular, gray hairs. The flowers are in clusters of one or more at the end of the branch, fragrant, pure white; occasionally an odd branch may have rose-colored flowers, or flowers striped with rose. The calyx is green and ample, the calyx-lobes are erect, lance-shaped, about 1 cm. long, often somewhat erose and more or less densely glandular-pubescent. The corolla is wide-funnel shape from 4 to 5 cm. long and broad; the stamens are usually 10 (occasionally 8) as long as the corolla or nearly so, and are exceeded by the pistil which is slightly exserted. The fruit is conic, with appressed setose hairs, and is shorter than the persistent calyx-lobes which surround it.

This Azalea is a favorite plant in Japan and has a place in almost every garden more especially from Tokyo southward, but its wild prototype was discovered only quite recently.¹ It is the "Jedogawa-tsutsuji with white flowers" mentioned by Kaempfer (*Amoen. Exot.* 848 [1712]), and by Thunberg was included under his *Azalea indica* as "Yedogawa Satsuke." It was cultivated in Java in the days of Burmann and without doubt was brought there from Nagasaki, Japan, by Dutch trading ships. In China it is grown in the gardens of the wealthy in many parts of the Empire, but how it got there we do not know. According to Lindley (*Bot. Reg.* X. t. 811 [1824]) it was first introduced into England to the nurseries of Mr. Brookes at Ball's Pond through his collector Joseph Poole who sent it from China in 1819. Like all introductions of this period it came, in all probability, either from Canton or Macao. Probably there were other importations from China in early times and during the last thirty years hundreds of thousands of plants have been sent from Japan to America and Europe. Its fragrant, white flowers and good constitution have made it a favorite and common plant in the greenhouses of the west. In fairly recent years it has been found to be more hardy than was once supposed, and it is now quite frequently seen as an outdoor garden shrub. I have not been able to ascertain the exact date of its introduction into America, but Hovey tells us that it was growing in the garden of J. P. Cushing, Belmont Place, Watertown, Massachusetts, in February, 1838. It is scarcely hardy near Boston but on Long Island, New York, and southward it is perfectly hardy. In Japan it is known as the "Shiro-yodogawa," that is "White Yodogawa."

I am familiar with this plant in the gardens of Japan and China as well as in those of Europe and America and in addition to the material enumerated I have seen the many old specimens preserved in Herb. Kew and Herb. Gray, which include several sent from Java many years ago. Every specimen in herbaria that I have seen is definitely stated to have come from cultivated plants. That it is an albino form of some species has long been suspected and it is satisfactory to be able to identify it at last. After an exhaustive study I am of the definite opinion that this old favorite garden plant is a white-flowered form of a plant wild near the banks of rivers on the island of Shikoku, which has lilac-purple flowers and to which Makino (in *Tokyo Bot. Mag.* XXII. 55 (1908) has given the name *R. ripense*.

¹ Millais states that "Carles collected it on the Seoul mountains in Korea." This is incorrect; the plant is *R. yedoense* var. *poukhanense* Nakai.

The specific name here employed to designate this species is the oldest valid name but it seems to have been quite overlooked. Burmann named the plant *Azalea rosmarinifolia* in 1768. This is the oldest name but unfortunately it was neglected for over a century and became invalidated under *Rhododendron* by the *R. rosmarinifolium* Vidal, a Philippine species, some years before it was taken up for Burmann's plant by Dippel.

A form with larger flowers than the type is: —

***Rhododendron mucronatum* f. *Noordtianum* Wilson, n. comb.**

Azalea japonica alba grandiflora van Noordt Pynaert in *Rev. Hort. Belg.* XXX. 133, t. (1904). — Reuter in *Möller's Deutsch. Gärtner-Zeit.* XX. 73, fig. (1905).

Azalea ledifolia var. *Noordtiana* Wittmack in *Gartenfl.* LIX. 310, fig. 32 (1910).

Rhododendron ledifolium var. *Noordtianum* Rehder in *Mitt. Deutsch. Ges.* XXIV. 225 (1916).

Rhododendron rosmarinifolium f. *Oriuki* Komatsu in *Tokyo Bot. Mag.* XXXII. [35] (1918).

Japan: Hondo, prov. Kawachi, Ikeda, cultivated, May 8, 1918, *E. H. Wilson*.

Pynaert states that this form was raised in the nursery of P. van Noordt & Sons, Boskoop, from seeds collected from some plants introduced from Japan. I can find no difference between it and the form long cultivated in Japan under the name of *O-riuki-tsutsumi* (large *Liuki* Azalea). It is considered in Europe to be a very hardy form of the type.

Very similar to the last-named form is: —

***Rhododendron mucronatum* "Mattapan" Wilson, n. f.**

Azalea "Mattapan" C. M. Hovey, *Mag. Hort.* XXVII. 190 (1861).

This plant was exhibited for the first time before the Massachusetts Horticultural Society on March 23, 1861, by C. M. Hovey, and described as "a new seedling, flowers white with a yellowish blotch on the upper petals which are fringed in the way of *A. crispiflora*." This handsome form is still in the Holm Lea collection and in other gardens round Boston, Mass. In some gardens this Azalea is known by the name of "America."

A form with double white flowers is: —

***Rhododendron mucronatum* f. *narcissiflorum* Wilson, n. comb.**

Rhododendron narcissiflorum Planchon in *Fl. des Serr.* IX. 82 (1854); in *Rev. Hort.* 1854, 67. — Hemsley in *Jour. Linn. Soc.* XXVI. 27 (1889). — Millais, *Rhodod.* 216 (1917).

Azalea narcissiflora Fortune apud Planchon in *Fl. des Serr.* IX. 82 (1854), as a synonym. — Hovey, *Mag. Hort.* XXII. 9 (1856).

Rhododendron ledifolium γ . *narcissiflorum* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 36 (*Rhodod. As. Or.*) (1870), in part. — Franchet & Savatier, *Enum. Pl. Jap.* I. 291 (1875). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 462 (1912). — Bean, *Trees and Shrubs Brit. Isl.* II. 366 (1914). — Rehder in *Bailey, Stand. Cycl. Hort.* V. 2944 (1916).

Azalea rosmarinifolia var. *narcissiflora* Rehder in *Bailey, Cycl. Am. Hort.* 123 (1900).

Rhododendron rosmarinifolium var. *narcissiflorum* Schneider, *Ill. Handb. Laubholz.* II. 504 (1911).

Rhododendron rosmarinifolium f. *narcissiflorum* a. *Shiromanyo* Komatsu in *Tokyo Bot. Mag.* XXXII. [36] (1918).

Japan: Kyushu, prov. Chikugo, Kurume, cultivated, May 3, 1918, *E. H. Wilson*.

Under the name of "Shiro-manyo-tsutsuji" this double-flowered form has been cultivated in Japan for a long time — probably centuries. It was introduced into England by Robert Fortune who sent plants from China to Messers Standish & Noble about 1850 or 1851. It was imported into this country by C. M. Hovey in 1855. Hovey (*Mag. Hort.* V. 25 [1839]) speaks of an *Azalea ledifolia alba pleno* having been introduced into Boston in 1838. The name suggests that the form *narcissiflorum* may have been in gardens prior to Fortune's introduction but I have been unable to find any further reference to the plant mentioned by Hovey. Millais suggests this may be a hybrid *R. indicum* × *R. linearifolium*? but it is difficult to understand why it should be regarded as anything but the common *R. mucronatum* G. Don (*Azalea ledifolia* Hook.) with double flowers. The habit, fragrance, calyx and pubescence are identical. Maximowicz refers the double rose-colored form (f. *plenum* Wils.) to this f. *narcissiflorum* and Millais does the same but for horticultural purposes it is best kept distinct.

A fine garden form of the type is: —

***Rhododendron mucronatum* f. *sekidera* Wilson, n. comb.**

Rhododendron rosmarinifolium f. *Sekidera* Komatsu in *Tokyo Bot. Mag.* XXXII. [35] (1918).

Rhododendron Sekidera (*Rhod. indicum* × *Rhod. Kaempferi*) Komatsu in *Tokyo Bot. Mag.* XXXII. [46] (1918).

Azalea magnifica Hort. Yokohama Nursery Co. *Cat.* 1918-19, p. 19.

Azalea indica roseum Hort. Andorra Nurseries, May, 1919.

Azalea japonica grandiflora superba Hort. Andorra Nurseries, May, 1919.

Azalea "Damask Rose" Hort. Andorra Nurseries, May, 1919.

Japan: prov. Musashi, cultivated, May 21, 1914, *E. H. Wilson* (No. 6670).

This very handsome plant I saw cultivated in the grounds of the Yokohama Nursery Company. The flowers are as large as those of f. *Noordtianum*, pure white with the dorsal lobes spotted and splashed with rose, occasionally a flower is rose-colored; the stamens vary in number from 6 to 10 but are usually 10. Under the names *Azalea indica roseum* and *Azalea* "Damask Rose" slightly inferior forms were sent to us for determination last year. The true variety is cultivated by Mr. John S. Ames, North Easton, Mass., who imported it in 1917. In 1919 I brought back plants from Japan to the Arnold Arboretum. This most desirable addition to our gardens will probably prove to be as hardy as the typical white-flowered *R. mucronatum* G. Don.

A spontaneous variety and undoubtedly the phylogenetic type of the species is: —

***Rhododendron mucronatum* var. *ripense* Wilson, n. comb.**

Rhododendron ripense Makino in *Tokyo Bot. Mag.* XXII. 55 (1908). — Schneider, *Ill. Handb. Laubholz.* II. 503 (1911). — Komatsu in *Icon. Pl. Koisi-kav.* II. 71, t. 120 (1914); in *Tokyo Bot. Mag.* XXXII. [12] (1918).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

in western gardens, having been sent by Joseph Poole from China to Samuel Brookes, nurseryman, Ball's Pond, Newington Green, England, in 1819. The flower is rose-purple, fragrant, quite, or only partially double, and with or without green rudimentary leaves showing in the centre of the flower. Hovey tells us that this plant was cultivated by Mrs. Hibbert and exhibited before the Pennsylvania Horticultural Society on February 19, 1839.

Rhododendron linearifolium Siebold & Zuccarini in *Abh. Akad. Münch.* IV. pt. III. 131 (*Fl. Jap. Nat.* II. 7) (1846).—Miquel in *Ann. Mus. Lugd.-Bat.* I. 34 (1863); II. 165 (1864–66); *Prol. Fl. Jap.* 97 (1866–67). — Maximowicz in *Mém. Acad. Sci. Pétersbourg*, sér. 7, XVI. No. 9, 34 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 290 (1875). — Boissier in *Bull. Herb. Boiss.* V. 918 (1897). — Schneider, *Ill. Handb. Laubholz.* II. 504, figs. 330 g–k, 331 f. (1911). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463 (1912). — Bean, *Trees and Shrubs Brit. Isl.* II. 366 (1914). — Millais, *Rhodod.* 203 (1917).

Azalea linearifolia Hooker f. in *Bot. Mag.* XCV. t. 5769 (1869).

Rhododendron macrosepalum var. *linearifolium* Makino in *Tokyo Bot. Mag.* XXII. 55 (1908). — Komatsu in *Tokyo Bot. Mag.* XXXII. [34] (1918).

Rhododendron linearifolium a. *linearifolium* Makino in *Tokyo Bot. Mag.* XXVII. 108 (1913).

Cultivated: November, 1867, February, 1869, ex Hort. Standish (Herb. Kew); 1876, ex Hort. Hunnewell (Herb. Gray).

This is a monstrous garden form cultivated in Japan but rather rare. The leaves are narrow, strap-shaped, and the corolla is divided into narrow, ligulate segments. An analogous plant is *R. indicum* f. *polypetalum* Wils. Maximowicz says it was introduced into the gardens of Tokyo from the alpine regions of Sendai in northern Japan. This is a mistake. It is unknown in a wild state and was first known from the neighborhood of Nagasaki in Kyushu the southern island of Japan. Its Japanese name is "Seigai-tsutsuji." It was introduced into England by Messers Standish with whom it flowered in November, 1867, and again in February, 1869. It is well figured in the *Botanical Magazine*. From England it appears to have been sent to this country for it flowered in the garden of Mr. H. H. Hunnewell, Wellesley, Mass., in 1876. According to Bean it is quite hardy at Kew. It is not hardy in the Arnold Arboretum and I do not know whether it is now cultivated in America.

Rhododendron linearifolium var. **macrosepalum** Makino in *Tokyo Bot. Mag.* XXVII. 108 (1913).

Rhododendron macrosepalum Maximowicz in *Gartenfl.* XIX. 258, t. 662 (1870); in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 31 (*Rhodod. As. Or.*) (1870); in *Bull. Acad. St. Pétersbourg*, sér. III. XV. 227 (*Mél. Biol.* VII. 335) (1871). — Franchet & Savatier, *Enum. Pl. Jap.* I. 290 (1875). — Dippel, *Handb. Laubholz.* I. 420, fig. 271 (1889). — Makino in *Tokyo Bot. Mag.* IV. 151 (1890); XXII. 55 (1908). — Boissier in *Bull. Herb. Boiss.* V. 918 (1897). — Schneider, *Ill. Handb. Laubholz.* II. 503, figs. 330 e, 331 a–b

- (1911). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463 (1912). — Millais, *Rhodod.* 205 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [13] (1918).
Azalea macrosepala K. Koch, *Dendr.* II. pt. 1, 180 (1872). — O. Kuntze, *Rev. Gen.* II. 387 (1891).
Rhododendron linearifolium β . *macrosepalum* forma *a. genuinum* Makino in *Tokyo Bot. Mag.* XXVII. 108 (1913).

Japan: Kyushu, prov. Hizen, roadside, Nagasaki to Mogi, November 12, 1903, *C. S. Sargent*. Shikoku, prov. Tosa, thickets up to 300 m. alt. November 20, 1914, *E. H. Wilson* (No. 7794). Hondo, prov. Settsu, near Kobe, April, 1875, *H. N. Moseley* (Herb. Kew); prov. Kawachi, Yomomoto and Ikeda, May 8, 1918, *E. H. Wilson* (No. 10,355); prov. Yamato, Koyasan, August 26, 1907, *U. Faurie* (No. 124); Yoshino, April 22, 1914, *E. H. Wilson* (No. 6574, in part); prov. Yamashiro, Kyoto, Temple Garden, October 16, 1892, *C. S. Sargent*; Kyoto, *J. Rein*, ex Mus. Bot. Berol. (Herb. Bur. Sci. Manila); prov. Omi, mountains round Lake Biwa, April 30, 1892, *U. Faurie* (No. 7820, Herb. Kew); prov. Mino, Pine-barrens near Gifu, October 20, 1892, *C. S. Sargent*; prov. Mikawa, Pine-woods, Futagawa, May 9, 1918, *E. H. Wilson* (Nos. 10,346, 10,348, 10,349); prov. Shimotsuke "alpe Niko," 1866–74, *L. Savatier* (No. 766, Herb. Kew).

This is the Mochi-tsutsuji (Glandular Azalea) of Japan, where it is common in central Hondo at low altitudes from near Osaka north to Hamamatsu in Totomi province; also in Tosa province, Shikoku. It is partial to gravelly soil and dry situations such as Pine-woods and open thickets afford. A social plant it occurs in great numbers either alone or with *R. obtusum* var. *Kaempferi* Wils. At Futagawa, for instance, in Pine-woods these two species grow in thousands and in early May when in full blossom they are conspicuous from the railway carriage windows. The specimen from Nagasaki is probably from an escaped plant and there is no other record of its growing in Kyushu. North of Totomi province I have not seen it wild, and I think there is some mistake about it being indigenous in the Nikko region, Maximowicz's statements and Savatier's specimens notwithstanding. It is not mentioned by Matsumura in his *Plants of Nikko* nor by Hayata in his *Vegetation of Mt. Fuji*.

As usually seen *R. linearifolium* var. *macrosepalum* is a low, rather laxly branched shrub spreading in habit and less than one metre high but in Shikoku I saw it full three metres tall. It has the usual dimorphic leaves and these are mostly deciduous. The shoots are terete and with the petioles are densely clothed with gray to gray-brown pilose (some of them gland-tipped) hairs and a few flattened, spreading bristles which disappear in the second season; the winter-buds are ovoid, acute, densely covered with strigose, yellow-brown hairs. The leaves are membranous, scattered on the free-growing shoots and crowded on others, short-petioled, lanceolate-ovate to ovate-elliptic, sometimes lanceolate or ovate, short acuminate, acute or merely mucronulate; those immediately beneath the winter-buds oblanceolate to oblanceolate-oblong obtuse, or rounded, usually mucronulate, the under surface clothed with soft, spreading, gray to gray-brown glandular hairs, the upper surface sparsely glandular hairy, slightly rugulose and hispid; dull green during summer changing to rich vinous purple in the autumn. The flowers are

fragrant, from 2 to 10 in terminal umbels, very variable in size and form. The pedicels are from 2 to 4 cm. long, straight or spreading, densely glandular and pilose. The calyx is green, glandular-pubescent and very variable in size, the strap-shaped segments being from 1 to 3 cm. long and occasionally exceeding the rose-pink to red-purple corolla, which is broad funnel-shape, sometimes 5 cm. but often not more than 3 cm. across; it is glabrous without, and normally larger than the calyx. Stamens normally 5, but at Ikeda I gathered specimens with from 6 to 10 stamens; they are shorter than the pistil and included within the corolla except in abnormally small flowers. The fruit is narrow-ovoid, about 1 cm. long, glandular-pubescent, purplish-brown in color and subtended by the persistent calyx the lobes of which are reflexed, brown without and wine-purple on the inside. There can be no question of this being the phylogenetic type of the species and it is unfortunate that it should have to rank as a variety. Even in a wild state it is a very variable plant, and a number of forms have been detected and are cultivated in Japanese gardens; these are dealt with hereafter. The calyx, pedicels and innermost scales of the flower-buds are always glandular, usually densely so and commonly sufficiently viscid to hold gnats and small flies captive, but this viscosity like everything else in this plant is variable. The flowers are pleasantly fragrant and some of the color forms are very handsome. When the flowers have a comparatively small corolla and very long calyx-lobes the effect is curious. This *Azalea* was introduced into Petrograd by Maximowicz who sent seeds in 1863. It flowered for the first time in 1870 and is well figured in *Gartenflora*. Maximowicz claims that the seeds and his two plants came from the Nikko alps but if this is not a mistake it is strange that no other collector has seen the plant in this locality which is fully two hundred miles north of where it has been seen wild by subsequent observers. Further, no one else has found it in alpine regions anywhere in Japan. However, there is no doubt that the plant under discussion is Maximowicz's species. What happened to the original introduction I do not know but neither Bean nor Millais mention its cultivation in England. In 1914 I sent to the Arnold Arboretum seeds of this plant from Shikoku some of which were sent to England. Plants raised from this seed in the Arnold Arboretum have not proved very hardy.

A form with double flowers is: —

Rhododendron linearifolium* var. *macrosepalum* f. *dianthiflorum
Wilson, n. comb.

Azalea dianthiflora Carrière in *Rev. Hort.* 1889, 391; 1891, 60, fig. 18, t.
Rhododendron dianthiflorum Millais, *Rhodod.* 155 (1917).

Japan: Hondo, prov. Mikawa, Pine-woods, Futagawa, May 9, 1918, *E. H. Wilson* (No. 10,350).

Cultivated: ex Hort. James Veitch & Sons, June 28, 1904 (Herb. Kew).

I have seen at Futagawa one or two plants growing wild with double flowers. The pedicels are more spreading than in the type and the calyx-lobes are variable in length. Carrière's figures give a good idea of this form except that the color is brighter and more pleasing than the plate shows. According to Carrière it was introduced into France from Japan in 1889 by M. Wiesener, of Fontenay-aux-Roses. The specimen in Herb. Kew shows that it was cultivated by Messers Veitch in 1904. Bean does not mention it and Millais's account is abbreviated from Carrière's description. I have no knowledge of its being in the gardens of this country.

Another spontaneous form is: —

Rhododendron linearifolium var. **macrosepalum** f. **decandrum** Wilson, n. f.

Japan: Hondo, prov. Kawachi, Yomomoto, roadside, Pine-woods, May 8, 1918, *E. H. Wilson* (No. 10,355a, type); Ikeda, cultivated, May 8, 1918, *E. H. Wilson*; prov. Yamato, near Yoshino, dry woods, April 22, 1914, (No. 6574, in part).

This differs from the type in having 10 stamens; the Ikeda specimens have from 6 to 10 stamens and that from Yoshino (a fragment) has 6 only. The typical var. *macrosepalum* is common round Yomomoto where it is known as the Yama-tsutsuji (Mountain Azalea) and growing with it I found occasional plants with 10 or from 6 to 10 stamens in each flower. I have seen none with from 5 to 10 stamens and I do not remember whether the fragment from near Yoshino came from a bush of the ordinary 5-stamen type.

An anomalous form is: —

Rhododendron linearifolium var. **macrosepalum** f. **rhodoroides** Makino in *Tokyo Bot. Mag.* XXVII. 109 (1913).

Rhododendron macrosepalum β . *rhodoroides* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 31 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 290 (1875). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463 (1912). — Komatsu in *Tokyo Bot. Mag.* XXXII. [34] (1918). *Rhododendron ledifolium* var. *Kochozoro* Komatsu apud Makino in *Tokyo Bot. Mag.* XXVII. 109 (1913), as a synonym.

Japan: Hondo, prov. Kawachi, cultivated, Yamomoto and Ikeda, May 8, 1918, *E. H. Wilson*; prov. Musashi, Okbo, cultivated, May 18, 1909, *K. Ōnuma*; Hatagaya, cultivated, April 29, 1914, *E. H. Wilson* (No. 6596).

This is a curious form often seen in Japanese gardens. It has a deeply cleft 5-partite corolla only slightly longer than the calyx-lobes, and greenish-white shaded with rose, exserted stamens with pale purple anthers exceeded in length by the style. Maximowicz raised seedlings from this plant and states that in every instance they reverted to typical var. *macrosepalum*. Makino gives "Kocho-zoro" as the Japanese name of this plant; in the nursery district near Osaka it is known as "Seikan-tsutsuji."

Another anomalous form is: —

Rhododendron linearifolium var. **macrosepalum** f. **hanaguruma** Makino in *Tokyo Bot. Mag.* XXVII. 109 (1913).

Rhododendron ledifolium var. *leucanthum* f. *Hanaguruma* Komatsu apud Makino in *Tokyo Bot. Mag.* XXVII. 109 (1913), as a synonym. *Rhododendron macrosepalum* var. *Hanaguruma* Makino in *Tokyo Bot. Mag.* XXVII. 109 (1913), as a synonym. — Komatsu in *Tokyo Bot. Mag.* XXXII. [35] (1918). *Rhododendron linearifolium* var. *macrosepalum* f. *polypetalum* Makino in *Tokyo Bot. Mag.* XXVII. 109 (1913), as a synonym.

Japan: Hondo, prov. Kawachi, cultivated at Yomomoto and Ikeda, May 8, 1918, *E. H. Wilson*; prov. Rikuzen, Sendai, cultivated, May 27, 1888, *U. Faurie* (No. 2300, Herb. Kew).

Cultivated: ex Hort. Ball, March, 1879 (Herb. Kew).

In this form the corolla is rose-purple in color, deeply cleft and 5-partite, and the stamens are usually shorter than the corolla-lobes which are from 2.5 to 4 cm. long. Makino gives "Hana-guruma" as the Japanese name; in the nursery district near Osaka it is known as "Oyeyama-tsutsuji." The specimen in Herb. Kew proves that at one time it was in cultivation in England.

Komatsu (in *Tokyo Bot. Mag.* XXXII. [34] [1918]) enumerates three other forms ("Usuyo," "Amaga-shita" and "Suruga-momyo") of this species, but these are unknown to me.

These are all the species of the section Tsutsutsi that are named or of which I have seen sufficient material. There is, however, in this herbarium specimens (ex Herb. Hongkong Nos. 7030, 5612) from Taimo-shan, Kowloon, Kwangtung province, of a curious Azalea. It has narrow, linear-lanceolate, acuminate leaves and inconspicuous subsessile flowers with the corolla deeply divided into five narrow segments. It is in every way anomalous, and probably a monstrous condition of some species not yet described, or possibly of the widespread *R. Simsii* Planch. We have, also, a leafy fragment, collected in Kweichow province by Dr. Handel-Mazzetti (No. 221, July 11, [1917]) which probably belongs to a new species related to *R. Oldhamii* Maxim.

SECT. II. SCIADORHODION REHD. & WILS., n. sect.¹

Rhododendron section *Tsutsutsi* G. Don, *Gen. Syst.* III. 847 (1834), as to species nos. 30, 31. — Rehder & Wilson in Sargent, *Pl. Wilson.* I. 547 (1913), in part.

Rhododendron subgenus *Tsutsia* Planchon in *Fl. des Serr.* XI. 75 (1854); in *Rev. Hort.* 1854, 43, as to *R. Farrerae*.

Rhododendron section *Azalea* Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 24 (*Rhodod. As. Or.*) (1870), in part.

Azalea subgenus *Euazalea* K. Koch, *Dendr.* II. pt. 1, 179 (1872), as to *A. reticulata*.

Rhododendron section *Rhodora* Dippel, *Handb. Laubholz.* I. 418 (1889), in part.

Shrubs with rigid, usually verticillate, often ascending-spreading branches and glabrous or villose shoots without bristle-like hairs. Flowers from terminal buds, with leafy shoots from axils of lower scales of the same bud. Leaves deciduous, in whorls of 3 to 5 at the end of the branchlets, on vigorous shoots scattered, not dimorphic, usually rhombic to obovate, rarely ovate, without appressed bristle-like hairs. Corolla rotate-campanulate, rarely rotate-funnel-form; stamens from 6 to 10 (rarely 5 by abortion), unequal, sometimes declinate. Fruit oblong-ovoid to cylindrical, rarely conic-ovoid, often furrowed, villose or glandular, never strigose.

This section is found only in eastern Asia, where it is represented by 6 species.

Rhododendron Farrerae Tate apud Sweet, *Brit. Flow. Gard.* ser. 2, I. t. 95 (1831). — G. Don, *Gen. Syst.* III. 846 (1834). — De Candolle, *Prodr.* VII. 725, pt. 2 (1839). — Planchon in *Fl. des Serr.* XIX. 80 (1854); in *Rev. Hort.* 1854, 64. — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 25 (*Rhodod. As. Or.*) (1870). — Hance in *Jour. Linn. Soc.* XIII. 110 (1873). — Hemsley in *Jour. Linn. Soc.*

¹ Sectio inter sect. *Tsutsutsi* et *Rhodoram* medium tenens, a priori indumento piloso vel villoso pallido, non strigoso et fusco, foliis semper deciduis satis magnis, corolla rotato vel rotato-campanulata, ovario villoso vel glabro, non strigoso, a posteriori gemmis terminalibus flores et folia simul proferentibus bene differt, ab utraque sectione foliis in apice ramulorum subverticillatim congestis quasi umbellam vel umbraculum (graece *σκιάς*, *σκιάδος*, unde nomen) formantibus satis magnis plerumque late obovatis vel rhombico-ovatis distat.

In Sargent, *Plantae Wilsonianae*, I. 548 (1913), we suggested that with their whorled leaves the species here enumerated formed a well-marked group easily distinguished from *Tsutsutsi*, although agreeing in the character of the winter-buds. Further investigation has brought to light other differences and we are of the opinion that it is best to place them in a distinct section.

A. R. & E. H. W.

XXVI. 23 (1889); in *Kew Bull. Misc. Inform.* 1907, 245. — Dunn & Tutcher in *Kew Bull. Misc. Inform.* add. ser. X. 155 (*Fl. Kwangtung and Hongk.*) (1912). — Millais, *Rhodod.* 163 (1917).

Azalea squamata Lindley in *Jour. Hort. Soc. Lond.* I. 152 (1846); in *Bot. Reg.* XXXIII, t. 3 (1847). — Walpers, *Ann.* I. 481 (1848-49). — Bentham, *Fl. Hongk.* 201 (1861).

Azalea Farrerae K. Koch, *Dendr.* II. pt. 1, 178 (1872). — O. Kuntze, *Rev. Gen.* pt. 2, 387 (1891).

Azalea squamosa O. Kuntze, *Rev. Gen.* pt. 2, 387 (1891), as a synonym.

Rhododendron Farrerae δ . *typicum* Diels in *Bot. Jahrb.* XXIX. 514 (1900).

China: prov. Kwangtung, Loh-fau-san, August 21, 1917, C. O. Levine (ex Canton Christian College Herb. No. 1533).

This species seems to be peculiar to Hongkong and the neighboring islands and to the adjacent parts of Kwangtung province. It is a low, densely branched shrub with short, rigid, shining brown, verticillate branchlets clothed with appressed, straight and villose hairs when young, becoming glabrous and gray in their second year. The leaves are crowded, usually in threes at the end of the branches, subcoriaceous, deciduous or semi-persistent, glabrescent, ovate, from 2 to 3 cm. long and from 1.2 to 2 cm. wide, acute or sub-acute, with a short mucro, and a rounded base; they are dark green above and pallid below, reticulate, with the principal veins impressed above and raised below. The petioles are from 1 to 3 mm. long and densely villose. The winter-buds are small, broadly ovoid, and are densely clothed with gray or rufous pubescence. The flowers appear before the leaves unfold, either solitary or in pairs at the end of the branches. The corolla is pale to deep rose-color with red-purple spots, from 4 to 5 cm. in diameter; it has a short, narrow, funnel-shape tube and spreading, undulate lobes. The ring-like calyx is minutely 5-toothed, and pubescent; and the pedicel and ovary are thickly clothed with appressed and villose, ferruginous or gray hairs. The stamens are from 8 to 10 of unequal length and shorter than the corolla, and are equalled or slightly overtopped by the slender, curved style. The fruit is conic-ovoid, from 0.5 to 1.5 cm. long and from 0.8 to 1 cm. in diameter and densely villose, and is borne on a stout, curved, villose pedicel about 1 cm. long.

I have seen this *Azalea* growing wild in Hongkong and as Hemsley has pointed out it is a very distinct species. It is characterized by its small, ovate leaves, its very short villose petioles and by its relatively large fruit. It was first introduced into England in 1829, by Captain Farrer of the East India Company's ship "Orwell." It was reintroduced by Robert Fortune who in 1844 sent it to the garden of the Horticultural Society of London where it flowered and was renamed by Lindley. I have no knowledge of its having been in cultivation in America, and Millais states that it is a rare plant in English gardens, and hardy only in the south.

Rhododendron Mariesii Hemsley and Wilson in *Kew Bull. Misc. Inform.* 1907, 244. — Hutchinson in *Bot. Mag.* CXXXIV. t. 8206 (1908). — Schneider, *Ill. Handb. Laubholz.* II. 496, fig. 327 l-m (1909). — Rehder & Wilson in Sargent, *Pl. Wilson.* I. 548 (1913). — Bean, *Trees and Shrubs Brit. Isl.* II. 377 (1914). — Millais, *Rhodod.* 207 (1917).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

ovate-lanceolate, with an acute mucronate apex and cuneate base, but occasionally they are broad-ovate or elliptic. The petioles are from 0.4 to 1.2 cm. in length, glabrescent, purplish, and often slightly glaucescent on the under-side. The flowers are usually in pairs at the end of the naked branchlets, but are sometimes solitary, and occasionally are in 4- or 5-flowered clusters; the corolla is from 4 to 5 cm. across, spotted with red-purple on the posterior lobes; it has a short, narrow, funnel-shape tube and deeply cleft, spreading lobes. The pedicels are villose; the minute, 5-lobed calyx and the ovary are clothed with gray or yellow-brown, straight appressed hairs. The 10 stamens are of nearly equal length and as long as the corolla, the filaments are glabrous, curved and the anthers purple. The style is longer than the stamens and is curved upward and has a capitate, slightly lobed stigma. The fruit is cylindric, from 1 to 1.5 cm. long, slightly furrowed, and clothed with gray or yellow-brown, villose pubescence, and is borne on a straight pedicel.

From those species with which *R. Mariesii* has been confused it is readily distinguished by its leaves. The flowers are of a different color from those of the Japanese species, the corolla is differently shaped and there are other distinctions of importance. In Tokyo I saw the types of Hayata's two species cited and could find nothing by which to distinguish them from *R. Mariesii*.

This Chinese plant was discovered by Fortune, probably in Chekiang province, and afterward by Charles Maries on the Lushan mountains behind Kiukiang in the spring of 1878. It was introduced into Kew gardens through seeds sent from the neighborhood of Ichang in 1886 by Augustine Henry; in 1900 I sent seeds to Messers Veitch from south of Ichang. This Azalea flowered for the first time in the Temperate House at Kew in April, 1907. Bean does not consider it to be hardy. I have not heard of its being cultivated in America.

Rhododendron Weyrichii Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 26, t. 2, fig. 1-6 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 288 (1875). — Makino in *Tokyo Bot. Mag.* XVIII. 48 (1904). — Schneider, *Ill. Handb. Laubholz.* II. 495 (1909), in a note. — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 464 (1912). — Nakai, *Rep. Fl. Quelpaert Isl.* No. 999 (1915); in *Tokyo Bot. Mag.* XXXI. 243 (1917); *Fl. Syl. Kor.* VIII. 44, t. 16 (1919). — Millais, *Rhodod.* 261 (1917), excluding the Chinese plant. — Komatsu in *Tokyo Bot. Mag.* XXXII. [10] (1918).

Azalea Weyrichii O. Kuntze, *Rev. Gen.* pt. 2, 387 (1891).

Rhododendron shikokianum Makino in *Tokyo Bot. Mag.* VI. 43 (1892); in IX. 111 (1895).

Japan: Shikoku, prov. Tosa, round Kochi, sea-level to 800 m. altitude, November 17, 1914, *E. H. Wilson* (No. 7813). Kyushu, prov. Hizen, near Nagasaki, 1862, *R. Oldham* (No. 511, Herb. Gray).

Korea: Quelpaert Island, between Saishu and the Monastery, 400-650 m. October 30, 1917, *E. H. Wilson* (No. 9411); same island, Hallai-san, October, 1906, May, 1907, *U. Faurie* (Nos. 661, 1863 in part, 1864); same place, May, 1909, April 14, June 6, 1908, *E. Taquet* (Nos. 2970, 4678, 1089).

This little known species has a very curious distribution. On the island of Shikoku it is common round Kochi in Tosa province but does not grow on the mainland of Kyushu, but it is found on the adjacent islands of Amakusa and Goto south and west of Nagasaki. I do not know whether it grows on the island of Tsushima in the Japan Sea — probably it does — but it is common on the Korean island of Quelpaert although unknown on the mainland of Korea. In Shikoku I found it to be fairly plentiful from near sea-level up to an altitude of 800 m. growing in thickets and open woods. On Hallai-san, an extinct volcano on Quelpaert Island, between 200 and 1000 m. altitude, and especially in thickets by the sides of water-courses and on old lava flows, it is common.

This Azalea is a vigorous growing shrub from 1 to 5 m. tall, often decidedly tree-like in habit, with moderately stout ascending to spreading often whorled branches. These are yellow-brown, clothed with ferrugineous tomentum when young, becoming glabrous, gray and finally dark purple in color. The winter-buds are characteristic, being conic and sub-acute and clothed with yellow-brown, matted and straight hairs and are unlike those of other species. The leaves are distinctly petiolate, chartaceous, shining green and deciduous, changing to vinous purple in the autumn; on the vigorous shoots the leaves are scattered and in pairs or whorls but on the branchlets they are clustered in whorls of 3, or in pairs; they vary from broad-ovate to semiorbicular or to rhombic, and in size from 3.5 to 8 cm. long and from 2 to 6 cm. wide; they are abruptly subacute or obtuse at the apex which is mucronate, the base is broadly cuneate or rounded. The principal veins are impressed above and slightly raised on the lower surface which is venulose. When young the petioles and both surfaces of the leaf are covered with rufous-brown pilose hairs but these soon disappear. The flowers are from 2 to 4 in terminal umbels, and open before or at the same time as the leaves unfold. The pedicels, calyx and ovary are densely clad with rufous-brown pilose hairs. The corolla is red, almost brick-red, from 3.5 to 6 cm. across, funnel-shape, with a rather narrow expanding tube and spreading lobes. The calyx is merely a ring with five minute teeth. The stamens vary from 6 to 10, and are of unequal length, and like the curved style and its capitate stigma are included. The fruit is cylindric or oblong-ovoid, from 1.5 to 2 cm. long, oblique, furrowed and obtusely angled and clothed with appressed, villose pubescence. This species was discovered on the Goto Islands in 1853 by Dr. Heinrich Weyrich, a surgeon on the Russian warship "Vostok" which formed part of Admiral Putiatin's fleet. On Oldham's specimen the locality cited is Nagasaki but this is a mistake; it probably came from the Amakusa Islands south by west of that city. Makino first found it in Shikoku and Père Faurie on Quelpaert Island. It was unknown in gardens until 1914 when I sent the seeds from Shikoku to the Arnold Arboretum which were distributed in America and Europe. Plants raised in this Arboretum have not proved hardy but in another Massachusetts garden where they have received the protection of a cold frame they have flowered. In 1917 I collected seeds on Quelpaert and hope that they will produce a hardier type. I have never seen this Rhododendron in flower in a wild state and only a late flower or two on cultivated plants. The color of the flowers is good and where it proves hardy this species should be a welcome addition to gardens.

Rhododendron reticulatum D. Don apud G. Don, *Gen. Syst.* III. 846 (1834). — De Candolle, *Prodr.* VII. pt. 2, 727 (1839).

Rhododendron dilatatum Miquel in *Ann. Mus. Lugd.-Bat.* I. 34 (1863-64); in II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67).

Rhododendron rhombicum Miquel in *Ann. Mus. Lugd.-Bat.* II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67). — Regel in *Gartenfl.* X. XVII. 225, t. 586

(1868). — Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7 XVI. No. 9, 26 (*Rhod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 288. (1875). — Hooker f. in *Bot. Mag.* CXIII. t. 6972 (1887). — Bean in *Gard. Chron.* ser. 3, XX. 38, fig. 9 (1896); *Trees and Shrubs Brit. Isl.* II. 375, fig. (1914). — Boissier in *Bull. Herb. Boiss.* V. 918 (1897). — Dallimore in *Garden*, LXXII. 267, fig. (1908). — Mottet in *Rev. Hort.* 1909, 79, fig. 28. — Schneider, *Ill. Handb. Laubholz.* II. 497, figs. 327 f–g, 328 d–e (1911). — W. Watson, *Rhodod. and Azaleas*, 111 (1911). — Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463. (1912). — Rehder in *Bailey, Stand. Cycl. Hort.* V. 2943 (1916). — Millais, *Rhodod.* 235 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [10] (1918).

Azalea reticulata K. Koch, *Dendr.* II. pt. 1, 179 (1872).

Azalea dilatata O. Kuntze, *Rev. Gen.* pt. 2, 387 (1891).

Azalea rhombica O. Kuntze, *Rev. Gen.* pt. 2, 387 (1891). — Rehder in *Bailey, Cycl. Am. Hort.* I. 122 (1900).

Rhododendron dilatatum β . *decandrum* Makino in *Tokyo Bot. Mag.* VII. 134 (1893).

Rhododendron decandrum Makino in *Jour. Jap. Bot.* I. 21 (1917).

Rhododendron Wadanum Makino in *Jour. Jap. Bot.* I. 21 (1917).

Japan: Kyushu, prov. Satsuma, Kagoshima in gardens, March 4, 1914, *E. H. Wilson* (No. 6158); Togo, in a garden, March 14, 1914, *E. H. Wilson* (No. 6266); prov. Osumi, Mt. Kirishima, alt. 100–1000 m. 1917, *Z. Tashiro*; Nishi-Kirishima, alt. 1000 m. March 11, 1914, *E. H. Wilson* (No. 6241); prov. Hizen, Nagasaki, Mt. Yuwaja, 1863, *C. Maximowicz* (Herb. Gray). Shikoku, prov. Tosa, Nano-kawa, May 12, 1889, *K. Watanabe* (Herb. Gray). Hondo, prov. Choshu, between Habu and Asa, May 7, 1918, *E. H. Wilson* (No. 10,369); prov. Aki, Miyajima, November 28, 1914, *E. H. Wilson* (No. 7842); prov. Yamato, Yoshino, April 22, 1914, *E. H. Wilson* (No. 6573); prov. Yamashiro, Kyoto, April 22, 1914, *E. H. Wilson* (No. 6568); prov. Shinano, Ogawa, September 5, 1905, *J. G. Jack*; Nojiri to Nakatsugawa, September 6, 1905, *J. G. Jack*; slopes of Mt. Komaga-take, October 24, 1892, *C. S. Sargent*; prov. Suruga, lower slopes of Mt. Fuji, alt. 800–1000 m. May 8, 1914, *E. H. Wilson* (No. 6640); prov. Sagami, Hakone Mts. Kintoki, May, 1876, *James Bissett* (No. 237, Herb. Kew); prov. Shimotsuke, Nikko region, alt. 1000–1800 m. May 18, October 24, 1914, *E. H. Wilson* (Nos. 6705, 7694), shores of Lake Chuzenji, September 3, November 7, 1892, *C. S. Sargent*; same locality, October 25, 1905, *J. G. Jack*. Hokkaido, prov. Hidaka, Shoya village, August 17, *E. Tokubuchi*. Province unknown, “Mimasaka,” May 10, 1904, *S. Arimoto* (Herb. Gray).

Cultivated: Arnold Arboretum (No. 3427–1).

This is a common plant throughout the greater part of Japan. In Kyushu, Shikoku and Hondo it is everywhere abundant; in Hokkaido it is rather rare and

finds the northern limits of its range in Hidaka province. On volcanic ash, lava-flows and on cliffs in many places, as in the Nikko region, lower slopes of Mt. Fuji and on the Hakone mountains for example, it is extraordinarily abundant both in open country, in thickets, on the margins of woods and as undergrowth in thin forests. It is in flower from early in March to late in June according to latitude and altitude and produces in great abundance its rose- to red-purple or rich magenta-colored blossoms. The richest colors are on plants in the open, and in dense shade the color is quite pale. Nowadays magenta is not a popular color in the west but massed and alone save for the varying shades of green the flowers of this *Azalea* are impressive. The plant forms a much-branched bush or bushy tree from 1 to 8 metres tall, with numerous erect or erect-spreading verticillate, slender but rigid branches, yellow-brown and sparsely or densely hairy when young, usually soon becoming glabrous, and in the second year pale gray. In the shade the branches are more sparse and the plants less compact and shapely in habit. The winter-buds are conic and acute, with rufous- or gray-brown, villose and ciliate scales, the inner ones being viscid. The leaves unfold as the corollas fall, and are clustered in threes or in pairs at the end of the branchlets, and vary in shape from broad-ovate to rhombic or occasionally to ovate, and terminate in a short mucro. In size they vary from 3 to 6 cm. in length and from 2.5 to 5.5 cm. in width, with petioles from 0.5 to 1.5 long. In the shade the leaves are membranous but in the open they are firm and the reticulation is more prominent; they are dark green above, pale even sub-glaucous below, and in the autumn change to vinous or blackish purple or occasionally to yellow with splashes of red-purple. The principal veins are impressed above and raised and usually more or less villose on the under surface. When young the leaves are densely clothed with long, or with short and long, appressed, gray to yellow-brown pilose hairs which disappear early from the upper and entirely or partially from the lower surface. The petiole is flattened and usually villose, but may be glabrous and glaucescent. The flowers are borne at the end of the shoots in pairs or solitary or in clusters of three or four. The corolla is from 3.5 to 5 cm. across usually unspotted, with a short tube and spreading lobes which are often divergent and give the corolla a two-lipped appearance. The pedicels are clothed with gray or brownish appressed hairs, and the minute, 5-toothed calyx is ciliate, villose or glabrescent. The stamens are normally 10, in two series, the longer equalling the corolla in length, the shorter half as long, the filaments curving upward; the anthers are purple. The ovary is covered with short straight hairs, or is lepidote, or lepidote and hairy, and the style is hairy or glabrous; the stigma capitate and minutely 5-lobed. The fruit is curved, cylindrical, from 1 to 1.5 cm. long, furrowed, lepidote, glabrous, or clothed with gray or brown soft hairs.

I have gone very minutely into the degree of variation exhibited by this plant for I realize that those who compare the two figures in the *Botanical Magazine* will find it difficult to appreciate that they represent the same species and really only differ in the number of stamens. In the array of specimens before me and cited above the variations mentioned are all shown and neither among them nor in the field have I been able to distinguish two species. The distinguishing characters emphasized by Maximowicz are inconstant except the number of stamens and I am doubtful if careful examination among many plants would not reveal flowers with five and others with ten stamens on the same plant. I have 10-stamened flowers with a lepidote ovary and style, with a hirsute and lepidote ovary and glabrous style; I have 5-stamened flowers which show exactly the same variation. On both forms I find glabrous leaves and leaves more or less pubescent below especially on the principal veins.

Miquel had only leaf specimens and founded his two species chiefly on the difference in shape of the leaves. On every plant of this species as here understood

the leaves tend toward a rhombic shape, in fact this predominates and from this they vary to broad-ovate and occasionally to ovate. An attempt to make leaf-distinctions in this plant of specific value breaks down on the first bush examined. Maximowicz who had flowering specimens and young leaves added to Miquel's description of *R. dilatatum* the 5 stamens and glandular ovary which subsequent authors have used to distinguish Miquel's two species. That Maximowicz was not justified in this is abundantly proved by the material before me. As to Makino's *R. Wadanum* I have specimens from Shikoku with slightly spotted flowers but can find no other difference and do not consider that they can be separated even as a variety. It is unfortunate that both Miquel's names should have to be dropped but G. Don's description, brief as it is, leaves no doubt that the plant he had was the same as that later described by Miquel. It has long been accepted that Don's name belonged to one of Miquel's species but the question remained to which? With the acceptance of the view first suggested by K. Koch that both represent one species the question disappears and Don's name becomes the valid, as it is the oldest, name. I have taken the form with ten stamens as representing the type since it is by far the more common.

According to G. Don this Azalea was first introduced into England from Japan by Messrs Knight of Chelsea and evidently about 1832-33, but it must have been lost. It was introduced by Maximowicz into the Petrograd Botanic Gardens in 1865, and is figured in the *Gartenflora* for 1868 but the color is poor. It was introduced into the Arnold Arboretum by Professor Sargent who sent seeds from the Nikko region in the autumn of 1892. The plants have grown slowly but have proved perfectly hardy and each spring for a number of years past have flowered profusely. Apparently it is rare in Europe and is one of the plants which will probably thrive better in eastern North America than in England.

The specimen cited from Kintoki on the Hakone mountains is simply an abnormal condition of typical *R. reticulatum* G. Don in which the flowers have been retarded and are open when the leaves are well grown. There are two flowers and the pistil of a third, and a small deformed flower on the specimen, and one flower with two unfolding leaves. The leaves have the characteristic rhombic shape and the flowers have from 8 to 10 stamens. Branches bearing late flowers and leaves together may be found occasionally on any species of precocious flowering Rhododendron.

A variety with white flowers is: —

Rhododendron reticulatum var. **albiflorum** Wilson, n. comb.

Rhododendron rhombicum var. *albiflorum* Makino in *Tokyo Bot. Mag.* XVIII. 66 (1904). — Rehder in *Bailey, Stand. Cycl. Hort.* V. 2943 (1916).

Makino describes this plant from Tosa province in Shikoku and states that it is very rare. It is unknown to me and is not cultivated in western gardens.

The form adopted by Maximowicz as the typical *R. dilatatum* Miquel, and so considered by subsequent authors, and which is characterised by having 5 stamens only may be distinguished as: —

Rhododendron reticulatum f. **pentandrum** Wilson, n. nom.

Rhododendron dilatatum Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 27 (*Rhodod. As. Or.*) (1870), in part, not Miquel. — Franchet & Savatier, *Enum. Pl. Jap.* I. 289 (1875). — Matsumura, *Pl. Nikko* 71 (1894); *Ind. Pl. Jap.* II. pt. 2, 459 (1912). — Boissier in *Bull.*

Herb. Boiss. V. 918 (1897). — Hooker f. in *Bot. Mag.* CXXV. t. 7681 (1899). — J. H. Veitch, *Hortus Veitchii*, 373 (1906). — Shirisawa, *Icon. Ess. For. Jap.* II. t. 61, figs. 23–32 (1908). — Schneider, *Ill. Handb. Laubholz.* II. 497, figs. 327 h, 328 a–b (1911). — Bean, *Trees and Shrubs Brit. Isl.* II. 353 (1914). — Millais, *Rhodod.* 155 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [9] (1918).

Rhododendron dilatatum a. typicum Makino in *Tokyo Bot. Mag.* VII. 134 (1893).

Japan: Hondo, prov. Shinano, 1864, *Tschonoski* (Herb. Gray); prov. Suruga, Mt. Fuji, May 12, 1907, (flowers), August 15, 1905, (leaves); prov. Sagami, Hakone Mts., alt. 600–1000 m. April 16, 1914, *E. H. Wilson* (No. 6448); in a garden, Miyanoshita, April 17, 1914, *E. H. Wilson* (No. 6429).

Cultivated: Royal Gardens, Kew, May, 1880, *G. Nicholson*.

Of the characters proposed by Maximowicz only that of the 5 stamens seems to hold and, as I have already mentioned, I am not convinced that this is constant. This form is believed to have been introduced to England by Messers Veitch in 1883, but that it was in cultivation in Kew as early as 1880 is shown by Nicholson's specimens preserved in this herbarium. I have no record of its being cultivated in America though it will probably appear among the seedlings raised from Japanese seeds collected by me in 1914 and distributed by the Arnold Arboretum.

Rhododendron quinquefolium Bisset & Moore in *Jour. Bot.* XV 292 (1877). — Franchet & Savatier, *Enum. Pl. Jap.* II. 653 (1879). — Matsumura, *Pl. Nikko*, 71 (1894); in *Tokyo Bot. Mag.* XIV. 69 (1900); *Ind. Pl. Jap.* II. pt. 2, 463 (1912). — Komatsu in *Icon. Pl. Koisikav.* I. 59, t. 30 (1912); in *Tokyo Bot. Mag.* XXXII. [8] (1918). — Bean, *Trees and Shrubs Brit. Isl.* II. 374 (1914). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2947 (1916). — Millais, *Rhodod.* 233 (1917).

Japan: Hondo, prov. Shimotsuke, Nikko region, alt. 600–1600 m. May 14, May 18, October 20, 1914, *E. H. Wilson* (No. 6683, type, 6683A, 7676); Lake Chuzenji, October 21, 1914, *E. H. Wilson* (No. 7676A); same locality, August 11, October 25, 1905, *J. G. Jack*.

This very distinct species is well characterised by its corky, brown bark, its broad-elliptic to obovate verticillate leaves and by its snow white, rotate-campanulate corolla. In habit of growth, arrangement and shape of leaves and in the shape of the flowers it superficially resembles *R. pentaphyllum* Maxim. but the two belong to different sections. In Maximowicz's species the flowers and leafy shoots originate from different buds, the flower-bud being terminal and the leaf-buds lateral below it. In this new species both flowers and leafy shoots issue from the same bud. This Azalea is abundant in shady, rocky ravines in the Nikko region but is elsewhere unknown to me. It is a bush or small tree from 1.5 to 8 m. tall with gray-brown, corky bark, fissured into thick irregular plates; the shoots are glabrous, terete, shining brown the first year and arranged partly in verticils and partly alternate. The winter-buds are narrow-ovoid, acute, with

purple-brown ciliate scales, the basal ones having long aristate points. The leaves are deciduous and unfold at the same time as the flowers open; they are clustered 4 or 5 together at the end of the shoots and vary from broad-elliptic to obovate, the apex is mucronate and the base cuneate, the edges are ciliate and often margined with red-purple; they are more or less villose, especially the lower half of the midrib; the petiole is very short and villose. The flowers are either solitary, or two or three together on slender, glandular-hairy pedicels; the corolla is pure white with green spots within, rotate-campanulate, and the calyx is membranous, with 5 lanceolate or deltoid, glandular-ciliolate segments. The stamens are ten, with greenish filaments of unequal length, dilated and villose at the base. The pistil much exceeds the stamens and is glabrous except at the summit of the ovary. The fruit is stout, cylindric, and from 1 to 1.5 cm. long.

According to Bean, this *Azalea* was introduced into English gardens by Lord Redesdale about 1896, but it appears to be rare in gardens. To America it has been sent by the Yokohama Nursery Company under the name of "*Azalea quinquefolia* white" and I have seen it growing fairly well in the garden of Mr. T. A. Havemeyer, on Long Island. It is a woodland plant and needs shade and a cool situation if it is to thrive. In 1905 Mr. J. G. Jack sent seeds from Lake Chuzenji to this Arboretum but the resultant plants failed to prove hardy. The plants raised from seeds sent from the same locality in 1914 have also proved tender. Like many other woodland plants this *Azalea* is difficult to manage when young.

Rhododendron Schlippenbachii Maximowicz in *Bull. Acad. Sci. St. Pétersbourg*, sér. 3, XV. 226 (*Mél. Biol.* VII, 333) (1870); in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 29, t. 2, figs. 7-13 (*Rhodod. As. Or.*) (1870).—Herder in *Act. Hort. Petrop.* I. 347 (1871).—Franchet & Savatier, *Enum. Pl. Jap.* 289 (1875).—Hemsley in *Jour. Linn. Soc.* XXVI. 30 (1889).—Hooker f. in *Bot. Mag.* CXX. t. 7373 (1894).—*Gard. Chron.* ser. 3, XV. 462, fig. 58 (1894); XIX. 561, fig. 87 (1896); LV. 9, t. (1914).—W. Watson in *Garden* XLVI. 80, t. 972 (1894); *Rhodod. and Azaleas*, 112 (1911).—Palibin in *Act. Hort. Petrop.* XVIII. 150 (*Consp. Fl. Kor.* II.) (1900).—J. H. Veitch, *Hortus Veitchii*, 408 (1906).—Komarov in *Act. Hort. Petrop.* XXIV. 206 (*Fl. Mandsh.* III.) (1907).—Schneider, *Ill. Handb. Laubholz.* II. 494, figs. 325 i-l 327 a-b (1909).—Nakai in *Jour. Coll. Sci. Tokyo*, XXXI. 75 (*Fl. Kor.* pt. 2) (1911); *Fl. Syl. Kor.* VIII. 43, t. 15 (1919).—Matsumura, *Ind. Pl. Jap.* II. pt. 2, 464 (1912).—W. Taylor in *Garden*, LXXVII. 136, fig. (1913).—Bean, *Trees and Shrubs Brit. Isl.* II. 378 (1914).—Rehder in Bailey, *Stand. Cycl. Hort.* V. 2943 (1916).—Millais, *Rhodod.* 239 (1917).—Komatsu in *Tokyo Bot. Mag.* XXXII. [9] (1918).

Azalea Schlippenbachii O. Kuntze, *Rev. Gen.* pt. 2, 387 (1891).

Korea: prov. South Keisho, Herschel Island, 1863, *R. Oldham* (No. 510, Herb. Gray); Chirisan, alt. 600-1845 m. November 16,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

This species was discovered by *Baron A. von Schlippenbach* of the Russian navy on the shores of northeast Korea in 1854, and appears to have been cultivated sparingly in Japan for many years under the name of "Kurofune-tsutsuji." *James H. Veitch* in 1893 saw it in a garden in Japan and sent it to England and this was its first appearance in the west. Since then Japanese nurserymen have exported it in small quantities to Europe and America but to this time it is by no means so well known as its beauty warrants. The plants in this Arboretum were raised from seeds collected in Korea in 1905 by *Mr. J. G. Jack*; they have grown slowly but have been unaffected by winter-cold and summer-drought. For two to three years past they have flowered freely and their large, pure pink blossoms are lovely. This species has the second largest leaves and the most beautiful flowers of its group and is worthy of a place in every garden. I collected a quantity of seed in Korea in the autumns of 1917 and 1918, and this has been widely distributed by the Arnold Arboretum and in a few years this *Azalea* should be a prominent feature in New England gardens.

SECT. III. RHODORA G. DON¹

Flowers from a terminal bud, the leafy shoots arising from separate, lower, lateral buds; corolla rotate-campanulate, rarely campanulate or two-lipped; stamens 10, rarely from 5 to 7. Leaves deciduous, scattered or whorled. Fruit more or less cylindrical, rarely conic-ovoid. Shrubs often of large size, usually with rigid ascending or spreading branches; shoots glabrous or hairy.

This section contains 5 species; three of these grow in eastern Asia and two (*R. canadense* Zabel and *R. Vaseyi* A. Gray) in eastern North America.

Rhododendron Albrechtii Maximowicz in *Bull. Acad. Sci. St. Pétersbourg*, sér. 3, XV. 227 (*Mél. Biol.* VII. 335) (1870); in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7. XVI. No. 9, 30, t. 2, figs. 14–20 (*Rhodod. As. Or.*) (1870). — Franchet & Savatier, *Enum. Pl. Jap.* I. 290 (1875). — Matsumura, *Pl. Nikko*, 71 (1894); *Ind. Pl. Jap.* II. pt. 2, 458 (1912). — Boissier in *Bull. Herb. Boiss.* V. 917 (1897). — Schneider, *Ill. Handb. Laubholz.* II. 496, figs. 325 m–n, 327 e (1909). — Millais, *Rhodod.* 112 (1917). — Komatsu in *Tokyo Bot. Mag.* XXXII. [8] (1918).

Azalea Albrechtii O. Kuntze, *Rev. Gen.* pt. 2, 387 (1892). — Rehder in Bailey, *Cycl. Am. Hort.* I. 122 (1900).

Japan: Hondo, prov. Shinano, 1864, *Tschonoski* (Herb. Gray); prov. Musashi, cultivated, *Bot. Gard. Tokyo*, April 2, 1914, *E. H. Wilson* (No. 6363); prov. Shimotsuke, Nikko region, alt. 1600–2000 m., June 23, October 15, 1914, *E. H. Wilson* (Nos. 6858, 7638); prov. Uzen, lower slopes of Adzuma-san, alt. 800–1100 m., July 19, October 24, 1914, *E. H. Wilson* (Nos. 7208, 7721); prov. Mutsu, Hirosaki, May, 1905, *U. Faurie* (No. 6785); Hakkoda-yama, alt. 1000–1600 m., July 5, 1914, *E. H. Wilson* (No. 7083); same locality, October 2, 1892, *C. S. Sargent*. Hokkaido, prov. Oshima, *Fuku-yama*, June 3, 1889, *U. Faurie* (No. 3831, Herb. Kew); Kakumi Hot Springs, September 29, 1892, *C. S. Sargent*; prov. Iburi, Nobori-betsu, July 2, 1914, *E. H. Wilson*; prov. Ishikari, Sapporo, May 17, October 10, 1903, *S. Arimoto* (Herb. Gray); same locality, May 29, 1890, *E. Tokubuchi* (Herb. Gray); *Moiwa-dake*, June 15, 1885, *K. Miyabe* (Herb. Gray); same locality, June 23, 1891, *E. Tokubuchi*; cultivated, *Bot. Gard. Sapporo*, August 22, 1905, *J. G. Jack*.

¹ For synonyms and further details see page 119.

On the margin of forests and in thickets this species is widely distributed in northern Japan from the mountains round Sapporo in central Hokkaido southward to those of Shinano in central Hondo. I am familiar with it from the Nikko region northward, but do not consider it a common plant. It is scattered here and there and is partial to dense thickets on steep slopes of loose soil. It is a shrub of loose and sparse habit, from 1 to 1.5 m. tall, with slender, smooth, purple-brown branches which are brown and furnished when young with curled, partly viscid, gray hairs. The leaves are deciduous, membranous, dark green and change to yellow before they fall; they are scattered on the vigorous shoots, and crowded, forming false whorls of fives, at the end of the branchlets; they are obovate to oblanceolate, from 4 to 12 cm. long and from 1.5 to 6 cm. wide, acute or subacute, with a glandular mucro; the base is narrowed to a short, winged petiole; the margins are serrulate and ciliate, the upper surface is sparsely clad with appressed, flattened, hispid hairs and the under surface is more or less densely clothed with short, pale gray tomentum. The flowers are about 5 cm. across, and open in June and early July, according to climate, and immediately before or as the leaves unfold; they are borne in terminal umbels, each containing from 3 to 5 flowers. The pedicels are moderately stout, from 1 to 2 cm. long, and are clothed with curled, glandular, yellowish hairs; the calyx is small, with 5 purple and ciliate lobes; the corolla is rich red-purple, rotate-campanulate, with a short, wide tube and spreading, rounded lobes; the stamens are 10 in number in two series of unequal length, the longest equalling the corolla and overtopped by the curved style, which has a bifid stigma. The fruit is erect, conic-ovoid, from 1 to 1.2 cm. long, purple-brown, and clothed with yellow, viscid, pilose hairs.

This species was discovered about 1860 by Dr. Michael Albrecht of the Russian Consulate in Hakodate, and was found by Maximowicz also in the vicinity of Hakodate in 1861. However, it does not appear to have been introduced into western gardens until 1892, when Professor Sargent sent seeds to the Arnold Arboretum from two localities in Hokkaido. Plants raised from these seeds grew for several years in this Arboretum, but never really flourished. I sent seeds from northern Hondo in 1914, and these were distributed by the Arboretum. Plants raised from these seeds have grown fairly well, but it is too early to say whether they will make themselves at home here. There is no apparent reason why this *Azalea* should not thrive here. The habit is sparse and the flowers are not large, but the color is intense and the plant is graceful and attractive.

Rhododendron pentaphyllum Maximowicz in *Bull. Acad. Sci. St. Pétersbourg*, sér. 3, XXXI. 65 (Mél. Biol. XII. 491) (1887).—Matsumura, *Ind. Pl. Jap.* II. pt. 2, 463 (1912).—Millais, *Rhodod.* 225 (1917).—Komatsu in *Tokyo Bot. Mag.* XXXII. [8] (1918).

Rhododendron pentaphyllum var. *nikoense* Komatsu in *Icon. Pl. Koisikav.* III. 45, t. 168 (1916).

Rhododendron quinquefolium var. *roseum* Rehder in *Bailey, Stand. Cycl. Hort.* V. 2947 (1916).

Japan: Kyushu, prov. Osumi, Tarumizu village, April and May, 1911, S. Kawagoe. Hondo, prov. Kozuki, Mt. Akagi, June 13, 1911, as to flowers, May 17, 1921, K. Sakurai; prov. Musashi, Yokohama, cultivated, April 11, 1914, E. H. Wilson (No. 6396); prov. Shimot-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Japan: Hondo, prov. Uzen, near Toge, alt. 1000 m., July 20, October 23, 1914, *E. H. Wilson* (Nos. 7219, 7191).

This little known Japanese species is confined to the mountains of north-central Hondo. It is very distinct in its flowers and fruit and also in its papery, cinnamon-brown bark which shreds off and leaves polished brown stems and branches. No other species with deciduous leaves has such attractive bark. In habit it forms an upright, bushy shrub from 1 to 2 m. high, with rigid, red-brown branches clothed with pilose and glandular hairs. The winter-buds are pale straw-color tinged with purple, large, ovoid, with ciliate, eglandular bud-scales. The leaves are scattered, sessile, membranous and deciduous; they have appressed, hispid hairs on both surfaces and the margins are ciliate; in shape they are panduriform, from 6 to 18 cm. long and from 3.5 to 8.5 cm. wide, with the apex rounded or truncate, usually emarginate, rarely obtuse, and are narrowed to the base, which is firmly appressed to the stem and often has a curious shield-like protuberance continued below the point of insertion; the veins are prominent and reticulate on the under side and many are furnished with strigose hairs. In the autumn the leaves are tinted from orange to crimson. The flowers are small and not conspicuous, and open in late June and July with, or even after, the leaves. They are borne from 6 to 15 together in terminal, umbellate corymbs. The corolla is white, tubular or campanulate, from 1.5 to 2 cm. long and from 0.8 to 1 cm. broad, with 5 short, very slightly spreading lobes. The calyx is small, membranous, with lobes of unequal size, and like the slender pedicel is covered with viscid hairs. The stamens are of unequal length, 10 in number and are included, as is the style, which is straight and slightly thickened below the capitate stigma. The fruit is pendent, oval, from 8 to 10 mm. long, and has thin walls which open to the base and have wavy margins.

This species in foliage suggests *R. Schlippenbachii* Maxim., but otherwise is very distinct. On the hills round Toge, near the base of Adzuma-san, in 1914, I found this species to be common in rather open country. The season of flowering was almost past. In the autumn I gathered seeds for the Arnold Arboretum, which distributed them in America and Europe. The plants raised from these seeds have not yet flowered in the Arboretum. This Azalea was unknown to Japanese nurserymen until I gave plants to the Yokohama Nursery Company in 1914. Its bark and foliage is striking, but the size of the flowers and the manner in which they are hidden by the leaves and young shoots detracts from the plant. As a flowering shrub it is one of the least desirable of Japanese Azaleas.

SECT. IV. PENTANTHERA G. DON¹

Flowers from a terminal bud, the leafy shoots from separate, lower, lateral buds; corolla funnel-form-campanulate to funnel-form; stamens 5. Leaves deciduous, scattered on the shoots. Fruit more or less cylindrical. Shrubs usually with stiff, ascending branches. Shoots villose or glabrous.

This section contains some 15 species and is the most widely distributed of the four sections which we are discussing. Two species grow in eastern Asia, one in Japan and the other in China; one species, *R. luteum* Sweet, grows on the Caucasus, in the Pontus region in Asia Minor and in parts of eastern Europe. The others are confined to North America, where they are widely distributed.

Rhododendron molle G. Don, *Gen. Syst.* III. 846 (1834).

Azalea mollis Blume, *Cat. Gewass. Buitenz.* 44 (1823); *Bijd. Fl. Ned. Ind.* 853 (1825).

Azalea sinensis Loddiges, *Bot. Cab.* IX. t. 885 (1824). — De Candolle, *Prodr.* VII. pt. 2, 718 (1839). — Fortune, *Wanderings in China*, 154 (1847); *Tea Countries of China* 154 (1852); *Residence among Chinese*, 28 (1857); Suringar in *Gartenfl.* LVII. 505 (1908).

Rhododendron sinense β . *flavescens* Sweet, *Brit. Flow. Gard.* ser. 1, III. t. 290 (1829).

Rhododendron sinense Sweet, *Brit. Flow. Gard.* ser. 1, III. sub. t. 290 (1829). — Hovey, *Am. Gard. Mag.* II. 142, 179 (1836). — Hance in *Journ. Bot.* XVI. 109 (1878). — Hemsley in *Jour. Linn. Soc.* XXVI. 30 (1889), as to the Chinese plant. — Suringar in *Gartenfl.* LVII. 516 (1908). — W. Watson, *Rhodod. and Azaleas*, 112 (1911), in part. — Schneider, *Ill. Handb. Laubholz.* II. 497, figs. 328 g-h, 329 a-b (1911). — Rehder & Wilson in Sargent, *Pl. Wilson.* I. 549 (1913). — Bean, *Trees and Shrubs Brit. Isl.* II. 379 (1914). — Millais, *Rhodod.* (1917), in part.

Rhododendron sinense α . *flammeum* Sweet, *Brit. Flow. Gard.* ser. 1, III. sub. t. 290 (1829).

Azalea pontica var. *sinensis* Lindley in *Bot. Reg.* XV. t. 1253 (1829), poor figure.

Azalea sinensis α . *genuina* Maximowicz in *Suppl. Ind. Sem. Hort. Petrop.* 1869, 11.

China: prov. Hupeh, stony hills and Pine-woods 15 miles below Ichang on right bank of Yangtze River, alt. 30–300 m., April 24, 1907 (flowers), January 18, 1908 (fruit), *E. H. Wilson* (No. 800); same locality, *A. Henry* (No. 268); prov. Hunan, near Changsha, among shrubs, alt. 70–400 m., April 7, 1918, *Dr. Handel-Mazzetti* (No. 2342); prov. Chekiang, vicinity of Ningpo, 1908, *D. MacGregor*; near Changhua, rocky banks, alt. 300 m., July 12, 1915, *F. N. Meyer* (No. 1543);

¹ For synonyms and further details see page 124.

prov. Kiangsii, Nanking, April 23, 1914, *Mary Strong Clemens* (4226, Herb. Bur. Sci. Manila).

This species is abundant in eastern China, especially on the mountains of the Chekiang province, and appears to have the western limits of its range on the conglomerate hills a little to the east of the city of Ichang where Henry and I collected it. The plant is evidently common in the neighborhood of Ningpo, where every collector since *Fortune* has gathered it. It grows among coarse grasses and shrubs and in thin pine-woods. It is a sturdy, sparingly branched shrub from half a metre to a metre and a half high, taller than broad, and has rich golden-yellow flowers produced in large trusses before the leaves unfold. In habit and floral characters in general it closely resembles its Japanese relative (*R. japonicum* Suring.), but it has a smaller and less bristly calyx, flower-stalks without or with few bristles, stamens as long or longer than the corolla and winter-buds densely clothed with a short, nearly white, velvety pubescence. The leaves of the two species are conspicuously different. Those of the Chinese plant are densely clothed on the lower surface with soft, nearly white, matted pubescence which persists through the life of the leaf; the upper surface is also softly pubescent but much of this disappears before autumn. In late summer and autumn the leaves appear glaucous below on account of the density of the pale gray pubescence. The leaves of the Chinese species are also larger, up to 15 cm. long and 5.5 cm. wide, more decidedly oblong-lanceolate, and only occasionally broadest above the middle. Sweet's figure (*Brit. Flow. Gard.* ser. 1, III. t. 290 [1892]) clearly shows these essential characters, and if it is compared with Regel's figure (*Gartenfl.* XVI. 299, t. 556 [1867]) of the Japanese species the two plants cannot be confused. Since the two may easily be distinguished by the leaves and other characters, behave differently under cultivation and grow wild in widely separated geographical areas, I consider that they are best kept under separate names without discussing the vexed question of what constitutes a species.

I have not seen this *Azalea* in Chinese gardens but it must long ago have been grown in those of Canton, Hanchow and Soochow. Loddiges received the plant from China in 1823 and figured it in the *Botanical Cabinet*, IX. t. 885 (1824). This plant in all probability was brought by some ship of the East India Company from Canton. Others were introduced through the same agency during the succeeding years but the species appears to have been lost, or nearly so, when it was sent to the gardens of the Horticultural Society of London by Robert Fortune in 1845. Fortune found it in gardens at Canton and wild on the hills near Ningpo, and in his books he mentions this *Azalea* several times. According to Blume (*Cat. Gewass. Buitenz.* 44 [1823]) it was introduced into gardens in Java before 1823. Of its first introduction into America I can find no record, but Hovey tells us that it flowered with him, and also in the garden of Mr. J. P. Cushing, Watertown, Mass., in April, 1836. Later it became lost and it was not cultivated in America when in January, 1908, I sent seeds from China to the Arnold Arboretum. Some of these seeds were sent to England, where the species appears again to have been lost. Indeed, in western gardens the pure species never seems to have established itself, and like its Japanese relative soon got lost under a mass of hybrid and seminal forms. In this Arboretum the Chinese *Azalea* is doubtfully hardy, but it is an excellent pot plant and forces well. This tenderness, which is to be expected when the climate of the region in which it grows is taken into consideration, explains its frequent disappearance from gardens. The habit and fine yellow flowers make it a very decorative garden plant but it lacks stamina. Its great use has been in hybridising, notably with the Japanese species, from which has sprung a race of hybrids popularly known as "Mollis Azaleas," correctly \times *R. Kosterianum*.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Dendr. Ges. XI. 30 (1902). — Schneider, *Ill. Handb. Laubholz.* II. 499 (1911).

Rhododendron occidentale × *sinense* Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz-Ben.* 380 (1903), as a synonym.

Rhododendron molle × *occidentale* Schneider, *Ill. Handb. Laubholz.* II. 499 (1911), as a synonym.

Azalea occidentalis "Hybrids" Hort. Wezelenburg, *Cat.* 9 (1915).

Rhododendron occidentale × *R. molle* × "Ghent Azaleas" Millais, *Rhodod.* 220, fig. (1917).

Many varieties of this hybrid have been raised by Anthony Waterer and by Koster & Sons, among the best being "Superba," "Graciosa," "Exquisite" and "Magnifica." The parentage is, apparently, very mixed; a majority have large white flowers with the usual yellow blotch characteristic of the flowers of the American species, but some have cream and others yellow flowers with a darker blotch. They flower between early June and mid-July and are very hardy.

In quite recent years × *R. Kosterianum* has been hybridised with a variety of forms of × *R. gandavense* and the result is a race with double flowers, for which I propose the name of:—

× ***Rhododendron mixtum*** Wilson, n. nom.

Azalea rustica flore-pleno Hort. ex Vandervoorde in *Rev. Hort. Belg.* XIX. 232, t. (1893), not Van Houtte. — *Cat. A. Waterer*, 10 (1911-12.) — *Cat. Wezelenburg*, 10 (1915). — Millais, *Rhodod.* 213 (1917).

It appears to be safer to give a new name to this race than to risk confusing it with the double-flowered "Gandavensis hybrids" (Azaleas rustique de Gand) first raised by Louis van Houtte, who figures four varieties in *Flore des Serres* XIX. tt. 2021-2024 (1873), and of which "Graf von Meran," "Narcissiflora" and "Louis A. van Houtte," may be considered types. Typical forms of × *R. mixtum* are "Murillo" with rose-colored flowers, "Virgile" with yellow flowers, "Il Tasso" with red flowers, "Milton" with white flowers, and "Racine" with pale pink flowers. According to Millais (*Rhodod.* 213 [1917]) there are hybrids raised by Vuylsteke between *R. molle* and *Azalea altaclarensis*. These are unknown to me.

Rhododendron molle G. Don has been crossed with one or more of the broad-leafed Rhododendrons and the oldest of these is:—

× ***Rhododendron "Smithii aureum"*** Paxton, *Mag. of Bot.* IX. 79, t. and fig. (1842). — *Gard. Chron.* II. 352 (1842). — Robinson, *Fl. and Sylva* II. 152, t. (1904). — Bean, *Trees and Shrubs Brit. Isl.* II. 346 (1914).

Rhododendron "Hardy Seedling" × *Azalea sinensis* Paxton, *Mag. of Bot.* IX. 79 (1842.)

"Norbiton hybrids" Hort. ex Masters in *Gard. Chron.* ser. 3, XIII. 665 (1893).

Rhododendron sinense × *Eurhododendron spec.* Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz.-Ben.* 379 (1903).

I have given the parentage as set down by Paxton but Robinson says "between a yellow form of the Chinese Azalea and *Rhododendron caucasicum*, the latter of

which it resembles in habit." This interesting *Azaleodendron* was raised by a nurseryman at Norbiton, Surrey, named W. Smith, and exhibited before the Horticultural Society of London in the spring of 1842. It is not now in cultivation in this Arboretum, not being sufficiently hardy to withstand the winter's cold. A specimen in this herbarium under the name of *Rhododendron* "Victoria regina" and from an exhibition held by the Massachusetts Horticultural Society on April 19, 1889, evidently belongs here, but whether the name it was exhibited under has any standing I do not know.

Another hybrid between a broad-leaf evergreen *Rhododendron* and *R. molle* G. Don is:—

× *Rhododendron* "broughtonii aureum" Hort. apud W. Watson, *Rhodod. and Azaleas* 93 (1911). — Osborne in *Gard. Chron.* ser. 3, LI. 53. fig. 29 (1912). — Bean, *Trees and Shrubs Brit. Isl.* II. 345, fig. (1914).

Osborne in his account of this plant says it "obtains its name from the village of Broughton in Peebleshire, where it was raised by crossing a yellow *Azalea* and a seedling evergreen *Rhododendron*." Both Osborne and Bean regard this as the best of its class. It has been in cultivation in this Arboretum since 1910 but does not thrive. More recently M. G. Vander Meulen has raised hybrids between *R. molle* and the evergreen *Rhododendrons* "Prince Camille de Rohan" and "Leopard" which he names "Édouard André," "Director Rodigas" and "Dr. Masters." Mr. White of Sunningdale has raised similar hybrids. Millais (*Rhodod.* 214 [1917]) says that they retain their leaves in winter, and are quite hardy and very floriferous.

Rhododendron japonicum Suringar in *Gartenfl.* LVII. 516 (1908). — Schneider, *Ill. Handb. Laubholz.* II. 1046 (1912). — Rehder & Wilson in Sargent, *Pl. Wilson.* I. 549 (1913), in a note. — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2942 (1916). — Arnold *Arb. Bull. Popular Inform.* n. ser. III. 26 (1917); IV. 20, 28 (1918). — Komatsu in *Tokyo Bot. Mag.* XXXII. [6] (1918). — E. G. Loder in *Rhodod. Soc. Notes*, I. No. 4, 197 (1918).

Rhododendron molle Siebold and Zuccarini in *Abh. Akad. Münch.* IV. pt. 3, 131 (*Fl. Jap. Fam. Nat.* II. 7) (1846), not G. Don. — Miquel in *Ann. Mus. Lugd.-Bat.* I. 33 (1864); II. 164 (1865-66); *Prol. Fl. Jap.* 96 (1866-67). — Schneider, *Ill. Handb. Laubholz.* III. 499, figs. 328 i-k, 329 c-d (1911). — Bean, *Trees and Shrubs Brit. Isl.* II. 368 (1914). — Millais, *Rhodod.* 211 (1917), in part.

Azalea japonica A. Gray in *Mem. Am. Acad.* n. ser. VI. 400 (1858-59). — Hemsley in *Garden*, XI. 420, t. (1877), in part.

Rhododendron molle var. *glabrior* Miquel in *Ann. Mus. Lugd.-Bat.* I. 33 (1864).

Azalea mollis β. *glabrior* Regel in *Gartenfl.* XVI. 289, t. 556 (1867). — Van Houtte in *Fl. des Serr.* XIX. 177 tt. 2034-2035 (1873).

Azalea sinensis β. *glabrior* Maximowicz in *Suppl. Ind. Sem. Hort. Petrop.* (1869) 11.

Rhododendron Sinense Maximowicz in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 28 (*Rhodod. As. Or.*) (1870), not Sweet. — Hooker f. in

Bot. Mag. XCVII. t. 5905 (1871). — Franchet & Savatier, *Enum. Pl. Jap.* I. 289 (1875). — Matsumura & Yatabe, *Nippon Shokub.* 163 (1884). — Matsumura, *Pl. Nikko*, 71 (1894); *Shokub.* 250 (1895); *Ind. Pl. Jap.* II. pt. 2, 464 (1912). — Shirasawa, *Icon. Ess. For. Jap.* II. t. 62, figs. 19–27 (1908). — Hayata, *Veget. Mt. Fuji*, 66 (1911). — Ito, *Icon. Pl. Jap.* I. No. 5, 1, t. 17 (1913). — W. Watson, *Rhodod. and Azaleas*, 112 (1911), in part.

Azalea mollis André in *Ill. Hort.* XVIII. 132, t. 68 (1871), not Blume. — Suringar in *Gartenfl.* LVII. 505 (1908).

Japan: Hondo, prov. Suruga, Mt. Fuji, alt. 600–1000 m., open country, May 8, 1914, *E. H. Wilson* (No. 6657); same locality, July 28, 1891, *K. Watanabe* (Herb. Gray); prov. Shinano, Karuizawa, alt. 1000 m. May, 1903, *B. Hayata*; prov. Shimotsuke, between Lake Chuzenji and Yumoto, open country, alt. 1600–1800 m., June 24, 1914, *E. H. Wilson* (No. 6872); same locality, September 5, 1892, *C. S. Sargent*; same locality, August 11, 1905, *J. G. Jack*; Nikko region, alt. 500–1800 m., May 22, flowers, October 19, fruit, 1914, *E. H. Wilson* (Nos. 6727, 7670); prov. Uzen, foot of Adzuma-san, grassy slopes, alt. 1000 m., October 23, 1914, *E. H. Wilson* (No. 7192); without locality, June 7, 1913, *K. Sakurai*; prov. Rikuchu, round base of Hayachine-san, June 4, 1905, *U. Faurie* (No. 6784). Hokkaido, prov. Oshima, Hakodate, cultivated, 1855, *C. Wright* (Herb. Gray).

Cultivated: Arnold Arboretum (Nos. 861–1, 861–4, 861–5, 861–6); Hort. Kew, May 18 (flowers), August 7 (leaves), 1880, *G. Nicholson* (No. 807, Herb. Arnold Arboretum).

This handsome species is common over a great part of Hondo, the main island of Japan. I am familiar with it from the neighborhood of Kamo, on the Kwan-sai railroad beyond Nara, northward to the foot-hills of Hayachine-san. It is a special feature of the moorlands round the base of sacred Fuji-san and of Ontake-san, and of open grass and scrub-clad places in the Nikko region and elsewhere. Matsumura reports it from Higo province in Kyushu but I have not seen it wild anywhere in this southern island. It is cultivated in and around Hakodate in Hokkaido but I can find no record of its growing wild in that island. Wherever I have seen this *Azalea* wild it was scattered in open grass and scrub-clad country, never in woods or dense thickets and obviously it is sun-loving. It is a sturdy-growing plant, with fairly stout, erect shoots, as much as two metres high and a metre through; usually, however, it is less than a metre high and broad. The flowering season is from about the last week in April to the end of June, according to locality. The flowers appear before the leaves, and the corolla varies in color from orange-red to flame-red or almost red; they are broad funnel-shape, from 5 to 6 cm. across and the flowers are produced 6 to 12 together in umbellate corymbs at the end of shoots. The calyx is small, obtusely 5-lobed with many (rarely few) straight, bristle-like gray hairs; the flower-stalk and annual shoot are also clothed with similar bristles, though occasionally these are wanting on the foot-stalks. The outside of the corolla is velutinous. The stamens, five in number and of unequal length, are shorter than the corolla, with pilose filaments and dark brown anthers. The ovary is villose, and the fruit is erect, oblong-ovoid to cylindrical, from 2 to 2.5

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

the nurseries round Tokyo and Osaka for export it is but little used by the Japanese in their own gardens, probably because it is not amenable to clipping.

In western gardens the first to raise and name varieties of *R. japonicum* from seed was Louis van Houtte of Ghent, Belgium, who published (in the *Fl. des Serr.* XIX. 177 [1873]) an article, accompanied by four plates. Enthusiastic on the future possibilities of this plant, he named and described twenty kinds. Among these are some of the best seminal kinds of this Azalea yet produced. The first of them to attain popularity was "Comte de Gomer," which is figured in the *Revue Horticulture Belge*, I. 97, t. (1875), by Kerchove de Denterghen under the name of *Azalea mollis* "Comte de Gomer." In 1879, Messers James Veitch & Sons exhibited it before the Royal Horticultural Society under the same name and it received a First Class certificate. It is figured by Richard Dean in the *Floral Magazine*, XVIII. t. 367 (1879), and is still among the best kinds raised. Others of the same stock are "Alphonse Lavallée," "Baron Edmond de Rothschild," "Comte Papadopoli," "Consul Pêcher" and "Charles Kékulé." In his article Van Houtte makes no mention of using the Chinese plant but his figure 2033 suggests that plant. His t. 2035, depicting white flowers with yellow blotch on upper petal is presumably the variety "Chevalier A. de Réali," and suggests an albino form of the Chinese plant rather than of the Japanese species. I have seen no specimen of this, but the leaves should tell its origin. In this connection, however, it is well to remember that although only recently definitely known, the yellow-flowered form (f. *aureum*) of the Japanese species might have been raised from the seedlings distributed by Groenewegen or from the Petrograd stock. I saw the f. *aureum* in the British Consulate garden in Hakodate in 1914 and the plant might well have been growing somewhere in the town when Maximowicz was there in 1861. In reading Van Houtte's account the inference certainly is that all the forms he mentions and figures are of the Japanese plant, and Van Houtte, of course, knew the Chinese plant well. At this late date it is not possible to be sure, but I incline to the belief that the yellow-flowered form of the Japanese species appeared among the seedlings raised from Groenewegen's or Maximowicz's seeds.

Rhododendron japonicum f. aureum Wilson, n. f.

Rhododendron sinense Tanaka, *Useful Pl. Japan*, 74, No. 652, fig. (1891), not Sweet.

Rhododendron sinense var. *flavescens* Ito, *Icon. Pl. Jap.* I. No. 5, p. 4 (1913), not Sweet.

Japan: Hondo, prov. Musashi, brought from Chichibu mountains, cultivated, Botanic Gardens, Tokyo, May 11, 1917, *E. H. Wilson* (No. 8414, type); prov. Kawachi, between Yomomoto and Ikedo, cultivated, said to have been brought from round Tokyo, May 8, 1918, *E. H. Wilson* (No. 10,351).

Except for its rich, deep yellow flowers there is nothing to distinguish this form from the typical *R. japonicum*. I was informed by Dr. Nakai that this form grows wild mixed with the type on Chichibu mountains in the northwestern part of Musashi province and that the plant my specimens are from came from there. I did not see it wild but Dr. Nakai's statement was confirmed to me by various horticulturists. I saw it in gardens in Hakodate in 1914, and in nurseries not far from Osaka in 1918. The winter-buds are pale colored and afford an easy means of distinguishing it from the type. The Japanese name for this plant is "Ki-tsutsuji" or "Ki-rengé-tsutsuji" and Tanaka's figure cited above is quite good. As far as I

can discover it was first introduced to America in the spring of 1917 among plants of the type imported from the Yokohama Nursery Company by Mr. J. S. Ames, North Easton, Mass. In 1919 I brought back plants from Japan to the Arnold Arboretum. Whether this plant has reached Europe I do not know, but there is no mention of it in Millais's recent work on Rhododendrons. In this connection it should be mentioned that Maximowicz (in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. No. 9, 29 (*Rhodod. As. Or.*) [1870]) speaks of the form with yellow flowers as rare, but it must be remembered that he confuses the Japanese and Chinese species and calls both *R. sinense*. Maximowicz identifies this plant with the "Rjuku Tsutsusi, lillifera Insulae Luconiae, flore languidé luteo, petalis, surrectis, punctis saturatoribus variegatis" and with "Rjuku Tsutsusi flore igneo, punctis petalorum croceis, apicibus rufis" (Kaempfer, *Amoen. Exot.* 848 [1712]). What the first named of Kaempfer's plant is I do not know, neither can I find out the "Insulae Luconiae;" the description, however, would fit our new form (*aureum*) or the Chinese species. The "Rjuku Tsutsusi fl. igneo" is *R. scabrum* G. Don (*R. sublanceolatum* Miquel). Personally I doubt if Kaempfer saw or heard of *R. japonicum* or its form *aureum*, since they are not favorite garden plants in Japan, neither do they grow wild by the road he traversed from Nagasaki to Tokyo.

A recently raised hybrid between *R. japonicum* Suring. and *R. canadense* Zabel is: —

× *Rhododendron Fraseri* W. Watson in *Gard. Chron.* ser. 3, LXVII. 225 (1920).¹

Rhododendron luteum Sweet, *Hort. Brit.* ed. 2, 343 (1830), not *Azalea lutea* L. — Schneider, *Ill. Handb. Laubholz.* II. 1046 (1912), not II. 500 (1911). — Rehder in Bailey, *Stand. Cycl. Hort.* V. 2942 (1916).

Azalea pontica Linnaeus, *Spec.* 150 (1753), not *R. ponticum* Linnaeus, — Pallas, *Fl. Ross.* II. 51, t. 19 (1788). — Andrews, *Bot. Rep.* I. t. 16 (1797). — Schmidt, *Öster. Baumz.* III. 44, t. 169 (1800). — Marschall & Bieberstein, *Fl. Taur.-Caucas.* I. 144 (1808); III. 136 (1819). — Hovey, *Mag. Hort.* IV. 276, 367 (1838). — De Candolle, *Prodr.* VII. pt. 2, 718 (1839).

? *Azalea flava* Hoffmannsegg, *Verz. Pflanzenkult.* 61 (1826).

Anthodendron ponticum Reichenbach in Mösseler, *Handb. Gewächsk.* II. Aufl. 1. 309 (1827); *Fl. Germ. Excur.* I. 416 (1830).

Rhododendron flavum G. Don, *Gen. Syst.* III. 847 (1834), not Pallas. — Nyman, *Consp. Fl. Europ.* 491 (1878–82). — Schneider, *Ill. Handb. Laubholz.* II. 499, figs. 328 n–q, 329 e–h (1911). — Bean, *Trees and Shrubs Brit. Isl.* II. 356 (1914). — Millais, *Rhodod.* 165 t. (1917). — Boissier, *Fl. Or.* III. 972 (1875). — Kuzentzof, *Fl. Caucas. Crit.* IV. pt. 1, 31, 488 (1901, 1906).

Rhododendron flavum var. *coronarium* Sweet, *Brit. Flow. Gard.* ser. 2, IV. t. 331 (1836).

Azalea arborea Linnaeus apud De Candolle, *Prodr.* VII. pt. 2, 718 (1839), as a synonym.

Rhododendron ponticum Schreber apud De Candolle, *Prodr.* VII. pt. 2, 718 (1839), not Linnaeus, as a synonym.

Azalea Pontica a. flava De Candolle, *Prodr.* VII. pt. 2, 718 (1839).

Azalea pontica var. *autumnalis* K. Koch in *Linnaea* XVII. 281 (1843).

Anthodendron flavum Reichenbach apud K. Koch, *Dendr.* II. pt. 2, 184 (1872), as a synonym. — Schneider, *Ill. Handb. Laubholz.* II. 499 (1911).

Azalea pontica "Goldlack" Hesse in *Gartenfl.* LII. 48 (1903).

¹ See page 185.

Caucasus, near summit of Dariel Pass, south side 2000 m. alt., July 24, 1903, *C. S. Sargent*; Vladikavkaz, April 22, 1902, *A. Fomine*; Mt. Beschtou, June, July, *R. F. Hohenacker*; prov. Ossetia, Alagir, swamps, May 10, Sept. 12, 1901, *R. Marcowicz*. Pontus region, "Ciganadagh," July 20, 1889, *P. Sintenis* (No. 1357); Sumila, in woods, July 31, 1889, *P. Sintenis* (No. 2017).

Cultivated: Arnold Arboretum (No. 3159-1) from seeds collected on the Caucasus by *C. Schneider*, June, August 28, 1917, *A. Rehder*; Hort. Holm Lea, May and September, 1880; Bot. Gard. Cambridge, Mass., May 31, 1916, *A. Rehder*; Hort. Kew, May 19, August 9, 1880, *G. Nicholson* (No. 835); Breslau, Schneitniger Park, June 13, July 23, 1902, *C. Baenitz*; Hort. Bot. Göttingen, May 22, 1893, *A. Rehder* (No. 1574).

This well-known species is geographically isolated from its relatives and has a rather remarkable distribution. It is native of the Caucasus Mountains, the Pontus region of the Black Sea, of Cilicia, and also of eastern Europe (Lithuania, southeastern Poland, Volhynia and in southwestern Russia). It is a broad, densely branched shrub from 2.5 to 4 m. high and as much as 6 m. in diameter. The branches are twiggy and clothed when young with glandular, villose hairs. The leaves are scattered, oblong-lanceolate often broadest above the middle, from 4 to 12 cm. long and from 1.5 to 3.5 cm. long, acute or obtuse, mucronate, narrowed to the base, the margin is serrulate and ciliate, and on both surfaces are scattered, appressed, short glandular bristles and short curled pale gray pubescence which later disappears. The petioles, calyx and outside of the corolla tube are glandular. The richly fragrant flowers are borne from 10 to 12 together in clusters at the end of the naked shoots. The calyx is green, deeply divided into 5 oblong or ovate, glandular-ciliate lobes from 2 to 5 mm. long. The corolla is yellow, from 4.5 cm. in diameter with a narrow, cylindric tube from 1.5 to 2 cm. long. The stamens are long exserted but overtopped by the pistil which has a glandular ovary, slender style and capitate stigma. The fruit is cylindric, from 2 to 2.5 cm. long and from 0.5 to 0.8 cm. wide, furrowed and glabrescent.

This Azalea was discovered by Tournefort on the eastern side of the Black Sea during his voyage to the Levant (1700-02) and is described in his *Corollarium Institutionum rei herbariae*, p. 42 (1703), as "*Chamaerhododendros Pontica, maxima, Mespili folio, flore luteo*"; he also figures it in *Mémoires de l'Académie royale des sciences*, 348 (1704). A long account of it is given in his *Voyage du Levant*, II. 99 (1718). Buxbaum in his *Plantarum minus cognitarum centuria V.* 36, fig. 69 (1740) gives a crude figure of it under Tournefort's name. It was on this description and figure that Linnaeus based his *Azalea pontica*. The plant was introduced into England from the Caucasus by Pallas, who sent seeds to Messers Lee & Kennedy of Hammersmith in 1792, with whom they germinated the following year. In 1796, Mr. Anthony Hove of Warsaw found this Azalea on the north side of the Black Sea near Oczakow in marshy places, on the banks of the rivers Dnieper and Dniester, and on the Asia Minor side of the Black Sea round Trebizond. Mr. Hove sent seeds to Watson, a nurseryman of Islington, and plants flowered in a greenhouse in 1798. According to Schmidt it was found in Volhynia by Dr. Besser, and was introduced to the Botanic Garden at Vienna before 1800. I have not been able to ascertain exactly when it was introduced into this country, but according to Hovey

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

THE AZALEAS OF NORTH AMERICA

BY

ALFRED REHDER

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

first American species recorded is *Rhododendron viscosum*, of which we find a figure published in 1691 by Plukenet (*Phytogr.* I. t. 161, fig. 4) under the name *Cistus virginiana flore et odore Periclymeni* after a drawing made from nature by John Banister, an English missionary in Virginia who took great interest in botany and sent in 1680 a catalogue of Virginian plants to Ray. A few years later Plukenet (*Mant.* 49) mentioned the plant now known as *R. nudiflorum* under the name *Cistus virginiana Periclymeni flore ampliore minus odorato*. Both species were introduced into England by Peter Collinson before 1730 and of *R. nudiflorum* a good colored plate was published by Trew in 1750 (*Pl. Select. Ehret.* t. 48), while the first colored plate of *R. viscosum* was published by Meerburgh in 1798 (*Pl. Select. Icon.* t. 9). Linnaeus when publishing his *Species plantarum* in 1753 knew only these two species which he referred to the genus *Azalea* and named *A. viscosa* and *A. lutea*, changing the latter name to *A. nudiflora* in the second edition of his work. In the same edition, in 1762, he described *Rhodora canadensis* introduced some time before the year 1756 from Canada to Paris and figured first by Duhamel in 1760 (*Sem. Pl. Add.* 10, t. 27, fig. 2). In 1763 Adanson (*Fam. Pl.* 164) proposed the name *Tsutsusi* based on Kaempfer's t. 846 which is *R. obtusum* var. *Kaempferi*, and cites "Azalea L." as a synonym; though not an American plant, this is mentioned here, because Small in the *North American Flora* considers *A. indica* L., under which Linnaeus cites Kaempfer's tab. 846, the type of the genus *Azalea*. In 1766 Crantz (*Inst.* II. 468) proposed the new name *Hochenwartia* for *Rhodora*, because it was not the *Rhodora* of Pliny. Gmelin (*Syst. Nat.* II. pt. 1, 694) in 1791 united *Rhodora* with *Rhododendron*, and Salisbury in 1796 (*Prodr.* 286) first recognized the close relationship of most Azaleas to the genus *Rhododendron*; he referred the American Azaleas and *Rhodora* to the genus *Rhododendron*, retaining the name *Azalea* for *A. procumbens*, one of the two species upon which Linnaeus founded the genus. In 1803 Michaux (*Fl. Bor.-Am.* I. 150) described two new species, *A. calendulacea* and *A. canescens*; the former had been mentioned as long before as 1749 by Colden but without sufficient description. It was mentioned again in 1791 by W. Bartram, who discovered it in 1776 and describes its beauty in glowing terms. The second species is possibly identical with one of Aiton's varieties of *A. nudiflora*

and its introduction would be in that case prior to 1789. In 1811 Willdenow (*Berl. Baumz.* ed. 2, 49) described from cultivated plants *A. speciosa* identical with one of Aiton's varieties of *A. nudiflora* and the following year Loiseleur Deslongchamps (*Nouv. Duhamel.* V. 224) described *A. rosea* also from cultivated plants. Pursh in 1814 describes as a new species *A. arborescens* and raises three of Aiton's varieties of *A. viscosa* and one of *A. nudiflora* to specific rank. In 1822 D. Don (in *Edinb. Philos. Jour.* VI. 47) transferred, like Salisbury twenty-six years before, the species of Azalea except *A. procumbens* to *Rhododendron*. In this he was followed by Torrey, G. Don, Endlicher, Maximowicz, Bentham & Hooker and most recent authors, except that Maximowicz, Bentham & Hooker and later authors took up Desvaux's name *Loiseleuria* for *Azalea procumbens*. In 1827 Reichenbach, who like the two Dons and Endlicher retained the name *Azalea* for *A. procumbens*, proposed the genus *Anthodendron* (in Moessler, *Handb. Gewächsk.* I. 244) for *A. pontica*, *A. nudiflora* and *A. viscosa*. G. Don (*Gen. Syst.* III. 84) in 1834 was the first botanist to subdivide the genus *Rhododendron* into sections and he proposed the sections *Pentanthera* and *Rhodora* with which we are here concerned. After 1814 no new species were added until the year 1855, when *A. californica* Torr. & Gray (*Rh. occidentale* Gray) from California was described; it was introduced by Lobb about 1850. The very distinct *R. Vaseyi* was discovered in 1878 by George Vasey and described the following year by A. Gray; it was introduced into cultivation about the same time. In 1891 Greene (in *Pittonia*, II. 172) described *R. sonomense* as a new species, but this is only a form of *R. occidentale*. In 1892 Otto Kuntze, who took the year 1737 as the starting point for generic nomenclature and consequently considered *Azalea* the older generic name, transferred all the species of *Rhododendron* known to him to *Azalea*. In 1901 Small (in *Bull. Torr. Bot. Club*, XXVIII. 360) described as a new species *A. candida* now referred to *R. canescens* as a variety, and in 1903 (*Fl. S. E. U. S.* 883) he described *A. oblongifolia* and *A. serrulata* and separated *R. Vaseyi* as a distinct genus under the name *Biltia*. In 1913 (*l. c.* ed. 2, 1356) he adds two new species *A. austrina* and *A. prunifolia*, and in 1914 (in *N. Am. Fl.* XXIX. 42) *A. prinophylla* identical with *A. rosea* Loisel. (*R. roseum* Rehd.). In 1917 Ashe

(in *Bull. Charleston Mus.* XIII. 26) published *A. atlantica* as a new species and three years later *A. neglecta* (in *Bull. Torr. Bot. Club*, XLVII. 581 [1920]) here referred to *R. atlanticum* as a form. One new species and several new varieties are described in this paper. Of these recent species *R. austrinum* was introduced into cultivation in 1914, *R. oblongifolium* in 1917 and *R. serrulatum* and *R. prunifolium* in 1918, all through the agency of the Arnold Arboretum. *Rhododendron atlanticum* has been cultivated for several years by W. W. Ashe in Washington and by Professor Coker in the Arboretum of the University of North Carolina at Chapel Hill. *Rhododendron alabamense* is the only species not yet in cultivation.

Soon after the introduction of the first American Azaleas to England about 1730 they become favorite garden plants and at the beginning of the last century there were already several species and many varieties in cultivation which played an important part among the so-called American plants. By hybridization between themselves and with the *Rhododendron luteum* introduced at the end of the 18th century and also with evergreen Rhododendrons the garden forms began rapidly to increase in numbers and in 1836 Loddiges cultivated in his nursery near London, according to his catalogue, 107 varieties and hybrids of *R. viscosum* and 43 varieties of *R. nudiflorum* and several other species and varieties. The delicate coloring of some and the brilliancy of others, combined with a delicate fragrance in several species and the hardiness of almost all these Azaleas, made them very desirable and favorite garden plants both in England and on the Continent. Hybridization in the seventies with *R. japonicum* added a new race of hybrids which are perhaps even more showy in some respects but weaker in constitution than the older hybrids generally known as Ghent Azaleas. It is, however, doubtful if any of these hybrids surpass in the brilliancy of their flowers the true species like *R. calendulaceum* and *R. speciosum* which had almost disappeared from European gardens until recently reintroduced.

The American Azaleas are chiefly confined to the Appalachian Mountains and the adjoining regions and to the Atlantic and Gulf coast region; only one species, *Rhododendron occidentale*, is found in the Pacific states from southern Oregon to southern California. No Azalea occurs south of the United States bound-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

The present study of American Azaleas is based chiefly on the collections of the Arnold Arboretum. For many years Professor Sargent has taken special interest in Azaleas and has succeeded in assembling a large amount of material, particularly from the southern states usually poorly represented in most herbaria. Besides having this material at my disposal, I have consulted the collections of the Gray Herbarium at Cambridge and of the National Herbarium at Washington and the herbaria of the New York Botanic Garden, the Academy of Natural Sciences of Philadelphia, and of the New England Botanical Club; specimens for further study were kindly loaned to me from most of these institutions and from the herbaria of the State University of Ohio, of the Missouri Botanical Garden and of the Academy of Sciences at Rochester. I also am under obligation for specimens to Professor W. C. Coker of the University of North Carolina, to Dr. K. M. Wiegand of Cornell, Mr. W. W. Ashe of Washington and Mr. John Dunbar of Rochester. I have further had the opportunity to see most of the species in their native habitats during various journeys, and the living collections of Azaleas of this Arboretum supplied me with material to study many of the species and a large number of garden forms and hybrids in the living state. The great variability of most species together with the absence of strong morphological characters within this group makes it difficult to assemble the numerous forms into well defined species and varieties. The difficulty is considerably increased if one tries to elucidate the innumerable garden forms and hybrids, as few of these forms are adequately described or figured and herbarium or living material of only a small percentage of them is available.

The Azaleas form a distinct subgenus of the *Rhododendron* for which we propose to use Endlicher's subgeneric name *Anthodendron*, as we do not believe that the name *Azalea* used for it by most authors can be correctly applied to this subgenus, as will be seen from the following enumeration of synonyms and the appended remarks.

THE SUBGENUS ANTHODENDRON ENDL.

Rhododendron subgen. **Anthodendron** Endlicher, *Gen.* 759 (1839), emended.

Azalea Linnaeus, *Spec.* 150 (1753), as to species 1-4. — Desvaux in *Jour. Bot. Appl.* I. 35 (1813). — Roemer & Schultes, *Syst.* IV. 374 (1819), excluding species 12 and 13. — Gray, *Man.* 268 (1848). — K. Koch, *Dendr.* II. 1, 171 (1872). — Rehder in *Bailey, Cycl. Am. Hort.* I. 119 (1900), excluding *A. albiflora*, *semibarbata*, *kamschatica*. — Britton & Brown, *Ill. Fl.* II. 558 (1897). — Britton, *Man.* 698 (1901). — Small in *N. Am. Fl.* XXIX. 41 (1914).

Anthodendron Reichenbach in Moessler, *Handb. Gewächsk.* I. 244, 308 (1827); *Fl. Germ. Exc.* 416 (1831).

Rhododendron sect. *Pentanthera* G. Don, *Gen. Syst.* III. 846 (1834). — See also page 124.

Rhododendron a. *Anthodendron*, Endlicher, *Gen.* 759 (1839).

Rhododendron subgen. *Azalea* Planchon in *Rev. Hort.* 1854, 43; with *Tsusia* as distinct subgenus. — Maximowicz in *Bull. Acad. Sci. St. Pétersbourg*, XV. 229 (1870); in *Mém. Acad. Sci. St. Pétersbourg*, sér. 7, XVI. 9, 24 (1870), as section, with *Tsusia* as distinct section. — Bentham & Hooker, *Gen.* II. 601 (1876), as series, with *Tsusia* as distinct group. — Gray, *Syn. Fl. N. Am.* II. 1, 40 (1878). — Dippel, *Handb. Laubholz.* I. 410 (1889), with *Tsusia* as distinct subgenus. — Drude in Engler & Prantl, *Nat. Pflanzenfam.* IV. 1, 37 (1889). — Koehne, *Deutsch. Dendr.* 454 (1893), excluding *Azaleastrum* and *Therorhodion*. — Rehder & Wilson in Sargent, *Pl. Wilson.* I. 516 (1913), excluding *Chionastrum* and *Azaleastrum*. — Rehder in *Bailey, Stand. Cycl. Hort.* V. 2937 (1916), excluding *Azaleastrum*.

Shrubs or occasionally small trees with deciduous or half evergreen rarely evergreen leaves, usually more or less strigose- or villose-pubescent, sometimes glabrous, never lepidote-glandular. Flowers from a terminal bud, one or several to many in umbel-like racemes. Corolla funnel-form to rotate, rarely campanulate, usually with a distinct often long and slender tube, rarely two-lipped and divided partly almost to the base. Stamens 5 to 10, of unequal length. Ovary 5-celled, usually strigose and pubescent, rarely glabrous, never lepidote.

The type section of the subgenus is *Pentanthera* and the type species *Rhododendron luteum* Sweet (*Azalea pontica* Linnaeus). In North America this subgenus is represented by sixteen species.

Our conception of the subgenus coincides exactly with that of Drude (*l. c.*), while all other authors draw its limits narrower or wider, as shown in the synonymy given above. In former publications I had included *Azaleastrum* and *Therorhodion* in this subgenus, but as the inflorescences of these two groups is entirely different from that of the type of this subgenus, I prefer to consider these two groups as coördinate subgenera and restrict the sections belonging to *Anthodendron* to *Tsutsutsi* G. Don, *Sciadorhodion* Rehd. & Wils., *Rhodora* G. Don and *Pentanthera* G. Don, of which the first two are exclusively Asiatic, while the last two contain American and Asiatic species. It does not seem advisable to

maintain *Tsutsutsi* as a distinct subgenus, since it is closely connected with *Rhodora* by the new section *Sciadorhodion*.

As the name *Azalea* used first by Planchon as a subgeneric name is not the oldest subgeneric name for the group, and as, moreover, the generic name *Azalea* upon which the name of the subgenus is based properly belongs as a synonym to *Loiseleuria*,¹ either Don's *Penthanthera* or Endlicher's *Anthodendron* must take its place. As *Penthanthera* has been proposed originally as a sectional name and used as such by later authors it may be retained as the name for the section to which it should apparently be restricted by the meaning of its name, and *Anthodendron* first published in 1827 as a genus may be adopted in an enlarged sense as the proper name for the subgenus.

¹ See *Jour. Arnold Arboretum*, II. 156 (1921).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Flower-buds with prominent rows of conspicuous long-stipitate glands at apex; corolla glandular-pilose, not or slightly villose outside, the tube abruptly expanded at apex; low stoloniferous shrub with simple or little branched stems *R. atlanticum*, p. 147.

Flowers expanding after the leaves are fully developed; the flower-buds with prominent rows of stipitate glands at apex, rarely quite glabrous.

Branchlets and leaves more or less strigose or pubescent; style usually pubescent.

Calyx-lobes oblong to oblong-lanceolate, rarely short; corolla finely villose; leaves usually large, 4 to 10 cm. long; winter-buds pubescent; flowers appearing before the terminal buds are formed.

R. oblongifolium, p. 150.

Calyx-lobes short, semi-orbicular to ovate; corolla villose and glandular-pilose; leaves usually 2 to 6 cm. long; winter-buds glabrous or pubescent; flowers appearing usually when the terminal buds are forming.

Winter-buds with many (usually more than 15) aristate-mucronate scales, usually pale with conspicuous dark margin; branchlets bright red-brown, usually densely strigose, specially toward the apex; leaves often pubescent beneath, particularly at the end of shoots, serrulate-ciliate *R. serrulatum*, p. 153.

Winter-buds with fewer (usually less than 15) obtusish or acutish, usually pale brown scales; branchlets usually pale, strigillose, hispid or glabrescent; leaves glabrous, strigose on midrib beneath, pubescent only in one rare variety *R. viscosum*, p. 157.

Branchlets and leaves glabrous or the leaves sometimes strigillose on midrib beneath; style usually glabrous; winter-buds glabrous.

Corolla white or pinkish, tube glandular-pilose outside; leaves glaucous beneath, generally obovate, acute or obtusish.

R. arborescens, p. 166.

Corolla crimson, tube glabrous or nearly glabrous outside; leaves light green beneath, elliptic to oblong-lanceolate, acuminate.

R. prunifolium, p. 169. †

ENUMERATION OF THE SPECIES

SECT. I. RHODORA G. DON¹

Rhododendron sect. **Rhodora** G. Don, *Gen. Syst.* III. 848 (1834). — Endlicher, *Gen.* 759 (1839). — Dippel, *Handb. Laubholz.* I. 417 (1889), as section of subgenus *Azalea*, excluding his Nos. 29 and 31. — Koehne, *Deutsch. Dendr.* 455 (1893), as section of subgenus *Azalea*, excluding his No. 36. — Zabel in *Mitt. Deutsch. Dendr. Ges.* XI. 26 (1902), in part.

Rhodora Linnaeus, *Spec.* ed. 2, 561 (1762); *Gen.* ed. 6, 218 (1764). — Willdenow, *Spec.* I. pt. 2, 866 (1798). — De Candolle, *Prodr.* VII. 719 (1839). — Gray, *Man.* 267 (1848); ed. 5, 309 (1867). — Britton & Brown, *Ill. Fl.* II. 560 (1897). — Britton, *Man.* 699 (1901). — Small in *N. Am. Fl.* XXIX. 44 (1914).

Hochenwartia Crantz, *Inst.* II. 468 (1766).

Rhododendron sect. *Azalea*, ser. 2, [subser.] b, Maximowicz in *Mém. Acad. Sci. St. Pétersb.*, sér. 7, XVI. No. IX. 28 (1870). — Bentham & Hooker, *Gen. Pl.* II. 601 (1876), as subser. 2, in part. — Drude in Engler & Prantl, *Nat. Pflanzenfam.* IV. 1, 37 (1889), as subgenus *Azalea*, § 4.

Biltia Small, *Fl. S. E. U. S.* 884 (1903); in *N. Am. Fl.* XXIX. 48 (1914).

Deciduous upright shrubs without strigose pubescence, densely or loosely branched, sometimes irregularly dichotomous, but not whorled; branchlets glabrous or pilose to hirsute, winter-buds of many imbricate scales, glabrous or puberulous. Leaves scattered or crowded at the end of the branchlets. Flowers from a terminal bud, the leafy shoots from separate lateral buds; corolla two-lipped, or rotate-campanulate, rarely campanulate; stamens 10, or in one species 7 or occasionally 5 or 6, not or only little longer than the corolla, usually declinate; ovary pubescent or glabrous and stipitate-glandular. Capsule usually with rather thin valves.

The type species of this section is *R. canadense* Torr. The section contains five species, of which two are native of eastern North America and three are Japanese. All these species are very distinct from each other and may be divided into three groups. The first group, containing only *R. canadense*, is characterized by the deeply two-lipped corolla, the lower lip divided nearly to the base, the glandular-pubescent and sparingly setose ovary, the puberulous branchlets and the tomentulose under side of the usually oblong leaves and the glandular-puberulous narrowly oblong capsule. The second group with three species, of which *R. Vaseyi* is one, has a rotate-campanulate slightly two-lipped corolla, a stipitate-glandular but otherwise glabrous or sometimes quite glabrous ovary, an ovoid to oblong glandular-hirsute or stipitate-glandular or glabrous capsule, and glabrous or pilose leaves and branchlets. The third group, consisting only of *R. nipponicum*, has a campanulate nearly regular corolla, with the lobes shorter than the tube, a glandu-

¹ See page 91 for the Old World species of this section.

lar-pilose but otherwise glabrous ovary, hirsute branchlets and hirsute large obovate leaves, a very thin-valved oblong-ovoid capsule almost destitute of glands at maturity; its flowers appear after the full development of the leaves, while in all the other species of this group they open before the leaves unfold. The petiole of the shoot-leaves in *R. Vaseyi*, *R. nipponicum* and *R. Albrechtii* is somewhat decurrent on the stem, leaving an elongated scar distinctly longer than broad.

In the section *Rhodora* I have included *R. Vaseyi* as suggested by Gray and as done by Dippel and by Koehne; by Small this has been made the type of a new genus *Biltia*, chiefly distinguished by the exterior aestivation of the middle lobe of the upper lip of the corolla. This position of the posterior lobe is without doubt unusual in the genus, but can be observed at least occasionally in species related to *R. Vaseyi*. Among flowers of *R. canadense* I have found a small percentage with the middle lobe of the upper lip exterior and in *R. Schlippenbachii*, which belongs to the section nearest to *Rhodora* and is very similar in its flowers to *R. Albrechtii* and *R. quinquefolium*, both closely related to *R. Vaseyi*, I have found after examining numerous fresh flowers just before opening that about 15 per cent of the flowers had the middle lobe of the upper lip completely exterior and 25 to 30 per cent had one side exterior and the other side covered. This shows that the position of the upper lobe cannot be used as a reliable character to separate *Biltia* generically. As there is no other character to separate *R. Vaseyi* from the other species of the section *Rhodora* except the number of stamens, which also cannot be considered a reliable character in the genus *Rhododendron*, it seems best to retain it in this section.

Rhododendron canadense Torrey, *Cat. Pl.* 151 (in *Geol. Surv. New York Assembly*, No. 50) (1839). — Oakes, *Cat. Vermont Pl.* 190 (in Thompson, *Hist. Vermont*) (1842). — Wood, *Classb. Bot.* 235 (1845). — Zabel, *Syst. Verz. Muenden*, 26 (1878). — Britton, Sterns, Poggenburg, *Prelim. Cat. New York*, 33 (1888). — Britton, *Cat. Pl. N. Jersey*, 162 (1889). — Dippel, *Handb. Laubholz.* I. 418 (1889). — Robinson & Fernald, *Gray's New Man.* ed. 7, 631 (1908). — Schneider, *Ill. Handb. Laubholz.* I. 494, fig. 327 i-k, 328 f (1909). — Jackson, *Cat. Fl. Worcester Co., Mass.* 76 (1909). — Graves & others, *Cat. Flow. Pl. Conn.* 308 (1910). — Twining, *Fl. N. E. Pennsylv.* 56 (1917).

Rhodora canadensis Linnaeus, *Spec. ed.* 2, 561 (1762). — Willdenow, *Spec. I.* pt. 2, 866 (1798). — Michaux, *Fl. Bor.-Am.* I. 259 (1803). — Pursh, *Fl. Am. Sept.* I. 298 (1814). — Bigelow, *Fl. Boston*, 104 (1814). — De Candolle, *Prodr.* VII. 719 (1839). — Gray, *Man.* 267 (1848); ed. 5, 309 (1867). — Tracy, *Stud. Essex Fl.* 49 (1858). — Britton & Brown, *Ill. Fl.* 560, fig. 2748 (1897). — Clute, *Fl. Upp. Susquehanna*, 70 (1898). — Britton, *Man.* 699 (1901). — Small, *Fl. Pennsylv.* 238 (1903); in *N. Am. Fl.* XXIX. 44 (1914).

Hohenwartia canadensis Crantz, *Inst.* II. 469 (1766).

Rhododendron Rhodora G. F. Gmelin, *Syst. Nat.* II. pt. 1, 694 (1791). — Sweet, *Hort. Brit.* ed. 2, 343 (1830). — G. Don, *Gen. Syst.* III. 848 (1834). — Gray, *Syn. Fl.* II. 1, 41 (1878). — J. Robinson, *Fl. Essex Co., Mass.* 73 (1880). — Macoun, *Cat. Canad. Pl.* II. 302 (1884). — Dame & Collins, *Fl. Middlesex Co., Mass.* 63 (1888). — Perkins, *Cat. Fl. Vermont*, 33 (1888). — Bennett, *Pl. Rhode Isl.* 26 (1888). — Coulter & Watson, *Gray's Man.* ed. 6, 321 (1890). — Rand & Redfield, *Fl. Mt. Desert Isl.* 127 (1894). —

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

relation to *Rhododendron* was perceived as early as 1791 by Gmelin who referred it to that genus. The species was first described and figured by Duhamel from a cultivated plant which flowered the first time in March, 1756, in the Botanic Garden at Paris, where it had been brought from Canada a few years earlier. Into England it was introduced by Joseph Banks in 1767.

Observations on the pollination of this species have been made by F. Hildebrand (*Flora*, LXIV. 501, t. 10, fig. 6, 7 (1881)), who describes the peculiar arrangement to avoid self-fertilization. When the flower opens, the apex of the style with the stigma remains enclosed in a little hood formed by the lobes of the upper lip, while visiting insects carry away pollen from the anthers; further elongation of the style forces it out of the hood and the bent style stretches itself, extending the stigma beyond the stamens, and is then in a position to be fertilized by insects carrying pollen from other flowers.

***Rhododendron canadense* f. *albiflorum* Rehder, comb. nov.**

Rhododendron Rhodora f. *albiflora* Rand & Redfield, *Fl. Mt. Desert Isl.* 127 (August, 1894).

Rhododendron canadense f. *album* Voss, *Vilmorin's Blumengärt.* I. 587 (Sept. or Oct., 1894). — Zabel in *Mitt. Deutsch. Dendr. Ges.* XI. 31 (1902).

A form with white flowers, occasionally found wild under the typical form, e.g. Maine, Mt. Desert, Southwest Harbor.

***Rhododendron canadense* f. *viridifolium* Fernald, n. forma.**

Ramulis glabriusculis; foliis subtus sparse pilosis supra atroviridibus lucidis nec glaucis.

NOVA SCOTIA: boggy thickets bordering Trefry's Lake, Arcadia, July 29, 1920, *Fernald & Long*, No. 22,150 (TYPE in Gray Herb.).

This shrub occurs in scattered clumps on the shore of Trefry's Lake, the only place we met it, the ordinary form of the species with glaucous foliage being generally common in the province. — M. L. FERNALD.

***Rhododendron Vaseyi* A. Gray** in *Proc. Am. Acad.* XV. 48 (1879); in *Bot. Gaz.* VIII. 282 (1883); *Syn. Fl.* ed. 2, II, 1, 398 (1886). — Donnell Smith in *Bull. Torr. Bot. Club*, XV. 164 (1888). — Sargent in *Gard. & For.* I. 376, fig. 60 (1888). — Dippel, *Handb. Laubholz.* I. 419 (1889). — Skan in *Bot. Mag.* CXXXII. t. 8081 (1906). — Schneider, *Ill. Handb. Laubholz.* II, 495, fig. 325 o, 327 c-d (1909). — Millais, *Rhodod.* 257, t. opp. p. 10 and 20 (1917).

Azalea Vaseyi Rehder in *Möller's Deutsch. Gärt.-Zeit.* XIV. 332, fig. (1899); in *Bailey, Cycl. Am. Hort.* I. 122 (1900).

Biltia Vaseyi Small, *Fl. S. E. U. S.* 884 (1903); in *N. Am. Fl.* XXIX. 49 (1914).¹

¹ ADDITIONAL ILLUSTRATIONS. COLORED PLATES: *Meehan's Monthl.* VII. 121, t. 7 (1897). — *Garden*, LIV. t. 1191, opp. p. 282 (1898). — BLACK FIGURES: *Gard. Chron.* sér. 3, XX. 71 (1896); LI. 313 (1912). — *Möller's Deutsch. Gärt.-Zeit.* XIV. 332 (1899). — *Gard.* LVI. 122 (1899); LXXVI. 332 (1912); LXXXIV. 239 (1920). — *Gard. Mag.* V. 219 (1907). — *Country Life Am.* XI. 499 (1907). — *Bean, Trees & Shrubs Gt. Brit.* II. 384 (1914). — *Bailey, Stand. Cycl. Hort.* V. fig. 3391 (1916), after *Gard. & For.* — HABIT FIGURES: *Möller's Deutsch. Gärt.-Zeit.* XIV. 333 (1899). —

An upright, rather irregularly branched shrub with spreading branches, occasionally 5 m. tall; young branchlets puberulous and sparingly pilose at first, soon glabrous, light red-brown at the end of the season, becoming brown or grayish brown the second year; older branches grayish brown with flaky or shredding bark; floral winter-buds large, broadly ovoid, obtuse, with 7 to 10 orbicular-ovate scales, rounded or the upper ones truncate or slightly emarginate at apex, mucronulate, the basal ones acuminate, glabrous or the upper and inner ones minutely puberulous, white-ciliate and sometimes glandular on the margin. Leaves elliptic or elliptic-oblong, acuminate, cuneate at base, 5 to 12 cm. long and 2 to 5 cm. broad, usually slightly undulate on the margin, ciliate, dark green and glabrous above except the sparingly and finely villose midrib, rarely sparingly strigillose, light green beneath and glabrous or sparingly pilose on the midrib; petioles slender, 3 to 7 mm. long, glabrous or sparingly pilose. Flowers appearing in May before the leaves, in 5- to 8-flowered umbel-like racemes; pedicels 0.5 to 1.5 cm. long, with short-stipitate glands, sometimes sparingly so; calyx oblique, with shallow rounded lobes, erose-glandular on the margin; corolla rotate-campanulate 2.5 to 3 cm. long, two-lipped, light rose-colored with orange or red-orange dots at the base of the upper lobes, glabrous, tube very short, about 5 mm. long, the lobes oblong, rounded at apex, the upper lip less deeply divided, with the middle lobe exterior in bud; stamens usually 7, sometimes 5 or 6, unequal, the longer ones exceeding the corolla; filaments glabrous; anthers ellipsoid, 1 to 2 mm. long, pinkish; style 2.5 to 3 cm. long, longer than stamens, glabrous or with few stipitate glands near the base. Capsule narrow-oblong, 1.2 to 1.5 cm. long, with a thin narrow keel on back of the rather thin valves.

This species is very distinct from all other American Azaleas and seems most nearly related to the Japanese *R. Albrechtii* Maxim. which, however, differs considerably in its 10 stamens, the more pubescent obovate leaves and the shorter and thicker more woody, hispid capsule. It is restricted to the higher mountains of western North Carolina, where it grows at an altitude of from 3000 to 5500 feet on mountain slopes, in ravines and sometimes in swamps. It was first discovered in 1878 on Balsam Mountain near Webster, Jackson County, by George Vasey and soon afterward introduced into cultivation, for in 1880 the Arnold Arboretum received a living plant from G. C. Woolson & Co. of Passaic, New Jersey. It has proved perfectly hardy near Boston and is now well known in gardens in this country as well as in Europe and valued for its handsome pale pink flowers appearing in great profusion in spring before the leaves unfold. It has been found escaped from cultivation near Halifax, Mass., as shown by the following specimens in the herbarium of the New England Botanical Club: swamp near an abandoned nursery, May 30, 1907, A. S. Pease (No. 9998); sandy soil, probably part of old nursery, May 30, 1907, J. A. Cushman (No. 653).

Rhododendron Vaseyi f. **album** Nicholson in *Kew Hand-list Arb.* II. 57 (1896), as var., name only. — Bean, *Trees & Shrubs Gt. Brit.* II. 384 (1914), as var. — Rehder in *Mitt. Deutsch. Dendr. Ges.* XXIV. (1915), 225 (1916).

This form with white flowers was found at the Arnold Arboretum and also in Kew Gardens among seedlings of the typical form; all transitions from pure pink to white can be found among seedlings of this species.

Gard. LVI. 119 (1899). — *Gartenwelt*, XVI. 231 (1912); XXII. 363 (1918). — Silva-Tarouca, *Uns. Freil.-Laubgeh.* fig. 64 (1913). — *Gard.* LXXIX. 302 (1915). — *Gard. Mag.* XXX. 259 (1920). — *Gartenschönheit*, I. 33 (1920).

SECT. II. PENTANTHERA G. DON

Rhododendron sect. **Pentanthera** G. Don, *Gen. Syst.* III. 846 (1834). — Rehder & Wilson in Sargent, *Pl. Wilson.* I. 549 (1913).

Azalea Linnaeus, *Spec.* 150 (1753), as to species 2 to 4. — De Candolle, *Prodr.* VII. 715 (1838), excluding species 13, 14, 16. — Britton & Brown, *Ill. Fl.* II. 558 (1897).

Rhododendron a. *Anthodendron* Endlicher, *Gen.* 759 (1839). — Zabel in *Mitt. Deutsch. Dendr. Ges.* XI. 28 (1902), as section.

Theis "Salisb. in coll. 1817" ex De Candolle, *Prodr.* VII. 715 (1839), synonym.

Rhododendron sect. *Azalea* Regel in *Act. Hort. Petrop.* I. 162 (1871), excluding species 5 and 8.

Azalea subgen. *Pentanthera* K. Koch, *Dendr.* II. 1, 186 (1872).

Azalea subgen. *Euazalea* K. Koch, *Dendr.* II. 1, 179 (1872), as to species 11, 12.

Rhododendron sect. *Euazalea* Dippel, *Handb. Laubholz.* I. 411 (1889). — Koehne, *Deutsch. Dendr.* 458 (1893).

Deciduous upright shrubs, rarely almost tree-like, sometimes stoloniferous, with usually irregularly whorled branches; branchlets usually more or less strigose or hirsute, sometimes glabrous. Winter-buds with many imbricate scales. Leaves scattered. Flowers several to many from a terminal bud, the leafy shoots from separate lateral buds; corolla funnel-form, subregular, the tube about as long or longer than the limb; stamens 5, usually much exserted; ovary more or less strigose or setose. Capsule with woody valves.

The type species of this section is *Rhododendron luteum* Sweet. The section contains besides the sixteen American species enumerated below, two species from eastern Asia, *R. molle* Miq. and *R. japonicum* Suring., and one from southeastern Europe and western Asia, *R. luteum* Sweet. In America it occupies two distinct geographical areas: one on the Atlantic coast, extending from southern Quebec to central Florida and west to Vermont, eastern Ohio, southeastern Missouri, central Alabama, extreme eastern Oklahoma and eastern Texas; and one on the Pacific coast from southern Oregon to southern California.

From the section *Rhodora* this section is easily distinguished by the funnel-form subregular corolla with a distinct cylindrical tube longer or little shorter than the lobes and by the 5 stamens; the ovary is always densely villose and strigillose, with or without glands. Most species show a wide range of variation particularly in the pubescence of the leaves and in the color of the flowers. The leaves vary from nearly glabrous to pubescent in *R. oblongifolium*, *R. serrulatum*, *R. canescens*, *R. roseum*, *R. occidentale* and *R. atlanticum*; they are always glabrous except the midrib in *R. nudiflorum*, *R. arborescens* and *R. prunifolium* and usually so in *R. viscosum*; they are always pubescent in *R. alabamense* and, though usually only slightly so, in *R. calendulaceum* and *R. speciosum*. The pubescence of the winter-buds which affords a character hitherto neglected, shows less variation; they vary from glabrous to finely pubescent in *R. nudiflorum*, *R. viscosum*, *R. serrulatum*, *R. atlanticum* and *R. occidentale*; they are always glabrous in *R. calendulaceum*, *R. speciosum*, *R. arborescens*, *R. alabamense* and *R. prunifolium* and always pubescent in *R. canescens*, *R. roseum* and *R. oblongifolium*. The branchlets are in most species more or less pubescent, usually villose and strigillose, but are always

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Handb. Laubholz. I. 415, fig. 268 (1889). — Jepson, *Fl. W. Middle Cal.* 369 (1901); ed. 2, 311 (1911). — Schneider, *Ill. Handb. Laubholz.* II. 501, fig. 328 l-m, 329 i-k (1911). — Millais, *Rhodod.* 220 (1917).

Azalea calendulacea Hooker & Arnott, *Bot. Capt. Beechey Voy.* 362 (1841), not Michaux. — Bentham, *Pl. Hartweg.* 321 (1849).

Azalea nudiflora var. *ciliata* Kellogg in *Proc. Calif. Acad. Sci.* I. 60 (1855).

Azalea californica Torrey & Gray apud Durand in *Jour. Acad. Philadelphia*, ser. 2, III. 94 (1855). — Small in *N. Am. Fl.* XXIX. 44 (1914).

Azalea occidentalis Torrey & Gray in *Rep. Explor. Surv. Miss. Pacif. Ocean*, IV. 116 (1856). — Hooker in *Bot. Mag.* LXXXIII. t. 5005 (1857). — Howell, *Fl. N. W. Coast*, I. 422 (1901).¹

A shrub to 3 m.; young branchlets soft-pubescent or glabrous, brown at the end of the first season; floral winter-buds finely pubescent or sometimes nearly glabrous with numerous ovate to ovate-lanceolate, acuminate and aristate-mucronate or mucronulate scales. Leaves elliptic to oblong-lanceolate, acute or obtusish, gland-tipped, cuneate or attenuate at base, 3 to 9 cm. long, ciliate, thinly pubescent on both surfaces, rarely nearly glabrous, at maturity chartaceous, glabrous or nearly glabrous except pubescent on the midrib above and sparingly strigose on the midrib and sometimes on the veins beneath; petioles rather stout, 1 to 5 mm. long. Flowers appearing with or after the leaves or rarely before the leaves, 6 to 12 in umbel-like racemes; pedicels 0.5 to 1.5 cm. long, pubescent and more or less glandular, sometimes pilose and glandless, rarely nearly glabrous; sepals broadly ovate to oblong-ovate, obtuse, 1.5 to 4 or sometimes 5 mm. long, densely ciliate with usually setose, gland-tipped or glandless hairs; corolla white or sometimes pink with a yellow blotch on the upper lobe or on the three upper lobes, the tube about 2 cm. long, or sometimes shorter, gradually dilated upwards, villose and glandular-pilose outside, the lobes about as long as the tube, broad-ovate, acute or acutish; stamens exserted, more than twice as long as the tube; filaments pilose below the middle; anthers 2.5 to 3 mm. long, whitish; style about as long or longer than stamens, finely pubescent or sometimes villose above the base, rarely nearly glabrous; ovary pilose and glandular. Capsule ovate-oblong, 1 to 2 cm. long, setosely pilose or sometimes furnished with short gland-tipped hairs.

This species is distributed from southern Oregon to southern California, growing by brooks among the mountains up to 5000 or sometimes 5800 feet altitude. It is the only *Azalea* west of the Rocky Mountains and was first discovered during the Expedition of Captain Beechey in 1827 and was also collected later by Douglas, Hartweg and Burke. Hooker as well as Bentham referred it to *A. calendulacea*, and it was not recognized as a distinct species until named in 1855 by Torrey & Gray *A. californica* and a year later by them *A. occidentale*.

Rhododendron occidentale seems most closely related to *R. calendulaceum*, from which it is easily and chiefly distinguished by the color of its flowers, which otherwise in size and shape resemble those of that species more than those of other American species. It is very variable in the shape and size of its leaves, in the pubescence of all its parts and also to some extent in the size of the flowers; the sepals are apparently always well developed, never nearly obsolete, as is often the case in the Eastern

¹ ADDITIONAL ILLUSTRATIONS. COLORED PLATES: *Fl. des Serres*, XIV. t. 1432 (1861). — *Gard.* XXXIV. 416, t. 673 (1888). — BLACK FIGURE: *Gard. Chron.* ser. 2, XXVI. 105, fig. 21 (1886).² — HABIT FIGURES: *Gartenwelt*, XI. 8 (1906); XV. 650 (1911).

² The legend under the wood-cut reads "flowers orange and crimson, fragrant" which is hardly correct; in the text the flowers are described as deep rosy-red.

species. It exhibits a remarkable latitude in the time of blooming; I have before me a specimen collected by Orcutt in San Diego County in April without any leaf buds developed and in northern California I collected it myself in full bloom as late as August 25 with the winter-buds of the new growth fully developed; generally it flowers between the middle of June and the middle of July, just when the young leaves are fully grown. The autumnal tints of this species are very brilliant. In *Garden and Forest* (III. 626 [1890]), E. W. Hammond says "in the fall of the year its leaves glow with such a profusion of scarlet and crimson mingled with the gold and green, that it produces the most delightful effects."

Behr (*Fl. Vicin. San Francisco*, 173) states that the root contains a powerful narcotic, and according to Jepson (*Fl. W. Middle Calif.* ed. 2, 311) the plant is feared by sheepmen as poisonous.

Rhododendron occidentale was introduced into cultivation by William Lobb, who sent seeds from California about 1850 to the Veitchian Nurseries, where it flowered in 1857. It is tenderer than many other American Azaleas and has not proved hardy at the Arnold Arboretum.

***Rhododendron occidentale* var. *sonomense* Rehder, var. nov.**

Rhododendron sonomense Greene in *Pittonia*, II. 172 (1891). — Zabel in *Mitt. Deutsch. Dendr. Ges.* XI. 29 (1902). — Millais, *Rhodod.* 245 (1917).

This variety differs from the type according to Greene chiefly in its smaller narrower leaves, smaller rose-colored and very fragrant flowers; the odor of typical *R. occidentale* is mephitic according to Greene. It occurs on dry slopes in the mountains of Sonoma County. It does not seem to be in cultivation outside of California.

***Rhododendron calendulaceum* Torrey, *Fl. U. S.* 425 (1824). — G. Don, *Gen. Syst.* III. 847 (1834). — Bigelow & Hor, *Fl. Lancast.* 56 (in *Trans. Med. Convent. Ohio*) (1841); Reprint, p. 10 (1841). — Gray, *Syn. Fl. N. Am.* II. 1, 41 (1878). — Dippel, *Handb. Laubholz.* I. 413, fig. 266 (1889). — Millspaugh, *Prelim. Cat. Fl. W. Va.* 402 (1892). — Blake in *Rhodora*, XX, 53 (1918).**

Azalea lutea Linnaeus, *Spec.* 150 (1753), only as to Colden's synonym. — Weston, *Bot. Univ.* I. 16 (1770). — Michaux, *Jour.* ed. C. S. Sargent, 111, 116 (in *Proc. Am. Philos. Soc.* XXVI.) (1889). — Britton & Brown, *Ill. Fl.* II. 559, fig. 2745 (1897). — Gattinger, *Fl. Tenn.* 131 (1901). — Britton, *Man.* 698 (1901). — Porter, *Fl. Pennsylv.* 238 (1903). — Millspaugh, *Liv. Fl. W. Va.* 323 (*W. Va. Geol. Surv.*) (1913). — Small, *Fl. S. E. U. S.* 883 (1903); in *N. Am. Fl.* XXIX. 41 (1914). — Shreve & others, *Pl. Life Maryland*, 466 (1910). — Schaffner, *Cat. Ohio Vasc. Pl.* 204 (in *Ohio Biol. Surv.* I.) (1914). — Griggs, *Bot. Surv. Sugar Grov. Reg.* 328 (*Ohio Biol. Surv.* I.) (1914). — Taylor, *Fl. Vicin. N. Y.* 489 (in *Mem. N. Y. Bot. Gard.* V.) (1915).

Azalea nudiflora Linnaeus, *Spec.* ed. 2, 214 (1762), as to Colden's synonym only.

Azalea flammea Bartram, *Travels*, I. 327 (1791), name only.

Azalea calendulacea Michaux, *Fl. Bor.-Am.* I. 151 (1803), excluding var. a. — Pursh, *Fl. Am. Sept.* 151 (1814). — Elliott, *Sketch Bot.* I. 238 (1821), excluding var. a. and c. — De Candolle, *Prodr.* VII. 717 (1839). — Gray, *Man.* 268 (1848); ed. 5, 299 (1872). — M. A. Curtis, *Descript. Trees & Shrubs*, 98 (in *Geol. Nat. Hist. Surv. N. Car.* III. Bot.) (1860).

Azalea aurantiaca Dietrich, *Darst. Vorz. Zierpfl.* 4, t. 1 (1803).¹

Azalea pontica β. *A. calendulacea* Persoon, *Syn.* I. 212 (1805).

? *Azalea nudiflora* var. *h. lutea* Elliott, *Sketch Bot.* I. 241 (1821).

? *Azalea flava* Michaux, *Jour.* ed. C. S. Sargent, 111 (in *Proc. Am. Philos. Soc.* XXVI.) (1889), name only.

Rhododendron luteum Schneider, *Ill. Handb. Laubholz.* II. 500, fig. 329 q-r, 320 a (1911), not Sweet.²

A shrub with erect and spreading branches, usually 1.5 to 3, rarely to 5 m. tall; young branchlets pubescent and strigillose; floral winter-buds with broadly ovate scales, rounded or even emarginate at the apiculate apex, brown, glabrous except the densely white-ciliolate margin. Leaves broadly elliptic to elliptic-oblong or obovate-oblong, rarely obovate, 4 to 8 cm. long, and 1.5 to 3.5 cm. broad, acute and gland-tipped, broadly cuneate at base, finely pubescent above and more densely so beneath when young, particularly on and along the midrib; petioles 2 to 5 mm. long, pubescent. Flowers in May and June, in usually 5- to 7-flowered umbels, expanding with or shortly after the leaves; pedicels setulose and usually glandular; calyx-lobes usually oblong and 3 to 4 mm. long or ovate and 1 to 2 mm. long, obtuse, setulose outside, ciliate with long gland-tipped hairs; corolla funnel-form, yellow or orange to scarlet, with orange blotch on the upper lip, the cylindrical tube gradually dilated above the middle, 1.5 to 2 cm. long, glandular-pilose, and pubescent outside, the lobes ovate, about 2 cm. long and 1.5 cm. broad, usually wavy at the margin, abruptly contracted into a short point, slightly pubescent outside; stamens nearly three times as long as corolla-tube; the filaments pubescent below the middle or on the lower third, the anthers about 3 mm. long; style pilose toward the base, sometimes glabrous, 6 to 7 cm. long, about as long or longer than stamens; ovary covered with long setose, partly gland-tipped hairs. Capsule ovoid-oblong, 1.5 to 2 cm. long, setose and pubescent.

This species is confined to the Appalachian Mountain region from Pennsylvania to northern Georgia; according to the herbarium material seen, its western limit runs from western Pennsylvania (Venango County) through southeastern Ohio (Fairfield County),³ eastern Kentucky (Rockcastle County), eastern Tennessee (Knox County) to northern Georgia (Gwinnett County); its eastern limit

¹ Dippel and Schneider cite 1801 as date of publication, which would give Dietrich's name priority over *A. calendulacea*, but the part which contains *A. aurantiaca* is dated 1803 as well on the title-page as on the cover. The cover bears the title "Modeblumen für Botaniker . . . Sechstes Heft." The first part of the latter publication is the one which is dated 1801. There is no technical description given, and this together with the fact that the part was published during the second half of the year, as it contains a letter dated July 1, 1803, makes it advisable to give Michaux's name the preference.

² PRELINNEAN NAME: *Azalea erecta foliis ovatis integris alternis flore luteo piloso praecoci* Colden in *Act. Soc. Sci. Upsala*, IV. 93 (1749).

ADDITIONAL ILLUSTRATIONS. BLACK FIGURES: Loudon, *Arb. Brit.* II. 1142, fig. 946 (1838); reproduced in *Gard.* XXIX. 550 (1886).—Britton & Brown, *Ill. Fl.* II. fig. 2745 (1897); ed. 2, II. fig. 3217 (1913).—Keeler, *Our North. Shrubs*, 351 (1903).—*Gard. Mag.* V. 218 (1907).—*Country Life Am.* XI. 496 (1907).—Stevens, *Ill. Guide Flow. Pl.* t. 114, fig. 3 (1910).—Schneider, *Ill. Handb. Laubholz.* I. fig. 329 q-r, 330 a (1911).

³ This station, interesting on account of its isolation, was first discovered in 1841 according to Bigelow & Hor, *Fl. Lancast.* 56 (1841): "For the discovery of this splendid plant . . . we are indebted to the hospitality of Mr. Jacob Greene, who had found it a few days previously at the mouth of Clear Creek."

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Small in *North American Flora* gives "Savannah River" as the type locality for his *A. lutea*, but if *A. lutea* of Linnaeus is taken up as the name for *R. calendulaceum* it could be based only on Cadwallader Colden's enumeration of the plant from southern New York and not on the locality given by Michaux for var. *a.* of his *A. calendulacea*. Unfortunately Michaux's specimen collected on the Savannah River near Two-Sisters Ferry and cited by him under *a. flammea* does not belong to *A. calendulacea* as described by him, but to *R. speciosum*, and therefore the specimen labeled in his herbarium *β. crocea* must be considered the type specimen and the high mountains of Carolina or more particularly the Blue Ridge Mountains of North Carolina, the type locality. Though it is against general usage to take var. *β.* as the type of a species which is divided into varieties, the specific description in this case appears to be chiefly based on the specimen designated var. *β.*, particularly the character "corolla tubo hirsuto, laciniis brevior" must have been taken from var. *β.*, for in var. *a.* (*R. speciosum* Sweet) the tube is distinctly longer than the lobes and less conspicuously hirsute, also the color "floribus . . . calendulaceis" referring to *Calendula* which has yellow to orange flowers, indicates the mountain form, var. *β.*, rather than var. *a.*, which has bright red to scarlet flowers.

Rhododendron calendulaceum does not show much variation except in the color of the flowers and in the length of the sepals, which are usually oblong, but sometimes rather small; the style is normally pilose near the base, but sometimes glabrous, as in Harbison's No. 171 from Macon County, North Carolina, in Small's specimen from White Rock Mountains, Smyth County, Va., and in Britton's & A. M. Vail's specimen from the same locality. There are also forms with partly salmon-colored flowers of which I collected a specimen myself on Roan Mountain of North Carolina, and I have before me a specimen from North Carolina stated to have pink flowers. The two following forms have been described, of which the first represents the typical form, though probably more color forms could be distinguished but this cannot be done satisfactorily without copious living material of undoubtedly spontaneous origin.

***Rhododendron calendulaceum* f. *croceum* Sweet, *Hort. Brit.* ed. 2, 343 (1830), as var. *a.* — Rehder in *Mitt. Deutsch. Dendr. Ges.* XXIV. (1915), 225 (1916), as forma; in Bailey, *Stand. Cycl. Hort.* V. 2942 (1916), as var.**

Azalea calendulacea *β. crocea* Michaux, *Fl. Bor.-Am.* I. 151 (1803). — Sims in *Bot. Mag.* XLI. t. 1721 (1815). — Rehder in Bailey, *Cycl. Am. Hort.* I. 121, (1900).¹

Flowers yellow to orange-yellow with a darker blotch, sometimes partly salmon-colored. As stated before there are all kinds of intermediate shades between the plants with scarlet flowers and those with yellow flowers.

***Rhododendron calendulaceum* f. *aurantium* Rehder, comb. nov.**

Azalea calendulacea *a.* Ker in *Bot. Reg.* II. t. 145 (1816), not var. *a. flammea* Michaux.

Azalea aurantiaca Dietrich, *Darst. Vorz. Zierpfl.* 4, t. 1 (1803). — Hort. apud Dumont de Courset, *Bot. Cult.* ed 2, III. 333 (1811), as var. of *A. nudiflora*.
? *Azalea coccinea major* Dumont de Courset, *l. c.* (1811), as var. of *A. nudiflora*.

¹ ADDITIONAL REFERENCE TO COLORED PLATE: *Mag. Bot. & Gard.* II. 119, t. 36, fig. 7 (1834).

- Azalea speciosa* f. *aurantia* Sweet, *Hort. Brit.* 265. (1826), name only.
Azalea coccinea aurantia Loddiges, *Bot. Cab.* XIII. t. 1255 (1827).
Rhododendron calendulaceum var. *a. flammeum* Sweet, *Hort. Brit.* ed. 2, 343 (1830). — Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz.-Ben.* 380 (1903).
Rhododendron speciosum f. *aurantium* Sweet, *Hort. Brit.* ed. 2, 343 (1830).
Azalea speciosa β. *aurantia* De Candolle, *Prodr.* VII. 117 (1838).
Azalea speciosa α. *aurantiaca* Kirchner in Petzold & Kirchner, *Arb. Musc.* 479 (1864).
Azalea calendulacea var. *flammea* Rehder in Bailey, *Cycl. Am. Hort.* I. 121 (1900), in part.
Rhododendron calendulaceum aurantiacum Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz.-Ben.* 380 (1903).
Rhododendron luteum var. *flammeum* Schneider, *Ill. Handb. Laubholzk.* II. 500 (1911), in part.

Flowers deep orange-red to scarlet, with large orange blotch.

Rhododendron speciosum Sweet, *Hort. Brit.* ed. 2, 343 (1830), not Salisbury.¹ — G. Don, *Gen. Syst.* III. 847 (1834).

- Azalea nudiflora* α. *coccinea* Aiton, *Hort. Kew.* I. 202 (1789). — Sims in *Bot. Mag.* V. t. 180 (1792).
Azalea coccinea Sims in *Bot. Mag.* text to t. 180 (1792), as synonym. — Michaux, *Jour.* ed. C. S. Sargent, 9 (in *Proc. Am. Philos. Soc.* XXVI.) (1889).
Azalea fulva Michaux in *Jour. Hist. Nat.* I. 410 (1792), name only.²
Azalea calendulacea α. *flammea* Michaux, *Fl. Bor.-Am.* I. 151 (1803). — Pursh, *Fl. Am. Sept.* 152 (1814).
Azalea speciosa Willdenow, *Berl. Baumz.* ed. 2, 49 (1811); *Enum. Hort. Berol. Suppl.* 10 (1813). — Guimpel, Otto & Hayne, *Abb. Fremd. Holzart.* I. 37, t. 31 (1825). — Watson, *Dendr. Brit.* II. t. 116 (1825).
Azalea periclymenoides var. *coccinea* Pursh, *Fl. Am. Sept.* I. 152 (1814).
Azalea nudiflora Loiseleur-Deslongchamps, *Herb. Amat.* IV. 213, t. (1820), not Linnaeus.
Azalea calendulacea var. *a.* Elliott, *Sketch Bot.* I. 239 (1821).
Azalea coccinea major Loddiges, *Bot. Cab.* VII. t. 624 (1822).
Azalea speciosa α. *major* Sweet, *Hort. Brit.* 265 (1826).

¹ *Rhododendron speciosum* Salisbury, *Prodr.* 287 (1796) cannot invalidate Sweet's combination, as it is a straight synonym, being only a new name for *R. ponticum* L.

² This is apparently the *Azalea* which he calls *Azalea coccinea* in his *Journal* (ed. C. S. Sargent, p. 9) and describes under the entry of the following day (p. 10) as "l'Azalea couleur de feu. La couleur de cet Azal. qui est dans toutes les parties de la fl. aussi foncée, Corolla, Etam. et Pistille, est celle de *Hemerocallis fulva* mais dans les lieux plus découverts et moins ombragés cet couleur est encore plus forte." This *Azalea* Michaux collected on April 26 and 27, 1787, after having crossed the Savannah River at Two-Sisters Ferry; in his *Flora* it is referred to under *A. calendulacea* α. *flammea*. In comparing the color of this plant with that of *Hemerocallis fulva* Michaux must have had in mind the color form figured by Redouté in his *Liliaceae*, tab. 16, which has red-purple flowers and was probably at that time well known in France. The color of this form agrees very well with that of *R. speciosum*, while the plant now generally cultivated as *Hemerocallis fulva* has orange flowers.

? *Azalea nudiflora* var. *thyrsiflora* Gowen apud Lindley in *Bot. Reg.* XVI. t. 1367 (1830).

Rhododendron speciosum a. *major* Sweet, *Hort. Brit.* ed. 2, 343 (1830).

Rhododendron nudiflorum ζ. *coccineum* Sweet, *Hort. Brit.* ed. 2, 343 (1830). — G. Don, *Gen. Syst.* III. 847 (1834).

Azalea speciosa a. *coccinea* De Candolle, *Prodr.* VII. 717 (1839).

Azalea calendulacea Darby, *Bot. S. States*, 422 (1855), in part.

Rhododendron calendulaceum Chapman, *Fl. S. U. S.* 265 (1860), in part.

Rhododendron calendulaceum f. *speciosum* Voss, *Vilmorin's Blumengärt.* I. 588 (1894). — Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz-Ben.* 380 (1903).

Rhododendron flammeum Sargent in *Rhodod. Soc. Notes*, I. 120 (1918), name only, not *Azalea flammea* Bartr.¹

A shrub 0.3 to 2 m. tall with slender, irregularly whorled branches; the young branchlets finely pubescent and rather densely strigillose, becoming grayish brown the second year; floral winter-buds glabrous, with generally ovate, ciliolate scales, usually rounded and mucronate at apex. Leaves obovate or elliptic to oblong, acute or obtusish, mucronulate, broad-cuneate at base, 3 to 6 cm. long and 1.2 to 2 or sometimes to 3 cm. broad, strigillose above, finely pubescent beneath, more densely on the leaves of the shoots and particularly on the veins, often strigose on the midrib, sometimes nearly glabrous except on the midrib beneath, setosely ciliate, subchartaceous at maturity, with prominent lateral veins; petioles 3 to 5 mm. long, pubescent and strigillose. Flowers appearing with the leaves end of April and beginning of May in 6- to 15-flowered umbel-like racemes, scarlet or bright red, with a large orange blotch on the upper lobe; pedicels strigillose, not glandular; sepals roundish ovate to oblong, 0.5 to 3 mm. long, long-ciliate, pubescent or nearly glabrous outside; corolla funnel-form, tube cylindric, rather slender, 2 to 2.5 cm. long, longer than the limb, rather abruptly dilated at the apex, covered outside with fine villose pubescence interspersed with long pilose not gland-tipped hairs, pubescent inside, lobes ovate, abruptly acuminate, 1.5 to 2 cm. long; stamens much exserted, more than twice as long as tube and slightly shorter than the style, the filaments pubescent below the middle; style 5 to 6 cm. long, finely pubescent at the lower third, usually purplish above; ovary covered with long setose hairs, not glandular. Fruit ovoid to narrow-oblong narrowed toward the apex, 2 to 3 cm. long, strigose.

This species is distributed through central Georgia from the Savannah River to the Chattahoochee River, its range extending north to Gwinnett County and to Oconee County of South Carolina, south to Screven County in the east and to Muscogee County in the west.² It is usually a low shrub and grows in open dry woods and on sand hills, often covering large stretches and is very conspicuous when in bloom on account of the brilliant color of its flowers.

¹ ADDITIONAL ILLUSTRATIONS. COLORED PLATES: Drapiez, *Herb. Amat.* I. t. 51 (1828), as *A. nudiflora*. — ? Audubon, *Birds Am.* II. t. 198 (1831); Quarto ed. II. 86 & 104 (1841), poorly colored. — BLACK FIGURES: Dippel, *Handb. Laubholz.* I. fig. 266 (1889), after Watson, *Dendr. Brit.*, as *R. calendulaceum*. — Voss, *Vilmorin's Blumengärt.* I. 588 (1894), same as preceding.

² Mohr mentions under *A. nudiflora* (*Pl. Life Ala.* 653 [1901]) a form with flame or deep orange-colored flowers from Springhill, Mobile County, which he refers to *A. nudiflora* var. *rutilans* Pursh; this may be either an undescribed color form of *R. canescens* or it may be *R. speciosum* and in this case the range of *R. speciosum* would extend to southwestern Alabama. I have not seen any specimen of this form.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Of Michaux's specimens of var. *a. flammeum* collected according to his *Journal* near Two-Sisters Ferry, Georgia, on April 27, 1787, and of his specimens of var. *β. croceum* I have photographs before me which show that the latter variety is *R. calendulaceum*, while *a. flammeum* differs chiefly in its slenderer corolla-tube distinctly longer than the limb and not conspicuously glandular-hairy and in the smaller corolla-lobes and thus agrees with the plant here called *R. speciosum*.

Rhododendron speciosum has been usually confused with *R. calendulaceum*, and particularly with the scarlet-flowered form of it, from which it may be chiefly distinguished by the slenderer corolla-tube more abruptly enlarged at the apex into the limb and longer than the limb, by the absence of glandular hairs on the corolla-tube which is covered with a short villous pubescence interspersed with pilose hairs and by the generally smaller and comparatively broader leaves. The color of the flowers is always scarlet or bright red and never varies to yellow, as it does in *R. calendulaceum* and the flowers are more numerous in each inflorescence. It shows little variation except in shape and size of sepals, which may be oblong and up to 3 mm. long or nearly obsolete, and in the shape and size of the capsules. Geographically the two species are well separated; *R. calendulaceum* is essentially an Alleghanian plant, while *R. speciosum* belongs to the Piedmont region of Georgia, but in northern Georgia the ranges of the two species touch and at Thompson's Mill, Gwinnett County, both species have been collected by H. A. Allard.

Though the species was well known in cultivation at the beginning of the last century, it later almost disappeared from the gardens until it was reintroduced in 1916 through the Arnold Arboretum, though in 1881 it was still cultivated at Kew as *Azalea coccinea major* according to a specimen before me. There is also in the herbarium of the Arnold Arboretum a specimen collected by H. Zabel in the Botanic Garden at Muenden, Hanover, from a plant received as *Azalea coccinea* from Booth's nursery at Flottbeck near Hamburg in 1872. It is certainly as handsome as *R. calendulaceum*, but probably more tender and not suited for cultivation in the open ground in the North.

Rhododendron nudiflorum, Torrey, *Fl. U. S.* 424 (1824); *Fl. N. Y.* I. 438 (1843). — Gray, *Syn. Fl.* II. 1, 41 (1878). — J. Robinson, *Fl. Essex Co. Mass.* 73 (1880). — Dame & Collins, *Fl. Middlesex Co. Mass.* 63 (1888). — Britton, *Cat. Pl. N. Jersey*, 162 (1889). — Dippel, *Handb. Laubholz.* I. 414, fig. 267 (1889). — Coulter & Watson, *Gray's Man.* ed. 6, 320 (1890). — Dudley & Thurston, *Cat. Pl. Lackawanna & Wyoming Valleys*, 38 (1892). — Beckwith & Macauley, *Pl. Monroe Co. N. Y.* 83 (in *Proc. Rochester Acad. Sci.* III) (1894), in part. — Deane in *Rhodora*, I. 94 (1899); III. 196 (1901). — Robinson & Fernald, *Gray's New Man.* 631 (1908). — Graves & others, *Cat. Flow. Pl. Conn.* 308 (1910). — Schneider, *Ill. Handb. Laubholz.* 502, fig. 329 s-t (1911).

Azalea lutea Linnaeus *Spec.* I. 150 (1753), excluding the synonyms of Colden and Gronovius.

Azalea nudiflora Linnaeus, *Spec.* ed. 2, 214 (1762), excluding the synonyms of Colden and Gronovius. — Wangenheim, *Beytr. Forstwiss.* 67 (1787). — Willdenow, *Spec.* I. pt. 2, 831 (1798). — Eaton, *Man. Bot.* ed. 2, 165 (1818). — Elliot, *Sketch Bot.* I. 241 (1821), in part. — Bigelow, *Fl. Boston*, ed. 2, 82

(1824). — De Candolle, *Prodr.* VII. 716 (1839). — Gray, *Man.* 268 (1848); ed. 5, 299 (1872). — M. A. Curtis, *Descript. Trees & Shrubs*, 98 (*Geol. Nat. Hist. Surv. N. Car.* III. Bot.) (1860). — Willis, *Cat. Pl. N. Jersey*, 33 (1874). — Kellermann, *Cat. Ohio Pl.* 129 (in *Rep. Geol. Surv. Ohio*, VII. pt. 2) (1893), in part. — Britton & Brown, *Ill. Fl.* II. 558, fig. 2743 (1897). — Clute, *Fl. Upp. Susquehanna*, 69 (1898). — Britton, *Man.* 698 (1901). — Gattinger, *Fl. Tennessee*, 131 (1901). — Porter, *Fl. Pennsylv.* 238 (1903). — Small, *Fl. S. E. U. S.* 883 (1903); in *N. Am. Fl.* XXIX. 42 (1914). — Shreve & others, *Pl. Life Maryland*, 466 (1910). — Stone in *Ann. Rep. N. J. State Mus.* 1910, 612, t. 95 (1911). — Millsbaugh, *Liv. Fl. W. Virginia*, 323 (in *W. Va. Geol. Surv.*) (1913). — Schaffner, *Cat. Ohio Vasc. Pl.* 204 (in *Ohio Biol. Surv.* I.) (1914), in part. — Taylor, *Fl. Vicin. N. Y.* 488 (in *Mem. N. Y. Bot. Gard.* V.) (1915).

Rhododendron venustum Salisbury, *Prodr.* 287 (1796), in part.

Azalea periclymenoides Michaux, *Fl. Bor.-Am.* I. 151 (1803). — Pursh, *Fl. Am. Sept.* I. 152 (1814).

Azalea periclymena Persoon, *Syn.* I. 212 (1805).

Azalea nudiflora β. *periclymenoides*, Heynhold, *Nomencl. Bot. Hort.* I. 108 (1840).

Anthodendron nudiflorum Reichenbach in Moessler, *Handb. Gewächsk.* ed. 2, I. 309 (1827).¹

An upright shrub, usually 1 to 2, rarely to 3 m. tall; young branchlets slightly pubescent and sparingly strigose or nearly glabrous, becoming dull brown or grayish brown; winter-buds brown, glabrous or slightly pubescent, with ovate abruptly acuminate finely ciliate scales; leaves elliptic to oblong or obovate to oblong-obovate, acute or abruptly acuminate, rarely obtusish, gland-tipped, cuneate at base, 3 to 8 cm. long and 1.5 to 3 cm. wide, finely pubescent on the midrib above, strigillose and sometimes sparingly pubescent on the midrib beneath, finely ciliate on the margins, bright green above and scarcely paler beneath; petioles 2 to 4 mm. long, strigillose and sometimes slightly pubescent. Flowers

¹ PRELINNEAN SYNONYMS: *Azalea scapo nudo floribus confertis terminatricibus staminibus declinatis*, Linnaeus, *Hort. Cliff.* 69 (1737). — *Chamaerhododendros Virginiana Periclymeni flore ampliori minus odorato*, Duhamel, *Traité Arb.* I. 160 (1755) and doubtfully *Cistus Virginiana Periclymeni flore ampliori et minus odorato* [sic] Plukenet, *Mant.* 49 (1700).

ADDITIONAL ILLUSTRATIONS. COLORED PLATES: Trew, *Pl. Sel. Ehret.* t. 48 (1750). — Schmidt, *Oester. Baumz.* III. t. 170 (1800), good plate of the typical form. — *Bot. Reg.* II. t. 120 (1816), as *A. nudiflora* var. *carnea*. — Loddiges, *Bot. Cab.* I. t. 51 (1818), as *A. nudiflora* var. *rubra*. — Guimpel, Otto & Hayne, *Abb. Fremd. Holzgew.* t. 110 (1825), apparently var. *glandifera* with white flowers. — Emerson, *Trees & Shrubs Mass.* ed. 2, II. 440 t. (1875), pink flowers. — *Jour. N. Y. Bot. Gard.* XIV. 79, pl. (1913). — House, *Wild Fl. N. Y.* II. 200, pl. 154 (1920). — BLACK FIGURES: Loudon, *Arb. Brit.* II. 1140, fig. 943 (1838), after *Bot. Reg.* t. 120. — André, *Pl. Terre de Bruyère*, 179, fig. 6 (1864). — *Gard.* XXIX. 550 (1886), same as Loudon's. — Mottet, *Dict. Prat. Hort.* I. 303, f. 359 (1892), after André. — Newhall, *Shrubs N. E. Am.* fig. 83 (1893). — Miller & Whiting, *Wild Flow. N. E. St.* 379 (1895). — Bailey, *Cycl. Am. Hort.* I. fig. 172, 173 (1900); *Stand. Cycl. Hort.* V. fig. 3389, 3390 (1916). — Keeler, *Our North. Shrubs*, 347 (1903). — Stevens, *Ill. Guide Flow. Pl.*, t. 114, fig. 2 (1910). — *Rep. N. Jersey St. Mus.* 1910, t. 15 (1911). — *Amer. For.* XXII. 655, fig. (1916). — HABIT FIGURES: *Gard. Mag.* V. 218, fig. (1907). — *Country Life Am.* XI. 496 (1907).

expanding in April and May just before the leaves, scentless or nearly scentless,¹ in 6- to 12-flowered umbel-like racemes; pedicels 0.5 to 1 cm. long, strigose-pilose and sometimes finely pubescent, rarely sparingly glandular; calyx-lobes semi-orbicular or ovate, rarely oblong, 0.5 to 2 mm. long, long-ciliate; corolla usually light pink or whitish with pink tube, funnel-form, tube 1.5 to 2 cm. long, rather gradually dilated above, finely pubescent and pilose or strigose-pilose outside; lobes ovate to oblong-ovate, abruptly acuminate, 1.2 to 1.6 cm. long, finely pubescent outside or glabrescent toward the apex, pilose along the middle; stamens nearly three times as long as the corolla-tube, declinate, the filaments pubescent below the middle; anthers 2 to 2.5 mm. long, usually brownish yellow; style 5 to 6 mm. long, somewhat exceeding the stamens, finely pubescent at the lower third, usually purple or purplish above; ovary usually densely covered with long setose glandless hairs, rarely sparingly setose and nearly entirely white-tomentose. Capsule oblong to narrow-oblong, narrowed upward, 1 to 2 cm. long, finely pubescent and usually strigose.

This species has the center of its distribution in the Appalachian mountains from Massachusetts to northwestern North Carolina and ascends to an altitude of 3800 feet; its range extends east to the coast from Massachusetts to southern North Carolina and west to central New York (Monroe County), central Pennsylvania, southern Ohio, eastern Kentucky and eastern Tennessee. According to Small it extends into northern Florida, but I have seen no specimens from south of North Carolina. It inhabits chiefly dry open woods and is also found along mountain streams.

Rhododendron nudiflorum in its typical form is chiefly characterized by the glabrous bright green, usually elliptic or obovate leaves pubescent only on the midrib above, strigose along the midrib below and ciliate, by the glabrous winter-buds, by the pink, scentless flowers with a cylindrical tube slightly longer than the lobes rather gradually dilated above the middle into the spreading limb and covered outside with a fine pubescence and pilose glandless hairs. The species shows, however, considerable variation and in some forms it approaches *R. roseum*. The most striking variation is that with the hairs of the corolla gland-tipped instead of glandless; this is separated below as a distinct variety. There are slighter variations in the shape of the leaves, in the pubescence of the winter-buds which are sometimes, particularly in the form with glandular corolla, finely pubescent, in the color of the flowers which varies from almost white to deep carmine. The sepals are occasionally well developed as in specimens of var. *glandifera* from Somerset, Massachusetts, collected by J. G. Jack, May 21, 1903, where they are up to 5 mm. long. In one specimen, Harbison's No. 35 (1919) from Wilmington, North Carolina, the filaments are pubescent beyond the middle and the ovary is only sparingly setose.

Rhododendron nudiflorum was first discovered probably by Banister and mentioned by Plukenet as "Cistus virginiana Periclymeni flore ampliori et minus odorato." It was introduced into England together with *R. viscosum* in the first half of the 18th century, probably between 1725 and 1730 by Peter Collinson, who had received it from John Bartram, for in a letter to Dr. Colden dated Ridge Way House, Feb. 25, 1764, speaking of the plants in his garden he says, "Regard the variety of trees and shrubs in this plantation, as Mountain Magnolia, Sarsifax, Rhododendrons, Kalmias and Azaleas, etc. etc.; all are the bounty of my curious botanic friend, J. Bartram of Philadelphia."² According to Aiton it was

¹ Clute (*Fl. Upp. Susquehanna*, 69), however, calls the flowers very fragrant.

² Gray, Selections from the Scientific Correspondence of Cadwallader Colden with Gronovius, Linnaeus, Collinson, p. 48 (1843); reprinted from *Ann. Jour. Sci. Arts*, LXIV.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Rhododendron nudiflorum var. **glandiferum** Rehder, comb. nov.¹

Azalea nudiflora glandifera Porter in *Bull. Torr. Bot. Club*, XXVII. 508 (1900); *Fl. Pennsylv.* 238 (1903). — Keller & Brown, *Handb. Fl. Phila.* 246 (1905). — Stone in *Ann. Rep. N. Jersey State Mus.* 1910, 613 (1911).

This variety differs from the type in having the pedicels and the outside of the corolla-tube furnished with more or less numerous gland-tipped hairs of usually nearly equal length, and the color of the corolla is usually of a deeper pink or carmine color. The variety was based originally on specimens collected in eastern Pennsylvania, but it is found frequently within the range of the type from Massachusetts to northwestern South Carolina and often grows together with the typical form. It seems, however, nearly absent from the southeastern part of the range, that is from the coastal plain from New Jersey to North Carolina and more common West, for the plants I have seen from the extreme Western localities, western Tennessee and southern Ohio, represent this variety though from the last named locality (Lawrence County) and from western New York (Monroe County) I have seen specimens of the typical form.

A form with double flowers is mentioned by Pursh (*Fl. Am. Sept.* 152 [1814]) under his *A. periclymenoides*, but he does not state where he saw it. In 1826 Sweet enumerated a double white form (*Azalea nudiflora* β . *albo-plena* Sweet, *Hort. Brit.* 265 [1826]), name only. — *Rhododendron nudiflorum* β . *albo-plenum* Sweet, *Hort. Brit.* ed. 2, 343 (1830) and a semi-double white form (*Azalea nudiflora* $\nu\nu$. *semiduplex* Sweet, *Hort. Brit.* 265 [1826]. — *Rhododendron nudiflorum* $\nu\nu$. *semiduplex* Sweet, *l. c.* ed. 2, 344 [1830]), also a double purple form (*Azalea nudiflora* $\delta\delta$. *purpureo-plena* Sweet, *l. c.* 265 [1826]. — *Rhododendron nudiflorum* $\delta\delta$. *purpureo-plenum* Sweet, *l. c.* ed. 2, 344 [1830]) and a double red form (*Azalea nudiflora* $\kappa\kappa$. *rubro-plena* Sweet, *l. c.* 265 [1826]. — *Rhododendron nudiflorum* $\kappa\kappa$. *rubro-plenum* Sweet, *l. c.* ed. 2, 344 [1830]). Probably *Azalea nudiflora flore pleno* Loddiges Cat. 1836 (ex Loudon, *Arb. Brit.* II. 1142 [1838]) is the same as the double white form of Sweet. Two different forms *Azalea periclymenoides* "alba pleno" and "rubra pleno" were offered in 1831 by R. Carr (*Cat. Am. Trees Bartram Bot. Gard.* 64). All the double-flowered forms in cultivation which I have seen seem to be of hybrid origin, but from Kew Gardens I have seen a specimen collected by G. Nicholson in 1880 as *Azalea* "Double Blush" which is probably a double-flowered *R. nudiflorum*; it has the leaves perfectly glabrous beneath except the strigose midrib and the corolla is pilose and villose outside but not glandular.

Several other varieties and forms have been described or mentioned by Aiton, Loddiges and others, but as the descriptions are inadequate and I have not seen the type specimen or other reliable material of them, I am in doubt whether they belong here or to *R. roseum* or to *R. canescens*; they will therefore be enumerated under the doubtful names.

Rhododendron roseum Rehder, comb. nov.

Azalea rosea Loiseleur-Deslongchamps in Duhamel, *Traité Arb. Arbust.* ed. 2, V. 224, t. 64 (1812). — ? Michaux, *Journal*, ed. C. S. Sargent, 111 (in *Proc. Am. Philos. Soc.* XXVI. (1889)).

¹ *Azalea nudiflora rubra* Loddiges, *Bot. Cab.* I. t. 51 (1817). — De Candolle, *Prodr.* VII. 717 (1839), as var. — *Rhododendron nudiflorum* var. *rubrum* Sweet, *Hort. Brit.* ed. 2, 343 (1830); G. Don, *Gen. Syst.* III. 847 (1834) is probably identical with this variety, judging from Loddiges' figure, but as long as one cannot be positively sure I prefer to keep Porter's name.

Azalea canescens Pursh, *Fl. Am. Sept.* I. 152 (1814), in part, as to the specimen from Virginia,¹ not Michaux. — Eaton, *Man. Bot.* 165 (1818). — Britton & Brown, *Ill. Fl.* II. fig. 2744 (1897). — Clute, *Fl. Upp. Susquehanna*, 70 (1898). — Britton, *Man.* 698 (1901). — Porter, *Fl. Pennsylv.* 238 (1903). — Millspaugh, *Liv. Fl. W. Virginia*, 322 (in *W. Va. Geol. Surv.*) (1913). — House, *Wild Fl. N. Y.* II. 201 (1920), excl. figure.

Azalea nudiflora $\epsilon\epsilon$. *rosea* Sweet, *Hort. Brit.* 265 (1826), name only. — Loddiges apud Loudon, *Arb. Brit.* II, 1142, fig. 945 (1838), name and figure only.

?*Azalea calendulacea* δ . *cupraea* Sweet, *Hort. Brit.* 265 (1826), name only. — Loddiges, *Bot. Cab.* XIV. t. 1394 (1828).

?*Rhododendron calendulaceum* δ . *cupreum* Sweet, *Hort. Brit.* ed. 2, 343 (1830), name only.

Rhododendron nudiflorum $\epsilon\epsilon$. *roseum* Sweet, *Hort. Brit.* ed. 2, 344 (1830).

Rhododendron nudiflorum Darlington, *Fl. Cestricea*, ed. 2, 262 (1837), not Torrey.² — Oakes, *Cat. Vermont Pl.* 190 (in Thompson, *Hist. Vermont*) (1842); reprint p. 18 (1842). — Dudley, *Cayuga Fl.* 59 (in *Bull. Cornell Univ.* II.) (1886). — Perkins, *Cat. Fl. Vermont*, 33 (1888). — Beckwith & Macauley, *Pl. Monroe Co. N. Y.* 83 (in *Proc. Rochester Acad. Sci.* III.) (1894), in part.

Azalea nudiflora Darlington, *Fl. Cestricea*, 26 (1826).² — Wright & Hall, *Cat. Pl. Troy*, 8 (1836). — Clinton, *Prelim. List Pl. Buffalo*, 7 (1864). — Payne, *Cat. Pl. Oneida Co.* 51 (in *Rep. Regents Univ. N. Y.*) (1865). — Patterson, *Cat. Pl. Illinois*, 25 (1876). — D. F. Day, *Pl. Buffalo*, 115 (in *Bull. Buffalo Nat. Sci.* IV.) (1882). — Meehan in *Meehan's Monthly*, II. 17, t. 2 (1892).³ — Kellerman, *Cat. Ohio Pl.* 129 (in *Rep. Geol. Surv.* I.) (1914), in part. — Schaffner, *Cat. Ohio Vasc. Pl.* (in *Ohio Biol. Surv.* I.) (1914), in part.

Rhododendron canescens Porter in *Bull. Torr. Bot. Club* XVI. 220 (1889), not Sweet. — Dudley & Thurston, *Cat. Pl. Lackawanna & Wyoming Valleys*, 38 (1892). — Deane in *Rhodora*, III. 196 (1901). — Fernald & Robinson, *Gray's New Man.* 631 (1908). — Jackson, *Cat. Pl. Worcester Co. Mass.* 76 (1909). — Graves & others, *Cat. Flow. Pl. Conn.* 308 (1910). — Stone, *List Pl. Frankl. Hampshire, Hampden Cos. Mass.* 48 (1913). — Rehder in *Bailey, Stand. Cycl. Hort.* V. 2942 (1916). — Twining, *Fl. N. E. Pennsylv.* 56 (1917).

Azalea prinophylla Small in *N. Am. Fl.* XXIX. 42 (1914). — Taylor, *Fl. Vicin. N. Y.* 488 (in *Mem. N. T. Bot. Gard.* V.) (1915).

Rhododendron prinophyllum Millais, *Rhodod.* 229 (1917). — Millais & Williams in *Rhodod. Soc. Notes*, I. 125 (1918).⁴

A shrub 1 to 3, rarely to 5 m. tall, with irregularly whorled branches; the young branchlets finely pubescent, and usually sparingly strigose, becoming grayish brown or light brown the second year; buds grayish pubescent with numerous,

¹ The phrase in the description "staminibus vix exertis" also refers to *R. roseum* rather than *R. canescens*.

² As the leaves are described as pubescent beneath, the species must be *R. roseum* rather than *R. nudiflorum*.

³ As Meehan describes the leaves as pubescent beneath and the corolla as glandular, the plant figured must be *R. roseum* rather than *R. nudiflorum*.

⁴ ADDITIONAL ILLUSTRATIONS. COLORED PLATES: Goodale, *Wild Flow. Am.* t. 36 (1882). — *Meehan's Monthly*, II. t. 2 (1892). — BLACK FIGURES: Loudon, *Arb. Brit.* II. fig. 945 (1838), as *R. nudiflorum* var. *rosea*, after *Nowv. Duhamel.* — Keeler, *Our North. Shrubs*, 349, fig. (1903), as *Azalea canescens*. — Stevens, *Ill. Guide Flow. Pl.* t. 114, fig. 1 (1910), as *A. canescens*.

generally ovate, obtusish and mucronulate scales. Leaves elliptic or obovate to obovate-oblong, acute or short-acuminate, gland-tipped, cuneate at the base, 3 to 7 cm. long and 1.2 to 3.3 cm. broad, sparingly pubescent above, densely grayish villose beneath or sometimes thinly villose rarely nearly glabrous except along the midrib, sparingly strigillose on the midrib, ciliolate, thinnish at maturity, usually more or less bluish green or dull green; petioles 2 to 5 mm. long, soft-pubescent and sparingly strigose. Flowers expanding in May with the leaves in usually 5- to 9-flowered umbel-like racemes, fragrant; pedicels 0.5 to 1.5 cm. long, finely villose and glandular-setose; calyx-lobes semiorbicular to ovate, unequal, scarcely exceeding 1 mm., finely pubescent, glandular-ciliate; corolla funnel-form with oblique limb, usually bright pink, rarely whitish, the tube 1.5 to 2 cm. long, cylindrical, rather gradually dilated toward the apex, covered with a thin villose tomentum interspersed with numerous gland-tipped hairs of unequal length, pubescent inside, the lobes about as long or slightly shorter than tube, ovate, abruptly pointed; stamens little more than twice as long as tube, filaments pubescent below the middle, the anthers about 2 mm. long, ochraceous; style exceeding the stamens, 4 to 5 cm. long, sparingly or sometimes rather densely pubescent below, purple above; ovary finely covered with a villous pubescence overlaid with numerous appressed white, setose, partly gland-tipped hairs. Capsule oblong, narrowed toward the apex, 1.5 to 2 cm. long, sparingly puberulous and setosely glandular.

Rhododendron roseum is distributed from southwestern New Hampshire, central Vermont and southwestern Quebec¹ through western New York, eastern Ohio, southern Illinois to southeastern Missouri, and through western and central Massachusetts, northwestern Connecticut, northern Rhode Island, New York, Pennsylvania, and northern New Jersey to southwestern Virginia and western Tennessee. Its range lies chiefly west of the Appalachian Mountains, except in the north, and does not extend into the coastal plain.

This species is closely related to *R. nudiflorum* and particularly to its variety *glandiferum* and I was for some time inclined to refer it as a variety to the latter species, as it seems to be connected with it by intermediate forms. It differs, however, in several important characters from typical *R. nudiflorum*, as in the pubescent winter-buds, the more or less densely villous branchlets, the pubescent bluish-green leaves, in the shorter stamens and in the glandular corolla with larger and broader lobes and a wider tube, and as the two are easily distinguished in their typical forms, I think it better to keep them as distinct species. Glabrescent forms with the leaves pubescent only along the midrib might be confused with *R. nudiflorum*, but can be distinguished from it by the flowers, the pubescent winter-buds and the finely villous branchlets. *Rhododendron roseum* has also been confused with *R. canescens* Don, with which it agrees in the pubescence of the leaves, but differs in the shorter, gradually dilated pilose-glandular corolla-tube, in the shorter stamens and in the thinner, not reticulate leaves.

This species was first mentioned in 1787 as a variety of *Azalea nudiflora* "with broad rough leaves" by Wangenheim (*Beytr. Forstwiss.* 68), who observed it in the state of New York. When it was introduced into cultivation cannot be stated with certainty. It may be one of the varieties mentioned by Aiton in 1789, and without doubt Pursh's specimen from Cacapoon Springs near Winchester, West Virginia, referred by him to *R. canescens* belongs here. According to a note by Loudon (*Arb. Brit.* II. 1143) this has been found in Virginia "on the mountains of the Cacapoon Springs, near Winchester, where it forms a shrub growing 3 feet or 4 feet high and producing its rose-coloured flowers in May and June. Introduced in 1812 and

¹ Near Hull, Quebec, June 5, 1916, *John Dunbar* (Herb. Park Dept., Rochester, N. Y.)

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

lose, short. Flowers fragrant, appearing with the leaves, in April and May, in usually 6- to 10-flowered umbel-like clusters; pedicels 0.5 to 1 cm. long, sparingly villose and rather densely hirsute, glandless or glandular; sepals unequal, roundish to ovate, 0.5 to 1.5 mm. long, densely setose, glandless; corolla funnel-form, white, tube cylindric, gradually dilated at apex, 2 to 2.8 cm. long, rarely slightly shorter, outside sparingly and more or less glandular-pilose or sometimes hirsute without glands, lobes ovate or sometimes ovate-oblong, acute, 1.5 to 1.8 cm. long, distinctly shorter than the tube; stamens twice as long as tube, scarcely declinate, the filaments villose below the middle, the anthers yellowish, 2 to 3 mm. long; style scarcely exceeding the longer stamens, 5 to 6 cm. long, finely villose toward the base, whitish; ovary densely setose-strigose, glandular or nearly glandless. Capsule (immature) cylindric-oblong, about 1.5 cm. long, sparingly and finely villose and glandular-setulose or nearly glandless.

ALABAMA. Cullman County: Pine woods, May 21, 1902, Biltmore Herb. H/6292¹ (Nat. Herb. No 980,507, type); on rocky slope in woods of Oak, Hickory and Pine, May 21, 1902, Biltmore Herb. H/6293 (Nat. Herb. No. 980,506); dry rocky hillside, May 19, 1902, Biltmore Herb. H/6258 (Nat. Herb. No. 980,524); dry woods, May 19, 1902, Biltmore Herb. H/6267, 6268 (Nat. Herb. Nos. 980,528, 980,511); without special locality, May 19, 1902, Biltmore Herb. H/6259, 6262-6265 (Nat. Herb. Nos. 980,525, 980,510, 980,523, 980,509, 980,527); dry rocky open woods on the high tablelands, May 2, 1883, *C. Mohr* (as *R. (Azalea) nudiflora alba*; "flowers snowy white, highly fragrant"). Marshall County: Albertville, May 8, 1901, Biltmore Herb. H/4290, 4292, 4293, 4295, 4297 (Nat. Herb. 969,632, 969,628, 969,682, 969,634, 969,658); dry woods, August 27, Biltmore Herb. H/2452 (Nat. Herb. 969,617); October 12, 1900, Biltmore Herb. H/3321 (Nat. Herb. 969,649). Henry County: Abbeville, April 15, 1902, Biltmore Herb. H/6112 (Nat. Herb. 980,513).

As it appears from these specimens this species is known to me only from two counties in northern and from one county in southeastern Alabama, but probably it is more widely distributed through the northern and central part of the state; it inhabits dry open woods and rocky hillsides and thus agrees in its habitat with the northern *R. roseum*, while *R. canescens*, the other common Alabama species, prefers damp woods and swamps.

This new species is closely related to *R. nudiflorum* Torr., *R. roseum* Rehd. and *R. canescens* G. Don. From the first species it is easily distinguished by the much longer and slenderer corolla-tube, the pure white fragrant flowers, less long-exserted scarcely declinate stamens and the pubescent leaves; from *R. roseum* it differs in the glabrous winter-buds, the longer and slenderer corolla-tube, the glandless setae of the sepals, in the white color of the flowers and in the glaucous under side of the leaves; from *R. canescens* it is distinguished by the glabrous winter-buds, the much less villose and more or less pilose corolla-tube, less slender and more gradually dilated at the apex, by the less exserted stamens, the larger limb, the densely strigose ovary, the less villose and more setose capsules,

¹ "H" stands for T. G. Harbison who collected for the Biltmore Herbarium.

and the generally broader leaves. *Rhododendron alabamense* shows some variation chiefly in the size and pubescence of its flowers. The type specimen has large flowers with a glandular-pilose tube, while in Nos. 6258 and 6259 the tube has longer and quite glandless hairs; the specimens from Albertville have generally smaller flowers, the tube sometimes scarcely 2 cm. long and the lobes longer and narrower, sometimes nearly oblong and acuminate, the leaves are also smaller and less glaucescent and often less pubescent beneath. The Abbeville specimen from southeastern Alabama is again much like the type in its flowers, but the leaves are less densely pubescent and less glaucescent beneath. In the National Herbarium is a sheet (No. 770,587) labeled "*R. viscosum* Torr., borders of stream in thickets, Pointclear (not quite legible), C. Mohr, June 16, 1889," which contains a specimen of *R. arborescens* and two specimens which suggest a glabrous form of *R. alabamense*, but as apparently the locality and the date do not belong to these two specimens, I consider them for the present as doubtful.

Rhododendron alabamense was apparently first observed in 1883 by Dr. C. Mohr, who recognized it as distinct from the more widely distributed *R. canescens* which he included under his *A. nudiflora*; this probably induced him to identify this particular form with *A. nudiflora* var. *alba* Aiton. The species has not yet been introduced into cultivation; it may be hardy North, and for southern gardens it would be superior to *R. canescens* on account of its very fragrant larger flowers, which are "snowy white" according to Dr. Mohr.

***Rhododendron canescens* Sweet, Hort. Brit. ed. 2, 343 (1830). — G. Don, Gen. Syst. III, 848 (1834), excl. the reference to plant of Winchester. — Loudon, Arb. Brit. II, 1143 (1838), in part. — Millais, Rhodod. 136 (1917).**

Azalea lutea Linnaeus, *Spec.* 150 (1753), as to the synonym of Gronovius only.

Azalea nudiflora Linnaeus, *Spec. ed.* 2, 214 (1762), as to the synonym of Gronovius only. — Walter, *Fl. Car.* 97 (1788). — Elliott, *Sketch Bot.* I. 240 (1821), in part. — Mohr, *Pl. Life Ala.* 653 (1901).

? *Azalea nudiflora* ε. *bicolor* Aiton, *Hort. Kew.* I. 203 (1789).

Azalea canescens Michaux, *Fl. Bor.-Am.* I. 150 (1803). — Pursh, *Fl. Am. Sept.* 152 (1814), in part. — Elliott, *Sketch Bot.* I. 239 (1821). — De Candolle, *Prodr.* VIII. 717 (1838). — Small, *Fl. S. E. U. S.* 883 (1903); *Shrubs Fla.* 93 (1913); in *N. Am. Fl.* XXIX. 42 (1914).

Azalea bicolor Pursh, *Fl. Am. Sept.* I. 153 (1814). — Elliott, *Sketch Bot.* I. 239 (1821). — De Candolle, *Prod.* VII. 717 (1839).

Rhododendron bicolor Sweet, *Hort. Brit. ed.* 2, 344 (1830). — G. Don, *Gen. Syst.* III. 847 (1834). — Loudon, *Arb. Brit.* II. 1142 (1838).

Rhododendron nudiflorum Chapman, *Fl. S. U. S.* 265 (1865), not Torrey.

Azalea nudiflora var. *canescens* Rehder in Bailey, *Cycl. Am. Hort.* I. 121 (1900), in part.¹

A shrub to 4 or 5 m. tall with irregularly whorled upright branches; young branchlets finely villous-pubescent and usually sparingly strigose; winter-buds with broadly ovate, acuminate and densely grayish pubescent scales. Leaves usually

¹ PRELINNEAN SYNONYM: *Cistus Virginiana flore et odore Periclymeni*, Catesby, *Nat. Hist. Car.* I. 57, t. 57 (1731), not Banister apud Plukenet.

ADDITIONAL ILLUSTRATIONS. COLORED PLATES: Abbot, *Nat. Hist. Lepidopt. Georgia*, I. t. 27 (1797). — Audubon, *Birds of Am.* IV. t. 398 (1835–38).

oblong-obovate to oblanceolate or oblong, rarely elliptic or obovate, acute, rarely obtusish, gland-tipped, cuneate at base, 4 to 9 cm. long and 1.5 to 3.5 cm. broad, setosely ciliate, sparingly pubescent above or glabrescent except at the finely villose midrib, densely pubescent or grayish tomentulose beneath particularly on the veins and midrib, usually sparingly strigose on the midrib toward the base, of firm texture and slightly reticulate beneath at maturity; petioles 2 to 7 mm. long, finely pubescent and sparingly strigose. Flowers expanding before or with the leaves in April and May, in 6- to 15-flowered clusters, slightly fragrant; pedicels 0.5 to 1 cm. long, villose-pubescent and hirsute, sometimes stipitate-glandular; calyx-lobes, unequal, semi-orbicular to ovate, scarcely exceeding 1 mm., ciliate or sometimes glandular-ciliate; corolla-tube cylindrical, abruptly dilated at apex, 1.5 to 2.3 cm. long, distinctly longer, sometimes nearly twice as long as lobes, densely and finely villose outside and stipitate-glandular, rarely somewhat pilose, pubescent inside; lobes ovate, acute or obtusish, 1.2 to 1.5 cm. long; stamens much exserted, nearly 3 times as long as tube, pubescent below the middle, anthers 1.5 to 2 mm. long, ochraceous; style 4 to 6 cm. long, finely villose toward the base, slightly longer or about as long as stamens; ovary densely white pubescent with appressed silky hairs and sparingly setulose, without glands. Capsule cylindrical-oblong, narrowed upward, slender, 1.5 to 2 cm. long, pubescent and sparingly setose.

This species is chiefly a plant of the southern Atlantic and the Gulf coastal plain, but extends into the Piedmont region of North Carolina, Georgia and Tennessee; it ranges from southwestern Tennessee and southern central North Carolina to eastern South Carolina and northeastern Florida to extreme southeastern Texas, central Louisiana and northeastern Alabama. It usually grows in woods on moist sandy soil, often along streams, rarely in swampy soil.

Rhododendron canescens is chiefly characterized by the more or less pubescent leaves firm at maturity and slightly reticulate beneath, by the slender corolla-tube abruptly enlarged into a comparatively small limb and clothed outside with a fine dense villose tomentum interspersed with rather short and small stipitate glands, by the long exserted stamens being nearly three times as long as the tube and by the grayish pubescent eglandular and not strongly setose ovary and capsule. The color of the flowers is usually pink or perhaps more often the tube only is pink and the limb white or only slightly pinkish; a form with nearly purple flowers differing further from the type in the comparatively short and wide corolla-tube has been collected by T. G. Harbison at Valleyhead, Alabama (No. 530; May 2, 1911). There is considerable variation in the pubescence of the leaves; in one extreme they are nearly white tomentose beneath, while in the other extreme they are nearly glabrous. The glabrous or glabrescent form has been confused with *R. nudiflorum*, but it is easily distinguished from that species by the shape and pubescence of the corolla and by the pubescent winter-buds.

Rhododendron canescens was discovered by Catesby, who published a characteristic and fairly good figure of it in 1731. In 1797 an excellent colored plate of it, drawn by J. Abbot, was published as *A. nudiflora* in his work on the Lepidopterous insects of Georgia. It was also collected by Michaux in South Carolina between 1784 and 1796, and first recognized by him as a distinct species and described in his *Flora* as *R. canescens*. Pursh in 1814 confused Michaux's *A. canescens* with *R. roseum*; he identified a plant collected near Winchester, West Virginia, which apparently was *R. roseum*, with Michaux's *A. canescens* and redescribed the true *A. canescens* under the name *A. bicolor*.

It was probably introduced into cultivation about the middle of the 18th century and it may be one of the varieties enumerated by Aiton in 1789 and possibly his var. *bicolor*, cited by Pursh as a synonym of his *A. bicolor* which is identical with *R. canescens*, but it apparently soon disappeared again from the gardens.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Rhododendron canescens var. **candidum** Rehder, var. nov.

Azalea candida Small in *Bull. Torr. Bot. Club* XXVIII. 360 (1901); *Fl. S. E. U. S.* 883 (1903); *Shrubs of Fla.* 93 (1913); in *N. Am. Fl.* XXIX. 43 (1914).
Rhododendron candidum Rehder in *Bailey, Stand. Cycl. Hort.* V. 2945 (1916); VI. 3574 (1917). — Millais, *Rhodod.* 136 (1917).

This variety differs from the typical form in the glaucous or glaucescent, densely pubescent under side of the leaves and in the mostly denser pubescence of the branchlets usually interspersed with numerous short gland-tipped setae. The white or whitish color of the under side is not caused by the dense tomentum, as one might suppose, but by the glaucous or glaucescent epidermis, which is coated with a waxy exudation as in glaucous forms of *R. viscosum*.

GEORGIA. Decatur County: Climax, Nov. 6, 1917, *T. G. Harbison* (No. 14). Brooks County: Quitman, April 2, 1918, *T. G. Harbison* (No. 3). Lowndes County: near Valdosta along the Withlacoochee River, June 6–12, 1895, *J. K. Small* (type); in low sandy soil along the Withlacoochee River, April 2, 1918, *T. G. Harbison* (Nos. 1, 2); near Ousley, Oct. 21, 1910, *T. G. Harbison* (Nos. 247, 250). Charlton County: Folkston, July 16, 1918, *T. G. Harbison* (No. 76). Tattnall County: rock outcrop near Choopee River, June 24, 1903, *R. M. Harper* (No. 1858).

FLORIDA. Hamilton County: White Springs, May 11, 1901, *H. H. Blume*.

This variety is restricted to southern Georgia and northern Florida and I have seen no specimen farther north or south than about twenty miles from the border line except a specimen from Tattnal County about one hundred miles north of Florida, but that specimen is not a typical var. *candidum*, as it lacks the glandular pubescence of the branchlets, and the under side of the leaves though densely villous or almost whitish tomentose, is not very glaucous. Besides the character already mentioned I can find no difference between this variety and *R. canescens* and I therefore prefer to consider it only a variety of that species. I do not understand why Small places it among the species with the flowers appearing after the leaves; the flowering specimens before me, collected April 2, differ in no way as regards the development of leaves from typical *R. canescens*.

The variety seems to have been first observed and collected in 1895 by Dr. Small on the Withlacoochee River. It is doubtful whether it is in cultivation; the plant introduced under this name a few years ago raised from seed collected near Folkston, Georgia, is possibly typical *R. canescens*.

Rhododendron austrinum Rehder in *Bailey, Stand. Cycl. Hort.* VI. 3571, 3574 (1917). — Millais, *Rhodod.* 125 (1917).

Rhododendron nudiflorum var. *luteum* Curtiss in herb. sub No. 1718.
Azalea austrina Small, *Fl. S. E. U. S.* ed. 2, 1356 (1913); *Shrubs Fla.* 94 (1913); in *N. Am. Fl.* XXIX. 42 (1914).

Slender branched shrub to 3 m. tall; young branchlets covered with a short soft pubescence interspersed with usually copious gland-tipped hairs and sparingly strigose, particularly toward the apex, becoming red-brown in autumn and grayish brown the second year; floral winter-buds with ovate scales, rounded to

acutish and acuminate at the apex and densely grayish pubescent. Leaves elliptic to obovate or oblong-obovate, 3 to 9 cm. long and 1.5 to 3.5 cm. broad, acute or obtusish and gland-tipped at apex, cuneate at base, setosely ciliate, finely pubescent on both sides, more densely beneath, particularly on the veins, rarely glabrescent above at maturity; petioles 3 to 8 mm. long, pubescent and glandular and usually sparingly strigose. Flowers appearing shortly before or with the leaves, end of March or in April in 8- to 15-flowered umbel-like racemes; pedicels 0.5 to 1 cm. long, pubescent and glandular-setose; sepals broadly ovate to oblong, unequal, 1 to 2.5 mm. long, glandular-ciliate, pubescent outside; corolla yellow and orange, the tube usually more or less purplish or with 5 longitudinal purplish stripes, slightly fragrant, the tube cylindric, abruptly dilated at the apex, 1.5 to 2 cm. long, finely pubescent and stipitate-glandular, the lobes ovate 1 to 1.5 cm. long and 0.6 to 1 cm. broad, short-acuminate with recurved point, finely pubescent outside; stamens nearly three times as long as tube, the filaments pubescent below the middle, the anthers ochraceous, 2 to 3 mm. long; style slightly exceeding the stamens, 5 to 6 cm. long, short-pilose near the base; ovary covered with long whitish strigose, partly gland-tipped hairs. Capsule oblong-cylindric, slightly narrowed toward the apex, 1.5 to 2.5 cm. long, covered with a thin fine pubescence interspersed with long strigose partly gland-tipped hairs.

This species has a very restricted distribution; it has been found only in Gadsden County, in northern Florida, near River Junction and Chattahoochee, close to the Georgia state line and about ten miles farther south at Aspalaga near Rock Bluff, Liberty County (Biltmore Herb. 6088). It was apparently first discovered by Dr. Chapman some time before 1865, as he refers to it in his *Flora* under *R. nudiflorum* as a color variation of that species; there is also a specimen collected by him in the National Herbarium without date and locality with the note "flowers orange." The species was again collected in 1880 by A. W. Curtiss and distributed as *R. nudiflorum* var. *luteum*, but that name was never published and this *Azalea* was not recognized as distinct until described as a species by Dr. Small in 1913.

Rhododendron austrinum is so closely related to *R. canescens* that one might be induced to consider it only a color form of that species, but aside from the fact that the yellow color is a disconnected deviation from the pink color tones of *R. canescens*, *R. austrinum* differs further in the glandular ovary and capsule, entirely destitute of glands in *R. canescens*, and in the copious glandular pubescence of the calyx and pedicels which are usually, though not always, glandless in *R. canescens*. In their general appearance, however, I could see no difference except the color between the two species, when I saw them growing side by side in Mr. H. H. Hume's garden in Jacksonville, who had brought the plants of both species from River Junction the year before.

The species was first introduced into cultivation through the Arnold Arboretum, where plants were raised from seed collected in 1914 by Mr. T. G. Harbison near River Junction. It cannot be expected to thrive in northern gardens, but for southern gardens it may be desirable on account of its yellow flowers.

***Rhododendron atlanticum* Rehder, comb. nov.**

Azalea nudiflora M. A. Curtis in *Boston Jour. Nat. Hist.* I. 100 (*Enum. Pl. Wilmington*) (1834), not Linnaeus.¹—Croom, *Cat. Pl. Newbern*, 21 (1837).—Wood & McCarthy, *Wilmington Fl.* 34 (1886).

¹ I have seen in the herbarium of the Academy of Natural Science at Philadelphia a specimen collected by Curtis at Wilmington and named by him *A. nudiflora*, which proves to be *R. atlanticum*. The true *R. nudiflorum* has not been found near Wilmington.

Azalea atlantica Ashe in *Bull. Charleston Mus.* XIII. 26 (1917). — Coker in *Jour. Elisha Mitchell Soc.* XXXVI. 97, pl. 1, 7 (1920).

Low stoloniferous shrub rarely more than 0.5 m. tall, with slender upright usually sparingly branched or simple stems; young branchlets sparingly strigillose and sometimes glandular hirsute, otherwise glabrous, occasionally slightly glaucescent, reddish brown at the end of the season, becoming grayish brown and glabrous the second year; floral winter-buds with ovate, mucronulate, light brown and white-ciliate, glabrous or slightly silky-pubescent scales. Leaves obovate to oblong-obovate, rarely elliptic, rounded or acutish and gland-tipped at apex, cuneate at base, 3 to 6 cm. long and 1.5 to 2 cm. broad, setosely ciliate or minutely serrulate-ciliate, glabrous above except the villose midrib and bright green or bluish green, glabrous and sparingly strigillose and sometimes pubescent on the midrib beneath, rarely strigillose on the surface, and glaucescent or bright green, chartaceous at maturity, the secondary veins scarcely prominent beneath and the veinlets obscure; petioles 1 to 5 mm. long, sparingly strigillose. Flowers appearing before or with the leaves in April or beginning of May in clusters of 4 to 10; pedicels about 1 cm. long, hirsute or glandular-hirsute, otherwise glabrous, rarely minutely and slightly villose; sepals broadly ovate, small or occasionally oblong and 2 to 4 mm. long, glandular-ciliate or sometimes eglandular; corolla funnel-form, white, usually flushed pink or purple outside, very fragrant, the tube cylindric, rather gradually dilated at apex, 2 to 2.5 cm. or rarely to nearly 3 cm. long, outside with numerous short gland-tipped hairs, otherwise glabrous or only slightly villose, glabrous or nearly so inside, the lobes ovate to ovate-oblong, acute, 1.3 to 2 cm. long, stipitate-glandular along the middle; stamens exerted, about twice as long as the tube, the filaments hairy below the middle, the anthers 2 to 3 mm. long; style exceeding the stamens, 4.5 to 5.5 cm. long, pubescent below the middle, usually purplish above; ovary with setose, gland-tipped or glandless hairs. Capsule ovoid-oblong, 1.5 to 2 cm. long, setose, the setae glandless or gland-tipped.

DELAWARE. Kent County: Choptank Mills, May 29–30, 1904, *Witmer Stone* (Herb. Acad. Nat. Sci. Phila.).

DISTRICT OF COLUMBIA. Washington, Takoma Park, May 11, 1900, *T. A. Williams*.

VIRGINIA. Princess Anne County: May 9, 1898, *Th. H. Kearney* (No. 1062; Nat. Herb. No. 355,802). Nansemond County: near Suffolk, April 30, 1898, *Fr. Coville & Th. H. Kearney* (No. 49; Nat. Herb. No. 355,681).

NORTH CAROLINA. Johnston County: Selma, April 23, 1918, and November 19, 1919, *T. G. Harbison* (Nos. 1–8, 10). Craven County: Newbern, April 7 and 18, 1918, April 20 and Nov. 20, 1919, *T. G. Harbison* (Nos. 7, 14, 17, 23, 30, 101, 102, 115, 116, 117, 121, 131, 132). New Hanover County: Wilmington, April 2, 1880, *C. S. Sargent*; April 14 and 15, 1911, *C. S. Williamson*; April 18 and 19, 1919, *T. G. Harbison* (Nos. 36–41, 47, 48), without date, *M. A. Curtis*. Eastern N. Carolina, without precise locality, April, 1888, *G. McCarthy* (Nat. Herb. 42,633).

SOUTH CAROLINA. Florence County: Ebenezer, wet Pine barren,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

by Mr. Ashe, show well the purplish pink color of the flowers; the corolla tube is slightly shorter and wider than in the typical *R. atlanticum* and is furnished outside with numerous gland-tipped stiff hairs, which are more numerous in one of the specimens sent and less numerous in the other, but in the latter the tube is also thinly covered with a fine villose pubescence, which in typical *R. atlanticum* is absent or nearly absent, though occasionally, as in Harbison's No. 14 from Georgetown, South Carolina, the type locality of the species, and in his Nos. 38 and 41 from Wilmington, North Carolina, this villose pubescence is present. To f. *neglectum* I also refer Harbison's No. 14 from Newbern, North Carolina, collected April 17, 1918, though this differs in its narrow-oblong calyx-lobes, up to 5 mm. long, and possibly also his Nos. 7, 101, 102 from the same locality, Nos. 37, 40 and 47 from Wilmington, and his Nos. 3 and 5 from Selma, North Carolina, may belong to this form, if we consider the color its chief distinguishing character.

Rhododendron atlanticum var. luteo-album Rehder, comb. nov.

Azalea atlantica var. *luteo-alba* Coker in *Jour. Elisha Mitchell Sci. Soc.* XXXVI. 98, pl. 1, fig. (at right) (1920).

This variety differs from the type in the dull bluish green leaves finely pubescent on both surfaces and strigillose above, in the finely villose petioles and in the stipitate glandular young branchlets; the flowers white, when open, are sometimes yellowish in bud without any trace of pink.

SOUTH CAROLINA. Darlington County: Ditch bank southeast of Hartsville plantation of J. L. Coker & Co., spring of 1918, *J. L. Coker, Jr.* (co-type); flat woods south of Hartsville plantation, April 29, 1915, *W. C. Coker*; without precise locality, in sandy soil (No. 8), in sandy soil near roadside (No. 14), in cultivated field (No. 13), April 24, 1918, *T. G. Harbison*; in low sandy pine woods, November 22, 1919, *T. G. Harbison* (Nos. 31, 32); Society Hill, no date, *M. A. Curtis* (Gray Herb., partly typical *R. atlanticum*).

This variety was based originally on the color of the flowers, but as a specimen kindly sent me by the author of the variety shows, it is at the same time rather densely pubescent and differs in this respect from the type of the species. As the pubescence seems to me the more important character, I prefer to include in this variety all the specimens with pubescent leaves without regard to the difference in the color of the flowers, which, moreover, seems to be not very pronounced, for I can hardly see any difference between the two forms as represented on the colored plate cited above.

Rhododendron oblongifolium Millais, *Rhodod.* 219 (1917). — Millais & Williams in *Rhodod. Soc. Notes*, I. 125 (1918).

Azalea oblongifolia Small, *Fl. S. E. U. S.* 883, 1336 (1903); in *N. Am. Fl.* XXIX. 43 (1914).

Shrub to 2 m. high or less, with irregularly whorled branches; young branchlets with short and rather sparse villose pubescence and usually with sparse strigose hairs or sometimes nearly glabrous; floral winter-buds with ovate, acuminate or obtuse, grayish pubescent scales. Leaves obovate or elliptic-obovate to oblong-ob lanceolate, rarely oblong, 4 to 10 cm. long and 1.5 to 3.8 cm. broad, acutish or acute, gland-tipped, cuneate, ciliate, dull green and glabrous above except the vil-

lose midrib or sometimes strigillose and minutely pubescent, light green, or rarely glaucescent beneath, pilose and strigillose on the midrib, more or less pubescent on the whole under surface or nearly glabrous; petioles 2 to 3 mm. long, finely pubescent. Flowers expanding at the middle or end of May or in southeastern Texas at the end of April, after the leaves are fully formed, and borne in 7- to 12-flowered clusters; pedicels 1.5 to 2 cm. long, sparingly villose and glandular-hirsute; calyx-lobes unequal, ovate to oblong or lanceolate, obtuse to acute, 1 to 3 mm. long, long-ciliate and sparingly villose outside; corolla funnel-form, white, the tube cylindrical, abruptly dilated at the apex, 2.5 to 3 cm. long, thinly villose and sparingly glandular-hirsute outside or more densely villose and destitute of stipitate glands except on the outside of the lobes, sparingly pubescent above the middle or glabrous inside, the lobes oblong-ovate, acute, 1.8 to 2 cm. long, glandular-hairy along the middle outside; stamens about twice as long as the tube, slightly exceeding the corolla-lobes, villose below the middle, the anthers ochraceous, 2.5 to 3 mm. long; style exceeding the stamens, 5 to 6 cm. long, finely pubescent at the lower third, or sometimes glabrous; ovary usually not or little longer than the sepals, covered with upright setose hairs partly glandular. Capsule oblong or ovoid-oblong to narrow-oblong 1.2 to 1.8 cm. long, finely villose and glandular-setose.

ARKANSAS. Newton County: rocky woods, Jasper, October 28, 1914, *E. J. Palmer* (No. 6932). Cleburne County: sandy woods, Heber Springs, October 31, 1914, *E. J. Palmer* (No. 6973); Pulaski County: Little Rock, *H. E. Hasse*. Pike County: Delight, May 20, 1910, *A. H. Howell* (No. 659; Nat. Herb. No. 514,206). Hempstead County: Fulton, sandy woods, May 21 and Oct. 5, 1909, *B. F. Bush* (Nos. 5684, 5955); Washington, moist sandy woods, Sept. 4, 1916, *E. J. Palmer* (No. 10,583); Columbus, along sandy streams, July 17, 1916, *E. J. Palmer* (No. 10,505). Ouachita County: Camden, 1850, *A. Fendler*. Miller County: Doddridge, moist sandy woods, July 19, 1916, *E. J. Palmer* (No. 10,515). "National Forest," 1907 (Nat. Herb. No. 615,731).

OKLAHOMA. Leflore County: near Page, near mountain creek, April 20 and September 7, 1913, *G. W. Stevens* (Nos. 1390, 2658), on low creek branch, June 20, 1914, *D. W. Blakeley* (*G. W. Stevens*, No. 1418; Nat. Herb. No. 589,543); rocky woods, valleys in limestone and lower levels of sandstone hills, July 27, 1917, *E. J. Palmer* (No. 12,644). McCurtain County: Idabel, May 29, 1916, *H. W. Houghton* (*G. W. Stevens*, No. 3925).

TEXAS. Harrison County: Marshall, common in swamp, Oct. 5, 1901, *B. F. Bush* (No. 983). San Augustine County: San Augustine, margin of sandy swamps, September 10, 1917, *E. J. Palmer* (No. 12,710). Houston County: Grapeland, 1879, *Frank Tweedy*¹;

¹ This is the type specimen collected at "Grapelands," as stated in a letter by *F. Tweedy* accompanying the specimen and dated March 15, 1899. On the label, which is not in Mr. Tweedy's handwriting, however, Tom Green County is given as the locality, which is apparently a mistake, for there are no *Rhododendrons*

sandy bogs, May 28, 1917, September 22 and 28, 1917 (Nos. 12,830, 12,831), March 26 and April 2, 1918, *E. J. Palmer* (Nos. 13,181, 13,265). Hardin County: Silsbee, sandy pine lands, April 25 and September 10, 1916, *E. J. Palmer* (No. 9568; shrub 4–6 cm. high, flowers pure white, No. 10,689). Without definite locality and date; *C. Wright* (in Gray Herb.).

This is the most western species of the eastern group of *Rhododendrons* and is distributed from northern Arkansas (Newton and Cleburne Counties) to southeastern Oklahoma and through southeastern Arkansas to southeastern Texas (Houston and Hardin Counties). It is a low shrub growing in moist sandy woods or on the margins of sandy bogs and along streams.

Rhododendron oblongifolium is most nearly related to *R. viscosum* and *R. serrulatum*, but differs chiefly in the finely villose, sparingly strigose branchlets, the pubescent winter-buds, in the larger leaves of thinner texture and usually more or less pubescent, in the larger corolla finely villose and glandular outside and in the usually longer calyx-lobes; the flowers appear earlier, in May or even at the end of April, while *R. serrulatum* flowers in July and August and *R. viscosum* from the end of May to the beginning of July. Without flowers the pubescent forms of it have some resemblance to *R. canescens* and have been sometimes confused with that species. It shows considerable variability in some characters, particularly in the pubescence of the leaves, which may be glabrous or more or less pubescent beneath; the extreme forms look very distinct but they are closely connected by numerous intermediates and both forms frequently occur at the same locality and have been distributed by the collectors even under the same number as Bush's No. 5955¹ from *Fulton*, Arkansas, and Palmer's No. 12,644 from *Page*, Oklahoma. The upper surface of the leaves also varies from being quite glabrous, except the pubescent midrib, to strigillose and with or without a minute villose pubescence. Sometimes the under side of the leaves is more or less covered with a glaucous bloom, as in part of Palmer's No. 12,644 from Oklahoma and in his 12,062 and part of 12,830 from Texas. The young branchlets may be puberulous all over and sparingly strigose toward the apex or nearly glabrous; the winter-buds are always densely pubescent. The calyx-lobes are sometimes very short and semi-orbicular or ovate, as in Palmer's Nos. 9568, 12,062 and 12,063 from Texas and in his No. 10,583 from Arkansas. The outside of the corolla is usually covered with a thin villose pubescence interspersed with numerous stipitate glands but sometimes, as in Palmer's Nos. 9568 and 12,063, the tube is more densely villose and destitute of glands, which are present only toward the apex of the corolla on the outside of the lobes; in the same numbers the style and the inside of the tube are glabrous. These characters together with the rather dense villose pubescence of the under side of the leaves of No. 12,063 might induce one to make this number the type of a distinct variety, but considering the great variability of the species even in the same locality and the apparent absence of any correlation of the different forms with their geographical distribution, I think it wiser to defer a subdivision of the species into varieties or forms until a careful field study has been made and more copious material is available.

known growing as far west as Tom Green County, Houston County marking the farthest western limit of the genus in the Eastern States.

¹ The glabrous form of this number looks very distinct, the leaves are broad and generally obovate, quite glabrous except sparingly strigillose on the midrib beneath, even the branchlets are quite glabrous and the specimen, which is in fruit, might be taken for *R. arborescens* but for the pubescent winter-buds.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

July 15–17 (Nos. 62, 63, 64, 65, 67, 70, 81, 82, 84, 87) and December 15, 1918 (No. 93), June 16 and 18 (Nos. 97, 99) and November 26 and 27, 1919 (Nos. 100, 102, 103, 110, 111, 112, 114, 115), *T. G. Harbison*; without locality, May 17, 1918, *T. G. Harbison* (Nos. 29, 33, 34, 46, 57). Lowndes County: Ousley, October 21, 1910, *T. G. Harbison* (Nos. 242, 246, 251); near Valdosta, June 6–12, 1895, *J. K. Small*. Brooks County: Quitman, April 2, 1918, *T. G. Harbison* (No. 5).

FLORIDA. Lake County: Eustis, June 1–15, 1894, *G. V. Nash* (No. 967, type); June 16–31, *G. V. Nash* (No. 1306); same locality, December 12, 1919, *T. G. Harbison* (Nos. 21, in part, 26, 27). Seminole County: near Wekiva Springs, May 9, 1918, *C. H. Baker* (No. 499 a–c). Orange County: near Maitland, February 27 (Nos. 478 a–b), June 20 (No. 529 a) and July 12, 1918, *C. H. Baker* (Nos. 529 b, 478 c, d, f); swamps, July 10, 1902, *A. Fredholm* (No. 5379; in Gray Herb.). Polk County: Lakeland, August 1, 1918, *T. G. Harbison*. De Soto County: Sebring, July 28, 30, 1918, *T. G. Harbison* (Nos. 19, 20, 21, 47). Escambia County: Pensacola, April 18, 1899, Biltmore Herb. (Nat. Herb. No. 969,619). Calhoun County: Apalachicola, *T. Drummond* (in Gray Herb.). Without precise locality: 1842–49, *F. Rugel* (No. 237; Nat. Herb. 512,161); without date, Herb. Chapman (Nat. Herb. No. 955,093).

ALABAMA. Mobile County: near Mobile, July, 1887, *C. Mohr*; same locality, December 4, 1919, *T. G. Harbison* (Nos. 1, 2); Spring Hill, August 6, 1897, *B. F. Bush* (No. 12); same locality, December 4, 1919, *T. G. Harbison* (Nos. 1–4).

MISSISSIPPI. Jackson County: Ocean Springs, June 19, 1895, *J. Skehan*. Harrison County: Biloxi, July 29, 1896, *C. L. Pollard* (No. 1097; Nat. Herb. 271,755); August 21, 1898, *S. M. Tracy* (No. 4395; Nat. Herb. 770,586); October 30, 1917, *T. G. Harbison* (No. 6), April 1 (No. 1), May 6 (Nos. 2, 3, 24) and July 19, 1918, *T. G. Harbison* (No. 25–33).

LOUISIANA. St. Tammany Parish: Pearl River, July 9, 1909, *R. S. Cocks* (No. 2222); New Orleans (Covington), 1832, *T. Drummond* (No. 190).¹

This Azalea is distinctly a species of the southern coastal plain region and extends from southern Georgia or, including the variety *georgianum*, from middle Georgia to central Florida (De Soto County) and westward along the coast to southeastern Louisiana. Its type locality is Eustis, Lake County, Florida. It inhabits swamps or moist sandy soil and is usually a tall shrub 2 to 7 m. high.

Rhododendron serrulatum is closely related to *R. viscosum* and takes its place south

¹ See footnote on page 153.

of central Georgia. It is chiefly distinguished by the brown-red branchlets usually conspicuously and densely covered toward the ends with strigose red-brown hairs, by the aristate or at least distinctly mucronate more numerous scales of the winter-buds, usually glabrous and with a conspicuous dark band along the margin, by the long and slender corolla-tube about twice as long as the comparatively small lobes, and by the stamens considerably exceeding the lobes; the leaves are of firmer texture and more distinctly serrulate on the margin and are not unfrequently, particularly those toward the end of vigorous shoots, more or less densely pubescent on the under side; it is usually a tall shrub, while *R. viscosum*, particularly in the southern part of its range, is usually a low, sometimes even stoloniferous shrub. Small in his description and keys lays much stress on the oval shape of the leaves, but though the type specimen which has only short and weak branchlets and no vigorous shoots happens to have most of its leaves of elliptic shape, this is the more unusual shape, while the prevailing shape of the leaves varies from obovate to oblong-obovate or sometimes oblanceolate. Until Small described it as a new species, *R. serrulatum* had been always confused with *R. viscosum* and even he understood under that name only the uncommon form with oval or elliptic leaves and referred the forms with obovate leaves to *R. viscosum*, making it a very local species, while really it has a rather wide distribution. Like the other related species it varies in the pubescence of its leaves, which may be glabrous except strigose hairs on the midrib beneath, as in the type, or more or less, sometimes densely pubescent on their under side and strigillose and puberulous above; the pubescence may appear either on all the leaves or chiefly on upper leaves of the more vigorous shoots, while the leaves appearing earlier in the season are glabrous or nearly glabrous. The form with the leaves always pubescent is described below as a distinct form. Sometimes the leaves of the glabrous form are glaucescent beneath, as in some specimens from Folkston, Georgia. Also the winter-buds vary in their pubescence; typically they are perfectly glabrous, but sometimes they are more or less densely pubescent and these specimens I have referred to a distinct variety, described below. In the shape of the corolla, in its pubescence and in its color, which is apparently always white or nearly white, I find little variation except that the tube varies somewhat in length.

This species was apparently first collected by T. Drummond about 1830 near New Orleans in Louisiana and near Appalachicola in Florida; it also was collected in Florida by T. Rugel in the forties and by Chapman in the fifties or sixties, but it was not distinguished until Dr. Small in 1903 described his *Azalea serrulata*, based on a specimen collected by G. V. Nash near Eustis, Florida, which had most of the leaves of distinctly oval or elliptic shape. The shape of the leaves, however, cannot be considered a reliable character, as already stated.

Rhododendron serrulatum was introduced into cultivation through the Arnold Arboretum, where plants were raised in 1919 from seeds collected by T. G. Harbison in December, 1918, near Folkston, Georgia. It will probably not be hardy north and as an ornamental shrub it will not be superior to *R. viscosum*, but for southern gardens it may be valuable on account of its late appearing fragrant flowers.

***Rhododendron serrulatum* f. *molliculum* Rehder, forma nova.**

A typo recedit ramulis annotinis apicem versus densius villosulis v. glabrius culis minus strigosis, foliis subtus molliter subaccumbenti-villosis supra sparsius villosulis et saepe praecipue in foliis superioribus strigillosis, pedicellis magis villosulis.

FLORIDA. Lake County: Eustis, June 23, 1919, T. G. Harbison (No. 17, type, 19, 20, 21, partly); December 12, 1919 (No. 25); June 16-30, 1894, G. V. Nash (No. 1104; Nat. Herb. 223,237).

GEORGIA. Charlton County: *Folkston*, June 16 and July 16, 1918, *T. G. Harbison* (No. 69, 98); without precise locality, May 18, 1918, *T. G. Harbison* (No. 58).

This form differs from the type in all the leaves being densely soft-pubescent beneath and slightly puberulous and often strigillose above; also the branchlets are usually more or less puberulous and the pedicels are more densely villose, while the winter-buds are glabrous as in the type. Pubescent leaves are also found occasionally in the typical form, but they are restricted to the upper part of the more vigorous branchlets, while the leaves of the short branchlets which are usually terminated by floral winter-buds and the lower leaves on the shoots are glabrous or nearly so. Such pubescent leaves are present, e. g. in Chapman's specimen, in Harbison's No. 6 from Lakeland, Florida, in his Nos. 21 (partly) and 27 from Eustis, in Nash's No. 1306 from the same locality, in Baker's No. 529 a from Maitland and in his No. 499 a from Wekiva Springs, in Harbison's No. 20 from Sebring, Florida, in his No. 84 and 100 from *Folkston*, Small's specimen from Valdosta, Harbison's No. 2 from Mobile, Alabama, and his Nos. 24, 27, 28 from *Biloxi*, Mississippi, and Cock's No. 2222 from Pearl River, Louisiana.

***Rhododendron serrulatum* var. *georgianum* Rehder, var. nov.**

A typo recedit gemmis dense sericeo-pubescentibus perulis paucioribus et tantum mucronatis vix aristatis.

GEORGIA. Charlton County: *Folkston*, July 16, 1918, *T. G. Harbison* (No. 68, type; shrub 10 ft. tall; No. 77), July 15 (No. 66), July 17, 1918 (No. 86, shrub 18 ft. tall) and Nov. 27, 1919, *T. G. Harbison* (No. 118, low shrub); without precise locality May 17 and 18, 1918, *T. G. Harbison* (Nos. 34 and 57; tall late-blooming species). Richmond County: *Augusta*, sand hills, swamp, July, 1904, *A. Cuthbert*. Bryan County: boggy places in sand hills of Canoochee River near *Groveland*, June 22, 1903, *R. M. Harper* (No. 1850). Screven County: near *Oliver*, July 9, 1901, *A. H. Curtiss* (No. 6837; 8 to 12 feet high). Appling County: swamps near *Baxley*, August 10, 1901, *Biltmore Herb.* No. 4823 (*Nat. Herb.* 980,173). Lowndes County: *Ousley*, October 21, 1910, *T. G. Harbison* (Nos. 244, 248). Randolph County: *Cuthbert*, December 1, 1919, *T. G. Harbison* (Nos. 31, 32; low shrub). Decatur County: moist pine barrens near *Climax*, August 13, 1903, *R. M. Harper* (No. 1929).

SOUTH CAROLINA. Beaufort County: *Bluffton*, 1893, *J. H. Mellichamp*.

This variety differs from the type only in the densely grayish pubescent winter-buds with usually less numerous and only mucronate, not aristate scales from the type. It approaches thus in its winter-buds *R. viscosum* and may be considered an intermediate form not only on account of its characters but also in regard to its distribution, occupying as it does an area between *R. viscosum* and typical *R. serrulatum*. I have not yet seen any specimens of this variety from Florida or from localities west of Georgia, while typical *R. serrulatum* does not seem to extend farther

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Anthodendron viscosum Reichenbach in Moessler, *Handb. Gewächsk.* ed. 2, I. 309 (1827).¹

Shrub to 3 or nearly 5 m. tall, sometimes low, rarely stoloniferous, with irregularly whorled branches; young branchlets, particularly toward the apex, with hirsute or strigose loosely appressed, rarely spreading hairs, otherwise glabrous, yellowish, or grayish brown, rarely red-brown at the end of the first season, the older branches light grayish brown; winter-buds glabrous sometimes pubescent, usually brown, with 8 to 12 broadly ovate rounded at apex and usually mucronate or sometimes without mucro, the basal scales sometimes long-pointed, ciliate on the margin. Leaves ovate or elliptic-obovate to oblong-ob lanceolate, acute or rounded and gland-tipped at apex, cuneate at base, 2 to 6 cm. long and 0.7 to 2 cm. broad, ciliate, glabrous on both sides except the midrib slightly villose above and more or less strigose beneath, sometimes strigillose above, particularly on leaves at the end of the shoots, usually dark green above, lighter green or glaucescent beneath, at maturity firm but rather thin; petioles 1 to 3 mm. long, strigillose. Flowers appearing after the leaves in June or July or farther south in May, in clusters of 4 to 9; pedicels 0.5 to 1, rarely to 1.5 cm. long, pubescent and glandular-hirsute; calyx-lobes semi-orbicular to ovate, about 1 mm. long, setosely glandular-ciliate; corolla white or more or less suffused with pink, rarely deep pink, cylindric, the tube somewhat dilated near apex, 1.5 to 2.5 cm. long, about one and one half times as long as the lobes or sometimes little longer than lobes, rarely almost twice as long, finely villose and glandular-hirsute outside, slightly pubescent inside above the middle, the lobes ovate-oblong, acuminate, about 1.5 cm. long, glandular-hirsute along the middle outside and finely villose; stamens exserted, somewhat longer than the lobes; the filaments villose except the upper third; style 4 to 5 cm. long, exceeding the stamens, finely pubescent below the middle or on the lower third, pale or sometimes purplish toward the stigma; ovary glandular-setose or sometimes setose and glandless. Capsule oblong-ovoid, 1.3 to 2 cm. long, glandular-setose and finely pubescent or sometimes glandless.

The species is distributed from southwestern Maine through the coastal plain to southeastern South Carolina, extending inland to central Massachusetts, northwestern Connecticut and up the valley of the Hudson River to the neighborhood of Albany,² northern New Jersey, central and southwestern Pennsylvania, eastern

¹ PRELINNEAN SYNONYMS: *Cistus Virginiana flore et odore Periclymeni* Banister apud Plukenet, *Phytog.* t. 161, fig. 4 (1691); *Almag.* 106 (1696). — *Azalea ramis infra foliosis* Gronovius, *Fl. Virg.* 21 (1739). — Colden in *Act. Soc. Sci. Upsala*, IV. 92 (*Pl. Coldengh.*) (1749).

ADDITIONAL ILLUSTRATIONS. COLORED PLATES: Kerner, *Darst. Vorz. Ausl. Bäum.* t. 57 (1796). — Meerburgh, *Pl. Select. Icon.* t. 9 (1798). — Schmidt, *Oesterr. Baumz.* III. t. 171 (1800). — Savi, *Fl. Ital.* II. t. 66 (1822). — Audubon, *Birds Am.* II. 115 (1831-34); Quarto ed. I. t. 64 (1840). — Emerson, *Trees & Shrubs Mass.* ed. 2, II. t. opp. p. 438 (1875). — *Meehan's Monthl.* X. 81, t. 6 (1900). — BLACK FIGURES: Loudon, *Arb. Brit.* II. figs. 947, 949 (1838), 947 after Loddiges t. 44, and 949 after *Bot. Reg.* t. 414. — Newhall, *Shrubs N. E. Am.* fig. 82 (1893). — Miller & Whiting, *Wild Flow. N. E. St.* 381 (1895). — Keeler, *Our North. Shrubs*, 355 (1903). — Stevens, *Ill. Guide Flow. Pl.* t. 114, fig. 5 (1910). — Schneider, *Ill. Handb. Laubholz.* II. figs. 329 n-p, 330 c-d (1911). — *Country Life Am.* XI. 496 (1907). — House, *Wild. Fl. N. Y.* II. fig. 22 (1920), as *A. canescens*.

² In the state of New York *R. viscosum* has been reported as far west as Oneida County by Payne, *Cat. Pl. Oneida Co.* 50 (in *Rep. Regents Univ. N. Y.*) (1865), but I have seen no specimens from that locality and Professor K. M. Wiegand writes me that he does not know it west of the Albany region and the Catskills.

West Virginia, western North Carolina and southeastern Tennessee and southwestern South Carolina (Oconee County). It also has been reported from northeastern Ohio.¹ It forms a densely-branched shrub from 1 to 5 meters tall and grows almost exclusively in swamps together with other swamp-loving shrubs and trees; it is rather common in the swamps of the coastal plain from southwestern Maine to South Carolina, but it also is found in swamps on the higher mountains.

Rhododendron viscosum is a very variable species, but it is never pubescent except in one rare variety and in this respect differs from the otherwise very closely related *R. serrulatum* from which it is often difficult to distinguish. With the latter species and with *R. arborescens*, *R. oblongifolium* and *R. atlanticum*, it forms a group of very closely connected species which have been frequently confused and are sometimes not easily distinguished from each other. The most distinct and least variable is *R. arborescens*, which differs from all these related species in the glabrous branchlets, glabrous leaves and usually glabrous style; *R. atlanticum* differs in its early flowers appearing before or with the leaves, in the absence of the villose pubescence of the corolla and in its low stature and stoloniferous habit; *R. oblongifolium* differs chiefly in its usually more or less pubescent, larger and narrower, acute or sometimes almost acuminate leaves, in the usually larger calyxlobes, the larger more villose and less glandular corolla and in the always pubescent winter-buds. The most closely related species is *R. serrulatum* which might be considered only a southern variety of *R. viscosum* and in fact was not separated at all from the latter species until recently; it may be chiefly distinguished by the more chartaceous, setosely serrulate, somewhat reticulate, often pubescent leaves never glaucous beneath, the more numerous always distinctly mucronate, often aristate scales of its floral winter-buds, the longer and slenderer corolla-tube about twice as long as the lobes and the bright red-brown branchlets usually densely strigose toward the apex. *Rhododendron viscosum* varies in the pubescence of its winter-buds, in the amount and character of the strigose or hirsute pubescence on the branchlets and leaves, in the color and shape of the leaves which may be bright green or more or less glaucous and vary from obovate to oblong-lanceolate, in the size and color of the flowers which may be almost pure white or more or less pink, rarely deep pink. Sometimes it has the low stoloniferous habit of *R. atlanticum*; e. g. I. Tidestrom's No. 7080, from between Berwyn and Lanham, Maryland, June 8, 1914, L. F. Ward's specimen from Terra Cotta Swamp, Washington, D. C., June 13, 1880 (Nat. Herb. No. 14,480), E. S. Steele's specimen from Hyattsville, Maryland, May 31, 1915 (Nat. Herb. No. 835,633) and Witmer Stone's No. 571, from Tyndalls Mill, south of Manning, South Carolina, May 5, 1917; also P. C. Standley's No. 11,547 from Hyattsville Swamp, May 27, 1915,

¹ Newberry, *Cat. Pl. Ohio*, 24 (1860). — Kellerman, *Cat. Ohio Pl.* 129 (in *Rep. Geol. Surv. Ohio*, VII. pt. 2) (1893). — Schaffner, *Cat. Ohio Vasc. Pl.* 204 (in *Ohio Biol. Surv.* I.) (1914). — *Rhododendron viscosum* was first reported from Ohio by Newberry in 1860 without definite locality except "East." Kellerman in 1893 quotes it from Beardslee's Catalogue of 1874, where it is recorded from near Painesville. Schaffner in 1914 gives Ashtabula County as the only locality for the species. Among the Azaleas from the herbarium of the Ohio State University received through the kindness of Professor J. H. Schaffner I found only one specimen labeled *Azalea viscosa* L., Pymatuning Swamp, Ashtabula County, August 11, 1900, C. A. Davis, A. Dachnowski, F. Detmers. This specimen has neither flowers nor fruits and is apparently some form of *R. roseum*; it is certainly not *R. viscosum*. Beardslee's record is probably based on a similar form; Professor Schaffner writes me that there are a few of Beardslee's specimens in the herbarium of the Ohio State University, but none labeled *Azalea viscosa*, and that Newberry's Catalogue was not based on specimens that could be verified.

which belongs to var. *glaucum* and a specimen from Fayetteville, North Carolina, Biltmore Herb. (Nat. Herb. No. 969,586) which belongs to var. *nitidum*. Several forms and varieties have been distinguished but on account of their mostly insufficient description and in the absence of type specimens it is not always possible to recognize them with certainty and the numerous intermediate forms make their distinction still more difficult.

Rhododendron viscosum was apparently first observed by John Banister, an English missionary who sent a drawing of it to Dr. Compton, Bishop of London. Dr. Compton communicated it to Plukenet, who published it in 1691 in his *Phytographia*. The plant was introduced into England apparently in the first half of the 18th century by John Bartram, who sent seeds or plants to Peter Collinson,¹ but the exact date is not known. According to Aiton it was introduced in 1734, but Catesby in 1731 speaks of this shrub in the text to the figure of his *Cistus virginiana*, which is *R. canescens*, as "having for some years past produced its beautiful and fragrant blossoms at Mr. Bacon's at Horton and at Mr. Collinson's at Peckham." I suppose Catesby confused several species, as appears from the fact that he applied Plukenet's name belonging to *R. viscosum* to his figure which represents *R. canescens*. His reference to fragrant blossoms seems to indicate that the shrub in cultivation was *R. viscosum*. In any case this species must have been well known about 1789 when Aiton enumerated five different varieties in cultivation. Wangenheim in 1787 published the first detailed description and characteristic figure; he mentions three varieties observed by him in the state of New York.

From the type (*R. viscosum a. odoratum* Sweet, *Hort. Brit.* ed. 2, 344 [1830]. — *Azalea viscosa a. odorata* Aiton, *Hort. Kew.* I. 203 [1789]. — *Azalea viscosa a. virens* Michaux, *Fl. Bor.-Am.* I. 151 [1803]), which has green leaves and the flowers white or sometimes striped pink outside along the middle of the lobes (*Azalea viscosa β. vittata* Aiton, *Hort. Kew.* 203 [1789]. — *Rhododendron viscosum β. vittatum* Sweet, *Hort. Brit.* ed. 2, 344 [1830]), the following varieties and forms may be distinguished:

***Rhododendron viscosum* f. *rhodanthum* Rehder, forma nova.**

A typo recedit floribus intense roseis.

MARYLAND. Prince Georges County: opposite Hyattsville, eastward, May 31, 1915, *E. S. Steele* (Nat. Herb. No. 835,633).

DISTRICT OF COLUMBIA. North of Washington, June 15, 1909, Professor *Cook* (Nat. Herb. No. 615,776).

This is a form of the typical variety with bright pink flowers very glandular outside. In the type specimen, which is from a low stoloniferous shrub, the very young winter-buds are glabrous outside; in Professor Cook's specimen they are silky-pubescent. This bright pink form occurs occasionally with the type, but seems to be less common than the pink form of var. *glaucum*.

Rhododendron viscosum* var. *glaucum* Torrey, *Fl. U. S.* I. 425 (1824). — Gray, *Syn. Fl.* II. 1, 41 (1878). — J. Robinson, *Fl. Essex Co. Mass.* 73 (1880). — Dame & Collins, *Fl. Middlesex Co. Mass.* 62 (1888). — Britton, *Cat. Pl. N. Jersey*, 162 (1889). — Coulter & Watson, *Gray's Man.* ed. 6, 320 (1890). — Dudley & Thurston, *Cat. Pl.

¹ See *R. nudiflorum* on page 136.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

region, but I have seen no specimens of it from Maine, New Hampshire, nor from North and South Carolina. A specimen from Lee County, Alabama, seems doubtful.

Rhododendron viscosum var. **nitidum** A. Gray, *Syn. Fl.* II. 1, 41 (1878). — Ward, *Guide Fl. Wash.* 95 (1881). — Owen, *Pl. Nantucket*, 39 (1888). — Britton, *Cat. Pl. N. Jersey*, 162 (1889). — Coulter & Watson, *Gray's Man.* ed. 6, 320 (1890). — Voss, *Vilmorin's Blumengärt.* I. 588 (1894), as forma. — Deane in *Rhodora*, I. 94 (1899); III. 197 (1901). — Fernald & Robinson, *Gray's New Man.* 631 (1908). — Graves & others, *Cat. Fl. Pl. Conn.* 308 (1910).

? *Azalea lucida* Meerburgh, *Pl. Select. Icon.* t. 7 (1798).

? *Azalea spathulata* Dumont de Courset, *Bot. Cult.* III. 335 (1811), as "sous-var." of *A. viscosa*.

Azalea nitida Pursh, *Fl. Am. Sept.* I. 153 (1814). — De Candolle, *Prodr.* VII. 716 (1839).

Rhododendron nitidum Torrey, *Fl. U. S.* I. 425 (1824). — G. Don, *Gen. Syst.* III. 847 (1834).

Azalea viscosa var. *nitida* Gray, *Man.* ed. 2, 257 (1856). — Wood, *Classb. Bot.* 489 (1870); *Am. Bot. Fl.* 203 (1870). — Nicholson, *Ill. Dict. Gard.* I. 150 (1887). — Britton in *Mem. Torr. Bot. Club*, V. 248 (1894); *Man.* 699 (1901). — Britton & Brown, *Ill. Fl.* 560 (1897). — Jelliffe, *Fl. Long Island*, 124 (1899). — Porter, *Fl. Pennsylv.* 239 (1903). — Keller & Brown, *Handb. Fl. Phila.* 246 (1905). — Millspaugh, *Liv. Fl. W. Virginia*, 323 (in *W. Va. Geol. Surv.*) (1913).

This form differs from the type chiefly in its dwarfer habit and in its smaller leaves obovate-oblong or narrowly elliptic to oblanceolate, acute at the ends or abruptly acuminate, green on both sides, sometimes strigose above. The flowers are usually more or less pinkish.

This form occurs with the type from Massachusetts to eastern North Carolina and is apparently a coastal plain form; the most southern specimens I have seen are from Newbern, Craven County, North Carolina (*T. G. Harbison*, Nos. 45, 46, 47, 49, 53, 55, 56, June 6, 1918); a specimen collected near the same locality (May 26, 1919; No. 129) by the same collector is typical *R. viscosum*.

Rhododendron viscosum var. **tomentosum** Rehder, comb. nov.

Azalea tomentosa Dumont de Courset, *Bot. Cult.* ed. 2, III. 336 (1811), as var. of *A. viscosa*. — Bosse, *Vollst. Handb. Blumengärt.* I. 341 (1840).

? *Azalea tomentosa* Dur. ex Steudel, *Nomencl.* I. 96 (1821), as var. of *A. procumbens*.¹

Azalea viscosa tomentosa Hort. Angl. ex Bosse, *l. c.* (1840), as synonym.

MARYLAND. Prince Georges County: Hyattsville, June 27, *F. H. Knowlton* (Nat. Herb. 335,818); June 4, 1905, *H. D. House* (No. 925 a, Nat. Herb. No. 492,824, in part).

¹ The author citation "Dur." is possibly a mistake for "Dum." Duroi describes only *Azalea viscosa* in his *Harbkesche Baumzucht*. Why Steudel refers *A. tomentosa* to *A. procumbens* is not clear to me.

DISTRICT OF COLUMBIA. Bancroft, *Four Mile Run*, June 6, 1915, *E. S. Steele* (Nat. Herb. No. 835,632).

NORTH CAROLINA. Cumberland County: Manchester, June 25, 1902, Biltmore Herb. No. 6610 (Nat. Herb. No. 969,584; shrub 6–8 ft.).

Cultivated. A specimen in the Gray Herbarium without locality labeled "28. *A. tomentosa* Loddiges."

This variety differs from the type chiefly in its leaves being finely but rather densely pubescent beneath and more or less so above. No reference to such a form appears in any American flora, but the plant must have been introduced to Europe more than a hundred years ago and was first described by Dumont de Courset, who was in doubt whether it was a distinct species or a variety of *A. viscosa* and describes it as having wooly branches, lanceolate-oval leaves soft-pubescent on both sides and late appearing white, fragrant flowers. Of the plant cultivated in England as *A. tomentosa* there is a sterile specimen in the Gray Herbarium which was probably incorporated in the herbarium between 1840 and 1850, as "Torr. and Gray, *Flora N. Am.*" appears on the label. This specimen agrees exactly with the spontaneous specimens cited above and especially with the specimen from Hyattsville. The leaves are generally oblong-obovate, 2.5 to 5.5 cm. long, acute or acutish, sparingly strigillose above, sparingly strigose on the midrib beneath; the petioles and young branchlets are puberulous and sparingly strigose. The leaves in the specimen from Manchester are slightly shorter and broader and more densely pubescent above and not strigillose; the corolla-tube in this specimen is slender, about 2.8 cm. long, rather densely villose and sparingly glandular-pilose. The specimen from the District of Columbia has less densely pubescent leaves and a more glandular and less villose corolla as has the specimen from Hyattsville. This is a rather remarkable variety, as I have not observed in any of the numerous specimens of *R. viscosum* that I have seen any tendency to have the leaves pubescent, although the midrib may be sometimes slightly pubescent below.

Rhododendron viscosum* var. *hispidum Voss, *Vilmorin's Blumen-gärt.* I. 588 (1894), as forma. — Schneider, *Ill. Handb. Laubholz.* II. 502 (1911), as var. — Rehder in *Bailey, Stand. Cycl. Hort.* V. 2942 (1916).

? *Azalea scabra* Dumont de Courset, *Bot. Cult.* ed. 2, III. 335 (1811), as var. of *A. viscosa*.

Azalea hispida Pursh, *Fl. Am. Sept.* I. 154 (1814). — Watson, *Dendr. Brit.* t. 6 (1825). — De Candolle, *Prodr.* VII. 2, 716 (1839).

Rhododendron hispidum Torrey, *Fl. U. S.* I. 425 (1824). — Loudon, *Arb. Brit.* II. 1144, fig. 948 (1838). — G. Don, *Gen. Syst.* III. 847 (1834).

Azalea glauca β . *hispida* Heynhold, *Nomencl. Bot. Hort.* I. 108 (1840).

Azalea viscosa var. *hispida* Wood, *Classb. Bot.* 490 (1870); *Am. Bot. Flor.* 203 (1870). — Britton in *Mem. Torr. Bot. Club*, V. 248 (1894); *Man.* 699 (1901). — Britton & Brown, *Ill. Fl.* II. 559 (1897). — Porter, *Fl. Pennsylv.* 239 (1903).

This form differs from the type chiefly in its hispid young branchlets, the glaucous rather narrow leaves usually strigose above and in the hispid pedicels. The flowers are usually more or less carmine and have according to Pursh frequently ten stamens. The habitat of this form which is described as a shrub 3 to 5 m. tall

is given by Pursh as occurring "on the borders of lakes on the highest parts of the Blue Mountains: New York and Pennsylvania"; by Britton it has been reported from Shawangunk Mountain, Ulster County, New York, and Montauk Point, Long Island, and by Porter from Monroe County, Pennsylvania, but I have not yet seen a specimen which agrees with Pursh's description. The specimen from Montauk Point I consider a rather copiously strigose specimen of typical *R. viscosum*.

Rhododendron viscosum var. montanum Rehder, var. nov.

A typo recedit gemmis floriferis perulis paucioribus sericeo-pubescentibus praeditis, ramulis minute-villosulis et praesertim apicem versus pilis longis patentibus pallidis instructis ut petioli, foliis subtus ad costam sericeo-strigillosis v. subpatentibus hirsutis et satis dense v. interdum sparsissime minute villosulis vel interdum fere glabris, pedicellis dense glanduloso-pilosis.

NORTH CAROLINA. Macon County: Highlands, alt. 3825 and 3900 ft., July 9 and September 30, 1918 (No. 58, type, No. 60); alt. 3900 ft., June 20 and September 30, 1918 (Nos. 40, 41), July 9, 1918 (No. 63), June 13, 1918 (No. 38), June 20, 1918 (Nos. 31, 36), September 2, 1910 (No. 117), June 10, 1911 (No. 650), July 2, 1911 (No. 656), *T. G. Harbison*; Mt. Satuloh, June 21, 1898, *A. P. Anderson* (No. 1507; Nat. Herb. No. 344,433). Buncombe County: Biltmore, June 12, 1897 (Biltmore Herb. No. 1399 a), vicinity of Black Mountain, Sept. 8, 1913, *P. C. Standley & H. C. Bollman* (No. 10,453; Nat. Herb. No. 689,191). Wake County: Raleigh, June 11, 1918, *T. G. Harbison*.

This variety differs from the type chiefly in its pubescent winter-buds with comparatively few scales, in the spreading hairs and the minute pubescence of the branchlets and pedicels and of the midrib of the under side of the leaves. It is a low densely branched shrub, usually not exceeding 1.25 m. in height. The silky-pubescent winter-buds have only 5 to 7 larger broad rounded scales with 3 to 4 small acuminate scales at the base. The young branchlets are minutely puberulous and toward the apex more or less densely furnished with long spreading pale hairs not brown and loosely accumbent as on the otherwise glabrous branchlets of the typical plant. The leaves are obovate to oblong-obovate, 2.5 to 4 cm. long, glabrous and bright green somewhat lustrous above, the terminal leaves very rarely strigose, lighter or paler green beneath, the midrib with long pale subaccumbent or sometimes more or less spreading hairs and besides usually minutely villose particularly toward the base; the very short petioles have a pubescence similar to that of the branchlets. The flowers appear from the middle of June to the beginning of July; their pedicels are finely villose and rather densely glandular-pilose; the corolla is finely villose and furnished with numerous red long-stipitate glands and seems to be usually more or less suffused with pink; the calyx-lobes are very short; the style is purplish above, with purple stigma. The capsule is oblong-ovoid, 0.8 to 1.5 cm. long, finely villose and glandular-setose. The variety seems to be similar to and may be identical with *Azalea hispida* of Pursh, which is also a mountain form of *R. viscosum*; but Pursh describes his plant as 10 to 15 feet tall, the leaves as hispid above and the calyx-teeth as oblong, while this variety is a low shrub, with the leaves smooth above or only very exceptionally strigillose above, and the calyx-teeth very short.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

strigillose midrib, and slightly reticulate. The flower-clusters are large and many-flowered; the sepals semiorbicular and minute; the corolla-tube is about 3 cm. long, villose and glandular-hirsute; the setose hairs of the ovary are tipped with purple glands. Part of No. 36 consists of a branch with the leaves narrower and puberulous above and short-villose beneath. Whether the numbers 17 and 18 collected in bloom on March 29 belong here is doubtful. The leaves are half-grown, glaucescent on both sides and glabrous except the strigillose midrib beneath, the corolla is less pubescent and the tube somewhat more gradually dilated at the apex. It has some resemblance to *R. atlanticum*, but the corolla is less glandular and more villose, the winter-buds are pubescent and the habit is apparently different. Some specimens from Lee and Dallas Counties, Alabama, which I have referred to typical *R. viscosum*, may belong rather to this variety than to *R. viscosum*.

Aiton's *Azalea viscosa* γ . *fissa* (Hort. Kew. I. 203 [1789]. — *Rhododendron viscosum* ν . *fissum* Sweet, Hort. Brit. ed. 2, 344 [1830]) described as "floribus albis ad basin usque divisis, foliis saturate viridibus lucidis" is apparently not a form with a regularly five-parted corolla like his *A. nudiflora* η . *partita*, for Dumont de Courset describes the same form under *Azalea serotina* N., *A. fissa* H. K. (*Bot. Cult.* ed. 2, III. 336 [1811]), as follows: "Le tube de la corolla est plus court; le limbe se fend quelquefois dans une des ses divisions jusqu'au tiers ou la moitié du tube, même jusqu'à sa base; dans plusieurs corolles il ne se fend pas." Such split corollas may be seen occasionally and seem to occur in certain individuals more frequently than in others; for instance Harbison's No. 656 of var. *montanum* shows a large percentage of more or less deeply split corollas, but this can be considered only as an abnormality and not as a taxonomic form.

A form with double flowers is mentioned by Dumont de Courset in 1814 (*Bot. Cult.* VII. 166) as "Azalea visqueuse à fleurs doubles," and in 1864 Kirchner (in Petzold & Kirchner, *Arb. Musc.* 479 [1864]) enumerates an *Azalea viscosa flore pleno*. I have seen neither a plant nor a specimen of a double-flowered true *R. viscosum*; the forms I have seen under this name were apparently of hybrid origin. There is also an *Azalea glauca flore semipleno* Jäger (*Ziergeh.* 118 [1865]). — *Azalea viscosa glauca* sousvar. *flore pleno* Mouillefert, *Arb. Arbriss.* II. 1021 [1897]. — ? *Azalea glauca plena* Hort. Kew. [in herb. Arnold Arb.]. Under the last named synonym a specimen was collected by G. Nicholson in Kew Gardens in 1880; the leaves are glaucescent beneath and the corolla has the shape of that of *R. viscosum*, but it is almost destitute of villose pubescence, the sepals are well developed and oblong and the young branchlets are almost glabrous except stipitate glands. These characters, together with the early flowering time (June 1) when the leaves were only partly developed, suggest *R. atlanticum*.

Rhododendron arborescens Torrey, *Fl. U. S.* 425 (1824). — G. Don, *Gen. Syst.* III. 847 (1834). — Loudon, *Arb. Brit.* IV. 1145 (1838). — Gray, *Syn. Fl.* II. 1, 40 (1878). — Sargent, *Gard. & For.* I. 400, fig. 64 (1888). — Dippel, *Handb. Laubholz.* I. 416, fig. 269 (1889). — Coulter & Watson, *Gray's Man.*, ed. 320 (1890). — Robinson & Fernald, *Gray's New Man.* ed. 7, 631 (1908). — Schneider, *Ill. Handb. Laubholz.* II. 500, fig. 329 l-m, 330 b (1911). — Millais, *Rhodod.* 115 (1917).

? *Azalea viscosa* Marshall, *Arbust. Am.* 15 (1785), not Linnaeus.

Azalea arborescens Pursh, *Fl. Am. Sept.* 152 (1814). — De Candolle, *Prodr.* VII. 2, 716 (1839). — Gray, *Man.* 268 (1848); ed. 5, 299 (1867). — M. A.

Curtis, *Descript. Trees & Shrubs* (in *Geol. Nat. Hist. Surv. N. Car.* III. Bot.) (1860). — Britton & Brown, *Ill. Fl.* II. 559, fig. 2746 (1897). — Britton, *Man.* 698 (1901). — Gattinger, *Fl. Tenn.* 131 (1901). — Porter, *Fl. Pennsylv.* 238 (1903). — Keller & Brown, *Handb. Fl. Phila.* 246 (1905). — Small, *Fl. S. E. U. S.* 883 (1903); in *N. Am. Fl.* XXIX. 43 (1914). — Millspaugh, *Liv. Fl. W. Virginia*, 322 (in *W. Va. Geol. Surv.*) (1913).

Azalea verticillata Carr, *Cat. Trees & Shrubs Bartram's Bot. Gard.* 11 (1814), name only. — Loddiges, *Bot. Cab.* XVII. t. 1632 (1830).

Azalea arborea Bartram ex Pursh, *Fl. Am. Sept.* 152 (1814), as synonym.

Azalea fragrans Rafinesque, *Ann. Nat.* 12 (1820), not Adams.¹

Upright shrub to 3 m., rarely a small tree 6 m. tall; young branchlets glabrous or rarely with a few scattered strigose hairs, often slightly bloomy, yellowish brown or reddish brown at the end of the first season, becoming light gray or light grayish brown the second year; floral winter-buds glabrous, light brown, with broadly ovate rounded and usually mucronulate ciliate scales. Leaves obovate or sometimes elliptic to oblong-oblong, acute or obtuse and gland-tipped at the apex, cuneate at base, 3 to 8 cm. long and 1.3 to 3 cm. broad, ciliate, lustrous bright green above, glaucous or glaucescent or sometimes nearly green beneath, glabrous except sparingly short-villose on the midrib above and sparingly strigose on the midrib beneath or sometimes quite glabrous beneath; petioles slender, 3 to 7 mm. long, glabrous or rarely with a few strigose hairs. Flowers very fragrant, appearing the end of June or beginning of July in 3- to 6-flowered, umbel-like clusters; pedicels slender, 0.8 to 2 cm. long, glandular-hirsute or without glands, sometimes glabrous; sepals ovate to linear-oblong, usually obtuse, 3 to 6 mm. long, glandular-ciliate, otherwise glabrous or sometimes villose; corolla funnel-form, white or pinkish, the tube 2.5 to 3 cm. long, slightly dilated at the apex, usually sparingly, rarely densely glandular-hirsute outside, otherwise glabrous or slightly, rarely rather densely villose, pubescent inside except on the lower third, lobes ovate-oblong, acuminate 1.5 to 2 cm. long; stamens exerted about twice as long as the tube, the filaments purple above, pubescent below the middle; anthers 2 to 3 mm. long, ochraceous; style as long or somewhat longer than the stamens, 5.5 to 7 cm. long, purple above, glabrous or sometimes finely pubescent near the base, rarely villose on the lower third; ovary glandular-setose, with reddish glands. Capsule oblong-ovoid, 0.8 to 1.7 cm. long, densely glandular-hispid.

This species is distinctly a species of the Appalachian Mountain regions and does not occur in the coastal plane but along the Susquehanna it descends to the southern part of Lancaster County, Pennsylvania, along the Potomac to Fairfax County, Virginia, along the Savannah River to Richmond County, Georgia, and along the Chattahoochee River to Decatur County, Georgia; it is distributed from southern Pennsylvania through western Maryland, northern and western Virginia, eastern West Virginia, and eastern Kentucky, western North Carolina, northwestern South Carolina, eastern Tennessee to northern (Winston County) and eastern Alabama (Talladega and Clay Counties)² and to eastern (Richmond County) and southwestern Georgia (Decatur County). It grows chiefly on the banks of mountain streams, rarely on the borders of swamps and ascends to an altitude of 5200 feet on the mountains of North Carolina.

¹ ADDITIONAL ILLUSTRATIONS. COLORED PLATE: Lounsberry, *Guide to Trees*, t. 75 (1900). — BLACK FIGURES: Keeler, *Our North. Shrubs*, 353, fig. (1903). — *Garden Mag.* V. 219, fig. (1907). — *Country Life Am.* XI. 496 (1907). — Bailey, *Stand. Cycl. Hort.* V. f. 3388 (1916), after Gard. & For.

² Sheet No. 770,587 in the National Herbarium labeled "Rhododendron viscosum, borders of stream in thickets, Pointclear, C. Mohr, June 16, 1889," contains

Rhododendron arborescens is most nearly related to *R. viscosum*, but is easily distinguished from this and other Azaleas by the glabrous branchlets, glabrous leaves glaucous beneath and the large white or pinkish flowers. It shows little variation except in the length and shape of the sepals, in the amount of pubescence on the outside of the corolla, in the pubescence of the pedicels occasionally quite or nearly glabrous, in the pubescence of the style, which is glabrous or sometimes more or less pubescent below.

Rhododendron arborescens was probably discovered by John or William Bartram and appeared, according to Pursh, in Bartram's catalogue as *Azalea arborea*, while in the catalogue issued by Robert Carr in 1814 it appears as *A. verticillata*; also "*Azalea caroliniana*, Sweet Honeysuckle," in an undated "Catalogue of American trees and shrubs . . . in John Bartram's Garden" may possibly be the same species. Also Humphrey Marshall seems to have known it, for his *Azalea viscosa* (*Arbust. Am.* 15 [1785]), which he describes as a shrub five or six feet high growing in rich rocky places near streams of water, is probably *R. arborescens*, while his *A. viscosa palustris* is the true *R. viscosum*.

Michaux observed it in May, 1795 (see his *Jour.* ed. Sargent, 116), on the Blue Ridge in North Carolina, but there seems to be no specimen of it in his herbarium. Loddiges, who published an excellent colored plate of this species under the name *Azalea verticillata*, states that it was received "many years since from our old friend Bartram under this name," which tends to show that it was introduced before the plant was known as *A. arborescens* and before 1818, the date given by Sweet in his *Hortus Britannicus* under *A. arborescens* as the date of its introduction to England, but neither Sweet nor Loudon say who introduced it at that date; probably it was John Lyon, who was an active collector in North America up to that time. It was first described in 1814 as a new species by Pursh, who discovered it on the Blue Mountains of Pennsylvania, and also saw it cultivated in John Bartram's garden at Philadelphia; he called it the finest ornamental shrub he knew. It certainly is a very handsome shrub and it seems strange that it is still so little known in cultivation; the dry leaves have a very pleasing smell of coumarin, like *Melilotus* and *Anthoxanthum*, and retain it for many years in herbarium specimens.

***Rhododendron arborescens* var. *Richardsonii* Rehder, var. nov.**

A typo praecipue recedit statu humiliore plerumque 0.60–1.5, rarius ad 2.5 m. altum, foliis paulo minoribus et magis glaucescentibus, corolla densius villosa et magis glanduloso-hirsuta, sepalis plerisque brevibus ovatis vel semiorbicularibus extus villosis, stylo plerumque in triente inferiore dense villosa.

NORTH CAROLINA. Macon County: on Wayoh Bald, alt. 5200 feet, June 7, 1919, *T. G. Harbison* (No. 170, type, compact shrub, 4 feet tall, growing in dense patches), alt. 4800 feet (No. 176); Highlands, on mountain side in moist soil, alt. 4400 or 4500 feet, June 23, 1918, *T. G. Harbison* (Nos. G 34, D 42); alt. 4000 feet, July 9, 1918, *T. G. Harbison* (No. 62); without precise locality, alt. 3800 feet, July 2, 1911 (No. 655); high mountains, Sept. 2 and 6, 1910 (Nos. 114, 132),

two different species, one being *R. arborescens*. If the locality, which is not clearly written, is Pointclear, Baldwin County, the range of *R. arborescens* would extend to the coast of Alabama, but until the occurrence of the species there is confirmed by other collections I consider it doubtful.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

This plant is known only in southwestern Georgia and eastern Alabama and grows in shady ravines on the banks of streams. The type locality is about two miles northwest of Cuthbert, where it was collected by R. M. Harper in 1903 and first described by Small in 1913. It is the most distinct of the American species of the section *Pentanthera* and shows no very close relation to any species of this section. It is the most glabrous of all American species and entirely destitute of any glandular pubescence except occasionally some glands on the outside of the corolla-lobes; in its glabrousness it comes nearest to the preceding species, but that has generally obovate smaller leaves distinctly glaucous beneath and a glandular and pubescent corolla; in the leaves, the winter-buds and in the branchlets it resembles *R. Vaseyi* so much that it is not easy to distinguish sterile specimens, but the flowers are entirely different. The shape of the corolla of *R. prunifolium* is much like that of *R. calendulaceum* and *R. occidentale*, but these species differ widely from *R. prunifolium* in most other characters.

Rhododendron prunifolium was introduced into cultivation through the Arnold Arboretum, where plants were raised in 1918 from seeds collected by T. G. Harbison near Cuthbert, Georgia, in November, 1917.

ENUMERATION OF THE SPECIES

SPONTANEOUS HYBRIDS

No spontaneous Azalea hybrids have been as yet recorded, but I have before me several specimens which may possibly be of hybrid origin. There are some specimens collected by Professor C. S. Sargent near Bath, Georgia, on April 29, 1914, which appear intermediate between *R. canescens* and *R. speciosum*. They differ from the former in the red flowers with shorter more dilated corolla-tube destitute of glands and with larger limb and from *R. speciosum* in their smaller flowers, more densely pubescent leaves and more or less pubescent winter-buds. As the two supposed parent species have been collected at the same locality, and flower at about the same time, the occurrence of such a hybrid is not improbable. Possibly of the same origin is T. G. Harbison's No. 900 from Stone Mountain, De Kalb County, April 29, 1912; it much resembles *R. speciosum*, but the flowers are apparently of much lighter color and the winter-buds are pubescent; both supposed parent species occur in the region.

On the Nantahala Mountains, Macon County, North Carolina, T. G. Harbison collected a specimen (No. 186; June 7, 1919) similar to *R. calendulaceum* but with narrower and longer corolla-tube and pink flowers; it may possibly be a hybrid between *R. calendulaceum* and *R. nudiflorum*, but I am not aware whether the latter species is found in that particular locality, though it occurs in Macon County. On the Chilhowee Mountain, Tennessee, A. H. Curtiss collected an Azalea (No. 1719) which resembles closely *R. arborescens*, but has pubescent branchlets; whether this is a hybrid possibly with *R. viscosum*, I am not prepared to say.

GARDEN HYBRIDS

There exists in gardens, especially in Europe, on the Continent as well as in England, a bewildering variety of Azaleas which are undoubtedly of hybrid origin; in fact almost all of the so-called hardy Azaleas in cultivation are hybrids, the true species having almost entirely disappeared from gardens. The parentage of most of these forms is so mixed that it is impossible to recognize their origin with certainty, particularly if one is not sure of at least one of the parents. The difficulty is increased, as pointed out before, by the absence of strong morphological characters between the different species of this group, so that it is sometimes difficult to separate even the spontaneous forms satisfactorily. Many of the older hybrids have been figured, but usually without adequate description and with often vague and indefinite indication of the parentage; and as no herbarium material is available to supplement the illustrations, we have to depend on the figures and the indication of the parentage as given. Some of these older forms may still be in cultivation, and if one were able to observe and compare a large collection of them in cultivation, many of the problems might be solved, though the only sure way to elucidate the origin of these hybrids would be by raising plants from seed of self-fertilized flowers, that is from flowers carefully protected from foreign pollination and fertilized by pollen from the same plant. In some cases it may be necessary to raise more than one generation to arrive at satisfactory results. All I can do here is to assemble the more important literature relating to these hybrids as a basis for further studies.

It seems rather strange that the first hybrids known to have been raised in this genus are not between species of the Azalea group, but between Azaleas and true Rhododendrons. The very first hybrid which we know is one raised between *Rhododendron ponticum* and some form of Azalea; it originated at Thompson's Nursery at Mile End near London before 1800 and was figured by Andrews in 1801 as *R. ponticum* var. *deciduum*. This is possibly the plant distributed afterwards as *R. azaleoides* by English nurserymen, for in 1811 and 1814 Dumont de Courset describes an identical or very similar form under that name, giving "Hort. lond." as author citation. Similar crosses have

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

R. calendulaceum, *R. nudiflorum*, *R. speciosum*, *R. viscosum* and the Old World *R. luteum*. An interesting hybrid was raised by the Rev. William Herbert at Spofforth by crossing *R. luteum* and *R. canadense*, the species representing two different sections; it was figured by Lindley in 1838 as *Azalea Seymouri*, but apparently disappeared soon from cultivation, as it had no particular ornamental merits.

After *R. molle* G. Don (*Azalea sinensis* Lodd.), introduced in 1823 from China, had found its way into the gardens it also was used to produce new hybrids, and a hybrid with *R. viscosum* was figured as *Azalea altaclarensis* in 1842, one with *R. speciosum* as *Azalea marginata* in 1835 and one with a form of *R. Mortieri* was figured in 1845 as *A. sinensis* var. *Bylsiana*. The influence of *R. molle*, however, does not seem to have left a perceptible trace among the older *Azalea* hybrids, but when crossed with *R. occidentale* A. Gray introduced about 1850, which was used by Anthony Waterer of Knap Hill and by others in hybridization work with this and other hardy *Azaleas*, a very handsome race of hardy free-flowering *Azaleas* with large flowers of prevalently light colors was developed, classed here under *R. albicans*. A still greater influence on the development of hybrid *Azaleas* took place when *R. japonicum* was introduced into the gardens in the early sixties. It became soon widely known under the name *Azalea mollis* and by hybridizing it with other hardy *Azaleas* a distinct race known as "Azalea mollis hybrids" (*R. mixtum*) was raised, characterized by low compact habit and dense clusters of large scentless flowers with a wide limb and short tube. Among the "Mollis hybrids" are often included the hybrids of *R. japonicum* with *R. sinense* for which the name *R. Kosterianum* has been proposed; the first hybrids of this kind appear to have been raised in the seventies (see p. 97). About 1890 another race of hybrids was raised by crossing *R. japonicum* with evergreen *Rhododendrons* (see p. 99); these were called by André "Azaleodendron." Recently an interesting hybrid between two different sections was raised by G. Fraser of Ucluelet, Vancouver, B. C., by crossing *R. japonicum* with *R. canadense*. This promises to be a very ornamental plant unlike the older *R. Seymouri*, the first known hybrid of *R. canadense*.

In the following enumeration the different garden forms are placed under their supposed parents, though in many cases it is not certain whether the parentage given in the original description or the opinion based on the evidence presented is correct. If an available binomial exists, it follows the indication of the parentage and is printed in

heavy type. No attempt has been made to have binomial designations for all the different crosses; only when the plant is known to be still in cultivation and had no available name under *Rhododendron*, new combinations have been formed. The hybrids between Old World species have been included, as far as they have not been mentioned in the preceding paper by my colleague, E. H. Wilson, to make the enumeration of the *Azalea* hybrids as complete as possible.

SECT. PENTANTHERA

Rhododendron calendulaceum × **nudiflorum** (and partly × **speciosum**) = **Rhododendron Mortieri** Sweet, *Hort. Brit.* ed. 2, 343 (1830), as *R. Morterii*, name only.

Rhododendron Morterii var. *a. carnea* Sweet, *Hort. Brit.* ed. 2, 343 (1830), name only; *Brit. Flow. Gard.* ser. 2, I. t. 10 (1831).

Azalea calendulacea v. *Morterii* De Candolle, *Prodr.* VII. 718 (1839).

Azalea Mortieriana Spae in *Ann. Soc. Agric. Bot. Gand*, II. 325, t. 81 (1846), in part.

“*Azaleae Mortierianae*” Morren in *Ann. Soc. Agric. Bot. Gand*, III. 9, t. 106 (1847), in part.

Azalea Mortieri K. Koch, *Dendr.* II. 190 (1872), as synonym.

Rhododendron calendulaceum var. *Mortieri* Nicholson in *Kew Hand-list Arb.* ed. 2, 497 (1902).

Rhododendron Mortieri Zabel in *Mitteil. Deutsch. Dendr. Ges.* XI. 30 (1902).

This is stated by Sweet to be a hybrid of and intermediate between *R. calendulaceum* and *R. nudiflorum* and judging from the colored plate there is no reason to doubt that statement. The pink flowers are large with spreading limb consisting of broad lobes of which the upper is almost covered by an orange-yellow blotch. From this form named *a. carnea*, which must be considered the type, the form *praestans* differs in its pale copper-colored flowers tinged with blush and with the upper lobe of the limb coppery yellow. Sweet states that this plant was received from the Continent under the adopted specific name.

Of the same parentage and probably referable here as synonyms or varieties are the following forms:

Azalea calendulacea v. *flammea* Sims in *Bot. Mag.* XLVII. t. 2143 (1820), not Michaux.

The plant figured is apparently neither *R. speciosum* nor a true *R. calendulaceum*; the pink corolla tube and the partly pink, partly orange-colored lobes, except the upper one which is entirely orange, may be taken to indicate the influence of *R. nudiflorum*, though there is a possibility that a variation of *R. calendulaceum* with more or less pinkish flowers exists.

Azalea nudiflora ππ. *tricolor* Sweet, *Hort. Brit.* 265 (1826), name only. — Loddiges, *Bot. Cab.* XIV. t. 1382 (1828).

? *Azalea tricolor* Carr, *Cat. Trees & Shrubs Bartram's Bot. Gard.* 10 (1814), name only.

Rhododendron nudiflorum ππ. *tricolor* Sweet, *Hort. Brit.* ed. 2, 344 (1830).

Rhododendron tricolor Sweet, *Hort. Brit.* ed. 2, 343 (1830), “Hybrid 1816.” — Millais, *Rhodod.* 255 (1917).

This form is stated to have been received from Philadelphia in 1821, but it belongs apparently to this group of hybrids; the rather slender deep red tube and

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

Rhododendron luteum × speciosum.

Azalea nudiflora var. *scintillans*¹ Lindley in *Bot. Reg.* XVII. t. 1461 (1831).

Rhododendron nudiflorum var. *scintillans* Hooker in *Bot. Mag.* LXV. t. 3667 (1839).

In the *Botanical Register* Mr. Gowen says: "This Azalea was raised at Highclere in the same year with those already figured in previous parts of this work and is a seedling from *Azalea coccinea major* impregnated by the pollen of *Azalea pontica*. The flowers are bright pink with the narrow lobes of *R. nudiflorum* and with the upper lobe of a pale buff color or nearly whitish."

The following are probably of the same parentage:

Rhododendron flavum var. *ardens* Sweet, *Brit. Flow. Gard.* ser. 2, IV. t. 306 (1835).

Azalea pontica * *coronaria* β. *ardens* De Candolle, *Prodr.* VII. 718 (1839).

This form was introduced from Holland, according to Sweet, and he says that it "is doubtless a production between *R. flavum* and one of the varieties of *R. nudiflorum*." Judging from its bright scarlet flowers borne in a very large dense head it is near *R. speciosum* and if not simply a form of that species it may very well be a hybrid between it and *R. luteum*.

Rhododendron flavum var. *coronarium* Sweet, *Brit. Flow. Gard.* ser. 2, IV. t. 331 (1836).

Azalea pontica * *coronaria* α. *flavum* De Candolle, *Prodr.* VII. 718 (1839).

This is very similar to the preceding except that the flowers are orange-yellow with a red tinge. It also was imported from Holland.

Rhododendron luteum × viscosum = Rhododendron ornatum Sweet
Brit. Flow. Gard. ser. 2, II. t. 137 (1832).

Azalea pontica-rubescens Gowen ex Sweet, *l. c.* (1832) as synonym.

Rhododendron viscosum β. *ornatum* G. Don, *Gen. Syst.* III. 847 (1834).

Azalea viscosa * *ornata* De Candolle, *Prodr.* VII. 617 (1839).

Sweet states that he received the specimen figured the end of May, 1830, and that "the seeds were raised by Lord Carnarvon's gardener from *Azalea viscosa* β. *rubescens*, fertilized by *A. pontica* under Mr. Gowen's own inspection."

Four varieties are distinguished and figured:

α. *speciosum*, with reddish orange flowers (fig. 1).

β. *incarnatum*, with pale pink flowers tipped with white, the upper lobe yellow (fig. 3; in the text by mistake as var. and fig. 2).

γ. *luteum* with pale yellow flowers slightly tinged with blush, the upper lobe deep yellow (fig. 2; in the text by mistake as var. 3 and fig. 3).

δ. *roseum* with pale rose flowers, the upper lobe with a large orange blotch (fig. 4).

Rhododendron Mortieri × luteum (and partly viscosum or arborescens) = Rhododendron gandavense Rehder, comb. nov.

¹ This name should not be used for this plant as a binomial designation under *Rhododendron*, as there is already a *R. scintillans* Balf. f. & W. W. Sm.

Azalea Mortieriana Spae in *Ann. Soc. Agric. Bot. Gand*, II. 325, t. 81 (1846), in part (twelve named varieties are figured).

"*Azaleae Mortierianae*" Morren in *Ann. Soc. Agric. Bot. Gand*, 9, t. 106 (1847), in part.

Azalea pontica et var. Witte, *Flora*, 290, t. 73 (1868).

Azaleae Gandavenses K. Koch, *Dendr.* II. 185 (1872), as synonyms.

Azalea Gandavensis Hort. apud K. Koch, *Dendr.* II. 2, 367 (1873), in the index.—Rehder in *Bailey, Cycl. Am. Hort.* I. 121 (1900), in part.

This collective group is intended to include all the hybrids between the different species of American Azaleas (exclusive of *R. occidentale*) and *R. luteum* which cannot be traced to any of the binary hybrids but are apparently of more mixed and uncertain parentage. It comprises the garden forms known as Ghent Azaleas or *Azalea pontica* varieties, also sometimes called *Azalea Mortieriana*, though *Rhododendron Mortieri* as first figured and described by Sweet is apparently a binary hybrid between *R. calendulaceum* and *R. nudiflorum*. As it is desirable for practical reasons to have a definite name for this group of hybrids, I have adopted the name *gandavense*, which has been used in some nursery catalogues in the combination "*Azaleae gandavenses*" for this group of hybrids. It is one of those cultigen groups which as L. H. Bailey points out (in *Science*, XLVII. 306 [1918]) need names in the taxonomic treatment of cultivated plants, though such groups cannot be pinned down to a type and are more or less heterogenous and therefore not exactly in accordance with present taxonomic usage.

The following forms seem also to belong here:

"Azaleas de Jardins" Lemaire in *Ill. Hort.* II. pl. 75 (1855).

Sixteen different named varieties are figured, a flower of each; most of them are similar to the forms of *R. Mortieri*, some suggest the influence of *A. speciosum* and some of *R. luteum*.

"Azaleas rustiques de Gand" Van Houtte in *Rev. Hort. Belge*, III. 3 tab. (1877).

Two named varieties of the *R. Mortieri* type.

"Hardy Azaleas" Osborn in *Gard.* XV. 318 t. 176 (1879).

Four varieties, probably all representing hybrids of *R. Mortieri*.

***Rhododendron gandavense* var. *plenum* Rehder, nom. nov.**

Under this name all the double-flowered forms of *R. gandavense* may be classed. I refer here the following forms, of which the first may serve as a type of this group.

Azalea Van Houttei flore pleno Van Houtte in *Fl. des Serres*, XIII. t. 1298 (1858).

Raised from seed of a form of *R. Mortieri*. The flowers are semidouble with 5 to 10 stamens, and of pink color with the 3 or 4 upper lobes orange.

"Azaleas rustiques de Gand" Van Houtte in *Fl. des Serres*, XIX. t. 2019–2024 (1873), excl. t. 2019, 2020.

The plants were bought by Van Houtte from the widow and sons of Louis Hellebuyck, a contemporary of Mortier. Six forms are figured; four of them are double-flowered, two of them (t. 2019 and 2020) are single.

"Hardy double Azaleas" Moore in *Flor. & Pomol.* 1880, 89. t. 516.

Three forms are figured: "Narcissiflora," "Graf von Meran" and "Sappho." The two first named are in cultivation at the Arnold Arboretum. Of the form "Narcissiflora" *R. luteum* seems to be the dominant parent, while Graf von Meran, though similar, shows in its color and the less glandular pubescence the influence of *R. nudiflorum*.

Rhododendron molle × occidentale = R. albicans Waterer apud Zabel in *Mitteil. Deutsch. Dendr. Ges.* XI. 30 (1902). — For further citations see page 97.

The following references may be added to those given on pages 97 and 98.

Azalea "Waterer's New Hardy" Moore in *Flor. & Pomol.* 1883, 177, t. in part.

Three forms are figured, stated to have been raised by Anthony Waterer by crossing *A. occidentalis* with various hybrids. Two of them, "C. S. Sargent" with yellow flowers and "Henrietta Sargent" with bright rose-colored flowers, are probably hybrids with *R. Kosterianum* forms and may be referred to *R. albicans*, while "Rubra plena" is probably a hybrid of *R. occidentale* with one of the double-flowered *R. gandavense* and does not belong here.

"New Hybrid Azaleas" W. G. in *Gard.* XXIX. 550, t. 548 (1886), in part.

Two unnamed varieties are figured of which the white-flowered one apparently belongs here.

Azalea albicans [Waterer] apud *Gard.* XLV. 229, 322 (1894); apud *Jour. Roy. Hort. Soc.* XVII. XLVI (1895).

Described as a hybrid between *Azalea mollis* and *A. occidentalis* with white flowers just touched with yellow and with an orange suffusion on the upper lobes.

I am in favor of restricting the name *R. albicans* to the hybrid forms between *R. molle* (including perhaps × *R. Kosterianum*) and *R. occidentale*, for which it was originally intended, and to refer the last reference given on page 98 to *R. mixtum*. Of the varieties cited by Wilson, "Graciosa" and "Exquisite" belong certainly to *R. albicans*, of the other two I am not sure. The forms of *R. albicans* are characterized by large light-colored flowers with rather short tube, puberulous and scarcely glandular outside and with the orange blotch consisting of more or less separate dots and with not or slightly exerted stamens; from *R. molle* and *R. Kosterianum* they differ chiefly in the longer, more cylindrical tube and the generally pinkish white color of the flowers.

Rhododendron gandavense × occidentale.

Azalea "Waterer's New Hardy" Moore in *Flor. & Pomol.* 1883, 177, t.; as to var. "Rubra plena."

"New Hybrid Azaleas" W. G. in *Gard.* XXIX. 550, t. 548 (1886), as to the red-flowered variety.

To this group I am inclined to refer such forms as "Emelie" and "Roi des Belges." They are rather difficult to distinguish from the forms of *R. gandavense*, but may be recognized by the larger flowers with broad open limb, the comparatively short tube, the many-flowered dense inflorescence.

Rhododendron occidentale × (arborescens × japonicum).

Azalea occidentalis × *arborescens* × *mollis* Rehnelt in *Gartenwelt*, XX. 8, t. (1916).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Rhododendron calendulaceum × viscosum.

Such a cross is mentioned by Herbert (*Amaryll.* 360) in 1837: "I have raised seedlings from *Azalea Pontica-viscosa* var. *alba* and *Calendulacea-viscosa* var. *rubescens*." Nothing more, however, seems to have been published about the latter cross; *R. luteum* × *viscosum*, see on page 178.

Rhododendron molle × viscosum = Rhododendron viscosepalum
Rehder, comb. nov.

Azalea viscosepala R. A. in *Gard. Chron.* II. 348 (1842).

Azalea viscocephala Arb. Kew. in *Herb. Arnold Arb.*

In *Gardener's Chronicle* (*l. c.*) this is stated to have been raised by Mr. Waterer at Knap Hill and is described as "*Azalea viscosepala*, white with yellow throat. This is a cross between *viscosa* and *Indica alba* which latter it very much resembles in foliage."

I have before me specimens collected at Kew in 1880 as *Azalea viscocephala* and specimens from plants growing at the Arnold Arboretum which probably represent the same plant. They appear to be intermediate between *R. viscosum* var. *glaucum* and *R. molle*. From *R. viscosum* they differ chiefly in the earlier larger flowers with a yellow blotch, slightly yellowish in bud, with a stouter tube more dilated above the middle, in the shorter stamens not or slightly exceeding the lobes, in the more densely puberulous tube, in the tomentulose pedicels and calyx and in the thinly pubescent under side of the leaves. From *R. molle* they differ in the fragrant white flowers with a longer, more cylindrical tube glandular-pilose outside, in the glandular-pilose pedicels and in the smaller and broader usually elliptic leaves glaucous beneath and only slightly pubescent. There is not the slightest trace of *Azalea indica alba*, which is *R. mucronatum*, in the plant.

Rhododendron viscosepalum var. Daviesii Rehder, comb. nov.

Azalea Daviesii R. D. in *Gard. Chron.* ser. 2, XII. 200 (1879).¹

Azalea hybrida Daviesii Hort. apud Brettschneider in *Gartenfl.* XLII. 65, t. 1387 (1893).

This form differs from *R. viscosepalum* chiefly in the larger flowers with a more conspicuous yellow blotch and with a shorter, wider tube. The leaves are almost identical in shape, color and pubescence.

This hybrid was raised by Isaac Davies of Brook Lane Nursery, Ormskirk, and is stated to have been "produced from *A. sinensis* crossed with a hardy white *Azalea*" and the flowers are described as pale sulphur yellow, changing to white immediately they are fully expanded, and deliciously sweet-scented. A form "Avalanche" is mentioned, produced from the same cross and similar but with pure white smaller flowers. This must be very similar to *A. viscosepala*.

Very similar to var. *Daviesii* are the following forms:

Azalea altaclarensis Gowen apud Lindley in *Bot. Reg.* XXVIII. t. 27 (1842), not *Rhododendron altaclarensis* Lindl.

This handsome hybrid was raised at Highclere by J. R. Gowen, who states that "it was produced by fertilizing the flowers of *Azalea sinensis* with the pollen

¹ Not to be confused with *R. Daviesii* Dean, *Floral Mag.* n. ser. XX. t. 474 (1881), a hybrid of *R. javanicum* and *R. retusum*.

of the late flowering variety of *Azalea viscosa*, called by nurserymen *A. rubescens major*." The flowers are large white, slightly bordered pink, with a faint yellow tinge at the base of the lobes, and with the upper lobe distinctly dotted yellow and the tube pinkish outside. The gradually dilated tube and the yellow tinge and the spotted upper lobe indicate the influence of *R. sinense* and the white color and hirsute tube of the corolla and the delicious fragrance the influence of *R. viscosum*.

Azalea sinensis alba Van Houtte in *Fl. des Serres*, XX. 151. t. 2143-44 (1874).

Van Houtte gives no description and no hint of the origin of the plant save the remark that it is far removed from the true *A. sinensis*. It is possibly the same form of which a plant was received by the Arnold Arboretum from Koster and Son in 1904 under the name *Azalea chinensis alba grandiflora*.

Rhododendron ? gandavensis × molle.

Azalea sinensis var. *Bylsiana* Morren in *Ann. Soc. Agric. Bot. Gand*, I. 278, t. 27 (1845).

This form was raised by the Ghent horticulturist Byls from seeds of *A. sinensis*. It is apparently a hybrid and has large white flowers edged pink and the upper lobe with a large yellow blotch. This is not to be confused with *R. Bylsianum* Henderson, *Ill. Bouquet*, I. t. 18 (1857-59), a hybrid form of *Eurhododendron*.

Azalea sinensis var. *macrantha* Spae in *Ann. Soc. Agric. Bot. Gand*, III. 127, t. 122 (1847).

This form is of the same origin as the preceding; its flowers are white tinged with pale rose and marked with a large yellow blotch dotted orange. As a similar form is mentioned *A. sinensis* var. *versicolor* raised by Buyck-Vander Meersch.

Rhododendron japonicum × nudiflorum.

Azalea nudiflora × *sinensis* Van Fleet in *U. S. Dept. Agric. Bur. Pl. Indust.* CCXXIII. 17 (1911).

"Vigorous hybrids with profuse cream, rose and salmon colored blooms," raised by Dr. W. Van Fleet at Little Silver, N. J., and presented to the Plant Introduction Garden in Chico, California, in 1910. The parent species called *Azalea sinensis* is probably *R. japonicum*, for the former is not known to have been in cultivation in this country before 1910 except young seedlings at the Arnold Arboretum, raised in 1908.

Rhododendron japonicum × molle = Rhododendron Kosterianum Schneid. See page 97.

Rhododendron gandavense × Kosterianum = Rhododendron mixtum Wilson. See page 98.

Azalea rustica flore-pleno Vandevoorde in *Rev. Hort. Belge*, XIX. 232 t. (1893).

Azalea mollis fl. pleno "Virgil" Reiter in *Gartenwelt*, XV. 493, fig. (1911), habit figure.

Rhododendron occidentale × *molle* × "Ghent Azaleas" Millais, *Rhodod.* 220, fig. (habit) (1917).

Three forms are represented in the colored plate with pink, yellow and orange flowers. They are called double-flowered *A. mollis*, but are probably hybrids of *R. gandavense* and *R. Kosterianum*. I have seen neither plants nor specimens of these double-flowered forms.

Wilson has based the name *R. mixtum* on the double-flowered *Azalea rustica flore-pleno*, but we may include under this name all the different forms probably of the same parentage as given by Wilson.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

hairs; corolla deeply two-lipped, rosy-lilac, the upper lip spotted, slightly villose and short-pilose and sparingly glandular outside, tube funnel-form, 6 to 8 mm. long, sparingly pubescent inside, the two lower lobes separated nearly to the base of the limb, oblong, 2 to 2.5 cm. long and 1 to 1.2 cm. broad, obtuse or obtusish, upper lip as long and about 3 cm. broad, divided to about the middle into 3 ovate lobes; stamens 10, unequal, the longer about two thirds as long as corolla; the filaments pilose on the lower fourth; the anthers about 1 mm. long; style finely pilose at base, with green stigma; ovary setosely pilose, eglandular.

SECT. TSUTSUTSI

Rhododendron ? indicum × linearifolium = Rhododendron Eeckhautei Rehder in Bailey, *Stand. Cycl. Hort.* V. 2945 (1916).

Azalea Eeckhautei Pynaert in *Rev. Hort. Belge*, XXXVII. 117, t. (1911).

This very distinct hybrid was raised by Eeckhaute of St. Denis Westrem, Belgium, from “*Azalea linearifolia* fécondé par l’*A. indica* var. *Souvenir du Recteur Kickx*.” The shape of the corolla and of the leaves is similar to that of the mother plant, but the flowers are larger and bright rose-colored.

Rhododendron Simsii × obtusum = Rhododendron Sanderi Wilson. See page 43.

There are probably other hybrids in this section as *e. g.* *R. phoeniceum* (see page 61) which according to Wilson is possibly a hybrid between *R. mucronatum* G. Don and *R. scabrum* G. Don; and some of the varieties cultivated as Indian Azaleas are in all probability hybrids between *R. indicum* and *R. Simsii* (see page 25). *Rhododendron pulchrum* Sweet, which is considered by Sweet a hybrid between *R. indicum* and *R. ledifolium* (= *mucronatum*), is according to Wilson probably a form of *R. phoeniceum* (see page 62).

SUBGEN. ANTHODENDRON × EURHODODENDRON

SECT. PENTANTHERA × LEIORHODION

Rhododendro-Azalea Herbert, *Amaryllid.* 360 (1837).

Azaleodendron Rodigas in *Ill. Hort.* XXXIX. 53 (1892). — Masters in *Gard. Chron.* ser. 3, XIII. 665 (1893). — André in *Rev. Hort.* 1893, 369.

Rhodazalea Croux & fils in *Jour. Soc. Hort. France*, sér. 4, IX. 328 (1908). — Gérôme in *Jardin*, XXIV. 279 (1910).

The first hybrid known to have been raised in the genus *Rhododendron* is one between species of the two subgenera *Anthodendron* and *Eurhododendron*, which originated before 1800 in Thompson's Nursery at Mile End near London. Later other hybrids of similar origin were raised at Highclere in the gardens of the Earl of Carnarvon, by William Herbert at Spofforth, by Wm. Smith at Norbiton, who is particularly known for forms with yellow flowers produced by crossing *R. molle* with various evergreen *Rhododendrons*, by Isaac Davies of Brook Lane Nursery, Ormskirk, and others. Another race consisting of the hybrids between evergreen *Rhododendrons* and *R. japonicum* was raised by Vander Meulen of Ghent about 1890 and more recently by Croux & fils, Val d'Aulnay, at Châteney (Seine).

By those who keep *Azalea* and *Rhododendron* distinct new generic names have been coined. The oldest of them is *Rhododendro-Azalea* of Herbert, mentioned only incidentally by the author and not formally proposed. The second was proposed by Rodigas for the hybrids between *R. japonicum* and evergreen *Rhododendrons*. The third name was given for hybrids of similar origin, but *Rhodazalea Crouxii*, the type of this group, has only five stamens and thus differs from *Azaleodendron* which has ten stamens.

Rhododendron* ? *nudiflorum* × *ponticum* = *Rhododendron azaleoides Hort. Lond. apud Dumont de Courset, *Bot. Cult.* III. 327 (1811); VII. 165 (1814). — Desfontaines, *Tabl. Jard. Paris*, 272 (1815), as *R. azaloides*, name only.

Rhododendron ponticum var. *deciduum* Andrews, *Bot. Rep.* VI. t. 379 (1801).

Rhododendron ponticum β. *subdeciduum* "A. R." apud Sweet, *Hort. Brit.* 264 (1826).

Rhododendron ponticum b. *azaleoides* Loudon, *Arb. Brit.* II. 1131 (1838). — De Candolle, *Prodr.* VII. 722 (1839).

Rhododendron azaleoides inodorum André, *Pl. Terre Bruy.* 162 (1864).

According to Andrews this hybrid originated in Thompson's Nursery at Mile End near London and its supposed mother plant is *Rhododendron ponticum* fertilized by some *Azalea*. The leaves are rather large, oblong and acuminate, and the corolla is pale purple, not spotted or blotched, with the corolla-lobes very wavy at the margin; the stamens have purple anthers. It is likely that a form of *R. nudiflorum* is the male parent, as there is no indication of *R. viscosum* in the hybrid and these two species are the only *Azaleas* well known before 1800. The plant described by Dumont de Courset seems very similar, the flowers are about the same color and nothing is said about fragrance, while the plants described and figured by later authors as *R. azaleoides* differ in their usually white or nearly white very fragrant flowers and are apparently of different origin. The *R. azaleoides* of Sprengel (*Neu*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

Rhododendron ponticum azaleoides 2. *odoratum* Loudon, *Arb. Brit.* II. 1131 (1838).

Rhododendron odoratum Loddiges apud Steudel, *Nomencl.* ed. 2, II. 450 (1841).¹

Rhododendron azaleoides odoratum André, *Pl. Terr. Bruy.* 162 (1864).

Azalea odorata Hort. apud Goldring in *Gard. Chron.* ser. 3, VIII. 25 (1890). — H. P. & W. G. in *Gard.* XL. 34 (1891).

Rhododendron fragrans in *Jour. Hort.* ser. 3, XLIX. 489, fig. (1904)? not Paxton.

Rhododendron azaleoides Bean, *Trees & Shrubs Brit. Isl.* II. 344 (1914), not Dumont de Courset.

As stated by Ker this hybrid was "raised by Mr. Herbert at Spofforth near Weatherby, from seed ripened on the common white glaucous-leaved Azalea in the flower of which the pollen or dust from the anthers of *Rhododendron maximum* had been purposely substituted for that of its own flower." The evergreen leaves are oblong, acute at both ends and glaucous beneath; the flowers are rather small, whitish, the lobes with a pinkish border and the upper lobe spotted yellow, the tube cylindrical. I am referring to *R. hybridum*, though not without some doubt, the *R. azaleoides* as described by Mr. Bean, which according to his opinion possibly is a hybrid of *R. viscosum* and *R. maximum* and Mr. Bean's judgment can be depended upon. Also a specimen I collected nearly twenty years ago at Kew as *R. odoratum* agrees much better with Ker's figure than with that of *R. fragrans* Paxton, though the corolla-tube is not so cylindrical as represented in that figure.

***Rhododendron viscosum* × *ponticum* = *Rhododendron enneandrum* De Candolle, *Prodr.* VII. 728 (1839).**

Azalea hybrida enneandra Sims in *Bot. Mag.* XLIX. t. 2308 (1822).

Rhododendron hybridum enneandrum Herbert in *Trans. Hort. Soc.* III. ex Sims, *l. c.*

Rhododendron azaleoides Reider in *Ann. Blumist.* V. 84 (1829); VII. t. 11 (1831), the plate bearing the name *R. azaleoides alba odorata*.

Sims states that this is a hybrid raised by Wm. Herbert from seed of a white-flowered Azalea fertilized with the pollen of *R. ponticum*; the flowers are very fragrant, rather small, funnel-form, white, slightly tinged lilac, the upper lobe dotted green. Reider's *R. azaleoides* may belong here though the upper lobe is not dotted green.

***Rhododendron viscosum* × ? *catawbiense* = *Rhododendron fragrans* Paxton in *Paxton's Bot. Mag.* X. 147, t. (1843).²**

¹ A specimen collected by me at Kew Gardens in 1901 as *R. odoratum* apparently belongs here and certainly is the hybrid described by Bean as *R. azaleoides*, while another specimen in the herbarium of the Arnold Arboretum collected by H. Zabel in the Botanic Garden at Muenden, Hanover, from a plant received in 1882 as *R. odoratum* from T. Smith in Bergedorf near Hamburg, is not the same and similar to *R. azaleoides*.

² As *R. fragrans* Paxton must be considered a duly published name, it invalidates *R. fragrans* Maximowicz of 1870, for which I propose the following new name:

***Rhododendron Adamsii* Rehder, nom. nov.**

Azalea fragrans Adams in *Mém. Acad. Sci. St.-Pétersb.* II. 332, t. 14 (1808).

Azalea pallida Turczaninow in *Bull. Soc. Nat. Mosc.* 1838, 96 (1838), name only.

Paxton observed this hybrid in the nursery of Chandler & Sons, Vauxhall, and states that it was "raised accidentally from seed of a variety of *R. catawbiense* twenty-five or thirty years ago." He describes it as "a very compact dwarf shrub, decidedly evergreen with small and dense foliage and numerous clusters of pretty pale pinkish lilac blossoms in which there is a variety of delicate tints approaching to white in the center. It has quite the habit of a *Rhododendron* and looks like a small close-growing, pale-flowered variety of *R. ponticum*." Herbert had apparently this hybrid in mind when he says (*Amaryll.* 360): "The fragrance of the Azaleas has been communicated to the *Rhododendrons* both by the Mitcham crosses and that of Mr. Chandler." To the Mitcham crosses, however, I have found no other references.

Here may belong:

Rhododendron elegantissimum Davies apud R. D. in *Gard. Chron.* ser. 2, XII. 200 (1879).

"Early White Azalea" × *R. Hendersonii* R. D., *l. c.*

This is said to have white flowers tipped with deep lilac and of agreeable fragrance. I assume that under Early White Azalea an early flowering form of *R. viscosum* is understood, as the flowers are very fragrant and *R. Hendersonii*, a name for which I can find no other reference except Millais, *Rhodod.* 38, and which is possibly one of the *R. catawbiense* hybrids.

Rhododendron roseum odoratum Davies apud R. D. in *Gard. Chron.* ser. 2, XII. 200 (1879). — Vilmorin & Bois, *Frut. Vilmor.* 175 (1904).

"Early White Azalea" × *R. Brayanum* R. D., *l. c.*

This is described as similar to the preceding, but with rose-colored flowers. *Rhododendron Brayanum* is mentioned by Millais, *Rhodod.* 35.

***Rhododendron* ? *calendulaceum* × ? *catawbiense*.**

Rhododendron ponticum var. *azaleoides* Torloniana Morren in *Ann. Soc. Agric. Bot. Gand*, I. 21, t. 4 (1845).

? *Rhododendron Tolerianum* hort. apud Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz-Ben.* 379 (1903), name only.

Nothing is said of the origin or parentage of this hybrid form. The flowers much resemble in size, shape and color the plant figured by Guimpel as *R. azaleoides* mentioned below, but the upper lobe shows a large orange blotch while Guimpel's figure does not show the slightest trace of orange or yellow; this and the shape of the corolla seem to indicate the influence of *R. calendulaceum*.

I am not aware that this plant is at present in cultivation, though it is certainly one of the handsomest of this group of hybrids.

Rhododendron azaleoides Guimpel, Otto & Hayne, *Abb. Fremd. Holzgew.* 15, t. 15 (1825), not Dumont de Courset.

Osmothamnus fragrans De Candolle, *Prodr.* VII. 715 (1839).

Osmothamnus pallidus De Candolle, *l. c.* (1839).

Rhododendron fragrans Maximowicz in *Mém. Acad. Sci. St.-Pétersb.* XVI. No. 9, 16 (1870).

Rhododendron pallidum Dümmer in *Gard. Chron.* ser. 3, LIII. 264 (April 26, 1913), not *R. pallidum* W. Watson in *Gard. Chron.* *l. c.* 230 (April 12, 1913).

Dümmer seems to consider *Azalea pallida* Turcz. as specifically distinct, but even if we unite it with *Azalea fragrans* the combination *R. pallidum* cannot be used in place of *R. fragrans*, as it is antedated by W. Watson's name which, though insufficiently described, should stand as the name of the hybrid for which he intended it. (See also C. P. Raffle's remarks in *Gard. Chron.* *l. c.* 332.)

If the colored plate is a true representative of the plant it is certainly, with the form just mentioned, one of the handsomest of this group of hybrids on account of its large flowers and the pure pink color without any trace of the lilac or purple found in most related forms. The leaves are almost acuminate, to 8 cm. long and green on both sides; the flowers are broadly funnelform-campanulate, about 6 cm. wide, with broad lobes wavy at the margin, white and more or less tinged and striped pink. Nothing is said of the origin of the plant, but it was apparently English.

The following is a hybrid probably of similar origin :

Azalea (var.) *magnifica* (Rollisson's) Van Houtte in *Fl. des Serr.* XIII, 31, t. 1306-7 (1858). — Lemaire in *Ill. Hort.* VI. t. 209 (1859).

Nothing is said about the origin of this plant except that it was received from W. Rollisson. According to the plate it is a very handsome plant with large white flowers slightly tinged pink, the upper lobe with a large orange blotch and the other lobes marked with pink veins near the margin; the number of stamens is 6 to 8.

Rhododendron gandavense × ? **catawbiense**.

Rhododendron "Deception" Standish and Noble in *Jour. R. Hort. Soc.* V. 274 (1850). — Burbidge, *Cult. Pl.* 298 (1876).

Described as having pink spotted flowers and fine foliage and stated to be a hybrid between a "large yellow Ghent Azalea and *R. catawbiense*."

Rhododendron discolor × **occidentale** Bean in *Rhodod. Soc. Notes*, I. 191 (1919), name only.

Flowers not yet known.

Rhododendron luteum × **ponticum** = **Rhododendron Laetitia** Herbert apud Lindley in *Bot. Reg.* XXXI. t. 51 (1845).

This hybrid was raised by W. Herbert of Spofforth from seed of *Rhododendron ponticum* fertilized by *Azalea pontica*. The leaves are described as "rather more durable, broader and blunter than the leaves of *Azalea Pontica*"; the flowers are large, white and fragrant, the three upper lobes with large orange blotches and smaller dots. The plant is named after Mrs. W. Herbert. This is apparently one of the seedlings mentioned in 1837 by Herbert in his *Amaryllidaceae*, page 359: "Four evergreen seedlings obtained from the seed of *Rhododendron Ponticum* which I had fertilized with pollen of *Azalea Pontica* have flowered at Highclere."

Of similar origin ("*Azalea pontica* × evergreen *Rhododendron*") were the hybrids raised by Adam of Altenburg, and offered to the trade by Rinz of Frankfort, Germany, probably about 1850; eleven varieties are enumerated by André (*Pl. Terre Bruy.* 164 [1864]).

Rhododendron speciosum × **arboreum**.

Rhododendron nudiflorum var. *eximium* Sweet, *Brit. Flow. Gard.* ser. 2, III. t. 291 (1835).

Rhododendron eximium D. Don apud Zabel in Beissner, Schelle & Zabel, *Handb. Laubholz-Ben.* 379 (1903), not Nuttall.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

should be discarded, as it is not only against all rules of nomenclature but also misleading and confusing.

Rhododendron norbitonense var. **broughtonianum** Rehder, comb. nov.

Rhododendron broughtonianum André, *Pl. Terr. Bruy.* 166 (1864); in *Rev. Hort.* 1893, 370.

Rhododendron broughtonii aureum Hort. apud W. Watson, *Rhodod. & Azaleas*, 93 (1911). — For further literature and remarks, see page 99.

I have little doubt that *R. broughtonianum* and *R. broughtonii aureum* are identical, and though the plant is well known under the latter name, its use should be discontinued, as it is, like *R. Smithii aureum*, both incorrect and misleading, since there is already a *R. broughtonii* Hort. which is a hybrid between two species or forms of *Eurhododendron* with rosy-red flowers; moreover, the name *broughtonii* suggests the proper name of a person and not of a place. A good habit picture is given by Millais in his *Rhododendrons* (pl. opposite page 184).

Other forms of similar parentage were raised by Standish and Noble and later by Anthony Waterer and by Mr. White of Sunningdale. The following also belongs here:

Rhododendron Williamsii Millais, *Rhodod.* 262 (1917).

A hybrid raised by Williams of Holloway about 1885 between a hardy *Rhododendron* and *R. sinense*.

Rhododendron japonicum × **Eurhododendron** “hybrids” = **Rhododendron azaleodendron** Vilmorin & Bois, *Frut. Vilmor.* 175 (1904).

Azaleodendron Rodigas in *Ill. Hort.* XXXIX. 53 (1893). — (See also page 136).

Azaleodendron “Comte de Kerchove,” De Bosschere in *Rev. Hort. Belge*, XIX. 177, t. (1893).

Here belong the hybrids raised by G. Vander Meulen between “*Azalea mollis*” (= *R. japonicum*) and *R.* “Prince Camille de Rohan” and “Leopard.” As the type of this group of hybrids “Comte de Kerchove” may be taken; this form is described in detail by Masters (in *Gard. Chron.* ser. 3, XIII. 667 [1893]) and figured in *Gard. Chron.* l. c. 663 and also in *Rev. Hort. Belge*, l. c. (colored pl.), and in *Rev. Hort.* 1893, 369. Besides “Comte de Kerchove” five other varieties were exhibited at Ghent in 1892 by Pynaert and Van Gaert who acquired the entire stock of Vander Meulen’s hybrids; their color ranges between white and rosy-red or rose-lilac with conspicuous dark spots on the upper lobes. According to André (in *Rev. Hort.* 1893, 371) similar hybrids between “*Rhododendron* and *Azalea mollis*” had been shown in Ghent by Ad. Rosseel in 1888.

Rhododendron? **Kosterianum** × **Eurhododendron** hybrids = **Rhododendron Crouxii** Rehder, comb. nov.

Rhodazalea Crouxii Croux & fils in *Jour. Soc. Hort. France*, sér. 4, IX. 328, fig. (1908). — Gérôme in *Jardin*, XXIV. 279, fig. 157 (1910).

This hybrid was obtained by Croux & fils of Val d’Aulnay at Châtenay (Seine), according to their statement, by crossing a *Rhododendron* with a variety of *Azalea mollis*, the exact name of neither of the parents being given. The flowers are large,

their color is Naples yellow tinged with rose, the upper lobes conspicuously spotted. The number of stamens is 5, which distinguishes this hybrid from the similar *R. azaleodendron*.

Rhododendron Crouxii f. plenum Rehder, nom. nov.

Rhodazalea Crouxii flore pleno, Croux & fils in *Jour. Soc. Hort. France*, sér. 4, IX. 328 (1908). — Gérôme in *Jardin*, XXIV. 279, fig. 158, 159 (1910).

Azalea mollis × *Rhododendron* "Marie Van Houtte" Croux & fils, *l. c.*

This form has semi-double yellow flowers tinged with rose.

SECT. TSUTSUTSI × LEIORHODION

The following forms have been supposed to be hybrids between the sections Tsutsutsi and Leiorhodium.

Rhododendron azaleoides var. *crispiflorum* Lemaire in *Ill. Hort.* V. t. 181 (1858).¹

This handsome and interesting hybrid was raised, according to Lemaire, by a Ghent horticulturist, Louis Demotte, by crossing a hybrid *Rhododendron* with an Indian Azalea, probably the var. "*Reyndersiana*." Judging from the figure and the characters given in the description this plant seems indeed to be a hybrid of a species or form of the section Tsutsutsi and a form of *Eurhododendron*, the only one known to me, for the two following forms which are supposed to be hybrids of an Indian Azalea and an evergreen *Rhododendron* are apparently only forms of *R. phoeniceum*.

Azalea Rawsonii Paxton in *Paxton's Flow. Gard.* III. 123 t. (1836).

This Azalea was raised by John Menzies, gardener to Christopher Rawson of Hope House near Halifax. In a letter Mr. Menzies "states as his belief that it is produced between *A. phoenica* and *Rhododendron dauricum atrovirens*." Judging from the colored plate it is a simple form of *R. phoeniceum*. See also pages 61 and 62.

Rhododendron phoeniceum var. *splendens* Sweet, *Brit. Flow. Gard.* ser. 2, IV. t. 385 (1837).

This is said to have been "raised by Mr. Wood, gardener to Mrs. May at Sydenham, from a plant of *Rhododendron phoeniceum* that had been fecundated by *R. catawbiense*." It appears to be only a form of *R. phoeniceum*; not the slightest influence of *R. catawbiense* is noticeable in the plant figured. See also page 63.

¹ Not *R. indicum* var. *crispiflorum* Schneid.—See page 27.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

AZALEA CALENDULACEA η . TRIUMPHANS Sweet, *l. c.* 265 (1826), name.

Rhododendron calendulaceum η . *triumphans* Sweet, *l. c.* 343 (1830), "copper-yellow."

AZALEA FASCICULATA Dumont de Courset, *Bot. Cult.* VII. 166 (1814).

Azalea fastigiata Dumont de Courset, *l. c.* (1814).

Described as having the flowers fascicled in a kind of an umbel. Probably the same as *Azalea nudiflora* λ . *fastigiata* Sweet (see below).

AZALEA LUCIDA Meerburgh, *Pl. Selec. Icon.* t. 7 (1798).

This species is described as with "foliis lanceolatis, flore luteo piloso, staminibus brevioribus. Cl. Linnaeus hanc descripsit pro *Azalea nudiflora*." The illustration resembles in general appearance *R. nudiflorum*, but the leaves are narrow-lanceolate, and the flowers have more the shape of those of *R. viscosum* and are colored light yellow. It possibly represents *R. viscosum* var. *nitidum*, and the yellowish tinge of the flowers may be a mistake of the artist.

AZALEA NUDIFLORA AMOENA Loddiges Cat. 1836, apud Loudon, *Arb. Brit.* II. 1142 (1838), name only.

AZALEA NUDIFLORA γ . BLANDA Sweet, *Hort. Brit.* 265 (1826), name. — Loddiges, *l. c.*

Rhododendron nudiflorum γ . *blandum* Sweet, *Hort. Brit.* ed. 2, 343 (1830), "blush."

AZALEA NUDIFLORA β . CALYCOSA Wood, *Classb. Bot.* 490 (1870).

This is a form from Georgia described as having one calyx-lobe subulate and three or four times longer than the others.

AZALEA NUDIFLORA ϵ . CAROLINIANA Sweet, *Hort. Brit.* 265 (1826). — Loddiges Cat. 1836 apud Loudon, *Arb. Brit.* II. 1142 (1838), name only.

? *Azalea caroliniana* in *Cat. Am. Trees & Shrubs in John Bartram's Garden* (179?), name only.

Rhododendron nudiflorum ϵ . *carolinianum* Sweet, *Hort. Brit.* ed. 2, 343 (1830).

In Bartram's Catalogue it is called "Sweet Honeysuckle," which would suggest that it is one of the fragrant species.

AZALEA NUDIFLORA COBURGIA Loddiges, *l. c.* 1142 (1838).

AZALEA NUDIFLORA COLORATA Loddiges, *l. c.* 1142 (1838).

AZALEA NUDIFLORA CONSPICUA Loddiges, *l. c.* 1142 (1838), name.

AZALEA NUDIFLORA η . CORYMBOSA Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum η . *corymbosum* Sweet, *l. c.* 343 (1830), "white."

AZALEA NUDIFLORA θ . CRISPA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum θ . *crispum* Sweet, *l. c.* 343 (1830), "blush."

AZALEA NUDIFLORA ι . CUMULATA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838), as var. *cumula*.

Rhododendron nudiflorum ι . *cumulatum* Sweet, *l. c.* 344 (1830), "white."

AZALEA NUDIFLORA κ . DISCOLOR Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum κ . *discolor* Sweet, *l. c.* 344 (1830), "red, white."

AZALEA NUDIFLORA λ . FASTIGIATA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum λ . *fastigiatum* Sweet, *l. c.* 344 (1830), "blush."

AZALEA NUDIFLORA μ . FLORIDA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum μ . *floridum* Sweet, *l. c.* 344 (1830), "blush."

AZALEA NUDIFLORA ν . GLOBOSA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum ν . *globosum* Sweet, *l. c.* 344 (1830) "blush."

AZALEA NUDIFLORA ξ . GLOMERATA Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum ξ . *glomeratum* Sweet, *l. c.* 344 (1830) "blush."

AZALEA NUDIFLORA GRANDIFLORA Loddiges, *l. c.* 1142 (1838).

AZALEA NUDIFLORA \omicron . INCANA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* (1838).

Rhododendron nudiflorum \omicron . *incanum* Sweet, *l. c.* 344 (1830) "white."

AZALEA NUDIFLORA π . INCARNATA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum π . *incarnatum* Sweet, *l. c.* 344 (1830) "flesh-colored."

AZALEA NUDIFLORA ρ . MIRABILIS Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum ρ . *mirabile* Sweet, *l. c.* 344 (1830), "red."

AZALEA NUDIFLORA ϵ . ~~MONTANA~~ De Candolle, *Prodr.* VII. ? 717 (1839).

Described as having pale rose-colored flowers with the lower lobe inside yellowish and with a small calyx.

AZALEA NUDIFLORA σ . MONTANA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum σ . *montanum* Sweet, *l. c.* 265 (1830), "flesh-colored."

AZALEA NUDIFLORA OCHROLEUCA Loddiges, *l. c.* 1142 (1838), name.

AZALEA NUDIFLORA τ . PALLIDA Sweet, *l. c.* 265 (1826).

Rhododendron nudiflorum τ . *pallidum* Sweet, *l. c.* 344 (1826), "lilac white."

AZALEA NUDIFLORA υ . PALUDOSA Sweet, *l. c.* 265 (1826), "lilac white." — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum υ . *paludosum* Sweet, *l. c.* 344 (1830), as var. *pallidosa*.

AZALEA NUDIFLORA ζ . PAPILIONACEA Aiton, *Hort. Kew.* I. 203 (1789). — De Candolle, *Prodr.* VII. 717 (1838).

Azalea papilionacea Dumont de Courset, *Bot. Cult.* ed. 2, III. 333 (1811), as var. of *A. nudiflora*.

Azalea periclymenoides var. ϵ . *papilionacea* Pursh, *Fl. Am. Sept.* 152 (1814).

Rhododendron nudiflorum ϕ . *papilionaceum* Sweet, *Hort. Brit.* ed. 2, 344 (1830). —

G. Don, *Gen. Syst.* III. 847 (1834).

This form is described by Aiton as having pale reddish flowers with the lower lobe white, and a foliaceous calyx.

AZALEA NUDIFLORA η . PARTITA Aiton, *Hort. Kew.* I. 203 (1789). — De Candolle, *Prodr.* VII. 717 (1838).

Azalea periclymenoides ζ . *partita* Pursh, *Fl. Am. Sept.* 152 (1814).

Rhododendron nudiflorum χ . *partitum* Sweet, *Hort. Brit.* ed. 2, 344 (1830). —

G. Don, *Gen. Syst.* III. 847 (1834).

This form is described by Aiton as having pink flowers 5-parted to the base. This is apparently an abnormal form corresponding to *Kalmia latifolia* f. *monstrosa*. It seems to have soon disappeared from cultivation.

AZALEA NUDIFLORA ψ . PARVIFLORA Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum ψ . *parviflorum* Sweet, *l. c.* 344 (1830), "blush."

AZALEA NUDIFLORA PERICLYMENOIDES Loddiges, *l. c.* (1838), name.

AZALEA NUDIFLORA λ . POLYANDRA De Candolle, *Prodr.* VII. 717 (1838). — Wood, *Classb. Bot.* 490 (1870).

Azalea periclymenoides η . *polyandra* Pursh, *Fl. Am. Sept.* 152 (1814).

Rhododendron nudiflorum η. *polyandrum* G. Don, *Gen. Syst.* III. 847 (1834).

? *Azalea octandra* Michaux, *Journal*, ed. C. S. Sargent, 91 (in *Proc. Am. Philos. Soc.* (1889)), name only.

This form is described by Pursh as having short rose-colored flowers with 10 to 20 stamens and is said to have been found near Philadelphia. Michaux collected his *Azalea octandra* in or in the neighborhood of Perry County, Pennsylvania.

AZALEA NUDIFLORA ω. PROLIFERA Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum ω. *proliferum* Sweet, *l. c.* 344 (1830), "blush."

AZALEA NUDIFLORA αα. PUMILA Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum αα. *pumilum* Sweet, *l. c.* 344 (1830), "blush."

AZALEA NUDIFLORA ββ. PURPURASCENS Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum ββ. *purpurascens* Sweet, *l. c.* 344 (1830), "flesh-colored."

AZALEA NUDIFLORA γγ. PURPUREA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* (1838).

Rhododendron nudiflorum γγ. *purpureum* Sweet, *l. c.* 344 (1830), "purple."

AZALEA NUDIFLORA ζζ. RUBERRIMA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum ζζ. *ruberrimum* Sweet, *l. c.* 344 (1830), "red."

AZALEA NUDIFLORA ηη. RUBESCENS, Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum ηη. *rubescens* Sweet, *l. c.* 344 (1830), "pink."

AZALEA NUDIFLORA λλ. RUFΑ Sweet, *l. c.* 265 (1826), name. — Loddiges *l. c.* 1142 (1838).

Rhododendron nudiflorum λλ. *rufum* Sweet, *l. c.* 344 (1830), "red."

AZALEA NUDIFLORA νν. SEMI-DUPLEX Sweet, *l. c.* 265 (1826), name.

Rhododendron nudiflorum νν. *semiduplex* Sweet, *l. c.* 344 (1830), "white."

AZALEA NUDIFLORA β. RUTILANS Aiton, *Hort. Kew.* I. 203 (1789). — De Candolle, *Prodr.* VII. 717 (1838).

Azalea rubicunda Dumont de Courset, *Bot. Cult.* ed. 2, III. 332 (1811), as var. of *A. nudiflora*.

Azalea periclymenoides β. *rutilans* Pursh, *Fl. Am. Sept.* 152 (1814).

Rhododendron nudiflorum μμ. *rutilans* Sweet, *Hort. Brit.* ed. 2, 343 (1830). — G. Don, *Gen. Syst.* III. 847 (1834).

Azalea nudiflora rubicunda Loddiges *Cat.* 1836 apud Loudon, *Arb. Brit.* II. 1142 (1838).

Azalea nudiflora θθ. *rubicunda* Sweet, *Hort. Brit.* 265 (1826).

Rhododendron nudiflorum θθ. *rubicundum* Sweet, *Hort. Brit.* ed. 2, 343 (1830).

This form is described by Aiton as having deep red flowers with small calyx. Dumont de Courset says that the young branchlets and the leaves are pubescent, the calyx greenish brown and the winter-buds pale yellow in autumn.

AZALEA NUDIFLORA SEROTINA Loddiges, *l. c.* 1142 (1838), name.

AZALEA NUDIFLORA ξξ. STAMINEA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* 1142 (1838).

Rhododendron nudiflorum ξξ. *stamineum* Sweet, *l. c.* 344 (1830), "white."

AZALEA NUDIFLORA οο. STELLATA Sweet, *l. c.* 265 (1826), name. — Loddiges, *l. c.* (1838).

Rhododendron nudiflorum οο. *stellatum* Sweet, *l. c.* 344 (1830), "white."

AZALEA NUDIFLORA VARIA Loddiges, *l. c.* 1142 (1838), name.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

AZALEA SPECIOSA κ . UNDULATA Sweet, *l. c.* 265 (1826), name.

Rhododendron speciosum κ . *undulatum* Sweet, *l. c.* 343 (1830), "red."

AZALEA VISCOSA β . CRISPA Sweet, *Hort. Brit.* 266 (1826). — Loddiges, *Cat. apud Loudon, Arb. Brit.* II. 1143 (1838).

Rhododendron viscosum β . *crispum* Sweet, *Hort. Brit.* ed. 2, 344 (1830), "white."

AZALEA VISCOSA γ . DEALBATA Sweet, *l. c.* 266 (1826). — Loddiges, *l. c.* 1143 (1838).

Rhododendron viscosum γ . *dealbatum* Sweet, *l. c.* 344 (1830), "white."

AZALEA VISCOSA ϵ . PENICILLATA Sweet, *l. c.* 266 (1826). — Loddiges, *l. c.* 1143 (1838).

Rhododendron viscosum ϵ . *penicillatum* Sweet, *l. c.* 344 (1830), "red, white."

AZALEA VISCOSA PRAECOX Loddiges, *l. c.* 1143 (1838), name.

AZALEA VISCOSA STRICTA Carrière in *Rev. Hort.* 1872, p. 299.

Described as a very dwarf compact plant, with oval leaves glaucous beneath and small white slightly pinkish flowers appearing at the beginning of May. The early flowering time makes it improbable that this form belongs to *R. viscosum*.

AZALEA VISCOSA ζ . VARIEGATA Sweet, *l. c.* 265 (1826). — Loddiges, *l. c.* 1143 (1838).

Rhododendron viscosum ζ . *variegatum* Sweet, *l. c.* 344 (1830), "red, white."

AZALEA VISCOSA VIOLAE ODORE Loddiges, *l. c.* 1143 (1838), name.

Besides the doubtful names enumerated above, a large number of names without or with very inadequate description have appeared in horticultural publications taken in about all cases from various nursery catalogues. The following are some of the more important references to lists of garden forms of Azaleas (exclusive of Indian Azaleas):

Loudon, *Arb. Brit.* II. 1142–1144 (1838).

Bosse, *Vollst. Handb. Blumengärt.* I. 337, 340–342 (1840).

Heynhold, *Nomencl. Bot. Hort.* 108 (1840); II. 55–56 (1846).

Steudel, *Nomencl.* I. 175 (1841).

Henderson, *Ill. Bouquet*, text to t. 46 (1859–61), forms of *R. norbitonense*.

André, *Pl. Terr. Bruy.* 164, 166, 187–189 (1864).

Millais, *Rhodod.* 166–167, 212–214 (1917).

ILLUSTRATIONS

References to illustrations of garden forms (exclusive of Indian Azaleas) not mentioned elsewhere in this publication. They are mostly forms of *R. gandavense* or *R. Mortieri* if not stated otherwise.

ANTHONY KOSTER. *Gard. Chron.* ser. 3, XIII. 513, fig. 76 (1893). — Watson, *Rhodod. & Azaleas*, frontisp. (1912) = *R. Kosterianum*.

ATORRUBENS NOVA. Lemaire in *Ill. Hort.* II. pl. 75, fig. 15 (1855).

BARON G. PYCKE. Lemaire, *l. c.* fig. 16 (1855).

BEAUTÉ DE FLANDRE. Lemaire, *l. c.* fig. 9 (1855).

BIJOU DE GENDBRUGGE. Van Houtte in *Fl. des Serr.* XIX. t. 2024 (928) (1873) = *R. gandavense plenum*.

BONNEUR DE LA BELGIQUE. Lemaire in *Ill. Hort.* II. pl. 75, fig. 4 (1855).

BOUQUET DE FLORE. Ottolander, *Nederl. Fl. Pom.* 53, t. 16 (1876).

CARDINAL. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 9 (1846).

COCCINEA SPECIOSA. Ottolander, *Nederl. Fl. Pom.* 5, t. 15, fig. 2 (1876).

COMTE DE GOMER. *Rev. Hort. Belge*, I. 97, t. 6 (1875). — Dean, *Floral Mag.* n. ser. XVIII. t. 367 (1879) = *R. japonicum* var.

COMTESSE DE KERCHOVE. André in *Ill. Hort.* XXV. 59, t. 311 (1878) = *R. japonicum* var.

CUPREA EXIMIA NOVA. Osborn in *Gard.* XV. 318, t. 176 (1879).

CZARINE. Pynaert & Van Geert in *Rev. Hort. Belg.* XX. 277, t. (1894) = *R. Kosterianum*.

DELICATA NOVA. Lemaire in *Ill. Hort.* II. pl. 75, fig. 10 (1855).

DUC D'URSEL. Lemaire, *l. c.* fig. 1 (1855).

ÉTENDARD. Morren in *Ann. Soc. Agric. Bot. Gand*, III. 9, t. 106, fig. 14 (1847).

EUGÉNIE. Lemaire in *Ill. Hort.* II. pl. 75, fig. 6 (1855).

EUGÉNIE. Ottolander, *Nederl. Fl. Pom.* 51, t. 15, fig. 3 (1876), not the same as the preceding variety.

FLORENTINA. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 6 (1846).

FRANÇOIS DE TAYE. Van Houtte in *Fl. des Serr.* XIX. t. 2023 (927) (1873) = *R. gandavense plenum*.

GÉANT DES BATAILLES. Ottolander, *Nederl. Fl. Pom.* 51, t. 15, fig. 1 (1876).

GLOIRE DE VERSCHAFFELT. Morren in *Ann. Soc. Agric. Bot. Gand*, III. 9, t. 106, fig. 16 (1847).

GRAF VON MERAN. Rinz in *Gartenfl.* III. 171, t. 88 (1854). — Moore in *Fl. Pomol.* 1880, 89, t. 516, fig. 2, = *R. gandavense plenum*.

GRAND-DUC. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 7 (1846).

IGNAEA NOVA. Ottolander, *Nederl. Fl. Pom.* 51, t. 15, fig. 5 (1876).

JAS. H. LAING. Pynaert & Van Geert, *Rev. Hort. Belg.* XVII. 121, t. (1891) = *R. Kosterianum*.

- LOUIS AIMÉ VAN HOUTTE. Van Houtte in *Fl. des Serr.* XIX. t. 2022 (929) (1873) = *R. gandavense plenum*.
- LOUIS HELLEBUYCK. Van Houtte in *Fl. des Serr.* XIX. t. 2019 (931) (1873).
- MADAME ALEX. HARDY. Van Houtte in *Fl. des Serr.* XIX. t. 2020 (930) (1873).
- MAGNIFICA ALBICANS. Lemaire in *Ill. Hort.* II. pl. 75, fig. 14 (1855).
- MARIE-DOROTHÉE. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 5 (1846).
- MINA VAN HOUTTE. Van Houtte in *Fl. des Serr.* XIX. t. 2021 (926) (1873) = *R. gandavense plenum*.
- MINERVE. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 10 (1846).
- MINIATA FLORIBUNDA. Lemaire in *Ill. Hort.* II. pl. 75, fig. 13 (1855).
- MOLLIS VAR. ALBICANS. André in *Ill. Hort.* XXV. 59, t. 311 (1878) = *R. japonicum* var.
- MOLLIS VAR. AUREA FLORIBUNDA. André in *Ill. Hort.* XXV. 91, t. 317 (1878) = *R. japonicum* var.
- MURILLO. Vandevoorde in *Rev. Hort. Belg.* XIX, 232, t. fig. 1 (1893) = *R. mixtum*.
- NARCISSIFLORA. Moore in *Fl. Pomol.* 1880, 89, t. 516, fig. 1 (1880) = *R. gandavense plenum*.
- NE PLUS ULTRA. Osborn in *Gard.* XV. 318, t. 176 (1879).
- NÉRON. Lemaire in *Ill. Hort.* II. t. 75, fig. 7 (1855).
- NICOLAS II. Pynaert & Van Geert in *Rev. Hort. Belge*, XX. 277, t. (1894) = *R. Kosterianum*.
- NOBILIS. Lemaire in *Ill. Hort.* II. pl. 75, fig. 12 (1855).
- ORANGE PEINTE. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 2 (1846).
- OSCAR PREMIER. Morren in *Ann. Soc. Agric. Bot. Gand*, III. 9, t. 106, fig. 13 (1847).
- PERLE DU PRINTEMPS. Morren in *Ann. Soc. Agric. Bot. Gand*, II. 9, t. 106, fig. 19 (1847).
- PHÉBÉ. Vandervoorde in *Rev. Hort. Belg.* XIX. 232, t. fig. 3 (1893) = *R. mixtum*.
- PONTICA MACRANTHA. Osborn in *Gard.* XV. 318, t. 176 (1879).
- PONTICA VERSICOLOR. Osborn in *Gard.* XV. 318, t. 176 (1879).
- PRINCE HENRI DES PAYS-BAS. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 1 (1846).
- PRINCE TROUBETZKOY. André in *Ill. Hort.* XXV. 91, t. 317 (1878) = *R. japonicum* var.
- QUADRICOLORE. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 8 (1846).
- QUEEN VICTORIA. Van Hulle in *Rev. Hort. Belge*, III. 4, t. fig. 1 (1877).
- RAYON DU MATIN. Morren in *Ann. Soc. Agric. Bot. Gand*, II. 9, t. 106, fig. 18 (1847).
- REINE D'ANGLETERRE. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 4 (1846).
- RICHARDII. Lemaire in *Ill. Hort.* II. pl. 75, fig. 11 (1855).
- ROSALIE. Morren in *Ann. Soc. Agric. Bot. Gand*, III. 9, t. 106, fig. 15 (1847).
- ROSEA FORMOSISSIMA. Lemaire in *Ill. Hort.* II. pl. 75, fig. 3 (1855).
- ROSEA LINEATA. Lemaire, *l. c.* fig. 8 (1855). — Ottolander, *Nederl. Fl. Pom.* 51, t. 15, fig. 4 (1876).
- ROSEA ROTUNDIFLORA. Lemaire, *l. c.* fig. 5 (1855).
- RUBENS. Spae in *Ann. Soc. Agric. Bot. Gand*, II. 326, t. 81, fig. 12 (1846).
- RUBRA SPLENDIDISSIMA. Lemaire in *Ill. Hort.* II. pl. 75, fig. 2 (1855).

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

ADDITIONS

- P. 29. Add to the synonyms of **R. obtusum**:
Rhododendron amoenum var. *obtusum* Zabel in *Mitteil. Deutsch. Dendr. Ges.* XI. 35 (1902).
- P. 32. Add to the synonyms of **R. obtusum f. amoenum**:
Rhododendron amoenum var. *euamoenum* Zabel in *Mitteil. Deutsch. Dendr. Ges.* XI. 35 (1902).
Rhododendron amoenum subvar. *normale* Zabel, *l. c.* (1902).
Rhododendron amoenum subvar. *genuinum* Zabel, *l. c.* (1902).
Rhododendron amoenum subvar. *Buergeri* Zabel, *l. c.* (1902).
- P. 33. Add to the synonyms of **R. obtusum f. japonicum**:
Rhododendron amoenum subvar. *japonicum*, Zabel in *Mitteil. Deutsch. Dendr. Ges.* XI. 35 (1902).
- P. 105. Add as a doubtful name:
Rhododendron boninense Nakai in *Tokyo Bot. Mag.* XXXIV. (324) (1920), name only.

INDEX

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

- Azalea* Comte Papadopoli, 102
 — Comtesse de Kerchove, 203
 — Consul Pécher, 102
 — *crispiflora*, 27
 — *Cuprea eximia nova*, 203
 — Czarine, 203
 — Damask Rose, 72
 — *Danielsiana*, 23
 — *Daviesii*, 182
 — Decora, 44
 — *decumbens*, 23
 — *Delicata nova*, 203
 — *dianthiflora*, 76
 — *dilatata*, 84
 — Director Rodigas, 99
 — Double Blush, 138
 — Dr. Masters, 99
 — Duc de Nassau, 44
 — Duc d'Ursel, 203
 — Ebenezer Pyke, 97
 — Édouard André, 99
 — *Eeckhautei*, 187
 — *erecta foliis ovatis integris alternis flore luteo piloso praecoci*, 128
 — Étendard, 203
 — Eugénie, 203
 — Exquisite, 98, 180
 — *fasciculata*, 198
 — *fastigiata*, 198
 — *Farrerae*, 80
 — *fimbriata*, 105
 — Firefly, 33, 44
 — *fissa*, 166
 — Flambeau, 44
 — *flammea*, 127
 — *flava*, 103, 128
 — Florentina, 203
 — *floribunda*, 161
 — *fragrans*, 167, 190
 — François de Taye, 203
 — Fudesute Yama, 37
 — Fuji-manyo, 64
 — *fulva*, 131
 — Fürstin Baro Trysky, 49
 — *Gandavensis*, 179
 — Garnet, 43, 44
 — Géant des batailles, 203
 — *glauca*, 161
 — — *flore semipleno*, 166
 — — *hispida*, 163
 — — *plena*, 166
 — — *purpurascens*, 161
 — — *rosea*, 161
 — Gledstanesii, 27
 — Gloire de Verschaffelt, 203
 — Goldlack, 103
 — Gosho Zakura, 38
 — Graciosa, 98, 180
 — Graf von Meran, 98, 203
 — Grand-Duc, 203
 — *Azalea* Hachika Tsugi, 37
 — Hana Asobi, 38
 — Hana-guruma, 78
 — Hatsu-giri, 33
 — Havemeyer, 44
 — Hebe, 44
 — Helena, 44
 — Hermione, 44
 — Hexe, 33, 44
 — Hilda Hedlund, 44
 — Hinodegiri, 38
 — Hinode No Taka, 38
 — Hino-tsukasa, 30
 — Hiryu, 30
 — *hispida*, 163
 — Hiwoqe-yodogawa, 60
 — Holm Lea, 44
 — Hoō, 37
 — *hybrida Daviesii*, 182
 — — *enneandra*, 190
 — *Ignaea nova*, 203
 — *indica*, 22, 29, 38, 45
 — — *alba*, 68
 — — *amoena*, 32
 — — *angustifolia*, 23
 — — *calycina*, 63
 — — *crispiflora*, 27
 — — *exquisita*, 26
 — — *floribunda*, 23
 — — *grandiflora*, 27
 — — Hexe, 44
 — — *hybrida*, 62
 — — *ignescens*, 45
 — — *Iveryana albo-cincta*, 26
 — — *Kaempferi*, 38
 — — *lateritia*, 23
 — — *obtusata*, 29
 — — *obtusata alba*, 31
 — — *phoenicea*, 61
 — — *plena*, 73
 — — *punctulata*, 25
 — — *punicea*, 45
 — — *purpurea*, 73
 — — *purpurea plena*, 73
 — — *Rawsonii*, 61
 — — *roseum*, 72
 — — *rosiflora*, 26
 — — *Simsii*, 45
 — — *spatulata*, 23
 — — var. γ ., 61
 — — *variegata*, 26
 — — *vittata*, 48
 — — *vittata punctata*, 49
 — Ima Shōjō, 37
 — Irohayama, 37
 — Iso-tsutsuji, 73
 — Iveryana, 27
 — *japonica*, 99
 — — *alba grandiflora Van Noordt*, 71
 — — *grandiflora superba*, 72

- Azalea* Jas. H. Laing, 203
 — Jedogawa-tsutsuji, 70
 — J. T. Lovett, 25
 — Jupiter, 44
 — *Kaempferi*, 38
 — Kasane Kagaribi, 37
 — Kasumi Gaseki, 37
 — Katsura No Hana, 37
 — Kimigayo, 37
 — Kin-no-zai, 27
 — Kinshibe-tsutsuji, 42
 — Ki-rengé-tsutsuji, 102
 — Kirin, 37
 — Kirishima-tsutsuji, 30
 — Kiritsubo, 37
 — Ki-tsutsuji, 102
 — Kochonomai, 33
 — Kocho-zoroi, 77
 — Komanyo-satsuki, 26
 — Kumagai-yodogawa, 64
 — Kumo No Uye, 37
 — Kurai No Himo, 38
 — Kureno Yuki, 37
 — Kurofune-tsutsuji, 90
 — *lateritia*, 23
 — *ledifolia*, 68
 — — *alba pleno*, 72
 — — *Noordtiana*, 71
 — — *phoenicea*, 62
 — *liliflora*, 68
 — *linearifolia*, 74
 — Louis Aimé Van Houtte, 98, 204
 — Louis Hellebuyck, 204
 — *lucida*, 198
 — *lucida*, 162
 — *lugens*, 105
 — *lutea*, 127, 134, 143
 — *macrantha*, 23
 — *macrosepala*, 75
 — Madame Alex. Hardy, 204
 — Madame Camille van Langenhone, 97
 — Magnifica, 98
 — *magnifica*, 72, 192
 — Magnifica albicans, 204
 — *marginata*, 181
 — Marie-Dorothée, 204
 — Mars, 44
 — *Mattapan*, 71
 — *Maxwellii*, 64
 — Milton, 98
 — Mina Van Houtte, 204
 — Minerve, 204
 — Miniata floribunda, 204
 — Miss Louisa Hunnewell, 97
 — Mochi-tsutsuji, 75
 — *mollis*, 95, 100
 — Mollis var. albicans, 204
 — Mollis var. aurea floribunda, 204
 — *mollis flore-pleno*, 183
 — *Azalea mollis glabrior*, 99
 — — × *sinensis*, 97
 — Momo-iro-yodogawa, 73
 — *Mortieri*, 176
 — *Mortieriana*, 176, 179
 — Mrs. Carmichael, 33
 — *mucronata*, 68
 — Murasaki-botan-tsutsuji, 73
 — Murasaki Yama-tsutsuji, 43
 — Muriel, 44
 — Murillo, 98, 204
 — Nani Wagata, 37
 — *narcissiflora*, 71, 98, 204
 — Natalie, 44
 — *neglecta*, 149
 — Ne plus ultra, 204
 — Néron, 204
 — Nicolas II, 204
 — *nitida*, 157, 162
 — Nobilis, 204
 — Nonpareil, 97
 — *nudiflora*, 127, 131, 134, 139, 143, 147
 — — *alba*, 137, 141
 — — *albo-plena*, 138
 — — *amoena*, 198
 — — *bicolor*, 143
 — — *blanda*, 198
 — — *calycosa*, 198
 — — *canescens*, 143
 — — *carnea*, 137
 — — *caroliniana*, 198
 — — *ciliata*, 126
 — — Coburgia, 198
 — — *coccinea*, 131
 — — *colorata*, 198
 — — *conspicua*, 198
 — — *corymbosa*, 198
 — — *crispa*, 198
 — — *cumulata*, 198
 — — *discolor*, 198
 — — *fastigiata*, 198
 — — *flore pleno*, 138
 — — *florida*, 198
 — — *glandifera*, 138
 — — *globosa*, 199
 — — *glomerata*, 199
 — — *grandiflora*, 199
 — — *incana*, 199
 — — *incarnata*, 199
 — — *lutea*, 128
 — — *mirabilis*, 199
 — — *mixta*, 199
 — — *montana*, 199
 — — *ochroleuca*, 199
 — — *pallida*, 199
 — — *paludosa*, 199
 — — *papilionacea*, 199
 — — *partita*, 199
 — — *parviflora*, 199
 — — *periclymenoides*, 199

- Azalea nudiflora periclymenoides*, 135
 — — *polyandra*, 199
 — — *prolifera*, 200
 — — *pumila*, 200
 — — *purpurascens*, 200
 — — *purpurea*, 200
 — — *purpureo-plena*, 138
 — — *rosea*, 139
 — — *ruberrima*, 200
 — — *rubescens*, 200
 — — *rubicunda*, 200
 — — *rubra*, 138
 — — *rubro-plena*, 138
 — — *rufa*, 200
 — — *rutilans*, 200
 — — *rutilans*, 132
 — — *scintillans*, 178
 — — *semi-duplex*, 200
 — — *semi-duplex*, 138
 — — *serotina*, 200
 — — *staminea*, 200
 — — *stellata*, 200
 — — *thyrsiflora*, 132
 — — *tricolor*, 176
 — — *varia*, 200
 — — *variabilis*, 201
 — — *variegata*, 201
 — — *versicolor*, 201
 — — *violacea*, 201
 — — × *sinensis*, 183
 — *oblongifolia*, 150
 — *obtusa*, 29
 — — *alba*, 31
 — *occidentalis*, 126
 — — "Hybrids," 98
 — — × *arborescens* × *mollis*, 180
 — *octandra*, 200
 — *odorata*, 190
 — Oinō Mezame, 37
 — *Oldhamii*, 67
 — Omoine, 37
 — Omurasaki-tsutsuji, 62, 63
 — Optima, 48
 — Orange peinte, 204
 — Ō-riukiu-tsutsuji, 71
 — Osaraku, 37
 — Osaraku Seedling, 38
 — Oscar premier, 204
 — Otome, 37
 — Oyeyama-tsutsuji, 78
 — *pallida*, 190
 — *papilionacea*, 199
 — *periclymena*, 135
 — *periclymenoides*, 135
 — — *alba pleno*, 138
 — — *carnea*, 137
 — — *coccinea*, 131
 — — *papilionacea*, 199
 — — *partita*, 199
 — — *polyandra*, 199
 — — *rutilans*, 200
 — Perle du printemps, 204
 — Phébé, 204
 — *phoenicea*, 61
 — *pontica*, 103
 — — *autumnalis*, 103
 — — *calendulacea*, 128
 — — *coronaria* β. *ardens*, 178
 — — *coronaria* α. *flavum*, 178
 — — *flava*, 103
 — — Goldlack, 103
 — — *macrantha*, 204
 — — *sinensis*, 95
 — — et var., 179
 — — *versicolor*, 177, 204
 — — × evergreen *Rhododendron*, 192
 — *pontica-rubescens*, 178
 — Prince Albert, 48
 — Prince Henri des Pays-Bas, 204
 — Prince Troubetzkoy, 204
 — *prinophylla*, 139
 — *prolifera*, 73
 — *prunifolia*, 169
 — *pulchella carminea*, 201
 — — *roseola*, 201
 — *punctata*, 105
 — *punctulata omnicolor*, 25
 — — *variegata*, 25
 — *punicea*, 61
 — *purpurea pleno*, 73
 — Quadricolore, 204
 — Queen Victoria, 204
 — *quinquefolia pink*, 93
 — Racine, 98
 — *ramentacea*, 31
 — *ramis infra foliosis*, 158
 — Rashō Mon, 37
 — *Rawsonii*, 61, 196
 — Rayon du Matin, 204
 — Reine d'Angleterre, 204
 — Reine Louise, 27
 — Renge-tsutsuji, 101
 — *reticulata*, 84
 — *Reyndersiana*, 196
 — *rhombica*, 84
 — Rice Azalea, 54
 — *Richardii*, 204
 — *Rjuku Tsutsusi fl. igneo*, 103
 — *Rollissoni*, 26
 — *rosaeiflora*, 26
 — Rosalie, 204
 — *rosea*, 138
 — *Rosea formosissima*, 204
 — — *lineata*, 204
 — — *rotundiflora*, 204
 — Rose Queen, 44
 — *rosmarinifolia*, 68
 — — *alba*, 69
 — — *narcissiflora*, 71

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

DELVE INTO FANTASY, MAGIC, MYTHOLOGY & FOLKLORE

Forgotten Books'
Full Membership gives
access to 797,885 ancient
and modern, fiction and
non-fiction books.

Continue

*Fair usage policy applies

- Azalea vittata Bealii*, 49
 — *vittato-punctata*, 49
 — Vivid, 44
 — Vulcan, 44
 — Vuylestekeana, 44
 — Waka Kayede, 37
 — *Weyrichii*, 82
 — Yama tsutsuji, 40
 — Yayegiri, 30
 — Yayehiryu, 37
 — Yedogawa Satsuke, 70
 — Yin-shan-hung, 47
 — Yodogava, 64
 — Yorozuyo, 37
Azaleae Gandavenses, 179
 — *Mortierianae*, 176, 179
Azaleodendron, 188, 194
 — "Comte de Kerchove," 194
- Beverinckia*, 22
Biltia, 119
 — *Vaseyi*, 122
- Chamaerhodendron exoticum amplissimis floribus liliaceis*, 24
Chamaerhododendros, 121
 — *Pontica maxima Mespili folio flore luteo*, 104
 — *Virginiana Periclymeni flore ampliori minus odorato*, 135
Cistus Indicus Ledi Alpini foliis & floribus amplis, 25
 — *Virginiana flore et odore Periclymeni*, 143
 — — *flore et odore Periclymeni*, 158
 — — *Periclymeni flore ampliori et minus odorato*, 135
- Hochenwartia*, 119
 — *canadensis*, 120
- Loiseleria*, 22
- Osmothamnus fragrans*, 191
 — *pallidus*, 191
- Rhodazalea*, 188
 — *Crouxii*, 194
 — — *flore pleno*, 195
Rhododendro-Azalea, 188
Rhododendron a. *Anthodendron*, 115, 124
 — sect. *Azalea*, 79, 119, 124
 — sect. *Euazalea*, 124
 — sect. *Pentanthera*, 95, 124
 — sect. *Pentanthera*, 115
 — sect. *Pentanthera* × *Leiorhodion*, 188
 — sect. *Pentanthera* × *Rhodora*, 185
 — sect. *Rhodora*, 91, 119
 — sect. *Rhodora*, 79
- Rhododendron* sect. *Sciadorhodion*, 79
 — sect. *Toutousi*, 22
 — sect. *Tsusia*, 22
 — sect. *Tsutsugi*, 22
 — sect. *Tsutsutsi*, 22, 79, 187
 — subgen. *Anthodendron*, 115
 — subgen. *Anthodendron* × *Eurhodendron*, 188
 — subgen. *Azalea*, 115
 — subgen. *Tsutsia*, 22, 79
 — *Adamsii*, 190
 — *alabamense*, 141
 — *albicans*, 97, 180
 — *Albrechtii*, 91
 — *amoenum*, 32
 — — *Buergeri*, 206
 — — *euamoenum*, 206
 — — *genuinum*, 206
 — — *japonicum*, 33, 206
 — — *normale*, 206
 — — *obtusum*, 206
 — *Anneliesae*, 181
 — *arborescens*, 166
 — — *Richardsonii*, 168
 — — × *calendulaceum*, 181
 — — × *Mortieri*, 178
 — *atlanticum*, 147
 — — *luteo-album*, 150
 — — *neglectum*, 149
 — *atrovirens*, 50
 — *austrinum*, 146
 — *azaleodendron*, 194
 — *azaleoides*, 188
 — *azaleoides*, 190, 191
 — — *alba odorata*, 190
 — — *crispiflorum*, 196
 — — *inodorum*, 188
 — — *odoratum*, 190
 — *bicolor*, 143
 — *bigener*, 189
 — *boninense*, 206
 — *breviperulatum*, 28
 — *Breynii*, 23
 — *broughtonianum*, 194
 — *broughtonii aureum*, 99
 — *broughtonii aureum*, 194
 — *Buergeri*, 32
 — *Burmanni*, 68
 — *calendulaceum*, 127
 — *calendulaceum*, 132
 — — *aurantiacum*, 131
 — — *aurantium*, 130
 — — *chrysolectum*, 197
 — — *croceum*, 130
 — — *cupreum*, 139
 — — *flammeum*, 131
 — — *fulgidum*, 177
 — — *grandiflorum*, 197
 — — *ignescens*, 197
 — — *lepidum*, 177

- Rhododendron calendulaceum* Morteri, 176
 — — *speciosum*, 132
 — — *splendens*, 197
 — — *Stapletonianum*, 177
 — — *subcupreum*, 177
 — — *triumphans*, 198
 — — × ? *catawbiense*, 191
 — — × *nudiflorum*, 176
 — — × *occidentale*, 181
 — — × *speciosum*, 176
 — — × *viscosum*, 182
 — *Calleryi*, 45
 — *calycinum*, 63
 — *canadense*, 120
 — — *albiflorum*, 122
 — — *album*, 122
 — — *viridifolium*, 122
 — — × *japonicum*, 103, 185
 — — × *luteum*, 185
 — *candidum*, 146
 — *canescens*, 143
 — *canescens*, 139
 — — *candidum*, 146
 — — *subglabrum*, 145
 — *carneum elegantissimum*, 193
 — *Cartoni*, 189
 — *Cartonianum*, 189
 — *caryophyllum*, 52
 — Comte de Kerchove, 194
 — *coreanum*, 65
 — *crispiflorum*, 27
 — *Crouxii*, 194
 — — *plenum*, 195
 — *Danielsianum*, 23
 — *decandrum*, 84
 — Deception, 192
 — *decumbens*, 23
 — *dianthiflorum*, 76
 — *dilatatum*, 83, 86
 — — *decandrum*, 84
 — — *typicum*, 87
 — *discolor* × *occidentale*, 192
 — *Eeckhouttei*, 187
 — *elegantissimum*, 191
 — *enneandrum*, 190
 — *eximium*, 192
 — *Farrerae*, 79
 — *Farrerae*, 81
 — — *leucotrichum*, 81
 — — *mediocre*, 81
 — — *typicum*, 80
 — — *Weyrichii*, 81
 — *flammeum*, 132
 — *flavum*, 103
 — — *ardens*, 178
 — — *coronarum*, 103, 178
 — — *macranthum*, 105
 — — *Seymourii*, 185
 — — *versicolor*, 177
 — *fragrans*, 190
- Rhododendron fragrans*, 190, 191
 — *Fraseri*, 103, 185
 — *gandavense*, 178
 — — *plenum*, 179
 — — × ? *catawbiense*, 192
 — — × *Kosterianum*, 183
 — — × *molle*, 183
 — — × *occidentale*, 180
 — *gemmaferum*, 189
 — *glaucum*, 161
 — *gnaphalocarpum*, 81
 — *Gowenianum*, 189
 — *hallaisanense*, 65
 — *hangnoense*, 23
 — *hispidum*, 163
 — *hybridum*, 189
 — *hybridum enneandrum*, 190
 — — *glaucum*, 189
 — *indicum*, 22
 — *indicum*, 45, 59
 — — *amoenum Buergeri*, 32
 — — *amoenum genuinum*, 32
 — — *amoenum japonicum*, 33
 — — *amoenum normale*, 32
 — — *balsaminaeflorum*, 26
 — — *brachytrichum*, 23
 — — *Buergeri*, 32
 — — *crispiflorum*, 27
 — — *eriocarpum*, 49
 — — *exquisitum*, 26
 — — *formosanum*, 45
 — — *Hakatajiro*, 26
 — — *hakatashiro*, 25
 — — *Hexe*, 44
 — — *incarnatum*, 73
 — — *ignescens*, 45
 — — *japonicum*, 33
 — — *Kaempferi*, 38
 — — *Kaempferi subalpina*, 32
 — — *kinnozai*, 26
 — — *Kirishima*, 29
 — — *laciniatum*, 27
 — — *laciniatum*, 27
 — — *lateritium*, 23
 — — *macranthum*, 23
 — — *macranthum genuinum*, 23
 — — *macrostemon*, 31
 — — *Matsushima*, 26
 — — *mikawanum*, 43
 — — *obtusum*, 29
 — — *obtusum album*, 31
 — — *phoeniceum*, 61
 — — *polypetalum*, 27, 105
 — — *polypetalum*, 27
 — — *pulchrum*, 62
 — — *puniceum*, 45
 — — *purpureum*, 61
 — — *Rollisoni*, 26
 — — *rosiflorum*, 26
 — — *Simsii*, 45, 65

- Rhododendron indicum sinensis*, 59
 — — *Smithii*, 62
 — — *speciosum*, 61
 — — *subalpina*, 32
 — — *sublanceolatum*, 59
 — — *Tamurai*, 45
 — — *tanimanoyuki*, 26
 — — var. 45
 — — *variegatum*, 26, 105
 — — × *Kaempferi*, 72
 — — × *linearifolium*, 187
 — — × *mucronatum*, 187
 — — × *Simsii*, 187
 — *japonicum*, 99
 — — *aureum*, 102
 — — × *Eurhodendron* "hybrids,"
 194
 — — × *molle*, 183
 — — × *nudiflorum*, 83
 — *Jodogawa*, 59
 — *Kaempferi*, 38
 — — *album*, 41
 — — *amoenum*, 32
 — — *japonicum*, 33
 — — *Kinshibe*, 42
 — — *Komatsui*, 42
 — — *macrostemon*, 31
 — — *mikawanum*, 43
 — — *plenum*, 41
 — — × *rosmarinifolium*, 64
 — *Kanehirai*, 28
 — *kiusianum*, 33
 — *Kosterianum*, 97, 183
 — — × *Eurhodendron* hybrids, 194
 — *Laetitia*, 192
 — *lasiostylum*, 50
 — *lateritium*, 23
 — — *brachytrichum*, 23
 — *ledifolium*, 68
 — — *album*, 69
 — — *cryptopetalum*, 42
 — — *Hanaguruma*, 77
 — — *Kochozoroi*, 77
 — — *leucanthum*, 69
 — — *narcissiflorum*, 71, 73
 — — *Noordtianum*, 71
 — — *phoeniceum*, 61, 62
 — — *plena purpurea*, 73
 — — *purpureum*, 59, 63
 — *leucanthum*, 68
 — *liliflorum grandiflorum souchetianum*,
 69
 — *linearifolium*, 74, 105
 — — *decandrum*, 77
 — — *dianthiflorum*, 76
 — — *hanaguruma*, 77
 — — *linearifolium*, 74
 — — *macrosepalum*, 74
 — — *macrosepalum genuinum*, 75
 — — *polypetalum*, 77
- Rhododendron linearifolium rhodoroides*,
 77
 — *liukinense*, 59
 — *longiperulatum*, 67
 — *Loureiriana*, 105
 — *luteum*, 103
 — *luteum*, 128
 — — *flammeum*, 131
 — — *macranthum*, 105
 — — × *nudiflorum*, 177
 — — × *ponticum*, 192
 — — × *speciosum*, 178
 — — × *viscosum*, 178
 — *macranthum*, 23
 — — *albiflorum Souchetianum*, 68
 — — *polypetalum*, 105
 — — *semperflorens*, 105
 — *macrosepalum*, 74
 — — *Amaga-shita*, 78
 — — *Hanaguruma*, 77
 — — *linearifolium*, 74
 — — *Osuyo*, 78
 — — *rhodoroides*, 77
 — — *Suruga-momyo*, 78
 — *macrostemon*, 31
 — *Mariae*, 58
 — *Mariesii*, 80
 — *Matsumurai*, 64
 — *Maxwellii*, 64
 — *microphyton*, 57
 — *mixtum*, 98, 183
 — *molle*, 95
 — *molle*, 97, 99
 — — *glabrior*, 99
 — — × *arboreum*, 193
 — — × (*maximum* × *ponticum*), 193
 — — × *occidentale*, 98, 180
 — — × *sinense*, 97
 — — × *speciosum*, 181
 — — × *viscosum*, 182
 — *Morterii*, 176
 — — *carnea*, 176
 — *Mortieri*, 176
 — — × *luteum*, 178
 — — × *viscosum*, 178
 — *mucronatum*, 68
 — — *amethystinum*, 73
 — — *Mattapan*, 71
 — — *narcissiflorum*, 71
 — — *Noordtianum*, 71
 — — *plenum*, 73
 — — *ripense*, 72
 — — *sekidera*, 72
 — — × *scabrum*, 187
 — *Nakaharai*, 54
 — — *linearifolium*, 105
 — *narcissiflorum*, 71
 — *nipponicum*, 93, 119
 — *nitidum*, 162
 — *Norbiton hybrids*, 98, 193

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

Get Smart

Over 2,000 years of
human knowledge in
797,885 volumes

Instant access
\$8.99/month

Continue

*Fair usage policy applies

- Rhododendron puniceum*, 61
 — *purpureum*, 43
 — *quinquefolium*, 87
 — — *roseum*, 92
 — *ramentaceum*, 31
 — *reticulatum*, 83
 — — *albiflorum*, 86
 — — *pentandrum*, 86
 — *Rhodora*, 120
 — — *albiflora*, 122
 — *rhombicum*, 81, 83
 — — *albiflorum*, 86
 — *ripense*, 72
 — *roseum*, 138
 — *roseum odoratum*, 191
 — *rosmarinifolium*, 69
 — — *album*, 69
 — — *Fujimanyo*, 73
 — — *narcissiflorum*, 72
 — — *Oriuki*, 71
 — — *purpureum*, 63
 — — *Sekidera*, 72
 — — *Shiromanyo*, 72
 — — *speciosum*, 63
 — *rubropilosum*, 52
 — *Sanderi*, 43, 187
 — *scabrum*, 59
 — *scabrum*, 43
 — — *angustifolium*, 38
 — — *coccineum*, 61
 — — *Kaempferi*, 38
 — — *latifolium*, 38
 — — *purpureum*, 43
 — — *tubiflorum*, 38
 — *Schlippenbachii*, 88
 — *Seniavinii*, 57
 — *Sekidera*, 72
 — *serpyllifolium*, 53
 — *serpyllifolium*, 54
 — — *albiflorum*, 54
 — *serrulatum*, 153
 — — *georgianum*, 156
 — — *molliculum*, 155
 — *Seymouri*, 185
 — *shikokianum*, 82
 — *shojoense*, 81
 — *Sieboldii*, 38
 — — *ellipticum*, 60
 — — *minor*, 29
 — — *serrulatum*, 23
 — *Simsii*, 45
 — — *Bealii*, 49
 — — *eriocarpum*, 49
 — — *vittatum*, 48
 — — × *R. obtusum*, 43, 187
 — *sinense*, 95, 97, 99, 102
 — — *flammeum*, 95
 — — *flavescens*, 95, 102
 — — × *Eurhodendron*, 98, 193
 — — × *molle*, 97
- Rhododendron Smithii aureum*, 98
 — *Smithii aureum*, 193
 — *sonomense*, 127
 — *speciosum*, 131
 — *speciosum*, 131
 — — *acutifolium*, 201
 — — *aurantium*, 131
 — — *ciliatum*, 201
 — — *crispum*, 201
 — — *cucullatum*, 201
 — — *major*, 132
 — — *obliquum*, 201
 — — *prunifolium*, 201
 — — *revolutum*, 201
 — — *tortuifolium*, 201
 — — *tortulifolium*, 201
 — — *undulatum*, 202
 — — × *arboreum*, 192
 — — × *molle*, 181
 — *sublanceolatum*, 59
 — *sublateritium*, 59
 — *subsessile*, 51
 — *Tebotan*, 64
 — *Thunbergii*, 29
 — *Tolerianum*, 191
 — *tosaense*, 52
 — *tricolor*, 176
 — *trinerve*, 55
 — *Tschonoskii*, 55
 — — *pentamerum*, 55
 — — *tetramerum*, 55
 — — *trinerve*, 55
 — — *typicum*, 55
 — *Vaseyi*, 122
 — — *album*, 123
 — *venustum*, 135, 157
 — *viscosepalum*, 182
 — — *Daviesii*, 182
 — *viscosum*, 157
 — *viscosum*, 153
 — — *aemulans*, 165
 — — *coerulescens*, 165
 — — *crispum*, 202
 — — *dealbatum*, 202
 — — *fissum*, 166
 — — *glaucum*, 160
 — — *hispidum*, 163
 — — *montanum*, 164
 — — *nitidum*, 162
 — — *ornatum*, 178
 — — *penicillatum*, 202
 — — *pubescens*, 161
 — — *rhodanthum*, 160
 — — *roseum*, 161
 — — *rubescens*, 161
 — — *rubescens*, 161
 — — *tomentosum*, 162
 — — *variegatum*, 202
 — — × ? *catawbiense*, 190
 — — × *maximum*, 189

- Rhododendron *viscosum* × *ponticum*, 190
— *vittatum*, 49
— — *Bealii*, 49
— — *punctata*, 49
— *Wadanum*, 84
— *Weyrichii*, 82
— *Weyrichii*, 81, 89
— *Williamsii*, 194
- Rhododendron *yedoense*, 64
— — *poukhanense*, 65
— *Yodogawa*, 64
Rhodora, L., 119
— *canadensis*, 120
- Theis*, 124
Tsutsutsi, 22