




|   |                            |
|---|----------------------------|
| Number of surveys caught 1992–2010 (out of 19): | 19                         |
| Total catch weight (kg):                        | 2 326.7                    |
| Number measured                                 | 1 719                      |
| Length range (mean) (cm, FL)                    | 27–52 (41.2)               |
| Number weighed                                  | 300                        |
| Length-weight parameters a, b ( $r^2$ )         | 0.065605, 2.644337 (81.75) |


The core survey area and depth range **is not** appropriate for this species. It is found **shallower than 200 m**. Biomass of this species is **poorly** estimated in the core survey area. Biomass **shows no clear trend** since the start of the time series. Catch rates are highest in the **east**. Length frequencies are usually **unimodal**. Mean length has **decreased** since the start of the time series. Gonad stage data indicate that most fish are **maturing or spawning**.




#### Relative biomass estimates and length summary




| Year | Biomass (t) | cv (%) | Length (cm) |      |      | No. measure<br>d |
|------|-------------|--------|-------------|------|------|------------------|
|      |             |        | Min.        | Max. | Mean |                  |
| 1992 | 343         | 30     | 35          | 49   | 41.7 | 128              |
| 1993 | 205         | 65     | 34          | 49   | 42.7 | 93               |
| 1994 | 165         | 55     | 30          | 52   | 42.9 | 68               |
| 1995 | 110         | 56     | 38          | 47   | 42.2 | 28               |
| 1996 | 359         | 56     | 37          | 48   | 43.2 | 147              |
| 1997 | 224         | 54     | 27          | 51   | 41.3 | 135              |
| 1998 | 95          | 72     | 38          | 48   | 42.3 | 35               |
| 1999 | 91          | 41     | 33          | 50   | 43.4 | 42               |
| 2000 | 227         | 52     | 31          | 49   | 42.2 | 145              |
| 2001 | 218         | 26     | 33          | 48   | 40.4 | 110              |
| 2002 | 88          | 67     | 35          | 49   | 40.1 | 19               |
| 2003 | 633         | 36     | 32          | 48   | 39.7 | 343              |
| 2004 | 252         | 51     | 33          | 47   | 40.5 | 98               |
| 2005 | 238         | 66     | 32          | 45   | 40.4 | 63               |
| 2006 | 72          | 56     | 36          | 45   | 41.2 | 18               |
| 2007 | 19          | 79     | 37          | 43   | 40.2 | 4                |
| 2008 | 483         | 26     | 33          | 49   | 40.5 | 144              |
| 2009 | 74          | 41     | 36          | 44   | 41.0 | 21               |
| 2010 | 9           | 58     | 37          | 43   | 40.3 | 3                |








### Distribution































# Length Frequencies









## Gonad Stage Information

### Males

| Year | p_M1 | p_M2 | p_M3 | p_M4 | p_M5 | p_M6 | p_M7 | n_allM |
|------|------|------|------|------|------|------|------|--------|
| 1992 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1993 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1994 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1995 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1996 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1997 | 0    | 0.03 | 0.47 | 0.42 | 0.06 | 0.03 | 0    | 36     |
| 1998 | 0    | 0    | 0.05 | 0.95 | 0    | 0    | 0    | 20     |
| 1999 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2000 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2001 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2002 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2003 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| 2004 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2005 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| 2006 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2007 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2008 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2009 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 8      |
| 2010 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| ALL  | 0    | 0.17 | 0.27 | 0.52 | 0.03 | 0.02 | 0    | 66     |

### Females

| Year | p_F1 | p_F2 | p_F3 | p_F4 | p_F5 | p_F6 | p_F7 | n_allF |
|------|------|------|------|------|------|------|------|--------|
| 1992 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1993 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1994 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1995 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1996 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1997 | 0    | 0    | 0.75 | 0.25 | 0    | 0    | 0    | 4      |
| 1998 | 0    | 0    | 0.8  | 0.2  | 0    | 0    | 0    | 5      |
| 1999 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2000 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2001 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2002 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2003 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2004 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2005 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2006 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2007 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2008 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2009 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| 2010 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| ALL  | 0    | 0.18 | 0.64 | 0.18 | 0    | 0    | 0    | 11     |

Toadfishes

TOA



**Coded as COT**

|   |      |
|---|------|
| Number of surveys caught 1992–2010 (out of 19): | 9    |
| Total catch weight (kg):                        | 26.4 |

**Coded as PSY**

|   |      |
|---|------|
| Number of surveys caught 1992–2010 (out of 19): | 7    |
| Total catch weight (kg):                        | 64.6 |
| Number measured                                 | 6    |
| Number weighed                                  | 6    |

**Coded as TOA**

|   |      |
|---|------|
| Number of surveys caught 1992–2010 (out of 19): | 3    |
| Total catch weight (kg):                        | 27.3 |

**Coded as TOD**

|   |       |
|---|-------|
| Number of surveys caught 1992–2010 (out of 19): | 8     |
| Total catch weight (kg):                        | 114.4 |
| Number measured                                 | 1     |


**Coded as TOP**

|   |         |
|---|---------|
| Number of surveys caught 1992–2010 (out of 19): | 19      |
| Total catch weight (kg):                        | 2 770.9 |
| Number measured                                 | 24      |
| Number weighed                                  | 3       |

The core survey area and depth range **is** appropriate for this group. Biomass of this group is **well** estimated in the core survey area. Biomass has **increased and then decreased** since the start of the time series. Catch rates are highest in the **west**.


### Relative biomass estimates

| Year | Biomass (t) | cv (%) |
|------|-------------|--------|
| 1992 | 88          | 22     |
| 1993 | 136         | 19     |
| 1994 | 109         | 24     |
| 1995 | 90          | 33     |
| 1996 | 231         | 22     |
| 1997 | 249         | 19     |
| 1998 | 176         | 21     |
| 1999 | 386         | 17     |
| 2000 | 313         | 17     |
| 2001 | 470         | 15     |
| 2002 | 769         | 11     |
| 2003 | 410         | 19     |
| 2004 | 354         | 16     |
| 2005 | 197         | 24     |
| 2006 | 308         | 21     |
| 2007 | 156         | 28     |
| 2008 | 81          | 31     |
| 2009 | 167         | 24     |
| 2010 | 119         | 38     |





## Distribution






|   |              |
|---|--------------|
| Number of surveys caught 1992–2010 (out of 19): | 11           |
| Total catch weight (kg):                        | 97.5         |
| Number measured                                 | 12           |
| Length range (mean) (cm, FL)                    | 60–74 (67.5) |
| Number weighed                                  | 6            |
| Length-weight parameters a, b ( $r^2$ )         | –            |


There were **too few fish caught to determine whether the core survey area is appropriate for this species**. Biomass of this species is **poorly** estimated in the core survey area. Biomass **shows no clear trend** since the start of the time series.

**Relative biomass estimates**

| Year | Biomass (t) | cv (%) |
|------|-------------|--------|
| 1992 | 0           | -      |
| 1993 | 10          | 75     |
| 1994 | 6           | 74     |
| 1995 | 0           | -      |
| 1996 | 6           | 100    |
| 1997 | 0           | -      |
| 1998 | 9           | 100    |
| 1999 | 0           | -      |
| 2000 | 5           | 100    |
| 2001 | 17          | 100    |
| 2002 | 22          | 100    |
| 2003 | 0           | -      |
| 2004 | 0           | -      |
| 2005 | 8           | 100    |
| 2006 | 43          | 82     |
| 2007 | 0           | -      |
| 2008 | 23          | 100    |
| 2009 | 13          | 100    |
| 2010 | 0           | -      |



### Distribution






|   |       |
|---|-------|
| Number of surveys caught 1992–2010 (out of 19): | 8     |
| Total catch weight (kg):                        | 112.6 |
| Number measured                                 | 0     |
| Length range (mean) (cm)                        | –     |
| Number weighed                                  | 0     |
| Length-weight parameters a, b ( $r^2$ )         | –     |


There were **too few fish caught to determine whether the core survey area is appropriate for this species**. Biomass of this species is **poorly** estimated in the core survey area. Biomass **shows no clear trend** since the start of the time series.

**Relative biomass estimates**

| <b>Year</b> | <b>Biomass (t)</b> | <b>cv (%)</b> |
|-------------|--------------------|---------------|
| 1992        | 0                  | -             |
| 1993        | 4                  | 100           |
| 1994        | 0                  | -             |
| 1995        | 66                 | 30            |
| 1996        | 0                  | -             |
| 1997        | 65                 | 52            |
| 1998        | 0                  | -             |
| 1999        | 0                  | -             |
| 2000        | 1                  | 100           |
| 2001        | 0                  | -             |
| 2002        | 0                  | -             |
| 2003        | 46                 | 47            |
| 2004        | 0                  | -             |
| 2005        | 0                  | -             |
| 2006        | 1                  | 100           |
| 2007        | 15                 | 75            |
| 2008        | 0                  | -             |
| 2009        | 0                  | -             |
| 2010        | 1                  | 100           |



### Distribution






|   |            |
|---|------------|
| Number of surveys caught 1992–2010 (out of 19): | 19         |
| Total catch weight (kg):                        | 77.7       |
| Number measured                                 | 26         |
| Length range (mean) (cm, TL)                    | 24–34 (29) |
| Number weighed                                  | 16         |
| Length-weight parameters a, b ( $r^2$ )         | –          |


The core survey area and depth range **is not** appropriate for this species. It is found **deeper than 800 m**. Biomass of this species is **well** estimated in the core survey area. Biomass **shows no clear trend** since the start of the time series.

#### Relative biomass estimates

| Year | Biomass (t) | cv (%) |
|------|-------------|--------|
| 1992 | 4           | 23     |
| 1993 | 6           | 25     |
| 1994 | 8           | 32     |
| 1995 | 0           | 100    |
| 1996 | 17          | 26     |
| 1997 | 8           | 36     |
| 1998 | 12          | 20     |
| 1999 | 6           | 33     |
| 2000 | 10          | 19     |
| 2001 | 14          | 25     |
| 2002 | 11          | 15     |
| 2003 | 14          | 22     |
| 2004 | 10          | 21     |
| 2005 | 5           | 33     |
| 2006 | 19          | 15     |
| 2007 | 18          | 18     |
| 2008 | 7           | 26     |
| 2009 | 12          | 26     |
| 2010 | 15          | 16     |



### Distribution






|   |            |
|---|------------|
| Number of surveys caught 1992–2010 (out of 19): | 19         |
| Total catch weight (kg):                        | 2 643.7    |
| Number measured                                 | 489        |
| Length range (mean) (cm, TL)                    | 40–96 (64) |
| Number weighed                                  | 319        |
| Length-weight parameters a, b ( $r^2$ )         | –          |

The core survey area and depth range **is not** appropriate for this species. It is found **deeper than 800 m**. Biomass of this species is **poorly** estimated in the core survey area. Biomass **shows no clear trend** since the start of the time series.


#### Relative biomass estimates

| Year | Biomass (t) | cv (%) |
|------|-------------|--------|
| 1992 | 267         | 16     |
| 1993 | 167         | 40     |
| 1994 | 355         | 32     |
| 1995 | 157         | 47     |
| 1996 | 82          | 74     |
| 1997 | 148         | 71     |
| 1998 | 140         | 77     |
| 1999 | 283         | 56     |
| 2000 | 35          | 65     |
| 2001 | 164         | 49     |
| 2002 | 65          | 74     |
| 2003 | 238         | 51     |
| 2004 | 35          | 51     |
| 2005 | 132         | 34     |
| 2006 | 166         | 39     |
| 2007 | 161         | 45     |
| 2008 | 264         | 25     |
| 2009 | 139         | 37     |
| 2010 | 125         | 46     |





### Distribution



Witch (*Arnoglossus scapha*)

WIT




|   |              |
|---|--------------|
| Number of surveys caught 1992–2010 (out of 19): | 19           |
| Total catch weight (kg):                        | 200.1        |
| Number measured                                 | 17           |
| Length range (mean) (cm, TL)                    | 24–33 (28.2) |
| Number weighed                                  | 12           |
| Length-weight parameters a, b ( $r^2$ )         | –            |


The core survey area and depth range **is not** appropriate for this species. It is found **shallower than 200 m**. Biomass of this species is **moderately well** estimated in the core survey area. Biomass **shows no clear trend** since the start of the time series. Catch rates are highest in the **north**.

**Relative biomass estimates**

| Year | Biomass (t) | cv (%) |
|------|-------------|--------|
| 1992 | 23          | 37     |
| 1993 | 12          | 22     |
| 1994 | 40          | 29     |
| 1995 | 17          | 64     |
| 1996 | 10          | 34     |
| 1997 | 8           | 44     |
| 1998 | 12          | 51     |
| 1999 | 23          | 25     |
| 2000 | 31          | 31     |
| 2001 | 16          | 24     |
| 2002 | 17          | 26     |
| 2003 | 14          | 26     |
| 2004 | 10          | 32     |
| 2005 | 23          | 36     |
| 2006 | 25          | 49     |
| 2007 | 10          | 47     |
| 2008 | 15          | 38     |
| 2009 | 15          | 33     |
| 2010 | 8           | 34     |



**Distribution**






|   |                            |
|---|----------------------------|
| Number of surveys caught 1992–2010 (out of 19): | 19                         |
| Total catch weight (kg):                        | 36 293                     |
| Number measured                                 | 16 462                     |
| Length range (mean) (cm, FL)                    | 12–65 (37.1)               |
| Number weighed                                  | 5 055                      |
| Length-weight parameters a, b ( $r^2$ )         | 0.018314, 3.049201 (97.94) |


The core survey area and depth range is appropriate for this species. Biomass of this species is **moderately well** estimated in the core survey area. Biomass has **increased and then decreased** since the start of the time series. Catch rates are highest in the **west**. Length frequencies **have multiple modes which may contain information about year-class strength**. Mean length has **decreased and then increased** since the start of the time series. Gonad stage data indicate that most fish are **immature or resting**.




#### Relative biomass estimates and length summary




| Year | Biomass (t) | cv (%) | Length (cm) |      |      | No. measured |
|------|-------------|--------|-------------|------|------|--------------|
|      |             |        | Min.        | Max. | Mean |              |
| 1992 | 2 227       | 30     | 18          | 65   | 40.8 | 863          |
| 1993 | 2 939       | 46     | 20          | 63   | 42.2 | 956          |
| 1994 | 1 606       | 27     | 20          | 63   | 34.5 | 863          |
| 1995 | 734         | 25     | 19          | 62   | 37.1 | 295          |
| 1996 | 533         | 24     | 18          | 56   | 39.1 | 162          |
| 1997 | 2 287       | 20     | 14          | 62   | 35.6 | 789          |
| 1998 | 1 009       | 24     | 16          | 60   | 38.2 | 342          |
| 1999 | 3 136       | 41     | 15          | 62   | 37.7 | 825          |
| 2000 | 2 385       | 24     | 16          | 61   | 32.8 | 1 230        |
| 2001 | 4 262       | 36     | 13          | 65   | 32.7 | 1 387        |
| 2002 | 6 881       | 45     | 12          | 63   | 34.9 | 1 076        |
| 2003 | 3 685       | 34     | 24          | 60   | 34.2 | 1 368        |
| 2004 | 7 932       | 44     | 19          | 63   | 38.3 | 1 125        |
| 2005 | 4 542       | 25     | 23          | 61   | 37.5 | 1 102        |
| 2006 | 2 929       | 29     | 17          | 62   | 40.1 | 682          |
| 2007 | 2 853       | 21     | 23          | 61   | 39.1 | 874          |
| 2008 | 1 899       | 28     | 23          | 59   | 40.5 | 558          |
| 2009 | 3 667       | 33     | 26          | 62   | 43.0 | 756          |
| 2010 | 983         | 21     | 15          | 56   | 39.3 | 245          |





**Distribution**


































## Length Frequencies











## Gonad Stage Information

### Males

| Year | p_M1 | p_M2 | p_M3 | p_M4 | p_M5 | p_M6 | p_M7 | n_allM |
|------|------|------|------|------|------|------|------|--------|
| 1992 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1993 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1994 | 0.38 | 0.53 | 0    | 0    | 0    | 0    | 0.09 | 34     |
| 1995 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| 1996 | 0.41 | 0.59 | 0    | 0    | 0    | 0    | 0    | 44     |
| 1997 | 0.46 | 0.36 | 0.04 | 0    | 0    | 0    | 0.13 | 67     |
| 1998 | 0.45 | 0.33 | 0.01 | 0    | 0    | 0    | 0.22 | 138    |
| 1999 | 0.42 | 0.58 | 0    | 0    | 0    | 0    | 0    | 45     |
| 2000 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2001 | 1    | 0    | 0    | 0    | 0    | 0    | 0    | 2      |
| 2002 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2003 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2004 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2005 | 0.6  | 0.38 | 0.02 | 0    | 0    | 0    | 0    | 90     |
| 2006 | 1    | 0    | 0    | 0    | 0    | 0    | 0    | 4      |
| 2007 | 1    | 0    | 0    | 0    | 0    | 0    | 0    | 2      |
| 2008 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2009 | 0.08 | 0.91 | 0    | 0    | 0    | 0    | 0.01 | 150    |
| 2010 | 0.35 | 0.65 | 0    | 0    | 0    | 0    | 0    | 40     |
| ALL  | 0.37 | 0.55 | 0.01 | 0    | 0    | 0    | 0.07 | 617    |

### Females

| Year | p_F1 | p_F2 | p_F3 | p_F4 | p_F5 | p_F6 | p_F7 | n_allF |
|------|------|------|------|------|------|------|------|--------|
| 1992 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1993 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 1994 | 0.37 | 0.53 | 0    | 0    | 0    | 0    | 0.11 | 19     |
| 1995 | 0    | 0    | 1    | 0    | 0    | 0    | 0    | 2      |
| 1996 | 0.32 | 0.65 | 0.03 | 0    | 0    | 0    | 0    | 34     |
| 1997 | 0.31 | 0.52 | 0.17 | 0    | 0    | 0    | 0    | 58     |
| 1998 | 0.56 | 0.29 | 0.08 | 0    | 0    | 0    | 0.07 | 118    |
| 1999 | 0.44 | 0.56 | 0    | 0    | 0    | 0    | 0    | 25     |
| 2000 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| 2001 | 0.5  | 0    | 0.5  | 0    | 0    | 0    | 0    | 2      |
| 2002 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2003 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2004 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2005 | 0.22 | 0.54 | 0.24 | 0    | 0    | 0    | 0    | 82     |
| 2006 | 1    | 0    | 0    | 0    | 0    | 0    | 0    | 2      |
| 2007 | 0    | 1    | 0    | 0    | 0    | 0    | 0    | 1      |
| 2008 | NA   | NA   | NA   | NA   | NA   | NA   | NA   | 0      |
| 2009 | 0.07 | 0.92 | 0.01 | 0    | 0    | 0    | 0    | 74     |
| 2010 | 0.29 | 0.64 | 0.07 | 0    | 0    | 0    | 0    | 28     |
| ALL  | 0.33 | 0.54 | 0.11 | 0    | 0    | 0    | 0.02 | 446    |