

NOVEMBER 2017



*Providing sustainable environmental strategies,
management and monitoring solutions
to industry and government.*



**WIND PROSPECT PTY LTD
YANDIN WIND FARM
FLORA, VEGETATION AND AVIFAUNA ASSESSMENT**

This page has been left blank intentionally.

Document Status						
Rev.	Author(s)	Reviewer	Date	Approved for Issue		
				Name	Distributed To	Date
1	A Craigie A Schmitz	S Grein	30/10/2017	S. Grein	R. Baker	30/10/2017

ecologia Environment (2017). Reproduction of this report in whole or in part by electronic, mechanical or chemical means, including photocopying, recording or by any information storage and retrieval system, in any language, is strictly prohibited without the express approval of Wind Prospect Pty Ltd

ecologia Environment
45 Gladstone Street
East Perth WA 6004
Phone: 08 6168 7200
Email: admin@ecologia.com.au

This page has been left blank intentionally.

TABLE OF CONTENTS

1	INTRODUCTION.....	1
1.1	PROJECT BACKGROUND.....	1
1.2	SURVEY OBJECTIVES.....	1
1.3	LEGISLATIVE FRAMEWORK	1
2	DEFINITIONS	4
2.1	SIGNIFICANT FLORA	4
2.2	SIGNIFICANT VEGETATION.....	4
3	METHODOLOGY	6
3.1	DESKTOP STUDY.....	6
3.2	FLORA AND VEGETATION.....	7
3.3	AVIFAUNA	11
3.4	STUDY TEAM AND LICENCES.....	11
4	DESKTOP RESULTS.....	12
4.1	CLIMATE	12
4.2	IBRA 7 BIOGEOGRAPHIC SUBREGIONS	12
4.3	SOILS	13
4.4	VASCULAR FLORA.....	16
4.5	VEGETATION	23
4.6	SIGNIFICANT ECOLOGICAL COMMUNITIES.....	25
4.7	PREVIOUS SURVEYS WITHIN THE STUDY AREA.....	27
5	FIELD SURVEY RESULTS.....	28
5.1	FLORA.....	28
5.2	VEGETATION	30
5.3	AVIFAUNA	45
5.4	SURVEY LIMITATIONS AND CONSTRAINTS	50
6	DISCUSSION	51
6.1	FLORA.....	51
6.2	VEGETATION	51
6.3	AVIFAUNA	52
7	REFERENCES.....	53
8	APPENDICES.....	55

TABLES

Table 3.1 – Databases searched for the literature review	6
Table 3.2 – Criteria used to assess the likelihood of occurrence of significant fauna, flora and vegetation	7
Table 3.3 – Vegetation condition scale (Keighery 1994)	10
Table 3.4 – Study team and licences	11
Table 4.1 – Threatened and Priority Flora records from database searches	17
Table 4.2 – Pre-European vegetation association extent	23
Table 4.3 – TECs in the vicinity of the study area.....	25
Table 5.1 – <i>Hypocalymma tetrapterum</i> (P3) records within the study area.....	28
Table 5.2 – Summary of ‘Banksia Woodlands of the Swan Coastal Plain’ patches within the study area	33
Table 5.3 – Summary of vegetation units at the study area	31
Table 5.4 – Avifauna records within sites and habitats.....	45
Table 5.5 – Avifauna records within flight zones	47
Table 5.6 – Regional records of avifauna of conservation significance	49
Table 5.7 – Flora and vegetation survey limitations	50

FIGURES

Figure 1.1 – Regional location of the study area.....	2
Figure 1.2 – The study area	3
Figure 4.1 – Climate data from Badgingarra BOM weather station (Station No. 009037)	12
Figure 4.2 – IBRA 7 subregions.....	14
Figure 4.3 – Soils.....	15
Figure 4.4 – Threatened Flora records from DBCA databases	20
Figure 4.5 – Priority 1 and 2 Flora records from DBCA databases	21
Figure 4.6 – Priority 3 and 4 Flora records from DBCA databases	22
Figure 4.7 – Pre-European vegetation mapped at the study area	24
Figure 4.8 – TECs in the vicinity of the study area	26
Figure 5.1 – <i>Hypocalymma tetrapterum</i> (P3) records within the study area	29
Figure 5.2 – Vegetation units within the study area (map index).....	33
Figure 5.3 – Vegetation units within the study area (Map A1)	35
Figure 5.4 – Vegetation units within the study area (Map A2)	36
Figure 5.5 – Vegetation units within the study area (Map A3)	37
Figure 5.6 – Vegetation units within the study area (Map B1)	38

Figure 5.7 – Vegetation units within the study area (Map B2)39
Figure 5.8 – Vegetation units within the study area (Map B3)40
Figure 5.9 – Vegetation units within the study area (Map C1)41
Figure 5.10 – Vegetation units within the study area (Map C2)42
Figure 5.11 – Vegetation condition within the study area.....43
Figure 5.12 – Banksia Woodlands of the Swan Coastal Plain TEC within the study area44

APPENDICES

- Appendix A Definitions
- Appendix B Vascular flora records - NatureMap
- Appendix C EPBC protected matters search tool
- Appendix D Significant flora records - TPFL and WAHERB
- Appendix E Regional Bird Records - NatureMap
- Appendix F Conservation Significant Bird Records - NatureMap
- Appendix G Study area plant species list
- Appendix H Flora Survey site data

ACRONYMS

BAM Act	<i>Biosecurity and Agriculture Management Act 2007</i>
BOM	Bureau of Meteorology
BIF	Banded Ironstone Formation
CALM	Department of Conservation and Land Management (now DBCA and DER)
CAMBA	China – Australia Migratory Bird Agreement
CSIRO	Commonwealth Scientific and Industrial Research Organisation
DAFWA	Department of Agriculture and Food Western Australia
DBCA	Department of Biodiversity, Conservation and Attractions (previously DPaW)
DEC	Department of Environment and Conservation (now DBCA)
DER	Department of Environmental Regulation
DoEE	Department of the Environment and Energy (Previously DSEWPaC)
DPaW	Department of Parks and Wildlife (now DBCA)
DSEWPaC	Department of Sustainability, Environment, Water, Population and Communities (now DoEE)
EPA	Environment Protection Authority
EP Act	<i>Environment Protection Act 1986</i>
EPBC Act	<i>Environment Protection and Biodiversity Conservation Act 1999</i>
ESCAVI	Executive Steering Committee for Australian Vegetation Information
IA	International Agreement
IBRA	Interim Biogeographic Regionalisation for Australia
ICE	Incidence-based Coverage Estimators
IPA	Indigenous Protected Area
IUCN	International Union for Conservation of Nature
NVIS	National Vegetation Information System
PEC	Priority Ecological Community
TEC	Threatened Ecological Community
TPFL	Threatened and Priority Flora database
TP List	Threatened and Priority Flora List
WA	Western Australia
WAHERB	Western Australian Herbarium Specimen Database
WAOL	Western Australian Organism List
WC Act	<i>Wildlife Conservation Act 1950</i>
WONS	Weeds of National Significance

EXECUTIVE SUMMARY

ecologia Environment (*ecologia*) was commissioned by Wind Prospect Pty Ltd to undertake revised flora and fauna assessments of the Yandin Wind Farm project area located to the southeast of Cataby, Western Australia. A reconnaissance flora and vegetation survey, Threatened Ecological Community assessment, and avifauna survey were undertaken over the approximately 15,360 ha study area between September 18th and 20th 2017.

Flora and Vegetation

A total of 117 sub-generic vascular plant taxa from 35 families were recorded from the study area during the current survey. The most diverse families recorded were the Proteaceae (22 taxa), Myrtaceae (15 taxa), Fabaceae (10 taxa), and Poaceae (9 taxa).

DBCAs database searches identified 61 conservation significant plant species that have been recorded within or have potential to occur within the study area. Only one Priority listed species (*Hypocalymma tetrapterum* P3) was recorded during the current survey, with targeted searches focussed on native vegetation patches in the vicinity of proposed development areas.

Vegetation data collected from 52 sites from across the study area were used to describe and map 16 vegetation units. Vegetation over 91.2% of the study area comprised scattered *Corymbia calophylla* or *Eucalyptus todtiana* trees over pasture weeds in ‘completely degraded’ condition. Low-lying areas dominated by *Juncus acutus* were also completely degraded. Two species rich shrubland units in good to excellent condition, five eucalypt woodland units in degraded condition, and seven eucalypt or *Banksia* woodland units in good to excellent condition, were described from remnant native vegetation patches.

Five patches of the ‘Banksia Woodlands of the Swan Coastal Plain’ ecological community were identified within the study area, in addition to two previously mapped patches. Based on vegetation structure and composition, patch size, and condition, each of these patches are considered to qualify as the TEC according to ‘Approved Conservation Advice (incorporating listing advice) for the Banksia Woodlands of the Swan Coastal Plain Ecological Community’ (Threatened Species Scientific Committee 2016a).

Avifauna

Results of the current study verify those of a more comprehensive survey, impact assessment and risk assessment of potential bird strike with wind turbines undertaken by RPS Environment and Planning in 2010 (RPS 2010). That is, that there was a relatively low diversity of bird species recorded from open pasture areas characterising the proposed wind turbine positions, with the greatest diversity limited to stands of intact native vegetation comprised mainly of shrubland and heath. The presence of isolated trees within cleared areas often attracted birds, albeit in small numbers and at low diversities.

Although there are a number of regional wetlands in the wider locality wetland habitats at the study area were limited to small farm dams or degraded creeklines.

Forty-four species, within seven habitat types, were recorded during the survey. All species had previously been recorded by RPS (2010), apart from the Australasian Pipit and Emu. Birds recorded during the survey with at least the potential to fly within the rotor-swept area (RSA) potential collision zone were; Straw-necked Ibis, Black-shouldered Kite, Carnaby's Black-Cockatoo, Butler's Corella, Galah, and Australian Kestrel. All these species were similarly identified by RPS (2010).

RPS (2010) recorded species flying at RSA elevations ‘on more than a rare occasion’ as; Australian Kestrel, Wedge-tailed Eagle, Brown Falcon, White-backed Swallow, Black-shouldered Kite and Fairy

Martin. Of these, the Australian Kestrel and Black-shouldered Kite were also recorded within RSA elevations during this study.

Proteaceous heath and shrubland over the study area, as well as plantations of pine, provide foraging habitat for the threatened Carnaby's Black-Cockatoo. Flight movements are likely to follow movements to and from these food resources as well as roosting and/or breeding trees. RPS (2010) reported that this species primarily frequented lowland areas with movements tending to follow valleys with woodland vegetation.

RPS (2010) provide a comprehensive survey, impact assessment, and risk assessment of potential bird strike, with data obtained from this study corroborating the results presented there.

This page has been left blank intentionally.

1 INTRODUCTION

1.1 PROJECT BACKGROUND

Ecologia Environment (*ecologia*) was commissioned by Wind Prospect Pty Ltd to undertake revised flora and fauna assessments of the Yandin Wind Farm project area located to the southeast of Cataby, Western Australia to support a native vegetation clearing permit (NVCP) application and planning approvals for the proposed development (Figure 1.1). A reconnaissance flora and vegetation survey, Threatened Ecological Community assessment, and avifauna survey were undertaken over the approximately 15,360 ha study area (Figure 1.2) between September 18 and 20 2017.

1.2 SURVEY OBJECTIVES

The EPA's environmental objectives for the factors *Flora and Vegetation* (EPA 2016a) and *Terrestrial Fauna* (EPA 2016b) are: "To protect flora and vegetation and terrestrial fauna so that biological diversity and ecological integrity are maintained." In the context, 'ecological integrity' is the composition, structure, function and processes of ecosystems, and the natural range of variation of these elements.


The primary objective of this flora and avifauna assessment was to provide sufficient information for the EPA to assess the impact of the proposed development on the flora, vegetation and avifauna of the study area, thereby ensuring that the EPA objectives will be upheld. To this end, the following were provided as part of this assessment:


- A desktop study to evaluate biological values of the study area and surrounds, including a review of existing environmental values, threatened and priority flora and vegetation databases, and other relevant available literature;
- A reconnaissance survey to confirm the findings of the desktop study and to verify the vegetation types described and mapped during a previous flora and vegetation survey (Outback Ecology 2009) and the assemblages and conservation status of bird species identified in the previous avifauna assessment (RPS 2010);
- An assessment of the EPBC Listed 'Banksia Woodlands of the Swan Coastal Plain' Threatened Ecological Community (TEC) within the study area;
- A targeted flora survey for Threatened and Priority listed plant species within remnant patches of native vegetation within the study area, with a focus on areas adjacent to proposed development;
- Vegetation community and condition mapping; and
- An up-to-date account of the birds of conservation significance potentially occurring, and augmentation and verification of the bird records obtained by RPS (2010) with an emphasis on flight patterns and movements.

1.3 LEGISLATIVE FRAMEWORK

The surveys were designed and undertaken to comply with the following guidance documents:


- Environmental Factor Guideline: Flora and Vegetation (EPA 2016a);
- Environmental Factor Guideline: Terrestrial Fauna (EPA 2016b);
- Technical Guidance: Flora and Vegetation Surveys for Environmental Impact Assessment (EPA 2016c); and
- Technical Guidance: Terrestrial Fauna Surveys (EPA 2016d).



 Yandin Wind Farm Study Area


ecologia
ENVIRONMENT

Drawn: AC Project ID: 1713
Date: 17 August 2017 A4



Regional location of the study area

Figure:
1.1



Yandin Wind Farm Study Area

ecologia
ENVIRONMENT

Drawn: AC Project ID: 1713
Date: 17 August 2017 A4



0 1.5 3
Kilometers
Scale: 1:150,000
MGA94 (Zone 50)

Yandin Wind Farm study area

Figure:
1.2

2 DEFINITIONS

2.1 SIGNIFICANT FLORA

According to the *EPA Factor Guideline: Flora and Vegetation* (EPA 2016a), plant species (or records) may be considered significant for a number of reasons including, but not restricted to, the following:

- Being identified as Threatened or Priority species;
- Locally endemic species or those associated with a restricted habitat type (e.g. surface water or groundwater dependent ecosystems);
- New species or those having anomalous features that indicate a potential new species;
- Being representative of the range of a species (particularly, at the extremes of range, recently discovered range extensions, or isolated outliers of the main range);
- Unusual species, including restricted subspecies, varieties or naturally occurring hybrids; and
- Being representative of taxonomic groups that no longer occur widely in the broader landscape (relictual species/populations).

Environment Protection and Biodiversity Conservation Act 1999 (Commonwealth of Australia)

At a Commonwealth level, Threatened Flora are protected under the EPBC Act, which lists species that are considered Critically Endangered, Endangered, Vulnerable, Conservation Dependant, Extinct, or Extinct in the Wild (refer to Appendix A for category definitions).

Wildlife Conservation Act 1950 (Western Australia)

At State level, Threatened Flora species are protected under the WC Act. These are taxa which have been adequately surveyed and are deemed to be either rare, in danger of extinction, or otherwise in need of special protection in the wild, and are gazetted as Threatened (Declared Rare) Flora. Threatened Flora are further categorised by DBCA according to their level of threat using the International Union for Conservation of Nature (IUCN) red list criteria (IUCN 2001) (Appendix A). These taxa are legally protected and their removal or impact to their surroundings cannot be conducted without Ministerial approval, obtained specifically on each occasion for each population.

Priority Flora (DCBA)

DBCA maintains a list of Priority Flora species, which are considered poorly known, uncommon or under threat but for which there is insufficient justification to be listed as Threatened, based on known distribution and population sizes. Priority Flora species are assigned to one of four categories, described in Appendix A.

2.2 SIGNIFICANT VEGETATION

According to *EPA Factor Guideline: Flora and Vegetation* (EPA 2016a), vegetation may be considered significant for a number of reasons including, but not restricted to, the following:

- Being identified as Threatened or Priority Ecological Communities;
- Having a restricted distribution;
- The degree of historical impact from threatening processes;
- Playing a role as a refuge;
- Providing an important function required to maintain ecological integrity of a significant ecosystem.

Threatened Ecological Communities (Nationally Listed)

Ecological communities are naturally occurring biological assemblages associated with a particular type of habitat (DEC 2010). At a national level, Threatened Ecological Communities (TECs) are protected under the Commonwealth EPBC Act. An ecological community may be categorised into one of three sub-categories: Critically Endangered, Endangered, and Vulnerable (Appendix A).

Threatened Ecological Communities (State Listed)

DBCA maintains a list of state listed TECs which are further categorised into three subcategories, reflecting those of the EPBC Act. Within the Western Australian classification, an ecological community will be listed as Vulnerable "when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing a high risk of total destruction or significant modification in the medium to long-term future".

Priority Ecological Communities

DBCA maintains a list of Priority Ecological Communities (PEC). PECs include potential TECs that do not meet survey criteria, or that are not adequately defined. DBCA categorises PECs into five categories, P1 to P5, depending on the level of threat to the community (Appendix A).

Regional and Local Significance

Regional significance addresses the representation of habitats at a biogeographic regional level. Vegetation communities that are restricted or uncommon in a regional context are considered regionally significant. Vegetation communities supporting Threatened Flora species may also be considered regionally significant. Accurate assessment of regional significance requires sufficient regional vegetation community data to be available, and described at a similar level to the current study.

Locally significant vegetation may include vegetation communities that are locally restricted, contain comparatively high structural or species diversity, or contain Priority Flora species that are restricted to these vegetation communities.

3 METHODOLOGY

3.1 DESKTOP STUDY

The methodology adopted for this desktop study was consistent with that recommended by EPA (2016c) and EPA (2016d). A review of background environmental information for the study area was conducted including previous flora, vegetation, and fauna surveys, climate (BoM), biogeography (IBRA 7) (USE 2012), soils (Northcote *et al.* 1960-1968), and pre-European vegetation (Shepherd *et al.* 2001).

A search and review of all available relevant reports in the vicinity of the study area was undertaken, as well as searches of mapping resources and databases listed in Table 3.1 to determine conservation significant species and communities previously recorded within the study area or vicinity.

All results were reviewed on the basis of the likelihood of occurrence of relevant conservation significant species occurring within the study area with consideration given to previous records, habitat requirements, and landform.

The database searches and literature review resulted in an inventory of flora, vegetation and avifauna of conservation significance with at least the potential to occur within the study area.

Table 3.1 – Databases searched for the literature review

Database	Search Details
EPBC Act Protected Matters Database	Records of matters of national significance under the EPBC Act within 10 km of the study area
DBCA Threatened and Priority Ecological Communities Database	Records of TEC/PECs within 10km of the study area
DBCA Threatened and Priority flora Database	Records of significant flora within 10 km of the study area
Threatened and Priority flora List (TPList)	Records of significant flora by place names within 10 km of the study area
Western Australian Herbarium Specimen Database (WAHERB)	Records of significant flora within 10 km of the study area
DBCA NatureMap	All flora records within 10 km of the study area and avifauna records within 20 km of the study area

The database searches and literature review resulted in an inventory of flora, vegetation and avifauna of conservation significance with at least the potential to occur within the study area. The criteria listed in Table 3.2 were then applied to determine the likelihood of occurrence of significant species and vegetation occurring within the study area given the likely landforms and broad habitats present.

Table 3.2 – Criteria used to assess the likelihood of occurrence of significant fauna, flora and vegetation

Rating	Criteria (significant flora and fauna)	Criteria (TEC/PEC)
Recorded	The taxon has previously been recorded in the study area.	The TEC/PEC (not including buffer) has previously been recorded in the study area.
Possible (1)	The habitat preference of the taxon is well defined or broadly defined and this habitat likely occurs within the study area, and there are previous records in the vicinity of the study area.	Due to the proximity of previous records and the likely presence of suitable habitat/geology within the study area, the TEC/PEC possibly occurs within the study area.
Possible (2)	The habitat preference of the taxon is broadly defined or undefined and suitable habitat possibly occurs at the study area, but there are no records in the vicinity of the study area; or there is otherwise insufficient information available to exclude the possibility of occurrence at the study area.	The community is broadly defined and could possibly occur at the study area and there are records in the vicinity of the study area; or there is insufficient information available to exclude the possibility of occurrence at the study area.
Unlikely (3)	The habitat preference of the taxon is well defined and suitable habitat is considered unlikely to be present within the study area.	The community is well defined and suitable habitat/geology is considered unlikely to be present within the study area.

3.2 FLORA AND VEGETATION

3.2.1 Field Methodology

The flora and vegetation survey was conducted at the study area between 18 and 20 September 2017, and was consistent with guidelines for a reconnaissance flora and vegetation survey (EPA 2016c) and *'Approved Conservation Advice (incorporating listing advice) for the Banksia Woodlands of the Swan Coastal Plain Ecological Community'* (Threatened Species Scientific Committee 2016a).

Reconnaissance Survey

As part of the reconnaissance flora and vegetation assessment, 48 'check sites' were surveyed across the study area to provide ground-truthed data for vegetation community and condition mapping. Check sites are an unmarked area in which floristic data are collected, and are a low-intensity survey technique used to confirm the presence of vegetation communities and assess vegetation condition. The following data were recorded for each site:

- Site number and location (GPS co-ordinate of the north-west corner);
- Photograph of the vegetation;
- Dominant growth form, height, cover and up to three species for the three traditional strata (upper, mid and ground) compatible with NVIS Level V (ESCAVI 2003);
- Additional information to assist vegetation classification, including landform, soils, slope, aspect, rock type and abundance, litter, and fire history; and
- Vegetation condition (Table 3.3) and description of disturbance.

‘Banksia woodland of the Swan Coastal Plain’ TEC Survey

To assess presence of the ‘Banksia woodlands of the Swan Coastal Plain’ TEC, 10 m x 10 m quadrats were established and sampled within remnant vegetation patches that were considered to potentially represent the TEC. The following data were recorded at each quadrat:

- Site number and location (GPS co-ordinate of the north-west corner);
- Photograph from the north-west corner;
- Size and shape of quadrat;
- Dominant growth form, height, cover and up to three species for the three traditional strata (upper, mid and ground) compatible with NVIS Level V (ESCAVI 2003);
- A comprehensive species list (including weeds);
- Additional information to assist vegetation classification, including landform, soils, slope, aspect, rock type and abundance, litter, and fire history; and
- Vegetation condition (Table 3.3) and description of disturbance.

Community characteristics given in the *Banksia Woodlands of the Swan Coastal Plain Approved Conservation Advice* (Threatened Species Scientific Committee 2016a) were used to identify the TEC, primarily:

1. Key Diagnostic Features

- Location
 - Swan Coastal Plain IBRA Bioregion (SWA01, SWA02), and extending into the Jarrah Forest Bioregion (JAF01, JAF02).
- Soils and Landform
 - Well drained, low nutrient soils on sandplains - Bassendean and Spearwood sands and occasionally on Quindalup sands.
- Structure and Composition
 - A distinctive upper sclerophyllous layer of low trees typically dominated or co-dominated by *Banksia attenuata*, *Banksia menziesii*, *Banksia prionotes*, or *Banksia ilicifolia*;
 - Emergent trees of medium or tall (>10 m) eucalypts (typically *Corymbia calophylla*, *Eucalyptus gomphocephala*, or *Eucalyptus marginata*) or *Allocasuarina* may be present above the *Banksia* canopy;
 - An often highly species-rich understorey of shrubs and herbs.

2. Condition Thresholds

To be considered as part of the TEC, a patch should meet at least the ‘Good’ condition category (Keighery 1994), typically with at least ‘low’ native plant species diversity and weeds at 5-50% cover.

3. Minimum Patch Size

Minimum patch sizes are apply based on condition:

- ‘Pristine’ – no minimum patch size applies
- ‘Excellent’ – 0.5 ha
- ‘Very Good’ – 1 ha
- ‘Good’ – 2 ha

4. Additional Considerations

- A patch is a discrete and mostly continuous area of the ecological community and may include small-scale (<30 m) variations, gaps and disturbances, such as tracks, paths or breaks
- Restored (revegetated or replanted) sites are not excluded from the listed community.

Conservation Significant Flora Survey

Threatened and Priority Flora species identified from the database searches were targeted during the field survey, informed by previous record locations and known habitat preferences. Searches for conservation significant species involved searches of potential suitable habitat and opportunistic records taken during traverses walked between sites, specifically targeting areas that would potentially be impacted by the proposed development, or in close vicinity to impact areas. Where conservation significant species were observed the following data were recorded:

- Recorder and date;
- Individual GPS coordinates (GDA94) (for individual or localised plants), or GPS coordinates of population extent (for more extensive populations);
- Number of plants (count, for individual or localised plants) or estimated number of plants for more extensive populations;
- Reproductive state;
- Vegetation type; and
- Landform.

Specimen Identification

Plant specimen identification was undertaken with reference to current taxonomic literature and herbarium reference specimens. Scientific names used in this report follows the species nomenclature currently adopted by the Western Australian Herbarium. Specimens that were believed to differ significantly from typical material were indicated with 'affinity' (aff.). Specimens that could not be adequately identified to genus or species level due to the absence of flowering or fruit material required for positive identification were indicated with a question mark, but were not considered to be otherwise anomalous.

3.2.2 Vegetation Mapping

Vegetation units were characterised by determining diagnostic or characteristic combinations of dominant species. Vegetation communities are naturally variable across wide geographic areas, and units here are delineated based on the overall floristic similarity of sites with various spatial coverage. Therefore, species used in descriptions are those that are most dominant or characteristic of the vegetation type as a whole, but were not necessarily recorded in the same combination at all sites. Species that are recorded as sometimes occurring in a vegetation unit are indicated in the description by "±" (ESCAVI 2003). Extrapolative vegetation mapping based on aerial imagery and ground-truth data provided by site assessments was then used to map the described vegetation units within study area.

Vegetation condition was mapped across the study area based on vegetation condition ratings recorded at quadrats and check sites, as well as additional observations made during the survey.

Table 3.3 – Vegetation condition scale (Keighery 1994)

Vegetation Condition	Criterion
Pristine	Pristine or nearly so, no obvious signs of disturbance or damage caused by human activities since European settlement.
Excellent	Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species. Damage to trees caused by fire, the presence of non-aggressive weeds and occasional vehicle tracks.
Very good	Vegetation structure altered, obvious signs of disturbance. Disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing.
Good	Vegetation structure significantly altered by very obvious signs of multiple disturbances. Retains basic vegetation structure or ability to regenerate it. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.
Degraded	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds at high density, partial clearing, dieback and grazing.
Completely Degraded	The structure of the vegetation is no longer intact and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees and shrubs.

3.3 AVIFAUNA

RPS (2010) previously undertook targeted and systematic avifauna field surveys within the study area, utilising 27 bird census points during three separate survey periods; from 29th October to 7th November 2008, from 18th to 26th November 2008, and from 15th to 16th January 2009. In addition, RPS (2010) also conducted comprehensive surveys at the nearby Waddi Wind Farm, as well as around wetland habitats in the wider locality to determine regional status, and movements of local waterbird populations and to assess habitat potential for migratory wading bird species.

The flight characteristics of all individual species throughout all bird surveys were recorded by RPS (2010), with flight height ranges split into three zones in consideration of the potential collision zone of the rotor-swept area (RSA), that is, from 40 m to 152 m above ground level:

- Zone 1 – 0 to 40 m (that is, below the tip of turbine blade).
- Zone 2 – 40 to 152 m (RSA and potential collision zone).
- Zone 3 – >152 m (above the tip of turbine blade).

Sixty-one species were recorded within the Yandin Study area with flight activity and behaviour recorded in respect to the three zones above. A comprehensive impact assessment including a risk assessment of bird strike with turbines was undertaken.

The objectives of the current survey were to provide an up-to-date account of the birds of conservation significance potentially occurring over the Yandin Study area (given status changes since the RPS, 2010 study), and to augment the avian records obtained by RPS (2010).

Avifauna were recorded opportunistically during the current survey at 22 census sites, as part of the associated flora and vegetation study, in close proximity to the proposed wind turbine locations as well as over the entire study area. Heights and flight patterns were recorded for each species in consideration of the three zones above, and compared with the results of RPS (2010). The NatureMap database was scrutinised to obtain a current list of birds of conservation significance potentially occurring over the study area and a likelihood assessment undertaken as to their potential to occur, given the results of regional records, the previous work of RPS (2010) and the results of the current survey.

3.4 STUDY TEAM AND LICENCES

The flora, vegetation, and fauna assessments undertaken by *ecologia* was planned, coordinated, executed and reported by those summarised below in Table 3.4. DBCA licences to take flora and fauna for scientific purposes are also provided.

Table 3.4 – Study team and licences

Project Staff			
Name	Qualification	Role	Project role
Dr Andrew Craigie	B.Sc (Hons.), PhD (Botany)	Senior Botanist	Flora & vegetation assessment, plant specimen identification, reporting (flora)
Andre Schmitz	B.Sc. Env. Man.	Principal Ecologist	Vegetation and avifauna assessment, reporting (fauna)
Licences - "Licence to Take Flora for Scientific Purposes"			
The flora, vegetation assessment and fauna assessment described in this report was conducted under the authorisation of the following licences issued by DBCA:			
Name	Licence Number	Valid until	
Dr Andrew Craigie	SL012097	30/04/2018	

4 DESKTOP RESULTS

4.1 CLIMATE

The closest Bureau of Meteorology (BOM) weather station to the study area is the Badgingarra Research Station, approximately 40 km to the north. Based on these data, the study area experiences a typical dry Mediterranean climate with a hot dry period from December to March and a mild winter from June to August (Figure 4.1). The current survey was conducted in mid-September 2016, following a higher than average period of rainfall during August (Figure 4.1).




Figure 4.1 – Climate data from Badgingarra BOM weather station (Station No. 009037)

4.2 IBRA 7 BIOGEOGRAPHIC SUBREGIONS

The Interim Biogeographic Regionalisation for Australia (IBRA) (Version 7) classifies the Australian continent into regions (bioregions) of similar geology, landform, vegetation, fauna and climate characteristics, and has currently 89 recognised regions (DSEWPaC 2012). The study area is located on the boundary of two Bioregions; Swan Coastal Plain and Geraldton Sandplains.

Bioregions are further delineated into subregions, with the study area located on the junction of the Dandgaragan Plateau (SWA01), Swan Coastal Plain (SWA02), and Lesueur Sandplain (DES02) subregions (Figure 4.2).

The Dandgaragan Plateau subregion is bordered by the Derby and Dandaragan Faults, covering an area of approximately 447, 862 ha. Vegetation is typically dominated by low *Banksia* woodland, Jarrah-Marri woodlands, and scrub-heaths on laterite pavement and on gravelly sandplains (Desmond 2001). The subregion hosts a large number of rare plant and animal species, and supports a number of significant wetlands (Wannamal Lake System) and ecosystems (Desmond 2001), including the 'Banksia Woodlands of the Swan Coastal Plain' TEC.

The Swan Coastal Plain subregion is characterised by a low lying coastal plain of approximately 1,333,901 ha, mainly covered by woodlands dominated by *Banksia* or Tuart on sandy soils. *Casuarina obesa* is characteristic on outwash plains and *Melauleca* species are typical of swampy areas. In the


east, Jarrah woodland occur on elevated plains (Mitchell *et al.* 2002). The subregion exhibits very high species and ecosystem diversity, and supports numerous significant ecological communities, wetlands and other landscape features (Mitchell *et al.* 2002).


The Lesueur Sandplains subregion covers a total area of approximately 1,358,915 ha, comprising coastal aeolian sands and limestones, siltstones and sandstones, with extensive yellow sandplains in the south and east. Mosaics of lateritic mesas, sandplains, and coastal sands and limestones support shrub-heaths that are rich in endemic plant species, with heath occurring on the lateritised sandplains of the north and east eastern margins (Desmond and Chant 2003).

4.3 SOILS

Five soil units have been mapped within the study area using the Digital Atlas of Australian Soils (Northcote *et al.* 1960-1968) (Figure 4.3), none of which are considered to be restricted:

- AB3 'Undulating to low hilly dissected plateau slopes often flanking areas of unit AC2, or occupying a zone between units AC2 and AB4: chief soils on the slopes are red earthy sands (Uc5.21). Associated are (Uc5.22) soils on ridges and (Uc2.21) soils in the centre of valleys, apart from limited areas of swamps in which occur diatomaceous earths' (Northcote *et al.* 1960-1968).
- AB4 'Slopes flanking main trunk valleys; breakaways are common. There are two common sequences of soils: (i) on smooth slopes below breakaways red earthy sands (Uc5.21) occur, occasionally with (Gn2.12) soils and sometimes (Uc2.21) soils in the adjacent valley floors; (ii) on dissected slopes below breakaways red earthy sands (Uc5.21) occur but with some (Um6.21) soils on chalk outcrops, and further down slope are areas of (Dr), (Db), (Dy), and (Dd) soils, such as (Dr4.22), (Db2.22), (Dy3.12), and (Dd2.43) while adjacent valley floors also have (Dr) and (Db) soils. Areas of diatomaceous earths occur on some valley floors' (Northcote *et al.* 1960-1968).
- AC2 'Gently undulating plateau underlain by sedimentary rocks: chief soils are yellow earthy sands (Uc5.22) with siliceous sands (Uc1.22). Associated are patches of (KS-Uc2.12) and (Dy5.84) soils; and (Uc2.21) soils in some shallow valley floors' (Northcote *et al.* 1960-1968).
- Wd9 'Broad valleys and undulating interfluvial areas with some discontinuous breakaways and occasional mesas; lateritic materials mantle the area: chief soils are sandy acidic yellow mottled soils, (Dy5.81) containing much ironstone gravel in the A horizons, and (Dy5.84), both forming a complex pattern with each other and with lateritic sandy gravels (KS-Uc2.12). Associated are leached sands (Uc2.21) underlain by lateritic gravels and mottled clays that occur at a progressively greater depth down slope' (Northcote *et al.* 1960-1968).
- Wd10 'Broad valleys and undulating interfluvial areas; some evenly sloping pediments with exposures of sandstone and shale: chief soils are sandy acidic yellow mottled soils, (Dy5.81) containing much ironstone gravel in the A horizons, and (Dy5.84), both forming a complex pattern with each other and with lateritic sandy gravels (KS-Uc2.12). Associated are leached sands (Uc2.21) underlain by lateritic gravels, and mottled clays that occur about 3 ft in depth and are shallower than in unit Wd9. Other soils include (Dy3.71), (Dy3.81), (Dy5.41) as well as (Uc2.21 and Uc2.22) on the pediments; and (Dr3.32), (Dy3.32), and (Dy3.22) in areas where country rock has been exposed' (Northcote *et al.* 1960-1968).






Yandin Wind Farm Study Area

Map Unit

- AB3
- AB4
- AC2
- AC5
- AC6
- Ca27
- Cb39
- JK9
- Ub97
- Wd10
- Wd9



Drawn: AC Project ID: 1713
Date: 17 August 2017 A4

Soil units (Northcote et al. 1960-1968)

4.4 VASCULAR FLORA

4.4.1 Floristic Diversity

A total of 485 vascular plant taxa (including species, infraspecific taxa, and phrase name taxa) have been recorded within 10 km of the study area (NatureMap) representing 59 families and 189 genera (Appendix B). The most diverse families were the Fabaceae (62 taxa), Proteaceae (58 taxa), Myrtaceae (57 taxa), and Cyperaceae (25 taxa). The most diverse genus was *Acacia* (19 taxa), *Eucalyptus* (15 taxa), *Stylidium* (15 taxa), *Hakea* (14 taxa), *Banksia* (13 taxa), and *Grevillea* (13 taxa).

4.4.2 Significant Plant Species

The TPFL and WAHERB database searches identified 61 conservation significant plant species occurring within a 10 km buffer of the study area (Figure 4.4; Figure 4.5, Figure 4.6). Eighteen of these, including two Threatened taxa and 14 Priority listed taxa have been recorded within the study area (Table 4.1). For the remaining taxa, due to the close proximity of previous records and habitat preferences, all are considered to have potential to occur within the study area.


Habitat preferences and flowering times, as indicated in Table 4.1 were derived, where available, from relevant taxonomic literature, FloraBase (Western Australian Herbarium 1998-2016), Threatened species profiles (SPRATs) (Threatened Species Scientific Committee 2016b), or specimen data from Australia's Virtual Herbarium (AVH) (CHAH 2017). Herbarium catalogue numbers are provided if habitat information were derived from specimen data (AVH).

Table 4.1 – Threatened and Priority Flora records from database searches

Status	Taxon	Habitat	Flowering Time	Recorded within the study area
T	<i>Acacia forrestiana</i>	Lateritic gravelly soils, clay loam over sandstone. Gullies, hills, breakaways.	Nov to Dec	Possible
T	<i>Acacia splendens</i>	White sand over clay, pale brown loam, cracked brown soil, gravel, laterite, ironstone. Slopes of breakaways, especially southern slopes, hills.	May	Possible
T	<i>Chamelaucium</i> sp. Cataby (G.J. Keighery 11009)	Growing amongst laterite rocks on W-facing edge of breakaway (MEL 2013770A)	–	Recorded
T	<i>Drakaea elastica</i>	White or grey sand. Low-lying situations adjoining winter-wet swamps.	Oct to Nov	Possible
T	<i>Eleocharis keigheryi</i>	Clay, sandy loam. Emergent in freshwater: creeks, claypans.	Aug to Nov	Possible
T	<i>Eucalyptus dolorosa</i>	Laterite. Hillsides.	Feb to Mar	Possible
T	<i>Grevillea calliantha</i>	Grey or yellow sand over laterite, with gravel.	Apr or Jun or Aug	Recorded
T	<i>Macarthuria keigheryi</i>	White or grey sand.	Sep to Dec or Feb to Mar	Possible
T	<i>Ptychosema pusillum</i>	Sand. Rises.	Aug to Oct.	Possible
T	<i>Thelymitra stellata</i>	Sand, gravel, lateritic loam.	Oct to Nov	Possible
P1	<i>Babingtonia delicata</i>	Valley flat, slope. Pale grey sand / brown sandy clay. Winter wet area (PERTH 3255166)	–	Possible
P1	<i>Banksia prionophylla</i>	Dry grey sand over laterite with surface boulders. Rises.	Jul	Recorded
P1	<i>Eucalyptus annuliformis</i>	Shallow sandy soils. Rocky laterite slope.	May to Sep	Possible
P1	<i>Grevillea synapheae</i> subsp. A Flora of Australia (S.D. Hopper 6333)	Gravelly loam.	Sep	Possible
P1	<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	Gravel, laterite.	Aug to Sep	Possible
P1	<i>Hypocalymma</i> sp. Dandaragan (C.A. Gardner 9014)	In grey sand with lateritic pebbles (PERTH 2353539)	–	Possible
P1	<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	Hilltop, dry brown loam over laterite boulder (PERTH 3022080)	–	Possible
P1	<i>Rhedinocarpha suffruticosa</i>	Red-brown loamy clay, gravelly loam or clay loam over laterite. Slopes, small ridges.	–	Possible
P2	<i>Anigozanthos humilis</i> subsp. Badgingarra (S.D. Hopper 7114)	Grey-white sand, rich brown sandy loam, sandy clay, alluvial soils. Low plains, river-banks, winter-wet swamps.	–	Recorded

Status	Taxon	Habitat	Flowering Time	Recorded within the study area
P2	<i>Boronia scabra</i> subsp. <i>condensata</i>	Sandy clay or gravel. Upper slopes, edges of lateritic breakaways.	Aug	Possible
P2	<i>Dampiera tephrea</i>	Sand, gravelly loam.	Jul	Recorded
P2	<i>Desmocladius microcarpus</i>	–	–	Possible
P2	<i>Eucalyptus abdita</i>	Laterite, sandy clay with gravel over laterite. Slopes, breakaways.	–	Recorded
P2	<i>Gastrolobium nudum</i>	Red-brown clay, brown loam, gravel, laterite, granite. Flats, slopes, hilltops, ridges, valleys, breakaways.	Feb	Recorded
P2	<i>Hemigenia curvifolia</i>	Sandy soils.	Sep to Oct	Possible
P2	<i>Hypocalymma</i> sp. Cataby (G.J. Keighery 5151)	Grey sand.	Aug	Possible
P3	<i>Acacia cummingiana</i>	Grey or yellow sand, lateritic gravel. Sandplains, lateritic breakaways.	May to Jun or Aug	Possible
P3	<i>Acacia epacantha</i>	Lateritic gravelly loam or clay.	Jul to Aug	Possible
P3	<i>Acacia plicata</i>	Loamy & clayey soils, often over sandstone or siltstone. Along drainage lines.	Aug to Oct	Possible
P3	<i>Banksia dallanneyi</i> subsp. <i>pollostata</i>	Grey/yellow sand. Flats, lateritic rises.	Aug to Sep	Possible
P3	<i>Banksia kippistiana</i> var. <i>paenepeccata</i>	Lateritic gravelly soils.	Oct to Nov	Possible
P3	<i>Banksia pteridifolia</i> subsp. <i>vernalis</i>	White/grey sand over laterite	Sep to Oct	Possible
P3	<i>Beaufortia eriocephala</i>	Lateritic sandy soils. Slopes.	Sep to Nov	Possible
P3	<i>Calytrix ecalycata</i> subsp. <i>brevis</i>	Dry yellow sand. Sandplains, low rises.	Aug to Sep	Possible
P3	<i>Drosera marchantii</i> subsp. <i>prophylla</i>	Laterite-silica sand soils. Hilltops.	Jun to Jul	Possible
P3	<i>Grevillea florida</i>	Sand, sandy clay, gravel, laterite. Sandplain, slopes, road verges.	Jul to Sep	Possible
P3	<i>Grevillea thyrsoides</i> subsp. <i>thyrsoides</i>	Sand or sandy lateritic gravel.	Feb or Aug to Sep	Possible
P3	<i>Guichenotia alba</i>	Sandy & gravelly soils. Low-lying flats, depressions.	Jul to Aug	Possible
P3	<i>Haemodorum loratum</i>	Grey or yellow sand, gravel.	Nov	Possible
P3	<i>Hakea longiflora</i>	White sand, loam, gravel, laterite. Breakaway.	Jun to Sep	Recorded
P3	<i>Hypocalymma tetrapterum</i>	Grey sand, loam, lateritic gravel. Riverbanks, breakaways.	Aug	Recorded
P3	<i>Jacksonia carduacea</i>	Grey sand, sandy clay.	Aug to Dec	Recorded
P3	<i>Lechenaultia galactites</i>	Yellow sand, clay, gravel, laterite. Sandplains.	Jun to Oct	Possible
P3	<i>Lepidobolus quadratus</i>	Lateritic gravel, grey/white sand. Dry kwongan.	Aug to Sep	Possible
P3	<i>Leucopogon foliosus</i>	Dry grey/yellow sand and laterite (PERTH 5984475)	–	Possible

Status	Taxon	Habitat	Flowering Time	Recorded within the study area
P3	<i>Podotheca pritzelii</i>	Sand ridges in salt flats.	Sep to Oct	Recorded
P3	<i>Stylidium periscelanthum</i>	Loamy clay, moist soils pockets. Wet flats, low granitic hills.	Sep to Oct	Possible
P3	<i>Verticordia huegelii</i> var. <i>tridens</i>	Sandy or gravelly loam. Winter-wet areas, low hills.	Sep to Nov	Possible
P3	<i>Verticordia insignis</i> subsp. <i>eomagis</i>	Sandy soils over laterite. Sandplains, rocky rises.	Aug to Nov	Possible
P4	<i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i>	Grey or yellow sand.	Jul to Oct	Recorded
P4	<i>Asterolasia drummondii</i>	Lateritic gravel & sand or loam. Lateritic hills & sandplains, breakaways.	Jul to Sep	Recorded
P4	<i>Conostephium magnum</i>	White-grey sands sometimes associated with laterite gravels. Sand dunes, swampland, disturbed roadside, drainage channels, open woodland.	Jul to Sep	Possible
P4	<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	White or grey sand over laterite. Hillslopes, ridges, sandplains.	Aug to Sep or Nov to Dec	Recorded
P4	<i>Grevillea drummondii</i>	Lateritic soils (sandy clay, gravel, loam, sand), sand over granite. Rocky hillsides, boulders, granite outcrops.	Jun to Sep	Recorded
P4	<i>Grevillea olivacea</i>	White or grey sand. Coastal dunes, limestone rocks.	Jun to Sep	Possible
P4	<i>Grevillea saccata</i>	Yellow or brown sand, often with lateritic gravel.	Apr or Jun to Nov	Possible
P4	<i>Hypolaena robusta</i>	White sand. Sandplains.	Sep to Oct	Recorded
P4	<i>Stylidium aeonioides</i>	Sandy clay loam over laterite. Hillsides and breakaways. Low heath, open woodland.	Sep to Nov	Recorded
P4	<i>Thelymitra apiculata</i>	Grey sand, lateritic gravel.	May to Jul	Recorded
P4	<i>Thysanotus glaucus</i>	White, grey or yellow sand, sandy gravel.	Oct to Dec or Jan to Mar	Possible
P4	<i>Verticordia lindleyi</i> subsp. <i>lindleyi</i>	Sand, sandy clay. Winter-wet depressions.	May or Nov to Dec or Jan	Possible



Taxon


- *Acacia forrestiana*
- *Acacia splendens*
- *Chamelaucium* sp. *Cataby* (G.J. Keighery 11009)
- *Drakaea elastica*
- *Eleocharis keigheryi*
- ▲ *Eucalyptus dolorosa*
- ▲ *Grevillea calliantha*
- ▲ *Macarthuria keigheryi*
- ▲ *Ptychosema pusillum*
- ▲ *Thelymitra stellata*
- Yandin Wind Farm Study Area

Drawn: AC Project ID: 1713
Date: 17 August 2017 A4


0 1.5 3
Kilometers
Scale: 1:150,000
MGA94 (Zone 50)

DBCA Threatened Flora records

Figure:
4.4



- Taxon**
- *Anigozanthos humilis* subsp. *Badgingarra* (S.D. Hopper 7114)
 - *Babingtonia delicata*
 - *Banksia prionophylla*
 - *Boronia scabra* subsp. *condensata*
 - *Dampiera tephrea*
 - *Desmocladius microcarpus*
 - ▲ *Eucalyptus abdita*
 - ▲ *Eucalyptus annuliformis*
 - ▲ *Gastrolobium nudum*
 - ▲ *Grevillea synapheae* subsp. *A Flora of Australia* (S.D. Hopper 6333)
 - ▲ *Grevillea synapheae* subsp. *minyulo*
 - ▲ *Hemigenia curvifolia*
 - ★ *Hypocalymma* sp. *Cataby* (G.J. Keighery 5151)
 - ★ *Hypocalymma* sp. *Dandaragan* (C.A. Gardner 9014)
 - ★ *Lasiopetalum* sp. *Hill River* (T.N. Stoate 5)
 - ★ *Rhetinocarpha suffruticosa*
 - Yandin Wind Farm Study Area



- Taxon**
- Acacia cummingiana
 - Acacia epacantha
 - Acacia plicata
 - Anigozanthos humilis subsp. chrysanthus
 - Asterolasia drummondii
 - Banksia dallanneyi subsp. pollostia
 - Banksia kippistiana var. paenepeccata
 - Banksia pteridifolia subsp. vernalis
 - Beaufortia eriocephala
 - Calytrix ecalycata subsp. brevis
 - Conostephium magnum
 - Drosera marchantii subsp. prophylla
 - ▲ Eucalyptus macrocarpa subsp. elachantha
 - ▲ Grevillea drummondii
 - ▲ Grevillea florida
 - ▲ Grevillea olivacea
 - ▲ Grevillea saccata
 - ▲ Grevillea thyrsoides subsp. thyrsoides
 - + Guichenotia alba
 - + Haemodorum loratum
 - + Hakea longiflora
 - + Hypocalymma tetrapterum
 - + Hypolaena robusta
 - + Jacksonia carduacea
 - ✱ Lechenaultia galactites
 - ✱ Lepidobolus quadratus
 - ✱ Leucopogon foliosus
 - ✱ Podotheca pritzelii
 - ✱ Styliidium aeonioides
 - ✱ Styliidium periscelanthum
 - ✕ Thelymitra apiculata
 - ✕ Thysanotus glaucus
 - ✕ Verticordia huegelii var. tridens
 - ✕ Verticordia insignis subsp. eomagis
 - ✕ Verticordia lindleyi subsp. lindleyi
 - Yandin Wind Farm Study Area

4.5 VEGETATION

4.5.1 Pre-European Vegetation


The pre-European vegetation of Western Australia was mapped at the 1:1,000,000 scale by Beard (1976), and was subsequently reinterpreted and updated to reflect the National Vegetation Information System (NVIS) standards (Shepherd *et al.* 2001). Five vegetation associations of Beard (1976) have been mapped within the area (Figure 4.7):

- Association 4: *Corymbia calophylla*, *Eucalyptus wandoo*, *Acacia acuminata* mid woodland over *Acacia nervosa*, *Gastrolobium* sp., *Hakea lissocarpha* tall open shrubland.
- Association 999: *Corymbia calophylla*, *Eucalyptus loxophleba*, *Acacia cyanophylla* mid woodland, over *Acacia pulchella*, *Boronia scabra*, *Bossiaea* sp. tall open shrubland, over *Hibbertia hypericoides*, *Hybanthus calycinus*, *Lechenaultia biloba* low open shrubland/forbland.
- Association 1030: *Banksia attenuata*, *Banksia menziesii*, *Eucalyptus todtiana* low woodland, over *Adenanthos cygnorum*, *Allocasuarina humilis*, *Jacksonia furcellata* tall open shrubland, over *Anigozanthos humilis*, *Conostylis aculeata*, *Eremaea fimbriata* low open shrubland.
- Association 1031: *Xanthorrhoea reflexa* tall open shrubland, over *Dryandra (Banksia) bipinnatifida*, *Hakea auriculata*, *Dryandra (Banksia) shuttleworthiana* mid shrubland, over *Banksia* sp., *Burchardia umbellata*, *Calectasia cyanea* low open shrubland/forbland.
- Association 1035: *Corymbia calophylla* mid open woodland.

The pre-European and current extent of each of these vegetation associations in the Swan Coastal Plain and Geraldton Sandplains bioregions, and within the study area, is presented in Table 4.2. The vegetation over most of the study area has been historically cleared, and therefore the area of each mapped within it does not accurately reflect current extent. A significant proportion (84.7%) of vegetation association 1035 within the Geraldton Sandplains Bioregion (pre-European) occurs within the study area; however, this association is very broadly defined as *Corymbia calophylla* mid open woodland.

Table 4.2 – Pre-European vegetation association extent

Vegetation Association	IBRA region	Pre-European extent (ha)	Current extent (ha)	% remaining	Area mapped in study area (ha)	Proportion of pre-European extent within study area (%)
4	GES	5,336.70	2,130.04	39.91	40.8	0.8
999		1,095.04	390.37	35.65	25.3	2.3
1030		3,848.52	2,790.59	72.51	NA	NA
1031		241,349.97	83,154.99	34.45	2270.6	0.9
1035		1,582.96	133.16	8.41	1340.9	84.7
4	SWA	15,897.08	3,010.45	18.94	NA	NA
999		102,939.79	9,609.84	9.34	10967.4	10.7
1030		134,788.56	86,061.30	63.85	606.0	0.4
1031		27,729.97	5,352.64	19.30	NA	NA
1035		3,435.37	360.96	10.51	NA	NA



Pre-European vegetation (Beard 1976)

Figure:

4.7




4.6 SIGNIFICANT ECOLOGICAL COMMUNITIES

The EPBC Act listed (Endangered) ‘Banksia woodlands of the Swan Coastal Plain’ threatened ecological community has been recorded within the south-western boundary of the study area, and also in surrounding areas to the south-west and east (Table 4.3; Figure 4.8). This community is Priority 3 listed at a State level (DPaW 2017). No other TECs or PECs have been recorded within 10 km of the study area.


Table 4.3 – TECs in the vicinity of the study area

Name	EPBC Act Listing Status	WA Status	Description
Banksia woodlands of the Swan Coastal Plain	Endangered	Priority 3	‘Canopy is most commonly dominated or co-dominated by <i>Banksia attenuata</i> and/or <i>B. menziesii</i> . Other <i>Banksia</i> species that can dominate in the community are <i>B. prionotes</i> or <i>B. ilicifolia</i> . It typically occurs on well drained, low nutrient soils on sandplain landforms, particularly deep Bassendean and Spearwood sands and occasionally on Quindalup sands; it is also common on sandy colluvium and aeolian sands of the Ridge Hill Shelf, Whicher Scarp and Dandaragan Plateau and, in other less common scenarios’ (DPaW 2017).



Yandin Wind Farm Study Area
 Banksia Woodlands of the Swan Coastal Plain (DBCA)

ecologia
 ENVIRONMENT
 Drawn: AC Project ID: 1713
 Date: 17 August 2017 A4



Banksia Woodlands of the Swan Coastal Plain TEC (DBCA September 2017)

Figure:
4.8

4.7 PREVIOUS SURVEYS WITHIN THE STUDY AREA

4.7.1 Vegetation and Flora Assessments

Outback Ecology (2009)

Outback Ecology (2009) undertook a Level 1 Vegetation and Flora assessment at the Yandin Wind Farm project area (not corresponding exactly to the current study area) in November 2008 and January 2009, the purpose of which was to verify the findings of a desktop study, and to characterise the flora and vegetation communities present, including a targeted search for conservation significant plant species.

48 plant taxa were recorded within the study area from 16 families and 32 genera, one of which was a Priority Flora species (*Acacia plicata*), and four were introduced (*Cyperus congestus*, *Juncus acutus* subsp. *acutus*, *Typha orientalis*, and *Zantedeschia aethiopica*).

35 sampling points were assessed for vegetation condition. At all points, vegetation was considered degraded, occurring in 'parkland cleared' areas, occasionally with scattered native trees. An additional 14 relevés were assessed within remnant patches of native vegetation, within which 12 vegetation communities were described. Native vegetation patches were primarily dominated by eucalypt woodland communities, and were in degraded to excellent condition. An additional 11 relevés were surveyed along proposed access tracks, cable routes, and overhead transmission lines. At these sites, seven vegetation communities were described, which were primarily in degraded condition.

4.7.2 Avifauna Assessment

RPS (2010)

RPS Environment and Planning Pty Ltd undertook an avifauna assessment proposed Yandin Wind Farm Development in 2010 (RPS 2010). RPS (2010) undertook three separate targeted and systematic avifauna field surveys within the study area, utilising 27 bird census points. In addition, RPS (2010) also conducted comprehensive surveys at the nearby Waddi Wind Farm, as well as around wetland habitats in the wider locality to determine regional status, and movements of local waterbird populations and to assess habitat potential for migratory wading bird species.

Sixty-one species were recorded within the Yandin Study area by RPS (2010) with flight activity and behaviour recorded in respect to potential impact with wind turbines. A comprehensive impact assessment was undertaken that included a risk assessment of bird strike.

5 FIELD SURVEY RESULTS

5.1 FLORA

5.1.1 Flora of the Study Area

A total of 117 sub-generic vascular plant taxa (including species, infraspecific taxa, and phrase names) from 35 families were recorded from the study area during the current survey. A list of taxa recorded is included in Appendix G. This list is not considered to be comprehensive for the study area, and includes only taxa recorded within quadrats, dominant species at check sites, and some taxa recorded opportunistically. The most diverse families recorded were the Proteaceae (22 taxa), Myrtaceae (15 taxa), Fabaceae (10 taxa), and Poaceae (9 taxa).


5.1.2 Conservation Significant Plant Species

No EPBC Act (1999) listed Threatened Flora taxa or any WC Act (1950) listed Threatened Flora taxa were recorded within the study area during the current survey. One State Listed Priority Flora species (*Hypocalymma tetrapterum* P3) was recorded during the current survey, with targeted searches focussed on native vegetation patches in the vicinity of proposed development areas.

Hypocalymma tetrapterum (Myrtaceae) is a low-growing perennial shrub primarily restricted to the northern Swan Coastal Plain and southern Geraldton Sandplains regions. It was recorded from two locations in the west of the study area, on loamy soils in degraded *Eucalyptus wandoo* woodland, and on an adjacent lateritic rise dominated by *Xanthorrhoea priessii* and mixed low shrubs. Individual locations are provided in Table 5.1, and mapped in Figure 5.1.


Table 5.1 – *Hypocalymma tetrapterum* (P3) records within the study area

Taxon	Easting	Northing	Estimated Abundance
<i>Hypocalymma tetrapterum</i> (P3)	365629.108	6597181.477	5
	366021.159	6596807.245	20-50
	366025.597	6596829.052	20-50
	366039.707	6596835.324	20-50
	366055.643	6596834.752	20-50



Yandin Wind Farm Study Area
● *Hypocalymma tetrapterum* (P3)

ecologia
 ENVIRONMENT
 Drawn: AC Project ID: 1713
 Date: 19 October 2017 A4



Priority Flora Locations (*ecologia* 2017)

5.2 VEGETATION

5.2.1 Vegetation Units and Vegetation Condition

Sixteen vegetation units within eight broader community types were described within the study area, which are summarised in Table 5.2 and mapped in Figure 5.2 to Figure 5.10. Vegetation condition is mapped in Figure 5.11.

Vegetation over 91.2% (14,015 ha) of the study area comprised scattered *Corymbia calophylla* (marri) or *Eucalyptus tottiana* (coastal blackbutt) trees over pasture weeds with no native understorey (vegetation unit W3) in ‘completely degraded’ condition. Low-lying areas occupied by **Juncus acutus* (spiny rush) rushland (R1), were also completely degraded, accounting for 0.6% (98.6 ha) of the study area. Plantations account for a further 3% (454.3 ha) of the study area.

Two species rich shrubland units in good to excellent condition were described from remnant vegetation patches, occurring primarily on lateritic rises and hills: *Banksia hewardiana* mid shrubland over mixed low shrubland (S1) accounts for 0.4% (56.8 ha) of the study area, and *Xanthorrhoea preissii* mid sparse shrubland over mixed low shrubland (S2) accounts for 0.6% (87.1 ha) of the study area.

Five eucalypt woodland units (*Eucalyptus rudis* woodland in creek lines [W11, W12], *Corymbia calophylla* and *Banksia attenuata* open woodland over pasture weeds [W6], *Eucalyptus wandoo* woodland [W13], and *Eucalyptus* sp. open mallee woodland [W10]), were in degraded condition due to historical clearing, heavy grazing, and/or the presence of abundant weed species. These account for 1% (156 ha) of the study area.

The remaining seven woodland units occurring in remnant vegetation patches broadly comprise *Corymbia calophylla* woodland, *Eucalyptus tottiana* woodland, and *Banksia attenuata* woodland, and are in good to excellent condition. These account for 3% (467.1 ha) of the study area, and include two units identified as the ‘Banksia Woodlands of the Swan Coastal Plain’ TEC (W1, W2).

Table 5.2 – Summary of vegetation units at the study area

Broad Community Type	Map Code	Vegetation Unit	Vegetation Condition	Banksia Woodlands TEC	Landforms	Sites	Area within study area (ha)	% of study area
* <i>Juncus acutus</i> rushland	R1	* <i>Juncus acutus</i> rushland with isolated <i>Acacia saligna</i> shrubs.	Completely Degraded		Valley	C2b	98.68	0.6%
<i>Banksia hewardiana</i> shrubland	S1	<i>Banksia hewardiana</i> (± <i>Banksia sessilis</i> , ± <i>Allocasuarina humilis</i> , ± <i>Xanthorrhoea preissii</i>) mid shrubland over mixed low shrubland.	Good - Excellent		Undulating Plains, Midslopes, Ridgetops	C17, C18, C31, C36, C42	56.88	0.4%
<i>Xanthorrhoea preissii</i> shrubland	S2	<i>Xanthorrhoea preissii</i> mid sparse shrubland over mixed low shrubland (± <i>Acacia pulchella</i> , <i>Hibbertia hypericoides</i> , <i>Leucopogon sprengeioides</i> , <i>Hakea incrassata</i> , <i>Hakea lissocarpha</i> , <i>Isopogon asper</i> , <i>Melaleuca ciliosa</i>).	Excellent		Midslopes, Ridgetops	C8, C9b, C10a, C16	87.19	0.6%
<i>Banksia attenuata</i> woodland	W1	<i>Banksia attenuata</i> (± <i>Corymbia calophylla</i> , <i>Eucalyptus tottiana</i> , <i>Banksia menziesii</i>) woodland to open woodland over <i>Hibbertia hypericoides</i> low sparse shrubland over <i>Mesomelaena pseudostygia</i> and <i>Desmocladius flexuosus</i> sparse sedgeland/herbland.	Good - Excellent	YES	Plains, Foothslopes, Midslopes	Q1, Q4, Q8, Q33, C41	201.91	1.3%
	W2	<i>Banksia attenuata</i> and <i>Banksia menziesii</i> woodland over <i>Mesomelaena pseudostygia</i> and <i>Schoenus grandiflorus</i> open sedgeland.	Good	YES	Plains, Foothslopes, Midslopes	C14, C15	10.10	0.1%
<i>C. calophylla</i> / <i>E. tottiana</i> woodland	W3	<i>Corymbia calophylla</i> and/or <i>Eucalyptus tottiana</i> open woodland/isolated trees over pasture weeds.	Completely Degraded - Degraded		Undulating Plains, Foothslopes, Midslopes	C4, C9a, C10b, C11b, C24, C26, C28, C29, C30, C35, C38, C39	14015	91.2%
	W4	<i>Corymbia calophylla</i> mid open woodland over <i>Banksia hewardiana</i> and <i>Xanthorrhoea preissii</i> mid open shrubland.	Good - Excellent		Midslopes, Ridgetops	C5b, C19, C23, C25, C32	108.33	0.7%
	W5	<i>Corymbia calophylla</i> woodland over <i>Xanthorrhoea preissii</i> and <i>Macrozamia fraseri</i> mid sparse shrubland over <i>Hibbertia racemosa</i> low sparse shrubland.	Good - Very Good		Midslopes, Ridgetops	C5c, C6, C7	82.15	0.5%
	W6	<i>Corymbia calophylla</i> and <i>Banksia attenuata</i> open woodland over pasture weeds.	Degraded		Midslopes	C11a, C12b, C13	8.47	0.1%
	W7	<i>Corymbia calophylla</i> mid open woodland over <i>Hibbertia hypericoides</i> low open shrubland over <i>Mesomelaena pseudostygia</i> open sedgeland	Excellent		Midslopes	C3a, C98	29.52	0.2%
	W8	<i>Eucalyptus tottiana</i> open woodland over <i>Allocasuarina humilis</i> and <i>Adenanthos cygnorum</i> open shrubland over <i>Mesomelaena tetragona</i> sparse sedgeland.	Excellent		Floodplains, Midslopes	C3b, C37	22.90	0.1%
	W9	<i>Corymbia calophylla</i> mid open woodland over <i>Xanthorrhoea preissii</i> mid sparse shrubland over <i>Calothamnus quadrifidus</i> , <i>Hibbertia hypericoides</i> , and <i>Leucopogon polymorphus</i> low shrubland.	Excellent		Ridgetop	C34	12.23	0.1%

Broad Community Type	Map Code	Vegetation Unit	Vegetation Condition	Banksia Woodlands TEC	Landforms	Sites	Area within study area (ha)	% of study area
<i>Eucalyptus</i> sp. mallee woodland	W10	<i>Eucalyptus</i> sp. open mallee woodland over pasture weeds.	Degraded		Ridgetop	C40	8.43	0.1%
<i>Eucalyptus rudis</i> woodland	W11	<i>Eucalyptus rudis</i> mid woodland over <i>Juncus acutus</i> open rushland.	Degraded		Valley	C1	38.68	0.3%
	W12	<i>Eucalyptus rudis</i> and <i>Melaleuca raphiophylla</i> mid woodland over <i>Zantedeschia aethiopica</i> herbland.	Degraded		Floodplains, Valleys	C37, C3b	46.27	0.3%
<i>Eucalyptus wandoo</i> woodland	W13	<i>Eucalyptus wandoo</i> woodland.	Degraded - Good		Floodplain, Midslopes	C5a, C12a	54.61	0.4%
Rehab	Rehab	Roadside Rehabilitation (<i>Eucalyptus</i> spp., <i>Acacia pulchella</i> , <i>Acacia saligna</i>).	NA		Plain	C99	24.41	0.2%
Plantation	Plantation	Plantation	Completely Degraded		Plain	NA	454.38	3.0%

5.2.2 ‘Banksia Woodlands of the Swan Coastal Plain’ TEC

Five patches of the ‘Banksia Woodlands of the Swan Coastal Plain’ Threatened Ecological Community were identified within the study area (Figure 5.12), in addition to two previously identified patches (Figure 4.8, DBCA database, Sept. 2017) (Table 5.3). Patches ECO1, ECO2, and ECO3 (vegetation unit W1) occur in remnant vegetation patches and are in ‘excellent’ condition (Keighery 1994), with few non-aggressive weed species present. Linear patches ECO4 (W1) and ECO5 (W2) occur along the verge of Yandin road and an adjacent fence line, and are in ‘good’ condition (Keighery 1994), due to partial clearing, heavy grazing, and weed invasion, although basic vegetation structure is intact. Based on vegetation structure and composition, patch size, and condition (Keighery 1994), each of these patches should qualify as the TEC.

Table 5.3 – Summary of ‘Banksia Woodlands of the Swan Coastal Plain’ patches within the study area


Patch Name	Area (ha)	Vegetation Unit	Vegetation Condition	Site
DBCA 1	139.19	W1	Excellent	Q1
DBCA 2	10.11	W1	Good - Excellent	NA
ECO 1	14.49	W1	Excellent	Q8
ECO 2	11.13	W1	Excellent	Q33
ECO 3	19.43	W1	Excellent	Q4
ECO 4	2.89	W1	Good	C41
ECO 5	5.84	W2	Good	C14, C15

36000

368000

376000

384000



A1

A2

A3

B1

B2

B3

C1

C2

6600000

6592000

ecologia
ENVIRONMENT



0 0.5 1
Kilometers
Scale: 1:70,000
MGA94 (Zone 50)

Drawn: AC Project ID: 1713
Date: 19 October 2017 A3

Vegetation units - Map Index

Figure:

5.2



6604000

6602000

6600000

Vegetation units - A1



Study area	W2
Flora survey site	W3
WTG Location	W4
Grid Connection Route	W5
Access Track	W6
Underground Cable	W7
Onsite Substation	W8
Iluka Substation	W9
Banksia Woodland TEC	W10
Vegetation Unit	W11
R1	W12
S1	W13
S2	Rehab
W1	Plantation

ecologia
ENVIRONMENT

Drawn: AC Project ID: 1713
Date: 19 October 2017 A3


0 0.25 0.5
Kilometers
Scale: 1:26,000
MGA94 (Zone 50)


Vegetation units - A2



376000 378000 380000 382000 384000
 6604000
 6602000
 6600000
 W12
 C38
 C35
 C30
 C1
 C2b
 W11
 W3
 R1
 R1
 R1
 R1
 Plantation
 Plantation
 R1


Vegetation units - A3






- Study area
 - Flora survey site
 - WTG Location
 - Grid Connection Route
 - Access Track
 - Underground Cable
 - Onsite Substation
 - Iluka Substation
 - Banksia Woodland TEC
- | | |
|--|--|
| <p>Vegetation Unit</p> <ul style="list-style-type: none"> R1 S1 S2 W1 W2 W3 W4 W5 W6 W7 W8 W9 W10 W11 W12 W13 Rehab Plantation | <ul style="list-style-type: none"> W2 W3 W4 W5 W6 W7 W8 W9 W10 W11 W12 W13 Rehab Plantation |
|--|--|

Vegetation units - B2



- | | |
|------------------------|------------|
| Study area | W2 |
| Flora survey site | W3 |
| WTG Location | W4 |
| Grid Connection Route | W5 |
| Access Track | W6 |
| Underground Cable | W7 |
| Onsite Substation | W8 |
| Iluka Substation | W9 |
| Banksia Woodland TEC | W10 |
| Vegetation Unit | |
| R1 | W11 |
| S1 | W12 |
| S2 | W13 |
| W1 | Rehab |
| | Plantation |

Vegetation units - B3




- Study area
 - Flora survey site
 - WTG Location
 - Grid Connection Route
 - Access Track
 - Underground Cable
 - Onsite Substation
 - Iluka Substation
 - Banksia Woodland TEC
- | | |
|--|--|
| <ul style="list-style-type: none"> W2 W3 W4 W5 W6 W7 W8 W9 W10 W11 W12 W13 Rehab Plantation | <ul style="list-style-type: none"> R1 S1 S2 W1 |
|--|--|


Drawn: AC Project ID: 1713
Date: 19 October 2017 A3

0 0.25 0.5
Kilometers
Scale: 1:26,000
MGA94 (Zone 50)




Vegetation units - C1



Vegetation units - C2




Study area
Average Condition
 Excellent
 Very Good
 Good
 Degraded
 Completely Degraded




 Scale: 1:70,000
 MGA94 (Zone 50)

Drawn: AC Project ID: 1713
 Date: 19 October 2017 A3

Vegetation condition



Banksia Woodlands of the Swan Coastal Plain TEC within the study area

5.3 AVIFAUNA

One hundred and twenty-six bird species have been recorded within 20 km of the study area (Appendix E). Of these regional records the most diverse families were the ducks and swans (Anatidae - 13 species), the parrots (Psittacidae - 12 species) and the honeyeaters (Meliphagidae - 12 species).

Sixty-one species were recorded within the Yandin study area by RPS (2010) and the current survey recorded 44 species from 22 census sites within seven habitat types as well as the overall study area (Table 5.4).

All birds recorded had previously been recorded by RPS (2010) during the regional study, apart from the Australasian Pipit (*Anthus novaeseelandiae*) and the Emu (*Dromaius novaehollandiae*)(one individual sighted).

Table 5.4 – Avifauna records within sites and habitats

Species	Parkland Cleared	Marri Woodland	Proteraceous Heath	Dryandra-Allocastraria Shrubland	<i>E. Todtiana</i> Woodland	Banksia Woodland	Creek-line	Overall study area
	C04, C09A, C28, C29, C35, C38, C40	C05, C09b, C13, C32	C10, C16, C18	C17, C31, C42	Q11, C19, C30	C41	C37	
Singing Honeyeater			2					4
Brown Honeyeater		3		4				
White-cheeked Honeyeater			6					12
New Holland Honeyeater				4				
Red Wattlebird		2		1				3
Western Little Wattlebird			1					
Yellow-throated Miner								3
Inland Thornbill		2						
Western Thornbill		1						
Yellow-rumped Thornbill					5			
Weebill	2	3						3
White-browed Scrubwren								
Striated Pardalote	2	1						
Western Gerygone		1						
Variiegated Fairy-wren			3					
Splendid Fairy-wren				4				
Grey Fantail		1						
Willie Wagtail	2							4
Silvereye					3			
Australasian Pipit								6
Rufous Whistler	1	2		1				
Grey Shrike-thrush								1
Black-faced Cuckoo-shrike								3
Black-faced Woodswallow								6
Magpie-lark	2					1	2	4
Australian Magpie	3							
Grey Butcherbird			1					
Australian Raven								7
Tree Martin	5							
Emu								1
Australian Ringneck	4	6	5		4		2	11

Species	Parkland Cleared	Marri Woodland	Proteraceous Heath	Dryandra-Allocasuarina Shrubland	<i>E. Todtiana</i> Woodland	Banksia Woodland	Creek-line	Overall study area
	C04, C09A, C28, C29, C35, C38, C40	C05, C09b, C13, C32	C10, C16, C18	C17, C31, C42	Q11, C19, C30	C41	C37	
Butler's Corella	21	10		6				30
Galah	6	4	6		5			25
Carnaby's Black-Cockatoo			22	16				17
Common Bronzewing								2
Fan-tailed Cuckoo								1
*Laughing Kookaburra								2
Stubble Quail								2
Square-tailed Kite								1
Black-shouldered Kite								2
Australian Kestrel		1						3
Australian Hobby								1
White-faced Heron								1
Straw-necked Ibis								8

Bird flight patterns and activity were assessed using the same parameters as RPS (2010) within Zones 1 to 3 (Table 5.5). Birds recorded with at least the potential to fly within the RSA potential collision zone (Zone 2) were; Straw-necked Ibis, Black-shouldered Kite, Carnaby's Black-Cockatoo, Butler's Corella, Galah, and Australian Kestrel.

All these species were similarly identified by RPS (2010). However the percentages of recordings within individual Zones differed for some species. This can be attributed to the smaller scale of the current study, and fewer observations, with an associated less robust data set. For example, just one observational record correlates with the highest percentage recording – “*Species commonly recorded at this height range during study (25% and >25%)*” whereas the record may potentially represent an aberration. Nevertheless, species recorded during this survey were generally observed within the same, or very similar zones, as RPS (2010) thereby verifying those results.

Table 5.5 – Avifauna records within flight zones

Species	Total Records	Zone 1	Zone 2	Zone 3	Habitat (of RPS 2010)	Movements	Notes
		0m - 40m	40m - 152m	>152m			
		Below tip of turbine	RSA: Potential collision	Above tip of turbine			
Butler's Corella	62				Woodland, Grassland	Locally nomadic	Large flocks at medium speed
Carnaby's Black-Cockatoo	38				Forest/Woodland/Heath	Seasonally Nomadic	Large flocks at slow speed
Australian Ringneck	21				Forest/Woodland	Sedentary	Swift
Galah	21				Woodland/Grassland	Locally nomadic	Medium flocks at medium speed
Straw-necked Ibis	8				Wetlands/Grassland	Locally nomadic	Large flocks at slow speed
Brown Honeyeater	7				Forest/Woodland/Shrubland/Heath	Nomadic-Dispersive	Slow-Medium speed
Australian Raven	7				Woodland/Grassland	Locally nomadic	Slow-Medium speed
White-cheeked Honeyeater	6				Shrubland/Heath	Nomadic-Dispersive	Slow-Medium speed
Australasian Pipit	6				Grassland	Seasonally Nomadic	Slow-Medium speed
Black-faced Woodswallow	6				Forest/Woodland	Locally nomadic	Slow-Medium speed
Yellow-rumped Thornbill	5				Woodland/Grassland	Sedentary	Slow speed
Weebill	5				Forest/Woodland	Locally nomadic	Slow speed
Magpie-lark	5				Forest/Woodland	Sedentary	Slow speed
Tree Martin	5				Aerial proximate to nesting sites	Seasonally Nomadic	Swift
New Holland Honeyeater	4				Woodland/Shrubland/Heath	Nomadic-Dispersive	Medium speed
Splendid Fairy-wren	4				Shrubland/Heath	Sedentary	Slow speed
Rufous Whistler	4				Forest/Woodland/Shrubland	Locally nomadic	Medium speed
Australian Kestrel	4				Grassland/Woodland edges	Sedentary/Dispersive	Slow to swift
Red Wattlebird	3				Forest/Woodland/Shrubland/Heath	Seasonally Nomadic	Slow-Medium speed
Yellow-throated Miner	3				Woodland	Seasonally Nomadic	Slow-Medium speed
Striated Pardalote	3				Forest/Woodland	Seasonally Nomadic	Slow-Medium speed
Variegated Fairy-wren	3				Shrubland/Heath	Sedentary	Slow speed
Silvereye	3				Forest/Woodland/Shrubland/Heath	Seasonally Nomadic	Slow-Medium speed
Black-faced Cuckoo-shrike	3				Forest/Woodland	Seasonally Nomadic	Medium speed
Australian Magpie	3				Woodland/Grasslands	Sedentary	Medium speed
Singing Honeyeater	2				Forest/Woodland/Shrubland/Heath	Nomadic-Dispersive	Medium speed
Inland Thornbill	2				Forest/Woodland/Shrubland	Sedentary	Slow speed
White-browed Scrubwren	2				Shrubland/Heath	Sedentary	Slow speed

Species	Total Records	Zone 1	Zone 2	Zone 3	Habitat (of RPS 2010)	Movements	Notes
		0m - 40m	40m - 152m	>152m			
		Below tip of turbine	RSA: Potential collision	Above tip of turbine			
Willie Wagtail	2				Woodland/Grasslands	Sedentary	Slow speed
Common Bronzewing	2				Forest/Woodland	Sedentary	Medium to swift speed
*Laughing Kookaburra	2				Forest/Woodland	Sedentary	Medium speed
Stubble Quail	2				Woodland, Grassland	Nomadic/Irruptive	Medium to swift speed
Black-shouldered Kite	2				Grassland/Woodland edges	Locally nomadic/dispersive	Slow to swift
Western Little Wattlebird	1				Forest/Woodland/Shrubland/Heath	Seasonally Nomadic	Medium speed
Western Thornbill	1				Forest/Woodland	Sedentary	Slow speed
Western Gerygone	1				Forest/Woodland	Sedentary	Slow speed
Grey Fantail	1				Forest/Woodland	Seasonal Migrant	Slow speed
Grey Shrike-thrush	1				Forest/Woodland	Sedentary	Medium speed
Grey Butcherbird	1				Forest/Woodland	Sedentary	Medium speed
Emu	1				Woodland, Grassland	Seasonally Nomadic	Medium speed
Fan-tailed Cuckoo	1				Forest/Woodland	Seasonal Migrant	Medium speed
Square-tailed Kite	1				Grassland/Woodland edges	Seasonal Migrant	Medium speed
Australian Hobby	1				Woodland edges	Sedentary/Dispersive	Medium to swift
White-faced Heron	1				Grasslands/Wetlands	Nomadic	Medium speed

Legend

	Species commonly recorded at this height range during study (25% and >25%)
	Species occasionally recorded at this height range during study (>5% and < 25%)
	Species rarely recorded at this height range during study (> 0% and up to 5%)
	Species not recorded at this height range during study (0%)

Fifteen birds of conservation significance were identified by NatureMap (2017) as occurring within 20 km of the study area. A likelihood of occurrence over the study area was undertaken using the criteria presented in Table 3.2.

All species identified by NatureMap (Table 5.6) were also assessed by RPS (2010), apart from the inland sub-species of the Western Rosella (P4). The Western Rosella was not recorded during this survey or by RPS (2010) however a historical record from the vicinity is included in NatureMap (2017). Although habitat at the study area is fragmented, and no recent records have been obtained, there is the possibility that this species may occur. However it is unlikely that the Western Rosella would fly within the RSA of Zone 2 and is therefore at little risk from the project.

Apart from the Western Rosella, RPS (2010) undertook a significant risk assessment of all the species listed in Table 5.6, and results conform with the likelihood of occurrence table presented here.

Note that the likelihood of occurrence table (Table 5.6) refers a species likelihood of occurrence at the study area itself, however, birds may still be at risk flying over the study area to and from suitable habitat. For example migratory wading birds have been recorded from wetland habitats around Lake Guraga to the south-west, Lake Thetis near Cervantes and from the Upper Moore River area. Movements to and from these wetland habitats by migratory wading birds would occur, and RPS (2010) consider these factors in their impact assessment and risk assessment of bird strike.

Table 5.6 – Regional records of avifauna of conservation significance

Species	WA Status	EPBC Status	Notes	Likelihood of occurrence
Carnaby's Black-Cockatoo	EN	EN	Recorded regularly in the Study area and region	Recorded
Western Rosella (inland)	P4	-	Not recorded - historical record from Dandaragan	Possible
Malleefowl	VU	VU	Not recorded. Habitat not present.	Low
Peregrine Falcon	OS	-	Recorded in the region by RPS (2010)	Recorded
Rainbow Bee-eater	IA	-	Recorded regularly in the Study area and region	Recorded
Wetland/Estuarine Species				
Blue-billed Duck	P4	-	Not recorded. Deep water habitat not present	Low
Great Egret	IA	-	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Glossy Ibis	IA	-	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Red-necked Stint	IA	IA	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Curlew Sandpiper	VU	CR	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Sharp-tailed Sandpiper	IA	IA	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Pacific Golden Plover	IA	-	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Hooded Plover	P4	-	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Wood Sandpiper	IA	IA	Not recorded. Suitable wetland or estuarine habitat not present.	Low
Common Greenshank	IA	IA	Not recorded. Suitable wetland or estuarine habitat not present.	Low

5.4 SURVEY LIMITATIONS AND CONSTRAINTS

5.4.1 Flora Survey Limitations and Constraints

An assessment of survey-specific issues and limitations (EPA 2016c) is detailed in Table 5.7.

Table 5.7 – Flora and vegetation survey limitations

Constraint	Impact	Comment
Availability of contextual information at a regional and local scale	Nil	Broad scale vegetation, soil, and geology mapping data were available for the study area, in addition to flora database records, and conservation significant vegetation community records. This information is considered to be adequate to provide appropriate contextual information for the current survey.
Competency/experience of the team carrying out the survey, including experience in the bioregion surveyed	Nil	The Senior Botanist undertaking the field work and specimen identification for the survey has conducted numerous botanical surveys in Western Australia, including the Swan Coastal Plain and Geraldton Sandplains bioregions.
Proportion of flora recorded and/or collected, any identification issues	Nil	Species were only recorded within quadrats, and dominant species were recorded at check sites. Only some taxa were recorded opportunistically. Providing a comprehensive species inventory for the study area was not part of the scope of this reconnaissance survey.
Was the appropriate area fully surveyed (effort and extent)	Nil	4 quadrats and 48 check sites were surveyed across the study area. This level of survey effort was considered to be sufficient for a reconnaissance survey of the study area.
Access restrictions within the survey area	Nil	All parts of the study area were accessible by walking from existing vehicle tracks.
Survey timing, rainfall, season of survey	Nil	Seasonal conditions were considered to be good for a flora and vegetation assessment.
Disturbance that may have affected the results of survey such as fire, flood or clearing	Nil	There were no natural or human interventions that constrained the survey of the study area.

6 DISCUSSION

6.1 FLORA

A total of 117 sub-generic vascular plant taxa from 35 families were recorded from the study area during the current survey. This list is not considered to be comprehensive, and includes only taxa recorded within quadrats, dominant species at check sites, and some taxa recorded opportunistically. The most diverse families recorded were the Proteaceae (22 taxa), Myrtaceae (15 taxa), Fabaceae (10 taxa), and Poaceae (9 taxa).

TPFL and WAHERB database searches identified 61 conservation significant plant species that have been recorded within or have potential to occur within the study area. One Priority listed species (*Hypocalymma tetrapterum* P3) was recorded within the study area during the current survey, where it occurred in degraded *Eucalyptus wandoo* woodland and on an adjacent lateritic hill within a species-rich low shrubland. These diverse shrubland communities (S1 and S2) are likely to support additional conservation significant plant species that were not observed during the current survey due to time constraints. However, the proposed development is unlikely to have any impact on any of these communities.

6.2 VEGETATION

6.2.1 Vegetation Units

The Yandin Wind Farm study area occupies an area of approximately 15,360 ha in the Swan Coastal Plain (Swan Coastal Plain and Dandaragan Plateau subregions) and Geraldton Sandplains (Lesueur Sandplain subregion) bioregions of Western Australia, within which five vegetation associations of Beard (1975) (4, 999, 1030, 1031, 1035) have been previously mapped. The vegetation units described here are broadly consistent with those of Beard (1975).

In a previous reconnaissance survey of the Yandin Wind Farm project area (not corresponding exactly to the current study area) (Outback Ecology 2009), nine woodland communities (primarily *C. calophylla*, *E. todtiana*, and other eucalypt woodlands), two shrubland communities (*Xanthorrhoea preissii* shrublands and mixed low heaths), and one sedgeland community (Cyperaceae spp. sedgelands), were described for several remnant vegetation patches. These communities broadly correspond to the sixteen vegetation units described as part of this survey.

Vegetation over 91.2% of the study area comprised scattered *Corymbia calophylla* or *Eucalyptus todtiana* trees over pasture weeds with no native understorey (W3) in 'completely degraded' condition. Low-lying areas occupied by *Juncus acutus* rushland (R1), were also completely degraded, accounting for 0.6% of the study area. Plantations account for a further 3% of the study area.

Two species rich shrubland units in good to excellent condition were occur in remnant vegetation patches, occurring primarily on lateritic rises and hills: *Banksia hewardiana* mid shrubland over mixed low shrubland (S1) accounts for 0.4% of the study area, and *Xanthorrhoea preissii* mid sparse shrubland over mixed low shrubland (S2) accounts for 0.6% of the study area.

Five eucalypt woodland units (*Eucalyptus rudis* woodland in creek lines [W11, W12], *Corymbia calophylla* and *Banksia attenuata* open woodland over pasture weeds [W6], *Eucalyptus wandoo* woodland [W13], and *Eucalyptus* sp. open mallee woodland [W10]), were in degraded condition due to historical clearing, heavy grazing, and/or the presence of abundant weed species. These account for 1% of the study area.

The remaining seven woodland units occurring in remnant vegetation patches broadly comprise *Corymbia calophylla* woodland, *Eucalyptus todtiana* woodland, and *Banksia attenuata* woodland, and are in good to excellent condition. These account for 3% of the study area.

A section of road reserve along Mimegarra Road to the west of the Brand Highway comprised a rehabilitated area dominated primarily by planted *Acacia pulchella*, *A. saligna*, and eucalypt species.

6.2.2 'Banksia Woodlands of the Swan Coastal Plain' TEC

Five patches of the 'Banksia Woodlands of the Swan Coastal Plain' Threatened Ecological Community were identified within the study area, in addition to two previously mapped patches (DBCAs database, Sept. 2017). Three patches corresponding to vegetation unit W1 occur in remnant vegetation patches and are in 'excellent' condition (Keighery 1994), with few non-aggressive weed species present.

Two patches are linear (ECO4 [W1], ECO5 [W2]), occurring along the verge of Yandin road and an adjacent fence line, and are in 'good' condition due to partial clearing, heavy grazing, and weed invasion, although basic vegetation structure is intact. Existing vehicle tracks, roads, and fence lines intersect both of these patches.

Based on vegetation structure and composition, patch size, and condition, each of these patches are considered to qualify as the TEC according to 'Approved Conservation Advice (incorporating listing advice) for the Banksia Woodlands of the Swan Coastal Plain Ecological Community' (Threatened Species Scientific Committee 2016a).

6.3 AVIFAUNA

Results of the current study verify those of a more comprehensive survey, impact assessment and risk assessment of potential bird strike undertaken by RPS (2010). That is, that there was a relatively low diversity of bird species recorded from open pasture areas characterising the proposed wind turbine positions, with the greatest diversity limited to stands of intact native vegetation comprised mainly of shrubland and heath. The presence of isolated trees within cleared areas often attracted birds, albeit in small numbers and at low diversities.

Although there are a number of regional wetlands in the wider locality, most wetland habitats at the study area were limited to small farm dams or degraded creeklines.

RPS (2010) recorded species flying at RSA elevations 'on more than a rare occasion' as; Australian Kestrel, Wedge-tailed Eagle, Brown Falcon, White-backed Swallow, Black-shouldered Kite and Fairy Martin. Of these, the Australian Kestrel and Black-shouldered Kite were also recorded within RSA elevations during this study.

The threatened Carnaby's Black-Cockatoo was found by RPS (2010) to primarily frequent low-land areas and movements tended to follow valleys with woodland vegetation. Proteaceous heath and shrubland, as well as plantations of pine, provide foraging habitat for this species and flight movements are likely to follow movements to and from these food resources as well as roosting and/or breeding trees.

RPS (2010) provide a comprehensive survey, impact assessment, and risk assessment of potential bird strike, with data obtained from this study corroborating the results presented there.

7 REFERENCES

- Beard, J. S. 1975. The vegetation of the Pilbara region. Explanatory notes to map sheet 5 of vegetation survey of Western Australia: Pilbara. University of Western Australia Press, Perth.
- Beard, J. S. 1976. Vegetation survey of Western Australia - Murchison 1:1 000 000 vegetation series. University of Western Australia Press, Perth.
- CHAH. 2017. Australia's Virtual Herbarium. Available at <http://avh.chah.org.au>.
- DEC. 2010. Definitions, categories and criteria for Threatened and Priority Ecological Communities. Department of Environment and Conservation. Government of Western Australia.
- Desmond, A. 2001. Swan Coastal Plain 1 (SWA1 - Dandaragan Plateau subregion). Department of Conservation and Land Management, Perth, Western Australia.
- Desmond, A. J. and Chant, A. 2003. Geraldton Sandplains 3 (GS3 - Lesueur Sandplain subregion). pp. 293-313 in May, J. E., and McKenzie, N. L., eds. A Biodiversity Audit of Western Australia's 53 Biogeographical Subregions in 2002. Department of Environment and Conservation, Perth.
- DPaW. 2017. Priority Ecological Communities for Western Australia version 27 (30 June 2017). Species and Community Branch, Department of Parks and Wildlife, Western Australia.
- DSEWPaC. 2012. Interim Biogeographic Regionalisation for Australia (IBRA), Version 7. Australian Government Department of Sustainability, Environment, Water, Population and Communities.
- EPA. 2016a. Environmental Factor Guideline: Flora and Vegetation, Environmental Protection Authority, Western Australia.
- EPA. 2016b. Environmental Factor Guideline: Terrestrial Fauna, Environmental Protection Authority, Western Australia.
- EPA. 2016c. Technical Guidance: Flora and Vegetation Surveys for Environmental Impact Assessment, Environmental Protection Authority, Western Australia. December 2016.
- EPA. 2016d. Technical Guidance: Terrestrial Fauna Surveys. December 2016. Environmental Protection Authority, Western Australia. .
- ESCAVI. 2003. Australian Vegetation Attribute Manual: National Vegetation Information System (NVIS), Version 6.0. Department of the Environment and Heritage, Canberra.
- IUCN. 2001. IUCN Red List Categories and Criteria: Version 3.1. IUCN, Gland, Switzerland and Cambridge, UK.
- Mitchell, D., Williams, K., and Desmond, A. 2002. Swan Coastal Plain 2 (SWA2 - Swan Coastal Plain subregion). Department of Conservation and Land Management, Perth, Western Australia.
- Northcote, K. H., Beckmann, G. G., Bettenay, E., Churchward, H. M., Van Dijk, D. C., Dimmock, G. M., Hubble, G. D., Isbell, R. F., McArthur, W. M., Murtha, G. G., Nicolls, K. D., Paton, T. R., Thompson, C. H., Webb, A. A., and Wright, M. J. 1960-1968. Atlas of Australian Soils, Sheets 1 to 10. With explanatory data. CSIRO Australia and Melbourne University Press, Melbourne.
- Outback Ecology. 2009. Yandin Wind Farm Targeted Level 1 Vegetation and Flora Assessment - November 2008 and January 2009. Unpublished report for RPS.
- RPS. 2010. Avifauna Assessment. Proposed Yandin Wind Farm Development, Dandaragan Shire. RPS Environment and Planning Pty Ltd. November 2010.
- Shepherd, D. P., Beeston, G. R., and Hopkins, A. J. M. 2001. Native vegetation in Western Australia: Extent, type and status. Technical Report 249. Department of Agriculture, South Perth, Western Australia.

- Threatened Species Scientific Committee. 2016a. Approved Conservation Advice (incorporating listing advice) for the Banksia Woodlands of the Swan Coastal Plain Ecological Community. Canberra: Department of the Environment and Energy. Available from: <http://www.environment.gov.au/biodiversity/threatened/communities/pubs/131-conservation-advice.pdf>. In effect under the EPBC Act from 16-Sep-2016.
- Threatened Species Scientific Committee. 2016b. Species Profile and Threats Database (retrieved from <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>). Threatened Species Scientific Committee.
- USE, D. D. N. 2012. Interim Biogeographic Regionalisation for Australia (IBRA) Version 7. Australian Government Department of Sustainability, Environment, Water, Population and Communities, Canberra.
- Western Australian Herbarium. 1998-2016. FloraBase - The Western Australian Flora. Government of Western Australia Department of Parks and Wildlife. Available at: <http://florabase.dpaw.wa.gov.au>.

8 APPENDICES

APPENDIX A DEFINITIONS

Threatened (WC Act) and Priority flora Categories

Code	Definition
T	Threatened flora – (Declared Rare Flora – Extant) Taxa which have been adequately searched for and are deemed to be in the wild either rare, in danger of extinction, or otherwise in need of special protection and have been gazetted as such (Schedule 1 under the <i>Wildlife Conservation Act 1950</i>).
X	Presumed Extinct Flora (Declared Rare Flora - Extinct) Taxa which have been adequately searched for and there is no reasonable doubt that the last individual has died, and have been gazetted as such Schedule 2 under the <i>Wildlife Conservation Act 1950</i> .
P1	Priority One – Poorly Known Species Species that are known from one or a few collections or sight records (generally less than five), all on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, Shire, Westrail and Main Roads WA road, gravel and soil reserves, and active mineral leases and under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes.
P2	Priority Two – Poorly Known Species Species that are known from one or a few collections or sight records, some of which are on lands not under imminent threat of habitat destruction or degradation, e.g. national parks, conservation parks, nature reserves, State forest, vacant Crown land, water reserves, etc. Species may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes.
P3	Priority Three – Poorly Known Species Species that are known from collections or sight records from several localities not under imminent threat, or from few but widespread localities with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and known threatening processes exist that could affect them.
P4	Priority Four – Rare, Near Threatened and other species in need of monitoring (a) Rare. Species that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These species are usually represented on conservation lands. (b) Near Threatened. Species that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable. (c) Species that have been removed from the list of threatened species during the past five years for reasons other than taxonomy.
P5	Priority Five - Conservation Dependent species Species that are not threatened but are subject to a specific conservation program, the cessation of which would result in the species becoming threatened within five years.

Threatened flora (EPBC Act) Categories

Code	Definition
Ex	Extinct Taxa which at a particular time if, at that time, there is no reasonable doubt that the last member of the species has died.
ExW	Extinct in the Wild Taxa which is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; or it has not been recorded in its known and/or expected habitat, at appropriate seasons, anywhere in its past range, despite exhaustive surveys over a time frame appropriate to its life cycle and form.
CE	Critically Endangered Taxa which at a particular time if, at that time, it is facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria.
E	Endangered Taxa which is not critically endangered and it is facing a very high risk of extinction in the wild in the immediate or near future, as determined in accordance with the prescribed criteria.
V	Vulnerable Taxa which is not critically endangered or endangered and is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria.
CD	Conservation Dependent Taxa which at a particular time if, at that time, the species is the focus of a specific conservation programme, the cessation of which would result in the species becoming vulnerable, endangered or critically endangered within a period of 5 years.

Definition of codes for Threatened Ecological Communities

Code	Definition
PD: Presumed Totally Destroyed	An ecological community that has been adequately searched for but for which no representative occurrences have been located. The community has been found to be totally destroyed or so extensively modified throughout its range that no occurrence of it is likely to recover its species composition and/or structure in the foreseeable future. An ecological community will be listed as presumed totally destroyed if there are no recent records of the community being extant
CR: Critically Endangered	An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or that was originally of limited distribution and is facing severe modification or destruction throughout its range in the immediate future, or is already severely degraded throughout its range but capable of being substantially restored or rehabilitated. An ecological community will be listed as Critically Endangered when it has been adequately surveyed and is found to be facing an extremely high risk of total destruction in the immediate future.
EN: Endangered	An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or was originally of limited distribution and is in danger of significant modification throughout its range or severe modification or destruction over most of its range in the near future. An ecological community will be listed as Endangered when it has been adequately surveyed and is not Critically Endangered but is facing a very high risk of total destruction in the near future.
VU: Vulnerable	An ecological community that has been adequately surveyed and is found to be declining and/or has declined in distribution and/or condition and whose ultimate security has not yet been assured and/or a community that is still widespread but is believed likely to move into a category of higher threat in the near future if threatening processes continue or begin operating throughout its range. An ecological community will be listed as Vulnerable when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing a high risk of total destruction or significant modification in the medium to long-term future.

Definition of codes for Priority Ecological Communities

Code	Definition
P1: Priority One	Ecological communities with apparently few, small occurrences, all or most not actively managed for conservation (e.g. within agricultural or Pastoral lands, urban areas, active mineral leases) and for which current threats exist. Communities may be included if they are comparatively well-known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under immediate threat from known threatening processes across their range.
P2: Priority Two	Communities that are known from few small occurrences, all or most of which are actively managed for conservation (e.g. within national parks, conservation parks, nature reserves, State forest, unallocated Crown land, water reserves, etc.) and not under imminent threat of destruction or degradation. Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under threat from known threatening processes.
P3: Priority Three	<p>(i) Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or:</p> <p>(ii) Communities known from a few widespread occurrences, which are either large or within significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat, or;</p> <p>(iii) Communities made up of large, and/or widespread occurrences that may or not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing by domestic and/or feral stock, and inappropriate fire regimes.</p> <p>Communities may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and/or are not well defined, and known threatening processes exist that could affect them.</p>
P4: Priority Four	<p>Ecological communities that are adequately known, Rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list. These communities require regular monitoring.</p> <p>(a) Rare. Ecological communities known from few occurrences that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These communities are usually represented on conservation lands.</p> <p>(b) Near Threatened. Ecological communities that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable.</p> <p>(c) Ecological communities that have been removed from the list of threatened communities during the past five years.</p>
P5: Priority Five	Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.

Threatened (WC Act) Fauna Categories

Category	Code	Definition	Schedule
Critically Endangered	CR	Threatened species considered to be facing an extremely high risk of extinction in the wild. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 1 of the Wildlife Conservation (Specially Protected Fauna) Notice for Threatened Fauna and Wildlife Conservation (Rare Flora) Notice for Threatened Flora.	Schedule 1 Fauna that is rare or is likely to become extinct as critically endangered fauna
Endangered	EN	Threatened species considered to be facing a very high risk of extinction in the wild. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 2 of the Wildlife Conservation (Specially Protected Fauna) Notice for Threatened Fauna and Wildlife Conservation (Rare Flora) Notice for Threatened Flora.	Schedule 2 Fauna that is rare or is likely to become extinct as endangered fauna
Vulnerable	VU	Threatened species considered to be facing a high risk of extinction in the wild. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 3 of the Wildlife Conservation (Specially Protected Fauna) Notice for Threatened Fauna and Wildlife Conservation (Rare Flora) Notice for Threatened Flora.	Schedule 3 Fauna that is rare or is likely to become extinct as vulnerable fauna
Presumed Extinct	EX	Species which have been adequately searched for and there is no reasonable doubt that the last individual has died. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 4 of the Wildlife Conservation (Specially Protected Fauna) Notice for Presumed Extinct Fauna and Wildlife Conservation (Rare Flora) Notice for Presumed Extinct Flora.	Schedule 4 Fauna presumed to be extinct
Migratory	IA	Birds that are subject to an agreement between the government of Australia and the governments of Japan (JAMBA), China (CAMBA) and The Republic of Korea (ROKAMBA), and the Bonn Convention, relating to the protection of migratory birds. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 5 of the Wildlife Conservation (Specially Protected Fauna) Notice.	Schedule 5 Migratory birds protected under an international agreement
Conservation Dependent	CD	Fauna of special conservation need being species dependent on ongoing conservation intervention to prevent it becoming eligible for listing as threatened. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 6 of the Wildlife Conservation (Specially Protected Fauna) Notice.	Schedule 6 Fauna that is of special conservation need as conservation dependent fauna
Special Protection	OS	Fauna otherwise in need of special protection to ensure their conservation. Published as Specially Protected under the <i>Wildlife Conservation Act 1950</i> , in Schedule 7 of the Wildlife Conservation (Specially Protected Fauna) Notice.	Schedule 7 Other specially protected fauna

Fauna (EPBC Act) Categories

Category	Code	Definition
Extinct	Ex	Fauna not definitely located in the wild during the past 50 years
Extinct in the Wild	EW	Fauna which is known only to survive in captivity
Critically Endangered	CR	Fauna that is considered to be facing an extremely high risk of extinction in the wild in the immediate future
Endangered	EN	Fauna that is considered to be facing a very high risk of extinction in the wild in the near future
Vulnerable	VU	Fauna that is considered to be facing a high risk of extinction in the wild in the medium-term future
Conservation Dependent	CD	Fauna whose survival depends upon ongoing conservation measures. Without these measures, a conservation dependent taxon would be classified as Vulnerable or more severely threatened.
Migratory	IA	Fauna that migrates to, over and within Australia and its external territories.

Definition of codes for Priority Fauna

Code	Definition
P1: Priority One	<p>Poorly-known species Species that are known from one or a few locations (generally five or less) which are potentially at risk. All occurrences are either: very small; or on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, road and rail reserves, gravel reserves and active mineral leases; or otherwise under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes. Such species are in urgent need of further survey.</p>
P2: Priority Two	<p>Poorly-known species Species that are known from one or a few locations (generally five or less), some of which are on lands managed primarily for nature conservation, e.g. national parks, conservation parks, nature reserves and other lands with secure tenure being managed for conservation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes. Such species are in urgent need of further survey.</p>
P3: Priority Three	<p>Poorly-known species Species that are known from several locations, and the species does not appear to be under imminent threat, or from few but widespread locations with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several locations but do not meet adequacy of survey requirements and known threatening processes exist that could affect them. Such species are in need of further survey.</p>
P4: Priority Four	<p>Rare, Near Threatened and other species in need of monitoring</p> <p>(a) Rare. Species that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These species are usually represented on conservation lands.</p> <p>(b) Near Threatened. Species that are considered to have been adequately surveyed and that are close to qualifying for Vulnerable, but are not listed as Conservation Dependent.</p> <p>(c) Species that have been removed from the list of threatened species during the past five years for reasons other than taxonomy.</p>

APPENDIX B VASCULAR FLORA RECORDS - NATUREMAP

NatureMap Species Report

Created By Guest user on 13/09/2017

Kingdom Plantae
Current Names Only Yes
Core Datasets Only Yes
Method 'By Circle'
Centre 115° 36' 41" E, 30° 44' 52" S
Buffer 10km

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
1.	3200 <i>Acacia acuminata</i> (Jam, Mangard)			
2.	15466 <i>Acacia appplanata</i>			
3.	3231 <i>Acacia auronitens</i>			
4.	3242 <i>Acacia blakelyi</i>			
5.	14061 <i>Acacia clydonophora</i>			
6.	3271 <i>Acacia costata</i>			
7.	20435 <i>Acacia daphnifolia</i>			
8.	11926 <i>Acacia drewiana</i> subsp. <i>drewiana</i>			
9.	3341 <i>Acacia forrestiana</i> (Forrest's Wattle)		T	
10.	11611 <i>Acacia lasiocarpa</i> var. <i>lasiocarpa</i>			
11.	3442 <i>Acacia microbotrya</i> (Manna Wattle, Kalyang)			
12.	3493 <i>Acacia plicata</i>		P3	
13.	15481 <i>Acacia pulchella</i> var. <i>glaberrima</i>			
14.	15483 <i>Acacia pulchella</i> var. <i>pulchella</i>			
15.	30033 <i>Acacia saligna</i> subsp. <i>lindleyi</i>			
16.	3532 <i>Acacia scirpifolia</i>			
17.	3541 <i>Acacia sessilis</i>			
18.	15486 <i>Acacia sphacelata</i> subsp. <i>verticillata</i>			
19.	3557 <i>Acacia stenoptera</i> (Narrow Winged Wattle)			
20.	6205 <i>Actinotus leucocephalus</i> (Flannel Flower)			
21.	1779 <i>Adenanthos drummondii</i>			
22.	185 <i>Aira cupaniana</i> (Silvery Hairgrass)	Y		
23.	1056 <i>Alexgeorgea nitens</i>			
24.	1732 <i>Allocasuarina humilis</i> (Dwarf Sheoak)			
25.	1734 <i>Allocasuarina microstachya</i>			
26.	1739 <i>Allocasuarina thuyoides</i> (Horned Sheoak)			
27.	12025 <i>Amphipogon caricinus</i> var. <i>caricinus</i>			
28.	199 <i>Amphipogon strictus</i> (Greybeard Grass)			
29.	1058 <i>Anarthria gracilis</i>			
30.	1059 <i>Anarthria humilis</i>			
31.	6311 <i>Andersonia heterophylla</i>			
32.	40908 <i>Androcalva pulchella</i>			
33.	1409 <i>Anigozanthos humilis</i> (Catspaw)			
34.	29437 <i>Anigozanthos humilis</i> subsp. <i>Badgingarra</i> (S.D. Hopper 7114)		P2	
35.	11957 <i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i> (Golden Catspaw)		P4	
36.	11434 <i>Anigozanthos humilis</i> subsp. <i>humilis</i>			
37.	1414 <i>Anigozanthos pulcherrimus</i> (Yellow Kangaroo Paw)			
38.	<i>Aristida</i> sp.			
39.	1265 <i>Arthropodium curvipes</i>			
40.	4397 <i>Asterolasia drummondii</i> (Gairdner Range Starbush)		P4	
41.	6328 <i>Astroloma glaucescens</i>			
42.	6332 <i>Astroloma microdonta</i> (Sandplain Cranberry)			
43.	6334 <i>Astroloma pallidum</i> (Kick Bush)			
44.	14501 <i>Astroloma</i> sp. <i>Eneabba</i> (N. Marchant s.n. PERTH 01291777)			
45.	6339 <i>Astroloma xerophyllum</i>			
46.	17237 <i>Austrostipa elegantissima</i>			
47.	17244 <i>Austrostipa macalpinei</i>			
48.	17257 <i>Austrostipa variabilis</i>			
49.	233 <i>Avena barbata</i> (Bearded Oat)	Y		
50.	45397 <i>Babingtonia cherticola</i>		P3	
51.	45395 <i>Babingtonia delicata</i>		P1	
52.	45416 <i>Babingtonia grandiflora</i> (Large-flowered Babingtonia)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
53.	1800 <i>Banksia attenuata</i> (Slender Banksia, Piara)			
54.	32679 <i>Banksia bipinnatifida</i> subsp. <i>multifida</i>			
55.	32576 <i>Banksia dallanneyi</i> (Couch Honeypot)			
56.	32580 <i>Banksia dallanneyi</i> var. <i>dallanneyi</i>			
57.	32577 <i>Banksia dallanneyi</i> var. <i>melliculca</i>			
58.	32521 <i>Banksia fraseri</i>			
59.	1820 <i>Banksia grossa</i>			
60.	32518 <i>Banksia hewardiana</i>			
61.	32216 <i>Banksia kippistiana</i> var. <i>paenepeccata</i>		P3	
62.	32157 <i>Banksia prionophylla</i>		P1	Y
63.	32086 <i>Banksia sclerophylla</i>			
64.	32074 <i>Banksia shuttleworthiana</i> (Bearded Dryandra)			
65.	32033 <i>Banksia tortifolia</i>			
66.	5384 <i>Beaufortia eriocephala</i> (Woolly Bottlebrush, Woolly Beaufortia)		P3	
67.	1417 <i>Blancoa canescens</i> (Winter Bell)			
68.	11274 <i>Boronia coerulescens</i> subsp. <i>spinescens</i>			
69.	11381 <i>Boronia ramosa</i> subsp. <i>anethifolia</i>			
70.	4440 <i>Boronia scabra</i> (Rough Boronia)			
71.	16637 <i>Boronia scabra</i> subsp. <i>condensata</i>		P2	
72.	16639 <i>Boronia scabra</i> subsp. <i>scabra</i>			
73.	1273 <i>Borya sphaerocephala</i> (Pincushions)			
74.	3710 <i>Bossiaea eriocarpa</i> (Common Brown Pea)			
75.	1383 <i>Burchardia bairdiae</i>			
76.	12770 <i>Burchardia congesta</i>			
77.	29439 <i>Caesia</i> sp. <i>Wongan</i> (K.F. Kenneally 8820)			
78.	1586 <i>Caladenia discoidea</i> (Dancing Orchid)			
79.	15348 <i>Caladenia flava</i> subsp. <i>flava</i>			
80.	15360 <i>Caladenia longicauda</i> subsp. <i>borealis</i>			
81.	15369 <i>Caladenia lorea</i>			
82.	2848 <i>Calandrinia corrigioloides</i> (Strap Purslane)			
83.	19309 <i>Calectasia narragara</i>			
84.	35816 <i>Calothamnus quadrifidus</i> subsp. <i>quadrifidus</i>			
85.	5429 <i>Calothamnus sanguineus</i> (Silky-leaved Blood flower, Pindak)			
86.	5431 <i>Calothamnus torulosus</i>			
87.	5439 <i>Calytrix angulata</i> (Yellow Starflower)			
88.	5441 <i>Calytrix aurea</i>			
89.	5465 <i>Calytrix leschenaultii</i>			
90.	760 <i>Caustis dioica</i>			
91.	41564 <i>Cenchrus clandestinus</i> (Kikuyu Grass)	Y		
92.	1121 <i>Centrolepis aristata</i> (Pointed Centrolepis)			
93.	1125 <i>Centrolepis drummondiana</i>			
94.	1134 <i>Centrolepis polygyna</i> (Wiry Centrolepis)			
95.	17685 <i>Chaetanthus aristatus</i>			
96.	11299 <i>Chamaescilla corymbosa</i> var. <i>corymbosa</i>			
97.	35619 <i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)		T	Y
98.	5498 <i>Chamelaucium uncinatum</i> (Geraldton Wax)			
99.	17706 <i>Chordifex sinuosus</i>			
100.	4566 <i>Comesperma volubile</i> (Love Creeper)			
101.	40922 <i>Commersonia densiflora</i>			
102.	1882 <i>Conospermum stoechadis</i> (Common Smokebush)			
103.	15611 <i>Conospermum stoechadis</i> subsp. <i>stoechadis</i> (Common Smokebush)			
104.	19026 <i>Conostephium magnum</i>		P4	
105.	6347 <i>Conostephium minus</i> (Pink-tipped Pearl flower)			
106.	6348 <i>Conostephium pendulum</i> (Pearl Flower)			
107.	1418 <i>Conostylis aculeata</i> (Prickly Conostylis)			
108.	11414 <i>Conostylis aculeata</i> subsp. <i>breviflora</i>			
109.	1420 <i>Conostylis androstemma</i> (Trumpets)			
110.	1421 <i>Conostylis angustifolia</i>			
111.	1423 <i>Conostylis aurea</i> (Golden Conostylis)			
112.	11438 <i>Conostylis candicans</i> subsp. <i>candicans</i>			
113.	11773 <i>Conostylis crassinervia</i> subsp. <i>absens</i>			
114.	11695 <i>Conostylis festucacea</i> subsp. <i>festucacea</i>			
115.	1436 <i>Conostylis juncea</i>			
116.	11870 <i>Conostylis teretifolia</i> subsp. <i>teretifolia</i>			
117.	1458 <i>Conostylis teretiuscula</i>			
118.	5502 <i>Conothamnus trinervis</i>			
119.	13354 <i>Craspedia variabilis</i>			
120.	31571 <i>Cryptandra intermedia</i>			
121.	9076 <i>Cryptandra myriantha</i>			
122.	4809 <i>Cryptandra pungens</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
123.	768 <i>Cyathochaeta avenacea</i>			
124.	783 <i>Cyperus congestus</i> (Dense Flat-sedge)	Y		
125.	816 <i>Cyperus tenuiflorus</i> (Scaly Sedge)	Y		
126.	7428 <i>Dampiera coronata</i> (Wedge-leaved Dampiera)			
127.	7451 <i>Dampiera lavandulacea</i>			
128.	7475 <i>Dampiera spicigera</i> (Spiked Dampiera)			
129.	7481 <i>Dampiera tephrea</i>		P2	
130.	7482 <i>Dampiera teres</i> (Terete-leaved Dampiera)			
131.	5518 <i>Darwinia neildiana</i> (Fringed Bell)			
132.	5524 <i>Darwinia pinifolia</i>			
133.	1220 <i>Dasyogon obliquifolius</i>			
134.	3793 <i>Daviesia angulata</i>			
135.	19747 <i>Daviesia decurrens</i> subsp. <i>decurrens</i>			
136.	18560 <i>Daviesia divaricata</i> subsp. <i>divaricata</i>			
137.	11879 <i>Daviesia hakeoides</i> subsp. <i>hakeoides</i>			
138.	15505 <i>Daviesia incrassata</i> subsp. <i>incrassata</i>			
139.	15506 <i>Daviesia incrassata</i> subsp. <i>teres</i>			
140.	12329 <i>Daviesia nudiflora</i> subsp. <i>hirtella</i>			
141.	16585 <i>Daviesia nudiflora</i> subsp. <i>nudiflora</i>			
142.	3833 <i>Daviesia podophylla</i>			
143.	3835 <i>Daviesia preissii</i>			
144.	3845 <i>Daviesia triflora</i>			
145.	17662 <i>Desmocladus lateriticus</i>			
146.	46364 <i>Desmocladus microcarpus</i>		P2	
147.	29078 <i>Dillwynia</i> sp. Northern Sandplains (M. Hislop 3278)			
148.	15275 <i>Diplolaena obovata</i>			
149.	3011 <i>Diplotaxis muralis</i> (Wall Rocket)	Y		
150.	42182 <i>Diuris perialla</i>			
151.	43300 <i>Diuris refracta</i>			
152.	4761 <i>Dodonaea ericoides</i>			
153.	13201 <i>Drosera eneabba</i>			
154.	13212 <i>Drosera erythrorhiza</i> subsp. <i>magna</i>			
155.	3098 <i>Drosera glanduligera</i> (Pimpernel Sundew)			
156.	3101 <i>Drosera heterophylla</i> (Swamp Rainbow)			
157.	13199 <i>Drosera leioblastus</i>			
158.	14298 <i>Drosera macrantha</i> subsp. <i>macrantha</i>			
159.	13208 <i>Drosera marchantii</i> subsp. <i>prophylla</i>		P3	
160.	3118 <i>Drosera pallida</i> (Pale Rainbow)			
161.	3119 <i>Drosera parvula</i> (Small Sundew)			
162.	29178 <i>Drosera porrecta</i>			
163.	13185 <i>Drosera spilos</i>			
164.	1066 <i>Ecdiocolea monostachya</i>			
165.	11105 <i>Echinochloa crus-galli</i>	Y		
166.	347 <i>Ehrharta calycina</i> (Perennial Veldt Grass)	Y		
167.	349 <i>Ehrharta longiflora</i> (Annual Veldt Grass)	Y		
168.	17605 <i>Eleocharis keigheryi</i>		T	
169.	1643 <i>Elythranthera brunonis</i> (Purple Enamel Orchid)			
170.	13950 <i>Eremaea asterocarpa</i> subsp. <i>asterocarpa</i>			
171.	5541 <i>Eremaea pauciflora</i>			
172.	45244 <i>Ericomyrtus serpyllifolia</i>			
173.	45215 <i>Ericomyrtus tenuior</i>			
174.	12898 <i>Eucalyptus abdita</i>		P2	
175.	12895 <i>Eucalyptus arachnaea</i> subsp. <i>arachnaea</i>			
176.	13546 <i>Eucalyptus dolorosa</i>		T	Y
177.	5628 <i>Eucalyptus drummondii</i> (Drummond's Gum)			
178.	5658 <i>Eucalyptus gittinsii</i> (Northern Sandplain Mallee)			
179.	18292 <i>Eucalyptus gittinsii</i> subsp. <i>illucida</i>			
180.	5690 <i>Eucalyptus lane-poolei</i> (Salmon White Gum)			
181.	13531 <i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i> (Small-leaved Mottlecah)		P4	
182.	13530 <i>Eucalyptus macrocarpa</i> subsp. <i>macrocarpa</i> (Mottlecah)			
183.	5717 <i>Eucalyptus myriadena</i>			
184.	12866 <i>Eucalyptus pluricaulis</i> subsp. <i>pluricaulis</i>			
185.	5763 <i>Eucalyptus rudis</i> (Flooded Gum, Kulurda)			
186.	13511 <i>Eucalyptus rudis</i> subsp. <i>rudis</i>			
187.	5790 <i>Eucalyptus todtiana</i> (Coastal Blackbutt)			
188.	12905 <i>Eucalyptus wandoo</i> subsp. <i>pulverea</i>			
189.	20515 <i>Gastrolobium axillare</i>			
190.	20475 <i>Gastrolobium capitatum</i>			
191.	3906 <i>Gastrolobium ilicifolium</i>			
192.	20482 <i>Gastrolobium nervosum</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
193.	20514 <i>Gastrolobium nudum</i>		P2	
194.	3912 <i>Gastrolobium oxylobioides</i> (Champion Bay Poison)			
195.	3915 <i>Gastrolobium plicatum</i>			
196.	3916 <i>Gastrolobium polystachyum</i> (Horned Poison)			
197.	3924 <i>Gastrolobium spinosum</i> (Prickly Poison)			
198.	6143 <i>Glischrocaryon aureum</i> (Common Popflower)			
199.	8002 <i>Gnephosis tenuissima</i>			
200.	10909 <i>Gompholobium confertum</i>			
201.	3950 <i>Gompholobium knightianum</i>			
202.	3955 <i>Gompholobium preissii</i>			
203.	3957 <i>Gompholobium tomentosum</i> (Hairy Yellow Pea)			
204.	6149 <i>Gonocarpus cordiger</i>			
205.	6161 <i>Gonocarpus pithyoides</i>			
206.	7495 <i>Goodenia berardiana</i>			
207.	29362 <i>Goodenia coerulea</i>			
208.	12516 <i>Goodenia convexa</i>			
209.	12520 <i>Goodenia fasciculata</i>			
210.	7538 <i>Goodenia pulchella</i>			
211.	15763 <i>Grevillea biformis</i> subsp. <i>biformis</i>			
212.	12221 <i>Grevillea calliantha</i>		T	
213.	1994 <i>Grevillea drummondii</i> (Drummond's Grevillea)		P4	
214.	2001 <i>Grevillea eriostachya</i> (Flame Grevillea, Kaliny-kalinyapa)			
215.	19567 <i>Grevillea florida</i>		P3	
216.	2087 <i>Grevillea saccata</i> (Pouched Grevillea)		P4	
217.	19569 <i>Grevillea synapheae</i> subsp. <i>A Flora of Australia</i> (S.D. Hopper 6333)		P1	Y
218.	17450 <i>Grevillea synapheae</i> subsp. <i>minyulo</i>		P1	
219.	14420 <i>Grevillea synapheae</i> subsp. <i>pachyphylla</i>			
220.	14423 <i>Grevillea thyrsoidea</i> subsp. <i>thyrsoidea</i>		P3	
221.	2116 <i>Grevillea uncinulata</i> (Hook-leaf Grevillea)			
222.	19231 <i>Grevillea uncinulata</i> subsp. <i>Coomallo</i> (S.J. Patrick 719)			
223.	12824 <i>Grevillea vestita</i> subsp. <i>vestita</i>			
224.	13233 <i>Guichenotia alba</i>		P3	
225.	5014 <i>Guichenotia sarotes</i>			
226.	2783 <i>Gyrostemon racemiger</i>			
227.	2784 <i>Gyrostemon ramulosus</i> (Corkybark)			
228.	1469 <i>Haemodorum loratum</i>		P3	
229.	1473 <i>Haemodorum simulans</i>			
230.	1475 <i>Haemodorum spicatum</i> (Mardja)			
231.	17670 <i>Hakea anadenia</i>			
232.	2131 <i>Hakea auriculata</i>			
233.	2136 <i>Hakea candolleana</i>			
234.	2143 <i>Hakea conchifolia</i> (Shell-leaved Hakea)			
235.	2164 <i>Hakea gilbertii</i>			
236.	2175 <i>Hakea lissocarpha</i> (Honey Bush)			
237.	12230 <i>Hakea longiflora</i>		P3	
238.	45333 <i>Hakea neospathulata</i>			
239.	2197 <i>Hakea prostrata</i> (Harsh Hakea)			
240.	12233 <i>Hakea psilorrhyncha</i>			
241.	2205 <i>Hakea smilacifolia</i>			
242.	2206 <i>Hakea stenocarpa</i> (Narrow-fruited Hakea)			
243.	2214 <i>Hakea trifurcata</i> (Two-leaf Hakea)			
244.	2215 <i>Hakea undulata</i> (Wavy-leaved Hakea)			
245.	6838 <i>Hemiandra linearis</i> (Speckled Snakebush)			
246.	6839 <i>Hemiandra pungens</i> (Snakebush)			
247.	38320 <i>Hemiandra</i> sp. <i>Jurien</i> (B.J. Conn & M.E. Tozer BJC 3885)			
248.	6856 <i>Hemigenia incana</i> (Silky Hemigenia)			
249.	41020 <i>Hemiphora bartlingii</i> (Woolly Dragon)			
250.	5108 <i>Hibbertia acerosa</i> (Needle Leaved Guinea Flower)			
251.	5112 <i>Hibbertia aurea</i>			
252.	5116 <i>Hibbertia crassifolia</i>			
253.	20046 <i>Hibbertia hibbertioides</i> var. <i>hibbertioides</i>			
254.	45534 <i>Hibbertia hypericoides</i> subsp. <i>hypericoides</i>			
255.	5148 <i>Hibbertia mylnei</i>			
256.	5157 <i>Hibbertia polystachya</i>			
257.	5162 <i>Hibbertia racemosa</i> (Stalked Guinea Flower)			
258.	<i>Hibbertia</i> sp.			
259.	46435 <i>Hibbertia</i> sp. <i>Geraldton Sandplains</i> (R. Edmiston E 421)			
260.	48381 <i>Hibbertia striata</i>			
261.	5173 <i>Hibbertia subvaginata</i>			
262.	6222 <i>Homalosciadium homalocarpum</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
263.	3966 <i>Hovea pungens</i> (Devil's Pins, Puyenak)			
264.	3967 <i>Hovea stricta</i>			
265.	12741 <i>Hyalosperma cotula</i>			
266.	5216 <i>Hybanthus calycinus</i> (Wild Violet)			
267.	5221 <i>Hybanthus floribundus</i>			
268.	20044 <i>Hypocalymma hirsutum</i>			
269.	14493 <i>Hypocalymma</i> sp. <i>Cataby</i> (G.J. Keighery 5151)		P2	
270.	5828 <i>Hypocalymma tetrapterum</i>		P3	
271.	5829 <i>Hypocalymma xanthopetalum</i>			
272.	1070 <i>Hypolaena exsulca</i>			
273.	17622 <i>Hypolaena robusta</i>		P4	
274.	20200 <i>Isolepis cernua</i> var. <i>setiformis</i>			
275.	919 <i>Isolepis oldfieldiana</i>			
276.	2221 <i>Isopogon asper</i>			
277.	2227 <i>Isopogon divergens</i> (Spreading Coneflower)			
278.	2232 <i>Isopogon linearis</i>			
279.	7396 <i>Isotoma hypocrateriformis</i> (Woodbridge Poison)			
280.	3992 <i>Isotropis cuneifolia</i> (Granny Bonnets)			
281.	19700 <i>Isotropis cuneifolia</i> subsp. <i>cuneifolia</i>			
282.	4003 <i>Jacksonia carduacea</i>		P3	
283.	4010 <i>Jacksonia floribunda</i> (Holly Pea)			
284.	4019 <i>Jacksonia macrocalyx</i>			
285.	14778 <i>Jacksonia nutans</i>			
286.	4025 <i>Jacksonia restioides</i>			
287.	19632 <i>Johnsonia pubescens</i> subsp. <i>pubescens</i>			
288.	20454 <i>Juncus acutus</i> subsp. <i>acutus</i>	Y		
289.	1188 <i>Juncus pallidus</i> (Pale Rush)			
290.	4044 <i>Kennedia prostrata</i> (Scarlet Runner)			
291.	6780 <i>Lachnostachys eriobotrya</i> (Lambswool)			
292.	468 <i>Lamarckia aurea</i> (Goldentop)	Y		
293.	15528 <i>Lambertia multiflora</i> var. <i>multiflora</i>			
294.	5031 <i>Lasiopetalum drummondii</i>			
295.	15552 <i>Lasiopetalum</i> sp. <i>Hill River</i> (T.N. Stoate 5)		P1	
296.	1305 <i>Laxmannia omnifertilis</i>			
297.	11464 <i>Laxmannia sessiliflora</i> subsp. <i>australis</i>			
298.	1309 <i>Laxmannia squarrosa</i>			
299.	7568 <i>Lechenaultia biloba</i> (Blue Leschenaultia)			
300.	7586 <i>Lechenaultia stenosepala</i> (Narrow-sepaled Leschenaultia)			
301.	1073 <i>Lepidobolus chaetocephalus</i> (Bristle-headed Chaff Rush)			
302.	1075 <i>Lepidobolus preissianus</i>			
303.	13775 <i>Lepidobolus quadratus</i>		P3	
304.	930 <i>Lepidosperma costale</i>			
305.	937 <i>Lepidosperma longitudinale</i> (Pithy Sword-sedge)			
306.	944 <i>Lepidosperma scabrum</i>			
307.	945 <i>Lepidosperma squamatum</i>			
308.	1078 <i>Leptocarpus coangustatus</i>			
309.	5847 <i>Leptospermum erubescens</i> (Roadside Teatree)			
310.	5857 <i>Leptospermum spinescens</i>			
311.	48179 <i>Leucopogon foliosus</i>		P3	
312.	6400 <i>Leucopogon gracillimus</i>			
313.	6420 <i>Leucopogon oldfieldii</i>			
314.	6421 <i>Leucopogon oliganthus</i>			
315.	6434 <i>Leucopogon polymorphus</i>			
316.	39501 <i>Leucopogon</i> sp. <i>Coomallo</i> (R.J. Cranfield 1457)			
317.	20086 <i>Leucopogon</i> sp. <i>Northern Scarp</i> (M. Hislop 2233)			
318.	6444 <i>Leucopogon sprengelioides</i>			
319.	48184 <i>Leucopogon stenophyllus</i>			
320.	7677 <i>Levenhookia stipitata</i> (Common Stylewort)			
321.	9289 <i>Lobelia anceps</i> (Angled Lobelia)			
322.	7406 <i>Lobelia rhombifolia</i> (Tufted Lobelia)			
323.	475 <i>Lolium multiflorum</i> (Italian Ryegrass)	Y		
324.	1239 <i>Lomandra preissii</i>			
325.	1243 <i>Lomandra sericea</i> (Silky Mat Rush)			
326.	8564 <i>Lotus subbiflorus</i>	Y		
327.	1097 <i>Lyginia barbata</i>			
328.	18049 <i>Lyginia imberbis</i>			
329.	34736 <i>Lysinema pentapetalum</i>			
330.	5281 <i>Lythrum hyssopifolia</i> (Lesser Loosestrife)	Y		
331.	1477 <i>Macropidia fuliginosa</i> (Black Kangaroo Paw)			
332.	18119 <i>Macrozamia fraseri</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
333.	17633 <i>Marianthus erubescens</i>			
334.	19380 <i>Melaleuca calyptroides</i>			
335.	17982 <i>Melaleuca carrii</i>			
336.	19387 <i>Melaleuca clavifolia</i>			
337.	5893 <i>Melaleuca concreta</i>			
338.	5958 <i>Melaleuca radula</i> (Graceful Honeymyrtle)			
339.	18278 <i>Melaleuca tinkeri</i>			
340.	5983 <i>Melaleuca trichophylla</i>			
341.	954 <i>Mesomelaena preissii</i>			
342.	955 <i>Mesomelaena pseudostygia</i>			
343.	957 <i>Mesomelaena tetragona</i> (Semaphore Sedge)			
344.	15456 <i>Microcorys</i> sp. Coomallo (L. Haegi 2677)			
345.	4100 <i>Mirbelia spinosa</i>			
346.	29418 <i>Monoculus monstrosus</i>	Y		
347.	37440 <i>Monopsis debilis</i> var. <i>depressa</i>	Y		
348.	2401 <i>Nuytsia floribunda</i> (Christmas Tree, Mudja)			
349.	16347 <i>Oenothera laciniata</i>	Y		
350.	32716 <i>Olearia lehmanniana</i>			
351.	18256 <i>Opercularia spermacocea</i>			
352.	18255 <i>Opercularia vaginata</i> (Dog Weed)			
353.	46255 <i>Orianthera campanulata</i>			
354.	46254 <i>Orianthera spermacocea</i>			
355.	1537 <i>Orthrosanthus laxus</i> (Morning Iris)			
356.	4349 <i>Oxalis corniculata</i> (Yellow Wood Sorrel)	Y		
357.	1546 <i>Patersonia juncea</i> (Rush Leaved Patersonia)			
358.	1550 <i>Patersonia occidentalis</i> (Purple Flag, Koma)			
359.	40423 <i>Pentameris airoides</i> (False Hairgrass)	Y		
360.	13911 <i>Persicaria decipiens</i>			
361.	2258 <i>Persoonia comata</i>			
362.	20368 <i>Petrophile axillaris</i>			
363.	2286 <i>Petrophile brevifolia</i>			
364.	16874 <i>Petrophile recurva</i>			
365.	2306 <i>Petrophile rigida</i>			
366.	2309 <i>Petrophile serruriae</i>			
367.	2310 <i>Petrophile shuttleworthiana</i>			
368.	2312 <i>Petrophile striata</i>			
369.	18529 <i>Philothea spicata</i> (Pepper and Salt)			
370.	1173 <i>Philydrella pygmaea</i> (Butterfly Flowers)			
371.	1478 <i>Phlebocarya ciliata</i>			
372.	4 <i>Phylloglossum drummondii</i> (Pigmy Clubmoss)			
373.	6985 <i>Physalis pubescens</i>	Y		
374.	5231 <i>Pimelea angustifolia</i> (Narrow-leaved Pimelea)			
375.	5244 <i>Pimelea floribunda</i>			
376.	11402 <i>Pimelea imbricata</i> var. <i>piligera</i>			
377.	5254 <i>Pimelea leucantha</i>			
378.	18353 <i>Pithocarpa pulchella</i> var. <i>pulchella</i>			
379.	6262 <i>Platysace xerophila</i>			
380.	45237 <i>Podolepis aristata</i> subsp. <i>aristata</i>			
381.	8177 <i>Podolepis lessonii</i>			
382.	8182 <i>Podotheca angustifolia</i> (Sticky Longheads)			
383.	8184 <i>Podotheca gnaphalioides</i> (Golden Long-heads)			
384.	12733 <i>Podotheca pritzelii</i>		P3	
385.	29919 <i>Polianthion wichurae</i>			
386.	582 <i>Polypogon monspeliensis</i> (Annual Beardgrass)	Y		
387.	1680 <i>Prasophyllum parvifolium</i> (Autumn Leek Orchid)			
388.	10853 <i>Prasophyllum plumiforme</i>			
389.	13255 <i>Pterochaeta paniculata</i>			
390.	1687 <i>Pterostylis dilatata</i>			
391.	45343 <i>Pterostylis platypetala</i>			
392.	1698 <i>Pterostylis vittata</i> (Banded Greenhood)			
393.	2742 <i>Ptilotus manglesii</i> (Pom Poms, Mulamula)			
394.	2751 <i>Ptilotus polystachyus</i> (Prince of Wales Feather)			
395.	4172 <i>Pultenaea ericifolia</i>			
396.	3061 <i>Raphanus raphanistrum</i> (Wild Radish)	Y		
397.	13300 <i>Rhodanthe citrina</i>			
398.	13234 <i>Rhodanthe manglesii</i>			
399.	2430 <i>Rumex brownii</i> (Swamp Dock)	Y		
400.	2433 <i>Rumex crispus</i> (Curled Dock)	Y		
401.	7595 <i>Scaevola anchusifolia</i>			
402.	7603 <i>Scaevola canescens</i> (Grey Scaevola)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
403.	7613 <i>Scaevola glandulifera</i> (Viscid Hand-flower)			
404.	12585 <i>Scaevola repens</i>			
405.	12588 <i>Scaevola virgata</i>			
406.	979 <i>Schoenus caespititius</i>			
407.	982 <i>Schoenus clandestinus</i>			
408.	986 <i>Schoenus efoliatus</i>			
409.	17617 <i>Schoenus insolitus</i>			
410.	1000 <i>Schoenus minutulus</i>			
411.	1011 <i>Schoenus rigens</i>			
412.	1018 <i>Schoenus subfascicularis</i>			
413.	16252 <i>Schoenus subflavus</i> subsp. <i>subflavus</i>			
414.	1026 <i>Schoenus unispiculatus</i>			
415.	6033 <i>Scholtzia involucrata</i> (Spiked Scholtzia)			
416.	6037 <i>Scholtzia parviflora</i>			
417.	25884 <i>Senecio pinnatifolius</i> var. <i>latilobus</i>			
418.	7013 <i>Solanum hoplopetalum</i> (Thorny Solanum)			
419.	7018 <i>Solanum lasiophyllum</i> (Flannel Bush, Mindjulu)			
420.	7022 <i>Solanum nigrum</i> (Black Berry Nightshade)	Y		
421.	8230 <i>Sonchus asper</i> (Rough Sowthistle)	Y		
422.	17551 <i>Sphaerolobium drummondii</i>			
423.	10800 <i>Sphaerolobium pulchellum</i>			
424.	4713 <i>Stachystemon axillaris</i> (Leafy Stachystemon)			
425.	4733 <i>Stackhousia monogyna</i>			
426.	9070 <i>Stackhousia pubescens</i> (Downy Stackhousia)			
427.	15065 <i>Stenanthemum notiale</i> subsp. <i>notiale</i>			
428.	14240 <i>Stenanthemum reissekii</i>			
429.	7680 <i>Stylidium aeonioides</i>		P4	
430.	7709 <i>Stylidium crossocephalum</i> (Posy Triggerplant)			
431.	7710 <i>Stylidium cygnorum</i>			
432.	7716 <i>Stylidium diuroides</i> (Donkey Triggerplant)			
433.	18420 <i>Stylidium flagellum</i>			
434.	7762 <i>Stylidium miniatum</i> (Pink Butterfly Triggerplant)			
435.	7768 <i>Stylidium obtusatum</i> (Pinafore Triggerplant)			
436.	7771 <i>Stylidium periscelanthum</i> (Pantaloon Triggerplant)		P3	
437.	25837 <i>Stylidium purpureum</i> (Purple Fountain Triggerplant)			
438.	20521 <i>Stylidium rigidulum</i>			
439.	7798 <i>Stylidium schoenoides</i> (Cow Kicks)			
440.	<i>Stylidium</i> sp.			
441.	17510 <i>Stylidium</i> sp. <i>Kalbarri</i> (A. Carr 145)			
442.	25836 <i>Stylidium spiciforme</i> (Spiciform Triggerplant)			
443.	17578 <i>Stylidium udusicola</i>			
444.	16882 <i>Synaphea aephyrsa</i>			
445.	2329 <i>Synaphea spinulosa</i>			
446.	15532 <i>Synaphea spinulosa</i> subsp. <i>spinulosa</i>			
447.	1036 <i>Tetraria octandra</i>			
448.	4539 <i>Tetratheca paucifolia</i>			
449.	11032 <i>Thelymitra apiculata</i>		P4	
450.	10862 <i>Thelymitra stellata</i> (Star Orchid)		T	
451.	6060 <i>Thryptomene mucronulata</i>			
452.	1334 <i>Thysanotus glaucus</i>		P4	
453.	1343 <i>Thysanotus patersonii</i>			
454.	1351 <i>Thysanotus sparteus</i>			
455.	1357 <i>Thysanotus thyrsoideus</i>			
456.	1358 <i>Thysanotus triandrus</i>			
457.	6280 <i>Trachymene pilosa</i> (Native Parsnip)			
458.	1481 <i>Tribonanthes australis</i>			
459.	1361 <i>Tricoryne elatior</i> (Yellow Autumn Lily)			
460.	29481 <i>Tricoryne</i> sp. <i>Eneabba</i> (E.A. Griffin 1200)			
461.	1038 <i>Tricostularia neesii</i>			
462.	4292 <i>Trifolium campestre</i> (Hop Clover)	Y		
463.	4298 <i>Trifolium hirtum</i> (Rose Clover)	Y		
464.	148 <i>Triglochin muelleri</i>			
465.	151 <i>Triglochin striata</i>			
466.	13479 <i>Trymalium ledifolium</i> var. <i>rosmarinifolium</i>			
467.	7665 <i>Velleia trinervis</i>			
468.	15725 <i>Verbesina encelioides</i>	Y		
469.	7666 <i>Verreauxia reinwardtii</i> (Common Verreauxia)			
470.	12411 <i>Verticordia densiflora</i> var. <i>cespitosa</i>			
471.	15432 <i>Verticordia densiflora</i> var. <i>densiflora</i>			
472.	6077 <i>Verticordia drummondii</i> (Drummond's Featherflower)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
473.	14714 <i>Verticordia lindleyi</i> subsp. <i>lindleyi</i>		P4	
474.	10822 <i>Verticordia nobilis</i>			
475.	6107 <i>Verticordia pennigera</i>			
476.	6109 <i>Verticordia picta</i> (<i>Painted Featherflower</i>)			
477.	7386 <i>Wahlenbergia gracilentia</i> (<i>Annual Bluebell</i>)			
478.	8282 <i>Waitzia suaveolens</i> (<i>Fragrant Waitzia</i>)			
479.	13333 <i>Waitzia suaveolens</i> var. <i>suaveolens</i>			
480.	1252 <i>Xanthorrhoea drummondii</i>			
481.	1256 <i>Xanthorrhoea preissii</i> (<i>Grass tree, Palga</i>)			
482.	6285 <i>Xanthosia ciliata</i>			
483.	6287 <i>Xanthosia fruticulosa</i>			
484.	6289 <i>Xanthosia huegelii</i>			
485.	1049 <i>Zantedeschia aethiopica</i> (<i>Arum Lily</i>)	Y		

Conservation Codes

- T - Rare or likely to become extinct
- X - Presumed extinct
- IA - Protected under international agreement
- S - Other specially protected fauna
- 1 - Priority 1
- 2 - Priority 2
- 3 - Priority 3
- 4 - Priority 4
- 5 - Priority 5

¹ For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.

APPENDIX C EPBC PROTECTED MATTERS SEARCH TOOL



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 20/10/17 14:21:46

[Summary](#)

[Details](#)


[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)

[Acknowledgements](#)



This map may contain data which are
©Commonwealth of Australia
(Geoscience Australia), ©PSMA 2010

[Coordinates](#)

Buffer: 10.0Km



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

World Heritage Properties:	None
National Heritage Places:	None
Wetlands of International Importance:	None
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	None
Listed Threatened Ecological Communities:	1
Listed Threatened Species:	29
Listed Migratory Species:	9

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	None
Commonwealth Heritage Places:	None
Listed Marine Species:	15
Whales and Other Cetaceans:	None
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Commonwealth Reserves Marine:	None

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	2
Regional Forest Agreements:	None
Invasive Species:	16
Nationally Important Wetlands:	None
Key Ecological Features (Marine)	None

Details

Matters of National Environmental Significance

Listed Threatened Ecological Communities [\[Resource Information \]](#)

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Name	Status	Type of Presence
Banksia Woodlands of the Swan Coastal Plain ecological community	Endangered	Community likely to occur within area

Listed Threatened Species [\[Resource Information \]](#)

Name	Status	Type of Presence
------	--------	------------------

Birds

Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
---	-----------------------	--

Calyptorhynchus latirostris Carnaby's Cockatoo, Short-billed Black-Cockatoo [59523]	Endangered	Species or species habitat known to occur within area
--	------------	---

Leipoa ocellata Malleefowl [934]	Vulnerable	Species or species habitat likely to occur within area
---	------------	--

Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat may occur within area
---	-----------------------	--

Rostratula australis Australian Painted Snipe [77037]	Endangered	Species or species habitat may occur within area
--	------------	--

Mammals

Dasyurus geoffroii Chuditch, Western Quoll [330]	Vulnerable	Species or species habitat likely to occur within area
---	------------	--

Parantechinus apicalis Dibbler [313]	Endangered	Species or species habitat may occur within area
---	------------	--

Plants

Acacia forrestiana Forest's Wattle [17235]	Vulnerable	Species or species habitat known to occur within area
---	------------	---

Andersonia gracilis Slender Andersonia [14470]	Endangered	Species or species habitat likely to occur within area
---	------------	--

Anigozanthos viridis subsp. terraspectans Dwarf Green Kangaroo Paw [3435]	Vulnerable	Species or species habitat likely to occur within area
--	------------	--

Name	Status	Type of Presence
Caladenia huegelii King Spider-orchid, Grand Spider-orchid, Rusty Spider-orchid [7309]	Endangered	Species or species habitat may occur within area
Chamelaucium sp. Cataby (G.J.Keighery 11009) Griffin's Waxflower [82509]	Vulnerable	Species or species habitat known to occur within area
Chamelaucium sp. Gingin (N.G.Marchant 6) Gingin Wax [88881]	Endangered	Species or species habitat may occur within area
Conospermum densiflorum subsp. unicephalum One-headed Smokebush [64871]	Endangered	Species or species habitat likely to occur within area
Drakaea elastica Glossy-leaved Hammer Orchid, Glossy-leaved Hammer Orchid, Warty Hammer Orchid [16753]	Endangered	Species or species habitat likely to occur within area
Eleocharis keigheryi Keighery's Eleocharis [64893]	Vulnerable	Species or species habitat known to occur within area
Eucalyptus absita Badgingarra Box [24260]	Endangered	Species or species habitat may occur within area
Eucalyptus dolorosa Dandaragan Mallee, Mount Misery Mallee [56709]	Endangered	Species or species habitat known to occur within area
Eucalyptus impensa Eneabba Mallee [56711]	Endangered	Species or species habitat may occur within area
Eucalyptus leprophloia Scaly Butt Mallee, Scaly-butt Mallee [56712]	Endangered	Species or species habitat may occur within area
Eucalyptus recta Silver Mallet [56430]	Endangered	Species or species habitat may occur within area
Eucalyptus x balanites Cadda Road Mallee, Cadda Mallee [87816]	Endangered	Species or species habitat may occur within area
Grevillea calliantha Foote's Grevillea, Cataby Grevillea, Black Magic Grevillea [56339]	Endangered	Species or species habitat known to occur within area
Hakea megalosperma Lesueur Hakea [10505]	Vulnerable	Species or species habitat likely to occur within area
Hemiandra gardneri Red Snakebush [7945]	Endangered	Species or species habitat may occur within area
Leucopogon obtectus Hidden Beard-heath [19614]	Endangered	Species or species habitat may occur within area
Paracaleana dixonii Sandplain Duck Orchid [86882]	Endangered	Species or species habitat may occur within area
Thelymitra dedmaniarum Cinnamon Sun Orchid [65105]	Endangered	Species or species habitat may occur within area

Name	Status	Type of Presence
Thelymitra stellata Star Sun-orchid [7060]	Endangered	Species or species habitat may occur within area

Listed Migratory Species [Resource Information]

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Migratory Marine Birds		
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area

Migratory Terrestrial Species

Motacilla cinerea Grey Wagtail [642]		Species or species habitat may occur within area
---	--	--

Migratory Wetlands Species

Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat may occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat may occur within area
Pandion haliaetus Osprey [952]		Species or species habitat known to occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat likely to occur within area

Other Matters Protected by the EPBC Act

Listed Marine Species [Resource Information]

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Birds		
Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat may occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea alba Great Egret, White Egret [59541]		Species or species habitat likely to occur within area

Name	Threatened	Type of Presence
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Haliaeetus leucogaster White-bellied Sea-Eagle [943]		Species or species habitat likely to occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Motacilla cinerea Grey Wagtail [642]		Species or species habitat may occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat may occur within area
Pandion haliaetus Osprey [952]		Species or species habitat known to occur within area
Rostratula benghalensis (sensu lato) Painted Snipe [889]	Endangered*	Species or species habitat may occur within area
Thinornis rubricollis Hooded Plover [59510]		Species or species habitat may occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat likely to occur within area

Extra Information

State and Territory Reserves	[Resource Information]
Name	State
Unnamed WA27993	WA
Unnamed WA39571	WA

Invasive Species**[Resource Information]**

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resources Audit, 2001.

Name	Status	Type of Presence
Birds		
Anas platyrhynchos Mallard [974]		Species or species habitat likely to occur within area
Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
Streptopelia senegalensis Laughing Turtle-dove, Laughing Dove [781]		Species or species habitat likely to occur within area
Sturnus vulgaris Common Starling [389]		Species or species habitat likely to occur within area
Mammals		
Felis catus Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area
Oryctolagus cuniculus Rabbit, European Rabbit [128]		Species or species habitat likely to occur within area
Rattus rattus Black Rat, Ship Rat [84]		Species or species habitat likely to occur within area
Vulpes vulpes Red Fox, Fox [18]		Species or species habitat likely to occur within area
Plants		
Asparagus asparagoides Bridal Creeper, Bridal Veil Creeper, Smilax, Florist's Smilax, Smilax Asparagus [22473]		Species or species habitat likely to occur within area
Brachiaria mutica Para Grass [5879]		Species or species habitat may occur within area
Cenchrus ciliaris Buffel-grass, Black Buffel-grass [20213]		Species or species habitat may occur within area
Chrysanthemoides monilifera Bitou Bush, Boneseed [18983]		Species or species habitat may occur within area
Genista sp. X Genista monspessulana Broom [67538]		Species or species habitat may occur within area
Olea europaea Olive, Common Olive [9160]		Species or species habitat may occur within area
Pinus radiata Radiata Pine Monterey Pine, Insignis Pine, Wilding Pine [20780]		Species or species habitat may occur within area
Tamarix aphylla Athel Pine, Athel Tree, Tamarisk, Athel Tamarisk, Athel Tamarix, Desert Tamarisk, Flowering		Species or species habitat likely to occur

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-30.74778 115.61139

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [Office of Environment and Heritage, New South Wales](#)
- [Department of Environment and Primary Industries, Victoria](#)
- [Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [Department of Environment, Water and Natural Resources, South Australia](#)
- [Department of Land and Resource Management, Northern Territory](#)
- [Department of Environmental and Heritage Protection, Queensland](#)
- [Department of Parks and Wildlife, Western Australia](#)
- [Environment and Planning Directorate, ACT](#)
- [Birdlife Australia](#)
- [Australian Bird and Bat Banding Scheme](#)
- [Australian National Wildlife Collection](#)
- Natural history museums of Australia
- [Museum Victoria](#)
- [Australian Museum](#)
- [South Australian Museum](#)
- [Queensland Museum](#)
- [Online Zoological Collections of Australian Museums](#)
- [Queensland Herbarium](#)
- [National Herbarium of NSW](#)
- [Royal Botanic Gardens and National Herbarium of Victoria](#)
- [Tasmanian Herbarium](#)
- [State Herbarium of South Australia](#)
- [Northern Territory Herbarium](#)
- [Western Australian Herbarium](#)
- [Australian National Herbarium, Canberra](#)
- [University of New England](#)
- [Ocean Biogeographic Information System](#)
- [Australian Government, Department of Defence Forestry Corporation, NSW](#)
- [Geoscience Australia](#)
- [CSIRO](#)
- [Australian Tropical Herbarium, Cairns](#)
- [eBird Australia](#)
- [Australian Government – Australian Antarctic Data Centre](#)
- [Museum and Art Gallery of the Northern Territory](#)
- [Australian Government National Environmental Science Program](#)
- [Australian Institute of Marine Science](#)
- [Reef Life Survey Australia](#)
- [American Museum of Natural History](#)
- [Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
- [Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
- Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.

APPENDIX D

SIGNIFICANT FLORA RECORDS - TPFL AND WAHERB

SIGNIFICANT FLORA RECORDS - TPFL AND WAHERB

Taxon	ConsStatus	WARank	Easting	Northing	Source
<i>Acacia cummingiana</i>	3		375470.0561	6605694.508	WAHERB
<i>Acacia epacantha</i>	3		375470.0561	6605694.508	WAHERB
<i>Acacia epacantha</i>	3		375470.0561	6605694.508	WAHERB
<i>Acacia forrestiana</i>	T	VU	367554.1158	6603795.499	TPFL
<i>Acacia forrestiana</i>	T		367826.4169	6603940.519	WAHERB
<i>Acacia forrestiana</i>	T		375459.3649	6606618.082	WAHERB
<i>Acacia forrestiana</i>	T		365901.4586	6604655.735	WAHERB
<i>Acacia forrestiana</i>	T		375480.75	6604770.933	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361207.2789	6597205.74	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		359587.5841	6599032.311	WAHERB
<i>Acacia plicata</i>	3		359587.5841	6599032.311	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		361231.1935	6595358.402	WAHERB
<i>Acacia plicata</i>	3		362414.1884	6597664.715	WAHERB
<i>Acacia splendens</i>	T		375480.75	6604770.933	WAHERB
<i>Acacia splendens</i>	T		375470.0561	6605694.508	WAHERB
<i>Anigozanthos humilis</i> subsp. <i>Badgingarra</i> (S.D. Hopper 7114)	2		364328.1778	6602788.398	TPFL
<i>Anigozanthos humilis</i> subsp. <i>Badgingarra</i> (S.D. Hopper 7114)	2		365970.623	6599113.932	TPFL
<i>Anigozanthos humilis</i> subsp. <i>Badgingarra</i> (S.D. Hopper 7114)	2		365970.623	6599113.932	WAHERB
<i>Anigozanthos humilis</i> subsp. <i>Badgingarra</i> (S.D. Hopper 7114)	2		366100.0432	6585231.98	WAHERB
<i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i>	4		361165.255	6594847.164	TPFL
<i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i>	4		367225.1688	6601482.936	TPFL
<i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i>	4		366369.2466	6594660.026	TPFL
<i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i>	4		367520.8218	6602828.234	WAHERB
<i>Anigozanthos humilis</i> subsp. <i>chrysanthus</i>	4		365924.5025	6602808.435	WAHERB
<i>Asterolasia drummondii</i>	4		366964.2488	6594643.708	TPFL
<i>Asterolasia drummondii</i>	4		366964.2488	6594643.708	TPFL
<i>Asterolasia drummondii</i>	4		366964.2488	6594643.708	TPFL
<i>Asterolasia drummondii</i>	4		367385.4984	6603578.81	TPFL
<i>Asterolasia drummondii</i>	4		367498.0524	6604675.528	WAHERB
<i>Asterolasia drummondii</i>	4		367612.0101	6595439.125	WAHERB
<i>Asterolasia drummondii</i>	4		367612.0101	6595439.125	WAHERB
<i>Asterolasia drummondii</i>	4		364965.3675	6594451.532	WAHERB
<i>Asterolasia drummondii</i>	4		361207.2789	6597205.74	WAHERB
<i>Asterolasia drummondii</i>	4		358016.3347	6597163.967	WAHERB
<i>Asterolasia drummondii</i>	4		366016.7902	6595419.3	WAHERB
<i>Asterolasia drummondii</i>	4		366016.7902	6595419.3	WAHERB
<i>Asterolasia drummondii</i>	4		367536.4792	6594595.371	WAHERB
<i>Asterolasia drummondii</i>	4		366981.5467	6594815.524	WAHERB
<i>Asterolasia drummondii</i>	4		366981.1641	6594846.337	WAHERB
<i>Asterolasia drummondii</i>	4		367498.0524	6604675.528	WAHERB
<i>Asterolasia drummondii</i>	4		364958.3829	6595005.719	WAHERB
<i>Babingtonia delicata</i>	1		364173.3102	6585634.666	TPFL
<i>Babingtonia delicata</i>	1		365927.2605	6585595.194	TPFL
<i>Babingtonia delicata</i>	1		365927.2605	6585595.194	TPFL
<i>Babingtonia delicata</i>	1		361096.646	6603701.688	WAHERB
<i>Babingtonia delicata</i>	1		365972.0255	6585609.945	WAHERB
<i>Babingtonia delicata</i>	1		364321.8392	6585792.752	WAHERB
<i>Babingtonia delicata</i>	1		366062.8753	6585747.77	WAHERB
<i>Babingtonia delicata</i>	1		364786.5943	6585457.651	WAHERB

Taxon	ConsStatus	WARank	Easting	Northing	Source
<i>Banksia dallanneyi</i> subsp. <i>pollostata</i>	3		367699.0244	6584571.039	WAHERB
<i>Banksia kippistiana</i> var. <i>paenepeccata</i>	3		364684.8739	6592507.803	WAHERB
<i>Banksia prionophylla</i>	1		364331.4475	6592651.434	TPFL
<i>Banksia prionophylla</i>	1		364217.024	6592656.08	WAHERB
<i>Banksia prionophylla</i>	1		364444.9394	6593552.02	WAHERB
<i>Banksia prionophylla</i>	1		364317.825	6592842.819	WAHERB
<i>Banksia prionophylla</i>	1		364176.3589	6592723.297	WAHERB
<i>Banksia pteridifolia</i> subsp. <i>vernalis</i>	3		375470.0561	6605694.508	WAHERB
<i>Beaufortia eriocephala</i>	3		364600.8352	6592304.54	TPFL
<i>Beaufortia eriocephala</i>	3		362826.4298	6595378.94	WAHERB
<i>Beaufortia eriocephala</i>	3		366667.4176	6602264.301	WAHERB
<i>Beaufortia eriocephala</i>	3		364539.1616	6592413.84	WAHERB
<i>Beaufortia eriocephala</i>	3		362850.0795	6593531.715	WAHERB
<i>Boronia scabra</i> subsp. <i>condensata</i>	2		367498.0524	6604675.528	WAHERB
<i>Calytrix ecalycata</i> subsp. <i>brevis</i>	3		366903.028	6583772.073	WAHERB
<i>Calytrix ecalycata</i> subsp. <i>brevis</i>	3		375674.0558	6588114.919	WAHERB
<i>Calytrix ecalycata</i> subsp. <i>brevis</i>	3		362524.0226	6588501.953	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T	VU	366800.1175	6595011.145	TPFL
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T	VU	366421.0031	6595560.696	TPFL
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T	VU	366166.7449	6594633.775	TPFL
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T	VU	365658.1427	6594904.547	TPFL
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		366662.9428	6594780.747	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		366016.7902	6595419.3	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		365993.7009	6597266.618	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		366166.7449	6594633.775	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		366016.7902	6595419.3	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		366028.3384	6594495.695	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		365036.4205	6597254.614	WAHERB
<i>Chamelaucium</i> sp. <i>Cataby</i> (G.J. Keighery 11009)	T		365036.4205	6597254.614	WAHERB
<i>Conostephium magnum</i>	4		361180.6663	6597205.395	WAHERB
<i>Dampiera tephrea</i>	2		362006.4529	6597660.253	TPFL
<i>Dampiera tephrea</i>	2		369223.6728	6605602.863	TPFL
<i>Dampiera tephrea</i>	2		360885.9842	6594821.037	WAHERB
<i>Dampiera tephrea</i>	2		375695.1891	6586298.427	WAHERB
<i>Dampiera tephrea</i>	2		375470.0561	6605694.508	WAHERB
<i>Dampiera tephrea</i>	2		375470.0561	6605694.508	WAHERB
<i>Dampiera tephrea</i>	2		361207.2789	6597205.74	WAHERB
<i>Desmocladus microcarpus</i>	2		359587.5841	6599032.311	WAHERB
<i>Drakaea elastica</i>	T	CR	365518.2185	6585374.546	TPFL
<i>Drakaea elastica</i>	T		364216.3401	6584341.846	WAHERB
<i>Drosera marchantii</i> subsp. <i>prophylla</i>	3		364528.3456	6592429.112	TPFL
<i>Drosera marchantii</i> subsp. <i>prophylla</i>	3		364528.8156	6592429.783	WAHERB
<i>Eleocharis keigheryi</i>	T	VU	362481.9588	6597912.682	TPFL
<i>Eleocharis keigheryi</i>	T		362455.7382	6597881.638	WAHERB
<i>Eleocharis keigheryi</i>	T		361245.9452	6594219.318	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367634.8342	6593591.919	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367498.0524	6604675.528	WAHERB
<i>Eucalyptus abdita</i>	2		367511.9023	6603551.769	WAHERB
<i>Eucalyptus annuliformis</i>	1		383604.697	6591929.89	WAHERB
<i>Eucalyptus dolorosa</i>	T	CR	367515.5066	6603826.727	TPFL
<i>Eucalyptus dolorosa</i>	T		367611.2481	6603636.107	WAHERB
<i>Eucalyptus dolorosa</i>	T		367523.8218	6602828.234	WAHERB

Taxon	ConsStatus	WARank	Easting	Northing	Source
<i>Eucalyptus dolorosa</i>	T		367502.0524	6604675.528	WAHERB
<i>Eucalyptus dolorosa</i>	T		367503.0524	6604675.528	WAHERB
<i>Eucalyptus dolorosa</i>	T		367504.0524	6604675.528	WAHERB
<i>Eucalyptus dolorosa</i>	T		367505.0524	6604675.528	WAHERB
<i>Eucalyptus dolorosa</i>	T		367506.0524	6604675.528	WAHERB
<i>Eucalyptus dolorosa</i>	T		367657.927	6603230.101	WAHERB
<i>Eucalyptus dolorosa</i>	T		367619.2481	6603636.107	WAHERB
<i>Eucalyptus dolorosa</i>	T		367509.0524	6604675.528	WAHERB
<i>Eucalyptus dolorosa</i>	T		367510.0524	6604675.528	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		362155.2572	6596430.445	TPFL
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364202.2152	6592299.491	TPFL
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364475.4327	6591717.857	TPFL
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		362839.7455	6594899.547	TPFL
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364434.5649	6595399.238	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		361245.1935	6595358.402	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		359626.7618	6597184.971	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		361247.1935	6595358.402	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364438.5649	6595399.238	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		362891.7421	6591684.375	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		363433.4651	6595140.129	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		361251.1935	6595358.402	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		359608.5841	6599032.311	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364443.5649	6595399.238	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364498.0425	6591748.67	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364445.5649	6595399.238	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		364423.2006	6597246.562	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		362784.3481	6595021.666	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		362620.4656	6594883.311	WAHERB
<i>Eucalyptus macrocarpa</i> subsp. <i>elachantha</i>	4		363421.3532	6594708.742	WAHERB
<i>Gastrolobium nudum</i>	2		366045.7902	6595419.3	WAHERB
<i>Grevillea calliantha</i>	T	CR	367573.8436	6601332.379	TPFL
<i>Grevillea calliantha</i>	T	CR	366899.9282	6601179.274	TPFL
<i>Grevillea calliantha</i>	T	CR	363873.0871	6603565.606	TPFL
<i>Grevillea calliantha</i>	T	CR	363158.1064	6604824.368	TPFL
<i>Grevillea calliantha</i>	T	CR	364119.3698	6602468.606	TPFL
<i>Grevillea calliantha</i>	T	CR	364378.3839	6600504.217	TPFL
<i>Grevillea calliantha</i>	T	CR	366272.4553	6605208.411	TPFL
<i>Grevillea calliantha</i>	T		365977.5578	6600961.13	WAHERB
<i>Grevillea calliantha</i>	T		365978.5578	6600961.13	WAHERB
<i>Grevillea calliantha</i>	T		362740.3503	6604615.439	WAHERB
<i>Grevillea calliantha</i>	T		367553.8218	6602828.234	WAHERB
<i>Grevillea calliantha</i>	T		365981.5578	6600961.13	WAHERB
<i>Grevillea calliantha</i>	T		365982.5578	6600961.13	WAHERB
<i>Grevillea calliantha</i>	T		363568.1727	6602612.037	WAHERB
<i>Grevillea calliantha</i>	T		363569.1727	6602612.037	WAHERB
<i>Grevillea calliantha</i>	T		363271.7947	6603476.566	WAHERB
<i>Grevillea calliantha</i>	T		364613.0224	6600741.587	WAHERB
<i>Grevillea calliantha</i>	T		364414.8477	6599093.883	WAHERB
<i>Grevillea calliantha</i>	T		363232.3378	6604720.144	WAHERB
<i>Grevillea calliantha</i>	T		362797.5478	6600920.808	WAHERB
<i>Grevillea drummondii</i>	4		375308.8219	6594990.491	TPFL
<i>Grevillea drummondii</i>	4		366828.6376	6594857.523	TPFL
<i>Grevillea drummondii</i>	4		375334.7122	6595052.314	TPFL
<i>Grevillea drummondii</i>	4		358059.3347	6597163.967	WAHERB
<i>Grevillea drummondii</i>	4		365568.3286	6596521.677	WAHERB
<i>Grevillea drummondii</i>	4		366038.7009	6597266.618	WAHERB
<i>Grevillea drummondii</i>	4		367658.0101	6595439.125	WAHERB
<i>Grevillea drummondii</i>	4		362873.4298	6595378.94	WAHERB
<i>Grevillea drummondii</i>	4		365006.3829	6595005.719	WAHERB

Taxon	ConsStatus	WARank	Easting	Northing	Source
<i>Grevillea drummondii</i>	4		365007.3829	6595005.719	WAHERB
<i>Grevillea florida</i>	3		359587.5841	6599032.311	TPFL
<i>Grevillea florida</i>	3		359637.5841	6599032.311	WAHERB
<i>Grevillea olivacea</i>	4		376171.6877	6588551.202	WAHERB
<i>Grevillea olivacea</i>	4		376172.6877	6588551.202	WAHERB
<i>Grevillea saccata</i>	4		375523.0561	6605694.508	WAHERB
<i>Grevillea saccata</i>	4		386690.9265	6606741.649	WAHERB
<i>Grevillea saccata</i>	4		386691.9265	6606741.649	WAHERB
<i>Grevillea synapheae</i> subsp. <i>A Flora of Australia (S.D. Hopper 6333)</i>	1		367503.9577	6603761.071	TPFL
<i>Grevillea synapheae</i> subsp. <i>A Flora of Australia (S.D. Hopper 6333)</i>	1		367554.0524	6604675.528	WAHERB
<i>Grevillea synapheae</i> subsp. <i>A Flora of Australia (S.D. Hopper 6333)</i>	1		365958.4586	6604655.735	WAHERB
<i>Grevillea synapheae</i> subsp. <i>A Flora of Australia (S.D. Hopper 6333)</i>	1		367556.0524	6604675.528	WAHERB
<i>Grevillea synapheae</i> subsp. <i>A Flora of Australia (S.D. Hopper 6333)</i>	1		367557.0524	6604675.528	WAHERB
<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	1		362764.0341	6602753.127	TPFL
<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	1		361126.1458	6603064.65	TPFL
<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	1		360919.5772	6601550.017	TPFL
<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	1		361219.4866	6600900.29	WAHERB
<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	1		359648.5841	6599032.311	WAHERB
<i>Grevillea synapheae</i> subsp. <i>minyulo</i>	1		359649.5841	6599032.311	WAHERB
<i>Grevillea thyrsoides</i> subsp. <i>thyrsoides</i>	3		361867.9744	6596763.05	WAHERB
<i>Grevillea thyrsoides</i> subsp. <i>thyrsoides</i>	3		363383.7857	6590058.053	WAHERB
<i>Grevillea thyrsoides</i> subsp. <i>thyrsoides</i>	3		364446.9953	6592227.053	WAHERB
<i>Grevillea thyrsoides</i> subsp. <i>thyrsoides</i>	3		364376.4718	6592146.885	WAHERB
<i>Grevillea thyrsoides</i> subsp. <i>thyrsoides</i>	3		362794.5865	6594792.687	WAHERB
<i>Guichenotia alba</i>	3		362918.0795	6593531.715	WAHERB
<i>Haemodorum loratum</i>	3		365902.2941	6588163.851	TPFL
<i>Haemodorum loratum</i>	3		366178.258	6588030.204	WAHERB
<i>Hakea longiflora</i>	3		366063.7009	6597266.618	WAHERB
<i>Hakea longiflora</i>	3		375530.3649	6606618.082	WAHERB
<i>Hakea longiflora</i>	3		361989.0176	6597050.946	WAHERB
<i>Hakea longiflora</i>	3		366066.7009	6597266.618	WAHERB
<i>Hemigenia curvifolia</i>	2		375554.75	6604770.933	WAHERB
<i>Hemigenia curvifolia</i>	2		375555.75	6604770.933	WAHERB
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		362145.6012	6597591.205	TPFL
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		362067.3602	6597473.247	TPFL
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		362145.6012	6597591.205	TPFL
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		368539.4129	6606351.083	TPFL
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		362017.0132	6597195.035	WAHERB
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		361284.2789	6597205.74	WAHERB
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		361285.2789	6597205.74	WAHERB
<i>Hypocalymma</i> sp. <i>Cataby (G.J. Keighery 5151)</i>	2		361274.3268	6598129.351	WAHERB
<i>Hypocalymma</i> sp. <i>Dandaragan (C.A. Gardner 9014)</i>	1		375539.3649	6606618.082	WAHERB
<i>Hypocalymma tetrapterum</i>	3		362143.3834	6597354.056	TPFL
<i>Hypocalymma tetrapterum</i>	3		362143.3834	6597354.056	TPFL
<i>Hypocalymma tetrapterum</i>	3		366565.4492	6594638.748	TPFL
<i>Hypocalymma tetrapterum</i>	3		361288.2789	6597205.74	WAHERB
<i>Hypocalymma tetrapterum</i>	3		361289.2789	6597205.74	WAHERB
<i>Hypocalymma tetrapterum</i>	3		366076.7009	6597266.618	WAHERB
<i>Hypocalymma tetrapterum</i>	3		361291.2789	6597205.74	WAHERB
<i>Hypocalymma tetrapterum</i>	3		360994.7859	6595539.034	WAHERB
<i>Hypocalymma tetrapterum</i>	3		359673.5841	6599032.311	WAHERB
<i>Hypocalymma tetrapterum</i>	3		359674.5841	6599032.311	WAHERB
<i>Hypocalymma tetrapterum</i>	3		359675.5841	6599032.311	WAHERB
<i>Hypocalymma tetrapterum</i>	3		359676.5841	6599032.311	WAHERB
<i>Hypocalymma tetrapterum</i>	3		359677.5841	6599032.311	WAHERB
<i>Hypocalymma tetrapterum</i>	3		364489.2006	6597246.562	WAHERB
<i>Hypocalymma tetrapterum</i>	3		361937.5034	6597213.981	WAHERB
<i>Hypolaena robusta</i>	4		364514.5649	6595399.238	WAHERB
<i>Hypolaena robusta</i>	4		367511.3078	6594556.872	WAHERB

Taxon	ConsStatus	WARank	Easting	Northing	Source
<i>Hypolaena robusta</i>	4		367512.3078	6594556.872	WAHERB
<i>Jacksonia carduacea</i>	3		360096.0065	6594802.957	WAHERB
<i>Jacksonia carduacea</i>	3		364028.4491	6593009.854	WAHERB
<i>Jacksonia carduacea</i>	3		364167.8109	6593147.852	WAHERB
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		367739.6795	6603850.106	TPFL
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		367597.0524	6604675.528	WAHERB
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		367598.0524	6604675.528	WAHERB
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		366002.4586	6604655.735	WAHERB
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		366003.4586	6604655.735	WAHERB
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		367625.4842	6603773.603	WAHERB
<i>Lasiopetalum</i> sp. Hill River (T.N. Stoate 5)	1		367626.4842	6603773.603	WAHERB
<i>Lechenaultia galactites</i>	3		375575.0561	6605694.508	WAHERB
<i>Lepidobolus quadratus</i>	3		364597.7255	6589857.349	WAHERB
<i>Lepidobolus quadratus</i>	3		362135.7645	6595368.7	WAHERB
<i>Lepidobolus quadratus</i>	3		364805.406	6592508.183	WAHERB
<i>Lepidobolus quadratus</i>	3		364600.7255	6589857.349	WAHERB
<i>Lepidobolus quadratus</i>	3		364672.0016	6592371.006	WAHERB
<i>Lepidobolus quadratus</i>	3		364673.0016	6592371.006	WAHERB
<i>Leucopogon foliosus</i>	3		364545.2994	6592521.027	WAHERB
<i>Leucopogon foliosus</i>	3		364314.9666	6592718.966	WAHERB
<i>Leucopogon foliosus</i>	3		364791.9121	6592521.018	WAHERB
<i>Macarthuria keigheryi</i>	T		362635.0401	6588528.951	WAHERB
<i>Podotheca pritzelii</i>	3		366063.5578	6600961.13	WAHERB
<i>Ptychosema pusillum</i>	T	VU	365308.2289	6585599.709	TPFL
<i>Ptychosema pusillum</i>	T		364333.3401	6584341.846	WAHERB
<i>Rhedinocarpus suffruticosus</i>	1		375598.75	6604770.933	WAHERB
<i>Stylidium aeoniooides</i>	4		367826.4169	6603940.519	TPFL
<i>Stylidium aeoniooides</i>	4		366978.4893	6595061.804	TPFL
<i>Stylidium aeoniooides</i>	4		364217.024	6592656.08	TPFL
<i>Stylidium aeoniooides</i>	4		364217.024	6592656.08	TPFL
<i>Stylidium aeoniooides</i>	4		364517.2006	6597246.562	WAHERB
<i>Stylidium aeoniooides</i>	4		367618.0524	6604675.528	WAHERB
<i>Stylidium aeoniooides</i>	4		363328.0432	6592643.224	WAHERB
<i>Stylidium aeoniooides</i>	4		366138.7902	6595419.3	WAHERB
<i>Stylidium aeoniooides</i>	4		362109.9711	6597105.722	WAHERB
<i>Stylidium aeoniooides</i>	4		364479.3844	6592794.075	WAHERB
<i>Stylidium aeoniooides</i>	4		366946.6222	6594844.355	WAHERB
<i>Stylidium aeoniooides</i>	4		362043.0176	6597050.946	WAHERB
<i>Stylidium aeoniooides</i>	4		367054.5703	6594876.378	WAHERB
<i>Stylidium aeoniooides</i>	4		366954.1028	6594915.136	WAHERB
<i>Stylidium periscelanthum</i>	3		361309.6663	6597205.395	WAHERB
<i>Thelymitra apiculata</i>	4		362706.1034	6590017.706	WAHERB
<i>Thelymitra apiculata</i>	4		364430.9357	6592978.169	WAHERB
<i>Thelymitra apiculata</i>	4		364688.8987	6592418.831	WAHERB
<i>Thelymitra stellata</i>	T	EN	364589.6971	6592617.677	TPFL
<i>Thysanotus glaucus</i>	4		364208.0744	6591837.629	TPFL
<i>Thysanotus glaucus</i>	4		375332.9211	6595206.263	TPFL
<i>Thysanotus glaucus</i>	4		364624.7255	6589857.349	WAHERB
<i>Thysanotus glaucus</i>	4		363007.7421	6591684.375	WAHERB
<i>Thysanotus glaucus</i>	4		363008.7421	6591684.375	WAHERB
<i>Verticordia huegelii</i> var. <i>tridens</i>	3		375606.0561	6605694.508	WAHERB
<i>Verticordia insignis</i> subsp. <i>eomagis</i>	3		385133.0531	6598410.9	WAHERB
<i>Verticordia lindleyi</i> subsp. <i>lindleyi</i>	4		359516.4709	6594856.492	TPFL

APPENDIX E REGIONAL BIRD RECORDS - NATUREMAP

Name ID	Species Name	Status
24260	<i>Acanthiza apicalis</i> (Inland Thornbill)	
24261	<i>Acanthiza chrysorrhoa</i> (Yellow-rumped Thornbill)	
24262	<i>Acanthiza inornata</i> (Western Thornbill)	
24560	<i>Acanthorhynchus superciliosus</i> (Western Spinebill)	
25536	<i>Accipiter fasciatus</i> (Brown Goshawk)	
25544	<i>Aegotheles cristatus</i> (Australian Owlet-nightjar)	
24310	<i>Anas castanea</i> (Chestnut Teal)	
24312	<i>Anas gracilis</i> (Grey Teal)	
24313	<i>Anas platyrhynchos</i> (Mallard)	
24315	<i>Anas rhynchotis</i> (Australasian Shoveler)	
24316	<i>Anas superciliosa</i> (Pacific Black Duck)	
47414	<i>Anhinga novaehollandiae</i> (Australasian Darter)	
24561	<i>Anthochaera carunculata</i> (Red Wattlebird)	
24562	<i>Anthochaera lunulata</i> (Western Little Wattlebird)	
24285	<i>Aquila audax</i> (Wedge-tailed Eagle)	
41324	<i>Ardea modesta</i> (Great Egret)	IA
24340	<i>Ardea novaehollandiae</i> (White-faced Heron)	
24341	<i>Ardea pacifica</i> (White-necked Heron)	
25566	<i>Artamus cinereus</i> (Black-faced Woodswallow)	
24356	<i>Artamus personatus</i> (Masked Woodswallow)	
24318	<i>Aythya australis</i> (Hardhead)	
24319	<i>Biziura lobata</i> (Musk Duck)	
24723	<i>Cacatua pastinator</i> subsp. <i>butleri</i> (Butler's Corella)	
25716	<i>Cacatua sanguinea</i> (Little Corella)	
42307	<i>Cacomantis pallidus</i> (Pallid Cuckoo)	
24779	<i>Calidris acuminata</i> (Sharp-tailed Sandpiper)	IA
24784	<i>Calidris ferruginea</i> (Curlew Sandpiper)	T
24788	<i>Calidris ruficollis</i> (Red-necked Stint)	IA
24734	<i>Calyptorhynchus latirostris</i> (Carnaby's Black-Cockatoo)	T
24377	<i>Charadrius ruficapillus</i> (Red-capped Plover)	
24321	<i>Chenonetta jubata</i> (Australian Wood Duck)	
47909	<i>Cheramoeca leucosterna</i> (White-backed Swallow)	
24288	<i>Circus approximans</i> (Swamp Harrier)	
24774	<i>Cladorhynchus leucocephalus</i> (Banded Stilt)	
25675	<i>Colluricincla harmonica</i> (Grey Shrike-thrush)	
25568	<i>Coracina novaehollandiae</i> (Black-faced Cuckoo-shrike)	
24416	<i>Corvus bennetti</i> (Little Crow)	
25592	<i>Corvus coronoides</i> (Australian Raven)	
24671	<i>Coturnix pectoralis</i> (Stubble Quail)	
24420	<i>Cracticus nigrogularis</i> (Pied Butcherbird)	
25595	<i>Cracticus tibicen</i> (Australian Magpie)	
25596	<i>Cracticus torquatus</i> (Grey Butcherbird)	
24322	<i>Cygnus atratus</i> (Black Swan)	

Name ID	Species Name	Status
30901	* Dacelo novaeguineae (Laughing Kookaburra)	
25673	Daphoenositta chrysoptera (Varied Sittella)	
24606	Daphoenositta chrysoptera subsp. pileata (Varied Sittella)	
25607	Dicaeum hirundinaceum (Mistletoebird)	
24470	Dromaius novaehollandiae (Emu)	
24290	Elanus caeruleus subsp. axillaris (Black-shouldered Kite)	
47937	Elseyornis melanops (Black-fronted Dotterel)	
24567	Epthianura albifrons (White-fronted Chat)	
24570	Epthianura tricolor (Crimson Chat)	
24379	Erythrogonys cinctus (Red-kneed Dotterel)	
25621	Falco berigora (Brown Falcon)	
25622	Falco cenchroides (Australian Kestrel)	
25623	Falco longipennis (Australian Hobby)	
25624	Falco peregrinus (Peregrine Falcon)	S
25727	Fulica atra (Eurasian Coot)	
24763	Gallinula tenebrosa subsp. tenebrosa (Dusky Moorhen)	
42314	Gavicalis virescens (Singing Honeyeater)	
24404	Geophaps plumifera (Spinifex Pigeon)	
25530	Gerygone fusca (Western Gerygone)	
47962	Glyciphila melanops (Tawny-crowned Honeyeater)	
24443	Grallina cyanoleuca (Magpie-lark)	
24295	Haliastur sphenurus (Whistling Kite)	
47965	Hieraaetus morphnoides (Little Eagle)	
25734	Himantopus himantopus (Black-winged Stilt)	
24491	Hirundo neoxena (Welcome Swallow)	
24511	Larus novaehollandiae subsp. novaehollandiae (Silver Gull)	
24557	Leipoa ocellata (Malleefowl)	T
25661	Lichmera indistincta (Brown Honeyeater)	
24582	Lichmera indistincta subsp. indistincta (Brown Honeyeater)	
24326	Malacorhynchus membranaceus (Pink-eared Duck)	
25651	Malurus lamberti (Variegated Fairy-wren)	
25652	Malurus leucopterus (White-winged Fairy-wren)	
25654	Malurus splendens (Splendid Fairy-wren)	
24583	Manorina flavigula (Yellow-throated Miner)	
25758	Megalurus gramineus (Little Grassbird)	
24736	Melopsittacus undulatus (Budgerigar)	
24598	Merops ornatus (Rainbow Bee-eater)	IA
25542	Milvus migrans (Black Kite)	
25610	Myiagra inquieta (Restless Flycatcher)	
24407	Ocyphaps lophotes (Crested Pigeon)	
24618	Oreoica gutturalis (Crested Bellbird)	
24328	Oxyura australis (Blue-billed Duck)	P4
25680	Pachycephala rufiventris (Rufous Whistler)	

Name ID	Species Name	Status
24624	Pachycephala rufiventris subsp. rufiventris (Rufous Whistler)	
	Pandion cristatus (Osprey)	
25682	Pardalotus striatus (Striated Pardalote)	
24648	Pelecanus conspicillatus (Australian Pelican)	
48061	Petrochelidon nigricans (Tree Martin)	
48066	Petroica boodang (Scarlet Robin)	
24659	Petroica goodenovii (Red-capped Robin)	
25698	Phalacrocorax melanoleucos (Little Pied Cormorant)	
24667	Phalacrocorax sulcirostris (Little Black Cormorant)	
24409	Phaps chalcoptera (Common Bronzewing)	
48071	Phylidonyris niger (White-cheeked Honeyeater)	
24596	Phylidonyris novaehollandiae (New Holland Honeyeater)	
24841	Platalea flavipes (Yellow-billed Spoonbill)	
24746	Platycercus icterotis subsp. xanthogenys (Western Rosella -inland)	P4
25721	Platycercus zonarius (Australian Ringneck)	
24843	Plegadis falcinellus (Glossy Ibis)	IA
24382	Pluvialis fulva (Pacific Golden Plover)	IA
24681	Poliocephalus poliocephalus (Hoary-headed Grebe)	
25722	Polytelis anthopeplus (Regent Parrot)	
25731	Porphyrio porphyrio (Purple Swamphen)	
24767	Porphyrio porphyrio subsp. bellus (Purple Swamphen)	
24776	Recurvirostra novaehollandiae (Red-necked Avocet)	
48096	Rhipidura albiscapa (Grey Fantail)	
25614	Rhipidura leucophrys (Willie Wagtail)	
25534	Sericornis frontalis (White-browed Scrubwren)	
30948	Smicronis brevirostris (Weebill)	
24329	Stictonetta naevosa (Freckled Duck)	
25590	* Streptopelia senegalensis (Laughing Turtle-Dove)	
25705	Tachybaptus novaehollandiae (Australasian Grebe)	
24331	Tadorna tadornoides (Australian Shelduck)	
48135	Thinornis rubricollis (Hooded Plover)	P4
24845	Threskiornis spinicollis (Straw-necked Ibis)	
25549	Todiramphus sanctus (Sacred Kingfisher)	
48141	Tribonyx ventralis (Black-tailed Native-hen)	
24806	Tringa glareola (Wood Sandpiper)	IA
24808	Tringa nebularia (Common Greenshank)	IA
24851	Turnix velox (Little Button-quail)	
24386	Vanellus tricolor (Banded Lapwing)	
25765	Zosterops lateralis (Silvereye)	

APPENDIX F

**CONSERVATION SIGNIFICANT BIRD RECORDS -
NATUREMAP**

NatureMap Species Report

Created By Andre Schmitz on 22/09/2017

Conservation Status Conservation Taxon (T, X, IA, S, P1-P5)
Current Names Only Yes
Core Datasets Only Yes
Species Group Birds
Method 'By Circle'
Centre 115° 40' 26" E, 30° 45' 16" S
Buffer 20km

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
1.	41324 <i>Ardea modesta</i> (great egret, white egret)		IA	
2.	24779 <i>Calidris acuminata</i> (Sharp-tailed Sandpiper)		IA	
3.	24784 <i>Calidris ferruginea</i> (Curlew Sandpiper)		T	
4.	24788 <i>Calidris ruficollis</i> (Red-necked Stint)		IA	
5.	24734 <i>Calyptorhynchus latirostris</i> (Carnaby's Cockatoo (short-billed black-cockatoo), Carnaby's Cockatoo)		T	
6.	25624 <i>Falco peregrinus</i> (Peregrine Falcon)		S	
7.	24557 <i>Leipoa ocellata</i> (Malleefowl)		T	
8.	24598 <i>Merops ornatus</i> (Rainbow Bee-eater)		IA	
9.	24328 <i>Oxyura australis</i> (Blue-billed Duck)		P4	
10.	24746 <i>Platycercus icterotis</i> subsp. <i>xanthogenys</i> (Western Rosella (inland))		P4	
11.	24843 <i>Plegadis falcinellus</i> (Glossy Ibis)		IA	
12.	24382 <i>Pluvialis fulva</i> (Pacific Golden Plover)		IA	
13.	48135 <i>Thinornis rubricollis</i> (Hooded Plover, Hooded Dotterel)		P4	
14.	24806 <i>Tringa glareola</i> (Wood Sandpiper)		IA	
15.	24808 <i>Tringa nebularia</i> (Common Greenshank, greenshank)		IA	

Conservation Codes

T - Rare or likely to become extinct
X - Presumed extinct
IA - Protected under international agreement
S - Other specially protected fauna
1 - Priority 1
2 - Priority 2
3 - Priority 3
4 - Priority 4
5 - Priority 5

¹ For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.

APPENDIX G STUDY AREA PLANT SPECIES LIST

STUDY AREA PLANT SPECIES LIST

Family	Taxon
Anarthriaceae	<i>Lyginia barbata</i>
Apiaceae	<i>Xanthosia huegelii</i>
Araceae	* <i>Zantedeschia aethiopica</i>
Asparagaceae	<i>Sowerbaea laxiflora</i>
Asteraceae	* <i>Arctotheca calendula</i>
Asteraceae	<i>Brachyscome</i> sp.
Asteraceae	<i>Hyalosperma cotula</i>
Asteraceae	* <i>Hypochaeris glabra</i>
Asteraceae	<i>Lagenophora huegelii</i>
Asteraceae	<i>Podotheca gnaphalioides</i>
Asteraceae	* <i>Ursinia anthemoides</i>
Caryophyllaceae	* <i>Petrorhagia dubia</i>
Casuarinaceae	<i>Allocasuarina humilis</i>
Colchicaceae	<i>Burchardia umbellata</i>
Cyperaceae	? <i>Schoenus</i> sp.
Cyperaceae	<i>Caustis dioica</i>
Cyperaceae	<i>Isolepis</i> sp.
Cyperaceae	<i>Lepidosperma longitudinale</i>
Cyperaceae	<i>Mesomelaena pseudostygia</i>
Cyperaceae	<i>Mesomelaena tetragona</i>
Cyperaceae	<i>Schoenus grandiflorus</i>
Dasygongonaceae	<i>Calectasia hispida</i>
Dennstaedtiaceae	<i>Pteridium esculentum</i>
Dilleniaceae	<i>Hibbertia huegelii</i>
Dilleniaceae	<i>Hibbertia hypericoides</i>
Dilleniaceae	<i>Hibbertia mylnei</i>
Dilleniaceae	<i>Hibbertia racemosa</i>
Droseraceae	<i>Drosera ?erythrorhiza</i> subsp. <i>magna</i>
Droseraceae	<i>Drosera subhirtella</i>
Elaeocarpaceae	<i>Tetratheca hirsuta</i>
Ericaceae	<i>Astroloma glaucescens</i>
Ericaceae	<i>Conostephium minus</i>
Ericaceae	<i>Conostephium ?pendulum</i>
Ericaceae	<i>Leucopogon ?polymorphus</i>
Ericaceae	<i>Leucopogon</i> sp.
Ericaceae	<i>Leucopogon sprengelioides</i>
Fabaceae	<i>Acacia pulchella</i>
Fabaceae	<i>Acacia saligna</i>
Fabaceae	<i>Bossiaea eriocarpa</i>
Fabaceae	<i>Daviesia decurrens</i>
Fabaceae	<i>Daviesia divaricata</i>
Fabaceae	<i>Gastrolobium nervosum</i>
Fabaceae	<i>Gompholobium knightianum</i>
Fabaceae	<i>Gompholobium tomentosum</i>

Family	Taxon
Fabaceae	<i>Jacksonia sternbergiana</i>
Fabaceae	* <i>Trifolium</i> sp.
Geraniaceae	* <i>Erodium cicutarium</i>
Goodeniaceae	<i>Leschenaultia biloba</i>
Haemodoraceae	<i>Anigozanthos humilis</i>
Haemodoraceae	<i>Conostylis candicans</i> subsp. <i>candicans</i>
Haemodoraceae	<i>Conostylis teretifolia</i> subsp. <i>planescens</i>
Hemerocallidaceae	<i>Johnsonia pubescens</i> subsp. <i>pubescens</i>
Iridaceae	* <i>Romulea rosea</i>
Juncaceae	* <i>Juncus acutus</i>
Juncaceae	<i>Juncus pallidus</i>
Loranthaceae	<i>Nuytsia floribunda</i>
Myrtaceae	? <i>Scholtzia</i> sp.
Myrtaceae	<i>Calothamnus quadrifidus</i>
Myrtaceae	<i>Corymbia calophylla</i>
Myrtaceae	<i>Eremaea asterocarpa</i>
Myrtaceae	<i>Eremaea pauciflora</i>
Myrtaceae	<i>Eucalyptus</i> ? <i>subangusta</i> subsp. <i>subangusta</i>
Myrtaceae	<i>Eucalyptus rudis</i>
Myrtaceae	<i>Eucalyptus todtiana</i>
Myrtaceae	<i>Eucalyptus wandoo</i>
Myrtaceae	<i>Hypocalymma tetrapterum</i> (P3)
Myrtaceae	<i>Hypocalymma xanthopetalum</i>
Myrtaceae	<i>Leptospermum erubescens</i>
Myrtaceae	<i>Melaleuca</i> ? <i>trichophylla</i>
Myrtaceae	<i>Melaleuca ciliosa</i>
Myrtaceae	<i>Melaleuca raphiophylla</i>
Orchidaceae	<i>Caladenia flava</i>
Orchidaceae	<i>Elythranthera brunonis</i>
Phyllanthaceae	<i>Phyllanthus calycinus</i>
Poaceae	* <i>Briza maxima</i>
Poaceae	* <i>Briza minor</i>
Poaceae	* <i>Ehrharta longiflora</i>
Poaceae	* <i>Eragrostis falcata</i>
Poaceae	* <i>Hordeum</i> sp.
Poaceae	* <i>Hordeum vulgare</i>
Poaceae	* <i>Lolium perenne</i>
Poaceae	<i>Neurachne alopecuroidea</i>
Poaceae	* <i>Vulpia bromoides</i>
Primulaceae	* <i>Lysimachia arvensis</i>
Proteaceae	<i>Adenanthos cygnorum</i>
Proteaceae	<i>Banksia attenuata</i>
Proteaceae	<i>Banksia hewardiana</i>
Proteaceae	<i>Banksia menziesii</i>
Proteaceae	<i>Banksia sessilis</i>
Proteaceae	<i>Banksia shuttleworthiana</i>

Family	Taxon
Proteaceae	<i>Conospermum stoechadis</i>
Proteaceae	<i>Grevillea ?vestita</i>
Proteaceae	<i>Hakea ?stenocarpa</i>
Proteaceae	<i>Hakea conchifolia</i>
Proteaceae	<i>Hakea flabellifolia</i>
Proteaceae	<i>Hakea incrassata</i>
Proteaceae	<i>Hakea lissocarpha</i>
Proteaceae	<i>Hakea trifurcata</i>
Proteaceae	<i>Hakea undulata</i>
Proteaceae	<i>Isopogon asper</i>
Proteaceae	<i>Lambertia multiflora</i>
Proteaceae	<i>Petrophile brevifolia</i>
Proteaceae	<i>Petrophile linearis</i>
Proteaceae	<i>Petrophile recurva</i>
Proteaceae	<i>Stirlingia latifolia</i>
Proteaceae	<i>Synaphea spinulosa</i>
Restionaceae	<i>Desmocladius fasciculatus</i>
Restionaceae	<i>Desmocladius flexuosus</i>
Restionaceae	<i>Hypolaena exsulca</i>
Restionaceae	<i>Lepidobolus preissianus</i>
Rutaceae	? <i>Boronia</i> sp.
Rutaceae	<i>Boronia ramosa</i>
Sapindaceae	<i>Diplopeltis huegelii</i>
Stylidiaceae	<i>Levenhookia stipitata</i>
Stylidiaceae	<i>Stylidium</i> sp.
Xanthorrhoeaceae	<i>Xanthorrhoea preissii</i>
Zamiaceae	<i>Macrozamia fraseri</i>

APPENDIX H FLORA SURVEY SITE DATA

Site C1

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	378338.1029 mE 6600397.269 mN
Landform	Valley (Minor Creek (< 5 m))
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Loam
Rock Type	
Vegetation Condition	Degraded
Disturbance Type	Weeds; Grazing; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus rudis</i>	Tree (10-30 m)	50
<i>Ehrharta longiflora</i>	Grass	5
<i>Zantedeschia aethiopica</i>	Herb	5
<i>Isolepis</i> sp.	Herb	1
<i>Juncus acutus</i>	Herb	30



Site C10a

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366425 mE 6596890 mN
Landform	Midslope
Slope & Aspect	Slope - Steep
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	No obvious disturbance
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1
<i>Acacia pulchella</i>	Shrub (0-1 m)	5
<i>Hakea lissocarpha</i>	Shrub (0-1 m)	10
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	10
<i>Leucopogon sprengeioides</i>	Shrub (0-1 m)	15
<i>Melaleuca ?trichophylla</i>	Shrub (0-1 m)	10
<i>Gastrolobium nervosum</i>	Shrub (0-1 m)	10
<i>Diplopeltis huegelii</i>	Shrub (0-1 m)	5



Site C10b

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	374600.5827 mE 6599163.918 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; Grazing; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus tottiana</i>	Mallee (3-10 m)	20
<i>Vulpia bromoides</i>	Grass	10
<i>Ehrharta longiflora</i>	Grass	5
<i>Arcotheca calendula</i>	Herb	5



Site C11a

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 366984 mE 6591765 mN
Landform Midslope
Slope & Aspect Slope - Gentle
Soil Colour Grey
Soil Texture Sand
Rock Type Latereite
Vegetation Condition Completely Degraded
Disturbance Type Grazing; Vegetation Clearing; Animal Tracks
Time since Fire > 5 Years
Leaf Litter Distribution and Cover Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	15
<i>Eucalyptus tottiana</i>	Tree (<10 m)	15



Site C11b

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 365333.118875 mE 6594959.755513 mN
Landform Midslope
Slope & Aspect Slope - Gentle
Soil Colour Brown
Soil Texture Sandy Loam
Rock Type Latereite
Vegetation Condition Completely Degraded
Disturbance Type Weeds; Vegetation Clearing
Time since Fire > 5 Years
Leaf Litter Distribution and Cover Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus tottiana</i>	Mallee (10-30 m)	5
<i>Hordeum vulgare</i>	Grass	70



Site C12a

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	361069.3507 mE 6594838.41 mN
Landform	Floodplain (Drainage Line)
Slope & Aspect	Slope - Negligible
Soil Colour	Orange
Soil Texture	Clay Loam
Rock Type	
Vegetation Condition	Good
Disturbance Type	Weeds; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus wandoo</i>	Tree (10-30 m)	40
<i>Acacia saligna</i>	Shrub (1-2 m)	1
<i>Grevillea ?vestita</i>	Shrub (1-2 m)	5
<i>Hibbertia mylnei</i>	Shrub (0-1 m)	1
<i>Briza minor</i>	Grass	5
<i>Eragrostis falcata</i>	Grass	30
<i>Desmocladius flexuosus</i>	Sedge	5



Site C12b

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366235.2475 mE 6596780.537 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	
Vegetation Condition	Degraded
Disturbance Type	Weeds; Vegetation Structure Altered; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	5
<i>Corymbia calophylla</i>	Mallee (3-10 m)	20
<i>Jacksonia sternbergiana</i>	Shrub (>2 m)	1
<i>Daviesia divaricata</i>	Shrub (1-2 m)	5
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	5
<i>Briza maxima</i>	Grass	1
<i>Neurachne alopecuroidea</i>	Grass	5
<i>Schoenus grandiflorus</i>	Sedge	5



Site C13

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	367301 mE 6596364 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Grey
Soil Texture	Loamy Sand
Rock Type	Other
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	30
<i>Banksia attenuata</i>	Tree (<10 m)	5
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1



Site C14

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368183.7627 mE 6596367.406 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Good
Disturbance Type	Weeds; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	15
<i>Banksia attenuata</i>	Tree (<10 m)	10
<i>Banksia menziesii</i>	Tree (<10 m)	10
<i>Acacia pulchella</i>	Shrub (0-1 m)	1
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	1
<i>Mesomelaena pseudostygia</i>	Sedge	5
<i>Schoenus grandiflorus</i>	Sedge	10



Site C15

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368197.8348 mE 6595510.8 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Good
Disturbance Type	Weeds; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	10
<i>Banksia menziesii</i>	Tree (<10 m)	10
<i>Nuytsia floribunda</i>	Tree (<10 m)	1
<i>Eucalyptus tottiana</i>	Mallee (3-10 m)	10
<i>Acacia pulchella</i>	Shrub (0-1 m)	1
<i>Neurachne alopecuroidea</i>	Grass	5
<i>Mesomelaena pseudostygia</i>	Sedge	10



Site C16

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366836 mE 6592107 mN
Landform	Midslope
Slope & Aspect	Slope - Moderate
Soil Colour	Grey
Soil Texture	Sand
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	Grazing; No obvious disturbance
Time since Fire	2-5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	15
<i>Hakea incrassata</i>	Shrub (0-1 m)	10
<i>Hakea ?stenocarpa</i>	Shrub (0-1 m)	5
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	40
<i>Isopogon asper</i>	Shrub (0-1 m)	5
<i>Lambertia multiflora</i>	Shrub (0-1 m)	5
<i>Leschenaultia biloba</i>	Shrub (0-1 m)	1
<i>Melaleuca ciliosa</i>	Shrub (0-1 m)	5
<i>Burchardia umbellata</i>	Herb	1



Site C17

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	367352 mE 6592737 mN
Landform	Ridgetop
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	Other; No obvious disturbance
Time since Fire	2-5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Banksia sessilis</i>	Shrub (>2 m)	10
<i>Banksia hewardiana</i>	Shrub (1-2 m)	30
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	5
<i>Daviesia decurrens</i>	Shrub (0-1 m)	1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	15
<i>Petrophile brevifolia</i>	Shrub (0-1 m)	5
<i>Lambertia multiflora</i>	Shrub (0-1 m)	1



Site C18

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	365849 mE 6591112 mN
Landform	Ridgetop
Slope & Aspect	Slope - Gentle
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	Vehicle Tracks; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	10
<i>Astroloma glaucescens</i>	Shrub (0-1 m)	20
<i>Allocasuarina humilis</i>	Shrub (0-1 m)	20
<i>Calothamnus quadrifidus</i>	Shrub (0-1 m)	15
<i>Calectasia hispida</i>	Shrub (0-1 m)	5
<i>Conospermum stoechadis</i>	Shrub (0-1 m)	5
<i>Banksia hewardiana</i>	Shrub (0-1 m)	15
<i>Banksia shuttleworthiana</i>	Shrub (0-1 m)	10
<i>Hakea conchifolia</i>	Shrub (0-1 m)	5
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	10
<i>Lambertia multiflora</i>	Shrub (0-1 m)	10
<i>Synaphea spinulosa</i>	Shrub (0-1 m)	10



Site C19

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366909 mE 6591597 mN
Landform	Ridgetop
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Good
Disturbance Type	Grazing; Vegetation Clearing
Time since Fire	2-5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus tottiana</i>	Tree (<10 m)	10
<i>Banksia hewardiana</i>	Shrub (>2 m)	60
<i>Acacia pulchella</i>	Shrub (0-1 m)	5
<i>Daviesia decurrens</i>	Shrub (0-1 m)	10
<i>Schoenus grandiflorus</i>	Sedge	20



Site C23

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	367149.3678 mE 6598014.941 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Very Good
Disturbance Type	Weeds; Grazing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Mallee (3-10 m)	5
<i>Banksia hewardiana</i>	Shrub (1-2 m)	20
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1
<i>Acacia pulchella</i>	Shrub (0-1 m)	1
<i>Daviesia decurrens</i>	Shrub (0-1 m)	1
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	1
<i>Schoenus grandiflorus</i>	Sedge	5



Site C24

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	367951.2133 mE 6591827.148 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; Vegetation Clearing; Grazing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus tottiana</i>	Mallee (3-10 m)	10
<i>Xanthorrhoea preissii</i>	Shrub (>2 m)	1
<i>Hordeum vulgare</i>	Grass	70
<i>Arctotheca calendula</i>	Herb	20



Site C25

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368356.3838 mE 6592175.233 mN
Landform	Midslope
Slope & Aspect	Slope - Moderate
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Good
Disturbance Type	Weeds; Grazing; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Mallee (3-10 m)	10
<i>Banksia hewardiana</i>	Shrub (>2 m)	10
<i>Acacia pulchella</i>	Shrub (1-2 m)	1
<i>Hibbertia mylnei</i>	Shrub (0-1 m)	5
<i>Briza maxima</i>	Grass	30
<i>Burchardia umbellata</i>	Herb	1
<i>Ursinia anthemoides</i>	Herb	1



Site C26

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	370372.2013 mE 6595927.555 mN
Landform	Footslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; Vegetation Structure Altered; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	50
<i>Nuytsia floribunda</i>	Tree (<10 m)	1
<i>Ehrharta longiflora</i>	Grass	30
<i>Arctotheca calendula</i>	Herb	1
<i>Hypochaeris glabra</i>	Herb	1
<i>Trifolium sp.</i>	Herb	5



Site C27

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	370714.8675 mE 6596219.73 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; Vegetation Structure Altered; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	60
<i>Ehrharta longiflora</i>	Grass	60
<i>Lolium perenne</i>	Grass	1
<i>Arctotheca calendula</i>	Herb	1
<i>Hypochaeris glabra</i>	Herb	1
<i>Romulea rosea</i>	Herb	1



Site C28

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	371847 mE 6597080 mN
Landform	Undulating Plain
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	30



Site C29

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	370123.2922 mE 6596763.455 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; Vegetation Structure Altered; Vegetation Clearing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	20
<i>Ehrharta longiflora</i>	Grass	50
<i>Arctotheca calendula</i>	Herb	1
<i>Hypochaeris glabra</i>	Herb	1
<i>Romulea rosea</i>	Herb	1



Site C2a

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 369100.1329 mE 6590680.485 mN
Landform Plain
Slope & Aspect Slope - Gentle
Soil Colour White
Soil Texture Sand
Rock Type
Vegetation Condition Excellent
Disturbance Type Weeds
Time since Fire
Leaf Litter Distribution and Cover

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus tottiana</i>	Mallee (3-10 m)	10
<i>Adenanthos cygnorum</i>	Shrub (>2 m)	5
<i>Allocasuarina humilis</i>	Shrub (>2 m)	5
<i>Hibbertia racemosa</i>	Shrub (1-2 m)	10
<i>Cautis dioica</i>	Sedge	5
<i>Mesomelaena tetragona</i>	Sedge	5
? <i>Schoenus sp.</i>	Sedge	5



Site C2b

Date	September 2017
Site Type	10 x 10 m
Location (GDA94 Zone 50)	378762.6069 mE 6600273.352 mN
Landform	Valley (Minor Creek (< 5 m))
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Loam
Rock Type	
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; Grazing; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Acacia saligna</i>	Tree (<10 m)	1
<i>Ehrharta longiflora</i>	Grass	5
<i>Arctotheca calendula</i>	Herb	1
<i>Hordeum vulgare</i>	Herb	1
<i>Juncus acutus</i>	Herb	80



Site C30

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	376857 mE 6600828 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Grey
Soil Texture	Sand
Rock Type	Other
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey; Grazing
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus tottiana</i>	Tree (<10 m)	40



Site C31

Date	September 2017
Site Type	10 x 10 m
Location (GDA94 Zone 50)	369303 mE 6592779 mN
Landform	Ridgetop
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sand
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	Vehicle Tracks; No obvious disturbance
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Allocasuarina humilis</i>	Shrub (>2 m)	10
<i>Banksia hewardiana</i>	Shrub (>2 m)	30
<i>Banksia sessilis</i>	Shrub (1-2 m)	5
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	10
<i>Lambertia multiflora</i>	Shrub (0-1 m)	10
<i>Leucopogon sprengeioides</i>	Shrub (0-1 m)	15



Site C32

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	369391 mE 6592524 mN
Landform	Midslope
Slope & Aspect	Slope - Steep
Soil Colour	Grey
Soil Texture	Sand
Rock Type	Granite
Vegetation Condition	Excellent
Disturbance Type	No obvious disturbance
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	15
<i>Banksia hewardiana</i>	Shrub (1-2 m)	40
<i>Hakea trifurcata</i>	Shrub (1-2 m)	5
<i>Macrozamia fraseri</i>	Shrub (1-2 m)	1
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1
<i>Hakea undulata</i>	Shrub (0-1 m)	5
<i>Leucopogon sprengeloides</i>	Shrub (0-1 m)	10
<i>Tetradlea hirsuta</i>	Shrub (0-1 m)	5



Site C34

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368076.1227 mE 6599318.261 mN
Landform	Ridgetop
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Clay Loam
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	Weeds
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Mallee (3-10 m)	5
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	5
<i>Calothamnus quadrifidus</i>	Shrub (0-1 m)	20
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Leucopogon ?polymorphus</i>	Shrub (0-1 m)	20



Site C35

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	377090 mE 6599475 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	40
<i>Caladenia flava</i>	Herb	< 1



Site C36

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368912.957 mE 6591581.081 mN
Landform	Midslope
Slope & Aspect	Slope - Moderate
Soil Colour	Brown
Soil Texture	Clay Loam
Rock Type	Latereite
Vegetation Condition	Good
Disturbance Type	Weeds; Vegetation Structure Altered; Grazing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Nuytsia floribunda</i>	Tree (<10 m)	5
<i>Allocasuarina humilis</i>	Shrub (>2 m)	5
<i>Banksia sessilis</i>	Shrub (>2 m)	20
<i>Banksia hewardiana</i>	Shrub (>2 m)	10
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	5
<i>Arctotheca calendula</i>	Herb	10
<i>Erodium cicutarium</i>	Herb	1
<i>Ursinia anthemoides</i>	Herb	1



Site C37

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	374867 mE 6597078 mN
Landform	Floodplain (Minor Creek (< 5 m))
Slope & Aspect	Slope - Moderate
Soil Colour	Grey
Soil Texture	Loamy Sand
Rock Type	Other
Vegetation Condition	Degraded
Disturbance Type	Weeds; No Native Understorey
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus rudis</i>	Tree (10-30 m)	30
<i>Melaleuca raphiophylla</i>	Tree (<10 m)	10
<i>Zantedeschia aethiopica</i>	Herb	80



Site C38

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 375591 mE 6599257 mN
Landform Midslope
Slope & Aspect Slope - Gentle
Soil Colour Grey
Soil Texture Loamy Sand
Rock Type Latereite
Vegetation Condition Completely Degraded
Disturbance Type Weeds; No Native Understorey
Time since Fire No Evidence
Leaf Litter Distribution and Cover High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	60



Site C39

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 371852.8296 mE 6598381.667 mN
Landform Midslope
Slope & Aspect Slope - Gentle
Soil Colour Brown
Soil Texture Sandy Loam
Rock Type Latereite
Vegetation Condition Completely Degraded
Disturbance Type Weeds; Grazing; Vegetation Clearing
Time since Fire > 5 Years
Leaf Litter Distribution and Cover High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Brassica tournefortii</i>	Herb	80



Site C3a

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	369765.1385 mE 6593375.173 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	
Vegetation Condition	Excellent
Disturbance Type	No obvious disturbance
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Nuytsia floribunda</i>	Tree (<10 m)	1
<i>Corymbia calophylla</i>	Mallee (10-30 m)	30
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	5
<i>Acacia pulchella</i>	Shrub (0-1 m)	1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Mesomelaena pseudostygia</i>	Sedge	1
<i>Desmodcladus flexuosus</i>	Herb	10



Site C3b

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	375554.7662 mE 6598633.753 mN
Landform	Valley (Minor Creek (< 5 m))
Slope & Aspect	Slope - Negligible
Soil Colour	Brown
Soil Texture	Loam
Rock Type	
Vegetation Condition	Degraded
Disturbance Type	Weeds; Grazing; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus rudis</i>	Tree (10-30 m)	70
<i>Acacia saligna</i>	Tree (<10 m)	5
<i>Melaleuca raphiophylla</i>	Tree (<10 m)	5
<i>Lepidosperma longitudinale</i>	Sedge	50
<i>Zantedeschia aethiopica</i>	Herb	10
<i>Pteridium esculentum</i>	Herb	20
<i>Juncus pallidus</i>	Herb	5



Site C4

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	373696 mE 6599311 mN
Landform	Undulating Plain
Slope & Aspect	Slope - Gentle
Soil Colour	Grey
Soil Texture	Sand
Rock Type	
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey; Grazing
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	20



Site C40

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368534 mE 6599940 mN
Landform	Ridgetop
Slope & Aspect	Slope - Steep
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Ironstone
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey; Grazing
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus sp.</i>	Mallee (3-10 m)	50



Site C41

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	369103 mE 6594970 mN
Landform	Undulating Plain
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sand
Rock Type	Other
Vegetation Condition	Good
Disturbance Type	Weeds; Vehicle Tracks
Time since Fire	2-5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	15
<i>Eucalyptus tottiana</i>	Tree (<10 m)	10
<i>Adenanthos cygnorum</i>	Shrub (1-2 m)	30
<i>Hibbertia racemosa</i>	Shrub (1-2 m)	5
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	1
<i>Stirlingia latifolia</i>	Shrub (0-1 m)	1
? <i>Boronia</i> sp.	Shrub (0-1 m)	1



Site C42

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	370119 mE 6594980 mN
Landform	Undulating Plain
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sand
Rock Type	Other
Vegetation Condition	Very Good
Disturbance Type	Vehicle Tracks
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Allocasuarina humilis</i>	Shrub (1-2 m)	20
<i>Banksia hewardiana</i>	Shrub (1-2 m)	15
<i>Astroloma glaucescens</i>	Shrub (0-1 m)	10
<i>Banksia shuttleworthiana</i>	Shrub (0-1 m)	1
<i>Hakea incrassata</i>	Shrub (0-1 m)	1
<i>Hakea trifurcata</i>	Shrub (0-1 m)	1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	10
<i>Leschenaultia biloba</i>	Shrub (0-1 m)	5
<i>Leucopogon sprengeloides</i>	Shrub (0-1 m)	5



Site C5a

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	365646.4148 mE 6597191.339 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Clay Loam
Rock Type	Latereite
Vegetation Condition	Degraded
Disturbance Type	Weeds; Vegetation Structure Altered; Vegetation Clearing
Time since Fire	2-5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Eucalyptus wandoo</i>	Tree (10-30 m)	30
<i>Lagenophora huegelii</i>	Herb	1
<i>Lysimachia arvensis</i>	Herb	1
<i>Ursinia anthemoides</i>	Herb	1



Site C5b

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 368118.232 mE 6599378.829 mN
Landform Ridgetop
Slope & Aspect Slope - Gentle
Soil Colour Brown
Soil Texture Clay Loam
Rock Type Latereite
Vegetation Condition Excellent
Disturbance Type Weeds
Time since Fire > 5 Years
Leaf Litter Distribution and Cover

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Mallee (3-10 m)	20
<i>Banksia hewardiana</i>	Shrub (>2 m)	20
<i>Calothamnus quadrifidus</i>	Shrub (1-2 m)	10
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	1
<i>Leucopogon ?polymorphus</i>	Shrub (0-1 m)	5



Site C5c

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368103 mE 6599565 mN
Landform	Ridgetop
Slope & Aspect	Slope - Negligible
Soil Colour	Grey
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Good
Disturbance Type	Weeds; Grazing
Time since Fire	1-2 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	40
<i>Macrozamia fraseri</i>	Shrub (1-2 m)	1
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	10
<i>Phyllanthus calycinus</i>	Shrub (0-1 m)	10



Site C6

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	368733.3156 mE 6599661.576 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	
Vegetation Condition	Very Good
Disturbance Type	Weeds
Time since Fire	2-5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Mallee (10-30 m)	50
<i>Macrozamia fraseri</i>	Shrub (1-2 m)	1
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	5
<i>Phyllanthus calycinus</i>	Shrub (0-1 m)	5
<i>Sowerbaea laxiflora</i>	Herb	1



Site C7

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	369017.6526 mE 6599282.258 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Brown
Soil Texture	Sandy Loam
Rock Type	Latereite
Vegetation Condition	Good
Disturbance Type	Weeds; Grazing; Vegetation Structure Altered
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	50
<i>Xanthorrhoea preissii</i>	Shrub (>2 m)	1
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	5
<i>Phyllanthus calycinus</i>	Shrub (0-1 m)	1
<i>Hordeum sp.</i>	Grass	5



Site C8

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366033.9693 mE 6596715.222 mN
Landform	Ridgetop
Slope & Aspect	Slope - Moderate
Soil Colour	Brown
Soil Texture	Clay Loam
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	Weeds
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	10
<i>Acacia pulchella</i>	Shrub (0-1 m)	5
<i>Hakea incrassata</i>	Shrub (0-1 m)	1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Isopogon asper</i>	Shrub (0-1 m)	1
<i>Leucopogon sprengeioides</i>	Shrub (0-1 m)	15
<i>Desmocladus flexuosus</i>	Herb	5



Site C98

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	369572.9238 mE 6593095.21 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Excellent
Disturbance Type	No obvious disturbance
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Mallee (3-10 m)	5
<i>Eucalyptus tottiana</i>	Mallee (3-10 m)	5
<i>Allocasuarina humilis</i>	Shrub (1-2 m)	5
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Leucopogon sprengeioides</i>	Shrub (0-1 m)	5
<i>Mesomelaena pseudostygia</i>	Sedge	1



Site C99

Date September 2017
Site Type Check Site
Location (GDA94 Zone 50) 361907.492 mE 6594848.232 mN
Landform Plain
Slope & Aspect Slope - Gentle
Soil Colour Orange
Soil Texture Clay Loam
Rock Type
Vegetation Condition Good
Disturbance Type Rehab; Weeds; Vegetation Clearing
Time since Fire > 5 Years
Leaf Litter Distribution and Cover Low (< 10%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Acacia saligna</i>	Shrub (1-2 m)	5
<i>Acacia pulchella</i>	Shrub (0-1 m)	20
<i>Hakea flabellifolia</i>	Shrub (0-1 m)	1
<i>Eragrostis falcata</i>	Grass	10



Site C9a

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366607 mE 6595939 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	Grey
Soil Texture	Loamy Sand
Rock Type	Latereite
Vegetation Condition	Completely Degraded
Disturbance Type	Weeds; No Native Understorey
Time since Fire	No Evidence
Leaf Litter Distribution and Cover	High (> 40 %)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	50



Site C9b

Date	September 2017
Site Type	Check Site
Location (GDA94 Zone 50)	366568 mE 6595959 mN
Landform	Midslope
Slope & Aspect	Slope - Steep
Soil Colour	Grey
Soil Texture	Loamy Sand
Rock Type	Latereite
Vegetation Condition	Excellent
Disturbance Type	No obvious disturbance
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)

Dominant Species

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	20
<i>Xanthorrhoea preissii</i>	Shrub (1-2 m)	15
<i>Acacia pulchella</i>	Shrub (0-1 m)	15
<i>Hakea lissocarpha</i>	Shrub (0-1 m)	5
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	15
<i>Leucopogon sprengeioides</i>	Shrub (0-1 m)	10



Site Q1

Date	September 2017
Site Type	10 x 10 m
Location (GDA94 Zone 50)	368106.9246 mE 6590001.358 mN
Landform	Plain
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Excellent
Disturbance Type	Weeds
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)



Stratum Table

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	5
<i>Banksia sessilis</i>	Tree (<10 m)	5
<i>Xanthorrhoea preissii</i>	Shrub (>2 m)	1
<i>Petrophile recurva</i>	Shrub (1-2 m)	1
<i>Melaleuca ciliosa</i>	Shrub (1-2 m)	< 1
? <i>Scholtzia</i> sp.	Shrub (1-2 m)	< 1
<i>Astroloma glaucescens</i>	Shrub (0-1 m)	< 1
<i>Banksia shuttleworthiana</i>	Shrub (0-1 m)	< 1
<i>Bossiaea eriocarpa</i>	Shrub (0-1 m)	< 1
<i>Conospermum stoechadis</i>	Shrub (0-1 m)	1
<i>Conostephium ?pendulum</i>	Shrub (0-1 m)	< 1
<i>Gompholobium tomentosum</i>	Shrub (0-1 m)	< 1
<i>Gompholobium knightianum</i>	Shrub (0-1 m)	< 1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Hibbertia huegelii</i>	Shrub (0-1 m)	< 1
<i>Hypocalymma xanthopetalum</i>	Shrub (0-1 m)	< 1
<i>Leptospermum erubescens</i>	Shrub (0-1 m)	< 1
<i>Leucopogon sprengeloides</i>	Shrub (0-1 m)	1

Taxon	Stratum	Cover (%)
<i>Leucopogon</i> sp.	Shrub (0-1 m)	< 1
<i>Petrophile brevifolia</i>	Shrub (0-1 m)	< 1
<i>Synaphea spinulosa</i>	Shrub (0-1 m)	< 1
<i>Neurachne alopecuroidea</i>	Grass	< 1
<i>Mesomelaena pseudostygia</i>	Sedge	1
<i>Anigozanthos humilis</i>	Herb	< 1
<i>Hyalosperma cotula</i>	Herb	< 1
<i>Boronia ramosa</i>	Herb	< 1
<i>Burchardia umbellata</i>	Herb	< 1
<i>Caladenia flava</i>	Herb	< 1
<i>Caustis dioica</i>	Herb	< 1
<i>Drosera subhirtella</i>	Herb	< 1
<i>Drosera ?erythrorhiza</i> subsp. <i>magna</i>	Herb	< 1
<i>Elythranthera brunonis</i>	Herb	< 1
<i>Levenhookia stipitata</i>	Herb	< 1
<i>Petrorhagia dubia</i>	Herb	< 1
<i>Podotheca gnaphalioides</i>	Herb	< 1
<i>Ursinia anthemoides</i>	Herb	< 1

Site Q33

Date	September 2017
Site Type	10 x 10 m
Location (GDA94 Zone 50)	369649.9081 mE 6593302.207 mN
Landform	Footslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Excellent
Disturbance Type	Weeds
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)



Stratum Table

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	10
<i>Banksia menziesii</i>	Tree (<10 m)	10
<i>Stirlingia latifolia</i>	Shrub (1-2 m)	1
<i>Bossiaea eriocarpa</i>	Shrub (0-1 m)	< 1
<i>Conostephium ?pendulum</i>	Shrub (0-1 m)	< 1
<i>Conostephium minus</i>	Shrub (0-1 m)	< 1
<i>Eremaea asterocarpa</i>	Shrub (0-1 m)	< 1
<i>Eremaea pauciflora</i>	Shrub (0-1 m)	< 1
<i>Gompholobium tomentosum</i>	Shrub (0-1 m)	< 1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	1
<i>Hypocalymma xanthopetalum</i>	Shrub (0-1 m)	< 1
<i>Leucopogon sprengeioides</i>	Shrub (0-1 m)	10
<i>Leucopogon sp.</i>	Shrub (0-1 m)	< 1
<i>Melaleuca ciliosa</i>	Shrub (0-1 m)	< 1
<i>Petrophile linearis</i>	Shrub (0-1 m)	< 1
<i>Xanthorrhoea preissii</i>	Shrub (0-1 m)	< 1
<i>Daviesia decurrens</i>	Sedge	< 1

Taxon	Stratum	Cover (%)
<i>Conostylis candicans subsp. candicans</i>	Herb	< 1
<i>Desmocladus flexuosus</i>	Herb	5
<i>Drosera subhirtella</i>	Herb	< 1
<i>Drosera ?erythrorhiza subsp. magna</i>	Herb	< 1
<i>Xanthosia huegelii</i>	Herb	< 1
<i>Johnsonia pubescens subsp. pubescens</i>	Herb	< 1
<i>Lepidobolus preissianus</i>	Herb	< 1
<i>Lyginia barbata</i>	Herb	< 1
<i>Hypolaena exsulca</i>	Herb	1
<i>Stylidium sp.</i>	Herb	< 1
<i>Ursinia anthemoides</i>	Herb	< 1

Site Q4

Date	September 2017
Site Type	10 x 10 m
Location (GDA94 Zone 50)	370718.0377 mE 6594856.118 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Excellent
Disturbance Type	Weeds
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Moderate (10-40%)



Stratum Table

Taxon	Stratum	Cover (%)
<i>Corymbia calophylla</i>	Tree (10-30 m)	10
<i>Banksia attenuata</i>	Tree (<10 m)	10
<i>Acacia pulchella</i>	Shrub (0-1 m)	< 1
<i>Allocasuarina humilis</i>	Shrub (0-1 m)	1
<i>Bossiaea eriocarpa</i>	Shrub (0-1 m)	< 1
<i>Daviesia decurrens</i>	Shrub (0-1 m)	< 1
<i>Drosera subhirtella</i>	Shrub (0-1 m)	< 1
<i>Gompholobium knightianum</i>	Shrub (0-1 m)	< 1
<i>Gompholobium tomentosum</i>	Shrub (0-1 m)	< 1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	5
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	< 1
<i>Melaleuca ciliosa</i>	Shrub (0-1 m)	< 1
? <i>Scholtzia</i> sp.	Shrub (0-1 m)	< 1
<i>Petrophile linearis</i>	Shrub (0-1 m)	< 1
<i>Xanthorrhoea preissii</i>	Shrub (0-1 m)	5
<i>Briza maxima</i>	Grass	< 1
<i>Mesomelaena pseudostygia</i>	Sedge	5

Taxon	Stratum	Cover (%)
<i>Lagenophora huegelii</i>	Herb	< 1
<i>Caladenia flava</i>	Herb	< 1
<i>Calectasia hispida</i>	Herb	< 1
<i>Conostylis teretifolia</i> subsp. <i>planescens</i>	Herb	< 1
<i>Conostylis candicans</i> subsp. <i>candicans</i>	Herb	< 1
<i>Desmocladius fasciculatus</i>	Herb	< 1
<i>Desmocladius flexuosus</i>	Herb	5
<i>Drosera ?erythrorhiza</i> subsp. <i>magna</i>	Herb	< 1
<i>Hypochaeris glabra</i>	Herb	< 1
<i>Lepidobolus preissianus</i>	Herb	5
<i>Lyginia barbata</i>	Herb	< 1
<i>Hypolaena exsulca</i>	Herb	1
<i>Ursinia anthemoides</i>	Herb	< 1

Site Q8

Date	September 2017
Site Type	10 x 10 m
Location (GDA94 Zone 50)	368989.9645 mE 6592092.794 mN
Landform	Midslope
Slope & Aspect	Slope - Gentle
Soil Colour	White
Soil Texture	Sand
Rock Type	
Vegetation Condition	Excellent
Disturbance Type	Weeds; Grazing
Time since Fire	> 5 Years
Leaf Litter Distribution and Cover	Low (< 10%)



Stratum Table

Taxon	Stratum	Cover (%)
<i>Banksia attenuata</i>	Tree (<10 m)	5
<i>Eucalyptus tottiana</i>	Tree (<10 m)	5
<i>Jacksonia sternbergiana</i>	Shrub (>2 m)	1
<i>Xanthorrhoea preissii</i>	Shrub (>2 m)	5
<i>Acacia pulchella</i>	Shrub (0-1 m)	1
<i>Banksia shuttleworthiana</i>	Shrub (0-1 m)	< 1
<i>Hibbertia hypericoides</i>	Shrub (0-1 m)	10
<i>Hibbertia racemosa</i>	Shrub (0-1 m)	1
<i>Leucopogon</i> sp.	Shrub (0-1 m)	< 1
<i>Melaleuca ciliosa</i>	Shrub (0-1 m)	< 1
<i>Petrophile linearis</i>	Shrub (0-1 m)	< 1
<i>Briza maxima</i>	Grass	< 1
<i>Caustis dioica</i>	Sedge	1
<i>Mesomelaena pseudostygia</i>	Sedge	< 1
<i>Anigozanthos humilis</i>	Herb	< 1
<i>Hyalosperma cotula</i>	Herb	< 1
<i>Brachyscome</i> sp.	Herb	< 1

Taxon	Stratum	Cover (%)
<i>Caladenia flava</i>	Herb	< 1
<i>Desmocladius flexuosus</i>	Herb	< 1
<i>Drosera subhirtella</i>	Herb	< 1
<i>Drosera ?erythrorhiza subsp. magna</i>	Herb	< 1
<i>Stylidium sp.</i>	Herb	< 1
<i>Ursinia anthemoides</i>	Herb	< 1
<i>Hyalosperma cotula</i>	Herb	< 1