



HAL
open science

A New Species of *Argophyllum* (Argophyllaceae) with Notes on the Species from New Caledonia and Nickel Hyperaccumulation

Yohan Pillon, Vanessa Hequet

► **To cite this version:**

Yohan Pillon, Vanessa Hequet. A New Species of *Argophyllum* (Argophyllaceae) with Notes on the Species from New Caledonia and Nickel Hyperaccumulation. *Plants*, 2021, 10, 10.3390/plants10040701 . hal-03238655

HAL Id: hal-03238655

<https://hal.inrae.fr/hal-03238655>

Submitted on 27 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Article

A New Species of *Argophyllum* (Argophyllaceae) with Notes on the Species from New Caledonia and Nickel Hyperaccumulation

Yohan Pillon ^{1,*} and Vanessa Hequet ²¹ LSTM, IRD, INRAE, CIRAD, Institut Agro, Univ. Montpellier, 34398 Montpellier, France² AMAP, IRD, Herbier NOU, 98848 Nouméa, New Caledonia; vanessa.hequet@ird.fr

* Correspondence: yohan.pillon@ird.fr

Abstract: The taxonomy of *Argophyllum* (Argophyllaceae) in New Caledonia is reviewed here. All names validly published in *Argophyllum* in this archipelago are discussed and lectotypified when necessary. A new species is described, *Argophyllum riparium* (The LSID for the name *Argophyllum riparium* is: 77216335-1) Pillon and Hequet sp. nov. *Argophyllum grunowii* and *A. ellipticum* are both species complexes in which several species previously recognized are included here as well. Seven species are recognized in New Caledonia: *A. brevipetalum*, *A. ellipticum*, *A. grunowii*, *A. montanum*, *A. nitidum*, *A. riparium* and *A. vernicosum*, all endemic. Leaf nickel content of *A. riparium* can exceed 1000 $\mu\text{g}\cdot\text{g}^{-1}$, which makes this species a nickel hyperaccumulator. Measurements with a handheld X-Ray Fluorescence (XRF) spectrometer confirmed that this was also the case for all other species from New Caledonia, except *A. nitidum*. An identification key of New Caledonian species is provided.

Keywords: island; metal hyperaccumulation; Pacific; serpentine; systematics; ultramafic



Citation: Pillon, Y.; Hequet, V. A New Species of *Argophyllum* (Argophyllaceae) with Notes on the Species from New Caledonia and Nickel Hyperaccumulation. *Plants* **2021**, *10*, 701. <https://doi.org/10.3390/plants10040701>

Academic Editor:
Celestino Quintela-Sabaris

Received: 25 February 2021
Accepted: 27 March 2021
Published: 5 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Argophyllum J.R.Forst. and G.Forst. [1] (p. 29) is a genus of shrubs and small trees that has traditionally been placed in Saxifragaceae, or sometimes in Escalloniaceae or Grossulariaceae. Molecular phylogenetic studies [2] have indicated affinities with the genus *Corokia* A. Cunn. [3] (p. 248) and the two genera form the family Argophyllaceae [4] in the order Asterales [5] (APG I and subsequent others). Argophyllaceae are closely related to two other small families [2,6,7]: Phellinaceae, with a single genus of ten species all endemic to New Caledonia [8–10], and Alseuosmiaceae, with four to five genera and ten species from eastern Australia, New Guinea, New Zealand and New Caledonia [11–13]. The exact relationships between Argophyllaceae, Phellinaceae and Alseuosmiaceae, which form the “APA clade” or Alseuosmiineae Shipunov [14] (p. 63), are still not clear [15].

The genus *Argophyllum* was described by Johann and Georg Forster, two German botanists who travelled with James Cook during his second voyage around the world. They collected material in New Caledonia that they subsequently described as *Argophyllum nitidum* J. R. Forst. and G. Forst. [1] (p. 29). A second species, *A. ellipticum* Labill. [16] (p. 39), was collected and described by Labillardière, a member of Bruni d’Entrecasteaux’s voyage in search of the La Pérouse expedition. The genus was last revised in its entirety by Zemmann [17], who recognized ten species in Australia and New Caledonia. Leaf cuticle attributed to *Argophyllum* have been described from the Miocene of New Zealand [18]. The species in Australia have recently been revised with a total of eleven endemic species [19,20]. Guillaumin [21] revised the New Caledonia species as part of his revision of New Caledonian “Saxifragaceae” and recognised seven species, to which he subsequently added three new species [22]. Here, we provide a taxonomic update of New Caledonian *Argophyllum* to complete work on the genera from New Caledonia previously included in “Saxifragaceae” [23,24].

High nickel content was previously reported in the leaves of New Caledonian *Argophyllum* [25,26]. However, because the taxonomy of the genus was not clear at that time, there was uncertainty regarding the number and the names of the species that can actually be considered as nickel hyperaccumulators [27,28]. Here, we provide new analyses of herbarium specimens [29] and field-collected material to update the list of nickel hyperaccumulators in the genus *Argophyllum*.

2. Materials and Methods

Measurements, shapes and colours of the different organs are based on the examination of herbarium material and field observations. All herbarium specimens of *Argophyllum* present at NOU and P were examined (for herbarium acronyms: [30]). Additional type material was viewed online.

Nickel content in herbarium specimens was measured with a handheld X-Ray Fluorescence (XRF) spectrometer [28,31]. For each species, we selected specimens to represent their morphological, geographical and ecological range. For a putative new species, one leaf from five individuals was collected in the field, dried, ground to powder, digested in HNO₃/H₂O₂ and analysed by Microwave Plasma-Atomic Emission Spectrometer (MP-AES) [32].

3. Results

Examination of herbarium specimens and observations in the field suggests that seven endemic species should be recognized in the genus. Some material resembling *A. vernicosum* Däniker [33] (p. 165) but with larger leaves, thicker twigs, sparser branching, sepals with a different shape, and a riparian ecology are placed in a new taxon described below, *A. riparium* Pillon and Hequet sp. nov.

Measurements with an XRF spectrometer indicate that the nickel content in dried leaves exceeds 1000 µg·g⁻¹ in at least some individuals of six species: *A. brevipetalum*, *A. ellipticum*, *A. grunowii*, *A. montanum*, *A. riparium* sp. nov. and *A. vernicosum* (Table 1). The range of values observed between these six species are similar. The median value for all species is below 1000 µg·g⁻¹ and the highest value, 6670 µg·g⁻¹, is observed in *A. ellipticum*. The only exception is *A. nitidum*, where values were consistently below the level of detection. Values observed with MP-AES on five individuals of *Argophyllum riparium* sp. nov. confirmed that it is a nickel hyperaccumulator: 1498, 1520, 1678, 2111 and 2214 µg·g⁻¹ nickel in dried leaves.

Table 1. Nickel content (µg·g⁻¹) measured from herbarium specimens with a handheld X-Ray Fluorescence (XRF) spectrometer. Values are in µg·g⁻¹. LOD: Level of Detection. See Appendix A for individual values.

Species	Number of Specimens	Min (Ni)	Max (Ni)	Mean (Ni)	Median (Ni)
<i>A. brevipetalum</i>	4	<LOD	1675	483	129
<i>A. ellipticum</i>	12	<LOD	6670	1215	371
<i>A. grunowii</i>	40	<LOD	5694	1027	538
<i>A. montanum</i>	10	<LOD	1110	316	89
<i>A. nitidum</i>	9	<LOD	<LOD	<LOD	<LOD
<i>A. riparium</i>	4	173	1667	708	497
<i>A. vernicosum</i>	10	116	1872	608	387

4. Discussion

Seven species of *Argophyllum*, all endemic, can be recognized in New Caledonia. Two of them are variable and appear as species complexes: *A. ellipticum* or *A. grunowii*. It is not clear if this complexity may be caused by, e.g., hybridization [34,35] or cryptic species [36,37].

The obscure taxonomy of the genus led to uncertainty and discrepancy regarding the number and identity of the nickel hyperaccumulating species in different publications: *A. grunowii* and *A. laxum* [25,26,38], *A. ellipticum* [28], *A. brevipetalum*, *A. grunowii*, *A. latifolium*, *A. laxum*, *A. montanum* and *A. vernicosum* [27]. Novel measurements associated with a new taxonomy indicate that the nickel content in dried leaves exceeds $1000 \mu\text{g}\cdot\text{g}^{-1}$ in at least some specimens of six species that can, therefore, be considered as nickel hyperaccumulators [39]: *A. brevipetalum*, *A. ellipticum*, *A. grunowii*, *A. montanum*, *A. riparium* sp. nov. and *A. vernicosum*. These six species are ultramafic obligates, except *A. ellipticum*, which can also occur on other substrates. The seventh species of the genus in New Caledonia, *A. nitidum*, never occurs on ultramafic substrates and consistently has nickel content below the level of detection. The genus *Argophyllum* is therefore similar to *Geissois* Labill. [16] (p. 50) (Cunoniaceae): Out of the 13 species endemic to New Caledonia, seven of the eight species occurring on ultramafic substrates are nickel hyperaccumulators [40].

Although six species of *Argophyllum* qualify as nickel hyperaccumulators, their nickel content do not reach the high values observed in some other New Caledonian species [27]. The genus could nevertheless be used for phytoremediation, phytoextraction or ecological restoration, but its multiplication and cultivation has apparently not yet been tested in New Caledonia [41]. The cultivation of the Australian species is considered easy [42].

5. Taxonomic Section

- ARGOPHYLLUM J. R. Forst. and G. Forst. [1] (p. 29). Type:—*Argophyllum nitidum* J. R. Forst. & G. Forst.

non *Argophilum* Blanco [43] (p. 186) = *Aglaiia* Lour. [44] (p. 173)

- Sectional names

1. Sect. *Argophyllum*

sect. *Brachycalyx* Zemann [17] (p. 271) nom. inval.

- 2. Sect. *Dolichocalyx* Zemann [17] (p. 271). Type [17] (p. 289):—*Argophyllum laxum* Schltr.

Zemann [17] did not provide a formal Latin diagnosis for either of the two sections. Section *Brachycalyx* included species from Australia and New Caledonia: *Argophyllum ellipticum*, *A. lejourdani* F.Muell. [45] (p. 33), *A. nitidum*, *A. nullumense* Baker [46] (p. 439) and *A. cryptophlebium* Zemann [17] (p. 283), whereas section *Dolichocalyx* included only New Caledonian species: *A. montanum*, *A. schlechterianum*, *A. grunowii*, *A. laxum* and *A. latifolium*. Because it includes the type of the genus, section *Brachycalyx* is not a valid name [47] (article 22.2). Zemann distinguished the two sections based on the length of the petals: calyx much shorter than the corolla in section *Brachycalyx*, calyx half as long but generally equal to the corolla in sect. *Dolichocalyx*. These sections are not maintained. They may not be reciprocally monophyletic groups, or this would imply that Australian and New Caledonian *Argophyllum* are not monophyletic groups. Additionally, calyx length is a labile character, i.e., it is variable within *A. ellipticum*.

- *Argophyllum brevipetalum* Guillaumin [22] (p. 25). Type:—New Caledonia, cours moyen de la Tontouta, le long des rives, ± 50 m, 14 December 1940, Virot 375 (holo-: P00537620!)

Argophyllum brevipetalum is very similar to *A. ellipticum*, with which it shares the combination of yellow petals and hairy reddish coriaceous leaves, with the entire margin and inflorescences covered with red hairs. However, it occurs at a much lower elevation and differs in its smaller proportions (smaller leaves, slender petiole and stem). Guillaumin [22] (p. 25) segregated *A. brevipetalum* from *A. ellipticum* because of its calyx that is as long as its corolla, but this character is variable within *A. ellipticum* and is therefore not a reliable character alone on which to separate them.

- *Argophyllum ellipticum* Labill. [16] (p. 39). Lectotype (designated here):—New Caledonia, *Labillardière s.n.* (P00537623!, iso-: P00537624!, P00537625!)

Argophyllum ellipticum var. *comptonii* Baker f. [48] (p. 299), syn. nov. Type:—New Caledonia, Tonine, 30 September 1914, *Compton* 1938 (holo-: BM001254561!)

Argophyllum ellipticum var. *ovatum* Pamp. [49] (p. 81), syn. nov. Type:—New Caledonia, Labillardière s.n. (holo-: FI018226!)

Argophyllum ellipticum var. *rigidum* Däniker [50] (p. 160), syn. nov. Type:—New Caledonia, auf des sud ost crête des Mt Humboldt, 7 November 1924, *Däniker* 547 (holo-Z000015794!, iso-: Z000015795!)

Argophyllum rufum Vieill. ex Zemann [17] (p. 282) nom. nud.

Argophyllum obovatum Brongn. & Gris ex Guillaumin [51] (p. 60) [52] (p. 134) nom. nud. *Argophyllum vernicosum* var. *obovatum* (Brongn. & Gris ex Guillaumin) Guillaumin [21] (p. 277) nom. nud.

Argophyllum ellipticum is characterized by the combination of yellow petals, sepals much shorter or as long as petals, hairy coriaceous leaves (generally red and obovate in the most typical forms) with the entire margin and inflorescences generally covered with red hairs. *Argophyllum ellipticum* is widespread across the main island of New Caledonia, at mid- to high elevation, generally on ultramafic substrates, but occasionally on other, mostly poor, soils. It is a variable taxon, more robust in its dimension than the rare low-elevation *A. brevipetalum*. Specimens collected in the northeast of the island on non-ultramafic substrates, including the type of the species and of var. *comptonii*, have particularly short sepals (three times shorter than petals) and inflorescences with a long basal peduncle ramifying very distally. Most material collected on ultramafic substrates from Mont Humboldt and Mont Kouakoué, including the type of *A. ellipticum* var. *rigidum*, have leaves that are often revolute and whitish underneath, slender sepals (almost as long as petals), and shorter inflorescence ramified nearer the base. In locations in between these, generally on ultramafic substrates, the material displays a range of intermediate characters, which make this taxon difficult to divide further.

The name *Argophyllum obovatum* was published without a description and the specimen cited by Guillaumin [51] (p. 60), *Balansa* 1814 (P00537626!, P00537627!) belongs to *A. ellipticum*. *Argophyllum rufum* has never been published with a description and was treated by Zemann [17] as a synonym of *A. ellipticum*.

- *Argophyllum grunowii* Zahlbr. [53] (p. 278). Type:—New Caledonia Thio, auf serpentenberg, September 1884, *Grunow* s.n. (holo-: W1887-0007474!, iso-: NSW923234!)

Argophyllum acinetochromum Guillaumin [22] (p. 24), **syn. nov.** Type:—New Caledonia, pentes sud du Mont Kaala, ± 500 m, 2 November 1943, *Viro*t 1320 (holo-: P00537619!; iso-: NOU008828!)

Argophyllum amoenum var. *ovatum* Vieill. ex Guillaumin [52] (p. 133) nom. nud.

Argophyllum brevistylum Guillaumin [22] (p. 25), **syn. nov.** Type:—New Caledonia, près du sommet sud du Mont Kaala, ± 1000 m, 2 November 1943, *Viro*t 1318 (holo-: P00537621!; iso-: A01154254!, P00537622!)

Argophyllum latifolium Vieill. ex Zemann [17] (p. 285), **syn. nov.** Lectotype (designated here):—New Caledonia, ad montes prope Wagap, 1861-1867, *Vieillard* 2199 (P00537630!, iso-: P00537631!, P00537632!). Remaining syntypes:—New Caledonia Kanala, *Deplanche* 61 (P00537634!, P00537635!, P00537636!)

Argophyllum laxum Schltr. [54] (p. 118), syn. nov. Lectotype (designated here):?New Caledonia, Auf den Bergen bei Paita, 400 m, 7 October 1902, *Schlechter* 14962 (P00537637!, iso-: BR699808!, E00346922!, HBG515583!, K000739390!, W1907-0011817!). Remaining syntypes:—New Caledonia, Auf den Bergen am Ngoye, 150 m, 30 November 1902, *Schlechter* 15149 (B100715954!, BR699807!, E00346923!, HBG515582!, P00537638!, K000739391!, W1907-0011659!)

Argophyllum laxum var. *subintegrifolium* Baker f. [48] (p. 300), syn. nov. Type:—New Caledonia, riv. Ngoye, 20 October 1914, *Compton* 2095 (holo-: BM000600291!)

Argophyllum schlechterianum var. *vestitum* Baker f. [48] (p. 299), syn. nov. Type:—New Caledonia, Taom, 2 December 1914, *Compton* 2298 (holo-: BM001254563!)

Argophyllum splendens Vieill. ex Guillaumin [52] (p. 134) nomen. nud.

Argophyllum grunowii is characterized by the combination of yellow petals, and leaves and inflorescences covered with greyish hairs. Plants from southwest New Caledonia (e.g., type of *A. laxum*) have large leaves and inflorescences much longer than the leaves. Plants from the eastern coast (e.g., type of *A. latifolium*) have large leaves but inflorescences shorter than the leaves. Plants from the northwest (e.g., the types of *A. acinetochromum* and *A. brevistylum*) have medium-sized leaves with a very dense indumentum underneath and inflorescences about the length of the leaves. In spite of these apparent geographical trends, there are many intermediate forms and variability found within populations. Although not entirely satisfactory, a broad concept is adopted here. *Argophyllum grunowii* is widespread across the main island of New Caledonia, at low to high elevation, only on ultramafic substrates.

The name *Argophyllum amoenum* (published as “*amœnum*”) apparently never appeared in print except in combination with var. *ovatum*. The name was published without a description and the single specimen cited, *Vieillard* 2639 (P06309317!, P06309318!, P06309319!, P06309320!, P06309321!, P06309322!) belongs to *A. grunowii*. The collection number for the type collection of *Argophyllum laxum* var. *subintegrifolium* is 2095 on the herbarium label but was printed as 2905 in Rendle et al. [48]. The name *A. splendens* was published without a description and was treated by Guillaumin [52] as a synonym of *A. laxum*. Zemann [17] cited two collections for *A. latifolium*, *Vieillard* 2199 and *Deplanche* 61; one sheet of *Vieillard* 2199 is chosen here as a lectotype. Schlechter [54] cited two of his own collections for *A. laxum*: *Schlechter* 14962 and 15149. The first one (from Païta) is chosen over the second one (from Ngoye), as it was collected in the area where most of the typical forms have been collected.

- *Argophyllum montanum* Schlechter [54] (p. 118). Lectotype (designated here):—New Caledonia, Auf den Bergen bei Yaouhé, 700 m, 15 October 1902, *Schlechter* 15032 (B100715955!; iso:-P00537639!, K000739388!)

Argophyllum schlechterianum Bonati & Petitmangin [55] (p. 650). Type:—New Caledonia, forêt du mont Dzumacs, 900 m, October 1906, *Franc* 566 (holo:- Z00054678!)

Argophyllum montanum is characterized by the combination of yellow petals, glabrous leaves and inflorescences covered by greyish hairs. *Argophyllum montanum* occurs in the southern part of New Caledonia, at low to mid elevation, only on ultramafic substrates.

The type specimen is unusual in having some toothed leaves. We follow Guillaumin [21] in treating *A. schlechterianum* as a synonym of *A. montanum*.

- *Argophyllum nitidum* J.R.Forst. and G.Forst. [1] (p. 30). *Argophyllum nitidum* var. *nitidum* J.R.Forst. & G.Forst. Lectotype (designated here): New Caledonia, *Forster s.n.* (BM000600294!, iso:- BM000600292!, BM000600293!, K000739389! B100296104!, BW04962010!, W0017375!). See Nicolson and Fosberg [56] for a complete list of putative islectotypes.

Argophyllum sericeum Poir. [57] (p. 449) nomen. nud.

This species here is considered as endemic to New Caledonia, although some of the Australian species have sometimes been treated as varieties of *A. nitidum*. It is the only species from New Caledonia with white petals. Its calyx is particularly short (<2 mm). Its leaves are more chartaceous with a finer silvery indumentum than other species. Its inflorescences are silvery too, sometimes tinted with red. *Argophyllum nitidum* is widespread across the main island of New Caledonia, from Mount Koghis to Balade, and it is most common in the northern part, on non-ultramafic substrates and at low- to mid-elevation.

- *Argophyllum vernicosum* Däniker [33] (p. 165). Type:—New Caledonia, im dichten hohen Gebüsch der Crêten in 600 m Meereshöhe am Mt Koghi, 1 February 1926, *Däniker* 2725 (holo:- Z000015797!, iso:- Z000015798!, Z000015799!)

Argophyllum ellipticum var. *oblongifolium* Brongn. and Gris ex Guillaumin [52] (p. 133). nom. nud.

Argophyllum vernicosum is characterized by the combination of yellow petals, short calyx, glabrous shiny leaves and inflorescences covered with reddish hairs. *Argophyllum*

vernicosum occurs in the southern part of New Caledonia, at mid- to high elevation, on ultramafic substrates.

The name *Argophyllum ellipticum* var. *oblongifolium* (originally “*oblongifolia*”) was published without a description; the single specimen cited, *Balansa 1816* (P0537640!), belongs to *A. vernicosum*, and we therefore follow Guillaumin [21] in considering *A. ellipticum* var. *oblongifolium* as a synonym of *A. vernicosum*.

- *Argophyllum riparium* Pillon and Hequet, sp. nov (Figure 1)

Diagnosis: A species most similar to *Argophyllum vernicosum* Däniker, from which it differs by its larger leaves (generally ≥ 10 cm long), thicker twigs (at least 3–4 mm in diameter) and sparse branching. Its sepals are narrowly attenuated, nearly as long as petals, rather than shortly acute and not more than half the size of petals as in *A. vernicosum*.

Type: New Caledonia, Forêt de la rivière bleue près d’un creek, 23 April 1970, *Veillon 2137* (holo-: P04445141!, iso-: NOU024016 !).

Description: Shrub 1–2 m tall. Stipules absent. Hairs on new growth brown or rusty. Branchlet terete, thick, 3–4 mm in diameter at the insertion of inflorescences, reddish-brown. Leaves alternate, discolorous, at the end of the twigs. Petiole 14–41 mm \times 2–2.5 mm. Leaf blade elliptic 9–18 cm \times 2.5–6.5 cm, base acute, apex acute or retuse, 5–8 secondary veins on either side of midrib, margin entire to slight wavy, sometimes with a few teeth, few hairs invisible to the naked eyes (binocular $\times 4$), glabrous and shiny on both sides. Inflorescence axillary, paniculate, densely reddish tomentose, 8–14 cm long, 15–60 flowers, primary axis 1/2–2/3 of the length of the inflorescence, secondary axis 10–50 mm long, branching spreading at 30–60° from adjacent branch. Bracts subtending primary ramifications of the inflorescence 4 \times 0.5 mm, persistent. Floral bracts 2–3 mm, (?1–)2 per flower. Flower apparently bisexual, actinomorphic. Pedicels 2 mm long, flowering hypanthium cupular, 2.5 mm diameter. Calyx lobes 2.5 mm long; petals 3 mm long, yellow, corolla appendage yellow, 1 mm long. Stamen antesealous, staminal filament 0.8 mm long, anthers 0.4 mm long. Style simple, 1.2 mm long, stigma swollen, entire, globose. Seed unknown.

Ecology and conservation: This species has been found in two locations (Figure 2). Most collections came from within the Rivière Bleue Provincial Park, where it grows near the rainforest edge on the banks of “rivière bleue”. All these collections may have been made from the same population. It appears in small, scattered groups of individuals, and its population size may be small. It was found recently in a second location in Plaine des lacs, growing along a stream close to the Kuebini river, 35 km apart from the type locality. Its ecosystem is undisturbed, and one of the two sites occurs within a protected area where the most likely disturbance might be floods, which can be severe during tropical storms. Its conservation status will be evaluated by the Plant Red List Authority–New Caledonia, and it could qualify as Vulnerable (VU D2).

Paratypes: New Caledonia, haute rivière bleue, 200m, 6 August 1951, *Baumann 15031* (P03609388!); *ibid.*, 5 August 1951, *Baumann 15034* (P03609386!); Rivière Bleue S 22°5′–22°7′, E 166°37′–166°44′, 150 m, 1965, *Bernardi 9317* (P06234104!); Parc Provincial de la Rivière Bleue, S 22°05′47.4″, E 166°38′16.4″, 200 m, 8 November 2018, *Pillon & Isnard 1480* (P, NOU); Plaine des lacs: Kuebini S 22°13′9.89″, E 166°57′15.24″, 350 m, 23 September 2020, *Hequet 4668* (NOU107206!); Parc de la rivière bleue, S 22°05′49″, E 166°38′16″, 200 m, 18 March 2020, *Hequet 4662* (NOU106963 !).




Figure 1. *Argophyllum riparium* Pillon and Hequet. (A) flowering branch, scale = 2 cm; (B) portion of an inflorescence, scale = 1 mm. Drawn from Hequet 4662 by Thierry Sanchiz.




Figure 2. Distribution of *Argophyllum riparium* Pillon and Hequet (squares). Grey areas indicate ultramafic rocks.

6. Identification Key to the Species of *Argophyllum* in New Caledonia

- | | |
|---|------------------------|
| 1. Leaf chartaceous, apex pointed, only primary and secondary veins distinct, leaf and inflorescences silky silvery (rarely tinted in red), petals white, calyx short (<2 mm), non-ultramafic substrates | <i>A. nitidum</i> |
| 1'. Leaf coriaceous, apex pointed or rounded, venation network distinct, leaves and inflorescences glabrous or covered with silver or red hairs, petals yellow, calyx short or as long as petals, generally on ultramafic substrate (occasionally on non-ultramafic but then on poor substrate for <i>A. ellipticum</i>) | 2 |
| 2. Inflorescences covered with red hairs, leaf margin generally entire | 3 |
| 2'. Inflorescences covered with white hairs, leaf margin entire to markedly toothed | 6 |
| 3. Leaf blade covered with red hairs underneath, occasionally white when mature | 4 |
| 3'. Leaf blade glabrous, shiny underneath | 5 |
| 4. Overall plant slender, twig slender (diameter 2.5–4 mm), leaf 4–10 cm long, petiole 1.5 mm in diameter, petals and sepals about the same length, low elevation in Tontouta valley (<300 m) | <i>A. brevipetalum</i> |
| 4'. Plant robust, twigs thicker (diameter 4–6 mm), leaf 5–15 cm long, petiole 2 mm in diameter, sepals much shorter or as long as petals, widespread, mid- to high elevation (400–) 800–1600 m | <i>A. ellipticum</i> |
| 5. Leaves small (mostly < 10 cm long), twig thin (c. 2 mm in diameter), sepals shortly acute, never more than half the size of petals, well-ramified shrub | <i>A. vernicosum</i> |
| 5'. Leaves large (mostly > 10 cm long), twig thicker (3–4 mm in diameter), sepals narrowly attenuate, nearly as long as petals, sparsely branched shrub | <i>A. riparium</i> |
| 6. Leaf glabrous, leaf margin entire, mostly south of Dzumacs-Yaté | <i>A. montanum</i> |
| 6'. Leaf blade covered with pale to red hairs, leaf margin entire to toothed, widespread and variable | <i>A. grunowii</i> |

Author Contributions: Conceptualization, Y.P.; validation, Y.P. and V.H.; writing—original draft preparation, Y.P.; writing—review and editing, Y.P. and V.H. All authors have read and agreed to the published version of the manuscript.

Funding: This work was partly funded by the X-TreM project of the CNRS MITI X-Life program (PI: Sylvain Merlot).

Acknowledgments: We thank the curators of herbarium NOU and P for providing access to their collections; Chiara Nepi (herbarium FI), Robert Vogt (herbarium B), Martin Callmender (G), Raneer Prakash (BM) for sending digital images of specimens; Cécile Nowak for help with bibliographical searches; and Thierry Sanchiz for the line drawing. Nickel measurements were performed with the help of Sylvain Merlot, Pierre Jouannais and Anaïs da Costa. Two anonymous reviewers provided insightful comments on an earlier version of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

List of herbarium specimens analysed with an XRF spectrometer with herbarium barcodes and Ni content. Specimens with value above 1000 $\mu\text{g}\cdot\text{g}^{-1}$ are underlined.

Argophyllum brevipetalum: P03609387 (157 $\mu\text{g}\cdot\text{g}^{-1}$), P03609438 (101), P04351395 (<LOD), P06234113 (1676); *A. ellipticum*: P00354794 (309), P03609379 (<LOD), P03609382 (2253), P03609437 (967), P03609439 (3223), P04284932 (608), P04445120 (118), P04445184 (6670), P04445187 (<LOD), P04445188 (<LOD), P06234110 (433), P06234114 (<LOD); *A. grunowii*: P00217129 (167), P00217183 (1534), P00354673 (1351), P00537619 (207), P00537622 (3879), P00537637 (246), P00537638 (960), P03609286 (41), P03609287 (1581), P03609305 (737), P03609307 (485), P03609314 (320), P03609351 (437), P03609383 (2649), P03609384 (1238), P03609385 (5694), P04284517 (373), P04284521 (796), P04284933 (347), P04284938 (<LOD), P04284939 (489), P04351389 (587), P04351394 (1280), P04351397 (2514), P04351401 (752), P04351404 (782), P04445122 (127), P04445128 (461), P04445152 (229), P04445157 (73), P04445160 (<LOD), P04445164 (896), P04445165 (221), P04445166 (2778), P04445172 (672), P04445175 (1775), P04445179 (213), P04445180 (178), P06234106 (133), P06234119 (3896); *A. montanum*:

P03609289 (1110), P03609293 (1076), P03609294 (34), P04284918 (103), P04284920 (<LOD), P04351398 (511), P04445124 (33), P04445145 (50), P04445158 (165), P05563751 (75); *A. nitidum*: P04284934 (<LOD), P04284935 (<LOD), P04351380 (<LOD), P04445131 (<LOD), P04445133 (<LOD), P04445173 (<LOD), P05563784 (<LOD), P05563794 (<LOD), P06234107 (<LOD); *A. riparium*: P03609386 (1667), P03609388 (591), P04445141 (402), P06234104 (173); *A. vernicosum*: P04284916 (132), P04284936 (402), P04351382 (345), P04445117 (1023), P04445162 (116), P05563809 (1872), P05563811 (217), P05563824 (387), P05563829 (977).

References

- Forster, J.R.; Forster, G. *Characteris Generum Plantarum*; White, B., Cadell, T., Elmsly, P., Eds.; Londini: London, UK, 1776.
- Kårehed, J.; Lundberg, J.; Bremer, B.; Bremer, K. Evolution of the Australasian Families Alseuosmiaceae, Argophyllaceae, and Phellinaceae. *Syst. Bot.* **1999**, *24*, 660–682. [[CrossRef](#)]
- Cunningham, A. XXXI.-Florae Insularum Novae Zelandiae Precursor; or a Specimen of the Botany of the Islands of New Zealand. *Ann. Nat. Hist.* **1839**, *3*, 244–250. [[CrossRef](#)]
- Kårehed, J. Argophyllaceae. In *The Families and Genera of Vascular Plants*; Kadereit, J.W., Jeffrey, C., Eds.; Springer: Berlin/Heidelberg, Germany, 2007; Volume 8, pp. 13–18.
- Angiosperm Phylogeny Group An Ordinal Classification for the Families of Flowering Plants. *Ann. Mo. Bot. Gard.* **1998**, *85*, 531–553. [[CrossRef](#)]
- Bremer, B.; Bremer, K.; Heidari, N.; Erixon, P.; Olmstead, R.G.; Anderberg, A.A.; Källersjö, M.; Barkhordarian, E. Phylogenetics of Asterids Based on 3 Coding and 3 Non-Coding Chloroplast DNA Markers and the Utility of Non-Coding DNA at Higher Taxonomic Levels. *Mol. Phylogenet. Evol.* **2002**, *24*, 274–301. [[CrossRef](#)]
- Tank, D.C.; Donoghue, M.J. Phylogeny and Phylogenetic Nomenclature of the Campanulidae Based on an Expanded Sample of Genes and Taxa. *Syst. Bot.* **2010**, *35*, 425–441. [[CrossRef](#)]
- Barriera, G. Novitates Neocaledonicae. VIII. Taxonomie et Nomenclature du Genre *Phelline* (Phellinaceae) avec la Description de la Nouvelle Espèce *Phelline Barrierei*. *Candollea* **2017**, *72*, 361–370. [[CrossRef](#)]
- Barriera, G. Phellinaceae. In *Flore de la Nouvelle-Calédonie*; Faune et Flore Tropicales; Publications scientifiques du Muséum; IRD: Paris, France; Marseille, France, 2020; Volume 27, pp. 234–295.
- Barriera, G.; Savolainen, V.; Spichiger, R. Phellinaceae. In *The Families and Genera of Vascular Plants*; Kadereit, J.W., Jeffrey, C., Eds.; Springer: Berlin/Heidelberg, Germany, 2007; Volume 8, pp. 608–610.
- Kårehed, J. Alseuosmiaceae. In *The Families and Genera of Vascular Plants*; Kadereit, J.W., Jeffrey, C., Eds.; Springer: Berlin/Heidelberg, Germany, 2007; Volume 8, pp. 7–12.
- Van Steenis, C.G.G.J. A Synopsis of Alseuosmiaceae in New Zealand, New Caledonia, Australia, and New Guinea. *Blumea* **1984**, *29*, 387–394.
- Shepherd, L.D.; de Lange, P.J.; Townsend, A.; Perrie, L.R. A Biological and Ecological Review of the Endemic New Zealand Genus *Alseuosmia* (Toropapa; Alseuosmiaceae). *N. Z. J. Bot.* **2020**, *58*, 2–18. [[CrossRef](#)]
- Shipunov, A.; Reveal, J.L. Validation of Subordinal Names for “Systema Angiospermarum”. *Phytotaxa* **2011**, *16*, 63–64. [[CrossRef](#)]
- Soltis, D.E.; Smith, S.A.; Cellinese, N.; Wurdack, K.J.; Tank, D.C.; Brockington, S.F.; Refulio-Rodriguez, N.W.J.B.; Moore, M.J.; Carlswald, B.S.; Bell, C.D.; et al. Angiosperm Phylogeny: 17 Genes, 640 Taxa. *Am. J. Bot.* **2011**, *98*, 704–730. [[CrossRef](#)]
- Labillardière, J.-J.H. *Sertum Austrocaledonicum*; Dominae Huzard: Paris, France, 1824.
- Zemann, M. Studien Zu Einer Monographie Der Gattung *Argophyllum* Forst. *Ann. Des K.K. Nat. Hofmuseums* **1907**, *22*, 270–292.
- Pole, M.S. Dispersed Leaf Cuticle from the Early Miocene of Southern New Zealand. *Palaeontol. Electron.* **2008**, *11*, 15A.
- Bean, A.R.; Forster, P.I. A Taxonomic Revision of *Argophyllum* J.R.Forst. & G.Forst. (Argophyllaceae) in Australia. *Austrobaileya* **2018**, *10*, 207–235.
- Forster, P.I. *Argophyllum Verae* (Saxifragaceae), a New Species from Northern Queensland. *Austrobaileya* **1990**, *3*, 173–176.
- Guillaumin, A. Matériaux pour la Flore de La Nouvelle-Calédonie. LV.—Révision des Saxifragacées. *Bull. De La Société Bot. De Fr.* **1939**, *86*, 275–278. [[CrossRef](#)]
- Guillaumin, A.; Viro, R. Contributions à la Flore de la Nouvelle-Calédonie CII.—Plantes Récoltées par M. R. Viro. *Mémoires Du Muséum Natl. D’histoire Nat. Ser. Bbotanique* **1953**, *4*, 1–82.
- Pillon, Y. Nomenclature and Typification in *Polyosma* (Escalloniaceae) from New Caledonia, with the Description of a New Species. *Phytotaxa* **2018**, *371*, 267–272. [[CrossRef](#)]
- Pillon, Y.; Hequet, V. Two New Species of *Quintinia* (Paracryphiaceae) with Notes on the Species from New Caledonia and Vanuatu. *Phytotaxa* **2019**, *397*, 45–54. [[CrossRef](#)]
- Jaffré, T. Composition Chimique Élémentaire des Tissus Foliaires des Espèces Végétales Colonisatrices des Anciennes Mines de Nickel en Nouvelle-Calédonie. *Cah. Orstomsérie Biol.* **1977**, *12*, 323–330.
- Jaffré, T. *Etude Écologique du Peuplement Végétal des Sols Dérivés de Roches Ultrabasiqes en Nouvelle-Calédonie*; Travaux et Documents de l’ORSTOM; ORSTOM: Paris, France, 1980.
- Gei, V.; Isnard, S.; Erskine, P.D.; Echevarria, G.; Fogliani, B.; Jaffré, T.; van der Ent, A. A Systematic Assessment of the Occurrence of Trace Element Hyperaccumulation in the Flora of New Caledonia. *Bot. J. Linn. Soc.* **2020**, *194*, 1–22. [[CrossRef](#)]

28. Jaffré, T.; Pillon, Y.; Thomine, S.; Merlot, S. The Metal Hyperaccumulators from New Caledonia Can Broaden Our Understanding of Nickel Accumulation in Plants. *Front. Plant Sci.* **2013**, *4*, 279. [[CrossRef](#)]
29. Van der Ent, A.; Echevarria, G.; Pollard, A.J.; Erskine, P.D. X-Ray Fluorescence Ionomics of Herbarium Collections. *Sci. Rep.* **2019**, *9*, 4746. [[CrossRef](#)]
30. Thiers, B. *Index Herbariorum: A Global Directory of Public Herbaria and Associated Staff*; New York Botanical Garden: New York, NY, USA, 2020.
31. Gei, V.; Erskine, P.D.; Harris, H.H.; Echevarria, G.; Mesjasz-Przybyłowicz, J.; Barnabas, A.D.; Przybyłowicz, W.J.; Kopittke, P.M.; van der Ent, A. Tools for the discovery of hyperaccumulator plant species and understanding their ecophysiology. In *Agromining: Farming for Metals*; Van der Ent, A., et al., Eds.; Springer International Publishing: Cham, Switzerland, 2018; pp. 117–133, ISBN 978-3-319-61898-2.
32. García de la Torre, V.S.; Majorel-Loulergue, C.; Rigail, G.J.; Alfonso-González, D.; Soubigou-Taconnat, L.; Pillon, Y.; Barreau, L.; Thomine, S.; Fogliani, B.; Burtet-Sarramegna, V.; et al. Wide Cross-species RNA-Seq Comparison Reveals Convergent Molecular Mechanisms Involved in Nickel Hyperaccumulation across Dicotyledons. *New Phytol.* **2021**, *229*, 994–1006. [[CrossRef](#)] [[PubMed](#)]
33. Däniker, A.U. Ergebnisse Der Reise von Dr. A.U. U. Däniker Nach Neu-Caledonien Und Den Loyalitäts-Inseln (1924/25). 2. Neue Phanerogamen von Neu-Caledonien Und Den Loyalitäts-Inseln. *Vierteljahrsschr. Der Nat. Ges. Zürich* **1931**, *76*, 160–170.
34. Pillon, Y.; Hopkins, H.C.F.; Bradford, J.C. Two New Species of *Cunonia* (Cunoniaceae) from New Caledonia. *Kew Bull.* **2008**, *63*, 419–431. [[CrossRef](#)]
35. Pillon, Y.; Munzinger, J.; Amir, H.; Hopkins, H.C.F.; Chase, M.W. Reticulate Evolution on a Mosaic of Soils: Diversification of the New Caledonian Endemic Genus *Codia* (Cunoniaceae). *Mol. Ecol.* **2009**, *18*, 2263–2275. [[CrossRef](#)] [[PubMed](#)]
36. Pillon, Y.; Hopkins, H.C.F.; Munzinger, J.; Amir, H.; Chase, M.W. Cryptic Species, Gene Recombination and Hybridization in the Genus *Spiraeanthemum* (Cunoniaceae) from New Caledonia. *Bot. J. Linn. Soc.* **2009**, *161*, 137–152. [[CrossRef](#)]
37. Swenson, U.; Munzinger, J.; Lowry, P.P.; Cronholm, B.; Nylander, S. Island Life—Classification, Speciation and Cryptic Species of *Pycnanandra* (Sapotaceae) in New Caledonia. *Bot. J. Linn. Soc.* **2015**, *179*, 57–77. [[CrossRef](#)]
38. Brooks, R.R. Geobotany and hyperaccumulators. In *Plants That Hyperaccumulate Heavy Metals. Their Role in Phytoremediation, Microbiology, Archaeology, Mineral Exploitation and Phytomining*; Brooks, R.R., Ed.; CAB International: Wallingford, UK, 1998; pp. 55–94.
39. Van der Ent, A.; Baker, A.J.M.; Reeves, R.D.; Pollard, A.J.; Schat, H. Hyperaccumulators of Metal and Metalloid Trace Elements: Facts and Fiction. *Plant Soil* **2013**, *362*, 319–334. [[CrossRef](#)]
40. Pillon, Y.; Hopkins, H.C.F.; Rigault, F.; Jaffré, T.; Stacy, E.A. Cryptic Adaptive Radiation in Tropical Forest Trees in New Caledonia. *New Phytol.* **2014**, *202*, 521–530. [[CrossRef](#)] [[PubMed](#)]
41. Gâteblé, G. *Flore Ornementale de Nouvelle-Calédonie*; Au vent des îles: Tahiti, French Polynesia, 2016; ISBN 978-2-36734-078-4.
42. Kupsch, K. The Silver Leaves of the Rainforest Understorey. *Assoc. Soc. Grow. Aust. Plants. Rainfor. Study Group Newsl.* **2005**, *61*, 6.
43. Blanco, M. *Flora de Filipinas*; Sto. Thomas, por Candido Lopez: Manila, Philippines, 1837.
44. De Loureiro, J. *Flora Cochinchinensis*; Ulyssipone: Lisbon, Portugal, 1790.
45. Von Mueller, F. *Fragmenta Phytographiae Australiae*; Melbourne Auctoritate: Melbourne, Australia, 1863; Volume 4.
46. Baker, R.T. Contributions to a Knowledge of the Flora of Australia. No II. *Proc. Linn. Soc. N. S. W.* **1899**, *24*, 437–447. [[CrossRef](#)]
47. *International Code of Nomenclature for Algae, Fungi, and Plants*; Turland, N.; Wiersema, J.; Barrie, F.; Greuter, W.; Hawksworth, D.; Herendeen, P.; Knapp, S.; Kusber, W.-H.; Li, D.-Z.; Marhold, K.; et al., Eds.; Regnum Vegetabile; Koeltz Botanical Books: Oberreifenberg, Germany, 2018; Volume 159, ISBN 978-3-946583-16-5.
48. Rendle, A.B.; Baker, E.G.; Moore, S.L.M. A Systematic Account of the Plants Collected in New Caledonia and the Isle of Pines by Prof. R. H. Compton, M.A., in 1914.-Part I. Flowering Plants (Angiosperms). *J. Linn. Soc. Lond. Bot.* **1921**, *45*, 245–417. [[CrossRef](#)]
49. Pampanini, R. Sassafragacee Dell’Erbario Webb. *Nuovo G. Bot. Ital.* **1904**, *11*, 79–82.
50. Däniker, A.U. Ergebnisse Der Reise von Dr. A. U. Däniker Nach Neu-Caledonien Und Den Loyalty-Inseln (1924/6). 4. Katalog Der Pteridophyta Und Embryophyta Siphonogama. II. *Vierteljahr. Der Nat. Ges. Zürich* **1932**, *19*, 116–235.
51. Guillaumin, A. Contribution à la Flore de Bourail (Nouvelle-Calédonie). *Ann. Du Musée Colonial De Marseille* **1911**, *9*, 55–75.
52. Guillaumin, A. Catalogue des Plantes Phanérogames de la Nouvelle-Calédonie et Dépendances (Île Des Pins et Loyalty). *Ann. Du Musée Colonial De Marseille* **1911**, *9*, 77–290.
53. Zahlbruckner, A. Beitrag Zur Flora von Neu-Caledonien, Enthaltend Die von A. Grunow Im Jahre 1884 Dasselbst Gesammelten Pflanzen. *Ann. Des. K.K. Nat. Hofmuseums* **1888**, *3*, 271–292.
54. Schlechter, R. Beiträge Zur Kenntnis der Flora von Neu-Kaledonien. *Bot. Jahrbücher Für Syst. Pflanz. Und Pflanz.* **1906**, *39*, 1–274.
55. Bonati, G.; Petitmengin, M. Sur Quelques Plantes de la Nouvelle-Calédonie. *Bull. De L’herbier Boissier* **1907**, *7*, 647–652.
56. Nicolson, D.H.; Fosberg, F.R. *The Forsters and the Botany of the Second Cook Expedition (1772–1775)*; Regnum vegetabile; Gantner: Ruggell, Liechtenstein, 2003; ISBN 978-3-906166-02-5.
57. Poiret, J.L.M. *Encyclopédie Méthodique. Botanique. Supplément, Tome I*; H. Agasse: Paris, France, 1810.