



**HAL**  
open science

**Molecular and morphological recognition of species boundaries in the neglected ant genus *Brachymyrmex* (Hymenoptera: Formicidae): toward a taxonomic revision**

Claudia M Ortiz-Sepulveda, Bert Van Bocxlaer, Andrés D Meneses, Fernando Fernandez

► **To cite this version:**

Claudia M Ortiz-Sepulveda, Bert Van Bocxlaer, Andrés D Meneses, Fernando Fernandez. Molecular and morphological recognition of species boundaries in the neglected ant genus *Brachymyrmex* (Hymenoptera: Formicidae): toward a taxonomic revision. *Organisms Diversity and Evolution*, 2019, 19 (3), pp.447-542. 10.1007/s13127-019-00406-2 . hal-02323851

**HAL Id: hal-02323851**

**<https://hal.science/hal-02323851>**

Submitted on 16 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Molecular and morphological recognition of species boundaries in the**  
2 **neglected ant genus *Brachymyrmex* (Hymenoptera: Formicidae): towards a**  
3 **taxonomic revision**

4  
5 Claudia M. Ortiz-Sepulveda<sup>1,2,\*</sup>, Bert Van Bocxlaer<sup>1</sup>, Andrés D. Meneses<sup>2</sup>, Fernando Fernández<sup>2</sup>

6  
7 <sup>1</sup>Université de Lille, CNRS, UMR 8198 - Evo-Eco-Paleo, F-59000 Lille, France.  
8 <sup>2</sup>Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Carrera 30 # 45 – 03, Bogotá,  
9 Colombia.

10 ORCID: CMOS: 0000-0003-0072-719X; BVB: 0000-0003-2033-326X.

11 \* Corresponding author. E-mail: claudia.ortiz-sepulveda@univ-lille.fr; Tel.: +33 3 20 43 40 15.

12  
13 **ABSTRACT**  
14 *Brachymyrmex* is a neglected genus of Formicinae because of its small body size, soft mesosoma  
15 and superficially monotonous external morphology. These features have complicated the  
16 documentation of morphological variation, resulting in poorly-defined and incompletely  
17 described species. Consequently, the taxonomy of the genus is complex and problematic, which  
18 has impeded research and conservation efforts. Here we integrate molecular and morphological  
19 data to recognize species boundaries in *Brachymyrmex* and to guide its long-overdue revision.  
20 Specifically, we (1) redefine the limits of all described species, subspecies and varieties based on  
21 intra- and interspecific morphological variation in workers; (2) document this variation  
22 quantitatively by constructing morphospace occupation and statistically analyzing measurements;  
23 (3) synthesize our findings on diagnostic traits in a dichotomous, illustrated identification key,

24 and (4) examine the significance of our morphological identification system with molecular  
25 evidence from four genes (EF1aEF1, EF1aEF2, WG and COI). We recognize 40 species, of  
26 which four are new to science: *B. bahamensis*, *B. bicolor*, *B. iridescens* and *B. sosai*.  
27 Furthermore, *Brachymyrmex attenuatus* and *B. bonariensis* are raised to species, and we propose  
28 25 new synonyms. Morphometrics indicated that even poorly distinguishable species pairs show  
29 statistically significant differences in some traits, and that taxonomically problematic cases relate  
30 to taxa that demonstrate large intraspecific trait variance. Our molecular analysis supports the  
31 monophyly of the genus based on increased taxon sampling, and of the 19 species that were  
32 included 18 were retrieved as monophyletic. The single case of incongruence was also flagged in  
33 morphological analyses and requires extended geographic sampling before it can be resolved. In  
34 conclusion, the molecular work corroborates the morphologically-recognized species boundaries.  
35 We also document the presence of worker dimorphism and putative worker-queen intercastes in  
36 several *Brachymyrmex* species, which indicates that the genus may present a promising study  
37 system to understand caste evolution in ants.

38

39 **Keywords.** *Brachymyrmex*, Formicinae, phylogeny, taxonomy, neotropics, morphometrics.

40

## 41 INTRODUCTION

42 *Brachymyrmex* is a neglected genus of Formicinae that consists of minute ants (maximum length  
43 ~3 mm), which are morphologically diagnosed by the presence of an acidopore and antennae  
44 with 9 segments lacking a club (Bolton 2003). The combination of their small body size, soft  
45 metasoma, and at least superficially monotonous external morphology complicate the observation  
46 and interpretation of morphological variation. *Brachymyrmex* is native to America and  
47 predominantly Neotropical. It ranges from the south of Canada to Argentina and Chile, including

48 the Caribbean islands (Kempf 1972; Brandão 1991; Bolton 1995; Bolton 2007). Creighton (1950)  
49 pointed out that these tiny ants are easily transported with living plants, and beyond the native  
50 distribution some species have been introduced to, among others, various places in Africa (Forel  
51 1895a; Dejean et al. 2010), Europe (Forel 1874), and Asia (Guénard 2018; Yoshimura pers.  
52 comm.). *Brachymyrmex heeri* and *B. longicornis*, for example, were described by Forel (1874;  
53 1907) from ant colonies in European greenhouses and *B. cordemoyi* was described from Réunion  
54 (Forel 1895a). Some *Brachymyrmex* species, like *B. patagonicus*, are notorious invaders which  
55 are considered pests in the southern United States (MacGown et al. 2007) and probably beyond.

56  
57 The only complete taxonomic treatment of *Brachymyrmex* was published by Santschi (1923a)  
58 and included 27 species and 15 subspecies and varieties. The work was based on worker  
59 morphology, but unfortunately the identification key is difficult to use because it includes  
60 polytomous steps with strongly overlapping character suites. Furthermore, character descriptions  
61 are regularly ambiguous and contain contradictions. As a result, species, subspecies and varieties  
62 are often poorly defined and incompletely described (De Zolessi et al. 1978). The small size and  
63 taxonomic ambiguity prompted Creighton (1950) to label *Brachymyrmex* as a “miserable little  
64 genus” in his treatment of the ants of North America, and for more than a century colleagues  
65 (Wheeler 1903; Kusnezov 1959; Wilson and Taylor 1967) have raised warnings on the  
66 taxonomic challenges in this genus. Since Santschi (1923a), Alayo (1974) examined the species  
67 from Cuba and Wheeler and Wheeler (1978) those from the United States. More recently, Quirán  
68 and collaborators (Quirán et al. 2004; Quirán 2005, 2007) reported on the *Brachymyrmex* species  
69 from Argentina, and Ortiz and Fernández (2014) reviewed the species with tumuliform  
70 metathoracic spiracles. Additionally, Wilson et al. (2016) documented the male genitalia of  
71 *Brachymyrmex*. Currently 44 species with 17 subspecies and varieties are attributed to the genus

72 in the online catalog of the ants of the world (Bolton 2018). However, the biology, diversity and  
73 phylogeny of the genus remain poorly understood and a comprehensive revision is long-overdue  
74 (see Wilson and Taylor 1967).

75  
76 A detailed account of opinions on the phylogenetic position of *Brachymyrmex* within Formicinae  
77 is provided by Wilson et al. (2016). Agosti (1991) divided the subfamily in four groups based on  
78 morphological characters, with *Brachymyrmex* included in the ‘*Pseudolasius* genus group’ based  
79 on the widely separated hind coxae, the petiole that is ventrally u-shaped and the simple helcium  
80 that is antero-ventrally often concealed by the anteriorly-fused sternite and tergite, which meet  
81 laterally. Bolton (2003), also based on morphology, assigned *Brachymyrmex* (and *Pseudolasius*)  
82 to the Plagiolepidini, which is one of three tribes of the lasiine group. More recently, Blaimer et  
83 al. (2015) obtained strong support for a sister group relationship between *Brachymyrmex* and  
84 *Myrmelachista* upon analysis of ultraconserved elements (UCEs), and these genera form a well-  
85 supported sister group to all other formicines. Therefore, Ward et al. (2016) resurrected the tribe  
86 Myrmelachistini (= Brachymyrmicini) for these two genera. This tribe is morphologically  
87 characterized by 9-10 antennal segments, five mandibular teeth, an anteriorly inclined petiole  
88 with a long posterior peduncle, and an anterior tergo-sternal fusion of the third abdominal  
89 segment.

90  
91 Here, we work towards a comprehensive revision of *Brachymyrmex* by 1) redefining the limits of  
92 all of the described species, subspecies and varieties in light of intra- and interspecific  
93 morphological variation in workers; 2) documenting this variation both qualitatively and  
94 quantitatively; 3) summarizing these findings on diagnostic traits with a new, dichotomous,  
95 illustrated identification key to increase reproducibility and to make the diversity of

96 *Brachymyrmex* more accessible for future research; and 4) examining the significance of our  
97 morphological identification system and the monophyly of the genus in light of molecular  
98 evidence. Finally, we also report on the biogeographical distribution of the recognized species  
99 and how our taxonomic framework compares with previous studies.

100

101 In summary, we recognize a total of 40 species, four of which are newly described here. We also  
102 synonymize 25 previously described species/subspecies and raise two former subspecies to  
103 species status. The proposed species delimitations follow a new, dichotomous identification key  
104 that is supported by quantitative morphological studies. More importantly, we tested our  
105 morphological identification system with molecular data for half of the recognized species and  
106 found strong congruence (18 of the 19 included species were retrieved as monophyletic),  
107 indicating its overall validity. During our studies, it also became clear that several samples  
108 contain specimens that presumably belong to undescribed species, but we prefer to await more  
109 material before formal description. This remark includes, but is not limited to, several potentially  
110 new species from Central America. We also observed that some species have dimorphic workers  
111 and others a possible intercaste between worker and queen. However, confirming the presence of  
112 an intercaste necessitates distinguishing it from ergatoid queens, which requires dissections of the  
113 ovaries and demographic data (Peeters 1991). Unfortunately, such confirmation is not usually  
114 possible based on the museum specimens studied here, but we discuss the issue where relevant.  
115 In general, it warrants further study and for now, we highlight such specimens as putative  
116 intercastes. If intercastes would be confirmed in the future beyond dimorphic workers,  
117 *Brachymyrmex* would present a promising study system to understand caste evolution in ants  
118 (Ortiz and Fernández 2014).

119

120 **MATERIAL AND METHODS**

121  
122 **Material and repositories.**  
123 Authors of previous taxonomic studies of *Brachymyrmex* (e.g. Santschi 1923a; Creighton 1950;  
124 Quirán et al. 2004; Quirán 2005, 2007) have mainly or exclusively focused on the morphology of  
125 workers, for which abundant material is available in existing museum collections. Consequently,  
126 we adopt the same focus here. A comparative framework is largely lacking for queens and males,  
127 because they are not available for all species, and even when collections exist they are often  
128 poorly preserved. Nevertheless, we provide a genus-level diagnosis of queens and males with  
129 selected pictures for illustrative purposes. The morphological terminology used follows Bolton  
130 (1994), that for hair inclination Kugler (1994), and for sculpture Harris (1979).

131 We examined a total of 1303 *Brachymyrmex* samples. This material belongs to the following  
132 institutions, and it includes all relevant types and many additional specimens; most collection  
133 acronyms follow Ward (1989). In some collections not all specimens have individual voucher  
134 numbers. In such cases, we assigned an identifier (either a personal code or number, such as  
135 CMOS 000032, or a Smithsonian database reference number, such as USNMENT00757197) to  
136 the relevant specimen, preceded by the acronym of the proprietary institution. These unique  
137 identifiers are used here for traceability.

138  
139 **ALWC** Alex L. Wild Personal Collection, University of Texas, Austin,, TX, USA.  
140 **CASC** California Academy of Sciences, San Francisco, CA, USA.  
141 **CPDC** Laboratório de Mirmecologia do Centro de Pesquisas do Cacau, Comissão  
142 Executiva do Plano da Lavoura Cacaueira (CEPLAC), Itabuna, Bahia, Brazil.  
143 **IAvH** Instituto Humboldt, Claustro San Agustín, Villa de Leyva, Boyacá, Colombia.

- 144 **ICN** Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá D.C.,  
145 Colombia.
- 146 **INBC** Instituto Nacional de Biodiversidad, Santo Domingo de Heredia, Costa Rica.
- 147 **INSUE** Instituto Superior de Entomología, Universidad Nacional de Tucumán, San Miguel  
148 de Tucumán, Argentina.
- 149 **JTLC** John Longino Collection, the University of Utah, Salt Lake City, UT, USA.
- 150 **MACN** Museo Argentino de Ciencias Naturales Bernardino Rivadavia, Buenos Aires,  
151 Argentina.
- 152 **MCZC** Museum of Comparative Zoology, Harvard University, Cambridge, MA, USA.
- 153 **MCSN** Museo Civico di Storia Naturale “Giacomo Doria”, Genoa, Italy.
- 154 **MfNB** Museum für Naturkunde, Berlin, Germany.
- 155 **MLP** Museo de La Plata, Buenos Aires, Argentina.
- 156 **MHNG** Muséum d’Histoire Naturelle, Genève, Switzerland.
- 157 **MPEG** Museu Paraense “Emílio Goeldi”. Belém, Pará, Brazil.
- 158 **MZSP** Museu de Zoologia da Universidade de São Paulo, São Paulo, Brazil.
- 159 **NHMB** Naturhistorisches Museum, Basel, Switzerland.
- 160 **NHMW** Naturhistorisches Museum, Wien, Austria.
- 161 **PSWC** Philip S. Ward Collection, University of California, Davis, CA USA.
- 162 **RBINS** Royal Belgium Institution of Natural Sciences, Bruxelles, Belgium.
- 163 **UFUC** Universidade Federal de Uberlândia, Uberlândia, Minas Gerais, Brazil.
- 164 **UNMSM** Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima,  
165 Peru.
- 166 **USNM** Department of Entomology, National Museum of Natural History Smithsonian  
167 Institution, Washington DC, USA.



168 **WEMC** William and Emma MacKay, Personal Collection, El Paso Texas, TX, USA.

169

### 170 **Georeferencing and mapping.**

171 Although we tried to georeference all studied samples, some were excluded because locality  
172 information was too ambiguous for georeferencing (e.g., when only a country name was  
173 available). Furthermore, specimens from the same collecting event were sometimes separated  
174 over replicate samples. After removing such ‘duplicates’, 747 georeferenced localities remained,  
175 of which 736 represented specimens from the native range. These were mapped in R v3.2.1. (R  
176 Core Team 2015) using the packages maps v3.0.1. (Brownrigg et al. 2015) and mapdata v2.2-5  
177 (Brownrigg 2015), and subsequently projected on the ETOPO1 global topographic map of  
178 Amante and Eakins (2009).

179

### 180 **Images.**

181 Photographs were taken in dorsal, lateral and full-face view. At the MCZC we used an imaging  
182 system that consisted of a Leica MZ16 stereomicroscope equipped with a Leica DCF 420 digital  
183 camera, software from Leica Application Suite 3.7 and Helicon Focus 5.1 for auto-montage; at  
184 the USNM the imaging system consisted of a Leica Z16APO stereomicroscope with a JVC KY-  
185 F75U digital camera mounted to the Leica motor-focus system. Composite images made with this  
186 system were assembled using Auto-Montage Pro Version 5.03.0018 BETA (Synoptics Ltd.); at  
187 the MZSP the imaging system consisted of a Leica M250c stereomicroscope and Auto-Montage  
188 Professional software LAS3.6.0. Some images were obtained from [www.antweb.org](http://www.antweb.org), which is  
189 specified in the figure captions. Images were processed with Adobe Photoshop CS5.

190

### 191 **Analysis of measurements and indices.**

192 Measurements were made using an Advanced Optical microscope, a Leica Z16 APO microscope,  
193 and a Zeiss StereoDiscovery V20 in combination with an ocular micrometer. All measurements  
194 were taken at 80-120× magnification and are reported in mm to an accuracy of 2 decimal places.  
195 Indices were calculated from these measurements following Ortiz and Fernández (2014) (Fig. 1).

196  
197 *Head Length<sub>1</sub>* (HL<sub>1</sub>): The maximum length of the head excluding the mandibles in full-face view.  
198 HL<sub>1</sub> is measured as the straight-line distance from the mid-point of the anterior margin of the  
199 clypeus to the mid-point of the posterior (= vertexal) margin of the head (for major workers the  
200 posterior mid-point is located at the middle of the virtual line between the posterior apices of the  
201 head).

202  
203 *Head Length<sub>2</sub>* (HL<sub>2</sub>): Distance from the posterior margin of the frontal triangle (see Bolton, 1994,  
204 p. 192) to the vertexal margin in full-face view.

205  
206 *Head Length<sub>3</sub>* (HL<sub>3</sub>): Measurement of the gena in lateral view; this measurement equals the  
207 distance from the anterior margin of the eye to the posterior edge of the clypeus, parallel to the  
208 longest axis of the eye.

209  
210 *Head Width* (HW): The maximum width of the head measured in full-face view. Eyes are  
211 included in the measurement if they project laterally from the head.

212  
213 *Scape Length* (SL): The maximum length of the scape, excluding the basal constriction just distal  
214 to the condylar bulb.

215

216 *Eye Length* (EL): Maximum diameter of the compound eye.

217

218 *Weber's Length* (WL): The diagonal length of the mesosoma in lateral view, i.e. from the  
219 anterior-most point of the pronotum to the posterior-most basal angle of the metapleuron (this  
220 measurement excludes the cervical neck of the pronotum).

221

222 *Pronotum Length* (PnL): The length along the midline between the anterior and posterior edges  
223 of the pronotum in dorsal view (this measurement excludes the cervical neck of the pronotum).

224

225 *Pronotum Width* (PnW): The maximum width of the pronotum in dorsal view.

226

227 *Mesonotum Length* (ML): The length between the anterior edge of the mesonotum and the  
228 mesometanotal suture in dorsal view.

229

230 *Mesonotum Width* (MW): The maximum width of the mesonotum in dorsal view.

231

232 *Cephalic Index* (CI):  $(HW/HL_1) \times 100$ .

233 *Scape Index<sub>1</sub>* (SI<sub>1</sub>):  $(SL/HW) \times 100$ .

234 *Scape Index<sub>2</sub>* (SI<sub>2</sub>):  $(SL/HL_2) \times 100$ .

235 *Ocular Index<sub>1</sub>* (OI<sub>1</sub>):  $(EL/HW) \times 100$ .

236 *Ocular Index<sub>2</sub>* (OI<sub>2</sub>):  $(HL_3/HL_1) \times 100$ .

237

238 *Ommatidia*: The number of facets in the compound eye along its maximal diameter.

239

240 In total 347 specimens of 38 species were measured. In some cases, it was not possible to reliably  
241 measure all features, e.g. because of the preservation of the specimen or the way it was mounted.  
242 The ranges of the obtained measurements are described in the systematic treatment, but we also  
243 performed a statistical analysis of morphometric variables.

244 First, we ordinated these data with non-metric multidimensional scaling (nmMDS) using  
245 functions of *vegan* v2.3-0 (Oksanen et al. 2015) and *MASS* v7.3-41 (Venables and Ripley 2002).  
246 As this rank-based method does not allow missing data, we selected only specimens for which all  
247 measurements were taken, i.e. a subset of 240 individuals for 38 species. We converted this  
248 dataset into a Euclidean distance matrix and ordinated it in two dimensions using 1,000 random  
249 starting configurations to find the solution with minimal stress without getting trapped in local  
250 minima. The resulting stress value obtained, i.e. the goodness-of-fit, was multiplied by 100 and  
251 evaluated using the criteria of Kruskal (1964) and Clarke (1993). We also examined how  
252 individual morphometric variables (i.e. the measurements, indices and counts) contributed to the  
253 morphospace occupation with the ‘*envfit*’ function of *vegan* using 1,000 permutations.

254  
255 Subsequently, we conducted statistical tests for the univariate morphometric variables on all  
256 species that were represented with at least 5 specimens, resulting in a subset of 286 specimens for  
257 20 species. (Specimens with missing data were allowed for these tests.) Given that the data of  
258 several species differed significantly from a normal distribution, we used non-parametric Dunn’s  
259 tests to test pairwise differences between species for each measurement and index. These tests  
260 were performed in R using functions of the package *dunn.test* v1.3.4. (Dinno 2017), and the  
261 resulting *p*-values were adjusted with a Benjamini-Hochberg correction, i.e. using the false  
262 discovery rate (Benjamini and Hochberg 1995). These results were represented with boxplots,  
263 featuring letters to indicate significance levels of comparisons.

264

265 **Molecular phylogenetics.**

266 We examined the monophyly of the genus with a dataset that has substantially enhanced taxon  
267 sampling compared to previous efforts (Brady et al. 2006; Moreau et al. 2006; Blaimer et al.  
268 2015), and we examined the molecular support of the here proposed morphological identification  
269 system. The specimens used for genetics are indicated in Supplementary material table S1, i.e. 82  
270 specimens covering 19 *Brachymyrmex* species and 6 specimens of 5 *Myrmelachista* species (the  
271 sister-genus of *Brachymyrmex* [Blaimer et al. 2015]). *Acanthoponera minor*, *Manica rubida* and  
272 *Rhytidoponera metallica* were used as outgroups.

273

274 DNA extraction, amplification, and sequencing were carried out at the Laboratories of Analytical  
275 Biology (LAB) of the Smithsonian National Museum of Natural History, Washington, DC.

276 Genomic DNA was extracted using the Qiagen DNEasy Tissue Kit. Fragments of four protein-  
277 coding genes were amplified, i.e. one fragment for each of the nuclear genes elongation factor 1-  
278 alpha paralog F1 (EF1 $\alpha$  F1), elongation factor 1-alpha paralog F2 (EF1 $\alpha$  F2) and wingless (wg),  
279 and two of the mitochondrial gene cytochrome oxidase subunit 1 (COI). Primer sequences used  
280 for polymerase chain reaction (PCR) amplification are those used by LaPolla et al. (2010). PCR  
281 products were sequenced on an ABI sequencer (ABI 377 or ABI 3100) using Big Dye Cycle  
282 Sequencing chemistry. Fragments were sequenced bidirectionally, and the resulting chromatograms  
283 were assembled and edited with SEQUENCHER v.4.8.

284

285 Furthermore, our dataset includes unpublished sequences that are available in GenBank by the  
286 International Barcode of Life Consortium. These sequences are provided without species  
287 identification, but they are linked to an image database, and we included specimens for which

288 unambiguous identification was possible based on the available images. Additional sequences  
289 with specimen images were kindly provided by David Donoso and John Longino, and we used  
290 the same criteria for inclusion as for GenBank sequences.

291  
292 Sequences for each gene fragment were aligned using MAFFT v.7 (Kato and Standley 2013)  
293 and results were visually inspected in MESQUITE v.2.10 (Maddison and Maddison 2017) to  
294 determine codon positions. We tested for substitutional saturation using DAMBE v.5.5.9 (Xia  
295 2013) but none of the gene fragments used were saturated. Models of sequence evolution were fit  
296 with PARTITION FINDER v.1.1.1 (Lanfear et al. 2012) to individual gene fragments accounting  
297 for potential differences between codon positions. The resulting model fit was examined with a  
298 corrected Akaike Information Criterion (AICc). Subsequently, the data for the individual  
299 fragments was concatenated into a total dataset with seven partitions (Supplementary material  
300 table S2) and phylogenetically analysed with maximum parsimony (MP), maximum likelihood  
301 (ML) and Bayesian inference (BI). MP analyses were performed in PAUP\* v.4.0b0.10 for  
302 Windows (Swofford 2002) with gaps treated as fifth state, 10 000 bootstrap replicates and tree-  
303 bisection-reconnection branch swapping. ML analyses were performed with the RAxML  
304 BlackBox (Stamatakis et al. 2008) with 100 replicates and the implemented GTR +  $\Gamma$  model,  
305 individually parameterized for each of the 7 partitions. BI analyses were executed in MrBayes  
306 v.3.2.6 (Ronquist et al. 2012) as implemented in CYPRES v.3.3 (Miller et al. 2010). Two  
307 independent Markov chain Monte Carlo (MCMC) runs were conducted for 20 million  
308 generations and sampled every 1,000 generations. Each run was distributed across four chains  
309 with a heating parameter of 0.2 and 25% of the samples were discard as burnin. Convergence  
310 between runs was examined using the sump command and by inspecting effective sample sizes  
311 for the parameters in TRACER v.1.6 (Rambaut et al. 2013). The maximum clade credibility tree

312 was visualized with FigTree v.1.4.0 (Rambaut 2012) and the bootstrap support for clades  
313 retrieved under MP and ML was added. Sequences are deposited in GenBank and accession  
314 numbers are indicated in Supplementary table S1.

315

### 316 **Automated species delimitation.**

317 We identified hypothetical species entities from sequence data with an automated procedure. Our  
318 specific aim was to evaluate the congruence of automated species delimitation and our  
319 morphological identification system, and thus to test the reliability of that identification system.  
320 Multiple such methods exist (Pons et al. 2006; Leliaert et al. 2014; Da Silva et al. 2018) and  
321 results may vary considerably among methods (Da Silva et al. 2018) related to the size of the  
322 dataset, the methodological procedures adopted, variation in underlying population genetic  
323 parameters and evolutionary processes. Many of the potentially influencing biological factors are  
324 poorly known for *Brachymyrmex*. We used the automatic barcode gap discovery method (ABGD;  
325 Puillandre et al. 2011), i.e. a fast single-locus method, on the barcoding fragment of COI, because  
326 of the exceptional suitability of this fragment for species identification as well as species  
327 delimitation and discovery in Metazoa (Hebert et al. 2003). The ABGD method is generally  
328 considered to be conservative as to the number of hypothetical species lineages it detects (Da  
329 Silva et al. 2018). We performed the analysis on the ABGD website  
330 (<http://www.wabi.snv.jussieu.fr/public/abgd/abgdweb.html>, accessed 8 October 2018) using default  
331 parameters except for relative gap width, which was set to 1.0.

332

## 333 **RESULTS AND DISCUSSION**

### 334 **Systematic treatment.**

335 *Brachymyrmex* Mayr

336 *Plagiolepis* (in part): Roger (1863: 162).

337 *Brachymyrmex* Mayr, 1868: 163. Type species: *B. patagonicus*, by monotypy.

338 *Brachymyrmex* subgenus *Brysha* Santschi, 1923a: 652.

339 *Brachymyrmex* senior synonym of *Brysha*: Smith (1979: 1424).

340 *Brachymyrmex*: Kempf (1972), Bolton (1995; 2003; 2018).

341  
342 **Diagnosis.** *Brachymyrmex* differs from most other formicine genera by having workers with nine  
343 antennal segments. Some species of *Myrmelachista* also have nine antennal segments, but these  
344 have a well-defined antennal club, whereas such a club is absent in *Brachymyrmex*. Some  
345 *Agraulomyrmex* species from Africa also have 9 antennal segments without club (unpublished  
346 results), but *Brachymyrmex* differs from these by the presence of a mesometanotal suture.  
347 Workers are monomorphic to dimorphic; some species have a putative worker-queen intercaste.

348  
349 **Worker. Head.** Usually longer than wide, cordate in some species, with sparse to dense  
350 pubescence and hairs in variable orientation (appressed, decumbent, erect). Mandibles with five  
351 teeth, of which intercalar (central) and basal teeth are smaller than the others and the apical tooth  
352 is largest. Maxillary palps and labial palps with 6 and 4 segments, respectively. Maxillary palps  
353 usually reach the occiput and bear several long ventral hairs. The clypeus has a rounded anterior  
354 margin or in some taxa, notably *B. nebulosus*, its medio-anterior portion forms a “lip”. In  
355 monomorphic species the clypeus bears five long, erect hairs of which one usually conspicuous  
356 hair is near the anterior margin, two are in mediolateral position and the other two close to the  
357 toruli; other hairs are markedly shorter and appressed or decumbent. In dimorphic species the  
358 clypeus is larger and with a row of long thick hairs near the anterior margin. Toruli either touch



359 the posterior clypeal margin in oblique anterodorsal view or surpass it. Compound eye  
360 conspicuous, positioned usually on the cephalic midline or anterior to it; with 3-14 ommatidia  
361 along its maximal diameter. Number of ocelli either 0, 1 or 3, but when present often  
362 inconspicuous. Antennae with 9 segments, without antennal club; flagelomeres sometimes  
363 gradually increasing apically in diameter; scapes variable in length, with appressed, decumbent or  
364 erect hairs.

365 **Mesosoma.** With sparse or dense pubescence and hairs in variable orientation. The pronotum and  
366 mesonotum typically bear two erect hairs each, but sometimes additional suberect hairs on one or  
367 both are present, or erect hairs may be absent from the mesonotum. The pronotum is slightly to  
368 strongly convex, and the promesonotal suture, i.e. the line of junction between the pronotum and  
369 mesonotum, is always present. The mesonotum may bulge dorsally above the propodeum, and  
370 the mesometanotal suture, i.e. the line of junction between the mesonotum and the metanotum, is  
371 usually conspicuous, although the mesonotum and metanotum appear fused in some species. The  
372 metanotum is reduced to a transverse groove, the metanotal groove, which separates the  
373 mesonotum from the propodeum on the mesosomal dorsum. The metanotal groove is variable,  
374 from absent to wide and deep. The metathoracic spiracles are dorsal near the midline or  
375 dorsolateral, and not, slightly or very strongly protruding, i.e. tumiliform. The propodeal suture,  
376 i.e. the line of junction between either the mesonotum (if the metanotal groove is absent) or the  
377 metanotal groove anteriorly and the propodeum posteriorly, is present as a dorsal fold with  
378 variable lateral extension. Dorsum of the propodeum flat or convex and usually shorter than the  
379 propodeal slope. Propodeal spiracles circular and positioned near the posterior propodeal margin.  
380 Petiole usually with a low scale, reduced to a narrow subcylindrical segment that is overhung  
381 from behind by the gaster, but in dimorphic species the scale of the petiole may be high and  
382 visible in dorsal view. Hairs on the legs may be appressed, decumbent or erect.

383 **Gaster.** Of variable size, with five segments that bear sparse or dense pubescence and usually  
384 erect hairs, mainly but not exclusively along the posterior edges of the segments.

385 **Color and sculpture.** Body color ranges from light yellow to dark brown and black; most often it  
386 is uniform, but some species display markedly contrasting patterns, e.g. with the head and/or the  
387 gaster darker than the rest of the body. Body usually smooth and shiny, but in some species the  
388 head and/or mesonotum bear microsculpture.

389

390 **Queen (Fig. 2)** Head wider than long, with abundant, fine pubescence, and with long erect hairs;  
391 eyes large, located laterally along the cephalic midline; 3 ocelli present; frontal lobes well-  
392 developed; scapes usually extending beyond the posterior margin of the head; palpal formula:  
393 6,4. Mesosoma with moderately dense, fine pubescence and several erect hairs; anepisternum and  
394 katepisternum separated by a distinct suture. Anterior wing with a single dark brown cell, i.e.  
395 pterostigma, the first submarginal cell is closed, others open. Posterior wing with five to seven  
396 hamuli. Gaster with moderately dense, fine pubescence, and erect hairs along the posterior  
397 edges of the segments. Body color ranges from yellow to dark brown, and it is uniform or  
398 sometimes with the head and/or gaster darker than the rest of the body.

399

400 **Male (Fig. 3).** Head wider than long, with fine, sparse pubescence, lacking erect hairs except on  
401 mouthparts, and with smooth, shiny integument; maxillary palps with four segments, labial palps  
402 with two; mandibles unidentate; frontal lobes reduced; ocelli and eyes well-developed; antennae  
403 with 10 segments. Mesosoma with sparse pubescence and shiny integument, without erect hairs.  
404 Gaster shiny, lacking pubescence, with scattered erect hairs on the last few segments. Head dark  
405 brown to almost black, rest of body, including appendages, very light brown or concolorous.  
406 Wilson et al. (2016) described the morphology of the male genitalia in detail.

407

408 **Distribution.** Neotropical and Nearctic, with introductions elsewhere. The native distribution of  
409 *Brachymyrmex* is illustrated in Fig. 4.

410

411 **Biology.** *Brachymyrmex* is commonly collected from leaf-litter and some species occur in  
412 association with epiphytes; nests are found under stones, among plant roots, in trees, in rotten  
413 wood (Wheeler 1942; LaPolla and Longino 2006), and in urban buildings (MacGown et al.  
414 2007). The biology and natural history of the genus are poorly known although habitat  
415 information exists for some species, such as the arboreal *B. nebulosus* (LaPolla and Longino  
416 2006). As mentioned, some *Brachymyrmex* species are notorious invaders which are considered  
417 pests (MacGown et al. 2007).

418 Interestingly, *Brachymyrmex* species occur sometimes in association with other insects. Santschi  
419 (1923a) mentioned associations of *Brachymyrmex depilis*, *B. giardi* and *B. heeri* with mealybugs  
420 (Hemiptera: Coccidae) and observed that some species live in or very close to termite nests (*B.*  
421 *fiebrigi*, *B. modestus*, *B. myops*, *B. termitophilus*). Moretti et al. (2011), suggested a possible  
422 association between *B. cordemoyi* and the cockroach *Pycnoscelus surinamensis* (Blaberoidea:  
423 Blaberidae), whereas Delssine (pers. comm.) found a staphylinid beetle in a nest of *B. modestus*  
424 in Ecuador.

425

426 **Synonymy of species.**

427 ***B. admotus* Mayr, 1887**

428 = *B. longicornis* var. *immunis* Forel, 1908 **n. syn.**

429 ***B. antennatus* Santschi, 1929**

430 ***B. aphidicola* Forel, 1909**

- 431 = *B. heeri* var. *fallax* Santschi, 1923a
- 432 = *B. longicornis* var. *hemiops* Santschi, 1923a **n. syn.**
- 433 ***B. attenuatus* Santschi, 1929 n. st.**
- 434 ***B. australis* Forel, 1901**
- 435 = *B. australis* var. *curta* Santschi, 1922 **n. syn.**
- 436 = *B. longicornis* Forel, 1907 **n. syn.**
- 437 ***B. bahamensis* n. sp.**
- 438 ***B. bicolor* n. sp.**
- 439 ***B. bonariensis* Santschi, 1933 n. st.**
- 440 ***B. brasiliensis* Ortiz & Fernández, 2014**
- 441 ***B. bruchi* Forel, 1912a**
- 442 = *B. bruchi* var. *rufipes* Forel, 1912a
- 443 = *B. giardi* var. *nitida* Santschi, 1922 **n. syn.**
- 444 = *B. laevis* var. *andina* Santschi, 1923a **n. syn.**
- 445 ***B. cavernicola* Wheeler, 1938**
- 446 ***B. coactus* Mayr, 1887**
- 447 = *B. coactus* var. *nictitans* Emery, 1906 **n. syn.**
- 448 = *B. constrictus* Santschi, 1923a **n. syn.**
- 449 = *B. coactus* var. *robustus* Santschi, 1923b **n. syn.**
- 450 ***B. cordemoyi* Forel, 1895a**
- 451 = *B. laevis* var. *fuscula* Emery, 1906 **n. syn.**
- 452 = *B. brevicornis* Emery, 1906 **n. syn.**
- 453 = *B. patagonicus* var. *brevicornoeides* Forel, 1914 **n. syn.**
- 454 = *B. cordemoyi* var. *nigricans* Santschi, 1916

- 455 = *B. cordemoyi* var. *distinctus* Santschi, 1923a **n. syn.**
- 456 ***B. degener* Emery, 1906**
- 457 = *B. admotus* r. *niger* Forel, 1912a **n. syn.**
- 458 = *B. incisus* Forel, 1912a **n. syn.**
- 459 = *B. luederwaldti* Santschi, 1923a **n. syn.**
- 460 ***B. delabiei* Ortiz & Fernández, 2014**
- 461 ***B. depilis* Emery, 1893**
- 462 = *B. depilis* subsp. *flavescens* Grundmann, 1952
- 463 = *B. nanellus* Wheeler, 1903.
- 464 ***B. donisthorpei* Santschi, 1939**
- 465 ***B. feitosai* Ortiz & Fernández, 2014**
- 466 ***B. fiebrigi* Forel, 1908**
- 467 = *B. fiebrigi* var. *funicularis* Santschi, 1922 **n. syn.**
- 468 = *B. fiebrigi* var. *fumida* Santschi, 1923a **n. syn.**
- 469 ***B. flavidulus* Roger, 1863**
- 470 ***B. gagates* Wheeler, 1934**
- 471 ***B. gaucho* Santschi, 1917**
- 472 ***B. giardi* Emery, 1895**
- 473 = *B. melensis* De Zolessi, Abenante & Gonzalez, 1978 **n. syn.**
- 474 ***B. heeri* Forel, 1874**
- 475 = *B. goeldii* Forel, 1912a **n. syn.**
- 476 = *B. giardi* var. *cordobensis* Santschi, 1929 **n. syn.**
- 477 = *B. physogaster* Kusnezov, 1960 **n. syn.**
- 478 ***B. iridescens* n.sp.**

- 479 *B. micromegas* Emery in Santschi, 1923a  
480 *B. minutus* Forel, 1893  
481 *B. modestus* Santschi, 1923b  
482 *B. musculus* Forel, 1899  
483 *B. myops* Emery, 1906  
484 *B. nebulosus* LaPolla & Longino, 2006  
485 *B. obscurior* Forel, 1893  
486 *B. oculatus* Santschi, 1919  
487 *B. patagonicus* Mayr, 1868  
488 = *B. laevis* Emery, 1895 n. syn.  
489 = *B. patagonicus* var. *atratura* Santschi, 1923a  
490 *B. pictus* Mayr, 1887  
491 = *B. heeri* var. *basalis* Wheeler, 1921. n. syn.  
492 = *B. pictus* subsp. *balboae* Wheeler, 1942 n. syn.  
493 *B. pilipes* Mayr, 1887  
494 *B. santschii* Menozzi, 1927  
495 *B. sosai* n. sp.  
496 *B. termitophilus* Forel, 1895b  
497 *B. tristis* Mayr, 1870

498

499 **Identification key to *Brachymyrmex* species.**

- 500 1 Clypeus with a single long apical hair near the anterior margin, two lateral hairs medially and  
501 two hairs near the toruli (**Fig. 5.a1**); monomorphic ..... 2

502 - Clypeus with a row of long thick hairs near the anterior margin (**Fig. 5.a2**), remaining pilosity not  
503 as above; dimorphic ..... **39**  
504

505 **2(1)** Metathoracic spiracles tumuliform (i.e. strongly protruding dorsally) (**Fig. 6a1**); known only  
506 from Brazil ..... **3**

507 - Metathoracic spiracles not (**Fig. 6.a2**) or slightly protruding but not tumiliform (**Fig. 6.a3**);  
508 naturally occurring throughout the Neotropics ..... **5**  
509

510 **3(2)** Toruli surpassing the posterior clypeal margin in oblique anterodorsal view (**Fig. 5.a3**); head  
511 and mesosoma smooth and shiny ..... **4**

512 – Toruli touching the posterior clypeal margin but never surpassing it in oblique anterodorsal view  
513 (**Fig. 5.a2**); head and mesosoma finely punctate and opaque ..... *B. brasiliensis*  
514

515 **4(3)** Mesosoma without erect hairs; gaster with scattered long erect hairs, except for the first  
516 segment which has dense yellowish pubescence ..... *B. feitosa*

517 – Mesosoma with two erect hairs on pronotum and two on mesonotum; gaster with scattered long  
518 erect hairs, also on the first segment ..... *B. delabiei*  
519

520 **5(2)** Dorsum of the head, mesosoma and gaster with thick erect black hairs (as in *Nylanderia*) that  
521 contrast with the body color (head and gaster may be darker than mesosoma) *B. cavernicola*

522 - Dorsum of the head, mesosoma and gaster without hairs, or with thin hairs that do not contrast  
523 with the body color..... **6**  
524

525	<b>6(5)</b> Eyes positioned below the cephalic midline ( <b>Fig. 5.b1</b> ), with 3 or 4 ommatidia along the	
526	maximal diameter of the eye (EL) ( <b>Fig. 5.c1</b> ).....	7
527	- Eyes usually positioned on the cephalic midline ( <b>Fig. 5.b2</b> ), with more than 4 ommatidia along	
528	the maximal diameter of the eye ( <b>Fig. 5.c2</b> ) .....	9
529		
530	<b>7(6)</b> Mesonotum not bulging dorsally above the pronotum in lateral view ( <b>Fig. 6.b1</b> ).....	8
531	- Mesonotum bulging dorsally above the pronotum in lateral view ( <b>Fig. 6.b2</b> ) .....	
532	.....	<i>B. modestus</i>
533		
534	<b>8(7)</b> Scapes short, just reaching the posterior margin of the head or surpassing it by a length shorter	
535	than the maximal diameter of the eye ( <b>Fig.5.d1,d2</b> ).....	
536	.....	<i>B. donisthorpei</i>
537	- Scapes long, surpassing the posterior margin of the head by a length approximately equal to the	
538	maximal diameter of the eye ( <b>Fig. 5.d3</b> ).....	<i>B. myops</i>
539		
540	<b>9(6)</b> Two erect hairs between the metathoracic spiracles .....	10
541	- Without erect hairs between the metathoracic spiracles .....	11
542		
543	<b>10(9)</b> Scapes surpass the posterior cephalic margin by a length of approximately 1.5× the maximal	
544	diameter of the eye ( <b>Fig. 5.d3: 2A ≤ B</b> ); hairs on scapes decumbent; body uniform in color (usually	
545	dark brown) .....	<i>B. admotus</i>
546	- Scapes surpass the posterior cephalic margin by a length of approximately 1.0× the maximal	
547	diameter of the eye ( <b>Fig. 5.d3: 2A &gt; B</b> ); hairs on scapes appressed; head and mesosoma light brown,	
548	gaster darker .....	<i>B. bonariensis n. st.</i>



549

550 **11(9)** Dorsal margin of the mesosoma having a marked sinusoidal shape (**Fig. 6.c**) ..... **12**

551 - Dorsal margin of the mesosoma not sinusoidal or only of sub-sinusoidal shape (**Figs.**

552 **6.a2,a3,b1,b2,d1,d2,e1,e2**) ..... **13**

553

554 **12(11)** Clypeus with its medial anterior portion forming a “lip” (**Fig. 5.e1**); head and mesosoma

555 partially or completely alveolate (sometimes alveolate-strigate); dorsum of the mesosoma with

556 many erect hairs; body uniform in color ..... *B. nebulosus*

557 -Clypeus without anteromedial “lip” (**Fig. 5.e2**); entire body non-alveolate; dorsum of the

558 mesosoma without erect hairs; head and gaster black; mesosoma yellowish

559 ..... *B. bicolor n. sp.*

560

561 **13(11)** Head with strong alveolate sculpture ..... **14**

562 - Head without alveolate sculpture ..... **15**

563

564 **14(13)** Metanotal groove wider than the diameter of the metathoracic spiracles (**Fig. 6.f1**:  $A \leq B$ );

565 scapes surpassing the posterior margin of the head by approximately 1.0× the maximal diameter of

566 the eye (**Fig. 5.d3**); gaster with scattered pubescence (**Fig. 6.g1**) ..... *B. santschii*

567 - Metanotal groove narrower than the diameter of the metathoracic spiracles (**Fig. 6.f2**:  $A > B$ );

568 scapes just reaching the posterior margin of the head (**Fig. 5.d2**); gaster with dense pubescence

569 (**Fig. 6.g2**) ..... *B. iridescens n. sp.*

570

571 **15(13)** Mesometanotal suture inconspicuous (**Fig. 6.d1**) ..... **16**

572 - Mesometanotal suture readily visible (**Fig. 6.d2**) ..... **17**

573

574 **16(15)** Pronotum without erect hairs; scapes short or reaching the posterior margin of the head  
575 **(Fig. 5.d1,d2)**; gaster with dense pubescence **(Fig. 6.g2)** ..... *B. flavidulus*  
576 - Pronotum with two erect hairs **(Fig. 6.d1)**; scapes surpassing the posterior margin of the head  
577 **(Fig. 5.d3)**; gaster without dense pubescence, but with scattered appressed hairs **(Fig. 6.g1)** ..  
578 ..... *B. minutus*  
579

580 **17(15)** Gaster with dense appressed or decumbent pubescence **(Fig. 6.g2)** ..... **18**  
581 - Gaster with sparse pubescence, but with scattered, appressed hairs **(Fig. 6.g1)** ..... **28**  
582

583 **18(17)** Metanotal groove absent or when present shallow and narrower than the diameter of the  
584 metathoracic spiracles **(Fig. 6.f2: A > B)**..... **19**  
585 - Metanotal groove deep and wider than the diameter of the metathoracic spiracles **(Fig. 6.f1: A ≤**  
586 **B)**..... **26**  
587

588 **19(18)** Mesonotum bulging dorsally above the pronotum in lateral view **(Fig. 6.b1)** ..... **20**  
589 - Mesonotum not bulging dorsally above the pronotum in lateral view **(Fig. 6.b2)** ..... **22**  
590

591 **20(19)** Scapes just reaching the posterior margin of the head or surpassing it by a length of less  
592 than 1.0× the maximal diameter of the eye **(Fig. 5.d2,d3: A > B)** ..... **21**  
593 - Scapes surpassing the posterior margin of the head by a length of approximately 1.0× the maximal  
594 diameter of the eye **(Fig. 5.d3: A ≈ B)**..... *B. heeri*  
595

596 **21(20)** Body usually dark brown; eye with on average 9 ommatidia along its maximal diameter;  
597 scapes on average >0.5 mm; known only from South America ..... *B. giardi*  
598 - Body yellowish; eye with on average 6 ommatidia along its maximal diameter; scapes on average  
599 <0.5 mm; known only from Canada, Mexico, USA ..... *B. depilis*  
600  
601 **22(19)** Body yellowish ..... **23**  
602 - Body dark brown..... **25**  
603  
604 **23(22)** Scapes not or barely reaching the posterior margin of the head (**Fig. 5.d1,d2**)  
605 ..... *B. fiebrigi*  
606 - Scapes surpassing the posterior margin of the head (**Fig. 5.d3**) ..... **24**  
607  
608 **24(23)** About 6 erect hairs on the pronotum and two on the mesonotum, each hair with a length of  
609 about 2.0× the maximal diameter of the eye; known only from the Bahamas. *B. bahamensis n. sp.*  
610 - Two erect hairs on the pronotum and two on the mesonotum, each with a length shorter than the  
611 maximal diameter of the eye; widespread..... *B. termitophilus*  
612  
613 **25(22)** Dorsum of the head and mesosoma with light-colored, dense pubescence; gaster with dense  
614 appressed pubescence; eye with on average 11 ommatidia along its maximal diameter, head on  
615 average long (HL1 > 0.5 mm) and wide (HW > 0.4 mm)..... *B. cordemoyi*  
616 - Dorsum of the head and mesosoma with less conspicuous dense pubescence; gaster with dense  
617 decumbent pubescence; eye with on average 9 ommatidia along its maximal diameter, head on  
618 average short (HL1 < 0.5 mm) and narrow (HW < 0.4 mm) ..... *B. obscurior*  
619

620 **26(18)** Dorsum of the mesosoma without conspicuous sculpture; metathoracic spiracles fully dorsal  
621 in position; dorsal margin of the mesonotum strongly antero-posteriorly inclined (**Fig. 6.e1**) **B.**

622 *sosai* **n.sp.**

623 - Dorsum of the mesosoma with imbricate sculpture; metathoracic spiracles in dorsolateral  
624 position; dorsal margin of the mesonotum not or slightly antero-posteriorly inclined (**Fig. 6.e2**)

625 ..... **27**

626  
627 **27(26)** Second segment of the antennal funiculus shorter than the first antennal segment (**Fig. 5.g1:**  
628  $S2 < S1$ ); scapes with appressed hairs; metathoracic spiracles protruding slightly dorsally, but not  
629 tumiliform (**Fig. 6.a3**); hairs lighter in color than the body, which is brownish .....

630 ..... ***B. attenuatus* n.st**

631 - Second segment of the antennal funiculus as long or longer than the first antennal segment (**Fig.**  
632 **6.g2:**  $S2 \geq S1$ ); scapes with decumbent hairs; methatoracic spiracles not protruding (**Fig. 6.a2**)  
633 hairs darker in color than the body, which is yellowish ..... ***B. antennatus***

634  
635 **28(17)** Eyes large, with a maximal diameter  $>1/4$ th of the length of the head ( $HL_1$ ), usually with  
636  $>14$  ommatidia along their maximal diameter ..... ***B. oculus***

637 - Eyes small, with a maximal diameter of approximately  $1/4$ th the length of  $HL_1$ , typically with  
638  $<14$  ommatidia along their maximal diameter ..... **29**

639  
640 **29(28)** Metanotal groove absent, or, when present, shallow and narrower than the diameter of the  
641 metathoracic spiracles (**Fig. 6.f2:**  $A > B$ )..... **30**

642 - Metanotal groove deep and wider than the diameter of the metathoracic spiracles (**Fig. 6.f1:**  $A \leq$   
643 **B.**) ..... **34**

644

645 **30(29)** Head and thorax yellowish; gaster black or yellowish with a black spot,  $OI_2$  usually  $>27$

646 *B. pictus*

647 - Body of uniform color,  $OI_2$  usually  $<25$  ..... **31**

648

649 **31(30)** Body yellowish, usually with a narrow mesonotum (MW ~ 16) and 8-9 ommatids along the

650 maximum diameter of the eye ..... **32**

651 - Body brownish or dark brown, usually with a wide mesonotum (MW ~ 20 or more) and 10 or

652 more ommatids along the maximum diameter of the eye ..... **33**

653

654 **32(31)** Scapes surpassing the posterior margin of the head by a length exceeding the maximal

655 diameter of the eye (**Fig. 5.d3**:  $A < B$ ) ..... *B. aphidicola*

656 -Scapes surpassing the posterior margin of the head by a length smaller than or equal to the maximal

657 diameter of the eye (**Fig. 5.d3**:  $A \geq B$ ) ..... *B. australis*

658

659 **33(31)** Scapes surpassing the posterior margin of the head by a length smaller than the maximal

660 diameter of the eye (**Fig. 5.d3**:  $A > B$ ); usually with two erect hairs on the pronotum and two on

661 the mesonotum ..... *B. patagonicus*

662 - Scapes surpassing the posterior margin of the head by a length approximately equal to the

663 maximal diameter of the eye (**Fig. 5.d3**:  $A \approx B$ ); usually with more than two erect or decumbent

664 hairs on the pronotum and two erect hairs on the mesonotum ..... *B. bruchi*

665

666 **34(29)** Legs and antennae with erect hairs; second segment of the antennal funiculus as long as or

667 longer than the first (**Fig. 5.g2**:  $S_2 \geq S_1$ ) ..... *B. gaucho*

668 - Legs and antennae with decumbent or appressed hairs; second segment of the antennal funiculus  
669 shorter than the first (**Fig. 5.g1**: S2 < S1) ..... **35**  
670  
671 **35(34)** Mesonotum not bulging dorsally above the pronotum in lateral view (**Fig. 6.b2**);  
672 metathoracic spiracles low, not protruding dorsally (**Fig. 6.a2**) ..... *B. musculus*  
673 - Mesonotum bulging dorsally above the pronotum in lateral view (**Fig. 6.b1**); metathoracic  
674 spiracles protruding slightly in lateral view but not tumiliform in shape (**Fig. 6.a3**) ..... **36**  
675  
676 **36(35)** Head and thorax yellow or brown, gaster darker ..... *B. coactus*  
677 - Body uniform in color..... **37**  
678  
679 **37(36)** Head with dense decumbent pubescence (**Fig. 5.f1**) ..... *B. tristis*  
680 - Head with sparse decumbent pubescence (**Fig. 5.f2**)..... **38**  
681  
682 **38(37)** Mesonotum laterally extended and therefore oval in dorsal view (**Fig. 6.h1**); body light  
683 brown ..... *B. degener*  
684 -Mesonotum almost circular in dorsal view (**Fig. 6.h2**); body dark brown or black .....  
685 ..... *B. gagates*  
686  
687 **39(1)** Mesosoma mostly smooth and shiny, except for longitudinal striations restricted to the  
688 metapleura; body uniform light brown..... *B. micromegas*  
689 -Mesosoma entirely covered with fine longitudinal striations; gaster darker than the rest of the  
690 body..... *B. pilipes*  
691

692 **Species accounts.**

693 *Brachymyrmex admotus* Mayr

694 (Fig. 7, supplementary material Fig. S1)

695 *Brachymyrmex admotus* Mayr, 1887: 523 (w.q.). **Lectotype worker** (NHMW:  
696 USNMENT00757197) and **paralectotype workers, queen** (NHMW: USNMENT00757196,  
697 00757198-00757200; **here designated**): 5 workers, 1 queen [examined]. **BRAZIL: Santa**  
698 **Catharina**. Other relevant descriptions: Wheeler and Wheeler (1982: 178) (l.). See also: Santschi  
699 (1923a: 669); Quirán (2005: 762).  
700 =*Brachymyrmex longicornis* var. *immunis* Forel, 1908: 400 (w.q.m.). (MHNG:  
701 USNMENT00757148): 2 workers [examined]. **BRAZIL: São Paulo**. See also: Forel (1911: 308);  
702 Santschi (1923a: 668) **n. syn.**

703  
704 **Additional material examined. ARGENTINA: Misiones:** Est. Ex. Loreto, A.A. Oglobin, 3  
705 workers (NHMB: USNMENT00758065 - 00758067). **BRAZIL: Bahia:** Itacaré, -14.30917 -  
706 39.01944, 26 June 1998, Santos, J.R.M. dos, 2 workers (CPDC: USNMENT00757769); Ituberá,  
707 08 May 1994, 4815, J.H.C. Delabie, 3 workers (CPDC: USNMENT00757772); **Mato Grosso do**  
708 **Sul:** 8 km SE Ponta Pora, 15 Oct. 1989, W.P. MacKay #12523, 2 workers (WEMC:  
709 USNMENT00759009); **Minas Gerais:** Alfenas, 05 May 2011, 1 worker (ICN:  
710 USNMENT00759050); Cristina, Luederwaldt, 9 workers (NHMB: USNMENT00758053,  
711 00758059, 00758061); Cristina, MP17192, 2 workers (MZSP: USNMENT00757765, 00757819);  
712 Cocais das Estrelas, -19.73333 -43.41667, 19-22 June 2007, D.L. Braga #5512, 1 worker (CPDC:  
713 USNMENT00757768); Serra Caraça, 1380 m, Nov. 1961, Kloss, Lemko, 2713, Martins & Silva,  
714 9 workers, 3 males (MCZC: USNMENT00757252, 00757253, 00757764); Viçosa, Mata do

715 Paraiso, Dec. 1993-1994, P.S.F. Ferreira, 3 workers (CPDC: USNMENT00757770); **Paraná:**  
716 Antonina, Parque Estadual do Pauôco, -25.57597 -48.88875, 6-11 May 2002, R.R. Silva & B.H.  
717 Dietz, 24 workers (ICN: MZSP016, 018, 019); Río Negro, Goeldi, 2 workers (MCZC:  
718 USNMENT00757235); **Rio de Janeiro:** Reischensperger, 8 workers (NHMB:  
719 USNMENT00758056-00758058); Goeldi, 1 worker (NHMB: USNMENT00758050); Floresta de  
720 Tijuca, D. Federal, 16 Dec. 1959, C.A.C Seabra, 5 workers (MZSP: USNMENT00757766);  
721 Itatiaia, 17 Oct. 1933, 1 worker (NHMB: USNMENT00758068); Petropolis 77 9, T. Borgmeier,  
722 5 workers, 1 queen (MCZC: USNMENT00757233, 00757234, 00757236); **Santa Catarina:**  
723 Blumenau, Reichensperger, 9 workers (NHMB: USNMENT00758055, 00758060, 00758064);  
724 Blumenau, Rev PM Witte, 2 workers (NHMB: USNMENT00758063); Blumenau, Rev Wittz, 19  
725 workers (NHMB: USNMENT00758051, 00758052, 00758062); Palhoça, PE Serra do Tabuleiro,  
726 -27.74111 -48.69722, 02-10 June 2003, R.R. Silva, B.H. Dietz & A. Tavares, 25 workers (ICN:  
727 MZP030, 031, 035, 040); São Bento do Sul, APA Rio Vermelho, -26.36417 -49.27111, 30 Mar.-  
728 04 Apr. 2001, R.R Silva & Eberhardt, 27 workers (ICN: MZP043, 134, 137); São Bento do Sul,  
729 APA Rio Vermelho, -26.36417 -42.27111, 30 Mar.-04 Apr. 2001, R.R. Silva & R.M. Feitosa, 5  
730 workers (ICN: MZP044); **São Paulo:** Agudos, 24 Jan. 1955, W.S. Kempf leg 1337, 3 workers  
731 (MZSP: USNMENT00757767); Barueri, n 297, 17 Dec. 1957, K. Lenko, 5 workers (MZSP:  
732 USNMENT00757775); Iguape, EE Jureia-Itatins, Nucleo Rio Verde, -24.54417 -47.23556, 5-14  
733 Mar. 2001, A.A. Tavares, 13 workers (ICN: MZP157, 158); Ipiranga Ihering, 4 workers  
734 (MHNG); Ipiranga (x.60), Ihering, 2 workers (MHNG); Jardim Botânico, Agua Funda, wet  
735 Forest, 08 Feb. 1967, W.L. Brown, 5 workers, 1 queen, 1 male (MCZC: USNMENT00757961,  
736 00757771, 00757773, 00757774, CMOS00148, 00153); Jundiai, Serra Do Japi, 03 Jan. 2009, S.  
737 Diniz, 4 workers (ICN: USNMENT00759039); Miracatu, Serra do Mar, Clube pesca & Cia, 04-  
738 07 Sep. 2004, R.M. Feitosa, 11 workers (ICN: MZP092, 097); Picinguaba, PE Serra do Mar, -



739 23.33611 -44.83758, 30 Mar.-04 Apr. 2001, Brandão C.R.F. & Eq, 52 workers, 1 queen (ICN:  
740 MZP060 - 062, 064); Picinguaba, PE Serra do Mar, -23.33611 -44.83758, 30 Mar.-04 Apr. 2001,  
741 Brandão, Albuquerque & Silva, 15 workers (ICN: MZP063); Piedade, Floresta Atlantica  
742 "Theomar", Mar 2010, G. Bieber, 1 worker (ICN: USNMENT00759040); Piedade, Floresta  
743 Atlantica, Jurupará, Apr. 2009, G. Bieber, 1 worker (ICN: USNMENT00759041); Serra du  
744 Cantareira, Horto Florestel, 20 Feb. 1967, R. Crozier, 9 workers (MCZC: USNMENT00757774,  
745 CMOS000089, 000090); **PANAMA: Colon Province:** San Lorenzo Forest, 9-16 Feb. 2004,  
746 Springate & Pinzon, 1 worker, (PSWC: USNMENT00757747). **PARAGUAY: Canindeyú:**  
747 Reserva Natural, Bosque Mbaracayú, Jejuimi, -24.10000 -55.50421, 15 Aug. 1996, A. Wild  
748 #AW0295, 3 workers (ALWC: USNMENT00757763).

749  
750 **Diagnosis.** *Brachymyrmex admotus* morphologically resembles *B. bonariensis* **n. st.**, because  
751 both species have long scapes, a conspicuous metanotal groove, a pair of thin erect hairs between  
752 the metathoracic spiracles, and a gaster with scarce pubescence. However, *B. admotus* is usually  
753 more uniform brownish in color, it has longer scapes (i.e. the scapes surpass the posterior  
754 cephalic margin with a length of approximately 1.5× the maximal diameter of the eye) with  
755 decumbent hairs, and its metathoracic spiracles are positioned more dorsally. *Brachymyrmex*  
756 *admotus* also resembles *B. cavernicola* in having a pair of erect hairs between the dorsal  
757 metathoracic spiracles, but these hairs are thinner in *B. admotus* and they are not darker in color  
758 than the body.

759  
760 *Lectotype measurements* (mm). HL<sub>1</sub> 0.51; HL<sub>2</sub> 0.35; HL<sub>3</sub> 0.12; HW 0.45; SL 0.49; EL 0.10; WL  
761 0.49; PnL 0.14; PnW 0.31; ML 0.08; MW 0.18; *Indices* CI 88.46; SI<sub>1</sub> 108.70; SI<sub>2</sub> 138.89; OI<sub>1</sub>  
762 21.74; OI<sub>2</sub> 23.08.

763  
764 *Paralectotypes measurements* (mm) (n=3). HL<sub>1</sub> 0.51-0.57; HL<sub>2</sub> 0.35-0.39; HL<sub>3</sub> 0.12-0.14; HW  
765 0.45-0.49; SL 0.55-0.59; EL 0.10; WL 0.53-0.59; PnL 0.14-0.21; PnW 0.31-0.35; ML 0.08-0.12;  
766 MW 0.18-0.21; *Indices* CI 86.21-92.59; SI<sub>1</sub> 112.00-121.74; SI<sub>2</sub> 140.00-155.60; OI<sub>1</sub> 20.00-21.74;  
767 OI<sub>2</sub> 23.08-25.93.

768  
769 *Additional material examined measurements* (mm) (n=16). HL<sub>1</sub> 0.46-0.57; HL<sub>2</sub> 0.30-0.43; HL<sub>3</sub>  
770 0.11-0.14; HW 0.43-0.51; SL 0.47-0.57; EL 0.09-0.13; WL 0.46-0.61; PnL 0.16-0.22; PnW 0.29-  
771 0.34; ML 0.09-0.13; MW 0.17-0.21; *Indices* CI 87.72-96.6; SI<sub>1</sub> 105.26-120.00; SI<sub>2</sub> 131.91-  
772 155.88; OI<sub>1</sub> 18.87-26.92; OI<sub>2</sub> 21.43-28.30.

773  
774 **Description. Head.** Slightly longer than wide in full face view, with scattered appressed hairs  
775 except for two frontal rows of erect hairs; posterior cephalic border slightly concave. Dorsum of  
776 the head with sparse appressed pubescence. Clypeus with a rounded anterior margin and five  
777 long, erect hairs of which a single, usually conspicuous apical hair is near the anterior margin,  
778 two lateral hairs in medial position and two more near the toruli; other hairs on the clypeus are  
779 markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in  
780 oblique anterodorsal view. Scapes extend beyond the posterior cephalic margin by a length of  
781 ~1.5× the maximum eye diameter or more. The scapes typically have appressed, sometimes  
782 decumbent, but never erect hairs. A single central ocellus is present, but sometimes  
783 inconspicuous. Eyes are positioned on the cephalic midline and have 7-9 ommatidia along their  
784 maximal diameter.

785 **Mesosoma.** Usually with two erect hairs on the pronotum and two on the mesonotum; sometimes  
786 with additional suberect hairs on both. In lateral view, the mesonotum is not or slightly inflated

787 and it does not or only slightly bulge dorsally above the pronotum. Metanotal groove deep and  
788 wider than the diameter of the metathoracic spiracles. Metathoracic spiracles fully dorsal and  
789 slightly protruding, they are closer to the propodeal than to the mesometanotal suture, but not  
790 touching any suture. Between the metathoracic spiracles two simple erect hairs are present, which  
791 are shorter than those on the pronotum and mesonotum. Dorsum of the propodeum flat and  
792 ~1/3th of the length of the propodeal slope. Propodeal spiracles circular, positioned ventrally of  
793 the posterior propodeal margin slightly posterior of the middle of the propodeal slope. Legs with  
794 appressed and scattered hairs. Petiole short and inclined forward.

795 **Gaster.** With scattered pubescence and scattered long suberect hairs.

796 **Color and sculpture.** Body overall smooth and shiny, except for the sometimes slightly imbricate  
797 sculpture on the dorsum of the mesosoma; body typically uniform dark brown in color, although  
798 in some specimens the head and mesosoma may be light brownish and the gaster darker brown.

799

800 **Distribution (Supplementary material Fig, S1).** *Brachymyrmex admotus* is mainly known  
801 from Argentina, Brazil, and Paraguay, but we also examined a specimen from Panamá that  
802 appears to belong to this species.

803

804 **Biology.** This species makes nests in rotting wood [USNMENT00757763] and it has been  
805 collected from below rocks [USNMENT00759009].

806

807 **Remarks.** We synonymize *B. longicornis* var. *immunis* Forel, 1908 under *B. admotus*, because  
808 the workers have all morphological characteristics of *B. admotus*, although they are slightly  
809 larger and of somewhat darker color. Forel (1908) did not specify a holotype for *B. longicornis*  
810 var. *immunis* but considered it to differ from *B. longicornis* in color, size and the position of the

811 metathoracic spiracles. The similarity between *B. admotus* and *B. longicornis* var. *immunis* was  
812 previously pointed out by Santschi (1923a) and Quirán (2005), who suggested that the main  
813 difference between both relates to the size and position of the metathoracic spiracles. However,  
814 these traits appear to be variable among populations and we have not found consistent differences  
815 between both taxa. For example, Santschi's collection (MHNB) includes syntype specimens of *B.*  
816 *longicornis* var. *immunis* that match the diagnostic traits of *B. admotus* entirely. Hence,  
817 synonymization seems appropriate.  
818 Quirán (2005) indicated that *B. admotus* has 3 small ocelli, but in the samples that we studied we  
819 only observed one central ocellus, although this trait is inconspicuous.

820

821

822 *Brachymyrmex antennatus* Santschi

823 (Fig. 8, supplementary material Fig. S2)

824 *Brachymyrmex (Brysha) antennatus* Santschi, 1929: 312 (w.q.). **Lectotype worker** (NHMB:  
825 USNMENT00758161) and **paralectotype worker, queen** (NHMB: USNMENT00758161; **here**  
826 **designated**): 2 workers, 1 queen (without USNMENT number) [examined]. **Brazil:** Paraná, Rio  
827 Negro.

828

829 **Additional material examined. ARGENTINA: Misiones:** loc. IGUAZU: PNI, Garganta, 28  
830 Feb.-03 Mar. 2009, -25.70323 -54.42992, P.E. Hanisch & C.I Paris, Bait T4S10a, 1 worker  
831 (MACN-Bar-Ins-3120). **BRAZIL: Ceará:** Meruoca (Baixa, Gnd), ±970 m, -3.550 -40.467, July  
832 2003, Y. Quinet, 3 workers (CPDC: USNMENT00757781). **São Paulo:** Ubatuba, P.E. Serra do  
833 Mar, N. Picinguaba, -23.297 -44.789, 800 m, 03-14 Mar. 2008, armadilha subterrânea #4, F.A.  
834 Esteves & R.M. Feitosa, 3 workers (MZSP: USNMENT00757777, 00757591). **COLOMBIA:**

835 **Quindío:** Génova, Vereda El Cedral, Finca Buenos Aires, 1600 m, 4.235 -75.775, 26 Oct. 1999,  
836 E. Gonzales & J. Sossa, 1 worker (IAvH-E 74166). **ECUADOR: Napo:** Carlos Julio Arosemena  
837 Tola, -1.150 -77.883, 500 m, 11 Dec. 2003. A. L. Wild #AW2300, 1 worker (ALWC:  
838 USNMENT00757782). **FRENCH GUIANA:** Petit, Satn Basse vie, June-July 2000, S. Durou, J.  
839 Delabie, A. Dejean & A. Gibernau, 2 workers (CPDC: USNMENT00757779, 00757780).  
840 **PERU: Madre de Dios:** Reserva Nacional Tambopata, Centro Sachacavayoc, -12.85583 -  
841 69.36194, 210 m, 19-31 July 2012, 2 workers (ICN: USNMENT00757627). Prov. Tambopata,  
842 Cuzco Amazónico, 15 km NE Puerto Maldonado, 24 June 1989, 200 m, S.P. Cover & J.E. Tobin,  
843 CA-115, 1 worker, 1 queen (MCZC: USNMENT00757630). **SURINAME:** Maripahewel, IX-  
844 1959 14-XX-29 I.v.d. Drift, 1 worker (MZSP: USNMENT00757778).

845  
846 **Diagnosis.** *Brachymyrmex antennatus* morphologically resembles *B. gaucho*, because both  
847 species have legs and antennae with suberect hairs and both have an antennal funiculus with the  
848 second segment as long as or longer than the first. However, they differ from one another because  
849 *B. antennatus* has abundant, suberect hairs on the dorsum of the head and mesosoma, its gaster  
850 has dense pubescence and its body is lighter and yellowish. *Brachymyrmex antennatus* also  
851 resembles *B. cavernicola* in having suberect hairs on the mesosoma that are generally darker in  
852 color than the tegument, but *B. antennatus* has a more elongated head, a longer second segment  
853 of the antennal funiculus, as mentioned above, thinner hair on its body and denser pubescence on  
854 the gaster.

855  
856 *Lectotype and paralectotype measurements* (mm) (n=2). HL<sub>1</sub> 0.68-0.71; HL<sub>2</sub> n.a.; HL<sub>3</sub> 0.19-0.23;  
857 HW 0.68-0.71; SL 0.68-0.80; EL 0.15; WL 0.87; PnL 0.31-0.33; PnW 0.45; ML 0.21; MW 0.27;  
858 *Indices* CI 95.50-95.80; SI<sub>1</sub> 100.00-123.00; SI<sub>2</sub> n.a.; OI<sub>1</sub> 22.22-23.25; OI<sub>2</sub> 27.94-31.41.

859  
860 *Additional material examined measurements* (mm) (n=5). HL<sub>1</sub> 0.54-0.60; HL<sub>2</sub> 0.38-0.41; HL<sub>3</sub>  
861 0.12-0.16; HW 0.54-0.58; SL 0.52-0.63; EL 0.09-0.14; WL 0.60-0.71; PnL 0.15-0.22; PnW 0.35-  
862 0.40; ML 0.14-0.18; MW 0.20-0.24; *Indices* CI 92.31-100.00; SI<sub>1</sub> 93.33-130.23; SI<sub>2</sub> 130.23-  
863 155.56; OI<sub>1</sub> 15.38-26.67; OI<sub>2</sub> 21.67-29.03.

864  
865 **Description. Head.** Longer than wide in full face view; posterior cephalic border concave.  
866 Dorsum of the head with scattered decumbent hairs. Clypeus with a rounded anterior margin and  
867 five long, erect hairs of which a single, usually conspicuous apical hair is near the anterior  
868 margin, two lateral hairs in medial position and two more near the toruli; other hairs on the  
869 clypeus are conspicuously shorter and appressed or decumbent. Toruli surpassing the posterior  
870 clypeal margin (in oblique anterodorsal view). Scapes surpass the posterior cephalic margin by a  
871 length of 1.5× the maximum eye diameter or more. The second segment of the antennal funiculus  
872 is as long as the first or longer. The scapes typically have suberect and erect hairs. Three ocelli  
873 present. Eyes are positioned on the cephalic midline and have 7-9 ommatidia along their maximal  
874 diameter.

875 **Mesosoma.** With conspicuous, thin erect hairs of darker color than the tegument. Pronotum  
876 anteroposteriorly elongated. The mesonotum is slightly antero-posteriorly inclined, weakly  
877 inflated, and it does not bulge dorsally above the pronotum in lateral view. Metanotal groove  
878 deep and wider than the diameter of the metathoracic spiracles. Metathoracic spiracles in  
879 dorsolateral position and slightly protruding, closer to the propodeal than to the mesometanotal  
880 suture, but not touching any suture. Dorsum of propodeum flat and equal in length to the  
881 propodeal slope. Propodeal spiracles circular, situated ventral of the posterior propodeal margin.  
882 Legs with suberect and erect hairs. Petiole short and inclined forward.

883 **Gaster.** With dense pubescence and scattered long suberect hairs, mainly at the edges of the  
884 segments.

885 **Color and sculpture.** Body typically uniformly light brown, although some specimens may be  
886 light brownish with the head and gaster darker brown.

887

888 **Distribution (Supplementary material Fig. S2).** Known from Argentina, Brazil, Colombia,  
889 Ecuador, French Guiana, Perú and Suriname.

890

891 **Biology.** Unknown.

892

893 **Remarks.** The ant at the top of pin USNMENT00758161 is designated here as lectotype and the  
894 one immediately below as paralectotype. Santschi's collection (MHNB) contains three additional  
895 pins with four workers from the same locality but they are not considered to be part of the type  
896 collection as they have no type label. Santschi (1929) described the queen from a sample that  
897 does not contain any workers but expressed confidence that it belongs to *B. antennatus*; the issue  
898 may require verification from independent material. Substantial variation exists in the body size  
899 of workers of *B. antennatus* from various locations that were studied here, and the cause of this  
900 variation remains uncertain.

901

902

903 ***Brachymyrmex aphidicola* Forel**

904 **(Fig. 9, supplementary material Fig. S3)**

905 *Brachymyrmex heeri* var. *aphidicola* Forel, 1909: 263 (w.). **Lectotype worker** (MHNG:

906 USNMENT00757130) and **paralectotype workers** (MHNG: USNMENT00757129, 00757130,

907 00758121-00718123; **here designated**): 11 workers [examined]. **PARAGUAY**: San Bernardino.  
908 Other relevant descriptions: Forel (1912a: 62) (q.). (MHNG: USNMENT00757128): 1 queen.  
909 **BRAZIL: Santa Catarina**: Blumenau. Raised to species: Wild (2007: 43).  
910 = *B. heeri* var. *fallax* Santschi, 1923a: 665 (w.). (NHMB: USNMENT00757697): 1 worker  
911 [examined]. **PARAGUAY**. Junior synonym of *B. aphidicola*: Wild (2007: 43).  
912 = *B. longicornis* var. *hemiops* Santschi 1923a: 668 (w.). (NHMB: USNMENT00757188-  
913 00757190) 11 workers, 3 queens [examined]. **BRAZIL**: São Paulo, Ypiranga. **n. syn.**  
914  
915 **Additional material examined. ARGENTINA**: Entre Ríos: 8.63 km W Concordia, -31.42048 -  
916 58.11700, 16 m, 27 Dec. 2007, W. & E. MacKay, 1 worker, 1 male (WEMC:  
917 USNMENT00757975). **Misiones**: Parque Provincial Cañadón de Profundidad, -27.56020 -  
918 55.70988, 160 m, 29 Dec. 2007, W. & E. MacKay leg #22710, #22711, #22712, #22724, #22732,  
919 10 workers (WEMC: USNMENT00757617, 00757897, 00757901, 00757924, 00757925,  
920 00757929, 00757930, 00757956, 00757992). **BOLIVIA: Santa Cruz**: Parque Nacional Noel  
921 Kempff, Mercado, -18.800 -60.383, 700 m, 04 Dec. 1993, P.S. Ward #12285-46, 2 workers, 1  
922 queen (PSWC: USNMENT00757910). **BRAZIL: Amazonas**: 61 km N Manaus, om Caracá Rd,  
923 “caatinga”, 10 June 1972, W.L. & D.E. Brown, 3 workers (MCZC: UNSMENT00757619).  
924 **Goias**: Anapolis, 12 Feb. 1958, W. Kempf, 2 workers (MZSP: USNMENT00757921). Ouro  
925 Verde, Faz Boa Vista, -16.29847 -49.21183, 01-07 Aug. 2005, R.R. Silva & R.M. Feitosa, 5  
926 workers (ICN: MZSP123). **Mato Grosso**: Chapada dos Guimarães, -15.43333 -55.44874, 740 m,  
927 03 Sep. 1996, P.S. Ward leg #13203-7, 3 workers (PSWC: USNMENT00757911). **Mato Grosso**  
928 **du Sud**: 24 km W Campo Grande, 07 Oct. 1989, W. MacKay, 2 workers (WEMC:  
929 USNMENT00758000). 48 km E Campo Grande, 12 Oct. 1989, S. Porter #12791, 2 workers  
930 (WEMC: USNMENT00759011). 8 km SE Punta Bora, 15 Oct. 1989, W.P. MacKay #12508, 2



931 workers (WEMC: USNMENT00759003). Río de Alegría, 17 Oct. 1989, W.P. MacKay #12950, 2  
932 workers (WEMC: USNMENT00759022). **Pará:** Melgaço, Caxiuanã-ECFPn, 27 June-03 Dec.  
933 2001, I. Andrade, 5 workers (MPEG: USNMENT00757674, 00757927, 00758030, 00759030).  
934 Melgaço, Caxiuanã-ECFPn, -1.78155 -51.59758, 30 Oct. 2003, A.Y. Harada, E.P. Fagundes,  
935 C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 2 workers (MPEG:  
936 AYH112, 127). Serra Norte, Calderião, 20 Oct. 1980, 2 workers (MPEG:  
937 MPEG\_HYM11505158, USNMENT00757902). **Rio de Janeiro:** Teresópolis, P.N. Serra dos  
938 Orgãos, -22.45333 -42.99806, 23-28 Nov. 1999, Dietz, Silva & Rocha, 8 workers (ICN:  
939 MZSP130). **Rondônia:** Ouro Preto do Oeste, 03 Apr. 1985, Linha 212 N°0375, 339, W. França,  
940 4 workers (MPEG: USNMENT00757899,00757914, 00757915, 00758999,). Ouro Preto do  
941 Oeste, 04 May 1985, Linha 212 N°0375, 0413, W. França, 3 workers (MPEG:  
942 USNMENT00757913, 00757938, 00758038). Ouro Preto do Oeste, 25 Mar. 1985,  
943 ResINPA0035, J. Dias, 2 workers (MPEG: USNMENT00757914, 00757936). **São Paulo:**  
944 Ipiranga, 2371, 2 workers (MZSP: USNMENT00757926]. Itirapina, Dec. 2008, D.P. Silva, 1  
945 worker (MPEG: AYH008). **COLOMBIA: Cundinamarca:** Bogotá-Villavicencio Km 88  
946 (Susumuko), 1100 m, 28 June 1976, W.L. & D.E. Brown, 1 worker, 1 queen (MCZC:  
947 USNMENT00757746). Caqueza, 29 Dec. 1975, W. & E. MacKay #945, 2 workers (WEMC:  
948 USNMENT00757686). **Huila:** 4 km NE Rivera, 30 Dec. 1986, W. & E. MacKay, 3 workers  
949 (WEMC: USNMENT00757903). La Vega, A280, 14-17 Jul 1975, W. MacKay, 1 worker, 1 male  
950 (WEMC: USNMENT00757912). **Magdalena:** 4 km San Pedro, 10.95 -74.05, 550 m, 14 Aug.  
951 1985, P.S. Ward #17912-36, 3 workers (PSWC: USNMENT00757585). **Meta:** San Juan de  
952 Arama, RNN La Macarena, Caño La Curía, 580 m, 13 July 1992, Est. U. Nacional, 1 worker  
953 (ICN: USNMENT00758035). **Tolima:** Mendez, 15 Nov. 1995, F. Fernández, 1 worker (IAvH:  
954 USNMENT00759058). **Valle del Cauca:** Medio Calima, 24 June 1989, E. MacKay #11746

955 (WEMC: USNMENT00758041). **COSTA RICA: Heredia:** Estación biológica La Selva, 10.417  
956 -84.000, 50 m, 21 Oct. 1991, J. Longino #3126-s, 1 worker, 1 queen (JTLC:  
957 INBIOCRI001238064). **ECUADOR: Loja:** Estación San Francisco, 2200 m, 14 Sep. 2011, 14  
958 workers (ICN: USNMENT00759036, 00759037). Estación San Francisco, 17 sep. 2011, 2  
959 workers (ICN: USNMENT00759034). **Napo:** 11 km SE Consaga, -0.66667 -77.80000, 1640 m,  
960 09 Dec. 2003, A.L. Wild #AW2263, 4 workers (ALWC: USNMENT00757586, 00757928).  
961 **GUATEMALA:** Alta Vera Paz, Parque Nacional Las Victorias, 15.47492 -90.37528, 206 m, 18  
962 July 2004, W. & E. MacKay, 1 worker (WEMC: USNMENT00758018). **GUYANA: Karto:** Pt.  
963 Mazaruni-Potaru Dist. J. Weintraub, 2 workers, 1 queen (MCZC: USNMENT00757931).  
964 **MEXICO: Guanajuato:** Highway 57, Km 306, Rancho Jardin, 10 Aug. 1965, Cornell  
965 University, 2 workers, 1 queen (MCZC: USNMENT00759002). Yuriria, 03 Feb. 1964, P. Reyes  
966 C. & H. Romero, 1 worker (MZSP: USNMENT00757616). **NICARAGUA: Río San Juan:**  
967 Bartola, 8 km SE El Castillo, 10.97303 -84.33897, 47 m, 12 July 2003, W. & E. MacKay #20187,  
968 2 workers (WEMC: USNMENT00758029). **PANAMA: San Blas:** Nusegandi, 08 June 1992,  
969 L.E. Tennant, 1 worker, 1 queen (MZSP: USNMENT00757923). **PARAGUAY: Amambay:**  
970 Parque Nacional Cerro Corá, -22.650 -56.017, 13 May 1997, A. Wild #AW0563, 1 worker  
971 (ALWC: USNMENT00757625). **Canindeyu:** Reserva Natural del Bosque, Mbaracayù, Jejuimi, -  
972 24.1 -55.5, 19 Sep. 1996, A. Wild #AW0563, 6 workers (ALWC: USNMENT00757584,  
973 00757895). Reserva Natural del Bosque, Mbaracayù, Jejuimi, -24.1 -55.5, 16-23 May 1996,  
974 A.C.F. Costa, 1 worker (ALWC: USNMENT00757868). Reserva Natural del Bosque,  
975 Mbaracayù, Jejuimi, -24.1 -55.5, 11 Oct 1996, A. Wild #AW0334, 1 worker (ALWC:  
976 USNMENT00757893). Reserva Natural del Bosque, Mbaracayù, Jejuimi, -24.1 -55.5, 28 Jan.  
977 1997, A. Wild #AW384, 3 workers (ALWC: USNMENT00757894). Reserva Natural del  
978 Bosque, Mbaracayù, Jejuimi, -24.1 -55.5, 12 Mar. 1997, A. Wild #AW0490, 3 workers (ALWC:

979 USNMENT00757896). **PERU: Huanuco:** 42 km E. Tingo Maria, 1100 m, 10 Dec. 1954, 1  
980 worker (CASC: CASENT0196020). **Madre de Dios:** Reserva Nacional Tambopata, Centro  
981 Sachavacayoc, -12.85583 -69.36194, 19-31 July 2012, Curso de hormigas, 7 workers (ICN:  
982 CAB-120725-1). Reserva Nacional Tambopata, Centro Sachavacayoc, casa camping, -12.85583 -  
983 69.36194, 198 m, 26 July 2012, GSNMBU, 1 worker (ICN: USNMENT00757613). **USA:**  
984 **Lousiana:** East Baton Rouge Par. Baton Rouge, Kennilworth & Perkins Rd. BREC Perkins Park,  
985 03 Apr. 2003, S.T. Dash, 1 worker (WEMC: USNMENT00759023). **VENEZUELA: Bolivar:**  
986 Canaima, Orchid Is, 14 Oct. 1988, W. MacKay #11159, 1 worker (WEMC:  
987 USNMENT00757906).

988  
989 **Diagnosis.** *Brachymyrmex aphidicola* is morphologically similar to *B. australis*, *B. minutus* and  
990 *B. termitophilus*, because all of them typically have smooth and shiny yellowish bodies, their  
991 mesonotum does not bulge dorsally above the pronotum, their eyes are positioned on the cephalic  
992 midline, and the metanotal groove is either absent or narrower than the diameter of the  
993 metathoracic spiracles. However, *B. aphidicola* differs from *B. australis* by having scapes that  
994 surpass the posterior margin of the head by a length longer than the maximal diameter of the eye;  
995 from *B. minutus* by having a well-differentiated mesometanotal suture and by the presence of two  
996 erect hairs on the pronotum and two on the mesonotum; and from *B. termitophilus* by having  
997 scattered pubescence on the gaster.

998  
999 *Lectotype measurements* (mm). HL<sub>1</sub> 0.41; HL<sub>2</sub> 0.29; HL<sub>3</sub> 0.10; HW 0.37; SL 0.39; EL 0.10; WL  
1000 0.41; PnL 0.10; PnW 0.25; ML 0.08; MW 0.18; *Indices* CI 90.48; SI<sub>1</sub> 105.26; SI<sub>2</sub> 133.33; OI<sub>1</sub>  
1001 26.32; OI<sub>2</sub> 23.81.

1002

1003 *Paralectotype measurements* (mm) (n=4). HL<sub>1</sub> 0.43-0.45; HL<sub>2</sub> 0.27-0.31; HL<sub>3</sub> 0.10; HW 0.37; SL  
1004 0.39-0.41; EL 0.10; WL 0.41-0.43; PnL 0.14-0.17; PnW 0.23-0.29; ML 0.08-0.10; MW 0.16-  
1005 0.18; *Indices* CI 82.61-86.36; SI<sub>1</sub> 105.26-110.53; SI<sub>2</sub> 125.00-142.86; OI<sub>1</sub> 26.32; OI<sub>2</sub> 21.74-22.73.

1006  
1007 *Additional material examined measurements* (mm) (n=20). HL<sub>1</sub> 0.36-0.55; HL<sub>2</sub> 0.26-0.37; HL<sub>3</sub>  
1008 0.08-0.13; HW 0.33-0.51; SL 0.32-0.54; EL 0.09-0.12; WL 0.35-0.58; PnL 0.09-0.18; PnW 0.24-  
1009 0.36; ML 0.07-0.12; MW 0.14-0.20; *Indices* CI 84.21-95.65; SI<sub>1</sub> 94.74-112.5; SI<sub>2</sub> 117.14-157.89;  
1010 OI<sub>1</sub> 21.82-28.57; OI<sub>2</sub> 20.62-28.89.

1011  
1012 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin flat or  
1013 slightly concave. Dorsum of the head with scattered appressed hairs. Clypeus with a rounded  
1014 anterior margin and five long, erect hairs of which a single, usually conspicuous hair is near the  
1015 anterior margin, two hairs are in mediolateral position and two more near the toruli; other hairs  
1016 on the clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior  
1017 clypeal margin in oblique anterodorsal view. The scapes surpass the posterior margin of the head  
1018 by a length larger than the maximal diameter of the eye, and typically bear appressed, sometimes  
1019 decumbent, but never erect hairs. Three ocelli usually present, but sometimes inconspicuous.  
1020 Eyes are positioned on the cephalic midline and have 7-10 ommatidia along their maximal  
1021 diameter.

1022 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum. In lateral  
1023 view the mesonotum is not or weakly inflated and does not bulge dorsally above the pronotum.  
1024 Metanotal groove absent or shallow and narrower than the diameter of the metathoracic spiracles.  
1025 Metathoracic spiracles in dorsolateral position, not protruding, and usually touching the  
1026 propodeal suture. Dorsum of propodeum slightly convex and ~1/3th of the length of the

1027 propodeal slope. Propodeal spiracles circular, positioned ventrally of the posterior propodeal  
1028 margin, and slightly posterior of the middle of the propodeal slope. Legs with appressed and  
1029 scattered hairs. Petiole short and inclined forward.

1030 **Gaster.** With scattered pubescence and scattered long suberect hairs, especially along the  
1031 posterior edges of the segments.

1032 **Color and sculpture.** Body smooth and shiny, yellowish.

1033  
1034 **Distribution (Supplementary material Fig. S3).** *Brachymyrmex aphidicola* is widely  
1035 distributed and known from Argentina, the Bermudas, Bolivia, Brazil, Colombia, Costa Rica,  
1036 Ecuador, Guatemala, Guyana, Mexico, Nicaragua, Panama, Paraguay, Peru, the United States, and  
1037 Venezuela.

1038  
1039 **Biology.** Specimens of this species have been found under stones, in rotten wood, on epiphytes  
1040 (USNMENT00757619), and on *Conostegia setosa* (USNMENT00757923). *Brachymyrmex*  
1041 *aphidicola* nests underground or in organic litter, and it appears to be abundant in Paraguayan  
1042 forests (Wild, 2007).

1043  
1044 **Remarks.** Some specimens from Argentina have expanded gasters and Forel (1912a) highlighted  
1045 a worker identified as *B. aphidicola* from Santa Catharina (Brazil) that also has a somewhat  
1046 expanded gaster, but this specimen has not been studied and its identification remains to be  
1047 confirmed. The original description of Forel (1909) indicates *B. aphidicola* to occur in both  
1048 Paraguay and the Bermudas, however, a type locality is not designated. In the type series of  
1049 Forel's collection (NHMG) only specimens from Paraguay are present. Nevertheless, Santschi's  
1050 collection (NHMB) contains a decapitated specimen from Bermudas that is labeled as the type of

1051 *B. aphidicola*. Given this complication, only the specimens from Forel's collection are designated  
1052 here as lectotype and paralectotypes.

1053 We concur with Wild (2007) that *B. heeri* var. *fallax* is a junior synonym of *B. aphidicola*. The  
1054 workers of the type series of this variation have all the diagnostic characteristics of *B. aphidicola*.  
1055 The description of *B. longicornis* var. *hemiops* (Santschi, 1923a) only specifies color and the  
1056 smaller size of the body and eyes of this variation in comparison to *B. longicornis*, however,  
1057 detailed study of the syntype renders it indistinguishable from *B. aphidicola*.

1058

1059

1060 ***Brachymyrmex attenuatus* Santschi NEW STATUS**

1061 **(Fig. 10, supplementary material Fig. S4)**

1062 *Brachymyrmex luederwaldti* st. *attenuatus* Santschi, 1929: 310 (w.). **Lectotype worker** (NHMB:  
1063 USNMENT00757177) and **Paralectotype worker** (NHMB: USNMENT00757177; **here**  
1064 **designated**) 2 workers [examined]. **BRAZIL**, Santa Catarina, Blumenau.

1065

1066 **Diagnosis.** *Brachymyrmex attenuatus* **n. st.** morphologically resembles *Brachymyrmex degener*,  
1067 because both have long scapes that extend beyond the posterior margin of the head, they have  
1068 faint sculpture on the mesosomal dorsum, and dorsally positioned, slightly protruding  
1069 metathoracic spiracles. *Brachymyrmex attenuatus* **n. st.** differs from *B. degener*, however, by  
1070 having a gaster with dense pubescence.

1071

1072 *Lectotype worker measurements* (mm) HL<sub>1</sub> 0.47; HL<sub>2</sub> 0.28; HL<sub>3</sub> 0.11; HW 0.43; SL n.a.; EL  
1073 0.12; WL 0.52; PnL 0.09; PnW 0.31; ML 0.12; MW 0.21; *Indices* CI 90.74; SI<sub>1</sub> n.a; SI<sub>2</sub> n.a; OI<sub>1</sub>  
1074 28.57; OI<sub>2</sub> 23.15.

1075  
1076 *Paralectotype worker measurements* (mm) HL<sub>1</sub> 0.43; HL<sub>2</sub> 0.31; HL<sub>3</sub> 0.11; HW 0.39; SL 0.45; EL  
1077 0.13; WL 0.46; PnL 0.14; PnW 0.29; ML 0.11; MW 0.15; *Indices* CI 91.84; SI<sub>1</sub> 113.33; SI<sub>2</sub>  
1078 145.71; OI<sub>1</sub> 33.33; OI<sub>2</sub> 25.51.

1079  
1080 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1081 concave. Dorsum of head with scattered appressed hairs. Clypeus with a rounded anterior margin  
1082 and five long, erect hairs of which a single, usually conspicuous hair is near the anterior margin,  
1083 two hairs are in mediolateral position and two more near the toruli; other hairs on the clypeus are  
1084 substantially shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin  
1085 in oblique anterodorsal view. The scapes extend beyond the posterior cephalic margin by a length  
1086 approximately equal to the maximum diameter of the eye (and not more than 1.5× this diameter),  
1087 and have decumbent hairs. A single central ocellus seems to be present but is inconspicuous.  
1088 Eyes are positioned on the cephalic midline and have 7-9 ommatidia along their maximal  
1089 diameter.

1090 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum,  
1091 sometimes with additional suberect hairs, mainly on pronotum. In lateral view, the mesonotum is  
1092 inflated, but it does not bulge dorsally above the pronotum. Metanotal groove deep and wider  
1093 than the diameter of the metathoracic spiracles. Metathoracic spiracles fully dorsal in position,  
1094 slightly protruding and not touching the mesometanotal or propodeal sutures. Dorsum of  
1095 propodeum slightly convex and shorter than the propodeal slope. Propodeal spiracles circular,  
1096 positioned ventrally of the posterior propodeal margin, and slightly posteriorly of the middle of  
1097 the propodeal slope. Legs with appressed and scattered hairs. Petiole short and inclined forward.  
1098 **Gaster.** With appressed dense pubescence and several scattered long erect hairs.

1099 **Color and sculpture.** Body overall smooth and shiny, with faint sculpture on mesosomal dorsum.

1100 Body uniformly light or dark brownish, but the legs and antennae are yellowish.

1101

1102 **Distribution (Supplementary material Fig. S4).** *Brachymyrmex attenuatus* is currently only  
1103 known from Brazil.

1104

1105 **Biology.** Unknown.

1106

1107 **Remarks.** Here we designate the lectotype as the specimen closest to the pin

1108 (USNMENT00757177); the paralectotype has lighter brownish color in comparison with the

1109 lectotype. This species was described by Santschi (1929) as a subspecies of *B. luederwaldti* that

1110 has a smaller body size in comparison with *B. luederwaldti*. Additionally, *B. attenuatus* has

1111 weaker sculpture, a shinier body, especially on pronotum, a more concave posterior cephalic

1112 border, smaller eyes and a more convex mesonotum than *B. luederwaldti*. All these

1113 characteristics are somewhat subjective, because they represent differences in intensity rather

1114 than state and as such it is difficult to determine clear boundaries. A more marked difference is

1115 the presence of pubescence on the gaster, which is clearly present in *B. attenatus n. st.*, but absent

1116 in *B. luederwaldti*. The presence or absence of pubescence is an important trait to delimit other

1117 *Brachymyrmex* species, and hence we raise *B. attenatus n. st.* to species level.

1118

1119

1120 *Brachymyrmex australis* Forel

1121 (Fig. 11, supplementary material Fig. S5)

1122



1123 *Brachymyrmex minutus* r. *australis* Forel, 1901b: 302 (w.). **Lectotype worker** (MHNG:  
1124 USNMENT00757156) and **paralectotype worker** (MHNG: USNMENT00758102; **here**  
1125 **designated**): 2 workers [examined]. **BRAZIL**, Rio Grande do sul, Pelotas. Raised to species:  
1126 Santschi (1922: 260). See also: Santschi (1923a: 662).  
1127 = *Brachymyrmex australis* var. *curta* Santschi, 1922: 260 (w.q.m.). (NHMB:  
1128 USNMENT00757700 - 00757703, 00758069 - 00758071): 17 workers, 2 queens, 7 males  
1129 [examined]. **ARGENTINA**: Cordoba, Alta Gracia [Also described as a new variety in Santschi  
1130 (1923a: 663)]. **n. syn.**  
1131 = *Brachymyrmex longicornis* Forel, 1907: 9 (w.). (MHNG: USNMENT00757144): 2 workers  
1132 [examined]. **BRAZIL**: Porto Alegre. Other relevant descriptions: Forel (1912a: 62) (q.).  
1133 (MHNG: USNMENT00757145): 1 queen [examined]. **BRAZIL**: São Paulo. **n. syn.**  
1134  
1135 **Additional material examined. ARGENTINA: Santa Cruz**: O. Bondensköld, 3 workers  
1136 (MCZC: USNMENT00759000). **Tucumán**: 11 km N Tafí Viejo, -26.63333 -65.23333, 820 m,  
1137 01 Feb. 1995, P.S. Ward #12826-25, 3 workers (PSWC: USNMENT00757628). **BAHAMAS**:  
1138 Exumas: unnamed cay, 175 m N of NW tips of Obrien's Cay, 05 May 1995, J. W. Morrison 321-  
1139 92, 2 workers (PSWC: USNMENT00758991). **BRAZIL: Bahia**: Ilheus, CEPEC, Antonio 455E,  
1140 2 workers, 2 queens (CPDC: USNMENT00757922). Lençóis, Chap. Diamantina, -12.55 -41.38,  
1141 25 Mar. 2001, Santos, J.R.M. dos, 2 workers, 1 queen (CPDC: USNMENT00757909). **Goiás**:  
1142 Ouro verde, Faz Boa Vista, -16.29847 -49.21183, 01-07 July 2005, R.R. Silva & R.M. Feitosa, 4  
1143 workers (ICN: MZSP122). **Minas Gerais**: Alfenas, Porto, 06 Oct. 2011, I.A. Dos Santos, 6  
1144 workers (ICN: USNMENT00759048, 00759049). **Pará**: Melgaço, Caxiuanã, ECFPn, -1.73584 -  
1145 51.48762, II: transecto (4-600), 23-25 Oct. 2005, Equipe A.Y. Harada, 3 workers (MPEG:  
1146 AYH036). **Rio de Janeiro**: Teresópolis, P.N. Serra dos Orgãos, -22.45333 -42.99806, 23-28

1147 Nov. 1999, Dietz, Silva & Rocha, 6 workers (ICN: MZSP130). **Santa Catarina:** São Bento do  
1148 Sul, APA Rio Vermelho, -26.36417 -42.99806, 30 Mar.–04 Apr. 2001, R. Silva & Eberhardt, 5  
1149 workers (ICN: MZSP134 135). **São Paulo:** Ilha da Vitória, 16-27 Mar. 1964, Exp. Dep. Zool.  
1150 3592, 5 workers, 1 queen (MCZC: USNMENT00757932); Itirapina, 10 Feb. 2009, S. Sendoya,  
1151 20 workers (ICN: USNMENT00759046); Jundiai, Sierra Do Japi, Apr. 2009, S. Diniz, 3 workers  
1152 (ICN: USNMENT00759043); Piedade, Floresta Atlantica, "Cristo", Mar. 2010, G. Bieber, 3  
1153 workers (ICN: USNMENT00759045); Tapirai, -24.03208 -47.46556, 08-14 Jan. 2001, R. Silva  
1154 & Eberhard, 4 workers (ICN: MZSP170). **COLOMBIA: Bolivar:** Zambrano, Hacienda  
1155 Monterrey, 9.617 -74.900, 9-75 m, 04 Aug. 1992, A. Molano, 3 workers (ICN:  
1156 USNMENT00757898). **Caldas:** Municipio Aranzazu, Vereda Alegrias, Finca Betania, La  
1157 Esperanza, 5.29811 -75.49047, 1990 m, L.E. Franco & J. Cruz, 2 workers (IAvH: IAvH27305);  
1158 Municipio Aranzazu, Vereda Alegrias, Finca Villa Rosita, 5.30603 -75.48492, 1825 m, 06-08  
1159 Aug. 2003, L.E. Franco & J. Cruz, 1 worker (IAvH: IAvH25467). **Caquetá:** PNN Serranía de  
1160 Chiribiquete, Cuñané-Anuí, 26 Feb. 2001, 2 workers (IAvH: IAvH-E71471). **Cauca:** El  
1161 Hortigal, Holanda, Mar. 2002, Valderrama, 1 worker (ICN: USNMENT00757937).  
1162 **Cundinamarca:** Fusagasugá, 08 Dec. 1975, W. & E. MacKay, 2 workers (WEMC:  
1163 USNMENT00757907). **Guajira:** Serrania de Macuira, 6-8 km S Nazareth, 70-200 m, 13 June  
1164 1957, W.L. Brown & Kugler, 2 workers (IAvH: IAvH-E74171). **Huila:** 21 Km W La Plata  
1165 Gallego, 03 Jan. 1984, W.P. MacKay #7153, 6 workers (WEMC: USNMENT00757623,  
1166 00757624, 00759014); Neiva, 05 Dec 1975, W. & E. MacKay, 2 workers (WEMC:  
1167 USNMENT00757620). **Magdalena:** PNN Tayrona, Cañaveral, 11.33 -74.03, 30 m, 20-27 Apr.  
1168 2000, C. Sarmiento, 1 worker (IAvH: USNMENT00759056). **Meta and Cundinamarca**  
1169 **border:** 28 Dec. 1975, W. & E. Mackay, 4 workers (WEMC: USNMENT00757917, 00757939).  
1170 **Meta:** Puerto Gaitán, 21 Dec. 1975, W. & E. MacKay #783, 2 workers (WEMC:

1171 USNMENT00757672). **Quindío:** Buenavista, Vereda El Infierno, Finca Guadalajara, 4.3767 -75.  
1172 7694, 1160 m, 16 Nov. 1999, E. Gonzalez, 2 workers (IAvH: IAvH-E74165); Municipio  
1173 Filandia, Vereda Cruces, Finca Los Micos, 4.70424 75.65917, 12-13 July 2002, E. Jimenez &  
1174 L.E. Franco, 1 worker (IAvH: IAvH27232). **Risaralda:** Apia La Felisa, Cafetal de sol (S-I), 5.13  
1175 -75.95, 1480 m, 29 Oct. 2001, 1 worker (IAvH: IAvH-E74174). **Valle del Cauca:** Dagua, 07 Jan.  
1176 1976, W. & E. MacKay, 5 workers (WEMC: USNMENT00758993); Medio Calima, 24 June  
1177 1989, E. MacKay #11740 #11743 #11744, 2 workers (WEMC: USNMENT00757738, 00759005,  
1178 00757908, 00759012). **Vichada:** Municipio Cumaribo, Corregimiento Santa Rita, PNN El  
1179 Tuparro, 5.3075 -67.9500, 135 m, 14-16 Feb. 2004, I. Quintero & E. Gonzalez, 2 workers (IAvH:  
1180 USNMENT00759057). **COSTA RICA: Guanacaste:** Provincia Maritza field Station, 03 May  
1181 1995, R. Anderson #17716, 3 workers (WEMC: USNMENT00757671). **Puntarenas:** Pen. Osa.  
1182 Par. Nat. Corcovado, Llorona, 8.58 -83.70, 5 m, 30 Dec. 1981, J. Longino, 1 worker (JTLC:  
1183 JTLC000005948). **CUBA:** Holguín: 6 km S Yamanigüey, 20.55 -74.73, 25 m, 23 Aug. 2001,  
1184 P.S. Ward #14437-19, 3 workers (PSWC: USNMENT00757919). **DOMINICAN REPUBLIC:**  
1185 16 km ENE Pedernales, 18.11667 -71.62361, 800 m, 9 Sep. 1992, P.S. Ward #11726-22, 3  
1186 workers (PSWC: USNMENT00757959); Prov. La Vega, Jarabacoa to El Rio, shady ravine, 80-  
1187 1500 m, Feb. 1975, W.L. & D.E. Brown, 2 workers (MCZC: USNMENT00757736).  
1188 **ECUADOR:** Guayaquil, 10 m, Dec. 1997, Forero, 2 workers (IAvH: USNMENT00759054).  
1189 **FRENCH GUIANA:** Petit Saut Basse Vie II/III, 2001, A. Dejean, 1 worker (CPDC:  
1190 USNMENT00757734); Reserve Naturel de Nouragues- Inselbery forest, Oct. 2009, Sara Groc, 4  
1191 workers (ICN: USNMENT00759033). **GUATEMALA: El progreso:** 5 km W El Rancho,  
1192 14.91667 -90.06666, 400 m, 17 Nov. 2003, A.L. Wild #AW2002, 3 workers (ALWC:  
1193 USNMENT00757957). **Suchitepéquez:** Cocales (Mpio. San Antonio), 14.39206 -91.19347, 242  
1194 m, 31 Aug. 2004, W & E. MacKay #20820, 1 worker (WEMC: USNMENT00758995).

1195 **GUYANA:** Demerara-Mahaica: Wales, 6.67 -58.25, 50 m, 23 Jan. 1981, 2 workers, 1 queen  
1196 (JTLC: JTLC000005920); Rupununi, Karanambo, 3.75 -59.3, 100 m, 01 Jan. 1981, 1 worker  
1197 (JTLC: JTLC000005926). **MAURITIUS:** Mgne. Brise Fer, -20.37 57.43, 600 m, 07 May 1989,  
1198 P.S. Ward #10518-2, 3 workers (PSWC: USNMENT00757934). **MEXICO: Chiapas:** 10 km S  
1199 Palenque, 30 May 1988, 31 May 1988, W. MacKay #10611 #10613, 8 workers (WEMC:  
1200 USNMENT00757588, 00757677, 00757678). **Veracruz:** Los Tuxtlas, 26 July 1974, R.L. Jeanne,  
1201 1 worker (MCZC: USNMENT00757735). **Yucatan:** 25.7 km E Progreso, 12 Apr. 1982, Smalley  
1202 Thien & Bradburn, 1 worker (MCZC: USNMENT00757618). **PARAGUAY: Boquerón:** Enciso,  
1203 -21.20 -61.67, 03-06 Nov 2001, M. LePonce & T. Delsinne, 2 workers (ALWC:  
1204 USNMENT00757904). **Central:** Areguá, CHP center, -25.30 -57.38, 01 Oct. 1995, A. Wild  
1205 #AW 0059, 1 worker (ALWC: USNMENT00757905). **Presidente Hayes:** Monte Lindo, -  
1206 23.86667 -58.46667, 800-1500 m, Feb. 1975, W.L. & D.E. Brown, 2 workers (MCZC:  
1207 USNMENT00757736). **PERU: Madre de Dios:** Reserva Nacional Tambopata, Centro  
1208 Sachavacayoc, Centre, -12.85583 -69.36194, 209 m, 19-31 Jul 2012, R. Feitosa, 1 worker (ICN:  
1209 USNMENT00757611). **URUGUAY:** Montevideo, L. Pastre, 1 worker (CPDC:  
1210 USNMENT00757684).

1211

1212 **Diagnosis.** *Brachymyrmex australis* is very similar in morphology to *B. aphidicola*, *B. minutus*  
1213 and *B. termitophilus*, because all these species have a mesonotum that does not bulge dorsally  
1214 above the pronotum in lateral view, their bodies are smooth, shiny and yellowish, and their eyes  
1215 are positioned on the cephalic midline. However, *B. australis* differs from *B. aphidicola* by  
1216 somewhat shorter scapes, although they still reach to the posterior margin of the head or surpass  
1217 it by a length equal to or smaller than the maximal diameter of the eye; it differs from *B. minutus*  
1218 by having a well-marked mesometanotal suture and two erect hairs on the pronotum and two on

1219 the mesonotum; finally, it has scattered pubescence on the gaster whereas that of *B. termitophilus*  
1220 is dense.

1221

1222 *Lectotype worker measurements* (mm) HL<sub>1</sub> 0.37; HL<sub>2</sub> 0.27; HL<sub>3</sub> 0.10; HW 0.29; SL 0.29; EL  
1223 0.10; WL 0.35; PnL n.a.; PnW 0.23; ML 0.08; MW 0.17; *Indices* CI 78.95; SI<sub>1</sub> 100.00; SI<sub>2</sub>  
1224 107.14; OI<sub>1</sub> 33.33; OI<sub>2</sub> 26.31.

1225

1226 *Additional material examined measurements* (mm) (n=13). HL<sub>1</sub> 0.32-0.54; HL<sub>2</sub> 0.21-0.38; HL<sub>3</sub>  
1227 0.08-0.15; HW 0.29-0.53; SL 0.26-0.48; EL 0.08-0.14; WL 0.29-0.55; PnL 0.08-0.20; PnW 0.21-  
1228 0.32; ML 0.06-0.11; MW 0.14-0.20; *Indices* CI 82.61-97.09; SI<sub>1</sub> 89.09-104.54; SI<sub>2</sub> 114.29-  
1229 135.29; OI<sub>1</sub> 15.38-30.91; OI<sub>2</sub> 19.35-28.33.

1230

1231 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1232 concave. Dorsum of the head with scattered appressed hairs. Clypeus with a rounded anterior  
1233 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
1234 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
1235 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
1236 margin in oblique anterodorsal view. The scapes extend beyond the posterior margin of the head  
1237 by a length equal to or smaller than the maximal diameter of the eye; they typically have  
1238 appressed, sometimes decumbent, but never erect hairs. Three inconspicuous ocelli. Eyes on the  
1239 cephalic midline, with 7-14 ommatidia along their maximal diameter.

1240 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum. The  
1241 mesonotum is not inflated and it does not bulge dorsally above the pronotum in lateral view.  
1242 Metanotal groove absent or shallow and narrower than the diameter of the metathoracic spiracles.

1243 Metathoracic spiracles in dorsolateral position, not protruding, and usually touching both the  
1244 mesometanotal and propodeal sutures. Dorsum of the propodeum flat or weakly convex and  
1245 ~1/3th of the length of the propodeal slope. Propodeal spiracles circular, positioned ventrally of  
1246 the posterior propodeal margin slightly posterior of the middle of the propodeal slope. Legs with  
1247 appressed and scattered hairs. Petiole short and inclined forward.

1248 **Gaster.** With scattered pubescence and scattered long suberect hairs.

1249 **Color and sculpture.** Body smooth and shiny, uniform yellowish in color.

1250

1251 **Distribution (Supplementary material Fig. S5).** *Brachymyrmex australis* is known from  
1252 Argentina, the Bahamas, Brazil, Colombia, Costa Rica, Cuba, the Dominican Republic, Ecuador,  
1253 French Guiana, Guatemala, Guyana, Mexico, Paraguay, Peru and Uruguay. It has also been  
1254 introduced in Mauritius.

1255

1256 **Biology.** Some specimens have been found under stones and among leaf litter. The type material  
1257 of *B. longicornis* (here considered a junior synonym of *B. australis*) was collected from orchids.

1258

1259 **Remarks.** Forel (1901b) described *B. australis* as a variety of *B. minutus* but did not indicate  
1260 diagnostic traits to separate it from typical *B. minutus*. Subsequently, Santschi (1922) raised *B.*  
1261 *australis* to species level, described a new variety to it (*B. australis* var. *curta*), again without  
1262 clear motivation, although he pointed out morphological similarities between the males of *B.*  
1263 *australis* and *B. fiebrigi*. Later, Santschi (1923a) indicated that *B. australis* has a conspicuous  
1264 mesometanotal suture, and *B. minutus* does not, and that *B. australis* var. *curta* is smaller and  
1265 shinier than typical *B. australis*.

1266 The type material of *B. australis* var. *curta* and *B. longicornis* share the same diagnostic traits and  
1267 display only minor variation in body size and the length of the scapes compared with *B. australis*.  
1268 Most of the specimens of *B. longicornis* we studied are yellowish, but one was brownish, and the  
1269 nature of this variation remains to be documented. In any case, Forel (1907) originally described  
1270 *B. longicornis* as “yellowish brown”. Considering our observations *B. australis* var. *curta* and *B.*  
1271 *longicornis* are synonymized here to *B. australis*.  
1272 Santschi (1923a) identified a specimen (1 worker, NHMB) from a termite nest in Sao Leopoldo,  
1273 Rio Grande do Sul, Brazil, i.e. the type locality of *B. termitophilus* (which has also been recored  
1274 from termite nests), as *B. australis* but this specimen has the diagnosis traits of *B. fiebrigi*;  
1275 additional specimens (2 workers, NHMB) from Uruguay, Nueva Helvetia (Mme.v. Steiger) that  
1276 he identified as *B. australis* var. *curta* have the diagnostic traits of *B. termitophilus*.

1277

1278

1279 ***Brachymyrmex bahamensis* NEW SPECIES**

1280 **(Fig. 12, supplementary material S6)**

1281 **Holotype worker** (MCZC: USNMENT00757689) and **Paratype workers** (MCZC:  
1282 USNMENT00757689, PSWC: USNMENT00757726): 5 workers. **BAHAMAS:** Exuma,  
1283 unnamed cay, 175 m S of Staniel Cay, 21 May 1990, L.W. Morrison 101-90.

1284

1285 **Additional material examined. BAHAMAS,** Andros Island, May-June 1904, col. W.M.  
1286 Wheeler, 13 workers, 2 queens (MCZC: USNMENT00757690).

1287

1288 **Etymology:** In reference of the type locality.

1289

1290 **Diagnosis.** *Brachymyrmex bahamensis* resembles *B. termitophilus* because both species have  
1291 scapes that are surpass the posterior margin of the head by a length smaller than the maximal  
1292 diameter of the eye, their mesonotum does not bulge dorsally above the pronotum, they have  
1293 erect or suberect hairs on the mesosoma, a gaster with dense pubescence, and yellowish body  
1294 color. However, the unique feature of *B. bahamensis* is that it has approximately 6 erect hairs on  
1295 the pronotum and two on the mesonotum that are very long, i.e. about twice the length of the  
1296 maximal diameter of the eye. *Brachymyrmex bahamensis* also resembles *B. heeri*, but this latter  
1297 species has a mesonotum that bulges out dorsally above the pronotum.

1298  
1299 *Holotype measurements* (mm). HL<sub>1</sub> 0.46; HL<sub>2</sub> 0.31; HL<sub>3</sub> 0.13; HW 0.41; SL 0.38; EL 0.09; WL  
1300 0.45; PnL 0.13; PnW 0.29; ML 0.09; MW 0.19; *Indices* CI 88.46; SI<sub>1</sub> 91.30; SI<sub>2</sub> 120.00; OI<sub>1</sub>  
1301 21.74; OI<sub>2</sub> 28.85.

1302  
1303 *Paratype measurements* (mm) (n=2). HL<sub>1</sub> 0.47-0.48; HL<sub>2</sub> 0.32; HL<sub>3</sub> 0.14; HW 0.43-0.44; SL  
1304 0.39-0.40; EL 0.09-0.10; WL 0.47-0.49; PnL 0.13-0.16; PnW 0.30-0.31; ML 0.09-0.12; MW  
1305 0.21; *Indices* CI 90.57-90.74; SI 91.67-91.84; SI<sub>2</sub> 122.22-125.00; OI<sub>1</sub> 22.45-25; OI<sub>2</sub> 27.78-28.30.

1306  
1307 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1308 concave. Dorsum of head with appressed hairs and with two rows of erect hairs. Clypeus with a  
1309 rounded anterior margin and five long, erect hairs of which a single, usually conspicuous hair is  
1310 near the anterior margin, two hairs are in mediolateral position and two more near the toruli;  
1311 other hairs on the clypeus are markedly shorter and appressed or decumbent. Toruli surpassing  
1312 the posterior clypeal margin in oblique anterodorsal view. The scapes surpass the posterior  
1313 cephalic margin by a length smaller than the maximal diameter of the eye; they typically have



1314 appressed, sometimes decumbent but never erect hairs. Ocelli absent. Eyes are positioned on the  
1315 cephalic midline and have 8–9 ommatidia along their maximal diameter.

1316 **Mesosoma.** Approximately 6 long, erect hairs on the pronotum and two on the mesonotum, each  
1317 having a length of about twice the maximal diameter of the eye. In lateral view, the mesonotum is  
1318 not inflated and it does not bulge dorsally above the pronotum. Metanotal groove absent or  
1319 shallow and narrower than the diameter of the metathoracic spiracles. Dorsum of the propodeum  
1320 is flat and ~1/3th of the length of the propodeal slope. Metathoracic spiracles in dorsolateral  
1321 position, not protruding, and usually touching the propodeal suture, but not the mesometanotal  
1322 suture. Propodeal spiracles circular, positioned ventrally of the posterior propodeal margin,  
1323 posterior of the middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole  
1324 short and inclined forward.

1325 **Gaster.** With dense pubescence and several scattered conspicuous long erect hairs.

1326 **Color and sculpture.** Body usually smooth and shiny, yellowish.

1327

1328 **Distribution (Supplementary material S6).** Currently exclusively known from the Bahamas.

1329

1330 **Biology.** Unknown.

1331

1332 **Remarks.** The holotype is located at the top of pin USNMENT00757689, with the two paratypes  
1333 below.

1334

1335

1336 ***Brachymyrmex bicolor* NEW SPECIES**

1337 **(Fig. 13, supplementary material S7)**

1338 **Holotype worker** (USNM: CASENT0615272) and **Paratypes** (USNM: CASENT0615274  
1339 (putative worker-queen intercaste); CASENT0615277, 0615294, 0617077 (destroyed) (3  
1340 workers); ANTWEB: CASENT0615273 (putative worker-queen intercaste), 0615276 (1 worker),  
1341 0615292 (1 queen)): 5 workers, 1 queen, 2 putative worker-queen intercastes. **HONDURAS:**  
1342 **Comayagua:** PN Cerro Azul Meambar, 14.87092, -87.89917, 1120 m, 20 May 2010,  
1343 LLAMA#Wa-C-04-1-31.

1344

1345 **Etymology:** The epithet *bicolor* reflects the conspicuous body coloration with black head and  
1346 gaster and yellow mesosoma.

1347

1348 **Diagnosis.** The conspicuous color pattern allows distinguishing *B. bicolor* from any other  
1349 *Brachymyrmex* species.

1350

1351 *Holotype measurements* (mm). HL<sub>1</sub> 0.43; HL<sub>2</sub> 0.30; HL<sub>3</sub> 0.10; HW 0.38; SL 0.44; EL 0.11; WL  
1352 0.48; PnL 0.12; PnW 0.28; ML 0.10; MW 0.16; *Indices* CI 88.78; SI<sub>1</sub> 114.94; SI<sub>2</sub> 147.06; OI<sub>1</sub>  
1353 27.59; OI<sub>2</sub> 22.45.

1354

1355 *Paratype measurements* (mm). HL<sub>1</sub> 0.43; HL<sub>2</sub> n.a.; HL<sub>3</sub> 0.10; HW 0.41; SL 0.45; EL 0.11; WL  
1356 0.48; PnL 0.12; PnW 0.28; ML 0.11; MW 0.15; *Indices* CI 94.90; SI<sub>1</sub> 109.68; SI<sub>2</sub> n.a.; OI<sub>1</sub> 25.81;  
1357 OI<sub>2</sub> 22.45.

1358

1359 **Worker description. Head.** Slightly longer than wide in full face view; posterior cephalic  
1360 margin slightly convex. Dorsum of the head with appressed hairs. Clypeus with a rounded  
1361 anterior margin and five long, erect hairs of which a single, usually conspicuous hair is near the

1362 anterior margin, two hairs are in mediolateral position and two more near the toruli; other hairs  
1363 on the clypeus are clearly shorter and appressed or decumbent. Toruli surpassing the posterior  
1364 clypeal margin in oblique anterodorsal view. The scapes surpass the posterior margin of the head  
1365 by a length larger than the maximal diameter of the eye and have appressed pubescence. Three  
1366 conspicuous ocelli. Eyes are positioned on the cephalic midline and have 8 ommatidia along their  
1367 maximal diameter.

1368 **Mesosoma.** Without erect hairs and in lateral view approximately hour-glass shaped (this  
1369 condition is absent in the presumed intercast) with a constriction between the bulging  
1370 promesonotum and propodeum. In lateral view, the mesonotum is anteriorly inclined, but it does  
1371 not bulge dorsally above the pronotum. Metanotal groove present and wider than the diameter of  
1372 the metathoracic spiracles. Dorsum of the propodeum is convex and shorter than the propodeal  
1373 slope. Metathoracic spiracles in dorsal position, not protruding, not touching any sutures.  
1374 Propodeal spiracles circular, positioned just ventrally of the posterior propodeal margin slightly  
1375 posterior of the middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole  
1376 short and inclined forward.

1377 **Gaster.** With scarce pubescence and several scattered long erect hairs at the edge of the segments.

1378 **Color and sculpture.** Body smooth and shiny, with a conspicuous bicolored pattern. The head  
1379 and gaster are blackish in color, however, the mandibles, the labial and maxillary palps, the bulbi  
1380 and bases of the antennae, the terminal antennomeres and hairs are conspicuously yellow in  
1381 color. Additionally, the mesosoma and legs are yellowish, with the tibia of the second and third  
1382 pairs of legs being dark brownish, like most of the scape.

1383

1384 **Intercaste description.** The presumed worker-queen intercaste differs from the worker mainly  
1385 by its larger body size, the shape of the mesosoma in lateral view, and its dense pubescence on

1386 the gaster. The dorsum of the head bears two rows of erect hairs. Eyes have around 10 ommatidia  
1387 along their maximal diameter; the promesonotum is bluntly angular, with the mesonotum being  
1388 not inflated or bulging out dorsally above the pronotum in lateral view, mesonotum in dorsal  
1389 view posteriorly extended along the midline. Metanotal groove absent. Dorsum of the propodeum  
1390 is flat and shorter than the length of the propodeal slope. Metathoracic spiracles in dorsolateral  
1391 position, not protruding, not touching any sutures. Gaster with dense pubescence and several  
1392 scattered long erect hairs at the edges of the segments.

1393

1394 **Distribution (Supplementary material S7).** Currently exclusively known from Honduras.

1395

1396 **Biology.** Specimens were collected from leaf litter in cloud forest.

1397

1398 **Remarks.** Further comments on the putative worker-queen intercaste in *Brachymyrmex* are  
1399 provided in the remarks on *B. giardi*.

1400

1401

1402 *Brachymyrmex bonariensis* Santschi NEW STATUS

1403 (Fig. 14, supplementary material Fig. S8)

1404 *Brachymyrmex constrictus* st. *bonariensis* Santschi, 1933: 122 (w.). **Lectotype worker** (NHMB:

1405 USNMENT00757706) and **paralectotype worker** (NHMB: USNMENT00757705; **here**

1406 **designated**): 2 workers [examined]. **ARGENTINA: Buenos Aires:** Buenos Aires, 08 Mar. 1803,

1407 C. Bruch. **n. st.**

1408

1409 **Diagnosis.** *Brachymyrmex bonariensis* n. st. resembles *B. admotus* because they both have  
1410 scapes that surpass the posterior margin of the head, a pair of simple erect hairs between the  
1411 dorsal methathoracic spiracles, a wide metanotal groove, and a gaster with scarce pubescence.  
1412 However, in *B. bonariensis* n. st. the head and mesosoma are light brownish in color, and the  
1413 gaster is darker, whereas the body of *B. admotus* is uniformly colored. The scapes of *B.*  
1414 *bonariensis* are shorter than those of *B. admotus* and surpass the posterior margin of the head  
1415 with a length approximately equal to the maximal diameter of the eyes. The metathoracic  
1416 spiracles of *B. bonariensis* are furthermore positioned more laterally and are not protruding. Like  
1417 *B. admotus*, *B. bonariensis* could be confused with *B. cavernicola* because this latter species also  
1418 has a pair of erect hairs between the methathoracic spiracles, however in *B. cavernicola* these  
1419 hairs are very thick, and they are darker in color than the body.

1420  
1421 *Lectotype measurements* (mm) HL<sub>1</sub> 0.53; HL<sub>2</sub> 0.36; HL<sub>3</sub> 0.15; HW 0.48; SL 0.49; EL 0.13; WL  
1422 0.50; PnL 0.14; PnW 0.31; ML 0.12; MW 0.20; *Indices* CI 90.00; SL<sub>1</sub> 102.78; SL<sub>2</sub> 134.55; OI<sub>1</sub>  
1423 22.78; OI<sub>2</sub> 27.50.

1424  
1425 *Paralectotypes measurements* (mm). HL<sub>1</sub> 0.53; HL<sub>2</sub> 0.35; HL<sub>3</sub> 0.15; HW 0.50; SL 0.50; EL 0.13;  
1426 WL 0.53; PnL 0.18; PnW 0.34; ML 0.11; MW 0.18; *Indices* CI 92.59; SL<sub>1</sub> 101.33; SL<sub>2</sub> 143.40;  
1427 OI<sub>1</sub> 26.67; OI<sub>2</sub> 27.16.

1428  
1429 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1430 concave. Dorsum of head with scattered, appressed hairs and usually two rows of erect hairs.  
1431 Clypeus with a rounded anterior margin and five long, erect hairs of which a single, usually  
1432 conspicuous hair is near the anterior margin, two hairs are in mediolateral position, and two more

1433 near the toruli; other hairs on the clypeus are markedly shorter and appressed or decumbent.

1434 Toruli surpassing the posterior clypeal margin in oblique anterodorsal view. The scapes surpass  
1435 the posterior margin of the head by a length approximately equal to the maximal diameter of the  
1436 eyes. Ocelli appear to be present but are inconspicuous. Eyes are positioned on the cephalic  
1437 midline and have 8-9 ommatidia along their maximal diameter.

1438 **Mesosoma.** With two erect hairs on the pronotum and two on the mesonotum. In lateral view, the  
1439 mesonotum is somewhat inflated, but it does not bulge dorsally above the pronotum. Metanotal  
1440 groove wider than the diameter of the metathoracic spiracles. Metathoracic spiracles in  
1441 dorsolateral position, not protruding, but touching the propodeal suture. Between the metathoracic  
1442 spiracles two thin erect hairs are present, but they are shorter than those on the pronotum and  
1443 mesonotum. Dorsum of the propodeum flat and  $\sim 1/3$ th of the length of the propodeal slope.  
1444 Propodeal spiracles circular, positioned on the posterior propodeal margin slightly posterior of  
1445 the middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole short and  
1446 inclined forward.

1447 **Gaster.** With scattered pubescence, and scattered suberect hairs, mainly along the edges of the  
1448 segments.

1449 **Color and sculpture.** Body overall smooth and shiny, except for the slightly imbricate sculpture  
1450 on the dorsum of the mesosoma in some specimens. Head and mesosoma light brown, gaster  
1451 darker in color.

1452

1453 **Distribution (Supplementary material Fig. S8).** Exclusively known from Argentina.

1454

1455 **Biology.** Unknown.

1456

1457 **Remarks.** *Brachymyrmex bonariensis* was first described by Santschi (1933) as a variety of *B.*  
1458 *constrictus* because it has the thorax less strongly strangled, a little wider head and more concave  
1459 posterior margin of the head in comparison with *B. constrictus*. However, in our opinion *B.*  
1460 *bonariensis* is morphologically very different than *B. constrictus*: it has a mesonotum that does  
1461 not bulge dorsally above the pronotum in lateral view whereas that of *B. constrictus* does; *B.*  
1462 *constrictus* does not have erect hairs between the metathoracic spiracles and moreover it has a  
1463 uniformly dark brownish body. As mentioned before, *B. bonariensis* resembles *B. admotus* more  
1464 closely (see diagnosis).

1465

1466

1467 ***Brachymyrmex brasiliensis* Ortiz & Fernández**

1468 **(Fig. 15, supplementary material Fig. S9)**

1469 *Brachymyrmex brasiliensis* Ortiz and Fernández, 2014: 22, Figs. 19, 20, 21 (w). **Holotype**  
1470 **worker** (MZSP: USNMENT00757748) and **paratype worker** (UFUC: USNMENT00757833): 2  
1471 workers. **BRAZIL: Rio de Janeiro:** Nova Friburgo, Fazenda Barreto, -22.161242 -42.524302,  
1472 1068 m, 11–12 June 2011, T.M.S. Mesquita

1473

1474 *Additional material examined.* **BRAZIL: Goiás:** Anapolis, 12 Feb. 1958, W. Kempf, 1 worker  
1475 (MZSP: USNMENT00757820). **ECUADOR: Zamora:** Chinchipe, -3.98228 -79.083528, 1  
1476 worker (RBINS: 4048410).

1477

1478 **Diagnosis.** *Brachymyrmex brasiliensis* differs from other *Brachymyrmex* species by having  
1479 tumuliform metathoracic spiracles, in combination with a smooth and shiny gaster as well as an  
1480 opaque head and mesosoma.

1481

1482 **Description.** See Ortiz and Fernández (2014).

1483

1484

1485 *Brachymyrmex bruchi* Forel

1486 (Figs. 16, 17, supplementary material Fig. S10)

1487 *Brachymyrmex bruchi* Forel, 1912a: 64 (w.m.). **Lectotype worker** (MHNG:

1488 USNMENT00757159), and **paralectotype workers, males** (MHNG: USNMENT007157-

1489 007159, 00758104, 00758149-00758181; **here designated**): 21 workers, 3 males [examined].

1490 **ARGENTINA: Catamarca:** Aconquija, Filo blanco, 4300 m, Bruch. Santschi (1929: 309) (q.).

1491 = *Brachymyrmex bruchi* var. *rufipes* Forel, 1912a: 65 (w.). (MHNG: USNMENT00757160,

1492 00757161): 3 workers [examined]. **ARGENTINA: Catamarca:** Huasan; synonymy by Quirán et

1493 al. (2004: 279). See also: Santschi (1923a: 660).

1494 = *Brachymyrmex giardi* var. *nitida* Santschi, 1922: 261 (w.). (NHMB: USNMENT00757182): 1

1495 worker [examined]. **CHILE:** Los Lagos, Petrohué, 1922, Schiller. Snelling and Hunt (1975: 114)

1496 as junior synonym of *Brachymyrmex giardi*. **n.syn.**

1497 = *B. laevis* var. *andina* Santschi, 1923a: 659 (w.). (NHMB: USNMENT00758161, 00757186,

1498 00757187; MHNG: USNMENT00758129): 16 workers [examined]. **ARGENTINA: Jujuy:**

1499 Puna, 4000 m, D. Witter. **n. syn.**

1500

1501 **Additional material examined. ARGENTINA: Entre Ríos:** 8.63 km W Concordia, -31.42303 -

1502 58.11672, 16 m, 26 Dec. 2007, W. & E. MacKay, 7 workers, 1 male, 1 queen (WEMC:

1503 USNMENT00757969, 00757997, 00758001, 00758003, 00758004, 00758013, 00759019) **Santa**

1504 **Fe:** 10 km E Santa Fe, -31.6666 -60.5833, 12 Oct. 2002, A.L. Wild & N. Helle, 1 worker



1505 (ALWC: USNMENT00757998). **Tucumán:** Lara, 4000 m, Feb. 2003, G.A. Baer, 2 workers  
1506 (MCSN: USNMENT00757709). **BOLIVIA: Santa Cruz:** Perforación, 68 km ESE Charagua, -  
1507 19.91667 -62.56667, 470 m, 11 Dec. 1993, P.S. Ward, 3 workers (PSWC:  
1508 USNMENT00758008). **BRAZIL:** Brasilia D.F., Aug. 1996, R.M. Oliveira, 6 workers (CPDC:  
1509 USNMENT00758011). **CHILE:** Temuco, 24 Nov. 1967, W.W. Kempf, 6 workers (PSWC:  
1510 USNMENT00758015-00758022). **COLOMBIA: Quindío:** Génova, Vereda El Cedral, Finca  
1511 Venecia, 4.2275 -75.7586, 1800 m, 19 Oct. 1999, E. González & J. Sossa, 1 worker (IAvH:  
1512 IAvH-E74162). **COLOMBIA: Risaralda:** Apia, La Clarita, 3.13 -75.95, 1550 m, 26 Oct. 2001,  
1513 I. Armbrecht, 1 worker (IAvH: IAvH-E74173). **DOMINICAN REPUBLIC:** 28 km SSE  
1514 Constanza, 18.7 -70.9, 2220 m, 11 Nov. 1992, P.S. Ward #11757, 3 workers (PSWC:  
1515 USNMENT00758034). **La Vega:** Reserva Valle Nuevo, 18.81667 -70.68333, 2240 m, 01 Sep.  
1516 2001, A.L. Wild #AW1348, 2 workers, 1 queen (ALWC: USNMENT00757682); Cervantía,  
1517 18.85 -70.70, 1730 m, 01 Sep. 2001, A.L. Wild, 2 workers (ALWC: USNMENT00757988).  
1518 **ECUADOR: Napo:** near Dureno, 0.0780 -76.7307, 287 m, 20 July 2005, W. & E. MacKay  
1519 #21277, 2 workers (WEMC: USNMENT00759007). **GUATEMALA: Sacatepéquez:** Finca El  
1520 Pilar, near Antigua, 14.55 -90.72, 1700 m, 13 Nov. 2003, A.L. Wild, 3 workers (ALWC:  
1521 USNMENT00757963). **PARAGUAY: Boquerón:** Filadelfia, -22.35 -60.03, 22 Sep. 1994, B.  
1522 Garcete, 1 worker (ALWC: USNMENT00758005). **Canindeyú:** Reserva Natural Bosque  
1523 Mbaracayú Lagunita, -24.13 -55.43, 12 Feb. 1997, A. Wild, 3 workers (ALWC:  
1524 USNMENT00757999). **Concepción:** Concepción centro, -23.42 -57.35, 7 Feb. 1998, A. Wild, 3  
1525 workers (ALWC: USNMENT00757976). **Presidentes Hayes:** Villa Hayes, -25.10 -57.57, 21  
1526 Sep. 1994, B. Garcete, 2 workers (ALWC: USNMENT00757996). **UNITED STATES:**  
1527 **Arizona:** Pima Co. Tucson International Airport, 32.11667 -110.93333, 800 m, 07 Aug. 2001,  
1528 P.S. Ward #14412, 2 workers (PSWC: USNMENT00757972). **Florida:** Florida Gulf Co.

1529 Wewahitchka Steele Rd/GCI Bond, 30.1 -85.2, 13.6 m, 23 Dec. 2000, Corrie Saux, 1 worker  
1530 (MCZC: USNMENT00758014). **Louisiana:** Audubon Park, New Orleans, 29 Apr. 1995, A.L.  
1531 Wild, 2 workers (ALWC: USNMENT00757979); Baton Rouge, 01 Oct. 2000, B. Raphaël, 5  
1532 workers (CPDC: USNMENT00757980); E. Baton Rouge Par. Baton Rouge. Kenniloworth &  
1533 Perkins Rd. BREC Perkins Park, 03 Apr. 2003, J. Rosson, 1 worker (CPDC:  
1534 USNMENT00757967). Plaquemines Co. St. Bernard St. Pk., 22 Aug. 1987, W. MacKay, 22  
1535 Aug. 1987, W. MacKay, 2 workers (WEMC: USNMENT00757970). **Texas:** Austin, Travis Co.,  
1536 30.25167 -97.76722, 160 m, 21 Nov. 2006, A.L. Wild, 3 workers (ALWC:  
1537 USNMENT00757977). **URUGUAY:** Montevideo, Nov. 2000, L. Pastre, 2 workers (CPDC:  
1538 USNMENT00757978). Salto: Salto Parque Municipal Benito Solari, 25 Dec. 2007, W. & E.  
1539 MacKay, 1 worker (WEMC: USNMENT00758047).

1540  
1541 **Diagnosis.** *Brachymyrmex bruchi* is morphologically most similar to *B. patagonicus* and *B.*  
1542 *oculatus* because these species have scapes that surpass the posterior margin of the head,  
1543 typically two erect hairs on the mesonotum, their metanotal groove is either absent or narrower  
1544 than the diameter of the metathoracic spiracles, their mesonotum does not bulge dorsally above  
1545 the pronotum in lateral view, their gaster has several scattered long erect hairs and sparse  
1546 pubescence, and they have brownish bodies. However, *B. bruchi* differs from *B. patagonicus* by  
1547 having a larger body size, abundant suberect hairs on the dorsum of the pronotum, and scapes  
1548 that surpass the posterior margin of the head by a length approximately equal to the maximal  
1549 diameter of the eye. It differs from *B. oculatus* by having smaller eyes with less than 14  
1550 ommatidia along the maximal diameter, which approximately equal only a quarter of the length  
1551 of the head (HL<sub>1</sub>).

1552

1553 *Lectotype and paralectotypes measurements* (mm) (n= 4). HL<sub>1</sub> 0.58-0.64; HL<sub>2</sub> 0.41-0.45; HL<sub>3</sub>  
1554 0.14-0.18; HW 0.57-0.60; SL 0.55-0.62; EL 0.16-0.18; WL 0.64-0.78; PnL 0.20-0.25; PnW 0.39-  
1555 0.45; ML 0.18-0.21; MW 0.27-0.31; *Indices* CI 90.63-100.00; SI<sub>1</sub> 93.33-103.45; SI<sub>2</sub> 130.43-  
1556 142.86; OI<sub>1</sub> 26.67-31.03; OI<sub>2</sub> 21.88-28.13.

1557  
1558 *Additional material examined measurements* (mm) (n=25). HL<sub>1</sub> 0.41-0.61; HL<sub>2</sub> 0.29-0.42; HL<sub>3</sub>  
1559 0.09-0.16; HW 0.37-0.60; SL 0.31-0.59; EL 0.10-0.20; WL 0.40-0.72; PnL 0.10-0.20; PnW 0.25-  
1560 0.44; ML 0.09-0.20; MW 0.16-0.29; *Indices* CI 70.98-96.97; SI<sub>1</sub> 78.13-136.36; SI<sub>2</sub> 106.67-  
1561 156.76; OI<sub>1</sub> 23.91-45.45; OI<sub>2</sub> 19.23-28.00.

1562  
1563 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1564 concave. Dorsum of the head with sparse and appressed hairs. Clypeus with a rounded anterior  
1565 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
1566 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
1567 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
1568 margin in oblique anterodorsal view. The scapes surpass the posterior cephalic margin with a  
1569 length that approximately equals the maximal diameter of the eye; they typically have appressed  
1570 and decumbent hairs. Three inconspicuous ocelli present. Eyes are positioned on the cephalic  
1571 midline and have 8–13 ommatidia along their maximal diameter.

1572 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum,  
1573 sometimes with additional suberect hairs, mainly on the pronotum. The mesonotum is not inflated  
1574 and does not bulge dorsally above the pronotum in lateral view. Metanotal groove absent or  
1575 narrower than the diameter of the metathoracic spiracles. Metathoracic spiracles in dorsolateral  
1576 position, not protruding, but touching the propodeal suture. Dorsum of the propodeum slightly

1577 convex and shorter than the posterior slope. Propodeal spiracles circular, positioned on the  
1578 posterior propodeal margin at the middle of the propodeal slope. Legs with appressed and  
1579 scattered hairs. Petiole short and inclined forward.

1580 **Gaster.** With sparse pubescence and several scattered long erect hairs.

1581 **Color and sculpture.** Body overall smooth and shiny, except for the sometimes slightly  
1582 imbricate sculpture on the dorsum of the mesosoma; typically brownish.

1583

1584 **Distribution (supplementary material Fig. S10).** *Brachymyrmex bruchi* is known from  
1585 Argentina, Bolivia, Brazil, Chile, Colombia, the Dominican Republic, Ecuador, Guatemala,  
1586 Paraguay, Uruguay, and the United States.

1587

1588 **Biology.** Unknown.

1589

1590 **Remarks.** The here designated lectotype is the specimen at the top of pin MHNG:  
1591 USNMENT00757159, and the ants below are paralectotypes. The type material of *B. bruchi*, *B.*  
1592 *giardi* var. *nitida*, *B. laevis* var. *andina*, and *B. bruchi* var. *rufipes* shares a common set of  
1593 diagnostic characters, i.e. the brownish body color, scapes that surpass the posterior cephalic  
1594 margin by a length that approximately equals the maximal diameter of the eye, the metanotal  
1595 groove is lacking or narrow, and scattered pubescence on the gaster. As such, these species and  
1596 varieties are synonymized here.

1597 *Brachymyrmex giardi* var. *nitida* was considered to be a junior synonym of *B. giardi* by Snelling  
1598 and Hunt (1975), however we disagree with this synonymization taking in account both the  
1599 description of Santschi (1922) and our own observations of important differences in diagnostic

1600 traits: *B. giardi* var. *nitida* differs from *B. giardi* by having erect hairs on the pronotum, a darker  
1601 body color and scarce pubescence on the body.

1602 Quirán et al. (2004) suggested *B. bruchi* var. *rufipes* to be a junior synonym of *B. bruchi* based on  
1603 the original description by Forel (1912a), which only specifies a difference in body color. Quirán  
1604 et al. (2004) argued that this difference is not taxonomically informative and therefore they  
1605 proposed synonymization. We agree that body color is variable in several *Brachymyrmex* species,  
1606 and therefore we follow the suggestion of Quirán et al. (2004) here. Nevertheless, it is  
1607 noteworthy that some individuals of *B. bruchi* var. *rufipes* have somewhat denser pubescence on  
1608 the gaster than pointed out in our description above (see Fig. 40 d, f). Such moderately dense  
1609 pubescence on the gaster has been also observed in some of the examined specimens of *B. laevis*  
1610 var. *andina*. Future studies on *B. bruchi* and its geographical variation is required.

1611 Forel (1912a) indicated that *B. bruchi* and *B. patagonicus* are very similar as to their mesosoma,  
1612 and Santschi (1923a) likewise compared *B. laevis* var. *andina* with *B. patagonicus* var. *atrátula*,  
1613 which has been synonymized with *B. patagonicus* (Quirán et al. 2004). We concur with these  
1614 authors that *B. bruchi* and *B. patagonicus*, including their type material, are morphologically very  
1615 similar, but as noted in the diagnosis above, consistent differences also exist between both  
1616 species. Furthermore, our morphometric and phylogenetic analyses tentatively confirm these taxa  
1617 to be distinct, although further studies on the morphology and phylogenetics of these species as  
1618 well as their ecology and biology are admittedly needed.

1619 Santschi (1929) and Quirán et al. (2004) also referred specimens from Jujuy: Pueblo Viejo  
1620 (Weiser), Catamarca: Aconquija and Tucuman (Argentina) to *B. bruchi*, however, this material  
1621 was not studied here.

1622

1623

1624 *Brachymyrmex cavernicola* Wheeler  
1625 (Figs. 18, supplementary material Fig. S11)  
1626 *Brachymyrmex cavernicola* Wheeler, 1938: 252 (w.m.). **Lectotype worker** (USNM:  
1627 USNMENT00529073) and **Paralectotype workers, male** (USNM: USNMENT00529073;  
1628 MCZC: M.C.Z. Cotype 17-19 22428, M.C.Z. Cotype 11-13 22428, M.C.Z. Cotype 23-25 22428,  
1629 M.C.Z. Cotype 14-16 22428, M.C.Z. Cotype 1-3 22428, M.C.Z. Cotype 5-7 22428, M.C.Z.  
1630 Cotype 4 22428; **here designated**): 21 workers, 1 male [examined]. **MEXICO: Yucatan,**  
1631 Chichenitza, Balaam Canche Cave, H.S. Pearse, 13 June 1936.  
1632  
1633 **Additional material examined. BRAZIL: Amazonas:** Manaus, BR.174 km 45 EEST-S1, 12  
1634 Nov. 1990, Eq. A. Y. Harada, A. G Baineira, 1 worker (MPEG: USNMENT00757857]. **Pará:**  
1635 Melgaço, Caxiuanã ECFPn, -1.73584 -51.48762, 12-14 Oct. 2006, Equipe A.Y. Harada, 1 worker  
1636 (MPEG: AYH051). Melgaço, Caxiuanã ECFPn, -1.75444 -51.52241, 28 Oct. 2003, A.Y. Harada,  
1637 E.P. Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 1  
1638 worker (MPEG: AYH067). Melgaço, Caxiuanã ECFPn, -1.73584 -51.48762, 30 Oct. 2003, A.Y.  
1639 Harada, E.P. Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato,  
1640 3 workers (MPEG: AYH086, AYH088, AYH126). Serra Norte, Est. Do. Mang, 6 Sep. 1983, 12  
1641 Sep. 1983, 29 Feb. 1984, 12 May 1984, 15 May 1984, 22 May 1984, Lote: 2105, 2108, 2195,  
1642 2197, 2208, 2213, 2214, 2223, 2227, 2231, 2232, 2235, 11 workers (MPEG:  
1643 MPEG\_HYM11505683, 11505907, 11505913, 11505945, 11505960, 11505969, 11505999,  
1644 11506007, 11506023, 11506030, 11506036). **COLOMBIA: Cauca:** Isla Gorgona, 17 Sep. 1989,  
1645 M. Baena #GQA-05, 1 worker (WEMC: USNMENT00757854). Isla Gorgona, 16 Jan. 1990, M.  
1646 Baena #Gacd-19, 2 workers (WEMC: USNMENT00757855, 00757856). **Cundinamarca: La**  
1647 Vega, R.N. Natautá, 5.00 -74.33, 1040 m, 10 Nov. 2010, F. Fernández, 2 workers (IAvH:

1648 USNMENT00757859). **Nariño:** Barbacoas, Tajadas, 100 m, 1 worker (ICN:  
1649 USNMENT00757858). **Quindío:** Buenavista, Finca Ceilán, Bs., 4.35833 -75.78472, 1100 m, 15  
1650 Nov. 1999. E. Gonzalez, 1 worker (IAvH: IAvH-E744170). **COSTA RICA: Guanacaste:** Prov.  
1651 Maritza field, Sta., 800 m, 03 May 1995, R. Anderson #17714, 4 workers (WEMC:  
1652 USNMENT00757844, 00757845). **Heredia:** Estación Biológica, La Selva, 10.433 -84.017, 50-  
1653 150 m, 01 June 1993, INBio-OET, 1 worker (JTLC: INBIOCRI001276875). Puerto Viejo, #733,  
1654 25 June 1979, J. Raich, 7 workers (MCZC: USNMENT00757273, 00757275). **ECUADOR:**  
1655 **Endesa:** Forest Reserve Pichincha Province, 25 Jan. 1994, L.E. Tennant, 1 worker (MCZC:  
1656 USNMENT00757274). **Pichincha:** La Unión del Río Toachi, -0.31889 -78.95442, 770 m, 15  
1657 Jul. 2005, W. & E. MacKay #21169, 2 workers (WEMC: USNMENT00757841, 00757842).  
1658 **MEXICO: Chiapas:** 8.8 km SE Salto de Agua, 17.51328 -92.29515, 50 m, 14 July 2007, J.L.  
1659 Cozar ANTC#4225, 1 worker (JTLC: CASENT0600011). 10 km S Palenque, 30 May 1988, A.  
1660 Rabeles, 2 workers (WEMC: USNMENT00757848). 10 km S Palenque, 30 May 1988, W.  
1661 MacKay #10563, #10571, #10627, #10674, 15 workers (WEMC: USNMENT00757849,  
1662 00757850, 00757851, 00757852, 00757853, 00758028). 10 km S Palenque, 30 May 1988, VIAL,  
1663 D. Gonzalez, 1 worker (WEMC: USNMENT00757846). **Veracruz:** Los Tuxtlas, 10 km NNW  
1664 Sontecomapan, 18.583 -95.083, 500 m, 21 Mar. 1985, P.S. Ward #7366, 3 workers (PSWC:  
1665 USNMENT00757843). **PERU: Madre de Dios:** Prov. Tambopata, Cuzco Amazónico, 15 km  
1666 NE Puerto Maldonado, CA-130, 200 m, 13 June 1989, S.P. Cover & J.E. Tobin, 6 workers  
1667 (MCZC: USNMENT00757260, 00757269, 00757270). Prov. Tambopata, Cuzco Amazónico, 15  
1668 km NE Puerto Maldonado, CA-601 JT79, CA-601 JT80, CA-365, CA-659 JT138, CA-116, CA-  
1669 141, June 1989, S.P. Cover & J.E. Tobin, 17 workers (MCZC: USNMENT00757260-00757272).  
1670

1671 **Diagnosis.** The feature that allows distinguishing *B. cavernicola* from all other *Brachymyrmex*  
1672 species is the presence of conspicuous thick black hairs on the head, mesosoma and gaster, which  
1673 contrast strongly with the yellowish body, a condition reminiscent of *Nylanderia*. *Brachymyrmex*  
1674 *antennatus* also has erect hairs on the mesosoma that are darker than the tegument, however these  
1675 are not as thick as those of *B. cavernicola*, and in other features these species are very different.

1676  
1677 *Additional material examined measurements* (mm) (n=10). HL<sub>1</sub> 0.51-0.57; HL<sub>2</sub> 0.35-0.41; HL<sub>3</sub>  
1678 0.13-0.18; HW 0.45-0.51; SL 0.54-0.63; EL 0.09-0.10; WL 0.51-0.63; PnL 0.19-0.21; PnW 0.29-  
1679 0.35; ML 0.10-0.13; MW 0.18-0.22; *Indices* CI 83.33-91.67; SI<sub>1</sub> 118.18-134.62; SI<sub>2</sub> 139.53-  
1680 166.67; OI<sub>1</sub> 17.54-20.83; OI<sub>2</sub> 25.00-31.25.

1681  
1682 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1683 concave. Thick hairs cover the front of the head. Clypeus with a rounded anterior margin and five  
1684 long, erect hairs of which a single hair is near the anterior margin, two hairs are in mediolateral  
1685 position and two more near the toruli; other hairs on the clypeus are markedly shorter and  
1686 appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique anterodorsal  
1687 view. The scapes surpass the posterior cephalic margin by a length of approximately 2.0× the  
1688 maximal diameter of the eyes and bear appressed or decumbent hairs. Ocelli are absent or one  
1689 central ocellus is present. Eyes are positioned on the cephalic midline and have 7-8 ommatidia  
1690 along their maximal diameter.

1691 **Mesosoma.** With several thick erect hairs on the promesonotum (>2), and two between the  
1692 metathoracic spiracles, but none on the propodeum. The mesonotum is not inflated and does not  
1693 bulge dorsally above the pronotum in lateral view. Metanotal groove wider than the diameter of  
1694 the metathoracic spiracles. Metathoracic spiracles in dorsolateral position, not protruding, and not



1695 touching any suture. Dorsum of the propodeum is weakly convex and shorter than the posterior  
1696 slope. Propodeal spiracles circular, positioned just ventrally of the posterior propodeal margin  
1697 slightly posterior of the middle of the propodeal slope. Legs with appressed and scattered  
1698 suberect hairs. Petiole short and inclined forward.

1699 **Gaster.** With scarce pubescence but densely covered by thick, erect hairs.

1700 **Color and sculpture.** Body smooth and shiny, except for the dorsum of the mesosoma which  
1701 sometimes has slightly imbricate sculpture. The body color is typically yellowish, although the  
1702 head and gaster are sometimes darker than the mesosoma. Tegument color contrasts with the  
1703 thick black hairs.

1704

1705 **Distribution (Supplementary material Fig. S11).** *Brachymyrmex cavernicola* is known from  
1706 Argentina, Brazil, Colombia, Costa Rica, Ecuador, Mexico and Peru.

1707

1708 **Biology.** Nests of this species have been found in the soil under stones.

1709

1710 **Remarks.** The lectotype is the top specimen on pin USNMENT00759073, whereas the  
1711 specimens below are the paralectotypes. As indicated in the diagnosis and pointed out before by  
1712 Wheeler (1938), *B. cavernicola* resembles *Nylanderia* species by its thick hairs that cover the  
1713 entire body.

1714

1715

1716 *Brachymyrmex coactus* Mayr

1717 (Figs. 19, 20, supplementary material Fig. S12)

1718 *Brachymyrmex coactus* Mayr, 1887: 523 (w.q.m.). **Lectotype worker** (NHMW:  
1719 USNMENT00757191) and **paralectotype workers, males, queens** (NHMW:  
1720 USNMENT00757191- 00757195; **here designated**): 3 workers, 3 males, 2 queens [examined].  
1721 **BRAZIL: Santa Catharina** (Hetscko). See also: Santschi (1923a: 669); Santschi (1923b: 272).  
1722 = *Brachymyrmex coactus* var. *nictitans* Emery 1906: 178 (w.). (MCSN: USNMENT 00757209):  
1723 1 worker [examined]. **COSTA RICA**. See also: Santschi (1923a: 670). **n. syn.**  
1724 = *Brachymyrmex constrictus* Santschi 1923a: 671, Figs. 5, 38, 61 (w.). (NHMB:  
1725 USNMENT00758087): 1 worker [examined]. **BOLIVIA: La Paz**: Mapiri. **n. syn.**  
1726 = *Brachymyrmex coactus* var. *robustus* Santschi 1923b: 272 (w.). (NHMB:  
1727 USNMENT00757224): 4 workers [examined]. **BRAZIL: Santa Catharina**: Encano alto;  
1728 (NHMB: USNMENT00758085, 00758086): 6 workers [examined]. **BRAZIL: Santa Catharina**:  
1729 Blumenau. **n. syn.**  
1730  
1731 **Additional material examined. ARGENTINA**: “Fives Lile”, 4 workers, 5 queens, 1 male  
1732 (NHMB: USNMENT00758083, 00758084). **BRAZIL: Alagoas**: Maceió – Emilia Flores, Hm  
1733 For 68, For 91, 18 Mar. 2005, 04 June 2005, #5460, M.C.C. Diniz, 8 workers (CEPLAC:  
1734 USNMENT00757552, 00757553, 00757555); **Bahia**: Barreiras, Serra do Mimo, 24 Apr. 2010, S.  
1735 Souza & B. Santos, 3 workers (CEPLAC: USNMENT00757564); Esplanada, Baixio, -12.11444 -  
1736 37.69944, June– Oct. 2010, M.L.O. Travassos, 1 worker (CEPLAC: USNMENT00757556);  
1737 Porto Seguro, Troncoso, 12 June 1991, J. Delabie 4451, 3 workers (CEPLAC:  
1738 USNMENT00757559); UNA-ESMAI, Estação Experimental Lemos Maia, Em coqueiro-anão,  
1739 Oct. 2005, J.R.M. Santos, 8 workers (CEPLAC: USNMENT00757558, 00757559, 00757562);  
1740 **Goiás**: Ouro Verde, Faz Boa Vista, -16.29847 -49.21183, 01-07 July 2005, R.R. Silva & R.M.  
1741 Feitosa, 3 workers (ICN: MZSP123); **Santa Catharina**: Blumenau, M. Witte #150, 9 workers

1742 (MCZC: USNMENT00757238, 00757239, 00757251); **Paraíba:** Independencia, Mann & Heath,  
1743 -7.15194 -34.90556, 3 workers, 1 queen (MZUSP: USNMENT00757240); **São Paulo:**  
1744 ANHEMBI, Faz B. Rio, 14 Feb. 1969, W. Kemf, J.C. Mahalhães, L.T.F., M. Kulmann, 2  
1745 workers, 1 queen (MZUSP: USNMENT00757563); Sete Barras, PE Carlos Bothelo, -24.20833 -  
1746 47.97056, 200 m, 11-15 May 2009, F. Esteves leg, 2 workers (MZUSP: USNMENT00757560).  
1747 **COSTA RICA: Puntarenas:** Sirena, Corcovado National Park, 8.48333 -83.60000, 10 m, 24  
1748 Dec. 1981, J. Longino, 1 worker, 1 male (JTLC: JTLC000005905); Peninsula Osa, 8.46667 -  
1749 83.58333, 50 m, 24 Dec. 1981, J. Longino, 1 worker, 1 male (MCZC: USNMENT00757243).  
1750 **ECUADOR: Zamora-Chinchi:** Copalinga, -4.09122 -69.93591, Jacquemin, Col id 5087, 1  
1751 worker (RBINS) **GUYANA: Rupunini:** Kananambo, 16 Jan. 1981, 3.75 -59.3, 100 m, J.  
1752 Longino, 1 worker (JTLC: JTLC000005907). **PANAMA: Barro Colorado I:** Canal Zone, B50,  
1753 Jan 1960, W.L. Brown, E.S. McCluskey, 3 workers (MCZC: CMOS0000097). **PARAGUAY:**  
1754 Fortin mayor infante Rd, trans Chaco locality 1, 01 Oct. 2004, T. Delsinne, 1 worker (RBINS:  
1755 Coll.RIScNB SID SPM\_ID 30833); **Boquerón:** Enciso, -21.20 -61.67, 3-6 Nov. 2001, M.  
1756 LePonce & T. Delsinne, Dry Chaco, Pitfall trap, 3 workers (ALWC: USNMENT00757554);  
1757 **Boquerón:** Enciso, -21.20609 -61.65748, 04-06 Nov. 2001, 23-25 Sep. 2004, M. LePonce, T.  
1758 Delsinne, Col ID4132, Col ID 13623, 2 workers (RBINS: Coll.RIScNB SID SPM\_ID 22607,  
1759 ID27462); Nueva Asunción, -20.69190 -61.92925, 02-06 Nov 2001, M. Leponce, Col ID 3948, 1  
1760 worker (RBINS: Coll.RIScNB SID SPM\_ID 30542); **Canindeyú:** Reserva Natural Bosque  
1761 Mbaracayú, Jejuimi, -24.10 -61.67, 02 Apr. 1996, A.L. Wild, 3 workers (ALWC:  
1762 USNMENT00757561); **Cordillera:** Caacupé, Camp. J. Noment, -25.36667 -57.08333, 19 Jan.  
1763 1994, B. Garcete #AW0395, 1 worker (ALWC: USNMENT00757567); **Misiones:** 8 km SE San  
1764 Juan Bautista, -26.71666 -57.06667, 150 m, 10 Dec. 2002, A.L. Wild & B. Garcete #AV1781, 1  
1765 worker (ALWC: USNMENT00757570). **PERU: Madre de Dios:** Reserva Nacional Tambopata,

1766 Centro Sachavacayoc, -12.85583 -69.36194, 210 m, 19-31 July 2012, R.M. Feitosa, 2 workers  
1767 (ICN: USNMENT00757614, 00757612); Reserva Nacional Tambopata, Centro Sachavacayoc, -  
1768 12.82667 -69.37056, 198 m, 26 July 2012, GSNMBU, 2 workers (ICN: USNMENT00757615,  
1769 00757838).

1770  
1771 **Diagnosis.** *Brachymyrmex coactus* is morphologically very similar to *B. degener* as both species  
1772 have scapes that surpass the posterior margin of the head, faint sculpture on the mesosoma, a  
1773 mesonotum that is inflated and bulges dorsally above the pronotum in lateral view, a wide  
1774 metanotal groove, metathoracic spiracles that are slightly protruding dorsally, and a gaster with  
1775 scarce pubescence. However, *B. coactus* has a brown yellowish head and mesosoma, but a darker  
1776 gaster, whereas *B. degener* has a uniformly brownish body.

1777  
1778 *Lectotype and paralectotypes measurements* (mm) (n=3). HL<sub>1</sub> 0.72-0.84; HL<sub>2</sub> 0.44-0.55; HL<sub>3</sub>  
1779 0.21-0.25; HW 0.64-0.82; SL 0.68-0.80; EL 0.16-0.21; WL 0.53-0.88; PnL 0.21; PnW 0.43-0.55;  
1780 ML 0.14-0.20; MW 0.23-0.35; *Indices* CI 94.29-97.67; SI<sub>1</sub> 97.62-106.06; SI<sub>2</sub> 144.00-152.17; OI<sub>1</sub>  
1781 24.24-28.57; OI<sub>2</sub> 29.73-31.43.

1782  
1783 *Additional material examined measurements* (mm) (n=10). HL<sub>1</sub> 0.52-0.88; HL<sub>2</sub> 0.34-0.60; HL<sub>3</sub>  
1784 0.16-0.25; HW 0.50-0.82; SL 0.51-0.82; EL 0.13-0.21; WL 0.53-0.98; PnL 0.18-0.25; PnW 0.36-  
1785 0.57; ML 0.13-0.23; MW 0.18-0.35; *Indices* CI 93.33-101.45; SI<sub>1</sub> 92.86-105.36; SI<sub>2</sub> 126.67-  
1786 165.38; OI<sub>1</sub> 23.53-35.71; OI<sub>2</sub> 26.67-32.5.

1787  
1788 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin flat or  
1789 slightly concave. Dorsum of the head has scattered appressed hairs. Clypeus with a rounded

1790 anterior margin and five long, erect hairs of which a single, usually conspicuous hair is near the  
1791 anterior margin, two hairs are in mediolateral position and two more near the toruli; other hairs  
1792 on the clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior  
1793 clypeal margin in oblique anterodorsal view. The scapes extend beyond the posterior margin of  
1794 the head by a length that is equal to the maximal diameter of the eye or larger, and they bear  
1795 appressed and decumbent hairs. Three ocelli are present. Eyes are positioned on the cephalic  
1796 midline and have 10-14 ommatidia along their maximal diameter.

1797 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum;  
1798 sometimes with additional suberect hairs, mainly on pronotum. Dorsum of the mesosoma with  
1799 imbricate sculpture. The mesonotum is inflated and bulges dorsally above the pronotum in lateral  
1800 view. Metanotal groove wider than the diameter of the metathoracic spiracles. Metathoracic  
1801 spiracles in dorsolateral position, slightly protruding and not touching any sutures. Dorsum of the  
1802 propodeum strongly convex and shorter than the propodeal slope. Propodeal spiracles  
1803 conspicuous and circular, positioned on the propodeal margin, anterior of the middle of the  
1804 propodeal slope. Legs with appressed hairs. Petiole short and inclined forward.

1805 **Gaster.** With scattered pubescence and several scattered long erect hairs.

1806 **Color and sculpture.** Head and gaster are smooth and shiny, but the dorsum of the mesosoma  
1807 usually has imbricate sculpture. Head and mesosoma are brown yellowish, whereas the gaster is  
1808 darker.

1809  
1810 **Distribution (Supplementary material Fig. S12).** *Brachymyrmex coactus* occurs in Argentina,  
1811 Bolivia, Brazil, Costa Rica, Guyana, Panama, Paraguay and Peru.

1812

1813 **Biology.** The specimens from UNA-ESMAI in Bahia (Brazil) were found in dwarf coconuts  
1814 (CEPLAC: USNMENT00757557, 00757558), those from Canindeyú (Paraguay) on shrubs  
1815 (ALWC: USNMENT00757561), and those from St. Catharina underneath bark in association with  
1816 a beetle of the genus *Claviger* (Mayr 1887).

1817

1818 **Remarks.** The worker on pin USNMENT00757191 is designated here as lectotype. Emery  
1819 (1906) considered *B. coactus* var. *nictitans* to be a separate variety because it has smaller eyes  
1820 than *B. coactus*, but he also expressed doubt on the level of consistency of this difference.

1821 Santschi (1923a) did not provide a motivation to distinguish *B. constrictus* from *B. coactus* but  
1822 indicated that *B. constrictus* has more finely imbricate sculpture on the mesosoma and smaller  
1823 metathoracic spiracles. Subsequently, he (Santschi 1923b) reported that his original description  
1824 (Santschi 1923a) of *B. coactus* refers to *B. coactus* var. *robusta*. This variety has a larger body  
1825 size, more sculpture on the mesosoma, and a somewhat bigger head than the ‘typical’ form of *B.*  
1826 *coactus*. Both varieties of *B. coactus* were described from the same type locality, however.

1827 After examining specimens of all these varieties and *B. constrictus*, we consider the main  
1828 morphological differences to relate to variation in body size. This trait is, however, very variable  
1829 even within localities (including among specimens mounted on the same pin). As these taxa all  
1830 have the same diagnostic features we here synonymize *B. coactus* var. *nictitans*, *B. coactus* var.  
1831 *robustus*, and *B. constrictus* with *B. coactus*.

1832

1833

1834 *Brachymyrmex cordemoyi* Forel

1835 (Figs. 21, 22, 23, supplementary material Fig. S13)

- 1836 *Brachymyrmex patagonicus* var. *cordemoyi* Forel, 1895a: 49 (w.). (MHNG) [not examined].
- 1837 **REUNION.** See also: Emery (1906: 180) (q.m.). Raised to species: Emery (1906: 179).
- 1838 Subspecies of *Brachymyrmex patagonicus*: Forel (1908: 399); Forel (1912b: 165); Santschi
- 1839 (1912: 533). Revived status as species: Wheeler (1922: 1036); Emery (1925: 41). See also Forel
- 1840 (1912a: 62).
- 1841 = *Brachymyrmex laevis* var. *fuscula* Emery, 1906: 178 (w.q.). (MCSN: USNMENT00757216,
- 1842 00757217; MHNG00758131 - 00758133): 10 workers, 1 queen [examined]. **ARGENTINA:**
- 1843 **Mendoza:** Mendoza, Punta de vacas; (MCSN: USNMENT00757215): 8 workers [examined].
- 1844 **ARGENTINA: Buenos Aires:** Buenos Aires. **n. syn.**
- 1845 = *Brachymyrmex brevicornis* Emery, 1906: 180, Figs. 38, 40, 41 (w.q.m.). (MCSN:
- 1846 USNMENT00757210 - 00757214): 16 workers, 1 queen, 1 male [examined]. **ARGENTINA:**
- 1847 **Buenos Aires:** Santa Catalina. See also Quirán (2005: 765). **n. syn.**
- 1848 = *Brachymyrmex patagonicus* var. *brevicornoides* Forel, 1914: 287 (w.q.m.). (MHNG:
- 1849 USNMENT00758141 - 00758143): 4 workers, 3 males [examined]. **ARGENTINA: Buenos**
- 1850 **Aires:** Tapalquén. Junior synonym of *Brachymyrmex nigricans*: Santschi (1923a: 657).
- 1851 [*Brachymyrmex brevicornoides* has priority as senior name, *Brachymyrmex nigricans* is its
- 1852 junior synonym: Bolton (1995: 81)]. **n. syn.**
- 1853 = *Brachymyrmex cordemoyi* var. *nigricans* Santschi, 1916: 395 (w.). (NHMB:
- 1854 USNMENT00758081): 3 workers [examined]. **ARGENTINA:** Río de la plata, Isla Martín
- 1855 García; (NHMB: USNMENT00758078, 00758080): 10 workers. **ARGENTINA: Buenos Aires:**
- 1856 Buenos Aires. [First available use of *Brachymyrmex patagonicus cordemoyi nigricans* Santschi
- 1857 (1912: 533) unavailable name]. Raised to species: Santschi (1923a: 657). Junior synonym of
- 1858 *Brachymyrmex nigricans*: Santschi (1923a: 657) [As mentioned above *Brachymyrmex*
- 1859 *brevicornoides* has priority over *B. nigricans*].

1860 = *Brachymyrmex cordemoyi* var. *distincta* Santschi, 1923a: 658, Figs. 6, 50, 59 (w.q.). (NHMB:  
1861 USNMENT00757178, 00757179): 3 workers [examined]. **ARGENTINA: Santa Cruz;**  
1862 (NHMB: USNMENT00758089): 9 workers [examined]. **ARGENTINA: Delta del Paraná. n.**  
1863 **syn.**  
1864  
1865 **Additional material examined. ARGENTINA: Buenos Aires:** Buenos Aires, E.V. Steigen, 3  
1866 workers (MZUSP: USNMENT00759023); Buenos Aires, Universidad de Buenos Aires, 18  
1867 workers (ICN: USNMENT00759032); La Plata, Silvestri, 4 workers (NHMB:  
1868 USNMENT00758088), 8 workers (MCZC: USNMENT00757244); **Entre Ríos:** 8.63 km  
1869 Concordia, -31.41667 -58.11667, 16 m, 26 Dec. 2007, W. & E. MacKay #22670, 1 worker  
1870 (WEMC: USNMENT00757638); Isla frente Puerto Victoria, -32.63333 -60.16667, 10 m, 29 Oct.  
1871 2002, A.L. Wild & N. Heller, 1 worker (ALWC: USNMENT00757966); **Misiones:** 48.93 km N  
1872 Campinas de America, -25.8565 -53.9939, 360 m, 03 Jan. 2008, W. & E. MacKay #22794, 1  
1873 worker (WEMC: USNMENT00757728); **San Juan:** 8.59 km S Villa Aberastain, -31.72528 -  
1874 68.55447, 592 m, 10 Jan. 2008, W. MacKay #22879, 1 worker (WEMC: USNMENT00757737).  
1875 **BRAZIL: Bahia:** Boa Vista do Tupim, 06 Dec. 2010, V.M.S. Cameiro & J.J. Resende, 1 worker  
1876 (CEPLAC: USNMENT00757887); **Mato Grosso do Sul:** ~70 km E Corumbá, Faz. Maria  
1877 Bonita, -19.16666 -57.16666, 22 Aug. 1998, A. L. Wild #AW0657, 1 worker (ALWC:  
1878 USNMENT00759025); 10 km Posto Chapadao, 18 Oct. 1989, S. Porter, 3 workers (WEMC:  
1879 USNMENT00758994); 3 km Anastácio, 17 Oct. 1989, W. MacKay #12605, 3 workers (WEMC:  
1880 USNMENT00757652); Passo da Lontra, -19.53333 -57.01667, 80 m, 08 Sep. 1996, P.S. Ward  
1881 #13222, 1 worker, 1 male (MCZC: CMOS000020); **Para:** Santarem, Taperinha, -2.9 -54.3, July  
1882 1975, R.L. Jeanne, 440, 4 workers (MCZC: CMOS000015, CMOS000016); **Rio Grande do**  
1883 **Norte:** Ceará, Mirim, W.M. Mann, 10 workers (MCZC: CMOS000124, CMOS000125);



1884 **Rondonia:** Ji Parana, 27 Aug. 1984, W. Overal, 2 workers (MPEG: USNMENT00757583,  
1885 00757964); Ouro Preto do Oeste, 25 Mar. 1985, W. França, Res INPA 0050, 3 workers (MPEG:  
1886 USNMENT00757984, 00758043); **São Paulo:** Aguas de São Pedro, May-June 1985, S. Silva, 1  
1887 worker (ICN: USNMENT00757670); Caraguatatuba, Res. Florest rain for, 40-80 m 18-22 May  
1888 1971, W.L. & D.E. Brown, 6 workers (MCZC: CMOS000017, CMOS000019, CMOS000021);  
1889 José Bonifacio, 17 Nov. 1970, J. Diniz, 2 workers, 1 queen (MZUSP: USNMENT00757582).  
1890 **COLOMBIA: Caqueta:** Florencia, 2 workers (ICN: LEV127); **Huila:** 17 km NW, La Plata, 03  
1891 Jan. 1984, W. & E. MacKay #7138, 2 workers (WEMC: USNMENT00757673); **Meta:** 65 km E  
1892 Puerto Lopez, 30 Jan. 1973, W.P. MacKay #7365, 3 workers (WEMC: USNMENT00757636);  
1893 Villavicencio, 17 Dec. 1975, W. & E. MacKay, 2 workers (WEMC: USNMENT00757648);  
1894 Vista Hermosa, 25 Dec. 1975, W. & E. MacKay #75812, #815, 7 workers, 1 queen, 1 male  
1895 (WEMC: USNMENT00757653, 00757675, 00757985). **COSTA RICA: Limón:** Guapiles, R.  
1896 Toro Amarillo vic., 15 Feb.-09 Mar. 1966, W.L. Brown, 3 workers (MCZC:  
1897 USNMENT00757647); **Puntarenas:** 8 km WNW Potrero Grande, 9.03 -85.26, 200 m, 01 Aug.  
1898 1985, P.S. Ward #7791, 3 workers (PSWC: USNMENT00757877). **CUBA: Guantanamo:**  
1899 Baracoa, 20.35 -74.5, 20 m, 26 Aug. 2001, P.S. Ward #14462-14, 3 workers (PSWC:  
1900 USNMENT00757881); **Pinar del Río:** Viñales del Río, 14 June 1953, E.O. Wilson #11, 3  
1901 workers (MCZC: CMOS000018). **DOMINICAN REPUBLIC:** 28 km SSE Constanza, -9.29576  
1902 -75.99786, 11 Sep. 1992, P.S. Ward #11757, 1 worker, 1 queen (PSWC: USNMENT00758016).  
1903 **ECUADOR: Loja:** Estación San Francisco, 2200 m, 11 & 14 Sep. 2011, F. Fernandez, 46  
1904 workers (ICN: USNMENT00759034, 00759036, 00759037); **Napo:** near Dureno, 0.07778 -  
1905 76.73056, 287 m, 20 July 2005, W. & E. MacKay #21273 #21277, 4 workers (WEMC:  
1906 USNMENT00757581, 00757637); **Pichincha:** Mitad del Mundo, 00.00 -78.45, 2483 m, 07 Dec.  
1907 2003, A.L. Wild & J.M. Vieira #AW 2235, 1 worker (ALWC: USNMENT00757888). **EL**

1908 **SALVADOR: La Libertad:** Quezaltepeque, 500 m, 19 June 1963, D.Q. Cavagnaro & M.E.  
1909 Irwin, ANTC 10258, 1 worker (CASENT: CASENT0196000). **GUATEMALA: Suchitepéquez:**  
1910 Finca Tarrales, 12.3 km N Patulul, 14.52256 -91.13642, 740 m, 30 July 2004, W. & E. Mackay  
1911 #20782, 3 workers (WEMC: USNMENT00758045, 00758046). **GUYANA: Kartabo,** July-Aug.  
1912 1920, W.M. Wheeler, 16 workers (MCZC: CMOS000022-000027). **MEXICO: Morelos:**  
1913 Cuernavaca, 25 May 1989, W. MacKay #11418, 2 workers (WEMC: USNMENT00757981);  
1914 **Oaxaca:** 45 km N. San Pedro Pochutla, 1000 m, 03 June 1988, W. MacKay #10755, 6 workers  
1915 (WEMC: USNMENT00758032, 00757649); **Veracruz:** Los Tuxtlas, 10 June 1994, L. Quiroz, 2  
1916 workers (ICN: USNMENT00757661). **NEW CALEDONIA: Kuto Penin. Ile des Pins,** -22.6666  
1917 167.4333, 5 m, 11 May 1980, P.S. Ward #4294-9, 3 workers (PSWC: USNMENT00757882);  
1918 Noumea, 0-100 m, N.L.H. Krauss, 1 worker (CASENT: CASENT0196015). **PARAGUAY:**  
1919 **Central:** Guarambaré, -25.48 -57.45, 25 Apr. 1997, A. Wild #AW0514, 1 worker (ALWC:  
1920 USNMENT00757645); **Guairá:** Roque Gonzalez, -25.88333 -57.28333, 14 Jan. 1995, B.  
1921 Garcete #AW0457, 1 worker (ALWC: USNMENT00759028). **PERU: Madre de Dios:**  
1922 Tambopata, Cuzco Amazónico, 15 km NE Puerto Maldonado, June 1989, S.P. Cover & J.E.  
1923 Tobin, CA-275, 6 workers, 1 queen (MCZC: USNMENT00757276-00757279); **San Martín:**  
1924 Con. Mun. Zona Barreal 23 km S Picota, -7.09111 -76.31361, 335 m, 06-15 Mar. 2005, M.E.  
1925 Irwin & J.D. Vasquez, ANTC1447, 1 worker (CASENT: CASENT0066404). **SEYCHELLES:**  
1926 La Dique Island, 1 m, 09 Nov 1993, Alpert et al., 2 workers (MCZC: USNMENT00757245).  
1927 **SOLOMON ISLAND:** Guadalcanal, Honiara, 0-100 m, Mar 1986, N.L.H. Krauss, ANTC  
1928 10277, 1 worker (CASENT: CASENT0196019). **SURINAME:** Dirkshoop, May 1959, I. V. d.  
1929 Drift, 3 workers (MPEG: USNMENT00757580). **USA: Arizona:** Pima Co. Tucson, 32.23417 -  
1930 110.96666, A.L. Wild #AW2826 (ALWC: USNMENT00757958). **VANUATU:** Tafea, Tanna, 0-  
1931 100 m, Dec. 1985, N.L.H. Krauss, ANTC 10270, 10271, 2 workers (CASENT:

1932 CASENT0196012, 0196013). **VENEZUELA: Lara:** Barquisimeto, to Carora km 19, 29 June  
1933 1971, W.L.& D.E. Brown, 2 workers (MCZC: USNMENT00757884); **Miranda:** D.F Inst Estud.  
1934 Avan. Caracas, 10 Oct 1988, W. MacKay #11144-2 #11146-6, 4 workers (WEMC:  
1935 USNMENT00757654, 00757744).

1936  
1937 **Diagnosis.** *Brachymyrmex cordemoyi* strongly resembles *B. obscurior* and to some extent also *B.*  
1938 *patagonicus*. All these species have scapes that reach or surpass the posterior cephalic margin,  
1939 but by less than the maximal diameter of the eye; their mesonotum does not bulge dorsally above  
1940 the pronotum in lateral view, and the metanotal groove is absent or narrower than the diameter of  
1941 methathoracic spiracles. In general, *B. cordemoyi* has a longer pronotum and mesonotum than *B.*  
1942 *obscurior* and *B. patagonicus*, but these characters show important intraspecific variation.  
1943 Furthermore, it differs from *B. patagonicus* by having considerably denser pubescence on the  
1944 gaster, and from *B. obscurior* by having a larger head, more ommatidia along the maximal  
1945 diameter of the eye, and lighter-colored pubescence which is denser on the dorsum of the entire  
1946 body and appressed on the gaster instead of decumbent in *B. obscurior*.

1947  
1948 *Additional material examined measurements* (mm) (n=20). HL<sub>1</sub> 0.39-0.62; HL<sub>2</sub> 0.27-0.41; HL<sub>3</sub>  
1949 0.10-0.16; HW 0.33-0.59; SL 0.27-0.53; EL 0.08-0.16; WL 0.37-0.60; PnL 0.10-0.20; PnW 0.23-  
1950 0.39; ML 0.08-0.18; MW 0.16-0.29; *Indices* CI 84.38-96.78; SI<sub>1</sub> 82.35-106.38; SI<sub>2</sub> 100.00-  
1951 142.86; OI<sub>1</sub> 23.33-34.69; OI<sub>2</sub> 20.00-29.63.

1952  
1953 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
1954 concave. Dorsal hairs dense and appressed. Clypeus with a rounded anterior margin and five  
1955 long, erect hairs of which a single, usually conspicuous hair is near the anterior margin, two hairs

1956 are in mediolateral position and two more near the toruli; other hairs on the clypeus are markedly  
1957 shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique  
1958 anterodorsal view. Scapes reach the posterior cephalic margin or surpass it by a length up to the  
1959 maximal diameter of the eye; they have appressed hairs. Three inconspicuous ocelli are usually  
1960 present. Eyes are positioned on the cephalic midline and have 10-12 ommatidia along their  
1961 maximal diameter.

1962 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum;  
1963 sometimes with additional suberect hairs, mainly on the pronotum. The mesonotum is not inflated  
1964 and does not bulge dorsally above the pronotum in lateral view. Metanotal groove absent or  
1965 narrower than the diameter of the metathoracic spiracles. Metathoracic spiracles in dorsolateral  
1966 position, not protruding, and typically touching the mesometanotal and propodeal sutures.  
1967 Dorsum of the propodeum slightly convex and shorter than the posterior slope. Propodeal  
1968 spiracles circular, positioned on the posterior propodeal margin, slightly posterior of the middle  
1969 of the propodeal slope. Legs with appressed hairs. Petiole short and inclined forward.

1970 **Gaster.** With dense yellowish pubescence and several scattered and sub-erect hairs, mainly but  
1971 not exclusively along the edges of the segments.

1972 **Color and sculpture.** Body smooth, shiny, and brownish in color.

1973  
1974 **Distribution (Supplementary material Fig. S13).** *Brachymyrmex cordemoyi* is widespread and  
1975 known from Argentina, Brazil, Colombia, the Dominican Republic, Ecuador, El Salvador,  
1976 Guatemala, Guyana, Mexico, Paraguay, Peru, Suriname, the United States, Venezuela and it has  
1977 been introduced in New Caledonia, Seychelles, Vanuatu, Solomon Island, Saudi Arabia (Sharaf  
1978 et al., 2016), Europe and Asia (Ortiz-Sepulveda, pers. obs.).

1979

1980 **Biology.** Some specimens were collected from under stones (PSWC: USNMENT00757877,  
1981 00757881; MCZC: CMOS000020), or on cacti (WEMC: USNMENT00757981).

1982

1983 **Remarks.** We refrain from designating a lectotype because we did not come across the type  
1984 series of *B. cordemoyi* at the MHNG. However, we studied the original description and the type  
1985 series of its varieties (i.e. *distincta*, *nigricans*) (Santschi 1916; Santschi 1923a). Hence, the  
1986 taxonomic decisions made here are based on these data together with the overall morphological  
1987 framework we developed for the genus.

1988 The type series of *B. cordemoyi* was collected in Reunion, and Forel (1895a) already suggested  
1989 that it represents an introduction from the Neotropics. Forel (1895a) indicated that *B. cordemoyi*  
1990 resembles *B. patagonicus*, and originally described it as a variety of the latter species that has  
1991 more pubescence. Such dense pubescence is also observed in *B. patagonicus* var.

1992 *brevicornoides*. Forel (1914) did not provide diagnostic features to distinguish *B. patagonicus*  
1993 var. *brevicornoides* and typical *B. patagonicus*. However, he suggested that the scapes in *B.*  
1994 *patagonicus* var. *brevicornoides* resemble those of *B. brevicornis*, but are slenderer, and that *B.*  
1995 *patagonicus* var. *brevicornoides* has somewhat larger eyes than *B. brevicornis*. Before, Emery  
1996 (1906) had suggested that *B. brevicornis* is closely related to *B. cordemoyi* because they have  
1997 similar integument and pubescence, although the integument is slightly more lucid in *B.*

1998 *brevicornis*. Furthermore, the head and antennal funiculi of *B. brevicornis* are somewhat longer  
1999 than those of *B. cordemoyi*, the clypeus slightly more prominent, and the eyes smaller. However,  
2000 many of these differences could represent geographic variation rather than specific differences. It  
2001 is noteworthy that *B. cordemoyi* is variable in most of these features, and therefore we  
2002 synonymize *B. patagonicus* var. *brevicornoides* and *B. brevicornis* to it here. Quirán (2005)  
2003 redescribed *B. brevicornis* but did not compare or relate it to other *Brachymyrmex* species.

2004 Another example of the variability within *B. cordemoyi* represents *Brachymyrmex cordemoyi* var.  
2005 *distinta*, which was obtained from various places in Argentina, and which has somewhat shorter  
2006 scapes than specimens of the typical *B. cordemoyi* (Santschi 1923a). Beyond this feature, the only  
2007 difference that Santschi (1923a) remarked is the body color of queens. Santschi (1923a) also  
2008 synonymized *B. patagonicus* var. *brevicornoeides* with *B. nigricans*, which he previously  
2009 (Santschi 1916) considered a variety of *B. cordemoyi*. Studying the type material of *B. nigricans*  
2010 we agree with this taxonomic decision, but *brevicornoeides* has taxonomic priority over *nigricans*  
2011 (Bolton 1995: 81), so that *B. nigricans* is also synonymized to *B. cordemoyi* here.

2012 The type specimens of *B. laevis* var. *fuscula* are morphologically more similar to *B. cordemoyi*  
2013 than they are to *B. laevis*, and the only difference Emery (1906) described between *B. laevis* and  
2014 *B. laevis* var. *fuscula* is body color, however, from our observations both differ also in other  
2015 traits, notably the pubescence of the gaster, resulting in their synonymization to *B. patagonicus*  
2016 and *B. cordemoyi*, respectively.

2017 In summary, *B. cordemoyi* has several diagnostic features, however, we also observed  
2018 considerable intraspecific variation in various traits. This variation may hint to a potential species  
2019 complex. *Brachymyrmex cordemoyi* is very widespread and a more comprehensive study of the  
2020 variation within and between its populations would be required to fully resolve the taxonomic  
2021 status of this species.

2022  
2023  
2024 *Brachymyrmex degener* Emery

2025 (Figs. 24, 25, supplementary material Fig. S14)

2026 *Brachymyrmex coactus* subs. *degener* Emery, 1906: 177 (w.). **Lectotype worker** (MCSN:  
2027 USNMENT00757208) and **paralectotype workers** (MCSN: USNMENT 00757207, MCZC:  
2028 M.C.Z. Cotype 01435; **here designated**): 4 workers [examined]. **BRAZIL: Matto Grosso:**  
2029 Cuiaba. Raised to species: Santschi (1923a: 670).  
2030 = *Brachymyrmex admotus* r. *niger* Forel, 1912a: 62 (w.). (MHNG: USNMENT00757162,  
2031 00757163, 00758155): 7 workers [examined]. **BRAZIL: Ceara.** Assigned as *B. degener* st. *niger*  
2032 by Santschi (1923a: 671). **n. syn.**  
2033 = *Brachymyrmex incisus* Forel, 1912a: 63 (w.m.). (MHNG: USNMENT00758134 -00758139,  
2034 00757141-00757143; NHMB: USNMENT00758096): 24 workers, 1 male [examined].  
2035 **COLOMBIA: Naranjo. n. syn.**  
2036 = *Brachymyrmex luederwaldti* Santschi, 1923a: 672, Figs. 36, 66 (w.). (NHMG:  
2037 USNMENT00758140; NHMB: USNMENT00758097, 00758098): 6 workers [examined].  
2038 **BRAZIL: São Paulo: Alcatrazes. n. syn.**  
2039  
2040 **Additional material examined. BRAZIL: Bahia:** Canavieiras, -15.69028 -39.00722, 17 July  
2041 1998, J.C.S. Carmo & J.R.M. Santos, 4 workers (CEPLAC: USNMENT00757566); **Para:** Serra  
2042 Norte, Serraria, -6.08276 -50.16666, 22 Oct. 1984 (MPEG: MPEG\_HYM11506088).  
2043 **COLOMBIA: Caldas:** Aguadas, Cañón del Río Arma, 5.61472 -75.45972, 1995, C. Sarmiento  
2044 CES096, 3 workers (IAvH: USNMENT00757575); **Guajira:** Quebrada Guacoche, nr. Don  
2045 Diego, forest, 10.72305 -72.96972, 10 m, 22 July 1976, W.L. Brown & R.C. Kugler, 8 workers  
2046 (MCZC: USNMENT00757565, CMOS000094-CMOS000096); **Huila:** 4 km NE Rivera, 30 Dec.  
2047 1986, W. MacKay #9023, 3 workers (WEMC: USNMENT00758026); **Tolima:** Cunday, vereda  
2048 “El Eden”, 4.08333 -74.66667, 450 m, 01 Oct. 1999, Mejia et al., 2 workers (ICN:  
2049 MPUJ\_ENT0000416); **Valle del Cauca:** 08 Jan. 1976, W. & E. MacKay, 2 workers (WEMC:

2050 USNMENT00758162). **GUATEMALA: El Progreso:** 5 km W Morazan, 14.93 -90.20, 800 m,  
2051 19 Nov. 2003, A.L. Wild #AW2121, 2 workers (ALWC: USNMENT00757576). **FRENCH**  
2052 **GUIANA:** Basse Vie-foret, 04 July 1999, S. Durou, 2 workers, 1 queen (CEPLAC:  
2053 USNMENT00757568); Petit Saut, May 2003, J. Orivel & J. Le Breton, 3 workers (CEPLAC:  
2054 USNMENT00757573). **PANAMA: Barro Colorado,** Canal Zone, Jan. 1960, W.L. Brown &  
2055 E.S. McCluskey, 3 workers (MCZC: USNMENT00758033). **PARAGUAY: Amambay:** Parque  
2056 Nacional Cerro Corá, -22.65 -56.05, 13 May 1997, A. Wild #AW0576, 3 workers (ALWC:  
2057 USNMENT00757569); **Boqueron:** Enciso N.P. (Southern side), -21.20609 -61.65748, 01-02  
2058 Oct. 2002, T. Delsinne, 2 workers (RBINS: Coll.RIScNB SID SPM\_ID11523); Enciso N.P.  
2059 (Southern side), -21.20609 -61.65748, 01-02 Oct. 2002, M. Leponce, 1 worker (RBINS:  
2060 Coll.RIScNB SID SPM\_ID31985); Estancia Maria Vicenta, -20.92130 -61.39321, T. Delsinne, 1  
2061 worker (RBINS: Coll.RIScNB SID SPM\_ID26822); **Canindeyú:** Residencias, 6 km N Ygatimi, -  
2062 24.06667 -55.63333, 21 Feb. 1997, A. Wild #AW0451, 1 worker (AWLC:  
2063 USNMENT00757577). **TRINIDAD AND TOBAGO:** Cumuto, 24 Apr. 1929, Darlinhton, 1  
2064 worker (MCZC: USNMENT00757578).

2065

2066 **Diagnosis.** *Brachymyrmex degener* morphologically resembles *B. coactus* as both species have  
2067 scapes that surpass the posterior margin of the head, they have faint sculpture on the mesosoma, a  
2068 mesonotum that is inflated and that bulges dorsally above the pronotum in lateral view, a wide  
2069 metanotal groove, metathoracic spiracles that are slightly protruding dorsally, and their gasters  
2070 have sparse pubescence. However, *B. degener* has a uniformly brownish body, whereas the gaster  
2071 is conspicuously darker than the rest of the body in *B. coactus*.

2072



2073 *Lectotype and paralectotypes measurements* (mm) (n=3). HL<sub>1</sub> 0.51-0.55; HL<sub>2</sub> 0.35-0.41; HL<sub>3</sub>  
2074 0.16; HW 0.49-0.55; SL 0.37-0.53; EL 0.12-0.14; WL 0.55-0.68; PnL 0.16-0.20; PnW 0.31-0.37;  
2075 ML 0.12-0.16; MW 0.20-0.23; *Indices* CI 96.15-100.00; SI<sub>1</sub> 76.00-96.43; SI<sub>2</sub> 105.56-135.00; OI<sub>1</sub>  
2076 21.43-25.93; OI<sub>2</sub> 28.57-30.77.

2077  
2078 *Additional material examined measurements* (mm) (n=24). HL<sub>1</sub> 0.53-0.70; HL<sub>2</sub> 0.29-0.49; HL<sub>3</sub>  
2079 0.12-0.20; HW 0.51-0.73; SL 0.55-0.68; EL 0.12-0.20; WL 0.60-0.79; PnL 0.14-0.23; PnW 0.33-  
2080 0.50; ML 0.12-0.21; MW 0.20-0.31; *Indices* CI 87.50-112.50; SI<sub>1</sub>82.22-117.86; SI<sub>2</sub>137.04-  
2081 233.33; OI<sub>1</sub> 20.00-30.30; OI<sub>2</sub> 22.22-33.33.

2082  
2083 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
2084 concave. Dorsum of the head has scattered appressed hairs. Clypeus with a rounded anterior  
2085 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
2086 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
2087 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
2088 margin in oblique anterodorsal view. The scapes surpass the posterior cephalic margin by a  
2089 length smaller or equal to the maximal diameter of the eye, and they bear appressed and  
2090 decumbent hairs. Three ocelli are present. The eyes are positioned on the cephalic midline and  
2091 have 8-14 ommatidia along their maximal diameter.

2092 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum;  
2093 sometimes with additional suberect hairs, mainly on the pronotum. The mesonotum is inflated  
2094 and bulges dorsally above the pronotum in lateral view. Metanotal groove wider than the  
2095 diameter of the metathoracic spiracles. Metathoracic spiracles in dorsolateral position, slightly  
2096 protruding, and not touching any sutures. Dorsum of the propodeum strongly convex and shorter

2097 than the posterior slope. Propodeal spiracles conspicuous and circular, positioned on the  
2098 propodeal margin or just dorsal of it, at the anterior margin of the propodeal slope. Legs with  
2099 appressed hairs. Petiole short and inclined forward.

2100 **Gaster.** With scattered pubescence and several scattered long erect hairs.

2101 **Color and sculpture.** Body shiny and uniformly brownish in color. Head and gaster smooth  
2102 whereas the dorsum of the mesosoma usually bears imbricate sculpture.

2103

2104 **Distribution (supplementary material Fig. S14).** *Brachymyrmex degener* occurs in Brazil,  
2105 Colombia, Guatemala, French Guiana, Trinidad and Tobago.

2106

2107 **Biology.** Unknown.

2108

2109 **Remarks.** The lectotype of *B. degener* is the top specimen on pin USNMENT00758155, whereas  
2110 the others on that pin are paralectotypes.

2111 *Brachymyrmex degener* was first described as a subspecies of *B. coactus* by Emery (1906) and  
2112 Santschi (1923a) subsequently raised it to species. *Brachymyrmex degener* indeed resembles *B.*  
2113 *coactus* closely, and in retrospect we are uncertain that it represents a separate species. However,  
2114 the issue of *B. degener* and *B. coactus* is taxonomically very complex and involves several other  
2115 previously described species and subspecies that warrant synonymization too. Perhaps this  
2116 process is best performed incrementally, which is the approach taken here.

2117 One of these other taxa involved is *B. admotus* r *niger* which was described by Forel (1912a)  
2118 with the following diagnostic traits: the metanotal groove is deep, the mesonotum bulges out  
2119 dorsally above the pronotum (in lateral view), the body is shiny and the head and gaster smooth  
2120 whereas faint sculpture is present on the mesosoma. Upon examination, the type specimens of *B.*

2121 *admotus* r *niger* have the diagnostic traits of *B. degener*, but lack some of those for *B. admotus*,  
2122 such as the presence of a pair of thin erect hairs between the methathoracic spiracles, which are  
2123 positioned fully dorsal instead of dorsolaterally, and the absence of sculpture on the mesosoma.  
2124 These criteria that motivated Santschi (1923a) to reclassify the race as *B. degener* st. *niger*, and  
2125 whereas we agree with this reclassification, we do not consider there to be sufficient differences  
2126 to prevent synonymization of *niger* to *B. degener*.

2127 In the description of *B. incisus* Forel (1912a) likewise indicated similarities to *B. coactus* and *B.*  
2128 *admotus*, but again some of the diagnostic traits of *B. admotus* are absent. Moreover, whereas the  
2129 specimens resemble *B. coactus* closely they do not have a gaster that is conspicuously darker in  
2130 color than the head and mesosoma, and hence *B. incisus* is here synonymized to *B. degener*.

2131 As to *B. luederwaldti*, Santschi (1923a) indicated similarities to *B. coactus*, and even more so to  
2132 *B. admotus* r. *niger* (Forel 1912a), from which he distinguished *B. luederwaldti* mainly by its  
2133 fainter propodeal suture. We consider this variation to be intraspecific here and consequently also  
2134 synonymize *B. luederwaldti* to *B. degener*.

2135 Upon describing *degener* as a variety of *B. coactus*, Emery (1906) indicated that this variety  
2136 differs from typical *B. coactus* mainly in body and eye size, which are traits with a lot of  
2137 intraspecific variation as we already indicated in the remarks of *B. coactus*. We did not find  
2138 consistent differences in body size, nor in the number of ommatidia along the maximal diameter  
2139 of the eye between both taxa. Emery (1906) further emphasized that the medial antenomereres are  
2140 somewhat longer than wide in *B. degener*, and vice versa in *B. coactus*. However, we cannot  
2141 confirm this putative difference from the type material of both species. Although some putatively  
2142 diagnostic differences between *B. coactus* and *B. degener* are indicated in the diagnosis, the  
2143 taxonomic importance of these differences remains to be examined. Our morphometric  
2144 measurements confirm the difficulty to establish consistent differences between both taxa, and

2145 furthermore our phylogenetic analyses (see below) recovered *B. degener* nested within *B.*  
2146 *coactus*. However, a deep phylogenetic branch separates *B. degener* from *B. coactus* as is also  
2147 recognized by in the ABGD analysis (see below), and given the taxonomic complexity  
2148 surrounding *B. coactus* it is possible that the one specimen identified as *B. coactus* that renders  
2149 the species paraphyletic in fact belongs to a distinct taxon. With the sampling that is currently  
2150 available, this issue cannot be resolved and therefore we do not further synonymize *B. coactus*  
2151 and *B. degener* here.

2152

2153

2154 *Brachymyrmex delabiei* Ortiz & Fernández

2155 (Fig. 26, supplementary material Fig. S15)

2156 *Brachymyrmex delabiei* Ortiz and Fernández, 2014: 24, Figs. 22, 23, 24 (w). **Holotype worker**  
2157 (MZSP: USNMENT00757718) and **paratype workers** (CPDC: USNMENT00757719, ICN:  
2158 USNMENT00757720, USNM: USNMENT00757721): 4 workers. **BRAZIL: São Paulo:**  
2159 Tapiraí, -24.03208 -47.65556, 08-14 Jan. 2001, R.R. Silva & Eberhardt.

2160

2161 *Additional material examined.* **BRAZIL: Bahia:** Boa Nova, João Mata, 13 Aug. 2003, J.R.M.  
2162 Santos & J.C.S. Carmo, 1 worker (CPDC: USNMENT00757610); A61 Camacan, 27 Aug. 1999,  
2163 -15.60111 -39.52111, col. J.R.M. dos Santos, 1 worker (CPDC: USNMENT00757837); **Santa**  
2164 **Catharina:** Palhoça, PE Serra do Tabuleiro, 02–10 Nov. 2003, -27.74111 -48.69722, R.R. Silva,  
2165 B.H. Dietz and A. Tavares, 1 worker (MZSP: USNMENT00757725); **São Paulo:** São Bernardo  
2166 do Campo, 01 June 1971, W.L. & D.E. Brown, 1 worker (MCZC: USNMENT00757835).

2167

2168 **Diagnosis.** *Brachymyrmex delabiei* is most similar in morphology to *B. brasiliensis* and *B.*  
2169 *feitosai*, because they all have tumuliform metathoracic spiracles, however, it differs from *B.*  
2170 *brasiliensis* by its entirely smooth and shiny body, and from *B. feitosai* by the presence of two  
2171 erect hairs on the pronotum and two on the mesonotum, the lack of dense pubescence on the first  
2172 segment of the gaster and the yellowish body.

2173

2174 **Description.** See Ortiz and Fernández (2014).

2175

2176

2177 *Brachymyrmex depilis* Emery

2178 (Fig. 27, supplementary material Fig. S16)

2179 *Brachymyrmex heeri* subsp. *depilis* Emery, 1893: 635 (w.q.). **Lectotype worker** (MCSN:  
2180 USNMENT00757228) and **paralectotype workers, queen, male** (MCSN: USNMENT00757225  
2181 - 00757232; **here designated**): 37 workers, 1 queen, 10 males [examined]. **USA: District of**  
2182 **Columbia:** Georgetown College, 10 Aug. 1885, leg. Pergrande. Wheeler and Wheeler (1953:  
2183 139) (l.). Raised to species: Santschi (1923a: 663).  
2184 = *Brachymyrmex nanellus* Wheeler, 1903: 102, Fig. 7b (w.m.). (MCZC: MCZ Cotype 22939): 5  
2185 workers [examined]. **USA: Texas:** Austin, 25 May 1901. Synonymy by Creighton (1950: 359).  
2186 = *Brachymyrmex depilis* subsp. *flavescens* Grundmann, 1952: 117 (w.). (USNM:  
2187 USNMENT00529204): 3 workers [examined]. **USA: Utah:** near Salt Lake City. Lower portion  
2188 of Big Cottonwood Canyon, 24 June 1947. Synonymy by Cole (1953: 266).

2189

2190 **Additional material examined. CANADA: Nova Scotia:** Halifax, 15 m, 44.63333 -63.61667,  
2191 25 Oct. 1996, P.S. Ward #13234, 2 workers, 1 queen (PSWC: USNMENT00757818). **MEXICO:**  
2192 **Tamaulipas:** Gomez Parias, 25 Sep. 1987, W. MacKay #10073, 2 workers (WEMC:  
2193 USNMENT00757816); **Veracruz:** Las Hamacas, 17 km. N Santiago Tuxtla, 26-28 Aug. 1953,  
2194 E.O. Wilson, 5 workers, 1 queen (MCZC: CMOS000114 - 000115); Los Tuxtlas, 10 km NNW  
2195 Sontecomapan, 18.58333 -95.08333, 200 m, 20 Mar. 1985, P.S. Ward #7333-55, 3 workers  
2196 (PSWC: USNMENT00757815). **UNITED STATES: Alabama:** Marshall Co. JCT 420 7 km S  
2197 Morgna city, 34.41111 -86.52361, 09 June 1998, MacKay fam. #188203, 2 workers (WEMC:  
2198 USNMENT00757813); **Arkansas:** Cross Co. Village Cr. St. Pk. 14 Aug. 1988, R. Anderson, 3  
2199 workers (WEMC: USNMENT00757805 - 00757807); **California:** 8 km S, Brans. Wiask, 10 Feb.  
2200 1943, W.S. Ross, ANTC10266, 4 workers (CASENT: CASENT0196008); Santa Barbara Co,  
2201 Figueroa Crk., Sedgwick Ranch, 34.71667 -120.03333, 350 m, 02 Mar. 1996, P.S. Ward #12963,  
2202 3 workers (PSWC: USNMENT00757590); **District of Columbia:** Washington D.C. 25 May  
2203 1948, F. Bonet #1718, 3 workers (MZUSP: USNMENT00757798). **Florida:** Highlands Co.  
2204 Archbold Biol. Station, 22 Aug. 1995, A. Wild, 5 workers (ALWC: USNMENT00757817);  
2205 **Kentucky:** Floyd Co. Jennie Wiley St. Pk., 07 July 1968, S. Peck Ber #134, 1 worker (MCZC:  
2206 CMOS000028); **Louisiana:** Tammany Par. Abita, Springs, Money Hills Golf Course, 30.55156 -  
2207 89.95488, 08 Sep. 2000, A.M. Pranschke, 2 workers (CEPLAC: USNMENT00757801); **New**  
2208 **Mexico:** Sandoval Co, Bandelier, Nat. Mon, 21 Aug. 1986, W. & E. MacKay #8784, 2 workers  
2209 (WEMC: USNMENT00757814); **New York:** Newark, Morris Farm, U. Delaware,  
2210 Liriadendrofagus, 18 Apr. 1976, S. Handel, 2 workers (MCZC: CMOS000116, 000117); Ontario  
2211 Co. Gannet Hill, 42.7 -77.4, 640 m, 27-29 Aug 2003, A.L. Wild #AW1970, 2 workers, 1 queen  
2212 (ALWC: USNMENT00757799); Ontario Co. Gannet Hill, 42.7 -77.4, 640 m, 10 Sep. 1995, A.L.  
2213 Wild #AW0719, 3 workers (ALWC: USNMENT00757799); **Texas:** 16 km S San Antonio, 18

2214 Feb. 1942, E.S. Ross, ANTC10267, 3 workers (CASENT: CASENT0196009); Houston Co. Big  
2215 Stough Wild Area, 09 May 1988, R. Anderson #12760, 4 workers (WEMC:  
2216 USNMENT00757811 - 00757812, 00758040); Sabino Co. 14.5 K E Nerwphill, 11 May 1988, R.  
2217 Anderson #12763, #12763, 6 workers (WEMC: USNMENT00757808 - 00757810); **Vermont:**  
2218 Nr. Burlington, Temperate Forest, Nov 2001, R. Blatrix, 9 workers (CEPLAC:  
2219 USNMENT00757802 - 00757804).

2220  
2221 **Diagnosis.** *Brachymyrmex depilis* resembles *B. heeri* and *B. giardi* as all three taxa have the  
2222 mesonotum bulging dorsally above the pronotum in lateral view, and a gaster with dense  
2223 pubescence. However, *B. depilis* differs from *B. heeri* by its shorter scapes and the lack of erect  
2224 hairs on the mesosoma, and from *B. giardi* by its smaller eyes, its appressed hairs on the dorsum  
2225 of the mesosoma, its yellowish color and its Nearctic distribution, i.e from the South of Canada to  
2226 the North of Mexico.

2227  
2228 *Lectotype and paralectotypes measurements* (mm) (n=10). HL<sub>1</sub> 0.47-0.49; HL<sub>2</sub> 0.33-0.37; HL<sub>3</sub>  
2229 0.10-0.18; HW 0.39-0.45; SL 0.35-0.41; EL 0.08-0.12; WL 0.39-0.51; PnL 0.12-0.20; PnW 0.27-  
2230 0.33; ML 0.10-0.14; MW 0.20-0.21; *Indices* CI 80.00-92.00; SI<sub>1</sub> 85.71-100.00; SI<sub>2</sub> 100.00-  
2231 117.65; OI<sub>1</sub> 18.18-30.00; OI<sub>1</sub> 20.00-36.00.

2232  
2233 *Additional material examined measurements* (mm) (n=10). HL<sub>1</sub> 0.31-0.50; HL<sub>2</sub> 0.19-0.35; HL<sub>3</sub>  
2234 0.05-0.14; HW 0.29-0.46; SL 0.27-0.42; EL 0.08-0.10; WL 0.31-0.46; PnL 0.08-0.18; PnW 0.22-  
2235 0.31; ML 0.07-0.13; MW 0.17-0.22; *Indices* CI 88.89-94.74; SI<sub>1</sub> 85.71-95.74; SI<sub>2</sub> 110.71-142.86;  
2236 OI<sub>1</sub> 19.57-30.30; OI<sub>2</sub> 17.14-30.00.

2237

2238 **Description. *Head.*** Slightly longer than wide in full face view; posterior cephalic margin slightly  
2239 concave. Clypeus with a rounded anterior margin and five long, erect hairs of which a single,  
2240 usually conspicuous hair is near the anterior margin, two hairs are in mediolateral position and  
2241 two more near the toruli; other hairs on the clypeus are markedly shorter and appressed or  
2242 decumbent. Toruli surpassing the posterior clypeal margin in oblique anterodorsal view. The  
2243 scapes are short, usually barely reaching the posterior margin of the head, and never surpassing it  
2244 by a length that equals the maximal diameter of the eye. Ocelli are absent. Eyes are small and  
2245 positioned on the cephalic midline; they have 6-8 ommatidia along their maximal diameter.

2246 ***Mesosoma.*** Not bearing any erect hairs. The mesonotum is inflated and bulges dorsally above the  
2247 pronotum in lateral view. Metanotal groove absent or narrower than the diameter of the  
2248 metathoracic spiracles. Metathoracic spiracles are small, in dorsolateral position, not protruding,  
2249 but touching the propodeal suture. Dorsum of the propodeum is weakly convex and much shorter  
2250 than the propodeal slope. Propodeal spiracles are circular, positioned on the posterior propodeal  
2251 margin at the middle of the propodeal slope. Legs with appressed hairs. Petiole short and inclined  
2252 forward.

2253 ***Gaster.*** With dense pubescence and scattered long erect hairs at the edges of the segments.

2254 ***Color and sculpture.*** Body opaque with inconspicuous sculpture. Body yellowish, sometimes  
2255 with the gaster a bit darker than the mesosoma.

2256

2257 **Distribution (supplementary material Fig. S16).** *Brachymyrmex depilis* is known from Canada,  
2258 México, and the United States.

2259

2260 **Biology.** Grundmann (1952) collected a nest of *B. depilis* subsp. *flavescens* from among the roots  
2261 of the scrub oak *Quercus gambelii* and suggested that this species is subterranean and tends



2262 aphids and coccids on the roots of plants. This association was also highlighted by Yensen et al.  
2263 (1980) and Wheeler and Wheeler (1986). Small colonies of *B. depilis* were found in the soil  
2264 under stones or in rotting wood in a wide variety of habitats: open forest, dense moist forest,  
2265 grass lands, and dry fields (Wheeler and Wheeler 1986). Surprisingly, Yensen et al. (1980)  
2266 reported *B. depilis* from an intertidal halophyte-covered mud flat in the Gulf of California in  
2267 Mexico, where colonies are regularly inundated by sea water. The authors suggest that the  
2268 mechanisms that allow the species to survive heavy rains elsewhere may have preadapted their  
2269 survival in this unusual habitat. Page (1982) reported on copulatory behavior and observed a  
2270 queen of *B. depilis* with attached to her abdomen three motionless males, that she dragged around.  
2271 As such he suggested that *B. depilis* seems to have multiple copulations but whether insemination  
2272 occurs by several partners is unknown.

2273

2274 **Remarks.** The lectotype is designated here as the worker on pin USNMENT00757229 and the  
2275 other specimens are paralectotypes.

2276 *Brachymyrmex depilis* was originally described as a subspecies of *B. heeri* by Emery (1893), and  
2277 he distinguished it from typical *B. heeri* because *B. depilis* lacks erect hairs on the mesosoma.  
2278 Santschi (1923a) raised the subspecies to species but did not provide criteria to support his  
2279 decision. Nevertheless, we consider this decision justified given the differences we mention here  
2280 in the diagnosis.

2281 In the original description of *B. nanellus* Wheeler (1903) reported a comparison of his material to  
2282 alleged specimens of *B. depilis*, and he described a series of differences to support the status of *B.*  
2283 *nanellus* as a separate species. Santschi (1923a) accepted this taxonomic decision, but as  
2284 Creighton (1950) pointed out the comparative material unlikely belonged to *B. depilis*, and after a

2285 further comparison of both taxa he synonymized *B. nanellus* with *B. depilis*, which we support  
2286 here after re-examining the material.  
2287 *Brachymyrmex depilis* subsp. *flavescens* was originally distinguished from *B. depilis* by having a  
2288 lighter body color, smaller eyes, an opaquer body due to its shriveled integument, scarcer  
2289 pubescence and hairs (Grundmann 1952). However, after examining this material we agree with  
2290 the conclusion of Cole (1953) that these specimens appear to be part of an incipient colony,  
2291 which adequately explains all these morphological differences outlined by Grundmann (1952).  
2292 Fisher and Cover (2007) suggested that *B. depilis* may constitute a complex of several species.  
2293 The material studied here is perhaps to limited to accurately comment on this issue, however, we  
2294 did not find consistent morphological differences between samples, except perhaps in body size.

2295

2296

2297 *Brachymyrmex donisthorpei* Santschi

2298 (Fig. 28, supplementary material Fig. S17)

2299 *Brachymyrmex donisthorpei* Santschi, 1939: 320, Figs. 4, 5 (w.). **Lectotype worker** (NHMB:  
2300 USNMENT00757183) and **paralectotype workers** (NHMB: USNMENT00757184-00757185;  
2301 **here designated**): 3 workers [examined]. **COLOMBIA**, Mar. 1937, Paul Robá, leg.

2302

2303 **Additional material examined. BRAZIL: Bahia:** Vargito, -15.40 -39.55, 22 Mar. 1999, J.R.M.  
2304 dos Santos, 1 worker (CEPLAC: USNMENT00757839); **São Paulo:** Iguape, E.E. Jureia-Itatins,  
2305 Nucleo Rio verde, -24.54417 -47.23556, 5-14 Mar. 2001, A.A. Tavares, 1 worker (ICN:  
2306 MZSP158). **COLOMBIA: Magdalena:** El Campano, 11.12 -74.10, 1300 m, 13 Aug. 1985. P.S.  
2307 Ward #7891-23, 2 workers, 1 queen (PSWC: USNMENT00757840); **Nariño:** territorio Kofan,

2308 0.47481 -77.17913, 1000 m, 28 Sep. 1998, 1 worker (IAvH). **ECUADOR: Zamora-Chinchi:**  
2309 Copalinga, -4.09122 -78.96069, 17-19 Oct. 2009, Jacquemin, 1 worker (RBIN: Coll.RIScNB SID  
2310 SPM\_ID3753921). **PARAGUAY: Boquerón:** Enciso N.P. (Southern side), -21.20298 -  
2311 61.65909, 04-06 Nov. 2001, M. Leponce.

2312

2313 **Diagnosis.** *Brachymyrmex donisthorpei* morphologically resembles *B. modestus* and *B. myops*  
2314 because they all have dense, short pubescence over the entire body, scapes with short suberect  
2315 hairs, eyes that are positioned below the cephalic midline, a metanotal groove that is either  
2316 absent or narrower than the diameter of the metathoracic spiracles, and yellowish body color.  
2317 *Brachymyrmex donisthorpei* differs from *B. modestus* and *B. myops* by its short scapes that  
2318 approximately reach the posterior margin of the head or surpass it by less than the maximal  
2319 diameter of the eye.

2320

2321 *Lectotype and paralectotype measurements* (mm) (n=2). HL<sub>1</sub> 0.39-0.41; HL<sub>2</sub> 0.25-0.27; HL<sub>3</sub>  
2322 0.08; HW 0.33-0.35; SL 0.27-0.29; EL 0.05; WL 0.39; PnL 0.10; PnW 0.27-0.29; ML 0.10; MW  
2323 0.20; *Indices* CI 85.00-85.71; SI<sub>1</sub> 82.35-83.33; SI<sub>2</sub> 107.14-107.69; OI<sub>1</sub> 13.89-14.71; OI<sub>2</sub> 19.05-  
2324 20.00.

2325

2326 *Additional material examined measurements* (mm) (n=4). HL<sub>1</sub> 0.32-0.38; HL<sub>2</sub> 0.22-0.29; HL<sub>3</sub>  
2327 0.07-0.09; HW 0.26-0.33; SL 0.24-0.29; EL 0.04; WL 0.27-0.38; PnL 0.11-0.12; PnW 0.20-0.23;  
2328 ML 0.06-0.09; MW 0.13-0.16; *Indices* CI 80.55-86.05; SI<sub>1</sub> 86.49-96.55; SI<sub>2</sub> 96.97-112.00; OI<sub>1</sub>  
2329 13.51-16.67; OI<sub>2</sub> 22.22-23.26.

2330

2331 **Description. Head.** Substantially longer than wide in full face view; posterior cephalic margin  
2332 slightly concave. Clypeus with a rounded anterior margin and five long, erect hairs of which a  
2333 single, usually conspicuous hair is near the anterior margin, two hairs are in mediolateral position  
2334 and two more near the toruli; other hairs on the clypeus are markedly shorter and appressed or  
2335 decumbent. Toruli surpassing the posterior clypeal margin in oblique anterodorsal view. Dorsum  
2336 of the head has conspicuous appressed pubescence and several suberect hairs. The scapes  
2337 approximately reach the posterior margin of the head or surpass it by a length smaller than the  
2338 maximal diameter of the eye, and they have appressed and decumbent hairs. Ocelli absent. The  
2339 eyes are positioned below the cephalic midline and have only 3-4 ommatidia along their maximal  
2340 diameter.

2341 **Mesosoma.** With several short appressed and sub-erect hairs. The mesonotum is not inflated and  
2342 does not bulge dorsally above the pronotum in lateral view. Metanotal groove absent or narrower  
2343 than the diameter of the metathoracic spiracles. Metathoracic spiracles dorsolateral in position,  
2344 not protruding, and touching the propodeal suture. Dorsum of the propodeum flat and much  
2345 shorter than propodeal declivity. Propodeal spiracles circular, small and inconspicuous,  
2346 positioned on the posterior propodeal margin, slightly posterior of the middle of the propodeal  
2347 slope. Legs with appressed and sub-erect hairs. Petiole short and inclined forward.

2348 **Gaster.** With appressed dense pubescence and some sub-erect hairs near the edges of the  
2349 segments.

2350 **Color and sculpture.** Body yellowish, with imbricate sculpture on the dorsum of the mesosoma.

2351

2352 **Distribution (supplementary material Fig. S17).** *Brachymyrmex donisthorpei* is known from  
2353 Brazil, Colombia, Ecuador and Paraguay.

2354

2355 **Biology.** Unknown.

2356

2357 **Remarks.** No specific geographic information is available on the type material beyond

2358 Colombia.

2359

2360

2361 *Brachymyrmex feitosai* Ortiz & Fernández

2362 (Figs. 29, supplementary material Fig. S18)

2363 *Brachymyrmex feitosai* Ortiz and Fernández, 2014: 27, Figs. 25, 26, 27 (w). **Holotype worker**

2364 and **paratype workers** (MZSP: USNMENT00757694): 3 workers. **BRAZIL: Rio de Janeiro:**

2365 Floresta de Tijuca, D. Federal. 16 Dec. 1959, C.A: Campos Seabra.

2366

2367 **Additional material examined. BRAZIL: Minas Gerais:** Lavras, Ijaci e Perdões, -21.24528 -

2368 44.99972, Fragmento, 06 Dec. 2002, M.S. Santos & N.S. Dias, 4 workers (CPDC:

2369 USNMENT00757836, 00759008); **São Paulo:** Piedade, Floresta Atlantica "Theomar", -23.73846

2370 -47.38957, 16 Nov. 2008, G. Bieber, 3 workers (ICN: USNMENT00759038); Sete Barras, PE

2371 Carlos Botelho, 600 m, -24.20833 -47.97056, 11–15 May 2009, armadilha subterrânea #18, F.

2372 Esteves et al., 1 worker (MZSP: ANTWEB CASENT0217326).

2373

2374 **Diagnosis.** *Brachymyrmex feitosai* resembles *B. brasiliensis* and *B. delabiei* because they all have

2375 tumuliform metathoracic spiracles. However, *B. feitosai* differs from *B. brasiliensis* by its

2376 entirely smooth and shiny body, that is more brownish, and by the dense yellowish pubescence

2377 on the first gastral segment. It differs from *B. delabiei* by the presence of many suberect hairs on  
2378 the pronotum and mesonotum and its dense yellowish pubescence on the first gastral segment.

2379

2380 **Description.** See Ortiz and Fernández (2014).

2381

2382

2383 *Brachymyrmex fiebrigi* Forel

2384 (Figs. 30, 31, supplementary material Fig. S19)

2385 *Brachymyrmex fiebrigi* Forel, 1908: 400 (w.). **Lectotype worker** (MHNG:

2386 USNMENT00757164) and **paralectotype workers** (MHNG: USNMENT00757164-00757165;

2387 **here designated**): 4 workers [examined]. **PARAGUAY:** San Bernardino, Fiebrig leg. Santschi

2388 (1922: 260) (q.m.). See also: Santschi (1923a: 661).

2389 = *Brachymyrmex fiebrigi* var. *funicularis* Santschi, 1922: 260 (w.). (NHMB:

2390 USNMENT00757180-007581, 00758094): 22 workers [examined]. **ARGENTINA: Córdoba:**

2391 Alta Gracia. **n. syn.**

2392 = *Brachymyrmex fiebrigi* var. *fumida* Santschi, 1923a: 661 (w.). (MHNB: USNMENT00757704,

2393 00758157): 4 workers [examined]. **ARGENTINA: Buenos Aires:** Cerro “Ruinas”; (MHNB:

2394 USNMENT00758093, NHMG: USNMENT00758153): 3 workers [examined]. **ARGENTINA:**

2395 **Cordoba:** La Cabana. **n. syn.**

2396

2397 **Additional material examined. ARGENTINA: Cordoba:** Alta Gracia, Bruchi, 2 workers

2398 (MZUSP: USNMENT00757548). **BRAZIL: Bahia:** Canavieiras, -15.56361 -39.01722, 24 Aug.

2399 1998, J.C.S. Carmo & J.R.M. Santos, 1 worker (CEPLAC: USNMENT00757962; Esplanada

2400 Baixio, -12.11444 -37.69694, June-Oct. 2010, M.L.O. Travassos, #5644, 1 worker (CEPLAC:  
2401 USNMENT00757545). **Minas Gerais:** Lavras, 06-12 2002, M.S. Santos, N.S. Dias, 2 workers  
2402 (CEPLAC: USNMENT00759008). **São Paulo:** Iguape, EE Jureia-Itatins, Nucleo Rio verde, -  
2403 24.54417 -47.23556, 05-14 Mar. 2001, A.A. Tavares, 1 worker (ICN: MZSP158). **COSTA**  
2404 **RIKA: Heredia:** Cantarrana, 11 km ESE La Virgen, 10.33516 -84.04856, 300 m, 26 Feb. 2007,  
2405 Marcos-Deimer-Joel, 1 worker (JTLC: INBIO0003646597); **Limón:** Casa Verde, Tortuguero,  
2406 10.58333 -83.51667, 5 m, 24 June 1988, J. Longino #2154, 3 workers (JTLC:  
2407 INBIOCRI001280321, 001280326, 001280331); **Puntarenas:** 8 km WNW Potrero grande, 9.03 -  
2408 85.25, 200 m, 01 Aug. 1985, P.S. Ward #7792, 3 workers (PSWC: USNMENT00757549); La  
2409 Pita, rd. To Monteverde 10.16667 -84.91667, 120 m, 13 July 1984, J. Longino, 1 worker (JTLC:  
2410 JTLC000005902). **CUBA: Viñales:** Pinar del Rio, 14 June 1953, E.O. Wilson #10, 2 workers, 1  
2411 male (MCZC: USNMENT00757546). **MEXICO: Quintana Roo:** Municipio Leona Vicario,  
2412 Reserva Ecológica "El Edén", 21.21667 -87.18333, 03 July 1997, G.M. Daniel, 1 worker (ICN:  
2413 USNMENT00757626); **PARAGUAY: Boquerón:** Garrapatal, -21.44306 -61.87500, 04-06 Nov.  
2414 2001, M. Leponce, 1 worker (RBIN: Coll.RIScNB SID SPM\_ID14544); Garrapatal, -21.43965 -  
2415 61.48899, 05-06 Nov. 2001, M. Leponce, 1 worker (RBIN: Coll.RIScNB SID SPM\_ID25159);  
2416 Enciso, -21.20 -61.67, 03-06 Nov. 2001, M. Leponce & T. Delsinne #4123-9/3, 3 workers  
2417 (ALWC: ANTWEB CASENT0173481); Enciso N.P. (Southern side), -21.19978 -61.66084, 17-  
2418 18 Sep. 2003, T. Delsinne, 1 worker (RBIN: Coll.RIScNB SID SPM\_ID31851); Enciso N.P.  
2419 (Southern side), -21.19978 -61.66084, 04-06 Nov. 2003, M. Leponce, 1 worker (RBIN:  
2420 Coll.RIScNB SID SPM\_ID32154); Fortín Mayor Infante Rivarola, -21.67146 -62.41312, 02-06  
2421 Nov. 2001, M. Leponce, 1 worker (RBIN: Coll.RIScNB SID SPM\_ID30618); Mister Long, -  
2422 20.60386 -62.05053, 05-06 Nov. 2001, M. Leponce, 1 worker (RBIN: Coll.RIScNB SID  
2423 SPM\_ID25477); Mister Long, -20.60386 -62.05053, 17-18 Sep. 2003, T. Delsinne, 2 workers

2424 (RBIN: Coll.RIScNB SID SPM\_ID 26023, Coll.RIScNB SID SPM\_ID27108); Mister Long, -  
2425 20.60386 -62.05053, 01-04 Oct 2004, T. Delsinne, 1 worker (RBIN: Coll.RIScNB SID  
2426 SPM\_ID30953); Nueva Asunción, -20.68896 -61.92886, 17-18 Sep. 2003, Delsinne (RBIN:  
2427 Coll.RIScNB SID SPM\_ID27184); **Central:** Capiata, -25.35 -57.42, 22 Feb. 1994, B. Garcete  
2428 #ibn 197, 1 worker (ALWC: USNMENT00757544); **Itapúa:** Isla Yacyretá E Melgarejo, -22.42 -  
2429 56.50, 11 Nov. 1997, B. Barrios #ibn 217, 2 workers (ALWC: USNMENT00757891).  
2430 **SURINAME:** Paramaribo, Apr. 1959, I.v.d. Drif, 3 workers (MZUSP: USNMENT00757547).

2431  
2432 **Diagnosis.** *Brachymyrmex fiebrigi* morphologically resembles *B. depilis*, because they both have  
2433 short scapes that do not or just reach the posterior margin of the head, a gaster with dense  
2434 pubescence, a yellowish body and eyes that are positioned on the cephalic midline.  
2435 *Brachymyrmex fiebrigi* differs from *B. depilis* by its mesosoma, which usually bears several erect  
2436 hairs, two on the pronotum and two on the mesonotum and by its geographic distribution, which  
2437 ranges from the South of Mexico until Paraguay, including Cuba.

2438  
2439 *Lectotype and paralectotypes measurements* (mm) (n=3). HL<sub>1</sub> 0.35-0.37; HL<sub>2</sub> 0.23; HL<sub>3</sub> 0.08-  
2440 0.10; HW 0.31; SL 0.25-0.29; EL 0.08-0.10; WL 0.27-0.31; PnL 0.12; PnW 0.20-0.25; ML 0.06;  
2441 MW 0.16-0.20; *Indices* CI 87.21-88.89; SI<sub>1</sub> 81.25-93.75; SI<sub>2</sub> 108.33-125.00; OI<sub>1</sub> 25.00-31.25;  
2442 OI<sub>2</sub> 21.05-27.78.

2443  
2444 *Additional material examined measurements* (mm) (n=12). HL<sub>1</sub> 0.32-0.46; HL<sub>2</sub> 0.22-0.34; HL<sub>3</sub>  
2445 0.07-0.13; HW 0.27-0.41; SL 0.22-0.36; EL 0.08-0.10; WL 0.26-0.40; PnL 0.09-0.14; PnW 0.20-  
2446 0.28; ML 0.06-0.11; MW 0.15-0.20; *Indices* CI 75.71-88.46; SI<sub>1</sub> 80.00-93.75; SI<sub>2</sub> 96.77-115.38;  
2447 OI<sub>1</sub> 21.74-33.33; OI<sub>2</sub> 20.00-28.85.



2448

2449 **Description. *Head.*** Slightly longer than wide in full face view; posterior cephalic margin flat.

2450 Dorsum of the head with appressed hairs. Clypeus with a rounded anterior margin and five long,

2451 erect hairs of which a single, usually conspicuous hair is near the anterior margin, two hairs are in

2452 mediolateral position and two more near the toruli; other hairs on the clypeus are markedly

2453 shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique

2454 anterodorsal view. The scapes are short, usually approximately reaching the posterior margin of

2455 the head, and they bear appressed and decumbent hairs. Ocelli apparently absent. Eyes are

2456 positioned on the cephalic midline and have 6-9 ommatidia along their maximal diameter.

2457 ***Mesosoma.*** Typically with two erect hairs on the pronotum and two on the mesonotum. The

2458 mesonotum is not inflated and does not bulge dorsally above the pronotum in lateral view.

2459 Metanotal groove absent or narrower than the diameter of the metathoracic spiracles.

2460 Metathoracic spiracles dorsolateral in position, not protruding, and touching the propodeal suture.

2461 Dorsum of the propodeum flat and much shorter than the posterior slope. Propodeal spiracles

2462 circular, small and inconspicuous, positioned on the posterior propodeal margin slightly posterior

2463 of the middle of the propodeal slope. Legs with appressed hairs. Petiole short and inclined

2464 forward.

2465 ***Gaster.*** With dense pubescence and scattered suberect hairs at the edges of the segments.

2466 ***Color and sculpture.*** Body usually smooth, shiny and yellowish.

2467

2468 **Distribution (supplementary material Fig. S19).** *Brachymyrmex fiebrigi* is known from

2469 Argentina, Brazil, Costa Rica, Cuba, Mexico, Paraguay and Suriname.

2470

2471 **Biology.** Some specimens have been collected from below stones (PSWC:  
2472 USNMENT00757549); Forel (1908) suggested that this species nests in dry branches of bushes.  
2473  
2474 **Remarks.** The lectotype is the second ant from the top on pin USNMENT00757164, whereas the  
2475 other specimens on that pin are paralectotypes. Santschi (1922) considered *B. fiebrigi* var.  
2476 *funicularis* as a variety of *B. fiebrigi* mainly based on its darker-colored funiculus and posterior  
2477 segments of the gaster, but otherwise the type specimens of this variety are very similar in  
2478 measurements, head shape, and gastric pubescence compared to the type material of *B. fiebrigi*.  
2479 Overall, we consider these differences to represent intraspecific variation.  
2480 Santschi (1923a) considered *B. fiebrigi* var. *fumida* as a variety that only differs from the typical  
2481 *B. fiebrigi* by its somewhat darker body color, and the overall light yellowish scapes and tibia. As  
2482 for *B. fiebrigi* var. *funicularis* we consider these differences to represent intraspecific variation  
2483 and both varieties are synonymized here.

2484

2485

2486 *Brachymyrmex flavidulus* (Roger)

2487 (Supplementary material Fig. S20)

2488 *Plagiolepis flavidula* Roger, 1863: 162 (w.). **Lectotype worker** (MfNB: 19185: GBIF-  
2489 D/FoCol2900; GBIF-D/FoCol2910; **here designated**): 1 worker [examined]. **CUBA.** Attributed  
2490 to *Brachymyrmex* by Smith (1955: 99).

2491

2492 **Additional material examined. COLOMBIA: Valle del Cauca:** Bosque Yotoco, 1575 m, 25  
2493 June 1989, W.P. MacKay #11562, 2 workers (WEMC: USNMENT00757634). **COSTA RICA:**  
2494 **Puntarenas:** Monteverde, 10.29564 -84.79009, 1540 m, 10 Dec. 1987, J. Longino #1975-s, 1

2495 worker, 1 queen (JTLC: JTLC0000005251). **JAMAICA: Trelawny:** 5 km N Quick Step,  
2496 18.26667 -77.71667, 360 m, A.L. Wild #AW1382, 1 worker (ALWC: USNMENT00757658).

2497  
2498 **Diagnosis.** *Brachymyrmex flavidulus* resembles *B. fiebrigi*, *B. giardi* and *B. depilis* in that they all  
2499 have short scapes that approximately reach the posterior margin of the head or surpass it by less  
2500 than one maximal diameter of the eye, their gaster bears dense pubescence, their eyes are located  
2501 on the cephalic midline and their bodies are yellowish. *Brachymyrmex flavidulus* differs from *B.*  
2502 *depilis* and *B. giardi* by its mesonotum that does not bulge dorsally above the pronotum in lateral  
2503 view, and from *B. fiebrigi* by the absence of erect hairs on the pronotum and mesonotum.

2504  
2505 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
2506 convex. Clypeus with a rounded anterior margin and five long, erect hairs of which a single,  
2507 usually conspicuous hair is near the anterior margin, two hairs are in mediolateral position and  
2508 two more near the toruli; other hairs on the clypeus are markedly shorter and appressed or  
2509 decumbent. Toruli surpassing the posterior clypeal margin in oblique anterodorsal view. The  
2510 scapes are short and barely reach the posterior margin of the head. Ocelli apparently absent. Eyes  
2511 are positioned on the cephalic midline and have 7-9 ommatidia along their maximal diameter.

2512 **Mesosoma.** Without erect hairs. The mesonotum does not bulge dorsally above the pronotum in  
2513 lateral view. Metanotal groove absent. Metathoracic spiracles dorsolateral in position, not  
2514 protruding, and touching the propodeal suture. Dorsum of the propodeum shorter than posterior  
2515 slope. Propodeal spiracles circular, positioned on the posterior propodeal margin, slightly  
2516 posterior of the middle of the propodeal slope. Legs with appressed hairs. Petiole short and  
2517 inclined forward.

2518 **Gaster.** With dense pubescence.

2519 **Color and sculpture.** Body usually smooth, shiny and yellowish.

2520

2521 **Distribution (Supplementary material Fig. S20).** This species is known from Colombia, Costa  
2522 Rica, Cuba and Jamaica.

2523

2524 **Biology.** Unknown.

2525

2526 **Remarks.** *Brachymyrmex flavidulus* is a problematic species for several reasons. It was described  
2527 by Roger (1863) as a species of *Plagiolepis*. Smith (1955) transferred it to *Brachymyrmex*,  
2528 seemingly based in geographic reasons, i.e. that *Plagiolepis* is not native to the neotropics. We  
2529 agree with the attribution to *Brachymyrmex* because the type of *Brachymyrmex flavidulus* has 9  
2530 antennal segments, which is a diagnostic trait of *Brachymyrmex*, whereas *Plagiolepis* has 11  
2531 antennal segments (Bolton 2003). The type series consist of a single individual of which the  
2532 mesosoma and gaster are mounted on a pin, and the head is prepared on a microscope slide. This  
2533 preservation hampers us to document the arrangement of hairs on scapes, head, and clypeus as  
2534 well as the number of ommatidia in the maximal diameter of the eye.

2535 The worker on pin JTLC0000005251 is unusual in comparison to the other specimes of *B.*

2536 *flavidulus* in having sparser gastral pubescence and somewhat longer scapes. Additional material  
2537 from Costa Rica would be required to adequately characterise the morphological variation in  
2538 these populations, and to verify its species attribution.

2539 This species is morphologically very similar to *B. fiebrigi*. *Brachymyrmex flavidulus* lacks the  
2540 erect hairs on the mesosoma that are present in *B. fiebrigi*, but as the number of specimens  
2541 available of *B. flavidulus* is very limited we cannot currently comment on the consistency of this

2542 difference. An in-depth comparison with *B. fiebrigi* is required when more specimens of *B.*  
2543 *flavidulus* become available, especially from Cuba, where both species occur.

2544

2545

2546 *Brachymyrmex gagates* Wheeler

2547 (Fig. 32, supplementary material Fig. S21)

2548 *Brachymyrmex gagates* Wheeler, 1934: 206 (w.). **Lectotype worker** (USNM:

2549 USNMENT00529454) and **paralectotype workers** (USNM: USNMENT00529454; MCZC:

2550 M.C.Z. Cotype 1-3 21436, M.C.Z. Cotype 4-6 21436; **here designated**): 9 workers [examined].

2551 **MEXICO: Veracruz:** Mirador, 20 Apr. 1929.

2552

2553 **Additional material examined. PANAMA: Colon:** San Lorenzo Forest, 9.28333 -79.97194, J.

2554 Schmidt & J. Bail, fogging, 2 workers (ICN: USNMENT00759031).

2555

2556 **Diagnosis.** *Brachymyrmex gagates* resembles *B. degener* and *B. gaucho* in morphology, because

2557 they all have smooth, shiny, and dark brown or black bodies, scapes that surpass the posterior

2558 margin of the head, and a gaster with scarce pubescence. *Brachymyrmex gagates* differs from *B.*

2559 *degener* by its darker body and by having a mesonotum that is almost circular in dorsal view and

2560 that does not bulge above the pronotum in lateral view. It differs from *B. gaucho* by having a

2561 slightly concave posterior cephalic margin, scapes with decumbent hairs, a second segment of the

2562 antennal funiculus that is conspicuously shorter than the first antennal segment, and its almost

2563 circular mesonotum in dorsal view that does not bulge above the pronotum in lateral view.

2564

2565 *Lectotype measurements* (mm). HL<sub>1</sub> 0.59; HL<sub>2</sub> 0.39; HL<sub>3</sub> 0.18; HW 0.55; SL 0.51; EL 0.16; WL  
2566 0.59; PnL 0.18; PnW 0.39; ML 0.16; MW 0.20; *Indices* CI 93.33; SI<sub>1</sub> 92.86; SI<sub>2</sub> 130.00; OI<sub>1</sub>  
2567 28.57; OI<sub>2</sub> 30.00.

2568  
2569 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic border slightly  
2570 concave. Dorsum of the head with scattered appressed hairs. Clypeus with a rounded anterior  
2571 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
2572 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
2573 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
2574 margin in oblique anterodorsal view. The scapes bear decumbent hairs and surpass the posterior  
2575 margin of the head by a length smaller than the maximal diameter of the eye. Three ocelli  
2576 present. Eyes are positioned slightly posteriorly to the cephalic midline and have 10-12  
2577 ommatidia along their maximal diameter.

2578 **Mesosoma.** Typically bearing two erect hairs on the pronotum and two on the mesonotum,  
2579 sometimes with some additional appressed hairs on the dorsum of the mesonotum. The  
2580 mesonotum is inflated but does not bulge dorsally above the pronotum in lateral view; it is almost  
2581 circular in dorsal view. Metanotal groove wider than the diameter of the metathoracic spiracles.  
2582 Metathoracic spiracles in dorsolateral position, protruding slightly, and either just or just not  
2583 touching the propodeal suture. Dorsum of the propodeum convex and somewhat shorter than the  
2584 posterior declivity. Propodeal spiracles circular, positioned on the posterior propodeal margin,  
2585 anterior of the middle of the propodeal slope. Legs have appressed hairs. Petiole short and  
2586 inclined forward.

2587 **Gaster.** With scattered pubescence and several scattered long erect hairs.

2588 **Color and sculpture.** Head and gaster are smooth and shiny, whereas the dorsum of the  
2589 mesosoma is slightly imbricate. Body uniformly dark brown, apart from the terminal segments  
2590 of the tarsus and the hairs, which are lighter.

2591

2592 **Distribution (Supplementary material Fig. S21).** *Brachymyrmex gagates* is known from  
2593 Mexico and Panama.

2594

2595 **Biology.** The type specimens were collected from an epiphytic bromelia (*Tillandsia*  
2596 *streptophylla*) (Wheeler 1934).

2597

2598 **Remarks.** The lectotype is the ant at the top of pin USNM: USNMENT00529454, whereas the  
2599 others on that pin are paralectotypes. Wheeler (1934) pointed out that *B. gagates* is similar to but  
2600 nevertheless different from *B. incisus* (which is here synonymized to *B. degener*) of which he had  
2601 specimens from Panama in his collection. He reported that *B. gagates* has a wider head, a much  
2602 more prominent mesonotum, a more distinct and impressed promesonotal suture, longer funicular  
2603 joints and darker body color. We are uncertain as to what he exactly implied about the  
2604 promesonotal suture, because it is very distinctive in all *Brachymyrmex* species, and the  
2605 mesonotum bulges dorsally above the pronotum in lateral view in *B. degener* whereas it does not  
2606 in *B. gagates* (see diagnosis). We agree that generally *B. incisus* (and thus *B. degener*) generally  
2607 have much lighter body color than *B. gagates*, but as mentioned in the remarks of *B. degener* and  
2608 *B. coactus* the variation in body color in these species requires more detailed documentation.

2609

2610

2611 *Brachymyrmex gaucho* Santschi

2612 (Fig. 33, supplementary material Fig. S22)

2613 *Brachymyrmex gaucho* Santschi, 1917: 283 (w.). (NHMB) [examined, but the type is severely

2614 damaged]. **ARGENTINA: Córdoba:** Unquillo, M. Birabén. Combination in *Brachymyrmex*

2615 (*Brysha*) by Santschi (1923a: 674). See also: Quirán (2005: 767).

2616

2617 **Diagnosis.** *Brachymyrmex gaucho* is morphologically similar to *B. antennatus* because both have

2618 legs and antennae with erect hairs and a second segment of the antennal funiculus that is as long

2619 as or longer than the first. However, *B. gaucho* differs from *B. antennatus* by having a flat

2620 posterior cephalic margin, a dark brown body, erect hairs on the scape and the dorsum of the

2621 head, a mesonotum that bulges dorsally above the pronotum, and a gaster with scarce

2622 pubescence.

2623

2624 **Description. Head.** Almost equally wide as long in full face view; the posterior cephalic margin

2625 is flat and the posterior side of the head is wider than the anterior side. Dorsum of the head bears

2626 scattered erect hairs. Clypeus with a rounded anterior margin and five long, erect hairs of which a

2627 single, usually conspicuous hair is near the anterior margin, two hairs are in mediolateral position

2628 and two more near the toruli; other hairs on the clypeus are clearly shorter and decumbent. Toruli

2629 surpassing the posterior clypeal margin in oblique anterodorsal view. The scapes surpass the

2630 posterior cephalic margin by a length smaller than 1.5× the maximal diameter of the eye and have

2631 erect hairs. The second segment of the antennal funiculus is as long as or longer than the first.

2632 Three ocelli are present. Eyes are positioned on the cephalic midline and have 13-14 ommatidia

2633 along their maximal diameter.



2634 **Mesosoma.** With several thin, erect hairs. The mesonotum is inflated, anteroposteriorly inclined  
2635 and bulges dorsally above the pronotum in lateral view. Metanotal groove usually absent, but  
2636 when present narrower than the diameter of the metathoracic spiracles. Metathoracic spiracles in  
2637 dorsolateral position, slightly protruding, and near the propodeal suture. Dorsum of the  
2638 propodeum flat and equal in length to the propodeal slope. Propodeal spiracles circular,  
2639 positioned just dorsally of the posterior propodeal margin and slightly posterior of the middle of  
2640 the propodeal slope. Legs have suberect and erect hairs. Petiole short and inclined forward.

2641 **Gaster.** With scattered pubescence and several scattered long erect hairs.

2642 **Color and sculpture.** Head and gaster are smooth and shiny, whereas the dorsum of the  
2643 mesosoma is imbricate. Body uniformly dark brown.

2644

2645 **Distribution (Supplementary material Fig. S22).** *Brachymyrmex gaucho* is currently only  
2646 known from Argentina.

2647

2648 **Biology.** Unknown.

2649

2650 **Remarks.** The type specimen in the NHMB is damaged but Quirán deposited 3 workers  
2651 belonging to this species from Argentina, Córdoba II-2001, E. Quirán, leg. at the NHMB, which  
2652 we also examined. These specimens come from the same state as the type material, but they are  
2653 no types. We studied these specimens, one of which is illustrated in Fig. 33. More type material  
2654 would exist at the MACN (5 workers) and the MLP (1 worker) (Quirán 2005), but it is not  
2655 studied here.

2656

2657

2658 *Brachymyrmex giardi* Emery  
2659 (Fig. 34, supplementary material Fig. S23)  
2660 *Brachymyrmex giardi* Emery, 1895: 215 (w.q.). **Lectotype worker** (MSNG:  
2661 USNMENT00757220) and **paralectotype workers, putative worker-queen intercaste, queen,**  
2662 **male** (MSNG: USNMENT00757218 - 00757220, MHNG: USNMENT00758105-00758109;  
2663 **here designated**): 6 workers, 9 putative worker-queen intercastes, 1 queen, 1 male [examined].  
2664 **CHILE:** Santiago de Chile. Emery (1906: 178) (m.).  
2665 = *Brachymyrmex melensis* De Zolessi et al., 1978: 26 (w.q.l.). **URUGUAY: Cerro Largo: Melo.**  
2666 [not examined]. **n. syn.**  
2667  
2668 **Additional material examined. CHILE:** Valparaiso, 2 workers (MCZC: M.C.Z. Cotype22940).  
2669  
2670 **Diagnosis.** *Brachymyrmex giardi* resembles *B. depilis* and *B. heeri* as these species have a  
2671 mesonotum that bulges dorsally above the pronotum in lateral view, and a gaster with dense  
2672 pubescence. Additionally, *B. giardi* and *B. depilis* have bodies without erect hairs; they can be  
2673 distinguished because *B. giardi* has dense decumbent pubescence on the head and mesosoma,  
2674 usually a dark brownish body, and it is geographically restricted to Chile and Uruguay.  
2675 Furthermore, *B. giardi* differs from *B. heeri* by having short scapes, and no erect hairs on the  
2676 pronotum or mesonotum.  
2677  
2678 *Lectotype and paralectotype measurements* (mm) (n=8). HL<sub>1</sub> 0.45-0.68; HL<sub>2</sub> 0.29-0.42; HL<sub>3</sub>  
2679 0.10-0.32; HW 0.35-0.68; SL 0.33-0.59; EL 0.10-0.18; WL 0.39-0.89; PnL 0.10-0.21; PnW 0.25-  
2680 0.57; ML 0.08-0.35; MW 0.16-0.52; *Indices* CI 78.26-105.00; SL<sub>1</sub> 84.85-100.00; SL<sub>2</sub> 113.33-  
2681 142.86; OI<sub>1</sub> 23.53-33.33; OI<sub>2</sub> 21.74-50.00.

2682  
2683 **Worker description. Head.** Slightly longer than wide in full face view; posterior cephalic  
2684 margin is flat. Dorsum of the head has dense appressed hairs. Clypeus with a rounded anterior  
2685 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
2686 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
2687 clypeus are clearly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
2688 margin in oblique anterodorsal oblique view. The scapes surpass the posterior cephalic margin by  
2689 a length smaller than the maximal diameter of the eye and have decumbent hairs. Three ocelli  
2690 appear to be present. Eyes are positioned on the cephalic midline and have 7-9 ommatidia along  
2691 their maximal diameter.

2692 **Mesosoma.** Without erect hairs. The mesonotum is inflated and bulges dorsally above the  
2693 pronotum in lateral view. Metanotal groove usually absent, or narrower than the diameter of the  
2694 metathoracic spiracles. Metathoracic spiracles in dorsolateral position, not protruding, and  
2695 touching the propodeal suture. Dorsum of the propodeum slightly convex and shorter than the  
2696 posterior propodeal margin. Propodeal spiracles circular, positioned on the posterior propodeal  
2697 margin slightly posterior of the middle of the propodeal slope. Legs with appressed hairs. Petiole  
2698 short and inclined forward.

2699 **Gaster.** With dense pubescence and scattered long erect hairs along the edges of the segments.  
2700 Some specimens have the same morphology as a regular worker as to the head and mesosoma,  
2701 but have a gaster that is somewhat expanded, i.e. they are somewhat physogastric.

2702 **Color and sculpture.** Body smooth and shiny, usually dark brownish with yellowish legs and  
2703 pubescence.

2704

2705 **Intercaste description.** The morphology of the putative worker-queen intercaste differs from  
2706 that of the worker by its larger size, the eyes that have around 9 ommatidia along their maximal  
2707 diameter, the pronotum that bears several semi-erect hairs, the enlarged mesonotum that does  
2708 not bulge dorsally above the pronotum in lateral view, and that does not bear erect but several  
2709 subdecumbent hairs, the sharper posterior ending of the mesonotum in dorsal view, the deep  
2710 metanotal groove that is wider than the metathoracic spiracles, the almost laterally positioned  
2711 metathoracic spiracles, that do not protrude and do not touch any suture, and the uniform  
2712 yellowish body color (albeit with lighter-colored legs).

2713

2714 **Distribution (Supplementary material Fig. S23).** *Brachymyrmex giardi* is known to occur in  
2715 Chile and Uruguay.

2716

2717 **Biology.** The biology of *B. giardi* has not recently been studied, however Emery (1895) indicated  
2718 that Prof. Giard observed an association between *B. giardi* and the coccid *Margadores vitium*. As  
2719 to the biology of *B. melensis*, which we synonymize here with *B. giardi*, De Zolessi et al. (1978:  
2720 39) provided detailed habitat information including specifications on the landscape, soil and  
2721 vegetation. Their nest was subterranean, with several chambers between 15 and 50 cm deep, each  
2722 chamber being about 3 cm high and 3 to 5 cm long and wide. Some repletes were found hanging  
2723 from the roof together with normal workers as is observed in *Mymecocystus hortideorum*.

2724

2725 **Remarks.** The second ant from the top in pin MSNG: USNMENT00757220 is designated here as  
2726 lectotype, whereas the other specimens are paralectotypes. In the original description of *B. giardi*  
2727 Emery (1895: 215) described a worker, a replete and a queen, and the replete is what we consider  
2728 here as a putative worker-queen intercaste, because a regular queen was also reported by Emery

2729 (1895). Note that this queen was indicated to be wingless, however, after studying the material  
2730 we confirm that it represents a real queen rather than an ergatoid, and the replete has, as  
2731 mentioned above, a hybrid morphology between queen and worker. Upon dissection of the  
2732 abdomen of the replete Emery (1895: 215) reported that the crop is full of honey-like liquid, but  
2733 also that the ovaries are more developed than in normal workers, and that these repletes likely  
2734 have a reproductive function. Nevertheless, he considered nourishment their primary function, as  
2735 is confirmed by De Zolessi et al. (1978). In summary, the exact affinity of these repletes is  
2736 uncertain: if it were ergatoid queens we would not expect a regular queen to be present (Peeters  
2737 1991), which points to an intercaste, because intercastes co-exist with a regular queen. However,  
2738 intercastes do not usually participate in reproduction (Peeters 1991). Given all the available data,  
2739 we consider these specimens for now to be a putative worker-queen intercaste, as mentioned  
2740 before, but the intriguing issue of the repletes in *B. giardi* requires further study.

2741 We have not been able to locate the type material of *B. melensis* and have therefore studied it  
2742 from the detailed work of De Zolessi et al. (1978). These authors subdivided the putative worker-  
2743 queen intercaste into two categories: the first for specimens that resemble normal workers but  
2744 have the gaster somewhat expanded, and the second for the putative intercaste, which displays a  
2745 strongly enlarged gaster with the ability to store liquids. De Zolessi et al. (1978) indicate that  
2746 *B. melensis* resembles *B. physogaster* Kusnezov (1960) most, a species here synonymized to *B.*  
2747 *heeri*, but that both differ in size and in the number of ocelli (see additional differences in the  
2748 diagnosis above). These authors did not compare *B. melensis* and *B. giardi*, but upon doing so we  
2749 did not find any trait that allows distinguishing these taxa and hence we synonymize *B. melensis*  
2750 here. *Brachymyrmex giardi* and *B. heeri* are indeed quite similar, not in the least by the presence  
2751 of a putative worker-queen intercaste, and further in-depth study of both species is required.

2752 *Brachymyrmex giardi* var. *nitida* was previously suggested to be a junior synonym of *B. giardi*  
2753 (Snelling and Hunt 1975), but in our opinion, it is a junior synonym of *B. bruchi* (see above).  
2754 *Brachymyrmex giardi* var. *cordobensis* on the other hand appears to be a junior synonym of *B.*  
2755 *heeri* (see below).

2756  
2757

2758 ***Brachymyrmex heeri* Forel**

2759 **(Figs. 35, 36, supplementary material Fig. S24)**

2760 *Brachymyrmex heeri* Forel, 1874: 91, Figs. 16, 20 (w.). **Lectotype worker** (MHNG:  
2761 USNMENT00757169) and **paralectotype workers, males, queen** (MHNG:  
2762 USNMENT00757167-00757171, USNMENT00758116-00758120); **here designated**): 15  
2763 workers, 3 males, 1 queen [examined]. **SWITZERLAND: Zurich:** Serra des orchidiées. Forel  
2764 (1876: 52) (q.m.). See also: Santschi (1923a: 664).  
2765 = *Brachymyrmex goeldii* Forel, 1912a: 65 (w.). (MHNG: USNMENT00757166): 1 worker  
2766 [examined]. **BRAZIL: São Paulo:** Botucatu. **n. syn.**  
2767 = *Brachymyrmex giardi* var. *cordobensis* Santschi, 1929: 309 (w.). (NHMB:  
2768 USNMENT00757698, 00757699, CASENT0911600): 23 workers [examined]. **ARGENTINA:**  
2769 **Cordoba:** Alta Gracia. **n. syn.**  
2770 = *Brachymyrmex physogaster* Kusnezov, 1960: 382, Figs.1-4 (w.). (INSUE): 7 workers  
2771 [examined]. **ARGENTINA: Salta:** National park Estancia El Rey. **n. syn.**

2772

2773 **Additional material examined. ARGENTINA: Misiones:** Loreto, C. Bruch, 1 worker (NHMB:  
2774 USNMENT00758095). **BOLIVIA: Santa Cruz:** 10 km NW Terevinto, -17.67 -63.45, 380 m, 09  
2775 Dec. 1993, P.S. Ward #12314-61, 2 workers (MCZC: USNMENT00757940); Buena Vista, -

2776 17.45 -63.67, 350 m, 18 Dec. 1993, P.S. Ward #12438-79, 3 workers (PSWC:  
2777 USNMENT00757745); Las Gamas, Parque Nacional Noel Kempff Mercado, -14.80 -60.38, 700  
2778 m, 04 Dec. 1993, 6 workers (PSWC: USNMENT00757941, 00758024). **BRAZIL: Goiás:**  
2779 Campo Limpo, faz conceição, -16.33083 -49.16367, 01-07 July 2005, R.R. Silva & R.M. Feitosa,  
2780 8 workers (ICN: MZSP120, 121); **Minas Gerais:** Serra Caraça, 1380 m, Oct. 1961, Martins &  
2781 Silva, 2 workers, 3 putative worker-queen intercastes (MZSP: USNMENT00757603); Serra  
2782 Caraca, Kloss, Lenko, Nov. 1961, Martins & Silva, 3 workers, 1 putative worker-queen intercaste  
2783 (MCZC: USNMENT00757598); **Pará:** Melgaço, Caixiuanã, ECFPn, -1.77803 -51.42694, 27  
2784 Nov.- 03 Dec. 2001, 2 workers (MPEG: USNMENT00757592, 00757550); Melgaço, Caixiuanã,  
2785 ECFPn, -1.70661 -51.45909, 25-27 Oct. 2005, Equipe A.Y. Harada, 11 workers (MPEG:  
2786 AYH057); Melgaço, Caixiuanã, ECFPn, -1.75444 -51.52241, 24-26 Jan. 2006, Equipe A.Y.  
2787 Harada, 5 workers (MPEG: AYH023); Melgaço, Caixiuanã, ECFPn, -1.75444 -51.52241, 28 Oct.  
2788 2003, A.Y. Harada, E.P. Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P.  
2789 Souza, C. Renato, 8 workers (MPEG: AYH083); **Santa Catarina:** São Bento do Sul, APA Rio  
2790 Vermelho, -26.36417 -49.27111, 30 Mar.-04 Apr. 2001, R.R. Silva & Everhardt, 2 workers (ICN:  
2791 MZSP043); **São Paulo:** Barueri, K. Lenko, 4 workers, 1 queen (MZSP: USNMENT00757602);  
2792 Caraguatatuba, Reserva Florestal, 13 July 1965, Exp. Dep. Zool. 3487, 4 workers, 1 queen  
2793 (MZSP: USNMENT00757597); Cunha, PE Serra do Mar, Nucleo Cunha-Indara, -23.25083 -  
2794 45.00722, 21-22 Apr. 2001, A.A. Tavares & R.R. Silva, 15 workers (ICN: MZSP151).  
2795 **COLOMBIA: Caldas:** Aranzazu, Vereda Cuatro Esquinas, Finca Tres Esquinas, 5.31870 -  
2796 75.48947, 1837 m, 06-08 Aug. 2003, L.E. Franco & J. Cruz, 2 workers (IAvH: IAvH27322);  
2797 Aranzazu, Vereda La Guaira, Finca Alto Bonito, 5.27883 -75.48461, 2056 m, 25-26 July 2003,  
2798 L.E. Franco & J. Cruz, 1 worker (IAvH: IAvH27303); Aranzazu, Vereda Sabana Larga, Finca  
2799 Las Colinas de Zega, 5.31713 -75.47556, 2000 m, 25-27 July 2003, L.E. Franco & J. Cruz, 5

2800 workers, 1 queen (IAvH: IAvH25458, 25459); Salamina, Vereda El Cedrito, Finca El Cedrito,  
2801 5.33117 -75.46744, 1960 m, 27-29 Aug. 2002, L.E. Franco & J. Cruz, 1 worker (IAvH:  
2802 IAvH25465); **Huila**: 8 km S. Neiva, 20 Mar. 1976, W. & E. MacKay, 4 workers (WEMC:  
2803 USNMENT00757948, 00757949); Neiva: 23 Mar. 1976, W. & E. MacKay, 4 workers, 1 queen, 1  
2804 male (WEMC: USNMENT00757733, 00757946, 00757947); **Nariño**: Altaquer, Barro Ñambi, C.  
2805 Sanda #22, 2 workers (ICN: USNMENT00757633); **Quindio**: Calarcá, Vereda Pradera Baja,  
2806 Finca La Holanda, 4.55694 -75.63917, 1575 m, 29 Nov. 1999, E. Gonzalez & J. Sossa, 1 worker  
2807 (IAvH: IAvH-E74153); Calarcá, Vereda Santo Domingo, Finca Santa Librada, 4.55694 -  
2808 75.63917, 1575 m, 16 Mar. 2000, J. Sossa, 1 worker (IAvH: IAvH-E74154); Filandia, Vereda  
2809 Cruces, Finca Pavas, 4.70422 -76.63250, 1900 m, 04-06 June 2002, E. Jimenez & M.F. Reina, 2  
2810 workers (IAvH: IAvH27228); Génova, Vereda El Cedral, Finca Buenos Aires, 4.235 -75.77556,  
2811 1600 m, 26 Oct. 1999, E. Gonzalez & J. Sossa, 2 workers (IAvH: IAvH-E74166, 74167);  
2812 **Risaralda**: Apia, La María, Cafetal de sol (S-III), 3.13 -75.95, 1405 m, 28 Jan. 2002, L. Rivera, 1  
2813 worker (IAvH: IAvH-E74175); Pereira Vereda La Suiza, SFF Otún Quimbaya, 4.72800 -  
2814 75.57744, 1900 m, 24-26 Nov. 2002, M. Reina & L.E. Franco, 2 workers, 1 putative worker-queen  
2815 intercaste (IAvH: IAvH27279); Pereira, Vereda La Suiza, Finca Pez Fresco, 1890 m, 22-24 Nov.  
2816 2002, E. Jimenez & M.F. Reina, 1 worker (IAvH: IAvH27285); **Valle del Cauca**: Cairo, Vereda  
2817 Llano Grande, Finca Encanto, 4.73603 -76.21698, 1650 m, 03 Apr. 2003, J. Henao, 1 worker  
2818 (IAvH: IAvH25147); Medio Calima Campamento DR., C.H.M. Aldana, 1 worker (ICN:  
2819 USNMENT00757551). **COSTA RICA: Guanacaste**: Cerro Cacao, 10.92682 -85.46823, 1100  
2820 m, 09 Feb. 1989, J. Longino #2342, 1 worker (INBIOCRI001280503); **Heredia**: 16 km N Vol.  
2821 Barba, 10.283 -84.083, 950 m, 12 July 1986, J. Longino #1367, 1 worker, 1 queen (JTLC:  
2822 JTLC000005274); 16 km SSE La Virgen, 10.26871 -84.08572, 1100 m, 09 Mar. 2001, 1 worker  
2823 (JTLC: INBIO0003205132); coffee farms vic. Heredia, 1100 m, 01 July 1991, I. Perfecto, 1



2824 worker (JTLC: LACM ENT 139924); **Puntarenas:** Monteverde, 10.3 -84.8, 1500 m, 17 May  
2825 2001, S. Yanoviak & J. Gering, 1 worker, 1 queen (JTLC: JTLC000002089, JTLC000002253);  
2826 Monteverde, 10.301 -84.806, 1500 m, 14 July 1984, J. Longino 1556, 1 worker, 1 queen (JTLC:  
2827 JTLC000005268); 3 km SE Monteverde, 10.283 -84.783, 1200 m, 02 Mar. 1994, J. Longino  
2828 #3578-s, 1 worker (JTLC: INBIO CRI001282749); 4 km S San Vito, 8.783 -82.967, 1200 m, 30  
2829 June 1995, J. longino #3702-s, 1 worker (JTLC: INBIOCRI001280752); Sirena, Parque Nacional  
2830 Corcovado, 8.467 -83.583, 0-100 m, 23 Sep. 1982, J. Longino, 1 worker (JTLC:  
2831 JTLC000005266); **ECUADOR:** 3.2-13 km N of Puyo, Napo, Pastaza, 953 m, 09 Feb. 1955, R.I.  
2832 Schlinger & E.S. Ross, ANTC10200, 1 worker (CASENT: CASENT0196022); **Pichincha:**  
2833 Maquipucuna, 5km ESE Nanegal, 0.116 -78.633, 1500 m, 17 Aug. 1991, P.S. Ward #11503-19, 2  
2834 workers, 1 queen (PSWC: USNMENT00757596); **Zamora-Chinchipe:** Copalinga, -4.09122 -  
2835 78.96069, 1000 m, 01-03 Oct. 2009, Delsinne & Arias. **FRENCH GUIANA:** Basse Vie (Petit  
2836 Saut), Aug. 1999, S. Dorou, 3 workers, 2 males (CPDC: USNMENT00757952).  
2837 **GUATEMALA: El Progreso,** 20 km N Estancia de la Virgen, 1800-1900 m, 08 June 1991, R.S.  
2838 Anderson, 1 worker (JTLC: CASENT0601427); Sololá, 1 km N San Andrés, Semetabaj, 14.75 -  
2839 91.13, 1840 m, 16 Nov. 2003, A.L. Wild #AW2059, A.L. Wild #AW 2059, 3 workers (ALWC:  
2840 USNMENT00757942). **MEXICO: Chiapas :** 15.1 km N.W. Bochil, 17.09120 -92.99138, 1930  
2841 m, 24 Sep. 1992, R.S. Anderson, 1 worker (JTLC: CASENT0603200); 29 km E La Trinitaria,  
2842 16.106 -91.772, 1520 m, 21 July 2007, J. Longino #6100, 1 worker (JTLC: JTLC000010323); 3.8  
2843 km ESE Custepec, 15.71205 -92.93387, 1900 m, 18 July 2007, J. Longino #6072-2, 2 workers, 1  
2844 queen, 1 male (JTLC: JTLC000010342, 000010343); **Jalisco:** 6.76 km SW Mazamitla, 19.89222  
2845 -103.07722, 1997 m, 22 June 2000, W. & E. MacKay, 1 worker, 1 male (WEMC:  
2846 USNMENT00757739); **Nuevo León:** nr Monterrey, Mesa de Chipinque, 1365 m, 16-18 July  
2847 1965, 2 workers, 1 queen (MCZC: USNMENT00757666); **Veracruz:** 2.7 km N Teocelo, 1128

2848 m, 22-24 July 1973, A. Newton, 2 workers (WEMC: USNMENT00757945, 00757607); Km 38  
2849 on Fortin-Huatusco road, Cornell University, 1965, 2 workers, 1 queen (MCZC:  
2850 USNMENT00757954); Las Hamacas, 17 km N Santiago Tuxtla, 26-28 Aug. 1853, E.O. Wilson  
2851 #357, 2 workers (MCZC: USNMENT00757951); Los Tuxtlas, 10 km NNW Sontecomapan,  
2852 18.58333 -95.08333, 500 m, 21 Mar. 1985, P.S. Ward #7364; 5.5 km NE Coscomatepec, 05 June  
2853 1988, W. MacKay #10844, 2 workers, 1 male (PSWC: USNMENT00757950); 5.5 km NE  
2854 Coscomatepec, 05 June 1988, W. MacKay #10844, 2 workers, 1 male (WEMC:  
2855 USNMENT00757944). **NICARAGUA: Granada:** Mombacho Volcano, 11.93394 -85.97858,  
2856 1150 m, 18 July 2003, W. & E. Mackay, 2 workers, 1 queen (WEMC: USNMENT00758042).  
2857 **PANAMA: Canal zone:** 3 km NW Gamboa, 9.13333 -79.71667, 40 m, 10 Dec. 1983, P.S. Ward  
2858 #6391-14, 3 workers (PSWC: USNMENT00757542); **Chiriqui:** Parque Nacional Volcan Baru  
2859 Boquete, 1850 m, 18 June 1995, R. Anderson #17810, 1 worker (WEMC:  
2860 USNMENT00757943). **PARAGUAY: Canindeyú,** Reserva Natural Bosque Mbaracayú,  
2861 Jejuimini, -24.1 -55.5, 24 July 1996, A. Wild #AW0235, 2 workers, 1 queen (ALWC:  
2862 USNMENT00758025); Reserva Natural Bosque Mbaracayú, Jejuimini, -24.1 -55.5, 11 Mar.  
2863 1997, A. Wild #AW0477, #AW0478, 3 workers, 1 queen (ALWC: USNMENT00757543,  
2864 00757955). **VENEZUELA: Aragua:** Parque Nacional Henri Pittier, La Toma, 10.59233 -  
2865 68.14031, 1169 m, 09-19 Aug. 2008, 1 worker (ICN: USNMENT00757740); **Lara:** 9 km SE  
2866 Barbacoas, 9.77 -70.06, 2000 m, 22 Aug. 1987, P.S. Ward #8922, 2 workers, 1 queen (PSWC:  
2867 USNMENT00757953).  
2868  
2869 **Diagnosis.** *Brachymyrmex heeri* resembles *B. depilis* and *B. giardi* closely, because all three  
2870 species have a mesonotum that bulges above the pronotum in lateral view, and a gaster with

2871 dense pubescence. However, *B. heeri* differs from *B. depilis* and *B. giardi* by having scapes that  
2872 surpass the posterior margin of the head.

2873  
2874 *Lectotype and paralectotypes measurements* (mm) (n=5). HL<sub>1</sub> 0.39-0.43; HL<sub>2</sub> 0.27-0.30; HL<sub>3</sub>  
2875 0.10; HW 0.35-0.41; SL 0.37-0.39; EL 0.10-0.11; WL 0.39-0.45; PnL 0.10; PnW 0.25-0.31; ML  
2876 0.08-0.12; MW 0.16-0.20; *Indices* CI 90.00-100.00; SI<sub>1</sub> 90.48-111.11; SI<sub>2</sub> 126.67-142.86; OI<sub>1</sub>  
2877 23.81-27.78; OI<sub>2</sub> 22.72-25.00.

2878  
2879 *Additional material examined measurements* (mm) (n=5). HL<sub>1</sub> 0.41-0.60; HL<sub>2</sub> 0.27-0.39; HL<sub>3</sub>  
2880 0.10-0.13; HW 0.40-0.66; SL 0.36-0.52; EL 0.09-0.19; WL 0.35-0.68; PnL 0.11-0.16; PnW 0.27-  
2881 0.45; ML 0.09-0.21; MW 0.18-0.35; *Indices* CI 93.75-110.81; SI<sub>1</sub> 78.05-93.62; SI<sub>2</sub> 114.29-  
2882 135.71; OI<sub>1</sub> 21.74-29.27; OI<sub>2</sub> 16.22-28.00.

2883  
2884 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
2885 concave. Clypeus with a rounded anterior margin and five long, erect hairs of which a single,  
2886 usually conspicuous hair is near the anterior margin, two hairs are in mediolateral position and  
2887 two more near the toruli; other hairs on the clypeus are markedly shorter and appressed or  
2888 decumbent. Toruli surpassing the posterior clypeal margin in oblique anterodorsal view. The  
2889 scapes surpass the posterior margin of the head by a length smaller than the maximal diameter of  
2890 the eye; they bear decumbent hairs. Ocelli absent. Eyes are positioned on the cephalic midline  
2891 and have 6-7 ommatidia along their maximal diameter.

2892 **Mesosoma.** With several decumbent hairs and usually two erect hairs on the pronotum and two  
2893 on the mesonotum, but sometimes those on the mesonotum or on both are absent. The  
2894 mesonotum is inflated and bulges dorsally above the pronotum in lateral view. Metanotal groove

2895 absent or narrower than the diameter of the metanotal spiracles. Metathoracic spiracles in  
2896 dorsolateral position, not protruding, and touching the propodeal suture. Dorsum of the  
2897 propodeum convex and shorter than the propodeal slope. Propodeal spiracles circular, positioned  
2898 on the posterior propodeal margin, slightly posterior of the middle of the propodeal slope. Legs  
2899 with appressed hairs. Petiole short and inclined forward.

2900 **Gaster.** With dense pubescence and scattered long erect hairs at the edges of the segments. Some  
2901 specimens of *B. heeri* resemble the regular worker in head and mesosoma, but they have a  
2902 strongly expanded gaster (physogastry).

2903 **Color and sculpture.** Head and gaster smooth, dorsum of the mesosoma with imbricate sculpture,  
2904 body opaque and yellowish, sometimes with a somewhat darker gaster.

2905  
2906 **Distribution (Supplementary material Fig. S24).** We have studied *B. heeri* from localities in  
2907 Argentina, Bolivia, Brazil, Colombia, Costa Rica, Ecuador, French Guiana, Guatemala, Mexico,  
2908 Nicaragua, Panama, Paraguay, Uruguay, Venezuela and from populations that were introduced in  
2909 Switzerland. Santschi (1923a) also reported this species from Puerto Rico, Jamaica, Haiti and  
2910 Guyana, but we have not studied this material.

2911  
2912 **Biology.** *Brachymyrmex heeri* was originally described from specimens collected in a tropical  
2913 orchid greenhouse in Switzerland (Forel 1874). The species was found climbing on various  
2914 plants, and was associated with aphids, but the nest was not located. Santschi (1929) reported that  
2915 the nest of the type specimens of *B. giardi* var. *cordobensis* was found at the base of a tree, and *B*  
2916 *physogaster* was collected in the heavily humic, finely loamy soil in the shadow of trees, where  
2917 both excessive moisture and eventual drought are unlikely (Kusnezov 1960).

2918

2919 **Remarks.** The lectotype is designated here as the ant in the middle of their holder MHNG:  
2920 USNMENT00757169, whereas the other specimens are paralectotypes. Forel (1874) described  
2921 the worker of *B. heeri*, but did not indicate physogastry in his description, and Santschi (1923a)  
2922 did not comment on this issue neither. In the description of *B. giardi* var. *cordobensis* (which is  
2923 here synonymized under *B. heeri*) Santschi (1929) referred to physogastric specimens, and also to  
2924 ‘worker-queens’ with a strongly expanded gaster. He does not provide any additional description  
2925 of these worker-queen specimens, but upon examination of this material and other samples of *B.*  
2926 *heeri* we exclusively found worker-like specimens with a normal or robust mesosoma and an  
2927 expanded gaster. These specimens do not have morphological features reminiscent of a queen,  
2928 but rather differ from regular workers by having a larger body size, a subquadrate head, with the  
2929 posterior cephalic margin slightly concave to almost flat, scapes that barely reach the posterior  
2930 margin of the head, three ocelli, eyes that have approximately 11 ommatidia along their maximal  
2931 diameter, a deep metanotal groove that is wider than the diameter of the metathoracic spiracles,  
2932 and as already mentioned, a strongly expanded gaster. To exclude the possibility that the museum  
2933 samples we studied reflect a mixture of two species, we sequenced specimens of both morphs,  
2934 and found them to be genetically very similar. In summary, *B. heeri* either has dimorphic  
2935 workers, or a putative worker-queen intercaste, and whereas we consider the first scenario more  
2936 likely (because we did not find specimens with a hybrid worker-queen morphology), the issue  
2937 requires more study. These observations also apply to *B. physogaster*, and this species and *B.*  
2938 *giardi* var. *cordobensis* are synonymized here with *B. heeri* because they contain all its  
2939 morphological features. Forel (1912a) distinguished *B. goeldii* from other *Brachymyrmex* species  
2940 principally based on the form of the head, and he mainly compared it with *B. bruchi*, however,  
2941 this material appears to belong to a ‘robust worker’ of *B. heeri*. Interestingly, as for *B. heeri*  
2942 before, Forel (1912a) does not mention physogastry.

2943 Two varieties have been assigned to *B. heeri*, namely *B. heeri* var. *fallax*, and *B. heeri* var.  
2944 *basalis*. As indicated before, *B. heeri* var. *fallax* is attributed here to *B. aphidicola*, and *B. heeri*  
2945 var. *basalis* is synonymized under *B. pictus* (see below) here.

2946 It is noteworthy that *B. heeri* has a wide geographic distribution and a study of the morphological  
2947 variation within the species in a molecular and geographic context would be required. Likewise,  
2948 it would be interesting to study the biology of the various morphs within a colony, including the  
2949 underlying developmental processes and their distribution and behavior within the colony.

2950 Indeed, some colleagues (Kusnezov 1960) have labeled physogastric specimens here assigned to  
2951 *B. heeri* as honey pot workers and have observed them hanging from the ceiling of the chambers  
2952 of the nest, as in *Myrmecocystus*, and *B. melensis* (see *B. giardi* above). However, why  
2953 physogastry is present in some *Brachymyrmex* species, but not in others remains unclear.

2954 Kusnezov (1960) suggested that it may relate to a trophobiotic lifestyle, whereas others suggested  
2955 it is an adaptation to periodically arid conditions or food scarcity (Forel 1902; Wheeler 1910).  
2956 However, Creighton (1950), argued exactly the opposite by suggesting that physogastry may  
2957 develop when a xerophyte species encounters and adapts to less arid and more resource-rich  
2958 habitats. Clearly, more work on this issue is required, but as mentioned above, the habitat from  
2959 which *B. physogaster* was reported by Kusnezov (1960) does not appear to confirm the ‘scarcity  
2960 hypothesis’. This future work should perhaps also consider potential intrinsic causes of  
2961 physogastry beyond the putative environmental drivers discussed above.

2962

2963

2964

***Brachymyrmex iridescens* NEW SPECIES**

2965

**(Fig. 37, supplementary material Fig. S25)**

2966 **Holotype worker** (MZSP: USNMENT00757758) and **paratype workers** (MZSP:  
2967 USNMENT00757757): 5 workers. **BRAZIL: Santa Catharina**, Chapecó, July 1960, F.  
2968 Plaumann 9878.  
2969  
2970 **Additional material examined. ARGENTINA: Misiones:** Parque Nacional Iguazú, -25.71847 -  
2971 54.43319, 23 Sep. 1999, Leponce, Roisin & Theunis, 1 worker (MACN: MACN\_En 18283,  
2972 MACN\_En 18201, MACN\_En 18242); Parque Nacional Iguazú, 1 worker (RBINS: Coll.  
2973 RISCNB SID SPM\_ID01612). **BRAZIL: Parana:** Londrina, Nov. 2004, D.T. Lopes, 1 worker  
2974 (CPDC: USNMENT00757755); **Santa Catharina:** Teutônia, June 1961, F. Plaumann, 8219, 3  
2975 workers (MZSP: USNMENT00757756).  
2976  
2977 **Etymology:** The epithet *iridescens* refers to the conspicuous iridescent appearance of this species  
2978 under stereoscope illumination.  
2979  
2980 **Diagnosis.** *Brachymyrmex iridescens* morphologically resembles *B. santschii*, because both have  
2981 conspicuous, strongly alveolate sculpture on the head and mesosoma. However, they differ from  
2982 one another because *B. iridescens* has a metanotal groove that is narrower than the diameter of  
2983 the metathoracic spiracles, scapes that approximately reach the posterior margin of the head, and  
2984 a gaster with dense pubescence.  
2985  
2986 *Holotype and paratype measurements* (mm) (n=2). HL<sub>1</sub> 0.40-0.43; HL<sub>2</sub> 0.29 - 0.30; HL<sub>3</sub> 0.09;  
2987 HW 0.35 - 0.37; SL 0.30-0.34; EL 0.09-0.10; WL 0.39-0.40; PnL 0.13; PnW 0.22-0.25; ML 0.09  
2988 - 0.11; MW 0.13-0.16; *Indices* CI 85.42-86.67; SI<sub>1</sub>80.95-82.5; SI<sub>2</sub> 100.00; OI<sub>1</sub> 23.81-25.00; OI<sub>2</sub>  
2989 23.81-25.00.

2990  
2991 *Additional material examined measurements* (mm) (n=2). HL<sub>1</sub> 0.37-0.42; HL<sub>2</sub> 0.26-0.30; HL<sub>3</sub>  
2992 0.07-0.11; HW 0.29-0.38; SL 0.28-0.30; EL 0.09; WL 0.39-0.42; PnL 0.12 - 0.13; PnW 0.22-  
2993 0.22; ML 0.09-0.11; MW 0.13-0.18; *Indices* CI 78.57-89.36; SI<sub>1</sub> 80.95-93.94; SI<sub>2</sub> 100.00-106.90;  
2994 OI<sub>1</sub> 23.81-30.30; OI<sub>2</sub> 19.05-25.53.

2995  
2996 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
2997 concave or flat. Dorsum of the head with decumbent hairs. Clypeus with a rounded anterior  
2998 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
2999 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
3000 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
3001 margin in oblique anterodorsal view. The scapes are short and approximately reach the posterior  
3002 margin of the head; they have decumbent and suberect hairs. Three ocelli are present. Eyes are  
3003 positioned on the cephalic midline and have 8-9 ommatidia along their maximal diameter.

3004 **Mesosoma.** With several decumbent hairs, and on the pronotum some suberect hairs. The  
3005 mesonotum is inflated and bulges dorsally above the pronotum. The metanotal groove is  
3006 narrower than the diameter of the metathoracic spiracles. Metathoracic spiracles dorsal, slightly  
3007 protruding, and not touching the mesonotal nor propodeal suture. Dorsum of the propodeum  
3008 slightly convex and shorter than the propodeal slope. Propodeal spiracles circular, positioned on  
3009 the posterior propodeal margin at the middle of the propodeal slope. Legs with appressed hairs.  
3010 Petiole short and inclined forward.

3011 **Gaster.** With decumbent, dense pubescence and several scattered long erect hairs.

3012 **Color and sculpture.** Head and thorax finely alveolate, gaster smooth and shiny. The body is  
3013 uniformly brownish.



3014  
3015 **Distribution (supplementary material Fig. S25).** *Brachymyrmex iridescens* is known from  
3016 Argentina and Brazil.

3017  
3018 **Biology.** Unknown.

3019  
3020 **Remarks.** *Brachymyrmex iridescens* resembles *B. santschii*, but their distributions are strongly  
3021 disjunct: *B. iridescens* is only known from Brazil and Argentina, whereas *B. santschii* from Costa  
3022 Rica and Panama.

3023  
3024

3025 *Brachymyrmex micromegas* Emery

3026 (Fig. 38, supplementary material Fig. S26)

3027 *Brachymyrmex (Brysha) micromegas* Emery in Santschi, 1923a: 675, Figs. 30, 32 (w.).

3028 **Lectotype minor worker** (MCSN: USNMENT00757222) and **paralectotype minor workers,**  
3029 **major workers** (MCSN: USNMENT00757222-00757223; MZSP: USNMENT00758145-  
3030 00758146, 00757825-00757827, 00757830; NHMG: USNMENT00758145-00758146): 4 major  
3031 workers, 5 minor workers [examined]. **BRAZIL: São Paulo:** São Paulo city, Ipiranga. See also:  
3032 Ortiz and Fernández (2014: 16, Figs. 1-6).

3033  
3034 *Additional material examined.* **BRAZIL: São Paulo:** Agudos, 05 Nov. 1967, W. Kempf, 1 minor  
3035 worker (MZSP: USNMENT00757830); Anhembi, Faz B. Rico, 14 Feb. 1969, W. Kempf, J.C.  
3036 Magalhães, L.T.F.M. Kulman, 1 minor worker (MZSP: USNMENT00757834).

3037

3038 **Diagnosis.** *Brachymyrmex micromegas* morphologically resembles *B. pilipes* because both  
3039 species have a dimorphic worker caste, a clypeus with a row of long thick hairs near the anterior  
3040 margin, toruli that touch the posterior clypeal margin, but never surpass it in oblique anterodorsal  
3041 view, and tumuliform metathoracic spiracles. However, *B. micromegas* differs from *B. pilipes* by  
3042 having a smooth and shiny body, with very fine longitudinal striations restricted to the  
3043 metapleura.

3044  
3045 **Description.** See Ortiz and Fernández (2014).

3046  
3047 *Brachymyrmex minutus* Forel

3048 (Fig. 39, supplementary material Fig. S27)

3049 *Brachymyrmex minutus* Forel, 1893: 346 (w.q.). **Lectotype worker** (MHNG:  
3050 USNMENT00757150) and **paralectotype workers, queen** (MHNG: USNMENT00757149-  
3051 00757151; USNMENT00758110-00758112; **here designated**): 15 workers, 1 queen [examined].  
3052 **ANTILLES:** Saint Vincent. See also: Santschi (1923a: 667).

3053  
3054 **Additional material examined. BOLIVIA: Santa Cruz:** Aserradero Moira, -14.57 -61.20, 180  
3055 m, 27 Nov. 1993, P.S. Ward #12174-49, 3 workers (PSWC: USNMENT00757587), 3 workers  
3056 (MCZC: USNMENT00757305); 10 km NW Terevinto, -17.67 -63.45, 380 m, 09 Dec. 1993, P.S.  
3057 Ward #12314-62, #12314-63, 2 workers (PSWC: USNMENT00757870), 5 workers (MCZC:  
3058 USNMENT00757303-00757304); 35 km SSE Flor de Oro, -13.833 -60.867, 450 m, 29 Nov.  
3059 1993, P.S. Ward #12199-73, 7 workers (MCZC: USNMENT00757301 - 00757302); Las Gamas,  
3060 Parque Natural Noel Kempff Mercado, -14.80 -60.38, 700 m, 03 Dec. 1993, P.S. Ward #12266-  
3061 43, 4 workers (MCZC: USNMENT00757299-00757300); Las Gamas, Parque Natural Noel

3062 Kempff Mercado P, -14.80 -60.38, 700 m, 04 Dec. 1993, P.S. Ward #12266-45, 6 workers  
3063 (MCZC: USNMENT00757297-00757298). **BRAZIL: Ceará:** Guaramiranga (Pq. Tr.), -4.267 -  
3064 38.933, 900 m, 22 Feb. 2002, Y. Quinet, 1 worker (CPDC: USNMENT00757874); **Goias:**  
3065 Campo Limpo, faz conceição, -16.33083 -49.16367, 20-24 Jan. 2005, R.R. Silva, 9 workers  
3066 (ICN: MZSP071); Campo Limpo, faz conceição, -16.33083 -49.16367, 01-07 July 2005, R.R.  
3067 Silva & R.M. Feitosa, 15 workers (ICN: MZSP120 - MZSP121); **Pará:** Belem, 07 Aug. 1962,  
3068 P.F. Doulington. B.F. 14, 1 worker (MCZC: USNMENT00757256); Melgaço, Caxiuanã ECFPn,  
3069 -1.70661 -51.45909, 10-12 Oct. 2006, Equipe A.Y. Harada, 1 worker (MPEG: AYH018); 10-11  
3070 Oct. 2006, 1 worker (MPEG: AYH019); 10-12 Oct. 2005, 7 workers (MPEG: AYH041); 24-26  
3071 Jan. 2007, 6 workers (MPEG: AYH003, 069, 115, 122, 124); 28-30 Jan. 2006, 5 workers  
3072 (MPEG: AYH001, 007); 23-25 Apr. 2007, 1 worker (MPEG: AYH005); A.Y. Harada, E.P.  
3073 Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 3 workers  
3074 (MPEG: AYH076); -1.72484 -51.42979, 26 Oct. 2006, A.Y. Harada, E.P. Fagundes, C.J.M.  
3075 Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 45 workers 1 Queen (MPEG:  
3076 AYH005, 016, 032, 052, 053, 068, 070, 078, 087, 093, 097, 100, 102, 106, 113, 136); 12-14 Oct.  
3077 2006, Equipe A.Y. Harada, 1 worker (MPEG: AYH045); 20-22 July 2006, 2 workers (MPEG:  
3078 AYH095); 20-22 Oct. 2005, 2 workers (MPEG: AYH082, 105); 21-23 Oct. 2005, 1 workers  
3079 (MPEG: AYH006); 22-24 Apr. 2005, 2 workers (MPEG: AYH082, 105); 23-25 Feb. 2006, 1  
3080 worker (MPEG: AYH033); 24-26 July 2003, 10 workers (MPEG: AYH066, 084, 108, 119, 133,  
3081 135); 26-28 Nov. 2003, A.Y. Harada, E.P. Fagundes, C. Renato, 1 worker (MPEG: AYH117);  
3082 07-09 Feb. 2003, 1 worker (MPEG: AYH060); Melgaço, Caxiuanã ECFPn, -1.73360 -51.51054,  
3083 27 Oct. 2003, Equipe A.Y. Harada, 3 workers (MPEG: AYH014, 061); Melgaço, Caxiuanã  
3084 ECFPn, -1.73360 -51.51053, 27 Oct. 2003, A.Y. Harada, E.P. Fagundes, C.J.M. Ribeiro, C.E.D.  
3085 Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 3 workers (MPEG: AYH101, 035, 050); 14-16

3086 Oct. 2006, Equipe A.Y. Harada, 15 workers, 2 queens (MPEG: AYH107, 081, 116); 23-25 July  
3087 2006, 1 worker (MPEG: AYH104); 12-14 Oct. 2006, Equipe A.Y. Harada, 3 workers (MPEG:  
3088 AYH051); Melgaço, Caxiuanã ECFPn, -1.73584 -51.48762, 30 Oct. 2003, A.Y. Harada, E.P.  
3089 Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 1 worker  
3090 (MPEG: AYH091); 13-15 Oct. 2005, Equipe A.Y. Harada, 5 workers (MPEG: AYH031, 056);  
3091 15-17 Jan. 2006, 1 worker (MPEG: AYH132); 21 Apr. 2006, 1 worker (MPEG: AYH080); 22-24  
3092 July 2006, 1 worker (MPEG: AYH121); 23-25 Oct. 2005, 3 workers (MPEG: AYH017, 027,  
3093 030); 23-28 Oct. 2005, 1 worker (MPEG: AYH022); 25-27 Jan. 2006, 1 worker (MPEG:  
3094 AYH128); 26-28 Jan. 2007, 2 workers (MPEG: AYH075); -1.75444 -51.52241, 28 Jan. 2003,  
3095 AY Harada, E.P. Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C.  
3096 Renato, 1 worker (MPEG: AYH011); 28 Oct. 2003, 10 workers (MPEG: AYH028, 079, 089,  
3097 092, 118); 01-03 Nov. 2004, Equipe A.Y. Harada, 1 worker (MPEG: AYH020); 01-05 Nov.  
3098 2004, 1 worker (MPEG: AYH004); 19-21 Oct. 2007, 1 worker (MPEG: AYH026); 20-22 July  
3099 2007, 3 workers (MPEG: AYH025, 065); 22-24 Oct. 2003, 1 worker (MPEG: AYH085); 22-24  
3100 Oct. 2005, 2 workers (MPEG: AYH110, 123); 24-26 Oct. 2006, 1 worker (MPEG: AYH040);  
3101 Melgaço, Caxiuanã ECFPn, -1.78155 -51.59197, 24-26 Oct. 2007, 1 worker (MPEG: AYH009);  
3102 30 July-01 Aug. 2003, 2 workers (MPEG: AYH010); 02 Oct. 2006, A.Y. Harada, E.P. Fagundes,  
3103 C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato, 2 workers (MPEG:  
3104 AYH077); 30 Oct. 2003, 2 workers (MPEG: AYH125, 130); 02 Nov. 2003, 1 worker (MPEG:  
3105 AYH120); 12-14 Oct. 2006, Equipe A.Y. Harada, 1 worker (MPEG: AYH054); 14-16 Oct. 2006,  
3106 1 worker (MPEG: AYH043); 15-17 Oct. 2006, 1 worker (MPEG: AYH055); 15-17 Oct. 2010, 16  
3107 workers (MPEG: AYH013, 024); 23-25 Apr. 2003, 1 worker (MPEG: AYH096); 23-25 July  
3108 2007, 6 workers (MPEG: AYH002, 042, 072); 23-25 May 2003, 1 worker (MPEG: AYH114);  
3109 24-26 July 2006, 2 workers (MPEG: AYH015); 25-27 Jan. 2007, 2 workers (MPEG: AYH064,

3110 090); 25-27 June 2007, 1 worker (MPEG: AYH012); 30 July-01 Aug. 2003, 3 workers (MPEG:  
3111 AYH103, 131); 30 July-01 Aug. 2003, A.Y. Harada, E.P. Fagundes, C. Renato, 2 workers  
3112 (MPEG: AYH098, 109); **Rondônia:** Ouro Preto do Oeste, 04 Apr. 1985, F.F. Ramos, Res. INPA  
3113 N°0388, 2 workers (MPEG: USNMENT00757865, 00757866); São Paulo, Jundiá, Serra Do  
3114 Japi, Jan. 2009, S. Diniz, 3 workers (ICN: USNMENT00759044); **São Paulo:** Tapirai, -24.03208  
3115 -47.46556, 08-14 Jan. 2001, R.R. Silva & Everhardt, 1 worker, 2 males (ICN: MZSP172).  
3116 **COLOMBIA: Amazonas:** Leticia, 1 worker (ICN); Parque Nacional Natural Amacayacu,  
3117 Matamata, -3.68 -70.25, 150 m, 23 Oct. 2000, 1 worker (ICN); **Cundinamarca:** La Vega,  
3118 Reserva Natural Natautá, 5.00 -74.33, 1040 m, 10 Nov. 2010, F. Fernández, 1 worker (ICN:  
3119 USNMENT00757873); **Magdalena:** 4 km N San Pedro, 10.95 -74.05, 550 m, 14 Aug. 1985, P.S.  
3120 Ward #7912-35, 3 workers (PSWC: USNMENT00757863); **Norte de Santander:** 2 km N  
3121 Barrancabermeja, Dec. 1962, leafmodd Berles. J. Archer; **Quindío:** Buenavista, Vereda El  
3122 infierno, Finca Guadalajara, 4.37667 -75.76944, 1160 m, 16 Nov. 1999, E. Gonzalez, 3 workers  
3123 (IAvH: IAvH-E74164); Calarca, Vereda Santo Domingo, Finca Santa Librada, 4.55694 -  
3124 75.63917, 1575 m, 16 Jan. 2000, J. Sosa, 5 workers, 1 queen (IAvH: IAvH-E74163, 74157);  
3125 Circasia, Vereda Buenavista, Finca Calamar, 4.59806 -75.69861, 1450 m, 12 Oct. 1999, E.  
3126 Gonzalez, 2 workers (IAvH: IAvH-E74168); **Risaralda:** La Celia, 5.00361 -76.00444, 1900 m,  
3127 27 Jan. 2011, Gustavo Zabala, 2 workers (ICN); **Valle del Cauca:** Cairo, Vereda Vallecitos,  
3128 Finca El Maladero, 4.75803 -76.22732, 1850 m, 29 Mar. 2003, J. Henao, 1 worker (IAvH:  
3129 IAvH25152); **Vichada:** Cumaribo, Corregimiento Santa Rita, PNN El Tuparro, 5.3075 -67.9500,  
3130 135 m, 14-16 Feb. 2004, I. Quintero & E. Gonzalez, 1 worker (IAvH: USNMENT00759055).  
3131 **COSTA RICA: Heredia:** Estación Biológica La Selva, 10.43691 -84.01374, 50 m, 19 Mar. 93,  
3132 J. Longino, 1 worker (JTLC: JTLC000007845); La Selva, 12 June 1991, L.E. Tennant, 2 workers  
3133 (MCZC: USNMENT00757284, 00757285); **Puntarenas:** Estación Biológica Los Llanos,

3134 10.30487 -84.83735, 1150 m, 28 Feb. 2004, J. Longino #5249-s, 1 worker (JTLC:  
3135 JTLC000004545); Peninsula Osa, Parque Natural Corcovado, Sirena, 8.467 -83.583, 11 Aug.  
3136 1980, J. Longino, 1 worker (MCZC: USNMENT00757289); Parque Natural Corcovado, Sirena,  
3137 8.48333 -83.60000, 10 m, 27 June 1982, J. Longino, 1 worker (JTLC: JTLC000005929); **San**  
3138 **José**: San Jose, 9.933 -84.083, 1100 m, 14 June 1999, J. Longino #4040-s, 1 worker (JTLC:  
3139 LACM ENT 143521). **CUBA: Holguín**: 2 km N La Melba, 20.467 -74.817, 400 m, 22 Aug.  
3140 2001, P.S. Ward #14424-16, 3 workers (PSWC: USNMENT00757862); **Manicaragua**: Trinidad  
3141 Mts, 01 Aug. 1953, E. O. Wilson #65, 3 workers, 1 queen (MCZC: USNMENT00757295).  
3142 **DOMINICAN REPUBLIC**: 16 km ENE Pedernales, 18.1167 -71.6167, 800 m, 10 Sep. 1992,  
3143 P.S. Ward #11751-16, 3 workers (PSWC: USNMENT00757872). **ECUADOR: Pichincha**:  
3144 Reserva Forestal ENDESA, 0.083 -79.033, 25 Jan. 1994, L.E. Tennant, 3 workers MCZC:  
3145 USNMENT00757293, 00757869); Reserva Forestal ENDESA, 0.13 -79.05, 600 m, 05 Dec.  
3146 2003, A. L. Wild #AW2195, 1 worker (ALWC: USNMENT00757861); **Zamora-Chinchipe**:  
3147 Copalinga, -4.09122 -78.96069, 1000 m, 28-30 Sep. 2009, Delsinne/Arias, col id 35155, 35161,  
3148 15 workers (RBINS: Coll.RIScNB SID SPM\_ID3515505, 3516111); 30 Sep.– 02 Oct. 2009,  
3149 Delsinne/Arias, col id 34651, 39654, 3 workers (RBINS: Coll.RIScNB SID SPM\_ID3465109,  
3150 3465404); 01-03 Oct. 2009, Delsinne/Arias, col id34663, 34671, 34673, 14 workers (RBINS:  
3151 Coll.RIScNB SID SPM\_ID3466304, 3467107, 3467305); 07 Oct. 2009, Delsinne/Arias, col Id  
3152 34715, 34733, 29 workers (RBINS: Coll.RIScNB SID SPM\_ID3471501, 3473311); 04-06 Oct.  
3153 2009, Delsinne/Arias, col id34695, 34703-34705, 34709), 19 workers (RBINS: Coll.RIScNB SID  
3154 SPM\_ID3469506, 3470305, 3470405, 3470504, 3470909); 05-07 Oct. 2009, Delsinne/Arias, col  
3155 id34746, 34751, 34753, 5 workers (RBINS: Coll.RIScNB SID SPM\_ID3474605, 3475106,  
3156 3475305). **FRENCH GUIANA**: Araguez, Inselbery forest, Sara Groc, 3 workers (ICN:  
3157 USNMENT00757864); Kaw Mountains, Oct. 2008, Sara Groc, 2 workers (ICN:

3158 USNMENT00757867); Oct. 2009, Sara Groc, 6 workers (ICN: USNMENT00759035).

3159 **GUATEMALA: Zacapa:** 8.5 km NE Tuculután, 15.058717 -89.67638, 1100 m, 06 July 2001, J.

3160 Longino #6016-s, 1 worker (JTLC: JTLC000009864). **MEXICO: Oaxaca:** 7.2 km S Valle

3161 Nacional, 490 m, 11-18 Aug. 1973, A. Newton, 2 workers (MCZC: USNMENT00757286,

3162 00757287); **Veracruz:** Los Tuxtlas, 10 km NNW Sontecomapan, 18.583 -95.083, 500 m, 21

3163 Mar. 1995, P.S. Ward #1369-39, 2 workers (PSWC: USNMENT00757871); Sa. Teviscocla, nr.

3164 Cuichapa to 1600 m trop. For w. coffe, 04 Aug. 1965, Cornell Univ. Mexico Field Party, 6

3165 workers, 1 queen (MCZC: USNMENT00757290, 00757291, 00757296). **PANAMA:** Barro

3166 Colorado I., Canal Zone, Jan. 1960, m-16 Strays, W.L. Brown, E.S. McCluskey, 1 worker

3167 (MCZC: USNMENT00757288); **PARAGUAY: Canindeyú:** Reserva Natural del Bosque,

3168 Mbaracayú, Jejuimi, -24.1 -55.5, 02 May 1996, A. Wild #0132, 3 workers, 1 queen (ALWC:

3169 USNMENT00757601); 11 Feb. 1997, A. Wild #AW 0409, 1 worker (ALWC:

3170 USNMENT00757860). **PERU: Madre de Dios:** Reserva Nacional Tambopata, Centro

3171 Sachavacayoc, -12.85583 -69.36194, 19-31 July 2012, curso de hormigas, 7 workers (ICN:

3172 JSC120726-LS01). **SURINAME:** Tambahredjo, June 1959, I. V. d. Drift, 1 worker (MPEG:

3173 USNMENT00757876). **VENEZUELA: Aragua:** Parque Nac. Henri Pittier, La Toma, 10.34924

3174 -67.68251, 1169 m, 09-19 Aug. 2008, Ant Course, 3 workers, 1 queen (ICN:

3175 USNMENT00757600, 00758023); Rancho Grande, 1100 m, 23-27 June 1967, W.L. Brown, 1

3176 worker (MCZC: USNMENT00757294); 12 Aug. 1967, R.W. Poole, 3 workers (MCZC:

3177 USNMENT00757875).

3178

3179 **Diagnosis.** *Brachymyrmex minutus* morphologically resembles *B. australis*, *B. aphidicola* and *B.*

3180 *termitophilus*, because all these species have eyes located on the cephalic midline, a mesonotum

3181 that does not bulge dorsally above the pronotum in lateral view, and yellowish body color.

3182 *Brachymyrmex minutus* differs from the three other species, or any other *Brachymyrmex* species  
3183 for that matter, by having a very inconspicuous mesometanotal suture, giving the impression that  
3184 the mesonotum and metanotum are fused. Additionally, *B. minutus* differs from *B. termitophilus*  
3185 by having scattered pubescence on the gaster, from *B. aphidicola* by the presence of only two  
3186 erect hairs on the pronotum and from *B. australis* by having scapes that surpass the posterior  
3187 margin of the head by a length that is smaller than the maximal diameter of the eye.

3188  
3189 *Lectotype and paralectotypes worker measurements* (mm) (n=4). HL<sub>1</sub> 0.37-0.40; HL<sub>2</sub> 0.25-0.31;  
3190 HL<sub>3</sub>0.07-0.10; HW 0.32-0.36; SL 0.32-0.36; EL 0.08-0.11; WL n.a.; PnL n.a.; PnW 0.20-0.24;  
3191 ML 0.06-0.07; MW 0.14-0.15; *Indices* CI 85.88-89.29; SI<sub>1</sub> 92.59-108.00; SI<sub>2</sub> 107.14-126.67; OI<sub>1</sub>  
3192 26.03-29.63; OI<sub>2</sub> 20.00-25.27.

3193  
3194 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin weakly  
3195 convex. Dorsum of the head with sparse hairs. Clypeus with a rounded anterior margin and five  
3196 long, erect hairs of which a single, usually conspicuous hair is near the anterior margin, two hairs  
3197 are in mediolateral position and two more near the toruli; other hairs on the clypeus are markedly  
3198 shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique  
3199 anterodorsal oblique view. The scapes surpass the posterior margin of the head by a length that is  
3200 smaller than the maximal diameter of the eye; they have appressed pubescence. Ocelli absent.  
3201 Eyes are positioned on the cephalic midline and have 8-9 ommatidia along their maximal  
3202 diameter.

3203 **Mesosoma.** Typically with two erect hairs on the pronotum and without erect hairs on the  
3204 mesonotum. The mesonotum is not inflated and does not bulge dorsally above the pronotum in  
3205 lateral view. Mesometanotal suture inconspicuous, giving the impression that the mesonotum and



3206 metanotum are fused. Metanotal groove absent or narrower than the diameter of the metathoracic  
3207 spiracles. Metathoracic spiracles in dorsolateral position, not protruding, and touching the  
3208 propodeal suture. Dorsum of the propodeum slightly convex and shorter than the propodeal slope.  
3209 Propodeal spiracles circular, positioned on the posterior propodeal margin, just dorsally at the  
3210 middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole short and inclined  
3211 forward.

3212 **Gaster.** With scarce pubescence and several scattered long suberect hairs.

3213 **Color and sculpture.** Body smooth, or finely granulated and shiny, sometimes with some  
3214 imbricate sculpture on the dorsum of the mesosoma. Body usually yellowish, but sometimes  
3215 reddish.

3216  
3217 **Distribution (Supplementary material Fig. S27).** *Brachymyrmex minutus* is known to occur in  
3218 Bolivia, Brazil, Colombia, Costa Rica, Cuba, the Dominican Republic, Ecuador, French Guiana,  
3219 Guatemala, Mexico, Panama, Paraguay, Peru, Suriname and Venezuela.

3220  
3221 **Biology.** Forel (1893) found several colonies subterraneous at the roots of sod, usually where it  
3222 overgrows rocks, typically within the forest and/or nearby streams. The nest of *B. minutus*  
3223 consists of large tunnels with small chambers at intervals. Colonies contain ~50 to 100  
3224 individuals, with a single queen.

3225  
3226 **Remarks.** The lectotype is designated here as the second ant counting from the top of pin  
3227 MHNG: USNMENT00757150, whereas the other specimens on that pin are paralectotypes.

3228 Forel (1893) admitted having confused this species for *B. heeri* before but provided several  
3229 morphological differences. However, *B. minutus* is morphologically more similar to the species  
3230 indicated here in the diagnosis.

3231

3232 *Brachymyrmex modestus* Santschi

3233 (Fig. 40, supplementary material Fig. S28)

3234 *Brachymyrmex modestus* Santschi, 1923b: 271 (w.). **Lectotype worker** (NHMB:  
3235 USNMENT00758099) and **paralectotype workers** (NHMB: USNMENT00758099, 00758100;  
3236 **here designated**): 3 workers [examined]. **BRAZIL: Santa Catarina:** Blumerau, A.  
3237 Reinchensperguer leg.

3238

3239 **Additional material examined. BRAZIL: Santa Catarina:** Palhoça, PE Serra do Tabuleiro, -  
3240 27.74111 -48.69722, 02-10 June 2003, R.R. Silva, B.H. Dietz & A. Tavares, 1 worker (ICN:  
3241 MZSP033); São Bento do Sul, APA Rio Vermelho, -26.36417 -49.27111, 30 Mar.-04 Apr. 2004,  
3242 R. Silva & Eberhardt, 8 workers (ICN: MZSP138); **São Paulo:** Cunha, PE Serra do Mar, Nucleo  
3243 Cunha-Indara, -23.25083 -45.00722, 21-22 Apr. 2001, A.A. Tavares & R.R. Silva, 12 workers  
3244 (ICN: MZSP149); Praia grande, PE Serra do Mar, nucleo pilhões-Cubatão, -23.9753 -46.5400,  
3245 26-27 May 2001, A.A. Tavares & R.R. Silva, 5 workers (ICN: MZSP165). **COLOMBIA:**  
3246 **Amazonas:** Leticia, Reserva Forestal del Rio Calderón, Estación Biológica El Zafre, -4.00583, -  
3247 69.89861, 150 m, 02-04 Dec. 2007, L.E. Franco & S. Florez, 2 workers (IAvH); **Caldas:**  
3248 Aranzazu, Vereda Chambery, Finca Los Planes, 5.29231 -75.47283, 1910 m, 01-03 Aug. 2003,  
3249 L.E. Franco & Cruz, 1 worker (IAvH: IAvH27307); Finca maranduba, 5.30731 -75.47250, 2050  
3250 m, 30 July–01 Aug. 2003), L.E. Franco & Cruz, 2 workers (IAvH: IAvH27288); Vereda La  
3251 Guaira, Finca Villa Ofelia, 5.286 -75.465, 1965 m, 01-03 Aug. 2003, L.E. Franco & Cruz, 1

3252 worker, 1 queen (IAvH: IAvH27320); Vereda Sabana Larga, Finca Cañada Brujas, 5.30883 -  
3253 75.47281, 1920 m, 31 July-02 Aug. 2003, L.E. Franco & Cruz, 3 workers (IAvH: IAvH55442);  
3254 **Quindío:** Circasia, Finca Calamar, 4.60 -75.70, 1450 m, 12 Oct. 1999, E. Gonzalez, 1 worker  
3255 (IAvH: IAvH 74169); Filandia, Vereda Cruces, Finca Aranjuez, 4.70826 -75.64679, 1750 m, 13-  
3256 15 July 2002, E. Jiménez & L.E. Franco, 2 workers (IAvH: IAvH27261); Finca La Cha, 4.70468  
3257 -75.60982, 1920 m, 28-30 Aug. 2002, E. Jiménez & L.E. Franco, 1 worker, 1 queen (IAvH:  
3258 IAvH27233); 4.69617 -75.61056, 1920 m, 28-30 Aug. 2002, E. Jiménez & L.E. Franco, 2  
3259 workers (IAvH: IAvH27239); **Risaralda:** Pereira, Vereda El Manzano, Finca Santa Isabel,  
3260 4.70515 -75.62377, 1860 m, 15-17 July 2002, E. Jiménez & L.E. Franco, 3 workers (IAvH:  
3261 IAvH27234); Vereda La Suiza, Finca El Amparo de Niños, 4.74624 -75.59830, 1810 m, 28-30  
3262 Nov. 2002, L.E. Franco, 3 workers (IAvH: IAvH27273); 4.75013 -75.60278, 1780 m, 26-28 Nov.  
3263 2002, L.E. Franco & E. Londoño, 2 workers (IAvH: IAvH27281); Finca Pez Fresco, 4.73838 -  
3264 75.58016, 1910 m, 22-24 Nov. 2002, E. Jiménez & M.F. Reina, 1 worker, 1 queen (IAvH:  
3265 IAvH27275); Finca Tesorito, 4.72141 -75.56186, 1940 m, 27-29 Nov. 2002, E. Jiménez, L.E.  
3266 Franco & E. Londoño, 2 workers, 1 queen (IAvH: IAvH27280); Santuario, 26 Feb. 2003, 1  
3267 worker (IAvH: IAvH27286). **ECUADOR: Pichincha:** Maquipucuna, 5 km, ESE Nanegal,  
3268 0.11667 -78.63333, 1500 m, 17 Aug. 1991, P.S. Ward #11503-19, 5 workers (MCZC:  
3269 USNMENT00757255, 00757283).

3270  
3271 **Diagnosis.** *Brachymyrmex modestus* morphologically resembles *B. donisthorpei* and *B. myops*  
3272 because these species have short dense hairs on the head and the mesosoma, short suberect hairs  
3273 on the scapes, eyes below the cephalic midline of the head, the metanotal groove is either absent  
3274 or narrower than the diameter of the metathoracic spiracles, their gaster bears dense pubescence,  
3275 and their bodies are yellowish. *Brachymyrmex modestus* differs from *B. donisthorpei* by having

3276 scapes that surpass the posterior margin of the head, and from *B. myops* by having a mesonotum  
3277 that bulges dorsally above the pronotum in lateral view.

3278  
3279 *Lectotype and paralectotypes measurements* (mm) (n=3) HL<sub>1</sub> 0.49-0.51; HL<sub>2</sub> n.a.; HL<sub>3</sub> 0.14-0.16;  
3280 HW 0.45; SL 0.43; EL 0.08-0.09; WL 0.49-0.55; PnL 0.10-0.14; PnW 0.29-0.39; ML 0.08-0.16;  
3281 MW 0.20-0.25; *Indices* CI 88.46 - 92.00; SI<sub>1</sub> 95.65; SI<sub>2</sub> n.a.; OI<sub>1</sub> 17.39-19.57; OI<sub>2</sub> 20.00-25.27.

3282  
3283 *Additional material examined measurements* (mm) (n=3). HL<sub>1</sub> 0.41-0.46; HL<sub>2</sub> 0.29-0.32; HL<sub>3</sub>  
3284 0.11-0.12; HW 0.38-0.45; SL 0.38-0.40; EL 0.06-0.07; WL 0.45-0.49; PnL 0.12-0.13; PnW 0.22-  
3285 0.28; ML 0.10-0.12; MW 0.18-0.22; *Indices* CI 91.49-100.00; SI<sub>1</sub> 90.00-97.78; SI<sub>2</sub> 120.00-  
3286 133.33; OI<sub>1</sub> 15.56-18.61; OI<sub>2</sub> 23.53-27.66.

3287  
3288 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin flat.  
3289 Dorsum of the head with appressed and several suberect hairs. Clypeus with a rounded anterior  
3290 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
3291 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
3292 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
3293 margin in oblique anterodorsal view. The scapes surpass the posterior margin of the head by a  
3294 length approximately equal to the maximal diameter of the eye and bear decumbent hairs. Ocelli  
3295 appear to be absent. Eyes are positioned below the cephalic midline and have 3-4 ommatidia  
3296 along their maximal diameter.

3297 **Mesosoma.** Usually with two erect hairs on the pronotum and two on the mesonotum; sometimes  
3298 additional decumbent hairs are present, mainly on the pronotum. The mesonotum is inflated,  
3299 somewhat anteriorly inclined, and bulges dorsally above the pronotum in lateral view. Metanotal

3300 groove usually absent or narrower than the diameter of the metathoracic spiracles. Metathoracic  
3301 spiracles small and in dorsolateral position, not protruding, and touching the proprodeal suture.  
3302 Dorsum of the propodeum convex and shorter than the propodeal slope. Propodeal spiracles  
3303 circular and positioned on the posterior propodeal margin, ventrally and slightly posterior of the  
3304 middle of the propodeal slope. Legs with appressed hairs. Petiole short and inclined forward.

3305 **Gaster.** With dense pubescence and some long erect hairs at the edges of the segments.

3306 **Color and sculpture.** Head and gaster smooth and shiny, dorsum of the mesosoma slightly  
3307 imbricate. Body yellowish, with the gaster sometimes being darker in color.

3308

3309 **Distribution (Supplementary material Fig. S28).** *Brachymyrmex modestus* is known from  
3310 Brazil, Colombia and Ecuador.

3311

3312 **Biology.** The type material of *B. modestus* was collected in association with termites (Santschi  
3313 1923b).

3314

3315 **Remarks.** The here designated lectotype is the specimen without expanded gaster on pin NHMB:  
3316 USNMENT00758099, whereas the other specimen is one of the paralectotypes.

3317 Santschi (1923b) mentioned the presence of physogastric workers in *B. modestus*, and we also  
3318 observed some physogastric individuals in other samples (ICN: MZSP138, 149). These  
3319 specimens are characterized by having a larger body in comparison to regular workers, and they  
3320 have shorter scapes that barely reach the posterior margin of the head.

3321

3322

3323 *Brachymyrmex musculus* Forel

3324 (Fig. 41, supplementary material Fig. S29)

3325 *Brachymyrmex tristis* r. *musculus* Forel, 1899: 124 (w.). **Lectotype worker** (MHNG:

3326 USNMENT00757155) and **paralectotype workers** (MHNG: USNMENT00757153-00757155;

3327 USNMENT00758113-00758115; MCSN: USNMENT00757152; **here designated**): 16 workers

3328 [examined]: **COSTA RICA:** Pittier; Forel (1901a: 135) (q.). Raised to species: Forel (1901a:

3329 135). See also: Santschi (1923a: 673).

3330

3331 **Additional material examined. ARGENTINA: Mendoza:** 22.81 km W Villa Seca, -33.58515 -

3332 69.41708, 1835 m, 06 Jan. 2008, W. MacKay, 1 worker (WEMC: USNMENT00757974); **Santa**

3333 **Cruz:** 12 km S Bajo Caracoles, 640 m, E.I. Schlinger & M.E. Trwin, ANTC10276, 1 worker

3334 (CASC: CASENT0196018). **COLOMBIA: Quindio:** Filandia, Vereda Cruces, Finca Los

3335 Micos, 4.70239 -75.64665, 1800 m, 12-14 July 2002, E. Jimenez & L.E. Franco, 1 worker

3336 (IAvH: IAvH27235). **COSTA RICA: Puntarenas:** Monteverde, 10.301 -84.806, 1500 m, 27

3337 June 1984, J. Longino, 2 workers, 1 queen (JTLC: JTLC000005951, JTLC000005953).

3338 **MEXICO:** Mexico, 6.5 km E Chalma, 26 May 1988, W.P. MacKay #10386, 1 worker, 2 males,

3339 1 queen (WEMC: USNMENT00759016). **PARAGUAY: Amambay:** Pedro Juan Caballero, -

3340 22.567 -55.617, 20 Aug. 1998, A. Wild #AW0642, 1 worker (ALWC: USNMENT00757965).

3341 **VENEZUELA: Lara:** 9 km SE Barbacoas, 9.77 -71.06, 2000 m, 22 Aug. 1987, P.S. Ward

3342 #8923, 3 workers (PSWC: USNMENT00757589).

3343

3344 **Diagnosis.** *Brachymyrmex musculus* morphologically resembles *B. bruchi*, *B. patagonicus* and *B.*

3345 *oculatus*, because all four species have scapes that surpass the posterior margin of the head, a

3346 mesonotum that does not bulge dorsally above the pronotum in lateral view, a gaster with

3347 scattered pubescence and a brownish body. However, *B. musculus* differs from the three other

3348 species by having a metanotal groove that is wider than the diameter of the metathoracic  
3349 spiracles.

3350  
3351 *Lectotype and paralectotype measurements* (mm) (n=5). HL<sub>1</sub> 0.43-0.48; HL<sub>2</sub> 0.30-0.33; HL<sub>3</sub>  
3352 0.10-0.11; HW 0.39-0.42; SL 0.39-0.42; EL 0.10-0.12; WL 0.42-0.46; PnL 0.11; PnW 0.25-0.29;  
3353 ML 0.08-0.10; MW 0.18-0.20; *Indices* CI 87.16-91.26; SI<sub>1</sub> 98.92-100.00; SI<sub>2</sub> 117.65-136.76; OI<sub>1</sub>  
3354 23.16-30.00; OI<sub>2</sub> 21.10-23.81.

3355  
3356 *Additional material examined measurements* (mm) (n=3) HL<sub>1</sub> 0.48-0.51; HL<sub>2</sub> 0.33-0.37; HL<sub>3</sub>  
3357 0.10-0.12; HW 0.44-0.47; SL 0.43-0.47; EL 0.12-0.14; WL 0.35-0.51; PnL 0.10-0.14; PnW 0.30-  
3358 0.32; ML 0.10-0.12; MW 0.19-0.20; *Indices* CI 89.09-92.59; SI<sub>1</sub> 97.87-102.04; SI<sub>2</sub> 125.00-  
3359 131.58; OI<sub>1</sub> 26.00-30.61; OI<sub>2</sub> 21.15-24.07.

3360  
3361 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
3362 concave. Dorsum of the head has sparse appressed hairs. Clypeus with the medial anterior margin  
3363 somewhat forming a lip and five long, erect hairs of which a single, usually conspicuous hair is  
3364 near the anterior margin, two hairs are in mediolateral position and two more are near the toruli;  
3365 other hairs on the clypeus are markedly shorter and appressed or decumbent. Toruli surpassing  
3366 the posterior clypeal margin in oblique anterodorsal view. The scapes surpass the posterior  
3367 cephalic margin with a length up to the maximal diameter of the eye; with appressed hairs. Three  
3368 ocelli are present but inconspicuous. Eyes are positioned on the cephalic midline with 9-11  
3369 ommatidia along their maximal diameter.

3370 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum;  
3371 sometimes with additional suberect hairs mainly on the pronotum. The mesonotum is not inflated

3372 and does not bulge dorsally above the pronotum in lateral view. Metanotal groove wider than the  
3373 diameter of the metathoracic spiracles. Metathoracic spiracles in dorsolateral position, not  
3374 protruding, and close to the propodeal suture. Dorsum of the propodeum convex and shorter than  
3375 the posterior propodeal slope. Propodeal spiracles circular, positioned on the posterior propodeal  
3376 margin at the middle of the propodeal slope. Legs with appressed hairs. Petiole short and inclined  
3377 forward.

3378 **Gaster.** With scattered pubescence and several scattered long erect hairs.

3379 **Color and sculpture.** Body overall smooth and shiny, except for the sometimes slightly imbricate  
3380 sculpture on the dorsum of the mesosoma. Body typically brownish, however, the bulbi of the  
3381 antennae are whitish, and the antenna and legs, especially the tarsi, are somewhat lighter in color  
3382 than the body.

3383

3384 **Distribution (Supplementary material Fig. S29).** *Brachymyrmex musculus* is known from  
3385 Argentina, Colombia, Costa Rica, Mexico, Paraguay and Venezuela.

3386

3387 **Biology.** Unknown.

3388

3389 **Remarks.** The lectotype is here designated as the ant at the top of pin MHNG:

3390 USNMENT00757155, whereas the other specimens in the pin are paralectotypes.

3391 Forel (1899) originally described *B. musculus* as a race of *B. tristis*, but later indicated that the  
3392 difference in size urged him to elevate it to a separate species (Forel 1901a). We agree with this  
3393 decision, because *B. tristis* differs from *B. musculus* by having dense decumbent hairs on the  
3394 head and the dorsum of the mesosoma, a mesonotum that bulges dorsally above the pronotum,  
3395 and metathoracic spiracles that protrude slightly.



3396

3397

3398

*Brachymyrmex myops* Emery

3399

(Fig. 42, supplementary material Fig. S30)

3400 *Brachymyrmex myops* Emery, 1906: 182, footnote, Fig. 42 (w.m.). **Lectotype worker** (NHMB:

3401 USNMENT00757221) and **paralectotype male** (NHMB: USNMENT00757221; **here**

3402 **designated**): 1 worker, 1 male [examined]. **BRAZIL: Santa Catarina:** Joinville, J. P. Schmalz,

3403 leg. See also: Santschi (1923a: 663).

3404

3405 **Additional material examined. BOLIVIA: Santa Cruz:** 35 km SSE Flor de Oro, -13.833 -

3406 60.867, 450 m, 29 Nov. 1993, P.S. Ward #12199-74, 5 workers (MCZ: USNMENT00757254,

3407 00757892), 3 workers (PSWC: USNMENT00758027); **BRAZIL: Amazonia:** Rondonia, Jerau,

3408 R.M. Feitosa, 5 workers (ICN: MZSP178); **Pará:** Belém, 07 Aug. 1962, P.F. Doulington. B.F.

3409 14, 1 worker (MCZC: USNMENT00757257); 14 Aug. 1962, P.F. Doulington. B.F. 19, 2 workers

3410 (MCZC: USNMENT00757258); 14 Aug. 1962, P.F. Doulington. B.F. 19, 1 worker (MCZC:

3411 USNMENT00757259); Melgaço, Caxiuanã ECFPn, -1.70661 -51.45909, 01 Nov. 2003, A.Y.

3412 Harada, E.P. Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R. Moura, J.L.P. Souza, C. Renato,

3413 1 worker (MPEG: AYH094); 10-12 Oct. 2006, Equipe A.Y. Harada, 1 worker (MPEG:

3414 AYH046); 10-12 Oct. 2007, Equipe A.Y. Harada, 2 workers (MPEG: AYH038); 20-22 Oct.

3415 2006, Equipe A.Y. Harada, 8 workers (MPEG: AYH058); 20-22 Oct. 2007, Equipe A.Y. Harada,

3416 4 workers (MPEG: AYH021, 047, 059); 25-27 Oct. 2007, Equipe A.Y. Harada, 3 workers

3417 (MPEG: AYH039, 048, 099); -1.73584 -51.48762, 13-15 Oct. 2005, Equipe A.Y. Harada, 5

3418 workers (MPEG: AYH037); 25-27 Jan. 2007, Equipe A.Y. Harada, 1 worker (MPEG: AYH111);

3419 -1.78155 -51.59753, 15-17 Oct. 2007, Equipe A.Y. Harada, 3 workers (MPEG: AYH013, 044);

3420 24-26 Oct. 2007, Equipe A.Y. Harada, 1 worker (MPEG: AYH029); 25-27 Jan. 2007, Equipe  
3421 A.Y. Harada, 1 worker (MPEG: AYH062); 27-29 Jan. 2006, Equipe A.Y. Harada, 5 workers  
3422 (MPEG: AYH034); 30 July-01 Aug. 2003, Equipe A.Y. Harada, 13 workers (MPEG: AYH134);  
3423 A.Y. Harada, E.P. Fagundes, C. Renato, 3 workers (MPEG: AYH129). **COLOMBIA: Cauca:**  
3424 PNN Gorgona, Mancora, 2.967 -78.183, 60 m, 02 Feb. 2000, D. Campos, 16 workers (IAvH).

3425  
3426 **Diagnosis.** *Brachymyrmex myops* morphologically resembles *B. donisthorpei* and *B. modestus*  
3427 because all three species have dense short hairs on the head and mesosoma, scapes with short  
3428 suberect hairs, eyes that are positioned below the cephalic midline of the head, a metanotal  
3429 groove that is absent or narrower than the diameter of the metathoracic spiracles, a gaster with  
3430 dense pubescence, and yellowish body color. However, *B. myops* differs from *B. donisthorpei* by  
3431 having scapes that surpass the posterior margin of the head and from *B. modestus* by having a  
3432 mesonotum that does not bulge dorsally above the pronotum.

3433  
3434 *Lectotype measurements* (mm) HL<sub>1</sub> 0.41; HL<sub>2</sub> 0.29; HL<sub>3</sub> 0.12; HW 0.37; SL 0.35; EL 0.06; WL  
3435 0.41; PnL 0.14; PnW 0.25; ML 0.12; MW 0.20; *Indices* CI 90.48; SI<sub>1</sub> 94.74; SI<sub>2</sub> 120.00; OI<sub>1</sub>  
3436 15.79; OI<sub>2</sub> 28.57.

3437  
3438 *Additional material examined measurements* (mm) (n=3). HL<sub>1</sub> 0.37-0.38; HL<sub>2</sub> 0.27-0.30; HL<sub>3</sub>  
3439 n.a.; HW 0.33-0.34; SL 0.34-0.36; EL 0.05-0.06; WL 0.36; PnL 0.09-0.10; PnW 0.22-0.23; ML  
3440 0.06-0.07; MW 0.15-0.18; *Indices* CI 88.10-88.37; SI<sub>1</sub> 102.70-105.40; SI<sub>2</sub> 111.76-130.00; OI<sub>1</sub>  
3441 15.79-18.92; OI<sub>2</sub> n.a.

3442

3443 **Description. *Head.*** Slightly longer than wide in full face view; posterior cephalic margin slightly  
3444 concave. Dorsum of the head with appressed and several suberect hairs. Clypeus with a rounded  
3445 anterior margin and five long, erect hairs of which a single, usually conspicuous hair is near the  
3446 anterior margin, two hairs are in mediolateral position and two more are near the toruli; other  
3447 hairs on the clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the  
3448 posterior clypeal margin in oblique anterodorsal view. The scapes surpass the posterior margin of  
3449 the head by a length approximately equal to the maximal diameter of the eye and have decumbent  
3450 and suberect hairs. Ocelli absent. Eyes are positioned below the cephalic midline and have 3-4  
3451 ommatidia along their maximal diameter.

3452 ***Mesosoma.*** Usually two erect hairs on the pronotum and two on the mesonotum; sometimes  
3453 decumbent hairs are present, mainly on the pronotum. The mesonotum is somewhat inflated, but  
3454 it does not bulge dorsally above the pronotum in lateral view. Metanotal groove absent or  
3455 narrower than the diameter of the metathoracic spiracles. Metathoracic spiracles inconspicuous,  
3456 in dorsolateral position, not protruding, and not touching any suture. Dorsum of the propodeum  
3457 weakly convex and much shorter than the propodeal slope. Propodeal spiracles circular but  
3458 inconspicuous, positioned on the posterior propodeal margin, slightly posterior of the middle of  
3459 the propodeal slope. Legs with appressed hairs. Petiole short and inclined forward.

3460 ***Gaster.*** With dense pubescence and several long erect hairs, mainly along the edges of the  
3461 segments.

3462 ***Color and sculpture.*** Head and gaster smooth and shiny, dorsum of the mesosoma slightly  
3463 imbricate. Body usually uniformly yellowish, but sometimes with a darker gaster

3464

3465 **Distribution (Supplementary material Fig. S30).** *Brachymyrmex myops* is known from Bolivia,  
3466 Brazil and Colombia.

3467  
3468 **Biology.** Emery (1906) described *B. myops* from specimens that were collected in the nest of the  
3469 termite *Anoplotermes ater*.

3470  
3471 **Remarks.** The worker on pin NHMB: USNMENT00757221 is designated here as the lectotype.  
3472 *Brachymyrmex myops* has been described from one worker and a male, which makes it currently  
3473 impossible to determine the intraspecific variation. *Brachymyrmex modestus* has also been  
3474 collected from termite nests, has a very similar geographic distribution as *B. myops*, and both  
3475 species resemble one another morphologically (see diagnosis). They differ in whether the  
3476 mesonotum bulges dorsally above the pronotum in lateral view, which is a trait of diagnostic  
3477 value to delimit several *Brachymyrmex* species, however, the conditions in *B. myops* and *B.*  
3478 *modestus* are not strongly different, and both species may be conspecific.

3479

3480

3481 ***Brachymyrmex nebulosus* LaPolla & Longino**

3482 **(Fig. 43, supplementary material Fig. S31)**

3483 *Brachymyrmex nebulosus* LaPolla and Longino, 2006: 299, Fig. 1 (w.). **COSTA RICA:**

3484 **Puntarenas:** 6 km south of Monteverde, 10.25 -84.82, 800 m, 22 June 1999, J. Longino #4050,  
3485 LACM ENT 143550 (INBC).

3486

3487 **Paratype examined. COSTA RICA: Puntarenas:** Ojo de agua, rd to Monteverde, 800 m, 05  
3488 July 1991, J. Longino, leg. (JTLC #2965), 2 workers (INBIO).

3489

3490 **Additional material examined. MEXICO: Chiapas:** Sierra Morena, 16.15427 -93.58961, 1150  
3491 m, 11 May 2008, J. Longino #6218-s, 2 workers (JTLC: JTLC0000007379; CASENT0609689).

3492  
3493 **Diagnosis.** *Brachymyrmex nebulosus* differs from other *Brachymyrmex* species in having a  
3494 clypeus with its medial portion forming a conspicuous “lip”, its hour-glass shaped mesosoma and  
3495 it has portions of the head and mesosoma that bear alveolate sculpture. *Brachymyrmex musculus*  
3496 is the only other *Brachymyrmex* species known to date that has a clypeus with a somewhat  
3497 developed medial lip, but it is less conspicuous than in *B. nebulosus*.

3498  
3499 *Paratype measurements* (mm). HL<sub>1</sub> 0.67; HL<sub>2</sub> 0.40; HL<sub>3</sub> 0.24; HW 0.63; SL 0.62; EL 0.18; WL  
3500 0.80; PnL 0.29; PnW 0.45; ML 0.26; MW 0.23; *Indices* CI 94.67; SI<sub>1</sub> 97.18; SI<sub>2</sub> 153.33; OI<sub>1</sub>  
3501 28.17; OI<sub>2</sub> 36.00.

3502  
3503 *Additional material examined measurements* (mm) (n=1). HL<sub>1</sub> 0.80; HL<sub>2</sub> 0.49; HL<sub>3</sub> 0.23; HW  
3504 0.73; SL 0.72; EL 0.20; WL 0.92; PnL 0.31; PnW 0.50; ML 0.12; MW 0.27; *Indices* CI 91.11;  
3505 SI<sub>1</sub> 98.78; SI<sub>2</sub> 111.76; OI<sub>1</sub> 15.79; OI<sub>2</sub> n.a.

3506  
3507 **Description.** See LaPolla and Longino (2006).

3508  
3509 **Distribution** (supplementary material Fig. S31). This species is known from Costa Rica and  
3510 Mexico.

3511  
3512 **Biology.** *Brachymyrmex nebulosus* was collected from sites at about 800 m elevation in moist  
3513 forest at the transition between lowland dry forest and cloud forest. All collected workers were

3514 obtained from open scrubby vegetation. In the field, these ants look and behave remarkably like  
3515 *Crematogaster* (LaPolla and Longino 2006).

3516  
3517 **Remarks.** *Brachymyrmex nebulosus* has been originally described from Costa Rica, but during  
3518 our studies we came across two specimens from Mexico that very strongly resemble this species.  
3519 The only differences are that the Mexican specimens have a more squared head, and stronger  
3520 alveolate sculpture on the head and the dorsum of the mesosoma. Considering these differences  
3521 and the geographical distribution, these Mexican specimens may be a variety of *B. nebulosus*, or  
3522 potentially a different species, although more material and further study would be required to  
3523 resolve this issue.

3524

3525 *Brachymyrmex obscurior* Forel

3526 (Fig. 44, supplementary material Fig. S32)

3527 *Brachymyrmex heeri* var. *obscurior* Forel, 1893: 345 (w.q.m.). **Lectotype worker** (MHNG:  
3528 USNMENT00757132) and **paralectotype workers, queens, males** (MHNG:  
3529 USNMENT00757132-00757135; USNMENT00758124-00758128; **here designated**): 16  
3530 workers, 3 queens, 3 males [examined]. **ANTILLES:** Saint Vincent. Subspecies of  
3531 *Brachymyrmex heeri*: Forel (1897: 298); Forel (1912a: 62). Raised to species: Wilson and Taylor  
3532 (1967: 92). See also Santschi (1923a: 666).

3533

3534 **Additional material examined. ARGENTINA: Chubut:** 3 km N. Puerto Lobos, 20 m, 14 Dec.  
3535 1966, E.I. Schlinger & M. Irwin, ANTC 10275, 1 worker (CASC: CASENT0196017). **BRAZIL:**  
3536 **Bahia:** CEPEC, 11 Nov. 1997, L.S. Ramos, 1 worker (CPDC: USNMENT00757668); **São**  
3537 **Paulo:** Caraguatatuba, Reserva Florestal, 40-80 m, 12-22 May 1971, W.L. & D.E. Brown, 2

3538 workers, 1 queen (MCZC: USNMENT00757659); Itirapina, cerrado, Dec. 2008, D.P. Silva, 3  
3539 workers (ICN: USNMENT00759042); Ubatuba, Picinguaba, July 2011, 2 workers (ICN:  
3540 USNMENT00759051). **CHILE:** Santiago, Nov. 1996, C. Errard, 3 workers (CPDC:  
3541 USNMENT00757683). **COLOMBIA: Córdoba:** Monteria, Finca Betania, 29 June 2009, Juan  
3542 C. Abadia, 4 workers (IAvH); Valencia, Villa Mary, 02 June 2009, Juan C. Abadia, 4 workers  
3543 (IAvH); **Huila:** 17 km NW La Plata, 03 Jan. 1984, W.P. MacKay #7133, 1 worker (WEMC:  
3544 USNMENT00759021); 17km N La Plata, 03 Jan. 1984, W.P. MacKay #7139, 1 worker 1 queen  
3545 (WEMC: USNMENT00758996); **Magdalena:** Parque Nacional Natural Tayrona, Cañaveral,  
3546 11.33 -74.03, 30 m, 18-20 Aug. 2002, M. Sharkey, P. Arias & E. Torres, 7 workers (IAvH);  
3547 **Valle del Cauca:** Cairo, vereda Llano Grande, Finca Encanto, 4.73620 -76.21698, 1550 m, 31  
3548 Mar. 2003, R. Garcia, 1 worker (IAvH: IAvH25144). **DOMINICAN REPUBLIC: La Vega:** 12  
3549 km NW Bona, 19.03333 -70.48333, 890 m, 31 Aug. 2001, A.L. Wild #AW1324, 2 workers  
3550 (ALWC: USNMENT00757987); 5 km N El Río, 19.02 -70.60, 1230 m, 01 Sep. 2001, A.L. Wild  
3551 #AW1339, 1 worker (ALWC: USNMENT00757657); Casabito For. El Rio - Bona Km 8, 07 Feb.  
3552 1975, W.L.& D.E. Brown, 2 workers (MCZC: USNMENT00757664). **FRENCH GUIANA:** 20  
3553 km Sinnamary, 12 Feb 1994, A. Dejean #17025, 2 workers (ICN: USNMENT00757660).  
3554 **GUATEMALA: El Progreso:** 5 km El Rancho, 14.9167 -90.0667, 400 m, 17 Nov. 2003, P.S.  
3555 Ward #15076-3, 3 workers (PSWC: USNMENT00757667); **Escuintla:** Escuintla, 30 Dec. 1911,  
3556 W.M. Wheeler, 3 workers (MCZC: USNMENT00759001). **JAMAICA:** Manchester, Gourie  
3557 Forest Res., 18.20 -77.52, 860 m, 10 Sep. 2001, A.L. Wild #AW1375, 1 worker (ALWC:  
3558 USNMENT00757669). **MEXICO: Chiapas:** Custepec, 15.72196 -92.95037, 1530 m, 19 May  
3559 2008, J. Longino #6280, 1 worker, 1 queen (JTLC: JTLC000007437, JTLC 000007438); **Jalisco:**  
3560 7 km SW Tamazula, 19.68056 -103.32194, 992 m, 22 June 2000, W. & E. Mackay, 2 workers  
3561 (WEMC: USNMENT00757729). **NEW CALEDONIA:** Road to My. Koghi, Dec. 1985, N.L.H.

3562 Krauss, ANTC10279, 1 worker (CASC: CASENT0196021). **UNITED STATES: Florida:**  
3563 Sarasota Co. Longino Ranch. T38S, R22E Sect, 27.15 -82.12, 20 m, 07 June 1981, J. Longino, 2  
3564 workers (JTLC: JTLC0000005943); **Texas:** Del Mar, 26.01167 -97.31861, 26 Sep. 1972, W.S.  
3565 Ross, ANTC 10263 10264, 2 workers (CASC: CASENT0196005, 0196006).

3566  
3567 **Diagnosis.** *Brachymyrmex obscurior* morphologically resembles *B. cordemoyi* and *B.*  
3568 *patagonicus* because all three species have a metanotal groove that is absent or narrower than the  
3569 diameter of metathoracic spiracles, their mesonotum does not bulge dorsally above the pronotum,  
3570 their scapes usually surpass the posterior margin of the head, and their bodies are brownish.  
3571 *Brachymyrmex obscurior* and *B. cordemoyi* differ from *B. patagonicus*, however, because they  
3572 have dense pubescence on the gaster. *Brachymyrmex obscurior* differs from *B. cordemoyi* by  
3573 having less conspicuous dense pubescence on the dorsum of the head and the mesosoma, dense  
3574 decumbent pubescence on the gaster, and eyes with fewer ommatidia along their maximal  
3575 diameter (on average 9 instead of 10-12).

3576  
3577 *Lectotype and paralectotypes workers measurements* (mm) (n=8). HL<sub>1</sub> 0.39-0.47; HL<sub>2</sub> 0.27-0.35;  
3578 HL<sub>3</sub> 0.10-0.12; HW 0.35-0.48; SL 0.35-0.45; EL 0.10-0.15; WL 0.31-0.53; PnL 0.09-0.15; PnW  
3579 0.23-0.30; ML 0.06-0.14; MW 0.15-0.23; *Indices* CI 79.17-106.67; SI<sub>1</sub> 96.59-115.79; SI<sub>2</sub> 125.71-  
3580 1146.67; OI<sub>1</sub> 25.00-31.25.

3581  
3582 *Additional material examined measurements* (mm) (n=2). HL<sub>1</sub> 0.44-0.47; HL<sub>2</sub> 0.30-0.32; HL<sub>3</sub>  
3583 0.11-0.12; HW 0.37-0.42; SL 0.38-0.41; EL 0.11-0.12; WL 0.42-0.47; PnL 0.11-0.13; PnW 0.26-  
3584 0.30; ML 0.11; MW 0.18-0.19; *Indices* CI 84.34-88.89; SI<sub>1</sub> 96.25-102.86; SI<sub>2</sub> 126.32-128.33; OI<sub>1</sub>  
3585 28.57-28.75; OI<sub>2</sub> 24.10-24.44.



3586

3587 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
3588 concave. Dorsum of the head with sparse appressed hairs. Clypeus with a rounded anterior  
3589 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
3590 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
3591 clypeus are clearly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
3592 margin in oblique anterodorsal view. The scapes surpass the posterior cephalic margin by a  
3593 length up to the maximal diameter of the eye, and they have appressed hairs. Ocelli absent. Eyes  
3594 are positioned on the cephalic midline and have 8-10 ommatidia along their maximal diameter.

3595 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum. The  
3596 mesonotum is not inflated and does not bulge dorsally above the pronotum in lateral view.  
3597 Metanotal groove absent or narrower than the diameter of the metathoracic spiracles.  
3598 Metathoracic spiracles in dorsolateral position, not protruding, and typically touching the  
3599 mesometanotal and propodeal sutures. Dorsum of the propodeum convex and shorter than the  
3600 propodeal slope. Propodeal spiracles circular and positioned on the posterior propodeal margin,  
3601 slightly anterior of the middle of the propodeal slope. Legs with appressed hairs. Petiole short and  
3602 inclined forward.

3603 **Gaster.** With decumbent dense pubescence and several scattered long erect hairs.

3604 **Color and sculpture.** Body smooth and shiny, except for the dorsum of the mesosoma which is  
3605 sometimes slightly imbricate. Body uniformly brownish, but with slightly lighter antenna and  
3606 legs.

3607

3608 **Distribution (supplementary material Fig. S32).** *Brachymyrmex obscurior* is widespread and  
3609 known from the Antilles, Argentina, Brazil, Chile, Colombia, the Dominican Republic, French

3610 Guiana, Guatemala, Hawaii, Jamaica and Mexico. It has also been introduced in New Caledonia,  
3611 the United States and other colleagues have reported it from Samoa and Hawaii (Wilson and  
3612 Taylor, 1967), but we did not examine this material.

3613  
3614 **Biology.** In the original description, Forel (1893) mentioned that this species forms colonies of a  
3615 few hundred individuals. Nests are usually constructed on open ground, typically under a stone,  
3616 or at the roots of grass and weeds. The nest consists of only one or two simple chambers that are  
3617 connected with a short passage. Forel (1893) indicated that *B. obscurior* occurs from sea level up  
3618 to 800 m of altitude, but here we report new records from localities above 800 m.

3619  
3620 **Remarks.** The specimen at the top of pin MHNG: USNMENT00757132 is designated here as the  
3621 lectotype, whereas the other ants are paralectotypes. Forel (1893) described *B. obscurior* as a  
3622 variety of *B. heeri* and indicated that it differs from typical *B. heeri* by having a brownish instead  
3623 of yellowish body and slightly denser pubescence on the gaster. We observed that both species  
3624 can readily be distinguished as to whether or not the mesonotum bulges dorsally above the  
3625 pronotum in lateral view. Forel (1893) also reported that *B. obscurior* resembles *B. patagonicus*  
3626 but that both taxa differ as to the presence or absence of ocelli, in body size, as to pubescence and  
3627 the length of the scapes. He further considered *B. obscurior* to be a difficult “form” that  
3628 represents a morphological transition between *B. patagonicus* and *B. heeri* (Forel 1912a). We  
3629 agree that *B. obscurior* and *B. patagonicus* are morphologically very similar (see diagnosis), and  
3630 molecular studies of both taxa will be required to examine whether they are distinct species.  
3631 Wilson and Taylor (1967) recognized *B. heeri* var. *obscurior* as a distinct species as a provisional  
3632 measure in anticipation of a full-scale revision of the genus. We agree with this decision, based  
3633 on the morphological differences indicated above, but we cannot for now comment on their

3634 proposed synonymization of *B. heeri* var. *aphidicola* Wheeler, 1934 to *B. obscurior*, as this  
3635 material from Hawaii was not available to us.

3636

3637 ***Brachymyrmex oculatus* Santschi**

3638 **(Fig. 42, supplementary material Fig. S33)**

3639 *Brachymyrmex oculatus* Santschi, 1919: 55 (w.). **Lectotype worker** (NHMB:  
3640 USNMENT00758101) and **paralectotype workers** (NHMB: USNMENT00758101; **here**  
3641 **designated**): 6 workers [examined]. **ARGENTINA: Buenos Aires:** Sierra de las Ventanas,  
3642 Bruch leg. Quirán et al. (2004: 282) (m).

3643

3644 **Additional material examined. ARGENTINA: Entre Ríos:** Vilcaguay, Bruchi, 3 workers  
3645 (MZSP: USNMENT00757776); **Santa Fé:** Fives Lilles, Wisser, 6 workers (MCZC:  
3646 USNMENT00757250).

3647

3648 **Diagnosis.** *Brachymyrmex oculatus* morphologically resembles *B. bruchi* and *B. patagonicus*,  
3649 because all three species have scapes that surpass the posterior margin of the head, they usually  
3650 have two erect hairs on the mesonotum, their mesonotum does not bulge dorsally above the  
3651 pronotum in lateral view, a metanotal groove is absent or narrower than the diameter of the  
3652 metathoracic spiracles, their gaster has scarce pubescence and several scattered long erect hairs  
3653 and their body is brownish. *Brachymyrmex oculatus* differs from *B. bruchi* and *B. patagonicus*,  
3654 however, by having larger eyes, with a maximal diameter that approximates a third of the length  
3655 of the head (HL<sub>1</sub>). They usually have more than 14 ommatidia along their maximal diameter.

3656

3657 *Lectotype and paralectotypes worker measurements* (mm) (n=4). HL<sub>1</sub> 0.45-0.50; HL<sub>2</sub> 0.35-0.41;  
3658 HL<sub>3</sub> 0.11-0.14; HW 0.48-0.58; SL 0.45-0.50; EL 0.15-0.18; WL 0.53-0.61; PnL 0.15-0.17; PnW  
3659 0.30-0.39; ML 0.14-0.20; MW 0.23-0.27; *Indices* CI 106.67-115.15; SI<sub>1</sub> 86.84-93.75; SI<sub>2</sub> 122.22-  
3660 133.33; OI<sub>1</sub> 31.25-34.29; OI<sub>2</sub> 21.88-27.27.

3661  
3662 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic border flat.  
3663 Dorsum of the head with sparse appressed hairs. Clypeus with a rounded anterior margin and five  
3664 long, erect hairs of which a single, usually conspicuous hair is near the anterior margin, two hairs  
3665 are in mediolateral position and two more near the toruli; other hairs on the clypeus are markedly  
3666 shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique  
3667 anterodorsal view. The scapes surpass the posterior cephalic margin by a length that is shorter  
3668 than the maximal diameter of the eye; they have appressed hairs. Ocelli absent. Eyes are  
3669 positioned on the cephalic midline and usually have more than 14 ommatidia along their maximal  
3670 diameter.

3671 **Mesosoma.** Dorsum of the mesosoma with sparse appressed hairs, and typically with two erect  
3672 hairs on the pronotum and two on the mesonotum; sometimes with additional suberect hairs,  
3673 mainly on pronotum. The mesonotum is not inflated and does not bulge dorsally above the  
3674 pronotum in lateral view. Metanotal groove absent or narrower than the diameter of the  
3675 metathoracic spiracles. Metathoracic spiracles in dorsolateral position, not protruding, and  
3676 touching the propodeal suture. Dorsum of the propodeum slightly convex and shorter than the  
3677 propodeal slope. Propodeal spiracles circular, positioned on the posterior propodeal margin, at the  
3678 middle of the propodeal slope. Legs with scattered appressed hairs. Petiole short and inclined  
3679 forward.

3680 **Gaster.** With scattered pubescence and several scattered long erect hairs, mainly at the edges of  
3681 the segments.

3682 **Color and sculpture.** Body overall smooth and shiny, with the dorsum of the mesosoma slightly  
3683 imbricate. Body uniformly brownish.

3684  
3685 **Distribution (Supplementary material Fig. S33).** *Brachymyrmex oculatus* is exclusively known  
3686 from Argentina. Quirán et al. (2004) examined material from additional localities throughout the  
3687 country.

3688  
3689 **Biology.** Unkown.

3690  
3691 **Remarks.** The specimen at the top of pin NHMB: USNMENT00757132 is here designated as  
3692 lectotype, whereas the other ants in that pin are paralectotypes. Santschi (1919) suggested that *B.*  
3693 *oculatus* can be confused with *B. bruchi* based on overall similarity. *Brachymyrmex patagonicus*  
3694 is also very similar, and as mentioned in the diagnosis *B. oculatus* differs mainly from these  
3695 species by its larger eyes. Whereas *B. patagonicus* has two erect hairs on the pronotum and *B.*  
3696 *bruchi* usually more than two, the number of erect hairs on the pronotum is variable in *B.*  
3697 *oculatus*.

3698 Quirán et al. (2004) designated a male to be the lectotype of *B. oculatus*, because it was labelled  
3699 as “typus”, and the associated workers were designated to be paralectotypes. However, the  
3700 original description by Santschi (1919) exclusively described the worker morphology, and  
3701 therefore logic dictates that the lectotype should be a worker. As such, we redesignated a worker  
3702 of Santschi (1919) type series as lectotype here.

3703

3704

3705

*Brachymyrmex patagonicus* Mayr

3706

(Figs. 46, 47, supplementary material Fig. S34)

3707 *Brachymyrmex patagonicus* Mayr, 1868: 164 (w.m.). (NHMW), Emery (1906: 178) (q.).

3708 **ARGENTINA: Buenos Aires:** Rio Negro. See also: Santschi (1923a: 657).

3709 = *Brachymyrmex patagonicus* var. *atrátula* Santschi, 1923a: 657, Fig. 3 (w.). (NHMB:

3710 USNMENT00757695): 2 workers [examined] **ARGENTINA: Jujuy:** Alfarito; synonym

3711 proposed by Quirán et al. (2004: 275). See also: Santschi (1923a: 657).

3712 = *Brachymyrmex laevis* Emery, 1895: 216 (w.). (MSNG: USNMENT00757205, 00757206;

3713 MHNG: USNMENT00758130): 4 workers [examined] **CHILE:** Valdivia. See also: Forel (1908:

3714 400); Forel (1912a: 62); Santschi (1923a: 659). **n. syn.**

3715

3716

3717 **Additional material examined. ARGENTINA: Entre Rios:** 8.63 km W Concordia, -31.42303 -

3718 58.11672, 16 m, 26 Dec. 2007, W. MacKay #22667, 1 worker (WEMC: USNMENT00758002);

3719 **Las Heras:** Agua de las zorras, Paramillo de Uspallata, 13 km NW of Villavicencio, -32.48011 -

3720 69.16467, 2750 m, 3 workers 1 queen (MCZC: USNMENT00759011); **Mendoza:** 22.81 km W

3721 Villa Seca, -33.58515 -69.41708, 1835 m, 06 Jan. 2008, W. MacKay, 4 workers, 1 queen

3722 (WEMC: USNMENT 00757968, 00757973, 00759018, 00759020); **Santiago del Estero:** 2

3723 workers (ICN: USNMENT00759047). **BRAZIL: Pará:** Melgaço, Caxiuanã ECFPn, -1.73584 -

3724 51.48762, 23-25 Oct. 2005, Equipe A.Y. Harada, 5 workers (MPEG: AYH036); -1.72484 -

3725 51.42979, 26 Oct. 2003, A.Y. Harada, E.P. Fagundes, C.J.M. Ribeiro, C.E.D. Sanhudo, C.A.R.

3726 Moura, J.L.P. Souza, C. Renato, 2 workers (MPEG: AYH053); Santarem, Taperinha, -2.90 -

3727 54.33, July 1975, R. L. Jeanne, 3 workers (MCZC: USNMENT00757990); **São Paulo:**

3728 Cananéia, P.E. Ilha Do Cardoso, -22.30 -47.88, 02 Nov. 2007, C. Bottcher & E.R. Pereira, 1  
3729 worker (ICN: USNMENT00757730); Itirapina: 11 Feb. 2009, S. Sendoya, 23 workers (ICN:  
3730 USNMENT00759052). **COLOMBIA: Caldas:** Aranzazu, Vereda Alegrias, Finca Betania, La  
3731 Esperanza, 5.29831 -75.49053, 1990 m, 08-09 Aug. 2003, L.E. Franco & J. Cruz, 3 workers  
3732 (IAvH: 55576); Finca Villa Rosita, 5.30603 -75.48489, 1825 m, 06-08 Aug. 2003, L.E. Franco &  
3733 J. Cruz, 2 workers (IAvH: IAvH25467, 25468); Vereda Buenavista, Naranjal Finca Bizerta,  
3734 5.28431 -75.48942, 2065 m, 25-27 July 2003, L.E. Franco & J. Cruz, 1 worker (IAvH:  
3735 IAvH25471); 4.285 -75.489, 2065 m, 25-27 July 2003, L.E. Franco & J. Cruz, 2 workers, 1  
3736 queen (IAvH: IAvH27267); Finca La Herradura, 5.27936 -75.49744, 2020 m, 5-7 Aug. 2003,  
3737 L.E. Franco & J. Cruz, 1 worker (IAvH: IAvH27291); Vereda Chambery, Finca Las Garzas,  
3738 5.29481 -75.47292, 1980 m, 5-6 July 2003, L.E. Franco & J. Cruz, 3 workers (IAvH:  
3739 IAvH55563); Vereda Chupaleros, Finca Alegrias, 5.30633 -75.50028, 1960 m, 27-29 July 2003,  
3740 L.E. Franco & J. Cruz, 3 workers (IAvH: IAvH25466); Veredada El Eden, Finca Los  
3741 Guayacanes, 5.29606 -75.49428, 1984 m, 26 July 2003, L.E. Franco & J. Cruz, 1 worker, 1 male  
3742 (IAvH: IAvH27294); Vereda San José, Finca Casa Roja 5.33348 -75.48892, 1777 m, 07-09 Aug.  
3743 2003, L.E. Franco & J. Cruz, 2 workers (IAvH: IAvH25462, 25463); Salamina, Vereda El  
3744 Cedrito, Finca El Cedrito, 5.33197 -75.46785, 1960 m, 27-28 July 2003, L.E. Franco & J. Cruz, 3  
3745 workers (IAvH: IAvH25470, 25472); **Córdoba:** Monteria, 29 June 2009, Juan C. Abadia, 3  
3746 workers (ICN); **Cundinamarca:** Villeta, Conjunto Residencial Las Acacias, 5.01361 -74.47306,  
3747 11 Jan. 2010, C.M. Ortiz, 8 workers (ICN); **Quindío:** Filandia, Vereda Cruces, Finca Los Micos,  
3748 4.70452 -75.64665, 1800 m, 12-14 July 2002, E. Jimenez & L.E. Franco, 2 workers (IAvH:  
3749 IAvH27227, 27240); **Valle del Cauca:** bosque Yotoco, 1575 m, 23 June 1989, W.P. MacKay  
3750 #11720, 1 worker (WEMC: USNMENT00757995); Medio Calima, 24 June 1989, E. MacKay  
3751 #11744, 3 workers, 1 queen (WEMC: USNMENT00759012, 00759013). **COSTA RICA:**

3752 **Alajuela:** Juan Santa Maria airport, 9.98 -84.20, 900 m, 09 Jan. 1999, J. Longino #3958-s, 1  
3753 worker (JTLC: LACM ENT 142311); **Heredia:** Estación Biológica la Selva, 10.433 -84.017,  
3754 May 1994, J. Longino #3625, 1 worker (JTLC: INBIOCRI001260979); 10.423 -84.001, 50 m, 04  
3755 Aug. 2004, W. & E. Mackay #20890, 1 worker (WEMC: USNMENT00758039); 06 Aug. 2004,  
3756 W. & E. Mackay #20917, 1 worker (WEMC: USNMENT00757982); **Limon:** 3 km SSE Cahuita,  
3757 9.71667 -82.83333, 70 m, 24 Dec. 1983, P.S. Ward #6530-40, 3 workers (PSWC:  
3758 USNMENT00758007); **Puntarenas:** Estación Biológica Los Llanos, 10.30487 -84.83735, 1150  
3759 m, 28 Feb. 2004, J. Longino, 1 worker (JTLC: JTLC000005287); 6 km S Monteverde, 10.25 -  
3760 84.82, 800 m, 22 June 1999, J. Longino, 1 worker, 1 queen (JTLC: LACM ENT 143543).  
3761 **ECUADOR: Chimborazo:** Huigra, -2.29417 -78.98861, 1200 m, 18 Feb. 2004, Roger Vila I-  
3762 457, 2 workers (ICN: USNMENT00758037). **GUATEMALA: El Progreso:** 3.8 km E. San  
3763 Cristobal, 14.91850 -90.04075, 302 m, 19 July 2004, W. & E. MacKay #20586, 3 workers  
3764 (WEMC: USNMENT00759006); **Santa Rosa:** 5 km SW Cuilapa, 14.23333 -90.33333, 575 m,  
3765 14 Nov. 2003, A.L. Wild #AW2030, 1 worker (ALWC: USNMENT00758009); **Suchitepéquez:**  
3766 Finca Tarrales, 12.3 km N Patulul, 14.52256 -91.13642, 740 m, 30 July 2004, W. & E. Mackay  
3767 #20782, 2 workers (WEMC: USNMENT00757688, 00757983). **HONDURAS:** La Lima, 23 Jan.  
3768 1960, C. Evers, UFC-217-35 (6871), 5 workers, 1 male (MZSP: USNMENT00757621,  
3769 00757622). **JAPAN:** Hyogo, Kob, Port Island, 34.67 135.20, 18 Sep. 2007, M. Yoshimura, 1  
3770 worker (ICN: MY1862-12). **MEXICO: Federal:** Mexico City, 6.5 km E Chalma, 26 May 1998,  
3771 W. MacKay #10386, 1 worker, 2 males (WEMC: USNMENT00757993); **Guanajuato:** Highway  
3772 57, km 306, Rancho Jardin, 21.14224 -100.95341, 10 Aug. 1965, Cornell University, 1 worker  
3773 (MCZC: USNMENT00759002); **Jalisco:** 6 km N El Tuito, 20.3667 -105.3167, 730 m, 31 Dec.  
3774 1987, P.S. Ward #9327-11, 2 workers (PSWC: USNMENT00757679); **Nuevo Leon:** 8 km W.  
3775 Iturbide, 09 Nov. 1946, W.S. Ross, ANTC10261, 2 workers (CPDC: CASENT0196003);



3776 **Nayarit:** 19.3 km S Rosamorada, 21.94389 -105.20639, 51 m, 19 June 2000, W. & E. MacKay  
3777 #19126, 2 workers (WEMC: USNMENT00757983); **Oaxaca:** 148 km NE Oaxaca Rt175,  
3778 17.02647 -96.71947, 1210 m, 04 June 1988, W. MacKay #10825, 3 workers (WEMC:  
3779 USNMENT00757732); 1.6 km E Reforma, near Tuxtepec, 18.08078 -96.13677, 12-15 Aug.  
3780 1973, A. Newton, 1 worker (MCZC: USNMENT00757680); **Puebla:** 6.5 km W Izucar  
3781 Matamoros, 1220 m, 26 May 1988, W. MacKay, 1 worker, 1 male (WEMC:  
3782 USNMENT00757991); **San Luis de Potosi:** 10 km S San Luis Potosi, 21 May 1988, W.P.  
3783 MacKay #10307 #10308, 6 workers, 2 males, 1 queen (WEMC: USNMENT00757685,  
3784 00759015, 00759017); 16 km S San Luis Potosi, 21 May 1988, W.P. MacKay #10307, 2  
3785 workers, 1 male (WEMC: USNMENT00758006); 11 km N Cardenas, 1720 m, 09 June 1988, W.  
3786 MacKay #1095, 1 worker, 2 males (WEMC: USNMENT00759027); **Tamaulipas:** 32.3 km SE  
3787 Ciudad Victoria, 23.49161 -96.97775, 289 m, 24 Mar. 2008, W. & E. MacKay #22930, 2 workers  
3788 (WEMC: USNMENT00757681, 00757687); Gomez Farias, 25 Sep. 1987, A. Rebeles #10096, 1  
3789 worker (WEMC: USNMENT00757994); **Veracruz:** Los Tuxlas, July 2001, A. Pezon, 1 worker  
3790 (CPDC: USNMENT00757989); Xalapa, V. Rico Gray #17209, 4 workers (MCZC:  
3791 USNMENT00759026, 00758044). **PARAGUAY: Canindeyú:** Reserva Natural Bosque  
3792 Mbaracayú, Aguara Ñu, -24.18333 -55.28333, 240 m, 16 Nov. 2002, A. L. Wild, 1 worker  
3793 (ALWC: USNMENT00757971); **Itapúa:** Isla Yacyretá, -27.41667 -56.75417, 25 Sep. 1997, B.  
3794 Barrios #ibn 216, 1 worker (ALWC: USNMENT00757683); **Presidente Hayes:** 5 km SSE Pozo  
3795 Colorado, -23.55 -58.77, 140 m, 05 Dec. 2002, A.L. Wild #AW1764, 1 worker (ALWC:  
3796 USNMENT00758010). **URUGUAY: Salta:** Salta, Parque Municipal Benito Solari, 25 Dec.  
3797 2007, W. & E. MacKay #22634, 1 worker (WEMC: USNMENT00757731).  
3798

3799 **Diagnosis.** *Brachymyrmex patagonicus* morphologically resembles *B. bruchi* and *B. oculatus*,  
3800 because all three species have scapes that surpass the posterior margin of the head by a length  
3801 approximately equal to the maximum diameter of the eye or less, they usually have two erect  
3802 hairs on the mesonotum, which does not bulge dorsally above the pronotum in lateral view, the  
3803 metanotal groove is absent or narrower than the diameter of the metathoracic spiracles, their  
3804 gaster has scarce pubescence and several scattered long erect hairs, the body is uniformly  
3805 brownish. *Brachymyrmex patagonicus* differs from *B. bruchi*, however, by usually having two  
3806 erect hairs on the pronotum and two on the mesonotum and from *B. oculatus* by having smaller  
3807 eyes, with a maximal diameter of approximately 1/4<sup>th</sup> of the length of the head (HL<sub>1</sub>) and usually  
3808 with less than 14 ommatidia along their maximal diameter.

3809  
3810 *Types measurements* (mm) (n=2). HL<sub>1</sub> 0.45-0.53; HL<sub>2</sub> 0.33-0.37; HL<sub>3</sub> 0.10-0.13; HW 0.38-0.49;  
3811 SL 0.40-0.48; EL 0.14-0.17; WL 0.38-0.51; PnL 0.11-0.15; PnW 0.26-0.33; ML 0.07-0.11; MW  
3812 0.17-0.24; *Indices* CI 85.29-92.50; SI<sub>1</sub> 97.30-103.45; SI<sub>2</sub> 120.00-128.57; OI<sub>1</sub> 33.78-36.21; OI<sub>2</sub>  
3813 22.06-25.00.

3814  
3815 *Additional material examined measurements* (mm) (n=13). HL<sub>1</sub> 0.40-0.59; HL<sub>2</sub> 0.28-0.39; HL<sub>3</sub>  
3816 0.07-0.16; HW 0.33-0.51; SL 0.35-0.49; EL 0.09-0.14; WL 0.35-0.55; PnL 0.09-0.20; PnW 0.23-  
3817 0.35; ML 0.07-0.14; MW 0.15-0.23; *Indices* CI 81.40-93.33; SI<sub>1</sub> 92.00-128.95; SI<sub>2</sub> 116.22-  
3818 163.33; OI<sub>1</sub> 22.22-32.5; OI<sub>2</sub> 18.18-28.57.

3819  
3820 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
3821 concave. Dorsum of head with sparse appressed hairs. Clypeus with a rounded anterior margin  
3822 and five long, erect hairs of which a single, usually conspicuous hair is near the anterior margin,

3823 two hairs are in mediolateral position and two more near the toruli; other hairs on the clypeus are  
3824 markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in  
3825 oblique anterodorsal view. The scapes surpass the posterior cephalic margin by a length that is  
3826 shorter than the maximal diameter of the eye; they have appressed hairs. At least one central  
3827 ocellus is present. Eyes are positioned on the cephalic midline and have 8-12 ommatidia along  
3828 their maximal diameter.

3829 **Mesosoma.** Dorsum of the mesosoma with sparse appressed hairs, typically with two erect hairs  
3830 on the pronotum and two on the mesonotum. The mesonotum is not inflated and does not bulge  
3831 dorsally above the pronotum. Metanotal groove absent or narrower than the diameter of the  
3832 metathoracic spiracles. Metathoracic spiracles in dorsolateral position, not protruding, and  
3833 touching the propodeal suture. Dorsum of the propodeum slightly convex and shorter than the  
3834 propodeal slope. Propodeal spiracles circular, situated on the posterior propodeal margin, at the  
3835 middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole short and inclined  
3836 forward.

3837 **Gaster.** With scattered pubescence and several scattered long erect hairs, mainly at the edges of  
3838 the segments.

3839 **Color and sculpture.** Body overall smooth and shiny, except for the sometimes slightly imbricate  
3840 sculpture on the dorsum of the mesosoma, and typically uniformly brownish.

3841  
3842 **Distribution (supplementary material Fig. S34).** *Brachymyrmex patagonicus* is widespread and  
3843 known from Argentina, Brazil, Colombia, Chile, Costa Rica, Ecuador, Guatemala, Honduras,  
3844 Mexico, Paraguay, Uruguay and it has been introduced in Japan (Ortiz-Sepulveda pers. obs.) and  
3845 the United States (MacGown et al. 2007).

3846

3847 **Biology.** MacGown et al. (2007) indicated that *B. patagonicus* nests in a variety of habitats, both  
3848 natural and disturbed, ranging from pine forests over mixed forest and prairie to urban  
3849 environments. Colonies may contain many hundreds of workers packed into a small sheltered  
3850 area. Where this species is found, colonies are often abundant and within a few centimeters from  
3851 one another. Nests can be found in loose tree bark, at the base of plants, in soil, dead wood and  
3852 organic litter, or below natural and man-made objects. The species is considered a nuisance pest,  
3853 as both alates and foraging workers may enter man-made structures to forage and/or nest  
3854 (MacGown et al. 2007).

3855  
3856 **Remarks.** As Quirán et al. (2004) already indicated Mayr (1868) described *B. patagonicus* based  
3857 on specimens from Argentina. However, the specimens in the NHMW: USNMENT00757201-  
3858 00757204 that were examined and identified as types by Mayr are from Chile. Either the locality  
3859 indicated in the original description may be wrong, or specimens from Argentina may be lacking  
3860 from the collection, and as such we do not designate a lectotype here.

3861 Emery (1895), upon describing *B. laevis* (which is sometimes misspelled as *B. levis*, e.g. Emery  
3862 (1906: 178); Santschi (1923a: 659)), indicated that it is closely related to *B. patagonicus* and to  
3863 the dark forms of *B. heeri*, but he considered *B. laevis* distinct by having a smooth and shiny  
3864 tegument of the head. However a description of the species is not provided, and after examining  
3865 the type material of *B. laevis* we have not identified any consistent morphological differences  
3866 compared to *B. patagonicus* so that we synonymize the species here.

3867 *Brachymyrmex patagonicus* displays variation in color from light to dark brown; *B. patagonicus*  
3868 var. *atrátula* was described by Santschi (1923a) as a variety with darker, almost black tegument,  
3869 and a smooth and shiny body. Evaluating these morphological differences Quirán et al. (2004)

3870 suggested that *B. patagonicus* var. *atratura* is a junior synonym of *B. patagonicus*, and we agree  
3871 with this decision.

3872 Guénard (2018) reports the first record of *B. patagonicus* from continental Asia (Hong Kong),  
3873 however, the specimen illustrated in the paper does not display the diagnostic features of this  
3874 species, but rather those of *B. cordemoyi*. It is noteworthy that *B. patagonicus* is very abundant  
3875 and geographically widespread, but its morphological variation and genetic diversity as well as  
3876 other biological features remain poorly studied. An in-depth study of these features in a  
3877 geographic context would be required to determine if *B. patagonicus* is a distinct species, a  
3878 species complex, or conspecific with some other taxa, e.g. *B. bruchi* and *B. obscurior*.

3879

3880

3881 *Brachymyrmex pictus* Mayr

3882 (Figs. 48, supplementary material Fig. S35)

3883 *Brachymyrmex pictus* Mayr, 1887: 522 (w.q.). **Lectotype worker** (NHMW: ANTWEB

3884 CASENT0915735) and **para lectotype worker** (NHMW; MHNG: USNMENT00758144; **here**  
3885 **designated**): 3 workers [examined]. **BRAZIL: Santa Catharina.**

3886 = *B. heeri* var. *basalis* Wheeler, 1921: 166 (w.). [not examined]. **GUYANA:** Kartabo, Puruni  
3887 trail. **n. syn.**

3888 = *B. pictus* subsp. *balboae* Wheeler, 1942: 253 (w.q.m.). (MCZC: M.C.Z. Cotype 1-3, 4-6, 7-9,  
3889 21438): 2 workers, 8 queens, 2 males [examined]. **PANAMA:** Balboa. **n. syn.**

3890

3891 **Additional material examined. BOLIVIA: Santa Cruz:** 35 km SSE Flor de Oro, -13.83333 -  
3892 60.87763, 450 m, 01 Dec. 1993, P.S. Ward #12232, 3 workers, 1 male (MCZC: CMOS000012,  
3893 CMOS000013). **BRAZIL: Amazonas:** 11 Sep. 1962, W.L. Brown, 3 workers (MCZC:

3894 CMOS000002); Aleixo nr. Manaus, 11 Sep. 1962, W. L. Brown, 8 workers, 2 males, 6 queens  
3895 (MCZC: CMOS000004, CMOS000006, CMOS000009-0000011); Peredão Rd. S. of Manaus, 02  
3896 Sep. 1962; W. L. Brown, 12 workers (MCZC: CMOS000003, CMOS000005, CMOS000007,  
3897 CMOS000008); **Bahia:** Ilheus, 27 Mar. 1997, C.S.F. Mariano, 4 workers (CPDC:  
3898 USNMENT00757794); **Espirito Santo:** Nov. 1977, M. Alvarenga, 4 workers (MZSP:  
3899 USNMENT00757785); **São Paulo:** Caraguatatuba, Reserva Florestal, 40 m, 22 May-01 June  
3900 1962, Exp. Dep. Zool. 2056, 8 workers (MZSP: USNMENT00757676, 00757783); Ilha dos  
3901 Pescadores (Ilha da Vitoria), 24 Mar. 1964, 2 workers, 2 queens (MZSP: USNMENT00757604);  
3902 Ubatuba, Picinguaba, July 2011, 9 workers (ICN: USNMENT00759053). **COLOMBIA: Cauca:**  
3903 Isla Gorgona, 11 Sep. 1989, M. Baena #GQA-05, 3 workers (WEMC: USNMENT00757796,  
3904 00757797); **Magdalena:** 2 km ESE Minca, 11.13 -74.10, 780 m, 13 Aug. 1985, P.S. ward #7895,  
3905 2 workers, 1 queen (PSWC: USNMENT00757792); **Putumayo:** Parque Nacional Natural La  
3906 Paya, Cabaña La Paya, -0.03, -75.20, 330 m, 15-30 Sep. 2002, A. Morales, 1 worker (IAvH).  
3907 **COSTA RICA: Puntarenas:** Parque Natural Corcovado, Sirena, 8.483 -83.583; 23 Apr. 1981, J.  
3908 Longino, 1 worker, 1 queen (JTLC000005913, 000005914); 14 June 1982, J. Longino, 2 workers  
3909 (MCZC: USNMENT00757793); Reserva Biologica Carara, 9.78 -84.60, 30 m, 24 July 1985, P.S.  
3910 Ward #7615, 2 workers, 1 queen (PSWC: USNMENT00757784); Reserva Biologica Carara,  
3911 Estación Quebrada Bonita, 9.78 -84.60, 30 m, 24 July 1985, J. Longino #0562, 1 worker, 1 queen  
3912 (JTLC: JTLC000006051). **ECUADOR: Esmeraldas:** 6.1 km E Rio Verde, 1.07694, -79.41389,  
3913 13 July 2005, W. & E. MacKay #21098, 1 worker (WEMC: USNMENT00757791); **Manabí:** 20  
3914 km NE Chone, 300 m, 1976, S. & J. Peck, 2 workers (MCZC: CMOS000014); **Pichincha:**  
3915 Cotopaxi, 19 km ENE La Maná, -0.88 -79.05, 1100 m, 10 Aug. 1991, P.S. Ward #11418-23, 1  
3916 worker (MCZC: USNMENT00758017). **FRENCH GUIANA:** Saint Elie-K, 4.82261, -53.27649,  
3917 Apr. 2002, J. Orivel & J. Le Breton, 6 workers, 1 queen (CPDC: USNMENT00757786,

3918 00757787). **GUATEMALA: Petén:** Finca Ixobel, 83 km SE Flores, 16.30367, -89.42353, 365  
3919 m, 25 July 2004, W. & E. MacKay #20694, 1 worker, 1 male, 1 queen (WEMC:  
3920 USNMENT00758998); **GUYANA:** Rupununi, Apoteri, 4.08333 -58.58333, 100 m, 12 Jan. 1981,  
3921 J. Longino, 1 worker (JTLC: JTLC000005918). **JAMAICA: Saint Andrew:** Cinchona, 18.067 -  
3922 76.650, 1450 m, 19 Mar. 1984, 1 worker (JTLC: JTLC000005924). **PANAMA:** Gamboa,  
3923 Parque, 9.11722 -79.69972, 24 Apr. 1988, D. Quintero #1, 2 workers (WEMC:  
3924 USNMENT00757795). **PERU: Madre de Dios:** Tambopata, 15 km NE Puerto Maldonado, June  
3925 1989, S.P. Cover & J.E. Tobin, JT 219 CA-740, 2 workers (MCZC: CMOS000032).  
3926 **VENEZUELA: Bolívar:** Canaima, Orchid Is., 14 Oct. 1988, W. MacKay #11165, 8 workers, 3  
3927 males, 1 queen (WEMC: USNMENT0075778800757790, 00757960, 00758997).

3928  
3929 **Diagnosis.** The unique feature for *B. pictus* is a conspicuous color difference between the head  
3930 and thorax, which are yellow and the gaster, which is black, or yellow with (a) black spot(s).

3931  
3932 *Additional material examined measurements* (mm) (n=1). HL<sub>1</sub> 0.43; HL<sub>2</sub> n.a.; HL<sub>3</sub> 0.08; HW  
3933 0.38; SL 0.38; EL 0.11; WL 0.44; PnL n.a.; PnW n.a.; ML 0.09; MW 0.18; *Indices* CI 89.58; SI<sub>1</sub>  
3934 100.00; SI<sub>2</sub> n.a.; OI<sub>1</sub> 26.83; OI<sub>2</sub> 28.57.

3935  
3936 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin flat.  
3937 Dorsum of the head has sparse appressed hairs. Clypeus with a rounded anterior margin and five  
3938 long, erect hairs of which a single, usually conspicuous hair is near the anterior margin, two hairs  
3939 are in mediolateral position and two more near the toruli; other hairs on the clypeus are markedly  
3940 shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique  
3941 anterodorsal view. The scapes surpass the posterior cephalic margin by a length that is smaller

3942 than the maximal diameter of the eye; they have appressed hairs. Three ocelli present. Eyes are  
3943 positioned on the cephalic midline and have 7-10 ommatidia along their maximal diameter.

3944 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum. The  
3945 mesonotum is not inflated and does not bulge dorsally above the pronotum in lateral view. The  
3946 mesometanotal suture is directly visible, however, there is no marked constriction between the  
3947 mesonotum and metanotum, and as a result the metanotal groove is absent. Metathoracic  
3948 spiracles widely separated in dorsolateral position, not protruding, and touching the propodeal  
3949 suture. Dorsum of the propodeum flat and shorter than the propodeal slope. Propodeal spiracles  
3950 circular, situated slightly ventral to the posterior propodeal margin, and slightly posterior of the  
3951 middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole short and inclined  
3952 forward.

3953 **Gaster.** With scattered pubescence and several scattered long erect hairs.

3954 **Color and sculpture.** Body smooth and shiny. The head and thorax are yellowish whereas the  
3955 gaster is either totally black or yellowish with one or more black spots.

3956  
3957 **Distribution (Supplementary material Fig. S35).** *Brachymyrmex pictus* is known to occur in  
3958 Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guatemala, Guyana, Jamaica, Panama, Peru  
3959 and Venezuela.

3960  
3961 **Biology.** The colony of *B. heeri* var. *basalis* reported by Wheeler (1921) and Wheeler (1942) was  
3962 small but nevertheless contained brood. It was found in hollow petioles of a small *Tachigalia*  
3963 *paniculata* tree on the Puruni trail at Kartabi, Guyana. Several colonies of *B. pictus* subsp.  
3964 *balboae* were found to be nesting in hollow twigs of *Tripalis americana* at Balboa, Panama



3965 (Wheeler 1942). No biological information exists on typical *B. pictus*, but it seems this species is  
3966 arboreal.

3967

3968 **Remarks.** One of the specimens determined as a syntype of *B. pictus* (NHMW:  
3969 CASENT0915734) displays the diagnostic features of *B. admotus*. Given that both species were  
3970 described by Mayr (1887) in the same publication from the same type locality, we consider the  
3971 identification of this specimen to be a labeling mistake. Upon describing *B. pictus*, Mayr (1887)  
3972 did not provide any information on the diagnostic features he considered relevant to distinguish  
3973 *B. pictus* from other *Brachymyrmex* species. Wheeler (1921) described *B. heeri* var. *basalis*, but  
3974 he did not provide any morphological description for his typical *B. heeri* specimens nor for the  
3975 variety. The only feature described for *B. heeri* var. *basalis* is its yellow body with a black first  
3976 segment of the gaster (which to our knowledge only fits with the characters of *B. pictus*). Other  
3977 complications are that the material has not been illustrated, that both the typical form and the  
3978 variety were found on the same tree species and the same locality, and that we have not been able  
3979 to locate the material. Later, Wheeler (1942) continued to consider his specimens of *B. heeri* and  
3980 *B. heeri* var. *basalis* to be distinct of *B. pictus*. Given the information available we consider it  
3981 likely that *B. heeri* var. *basalis* belongs to *B. pictus*, but we cannot comment on the taxonomic  
3982 status of his typical *B. heeri* specimens for now.

3983 Fortunately, Wheeler (1942) provided a description of *B. pictus* subsp. *balboae*, which indicates  
3984 that it differs from the typical *B. pictus* only by being smaller. It is indeed somewhat smaller (1.0-  
3985 1.2 mm) than the typical form described by Mayr (1887; 1.3-1.6 mm), however, after studying  
3986 the material we consider this difference to likely represent geographic variation, and we  
3987 synonymize this subspecies here, although a better characterization of the variation in *B. pictus* is  
3988 required.

3989

3990

3991

*Brachymyrmex pilipes* Mayr

3992

(Fig. 49, supplementary material Fig. S36)

3993 *Brachymyrmex pilipes* Mayr, 1887: 524 (q.m.). **Lectotype queen** (NHMW) and **paralectotype**

3994 **queen, male** (NHMW): 2 queens, 1 male [examined]. **BRAZIL: Santa Catharina.** Santschi

3995 (1929: 310) (w.). (NHMB): 2 major workers, 2 minor workers, 1 queen [examined]. **BRAZIL:**

3996 **Parana:** Río Negro. Combination in *Brachymyrmex (Brysha)*: Santschi (1923a: 674). See also:

3997 Ortiz and Fernández (2014: 19, Figs. 7-12).

3998

3999 **Diagnosis.** *Brachymyrmex pilipes* morphologically resembles *B. micromegas* because both

4000 species have a dimorphic worker caste, tumuliform metathoracic spiracles, toruli that touch the

4001 posterior clypeal margin, but never surpass it (best observed in oblique anterodorsal view) and a

4002 clypeus with a row of long thick hairs near the anterior margin. However, *B. pilipes* differs from

4003 *B. micromegas* by the fine, longitudinal striations on most of the mesosoma, and by usually

4004 having a gaster of darker color than the rest of the body.

4005

4006 **Description.** See Ortiz and Fernández (2014).

4007

4008

4009

*Brachymyrmex santschii* Menozzi

4010

(Fig. 50, supplementary material Fig. S37)

4011 *Brachymyrmex santschii* Menozzi, 1927: 338, Fig. 5 (w.). [not examined]. **COSTA RICA: San**

4012 **José.**

4013  
4014 **Additional material examined. COSTA RICA: Cartago:** 2 km N Cervantes, 1600 m, Jan.  
4015 1973, W.L. Brown, 4 workers, 2 putative worker-queen intercastes (MCZC: CMOS000098,  
4016 USNMENT00757750-00757751); **Guanacaste:** Rincon de la Vieja, Las Pailas 7676, 10.77556 -  
4017 85.34528, 1400 m, 18 Feb. 1996, R. Anderson, 2 workers (WEMC: USNMENT00757753,  
4018 00757754); **Puntarenas:** Monteverde, 10.30 -84.83, 1400 m, Apr.-May 1987, S. Little, 1 worker  
4019 (JTLC: JTLC000005243); Monteverde, 10.2964 -84.7831, 1550 m, 18 Jan. 2003, L.A.  
4020 Schonberg, 1 worker (JTLC: JTLC000005055); **San José:** Cerros de Escazu, 2 km S Antonio,  
4021 1650 m, 13 June 1997, R. Anderson #186880C, 1 worker (WEMC: USNMENT00757593).  
4022 **PANAMA: Chiriqui:** Volcan Hartman`s, Finca #17815, 1450 m, 14 June 1996, R.S. Anderson,  
4023 1 worker (WEMC: USNMENT00757752).

4024  
4025 **Diagnosis.** *Brachymyrmex santschii* morphologically resembles *B. iridescens*, because both  
4026 species have the head and the mesosoma with strongly alveolate sculpture. However, they can be  
4027 distinguished from one another because *B. santschii* has a metanotal groove that is wider than the  
4028 diameter of the metathoracic spiracles, scapes that surpass the posterior margin of the head, and a  
4029 gaster with scattered pubescence.

4030  
4031 *Additional material examined measurements* (mm) (n=4). HL<sub>1</sub> 0.40-0.44; HL<sub>2</sub> 0.29-0.30; HL<sub>3</sub>  
4032 0.08-0.10; HW 0.37-0.40; SL 0.39-0.42; EL 0.09-0.11; PnL 0.11-0.13; PnW 0.24-0.28; ML 0.09;  
4033 MW 0.13-0.17; *Indices* CI 89.80-91.82; SI<sub>1</sub> 102.27-109.76; SI<sub>2</sub> 136.36-138.24; OI<sub>1</sub> 24.39-26.84;  
4034 OI<sub>2</sub> 20.00-24.44.

4035

4036 **Worker description. Head.** Longer than wide in full face view; posterior cephalic margin flat or  
4037 slightly concave. Dorsum of the head with subdecumbent hairs. Clypeus with a rounded anterior  
4038 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
4039 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
4040 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
4041 margin in oblique anterodorsal view. The scapes surpass the posterior margin of the head by a  
4042 length approximately equal to the maximal diameter of the eye; they bear appressed hairs. Three  
4043 conspicuous ocelli are present. Eyes are positioned on the cephalic midline and have 8-9  
4044 ommatidia along their maximal diameter.

4045 **Mesosoma.** Dorsum subsinusoidal in lateral view. Without erect hairs, but with decumbent hairs  
4046 on the promesonotum. The mesonotum is variable, typically not or weakly inflated, and not or  
4047 slightly bulging dorsally above the pronotum in lateral view. Metanotal groove wider than the  
4048 diameter of the metathoracic spiracles. Metathoracic spiracles in dorsal position, not protruding  
4049 and not touching any sutures. Dorsum of the propodeum slightly convex and shorter than the  
4050 propodeal slope. Propodeal spiracles circular, positioned on the posterior propodeal margin, at the  
4051 middle of the propodeal slope. Legs with appressed hairs. Petiole short and inclined forward.

4052 **Gaster.** With scattered pubescence and long erect hairs at the edges of the segments.

4053 **Color and sculpture.** Head and dorsum of the mesosoma finely alveolate, those parts that are not  
4054 sculptured, including the gaster, are smooth and shiny. The body is brownish, but sometimes the  
4055 antennae, tarsi, and articulations of the legs are more yellowish.

4056

4057 **Intercaste description.** The morphology of the putative worker-queen intercaste differs from  
4058 that of the worker by its larger body size, the eyes that have around 10 ommatidia along their  
4059 maximal diameter, its strongly expanded mesonotum, the absence of a metanotal groove, the

4060 dorsolateral position of the metathoracic spiracles, the less convex dorsum of the propodeum, and  
4061 a markedly expanded gaster with dense pubescence.

4062

4063 **Distribution (Supplementary material Fig. S37).** *Brachymyrmex santschii* is known from  
4064 Costa Rica and Panama.

4065

4066 **Biology.** Unknown.

4067

4068 **Remarks.** Menozzi (1927) considered *B. santschii* to differ from any other *Brachymyrmex*  
4069 species by its sculpture and pubescence. He described the sculpture as strongly punctuate-  
4070 reticulate, however, following the terminology of Harris (1979) we consider it rather alveolate.  
4071 We do not designate a lectotype here, as we have not studied the type series, which would be  
4072 deposited at the German Entomological Institute in Berlin-Dahlem.

4073

4074

4075 ***Brachymyrmex sosai* NEW SPECIES**

4076 **(Fig. 51, supplementary material Fig. S38)**

4077 **Holotype worker** (UNMSM: USNMENT00757760) and **paratype workers** (USNM:  
4078 USNMENT00759061, 00759062): 3 workers. **PERU: Cusco:** Paucartambo, Kcosñipata, Predio  
4079 Los Wayqechas, ACCA [Asociacion para la Conservacion de la Cuenca Amazonica], -13.17956 -  
4080 71.60556, 2825 m, Andean Forest, J. Sosa-Calvo, JSC040920-04.

4081

4082 **Additional material examined. BOLIVIA: Santa Cruz:** 32.8 km N Comparapa, Kara Huasi, -  
4083 18.05972 -63.91056, 21 Jan. 1999, R. Anderson #18567, 1 worker (WEMC:

4084 USNMENT00759024); **PERU: Lima:** Zárate forest, 2850 m, N. Valencia, I. Frank, 16 workers  
4085 (MCZC: USNMENT00757314-00757320).

4086

4087 **Etymology:** In honor of Dr. Jeffrey Sosa-Calvo, the collector, for his unconditional support and  
4088 friendship.

4089

4090 **Diagnosis.** *Brachymyrmex sosai* n. sp. does not have a specific unique feature but rather a unique  
4091 combination of features that render it distinct: its scapes surpass the posterior margin of the head  
4092 by a length approximately equal to the maximal diameter of the eye, the dorsum of the mesosoma  
4093 does not have conspicuous sculpture, a metanotal groove is present, the metathoracic spiracles are  
4094 in dorsal position, and the dorsal margin of the mesonotum is strongly antero-posteriorly inclined.  
4095 Some features of this species are reminiscent of *B. antennatus*, however, *B. sosai* differs from this  
4096 species in body color, the color of the hairs, the length of the scapes, and in having an antennal  
4097 funiculus with the second segment shorter than the first.

4098

4099 *Holotype measurements* (mm) HL<sub>1</sub> 0.57; HL<sub>2</sub> 0.35; HL<sub>3</sub> 0.16; HW 0.53; SL 0.59; EL 0.14; WL  
4100 0.68; PnL 0.21; PnW 0.33; ML 0.16; MW 0.21; *Indices* CI 93.10; SI<sub>1</sub> 111.11; SI<sub>2</sub> 166.67; OI<sub>1</sub>  
4101 25.93; OI<sub>2</sub> 27.59.

4102

4103 *Paratypes measurements* (n=3) HL<sub>1</sub> 0.60-0.62; HL<sub>2</sub> 0.41-0.43; HL<sub>3</sub> 0.16-0.20; HW 0.57-0.60; SL  
4104 0.59-0.62; EL 0.14-0.16; WL 0.68-0.72; PnL 0.20-0.23; PnW 0.41; ML 0.18-0.20; MW 0.14-  
4105 0.27; *Indices* CI 93.55-96.88; SI<sub>1</sub> 103.23-103.45; SI<sub>2</sub> 142.86-145.45; OI<sub>1</sub> 22.58-27.59; OI<sub>2</sub> 25.81-  
4106 31.25.

4107

4108 **Description. *Head.*** Slightly longer than wide in full face view; posterior cephalic margin flat or  
4109 slightly concave. Dorsum of the head with scattered appressed hairs. Clypeus with a rounded  
4110 anterior margin and five long, erect hairs of which a single, usually conspicuous hair is near the  
4111 anterior margin, two hairs are in mediolateral position and two more near the toruli; other hairs  
4112 on the clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior  
4113 clypeal margin in oblique anterodorsal view. The scapes surpass the posterior margin of the head  
4114 by a length that exceeds the maximal diameter of the eye. Ocelli typically appear to be absent but  
4115 some workers have a central ocellus. Eyes are positioned on the cephalic midline and have 9-10  
4116 ommatidia along their maximal diameter.

4117 ***Mesosoma.*** With several semi-erect hairs on the pronotum and scattered decumbent hairs on the  
4118 promesonotum. The mesonotum is slightly inflated, antero-posteriorly inclined, and it bulges  
4119 dorsally above the pronotum in lateral view. Metanotal groove present and wider than the  
4120 diameter of the metathoracic spiracles. Metathoracic spiracles in fully dorsal position, not  
4121 protruding, and not touching any sutures. Dorsum of the propodeum weakly convex and shorter  
4122 than the propodeal slope. Propodeal spiracles circular, positioned slightly ventral of the posterior  
4123 propodeal margin; they are posterior of the middle of the propodeal slope. Legs with appressed  
4124 hairs. Petiole short and inclined forward.

4125 ***Gaster.*** With dense pubescence and scattered long hairs at the edges of the segments.

4126 ***Color and sculpture.*** Body smooth and shiny, and usually dark brown, but with conspicuously  
4127 lighter hairs. Additionally, the bulbi of the antennae, the terminal funiculus, the tarsi and the  
4128 articulations of the legs are conspicuously yellowish.

4129

4130 **Distribution (Supplementary material Fig. S38).** *Brachymyrmex sosai* is known from Bolivia  
4131 and Peru.

4132

4133 **Biology.** This species was collected from sandy soil, from below a rock.

4134

4135 **Remarks.** The single specimen of *B. sosai* known from Bolivia (WEMC: USNMENT00759024)  
4136 differs in color from the specimens from Peru: its head and thorax are more yellowish than brown  
4137 and the gaster is darker than the rest of the body. We consider it to be part of *B. sosai* as all other  
4138 traits match. Currently *B. sosai* is only known from three localities, and more specimens from  
4139 additional localities will be required to characterize the intraspecific variation in body color.

4140

4141

4142 *Brachymyrmex termitophilus* Forel

4143 (Fig. 52, supplementary material Fig. S39)

4144 *Brachymyrmex heeri* var. *termitophilus* Forel, 1895b: 179 (w.). **Lectotype worker** (MHNG:  
4145 USNMENT00757137) and **paralectotype workers, queen** (MHNG: USNMENT00757136-  
4146 00757138; NHMB: USNMENT00758159; MSNG: USNMENT00757139; **here designated**): 6  
4147 workers, 1 queen [examined]. **BRAZIL: Rio Grande do Sul:** San Leopoldo, col. Wasmann.  
4148 Raised to species: Wild (2007: 44).

4149

4150 **Additional material examined. BRAZIL: São Paulo:** Tapirai, -24.03208 -47.46556, 08-14 Jan.  
4151 2001, R.R. Silva & Everhardt, 2 workers (ICN: MZSP170). **COLOMBIA: Norte de Santander:**  
4152 Parque Nacional Natural Tamá, Vereda El Diamante, Alto Herrera, 7.12278 -72.23472, 1000 m,  
4153 26 Nov. 1999, 1 worker (IAvH: USNMENT00759060). **COSTA RICA: Guanacaste:** Maritza



4154 field Station, 10.95694 -85.49389, 03 May 1995, R. Anderson #17716, 3 workers (WEMC:  
4155 USNMENT00757632); **DOMINICAN REPUBLIC: Pedemales:** Parque Nacional Sierra  
4156 Baoruco, "Las Abejas", 18.15 -71.62, 1320 m, 02 Sep. 2001, A.L. Wild #AW1359, 1 worker, 1  
4157 male, 1 queen (ALWC: USNMENT00757918). **MEXICO: Puebla:** 17 km NE Teztlutlán, 1940  
4158 m, 07 June 1988, W. MacKay #10879, 1 worker, 1 queen (WEMC: USNMENT00758036).  
4159 **PARAGUAY: Itapúa:** San Miguel Potrero, c/Villa Yacyreta, -27.03 -56.20, 17 Jan. 1996, N.E.  
4160 Lopez #ibn 227, 1 worker (ALWC: USNMENT00757662). **UNITED STATES: Texas:** Sabino  
4161 Co., 14.5 km E Nempfil, 11 May 1988, R. Anderson #12763, 1 worker (WEMC:  
4162 USNMENT00758031).

4163  
4164 **Diagnosis.** *Brachymyrmex termitophilus* morphologically resembles *B. aphidicola*, *B. australis*,  
4165 *B. cordemoyi* and *B. obscurior* because these species have scapes that are usually surpassing the  
4166 posterior cephalic margin, their eyes are positioned on the cephalic midline, they have two erect  
4167 hairs on the pronotum and two on the mesonotum, and their mesonotum does not bulge dorsally  
4168 above the pronotum in lateral view. *Brachymyrmex termitophilus* differs from *B. australis* and *B.*  
4169 *aphidicola*, however, by having dense pubescence on the gaster, and from *B. cordemoyi* and *B.*  
4170 *obscurior* by having a yellowish body instead of brownish. *Brachymyrmex termitophilus* also  
4171 resembles *B. bahamensis* somewhat, but *B. termitophilus* typically bears two erect hairs on the  
4172 pronotum whereas *B. bahamensis* approximately six that are moreover much longer.

4173  
4174 *Lectotype and paralectotype measurements* (mm) (n=2). HL<sub>1</sub> 0.45; HL<sub>2</sub> 0.29-0.31; HL<sub>3</sub> 0.10-  
4175 0.12; HW 0.39; SL 0.41-0.43; EL 0.10; Pnl 0.12; PnW 0.25-0.27; ML 0.08; MW 0.18; *Indices* CI  
4176 89.96; SI<sub>1</sub> 105.00-110.00; SI<sub>2</sub> 131.25-146.67; OI<sub>1</sub> 25.00; OI<sub>2</sub> 21.74-26.09.

4177

4178 *Additional material examined measurements* (mm) (n=2). HL<sub>1</sub> 0.43-0.44; HL<sub>2</sub> 0.30; HL<sub>3</sub> 0.10;  
4179 HW 0.38-0.39; SL 0.33-0.40; EL 0.10; WL 0.42; PnL 0.12; PnW 0.25-0.27; ML 0.08-0.09; MW  
4180 0.16-0.18; *Indices* CI 89.80-90.00; SI<sub>1</sub> 85.23-101.11; SI<sub>2</sub> 110.29-133.82; OI<sub>1</sub> 25.56-26.14; OI<sub>2</sub>  
4181 22.00-22.45.

4182  
4183 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin flat.  
4184 Dorsum of the head with sparse appressed hairs. Clypeus with a rounded anterior margin and five  
4185 long, erect hairs of which a single, usually conspicuous hair is near the anterior margin, two hairs  
4186 are in mediolateral position and two more near the toruli; other hairs on the clypeus are markedly  
4187 shorter and appressed or decumbent. Toruli surpassing the posterior clypeal margin in oblique  
4188 anterodorsal view. The scapes surpass the posterior margin of the head by a length equal to the  
4189 maximal diameter of the eye or less; they have appressed hairs. Ocelli inconspicuous. Eyes are  
4190 positioned on the cephalic midline and have 7-9 ommatidia along their maximal diameter.

4191 **Mesosoma.** Typically with two erect hairs on the pronotum and two on the mesonotum. The  
4192 mesonotum is not inflated and does not bulge dorsally above the pronotum in lateral view.  
4193 Metanotal groove absent or narrower than the diameter of the metathoracic spiracles.  
4194 Metathoracic spiracles in dorsolateral position, not protruding, and usually touching the  
4195 propodeal sutures. Dorsum of the propodeum is convex and shorter than the propodeal slope.  
4196 Propodeal spiracles circular, positioned slightly ventral of the posterior propodeal margin, at the  
4197 middle of the propodeal slope. Legs with appressed and scattered hairs. Petiole short and inclined  
4198 forward.

4199 **Gaster.** Usually with dense pubescence, and scattered long erect hairs, among others at the edges  
4200 of the segments.

4201 **Color and sculpture.** Body smooth, shiny and yellowish.

4202

4203 **Distribution (Supplementary material Fig. S39).** *Brachymyrmex termitophilus* is known from  
4204 Brazil, Colombia, Costa Rica, the Dominican Republic, Mexico, Paraguay and the United States.  
4205

4206 **Biology.** Forel (1895b) indicated that this species was collected in association with termites.  
4207

4208 **Remarks.** The type material of *B. termitophilus* at the MHNG is somewhat problematic and may  
4209 have caused confusion as to the diagnostic traits of the species (see below). This material consists  
4210 of specimens mounted on three pins of which one (USNMENT00757136) holds an undescribed  
4211 queen; the second (USNMENT00757137) holds a brownish worker with dense pubescence on  
4212 the gaster, which is here designated as lectotype, and the gaster of another worker of which the  
4213 rest of the body is missing; the third pin (USNMENT00757138) holds two workers with  
4214 yellowish heads and mesosoma, and a darker gaster which bears scarce pubescence.

4215 Originally, Forel (1895b) described *B. termitophilus* as a variety of *B. heeri* that is slightly  
4216 smaller than the typical form, that has longer scapes, and a somewhat sparser pubescence but  
4217 denser, thicker erect hairs, mainly on the gaster. Wild (2007) subsequently elevated *B.*  
4218 *termitophilus* to species level reporting two differences with *B. heeri*, i.e. the length of the scapes  
4219 and the lateral morphology of the mesosoma. The first trait is suspect, however, as he reports the  
4220 scapes of *B. termitophilus* to barely reach the posterior margin of the head, which contrasts  
4221 strongly with the original statement by Forel (1895b). We believe that this error caused Wild  
4222 (2007) to suggest that *B. termitophilus* may be conspecific with *B. fiebrigi*. The latter species has  
4223 indeed scapes that do not reach the posterior margin of the head. Rather than *B. fiebrigi*, *B.*  
4224 *termitophilus* resembles the species here mentioned in the diagnosis. The status of *B.*  
4225 *termitophilus* is unclear: Several of the specimens in the type series of *B. termitophilus*

4226 morphologically resemble *B. australis* and *B. aphidicola* in having scarce pubescence on the  
4227 gaster. Additionally, Santschi (1923a) mentioned that *B. termitophilus* and *B. australis* are both  
4228 found in association with termites. As mentioned in the diagnosis also the differences with *B.*  
4229 *cordemoyi* and *B. obscurior* are limited, and *B. termitophilus* may be conspecific with one or  
4230 several of these four species mentioned in the diagnosis. We tentatively preserve the current  
4231 status of *B. termitophilus* awaiting more material and study.

4232

4233

4234 *Brachymyrmex tristis* Mayr

4235 (Figs. 53, supplementary material Fig. S40)

4236

4237 *Brachymyrmex tristis* Mayr, 1870: 389 (w.). **Lectotype worker** (NHMW: ANTWEB  
4238 CASENT0915737; **here designated**): 1 worker [examined]. **COLOMBIA**: Santafé de Bogotá.  
4239 See also: Santschi (1923a: 673).

4240

4241 **Additional material examined. COLOMBIA: Boyacá:** Chinquinquirá, 07 Dec. 1975, W. & E.  
4242 MacKay #572, 2 workers (WEMC: USNMENT00757574); **Cundinamarca:** Mosquera to La  
4243 Mesa, km 8, >2600 m, arid slope, under rock, 30 June 1976, W.L. & D.E. Brown, 18 workers, 2  
4244 queens, 3 males (MCZC: USNMENT00757280-00757282, 00757306-00757311).

4245

4246 **Diagnosis.** *Brachymyrmex tristis* morphologically resembles *B. degener* and *B. coactus*, because  
4247 all three species have scapes that surpass the posterior margin of the head, they have faint  
4248 sculpture on the mesosoma, their mesonotum bulges dorsally above the pronotum in lateral view,  
4249 their metanotal groove is wider than the diameter of the metathoracic spiracles, the metathoracic

4250 spiracles slightly protrude, and the gaster has sparse pubescence. *Brachymyrmex tristis* differs  
4251 from *B. coactus* by having a uniform body color and dense decumbent hairs on the head. It differs  
4252 from *B. degener* by having many decumbent hairs on the head.

4253

4254 *Lectotype worker measurements* (mm). HL<sub>1</sub> 0.61; HL<sub>2</sub> 0.38; HL<sub>3</sub> 0.15; HW 0.56; SL 0.61; EL  
4255 0.15; WL 0.70; PnL 0.17; PnW 0.39; ML 0.17; MW 0.30; *Indices* CI 92.50; SI<sub>1</sub>108.11; SI<sub>2</sub>  
4256 160.00; OI<sub>1</sub> 27.03; OI<sub>2</sub> 25.00.

4257

4258 **Description. Head.** Slightly longer than wide in full face view; posterior cephalic margin slightly  
4259 concave. Dorsum of the head with dense decumbent pubescence. Clypeus with a rounded anterior  
4260 margin and five long, erect hairs of which a single, usually conspicuous hair is near the anterior  
4261 margin, two hairs are in mediolateral position and two more near the toruli; other hairs on the  
4262 clypeus are markedly shorter and appressed or decumbent. Toruli surpassing the posterior clypeal  
4263 margin in oblique anterodorsal view. The scapes surpass the posterior margin of the head by a  
4264 length up to 1.5× the maximal diameter of the eye; they have appressed hairs. Three ocelli are  
4265 present. Eyes are positioned on the cephalic midline and have 10-12 ommatidia along their  
4266 maximal diameter.

4267 **Mesosoma.** With two erect hairs on the pronotum and usually also two on the mesonotum. The  
4268 mesonotum is slightly inflated and bulges dorsally above the pronotum in lateral view. Metanotal  
4269 groove wider than the diameter of the metathoracic spiracles. Metathoracic spiracles in  
4270 dorsolateral position, slightly protruding, and not touching any sutures. Dorsum of the  
4271 propodeum slightly convex and shorter than propodeal slope. Propodeal spiracles circular,  
4272 positioned ventral of the posterior propodeal margin, at the middle of the propodeal slope. Legs  
4273 with appressed hairs. Petiole short and inclined forward.

4274 **Gaster.** With scattered pubescence and several scattered long erect hairs.

4275 **Color and sculpture.** Head and gaster are smooth and shiny, whereas the mesosoma has faint  
4276 sculpture; body uniformly dark brown apart from the tarsi of the legs which are lighter in color.

4277

4278 **Distribution (supplementary material Fig. S40).** *Brachymyrmex tristis* is only known from  
4279 Colombia.

4280

4281 **Biology.** Unknown.

4282

4283 **Remarks.** *Brachymyrmex tristis* is as mentioned morphologically very similar to *B. degener* and  
4284 *B. coactus* and further studies are required to assess whether these species are distinct or  
4285 conspecific. For now, we follow previous authors in maintaining them as separate species.  
4286 Forel (1899) initially considered *B. musculus* to be a race of *B. tristis*, however, we agree with his  
4287 later decision to consider *B. musculus* distinct (Forel 1901a). Santschi (1923a) also considers *B.*  
4288 *musculus* to be closely related to *B. tristis*, however, the first species has a mesonotum that does  
4289 not bulge dorsally above the pronotum in lateral view, and its metathoracic spiracles are not  
4290 protruding.

4291

4292 **Additional taxonomic remarks.**

4293 We could not include information on *B. longicornis* var. *pullus* Santschi, 1933 in the above  
4294 because the type series consists of a single, destroyed worker (NHMB). As such, we could only  
4295 re-examine the morphological descriptions of Santschi (1933), from which we conclude that *B.*  
4296 *longicornis* var. *pullus* seems to be morphologically similar to *B. patagonicus* and *B. bruchi*.  
4297 *Brachymyrmex longicornis* var. *pullus* has a shiny body that is black or dark brownish, the scapes

4298 surpass the posterior margin of the head, it has large eyes that occupy a third of the sides of the  
4299 head, and the thorax would have been similar to that of *B. longicornis* which indicates that the  
4300 mesonotum did not bulge dorsally above the pronotum in lateral view. Santschi (1933) did not  
4301 describe the pubescence on the gaster, and so in the above we consider the pubescence to be  
4302 similar to that of typical *B. longicornis* (here synonymized to *B. australis*), *B. longicornis* var.  
4303 *hemiops* (here synonymized to *B. aphidicola*) and *B. longicornis* var. *immunis* (here synonymized  
4304 to *B. admotus*). If this assumption were not true *B. longicornis* var. *pullus* would resemble *B.*  
4305 *cordemoyi* and *B. obscurior* more than *B. bruchi* and *B. patagonicus*.

4306

#### 4307 **Morphometric measurements.**

4308 Although the abovementioned identifications mention morphometric measurements, indices and  
4309 count data, our identification system and the key were constructed before analysis of quantitative  
4310 data, and thus somewhat independent from the quantitative comparison that follows. Here, we  
4311 examine how well measurements corroborate the established identification system.

4312 The morphospace occupation of the various species is displayed in Fig. 54. The stress associated  
4313 with nmMDS is small (5.70), indicating that this ordination is reliable and the risk of drawing  
4314 false inferences very limited. The contribution of the individual morphometric variables  
4315 (measurements, indices and counts) to the morphospace is indicated with a biplot. Permutation  
4316 tests revealed that all variables contributed significantly to the morphospace occupation of taxa,  
4317 apart from  $OI_1$ , and therefore we excluded this index from further statistical tests. Overall, many  
4318 of the 38 species included occupy very similar regions of the morphospace, which testifies to the  
4319 cryptic nature of morphological differences among these taxa, and therefore to the legacy of  
4320 difficulties with species identifications that have plagued workers of *Brachymyrmex* (see Wheeler

4321 1903; Santschi 1923a; Creighton 1950; Kusnezov 1959; Wilson and Taylor 1967). Indeed, the  
4322 observation that several species overlap in the central region of the morphospace suggests that the  
4323 genus is overall characterized by a large degree of morphological conservatism. However, upon  
4324 more detailed examination we also perceive that most taxa occupy rather restricted regions of the  
4325 morphospace, despite measurements typically deriving from specimens obtained from distant  
4326 localities. For example, even widespread species, such as *B. patagonicus* occupy a rather  
4327 restricted region of the morphospace. One notable exception is *B. bruchi*, which displays large  
4328 variation on both nmMDS1 and 2, and which is difficult to characterize morphologically  
4329 (although molecular analyses suggest our identification system works well for this species too  
4330 [see below]). Beyond the measured traits, the first nmMDS axis also reflects general differences  
4331 in body size, with small species (*B. depilis*, *donisthorpei*, *fiebrigi*, *iridescens*, and *feitosai*)  
4332 plotting along the most negative and large species (*B. admotus*, *cavernicola*, and *degener*) along  
4333 the most positive values. The overall restricted morphospace occupation of individual species  
4334 testifies to the possibility to distinguish many species in one-on-one comparisons, and we  
4335 examine this issue for univariate variables, because such univariate comparisons may be more  
4336 helpful than multivariate comparisons for colleagues aiming to identify specimens directly in the  
4337 field.

4338  
4339 Boxplots (Figs. 55, 56) highlight similarities and differences among 20 *Brachymyrmex* species  
4340 for the 16 univariate variable (after exclusion of  $OI_1$ ) with statistical pairwise comparisons. Here  
4341 we will not exhaustively compare all these species for each of the variables, as this would lead to  
4342 3040 pairwise comparisons. Rather, we will focus on comparing five species pairs that are  
4343 difficult to distinguish, i.e. *aphidicola-australis*, *bruchi-patagonicus*, *coactus-degener*,  
4344 *cordemoyi-obscurior* and *donisthorpei-modestus*, with the aim to find additional criteria that may



4345 allow differentiating these taxa. *Brachymyrmex aphidicola-australis* differ significantly in SI<sub>1</sub>  
4346 and SI<sub>2</sub>, but not in other variables. *Brachymyrmex bruchi-patagonicus* display significant  
4347 differences in HL<sub>1</sub>, HL<sub>2</sub>, HW, SL, EL, WL, PnL, and PnW suggesting that the main difficulty  
4348 differentiating these taxa relates to the very variable nature of *B. bruchi*, as already highlighted  
4349 above. *Brachymyrmex coactus* and *B. degener* are effectively very difficult to distinguish as the  
4350 only significant difference we found is in OI<sub>2</sub>, which was admittedly very variable for *B. degener*.  
4351 Although *B. cordemoyi* and *B. obscurior* overlapped strongly in morphospace occupation, they  
4352 can nevertheless be distinguished based on HL<sub>1</sub>, HL<sub>2</sub>, HL<sub>3</sub>, HW, EL, PnL, PnW, MI, MW, SI<sub>1</sub>,  
4353 and the number of ommatidia. For several of these variables, *B. obscurior* showed limited  
4354 variation, despite the inclusion of 10 specimens from 4 different countries, which may have  
4355 driven statistical significance. Finally, *B. donisthorpei* and *B. modestus* differed significantly in  
4356 HL<sub>1</sub>, HL<sub>3</sub>, HW, SL, WL, MW, CI, and OI<sub>2</sub>. In summary, the morphometric variables confirm  
4357 significant morphological differences for all five species pairs. Interestingly, as we will document  
4358 in the next section, the two species pairs with the most limited number of differences, i.e.  
4359 *aphidicola-australis* and especially *coactus-degener* are phylogenetically closely related (see  
4360 below).

4361

### 4362 **Phylogenetic inference.**

4363 Our phylogenetic analyses with maximum parsimony (MP), maximum likelihood (ML) and  
4364 Bayesian Inference (BI) on five gene fragments (three nuclear, two mitochondrial) for 19  
4365 *Brachymyrmex* species, 5 species of its sister clade *Myrmelachista* and outgroups retrieved  
4366 *Brachymyrmex* and *Myrmelachista* as a well-supported monophyletic clade (MP = 98, ML = 100,  
4367 BPP=1.00; Fig.57). This finding agrees with recent studies of the deep phylogenetic relationships  
4368 within the subfamily Formicinae based on UCEs (ultraconserved elements: Blaimer et al. 2015;

4369 Ward et al. 2016), however, this previous study contained only two *Brachymyrmex* and one  
4370 *Myrmelachista* species. Under expanded taxon sampling *Brachymyrmex* and *Myrmelachista* were  
4371 found to be reciprocally monophyletic, with high support for each genus (MP = 98, ML = 100,  
4372 BPP = 1.00 and MP = 96, ML = 82, BPP=1.00, respectively). This finding suggests that the  
4373 morphological criteria currently used to delimit these genera are unambiguous autapomorphies.  
4374 Examining *Brachymyrmex* in more detail, many of the nodes of intermediate depth are poorly  
4375 supported, indicating that more markers are required, or at least more complete sampling of  
4376 markers across taxa, to reveal the phylogenetic relationships between individual *Brachymyrmex*  
4377 species. Given that our analysis includes half of the currently recognized *Brachymyrmex* species,  
4378 increased taxon sampling may also help to resolve phylogenetic relationships among the species.  
4379  
4380 Despite ambiguity as to interspecific relationships, species-level nodes (indicated in bold in Fig.  
4381 57) are overall well-supported, and of the 14 *Brachymyrmex* species that were sampled with 2-15  
4382 specimens, 13 proved to be monophyletic. This finding largely confirms our assessment of  
4383 intraspecific and interspecific components of morphological variation, the phylogenetic value of  
4384 the morphological traits used, and thus the significance of our proposed morphological system of  
4385 species delimitations. The only species that was not retrieved as monophyletic is *B. coactus*,  
4386 which included the monophyletic *B. degener*. *Brachymyrmex coactus* and *B. degener* are  
4387 morphologically very similar (as indicated already above in the taxonomic treatment and  
4388 morphometrics), and they mainly differ in body color, which may be a trait with large  
4389 intraspecific variation. Significant differences between both species were also found in OI<sub>2</sub>,  
4390 although *B. degener* is very variable as to this index. On the other hand, the genetic  
4391 differentiation between *B. coactus* and *B. degener* is substantial, as indicated by the branch  
4392 lengths in Fig. 57, suggesting that both may be part of a larger clade with cryptic diversity, and

4393 that the observed bimodal distribution in body color may hint at interspecific differences.  
4394 Considering the substantial genetic differentiation, we do not synonymize both species but rather  
4395 postpone our assessment until more specimens become available, especially of *B. coactus* from  
4396 Brazil. Another possible indication of cryptic species diversity relates to *B. cavernicola*, which  
4397 contains two well-supported subclades, one with specimens from Central America, and the other  
4398 clade with specimens from South America. More in-depth studies are required to test whether this  
4399 split relates to different species, or rather variation between geographically-separated  
4400 populations. The hypothesis of cryptic diversity is furthermore supported by the ABGD analysis  
4401 (see below).

4402 For *B. feitosai*, *B. nebulosus*, *B. pilipes*, *B. brasiliensis* and *B. bicolor* **n. sp.** only a single  
4403 individual per species was included in the phylogenetic analyses so that it is difficult to make  
4404 conclusions on the integrity of these species, however, all these species are deeply split from  
4405 other *Brachymyrmex* species, suggesting that they are truly distinct. It was particularly important  
4406 for us to include *B. pilipes*, because this species has very distinctive and different morphological  
4407 traits (see species description) in comparison to most other *Brachymyrmex* species. Interestingly,  
4408 the species seems to have a basal position in our phylogeny and revealing its position in the  
4409 future may yield important insights into trait evolution within the genus. Nevertheless, our  
4410 phylogenetic analysis confirms that *B. pilipes* is part of *Brachymyrmex*, rather than an  
4411 independent lineage.

4412

#### 4413 **Automated species delimitation.**

4414 A total of 24 hypothetical species entities were retrieved within *Brachymyrmex* upon analyzing  
4415 the barcoding fragment of COI (658 basepairs) with ABGD. Overall these entities are in good  
4416 agreement with the morphologically-recognized species indicated in Fig. 57, and the differences

4417 are limited to the potential further subdivision of morphologically recognized species by ABGD.  
4418 Within *B. heeri* four groups are recognized, with each group containing the specimens from one  
4419 country in the phylogeny. *Brachymyrmex antennatus* is subdivided in two groups, with one group  
4420 consisting of specimen CX81 from Peru and the other contains the additional specimens.  
4421 *Brachymyrmex cavernicola* was subdivided in two groups also, along the main subdivision  
4422 observed in Fig. 57 and discussed above, indicating that differentiation and perhaps cryptic  
4423 speciation is taking place along a geographic gradient. The final difference relates to the *B.*  
4424 *coactus/degener* clade. The ABGD analysis recognized three groups: one group containing only  
4425 specimen CX37 from Brazil, the remainder of *B. coactus* as a second group, and the third group  
4426 contains the specimens identified as *B. degener*. In summary, the ABGD analysis corroborates  
4427 our morphological classification system. It suggests that this classification is conservative and  
4428 that more cryptic diversity may exist within *Brachymyrmex*.

4429

## 4430 **CONCLUSIONS**

4431 For over a century the ant genus *Brachymyrmex* has been in dire need of revision, and here we  
4432 present such a revision based on the morphology of workers, validated with morphometric and  
4433 molecular data. Currently the strategy to focus on workers is the most effective solution to revise  
4434 the genus, because other castes (queens and males) are poorly known for most *Brachymyrmex*  
4435 species. We studied 1303 samples that have been assigned to 40 species based on the established  
4436 morphological identification system as represented in a dichotomous identification key, which we  
4437 tested with previous and new material. Additionally, this key was tested independently by several  
4438 colleagues (Fabiana Cuezco (Argentina), John Lattke and Livia Pires do Prado (Brazil)), and the  
4439 obtained identifications were in good agreement with our own diagnoses, suggesting that it  
4440 effectively allows discerning interspecific differences from intraspecific variation. Beyond these

4441 qualitative tests we also complemented our identifications with measurements to reconstruct the  
4442 distribution of species in morphospace, and we statistically analyzed individual measurements as  
4443 univariate variables. These efforts suggest that even species pairs that are qualitatively difficult to  
4444 discern can be separated statistically, and they illustrated that taxonomically problematic cases  
4445 relate to taxa that have high intraspecific trait variance. The species pair that was most difficult to  
4446 discern based on measurements proved to be *B. coactus* and *B. degener*, and interestingly, these  
4447 taxa represent the only disagreement between our morphological identification system and our  
4448 phylogenetic analysis based on five gene fragments. *Brachymyrmex degener* was nested within *B.*  
4449 *coactus* but considering the deep phylogenetic splits in the *coactus-degener* clade and the results  
4450 from automated species delimitation, we await more material to resolve the status of the  
4451 morphospecies in this clade. In summary, 13 of the 14 morphologically-identified species that  
4452 were included in molecular work with 2-15 individuals each were recovered as monophyletic,  
4453 indicating the overall robustness of our proposed morphological identification system, and by  
4454 extension our taxonomic revision. Finally, we have reported dimorphic workers for some  
4455 *Brachymyrmex* species and the existence of a putative worker-queen intercaste in others. As such,  
4456 the genus altogether may represent a promising system to study caste evolution in ants.

4457

#### 4458 **ACKNOWLEDGMENTS**

4459 We thank Alex Wild (ALWC), Brian Fisher (CASC), Jacques Delabie (CPDC), Claudia Medina  
4460 (IAvH), John Longino (JTLC), Carlos R. F. Brandão, Rodrigo Feitosa, Flávia Esteves (MZSP),  
4461 Stefan Cover (MCZC), Heraldo Vasconcelos, Renata Pacheco, Gabriela Camacho (UFUC),  
4462 Maria Tavano and Roberto Poggi (MCSN), Daniel Burckhardt (NHMB), Dominique  
4463 Zimmermann (NHMW), Frank Koch (MfNB), Fabiana Cuzzo (INSUE), Priscila Hanisch  
4464 (MACN), Phil Ward (PSWC), Maurice Leponce and Thibaut Delsinne (RBINS), William and

4465 Emma MacKay (WEMC) for providing acces to collections and/or for the loan of critical  
4466 material. David Donoso and John Longino kindly provided sequencing data. Ted Schultz, Jeffrey  
4467 Sosa-Calvo, Eugenia Okonski (USNM) and Carlos Sarmiento (ICN) provided continuous support  
4468 in the development of this paper. John Lattke, Fabiana Cuezso, Lívia Pires do Prado and an  
4469 anonymous referee tested the identification key and providing invaluable comments and  
4470 suggestions. Comments of John Longino, an anonymous referee and the editors have strongly  
4471 improved this manuscript.

4472

#### 4473 **FUNDING INFORMATION**

4474 We are grateful for funding from the Division de Investigación de Bogotá (DIB), Facultad de  
4475 Ciencias, Universidad Nacional de Colombia and the Colciencias program “Proyectos de  
4476 Investigación Científica y Tecnológica” of 2010 (110152128319 CT 413–2011) (to CMOS and  
4477 FF), for an Ernst Mayr Grant in 2011 (MCZ Harvard University), a grant from the Colciencias  
4478 program “Jóvenes Investigadores e Innovadores – Virginia Gutiérrez de Pineda” in 2010–2011, a  
4479 grant of the Smithsonian National Museum of Natural History installed by Cristian Samper (to  
4480 CMOS), CPER CLIMIBIO funded by the Région Hauts-de-France, the French Ministry of  
4481 Higher Education and Research, and the European Fund for Regional Economic Development (to  
4482 CMOS and BVB), the Institut de Recherches Pluridisciplinaires en Sciences de l’Environnement  
4483 (IREPSE; to BVB) and ANR-JCJC-EVOLINK of the French Agence Nationale de la Recherche  
4484 (to BVB).

4485

#### 4486 **CONFLICT OF INTEREST**

4487 The authors declare that they have no conflict of interest.

4488

4489 **DATA AVAILABILITY**

4490 Sequence data is deposited in NCBI Genbank [upon acceptance of the manuscript], and Genbank  
4491 accession numbers are indicated in Supplementary Table S1. Measurement data are provided as an  
4492 online supplementary file.

4493

4494 **REFERENCES**

4495 Agosti, D. (1991). Revision of the oriental ant genus *Cladomyrma*, with an outline of the higher  
4496 classification of the Formicinae (Hymenoptera: Formicidae). *Systematic Entomology*, *16*,  
4497 293-310.

4498 Alayo, D. P. (1974). Introducción al estudio de los Himenopteros de Cuba. Superfamilia  
4499 Formicoidea. *Academia de Ciencias de Cuba*, *53*, 1-58.

4500 Amante, C., & Eakins, B. W. (2009). ETOPO1 1 Arc-Minute Global Relief Model: Procedures,  
4501 Data Sources and Analysis. *NOAA Technical Memorandum NESDIS NGDC-24*.

4502 Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and  
4503 powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series*  
4504 *B(57)*, 289-230.

4505 Blaimer, B. B., Brady, S. G., Schultz, T. R., Lloyd, M. W., & Ward, P. S. (2015). Phylogenomic  
4506 methods outperform traditional multi-locus approaches in resolving deep evolutionary  
4507 history: a case study of formicine ants. *BMC Evolutionary Biology*, *15*, 1-14,  
4508 doi:10.1186/s12862-015-0552-5.

4509 Bolton, B. (1994). *Identification guide to the ant genera of the world*. Cambridge, Massachusetts:  
4510 Harvard University Press.

4511 Bolton, B. (1995). *A new general catalogue of the ants of the world*. Cambridge, Massachusetts:  
4512 Harvard University Press.

- 4513 Bolton, B. (2003). Synopsis and classification of Formicidae. *Memoirs of the American*  
4514 *Entomological Institute*, 71, 1-370.
- 4515 Bolton, B. (2007). How to conduct large-scale taxonomic revisions in Formicidae. *Memoirs of*  
4516 *the American Entomological Institute*, 80, 52-71.
- 4517 Bolton, B. (2018). An online catalog of ants of the world. <http://www.antcat.org/>. Accessed  
4518 December 15 2016.
- 4519 Brady, S. G., Schultz, T. R., Fisher, B. L., & Ward, P. S. (2006). Evaluating alternative  
4520 hypotheses for the early evolution and diversification of ants. *Proceedings of the National*  
4521 *Academy of Sciences of the United States of America*, 103, 18172-18177.
- 4522 Brandão, C. R. F. (1991). Adendos ao catálogo abreviado das formigas da região Neotropical  
4523 (Hymenoptera: Formicidae). *Revista Brasileira de Entomologia*, 35, 319-412.
- 4524 Brownrigg, R. (2015). Mapdata: Extra Map Databases (original S code by Becker R. A. and  
4525 Wilks A. R). R package, Version 2.2-5.
- 4526 Brownrigg, R., Minka, T. P., & Deckmyn, A. (2015). Maps: draw geographical maps (original S  
4527 code by Becker R. A. and Wilks A. R.). R package, Version 3.0.1.
- 4528 Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community structure.  
4529 *Australian Journal of Ecology*, 18, 117-143.
- 4530 Cole, A. C. J. (1953). Studies of New Mexico ants. V. The genus *Pheidole* with synonymy  
4531 (Hymenoptera: Formicidae). *Journal of the Tennessee Academy of Science*, 28, 297-299.
- 4532 Creighton, W. S. (1950). The ants of North America. *Bulletin of the Museum of Comparative*  
4533 *Zoology*, 104, 1-585.
- 4534 Da Silva, R., Peloso, P. L. V., Sturaro, M. J., Veneza, I., Sampaio, I., Schneider, H., et al. (2018).  
4535 Comparative analyses of species delimitation methods with molecular data in snappers  
4536 (Perciformes: Lutjaninae) *Mitochondrial DNA Part A*, 29, 1108-1114.



- 4537 De Zolessi, L. C., Abenante, Y. P., & González, L. A. (1978). Descripción y observaciones  
4538 bioetológicas sobre una nueva especie de *Brachymyrmex* (Hymenoptera: Formicidae).  
4539 *Revista de Biología del Uruguay*, 4, 21-44.
- 4540 Dejean, A., Fisher, B. L., Corbara, B., Rarevohitra, R., Randrianaivo, R., Rajemison, B., et al.  
4541 (2010). Spatial distribution of dominant arboreal ants in a Malagasy coastal rainforest:  
4542 Gaps and presence of an invasive species. *PLoS ONE*, 5(2), e9319, doi:10.1371/journal.  
4543 pone.0009319.
- 4544 Dinno, A. (2017). dunn.test: Dunn's test of multiple comparisons using rank sums. R package  
4545 version 1.3.4.
- 4546 Emery, C. (1893). Beiträge zur Kenntniss der nordamerikanischen Ameisenfauna. Zoologische  
4547 Jahrbücher. *Abteilung für Systematik, Geographie und Biologie der Tiere*, 7, 633-682.
- 4548 Emery, C. (1895). Note sur les fourmis du Chili avec descriptions de deux espèces nouvelles.  
4549 *Actes de la Société Scientifique du Chili*, 4, 213-216.
- 4550 Emery, C. (1906). Studi sulle formiche della fauna Neotropica. XXVI. *Bullettino della Società*  
4551 *Entomologica Italiana*, 37, 107-194.
- 4552 Emery, C. (1925). Hymenoptera. Fam. Formicidae. Subfam. Formicinae. *Genera Insectorum*,  
4553 183, 1-302.
- 4554 Fisher, B. L., & Cover, S. P. (2007). *Ants of North America. A guide to genera*: Berkeley:  
4555 University of California Press.
- 4556 Forel, A. (1874). Les fourmis de la Suisse. Systématique, notices anatomiques et physiologiques,  
4557 architecture, distribution géographique, nouvelles expériences et observations de moeurs.  
4558 *Neue Denkschriften der Allgemeinen Schweizerischen Gesellschaft für die Gesamten*  
4559 *Naturwissenschaften*, 26, 1-452.

- 4560 Forel, A. (1876). Études myrmécologiques en 1875 avec remarques sur un point de l'anatomie  
4561 des coccides. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 14, 33-62.
- 4562 Forel, A. (1893). Formicides de l'Antille St. Vincent, récoltées par Mons. H. H. Smith.  
4563 *Transactions of the Entomological Society of London*, 333-418.
- 4564 Forel, A. (1895a). Nouvelles fourmis de diverses provenances, surtout d'Australie. *Annales de la*  
4565 *Société Entomologique de Belgique*, 39, 41-49.
- 4566 Forel, A. (1895b). Die Ameisen- und Termitengäste von Brasilien. Anhang. Beschreibung einiger  
4567 neuer brasilianischer Ameisenarten. . *Verhandlungen der Kaiserlich-Königlichen*  
4568 *Zoologisch-Botanischen Gesellschaft in Wien*, 45, 178-179.
- 4569 Forel, A. (1897). Quelques Formicides de l'Antille de Grenada récoltés par M. H. H. Smith.  
4570 *Transactions of the Entomological Society of London*, 297-300.
- 4571 Forel, A. (1899). Formicidae. [part]. *Biologia Centrali-Americana Hymenoptera*, 3, 105-136.
- 4572 Forel, A. (1901a). I. Fourmis mexicaines récoltées par M. le professeur W.-M. Wheeler. II. A  
4573 propos de la classification des fourmis. *Annales de la Société Entomologique de Belgique*,  
4574 45, 123-141.
- 4575 Forel, A. (1901b). Einige neue Ameisen aus Südbrasilien, Java, Natal und Mossamedes.  
4576 *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 10, 297-311.
- 4577 Forel, A. (1902). Fourmis nouvelles d'Australie. . *Revue Suisse de Zoologie*, 10, 405-548.
- 4578 Forel, A. (1907). Formiciden aus dem Naturhistorischen Museum in Hamburg. II. Teil.  
4579 Neueingänge seit 1900. *Mitteilungen aus dem Naturhistorischen Museum in Hamburg*,  
4580 24, 1-20.
- 4581 Forel, A. (1908). Ameisen aus Sao Paulo (Brasilien), Paraguay etc. gesammelt von Prof. Herm. v.  
4582 Ihering, Dr. Lutz, Dr. Fiebrig, etc. *Verhandlungen der Kaiserlich-Königlichen*  
4583 *Zoologisch-Botanischen Gesellschaft in Wien*, 58, 340-418.

- 4584 Forel, A. (1909). Ameisen aus Guatemala usw., Paraguay und Argentinien (Hym.). *Deutsche*  
4585 *Entomologische Zeitschrift*, 1909, 239-269.
- 4586 Forel, A. (1911). Ameisen des Herrn Prof. v. Ihering aus Brasilien (Sao Paulo usw.) nebst einigen  
4587 anderen aus Südamerika und Afrika (Hym.). *Deutsche Entomologische Zeitschrift*, 285-  
4588 312.
- 4589 Forel, A. (1912a). Formicides néotropiques. Part VI. 5me sous-famille Camponotinae Forel.  
4590 *Mémoires de la Société Entomologique de Belgique*, 20, 59-92.
- 4591 Forel, A. (1912b). The Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the  
4592 leadership of Mr. J. Stanley Gardiner, M.A. Volume 4. No. XI. Fourmis des Seychelles et  
4593 des Aldabras, reçues de M. Hugh Scott. *Transactions of the Linnean Society of London.*  
4594 *Zoology*, 15(2), 159-167.
- 4595 Forel, A. (1914). Formicides d'Afrique et d'Amérique nouveaux ou peu connus. *Bulletin de la*  
4596 *Société Vaudoise des Sciences Naturelles*, 50, 211-288.
- 4597 Guénard, B., Weiser, M., Gomez, K., Narula, N., & Economo, E. P. (2017). The Global Ant  
4598 Biodiversity Informatics (GABI) database: a synthesis of ant species geographic  
4599 distributions. *Myrmecological News* 24, 83-89.
- 4600 Guénard, B. (2018). First record of the emerging global pest *Brachymyrmex patagonicus* Mayr  
4601 1868 (Hymenoptera: Formicidae) from continental Asia. *Asian Myrmecology*, 10, 1-6  
4602 (doi: 10.20362/am.010012).
- 4603 Grundmann, A. W. (1952). A new *Brachymyrmex* from northern Utah *Journal of the Kansas*  
4604 *Entomological Society*, 25, 117.
- 4605 Harris, R. A. (1979). A glossary of surface sculpturing. California Department of Food and  
4606 Agriculture. Laboratory Services, Entomology. *Occasional Papers*, 28, 1-31.

- 4607 Hebert, P. D. N., Cywinska, A., Ball, S. L., & DeWaard, J. R. (2003). Biological identifications  
4608 through DNA barcodes *Proceedings of the Royal Society B-Biological Sciences*, 270, 313-  
4609 321.
- 4610 Janicki, J., Narula, N., Ziegler, M., Guénard, B., & Economo, E.P. (2016). Visualizing and  
4611 interacting with large-volume biodiversity data using client-server web-mapping  
4612 applications: The design and implementation of antmaps.org. *Ecological Informatics* 32,  
4613 185-193.
- 4614 Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:  
4615 Improvements in Performance and Usability. *Molecular Biology and Evolution*, 30, 778-  
4616 780.
- 4617 Kempf, W. W. (1972). Catálogo abreviado das formigas da região Neotropical. *Studia*  
4618 *Entomologica*, 15, 3-344.
- 4619 Kruskal, J. (1964). Multidimensional scaling by optimizing goodness of fit to a nonmetric  
4620 hypothesis *Psychometrika*, 29, 1-27.
- 4621 Kugler, C. (1994). A revision of the ant genus *Rogeria* with description of the sting apparatus  
4622 (Hymenoptera: Formicidae). *Journal of Hymenoptera Research*, 3, 17-89.
- 4623 Kusnezov, N. (1959). La fauna de hormigas en el oeste de la Patagonia y Tierra del Fuego. *Acta*  
4624 *Zoológica Lilloana. Tucumán. Argentina*, 17, 321-401.
- 4625 Kusnezov, N. (1960). *Brachymyrmex physogaster* n. sp. aus Argentinien und das Problem der  
4626 Physogastrie bei den Ameisen. *Zoologischer Anzeiger*, 165, 381-388.
- 4627 Lanfear, R., Calcott, B., & Ho, S. Y. W. (2012). Guinon S PartitionFinder: combined selection  
4628 of partitioning schemes and substitution models for phylogenetic analyses *Molecular*  
4629 *Biology and Evolution*, 29(6), 1695-1701, doi:<http://dx.doi.org/10.1093/molbev/mss020>.

- 4630 LaPolla, J. S., Brady, S. G., & Shattuck, S. O. (2010). Phylogeny and taxonomy of the *Prenolepis*  
4631 genus-group of ants (Hymenoptera: Formicidae). *Systematic Entomology*, 35, 118-131.
- 4632 LaPolla, J. S., & Longino, J. T. (2006). An unusual new *Brachymyrmex* Mayr (Hymenoptera:  
4633 Formicidae) from Costa Rica, with implications for the phylogeny of the Lasiine tribe  
4634 group. *Proceedings of the Entomological Society of Washington*, 108, 297-305.
- 4635 Leliaert, F., Verbruggen, H., Vanormelingen, P., Steen, F., López-Bautista, J. M., Zuccarello, G.  
4636 C., et al. (2014). DNA-based species delimitation in algae *European Journal of*  
4637 *Phycology*, 49, 179-196.
- 4638 MacGown, J. A., Hill, J. G., & Deyrup, M. A. (2007). *Brachymyrmex patagonicus*  
4639 (Hymenoptera: Formicidae), an emerging pest species in the southeastern United States.  
4640 *Florida Entomologist*, 90, 457-464.
- 4641 Maddison, W. P., & Maddison, D. R. (2017). Mesquite: a modular system for evolutionary  
4642 analysis. (2.10 ed.).
- 4643 Mayr, G. (1868). Formicidae novae Americanae collectae a Prof. P. de Strobel. *Annuario della*  
4644 *Società dei Naturalisti e Matematici, Modena*, 3, 161-178.
- 4645 Mayr, G. (1870). Formicidae novogranadenses. *Sitzungsberichte der Kaiserlichen Akademie der*  
4646 *Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Classe. Abteilung I*, 61,  
4647 370-417.
- 4648 Mayr, G. (1887). Südamerikanische Formiciden. *Verhandlungen der Kaiserlich-Königlichen*  
4649 *Zoologisch-Botanischen Gesellschaft in Wien*, 37, 511-632.
- 4650 Menozzi, C. (1927). Formiche raccolte dal Sig. H. Schmidt nei dintorni di San José di Costa Rica  
4651 (Schluss). *Entomologische Mitteilungen. Berlin-Dahlem*, 16, 336-345.

- 4652 Miller, M. A., Pfeiffer, W., & Schwartz, T. "Creating the CIPRES Science Gateway for inference  
4653 of large phylogenetic trees". In *Proceedings of the Gateway Computing Environments*  
4654 *Workshop (GCE), New Orleans, LA, 14 Nov. 2010 2010* (pp. 1-8)
- 4655 Moreau, C. S., Bell, C. D., Vila, R., Archibald, S. B., & Pierce, N. E. (2006). Phylogeny of the  
4656 ants: diversification in the age of angiosperms. *Science*, *312*, 101-104.
- 4657 Moretti, T. C., Quirán, E. M., Solis, D. R., Rossi, M. L., & Thyssen, P. J. (2011). *Pycnoscelus*  
4658 *surinamensis* (Linnaeus, 1758) (Blaberoidea: Blaberidae), a cockroach with a possible  
4659 association with the ant *Brachymyrmex cordemoyi* Forel, 1895 (Hymenoptera:  
4660 Formicidae) and which may be exhibiting a domiciliation trend. *Symbiosis*, *53*, 37-39,  
4661 doi:10.1007/s13199-010-0101-3.
- 4662 Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., et al. (2015).  
4663 vegan: Community Ecology Package. Version 2.3-0.
- 4664 Ortiz, C. M., & Fernández, F. (2014). *Brachymyrmex* species with tumuliform metathoracic  
4665 spiracles: description of three new species and discussion of dimorphism in the genus  
4666 (Hymenoptera, Formicidae). *ZooKeys*, *371*, 13-33.
- 4667 Page, R.E., Jr. (1982) Polyandry in *Brachymyrmex depilis* Emery (Hymenoptera: Formicidae).  
4668 *Pan-Pacific Entomologist*, *58*, 258.
- 4669 Peeters, C. P. (1991). Ergatoid queens and intercast in ants: two distinct adult forms wich look  
4670 morphologically intermediate between workers and winged queens. *Insectes Sociaux*,  
4671 *38*(1), 1-15, doi:doi: 10.1007/BF01242708.
- 4672 Pons, J., Barraclough, T. G., Gomez-Zurita, J., Cardoso, A., Duran, D. P., Hazell, S., et al.  
4673 (2006). Sequence-based species delimitation for the DNA taxonomy of undescribed  
4674 insects *Systematic Biology*, *55*, 595-609.

- 4675 Puillandre, N., Lambert, A., Brouillet, S., & Achaz, G. (2011). ABDG, Automatic Barcode Gap  
4676 Discovery for primary species delimitation *Molecular Ecology*, *21*, 1864-1877.
- 4677 Quirán, E. M. (2005). El género neotropical *Brachymyrmex* Mayr (Hymenoptera: Formicidae) en  
4678 la Argentina. II: Redescrición de las especies *B. admotus* Mayr, de *B. brevicornis* Emery  
4679 y *B. gaucho* Santschi. *Neotropical Entomology*, *34*, 761-768.
- 4680 Quirán, E. M. (2007). El género neotropical *Brachymyrmex* Mayr (Hymenoptera: Formicidae) en  
4681 la Argentina. III: Redescrición de las especies: *B. aphidicola* Forel, de *B. australis* Forel  
4682 y *B. constrictus* Santschi. *Neotropical Entomology*, *36*, 699-706.
- 4683 Quirán, E. M., Martínez, J. J., & Bachmann, A. O. (2004). The Neotropical genus *Brachymyrmex*  
4684 Mayr, 1868 (Hymenoptera: Formicidae) in Argentina. Redescription of the type species,  
4685 *B. patagonicus* Mayr, 1868; *B. bruchi* Forel, 1912 and *B. oculatus* Santschi, 1919. *Acta*  
4686 *Zoológica Mexicana*, *20*, 273-285.
- 4687 R Core Team (2015). R: A language and environment for statistical computing. Version 3.2.1.  
4688 Vienna, Austria: R Foundation for Statistical Computing.
- 4689 Rambaut, A. (2012). FigTree. Version 1.4.0.
- 4690 Rambaut, A., Suchard, M. A., Xie, D., & Drummond, A. J. (2013). Tracer. Version 1.6.
- 4691 Roger, J. (1863). Die neu aufgeführten Gattungen und Arten meines Formiciden-Verzeichnisses,  
4692 nebst Ergänzung einiger früher gegeben Beschreibungen. *Berliner Entomologische*  
4693 *Zeitschrift*, *7*, 131-214.
- 4694 Ronquist, F., Teslenko, M., Van der Mark, P., Ayres, D., Darling, A., Höhna, S., et al. (2012).  
4695 MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large  
4696 model space *Systematic Biology*, *61*(3), 539-542.
- 4697 Santschi, F. (1912). Quelques fourmis de l'Amérique australe. *Revue Suisse de Zoologie*, *20*, 519-  
4698 534.

- 4699 Santschi, F. (1916). Formicides sudaméricains nouveaux ou peu connus. *Physis (Buenos Aires)*,  
4700 2, 365-399.
- 4701 Santschi, F. (1917). Description de quelques nouvelles fourmis de la République Argentine.  
4702 *Anales de la Sociedad Científica Argentina*, 84, 277-283.
- 4703 Santschi, F. (1919). Nouveaux formicides de la République Argentine. *Anales de la Sociedad*  
4704 *Científica Argentina*, 87, 37-57.
- 4705 Santschi, F. (1922). Description de nouvelles fourmis de l'Argentine et pays limitrophes. *Anales*  
4706 *de la Sociedad Científica Argentina*, 94, 241-262.
- 4707 Santschi, F. (1923a). Revue des fourmis du genre *Brachymyrmex* Mayr. *Anales del Museo*  
4708 *Nacional de Historia Natural de Buenos Aires*, 31, 650-678.
- 4709 Santschi, F. (1923b). *Solenopsis* et autres fourmis néotropicales. *Revue Suisse de Zoologie*, 30,  
4710 245-273.
- 4711 Santschi, F. (1929). Nouvelles fourmis de la République Argentine et du Brésil. *Anales de la*  
4712 *Sociedad Científica Argentina*, 107, 273-316.
- 4713 Santschi, F. (1933). Fourmis de la République Argentine en particulier du territoire de Misiones.  
4714 *Anales de la Sociedad Científica Argentina*, 116, 105-124.
- 4715 Santschi, F. (1939). Études et descriptions de fourmis néotropiques. *Revista de Entomologia (Rio*  
4716 *de Janeiro)*, 10, 312-330.
- 4717 Sharaf, M. R., Salman, S., Aldhafer, H. M., Youysef, A. F. A., & Aldawood, A. S. (2016). First  
4718 occurrence of the ant genus *Brachymyrmex* Mayr, 1868 (Hymenoptera: Formicidae) from  
4719 the Kingdom of Saudi Arabia. *Sociobiology*, 63(2), 800-803,  
4720 doi:10.13102/sociobiology.v63i2.981.



- 4721 Smith, D. R. (1979). Superfamily Formicoidea. In K. V. Krombein, P. D. Hurd, D. R. Smith, &  
4722 B. D. Burks (Eds.), *Catalog of Hymenoptera in America north of Mexico. Volume 2.*  
4723 *Apocrita (Aculeata)* (pp. 1323-1467). Washington, D.C.: Smithsonian Institution Press.
- 4724 Smith, M. R. (1955). Ants of the genus Pheidole, subgenus Hendecapheidole (Hymenoptera,  
4725 Formicidae). *Proceedings of the Entomological Society of Washington*, 57, 301-305.
- 4726 Snelling, R. R., & Hunt, J. H. (1975). The ants of Chile (Hymenoptera: Formicidae). *Revista*  
4727 *Chilena de Entomología*, 9, 63-129.
- 4728 Stamatakis, A., Hoover, P., & Rougemont, J. (2008). A Rapid Bootstrap Algorithm for the  
4729 RAxML Web-Servers. *Systematic Biology*, 57(5), 758-771.
- 4730 Swofford, D. L. (2002). PAUP\*: Phylogenetic analysis using parsimony (\*and other methods).  
4731 Version 4.0b: Sinauer Associates; Sunderland, Massachusetts.
- 4732 Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4ed.). New York:  
4733 Springer.
- 4734 Ward, P. S. (1989). Systematic studies on pseudomyrmecine ants: revision of the *Pseudomyrmex*  
4735 *oculatus* and *P. subtilissimus* species groups, with taxonomic comments on other species.  
4736 *Quaestiones Entomologicae*, 25, 393-468.
- 4737 Ward, P. S., Blaimer, B. B., & Fisher, B. L. (2016). A revised phylogenetic classification of the  
4738 ant subfamily Formicinae (Hymenoptera: Formicidae), with resurrection of the genera  
4739 *Colobopsis* and *Dinomyrmex*. *Zootaxa*, 4072, 343-357.
- 4740 Wheeler, G. C., & Wheeler, J. (1953). The ant larvae of the myrmicine tribe Pheidolini  
4741 (Hymenoptera, Formicidae). *Proceedings of the Entomological Society of Washington*,  
4742 55, 49-84.
- 4743 Wheeler, G. C., & Wheeler, J. (1978). *Brachymyrmex musculus*, a new ant in the United States.  
4744 *Entomological News*, 89, 189-190.

- 4745 Wheeler, G. C., & Wheeler, J. (1982). Supplementary studies on ant larvae: Formicinae  
4746 (Hymenoptera: Formicidae). *Psyche*, 89, 175-181.
- 4747 Wheeler, G.C., & Wheeler, J. (1986) The ants of Nevada. Natural History Museum of Los  
4748 Angeles County, Los Angeles, vii + 138 pp.
- 4749 Wheeler, W. M. (1903). A decad of Texan Formicidae. *Psyche*, 10, 93-111.
- 4750 Wheeler, W. M. (1910). *Ants: their structure, development and behavior*. New York: Columbia  
4751 University Press.
- 4752 Wheeler, W. M. (1921). The Tachigalia ants. *Zoologica (New York)*, 3, 137-168.
- 4753 Wheeler, W. M. (1922). Ants of the American Museum Congo Expedition. A contribution to the  
4754 Myrmecology of Africa. *Bulletin of the American Museum of Natrural History*, 45, 1-582.
- 4755 Wheeler, W. M. (1934). Neotropical ants collected by Dr. Elisabeth Skwarra and others. *Bulletin*  
4756 *of the Museum of Comparative Zoology*, 77, 157-240.
- 4757 Wheeler, W. M. (1938). Ants from the caves of Yucatan. In A. S. Pearse (Ed.), *Fauna of the*  
4758 *caves of Yucatan* (Vol. 491, pp. 251-304): Carnegie Institution of Washington  
4759 Publication.
- 4760 Wheeler, W. M. (1942). Studies of Neotropical ant-plants and their ants. *Bulletin of the Museum*  
4761 *of Comparative Zoology*, 90, 1-262.
- 4762 Wild, A. L. (2007). A catalogue of the ants of Paraguay (Hymenoptera: Formicidae). *Zootaxa*,  
4763 1622, 1-55.
- 4764 Wilson, C. M., Smith-Herron, A., & Cook, J. L. (2016). Morphology of the male genitalia of  
4765 *Brachymyrmex* and their implications in the Formicinae phylogeny. *Journal of*  
4766 *Hymenoptera Research*, 50, 81-95, doi:DOI 10.3897/JHR.50.8697.
- 4767 Wilson, E. O., & Taylor, R. W. (1967). The ants of Polynesia (Hymenoptera: Formicidae).  
4768 *Pacific Insects Monograph*, 14, 1-109.

- 4769 Xia, X. (2013). DAMBE5: A Comprehensive Software Package for Data Analysis in Molecular  
4770 Biology and Evolution. *Molecular Biology and Evolution*, 30(7), 1720-1728,  
4771 doi:10.1093/molbev/mst064.
- 4772 Yensen, N., Yensen, E. & Yensen, D. (1980) Intertidal ants from the Gulf of California, Mexico.  
4773 *Annals of the Entomological Society of America*, 73, 266–269.
- 4774

4775 **FIGURE CAPTIONS**

4776

4777 **Fig. 1** Morphological measurements for *Brachymyrmex* workers. See text for details

4778 **Fig. 2** Habitus of a selection of queens of *Brachymyrmex*: head and lateral view of **(a,b)** *B.*

4779 *admotus*; **(c,d)** *B. antennatus*; **(e,f)** *B. aphidicola* and **(g,h)** *B. giardi*

4780 **Fig. 3** Habitus of a selection of males of *Brachymyrmex*: head and lateral view of **(a,b)** *B.*

4781 *coactus*; **(c,d)** *B. myops*; **(e,f)** *B. longicornis* var. *immunis* (junior synonym of *B. admotus*) and

4782 **(g,h)** *B. australis* var. *curta* (= *B. australis*)

4783 **Fig. 4** The native distribution range of *Brachymyrmex* as reconstructed from the unique

4784 georeferenced localities of the here studied material (black circles) and the *Brachymyrmex*

4785 records available in the Global Ant Biodiversity Informatics database (Guénard et al. 2017) as

4786 viewed in [www.antmaps.org](http://www.antmaps.org) (Janicki et al. 2016; shaded area)

4787 **Fig. 5** Morphological characteristics of the head of *Brachymyrmex*. **(a1)** clypeus with five hairs of

4788 which a single long apical hair is positioned near the anterior margin, two in mediolateral

4789 position and two near the toruli (black arrow); **(a2)** clypeus with a row of long, thick hairs near

4790 the anterior margin (black arrow), toruli touching but not surpassing the posterior clypeal margin

4791 in oblique anterodorsal view (grey arrow); **(a3)** toruli surpassing the posterior clypeal margin in

4792 oblique anterodorsal view (grey arrow); **(b1)** eyes below the cephalic midline; **(b2)** eyes on

4793 cephalic midline; **(c1)** eyes with 3 or 4 ommatidia along the maximal diameter of the eye; **(c2)**

4794 eyes with more than 4 ommatidia along the maximal diameter of the eye; **(d1)** scapes short and

4795 not reaching the posterior margin of the head; **(d2)** scapes just reaching the posterior margin of

4796 the head; **(d3)** scapes long and surpassing the posterior margin of the head; the length by which

4797 the scapes surpass this margin is compared to the length of the maximal diameter of the eye; **(e1)**

4798 anterior clypeal margin with the medial portion forming a “lip”; **(e2)** anterior clypeal margin

4799 evenly convex without antero-medial “lip”; (**f1**) head with dense pilosity; (**f2**) head with sparse  
4800 decumbent hairs; (**g1**) second segment of the antennal funiculus shorter than the first; (**g2**) second  
4801 segment of the antennal funiculus as long or longer than the first

4802 **Fig. 6** Morphological characteristics of the mesosoma and gaster of *Brachymyrmex*. (**a1**)

4803 metathoracic spiracles tumiliform; (**a2**) metathoracic spiracles not protruding; (**a3**) metathoracic

4804 spiracles slightly protruding, but not tumiliform; (**b1**) mesonotum inflated and bulging dorsally

4805 above the pronotum in lateral view; (**b2**) mesonotum not inflated or bulging dorsally above the

4806 pronotum in lateral view; (**c**) dorsal margin of the mesosoma of conspicuous sinusoidal shape;

4807 (**d1**) mesometanotal suture inconspicuous (dashed line); (**d2**) mesometanotal suture directly

4808 visible; (**e1**) mesonotum strongly antero-posteriorly inclined and thus elongated in lateral view;

4809 (**e2**) mesonotum weakly antero-posteriorly inclined in lateral view; (**f1**) metanotal groove deep

4810 and wider than the diameter of the metathoracic spiracles; (**f2**) metanotal groove shallow and

4811 narrower than the diameter of the metathoracic spiracles; (**g1**) gaster with scattered pubescence;

4812 (**g2**) gaster with dense pubescence, in both cases illustrated with long erect hairs near the edges of

4813 the segments; (**h1**) mesonotum laterally extended and oval in dorsal view; (**h2**) mesonotum

4814 almost circular in dorsal view

4815 **Fig. 7** *Brachymyrmex admotus*: (**a,c,e**) head, dorsal and lateral view of the lectotype worker;

4816 (**b,d,f**) = *B. longicornis* var. *immunis* **n. syn.**: head, dorsal and lateral view of a syntype worker

4817 **Fig. 8** *Brachymyrmex antennatus*: (**a,b,c**) head, dorsal and lateral view of the lectotype worker

4818 **Fig. 9** *Brachymyrmex aphidicola*: (**a,d,g**) head, dorsal and lateral view of the lectotype worker;

4819 (**b,e,h**) = *B. heeri* var. *fallax*: head, dorsal and lateral view of a syntype worker; (**c, f**) = *B.*

4820 *longicornis* var. *hemiops* **n. syn.**: head and dorsal view of a syntype worker

4821 **Fig. 10** *Brachymyrmex attenuatus*: head, dorsal and lateral view of the lectotype worker (**a,b,c**)

- 4822 **Fig. 11 *Brachymyrmex australis*: (a,d,g)** head, dorsal and lateral view of the lectotype worker;  
4823 **(b,e,h) =*B. australis* var. *curta* n. syn.:** head, dorsal and lateral view of a syntype worker; **(c,f)**  
4824 **=*B. longicornis* n. syn.:** head and dorsal view of a syntype worker
- 4825 **Fig. 12 *Brachymyrmex bahamensis* n.sp.:** **(a,b,c)** head, dorsal and lateral view of the holotype  
4826 worker
- 4827 **Fig. 13 *Brachymyrmex bicolor* n. sp.:** **(a,c,e)** head, dorsal and lateral view of a paratype worker;  
4828 **(b,d,f)** head, dorsal and lateral view of a syntype specimen of the putative worker-queen inter-  
4829 caste. (from [www.AntWeb.org](http://www.AntWeb.org); photographer: Ryan Perry)
- 4830 **Fig. 14 *Brachymyrmex bonariensis* n. st.:** **(a,b,c)** head, dorsal and lateral view of the lectotype  
4831 worker
- 4832 **Fig. 15 *Brachymyrmex brasiliensis*: (a,b,c)** head, dorsal and lateral view of the holotype worker
- 4833 **Fig. 16 *Brachymyrmex bruchi*: (a,c,e)** head, dorsal and lateral view of the lectotype worker;  
4834 **(b,d,f) =*B. giardi* var. *nitida*:** head, dorsal and lateral view of a syntype worker
- 4835 **Fig. 17 *Brachymyrmex bruchi*: (a,c,e) =*B. laevis* var. *andina*:** head, dorsal and lateral view of a  
4836 syntype worker; **(b,d,f) =*B. bruchi* var. *rufipes*:** head, dorsal and lateral view of a syntype worker
- 4837 **Fig. 18 *Brachymyrmex cavernicola*: (a,b,c)** head, dorsal and lateral view of a worker (from  
4838 [www.AntWeb.org](http://www.AntWeb.org); photographer: Estella Ortega)
- 4839 **Fig. 19 *Brachymyrmex coactus*: (a,c,e)** head, dorsal and lateral view of the lectotype worker,  
4840 **(b,d,f) =*B. coactus* var. *nictitans* n. syn.:** head, dorsal and lateral view of a syntype worker
- 4841 **Fig. 20 *Brachymyrmex coactus*: (a,c,e) =*B. constrictus* n. syn.:** head, dorsal and lateral view of a  
4842 syntype worker; **(b,d,f) =*B. robustus*:** head, dorsal and lateral view of a syntype worker
- 4843 **Fig. 21 *Brachymyrmex cordemoyi*: (a,c,e) =*B. laevis* var. *fuscula* n. syn.:** head, dorsal and lateral  
4844 view of a syntype worker; **(b,d,f) =*B. cordemoyi* var. *nigricans*:** head, dorsal and lateral view of a  
4845 syntype worker

4846 **Fig. 22 *Brachymyrmex cordemoyi*: (a,b,c) =*B. brevicornis* n. syn.**: head, dorsal and lateral view  
4847 of a syntype worker

4848 **Fig. 23 *Brachymyrmex cordemoyi*: (a,c,e) =*B. brevicornoides* n. syn.**: head, dorsal and lateral  
4849 view of a syntype worker (from [www.AntWeb.org](http://www.AntWeb.org); photographer: Zach Lieberman); **(b,d,f) =*B.***  
4850 ***cordemoyi* var. *distincta* n. syn.**: head, dorsal and lateral view of a syntype worker

4851 **Fig. 24 *Brachymyrmex degener*: (a,c,e)** head, dorsal and lateral view of the lectotype worker;  
4852 **(b,d,f) =*B. admotus* r. *niger* n. syn.** head, dorsal and lateral view of a syntype worker

4853 **Fig. 25 *Brachymyrmex degener*: (a,c,e) =*B. incisus* n. syn.**: head, dorsal and lateral view of a  
4854 syntype worker; **(b,d,f) =*B. luederwaldti* n. syn.**: head, dorsal and lateral view of a syntype  
4855 worker

4856 **Fig. 26 *Brachymyrmex delabiei*: (a,b,c)** head, dorsal and lateral view of the holotype worker

4857 **Fig. 27 *Brachymyrmex depilis*: (a,c,e)** head, dorsal and lateral view of the lectotype worker;  
4858 **(b,d,f) =*B. depilis* subsp. *flavescens*:** head, dorsal and lateral view of a syntype worker

4859 **Fig. 28 *Brachymyrmex donisthorpei*: (a,b,c)** head, dorsal and lateral view of the lectotype  
4860 worker

4861 **Fig. 29 *Brachymyrmex feitosai*: (a,b,c)** head, dorsal and lateral view of a worker (from  
4862 [www.AntWeb.org](http://www.AntWeb.org); photographer: Erin Prado)

4863 **Fig. 30 *Brachymyrmex fiebrigi*: (a,c,e)** head, dorsal and lateral view of the lectotype worker;  
4864 **(b,d,f) =*B. fiebrigi* var. *fumida* n. syn.**: head, dorsal and lateral view of a syntype worker

4865 **Fig. 31 *Brachymyrmex fiebrigi*: (a,b,c) =*B. fiebrigi* var. *funicularis* n. syn.**: head, dorsal and  
4866 lateral view of a syntype worker

4867 **Fig. 32 *Brachymyrmex gagates*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4868 **Fig. 33 *Brachymyrmex gaucho*: (a,b,c)** head, dorsal and lateral view of a worker

4869 **Fig. 34 *Brachymyrmex giardi*: (a,c,e)** head, dorsal and lateral view of the lectotype worker;  
4870 **(b,d,f)** head, dorsal and lateral view of a putative worker-queen intercaste

4871 **Fig. 35 *Brachymyrmex heeri*: (a,c,e)** head, dorsal and lateral view of the lectotype worker;  
4872 **(b,d,f)** =*Brachymyrmex* var. *goeldii* **n. syn.:** head, dorsal and lateral view of a syntype worker

4873 **Fig. 36 *Brachymyrmex heeri*: (a,c,e)** =*B. giardi* var. *cordobensis*: head, dorsal and lateral view  
4874 of a syntype worker; **(b,d,f)** head, dorsal and lateral view of a syntype of the putative worker-  
4875 queen intercaste

4876 **Fig. 37 *Brachymyrmex iridiscens* n. sp.:** **(a,b,c)** head, dorsal and lateral view of the holotype  
4877 worker

4878 **Fig. 38 *Brachymyrmex micromegas*: (a,c,e)** head, dorsal and lateral view of the lectotype  
4879 worker; **(b,d,f)** head, dorsal and lateral view of a soldier

4880 **Fig. 39 *Brachymyrmex minutus*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4881 **Fig. 40 *Brachymyrmex modestus*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4882 **Fig. 41 *Brachymyrmex musculus*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4883 **Fig. 42 *Brachymyrmex myops*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4884 **Fig. 43 *Brachymyrmex nebulosus*: (a,b,c)** head, dorsal and lateral view of a worker (from  
4885 [www.AntWeb.org](http://www.AntWeb.org); photographer: Ryan Perry)

4886 **Fig. 44 *Brachymyrmex obscurior*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4887 **Fig. 45 *Brachymyrmex oculatus*: (a,b,c)** head, dorsal and lateral view of the lectotype worker

4888 **Fig. 46 *Brachymyrmex patagonicus*: (a,c,e)** head, dorsal and lateral view of the lectotype  
4889 worker; **(b,d,f)** =*B. laevis* **n. syn.:** head, dorsal and lateral view of a syntype worker

4890 **Fig. 47 *Brachymyrmex patagonicus*: (a,b,c)** =*B. patagonicus* var. *atratura*: head, dorsal and  
4891 lateral view of a syntype worker



4892 **Fig. 48** *Brachymyrmex pictus*: (a,b,c) head, dorsal and lateral view of a syntype worker (from  
4893 [www.AntWeb.org](http://www.AntWeb.org); photographer: Zach Lieberman)

4894 **Fig. 49** *Brachymyrmex pilipes*: (a,c,e) head, dorsal and lateral view of the minor worker  
4895 lectotype; (b,d) head, dorsal and lateral view of a major worker

4896 **Fig. 50** *Brachymyrmex santschii*: (a,c,e) head, dorsal and lateral view of a worker (from  
4897 [www.AntWeb.org](http://www.AntWeb.org); photographer: Will Ericson); (b,d,f) head, dorsal and lateral view of a  
4898 putative worker-queen intercaste

4899 **Fig. 51** *Brachymyrmex sosai* n. sp.: (a,b,c) head, dorsal and lateral view of the holotype worker

4900 **Fig. 52** *Brachymyrmex termitophilus*: (a,b,c) head, dorsal and lateral view of the lectotype  
4901 worker (from [www.AntWeb.org](http://www.AntWeb.org); photographer: Zach Lieberman)

4902 **Fig. 53** *Brachymyrmex tristis*: (a,b,c) head, dorsal and lateral view of the lectotype worker


4903 **Fig. 54** Morphospace occupation of 38 of the here studied *Brachymyrmex* species as  
4904 reconstructed with non-metric multidimensional scaling. The limited stress (5.70) indicates that  
4905 the ordination is robust and the biplot displays how the various morphometric variables  
4906 contribute to the morphospace occupation. OI<sub>1</sub> is indicated in gray, as this variable did not  
4907 contribute significantly to the morphospace

4908 **Fig. 55** Boxplots representing intraspecific variation and interspecific differences for eight  
4909 morphometric traits. Interspecific differences are tested with Benjamini-Hochberg corrected  
4910 pairwise Dunn's tests, with significance levels indicated by letter codes (if species carry at least  
4911 one identical letter than observed differences are insignificant, if they carry no identical letter, the  
4912 observed differences for the studied trait are significant)

4913 **Fig. 56** Boxplots representing intraspecific variation and interspecific differences for eight  
4914 morphometric traits. Interspecific differences are tested with Benjamini-Hochberg corrected  
4915 pairwise Dunn's tests, with significance levels indicated by letter codes (if species carry at least

4916 one identical letter than observed differences are insignificant, if they carry no identical letter, the  
4917 observed differences for the studied trait are significant)


4918 **Fig. 57** Maximum clade credibility tree of *Brachymyrmex* and *Myrmelachista* based on five gene  
4919 fragments (see Supplementary material Table S1). Analyses were run under maximum parsimony  
4920 (MP), maximum likelihood (ML) and Bayesian inference (BI) with bootstrap support values and  
4921 Bayesian posterior probabilities indicated above nodes (MP/ML/BI). Support for species-level  
4922 clades is indicated in bold; specimens were assigned to clades based on the morphological  
4923 identification system, which proves to be overall in good agreement with the genealogy, apart  
4924 from *B. degener* and *B. coactus*



























































































































































*admotus* *antenattus* *aphidicola* *australis* *bruchi* *cavernicola* *coactus* *cordermoyi* *degener* *delabie* *depilis* *donisthorpei* *fiebrigi* *heeri* *iridiscens* *modestus* *musculus* *obscurior* *patagonicus* *santschii*

0.05

