Coprinopsis variegata (Peck) Redhead, Vilgalys & Moncalvo









OVERALL: Daniels, PVSP, 6/19/2009, *J. Solem.*GILLS and DELIQUESCING CAP: Schooley Mill Park,5/18//2009, *J. Solem.* 

CUT TOP: Daniels, PVSP, 6/19/2009, J. Solem.

DELIQUESCING GROUP: Savage Park, 7/19/2008, J. Solem.

J F M A M J J A S O N D +++++++++

**ID**: Cap grayish w/ dingy scales; egg-shaped when young. Deliquesces. Odor unpleasant (garlicky).

**Habitat**: Dense clusters. On decaying hardwood (often on buried wood in lawns).

**Cap**: 1.5 - 3" [4 - 7.5 cm]

Grayish-brown/gray w/ white/dingy yellow scales that wash off. Curved to egg-/bell-shaped in age. Flesh pale tan. Garlic odor (strong and unpleasant).

**Gills**: White; quickly gray, purplish-brown/black in age and deliquescing. Free. Very crowded.

**Spores**: *Black*. Ellipsoid w/ apical pore, smooth.

**Stalk**: 2 - 4.8"x 0.2 - 0.4" [5 - 12 cm x 5 - 10 mm]

Whitish. Scurfy/smooth near top; fibrillose from base to obscure ring area. Tapers to top. Whitish flesh. Hollow in age.

Frequency: Common.

**Locations**: BENBR, CASCT, DANPV, GONRA, MCHPV, MPENA, MTPLT, MURHL, PATBT, SLMLP, WINFR, WINTR.

**Notes**: Mycobank 474342. Schooley Mill Park specimen identified by examination of spores by R. Solem.

**Synonym**: Coprinus variegatus. Coprinus quadrifidus.

**References**: Bar 195. BBF 105, 281. BBH 180. Bni 331. BRBD 126fn. E&S 151fn. K&M 150. Lin 601, 703. Phi 236. Roo 206. Stu 220. W&L 278-9.



SPORES (8.0-8.6 X 5.2-5.6 μm): Schooley Mill Park, 5/18//2009, *R. Solem*.