



# Awel y Môr Offshore Wind Farm

## Category 6: Environmental Statement

### Volume 4, Annex 5.1: Benthic Ecology Subtidal Characterisation (Array)

**Date:** April 2022

**Revision:** A

Application Reference: 6.4.5.1

Pursuant to: APFP Regulation 5(2)(a)



| REVISION | DATE          | STATUS/<br>REASON<br>FOR ISSUE | AUTHOR: | CHECKED<br>BY: | APPROVED<br>BY: |
|----------|---------------|--------------------------------|---------|----------------|-----------------|
| A        | March<br>2022 | ES                             | Fugro   | RWE            | RWE             |
|          |               |                                |         |                |                 |
|          |               |                                |         |                |                 |
|          |               |                                |         |                |                 |

[www.awelymor.cymru](http://www.awelymor.cymru)

RWE Renewables UK  
Swindon Limited

Windmill Hill Business Park  
Whitehill Way  
Swindon  
Wiltshire SN5 6PB  
T +44 (0)8456 720 090  
[www.rwe.com](http://www.rwe.com)

Registered office:  
RWE Renewables UK  
Swindon Limited  
Windmill Hill Business Park  
Whitehill Way  
Swindon



---

# Fugro – WPM1 - Main Array – Benthic Ecology Monitoring Report

Offshore Site Investigation, Awel y Môr Offshore Wind Farm | UK, Irish Sea  
Survey Period: 16 to 30 August 2020

003616051-02 | 9 February 2021

Final

**Awel y Môr Offshore Windfarm Ltd**

# Document Control

## Document Information

|                        |   |
|------------------------|---|
| Project Title          | Offshore Site Investigation Awel y Môr Offshore Windfarm                              |
| Document Title         | Fugro – WPM1 - Main Array – Benthic Ecology Monitoring Report                         |
| Fugro Project No.      | 200145  |
| Client Document No.    | 003616051-02  |
| Fugro Document No.     | 200145-R-005(02)  |
| Issue Number           | 02  |
| Issue Status           | Final   |
| Fugro Legal Entity     | Fugro GB Marine Limited   |
| Issuing Office Address | Victory House, 16 Trafalgar Wharf, Hamilton Road, Portchester, Hampshire, PO6 4PX, UK |

## Client Information

|                     |   |
|---------------------|---|
| Client              | Awel y Môr Offshore Windfarm Ltd  |
| Client Address      | Trigonos, Windmill Hill Business Park, Whitehill Way, Swindon SN5 6PB, UK |
| Client Contact      | Mark Osola  |
| Client Document No. | 003616051-02  |

## Revision History


| Issue | Date             | Status   | Comments on Content | Prepared By | Checked By | Approved By |
|-------|------------------|----------|---------------------|-------------|------------|-------------|
| 01    | 18 December 2020 | Complete | First issue         | AOW/AIS     | GHB/SRB    | CAB         |
| 02    | 9 February 2021  | Final    | Final               | RVC         | JLL        | CAB         |

## Project Team

| Initials | Name                  | Role   |
|----------|-----------------------|--|
| CAB      | Craig Barrett         | Project Manager                                  |
| GHB      | Geraint Harris-Bryant | Chief Environmental Scientist                    |
| JLL      | Jo Lusted             | Environmental Group Survey and Reporting Manager |
| RVC      | Rachel Cox            | Marine Environmental Scientist                   |
| SRB      | Susan Brittain        | Principal Chemist                                |
| AIS      | Andrew Sutherland     | Senior Environmental Consultant                  |
| AOW      | Angus Williams        | Assistant Marine Environmental Chemist           |



# Frontispiece



---

# Executive Summary

## Introduction

On the instruction of Awel y Môr Offshore Windfarm Ltd, Fugro performed a geophysical and environmental site investigation at the proposed Awel y Môr Offshore Windfarm (OWF) site, located 10.6 km off the north coast of Wales. The proposed OWF development will consist of the main array and three export cable routes, East A and B and West C.

Environmental operations were conducted onboard the MV Mersey Guardian during the survey period 16 to 30 August 2020. This report details the results of the benthic characterisation survey of the main array survey area.

## Survey Strategy

A total of 62 grab sampling stations were selected across the survey area. Forty-two of the stations were selected following consultation with Natural Resources Wales (NRW) to increase the sampling spatial coverage. At each proposed station, one macrofaunal sample with one particle size distribution (PSD) subsample were to be acquired. Of these 62 stations, 10 were selected for additional sampling for sediment chemistry. At 22 of the stations, drop-down video (DDV) data were to be acquired for spatial coverage or to investigate boundaries between two sediment types. A complete suite of grab samples were successfully acquired from all 62 stations and video and stills data were successfully acquired along all 22 DDV locations.

## Sediment Characteristics

The sediments within the survey area comprised mainly sand, with varying proportions of gravel, and little to no fines. When the mean particle size was expressed using the Wentworth classification, sediments were described as medium sand to very coarse sand. Spatial patterns were evident in sediment composition and the clear linkage of these to the geophysical characteristics of the seafloor, suggest that particle size is likely to be influenced by the degree of sediment mobility.

## Sediment Chemistry

The median total 2 to 6 ring polycyclic aromatic hydrocarbon (PAH) concentration was broadly comparable to the median concentration recorded during the Strategic Environmental Assessment (SEA6 area) Irish Sea surveys. All individual PAH concentrations were below their respective effects range low (ERL) values.

All metals concentrations were less than their respective Cefas guideline action levels (AL1 and AL2) and Oslo and Paris Commission (OSPAR) ERL values.

## Macrofauna

The number of infaunal and solitary epifaunal taxa recorded from the grab samples was variable across the survey area. Three statistically significantly different communities, characterised by differing

infaunal taxa, were identified from multivariate statistical analysis. Variations in macrofaunal community were linked to variations in sediment composition, which could, in turn, be linked to variations in wave/tidal exposure.

## Seabed Habitats and Biotopes

One habitat, one benthic biotope complex and two biotopes were defined within the survey area; 'Circalittoral coarse sediment' (A5.14), '*Branchiostoma lanceolatum* in circalittoral coarse sand with shell gravel' (A5.145), 'Sublittoral sand' (A5.2) and '*Nephtys cirrosa* and *Bathyporeia* spp. in infralittoral sand' (A5.233).

These habitats are encompassed within 'Subtidal sands and gravels', a priority habitat within UK waters.

No other Annex I habitats or Annex II species, OSPAR threatened and/or declining species and habitats or UK Biodiversity Action Plan priority habitats and species observed within the survey area.

---

## Document Arrangement

|                 |  |
|-----------------|--|
| Volume 1        | WPM1 & WPM2 Array Area & ECR Acquisition/Operations Report - Mersey Guardian |
| Volume 2        | Fugro – WPM1- Main Array - Environmental Features Report                     |
| Volume 3        | Fugro – WPM2 - ECR West C – Environmental Features Report                    |
| Volume 4        | Fugro – WPM2 - ECR East A&B - Environmental Features Report                  |
| <b>Volume 5</b> | <b>Fugro – WPM1 – Main Array – Benthic Ecology Monitoring Report</b>         |
| Volume 6        | Fugro – WPM2 - ECR West C - Benthic Ecology Monitoring Report                |
| Volume 7        | Fugro – WPM2 - ECR East A&B - Benthic Ecology Monitoring Report              |

# Contents

|  |            |
|--|------------|
| <b>Executive Summary</b>   | <b>i</b>   |
| <b>Document Arrangement</b>  | <b>iii</b> |
| <b>1. Introduction</b>   | <b>1</b>   |
| 1.1 General Project Description  | 1          |
| 1.2 Scope of Work  | 1          |
| 1.2.1 Geophysical Survey   | 1          |
| 1.2.2 Environmental Survey   | 1          |
| 1.3 Environmental Legislation  | 1          |
| 1.4 Regional Habitats, Species and Protected Areas                       | 2          |
| 1.5 Environmental Quality Standards for Sediment Chemical Concentrations | 5          |
| 1.6 Coordinate Reference System  | 5          |
| <b>2. Survey Strategy</b>  | <b>6</b>   |
| <b>3. Methods</b>  | <b>11</b>  |
| 3.1 Survey Methods   | 11         |
| 3.1.1 Seabed Photography   | 11         |
| 3.1.2 Sediment Sampling  | 11         |
| 3.2 Analytical Methods   | 11         |
| 3.2.1 Sediment Characteristics   | 11         |
| 3.2.2 Sediment Hydrocarbons  | 11         |
| 3.2.3 Sediment Metals  | 12         |
| 3.2.4 Sediment Macrofauna  | 12         |
| 3.3 Data Analysis  | 12         |
| 3.3.1 Sediment Particle Size Distribution (PSD) Statistics               | 12         |
| 3.3.2 Sediment Macrofauna  | 13         |
| 3.3.3 Seabed Habitats and Biotopes                                       | 15         |
| <b>4. Results</b>  | <b>18</b>  |
| 4.1 Field Operations   | 18         |
| 4.1.1 Bathymetry and Seabed Features                                     | 18         |
| 4.1.2 Seabed Photography   | 18         |
| 4.1.3 Seabed Sampling  | 20         |
| 4.2 Sediment Characterisation  | 24         |
| 4.3 Sediment Chemistry   | 39         |
| 4.3.1 Sediment Hydrocarbons  | 39         |
| 4.3.2 Sediment Metals  | 41         |
| 4.4 Sediment Macrofauna  | 45         |
| 4.4.1 Infaunal and Solitary Epifauna                                     | 45         |

|           |  |           |
|-----------|--|-----------|
| 4.4.2     | Colonial Epifauna                        | 66        |
| 4.5       | Seabed Habitats and Biotopes             | 71        |
| 4.5.1     | Biotope Classifications                  | 71        |
| 4.5.2     | Potential Sensitive Habitats and Species | 80        |
| <b>5.</b> | <b>Discussion</b>                        | <b>81</b> |
| 5.1       | Sediment Characterisation                | 81        |
| 5.2       | Sediment Chemistry                       | 82        |
| 5.2.1     | Sediment Hydrocarbons                    | 82        |
| 5.2.2     | Sediment Metals                          | 83        |
| 5.3       | Macrofaunal Communities                  | 83        |
| 5.4       | Seabed Habitats and Biotopes             | 86        |
| <b>6.</b> | <b>Conclusions</b>                       | <b>88</b> |
| <b>7.</b> | <b>References</b>                        | <b>89</b> |

## Appendices

### Appendix A Guidelines on Use of Report

---

### Appendix B Methodologies

- B.1 Survey Methods
  - B.2 Laboratory Analysis for Sediment Samples
  - B.3 Statistical Analysis
- 

### Appendix C Logs

- C.1 Survey Log
  - C.2 Grab Log
  - C.3 Video and Photographic Log
- 

### Appendix D Sediment Particle Size and Grab Sample Photographs

---

### Appendix E Sediment Hydrocarbon Analysis

- E.1 United States Environmental Protection Agency (US EPA) 16 Polycyclic Aromatic Hydrocarbon (PAH) Concentrations
  - E.2 Total 2 to 6 Ring PAH Concentrations
- 

### Appendix F Macrofaunal Analysis

- F.1 Macrofaunal Abundance
  - F.2 Macrofaunal Biomass
-

## Figures in the Main Text

|   |    |
|---|----|
| Figure 1.1: Protected areas relevant to the survey area, Main Array   | 4  |
| Figure 2.1: Proposed environmental survey locations, Main Array   | 10 |
| Figure 4.1: Completed survey locations overlaid on bathymetry, Main Array   | 23 |
| Figure 4.2: Sediment fractional composition overlaid on bathymetry, Main Array  | 32 |
| Figure 4.3: Sediment mean particle size ( $\mu\text{m}$ ) overlaid on bathymetry, Main Array  | 33 |
| Slice at 1.5 resemblance based on Euclidean distance  | 35 |
| Figure 4.4: Dendrogram of hierarchical clustering of sediment characteristics data, Main Array  | 35 |
| Figure 4.5: Sediment groups identified in multivariate analysis overlaid on a side scan sonar mosaic, Main Array                      | 37 |
| Figure 4.6: Non-metric multidimensional scaling ordination of particle sizes ( $\mu\text{m}$ )/fractional composition (%), Main Array | 38 |
| Figure 4.7: PCA ordination of particle sizes ( $\mu\text{m}$ )/fractional composition (%), Main Array                                 | 39 |
| Figure 4.8: Relative (maximum normalised) elemental concentrations in sediments, Main Array   | 44 |
| Figure 4.9: Phyletic composition of macrofaunal taxa, Main Array  | 47 |
| Figure 4.10: Phyletic composition of macrofaunal individuals, Main Array  | 47 |
| Figure 4.11: Number of macrofaunal taxa per station/sample ( $0.1 \text{ m}^2$ ) overlaid on bathymetry, Main Array                   | 51 |
| Figure 4.12: Number of macrofaunal individuals per station/sample ( $0.1 \text{ m}^2$ ) overlaid on bathymetry, Main Array            | 52 |
| Figure 4.13: Dendrogram of hierarchical clustering of macrofaunal station ( $0.1 \text{ m}^2$ ) abundance data, Main Array            | 54 |
| Figure 4.14: Macrofaunal clusters per station ( $0.1 \text{ m}^2$ ) overlaid on bathymetry Main Array                                 | 55 |
| Figure 4.15: <i>Urothoe marina</i> abundance overlaid on macrofaunal community groupings and a bathymetric mosaic, Main Array         | 58 |
| Figure 4.16: <i>Pisione remota</i> abundance overlaid on macrofaunal community groupings and a bathymetric mosaic, Main Array         | 59 |
| Figure 4.17: <i>Bathyporeia elegans</i> abundance overlaid on macrofaunal community groupings and a bathymetric mosaic, Main Array    | 60 |
| Figure 4.18: Phyletic composition of biomass, Main Array  | 64 |
| Figure 4.19: Total biomass overlaid on bathymetry, Main Array   | 65 |
| Figure 4.20: Number of epifaunal taxa per station ( $0.1 \text{ m}^2$ ) overlaid on bathymetry, Main Array                            | 70 |
| Figure 4.21: The spatial distribution of EUNIS (EEA, 2019) habitat classifications, Main Array  | 79 |

## Tables in the Main Text

|  |    |
|--|----|
| Table 1.1: Gwynt y Môr OWF (CMACS, 2005) habitat classifications, Main Array | 3  |
| Table 1.2: Summary of nearby protected areas, Main Array                     | 3  |
| Table 1.3: Project geodetic and projection parameters, Main Array            | 5  |
| Table 2.1: Proposed sampling stations, Main Array                            | 6  |
| Table 2.2: Proposed transects, Main Array                                    | 9  |
| Table 3.1: Sediment particle size distribution statistics, Main Array        | 13 |
| Table 3.2: Macrofaunal univariate statistics, Main Array                     | 13 |



|  |    |
|--|----|
| Table 3.3: Macrofaunal standard biomass corrections by phyla, Main Array           | 14 |
| Table 3.4: Sediment particle size and classification terms, Main Array             | 16 |
| Table 3.5: EEA (2019) biotope classification hierarchy example, Main Array         | 17 |
| Table 4.1: Completed transects, Main Array   | 18 |
| Table 4.2: Completed sediment sampling stations, Main Array                        | 20 |
| Table 4.3: Summary of sediment characteristics, Main Array                         | 26 |
| Table 4.4: Summary of particle size distribution, Main Array                       | 29 |
| Table 4.5: Summary of particle size for multivariate analysis clusters, Main Array | 36 |
| Table 4.6: Summary of sediment aromatic hydrocarbon analysis, Main Array           | 41 |
| Table 4.7: Summary of sediment metals analysis, Main Array                         | 43 |
| Table 4.8: Taxonomic groups, Main Array  | 45 |
| Table 4.9: Macrofaunal community statistics (0.1 m <sup>2</sup> ), Main Array      | 48 |
| Table 4.10: Top ten most abundant macrofaunal taxa within cluster, Main Array      | 57 |
| Table 4.11: Phyletic composition of macrofaunal biomass, Main Array                | 62 |
| Table 4.12: Biomass within each grouping, Main Array                               | 66 |
| Table 4.13: Taxonomic groups of colonial epifauna, Main Array                      | 67 |
| Table 4.14: Number of epifaunal taxa (0.1 m <sup>2</sup> ), Main Array             | 67 |
| Table 4.15: Most frequently occurring colonial epifaunal taxa, Main Array          | 71 |
| Table 4.16: Habitat classifications, Main Array                                    | 72 |
| Table 4.17: Summary of EUNIS habitat classifications, Main Array                   | 76 |
| Table 5.1: Summary of sensitive habitats potentially present, Main Array           | 87 |



## Abbreviations

|         |   |
|---------|---|
| AFDW    | Ash free dry weight                                       |
| AL1/AL2 | Action Level 1 or 2                                       |
| ANOSIM  | Analysis of similarity                                    |
| BGS     | British Geological Survey                                 |
| BSL     | Below sea level   |
| Cefas   | Centre for Environment, Fisheries and Aquaculture Science |
| CEMP    | Coordinated Environmental Monitoring Programme            |
| CM      | Central meridian  |
| CS      | Chemistry sample  |
| DCM     | Dichloromethane   |
| DDV     | Drop-down video   |
| EC      | European Commission                                       |
| ED50    | European Datum 1950                                       |
| EMODnet | European Marine Observation Data Network                  |
| EEA     | European Environment Agency                               |
| EOL     | End of line   |
| EPSG    | European Petroleum Survey Group                           |
| ERL     | Effects range low   |
| EUNIS   | European Nature Information System                        |
| FA      | Faunal sample A   |
| FOCI    | Feature of Conservation Interest                          |
| GC      | Gas chromatography  |
| GC-MS   | Gas chromatography – mass spectrometry                    |
| HC      | Hydrocarbon sample  |
| HM      | Heavy metal sample  |
| ICP-MS  | Inductively coupled plasma-mass spectrometry              |
| ICP-OES | Inductively coupled plasma-optical emission spectrometry  |
| JNCC    | Joint Nature Conservation Committee                       |
| LAT     | Lowest Astronomical Tide                                  |
| MBES    | Multibeam echo sounder                                    |
| MCZ     | Marine Conservation Zone                                  |
| MPA     | Marine Protected Area                                     |
| MRV     | Minimum reporting value                                   |
| MV      | Motor vessel  |
| NF      | No fix  |
| NMBAQC  | National Marine Biological Association Quality Control    |
| nMDS    | Non-metric multi-dimensional scaling                      |
| NPD     | Naphthalene, phenanthrene/anthracene and dibenzothiophene |
| NRW     | Natural Resources Wales                                   |
| NS      | No sample   |
| OSPAR   | Oslo and Paris Commission                                 |
| OWF     | Offshore wind farm  |
| P       | Present   |
| PAH     | Polycyclic aromatic hydrocarbon                           |
| PC      | Physico-chemical sample                                   |
| PCA     | Principle component analysis                              |

|                  |  |
|------------------|--|
| <b>PRIMER</b>    | Plymouth Routines in Multivariate Ecological Research  |
| <b>PSA</b>       | Particle size analysis   |
| <b>PSD</b>       | Particle size distribution   |
| <b>RSD</b>       | Relative standard deviation  |
| <b>SAC</b>       | Special Area of Conservation   |
| <b>SACFOR</b>    | Superabundant, abundant, common, frequent, occasional and rare (semi-quantitative abundance scale) |
| <b>SD</b>        | Standard deviation   |
| <b>SEA</b>       | Strategic Environmental Assessment   |
| <b>SOL</b>       | Start of line  |
| <b>SPA</b>       | Special protected area   |
| <b>SSS</b>       | Side scan sonar  |
| <b>US EPA</b>    | United States Environmental Protection Agency  |
| <b>US EPA 16</b> | United States Environmental Protection Agency's 16 priority PAH pollutants                         |
| <b>UTC</b>       | Coordinated Universal Time   |
| <b>UTM</b>       | Universal Transverse Mercator  |
| <b>WGS84</b>     | World Geodetic System 1984   |
| <b>WoRMS</b>     | World Register of Marine Species   |
| <b>WP</b>        | Waypoint   |

---

# 1. Introduction

## 1.1 General Project Description

On the instruction of Awel y Môr Offshore Windfarm Ltd, Fugro performed a geophysical and environmental site investigation at the proposed Awel y Môr Offshore Wind Farm (OWF) site. The proposed OWF development will consist of the Main Array and three export cable route survey areas; East A & B and West C.

The Awel y Môr OWF is located in the Irish Sea, 10.6 km off the north coast of Wales, to the west of the existing Gwynt y Môr OWF, in water depths ranging between 20 m and 40 m Lowest Astronomical Tide (LAT) and covers an area of approximately 136 km<sup>2</sup>.

Environmental operations were conducted onboard the MV Mersey Guardian during the survey period 16 to 30 August 2020. The environmental site survey included a habitat assessment and a benthic monitoring report. This volume, the benthic monitoring report, details the results of the benthic characterisation survey of the main array.

Appendix A outlines the guidelines for use of this report.

## 1.2 Scope of Work

### 1.2.1 Geophysical Survey

The aim of the geophysical survey was to provide ultra-high resolution seismic, multibeam echosounder (MBES), side scan sonar (SSS), sub-bottom profiler, and magnetometer geophysical data acquisition to characterise the main array and to identify any features of conservation interest.

### 1.2.2 Environmental Survey

The purpose of the environmental survey was to provide a robust baseline characterisation of the site and to supplement the existing benthic ecology data that exists across the area of interest. The survey was also required to establish whether any sensitive habitats are present in the area, specifically habitats listed under Annex I of the EC Habitats Directive such as *Sabellaria* reefs and stony reefs. In addition, grab samples were collected to establish physical and biological properties of the sediment at key locations for characterisation purposes.

The results pertaining to the identification of seabed habitats and biotopes of potential conservation interest can be found in the environmental features report (Volume 2).

## 1.3 Environmental Legislation

Council Directive 92/43/EEC on the Conservation of natural habitats and of wild fauna and flora, commonly known as the EC Habitats Directive, was adopted in 1992, with the main aim of promoting maintenance of biodiversity at a European level. Member states are required to take measures to ensure that protection is afforded to habitats and species of European

importance, and that these features are restored to a 'favourable conservation status'. These habitats and species are listed under Annexes I and II of the Directive and may be selected as features for which Natura 2000 protected areas, namely Special Areas of Conservation (SACs) and Special Protected Areas (SPAs), may be designated. These together contribute to a European ecological network of protected sites. The Annexes include species and habitats identified by the Oslo-Paris commission (OSPAR) and detailed on the OSPAR list of threatened and/or declining species and habitats (OSPAR, 2008). This environmental site investigation was undertaken, in part, to satisfy the Habitats Directive requirement of surveillance for Annex I habitats, based on which SACs may be designated. 'Areas of search' for Annex I habitats have been determined by the Joint Nature Conservation Committee (JNCC) and indicate regions where these habitats are most likely to be encountered; Annex I habitats of relevance for the offshore wind farm industry include subtidal reefs (biogenic, or geogenic) and sandbanks slightly covered by seawater all the time.

This benthic monitoring report assesses the occurrence of relevant OSPAR listed threatened and/or declining species and habitats and EC Habitats Directive Annex I habitats within the survey area.

## 1.4 Regional Habitats, Species and Protected Areas

The main array is located approximately 7 km north-east of the Menai Strait and Conwy Bay Special Area of Conservation (SAC). The SAC covers an area of 265 km<sup>2</sup>, designated for the protection of the subtidal Annex I habitats 'Sandbanks which are slightly covered by sea water all the time' and 'Reefs' (JNCC, 2019a; JNCC 2019b). The main array is located approximately 21.5 km to the north-west of the Dee Estuary SAC, which is designated for the protection of the Annex I habitats 'Mudflats and sandflats not covered by seawater at low tide', '*Salicornia* and other annuals colonising mud and sand' and 'Atlantic salt meadows (*Glauco-Puccinellietalia maritima*)' and 'Reefs'.

The main array is also located approximately 26 km south-west of the Fylde Marine Conservation Zone (MCZ) which covers an area of 260 km<sup>2</sup>, designated for the protection of 'subtidal sediments'.

Based on the European Marine Observation and Data Network (EMODnet) seabed habitats map, the Awel y Môr main array survey area lies in an area likely to comprise a range of European Nature Information System (EUNIS) habitats (EMODnet, 2019), including:

- 'Deep circalittoral mixed sediments' (A5.45);
- 'Deep circalittoral coarse sediment' (A5.15);
- 'Deep circalittoral sand' (A5.27);
- 'Circalittoral fine sand' (A5.25);
- 'Circalittoral muddy sand' (A5.26).

Historic data from a benthic characterisation survey of the Gwynt y Môr OWF (Centre for Marine and Coastal Studies [CMACS], 2005) recorded three Joint Nature Conservation

Committee (JNCC) biotope complexes in the vicinity of the main array survey area. Table 1.1 presents these habitat biotopes along with the equivalent EUNIS classification.


Table 1.1: Gwynt y Môr OWF (CMACS, 2005) habitat classifications, Main Array

| JNCC (2015) Classification  | Equivalent EUNIS Classification (EEA,2019)   |
|---|--|
| SS.SCS.CCS.MedLumVen ( <i>Mediomastus fragilis</i> , <i>Lumbrineris</i> spp. and venerid bivalves in circalittoral coarse sand or gravel) | <i>Mediomastus fragilis</i> , <i>Lumbrineris</i> spp. and venerid bivalves in circalittoral coarse sand or gravel (A5.142) |
| SS.SCS.ICS.MoeVen ( <i>Moerella</i> spp. with venerid bivalves in infralittoral gravelly sand)  | <i>Moerella</i> spp. with venerid bivalves in infralittoral gravelly sand (A5.133)   |
| SS.SSA.IfSa.NcirBat ( <i>Nephtys cirrosa</i> and <i>Bathyporeia</i> spp. in infralittoral sand)   | <i>Nephtys cirrosa</i> and <i>Bathyporeia</i> spp. in infralittoral sand (A5.233)  |

Table 1.2 lists the nearby protected areas relevant to the main array survey area, summarising the sensitive habitats and species for which they were designated to protect. Figure 1.1 spatially displays the protected areas of relevance to the benthic subtidal and intertidal ecology relevant to the main array survey area.

Table 1.2: Summary of nearby protected areas, Main Array

| Protected Area  | Status                       | Distance* [m] | Direction* | Protected Habitats/Species  |
|---|------------------------------|---------------|------------|---|
| Menai Strait and Conwy Bay  | Special Area of Conservation | 3             | SW         | EC Habitats Directive Annex I 'Sandbanks which are slightly covered by sea water all the time', 'Mudflats and sandflats not covered by seawater at low tide' and 'Reefs'  |
| Dee Estuary   | Special Area of Conservation | 21.5          | SE         | EC Habitats Directive Annex I; 'Mudflats and sandflats not covered by seawater at low tide', ' <i>Salicornia</i> and other annuals colonising mud and sand' and 'Atlantic salt meadows ( <i>Glauco-Puccinellietalia maritima</i> )' and 'Reefs' |
| Fylde   | Marine Conservation Zone     | 37            | NE         | Subtidal sands and sediments  |
| <b>Notes</b><br>OSPAR = Oslo and Paris Commission<br>* = Distance and direction from the closest sampling station within the Main Array |                              |               |            |   |



Map Document: (S:\430-MGC-IT\Charting\E200145\_Innogy\_AyM\_Benthic\_Survey\3\_Plots\2\_Draft\Array\Q200145\_01\_Conservation.mxd) 18/01/2021 - 16:44:23

Figure 1.1: Protected areas relevant to the survey area, Main Array



## 1.5 Environmental Quality Standards for Sediment Chemical Concentrations

Selected data have been compared to the OSPAR effects range low (ERL) concentrations (OSPAR, 2014) where available. The ERL thresholds represent the low point (10th percentile) and are therefore indicative of concentrations below which adverse effects rarely occur (OSPAR, 2009; 2014).

The Centre for Environment, Fisheries and Aquaculture Science (Cefas) Guideline Action Levels for the disposal of dredged material are non-statutory guidelines for assessment of disposal of dredged materials to sea, against which reported contaminants concentrations were compared to. In general, concentrations below Cefas Action Level 1 (AL1) are of no concern, whilst concentrations above Action Level 2 (AL2) indicate that dredged material is unsuitable for disposal at sea. Values between AL1 and AL2 may require further investigatory work prior to a disposal decision (Cefas, 2003).

A Strategic Environmental Assessment (SEA) review of the contaminant status of the Irish Sea (SEA6 area) was published in 2005 (Cefas, 2005). Comparative data, specifically for polycyclic aromatic hydrocarbons (PAHs) from site 715 (Liverpool Bay) was utilised due to its spatial proximity to the current survey area. The PAH concentrations from the SEA6 survey are median concentrations of 10 PAHs in sediment samples collected between 1999 and 2002, and, although not directly comparable to total 2 to 6 ring PAH values derived for the current survey, allow the data to be placed into a regional context.

## 1.6 Coordinate Reference System

All coordinates detailed in this report are referenced to World Geodetic System 1984 (WGS84) Universal Transverse Mercator (UTM) projection Zone 30N central meridian 3° West (CM 3° W). Table 1.3 provided the detailed geodetic and projection parameters.

Table 1.3: Project geodetic and projection parameters, Main Array

| Global Positioning System Geodetic Parameters |                                     |
|---|-------------------------------------|
| Datum:  | World Geodetic System 1984 (WGS84)  |
| Spheroid:                                     | World Geodetic System 1984          |
| Semi major axis:                              | a = 6 378 137.000 m                 |
| Reciprocal flattening:                        | 1/f = 298.257 223 563               |
| Project Projection Parameters                 |                                     |
| Grid Projection:                              | Universal Transverse Mercator (UTM) |
| UTM Zone:                                     | 30N (EPSG: 32630)                   |
| Central Meridian:                             | 3° 00' 00" West                     |
| Latitude of Origin:                           | 00° 00' 00" North                   |
| False Easting:                                | 500 000 m                           |
| False Northing:                               | 0 m                                 |
| Scale factor on Central Meridian:             | 0.9996                              |
| Units:  | Metre                               |

## 2. Survey Strategy

Historic data from a benthic characterisation survey of the Gwynt y Môr OWF (CMACS, 2005) and side scan sonar (SSS) data acquired in 2020 were reviewed by Fugro in order to propose environmental survey locations within the proposed Awel y Môr OWF and export cable routes. Particular emphasis was placed on locating areas of potential conservation value (e.g. Annex I listed habitats), on boundaries between areas of differing sonic reflectivity and areas characteristic of the general background conditions of the site.

A total of 62 grab sampling stations were selected to ground-truth the different sediment types observed from the geophysical data, and to also provide spatial coverage across the survey area. Forty-two of the stations were selected following consultation with Natural Resources Wales. At each station, one macrofaunal sample was required, with one particle size distribution (PSD) subsampled from the same sample. A subset of 10 grab stations were selected for additional sampling for chemistry analysis, with second grab samples taken at these stations and subsampled for the required analysis.

At 22 of the stations, drop-down video data were to be acquired along a transect of approximately 50 m in length, centred on each station. At four stations, the orientation and distance of the transect were defined to investigate boundaries between two sediment types. Station MA\_ST25 was orientated to target an area of differing reflectivity crossing station MA\_ST26, station MA\_ST27 was orientated to target an area of higher reflectivity crossing station MA\_ST28, station MA\_ST54 was orientated to target an area of higher reflectivity within an overall area of mobile sediments crossing station MA\_ST53, and station MA\_ST61 was orientated to target a relatively featureless area besides an area of mobile sediments crossing station MA\_ST62.

Tables 2.1 and 2.2 provide the coordinates, data to be acquired and rationale for each proposed survey location. Figure 2.1 spatially displays the proposed survey locations overlaid on the SSS mosaic.

Table 2.1: Proposed sampling stations, Main Array

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |           |             |   |                                  |
|---|-----------|-------------|---|----------------------------------|
| Station   | Easting   | Northing    | Rationale   | Data and Sample Acquisition      |
| MA_ST01   | 432 960.8 | 5 927 193.8 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST02   | 432 777.0 | 5 926 144.4 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST03   | 435 317.0 | 5 927 335.0 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST04   | 434 714.0 | 5 926 692.0 | Station in the western extent of the main array, in a relatively featureless area | Video and stills<br>FA, PSD, CS, |
| MA_ST05   | 434 205.7 | 5 926 049.2 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST06   | 432 899.8 | 5 925 014.1 | Station selected for spatial coverage*  | FA, PSD                          |



| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |           |             |   |                                 |
|---|-----------|-------------|---|---------------------------------|
| Station   | Easting   | Northing    | Rationale   | Data and Sample Acquisition     |
| MA_ST07   | 436 777.5 | 5 926 461.9 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST08   | 435 698.0 | 5 925 604.7 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST09   | 434 372.4 | 5 924 396.8 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST10   | 439 327.0 | 5 927 350.9 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST11   | 438 522.0 | 5 926 615.0 | Station in the western extent of the main array, in a relatively featureless area. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD     |
| MA_ST12   | 437 529.0 | 5 925 669.0 | Station in the western extent of the main array, in a localised area of variable reflectivity   | Video and stills<br>FA, PSD, CS |
| MA_ST13   | 437 110.2 | 5 924 842.5 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST14   | 436 240.4 | 5 923 662.6 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST15   | 441 200.3 | 5 927 398.5 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST16   | 440 508.1 | 5 926 557.2 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST17   | 439 285.8 | 5 925 699.9 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST18   | 443 025.9 | 5 927 366.8 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST19   | 442 461.0 | 5 926 594.0 | Station in the western extent of the main array, in a relatively featureless area. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD     |
| MA_ST20   | 440 800.8 | 5 925 686.4 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST21   | 439 539.8 | 5 924 525.2 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST22   | 438 596.0 | 5 923 566.3 | Station selected for spatial coverage   | Video and stills<br>FA, PSD, CS |
| MA_ST23   | 441 118.0 | 5 925 220.2 | Station selected for spatial coverage   | Video and stills<br>FA, PSD     |
| MA_ST24   | 438 799.7 | 5 923 486.3 | Station selected for spatial coverage   | FA, PSD                         |
| MA_ST25   | 441 223.0 | 5 925 039.0 | Station in the western extent of the main array. In an area of differing reflectivity. The orientation of the drop-down video was predetermined (MA_ST26)   | Video and stills<br>FA, PSD, CS |
| MA_ST27   | 439 295.0 | 5 923 381.0 | Station in the western extent of the main array. In an area of higher reflectivity to the east of an area of mobile sediments. The orientation of the drop-down video was predetermined (MA_ST28) | Video and stills<br>FA, PSD     |
| MA_ST29   | 444 661.0 | 5 927 335.0 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST30   | 442 549.7 | 5 925 731.7 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST31   | 441 437.6 | 5 924 729.1 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST32   | 438 674.6 | 5 922 704.8 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST33   | 443 994.3 | 5 925 763.4 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST34   | 442 470.3 | 5 924 795.0 | Station selected for spatial coverage*  | FA, PSD                         |
| MA_ST35   | 441 706.4 | 5 924 346.0 | Station selected for spatial coverage*  | FA, PSD                         |

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |           |             |   |                                  |
|---|-----------|-------------|---|----------------------------------|
| Station   | Easting   | Northing    | Rationale   | Data and Sample Acquisition      |
| MA_ST36   | 446 474.6 | 5 927 357.1 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST37   | 445 413.5 | 5 926 398.4 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST38   | 441 672.7 | 5 923 361.5 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST39   | 444 076.2 | 5 924 638.8 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST40   | 443 343.9 | 5 923 890.9 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST41   | 441 095.5 | 5 921 805.2 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST42   | 448 252.6 | 5 927 341.2 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST43   | 447 038.0 | 5 926 464.0 | Station in the middle of the main array, in a relatively featureless area. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD, CS, |
| MA_ST44   | 445 277.0 | 5 924 515.0 | Station in the middle of the array, in a relatively featureless area between mobile sediments to the east and west  | Video and stills<br>FA, PSD      |
| MA_ST45   | 443 807.8 | 5 922 836.9 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST46   | 443 175.0 | 5 922 829.0 | Station in the middle of the main array, in an area of mobile sediments. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD      |
| MA_ST47   | 445 090.2 | 5 923 234.7 | Station selected for spatial coverage*  | FA, PSD, CS                      |
| MA_ST48   | 443 918.9 | 5 922 130.4 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST49   | 442 894.7 | 5 921 292.0 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST50   | 446 010.4 | 5 923 271.0 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST51   | 445 486.5 | 5 922 318.5 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST52   | 444 625.1 | 5 920 709.9 | Station selected for spatial coverage*  | FA, PSD                          |
| MA_ST54   | 447 949.0 | 5 923 456.0 | Station in the eastern extent of the main array. In an area of higher reflectivity within an overall area of mobile sediments. The orientation of the drop-down video was predetermined (MA_ST53) | Video and stills<br>FA, PSD      |
| MA_ST55   | 447 905.0 | 5 923 406.9 | Station selected for spatial coverage   | Video and stills<br>FA, PSD      |
| MA_ST56   | 448 166.2 | 5 923 311.0 | Station selected for spatial coverage   | Video and stills<br>FA, PSD      |
| MA_ST57   | 445 929.0 | 5 921 227.0 | Station in the eastern extent of the main array, in an area of mobile sediments. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD      |
| MA_ST58   | 452 286.0 | 5 926 081.0 | Station in the eastern extent of the main array, in a relatively featureless area. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD      |
| MA_ST59   | 450 387.0 | 5 924 424.0 | Station in the eastern extent of the main array, in an area of mobile sediments. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD, CS  |
| MA_ST60   | 446 905.8 | 5 919 910.8 | Station selected for spatial coverage*  | FA, PSD                          |

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m]   |           |             |  |                                 |
|---|-----------|-------------|--|---------------------------------|
| Station   | Easting   | Northing    | Rationale  | Data and Sample Acquisition     |
| MA_ST61   | 456 206.0 | 5 925 508.0 | Station in the eastern extents of the main array, in a relatively featureless area besides an area of mobile sediments. The orientation of the drop-down video was predetermined (MA_ST62) | Video and stills<br>FA, PSD, CS |
| MA_ST63   | 456 177.3 | 5 925 488.6 | Station selected for spatial coverage*   | Video and stills<br>FA, PSD     |
| MA_ST64   | 456 206.7 | 5 925 480.6 | Station selected for spatial coverage*   | Video and stills<br>FA, PSD     |
| MA_ST65   | 453 569.0 | 5 923 291.0 | Station in the eastern extents of the main array, in an area of mobile sediments. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD, CS |
| MA_ST66   | 450 568.0 | 5 920 636.0 | Station in the eastern extents of the main array, in an area of mobile sediments. Representative sediments, selected for spatial coverage  | Video and stills<br>FA, PSD, CS |
| <b>Notes</b><br>FA = Faunal sample<br>PSD = Particle size distribution subsample<br>CS = Chemistry sample<br>* = Stations selected by Natural Resources Wales |           |             |  |                                 |

Table 2.2: Proposed transects, Main Array

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m]      |     |           |             |  |                  |
|--|-----|-----------|-------------|--|------------------|
| Station  |     | Easting   | Northing    | Rationale  | Data Acquisition |
| MA_ST26  | SOL | 441 208.0 | 5 925 055.0 | Transect across a boundary of differing reflectivity                                       | Video, stills    |
|  | EOL | 441 235.0 | 5 925 021.0 |  |                  |
| MA_ST28  | SOL | 439 234.0 | 5 923 384.0 | Transect across a boundary of two sediment types (mobile sediments, and high reflectivity) | Video, stills    |
|  | EOL | 439 317.0 | 5 923 381.0 |  |                  |
| MA_ST53  | SOL | 447 911.0 | 5 923 474.0 | Transect across and area of high reflectivity and adjacent area of mobile sediments        | Video, stills    |
|  | EOL | 447 958.0 | 5 923 450.0 |  |                  |
| MA_ST62  | SOL | 456 168.0 | 5 925 503.0 | Transect across and area of high reflectivity and adjacent area of mobile sediments        | Video, stills    |
|  | EOL | 456 234.0 | 5 925 510.0 |  |                  |
| <b>Notes</b><br>SOL = Start of line<br>EOL = End of line |     |           |             |  |                  |




Figure 2.1: Proposed environmental survey locations, Main Array

---

## 3. Methods

### 3.1 Survey Methods

#### 3.1.1 Seabed Photography

Seabed photography was acquired using a ROV Tech subsea camera system mounted within a purpose-built camera frame, which included a pressure housing filled with freshwater. The purpose of the freshwater housing was to provide a column of clear water between the seabed and the camera to facilitate the capture of good quality stills in highly turbid waters. The system consisted of one high resolution video camera, a still image camera, and two high intensity LED lighting strips at the side of the frame. Two lasers were set up 10 cm apart to provide a scale.

Seabed video was displayed on a computer monitor and recorded directly onto the computer hard drive. A video overlay was used to overlay a navigation string from an online positioning system using a vessel offset for position, including the time, date and location. The survey location and station number were also displayed (manually updated). The stills camera imagery was visible on a second window of the computer. Video was viewed in real time via an umbilical, assisting in the control of the camera in the water.

#### 3.1.2 Sediment Sampling

Macrofaunal and PSD grab samples were collected using a 0.1 m<sup>2</sup> Hamon grab. Chemistry samples were acquired using a 0.1 m<sup>2</sup> Day grab. Manual position fixes were recorded when the grab reached the seabed using an offset from the vessel reference point.

Further details on survey methodology are available within Appendix B.

### 3.2 Analytical Methods

Brief analytical methodologies are described in the following subsections. Further descriptions of the analytical methodologies are detailed in Appendix B.

#### 3.2.1 Sediment Characteristics

Sediment samples were analysed for their PSD using a combination of two techniques; sieve analysis for all material retained by a 1.0 mm sieve followed by laser diffraction analysis of the finer material. The PSD parameters include the descriptive statistics derived in Gradistat (Blott, 2010), based on the Folk & Ward (1957) method. The sediment descriptions are based on the Wentworth (1922) scale and the British Geological Survey (BGS) modified Folk classification (Long, 2006).

#### 3.2.2 Sediment Hydrocarbons

The sediment samples were analysed for PAHs, specifically the United States Environmental Protection Agency's 16 priority PAH pollutants (US EPA 16 PAHs) and alkylated PAHs.

Samples were extracted by ultrasonication of wet sediments with mixed solvents. The sample extracts were then cleaned-up using absorption column chromatography. Aromatic hydrocarbons were analysed by gas chromatography-mass spectrometry (GC-MS).

The distributions and concentrations of 2 to 6 ring PAHs within the samples were determined by GC-MS. Standard solutions containing an appropriate range and concentration of aromatic hydrocarbons were run to calibrate the instrument and acquire response factors for quantification purposes. Individual aromatic compounds were quantified using a series of deuterated internal standards. The total 2 to 6 ring PAH values for each sediment sample was calculated by summing the concentrations of individual aromatic compounds.

### 3.2.3 Sediment Metals

The sediment samples were analysed using an aqua regia digest technique. This provides a strong partial digest, releasing into solution metals associated with the fines fraction within the sediments (but does not extract all trace elements associated with the coarse fraction). The concentrations of metals released by an aqua regia digest are typically considered indicative of those influencing biological interactions, as the released metals are not incorporated into the mineral matrix and are therefore potentially available for biological uptake.

The sediment samples underwent an aqua regia digest followed by multi-element analysis by inductively coupled plasma-mass spectrometry (ICP-MS) (arsenic, cadmium, chromium, copper, lead, mercury, nickel, tin and zinc) or by inductively coupled plasma-optical emission spectrometry (ICP-OES) (aluminium and barium).

### 3.2.4 Sediment Macrofauna

Macrofaunal samples were processed on a sieve mesh of 1 mm, with retained taxa identified and enumerated using a combination of stereo and compound microscopy.

Biomass (phylum level) was determined for infaunal invertebrates from grab samples; biomass was not determined for epifauna.

## 3.3 Data Analysis

Summary statistics (minimum, maximum, mean, standard deviation) for all reported datasets were derived in Excel.

### 3.3.1 Sediment Particle Size Distribution (PSD) Statistics

Table 3.1 summarises the sediment PSD statistics that were calculated using Gradistat V8 (Blott, 2010).

Table 3.1: Sediment particle size distribution statistics, Main Array

| Statistic | Definition  |
|-----------|---|
| Mean      | A measure of central tendency: sum of values, divided by number of observations |
| Median    | A measure of central tendency: central value                                    |
| Mode      | A measure of central tendency: most frequently observed value                   |
| Modality  | A measure of the number of peaks in the frequency distribution                  |
| Sorting   | A measure of the grain size range and magnitude of their spread around the mean |
| Skewness  | A measure of the degree of symmetry   |

### 3.3.2 Sediment Macrofauna

#### 3.3.2.1 Data Rationalisation

Prior to analysis, the macrofaunal dataset was rationalised. Damaged taxa were removed, and indeterminable taxa were merged to avoid spurious enhancement of the species list. Juvenile species were removed, as they represent an ephemeral stage and are, therefore, not considered part of the permanent benthic community. Meiofauna, fish, algae, eggs and foraminifera were also removed for this reason. Sessile colonial epifauna was recorded as present, and assessed separately from the enumerated fauna, which comprised infaunal and solitary epifaunal taxa.

#### 3.3.2.2 Univariate Analysis

Table 3.2 summarises the univariate statistics derived from Plymouth Routines in Multivariate Ecological Research (PRIMER) version 7 (v7).

Table 3.2: Macrofaunal univariate statistics, Main Array

| Statistic   | Definition   |
|---|--|
| Number of taxa (S)  | Count of taxa  |
| Abundance (N)   | Count of individuals   |
| Shannon-Wiener index of diversity ( $H' \log_2$ )             | <p>A measure of the number of taxa in a sample and the distribution of abundance across these taxa; results were assessed in line with the threshold values in Dauvin et al. (2012):</p> <ul style="list-style-type: none"> <li>■ High diversity (<math>H' \log_2 &gt; 4.00</math>);</li> <li>■ Good diversity (<math>3.00 &lt; H' \log_2 &lt; 4.00</math>);</li> <li>■ Moderate diversity (<math>2.00 &lt; H' \log_2 &lt; 3.00</math>);</li> <li>■ Poor diversity (<math>1.00 &lt; H' \log_2 &lt; 2.00</math>);</li> <li>■ Bad diversity (<math>H' \log_2 &lt; 1.00</math>).</li> </ul> |
| Pielou's index of evenness (J)                                | A measure of how evenly distributed the individuals are among the different species  |
| Complement of the Simpsons index of dominance ( $1-\lambda$ ) | A measure of evenness in which its highest value corresponds to assemblages with individuals evenly distributed across taxa, whilst its lowest value corresponds to those in which the total abundance is dominated by one or very few of the taxa present   |



### 3.3.2.3 Multivariate Analysis

Various multivariate statistical techniques were applied to the macrofauna and sediment characteristics data to investigate patterns of similarity in PRIMER v7. These included:

- Data transformation to reduce skewness of data, for optimal performance of multivariate analysis (square root for macrofaunal, fourth root for PSD);
- Hierarchical clustering analysis (subsequently referred to as 'Cluster' analysis) to group samples based on the nearest neighbour sorting of a matrix of sample similarities using Bray Curtis similarity (for biological datasets) or Euclidean distance measure (for environmental datasets);
- Non-metric multidimensional scaling (nMDS) ordination of Bray Curtis and Euclidean Distance similarity/distance matrices;
- SIMPROF algorithm was used to identify statistically significant clusters; in ecological terms the statistical relevance of similarity profile testing is assessed in line with the recommendation of Clarke et al. (2008), thus defining coarser grouping can be appropriate if the resulting groups are supersets of the similarity profile clusters;
- Similarity percentage analysis ('Simper' algorithm), to gauge the distinctiveness of each of the multivariate groups;
- ANOSIM algorithm was used to investigate significant differences between a priori defined groups;
- Principal component analysis (PCA), to identify spatial patterns and relationships between variables;
- BIOENV algorithm was used to indicate relationships between physical and biological variables.

### 3.3.2.4 Biomass Analysis

Biomass was assessed for infaunal invertebrates from grab samples; epifauna was not biomassed.

The macrofaunal blotted wet weight biomass dataset was converted to ash free dry weight (AFDW) by applying the appropriate standard corrections, as outlined in Eleftheriou and Basford (1989). Table 3.3 summarises the corrections applied.

Table 3.3: Macrofaunal standard biomass corrections by phyla, Main Array

| Phyla   | Standard Biomass Correction [%] |
|---|---------------------------------|
| Annelida  | 15.5                            |
| Arthropoda  | 22.5                            |
| Mollusca  | 8.5                             |
| Echinodermata   | 8.0                             |
| Other Taxa  | 15.5                            |
| Notes<br>Standard biomass corrections to convert blotted wet weight to ash free dry weight, from Eleftheriou & Basford (1989) |                                 |



### 3.3.3 Seabed Habitats and Biotopes

A habitat assessment was completed by Fugro detailing the analysis of completed transects (Volume 2). This will be summarised and refined with consideration of grab sample analysis.

#### 3.3.3.1 Seabed Habitat Classification

To assess the habitats present within the survey area, detailed analysis of video and still photographic data was undertaken, noting the locations of any observed changes in sediment type and/or associated faunal community.

Taxa were recorded to the lowest possible taxonomic level. It should be noted that many species cannot be identified from photographic data alone and, as such, higher taxonomic levels were used.

Descriptions of the substrate composition, corresponding to sediment changes, were undertaken to support the EUNIS habitat identification (Long, 2006). These descriptions were largely based on a reclassification of the Folk (1954) sediment classes, with the Wentworth (1922) classification also considered as the latter differentiates between pebbles, cobbles and boulders. The Folk (1954) sediment classification was reclassified into four categories, namely 'coarse sediment', 'mixed sediment', 'mud and sandy mud' and 'sand and muddy sand' (Long, 2006). Further sub-categories, namely 'mud', 'sandy mud' and 'muddy sand' are utilised to further account for differences in sediment in the 'mud to sandy mud' fraction (Kaskela et al., 2019). These categories are defined by the proportions of mud (the 'fines' fraction), sand and gravel. For example, a description of 'muddy sand' defines sediments that have sand as the principle component (50 % to 90 %) with a secondary component of mud (10 % to 50 %) and < 5 % gravel (Kaskela et al., 2019). The European Marine Observation and Data Network (EMODnet) Geology Consortium further revised these categories to include an additional category 'Rock and Boulders' (Kaskela et al., 2019), which includes the Wentworth (1922) categories 'boulders' and 'cobbles'. The presence of shell fragments and evident anthropogenic features were also noted.

Table 3.4 presents a summary of the sediment particle sizes and corresponding classifications.

Table 3.4: Sediment particle size and classification terms, Main Array

| Particle Size   | Wentworth (1922) | Folk (1954) | Folk, 5 classes (Kaskela et al., 2019)                                     |  |   |   |
|---|------------------|-------------|--|--|---|---|
| > 256 mm  | Boulder          | Gravel      | Rock & Boulders  |  |   |   |
| 64 mm to 256 mm   | Cobble           |             |  |  |   |   |
| 32 mm to < 64 mm  | Pebbles          |             | Coarse sediment:<br>(Gravel ≥ 80 %, or<br>Gravel ≥ 5 % and<br>Sand ≥ 90 %) | Mixed sediment:<br>(Mud ≥ 10 % - 95 %<br>Sand < 90 %<br>Gravel ≥ 5%) | Mud to sandy mud*:<br>(Mud 10 % -100 %<br>Sand < 90 %<br>Gravel < 5%) | Sand:<br>(Mud < 10 %<br>Sand ≥ 90 %<br>Gravel < 5%) |
| 16 mm to < 32 mm  |                  |             |  |  |   |   |
| 8 mm to < 16 mm   |                  |             |  |  |   |   |
| 4 mm to < 8 mm  |                  |             |  |  |   |   |
| 2 mm to < 4 mm  | Granules         |             |  |  |   |   |
| 1 mm to < 2 mm  | Very coarse sand | Sand        |  |  |   |   |
| 0.5 mm to < 1 mm  | Coarse sand      |             |  |  |   |   |
| 0.25 mm to < 0.5 mm   | Medium sand      |             |  |  |   |   |
| 0.125 mm to < 0.25 mm   | Fine sand        |             |  |  |   |   |
| 62.5 µm to 0.125 mm   | Very fine sand   |             |  |  |   |   |
| > 4 µm to 62.5 µm   | Silt             | Mud         | -  |  |   |   |
| > 1 µm to 4 µm  | Clay             |             |  |  |   |   |
| <b>Notes</b><br>* = Mud to sandy mud includes:<br>Mud (Mud ≥ 90 %, Sand <10 %, Gravel < 5%);<br>Sandy mud (Mud 50 % to 90 %, Sand 10 % to 50 %, Gravel < 5%);<br>Muddy sand (Mud 10 % to 50 %, Sand 50 % to 90 %, Gravel < 5%) (Kaskela et al., 2019) |                  |             |  |  |   |   |

Habitats within the survey area have been classified in accordance with the hierarchical EUNIS habitat classification (EEA, 2019), which has compiled habitat information from across Europe into a single database. Table 3.5 summarises the EUNIS hierarchy, with an example of the coding system. The equivalent classification from 'The Marine Habitat Classification for Britain and Ireland – Version 15.03' (JNCC, 2015) was also noted. The JNCC classification formed the basis of the marine section of the EUNIS habitat classification scheme (Davies & Moss, 2004), resulting in broad similarities, although there are some structural differences and habitat types. These classification systems are designed to incorporate small-scale temporal variations (e.g. seasonal) into the biotope/habitat categories. However, biological communities and marine environments can be highly dynamic and temporally variable, therefore the biotopes and habitats identified by the current assessment are representative of the survey area at the time of sampling only.

Table 3.5: EEA (2019) biotope classification hierarchy example, Main Array

| Level                            | Example Classification Name   | Example Classification Code |
|----------------------------------|---|-----------------------------|
| 1. Environment                   | Marine habitats   | A                           |
| 2. Broad habitat types           | Sublittoral sediments   | A5                          |
| 3. Main habitats                 | Sublittoral sand  | A5.2                        |
| 4. Biotope complexes             | Circolittoral muddy sand  | A5.26                       |
| 5 & 6. Biotopes and sub-biotopes | <i>Amphiura brachiata</i> with <i>Astropecten irregularis</i> and other echinoderms in circolittoral muddy sand | A5.262                      |

Classifications were assigned to each habitat type observed within the video and stills photography. Additional information from grab sampling, such as sediment particle size and macrofaunal communities, was used where applicable. Although, theoretically, a biotope can be assigned to any sized area of seabed, for the purposes of this assessment the commonly accepted minimum habitat size of 25 m<sup>2</sup> was adopted. For distinct areas of mixed habitats/biotopes (e.g. rock interspersed with coarse sediment) where the overall area was at least 25 m<sup>2</sup>, biotope mosaics were considered (Parry, 2019).

## 4. Results

### 4.1 Field Operations

#### 4.1.1 Bathymetry and Seabed Features

The following is summarised from the seafloor and shallow geophysical results report (Fugro, 2020). The water depths across the Main Array survey area ranged from 15.2 m LAT to 41.9 m LAT, generally increasing towards the north-west. The seafloor in the south-east was characterised by numerous sand waves and mega-ripples, whilst the west of the site was relatively flat and featureless. Lower SSS reflectivity areas in the east of the site were classified as sand with areas of sand waves, whilst the more homogeneous, medium reflectivity in the west were classified as gravelly sand.

During the survey operations it was observed that the sand waves were actively mobile and were migrating significantly in the time between adjacent survey lines.

#### 4.1.2 Seabed Photography

Photographic stills and video were successfully acquired at all 22 proposed stations in the main array. Station MA\_ST64 was repeated due to loss of the navigational overlay string. At stations MA\_ST46, MA\_ST57 and MA\_ST65, the tide caused the vessel to change heading during data acquisition. Waypoints indicating the point of direction change have therefore been included (Table 4.1).

Table 4.1: Completed transects, Main Array

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |     |           |             |               |            |                  |
|---|-----|-----------|-------------|---------------|------------|------------------|
| Station   |     | Easting   | Northing    | Depth [m BSL] | Length [m] | Data Acquisition |
| MA_ST04   | SOL | 434 692.7 | 5 926 680.4 | 44            | 39         | 2 min 7 sec      |
|   | EOL | 434 712.7 | 5 926 713.7 | 44            |            | 5 stills         |
| MA_ST11   | SOL | 438 495.6 | 5 926 605.4 | 42            | 54         | 6 min 4 sec      |
|   | EOL | 438 540.9 | 5 926 634.2 | 42            |            | 5 stills         |
| MA_ST12   | SOL | 437 501.2 | 5 925 669.5 | 38            | 63         | 6 min 49 sec     |
|   | EOL | 437 563.2 | 5 925 661.0 | 38            |            | 6 stills         |
| MA_ST19   | SOL | 442 434.2 | 5 926 597.3 | 39            | 51         | 8 min 32 sec     |
|   | EOL | 442 485.6 | 5 926 598.1 | 39            |            | 5 stills         |
| MA_ST22   | SOL | 438 618.0 | 5 923 551.4 | 36            | 51         | 6 min 8 sec      |
|   | EOL | 438 590.7 | 5 923 594.2 | 36            |            | 5 stills         |
| MA_ST23   | SOL | 441 092.9 | 5 925 229.2 | 36            | 50         | 4 min 59 sec     |
|   | EOL | 441 141.0 | 5 925 215.1 | 36            |            | 6 stills         |

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |     |           |             |               |            |                  |
|---|-----|-----------|-------------|---------------|------------|------------------|
| Station   |     | Easting   | Northing    | Depth [m BSL] | Length [m] | Data Acquisition |
| MA_ST26   | SOL | 441 196.7 | 5 925 054.8 | 37            | 61         | 5 min 0 sec      |
|   | EOL | 441 239.0 | 5 925 011.4 | 37            |            | 5 stills         |
| MA_ST28   | SOL | 439 205.4 | 5 923 380.7 | 39            | 84         | 15 min 45 sec    |
|   | EOL | 439 289.4 | 5 923 381.1 | 39            |            | 12 stills        |
| MA_ST43   | SOL | 447 059.3 | 5 926 476.0 | 35            | 44         | 5 min 43 sec     |
|   | EOL | 447 020.7 | 5 926 455.4 | 35            |            | 5 stills         |
| MA_ST44   | SOL | 445 298.2 | 5 924 530.8 | 32            | 54         | 6 min 35 sec     |
|   | EOL | 445 252.4 | 5 924 502.9 | 32            |            | 6 stills         |
| MA_ST46   | SOL | 443 152.3 | 5 922 810.2 | 30            | 84*        | 10 min 8 sec     |
|   | WP  | 443 172.8 | 5 922 844.0 | 30            |            | 5 stills         |
|   | EOL | 443 211.0 | 5 922 833.6 | 30            |            |                  |
| MA_ST53   | SOL | 447 909.5 | 5 923 476.5 | 29            | 65         | 4 min 42 sec     |
|   | EOL | 447 966.8 | 5 923 446.4 | 29            |            | 8 stills         |
| MA_ST55   | SOL | 447 878.7 | 5 923 416.6 | 28            | 53         | 4 min 35 sec     |
|   | EOL | 447 927.3 | 5 923 395.7 | 28            |            | 5 stills         |
| MA_ST56   | SOL | 448 144.9 | 5 923 327.0 | 31            | 53         | 4 min 22 sec     |
|   | EOL | 448 186.1 | 5 923 293.0 | 31            |            | 5 stills         |
| MA_ST57   | SOL | 445 939.0 | 5 921 251.2 | 28            | 70*        | 8 min 46 sec     |
|   | WP  | 445 932.8 | 5 921 249.4 | 28            |            | 6 stills         |
|   | EOL | 445 947.5 | 5 921 215.3 | 28            |            |                  |
| MA_ST58   | SOL | 452 258.9 | 5 926 079.5 | 35            | 47         | 4 min 51 sec     |
|   | EOL | 452 300.6 | 5 926 101.8 | 35            |            | 5 stills         |
| MA_ST59   | SOL | 450 361.3 | 5 924 431.9 | 35            | 46         | 5 min 39 sec     |
|   | EOL | 450 406.8 | 5 924 439.7 | 35            |            | 5 stills         |
| MA_ST62   | SOL | 456 142.3 | 5 925 497.9 | 31            | 95         | 6 min 43 sec     |
|   | EOL | 456 235.3 | 5 925 518.8 | 31            |            | 7 stills         |
| MA_ST63   | SOL | 456 148.4 | 5 925 482.0 | 31            | 45         | 3 min 6 sec      |
|   | EOL | 456 181.3 | 5 925 512.8 | 31            |            | 6 stills         |

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m]   |     |           |             |               |            |                          |
|---|-----|-----------|-------------|---------------|------------|--------------------------|
| Station   |     | Easting   | Northing    | Depth [m BSL] | Length [m] | Data Acquisition         |
| MA_ST64   | SOL | 456 125.1 | 5 925 474.3 | 31            | 96         | 8 min 0 sec              |
|   | EOL | 456 218.7 | 5 925 495.6 | 31            |            | 6 stills                 |
| MA_ST64(2)  | SOL | 456 179.2 | 5 925 492.0 | 28            | 47         | 6 min 11 sec             |
|   | EOL | 456 214.7 | 5 925 461.6 | 28            |            | 6 stills                 |
| MA_ST65   | SOL | 453 543.5 | 5 923 290.3 | 31            | 94*        | 7 min 31 sec<br>5 stills |
|   | WP  | 453 592.8 | 5 923 291.5 | 31            |            |                          |
|   | EOL | 453 552.5 | 5 923 275.5 | 31            |            |                          |
| MA_ST66   | SOL | 450 540.3 | 5 920 638.3 | 24            | 54         | 3 min 58 sec             |
|   | EOL | 450 594.3 | 5 920 635.3 | 24            |            | 5 stills                 |
| <b>Notes</b><br>BSL = Below sea level<br>SOL = Start of line<br>EOL = End of line<br>WP = Waypoint<br>* = Direction of the transect changed during data acquisition, the total distance is therefore calculated as the distance travelled rather than the distance between the recorded SOL and EOL |     |           |             |               |            |                          |

### 4.1.3 Seabed Sampling

Grab samples were successfully acquired at all proposed stations in the main array survey area. A complete suite of grab samples (one macrofauna and one PSD subsample) was retained at all 62 station (Table 4.2). A complete suite of chemical samples was retained at all 10 stations.

Table 4.2: Completed sediment sampling stations, Main Array

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |           |             |               |                    |
|---|-----------|-------------|---------------|--------------------|
| Station   | Easting   | Northing    | Depth [m BSL] | Sample Acquisition |
| MA_ST01   | 432 971.4 | 5 927 193.6 | 39            | FA, PSD            |
| MA_ST02   | 432 775.0 | 5 926 151.4 | 39            | FA, PSD            |
| MA_ST03   | 435 309.0 | 5 927 335.5 | 40            | FA, PSD            |
| MA_ST04   | 434 709.3 | 5 926 693.1 | 44            | FA, PSD, CS        |
| MA_ST05   | 434 210.4 | 5 926 047.8 | 39            | FA, PSD            |
| MA_ST06   | 432 894.4 | 5 925 022.5 | 37            | FA, PSD            |
| MA_ST07   | 436 761.7 | 5 926 443.9 | 44            | FA, PSD            |
| MA_ST08   | 435 704.2 | 5 925 604.3 | 38            | FA, PSD            |
| MA_ST09   | 434 384.2 | 5 924 402.5 | 35            | FA, PSD            |
| MA_ST10   | 439 328.6 | 5 927 348.2 | 43            | FA, PSD            |
| MA_ST11   | 438 525.0 | 5 926 611.7 | 42            | FA, PSD            |

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m] |           |             |               |                    |
|---|-----------|-------------|---------------|--------------------|
| Station   | Easting   | Northing    | Depth [m BSL] | Sample Acquisition |
| MA_ST12   | 437 531.5 | 5 925 677.0 | 38            | FA, PSD, CS        |
| MA_ST13   | 437 115.3 | 5 924 840.8 | 34            | FA, PSD            |
| MA_ST14   | 436 231.6 | 5 923 656.5 | 33            | FA, PSD            |
| MA_ST15   | 441 199.1 | 5 927 398.4 | 41            | FA, PSD            |
| MA_ST16   | 440 503.9 | 5 926 558.5 | 42            | FA, PSD            |
| MA_ST17   | 439 276.5 | 5 925 698.9 | 39            | FA, PSD            |
| MA_ST18   | 443 028.6 | 5 927 370.2 | 40            | FA, PSD            |
| MA_ST19   | 442 464.1 | 5 926 593.9 | 39            | FA, PSD            |
| MA_ST20   | 440 802.3 | 5 925 688.1 | 39            | FA, PSD            |
| MA_ST21   | 439 538.7 | 5 924 525.3 | 38            | FA, PSD            |
| MA_ST22   | 438 601.6 | 5 923 569.0 | 31            | FA, PSD, CS        |
| MA_ST23   | 441 122.5 | 5 925 222.9 | 36            | FA, PSD            |
| MA_ST24   | 438 792.6 | 5 923 498.0 | 32            | FA, PSD            |
| MA_ST25   | 441 235.7 | 5 925 039.3 | 38            | FA, PSD, CS        |
| MA_ST27   | 439 295.6 | 5 923 375.7 | 33            | FA, PSD            |
| MA_ST29   | 444 651.1 | 5 927 328.6 | 38            | FA, PSD            |
| MA_ST30   | 442 557.9 | 5 925 737.6 | 37            | FA, PSD            |
| MA_ST31   | 441 444.1 | 5 924 722.4 | 37            | FA, PSD            |
| MA_ST32   | 438 696.3 | 5 922 699.7 | 33            | FA, PSD            |
| MA_ST33   | 443 981.8 | 5 925 760.6 | 35            | FA, PSD            |
| MA_ST34   | 442 476.2 | 5 924 807.3 | 36            | FA, PSD            |
| MA_ST35   | 441 711.3 | 5 924 338.3 | 34            | FA, PSD            |
| MA_ST36   | 446 463.2 | 5 927 358.1 | 38            | FA, PSD            |
| MA_ST37   | 445 411.2 | 5 926 388.1 | 37            | FA, PSD            |
| MA_ST38   | 441 669.4 | 5 923 359.0 | 31            | FA, PSD            |
| MA_ST39   | 444 067.4 | 5 924 629.5 | 32            | FA, PSD            |
| MA_ST40   | 443 341.1 | 5 923 891.7 | 30            | FA, PSD            |
| MA_ST41   | 441 094.4 | 5 921 801.5 | 29            | FA, PSD            |
| MA_ST42   | 448 244.0 | 5 927 344.8 | 39            | FA, PSD            |
| MA_ST43   | 447 026.4 | 5 926 473.2 | 37            | FA, PSD, CS        |
| MA_ST44   | 445 272.5 | 5 924 509.3 | 32            | FA, PSD            |
| MA_ST45   | 443 817.3 | 5 922 832.6 | 29            | FA, PSD            |
| MA_ST46   | 443 177.1 | 5 922 838.5 | 30            | FA, PSD            |
| MA_ST47   | 445 086.7 | 5 923 234.0 | 31            | FA, PSD, CS        |
| MA_ST48   | 443 915.1 | 5 922 128.3 | 31            | FA, PSD            |

| Geodetic Parameters: ED50, UTM Zone 31N, CM 3°E [m]  |           |             |               |                    |
|--|-----------|-------------|---------------|--------------------|
| Station  | Easting   | Northing    | Depth [m BSL] | Sample Acquisition |
| MA_ST49  | 442 896.1 | 5 921 283.6 | 27            | FA, PSD            |
| MA_ST50  | 446 014.0 | 5 923 286.1 | 36            | FA, PSD            |
| MA_ST51  | 445 493.3 | 5 922 320.7 | 34            | FA, PSD            |
| MA_ST52  | 444 634.0 | 5 920 717.6 | 27            | FA, PSD            |
| MA_ST54  | 447 950.6 | 5 923 468.1 | 36            | FA, PSD            |
| MA_ST55  | 447 900.6 | 5 923 413.6 | 36            | FA, PSD            |
| MA_ST56  | 448 180.1 | 5 923 328.4 | 37            | FA, PSD            |
| MA_ST57  | 445 936.3 | 5 921 240.4 | 32            | FA, PSD            |
| MA_ST58  | 452 287.9 | 5 926 084.4 | 38            | FA, PSD            |
| MA_ST59  | 450 384.0 | 5 924 428.5 | 38            | FA, PSD, CS        |
| MA_ST60  | 446 917.0 | 5 919 909.6 | 23            | FA, PSD            |
| MA_ST61  | 456 198.9 | 5 925 502.3 | 29            | FA, PSD, CS        |
| MA_ST63  | 456 180.4 | 5 925 497.1 | 27            | FA, PSD            |
| MA_ST64  | 456 201.1 | 5 925 486.9 | 27            | FA, PSD            |
| MA_ST65  | 453 581.5 | 5 923 297.0 | 29            | FA, PSD, CS        |
| MA_ST66  | 450 568.4 | 5 920 641.0 | 24            | FA, PSD, CS        |
| <b>Notes</b><br>BSL = Below sea level<br>FA = Faunal sample<br>PSD = Particle size distribution subsample<br>CS = Chemistry sample |           |             |               |                    |






Figure 4.1: Completed survey locations overlaid on bathymetry, Main Array

## 4.2 Sediment Characterisation

### 4.2.1.1 Univariate Analysis

Table 4.3 presents a summary of sediment composition and Table 4.4 provides more detailed granulometric data. Figure 4.2 shows the spatial distribution of fractional composition across the survey area, while Figure 4.3 spatially displays the mean particle size of the stations sampled. Appendix B provides full details of the analytical techniques employed and Appendix D displays the histograms of particle size class summary for each station.

When fractional composition was considered, the sediment was fairly consistent across the main array survey area. All stations consisted of mainly sand with varying proportions of fines (mud; 0.00 % to 8.49 %) and gravel (0.00 % to 37.94 %) (Table 4.3 and Figure 4.3). The proportion of sand content ranged from 55.55 % at station MA\_ST61 to 99.89 % at station MA\_ST66, with a mean value of 80.55 % and low variation (RSD 14 %). The fines content ranged from 0.00 % at 14 stations to 8.49 % at station MA\_ST20, with a mean value of 3.61 % and high variation (standard deviation (SD) 2.80 %). The gravel content ranged from 0.11 % at station MA\_ST66 to 37.94 % at station MA\_ST61, with a mean value of 15.84 % and high variation (SD 9.19 %).

The Folk descriptions classify sediment by the relative proportion of sediment fractions (gravel, sand and fines). The Folk description (BGS modified) described the majority of sediment across the survey area as gravelly sand. Other Folk descriptions were also recorded: sand at ten stations, most of which were located in the eastern part of the survey, gravelly muddy sand at three stations located in the western part of the survey, muddy sandy gravel at station MA\_ST61 and sandy gravel at station MA\_ST02 (Table 4.3 and Figure 4.2).

Table 4.4 presents the physical composition of the sediments (Folk & Ward) at each station. The modality (or modal distribution) represents the number of peaks within the particle size frequency distribution. Distributions were varied across the survey area, with 23 stations displaying unimodal distributions, 22 displaying polymodal distributions and 17 displaying bimodal distributions.

Figure 4.3 presents the mean particle size ( $\mu\text{m}$ ) spatially across the survey area. The mean particle size ( $\mu\text{m}$ ) ranged from 481  $\mu\text{m}$  at station MA\_ST46 to 1315  $\mu\text{m}$  at station MA\_ST61, with a mean of 764  $\mu\text{m}$ . The median particle size ( $\mu\text{m}$ ) ranged from 473 at station MA\_ST46 to 875  $\mu\text{m}$  at station MA\_ST63, with a mean of 632  $\mu\text{m}$ .

The Wentworth description, assigned from mean particle size, categorised the main array survey area as very coarse to medium sand. Six stations were described as very coarse sand, two stations were described as medium sand and the remaining stations were all described as coarse sand.

The sorting coefficient of particle size indicates the degree of spread of individual size classes about the mean and provides the basis of a sorting index, in which low values indicate

sediments to be fairly homogeneous (well sorted) while high values suggest a relatively large scatter of particle sizes about the mean (poorly sorted). Across the survey area, 26 stations were described as poorly sorted, 18 stations were described as very poorly sorted, 8 were described as moderately sorted, 7 were described as moderately well sorted and 3 were described as well sorted.

Skewness indicates the tendency of particle size classes to be skewed about the mean, either towards finer sediment (negative skewed) or coarser sediment (positive skewedness). Across the survey area 27 stations were described as coarse skewed, 21 were described as symmetrical, 13 were described as very coarse skewed and 1 station (MA\_ST59) was described as fine skewed.

Table 4.3: Summary of sediment characteristics, Main Array

| Station | Fractional Composition |             |              | Fines       |             | Folk Description<br>(BGS modified) |
|---------|------------------------|-------------|--------------|-------------|-------------|------------------------------------|
|         | Gravel<br>[%]          | Sand<br>[%] | Fines<br>[%] | Silt<br>[%] | Clay<br>[%] |                                    |
| MA_ST01 | 28.76                  | 65.32       | 5.92         | 4.26        | 1.66        | Gravelly sand                      |
| MA_ST02 | 30.07                  | 66.86       | 3.07         | 2.45        | 0.62        | Sandy gravel                       |
| MA_ST03 | 25.53                  | 68.67       | 5.81         | 4.21        | 1.60        | Gravelly sand                      |
| MA_ST04 | 23.87                  | 69.85       | 6.28         | 4.75        | 1.54        | Gravelly sand                      |
| MA_ST05 | 22.68                  | 70.92       | 6.39         | 4.66        | 1.74        | Gravelly sand                      |
| MA_ST06 | 17.46                  | 80.49       | 2.04         | 1.65        | 0.39        | Gravelly sand                      |
| MA_ST07 | 19.51                  | 76.18       | 4.31         | 3.23        | 1.08        | Gravelly sand                      |
| MA_ST08 | 24.74                  | 70.63       | 4.63         | 3.48        | 1.15        | Gravelly sand                      |
| MA_ST09 | 23.35                  | 74.62       | 2.03         | 1.69        | 0.34        | Gravelly sand                      |
| MA_ST10 | 24.66                  | 70.67       | 4.67         | 3.54        | 1.13        | Gravelly sand                      |
| MA_ST11 | 23.35                  | 70.75       | 5.90         | 4.33        | 1.56        | Gravelly sand                      |
| MA_ST12 | 22.68                  | 69.08       | 8.24         | 6.03        | 2.21        | Gravelly muddy sand                |
| MA_ST13 | 16.30                  | 78.47       | 5.23         | 3.83        | 1.40        | Gravelly sand                      |
| MA_ST14 | 14.64                  | 78.50       | 6.86         | 4.96        | 1.90        | Gravelly sand                      |
| MA_ST15 | 28.79                  | 65.15       | 6.06         | 4.57        | 1.49        | Gravelly sand                      |
| MA_ST16 | 26.72                  | 69.11       | 4.18         | 3.22        | 0.95        | Gravelly sand                      |
| MA_ST17 | 19.90                  | 73.04       | 7.05         | 4.96        | 2.09        | Gravelly sand                      |
| MA_ST18 | 24.60                  | 69.87       | 5.52         | 4.10        | 1.43        | Gravelly sand                      |
| MA_ST19 | 23.87                  | 69.99       | 6.14         | 4.53        | 1.62        | Gravelly sand                      |
| MA_ST20 | 19.43                  | 72.08       | 8.49         | 6.18        | 2.31        | Gravelly muddy sand                |
| MA_ST21 | 25.50                  | 70.93       | 3.56         | 2.63        | 0.94        | Gravelly sand                      |
| MA_ST22 | 9.84                   | 85.62       | 4.54         | 3.38        | 1.16        | Gravelly sand                      |
| MA_ST23 | 22.85                  | 69.45       | 7.70         | 5.54        | 2.16        | Gravelly sand                      |
| MA_ST24 | 3.35                   | 93.42       | 3.23         | 2.55        | 0.68        | Sand                               |

| Station | Fractional Composition |             |              | Fines       |             | Folk Description<br>(BGS modified) |
|---------|------------------------|-------------|--------------|-------------|-------------|------------------------------------|
|         | Gravel<br>[%]          | Sand<br>[%] | Fines<br>[%] | Silt<br>[%] | Clay<br>[%] |                                    |
| MA_ST25 | 18.34                  | 76.36       | 5.30         | 3.83        | 1.47        | Gravelly sand                      |
| MA_ST27 | 12.98                  | 82.56       | 4.46         | 3.18        | 1.28        | Gravelly sand                      |
| MA_ST29 | 21.73                  | 75.36       | 2.90         | 2.24        | 0.66        | Gravelly sand                      |
| MA_ST30 | 29.34                  | 63.08       | 7.58         | 5.28        | 2.30        | Gravelly muddy sand                |
| MA_ST31 | 19.29                  | 73.95       | 6.76         | 4.97        | 1.79        | Gravelly sand                      |
| MA_ST32 | 5.75                   | 94.25       | 0.00         | 0.00        | 0.00        | Gravelly sand                      |
| MA_ST33 | 19.78                  | 75.91       | 4.31         | 3.15        | 1.16        | Gravelly sand                      |
| MA_ST34 | 7.49                   | 90.81       | 1.70         | 1.34        | 0.36        | Gravelly sand                      |
| MA_ST35 | 6.99                   | 93.01       | 0.00         | 0.00        | 0.00        | Gravelly sand                      |
| MA_ST36 | 13.38                  | 82.44       | 4.18         | 3.28        | 0.90        | Gravelly sand                      |
| MA_ST37 | 26.58                  | 67.13       | 6.29         | 4.56        | 1.73        | Gravelly sand                      |
| MA_ST38 | 17.01                  | 82.67       | 0.32         | 0.27        | 0.05        | Gravelly sand                      |
| MA_ST39 | 13.96                  | 86.04       | 0.00         | 0.00        | 0.00        | Gravelly sand                      |
| MA_ST40 | 25.15                  | 69.16       | 5.69         | 4.43        | 1.26        | Gravelly sand                      |
| MA_ST41 | 9.75                   | 84.92       | 5.34         | 4.08        | 1.26        | Gravelly sand                      |
| MA_ST42 | 17.09                  | 75.26       | 7.66         | 5.47        | 2.19        | Gravelly sand                      |
| MA_ST43 | 22.35                  | 70.42       | 7.22         | 5.22        | 2.00        | Gravelly sand                      |
| MA_ST44 | 16.40                  | 79.06       | 4.54         | 3.39        | 1.15        | Gravelly sand                      |
| MA_ST45 | 16.16                  | 82.80       | 1.04         | 0.81        | 0.23        | Gravelly sand                      |
| MA_ST46 | 7.12                   | 92.88       | 0.00         | 0.00        | 0.00        | Gravelly sand                      |
| MA_ST47 | 8.82                   | 90.62       | 0.56         | 0.53        | 0.03        | Gravelly sand                      |
| MA_ST48 | 9.31                   | 86.11       | 4.58         | 3.51        | 1.07        | Gravelly sand                      |
| MA_ST49 | 11.74                  | 87.18       | 1.08         | 0.93        | 0.15        | Gravelly sand                      |
| MA_ST50 | 7.37                   | 92.19       | 0.44         | 0.44        | 0.01        | Gravelly sand                      |
| MA_ST51 | 8.79                   | 90.90       | 0.31         | 0.27        | 0.04        | Gravelly sand                      |

| Station  | Fractional Composition |              |              | Fines       |             | Folk Description<br>(BGS modified) |
|--|------------------------|--------------|--------------|-------------|-------------|------------------------------------|
|  | Gravel<br>[%]          | Sand<br>[%]  | Fines<br>[%] | Silt<br>[%] | Clay<br>[%] |                                    |
| MA_ST52  | 1.96                   | 98.04        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST54  | 12.49                  | 86.11        | 1.39         | 1.12        | 0.27        | Gravelly sand                      |
| MA_ST55  | 1.80                   | 98.20        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST56  | 3.60                   | 96.40        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST57  | 0.99                   | 99.01        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST58  | 10.81                  | 83.51        | 5.68         | 4.33        | 1.35        | Gravelly sand                      |
| MA_ST59  | 0.97                   | 99.03        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST60  | 1.10                   | 98.90        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST61  | 37.94                  | 55.55        | 6.51         | 4.75        | 1.76        | Muddy sandy gravel                 |
| MA_ST63  | 8.40                   | 91.60        | 0.00         | 0.00        | 0.00        | Gravelly sand                      |
| MA_ST64  | 4.74                   | 95.26        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST65  | 1.94                   | 98.06        | 0.00         | 0.00        | 0.00        | Sand                               |
| MA_ST66  | 0.11                   | 99.89        | 0.00         | 0.00        | 0.00        | Sand                               |
| <b>Minimum</b>   | <b>0.11</b>            | <b>55.55</b> | <b>0.00</b>  | <b>0.00</b> | <b>0.00</b> | -                                  |
| <b>Maximum</b>   | <b>37.94</b>           | <b>99.89</b> | <b>8.49</b>  | <b>6.18</b> | <b>2.31</b> |                                    |
| <b>Median</b>  | <b>16.71</b>           | <b>78.78</b> | <b>4.31</b>  | <b>3.23</b> | <b>1.07</b> |                                    |
| <b>Mean</b>  | <b>15.84</b>           | <b>80.55</b> | <b>3.61</b>  | <b>2.68</b> | <b>0.93</b> |                                    |
| <b>Standard Deviation</b>  | <b>9.19</b>            | <b>11.38</b> | <b>2.80</b>  | <b>2.03</b> | <b>0.78</b> |                                    |
| <p><b>Notes</b><br/>                     KP = Kilometre point<br/>                     TOM = Total organic matter<br/>                     TOC = Total organic carbon<br/>                     Fines = silt and clay content<br/>                     Silt = +4.0 to +8.0 <math>\phi</math> units or 3.9 <math>\mu</math>m to 62.5 <math>\mu</math>m<br/>                     Clay = +8.0 to +10.0 <math>\phi</math> units or 0.98 <math>\mu</math>m to 3.9 <math>\mu</math>m<br/>                     BGS = British Geological Survey</p> |                        |              |              |             |             |                                    |

Table 4.4: Summary of particle size distribution, Main Array

| Station | Modality  | D10<br>[µm] | Median<br>[µm]* | D90<br>[µm] | Mean Particle Size |        |                                 | Sorting Coefficient |                    | Skewness |                    |
|---------|-----------|-------------|-----------------|-------------|--------------------|--------|---------------------------------|---------------------|--------------------|----------|--------------------|
|         |           |             |                 |             | [µm]*              | [phi]* | Wentworth (1922)<br>Description | [µm]*               | Description†       | [µm]*    | Description†       |
| MA_ST01 | Polymodal | 268         | 732             | 5880        | 1010               | -0.01  | Very coarse sand                | 4.30                | Very poorly sorted | 0.13     | Coarse skewed      |
| MA_ST02 | Polymodal | 289         | 763             | 8050        | 1138               | -0.19  | Very coarse sand                | 3.50                | Poorly sorted      | 0.39     | Very coarse skewed |
| MA_ST03 | Polymodal | 258         | 640             | 5500        | 919                | 0.12   | Coarse sand                     | 4.19                | Very poorly sorted | 0.19     | Coarse skewed      |
| MA_ST04 | Polymodal | 252         | 673             | 33800       | 1030               | -0.04  | Very coarse sand                | 6.19                | Very poorly sorted | 0.28     | Coarse skewed      |
| MA_ST05 | Polymodal | 238         | 644             | 4680        | 839                | 0.25   | Coarse sand                     | 4.28                | Very poorly sorted | 0.09     | Symmetrical        |
| MA_ST06 | Bimodal   | 290         | 662             | 3360        | 802                | 0.32   | Coarse sand                     | 2.55                | Poorly sorted      | 0.32     | Very coarse skewed |
| MA_ST07 | Bimodal   | 268         | 646             | 3700        | 805                | 0.31   | Coarse sand                     | 2.80                | Poorly sorted      | 0.30     | Coarse skewed      |
| MA_ST08 | Bimodal   | 263         | 674             | 4910        | 907                | 0.14   | Coarse sand                     | 3.50                | Poorly sorted      | 0.22     | Coarse skewed      |
| MA_ST09 | Polymodal | 279         | 688             | 4650        | 899                | 0.15   | Coarse sand                     | 2.91                | Poorly sorted      | 0.34     | Very coarse skewed |
| MA_ST10 | Polymodal | 266         | 645             | 5750        | 915                | 0.13   | Coarse sand                     | 3.38                | Poorly sorted      | 0.36     | Very coarse skewed |
| MA_ST11 | Bimodal   | 239         | 624             | 4530        | 825                | 0.28   | Coarse sand                     | 3.96                | Poorly sorted      | 0.13     | Coarse skewed      |
| MA_ST12 | Polymodal | 189         | 643             | 4360        | 804                | 0.31   | Coarse sand                     | 4.72                | Very poorly sorted | -0.01    | Symmetrical        |
| MA_ST13 | Bimodal   | 269         | 622             | 3650        | 757                | 0.40   | Coarse sand                     | 3.18                | Poorly sorted      | 0.14     | Coarse skewed      |
| MA_ST14 | Bimodal   | 196         | 603             | 2800        | 673                | 0.57   | Coarse sand                     | 3.84                | Poorly sorted      | -0.05    | Symmetrical        |
| MA_ST15 | Polymodal | 221         | 665             | 8210        | 1005               | -0.01  | Very coarse sand                | 4.99                | Very poorly sorted | 0.19     | Coarse skewed      |
| MA_ST16 | Polymodal | 258         | 682             | 5200        | 930                | 0.11   | Coarse sand                     | 3.31                | Poorly sorted      | 0.34     | Very coarse skewed |
| MA_ST17 | Bimodal   | 219         | 605             | 3810        | 764                | 0.39   | Coarse sand                     | 4.25                | Very poorly sorted | 0.03     | Symmetrical        |
| MA_ST18 | Polymodal | 260         | 657             | 6310        | 941                | 0.09   | Coarse sand                     | 4.15                | Very poorly sorted | 0.21     | Coarse skewed      |
| MA_ST19 | Polymodal | 252         | 650             | 4950        | 869                | 0.20   | Coarse sand                     | 4.29                | Very poorly sorted | 0.12     | Coarse skewed      |
| MA_ST20 | Bimodal   | 178         | 571             | 3750        | 716                | 0.48   | Coarse sand                     | 4.58                | Very poorly sorted | 0.00     | Symmetrical        |
| MA_ST21 | Bimodal   | 288         | 669             | 6050        | 1009               | -0.01  | Very coarse sand                | 3.21                | Poorly sorted      | 0.43     | Very coarse skewed |
| MA_ST22 | Unimodal  | 271         | 573             | 1970        | 583                | 0.78   | Coarse sand                     | 2.09                | Poorly sorted      | 0.12     | Coarse skewed      |
| MA_ST23 | Polymodal | 211         | 624             | 4800        | 819                | 0.29   | Coarse sand                     | 4.77                | Very poorly sorted | 0.04     | Symmetrical        |
| MA_ST24 | Unimodal  | 323         | 618             | 986         | 600                | 0.74   | Coarse sand                     | 1.63                | Moderately sorted  | -0.06    | Symmetrical        |

| Station | Modality  | D10<br>[µm] | Median<br>[µm]* | D90<br>[µm] | Mean Particle Size |        |                                 | Sorting Coefficient |                        | Skewness |                    |
|---------|-----------|-------------|-----------------|-------------|--------------------|--------|---------------------------------|---------------------|------------------------|----------|--------------------|
|         |           |             |                 |             | [µm]*              | [phi]* | Wentworth (1922)<br>Description | [µm]*               | Description†           | [µm]*    | Description†       |
| MA_ST25 | Polymodal | 254         | 606             | 3340        | 749                | 0.42   | Coarse sand                     | 3.35                | Poorly sorted          | 0.12     | Coarse skewed      |
| MA_ST27 | Bimodal   | 255         | 541             | 2380        | 650                | 0.62   | Coarse sand                     | 2.46                | Poorly sorted          | 0.29     | Coarse skewed      |
| MA_ST29 | Polymodal | 300         | 691             | 3910        | 874                | 0.19   | Coarse sand                     | 2.64                | Poorly sorted          | 0.34     | Very coarse skewed |
| MA_ST30 | Bimodal   | 206         | 718             | 5360        | 952                | 0.07   | Coarse sand                     | 5.24                | Very poorly sorted     | 0.02     | Symmetrical        |
| MA_ST31 | Bimodal   | 253         | 601             | 3410        | 765                | 0.39   | Coarse sand                     | 4.01                | Very poorly sorted     | 0.04     | Symmetrical        |
| MA_ST32 | Unimodal  | 401         | 705             | 1380        | 683                | 0.55   | Coarse sand                     | 1.60                | Moderately well sorted | 0.07     | Symmetrical        |
| MA_ST33 | Polymodal | 253         | 545             | 4120        | 743                | 0.43   | Coarse sand                     | 2.92                | Poorly sorted          | 0.40     | Very coarse skewed |
| MA_ST34 | Unimodal  | 285         | 577             | 1600        | 579                | 0.79   | Coarse sand                     | 1.88                | Moderately sorted      | 0.14     | Coarse skewed      |
| MA_ST35 | Unimodal  | 367         | 657             | 1590        | 664                | 0.59   | Coarse sand                     | 1.75                | Moderately sorted      | 0.15     | Coarse skewed      |
| MA_ST36 | Bimodal   | 257         | 613             | 2520        | 685                | 0.54   | Coarse sand                     | 2.44                | Poorly sorted          | 0.19     | Coarse skewed      |
| MA_ST37 | Polymodal | 223         | 637             | 5430        | 872                | 0.20   | Coarse sand                     | 4.61                | Very poorly sorted     | 0.13     | Coarse skewed      |
| MA_ST38 | Bimodal   | 294         | 554             | 3840        | 747                | 0.42   | Coarse sand                     | 2.55                | Poorly sorted          | 0.50     | Very coarse skewed |
| MA_ST39 | Bimodal   | 375         | 725             | 2400        | 827                | 0.27   | Coarse sand                     | 2.06                | Poorly sorted          | 0.28     | Coarse skewed      |
| MA_ST40 | Polymodal | 236         | 608             | 5830        | 866                | 0.21   | Coarse sand                     | 4.31                | Very poorly sorted     | 0.21     | Coarse skewed      |
| MA_ST41 | Unimodal  | 246         | 598             | 1960        | 603                | 0.73   | Coarse sand                     | 2.73                | Poorly sorted          | -0.06    | Symmetrical        |
| MA_ST42 | Polymodal | 197         | 639             | 3300        | 735                | 0.44   | Coarse sand                     | 4.22                | Very poorly sorted     | -0.05    | Symmetrical        |
| MA_ST43 | Polymodal | 207         | 670             | 5000        | 869                | 0.20   | Coarse sand                     | 4.77                | Very poorly sorted     | 0.03     | Symmetrical        |
| MA_ST44 | Polymodal | 265         | 585             | 3290        | 724                | 0.47   | Coarse sand                     | 2.64                | Poorly sorted          | 0.32     | Very coarse skewed |
| MA_ST45 | Bimodal   | 307         | 574             | 4300        | 752                | 0.41   | Coarse sand                     | 2.50                | Poorly sorted          | 0.49     | Very coarse skewed |
| MA_ST46 | Unimodal  | 290         | 473             | 933         | 481                | 1.06   | Medium sand                     | 1.86                | Moderately sorted      | 0.32     | Very coarse skewed |
| MA_ST47 | Unimodal  | 307         | 573             | 1770        | 600                | 0.74   | Coarse sand                     | 1.92                | Moderately sorted      | 0.26     | Coarse skewed      |
| MA_ST48 | Unimodal  | 267         | 519             | 1730        | 530                | 0.92   | Coarse sand                     | 2.09                | Poorly sorted          | 0.18     | Coarse skewed      |
| MA_ST49 | Unimodal  | 326         | 664             | 2410        | 721                | 0.47   | Coarse sand                     | 2.16                | Poorly sorted          | 0.27     | Coarse skewed      |
| MA_ST50 | Bimodal   | 303         | 569             | 1610        | 585                | 0.77   | Coarse sand                     | 1.82                | Moderately sorted      | 0.20     | Coarse skewed      |
| MA_ST51 | Unimodal  | 324         | 554             | 1210        | 568                | 0.82   | Coarse sand                     | 1.91                | Moderately sorted      | 0.29     | Coarse skewed      |



| Station        | Modality  | D10<br>[µm] | Median<br>[µm]* | D90<br>[µm]  | Mean Particle Size |              |                                 | Sorting Coefficient |                        | Skewness     |                    |
|----------------|-----------|-------------|-----------------|--------------|--------------------|--------------|---------------------------------|---------------------|------------------------|--------------|--------------------|
|                |           |             |                 |              | [µm]*              | [phi]*       | Wentworth (1922)<br>Description | [µm]*               | Description†           | [µm]*        | Description†       |
| MA_ST52        | Unimodal  | 298         | 481             | 807          | 484                | 1.05         | Medium sand                     | 1.45                | Moderately well sorted | 0.05         | Symmetrical        |
| MA_ST54        | Unimodal  | 315         | 599             | 3300         | 618                | 0.69         | Coarse sand                     | 2.18                | Poorly sorted          | 0.30         | Very coarse skewed |
| MA_ST55        | Unimodal  | 404         | 658             | 964          | 647                | 0.63         | Coarse sand                     | 1.41                | Moderately well sorted | -0.05        | Symmetrical        |
| MA_ST56        | Unimodal  | 458         | 719             | 982          | 703                | 0.51         | Coarse sand                     | 1.40                | Well sorted            | -0.04        | Symmetrical        |
| MA_ST57        | Unimodal  | 362         | 560             | 884          | 559                | 0.84         | Coarse sand                     | 1.42                | Moderately well sorted | -0.03        | Symmetrical        |
| MA_ST58        | Unimodal  | 260         | 606             | 2120         | 662                | 0.60         | Coarse sand                     | 2.94                | Poorly sorted          | -0.05        | Symmetrical        |
| MA_ST59        | Unimodal  | 488         | 720             | 958          | 703                | 0.51         | Coarse sand                     | 1.31                | Well sorted            | -0.20        | Fine skewed        |
| MA_ST60        | Unimodal  | 321         | 512             | 830          | 512                | 0.97         | Coarse sand                     | 1.43                | Moderately well sorted | 0.00         | Symmetrical        |
| MA_ST61        | Polymodal | 287         | 838             | 8690         | 1315               | -0.40        | Very coarse sand                | 5.47                | Very poorly sorted     | 0.13         | Coarse skewed      |
| MA_ST63        | Unimodal  | 528         | 875             | 1900         | 929                | 0.11         | Coarse sand                     | 1.66                | Moderately sorted      | 0.21         | Coarse skewed      |
| MA_ST64        | Unimodal  | 434         | 770             | 1540         | 786                | 0.35         | Coarse sand                     | 1.61                | Moderately well sorted | 0.10         | Coarse skewed      |
| MA_ST65        | Unimodal  | 343         | 533             | 859          | 532                | 0.91         | Coarse sand                     | 1.43                | Moderately well sorted | -0.01        | Symmetrical        |
| MA_ST66        | Unimodal  | 360         | 530             | 804          | 522                | 0.94         | Coarse sand                     | 1.37                | Well sorted            | -0.04        | Symmetrical        |
| <b>Minimum</b> | -         | <b>178</b>  | <b>473</b>      | <b>804</b>   | <b>481</b>         | <b>-0.40</b> | -                               | <b>1.31</b>         | -                      | <b>-0.20</b> | -                  |
| <b>Maximum</b> |           | <b>528</b>  | <b>875</b>      | <b>33800</b> | <b>1315</b>        | <b>1.06</b>  |                                 | <b>6.19</b>         |                        | <b>0.50</b>  |                    |
| <b>Median</b>  |           | <b>269</b>  | <b>630</b>      | <b>3350</b>  | <b>750</b>         | <b>0.41</b>  |                                 | <b>2.77</b>         |                        | <b>0.14</b>  |                    |
| <b>Mean</b>    |           | <b>289</b>  | <b>632</b>      | <b>3880</b>  | <b>764</b>         | <b>0.42</b>  |                                 | <b>3.00</b>         |                        | <b>0.16</b>  |                    |
| <b>SD</b>      |           | <b>70.6</b> | <b>77</b>       | <b>4340</b>  | <b>168</b>         | <b>0.316</b> |                                 | <b>1.26</b>         |                        | <b>0.157</b> |                    |

Notes  
SD = Standard deviation  
\* = Folk & Ward method (Gradistat statistics)  
† = Sorting and skewness based on geometric Folk & Ward (1957) graphical measures (Gradistat statistics)




Figure 4.2: Sediment fractional composition overlaid on bathymetry, Main Array






Figure 4.3: Sediment mean particle size ( $\mu\text{m}$ ) overlaid on bathymetry, Main Array

## 4.2.1.2 Investigation of Granular Similarities


### 4.2.1.2.1 Cluster Analysis

In PRIMER, the 'Cluster' algorithm was used to group stations according to their similarity. Figure 4.4 presents the dendrogram for fourth root transformed sample data. The 'SIMPROF' algorithm was used to identify statistically significant ( $P = 0.05$ ) differences between stations, with significant splits depicted as black lines and non-significant splits as red lines. Statistically significant splits may not be ecologically significant (Clarke et al., 2008), with ecological significance considered in subsequent sections of this report.

There was a high degree of similarity across the survey area with all stations recording a Euclidean distance of 2.54 or below. The 'SIMPROF' analysis ( $P \leq 0.05$ ) identified five clusters and one ungrouped station from the raw data. Following examination of the fractional sediment dataset this appeared to have over differentiated the dataset; as such a slice was positioned at a Euclidean distance of 1.5 to group three statistically significant clusters and one ungrouped station:

- Cluster A comprised 47 stations grouped together with an average squared distance of 1.46;
- Cluster B comprised six stations grouped together with an average squared distance of 1.22;
- Cluster C comprised eight stations grouped together with an average squared distance of 1.33;
- Ungrouped station MA\_ST04 was most similar to cluster A with an average squared distance of 2.07.

Table 4.5 summarises the mean physical characteristics of the sediment groups identified in multivariate analysis. Figure 4.5 spatially presents the sediment groups identified in multivariate analysis overlaid on a SSS mosaic. Sand is the most common fraction between all groups. Cluster A contains the highest percentage of gravel and the lowest percentage of sand out of the three clusters. Both clusters B and C lack fines content; their dissimilarity to each other is likely a result of a slightly higher gravel content in cluster C. Station MA\_ST04 was ungrouped as although the fractional composition is similar to those stations in cluster A, the higher mean particle size suggests the sediment at this station is coarser than those in cluster A.



Notes

Slice at 1.5 resemblance based on Euclidean distance

Figure 4.4: Dendrogram of hierarchical clustering of sediment characteristics data, Main Array

Table 4.5: Summary of particle size for multivariate analysis clusters, Main Array

| Sediment Group  | Stations   | Depth* [m]  | Mean Particle Size* [µm] | Fractional Composition [%] |              |             | Folk Description (BGS modified)†  |
|---|--|-------------|--------------------------|----------------------------|--------------|-------------|---|
|   |  |             |                          | Gravel*                    | Sand*        | Fines*      |   |
| A [■]   | ST16, ST61, ST15, ST37, ST40, ST08, ST30, ST43, ST01, ST19, ST05, ST18, ST03, ST13, ST42, ST14, ST31, ST20, ST11, ST23, ST17, ST12, ST27, ST25, ST58, ST41, ST48, ST22, ST21, ST10, ST02, ST07, ST33, ST44, ST09, ST29, ST06, ST36, ST24, ST50, ST47, ST49, ST34, ST51, ST54, ST45, ST38 | 36.1 ± 4.07 | 793 ± 160                | 19.12 ± 7.38               | 76.25 ± 8.70 | 4.63 ± 2.28 | Gravelly sand, sand, gravelly muddy sand, muddy sandy gravel and sandy gravel |
| B [▲]   | ST66, ST65, ST46, ST57, ST60, ST52   | 27.5 ± 3.51 | 515 ± 29.8               | 2.20 ± 2.51                | 97.80 ± 2.51 | 0.00 ± 0.00 | Gravelly sand and sand  |
| C [▼]   | ST59, ST55, ST56, ST64, ST63, ST35, ST39, ST32   | 33.0 ± 4.21 | 743 ± 97.2               | 5.78 ± 4.14                | 94.22 ± 4.14 | 0.00 ± 0.00 | Gravelly sand and sand  |
| ST04 [*]  | ST04   | 44          | 1030                     | 23.87                      | 69.85        | 6.28        | Gravelly sand   |
| <p>Notes</p> <p>* = Mean ± standard deviation within each sediment group</p> <p>† = Range of Folk descriptions (BGS modified) within each group</p> |  |             |                          |                            |              |             |   |






Figure 4.5: Sediment groups identified in multivariate analysis overlaid on a side scan sonar mosaic, Main Array

#### 4.2.1.2.2 Non-metric Multidimensional Scaling (nMDS)

Figure 4.6 displays the results of the non-metric multidimensional scaling (nMDS), which is an ordination technique that arranges stations on a two-dimensional plot, so that their relative distances from each other reflect similarities in sediment characteristics. The stress coefficient of 0.06 provides a good representation of the data (Clarke & Warwick, 2001).




Figure 4.6: Non-metric multidimensional scaling ordination of particle sizes ( $\mu\text{m}$ )/fractional composition (%), Main Array

#### 4.2.1.2.3 Principal Components Analysis

Figure 4.7 presents principal components analysis (PCA) ordination plot of the percentage of the sediment that comprised each sediment fraction; this analysis was used to identify the sediment fractions driving the variability of the sediment composition across the survey area. The first two principal components accounted for 63.6 % (PC1) and 27.7 % (PC2) of the variation present in the multivariate dataset, cumulatively accounting for 91.3 % of the variation and thereby indicating a good two-dimensional fit to the multivariate data. PC1 has strong positive loadings for granule and fine pebble (eigenvectors of 0.217 and 0.205, respectively), and a strong negative loading for coarse sand (eigenvector of -0.930). PC2 has a strong positive loading for medium sand (eigenvector of 0.901), and strong negative loadings for very coarse sand and granule (eigenvectors of -0.333 and -0.212, respectively). The PCA supports the preceding cluster and nMDS analyses by demonstrating clear separation of the multivariate clusters identified. The eigenvectors superimposed on the plot show that cluster B was differentiated from cluster A due to its higher proportion of medium sand and lower proportion of coarser material (very coarse sand, fine pebble and granule).



Cluster C was differentiated from cluster A due to its higher proportion of coarse sand material.




Figure 4.7: PCA ordination of particle sizes ( $\mu\text{m}$ )/fractional composition (%), Main Array

## 4.3 Sediment Chemistry

### 4.3.1 Sediment Hydrocarbons

#### 4.3.1.1 Sediment Aromatic Hydrocarbon Content

The distribution and concentration of aromatic compounds in seabed sediments were analysed by GC-MS. The aromatic compounds quantified were the naphthalenes (2 ring aromatics), 3 to 6 ring PAHs and the dibenzothiophenes (sulphur containing heteroaromatics). Table 4.6 summarises the total concentrations of aromatic hydrocarbons, including the US EPA 16 PAH and NPD.

Appendix E.1. presents the US EPA 16 PAH concentrations across the Main Array survey area with the individual aromatic hydrocarbons and their alkyl homologue concentrations presented in Appendix E.2., including threshold values where available.

Total 2 to 6 ring PAHs, total US EPA 16 PAHs and total naphthalenes, phenanthrenes and dibenzothiophenes (NPDs) are calculated as the sum of individual PAHs, some of which were less than the minimum reporting value (MRV). Consequently, the total concentration is

assigned as a less than value. However, the concentrations of the individual PAHs that were less than the MRV are unlikely to significantly influence the total concentrations. Therefore, for the purposes of this report, total 2 to 6 ring PAH concentrations, total US EPA 16 PAH concentrations and total NPDs are treated as absolute values to provide comparison between stations and with regional datasets.

Total 2 to 6 ring PAH concentrations ranged from < 0.004 µg/g at station MA\_ST65 to 1.42 µg/g at station MA\_ST12, with a mean concentration of 0.176 µg/g. Total 2 to 6 ring PAH concentrations were higher than the median concentration reported from the SEA6 Irish Sea surveys (0.0237 µg/g; Cefas, 2005) at six stations (stations MA\_ST04, MA\_ST12, MA\_ST22, MA\_ST25, MA\_ST43 and MA\_ST61).

Total US EPA 16 PAH concentrations ranged from < 1.9 ng/g at station MA\_ST65 to 820 ng/g at station MA\_ST12 with a mean of 92.9 ng/g. All US EPA 16 PAHs were below their respective ERL values (OSPAR, 2014) across the Main Array survey area (Appendix E.1).

The proportion of petrogenically derived NPD to total 2 to 6 ring PAH material ranged from 17 % at station MA\_ST12 to < 50 % at station MA\_ST59, with a mean of 35 %.

Table 4.6: Summary of sediment aromatic hydrocarbon analysis, Main Array

| Station   | Total 2 to 6 Ring PAH* | Total US EPA 16 PAH† | NPD                |                |
|---|------------------------|----------------------|--------------------|----------------|
|   |                        |                      | Total*             | [%]            |
| MA_ST04   | 0.0452                 | < 14.5               | 0.0155             | 34             |
| MA_ST12   | 1.42                   | 820                  | 0.247              | 17             |
| MA_ST22   | 0.0445                 | < 11.7               | 0.0207             | 47             |
| MA_ST25   | 0.0351                 | < 12.5               | 0.0120             | 34             |
| MA_ST43   | 0.0747                 | 23.6                 | 0.0295             | 39             |
| MA_ST47   | < 0.0154               | < 4.9                | < 0.0051           | < 33           |
| MA_ST59   | < 0.0159               | < 3.2                | < 0.0078           | < 50           |
| MA_ST61   | 0.0965                 | 33.7                 | 0.0294             | 30             |
| MA_ST65   | < 0.0043               | < 1.9                | < 0.0013           | < 30           |
| MA_ST66   | < 0.0104               | < 3.2                | < 0.0032           | < 31           |
| <b>Minimum</b>  | <b>&lt; 0.0043</b>     | <b>&lt; 1.9</b>      | <b>&lt; 0.0013</b> | <b>17</b>      |
| <b>Maximum</b>  | <b>1.42</b>            | <b>820</b>           | <b>0.247</b>       | <b>&lt; 50</b> |
| <b>Median</b>   | <b>0.0398</b>          | <b>12.1</b>          | <b>0.0138</b>      | <b>34</b>      |
| <b>Mean</b>   | <b>0.176</b>           | <b>92.9</b>          | <b>0.0372</b>      | <b>35</b>      |
| <b>Standard deviation</b>   | <b>0.438</b>           | <b>256</b>           | <b>0.0744</b>      | <b>9.3</b>     |
| <b>RSD [%]</b>  | <b>249</b>             | <b>275</b>           | <b>200</b>         | <b>27</b>      |
| <b>SEA6 (Irish Sea) (Cefas, 2005)‡</b>  |                        |                      |                    |                |
| <b>Mean</b>   | <b>0.0237</b>          | <b>-</b>             | <b>-</b>           | <b>-</b>       |
| <p>Notes</p> <p>For the summary statistics, less than values have been considered as absolute values</p> <p>Total 2 to 6 ring PAH = Total 2 to 6 ring polycyclic aromatic hydrocarbons (PAH), including alkyl homologues</p> <p>Total US EPA 16 PAH = Total United States Environmental Protection Agency's 16 (US EPA PAH) polycyclic aromatic hydrocarbons</p> <p>Total NPD = Total naphthalene, phenanthrene/anthracene and dibenzothiophene</p> <p>NPD [%] = Percentage of total 2 to 6 ring PAH concentration comprised of NPD</p> <p>RSD = Relative standard deviation</p> <p>SD = Standard deviation</p> <p>* = Concentrations expressed as µg/g of dry sediment</p> <p>† = Concentrations expressed as ng/g of dry sediment</p> <p>‡ = Value taken from site 715 (Liverpool Bay) from the Strategic Environmental Assessment 6 (SEA6) review of the Irish Sea (Cefas, 2005)</p> |                        |                      |                    |                |

### 4.3.2 Sediment Metals


Table 4.7 summarises the concentrations of the extractable metals in the sediment samples from an aqua regia digest.

All metals concentrations were less than their respective Cefas guideline action levels (AL1 and AL2) and OSPAR ERL values.

The overall trend in individual metals concentrations is presented in Figure 4.8, and is assessed by comparing the maximum normalised elemental concentrations. Normalising the elemental data for the sediment samples to the highest concentration for each element, the highest concentrations of most metals was seen at station MA\_ST04.

Table 4.7: Summary of sediment metals analysis, Main Array

| Station   | Al        | As   | Ba   | Cd       | Cr   | Cu   | Hg        | Ni   | Pb   | Sn    | Zn   |
|---|-----------|------|------|----------|------|------|-----------|------|------|-------|------|
| MA_ST04   | 5260      | 8.92 | 23.1 | < 0.0800 | 14.5 | 4.58 | < 0.0400  | 7.70 | 13.8 | 1.35  | 27.8 |
| MA_ST12   | 4700      | 12.3 | 17.7 | < 0.0800 | 10.1 | 3.14 | < 0.0400  | 7.57 | 15.5 | 2.46  | 24.7 |
| MA_ST22   | 2420      | 11.4 | 8.43 | < 0.0800 | 6.37 | 4.16 | < 0.0400  | 4.32 | 7.93 | 0.343 | 15.3 |
| MA_ST25   | 2800      | 9.78 | 8.34 | < 0.0800 | 7.39 | 2.53 | < 0.0400  | 4.95 | 9.77 | 0.295 | 16.6 |
| MA_ST43   | 4700      | 12.6 | 22.2 | < 0.0800 | 12.6 | 2.84 | < 0.0400  | 8.02 | 13.3 | 0.536 | 24.8 |
| MA_ST47   | 2480      | 16.5 | 8.01 | < 0.0800 | 7.33 | 1.86 | < 0.0400  | 5.97 | 7.97 | 0.305 | 18.9 |
| MA_ST59   | 2500      | 13.3 | 7.99 | < 0.0800 | 6.94 | 1.66 | < 0.0400  | 5.39 | 5.42 | 0.253 | 18.8 |
| MA_ST61   | 4800      | 10.4 | 16.8 | < 0.0800 | 11.5 | 3.07 | < 0.0400  | 7.17 | 11.6 | 5.11  | 26.1 |
| MA_ST65   | 1700      | 7.68 | 10.2 | < 0.0800 | 5.02 | 1.12 | < 0.0400  | 3.56 | 4.62 | 0.214 | 13.6 |
| MA_ST66   | 2120      | 14.3 | 6.08 | < 0.0800 | 5.75 | 1.48 | < 0.0400  | 4.29 | 4.85 | 0.430 | 14.0 |
| Minimum   | 1700      | 7.68 | 6.08 | < 0.0800 | 5.02 | 1.12 | < 0.0400  | 3.56 | 4.62 | 0.214 | 13.6 |
| Maximum   | 5260      | 16.5 | 23.1 | < 0.0800 | 14.5 | 4.58 | < 0.0400  | 8.02 | 15.5 | 5.11  | 27.8 |
| Median  | 2650      | 11.8 | 9.32 | -        | 7.36 | 2.68 | -         | 5.68 | 8.87 | 0.386 | 18.8 |
| Mean  | 3350      | 11.7 | 12.9 | -        | 8.75 | 2.64 | -         | 5.89 | 9.48 | 1.13  | 20.1 |
| Standard deviation  | 1340      | 2.64 | 6.42 | -        | 3.21 | 1.14 | -         | 1.63 | 3.95 | 1.57  | 5.34 |
| RSD   | 40        | 23   | 50   | -        | 37   | 43   | -         | 28   | 42   | 139   | 27   |
| <b>Cefas Guideline Action Levels</b>  |           |      |      |          |      |      |           |      |      |       |      |
| AL1   | -         | 20   | -    | 0.4      | 40   | 40   | 0.3       | 20   | 50   | -     | 130  |
| AL2   | -         | 100  | -    | 5        | 400  | 400  | 3         | 200  | 500  | -     | 800  |
| <b>CEMP Assessment Criteria (OSPAR, 2014)</b>   |           |      |      |          |      |      |           |      |      |       |      |
| ERL   | -         | -    | -    | 1.20     | 81.0 | 34.0 | 0.150     | -    | 47.0 | -     | 150  |
| <b>Notes</b>  |           |      |      |          |      |      |           |      |      |       |      |
| Concentrations expressed in µg/g dry sediment   |           |      |      |          |      |      |           |      |      |       |      |
| Al = Aluminium                      As = Arsenic                      Ba = Barium                      Cd = Cadmium                      Cr = Chromium                      Cu = Copper |           |      |      |          |      |      |           |      |      |       |      |
| Hg = Mercury                      Ni = Nickel                      Pb = Lead                      Sn = Tin                      Zn = Zinc   |           |      |      |          |      |      |           |      |      |       |      |
| RSD = Relative standard deviation   |           |      |      |          |      |      |           |      |      |       |      |
| Cefas = Centre for Environment, Fisheries and Aquaculture Science   |           |      |      |          |      |      |           |      |      |       |      |
| AL1 = Action level 1  |           |      |      |          |      |      |           |      |      |       |      |
| AL2 = Action level 2  |           |      |      |          |      |      |           |      |      |       |      |
| CEMP = Coordinated Environmental Monitoring Programme   |           |      |      |          |      |      |           |      |      |       |      |
| OSPAR = Oslo and Paris Commission   |           |      |      |          |      |      |           |      |      |       |      |
| ERL = Effects range low   |           |      |      |          |      |      |           |      |      |       |      |
| <b>Key:</b>   | Below ERL |      |      |          |      |      | Above ERL |      |      |       |      |



Notes

Al = Aluminium  
Pb = Lead

As = Arsenic  
Sn = Tin

Ba = Barium  
Zn = Zinc

Cr = Chromium

Cu = Copper

Ni = Nickel

Figure 4.8: Relative (maximum normalised) elemental concentrations in sediments, Main Array

## 4.4 Sediment Macrofauna

### 4.4.1 Infaunal and Solitary Epifauna

#### 4.4.1.1 Phyletic Composition

A full list of taxa identified and enumerated (individuals per 0.1 m<sup>2</sup>) from the survey area are presented in Appendix G.

A total of 336 taxa and 6946 individuals were identified within grab samples from the survey area. Of these taxa, 87 were recorded as juveniles, pelagic, colonial or solitary epifauna. To represent the permanent macrofaunal community and to avoid spurious enhancement of the species list, the dataset was rationalised and these taxa were removed prior to statistical analysis. Records of several indeterminable specimens were also merged with those of identifiable taxa (see Appendix G).

Table 4.8 summarises the abundance of taxonomic groups identified within the rationalised dataset across the survey area and Figures 4.9 and 4.1 display the data graphically.

The rationalised data comprised 247 benthic taxa, of which 124 (50.2 %) were annelids, 63 (25.5 %) were arthropods, 33 (13.4 %) were molluscs, 13 (5.3 %) were echinoderms and 14 (5.7 %) were members of other phyla (specifically cephalochordates, cnidarians, nemerteans, platyhelminths, phoronids, sipunculids and tunicates). A total of 6482 individuals was identified in the rationalised data, of which 3102 (47.9 %) were annelids, 2254 (34.8 %) were arthropods, 182 (2.8 %) were molluscs, 133 (2.1 %) were echinoderms and 811 (12.5 %) were other phyla (Table 4.8).

Table 4.8: Taxonomic groups, Main Array

| Taxonomic Group   | Number of Taxa | Composition of Taxa [%]* | Abundance   | Composition of Individuals [%]* |
|---|----------------|--------------------------|-------------|---------------------------------|
| Annelida  | 124            | 50.2                     | 3102        | 47.9                            |
| Arthropoda  | 63             | 25.5                     | 2254        | 34.8                            |
| Mollusca  | 33             | 13.4                     | 182         | 2.8                             |
| Echinodermata   | 13             | 5.3                      | 133         | 2.1                             |
| Other phyla   | 14             | 5.7                      | 811         | 12.5                            |
| <b>Total</b>  | <b>247</b>     | <b>100</b>               | <b>6482</b> | <b>100</b>                      |
| Notes<br>Macrofaunal samples were processed through a 1 mm sieve<br>Other phyla include: Cephalochordata, Cnidaria, Nemertea, Platyhelminthes, Phoronida, Sipuncula and Tunicata<br>* = Percentages expressed to 1 decimal place and, due to numerical rounding, values presented may not equate to 100 % |                |                          |             |                                 |

Figures 4.9 and 4.1 illustrate the phyletic composition of taxa and individuals for each station (per 0.1 m<sup>2</sup>), respectively, facilitating spatial comparison and highlighting the broad similarities between stations across the survey area. Annelids contributed the highest (or joint highest) number of taxa at all stations; at stations MA\_ST51 and MA\_ST66 annelids and

arthropods contributed the joint highest number of taxa. Annelids also contributed the highest (or joint highest) number of individuals at the majority of stations, although arthropods were the most abundant phylum at 19 of the 62 stations sampled. The most notable variations were in the proportional contributions of echinoderms and molluscs to both taxa and abundance of individuals, with their absence at 17 and 9 stations, respectively. Furthermore, arthropods, molluscs and echinoderms were absent from station MA\_ST59, and molluscs, echinoderms and other phylum were absent from station MA\_ST66.






Figure 4.9: Phyletic composition of macrofaunal taxa, Main Array




Figure 4.10: Phyletic composition of macrofaunal individuals, Main Array

#### 4.4.1.2 Community Statistics

Table 4.9 presents the number of taxa and individuals identified within the rationalised dataset from each station along with several commonly used diversity and evenness statistics. Figures 4.11 and 4.12 spatially presents the number of taxa and individuals across the survey area.

The number of taxa per station (0.1 m<sup>2</sup>) ranged from 2 at station MA\_ST59 to 65 at station MA\_ST61 with a mean of 37.

The number of individuals per station (0.1 m<sup>2</sup>) ranged from 2 at station MA\_ST59 to 244 at station MA\_ST11 with a mean of 105. A broad spatial trend in taxa and individuals was observed, where typically higher numbers were recorded in the west of the survey area (Figures 4.11 and 4.12).

When the Shannon-Wiener index was considered in the context of the diversity thresholds suggested by Dauvin et al. (2012; Table 3.2), diversity could be inferred to range from poor to high, with the majority of stations (40 of 62 the stations sampled) featuring high diversity (> 4.00). The minimum Shannon-Wiener of 1.00 (low diversity) was reported from station MA\_ST59 and the maximum of 5.49 from station MA\_ST61.

The Pielou's evenness index (J) is a measure of the distribution of individuals between taxa. It ranged from 0.716 at station MA\_ST63 to 1.000 at station MA\_ST59, with a mean of 0.866. All stations exhibited moderate to high evenness (> 0.700). The complement of the Simpsons index of dominance (1 - λ) largely mirrored the Pielou's evenness index.

Table 4.9: Macrofaunal community statistics (0.1 m<sup>2</sup>), Main Array

| Station | Numbers |             | Shannon-Wiener<br>(H'Log <sub>2</sub> ) | Evenness      |                    |
|---------|---------|-------------|---|---------------|--------------------|
|         | Taxa    | Individuals |   | Pielou<br>(J) | Simpson<br>(1 - λ) |
| MA_ST01 | 62      | 173         | 5.37                                    | 0.902         | 0.960              |
| MA_ST02 | 40      | 111         | 4.36                                    | 0.818         | 0.901              |
| MA_ST03 | 59      | 195         | 4.95                                    | 0.842         | 0.941              |
| MA_ST04 | 53      | 133         | 5.08                                    | 0.888         | 0.954              |
| MA_ST05 | 54      | 134         | 5.04                                    | 0.876         | 0.942              |
| MA_ST06 | 39      | 118         | 4.43                                    | 0.837         | 0.910              |
| MA_ST07 | 40      | 145         | 4.30                                    | 0.809         | 0.893              |
| MA_ST08 | 46      | 166         | 4.47                                    | 0.810         | 0.920              |
| MA_ST09 | 51      | 129         | 4.79                                    | 0.844         | 0.928              |
| MA_ST10 | 49      | 168         | 4.45                                    | 0.793         | 0.903              |
| MA_ST11 | 55      | 244         | 4.50                                    | 0.779         | 0.913              |
| MA_ST12 | 44      | 126         | 4.58                                    | 0.839         | 0.926              |
| MA_ST13 | 34      | 130         | 3.84                                    | 0.755         | 0.851              |

| Station | Numbers |             | Shannon-Wiener<br>(H'Log <sub>2</sub> ) | Evenness      |                    |
|---------|---------|-------------|---|---------------|--------------------|
|         | Taxa    | Individuals |   | Pielou<br>(J) | Simpson<br>(1 - λ) |
| MA_ST14 | 51      | 161         | 4.73                                    | 0.834         | 0.926              |
| MA_ST15 | 52      | 137         | 4.94                                    | 0.867         | 0.947              |
| MA_ST16 | 49      | 158         | 4.55                                    | 0.810         | 0.911              |
| MA_ST17 | 51      | 163         | 4.87                                    | 0.858         | 0.941              |
| MA_ST18 | 48      | 174         | 4.52                                    | 0.810         | 0.913              |
| MA_ST19 | 42      | 117         | 4.90                                    | 0.908         | 0.954              |
| MA_ST20 | 56      | 146         | 5.27                                    | 0.908         | 0.963              |
| MA_ST21 | 29      | 55          | 4.34                                    | 0.893         | 0.922              |
| MA_ST22 | 40      | 126         | 4.51                                    | 0.847         | 0.931              |
| MA_ST23 | 45      | 124         | 4.82                                    | 0.878         | 0.942              |
| MA_ST24 | 31      | 80          | 4.49                                    | 0.905         | 0.933              |
| MA_ST25 | 37      | 85          | 4.42                                    | 0.849         | 0.908              |
| MA_ST27 | 42      | 87          | 5.01                                    | 0.929         | 0.960              |
| MA_ST29 | 25      | 79          | 3.44                                    | 0.740         | 0.802              |
| MA_ST30 | 54      | 207         | 4.21                                    | 0.732         | 0.860              |
| MA_ST31 | 31      | 84          | 4.19                                    | 0.847         | 0.911              |
| MA_ST32 | 31      | 75          | 4.44                                    | 0.897         | 0.937              |
| MA_ST33 | 46      | 158         | 4.37                                    | 0.791         | 0.902              |
| MA_ST34 | 25      | 52          | 4.32                                    | 0.931         | 0.939              |
| MA_ST35 | 16      | 27          | 3.78                                    | 0.945         | 0.914              |
| MA_ST36 | 31      | 63          | 4.53                                    | 0.915         | 0.936              |
| MA_ST37 | 56      | 126         | 4.96                                    | 0.854         | 0.930              |
| MA_ST38 | 22      | 45          | 4.17                                    | 0.935         | 0.933              |
| MA_ST39 | 21      | 56          | 4.02                                    | 0.915         | 0.926              |
| MA_ST40 | 43      | 125         | 4.65                                    | 0.857         | 0.934              |
| MA_ST41 | 37      | 113         | 4.28                                    | 0.822         | 0.896              |
| MA_ST42 | 61      | 157         | 5.39                                    | 0.908         | 0.965              |
| MA_ST43 | 52      | 164         | 4.84                                    | 0.848         | 0.930              |
| MA_ST44 | 25      | 48          | 4.31                                    | 0.928         | 0.937              |
| MA_ST45 | 33      | 88          | 4.49                                    | 0.890         | 0.938              |
| MA_ST46 | 18      | 32          | 3.88                                    | 0.931         | 0.916              |
| MA_ST47 | 34      | 88          | 4.62                                    | 0.908         | 0.943              |
| MA_ST48 | 40      | 163         | 3.82                                    | 0.717         | 0.825              |
| MA_ST49 | 48      | 130         | 4.93                                    | 0.883         | 0.940              |
| MA_ST50 | 26      | 49          | 4.44                                    | 0.945         | 0.945              |

| Station                   | Numbers     |             | Shannon-Wiener<br>(H'Log <sub>2</sub> ) | Evenness      |                    |
|---------------------------|-------------|-------------|---|---------------|--------------------|
|                           | Taxa        | Individuals |   | Pielou<br>(J) | Simpson<br>(1 - λ) |
| MA_ST51                   | 24          | 53          | 4.10                                    | 0.895         | 0.917              |
| MA_ST52                   | 19          | 40          | 3.83                                    | 0.903         | 0.906              |
| MA_ST54                   | 43          | 93          | 4.87                                    | 0.897         | 0.950              |
| MA_ST55                   | 14          | 17          | 3.69                                    | 0.969         | 0.913              |
| MA_ST56                   | 12          | 14          | 3.52                                    | 0.982         | 0.908              |
| MA_ST57                   | 8           | 25          | 2.38                                    | 0.794         | 0.749              |
| MA_ST58                   | 34          | 95          | 4.43                                    | 0.870         | 0.927              |
| MA_ST59                   | 2           | 2           | 1.00                                    | 1.000         | 0.500              |
| MA_ST60                   | 11          | 17          | 3.29                                    | 0.952         | 0.886              |
| MA_ST61                   | 65          | 187         | 5.49                                    | 0.912         | 0.971              |
| MA_ST63                   | 25          | 122         | 3.32                                    | 0.716         | 0.830              |
| MA_ST64                   | 23          | 62          | 3.63                                    | 0.803         | 0.857              |
| MA_ST65                   | 14          | 31          | 3.58                                    | 0.940         | 0.905              |
| MA_ST66                   | 4           | 10          | 1.69                                    | 0.843         | 0.640              |
| <b>Minimum</b>            | <b>2</b>    | <b>2</b>    | <b>1.00</b>                             | <b>0.716</b>  | <b>0.500</b>       |
| <b>Maximum</b>            | <b>65</b>   | <b>244</b>  | <b>5.49</b>                             | <b>1.000</b>  | <b>0.971</b>       |
| <b>Median</b>             | <b>40</b>   | <b>115</b>  | <b>4.44</b>                             | <b>0.873</b>  | <b>0.926</b>       |
| <b>Mean</b>               | <b>37</b>   | <b>105</b>  | <b>4.30</b>                             | <b>0.866</b>  | <b>0.905</b>       |
| <b>Standard Deviation</b> | <b>15.6</b> | <b>56.2</b> | <b>0.790</b>                            | <b>0.0644</b> | <b>0.0744</b>      |






Figure 4.11: Number of macrofaunal taxa per station/sample (0.1 m<sup>2</sup>) overlaid on bathymetry, Main Array






Figure 4.12: Number of macrofaunal individuals per station/sample (0.1 m<sup>2</sup>) overlaid on bathymetry, Main Array


### 4.4.13 Investigation of Faunal Similarities

#### 4.4.13.1 Cluster Analysis

In PRIMER, the 'Cluster' algorithm was used to group stations according to their faunal similarity. Figure 4.13 presents the hierarchical agglomerative cluster dendrogram for square root transformed station data. The 'SIMPROF' algorithm was used to identify statistically significant ( $P = 0.05$ ) differences between stations, with significant splits depicted as black lines and non-significant splits as red lines. Statistically significant splits may not be ecologically significant (Clarke et al., 2008), with ecological significance considered in subsequent sections of this report.

There was a low to moderate degree of similarity across the Main Array survey area. The 'SIMPROF' analysis ( $P \leq 0.05$ ) identified 11 clusters and six ungrouped stations. Following analysis of the abundance dataset STIMPROF was deemed to have over differentiated the dataset; as such, a slice was positioned at 25 % Bray-Curtis similarity to define three statistically significant clusters and one ungrouped station:

- Cluster A comprised 45 stations which grouped together with a mean similarity of 42.9 %;
- Cluster B comprised 5 stations which grouped together with a mean similarity of 37.0 %;
- Cluster C comprised 11 stations which grouped together with a mean similarity of 38.4 %;
- Ungrouped station MA\_ST59 was statistically significantly different from clusters A, B and C, with an average dissimilarity of 99.0 %, 100.0 % and 98.8 %, respectively.



Note  
Slice at 25 % resemblance

Figure 4.13: Dendrogram of hierarchical clustering of macrofaunal station (0.1 m<sup>2</sup>) abundance data, Main Array






Figure 4.14: Macrofaunal clusters per station (0.1 m<sup>2</sup>) overlaid on bathymetry Main Array

#### 4.4.1.3.2 Similarity Percentage Analysis

Table 4.10 summarises the top 10 most abundant taxa within each cluster and at the ungrouped station. The average dissimilarity between the clusters ranged from 79.2 % (clusters A and B) to 100.0 % (cluster B and station MA\_ST59).


The top ten most abundant taxa for the clusters and ungrouped stations cumulatively comprised between 53.5 % (Cluster A) and 100.0 % (ungrouped station MA\_ST59) of the taxa present within them (Table 4.10). The most abundant characterising taxa within cluster A comprised the amphipod *Urothoe marina*, the horseshoe worm *Phoronis* sp. and the polychaete *Aonides paucibranchiata*. Cluster B was characterised by an abundance of polychaetes *Pisione remota* and *Polygordius*, followed by nemerteans. Cluster C was dominated by the amphipod crustacean *Bathyporeia elegans*, accompanied by the polychaetes *Spiophanes bombyx* and *Spio goniocephala*. The paucity of fauna at ungrouped station MA\_ST59 (as previously demonstrated in the analysis of community statistics; Table 4.9) lead to 100.0 % of the macrofaunal community being represented by two individuals of two different taxa, the polychaete *Syllis parapari* and the sipunculid *Phascolion strombus*. As a result, ungrouped station MA\_ST59 had high dissimilarity to the other clusters.

Figures 4.15 to 4.17 highlight the differences in abundance and distribution of key taxa across the survey area. *U. marina* were reported in greater abundances to the west of the survey area in deeper waters, whilst *B. elegans* showed the opposite trend with a greater abundance in the east/south of the survey area. *P. remota* showed no clear spatial distribution. The relationship between macrofauna communities and physical and chemical determinants will be discussed further in Sections 4.4.1.4 and 5.

Table 4.10: Top ten most abundant macrofaunal taxa within cluster, Main Array

| Group A (▲)  | Ind<br>[0.1 m <sup>2</sup> ] | Cum<br>[%] | Group B (■)                    | Ind<br>[0.1 m <sup>2</sup> ] | Cum<br>[%] |
|--|------------------------------|------------|--------------------------------|------------------------------|------------|
| <i>Urothoe marina</i>  | 22                           | 26.2       | <i>Pisone remota</i>           | 11                           | 23.5       |
| <i>Phoronis</i>  | 8                            | 31.3       | <i>Polygordius</i>             | 11                           | 33.5       |
| <i>Aonides paucibranchiata</i>   | 7                            | 39.8       | NEMERTEA                       | 4                            | 47.2       |
| <i>Poecilochaetus serpens</i>  | 5                            | 43.8       | <i>Spio symphyta</i>           | 3                            | 56.6       |
| NEMERTEA   | 5                            | 51.5       | <i>Eurydice pulchra</i>        | 2                            | 64.7       |
| <i>Spiophanes bombyx</i>   | 5                            | 57.8       | <i>Aonides paucibranchiata</i> | 2                            | 73.5       |
| <i>Ampelisca provincialis</i>  | 5                            | 60.9       | <i>Echinocyamus pusillus</i>   | 2                            | 76.0       |
| <i>Ampharete lindstroemi</i>   | 4                            | 64.8       | <i>Bathyporeia elegans</i>     | 1                            | 82.0       |
| <i>Ampelisca diadema</i>   | 3                            | 67.2       | <i>Goniadella gracilis</i>     | 1                            | 83.9       |
| <i>Ampelisca typica</i>  | 2                            | 69.3       | <i>Aricidea cerrutii</i>       | 1                            | 89.3       |
| <i>Notomastus</i>  | 2                            | 71.2       |                                |                              |            |
| Group C (●)  | Ind<br>[0.1 m <sup>2</sup> ] | Cum<br>[%] | Station MA_ST59 (*)            | Ind<br>[0.1 m <sup>2</sup> ] | Cum<br>[%] |
| <i>Bathyporeia elegans</i>   | 4                            | 19.4       | <i>Syllis parapari</i>         | 1                            | 50         |
| <i>Spiophanes bombyx</i>   | 3                            | 32.1       | <i>Phascolion strombus</i>     | 1                            | 100.0      |
| <i>Spio gonocephala</i>  | 3                            | 45.8       |                                |                              |            |
| <i>Echinocyamus pusillus</i>   | 3                            | 53.4       |                                |                              |            |
| <i>Bathyporeia gracilis</i>  | 3                            | 57.5       |                                |                              |            |
| <i>Nephtys cirrosa</i>   | 2                            | 71.8       |                                |                              |            |
| NEMERTEA   | 2                            | 77.4       |                                |                              |            |
| <i>Poecilochaetus serpens</i>  | 1                            | 80.5       |                                |                              |            |
| <i>Polycirrus</i>  | 1                            | 82.7       |                                |                              |            |
| <i>Glycera oxycephala</i>  | 1                            | 85.2       |                                |                              |            |
| <i>Spisula elliptica</i>   | 1                            | 87.7       |                                |                              |            |
| Notes  |                              |            |                                |                              |            |
| Ind = Number of individuals of each taxon within the station (0.1 m <sup>2</sup> )   |                              |            |                                |                              |            |
| Cum = Cumulative percentage of the individuals of each taxon relative to the total number of individuals recorded within the station (0.1 m <sup>2</sup> ) |                              |            |                                |                              |            |






Notes

Abundance of *Urothoe marina* expressed per station (0.1 m<sup>2</sup>)


Figure 4.15: *Urothoe marina* abundance overlaid on macrofaunal community groupings and a bathymetric mosaic, Main Array



Document Path: S:\430-MGC-ITV\Charting\E200145\_Innogy\_AyM\_Benthic\_Survey\3\_Plots\2\_Draft\Array\EBS\Q200145\_11\_Pisione.mxd  
30/11/2020 12:02:47

Notes  
Abundance of *Pisione remota* expressed per station (0.1 m<sup>2</sup>)  
Figure 4.16: *Pisione remota* abundance overlaid on macrofaunal community groupings and a bathymetric mosaic, Main Array





Document Path: S:\516-MSC-IT\Charling\EP00145\_Innogy\_AyM\_Benthic\_Survey\3\_Plot\2\_Draft\Array\EB5\Q00145\_12\_Bathyporeia.mxd  
30/11/2020 - 11:58:25

Coordinate System: WGS 1984 UTM Zone 30N

Notes

Abundance of *Bathyporeia elegans* expressed per station (0.1 m²)

Figure 4.17: *Bathyporeia elegans* abundance overlaid on macrofaunal community groupings and a bathymetric mosaic, Main Array

#### 4.4.1.4 Biomass

Table 4.12 summarises the total macrofaunal biomass and phyletic composition of the biomass by station across the survey area. Biomass is expressed as AFDW in g/0.1 m<sup>2</sup> grab sample. Figure 4.18 presents the phyletic composition of the biomass graphically, whilst Figure 4.19 spatially presents the total macrofaunal biomass across the survey area.

The total biomass of macrofauna ranged from < 0.001 g at station MA\_ST59 to 2.666 g at station MA\_ST42, with a median total biomass of 0.306 g and a mean of 0.451 g.

The biomass of annelids ranged from < 0.001 g at station MA\_ST59 to 0.537 g at station MA\_ST49. Annelids contributed the highest proportion of biomass at 23 of the 62 stations within the survey areas, accounting for over 90 % of the biomass at stations MA\_ST57 and MA\_ST59 (although total biomass at the latter was < 0.001 g).

Arthropods were absent from station MA\_ST59, but where recorded, contributed a biomass of between < 0.001 g at station MA\_ST59 to 1.054 g at station MA\_ST43. Arthropods contributed the highest proportion of biomass at 6 of the 62 stations within the survey areas, accounting for a maximum proportion of 82.57 % of the biomass (station MA\_ST43).

Molluscs were absent from six stations, but where recorded, contributed a biomass of between < 0.001 g at four stations to 1.334 g at station MA\_ST56. Molluscs contributed the highest proportion of biomass at 9 of the 62 stations within the survey areas, accounting for over 90 % of the biomass at station MA\_ST56.

Echinoderms were absent from twelve stations, but where recorded, contributed a biomass of between < 0.001 g at five stations to 0.923 g at station MA\_ST42. Echinoderms contributed the highest proportion of biomass at 7 of the 62 stations within the survey areas, accounting for over 90 % of the biomass at stations MA\_ST60 and MA\_ST64.

'Other phyla' were absent from seventeen stations, but where recorded, contributed a biomass of between < 0.001 g at three stations to 0.351 g at station MA\_ST04. 'Other phyla' contributed the highest proportion of biomass at 2 of the 62 stations within the survey areas, accounting for a maximum proportion of 45.26 % of the biomass (station MA\_ST04).


Neither total biomass, nor the biomass contributed by individual phyla, showed clear spatial distribution across the Main Array survey area (Figure 4.19).



Table 4.11: Phyletic composition of macrofaunal biomass, Main Array


| Station | Biomass (Ash Free Dry Weight [g]) |            |          |               |             | Total |
|---------|-----------------------------------|------------|----------|---------------|-------------|-------|
|         | Annelida                          | Arthropoda | Mollusca | Echinodermata | Other Phyla |       |
| MA_ST01 | 0.099                             | 0.061      | 0.054    | 0.182         | 0.046       | 0.442 |
| MA_ST02 | 0.100                             | 0.037      | 0.010    | 0.024         | 0.002       | 0.173 |
| MA_ST03 | 0.108                             | 0.124      | 0.089    | 0.006         | 0.022       | 0.349 |
| MA_ST04 | 0.060                             | 0.063      | 0.055    | 0.247         | 0.351       | 0.775 |
| MA_ST05 | 0.247                             | 0.073      | 0.296    | 0.300         | 0.007       | 0.922 |
| MA_ST06 | 0.110                             | 0.063      | 0.001    | 0.006         | 0.005       | 0.184 |
| MA_ST07 | 0.506                             | 0.080      | 0.005    | 0.051         | 0.002       | 0.644 |
| MA_ST08 | 0.040                             | 0.097      | 0.055    | -             | 0.002       | 0.195 |
| MA_ST09 | 0.080                             | 0.042      | 0.006    | 0.358         | 0.010       | 0.497 |
| MA_ST10 | 0.150                             | 0.132      | 1.138    | -             | 0.005       | 1.425 |
| MA_ST11 | 0.143                             | 0.141      | 0.068    | 0.055         | 0.057       | 0.464 |
| MA_ST12 | 0.095                             | 0.112      | 0.021    | 0.270         | 0.028       | 0.525 |
| MA_ST13 | 0.063                             | 0.048      | 0.001    | 0.027         | 0.006       | 0.145 |
| MA_ST14 | 0.103                             | 0.254      | 0.052    | 0.006         | 0.012       | 0.427 |
| MA_ST15 | 0.520                             | 0.091      | 0.718    | 0.294         | 0.008       | 1.632 |
| MA_ST16 | 0.192                             | 0.060      | 0.044    | 0.168         | 0.053       | 0.516 |
| MA_ST17 | 0.224                             | 0.058      | 0.023    | 0.007         | 0.028       | 0.339 |
| MA_ST18 | 0.145                             | 0.086      | 0.142    | 0.002         | 0.107       | 0.482 |
| MA_ST19 | 0.182                             | 0.076      | 0.055    | 0.201         | 0.033       | 0.547 |
| MA_ST20 | 0.215                             | 0.038      | 0.042    | 0.007         | 0.067       | 0.370 |
| MA_ST21 | 0.076                             | 0.009      | -        | -             | 0.013       | 0.098 |
| MA_ST22 | 0.073                             | 0.026      | 0.032    | 0.001         | 0.014       | 0.145 |
| MA_ST23 | 0.373                             | 0.132      | 0.189    | 0.005         | 0.018       | 0.716 |
| MA_ST24 | 0.209                             | 0.026      | 0.026    | 0.001         | < 0.001     | 0.262 |
| MA_ST25 | 0.054                             | 0.049      | 0.403    | 0.002         | 0.042       | 0.550 |
| MA_ST27 | 0.065                             | 0.064      | 0.049    | 0.342         | 0.009       | 0.529 |
| MA_ST29 | 0.053                             | 0.183      | -        | 0.001         | 0.007       | 0.244 |
| MA_ST30 | 0.097                             | 0.051      | 0.019    | 0.002         | 0.099       | 0.268 |
| MA_ST31 | 0.037                             | 0.020      | 0.015    | < 0.001       | 0.014       | 0.087 |
| MA_ST32 | 0.010                             | 0.020      | 0.091    | 0.261         | 0.002       | 0.383 |
| MA_ST33 | 0.073                             | 0.048      | < 0.001  | -             | 0.024       | 0.145 |
| MA_ST34 | 0.009                             | 0.022      | 0.053    | -             | 0.001       | 0.085 |
| MA_ST35 | 0.013                             | 0.002      | 0.005    | 0.001         | 0.001       | 0.021 |
| MA_ST36 | 0.054                             | 0.018      | 0.004    | -             | 0.028       | 0.104 |
| MA_ST37 | 0.139                             | 0.264      | 0.035    | 0.075         | 0.056       | 0.568 |

| Station  | Biomass (Ash Free Dry Weight [g]) |              |              |               |              | Total             |
|--|-----------------------------------|--------------|--------------|---------------|--------------|-------------------|
|  | Annelida                          | Arthropoda   | Mollusca     | Echinodermata | Other Phyla  |                   |
| MA_ST38  | 0.021                             | 0.005        | < 0.001      | 0.005         | < 0.001      | 0.031             |
| MA_ST39  | 0.013                             | 0.006        | < 0.001      | 0.004         | 0.003        | 0.026             |
| MA_ST40  | 0.419                             | 0.070        | -            | 0.002         | 0.301        | 0.792             |
| MA_ST41  | 0.089                             | 0.123        | 0.320        | 0.004         | 0.006        | 0.542             |
| MA_ST42  | 0.219                             | 0.059        | 1.203        | 0.923         | 0.263        | 2.666             |
| MA_ST43  | 0.121                             | 1.054        | 0.052        | 0.049         | 0.001        | 1.276             |
| MA_ST44  | 0.064                             | 0.005        | 0.017        | -             | -            | 0.085             |
| MA_ST45  | 0.144                             | 0.048        | 0.004        | -             | -            | 0.195             |
| MA_ST46  | 0.028                             | 0.002        | 0.004        | 0.002         | -            | 0.035             |
| MA_ST47  | 0.034                             | 0.013        | 0.067        | 0.010         | -            | 0.123             |
| MA_ST48  | 0.059                             | 0.051        | 0.098        | 0.011         | -            | 0.218             |
| MA_ST49  | 0.537                             | 0.057        | 1.083        | 0.003         | -            | 1.681             |
| MA_ST50  | 0.080                             | 0.004        | 0.002        | 0.009         | -            | 0.095             |
| MA_ST51  | 0.027                             | 0.026        | 0.010        | 0.011         | -            | 0.075             |
| MA_ST52  | 0.066                             | 0.006        | 0.013        | < 0.001       | -            | 0.086             |
| MA_ST54  | 0.061                             | 0.038        | 0.054        | 0.009         | -            | 0.162             |
| MA_ST55  | 0.163                             | 0.110        | -            | -             | -            | 0.273             |
| MA_ST56  | 0.003                             | 0.006        | 1.334        | < 0.001       | -            | 1.343             |
| MA_ST57  | 0.073                             | 0.002        | 0.001        | -             | -            | 0.076             |
| MA_ST58  | 0.105                             | 0.139        | 0.130        | < 0.001       | -            | 0.374             |
| MA_ST59  | < 0.001                           | -            | -            | -             | -            | < 0.001           |
| MA_ST60  | 0.032                             | < 0.001      | 0.004        | 0.508         | 0.000        | 0.544             |
| MA_ST61  | 0.391                             | 0.387        | 0.041        | 0.414         | 0.233        | 1.465             |
| MA_ST63  | 0.021                             | 0.003        | 0.071        | < 0.001       | 0.009        | 0.104             |
| MA_ST64  | 0.020                             | 0.006        | 0.001        | 0.347         | < 0.001      | 0.374             |
| MA_ST65  | 0.029                             | 0.002        | < 0.001      | 0.022         | -            | 0.053             |
| MA_ST66  | 0.004                             | 0.012        | -            | -             | -            | 0.016             |
| <b>Minimum</b>   | <b>&lt; 0.001</b>                 | <b>-</b>     | <b>-</b>     | <b>-</b>      | <b>-</b>     | <b>&lt; 0.001</b> |
| <b>Maximum</b>   | <b>0.537</b>                      | <b>1.054</b> | <b>1.334</b> | <b>0.923</b>  | <b>0.351</b> | <b>2.666</b>      |
| <b>Median</b>  | <b>0.078</b>                      | <b>0.049</b> | <b>0.029</b> | <b>0.006</b>  | <b>0.005</b> | <b>0.306</b>      |
| <b>Mean</b>  | <b>0.122</b>                      | <b>0.079</b> | <b>0.134</b> | <b>0.084</b>  | <b>0.032</b> | <b>0.451</b>      |
| <b>SD</b>  | <b>0.128</b>                      | <b>0.144</b> | <b>0.303</b> | <b>0.167</b>  | <b>0.072</b> | <b>0.503</b>      |
| Notes  |                                   |              |              |               |              |                   |
| Biomass expressed as ash free dry weight in g/0.1 m <sup>2</sup> grab sample |                                   |              |              |               |              |                   |
| SD = Standard deviation  |                                   |              |              |               |              |                   |
| - = Absent from samples  |                                   |              |              |               |              |                   |



Notes  
 Biomass expressed as ash free dry weight in g/0.1 m<sup>2</sup> grab sample

Figure 4.18: Phyletic composition of biomass, Main Array



**Notes**  
 Biomass expressed as ash free dry weight in g/0.1 m<sup>2</sup> grab sample  
 Figure 4.19: Total biomass overlaid on bathymetry, Main Array



Table 4.12 summarises the mean biomass of taxonomic groups identified within each macrofaunal group and ungrouped station. The highest average biomasses were in groups A and B (0.541 g and 0.496 g, respectively). Within the groups, the phyla contributing the highest biomass varied from annelids in station MA\_ST59 (100 %), molluscs in group B (60.4 %), echinoderms in group C (49.1 %) and a fairly equally split between annelids and molluscs in group A (28.4 % and 27.5 %, respectively) in group A. The lowest biomass was in station MA\_ST59 (< 0.001 g).

Table 4.12: Biomass within each grouping, Main Array

| Phyla   | Group A (▲)  |                            | Group B (■)         |                            |
|---|--------------|----------------------------|---------------------|----------------------------|
|   | Biomass      | Composition of Biomass [%] | Biomass             | Composition of Biomass [%] |
| Annelida  | 0.154        | 28.4                       | 0.044               | 8.8                        |
| Arthropoda  | 0.104        | 19.2                       | 0.029               | 5.8                        |
| Echinodermata   | 0.090        | 16.7                       | 0.122               | 24.6                       |
| Mollusca  | 0.149        | 27.5                       | 0.299               | 60.4                       |
| Other phyla   | 0.044        | 8.1                        | 0.002               | 0.5                        |
| <b>Total</b>  | <b>0.541</b> | <b>100</b>                 | <b>0.496</b>        | <b>100</b>                 |
| Phyla   | Group C (●)  |                            | Station MA_ST59 (★) |                            |
|   | Biomass      | Composition of Biomass [%] | Biomass             | Composition of Biomass [%] |
| Annelida  | 0.037        | 35.2                       | < 0.001             | 100.0                      |
| Arthropoda  | 0.007        | 6.4                        | 0.000               | 0.0                        |
| Echinodermata   | 0.052        | 49.1                       | 0.000               | 0.0                        |
| Mollusca  | 0.010        | 9.2                        | 0.000               | 0.0                        |
| Other phyla   | 0.000        | 0.1                        | 0.000               | 0.0                        |
| <b>Total</b>  | <b>0.105</b> | <b>100</b>                 | <b>&lt; 0.001</b>   | <b>100</b>                 |
| Notes<br>Biomass expressed as ash free dry weight in g/0.1 m <sup>2</sup> grab sample |              |                            |                     |                            |

## 4.4.2 Colonial Epifauna

### 4.4.2.1 Phyletic Composition

Table 4.13 summarises the epifaunal taxonomic groups identified across the survey area. A total of 33 epifaunal taxa were recorded across the survey area; bryozoans and hydrozoans accounted for the majority of the taxa reported (48.5 % and 45.5 % respectively), followed by the Cnidaria and Porifera, which equally contributed the remainder of the epifauna (3.0 % per phylum).

Table 4.13: Taxonomic groups of colonial epifauna, Main Array

| Taxonomic Group  | Number of Taxa | Composition of Taxa [%] |
|--|----------------|-------------------------|
| Bryozoan   | 16             | 48.5                    |
| Cnidarian  | 1              | 3.0                     |
| Hydrozoan  | 15             | 45.5                    |
| Porifera   | 1              | 3.0                     |
| <b>Total</b>   | <b>33</b>      | <b>100</b>              |
| Notes<br>Macrofaunal samples were processed through a 0.5 mm sieve |                |                         |

#### 4.4.2.2 Community Statistics

Table 4.14 presents the number of taxa and individuals identified from each station and Figure 4.20 spatially presents this data across the survey area. Station MA\_ST42 contained the highest number of colonial epifauna (12 taxa) and only two other stations had more than 10 taxa (MA\_ST03 and MA\_ST61 having 11 taxa and 10 taxa, respectively). No epifauna were recorded from 17 of the 62 stations sampled.

Table 4.14: Number of epifaunal taxa (0.1 m<sup>2</sup>), Main Array

| Station | Number of Taxa |
|---------|----------------|
| MA_ST01 | 4              |
| MA_ST02 | 6              |
| MA_ST03 | 11             |
| MA_ST04 | 7              |
| MA_ST05 | 5              |
| MA_ST06 | 5              |
| MA_ST07 | 3              |
| MA_ST08 | 5              |
| MA_ST09 | 8              |
| MA_ST10 | 8              |
| MA_ST11 | 6              |
| MA_ST12 | 1              |
| MA_ST13 | 5              |
| MA_ST14 | 4              |
| MA_ST15 | 6              |
| MA_ST16 | 4              |
| MA_ST17 | 3              |
| MA_ST18 | 3              |
| MA_ST19 | 2              |

| Station | Number of Taxa |
|---------|----------------|
| MA_ST20 | 6              |
| MA_ST21 | -              |
| MA_ST22 | 2              |
| MA_ST23 | 5              |
| MA_ST24 | -              |
| MA_ST25 | 3              |
| MA_ST27 | 6              |
| MA_ST29 | 3              |
| MA_ST30 | 6              |
| MA_ST31 | -              |
| MA_ST32 | 2              |
| MA_ST33 | 5              |
| MA_ST34 | 2              |
| MA_ST35 | -              |
| MA_ST36 | 2              |
| MA_ST37 | -              |
| MA_ST38 | 1              |
| MA_ST39 | -              |
| MA_ST40 | 3              |
| MA_ST41 | -              |
| MA_ST42 | 12             |
| MA_ST43 | 8              |
| MA_ST44 | 2              |
| MA_ST45 | 5              |
| MA_ST46 | 3              |
| MA_ST47 | -              |
| MA_ST48 | 5              |
| MA_ST49 | 5              |
| MA_ST50 | -              |
| MA_ST51 | 2              |
| MA_ST52 | 3              |
| MA_ST54 | 2              |
| MA_ST55 | -              |
| MA_ST56 | -              |
| MA_ST57 | -              |
| MA_ST58 | 1              |



| Station                                  | Number of Taxa |
|--|----------------|
| MA_ST59                                  | -              |
| MA_ST60                                  | 1              |
| MA_ST61                                  | 10             |
| MA_ST63                                  | -              |
| MA_ST64                                  | -              |
| MA_ST65                                  | -              |
| MA_ST66                                  | -              |
| <b>Minimum</b>                           | -              |
| <b>Maximum</b>                           | 12             |
| <b>Median</b>                            | 3              |
| <b>Mean</b>                              | 3              |
| <b>Standard Deviation</b>                | 3              |
| Notes<br>- = Epifauna absent from sample |                |




Figure 4.20: Number of epifaunal taxa per station (0.1 m<sup>2</sup>) overlaid on bathymetry, Main Array

### 4.4.2.3 Characteristic Taxa

Table 4.15 summarises the characteristic epifaunal taxa identified in grab samples across the survey area deduced from the infaunal and solitary epifauna analysis. Only the 11 taxa that occurred in 12.9 % or more of samples are displayed. Appendix F represents the full epifaunal dataset. The hydrozoan family *Campanulariidae* was ranked the most dominant colonial epifauna species across the survey area with a frequency of 43.6 % followed by the hydrozoan *Hydrallmania falcata* and bryozoan *Schizomavella*, which each had frequencies of 40.3 %. The remaining dominant epifaunal taxa, occurring at between 12.9 % and 21.0 % of the stations, comprised sponges (Porifera), four hydroid species, the cnidarian *Alcyonium digitatum*, and two bryozoans.

Table 4.15: Most frequently occurring colonial epifaunal taxa, Main Array

| Taxon   | Frequency Rank | Frequency [%] |
|---|----------------|---------------|
| <i>Campanulariidae</i>  | 1              | 43.5          |
| <i>Hydrallmania falcata</i>   | 2              | 40.3          |
| <i>Schizomavella</i>  | 2              | 40.3          |
| <i>Alcyonium digitatum</i>  | 3              | 21.0          |
| <i>Sertularia cupressina</i>  | 4              | 19.4          |
| PORIFERA  | 5              | 14.5          |
| <i>Abietinaria</i>  | 5              | 14.5          |
| <i>Escharella immersa</i>   | 5              | 14.5          |
| <i>Tubularia</i>  | 6              | 12.9          |
| <i>Phialella quadrata</i>   | 6              | 12.9          |
| <i>Vesicularia spinosa</i>  | 6              | 12.9          |
| Notes<br>Frequency rank is calculated based on frequency (i.e. percentage of stations from which each taxon was recorded) |                |               |

## 4.5 Seabed Habitats and Biotopes

### 4.5.1 Biotope Classifications

The physical and biological characteristics of the multivariate groups apparent within the macrofaunal community (see Section 4.4.1.3) were considered in conjunction with the photographic data for habitat classification. Soft sediment habitats are often defined on the sediment type and infaunal community composition. Therefore, soft sediments within the survey area have been predominantly classified using data from grab samples (specifically the PSD and macrofaunal data), with the photographic data analysis providing additional habitat information. Habitats comprising hard substrates, where grab sampling was not achieved, have been classified using photographic data only.

The seabed sediments observed across the survey area ranged from sandy gravel to gravelly sand. Within the western part of the survey area, where more homogenous, medium SSS

reflectivity was observed, the sediment consisted of gravelly sand and sandy gravel. This sediment type was classified as the EUNIS biotope complex 'Circalittoral coarse sediment (A5.14)', which was present at 50 stations; upon interpretation of physical and biological parameters from grab samples 5 of these stations were further classified as the biotope '*Branchiostoma lanceolatum* in circalittoral coarse sand with shell gravel (A5.145)'. Sand with varying proportions of shell fragments correlated to areas of low reflectivity on the SSS data, within the eastern part of the survey area. This sediment type has been classified as the EUNIS habitat 'Sublittoral sand' (A5.2) and was present at 12 stations. When grab sample data were considered it was possible to further refine the classification of 11 of these 12 stations to the biotope '*Nephtys cirrosa* and *Bathyporeia* spp. in infralittoral sand (A5.233)'.

Table 4.16 summarises the hierarchy of the assigned EEA (2019) classifications, and equivalent JNCC (2015) classifications. Figure 4.21 spatially presents the extrapolated distribution of the high level biotopes defined across the survey area. The survey area was heterogeneous with two habitats, two biotope complexes and two biotopes assigned to the transects and stations surveyed. Biotopes assigned are relevant to the time of year, with this survey completed in late summer.

Table 4.17 summarises the physical and biological parameters characteristics of the biotopes assigned, along with example photographs. Appendix G provides further example photographs. Sections 4.5.1.1 to 4.5.1.4 provide detailed descriptions of each biotope.

Table 4.16: Habitat classifications, Main Array

| EUNIS (EEA, 2019) Habitat Classification  |                            |                                     |  |   | Equivalent JNCC (2015) Classification  |
|---|----------------------------|-------------------------------------|--|---|--|
| Environment Level 1   | Broad Habitat Level 2      | Habitat Level 3                     | Biotope Complex Level 4                | Biotopes and sub-biotopes Level 5 & 6   |  |
| A<br>Marine   | A5<br>Sublittoral sediment | A5.1<br>Sublittoral coarse sediment | A5.14<br>Circalittoral coarse sediment | A5.145<br><i>Branchiostoma lanceolatum</i> in circalittoral coarse sand with shell gravel | SS.SCS.CCS.Blan<br><i>Branchiostoma lanceolatum</i> in circalittoral coarse sand with shell gravel |
|   |                            | A5.2<br>Sublittoral sand            | A5.23 Infralittoral fine sand          | A5.233 <i>Nephtys cirrosa</i> and <i>Bathyporeia</i> spp. in infralittoral sand           | SS.SSa.IFiSa.NcirBat<br><i>Nephtys cirrosa</i> and <i>Bathyporeia</i> spp. in infralittoral sand   |
| <p>Notes</p> <p>EUNIS = European Nature Information System</p> <p>EEA = European Environment Agency</p> <p>JNCC = Joint Nature Conservation Committee</p> |                            |                                     |  |   |  |

#### 4.5.1.1 Circalittoral Coarse Sediment (A5.14)

The biotope complex 'Circalittoral coarse sediment' (A5.14) is defined as tide-swept circalittoral coarse sands, gravel and shingle generally in depths of over 15 m depth. This habitat may be found in tidal channels of marine inlets, along exposed coasts and offshore. This habitat, as with shallower coarse sediments, may be characterised by robust infaunal

polychaetes, mobile crustacea and bivalves. Certain species of sea cucumber (e.g. *Neopentadactyla*) may also be prevalent in these areas along with the lancelet *Branchiostoma lanceolatum* (EEA, 2019).

This biotope complex was assigned to 45 stations (Table 4.17). It was the dominant habitat type defined in the western part of the survey area, but there was also an isolated occurrence of the habitat at station MA\_ST58, in the eastern part of the area (Figure 4.21). This sub-biotope was reported from depths ranging from 27 m BSL to 44 m BSL.

Sediment characteristics from grab samples described the seabed at these stations as gravelly muddy sand to sandy gravel, which corresponded with the video analysis descriptions (Table 4.17). The sediments of this biotope were grouped within cluster A of the multivariate analysis of PSD data (Figure 4.4 and Table 4.5).

This biotope complex also corresponded to cluster A of the multivariate analysis of infaunal data, which recorded species representative of 'Circalittoral coarse sediment'. The infauna identified from this biotope was typical of coarse sands and gravel, being dominated by *U. marina*, accompanied by *Phoronis* sp. and *A. paucibranchiata*. Cross-referencing of this community to the EUNIS habitat classification failed to identify an exact match of infaunal community at biotope or sub-biotope level.

The epibiota at these stations included hydrozoan/bryozoa, soft coral (*A. digitatum*), brittlestars (*O. fragilis*), starfish (*Asterias rubens*) and tube-building polychaetes (Serpulidae). Over a section of the transect undertaken at station MA\_ST12 a bed of brittlestars (*Ophiothrix fragilis*) was observed. This isolated habitat patch could be considered an example of the biotope '*Ophiothrix fragilis* and/or *Ophiocomina nigra* brittlestar beds on sublittoral mixed sediment' (A5.445). As this was only an isolated occurrence however and was deemed to not extend over the greater than 25 m<sup>2</sup> threshold for biotope classification (Parry, 2019) it has not been separately defined in this report.

#### 4.5.1.2 *Branchiostoma lanceolatum* in Circalittoral Coarse Sand with Shell Gravel (A5.145)

The EUNIS biotope '*Branchiostoma lanceolatum* in circalittoral coarse sand with shell gravel (A5.145)' is defined as gravel and coarse sand with shell gravel that may support a significant population of *Branchiostoma lanceolatum*. Other conspicuous infauna may include *Echinocyamus pusillus*, *Glycera lapidum*, *Polygordius* and *P. remota*. Sessile epifauna are typically a minor component of this community (EEA, 2019).

This biotope categorises the community associated with cluster B of the multivariate analysis of macrofaunal data, namely that found at stations MA\_ST32, MA\_ST55, MA\_ST56, MA\_ST63 and MA\_ST64 (Figure 4.21). These stations were distributed across the survey area, occurring in the west (stations MA\_ST32), near the middle (stations MA\_ST55 and MA\_ST56) and towards the eastern edge (stations MA\_ST63 and MA\_ST64) of the survey area. This biotope was identified in water depths of 27 m BSL to 36 m BSL.



Sediment characteristics from grab samples described the seabed at the stations as sand to gravelly sand (Table 4.3), with the seabed described from the video analysis as sand with varying proportions of gravel, shell and shell fragments (Table 4.17). The sediments of this biotope were grouped within clusters A and B of the multivariate analysis of PSD data (Figure 4.4 and Table 4.5).

Infaunal analysis showed that the cluster B stations were dominated by the polychaetes *P. remota* and *Polygordius*, having the same abundance (Table 4.10) followed by nemertea, which are characterising taxa of this biotope.

The epibiota at stations MA\_ST55, MA\_ST56, MA\_ST63 and MA\_ST64 were sparse, as expected for sandy sediments. The most regularly observed taxa included brittlestars (Ophiuroidea), hermit crabs (Paguridae) and starfish (Asteroidea).

#### 4.5.1.3 Sublittoral Sand (A5.2)

The EUNIS habitat 'Sublittoral sand (A5.2)' is defined as clean medium to fine sands or non-cohesive slightly muddy sands on open coasts, offshore or in estuaries and marine inlets. Such habitats are often subject to a degree of wave action or tidal currents which restrict the silt and clay content to less than 15 %. This habitat is characterised by a range of taxa including polychaetes, bivalve molluscs and amphipod crustacea (EEA, 2019).

This biotope categorises the infaunal community at station MA\_ST59 (Figure 4.21). This station was located in the east of the survey area in an area of mobile sediment and water depth of 38 m BSL. Sediment characteristics from the grab samples described the seabed at station MA\_ST59 as sand (Table 4.3).

Infaunal analysis from station MA\_ST59 recorded only two individuals and taxa (*S. parapari* and *P. strombus*).

The epibiota at this station included brittlestars (Ophiuridae), soft coral (*A. digitatum*) and faunal turf (Hydrozoa/Bryozoa).

#### 4.5.1.4 *Nephtys cirrosa* and *Bathyporeia* spp. in Infralittoral Sand (A5.233)

The EUNIS biotope '*Nephtys cirrosa* and *Bathyporeia* spp. in infralittoral sand (A5.233)' is defined as well-sorted medium and fine sands characterised by *Nephtys cirrosa* and *Bathyporeia* spp. which occur in the shallow sublittoral to at least 30 m depth. This biotope occurs in sediments subject to physical disturbance, as a result of wave action (and occasionally strong tidal streams). The magelonid polychaete may be frequent in this biotope in more sheltered, less tide swept areas. The faunal diversity of this biotope is considerably reduced compared to less disturbed biotopes (such as unit A5.242) and for the most part consists of the more actively swimming amphipods (EEA, 2019).

This biotope categorises the infaunal and epifaunal community recorded from stations within cluster C of the multivariate analysis of macrofaunal data (stations MA\_ST35, MA\_ST38, MA\_ST46, MA\_ST47, MA\_ST50, MA\_ST51, MA\_ST52, MA\_ST57, MA\_ST60, MA\_ST65 and




MA\_ST66) (Figure 4.21). These stations and transects were located towards the south and east of the survey area in water depths of 23 m BSL to 36 m BSL.



This biotope was recorded in association with areas of mobile sediment, as evident from sand waves and megaripples evident from MBES data, and low/medium SSS reflectivity. Sediment composition was variable with five stations described by the Folk description of sand and six stations described as gravelly sand (Table 4.3); from photographic data the seabed was described as sand with varying proportions of gravel, shells and shell fragments (Table 4.17).

Infaunal analysis showed that the cluster C stations were dominated by the amphipod *B. elegans*, with the polychaete *S. bombyx* the second most abundant taxon reported. Other prominent infauna in this cluster C were the annelid *S. goniocephala* and the echinoderm *E. pusillus* (Table 4.10).


The epibiota at these stations was sparse, as expected from sandy sediments. The most regularly observed taxa included faunal turf (Hydrozoa, including *Nemertesia antennina*, and Bryozoa), hermit crabs (Paguridae), starfish (Asteroidea, including *A. rubens*) and sea urchins (Echinoidea).

Table 4.17: Summary of EUNIS habitat classifications, Main Array

| Habitat Classification (EEA, 2019)     | Distribution   | Physical Characteristics                         | Biological Characteristics   | Example Photograph   |
|--|--|--|--|--|
| A5.14<br>Circalittoral coarse sediment | Cluster A (▲)*†:<br>MA_ST01, MA_ST02, MA_ST03, MA_ST04, MA_ST05, MA_ST06, MA_ST07, MA_ST08, MA_ST09, MA_ST10, MA_ST11, MA_ST12, MA_ST13, MA_ST14, MA_ST15, MA_ST16, MA_ST17, MA_ST18, MA_ST19, MA_ST20, MA_ST21, MA_ST22, MA_ST23, MA_ST24, MA_ST25, MA_ST27, MA_ST29, MA_ST30, MA_ST31, MA_ST33, MA_ST34, MA_ST36, MA_ST37, MA_ST39, MA_ST40, MA_ST41, MA_ST42, MA_ST43, MA_ST44, MA_ST45, MA_ST48, MA_ST49, MA_ST54, MA_ST58 and MA_ST61 | Mean gravel*:<br>19.88 %                         | Mean number of taxa*:<br>44 per 0.1 m <sup>2</sup>   |  |
|  |  | Mean sand*:<br>75.19 %                           | Mean number of individuals*:<br>21 per 0.1 m <sup>2</sup>  |  |
|  |  | Mean mud*:<br>4.94 %                             | Characteristic infaunal taxa*:<br><i>Urothoe marina</i> , <i>Phoronis</i> sp., <i>Aonides paucibranchiata</i> and <i>Poecilochaetus serpens</i>  |  |
|  |  | 'Gravelly sand with shells and shell fragments'† | Characteristic epifaunal taxa*:<br>Hydrozoan/bryozoa, soft coral ( <i>Alcyonium digitatum</i> ), brittlestars ( <i>Ophiothrix fragilis</i> ), starfish ( <i>Asterias rubens</i> ) and tube-building polychaetes (Serpulidae) |  |
|  |  | Bathymetry:<br>27 m to 44 m BSL                  | Mean Biomass*:<br>0.541  |  |

| Habitat Classification (EEA, 2019)  | Distribution   | Physical Characteristics   | Biological Characteristics  | Example Photograph   |
|---|--|--|---|--|
| A5.145<br><i>Branchiostoma lanceolatum</i> in circalittoral coarse sand with shell gravel | Cluster B (■)*†:<br>MA_ST32, MA_ST55,<br>MA_ST56, MA_ST63 and<br>MA_ST64 | Mean gravel*:<br>4.86 %  | Mean number of taxa*:<br>21 per 0.1 m <sup>2</sup>  |   |
|   |  | Mean sand*:<br>95.14 %   | Mean number of individuals*:<br>1 per 0.1 m <sup>2</sup>  |  |
|   |  | Mean mud*:<br>0.00 %   | Characteristic infaunal taxa*:<br><i>Pisone remota</i> , <i>Polygordius</i> ,<br>Nemertea and <i>Spio symphyta</i>          |  |
|   |  | †Sand with varying proportions of gravel, shell and shell fragments† | Characteristic epifaunal taxa*:<br>brittlestars (Ophiuroidea), hermit crabs (Paguridae) and starfish (Asteroidea)           |  |
|   |  | Bathymetry:<br>27 m to 36 m BSL                                      | Mean Biomass*:<br>0.496   |  |
| A5.2<br>Sublittoral sand  | Ungrouped station MA_ST59 (*)*:  | Mean gravel*:<br>0.97 %  | Mean number of taxa*:<br>1 per 0.1 m <sup>2</sup>   |  |
|   |  | Mean sand*:<br>99.03 %   | Mean number of individuals*:<br>1 per 0.1 m <sup>2</sup>  |  |
|   |  | Mean mud*:<br>0.00 %   | Characteristic infaunal taxa*:<br><i>Syllis paraparari</i> and <i>Phascolion strombus</i>                                   |  |
|   |  | †Sand with varying proportions of shells and shell fragments†        | Characteristic epifaunal taxa*:<br>Hydrozoan/bryozoa, soft coral ( <i>Alcyonium digitatum</i> ), brittlestars (Ophiuridae), |  |
|   |  | Bathymetry:<br>38 m BSL  | Mean Biomass*:<br>0.0004  |  |



| Habitat Classification (EEA, 2019)   | Distribution   | Physical Characteristics   | Biological Characteristics   | Example Photograph  |
|--|--|--|--|---|
| A5.233<br><i>Nephtys cirrosa</i> and <i>Bathyporeia</i> spp. in infralittoral sand   | Cluster C (●)*†:<br>MA_ST35, MA_ST38, MA_ST46, MA_ST47, MA_ST50, MA_ST51, MA_ST52, MA_ST57, MA_ST60, MA_ST65 and MA_ST66 | Mean gravel*:<br>5.65 %  | Mean number of taxa*:<br>18 per 0.1 m <sup>2</sup>   |  |
|  |  | Mean sand*:<br>94.20 %   | Mean number of individuals*:<br>2 per 0.1 m <sup>2</sup>   |   |
|  |  | Mean mud*:<br>0.15 %   | Characteristic infaunal taxa*:<br><i>Bathyporeia</i> spp.,<br><i>Spiophanes bombyx</i> ,<br><i>Spio goniocephala</i> and<br><i>Echinocyamus pusillus</i> |   |
|  |  | 'Sand with varying proportions of gravel, shell and shell fragments''† | Characteristic epifaunal taxa*:<br>Hermit crabs (Paguridae)  |   |
|  |  | Bathymetry:<br>23 m to 36 m BSL  | Mean Biomass*:<br>0.105  |   |
| <p><b>Notes</b></p> <p>Biomass expressed as ash free dry weight in g/0.1 m<sup>2</sup> grab sample</p> <p>EEA = European Environment Agency</p> <p>† = Data from grab analysis. Mean values generated for sample/station data within multivariate grouping of macrofaunal community</p> <p>‡ = Data from photographic analysis</p> |  |  |  |   |




Figure 4.21: The spatial distribution of EUNIS (EEA, 2019) habitat classifications, Main Array



## 4.5.2 Potential Sensitive Habitats and Species

### 4.5.2.1.1 e.g. Subtidal Sands and Gravels

The seabed within the survey area was classified as the EUNIS habitats 'Circalittoral coarse sediment (A5.14), 'Sublittoral sand' (A5.2), '*Nephtys cirrosa* and *Bathyporeia* spp. in infralittoral sand' (A5.233) and '*Branchiostoma lanceolatum* in circalittoral coarse sand with shell gravel' (A5.145). These habitats are encompassed within 'Subtidal sands and gravels', a priority habitat within UK waters.

### 4.5.2.1.2 Other Potentially Sensitive Habitats and Species

No other Annex I habitats or Annex II species, OSPAR threatened and/or declining species and habitats or UK Biodiversity Action Plan priority habitats and species (OSPAR, 2008; BRIG, 2011, JNCC, 2018; JNCC, 2019a; 2019b) were observed within the survey area.



---

## 5. Discussion

### 5.1 Sediment Characterisation

The general physical and chemical characteristics of sediment particles have a significant effect on how other chemical components and biological species interact with seabed sediments. For example, the silt/clay fraction is known to adsorb petroleum hydrocarbons/heavy metals from seawater and through this pathway, these chemicals become incorporated into the sediment system (Meyers & Quinn, 1973). Granulometry data can therefore be critical when interpreting chemical and biological data obtained in this type of benthic study. In addition, since waste discharges (such as drill cuttings) often possess significantly different physical characteristics from the natural sediments present in the area, such data may also provide some information on the spread of discharged material.

With regard to macrofaunal communities, the species distributions and community structure can be greatly influenced by the nature of the sediment, which represents the effects of a complex set of hydrological factors, such as water movement, turbulence and suspended load, at one particular point in time. Some animals have a behavioural preference for sediment of a particular grain size (Meadows, 1964; Gray, 1981), while this factor and organic matter content are closely associated with other properties of the sediment such as density, porosity, permeability, oxygenation and bacterial count (Buchanan, 1984), all of which affect animal functions such as locomotion, attachment, tube construction and feeding. Specifically, the proportion of fine (silt/clay) material often influences the distribution of macrofaunal communities.

Sediment descriptions using the Folk description (1954) categorised the seabed of the main array as predominantly gravelly sand (47 stations) or sand (10 stations), with the remaining stations described as gravelly muddy sand (3 stations), muddy sandy gravel (station MA\_ST61) or sandy gravel (station MA\_ST02). When the graphical mean particle size of the samples was considered, this was classified as Wentworth (1922) sediment descriptions which ranged from 'very coarse sand' to 'medium sand'. There was clear spatial pattern in sediment type across the survey area, which would be linked to variations in seabed acoustic character/morphology. Within the western part of the survey area, where more homogenous, medium SSS reflectivity was observed, sediments contained a moderate proportion of gravel and a small fines fraction. Sediments within this area were grouped within cluster A of the multivariate analysis of PSD data. Sand waves and mega ripples were evident from MBES data acquired in the eastern part of the survey area in association with generally low SSS reflectivity. Sediments in this area comprised clean sands (no fines) with a low gravel content (clusters B and C of the multivariate analysis of PSD data). Multivariate cluster analysis reinforced these results, with three significant clusters and one ungrouped station. PCA refined these results and demonstrated that the factors driving the differentiation of the clusters were the proportions of medium sand, coarse sand and coarse pebble sediment.

The spatial patterns evident in sediment composition and the clear linkage of these to the geophysical characteristics of the seafloor, suggest that particle size is likely to be influenced by the degree of sediment mobility. This is likely the result of a variable degree of exposure to tidal streams and wave action, which will be influenced by water depths across the area.

## 5.2 Sediment Chemistry

### 5.2.1 Sediment Hydrocarbons

#### 5.2.1.1 Aromatic Hydrocarbons

PAHs are widely spread in the environment (Butler et al., 1984) with natural sources occurring primarily through synthesis by plants (Neff, 1979; Sims & Overcash, 1983), related to natural seeps of petroleum (NRC, 1983; Kennicutt et al., 1988) and to formation during natural forest and prairie fires (Youngblood & Blumer, 1975; Wakeham et al., 1979). By far the greatest proportion of PAHs released into the environment are formed during fossil fuel combustion and anthropogenic forest and agricultural fires (Edwards, 1983; Sims & Overcash, 1983; Haritash & Kaushik, 2009). PAHs primarily enter marine sediments from atmospheric and riverine inputs; accumulation on the surface tends to adsorb into suspended inorganic and organic particulate matter, ultimately settling on the seabed where they accumulate to relatively high concentrations (Latimer & Zheng, 2003; Culotta et al., 2006).

Monitoring of aromatic hydrocarbon type and content is important due to the particularly toxic nature (mutagenic/carcinogenic) of several PAHs, particularly the heavier weight PAHs. The US EPA has identified 16 priority PAHs to be monitored (Keith, 2015) and the Coordinated Environmental Monitoring Programme (CEMP) specifies 9 PAHs of specific concern (OSPAR, 2014), which primarily reflect inputs from anthropogenic combustion sources.

Total 2 to 6 ring PAH concentrations were higher than the median concentration recorded from the SEA6 (Cefas, 2005) Irish Sea surveys (0.0237 µg/g) at six stations (MA\_ST04, MA\_ST12, MA\_ST22, MA\_ST25, MA\_ST43 and MA\_ST61); however, the median value from the current survey was broadly comparable to the SEA6 median value.

The individual US EPA 16 PAH concentrations were all below the CEMP ERL values, and therefore are unlikely to have any adverse effects on the macrofaunal community (Appendix E.1).

The source of the PAHs may be determined by investigation of the relative proportions of individual PAH concentrations (Neff, 1979; Budzinski et al., 1997; Yunker et al., 2002) as well as examining the overall distributions of parent and alkylated PAHs present. Pyrogenically (or pyrolytic) derived PAHs signatures (i.e. forest fires, etc.) are dominated by higher molecular weight compounds (mainly 4 to 6 ring) and are predominantly unsubstituted. In contrast, PAH formed during the slow geological maturation of petroleum, are dominated by alkylated, low molecular weight (mainly 1 to 3 ring) compounds (Neff, 1979; Stogiannidis & Laane, 2015). The proportion of petrogenically derived NPD to total aromatic

material present in these sediments (mean 35 %; Table 4.6) indicated a predominantly pyrolytic source for the aromatic compounds in the sediments.

## 5.2.2 Sediment Metals

### 5.2.2.1 Heavy and Trace Metals

Metals and metalloids occur naturally in the marine environment and are widely distributed in both dissolved and sedimentary forms. Some are essential to marine life while others have no biological function and therefore are toxic to numerous organisms at certain levels (Paez-Osuna & Ruiz-Fernandez, 1995; Boening, 1999). Metals can enter the environment via natural methods such as riverine transport, coastal discharges, geological weathering and atmospheric fallout (Brady et al., 2015). Other routes into marine sediments are from anthropogenic activities such as direct discharges from industrial activities.

Trace metal contaminants in the marine environment tend to form associations with the non-residual phases of mineral matter, such as iron and manganese oxides and hydroxides, metal sulphides, clays, organics and carbonates (Warren & Zimmerman, 1993; Dang et al., 2015; Wang et al., 2015). Non-residual trace metals are associated with more reactive and available sediment components through processes such as adsorption onto mineral surfaces and organic complexation. Metals associated with these more reactive phases are prone to various environmental interactions and transformations (physical, chemical and biological) potentially increasing their mobility and biological availability (Tessier et al., 1979; Warren & Zimmerman, 1993; Du Laing et al., 2009). Residual trace metals are defined as those that are part of the crystal structure of the component minerals and are generally unavailable to organisms (de Orte et al., 2018). Therefore, in monitoring trace metal contamination of the marine environment, it is important to distinguish the more mobile non-residual trace metals from the residual metals held tightly in the sediment lattice (Chester & Voutsinou, 1981), which are of comparatively lesser environmental significance because of their low reactivity and availability.

In this study, an analytical procedure involving the digestion of sediment in aqua regia was employed to analyse the elemental content of the sediments. The aqua regia digest releases for analysis the 'non-residual' heavy metals, which are not incorporated in the mineral matrix and are therefore potentially available for biological uptake.

The bioavailable metals concentrations in sediments were all below their respective Cefas action levels and the CEMP ERLs indicating that these metals are unlikely to have an adverse effect on the macrofaunal communities present.

## 5.3 Macrofaunal Communities

Seabed sediments provide support, protection and the food source for many macrofaunal species. The sediment macrofauna, most of which are infaunal (living within the sediment), are therefore particularly vulnerable to external influences that alter the sediments' physical, chemical or biological nature. Such infaunal animals are largely sedentary and are thus

unable to avoid unfavourable conditions. Each species has its own response and degree of sensitivity to changes in the physical and/or chemical environment and consequently the species composition and their relative abundance in a particular location provides a reflection of the health and condition of the immediate environment, both current and historical. The recognition that aquatic contaminant inputs may alter sediment characteristics, together with the relative ease of obtaining quantitative samples from specific locations, has led to the widespread use of infaunal communities in monitoring the impact of disturbances to the marine environment over a long period of time.

The phyletic composition for both taxa and abundance was variable throughout the survey area. Annelids were the most diverse and numerous phyla at all stations except for ten stations, where the dominant taxa were a combination of arthropods, molluscs, echinoderms and other phyla. Whilst annelids were the dominant phylum across the survey area, the most abundant taxon overall within the survey area was the arthropod *U. marina*.

The variation in the number of taxa and individuals was reflected in the diversity indices (Shannon-Wiener) and evenness (expressed as both as Pielou's (J) and Simpson's Index (1- $\lambda$ ), both of which showed high levels of variation across the survey area. A broad spatial trend in the number of taxa and individuals was observed, where typically higher numbers were observed in the west of the survey area.

The multivariate analysis of infauna showed three statistically significant clusters and one ungrouped station. Each of the infaunal groups were dominated by different key taxa, as well as having differing numbers of taxa and individuals. A spatial pattern was evident from the distribution of the infaunal groups across the survey area. Cluster A was predominately found in the west of the survey area, whereas clusters B and C were typically reported in the southern and eastern parts of the area (Figure 4.14). This was similar to the spatial trend observed in the sediment type across the survey area (Figure 4.5). Infaunal clusters broadly corresponded to the clusters produced from the multivariate analysis of the PSD data and both analyses showed clear association with the variations in seabed acoustic character/morphology evident from the MBES and SSS data. This suggested the infaunal community was likely to have been influenced by sediment type and the degree of sediment mobility, which may, in turn, have been linked to water depth, as this will likely determine the degree of exposure of benthic habitats to wave action and/or tidal streams.

The BIOENV algorithm in the PRIMER BEST routine was an additional technique used to identify the environmental variables that correlated significantly with the patterns outlined in community structures. BIOENV was run for a single and a combination of two and three variables. The single variable that correlated most strongly with the patterns in the macrofaunal community was granule (2000  $\mu\text{m}$  to 3000  $\mu\text{m}$ ) sediment ( $P \leq 0.05$ ;  $\rho = 0.685$ ). Two variables combined correlating with patterns in the macrofaunal community were very coarse sand and silt ( $P \leq 0.05$ ;  $\rho = 0.797$ ). When three variables were combined, granule, medium sand and silt ( $P \leq 0.05$ ;  $\rho = 0.836$ ) correlated with patterns in the macrofaunal community. Therefore, these results suggested that the variations in the macrofaunal

communities were being driven by sediment granulometry, which, as previously mentioned, was probably largely determined by exposure to wave action and/or tidal currents. BIOENV correlations run between chemical results and the patterns in the macrofaunal community provided no statistically significant results ( $P > 0.05$ ).

Stations within infaunal cluster A were described as poorly to very poorly sorted coarse sand and were dominated by fauna typical of this sediment type. Cross-referencing of this community to the EUNIS habitat classification failed to identify an exact match of infaunal community at biotope or sub-biotope level.

Infaunal cluster B was described as moderately well to well sorted coarse sand, containing fauna typical of circalittoral coarse sediment, characterised by the presence of the polychaetes *P. remota*, *Polygordius* and the echinoderm *E. pusillus*.

Stations within infaunal cluster C were dominated by faunal typical of well/moderately well sorted, medium to coarse sand sediments, with minimal mud content, which were present at these stations. These dominant taxa included the arthropods *B. elegans* and *B. gracilis* which are epistrate feeders (feed by scraping algae/bacteria off sand grains) and the polychaete *N. cirrosa* which is a psammophilous polychaete. These species show preference for wave exposed sand habitats and, by extension, any sediments subject to hydrodynamic disturbance (Tilin, 2016). The low mud content is typical of areas with moderate levels of exposure to tidal or wave action.

Ungrouped station MA\_ST59 was described as well sorted coarse sand. The paucity of fauna at this station (only one polychaete *S. parapari* and one sipunculid *P. strombus* were recorded) lead to this station being statistically differentiated from all other stations sampled. It is possible that the paucity of fauna identified may have resulted from recent disturbance of the sediments at this station, possibly as the result of natural hydrodynamic processes.

Biomass of the infauna showed variation in the samples at stations taken from across the survey area, likely due to the variation identified at stations. The overall biomass was highest within infauna cluster A, with annelids providing the highest contribution due to the high abundance of polychaetes. Cluster B had the second highest biomass, with molluscs providing the highest contribution. This was due to the presence of the mollusc *Asbjornsenia pygmaea* in an otherwise sparse station (MA\_ST56), which skewed the mean biomass towards molluscs even though they were not particularly abundant within cluster B.

Solitary epifauna were identified across four phyla; Bryozoa, Hydrozoa, Cnidaria and Porifera. Sediment cluster A had the highest diversity, containing taxa from all of the four phyla identified. The sediment associated with cluster A comprised sandy gravel to gravelly muddy sand with the highest gravel content (mean of 19.88 %) of all the sediment clusters identified. This gravel sized sediment, which includes shell and pebble material, will have provided suitable attachment substrata for the epifauna identified.

## 5.4 Seabed Habitats and Biotopes

When seabed photographic data, particle size data and macrofaunal data were considered, the EUNIS classifications; one habitat, two biotope complexes and two sub-biotopes were assigned to the station surveyed. The classifications were refined from the habitat classifications identified in the ECR Main array -Environmental Features Report (Volume 2 of this series).

The majority of the main array survey stations were classified as 'Circalittoral coarse sediment' (A5.14). The higher gravel content from the PSD results and the presence of species associated with coarse sediment (*Nemertea*, *S. bombyx* and *A. Lindstroemi*) are characteristic of this biotope complex. This community corresponded to cluster A of the multivariate analysis of macrofaunal data. Five stations, namely stations MA\_ST32, MA\_ST55, MA\_ST56, MA\_ST63 and MA\_ST64 (cluster B of the multivariate analysis,) had a higher sand content, with different dominant taxa, than those stations assigned to the 'Circalittoral coarse sediment' (A5.14) biotope complex. These stations were further classified into the biotope 'Branchiostoma lanceolatum in circalittoral coarse sand with shell gravel' (A5.145), due to characteristic taxa including the polychaetes *P. remota*, *Polygordius* and *H. elongata*, and the echinoderm *E. pusillus*.

Eleven stations were classified as the biotope complex 'Infralittoral fine sand' (A5.23) across the survey area, due to the high sand and low gravel/mud content and faunal assemblages being typical of clean sands with moderate exposure to wave or tidal action. Infaunal analysis showed similarities to the sub biotope complex 'Nephtys cirrosa and Bathyporeia spp. in infralittoral sand' (A5.233), whereby the dominant taxa at these stations included the annelid *N. cirrosa* and amphipods *B. gracilis* and *B. elegans* (cluster C of the multivariate analysis of infaunal data).

Ungrouped station MA\_ST59 was classified as the habitat 'Sublittoral sand' (A5.2) due to the sediment comprising of high sand content, minimal gravel and lacking fines content. The impoverished macrofaunal assemblage present at this station did not allow classification to biotope complex/biotope level.

As described for sediment characteristics and infaunal community structure, there was a clear spatial distribution in the habitat types present within the survey area and this could be linked to variations in SSS reflectivity and the seabed morphology evident from MBES data Figure 4.21.

The sediments observed throughout the survey area were identified as comprising the broadscale priority habitat 'subtidal sands and gravels'. However, this habitat is widely distributed and represented elsewhere in the UK Marine Protected Area (MPA) network (JNCC, 2019).



No other Annex I habitats or Annex II species, OSPAR threatened and/or declining species and habitats or UK Biodiversity Action Plan priority habitats and species observed within the survey area.

Table 5.1 provides a summary of the sensitive habitats that may occur in the survey area.

Table 5.1: Summary of sensitive habitats potentially present, Main Array

| Listed Feature  |                                | Relationship | Related Feature                     |                         |
|---|--------------------------------|--------------|-------------------------------------|-------------------------|
| Description   | Designation/Status             |              | Description                         | Designation/Status      |
| Subtidal sands and gravels  | Priority habitat; habitat FOCI | Contains     | Offshore subtidal sands and gravels | PMF; MPA search feature |
| <p>Notes</p> <p>FOCI = Feature of Conservation Interest</p> <p>PMF = Priority Marine Feature</p> <p>MPA = Marine Protected Area</p> |                                |              |                                     |                         |

## 6. Conclusions

The aim of this report has been to evaluate the existing physical, chemical and biological components in the marine environment within the survey area. A review of the environmental data in context with other cited studies from the region and estimated sediment effects threshold values (Cefas, 2005; Cefas, 2014; OSPAR, 2014) was also undertaken. Based on the overall assessment of the survey area, the following key conclusions can be stated:

The sediments within the survey area comprised mainly of sand, with varying proportions of gravel, and little to no fines. Using the BGS modified Folk (1954) sediment description, most stations were classified as gravelly sand (47 stations) or sand (10 stations). Multivariate cluster analysis demonstrated this variability in sediment fractions by grouping the stations into three significant clusters and one ungrouped station. The clear spatial pattern in sediment type, and its association with the seabed's geophysical characteristics, suggested that sediment type was influenced by hydrodynamic regime (variations in sediment mobility) and water depth.

The median total 2 to 6 ring PAH concentration was broadly comparable to the median concentration reported from the SEA6 Irish Sea surveys. All individual PAH concentrations were below their respective ERL values.

All metals concentrations were less than their respective Cefas guideline action levels (AL1 and AL2) and OSPAR ERL values.

The number of infaunal and solitary epifaunal taxa recorded from the grab samples varied across the Main Array survey area. Following multivariate statistical analysis, three separate benthic communities were differentiated. One of these was dominated by the amphipod *U. marina* and horseshoe worm *Phoronis* and was associated with coarse sediment (predominantly gravelly sand) areas. One was characterised by polychaetes typical of clean coarse sediments (*P. remota* and *Polygordius*) and nemerteans. The third community present was characterised by taxa typical of mobile sand sediments (the amphipods *B. elegans* and *B. gracilis* and polychaetes *S. bombyx* and *S. goniocephala*).

Four habitat types were identified within the survey area; 'Circalittoral coarse sediment' (A5.14), '*Branchiostoma lanceolatum* in circalittoral coarse sand with shell gravel' (A5.145), '*Nephtys cirrosa* and *Bathyporeia* spp. in infralittoral sand' (A5.233) and 'Sublittoral sand' (A5.2). The most widespread habitat being 'Circalittoral coarse sediment' (A5.14).

The UK priority habitat 'subtidal sands and gravels' was likely to be present within survey area, however this habitat is widely distributed within UK waters and is included within UK MPA networks. No other Annex I habitats or Annex II species, OSPAR threatened and/or declining species and habitats or UK Biodiversity Action Plan priority habitats and species were observed within the survey area.

## 7. References

- Balmer, J.E., Hung, H., Yu, Y., Letcher, R.J., & Muir, D.C. (2019). Sources and environmental fate of pyrogenic polycyclic aromatic hydrocarbons (PAHs) in the Arctic. *Emerging Contaminants*, 5, 128-142.
- Blott, S. (2010). *GRADISTAT Version 8.0: A grain size distribution and statistics package for the analysis of unconsolidated sediment by sieving or laser granulometer*. Berkshire: Kenneth Pye Associates.
- Boening, D.W. (1999). An evaluation of bivalves as biomonitors of heavy metals pollution in marine waters. *Environmental Monitoring and Assessment*, 55(3), 459-470.
- Brady, J.P., Ayoko, G.A., Martens, W.N., & Goonetilleke, A. (2015). Development of a hybrid pollution index for heavy metals in marine and estuarine sediments. *Environmental Monitoring and Assessment*, 187(5), 306.
- Buchanan, J.B. (1984). Sediment analysis. In: N.A. Holme and A.D. McIntyre (Eds). *Methods for the study of marine benthos*. 2nd edition. IBP Handbook No 16. Blackwell Scientific Publications, Oxford, London, Edinburgh, Boston, Palo Alto and Melbourne.
- Budzinski, H., Jones, I., Bellocq, J., Pierard, C., & Garrigues, P. (1997). Evaluation of sediment contamination by polycyclic aromatic hydrocarbons in the Gironde Estuary. *Marine Chemistry*, 58, 85-97.
- Butler, J.D., Butterworth, V., Kellow, S.C., & Robinson, H.G. (1984). Some observations on the polycyclic aromatic hydrocarbon (PAH) content of surface soils in urban areas. *The Science of the Total Environment*, 33, 75-85.
- Centre for Marine and Coastal Studies Limited [CMACS]. (2005). *Gwynt y Môr Offshore Wind farm. Marine ecology technical report*. Report Number: J3004/2005. Report to npower renewables.
- Centre for Environment, Fisheries and Aquaculture Science [Cefas]. (2005). *A review of the contaminant status of the Irish Sea*. CEFAS Contract report C2436.
- Chester, R., & Voutsinou, F.G. (1981). The initial assessment of trace metal pollution in coastal sediments. *Marine Pollution Bulletin*, 12, 84-91.
- Clarke, K.R., & Warwick, R.M. (2001). *Change in marine communities: an approach to statistical analysis and interpretation*. Natural Environmental Research Council.
- Clarke, K.R., & Gorley, R.N. (2015). *PRIMER v7: User Manual/Tutorial*, Plymouth: PRIMER-E.
- Clarke, K.R., Somerfield, P.J., & Gorley, R. N. (2008). Testing of null hypothesis in exploratory community analysis: similarity profiles and beta-environment linkage. *Journal of Experimental Marine Biology and Ecology*, 366, 56-69.

- Culotta, L., De Stefano, C., Gianguzza, A., Mannino, M.R., & Orecchio, S. (2006). The PAH composition of surface sediments from Stagnone coastal lagoon, Marsala (Italy). *Marine Chemistry*, 99(1-4), 117-127.
- Dang, D.H., Lenoble, V., Durrieu, G., Omanović, D., Mullot, J.U., Mounier, S., & Garnier, C. (2015). Seasonal variations of coastal sedimentary trace metals cycling: Insight on the effect of manganese and iron (oxy)hydroxides, sulphide and organic matter. *Marine Pollution Bulletin*, 92(1-2), 113-124.
- Dauvin, J.C., Alizier, S., Rolet, C., Bakalem, A., Bellan, G., Gesteira, J.G., Grimes, S., De-La-Ossa-Carretero, J.A., & Del-Pilar-Ruso, Y. (2012). Response of different benthic indices to diverse human pressures. *Ecological Indicators*, 12(1), 143-153.
- Davies, C.E., & Moss, D. (2004). *EUNIS habitat classification marine habitat types: revised classification and criteria*. Centre for Ecology & Hydrology (CEH) Dorset, Document No. C02492NEW.
- de Orte, M.R., Bonnail, E., Sarmiento, A.M., Bautista-Chamizo, E., Basallote, M.D., Riba, I., DelValls, Á., & Nieto, J.M. (2018). Metal fractionation in marine sediments acidified by enrichment of CO<sub>2</sub>: a risk assessment. *Marine Pollution Bulletin*, 131, 611-619.
- Department of Trade and Industry [DTI]. (1993). *Conditions for the discharge of oil contaminated cuttings resulting from offshore drilling*. London: Department of Trade and Industry, Oil and Gas Division.
- Du Laing, G., Rinklebe, J., Vandecasteele, B., Meers, E., & Tack, F.M.G. (2009). Trace metal behaviour in estuarine and riverine floodplain sediments: a review. *Science of the Total Environment*, 407, 3972-3985.
- Edwards, N.T. (1983). Polycyclic aromatic hydrocarbons (PAHs) in the terrestrial environment – a review. *Journal of Environmental Quality*, 12, 427-441.
- Eleftheriou, A., & Basford, D.J., 1989. The macrobenthic infauna of the offshore northern North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 69(1), 123-143.
- European Marine Observation Data Network [EMODnet]. (2020). Seabed habitats project.  
 [REDACTED]
- European Environment Agency [EEA]. (2019). The European Nature Information Service.  
 [REDACTED]
- Folk, R.L. (1954). The distinction between grain size and mineral composition in sedimentary rock nomenclature. *Journal of Geology*, 65(4), 344-359.
- Folk, R.L., & Ward, W.C. (1957). Brazos River bar (Texas); a study in the significance of grain size parameters. *Journal of Sedimentary Research*, 27(1), 3-26.

- Friocourt, M.P., Berthou, F., & Picart, D. (1982). Dibenzothiophene derivatives as organic markers of oil pollution. *Toxicological and Environmental Chemistry*, 5, 205-215.
- Fugro, 2020a. *Environmental features report, Awel Y Môr Offshore wind farm project, UK, Irish*. Fugro Document No.: 200145-R-004(02). Portchester: Fugro GB Marine Limited.
- Gray, J.S. (1981). *The ecology of marine sediments. an introduction to structure and function of benthic communities*. Cambridge University Press, 185.
- Haritash, A.K., & Kaushik, C.P. (2009). Biodegradation aspects of polycyclic aromatic hydrocarbons (PAHs): a review. *Journal of Hazardous Materials*, 169(1-3), 1-15.
- Howson, C.M., & Picton, B.E. eds. (1997). *The species directory of the marine fauna and flora of the British Isles and surrounding seas* (No. 276). Ulster Museum.
- Joint Nature Conservation Committee [JNCC]. (2015). *The Marine Habitat Classification for Britain and Ireland Version 15.03*. <https://mhc.jncc.gov.uk/about/>
- Joint Nature Conservation Committee [JNCC]. (2019a). *Annex I habitats list*. <https://sac.jncc.gov.uk/habitat/>
- Joint Nature Conservation Committee [JNCC]. (2019b). *Annex II species list*. <https://sac.jncc.gov.uk/species/>
- Kaskela, A. M., Kotilainen, A. T., Alanen, U., Cooper, R., Green, S., Guinan, J., van Heteren, S., Kihlman, S., Van Lancker, V., & Stevenson, A. (2019). Picking up the pieces—harmonising and collating seabed substrate data for European maritime areas. *Geosciences*, 9(2), 84.
- [REDACTED]
- Keith, L.H. (2015). The source of US EPA's sixteen PAH priority pollutants. *Polycyclic Aromatic Compounds*, 35(2-4), 147-160.
- Kennicutt II, M.C., Brooks, J.M., Bidgare, R.B., & Denoux, G.J. (1988). Gulf of Mexico hydrocarbon seep communities: Part II: Regional distribution of hydrocarbon seepage and associated fauna. *Deep-Sea Research*, 35, 1639–1651.
- Latimer, J.S., & Zheng, J. (2003). The sources, transport and fate of PAHs in the marine environment. In P.E.T. Douben (Ed). *PAHs: an ecotoxicological perspective*. John Wiley & Sons. 9-29
- Long, D. (2006). BGS Detailed explanation of seabed sediment modified Folk classification. MESH (Mapping European Seabed Habitats).
- [REDACTED]
- [REDACTED]
- Meadows, P.S. (1964). Substrate selection by *Corophium* species: the particle size of substrates. *Journal of Animal Ecology*, 33, 387-394.



Meyers, P.A., & Quinn, J.G. (1973). Association of hydrocarbons and mineral particles in saline solution. *Nature*, 224, 23-24.

National Research Council [NRC]. (1983). *Drilling discharges in the marine environment*. National Academy Press, Washington DC. 180.

Neff, J.M. (1979). *Polycyclic aromatic hydrocarbons in the aquatic environment. Sources, fates and biological effects*. Applied Science Publishers, London, 1-262.

Oslo and Paris Commission [OSPAR]. (2008). OSPAR List of threatened and/or declining species and habitats. Reference Number: 2008-06. [REDACTED]

Oslo and Paris Commission [OSPAR]. (2009). *Background document on CEMP assessment criteria for the QSR 2010*. Monitoring and Assessment Series. Report No. 978-1-907390-08-1.

Oslo and Paris Commission [OSPAR]. (2014). *Levels and trends in marine contaminants and their biological effects – CEMP assessment report 2013*. Monitoring and Assessment Series. Report No. 631/2014.

Paez-Osuna, F., & Ruiz-Fernandez, C. (1995). Comparative bioaccumulation of trace metals in *Penaeus stylirostris* in estuarine and coastal environments. *Estuarine, Coastal and Shelf Science*, 40, 35-44.

Parry, M.E.V. (2019). *Guidance on assigning benthic biotopes using EUNIS or the Marine Habitat Classification of Britain and Ireland (Revised 2019)*, JNCC Report No. 546, JNCC, Peterborough, ISSN 0963-8091.

Sims, R.C., & Overcash, M.R. (1983). Fate of polynuclear aromatic compounds (PNAs) I soil plant systems. *Residue Reviews*, 88, 1-68.

Stogiannidis, E., & Laane, R. (2015). Source characterization of polycyclic aromatic hydrocarbons by using their molecular indices: an overview of possibilities. *Reviews of Environmental Contamination and Toxicology*, 234, 49-133.

Tessier, A., Campbell, P.G.C., & Bisson, M. (1979). Sequential extraction procedure for the speciation of particulate trace metals. *Analytical Chemistry*, 51, 844-851.

Tillin, H.M. (2016). *Nephtys cirrosa and Bathyporeia spp. in infralittoral sand*.

Wakeham, S.G., Schaffner, C., & Giger, W. (1979). Polycyclic aromatic hydrocarbons in recent lake sediments – I. compounds having anthropogenic origins. *Geochimica et Cosmochimica Acta*, 44, 403-413.

Wang, Z., Wang, Y., Zhao, P., Chen, L., Yan, C., Yan, Y. & Chi, Q. (2015). Metal release from contaminated coastal sediments under changing pH conditions: Implications for metal mobilization in acidified oceans. *Marine Pollution Bulletin*, 101(2), 707-715.

- Warren, L.A., & Zimmerman, A.P. (1993). Trace metal-suspended particulate matter associations in a fluvial system: Physical and chemical influences. *Particulate and matter and aquatic contaminants*. Lewis Publishers, Boca Raton, 127-155.
- Wentworth, C.K. (1922). A scale of grade and class terms for clastic sediments. *Journal of Geology*, 30, 377-392.
- World Register of Marine Species [WoRMS] Editorial Board. (2019). *World register of marine species*. [REDACTED]
- Worsfold, T.M., Hall, D.J., & O'Reilly, M. (Ed). (2010). *Guidelines for processing marine macrobenthic invertebrate samples: a processing requirements protocol: Version 1.0, June 2010*. Report to the NMBAQC Committee.
- Youngblood, W.W., & Blumer, M. (1975). Polycyclic aromatic hydrocarbons in the environment: homologous series in soils and recent marine sediments. *Geochimica et Cosmochimica Acta*, 39, 1303–1314.
- Yunker, M.B., Macdonald, R.W., Vingarzan, R., Mitchell, R.H., Goyette, D., & Sylvestre, S. (2002). PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and composition. *Organic Geochemistry*, 33(4), 489-515.

# Appendices

---

## Appendix A Guidelines on Use of Report

---

### Appendix B Methodologies

- B.1 Survey Methods
  - B.2 Laboratory Analysis for Sediment Samples
  - B.3 Statistical Analysis
- 

### Appendix C Logs

- C.1 Survey Log
  - C.2 Grab Log
  - C.3 Video and Photographic Log
- 

### Appendix D Sediment Particle Size and Grab Sample Photographs

---

### Appendix E Sediment Hydrocarbon Analysis

- E.1 United States Environmental Protection Agency (US EPA) 16 Polycyclic Aromatic Hydrocarbon (PAH) Concentrations
  - E.2 Total 2 to 6 Ring PAH Concentrations
- 

### Appendix F Macrofaunal Analysis

- F.1 Macrofaunal Abundance
- F.2 Macrofaunal Biomass

# Appendix A

## Guidelines on Use of Report

This report (the "Report") was prepared as part of the services (the "Services") provided by Fugro GB Marine Limited ("Fugro") for its client (the "Client") under terms of the relevant contract between the two parties (the "Contract"). The Services were performed by Fugro based on requirements of the Client set out in the Contract or otherwise made known by the Client to Fugro at the time.

Fugro's obligations and liabilities to the Client or any other party in respect of the Services and this Report are limited in time and value as defined in Contract (or in the absence of any express provision in the Contract as implied by the law of the Contract) and Fugro provides no other representation or warranty whether express or implied, in relation to the Services or for the use of this Report for any other purpose. Furthermore, Fugro has no obligation to update or revise this Report based on changes in conditions or information which emerge following issue of this Report unless expressly required by the Contract.

The Services were performed by Fugro exclusively for the Client and any other party identified in the Contract for the purpose set out therein. Any use and/or reliance on the Report or the Services for purposes not expressly stated in the Contract, by the Client or any other party is that party's risk and Fugro accepts no liability whatsoever for any such use and/or reliance.



# Appendix B

## Methodologies

## B.1 Survey Methods

### B.1.1 Sediment Grab Sampling

Seabed samples were acquired using a 0.1 m<sup>2</sup> Hamon grab for macrofauna and particle size distribution (PSD) and a 0.1m<sup>2</sup> Day grab for chemistry sampling.

Operational procedures for grab sampling were as follows:

- The 0.1 m<sup>2</sup> Day grab or 0.1 m<sup>2</sup> Hamon grab was prepared for operations prior to arrival on station. Positioning was provided by a vessel offset from the vessel reference point. The Bridge communicated to the deck via a VHF radio when the vessel was steady and on location, and the grab was deployed from the starboard A-frame;
- When the deck crew operating the winch observed that the grab had reached the seabed (evidenced through a distinct slackening of the wire rope and snatch block), a positional fix was taken;
- On recovery to the deck, the sample was inspected and judged acceptable or otherwise (see below for rejection criteria);
- Once accepted, the grab sample was retained for faunal and PSD analysis (Hamon grab) and one grab sample was retained and subsampled for chemistry analysis (Day grab);
- Deck logs were completed for each sample acquired (including no samples) with date, time, sample number, fix number, sediment type, odour (i.e. H<sub>2</sub>S) and bioturbation or debris noted.

Samples were considered unacceptable in the following instances:

- Evidence of sediment washout caused through improperly closed grab jaws or inspection hatch;
- Sediment sample taken on an angle, where the grab jaws have not been parallel to the seabed when the grab fired;
- Disruption of the sample through striking the side of the vessel;
- Sample represented less than approximately 7 cm bite depth for the 0.1 m<sup>2</sup> Day grab or less than 5 L for the Hamon grab;
- Sample was more than 30 m from the target location.

### B.1.2 Chemical Sample Processing

- Hydrocarbon (HC) subsamples were collected using a metal scoop to a nominal depth of 2 cm. Subsamples collected were HCA1 and HCA2. The subsamples were preserved in glass jars at approximately –20 °C;
- Heavy metal (HM) subsamples were collected using a plastic scoop to a nominal depth of 2 cm. Subsamples collected were HMA1 and HMA2. The subsamples were preserved in polythene bags at approximately –20 °C.

### B.13 Macrofauna and Particle Size Distribution (PSD) Sample Processing

Macrofauna and PSD subsamples were processed as follows:

- Samples were processed in their entirety, by opening the grab to drop the sample into a container. All supernatant water was processed along with the sediment;
- A PSD subsample was collected prior to sieving of the sample and placed in a polythene bag and stored at approximately  $-20^{\circ}\text{C}$ ;
- The remaining sample was transferred to a 1 mm mesh sieve and transferred to the hand sieving table and sediment washed out;
- Once sieved samples were transferred to containers and fixed in 10 % buffered formal saline. The sample containers were then sealed, hazard labelled and stored securely on deck.

## B.2 Laboratory Analysis for Sediment Samples

### B.2.1 Particle Size Analysis

#### Dry Sieve Analysis

Particle size distribution (PSD) analysis was undertaken in accordance with FGBML in-house methods based on the National Marine Biological Association Quality Control scheme's (NMBAQC) best practice guidance document – Particle Size Analysis (PSA) for Supporting Biological Analysis, and BS1377: Parts 1: 2016 and 2: 1990.

Representative material > 1 mm was split from the bulk subsample and oven dried before sieving through a series of sieves with apertures corresponding to 0.5 phi intervals between 63 mm and 1 mm as described by the Wentworth scale (Wentworth, 1922). The weight of the sediment fraction retained on each mesh was subsequently measured and recorded.

#### Laser Diffraction

Particle size distribution (PSD) analysis was undertaken in accordance with FGBML in-house methods based on the NMBAQC scheme's best practice guidance document – Particle Size Analysis (PSA) for Supporting Biological Analysis, and BS ISO 13320: 2009.

Representative material < 1 mm was removed from the bulk subsample for laser analysis, a minimum of three triplicate analyses (mixed samples) or one triplicate analyses (sands) were analysed using the laser sizer at 0.5 phi intervals between < 1 mm to < 0.98 µm. Laser diffraction was carried out using a Malvern Mastersizer 2000 with a Hydro 2000G dispersion unit.

#### Sample Analysis Outputs and Deliverables

Sieve and laser data are merged and entered into GRADISTAT to derive statistics including mass and percentage retained within each size fraction, mean and median grain size, bulk sediment classes (percentage gravel, sand and silt/clay), skewness, sorting coefficients and Folk classification.

### B.2.2 Hydrocarbon Analysis in Sediments

Hydrocarbon analysis of sediments was carried out by Fugro GB Marine Limited.

### B.2.3 General Precautions

To effectively eliminate all possible sources of hydrocarbon contamination from the analysis the following precautionary measures were taken prior to sample work-up:

- All solvents were purchased as high purity grade. Each batch was checked for purity by concentrating approximately 400 mL down to a small volume (< 1 mL) and analysing by gas chromatography (GC);
- All water used was distilled through an all glass still and dichloromethane extracted to minimise contamination from plasticisers;

- All glassware was cleaned using an acid/base machine wash. The glassware was rinsed with acetone then finally with dichloromethane prior to use;
- Procedural blanks, replicate analyses and laboratory reference material were run with each batch.

### Ultrasonication Extraction for Hydrocarbons in Sediment

Sediment samples were thawed, homogenised and accurately weighed into a 250 mL conical flask. A solution containing an appropriate amount of the following internal standards was added to each sample using a microsyringe.

| Aromatic Standards           |
|------------------------------|
| D <sub>8</sub> Naphthalene   |
| D <sub>10</sub> Acenaphthene |
| D <sub>10</sub> Phenanthrene |
| D <sub>10</sub> Pyrene       |
| D <sub>12</sub> Chrysene     |
| D <sub>12</sub> Perylene     |

Methanol (50 mL) and solvent were mixed with the sediment. Dichloromethane (DCM) (60 mL) was then added and the sample mixed again. The flasks were then capped with solvent cleaned aluminium foil and ultrasonicated for 30 minutes.

After being allowed to settle the solvent was decanted through a GF-C filter paper into a 1 litre separating funnel. The extract was then partitioned with 100 mL of DCM extracted distilled water and the DCM layer run-off into a clean 500 mL round-bottomed flask. The ultrasonic extraction was repeated a further two times using 50 mL DCM and 15 minutes of ultrasonication. Each time the filtered extract was partitioned with the remaining methanol/water in the separating funnel. The DCM extracts were bulked and reduced in volume to approximately 2 mL using a rotary evaporator, then further reduced to approximately 1 mL under a gentle stream of nitrogen prior to clean-up.

Correction factors for wet/dry sediments were obtained by drying a subsample of the homogenised sediment to constant weight at 105 °C.

### Clean-up of Extracts by Column Chromatography

Removal of polar material, including lipids was carried out using a silica gel column. The silica gel used was 70 to 230 mesh which was heated at 400 °C for at least 4 hours to remove impurities and residual moisture and then stored at 200 °C prior to use. The sample extract was added to the silica gel column, containing 5 g of adsorbent and eluted with 35 mL of DCM/pentane (1:2). The eluant was reduced in volume using the evaporator to approximately 2 mL, with activated copper powder (for removal of free sulphur), before being further reduced under a gentle stream of nitrogen to an appropriate volume and analysed by and gas chromatography-mass spectrometry (GC-MS).

| Gas Chromatography-Mass Spectrometry [GC-MS] |   |
|--|---|
| Instrument                                   | HP 7890 Series GC with autoinjector and 5977A MSD   |
| Column                                       | (5 %phenyl)-methylpolysiloxane bonded fused silica, 60 m, 0.32 µm film thickness 0.25 mm internal diameter                      |
| Carrier Gas                                  | Hydrogen (constant flow 1.4 mL/min)   |
| Injector                                     | Splitless, 280 °C, split flow 40 mL/min, vent time 1.5 min (1 µL injection)   |
| Oven Temperature Programme                   | 60 °C – 1 min<br>60 °C to 180 °C at 11 °C/min<br>180 °C to 260 °C at 6 °C/min<br>260 °C to 320 °C at 6 °C/min<br>330 °C – 7 min |
| Source/Detector Temperature                  | 230 °C  |
| Electron Energy                              | 70 eV   |
| Selected Ion Monitoring (SIM)                | 9 groups - 6 ions per group   |
| Dwell Time (per ion)                         | 0.035 second  |

A full range of polycyclic aromatic hydrocarbon (PAH) and alkylated PAHs were quantified as specified by Department of Trade and Industry (DTI) regulations (DTI, 1993).

Calibration was undertaken using a range of PAH standard solutions, a number of alkylated PAH, dibenzothiophene and a range of suitable internal standards. Individual response factors were calculated for each of the compounds present in the calibration solution. Response factors for the non-calibrated alkylated PAHs were taken to be equivalent to closely related compounds. The MRV of individual and alkylated PAHs is 0.1 ng/g.

#### B.2.4 Metals Analysis

Sediment samples were dried at 40 °C and then sieved to the required size fraction (2000 µm). Samples were subjected to an aqua regia microwave digestion. This acid mixture allows a partial dissolution of metals, predominately releasing those associated with the sediment fines. The resulting digests were then analysed by inductively coupled plasma–mass spectrometry (ICP-MS) for arsenic, cadmium, chromium, copper, lead, mercury, nickel, tin and zinc; and inductively coupled plasma–optical emission spectrometry (ICP-OES) for aluminium and barium.

#### B.2.5 Macrofaunal Analysis

##### Benthic Infauna Analysis

Macrofauna analysis was carried out by FGBML benthic laboratories which are members of the NMBAQC scheme of external quality assurance.

On return to the laboratory, the samples were removed from formalin and washed through 0.1 mm mesh sieves. The material retained was then processed to remove fauna. The animals were separated by hand from the retained sediment by using a combination of stereo



microscopes for the fine sediments and in white trays for any coarser material. Processed sediment is stored in Phenoxetol (2 %) or returned to the original formalin.

Following extraction, the animals were identified and enumerated by specialist taxonomists. Identification was to species level where possible. Specimens which, due to their immaturity, damage incurred during processing or lack of suitable taxonomic literature, cannot be identified to species level are identified at higher taxonomic levels as appropriate. After identification, samples were stored in 70 % industrial denatured alcohol or a mixture of 70 % ethanol/1 % propylene glycol/29 % water. A minimum of 10 % of samples within the project were re-analysed (for extraction, species identification, enumeration and data entry) as per NMBAQC quality control guidelines (Worsfold, 2010). For biomass, identified macrofauna were blot dried and weighed at species/phyla level then returned to storage container.

Species abundances were entered on file in a spreadsheet package or the Unicorn database, both of which store and sort entries into taxonomic order and provide output files for numerical analysis. Nomenclature follows that given on the World Register of Marine Species (WoRMS Editorial Board, 2020). The taxonomic order is based on Species Directory codes (Howson & Picton, 1997) to give an idea of 'evolutionary rank'. Once all the entries had been checked, the resulting quantitative data were subjected to various statistical techniques to investigate community structure.

Prior to statistical analysis, the macrofaunal abundance data was manipulated to avoid spurious enhancement of community statistics. This involved the removal of all records of damaged, pelagic and juvenile taxa.

## B.3 Statistical Analysis

### B.3.1 Univariate and Multivariate Analysis of Macrofauna Data

Univariate analysis is used to extract features of communities which are not the function of specific taxa, i.e. these methods are species independent. They are not sensitive to spatio-temporal variations in species composition, so that assemblages with no species in common can theoretically have equal diversities. Univariate analyses were calculated using the Plymouth Routines in Multivariate Ecological Research (PRIMER) version (v)7 Diverse procedure and included number of individuals (N) and taxa (S), diversity employing the Shannon-Wiener index ( $H'_{Log_2}$ ) and evenness employing the Pielou's index (J).

#### Pielou's Equitability (J)

Pielou's index of evenness (also referred to as equitability) expresses how evenly distributed the individuals are among the different taxa. In general, the higher the evenness, the more balanced the sample is, as it indicates that the individuals are evenly distributed between the taxa recorded. It is expressed as:

$$J' = \frac{H'}{\text{Log } S}$$

Where:

$H'$  = Shannon-Wiener Index;

$S$  = total number of species.

#### Shannon-Wiener Diversity ( $H'_{Log_2}$ )

The Shannon-Wiener index of diversity incorporates richness and evenness as it expresses the number of species within a sample and the distribution of abundance across these species. In mathematical information theory, which is the context in which the Shannon-Wiener formula was originally devised, the Shannon-Wiener index of diversity measures the information content of a code in which one can write infinite messages. Analogously, the use of the Shannon-Wiener index of diversity as a measure of the diversity of a community, assumes that indefinitely samples can be taken from the community without depleting it. It is expressed as:

$$H' = - \sum_i P_i \text{Log}(P_i)$$

Where:

$P_i$  = proportion of the *ith* species.

### B.3.2 Multivariate Analysis

In the initial stage, multivariate analysis may involve transformation of data. For sediment analysis, transformation reduce the skewness allowing optimal performance of the

multivariate analysis. For macrofaunal analysis, transformation is applied where the fauna is numerically dominated by a few species which may mask the underlying community composition. Transformation reduces the influence of those more dominant species, with transformation ranging in severity from no transformation to the reduction of all data to presence absence only. If no transformation is applied to the data, greater emphasis is given to the most common species; a square root transformation allows the intermediate abundance species to play a role; a fourth root transformation results in a down-weighting of the dominant species, taking into much greater account the lowest abundant species, an allowing the underlying community composition to be assessed. An alternative transformation, with very similar effect to the fourth root, is the log transform  $\log(1+y)$ . The latter transformations are effectively equivalent in focusing attention on patterns within the whole community, mixing contribution from both common and rare species (Clarke & Warwick, 2001).

### Similarity Matrices

This analysis divides sites into groupings based on a measure of similarity or distance, depending on the nature of the data. For biological data, similarity based on the Bray-Curtis matrix is recommended, and for environmental data the Euclidean distance is recommended (Clarke & Warwick, 2001). The similarity/distance compares all samples with all other samples, producing a matrix.

### Hierarchical Agglomerative Clustering (CLUSTER) and Similarity Profile Testing (SIMPROF)

The hierarchical agglomerative clustering (CLUSTER) programme uses the similarity matrix to successively fuse samples into larger and larger groups according to their level of similarity. The results are displayed by means of a tree-like dendrogram with similarity (or distance) displayed on one axis and samples on the other. Similarity profile (SIMPROF) test was also performed in conjunction to cluster analysis. The test is a permutation of the null hypothesis that a set of specified samples, which are not a priori divided into groups, do not differ from each other in multivariate structure and looks for statistically significant evidence of "true" clusters in samples i.e. if the different sample groupings interpreted from the cluster analysis are significantly different. The results are displayed by colour convention on the dendrogram: samples connected by red lines constitute a significant group in statistical terms and cannot be separated. Conversely, samples connected by black lines, and therefore statistically different, may be interpreted as being ecologically not significantly different. The SIMPROF output was therefore always considered in terms of statistical and ecological significance, in line with Clarke et al. (2008) who indicate that, creating coarser groupings is entirely appropriate, provided that the resulting clusters are always supersets of the SIMPROF groups.

### Non-metric Multidimensional Scaling (nMDS)

Non-metric multidimensional scaling (nMDS) uses the similarity matrix to ordinate samples in a two-dimensional plane. This attempts to construct a map of the samples in which the more similar/close two samples are, the nearer they are on the map. The extent to which these

relations can be adequately represented in a two-dimensional map is expressed as the stress coefficient statistic or stress value. Stress values above 0.3 indicate near arbitrary points and the ordination should be considered unreliable. Stress values between 0.2 and 0.3 are poor representations of the data. Stress < 0.2 can show meaningful ordinations, while stress < 0.1 shows a good ordination of the data, with no real prospect of misleading interpretation. The combination of clustering and ordination analysis is a very effective way of checking the adequacy and mutual consistency of both representations (Clarke & Warwick, 2001).

#### Similarity Percentages Analysis (SIMPER)

This analysis can be applied to the data to gauge the faunal distinctiveness of each multivariate cluster, as identified by the clustering analysis. Similarity percentages analysis (SIMPER) provides a ranked list of taxa which contributes most to the similarity within clusters and the dissimilarity between clusters.

#### Principal Component Analysis (PCA)

The principal component analysis (PCA) identifies multidimensional patterns in datasets; once these multidimensional patterns have been found the data are compressed by reducing the number of dimensions without loss of information. The results of a PCA are graphically represented by the principal component (PC) axes, which are linear combinations of the values for each variable and represent the perpendicular distance in a multidimensional space along which the variance is maximised. The degree to which a 2D PCA succeeds in representing the full multidimensional information is in the percentage of the total variation expressed by the first two PCs. In general, a picture which accounts for as much as 70 % to 75 % of the original variation is likely to describe the overall structure rather well (Clarke & Warwick, 2001).

# Appendix C

## Logs

## C.1 Survey Log

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |                                      |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|--------------------------------------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes                                |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |                                      |
| 16/08/2020  | 12:13      | MA_ST64           | Video | SOL                   | 5       | 31                  | 456 206.7         | 5 925 480.6 | 456 125.1       | 5 925 474.3 | 81.8       | Station repeated as no video overlay |
| 16/08/2020  | 12:13      | MA_ST64           | Still | MA_ST64_01            | 6       | 31                  | 456 206.7         | 5 925 480.6 | 456 128.3       | 5 925 477.5 | 78.4       |                                      |
| 16/08/2020  | 12:17      | MA_ST64           | Still | MA_ST64_02            | 7       | 31                  | 456 206.7         | 5 925 480.6 | 456 182.1       | 5 925 474.7 | 25.3       |                                      |
| 16/08/2020  | 12:18      | MA_ST64           | Still | MA_ST64_03            | 8       | 31                  | 456 206.7         | 5 925 480.6 | 456 192.3       | 5 925 482.2 | 14.5       |                                      |
| 16/08/2020  | 12:19      | MA_ST64           | Still | MA_ST64_04            | 9       | 31                  | 456 206.7         | 5 925 480.6 | 456 202.0       | 5 925 484.8 | 6.3        |                                      |
| 16/08/2020  | 12:20      | MA_ST64           | Still | MA_ST64_05            | 10      | 31                  | 456 206.7         | 5 925 480.6 | 456 212.7       | 5 925 486.1 | 8.2        |                                      |
| 16/08/2020  | 12:21      | MA_ST64           | Still | MA_ST64_06            | 11      | 31                  | 456 206.7         | 5 925 480.6 | 456 218.7       | 5 925 495.6 | 19.2       |                                      |
| 16/08/2020  | 12:21      | MA_ST64           | Video | EOL                   | 11      | 31                  | 456 206.7         | 5 925 480.6 | 456 218.7       | 5 925 495.6 | 19.2       |                                      |
| 16/08/2020  | 12:35      | MA_ST63           | Video | SOL                   | 12      | 31                  | 456 177.3         | 5 925 488.6 | 456 148.4       | 5 925 482.0 | 29.7       |                                      |
| 16/08/2020  | 12:35      | MA_ST63           | Still | MA_ST63_01            | 13      | 31                  | 456 177.3         | 5 925 488.6 | 456 149.0       | 5 925 482.1 | 29.1       |                                      |
| 16/08/2020  | 12:36      | MA_ST63           | Still | MA_ST63_02            | 14      | 31                  | 456 177.3         | 5 925 488.6 | 456 157.3       | 5 925 492.8 | 20.5       |                                      |
| 16/08/2020  | 12:36      | MA_ST63           | Still | MA_ST63_03            | 15      | 31                  | 456 177.3         | 5 925 488.6 | 456 164.7       | 5 925 487.4 | 12.7       |                                      |
| 16/08/2020  | 12:37      | MA_ST63           | Still | MA_ST63_04            | 16      | 31                  | 456 177.3         | 5 925 488.6 | 456 171.6       | 5 925 493.9 | 7.8        |                                      |
| 16/08/2020  | 12:38      | MA_ST63           | Still | MA_ST63_05            | 17      | 31                  | 456 177.3         | 5 925 488.6 | 456 176.3       | 5 925 503.2 | 14.6       |                                      |
| 16/08/2020  | 12:38      | MA_ST63           | Still | MA_ST63_06            | 18      | 31                  | 456 177.3         | 5 925 488.6 | 456 181.5       | 5 925 510.8 | 22.6       |                                      |
| 16/08/2020  | 12:39      | MA_ST63           | Video | EOL                   | 19      | 31                  | 456 177.3         | 5 925 488.6 | 456 181.3       | 5 925 512.8 | 24.5       |                                      |
| 16/08/2020  | 12:48      | MA_ST62           | Video | SOL                   | 20      | 31                  | 456 168.0         | 5 925 503.0 | 456 142.3       | 5 925 497.9 | 26.3       |                                      |
| 16/08/2020  | 12:49      | MA_ST62           | Still | MA_ST62_01            | 21      | 31                  | -                 | -           | 456 163.0       | 5 925 506.1 | -          |                                      |
| 16/08/2020  | 12:50      | MA_ST62           | Still | MA_ST62_02            | 22      | 31                  | -                 | -           | 456 177.0       | 5 925 502.5 | -          |                                      |



| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |       |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|-------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |       |
| 16/08/2020  | 12:51      | MA_ST62           | Still | MA_ST62_03            | 23      | 31                  | -                 | -           | 456 193.9       | 5 925 506.8 | -          |       |
| 16/08/2020  | 12:51      | MA_ST62           | Still | MA_ST62_04            | 24      | 31                  | -                 | -           | 456 210.3       | 5 925 510.8 | -          |       |
| 16/08/2020  | 12:52      | MA_ST62           | Still | MA_ST62_05            | 25      | 31                  | -                 | -           | 456 216.6       | 5 925 512.6 | -          |       |
| 16/08/2020  | 12:53      | MA_ST62           | Still | MA_ST62_06            | 26      | 31                  | -                 | -           | 456 228.0       | 5 925 511.8 | -          |       |
| 16/08/2020  | 12:54      | MA_ST62           | Still | MA_ST62_07            | 27      | 31                  | -                 | -           | 456 233.9       | 5 925 511.7 | -          |       |
| 16/08/2020  | 12:54      | MA_ST62           | Video | EOL                   | 28      | 31                  | 456 234.0         | 5 925 510.0 | 456 235.2       | 5 925 518.8 | 8.9        |       |
| 16/08/2020  | 13:21      | MA_ST58           | Video | SOL                   | 29      | 35                  | 452 286.0         | 5 926 081.0 | 452 258.9       | 5 926 079.5 | 27.1       |       |
| 16/08/2020  | 13:21      | MA_ST58           | Still | MA_ST58_01            | 30      | 35                  | 452 286.0         | 5 926 081.0 | 452 261.1       | 5 926 079.6 | 24.9       |       |
| 16/08/2020  | 13:22      | MA_ST58           | Still | MA_ST58_02            | 31      | 35                  | 452 286.0         | 5 926 081.0 | 452 272.7       | 5 926 084.8 | 13.8       |       |
| 16/08/2020  | 13:23      | MA_ST58           | Still | MA_ST58_03            | 32      | 35                  | 452 286.0         | 5 926 081.0 | 452 282.3       | 5 926 084.6 | 5.2        |       |
| 16/08/2020  | 13:23      | MA_ST58           | Still | MA_ST58_04            | 33      | 35                  | 452 286.0         | 5 926 081.0 | 452 291.9       | 5 926 091.9 | 12.3       |       |
| 16/08/2020  | 13:24      | MA_ST58           | Still | MA_ST58_05            | 34      | 35                  | 452 286.0         | 5 926 081.0 | 452 299.6       | 5 926 096.1 | 20.3       |       |
| 16/08/2020  | 13:24      | MA_ST58           | Video | EOL                   | 35      | 35                  | 452 286.0         | 5 926 081.0 | 452 300.6       | 5 926 101.7 | 25.4       |       |
| 16/08/2020  | 13:45      | MA_ST59           | Video | SOL                   | 36      | 35                  | 450 387.0         | 5 924 424.0 | 450 361.3       | 5 924 431.9 | 26.9       |       |
| 16/08/2020  | 13:45      | MA_ST59           | Still | MA_ST59_01            | 37      | 35                  | 450 387.0         | 5 924 424.0 | 450 363.1       | 5 924 429.1 | 24.5       |       |
| 16/08/2020  | 13:46      | MA_ST59           | Still | MA_ST59_02            | 38      | 35                  | 450 387.0         | 5 924 424.0 | 450 370.8       | 5 924 421.7 | 16.3       |       |
| 16/08/2020  | 13:47      | MA_ST59           | Still | MA_ST59_03            | 39      | 35                  | 450 387.0         | 5 924 424.0 | 450 385.5       | 5 924 431.9 | 8.0        |       |
| 16/08/2020  | 13:48      | MA_ST59           | Still | MA_ST59_04            | 40      | 35                  | 450 387.0         | 5 924 424.0 | 450 394.2       | 5 924 430.0 | 9.4        |       |
| 16/08/2020  | 13:49      | MA_ST59           | Still | MA_ST59_05            | 41      | 35                  | 450 387.0         | 5 924 424.0 | 450 404.0       | 5 924 435.4 | 20.4       |       |
| 16/08/2020  | 13:49      | MA_ST59           | Video | EOL                   | 42      | 35                  | 450 387.0         | 5 924 424.0 | 450 406.8       | 5 924 439.7 | 25.2       |       |
| 16/08/2020  | 14:14      | MA_ST65           | Video | SOL                   | 43      | 31                  | 453 569.0         | 5 923 291.0 | 453 543.5       | 5 923 290.3 | 25.5       |       |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |                                |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|--------------------------------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes                          |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |                                |
| 16/08/2020  | 14:15      | MA_ST65           | Still | MA_ST65_01            | 44      | 31                  | 453 569.0         | 5 923 291.0 | 453 547.2       | 5 923 290.7 | 21.8       |                                |
| 16/08/2020  | 14:15      | MA_ST65           | Still | MA_ST65_02            | 45      | 31                  | 453 569.0         | 5 923 291.0 | 453 558.3       | 5 923 292.5 | 10.8       |                                |
| 16/08/2020  | 14:16      | MA_ST65           | Still | MA_ST65_03            | 46      | 31                  | 453 569.0         | 5 923 291.0 | 453 583.1       | 5 923 287.5 | 14.5       |                                |
| 16/08/2020  | 14:17      | MA_ST65           | Still | MA_ST65_04            | 47      | 31                  | 453 569.0         | 5 923 291.0 | 453 592.8       | 5 923 291.5 | 23.8       | Vessel turned after this point |
| 16/08/2020  | 14:19      | MA_ST65           | Still | MA_ST65_05            | 48      | 31                  | 453 569.0         | 5 923 291.0 | 453 574.1       | 5 923 285.1 | 7.9        |                                |
| 16/08/2020  | 14:20      | MA_ST65           | Video | EOL                   | NF      | 31                  | 453 569.0         | 5 923 291.0 | 453 552.5       | 5 923 275.5 | 22.6       |                                |
| 16/08/2020  | 14:42      | MA_ST66           | Video | SOL                   | 49      | 24                  | 450 568.0         | 5 920 636.0 | 450 540.3       | 5 920 638.3 | 27.8       |                                |
| 16/08/2020  | 14:42      | MA_ST66           | Still | MA_ST66_01            | 50      | 24                  | 450 568.0         | 5 920 636.0 | 450 546.0       | 5 920 639.9 | 22.4       |                                |
| 16/08/2020  | 14:42      | MA_ST66           | Still | MA_ST66_02            | 51      | 24                  | 450 568.0         | 5 920 636.0 | 450 555.8       | 5 920 634.6 | 12.3       |                                |
| 16/08/2020  | 14:43      | MA_ST66           | Still | MA_ST66_03            | 52      | 24                  | 450 568.0         | 5 920 636.0 | 450 564.6       | 5 920 637.9 | 3.9        |                                |
| 16/08/2020  | 14:43      | MA_ST66           | Still | MA_ST66_04            | 53      | 24                  | 450 568.0         | 5 920 636.0 | 450 576.1       | 5 920 637.3 | 8.2        |                                |
| 16/08/2020  | 14:44      | MA_ST66           | Still | MA_ST66_05            | 54      | 24                  | 450 568.0         | 5 920 636.0 | 450 590.5       | 5 920 637.6 | 22.5       |                                |
| 16/08/2020  | 14:44      | MA_ST66           | Video | EOL                   | 55      | 24                  | 450 568.0         | 5 920 636.0 | 450 594.3       | 5 920 635.3 | 26.3       |                                |
| 16/08/2020  | 15:06      | MA_ST56           | Video | SOL                   | 56      | 31                  | 448 166.2         | 5 923 311.0 | 448 144.9       | 5 923 327.0 | 26.7       |                                |
| 16/08/2020  | 15:06      | MA_ST56           | Still | MA_ST56_01            | 57      | 31                  | 448 166.2         | 5 923 311.0 | 448 147.9       | 5 923 324.9 | 23.0       |                                |
| 16/08/2020  | 15:06      | MA_ST56           | Still | MA_ST56_02            | 58      | 31                  | 448 166.2         | 5 923 311.0 | 448 157.9       | 5 923 323.3 | 14.9       |                                |
| 16/08/2020  | 15:07      | MA_ST56           | Still | MA_ST56_03            | 59      | 31                  | 448 166.2         | 5 923 311.0 | 448 169.3       | 5 923 305.7 | 6.1        |                                |
| 16/08/2020  | 15:08      | MA_ST56           | Still | MA_ST56_04            | 60      | 31                  | 448 166.2         | 5 923 311.0 | 448 175.4       | 5 923 299.4 | 14.8       |                                |
| 16/08/2020  | 15:08      | MA_ST56           | Still | MA_ST56_05            | 61      | 31                  | 448 166.2         | 5 923 311.0 | 448 184.5       | 5 923 294.0 | 24.9       |                                |
| 16/08/2020  | 15:08      | MA_ST56           | Video | EOL                   | 62      | 31                  | 448 166.2         | 5 923 311.0 | 448 186.1       | 5 923 293.0 | 26.8       |                                |
| 16/08/2020  | 15:18      | MA_ST55           | Video | SOL                   | 63      | 28                  | 447 905.0         | 5 923 406.9 | 447 878.7       | 5 923 416.6 | 28.0       |                                |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |                                |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|--------------------------------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes                          |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |                                |
| 16/08/2020  | 15:18      | MA_ST55           | Still | MA_ST55_01            | 64      | 28                  | 447 905.0         | 5 923 406.9 | 447 888.8       | 5 923 419.2 | 20.4       |                                |
| 16/08/2020  | 15:19      | MA_ST55           | Still | MA_ST55_02            | 65      | 28                  | 447 905.0         | 5 923 406.9 | 447 896.0       | 5 923 415.6 | 12.5       |                                |
| 16/08/2020  | 15:20      | MA_ST55           | Still | MA_ST55_03            | 66      | 28                  | 447 905.0         | 5 923 406.9 | 447 905.7       | 5 923 409.1 | 2.3        |                                |
| 16/08/2020  | 15:20      | MA_ST55           | Still | MA_ST55_04            | 67      | 28                  | 447 905.0         | 5 923 406.9 | 447 915.1       | 5 923 401.3 | 11.5       |                                |
| 16/08/2020  | 15:21      | MA_ST55           | Still | MA_ST55_05            | 68      | 28                  | 447 905.0         | 5 923 406.9 | 447 925.4       | 5 923 396.8 | 22.8       |                                |
| 16/08/2020  | 15:21      | MA_ST55           | Video | EOL                   | 69      | 28                  | 447 905.0         | 5 923 406.9 | 447 927.3       | 5 923 395.7 | 25.0       |                                |
| 16/08/2020  | 15:29      | MA_ST53           | Video | SOL                   | 70      | 29                  | 447 911.0         | 5 923 474.0 | 447 909.5       | 5 923 476.5 | 2.9        |                                |
| 16/08/2020  | 15:29      | MA_ST53           | Still | MA_ST53_01            | 71      | 29                  | -                 | -           | 447 911.7       | 5 923 474.3 | -          |                                |
| 16/08/2020  | 15:29      | MA_ST53           | Still | MA_ST53_02            | 72      | 29                  | -                 | -           | 447 915.6       | 5 923 463.7 | -          |                                |
| 16/08/2020  | 15:30      | MA_ST53           | Still | MA_ST53_03            | 73      | 29                  | -                 | -           | 447 930.3       | 5 923 461.6 | -          |                                |
| 16/08/2020  | 15:30      | MA_ST53           | Still | MA_ST53_04            | 74      | 29                  | -                 | -           | 447 937.2       | 5 923 454.7 | -          |                                |
| 16/08/2020  | 15:31      | MA_ST53           | Still | MA_ST53_05            | 75      | 29                  | -                 | -           | 447 944.4       | 5 923 453.7 | -          |                                |
| 16/08/2020  | 15:31      | MA_ST53           | Still | MA_ST53_06            | 76      | 29                  | -                 | -           | 447 951.7       | 5 923 451.6 | -          |                                |
| 16/08/2020  | 15:32      | MA_ST53           | Still | MA_ST53_07            | 77      | 29                  | -                 | -           | 447 960.3       | 5 923 451.4 | -          |                                |
| 16/08/2020  | 15:32      | MA_ST53           | Still | MA_ST53_08            | 78      | 29                  | -                 | -           | 447 964.0       | 5 923 447.2 | -          |                                |
| 16/08/2020  | 15:32      | MA_ST53           | Video | EOL                   | 79      | 29                  | 447 958.0         | 5 923 450.0 | 447 966.8       | 5 923 446.4 | 9.5        |                                |
| 16/08/2020  | 15:50      | MA_ST57           | Video | SOL                   | 80      | 28                  | 445 929.0         | 5 921 227.0 | 445 939.0       | 5 921 251.2 | 26.2       |                                |
| 16/08/2020  | 15:51      | MA_ST57           | Still | MA_ST57_01            | 81      | 28                  | 445 929.0         | 5 921 227.0 | 445 932.8       | 5 921 249.4 | 22.7       | Vessel turned after this point |
| 16/08/2020  | 15:55      | MA_ST57           | Still | MA_ST57_02            | 82      | 28                  | 445 929.0         | 5 921 227.0 | 445 914.1       | 5 921 238.4 | 18.7       |                                |
| 16/08/2020  | 15:56      | MA_ST57           | Still | MA_ST57_03            | 83      | 28                  | 445 929.0         | 5 921 227.0 | 445 923.5       | 5 921 236.3 | 10.8       |                                |
| 16/08/2020  | 15:56      | MA_ST57           | Still | MA_ST57_04            | 84      | 28                  | 445 929.0         | 5 921 227.0 | 445 928.9       | 5 921 229.3 | 2.3        |                                |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |                                |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|--------------------------------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes                          |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |                                |
| 16/08/2020  | 15:57      | MA_ST57           | Still | MA_ST57_05            | 85      | 28                  | 445 929.0         | 5 921 227.0 | 445 937.3       | 5 921 224.0 | 8.9        |                                |
| 16/08/2020  | 15:58      | MA_ST57           | Still | MA_ST57_06            | 86      | 28                  | 445 929.0         | 5 921 227.0 | 445 945.6       | 5 921 216.8 | 19.5       |                                |
| 16/08/2020  | 15:58      | MA_ST57           | Video | EOL                   | 87      | 28                  | 445 929.0         | 5 921 227.0 | 445 947.5       | 5 921 215.3 | 21.9       |                                |
| 16/08/2020  | 16:17      | MA_ST46           | Video | SOL                   | 88      | 30                  | 443 175.0         | 5 922 829.0 | 443 152.3       | 5 922 810.2 | 29.5       |                                |
| 16/08/2020  | 16:17      | MA_ST46           | Still | MA_ST46_01            | 89      | 30                  | 443 175.0         | 5 922 829.0 | 443 156.9       | 5 922 811.3 | 25.3       |                                |
| 16/08/2020  | 16:17      | MA_ST46           | Still | MA_ST46_02            | 90      | 30                  | 443 175.0         | 5 922 829.0 | 443 167.3       | 5 922 817.4 | 13.9       |                                |
| 16/08/2020  | 16:18      | MA_ST46           | Still | MA_ST46_03            | 91      | 30                  | 443 175.0         | 5 922 829.0 | 443 169.2       | 5 922 830.8 | 6.1        |                                |
| 16/08/2020  | 16:18      | MA_ST46           | Still | MA_ST46_04            | 92      | 30                  | 443 175.0         | 5 922 829.0 | 443 172.8       | 5 922 844.0 | 15.2       | Vessel turned after this point |
| 16/08/2020  | 16:25      | MA_ST46           | Still | MA_ST46_05            | 93      | 30                  | 443 175.0         | 5 922 829.0 | 443 205.0       | 5 922 832.7 | 30.3       |                                |
| 16/08/2020  | 16:25      | MA_ST46           | Video | EOL                   | 94      | 30                  | 443 175.0         | 5 922 829.0 | 443 211.0       | 5 922 833.6 | 36.3       |                                |
| 16/08/2020  | 16:45      | MA_ST44           | Video | SOL                   | 95      | 32                  | 445 277.0         | 5 924 515.0 | 445 298.2       | 5 924 530.8 | 26.4       |                                |
| 16/08/2020  | 16:45      | MA_ST44           | Still | MA_ST44_01            | 96      | 32                  | 445 277.0         | 5 924 515.0 | 445 293.8       | 5 924 531.2 | 23.3       |                                |
| 16/08/2020  | 16:46      | MA_ST44           | Still | MA_ST44_02            | 97      | 32                  | 445 277.0         | 5 924 515.0 | 445 286.3       | 5 924 526.3 | 14.6       |                                |
| 16/08/2020  | 16:46      | MA_ST44           | Still | MA_ST44_03            | 98      | 32                  | 445 277.0         | 5 924 515.0 | 445 278.2       | 5 924 520.3 | 5.4        |                                |
| 16/08/2020  | 16:47      | MA_ST44           | Still | MA_ST44_04            | 99      | 32                  | 445 277.0         | 5 924 515.0 | 445 271.0       | 5 924 514.7 | 6.0        |                                |
| 16/08/2020  | 16:48      | MA_ST44           | Still | MA_ST44_05            | 100     | 32                  | 445 277.0         | 5 924 515.0 | 445 259.9       | 5 924 508.6 | 18.3       |                                |
| 16/08/2020  | 16:48      | MA_ST44           | Still | MA_ST44_06            | 101     | 32                  | 445 277.0         | 5 924 515.0 | 445 253.9       | 5 924 503.9 | 25.6       |                                |
| 16/08/2020  | 16:48      | MA_ST44           | Video | EOL                   | 102     | 32                  | 445 277.0         | 5 924 515.0 | 445 252.4       | 5 924 502.9 | 27.4       |                                |
| 16/08/2020  | 17:06      | MA_ST43           | Video | SOL                   | 103     | 35                  | 447 038.0         | 5 926 464.0 | 447 059.3       | 5 926 476.0 | 24.4       |                                |
| 16/08/2020  | 17:06      | MA_ST43           | Still | MA_ST43_01            | 104     | 35                  | 447 038.0         | 5 926 464.0 | 447 057.2       | 5 926 475.3 | 22.3       |                                |
| 16/08/2020  | 17:07      | MA_ST43           | Still | MA_ST43_02            | 105     | 35                  | 447 038.0         | 5 926 464.0 | 447 049.1       | 5 926 470.9 | 13.1       |                                |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |       |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|-------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |       |
| 16/08/2020  | 17:07      | MA_ST43           | Still | MA_ST43_03            | 106     | 35                  | 447 038.0         | 5 926 464.0 | 447 041.6       | 5 926 467.4 | 5.0        |       |
| 16/08/2020  | 17:08      | MA_ST43           | Still | MA_ST43_04            | 107     | 35                  | 447 038.0         | 5 926 464.0 | 447 032.5       | 5 926 460.3 | 6.6        |       |
| 16/08/2020  | 17:09      | MA_ST43           | Still | MA_ST43_05            | 108     | 35                  | 447 038.0         | 5 926 464.0 | 447 021.4       | 5 926 457.0 | 18.1       |       |
| 16/08/2020  | 17:09      | MA_ST43           | Video | EOL                   | 109     | 35                  | 447 038.0         | 5 926 464.0 | 447 020.7       | 5 926 455.4 | 19.3       |       |
| 17/08/2020  | 09:25      | MA_ST22           | Video | SOL                   | 110     | 36                  | 438 596.0         | 5 923 566.3 | 438 618.0       | 5 923 551.4 | 26.6       |       |
| 17/08/2020  | 09:25      | MA_ST22           | Still | MA_ST22_01            | 111     | 36                  | 438 596.0         | 5 923 566.3 | 438 614.5       | 5 923 552.4 | 23.1       |       |
| 17/08/2020  | 09:26      | MA_ST22           | Still | MA_ST22_02            | 112     | 36                  | 438 596.0         | 5 923 566.3 | 438 596.8       | 5 923 558.3 | 8.0        |       |
| 17/08/2020  | 09:26      | MA_ST22           | Still | MA_ST22_03            | 113     | 36                  | 438 596.0         | 5 923 566.3 | 438 593.1       | 5 923 565.6 | 3.0        |       |
| 17/08/2020  | 09:28      | MA_ST22           | Still | MA_ST22_04            | 114     | 36                  | 438 596.0         | 5 923 566.3 | 438 582.3       | 5 923 575.6 | 16.5       |       |
| 17/08/2020  | 09:28      | MA_ST22           | Still | MA_ST22_05            | 115     | 36                  | 438 596.0         | 5 923 566.3 | 438 588.9       | 5 923 590.5 | 25.2       |       |
| 17/08/2020  | 09:28      | MA_ST22           | Video | EOL                   | 116     | 36                  | 438 596.0         | 5 923 566.3 | 438 590.7       | 5 923 594.2 | 28.4       |       |
| 17/08/2020  | 09:40      | MA_ST28           | Video | SOL                   | 117     | 39                  | 439 234.0         | 5 923 384.0 | 439 205.4       | 5 923 380.7 | 28.8       |       |
| 17/08/2020  | 09:40      | MA_ST28           | Still | MA_ST28_01            | 118     | 39                  | -                 | -           | 439 208.9       | 5 923 382.4 | -          |       |
| 17/08/2020  | 09:41      | MA_ST28           | Still | MA_ST28_02            | 119     | 39                  | -                 | -           | 439 228.2       | 5 923 385.5 | -          |       |
| 17/08/2020  | 09:41      | MA_ST28           | Still | MA_ST28_03            | 120     | 39                  | -                 | -           | 439 236.0       | 5 923 383.1 | -          |       |
| 17/08/2020  | 09:42      | MA_ST28           | Still | MA_ST28_04            | 121     | 39                  | -                 | -           | 439 259.6       | 5 923 382.1 | -          |       |
| 17/08/2020  | 09:44      | MA_ST28           | Still | MA_ST28_05            | 122     | 39                  | -                 | -           | 439 292.1       | 5 923 382.8 | -          |       |
| 17/08/2020  | 09:44      | MA_ST28           | Still | MA_ST28_06            | 123     | 39                  | -                 | -           | 439 301.9       | 5 923 382.5 | -          |       |
| 17/08/2020  | 09:45      | MA_ST28           | Still | MA_ST28_07            | 124     | 39                  | -                 | -           | 439 313.2       | 5 923 385.5 | -          |       |
| 17/08/2020  | 09:45      | MA_ST28           | Still | MA_ST28_08            | 125     | 39                  | -                 | -           | 439 325.4       | 5 923 380.0 | -          |       |
| 17/08/2020  | 09:52      | MA_ST28           | Still | MA_ST28_09            | 126     | 39                  | -                 | -           | 439 255.4       | 5 923 379.1 | -          |       |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |       |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|-------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |       |
| 17/08/2020  | 09:53      | MA_ST28           | Still | MA_ST28_10            | 127     | 39                  | -                 | -           | 439 264.4       | 5 923 379.3 | -          |       |
| 17/08/2020  | 09:54      | MA_ST28           | Still | MA_ST28_11            | 128     | 39                  | -                 | -           | 439 276.2       | 5 923 385.7 | -          |       |
| 17/08/2020  | 09:54      | MA_ST28           | Still | MA_ST28_12            | 129     | 39                  | -                 | -           | 439 286.1       | 5 923 385.0 | -          |       |
| 17/08/2020  | 09:55      | MA_ST28           | Video | EOL                   | 130     | 39                  | 439 317.0         | 5 923 381.0 | 439 289.4       | 5 923 381.1 | 27.6       |       |
| 17/08/2020  | 10:12      | MA_ST26           | Video | SOL                   | 131     | 37                  | 441 208.0         | 5 925 055.0 | 441 196.7       | 5 925 054.8 | 11.3       |       |
| 17/08/2020  | 10:12      | MA_ST26           | Still | MA_ST26_01            | 132     | 37                  | -                 | -           | 441 196.8       | 5 925 053.8 | -          |       |
| 17/08/2020  | 10:13      | MA_ST26           | Still | MA_ST26_02            | 133     | 37                  | -                 | -           | 441 204.1       | 5 925 045.4 | -          |       |
| 17/08/2020  | 10:14      | MA_ST26           | Still | MA_ST26_03            | 134     | 37                  | -                 | -           | 441 217.9       | 5 925 035.2 | -          |       |
| 17/08/2020  | 10:14      | MA_ST26           | Still | MA_ST26_04            | 135     | 37                  | -                 | -           | 441 223.0       | 5 925 022.6 | -          |       |
| 17/08/2020  | 10:15      | MA_ST26           | Still | MA_ST26_05            | 136     | 37                  | -                 | -           | 441 237.3       | 5 925 015.5 | -          |       |
| 17/08/2020  | 10:16      | MA_ST26           | Video | EOL                   | 137     | 37                  | 441 235.0         | 5 925 021.0 | 441 239.0       | 5 925 011.4 | 10.4       |       |
| 17/08/2020  | 10:27      | MA_ST23           | Video | SOL                   | 138     | 36                  | 441 118.0         | 5 925 220.2 | 441 092.9       | 5 925 229.2 | 26.6       |       |
| 17/08/2020  | 10:27      | MA_ST23           | Still | MA_ST23_01            | 139     | 36                  | 441 118.0         | 5 925 220.2 | 441 095.6       | 5 925 227.7 | 23.6       |       |
| 17/08/2020  | 10:28      | MA_ST23           | Still | MA_ST23_02            | 140     | 36                  | 441 118.0         | 5 925 220.2 | 441 103.7       | 5 925 221.0 | 14.3       |       |
| 17/08/2020  | 10:29      | MA_ST23           | Still | MA_ST23_03            | 141     | 36                  | 441 118.0         | 5 925 220.2 | 441 115.4       | 5 925 223.2 | 4.0        |       |
| 17/08/2020  | 10:30      | MA_ST23           | Still | MA_ST23_04            | 142     | 36                  | 441 118.0         | 5 925 220.2 | 441 118.4       | 5 925 221.6 | 1.5        |       |
| 17/08/2020  | 10:31      | MA_ST23           | Still | MA_ST23_05            | 143     | 36                  | 441 118.0         | 5 925 220.2 | 441 130.5       | 5 925 220.0 | 12.5       |       |
| 17/08/2020  | 10:31      | MA_ST23           | Still | MA_ST23_06            | 144     | 36                  | 441 118.0         | 5 925 220.2 | 441 137.9       | 5 925 215.3 | 20.5       |       |
| 17/08/2020  | 10:32      | MA_ST23           | Video | EOL                   | 145     | 36                  | 441 118.0         | 5 925 220.2 | 441 141.0       | 5 925 215.1 | 23.6       |       |
| 17/08/2020  | 10:49      | MA_ST19           | Video | SOL                   | 146     | 39                  | 442 461.0         | 5 926 594.0 | 442 434.2       | 5 926 597.3 | 27.0       |       |
| 17/08/2020  | 10:49      | MA_ST19           | Still | MA_ST19_01            | 147     | 39                  | 442 461.0         | 5 926 594.0 | 442 440.4       | 5 926 594.4 | 20.6       |       |



| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |       |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|-------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |       |
| 17/08/2020  | 10:50      | MA_ST19           | Still | MA_ST19_02            | 148     | 39                  | 442 461.0         | 5 926 594.0 | 442 450.5       | 5 926 588.2 | 12.0       |       |
| 17/08/2020  | 10:52      | MA_ST19           | Still | MA_ST19_03            | 149     | 39                  | 442 461.0         | 5 926 594.0 | 442 458.8       | 5 926 595.0 | 2.4        |       |
| 17/08/2020  | 10:52      | MA_ST19           | Still | MA_ST19_04            | 150     | 39                  | 442 461.0         | 5 926 594.0 | 442 469.2       | 5 926 596.7 | 8.6        |       |
| 17/08/2020  | 10:53      | MA_ST19           | Still | MA_ST19_05            | 151     | 39                  | 442 461.0         | 5 926 594.0 | 442 477.8       | 5 926 597.1 | 17.1       |       |
| 17/08/2020  | 10:54      | MA_ST19           | Video | EOL                   | 152     | 39                  | 442 461.0         | 5 926 594.0 | 442 485.6       | 5 926 598.1 | 24.9       |       |
| 17/08/2020  | 11:18      | MA_ST11           | Video | SOL                   | 153     | 42                  | 438 522.0         | 5 926 615.0 | 438 495.6       | 5 926 605.4 | 28.1       |       |
| 17/08/2020  | 11:18      | MA_ST11           | Still | MA_ST11_01            | 154     | 42                  | 438 522.0         | 5 926 615.0 | 438 501.6       | 5 926 606.1 | 22.2       |       |
| 17/08/2020  | 11:19      | MA_ST11           | Still | MA_ST11_02            | 155     | 42                  | 438 522.0         | 5 926 615.0 | 438 509.6       | 5 926 609.9 | 13.4       |       |
| 17/08/2020  | 11:20      | MA_ST11           | Still | MA_ST11_03            | 156     | 42                  | 438 522.0         | 5 926 615.0 | 438 511.8       | 5 926 619.0 | 11.0       |       |
| 17/08/2020  | 11:21      | MA_ST11           | Still | MA_ST11_04            | 157     | 42                  | 438 522.0         | 5 926 615.0 | 438 523.6       | 5 926 628.2 | 13.3       |       |
| 17/08/2020  | 11:22      | MA_ST11           | Still | MA_ST11_05            | 158     | 42                  | 438 522.0         | 5 926 615.0 | 438 536.2       | 5 926 632.5 | 22.6       |       |
| 17/08/2020  | 11:22      | MA_ST11           | Video | EOL                   | 159     | 42                  | 438 522.0         | 5 926 615.0 | 438 540.9       | 5 926 634.2 | 26.9       |       |
| 17/08/2020  | 11:39      | MA_ST12           | Video | SOL                   | 160     | 38                  | 437 529.0         | 5 925 669.0 | 437 501.2       | 5 925 669.5 | 27.8       |       |
| 17/08/2020  | 11:40      | MA_ST12           | Still | MA_ST12_01            | 161     | 38                  | 437 529.0         | 5 925 669.0 | 437 509.0       | 5 925 669.6 | 20.0       |       |
| 17/08/2020  | 11:40      | MA_ST12           | Still | MA_ST12_02            | 162     | 38                  | 437 529.0         | 5 925 669.0 | 437 521.0       | 5 925 669.6 | 8.0        |       |
| 17/08/2020  | 11:41      | MA_ST12           | Still | MA_ST12_03            | 163     | 38                  | 437 529.0         | 5 925 669.0 | 437 527.1       | 5 925 671.8 | 3.4        |       |
| 17/08/2020  | 11:42      | MA_ST12           | Still | MA_ST12_04            | 164     | 38                  | 437 529.0         | 5 925 669.0 | 437 536.4       | 5 925 667.9 | 7.5        |       |
| 17/08/2020  | 11:43      | MA_ST12           | Still | MA_ST12_05            | 165     | 38                  | 437 529.0         | 5 925 669.0 | 437 550.1       | 5 925 666.5 | 21.2       |       |
| 17/08/2020  | 11:44      | MA_ST12           | Still | MA_ST12_06            | 166     | 38                  | 437 529.0         | 5 925 669.0 | 437 560.8       | 5 925 666.3 | 31.9       |       |
| 17/08/2020  | 11:44      | MA_ST12           | Video | EOL                   | 167     | 38                  | 437 529.0         | 5 925 669.0 | 437 563.2       | 5 925 661.0 | 35.1       |       |
| 17/08/2020  | 12:51      | MA_ST04           | Video | SOL                   | 173     | 44                  | 434 714.0         | 5 926 692.0 | 434 692.7       | 5 926 680.4 | 24.2       |       |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |       |                       |         |                     |                   |             |                 |             |            |               |
|---|------------|-------------------|-------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|---------------|
| Date  | Time [UTC] | Transect/ Station | Type  | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes         |
|   |            |                   |       |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |               |
| 17/08/2020  | 12:51      | MA_ST04           | Still | MA_ST04_01            | 174     | 44                  | 434 714.0         | 5 926 692.0 | 434 701.5       | 5 926 689.3 | 12.8       |               |
| 17/08/2020  | 12:52      | MA_ST04           | Still | MA_ST04_02            | 175     | 44                  | 434 714.0         | 5 926 692.0 | 434 706.6       | 5 926 693.5 | 7.5        |               |
| 17/08/2020  | 12:52      | MA_ST04           | Still | MA_ST04_03            | 176     | 44                  | 434 714.0         | 5 926 692.0 | 434 708.1       | 5 926 703.6 | 13.1       |               |
| 17/08/2020  | 12:52      | MA_ST04           | Still | MA_ST04_04            | 177     | 44                  | 434 714.0         | 5 926 692.0 | 434 708.6       | 5 926 712.1 | 20.8       |               |
| 17/08/2020  | 12:53      | MA_ST04           | Still | MA_ST04_05            | 178     | 44                  | 434 714.0         | 5 926 692.0 | 434 710.7       | 5 926 714.5 | 22.7       |               |
| 17/08/2020  | 12:53      | MA_ST04           | Video | EOL                   | 179     | 44                  | 434 714.0         | 5 926 692.0 | 434 712.7       | 5 926 713.7 | 21.8       |               |
| 17/08/2020  | 14:19      | MA_ST64 (2)       | Video | SOL                   | 181     | 28                  | 456 206.7         | 5 925 480.6 | 456 179.2       | 5 925 492.0 | 29.8       |               |
| 17/08/2020  | 14:20      | MA_ST64 (2)       | Still | MA_ST64(2)_01         | 182     | 28                  | 456 206.7         | 5 925 480.6 | 456 188.4       | 5 925 494.2 | 22.8       |               |
| 17/08/2020  | 14:20      | MA_ST64 (2)       | Still | MA_ST64(2)_02         | 183     | 28                  | 456 206.7         | 5 925 480.6 | 456 192.3       | 5 925 487.4 | 15.9       |               |
| 17/08/2020  | 14:21      | MA_ST64 (2)       | Still | MA_ST64(2)_03         | 184     | 28                  | 456 206.7         | 5 925 480.6 | 456 202.3       | 5 925 483.1 | 5.1        |               |
| 17/08/2020  | 14:21      | MA_ST64 (2)       | Still | MA_ST64(2)_04         | 185     | 28                  | 456 206.7         | 5 925 480.6 | 456 204.3       | 5 925 477.7 | 3.8        |               |
| 17/08/2020  | 14:22      | MA_ST64 (2)       | Still | MA_ST64(2)_05         | 186     | 28                  | 456 206.7         | 5 925 480.6 | 456 209.3       | 5 925 472.1 | 8.9        |               |
| 17/08/2020  | 14:23      | MA_ST64 (2)       | Still | MA_ST64(2)_06         | 187     | 28                  | 456 206.7         | 5 925 480.6 | 456 214.4       | 5 925 464.2 | 18.1       |               |
| 17/08/2020  | 14:23      | MA_ST64 (2)       | Video | EOL                   | 188     | 28                  | 456 206.7         | 5 925 480.6 | 456 214.7       | 5 925 461.6 | 20.7       |               |
| 17/08/2020  | 15:05      | MA_ST61           | HG    | NS                    | 189     | 29                  | 456 206.0         | 5 925 508.0 | 456 202.1       | 5 925 508.7 | 3.9        | Not triggered |
| 17/08/2020  | 15:07      | MA_ST61           | HG    | FA/PSD                | 190     | 29                  | 456 206.0         | 5 925 508.0 | 456 198.9       | 5 925 502.3 | 9.1        |               |
| 17/08/2020  | 15:28      | MA_ST64           | HG    | FA/PSD                | 191     | 27                  | 456 206.7         | 5 925 480.6 | 456 201.1       | 5 925 486.9 | 8.4        |               |
| 17/08/2020  | 15:33      | MA_ST63           | HG    | NS                    | 192     | 27                  | 456 177.3         | 5 925 488.6 | 456 182.0       | 5 925 482.4 | 7.7        | Not triggered |
| 17/08/2020  | 15:34      | MA_ST63           | HG    | NS                    | 193     | 27                  | 456 177.3         | 5 925 488.6 | 456 174.6       | 5 925 479.1 | 9.9        | Not triggered |
| 17/08/2020  | 15:38      | MA_ST63           | HG    | FA/PSD                | 194     | 27                  | 456 177.3         | 5 925 488.6 | 456 180.4       | 5 925 497.1 | 9.0        |               |
| 17/08/2020  | 16:08      | MA_ST65           | HG    | FA/PSD                | 195     | 29                  | 453 569.0         | 5 923 291.0 | 453 581.5       | 5 923 297.0 | 13.9       |               |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |      |                       |         |                     |                   |             |                 |             |            |               |
|---|------------|-------------------|------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|---------------|
| Date  | Time [UTC] | Transect/ Station | Type | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes         |
|   |            |                   |      |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |               |
| 17/08/2020  | 16:28      | MA_ST66           | HG   | FA/PSD                | 196     | 24                  | 450 568.0         | 5 920 636.0 | 450 568.4       | 5 920 641.0 | 5.0        |               |
| 17/08/2020  | 16:47      | MA_ST60           | HG   | FA/PSD                | 197     | 23                  | 446 905.8         | 5 919 910.8 | 446 917.0       | 5 919 909.6 | 11.3       |               |
| 17/08/2020  | 16:58      | MA_ST52           | HG   | FA/PSD                | 198     | 27                  | 444 625.1         | 5 920 709.9 | 444 634.0       | 5 920 717.6 | 11.8       |               |
| 18/08/2020  | 09:48      | MA_ST57           | HG   | FA/PSD                | 199     | 32                  | 445 929.0         | 5 921 227.0 | 445 936.3       | 5 921 240.4 | 15.2       |               |
| 18/08/2020  | 10:09      | MA_ST51           | HG   | FA/PSD                | 200     | 34                  | 445 486.5         | 5 922 318.5 | 445 493.3       | 5 922 320.7 | 7.1        |               |
| 18/08/2020  | 10:19      | MA_ST50           | HG   | FA/PSD                | 201     | 36                  | 446 010.4         | 5 923 271.0 | 446 014.0       | 5 923 286.1 | 15.5       |               |
| 18/08/2020  | 10:32      | MA_ST55           | HG   | FA/PSD                | 202     | 36                  | 447 905.0         | 5 923 406.9 | 447 900.6       | 5 923 413.6 | 8.1        |               |
| 18/08/2020  | 10:41      | MA_ST54           | HG   | FA/PSD                | 203     | 36                  | 447 949.0         | 5 923 456.0 | 447 950.6       | 5 923 468.1 | 12.2       |               |
| 18/08/2020  | 10:49      | MA_ST56           | HG   | FA/PSD                | 204     | 37                  | 448 166.2         | 5 923 311.0 | 448 180.1       | 5 923 328.4 | 22.2       |               |
| 18/08/2020  | 11:06      | MA_ST59           | HG   | FA/PSD                | 205     | 38                  | 450 387.0         | 5 924 424.0 | 450 384.0       | 5 924 428.5 | 5.4        |               |
| 18/08/2020  | 11:25      | MA_ST58           | HG   | FA/PSD                | 206     | 38                  | 452 286.0         | 5 926 081.0 | 452 287.9       | 5 926 084.4 | 3.9        |               |
| 18/08/2020  | 12:29      | MA_ST42           | HG   | FA/PSD                | 207     | 39                  | 448 252.6         | 5 927 341.2 | 448 244.0       | 5 927 344.8 | 9.3        |               |
| 18/08/2020  | 12:46      | MA_ST43           | HG   | FA/PSD                | 208     | 37                  | 447 038.0         | 5 926 464.0 | 447 026.4       | 5 926 473.2 | 14.8       |               |
| 18/08/2020  | 12:59      | MA_ST36           | HG   | FA/PSD                | 209     | 38                  | 446 474.6         | 5 927 357.1 | 446 463.2       | 5 927 358.1 | 11.4       |               |
| 18/08/2020  | 13:15      | MA_ST37           | HG   | FA/PSD                | 210     | 37                  | 445 413.5         | 5 926 398.4 | 445 411.2       | 5 926 388.1 | 10.6       |               |
| 18/08/2020  | 13:28      | MA_ST29           | HG   | FA/PSD                | 211     | 38                  | 444 661.0         | 5 927 335.0 | 444 651.1       | 5 927 328.6 | 11.8       |               |
| 18/08/2020  | 13:51      | MA_ST33           | HG   | FA/PSD                | 212     | 35                  | 443 994.3         | 5 925 763.4 | 443 981.8       | 5 925 760.6 | 12.8       |               |
| 18/08/2020  | 14:08      | MA_ST39           | HG   | NS                    | 213     | 32                  | 444 076.2         | 5 924 638.8 | 444 064.3       | 5 924 635.9 | 12.2       | Not triggered |
| 18/08/2020  | 14:11      | MA_ST39           | HG   | FA/PSD                | 214     | 32                  | 444 076.2         | 5 924 638.8 | 444 067.4       | 5 924 629.5 | 12.9       |               |
| 18/08/2020  | 14:27      | MA_ST44           | HG   | FA/PSD                | 215     | 32                  | 445 277.0         | 5 924 515.0 | 445 272.5       | 5 924 509.3 | 7.3        |               |
| 18/08/2020  | 14:42      | MA_ST47           | HG   | FA/PSD                | 216     | 31                  | 445 090.2         | 5 923 234.7 | 445 086.7       | 5 923 234.0 | 3.6        |               |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |      |                       |         |                     |                   |             |                 |             |            |       |
|---|------------|-------------------|------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|-------|
| Date  | Time [UTC] | Transect/ Station | Type | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes |
|   |            |                   |      |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |       |
| 18/08/2020  | 14:57      | MA_ST40           | HG   | FA/PSD                | 217     | 30                  | 443 343.9         | 5 923 890.9 | 443 341.1       | 5 923 891.7 | 3.0        |       |
| 18/08/2020  | 15:13      | MA_ST46           | HG   | FA/PSD                | 218     | 30                  | 443 175.0         | 5 922 829.0 | 443 177.1       | 5 922 838.5 | 9.7        |       |
| 18/08/2020  | 15:23      | MA_ST45           | HG   | FA/PSD                | 219     | 29                  | 443 807.8         | 5 922 836.9 | 443 817.3       | 5 922 832.6 | 10.4       |       |
| 18/08/2020  | 15:33      | MA_ST48           | HG   | FA/PSD                | 220     | 31                  | 443 918.9         | 5 922 130.4 | 443 915.1       | 5 922 128.3 | 4.4        |       |
| 18/08/2020  | 15:51      | MA_ST49           | HG   | FA/PSD                | 221     | 27                  | 442 894.7         | 5 921 292.0 | 442 896.1       | 5 921 283.6 | 8.5        |       |
| 18/08/2020  | 16:08      | MA_ST41           | HG   | FA/PSD                | 222     | 29                  | 441 095.5         | 5 921 805.2 | 441 094.4       | 5 921 801.5 | 3.9        |       |
| 18/08/2020  | 16:21      | MA_ST38           | HG   | FA/PSD                | 223     | 31                  | 441 672.7         | 5 923 361.5 | 441 669.4       | 5 923 359.0 | 4.1        |       |
| 18/08/2020  | 16:41      | MA_ST27           | HG   | FA/PSD                | 224     | 33                  | 439 295.0         | 5 923 381.0 | 439 295.6       | 5 923 375.7 | 5.4        |       |
| 18/08/2020  | 16:50      | MA_ST24           | HG   | FA/PSD                | 225     | 32                  | 438 799.7         | 5 923 486.3 | 438 792.6       | 5 923 498.0 | 13.7       |       |
| 18/08/2020  | 16:58      | MA_ST22           | HG   | FA/PSD                | 226     | 31                  | 438 596.0         | 5 923 566.3 | 438 601.6       | 5 923 569.0 | 6.2        |       |
| 18/08/2020  | 17:09      | MA_ST32           | HG   | FA/PSD                | 227     | 33                  | 438 674.6         | 5 922 704.8 | 438 696.3       | 5 922 699.7 | 22.3       |       |
| 18/08/2020  | 17:29      | MA_ST14           | HG   | FA/PSD                | 228     | 33                  | 436 240.4         | 5 923 662.6 | 436 231.6       | 5 923 656.5 | 10.8       |       |
| 19/08/2020  | 08:46      | MA_ST35           | HG   | FA/PSD                | 229     | 34                  | 441 706.4         | 5 924 346.0 | 441 711.3       | 5 924 338.3 | 9.1        |       |
| 19/08/2020  | 08:56      | MA_ST34           | HG   | FA/PSD                | 230     | 36                  | 442 470.3         | 5 924 795.0 | 442 476.2       | 5 924 807.3 | 13.6       |       |
| 19/08/2020  | 09:06      | MA_ST30           | HG   | FA/PSD                | 231     | 37                  | 442 549.7         | 5 925 731.7 | 442 557.9       | 5 925 737.6 | 10.1       |       |
| 19/08/2020  | 09:16      | MA_ST19           | HG   | FA/PSD                | 232     | 39                  | 442 461.0         | 5 926 594.0 | 442 464.1       | 5 926 593.9 | 3.2        |       |
| 19/08/2020  | 09:26      | MA_ST18           | HG   | FA/PSD                | 233     | 40                  | 443 025.9         | 5 927 366.8 | 443 028.6       | 5 927 370.2 | 4.3        |       |
| 19/08/2020  | 09:44      | MA_ST15           | HG   | FA/PSD                | 234     | 41                  | 441 200.3         | 5 927 398.5 | 441 199.1       | 5 927 398.4 | 1.2        |       |
| 19/08/2020  | 10:15      | MA_ST10           | HG   | FA/PSD                | 235     | 43                  | 439 327.0         | 5 927 350.9 | 439 328.6       | 5 927 348.2 | 3.1        |       |
| 19/08/2020  | 10:30      | MA_ST16           | HG   | FA/PSD                | 236     | 42                  | 440 508.1         | 5 926 557.2 | 440 503.9       | 5 926 558.5 | 4.4        |       |
| 19/08/2020  | 10:41      | MA_ST20           | HG   | FA/PSD                | 237     | 39                  | 440 800.8         | 5 925 686.4 | 440 802.3       | 5 925 688.1 | 2.3        |       |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |      |                       |         |                     |                   |             |                 |             |            |   |
|---|------------|-------------------|------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|---|
| Date  | Time [UTC] | Transect/ Station | Type | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes   |
|   |            |                   |      |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |   |
| 19/08/2020  | 10:51      | MA_ST23           | HG   | FA/PSD                | 238     | 37                  | 441 118.0         | 5 925 220.2 | 441 122.5       | 5 925 222.9 | 5.3        |   |
| 19/08/2020  | 11:00      | MA_ST25           | HG   | FA/PSD                | 239     | 38                  | 441 223.0         | 5 925 039.0 | 441 235.7       | 5 925 039.3 | 12.7       |   |
| 19/08/2020  | 11:10      | MA_ST31           | HG   | FA/PSD                | 240     | 37                  | 441 437.6         | 5 924 729.1 | 441 444.1       | 5 924 722.4 | 9.3        |   |
| 19/08/2020  | 11:25      | MA_ST21           | HG   | FA/PSD                | 241     | 38                  | 439 539.8         | 5 924 525.2 | 439 538.7       | 5 924 525.3 | 1.1        |   |
| 19/08/2020  | 11:43      | MA_ST17           | HG   | FA/PSD                | 242     | 39                  | 439 285.8         | 5 925 699.9 | 439 276.5       | 5 925 698.9 | 9.3        |   |
| 19/08/2020  | 11:56      | MA_ST11           | HG   | FA/PSD                | 243     | 43                  | 438 522.0         | 5 926 615.0 | 438 525.0       | 5 926 611.7 | 4.4        |   |
| 19/08/2020  | 12:10      | MA_ST07           | HG   | FA/PSD                | 244     | 44                  | 436 777.5         | 5 926 461.9 | 436 761.7       | 5 926 443.9 | 24.0       |   |
| 19/08/2020  | 12:28      | MA_ST12           | HG   | FA/PSD                | 245     | 39                  | 437 529.0         | 5 925 669.0 | 437 531.5       | 5 925 677.0 | 8.3        |   |
| 23/08/2020  | 11:33      | MA_ST22           | DG   | NS                    | 308     | 36                  | 438 596.0         | 5 923 566.3 | 438 593.5       | 5 923 560.4 | 6.4        | Grab empty                                    |
| 23/08/2020  | 11:38      | MA_ST22           | DG   | NS                    | 309     | 36                  | 438 596.0         | 5 923 566.3 | 438 595.7       | 5 923 548.3 | 18.0       | Grab empty                                    |
| 23/08/2020  | 11:44      | MA_ST22           | DG   | NS                    | 310     | 36                  | 438 596.0         | 5 923 566.3 | 438 605.1       | 5 923 568.5 | 9.4        | 1 cm bite depth                               |
| 23/08/2020  | 11:50      | MA_ST22           | DG   | NS                    | 311     | 36                  | 438 596.0         | 5 923 566.3 | 438 611.8       | 5 923 572.0 | 16.9       | 1 cm bite depth                               |
| 23/08/2020  | 11:56      | MA_ST22           | DG   | CS                    | 312     | 36                  | 438 596.0         | 5 923 566.3 | 438 605.1       | 5 923 551.2 | 17.6       | 6 cm bite depth accepted due to sediment type |
| 23/08/2020  | 12:25      | MA_ST04           | DG   | NS                    | 313     | 45                  | 434 714.0         | 5 926 692.0 | 434 720.9       | 5 926 694.3 | 7.3        | 4 cm bite depth                               |
| 23/08/2020  | 12:33      | MA_ST04           | DG   | CS                    | 314     | 45                  | 434 714.0         | 5 926 692.0 | 434 719.9       | 5 926 677.4 | 15.7       | 6 cm bite depth accepted due to sediment type |
| 23/08/2020  | 12:48      | MA_ST12           | DG   | CS                    | 315     | 39                  | 437 529.0         | 5 925 669.0 | 437 520.5       | 5 925 669.3 | 8.5        |   |
| 23/08/2020  | 13:03      | MA_ST25           | DG   | CS                    | 316     | 38                  | 441 223.0         | 5 925 039.0 | 441 223.1       | 5 925 038.2 | 0.8        |   |
| 23/08/2020  | 13:20      | MA_ST47           | DG   | CS                    | 317     | 36                  | 445 090.2         | 5 923 234.7 | 445 082.4       | 5 923 224.0 | 13.3       |   |
| 23/08/2020  | 13:41      | MA_ST66           | DG   | CS                    | 318     | 30                  | 450 568.0         | 5 920 636.0 | 450 564.4       | 5 920 636.1 | 3.6        |   |
| 23/08/2020  | 13:59      | MA_ST59           | DG   | CS                    | 319     | 39                  | 450 387.0         | 5 924 424.0 | 450 392.2       | 5 924 435.9 | 12.9       |   |

| Geodetic Parameters: WGS84 UTM Zone 30N, CM 3°W [m] |            |                   |      |                       |         |                     |                   |             |                 |             |            |       |
|---|------------|-------------------|------|-----------------------|---------|---------------------|-------------------|-------------|-----------------|-------------|------------|-------|
| Date  | Time [UTC] | Transect/ Station | Type | Sample Rep/ Still No. | Fix No. | Water Depth [m BSL] | Proposed Location |             | Actual Location |             | Offset [m] | Notes |
|   |            |                   |      |                       |         |                     | Easting           | Northing    | Easting         | Northing    |            |       |
| 23/08/2020  | 14:21      | MA_ST43           | DG   | CS                    | 320     | 39                  | 447 038.0         | 5 926 464.0 | 447 044.7       | 5 926 466.5 | 7.2        |       |
| 23/08/2020  | 14:57      | MA_ST61           | DG   | CS                    | 321     | 34                  | 456 206.0         | 5 925 508.0 | 456 213.5       | 5 925 518.4 | 12.8       |       |
| 23/08/2020  | 15:17      | MA_ST65           | DG   | CS                    | 322     | 35                  | 453 569.0         | 5 923 291.0 | 453 573.0       | 5 923 303.1 | 12.8       |       |
| 24/08/2020  | 08:49      | MA_ST09           | HG   | FA/PSD                | 326     | 35                  | 434 372.4         | 5 924 396.8 | 434 384.2       | 5 924 402.5 | 13.0       |       |
| 24/08/2020  | 09:01      | MA_ST06           | HG   | FA/PSD                | 327     | 37                  | 432 899.8         | 5 925 014.1 | 432 894.4       | 5 925 022.5 | 9.9        |       |
| 24/08/2020  | 09:12      | MA_ST02           | HG   | FA/PSD                | 328     | 39                  | 432 777.0         | 5 926 144.4 | 432 775.0       | 5 926 151.4 | 7.3        |       |
| 24/08/2020  | 09:23      | MA_ST01           | HG   | FA/PSD                | 329     | 39                  | 432 960.8         | 5 927 193.8 | 432 971.4       | 5 927 193.6 | 10.6       |       |
| 24/08/2020  | 09:39      | MA_ST03           | HG   | FA/PSD                | 330     | 40                  | 435 317.0         | 5 927 335.0 | 435 309.0       | 5 927 335.5 | 8.0        |       |
| 24/08/2020  | 09:49      | MA_ST04           | HG   | FA/PSD                | 331     | 40                  | 434 714.0         | 5 926 692.0 | 434 709.3       | 5 926 693.1 | 4.8        |       |
| 24/08/2020  | 09:59      | MA_ST05           | HG   | FA/PSD                | 332     | 39                  | 434 205.7         | 5 926 049.2 | 434 210.4       | 5 926 047.8 | 4.8        |       |
| 24/08/2020  | 10:11      | MA_ST08           | HG   | FA/PSD                | 333     | 38                  | 435 698.0         | 5 925 604.7 | 435 704.2       | 5 925 604.3 | 6.2        |       |
| 24/08/2020  | 10:24      | MA_ST13           | HG   | FA/PSD                | 334     | 34                  | 437 110.2         | 5 924 842.5 | 437 115.3       | 5 924 840.8 | 5.4        |       |

**Notes**  
 UTC = Coordinated Universal Time  
 BSL = Below sea level  
 SOL = Start of line  
 EOL = End of line  
 FA = Fauna sample  
 PSD = Particle size distribution subsample  
 CS = Chemistry sample  
 NF = No fix



## C.2 Grab Log

| Date       | Time [UTC] | Station | Sample | Fix No. | Sample Volume [L] | Sediment Description (including stratigraphy) |                      | Comments (fauna, smell, bioturbation, debris) |
|------------|------------|---------|--------|---------|-------------------|---|----------------------|---|
|            |            |         |        |         |                   | Sediment Type                                 | Sediment Description |   |
| 17/08/2020 | 15:07      | MA_ST61 | FA/PSD | 190     | 8                 | msG   | Sandy gravel         | -   |
| 17/08/2020 | 15:28      | MA_ST64 | FA/PSD | 191     | 9                 | sG  | Sandy gravel         | -   |
| 17/08/2020 | 15:34      | MA_ST63 | FA/PSD | 193     | 9                 | sG  | Sandy gravel         | -   |
| 17/08/2020 | 16:08      | MA_ST65 | FA/PSD | 195     | 10                | gS  | Gravelly sand        | -   |
| 17/08/2020 | 16:28      | MA_ST66 | FA/PSD | 196     | 7                 | S   | Sand                 | -   |
| 17/08/2020 | 16:47      | MA_ST60 | FA/PSD | 197     | 7                 | S   | Sand                 | -   |
| 17/08/2020 | 16:58      | MA_ST52 | FA/PSD | 198     | 6                 | S   | Sand                 | -   |
| 18/08/2020 | 09:48      | MA_ST57 | FA/PSD | 199     | 7                 | gS  | Gravelly sand        | -   |
| 18/08/2020 | 10:09      | MA_ST51 | FA/PSD | 200     | 5                 | gS  | Coarse Sand          | -   |
| 18/08/2020 | 10:19      | MA_ST50 | FA/PSD | 201     | 6                 | gS  | Coarse Sand          | -   |
| 18/08/2020 | 10:32      | MA_ST55 | FA/PSD | 202     | 9                 | gS  | Coarse Sand          | -   |
| 18/08/2020 | 10:41      | MA_ST54 | FA/PSD | 203     | 6                 | gS  | Coarse Sand          | -   |
| 18/08/2020 | 10:49      | MA_ST56 | FA/PSD | 204     | 6                 | gS  | Coarse Sand          | -   |
| 18/08/2020 | 11:06      | MA_ST59 | FA/PSD | 205     | 7                 | gS  | Coarse Sand          | -   |
| 18/08/2020 | 11:25      | MA_ST58 | FA/PSD | 206     | 7                 | msG   | Sandy gravel         | -   |
| 18/08/2020 | 12:29      | MA_ST42 | FA/PSD | 207     | 7                 | msG   | Sandy gravel         | Soft coral ( <i>Alcyonium digitatum</i> )     |
| 18/08/2020 | 12:46      | MA_ST43 | FA/PSD | 208     | 8                 | msG   | Sandy gravel         | -   |
| 18/08/2020 | 12:59      | MA_ST36 | FA/PSD | 209     | 6                 | sG  | Sandy gravel         | -   |
| 18/08/2020 | 13:15      | MA_ST37 | FA/PSD | 210     | 8                 | msG   | Sandy gravel         | -   |
| 18/08/2020 | 13:28      | MA_ST29 | FA/PSD | 211     | 7                 | sG  | Sandy gravel         | -   |

| Date       | Time [UTC] | Station | Sample | Fix No. | Sample Volume [L] | Sediment Description (including stratigraphy) |                      | Comments (fauna, smell, bioturbation, debris) |
|------------|------------|---------|--------|---------|-------------------|---|----------------------|---|
|            |            |         |        |         |                   | Sediment Type                                 | Sediment Description |   |
| 18/08/2020 | 13:51      | MA_ST33 | FA/PSD | 212     | 7                 | sG  | Sandy gravel         | -   |
| 18/08/2020 | 14:11      | MA_ST39 | FA/PSD | 214     | 7                 | sG  | Sandy gravel         | -   |
| 18/08/2020 | 14:27      | MA_ST44 | FA/PSD | 215     | 8                 | sG  | Sandy gravel         | -   |
| 18/08/2020 | 14:42      | MA_ST47 | FA/PSD | 216     | 6                 | sG  | Sandy gravel         | -   |
| 18/08/2020 | 14:57      | MA_ST40 | FA/PSD | 217     | 9                 | sG  | Sandy gravel         | -   |
| 18/08/2020 | 15:13      | MA_ST46 | FA/PSD | 218     | 6                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 15:23      | MA_ST45 | FA/PSD | 219     | 7                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 15:33      | MA_ST48 | FA/PSD | 220     | 7                 | S   | Fine sand            | -   |
| 18/08/2020 | 15:51      | MA_ST49 | FA/PSD | 221     | 7                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 16:08      | MA_ST41 | FA/PSD | 222     | 5                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 16:21      | MA_ST38 | FA/PSD | 223     | 6                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 16:41      | MA_ST27 | FA/PSD | 224     | 9                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 16:50      | MA_ST24 | FA/PSD | 225     | 8                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 16:58      | MA_ST22 | FA/PSD | 226     | 6                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 17:09      | MA_ST32 | FA/PSD | 227     | 7                 | gS  | Coarse sand          | -   |
| 18/08/2020 | 17:29      | MA_ST14 | FA/PSD | 228     | 8                 | gS  | Coarse sand          | -   |
| 19/08/2020 | 08:46      | MA_ST35 | FA/PSD | 229     | 7                 | sG  | Coarse sediments     | -   |
| 19/08/2020 | 08:56      | MA_ST34 | FA/PSD | 230     | 7                 | sG  | Coarse sediments     | -   |
| 19/08/2020 | 09:06      | MA_ST30 | FA/PSD | 231     | 9                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 09:16      | MA_ST19 | FA/PSD | 232     | 6                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 09:26      | MA_ST18 | FA/PSD | 233     | 8                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 09:44      | MA_ST15 | FA/PSD | 234     | 6                 | msG   | Sandy gravel         | -   |

| Date       | Time [UTC] | Station | Sample | Fix No. | Sample Volume [L] | Sediment Description (including stratigraphy) |                      | Comments (fauna, smell, bioturbation, debris) |
|------------|------------|---------|--------|---------|-------------------|---|----------------------|---|
|            |            |         |        |         |                   | Sediment Type                                 | Sediment Description |   |
| 19/08/2020 | 10:15      | MA_ST10 | FA/PSD | 235     | 5                 | sG  | Coarse sediments     | -   |
| 19/08/2020 | 10:30      | MA_ST16 | FA/PSD | 236     | 6                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 10:41      | MA_ST20 | FA/PSD | 237     | 6                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 10:51      | MA_ST23 | FA/PSD | 238     | 7                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 11:00      | MA_ST25 | FA/PSD | 239     | 7                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 11:10      | MA_ST31 | FA/PSD | 240     | 7                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 11:25      | MA_ST21 | FA/PSD | 241     | 8                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 11:43      | MA_ST17 | FA/PSD | 242     | 7                 | msG   | Sandy gravel         | -   |
| 19/08/2020 | 11:56      | MA_ST11 | FA/PSD | 243     | 6                 | sG  | Coarse sediments     | -   |
| 19/08/2020 | 12:10      | MA_ST07 | FA/PSD | 244     | 5                 | sG  | Coarse sediments     | -   |
| 19/08/2020 | 12:28      | MA_ST12 | FA/PSD | 245     | 6                 | msG   | Sandy gravel         | -   |
| 23/08/2020 | 11:56      | MA_ST22 | CS     | 312     | 6*                | gS  | Coarse sand          | -   |
| 23/08/2020 | 12:33      | MA_ST04 | CS     | 314     | 6*                | gS  | Gravelly sand        | -   |
| 23/08/2020 | 12:48      | MA_ST12 | CS     | 315     | 7*                | msG   | Sandy gravel         | -   |
| 23/08/2020 | 13:03      | MA_ST25 | CS     | 316     | 7*                | msG   | Sandy gravel         | -   |
| 23/08/2020 | 13:20      | MA_ST47 | CS     | 317     | 10*               | sG  | Sandy gravel         | -   |
| 23/08/2020 | 13:41      | MA_ST66 | CS     | 318     | 8*                | S   | Sand                 | -   |
| 23/08/2020 | 13:59      | MA_ST59 | CS     | 319     | 7*                | gS  | Coarse Sand          | -   |
| 23/08/2020 | 14:21      | MA_ST43 | CS     | 320     | 7*                | msG   | Sandy gravel         | -   |
| 23/08/2020 | 14:57      | MA_ST61 | CS     | 321     | 7*                | msG   | Sandy gravel         | -   |
| 23/08/2020 | 15:17      | MA_ST65 | CS     | 322     | 11*               | gS  | Coarse Sand          | -   |
| 24/08/2020 | 08:49      | MA_ST09 | FA/PSD | 326     | 6                 | sG  | Sandy gravel         | -   |

| Date       | Time [UTC] | Station | Sample | Fix No. | Sample Volume [L] | Sediment Description (including stratigraphy) |                      | Comments (fauna, smell, bioturbation, debris) |
|------------|------------|---------|--------|---------|-------------------|---|----------------------|---|
|            |            |         |        |         |                   | Sediment Type                                 | Sediment Description |   |
| 24/08/2020 | 09:01      | MA_ST06 | FA/PSD | 327     | 7                 | sG  | Sandy gravel         | -   |
| 24/08/2020 | 09:12      | MA_ST02 | FA/PSD | 328     | 5                 | sG  | Sandy gravel         | -   |
| 24/08/2020 | 09:23      | MA_ST01 | FA/PSD | 329     | 5                 | gS  | Coarse sand          | -   |
| 24/08/2020 | 09:39      | MA_ST03 | FA/PSD | 330     | 7                 | sG  | Sandy gravel         | -   |
| 24/08/2020 | 09:49      | MA_ST04 | FA/PSD | 331     | 6                 | sG  | Sandy gravel         | -   |
| 24/08/2020 | 09:59      | MA_ST05 | FA/PSD | 332     | 6                 | msG   | Sandy gravel         | -   |
| 24/08/2020 | 10:11      | MA_ST08 | FA/PSD | 333     | 7                 | sG  | Coarse sediments     | -   |
| 24/08/2020 | 10:24      | MA_ST13 | FA/PSD | 334     | 6                 | msG   | Sandy gravel         | -   |

Notes

\* = Sample depth [cm]

UTC = Coordinated Universal Time

FA = Faunal sample




PSD = Particle size distribution subsample

CS = Chemistry sample




NS = No sample






### C3 Video and Photographic Log

| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W [m] |         |           |   |   |   |   |
|--|---------|-----------|---|---|---|---|
| Station/<br>Transect                                 | Easting | Northing  | Sediment Description                          | Fauna/Bioturbation/Debris   | Abundance<br>[SACFOR]   | Representative Image  |
| MA_ST04  | SOL     | 434 692.7 | Sandy gravel with pebbles and shell fragments | Faunal turf (Hydrozoa/Bryozoa)<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Hermit crab (Paguridae)<br>Anemones (Ceriantharia)<br>Sea urchin ( <i>Psammechinus miliaris</i> )<br>Polychaetes (Sabellidae)<br>Faunal tubes  | R<br>R<br>O<br>O<br>P<br>P<br>P                               |    |
|  | EOL     | 434 712.7 |   |   |   |   |
| MA_ST11  | SOL     | 438 495.6 | Sandy gravel with pebbles and shell fragments | Hermit crabs ( <i>Pagurus bernhardus</i> )<br>Hydroid ( <i>Nemertesia antennina</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Anemone (Sagartiidae)<br>Anemone (Actiniaria)<br>Anemone (Ceriantharia)<br>Brittlestars (Ophiuroidea)<br>Brittlestars ( <i>Ophiura albida</i> )<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Sea urchin ( <i>Psammechinus miliaris</i> )  | F<br>F<br>R<br>P<br>P<br>P<br>P<br>P<br>R<br>P                |   |
|  | EOL     | 438 540.9 |   |   |   |   |
| MA_ST12  | SOL     | 437 501.2 | Sandy gravel with pebbles and shell fragments | Brittlestars (Ophiuroidea)<br>Brittlestars ( <i>Ophiura albida</i> )<br>Brittlestars ( <i>Ophiothrix fragilis</i> )<br>Hermit crab (Paguridae)<br>Hermit crab ( <i>Pagurus prideaux</i> )<br>Anemone ( <i>Adamsia palliata</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Anemone (Ceriantharia)<br>Polychaetes (Serpulidae)<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Sea urchins ( <i>Psammechinus miliaris</i> ) | P<br>P<br>Locally A-S<br>O<br>O<br>P<br>R<br>P<br>P<br>R<br>P |  |
|  | EOL     | 437 563.2 |   |   |   |   |






| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W [m] |     |          |             |   |  |   |   |
|--|-----|----------|-------------|---|--|---|---|
| Station/<br>Transect                                 |     | Easting  | Northing    | Sediment Description  | Fauna/Bioturbation/Debris  | Abundance<br>[SACFOR]                               | Representative Image  |
| MA_ST19  | SOL | 442.4342 | 5.926.5973  | Sandy gravel with pebbles and shell fragments                         | Brittlestars (Ophiuroidea)<br>Brittlestars ( <i>Ophiura albida</i> )<br>Hermit crab (Paguridae)<br>Faunal turf (Hydrozoa/Bryozoa)<br>Hydroids ( <i>Nemertesia antennina</i> )<br>Polychaetes (Serpulidae)<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Faunal tubes<br>Faunal burrows   | P<br>P<br>O<br>R<br>F<br>P<br>R<br>P<br>P           |    |
|  | EOL | 442.4856 | 5.926.598.1 |   |  |   |   |
| MA_ST22  | SOL | 438.6180 | 5.923.5514  | Gravelly sand with a varying proportion of shells and shell fragments | Brittlestars ( <i>Ophiura ophiura</i> )<br>Brittlestars ( <i>Ophiura albida</i> )<br>Brittlestars ( <i>Ophiothrix fragilis</i> )<br>Hermit crabs (Paguridae)<br>Hydroids ( <i>Nemertesia antennina</i> )<br>Hydroids ( <i>Tubularia indivisa</i> )<br>Anemone (Actiniaria)<br>Anemone ( <i>Utricina</i> sp.)<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Starfish (Asteroidea)<br>Faunal burrows               | P<br>P<br>F<br>O<br>F<br>R<br>P<br>O<br>R<br>O<br>P |   |
|  | EOL | 438.5907 | 5.923.5942  |   |  |   |   |
| MA_ST23  | SOL | 441.0929 | 5.925.2292  | Sandy gravel with pebbles and shell fragments                         | Brittlestars (Ophiuroidea)<br>Brittlestars ( <i>Ophiura albida</i> )<br>Brittlestars ( <i>Ophiothrix fragilis</i> )<br>Hydroids ( <i>Nemertesia antennina</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Hermit crabs ( <i>Pagurus bernhardus</i> )<br>Anemone (Ceriantharia)<br>Polychaetes (Serpulidae)<br>Sea urchins ( <i>Psammechinus miliaris</i> )<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Faunal tubes | P<br>P<br>F<br>F<br>R<br>O<br>P<br>P<br>P<br>R<br>P |  |
|  | EOL | 441.1410 | 5.925.215.1 |   |  |   |   |






| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W [m] |     |           |             |  |   |   |   |
|--|-----|-----------|-------------|--|---|---|---|
| Station/<br>Transect                                 |     | Easting   | Northing    | Sediment Description   | Fauna/Bioturbation/Debris   | Abundance<br>[SACFOR]                               | Representative Image  |
| MA_ST26  | SOL | 441 196.7 | 5 925 054.8 | Gravelly sand with varying proportions of shells and shell fragments | Hermit crab ( <i>Paguridae</i> )<br>Faunal turf ( <i>Hydrozoa/Bryozoa</i> )<br>Starfish ( <i>Asterias rubens</i> )<br>Possible king scallop ( <i>Pecten maximus</i> )<br>Brittlestars ( <i>Ophiuroidea</i> )<br>Brittlestars ( <i>Ophiura albida</i> )<br>Brittlestars ( <i>Ophiothrix fragilis</i> )<br>Soft coral ( <i>Alcyonium digitatum</i> ),<br>Thornback ray ( <i>Raja clavata</i> )<br>Polychaetes ( <i>Serpulidae</i> )<br>Faunal burrows | O<br>R<br>O<br>O<br>P<br>P<br>F<br>R<br>F<br>P<br>P |    |
|  | EOL | 441 239.0 | 5 925 011.4 |  |   |   |   |
| MA_ST28  | SOL | 439 205.4 | 5 923 380.7 | Sandy gravel with pebbles and shell fragments                        | Brittlestars ( <i>Ophiuroidea</i> )<br>Brittlestars ( <i>Ophiura ophiura</i> )<br>Brittlestars ( <i>Ophiura albida</i> )<br>Anemone ( <i>Ceriantharia</i> )<br>Hydroids ( <i>Nemertesia antennina</i> )<br>Hermit crab ( <i>Pagurus prideaux</i> )<br>Anemone ( <i>Adamsia palliata</i> )<br>Sand mason ( <i>Lanice conchilega</i> )<br>Hydroids ( <i>Tubularia indivisa</i> )<br>Faunal turf ( <i>Hydrozoa/Bryozoa</i> )<br>Faunal burrows         | P<br>P<br>P<br>P<br>F<br>O<br>P<br>P<br>R<br>O<br>P |   |
|  | EOL | 439 289.4 | 5 923 381.1 |  |   |   |   |
| MA_ST43  | SOL | 447 059.3 | 5 926 476.0 | Sandy gravel with pebbles and shell fragments                        | Sea urchin ( <i>Psammechinus miliaris</i> )<br>Brittlestars ( <i>Ophiura albida</i> )<br>Hermit crab ( <i>Pagurus bernhardus</i> )<br>Hydroid ( <i>Hydractinia echinata</i> )<br>Faunal turf ( <i>Hydrozoa/Bryozoa</i> )<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Polychaetes ( <i>Serpulidae</i> )  | P<br>P<br>O<br>P<br>R<br>R<br>P                     |  |
|  | EOL | 447 020.7 | 5 926 455.4 |  |   |   |   |






| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W |     |          |            |  |  |   |   |
|--|-----|----------|------------|--|--|---|---|
| Station/<br>Transect                             |     | Easting  | Northing   | Sediment Description   | Fauna/Bioturbation/Debris  | Abundance<br>[SACFOR]   | Representative Image  |
| MA_ST44  | SOL | 445 2982 | 5 924 5308 | Gravelly sand with varying proportions of shells and shell fragments       | Sea urchin ( <i>Psammechinus miliaris</i> )<br>Soft coral ( <i>Alcyonium digitatum</i> )<br>Sea snail (Gastropoda)<br>Faunal turf (Hydrozoa/Bryozoa)<br>Hydroids ( <i>Tubularia indivisa</i> )<br>Faunal burrows   | P<br>R<br>P<br>R<br>R<br>P                                      |    |
|  | EOL | 445 2524 | 5 924 5029 |  |  |   |   |
| MA_ST46  | SOL | 443 1523 | 5 922 8102 | Rippled sand with varying proportions of gravel, shell and shell fragments | Starfish ( <i>Asterias rubens</i> )<br>Sea urchin ( <i>Psammechinus miliaris</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Flatfish (Pleuronectiformes)   | O<br>P<br>R<br>O  |   |
|  | EOL | 443 2110 | 5 922 8336 |  |  |   |   |
| MA_ST53  | SOL | 447 9095 | 5 923 4765 | Sand with varying proportions of gravel, shell and shell fragments         | Greater pipefish ( <i>Syngnathus acus</i> )<br>Dragonet ( <i>Callionymus</i> sp.)<br>Hermit crab ( <i>Pagurus prideaux</i> )<br>Anemone ( <i>Adamsia palliata</i> )<br>Anemone (Actiniaria)<br>Anemone (Ceriantharia)<br>Sea slug (Gastropoda)<br>Starfish ( <i>Asterias rubens</i> )<br>Sea urchin ( <i>Psammechinus miliaris</i> )<br>Hydroids ( <i>Nemertesia antennina</i> )<br>Hydroid ( <i>Tubularia indivisa</i> )<br>Thornback ray ( <i>Raja clavata</i> )<br>Pogge ( <i>Agonus cataphractus</i> ) | O<br>O<br>O<br>P<br>O<br>P<br>O<br>O<br>P<br>F-C<br>R<br>F<br>O |  |
|  | EOL | 447 9668 | 5 923 4464 |  |  |   |   |






| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W |     |           |             |   |  |                            |   |
|--|-----|-----------|-------------|---|--|----------------------------|---|
| Station/<br>Transect                             |     | Easting   | Northing    | Sediment Description  | Fauna/Bioturbation/Debris  | Abundance<br>[SACFOR]      | Representative Image  |
| MA_ST55  | SOL | 447 878.7 | 5 923 416.6 | Sand with varying proportions of gravel, shell and shell fragments            | Hydroids ( <i>Nemertesia antennina</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Hermit crab (Paguridae)<br>Starfish ( <i>Asterias rubens</i> )<br>Sea urchin (Spatangoida)<br>Polychaetes (Serpulidae) | F<br>R<br>O<br>O<br>P<br>P |    |
|  | EOL | 447 927.3 | 5 923 395.7 |   |  |                            |   |
| MA_ST56  | SOL | 448 144.9 | 5 923 327.0 | Sand with varying proportions of gravel, shell and shell fragments            | Hermit crab (Paguridae)<br>Brittlestar (Ophiuroidea)<br>Sea urchin ( <i>Psammechinus miliaris</i> )<br>Anemone (Actiniaria)<br>Colonial anemone (Zooantharia)  | O<br>P<br>P<br>O<br>P      |   |
|  | EOL | 448 186.1 | 5 923 293.0 |   |  |                            |   |
| MA_ST57  | SOL | 445 939.0 | 5 921 251.2 | Gravelly sand with varying proportions of pebbles, shells and shell fragments | Faunal turf (Hydrozoa/Bryozoa)<br>Starfish ( <i>Asterias rubens</i> )<br>Sea squirt (Ascidiacea)   | R<br>O<br>P                |  |
|  | EOL | 445 947.5 | 5 921 215.3 |   |  |                            |   |





| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W |     |          |            |   |  |  |   |
|--|-----|----------|------------|---|--|--|---|
| Station/<br>Transect                             |     | Easting  | Northing   | Sediment Description  | Fauna/Bioturbation/Debris  | Abundance<br>[SACFOR]                          | Representative Image  |
| MA_ST58  | SOL | 452 2589 | 5 926 0795 | Gravelly sand with varying proportions of shells and shell fragments          | Brittlestar ( <i>Ophiura ophiura</i> )<br>Anemone ( <i>Actinaria</i> )<br>Sea urchin ( <i>Psammechinus miliaris</i> )<br>Starfish ( <i>Asterias rubens</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Catshark ( <i>Scyliorhinus</i> sp.)<br>Hermit crab ( <i>Pagurus prideaux</i> )<br>Anemone ( <i>Adamsia palliata</i> )<br>Faunal burrows  | P<br>P<br>P<br>O<br>R<br>F<br>O<br>P<br>P      |    |
|  | EOL | 452 3006 | 5 926 1017 |   |  |  |   |
| MA_ST59  | SOL | 450 3613 | 5 924 4319 | Sand with varying proportions of gravel, shell and shell fragments            | Brittlestar ( <i>Ophiura ophiura</i> )<br>Brittlestars ( <i>Ophiura albida</i> )<br>Soft coral ( <i>Alyonium digitatum</i> )<br>Faunal turf (Hydrozoa/Bryozoa)   | P<br>P<br>R<br>R                               |   |
|  | EOL | 450 4068 | 5 924 4397 |   |  |  |   |
| MA_ST62  | SOL | 456 1423 | 5 925 4979 | Gravelly sand with varying proportions of pebbles, shells and shell fragments | Hermit crab ( <i>Paguridae</i> )<br>Starfish ( <i>Asterias rubens</i> )<br>Brittlestar ( <i>Ophiura ophiura</i> )<br>Brittlestar ( <i>Ophiura albida</i> )<br>Catshark ( <i>Scyliorhinus canicula</i> )<br>Anemone ( <i>Actinaria</i> )<br>Hydroids ( <i>Nemertesia antennina</i> ),<br>Polychaetes ( <i>Polychaeta</i> )<br>Crab ( <i>Inachus</i> sp.)<br>Anemone ( <i>Ceriantharia</i> ) | O<br>O<br>P<br>P<br>F<br>O<br>F<br>P<br>P<br>P |  |
|  | EOL | 456 2352 | 5 925 5188 |   |  |  |   |



| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3 <sup>rd</sup> W |     |           |             |   |   |   |   |
|--|-----|-----------|-------------|---|---|---|---|
| Station/<br>Transect   |     | Easting   | Northing    | Sediment Description  | Fauna/Bioturbation/Debris   | Abundance<br>[SACFOR]   | Representative Image  |
| MA_ST63  | SOL | 456 148.4 | 5 925 482.0 | Gravelly sand with varying proportions of pebbles, shells and shell fragments | Faunal turf (Hydrozoa/Bryozoa)<br>Possible starfish ( <i>Asteria rubens</i> )<br>Sea urchin ( <i>Psammochinus miliaris</i> )<br>Hermit crab (Paguridae)<br>Faunal tubes   | R<br>O<br>P<br>O<br>P   |    |
|  | EOL | 456 181.3 | 5 925 512.8 |   |   |   |   |
| MA_ST64  | SOL | 456 125.1 | 5 925 474.3 | Gravelly sand with varying proportions of pebbles, shells and shell fragments | Sea urchin ( <i>Psammochinus miliaris</i> )<br>Faunal turf (Hydrozoa/Bryozoa)<br>Brittlestars (Ophiuroidea)<br>Brittlestar ( <i>Ophiura albida</i> )<br>Anemone (Ceriantharia)<br>Ray (Rajidae)<br>Hermit crab (Paguridae)<br>Starfish ( <i>Asterias rubens</i> )<br>Faunal burrows   | P<br>R<br>P<br>P<br>P<br>F<br>O<br>O<br>P                               |   |
|  | EOL | 456 218.7 | 5 925 495.6 |   |   |   |   |
| MA_ST64(2)   | SOL | 456 179.2 | 5 925 492.0 | Gravelly sand with varying proportions of pebbles, shells and shell fragments | Brittlestar ( <i>Ophiura ophiura</i> )<br>Sea urchin ( <i>Echinus esculentus</i> )<br>Sea urchin ( <i>Psammochinus miliaris</i> )<br>Hermit crab (Paguridae)<br>Anemone (Ceriantharia)<br>Anemone (Actiniaria)<br>Anemone ( <i>Utricina</i> sp.)<br>Fish (Pisces)<br>Soft coral ( <i>Alcyonium digitatum</i> ),<br>Faunal turf (Hydrozoa/Bryozoa)<br>Starfish ( <i>Asterias rubens</i> )<br>Polychaetes (Serpulidae)<br>Polychaetes (Sabellidae)<br>Sea squirt (Ascidiacea)<br>Faunal tubes | P<br>F<br>P<br>O<br>P<br>P<br>O<br>P<br>R<br>O<br>O<br>P<br>P<br>P<br>P |  |
|  | EOL | 456 214.7 | 5 925 461.6 |   |   |   |   |



| Geodetic Parameters: WGS84, UTM Zone 30N, CM 3°W   |     |          |            |  |   |                                      |  |
|--|-----|----------|------------|--|---|--------------------------------------|--|
| Station/<br>Transect   |     | Easting  | Northing   | Sediment Description   | Fauna/Bioturbation/Debris   | Abundance<br>[SACFOR]                | Representative Image   |
| MA_ST65  | SOL | 453 5435 | 5 923 2903 | Sand with varying proportions of gravel, shell and shell fragments | Hermit crab ( <i>Paguridae</i> )<br>Hermit crab ( <i>Pagurus bernhardus</i> )<br>Catshark ( <i>Scyliorhinus</i> sp.)<br>Anemone ( <i>Ceriantharia</i> )<br>Anemone ( <i>Actiniaria</i> )<br>Anemone ( <i>Metridium</i> sp.)<br>Faunal turf ( <i>Hydrozoa/Bryozoa</i> )<br>Starfish ( <i>Asterias rubens</i> ) | O<br>O<br>F<br>O<br>O<br>R<br>R<br>O |   |
|  | EOL | 453 5525 | 5 923 2755 |  |   |                                      |  |
| MA_ST66  | SOL | 450 5403 | 5 920 6383 | Sand with varying proportions of gravel, shell and shell fragments | Flatfish ( <i>Pleuronectiformes</i> )<br>Starfish ( <i>Asterias rubens</i> )<br>Anemone ( <i>Actiniaria</i> )<br>Faunal turf ( <i>Hydrozoa/Bryozoa</i> )<br>Crab ( <i>Inachus</i> sp.)  | P<br>O<br>P<br>R<br>P                |  |
|  | EOL | 450 5943 | 5 920 6353 |  |   |                                      |  |
| <p>Notes</p> <p>SACFOR = Semi-quantitative abundance scale from Superabundant, Abundant, Common, Frequent, Occasional to Rare</p> <p>P = Present</p> <p>SOL = Start of line</p> <p>EOL = End of line</p> <p>Laser distance (green) = 10 cm</p> |     |          |            |  |   |                                      |  |



# **Appendix D**

Sediment Particle Size and Grab  
Sample Photographs


STATION: MA\_ST01



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 5.88           | 5.88           |
| 5600          | -2.50          | 4.77           | 10.65          |
| 4000          | -2.00          | 5.52           | 16.17          |
| 2800          | -1.50          | 5.82           | 21.99          |
| 2000          | -1.00          | 6.78           | 28.76          |
| 1400          | -0.50          | 4.99           | 33.75          |
| 1000          | 0.00           | 3.86           | 37.61          |
| *707.11       | *0.50          | 13.75          | 51.35          |
| *500.00       | *1.00          | 17.39          | 68.75          |
| *353.55       | *1.50          | 14.92          | 83.67          |
| *250.00       | *2.00          | 7.89           | 91.56          |
| *176.78       | *2.50          | 2.02           | 93.58          |
| *125.00       | *3.00          | 0.04           | 93.63          |
| *88.39        | *3.50          | 0.01           | 93.63          |
| *62.50        | *4.00          | 0.45           | 94.08          |
| *44.19        | *4.50          | 0.56           | 94.64          |
| *31.25        | *5.00          | 0.37           | 95.02          |
| *22.10        | *5.50          | 0.27           | 95.28          |
| *15.63        | *6.00          | 0.35           | 95.63          |
| *11.05        | *6.50          | 0.53           | 96.16          |
| *7.81         | *7.00          | 0.68           | 96.84          |
| *5.52         | *7.50          | 0.76           | 97.60          |
| *3.91         | *8.00          | 0.74           | 98.34          |
| *2.76         | *8.50          | 0.61           | 98.96          |
| *1.95         | *9.00          | 0.44           | 99.39          |
| *1.38         | *9.50          | 0.28           | 99.67          |
| *0.98         | *10.00         | 0.18           | 99.85          |
| * < 0.98      | * > 10.00      | 0.15           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Median [µm] <sup>†</sup>    | 732   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.45  |                    |
| Mean [µm] <sup>‡</sup>      | 1010  | Very coarse sand   |
| Mean [phi] <sup>‡</sup>     | -0.01 |                    |
| Sorting [µm] <sup>†</sup>   | 4.30  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.10  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.13  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.13 |                    |
| Gravel [%] <sup>#</sup>     | 28.76 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 65.32 |                    |
| Fines [%] <sup>#</sup>      | 5.92  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST02



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.08           | 1.08           |
| 8000          | -3.00          | 9.10           | 10.18          |
| 5600          | -2.50          | 5.22           | 15.41          |
| 4000          | -2.00          | 3.94           | 19.35          |
| 2800          | -1.50          | 5.90           | 25.25          |
| 2000          | -1.00          | 4.82           | 30.07          |
| 1400          | -0.50          | 4.51           | 34.58          |
| 1000          | 0.00           | 3.92           | 38.50          |
| *707.11       | *0.50          | 14.77          | 53.27          |
| *500.00       | *1.00          | 17.10          | 70.37          |
| *353.55       | *1.50          | 14.65          | 85.02          |
| *250.00       | *2.00          | 8.54           | 93.56          |
| *176.78       | *2.50          | 2.82           | 96.39          |
| *125.00       | *3.00          | 0.21           | 96.60          |
| *88.39        | *3.50          | 0.00           | 96.60          |
| *62.50        | *4.00          | 0.33           | 96.93          |
| *44.19        | *4.50          | 0.41           | 97.34          |
| *31.25        | *5.00          | 0.24           | 97.58          |
| *22.10        | *5.50          | 0.12           | 97.70          |
| *15.63        | *6.00          | 0.19           | 97.89          |
| *11.05        | *6.50          | 0.30           | 98.19          |
| *7.81         | *7.00          | 0.39           | 98.58          |
| *5.52         | *7.50          | 0.42           | 99.00          |
| *3.91         | *8.00          | 0.38           | 99.38          |
| *2.76         | *8.50          | 0.30           | 99.68          |
| *1.95         | *9.00          | 0.21           | 99.89          |
| *1.38         | *9.50          | 0.11           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| < 0.98        | > 10.00        | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Mode 3 [µm] <sup>†</sup>    | 3400  | Granule            |
| Median [µm] <sup>†</sup>    | 763   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.39  |                    |
| Mean [µm] <sup>‡</sup>      | 1138  | Very coarse sand   |
| Mean [phi] <sup>‡</sup>     | -0.19 |                    |
| Sorting [µm] <sup>†</sup>   | 3.50  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.81  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.39  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.39 |                    |
| Gravel [%] <sup>#</sup>     | 30.07 |                    |
| Sand [%] <sup>#</sup>       | 66.86 | Sandy gravel       |
| Fines [%] <sup>#</sup>      | 3.07  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST03



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 5.11           | 5.11           |
| 5600          | -2.50          | 4.60           | 9.71           |
| 4000          | -2.00          | 5.48           | 15.19          |
| 2800          | -1.50          | 6.05           | 21.24          |
| 2000          | -1.00          | 4.28           | 25.53          |
| 1400          | -0.50          | 3.75           | 29.27          |
| 1000          | 0.00           | 2.68           | 31.95          |
| *707.11       | *0.50          | 12.72          | 44.68          |
| *500.00       | *1.00          | 18.64          | 63.32          |
| *353.55       | *1.50          | 17.61          | 80.93          |
| *250.00       | *2.00          | 10.00          | 90.92          |
| *176.78       | *2.50          | 2.75           | 93.67          |
| *125.00       | *3.00          | 0.09           | 93.76          |
| *88.39        | *3.50          | 0.00           | 93.77          |
| *62.50        | *4.00          | 0.42           | 94.19          |
| *44.19        | *4.50          | 0.62           | 94.82          |
| *31.25        | *5.00          | 0.41           | 95.22          |
| *22.10        | *5.50          | 0.25           | 95.47          |
| *15.63        | *6.00          | 0.32           | 95.79          |
| *11.05        | *6.50          | 0.50           | 96.29          |
| *7.81         | *7.00          | 0.66           | 96.95          |
| *5.52         | *7.50          | 0.74           | 97.69          |
| *3.91         | *8.00          | 0.71           | 98.40          |
| *2.76         | *8.50          | 0.59           | 98.99          |
| *1.95         | *9.00          | 0.42           | 99.42          |
| *1.38         | *9.50          | 0.27           | 99.69          |
| *0.98         | *10.00         | 0.18           | 99.87          |
| * < 0.98      | * > 10.00      | 0.13           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                |       |                    |
|----------------|-------|--------------------|
| Mode 1 [µm]    | 604   | Coarse sand        |
| Mode 2 [µm]    | 3400  | Granule            |
| Mode 3 [µm]    | 9600  | Medium pebble      |
| Median [µm]    | 640   | Coarse sand        |
| Median [phi]   | 0.64  |                    |
| Mean [µm]      | 919   | Coarse sand        |
| Mean [phi]     | 0.12  |                    |
| Sorting [µm]   | 4.19  | Very poorly sorted |
| Sorting [phi]  | 2.07  |                    |
| Skewness [µm]  | 0.19  | Coarse skewed      |
| Skewness [phi] | -0.19 |                    |
| Gravel [%]     | 25.53 |                    |
| Sand [%]       | 68.67 | Gravelly sand      |
| Fines [%]      | 5.81  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST04



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 12.43          | 12.43          |
| 22 400        | -4.50          | 0.00           | 12.43          |
| 16 000        | -4.00          | 0.00           | 12.43          |
| 11 200        | -3.50          | 0.57           | 13.00          |
| 8000          | -3.00          | 0.54           | 13.54          |
| 5600          | -2.50          | 1.84           | 15.38          |
| 4000          | -2.00          | 2.28           | 17.65          |
| 2800          | -1.50          | 2.52           | 20.18          |
| 2000          | -1.00          | 3.69           | 23.87          |
| 1400          | -0.50          | 3.90           | 27.77          |
| 1000          | 0.00           | 4.01           | 31.78          |
| *707.11       | *0.50          | 15.58          | 47.36          |
| *500.00       | *1.00          | 18.26          | 65.62          |
| *353.55       | *1.50          | 15.64          | 81.26          |
| *250.00       | *2.00          | 9.00           | 90.26          |
| *176.78       | *2.50          | 2.84           | 93.10          |
| *125.00       | *3.00          | 0.15           | 93.25          |
| *88.39        | *3.50          | 0.01           | 93.26          |
| *62.50        | *4.00          | 0.46           | 93.72          |
| *44.19        | *4.50          | 0.59           | 94.31          |
| *31.25        | *5.00          | 0.41           | 94.72          |
| *22.10        | *5.50          | 0.31           | 95.02          |
| *15.63        | *6.00          | 0.41           | 95.43          |
| *11.05        | *6.50          | 0.61           | 96.04          |
| *7.81         | *7.00          | 0.78           | 96.83          |
| *5.52         | *7.50          | 0.85           | 97.68          |
| *3.91         | *8.00          | 0.79           | 98.46          |
| *2.76         | *8.50          | 0.62           | 99.09          |
| *1.95         | *9.00          | 0.42           | 99.51          |
| *1.38         | *9.50          | 0.26           | 99.77          |
| *0.98         | *10.00         | 0.16           | 99.93          |
| * < 0.98      | * > 10.00      | 0.07           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 38250 | Very coarse pebble |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule            |
| Median [µm] <sup>†</sup>    | 673   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.57  |                    |
| Mean [µm] <sup>‡</sup>      | 1030  | Very coarse sand   |
| Mean [phi] <sup>‡</sup>     | -0.04 |                    |
| Sorting [µm] <sup>†</sup>   | 6.19  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.63  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.28  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.28 |                    |
| Gravel [%] <sup>#</sup>     | 23.87 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 69.85 |                    |
| Fines [%] <sup>#</sup>      | 6.28  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)

STATION: MA\_ST05




No image available

FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 4.08           | 4.08           |
| 5600          | -2.50          | 3.42           | 7.50           |
| 4000          | -2.00          | 4.66           | 12.16          |
| 2800          | -1.50          | 4.86           | 17.02          |
| 2000          | -1.00          | 5.66           | 22.68          |
| 1400          | -0.50          | 4.36           | 27.04          |
| 1000          | 0.00           | 4.08           | 31.12          |
| *707.11       | *0.50          | 14.02          | 45.14          |
| *500.00       | *1.00          | 18.09          | 63.23          |
| *353.55       | *1.50          | 16.49          | 79.72          |
| *250.00       | *2.00          | 9.81           | 89.53          |
| *176.78       | *2.50          | 3.24           | 92.77          |
| *125.00       | *3.00          | 0.30           | 93.07          |
| *88.39        | *3.50          | 0.04           | 93.11          |
| *62.50        | *4.00          | 0.49           | 93.61          |
| *44.19        | *4.50          | 0.64           | 94.25          |
| *31.25        | *5.00          | 0.44           | 94.68          |
| *22.10        | *5.50          | 0.30           | 94.98          |
| *15.63        | *6.00          | 0.37           | 95.35          |
| *11.05        | *6.50          | 0.56           | 95.91          |
| *7.81         | *7.00          | 0.74           | 96.64          |
| *5.52         | *7.50          | 0.82           | 97.47          |
| *3.91         | *8.00          | 0.79           | 98.26          |
| *2.76         | *8.50          | 0.66           | 98.92          |
| *1.95         | *9.00          | 0.46           | 99.38          |
| *1.38         | *9.50          | 0.29           | 99.67          |
| *0.98         | *10.00         | 0.19           | 99.86          |
| * < 0.98      | * > 10.00      | 0.14           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Median [µm] <sup>†</sup>    | 644   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.63  |                    |
| Mean [µm] <sup>‡</sup>      | 839   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.25  |                    |
| Sorting [µm] <sup>†</sup>   | 4.28  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.10  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.09  | Symmetrical        |
| Skewness [phi] <sup>‡</sup> | -0.09 |                    |
| Gravel [%] <sup>#</sup>     | 22.68 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 70.92 |                    |
| Fines [%] <sup>#</sup>      | 6.39  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST06



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 2.16           | 2.16           |
| 8000          | -3.00          | 0.86           | 3.02           |
| 5600          | -2.50          | 1.85           | 4.87           |
| 4000          | -2.00          | 3.26           | 8.13           |
| 2800          | -1.50          | 3.81           | 11.94          |
| 2000          | -1.00          | 5.52           | 17.46          |
| 1400          | -0.50          | 4.80           | 22.27          |
| 1000          | 0.00           | 4.87           | 27.13          |
| *707.11       | *0.50          | 18.88          | 46.02          |
| *500.00       | *1.00          | 21.05          | 67.07          |
| *353.55       | *1.50          | 17.31          | 84.38          |
| *250.00       | *2.00          | 9.79           | 94.16          |
| *176.78       | *2.50          | 3.22           | 97.39          |
| *125.00       | *3.00          | 0.27           | 97.66          |
| *88.39        | *3.50          | 0.00           | 97.66          |
| *62.50        | *4.00          | 0.30           | 97.96          |
| *44.19        | *4.50          | 0.36           | 98.31          |
| *31.25        | *5.00          | 0.15           | 98.46          |
| *22.10        | *5.50          | 0.00           | 98.46          |
| *15.63        | *6.00          | 0.06           | 98.52          |
| *11.05        | *6.50          | 0.21           | 98.74          |
| *7.81         | *7.00          | 0.28           | 99.02          |
| *5.52         | *7.50          | 0.30           | 99.33          |
| *3.91         | *8.00          | 0.28           | 99.61          |
| *2.76         | *8.50          | 0.23           | 99.84          |
| *1.95         | *9.00          | 0.16           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| < 0.98        | > 10.00        | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 662   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.59  |                    |
| Mean [µm] <sup>‡</sup>      | 802   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.32  | Coarse sand        |
| Sorting [µm] <sup>†</sup>   | 2.55  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.35  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.32  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.32 |                    |
| Gravel [%] <sup>#</sup>     | 17.46 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 80.49 |                    |
| Fines [%] <sup>#</sup>      | 2.04  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST07



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 1.87           | 1.87           |
| 5600          | -2.50          | 3.57           | 5.44           |
| 4000          | -2.00          | 3.47           | 8.91           |
| 2800          | -1.50          | 4.95           | 13.86          |
| 2000          | -1.00          | 5.65           | 19.51          |
| 1400          | -0.50          | 6.12           | 25.63          |
| 1000          | 0.00           | 5.28           | 30.92          |
| *707.11       | *0.50          | 14.15          | 45.07          |
| *500.00       | *1.00          | 18.95          | 64.02          |
| *353.55       | *1.50          | 17.71          | 81.74          |
| *250.00       | *2.00          | 10.41          | 92.14          |
| *176.78       | *2.50          | 3.13           | 95.27          |
| *125.00       | *3.00          | 0.17           | 95.44          |
| *88.39        | *3.50          | 0.00           | 95.44          |
| *62.50        | *4.00          | 0.25           | 95.69          |
| *44.19        | *4.50          | 0.50           | 96.19          |
| *31.25        | *5.00          | 0.33           | 96.52          |
| *22.10        | *5.50          | 0.18           | 96.70          |
| *15.63        | *6.00          | 0.22           | 96.93          |
| *11.05        | *6.50          | 0.37           | 97.30          |
| *7.81         | *7.00          | 0.51           | 97.81          |
| *5.52         | *7.50          | 0.57           | 98.38          |
| *3.91         | *8.00          | 0.54           | 98.92          |
| *2.76         | *8.50          | 0.45           | 99.37          |
| *1.95         | *9.00          | 0.32           | 99.69          |
| *1.38         | *9.50          | 0.20           | 99.89          |
| *0.98         | *10.00         | 0.11           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                  |
|-----------------------------|-------|------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand      |
| Mode 2 [µm] <sup>†</sup>    | 1700  | Very coarse sand |
| Mode 3 [µm] <sup>†</sup>    | -     | -                |
| Median [µm] <sup>‡</sup>    | 646   | Coarse sand      |
| Median [phi] <sup>‡</sup>   | 0.63  |                  |
| Mean [µm] <sup>‡</sup>      | 805   | Coarse sand      |
| Mean [phi] <sup>‡</sup>     | 0.31  |                  |
| Sorting [µm] <sup>‡</sup>   | 2.80  | Poorly sorted    |
| Sorting [phi] <sup>‡</sup>  | 1.49  |                  |
| Skewness [µm] <sup>‡</sup>  | 0.30  | Coarse skewed    |
| Skewness [phi] <sup>‡</sup> | -0.30 |                  |
| Gravel [%] <sup>#</sup>     | 19.51 |                  |
| Sand [%] <sup>#</sup>       | 76.18 | Gravelly sand    |
| Fines [%] <sup>#</sup>      | 4.31  |                  |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST08



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.66           | 1.66           |
| 8000          | -3.00          | 1.73           | 3.39           |
| 5600          | -2.50          | 4.56           | 7.94           |
| 4000          | -2.00          | 5.25           | 13.19          |
| 2800          | -1.50          | 6.20           | 19.40          |
| 2000          | -1.00          | 5.34           | 24.74          |
| 1400          | -0.50          | 5.29           | 30.03          |
| 1000          | 0.00           | 4.05           | 34.08          |
| *707.11       | *0.50          | 13.44          | 47.52          |
| *500.00       | *1.00          | 17.76          | 65.28          |
| *353.55       | *1.50          | 16.36          | 81.64          |
| *250.00       | *2.00          | 9.79           | 91.43          |
| *176.78       | *2.50          | 3.26           | 94.69          |
| *125.00       | *3.00          | 0.30           | 94.98          |
| *88.39        | *3.50          | 0.01           | 94.99          |
| *62.50        | *4.00          | 0.38           | 95.37          |
| *44.19        | *4.50          | 0.53           | 95.90          |
| *31.25        | *5.00          | 0.34           | 96.24          |
| *22.10        | *5.50          | 0.20           | 96.45          |
| *15.63        | *6.00          | 0.26           | 96.71          |
| *11.05        | *6.50          | 0.41           | 97.12          |
| *7.81         | *7.00          | 0.55           | 97.67          |
| *5.52         | *7.50          | 0.61           | 98.27          |
| *3.91         | *8.00          | 0.58           | 98.85          |
| *2.76         | *8.50          | 0.47           | 99.32          |
| *1.95         | *9.00          | 0.33           | 99.65          |
| *1.38         | *9.50          | 0.21           | 99.86          |
| *0.98         | *10.00         | 0.13           | 99.99          |
| * < 0.98      | * > 10.00      | 0.01           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | 3400  | Granule       |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 674   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.57  |               |
| Mean [µm] <sup>†</sup>      | 907   | Coarse sand   |
| Mean [phi] <sup>†</sup>     | 0.14  | Coarse sand   |
| Sorting [µm] <sup>†</sup>   | 3.50  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.81  |               |
| Skewness [µm] <sup>†</sup>  | 0.22  | Coarse skewed |
| Skewness [phi] <sup>†</sup> | -0.22 |               |
| Gravel [%] <sup>#</sup>     | 24.74 |               |
| Sand [%] <sup>#</sup>       | 70.63 | Gravelly sand |
| Fines [%] <sup>#</sup>      | 4.63  |               |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST09



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 1.95           | 1.95           |
| 5600          | -2.50          | 5.70           | 7.65           |
| 4000          | -2.00          | 4.25           | 11.90          |
| 2800          | -1.50          | 5.63           | 17.52          |
| 2000          | -1.00          | 5.82           | 23.35          |
| 1400          | -0.50          | 5.38           | 28.73          |
| 1000          | 0.00           | 4.74           | 33.47          |
| *707.11       | *0.50          | 15.11          | 48.58          |
| *500.00       | *1.00          | 18.14          | 66.72          |
| *353.55       | *1.50          | 16.29          | 83.02          |
| *250.00       | *2.00          | 10.21          | 93.22          |
| *176.78       | *2.50          | 3.94           | 97.17          |
| *125.00       | *3.00          | 0.56           | 97.72          |
| *88.39        | *3.50          | 0.00           | 97.73          |
| *62.50        | *4.00          | 0.24           | 97.97          |
| *44.19        | *4.50          | 0.36           | 98.33          |
| *31.25        | *5.00          | 0.21           | 98.53          |
| *22.10        | *5.50          | 0.02           | 98.55          |
| *15.63        | *6.00          | 0.06           | 98.61          |
| *11.05        | *6.50          | 0.21           | 98.82          |
| *7.81         | *7.00          | 0.28           | 99.10          |
| *5.52         | *7.50          | 0.29           | 99.40          |
| *3.91         | *8.00          | 0.26           | 99.66          |
| *2.76         | *8.50          | 0.20           | 99.86          |
| *1.95         | *9.00          | 0.14           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 6800  | Fine pebble        |
| Median [µm] <sup>†</sup>    | 688   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.54  |                    |
| Mean [µm] <sup>‡</sup>      | 899   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.15  |                    |
| Sorting [µm] <sup>†</sup>   | 2.91  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.54  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.34  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.34 |                    |
| Gravel [%] <sup>#</sup>     | 23.35 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 74.62 |                    |
| Fines [%] <sup>#</sup>      | 2.03  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST10



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.04           | 1.04           |
| 8000          | -3.00          | 5.46           | 6.50           |
| 5600          | -2.50          | 3.78           | 10.28          |
| 4000          | -2.00          | 4.57           | 14.85          |
| 2800          | -1.50          | 4.73           | 19.58          |
| 2000          | -1.00          | 5.08           | 24.66          |
| 1400          | -0.50          | 3.91           | 28.56          |
| 1000          | 0.00           | 3.51           | 32.08          |
| *707.11       | *0.50          | 13.04          | 45.12          |
| *500.00       | *1.00          | 18.51          | 63.62          |
| *353.55       | *1.50          | 17.71          | 81.33          |
| *250.00       | *2.00          | 10.49          | 91.83          |
| *176.78       | *2.50          | 3.12           | 94.95          |
| *125.00       | *3.00          | 0.15           | 95.10          |
| *88.39        | *3.50          | 0.00           | 95.10          |
| *62.50        | *4.00          | 0.23           | 95.33          |
| *44.19        | *4.50          | 0.47           | 95.79          |
| *31.25        | *5.00          | 0.32           | 96.12          |
| *22.10        | *5.50          | 0.19           | 96.31          |
| *15.63        | *6.00          | 0.26           | 96.57          |
| *11.05        | *6.50          | 0.44           | 97.00          |
| *7.81         | *7.00          | 0.60           | 97.60          |
| *5.52         | *7.50          | 0.66           | 98.26          |
| *3.91         | *8.00          | 0.61           | 98.87          |
| *2.76         | *8.50          | 0.49           | 99.35          |
| *1.95         | *9.00          | 0.33           | 99.68          |
| *1.38         | *9.50          | 0.21           | 99.90          |
| *0.98         | *10.00         | 0.10           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule            |
| Median [µm] <sup>†</sup>    | 645   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.63  |                    |
| Mean [µm] <sup>‡</sup>      | 915   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.13  |                    |
| Sorting [µm] <sup>†</sup>   | 3.38  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.76  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.36  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.36 |                    |
| Gravel [%] <sup>#</sup>     | 24.66 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 70.67 |                    |
| Fines [%] <sup>#</sup>      | 4.67  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST11



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 2.47           | 2.47           |
| 5600          | -2.50          | 4.48           | 6.95           |
| 4000          | -2.00          | 4.85           | 11.79          |
| 2800          | -1.50          | 5.41           | 17.21          |
| 2000          | -1.00          | 6.14           | 23.35          |
| 1400          | -0.50          | 4.80           | 28.15          |
| 1000          | 0.00           | 3.41           | 31.56          |
| *707.11       | *0.50          | 12.12          | 43.68          |
| *500.00       | *1.00          | 17.48          | 61.15          |
| *353.55       | *1.50          | 17.33          | 78.49          |
| *250.00       | *2.00          | 11.03          | 89.52          |
| *176.78       | *2.50          | 3.88           | 93.40          |
| *125.00       | *3.00          | 0.39           | 93.79          |
| *88.39        | *3.50          | 0.00           | 93.79          |
| *62.50        | *4.00          | 0.31           | 94.10          |
| *44.19        | *4.50          | 0.58           | 94.68          |
| *31.25        | *5.00          | 0.44           | 95.12          |
| *22.10        | *5.50          | 0.29           | 95.41          |
| *15.63        | *6.00          | 0.34           | 95.75          |
| *11.05        | *6.50          | 0.52           | 96.26          |
| *7.81         | *7.00          | 0.69           | 96.95          |
| *5.52         | *7.50          | 0.76           | 97.71          |
| *3.91         | *8.00          | 0.72           | 98.44          |
| *2.76         | *8.50          | 0.59           | 99.03          |
| *1.95         | *9.00          | 0.42           | 99.44          |
| *1.38         | *9.50          | 0.27           | 99.71          |
| *0.98         | *10.00         | 0.17           | 99.88          |
| * < 0.98      | * > 10.00      | 0.12           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION






STATION: MA\_ST12



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.14           | 0.14           |
| 8000          | -3.00          | 0.29           | 0.43           |
| 5600          | -2.50          | 5.09           | 5.52           |
| 4000          | -2.00          | 6.02           | 11.54          |
| 2800          | -1.50          | 5.01           | 16.55          |
| 2000          | -1.00          | 6.13           | 22.68          |
| 1400          | -0.50          | 5.54           | 28.22          |
| 1000          | 0.00           | 4.32           | 32.54          |
| *707.11       | *0.50          | 12.96          | 45.50          |
| *500.00       | *1.00          | 16.45          | 61.95          |
| *353.55       | *1.50          | 15.29          | 77.24          |
| *250.00       | *2.00          | 9.75           | 86.99          |
| *176.78       | *2.50          | 3.72           | 90.71          |
| *125.00       | *3.00          | 0.48           | 91.19          |
| *88.39        | *3.50          | 0.07           | 91.25          |
| *62.50        | *4.00          | 0.51           | 91.76          |
| *44.19        | *4.50          | 0.68           | 92.44          |
| *31.25        | *5.00          | 0.53           | 92.98          |
| *22.10        | *5.50          | 0.43           | 93.40          |
| *15.63        | *6.00          | 0.54           | 93.94          |
| *11.05        | *6.50          | 0.78           | 94.72          |
| *7.81         | *7.00          | 0.99           | 95.70          |
| *5.52         | *7.50          | 1.08           | 96.78          |
| *3.91         | *8.00          | 1.01           | 97.79          |
| *2.76         | *8.50          | 0.81           | 98.60          |
| *1.95         | *9.00          | 0.56           | 99.16          |
| *1.38         | *9.50          | 0.35           | 99.51          |
| *0.98         | *10.00         | 0.23           | 99.74          |
| * < 0.98      | * > 10.00      | 0.26           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION




STATION: MA\_ST13



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.88           | 0.88           |
| 8000          | -3.00          | 1.31           | 2.19           |
| 5600          | -2.50          | 2.85           | 5.04           |
| 4000          | -2.00          | 3.98           | 9.02           |
| 2800          | -1.50          | 3.83           | 12.85          |
| 2000          | -1.00          | 3.45           | 16.30          |
| 1400          | -0.50          | 2.81           | 19.12          |
| 1000          | 0.00           | 3.33           | 22.45          |
| *707.11       | *0.50          | 19.09          | 41.53          |
| *500.00       | *1.00          | 23.02          | 64.55          |
| *353.55       | *1.50          | 18.28          | 82.83          |
| *250.00       | *2.00          | 9.05           | 91.88          |
| *176.78       | *2.50          | 2.25           | 94.14          |
| *125.00       | *3.00          | 0.06           | 94.20          |
| *88.39        | *3.50          | 0.07           | 94.27          |
| *62.50        | *4.00          | 0.50           | 94.77          |
| *44.19        | *4.50          | 0.53           | 95.30          |
| *31.25        | *5.00          | 0.31           | 95.62          |
| *22.10        | *5.50          | 0.21           | 95.83          |
| *15.63        | *6.00          | 0.31           | 96.14          |
| *11.05        | *6.50          | 0.48           | 96.62          |
| *7.81         | *7.00          | 0.62           | 97.24          |
| *5.52         | *7.50          | 0.69           | 97.93          |
| *3.91         | *8.00          | 0.67           | 98.60          |
| *2.76         | *8.50          | 0.55           | 99.15          |
| *1.95         | *9.00          | 0.39           | 99.55          |
| *1.38         | *9.50          | 0.25           | 99.80          |
| *0.98         | *10.00         | 0.17           | 99.97          |
| * < 0.98      | * > 10.00      | 0.03           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | 4800  | Fine pebble   |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 622   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.68  |               |
| Mean [µm] <sup>‡</sup>      | 757   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.40  |               |
| Sorting [µm] <sup>‡</sup>   | 3.18  | Poorly sorted |
| Sorting [phi] <sup>‡</sup>  | 1.67  |               |
| Skewness [µm] <sup>‡</sup>  | 0.14  | Coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.14 |               |
| Gravel [%] <sup>#</sup>     | 16.30 | Gravelly sand |
| Sand [%] <sup>#</sup>       | 78.47 |               |
| Fines [%] <sup>#</sup>      | 5.23  |               |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST14



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.72           | 0.72           |
| 8000          | -3.00          | 1.55           | 2.27           |
| 5600          | -2.50          | 1.41           | 3.68           |
| 4000          | -2.00          | 3.15           | 6.83           |
| 2800          | -1.50          | 3.16           | 9.99           |
| 2000          | -1.00          | 4.65           | 14.64          |
| 1400          | -0.50          | 5.15           | 19.79          |
| 1000          | 0.00           | 5.59           | 25.39          |
| *707.11       | *0.50          | 15.89          | 41.28          |
| *500.00       | *1.00          | 18.98          | 60.25          |
| *353.55       | *1.50          | 16.67          | 76.93          |
| *250.00       | *2.00          | 10.24          | 87.16          |
| *176.78       | *2.50          | 4.07           | 91.23          |
| *125.00       | *3.00          | 0.95           | 92.18          |
| *88.39        | *3.50          | 0.37           | 92.56          |
| *62.50        | *4.00          | 0.58           | 93.14          |
| *44.19        | *4.50          | 0.60           | 93.73          |
| *31.25        | *5.00          | 0.42           | 94.15          |
| *22.10        | *5.50          | 0.34           | 94.49          |
| *15.63        | *6.00          | 0.43           | 94.93          |
| *11.05        | *6.50          | 0.62           | 95.55          |
| *7.81         | *7.00          | 0.80           | 96.35          |
| *5.52         | *7.50          | 0.89           | 97.24          |
| *3.91         | *8.00          | 0.86           | 98.10          |
| *2.76         | *8.50          | 0.71           | 98.81          |
| *1.95         | *9.00          | 0.51           | 99.32          |
| *1.38         | *9.50          | 0.32           | 99.64          |
| *0.98         | *10.00         | 0.21           | 99.85          |
| * < 0.98      | * > 10.00      | 0.15           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | 4800  | Fine pebble   |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 603   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.73  | Coarse sand   |
| Mean [µm] <sup>‡</sup>      | 673   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.57  | Coarse sand   |
| Sorting [µm] <sup>†</sup>   | 3.84  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.94  | Poorly sorted |
| Skewness [µm] <sup>‡</sup>  | -0.05 | Symmetrical   |
| Skewness [phi] <sup>‡</sup> | 0.05  | Symmetrical   |
| Gravel [%] <sup>#</sup>     | 14.64 | Gravelly sand |
| Sand [%] <sup>#</sup>       | 78.50 |               |
| Fines [%] <sup>#</sup>      | 6.86  |               |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST15



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 3.51           | 3.51           |
| 8000          | -3.00          | 7.04           | 10.55          |
| 5600          | -2.50          | 4.52           | 15.07          |
| 4000          | -2.00          | 4.88           | 19.95          |
| 2800          | -1.50          | 4.55           | 24.50          |
| 2000          | -1.00          | 4.29           | 28.79          |
| 1400          | -0.50          | 3.82           | 32.61          |
| 1000          | 0.00           | 2.69           | 35.30          |
| *707.11       | *0.50          | 11.91          | 47.21          |
| *500.00       | *1.00          | 15.67          | 62.88          |
| *353.55       | *1.50          | 15.27          | 78.15          |
| *250.00       | *2.00          | 10.34          | 88.49          |
| *176.78       | *2.50          | 4.32           | 92.81          |
| *125.00       | *3.00          | 0.74           | 93.55          |
| *88.39        | *3.50          | 0.01           | 93.55          |
| *62.50        | *4.00          | 0.39           | 93.94          |
| *44.19        | *4.50          | 0.59           | 94.53          |
| *31.25        | *5.00          | 0.44           | 94.98          |
| *22.10        | *5.50          | 0.31           | 95.29          |
| *15.63        | *6.00          | 0.38           | 95.66          |
| *11.05        | *6.50          | 0.56           | 96.23          |
| *7.81         | *7.00          | 0.74           | 96.96          |
| *5.52         | *7.50          | 0.80           | 97.76          |
| *3.91         | *8.00          | 0.74           | 98.51          |
| *2.76         | *8.50          | 0.59           | 99.10          |
| *1.95         | *9.00          | 0.40           | 99.50          |
| *1.38         | *9.50          | 0.25           | 99.75          |
| *0.98         | *10.00         | 0.16           | 99.91          |
| * < 0.98      | * > 10.00      | 0.09           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Mode 3 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Median [µm] <sup>†</sup>    | 665   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.59  |                    |
| Mean [µm] <sup>‡</sup>      | 1005  | Very coarse sand   |
| Mean [phi] <sup>‡</sup>     | -0.01 |                    |
| Sorting [µm] <sup>†</sup>   | 4.99  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.32  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.19  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.19 |                    |
| Gravel [%] <sup>#</sup>     | 28.79 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 65.15 |                    |
| Fines [%] <sup>#</sup>      | 6.06  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST16



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.85           | 0.85           |
| 11 200        | -3.50          | 0.00           | 0.85           |
| 8000          | -3.00          | 4.25           | 5.10           |
| 5600          | -2.50          | 3.50           | 8.60           |
| 4000          | -2.00          | 6.31           | 14.92          |
| 2800          | -1.50          | 5.77           | 20.69          |
| 2000          | -1.00          | 6.03           | 26.72          |
| 1400          | -0.50          | 5.25           | 31.97          |
| 1000          | 0.00           | 3.70           | 35.67          |
| *707.11       | *0.50          | 12.65          | 48.32          |
| *500.00       | *1.00          | 16.40          | 64.72          |
| *353.55       | *1.50          | 15.77          | 80.49          |
| *250.00       | *2.00          | 10.45          | 90.94          |
| *176.78       | *2.50          | 4.15           | 95.09          |
| *125.00       | *3.00          | 0.53           | 95.62          |
| *88.39        | *3.50          | 0.00           | 95.62          |
| *62.50        | *4.00          | 0.20           | 95.82          |
| *44.19        | *4.50          | 0.47           | 96.29          |
| *31.25        | *5.00          | 0.35           | 96.64          |
| *22.10        | *5.50          | 0.21           | 96.84          |
| *15.63        | *6.00          | 0.24           | 97.08          |
| *11.05        | *6.50          | 0.38           | 97.47          |
| *7.81         | *7.00          | 0.51           | 97.97          |
| *5.52         | *7.50          | 0.56           | 98.53          |
| *3.91         | *8.00          | 0.52           | 99.05          |
| *2.76         | *8.50          | 0.41           | 99.46          |
| *1.95         | *9.00          | 0.28           | 99.74          |
| *1.38         | *9.50          | 0.18           | 99.92          |
| *0.98         | *10.00         | 0.08           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule            |
| Median [µm] <sup>†</sup>    | 682   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.55  |                    |
| Mean [µm] <sup>‡</sup>      | 930   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.11  |                    |
| Sorting [µm] <sup>†</sup>   | 3.31  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.73  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.34  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.34 |                    |
| Gravel [%] <sup>#</sup>     | 26.72 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 69.11 |                    |
| Fines [%] <sup>#</sup>      | 4.18  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST17



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.93           | 0.93           |
| 8000          | -3.00          | 0.42           | 1.35           |
| 5600          | -2.50          | 3.18           | 4.52           |
| 4000          | -2.00          | 4.79           | 9.31           |
| 2800          | -1.50          | 5.12           | 14.43          |
| 2000          | -1.00          | 5.47           | 19.90          |
| 1400          | -0.50          | 4.74           | 24.64          |
| 1000          | 0.00           | 3.87           | 28.51          |
| *707.11       | *0.50          | 13.23          | 41.74          |
| *500.00       | *1.00          | 18.42          | 60.16          |
| *353.55       | *1.50          | 17.66          | 77.82          |
| *250.00       | *2.00          | 10.81          | 88.63          |
| *176.78       | *2.50          | 3.59           | 92.21          |
| *125.00       | *3.00          | 0.32           | 92.53          |
| *88.39        | *3.50          | 0.00           | 92.54          |
| *62.50        | *4.00          | 0.41           | 92.95          |
| *44.19        | *4.50          | 0.65           | 93.60          |
| *31.25        | *5.00          | 0.48           | 94.08          |
| *22.10        | *5.50          | 0.33           | 94.41          |
| *15.63        | *6.00          | 0.39           | 94.80          |
| *11.05        | *6.50          | 0.58           | 95.38          |
| *7.81         | *7.00          | 0.78           | 96.16          |
| *5.52         | *7.50          | 0.89           | 97.04          |
| *3.91         | *8.00          | 0.87           | 97.91          |
| *2.76         | *8.50          | 0.74           | 98.65          |
| *1.95         | *9.00          | 0.54           | 99.18          |
| *1.38         | *9.50          | 0.35           | 99.53          |
| *0.98         | *10.00         | 0.23           | 99.76          |
| * < 0.98      | * > 10.00      | 0.24           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 605   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.72  |                    |
| Mean [µm] <sup>‡</sup>      | 764   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.39  |                    |
| Sorting [µm] <sup>†</sup>   | 4.25  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.09  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.03  | Symmetrical        |
| Skewness [phi] <sup>‡</sup> | -0.03 |                    |
| Gravel [%] <sup>#</sup>     | 19.90 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 73.04 |                    |
| Fines [%] <sup>#</sup>      | 7.05  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST18



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 6.70           | 6.70           |
| 5600          | -2.50          | 4.94           | 11.65          |
| 4000          | -2.00          | 4.12           | 15.77          |
| 2800          | -1.50          | 4.40           | 20.17          |
| 2000          | -1.00          | 4.44           | 24.60          |
| 1400          | -0.50          | 3.78           | 28.38          |
| 1000          | 0.00           | 3.17           | 31.56          |
| *707.11       | *0.50          | 14.48          | 46.04          |
| *500.00       | *1.00          | 18.71          | 64.75          |
| *353.55       | *1.50          | 16.73          | 81.48          |
| *250.00       | *2.00          | 9.56           | 91.04          |
| *176.78       | *2.50          | 2.88           | 93.92          |
| *125.00       | *3.00          | 0.18           | 94.11          |
| *88.39        | *3.50          | 0.00           | 94.11          |
| *62.50        | *4.00          | 0.37           | 94.48          |
| *44.19        | *4.50          | 0.55           | 95.03          |
| *31.25        | *5.00          | 0.38           | 95.41          |
| *22.10        | *5.50          | 0.25           | 95.66          |
| *15.63        | *6.00          | 0.32           | 95.98          |
| *11.05        | *6.50          | 0.50           | 96.48          |
| *7.81         | *7.00          | 0.66           | 97.14          |
| *5.52         | *7.50          | 0.73           | 97.88          |
| *3.91         | *8.00          | 0.69           | 98.57          |
| *2.76         | *8.50          | 0.56           | 99.13          |
| *1.95         | *9.00          | 0.39           | 99.52          |
| *1.38         | *9.50          | 0.25           | 99.77          |
| *0.98         | *10.00         | 0.16           | 99.93          |
| * < 0.98      | * > 10.00      | 0.07           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule            |
| Median [µm] <sup>†</sup>    | 657   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.61  |                    |
| Mean [µm] <sup>‡</sup>      | 941   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.09  |                    |
| Sorting [µm] <sup>†</sup>   | 4.15  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.05  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.21  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.21 |                    |
| Gravel [%] <sup>#</sup>     | 24.60 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 69.87 |                    |
| Fines [%] <sup>#</sup>      | 5.52  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST19



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 3.57           | 3.57           |
| 8000          | -3.00          | 1.42           | 4.99           |
| 5600          | -2.50          | 3.46           | 8.45           |
| 4000          | -2.00          | 4.20           | 12.65          |
| 2800          | -1.50          | 5.73           | 18.38          |
| 2000          | -1.00          | 5.49           | 23.87          |
| 1400          | -0.50          | 4.86           | 28.73          |
| 1000          | 0.00           | 3.65           | 32.38          |
| *707.11       | *0.50          | 13.30          | 45.69          |
| *500.00       | *1.00          | 17.92          | 63.60          |
| *353.55       | *1.50          | 16.67          | 80.27          |
| *250.00       | *2.00          | 9.90           | 90.18          |
| *176.78       | *2.50          | 3.13           | 93.31          |
| *125.00       | *3.00          | 0.23           | 93.54          |
| *88.39        | *3.50          | 0.00           | 93.54          |
| *62.50        | *4.00          | 0.31           | 93.86          |
| *44.19        | *4.50          | 0.56           | 94.42          |
| *31.25        | *5.00          | 0.42           | 94.84          |
| *22.10        | *5.50          | 0.30           | 95.14          |
| *15.63        | *6.00          | 0.37           | 95.50          |
| *11.05        | *6.50          | 0.56           | 96.06          |
| *7.81         | *7.00          | 0.74           | 96.79          |
| *5.52         | *7.50          | 0.82           | 97.61          |
| *3.91         | *8.00          | 0.77           | 98.38          |
| *2.76         | *8.50          | 0.62           | 99.01          |
| *1.95         | *9.00          | 0.43           | 99.44          |
| *1.38         | *9.50          | 0.27           | 99.71          |
| *0.98         | *10.00         | 0.17           | 99.88          |
| * < 0.98      | * > 10.00      | 0.12           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 13600 | Medium pebble      |
| Median [µm] <sup>†</sup>    | 650   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.62  |                    |
| Mean [µm] <sup>‡</sup>      | 869   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.20  |                    |
| Sorting [µm] <sup>†</sup>   | 4.29  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.10  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.12  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.12 |                    |
| Gravel [%] <sup>#</sup>     | 23.87 |                    |
| Sand [%] <sup>#</sup>       | 69.99 | Gravelly sand      |
| Fines [%] <sup>#</sup>      | 6.14  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST20



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 2.41           | 2.41           |
| 5600          | -2.50          | 1.94           | 4.35           |
| 4000          | -2.00          | 4.74           | 9.08           |
| 2800          | -1.50          | 5.13           | 14.21          |
| 2000          | -1.00          | 5.22           | 19.43          |
| 1400          | -0.50          | 4.92           | 24.35          |
| 1000          | 0.00           | 3.56           | 27.91          |
| *707.11       | *0.50          | 11.92          | 39.83          |
| *500.00       | *1.00          | 16.51          | 56.34          |
| *353.55       | *1.50          | 16.80          | 73.14          |
| *250.00       | *2.00          | 11.81          | 84.95          |
| *176.78       | *2.50          | 5.13           | 90.08          |
| *125.00       | *3.00          | 0.94           | 91.02          |
| *88.39        | *3.50          | 0.01           | 91.03          |
| *62.50        | *4.00          | 0.49           | 91.51          |
| *44.19        | *4.50          | 0.78           | 92.29          |
| *31.25        | *5.00          | 0.64           | 92.93          |
| *22.10        | *5.50          | 0.47           | 93.40          |
| *15.63        | *6.00          | 0.53           | 93.94          |
| *11.05        | *6.50          | 0.74           | 94.68          |
| *7.81         | *7.00          | 0.95           | 95.63          |
| *5.52         | *7.50          | 1.05           | 96.69          |
| *3.91         | *8.00          | 1.00           | 97.69          |
| *2.76         | *8.50          | 0.82           | 98.51          |
| *1.95         | *9.00          | 0.59           | 99.10          |
| *1.38         | *9.50          | 0.38           | 99.48          |
| *0.98         | *10.00         | 0.25           | 99.72          |
| < 0.98        | > 10.00        | 0.28           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                     |
|-----------------------------|-------|---------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand         |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule             |
| Mode 3 [µm] <sup>†</sup>    | -     | -                   |
| Median [µm] <sup>†</sup>    | 571   | Coarse sand         |
| Median [phi] <sup>†</sup>   | 0.81  |                     |
| Mean [µm] <sup>‡</sup>      | 716   | Coarse sand         |
| Mean [phi] <sup>‡</sup>     | 0.48  |                     |
| Sorting [µm] <sup>†</sup>   | 4.58  | Very poorly sorted  |
| Sorting [phi] <sup>†</sup>  | 2.20  |                     |
| Skewness [µm] <sup>‡</sup>  | 0.00  | Symmetrical         |
| Skewness [phi] <sup>‡</sup> | 0.00  |                     |
| Gravel [%] <sup>#</sup>     | 19.43 | Gravelly muddy sand |
| Sand [%] <sup>#</sup>       | 72.08 |                     |
| Fines [%] <sup>#</sup>      | 8.49  |                     |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST21



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.65           | 0.65           |
| 8000          | -3.00          | 3.56           | 4.21           |
| 5600          | -2.50          | 7.41           | 11.62          |
| 4000          | -2.00          | 5.57           | 17.19          |
| 2800          | -1.50          | 4.31           | 21.50          |
| 2000          | -1.00          | 4.00           | 25.50          |
| 1400          | -0.50          | 3.26           | 28.76          |
| 1000          | 0.00           | 2.82           | 31.58          |
| *707.11       | *0.50          | 15.24          | 46.82          |
| *500.00       | *1.00          | 20.12          | 66.94          |
| *353.55       | *1.50          | 17.50          | 84.44          |
| *250.00       | *2.00          | 9.38           | 93.83          |
| *176.78       | *2.50          | 2.40           | 96.23          |
| *125.00       | *3.00          | 0.04           | 96.27          |
| *88.39        | *3.50          | 0.00           | 96.27          |
| *62.50        | *4.00          | 0.17           | 96.44          |
| *44.19        | *4.50          | 0.37           | 96.81          |
| *31.25        | *5.00          | 0.24           | 97.05          |
| *22.10        | *5.50          | 0.12           | 97.17          |
| *15.63        | *6.00          | 0.17           | 97.34          |
| *11.05        | *6.50          | 0.31           | 97.65          |
| *7.81         | *7.00          | 0.44           | 98.08          |
| *5.52         | *7.50          | 0.50           | 98.58          |
| *3.91         | *8.00          | 0.48           | 99.06          |
| *2.76         | *8.50          | 0.40           | 99.46          |
| *1.95         | *9.00          | 0.29           | 99.76          |
| *1.38         | *9.50          | 0.19           | 99.95          |
| *0.98         | *10.00         | 0.05           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 6800  | Fine pebble        |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 669   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.58  |                    |
| Mean [µm] <sup>‡</sup>      | 1009  | Very coarse sand   |
| Mean [phi] <sup>‡</sup>     | -0.01 | Poorly sorted      |
| Sorting [µm] <sup>‡</sup>   | 3.21  |                    |
| Sorting [phi] <sup>‡</sup>  | 1.68  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.43  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.43 |                    |
| Gravel [%] <sup>#</sup>     | 25.50 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 70.93 |                    |
| Fines [%] <sup>#</sup>      | 3.56  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST22



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.33           | 0.33           |
| 8000          | -3.00          | 0.23           | 0.56           |
| 5600          | -2.50          | 1.62           | 2.18           |
| 4000          | -2.00          | 1.43           | 3.61           |
| 2800          | -1.50          | 2.63           | 6.24           |
| 2000          | -1.00          | 3.60           | 9.84           |
| 1400          | -0.50          | 3.39           | 13.23          |
| 1000          | 0.00           | 3.34           | 16.56          |
| *707.11       | *0.50          | 18.35          | 34.91          |
| *500.00       | *1.00          | 24.83          | 59.75          |
| *353.55       | *1.50          | 21.65          | 81.40          |
| *250.00       | *2.00          | 11.19          | 92.59          |
| *176.78       | *2.50          | 2.50           | 95.09          |
| *125.00       | *3.00          | 0.02           | 95.12          |
| *88.39        | *3.50          | 0.00           | 95.12          |
| *62.50        | *4.00          | 0.34           | 95.46          |
| *44.19        | *4.50          | 0.51           | 95.97          |
| *31.25        | *5.00          | 0.29           | 96.26          |
| *22.10        | *5.50          | 0.14           | 96.40          |
| *15.63        | *6.00          | 0.22           | 96.61          |
| *11.05        | *6.50          | 0.41           | 97.03          |
| *7.81         | *7.00          | 0.57           | 97.59          |
| *5.52         | *7.50          | 0.63           | 98.23          |
| *3.91         | *8.00          | 0.61           | 98.84          |
| *2.76         | *8.50          | 0.50           | 99.34          |
| *1.95         | *9.00          | 0.36           | 99.70          |
| *1.38         | *9.50          | 0.24           | 99.94          |
| *0.98         | *10.00         | 0.06           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | -     | -             |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 573   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.80  |               |
| Mean [µm] <sup>‡</sup>      | 583   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.78  |               |
| Sorting [µm] <sup>†</sup>   | 2.09  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.06  |               |
| Skewness [µm] <sup>‡</sup>  | 0.12  | Coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.12 |               |
| Gravel [%] <sup>#</sup>     | 9.84  | Gravelly sand |
| Sand [%] <sup>#</sup>       | 85.62 |               |
| Fines [%] <sup>#</sup>      | 4.54  |               |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST23



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 1.66           | 1.66           |
| 5600          | -2.50          | 6.12           | 7.78           |
| 4000          | -2.00          | 4.85           | 12.64          |
| 2800          | -1.50          | 4.43           | 17.07          |
| 2000          | -1.00          | 5.78           | 22.85          |
| 1400          | -0.50          | 4.52           | 27.37          |
| 1000          | 0.00           | 3.45           | 30.81          |
| *707.11       | *0.50          | 12.87          | 43.68          |
| *500.00       | *1.00          | 17.51          | 61.19          |
| *353.55       | *1.50          | 16.78          | 77.97          |
| *250.00       | *2.00          | 10.36          | 88.32          |
| *176.78       | *2.50          | 3.41           | 91.73          |
| *125.00       | *3.00          | 0.26           | 91.99          |
| *88.39        | *3.50          | 0.00           | 91.99          |
| *62.50        | *4.00          | 0.31           | 92.30          |
| *44.19        | *4.50          | 0.63           | 92.93          |
| *31.25        | *5.00          | 0.51           | 93.44          |
| *22.10        | *5.50          | 0.37           | 93.81          |
| *15.63        | *6.00          | 0.44           | 94.26          |
| *11.05        | *6.50          | 0.68           | 94.93          |
| *7.81         | *7.00          | 0.91           | 95.84          |
| *5.52         | *7.50          | 1.02           | 96.86          |
| *3.91         | *8.00          | 0.98           | 97.84          |
| *2.76         | *8.50          | 0.80           | 98.63          |
| *1.95         | *9.00          | 0.55           | 99.19          |
| *1.38         | *9.50          | 0.35           | 99.53          |
| *0.98         | *10.00         | 0.22           | 99.76          |
| < 0.98        | > 10.00        | 0.24           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION




STATION: MA\_ST24



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.54           | 0.54           |
| 5600          | -2.50          | 0.30           | 0.83           |
| 4000          | -2.00          | 0.45           | 1.28           |
| 2800          | -1.50          | 0.80           | 2.08           |
| 2000          | -1.00          | 1.27           | 3.35           |
| 1400          | -0.50          | 1.88           | 5.24           |
| 1000          | 0.00           | 3.52           | 8.76           |
| *707.11       | *0.50          | 29.49          | 38.25          |
| *500.00       | *1.00          | 30.33          | 68.57          |
| *353.55       | *1.50          | 19.62          | 88.19          |
| *250.00       | *2.00          | 7.05           | 95.24          |
| *176.78       | *2.50          | 0.77           | 96.01          |
| *125.00       | *3.00          | 0.00           | 96.01          |
| *88.39        | *3.50          | 0.16           | 96.17          |
| *62.50        | *4.00          | 0.60           | 96.77          |
| *44.19        | *4.50          | 0.46           | 97.23          |
| *31.25        | *5.00          | 0.12           | 97.35          |
| *22.10        | *5.50          | 0.08           | 97.43          |
| *15.63        | *6.00          | 0.18           | 97.61          |
| *11.05        | *6.50          | 0.35           | 97.97          |
| *7.81         | *7.00          | 0.44           | 98.40          |
| *5.52         | *7.50          | 0.47           | 98.87          |
| *3.91         | *8.00          | 0.45           | 99.32          |
| *2.76         | *8.50          | 0.37           | 99.69          |
| *1.95         | *9.00          | 0.21           | 99.90          |
| *1.38         | *9.50          | 0.08           | 99.98          |
| *0.98         | *10.00         | 0.02           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                   |
|-----------------------------|-------|-------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand       |
| Mode 2 [µm] <sup>†</sup>    | -     | -                 |
| Mode 3 [µm] <sup>†</sup>    | -     | -                 |
| Median [µm] <sup>†</sup>    | 618   | Coarse sand       |
| Median [phi] <sup>†</sup>   | 0.69  |                   |
| Mean [µm] <sup>‡</sup>      | 600   | Coarse sand       |
| Mean [phi] <sup>‡</sup>     | 0.74  |                   |
| Sorting [µm] <sup>†</sup>   | 1.63  | Moderately sorted |
| Sorting [phi] <sup>†</sup>  | 0.70  |                   |
| Skewness [µm] <sup>‡</sup>  | -0.06 | Symmetrical       |
| Skewness [phi] <sup>‡</sup> | 0.06  |                   |
| Gravel [%] <sup>#</sup>     | 3.35  |                   |
| Sand [%] <sup>#</sup>       | 93.42 | Sand              |
| Fines [%] <sup>#</sup>      | 3.23  |                   |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST25



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 1.02           | 1.02           |
| 5600          | -2.50          | 3.54           | 4.56           |
| 4000          | -2.00          | 3.07           | 7.63           |
| 2800          | -1.50          | 4.70           | 12.33          |
| 2000          | -1.00          | 6.01           | 18.34          |
| 1400          | -0.50          | 4.84           | 23.18          |
| 1000          | 0.00           | 3.75           | 26.93          |
| *707.11       | *0.50          | 14.51          | 41.44          |
| *500.00       | *1.00          | 19.23          | 60.67          |
| *353.55       | *1.50          | 18.37          | 79.04          |
| *250.00       | *2.00          | 11.46          | 90.50          |
| *176.78       | *2.50          | 3.80           | 94.30          |
| *125.00       | *3.00          | 0.26           | 94.56          |
| *88.39        | *3.50          | 0.00           | 94.56          |
| *62.50        | *4.00          | 0.15           | 94.70          |
| *44.19        | *4.50          | 0.47           | 95.17          |
| *31.25        | *5.00          | 0.38           | 95.56          |
| *22.10        | *5.50          | 0.23           | 95.79          |
| *15.63        | *6.00          | 0.27           | 96.06          |
| *11.05        | *6.50          | 0.44           | 96.50          |
| *7.81         | *7.00          | 0.62           | 97.12          |
| *5.52         | *7.50          | 0.71           | 97.84          |
| *3.91         | *8.00          | 0.69           | 98.53          |
| *2.76         | *8.50          | 0.58           | 99.11          |
| *1.95         | *9.00          | 0.41           | 99.52          |
| *1.38         | *9.50          | 0.27           | 99.79          |
| *0.98         | *10.00         | 0.18           | 99.97          |
| < 0.98        | > 10.00        | 0.03           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule       |
| Mode 3 [µm] <sup>†</sup>    | 6800  | Fine pebble   |
| Median [µm] <sup>†</sup>    | 606   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.72  |               |
| Mean [µm] <sup>‡</sup>      | 749   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.42  | Coarse sand   |
| Sorting [µm] <sup>†</sup>   | 3.35  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.75  |               |
| Skewness [µm] <sup>‡</sup>  | 0.12  | Coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.12 |               |
| Gravel [%] <sup>#</sup>     | 18.34 | Gravelly sand |
| Sand [%] <sup>#</sup>       | 76.36 |               |
| Fines [%] <sup>#</sup>      | 5.30  |               |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST27



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.44           | 0.44           |
| 5600          | -2.50          | 1.32           | 1.77           |
| 4000          | -2.00          | 2.74           | 4.50           |
| 2800          | -1.50          | 2.74           | 7.25           |
| 2000          | -1.00          | 5.74           | 12.98          |
| 1400          | -0.50          | 7.17           | 20.15          |
| 1000          | 0.00           | 4.95           | 25.11          |
| *707.11       | *0.50          | 9.89           | 35.00          |
| *500.00       | *1.00          | 19.37          | 54.37          |
| *353.55       | *1.50          | 22.22          | 76.59          |
| *250.00       | *2.00          | 14.30          | 90.89          |
| *176.78       | *2.50          | 4.30           | 95.20          |
| *125.00       | *3.00          | 0.21           | 95.40          |
| *88.39        | *3.50          | 0.00           | 95.40          |
| *62.50        | *4.00          | 0.14           | 95.54          |
| *44.19        | *4.50          | 0.52           | 96.06          |
| *31.25        | *5.00          | 0.39           | 96.45          |
| *22.10        | *5.50          | 0.17           | 96.62          |
| *15.63        | *6.00          | 0.16           | 96.78          |
| *11.05        | *6.50          | 0.32           | 97.10          |
| *7.81         | *7.00          | 0.49           | 97.58          |
| *5.52         | *7.50          | 0.57           | 98.15          |
| *3.91         | *8.00          | 0.57           | 98.72          |
| *2.76         | *8.50          | 0.49           | 99.21          |
| *1.95         | *9.00          | 0.37           | 99.58          |
| *1.38         | *9.50          | 0.25           | 99.82          |
| *0.98         | *10.00         | 0.16           | 99.98          |
| < 0.98        | > 10.00        | 0.02           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                  |
|-----------------------------|-------|------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand      |
| Mode 2 [µm] <sup>†</sup>    | 1700  | Very coarse sand |
| Mode 3 [µm] <sup>†</sup>    | -     | -                |
| Median [µm] <sup>†</sup>    | 541   | Coarse sand      |
| Median [phi] <sup>†</sup>   | 0.89  |                  |
| Mean [µm] <sup>‡</sup>      | 650   | Coarse sand      |
| Mean [phi] <sup>‡</sup>     | 0.62  | Coarse sand      |
| Sorting [µm] <sup>†</sup>   | 2.46  | Poorly sorted    |
| Sorting [phi] <sup>†</sup>  | 1.30  |                  |
| Skewness [µm] <sup>‡</sup>  | 0.29  | Coarse skewed    |
| Skewness [phi] <sup>‡</sup> | -0.29 |                  |
| Gravel [%] <sup>#</sup>     | 12.98 | Gravelly sand    |
| Sand [%] <sup>#</sup>       | 82.56 |                  |
| Fines [%] <sup>#</sup>      | 4.46  |                  |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST29



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.33           | 0.33           |
| 8000          | -3.00          | 2.70           | 3.03           |
| 5600          | -2.50          | 1.93           | 4.96           |
| 4000          | -2.00          | 4.73           | 9.69           |
| 2800          | -1.50          | 5.01           | 14.71          |
| 2000          | -1.00          | 7.03           | 21.73          |
| 1400          | -0.50          | 5.51           | 27.25          |
| 1000          | 0.00           | 4.40           | 31.65          |
| *707.11       | *0.50          | 16.98          | 48.63          |
| *500.00       | *1.00          | 20.33          | 68.95          |
| *353.55       | *1.50          | 16.84          | 85.79          |
| *250.00       | *2.00          | 8.88           | 94.67          |
| *176.78       | *2.50          | 2.27           | 96.94          |
| *125.00       | *3.00          | 0.04           | 96.98          |
| *88.39        | *3.50          | 0.00           | 96.98          |
| *62.50        | *4.00          | 0.12           | 97.10          |
| *44.19        | *4.50          | 0.27           | 97.36          |
| *31.25        | *5.00          | 0.17           | 97.53          |
| *22.10        | *5.50          | 0.09           | 97.62          |
| *15.63        | *6.00          | 0.16           | 97.78          |
| *11.05        | *6.50          | 0.30           | 98.08          |
| *7.81         | *7.00          | 0.41           | 98.49          |
| *5.52         | *7.50          | 0.44           | 98.93          |
| *3.91         | *8.00          | 0.41           | 99.34          |
| *2.76         | *8.50          | 0.33           | 99.66          |
| *1.95         | *9.00          | 0.23           | 99.89          |
| *1.38         | *9.50          | 0.11           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Median [µm] <sup>†</sup>    | 691   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.53  |                    |
| Mean [µm] <sup>‡</sup>      | 874   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.19  |                    |
| Sorting [µm] <sup>†</sup>   | 2.64  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.40  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.34  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.34 |                    |
| Gravel [%] <sup>#</sup>     | 21.73 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 75.36 |                    |
| Fines [%] <sup>#</sup>      | 2.90  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST30



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.40           | 1.40           |
| 8000          | -3.00          | 3.26           | 4.66           |
| 5600          | -2.50          | 4.36           | 9.03           |
| 4000          | -2.00          | 7.61           | 16.63          |
| 2800          | -1.50          | 6.63           | 23.26          |
| 2000          | -1.00          | 6.08           | 29.34          |
| 1400          | -0.50          | 5.27           | 34.61          |
| 1000          | 0.00           | 4.13           | 38.75          |
| *707.11       | *0.50          | 11.76          | 50.51          |
| *500.00       | *1.00          | 14.78          | 65.29          |
| *353.55       | *1.50          | 13.79          | 79.08          |
| *250.00       | *2.00          | 8.95           | 88.03          |
| *176.78       | *2.50          | 3.52           | 91.55          |
| *125.00       | *3.00          | 0.46           | 92.01          |
| *88.39        | *3.50          | 0.01           | 92.02          |
| *62.50        | *4.00          | 0.41           | 92.42          |
| *44.19        | *4.50          | 0.61           | 93.03          |
| *31.25        | *5.00          | 0.49           | 93.52          |
| *22.10        | *5.50          | 0.38           | 93.90          |
| *15.63        | *6.00          | 0.44           | 94.34          |
| *11.05        | *6.50          | 0.64           | 94.98          |
| *7.81         | *7.00          | 0.83           | 95.81          |
| *5.52         | *7.50          | 0.95           | 96.76          |
| *3.91         | *8.00          | 0.94           | 97.70          |
| *2.76         | *8.50          | 0.79           | 98.50          |
| *1.95         | *9.00          | 0.58           | 99.07          |
| *1.38         | *9.50          | 0.37           | 99.44          |
| *0.98         | *10.00         | 0.25           | 99.69          |
| * < 0.98      | * > 10.00      | 0.31           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                     |
|-----------------------------|-------|---------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand         |
| Mode 2 [µm] <sup>†</sup>    | 4800  | Fine pebble         |
| Mode 3 [µm] <sup>†</sup>    | -     | -                   |
| Median [µm] <sup>†</sup>    | 718   |                     |
| Median [phi] <sup>†</sup>   | 0.48  | Coarse sand         |
| Mean [µm] <sup>‡</sup>      | 952   |                     |
| Mean [phi] <sup>‡</sup>     | 0.07  | Coarse sand         |
| Sorting [µm] <sup>†</sup>   | 5.24  |                     |
| Sorting [phi] <sup>†</sup>  | 2.39  | Very poorly sorted  |
| Skewness [µm] <sup>‡</sup>  | 0.02  |                     |
| Skewness [phi] <sup>‡</sup> | -0.02 | Symmetrical         |
| Gravel [%] <sup>#</sup>     | 29.34 |                     |
| Sand [%] <sup>#</sup>       | 63.08 | Gravelly muddy sand |
| Fines [%] <sup>#</sup>      | 7.58  |                     |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST31



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.17           | 0.17           |
| 8000          | -3.00          | 1.50           | 1.67           |
| 5600          | -2.50          | 1.25           | 2.92           |
| 4000          | -2.00          | 4.36           | 7.28           |
| 2800          | -1.50          | 6.11           | 13.39          |
| 2000          | -1.00          | 5.90           | 19.29          |
| 1400          | -0.50          | 5.19           | 24.47          |
| 1000          | 0.00           | 4.15           | 28.63          |
| *707.11       | *0.50          | 12.05          | 40.67          |
| *500.00       | *1.00          | 19.84          | 60.51          |
| *353.55       | *1.50          | 19.38          | 79.89          |
| *250.00       | *2.00          | 10.51          | 90.39          |
| *176.78       | *2.50          | 2.43           | 92.82          |
| *125.00       | *3.00          | 0.04           | 92.86          |
| *88.39        | *3.50          | 0.00           | 92.86          |
| *62.50        | *4.00          | 0.37           | 93.24          |
| *44.19        | *4.50          | 0.63           | 93.86          |
| *31.25        | *5.00          | 0.42           | 94.28          |
| *22.10        | *5.50          | 0.26           | 94.54          |
| *15.63        | *6.00          | 0.37           | 94.92          |
| *11.05        | *6.50          | 0.63           | 95.55          |
| *7.81         | *7.00          | 0.85           | 96.40          |
| *5.52         | *7.50          | 0.94           | 97.34          |
| *3.91         | *8.00          | 0.87           | 98.21          |
| *2.76         | *8.50          | 0.69           | 98.90          |
| *1.95         | *9.00          | 0.47           | 99.37          |
| *1.38         | *9.50          | 0.29           | 99.65          |
| *0.98         | *10.00         | 0.19           | 99.84          |
| * < 0.98      | * > 10.00      | 0.16           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                |       |                    |
|----------------|-------|--------------------|
| Mode 1 [µm]    | 604   | Coarse sand        |
| Mode 2 [µm]    | 2400  | Granule            |
| Mode 3 [µm]    | -     |                    |
| Median [µm]    | 601   | Coarse sand        |
| Median [phi]   | 0.74  | Coarse sand        |
| Mean [µm]      | 765   | Coarse sand        |
| Mean [phi]     | 0.39  | Coarse sand        |
| Sorting [µm]   | 4.01  | Very poorly sorted |
| Sorting [phi]  | 2.00  | Very poorly sorted |
| Skewness [µm]  | 0.04  | Symmetrical        |
| Skewness [phi] | -0.04 | Symmetrical        |
| Gravel [%]     | 19.29 |                    |
| Sand [%]       | 73.95 | Gravelly sand      |
| Fines [%]      | 6.76  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST32



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.00           | 0.00           |
| 5600          | -2.50          | 1.14           | 1.14           |
| 4000          | -2.00          | 0.65           | 1.79           |
| 2800          | -1.50          | 1.35           | 3.14           |
| 2000          | -1.00          | 2.62           | 5.75           |
| 1400          | -0.50          | 3.98           | 9.74           |
| 1000          | 0.00           | 5.28           | 15.02          |
| *707.11       | *0.50          | 34.67          | 49.69          |
| *500.00       | *1.00          | 29.99          | 79.68          |
| *353.55       | *1.50          | 16.20          | 95.88          |
| *250.00       | *2.00          | 4.07           | 99.95          |
| *176.78       | *2.50          | 0.05           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 854   | Coarse sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>†</sup>    | 705   | Coarse sand            |
| Median [phi] <sup>†</sup>   | 0.51  |                        |
| Mean [µm] <sup>‡</sup>      | 683   | Coarse sand            |
| Mean [phi] <sup>‡</sup>     | 0.55  |                        |
| Sorting [µm] <sup>†</sup>   | 1.60  | Moderately well sorted |
| Sorting [phi] <sup>†</sup>  | 0.68  |                        |
| Skewness [µm] <sup>‡</sup>  | 0.07  | Symmetrical            |
| Skewness [phi] <sup>‡</sup> | -0.07 |                        |
| Gravel [%] <sup>#</sup>     | 5.75  | Gravelly sand          |
| Sand [%] <sup>#</sup>       | 94.25 |                        |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST33



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 1.16           | 1.16           |
| 5600          | -2.50          | 4.37           | 5.53           |
| 4000          | -2.00          | 4.89           | 10.43          |
| 2800          | -1.50          | 4.28           | 14.71          |
| 2000          | -1.00          | 5.07           | 19.78          |
| 1400          | -0.50          | 5.06           | 24.84          |
| 1000          | 0.00           | 3.06           | 27.90          |
| *707.11       | *0.50          | 8.66           | 36.56          |
| *500.00       | *1.00          | 17.89          | 54.45          |
| *353.55       | *1.50          | 21.56          | 76.01          |
| *250.00       | *2.00          | 14.57          | 90.58          |
| *176.78       | *2.50          | 4.68           | 95.26          |
| *125.00       | *3.00          | 0.29           | 95.55          |
| *88.39        | *3.50          | 0.00           | 95.55          |
| *62.50        | *4.00          | 0.14           | 95.69          |
| *44.19        | *4.50          | 0.54           | 96.22          |
| *31.25        | *5.00          | 0.41           | 96.64          |
| *22.10        | *5.50          | 0.17           | 96.81          |
| *15.63        | *6.00          | 0.15           | 96.96          |
| *11.05        | *6.50          | 0.30           | 97.26          |
| *7.81         | *7.00          | 0.47           | 97.74          |
| *5.52         | *7.50          | 0.56           | 98.29          |
| *3.91         | *8.00          | 0.55           | 98.84          |
| *2.76         | *8.50          | 0.46           | 99.30          |
| *1.95         | *9.00          | 0.34           | 99.65          |
| *1.38         | *9.50          | 0.23           | 99.87          |
| *0.98         | *10.00         | 0.13           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION






STATION: MA\_ST34



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.08           | 0.08           |
| 5600          | -2.50          | 1.15           | 1.23           |
| 4000          | -2.00          | 1.14           | 2.37           |
| 2800          | -1.50          | 1.98           | 4.34           |
| 2000          | -1.00          | 3.15           | 7.49           |
| 1400          | -0.50          | 4.02           | 11.51          |
| 1000          | 0.00           | 4.49           | 16.00          |
| *707.11       | *0.50          | 19.63          | 35.63          |
| *500.00       | *1.00          | 24.59          | 60.21          |
| *353.55       | *1.50          | 21.90          | 82.11          |
| *250.00       | *2.00          | 12.61          | 94.72          |
| *176.78       | *2.50          | 3.53           | 98.25          |
| *125.00       | *3.00          | 0.06           | 98.30          |
| *88.39        | *3.50          | 0.00           | 98.30          |
| *62.50        | *4.00          | 0.00           | 98.30          |
| *44.19        | *4.50          | 0.00           | 98.30          |
| *31.25        | *5.00          | 0.00           | 98.30          |
| *22.10        | *5.50          | 0.00           | 98.30          |
| *15.63        | *6.00          | 0.06           | 98.37          |
| *11.05        | *6.50          | 0.27           | 98.63          |
| *7.81         | *7.00          | 0.35           | 98.99          |
| *5.52         | *7.50          | 0.35           | 99.34          |
| *3.91         | *8.00          | 0.30           | 99.64          |
| *2.76         | *8.50          | 0.23           | 99.87          |
| *1.95         | *9.00          | 0.13           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                   |
|-----------------------------|-------|-------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand       |
| Mode 2 [µm] <sup>†</sup>    | -     | -                 |
| Mode 3 [µm] <sup>†</sup>    | -     | -                 |
| Median [µm] <sup>†</sup>    | 577   | Coarse sand       |
| Median [phi] <sup>†</sup>   | 0.79  |                   |
| Mean [µm] <sup>‡</sup>      | 579   | Coarse sand       |
| Mean [phi] <sup>‡</sup>     | 0.79  |                   |
| Sorting [µm] <sup>†</sup>   | 1.88  | Moderately sorted |
| Sorting [phi] <sup>†</sup>  | 0.91  |                   |
| Skewness [µm] <sup>‡</sup>  | 0.14  | Coarse skewed     |
| Skewness [phi] <sup>‡</sup> | -0.14 |                   |
| Gravel [%] <sup>#</sup>     | 7.49  |                   |
| Sand [%] <sup>#</sup>       | 90.81 | Gravelly sand     |
| Fines [%] <sup>#</sup>      | 1.70  |                   |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST35



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.38           | 0.38           |
| 11 200        | -3.50          | 0.00           | 0.38           |
| 8000          | -3.00          | 0.05           | 0.43           |
| 5600          | -2.50          | 0.33           | 0.76           |
| 4000          | -2.00          | 1.21           | 1.97           |
| 2800          | -1.50          | 2.01           | 3.98           |
| 2000          | -1.00          | 3.01           | 6.99           |
| 1400          | -0.50          | 4.63           | 11.62          |
| 1000          | 0.00           | 5.87           | 17.49          |
| *707.11       | *0.50          | 26.44          | 43.93          |
| *500.00       | *1.00          | 28.63          | 72.56          |
| *353.55       | *1.50          | 19.61          | 92.16          |
| *250.00       | *2.00          | 7.21           | 99.37          |
| *176.78       | *2.50          | 0.63           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                   |
|-----------------------------|-------|-------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand       |
| Mode 2 [µm] <sup>†</sup>    | -     | -                 |
| Mode 3 [µm] <sup>†</sup>    | -     | -                 |
| Median [µm] <sup>†</sup>    | 657   | Coarse sand       |
| Median [phi] <sup>†</sup>   | 0.61  |                   |
| Mean [µm] <sup>‡</sup>      | 664   | Coarse sand       |
| Mean [phi] <sup>‡</sup>     | 0.59  |                   |
| Sorting [µm] <sup>†</sup>   | 1.75  | Moderately sorted |
| Sorting [phi] <sup>†</sup>  | 0.81  |                   |
| Skewness [µm] <sup>‡</sup>  | 0.15  | Coarse skewed     |
| Skewness [phi] <sup>‡</sup> | -0.15 |                   |
| Gravel [%] <sup>#</sup>     | 6.99  | Gravelly sand     |
| Sand [%] <sup>#</sup>       | 93.01 |                   |
| Fines [%] <sup>#</sup>      | 0.00  |                   |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST36



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.00           | 0.00           |
| 5600          | -2.50          | 1.92           | 1.92           |
| 4000          | -2.00          | 2.53           | 4.45           |
| 2800          | -1.50          | 3.97           | 8.42           |
| 2000          | -1.00          | 4.96           | 13.38          |
| 1400          | -0.50          | 5.48           | 18.86          |
| 1000          | 0.00           | 4.82           | 23.68          |
| *707.11       | *0.50          | 17.79          | 41.48          |
| *500.00       | *1.00          | 20.69          | 62.17          |
| *353.55       | *1.50          | 17.87          | 80.04          |
| *250.00       | *2.00          | 10.87          | 90.91          |
| *176.78       | *2.50          | 4.06           | 94.96          |
| *125.00       | *3.00          | 0.51           | 95.47          |
| *88.39        | *3.50          | 0.00           | 95.47          |
| *62.50        | *4.00          | 0.35           | 95.82          |
| *44.19        | *4.50          | 0.49           | 96.31          |
| *31.25        | *5.00          | 0.32           | 96.63          |
| *22.10        | *5.50          | 0.19           | 96.83          |
| *15.63        | *6.00          | 0.25           | 97.08          |
| *11.05        | *6.50          | 0.40           | 97.48          |
| *7.81         | *7.00          | 0.53           | 98.01          |
| *5.52         | *7.50          | 0.57           | 98.58          |
| *3.91         | *8.00          | 0.53           | 99.10          |
| *2.76         | *8.50          | 0.41           | 99.52          |
| *1.95         | *9.00          | 0.28           | 99.80          |
| *1.38         | *9.50          | 0.17           | 99.97          |
| *0.98         | *10.00         | 0.03           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                  |
|-----------------------------|-------|------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand      |
| Mode 2 [µm] <sup>†</sup>    | 1700  | Very coarse sand |
| Mode 3 [µm] <sup>†</sup>    | -     | -                |
| Median [µm] <sup>†</sup>    | 613   | Coarse sand      |
| Median [phi] <sup>†</sup>   | 0.71  |                  |
| Mean [µm] <sup>‡</sup>      | 685   | Coarse sand      |
| Mean [phi] <sup>‡</sup>     | 0.54  |                  |
| Sorting [µm] <sup>†</sup>   | 2.44  | Poorly sorted    |
| Sorting [phi] <sup>†</sup>  | 1.29  |                  |
| Skewness [µm] <sup>‡</sup>  | 0.19  | Coarse skewed    |
| Skewness [phi] <sup>‡</sup> | -0.19 |                  |
| Gravel [%] <sup>#</sup>     | 13.38 | Gravelly sand    |
| Sand [%] <sup>#</sup>       | 82.44 |                  |
| Fines [%] <sup>#</sup>      | 4.18  |                  |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST37



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 6.35           | 6.35           |
| 5600          | -2.50          | 3.20           | 9.55           |
| 4000          | -2.00          | 4.89           | 14.44          |
| 2800          | -1.50          | 5.71           | 20.15          |
| 2000          | -1.00          | 6.43           | 26.58          |
| 1400          | -0.50          | 5.37           | 31.95          |
| 1000          | 0.00           | 3.54           | 35.48          |
| *707.11       | *0.50          | 9.95           | 45.44          |
| *500.00       | *1.00          | 15.14          | 60.58          |
| *353.55       | *1.50          | 16.42          | 77.01          |
| *250.00       | *2.00          | 11.53          | 88.53          |
| *176.78       | *2.50          | 4.50           | 93.03          |
| *125.00       | *3.00          | 0.53           | 93.56          |
| *88.39        | *3.50          | 0.00           | 93.56          |
| *62.50        | *4.00          | 0.15           | 93.71          |
| *44.19        | *4.50          | 0.54           | 94.25          |
| *31.25        | *5.00          | 0.49           | 94.74          |
| *22.10        | *5.50          | 0.32           | 95.06          |
| *15.63        | *6.00          | 0.35           | 95.41          |
| *11.05        | *6.50          | 0.53           | 95.94          |
| *7.81         | *7.00          | 0.72           | 96.66          |
| *5.52         | *7.50          | 0.82           | 97.48          |
| *3.91         | *8.00          | 0.79           | 98.27          |
| *2.76         | *8.50          | 0.64           | 98.91          |
| *1.95         | *9.00          | 0.45           | 99.36          |
| *1.38         | *9.50          | 0.29           | 99.65          |
| *0.98         | *10.00         | 0.18           | 99.83          |
| < 0.98        | > 10.00        | 0.17           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Median [µm] <sup>†</sup>    | 637   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.65  |                    |
| Mean [µm] <sup>‡</sup>      | 872   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.20  |                    |
| Sorting [µm] <sup>†</sup>   | 4.61  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.20  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.13  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.13 |                    |
| Gravel [%] <sup>#</sup>     | 26.58 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 67.13 |                    |
| Fines [%] <sup>#</sup>      | 6.29  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST38



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.09           | 1.09           |
| 8000          | -3.00          | 0.89           | 1.98           |
| 5600          | -2.50          | 3.29           | 5.27           |
| 4000          | -2.00          | 4.26           | 9.52           |
| 2800          | -1.50          | 4.15           | 13.67          |
| 2000          | -1.00          | 3.34           | 17.01          |
| 1400          | -0.50          | 1.52           | 18.54          |
| 1000          | 0.00           | 1.20           | 19.74          |
| *707.11       | *0.50          | 13.22          | 32.95          |
| *500.00       | *1.00          | 24.13          | 57.09          |
| *353.55       | *1.50          | 25.35          | 82.43          |
| *250.00       | *2.00          | 14.24          | 96.68          |
| *176.78       | *2.50          | 3.00           | 99.67          |
| *125.00       | *3.00          | 0.01           | 99.68          |
| *88.39        | *3.50          | 0.00           | 99.68          |
| *62.50        | *4.00          | 0.00           | 99.68          |
| *44.19        | *4.50          | 0.00           | 99.68          |
| *31.25        | *5.00          | 0.00           | 99.68          |
| *22.10        | *5.50          | 0.00           | 99.68          |
| *15.63        | *6.00          | 0.00           | 99.68          |
| *11.05        | *6.50          | 0.05           | 99.73          |
| *7.81         | *7.00          | 0.08           | 99.81          |
| *5.52         | *7.50          | 0.08           | 99.88          |
| *3.91         | *8.00          | 0.07           | 99.95          |
| *2.76         | *8.50          | 0.05           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand        |
| Mode 2 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 554   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.85  |                    |
| Mean [µm] <sup>‡</sup>      | 747   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.42  |                    |
| Sorting [µm] <sup>†</sup>   | 2.55  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.35  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.50  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.50 |                    |
| Gravel [%] <sup>#</sup>     | 17.01 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 82.67 |                    |
| Fines [%] <sup>#</sup>      | 0.32  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST39



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.00           | 0.00           |
| 5600          | -2.50          | 0.49           | 0.49           |
| 4000          | -2.00          | 2.36           | 2.85           |
| 2800          | -1.50          | 3.84           | 6.69           |
| 2000          | -1.00          | 7.27           | 13.96          |
| 1400          | -0.50          | 9.14           | 23.10          |
| 1000          | 0.00           | 7.73           | 30.83          |
| *707.11       | *0.50          | 20.67          | 51.49          |
| *500.00       | *1.00          | 24.24          | 75.74          |
| *353.55       | *1.50          | 17.13          | 92.86          |
| *250.00       | *2.00          | 6.48           | 99.35          |
| *176.78       | *2.50          | 0.65           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                  |
|-----------------------------|-------|------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand      |
| Mode 2 [µm] <sup>†</sup>    | 1700  | Very coarse sand |
| Mode 3 [µm] <sup>†</sup>    | -     |                  |
| Median [µm] <sup>†</sup>    | 725   | Coarse sand      |
| Median [phi] <sup>†</sup>   | 0.46  |                  |
| Mean [µm] <sup>‡</sup>      | 827   |                  |
| Mean [phi] <sup>‡</sup>     | 0.27  | Coarse sand      |
| Sorting [µm] <sup>‡</sup>   | 2.06  | Poorly sorted    |
| Sorting [phi] <sup>‡</sup>  | 1.04  |                  |
| Skewness [µm] <sup>‡</sup>  | 0.28  |                  |
| Skewness [phi] <sup>‡</sup> | -0.28 | Coarse skewed    |
| Gravel [%] <sup>#</sup>     | 13.96 |                  |
| Sand [%] <sup>‡</sup>       | 86.04 | Gravelly sand    |
| Fines [%] <sup>‡</sup>      | 0.00  |                  |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST40



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.71           | 1.71           |
| 8000          | -3.00          | 6.15           | 7.87           |
| 5600          | -2.50          | 2.41           | 10.27          |
| 4000          | -2.00          | 4.56           | 14.83          |
| 2800          | -1.50          | 4.76           | 19.59          |
| 2000          | -1.00          | 5.56           | 25.15          |
| 1400          | -0.50          | 4.35           | 29.50          |
| 1000          | 0.00           | 2.98           | 32.48          |
| *707.11       | *0.50          | 10.31          | 42.79          |
| *500.00       | *1.00          | 16.61          | 59.40          |
| *353.55       | *1.50          | 17.81          | 77.21          |
| *250.00       | *2.00          | 12.05          | 89.26          |
| *176.78       | *2.50          | 4.45           | 93.71          |
| *125.00       | *3.00          | 0.45           | 94.16          |
| *88.39        | *3.50          | 0.00           | 94.16          |
| *62.50        | *4.00          | 0.15           | 94.31          |
| *44.19        | *4.50          | 0.54           | 94.85          |
| *31.25        | *5.00          | 0.47           | 95.31          |
| *22.10        | *5.50          | 0.31           | 95.62          |
| *15.63        | *6.00          | 0.35           | 95.97          |
| *11.05        | *6.50          | 0.55           | 96.52          |
| *7.81         | *7.00          | 0.73           | 97.25          |
| *5.52         | *7.50          | 0.79           | 98.04          |
| *3.91         | *8.00          | 0.70           | 98.74          |
| *2.76         | *8.50          | 0.53           | 99.28          |
| *1.95         | *9.00          | 0.35           | 99.62          |
| *1.38         | *9.50          | 0.20           | 99.83          |
| *0.98         | *10.00         | 0.13           | 99.96          |
| * < 0.98      | * > 10.00      | 0.04           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand        |
| Mode 2 [µm] <sup>†</sup>    | 9600  | Medium pebble      |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule            |
| Median [µm] <sup>†</sup>    | 608   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.72  | Coarse sand        |
| Mean [µm] <sup>‡</sup>      | 866   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.21  | Coarse sand        |
| Sorting [µm] <sup>†</sup>   | 4.31  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.11  | Very poorly sorted |
| Skewness [µm] <sup>‡</sup>  | 0.21  | Coarse skewed      |
| Skewness [phi] <sup>‡</sup> | -0.21 | Coarse skewed      |
| Gravel [%] <sup>#</sup>     | 25.15 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 69.16 |                    |
| Fines [%] <sup>#</sup>      | 5.69  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST41



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.52           | 0.52           |
| 5600          | -2.50          | 1.88           | 2.40           |
| 4000          | -2.00          | 2.56           | 4.96           |
| 2800          | -1.50          | 2.06           | 7.02           |
| 2000          | -1.00          | 2.72           | 9.75           |
| 1400          | -0.50          | 4.04           | 13.78          |
| 1000          | 0.00           | 4.71           | 18.49          |
| *707.11       | *0.50          | 20.44          | 38.93          |
| *500.00       | *1.00          | 22.85          | 61.78          |
| *353.55       | *1.50          | 18.21          | 79.99          |
| *250.00       | *2.00          | 9.85           | 89.85          |
| *176.78       | *2.50          | 3.27           | 93.11          |
| *125.00       | *3.00          | 0.59           | 93.71          |
| *88.39        | *3.50          | 0.37           | 94.08          |
| *62.50        | *4.00          | 0.59           | 94.66          |
| *44.19        | *4.50          | 0.49           | 95.16          |
| *31.25        | *5.00          | 0.29           | 95.45          |
| *22.10        | *5.50          | 0.25           | 95.69          |
| *15.63        | *6.00          | 0.37           | 96.07          |
| *11.05        | *6.50          | 0.55           | 96.62          |
| *7.81         | *7.00          | 0.69           | 97.31          |
| *5.52         | *7.50          | 0.74           | 98.06          |
| *3.91         | *8.00          | 0.69           | 98.74          |
| *2.76         | *8.50          | 0.54           | 99.29          |
| *1.95         | *9.00          | 0.37           | 99.66          |
| *1.38         | *9.50          | 0.23           | 99.89          |
| *0.98         | *10.00         | 0.11           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | -     | -             |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 598   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.74  |               |
| Mean [µm] <sup>‡</sup>      | 603   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.73  |               |
| Sorting [µm] <sup>†</sup>   | 2.73  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.45  |               |
| Skewness [µm] <sup>‡</sup>  | -0.06 | Symmetrical   |
| Skewness [phi] <sup>‡</sup> | 0.06  |               |
| Gravel [%] <sup>#</sup>     | 9.75  | Gravelly sand |
| Sand [%] <sup>#</sup>       | 84.92 |               |
| Fines [%] <sup>#</sup>      | 5.34  |               |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST42



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 2.55           | 2.55           |
| 5600          | -2.50          | 2.08           | 4.63           |
| 4000          | -2.00          | 3.49           | 8.12           |
| 2800          | -1.50          | 3.48           | 11.61          |
| 2000          | -1.00          | 5.48           | 17.09          |
| 1400          | -0.50          | 5.54           | 22.62          |
| 1000          | 0.00           | 4.77           | 27.39          |
| *707.11       | *0.50          | 16.99          | 44.38          |
| *500.00       | *1.00          | 19.17          | 63.55          |
| *353.55       | *1.50          | 15.55          | 79.10          |
| *250.00       | *2.00          | 8.77           | 87.87          |
| *176.78       | *2.50          | 3.09           | 90.96          |
| *125.00       | *3.00          | 0.54           | 91.50          |
| *88.39        | *3.50          | 0.27           | 91.77          |
| *62.50        | *4.00          | 0.57           | 92.34          |
| *44.19        | *4.50          | 0.60           | 92.95          |
| *31.25        | *5.00          | 0.44           | 93.39          |
| *22.10        | *5.50          | 0.38           | 93.77          |
| *15.63        | *6.00          | 0.49           | 94.26          |
| *11.05        | *6.50          | 0.70           | 94.96          |
| *7.81         | *7.00          | 0.89           | 95.85          |
| *5.52         | *7.50          | 1.00           | 96.85          |
| *3.91         | *8.00          | 0.97           | 97.81          |
| *2.76         | *8.50          | 0.80           | 98.61          |
| *1.95         | *9.00          | 0.57           | 99.18          |
| *1.38         | *9.50          | 0.36           | 99.54          |
| *0.98         | *10.00         | 0.23           | 99.77          |
| * < 0.98      | * > 10.00      | 0.23           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Median [µm] <sup>†</sup>    | 639   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.65  |                    |
| Mean [µm] <sup>‡</sup>      | 735   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.44  |                    |
| Sorting [µm] <sup>†</sup>   | 4.22  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.08  |                    |
| Skewness [µm] <sup>‡</sup>  | -0.05 | Symmetrical        |
| Skewness [phi] <sup>‡</sup> | 0.05  |                    |
| Gravel [%] <sup>#</sup>     | 17.09 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 75.26 |                    |
| Fines [%] <sup>#</sup>      | 7.66  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST43



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.58           | 0.58           |
| 8000          | -3.00          | 2.87           | 3.45           |
| 5600          | -2.50          | 4.69           | 8.15           |
| 4000          | -2.00          | 5.59           | 13.74          |
| 2800          | -1.50          | 3.98           | 17.72          |
| 2000          | -1.00          | 4.64           | 22.35          |
| 1400          | -0.50          | 4.01           | 26.36          |
| 1000          | 0.00           | 3.63           | 29.99          |
| *707.11       | *0.50          | 17.14          | 47.14          |
| *500.00       | *1.00          | 18.45          | 65.59          |
| *353.55       | *1.50          | 14.49          | 80.08          |
| *250.00       | *2.00          | 8.21           | 88.28          |
| *176.78       | *2.50          | 3.12           | 91.40          |
| *125.00       | *3.00          | 0.69           | 92.09          |
| *88.39        | *3.50          | 0.25           | 92.34          |
| *62.50        | *4.00          | 0.44           | 92.78          |
| *44.19        | *4.50          | 0.49           | 93.27          |
| *31.25        | *5.00          | 0.40           | 93.67          |
| *22.10        | *5.50          | 0.37           | 94.04          |
| *15.63        | *6.00          | 0.47           | 94.51          |
| *11.05        | *6.50          | 0.67           | 95.18          |
| *7.81         | *7.00          | 0.88           | 96.06          |
| *5.52         | *7.50          | 0.99           | 97.05          |
| *3.91         | *8.00          | 0.95           | 98.00          |
| *2.76         | *8.50          | 0.77           | 98.77          |
| *1.95         | *9.00          | 0.53           | 99.30          |
| *1.38         | *9.50          | 0.33           | 99.63          |
| *0.98         | *10.00         | 0.21           | 99.83          |
| < 0.98        | > 10.00        | 0.17           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule            |
| Median [µm] <sup>†</sup>    | 670   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.58  | Coarse sand        |
| Mean [µm] <sup>‡</sup>      | 869   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.20  | Coarse sand        |
| Sorting [µm] <sup>†</sup>   | 4.77  | Very poorly sorted |
| Sorting [phi] <sup>†</sup>  | 2.25  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.03  | Symmetrical        |
| Skewness [phi] <sup>‡</sup> | -0.03 |                    |
| Gravel [%] <sup>#</sup>     | 22.35 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 70.42 |                    |
| Fines [%] <sup>#</sup>      | 7.22  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST44



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.74           | 0.74           |
| 8000          | -3.00          | 0.00           | 0.74           |
| 5600          | -2.50          | 2.96           | 3.70           |
| 4000          | -2.00          | 4.06           | 7.76           |
| 2800          | -1.50          | 4.12           | 11.87          |
| 2000          | -1.00          | 4.53           | 16.40          |
| 1400          | -0.50          | 4.31           | 20.71          |
| 1000          | 0.00           | 3.41           | 24.12          |
| *707.11       | *0.50          | 14.22          | 38.34          |
| *500.00       | *1.00          | 21.28          | 59.62          |
| *353.55       | *1.50          | 20.48          | 80.10          |
| *250.00       | *2.00          | 11.87          | 91.97          |
| *176.78       | *2.50          | 3.27           | 95.24          |
| *125.00       | *3.00          | 0.07           | 95.30          |
| *88.39        | *3.50          | 0.00           | 95.30          |
| *62.50        | *4.00          | 0.16           | 95.46          |
| *44.19        | *4.50          | 0.46           | 95.92          |
| *31.25        | *5.00          | 0.33           | 96.25          |
| *22.10        | *5.50          | 0.17           | 96.42          |
| *15.63        | *6.00          | 0.22           | 96.64          |
| *11.05        | *6.50          | 0.40           | 97.04          |
| *7.81         | *7.00          | 0.57           | 97.61          |
| *5.52         | *7.50          | 0.64           | 98.24          |
| *3.91         | *8.00          | 0.60           | 98.85          |
| *2.76         | *8.50          | 0.49           | 99.34          |
| *1.95         | *9.00          | 0.34           | 99.68          |
| *1.38         | *9.50          | 0.22           | 99.90          |
| *0.98         | *10.00         | 0.10           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 2400  | Granule            |
| Mode 3 [µm] <sup>†</sup>    | 4800  | Fine pebble        |
| Median [µm] <sup>†</sup>    | 585   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.77  |                    |
| Mean [µm] <sup>‡</sup>      | 724   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.47  |                    |
| Sorting [µm] <sup>†</sup>   | 2.64  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.40  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.32  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.32 |                    |
| Gravel [%] <sup>#</sup>     | 16.40 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 79.06 |                    |
| Fines [%] <sup>#</sup>      | 4.54  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST45



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.00           | 1.00           |
| 8000          | -3.00          | 0.83           | 1.82           |
| 5600          | -2.50          | 4.73           | 6.55           |
| 4000          | -2.00          | 4.37           | 10.92          |
| 2800          | -1.50          | 3.33           | 14.26          |
| 2000          | -1.00          | 1.90           | 16.16          |
| 1400          | -0.50          | 1.11           | 17.27          |
| 1000          | 0.00           | 1.48           | 18.75          |
| *707.11       | *0.50          | 15.47          | 34.22          |
| *500.00       | *1.00          | 26.26          | 60.47          |
| *353.55       | *1.50          | 24.69          | 85.16          |
| *250.00       | *2.00          | 11.91          | 97.07          |
| *176.78       | *2.50          | 1.89           | 98.96          |
| *125.00       | *3.00          | 0.00           | 98.96          |
| *88.39        | *3.50          | 0.00           | 98.96          |
| *62.50        | *4.00          | 0.00           | 98.96          |
| *44.19        | *4.50          | 0.00           | 98.96          |
| *31.25        | *5.00          | 0.00           | 98.96          |
| *22.10        | *5.50          | 0.00           | 98.96          |
| *15.63        | *6.00          | 0.00           | 98.96          |
| *11.05        | *6.50          | 0.16           | 99.12          |
| *7.81         | *7.00          | 0.23           | 99.35          |
| *5.52         | *7.50          | 0.23           | 99.57          |
| *3.91         | *8.00          | 0.20           | 99.77          |
| *2.76         | *8.50          | 0.16           | 99.93          |
| *1.95         | *9.00          | 0.07           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| < 0.98        | > 10.00        | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | 6800  | Fine pebble        |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 574   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.80  |                    |
| Mean [µm] <sup>‡</sup>      | 752   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.41  |                    |
| Sorting [µm] <sup>†</sup>   | 2.50  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.32  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.49  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.49 |                    |
| Gravel [%] <sup>#</sup>     | 16.16 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 82.80 |                    |
| Fines [%] <sup>#</sup>      | 1.04  |                    |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST46



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.48           | 0.48           |
| 8000          | -3.00          | 0.23           | 0.71           |
| 5600          | -2.50          | 2.20           | 2.91           |
| 4000          | -2.00          | 2.83           | 5.74           |
| 2800          | -1.50          | 0.95           | 6.68           |
| 2000          | -1.00          | 0.44           | 7.12           |
| 1400          | -0.50          | 0.65           | 7.78           |
| 1000          | 0.00           | 0.63           | 8.41           |
| *707.11       | *0.50          | 7.97           | 16.38          |
| *500.00       | *1.00          | 27.97          | 44.36          |
| *353.55       | *1.50          | 35.60          | 79.96          |
| *250.00       | *2.00          | 17.70          | 97.65          |
| *176.78       | *2.50          | 2.34           | 99.99          |
| *125.00       | *3.00          | 0.01           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| < 0.98        | > 10.00        | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand        |
| Mode 2 [µm] <sup>†</sup>    | -     | -                  |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 473   | Medium sand        |
| Median [phi] <sup>†</sup>   | 1.08  |                    |
| Mean [µm] <sup>‡</sup>      | 481   | Medium sand        |
| Mean [phi] <sup>‡</sup>     | 1.06  |                    |
| Sorting [µm] <sup>†</sup>   | 1.86  | Moderately sorted  |
| Sorting [phi] <sup>†</sup>  | 0.90  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.32  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.32 |                    |
| Gravel [%] <sup>#</sup>     | 7.12  | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 92.88 |                    |
| Fines [%] <sup>#</sup>      | 0.00  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST47



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.56           | 0.56           |
| 5600          | -2.50          | 0.90           | 1.46           |
| 4000          | -2.00          | 2.29           | 3.75           |
| 2800          | -1.50          | 2.43           | 6.18           |
| 2000          | -1.00          | 2.64           | 8.82           |
| 1400          | -0.50          | 3.42           | 12.24          |
| 1000          | 0.00           | 4.37           | 16.61          |
| *707.11       | *0.50          | 17.21          | 33.82          |
| *500.00       | *1.00          | 26.70          | 60.52          |
| *353.55       | *1.50          | 24.50          | 85.02          |
| *250.00       | *2.00          | 12.27          | 97.28          |
| *176.78       | *2.50          | 2.15           | 99.44          |
| *125.00       | *3.00          | 0.00           | 99.44          |
| *88.39        | *3.50          | 0.00           | 99.44          |
| *62.50        | *4.00          | 0.00           | 99.44          |
| *44.19        | *4.50          | 0.00           | 99.44          |
| *31.25        | *5.00          | 0.00           | 99.44          |
| *22.10        | *5.50          | 0.00           | 99.44          |
| *15.63        | *6.00          | 0.00           | 99.44          |
| *11.05        | *6.50          | 0.12           | 99.56          |
| *7.81         | *7.00          | 0.15           | 99.71          |
| *5.52         | *7.50          | 0.14           | 99.85          |
| *3.91         | *8.00          | 0.12           | 99.97          |
| *2.76         | *8.50          | 0.03           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| < 0.98        | > 10.00        | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                   |
|-----------------------------|-------|-------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand       |
| Mode 2 [µm] <sup>†</sup>    | -     | -                 |
| Mode 3 [µm] <sup>†</sup>    | -     | -                 |
| Median [µm] <sup>†</sup>    | 573   | Coarse sand       |
| Median [phi] <sup>†</sup>   | 0.80  |                   |
| Mean [µm] <sup>‡</sup>      | 600   | Coarse sand       |
| Mean [phi] <sup>‡</sup>     | 0.74  |                   |
| Sorting [µm] <sup>†</sup>   | 1.92  | Moderately sorted |
| Sorting [phi] <sup>†</sup>  | 0.94  |                   |
| Skewness [µm] <sup>‡</sup>  | 0.26  | Coarse skewed     |
| Skewness [phi] <sup>‡</sup> | -0.26 |                   |
| Gravel [%] <sup>#</sup>     | 8.82  | Gravelly sand     |
| Sand [%] <sup>#</sup>       | 90.62 |                   |
| Fines [%] <sup>#</sup>      | 0.56  |                   |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST48



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.53           | 0.53           |
| 8000          | -3.00          | 0.29           | 0.82           |
| 5600          | -2.50          | 1.81           | 2.64           |
| 4000          | -2.00          | 2.22           | 4.86           |
| 2800          | -1.50          | 2.29           | 7.15           |
| 2000          | -1.00          | 2.16           | 9.31           |
| 1400          | -0.50          | 1.69           | 11.00          |
| 1000          | 0.00           | 1.76           | 12.76          |
| *707.11       | *0.50          | 13.68          | 26.44          |
| *500.00       | *1.00          | 26.45          | 52.89          |
| *353.55       | *1.50          | 26.27          | 79.16          |
| *250.00       | *2.00          | 13.30          | 92.46          |
| *176.78       | *2.50          | 2.60           | 95.06          |
| *125.00       | *3.00          | 0.02           | 95.08          |
| *88.39        | *3.50          | 0.00           | 95.08          |
| *62.50        | *4.00          | 0.34           | 95.42          |
| *44.19        | *4.50          | 0.61           | 96.03          |
| *31.25        | *5.00          | 0.35           | 96.38          |
| *22.10        | *5.50          | 0.14           | 96.52          |
| *15.63        | *6.00          | 0.21           | 96.73          |
| *11.05        | *6.50          | 0.42           | 97.14          |
| *7.81         | *7.00          | 0.58           | 97.72          |
| *5.52         | *7.50          | 0.63           | 98.35          |
| *3.91         | *8.00          | 0.58           | 98.93          |
| *2.76         | *8.50          | 0.47           | 99.39          |
| *1.95         | *9.00          | 0.33           | 99.73          |
| *1.38         | *9.50          | 0.21           | 99.94          |
| *0.98         | *10.00         | 0.06           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | -     | -             |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 519   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.95  |               |
| Mean [µm] <sup>‡</sup>      | 530   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.92  |               |
| Sorting [µm] <sup>†</sup>   | 2.09  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.07  |               |
| Skewness [µm] <sup>‡</sup>  | 0.18  | Coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.18 |               |
| Gravel [%] <sup>#</sup>     | 9.31  |               |
| Sand [%] <sup>#</sup>       | 86.11 | Gravelly sand |
| Fines [%] <sup>#</sup>      | 4.58  |               |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST49



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.18           | 0.18           |
| 5600          | -2.50          | 3.47           | 3.64           |
| 4000          | -2.00          | 2.25           | 5.89           |
| 2800          | -1.50          | 2.74           | 8.63           |
| 2000          | -1.00          | 3.12           | 11.74          |
| 1400          | -0.50          | 5.16           | 16.90          |
| 1000          | 0.00           | 7.36           | 24.26          |
| *707.11       | *0.50          | 21.45          | 45.71          |
| *500.00       | *1.00          | 23.46          | 69.17          |
| *353.55       | *1.50          | 18.64          | 87.81          |
| *250.00       | *2.00          | 9.36           | 97.17          |
| *176.78       | *2.50          | 1.75           | 98.92          |
| *125.00       | *3.00          | 0.00           | 98.92          |
| *88.39        | *3.50          | 0.00           | 98.92          |
| *62.50        | *4.00          | 0.00           | 98.92          |
| *44.19        | *4.50          | 0.00           | 98.92          |
| *31.25        | *5.00          | 0.00           | 98.92          |
| *22.10        | *5.50          | 0.00           | 98.92          |
| *15.63        | *6.00          | 0.04           | 98.96          |
| *11.05        | *6.50          | 0.21           | 99.17          |
| *7.81         | *7.00          | 0.25           | 99.42          |
| *5.52         | *7.50          | 0.23           | 99.65          |
| *3.91         | *8.00          | 0.20           | 99.85          |
| *2.76         | *8.50          | 0.15           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | -     | -             |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 664   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.59  |               |
| Mean [µm] <sup>‡</sup>      | 721   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.47  |               |
| Sorting [µm] <sup>†</sup>   | 2.16  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.11  |               |
| Skewness [µm] <sup>‡</sup>  | 0.27  | Coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.27 |               |
| Gravel [%] <sup>#</sup>     | 11.74 |               |
| Sand [%] <sup>#</sup>       | 87.18 | Gravelly sand |
| Fines [%] <sup>#</sup>      | 1.08  |               |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST50



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.08           | 0.08           |
| 5600          | -2.50          | 1.10           | 1.18           |
| 4000          | -2.00          | 1.06           | 2.24           |
| 2800          | -1.50          | 1.69           | 3.92           |
| 2000          | -1.00          | 3.45           | 7.37           |
| 1400          | -0.50          | 4.34           | 11.71          |
| 1000          | 0.00           | 3.66           | 15.37          |
| *707.11       | *0.50          | 17.81          | 33.18          |
| *500.00       | *1.00          | 26.81          | 59.99          |
| *353.55       | *1.50          | 24.45          | 84.44          |
| *250.00       | *2.00          | 12.59          | 97.03          |
| *176.78       | *2.50          | 2.52           | 99.55          |
| *125.00       | *3.00          | 0.00           | 99.56          |
| *88.39        | *3.50          | 0.00           | 99.56          |
| *62.50        | *4.00          | 0.00           | 99.56          |
| *44.19        | *4.50          | 0.00           | 99.56          |
| *31.25        | *5.00          | 0.00           | 99.56          |
| *22.10        | *5.50          | 0.00           | 99.56          |
| *15.63        | *6.00          | 0.00           | 99.56          |
| *11.05        | *6.50          | 0.10           | 99.66          |
| *7.81         | *7.00          | 0.13           | 99.79          |
| *5.52         | *7.50          | 0.12           | 99.90          |
| *3.91         | *8.00          | 0.09           | 99.99          |
| *2.76         | *8.50          | 0.01           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                   |
|-----------------------------|-------|-------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand       |
| Mode 2 [µm] <sup>†</sup>    | 1700  | Very coarse sand  |
| Mode 3 [µm] <sup>†</sup>    | -     | -                 |
| Median [µm] <sup>†</sup>    | 569   | Coarse sand       |
| Median [phi] <sup>†</sup>   | 0.81  |                   |
| Mean [µm] <sup>‡</sup>      | 585   | Coarse sand       |
| Mean [phi] <sup>‡</sup>     | 0.77  |                   |
| Sorting [µm] <sup>‡</sup>   | 1.82  | Moderately sorted |
| Sorting [phi] <sup>‡</sup>  | 0.86  |                   |
| Skewness [µm] <sup>‡</sup>  | 0.20  | Coarse skewed     |
| Skewness [phi] <sup>‡</sup> | -0.20 |                   |
| Gravel [%] <sup>#</sup>     | 7.37  |                   |
| Sand [%] <sup>#</sup>       | 92.19 | Gravelly sand     |
| Fines [%] <sup>#</sup>      | 0.44  |                   |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST51



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| > 63 000      | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.43           | 0.43           |
| 8000          | -3.00          | 0.25           | 0.68           |
| 5600          | -2.50          | 1.83           | 2.51           |
| 4000          | -2.00          | 4.22           | 6.73           |
| 2800          | -1.50          | 1.33           | 8.06           |
| 2000          | -1.00          | 0.72           | 8.79           |
| 1400          | -0.50          | 0.78           | 9.56           |
| 1000          | 0.00           | 0.99           | 10.55          |
| *707.11       | *0.50          | 17.40          | 27.95          |
| *500.00       | *1.00          | 31.38          | 59.33          |
| *353.55       | *1.50          | 27.80          | 87.13          |
| *250.00       | *2.00          | 11.38          | 98.51          |
| *176.78       | *2.50          | 1.18           | 99.69          |
| *125.00       | *3.00          | 0.00           | 99.69          |
| *88.39        | *3.50          | 0.00           | 99.69          |
| *62.50        | *4.00          | 0.00           | 99.69          |
| *44.19        | *4.50          | 0.00           | 99.69          |
| *31.25        | *5.00          | 0.00           | 99.69          |
| *22.10        | *5.50          | 0.00           | 99.69          |
| *15.63        | *6.00          | 0.00           | 99.69          |
| *11.05        | *6.50          | 0.06           | 99.74          |
| *7.81         | *7.00          | 0.08           | 99.82          |
| *5.52         | *7.50          | 0.07           | 99.89          |
| *3.91         | *8.00          | 0.06           | 99.96          |
| *2.76         | *8.50          | 0.04           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                   |
|-----------------------------|-------|-------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand       |
| Mode 2 [µm] <sup>†</sup>    | -     | -                 |
| Mode 3 [µm] <sup>†</sup>    | -     | -                 |
| Median [µm] <sup>†</sup>    | 554   | Coarse sand       |
| Median [phi] <sup>†</sup>   | 0.85  | Coarse sand       |
| Mean [µm] <sup>‡</sup>      | 568   | Coarse sand       |
| Mean [phi] <sup>‡</sup>     | 0.82  | Coarse sand       |
| Sorting [µm] <sup>†</sup>   | 1.91  | Moderately sorted |
| Sorting [phi] <sup>†</sup>  | 0.93  | Moderately sorted |
| Skewness [µm] <sup>‡</sup>  | 0.29  | Coarse skewed     |
| Skewness [phi] <sup>‡</sup> | -0.29 | Coarse skewed     |
| Gravel [%] <sup>#</sup>     | 8.79  |                   |
| Sand [%] <sup>†</sup>       | 90.90 | Gravelly sand     |
| Fines [%] <sup>†</sup>      | 0.31  |                   |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST52



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.13           | 0.13           |
| 5600          | -2.50          | 0.18           | 0.31           |
| 4000          | -2.00          | 0.45           | 0.76           |
| 2800          | -1.50          | 0.51           | 1.27           |
| 2000          | -1.00          | 0.68           | 1.96           |
| 1400          | -0.50          | 0.66           | 2.62           |
| 1000          | 0.00           | 0.58           | 3.20           |
| *707.11       | *0.50          | 11.00          | 14.20          |
| *500.00       | *1.00          | 31.84          | 46.04          |
| *353.55       | *1.50          | 36.05          | 82.09          |
| *250.00       | *2.00          | 16.14          | 98.23          |
| *176.78       | *2.50          | 1.77           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 427   | Medium sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>†</sup>    | 481   | Medium sand            |
| Median [phi] <sup>†</sup>   | 1.05  |                        |
| Mean [µm] <sup>‡</sup>      | 484   | Medium sand            |
| Mean [phi] <sup>‡</sup>     | 1.05  |                        |
| Sorting [µm] <sup>‡</sup>   | 1.45  | Moderately well sorted |
| Sorting [phi] <sup>‡</sup>  | 0.53  |                        |
| Skewness [µm] <sup>‡</sup>  | 0.05  | Symmetrical            |
| Skewness [phi] <sup>‡</sup> | -0.05 |                        |
| Gravel [%] <sup>#</sup>     | 1.96  |                        |
| Sand [%] <sup>#</sup>       | 98.04 | Sand                   |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST54



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 2.57           | 2.57           |
| 8000          | -3.00          | 2.19           | 4.76           |
| 5600          | -2.50          | 2.37           | 7.13           |
| 4000          | -2.00          | 2.03           | 9.15           |
| 2800          | -1.50          | 1.56           | 10.71          |
| 2000          | -1.00          | 1.78           | 12.49          |
| 1400          | -0.50          | 2.11           | 14.60          |
| 1000          | 0.00           | 1.78           | 16.38          |
| *707.11       | *0.50          | 20.57          | 36.95          |
| *500.00       | *1.00          | 27.21          | 64.16          |
| *353.55       | *1.50          | 22.36          | 86.51          |
| *250.00       | *2.00          | 10.44          | 96.95          |
| *176.78       | *2.50          | 1.66           | 98.61          |
| *125.00       | *3.00          | 0.00           | 98.61          |
| *88.39        | *3.50          | 0.00           | 98.61          |
| *62.50        | *4.00          | 0.00           | 98.61          |
| *44.19        | *4.50          | 0.00           | 98.61          |
| *31.25        | *5.00          | 0.00           | 98.61          |
| *22.10        | *5.50          | 0.00           | 98.61          |
| *15.63        | *6.00          | 0.05           | 98.66          |
| *11.05        | *6.50          | 0.23           | 98.89          |
| *7.81         | *7.00          | 0.30           | 99.19          |
| *5.52         | *7.50          | 0.29           | 99.48          |
| *3.91         | *8.00          | 0.25           | 99.73          |
| *2.76         | *8.50          | 0.19           | 99.92          |
| *1.95         | *9.00          | 0.08           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                    |
|-----------------------------|-------|--------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand        |
| Mode 2 [µm] <sup>†</sup>    | -     | -                  |
| Mode 3 [µm] <sup>†</sup>    | -     | -                  |
| Median [µm] <sup>†</sup>    | 599   | Coarse sand        |
| Median [phi] <sup>†</sup>   | 0.74  |                    |
| Mean [µm] <sup>‡</sup>      | 618   | Coarse sand        |
| Mean [phi] <sup>‡</sup>     | 0.69  |                    |
| Sorting [µm] <sup>†</sup>   | 2.18  | Poorly sorted      |
| Sorting [phi] <sup>†</sup>  | 1.12  |                    |
| Skewness [µm] <sup>‡</sup>  | 0.30  | Very coarse skewed |
| Skewness [phi] <sup>‡</sup> | -0.30 |                    |
| Gravel [%] <sup>#</sup>     | 12.49 | Gravelly sand      |
| Sand [%] <sup>#</sup>       | 86.11 |                    |
| Fines [%] <sup>#</sup>      | 1.39  |                    |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST55



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.16           | 0.16           |
| 5600          | -2.50          | 0.41           | 0.58           |
| 4000          | -2.00          | 0.27           | 0.84           |
| 2800          | -1.50          | 0.38           | 1.23           |
| 2000          | -1.00          | 0.57           | 1.80           |
| 1400          | -0.50          | 1.39           | 3.19           |
| 1000          | 0.00           | 2.93           | 6.12           |
| *707.11       | *0.50          | 36.23          | 42.35          |
| *500.00       | *1.00          | 36.45          | 78.80          |
| *353.55       | *1.50          | 18.17          | 96.97          |
| *250.00       | *2.00          | 3.02           | 99.99          |
| *176.78       | *2.50          | 0.01           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| < 0.98        | > 10.00        | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>†</sup>    | 658   | Coarse sand            |
| Median [phi] <sup>†</sup>   | 0.60  |                        |
| Mean [µm] <sup>‡</sup>      | 647   | Coarse sand            |
| Mean [phi] <sup>‡</sup>     | 0.63  |                        |
| Sorting [µm] <sup>†</sup>   | 1.41  | Moderately well sorted |
| Sorting [phi] <sup>†</sup>  | 0.50  |                        |
| Skewness [µm] <sup>‡</sup>  | -0.05 | Symmetrical            |
| Skewness [phi] <sup>‡</sup> | 0.05  |                        |
| Gravel [%] <sup>#</sup>     | 1.80  | Sand                   |
| Sand [%] <sup>#</sup>       | 98.20 |                        |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes

Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST56



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 1.41           | 1.41           |
| 8000          | -3.00          | 0.00           | 1.41           |
| 5600          | -2.50          | 0.53           | 1.94           |
| 4000          | -2.00          | 0.68           | 2.62           |
| 2800          | -1.50          | 0.56           | 3.17           |
| 2000          | -1.00          | 0.43           | 3.60           |
| 1400          | -0.50          | 1.21           | 4.81           |
| 1000          | 0.00           | 2.84           | 7.65           |
| *707.11       | *0.50          | 44.50          | 52.16          |
| *500.00       | *1.00          | 34.74          | 86.90          |
| *353.55       | *1.50          | 12.13          | 99.03          |
| *250.00       | *2.00          | 0.97           | 100.00         |
| *176.78       | *2.50          | 0.00           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |             |
|-----------------------------|-------|-------------|
| Mode 1 [µm] <sup>†</sup>    | 854   | Coarse sand |
| Mode 2 [µm] <sup>†</sup>    | -     | -           |
| Mode 3 [µm] <sup>†</sup>    | -     | -           |
| Median [µm] <sup>†</sup>    | 719   | Coarse sand |
| Median [phi] <sup>†</sup>   | 0.48  |             |
| Mean [µm] <sup>‡</sup>      | 703   | Coarse sand |
| Mean [phi] <sup>‡</sup>     | 0.51  |             |
| Sorting [µm] <sup>†</sup>   | 1.40  | Well sorted |
| Sorting [phi] <sup>†</sup>  | 0.49  |             |
| Skewness [µm] <sup>‡</sup>  | -0.04 | Symmetrical |
| Skewness [phi] <sup>‡</sup> | 0.04  |             |
| Gravel [%] <sup>#</sup>     | 3.60  |             |
| Sand [%] <sup>#</sup>       | 96.40 | Sand        |
| Fines [%] <sup>#</sup>      | 0.00  |             |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST57



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.25           | 0.25           |
| 5600          | -2.50          | 0.00           | 0.25           |
| 4000          | -2.00          | 0.23           | 0.48           |
| 2800          | -1.50          | 0.11           | 0.59           |
| 2000          | -1.00          | 0.40           | 0.99           |
| 1400          | -0.50          | 0.67           | 1.66           |
| 1000          | 0.00           | 0.75           | 2.41           |
| *707.11       | *0.50          | 21.31          | 23.72          |
| *500.00       | *1.00          | 39.02          | 62.74          |
| *353.55       | *1.50          | 29.31          | 92.05          |
| *250.00       | *2.00          | 7.75           | 99.80          |
| *176.78       | *2.50          | 0.20           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>†</sup>    | 560   | Coarse sand            |
| Median [phi] <sup>†</sup>   | 0.84  |                        |
| Mean [µm] <sup>‡</sup>      | 559   | Coarse sand            |
| Mean [phi] <sup>‡</sup>     | 0.84  |                        |
| Sorting [µm] <sup>‡</sup>   | 1.42  | Moderately well sorted |
| Sorting [phi] <sup>‡</sup>  | 0.51  |                        |
| Skewness [µm] <sup>‡</sup>  | -0.03 | Symmetrical            |
| Skewness [phi] <sup>‡</sup> | 0.03  |                        |
| Gravel [%] <sup>#</sup>     | 0.99  | Sand                   |
| Sand [%] <sup>#</sup>       | 99.01 |                        |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST58



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.33           | 0.33           |
| 5600          | -2.50          | 0.92           | 1.26           |
| 4000          | -2.00          | 1.97           | 3.23           |
| 2800          | -1.50          | 2.84           | 6.07           |
| 2000          | -1.00          | 4.74           | 10.81          |
| 1400          | -0.50          | 6.52           | 17.32          |
| 1000          | 0.00           | 6.40           | 23.72          |
| *707.11       | *0.50          | 16.71          | 40.43          |
| *500.00       | *1.00          | 21.55          | 61.98          |
| *353.55       | *1.50          | 18.83          | 80.81          |
| *250.00       | *2.00          | 10.38          | 91.19          |
| *176.78       | *2.50          | 2.87           | 94.07          |
| *125.00       | *3.00          | 0.09           | 94.15          |
| *88.39        | *3.50          | 0.00           | 94.15          |
| *62.50        | *4.00          | 0.17           | 94.32          |
| *44.19        | *4.50          | 0.43           | 94.75          |
| *31.25        | *5.00          | 0.33           | 95.08          |
| *22.10        | *5.50          | 0.24           | 95.32          |
| *15.63        | *6.00          | 0.35           | 95.67          |
| *11.05        | *6.50          | 0.58           | 96.25          |
| *7.81         | *7.00          | 0.78           | 97.03          |
| *5.52         | *7.50          | 0.85           | 97.88          |
| *3.91         | *8.00          | 0.77           | 98.65          |
| *2.76         | *8.50          | 0.59           | 99.24          |
| *1.95         | *9.00          | 0.39           | 99.63          |
| *1.38         | *9.50          | 0.24           | 99.87          |
| *0.98         | *10.00         | 0.13           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |               |
|-----------------------------|-------|---------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand   |
| Mode 2 [µm] <sup>†</sup>    | -     | -             |
| Mode 3 [µm] <sup>†</sup>    | -     | -             |
| Median [µm] <sup>†</sup>    | 606   | Coarse sand   |
| Median [phi] <sup>†</sup>   | 0.72  |               |
| Mean [µm] <sup>‡</sup>      | 662   | Coarse sand   |
| Mean [phi] <sup>‡</sup>     | 0.60  |               |
| Sorting [µm] <sup>†</sup>   | 2.94  | Poorly sorted |
| Sorting [phi] <sup>†</sup>  | 1.56  |               |
| Skewness [µm] <sup>‡</sup>  | -0.05 | Symmetrical   |
| Skewness [phi] <sup>‡</sup> | 0.05  |               |
| Gravel [%] <sup>#</sup>     | 10.81 | Gravelly sand |
| Sand [%] <sup>#</sup>       | 83.51 |               |
| Fines [%] <sup>#</sup>      | 5.68  |               |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)








STATION: MA\_ST60



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.31           | 0.31           |
| 5600          | -2.50          | 0.24           | 0.56           |
| 4000          | -2.00          | 0.18           | 0.74           |
| 2800          | -1.50          | 0.15           | 0.89           |
| 2000          | -1.00          | 0.21           | 1.10           |
| 1400          | -0.50          | 0.29           | 1.39           |
| 1000          | 0.00           | 0.43           | 1.82           |
| *707.11       | *0.50          | 15.16          | 16.98          |
| *500.00       | *1.00          | 35.42          | 52.41          |
| *353.55       | *1.50          | 34.05          | 86.46          |
| *250.00       | *2.00          | 12.62          | 99.08          |
| *176.78       | *2.50          | 0.92           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>†</sup>    | 512   | Coarse sand            |
| Median [phi] <sup>†</sup>   | 0.97  |                        |
| Mean [µm] <sup>‡</sup>      | 512   | Coarse sand            |
| Mean [phi] <sup>‡</sup>     | 0.97  |                        |
| Sorting [µm] <sup>‡</sup>   | 1.43  | Moderately well sorted |
| Sorting [phi] <sup>‡</sup>  | 0.51  |                        |
| Skewness [µm] <sup>‡</sup>  | 0.00  | Symmetrical            |
| Skewness [phi] <sup>‡</sup> | 0.00  |                        |
| Gravel [%] <sup>#</sup>     | 1.10  | Sand                   |
| Sand [%] <sup>#</sup>       | 98.90 |                        |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST61



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 3.10           | 3.10           |
| 8000          | -3.00          | 9.15           | 12.25          |
| 5600          | -2.50          | 9.51           | 21.76          |
| 4000          | -2.00          | 6.47           | 28.23          |
| 2800          | -1.50          | 4.69           | 32.92          |
| 2000          | -1.00          | 5.02           | 37.94          |
| 1400          | -0.50          | 2.63           | 40.57          |
| 1000          | 0.00           | 1.36           | 41.93          |
| *707.11       | *0.50          | 15.86          | 57.79          |
| *500.00       | *1.00          | 17.77          | 75.56          |
| *353.55       | *1.50          | 11.84          | 87.40          |
| *250.00       | *2.00          | 4.29           | 91.69          |
| *176.78       | *2.50          | 0.55           | 92.24          |
| *125.00       | *3.00          | 0.03           | 92.27          |
| *88.39        | *3.50          | 0.49           | 92.76          |
| *62.50        | *4.00          | 0.73           | 93.49          |
| *44.19        | *4.50          | 0.53           | 94.02          |
| *31.25        | *5.00          | 0.31           | 94.33          |
| *22.10        | *5.50          | 0.29           | 94.62          |
| *15.63        | *6.00          | 0.44           | 95.06          |
| *11.05        | *6.50          | 0.64           | 95.70          |
| *7.81         | *7.00          | 0.81           | 96.51          |
| *5.52         | *7.50          | 0.89           | 97.40          |
| *3.91         | *8.00          | 0.84           | 98.24          |
| *2.76         | *8.50          | 0.67           | 98.91          |
| *1.95         | *9.00          | 0.46           | 99.37          |
| *1.38         | *9.50          | 0.27           | 99.64          |
| *0.98         | *10.00         | 0.17           | 99.82          |
| * < 0.98      | * > 10.00      | 0.18           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                     |
|-----------------------------|-------|---------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand         |
| Mode 2 [µm] <sup>†</sup>    | 9600  | Medium pebble       |
| Mode 3 [µm] <sup>†</sup>    | 2400  | Granule             |
| Median [µm] <sup>†</sup>    | 838   | Coarse sand         |
| Median [phi] <sup>†</sup>   | 0.25  | Coarse sand         |
| Mean [µm] <sup>†‡</sup>     | 1315  | Very coarse sand    |
| Mean [phi] <sup>†‡</sup>    | -0.40 | Very coarse sand    |
| Sorting [µm] <sup>†</sup>   | 5.47  | Very poorly sorted  |
| Sorting [phi] <sup>†</sup>  | 2.45  | Very poorly sorted  |
| Skewness [µm] <sup>†</sup>  | 0.13  | Coarse skewed       |
| Skewness [phi] <sup>†</sup> | -0.13 | Coarse skewed       |
| Gravel [%] <sup>#</sup>     | 37.94 | Muddy, sandy gravel |
| Sand [%] <sup>#</sup>       | 55.55 |                     |
| Fines [%] <sup>#</sup>      | 6.51  |                     |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)


STATION: MA\_ST63



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.08           | 0.08           |
| 5600          | -2.50          | 0.65           | 0.72           |
| 4000          | -2.00          | 0.58           | 1.31           |
| 2800          | -1.50          | 2.16           | 3.47           |
| 2000          | -1.00          | 4.93           | 8.40           |
| 1400          | -0.50          | 11.46          | 19.86          |
| 1000          | 0.00           | 16.45          | 36.31          |
| *707.11       | *0.50          | 35.53          | 71.84          |
| *500.00       | *1.00          | 21.53          | 93.37          |
| *353.55       | *1.50          | 6.41           | 99.78          |
| *250.00       | *2.00          | 0.22           | 100.00         |
| *176.78       | *2.50          | 0.00           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION




STATION: MA\_ST64



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.11           | 0.11           |
| 5600          | -2.50          | 0.14           | 0.26           |
| 4000          | -2.00          | 0.53           | 0.79           |
| 2800          | -1.50          | 0.95           | 1.74           |
| 2000          | -1.00          | 3.00           | 4.74           |
| 1400          | -0.50          | 7.19           | 11.93          |
| 1000          | 0.00           | 11.53          | 23.46          |
| *707.11       | *0.50          | 35.23          | 58.70          |
| *500.00       | *1.00          | 26.26          | 84.96          |
| *353.55       | *1.50          | 12.39          | 97.34          |
| *250.00       | *2.00          | 2.56           | 99.90          |
| *176.78       | *2.50          | 0.02           | 99.93          |
| *125.00       | *3.00          | 0.00           | 99.93          |
| *88.39        | *3.50          | 0.04           | 99.97          |
| *62.50        | *4.00          | 0.03           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 854   | Coarse sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>‡</sup>    | 770   | Coarse sand            |
| Median [phi] <sup>‡</sup>   | 0.38  |                        |
| Mean [µm] <sup>‡</sup>      | 786   | Coarse sand            |
| Mean [phi] <sup>‡</sup>     | 0.35  |                        |
| Sorting [µm] <sup>‡</sup>   | 1.61  | Moderately well sorted |
| Sorting [phi] <sup>‡</sup>  | 0.69  |                        |
| Skewness [µm] <sup>‡</sup>  | 0.10  | Coarse skewed          |
| Skewness [phi] <sup>‡</sup> | -0.10 |                        |
| Gravel [%] <sup>#</sup>     | 4.74  |                        |
| Sand [%] <sup>#</sup>       | 95.26 | Sand                   |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST65



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.72           | 0.72           |
| 11 200        | -3.50          | 0.00           | 0.72           |
| 8000          | -3.00          | 0.54           | 1.26           |
| 5600          | -2.50          | 0.02           | 1.28           |
| 4000          | -2.00          | 0.18           | 1.46           |
| 2800          | -1.50          | 0.18           | 1.64           |
| 2000          | -1.00          | 0.30           | 1.94           |
| 1400          | -0.50          | 0.28           | 2.22           |
| 1000          | 0.00           | 0.36           | 2.58           |
| *707.11       | *0.50          | 16.88          | 19.46          |
| *500.00       | *1.00          | 37.32          | 56.78          |
| *353.55       | *1.50          | 32.32          | 89.10          |
| *250.00       | *2.00          | 10.39          | 99.48          |
| *176.78       | *2.50          | 0.52           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |                        |
|-----------------------------|-------|------------------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand            |
| Mode 2 [µm] <sup>†</sup>    | -     | -                      |
| Mode 3 [µm] <sup>†</sup>    | -     | -                      |
| Median [µm] <sup>†</sup>    | 533   | Coarse sand            |
| Median [phi] <sup>†</sup>   | 0.91  |                        |
| Mean [µm] <sup>‡</sup>      | 532   | Coarse sand            |
| Mean [phi] <sup>‡</sup>     | 0.91  |                        |
| Sorting [µm] <sup>‡</sup>   | 1.43  | Moderately well sorted |
| Sorting [phi] <sup>‡</sup>  | 0.52  |                        |
| Skewness [µm] <sup>‡</sup>  | -0.01 | Symmetrical            |
| Skewness [phi] <sup>‡</sup> | 0.01  |                        |
| Gravel [%] <sup>#</sup>     | 1.94  |                        |
| Sand [%] <sup>#</sup>       | 98.06 | Sand                   |
| Fines [%] <sup>#</sup>      | 0.00  |                        |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)




STATION: MA\_ST66



FRACTIONAL DATA

| Aperture [µm] | Aperture [phi] | Fractional [%] | Cumulative [%] |
|---------------|----------------|----------------|----------------|
| 63 000        | -6.00          | 0.00           | 0.00           |
| 45 000        | -5.50          | 0.00           | 0.00           |
| 31 500        | -5.00          | 0.00           | 0.00           |
| 22 400        | -4.50          | 0.00           | 0.00           |
| 16 000        | -4.00          | 0.00           | 0.00           |
| 11 200        | -3.50          | 0.00           | 0.00           |
| 8000          | -3.00          | 0.00           | 0.00           |
| 5600          | -2.50          | 0.00           | 0.00           |
| 4000          | -2.00          | 0.01           | 0.01           |
| 2800          | -1.50          | 0.05           | 0.06           |
| 2000          | -1.00          | 0.05           | 0.11           |
| 1400          | -0.50          | 0.15           | 0.26           |
| 1000          | 0.00           | 0.32           | 0.58           |
| *707.11       | *0.50          | 14.92          | 15.50          |
| *500.00       | *1.00          | 41.44          | 56.94          |
| *353.55       | *1.50          | 34.79          | 91.73          |
| *250.00       | *2.00          | 8.10           | 99.83          |
| *176.78       | *2.50          | 0.17           | 100.00         |
| *125.00       | *3.00          | 0.00           | 100.00         |
| *88.39        | *3.50          | 0.00           | 100.00         |
| *62.50        | *4.00          | 0.00           | 100.00         |
| *44.19        | *4.50          | 0.00           | 100.00         |
| *31.25        | *5.00          | 0.00           | 100.00         |
| *22.10        | *5.50          | 0.00           | 100.00         |
| *15.63        | *6.00          | 0.00           | 100.00         |
| *11.05        | *6.50          | 0.00           | 100.00         |
| *7.81         | *7.00          | 0.00           | 100.00         |
| *5.52         | *7.50          | 0.00           | 100.00         |
| *3.91         | *8.00          | 0.00           | 100.00         |
| *2.76         | *8.50          | 0.00           | 100.00         |
| *1.95         | *9.00          | 0.00           | 100.00         |
| *1.38         | *9.50          | 0.00           | 100.00         |
| *0.98         | *10.00         | 0.00           | 100.00         |
| * < 0.98      | * > 10.00      | 0.00           | 100.00         |
| <b>Total</b>  |                | <b>100.00</b>  | <b>-</b>       |

PARTICLE SIZE DISTRIBUTION



SUMMARY STATISTICS

|                             |       |             |
|-----------------------------|-------|-------------|
| Mode 1 [µm] <sup>†</sup>    | 604   | Coarse sand |
| Mode 2 [µm] <sup>†</sup>    | -     | -           |
| Mode 3 [µm] <sup>†</sup>    | -     | -           |
| Median [µm] <sup>†</sup>    | 530   | Coarse sand |
| Median [phi] <sup>†</sup>   | 0.92  |             |
| Mean [µm] <sup>‡</sup>      | 522   | Coarse sand |
| Mean [phi] <sup>‡</sup>     | 0.94  |             |
| Sorting [µm] <sup>†</sup>   | 1.37  | Well sorted |
| Sorting [phi] <sup>†</sup>  | 0.46  |             |
| Skewness [µm] <sup>‡</sup>  | -0.04 | Symmetrical |
| Skewness [phi] <sup>‡</sup> | 0.04  |             |
| Gravel [%] <sup>#</sup>     | 0.11  |             |
| Sand [%] <sup>#</sup>       | 99.89 | Sand        |
| Fines [%] <sup>#</sup>      | 0.00  |             |

Notes  
 Particle Size Distribution by Dry Sieving (63 000 µm - 1000 µm) and Laser Diffraction\* (< 1000 µm - < 0.98 µm) at 0.5 phi Intervals  
 \* = Determinand not included in UKAS Accreditation  
 † = Particle size expressed in accordance with Wentworth (1922) scale  
 ‡ = Statistics calculated using Folk and Ward (1957) method  
 # = Description based on BGS modified Folk classification (Long, 2006)

# Appendix E

## Sediment Hydrocarbon Analysis

## E.1 United States Environmental Protection Agency (US EPA) 16 Polycyclic Aromatic Hydrocarbon (PAH) Concentrations

| PAH<br>[ng/g of Dry Sediment]  | MA_ST04          | MA_ST12    | MA_ST22          | MA_ST25          | MA_ST43     | MA_ST47         | MA_ST59         | MA_ST61     | MA_ST65         | MA_ST66         | CEMP Assessment Criteria<br>(OSPAR, 2014)<br>ERL |
|--|------------------|------------|------------------|------------------|-------------|-----------------|-----------------|-------------|-----------------|-----------------|--|
| Naphthalene  | 0.8              | 6.6        | 0.7              | 0.6              | 1.8         | 0.2             | 0.1             | 1.4         | < 0.1           | 0.1             | 160  |
| Acenaphthylene   | < 0.1            | 0.2        | < 0.1            | < 0.1            | 0.1         | < 0.1           | < 0.1           | 0.1         | < 0.1           | < 0.1           | -  |
| Acenaphthene   | 0.1              | 12.9       | 0.1              | 0.1              | 0.1         | < 0.1           | < 0.1           | 0.2         | < 0.1           | < 0.1           | -  |
| Fluorene   | 0.4              | 7.9        | 0.2              | 0.3              | 0.5         | 0.1             | 0.1             | 0.7         | < 0.1           | < 0.1           | -  |
| Phenanthrene   | 1.6              | 76.0       | 1.9              | 1.7              | 2.7         | 0.4             | 0.6             | 3.2         | 0.1             | 0.2             | 240  |
| Anthracene   | 0.2              | 24.0       | 0.1              | 0.1              | 0.3         | 0.1             | < 0.1           | 0.4         | < 0.1           | < 0.1           | 85   |
| Fluoranthene   | 1.4              | 137        | 1.2              | 1.3              | 2.2         | 0.5             | 0.3             | 3.3         | 0.2             | 0.3             | 600  |
| Pyrene   | 1.1              | 120        | 1.2              | 1.0              | 1.8         | 0.5             | 0.3             | 2.9         | 0.1             | 0.4             | 665  |
| Benzo(a)anthracene   | 0.7              | 61.4       | 0.7              | 0.6              | 1.2         | 0.3             | 0.1             | 1.9         | 0.1             | 0.2             | 261  |
| Chrysene   | 1.0              | 53.9       | 0.9              | 0.9              | 1.6         | 0.4             | 0.2             | 2.2         | 0.1             | 0.2             | 384  |
| Benzo(b)fluoranthene   | 2.3              | 84.3       | 1.6              | 2.1              | 3.5         | 0.8             | 0.5             | 5.3         | 0.3             | 0.6             | -  |
| Benzo(k)fluoranthene   | 0.7              | 33.1       | 0.4              | 0.6              | 1.1         | 0.2             | 0.1             | 1.7         | 0.1             | 0.1             | -  |
| Benzo(a)pyrene   | 0.8              | 77.5       | 0.6              | 0.6              | 1.4         | 0.2             | 0.1             | 2.3         | 0.1             | 0.1             | 430  |
| Indeno(1,2,3-cd)pyrene   | 1.6              | 59.7       | 0.9              | 1.2              | 2.5         | 0.4             | 0.2             | 3.7         | 0.1             | 0.2             | 240  |
| Benzo(ghi)perylene   | 1.4              | 54.5       | 0.9              | 1.1              | 2.3         | 0.5             | 0.2             | 3.6         | 0.1             | 0.3             | 85   |
| Dibenzo(a,h)anthracene   | 0.3              | 11.1       | 0.2              | 0.2              | 0.5         | 0.1             | < 0.1           | 0.8         | < 0.1           | 0.1             | -  |
| <b>Total US EPA 16</b>   | <b>&lt; 14.5</b> | <b>820</b> | <b>&lt; 11.7</b> | <b>&lt; 12.5</b> | <b>23.6</b> | <b>&lt; 4.9</b> | <b>&lt; 3.2</b> | <b>33.7</b> | <b>&lt; 1.9</b> | <b>&lt; 3.2</b> | <b>-</b>   |
| <p>Notes</p> <p>PAH = Polycyclic aromatic hydrocarbon</p> <p>OSPAR = Oslo and Paris Commission</p> <p>CEMP = Coordinated Environmental Monitoring Programme</p> <p>ERL = Effects range low</p> <p>US EPA 16 = United States Environmental Protection Agency's 16 priority polycyclic aromatic hydrocarbons</p> |                  |            |                  |                  |             |                 |                 |             |                 |                 |  |
| <b>Key:</b>  | Below ERL        |            |                  |                  |             |                 | Above ERL       |             |                 |                 |  |

## E.2 Total 2 to 6 Ring PAH Concentrations

| PAH   | Station     |             |             |             |             |                  |                  |             |                 |                  |
|---|-------------|-------------|-------------|-------------|-------------|------------------|------------------|-------------|-----------------|------------------|
|   | MA_ST04     | MA_ST12     | MA_ST22     | MA_ST25     | MA_ST43     | MA_ST47          | MA_ST59          | MA_ST61     | MA_ST65         | MA_ST66          |
| Naphthalene (128)                             | 0.8         | 6.6         | 0.7         | 0.6         | 1.8         | 0.2              | 0.1              | 1.4         | < 0.1           | 0.1              |
| C <sub>1</sub> 128                            | 1.3         | 8.7         | 1.3         | 1.1         | 3.5         | 0.2              | 0.2              | 2.5         | 0.1             | 0.1              |
| C <sub>2</sub> 128                            | 1.9         | 13.1        | 2.4         | 1.6         | 4.5         | 0.4              | 0.7              | 3.7         | 0.1             | 0.2              |
| C <sub>3</sub> 128                            | 2.0         | 15.1        | 3.3         | 1.6         | 4.4         | 0.7              | 1.7              | 4.2         | 0.1             | 0.4              |
| C <sub>4</sub> 128                            | 1.1         | 8.7         | 2.5         | 0.7         | 2.2         | 0.6              | 0.9              | 2.2         | 0.1             | 0.4              |
| <b>TOTAL 128</b>                              | <b>7.1</b>  | <b>52.2</b> | <b>10.2</b> | <b>5.6</b>  | <b>16.4</b> | <b>2.1</b>       | <b>3.6</b>       | <b>14.0</b> | <b>&lt; 0.5</b> | <b>1.2</b>       |
| Phenanthrene/anthracene (178)                 | 1.8         | 100         | 2.0         | 1.8         | 3.0         | 0.5              | 0.6              | 3.6         | 0.1             | 0.2              |
| C <sub>1</sub> 178                            | 1.9         | 29.0        | 2.4         | 1.5         | 3.1         | 0.6              | 1.1              | 3.5         | 0.1             | 0.4              |
| C <sub>2</sub> 178                            | 2.3         | 30.7        | 2.9         | 1.5         | 3.5         | 0.8              | 1.3              | 4.0         | 0.1             | 0.5              |
| C <sub>3</sub> 178                            | 1.7         | 23.7        | 2.4         | 1.0         | 2.5         | 0.7              | 0.8              | 3.0         | 0.1             | 0.5              |
| <b>TOTAL 178</b>                              | <b>7.7</b>  | <b>183</b>  | <b>9.7</b>  | <b>5.8</b>  | <b>12.1</b> | <b>2.6</b>       | <b>3.8</b>       | <b>14.1</b> | <b>0.4</b>      | <b>1.6</b>       |
| Dibenzothiophene (184)                        | 0.1         | 3.4         | 0.1         | 0.1         | 0.2         | < 0.1            | < 0.1            | 0.2         | < 0.1           | < 0.1            |
| C <sub>1</sub> 184                            | 0.2         | 2.6         | 0.2         | 0.2         | 0.3         | 0.1              | 0.1              | 0.4         | < 0.1           | < 0.1            |
| C <sub>2</sub> 184                            | 0.2         | 3.2         | 0.3         | 0.2         | 0.3         | 0.1              | 0.1              | 0.4         | < 0.1           | 0.1              |
| C <sub>3</sub> 184                            | 0.2         | 2.1         | 0.2         | 0.1         | 0.2         | 0.1              | < 0.1            | 0.3         | < 0.1           | 0.1              |
| <b>TOTAL 184</b>                              | <b>0.7</b>  | <b>11.3</b> | <b>0.8</b>  | <b>0.6</b>  | <b>1.0</b>  | <b>&lt; 0.4</b>  | <b>&lt; 0.4</b>  | <b>1.3</b>  | <b>&lt; 0.4</b> | <b>&lt; 0.4</b>  |
| Fluoranthene/pyrene (202)                     | 2.5         | 257         | 2.4         | 2.3         | 4.0         | 1.0              | 0.6              | 6.2         | 0.3             | 0.7              |
| C <sub>1</sub> 202                            | 1.6         | 58.7        | 1.8         | 1.2         | 2.4         | 0.6              | 0.7              | 3.6         | 0.1             | 0.5              |
| C <sub>2</sub> 202                            | 1.5         | 31.3        | 1.6         | 1.0         | 2.2         | 0.5              | 0.7              | 3.2         | 0.1             | 0.4              |
| C <sub>3</sub> 202                            | 1.4         | 27.1        | 1.4         | 0.8         | 2.1         | 0.5              | 0.6              | 2.8         | 0.1             | 0.4              |
| <b>TOTAL 202</b>                              | <b>7.0</b>  | <b>374</b>  | <b>7.2</b>  | <b>5.3</b>  | <b>10.7</b> | <b>2.6</b>       | <b>2.6</b>       | <b>15.8</b> | <b>0.6</b>      | <b>2.0</b>       |
| Benzenanthracenes/<br>benzphenanthrenes (228) | 2.7         | 150         | 2.5         | 2.4         | 4.2         | 1.1              | 0.8              | 6.3         | 0.3             | 0.7              |
| C <sub>1</sub> 228                            | 1.6         | 37.4        | 1.5         | 1.1         | 2.4         | 0.5              | 0.5              | 3.4         | 0.1             | 0.4              |
| C <sub>2</sub> 228                            | 1.9         | 33.9        | 1.8         | 1.2         | 2.7         | 0.6              | 0.7              | 3.8         | 0.2             | 0.5              |
| <b>TOTAL 228</b>                              | <b>6.2</b>  | <b>222</b>  | <b>5.8</b>  | <b>4.7</b>  | <b>9.3</b>  | <b>2.2</b>       | <b>2.0</b>       | <b>13.5</b> | <b>0.6</b>      | <b>1.6</b>       |
| m/z 252*                                      | 6.1         | 287         | 4.3         | 5.5         | 9.6         | 2.4              | 1.4              | 14.7        | 0.7             | 1.6              |
| C <sub>1</sub> 252                            | 2.3         | 54.8        | 1.7         | 1.8         | 3.4         | 0.8              | 0.7              | 5.0         | 0.3             | 0.6              |
| C <sub>2</sub> 252                            | 1.7         | 31.4        | 1.2         | 1.1         | 2.4         | 0.5              | 0.5              | 3.2         | 0.2             | 0.3              |
| <b>TOTAL 252</b>                              | <b>10.1</b> | <b>373</b>  | <b>7.2</b>  | <b>8.4</b>  | <b>15.4</b> | <b>3.7</b>       | <b>2.6</b>       | <b>22.9</b> | <b>1.2</b>      | <b>2.5</b>       |
| m/z 276 <sup>†</sup>                          | 4.5         | 157         | 2.5         | 3.4         | 6.9         | 1.2              | 0.5              | 10.4        | 0.4             | 0.7              |
| C <sub>1</sub> 276                            | 1.0         | 26.9        | 0.6         | 0.6         | 1.5         | 0.3              | 0.2              | 2.6         | 0.1             | 0.2              |
| C <sub>2</sub> 276                            | 0.9         | 18.9        | 0.5         | 0.7         | 1.4         | 0.3              | 0.2              | 1.9         | 0.1             | 0.2              |
| <b>TOTAL 276</b>                              | <b>6.4</b>  | <b>203</b>  | <b>3.6</b>  | <b>4.7</b>  | <b>9.8</b>  | <b>1.8</b>       | <b>0.9</b>       | <b>14.9</b> | <b>0.6</b>      | <b>1.1</b>       |
| <b>NPD<sup>‡</sup></b>                        | <b>15.5</b> | <b>247</b>  | <b>20.7</b> | <b>12.0</b> | <b>29.5</b> | <b>&lt; 5.1</b>  | <b>&lt; 7.8</b>  | <b>29.4</b> | <b>&lt; 1.3</b> | <b>&lt; 3.2</b>  |
| <b>NPD [%]</b>                                | <b>34</b>   | <b>17.0</b> | <b>47</b>   | <b>34</b>   | <b>39</b>   | <b>&lt; 33</b>   | <b>&lt; 50</b>   | <b>30</b>   | <b>&lt; 30</b>  | <b>&lt; 31</b>   |
| <b>Total 2 to 6 ring PAH</b>                  | <b>45.2</b> | <b>1420</b> | <b>44.5</b> | <b>35.1</b> | <b>74.7</b> | <b>&lt; 15.4</b> | <b>&lt; 15.9</b> | <b>96.5</b> | <b>&lt; 4.3</b> | <b>&lt; 10.4</b> |

### Notes

\* = m/z 252 - benzfluoranthenes/benzpyrenes/perylene

† = m/z 276 - anthanthrene/indenoxyrenes/benzperylene

‡ = NPD - naphthalenes, phenanthrenes and dibenzothiophenes (totals)

Concentrations expressed as ng/g dry sediment

# Appendix F

## Macrofaunal Analysis



## F.1 Macrofaunal Abundance

| Taxon                         | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                               |            |      |          |                       | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <b>CNIDARIA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cerianthus lloydii</i>     |            | D632 | 283798   | Gosse, 1859           | 1             | 1             |               | 1             |               |               |               |               |               |               | 1             |               |               |
| Actiniaria                    |            | D662 | 1360     |                       |               |               | 1             | 1             |               |               |               |               |               |               |               |               | 2             |
| <b>PLATYHELMINTHES</b>        |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| PLATYHELMINTHES               |            | F2   | 793      | Minot, 1876           | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>NEMERTEA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| NEMERTEA                      |            | G1   | 152391   |                       | 3             | 5             | 5             | 9             | 5             | 2             | 3             | 6             | 12            | 6             | 5             | 6             | 6             |
| <b>SIPUNCULA</b>              |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Golfingia elongata</i>     |            | N14  | 175026   | (Keferstein, 1862)    | 2             |               |               |               | 2             |               |               | 1             |               | 1             | 1             | 2             |               |
| <i>Golfingia vulgaris</i>     |            | N17  | 136050   | (de Blainville, 1827) |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Nephasoma minutum</i>      |            | N25  | 136060   | (Keferstein, 1862)    | 2             |               | 3             | 6             |               | 5             | 4             |               | 2             |               |               |               |               |
| <i>Phascolion strombus</i>    |            | N34  | 175043   | (Montagu, 1804)       |               |               |               | 1             |               |               | 1             |               | 1             | 1             | 1             |               |               |
| <b>POLYCHAETA</b>             |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pisione remota</i>         |            | P15  | 130707   | Southern, 1914        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aphrodita aculeata</i>     |            | P19  | 129840   | Malmgren, 1867        |               |               |               |               |               |               |               |               |               |               | 1             |               |               |
| <i>Enipo elisabethae</i>      |            | P43  | 130737   | McIntosh, 1900        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gattyana cirrhosa</i>      |            | P49  | 130749   | (Pallas, 1766)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Harmothoe</i>              |            | P50  | 129491   | Kinberg, 1855         | 3             |               | 4             |               | 1             |               | 2             |               |               | 1             | 1             | 1             |               |
| <i>Malmgrenia darbouxi</i>    |            |      | 863197   | (Pettibone, 1993)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Malmgrenia andreapolis</i> |            | P51  | 147008   | (McIntoch, 1874)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lepidonotus squamatus</i>  |            | P82  | 130801   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe inornata</i>        |            | P92  | 130601   | Johnston, 1839        | 2             | 1             |               |               |               |               |               | 1             |               |               |               | 1             |               |
| <i>Pholoe baltica</i>         |            | P95  | 130599   | Örsted, 1843          |               |               |               |               | 2             |               |               |               | 1             |               |               |               |               |
| <i>Sthenelais limicola</i>    |            | P109 | 131077   | (Ehlers, 1864)        |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                       | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|------|----------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |      |          |                                 | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Eteone longa</i>            | aggregate  | P117 | 130616   | (Fabricius, 1780)               |               |               | 1             | 1             | 1             |               |               |               |               |               |               |               | 1             |
| <i>Hesionura elongata</i>      |            | P122 | 130649   | (Southern, 1914)                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mysta picta</i>             |            | P127 | 147026   | (Quatrefages, 1866)             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phyllodoce groenlandica</i> |            | P141 | 334506   | (Ørsted, 1842)                  |               |               |               |               |               | 1             | 1             |               |               |               | 1             | 1             | 1             |
| <i>Phyllodoce longipes</i>     |            | P143 | 130763   | Kinberg, 1866                   |               | 1             |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Phyllodoce rosea</i>        |            | P146 | 334514   | (Mcintosh, 1877)                |               |               | 3             |               | 1             | 1             | 1             | 1             |               | 2             |               |               |               |
| <i>Eulalia bilineata</i>       |            | P152 | 130624   | (Johnston, 1840)                | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eulalia mustela</i>         |            | P155 | 130631   | Pleijel, 1987                   |               | 1             |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Eumida sanguinea</i>        |            | P167 | 130644   | (Ørsted, 1843)                  |               |               | 1             |               |               |               | 1             |               |               |               |               |               |               |
| <i>Glycera alba</i>            |            | P256 | 130116   | (O.F. Muller, 1788)             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera fallax</i>          |            | P259 | 336908   | Quatrefages, 1850               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera lapidum</i>         |            | P260 | 130123   | Quatrefages, 1866               | 3             | 1             |               | 1             | 4             | 2             | 1             | 1             | 2             | 3             |               |               |               |
| <i>Glycera oxycephala</i>      |            | P262 | 130126   | Ehlers, 1887                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycinde nordmanni</i>      |            | P268 | 130136   | (Malmgren, 1865)                |               | 1             |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Goniadella gracilis</i>     |            | P276 | 130145   | (Verrill, 1873)                 | 1             |               |               |               | 2             |               |               | 1             | 1             | 1             |               |               |               |
| <i>Psamathe fusca</i>          |            | P305 | 152249   | (Keferstein, 1862)              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Oxydromus pallidus</i>      |            | P317 | 340203   | (Claparede, 1864)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Podarkeopsis capensis</i>   |            | P319 | 130195   | Day, 1963                       |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Syllis garciai</i>          |            | P351 | 131431   | (Campoy, 1982)                  | 2             | 4             |               | 2             |               | 5             | 2             |               | 1             |               |               | 1             |               |
| <i>Syllis parapari</i>         |            |      | 196002   | San Martin & Lopez, 2000        |               |               | 1             |               |               |               |               |               |               | 1             |               |               |               |
| <i>Syllis pontxioi</i>         |            |      | 196003   | San Martín & López, 2000        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Syllis armillaris</i>       |            | P365 | 131415   | (O.F. Muller, 1776)             | 2             | 4             |               | 1             | 2             | 1             |               | 1             | 2             |               |               |               |               |
| <i>Syllis variegata</i>        |            | P371 | 131458   | (Grube, 1860)                   |               |               |               |               |               | 2             |               |               |               |               |               |               |               |
| <i>Eusyllis blomstrandii</i>   |            | P380 | 131290   | Malmgren, 1867                  |               |               | 2             |               |               |               | 1             |               |               |               |               |               | 1             |
| <i>Odontosyllis fulgurans</i>  |            | P387 | 131327   | (Audouin & Milne Edwards, 1833) |               |               | 1             |               |               |               |               |               |               | 1             |               |               | 2             |

| Taxon                             | Qualifiers | SDC  | APHIA ID | Authority                       | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
|-----------------------------------|------------|------|----------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|---|
|                                   |            |      |          |                                 | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |   |   |
| <i>Streptodonta pterochaeta</i>   |            | P391 | 238207   | Southern, 1914                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Streptosyllis campoyi</i>      |            |      | 238248   | Brito, Nunez & San Martin, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Parexogone hebes</i>           |            | P421 | 757970   | (Webster & Benedict, 1884)      |               |               |               |               |               |               |               |               |               |               |               | 1             |               |   |   |
| <i>Exogone verugera</i>           |            | P423 | 131307   | (Claparede, 1868)               |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Sphaerosyllis</i>              |            | P424 | 129677   | Claparède, 1863                 |               |               |               |               |               |               |               | 1             |               |               |               |               |               |   |   |
| Myrianida                         |            | P434 | 129659   | Milne Edwards, 1845             |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Eunereis longissima</i>        |            | P475 | 130375   | Johnston, 1840                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Nereis zonata</i>              |            | P478 | 130407   | Malmgren, 1867                  |               |               |               | 1             |               |               |               |               |               |               |               |               |               |   |   |
| <i>Nephtys assimilis</i>          |            | P495 | 130353   | Örsted, 1843                    |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |   |
| <i>Nephtys caeca</i>              |            | P496 | 130355   | (Fabricius, 1780)               | 2             |               |               |               |               | 2             | 2             | 2             |               |               |               | 1             |               |   |   |
| <i>Nephtys cirrosa</i>            |            | P498 | 130357   | Ehlers, 1868                    |               |               |               |               |               | 1             |               |               |               |               |               |               |               |   |   |
| <i>Nephtys longosetosa</i>        |            | P503 | 130364   | Örsted, 1843                    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Nothria conchylega</i>         |            | P545 | 130467   | (Sars, 1835)                    | 1             | 1             | 1             |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Paucibranchia bellii</i>       |            | P564 | 130072   | (Audouin & Milne-Edwards, 1833) |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Lysidice unicornis</i>         |            | P568 | 742232   | (Grube, 1840)                   | 2             | 2             | 2             | 1             | 1             | 1             | 3             | 3             | 1             | 1             | 3             |               |               |   |   |
| <i>Lumbrineris cf. cingulata</i>  |            |      | 130240   | (Ehlers, 1868)                  | 1             |               |               |               |               | 1             |               |               |               |               |               | 4             |               |   |   |
| <i>Lumbrineris futilis</i>        |            | P582 | 851788   | (Audouin & Milne Edwards, 1834) |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Drilonereis filum</i>          |            | P591 | 129856   | (Claparède, 1868)               |               |               |               |               |               |               |               |               |               |               |               |               | 1             |   |   |
| <i>Protodorvillea kefersteini</i> |            | P638 | 130041   | (McIntosh, 1869)                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Schistomeringos rudolphi</i>   |            | P643 | 154127   | Delle Chiaje, 1828              |               |               |               |               |               | 1             |               | 1             | 2             | 1             |               |               |               | 1 | 1 |
| <i>Orbinia sertulata</i>          |            | P665 | 334310   | (Savigny, 1820)                 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |   |
| <i>Scoloplos armiger</i>          |            | P672 | 334722   | (Muller, 1776)                  |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |   |
| <i>Aricidea catherinae</i>        |            | P684 | 333034   | (Laubier, 1967)                 |               |               |               |               |               | 1             |               | 1             | 1             |               |               |               |               |   | 1 |

| Taxon                              | Qualifiers | SDC  | APHIA ID | Authority                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|------|----------|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |      |          |                               | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Aricidea cerrutii</i>           |            | P685 | 525497   | Laubier, 1966                 | 2             | 1             | 1             | 1             | 2             | 3             |               | 4             | 2             | 4             | 1             |               |               |
| <i>Cirrophorus branchiatus</i>     |            | P689 | 130576   | Ehlers, 1908                  | 1             |               | 1             | 3             |               |               |               | 4             | 1             |               | 3             | 1             |               |
| <i>Paradoneis ilvana</i>           |            | P698 |          | Castelli, 1985                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Paradoneis lyra</i>             |            | P699 | 130585   | (Southern, 1914)              | 2             | 2             |               | 3             |               |               | 6             | 4             | 2             | 1             |               | 3             | 3             |
| <i>Poecilochaetus serpens</i>      |            | P718 | 130711   | Allen, 1904                   | 2             |               | 3             | 1             | 1             | 3             | 3             | 5             | 1             | 11            | 11            | 12            | 2             |
| <i>Aonides paucibranchiata</i>     |            | P723 | 131107   | Southern, 1914                | 7             | 7             | 18            | 9             | 9             | 10            | 10            | 10            | 12            | 6             | 9             | 5             | 14            |
| <i>Atherospio guillei</i>          |            |      | 478336   | (Laubier & Ramos, 1974)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Laonice bahusiensis</i>         |            | P733 | 131127   | Soderstrom, 1920              | 2             |               | 3             | 3             | 1             |               | 1             | 1             |               | 1             | 1             | 1             |               |
| <i>Dipolydora caulleryi/sp. A</i>  |            | P751 | 131116   | Mesnil, 1897                  | 4             | 1             |               |               | 2             |               |               |               |               |               |               |               | 1             |
| <i>Dipolydora flava</i>            |            | P754 | 131118   | Claparede, 1870               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora sp. B</i>            |            | P762 | 131124   | Verrill, 1881                 |               |               |               | 1             |               |               | 2             |               |               |               |               |               |               |
| <i>Pseudopolydora pulchra</i>      |            | P774 | 131169   | (Carazzi, 1895)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scolecopsis korsuni</i>         |            |      | 131174   | Sikorski, 1994                |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Spio gonocephala</i>            |            |      | 131184   | Thulin, 1957                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Spio symphyta</i>               |            |      | 596189   | Meisner, Bick & Bastrop, 2011 | 1             |               | 2             |               |               |               |               | 1             |               | 1             | 4             |               |               |
| <i>Spiophanes bombyx</i>           |            | P794 | 131187   | (Claparede, 1870)             | 4             | 3             | 4             | 8             | 3             | 7             | 4             | 14            | 3             | 8             | 8             | 3             | 12            |
| <i>Spiophanes kroyeri</i>          |            | P796 | 131188   | Grube, 1860                   |               |               |               | 1             |               |               |               |               |               | 1             |               |               | 1             |
| <i>Magelona johnstoni</i>          |            |      | 130269   | Fiege, Lichen & Mackie, 2000  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Chaetopterus variopedatus</i>   |            | P814 | 129914   | (Renier, 1804)                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phyllochaetopterus anglicus</i> |            | P815 |          | Potts, 1914                   | 2             |               | 2             | 1             | 3             | 3             |               | 1             |               | 3             | 4             |               |               |
| <i>Aphelochaeta sp. A</i>          |            | P823 | 129240   | Blake, 1991                   |               |               |               | 2             |               | 1             |               |               |               |               | 1             |               |               |
| <i>Caulleriella alata</i>          |            | P829 | 129943   | (Southern, 1914)              | 1             |               | 1             |               |               |               |               |               |               |               | 6             | 3             | 2             |
| <i>Chaetozone zetlandica</i>       |            | P831 | 129948   | (McIntosh, 1911)              | 1             |               |               |               | 1             |               |               |               |               | 1             | 3             |               |               |
| <i>Chaetozone christiei</i>        |            |      | 152217   | Chambers, 2000                |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |                          | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Cirratulus cirratus</i>        |            | P836  | 129959   | (O.F.Muller,1776)        |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Tharyx</i>                     |            | P847  | 129249   | Webster & Benedict, 1887 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Flabelligera affinis</i>       |            | P881  | 130103   | M.Sars, 1829             |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Mediomastus fragilis</i>       |            | P919  | 129892   | Rasmussen, 1973          |               |               |               |               |               |               |               | 1             |               |               |               |               | 1             |
| <i>Notomastus</i>                 |            | P920  | 129220   | Sars, 1850               | 1             |               | 3             | 2             | 3             | 1             | 4             | 2             |               | 5             | 3             | 2             | 2             |
| <i>Praxillura longissima</i>      |            | P944  | 130327   | Arwidsson, 1906          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leiochone</i>                  |            |       | 146991   | Grube, 1868              | 4             | 1             | 3             |               | 2             | 1             | 2             | 2             | 2             | 2             | 2             | 1             |               |
| <i>Euclymene lombricoides</i>     |            | P963  | 209899   | (Quatrefages, 1866)      |               |               |               |               |               |               |               |               |               |               | 1             |               |               |
| <i>Euclymene oerstedii</i>        |            | P964  | 130294   | (Claparède, 1863)        |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Praxillella affinis</i>        |            | P971  | 130322   | (M Sars, 1872)           | 2             |               | 6             | 3             | 1             |               | 3             | 4             | 1             |               | 6             | 2             |               |
| <i>Ophelia borealis</i>           |            | P999  | 130491   | Quatrefages, 1866        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelia celtica</i>            |            | P1000 | 130492   | Amoureux & Dauvin, 1981  |               |               |               |               | 2             |               |               |               |               |               |               |               |               |
| <i>Ophelina acuminata</i>         |            | P1014 | 130500   | Ørsted, 1843             |               |               |               |               |               |               |               |               | 1             |               | 1             | 1             |               |
| <i>Asclerocheilus intermedius</i> |            | P1022 | 130974   | (Saint-Joseph, 1894)     | 1             |               | 2             | 2             | 3             |               | 2             | 2             |               |               |               |               |               |
| <i>Scalibregma celticum</i>       |            | P1026 | 130979   | Mackie, 1991             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalibregma inflatum</i>       |            | P1027 | 130980   | Rathke, 1843             | 1             |               |               |               |               |               | 1             |               |               |               | 2             | 1             |               |
| <i>Polygordius</i>                |            | P1062 | 129472   | Schneider, 1868          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Galathowenia oculata</i>       |            | P1093 | 146950   | Zaks, 1922               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Owenia</i>                     |            | P1097 | 129427   | Delle Chiaje, 1844       |               |               | 1             | 1             | 2             | 1             |               |               | 1             | 1             |               | 1             |               |
| <i>Lagis koreni</i>               |            | P1107 | 152367   | Malmgren, 1866           | 3             |               | 2             |               | 1             | 1             |               | 1             | 2             | 1             |               | 1             |               |
| <i>Sabellaria spinulosa</i>       |            | P1117 | 130867   | Leuckart, 1849           |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Ampharete lindstroemi</i>      | aggregate  | P1139 | 129778   | M. Sars, 1864            | 4             | 1             | 3             | 1             | 1             |               | 6             | 2             | 6             | 4             | 4             | 2             | 2             |
| <i>Anobothrus gracilis</i>        |            | P1147 | 129789   | (Malmgren, 1866)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Terebellides stroemii</i>      |            | P1175 | 131573   | Sars, 1835               |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Lanice conchilega</i>          |            | P1195 | 131495   | (Pallas, 1766)           |               |               |               |               | 1             |               |               |               |               | 2             |               | 1             | 1             |



| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |   | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Phisidia aurea</i>             |            | P1215 | 131513   | Southward, 1956                               | 1             |               | 4             | 1             | 1             | 1             | 1             | 1             |               | 2             | 1             | 1             |               |
| <i>Pista mediterranea</i>         |            |       | 131519   | de Gaillande, 1970                            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pista bansei</i>               |            |       | 152254   | Saphronova, 1988                              |               |               |               |               |               | 1             |               | 1             |               | 1             | 2             | 1             | 1             |
| <i>Polycirrus</i>                 |            | P1235 | 129710   | Grube, 1850                                   | 4             | 2             | 4             |               | 4             | 1             | 2             | 3             | 1             | 4             | 3             | 1             | 2             |
| <i>Lysilla loveni</i>             |            | P1233 | 131500   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lysilla nivea</i>              |            | P1234 | 131501   | Langerhans, 1884                              | 2             |               | 1             | 2             | 2             |               | 1             |               |               |               | 1             |               | 1             |
| <i>Streblosoma intestinale</i>    |            | P1252 | 131540   | M. Sars in G.O. Sars, 1872                    | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thelepus cincinnatus</i>       |            | P1254 | 131543   | (Fabricius, 1780)                             |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Dialychone dunerificta</i>     |            |       | 558752   | (Tovar-Hernández, Licciano, Giangrande, 2007) |               | 2             | 1             | 3             |               |               | 1             | 1             | 2             |               | 3             | 1             |               |
| <i>Parasabella cambrensis</i>     |            | P1273 | 530920   | Knight-Jones & Walker, 1985                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Euchone pararosea</i>          |            |       | 390407   | Giangrande & Licciano, 2006                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudopotamilla reniformis</i> |            | P1316 | 130963   | (Bruguiere, 1789)                             |               | 1             |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Sabella pavonina</i>           |            | P1320 | 130967   | Savigny, 1822                                 |               |               |               |               |               | 1             |               |               |               |               | 1             |               |               |
| <i>Hydroides norvegica</i>        |            | P1334 | 131009   | Gunnerus, 1768                                | 3             | 1             | 2             |               |               | 1             |               |               |               |               |               |               |               |
| <i>Spirobranchus triqueter</i>    |            | P1341 | 555935   | (Linnaeus, 1758)                              | 3             | 3             |               | 5             | 2             |               | 5             | 1             |               |               | 2             | 3             | 4             |
| <b>OLIGOCHAETA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Grania</i>                     |            | P1524 | 137349   | Southern, 1913                                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>CHELICERATA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nymphon brevirostre</i>        |            | Q5    | 150520   | Hodge, 1863                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Anoplodactylus petiolatus</i>  |            | Q44   | 134723   | (Kroyer, 1884)                                |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <b>CRUSTACEA</b>                  |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalpellum scalpellum</i>      |            | R22   | 106204   | (Linnaeus, 1767)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Verruca stroemia</i>           |            | R41   | 106257   | O.F.Muller, 1776                              | 1             |               |               | 1             |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |  |
|--------------------------------|------------|------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                |            |      |          |                                   | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |  |
| <i>Nebalia reboreadae</i>      |            |      | 459311   | Moreira & Urgorri, 2009           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nebalia borealis</i>        |            | S7   | 156257   | Dahl, 1985                        |               |               |               |               |               |               |               | 1             |               |               |               |               |               |  |
| <i>Sarsinebalia urgorrhii</i>  |            |      | 388224   | Moreira, Gestoso & Troncoso, 2003 |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Heteromysis formosa</i>     |            | S93  | 120037   | (G. O. Sars, 1877)                |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Kroyera carinata</i>        |            | S125 | 547074   | Spence Bate, 1857                 |               |               | 1             |               |               |               |               |               |               |               |               |               |               |  |
| <i>Periculodes longimanus</i>  |            | S131 | 102915   | (Bate & Westwood, 1868)           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Pontocrates arenarius</i>   |            | S135 | 102918   | (Bate, 1858)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Synchelidium maculatum</i>  |            | S138 | 102928   | Stebbing, 1906                    |               |               |               |               |               |               | 1             |               |               |               |               |               |               |  |
| <i>Apolochus neapolitanus</i>  |            | S159 | 236495   | (Della Valle, 1893)               |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Leucothoe incisa</i>        |            | S177 | 102460   | Robertson, 1892                   |               |               |               |               |               | 2             |               |               |               |               |               |               |               |  |
| <i>Leucothoe procerca</i>      |            | S179 | 102466   | Bate, 1857                        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Stenothoe marina</i>        |            | S213 | 103166   | (Bate, 1856)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Urothoe elegans</i>         |            | S248 | 103228   | (Bate, 1856)                      |               |               | 1             |               |               |               |               |               |               |               |               |               |               |  |
| <i>Urothoe marina</i>          |            | S249 | 103233   | (Bate, 1857)                      | 25            | 30            | 33            | 17            | 26            | 30            | 42            | 33            | 27            | 44            | 44            | 25            | 45            |  |
| <i>Acidostoma neglectum</i>    |            |      | 102495   | (Spence Bate & Westwood, 1861)    |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Hippomedon denticulatus</i> |            | S296 | 102570   | (Bate, 1857)                      |               |               |               |               |               |               |               |               | 1             |               |               |               |               |  |
| <i>Tryphosa nana</i>           |            | S321 | 102691   | (Kroyer, 1846)                    |               |               |               |               |               |               |               |               |               |               | 1             |               |               |  |
| <i>Tmetonyx similis</i>        |            | S337 | 102742   | (G O Sars, 1891)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nototropis falcatus</i>     |            | S410 | 102139   | Metzger, 1871                     |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nototropis vedlomensis</i>  |            | S413 | 102132   | (Bate & Westwood, 1862)           | 1             | 1             | 2             | 1             | 1             |               |               | 1             | 1             | 1             | 1             |               |               |  |
| <i>Ampelisca diadema</i>       |            | S429 | 101896   | (Costa, 1853)                     | 5             | 10            | 12            | 1             | 3             | 1             |               | 3             | 4             | 5             | 12            | 2             | 3             |  |
| <i>Ampelisca provincialis</i>  |            | S434 | 101915   | Bellan-Santini & Kaim-Malka, 1977 | 7             | 2             | 15            | 10            | 9             | 4             | 11            | 13            | 4             | 19            | 21            | 5             |               |  |
| <i>Ampelisca spinipes</i>      |            | S438 | 101928   | Boeck, 1861                       | 1             |               | 2             | 2             | 4             | 1             | 1             | 3             |               | 1             | 3             | 5             |               |  |

| Taxon                              | Qualifiers | SDC         | APHIA ID      | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|-------------|---------------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |             |               |                         | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Ampelisca tenuicornis</i>       |            | S440        | 101930        | Lilljeborg, 1855        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca typica</i>            |            | S442        | 101933        | (Bate, 1856)            | 3             | 4             | 2             | 1             | 2             | 3             | 1             | 1             | 3             | 1             | 2             | 6             | 1             |
| <i>Bathyporeia elegans</i>         |            | S452        | 103058        | Watkin, 1938            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Bathyporeia gracilis</i>        |            | S453        | 103059        | G O Sars, 1891          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Bathyporeia guilliamsoniana</i> |            | S454        | 103060        | (Bate, 1857)            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Haustorius arenarius</i>        |            | S462        | 102317        | Slabber, 1769           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abludomelita obtusata</i>       |            | S498        | 102788        | (Montagu, 1813)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Cheirocratus</b>                |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b>     | 1             |               |               |               |               |               | 1             |               | 1             |               |               |               |               |
| <i>Othomaera othonis</i>           |            | S519        | 534781        | (Milne-Edwards, 1830)   | 1             |               |               | 2             |               |               |               |               |               |               |               |               |               |
| <i>Maerella tenuimana</i>          |            | S521        | 102831        | (Bate, 1862)            | 3             | 1             | 1             |               |               |               | 1             | 1             |               |               |               |               |               |
| <i>Megamphopus cornutus</i>        |            | S539        | 148545        | Norman, 1869            | 1             | 1             | 2             | 2             |               |               |               |               | 1             |               |               |               |               |
| <i>Gammaropsis maculata</i>        |            | S541        | 102364        | (Johnston, 1828)        |               |               | 1             |               |               |               |               |               | 1             | 1             |               |               |               |
| <i>Photis longicaudata</i>         |            | S552        | 102383        | (Bate & Westwood, 1862) | 12            |               |               |               |               |               |               | 21            | 3             |               | 46            |               | 1             |
| <b>Aoridae</b>                     |            | <b>S577</b> | <b>101368</b> | <b>Stebbing, 1899</b>   |               |               | 1             |               |               |               |               |               |               | 3             |               |               |               |
| <i>Leptocheirus hirsutimanus</i>   |            | S588        | 102036        | (Bate, 1862)            |               | 1             |               |               |               | 5             |               |               | 2             |               |               | 1             | 3             |
| <i>Crassikorophium crassicorne</i> |            | S611        | 397383        | Bruzelius, 1859         | 3             | 1             | 1             |               |               |               |               | 1             | 1             |               | 1             | 1             |               |
| <i>Unciola planipes</i>            |            | S622        | 102061        | Norman, 1867            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phtisica marina</i>             |            | S657        | 101864        | Slabber, 1769           |               |               |               |               |               |               |               |               | 2             |               |               |               |               |
| <b>Gnathia</b>                     |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>      | 6             |               | 1             |               |               |               | 5             |               | 1             |               | 1             |               | 3             |
| <i>Conilera cylindracea</i>        |            | S849        | 118842        | (Montagu, 1804)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eurydice pulchra</i>            |            | S854        | 118852        | Leach, 1815             | 3             | 1             | 3             | 1             | 3             | 4             | 4             | 1             |               |               | 2             |               | 5             |
| <i>Astacilla longicornis</i>       |            | S955        | 119024        | (Sowerby, 1806)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tanaopsis graciloides</i>       |            | S1142       | 136458        | (Lilljeborg, 1864)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Bodotria scorpioides</i>        |            | S1197       | 110445        | (Montagu, 1804)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Diastylis bradyi</i>            |            | S1248       | 110472        | Norman, 1879            |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|-------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |       |          |                                   | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Processa modica</i>         |            | S1366 | 108343   | Williamson & Rochanaburanon, 1979 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Callianassa subterranea</i> |            | S1415 | 107729   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Upogebia deltaura</i>       |            | S1419 | 107739   | (Leach, 1815)                     |               |               |               |               |               |               |               |               | 1             |               |               | 1             |               |
| <i>Pisidia longicornis</i>     |            | S1482 | 107188   | (Linnaeus, 1757)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ebalia tumefacta</i>        |            | S1509 | 107302   | (Montagu, 1808)                   |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Hyas coarctatus</i>         |            | S1519 | 107323   | Leach, 1815                       |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Inachus dorsettensis</i>    |            | S1526 | 107327   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia rostrata</i>     |            | S1532 | 107345   | (Linnaeus, 1761)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thia scutellata</i>         |            | S1559 | 107281   | (Fabricius, 1793)                 |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Liocarcinus pusillus</i>    |            | S1584 | 107393   | (Leach, 1816)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pinnotheres pisum</i>       |            | S1638 | 107473   | (Linnaeus, 1767)                  |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <b>MOLLUSCA</b>                |            |       |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptochiton asellus</i>     |            | W53   | 140199   | (Gmelin, 1791)                    | 4             | 6             |               | 2             |               |               | 1             | 1             |               | 1             |               |               |               |
| <i>Acanthochitona crinita</i>  |            | W86   | 138675   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gibbula magus</i>           | ?          | W159  | 141790   | (Linnaeus, 1758)                  |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Euspira nitida</i>          |            | W491  | 151894   | (Donovan, 1804)                   | 1             |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Melanella alba</i>          |            | W634  | 139832   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Colus gracilis</i>          |            | W715  | 138899   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Philine</i>                 |            | W1036 | 138339   | Ascanius, 1772                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tritonia plebeia</i>        |            | W1254 | 141738   | Johnston, 1828                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Onchidorididae                 |            | W1319 | 175      | Gray, 1827                        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Knoutsodonta depressa</i>   |            | W1323 | 845528   | (Alder & Hancock, 1842)           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nucula hanleyi</i>          |            | W1568 | 140588   | Winckworth, 1931                  |               |               |               |               | 1             | 1             |               | 1             |               | 1             | 1             |               |               |
| <i>Modiolus adriaticus</i>     |            | W1700 | 506025   | (Lamarck, 1819)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aequipecten opercularis</i> |            | W1773 | 140687   | (Linnaeus, 1758)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                         | Qualifiers | SDC   | APHIA ID | Authority            | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------|------------|-------|----------|----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                               |            |       |          |                      | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Heteranomia squamula</i>   |            | W1809 | 138749   | (Linnaeus, 1758)     |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Pododesmus squama</i>      |            | W1812 | 138752   | (Linnaeus, 1761)     |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Hemilepton nitidum</i>     |            | W1882 | 246148   | (W. Turton, 1822)    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Kurtiella bidentata</i>    |            | W1906 | 345281   | (Montagu, 1803)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Spisula elliptica</i>      |            | W1975 | 140300   | (Brown, 1827)        | 1             |               |               | 1             | 1             | 1             |               |               | 1             | 1             | 1             |               |               |
| <i>Ensis leei</i>             |            | W1997 | 876640   | M. Huber, 2015       |               |               |               |               | 1             |               |               |               | 1             |               |               |               |               |
| <i>Phaxas pellucidus</i>      |            | W2006 | 140737   | (Pennant, 1777)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Moerella donacina</i>      |            | W2021 | 147021   | Linnaeus, 1758       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Asbjornsenia pygmaea</i>   |            | W2023 | 879714   | (Lovén, 1846)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gari fervensis</i>         |            | W2051 | 140870   | (Gmelin, 1791)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Solecurtus scopula</i>     |            | W2054 | 141543   | (Turton, 1822)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abra alba</i>              |            | W2059 | 141433   | (W Wood, 1802)       |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Abra prismatica</i>        |            | W2062 | 141436   | (Montagu, 1808)      |               | 1             |               |               |               |               |               |               |               | 2             |               |               | 1             |
| <i>Clausinella fasciata</i>   |            | W2100 | 141909   | (da Costa, 1778)     |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Timoclea ovata</i>         |            | W2104 | 141929   | (Pennant, 1777)      | 1             | 1             |               | 1             |               |               |               | 1             |               | 1             |               | 1             |               |
| <i>Venerupis corrugata</i>    |            | W2124 | 181364   | (Gmelin, 1791)       |               |               | 1             |               |               |               |               |               |               |               | 1             | 1             |               |
| <i>Dosinia exoleta</i>        |            | W2130 | 141911   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hiatella arctica</i>       |            | W2166 | 140103   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thracia villosiuscula</i>  |            | W2233 | 141651   | (Macgillivray, 1827) |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Cochlodesma praetenuae</i> |            | W2239 | 181373   | (Pulteney, 1799)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>PHORONIDA</b>              |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phoronis</i>               |            | ZA3   | 128545   | Wright, 1856         |               |               | 14            | 3             |               | 2             |               | 1             |               | 1             | 1             | 13            | 2             |
| <b>ECHINODERMATA</b>          |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophiothrix fragilis</i>    |            | ZB124 | 125131   | (Abildgaard, 1789)   |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Amphipholis squamata</i>   |            | ZB161 | 125064   | (Chiaje, 1828)       |               | 1             |               | 1             | 1             |               |               |               | 1             |               |               |               |               |



| Taxon                                       | Qualifiers | SDC   | APHIA ID | Authority                           | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---|------------|-------|----------|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|   |            |       |          |                                     | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Ophiura albida</i>                       |            | ZB168 | 124913   | Forbes, 1839                        |               |               | 1             |               |               |               |               |               | 1             |               |               |               | 1             |
| <i>Psammechinus miliaris</i>                |            | ZB193 | 124319   | (Gmelin, 1778)                      | 1             |               | 2             | 3             | 1             |               |               |               | 1             |               |               |               |               |
| <i>Echinocyamus pusillus</i>                |            | ZB212 | 124273   | (O.F. Muller, 1776)                 |               |               |               |               | 1             | 4             |               |               | 3             |               | 3             |               | 1             |
| <i>Echinocardium cordatum</i>               |            | ZB223 | 124392   | (Pennant, 1777)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mesothuria intestinalis</i>              |            | ZB233 | 124568   | (Ascanius, 1805)<br>Östergren, 1896 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudothyone raphanus</i>                |            | ZB257 | 124661   | (Duben & Koren, 1845)               |               | 1             |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Thyone fusus</i>                         |            | ZB262 | 124670   | O.F. Muller, 1776                   |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Thyone roscovita</i>                     |            | ZB264 | 124676   | Hérouard, 1889                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Panningia hyndmani</i>                   |            | ZB272 | 848014   | (Thompson, 1840)                    |               |               |               |               | 1             |               |               |               |               |               | 1             |               |               |
| <i>Leptosynapta inhaerens</i>               |            | ZB296 | 124465   | (O.F. Muller, 1776)                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Oestergrenia digitata</i>                |            | ZB300 | 152547   | (Montagu, 1804)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>TUNICATA</b>                             |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidiella aspersa</i>                   |            | ZD84  | 103718   | O F Müller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidia conchilega</i>                   | ?          | ZD88  | 103702   | O F Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Polycarpa fibrosa</i>                    |            | ZD112 | 103902   | (Stimpson, 1852)                    |               |               |               |               |               |               |               |               | 2             |               |               |               |               |
| <i>Dendrodoa grossularia</i>                |            | ZD120 | 103882   | (van Beneden 1846)                  | 4             |               | 1             |               |               |               |               | 2             | 1             | 1             | 1             |               |               |
| <b>CEPHALOCHORDATA</b>                      |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Branchiostoma lanceolatum</i>            |            |       | 104906   | (Pallas, 1774)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                       |            |       |          |                                     | <b>62</b>     | <b>40</b>     | <b>59</b>     | <b>53</b>     | <b>54</b>     | <b>39</b>     | <b>40</b>     | <b>46</b>     | <b>51</b>     | <b>49</b>     | <b>55</b>     | <b>44</b>     | <b>34</b>     |
| <b>Abundance</b>                            |            |       |          |                                     | <b>173</b>    | <b>111</b>    | <b>195</b>    | <b>133</b>    | <b>134</b>    | <b>118</b>    | <b>145</b>    | <b>166</b>    | <b>129</b>    | <b>168</b>    | <b>244</b>    | <b>126</b>    | <b>130</b>    |
| The following taxa were merged for analysis |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aoridae</i>                              |            | S577  | 101368   | Stebbing, 1899                      |               |               | 1             |               |               |               |               |               |               | 3             |               |               |               |
| <i>Aoridae</i>                              | female     | S577  | 101368   | Stebbing, 1899                      |               |               | 1             |               |               |               |               |               |               | 2             |               |               |               |
| <i>Autonoe longipes</i>                     |            | S583  | 102021   | (Lilljeborg, 1852)                  |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Gnathia</i>                              |            | S793  | 118437   | Leach, 1814                         | 6             |               | 1             |               |               |               | 5             |               | 1             |               | 1             |               | 3             |

| Taxon   | Qualifiers | SDC         | APHIA ID      | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |          |
|---|------------|-------------|---------------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------|
|   |            |             |               |                       | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |          |
| <i>Gnathia</i>  | female     | S793        | 118437        | Leach, 1814           | 2             |               |               |               |               |               |               | 2             |               |               |               |               |               | 1        |
| <i>Gnathia oxyuraea</i>                               |            | S796        | 118995        | (Lilljeborg, 1855)    | 4             |               | 1             |               |               |               |               | 3             |               | 1             |               | 1             |               | 2        |
| <b>Cheirocratus</b>                                   |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b>   | <b>1</b>      |               |               |               |               |               |               | <b>1</b>      |               | <b>1</b>      |               |               |               |          |
| <i>Cheirocratus</i>                                   | female     | S503        | 101669        | Norman, 1867          | 1             |               |               |               |               |               |               | 1             |               | 1             |               |               |               |          |
| <i>Cheirocratus pseudosundevallii</i>                 | ?          |             |               | Gouillieux, 2019      |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <b>Number of taxa</b>                                 |            |             |               |                       | <b>2</b>      | <b>0</b>      | <b>2</b>      | <b>0</b>      | <b>0</b>      | <b>0</b>      | <b>0</b>      | <b>2</b>      | <b>0</b>      | <b>2</b>      | <b>1</b>      | <b>1</b>      | <b>0</b>      | <b>1</b> |
| <b>Abundance</b>                                      |            |             |               |                       | <b>7</b>      | <b>0</b>      | <b>2</b>      | <b>0</b>      | <b>0</b>      | <b>0</b>      | <b>0</b>      | <b>6</b>      | <b>0</b>      | <b>2</b>      | <b>3</b>      | <b>1</b>      | <b>0</b>      | <b>3</b> |
| <b>The following taxa were excluded from analysis</b> |            |             |               |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <b>Colonial</b>                                       |            |             |               |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| PORIFERA  |            | C1          | 558           | Grant, 1836           |               |               | P             | P             |               |               |               |               |               |               |               | P             |               |          |
| <i>Leuckartiara octona</i>                            |            | D240        | 117791        | (Fleming, 1823)       |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <i>Tubularia</i>                                      |            | D163        | 117258        | Linnaeus, 1758        |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <i>Eudendrium</i>                                     |            | D218        | 117093        | Ehrenberg, 1834       |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| Bougainvilliidae                                      |            | D246        | 1594          | Lütken, 1850          |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <i>Phialella quadrata</i>                             |            | D343        | 117804        | (Forbes, 1848)        |               |               |               |               |               |               |               |               |               |               | P             |               |               |          |
| <i>Calycella syringa</i>                              |            | D348        | 117402        | (Linnaeus, 1767)      |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <i>Lafoea dumosa</i>                                  |            | D386        | 117702        | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <i>Halecium</i>                                       |            | D390        | 117103        | Oken, 1815            |               |               |               |               |               |               |               |               |               |               |               | P             |               |          |
| <i>Abietinaria</i>                                    |            | D408        | 117225        | Kirchenpauer, 1884    |               |               | P             |               |               | P             | P             |               |               |               | P             |               |               | P        |
| <i>Diphasia</i>                                       |            | D413        | 117228        | L Agassiz, 1862       |               |               | P             |               |               |               |               |               |               | P             |               |               |               |          |
| <i>Hydrallmania falcata</i>                           |            | D424        | 117890        | (Linnaeus, 1758)      | P             | P             | P             | P             | P             |               |               |               | P             | P             | P             | P             |               | P        |
| <i>Sertularella gayi</i>                              |            | D429        | 117902        | (Lamouroux, 1821)     |               |               |               |               |               |               |               |               |               |               |               |               |               |          |
| <i>Sertularia cupressina</i>                          |            | D435        | 117913        | (Linnaeus, 1758)      |               | P             |               |               | P             |               |               |               |               |               | P             | P             |               |          |
| <i>Nemertesia</i>                                     |            | D462        | 117195        | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               | P             |               |               |               |          |
| Campanulariidae                                       |            | D491        | 1606          | Peron & Lesueur, 1810 |               | P             | P             | P             |               | P             |               |               | P             | P             | P             |               |               | P        |

| Taxon                           | Qualifiers | SDC   | APHIA ID | Authority            | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---------------------------------|------------|-------|----------|----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                 |            |       |          |                      | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| <i>Alcyonium digitatum</i>      |            | D597  | 125333   | Linnaeus, 1758       | P             | P             | P             | P             | P             |               | P             |               | P             | P             |               |               |               |
| <i>Crisia</i>                   |            | Y13   | 111032   | Lamouroux, 1812      |               |               | P             |               |               |               |               |               |               |               |               |               |               |
| <i>Tubulipora</i>               |            | Y27   | 111054   | Lamarck, 1816        | P             |               |               |               |               | P             |               |               |               |               |               |               |               |
| <i>Disporella hispida</i>       |            | Y66   | 111730   | (Fleming, 1828)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Alcyonidium parasiticum</i>  |            | Y81   | 111604   | (Fleming, 1828)      |               |               |               | P             |               |               |               |               | P             |               |               |               |               |
| <i>Vesicularia spinosa</i>      |            | Y131  | 111669   | (Linnaeus, 1758)     |               |               |               |               | P             |               |               | P             | P             |               |               |               | P             |
| <i>Amathia lendigera</i>        |            | Y135  | 111659   | (Linnaeus, 1758)     |               |               |               |               |               | P             |               |               |               |               |               |               |               |
| <i>Amathia</i>                  |            | Y137  | 111023   | Farre, 1837          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aetea</i>                    |            | Y153  | 110819   | Lamouroux, 1812      |               |               | P             |               |               |               |               |               |               |               |               |               |               |
| <i>Conopeum reticulum</i>       |            | Y172  | 111351   | (Linnaeus, 1767)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Electra pilosa</i>           |            | Y178  | 111355   | (Linnaeus, 1767)     |               |               | P             |               |               |               |               |               |               | P             |               |               |               |
| <i>Callopora</i>                |            | Y201  | 110851   | J E Gray, 1848       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Beania mirabilis</i>         |            | Y261  | 111072   | Johnston, 1840       |               |               | P             |               |               |               |               |               |               |               |               |               |               |
| <i>Chorizopora brongniartii</i> |            | Y344  | 111304   | (Audouin, 1826)      |               |               |               |               |               |               |               | P             |               |               |               |               |               |
| <i>Escharella immersa</i>       |            | Y364  | 111484   | (Fleming, 1828)      |               |               |               | P             |               |               |               |               |               |               | P             |               |               |
| <i>Porella concinna</i>         |            | Y385  | 111125   | (Busk, 1854)         |               | P             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Schizomavella</i>            |            | Y467  | 110829   | Canu & Bassler, 1917 | P             | P             | P             | P             | P             | P             | P             | P             | P             | P             | P             | P             | P             |
| <b>Damaged</b>                  |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eumida</i>                   | damaged    | P163  | 129446   | Malmgren, 1865       |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Aricidea</i>                 | damaged    | P675  | 129430   | Webster, 1879        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Paradoneis</i>               | damaged    | P695  | 129433   | Hartman, 1965        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Spionidae                       | damaged    | P720  | 913      | Grube, 1850          |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Dipolydora</i>               | damaged    | P748  | 129611   | Verrill, 1881        | 3             |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Terebellidae</i>             | damaged    | P1179 | 982      | Johnston, 1846       | 2             |               | 1             |               |               | 1             |               | 3             |               |               |               |               | 1             |
| Sabellidae                      | damaged    | P1257 | 985      | Latreille, 1825      |               |               |               | 1             |               |               |               |               |               |               |               |               |               |

| Taxon              | Qualifiers | SDC    | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------|------------|--------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                    |            |        |          |                         | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| Serpulidae         | damaged    | P1324  | 988      | Rafinesque, 1815        |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia</i>     | damaged    | S5     | 147031   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| AMPHIPODA          | damaged    | S97    | 1135     |                         |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Urothoe</i>     | damaged    | S246   | 101789   | Dana, 1852              |               | 1             |               |               | 1             | 3             | 1             | 3             | 1             |               |               | 1             |               |
| <i>Liocarcinus</i> | damaged    | S1577  | 106925   | Stimpson, 1870          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| GASTROPODA         | damaged    | W88    | 101      |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| BIVALVIA           | damaged    | W1560  | 151265   | Goldfuss, 1820          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| SOLENOIDEA         | damaged    | W1991  | 14635    | Lamarck, 1809           |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| Ophiuridae         | damaged    | ZB165  | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Fish</b>        |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Gobiesocidae       |            | ZG1180 | 125477   | Bleeker, 1859           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Juvenile</b>    |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| ANIMALIA           | eggs       |        | 2        |                         | 1             |               |               |               |               | 1             |               |               | 1             |               |               |               |               |
| SIPUNCULA          | juvenile   | N1     | 1268     |                         | 1             | 1             |               | 1             | 2             |               | 2             |               |               |               |               |               |               |
| <i>Golfingia</i>   | juvenile   | N12    | 136021   | Lankester, 1885         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera</i>     | juvenile   | P255   | 129296   | Lamarck, 1818           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Goniadidae         | juvenile   | P266   | 953      | Kinberg, 1866           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Nereididae         | juvenile   | P458   | 22496    | Blainville, 1818        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys</i>     | juvenile   | P494   | 129370   | Cuvier, 1817            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scolelepis</i>  | juvenile   | P778   | 129623   | Blainville, 1828        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| CRUSTACEA          | larva      | R1     | 1066     |                         | 1             |               |               |               |               |               |               |               |               |               | 1             |               |               |
| BALANOMORPHA       | juvenile   | R42    | 1082     | Burmeister, 1834        | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca</i>   | juvenile   | S423   | 101445   | Kroyer, 1842            |               | 1             |               |               |               |               |               | 3             |               | 1             | 2             |               |               |
| <i>Gnathia</i>     | juvenile   | S793   | 118437   | Leach, 1814             | 2             | 1             |               |               |               |               |               |               | 1             |               |               |               | 1             |
| GEBIIDEA           | juvenile   | S1403  | 477323   | de Saint Laurent, 1979  | 1             | 2             | 5             | 1             | 4             | 2             |               | 3             | 4             | 2             | 6             | 2             |               |

| Taxon                | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|----------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                      |            |       |          |                         | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| Paguridae            | juvenile   | S1445 | 106738   | Latreille, 1802         |               |               | 1             |               |               |               |               |               |               | 3             |               |               |               |
| DECAPODA             | larva      | S1276 | 1130     | Latreille, 1830         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ebalia</i>        | juvenile   | S1504 | 106889   | Leach, 1817             | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| Majidae              | juvenile   | S1512 | 106760   | Samouelle, 1819         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Inachinae            | juvenile   | S1520 | 148436   | MacLeay, 1838           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia</i>    | juvenile   | S1529 | 205077   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Buccinidae           | eggs       | W702  | 149      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tritonia</i>      | juvenile   | W1246 | 138580   | Cuvier, 1798            | 2             | 1             | 1             | 1             | 1             |               |               | 1             |               | 1             |               |               |               |
| <i>Nucula</i>        | juvenile   | W1565 | 138262   | Lamarck, 1799           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Mytilidae            | juvenile   | W1691 | 211      | Rafinesque, 1815        |               |               |               |               | 1             |               |               |               |               | 1             |               |               |               |
| <i>Modiolus</i>      | juvenile   | W1702 | 140467   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Pectinidae           | juvenile   | W1768 | 213      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Anomiidae            | juvenile   | W1805 | 214      | Rafinesque, 1815        | 1             |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Abra</i>          | juvenile   | W2058 | 138474   | Lamarck, 1818           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dosinia</i>       | juvenile   | W2126 | 138636   | Scopoli, 1777           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Ophiuridae           | juvenile   | ZB165 | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| ECHINOIDEA           | juvenile   | ZB181 | 123082   | Leske, 1778             | 1             |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Echinocardium</i> | juvenile   | ZB222 | 123426   | Gray, 1825              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| HOLOTHUROIDEA        | juvenile   | ZB229 | 123083   |                         |               |               | 2             | 2             |               |               |               |               |               |               |               |               |               |
| <i>Thyone</i>        | juvenile   | ZB261 | 146116   | Oken, 1815              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptosynapta</i>  | juvenile   | ZB291 | 123449   | Verrill, 1867           | 1             |               |               | 1             | 1             |               | 1             |               |               |               |               | 1             |               |
| ASCIDIACEA           | juvenile   | ZD2   | 1839     | Nielsen, 1995           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Meiofaunal</b>    |            |       |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| NEMATODA             |            | HD1   | 799      |                         |               |               |               |               |               |               |               | 1             |               |               |               |               | 2             |
| <b>Parasitic</b>     |            |       |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |



| Taxon                 | Qualifiers | SDC  | APHIA ID | Authority | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------|------------|------|----------|-----------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                       |            |      |          |           | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| COPEPODA              | parasitic  | R142 | 1080     |           |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <b>Number of taxa</b> |            |      |          |           | 17            | 13            | 16            | 14            | 13            | 9             | 6             | 12            | 14            | 14            | 9             | 4             | 9             |
| <b>Abundance</b>      |            |      |          |           | 19            | 8             | 10            | 8             | 12            | 7             | 4             | 15            | 9             | 9             | 9             | 4             | 5             |

| Taxon                         | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                               |            |      |          |                       | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <b>CNIDARIA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cerianthus lloydii</i>     |            | D632 | 283798   | Gosse, 1859           |               |               | 2             |               | 2             |               |               |               |               |               |               |               |               |
| Actiniaria                    |            | D662 | 1360     |                       |               | 1             |               |               |               |               |               | 1             | 1             |               | 1             |               |               |
| <b>PLATYHELMINTHES</b>        |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| PLATYHELMINTHES               |            | F2   | 793      | Minot, 1876           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>NEMERTEA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| NEMERTEA                      |            | G1   | 152391   |                       | 7             | 6             | 6             | 6             | 3             | 5             | 10            | 5             | 11            | 5             | 3             | 2             | 3             |
| <b>SIPUNCULA</b>              |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Golfingia elongata</i>     |            | N14  | 175026   | (Keferstein, 1862)    | 1             |               |               |               | 1             | 5             | 2             |               |               |               |               |               |               |
| <i>Golfingia vulgaris</i>     |            | N17  | 136050   | (de Blainville, 1827) |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephasoma minutum</i>      |            | N25  | 136060   | (Keferstein, 1862)    | 2             |               |               | 1             |               |               | 4             | 1             | 1             |               |               | 2             |               |
| <i>Phascolion strombus</i>    |            | N34  | 175043   | (Montagu, 1804)       |               |               |               | 1             |               |               |               |               |               | 2             |               | 1             |               |
| <b>POLYCHAETA</b>             |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pisione remota</i>         |            | P15  | 130707   | Southern, 1914        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aphrodita aculeata</i>     |            | P19  | 129840   | Malmgren, 1867        |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Enipo elisabethae</i>      |            | P43  | 130737   | McIntosh, 1900        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gattyana cirrhosa</i>      |            | P49  | 130749   | (Pallas, 1766)        |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Harmothoe</i>              |            | P50  | 129491   | Kinberg, 1855         |               |               | 2             |               | 1             | 1             | 1             |               | 1             |               |               |               |               |
| <i>Malmgrenia darbouxi</i>    |            |      | 863197   | (Pettibone, 1993)     | 1             |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Malmgrenia andreapolis</i> |            | P51  | 147008   | (McIntoch, 1874)      |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Lepidonotus squamatus</i>  |            | P82  | 130801   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe inornata</i>        |            | P92  | 130601   | Johnston, 1839        |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Pholoe baltica</i>         |            | P95  | 130599   | Örsted, 1843          | 1             | 1             | 2             | 1             | 1             |               | 1             |               |               |               |               | 1             |               |
| <i>Sthenelais limicola</i>    |            | P109 | 131077   | (Ehlers, 1864)        |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Eteone longa</i>           | aggregate  | P117 | 130616   | (Fabricius, 1780)     | 1             | 1             |               |               |               |               |               | 1             |               |               |               | 1             |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|--------------------------------|------------|------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                |            |      |          |                          | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |   |
| <i>Hesionura elongata</i>      |            | P122 | 130649   | (Southern, 1914)         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Mysta picta</i>             |            | P127 | 147026   | (Quatrefages, 1866)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Phyllodoce groenlandica</i> |            | P141 | 334506   | (Örsted, 1842)           |               |               | 1             | 1             | 1             | 1             |               |               |               |               |               |               | 1             |   |
| <i>Phyllodoce longipes</i>     |            | P143 | 130763   | Kinberg, 1866            |               |               |               |               |               |               |               |               |               | 1             |               |               |               |   |
| <i>Phyllodoce rosea</i>        |            | P146 | 334514   | (Mcintosh, 1877)         |               | 1             | 1             |               |               | 2             |               |               |               |               |               |               |               |   |
| <i>Eulalia bilineata</i>       |            | P152 | 130624   | (Johnston, 1840)         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Eulalia mustela</i>         |            | P155 | 130631   | Pleijel, 1987            |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Eumida sanguinea</i>        |            | P167 | 130644   | (Örsted, 1843)           |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Glycera alba</i>            |            | P256 | 130116   | (O.F. Muller, 1788)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Glycera fallax</i>          |            | P259 | 336908   | Quatrefages, 1850        | 1             | 1             |               |               |               |               |               |               |               | 1             |               |               |               |   |
| <i>Glycera lapidum</i>         |            | P260 | 130123   | Quatrefages, 1866        |               |               |               | 1             |               | 1             | 3             |               |               | 3             |               |               | 1             |   |
| <i>Glycera oxycephala</i>      |            | P262 | 130126   | Ehlers, 1887             |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Glycinde nordmanni</i>      |            | P268 | 130136   | (Malmgren, 1865)         |               |               |               |               |               |               |               |               |               |               |               |               | 1             |   |
| <i>Goniadella gracilis</i>     |            | P276 | 130145   | (Verrill, 1873)          | 3             |               |               |               |               | 2             |               | 1             | 1             |               |               |               |               |   |
| <i>Psamathe fusca</i>          |            | P305 | 152249   | (Keferstein, 1862)       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Oxydromus pallidus</i>      |            | P317 | 340203   | (Claparede, 1864)        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Podarkeopsis capensis</i>   |            | P319 | 130195   | Day, 1963                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Syllis garciai</i>          |            | P351 | 131431   | (Campoy, 1982)           | 1             | 1             |               |               | 1             |               | 1             |               | 3             |               |               |               |               | 1 |
| <i>Syllis parapari</i>         |            |      | 196002   | San Martin & Lopez, 2000 |               |               |               | 2             |               |               | 1             |               |               |               |               |               |               |   |
| <i>Syllis pontxioi</i>         |            |      | 196003   | San Martín & López, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Syllis armillaris</i>       |            | P365 | 131415   | (O.F. Muller, 1776)      | 1             | 1             |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Syllis variegata</i>        |            | P371 | 131458   | (Grube, 1860)            |               |               |               |               |               |               |               |               |               |               |               |               | 1             |   |
| <i>Eusyllis blomstrandii</i>   |            | P380 | 131290   | Malmgren, 1867           | 1             |               | 1             |               |               |               |               |               |               |               |               |               |               |   |

| Taxon                             | Qualifiers | SDC  | APHIA ID | Authority                       | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|-----------------------------------|------------|------|----------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                   |            |      |          |                                 | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |   |
| <i>Odontosyllis fulgurans</i>     |            | P387 | 131327   | (Audouin & Milne Edwards, 1833) | 1             | 1             |               |               |               |               |               | 1             |               |               |               |               |               |   |
| <i>Streptodonta pterochaeta</i>   |            | P391 | 238207   | Southern, 1914                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Streptosyllis campoyi</i>      |            |      | 238248   | Brito, Nunez & San Martin, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Parexogone hebes</i>           |            | P421 | 757970   | (Webster & Benedict, 1884)      |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <i>Exogone verugera</i>           |            | P423 | 131307   | (Claparede, 1868)               |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Sphaerosyllis</i>              |            | P424 | 129677   | Claparède, 1863                 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| Myrianida                         |            | P434 | 129659   | Milne Edwards, 1845             |               |               |               |               |               |               |               | 2             |               |               |               |               |               |   |
| <i>Eunereis longissima</i>        |            | P475 | 130375   | Johnston, 1840                  |               |               |               | 1             |               |               |               |               |               |               |               |               |               |   |
| <i>Nereis zonata</i>              |            | P478 | 130407   | Malmgren, 1867                  |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Nephtys assimilis</i>          |            | P495 | 130353   | Ørsted, 1843                    |               |               |               |               |               |               |               |               |               | 1             | 1             |               |               |   |
| <i>Nephtys caeca</i>              |            | P496 | 130355   | (Fabricius, 1780)               |               |               |               |               |               | 2             |               |               |               |               |               |               |               |   |
| <i>Nephtys cirrosa</i>            |            | P498 | 130357   | Ehlers, 1868                    |               |               |               |               |               |               |               | 1             | 1             |               |               |               |               |   |
| <i>Nephtys longosetosa</i>        |            | P503 | 130364   | Ørsted, 1843                    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Nothria conchylega</i>         |            | P545 | 130467   | (Sars, 1835)                    |               |               |               | 1             |               |               |               |               |               | 1             |               |               |               |   |
| <i>Paucibranchia bellii</i>       |            | P564 | 130072   | (Audouin & Milne-Edwards, 1833) |               |               |               |               |               |               |               | 1             |               |               |               |               |               |   |
| <i>Lysidice unicornis</i>         |            | P568 | 742232   | (Grube, 1840)                   | 2             |               | 1             |               | 3             | 3             | 3             | 1             |               | 2             |               |               | 2             |   |
| <i>Lumbrineris cf. cingulata</i>  |            |      | 130240   | (Ehlers, 1868)                  |               | 1             |               |               | 1             |               | 1             |               |               | 3             |               |               |               |   |
| <i>Lumbrineris futilis</i>        |            | P582 | 851788   | (Audouin & Milne Edwards, 1834) |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Drilonereis filum</i>          |            | P591 | 129856   | (Claparède, 1868)               |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Protodorvillea kefersteini</i> |            | P638 | 130041   | (McIntosh, 1869)                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Schistomeringos rudolphi</i>   |            | P643 | 154127   | Delle Chiaje, 1828              | 1             | 1             | 1             | 1             |               |               |               |               |               |               |               |               |               |   |
| <i>Orbinia sertulata</i>          |            | P665 | 334310   | (Savigny, 1820)                 |               | 1             |               |               |               |               |               |               |               |               |               |               |               |   |

| Taxon                              | Qualifiers | SDC  | APHIA ID | Authority                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|------|----------|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |      |          |                               | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Scoloplos armiger</i>           |            | P672 | 334722   | (Muller, 1776)                |               |               |               |               | 1             |               |               |               |               | 1             | 4             |               | 2             |
| <i>Aricidea catherinae</i>         |            | P684 | 333034   | (Laubier, 1967)               |               |               |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Aricidea cerrutii</i>           |            | P685 | 525497   | Laubier, 1966                 | 3             |               | 1             |               |               |               | 2             | 2             | 1             | 1             | 1             | 1             |               |
| <i>Cirrophorus branchiatus</i>     |            | P689 | 130576   | Ehlers, 1908                  |               | 1             | 1             |               |               |               |               | 1             |               |               |               |               |               |
| <i>Paradoneis ilvana</i>           |            | P698 |          | Castelli, 1985                |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Paradoneis lyra</i>             |            | P699 | 130585   | (Southern, 1914)              | 3             | 1             | 4             | 6             | 1             | 2             | 7             | 1             | 2             | 1             | 1             | 1             | 1             |
| <i>Poecilochaetus serpens</i>      |            | P718 | 130711   | Allen, 1904                   | 3             | 3             | 5             | 2             | 14            | 2             | 3             | 1             | 1             | 7             | 2             | 8             | 7             |
| <i>Aonides paucibranchiata</i>     |            | P723 | 131107   | Southern, 1914                | 21            | 4             | 9             | 7             | 7             | 6             | 7             | 4             | 13            | 8             | 3             | 1             | 1             |
| <i>Atherospio guillei</i>          |            |      | 478336   | (Laubier & Ramos,1974)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Laonice bahusiensis</i>         |            | P733 | 131127   | Soderstrom, 1920              | 4             |               |               |               |               |               | 2             |               | 1             |               |               |               |               |
| <i>Dipolydora caulleryi/sp. A</i>  |            | P751 | 131116   | Mesnil, 1897                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora flava</i>            |            | P754 | 131118   | Claparede, 1870               |               | 1             |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Dipolydora sp. B</i>            |            | P762 | 131124   | Verrill, 1881                 |               |               | 1             |               | 1             |               |               |               | 1             |               |               |               | 1             |
| <i>Pseudopolydora pulchra</i>      |            | P774 | 131169   | (Carazzi, 1895)               |               |               |               |               |               |               | 1             |               | 1             |               |               |               |               |
| <i>Scolelepis korsuni</i>          |            |      | 131174   | Sikorski, 1994                |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Spio goniocéphala</i>           |            |      | 131184   | Thulin, 1957                  |               |               |               |               |               |               | 2             |               |               | 2             |               |               |               |
| <i>Spio symphyta</i>               |            |      | 596189   | Meisner, Bick & Bastrop, 2011 | 1             | 3             | 3             | 1             |               | 1             |               | 2             | 1             | 1             | 1             | 1             | 4             |
| <i>Spiophanes bombyx</i>           |            | P794 | 131187   | (Claparede, 1870)             | 10            | 6             | 8             | 9             | 12            | 4             | 4             |               | 9             | 7             | 4             | 4             | 5             |
| <i>Spiophanes kroyeri</i>          |            | P796 | 131188   | Grube, 1860                   |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Magelona johnstoni</i>          |            |      | 130269   | Fiege, Lichen & Mackie, 2000  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Chaetopterus variopedatus</i>   |            | P814 | 129914   | (Renier, 1804)                |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Phyllochaetopterus anglicus</i> |            | P815 |          | Potts, 1914                   |               | 4             |               |               |               |               | 1             | 1             |               |               |               |               |               |
| <i>Aphelochaeta sp. A</i>          |            | P823 | 129240   | Blake, 1991                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Caulleriella alata</i>          |            | P829 | 129943   | (Southern, 1914)              | 3             | 1             | 2             |               | 1             | 1             | 3             | 1             | 2             | 3             | 3             |               | 3             |



| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |                          | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Chaetozone zetlandica</i>      |            | P831  | 129948   | (McIntosh, 1911)         |               |               | 1             | 2             |               |               | 2             |               | 1             |               | 1             |               |               |
| <i>Chaetozone christiei</i>       |            |       | 152217   | Chambers, 2000           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cirratulus cirratus</i>        |            | P836  | 129959   | (O.F.Muller,1776)        |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Tharyx</i>                     |            | P847  | 129249   | Webster & Benedict, 1887 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Flabelligera affinis</i>       |            | P881  | 130103   | M.Sars, 1829             |               |               |               |               | 1             |               |               |               |               |               |               |               | 1             |
| <i>Mediomastus fragilis</i>       |            | P919  | 129892   | Rasmussen, 1973          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Notomastus</i>                 |            | P920  | 129220   | Sars, 1850               | 2             |               |               | 3             | 3             | 3             | 2             | 1             | 3             | 4             |               | 1             |               |
| <i>Praxillura longissima</i>      |            | P944  | 130327   | Arwidsson, 1906          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leiochone</i>                  |            |       | 146991   | Grube, 1868              | 3             |               | 1             | 1             | 1             | 2             | 2             | 2             |               | 2             |               |               |               |
| <i>Euclymene lombricoides</i>     |            | P963  | 209899   | (Quatrefages, 1866)      |               | 1             | 1             | 1             | 1             |               | 1             |               |               |               |               |               |               |
| <i>Euclymene oerstedii</i>        |            | P964  | 130294   | (Claparède, 1863)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Praxillella affinis</i>        |            | P971  | 130322   | (M Sars, 1872)           | 1             | 3             | 1             | 2             | 1             | 5             |               |               | 2             | 1             |               | 1             |               |
| <i>Ophelia borealis</i>           |            | P999  | 130491   | Quatrefages, 1866        |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Ophelia celtica</i>            |            | P1000 | 130492   | Amoureux & Dauvin, 1981  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelina acuminata</i>         |            | P1014 | 130500   | Örsted, 1843             |               | 3             | 1             | 1             | 3             |               |               | 1             |               |               |               |               |               |
| <i>Asclerocheilus intermedius</i> |            | P1022 | 130974   | (Saint-Joseph, 1894)     | 1             | 1             | 3             | 3             | 1             | 1             |               |               |               | 1             |               | 1             |               |
| <i>Scalibregma celticum</i>       |            | P1026 | 130979   | Mackie, 1991             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalibregma inflatum</i>       |            | P1027 | 130980   | Rathke, 1843             |               | 2             | 2             | 1             | 2             | 1             | 2             |               |               |               | 1             |               | 2             |
| <i>Polygordius</i>                |            | P1062 | 129472   | Schneider, 1868          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Galathowenia oculata</i>       |            | P1093 | 146950   | Zaks, 1922               |               |               |               | 1             |               |               | 1             |               |               | 2             |               |               |               |
| <i>Owenia</i>                     |            | P1097 | 129427   | Delle Chiaje, 1844       |               | 1             | 1             |               |               |               | 1             |               | 1             | 1             | 1             | 2             | 2             |
| <i>Lagis koreni</i>               |            | P1107 | 152367   | Malmgren, 1866           | 2             |               | 1             |               |               |               |               |               | 3             |               | 3             | 2             | 2             |
| <i>Sabellaria spinulosa</i>       |            | P1117 | 130867   | Leuckart, 1849           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampharete lindstroemi</i>      | aggregate  | P1139 | 129778   | M. Sars, 1864            | 3             | 5             | 4             | 5             | 1             | 3             | 4             |               | 2             | 3             | 5             | 2             | 6             |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |   | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Anothenus gracilis</i>         |            | P1147 | 129789   | (Malmgren, 1866)                              |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Terebellides stroemii</i>      |            | P1175 | 131573   | Sars, 1835                                    |               | 1             |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Lanice conchilega</i>          |            | P1195 | 131495   | (Pallas, 1766)                                | 1             |               |               | 1             | 2             |               | 1             |               |               | 1             |               |               | 2             |
| <i>Phisidia aurea</i>             |            | P1215 | 131513   | Southward, 1956                               | 3             |               | 1             |               |               | 1             |               |               |               | 1             |               |               |               |
| <i>Pista mediterranea</i>         |            |       | 131519   | de Gaillande, 1970                            |               |               |               | 2             |               |               |               |               |               |               |               |               |               |
| <i>Pista bansei</i>               |            |       | 152254   | Saphronova, 1988                              | 1             |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Polycirrus</i>                 |            | P1235 | 129710   | Grube, 1850                                   | 3             |               | 2             | 2             | 3             | 4             | 2             |               | 2             |               | 3             |               | 1             |
| <i>Lysilla loveni</i>             |            | P1233 | 131500   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lysilla nivea</i>              |            | P1234 | 131501   | Langerhans, 1884                              | 2             | 2             |               | 2             |               |               | 2             |               |               | 5             |               | 1             |               |
| <i>Streblosoma intestinale</i>    |            | P1252 | 131540   | M. Sars in G.O. Sars, 1872                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thelepus cincinnatus</i>       |            | P1254 | 131543   | (Fabricius, 1780)                             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dialychone dunerificta</i>     |            |       | 558752   | (Tovar-Hernández, Licciano, Giangrande, 2007) | 2             |               |               | 1             | 4             |               | 1             |               |               |               |               |               |               |
| <i>Parasabella cambrensis</i>     |            | P1273 | 530920   | Knight-Jones & Walker, 1985                   |               |               |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Euchone pararosea</i>          |            |       | 390407   | Giangrande & Licciano, 2006                   |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Pseudopotamilla reniformis</i> |            | P1316 | 130963   | (Bruguiere, 1789)                             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Sabella pavonina</i>           |            | P1320 | 130967   | Savigny, 1822                                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hydroides norvegica</i>        |            | P1334 | 131009   | Gunnerus, 1768                                |               | 2             |               | 5             |               | 1             |               |               |               |               |               |               |               |
| <i>Spirobranchus triqueter</i>    |            | P1341 | 555935   | (Linnaeus, 1758)                              |               | 2             | 1             | 8             | 9             | 2             | 4             |               |               | 1             |               |               |               |
| <b>OLIGOCHAETA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Grania</i>                     |            | P1524 | 137349   | Southern, 1913                                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>CHELICERATA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nymphon brevirostre</i>        |            | Q5    | 150520   | Hodge, 1863                                   | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Anoplodactylus petiolatus</i>  |            | Q44   | 134723   | (Kroyer, 1884)                                |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|--------------------------------|------------|------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                |            |      |          |                                   | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |   |
| <b>CRUSTACEA</b>               |            |      |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Scalpellum scalpellum</i>   |            | R22  | 106204   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Verruca stroemia</i>        |            | R41  | 106257   | O.F.Muller, 1776                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Nebalia reboredae</i>       |            |      | 459311   | Moreira & Urgorri, 2009           |               |               | 1             |               | 1             |               |               |               |               |               |               |               |               |   |
| <i>Nebalia borealis</i>        |            | S7   | 156257   | Dahl, 1985                        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Sarsinebalia urgorrii</i>   |            |      | 388224   | Moreira, Gestoso & Troncoso, 2003 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Heteromysis formosa</i>     |            | S93  | 120037   | (G. O. Sars, 1877)                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Kroyera carinata</i>        |            | S125 | 547074   | Spence Bate, 1857                 |               |               |               |               | 1             |               |               |               |               |               |               |               |               |   |
| <i>Perioculodes longimanus</i> |            | S131 | 102915   | (Bate & Westwood, 1868)           |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Pontocrates arenarius</i>   |            | S135 | 102918   | (Bate, 1858)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Synchelidium maculatum</i>  |            | S138 | 102928   | Stebbing, 1906                    |               |               |               |               |               |               | 1             |               |               |               |               |               |               |   |
| <i>Apolochus neapolitanus</i>  |            | S159 | 236495   | (Della Valle, 1893)               |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Leucothoe incisa</i>        |            | S177 | 102460   | Robertson, 1892                   |               |               |               |               |               | 1             |               | 1             | 2             |               |               |               | 1             |   |
| <i>Leucothoe procera</i>       |            | S179 | 102466   | Bate, 1857                        |               |               |               | 1             |               |               |               |               |               |               |               |               |               |   |
| <i>Stenothoe marina</i>        |            | S213 | 103166   | (Bate, 1856)                      |               |               |               |               |               |               |               |               |               |               |               | 1             |               |   |
| <i>Urothoe elegans</i>         |            | S248 | 103228   | (Bate, 1856)                      |               | 1             | 1             |               | 1             |               |               |               |               |               |               |               | 2             | 2 |
| <i>Urothoe marina</i>          |            | S249 | 103233   | (Bate, 1857)                      | 33            | 17            | 40            | 29            | 42            | 15            | 13            | 3             | 17            | 22            | 16            | 22            | 6             |   |
| <i>Acidostoma neglectum</i>    |            |      | 102495   | (Spence Bate & Westwood, 1861)    |               |               | 2             |               |               |               |               |               |               |               |               |               |               |   |
| <i>Hippomedon denticulatus</i> |            | S296 | 102570   | (Bate, 1857)                      |               |               |               |               | 1             |               |               |               |               |               |               |               |               |   |
| <i>Tryphosa nana</i>           |            | S321 | 102691   | (Kroyer, 1846)                    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Tmetonyx similis</i>        |            | S337 | 102742   | (G O Sars, 1891)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Nototropis falcatus</i>     |            | S410 | 102139   | Metzger, 1871                     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Nototropis vedlomensis</i>  |            | S413 | 102132   | (Bate & Westwood, 1862)           | 1             | 1             | 1             | 4             | 3             |               | 5             |               |               |               | 3             |               |               |   |
| <i>Ampelisca diadema</i>       |            | S429 | 101896   | (Costa, 1853)                     | 2             | 7             | 4             | 6             | 4             | 7             | 7             |               | 2             |               |               | 3             | 1             |   |

| Taxon                              | Qualifiers | SDC         | APHIA ID      | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|-------------|---------------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |             |               |                                   | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Ampelisca provincialis</i>      |            | S434        | 101915        | Bellan-Santini & Kaim-Malka, 1977 | 1             | 16            | 11            | 9             | 14            | 9             | 2             |               |               | 6             |               | 1             | 2             |
| <i>Ampelisca spinipes</i>          |            | S438        | 101928        | Boeck, 1861                       | 3             | 1             |               | 1             | 1             |               | 1             |               |               | 6             | 2             | 2             | 5             |
| <i>Ampelisca tenuicornis</i>       |            | S440        | 101930        | Lilljeborg, 1855                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca typica</i>            |            | S442        | 101933        | (Bate, 1856)                      | 2             | 3             | 8             | 3             | 4             | 4             | 2             |               | 2             | 1             | 4             | 4             | 2             |
| <i>Bathyporeia elegans</i>         |            | S452        | 103058        | Watkin, 1938                      |               |               |               |               |               |               |               |               |               |               | 1             |               |               |
| <i>Bathyporeia gracilis</i>        |            | S453        | 103059        | G O Sars, 1891                    |               |               |               |               |               |               |               | 2             | 6             |               | 3             |               |               |
| <i>Bathyporeia guilliamsoniana</i> |            | S454        | 103060        | (Bate, 1857)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Haustorius arenarius</i>        |            | S462        | 102317        | Slabber, 1769                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abludomelita obtusata</i>       |            | S498        | 102788        | (Montagu, 1813)                   |               |               |               |               |               | 1             |               |               | 1             |               |               |               |               |
| <b>Cheirocratus</b>                |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b>               |               |               |               |               |               |               |               |               | 2             |               |               |               |               |
| <i>Othomaera othonis</i>           |            | S519        | 534781        | (Milne-Edwards, 1830)             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Maerella tenuimana</i>          |            | S521        | 102831        | (Bate, 1862)                      | 1             |               | 1             |               | 1             |               | 1             |               |               | 1             | 2             |               |               |
| <i>Megamphopus cornutus</i>        |            | S539        | 148545        | Norman, 1869                      |               | 1             | 2             |               |               |               | 1             |               |               |               | 2             |               |               |
| <i>Gammaropsis maculata</i>        |            | S541        | 102364        | (Johnston, 1828)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Photis longicaudata</i>         |            | S552        | 102383        | (Bate & Westwood, 1862)           |               |               | 10            |               |               |               | 4             |               |               |               |               |               |               |
| <b>Aoridae</b>                     |            | <b>S577</b> | <b>101368</b> | <b>Stebbing, 1899</b>             | 1             | 1             |               |               | 3             |               |               | 1             | 2             | 1             | 2             |               |               |
| <i>Leptocheirus hirsutimanus</i>   |            | S588        | 102036        | (Bate, 1862)                      | 4             | 1             |               |               |               | 2             |               | 1             | 1             |               |               | 1             |               |
| <i>Crassicorophium crassicorne</i> |            | S611        | 397383        | Bruzelius, 1859                   |               |               |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Unciola planipes</i>            |            | S622        | 102061        | Norman, 1867                      | 1             |               |               |               |               |               |               |               |               |               | 2             |               |               |
| <i>Phtisica marina</i>             |            | S657        | 101864        | Slabber, 1769                     |               |               |               |               | 1             | 1             |               |               |               |               |               |               |               |
| <b>Gnathia</b>                     |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>                |               |               |               | 1             |               | 2             |               |               |               | 1             |               |               |               |
| <i>Conilera cylindracea</i>        |            | S849        | 118842        | (Montagu, 1804)                   | 1             |               | 1             |               | 2             |               |               |               |               | 1             |               |               |               |
| <i>Eurydice pulchra</i>            |            | S854        | 118852        | Leach, 1815                       |               | 2             |               | 2             | 3             | 6             | 2             | 2             | 4             | 3             |               | 2             |               |
| <i>Astacilla longicornis</i>       |            | S955        | 119024        | (Sowerby, 1806)                   |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Tanaopsis graciloides</i>       |            | S1142       | 136458        | (Lilljeborg, 1864)                |               | 1             |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|-------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |       |          |                                   | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Bodotria scorpioides</i>    |            | S1197 | 110445   | (Montagu, 1804)                   |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Diastylis bradyi</i>        |            | S1248 | 110472   | Norman, 1879                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Processa modica</i>         |            | S1366 | 108343   | Williamson & Rochanaburanon, 1979 |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Callianassa subterranea</i> |            | S1415 | 107729   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Upogebia deltaura</i>       |            | S1419 | 107739   | (Leach, 1815)                     | 1             |               | 1             |               |               |               |               |               |               |               |               | 1             | 1             |
| <i>Pisidia longicornis</i>     |            | S1482 | 107188   | (Linnaeus, 1757)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ebalia tumefacta</i>        |            | S1509 | 107302   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hyas coarctatus</i>         |            | S1519 | 107323   | Leach, 1815                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Inachus dorsettensis</i>    |            | S1526 | 107327   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia rostrata</i>     |            | S1532 | 107345   | (Linnaeus, 1761)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thia scutellata</i>         |            | S1559 | 107281   | (Fabricius, 1793)                 |               |               |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Liocarcinus pusillus</i>    |            | S1584 | 107393   | (Leach, 1816)                     |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pinnotheres pisum</i>       |            | S1638 | 107473   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>MOLLUSCA</b>                |            |       |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptochiton asellus</i>     |            | W53   | 140199   | (Gmelin, 1791)                    |               | 1             | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Acanthochitona crinita</i>  |            | W86   | 138675   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gibbula magus</i>           | ?          | W159  | 141790   | (Linnaeus, 1758)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Euspira nitida</i>          |            | W491  | 151894   | (Donovan, 1804)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Melanella alba</i>          |            | W634  | 139832   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Colus gracilis</i>          |            | W715  | 138899   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Philine</i>                 |            | W1036 | 138339   | Ascanius, 1772                    |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Tritonia plebeia</i>        |            | W1254 | 141738   | Johnston, 1828                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Onchidorididae                 |            | W1319 | 175      | Gray, 1827                        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Knoutsodonta depressa</i>   |            | W1323 | 845528   | (Alder & Hancock, 1842)           |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Nucula hanleyi</i>          |            | W1568 | 140588   | Winckworth, 1931                  |               |               | 1             |               |               |               | 1             |               |               |               |               |               | 1             |



| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority            | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|--------------------------------|------------|-------|----------|----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                |            |       |          |                      | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |   |
| <i>Modiolus adriaticus</i>     |            | W1700 | 506025   | (Lamarck, 1819)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Aequipecten opercularis</i> |            | W1773 | 140687   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Heteranomia squamula</i>    |            | W1809 | 138749   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Pododesmus squama</i>       |            | W1812 | 138752   | (Linnaeus, 1761)     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Hemilepton nitidum</i>      |            | W1882 | 246148   | (W. Turton, 1822)    |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <i>Kurtiella bidentata</i>     |            | W1906 | 345281   | (Montagu, 1803)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Spisula elliptica</i>       |            | W1975 | 140300   | (Brown, 1827)        | 4             |               |               |               | 1             |               |               |               |               |               | 1             | 1             | 1             |   |
| <i>Ensis leei</i>              |            | W1997 | 876640   | M. Huber, 2015       |               | 1             |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Phaxas pellucidus</i>       |            | W2006 | 140737   | (Pennant, 1777)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Moerella donacina</i>       |            | W2021 | 147021   | Linnaeus, 1758       |               |               | 1             |               |               |               |               |               |               |               |               |               |               |   |
| <i>Asbjornsenia pygmaea</i>    |            | W2023 | 879714   | (Lovén, 1846)        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Gari fervensis</i>          |            | W2051 | 140870   | (Gmelin, 1791)       |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Solecurtus scopula</i>      |            | W2054 | 141543   | (Turton, 1822)       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Abra alba</i>               |            | W2059 | 141433   | (W Wood, 1802)       |               |               |               |               |               |               |               |               | 1             |               |               |               |               | 2 |
| <i>Abra prismatica</i>         |            | W2062 | 141436   | (Montagu, 1808)      |               |               |               | 1             |               |               |               |               |               |               | 1             | 1             |               |   |
| <i>Clausinella fasciata</i>    |            | W2100 | 141909   | (da Costa, 1778)     |               |               |               |               |               |               | 1             |               |               |               |               |               |               |   |
| <i>Timoclea ovata</i>          |            | W2104 | 141929   | (Pennant, 1777)      |               |               |               | 3             |               | 1             | 1             |               |               |               |               |               |               |   |
| <i>Venerupis corrugata</i>     |            | W2124 | 181364   | (Gmelin, 1791)       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Dosinia exoleta</i>         |            | W2130 | 141911   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <i>Hiatella arctica</i>        |            | W2166 | 140103   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Thracia villosiuscula</i>   |            | W2233 | 141651   | (Macgillivray, 1827) |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Cochlodesma praetenu</i>    |            | W2239 | 181373   | (Pulteney, 1799)     |               |               |               |               |               |               |               |               | 1             | 1             |               |               |               |   |
| <b>PHORONIDA</b>               |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Phoronis</i>                |            | ZA3   | 128545   | Wright, 1856         | 7             | 13            |               | 13            | 5             | 2             | 12            | 12            | 17            | 3             |               | 4             | 6             |   |
| <b>ECHINODERMATA</b>           |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |

| Taxon  | Qualifiers | SDC   | APHIA ID | Authority                           | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--|------------|-------|----------|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|  |            |       |          |                                     | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Ophiothrix fragilis</i>                         |            | ZB124 | 125131   | (Abildgaard, 1789)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Amphipholis squamata</i>                        |            | ZB161 | 125064   | (Chiaje, 1828)                      |               |               |               | 1             |               | 1             |               |               |               |               |               |               |               |
| <i>Ophiura albida</i>                              |            | ZB168 | 124913   | Forbes, 1839                        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Psammechinus miliaris</i>                       |            | ZB193 | 124319   | (Gmelin, 1778)                      |               | 1             |               |               | 2             |               |               |               |               |               |               |               | 1             |
| <i>Echinocyamus pusillus</i>                       |            | ZB212 | 124273   | (O.F. Muller, 1776)                 | 1             |               |               | 2             |               |               | 1             |               |               | 2             | 2             | 1             |               |
| <i>Echinocardium cordatum</i>                      |            | ZB223 | 124392   | (Pennant, 1777)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mesothuria intestinalis</i>                     |            | ZB233 | 124568   | (Ascanius, 1805)<br>Östergren, 1896 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudothyone raphanus</i>                       |            | ZB257 | 124661   | (Duben & Koren, 1845)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone fusus</i>                                |            | ZB262 | 124670   | O.F.Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone roscovita</i>                            |            | ZB264 | 124676   | Hérouard, 1889                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Panningia hyndmani</i>                          |            | ZB272 | 848014   | (Thompson, 1840)                    |               | 1             | 1             |               |               |               | 1             |               |               |               |               |               |               |
| <i>Leptosynapta inhaerens</i>                      |            | ZB296 | 124465   | (O.F.Muller, 1776)                  |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Oestergrenia digitata</i>                       |            | ZB300 | 152547   | (Montagu, 1804)                     |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <b>TUNICATA</b>                                    |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidiella aspersa</i>                          |            | ZD84  | 103718   | O F Müller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidia conchilega</i>                          | ?          | ZD88  | 103702   | O F Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Polycarpa fibrosa</i>                           |            | ZD112 | 103902   | (Stimpson, 1852)                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dendrodoa grossularia</i>                       |            | ZD120 | 103882   | (van Beneden 1846)                  | 1             | 1             | 1             |               |               |               |               |               |               | 1             |               |               |               |
| <b>CEPHALOCHORDATA</b>                             |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Branchiostoma lanceolatum</i>                   |            |       | 104906   | (Pallas, 1774)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                              |            |       |          |                                     | <b>51</b>     | <b>52</b>     | <b>49</b>     | <b>51</b>     | <b>48</b>     | <b>42</b>     | <b>56</b>     | <b>29</b>     | <b>40</b>     | <b>45</b>     | <b>31</b>     | <b>37</b>     | <b>42</b>     |
| <b>Abundance</b>                                   |            |       |          |                                     | <b>161</b>    | <b>137</b>    | <b>158</b>    | <b>163</b>    | <b>174</b>    | <b>117</b>    | <b>146</b>    | <b>55</b>     | <b>126</b>    | <b>124</b>    | <b>80</b>     | <b>85</b>     | <b>87</b>     |
| <b>The following taxa were merged for analysis</b> |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aoridae</i>                                     |            | S577  | 101368   | Stebbing, 1899                      | 1             | 1             |               |               | 3             |               |               | 1             | 2             | 1             | 2             |               |               |
| <i>Aoridae</i>                                     | female     | S577  | 101368   | Stebbing, 1899                      | 1             | 1             |               |               | 3             |               |               | 1             | 2             | 1             | 2             |               |               |

| Taxon   | Qualifiers | SDC         | APHIA ID      | Authority           | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|---|------------|-------------|---------------|---------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|   |            |             |               |                     | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |   |
| <i>Autonoe longipes</i>                               |            | S583        | 102021        | (Lilljeborg, 1852)  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <b>Gnathia</b>  |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>  |               |               |               | 1             |               | 2             |               |               |               | 1             |               |               |               |   |
| <i>Gnathia</i>  | female     | S793        | 118437        | Leach, 1814         |               |               |               |               |               | 1             |               |               |               |               |               |               |               |   |
| <i>Gnathia oxyuraea</i>                               |            | S796        | 118995        | (Lilljeborg, 1855)  |               |               |               | 1             |               | 1             |               |               |               | 1             |               |               |               |   |
| <b>Cheirocratus</b>                                   |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b> |               |               |               |               |               |               |               |               | 2             |               |               |               |               |   |
| <i>Cheirocratus</i>                                   | female     | S503        | 101669        | Norman, 1867        |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <i>Cheirocratus pseudosundevallii</i>                 | ?          |             |               | Gouillieux, 2019    |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <b>Number of taxa</b>                                 |            |             |               |                     | 1             | 1             | 0             | 1             | 1             | 1             | 0             | 1             | 2             | 2             | 1             | 0             | 0             | 0 |
| <b>Abundance</b>                                      |            |             |               |                     | 1             | 1             | 0             | 1             | 3             | 2             | 0             | 1             | 4             | 2             | 2             | 0             | 0             | 0 |
| <b>The following taxa were excluded from analysis</b> |            |             |               |                     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <b>Colonial</b>                                       |            |             |               |                     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| PORIFERA  |            | C1          | 558           | Grant, 1836         | P             |               | P             |               |               |               |               |               |               |               |               |               |               |   |
| <i>Leuckartiara octona</i>                            |            | D240        | 117791        | (Fleming, 1823)     |               |               |               |               |               |               |               |               |               |               |               |               | P             |   |
| <i>Tubularia</i>                                      |            | D163        | 117258        | Linnaeus, 1758      |               |               |               |               |               |               |               |               |               |               |               |               |               | P |
| <i>Eudendrium</i>                                     |            | D218        | 117093        | Ehrenberg, 1834     |               |               |               |               |               |               |               |               |               | P             |               |               |               |   |
| Bougainvilliidae                                      |            | D246        | 1594          | Lütken, 1850        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Phialella quadrata</i>                             |            | D343        | 117804        | (Forbes, 1848)      |               |               |               |               |               |               |               |               | P             |               |               |               | P             | P |
| <i>Calycella syringa</i>                              |            | D348        | 117402        | (Linnaeus, 1767)    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Lafoea dumosa</i>                                  |            | D386        | 117702        | (Fleming, 1828)     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Halcium</i>  |            | D390        | 117103        | Oken, 1815          |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Abietinaria</i>                                    |            | D408        | 117225        | Kirchenpauer, 1884  | P             |               |               |               |               |               |               |               |               |               |               |               |               | P |
| <i>Diphasia</i>                                       |            | D413        | 117228        | L Agassiz, 1862     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Hydrallmania falcata</i>                           |            | D424        | 117890        | (Linnaeus, 1758)    |               | P             | P             |               | P             |               | P             |               | P             |               |               |               |               |   |
| <i>Sertularella gayi</i>                              |            | D429        | 117902        | (Lamouroux, 1821)   | P             |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Sertularia cupressina</i>                          |            | D435        | 117913        | (Linnaeus, 1758)    |               | P             | P             |               | P             |               |               |               |               | P             |               |               |               |   |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|--------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                |            |      |          |                       | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |   |
| <i>Nemertesia</i>              |            | D462 | 117195   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| Campanulariidae                |            | D491 | 1606     | Peron & Lesueur, 1810 |               | P             |               |               | P             |               |               | P             |               |               |               |               |               | P |
| <i>Alcyonium digitatum</i>     |            | D597 | 125333   | Linnaeus, 1758        |               | P             |               | P             |               |               |               |               |               |               | P             |               |               |   |
| <i>Crisia</i>                  |            | Y13  | 111032   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Tubulipora</i>              |            | Y27  | 111054   | Lamarck, 1816         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Disporella hispida</i>      |            | Y66  | 111730   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Alcyonidium parasiticum</i> |            | Y81  | 111604   | (Fleming, 1828)       |               | P             |               |               |               |               |               | P             |               |               |               |               |               |   |
| <i>Vesicularia spinosa</i>     |            | Y131 | 111669   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               | P             |               |               |   |
| <i>Amathia lendigera</i>       |            | Y135 | 111659   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Amathia</i>                 |            | Y137 | 111023   | Farre, 1837           |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Aetea</i>                   |            | Y153 | 110819   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Conopeum reticulum</i>      |            | Y172 | 111351   | (Linnaeus, 1767)      |               |               |               |               |               |               |               | P             |               |               |               |               |               |   |
| <i>Electra pilosa</i>          |            | Y178 | 111355   | (Linnaeus, 1767)      |               |               |               |               |               |               | P             |               |               |               |               |               |               | P |
| <i>Callopora</i>               |            | Y201 | 110851   | J E Gray, 1848        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Beania mirabilis</i>        |            | Y261 | 111072   | Johnston, 1840        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Chorizopora brongiartii</i> |            | Y344 | 111304   | (Audouin, 1826)       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Escharella immersa</i>      |            | Y364 | 111484   | (Fleming, 1828)       |               |               |               |               | P             |               |               | P             |               |               |               |               | P             |   |
| <i>Porella concinna</i>        |            | Y385 | 111125   | (Busk, 1854)          |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Schizomavella</i>           |            | Y467 | 110829   | Canu & Bassler, 1917  | P             | P             | P             | P             |               |               | P             | P             |               |               | P             |               |               | P |
| <b>Damaged</b>                 |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Eumida</i>                  | damaged    | P163 | 129446   | Malmgren, 1865        |               |               |               |               |               |               |               |               |               |               | 1             |               |               |   |
| <i>Aricidea</i>                | damaged    | P675 | 129430   | Webster, 1879         |               |               |               |               |               | 1             |               |               |               |               |               |               |               |   |
| <i>Paradoneis</i>              | damaged    | P695 | 129433   | Hartman, 1965         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| Spionidae                      | damaged    | P720 | 913      | Grube, 1850           |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Dipolydora</i>              | damaged    | P748 | 129611   | Verrill, 1881         |               |               |               |               |               | 2             |               |               |               |               |               |               |               |   |

| Taxon               | Qualifiers | SDC    | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---------------------|------------|--------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                     |            |        |          |                         | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Terebellidae</i> | damaged    | P1179  | 982      | Johnston, 1846          | 3             |               |               |               | 1             |               |               |               |               | 2             |               |               |               |
| Sabellidae          | damaged    | P1257  | 985      | Latreille, 1825         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Serpulidae          | damaged    | P1324  | 988      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia</i>      | damaged    | S5     | 147031   | Leach, 1814             |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| AMPHIPODA           | damaged    | S97    | 1135     |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Urothoe</i>      | damaged    | S246   | 101789   | Dana, 1852              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Liocarcinus</i>  | damaged    | S1577  | 106925   | Stimpson, 1870          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| GASTROPODA          | damaged    | W88    | 101      |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| BIVALVIA            | damaged    | W1560  | 151265   | Goldfuss, 1820          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| SOLENOIDEA          | damaged    | W1991  | 14635    | Lamarck, 1809           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Ophiuridae          | damaged    | ZB165  | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <b>Fish</b>         |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Gobiesocidae        |            | ZG1180 | 125477   | Bleeker, 1859           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Juvenile</b>     |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| ANIMALIA            | eggs       |        | 2        |                         |               |               |               |               |               |               | 3             |               |               |               |               |               |               |
| SIPUNCULA           | juvenile   | N1     | 1268     |                         |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Golfingia</i>    | juvenile   | N12    | 136021   | Lankester, 1885         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera</i>      | juvenile   | P255   | 129296   | Lamarck, 1818           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Goniadidae          | juvenile   | P266   | 953      | Kinberg, 1866           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Nereididae          | juvenile   | P458   | 22496    | Blainville, 1818        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys</i>      | juvenile   | P494   | 129370   | Cuvier, 1817            | 1             |               |               |               |               |               |               |               | 3             |               | 1             | 1             |               |
| <i>Scolelepis</i>   | juvenile   | P778   | 129623   | Blainville, 1828        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| CRUSTACEA           | larva      | R1     | 1066     |                         |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| BALANOMORPHA        | juvenile   | R42    | 1082     | Burmeister, 1834        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca</i>    | juvenile   | S423   | 101445   | Kroyer, 1842            |               |               |               |               |               |               | 2             |               |               |               |               |               |               |



| Taxon                | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|----------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                      |            |       |          |                         | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| <i>Gnathia</i>       | juvenile   | S793  | 118437   | Leach, 1814             |               |               | 1             |               |               | 1             |               |               |               | 3             |               |               |               |
| GEBIIDEA             | juvenile   | S1403 | 477323   | de Saint Laurent, 1979  | 3             |               | 1             | 1             | 2             | 1             | 5             |               | 1             | 2             | 3             | 1             |               |
| Paguridae            | juvenile   | S1445 | 106738   | Latreille, 1802         |               |               |               | 1             | 1             | 1             |               |               | 1             | 1             |               |               |               |
| DECAPODA             | larva      | S1276 | 1130     | Latreille, 1830         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ebalia</i>        | juvenile   | S1504 | 106889   | Leach, 1817             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Majidae              | juvenile   | S1512 | 106760   | Samouelle, 1819         | 1             |               |               | 1             |               |               |               |               |               |               |               |               |               |
| Inachinae            | juvenile   | S1520 | 148436   | MacLeay, 1838           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia</i>    | juvenile   | S1529 | 205077   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Buccinidae           | eggs       | W702  | 149      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Tritonia</i>      | juvenile   | W1246 | 138580   | Cuvier, 1798            |               |               |               |               |               |               |               |               |               | 1             |               |               | 1             |
| <i>Nucula</i>        | juvenile   | W1565 | 138262   | Lamarck, 1799           |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| Mytilidae            | juvenile   | W1691 | 211      | Rafinesque, 1815        |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Modiolus</i>      | juvenile   | W1702 | 140467   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Pectinidae           | juvenile   | W1768 | 213      | Rafinesque, 1815        |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| Anomiidae            | juvenile   | W1805 | 214      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abra</i>          | juvenile   | W2058 | 138474   | Lamarck, 1818           |               |               |               |               | 1             |               |               |               |               |               |               |               | 1             |
| <i>Dosinia</i>       | juvenile   | W2126 | 138636   | Scopoli, 1777           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Ophiuridae           | juvenile   | ZB165 | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| ECHINOIDEA           | juvenile   | ZB181 | 123082   | Leske, 1778             | 1             |               |               | 1             |               | 1             | 1             |               |               |               |               |               |               |
| <i>Echinocardium</i> | juvenile   | ZB222 | 123426   | Gray, 1825              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| HOLOTHUROIDEA        | juvenile   | ZB229 | 123083   |                         |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone</i>        | juvenile   | ZB261 | 146116   | Oken, 1815              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptosynapta</i>  | juvenile   | ZB291 | 123449   | Verrill, 1867           | 1             |               |               | 1             | 1             |               |               |               |               |               |               |               |               |
| ASCIDIACEA           | juvenile   | ZD2   | 1839     | Nielsen, 1995           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Meiofaunal</b>    |            |       |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                 | Qualifiers | SDC  | APHIA ID | Authority | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------|------------|------|----------|-----------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                       |            |      |          |           | MA_ST14<br>FA | MA_ST15<br>FA | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| NEMATODA              |            | HD1  | 799      |           | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Parasitic</b>      |            |      |          |           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| COPEPODA              | parasitic  | R142 | 1080     |           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b> |            |      |          |           | <b>11</b>     | <b>8</b>      | <b>6</b>      | <b>9</b>      | <b>11</b>     | <b>6</b>      | <b>10</b>     | <b>0</b>      | <b>7</b>      | <b>13</b>     | <b>2</b>      | <b>5</b>      | <b>9</b>      |
| <b>Abundance</b>      |            |      |          |           | <b>12</b>     | <b>2</b>      | <b>2</b>      | <b>6</b>      | <b>10</b>     | <b>4</b>      | <b>11</b>     | <b>0</b>      | <b>7</b>      | <b>12</b>     | <b>4</b>      | <b>2</b>      | <b>3</b>      |

| Taxon                         | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                               |            |      |          |                       | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <b>CNIDARIA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cerianthus lloydii</i>     |            | D632 | 283798   | Gosse, 1859           |               | 1             |               |               | 1             |               |               |               | 1             |               |               | 2             |               |
| Actiniaria                    |            | D662 | 1360     |                       |               |               |               |               |               | 2             | 1             |               | 1             | 1             |               |               |               |
| <b>PLATYHELMINTHES</b>        |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| PLATYHELMINTHES               |            | F2   | 793      | Minot, 1876           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>NEMERTEA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| NEMERTEA                      |            | G1   | 152391   |                       | 3             | 4             | 4             | 8             | 7             | 4             | 3             | 5             | 5             | 2             | 6             | 8             | 5             |
| <b>SIPUNCULA</b>              |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Golfingia elongata</i>     |            | N14  | 175026   | (Keferstein, 1862)    | 1             | 2             |               |               |               |               |               | 1             |               |               |               | 4             | 1             |
| <i>Golfingia vulgaris</i>     |            | N17  | 136050   | (de Blainville, 1827) |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Nephasoma minutum</i>      |            | N25  | 136060   | (Keferstein, 1862)    | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phascolion strombus</i>    |            | N34  | 175043   | (Montagu, 1804)       |               |               |               |               |               |               |               | 2             |               |               | 1             |               |               |
| <b>POLYCHAETA</b>             |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pisione remota</i>         |            | P15  | 130707   | Southern, 1914        |               |               |               | 11            |               |               |               |               |               |               |               |               |               |
| <i>Aphrodita aculeata</i>     |            | P19  | 129840   | Malmgren, 1867        | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Enipo elisabethae</i>      |            | P43  | 130737   | McIntosh, 1900        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gattyana cirrhosa</i>      |            | P49  | 130749   | (Pallas, 1766)        |               |               |               | 1             |               |               |               |               |               |               |               | 1             |               |
| <i>Harmothoe</i>              |            | P50  | 129491   | Kinberg, 1855         |               | 1             |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Malmgrenia darbouxi</i>    |            |      | 863197   | (Pettibone, 1993)     | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Malmgrenia andreapolis</i> |            | P51  | 147008   | (McIntoch, 1874)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lepidonotus squamatus</i>  |            | P82  | 130801   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe inornata</i>        |            | P92  | 130601   | Johnston, 1839        |               | 1             |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Pholoe baltica</i>         |            | P95  | 130599   | Örsted, 1843          |               | 1             |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Sthenelais limicola</i>    |            | P109 | 131077   | (Ehlers, 1864)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eteone longa</i>           | aggregate  | P117 | 130616   | (Fabricius, 1780)     |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |      |          |                          | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Hesionura elongata</i>      |            | P122 | 130649   | (Southern, 1914)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mysta picta</i>             |            | P127 | 147026   | (Quatrefages, 1866)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phyllodoce groenlandica</i> |            | P141 | 334506   | (Örsted, 1842)           |               |               |               |               | 2             | 1             |               |               | 1             |               | 1             | 1             |               |
| <i>Phyllodoce longipes</i>     |            | P143 | 130763   | Kinberg, 1866            |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Phyllodoce rosea</i>        |            | P146 | 334514   | (Mcintosh, 1877)         |               |               |               |               | 2             |               |               |               |               |               |               | 1             |               |
| <i>Eulalia bilineata</i>       |            | P152 | 130624   | (Johnston, 1840)         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eulalia mustela</i>         |            | P155 | 130631   | Pleijel, 1987            |               |               |               | 1             |               |               |               |               |               |               |               | 1             |               |
| <i>Eumida sanguinea</i>        |            | P167 | 130644   | (Örsted, 1843)           |               | 3             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera alba</i>            |            | P256 | 130116   | (O.F. Muller, 1788)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera fallax</i>          |            | P259 | 336908   | Quatrefages, 1850        | 1             |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Glycera lapidum</i>         |            | P260 | 130123   | Quatrefages, 1866        | 1             | 1             | 3             |               |               |               |               |               |               | 1             | 1             |               | 2             |
| <i>Glycera oxycephala</i>      |            | P262 | 130126   | Ehlers, 1887             |               |               | 1             |               |               | 1             |               | 1             | 1             |               |               |               |               |
| <i>Glycinde nordmanni</i>      |            | P268 | 130136   | (Malmgren, 1865)         |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Goniadella gracilis</i>     |            | P276 | 130145   | (Verrill, 1873)          |               | 1             | 3             | 2             | 1             |               |               | 1             | 1             |               | 5             |               | 1             |
| <i>Psamathe fusca</i>          |            | P305 | 152249   | (Keferstein, 1862)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Oxydromus pallidus</i>      |            | P317 | 340203   | (Claparede, 1864)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Podarkeopsis capensis</i>   |            | P319 | 130195   | Day, 1963                |               |               |               |               |               | 1             |               |               | 1             |               |               |               |               |
| <i>Syllis garciai</i>          |            | P351 | 131431   | (Campoy, 1982)           | 2             |               | 1             | 1             |               |               | 1             |               |               |               | 5             | 1             |               |
| <i>Syllis parapari</i>         |            |      | 196002   | San Martin & Lopez, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Syllis pontxioi</i>         |            |      | 196003   | San Martin & López, 2000 |               |               |               | 2             |               |               |               |               |               |               |               |               |               |
| <i>Syllis armillaris</i>       |            | P365 | 131415   | (O.F. Muller, 1776)      |               |               |               |               |               |               | 1             | 1             | 1             |               |               |               |               |
| <i>Syllis variegata</i>        |            | P371 | 131458   | (Grube, 1860)            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eusyllis blomstrandii</i>   |            | P380 | 131290   | Malmgren, 1867           |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                             | Qualifiers | SDC  | APHIA ID | Authority                       | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|------|----------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |      |          |                                 | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Odontosyllis fulgurans</i>     |            | P387 | 131327   | (Audouin & Milne Edwards, 1833) |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Streptodonta pterochaeta</i>   |            | P391 | 238207   | Southern, 1914                  |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Streptosyllis campoyi</i>      |            |      | 238248   | Brito, Nunez & San Martin, 2000 | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Parexogone hebes</i>           |            | P421 | 757970   | (Webster & Benedict, 1884)      |               |               |               |               |               | 1             |               |               | 1             |               |               |               |               |
| <i>Exogone verugera</i>           |            | P423 | 131307   | (Claparede, 1868)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Sphaerosyllis</i>              |            | P424 | 129677   | Claparède, 1863                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Myrianida                         |            | P434 | 129659   | Milne Edwards, 1845             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eunereis longissima</i>        |            | P475 | 130375   | Johnston, 1840                  |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Nereis zonata</i>              |            | P478 | 130407   | Malmgren, 1867                  |               | 1             |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Nephtys assimilis</i>          |            | P495 | 130353   | Ørsted, 1843                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys caeca</i>              |            | P496 | 130355   | (Fabricius, 1780)               |               | 1             |               |               | 1             |               |               | 1             |               |               |               | 1             |               |
| <i>Nephtys cirrosa</i>            |            | P498 | 130357   | Ehlers, 1868                    |               |               | 1             |               |               |               | 2             |               |               | 2             |               | 1             |               |
| <i>Nephtys longosetosa</i>        |            | P503 | 130364   | Ørsted, 1843                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nothria conchylega</i>         |            | P545 | 130467   | (Sars, 1835)                    |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Paucibranchia bellii</i>       |            | P564 | 130072   | (Audouin & Milne-Edwards, 1833) |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lysidice unicornis</i>         |            | P568 | 742232   | (Grube, 1840)                   | 2             | 3             | 1             |               | 3             |               |               | 1             | 2             |               |               | 1             |               |
| <i>Lumbrineris cf. cingulata</i>  |            |      | 130240   | (Ehlers, 1868)                  |               |               |               |               | 3             |               |               |               |               |               |               |               |               |
| <i>Lumbrineris futilis</i>        |            | P582 | 851788   | (Audouin & Milne Edwards, 1834) |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Drilonereis filum</i>          |            | P591 | 129856   | (Claparède, 1868)               |               | 1             |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Protodorvillea kefersteini</i> |            | P638 | 130041   | (McIntosh, 1869)                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Schistomeringos rudolphi</i>   |            | P643 | 154127   | Delle Chiaje, 1828              |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Orbinia sertulata</i>          |            | P665 | 334310   | (Savigny, 1820)                 |               |               |               |               |               |               |               |               |               | 1             |               |               |               |



| Taxon                              | Qualifiers | SDC  | APHIA ID | Authority                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|------|----------|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |      |          |                               | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Scoloplos armiger</i>           |            | P672 | 334722   | (Muller, 1776)                |               |               |               |               |               |               |               |               |               | 2             |               | 2             |               |
| <i>Aricidea catherinae</i>         |            | P684 | 333034   | (Laubier, 1967)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aricidea cerrutii</i>           |            | P685 | 525497   | Laubier, 1966                 |               |               | 4             |               | 2             | 5             |               | 2             | 4             | 2             |               |               |               |
| <i>Cirrophorus branchiatus</i>     |            | P689 | 130576   | Ehlers, 1908                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Paradoneis ilvana</i>           |            | P698 |          | Castelli, 1985                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Paradoneis lyra</i>             |            | P699 | 130585   | (Southern, 1914)              |               |               | 1             |               | 2             |               |               | 1             |               |               |               | 1             |               |
| <i>Poecilochaetus serpens</i>      |            | P718 | 130711   | Allen, 1904                   | 2             | 6             |               |               | 2             | 1             |               |               | 3             | 1             | 1             | 15            | 3             |
| <i>Aonides paucibranchiata</i>     |            | P723 | 131107   | Southern, 1914                | 7             | 6             | 18            | 3             | 2             | 3             | 2             | 3             | 1             |               | 7             |               | 5             |
| <i>Atherospio guillei</i>          |            |      | 478336   | (Laubier & Ramos,1974)        | 1             | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Laonice bahusiensis</i>         |            | P733 | 131127   | Soderstrom, 1920              | 1             | 2             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora caulleryi/sp. A</i>  |            | P751 | 131116   | Mesnil, 1897                  |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Dipolydora flava</i>            |            | P754 | 131118   | Claparede, 1870               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora sp. B</i>            |            | P762 | 131124   | Verrill, 1881                 |               | 1             |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Pseudopolydora pulchra</i>      |            | P774 | 131169   | (Carazzi, 1895)               |               | 2             |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Scolecopsis korsuni</i>         |            |      | 131174   | Sikorski, 1994                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Spio goniocéphala</i>           |            |      | 131184   | Thulin, 1957                  |               | 1             |               |               |               |               | 1             |               |               | 3             |               |               |               |
| <i>Spio symphyta</i>               |            |      | 596189   | Meisner, Bick & Bastrop, 2011 |               | 1             | 1             | 2             | 1             |               |               |               | 1             | 1             | 4             | 1             |               |
| <i>Spiophanes bombyx</i>           |            | P794 | 131187   | (Claparede, 1870)             | 1             | 3             | 1             | 1             | 5             | 3             | 5             | 2             | 4             |               | 3             | 2             | 5             |
| <i>Spiophanes kroyeri</i>          |            | P796 | 131188   | Grube, 1860                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Magelona johnstoni</i>          |            |      | 130269   | Fiege, Lichen & Mackie, 2000  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Chaetopterus varipodatus</i>    |            | P814 | 129914   | (Renier, 1804)                |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Phyllochaetopterus anglicus</i> |            | P815 |          | Potts, 1914                   |               | 1             |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Aphelochaeta sp. A</i>          |            | P823 | 129240   | Blake, 1991                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Caulleriella alata</i>          |            | P829 | 129943   | (Southern, 1914)              |               |               |               |               |               | 1             |               |               | 1             |               |               | 2             | 4             |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |                          | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Chaetozone zetlandica</i>      |            | P831  | 129948   | (McIntosh, 1911)         | 1             |               |               |               |               |               |               | 1             | 1             | 1             |               |               | 1             |
| <i>Chaetozone christiei</i>       |            |       | 152217   | Chambers, 2000           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cirratulus cirratus</i>        |            | P836  | 129959   | (O.F.Muller,1776)        |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tharyx</i>                     |            | P847  | 129249   | Webster & Benedict, 1887 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Flabelligera affinis</i>       |            | P881  | 130103   | M.Sars, 1829             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mediomastus fragilis</i>       |            | P919  | 129892   | Rasmussen, 1973          |               |               |               | 1             | 1             |               |               |               |               |               |               |               |               |
| <i>Notomastus</i>                 |            | P920  | 129220   | Sars, 1850               |               | 3             | 1             |               | 1             |               |               | 2             | 2             |               |               | 6             | 2             |
| <i>Praxillura longissima</i>      |            | P944  | 130327   | Arwidsson, 1906          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leiochone</i>                  |            |       | 146991   | Grube, 1868              | 1             | 2             | 1             |               | 1             |               |               | 2             | 1             |               |               | 1             |               |
| <i>Euclymene lombricoides</i>     |            | P963  | 209899   | (Quatrefages, 1866)      |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Euclymene oerstedii</i>        |            | P964  | 130294   | (Claparède, 1863)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Praxillella affinis</i>        |            | P971  | 130322   | (M Sars, 1872)           |               | 1             |               | 1             | 3             |               |               |               | 1             |               |               |               |               |
| <i>Ophelia borealis</i>           |            | P999  | 130491   | Quatrefages, 1866        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelia celtica</i>            |            | P1000 | 130492   | Amoureux & Dauvin, 1981  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelina acuminata</i>         |            | P1014 | 130500   | Örsted, 1843             |               | 3             |               |               | 1             |               |               |               | 1             |               |               |               |               |
| <i>Asclerocheilus intermedius</i> |            | P1022 | 130974   | (Saint-Joseph, 1894)     |               |               |               |               | 1             |               |               |               | 2             |               |               | 1             | 3             |
| <i>Scalibregma celticum</i>       |            | P1026 | 130979   | Mackie, 1991             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalibregma inflatum</i>       |            | P1027 | 130980   | Rathke, 1843             |               | 5             |               |               | 3             |               |               |               | 1             |               |               | 3             | 1             |
| <i>Polygordius</i>                |            | P1062 | 129472   | Schneider, 1868          |               |               |               | 6             |               |               | 1             |               |               |               |               |               |               |
| <i>Galathowenia oculata</i>       |            | P1093 | 146950   | Zaks, 1922               |               | 1             |               |               |               |               |               |               | 3             |               |               |               |               |
| <i>Owenia</i>                     |            | P1097 | 129427   | Delle Chiaje, 1844       |               | 1             | 3             |               | 1             | 2             |               |               | 1             |               | 3             | 1             | 3             |
| <i>Lagis koreni</i>               |            | P1107 | 152367   | Malmgren, 1866           |               | 1             |               |               | 2             |               |               |               |               |               |               | 4             | 1             |
| <i>Sabellaria spinulosa</i>       |            | P1117 | 130867   | Leuckart, 1849           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampharete lindstroemi</i>      | aggregate  | P1139 | 129778   | M. Sars, 1864            |               | 7             |               |               | 15            | 1             |               | 2             | 2             |               |               | 3             | 6             |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|-----------------------------------|------------|-------|----------|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                   |            |       |          |   | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |   |
| <i>Anobothrus gracilis</i>        |            | P1147 | 129789   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Terebellides stroemii</i>      |            | P1175 | 131573   | Sars, 1835                                    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Lanice conchilega</i>          |            | P1195 | 131495   | (Pallas, 1766)                                |               |               | 1             |               | 1             |               |               |               |               |               |               | 2             |               |   |
| <i>Phisidia aurea</i>             |            | P1215 | 131513   | Southward, 1956                               |               | 1             |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Pista mediterranea</i>         |            |       | 131519   | de Gaillande, 1970                            |               |               |               |               |               |               |               | 1             |               |               |               |               |               | 1 |
| <i>Pista bansei</i>               |            |       | 152254   | Saphronova, 1988                              | 1             | 1             | 1             |               | 2             |               |               |               |               |               |               | 2             | 2             |   |
| <i>Polycirrus</i>                 |            | P1235 | 129710   | Grube, 1850                                   |               | 2             | 2             |               | 1             | 1             |               | 1             | 1             | 5             | 1             |               |               |   |
| <i>Lysilla loveni</i>             |            | P1233 | 131500   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Lysilla nivea</i>              |            | P1234 | 131501   | Langerhans, 1884                              | 4             |               | 2             |               | 3             |               |               | 2             | 3             |               |               |               | 1             | 1 |
| <i>Streblosoma intestinale</i>    |            | P1252 | 131540   | M. Sars in G.O. Sars, 1872                    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Thelepus cincinnatus</i>       |            | P1254 | 131543   | (Fabricius, 1780)                             |               |               |               |               |               |               |               | 1             |               |               |               |               |               |   |
| <i>Dialychone dunerificta</i>     |            |       | 558752   | (Tovar-Hernández, Licciano, Giangrande, 2007) |               | 1             |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Parasabella cambrensis</i>     |            | P1273 | 530920   | Knight-Jones & Walker, 1985                   |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Euchone pararosea</i>          |            |       | 390407   | Giangrande & Licciano, 2006                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Pseudopotamilla reniformis</i> |            | P1316 | 130963   | (Bruguiere, 1789)                             |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Sabella pavonina</i>           |            | P1320 | 130967   | Savigny, 1822                                 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Hydroides norvegica</i>        |            | P1334 | 131009   | Gunnerus, 1768                                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Spirobranchus triqueter</i>    |            | P1341 | 555935   | (Linnaeus, 1758)                              |               |               |               |               | 1             |               |               |               | 1             |               |               |               |               |   |
| <b>OLIGOCHAETA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Grania</i>                     |            | P1524 | 137349   | Southern, 1913                                |               |               |               | 1             |               |               |               |               |               |               |               |               |               |   |
| <b>CHELICERATA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Nymphon brevirostre</i>        |            | Q5    | 150520   | Hodge, 1863                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Anoplodactylus petiolatus</i>  |            | Q44   | 134723   | (Kroyer, 1884)                                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |      |          |                                   | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <b>CRUSTACEA</b>               |            |      |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalpellum scalpellum</i>   |            | R22  | 106204   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Verruca stroemia</i>        |            | R41  | 106257   | O.F.Muller, 1776                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia reboredae</i>       |            |      | 459311   | Moreira & Urgorri, 2009           |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Nebalia borealis</i>        |            | S7   | 156257   | Dahl, 1985                        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Sarsinebalia urgorrii</i>   |            |      | 388224   | Moreira, Gestoso & Troncoso, 2003 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Heteromysis formosa</i>     |            | S93  | 120037   | (G. O. Sars, 1877)                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Kroyera carinata</i>        |            | S125 | 547074   | Spence Bate, 1857                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Periculodes longimanus</i>  |            | S131 | 102915   | (Bate & Westwood, 1868)           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pontocrates arenarius</i>   |            | S135 | 102918   | (Bate, 1858)                      |               |               |               | 1             |               | 2             |               |               |               |               |               | 1             |               |
| <i>Synchelidium maculatum</i>  |            | S138 | 102928   | Stebbing, 1906                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Apolochus neapolitanus</i>  |            | S159 | 236495   | (Della Valle, 1893)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leucothoe incisa</i>        |            | S177 | 102460   | Robertson, 1892                   |               | 1             | 1             |               |               |               |               | 2             |               |               |               |               |               |
| <i>Leucothoe procera</i>       |            | S179 | 102466   | Bate, 1857                        |               |               |               |               |               |               |               |               |               |               | 1             |               |               |
| <i>Stenothoe marina</i>        |            | S213 | 103166   | (Bate, 1856)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Urothoe elegans</i>         |            | S248 | 103228   | (Bate, 1856)                      |               |               |               |               |               | 1             |               |               | 2             |               |               | 1             |               |
| <i>Urothoe marina</i>          |            | S249 | 103233   | (Bate, 1857)                      | 33            | 32            | 9             | 5             | 21            | 7             |               | 12            | 11            | 4             |               | 11            | 32            |
| <i>Acidostoma neglectum</i>    |            |      | 102495   | (Spence Bate & Westwood, 1861)    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hippomedon denticulatus</i> |            | S296 | 102570   | (Bate, 1857)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tryphosa nana</i>           |            | S321 | 102691   | (Kroyer, 1846)                    |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Tmetonyx similis</i>        |            | S337 | 102742   | (G O Sars, 1891)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nototropis falcatus</i>     |            | S410 | 102139   | Metzger, 1871                     |               |               |               |               |               | 2             | 2             |               |               |               | 1             |               |               |
| <i>Nototropis vedlomensis</i>  |            | S413 | 102132   | (Bate & Westwood, 1862)           |               | 4             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca diadema</i>       |            | S429 | 101896   | (Costa, 1853)                     |               | 3             |               |               | 4             |               |               | 3             | 2             |               | 1             | 2             |               |

| Taxon                              | Qualifiers | SDC         | APHIA ID      | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|-------------|---------------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |             |               |                                   | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Ampelisca provincialis</i>      |            | S434        | 101915        | Bellan-Santini & Kaim-Malka, 1977 | 1             | 5             |               |               | 8             |               |               | 2             | 4             |               |               | 4             |               |
| <i>Ampelisca spinipes</i>          |            | S438        | 101928        | Boeck, 1861                       |               | 1             |               |               |               | 1             |               |               | 3             |               |               | 1             | 9             |
| <i>Ampelisca tenuicornis</i>       |            | S440        | 101930        | Lilljeborg, 1855                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca typica</i>            |            | S442        | 101933        | (Bate, 1856)                      | 1             | 2             |               | 1             | 2             |               |               |               |               |               |               | 2             | 2             |
| <i>Bathyporeia elegans</i>         |            | S452        | 103058        | Watkin, 1938                      |               |               | 1             | 1             |               | 1             | 2             |               |               | 2             | 1             |               |               |
| <i>Bathyporeia gracilis</i>        |            | S453        | 103059        | G O Sars, 1891                    |               | 1             | 2             |               |               | 2             |               |               |               | 5             |               |               |               |
| <i>Bathyporeia guilliamsoniana</i> |            | S454        | 103060        | (Bate, 1857)                      | 1             |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <i>Haustorius arenarius</i>        |            | S462        | 102317        | Slabber, 1769                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abludomelita obtusata</i>       |            | S498        | 102788        | (Montagu, 1813)                   |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <b>Cheirocratus</b>                |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b>               |               |               |               |               |               |               |               |               | 1             | 1             |               |               |               |
| <i>Othomaera othonis</i>           |            | S519        | 534781        | (Milne-Edwards, 1830)             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Maerella tenuimana</i>          |            | S521        | 102831        | (Bate, 1862)                      |               | 1             |               | 1             | 1             |               |               | 1             |               |               |               |               |               |
| <i>Megamphopus cornutus</i>        |            | S539        | 148545        | Norman, 1869                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gammaropsis maculata</i>        |            | S541        | 102364        | (Johnston, 1828)                  |               |               |               | 2             |               |               |               |               |               |               |               |               |               |
| <i>Photis longicaudata</i>         |            | S552        | 102383        | (Bate & Westwood, 1862)           |               | 3             |               |               |               |               |               | 2             |               |               |               | 3             |               |
| <b>Aoridae</b>                     |            | <b>S577</b> | <b>101368</b> | <b>Stebbing, 1899</b>             |               | 1             |               |               | 1             | 1             |               |               |               | 1             |               | 1             |               |
| <i>Leptocheirus hirsutimanus</i>   |            | S588        | 102036        | (Bate, 1862)                      |               | 1             | 3             |               |               |               |               | 3             |               |               |               | 2             | 4             |
| <i>Crassikorophium crassicorne</i> |            | S611        | 397383        | Bruzelius, 1859                   |               |               |               |               |               |               |               | 1             |               |               |               |               | 1             |
| <i>Unciola planipes</i>            |            | S622        | 102061        | Norman, 1867                      |               |               |               |               |               | 3             | 1             |               |               |               |               |               | 1             |
| <i>Phtisica marina</i>             |            | S657        | 101864        | Slabber, 1769                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Gnathia</b>                     |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>                |               | 4             |               | 1             |               |               |               |               | 4             |               |               |               |               |
| <i>Conilera cylindracea</i>        |            | S849        | 118842        | (Montagu, 1804)                   |               |               |               |               | 1             |               |               |               | 1             |               |               |               | 1             |
| <i>Eurydice pulchra</i>            |            | S854        | 118852        | Leach, 1815                       | 5             | 2             | 1             | 5             | 2             |               |               |               | 3             | 1             | 3             |               | 2             |
| <i>Astacilla longicornis</i>       |            | S955        | 119024        | (Sowerby, 1806)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tanaopsis graciloides</i>       |            | S1142       | 136458        | (Lilljeborg, 1864)                |               |               |               |               |               |               |               |               |               |               |               |               |               |



| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|-------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |       |          |                                   | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Bodotria scorpioides</i>    |            | S1197 | 110445   | (Montagu, 1804)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Diastylis bradyi</i>        |            | S1248 | 110472   | Norman, 1879                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Processa modica</i>         |            | S1366 | 108343   | Williamson & Rochanaburanon, 1979 |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Callianassa subterranea</i> |            | S1415 | 107729   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Upogebia deltaura</i>       |            | S1419 | 107739   | (Leach, 1815)                     |               |               | 2             |               |               |               |               |               | 1             |               |               | 3             | 1             |
| <i>Pisidia longicornis</i>     |            | S1482 | 107188   | (Linnaeus, 1757)                  |               |               |               | 1             | 1             |               |               | 1             |               |               |               |               | 1             |
| <i>Ebalia tumefacta</i>        |            | S1509 | 107302   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hyas coarctatus</i>         |            | S1519 | 107323   | Leach, 1815                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Inachus dorsettensis</i>    |            | S1526 | 107327   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia rostrata</i>     |            | S1532 | 107345   | (Linnaeus, 1761)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thia scutellata</i>         |            | S1559 | 107281   | (Fabricius, 1793)                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Liocarcinus pusillus</i>    |            | S1584 | 107393   | (Leach, 1816)                     |               |               |               |               | 1             | 1             |               |               |               |               |               |               |               |
| <i>Pinnotheres pisum</i>       |            | S1638 | 107473   | (Linnaeus, 1767)                  |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <b>MOLLUSCA</b>                |            |       |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptochiton asellus</i>     |            | W53   | 140199   | (Gmelin, 1791)                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Acanthochitona crinita</i>  |            | W86   | 138675   | (Pennant, 1777)                   |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Gibbula magus</i>           | ?          | W159  | 141790   | (Linnaeus, 1758)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Euspira nitida</i>          |            | W491  | 151894   | (Donovan, 1804)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Melanella alba</i>          |            | W634  | 139832   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Colus gracilis</i>          |            | W715  | 138899   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Philine</i>                 |            | W1036 | 138339   | Ascanius, 1772                    |               |               |               |               |               | 1             |               |               | 1             | 1             | 1             |               |               |
| <i>Tritonia plebeia</i>        |            | W1254 | 141738   | Johnston, 1828                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Onchidorididae                 |            | W1319 | 175      | Gray, 1827                        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Knoutsodonta depressa</i>   |            | W1323 | 845528   | (Alder & Hancock, 1842)           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nucula hanleyi</i>          |            | W1568 | 140588   | Winckworth, 1931                  |               |               |               |               |               |               |               |               |               |               |               |               | 3             |

| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority            | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|-------|----------|----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |       |          |                      | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Modiolus adriaticus</i>     |            | W1700 | 506025   | (Lamarck, 1819)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aequipecten opercularis</i> |            | W1773 | 140687   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Heteranomia squamula</i>    |            | W1809 | 138749   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pododesmus squama</i>       |            | W1812 | 138752   | (Linnaeus, 1761)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hemilepton nitidum</i>      |            | W1882 | 246148   | (W. Turton, 1822)    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Kurtiella bidentata</i>     |            | W1906 | 345281   | (Montagu, 1803)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Spisula elliptica</i>       |            | W1975 | 140300   | (Brown, 1827)        |               |               | 1             | 2             |               | 4             | 2             | 1             | 1             |               |               |               |               |
| <i>Ensis leei</i>              |            | W1997 | 876640   | M. Huber, 2015       |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Phaxas pellucidus</i>       |            | W2006 | 140737   | (Pennant, 1777)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Moerella donacina</i>       |            | W2021 | 147021   | Linnaeus, 1758       |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Asbjornsenia pygmaea</i>    |            | W2023 | 879714   | (Lovén, 1846)        |               |               |               | 2             |               |               | 1             |               |               |               |               |               |               |
| <i>Gari fervensis</i>          |            | W2051 | 140870   | (Gmelin, 1791)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Solecurtus scopula</i>      |            | W2054 | 141543   | (Turton, 1822)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abra alba</i>               |            | W2059 | 141433   | (W Wood, 1802)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abra prismatica</i>         |            | W2062 | 141436   | (Montagu, 1808)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Clausinella fasciata</i>    |            | W2100 | 141909   | (da Costa, 1778)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Timoclea ovata</i>          |            | W2104 | 141929   | (Pennant, 1777)      |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Venerupis corrugata</i>     |            | W2124 | 181364   | (Gmelin, 1791)       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dosinia exoleta</i>         |            | W2130 | 141911   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hiatella arctica</i>        |            | W2166 | 140103   | (Linnaeus, 1758)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thracia villosiuscula</i>   |            | W2233 | 141651   | (Macgillivray, 1827) |               |               | 1             |               |               |               |               |               |               |               |               |               | 1             |
| <i>Cochlodesma praetenu</i>    |            | W2239 | 181373   | (Pulteney, 1799)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>PHORONIDA</b>               |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phoronis</i>                |            | ZA3   | 128545   | Wright, 1856         | 4             | 68            | 11            | 3             | 39            |               |               | 2             | 28            |               | 2             | 21            |               |
| <b>ECHINODERMATA</b>           |            |       |          |                      |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon  | Qualifiers | SDC   | APHIA ID | Authority                           | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|--|------------|-------|----------|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|  |            |       |          |                                     | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Ophiothrix fragilis</i>                         |            | ZB124 | 125131   | (Abildgaard, 1789)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Amphipholis squamata</i>                        |            | ZB161 | 125064   | (Chiaje, 1828)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophiura albida</i>                              |            | ZB168 | 124913   | Forbes, 1839                        |               |               |               |               |               |               |               |               | 1             | 2             |               |               |               |
| <i>Psammechinus miliaris</i>                       |            | ZB193 | 124319   | (Gmelin, 1778)                      |               |               |               |               |               |               |               |               | 2             |               |               |               |               |
| <i>Echinocyamus pusillus</i>                       |            | ZB212 | 124273   | (O.F. Muller, 1776)                 |               | 2             | 1             | 5             |               |               | 1             |               | 1             | 5             | 6             | 1             | 3             |
| <i>Echinocardium cordatum</i>                      |            | ZB223 | 124392   | (Pennant, 1777)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mesothuria intestinalis</i>                     |            | ZB233 | 124568   | (Ascanius, 1805)<br>Östergren, 1896 |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Pseudothyone raphanus</i>                       |            | ZB257 | 124661   | (Duben & Koren, 1845)               |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Thyone fusus</i>                                |            | ZB262 | 124670   | O.F.Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone roscovita</i>                            |            | ZB264 | 124676   | Hérouard, 1889                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Panningia hyndmani</i>                          |            | ZB272 | 848014   | (Thompson, 1840)                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptosynapta inhaerens</i>                      |            | ZB296 | 124465   | (O.F.Muller, 1776)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Oestergrenia digitata</i>                       |            | ZB300 | 152547   | (Montagu, 1804)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>TUNICATA</b>                                    |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Asciella aspersa</i>                            |            | ZD84  | 103718   | O F Müller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidia conchilega</i>                          | ?          | ZD88  | 103702   | O F Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Polycarpa fibrosa</i>                           |            | ZD112 | 103902   | (Stimpson, 1852)                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dendrodoa grossularia</i>                       |            | ZD120 | 103882   | (van Beneden 1846)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>CEPHALOCHORDATA</b>                             |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Branchiostoma lanceolatum</i>                   |            |       | 104906   | (Pallas, 1774)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                              |            |       |          |                                     | 25            | 54            | 31            | 31            | 46            | 25            | 16            | 31            | 56            | 22            | 21            | 43            | 37            |
| <b>Abundance</b>                                   |            |       |          |                                     | 79            | 207           | 84            | 75            | 158           | 52            | 27            | 63            | 126           | 45            | 56            | 125           | 113           |
| <b>The following taxa were merged for analysis</b> |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aoridae</i>                                     |            | S577  | 101368   | Stebbing, 1899                      |               | 1             |               |               | 1             | 1             |               |               |               | 1             |               | 1             |               |
| <i>Aoridae</i>                                     | female     | S577  | 101368   | Stebbing, 1899                      |               | 1             |               |               | 1             | 1             |               |               |               |               |               | 1             |               |

| Taxon   | Qualifiers | SDC         | APHIA ID      | Authority           | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---|------------|-------------|---------------|---------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|   |            |             |               |                     | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Autonoe longipes</i>                               |            | S583        | 102021        | (Lilljeborg, 1852)  |               |               |               |               |               |               |               |               |               | 1             |               |               |               |
| <b>Gnathia</b>  |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>  |               | <b>4</b>      |               | <b>1</b>      |               |               |               |               | <b>4</b>      |               |               |               |               |
| <i>Gnathia</i>  | female     | S793        | 118437        | Leach, 1814         |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gnathia oxyuraea</i>                               |            | S796        | 118995        | (Lilljeborg, 1855)  |               | 3             |               | 1             |               |               |               |               | 4             |               |               |               |               |
| <b>Cheirocratus</b>                                   |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b> |               |               |               |               |               |               |               |               | 1             | 1             |               |               |               |
| <i>Cheirocratus</i>                                   | female     | S503        | 101669        | Norman, 1867        |               |               |               |               |               |               |               |               | 1             | 1             |               |               |               |
| <i>Cheirocratus pseudosundevallii</i>                 | ?          |             |               | Gouillieux, 2019    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                                 |            |             |               |                     | <b>0</b>      | <b>2</b>      | <b>0</b>      | <b>1</b>      | <b>1</b>      | <b>1</b>      | <b>0</b>      | <b>0</b>      | <b>2</b>      | <b>2</b>      | <b>0</b>      | <b>1</b>      | <b>0</b>      |
| <b>Abundance</b>                                      |            |             |               |                     | <b>0</b>      | <b>5</b>      | <b>0</b>      | <b>1</b>      | <b>1</b>      | <b>1</b>      | <b>0</b>      | <b>0</b>      | <b>5</b>      | <b>2</b>      | <b>0</b>      | <b>1</b>      | <b>0</b>      |
| <b>The following taxa were excluded from analysis</b> |            |             |               |                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Colonial</b>                                       |            |             |               |                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| PORIFERA  |            | C1          | 558           | Grant, 1836         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leuckartiara octona</i>                            |            | D240        | 117791        | (Fleming, 1823)     |               |               |               |               | P             |               |               |               |               |               |               |               |               |
| <i>Tubularia</i>                                      |            | D163        | 117258        | Linnaeus, 1758      |               |               |               |               |               | P             |               |               |               |               |               |               |               |
| <i>Eudendrium</i>                                     |            | D218        | 117093        | Ehrenberg, 1834     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Bougainvilliidae                                      |            | D246        | 1594          | Lütken, 1850        |               | P             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phialella quadrata</i>                             |            | D343        | 117804        | (Forbes, 1848)      |               |               |               |               |               | P             |               |               |               |               |               |               |               |
| <i>Calycella syringa</i>                              |            | D348        | 117402        | (Linnaeus, 1767)    |               |               |               |               |               |               |               |               |               |               |               | P             |               |
| <i>Lafoea dumosa</i>                                  |            | D386        | 117702        | (Fleming, 1828)     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Halecium</i>                                       |            | D390        | 117103        | Oken, 1815          |               |               |               |               |               |               |               |               |               |               |               | P             |               |
| <i>Abietinaria</i>                                    |            | D408        | 117225        | Kirchenpauer, 1884  |               |               |               |               | P             |               |               |               |               |               |               |               |               |
| <i>Diphasia</i>                                       |            | D413        | 117228        | L Agassiz, 1862     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hydrallmania falcata</i>                           |            | D424        | 117890        | (Linnaeus, 1758)    | P             | P             |               | P             | P             |               |               | P             |               |               |               | P             |               |
| <i>Sertularella gayi</i>                              |            | D429        | 117902        | (Lamouroux, 1821)   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Sertularia cupressina</i>                          |            | D435        | 117913        | (Linnaeus, 1758)    |               | P             |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |  |
|--------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                |            |      |          |                       | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |  |
| <i>Nemertesia</i>              |            | D462 | 117195   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Campanulariidae                |            | D491 | 1606     | Peron & Lesueur, 1810 | P             | P             |               | P             | P             |               |               | P             |               | P             |               |               |               |  |
| <i>Alcyonium digitatum</i>     |            | D597 | 125333   | Linnaeus, 1758        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Crisia</i>                  |            | Y13  | 111032   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Tubulipora</i>              |            | Y27  | 111054   | Lamarck, 1816         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Disporella hispida</i>      |            | Y66  | 111730   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Alcyonidium parasiticum</i> |            | Y81  | 111604   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Vesicularia spinosa</i>     |            | Y131 | 111669   | (Linnaeus, 1758)      | P             |               |               |               | P             |               |               |               |               |               |               |               |               |  |
| <i>Amathia lendigera</i>       |            | Y135 | 111659   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Amathia</i>                 |            | Y137 | 111023   | Farre, 1837           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aetea</i>                   |            | Y153 | 110819   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Conopeum reticulum</i>      |            | Y172 | 111351   | (Linnaeus, 1767)      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Electra pilosa</i>          |            | Y178 | 111355   | (Linnaeus, 1767)      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Callopora</i>               |            | Y201 | 110851   | J E Gray, 1848        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Beania mirabilis</i>        |            | Y261 | 111072   | Johnston, 1840        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Chorizopora brongiartii</i> |            | Y344 | 111304   | (Audouin, 1826)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Escharella immersa</i>      |            | Y364 | 111484   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Porella concinna</i>        |            | Y385 | 111125   | (Busk, 1854)          |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Schizomavella</i>           |            | Y467 | 110829   | Canu & Bassler, 1917  |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <b>Damaged</b>                 |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Eumida</i>                  | damaged    | P163 | 129446   | Malmgren, 1865        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aricidea</i>                | damaged    | P675 | 129430   | Webster, 1879         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Paradoneis</i>              | damaged    | P695 | 129433   | Hartman, 1965         |               |               |               |               |               |               | 2             |               |               |               |               |               |               |  |
| Spionidae                      | damaged    | P720 | 913      | Grube, 1850           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Dipolydora</i>              | damaged    | P748 | 129611   | Verrill, 1881         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |



| Taxon               | Qualifiers | SDC    | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---------------------|------------|--------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                     |            |        |          |                         | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| <i>Terebellidae</i> | damaged    | P1179  | 982      | Johnston, 1846          |               |               | 1             |               |               |               |               |               | 3             |               |               | 1             |               |
| Sabellidae          | damaged    | P1257  | 985      | Latreille, 1825         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Serpulidae          | damaged    | P1324  | 988      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia</i>      | damaged    | S5     | 147031   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| AMPHIPODA           | damaged    | S97    | 1135     |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Urothoe</i>      | damaged    | S246   | 101789   | Dana, 1852              | 1             |               | 1             |               | 1             | 1             |               |               |               |               |               | 2             |               |
| <i>Liocarcinus</i>  | damaged    | S1577  | 106925   | Stimpson, 1870          |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| GASTROPODA          | damaged    | W88    | 101      |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| BIVALVIA            | damaged    | W1560  | 151265   | Goldfuss, 1820          |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| SOLENOIDEA          | damaged    | W1991  | 14635    | Lamarck, 1809           |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| Ophiuridae          | damaged    | ZB165  | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Fish</b>         |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Gobiesocidae        |            | ZG1180 | 125477   | Bleeker, 1859           |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <b>Juvenile</b>     |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| ANIMALIA            | eggs       |        | 2        |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| SIPUNCULA           | juvenile   | N1     | 1268     |                         | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Golfingia</i>    | juvenile   | N12    | 136021   | Lankester, 1885         |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Glycera</i>      | juvenile   | P255   | 129296   | Lamarck, 1818           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Goniadidae          | juvenile   | P266   | 953      | Kinberg, 1866           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Nereididae          | juvenile   | P458   | 22496    | Blainville, 1818        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys</i>      | juvenile   | P494   | 129370   | Cuvier, 1817            |               |               |               |               |               | 1             |               |               | 1             | 1             |               |               |               |
| <i>Scolelepis</i>   | juvenile   | P778   | 129623   | Blainville, 1828        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| CRUSTACEA           | larva      | R1     | 1066     |                         |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| BALANOMORPHA        | juvenile   | R42    | 1082     | Burmeister, 1834        |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Ampelisca</i>    | juvenile   | S423   | 101445   | Kroyer, 1842            |               | 1             |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |  |
|----------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                      |            |       |          |                         | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |  |
| <i>Gnathia</i>       | juvenile   | S793  | 118437   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| GEBIIDEA             | juvenile   | S1403 | 477323   | de Saint Laurent, 1979  | 1             | 3             | 2             |               | 4             |               |               |               | 3             |               |               | 1             |               |  |
| Paguridae            | juvenile   | S1445 | 106738   | Latreille, 1802         |               | 1             |               |               |               |               |               |               |               |               |               |               |               |  |
| DECAPODA             | larva      | S1276 | 1130     | Latreille, 1830         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Ebalia</i>        | juvenile   | S1504 | 106889   | Leach, 1817             |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Majidae              | juvenile   | S1512 | 106760   | Samouelle, 1819         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Inachinae            | juvenile   | S1520 | 148436   | MacLeay, 1838           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Macropodia</i>    | juvenile   | S1529 | 205077   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Buccinidae           | eggs       | W702  | 149      | Rafinesque, 1815        |               |               |               |               |               |               |               |               | 1             |               |               |               |               |  |
| <i>Tritonia</i>      | juvenile   | W1246 | 138580   | Cuvier, 1798            |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nucula</i>        | juvenile   | W1565 | 138262   | Lamarck, 1799           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Mytilidae            | juvenile   | W1691 | 211      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Modiolus</i>      | juvenile   | W1702 | 140467   | (Linnaeus, 1758)        |               |               |               | 1             |               |               |               |               |               |               |               |               |               |  |
| Pectinidae           | juvenile   | W1768 | 213      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Anomiidae            | juvenile   | W1805 | 214      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Abra</i>          | juvenile   | W2058 | 138474   | Lamarck, 1818           |               |               |               |               |               |               |               |               | 1             |               | 1             |               |               |  |
| <i>Dosinia</i>       | juvenile   | W2126 | 138636   | Scopoli, 1777           |               |               |               |               |               |               |               |               |               | 1             |               |               | 1             |  |
| Ophiuridae           | juvenile   | ZB165 | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| ECHINOIDEA           | juvenile   | ZB181 | 123082   | Leske, 1778             |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Echinocardium</i> | juvenile   | ZB222 | 123426   | Gray, 1825              |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| HOLOTHUROIDEA        | juvenile   | ZB229 | 123083   |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Thyone</i>        | juvenile   | ZB261 | 146116   | Oken, 1815              |               |               |               | 1             |               |               |               |               |               |               |               |               |               |  |
| <i>Leptosynapta</i>  | juvenile   | ZB291 | 123449   | Verrill, 1867           | 1             |               |               |               |               |               |               |               |               |               |               |               |               |  |
| ASCIDIACEA           | juvenile   | ZD2   | 1839     | Nielsen, 1995           |               |               |               |               |               | 1             |               |               |               |               |               |               |               |  |
| <b>Meiofaunal</b>    |            |       |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |

| Taxon                 | Qualifiers | SDC  | APHIA ID | Authority | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------|------------|------|----------|-----------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                       |            |      |          |           | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| NEMATODA              |            | HD1  | 799      |           |               | 1             |               |               |               |               |               |               | 3             |               |               | 2             |               |
| <b>Parasitic</b>      |            |      |          |           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| COPEPODA              | parasitic  | R142 | 1080     |           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b> |            |      |          |           | 7             | 12            | 3             | 4             | 7             | 5             | 1             | 3             | 6             | 3             | 2             | 8             | 3             |
| <b>Abundance</b>      |            |      |          |           | 4             | 8             | 4             | 2             | 5             | 3             | 2             | 1             | 12            | 2             | 2             | 7             | 3             |

| Taxon                         | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                               |            |      |          |                       | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <b>CNIDARIA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cerianthus lloydii</i>     |            | D632 | 283798   | Gosse, 1859           |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| Actiniaria                    |            | D662 | 1360     |                       |               |               |               | 1             |               |               |               | 2             |               |               | 1             |               |               |
| <b>PLATYHELMINTHES</b>        |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| PLATYHELMINTHES               |            | F2   | 793      | Minot, 1876           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>NEMERTEA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| NEMERTEA                      |            | G1   | 152391   |                       | 7             | 8             | 1             | 1             | 1             | 5             | 6             | 6             | 1             | 1             | 1             | 2             | 1             |
| <b>SIPUNCULA</b>              |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Golfingia elongata</i>     |            | N14  | 175026   | (Keferstein, 1862)    | 4             | 2             |               |               |               |               |               | 3             |               |               |               | 1             |               |
| <i>Golfingia vulgaris</i>     |            | N17  | 136050   | (de Blainville, 1827) | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephasoma minutum</i>      |            | N25  | 136060   | (Keferstein, 1862)    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phascolion strombus</i>    |            | N34  | 175043   | (Montagu, 1804)       |               |               |               |               |               |               |               |               | 1             |               | 1             |               |               |
| <b>POLYCHAETA</b>             |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pisione remota</i>         |            | P15  | 130707   | Southern, 1914        |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Aphrodita aculeata</i>     |            | P19  | 129840   | Malmgren, 1867        |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Enipo elisabethae</i>      |            | P43  | 130737   | McIntosh, 1900        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gattyana cirrhosa</i>      |            | P49  | 130749   | (Pallas, 1766)        | 1             |               |               |               |               |               | 1             |               |               |               |               | 1             |               |
| <i>Harmothoe</i>              |            | P50  | 129491   | Kinberg, 1855         | 3             |               |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Malmgrenia darbouxi</i>    |            |      | 863197   | (Pettibone, 1993)     |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Malmgrenia andreapolis</i> |            | P51  | 147008   | (McIntoch, 1874)      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lepidonotus squamatus</i>  |            | P82  | 130801   | (Linnaeus, 1758)      | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe inornata</i>        |            | P92  | 130601   | Johnston, 1839        | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe baltica</i>         |            | P95  | 130599   | Örsted, 1843          | 2             |               |               |               |               |               | 2             | 1             |               |               |               |               |               |
| <i>Sthenelais limicola</i>    |            | P109 | 131077   | (Ehlers, 1864)        |               |               |               |               |               |               |               |               | 1             |               | 1             | 1             |               |
| <i>Eteone longa</i>           | aggregate  | P117 | 130616   | (Fabricius, 1780)     |               |               |               |               |               |               | 1             |               | 1             |               |               | 1             |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|--------------------------------|------------|------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                |            |      |          |                          | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |   |
| <i>Hesionura elongata</i>      |            | P122 | 130649   | (Southern, 1914)         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Mysta picta</i>             |            | P127 | 147026   | (Quatrefages, 1866)      |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Phyllodoce groenlandica</i> |            | P141 | 334506   | (Örsted, 1842)           | 1             |               |               |               |               |               |               | 3             |               |               |               |               |               |   |
| <i>Phyllodoce longipes</i>     |            | P143 | 130763   | Kinberg, 1866            |               | 1             |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Phyllodoce rosea</i>        |            | P146 | 334514   | (Mcintosh, 1877)         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Eulalia bilineata</i>       |            | P152 | 130624   | (Johnston, 1840)         |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Eulalia mustela</i>         |            | P155 | 130631   | Pleijel, 1987            |               |               |               |               |               |               |               |               | 2             |               |               |               |               |   |
| <i>Eumida sanguinea</i>        |            | P167 | 130644   | (Örsted, 1843)           | 1             |               |               |               |               |               |               | 2             | 1             |               |               |               |               |   |
| <i>Glycera alba</i>            |            | P256 | 130116   | (O.F. Muller, 1788)      |               |               |               |               |               |               |               | 1             |               |               |               |               |               |   |
| <i>Glycera fallax</i>          |            | P259 | 336908   | Quatrefages, 1850        |               |               | 2             |               |               |               |               |               |               |               |               |               |               |   |
| <i>Glycera lapidum</i>         |            | P260 | 130123   | Quatrefages, 1866        |               |               | 1             |               |               |               |               |               | 4             |               |               |               |               |   |
| <i>Glycera oxycephala</i>      |            | P262 | 130126   | Ehlers, 1887             |               |               |               |               | 3             | 1             | 1             |               |               |               | 1             | 2             |               |   |
| <i>Glycinde nordmanni</i>      |            | P268 | 130136   | (Malmgren, 1865)         |               |               |               |               |               |               |               | 1             |               |               |               |               |               |   |
| <i>Goniadella gracilis</i>     |            | P276 | 130145   | (Verrill, 1873)          |               | 2             |               |               |               |               |               |               |               |               |               |               |               | 1 |
| <i>Psamathe fusca</i>          |            | P305 | 152249   | (Keferstein, 1862)       |               |               |               |               |               |               |               |               |               |               |               |               | 1             |   |
| <i>Oxydromus pallidus</i>      |            | P317 | 340203   | (Claparede, 1864)        |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Podarkeopsis capensis</i>   |            | P319 | 130195   | Day, 1963                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Syllis garciai</i>          |            | P351 | 131431   | (Campoy, 1982)           |               | 1             |               |               |               |               |               |               | 3             |               |               |               | 2             |   |
| <i>Syllis parapari</i>         |            |      | 196002   | San Martin & Lopez, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Syllis pontxioi</i>         |            |      | 196003   | San Martin & López, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Syllis armillaris</i>       |            | P365 | 131415   | (O.F. Muller, 1776)      | 5             |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <i>Syllis variegata</i>        |            | P371 | 131458   | (Grube, 1860)            | 4             |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Eusyllis blomstrandii</i>   |            | P380 | 131290   | Malmgren, 1867           |               |               |               |               |               |               |               |               |               |               |               |               |               |   |



| Taxon                             | Qualifiers | SDC  | APHIA ID | Authority                       | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
|-----------------------------------|------------|------|----------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|--|
|                                   |            |      |          |                                 | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |   |  |
| <i>Odontosyllis fulgurans</i>     |            | P387 | 131327   | (Audouin & Milne Edwards, 1833) |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Streptodonta pterochaeta</i>   |            | P391 | 238207   | Southern, 1914                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Streptosyllis campoyi</i>      |            |      | 238248   | Brito, Nunez & San Martin, 2000 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Parexogone hebes</i>           |            | P421 | 757970   | (Webster & Benedict, 1884)      |               |               |               |               |               |               |               | 1             |               |               |               | 1             |               |   |  |
| <i>Exogone verugera</i>           |            | P423 | 131307   | (Claparede, 1868)               |               |               | 1             |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Sphaerosyllis</i>              |            | P424 | 129677   | Claparède, 1863                 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| Myrianida                         |            | P434 | 129659   | Milne Edwards, 1845             |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Eunereis longissima</i>        |            | P475 | 130375   | Johnston, 1840                  |               |               |               | 1             |               |               |               |               |               |               |               |               |               |   |  |
| <i>Nereis zonata</i>              |            | P478 | 130407   | Malmgren, 1867                  | 4             |               |               |               |               |               |               |               | 1             |               |               |               |               |   |  |
| <i>Nephtys assimilis</i>          |            | P495 | 130353   | Ørsted, 1843                    |               |               |               | 1             |               |               |               |               |               |               |               |               | 1             |   |  |
| <i>Nephtys caeca</i>              |            | P496 | 130355   | (Fabricius, 1780)               | 1             |               |               | 2             |               |               |               |               |               | 3             |               |               |               |   |  |
| <i>Nephtys cirrosa</i>            |            | P498 | 130357   | Ehlers, 1868                    |               |               | 4             | 1             | 3             | 2             |               |               |               | 3             | 5             | 2             | 1             | 1 |  |
| <i>Nephtys longosetosa</i>        |            | P503 | 130364   | Ørsted, 1843                    |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Nothria conchylega</i>         |            | P545 | 130467   | (Sars, 1835)                    |               | 1             |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Paucibranchia bellii</i>       |            | P564 | 130072   | (Audouin & Milne-Edwards, 1833) |               |               |               |               |               |               | 1             |               |               |               |               |               |               |   |  |
| <i>Lysidice unicornis</i>         |            | P568 | 742232   | (Grube, 1840)                   | 2             | 1             |               |               |               |               |               |               |               | 1             |               |               |               |   |  |
| <i>Lumbrineris cf. cingulata</i>  |            |      | 130240   | (Ehlers, 1868)                  |               | 3             |               | 1             |               | 1             | 1             | 3             | 1             | 1             |               |               | 1             | 1 |  |
| <i>Lumbrineris futilis</i>        |            | P582 | 851788   | (Audouin & Milne Edwards, 1834) |               |               |               |               |               |               |               |               |               |               |               |               |               | 1 |  |
| <i>Drilonereis filum</i>          |            | P591 | 129856   | (Claparède, 1868)               |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Protodorvillea kefersteini</i> |            | P638 | 130041   | (McIntosh, 1869)                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Schistomeringos rudolphi</i>   |            | P643 | 154127   | Delle Chiaje, 1828              |               |               | 1             |               |               |               |               |               |               |               |               |               |               |   |  |
| <i>Orbinia sertulata</i>          |            | P665 | 334310   | (Savigny, 1820)                 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |  |

| Taxon                              | Qualifiers | SDC  | APHIA ID | Authority                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|------|----------|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |      |          |                               | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <i>Scoloplos armiger</i>           |            | P672 | 334722   | (Muller, 1776)                |               |               |               | 7             |               | 2             | 4             |               | 1             | 2             | 1             | 1             |               |
| <i>Aricidea catherinae</i>         |            | P684 | 333034   | (Laubier, 1967)               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aricidea cerrutii</i>           |            | P685 | 525497   | Laubier, 1966                 |               | 3             | 1             | 1             |               |               |               | 2             |               |               |               |               | 1             |
| <i>Cirrophorus branchiatus</i>     |            | P689 | 130576   | Ehlers, 1908                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Paradoneis ilvana</i>           |            | P698 |          | Castelli, 1985                |               |               |               |               |               |               |               |               |               |               |               |               | 1             |
| <i>Paradoneis lyra</i>             |            | P699 | 130585   | (Southern, 1914)              | 2             | 5             |               | 3             |               |               |               |               |               |               |               |               |               |
| <i>Poecilochaetus serpens</i>      |            | P718 | 130711   | Allen, 1904                   | 6             | 8             | 2             | 3             | 2             | 2             | 64            | 1             | 5             | 1             | 2             | 4             |               |
| <i>Aonides paucibranchiata</i>     |            | P723 | 131107   | Southern, 1914                | 6             | 7             | 7             |               |               | 3             | 1             | 3             | 3             |               |               | 5             | 1             |
| <i>Atherospio guillei</i>          |            |      | 478336   | (Laubier & Ramos,1974)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Laonice bahusiensis</i>         |            | P733 | 131127   | Soderstrom, 1920              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora caulleryi/sp. A</i>  |            | P751 | 131116   | Mesnil, 1897                  | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora flava</i>            |            | P754 | 131118   | Claparede, 1870               | 3             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora sp. B</i>            |            | P762 | 131124   | Verrill, 1881                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudopolydora pulchra</i>      |            | P774 | 131169   | (Carazzi, 1895)               | 1             | 1             |               |               | 1             |               |               |               |               |               |               |               |               |
| <i>Scolecopsis korsuni</i>         |            |      | 131174   | Sikorski, 1994                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Spio goniocéphala</i>           |            |      | 131184   | Thulin, 1957                  |               |               |               |               | 3             | 1             |               |               | 2             |               |               |               |               |
| <i>Spio symphyta</i>               |            |      | 596189   | Meisner, Bick & Bastrop, 2011 |               | 1             | 1             | 1             |               |               | 10            | 3             |               |               |               |               | 1             |
| <i>Spiophanes bombyx</i>           |            | P794 | 131187   | (Claparede, 1870)             | 7             | 7             | 2             | 14            | 6             | 1             | 1             | 2             | 6             | 4             | 2             | 2             |               |
| <i>Spiophanes kroyeri</i>          |            | P796 | 131188   | Grube, 1860                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Magelona johnstoni</i>          |            |      | 130269   | Fiege, Lichen & Mackie, 2000  |               |               |               |               |               | 1             |               |               |               |               |               |               |               |
| <i>Chaetopterus variopedatus</i>   |            | P814 | 129914   | (Renier, 1804)                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Phyllochaetopterus anglicus</i> |            | P815 |          | Potts, 1914                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Aphelochaeta sp. A</i>          |            | P823 | 129240   | Blake, 1991                   |               |               |               |               |               |               |               | 2             |               |               |               |               |               |
| <i>Caulleriella alata</i>          |            | P829 | 129943   | (Southern, 1914)              | 2             | 3             | 1             |               |               |               |               | 1             |               |               |               | 2             |               |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |                          | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <i>Chaetozone zetlandica</i>      |            | P831  | 129948   | (McIntosh, 1911)         | 2             |               |               |               |               |               | 1             | 1             | 2             |               |               |               |               |
| <i>Chaetozone christiei</i>       |            |       | 152217   | Chambers, 2000           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Cirratulus cirratus</i>        |            | P836  | 129959   | (O.F.Muller,1776)        |               |               |               |               |               |               |               | 1             |               |               |               | 3             |               |
| <i>Tharyx</i>                     |            | P847  | 129249   | Webster & Benedict, 1887 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Flabelligera affinis</i>       |            | P881  | 130103   | M.Sars, 1829             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Mediomastus fragilis</i>       |            | P919  | 129892   | Rasmussen, 1973          |               |               |               |               |               |               |               | 2             |               |               |               |               |               |
| <i>Notomastus</i>                 |            | P920  | 129220   | Sars, 1850               | 2             | 3             |               |               |               |               | 1             | 3             |               |               |               | 2             | 3             |
| <i>Praxillura longissima</i>      |            | P944  | 130327   | Arwidsson, 1906          |               |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Leiochone</i>                  |            |       | 146991   | Grube, 1868              | 2             |               | 1             |               |               |               |               |               | 1             |               |               |               |               |
| <i>Euclymene lombricoides</i>     |            | P963  | 209899   | (Quatrefages, 1866)      | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Euclymene oerstedii</i>        |            | P964  | 130294   | (Claparède, 1863)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Praxillella affinis</i>        |            | P971  | 130322   | (M Sars, 1872)           | 5             | 3             | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelia borealis</i>           |            | P999  | 130491   | Quatrefages, 1866        |               |               |               |               |               | 2             |               |               |               |               |               |               |               |
| <i>Ophelia celtica</i>            |            | P1000 | 130492   | Amoureux & Dauvin, 1981  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelina acuminata</i>         |            | P1014 | 130500   | Örsted, 1843             |               | 2             | 1             |               |               |               |               | 1             |               |               |               |               |               |
| <i>Asclerocheilus intermedius</i> |            | P1022 | 130974   | (Saint-Joseph, 1894)     | 1             | 1             |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Scalibregma celticum</i>       |            | P1026 | 130979   | Mackie, 1991             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalibregma inflatum</i>       |            | P1027 | 130980   | Rathke, 1843             | 2             | 3             |               | 3             |               |               | 5             |               | 2             | 1             | 1             |               |               |
| <i>Polygordius</i>                |            | P1062 | 129472   | Schneider, 1868          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Galathowenia oculata</i>       |            | P1093 | 146950   | Zaks, 1922               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Owenia</i>                     |            | P1097 | 129427   | Delle Chiaje, 1844       |               | 1             | 3             | 3             |               | 3             |               | 3             |               |               |               |               |               |
| <i>Lagis koreni</i>               |            | P1107 | 152367   | Malmgren, 1866           | 1             |               | 2             | 5             |               | 2             | 2             | 5             | 3             | 2             | 1             | 10            |               |
| <i>Sabellaria spinulosa</i>       |            | P1117 | 130867   | Leuckart, 1849           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ampharete lindstroemi</i>      | aggregate  | P1139 | 129778   | M. Sars, 1864            | 6             | 7             | 1             | 7             |               |               | 4             | 26            |               |               |               | 6             |               |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                                     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |   | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <i>Anobothrus gracilis</i>        |            | P1147 | 129789   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Terebellides stroemii</i>      |            | P1175 | 131573   | Sars, 1835                                    |               | 1             |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Lanice conchilega</i>          |            | P1195 | 131495   | (Pallas, 1766)                                |               | 2             |               |               |               |               | 1             | 1             |               |               |               |               |               |
| <i>Phisidia aurea</i>             |            | P1215 | 131513   | Southward, 1956                               |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pista mediterranea</i>         |            |       | 131519   | de Gaillande, 1970                            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pista bansei</i>               |            |       | 152254   | Saphronova, 1988                              | 2             | 1             | 1             |               |               | 1             |               | 1             |               |               |               | 1             |               |
| <i>Polycirrus</i>                 |            | P1235 | 129710   | Grube, 1850                                   | 1             |               |               | 7             | 1             | 5             |               | 2             |               |               | 1             | 1             |               |
| <i>Lysilla loveni</i>             |            | P1233 | 131500   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Lysilla nivea</i>              |            | P1234 | 131501   | Langerhans, 1884                              | 1             | 2             |               |               |               | 1             |               | 2             |               |               |               |               | 1             |
| <i>Streblosoma intestinale</i>    |            | P1252 | 131540   | M. Sars in G.O. Sars, 1872                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thelepus cincinnatus</i>       |            | P1254 | 131543   | (Fabricius, 1780)                             | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dialychone dunerificta</i>     |            |       | 558752   | (Tovar-Hernández, Licciano, Giangrande, 2007) | 1             | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Parasabella cambrensis</i>     |            | P1273 | 530920   | Knight-Jones & Walker, 1985                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Euchone pararosea</i>          |            |       | 390407   | Giangrande & Licciano, 2006                   | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudopotamilla reniformis</i> |            | P1316 | 130963   | (Bruguiere, 1789)                             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Sabella pavonina</i>           |            | P1320 | 130967   | Savigny, 1822                                 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hydroides norvegica</i>        |            | P1334 | 131009   | Gunnerus, 1768                                |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Spirobranchus triqueter</i>    |            | P1341 | 555935   | (Linnaeus, 1758)                              | 6             | 2             |               |               |               |               |               |               |               |               |               |               |               |
| <b>OLIGOCHAETA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Grania</i>                     |            | P1524 | 137349   | Southern, 1913                                |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>CHELICERATA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nymphon brevirostre</i>        |            | Q5    | 150520   | Hodge, 1863                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Anoplodactylus petiolatus</i>  |            | Q44   | 134723   | (Kroyer, 1884)                                |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |      |          |                                   | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA |
| <b>CRUSTACEA</b>               |            |      |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Scalpellum scalpellum</i>   |            | R22  | 106204   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Verruca stroemia</i>        |            | R41  | 106257   | O.F.Muller, 1776                  |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia reboredae</i>       |            |      | 459311   | Moreira & Urgorri, 2009           |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia borealis</i>        |            | S7   | 156257   | Dahl, 1985                        |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Sarsinebalia urgorrii</i>   |            |      | 388224   | Moreira, Gestoso & Troncoso, 2003 |               | 1             |               |               |               |               |               |               | 1             |               |               |               |
| <i>Heteromysis formosa</i>     |            | S93  | 120037   | (G. O. Sars, 1877)                |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Kroyera carinata</i>        |            | S125 | 547074   | Spence Bate, 1857                 |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Periculodes longimanus</i>  |            | S131 | 102915   | (Bate & Westwood, 1868)           |               |               |               |               |               | 2             |               |               |               |               |               |               |
| <i>Pontocrates arenarius</i>   |            | S135 | 102918   | (Bate, 1858)                      |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Synchelidium maculatum</i>  |            | S138 | 102928   | Stebbing, 1906                    |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Apolochus neapolitanus</i>  |            | S159 | 236495   | (Della Valle, 1893)               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leucothoe incisa</i>        |            | S177 | 102460   | Robertson, 1892                   |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leucothoe procera</i>       |            | S179 | 102466   | Bate, 1857                        |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Stenothoe marina</i>        |            | S213 | 103166   | (Bate, 1856)                      |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Urothoe elegans</i>         |            | S248 | 103228   | (Bate, 1856)                      |               | 1             |               | 3             |               |               | 7             | 2             | 2             |               |               |               |
| <i>Urothoe marina</i>          |            | S249 | 103233   | (Bate, 1857)                      | 17            | 36            | 4             | 5             |               | 5             |               | 8             | 2             |               | 10            |               |
| <i>Acidostoma neglectum</i>    |            |      | 102495   | (Spence Bate & Westwood, 1861)    |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Hippomedon denticulatus</i> |            | S296 | 102570   | (Bate, 1857)                      |               |               |               | 1             |               |               |               |               | 1             |               | 1             |               |
| <i>Tryphosa nana</i>           |            | S321 | 102691   | (Kroyer, 1846)                    |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tmetonyx similis</i>        |            | S337 | 102742   | (G O Sars, 1891)                  |               | 1             |               |               |               |               |               |               |               |               |               |               |
| <i>Nototropis falcatus</i>     |            | S410 | 102139   | Metzger, 1871                     |               |               |               |               |               | 2             |               |               | 1             |               | 1             |               |
| <i>Nototropis vedlomensis</i>  |            | S413 | 102132   | (Bate & Westwood, 1862)           | 1             | 1             |               |               |               |               |               |               | 1             |               | 1             |               |

| Taxon                              | Qualifiers | SDC         | APHIA ID      | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|-------------|---------------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |             |               |                                   | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <i>Ampelisca diadema</i>           |            | S429        | 101896        | (Costa, 1853)                     | 1             | 6             |               | 1             |               |               | 3             | 4             | 1             |               |               |               |               |
| <i>Ampelisca provincialis</i>      |            | S434        | 101915        | Bellan-Santini & Kaim-Malka, 1977 | 2             | 2             |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca spinipes</i>          |            | S438        | 101928        | Boeck, 1861                       |               | 2             |               | 2             |               |               | 4             | 5             |               |               |               | 2             |               |
| <i>Ampelisca tenuicornis</i>       |            | S440        | 101930        | Lilljeborg, 1855                  |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Ampelisca typica</i>            |            | S442        | 101933        | (Bate, 1856)                      | 1             | 2             |               | 4             | 1             |               | 3             |               |               |               |               | 1             |               |
| <i>Bathyporeia elegans</i>         |            | S452        | 103058        | Watkin, 1938                      |               |               |               |               | 1             | 4             |               |               | 2             | 11            | 8             | 1             | 2             |
| <i>Bathyporeia gracilis</i>        |            | S453        | 103059        | G O Sars, 1891                    |               |               | 3             | 1             | 2             | 13            |               |               | 2             | 5             | 2             |               |               |
| <i>Bathyporeia guilliamsoniana</i> |            | S454        | 103060        | (Bate, 1857)                      |               |               |               |               |               | 3             |               |               |               | 2             | 4             |               |               |
| <i>Haustorius arenarius</i>        |            | S462        | 102317        | Slabber, 1769                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abludomelita obtusata</i>       |            | S498        | 102788        | (Montagu, 1813)                   |               |               | 1             |               | 2             |               |               |               |               |               | 6             |               |               |
| <b>Cheirocratus</b>                |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b>               |               |               |               |               |               |               | <b>1</b>      |               |               |               |               |               |               |
| <i>Othomaera othonis</i>           |            | S519        | 534781        | (Milne-Edwards, 1830)             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Maerella tenuimana</i>          |            | S521        | 102831        | (Bate, 1862)                      | 1             | 2             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Megamphopus cornutus</i>        |            | S539        | 148545        | Norman, 1869                      |               | 1             |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Gammaropsis maculata</i>        |            | S541        | 102364        | (Johnston, 1828)                  | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Photis longicaudata</i>         |            | S552        | 102383        | (Bate & Westwood, 1862)           | 1             | 2             |               |               |               |               |               |               |               |               |               |               |               |
| <b>Aoridae</b>                     |            | <b>S577</b> | <b>101368</b> | <b>Stebbing, 1899</b>             | <b>1</b>      | <b>1</b>      |               | <b>2</b>      | <b>1</b>      | <b>1</b>      | <b>2</b>      |               |               | <b>1</b>      |               |               |               |
| <i>Leptocheirus hirsutimanus</i>   |            | S588        | 102036        | (Bate, 1862)                      |               |               |               | 1             |               |               |               | 4             |               |               |               | 3             |               |
| <i>Crassicorophium crassicorne</i> |            | S611        | 397383        | Bruzelius, 1859                   |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Unciola planipes</i>            |            | S622        | 102061        | Norman, 1867                      |               |               |               | 1             | 1             | 2             |               | 2             |               | 2             |               |               |               |
| <i>Phtisica marina</i>             |            | S657        | 101864        | Slabber, 1769                     |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <b>Gnathia</b>                     |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>                |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Conilera cylindracea</i>        |            | S849        | 118842        | (Montagu, 1804)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Eurydice pulchra</i>            |            | S854        | 118852        | Leach, 1815                       | 3             | 3             |               |               |               |               | 4             |               |               |               |               | 2             | 1             |



| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |   |
|--------------------------------|------------|-------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---|
|                                |            |       |          |                                   | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |   |
| <i>Astacilla longicornis</i>   |            | S955  | 119024   | (Sowerby, 1806)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Tanaopsis graciloides</i>   |            | S1142 | 136458   | (Lilljeborg, 1864)                |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Bodotria scorpioides</i>    |            | S1197 | 110445   | (Montagu, 1804)                   |               |               |               |               |               |               |               | 1             |               |               |               |               |               |   |
| <i>Diastylis bradyi</i>        |            | S1248 | 110472   | Norman, 1879                      |               |               |               |               |               | 1             |               |               |               | 2             |               |               |               |   |
| <i>Processa modica</i>         |            | S1366 | 108343   | Williamson & Rochanaburanon, 1979 |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Callianassa subterranea</i> |            | S1415 | 107729   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Upogebia deltaura</i>       |            | S1419 | 107739   | (Leach, 1815)                     |               | 3             |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Pisidia longicornis</i>     |            | S1482 | 107188   | (Linnaeus, 1757)                  | 4             |               |               | 1             |               |               |               |               |               |               |               |               |               |   |
| <i>Ebalia tumefacta</i>        |            | S1509 | 107302   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Hyas coarctatus</i>         |            | S1519 | 107323   | Leach, 1815                       |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Inachus dorsettensis</i>    |            | S1526 | 107327   | (Pennant, 1777)                   |               |               |               |               |               |               | 1             |               |               |               |               |               |               |   |
| <i>Macropodia rostrata</i>     |            | S1532 | 107345   | (Linnaeus, 1761)                  | 1             |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Thia scutellata</i>         |            | S1559 | 107281   | (Fabricius, 1793)                 |               |               |               |               |               |               |               |               |               |               |               |               | 1             | 1 |
| <i>Liocarcinus pusillus</i>    |            | S1584 | 107393   | (Leach, 1816)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Pinnotheres pisum</i>       |            | S1638 | 107473   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <b>MOLLUSCA</b>                |            |       |          |                                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Leptochiton asellus</i>     |            | W53   | 140199   | (Gmelin, 1791)                    |               |               |               |               |               |               | 1             | 1             |               |               |               |               |               |   |
| <i>Acanthochitona crinita</i>  |            | W86   | 138675   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Gibbula magus</i>           | ?          | W159  | 141790   | (Linnaeus, 1758)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Euspira nitida</i>          |            | W491  | 151894   | (Donovan, 1804)                   |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |
| <i>Melanella alba</i>          |            | W634  | 139832   | (da Costa, 1778)                  | 1             |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Colus gracilis</i>          |            | W715  | 138899   | (da Costa, 1778)                  | 1             |               |               |               |               |               |               |               |               |               |               |               |               |   |
| <i>Philine</i>                 |            | W1036 | 138339   | Ascanius, 1772                    |               |               |               |               |               |               | 1             |               |               |               |               |               |               |   |
| <i>Tritonia plebeia</i>        |            | W1254 | 141738   | Johnston, 1828                    | 1             |               |               |               |               |               |               |               |               |               |               |               |               |   |
| Onchidorididae                 |            | W1319 | 175      | Gray, 1827                        |               |               |               |               |               |               |               |               | 1             |               |               |               |               |   |

| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |  |
|--------------------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                |            |       |          |                         | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |  |
| <i>Knoutsodonta depressa</i>   |            | W1323 | 845528   | (Alder & Hancock, 1842) |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nucula hanleyi</i>          |            | W1568 | 140588   | Winckworth, 1931        |               |               |               |               |               |               |               |               |               |               | 1             |               | 1             |  |
| <i>Modiolus adriaticus</i>     |            | W1700 | 506025   | (Lamarck, 1819)         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aequipecten opercularis</i> |            | W1773 | 140687   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Heteranomia squamula</i>    |            | W1809 | 138749   | (Linnaeus, 1758)        | 1             |               |               |               |               |               | 1             |               |               |               |               |               |               |  |
| <i>Pododesmus squama</i>       |            | W1812 | 138752   | (Linnaeus, 1761)        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Hemilepton nitidum</i>      |            | W1882 | 246148   | (W. Turton, 1822)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Kurtiella bidentata</i>     |            | W1906 | 345281   | (Montagu, 1803)         | 1             |               |               |               |               |               |               | 15            |               |               |               |               | 1             |  |
| <i>Spisula elliptica</i>       |            | W1975 | 140300   | (Brown, 1827)           |               |               |               |               |               |               | 2             |               |               |               |               | 2             | 1             |  |
| <i>Ensis leei</i>              |            | W1997 | 876640   | M. Huber, 2015          |               |               |               |               |               |               | 1             |               |               |               | 1             |               |               |  |
| <i>Phaxas pellucidus</i>       |            | W2006 | 140737   | (Pennant, 1777)         |               |               |               | 1             |               |               |               | 2             |               |               |               |               |               |  |
| <i>Moerella donacina</i>       |            | W2021 | 147021   | Linnaeus, 1758          |               |               |               |               |               |               |               | 1             |               |               |               |               |               |  |
| <i>Asbjornsenia pygmaea</i>    |            | W2023 | 879714   | (Lovén, 1846)           |               |               |               |               | 1             | 2             |               |               | 1             | 1             |               |               |               |  |
| <i>Gari fervensis</i>          |            | W2051 | 140870   | (Gmelin, 1791)          |               |               | 1             |               |               |               |               | 1             |               |               |               |               | 1             |  |
| <i>Solecurtus scopula</i>      |            | W2054 | 141543   | (Turton, 1822)          |               |               |               |               |               |               |               |               |               |               |               |               | 1             |  |
| <i>Abra alba</i>               |            | W2059 | 141433   | (W Wood, 1802)          |               |               |               |               |               |               |               |               |               |               | 1             |               |               |  |
| <i>Abra prismatica</i>         |            | W2062 | 141436   | (Montagu, 1808)         |               |               |               |               | 1             | 1             |               |               |               |               |               |               |               |  |
| <i>Clausinella fasciata</i>    |            | W2100 | 141909   | (da Costa, 1778)        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Timoclea ovata</i>          |            | W2104 | 141929   | (Pennant, 1777)         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Venerupis corrugata</i>     |            | W2124 | 181364   | (Gmelin, 1791)          |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Dosinia exoleta</i>         |            | W2130 | 141911   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Hiatella arctica</i>        |            | W2166 | 140103   | (Linnaeus, 1758)        | 1             |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Thracia villosiuscula</i>   |            | W2233 | 141651   | (Macgillivray, 1827)    |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Cochlodesma praetenue</i>   |            | W2239 | 181373   | (Pulteney, 1799)        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <b>PHORONIDA</b>               |            |       |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |

| Taxon                                       | Qualifiers | SDC   | APHIA ID | Authority                           | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---|------------|-------|----------|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|   |            |       |          |                                     | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <i>Phoronis</i>                             |            | ZA3   | 128545   | Wright, 1856                        | 10            | 10            | 4             |               | 1             |               |               | 2             | 1             |               |               | 2             |               |
| <b>ECHINODERMATA</b>                        |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ophiothrix fragilis</i>                  |            | ZB124 | 125131   | (Abildgaard, 1789)                  | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Amphipholis squamata</i>                 |            | ZB161 | 125064   | (Chiaje, 1828)                      |               |               |               |               |               |               |               | 1             |               |               |               | 1             |               |
| <i>Ophiura albida</i>                       |            | ZB168 | 124913   | Forbes, 1839                        |               | 1             |               |               |               |               | 1             |               |               |               |               | 1             |               |
| <i>Psammechinus miliaris</i>                |            | ZB193 | 124319   | (Gmelin, 1778)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Echinocyamus pusillus</i>                |            | ZB212 | 124273   | (O.F. Muller, 1776)                 |               |               |               |               | 1             | 9             | 3             | 3             | 2             | 3             |               | 1             |               |
| <i>Echinocardium cordatum</i>               |            | ZB223 | 124392   | (Pennant, 1777)                     |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Mesothuria intestinalis</i>              |            | ZB233 | 124568   | (Ascanius, 1805)<br>Östergren, 1896 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudothyone raphanus</i>                |            | ZB257 | 124661   | (Duben & Koren, 1845)               |               | 1             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone fusus</i>                         |            | ZB262 | 124670   | O.F.Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone roscovita</i>                     |            | ZB264 | 124676   | Hérouard, 1889                      | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Panningia hyndmani</i>                   |            | ZB272 | 848014   | (Thompson, 1840)                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Leptosynapta inhaerens</i>               |            | ZB296 | 124465   | (O.F.Muller, 1776)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Oestergrenia digitata</i>                |            | ZB300 | 152547   | (Montagu, 1804)                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>TUNICATA</b>                             |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidella aspersa</i>                    |            | ZD84  | 103718   | O F Müller, 1776                    |               |               |               |               |               |               |               |               |               |               |               | 1             |               |
| <i>Ascidia conchilega</i>                   | ?          | ZD88  | 103702   | O F Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |               | 9             |               |
| <i>Polycarpa fibrosa</i>                    |            | ZD112 | 103902   | (Stimpson, 1852)                    | 1             | 1             |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Dendrodoa grossularia</i>                |            | ZD120 | 103882   | (van Beneden 1846)                  |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>CEPHALOCHORDATA</b>                      |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Branchiostoma lanceolatum</i>            |            |       | 104906   | (Pallas, 1774)                      |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                       |            |       |          |                                     | 61            | 52            | 25            | 33            | 18            | 34            | 40            | 48            | 26            | 24            | 19            | 43            | 14            |
| <b>Abundance</b>                            |            |       |          |                                     | 157           | 164           | 48            | 88            | 32            | 88            | 163           | 130           | 49            | 53            | 40            | 93            | 17            |
| The following taxa were merged for analysis |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon   | Qualifiers | SDC  | APHIA ID | Authority          | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---|------------|------|----------|--------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|   |            |      |          |                    | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| <b>Aoridae</b>  |            | S577 | 101368   | Stebbing, 1899     | 1             | 1             |               | 2             | 1             | 1             | 2             |               |               | 1             |               |               |               |
| <i>Aoridae</i>  | female     | S577 | 101368   | Stebbing, 1899     | 1             |               |               | 2             | 1             |               | 1             |               |               |               |               |               |               |
| <i>Autonoe longipes</i>                               |            | S583 | 102021   | (Lilljeborg, 1852) |               | 1             |               |               |               | 1             | 1             |               |               | 1             |               |               |               |
| <b>Gnathia</b>  |            | S793 | 118437   | Leach, 1814        |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Gnathia</i>  | female     | S793 | 118437   | Leach, 1814        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gnathia oxyuraea</i>                               |            | S796 | 118995   | (Lilljeborg, 1855) |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <b>Cheirocratus</b>                                   |            | S503 | 101669   | Norman, 1867       |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Cheirocratus</i>                                   | female     | S503 | 101669   | Norman, 1867       |               |               |               |               |               |               | 1             |               |               |               |               |               |               |
| <i>Cheirocratus pseudosundevallii</i>                 | ?          |      |          | Gouillieux, 2019   |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                                 |            |      |          |                    | 1             | 1             | 0             | 1             | 1             | 1             | 2             | 1             | 0             | 1             | 0             | 0             | 0             |
| <b>Abundance</b>                                      |            |      |          |                    | 1             | 1             | 0             | 2             | 1             | 1             | 3             | 1             | 0             | 1             | 0             | 0             | 0             |
| <b>The following taxa were excluded from analysis</b> |            |      |          |                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Colonial</b>                                       |            |      |          |                    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| PORIFERA  |            | C1   | 558      | Grant, 1836        | P             | P             |               |               |               |               |               | P             |               |               |               |               |               |
| <i>Leuckartiara octona</i>                            |            | D240 | 117791   | (Fleming, 1823)    | P             |               |               |               |               |               |               | P             |               |               |               |               |               |
| <i>Tubularia</i>                                      |            | D163 | 117258   | Linnaeus, 1758     |               |               | P             |               |               |               | P             | P             |               | P             |               |               |               |
| <i>Eudendrium</i>                                     |            | D218 | 117093   | Ehrenberg, 1834    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Bougainvilliidae                                      |            | D246 | 1594     | Lütken, 1850       |               |               | P             | P             |               |               |               |               |               |               | P             | P             |               |
| <i>Phialella quadrata</i>                             |            | D343 | 117804   | (Forbes, 1848)     |               |               |               |               | P             |               |               | P             |               |               | P             |               |               |
| <i>Calycella syringa</i>                              |            | D348 | 117402   | (Linnaeus, 1767)   | P             |               |               |               |               |               |               | P             |               |               |               |               |               |
| <i>Lafoea dumosa</i>                                  |            | D386 | 117702   | (Fleming, 1828)    | P             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Halecium</i>                                       |            | D390 | 117103   | Oken, 1815         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abietinaria</i>                                    |            | D408 | 117225   | Kirchenpauer, 1884 |               | P             |               |               |               |               |               |               |               |               |               |               |               |
| <i>Diphasia</i>                                       |            | D413 | 117228   | L Agassiz, 1862    |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Hydrallmania falcata</i>                           |            | D424 | 117890   | (Linnaeus, 1758)   | P             | P             |               |               |               |               |               |               | P             |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |  |
|--------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                |            |      |          |                       | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |  |
| <i>Sertularella gayi</i>       |            | D429 | 117902   | (Lamouroux, 1821)     | P             | P             |               |               |               |               |               |               |               |               |               |               | P             |  |
| <i>Sertularia cupressina</i>   |            | D435 | 117913   | (Linnaeus, 1758)      | P             | P             |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nemertesia</i>              |            | D462 | 117195   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| Campanulariidae                |            | D491 | 1606     | Peron & Lesueur, 1810 | P             | P             | P             | P             | P             |               | P             |               |               |               | P             | P             |               |  |
| <i>Alcyonium digitatum</i>     |            | D597 | 125333   | Linnaeus, 1758        | P             |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Crisia</i>                  |            | Y13  | 111032   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Tubulipora</i>              |            | Y27  | 111054   | Lamarck, 1816         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Disporella hispida</i>      |            | Y66  | 111730   | (Fleming, 1828)       | P             |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Alcyonidium parasiticum</i> |            | Y81  | 111604   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Vesicularia spinosa</i>     |            | Y131 | 111669   | (Linnaeus, 1758)      |               |               |               | P             |               |               |               |               |               |               |               |               |               |  |
| <i>Amathia lendigera</i>       |            | Y135 | 111659   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Amathia</i>                 |            | Y137 | 111023   | Farre, 1837           |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aetea</i>                   |            | Y153 | 110819   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Conopeum reticulum</i>      |            | Y172 | 111351   | (Linnaeus, 1767)      |               |               |               | P             | P             |               |               |               |               |               |               |               |               |  |
| <i>Electra pilosa</i>          |            | Y178 | 111355   | (Linnaeus, 1767)      |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Callopora</i>               |            | Y201 | 110851   | J E Gray, 1848        |               | P             |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Beania mirabilis</i>        |            | Y261 | 111072   | Johnston, 1840        |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Chorizopora brongiartii</i> |            | Y344 | 111304   | (Audouin, 1826)       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Escharella immersa</i>      |            | Y364 | 111484   | (Fleming, 1828)       | P             | P             |               |               |               |               |               | P             |               |               |               |               |               |  |
| <i>Porella concinna</i>        |            | Y385 | 111125   | (Busk, 1854)          |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Schizomavella</i>           |            | Y467 | 110829   | Canu & Bassler, 1917  | P             |               |               |               |               |               |               | P             |               |               |               |               |               |  |
| <b>Damaged</b>                 |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Eumida</i>                  | damaged    | P163 | 129446   | Malmgren, 1865        |               | 1             |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aricidea</i>                | damaged    | P675 | 129430   | Webster, 1879         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Paradoneis</i>              | damaged    | P695 | 129433   | Hartman, 1965         |               |               |               |               |               |               |               |               |               |               |               |               |               |  |

| Taxon               | Qualifiers | SDC    | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---------------------|------------|--------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                     |            |        |          |                         | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| Spionidae           | damaged    | P720   | 913      | Grube,1850              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Dipolydora</i>   | damaged    | P748   | 129611   | Verrill, 1881           | 1             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Terebellidae</i> | damaged    | P1179  | 982      | Johnston, 1846          |               | 1             |               | 1             |               |               |               |               |               | 1             |               | 1             |               |
| Sabellidae          | damaged    | P1257  | 985      | Latreille, 1825         |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| Serpulidae          | damaged    | P1324  | 988      | Rafinesque, 1815        | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia</i>      | damaged    | S5     | 147031   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| AMPHIPODA           | damaged    | S97    | 1135     |                         |               |               | 1             |               |               |               |               |               |               |               | 1             |               |               |
| <i>Urothoe</i>      | damaged    | S246   | 101789   | Dana, 1852              |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Liocarcinus</i>  | damaged    | S1577  | 106925   | Stimpson,1870           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| GASTROPODA          | damaged    | W88    | 101      |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| BIVALVIA            | damaged    | W1560  | 151265   | Goldfuss, 1820          |               |               |               |               |               |               |               |               |               |               |               |               |               |
| SOLENOIDEA          | damaged    | W1991  | 14635    | Lamarck, 1809           |               | 1             |               |               |               |               |               | 1             |               |               |               |               |               |
| Ophiuridae          | damaged    | ZB165  | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Fish</b>         |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Gobiesocidae        |            | ZG1180 | 125477   | Bleeker, 1859           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Juvenile</b>     |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| ANIMALIA            | eggs       |        | 2        |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| SIPUNCULA           | juvenile   | N1     | 1268     |                         | 1             |               |               |               |               |               | 2             | 1             |               |               |               |               |               |
| <i>Golfingia</i>    | juvenile   | N12    | 136021   | Lankester, 1885         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera</i>      | juvenile   | P255   | 129296   | Lamarck, 1818           |               |               |               |               | 1             |               |               |               |               |               |               |               |               |
| Goniadidae          | juvenile   | P266   | 953      | Kinberg, 1866           |               |               |               |               |               |               |               | 2             |               |               |               |               |               |
| Nereididae          | juvenile   | P458   | 22496    | Blainville, 1818        |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| <i>Nephtys</i>      | juvenile   | P494   | 129370   | Cuvier, 1817            |               |               |               | 1             |               |               |               | 2             |               | 1             |               |               |               |
| <i>Scoletepis</i>   | juvenile   | P778   | 129623   | Blainville, 1828        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| CRUSTACEA           | larva      | R1     | 1066     |                         |               |               |               |               |               | 1             |               |               | 1             | 1             | 1             | 1             |               |



| Taxon                | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|----------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                      |            |       |          |                         | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| BALANOMORPHA         | juvenile   | R42   | 1082     | Burmeister, 1834        |               |               |               |               | 1             | 1             |               |               |               |               |               |               |               |
| <i>Ampelisca</i>     | juvenile   | S423  | 101445   | Kroyer, 1842            |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Gnathia</i>       | juvenile   | S793  | 118437   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| GEBIIDEA             | juvenile   | S1403 | 477323   | de Saint Laurent, 1979  | 1             | 1             |               |               |               |               |               |               |               |               |               | 1             |               |
| Paguridae            | juvenile   | S1445 | 106738   | Latreille, 1802         | 1             |               |               |               |               |               | 1             |               |               |               |               |               |               |
| DECAPODA             | larva      | S1276 | 1130     | Latreille, 1830         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Ebalia</i>        | juvenile   | S1504 | 106889   | Leach, 1817             |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Majidae              | juvenile   | S1512 | 106760   | Samouelle, 1819         | 1             |               |               | 1             |               |               |               |               |               |               |               |               |               |
| Inachinae            | juvenile   | S1520 | 148436   | MacLeay, 1838           |               |               |               |               |               |               |               |               |               |               | 1             |               |               |
| <i>Macropodia</i>    | juvenile   | S1529 | 205077   | Leach, 1814             |               |               |               | 1             |               |               |               |               |               |               |               | 1             |               |
| Buccinidae           | eggs       | W702  | 149      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Tritonia</i>      | juvenile   | W1246 | 138580   | Cuvier, 1798            | 2             |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Nucula</i>        | juvenile   | W1565 | 138262   | Lamarck, 1799           |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Mytilidae            | juvenile   | W1691 | 211      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Modiolus</i>      | juvenile   | W1702 | 140467   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Pectinidae           | juvenile   | W1768 | 213      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| Anomiidae            | juvenile   | W1805 | 214      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Abra</i>          | juvenile   | W2058 | 138474   | Lamarck, 1818           |               |               |               | 1             |               |               |               |               |               |               |               |               |               |
| <i>Dosinia</i>       | juvenile   | W2126 | 138636   | Scopoli, 1777           |               |               |               | 2             |               | 1             |               | 1             |               | 1             |               |               |               |
| Ophiuridae           | juvenile   | ZB165 | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               | 1             |               |               |               |               |               |
| ECHINOIDEA           | juvenile   | ZB181 | 123082   | Leske, 1778             |               |               |               |               |               |               |               | 2             |               |               |               | 1             |               |
| <i>Echinocardium</i> | juvenile   | ZB222 | 123426   | Gray, 1825              |               |               |               |               | 1             |               |               |               | 1             |               | 1             |               |               |
| HOLOTHUROIDEA        | juvenile   | ZB229 | 123083   |                         |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone</i>        | juvenile   | ZB261 | 146116   | Oken, 1815              |               |               |               |               |               | 1             |               | 2             |               |               | 1             | 2             |               |
| <i>Leptosynapta</i>  | juvenile   | ZB291 | 123449   | Verrill, 1867           |               |               |               |               |               |               |               |               |               |               |               |               |               |

| Taxon                 | Qualifiers | SDC  | APHIA ID | Authority     | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------|------------|------|----------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                       |            |      |          |               | MA_ST42<br>FA | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| ASCIDIACEA            | juvenile   | ZD2  | 1839     | Nielsen, 1995 |               |               |               |               |               |               |               |               | 1             |               |               |               |               |
| <b>Meiofaunal</b>     |            |      |          |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| NEMATODA              |            | HD1  | 799      |               |               | 1             |               |               |               |               |               |               | 1             |               |               |               |               |
| <b>Parasitic</b>      |            |      |          |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| COPEPODA              | parasitic  | R142 | 1080     |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b> |            |      |          |               | 19            | 13            | 3             | 11            | 6             | 4             | 7             | 15            | 4             | 6             | 8             | 8             | 0             |
| <b>Abundance</b>      |            |      |          |               | 9             | 5             | 1             | 7             | 3             | 4             | 3             | 14            | 4             | 4             | 5             | 7             | 0             |

| Taxon                         | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |
|-------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                               |            |      |          |                       | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <b>CNIDARIA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |
| <i>Cerianthus lloydii</i>     |            | D632 | 283798   | Gosse, 1859           |               |               |               |               |               | 8             |               | 1             |               |               |
| Actiniaria                    |            | D662 | 1360     |                       |               |               |               |               |               |               |               |               | 3             |               |
| <b>PLATYHELMINTHES</b>        |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |
| PLATYHELMINTHES               |            | F2   | 793      | Minot, 1876           |               |               |               |               |               |               |               |               |               |               |
| <b>NEMERTEA</b>               |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |
| NEMERTEA                      |            | G1   | 152391   |                       | 1             | 1             | 3             |               |               | 7             | 5             | 7             | 2             |               |
| <b>SIPUNCULA</b>              |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |
| <i>Golfingia elongata</i>     |            | N14  | 175026   | (Keferstein, 1862)    |               |               | 1             |               |               | 6             |               |               |               |               |
| <i>Golfingia vulgaris</i>     |            | N17  | 136050   | (de Blainville, 1827) |               |               |               |               |               | 2             |               |               |               |               |
| <i>Nephasoma minutum</i>      |            | N25  | 136060   | (Keferstein, 1862)    |               |               |               |               |               |               |               |               |               |               |
| <i>Phascolion strombus</i>    |            | N34  | 175043   | (Montagu, 1804)       |               |               |               | 1             |               | 1             |               |               |               |               |
| <b>POLYCHAETA</b>             |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |
| <i>Pisione remota</i>         |            | P15  | 130707   | Southern, 1914        | 1             |               |               |               |               |               | 24            | 20            |               |               |
| <i>Aphrodita aculeata</i>     |            | P19  | 129840   | Malmgren, 1867        |               |               |               |               |               |               |               |               | 1             |               |
| <i>Enipo elisabethae</i>      |            | P43  | 130737   | McIntosh, 1900        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Gattyana cirrhosa</i>      |            | P49  | 130749   | (Pallas, 1766)        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Harmothoe</i>              |            | P50  | 129491   | Kinberg, 1855         |               |               | 1             |               |               | 1             |               |               |               |               |
| <i>Malmgrenia darbouxi</i>    |            |      | 863197   | (Pettibone, 1993)     |               |               |               |               |               | 6             |               |               |               |               |
| <i>Malmgrenia andreapolis</i> |            | P51  | 147008   | (McIntoch, 1874)      |               |               |               |               |               |               |               |               |               |               |
| <i>Lepidonotus squamatus</i>  |            | P82  | 130801   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe inornata</i>        |            | P92  | 130601   | Johnston, 1839        |               |               |               |               |               |               |               |               |               |               |
| <i>Pholoe baltica</i>         |            | P95  | 130599   | Örsted, 1843          |               |               |               |               |               | 2             |               |               |               |               |
| <i>Sthenelais limicola</i>    |            | P109 | 131077   | (Ehlers, 1864)        |               |               |               |               |               |               |               |               |               |               |
| <i>Eteone longa</i>           | aggregate  | P117 | 130616   | (Fabricius, 1780)     |               |               |               |               |               | 1             |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |      |          |                          | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Hesionura elongata</i>      |            | P122 | 130649   | (Southern, 1914)         |               |               |               |               |               |               | 12            | 1             |               |               |
| <i>Mysta picta</i>             |            | P127 | 147026   | (Quatrefages, 1866)      |               |               |               |               | 1             |               |               |               |               |               |
| <i>Phyllodoce groenlandica</i> |            | P141 | 334506   | (Örsted, 1842)           |               |               | 1             |               |               | 1             |               |               |               |               |
| <i>Phyllodoce longipes</i>     |            | P143 | 130763   | Kinberg, 1866            |               |               |               |               |               |               |               |               |               |               |
| <i>Phyllodoce rosea</i>        |            | P146 | 334514   | (Mcintosh, 1877)         |               |               |               |               |               |               |               |               |               |               |
| <i>Eulalia bilineata</i>       |            | P152 | 130624   | (Johnston, 1840)         |               |               |               |               |               |               |               |               |               |               |
| <i>Eulalia mustela</i>         |            | P155 | 130631   | Pleijel, 1987            |               |               | 1             |               |               |               |               |               |               |               |
| <i>Eumida sanguinea</i>        |            | P167 | 130644   | (Örsted, 1843)           |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera alba</i>            |            | P256 | 130116   | (O.F. Muller, 1788)      |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera fallax</i>          |            | P259 | 336908   | Quatrefages, 1850        |               |               |               |               |               |               |               |               |               |               |
| <i>Glycera lapidum</i>         |            | P260 | 130123   | Quatrefages, 1866        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Glycera oxycephala</i>      |            | P262 | 130126   | Ehlers, 1887             |               |               |               |               |               |               |               |               | 3             |               |
| <i>Glycinde nordmanni</i>      |            | P268 | 130136   | (Malmgren, 1865)         |               |               |               |               |               |               |               |               |               |               |
| <i>Goniadella gracilis</i>     |            | P276 | 130145   | (Verrill, 1873)          |               |               | 4             |               |               |               | 2             |               |               |               |
| <i>Psamathe fusca</i>          |            | P305 | 152249   | (Keferstein, 1862)       |               |               |               |               |               |               |               |               |               |               |
| <i>Oxydromus pallidus</i>      |            | P317 | 340203   | (Claparede, 1864)        |               |               | 1             |               |               |               |               |               |               |               |
| <i>Podarkeopsis capensis</i>   |            | P319 | 130195   | Day, 1963                |               |               |               |               |               | 1             |               | 1             |               |               |
| <i>Syllis garciai</i>          |            | P351 | 131431   | (Campoy, 1982)           |               |               | 1             |               |               |               |               |               |               |               |
| <i>Syllis parapari</i>         |            |      | 196002   | San Martin & Lopez, 2000 |               |               |               | 1             |               |               |               |               |               |               |
| <i>Syllis pontxioi</i>         |            |      | 196003   | San Martin & López, 2000 |               |               |               |               |               |               |               |               |               |               |
| <i>Syllis armillaris</i>       |            | P365 | 131415   | (O.F. Muller, 1776)      |               |               |               |               |               |               |               |               |               |               |
| <i>Syllis variegata</i>        |            | P371 | 131458   | (Grube, 1860)            |               |               |               |               |               |               |               |               |               |               |
| <i>Eusyllis blomstrandii</i>   |            | P380 | 131290   | Malmgren, 1867           |               |               |               |               |               |               |               |               |               |               |

| Taxon                             | Qualifiers | SDC  | APHIA ID | Authority                       | Sample        |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|------|----------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |      |          |                                 | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Odontosyllis fulgurans</i>     |            | P387 | 131327   | (Audouin & Milne Edwards, 1833) |               |               |               |               |               |               |               |               |               |               |
| <i>Streptodonta pterochaeta</i>   |            | P391 | 238207   | Southern, 1914                  |               |               |               |               |               |               |               |               |               |               |
| <i>Streptosyllis campoyi</i>      |            |      | 238248   | Brito, Nunez & San Martin, 2000 |               |               |               |               |               |               |               |               |               |               |
| <i>Parexogone hebes</i>           |            | P421 | 757970   | (Webster & Benedict, 1884)      |               |               |               |               |               | 1             |               |               |               |               |
| <i>Exogone verugera</i>           |            | P423 | 131307   | (Claparede, 1868)               |               |               |               |               |               |               |               |               |               |               |
| <i>Sphaerosyllis</i>              |            | P424 | 129677   | Claparède, 1863                 |               |               |               |               |               |               |               |               |               |               |
| Myrianida                         |            | P434 | 129659   | Milne Edwards, 1845             |               |               |               |               |               |               |               |               |               |               |
| <i>Eunereis longissima</i>        |            | P475 | 130375   | Johnston, 1840                  |               |               |               |               |               | 3             |               |               |               |               |
| <i>Nereis zonata</i>              |            | P478 | 130407   | Malmgren, 1867                  |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys assimilis</i>          |            | P495 | 130353   | Ørsted, 1843                    |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys caeca</i>              |            | P496 | 130355   | (Fabricius, 1780)               |               |               |               |               |               |               |               |               |               |               |
| <i>Nephtys cirrosa</i>            |            | P498 | 130357   | Ehlers, 1868                    | 1             | 2             |               |               | 1             |               |               |               | 3             | 1             |
| <i>Nephtys longosetosa</i>        |            | P503 | 130364   | Ørsted, 1843                    |               |               |               |               | 1             |               |               |               |               |               |
| <i>Nothria conchylega</i>         |            | P545 | 130467   | (Sars, 1835)                    |               |               |               |               |               |               |               |               |               |               |
| <i>Paucibranchia bellii</i>       |            | P564 | 130072   | (Audouin & Milne-Edwards, 1833) |               |               | 1             |               |               |               |               |               |               |               |
| <i>Lysidice unicornis</i>         |            | P568 | 742232   | (Grube, 1840)                   |               |               |               |               |               |               |               | 1             |               |               |
| <i>Lumbrineris cf. cingulata</i>  |            |      | 130240   | (Ehlers, 1868)                  |               |               |               |               |               | 1             |               |               |               |               |
| <i>Lumbrineris futilis</i>        |            | P582 | 851788   | (Audouin & Milne Edwards, 1834) |               |               |               |               |               |               |               |               |               |               |
| <i>Drilonereis filum</i>          |            | P591 | 129856   | (Claparède, 1868)               |               |               |               |               |               |               |               |               |               |               |
| <i>Protodorvillea kefersteini</i> |            | P638 | 130041   | (McIntosh, 1869)                |               |               |               |               |               |               | 1             | 3             |               |               |
| <i>Schistomeringos rudolphi</i>   |            | P643 | 154127   | Delle Chiaje, 1828              |               |               |               |               |               | 2             |               |               |               |               |
| <i>Orbinia sertulata</i>          |            | P665 | 334310   | (Savigny, 1820)                 |               |               |               |               |               |               |               |               |               |               |

| Taxon                              | Qualifiers | SDC  | APHIA ID | Authority                     | Sample        |               |               |               |               |               |               |               |               |               |  |
|------------------------------------|------------|------|----------|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                    |            |      |          |                               | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |  |
| <i>Scoloplos armiger</i>           |            | P672 | 334722   | (Muller, 1776)                |               | 1             |               |               |               |               | 11            | 1             |               |               |  |
| <i>Aricidea catherinae</i>         |            | P684 | 333034   | (Laubier, 1967)               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aricidea cerrutii</i>           |            | P685 | 525497   | Laubier, 1966                 | 1             |               |               |               |               |               | 1             | 1             |               |               |  |
| <i>Cirrophorus branchiatus</i>     |            | P689 | 130576   | Ehlers, 1908                  |               |               |               |               |               |               |               |               |               |               |  |
| <i>Paradoneis ilvana</i>           |            | P698 |          | Castelli, 1985                |               |               |               |               |               |               |               |               |               |               |  |
| <i>Paradoneis lyra</i>             |            | P699 | 130585   | (Southern, 1914)              |               |               |               |               |               | 1             |               |               |               |               |  |
| <i>Poecilochaetus serpens</i>      |            | P718 | 130711   | Allen, 1904                   |               |               | 3             |               |               | 2             |               | 1             | 1             |               |  |
| <i>Aonides paucibranchiata</i>     |            | P723 | 131107   | Southern, 1914                | 1             |               | 11            |               |               | 1             | 1             | 3             |               |               |  |
| <i>Atherospio guillei</i>          |            |      | 478336   | (Laubier & Ramos,1974)        |               |               | 2             |               |               |               |               |               |               |               |  |
| <i>Laonice bahusiensis</i>         |            | P733 | 131127   | Soderstrom, 1920              |               |               |               |               |               |               |               |               |               |               |  |
| <i>Dipolydora caulleryi/sp. A</i>  |            | P751 | 131116   | Mesnil, 1897                  |               |               |               |               |               |               |               |               |               |               |  |
| <i>Dipolydora flava</i>            |            | P754 | 131118   | Claparede, 1870               |               |               |               |               |               |               |               |               |               |               |  |
| <i>Dipolydora sp. B</i>            |            | P762 | 131124   | Verrill, 1881                 |               |               |               |               |               |               |               |               |               |               |  |
| <i>Pseudopolydora pulchra</i>      |            | P774 | 131169   | (Carazzi, 1895)               |               |               |               |               |               | 3             |               |               |               |               |  |
| <i>Scoelepis korsuni</i>           |            |      | 131174   | Sikorski, 1994                |               |               |               |               |               |               |               |               |               |               |  |
| <i>Spio goniocéphala</i>           |            |      | 131184   | Thulin, 1957                  |               | 9             |               |               | 3             |               |               |               | 4             | 3             |  |
| <i>Spio symphyta</i>               |            |      | 596189   | Meisner, Bick & Bastrop, 2011 | 1             |               | 1             |               |               | 2             | 9             | 3             |               |               |  |
| <i>Spiophanes bombyx</i>           |            | P794 | 131187   | (Claparede, 1870)             | 1             | 2             | 2             |               | 2             | 5             |               |               | 3             |               |  |
| <i>Spiophanes kroyeri</i>          |            | P796 | 131188   | Grube, 1860                   |               |               |               |               |               | 1             |               |               |               |               |  |
| <i>Magelona johnstoni</i>          |            |      | 130269   | Fiege, Lichen & Mackie, 2000  |               |               |               |               |               |               |               |               |               |               |  |
| <i>Chaetopterus variopedatus</i>   |            | P814 | 129914   | (Renier, 1804)                |               |               |               |               |               |               |               |               |               |               |  |
| <i>Phyllochaetopterus anglicus</i> |            | P815 |          | Potts, 1914                   |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aphelochaeta sp. A</i>          |            | P823 | 129240   | Blake, 1991                   |               |               |               |               |               |               |               |               |               |               |  |
| <i>Caulleriella alata</i>          |            | P829 | 129943   | (Southern, 1914)              |               |               | 3             |               |               | 6             | 1             | 2             |               |               |  |



| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                | Sample        |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |                          | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Chaetozone zetlandica</i>      |            | P831  | 129948   | (McIntosh, 1911)         |               |               | 2             |               |               |               |               |               |               |               |
| <i>Chaetozone christiei</i>       |            |       | 152217   | Chambers, 2000           |               |               |               |               |               |               |               |               | 1             |               |
| <i>Cirratulus cirratus</i>        |            | P836  | 129959   | (O.F.Muller,1776)        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Tharyx</i>                     |            | P847  | 129249   | Webster & Benedict, 1887 |               |               | 1             |               |               |               |               |               |               |               |
| <i>Flabelligera affinis</i>       |            | P881  | 130103   | M.Sars, 1829             |               |               |               |               |               |               |               |               |               |               |
| <i>Mediomastus fragilis</i>       |            | P919  | 129892   | Rasmussen, 1973          |               |               |               |               |               | 3             |               | 1             |               |               |
| <i>Notomastus</i>                 |            | P920  | 129220   | Sars, 1850               |               |               | 2             |               |               | 4             |               |               |               |               |
| <i>Praxillura longissima</i>      |            | P944  | 130327   | Arwidsson, 1906          |               |               |               |               |               |               |               |               |               |               |
| <i>Leiochone</i>                  |            |       | 146991   | Grube, 1868              |               |               |               |               | 1             | 1             |               |               |               |               |
| <i>Euclymene lombricoides</i>     |            | P963  | 209899   | (Quatrefages, 1866)      |               |               |               |               |               |               |               |               |               |               |
| <i>Euclymene oerstedii</i>        |            | P964  | 130294   | (Claparède, 1863)        |               |               |               |               |               |               |               |               |               |               |
| <i>Praxillella affinis</i>        |            | P971  | 130322   | (M Sars, 1872)           |               |               |               |               |               | 3             |               |               |               |               |
| <i>Ophelia borealis</i>           |            | P999  | 130491   | Quatrefages, 1866        |               | 1             |               |               |               |               |               |               | 1             |               |
| <i>Ophelia celtica</i>            |            | P1000 | 130492   | Amoureux & Dauvin, 1981  |               |               |               |               |               |               |               |               |               |               |
| <i>Ophelina acuminata</i>         |            | P1014 | 130500   | Örsted, 1843             |               |               |               |               |               |               |               |               |               |               |
| <i>Asclerocheilus intermedius</i> |            | P1022 | 130974   | (Saint-Joseph, 1894)     |               |               | 1             |               |               | 2             |               |               |               |               |
| <i>Scalibregma celticum</i>       |            | P1026 | 130979   | Mackie, 1991             |               |               |               |               |               | 1             |               |               |               |               |
| <i>Scalibregma inflatum</i>       |            | P1027 | 130980   | Rathke, 1843             |               |               |               |               |               | 10            |               |               |               |               |
| <i>Polygordius</i>                |            | P1062 | 129472   | Schneider, 1868          | 1             |               |               |               |               |               | 40            | 7             |               |               |
| <i>Galathowenia oculata</i>       |            | P1093 | 146950   | Zaks, 1922               |               |               |               |               |               |               |               |               |               |               |
| <i>Owenia</i>                     |            | P1097 | 129427   | Delle Chiaje, 1844       |               |               | 6             |               |               | 2             |               | 1             |               |               |
| <i>Lagis koreni</i>               |            | P1107 | 152367   | Malmgren, 1866           |               |               | 2             |               |               | 7             |               |               | 1             |               |
| <i>Sabellaria spinulosa</i>       |            | P1117 | 130867   | Leuckart, 1849           |               |               |               |               |               |               |               |               |               |               |
| <i>Ampharete lindstroemi</i>      | aggregate  | P1139 | 129778   | M. Sars, 1864            |               |               | 2             |               |               | 5             | 2             |               |               |               |

| Taxon                             | Qualifiers | SDC   | APHIA ID | Authority                                     | Sample        |               |               |               |               |               |               |               |               |               |
|-----------------------------------|------------|-------|----------|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                   |            |       |          |   | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Anobothrus gracilis</i>        |            | P1147 | 129789   | (Malmgren, 1866)                              |               |               |               |               |               |               |               |               |               |               |
| <i>Terebellides stroemii</i>      |            | P1175 | 131573   | Sars, 1835                                    |               |               |               |               |               |               |               |               |               |               |
| <i>Lanice conchilega</i>          |            | P1195 | 131495   | (Pallas, 1766)                                |               |               | 1             |               |               |               |               |               |               |               |
| <i>Phisidia aurea</i>             |            | P1215 | 131513   | Southward, 1956                               |               |               |               |               |               |               |               |               |               |               |
| <i>Pista mediterranea</i>         |            |       | 131519   | de Gaillande, 1970                            |               |               | 2             |               |               | 2             | 2             |               |               |               |
| <i>Pista bansei</i>               |            |       | 152254   | Saphronova, 1988                              |               |               | 3             |               |               | 1             | 1             |               |               |               |
| <i>Polycirrus</i>                 |            | P1235 | 129710   | Grube, 1850                                   |               |               |               |               | 1             |               |               |               |               |               |
| <i>Lysilla loveni</i>             |            | P1233 | 131500   | (Malmgren, 1866)                              |               |               |               |               |               | 1             |               |               |               |               |
| <i>Lysilla nivea</i>              |            | P1234 | 131501   | Langerhans, 1884                              |               |               | 4             |               |               | 1             |               |               |               |               |
| <i>Streblosoma intestinale</i>    |            | P1252 | 131540   | M. Sars in G.O. Sars, 1872                    |               |               |               |               |               |               |               |               |               |               |
| <i>Thelepus cincinnatus</i>       |            | P1254 | 131543   | (Fabricius, 1780)                             |               |               |               |               |               |               |               |               |               |               |
| <i>Dialychone dunerificta</i>     |            |       | 558752   | (Tovar-Hernández, Licciano, Giangrande, 2007) |               |               |               |               |               |               |               |               |               |               |
| <i>Parasabella cambrensis</i>     |            | P1273 | 530920   | Knight-Jones & Walker, 1985                   |               |               |               |               |               | 1             |               |               |               |               |
| <i>Euchone pararosea</i>          |            |       | 390407   | Giangrande & Licciano, 2006                   |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudopotamilla reniformis</i> |            | P1316 | 130963   | (Bruguiere, 1789)                             |               |               |               |               |               |               |               |               |               |               |
| <i>Sabella pavonina</i>           |            | P1320 | 130967   | Savigny, 1822                                 |               |               |               |               |               |               |               |               |               |               |
| <i>Hydroides norvegica</i>        |            | P1334 | 131009   | Gunnerus, 1768                                |               |               |               |               |               | 1             |               |               |               |               |
| <i>Spirobranchus triqueter</i>    |            | P1341 | 555935   | (Linnaeus, 1758)                              |               |               |               |               |               | 1             |               |               |               |               |
| <b>OLIGOCHAETA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |
| <i>Grania</i>                     |            | P1524 | 137349   | Southern, 1913                                |               |               |               |               |               |               | 8             |               |               |               |
| <b>CHELICERATA</b>                |            |       |          |   |               |               |               |               |               |               |               |               |               |               |
| <i>Nymphon brevirostre</i>        |            | Q5    | 150520   | Hodge, 1863                                   |               |               |               |               |               |               |               |               |               |               |
| <i>Anoplodactylus petiolatus</i>  |            | Q44   | 134723   | (Kroyer, 1884)                                |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |      |          |                                   | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <b>CRUSTACEA</b>               |            |      |          |                                   |               |               |               |               |               |               |               |               |               |               |
| <i>Scalpellum scalpellum</i>   |            | R22  | 106204   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Verruca stroemia</i>        |            | R41  | 106257   | O.F.Muller, 1776                  |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia reboredae</i>       |            |      | 459311   | Moreira & Urganri, 2009           |               |               |               |               |               |               |               |               |               |               |
| <i>Nebalia borealis</i>        |            | S7   | 156257   | Dahl, 1985                        |               |               |               |               |               |               |               |               |               |               |
| <i>Sarsinebalia urgorrii</i>   |            |      | 388224   | Moreira, Gestoso & Troncoso, 2003 |               |               |               |               |               |               |               |               |               |               |
| <i>Heteromysis formosa</i>     |            | S93  | 120037   | (G. O. Sars, 1877)                |               |               |               |               |               | 3             |               |               |               |               |
| <i>Kroyera carinata</i>        |            | S125 | 547074   | Spence Bate, 1857                 |               |               |               |               |               |               |               |               |               |               |
| <i>Periculodes longimanus</i>  |            | S131 | 102915   | (Bate & Westwood, 1868)           |               |               |               |               |               |               |               |               |               |               |
| <i>Pontocrates arenarius</i>   |            | S135 | 102918   | (Bate, 1858)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Synchelidium maculatum</i>  |            | S138 | 102928   | Stebbing, 1906                    |               |               |               |               |               | 1             |               |               |               |               |
| <i>Apolochus neapolitanus</i>  |            | S159 | 236495   | (Della Valle, 1893)               |               |               |               |               |               | 1             |               |               |               |               |
| <i>Leucothoe incisa</i>        |            | S177 | 102460   | Robertson, 1892                   |               |               |               |               |               |               |               | 1             |               |               |
| <i>Leucothoe procera</i>       |            | S179 | 102466   | Bate, 1857                        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Stenothoe marina</i>        |            | S213 | 103166   | (Bate, 1856)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Urothoe elegans</i>         |            | S248 | 103228   | (Bate, 1856)                      |               |               |               |               |               | 2             |               |               |               |               |
| <i>Urothoe marina</i>          |            | S249 | 103233   | (Bate, 1857)                      |               |               | 18            |               |               | 8             | 1             | 2             |               |               |
| <i>Acidostoma neglectum</i>    |            |      | 102495   | (Spence Bate & Westwood, 1861)    |               |               |               |               |               |               |               |               |               |               |
| <i>Hippomedon denticulatus</i> |            | S296 | 102570   | (Bate, 1857)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Tryphosa nana</i>           |            | S321 | 102691   | (Kroyer, 1846)                    |               |               |               |               |               |               |               |               |               |               |
| <i>Tmetonyx similis</i>        |            | S337 | 102742   | (G O Sars, 1891)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Nototropis falcatus</i>     |            | S410 | 102139   | Metzger, 1871                     |               |               |               |               |               |               |               | 1             |               |               |
| <i>Nototropis vedlomensis</i>  |            | S413 | 102132   | (Bate & Westwood, 1862)           |               |               |               |               |               |               |               |               |               |               |

| Taxon                              | Qualifiers | SDC         | APHIA ID      | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |
|------------------------------------|------------|-------------|---------------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                    |            |             |               |                                   | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Ampelisca diadema</i>           |            | S429        | 101896        | (Costa, 1853)                     |               |               |               |               |               | 1             |               |               |               |               |
| <i>Ampelisca provincialis</i>      |            | S434        | 101915        | Bellan-Santini & Kaim-Malka, 1977 |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca spinipes</i>          |            | S438        | 101928        | Boeck, 1861                       |               |               | 1             |               |               | 6             |               |               |               |               |
| <i>Ampelisca tenuicornis</i>       |            | S440        | 101930        | Lilljeborg, 1855                  |               |               |               |               |               |               |               |               |               |               |
| <i>Ampelisca typica</i>            |            | S442        | 101933        | (Bate, 1856)                      |               |               | 1             |               |               | 3             | 1             | 1             |               |               |
| <i>Bathyporeia elegans</i>         |            | S452        | 103058        | Watkin, 1938                      | 2             | 8             |               |               | 1             |               |               |               | 2             | 5             |
| <i>Bathyporeia gracilis</i>        |            | S453        | 103059        | G O Sars, 1891                    |               |               |               |               |               |               |               |               |               |               |
| <i>Bathyporeia guilliamsoniana</i> |            | S454        | 103060        | (Bate, 1857)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Haustorius arenarius</i>        |            | S462        | 102317        | Slabber, 1769                     |               |               |               |               |               |               |               |               |               | 1             |
| <i>Abludomelita obtusata</i>       |            | S498        | 102788        | (Montagu, 1813)                   |               |               |               |               |               |               |               |               |               |               |
| <b>Cheirocratus</b>                |            | <b>S503</b> | <b>101669</b> | <b>Norman, 1867</b>               |               |               |               |               |               |               |               |               |               |               |
| <i>Othomaera othonis</i>           |            | S519        | 534781        | (Milne-Edwards, 1830)             |               |               |               |               |               |               |               |               |               |               |
| <i>Maerella tenuimana</i>          |            | S521        | 102831        | (Bate, 1862)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Megamphopus cornutus</i>        |            | S539        | 148545        | Norman, 1869                      |               |               |               |               |               |               | 1             |               |               |               |
| <i>Gammaropsis maculata</i>        |            | S541        | 102364        | (Johnston, 1828)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Photis longicaudata</i>         |            | S552        | 102383        | (Bate & Westwood, 1862)           |               |               |               |               |               | 8             |               |               |               |               |
| <b>Aoridae</b>                     |            | <b>S577</b> | <b>101368</b> | <b>Stebbing, 1899</b>             |               |               |               |               |               |               |               |               |               |               |
| <i>Leptocheirus hirsutimanus</i>   |            | S588        | 102036        | (Bate, 1862)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Crassicorophium crassicorne</i> |            | S611        | 397383        | Bruzelius, 1859                   |               |               |               |               |               |               |               |               |               |               |
| <i>Unciola planipes</i>            |            | S622        | 102061        | Norman, 1867                      |               |               | 1             |               |               | 1             | 1             | 1             |               |               |
| <i>Phtisica marina</i>             |            | S657        | 101864        | Slabber, 1769                     |               |               |               |               |               | 1             |               |               |               |               |
| <b>Gnathia</b>                     |            | <b>S793</b> | <b>118437</b> | <b>Leach, 1814</b>                |               |               |               |               |               |               |               |               |               |               |
| <i>Conilera cylindracea</i>        |            | S849        | 118842        | (Montagu, 1804)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Eurydice pulchra</i>            |            | S854        | 118852        | Leach, 1815                       | 1             |               | 2             |               |               |               | 2             | 1             |               |               |

| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority                         | Sample        |               |               |               |               |               |               |               |               |               |
|--------------------------------|------------|-------|----------|-----------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                |            |       |          |                                   | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Astacilla longicornis</i>   |            | S955  | 119024   | (Sowerby, 1806)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Tanaopsis graciloides</i>   |            | S1142 | 136458   | (Lilljeborg, 1864)                |               |               |               |               |               |               |               |               |               |               |
| <i>Bodotria scorpioides</i>    |            | S1197 | 110445   | (Montagu, 1804)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Diastylis bradyi</i>        |            | S1248 | 110472   | Norman, 1879                      |               |               |               |               |               |               |               |               |               |               |
| <i>Processa modica</i>         |            | S1366 | 108343   | Williamson & Rochanaburanon, 1979 |               |               |               |               |               |               |               |               |               |               |
| <i>Callianassa subterranea</i> |            | S1415 | 107729   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Upogebia deltaura</i>       |            | S1419 | 107739   | (Leach, 1815)                     |               |               | 1             |               |               | 5             |               |               |               |               |
| <i>Pisidia longicornis</i>     |            | S1482 | 107188   | (Linnaeus, 1757)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Ebalia tumefacta</i>        |            | S1509 | 107302   | (Montagu, 1808)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Hyas coarctatus</i>         |            | S1519 | 107323   | Leach, 1815                       |               |               |               |               |               |               |               |               |               |               |
| <i>Inachus dorsettensis</i>    |            | S1526 | 107327   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia rostrata</i>     |            | S1532 | 107345   | (Linnaeus, 1761)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Thia scutellata</i>         |            | S1559 | 107281   | (Fabricius, 1793)                 |               |               |               |               |               |               |               |               |               |               |
| <i>Liocarcinus pusillus</i>    |            | S1584 | 107393   | (Leach, 1816)                     |               |               |               |               |               |               |               |               |               |               |
| <i>Pinnotheres pisum</i>       |            | S1638 | 107473   | (Linnaeus, 1767)                  |               |               |               |               |               |               |               |               |               |               |
| <b>MOLLUSCA</b>                |            |       |          |                                   |               |               |               |               |               |               |               |               |               |               |
| <i>Leptochiton asellus</i>     |            | W53   | 140199   | (Gmelin, 1791)                    |               |               |               |               |               |               |               |               |               |               |
| <i>Acanthochitona crinita</i>  |            | W86   | 138675   | (Pennant, 1777)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Gibbula magus</i>           | ?          | W159  | 141790   | (Linnaeus, 1758)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Euspira nitida</i>          |            | W491  | 151894   | (Donovan, 1804)                   |               |               |               |               |               |               |               |               |               |               |
| <i>Melanella alba</i>          |            | W634  | 139832   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Colus gracilis</i>          |            | W715  | 138899   | (da Costa, 1778)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Philine</i>                 |            | W1036 | 138339   | Ascanius, 1772                    |               |               |               |               |               |               |               |               |               |               |
| <i>Tritonia plebeia</i>        |            | W1254 | 141738   | Johnston, 1828                    |               |               |               |               |               |               |               |               |               |               |
| Onchidorididae                 |            | W1319 | 175      | Gray, 1827                        |               |               |               |               |               |               |               |               |               |               |

| Taxon                          | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |  |
|--------------------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                |            |       |          |                         | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |  |
| <i>Knoutsodonta depressa</i>   |            | W1323 | 845528   | (Alder & Hancock, 1842) |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nucula hanleyi</i>          |            | W1568 | 140588   | Winckworth, 1931        |               |               |               |               |               | 1             | 1             |               |               |               |  |
| <i>Modiolus adriaticus</i>     |            | W1700 | 506025   | (Lamarck, 1819)         |               |               |               |               |               | 1             |               |               |               |               |  |
| <i>Aequipecten opercularis</i> |            | W1773 | 140687   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Heteranomia squamula</i>    |            | W1809 | 138749   | (Linnaeus, 1758)        |               |               |               |               |               | 3             |               |               |               |               |  |
| <i>Pododesmus squama</i>       |            | W1812 | 138752   | (Linnaeus, 1761)        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Hemilepton nitidum</i>      |            | W1882 | 246148   | (W. Turton, 1822)       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Kurtiella bidentata</i>     |            | W1906 | 345281   | (Montagu, 1803)         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Spisula elliptica</i>       |            | W1975 | 140300   | (Brown, 1827)           |               | 1             |               |               | 2             |               | 2             |               |               |               |  |
| <i>Ensis leei</i>              |            | W1997 | 876640   | M. Huber, 2015          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Phaxas pellucidus</i>       |            | W2006 | 140737   | (Pennant, 1777)         |               |               |               |               |               |               |               |               | 1             |               |  |
| <i>Moerella donacina</i>       |            | W2021 | 147021   | Linnaeus, 1758          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Asbjornsenia pygmaea</i>    |            | W2023 | 879714   | (Lovén, 1846)           | 1             |               |               |               |               |               | 1             |               |               |               |  |
| <i>Gari fervensis</i>          |            | W2051 | 140870   | (Gmelin, 1791)          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Solecurtus scopula</i>      |            | W2054 | 141543   | (Turton, 1822)          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Abra alba</i>               |            | W2059 | 141433   | (W Wood, 1802)          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Abra prismatica</i>         |            | W2062 | 141436   | (Montagu, 1808)         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Clausinella fasciata</i>    |            | W2100 | 141909   | (da Costa, 1778)        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Timoclea ovata</i>          |            | W2104 | 141929   | (Pennant, 1777)         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Venerupis corrugata</i>     |            | W2124 | 181364   | (Gmelin, 1791)          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Dosinia exoleta</i>         |            | W2130 | 141911   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Hiatella arctica</i>        |            | W2166 | 140103   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Thracia villosiuscula</i>   |            | W2233 | 141651   | (Macgillivray, 1827)    |               |               |               |               |               |               |               | 1             |               |               |  |
| <i>Cochlodesma praetenue</i>   |            | W2239 | 181373   | (Pulteney, 1799)        |               |               |               |               |               |               |               |               |               |               |  |
| <b>PHORONIDA</b>               |            |       |          |                         |               |               |               |               |               |               |               |               |               |               |  |



| Taxon                                       | Qualifiers | SDC   | APHIA ID | Authority                           | Sample        |               |               |               |               |               |               |               |               |               |
|---|------------|-------|----------|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|   |            |       |          |                                     | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <i>Phoronis</i>                             |            | ZA3   | 128545   | Wright, 1856                        |               |               | 8             |               |               | 12            |               |               |               |               |
| <b>ECHINODERMATA</b>                        |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |
| <i>Ophiothrix fragilis</i>                  |            | ZB124 | 125131   | (Abildgaard, 1789)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Amphipholis squamata</i>                 |            | ZB161 | 125064   | (Chiaje, 1828)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Ophiura albida</i>                       |            | ZB168 | 124913   | Forbes, 1839                        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Psammechinus miliaris</i>                |            | ZB193 | 124319   | (Gmelin, 1778)                      |               |               |               |               |               |               |               |               |               |               |
| <i>Echinocyamus pusillus</i>                |            | ZB212 | 124273   | (O.F. Muller, 1776)                 | 2             |               |               |               | 3             |               | 1             |               | 5             |               |
| <i>Echinocardium cordatum</i>               |            | ZB223 | 124392   | (Pennant, 1777)                     |               |               |               |               | 1             |               |               |               |               |               |
| <i>Mesothuria intestinalis</i>              |            | ZB233 | 124568   | (Ascanius, 1805)<br>Östergren, 1896 |               |               |               |               |               |               |               |               |               |               |
| <i>Pseudothyone raphanus</i>                |            | ZB257 | 124661   | (Duben & Koren, 1845)               |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone fusus</i>                         |            | ZB262 | 124670   | O.F.Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |
| <i>Thyone roscovita</i>                     |            | ZB264 | 124676   | Hérouard, 1889                      |               |               |               |               |               |               |               | 1             |               |               |
| <i>Panningia hyndmani</i>                   |            | ZB272 | 848014   | (Thompson, 1840)                    |               |               |               |               |               | 2             |               |               |               |               |
| <i>Leptosynapta inhaerens</i>               |            | ZB296 | 124465   | (O.F.Muller, 1776)                  |               |               |               |               |               |               |               |               |               |               |
| <i>Oestergrenia digitata</i>                |            | ZB300 | 152547   | (Montagu, 1804)                     |               |               |               |               |               |               |               |               |               |               |
| <b>TUNICATA</b>                             |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |
| <i>Ascidella aspersa</i>                    |            | ZD84  | 103718   | O F Müller, 1776                    |               |               | 1             |               |               |               |               |               |               |               |
| <i>Ascidia conchilega</i>                   | ?          | ZD88  | 103702   | O F Muller, 1776                    |               |               |               |               |               |               |               |               |               |               |
| <i>Polycarpa fibrosa</i>                    |            | ZD112 | 103902   | (Stimpson, 1852)                    |               |               |               |               |               |               |               |               |               |               |
| <i>Dendrodoa grossularia</i>                |            | ZD120 | 103882   | (van Beneden 1846)                  |               |               |               |               |               | 1             |               |               |               |               |
| <b>CEPHALOCHORDATA</b>                      |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |
| <i>Branchiostoma lanceolatum</i>            |            |       | 104906   | (Pallas, 1774)                      |               |               |               |               |               |               | 1             |               |               |               |
| <b>Number of taxa</b>                       |            |       |          |                                     | 12            | 8             | 34            | 2             | 11            | 65            | 25            | 23            | 14            | 4             |
| <b>Abundance</b>                            |            |       |          |                                     | 14            | 25            | 95            | 2             | 17            | 187           | 122           | 62            | 31            | 10            |
| The following taxa were merged for analysis |            |       |          |                                     |               |               |               |               |               |               |               |               |               |               |

| Taxon  | Qualifiers | SDC  | APHIA ID | Authority          | Sample        |               |               |               |               |               |               |               |               |               |
|--|------------|------|----------|--------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|  |            |      |          |                    | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| <b>Aoridae</b>                                 |            | S577 | 101368   | Stebbing, 1899     |               |               |               |               |               |               |               |               |               |               |
| <i>Aoridae</i>                                 | female     | S577 | 101368   | Stebbing, 1899     |               |               |               |               |               |               |               |               |               |               |
| <i>Autonoe longipes</i>                        |            | S583 | 102021   | (Lilljeborg, 1852) |               |               |               |               |               |               |               |               |               |               |
| <b>Gnathia</b>                                 |            | S793 | 118437   | Leach, 1814        |               |               |               |               |               |               |               |               |               |               |
| <i>Gnathia</i>                                 | female     | S793 | 118437   | Leach, 1814        |               |               |               |               |               |               |               |               |               |               |
| <i>Gnathia oxyuraea</i>                        |            | S796 | 118995   | (Lilljeborg, 1855) |               |               |               |               |               |               |               |               |               |               |
| <b>Cheirocratus</b>                            |            | S503 | 101669   | Norman, 1867       |               |               |               |               |               |               |               |               |               |               |
| <i>Cheirocratus</i>                            | female     | S503 | 101669   | Norman, 1867       |               |               |               |               |               |               |               |               |               |               |
| <i>Cheirocratus pseudosundevallii</i>          | ?          |      |          | Gouillieux, 2019   |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b>                          |            |      |          |                    | 0             | 0             | 0             | 0             | 0             | 0             | 0             | 0             | 0             | 0             |
| <b>Abundance</b>                               |            |      |          |                    | 0             | 0             | 0             | 0             | 0             | 0             | 0             | 0             | 0             | 0             |
| The following taxa were excluded from analysis |            |      |          |                    |               |               |               |               |               |               |               |               |               |               |
| <b>Colonial</b>                                |            |      |          |                    |               |               |               |               |               |               |               |               |               |               |
| PORIFERA                                       |            | C1   | 558      | Grant, 1836        |               |               |               |               |               | P             |               |               |               |               |
| <i>Leuckartiara octona</i>                     |            | D240 | 117791   | (Fleming, 1823)    |               |               |               |               |               |               |               |               |               |               |
| <i>Tubularia</i>                               |            | D163 | 117258   | Linnaeus, 1758     |               |               | P             |               | P             |               |               |               |               |               |
| <i>Eudendrium</i>                              |            | D218 | 117093   | Ehrenberg, 1834    |               |               |               |               |               |               |               |               |               |               |
| Bougainvilliidae                               |            | D246 | 1594     | Lütken, 1850       |               |               |               |               |               | P             |               |               |               |               |
| <i>Phialella quadrata</i>                      |            | D343 | 117804   | (Forbes, 1848)     |               |               |               |               |               |               |               |               |               |               |
| <i>Calycella syringa</i>                       |            | D348 | 117402   | (Linnaeus, 1767)   |               |               |               |               |               |               |               |               |               |               |
| <i>Lafoea dumosa</i>                           |            | D386 | 117702   | (Fleming, 1828)    |               |               |               |               |               |               |               |               |               |               |
| <i>Halecium</i>                                |            | D390 | 117103   | Oken, 1815         |               |               |               |               |               | P             |               |               |               |               |
| <i>Abietinaria</i>                             |            | D408 | 117225   | Kirchenpauer, 1884 |               |               |               |               |               |               |               |               |               |               |
| <i>Diphasia</i>                                |            | D413 | 117228   | L Agassiz, 1862    |               |               |               |               |               |               |               |               |               |               |
| <i>Hydrallmania falcata</i>                    |            | D424 | 117890   | (Linnaeus, 1758)   |               |               |               |               |               | P             |               |               |               |               |

| Taxon                          | Qualifiers | SDC  | APHIA ID | Authority             | Sample        |               |               |               |               |               |               |               |               |               |  |
|--------------------------------|------------|------|----------|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                |            |      |          |                       | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |  |
| <i>Sertularella gayi</i>       |            | D429 | 117902   | (Lamouroux, 1821)     |               |               |               |               |               |               |               |               |               |               |  |
| <i>Sertularia cupressina</i>   |            | D435 | 117913   | (Linnaeus, 1758)      |               |               |               |               |               | P             |               |               |               |               |  |
| <i>Nemertesia</i>              |            | D462 | 117195   | Lamouroux, 1812       |               |               |               |               |               | P             |               |               |               |               |  |
| Campanulariidae                |            | D491 | 1606     | Peron & Lesueur, 1810 |               |               |               |               |               | P             |               |               |               |               |  |
| <i>Alcyonium digitatum</i>     |            | D597 | 125333   | Linnaeus, 1758        |               |               |               |               |               | P             |               |               |               |               |  |
| <i>Crisia</i>                  |            | Y13  | 111032   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Tubulipora</i>              |            | Y27  | 111054   | Lamarck, 1816         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Disporella hispida</i>      |            | Y66  | 111730   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Alcyonidium parasiticum</i> |            | Y81  | 111604   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Vesicularia spinosa</i>     |            | Y131 | 111669   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |  |
| <i>Amathia lendigera</i>       |            | Y135 | 111659   | (Linnaeus, 1758)      |               |               |               |               |               |               |               |               |               |               |  |
| <i>Amathia</i>                 |            | Y137 | 111023   | Farre, 1837           |               |               |               |               |               | P             |               |               |               |               |  |
| <i>Aetea</i>                   |            | Y153 | 110819   | Lamouroux, 1812       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Conopeum reticulum</i>      |            | Y172 | 111351   | (Linnaeus, 1767)      |               |               |               |               |               |               |               |               |               |               |  |
| <i>Electra pilosa</i>          |            | Y178 | 111355   | (Linnaeus, 1767)      |               |               |               |               |               |               |               |               |               |               |  |
| <i>Callopora</i>               |            | Y201 | 110851   | J E Gray, 1848        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Beania mirabilis</i>        |            | Y261 | 111072   | Johnston, 1840        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Chorizopora brongiartii</i> |            | Y344 | 111304   | (Audouin, 1826)       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Escharella immersa</i>      |            | Y364 | 111484   | (Fleming, 1828)       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Porella concinna</i>        |            | Y385 | 111125   | (Busk, 1854)          |               |               |               |               |               |               |               |               |               |               |  |
| <i>Schizomavella</i>           |            | Y467 | 110829   | Canu & Bassler, 1917  |               |               |               |               |               | P             |               |               |               |               |  |
| <b>Damaged</b>                 |            |      |          |                       |               |               |               |               |               |               |               |               |               |               |  |
| <i>Eumida</i>                  | damaged    | P163 | 129446   | Malmgren, 1865        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Aricidea</i>                | damaged    | P675 | 129430   | Webster, 1879         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Paradoneis</i>              | damaged    | P695 | 129433   | Hartman, 1965         |               |               |               |               |               |               |               |               |               |               |  |

| Taxon               | Qualifiers | SDC    | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |  |
|---------------------|------------|--------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                     |            |        |          |                         | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |  |
| Spionidae           | damaged    | P720   | 913      | Grube,1850              |               |               | 1             |               |               |               |               |               |               |               |  |
| <i>Dipolydora</i>   | damaged    | P748   | 129611   | Verrill, 1881           |               |               |               |               |               |               |               |               |               |               |  |
| <i>Terebellidae</i> | damaged    | P1179  | 982      | Johnston, 1846          |               |               |               |               |               | 3             |               | 2             |               |               |  |
| Sabellidae          | damaged    | P1257  | 985      | Latreille, 1825         |               |               |               |               |               |               |               |               |               |               |  |
| Serpulidae          | damaged    | P1324  | 988      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nebalia</i>      | damaged    | S5     | 147031   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |  |
| AMPHIPODA           | damaged    | S97    | 1135     |                         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Urothoe</i>      | damaged    | S246   | 101789   | Dana, 1852              |               |               | 1             |               |               |               |               |               |               |               |  |
| <i>Liocarcinus</i>  | damaged    | S1577  | 106925   | Stimpson,1870           |               |               |               |               |               |               |               |               |               |               |  |
| GASTROPODA          | damaged    | W88    | 101      |                         |               |               |               |               |               | 1             |               |               |               |               |  |
| BIVALVIA            | damaged    | W1560  | 151265   | Goldfuss, 1820          | 1             |               |               |               |               |               |               |               |               |               |  |
| SOLENOIDEA          | damaged    | W1991  | 14635    | Lamarck, 1809           |               |               | 1             |               |               |               |               |               |               |               |  |
| Ophiuridae          | damaged    | ZB165  | 123200   | Muller & Troschel, 1840 |               |               |               |               |               |               |               |               | 1             |               |  |
| <b>Fish</b>         |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |  |
| Gobiesocidae        |            | ZG1180 | 125477   | Bleeker, 1859           |               |               |               |               |               |               |               |               |               |               |  |
| <b>Juvenile</b>     |            |        |          |                         |               |               |               |               |               |               |               |               |               |               |  |
| ANIMALIA            | eggs       |        | 2        |                         |               |               |               |               |               | 1             |               |               |               |               |  |
| SIPUNCULA           | juvenile   | N1     | 1268     |                         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Golfingia</i>    | juvenile   | N12    | 136021   | Lankester, 1885         |               |               |               |               |               |               |               |               |               |               |  |
| <i>Glycera</i>      | juvenile   | P255   | 129296   | Lamarck, 1818           |               |               |               |               |               |               |               |               |               |               |  |
| Goniadidae          | juvenile   | P266   | 953      | Kinberg, 1866           |               |               |               |               |               |               |               |               |               |               |  |
| Nereididae          | juvenile   | P458   | 22496    | Blainville, 1818        |               |               |               |               |               |               |               |               |               |               |  |
| <i>Nephtys</i>      | juvenile   | P494   | 129370   | Cuvier, 1817            | 1             |               |               |               |               |               |               |               |               |               |  |
| <i>Scolelepis</i>   | juvenile   | P778   | 129623   | Blainville, 1828        |               |               |               |               |               |               |               |               | 1             |               |  |
| CRUSTACEA           | larva      | R1     | 1066     |                         |               |               |               |               |               |               |               |               | 1             |               |  |

| Taxon                | Qualifiers | SDC   | APHIA ID | Authority               | Sample        |               |               |               |               |               |               |               |               |               |
|----------------------|------------|-------|----------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                      |            |       |          |                         | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| BALANOMORPHA         | juvenile   | R42   | 1082     | Burmeister, 1834        |               |               |               |               |               | 1             |               |               |               |               |
| <i>Ampelisca</i>     | juvenile   | S423  | 101445   | Kroyer, 1842            |               |               |               |               |               |               |               |               |               |               |
| <i>Gnathia</i>       | juvenile   | S793  | 118437   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |
| GEBIIDEA             | juvenile   | S1403 | 477323   | de Saint Laurent, 1979  |               |               |               |               |               | 5             |               | 1             |               |               |
| Paguridae            | juvenile   | S1445 | 106738   | Latreille, 1802         |               |               | 1             |               |               |               |               |               | 1             |               |
| DECAPODA             | larva      | S1276 | 1130     | Latreille, 1830         |               | 1             |               |               |               |               |               |               |               |               |
| <i>Ebalia</i>        | juvenile   | S1504 | 106889   | Leach, 1817             |               |               |               |               |               |               |               |               |               |               |
| Majidae              | juvenile   | S1512 | 106760   | Samouelle, 1819         |               |               |               |               |               | 1             |               |               |               |               |
| Inachinae            | juvenile   | S1520 | 148436   | MacLeay, 1838           |               |               |               |               |               |               |               |               |               |               |
| <i>Macropodia</i>    | juvenile   | S1529 | 205077   | Leach, 1814             |               |               |               |               |               |               |               |               |               |               |
| Buccinidae           | eggs       | W702  | 149      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |
| <i>Tritonia</i>      | juvenile   | W1246 | 138580   | Cuvier, 1798            |               |               |               |               |               |               |               |               |               |               |
| <i>Nucula</i>        | juvenile   | W1565 | 138262   | Lamarck, 1799           |               |               |               |               |               |               |               |               |               |               |
| Mytilidae            | juvenile   | W1691 | 211      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |
| <i>Modiolus</i>      | juvenile   | W1702 | 140467   | (Linnaeus, 1758)        |               |               |               |               |               |               |               |               |               |               |
| Pectinidae           | juvenile   | W1768 | 213      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |
| Anomiidae            | juvenile   | W1805 | 214      | Rafinesque, 1815        |               |               |               |               |               |               |               |               |               |               |
| <i>Abra</i>          | juvenile   | W2058 | 138474   | Lamarck, 1818           |               |               |               |               |               |               |               |               |               |               |
| <i>Dosinia</i>       | juvenile   | W2126 | 138636   | Scopoli, 1777           |               |               |               |               |               |               |               |               |               |               |
| Ophiuridae           | juvenile   | ZB165 | 123200   | Muller & Troschel, 1840 |               |               | 1             |               |               |               |               |               |               |               |
| ECHINOIDEA           | juvenile   | ZB181 | 123082   | Leske, 1778             |               |               |               |               |               |               |               |               |               |               |
| <i>Echinocardium</i> | juvenile   | ZB222 | 123426   | Gray, 1825              |               |               |               |               |               |               |               |               |               |               |
| HOLOTHUROIDEA        | juvenile   | ZB229 | 123083   |                         |               |               |               |               |               |               |               |               | 2             |               |
| <i>Thyone</i>        | juvenile   | ZB261 | 146116   | Oken, 1815              |               |               |               |               |               | 2             |               |               |               |               |
| <i>Leptosynapta</i>  | juvenile   | ZB291 | 123449   | Verrill, 1867           |               |               |               |               |               |               |               |               |               |               |

| Taxon                 | Qualifiers | SDC  | APHIA ID | Authority     | Sample        |               |               |               |               |               |               |               |               |               |
|-----------------------|------------|------|----------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                       |            |      |          |               | MA_ST56<br>FA | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| ASCIDIACEA            | juvenile   | ZD2  | 1839     | Nielsen, 1995 |               |               |               |               |               |               |               |               |               |               |
| <b>Meiofaunal</b>     |            |      |          |               |               |               |               |               |               |               |               |               |               |               |
| NEMATODA              |            | HD1  | 799      |               |               |               |               |               |               | 1             | 38            | 13            |               |               |
| <b>Parasitic</b>      |            |      |          |               |               |               |               |               |               |               |               |               |               |               |
| COPEPODA              | parasitic  | R142 | 1080     |               |               |               |               |               |               |               |               |               |               |               |
| <b>Number of taxa</b> |            |      |          |               | 2             | 1             | 6             | 0             | 1             | 18            | 1             | 3             | 5             | 0             |
| <b>Abundance</b>      |            |      |          |               | 2             | 1             | 5             | 0             | 0             | 15            | 38            | 16            | 6             | 0             |



## F.2 Macrofaunal Biomass

| Taxon  | SDC   | Sample        |               |               |               |               |               |               |               |               |                |               |               |               |
|--|-------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|---------------|---------------|---------------|
|  |       | MA_ST01<br>FA | MA_ST02<br>FA | MA_ST03<br>FA | MA_ST04<br>FA | MA_ST05<br>FA | MA_ST06<br>FA | MA_ST07<br>FA | MA_ST08<br>FA | MA_ST09<br>FA | MA_ST10<br>FA  | MA_ST11<br>FA | MA_ST12<br>FA | MA_ST13<br>FA |
| CNIDARIA   | D1    | 0.2478        | 0.0016        |               | 0.0046        |               |               |               |               |               |                | 0.3354        |               |               |
| POLYCHAETA   | P2    | 0.6402        | 0.6448        | 0.6992        | 0.3854        | 1.5914        | 0.7088        | 3.2613        | 0.2604        | 0.5155        | 0.9680         | 0.9235        | 0.6097        | 0.4086        |
| OLIGOCHAETA  | P1402 |               |               |               |               |               |               |               |               |               |                |               |               |               |
| CRUSTACEA  | R1    | 0.2725        | 0.1652        | 0.5513        | 0.2803        | 0.3244        | 0.2780        | 0.3567        | 0.4322        | 0.1856        | 0.5851         | 0.6256        | 0.4970        | 0.2155        |
| MOLLUSCA   | W1    | 0.6305        | 0.1162        | 1.0426        | 0.6421        | 3.4819        | 0.0103        | 0.0598        | 0.6467        | 0.0745        | 13.3921        | 0.8033        | 0.2447        | 0.0129        |
| ECHINODERMATA  | ZB2   | 2.2758        | 0.3007        | 0.0797        | 3.0857        | 3.7452        | 0.0789        | 0.6434        |               | 4.4809        |                | 0.6815        | 3.3705        | 0.3372        |
| OSTEICHTHYES   | ZG1   |               |               |               |               |               |               |               |               |               |                |               |               |               |
| OTHERS   |       | 0.0491        | 0.0086        | 0.1409        | 2.2580        | 0.0465        | 0.0308        | 0.0114        | 0.0133        | 0.0668        | 0.0335         | 0.0324        | 0.1831        | 0.0357        |
| <b>Total</b>   |       | <b>4.1159</b> | <b>1.2371</b> | <b>2.5137</b> | <b>6.6561</b> | <b>9.1894</b> | <b>1.1068</b> | <b>4.3326</b> | <b>1.3526</b> | <b>5.3233</b> | <b>14.9787</b> | <b>3.4017</b> | <b>4.9050</b> | <b>1.0099</b> |
| Notes  |       |               |               |               |               |               |               |               |               |               |                |               |               |               |
| 'Blotted wet weight (g) prior to ash free dry weight conversion' |       |               |               |               |               |               |               |               |               |               |                |               |               |               |

| Taxon  | SDC   | Sample        |                |               |               |               |               |               |               |               |               |               |               |               |
|--|-------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|  |       | MA_ST14<br>FA | MA_ST15<br>FA  | MA_ST16<br>FA | MA_ST17<br>FA | MA_ST18<br>FA | MA_ST19<br>FA | MA_ST20<br>FA | MA_ST21<br>FA | MA_ST22<br>FA | MA_ST23<br>FA | MA_ST24<br>FA | MA_ST25<br>FA | MA_ST27<br>FA |
| CNIDARIA   | D1    |               |                | 0.3258        |               | 0.4274        |               |               |               |               |               |               |               |               |
| POLYCHAETA   | P2    | 0.6638        | 3.3554         | 1.2394        | 1.4438        | 0.9370        | 1.1774        | 1.3863        | 0.4904        | 0.4707        | 2.4050        | 1.3463        | 0.3504        | 0.3419        |
| OLIGOCHAETA  | P1402 |               |                |               |               |               |               |               |               |               |               |               |               |               |
| CRUSTACEA  | R1    | 1.1294        | 0.4047         | 0.2658        | 0.2571        | 0.3826        | 0.3380        | 0.1702        | 0.0397        | 0.1149        | 0.5848        | 0.1134        | 0.2178        | 0.8124        |
| MOLLUSCA   | W1    | 0.6098        | 8.4486         | 0.5140        | 0.2678        | 1.6695        | 0.6418        | 0.4991        |               | 0.3759        | 2.2248        | 0.3059        | 4.7396        |               |
| ECHINODERMATA  | ZB2   | 0.0786        | 3.6750         | 2.0960        | 0.0903        | 0.0276        | 2.5080        | 0.0909        |               | 0.0066        | 0.0575        | 0.0128        | 0.0260        | 0.0165        |
| OSTEICHTHYES   | ZG1   |               |                |               |               |               |               |               |               |               |               |               |               |               |
| OTHERS   |       | 0.0766        | 0.0535         | 0.0130        | 0.1780        | 0.2629        | 0.2140        | 0.4339        | 0.0839        | 0.0904        | 0.1162        | 0.0029        | 0.2701        | 0.0580        |
| <b>Total</b>   |       | <b>2.5582</b> | <b>15.9372</b> | <b>4.4540</b> | <b>2.2370</b> | <b>3.7070</b> | <b>4.8792</b> | <b>2.5804</b> | <b>0.6140</b> | <b>1.0585</b> | <b>5.3883</b> | <b>1.7813</b> | <b>5.6039</b> | <b>5.6157</b> |
| Notes  |       |               |                |               |               |               |               |               |               |               |               |               |               |               |
| 'Blotted wet weight (g) prior to ash free dry weight conversion' |       |               |                |               |               |               |               |               |               |               |               |               |               |               |

| Taxon         | SDC   | Sample        |               |               |               |               |               |               |               |               |               |               |               |               |
|---------------|-------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|               |       | MA_ST29<br>FA | MA_ST30<br>FA | MA_ST31<br>FA | MA_ST32<br>FA | MA_ST33<br>FA | MA_ST34<br>FA | MA_ST35<br>FA | MA_ST36<br>FA | MA_ST37<br>FA | MA_ST38<br>FA | MA_ST39<br>FA | MA_ST40<br>FA | MA_ST41<br>FA |
| CNIDARIA      | D1    |               | 0.3895        |               |               | 0.0018        |               |               |               | 0.2152        |               |               | 1.1121        |               |
| POLYCHAETA    | P2    | 0.3419        | 0.6255        | 0.2385        | 0.0635        | 0.4708        | 0.0598        | 0.0857        | 0.3511        | 0.8971        | 0.1367        | 0.0817        | 2.7048        | 0.5733        |
| OLIGOCHAETA   | P1402 |               |               |               | 0.0001        |               |               |               |               |               |               |               |               |               |
| CRUSTACEA     | R1    | 0.8124        | 0.2273        | 0.0891        | 0.0872        | 0.2121        | 0.0990        | 0.0071        | 0.0808        | 1.1724        | 0.0206        | 0.0276        | 0.3104        | 0.5463        |
| MOLLUSCA      | W1    |               | 0.2293        | 0.1786        | 1.0678        | 0.0016        | 0.6212        | 0.0556        | 0.0442        | 0.4143        | 0.0011        | 0.0024        |               | 3.7644        |
| ECHINODERMATA | ZB2   | 0.0165        | 0.0212        | 0.0051        | 3.2669        |               |               | 0.0084        |               | 0.9321        | 0.0625        | 0.0505        | 0.0247        | 0.0560        |
| OSTEICHTHYES  | ZG1   |               | 0.0068        |               |               |               |               |               |               |               |               |               |               |               |
| OTHERS        |       | 0.0454        | 0.2414        | 0.0909        | 0.0122        | 0.1512        | 0.0054        | 0.0077        | 0.1776        | 0.1438        | 0.0023        | 0.0216        | 0.8282        | 0.0378        |
| <b>Total</b>  |       | <b>1.2162</b> | <b>1.7410</b> | <b>0.6022</b> | <b>4.4977</b> | <b>0.8375</b> | <b>0.7854</b> | <b>0.1645</b> | <b>0.6537</b> | <b>3.7749</b> | <b>0.2232</b> | <b>0.1838</b> | <b>4.9802</b> | <b>4.9778</b> |

Notes

'Blotted wet weight (g) prior to ash free dry weight conversion'

| Taxon         | SDC   | Sample         |               |               |               |               |               |               |                |               |               |               |               |               |
|---------------|-------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|
|               |       | MA_ST42<br>FA  | MA_ST43<br>FA | MA_ST44<br>FA | MA_ST45<br>FA | MA_ST46<br>FA | MA_ST47<br>FA | MA_ST48<br>FA | MA_ST49<br>FA  | MA_ST50<br>FA | MA_ST51<br>FA | MA_ST52<br>FA | MA_ST54<br>FA | MA_ST55<br>FA |
| CNIDARIA      | D1    |                | 0.0075        |               |               |               |               |               |                |               |               |               |               |               |
| POLYCHAETA    | P2    | 1.4155         | 0.7779        | 0.4111        | 0.9272        | 0.1778        | 0.2198        | 0.3781        | 3.4641         | 0.5144        | 0.1764        | 0.4236        | 0.3948        | 1.0536        |
| OLIGOCHAETA   | P1402 |                |               |               |               |               |               |               |                |               |               |               |               |               |
| CRUSTACEA     | R1    | 0.2602         | 4.6834        | 0.0215        | 0.2133        | 0.0098        | 0.0572        | 0.2265        | 0.2540         | 0.0192        | 0.1169        | 0.0284        | 0.1705        | 0.4874        |
| MOLLUSCA      | W1    | 14.1518        | 0.6066        | 0.1946        | 0.0439        | 0.0423        | 0.7853        | 1.1496        | 12.7466        | 0.0217        | 0.1235        | 0.1585        | 0.6376        |               |
| ECHINODERMATA | ZB2   | 11.5337        | 0.6140        |               |               | 0.0224        | 0.1214        | 0.1342        | 0.0400         | 0.1185        | 0.1392        | 0.0057        | 0.1090        |               |
| OSTEICHTHYES  | ZG1   |                |               |               |               |               |               |               |                |               |               |               |               |               |
| OTHERS        |       | 1.6945         |               |               |               |               |               |               |                |               |               |               |               |               |
| <b>Total</b>  |       | <b>29.0557</b> | <b>6.6894</b> | <b>0.6272</b> | <b>1.1844</b> | <b>0.2523</b> | <b>1.1837</b> | <b>1.8884</b> | <b>16.5047</b> | <b>0.6738</b> | <b>0.5560</b> | <b>0.6162</b> | <b>1.3119</b> | <b>1.5410</b> |

Notes

'Blotted wet weight (g) prior to ash free dry weight conversion'

| Taxon  | SDC   | Sample         |               |               |               |               |                |               |               |               |               |
|--|-------|----------------|---------------|---------------|---------------|---------------|----------------|---------------|---------------|---------------|---------------|
|  |       | MA_ST56<br>FA  | MA_ST57<br>FA | MA_ST58<br>FA | MA_ST59<br>FA | MA_ST60<br>FA | MA_ST61<br>FA  | MA_ST63<br>FA | MA_ST64<br>FA | MA_ST65<br>FA | MA_ST66<br>FA |
| CNIDARIA   | D1    |                |               |               |               |               | 1.5008         |               | 0.0021        |               |               |
| POLYCHAETA   | P2    | 0.0210         | 0.4687        | 0.6771        | 0.0025        | 0.2069        | 2.5194         | 0.1324        | 0.1322        | 0.1896        | 0.0250        |
| OLIGOCHAETA  | P1402 |                |               |               |               |               |                | 0.0007        |               |               |               |
| CRUSTACEA  | R1    | 0.0262         | 0.0072        | 0.6172        |               | 0.0006        | 1.7219         | 0.0131        | 0.0255        | 0.0070        | 0.0552        |
| MOLLUSCA   | W1    | 15.6904        | 0.0156        | 1.5242        |               | 0.0450        | 0.4853         | 0.8387        | 0.0089        | 0.0009        |               |
| ECHINODERMATA  | ZB2   | 0.0034         |               | 0.0051        |               | 6.3493        | 5.1696         | 0.0046        | 4.3345        | 0.2745        |               |
| OSTEICHTHYES   | ZG1   |                |               |               |               |               |                |               |               |               |               |
| OTHERS   |       |                |               |               |               |               |                | 0.0592        |               |               |               |
| <b>Total</b>   |       | <b>15.7410</b> | <b>0.4915</b> | <b>2.8236</b> | <b>0.0025</b> | <b>6.6018</b> | <b>11.3970</b> | <b>1.0487</b> | <b>4.5032</b> | <b>0.4720</b> | <b>0.0802</b> |
| <b>Notes</b><br>'Blotted wet weight (g) prior to ash free dry weight conversion' |       |                |               |               |               |               |                |               |               |               |               |



RWE Renewables UK  
Swindon Limited

Windmill Hill Business Park  
Whitehill Way  
Swindon  
Wiltshire SN5 6PB  
T +44 (0)8456 720 090  
[www.rwe.com](http://www.rwe.com)

Registered office:  
RWE Renewables UK  
Swindon Limited  
Windmill Hill Business Park  
Whitehill Way  
Swindon