



FLORA OF KARNATAKA *A Checklist*

Volume - 2 : Gymnosperms & Angiosperms

KARNATAKA BIODIVERSITY BOARD

CITATION

Karnataka Biodiversity Board, 2019. FLORA OF KARNATAKA, A Checklist, Volume – 2: Gymnosperms and Angiosperms. 1 - 1002 (Published by Karnataka Biodiversity Board)

Published: December, 2019.

ISBN - 978-81-939228-1-1

© Karnataka Biodiversity Board, 2019

ALL RIGHTS RESERVED

- No part of this book, or plates therein, may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying recording or otherwise without the prior permission of the publisher.
- This book is sold subject to the condition that it shall not, by way of trade, be lent, re-sold, hired out or otherwise disposed of without the publisher's consent, in any form of binding or cover other than that in which it is published.
- The correct price of this publication is the price printed on this page. Any revised price indicated by a rubber stamp or by a sticker or by any other means is incorrect and should be unacceptable.

DISCLAIMER

- THE CONTENTS INCLUDING TEXT, PLATES AND OTHER INFORMATION GIVEN IN THE BOOK ARE SOLELY THE AUTHOR'S RESPONSIBILITY AND BOARD DOES NOT HOLD ANY LIABILITY.

PRICE:

₹ 2000/- (Two thousand rupees only).

Printed by :

Peacock Advertising India Pvt Ltd.

158 & 159, 3rd Main, 7th Cross, Chamarajpet,
Bengaluru – 560 018 | Ph: 080 - 2662 0566
Web: www.peacockgroup.in



B.S. YEDIYURAPPA
CHIEF MINISTER



VIDHANA Soudha
BENGALURU - 560 001

NO: CM/MS/84/2019

Date : 14.11.2019

MESSAGE

I am happy to know that Karnataka Biodiversity Board is bringing out scientific book titled "A Flora of Karnataka"- Volume-2: Checklist of Angiosperms and Gymnosperms as a part of implementation of Biodiversity Act 2002.

It is appreciable that Karnataka Biodiversity Board has taken good initiative to create bio resource database which would serve as ready reference in action towards conservation of Biological Diversity, sustainable utilization fair and equitable sharing of benefits arising out of the use of biological resource.

I extend my best wishes to the Karnataka Biodiversity Board and hope this publication would be helpful to all the stake holders involved in conservation of biodiversity.

(B.S.YEDIYURAPPA)

Sri Ananth Hegde AShisara
Chairman,
"Vana vikasa" 18th cross,
Malleshwaram,
Bengaluru – 560 003.



ಸಿ.ಸಿ. ಪಾಟೀಲ್

ಗಣೀ ಮತ್ತು ಭೂವಿಜ್ಞಾನ, ಅರಣ್ಯ, ಪರಿಸರ
ಮತ್ತು ಜೀವಿಶಾಸ್ತ ಹಾಗೂ ಗದಗ ಮತ್ತು
ವಿಜಯಪುರ ಜಿಲ್ಲೆ ಉಸ್ತುವಾರಿ ಸಚಿವರು



ಕೊಲಡಿ ಸಂಖ್ಯೆ: 244, 245

2ನೇ ಮಹಡಿ, ವಿಕಾಸಸೌಧ

ಬೆಂಗಳೂರು-560 001

ದೂರವಾಣಿ: 080-2225 6093

080-2203 4439

ಇ-ಮೆಲ್ಲೋ: ccpatilminister@gmail.com

ಸಂಖ್ಯೆ: ಗಭೂವಿಅಪಜೀನ/491A/2019

ದಿನಾಂಕ: 14-11-2019

ಸಂದೇಶ

ಜೀವವೈವಿಧ್ಯತೆಯ ಸಂರಕ್ಷಣೆ, ಸುಸ್ಥಿರ ಬಳಕೆ ಹಾಗೂ ಜೈವಿಕ ಸಂಪನ್ಮೂಲಗಳ ಬಳಕೆಯಿಂದ ಪಡೆದ ಲಾಭಾಂಶಗಳ ನ್ಯಾಯೋಚಿತ ಹಂಚಿಕೆಯಾಗಬೇಕು ಎಂಬ ಉದ್ದೇಶದಿಂದ ಜೈವಿಕ ವೈವಿಧ್ಯ ಅಧಿನಿಯಮ 2002 ನ್ನು ರಾಷ್ಟ್ರೀಯ ಅನುಷ್ಠಾನಗೊಳಿಸಲಾಗುತ್ತಿದೆ. ಈ ಮೂಲಕ ಪರಿಸರ ಸಮರ್ಪಣೆಯನ್ನು ಕಾಪಾಡುವಲ್ಲಿ ಜೈವಿಕ ವೈವಿಧ್ಯ ಅಧಿನಿಯಮವು ಪ್ರಮುಖ ಪಾತ್ರವಹಿಸುತ್ತದೆ.

ಜೀವವೈವಿಧ್ಯ ಸಂರಕ್ಷಣೆಗೆ ಸಂಬಂಧಿಸಿದ ವಿವಿಧ ತಾಂತ್ರಿಕ ಮತ್ತು ದತ್ತಾಂಶಗಳ ಮಾಹಿತಿಯನ್ನು ಪ್ರಕಟಣೆ ಮಾಡುವುದರಿಂದಾಗಿ ಜೈವಿಕ ಸಂಪನ್ಮೂಲಗಳ ಅತಿಯಾದ ಬಳಕೆಯನ್ನು ನಿಯಂತ್ರಣಾದಲ್ಲಿಡಲು ಸಹಕಾರಿಯಾಗುತ್ತದೆ. ಮಂಡಳಿಯು ಹೊರತರುತ್ತಿರುವ “Flora of Karnataka- A Checklist Vol-2, Angiosperms & Gymnosperms” ಎಂಬ ವೈಜ್ಞಾನಿಕ ಮಸ್ತಕವು ಜೀವವೈವಿಧ್ಯ ಸಂರಕ್ಷಣೆ ಹೊಣೆ ಹೊತ್ತಿರುವ ವಿವಿಧ ವಿಜ್ಞಾನ ಸಂಸ್ಥೆಗಳು ಮತ್ತು ವಿದ್ಯಾರ್ಥಿಗಳಿಗೆ ಅಧ್ಯಯನ ಮಾರ್ಗದರ್ಶಿಯಾಗಲಿದೆ.

ಈ ವೈಜ್ಞಾನಿಕ ಮಸ್ತಕವನ್ನು ಸಿದ್ಧಪಡಿಸಿ ಪ್ರಕಟಣೆಗೊಳಿಸಿದ ಕನಾರ್ಕ ಜೀವವೈವಿಧ್ಯ ಮಂಡಳಿಯನ್ನು ಅಭಿನಂದಿಸುತ್ತೇನೆ.

(ಸಿ.ಸಿ. ಪಾಟೀಲ್)

ಸದಸ್ಯ ಕಾರ್ಯದರ್ಶಿ,
ಕನಾರ್ಕ ಜೀವವೈವಿಧ್ಯ ಮಂಡಳಿ.
ಬೆಂಗಳೂರು.



Karnataka Biodiversity Board

ANANTH HEGDE ASHISARA

Chairman (State Minister Rank)



Shri Ananth Hegde Ashisara,
Chairman,
Karnataka Biodiversity Board,
Government of Karnataka.

'Vanavikas', Ground floor,
18th cross, Malleshwaram,
Bengaluru- 560003,
Ph: 080-23448783, 23448765

MESSAGE

Karnataka Biodiversity Board was established in the year 2004 and subsequently, the state of Karnataka notified the Karnataka Biological Diversity Rules in the year 2005. The Board has initiated a number of Biodiversity Conservation measures under the directions of Central and State authority. We have declared 4 Biodiversity Heritage Sites, initiated Biodiversity awards etc. The present work was commissioned under the provisions of Rule13(8) of Karnataka Biological Diversity Rules, 2005 which provides for collection, compilation and publishing of technical and statistical manuals or guides relating to conservation. The Board has been conducting various programmes, events, trainings, workshops which necessitate the requirement of outreach materials in the form of pamphlets, brochures and scientific books. This book is first of its kind in the entire country, will also serve as a checklist and field guide, especially for educational institutions and researchers. It is a commendable work which forms an instrument for measuring and conserving the plant wealth of Karnataka.

I appreciate the Member Secretary, Karnataka Biodiversity Board experts and its staff for bringing out such a useful publication.

(Ananth Hegde Ashisara)



ಡಾ. ಸಂದೀಪ್ ದವೆ, ಭಾ.ಆ.ಸೇ.
ಕರ್ನಾಟಕ ಸರ್ಕಾರದ ಅಧಿಕಾರಿ
ಅರಣ್ಯ, ಜೀವಿ ಪರಿಸ್ಥಿತಿ ಮತ್ತು ಪರಿಸರ ಇಲಾಖೆ



Dr. SANDEEP DAVE, I.A.S.
Additional Chief Secretary to Govt
Forest, Ecology and Environment
Department

FOREWORD

About 60% of the Western Ghats is present in Karnataka State, with this large part of the peninsular green cover coupled with inland forest plateau enriched by the seven river systems, the State of Karnataka showcases a wider floral wealth harboring highest number of apex predators all of which is conserved by a framework of various statutes.

The present book titled ‘Flora of Karnataka – A checklist of Angiosperms & Gymnosperms, Volume – 2’ gives a concise information on the present day seed-bearing plants - the Angiosperms - with highest number of species while also capturing information on the old world naked seed bearing plants – the Gymnosperms - which are the tallest and largest living plants on earth. These plants have the natural ability to produce seeds surrounded by nutritive tissue and coated with a seed coat. Gymnosperm plants were present 200 million years before the arrival of the Angiosperms, they have somehow acclimatized and managed to survive hitherto with a few countable specimens. On the other hand, the Angiosperms are the successors of Gymnosperms and are the modern day flower bearing plants, serving mankind with various uses especially as food and medicine. The contributions and services of experts engaged in formulating the methodology and compiling this huge data is praise worthy. The book not only provides an exhaustive list of plants from the above two groups, but also contains information on their taxonomy, synonyms, habitat including their distribution as well.

I would like to acknowledge Karnataka Biodiversity Board for its substantiating efforts in preparation of this book.



(Sandeep Dave)

Dr. Virender Singh
Member Secretary
Karnataka Biodiversity Board
Government of Karnataka.

'Vanavikas' Ground floor
18th Cross, Malleshwaram
Bengaluru – 560 003
Ph: 080-23448783, 23448765

PREFACE

The Biological Diversity Act, 2002 came into existence with three objectives i.e., 1) Conservation of Biological diversity, 2) Sustainable use of components and 3) fair and equitable sharing of benefits arising out of use of biological resources. Out of the three objectives, conservation is considered as the prime and foremost important since sustainable use and sharing of benefits is only possible if there is a resource database. The Board, under the provisions of Rule 13(11) of the Karnataka Biological Diversity Rules, 2005 which provides for building up database and to create information and documentation system for biological resources to ensure the effective management, promotion and sustainable use of biological resource and knowledge has considered that effective documentation is a key for future conservation. The Board felt the need to compile the baseline profile of plants of Karnataka which will serve as a ready reference for conservation. The work is an outcome of contributions made by some of the leading expert scientists in the country.

The present book titled 'Flora of Karnataka – A Checklist of Angiosperms & Gymnosperms, Volume – 2' with beautiful pictures is mainly focused on aiding the younger generation and scientific institutions with information on Gymnosperms - the magnificent huge trees and the Angiosperms – the modern flower bearing plants with highest recorded species.



(Virender Singh, IFS)

Authors

1. Dr. M. Sanjappa, INSA Senior Scientist, Mahatma Gandhi Botanical Garden, GVK, Bengaluru – 560 065.
2. Dr. A.N. Sringeswara, Curator, Mahatma Gandhi Botanical Garden, GVK, Bengaluru – 560 065.

Photograph Credits

Dr. M. Sanjappa, GVK.

Dr. A.N. Sringeswara, GVK.

Dr. S.R. Yadav, Shivaji University, Kollapur.

Acknowledgements

1. Sri R.M.N. Sahai, IFS (Retd.), Former Chairman, Karnataka Biodiversity Board for his vision which resulted in this book.
2. Sri R.K. Singh, IFS, APCCF (Working Plan) & Former Member Secretary, Karnataka Biodiversity Board for initiating the work.
3. Dr. M. Sanjappa, INSA Senior Scientist, Mahatma Gandhi Botanical Garden, GVK for taking the lead in gathering the information.
4. Dr. Virender Singh, IFS, APCCF & Member Secretary, Karnataka Biodiversity Board for compiling the information and publishing.
5. Smt Sahana S. Hegde, Deputy Director (Horticulture), Karnataka Biodiversity Board for facilitation and convening the meetings.
6. Sri Pritham S, Research Associate (Zoology), Karnataka Biodiversity Board for coordination and cover page design.
7. Members of Publication Committee, Karnataka Biodiversity Board for various inputs and suggestions:
 - a) Dr. M. Sanjappa, INSA Senior Scientist, Mahatma Gandhi Botanical Garden, GVK, Bengaluru – 560 065.
 - b) Prof. C. Kameswara Rao, Executive Secretary, FBAE, No. 1 Gupta's layout, Southend Road, Bengaluru – 560 004.
 - c) Prof. M. Narayana Swamy, Professor & Head, Dept. of Veterinary Physiology, Veterinary College, Hebbal, Bangalore – 560 024.
 - d) Dr. S.N. Hegde, Former Professor & Chairman, Dept. of Zoology, University of Mysore, No. 109, Vikasa, 7th Main, 3rd Cross, Bogadi II Stage, Mysore – 570 026.
 - e) Dr. B.N. Manjunatha, Professor of Agricultural Extension and Editor, Communication Centre, UAS, GVK, Bengaluru – 560 065.
 - f) Smt Sahana S. Hegde, Deputy Director (Horticulture), Karnataka Biodiversity Board, Malleshwaram, 18th Cross, Bengaluru – 560 003.
8. Dr. A.N Sringswara, Curator, Mahatma Gandhi Botanical Garden, GVK, Bengaluru – 560 065 for preparation of index, photo plates & proof reading

Composition of the Board

The present Board has been constituted vide order No:
FEE 203 ENV 2016, Bengaluru. Dated: 13.02.2017.

| Sl. No | Board Members | Designation |
|-------------------|---|-----------------------------------|
| 1 | Sri Ananth Hegde Ashisara, Chairman, Karnataka Biodiversity Board, Ground Floor, Vanavikas, 18 th Cross, Malleshwaram, Bengaluru – 560 003. | Chairman |
| 2 | Sri Sandeep Dave, IAS, Additional Chief Secretary to Government, Forest, Ecology and Environment Department, MS Building, Bengaluru – 560 001. | Ex-officio Member |
| 3 | Sri Vijay Kumar Gogi, IFS, Principal Secretary to Government, Forest, Ecology and Environment Department, MS Building, Bengaluru – 560 001. | Ex-officio Member |
| 4 | Sri Rajender Kumar Kataria, IAS, Secretary to Government, Department of Agriculture, MS Building – 560 001. | Ex-officio Member |
| 5 | Sri Punati Sridhar, IFS, Principal Chief Conservator of Forests (HoFF), Aranya Bhavan, 18 th Cross, Malleshwaram, Bengaluru – 560 003. | Ex-officio Member |
| 6 | Smt Meenakshi Negi, IFS, Commissioner, Department of AYUSH, Dhanvantari Road, Ananda Rao Circle, Bengaluru – 560 009. | Ex-officio Member |
| 7 | Dr. C. G. Kushalappa, Dean Forestry, University of Agriculture and Horticultural Sciences (Shivamogga), College of Forestry (Ponnampet), Kunda Road, Ponnampet – 571 216. | Non-officio Member |
| 8 | Dr. M. Mahadeva Murthy, Professor, Dept. of Forestry and Environmental Science, UAS, GKVK, Bengaluru – 560 065. | Non-officio Member |
| 9 | Dr. Jayakara Bhandary M, Associate Professor, Dept. of Botany, Government First Grade College, Car Street, Mangaluru – 575 001. | Non-officio Member |
| 10 | Prof. M. Narayana Swamy, Professor & Head, Dept. of Veterinary Physiology, Veterinary College, Hebbal, Bengaluru – 560 024. | Non-officio Member |
| 11 | Dr. Niranjana Murthy, Professor & Scheme Head, AICRP on Potential Crops, RIOF Building, UAS, GKVK, Bangalore – 560 065. | Non-officio Member |
| 12 | Smt Anita K. Arekal, IFS, Additional Principal Chief Conservator of Forests (Social Forestry), Aranya Bhavan, 18 th Cross, Malleshwaram, Bengaluru – 560 003. | Special Invitee |
| 13 | Sri B.T Khanapure, Drugs Controller, Drugs Control Department Palace Road, Near Karnataka Public Service Commission, Bengaluru – 560 001. | Special Invitee |
| 14 | Dr. Virender Singh, IFS, Member Secretary, Karnataka Biodiversity Board, Ground Floor, Vanavikas, 18 th Cross, Malleshwaram, Bengaluru – 560 003. | Member Secretary - Convener |

Flora of Karnataka, A Checklist
Volume 2 : Gymnosperms & Angiosperms.

CONTENTS

| | Page No. |
|---|------------|
| 1. Gymnosperms of Karnataka – A Checklist..... M. Sanjappa | 1 - 6 |
| 2. Angiosperms of Karnataka – A Checklist..... M. Sanjappa & A. N. Sringeswara | 7 - 815 |
| 3. Index..... | 816 - 1002 |

Gymnosperms of Karnataka

A Checklist

M. Sanjappa

Mahatma Gandhi Botanical Garden
University of Agricultural Sciences, GKVK, Bengaluru 560 065

Support by

Karnataka Biodiversity Board

INTRODUCTION

The angiosperms and gymnosperms are together called Spermatophytes or seed bearing plants. Gymnosperms (literally means ‘naked seeds’, in Greek *gymnos* meaning ‘naked’ and *sperma* ‘seed’) are a group of seed bearing plants and are called ‘naked seeded’ plants, unlike angiosperms in which the seeds are covered or enclosed by fruits. In gymnosperms seeds are generally borne in the form of cones or rarely solitary as in *Ginkgo biloba*. Characteristically the roots of some genera are known to be associated with mycorrhiza as in *Pinus* and in some *Cycas* species the specialised roots called corolloid roots have associated nitrogen fixing cyanobacteria. The gymnosperms bear a variety of characteristic leaves mostly needle-like with thick cuticles, flat triangular scale-like, broad strap-shaped. The leaves of Cycads appear like palm leaves and those of *Gnetum* like those of dicotyledonous angiosperms. The reproductive structures are called a strobilus or a cone. The male and female cones are always distinct, either borne on the same plant or on separate plants. The gymnosperms are represented by 12 families (under 4 subclasses and 8 orders), 83 known genera with ca 1080 known species (Christenhusz, M.J.M., Reveal, J.L., Farjon, A., Gardner, M.F., Mill, R.R. & Chase, M.W. 2011. A new classification and linear sequence of extant gymnosperms. *Phytotaxa* 19: 55–70.) distributed throughout the globe. Economically gymnosperms are very important, the conifers yield timber, resins, paper pulp, etc. and are used as raw materials in industry for producing a variety of products. Many species (of *Amentotaxus*, *Taxus*, *Ginkgo*, *Ephedra*, *Pinus*, *Cupressus*, *Juniperus*, etc.) are also used in various systems of medicine. Taxol from species of *Taxus* (including *Taxus wallichiana* from Indian Himalayas) is widely used for treating cancers.

In India, majority of the gymnosperms are confined to Himalayas. However, species of genera *Cycas*, *Gnetum*, *Nageia*, and a species of *Ephedra* are distributed in plains (including hot desert) and hills. About 70 species are recorded to occur wild in different parts of India and almost as many are cultivated as ornamentals. In Karnataka, 26 species under 8 are documented with majority of them cultivated as ornamentals and/or in botanical gardens. Only 5 species (*Cycas circinalis* L., *Cycas indica* A.Lindstr. & K.D.Hill, *Cycas swamyi* Rita Singh & P. Radha, *Gnetum scandens* Roxb. and *Gnetum ula* Brongn.) are known in wild. *Cycas indica* (from Arasikere hills, Hassan District) and *Cycas swamyi* (from Melkote hills, Mandya District) are exclusively endemic to Karnataka.

ANALYSIS OF GYMNOSPERMS

| Family | No. of Genera | No. of Species | No. of Subspecies | No. of Varieties | Endemic species |
|----------------------|---------------|----------------|-------------------|------------------|-----------------|
| Araucariaceae | 2 | 5 | - | - | - |
| Cupressaceae | 1 | 4 | - | - | - |
| Cycadaceae | 2 | 10 | - | - | 2 |
| Gnetaceae | 2 | 3 | - | - | - |
| Pinaceae | 1 | 4 | - | - | - |
| Total | 8 | 26 | - | - | - |

Araucariaceae

Araucaria Juss.

Araucaria bidwillii Hook, London J. Bot. 2: 503.1843.

Columbea bidwillii (Hook.) Carriere, Traite Gen. Conif. ed. 2: 601.1867.

Marywildea bidwillii (Hook.) A.V.Bobrov & Melikyan, Komarovia 4: 58.2006.

Distribution: Karnataka (cultivated as an ornamental), native to Australia.

Local names/Common names/Vernacular names: *Bidwill's araucaria*, *The bunya pine*, *Bunya Araucaria* (English).

Araucaria columnaris (G.Forst.) Hook., Bot. Mag. 78: t. 4635.1852.

Cupressus columnaris G.Forst., Fl. Ins. Austr. 67.1786.

Dombeya excelsa Lamb., Descr. Pinus 87.1806.

Araucaria excelsa (Lamb.) R.Br., Hortus Kew. 5: 412.1813.

Columbea excelsa (Lamb.) Spreng., Syst. Veg. 4(2.): 215.1827.

Eutacta excelsa (Lamb.) Link, Linnaea 15: 544.1841.

Araucaria cookii R.Br. ex Endl., Syn. Conif. 188.1847.

Distribution: Karnataka (cultivated as an ornamental), native to New Caledonia

Local names/Common names/Vernacular names: *Cook's pine*, *Cook's araucaria*, *Columnar araucaria*, *New Caledonia pine*, *The Coral reef araucaria*, (English).

Araucaria cunninghamii Aiton ex D.Don, Descr. Pinus ed. 2, 3: t. 79.1837.

Eutacta cunninghamii (Aiton ex D.Don) Link, Linnaea 15: 545.1841.

Eutassa cunninghamii (Aiton ex D.Don) G.Don, Hort. Brit. ed. 3: 623.1839.

Distribution: Karnataka (cultivated as an ornamental), native to Australia

Local names/Common names/Vernacular names: *Cunningham's Araucaria*, *Colonial Pine*, *Hoop Pine*, *Moreton Bay Pine*, *Queensland Pine* (English).

Juniperus

Juniperus chinensis L., Mant. Pl. 127.1767.

Juniperus cernua Roxb., Fl. Ind. ed. 2. 3: 839.1832.

Sabina dimorpha (Roxb.) Antoine, Cupress.-Gatt. 70.1857.

Distribution: Karnataka (cultivated as an ornamental), native to E Asia and S Russia.

Local names/Common names/Vernacular names: *Chinese Juniper* (English)

Juniperus communis L., Sp. Pl. 1040.1753.

Distribution: Karnataka (cultivated as an ornamental), wild in W Himalaya, cultivated elsewhere in India; Temperate regions of Africa, Asia, Europe, N America.

Local names/Common names/Vernacular names: *The common juniper* (English)

Cupressaceae

Cupressus L.

Cupressus funebris Endl., Syn. Conif. 58.1847.

Platycyparis funebris (Endl.) A.V.Bobrov & Melikyan, Komarovia 4: 73.2006.

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout India; native to China

Local names/Common names/Vernacular names: *Chinese Weeping Cypress* (English)

Cupressus lusitanica Mill., Gard. Dict. ed. 8, 3.1768.

Cupressus glauca Lam., Encycl. 2: 243.1786.

Callitropsis lusitanica (Mill.) D.P.Little, Syst. Bot. 31: 474.2006.

Distribution: Karnataka (cultivated as an ornamental), native to Mexico and C America.

Local names/Common names/Vernacular names: *White Cedar, Mexican White Cedar* (English)

Cupressus sempervirens L., Sp. Pl. 1002.1753.

Distribution: Karnataka (cultivated as an ornamental), native to the eastern Mediterranean region

Local names/Common names/Vernacular names: *The Mediterranean cypress, Italian cypress, Tuscan cypress, graveyard cypress, pencil pine, Persian cypress* (English)

Cupressus torulosa D.Don, Prodr. Fl. Nepal. 55.1825.

Cupressus nepalensis Loudon, Encycl. Trees Shrubs 1118.1842.

Cupressus sempervirens var. *indica* Parl., Prodr. 16(2): 471.1868.

Cupressus doniana Hook.f., Fl. Brit. India 5: 645.1888.

Cupressus lusitanica subsp. *torulosa* (D.Don) Silba & D.Z.Fu, Acta Bot. Yunnan. 28: 470.2006.

Cupressus lusitanica subsp. *torulosa* (D.Don) Silba & Brian Chen, J. Int. Conifer Preserv. Soc. 12: 85.2005.

Cupressus sempervirens subsp. *indica* (Parl.) Silba, J. Int. Conifer Preserv. Soc. 12: 97.2005.

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout India; native to Himalaya.

Local names/Common names/Vernacular names: *The Himalayan cypress, Bhutan cypress* (English)

Cycadaceae

Cycas L.

Cycas circinalis L., Sp. Pl. 1188.1753; B.D. Sharma et al., Fl. Karnataka Anal. 263.1984.

Plate 1 A, B & C

Distribution: Karnataka (Bengaluru, Hassan, Mandya, Mysore, Tumkur), Kerala, Maharashtra, Tamil Nadu, also cultivated as an ornamental), endemic.

Local names/Common names/Vernacular names: *Goddu ichala* (Kannada).

Cycas indica A.Lindstr. & K.D.Hill, Telopea 11: 481.2007.

Karnataka (Hassan), endemic.

Cycas revoluta Thunb., Nova Acta Regiae Soc. Sci. Upsal. 4: 40.1783.

Plate 1 D & E

Distribution: Karnataka (cultivated as an ornamental), native to China and Japan.

Local names/Common names/Vernacular names: *Sotetsu* (Japanese), *Japanese sago palm, King sago, Sago cycad, Sago palm* (English).

Cycas rumphii Miq., Bull. Sci. Phys. Nat. Néerl. 2: 45.1839.

Distribution: Karnataka (cultivated as an ornamental), native to SE Asia and Indian Ocean Islands

Cycas swamyi Rita Singh & P. Radha, Bot. J. Linn. Soc. 158: 431.2008.

Distribution: Karnataka (Mandya: Melkote), endemic.

Zamia L.

Zamia acuminata Oerst. ex Dyer, Biol. Cent.-Amer., Bot. 3: 194.1884.

Distribution: Karnataka (Cultivated as an ornamental), native to Central America (Costa Rica, Nicaragua, Panama).

Zamia angustifolia Jacq., Collectanea 3: 263.1791.

Palmifolium angustifolium (Jacq.) Kuntze, Revis. Gen. Pl. 2: 803.1891.

Distribution: Karnataka (Cultivated as an ornamental), native to Bahamas and Cuba.

Local names/Common names/Vernacular names: Eleuthera Coonti (English).

Zamia fischeri Miq. ex Lem., Hortus Van Houtteanus 1(1): 20.1845.

Palmifolium fischeri (Miq. ex Lem.) Kuntze, Revis. Gen. Pl. 2: 803.1891.

Distribution: Karnataka (Cultivated as an ornamental), native to Mexico.

Zamia furfuracea L.f. ex Aiton, Hort. Kew. 3: 477.1789.

Plate 1G

Palmifolium furfuraceum (L.f. ex Aiton) Kuntze, Revis. Gen. Pl. 2: 803.1891.

Distribution: Karnataka (Cultivated as an ornamental), native to Mexico.

Local names/Common names/Vernacular names: *Cardboard Cycad*, *Cardboard Plant*, *Cardboard Sago*, *Jamaican Sago*, *Mexican Cycad* (English).

Zamia pygmaea Sims, Bot. Mag. 42: t. 1741.1815.

Palmifolium pygmaeum (Sims) Kuntze, Revis. Gen. Pl. 2: 803.1891.

Distribution: Karnataka (Cultivated as an ornamental), native to Cuba

Local names/Common names/Vernacular names: *Guayiga*, *Comptie*, *Florida arrowroot* (English).

Gnetaceae

Gnetum L.

Gnetum scandens Roxb., Hort. Bengal. 66.1814.

Gnetum funiculare Brongn., Voy. Monde Phan.: 12.1829.

Distribution: Karnataka (Belgaum, Chickmagalur, Dakshina Kannada, Shimoga, Udupi?, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Maharashtra, Tamil Nadu, N E India; S E Asia.

Gnetum ula Brongn., Voy. Monde Phan.: 12.1829; B.D.Sharma et al., Fl. Karnataka Anal. 263.1984.

Plate 1 F

Distribution: Karnataka (Belgaum, Chickmagaluru, Dakshina Kannada, Shimoga, Udupi?, Uttara Kannada), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu, N E India; S E Asia.

Local names/Common names/Vernacular names: *Kodakan balli*, *Navaru katte* (Kannada), *Ombal* (Kunabi).

Ephedra L.

Ephedra foliata Boiss. ex C.A.Mey., Mém. Acad. Imp. Sci. Saint-Pétersbourg, Sér. 6, Sci. Math., Seconde Pt. Sci. Nat. VI, 6: 107.1846.

Distribution: Karnataka (cultivated), Gujarat, Punjab, Rajasthan; C Asia, Pakistan, NW Africa, West Asia.

Local names/Common names/Vernacular names: *Shrubby horse tail* (English).

Pinaceae

Pinus L.

Pinus caribaea var. **hondurensis** (Sénécl.) W.H.Barrett & Golfari, *Caribbean Forester* 23: 65.1962; Torvi, G.S.Kariyappa & B.K.M.Amanulla, Indian Forest. 587. 1998.

Pinus hondurensis Sénécl., *Conifères* 126.1868.

Pinus caribaea subsp. *hondurensis* (Sénécl.) Silba, *J. Int. Conifer Preserv. Soc.* 16: 17.2009.

Distribution: Karnataka (cultivated), native to Mexico, S America, The Carribeana.

Local names/Common names/Vernacular names: *Honduras pine* (English).

Pinus pinaster Aiton, *Hort. Kew.* 3: 367. 1789;Torvi, G.S.Kariyappa & B.K.M.Amanulla, Indian Forest. 587. 1998.

Distribution: Karnataka (cultivated), native to Mediterranean region.

Local names/Common names/Vernacular names: *The maritime pine*, *Cluster pine* (English).

Pinus kesiya Royle ex Gordon, *Gard. Mag. & Reg. Rural Domest. Improv.* 16: 8.1840;Torvi, G.S.Kariyappa & B.K.M.Amanulla, Indian Forest. 587. 1998.

Pinus khasyana Griff., *Itin. Pl. Khasyah Mts.* 58.1848.

Pinus kasya Royle ex Parl., *Prodr.* 16(2): 390.1868.

Pinus insularis var. *khasyana* (Griff.) Silba, *Phytologia* 68: 51.1990.

Distribution: Karnataka (cultivated), Arunachal Pradesh, Meghalaya;China, SE Asia.

Local names/Common names/Vernacular names: *Khasi pine*, *Benguet pine*, *Three-needed pine* (English).

Pinus oocarpa Schiede, *Linnaea* 12: 491.1838; Torvi, G.S.Kariyappa & B.K.M.Amanulla, Indian Forest. 588. 1998.

Pinus oocarpa var. *manzanoi* Martínez, *Anales Inst. Biol. Univ. Nac. México* 11: 70.1940.

Pinus oocarpoides Lindl. ex Loudon, *Encycl. Trees Shrubs* 1118.1842.

Pinus oocarpa var. *oocarpoides* (Lindl. ex Loudon) Endl., *Syn. Conif.* 152. 1847.

Pinus oocarpa subsp. *manzanoi* (Martínez) Silba, *J. Int. Conifer Preserv. Soc.* 16: 29.2009.

Distribution: Karnataka (cultivated), native to Mexico and C America.

Local names/Common names/Vernacular names: *Mexican yellow pine*, *Hazelnut pine* (English).



Plate 1: A. *Cycas circinalis* L.; B. *Cycas circinalis* L. - Female cone; C. *Cycas circinalis* L. - Male cone; D. *Cycas revoluta* Thunb.; E. *Cycas revoluta* Thunb. - Female cone; F. *Gnetum ula* Brongn.; G. *Zamia furfuracea* L.f. ex Aiton

Angiosperms of Karnataka

A Checklist

M. Sanjappa & A. N. Sringswara

With Assistance from
Sahana Vishwanath & M.D. Rajanna

Support by

Karnataka Biodiversity Board

Introduction

Plants are one of Earth's greatest resources. They are the main sources of food, medicines and industrial materials with great economic and cultural importance. They stabilize ecosystems and form the habitats that sustain the planet's human, animal and microbial life.

India is a signatory to CBD and, therefore, it is obligatory to document all plant resources of the country not only at national level but also at state, district and Panchayat level to evolve strategies for conservation, sustainable utilisation and for equitable sharing of benefits arising out of their commercial utilisation and associated traditional knowledge.

The Biodiversity Act 2002 (and Biodiversity Rules 2004) mandates establishment of National Biodiversity Authority, State Biodiversity Boards (SBBs) and Biodiversity Management Communities (BMC's) to regulate sustainable utilisation of bioresources and facilitate implementation of the said Act and Rules. Thus in order to sustainably utilise the bioresources and also facilitate equitable sharing of benefits and conservation all states need to document all bioresources. Karnataka state has been in the forefront to form State Biodiversity Board as well as BMCs to implement the Act and Rules. Most importantly, preparation of the *The Peoples Biodiversity Registers* by BMC's within their jurisdiction is basic requirement for conservation, sustainable utilisation and equitable sharing of benefits. To achieve this there is a need for preparing a check-list of all bioresources of the state documented so far, the present work is an effort in this direction by Karnataka Biodiversity Board.

Karnataka state (erstwhile State of Mysore) lies between 11° 36 and 18° 25 N latitude and 74° 10 and 78° 35 E longitudes. The state is bordered by the Arabian Sea to the west, Goa to the north west, Maharashtra to the north, Telangana to the North east Andhra Pradesh to the east, Tamil Nadu to the south east, and Kerala to the south west. The state covers an area of 1, 92, 204 square kilometres and thus it is the seventh largest state with 5.83 per cent of the total geographical area of India. The total area is divided into four administrative divisions comprising of 30 districts, 220 taluks and 6068 gram panchayats. Bengaluru city is the State capital. Belagavi with an area of 13, 415 square kilometres is the largest, whereas Bengaluru Rural, with an area of 2,259 square kilometres is the smallest. The longest North-South distance is about 750 km and widest East-West distance is about 400 km. According to the 2011 census the state has 61,130,704 inhabitants and thus it is the eighth largest state of India by population.

Physiographically, the state has three principal geographical regions viz. the coastal or Karavali region, the hilly/mountainous or Malenadu region and the plains or Bayaluseeme region. The Karavali region is about 320 km long and 5 – 65 km broad with an average elevation of 70 m covering western fringes of Dakshina Kannada, Udupi and Uttara Kannada districts. The Malenadu region comprising the Western Ghats with highest point of 1,929 metres (6,329 ft) in Karnataka is the Mullayanagiri hills in Chikkamagaluru district. The major rivers originate in Western Ghats are Kaveri (Cauveri), Tungabhadra, Krishna, Malaprabha and the Sharavathi. The plains or Bayaluseeme region of the Deccan Plateau occupies the bulk of the state (the second-largest arid region in India).

Karnataka consists of four main types of geological formations, viz., the Archean complex, the Proterozoic, the Deccan trappean and intertrappean deposits and recent laterites and alluvial deposits. About 60% of the state is composed of the Archean complex, which consists of gneisses, granites and charnockite rocks. Laterite cappings that are found in many districts over the Deccan traps formed after the cessation of volcanic activity in the early tertiary period. Raychaudhuri et al. (1963), divided the soils of the state into main groups viz., Coastal alluvium, Laterite soils, red soils, mixed red, yellow and grey soils, black soils (mainly in Bagalkote, Belagavi, Bidar, Dharwad, Gadag, Gulbarga, Koppala, Raichur, Vijayapura and Yadgir districts).

Karnataka state has tropical monsoon climate and experiences four seasons. The winter during January and February, summer during March and May, the monsoon season between June and September and the post-monsoon season from October and November. Metereorologically, the state is divided into three zones:

coastal, north interior and south interior. Of these, the coastal zone receives the heaviest rainfall with an average of about 3,638 mm per annum, far in excess of the state average of 1,139 mm. Agumbe in Shivamogga district receives the highest annual rainfall in the state and second highest in India. The climate is mild in the plains. The mean daily minimum temperature is least during January (14.5°C), which very gradually increases during hottest month May (22°C) and decreases during monsoon months to the lowest in January. While, the mean daily maximum temperature is highest in April (35°C) it decreases gradually during monsoon months of July-August (27.5°C) and rises slightly in January and rapidly thereafter to the highest in April again. However, the highest recorded temperature was 45.6°C at Raichur and the lowest recorded temperature was 2.8°C at Bidar. The relative humidity in the coastal region and Western Ghats is always much higher compared to plains.

About 20% of the state's geographic area (i.e., 38, 28, 430 square kilometres) of Karnataka is covered by forests. The forests are classified as reserved, protected, unclassed, village and private forests. The percentage of forested area is slightly less than the all-India average of about 21%, and significantly less than the 33.33% prescribed in the National Forest Policy. The Western Ghats, which covers about 60% of forest area of Karnataka, is recognized as one of the 18 mega diversity centres of the world occupying major part of the Indo-Sri Lankan hotspot. The extent and types of forest in Karnataka are: Evergreen and Semi-evergreen 5, 80, 000 ha. (15.11%), Moist Deciduous 5, 78, 000 ha. (14.96%), Dry Deciduous 7, 27, 000 ha. (18.64%), Scrub and Thorny 8, 18, 000 ha. (21.11%), Un-wooded 11, 25, 430 ha. (30.18%). According to their legal status, the forest areas in Karnataka are classified as: Reserve Forests 28,68,996 ha., Protected Forests 3,93,072 ha., Un-classed Forests 5,23,100 ha., Village Forests 12,420 ha. and Forests in private lands 30,842 ha. (Anonymous, *Annual plan of operation for utilisation of state CAMPA funds*, 2012).

The vegetation of Karnataka is diverse ranging from tropical wet evergreen to scrub forests depending upon the climate, altitude and edaphic factors. It can be broadly divided into Southern Tropical Moist Forests (Southern tropical wet evergreen forests, Southern tropical semi-evergreen forests, Southern tropical moist deciduous forests, littoral mangroves) Southern Tropical dry forests (Southern tropical dry deciduous forests), Southern Montane forests), which are further subdivided into 10 types and 9 subtypes. The composition of all these vegetation/forest types are discussed in detail both by Sharma et al. (1984) and Saldanha (1984). Although the basic vegetation types discussed in these works are mainly following Champion and Seth (1968), the latter also mentioned the types earlier recognised by Legris and Pascal (1964). However, the vegetation types of Champion and Seth (1968) are further modified by Forest Survey of India.

The Karnataka State government had taken steps to conserve biodiversity rich areas, pristine landscapes and many wildlife rich areas, which encompass a biosphere reserve, national parks, Wildlife sanctuaries, conservation reserves, community reserves, tiger reserves and elephant corridors. They are as given below:

Nilgiri Biosphere Reserve (Part of Wynad, Nagarhole, Bandipur and Mudumalai, Nilambur, Silent Valley and Siruvani Hills) spreading over 5520 sq. km shared among states of Karnataka, Kerala and Tamil Nadu was declared in 1986. Apart from this interstate Biosphere Reserve, Karnataka has 5 national parks (Anishi, Bandipur, Bannerghatta, Kederemukh and Rajiv Gandhi); 22 Wildlife Sanctuaries (Adichunchunagiri, Mandya; Arabithittu, Mysuru; Attiveri, Uttara Kannada; Bhadra, Chikkamagaluru & Shivamogga; Bhimgad, Belagavi; Biligiri Rangaswamy Temple, Chamarajanagar; Brahmagiri, Kodagu; Cauvery, Mysuru, Bengaluru Rural & Mandya; Dandeli, Uttara Kannada; Daroji Bear, Bellari; Ghataprabha Bird, Belagavi; Gudavi, Shivamogga; Melkote Temple, Mandya; Mookambika, Udupi; Nugu, Mysuru; Ranganathittu Bird Sanctuary, Mysuru; Pushpagiri, Kodagu & Dakshina Kannada; Ranebennur Black Buck, Dharwad; Sharavathi Valley, Shivamogga; Shettihalli, Shivamogga; Someshwara, Udupi; Talakaveri, Kodagu); 3 conservation reserves (Banakapur, Haveri and Jayamangali, Tumkur) and one Community Reserve (Kokkare Bellur, Mandya).

This work presents an up dated checklist of all seed plants occurring in Karnataka.

The entire list with distribution data and local/common names was extracted from secondary published and

internet sources. The basic list of species was prepared referring *Flora of Karnataka analysis* (Sharma et al. 1984) and *The Flora of Karnataka* (Saldanha, volume I, 1984 and volume II, 1996). The district-wise distribution data and local/common names were further refined from all accessible published Floras, research papers, unpublished Doctoral thesis, reports and miscellaneous publications from 1984 onwards till date.

Sharma et al. (1984) recorded 4120 species, subspecies and varieties under genera and families. Whereas, Saldanha (1984) recorded 4110 species (did not indicate the number of infraspecific ranks). All naturalised, invasive and cultivated species in the state are also listed here. There are many cultivated exotics particularly the species of horticultural importance, which need to be correctly identified before they are listed.

The nomenclatural details like family names, generic names, currently accepted species names, basionym and synonym/s are updated following *The Plant List* and taxonomic revisions and monographs. The names considered as undecided in *The Plant List* are treated as accepted here till a final taxonomic decision is taken them.

The data presented in this work is as given below:

Nomenclature: In this work the currently accepted names are given in bold letters. While, the basionyms (the epithet bringing names based on which the accepted is derived) and synonyms are given in italics. All basionyms and synonyms are arranged chronologically. The synonyms included here are generally limited to those relevant to those dealt in India Floras and publications. For exhaustive synonymy one can refer to up dated versions of *The Plant List* and other published taxonomic revisions and monographs. All names are given with literature citations following standard abbreviations as given in *The Plant List* (based on *Authors of Plant Names* by Brummitt & Powell; journal following Botanico-Periodica-Huntianum (BPH) and its supplements, and *Taxonomic Literature-TL2* and its supplements which are modified for improvement).

Distribution: All the geographical names of districts in Karnataka, states, broader regions in India, countries and continents are given in alphabetical sequence. All the distribution data is extracted from published sources. The distribution given under each species/subspecies/variety starts with recorded data in Karnataka giving known district wise distribution in brackets, followed by other states of India if they are limited in number or the regions if widely distributed. This is followed by countries (if they are few in number)/continents/subcontinents, if much wider as pantropical, paleotropical, cosmopolitan, etc. After giving the district wise distribution in bracket for Karnataka, if a species/subspecies/variety is exclusively confined to the state (considered as exclusively/localised endemic) the word endemic is given following a semicolon. In case, it is limited in distribution to a few states of India, the names of the states are given followed by semicolon and endemic. If it is endemic to India, but widely distributed in several states, broad regions like S India, Peninsular India, N India, W India, E India, NE India, Himalayas, Gangetic plain, Andaman & Nicobar Islands or Lakshadweep Islands with a semicolon followed by endemic is given. In some cases where name/s of district of Karnataka mentioned in brackets as the authors have simply given Karnataka in the distribution of the taxon.

Local/Common names: The names in English, Kannada, Konkani, Kunabi, Lambani, Sanskrit and Tulu are given here and extracted from published works and internet resources. All these names and their languages are given in alphabetical sequence taken from published resources and databases. After each name/names the language is given in brackets

Arrangement of species: All names of families, genera under each family, species (including infraspecific names under each species, i.e., subspecies and varieties) under each genus are given alphabetically. The families and genera are arranged following Angiosperm Phylogeny Group (APG III).

Recorded families, genera and species in Karnataka: This work presents recorded families, genera, species, subspecies and varieties within the present political boundaries of Karnataka. Species and varieties that are exclusively endemic to Karnataka and also those that extend to adjoining state/s and other states

within India are also given Based on literature.

We have included all families, genera, species, subspecies and varieties so far recorded from Karnataka state in this check-list. There will, of course, be errors and omissions, therefore, we would be most grateful if we are informed of these corrections as the data can be continually updated (including new species, new distributional records, and additional vernacular/common names from the state) in the database that the Karnataka Biodiversity Board is developing on biological resources.

During the compilation of this check-list, the authors received help of Drs. G. V. S. Murthy (MH). P. Lakshminarasimhan (CAL), S. K. Srivastava (BSD), C. Murugan (PBL) with literature and of J. V. Sudhakar (MH in checking genus *Ficus*) and G. Gnanasekaran (MH in checking *Andrographis*).

Summary of Karnataka Angiosperms Checklist

Total No. of Families: **191**; Total No. of Genera: **1535**; Total No. of Species including subspecies and varieties: **5009**

Table 1: No. of Families, Genera, Species, Subspecies and Varieties

| Sl. No. | Family | Genera | Species | Subsp. | Variety | Total |
|---------|-------------------|--------|---------|--------|---------|-------|
| 1 | Acanthaceae | 34 | 209 | 3 | 11 | 223 |
| 2 | Achariaceae | 1 | 3 | | | 3 |
| 3 | Acoraceae | 1 | 1 | | | 1 |
| 4 | Adoxaceae | 1 | 2 | | | 2 |
| 5 | Aizoaceae | 3 | 7 | | | 7 |
| 6 | Alismataceae | 4 | 5 | | | 5 |
| 7 | Amaranthaceae | 22 | 48 | 1 | 3 | 52 |
| 8 | Amaryllidaceae | 7 | 17 | | | 17 |
| 9 | Anacardiaceae | 11 | 24 | | | 24 |
| 10 | Ancistrocladaceae | 1 | 1 | | | 1 |
| 11 | Annonaceae | 17 | 38 | | | 38 |
| 12 | Apiaceae | 17 | 25 | | 1 | 26 |
| 13 | Apocynaceae | 54 | 139 | 1 | 6 | 146 |
| 14 | Aponogetonaceae | 1 | 4 | | | 4 |
| 15 | Aquifoliaceae | 1 | 3 | | | 3 |
| 16 | Araceae | 26 | 56 | 1 | 2 | 59 |
| 17 | Araliaceae | 3 | 11 | | | 11 |
| 18 | Arecaceae | 41 | 85 | | | 85 |
| 19 | Aristolochiaceae | 2 | 9 | | | 9 |
| 20 | Asparagaceae | 15 | 59 | 2 | 1 | 62 |
| 21 | Asteraceae | 123 | 299 | 3 | 9 | 311 |
| 22 | Balanophoraceae | 1 | 4 | | | 4 |
| 23 | Balsaminaceae | 1 | 49 | | 12 | 61 |
| 24 | Basellaceae | 1 | 1 | | | 1 |
| 25 | Begoniaceae | 1 | 9 | | | 9 |
| 26 | Bignoniaceae | 27 | 47 | | | 47 |
| 27 | Bixaceae | 2 | 2 | | | 2 |
| 28 | Boraginaceae | 9 | 41 | | 1 | 42 |
| 29 | Brassicaceae | 8 | 15 | 1 | | 16 |
| 30 | Bromeliaceae | 1 | 1 | | | 1 |
| 31 | Buddlejaceae | 1 | 3 | | | 3 |
| 32 | Burmanniaceae | 1 | 7 | | | 7 |
| 33 | Burseraceae | 5 | 7 | | | 7 |
| 34 | Buxaceae | 1 | 1 | | | 1 |
| 35 | Cactaceae | 4 | 7 | | | 7 |
| 36 | Calceolariaceae | 1 | 1 | | | 1 |
| 37 | Calophyllaceae | 4 | 8 | | | 8 |
| 38 | Campanulaceae | 5 | 13 | | | 13 |
| 39 | Cannabaceae | 4 | 6 | | | 6 |
| 40 | Cannaceae | 1 | 3 | | | 3 |

| | | | | | | |
|----|------------------|----|-----|----|---|-----|
| 41 | Capparaceae | 5 | 17 | | | 17 |
| 42 | Caprifoliaceae | 1 | 2 | | | 2 |
| 43 | Caricaceae | 1 | 1 | | | 1 |
| 44 | Caryophyllaceae | 7 | 8 | 1 | | 9 |
| 45 | Casurinaceae | 1 | 1 | | | 1 |
| 46 | Celastraceae | 14 | 34 | 1 | | 35 |
| 47 | Ceratophyllaceae | 1 | 1 | | | 1 |
| 48 | Cleomaceae | 2 | 9 | | | 9 |
| 49 | Clusiaceae | 2 | 9 | | 1 | 10 |
| 50 | Colchicaceae | 3 | 8 | | | 8 |
| 51 | Combretaceae | 6 | 20 | | | 20 |
| 52 | Commelinaceae | 9 | 58 | 1 | 1 | 60 |
| 53 | Connaraceae | 2 | 5 | | | 5 |
| 54 | Convolvulaceae | 25 | 102 | 1 | 2 | 105 |
| 55 | Cornaceae | 2 | 2 | 2 | | 4 |
| 56 | Costaceae | 4 | 3 | | | 3 |
| 57 | Crassulaceae | 4 | 10 | | | 10 |
| 58 | Cucurbitaceae | 22 | 53 | 1 | 1 | 55 |
| 59 | Cyperaceae | 21 | 181 | 11 | 4 | 196 |
| 60 | Daphniphyllaceae | 1 | 1 | | | 1 |
| 61 | Dichapetalaceae | 1 | 1 | | | 1 |
| 62 | Dilleniaceae | 1 | 4 | | | 4 |
| 63 | Dioscoreaceae | 2 | 16 | | | 16 |
| 64 | Dipterocarpaceae | 5 | 14 | | 1 | 15 |
| 65 | Droseraceae | 1 | 3 | | | 3 |
| 66 | Ebenaceae | 1 | 27 | | 1 | 28 |
| 67 | Elaeagnaceae | 1 | 4 | | | 4 |
| 68 | Elaeocarpaceae | 1 | 6 | | 1 | 7 |
| 69 | Elatinaceae | 2 | 3 | | | 3 |
| 70 | Ericaceae | 2 | 2 | | | 2 |
| 71 | Eriocaulaceae | 1 | 46 | | | 46 |
| 72 | Erythoxylaceae | 1 | 3 | | | 3 |
| 73 | Euphorbiaceae | 31 | 138 | 4 | 4 | 146 |
| 74 | Flagellariaceae | 1 | 1 | | | 1 |
| 75 | Gentianaceae | 8 | 26 | | | 26 |
| 76 | Gesneriaceae | 9 | 14 | | | 14 |
| 77 | Gisekiaceae | 1 | 1 | | | 1 |
| 78 | Goodeniaceae | 1 | 2 | | | 2 |
| 79 | Haloragaceae | 2 | 2 | | | 2 |
| 80 | Heliconiaceae | 1 | 4 | | | 4 |
| 81 | Hernandiaceae | 1 | 1 | | | 1 |
| 82 | Hydatellaceae | 1 | 1 | | | 1 |
| 83 | Hydrocharitaceae | 7 | 12 | | | 12 |
| 84 | Hydroleaceae | 1 | 1 | | | 1 |
| 85 | Hypericaceae | 1 | 4 | | | 4 |
| 86 | Hypoxidaceae | 3 | 4 | | | 4 |
| 87 | Icacinaceae | 4 | 4 | | | 4 |

| | | | | | | |
|-----|------------------|-----|-----|---|----|-----|
| 87 | Icacinaceae | 4 | 4 | | | 4 |
| 88 | Iridaceae | 1 | 1 | | | 1 |
| 89 | Juncaceae | 2 | 3 | | | 3 |
| 90 | Lamiaceae | 34 | 146 | | 3 | 149 |
| 91 | Lauraceae | 12 | 52 | | | 52 |
| 92 | Lecythidaceae | 3 | 4 | | | 4 |
| 93 | Leguminosae | 131 | 452 | 5 | 23 | 480 |
| 97 | Lemnaceae | 3 | 7 | | | 7 |
| 98 | Lentibulariaceae | 1 | 23 | | | 23 |
| 99 | Liliaceae | 2 | 3 | | 1 | 4 |
| 100 | Linaceae | 3 | 6 | | | 6 |
| 101 | Linderniaceae | 4 | 28 | | | 28 |
| 102 | Loganiaceae | 5 | 12 | | | 12 |
| 103 | Lophiocarpaceae | 1 | 1 | | | 1 |
| 104 | Loranthaceae | 8 | 22 | | | 22 |
| 105 | Lythraceae | 10 | 34 | | 1 | 35 |
| 106 | Magnoliaceae | 1 | 3 | | | 3 |
| 107 | Malpighiaceae | 4 | 8 | | | 8 |
| 108 | Malvaceae | 43 | 145 | 4 | 1 | 150 |
| 109 | Marantaceae | 4 | 9 | | | 9 |
| 110 | Martyniaceae | 1 | 1 | | | 1 |
| 111 | Melastomataceae | 5 | 29 | | 2 | 31 |
| 112 | Meliaceae | 16 | 24 | | | 24 |
| 113 | Menispermaceae | 10 | 14 | | | 14 |
| 114 | Menyanthaceae | 1 | 4 | | | 4 |
| 115 | Molluginaceae | 2 | 6 | | | 6 |
| 116 | Moraceae | 9 | 39 | 2 | 1 | 42 |
| 117 | Moringaceae | 1 | 1 | | | 1 |
| 118 | Muntingiaceae | 1 | 1 | | | 1 |
| 119 | Musaceae | 2 | 4 | | | 4 |
| 120 | Myristicaceae | 3 | 6 | 1 | | 7 |
| 121 | Myrtaceae | 7 | 43 | | | 43 |
| 122 | Nelumbonaceae | 1 | 1 | | | 1 |
| 123 | Nitrariaceae | 1 | 1 | | | 1 |
| 124 | Nyctaginaceae | 4 | 12 | | | 12 |
| 125 | Nymphaeaceae | 1 | 2 | | | 2 |
| 126 | Ochnaceae | 2 | 4 | | 1 | 5 |
| 127 | Olacaceae | 4 | 5 | | | 5 |
| 128 | Oleaceae | 6 | 26 | 2 | | 28 |
| 129 | Onagraceae | 2 | 8 | 1 | | 9 |
| 130 | Opiliaceae | 2 | 2 | | | 2 |
| 131 | Orchidaceae | 62 | 196 | | 1 | 197 |
| 132 | Orobanchaceae | 10 | 21 | | 1 | 22 |
| 133 | Oxalidaceae | 3 | 14 | | 1 | 15 |
| 134 | Pandanaceae | 2 | 11 | | | 11 |
| 135 | Papaveraceae | 1 | 1 | | | 1 |
| 136 | Passifloraceae | 4 | 10 | | | 10 |

| | | | | | | |
|-----|------------------|-----|-----|---|---|-----|
| 137 | Paulowniaceae | 1 | 1 | | | 1 |
| 138 | Pedaliaceae | 2 | 4 | | | 4 |
| 139 | Pentaphylacaceae | 1 | 1 | | | 1 |
| 140 | Phrymaceae | 2 | 2 | | | 2 |
| 141 | Phyllanthaceae | 13 | 60 | | 3 | 63 |
| 142 | Phytolaccaceae | 2 | 2 | | | 2 |
| 143 | Piperaceae | 2 | 22 | | | 22 |
| 144 | Pittosporaceae | 1 | 1 | | | 1 |
| 145 | Plantaginaceae | 17 | 36 | | | 36 |
| 146 | Plumbaginaceae | 1 | 3 | | | 3 |
| 147 | Poaceae | 132 | 452 | 1 | 7 | 460 |
| 148 | Podostemaceae | 6 | 11 | | | 11 |
| 149 | Polygalaceae | 3 | 13 | | | 13 |
| 150 | Polygonaceae | 7 | 21 | | 4 | 25 |
| 151 | Pontederiaceae | 1 | 3 | | | 3 |
| 152 | Portulacaceae | 1 | 7 | | | 7 |
| 153 | Potamogetonaceae | 2 | 3 | | | 3 |
| 154 | Primulaceae | 6 | 17 | | | 17 |
| 155 | Proteaceae | 2 | 2 | | | 2 |
| 156 | Putranjivaceae | 2 | 5 | | | 5 |
| 157 | Ranunculaceae | 5 | 16 | | | 16 |
| 158 | Rhamnaceae | 8 | 18 | | 2 | 20 |
| 159 | Rhizophoraceae | 6 | 7 | | | 7 |
| 160 | Rosaceae | 6 | 18 | | 1 | 19 |
| 161 | Rubiaceae | 52 | 170 | 1 | 8 | 179 |
| 162 | Rutaceae | 18 | 32 | | | 32 |
| 163 | Sabiaceae | 1 | 3 | 1 | | 4 |
| 164 | Salicaceae | 6 | 12 | | | 12 |
| 165 | Salvadoraceae | 2 | 2 | | | 2 |
| 166 | Santalaceae | 4 | 16 | | 1 | 17 |
| 167 | Sapindaceae | 11 | 16 | | 1 | 17 |
| 168 | Sapotaceae | 7 | 15 | | 1 | 16 |
| 169 | Saxifragaceae | 1 | 1 | | | 1 |
| 170 | Scrophulariaceae | 1 | 1 | | | 1 |
| 171 | Simaroubaceae | 2 | 3 | | | 3 |
| 172 | Smilacaceae | 1 | 4 | | | 4 |
| 173 | Solanaceae | 14 | 60 | | | 60 |
| 174 | Sphaenocleaceae | 1 | 1 | | | 1 |
| 175 | Staphyleaceae | 1 | 2 | | | 2 |
| 176 | Stemonuraceae | 1 | 2 | | | 2 |
| 177 | Strelitziaceae | 2 | 2 | | | 2 |
| 178 | Symplocaceae | 1 | 9 | | 1 | 10 |
| 179 | Talinaceae | 1 | 1 | | | 1 |
| 180 | Tamaricaceae | 1 | 1 | | | 1 |
| 181 | Tetramelaceae | 1 | 1 | | | 1 |
| 182 | Theaceae | 2 | 2 | | | 2 |
| 183 | Thymeliaceae | 1 | 1 | | | 1 |

| | | | | | | |
|--------------------|------------------|-------------|-------------|-----------|------------|-------------|
| 184 | Typhaceae | 1 | 2 | | | 2 |
| 185 | Ulmaceae | 1 | 1 | | | 1 |
| 186 | Urticaceae | 13 | 31 | 1 | 2 | 34 |
| 187 | Vahliaceae | 1 | 2 | | | 2 |
| 188 | Verbenaceae | 9 | 19 | | | 19 |
| 189 | Violaceae | 4 | 7 | | | 7 |
| 190 | Vitaceae | 7 | 32 | | | 32 |
| 191 | Xanthorrhoeaceae | 2 | 6 | | | 6 |
| 192 | Xyridaceae | 1 | 4 | | | 4 |
| 193 | Zingiberaceae | 10 | 51 | | | 51 |
| 194 | Zygophyllaceae | 3 | 6 | | | 6 |
| Grand Total | | 1535 | 4827 | 53 | 129 | 5009 |

Table 2: Twelve largest families with more than 100 species

| Sl. No. | Family | Genera | Species | Subsp. | Variety | Total |
|---------|----------------|--------|---------|--------|---------|-------|
| 1 | Leguminosae | 131 | 452 | 5 | 23 | 480 |
| 2 | Poaceae | 132 | 452 | 1 | 7 | 460 |
| 3 | Asteraceae | 123 | 299 | 3 | 9 | 311 |
| 4 | Acanthaceae | 34 | 209 | 3 | 11 | 223 |
| 5 | Orchidaceae | 62 | 196 | | 1 | 197 |
| 6 | Cyperaceae | 21 | 181 | 11 | 4 | 196 |
| 7 | Rubiaceae | 52 | 170 | 1 | 8 | 179 |
| 8 | Malvaceae | 43 | 145 | 4 | 1 | 150 |
| 9 | Lamiaceae | 34 | 146 | | 3 | 149 |
| 10 | Apocynaceae | 54 | 139 | 1 | 6 | 146 |
| 11 | Euphorbiaceae | 31 | 138 | 4 | 4 | 146 |
| 12 | Convolvulaceae | 25 | 102 | 1 | 2 | 105 |

Table 3: Number and distribution of endemic species exclusive to Karnataka and those with extended distribution in peninsular and rest of India

| Sl. No. | Family | Karnataka | Peninsular India | India |
|---------|-------------------|-----------|------------------|-------|
| 1 | Acanthaceae | 7 | 91 | 137 |
| 2 | Achariaceae | | 3 | 3 |
| 3 | Acoraceae | | | |
| 4 | Adoxaceae | | | |
| 5 | Aizoaceae | | | |
| 6 | Alismataceae | | 1 | 1 |
| 7 | Amaranthaceae | | 4 | 4 |
| 8 | Amaryllidaceae | | 1 | 2 |
| 9 | Anacardiaceae | | 10 | 11 |
| 10 | Ancistrocladaceae | | 1 | 1 |
| 11 | Annonaceae | 2 | 17 | 19 |
| 12 | Apiaceae | 1 | 9 | 12 |
| 13 | Apocynaceae | 4 | 39 | 51 |
| 14 | Aponogetonaceae | | 1 | 1 |
| 15 | Aquifoliaceae | 1 | 2 | 3 |
| 16 | Araceae | 1 | 15 | 18 |
| 17 | Araliaceae | | 2 | 3 |
| 18 | Arecaceae | 1 | 15 | 16 |
| 19 | Aristolochiaceae | | 2 | 2 |
| 20 | Asparagaceae | 4 | 15 | 19 |
| 21 | Asteraceae | | 20 | 39 |
| 22 | Balanophoraceae | | | |
| 23 | Balsaminaceae | 23 | 51 | 77 |
| 24 | Basellaceae | | | |
| 25 | Begoniaceae | | 5 | 6 |
| 26 | Bignoniaceae | | 3 | 5 |
| 27 | Bixaceae | | | |
| 28 | Boraginaceae | 1 | 10 | 13 |
| 29 | Brassicaceae | | | |
| 30 | Bromeliaceae | | | |
| 31 | Buddlejaceae | | | |
| 32 | Burmanniaceae | | 1 | 1 |
| 33 | Burseraceae | | 1 | 1 |
| 34 | Buxaceae | | | |
| 35 | Cactaceae | | | |
| 36 | Calceolariaceae | | | |
| 37 | Calophyllaceae | | 3 | 3 |
| 38 | Campanulaceae | | 1 | 1 |
| 39 | Cannabaceae | | | |
| 40 | Cannaceae | | | |
| 41 | Capparaceae | | 3 | 3 |
| 42 | Caprifoliaceae | | 1 | 1 |
| 43 | Caricaceae | | | |
| 44 | Caryophyllaceae | | | 1 |

| | | | | |
|----|------------------|---|----|----|
| 45 | Casurinaceae | | | |
| 46 | Celastraceae | 1 | 16 | 18 |
| 47 | Ceratophyllaceae | | | |
| 48 | Cleomaceae | | | |
| 49 | Clusiaceae | | 3 | 3 |
| 50 | Colchicaceae | 1 | 4 | 5 |
| 51 | Combretaceae | 2 | 3 | 6 |
| 52 | Commelinaceae | 1 | 12 | 17 |
| 53 | Connaraceae | | 3 | 3 |
| 54 | Convolvulaceae | | 15 | 17 |
| 55 | Cornaceae | | 1 | 1 |
| 56 | Costaceae | | | |
| 57 | Crassulaceae | | 2 | 2 |
| 58 | Cucurbitaceae | | 2 | 4 |
| 59 | Cyperaceae | 7 | 17 | 26 |
| 60 | Daphniphyllaceae | | | |
| 61 | Dichapetalaceae | | | |
| 62 | Dilleniaceae | | 1 | 1 |
| 63 | Dioscoreaceae | | | 1 |
| 64 | Dipterocarpaceae | 2 | 13 | 15 |
| 65 | Droseraceae | | | |
| 66 | Ebenaceae | | 6 | 9 |
| 67 | Elaeagnaceae | | | |
| 68 | Elaeocarpaceae | | 3 | 3 |
| 69 | Elatinaceae | | | |
| 70 | Ericaceae | | 1 | 1 |
| 71 | Eriocaulaceae | 8 | 24 | 38 |
| 72 | Erythroxylaceae | | | |
| 73 | Euphorbiaceae | 2 | 26 | 39 |
| 74 | Flagellariaceae | | | |
| 75 | Gentianaceae | 1 | 11 | 15 |
| 76 | Gesneriaceae | 1 | 5 | 6 |
| 77 | Gisekiaceae | | | |
| 78 | Goodeniaceae | | | |
| 79 | Haloragaceae | | | |
| 80 | Heliconiaceae | | | |
| 81 | Hernandiaceae | | | |
| 82 | Hydatellaceae | | 1 | 1 |
| 83 | Hydrocharitaceae | | 1 | 1 |
| 84 | Hydroleaceae | | | |
| 85 | Hypericaceae | | | |
| 86 | Hypoxidaceae | | | |
| 87 | Icacinaceae | | 2 | 2 |
| 88 | Iridaceae | | | |
| 89 | Juncaceae | | | |
| 90 | Lamiaceae | 3 | 37 | 45 |
| 91 | Lauraceae | 2 | 30 | 34 |

| | | | | |
|-----|------------------|---|----|----|
| 92 | Lecythidaceae | | | |
| 93 | Leguminosae | 1 | 41 | 85 |
| 94 | Lemnaceae | | | |
| 95 | Lentibulariaceae | | 9 | 10 |
| 96 | Liliaceae | | 1 | 1 |
| 97 | Linaceae | | 2 | 2 |
| 98 | Linderniaceae | | 6 | 6 |
| 99 | Loganiaceae | | 2 | 2 |
| 100 | Lophiocarpaceae | | | |
| 101 | Loranthaceae | | 12 | 12 |
| 102 | Lythraceae | 1 | 5 | 8 |
| 103 | Magnoliaceae | | | |
| 104 | Malpighiaceae | | 2 | 4 |
| 105 | Malvaceae | | 11 | 18 |
| 106 | Marantaceae | | | |
| 107 | Martyniaceae | | | |
| 108 | Melastomataceae | 1 | 16 | 17 |
| 109 | Meliaceae | | 2 | 2 |
| 110 | Menispermaceae | 1 | 1 | 2 |
| 111 | Menyanthaceae | | | |
| 112 | Molluginaceae | | | |
| 113 | Moraceae | | 3 | 4 |
| 114 | Moringaceae | | | |
| 115 | Muntingiaceae | | | |
| 116 | Musaceae | | 1 | 1 |
| 117 | Myristicaceae | | 6 | 6 |
| 118 | Myrtaceae | 4 | 16 | 21 |
| 119 | Nelumbonaceae | | | |
| 120 | Nitrariaceae | | | |
| 121 | Nyctaginaceae | | 1 | 1 |
| 122 | Nymphaeaceae | | | |
| 123 | Ochnaceae | | | 1 |
| 124 | Olacaceae | | | |
| 125 | Oleaceae | | 6 | 7 |
| 126 | Onagraceae | | | |
| 127 | Opiliaceae | | | |
| 128 | Orchidaceae | 1 | 68 | 78 |
| 129 | Orobanchaceae | 2 | 5 | 8 |
| 130 | Oxalidaceae | | | |
| 131 | Pandanaceae | 2 | 4 | 6 |
| 132 | Papaveraceae | | | |
| 133 | Passifloraceae | | 1 | 1 |
| 134 | Paulowniaceae | | | |
| 135 | Pedaliaceae | | 1 | 1 |
| 136 | Pentaphylacaceae | | | |
| 137 | Phrymaceae | | | |
| 138 | Phyllanthaceae | 1 | 11 | 18 |

| | | | | |
|-----|------------------|---|-----|-----|
| 139 | Phytolaccaceae | | | |
| 140 | Piperaceae | 2 | 6 | 8 |
| 141 | Pittosporaceae | | 1 | 1 |
| 142 | Plantaginaceae | | | 1 |
| 143 | Plumbaginaceae | | | |
| 144 | Poaceae | 5 | 108 | 121 |
| 145 | Podostemaceae | | 5 | 5 |
| 146 | Polygalaceae | | | 1 |
| 147 | Polygonaceae | 2 | 2 | 4 |
| 148 | Pontederiaceae | | | |
| 149 | Portulacaceae | 2 | 2 | 4 |
| 150 | Potamogetonaceae | | | |
| 151 | Primulaceae | 1 | 3 | 4 |
| 152 | Proteaceae | | 1 | 1 |
| 153 | Putranjivaceae | | 2 | 2 |
| 154 | Ranunculaceae | | 5 | 7 |
| 155 | Rhamnaceae | | 3 | 6 |
| 156 | Rhizophoraceae | | 1 | 1 |
| 157 | Rosaceae | 1 | 1 | 4 |
| 158 | Rubiaceae | 5 | 68 | 77 |
| 159 | Rutaceae | | 3 | 3 |
| 160 | Sabiaceae | | | |
| 161 | Salicaceae | | 3 | 4 |
| 162 | Salvadoraceae | | | |
| 163 | Santalaceae | 3 | 6 | 9 |
| 164 | Sapindaceae | | 2 | 2 |
| 165 | Sapotaceae | 1 | 6 | 7 |
| 166 | Saxifragaceae | | | |
| 167 | Scrophulariaceae | | | |
| 168 | Simaroubaceae | | | |
| 169 | Smilacaceae | | | |
| 170 | Solanaceae | | 1 | 1 |
| 171 | Sphaenocleaceae | | | |
| 172 | Staphyleaceae | | | |
| 173 | Stemonuraceae | | | |
| 174 | Strelitziaceae | | | |
| 175 | Symplocaceae | | 4 | 4 |
| 176 | Talinaceae | | | |
| 177 | Tamaricaceae | | | |
| 178 | Tetramelaceae | | | |
| 179 | Theaceae | | 1 | 1 |
| 180 | Thymeliaceae | | | |
| 181 | Typhaceae | | | |
| 182 | Ulmaceae | | | |
| 183 | Urticaceae | | 1 | 1 |
| 184 | Vahliaceae | | | |
| 185 | Verbenaceae | | | |

| | | | | |
|--------------------|------------------|------------|-------------|-------------|
| 186 | Violaceae | | | |
| 187 | Vitaceae | | 4 | 5 |
| 188 | Xanthorrhoeaceae | | | |
| 189 | Xyridaceae | | | |
| 190 | Zingiberaceae | 4 | 14 | 19 |
| 191 | Zygophyllaceae | | 1 | 1 |
| Grand Total | | 116 | 1053 | 1393 |

Acanthaceae

Acanthus L.

Acanthus ilicifolius L., Sp. Pl. 639. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Dilivaria ilicifolia (L.) Juss., Gen. Pl. 103. 1789

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), throughout coastal areas including Islands.

Local/Common/Vernacular names: *Hole chulli, Mullu chulli* (Kannada)

Andrographis Wall. ex Nees

Andrographis subg. *Indoneesiella* (Sreem.) L.H. Cramer

Indoneesiella Sreem.

Neesiella Sreem.

Andrographis affinis Nees in Wall., Pl. Asiat. Rar. 3: 116. 1832

Andrographis neesiana var. *affinis* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 504. 1884

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Andrographis alata (Vahl) Nees in A. DC., Prodr. 11: 516. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Justicia alata Vahl, Enum. Pl. 1: 139. 1804

Distribution: Karnataka (Chamarajanagar, Hassan, Kodagu, Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Andrographis atropurpurea (Dennst.) Alston, Taxon 26: 539. 1977

Justicia atropurpurea Dennst., Schlüssel Hortus Malab. 12, 22, 35. 1818 & in Fortsetz. Allg. Teutsch. Gart.-Mag. 3: 31, 40, 83. 1818

Andrographis gracilis Nees in A.DC., Prodr. 11: 516. 1847

Andrographis wightiana Arn. ex Nees in A.DC., Prodr. 11: 517. 1847 nom. illeg.; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Distribution: Karnataka (Chikkamagaluru, Mysuru, Shivamogga), Kerala, Tamil Nadu; endemic.

Andrographis ceylanica Nees in Hook., Compan. Bot. Mag. 2: 313. 1836

Andrographis macrobotrys Nees in A.DC., Prodr. 11: 516. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Andrographis macrobotrys var. *parvifolia* C.B.Clarke in Hook.f., Fl. Brit. India 4: 503. 1884

Andrographis macrobotrys var. *ceylanica* (Nees) L.H.Cramer in Dassan., Rev. Handb. Fl. Ceylon 12: 100. 1998

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Mysuru, Shivamogga, Udupi), Kerala, Tamil Nadu; Sri Lanka.

Andrographis echiooides (L.) Nees in Wall., Pl. Asiat. Rar. 3: 117. 1832

Justicia echiooides L., Sp. Pl. 16. 1753

Neesiella echiooides (L.) Sreem., Phytologia 15: 271. 1967

Indoneesiella echiooides (L.) Sreem., Phytologia 16: 466. 1968; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Erianthera echiooides (L.) M. R. Almeida, Fl. Maharashtra 4A: 40. 2003

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Dharwad, Gulbaraga, Raichur, Mysuru, Vijayapura, Yadgir), throughout tropical India.

Local/Common/Vernacular names: *Ativisha, Godi barasan, Peethumba* (Kannada)

Andrographis elongata (Vahl) T.Anderson, J. Linn. Soc., Bot. 9: 502. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Justicia elongata Vahl, Enum.Pl. 1: 130. 1804

Cryptophragmum cordifolium Nees, Pl. Asiat. Rar. 3: 100. 1832

Cryptophragmum elongatum (Vahl) Nees in Wall., Pl. Asiat. Rar. 3: 95. 1832

Distribution: Karnataka (Bengaluru), Andhra Pradesh, Kerala, Odisha, Tamil Nadu; endemic.

Andrographis explicata (C.B.Clarke) Gamble, Fl. Madras 1049. 1924; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Andrographis viscosa var. *explicata* C.B.Clarke in Hook.f., Fl. Brit. India 4: 504. 1884

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Andrographis glandulosa Nees in Wall., Pl. Asiat. Rar. 3: 115. 1832 & in A. DC., Prodr. 11: 518. 1847; C.B.Clarke in Hook.f., Fl. Brit. India 4: 505. 1884.

Justicia glandulosa Wall., Numer. List 2485. 1830

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Andrographis lawsonii Gamble, Bull. Misc. Inform. 1923: 375. 1923 & Fl. Madras 1050. 1924; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Distribution: Karnataka (Dakshina Kannada, Udupi), Kerala, Tamil Nadu; endemic.

Andrographis lineata Wall. ex Nees in Wall., Pl. Asiat. Rar. 3: 116. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

var. **lineata**

Distribution: Karnataka (Ballari, Chamarajanagar, Chitradurga, Hassan, Mysuru, Tumakuru), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Nerale nelabevu* (Kannada)

var. **lawii** C.B.Clarke in Hook.f., Fl. Brit. India 4: 505. 1884; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Distribution: Karnataka (Ballari, Chikkaballapur, Chikkamagaluru, Chitradurga); endemic.

Andrographis longipedunculata (Sreem.) L.H. Cramer, Kew Bull. 51: 555. 1996

Neesiella longipedunculata Sreem., Phytologia 15: 271. 1967

Indoneesiella longipedunculata (Sreem.) Sreem., Phytologia 16: 466. 1968; K.M. Matthew, Fl. Tamilnadu Carnatic 3: 1182. 1983.

Erianthera longipedunculata (Sreem.) M.R. Almeida, Fl. Maharashtra 4A: 41. 2003

Distribution: Karnataka (Ballari, Chitradurga, Dharwad), throughout India.

Andrographis neesiana Wight, Icon. Pl. Ind. Orient. 4: t. 1561. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Kaasinasarada gida* (Kannada)

Andrographis ovata (T.Anderson ex Bedd.) Benth. & Hook.f., Gen. Pl. 2(2): 1100. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Gymnostachyum ovatum T.Anderson ex Bedd., Icon. Pl. Ind. Or. 60, 61.t. 250. 1874

Distribution: Karnataka (Shivamogga), S to NE India.

Local/Common/Vernacular names: *Kalamegha, Nelabevina gida, Nelabevu, Oorakiriyathu, Urikiriyathu* (Kannada)

Andrographis paniculata (Burm.f.) Wall. ex Nees in Wall., Pl. Asiat. Rar. 3: 116. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Justicia paniculata Burm.f., Fl. indica 9. 1768

Andrographis paniculata var. *glandulosa* Trimen, J. Bot. 27: 165.1889

Distribution: Karnataka (throughout), throughout India; endemic.

Local/Common/Vernacular names: *Kaala megha, Nela bevina gida, Nela bevu, Oora kiriyathu* (Kannada)

Andrographis producta (C.B.Clarke) Gamble, Fl. Madras 1049. 1924; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Plate 1A

Andrographis neesiana var. *producta* C.B.Clarke in Hook.f., Fl. Brit. India 4: 504. 1884

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Kerala, Tamil Nadu; endemic.

Andrographis serpyllifolia (Vahl) Wight, Icon. Pl. Ind. Orient. t. 517. 1841; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Justicia serpyllifolia Rottler ex Vahl, Enum. Pl. 1: 169. 1804.

Distribution: Karnataka (Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kolar Mysuru, Shivamogga, Tumakuru), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Kasina sara, Sardali gedde, Sardali gida* (Kannada)

Andrographis viscosa Nees in Wall., Pl. Asiat. Rar. 3: 116. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Mysuru), Kerala, Tamil Nadu; endemic.

Asystasia Blume

Asystasia chelonoides Nees in Wall., Pl. Asiat. Rar. 3: 89. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Asystasia coromandeliana var. *variabilis* Nees in A.DC., Prodr. 11: 165.1847

Asystasia chelonoides var. *quadrangularis* B.Heyne ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 494.1884

Asystasia chelonoides var. *variabilis* (Nees) M.R.Almeida, Fl. Maharashtra 4A: 7.2003

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Mysuru, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Asystasia crispata Benth., Linnaea 24: 647. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Asystasia coromandeliana Wight, Icon. Pl. Ind. Orient. 4: t. 1506. 1850., auct. non Nees, 1832.

Distribution: Karnataka (Chikkamagaluru, Kodagu, Shivamogga), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Asystasia dalzelliana Santapau, Kew Bull. 1948: 276. 1948; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Asystasia violacea Dalzell ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 494. 1884.

Distribution: Karnataka (Belagavi, Ballari, Chikkmgaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru,Udupi,), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Asystasia gangetica (L.) T.Anderson in Thwaites, Enum. Pl. zeyl. 235. 1860; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Plate 1B

Justicia gangetica L., Cent. Pl. II : 3. 1756

Asystasia calycina Nees in Wall., Pl. Asiat. Rar. 3: 90. 1832

Asystasia coromandeliana Nees, Pl. Asiat. Rar. 3: 89. 1832., auct. non Wight, 1850.

Asystasia plumbaginea Nees in Wall., Pl. Asiat. Rar. 3: 89. 1832

Ruellia zeylanica K.D.Koenig ex Roxb., Fl. Ind. ed.2, 3: 42. 1832

Asystasia quarterna Nees in A.DC., Prodr. 11: 166. 1847

Ruellia coromandelina Nees in A.DC., Prodr. 11: 166. 1847

Ruellia filiformis Klein ex Nees in A.DC., Prodr. 11: 165. 1847

Ruellia nemoralis Vahl ex Nees in A.DC., Prodr. 11: 165. 1847

Ruellia secundiflora Bojer ex Nees in A.DC., Prodr. 11: 166. 1847

Ruellia variabilis Macrae ex Nees in A.DC., Prodr. 11: 165. 1847

Asystasia violacea Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 139. 1850.

Asystasia gangetica var. mendeliana S.R.Kundu & Mahua Pal, J. Econ. Taxon. Bot. 21: 486. 1997

subsp. *gangetica*

Distribution: Karnataka (throughout, sometimes cultivated), throughout India including Andaman & Nicobar Islands; Africa, S, SE & W Asia.

Local/Common/Vernacular names: *Lavanga valli*, *Madde soppu*, *Maithala Kaddi* (Kannada); *Maithaal*, *Maithala kaddi* (Tulu)

subsp. **micrantha** (Nees) Ensermu in J.H.Seyani & A.C.Chikuni, Proc. XIII Plenary Meet. AETFAT, Zomba, Malawi 1: 343. 1994; K.G.Bhat, Indian J. for. 39 (1): 67. 2016.

Asystasia coromandeliana var. *micrantha* Nees, Prodr. 11: 165. 1847.

Distribution: Karnataka (Udupi), West Bengal; Africa, Malaysia, Indonesia, Pacific Islands.

Asystasia mysorensis (Roth) T.Anderson, J. Linn. Soc., Bot. 9: 524. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Ruellia mysorensis Roth, Nov. Pl. Sp. 303. 1821

Asystasia lawiana Dalzell in Hooker's J. Bot. Kew Gard. Misc. 4: 344. 1852

Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Kodagu, Mysuru, Tumakuru, Uttara Kannada), Maharashtra; endemic.

Asystasia nemorum Nees, Pl. Asiat. Rar. 3: 90. 1832; Punekar & Lakshmin., Fl. Anshi Natl. Park 346. 2011

Asystasia blumei Nees A.DC., Prodr. 11: 167. 1847

Distribution: Karnataka (Uttara Kannada), Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; endemic.

Avicennia L.

Avicennia marina (Forssk.) Vierh., Denkschr. Kaiserl. Akad. Wiss., Wien. Math.-Naturwiss. Kl. 71: 435. 1907

Sceura marina Forssk., Fl. Aegypt.-Arab. 37. 1775

Avicennia alba Blume, Bijdr. 821. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Avicennia officinalis var. *alba* (Blume) C.B.Clarke in Hook.f., Fl. Brit. India 4: 604. 1885

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Goa, Maharashtra, Tamil Nadu; coastal S, SE & SW Asia

Avicennia officinalis L., Sp. Pl. 110. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Avicennia oepata Buch.-Ham., Trans. Linn. Soc. London 17: 221. 1837

Avicennia obovata Griff., Not. Pl. Asiat. 4: 189. 1854

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), throughout coastal areas of India; Australia, Asia.

Local/Common/Vernacular names: *Ipathi*, *Kandaale*, *Uppaathi* (Kannada)

Barleria L.

Barleria acuminata Wight ex Nees in Wall., Pl. Asiat. Rar. 3: 93. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.

Barleria tomentosa var. *acuminata* (Wight ex Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 485. 1884

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Davanagere, Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Barleria buxifolia L., Sp. Pl. 636. 1753; Wight, Icon. Pl. Ind. Orient. 3: t. 870. 1845; B.D.Sharma et al., Fl. Karnataka Anal. 201. 1984.

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Tumakuru), Andhra Pradesh, Kerala, Maharashtra, Punjab, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Chulli*, *Gubbachhi mullu*, *Gubbi mullu*, *Gundukothi mullu*, *Kari chulli*, *Kempu gorte*, *Krishna chora*, *Kuruvaka* (Kannada)

Barleria courtallica Nees in A.DC., Prodr. 11: 226. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.

Barleria nutans Nees in A. DC., Prodr. 11: 227. 1847

Barleria inaequalis Benth., Flora 32: 557. 1849

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Chethasahacharam* (Sanskrit)

Barleria cristata L., Sp. Pl. 636. 1753; Wight, Icon. Pl. Ind. Orient. t. 453. 1841; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.

Plate 1D

Barleria ciliata Roxb., Hort. Bengal.: 45. 1814

Barleria dichotoma Roxb., Fl. Ind. 3: 93. 1832

Barleria laciniata Nees in Wall., Pl. Asiat. Rar. 3: 91. 1832

Barleria napalensis Nees in Wall., Pl. Asiat. Rar. 3: 91. 1832

Barleria nuda Nees in Wall., Pl. Asiat. Rar. 3: 92. 1832

Barleria indica L. ex T.Anderson, J. Proc. Linn. Soc., Bot. 7: 115. 1864

Barleria cristata var. *dichotoma* (Roxb.) Prain, Bengal Pl. 812. 1903

Barleria cristata var. *albida* Haines, Bot. Bihar Orissa 4: 683. 1922

Distribution: Karnataka (throughout, often cultivated), throughout India; Bangladesh, China, Myanmar, Nepal, Pakistan, Philippines, Sri Lanka, Vietnam.

Local/Common/Vernacular names: Blue bell Barleria, Philippine Violet-white (English); Gorante, Gorte, Jhante, Jhinte, Kinkiraatha, Kuravaka, Mullu Gorate, Patikada hoo, Spatika, Spatikada Hoovu, Zanti (Kannada); Artagala, Bana, Dasi, Dasikurantaka, Jhinti, Kakubha, Mahasaha, Nilamlaainduka, Ragaprasava, Saereyakah-nilapuspa (Sanskrit)

Barleria cuspidata B. Heyne ex Nees in Wall., Pl. Asiat. Rar. 3: 93. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.

Distribution: Karnataka (Chamarajanagar, Davanagere, Mysuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Kerala, Maharashtra, Rajasthan, Tamil Nadu; endemic.

Barleria gibsonii Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 339. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.

Plate 13 A&B

Barleria mackenii Hook.f., Bot. Mag. 96: t. 5866. 1870

Barleria montana Wight & Nees ex C.B.Clarke, Fl. Brit. India 3: 487. 1882.

Barleria gibsonioides Blatt., J. Bombay Nat. Hist. Soc. 32: 733. 1928

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Kodagu), Andhra Pradesh, Gujarat, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Bachali*, *Gura*, *Kove mullu*, *Mullu ankole*, *Neeruppi gida* (Kannada)

Barleria grandiflora Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 339. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.
Barleria grandis auct. non Hochst. ex Nees, 1847; M.R.Almeida, Fl. Maharashtra 4A:13. 2003
Barleria grandifolia R. Br. ex T.Anderson, J. Proc.Linn. Soc., Bot. 7: 30. 1864
 Distribution: Karnataka (Belagavi, Shivamogga, Uttara kannada), Goa, Maharashtra; endemic.

Plate 1C&13C

Barleria involucrata Nees in Wall., Pl. Asiat. Rar. 3: 92. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.
Barleria pentandra Arn. ex Nees in A.DC., Prodr. 11: 232. 1847

var. involucrata

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

var. elata (Dalzell) C.B.Clarke in Hook.f., Fl. Brit. India 4: 486. 1884
Barleria elata Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 227. 1851
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Davanagere, Karwar, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Plate 1E&13D

Barleria lawii T.Anderson, J. Linn. Soc., Bot. 9: 492. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.
Barleria beddomei T.Anderson ex Bedd., Icon. Pl. Ind. Or.: 62, t. 258. 1874
Barleria sepalosa C.B.Clarke in Hook. f., Fl. Brit. India 4: 487. 1884
 Distribution: Karnataka (Uttara Kannada), Gujarat, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Barleria longiflora L.f., Suppl. Pl. 239. 1782; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.
 Distribution: Karnataka (Mysuru, Raichur), Andhra Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.
 Local/Common/Vernacular name: *Adyanda* (Sanskrit)

Barleria lupulina Lindl. in Edwards' Bot. Reg. 18: t. 1483. 1833; C.B.Clarke in Hook.f., Fl. Brit. India 4: 482. 1884.
 Distribution: Karnataka (cultivated, also as an escape); Bihar, Himachal Pradesh, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, Uttar Pradesh, West Bengal; Native to Mauritius.

Barleria montana Nees, Pl. Asiat. Rar. 3: 92. 1832.
Barleria morrisiana E.Barnes & C.E.C. Fisch., Bull. Misc. Inform. 1939: 659. 1939; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.
 Distribution: Karnataka (Mandy, Mysuru), Andhra Pradesh, Bihar, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.

Barleria mysorensis B.Heyne ex Roth, Nov. Pl. Sp. 313. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.
Barleria bispinosa Nees in Wall., Pl. Asiat. Rar. 3: 94. 1832
Barleria spina-ceylanica Nees in A.DC., Prodr. 11: 242. 1847
 Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Hassan, Kodagu, Mysuru, Ramanagara, Tumakuru), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Barleria nitida Nees in Wall., Pl. Asiat. Rar. 3: 91. 1832; Shendage & S. R. Yadav, Rheedea 21:106. 2011.
 Distribution: Karnataka (Mysuru), Tamil Nadu; Sri Lanka.

Barleria noctiflora L.f., Suppl. Pl. 290. 1782; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.
Barleriacanthus noctiflora (L.f.) Oerst., Vidensk. Med -del. Dansk Naturhist. Foren. Kjobenhavn: 136. 1854
 Distribution: Karnataka (cultivated in Mysuru), Andhra Pradesh, Tamil Nadu; Sri Lanka.

Barleria prattensis Santapau, Kew Bull. 1948: 487. 1949

Distribution: Karnataka (Belagavi), Peninsular India, Rajasthan.

Distribution: Karnataka (Bengaluru, Kolar, Mysuru, Shivamogga, Tumakuru), Andhra Pradesh, Dadra & Nagar Haveli, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu; endemic.

Barleria prionitis L., Sp. Pl. 636. 1753; Wight, Icon. Pl. Ind. Orient. t. 452. 1841; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Plate 1F

Barleria hystrix L., Mant. Pl. 1: 89. 1767

subsp. *prionitis*

Barleria prionitis subsp. *prionitis* var. *prionitis*; Bhandari, Fl. Ind. Desert: 299. 1978

Distribution: Karnataka (throughout), throughout tropical India; Malaysia, Pakistan, Philippines, Sri Lanka, tropical Africa, Yemen.

Local/Common/Vernacular names: *Porcupine flower* (English); *Gorate, Gotle hoo, Haladi gorte, Kuranta, Kurantike, Mullu corate, Mullu gorante, Mullu Gorate, Mullu jaji, Mullu kanakambra, Mullu madarangi, Semmuli* (Kannada)

subsp. *pubiflora* (Benth. ex Hohen.) Brummitt & J.R.I. Wood, Kew Bull. 38: 436. 1983

Barleria pubiflora Benth. ex Hohen. in Flora 32: 358. 1849

Barleria prionitis subsp. *Prionitis* var. *diacantha* Blatt. & Hallb., J. Bombay Nat. Hist. Soc. 26: 811. 1919 non Barleria *diacantha* Nees 1847

Distribution: Karnataka (Coastal and subcoastal areas), Diu & Daman, Goa, Gujarat, Madhya Pradesh, Maharashtra, Punjab, Rajasthan; Pakistan.

Barleria repens Nees in A.DC., Prodr. 11: 230. 1847; Shendage & S. R. Yadav, Rhedea 21: 95. 2011.

Distribution: Karnataka (Bengaluru), Maharashtra; native to tropical & subtropical SE Africa; cultivated elsewhere..

Barleria stocksii T.Anderson, J. Linn. Soc., Bot. 9: 493. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Distribution: Karnataka (Chikkamagaluru), Andhra Pradesh; endemic.

Barleria strigosa Willd., Sp. Pl. 3: 379. 1800 emend. S.K. Malhotra & Moorthy in Indian Forester 107: 449. 1981; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Barleria polytricha Wall., Pl. Asiat. Rar. 1: 72, t. 82. 1830

Barleria caerulea Roxb., Fl. Ind. 3: 39. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 202. 1984.

Barleria macrophylla B. Heyne ex C.B. Clarke in Hooker, Fl. Brit. India 4: 489. 1884

Barleria strigosa var. *subglabra* Kuntze, Revis. Gen. Pl. 1(2): 483. 1891

Distribution: Karnataka (Bengaluru, Ballari, Dakshina Kannada, Udupi, Uttara Kannada), throughout India; S & SE Asia.

Local/Common/Vernacular name: *Artagala* (Sanskrit)

Barleria terminalis Nees in A.DC., Prodr. 11: 225. 1847 emend. S.K. Malhotra & Moorthy, Indian Forester 107: 449. 1981; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Barleria polystachya Nees in A.DC., Prodr. 11: 226. 1847

Barleria strigosa var. *polystachya* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 490. 1884

Barleria strigosa var. *terminalis* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 490. 1884

Distribution: Karnataka (Belagavi), peninsular India; endemic.

Local/Common/Vernacular name: *Blue Barleria* (English)

Barleria tomentosa Roth, Nov. Pl. sp. 314. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Barleria paniculata Wall. ex Nees in Wall., Pl. Asiat. Rar. 3: 92. 1832

Barleria tomentosa var. *glandulosa* C.E.C.Fisch., Rec. Bot. Surv. India 9: 135. 1921

Distribution: Karnataka (Chamarajanagar, Chikmagaluru, Chitradurga, Koppala, Raichur, Vijayapura), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Blepharis Juss.

Blepharis integrifolia (L.f.) E. Mey. ex Schinz in Vierteljahrsschr. Naturf. Ges. Zürich 60: 416. 1915

Acanthus integrifolius L. f., Suppl. Pl. 294. 1782

Acanthus repens Vahl, Symb. Bot. 2: 76. 1791

Blepharis molluginifolia Pers., Syn. Pl. 2: 180. 1807

Acanthus molluginifolius (Pers.) Poir. in Lam., Encycl. Suppl. 1: 90. 1810

Blepharis repens (Vahl) Roth, Nov. Pl. Sp. 321. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Blepharis lawsonii G.S.Giri & R. N. Banerjee, Blumea 28: 363. 1983 p.p. quoad T.A. Rao 8115.

Distribution: Karnataka (throughout), throughout tropical India; Sri Lanka.

Local/Common/Vernacular names: *Haridachchga*, *Haridrata* (Kannada)

Blepharis maderaspatensis (L.) B. Heyne ex Roth, Nov. Pl. Sp. 320. 1821; B.D.Sharma et al., Fl.

Karnataka Anal. 203. 1984.

Acanthus maderaspatensis L., Sp. Pl. 639. 1753

Blepharis boerhaviifolia Pers., Syn. Pl. 2: 180. 1806 ('boerhaaviaefolia')

Blepharis boerhaviifolia var. *maderaspatensis* (L.) Nees in Wall., Pl. Asiat. Rar. 3: 97. 1832

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburgi, Kodagu, Kolar, Mysuru, Raichur, Vijayapura, Yadgir), throughout India; S Asia, tropical Africa.

Local/Common/Vernacular names: *Kodali soppu*, *Nanju naaraayani* (Kannada); *Utthangsa* (Sanskrit)

Calacanthus T.Anderson ex Benth.

Calacanthus grandiflorus (Dalzell) Radlk. in Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss.

München 13: 279. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

Plate 13 E

Lepidagathis grandiflora Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 138. 1850

Calacanthus dalzelliana T.Anderson ex Benth. in Benth. & Hook.f., Gen. Pl. 2: 1088.1876

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), peninsular India; endemic.

Crossandra Salisb.

Crossandra infundibuliformis (L.) Nees in Wall., Pl. Asiat. Rar. 3: 98. 1832; B.D.Sharma et al., Fl.

Karnataka Anal. 204. 1984.

Justicia infundibuliformis L., Syst. nat. ed. 10. 2: 850. 1759

Crossandra undulifolia Salisb., Parad. lond. t. 12. 1806; Sims in Curtis, Bot. Mag. 47: t. 2186. 1820; C.B.Clarke in Hook.f., Fl. Brit. India 4: 492. 1884.

Crassandra axillaris Nees in Wall., Pl. Asiat. Rar. 3: 98. 1832

Crassandra infundibuliformis var. *axillaris* (Nees) Trimen, Handb. Fl. Ceylon 3: 323. 1895

Distribution: Karnataka (cultivated & wild throughout), cultivated throughout India; native to S India.

Local/Common/Vernacular names: *Firecracker flower* (English); *Abballige*, *Abbollige*, *Kanakambra* (Kannada); *Amlaana* (Sanskrit); *Abbolige* (Tulu)

Cynarospermum Vollesen

Cynarospermum aspernum (Nees) Vollesen, Kew Bull. 54: 172, f. 1. 1999

Blepharis asperrima Nees in A.DC., Prodr11: 267. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 203. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Dicliptera Juss., nom. cons.

Dicliptera babui Karthik. & Moorthy, Fl. Pl. India 10.2009

Justicia baphica Spreng., Neue Entd. 3: 82. 1822

Peristrophe tinctoria auct. non Nees, 1832; C.B.Clarke in Hook.f., Fl. Brit. India 4: 556. 1885 p.p.

Peristrophe baphica (Spreng.) Bremek., Nova Guinea 8: 149. 1957 & Dansk. Bot. Ark. 20: 80. 1961 non

Dicliptera baphica Nees, 1847

Distribution: Karnataka (cultivated), cultivated elsewhere.

Dicliptera bupleuroides Nees in Wall., Pl. Asiat. Rar. 3: 111. 1832; Gamble, Fl. Madras 2: 1074. 1921.

Dicliptera roxburghiana var. *bupleuroides* (Nees) C.B.Clarke in Hook.f., Fl. Brit. India 4: 554. 1885

Distribution: Karnataka, Andhra Pradesh, Odisha; Afghanistan, Bhutan, Bangladesh, China, Indo-China, Nepal, Pakistan.

Dicliptera cuneata Nees in Wall., Pl. Asiat. Rar. 3: 111. 1832 & in DC., Prodr. 11: 481. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Davanagere, Hassan, Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Dicliptera foetida (Forssk.) Blatt., Rec. Bot. Surv. India 8: 361. 1921

Justicia foetida Forssk., Fl. Aegypt.-Arab. 5. 1775

Dicliptera bivalvis Nees in A.DC., Prodr. 11: 475. 1847

Dicliptera zeylanica Nees in A.DC., Prodr. 11: 474. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

var. **foetida**

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Dharwad, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil nadu; Sri Lanka.

var. **kanarana** M. R. Almeida, Fl. Maharashtra 4A: 29. 2003

Distribution: Karnataka (Uttara kannada), Maharashtra; endemic.

Dicliptera nasikensis Lakshmin. & B.D.Sharma, J. Econ. Taxon. Bot. 7: 481.1986; Nilpure, Fl Belagavi Dist. 2008.

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Dicliptera paniculata (Forssk.) I.Darbysh., Kew Bull. 62: 122. 2007

Dianthera paniculata Forssk., Fl. aegypt.-arab. 7. 1775

Dianthera malabarica L.f., Suppl. Pl. 85. 1782 nom. illegit.

Peristrophe bicalyculata var. *rivinoides* C.B.Clarke in Hook.f., Fl. Brit. India 4: 555. 1885

Peristrophe bicalyculata var. *subaequibracteata* Bennet, J. Bombay Nat. Hist. Soc. 66: 229.1969

Peristrophe paniculata (Forssk.) Brummitt, Kew Bull. 38: 451. 1983

Dicliptera paniculata var. *rivinoides* (C.B.Clarke) Karthik. & Moorthy, Fl. Pl. India 12.2009

Dicliptera paniculata var. *subaequibracteata* (Bennet) Karthik. & Moorthy, Fl. Pl. India 12.2009

Distribution: Karnataka (throughout), throughout subtropical & tropical India; SE Asia, tropical Africa.

Local/Common/Vernacular names: *Chibeera gida*, *Gibi gida* (Kannada)

Dicliptera verticillata (Forssk.) C. Chr., Dansk. Bot. Ark. 4 (3): 11. 1922

Dianthera verticillata Forssk., Fl. Aegypt.-Arab. 9. 1775

Dicliptera micranthes Nees in Wall., Pl. Asiat. Rar. 3: 112. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

Distribution: Karnataka (Belagavi, Chitradurga), Peninsular to E, N India; Afghanistan, Etiopia.

Dipteracanthus Nees

Dipteracanthus dejectus Nees in Wall., Pl. Asiat. Rar. 3: 82.1832; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

Distribution: Karnataka (Belagavi), Tamil Nadu; endemic.

Dipteracanthus prostratus (Poir.) Nees, Pl. Asiat. Rar. 3: 81.1832; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

Distribution: Karnataka (Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kalaburgi, Mysuru, Uttara Kannada), Daman, Goa, Kerala, Maharashtra, Nagar Haveli, Tamil Nadu; S & SE Asia.

Local/Common/Vernacular names: *Bheemana soppu, Silandi nayakana gida* (Kannada)

Dyschoriste Nees *Calophanes* D.Don

Dyschoriste dalzellii (T.Anderson ex Bedd.) Kuntze, Revis. Gen. Pl. 2: 486. 1891; T. Cooke, Fl. Bombay 2: 362. 1905.

Calophanes dalzellii T.Anderson ex Bedd., Icon. Pl. Ind. Or. t. 248. 1874

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Dyschoriste erecta (Burm.f.) Kuntze, Revis. Gen. Pl. 2: 485.1891

Ruellia erecta Burm.f., Fl. Indica 135.1768

Justicia Mysorensis Klein ex Schult., Mant. 1: 146.1822

Ruellia depressa Wall., Numer. List 2379.1830

Dyschoriste cernua Nees in Wall., Pl. Asiat. Rar. 3: 81.1832

Dyschoriste depressa (L.) Nees in Wall., Pl. Asiat. Rar. 3: 81.1832

Ruellia setigera Wight ex Nees in Wall., Pl. Asiat. Rar. 3: 81.1832

Calophanes depressus (L.) T.Anderson, Enum. Pl. Zeyl. 225.1860

Dyschoriste mysorensis (Klein ex Schult.) M.R.Almeida, Fl. Maharashtra 4A: 32.2003

Distribution: Karnataka (Bengaluru, Mysuru), Peninsular India; Sri Lanka.

Dyschoriste nagchana (Nees) Bennet, Indian Forester 109: 220. 1983

Dipteracanthus nagchana Nees in Wall., Pl. Asiat. Rar. 3: 82. 1832

Calophanes nagchana Nees in A.DC., Prodr.11: 109. 1847

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Uttara Kannada), Peninsular India; Sri Lanka.

Dyschoriste vagans (Wight) Kuntze, Revis. Gen. Pl. 2: 486. 1891

Calophanes vagans Wight, Icon. Pl. Ind. Orient. t. 1526. 1850

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Chitradurga), Peninsular to E India; endemic.

Ecbolium Kurz

Ecbolium ligustrinum (Vahl) Vollesen, Kew Bull. 44: 651. 1989

Justicia ecbolium L., Sp. Pl. 15. 1753

Justicia laetevirens Vahl, Enum. Pl. 1: 118. 1804

Justicia ligustrina Vahl, Enum. Pl. 1: 118. 1804

Justicia dentata Klein ex Link, Jahrb. Gewächsk. 1 (3): 48. 1820

Justicia ecbolium var. *mollis* Nees in A.DC., Prodr. 11: 427. 1847

Justicia emarginata Nees in A.DC., Prodr. 11: 427. 1847

Justicia livida Nees in A.DC., Prodr. 11: 427. 1847

Ecbolium linneanum Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 40: 75. 1875

Ecbolium linneanum var. *dentatum* (Klein ex Link) C.B. Clarke in Hook.f., Fl. Brit. India 4: 545. 1885

Ecbolium linneanum var. *laetevirens* (Vahl) C.B. Clarke in Hook.f., Fl. Brit. India 4: 545. 1885

Ecbolium viride var. *dentatum* (Klein ex Link) Santapau, Rec. Bot. Surv. India 16(1): 208. 1967

Ecbolium viride var. *laetevirens* (Vahl) Santapau, Rec. Bot. Surv. India 16 (1): 208. 1967; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Ecbolium viride subsp. *ligustrina* (Vahl) M.R.Almeida, Fl. Maharashtra 4A: 34. 2003

Ecbolium viride subsp. *ligustrina* var. *dentata* (Klein ex Link) M. R. Almeida, Fl. Maharashtra 4A: 35. 2003

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), E & NE to Peninsular India, Upper Gangetic Plains; S & SE Asia.

Ecbolium viride (Forssk.) Alston, in Trimen, Handb. Fl. Ceylon 6: 229. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984

Justicia viridis Forssk., Fl. aegypt.-arab. 5. 1775

Justicia rotundifolia Nees in Wall., Pl. Asiat. Rar. 3: 108. 1832

Ecbolium linneanum var. *rotundifolia* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 545. 1885

Ecbolium viride var. *rotundifolia* (Nees) Raizada, Indian Forester 84: 482. 1958

Distribution: Karnataka (Chitradurga, Davanagere, Shivamogga), E to Peninsular India.

Local/Common/Vernacular names: *Kappu karani*, *Kappubobbili*, *Nela abbolige*, *Pachambara*, *Sarpagenthige* (Kannada)

Elytraria Michx., nom. cons.

Elytraria acaulis (L.f.) Lindau in Engl. & Prantl, Nat. Pflanzenfam. Nachtr. 1: 304. 1897; Gamble, Fl. Madras 2: 1009. 1921.

Justicia acaulis L. f., Suppl. Pl. 84. 1782

Elytraria crenata Vahl, Enum. Pl. 1: 106. 1804.

Tubiflora acaulis (L.f.) Kuntze, Revis. Gen. Pl. 2: 500. 1891

var. *acaulis*

Distribution: Karnataka (), E to N India & E Himalaya, peninsular India; endemic.

var. *lyrata* (Vahl) Bremek., Reinwardtia 3: 251. 1955

Elytraria crenata sensu C.B.Clarke in Hook.f., Fl. Brit. India 4: 394. 1884.; non Vahl, 1804.

Elytraria lyrata Vahl, Enum. Pl. 1: 106. 1804

Elytraria crenata var. *lyrata* (Vahl) Nees in A.DC., Prodr. 11: 163. 1847

Distribution: Karnataka (), peninsular India; endemic.

Eranthemum L.

Daedalacanthus T.Anderson

Eranthemum capense L., Sp. Pl. 9. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Justicia fastigiata Lam., Tabl. Encycl. 1: 41.1791

Justicia montana Roxb., Pl. Coromandel2: 41, t. 176. 1805

Eranthemum fastigiatum (Lam.) R.Br. ex Roem. & Schult., Syst. Veg. 1: 174. 1817

Eranthemum montanum (Roxb.) Roxb., Fl. Ind. ed.1, 1: 110. 1820

Eranthemum wightianum Wall. ex Nees in Wall. Pl. Asiat. Rar. 3: 107. 1832

Daedalacanthus montanus (Roxb.) T.Anderson in Thwaites, Enum. Pl. zeyl. 229. 1860

Daedalacanthus montanus var. *wightianus* (Wall. ex Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 421. 1884

Daedalacanthus capensis (L.) Druce, Rep. Bot. Exch. Club Soc. Brit. Isles 3: 417. 1914

Eranthemum montanum var. *wightianum* (Wall ex Nees) Gamble, Fl. Madras 1025. 1924

Daedalacanthus fastigiatus (Lam.) Alston, Ann. Roy. Bot. Gard. (Peradeniya) 11: 299.1930

Eranthemum capense var. *wightianum* (Wall. ex Nees) Karthik. & Moorthy, Fl. Pl. India 1. 2009

var. *capense*

Distribution: Karnataka (Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Alpine eranthemum* (English)

var. **concanensis** (T.Anderson ex C.B.Clarke) Santapau, Bot. Mem. Univ. Bombay 2: 34. 1952; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Daedalacanthus montanus var. *concanensis* T.Anderson ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 421. 1884

Eranthemum montanum var. *concanensis* (T.Anderson ex C.B.Clarke) Gamble, Fl. Madras 1025. 1924

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Eranthemum pulchellum Andrews, Bot. Repos. 2, t. 88. 1800

Justicia nervosa Vahl, Enum. Pl. 1: 164. 1804

Eranthemum nervosum (Vahl) R. Br., Prodr. 477. 1810

Daedalacanthus nervosus (Vahl) T.Anderson, J. Linn. Soc., Bot. 9: 487. 1867

Distribution: Karnataka (throughout, also cultivated), throughout India; Pakistan; cultivated elsewhere.

Eranthemum purpurascens Nees in Wall., Pl. Asiat. Rar. 3:106. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Daedalacanthus purpurascens (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 488. 1867

Eranthemum pulchellum var. *purpurascens* (Nees) Kuntze, Revis. Gen. Pl. 1891

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), throughout India; endemic.

Eranthemum roseum (Vahl) R. Br., Prodr. 477. 1810; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Justicia rosea Vahl, Enum. Pl. 1: 165. 1804

Daedalacanthus roseus (Vahl) T.Anderson, J. Linn. Soc., Bot. 9: 487. 1867

Eranthemum pulchellum var. *roseum* (Vahl) Kuntze, Revis. Gen. Pl. 2: 489. 1891

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dakshina Kannada, Udupi, Uttara Kannada), Andhra Pradesh, E India, Goa, Kerala, Maharashtra, Nagar Haveli, Odisha, Tamil Nadu; endemic.

Graptophyllum Nees

Graptophyllum pictum (L.) Griff., Not. Pl. Asiat. 4: 139. 1854

Justicia picta L., Sp. Pl. ed. 2. 21. 1762

Graptophyllum hortense Nees in Wall., Pl. Asiat. Rar. 3: 102. 1832

Distribution: Karnataka (throughout in gardens), cultivated throughout India including Andaman & Nicobar Islands; SE Asia.

Gymnostachyum Nees

Gymnostachyum canescens (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 505. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Cryptophragmum canescens Nees in Wall., Pl. Asiat. Rar. 3: 100. 1832

var. **canescens**

Distribution: Karnataka (Dakshina Kannada, Uttara kannada), Kerala, Tamil Nadu; endemic.

var. **longifolium** C.B.Clarke in Hook.f., Fl. Brit. India 4: 509. 1884

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada); endemic.

Gymnostachyum febrifugum Benth., Flora 32: 558. 1849; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), Kerala; endemic.

Local/Common/Vernacular names: *Aneyadi*, *Bili agarada beru*, *Jwarahara soppu*, *Nela muchchala*, *Nelamuchhe* (Kannada); *Agaradaberu*, *Boldu* (Tulu)

Gymnostachyum glabrum (Dalzell) T.Anderson, J. Linn. Soc., Bot. 9: 506. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Cryptophragmum glabrum Dalzell in Hooker's J. Bot. Kew. Gard. Misc. 2: 338. 1850

Distribution: Karnataka (Belagavi, Uttara Kannada), Goa, Maharashtra; endemic.

Gymnostachyum latifolium (Dalzell) T.Anderson, J. Linn. Soc., Bot. 9: 505. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Cryptophragmum latifolium Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 137. 1850

Phlogacanthus latifolius Wight, Icon. Pl. Ind. Orient. t. 1537. 1850

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Gymnostachyum polyanthum Wight, Icon. Pl. Ind. Orient. t.1494.1849; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), Kerala; endemic.

Gymnostachyum pubescens (Lam.) M.R.Almeida, Fl. Maharashtra 4A: 45. 2003

Justicia pubescens Lam., Tabl. Encycl. 1: 40.1791

Cryptophragmum serrulatum Nees A.DC., Prodr. 11: 95.1847

Gymnostachyum serrulatum T.Anderson, J. Linn. Soc., Bot. 9: 505.1867

Gymnostachyum longifolium T.Anderson ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 509. 1884

Distribution: Karnataka (Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Haplanthodes Kuntze

Bremekampia Sreem.

Haplanthus sensu C.B.Clarke

Haplanthodes tentaculatus (L.) R.B.Majumdar, Bull. Bot. Soc. Bengal 25: 76.1971

Ruellia tentaculata L., Cent. Pl. 2: 22. 1756 & Amoen. Acad. 4: 320. 1759

Haplanthus tentaculatus (L.) Nees in A.DC., Prodr. 11: 513. 1847; C.B.Clarke in Hook.f., Fl. Brit. India 4: 507. 1884.

Haplanthus tentaculatus var. *neesianus* Santapau, Bot. Mem. Univ. Bombay 2: 52. 1952

Bremekampia tentaculata (L.) Sreem., Bull. Bot. Surv. India 6: 324. 1965

var. **tentaculatus**

Distribution: Karnataka (Belagavi), Peninsular India to Upper Gangetic Plains; endemic.

var. **neilgherrensis** (Wight) J.R.I.Wood, Novon 23(3): 386.2014

Haplanthus neilgherrensis Wight, Icon. Pl. Ind. Orient. t. 1556. 1850

Haplanthus plumosus T. Anderson, J. Linn. Soc., Bot. 9: 504.1867

Haplanthus tentaculatus var. *neilgherrensis* (Wight) C.B. Clarke in Hook.f., Fl. Brit. India 4: 507. 1884 ('nilgherrensis').

Haplanthus tentaculatus var. *plumosus* (T.Anderson) C. B. Clarke in Hook.f., Fl. Brit. India 4:507. 1884

Bremekampia neilgherryensis (Wight) Sreem., Bull. Bot. Surv. India 6: 323. 1965

Haplanthodes neilgherryensis (Wight) R.B. Majumdar, Bull. Bot. Soc. Bengal 25: 76. 1971; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Haplanthodes plumosa (T.Anderson) Panigrahi & G.C.Das, Bull. Bot. Surv. India 23: 200.1983

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Haplanthodes verticillatus (Roxb.) R.B. Majumdar, Bull. Bot. Soc. Bengal. 25: 76. 1971; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Plate 13F

Justicia verticillata Roxb., Fl. Ind. ed.1, 1: 135. 1820

Haplanthus verticillatus (Roxb.) Nees in A.DC., Prodr. 1: 513. 1847

Bremekampia verticillata (Roxb.) Sreem., Bull. Bot. Surv. India 6: 323. 1965

Distribution: Karnataka (Belagavi, Ballari, Dharwad, Uttara Kannada), Peninsular to E India & Upper Gangetic Plains; endemic.

Local/Common/Vernacular name: *Spiny bottle brush* (English)

Hygrophila R. Br.

Cardanthera Buch.-Ham. ex Nees

Nomaphila Blume

Santapaua N.P. Balakr. & Subr.

Hygrophila balsamica (L.f.) Raf., Fl. Tellur. 4: 66. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Ruellia balsamica L.f., Suppl. Pl. 289. 1782

Adenosma balsamica (L.f.) Spreng., Syst. veg. ed. 16. 2: 829. 1825 ('balsamea'); Wight, Icon. Pl. Ind. Orient. t. 446. 1841.

Adenosma thymus Nees in Wall., Pl. Asiat. Rar. 3: 79. 1832

Adenosma verticillata Nees in Wall., Pl. Asiat. Rar. 3: 79. 1832

Cardanthera balsamica (L.f.) Benth. in Benth. & Hook.f., Gen. Pl. 2: 1074. 1876

Cardanthera balsamica var. *thymus* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 404. 1884

Cardanthera verticillata (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 404. 1884

Synnema barbigera Kuntze, Revis. Gen. Pl. 2: 500. 1891

Synnema verticillatum (Nees) Kuntze, Revis. Gen. Pl. 2: 500. 1891

Synnema balsamicum (L.f.) Alston in Trimen, Handb. Fl. Ceylon 6 (Suppl.): 224. 1931

Dipteracanthus cernuus (Roxb.) Santapau, Indian Forester 79: 322. 1953

Distribution: Karnataka (Dakshina Kannada), Peninsular to E India; Sri Lanka.

Hygrophila erecta (Burm.f.) Hochr., Candollea 5: 230. 1934

Ruellia quadrivalvis Buch.-Ham., Trans. Linn. Soc. London 14: 291. 1824

Hygrophila bengalensis S.K. Mandal, A. Bhattach. & Nayek, J. Bombay Nat. Hist. Soc. 94: 546. 1997

Distribution: Karnataka (Uttara Kannad), Andaman & Nicobar Islands, E to Peninsular India.

Hygrophila incana Nees in A.DC., Prodr. 11: 91. 1847; C.B.Clarke in Hook.f., Fl. Brit. India 4: 408. 1884.

Distribution: Karnataka (), S India; endemic.

Hygrophila pinnatifida (Dalzell) Sreem., Bull. Bot. Surv. India 10: 222. 1968; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Nomaphila pinnatifida Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 38. 1851

Adenosma pinnatifida (Dalzell) T.Anderson, J. Linn. Soc., Bot. 9: 455. 1867

Cardanthera pinnatifida (Dalzell) Benth. in Benth. & Hook.f., Gen. Pl. 2: 1074. 1876

Synnema pinnatifidum (Dalzell) Kuntze, Revis. Gen. Pl. 2: 500. 1891

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Peninsular India to subhimalayan tracts; endemic.

Hygrophila polysperma (Roxb.) T.Anderson, J. Linn. Soc., Bot. 9: 456. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Hemiadelphus polyspermus (Roxb.) Nees in Wall., Pl. Asiat. Rar. 3: 80. 1832

Justicia polysperma Roxb., Fl. Ind. ed.2, 1: 119. 1832

Hemiadelphus polyspermus var. *joshianus* Prak.Rao & S.N. Biswas, Indian Forester 94: 657. 1968

Distribution: Karnataka (Belagavi, Dharwad, Shivamogga, Uttara Kannada), throughout India; Bhutan, China, Myanmar, Sri Lanka, Vietnam.

Hygrophila ringens (L.) R. Br. ex Steud., Nomencl. Bot. ed. 1. 1: 418. 1821

Ruellia ringens L., Sp. Pl. 635. 1753

Ruellia salicifolia Vahl, Symb. Bot. 3: 84. 1794

Hygrophila angustifolia R. Br., Prodr. 479. 1810

Ruellia obovata Roxb. ex Hornem., Suppl. Hort. Bot. hafn. 144. 1819

Hygrophila quadrivalvis (Buch.-Ham.) Nees in Wall., Pl. Asiat. Rar. 3: 80. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Hygrophila salicifolia (Vahl) Nees in Wall., Pl. Asiat. Rar. 3: 81. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Hygrophila obovata (Roxb.) Nees ex Wight, Icon. Pl. Ind. Orient. t. 1489. 1849

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Kodagu, Shivamogga, Udupi, Uttara Kannada), throughout India including Andaman & Nicobar Islands; E, S & SE Asia.

Local/Common/Vernacular name: *Indian Perry* (English); *Kokilaksha* (Sanskrit)

Hygrophila schullii (Buch.-Ham.) M.R.Almeida & S. M. Almeida, J. Bombay Nat. Hist. Soc. 83 (Suppl.): 221. 1987

Barleria longifolia L., Cent. Pl. II: 22. 1756 & Amoen. Acad. 4: 320. 1759 non *Hygrophila longifolia* Nees, 1847

Bahelschulli Buch.-Ham., Trans. Linn. Soc. London 14: 289. 1824

Barleria auriculata Schumach., Beskr. Guin. Pl. 285. 1827

Asteracantha longifolia (L.) Nees in Wall., Pl. Asiat. Rar. 3: 90. 1832

Hygrophila spinosa T.Anderson in Thwaites, Enum. Pl. zeyl. 225. 1860

Hygrophila longifolia (L.) Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 39: 78. 1870 nom. illegit.

Hygrophila auriculata (Schumach.) Heine, Kew Bull. 16: 172. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Kalaburgi, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada, Yadgir), throughout India; S & SE Asia, tropical Africa.

Local/Common/Vernacular names: Gokantaka, Golimadi, Goramidi, Gorividi, Gulimidi, Kokilaksha, Kolachulli, Kolagolike, Kolamulli, Kolase mullina gida, Kolavalike, Kolavanke, Koli kukkana gida, Kolikannina gida, Koravanke, Neeru gobli gida, Neeruppi gida (Kannada); Kolista (Konkani); Kokilaksha (Sanskrit); Mullankaana (Tulu)

Hygrophila serpyllum (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 456. 1867

Physichilus serpyllum Nees in Hook., Compan. Bot. Mag. 2: 310. 1836

Distribution: Karnataka (Belagavi, Uttar Kannada), Peninsular to E India, Upper Gangetic Plains; endemic.

Justicia L.

Adhatoda Mill.

Dianthera L.

Manathes Bremek.

Monechma Hochst.

Justicia adhatoda L., Sp. Pl. 15. 1753; Stearn, J. Arnold Arbor. 52: 640. 1971.

Adhatoda zeylanica Medik., Hist. & Commentat. Acad. Elect. Sci. Theod.-Palat. 6: 393. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Adhatoda vasica Nees in Wall., Pl. Asiat. Rar. 3: 103. 1832

Distribution: Karnataka (throughout, often cultivated); throughout India; S & SE Asia.

Local/Common/Vernacular names: *Adusoge*, *Vasaka* (Kannada); *Vasaka* (Sanskrit)

Justicia betonica L., Sp. Pl. 15. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

var. betonica

Distribution: Karnataka (throughout, often cultivated), throughout India; S & SE Asia, tropical Africa.

Local/Common/Vernacular names: *Kaadu kanakambra, Sanna adusoge* (Kannada); *Sahachara, Shweta sahachara* (Sanskrit)

var. ramosissima (Roxb.) C.B. Clarke in Hook.f., Fl. Brit. India 4: 525. 1885

Justicia ramosissima Roxb., Fl. Ind. ed. Carey & Wall., 1: 130. 1820 & ed. Carey, 1: 129. 1832

Distribution: Karnataka (throughout), throughout India; endemic.

Justicia concinna J.R.I. Wood, Novon 23(3): 386.2014

Rungia elegans Dalzell & A. Gibson, Bombay Fl. 196. 1861 non *Justicia elegans* P.Beauv. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Distribution: Karnataka (Belagavi, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Kalaburgi, Uttara Kannada), N, W & Peninsular India; endemic.

Local/Common/Vernacular name: *Koda gasale* (Kannada)

Justicia crenatifolia J.R.I. Wood, Novon 23(3): 387.2014

Rungia crenata T.Anderson, J. Linn. Soc., Bot. 9: 518. 1867 non *Justicia crenata* Phol ex Nees 1847; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Distribution: Karnataka (Belagavi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Justicia diffusa Willd., Sp. Pl. 4 : 87. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Rostellularia diffusa (Willd.) Nees in Wall., Pl. Asiat. Rar. 3: 100. 1832

Distribution: Karnataka (Belagavi, Chamarajanagar, Kalaburgi, Yadgir), throughout India; Bangladesh, India, Myanmar, Sri Lanka, Taiwan, Thailand, Vietnam.

Local/Common/Vernacular name: *Nandi nela bevu* (Kannada)

Justicia gendarussa Burm. f., Fl. Indica 10. 1768; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Gendarussa vulgaris Nees in Wall., Pl. Asiat. Rar. 3: 104. 1832

Distribution: Karnataka (throughout, often cultivated), throughout India; S & SE Asia, tropical Africa.

Justicia glabra J. König ex Roxb., Fl. Ind. ed. Carey & Wall., 1: 132. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Rhaphidospora glabra (J.Koenig ex Roxb.) Nees in Wall., Pl. Asiat. Rar. 3: 115. 1832

Rhaphidospora glabrituba Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 45 (2): 48. 1948

Distribution: Karnataka (Bengaluru, Chitradurga, Kodagu, Mysuru), Keral, Tamil Nadu; Sri Lanka, tropical Africa.

Justicia glauca Rottler in Ges. Naturf. Freunde Berlin Neue Schriften 4: 219. 1802; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.

Gendarussa tranquebariensis Wight, Icon. Pl. Ind. Orient. t. 462. 1841 non Nees, 1832

Distribution: Karnataka (Belagavi, Chamarajanagar, Mysuru), Peninsular to E India; endemic.

Justicia heyneana J.R.I. Wood, Novon 23(3): 387.2014

Rostellularia gracilis Wight, Icon. Pl. Ind. Orient. t. 1541. 1850

Justicia gracilis (Wight) T.Anderson, J. Linn. Soc., Bot. 9: 518. 1867 non *Justicia gracilis* Lyall ex Nees 1847

Rungia linifolia Nees in Wall., Pl. Asiat. Rar. 3: 110. 1832 non *Justicia linifolia* (Lindau)V.A.W.Graham 1988; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

var. *heyneana*

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Kerala, Maharashtra, Rajasthan, Tamil Nadu; endemic.

var. *saldanhae* (Mascar. & Janarth.) J.R.I.Wood, Novon 23(3): 387.2014

Rungia linifolia var. *saldanhae* Mascar. & Janarth., Novon 20(2):182.2010

Distribution: Karnataka (Chikkamagaluru); endemic.

***Justicia japonica* Thunb., Fl. jap. 20. 1784; V.S. Ramach. & V.J. Nair, Fl. Cannanore 340. 1988.**

Justicia simplex D.Don, Prodr. Fl. Nepal. 118. 1825.

Rostellularia japonica (Thunb.) J.L. Ellis, Bull. Bot. Surv. India 22: 196. 1982

Distribution: Karnataka (Belagavi, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Kalaburgi, Kodagu, Shivamogga, Uttara Kannada, Yadgir), Peninsular to N India; Africa, S & SE Asia.

***Justicia latispica* (C.B.Clarke) Gamble, Fl. Madras 1080. 1924**

Rostellularia procumbens Wight, Icon. Pl. Ind. Orient. t. 1539. 1850 non Nees, 1832

Justicia procumbens var. *latispica* C.B.Clarke in Hook.f., Fl. Brit. India 4: 539. 1885

Rostellularia latispica (C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 45 (2): 71. 1948

Distribution: Karnataka (Chamarajanagar, Mysuru), throughout except NE India; Sri Lanka.

***Justicia nagpurensis* V.A.W. Graham, Kew Bull. 43: 597. 1988**

Rostellularia procumbens var. *serpyllifolia* Benth. in Hohen., Pl. Ind. Orient. 571. 1847

Justicia simplex var. *serpyllifolia* (Benth.) C.B. Clarke in Hook.f., Fl. Brit. India 4: 539. 1885

Rostellularia serpyllifolia (Benth.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 45 (2): 62. 1948

Rostellularia mollissima var. *serpyllifolia* (C.B.Clarke) Raizada, Suppl. Duthie's Fl. Gangetic plain 209. 1976

Justicia hansenii Sivar. & P. Mathew, Taxon 38: 486. 1989

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Udupi, Uttara Kannada), Peninsular to E India, Upper Gangetic Plains; endemic.

***Justicia neesii* Ramamoorthy in C.J. Saldanha & Nicolson, Fl. Hassan Dist. 551. 1976**

Rostellularia crinita Nees in Wall., Pl. Asiat. Rar. 3: 101. 1832

Justicia micrantha Wall. ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 536. 1885 non Pohl ex Nees, 1847

Rostellularia neesii (Ramamoorthy) A.K. Mukh., Fl. Pachmarhi & Bori Res. 236. 1984

Distribution: Karnataka (Chitradurga, Davanagere, Kalaburgi, Yadgir), Kerala, Maharashtra; endemic.

***Justicia nepalensis* V.A.W. Graham, Kew Bull. 43: 597. 1988**

Rostellularia rotundifolia Nees in Wall., Pl. Asiat. Rar. 3: 100. 1832 non *Justicia rotundifolia* Nees, 1832

Rostellularia rotundifolia (Nees) Nees A.DC., Prodr. 11: 370. 1847

Distribution: Karnataka (), throughout India; Nepal.

***Justicia nilgherrensis* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 526. 1885; B.D.Sharma et al., Fl. Karnataka Anal. 207. 1984.**

Adhatoda nilgherrensis Nees in Wall., Pl. Asiat. Rar. 3: 103. 1832

Distribution: Karnataka (Chamarajanagar, Hassan,), Kerala, Tamil Nadu; endemic.

Justicia orbiculata Wall. ex T.Anderson, J. Linn. Soc., Bot. 9: 512. 1867; Gamble, Fl. Madras 2: 757. 1967 (Repr.).

Justicia diffusa var. *orbiculata* (Wall. ex T.Anderson) C.B. Clarke in Hook.f., Fl. Brit. India 4: 538. 1885

Rostellularia diffusa var. *orbiculata* (Wall. ex T.Anderson) J. L. Ellis, Biol. Mem. 2: 109. 1977

Rostellularia diffusa var. *rotundifolia* J. L. Ellis, Bull. Bot. Surv. India 30: 132. 1990

Distribution: Karnataka (Dakshina Kannada, Mysuru), Peninsular to E India, Gangetic Plains; Indonesia.

Justicia peploides (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 54. 1867

Justicia quinqueangularis J. König ex Roxb., Fl. Ind. ed.1, 1: 134. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Rostellularia peploides Nees in Wall., Pl. Asiat. Rar. 3: 101. 1832

Rostellularia quinqueangularis var. *peploides* (Nees) Raizada, Suppl. Duthie's Fl. Gangetic plain 209. 1976

Distribution: Karnataka (Belagavi, Uttara Kannada) Peninsular to E & N India, Himalayas.

Justicia procumbens L., Sp. Pl. 15. 1753

Rostellularia procumbens (L.) Nees in A.DC., Prodr. 11: 371. 1847

Rostellularia sundana Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 45 (2): 62. 1948

Distribution: Karnataka (throughout), throughout India including Andaman & Nicobar Islands; Australia, S & SE Asia.

Local/Common/Vernacular name: *Nucchu nela bevu* (Kannada)

Justicia prostrata (Roxb. ex C.B.Clarke) Gamble, Fl. Madras 1081. 1924; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Justicia diffusa var. *prostrata* Roxb. ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 538. 1885

Rostellularia prostrata (Roxb. ex C.B.Clarke) R.B. Majumdar, Bull. Bot. Soc. Bengal 25: 75. 1971

Rostellularia diffusa var. *prostrata* (Roxb. ex C.B.Clarke) J. L. Ellis, Bull. Bot. Surv. India 30: 131. 1990

Distribution: Karnataka (Uttara Kannada), throughout tropical India; Malaysia, Sri Lanka.

Local/Common/Vernacular name: *Chikka nekki soppu* (Kannada)

Justicia quinqueangularis K.D.König ex Roxb., Fl. Ind. 1: 134. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Rostellularia quinqueangularis (J.Koenig ex Roxb.) Nees in Wall., Pl. Asiat. Rar. 3: 101. 1832

Distribution: Karnataka(Belagavi, Chitradurga), throughout India; endemic.

Justicia salsoloides T.Anderson, J. Linn. Soc., Bot. 9: 514. 1867; C.B.Clarke in Hook.f., Fl. Brit. India 4: 530. 1885.

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Justicia santapaui Bennet, J. Bombay Nat. Hist. Soc. 67: 358. 1970; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Plate 13G

Hemichoriste montana Nees in Wall., Pl. Asiat. Rar. 3: 102. 1832

Justicia andersonii Ramamoorthy in C.J.Saldanha & Nicolson, Fl. Hassan dist. 551. 1976

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Justicia simplex sensu C. B. Clarke in Hook. f., Fl. Brit. India 4: 539. 1885 p.p.

Rostellularia mollissima Nees in Wall., Pl. Asiat. Rar. 3: 101. 1832

Rostellularia mollissima (Nees) Nees in A.DC., Prodr. 11: 373. 1847

Rostellularia pumila Nees in A.DC., Prodr. 11: 372. 1847

Rostellularia procumbens var. *simplex* (D.Don) T.Yamaz. in H. Hara, Fl. E. Himal. 302. 1966

Distribution: Karnataka (throughout), throughout India; S & SE Asia, tropical Africa.

Local/Common/Vernacular name: *Yelavu sandhi gida* (Kannada)

Justicia tranquebariensis L.f., Suppl. Pl. 85. 1782; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Gendarussa tranquebariensis (L.f.) Nees in Wall., Pl. Asiat. Rar. 3: 105. 1832

Distribution: Karnataka (Bengaluru, Mysuru), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Kaddi arakina gida, Shivanari balli* (Kannada)

Justicia trinervia Vahl, Enum. Pl. 1: 156. 1804; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Justicia vahlii Roth, Nov. Pl. sp. 14. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Rostellularia vahlii (Roth) Nees in Wall., Pl. Asiat. Rar. 3: 102. 1832

Justicia diffusa var. *vahlii* (Roth) C.B. Clarke in Hook.f., Fl. Brit. India 4: 538. 1885

Distribution: Karnataka (throughout), throughout India; endemic.

Justicia vallis-silentosae J.R.I.Wood, Novon 23(3):388. 2014

Rungia sisparensis T.Anderson, J. Proc. Linn. Soc., Bot. 7: 114.1864

Justicia sisparensis Lindau in Engl. & Prantl, Nat. Pflanzenfam. IV, heft 3b:350.1895

Distribution: Karnataka (cultivated throughout); Peninsular India; native to tropical Asia.

Justicia wynaedensis (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 515. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Gendarussa wynaedensis Nees in Wall., Pl. Asiat. Rar. 3: 104. 1832

Adhatoda wynaedensis (Nees) Nees in A.DC., Prodr. (Ed. A. DC.) 11: 406. 1847; Wight, Icon. Pl. Ind. Orient. t. 1545. 1850.

Ecbolium wynadense (B.Heyne) Kuntze, Revis. Gen. Pl. 2: 981.1891

Distribution: Karnataka (Belagavi, Chikamgluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Lepidagathis Willd.

Lepidagathis cristata Willd., Sp. Pl. 3: 400. 1800; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburgi, Kodagu, Raichur, Vijayapura, Yadgir), Peninsular to NE India; endemic.

Local/Common/Vernacular names: *Gantu kaalu gedde, Nari goodi* (Kannada)

Lepidagathis cuspidata Nees in Wall., Pl. Asiat. Rar. 3: 97. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 208. 1984.

Ruellia cuspidata Wall., Numer. List 2405. 1830

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Lepidagathis diffusa C.B.Clarke in Hook.f., Fl. Brit. India 4: 518. 1885; B.D.Sharma et al., Fl.

Karnataka Anal. 208. 1984.

Distribution: Karnataka (Ballari, Mysuru), Andhra Pradesh, Tamil Nadu; endemic.

Lepidagathis fasciculata (Retz.) Nees in Wall., Pl. Asiat. Rar. 3: 95. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Ruellia fasciculata Retz., Observ. Bot. 4: 28. 1786

Lepidagathis goensis Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 340. 1850

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), throughout tropical & subtropical India; endemic.

Lepidagathis hamiltoniana Wall. ex Nees in Wall., Pl. Asiat. Rar. 3: 96. 1832; C.B.Clarke in Hook.f., Fl. Brit. India 4: 516. 1885.

Distribution: Karnataka (Dharwad), E, N & S India; endemic.

Lepidagathis incurva Buch.-Ham. ex D. Don, Prodr. Fl. Nepal. 119.1825; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Lepidagathis hyalina Nees in Wall., Pl. Asiat. Rar. 3: 95. 1832

Lepidagathis mucronata Nees in Wall., Pl. Asiat. Rar. 3: 95. 1832

Lepidagathis hyalina var. *lophostachyoides* Nees in A.DC., Prodr. 11: 252. 1847

Lepidagathis hyalina var. *mucronata* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 521. 1885

Lepidagathis incurva var. *mucronata* (Nees) C.B. Clarke ex T. Cooke, Fl. Bombay 2: 397. 1904; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Lepidagathis incurva var. *lophostachyoides* (Nees) Abeyw., Ceylon J. Sci., Biol. Sci. 2(1): 85. 1959; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada); throughout plains of India & E Himalaya.

Lepidagathis mitis Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 226. 1851; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Lepidagathis prostrata Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 138. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), Goa, Maharashtra, subtropical W Himalaya; endemic.

Local/Common/Vernacular name: *Pare mullu* (Kannada)

Lepidagathis scariosa Nees in Wall., Pl. Asiat. Rar. 3: 95. 1832; C.B.Clarke in Hook.f., Fl. Brit. India 4: 520. 1885.

Distribution: Karnataka (Chamarajanagara ?), Kerala, Tamil Nadu; Ethiopia.

Lepidagathis spinosa Wight ex Nees in Wall., Pl. Asiat. Rar. 3: 97. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka (Chikkamagaluru, Tumakuru), Kerala, Tamil Nadu; endemic.

Lepidagathis subarmata (C.B.Clarke) Gamble, Fl. Madras 1067. 1924

Lepidagathis mitis var. *subarmata* C.B.Clarke in Hook.f., Fl. Brit. India 4: 516. 1885

Distribution: Karnataka (Ballari, Chitradurga, Raichur), Andhra Pradesh, Tamil Nadu; endemic.

Lepidagathis trinervis Nees in Wall., Pl. Asiat. Rar. 3: 96. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka (Belagavi), C, E & W India, Himalayas, Upper Gangetic Plains; endemic.

Meyenia Nees

Meyenia hawtayneana (Wall.) Nees in Wall., Pl. Asiat. Rar. 3: 78. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Thunbergia hawtayneana Wall., Tent. Fl. Napal. 49. 1826 in note; C.B.Clarke in Hook.f., Fl. Brit. India 4: 391. 1884.

Distribution: Karnataka(Belagavi, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Udupi, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Hawtayene's meyenia* (English); *Pera nela birudu* (Kannada)

Nelsonia R. Br.

Nelsonia canescens (Lam.) Spreng., Syst. veg. 1: 42. 1825

Justicia canescens Lam., Tabl. encycl. 1: 41. 1791

Nelsonia campestris R. Br., Prodr. 481. 1810; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka(Belagavi, Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), throughout India including Andaman & Nicobar Islands; Pantropical.

Neuracanthus Nees

Neuracanthus sphaerostachyus (Nees) Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 140. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Lepidagathis sphaerostachya Nees in A.DC., Prodr. 11: 254. 1847

Neuracanthus lawii Wight, Icon. Pl. Ind. Orient. t. 1531. 1850

Distribution: Karnataka (Belagavi, Uttara Kannada), Goa, Gujarat, Maharashtra, Rajasthan; endemic.

Neuracanthus tetragonostachyus

Nees, Pl. Asiat. Rar. 3: 97. 1832.

subsp. **trinervius** (Wight) Bidgood & Brummitt, Kew Bull. 53: 16. 1998

Neuracanthus trinervius Wight, Icon. Pl. Ind. Orient. t. 1532. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Dharwad, Uttara Kannada), Gujarat, Maharashtra; endemic.

Peristrophe Nees

Peristrophe bicalyculata (Retz.) Nees in Wall., Pl. Asiat. Rar. 3: 113 1832; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Dianthera bicalyculata Retz., Kongl. Vetensk. Acad. Handl. 1775: 297. 1775

Justicia malabarica Aiton, Hort. Kew. 1: 27. 1789

Justicia bicalyculata (Retz.) Vahl, Symb. Bot. 2: 13. 1791

Dicliptera bicalyculata (Retz.) Kostel., Allg. Med.-Pharm. Fl. 3: 934. 1834

Distribution: Karnataka (throughout), throughout drier areas of India; S Asia, tropical Africa.

Local/Common/Vernacular names: *Cheebi gida*, *Cibi gida*, *Cibira soppu*, *Yeluva sandaku* (Kannada)

Petalidium Nees

Petalidium barlerioides (Roth) Nees in Wall., Pl. Asiat. Rar. 3: 82. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Ruellia barlerioides Roth, Nov. Pl. sp. 319. 1821

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Shivamogga, Udupi, Uttara Kannada), throughout India; Nepal, Pakistan.

Phaulopsis Willd. ('Phaylopsis') nom. & orth. cons.

Phaulopsis imbricata (Forssk.) Sweet, Hort. brit. 327. 1826

Ruellia imbricata Forssk., Fl. Aegypt.-Arab. 113. 1775

Ruellia dorsiflora Retz., Observ. Bot. 6: 31. 1791

Micranthus oppositifolius J.C. Wendl., Bot. Beob. 39. 1798

Phaulopsis parviflora Willd., Sp. Pl. 3: 342. 1800

Aetheilema reniforme Nees in Wall., Pl. Asiat. Rar. 3: 94. 1832

Phaulopsis oppositifolia (J.C.Wendl.) Lindau in Engl. & Prantl, Nat. Pflanzenfam. I, 1, 148: 305. 1897

Micranthus dorsiflorus (Retz.) C.E.C. Fisch., Bull. Misc. Inform. 1932: 63. 1932

Phaulopsis dorsiflora (Retz.) Santapau, Kew Bull. 1948: 276. 1948; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (throughout), throughout India; Madagascar, S & SE Asia, tropical Africa.

Pseuderanthemum Radlk.*Eranthemum* sensu C.B.Clarke

Pseuderanthemum bicolor (Schrink) Radlk. ex Lindau in Engl. & Prantl, Nat. Pflanzenfam. IV, 3b: 330, f. 133, A-D. 1895; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Eranthemum bicolor Schrank, Pl. Rar. Hort. Monac. t. 8. 1819

Distribution: Karnataka (cultivated in Belagavi, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), W India; native to Philippines.

Pseuderanthemum carruthersii (Seem.) Guillaumin, Ann. Mus. Colon. Marseille VI, 5-6: 48 1948; K. G. Bhat, Fl. S. Kanara 792. 2014.

Eranthemum atropurpureum W.Bull, Gard. Chron. n.s., 3: 619. 1875

Pseuderanthemum atropurpureum (W.Bull) Radlk., Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 13: 286. 1884

Distribution: Karnataka (cultivated throughout), nativity to S Africa, cultivated throughout tropics.

Pseuderanthemum lapathifolium (Vahl) B. Hansen, Nord. J. Bot. 9: 214. 1989

Justicia lapathifolia Vahl, Enum. Pl. 1: 126. 1804

Eranthemum lapathifolium (Vahl) Nees in A.DC., Prodr. (Ed. A.DC.) 11: 454. 1847

Distribution: Karnataka (), Nicobar Islands, Peninsular India.

Pseuderanthemum latifolium (Vahl) B. Hansen, Nord. J. Bot. 9: 213. 1989

Justicia latifolia Vahl, Symb. Bot. 2: 4. 1791

Justicia palatifera Wall., Pl. Asiat. Rar. 1: 80, t. 92, f. 1-3. 1830

Eranthemum palatiferum (Wall.) Nees in Wall., Pl. Asiat. Rar. 3: 108. 1832; C.B.Clarke in Hook.f., Fl. Brit. India 4: 498. 1884.

Pseuderanthemum palatiferum (Wall.) Radlk., Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 13: 286. 1884

Antheliacanthus micranthus Ridl., J. Fed. Malay States Mus. 10: 109. 1920

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Goa, Maharashtra; endemic.

Pseuderanthemum malabaricum (C.B.Clarke) Gamble, Fl. Madras 1064. 1924; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Eranthemum malabaricum C.B.Clarke in Hook.f., Fl. Brit. India 4: 497. 1884

Distribution: Karnataka(Belagavi, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Rhinacanthus Nees

Rhinacanthus nasutus (L.) Kurz, J.Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 39: 79. 1870; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Justicia nasuta L., Sp. Pl. 16. 1753

Rhinacanthus communis Nees in Wall., Pl. Asiat. Rar. 3: 109. 1832 nom. illeg.

Distribution: Karnataka (throughout), throughout India; Madagascar, S & SE Asia.

Local/Common/Vernacular names: *Ringworm root, Snake jasmine* (English); *Dodda patika, Dodda pettika gida, Naaga mallige* (Kannada); *Yuthikaparani* (Sanskrit); *Naaga malle* (Tulu)

Ruellia L.

Ruellia beddomei C.B.Clarke, Fl. Brit. India 4: 413. 1884; C.E.C.Fischer, Bull. Misc. Inform. Kew 1940 (7): 301. 1940.

Distribution: Karnataka (Chamarajanagar), Madhya Pradesh, Chhattisgarh, Odisha, West Bengal, Jharkhand, Bihar, Uttarkhand, Tamil Nadu; endemic.

Ruellia patula Jacq., Misc. Austriac. 2: 358. 1781

Dipteracanthus patulus (Jacq.) Nees in Wall., Pl. Asiat. Rar. 3: 81. 1832

Petalidium patulum (Jacq.) Dalzell & A.Gibson, Bombay Fl. 185.1861

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Kalaburgi, Mysuru, Raichur, Uttara Kannada, Vijayapura), throughout India; Tropical Africa, Arabia, Pakistan, Sri Lanka.

Ruellia prostrata Poir. in Lam., Encycl. 6: 349. 1804

Ruellia prostrata var. *dejecta* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 412. 1884

Distribution: Karnataka (Mysuru), throughout India;Pakistan.

Ruellia tuberosa L., Sp. Pl. 635. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dakshina Kannada, Dharwad, Kalaburgi, Udupi, Uttara Kannada, Yadgir), throughout India; native to tropical America.

Local/Common/Vernacular name: *Minnie root* (Kannada)

Rungia Nees

Rungia muralis Nees in A.DC., Prodr. 11: 470. 1847

Rungia parviflora var. *muralis* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 550. 1885

Distribution: Karnataka (), Peninsular to NE India; Bangladesh, Myanmar.

Rungia parviflora (Retz.) Nees in Wall., Pl. Asiat. Rar. 3: 110. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Justicia parviflora Retz., Observ. Bot. 5: 9. 1788

Distribution: Karnataka (throughout), throughout India including Himalayas; S & SE Asia.

Rungia pectinata (L.) Nees in A.DC., Prodr. 11: 470. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 212. 1984.

Justicia pectinata L., Amoen. Acad. 4: 299. 1759

Dicliptera pectinata (L.) Juss., Ann. Mus. Hist. Nat. 9: 169.1807

Dianthera parviflora Roxb. ex Nees in A.DC., Prodr. 11: 471.1847

Rungia parviflora var. *pectinata* (L.) C.B. Clarke in Hook.f., Fl. Brit. India 4: 550. 1885

Diapodium pectinatum (L.) Kuntze, Revis. Gen. Pl. 2: 484.1891

Rungia pectinata var. *clarkeana* Hand.-Mazz., Symb. Sin. 7(4): 898.1936

Rungia parviflora subsp. *pectinata* (L.) L.H. Cramer in Dassan. & Clayton, Rev. Handb. Fl. Ceylon 12: 105. 1998

Distribution: Karnataka (Belagavi, Ballari, Bidar, Chamarajanagar, Chikkamagaluru, Chitraurga, Dakshina Kannada, Dharwad, Hassan, Kalaburgi, Kodagu, Shivamogga, Udupi, Uttara Kannada, Yadgir), throughout India including Andaman & Nicobar Islands; Bangladesh, Bhutan, Laos, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular name: *Comb rungia* (Kannada)

Rungia repens (L.) Nees in Wall., Pl. Asiat. Rar. 3: 110. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Justicia repens L., Sp. Pl. 15. 1753

Distribution: Karnataka (Belagavi, Ballari Bengaluru, Chamarajanagar, Chitradurga, Davanagere, Hassan, Kalaburgi, Mysuru, Shivamogga, Uttara Kannada, Vijayapura, Yadgir), throughout India; Sri Lanka.

Local/Common/Vernacular names: *Koda gasale gida*, *Sudaabu* (Kannada); *Parapaata* (Sanskrit)

Staurogyne Wall.

Ebermaiera Nees .

Staurogyne glutinosa (Wall. ex C.B. Clarke) Kuntze, Revis. Gen. Pl. 2: 497. 1891

Ebermaiera glutinosa Wall. ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 396. 1884

Distribution: Karnataka (), throughout except NW India, Himalayas & Islands; Bangladesh, Myanmar.

Staurogyne spatulata (Blume) Koord., Exkurs.-Fl. Java 3: 211. 1912

Adenosma spatulata Blume, Bijdr. 757. 1826

Ebermaiera glauca Nees, Compan. Bot. Mag. 2: 310. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Ebermaiera glauca var. *spatulata* (Blume) C. B. Clarke in Hook. f., Fl. Brit. India 4: 396. 1884

Staurogyne glauca (Nees) Kuntze, Revis. Gen. Pl. 2: 497. 1891

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), throughout except NW India, Himalayas & Islands; Myanmar.

Staurogyne zeylanica (Nees) Kuntze, Revis. Gen. Pl. 2: 497. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Ebermaiera zeylanica Nees in A.DC., Prodr. 11: 74. 1847

Distribution: Karnataka(Belagavi, Chamarajanagar, Dakshina Kannada, Hassan, Kalaburgi, Shivamogga, Udupi, Uttara Kannada, Yadgir), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Strobilanthes Blume

Baphiacanthus Bremek.

Buteraea Nees

Carvia Bremek.

Didyplosandra Wight ex Bremek.

Diffloglussa Bremek.

Ditrichospermum Bremek.

Dossifluga Bremek.

Goldfussia Nees

Hemigraphis Nees

Hymenochlaena Bremek.

Kanjarum Ramam.

Leptacanthus Nees

Listrobantnes Bremek.

Makenziea Nees

Niligirianthus Bremek.

Parasympagis Bremek.

Perilepta Bremek.

Phlebophyllum Nees

Pleocaulus Bremek.

Pseudaechmanthera Bremek.

Pseudostenosiphonium Lindau

Ptercanthus (Nees) Bremek.

Pyrrothrix Bremek.

Sericocalyx Bremek.

Stenosiphonium Nees

Strobilanthes subg. *Pteracanthus* Nees

Strobilanthes subg. *Sympagis* Nees

Supushpa Suryan.

Sympagis (Nees) Bremek.

Taeniandra Bremek.

Tarphochalmys Bremek.

Tetragoga Bremek.

Thelipaepale Bremek.

Triaenacacanthus Nees ('*Triaenanthus*')

Xenacanthus Bremek.

Strobilanthes alternata (Burm.f.) J.R.I.Wood, Novon 23(3): 389. 2014

Ruellia alternata Burm.f., Fl. Indica 135.1768

Ruellia colorata Blume, Bijdr. Fl. Ned. Ind. 795.1826

Hemigraphis alternata (Burm.f.) T.Anderson, J. Proc. Linn. Soc., Bot. 7: 114.1864

Hemigraphis colorata (Blume) Hallier f., Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur. 70(3): 199.1897; N.M.G.Babu, Fl. Ethnob. Chitradurga Dist. 437. 2011.

Distribution: Karnataka (cultivated ornamental), cultivated ornamental throughout India; native to Indonesia, Malaysia; widely cultivated, naturalised in tropics.

Strobilanthes anamallaica J.R.I.Wood, Kew Bull. 50: 10.1995; K.G.Bhat, Fl. S. Kanara 796. 2014.

Strobilanthes adenophorus auct. non Nees, 1847; Bedd.,Icon. Pl. Ind. Or. 53.1874.

Nilgirianthus beddomei Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2, 41(1): 172.1944

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Strobilanthes aurita J.R.I. Wood, Kew Bull. 50: 12. 1995

Mackenziea caudata Ramamoorthy in C. J. Saldanha & Nicolson, Fl. Hassan Dist. 553. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Udupi), Kerala; endemic.

Strobilanthes barbata Nees in Wall., Pl. Asiat. Rar. 3: 85. 1832

Strobilanthes tetraptera Dalzell in Hooker's J. Bot. Kew Gard.Misc. 2:342.1850

Nilgirianthus barbatus (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41(1): 172. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Strobilanthes barbata var. *bonaccordensis* E.S.S.Kumar & R. R. Vikram., J. Econ. Taxon. Bot. 25: 742. 2002

Distribution: Karnataka (Belagavi, Chamarajanagar, Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes callosa Nees in Wall., Pl. Asiat. Rar. 3: 85. 1832

Plate 14C

Strobilanthes grahamiana Wight, Icon. Pl. Ind. Orient. 4 : t. 1520. 1850

Strobilanthes callosa var. *hispida* C.B.Clarke in Hook.f., Fl. Brit. India 4: 451. 1884

Carvia callosa (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 187. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 204. 1984.

Carvia callosa var. *hispida* (C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 187. 1944

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Dharwad, Shivamogga, Uttara Kannada), Goa, Gujarat, Maharashtra, Rajasthan; endemic.

Local/Common/Vernacular name: *Karki* (Kannada); *Karvi* (Kunhi)

Strobilanthes campanulata Wight, Icon. Pl. Ind. Orient. 4: t. 1562. 1849

Strobilanthes involucrata var. β Nees in A.DC., Prodr. 11: 184. 1847

Strobilanthes heyneana var. *campanulata* (Wight) C.B. Clarke in Hook.f., Fl. Brit. India 4: 444. 1884

Nilgirianthus campanulatus (Wight) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 172. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Kodagu, Shivamogga, Uttara Kannada), Chattisgarh, Madhya Pradesh; endemic.

Strobilanthes canarica Bedd., Icon. Pl. Ind. Or. t. 215. 1874

Phlebophyllum canaricum (Bedd.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 169. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan), Kerala, Tamil Nadu; endemic.

Strobilanthes carinei J.R.I.Wood, Novon 23(3): 388. 2014

Stenosiphonium parviflorum T.Anderson, J. Linn. Soc., Bot. 9: 464. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984; non *Strobilanthes parviflora* Bedd. 1868-1874.

Stenosiphonium setosum T.Anderson, J. Linn. Soc., Bot. 9: 464. 1867 non

Strobilanthes setosa J.R.I.Wood, Edinburgh J. Bot. 51: 259 1994

Distribution: Karnataka (Bengaluru, Kodagu, Mysuru, Shivamogga), Kerala, Tamil Nadu; endemic.

Strobilanthes ciliata Nees in Wall., Pl. Asiat. Rar. 3: 85. 1832; C.B.Clarke in Hook.f., Fl. Brit. India 4: 439. 1884.

Goldfussia zenkeriana Wight, Icon. Pl. Ind. Orient. 4: t. 1517. 1850

Nilgirianthus ciliatus (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 172. 1944

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes consanguinea (Nees) T.Anderson in Thwites, Enum. Pl. Zeyl. 226 1860

Ruellia spicata Roth, Nov. Pl. Sp. 310. 1821 non *Strobilanthes spicata* T.Anderson

Endopogon consanguineus Nees, Pl. Asiat. Rar. 3: 99. 1832

Phlebophyllum spicatum (Roth) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2 41(1): 169. 1944

Distribution: Karnataka (Ballari, Bengaluru, Kolar, Mysuru), Kerala, Odisha, Tamil Nadu; endemic.

Strobilanthes cordifolia (Vahl) J.R.I.Wood, Novon 23(3): 390. 2014

Ruellia cordifolia Vahl, Symb. Bot. 3: 84. 1794

Stenosiphonium russelianum Nees in Wall., Pl. Asiat. Rar. 3: 84. 1832; N.M.G.Babu, Fl. Ethnob. Chitradurga Dist. 448. 2011.

Stenosiphonium subsericeum Nees in Wall., Pl. Asiat. Rar. 3: 84. 1832

Stenosiphonium confertum Nees in DC., Prodr. 11: 105. 1847

Stenosiphonium russelianum var. *subsericeum* (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 464. 1867

Stenosiphonium cordifolium (Vahl) Alston in Trimen, Handb. Fl. Ceylon 6(Suppl.): 226. 1931

Stenosiphonium cordifolium var. *subsericeum* (Nees) L.H.Cramer in M.D.Dasanayake & W.D.Clayton, Revised Handb. Fl.Ceylon 12:76. 1998

Distribution: Karnataka (Chitradurga), S India; endemic.

Strobilanthes crossandra (Steud.) J.R.I.Wood, Novon 23(3):391. 2014

Ruellia diffusa Nees in Wall., Pl. Asiat. Rar. 3:83. 1832 non *Ruellia diffusa* Vell. 1825

Ruellia crossandra Steud., Nomencl. Bot. ed.2, 2:481. 1841

Ruellia crenata Benth. ex Hohen. in Flora 32: 558. 1849

Hemigraphis elegans var. *crenata* (Benth. ex Hohen.) C.B. Clarke in Hook.f., Fl. Brit. India 4: 425. 1884

Hemigraphis crenata (Benth. ex Hohen.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 137. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Hemigraphis crossandra (Steud.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 137. 1944

Distribution: Karnataka (Ballari, Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada); throughout India; Myanmar.

Strobilanthes cuspidata (Benth.) T.Anderson, J. Linn. Soc., Bot. 9: 465. 1867; Bedd., Icon. Pl. Ind. Or. t. 221. 1874.

Endopogon versicolor Wight, Icon. Pl. Ind. Prient. 4: t. 1497. 1849

Endopogon cuspidatus Benth., Linnaea 24: 646. 1851

Phlebophyllum versicolor (Wight) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 169. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Dakshina Kannada), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Strobilanthes decurrens (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 470. 1867 non

Peristrophe montana auct. non Nees, 1832; V.S. Ramach. & V.J. Nair, Fl. Cannanore 346.1988

Goldfussia decurrens (Nees) Wight, Icon. Pl. Ind. Orient. 4: t. 1522. 1850

Nilgirianthus decurrens (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41(1): 173. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Kodagu, Shivamogga), Kerala, Tamil Nadu; endemic.

Strobilanthes dyeriana Mast., Gard. Chron. 1: 442. 1893

Perilepta dyeriana (Mast.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 194. 1944

Distribution: Karnataka (cultivated in gardens), cultivated throughout India; native to Myanmar

Strobilanthes foliosa (Wight) T.Anderson, J. Linn. Soc., Bot. 9: 467. 1867; C.B.Clarke in Hook.f., Fl. Brit. India 4: 433. 1884.

Endopogon capitatus Wight, Icon. Pl. Ind. Orient.4: t. 1499. 1849

Endopogon foliosus Wight, Icon. Pl. Ind. Orient. 4: t. 1501. 1849

Strobilanthes foliosa var. *capitata* (Wight) T.Anderson, J. Linn. Soc., Bot. 9: 467. 1867

Nilgirianthus foliosus (Wight) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Chamarajanagar, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Strobilanthes heteromalla T.Anderson ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 437. 1884

Xenacanthus heteromallus (T.Anderson ex C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 176. 1944

Distribution: Karnataka (Uttara Kannada), Kerala, Tamil Nadu; endemic.

Strobilanthes heyneana Nees in Wall., Pl. Asiat. Rar. 3: 85. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Strobilanthes asperrima Nees in A.DC., Prodr. 11: 183. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Strobilanthes tenera Nees in A.DC., Prodr. 11: 184.1847

Strobilanthes trifida Nees in Compan. Bot. Mag. 2: 312.1836 & in DC., Prodr. 11: 185.1847

Strobilanthes rugosa Wight, Icon. Pl. Ind. Orient. 4: t. 1619. 1850

Strobilanthes heyneana var. *fusca* C.B.Clarke in Hook.f., Fl. Brit. India 4: 444. 1884

Strobilanthes heyneana var. *viridis* C.B.Clarke in Hook.f., Fl. Brit. India 4: 444. 1884

Nilgirianthus heyneanus (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944

Nilgirianthus heyneanus var. *fuscus* (C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944

Nilgirianthus heyneanus var. *neesii* Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect.2, 41 (1): 173. 1944

Nilgirianthus heyneanus var. *viridis* (C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Odisha, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Gurgi* (Kannada)

Strobilanthes integrifolia (Dalzell) Kuntze, Revis. Gen. Pl. 2: 499. 1891

Endopogon integrifolius Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 324. 1850

Leptacanthus alatus Wight, Icon. Pl. Ind. Orient. 4: t. 1527. 1850

Strobilanthes perfoliata T.Anderson, J. Linn. Soc., Bot. 9: 471. 1867

Mackenziea integrifolia (Dalzell) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 182. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 209. 1984.

Distribution: Karnataka (Dakshina Kannad, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes ixiocephala Benth., Flora 32: 557. 1849

Strobilanthes neesiana Wight, Icon. Pl. Ind. Orient. 4: t. 1523. 1850

Thelepaepale ixiocephala (Benth.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 41 (1): 188. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Distribution: Karnataka (Dakshina Kannad, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes jogensis Gilli, Ann. Nat. Mus. Wien. 59:136. 1953; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Acanthopale jogensis Gilli, Ann. Naturhist. Mus. Wien 59: 135.1953; B.D.Sharma et al., Fl. Karnataka Anal. 200. 1984.

Distribution: Karnataka (Sivamogga); endemic.

Strobilanthes kunthiana (Nees) T.Anderson ex Benth., Fl. Hongk. 262. 1861

Plate 1G

Phlebophyllum kunthianum Nees in Wall., Pl. Asiat. Rar. 3: 83. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Chamarajanagara), Kerala, Tamil Nadu; endemic.

Strobilanthes lanata Nees in A.DC., Prodr. 11: 191. 1847

Strobilanthes gossypina T.Anderson, J. Linn. Soc., Bot. 9: 466. 1867

Phlebophyllum lanatum (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 169. 1944

Distribution: Karnataka (Chamarajanagara); endemic.

Strobilanthes lawsoni Gamble, Bull. Misc. Inform. 1923: 374. 1923 & Fl. Madras 1036. 1924

Phlebophyllum lawsonii (Gamble) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2 41(1): 169.1944

Strobilanthes pushpangadanii E.S.S.Kumar, Jabbar & A.E.S.Khan, Rhedea 10: 73.2002

Strobilanthes gamblei Carine, J.M. Alexander & Scotland, Kew Bull. 59: 5. 2004

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Strobilanthes lupulina T.Anderson, J. Linn. Soc. Bot. 9: 469. 1867, p.p.

Nilgirianthus lupulinus (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944

Distribution: Karnataka (Belagavi, Chamarajanagar, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes lurida Wight, Icon. Pl. Ind. Orient. 4: t. 1515.1850

Didyplosandra lurida (Wight) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2 41(1): 178.1944; Kesh. Murthy & Yogan., Fl Coorg. 330. 1990.

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Strobilanthes meeboldii Craib, Bull. Misc. Inform. 1910: 278. 1910; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Nilgirianthus meeboldii (Craib) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944

Distribution: Karnataka (Chamarajanagar, Uttara Kannada); endemic.

Strobilanthes membranacea Talbot, Syst. List Trees Bombay ed. 2. 61. 1902

Nilgirianthus membranaceus (Talbot) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 280. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Chikkamagaluru, Uttara Kannada), peninsular India; endemic.

Strobilanthes microstachya Benth. ex Hohen., Flora 32: 557. 1849

Ruellia punctata Nees in A.DC., Prodr. 11: 147. 1847

Strobilanthes anceps var. *microstachya* (Benth.) C.B. Clarke in Hook.f., Fl. Brit. India 4: 442. 1884

Nilgirianthus punctatus (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Dakshina Kannada, Shivamogga), Kerala, Tamil Nadu; endemic.

Strobilanthes minor Talbot, Syst. list. trees Bombay ed. 2. 262. 1902; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Distribution: Karnataka (Uttara Kannada); endemic.

Strobilanthes neilgherrensis Bedd., Icon. Pl. Ind. Orient. t. 196. 1874; C.B.Clarke in Hook.f., Fl. Brit. India 4: 438. 1884.

Nilgirianthus neilgherrensis (Bedd.) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Strobilanthes neoasper Venu & P. Daniel in P. Daniel & Sanjappa, Strobilanthes Penins. India 154. 2006

Strobilanthes asper auct. non Decne., 1834; Wight, Icon. Pl. Ind. Orient. 4: t. 1518. 1850.

Strobilanthes warreensis Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 341. 1850

Nilgirianthus warreensis auct. non Bremek., 1944; B.D.Sharma et al., Biol. Mem. 2: 108. 1977.

Nilgirianthus asper Santapau, Bot. Mem. Univ. Bombay 2: 42. 1952; B.D.Sharma et al., Fl. Karnataka Anal. 210. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga), peninsular India; endemic.

Strobilanthes newii Bedd. ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 464. 1884

Mackenziea newii (Bedd. ex C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 182. 1944

Distribution: Karnataka (Hassan), Kerala; endemic.

Strobilanthes papillosa T.Anderson, J. Linn. Soc., Bot. 9: 468. 1867

Nilgirianthus papillosum (T.Anderson) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 41(1): 173. 1944

Distribution: Karnataka (Chamarajanagar), Kerala, Tamil Nadu; endemic.

Strobilanthes pavala (Roxb.) J.R.I.Wood, Novon 23(3):392.2014

Ruellia latebrosa B. Heyne ex Roth, Nov. Pl. sp. 307. 1821

Ruellia pavala Roxb., Fl.Ind. ed.2, 3:47.1832

Ruellia elegans Hook. in Curtis's Bot. Mag. 62 (N.S. 9): t.3389.1835

Ruellia ebracteata Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 342.1850

Hemigraphis rupestris B. Heyne ex T.Anderson, J. Linn. Soc., Bot. 9: 462. 1867

Hemigraphis pavala (Roxb.) Kurz, J. Asiat. Soc. Bengal, Pt.2, Nat.Hist. 39(2): 78.1870

Ruellia satpoorensis Wawra, Oesterr. Bot. Z. 31:281.1881

Hemigraphis ebracteata (Dalzell) C.B.Clarke in Hook.f., Fl. Brit. India 4: 424. 1884

Hemigraphis latebrosa var. *beddomei* C.B.Clarke in Hook.f., Fl. Brit. India 4: 424. 1884; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Hemigraphis latebrosa var. *rupestris* (B.Heyne ex T.Anderson) C.B. Clarke in Hook.f., Fl. Brit. India 4: 424. 1884

Hemigraphis latebrosa var. *ebracteata* (Dalzell) Cooke, Fl. Bombay 2: 353. 1905 ('ebracteolata').

Hemigraphis latebrosa var. *incana* Gamble, Fl. Madras 2: 1019. 1921

Hemigraphis latebrosa (B.Heyne ex Roth) Nees in A.DC., Prodr. 11: 722. 1847 non Strobilanthes latebrosa Ridley 1922; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Hemigraphis latebrosa var. *heyneana* Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 139. 1944

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Shivamogga, Udupi, Uttara Kannada), throughout India; endemic.

Strobilanthes perrottetiana Nees in DC., Prodr. 11: 179. 1847

Nilgirianthus perrottetianus (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2, 41(1): 173. 1944

Distribution: Karnataka (Kodagu ?), Kerala, Tamil Nadu; endemic.

Strobilanthes pulneyensis C.B.Clarke in Hook.f., Fl. Brit. India 4: 438. 1884

Strobilanthes circarensis Gamble, Bull. Misc. Inform. 1923: 373. 1923 & Fl. Madras 1038. 1924

Nilgirianthus circarensis (Gamble) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944

Xenacanthus pulneyensis (C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41(1): 176. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Distribution: Karnataka (Chamarajanagar, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Strobilanthes reticulata Stapf, Bull. Misc. Inform. 1894: 347. 1894

Nilgirianthus reticulatus (Stapf) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41(1): 173. 1944

Pleocaulus reticulatus (Stapf) M.R.Almeida, Fl. Maharashtra 4A: 81. 2003

Distribution: Karnataka (Chitragurga ?), Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes rubicunda T.Anderson, J. Linn. Soc., Bot. 9: 483. 1867

Leptacanthus rubicundus Nees, Pl. Asiat. Rar. 3: 90. 1832

Distribution: Karnataka (Kodagu), Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes scrobiculata Dalzell ex C.B. Clarke in Hook.f., Fl. Brit. India 4: 445. 1884

Supushpa scrobiculata (Dalzell ex C.B.Clarke) Suryan., Proc. Kon. Ned. Akad. Wetensch., ser. C, 73: 428. 1970; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Maharashtra; endemic.

Strobilanthes sessilis Nees in Wall., Pl. Asiat. Rar. 3: 85. 1832

Pleocaulus sessilis (Nees) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 185. 1944

var. **sessilis**

Distribution: Karnataka (Belagavi, Chikkamagaluru, Kodagu, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

var. **ritchiei** C.B.Clarke in Hook.f., Fl. Brit. India 4: 452. 1884

Pleocaulus ritchiei (C.B.Clarke) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41(1): 185. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, kodagu, Shivamogga, Uttara Kannada), Maharashtra; endemic.

var. **sessiloides** Wight, Icon. Pl. Ind. Orient.4: t. 1512. 1850

Strobilanthes sessiloides Dalzell & A.Gibson, Bombay Fl. 187. 1861, p.p.

Pleocaulus sessiloides (Wight) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2, 41(1): 185. 1944

Pleocaulus sessilis var. *sessiloides* (Wight) Vajr., Fl. Palghat Dist. 358. 1990

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru), Kerala, Maharashtra, Tamil Nadu; endemic.

Strobilanthes sinuata J.R.I.Wood, Novon 23(3):393. 2015

Ruellia repanda L., Sp. Pl. ed. 2. 886. 1763 non *Strobilanthes repanda* (Blume) J.R.I. Wood 2003

Hemigraphis repanda (L.) Hallier f., Nov. Actorum Acad. Caes. Leop.-Carol. German Nat. Cur. 70: 198. 1897; Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 135. 1944.

Distribution: Karnataka (cultivated), cultivated in gardens of India; Indonesia.

Strobilanthes tristis (Wight) T.Anderson, J. Linn. Soc., Bot. 9: 470. 1867

Goldfussia tristis Wight, Icon. Pl. Ind. Orient. 4: t. 1508. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 205. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada), Kerala, Tamil Nadu; endemic.

Strobilanthes urceolaris Gamble, Bull. Misc. Inform. 1923: 374. 1923

Nilgirianthus urceolaris (Gamble) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 173. 1944; B.D.Sharma et al., Fl. Karnataka Anal. 211. 1984.

Distribution: Karnataka (Chamarajanagar), Kerala, Tamil Nadu; endemic.

Strobilanthes urens (Roth) J.R.I. Wood, Novon 23(3):393.2014

Ruellia crispa auct. non L., 1753, non Nees; B.D.Sharma et al., Fl. Karnataka Anal. 206. 1984.

Ruellia urens Roth, Nov. Pl. Sp. 302. 1821

Ruellia dura Nees, Hook., Compan. Bot. Mag. 2: 311. 1836 & in A.DC., Prodr. 11: 146. 1847

Hemigraphis dura (Nees) T.Anderson, J. Linn. Soc., Bot. 9: 461. 1867 (excl. syn. *Gantelbua urens* (Roth) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Tweede Sect. 2, 41 (1): 148. 1944; Santapau, Bot. Mem. Univ. Bombay 2: 30. 1952).

Hemigraphis urens (Roth) J.R.I. Wood, Edinburgh J. Bot. 51: 108. 1994

Distribution: Karnataka (Dharwad, Kalaburagi, Vijayapura), Peninsular India to Upper Gangetic Plains.

Strobilanthes warreensis Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 341.1850

Strobilanthes parviflorus Bedd., Icon. Pl. Ind. Orient. 1: t. 197. 1839

Nilgirianthus warreensis (Dalzell) Bremek., Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2 41(1): 173 1944.; B.D.Sharma et al., Biol. Mem. 2: 108. 1977.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Thunbergia Retz., nom. cons.***Thunbergia alata*** Bojer ex Sims in Curtis, Bot. Mag. 52: t. 2591. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Distribution: Karnataka (cultivated throughout, sometimes run wild), cultivated widely & naturalized; native to tropical Africa.

Local/Common/Vernacular name: *Black eyed susan* (English)

Thunbergia coccinea Wall., Tent. Fl. Napal. 49, 58, t. 37. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 213. 1984.

Distribution: Karnataka (cultivated in gardens), cultivated throughout India; native to tropical Asia Local names/Common names/Vernacular names: Scarlet Clock Vine, Scarlet thunbergia (English)

Thunbergia erecta (Benth.) T.Anderson, J. Linn. Soc., Bot. 7: 18.1864; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Meyenia erecta Benth. in Hook., Niger Fl. 476. 1849

Distribution: Karnataka (cultivated in gardens), cultivated throughout India; native to tropical Africa.

Local/Common/Vernacular names: *Bush clockvine, King's-mantle* (English); *Krishna hoovu* (Kannada)

Thunbergia grandiflora (Roxb. ex Rottl.) Roxb. in Bot. Reg. 6: 495. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Flemingia grandiflora Roxb. ex Rottl., Neue Schriften Ges. Naturf. Freunde Berlin 4: 202.1803

Pleuremidis grandiflora (Roxb. ex Rottl.) Raf., Fl. Tellur. 4: 67.1838

Thunbergia cordifolia Nees in A.DC., Prodr. 11: 35 1847

Thunbergia lacei Gamble, Bull. Misc. Inform. Kew 1913(3): 116.1913

Thunbergia clarkei T. Yamaz., J. Jap. Bot. 46: 52. 1971

Distribution: Karnataka (cultivated throughout in gardens & nurseries), throughout India; native to Sino-Indian region.

Thunbergia laevis Nees in Wall., Pl. Asiat. Rar. 3: 77. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Thunbergia fragrans var. *laevis* (Nees) C.B. Clarke in Hook.f., Fl. Brit. India 4: 391. 1884

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Udupi, Uttara Kannada), E, N & Peninsular India; endemic.

Local/Common/Vernacular name: *Sweet clock vine* (English)

Thunbergia laurifolia Lindl., Gard. Chron. 1856: 260. 1856; C.B.Clarke in Hook.f., Fl. Brit. India 4: 392. 1884.

Distribution: Karnataka (cultivated in gardens), Andaman & Nicobar Islands, E India; native to India.

Local/Common/Vernacular names: *Blue trumpet vine*, *Laurel clock vine* (English)

Thunbergia mysorensis (Wight) T.Anderson, J. Linn. Soc., Bot. 9: 448. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Plate 14D

Hexacentris mysorensis Wight, Icon. Pl. Ind. Orient. 3: t. 871. 1844

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada, wild in Western Ghats, often cultivated), also cultivated, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Muzerabad creeper* (English); *Kamana billu balli* (Kannada)

Achariaceae

Hydnocarpus Gaertn.

Hydnocarpus alpina Wight, Icon. Pl. Ind. Orient. 3: t. 942. 1845; C.J.Saldanha, Fl. Karnataka 1: 272. 1984

Distribution: Karnataka (Dakshina Kannada, Hassan), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Sanuasotti*, *Torathi* (Kannada)

Hydnocarpus pentandrus (Buch.-Ham.) Oken, Allg. Naturgesch. 3: 1381 1841; B.D.Sharma et al., Fl. Karnataka Anal. 13. 1984.

Plate 14E

Munnicksia laurifolia Dennst., Hort. Malab. 27. 1818

Chilmoria pentandra Buch.-Ham. in Trans. Linn. Soc. London 13: 501 1822

Distribution: Karnataka (Chikkamagaluru, Davanagere, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Hydnocarpus wightianus Blume, Rumphia 4: 22. 1848; B.D.Sharma et al., Fl. Karnataka Anal. 13. 1984.

Hydnocarpus laurifolius Sleumer in Engl., Bot. Jahrb. Syst. 69: 33. 1938

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Fish Poison tree* (English); *Chalmogra*, *Chaul moogra yenne mara*, *Garudaphala*, *Mirolhakai*, *Shunti mara*, *Surani*, *Suranti*, *Surti*, *Toratti* (Kannada); *Tuvaraka*, *Turveraka*, *Tuvrak*, *Kushtavairi* (Sanskrit)

Acoraceae

Acorus L.

Acorus calamus L., Sp. Pl. 324.1753; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Distribution: Karnataka (cultivated in wet places in most districts), cultivated throughout India except in desert area; tropical Asia.

Local/Common/Vernacular name: *Sweet flag* (English); *Baje* (Kannada)

Adoxaceae

Viburnum L.

Viburnum cylindricum Buch.-Ham. ex D. Don, Prodr. Fl. Nep. 142. 1825

Viburnum coriaceum Blume, Bijdr. 656.1826

Viburnum crassifolium Rehder, Trees & Shrubs 2(2): 112. 1908

Viburnum cylindricum var. *crassifolium* (Rehder) C.K. Schneid., Bot. Gaz. 64(1): 77.1917

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; China, S & SE Asia.

Viburnum punctatum Buch.-Ham. ex D. Don, Prodr. Fl. Nepal. 142.1825

Viburnum acuminatum Wall. ex DC., Prodr. 4: 325.1830

Distribution: Karnataka (Chamarajanagar, Hassan), Kerala, Tamil Nadu; China, S & SE Asia.

Local/Common/Vernacular names: *Kampari*, *Nonna*, *Yella sunde* (Kannada)

Aizoaceae

Sesuvium L.

Sesuvium portulacastrum (L.) L., Syst. Nat. (ed. 10) 1058. 1759; B.D.Sharma et al., Fl. Karnataka Anal. 117. 1984.

Portulaca portulacastrum L., Sp. Pl. 446. 1753

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada), Andhra Pradesh, Goa, Maharashtra, Odisha, Rajasthan, Tamil Nadu West Bengal; pantropical seashores.

Trianthema L.

Trianthema crystallina Vahl, Symb. Bot. 1: 32. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 117. 1984.

Distribution: Karnataka (Chikkamagaluru, Dharwad); NW India; E Africa, Middle East, Pakistan.

Trianthema portulacastrum L., Sp. Pl. 223. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.

Trianthema monogyna L., Mant. Pl. 1: 69. 1767

Trianthema obcordata Roxb., Fl. Ind. 1: 445. 1832

Distribution: Karnataka (Ballari, Chitradurga, Dharwad, Kalburgi, Mysuru, Vijayapura), throughout India; Pantropical weed

Local/Common/Vernacular name: *Balevadukana soppu* (Kannada)

Trianthema triquetra Rottler & Willd., Neue Schriften Ges. Naturf. Freunde Berlin 4: 181. 1803; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.

Distribution: Karnataka (Ballari, Dharwad, Mandya, Mysuru, Vijayapura), Andhra Pradesh, Goa, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu; Tropical region, especially in saline soil

Zaleya Burm. f.

Zaleya decandra (L.) Burm. f., Fl. Indica 110. 1768; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.

Trianthema decandra L., Mant. Pl. 1: 70. 1767

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Maharashtra, Rajasthan, Tamil Nadu; Australia, Malesia, Sri Lanka.

Local/Common/Vernacular name: *Kempuganaile* (Kannada)

Zaleya govindia (Buch.-Ham. ex G.Don) Nair, Bull. Bot. Surv. India 8: 86. 1966; C.J.Saldanha, Fl. Karnataka 1: 150. 1984

Trianthema govindia Buch.-Ham. ex G. Don, Gen. Syst. 3: 72. 1834

Trianthema pentandra auct., non L. Cooke, Fl. Bombay 1: 554. 1903

Distribution: Karnataka (Chitradurga, Vijayapura), Maharashtra, Rajasthan, Uttar Pradesh; Africa, Iran, Pakistan

Zaleya pentandra (L.) C.Jeffrey, Kew Bull. 14: 238 1960; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.

Distribution: Karnataka (Chikkamagaluru, Dharwad), NW India; Africa, Madagascar, Middle East, Pakistan.

Local/Common/Vernacular name: *African purslane* (English)

Alismataceae

Caldesia Parl.

Caldesia parnassifolia (L.) Parl., Fl. Ital. 3: 599.1860; B.D.Sharma et al., Fl. Karnataka Anal. 299. 1984.

Echinodorus parnassifolius (L.) Engelm., Fl. Brandenburg 1: 651.1864

Distribution: Karnataka (Bengaluru, Chamarajanagar, Davanagere, Hassan, Shivamogga); throughout plains; Africa, Australia, Asia, Europe.

Limnophyton Miq.

Limnophyton obtusifolium (L.) Miq., Fl. Ned. Ind. 3: 243.1856; B.D.Sharma et al., Fl. Karnataka Anal. 299. 1984.

Sagittaria obtusifolia L., Sp. Pl. 993.1753

Alisma obtusifolium (L.) Thwaites, Enum. Pl. Zeyl. 332.1864

Distribution: Karnataka (Belagavi, Bengaluru, Dharwad, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Shivamogga), Gujarat, Goa, Kerala, Maharashtra, Tamil Nadu; Africa, Madagascar, SE Asia, Sri Lanka.

Sagittaria L.

Sagittaria sagittifolia L., Sp. Pl. 993.1753; B.D.Sharma et al., Fl. Karnataka Anal. 299. 1984.

Distribution: Karnataka (Dharwad, Hassan), throughout plains; Australia, China, Europe, Russia, Siberia, The Caucasus Turkey, Ukraine, Vietnam.

Sagittaria trifolia L., Sp. Pl. 993.1753; B.D.Sharma et al., Fl. Karnataka Anal. 299. 1984.

Sagittaria sagittifolia var. *leucopetala* Miq., Ill. Fl. Archip. Ind. 2: 49.1870

Sagittaria sagittifolia subsp. *leucopetala* (Miq.) Hartog in Fl. Males. 5: 332.1957

Distribution: Karnataka (Dharwad, Hassan), Kerala, Maharashtra, Tamil Nadu; Asia, Europe, N America.

Wiesneria Micheli

Wiesneria triandra (Dalzell) Micheli, Monogr. Phan. 3: 82.1881

Sagittaria triandra Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 144. 1850

Distribution: Karnataka (Uttara Kannada), Kerala, Maharashtra; endemic.

Amaranthaceae

Achyranthes L.

Achyranthes aspera L., Sp. Pl. 204 1753; C.J.Saldanha, Fl. Karnataka 1: 161. 1984.

var. *aspera*

Plate 2A

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular name: *Uttarani* (Kannada)

var. **porphyristachya** (Wall. ex Moq.) Hook.f., Fl. Brit. India 4: 730 1885; B.D.Sharma et al., Fl. Karnataka Anal. 227. 1984.

Achyranthes porphyristachya Wall. ex Moq., Prodr. 13(2): 316 1849

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Shivamogga), Peninsular India; Sri Lanka.

var. **rubrofusca** (Wight) Hook.f., Fl. Brit. India 4: 730 1885; B.D.Sharma et al., Fl. Karnataka Anal. 227. 1984.

Achyranthes rubrofusca Wight, Icon. Pl. Ind. Orient. 5: t. 1778 1852

Achyranthes zaririi M.R.Almeida & S.M.Almeida, Fl. Maharashtra 4A: 196. 2003

Distribution: Karnataka (Chitradurga, Vijayapura), Peninsular India; China, Taiwan.

Achyranthes bidentata Blume, Bijdr. 545 1826; B.D.Sharma et al., Fl. Karnataka Anal. 227. 1984.

Distribution: Karnataka (Belagavi, Bidar, Chikkamagaluru, Hassan, Kalaburagi, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), throughout India; Paleotropical

Achyranthes coynei Santapau, Kew Bull. 1948: 488. 1949; Sandeep et al., J. Threat. Taxa. 3: 1875. 2011

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Aerva Forssk.

Aerva javanica (Burm.f.) Juss. ex Schult., Ann. Mus. Hist. Nat. 11: 131 1808; B.D.Sharma et al., Fl. Karnataka Anal. 227. 1984.

Iresine javanica Burm.f., Fl. Indica 212 1768

Iresine persica Burm.f., Fl. Indica 212.1768

Illecebrum javanicum (Burm.f.) L., Syst. Veg. ed. 13 206.1774

Aerva tomentosa Forssk., Fl. Aegypt.-Arab. 170 1775

Achyranthes javanica (Burm.f.) Pers., Syn. Pl. 1: 259.1805

Achyranthes incana Roxb., Fl. Ind. 1: 671.1820

Uretia persica (Burm.f.) Kuntze, Revis. Gen. Pl. 2: 544.1891

Aerva persica (Burm.f.) Merr. in Philipp. J. Sci. 19: 348. 1921

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dharwad, Kalaburagi, Mysuru, Raichur, Vijayapura), Andhra Pradesh, Maharashtra, Rajasthan, Uttar Pradesh; Paleotropical

Local/Common/Vernacular name: *Dodda bilihindi soppu* (Kannada)

Aerva lanata (L.) Juss., Ann. Mus. Hist. Nat. 11: 131 1808; B.D.Sharma et al., Fl. Karnataka Anal. 227. 1984. **Plate 2B**

Achyranthes lanata L., Sp. Pl. 204.1753

Celosia lanata L., Sp. Pl. 205.1753

Illecebrum lanatum (L.) L., Syst. Veg. ed. 13, 206.1774

Paronychia lanata Moench, Suppl. Meth. 112.1802

Aerva floribunda Wight, Icon. Pl. Ind. Orient. 5: t. 1776.1852

Uretia lanata Kuntze, Revis. Gen. Pl. 2: 544.1891

Distribution: Karnataka (throughout), throughout plains of India; paleotropical

Local/Common/Vernacular name: *Bilihindi soppu* (Kannada)

Aerva sanguinolenta (L.) Blume, Bijdr. 547 1826; B.D.Sharma et al., Fl. Karnataka Anal. 227. 1984.

Achyranthes sanguinolenta L., Sp. Pl. ed. 2,294.1762

Illecebrum sanguinolentum (L.) L., Mant. Pl. 2: 344.1771

Achyranthes scandens Roxb., Fl. Ind. 1: 676.1820

Aerva scandens (Roxb.) Wall., Icon. Pl. Orient. 2: t. 724.1840

Uretia sanguinolenta (L.) Kuntze, Revis. Gen. Pl. 2: 544.1891

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dharwad, Kalaburagi, Kolar, Uttara Kannada, Vijayapura, Yadgir), throughout India; Paleotropical

Allmania R. Br. ex Wight

Allmania nodiflora (L.) R.Br. ex Wight in J. Bot. (Hooker) 1: 226 1834; C.J.Saldanha, Fl. Karnataka 1: 164. 1984

Celosia nodiflora L., Sp. Pl. 205.1753

Celosia albida Willd., Sp. Pl. 1: 1197.1798

Celosia aspera Roth, Nov. Pl. Sp. 173.1821

Celosia dichotoma B.Heyne ex Roth, Nov. Pl. Sp. 172.1821

Achyranthes nodiflora (Willd.) Roxb., Fl. Ind. 2: 505.1824

Allmania aspera (Roth) Wight, Icon. Pl. Ind. Orient. 5: t. 1770.1852

Allmania dichotoma (B.Heyne ex Roth) Wight, Icon. Pl. Ind. Orient. 5: t. 1771.1852

Chamissoa aspera Wight, Icon. Pl. Ind. Orient. 5: t. 1772.1852

Chamissoa nodiflora sensu Wight, Icon. Pl. Ind. Orient. t. 1770 1852, non (L.) Mart. ex Moq. 1849

Allmania albida (Willd.) R.Br. ex Hook.f., Fl. Brit. India 4: 717.1885

Allmania nodiflora var. *angustifolia* Hook.f., Fl. Brit. India 4: 717.1885

Allmania nodiflora var. *aspera* (Roth) Hook.f., Fl. Brit. India 4: 717.1885

Allmania nodiflora var. *dichotoma* (B.Heyne ex Roth) Hook.f., Fl. Brit. India 4: 717.1885

Allmania nodiflora var. *esculenta* Hook.f., Fl. Brit. India 4: 717.1885

Allmania procumbens Wight ex Hook.f., Fl. Brit. India 4: 717.1885

Chamissoa angustifolia Buch.-Ham. ex Hook.f., Fl. Brit. India 4: 717 1885

Allmania nodiflora var. *procumbens* Ridl., Fl. Malay Penins. 3: 4.1924

Allmania longipedunculata (Trimen) Gamble, Fl. Madras 1168.1925

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalaburagi, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Vijayapura, Yadgir), Andhra Pradesh, Assam, Bihar, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, West Bengal; tropical Asia.

Local/Common/Vernacular name: *Budde soppu* (Kannada)

Alternanthera Forssk.

Alternanthera bettzickiana (Regel) G.Nicholson, Ill. Dict. Gard. 1: 59 1884

Telanthera bettzickiana Regel, Index Sem. (St. Petersburg) 1862: 28.1862

Alternanthera ficoidea var. *bettzickiana* (Regel) Backer in Fl. Males. 4: 93.1949

Alternanthera tenella var. *bettzickiana* (Regel) Veldkamp, Taxon 27: 313.1978

Alternanthera paronychioides var. *bettzickiana* (Regel) Fosberg, Willdenowia 20: 259. 1991

Distribution: Karnataka (throughout and colour varieties cultivated), throughout and colour varieties cultivated India; native of Brazil; pantropical.

Local/Common/Vernacular name: *Calico plant* (English)

Alternanthera ficoidea (L.) Sm., Cycl. 39(2 Suppl.): n. 8 1819

Gomphrena ficoidea L., Sp. Pl. 225. 1753; C.J.Saldanha, Fl. Karnataka 1: 165. 1984

Illecebrum ficoideum (L.) L., Sp. Pl. ed. 2 300.1762

Achyranthes ficoidea (L.) Lam., Encycl. 1: 548.1785

Paronychia ficoidea (L.) Desf., Tabl. École Bot. 44.1804

Alternanthera tenella Colla, Mem. Reale Accad. Sci. Torino 33: 131.1829

Steiremis ficoidea (L.) Raf., Fl. Tellur. 3: 41.1837

Telanthera ficoidea (L.) Moq., Prodr. 13(2): 363 1849; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout except NE India; pantropical.

Alternanthera littoralis P.Beauv., Fl. Oware 2: 72 1818

Alternanthera ficoidea Griseb., Fl. Brit. W. I. 67. 1859.

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Mysuru); pantropical.

Alternanthera paronychioides A.St.-Hil., Voy. Bres. 2: 439. 1833; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Hassan, Mysuru, Tumakuru), throughout India; native to tropical America; pantropical.

Alternanthera philoxeroides (Mart.) Griseb., Wiss. Gnett. 24: 36. 1879; C.J.Saldanha, Fl. Karnataka 1: 165. 1984

Plate 2C

Bucholzia philoxeroides Mart., Beitr. Amarantac. 107 1825

Distribution: Karnataka (throughout), throughout India; native to tropical S America, pantropical.

Local/Common/Vernacular name: *Aligator weed* (English)

Alternanthera pungens Kunth in HBK Nov. Gen. Sp. 2: 206. 1817; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Plate 2D

Achyranthes repens L., Sp. Pl. 205.1753

Illecebrum achyrantha L., Sp. Pl. ed. 2 299.1762

Alternanthera repens (L.) Link, Enum. Hort. Berol. Alt. 1: 154.1821

Illecebrum pungens (Kunth) Spreng., Syst. Veg. 1: 820.1824

Telanthera pungens (Kunth) Moq., Prodr. 13(2): 371.1849

Distribution: Karnataka (throughout), weed throughout tropical parts of India; native to tropical America; pantropical.

Local/Common/Vernacular names: *Creeing chaff weed*, *Khaki weed* (English); *Mirza mullu*, *Mullu honagone* (Kannada)

Alternanthera sessilis (L.) R.Br. ex DC., Cat. Hort. 4: 77. 1813; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Plate 3A

Gomphrena polygonoides L., Sp. Pl. 225.1753

Gomphrena sessilis L., Sp. Pl. 225.1753

Illecebrum sessile (L.) L., Sp. Pl. ed. 2 300.1762

Paronychia sessilis (L.) Desf., Tabl. École Bot. 44 1804

Achyranthes sessilis (L.) Besser, Cat. Jard. Bot. Krzemieniec 12.1810

Alternanthera ficoides P.Beauv., Fl. Oware 2: 66.1818

Achyranthes triandra Roxb., Fl. Ind. 1: 678.1820

Distribution: Karnataka (throughout), throughout India; pantropical.

Local/Common/Vernacular name: *Sessile joy weed* (English); *Honagone soppu* (Kannada)

Amaranthus L.

Amaranthus × catus B.Heyne ex Hook.f., Fl. Brit. India 4: 720.1885; V.Thakur, Kew Bull. 21: 63.1967.

Distribution: Karnataka (), Andhra Pradesh, Tamil Nadu; endemic.

Amaranthus blitum L., Sp. Pl. 990. 1753.

subsp. **oleraceus** (L.) Costea, Sida 19: 984.2001

Amaranthus lividus L., Sp. Pl. 990.1753

Amaranthus oleraceus L., Sp. Pl. 1403.1753

Blitum lividum (L.) Moench, Methodus 359.1794

Blitum oleraceum (L.) Moench, Methodus 359.1794

Glomeraria livida (L.) Cav., Elench. Pl. Horti Matr. 16.1803

Glomeraria oleracea (L.) Cav. in Elench. Pl. Horti Matr. 16.1803

Albersia oleracea (L.) Kunth, Fl. Berol. ed. 2, 2: 144.1823

Euxolus lividus Moq., Prodr. 13(2): 273 1849

Euxolus oleraceus Moq., Prodr. 13(2): 273 1849

Euxolus viridis (L.) Moq., Prodr. 13(2): 273 1849

Amaranthus blitum var. *oleraceus* (L.) Hook.f., Fl. Brit. India 4: 721.1885

Amaranthus lividus var. *oleraceus* (L.) Thell., Fl. Adv. Montpellier 215.1912

Amaranthus ascendens var. *oleraceus* (L.) Thell. ex Priszter, Agrártud. Egyet. Kert- Szölögazdaságtud. Karának Évk. 2: 221.1953

Amaranthus lividus subsp. *oleraceus* (L.) Soó, Acta Bot. Acad. Sci. Hung. 10: 376.1964

Distribution: Karnataka (cultivated throughout), throughout India ; Africa, Europe, Middle East, E & S Asia; native to N. America, pantropical.

Local/Common/Vernacular name: *Chilikaesoppu* (Kannada)

Amaranthus caudatus L., Sp. Pl. 990. 1753; C.J.Saldanha, Fl. Karnataka 1: 168. 1984; Seetharam, Kotress & Uploankar, Fl. Gulbarga Dist. 19. 2000.

Distribution: Karnataka (cultivated in gardens throughout and as crop in Kalburgi, Yadgir), Andhra Pradesh, Assam, Bihar, Goa, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh; cultivated particularly in Uttarakhand, Sikkim, West Bengal (Darjeeling); native to Americas

Local/Common/Vernacular names: *Foxtail amaranth*, *Love-lies-bleeding*, *Pendant amaranth*, *Quilette*, *Tasselflower*, *Velvetflower* (English)

Amaranthus cruentus L., Syst. Nat. ed. 10 2: 1269 1759

Amaranthus paniculatus L., Sp. Pl. ed. 2 1406 1763

Amaranthus hybridus subsp. *cruentus* (L.) Thell. in Fl. Adv. Montpellier 205. 1912

Amaranthus hybridus var. *paniculatus* (L.) Thell. in Asch. & Gr. Syn. 5.1: 247.1914

Distribution: Karnataka (cultivated throughout), throughout India, an escape near gardens, roadsides etc.; pantropical.

Local/Common/Vernacular names: *Kirel soppu*, *rajakere* (Kannada)

Amaranthus dubius Mart. ex Thell., Fl. Adv. Montpellier 203.1912

Amaranthus dubius var. *flexuosus* Thell., Syn. Mitteleur. Fl. 5: 266.1914

Amaranthus dubius var. *leptostachys* Thell., Syn. Mitteleur. Fl. 5: 266.1914

Amaranthus dubius var. *xanthostachys* Thell., Syn. Mitteleur. Fl. 5: 266.1914

Distribution: Karnataka (cultivated in throughout; also grows as an escape), cultivated throughout; native to tropical America

Local/Common/Vernacular names: *Chinese spinach*, *Red spinach*, *Spleen Amaranth* (English)

Amaranthus graecizans L., Sp. Pl. 990.1753; C.J.Saldanha, Fl. Karnataka 1: 166. 1984.

Amaranthus angustifolius Lam., Encycl. 1: 115.1783

Blitum graecizans (L.) Moench, Methodus 359.1794

Glomeraria graecizans (L.) Cav., Elench. Pl. Horti Matr. 16.1803

Amaranthus blitum var. *graecizans* (L.) Moq. in DC. Prodr. 13(2): 263. 1849

Amaranthus polygamus var. *angustifolia* (Lam.) Hook. f., Fl. Brit. India 4: 721. 1885

Galliaria graecizans (L.) Nieuwl., Amer. Midl. Naturalist 3: 278.1914

Amaranthus aschersonianus (Thell.) Chiov., Nuovo Giorn. Bot. Ital. n.s., 26: 103.1919

Amaranthus thellungianus Nevski, Act. Acad. Sc. U.R.S.S. 1(4): 311. 1937; C.J.Saldanha, Fl. Karnataka 1: 167. 1984.

Amaranthus graecizans subsp. *thellungianus* (Nevski) Gusev, Bot. Zhurn. 57:462.1972

Amaranthus roxburgianus var. *aschersonianus* (Thell.) N.C.Nair, J. Bombay Nat. Hist. Soc. 73: 61.1976

Distribution: Karnataka (throughout), throughout India; Africa, Middle East, West & SE Asia.

Amaranthus hybridus L., Sp. Pl. 990.1753; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Amaranthus chlorostachys Willd., Hist. Amaranth. 34.1790

Amaranthus frumentaceus Buch.-Ham. in Roxb., Fl. Ind. ed. 2.3: 609. 1832

Amaranthus retroflexus var. *hybridus* (L.) A.Gray, Manual ed. 5: 412.1867

Amaranthus chlorostachys var. *hybridus* (L.) S.Watson, Manual ed. 6: 428.1890

Amaranthus hybridus var. *chlorostachys* (Willd.) Beck, Icon. Fl. Germ. Helv. 24: 175.1908

Amaranthus hybridus subsp. *hypochondriacus* var. *chlorostachys* (Willd.) Thell., Fl. Adv. Montpellier 205.1912

Gallaria hybrida (L.) Nieuwl., Amer. Midl. Naturalist 3: 278.1914

Distribution: Karnataka (Belagavi, Bengaluru, Chikmagluru, , Mysuru, Yadgir), cultivated throughout India; native to N America.

Amaranthus hypochondriacus L., Sp. Pl. 991.1753; C.J.Saldanha, Fl. Karnataka 1: 168. 1984.

Amaranthus flavus L., Syst. Nat. ed. 10 2: 1269.1759

Amaranthus hybridus Vell., Fl. Flumin. 10: t. 28. 1831.

Amaranthus anardana Buch.-Ham. ex Moq., Prodr. 13(2): 256.1849

Amaranthus hypochondriacus var. *macrostachys* Moq., Prodr. 13(2): 256.1849

Amaranthus hypochondriacus var. *monstrosus* Moq., Prodr. 13(2): 256.1849

Amaranthus hypochondriacus var. *racemosus* Moq., Prodr. 13(2): 256.1849

Amaranthus hypochondriacus var. *tortuosus* Moq., Prodr. 13(2): 256.1849

Amaranthus macrostachyus Mérat ex Moq., Prodr. 13(2): 256.1849

Amaranthus monstrosus Moq., Prodr. 13(2): 256.1849

Amaranthus hybridus var. *hypochondriacus* (L.) H.Rob., Rhodora 10: 32.1908

Amaranthus hybridus subsp. *hypochondriacus* (L.) Thell., Fl. Adv. Montpellier 204.1912

Amaranthus polygonoides L., Pl. Jamaic. Pug. 27. 1759.

Distribution: Karnataka (Ballari, Bidar, Chikkamagaluru, Chitradurga, Kalaburagi, Raichur, Vijayapura), Maharashtra, Rajasthan; cultivated

Amaranthus spinosus L., Sp. Pl. 991. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Distribution: Karnataka (Throughout), throughout India; Pantropical.

Local/Common/Vernacular name: *Mulludantu* (Kannada)

Amaranthus tricolor L., Sp. Pl. ed. 1: 989. 1753; Manikand. & P. L. Narasim., Fl. Rajiv Gandhi Nat. Park 329. 2013.

Amaranthus melancholicus L., Sp. Pl. ed 1: 989. 1753

Amaranthus tristis L., Sp. Pl. ed. 1: 989. 1753

Amaranthus mangostanus L., Cent. Pl. I 32 1755

Amaranthus gangeticus L., Syst. Veg. ed. 10, 2: 1268.1759; Hook. f., Fl. Brit. India 4: 719. 1885.

Amaranthus lancifolius Roxb., Hort. Bengal. 67 1814

Amaranthus lanceolatus Roxb., Fl. Ind. ed. 1832 3: 607 1832

Amaranthus lividus Roxb., Fl. Ind. ed. 1832 3: 605. 1832.

Amaranthus oleraceus Roxb., Fl. Ind. ed. 1832 3: 605 1832.

Amaranthus polygamus Roxb., Fl. Ind. ed. 1832 3: 607. 1832.

Amaranthus tristis Wall., Numer. List 6905. 1832.

Amaranthus melancholicus var. *tricolor* (L.) Lam. ex Moq., Prodr. 13(2): 262 1849

Amaranthus polygamus Thwaites, Enum. Pl. Zeyl. 247. 1861.

Amaranthus tricolor var. *gangeticus* (L.) Fiori, Fl. Italia 1: 322.1898

Amaranthus tricolor var. *tristis* (L.) Thell. 1077 5 (1): 267. 1914

Amaranthus tricolor var. *mangostanus* (L.) Aellen, Ill. Fl. Mitt.-Eur. ed. 2, 3(2): 495.1959

Amaranthus tricolor var. *tristis* (Willd.) Mehrotra, Aswal & Bisht, Compan. Chopra's Gloss. Indian Medicin. Pl. 9 1987

Distribution: Karnataka (cultivated, occasionally run wild), throughout India; introduced in Africa, tropical America; considered native t India.

Local/Common/Vernacular names: *Chilkarive soppu*, *Dantina soppu*, *Dantu soppu*, *Hariye soppu* (Kannada)

Amaranthus viridis L., Sp. Pl. ed. 2: 1405.1763

Glomeraria viridis (L.) Cav., Descr. Pl. 319.1802

Amaranthus gracilis Desf., Tabl. Ecol. Bot. 43. 1804; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Amaranthus fasciatus Roxb., Fl. Ind. 3: 509.1832

Euxolus caudatus (Jacq.) Moq. in DC., Prodr. 13.2: 274.1849

Pyxidium viride (L.) Moq., Prodr. 13(2): 274.1849

Distribution: Karnataka (throughout), throughout India; native to Americas.

Local/Common/Vernacular names: *Green amaranth, Slender amaranth* (English); *chilateere, Cilakarive soppu, Dagli soppu* (Kannada)

Atriplex L.

Atriplex hortensis L., Sp. Pl. 1053. 1753

Distribution: Karnataka (Kodagu), cultivated elsewhere in India; native to Europe & Middle East

Local/Common/Vernacular name: *Garden orache* (English)

Beta L.

Beta vulgaris L., Sp. Pl. 222 1753; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Distribution: Karnataka (Chitradurga, Davanagere, Hassan, Kalaburagi), cultivated throughout India; native to Europe, cultivated as vegetable.

Local/Common/Vernacular name: *Beet root* (English)

Celosia L.

Celosia argentea L., Sp. Pl. 205.1753; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Celosia cristata L., Sp. Pl. 205.1753

Celosia castrensis L., Sp. Pl. ed. 2 297.1762

Celosia coccinea L., Sp. Pl. ed. 2 297.1762

Celosia margaritacea L., Sp. Pl. ed. 2 297. 1762

Celosia pyramidalis Burm.f., Fl. Indica 62. 1768

Celosia cernua Roxb., Fl. Ind. 1: 680.1820

Chamissoa margaritacea (L.) Schouw, Fortegn. Bot. Hav. Pl. 123.1847

Celosia argentea var. *cristata* (L.) Kuntze, Revis. Gen. Pl. 1: 541.1891

Distribution: Karnataka (throughout, ornamental varieties cultivated); throughout India, ornamental varieties cultivated; pantropical.

Local/Common/Vernacular names: *Silver cock's comb* (English); *Anne soppu, Hannehu, Kolani, Kollikondai* (Kannada)

Celosia polygonoides Retz., Observ. Bot. 2: 12.1781; B.D.Sharma et al., Fl. Karnataka Anal. 228. 1984.

Lophoxera polygonoides (Retz.) Raf., Fl. Tellur. 3: 42.1837

Distribution: Karnataka (Ballari, Chikmagluru, Mysuru, Tumakuru, Vijayapura), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Celosia pulchella Moq., Prodr. 13(2): 238.1849; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Distribution: Karnataka (Chikmagluru, Davanagere, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu, West Bengal; Sri Lanka.

Chenopodium L.

Chenopodium album L., Sp. Pl. 219.1753; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Atriplex alba (L.) Crantz, Inst. Rei Herb. 1: 206.1766

Botrys alba (L.) Nieuwl., Amer. Midl. Naturalist 3: 276.1914

Distribution: Karnataka (cultivated in Bengaluru, Chikkaballapur, Hassan, Kodagu, Kolar, Mysuru, Shimoga, Tumakuru, Yadgir), S and SE Asia; native to Europe.

Local/Common/Vernacular name: *Hunachikka* (Kannada)

Chenopodium murale L., Sp. Pl. 219.1753; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Atriplex muralis (L.) Crantz, Inst. Rei Herb. 1: 206.1766

Anserina muralis (L.) Montandon, Syn. Fl. Jura 263.1856

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Kodagu, Mysuru), Andhra Pradesh, Bihar, Delhi, Himachal Pradesh, Jammu & Kashmir, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh; cosmopolitan weed

Cyathula Blume

Cyathula prostrata (L.) Blume, Bijdr. 549.1826; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Achyranthes prostrata L., Sp. Pl. ed. 2 296.1762

Desmochaeta prostrata (L.) DC., Cat. Pl. Horti Monsp. 102.1813

Achyranthes repens B.Heyne ex Roth, Nov. Pl. Sp. 167. 1821.

Pupalia prostrata (L.) Mart., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 13: 321.1826

Distribution: Karnataka (Bengaluru, Chikmagluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shimoga, Uttara Kannada, Vijayapura), Arunachal Pradesh, Assam, Daman & Diu, Kerala, Maharashtra, Meghalaya, Nagaland, Sikkim, Tamil Nadu, Tripura, West Bengal; native to C America; Pantropical.

Local/Common/Vernacular names: *Kaadu dantu, Kempu uttarani* (Kannada)

Digera Forssk.

Digera muricata (L.) Mart., Beitr. Amarantac. 77. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Achyranthes muricata L., Sp. Pl. (ed. 2) 295. 1762

Achyranthes alternifolia L., Mant. Pl. 1: 50. 1767

Digera arvensis Forssk., Fl. Aegypt.-Arab. 65.1775

Digera alternifolia (L.) Asch., Beitr. Fl. Aethiop. 180. 1867

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Kalaburagi, Mysuru, Raichur, Vijayapura), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Goa, Gujarat, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh; tropical Africa, Madagascar, Middle East, S & SE Asia, West Asia.

Local/Common/Vernacular names: *False amaranth* (English); *Cenakali soppu, Gorje pallye* (Kannada)

Dysphania R. Br.

Dysphania ambrosioides (L.) Mosyakin & Clemants, Ukrayins'k. Bot. Zhurn. 59: 382 2002

Chenopodium ambrosioides L., Sp. Pl. 219.1753; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Atriplex ambrosioides (L.) Crantz, Inst. Rei Herb. 1: 207.1766

Orthosporum ambrosioides (L.) Kostel., Allg. Med. -Pharm. Fl. 4: 1433.1835

Vulvaria ambrosioides (L.) Bubani, Fl. Pyren. 1: 178.1897

Chenopodium album subsp. *ambrosioides* (L.) H.J.Coste & A.Reyn., Bull. Herb. Boissier II, 5: 979.1905

Blitum ambrosioides (L.) Beck, Icon. Fl. Germ. Helv. 24: 118.1908

Botrys ambrosioides (L.) Nieuwl., Amer. Midl. Naturalist 3: 275.1914

Teloxys ambrosioides (L.) W.A. Weber, Phytologia 58(7): 477.1985

Distribution: Karnataka (Belagavi, Bengaluru, Chikmagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru), native to N America; Pantropical.

Local/Common/Vernacular names: *Huli oma, Kaddu oma* (Kannada)

Gomphrena L.

Gomphrena celosioides Mart., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 13: 301.1826; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Distribution: Karnataka (Bengaluru, Chikmagluru, Chitradurga, Davanagere, Hassan, Kodagu, Mysuru, Raichur, Tumakuru, Vijayapura, Yadgir); native to C America; pantropical.

Local/Common/Vernacular names: *Gonde, Javalada malle, Neer bogada banta, Nela rudrakshi hu* (Kannada)

Gomphrena globosa L., Sp. Pl. 224.1753; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Xeraea globosa (L.) Kuntze, Revis. Gen. Pl. 2: 543.1891

Distribution: Karnataka (cultivated as an ornamental throughout), cultivated throughout India; native to tropical America.

Local/Common/Vernacular names: *Bachelor button, Globe amaranth* (English); *Adike huvu, Bili rudrakshi huvu, Kashi gonde, Kempu gonde, Nela rudraksha hoovu* (Kannada); *Bullamu Gamdo* (Konkani)

Gomphrena serrata L., Sp. Pl. 224.1753; Ramaswamy et al., Fl. Shimoga 489. 2001

Distribution: Karnataka (Chitradurga, Dharwad, Mysuru, Shivamogga), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Introduced, native to tropical America.

Local/Common/Vernacular names: *Jalavada malle, Nela rudrakshi* (Kannada)

Halosarcia Paul G. Wilson

Halosarcia indica (Willd.) Paul G. Wilson, Nuytsia 3: 63 1980

Salicornia indica Willd. in Neue Schriften Ges. Naturf. Freunde Berlin 2: 111 1799

Salicornia brachiata Roxb., Hort. Bengal. 79. 1814; C.J.Saldanha, Fl. Karnataka 1: 159. 1984

Arthrocnemum indicum (Willd.) Moq., Enum. 113 1840; C.J.Saldanha, Fl. Karnataka 1: 159. 1984

Distribution: Karnataka (Coastal areas of Dakshina Kannad, Udupi, Uttara Kannada), Coastal states of India; coastal areas Africa, Australia, Bangla desh, Sri Lanka.

Indobanalia A.N.Henry & B. Roy

Indobanalia thyrsiflora (Moq.) Henry & Roy, Bull. Bot. Surv. India 10: 274. 1968; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Banalia thyrsiflora Moq. in DC., Prodr. 13: 278. 1849

Distribution: Karnataka (Chikkaballapur, Dakshina Kannada, Hassan, Kalaburagi, Kodagu, Kolar, Mysuru, Yadgir), Kerala, Tamil Nadu; endemic.

Iresine P. Br.

Iresine herbstii Hook., Gard. Chron. 1864: 654. 1864

Iresine verschaffeltii Lem., Ill. Hort. 11: t. 418. t. 418. 1864

Distribution: Karnataka (cultivated as ornamental throughout), cultivated as ornamental throughout drier parts of India; native to Brazil.

Local/Common/Vernacular names: *Chicken Gizzard, Herbst's bloodleaf* (English)

Nothosaerva Wight

Nothosaerva brachiata (L.) Wight, Icon. Pl. Ind. Orient. 6: 1.1853; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Achyranthes brachiata L., Mant. Pl. 50.1767

Illecebrum brachiatum (L.) L., Mant. Pl. 2: 213.1771

Aerva brachiata (L.) Mart., Beitr. Amarantac. 83.1825

Pseudanthus brachiatus (L.) Wight, Icon. Pl. Ind. Orient. 5: 3.1852

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Dharwad, Kodagu, Mysuru), Andhra Pradesh, Dadra-Nagar Haveli, Daman & Diu, Goa, Gujarat, Haryana, Himachal Pradesh, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Africa, S & SE Asia.

Psilotrichum Blume

Psilotrichum nudum Moq., Prodr. 13(2): 280.1849; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Achyranthes nuda B.Heyne ex Wall., Fl. Ind. 2: 507.1824

Distribution: Karnataka (Belagavi), Tamil Nadu; endemic.

Pupalia Adans.

Pupalia lappacea (L.) Juss., Ann. Mus. Natl. Hist. Nat. 2: 132.1803; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Achyranthes lappacea L., Sp. Pl. 204.1753

Achyranthes atropurpurea Lam., Encycl. 1: 546.1785

Cadelari lappacea (L.) Medik., Malvenfam. 92.1787

Codivalia lappacea (L.) Raf., Fl. Tellur. 3: 40.1837

Pupalia atropurpurea (Lam.) Moq., Prodr. 13(2): 331.1849

Desmochaeta lappacea (L.) Griff., Not. Pl. Asiat. 4: 339.1854

var. **lappacea**

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Kalaburagi, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Uttara Kannada, Vijayapura, Yadgir), throughout India; Africa, S & SE Asia.

Local/Common/Vernacular names: *Antupurale gida, Haridacchaga* (Kannada)

var. **orbiculata** (B.Heyne ex Wall.) C.C. Towns., Kew Bull. 29:469. 1974

Achyranthes orbiculata Heyne ex Wall. in Roxb., Fl. Ind. 2: 507. 1824

Cyathula orbiculata Moq., Prodr. 13(2): 330.1849

Pupalia orbiculata Wight, Icon. Pl. Ind. Orient. 5: t. 1783.1852; B.D.Sharma et al., Fl. Karnataka Anal. 229. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Mysuru), Andhra Pradesh, Gujarat, Maharashtra, Rajasthan, Tamil Nadu; Sri Lanka.

Spinacia L.

Spinacia oleracea L., Sp. Pl. 1027.1753; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Distribution: Karnataka (cultivated in most districts), cultivated throughout India; native to C & W Asia; Pantropical.

Local/Common/Vernacular name: *Spinach* (English)

Trichuriella Bennet

Trichuriella monsoniae (L.f.) Bennet, Indian J. Forest. 8: 86.1985; B.D.Sharma et al., Fl. Karnataka Anal. 230. 1984.

Celosia monsoniae Retz., Observ. Bot. 2: 13 1781 'monsonia'

Illecebrum monsoniae L. f., Suppl. Pl. 161. 1781; C.J.Saldanha, Fl. Karnataka 1: 173. 1984.

Aerva monsoniae (L.f.) Mart., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 13(1): 291.1825

Trichurus monsoniae (L.f.) C.C. Towns., Kew Bull. 29(3): 466.1974

Distribution: Karnataka (Ballari, Benlgauru, Chitradurga, Hassan, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Bihar, Madhya Pradesh, Odisha, Tamil Nadu, West Bengal; China, Myanmar, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular name: *Majjigebern gida* (Kannada)

Amaryllidaceae

Crinum L.

Crinum × amabile Donn ex Ker Gawl., Bot. Mag. 39:t. 1605.1814

Crinum × augustum Roxb., J. Sci. Arts (London) 3: 113.1817

Crinum × amabile var. *augustum* (Roxb.) Ker Gawl., Bot. Reg. 8: 679. 1822

Crinum × superbum Roxb., Fl. Ind. 2: 133.1824

Distribution: Karnataka (cultivated in Bengaluru), cultivated in India; native to S Africa.

Local/Common/Vernacular names: *Giant/Grand Red/Pink Spider Lily, Pink Giant Spider Lily* (English)

Crinum amoenum Ker Gawl. ex Roxb., J. Sci. Arts (London) 3: 106.1817; Manikand. & Lakshminars., Fl. Rajiv Gandhi Nation. Park 398.2013.

Crinum himalense Royle, Ill. Bot. Himal. Mts. 374.1839

Distribution: Karnataka (Mysuru), Maharashtra; Bangladesh, Myanmar, Nepal.

Crinum asiaticum L., Sp. Pl. 292.1753; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Bulbine asiatica (L.) Gaertn., Fruct. Sem. Pl. 1: 42.1788

Distribution: Karnataka (Belagavi, Bengaluru, Chikkaballapur, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kalburgi, Kolar, Ramanagara, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout tropical parts of India; Australia, China, Indo-China, Mauritius, Pacific Islands, S and SE Asia.

Local/Common/Vernacular names: *Giant Crinum Lily, Grand Crinum Lily, Poison Bulb, Spider Lily* (English); *Kaadu irulli, Nagdali, Visha munguli* (Kannada)

Crinum latifolium L., Sp. Pl. 291.1753; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Crinum moluccanum Roxb. ex Ker Gawl., J. Sci. Arts (London) 3: 109.1817

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Dharwad, Kolar, Mysuru, Udupi, Uttara Kannada), China, Laos, Myanmar, Sri Lanka, Thailand, Vietnam.

Crinum lorifolium Roxb. ex Ker Gawl., J. Sci. Arts (London) 3: 110.1817

Crinum longifolium Roxb. ex Ker Gawl., J. Sci. Arts (London) 3: 107.1817

Crinum elegans Carey ex Herb., Bot. Mag. 47:t. 2121.1820

Crinum pratense Herb., Amaryllidaceae 256.1837; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Distribution: Karnataka (Belagavi), Goa, Gujarat, Maharashtra; Myanmar.

Crinum moorei Hook.f., Bot. Mag. 100:t. 6113. 1874

Distribution: Karnataka (cultivated in Dakshina Kannada, Udupi), cultivated in India; native to S Africa.

Local/Common/Vernacular names: *Moore's crinum, Natal Lily* (English)

Crinum viviparum (Lam.) R.Ansari & V.J.Nair, J. Econ. Taxon. Bot. 11: 205.1988

Amaryllis vivipara Lam., Encycl. 1: 123.1783

Crinum defixum Ker Gawl., J. Sci. Arts (London) 3: 105.1817; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Crinum ensifolium Roxb. ex Ker Gawl., J. Sci. Arts (London) 3: 106.1817

Crinum roxburghii Dalzell & Gibson, Bombay Fl. 275.1861

Crinum defixum var. *ensifolium* (Roxb. ex Ker Gawl.) Baker, Handb. Amaryll. 76.1888

Crinum viviparum var. *ensifolium* (Roxb. ex Ker Gawl.) R.Ansari & V.J.Nair, J. Econ. Taxon. Bot. 11: 205.1988

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Bangladesh, Bhutan, Indonesia, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *River crinum lily* (English); *Gajara, Gajina gida, Visha bidiru* (Kannada)

Haemanthus L.

Haemanthus coccineus L., Sp. Pl. 1: 325. 1753

Distribution: Karnataka (cultivated throughout), cultivated throughout in India; native to S Africa.
 Local/Common/Vernacular names: *Blood lily, Paintbrush lily, The bloodflower* (English)

Hippeastrum Herb.

Hippeastrum puniceum (Lam.) Voss, Vilm. Blumengärtn. ed. 3 1: 1033.1895

Amaryllis punicea Lam., Encycl. 1: 122.1783

Distribution: Karnataka (Dakshina Kannada, Udupi), cultivated in India; native to S America.
 Local/Common/Vernacular name: *Barbados Lily* (English)

Hippeastrum reginae (L.) Her., Appendix 31.1821

Amaryllis reginae L., Syst. Nat., ed. 10. 2: 977. 1759

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental in India; native to S America.

Hippeastrum vittatum (L'Hér.) Herb., Appendix 31.1821

Amaryllis vittata L'Hér., Sert. Angl. 13.1789

Distribution: Karnataka (Belagavi, Bengaluru), cultivated in India; native to Brazil.

Hymenocallis Salisb.

Hymenocallis littoralis (Jacq.) Salisb., Trans. Hort. Soc. London 1: 338.1812

Pancratium littorale Jacq., Select. Stirp. Amer. Hist. 99.1763

Hymenocallis tenuiflora Herb., Appendix 44. 1821

Troxistemon littorale (Jacq.) Raf., Fl. Tellur. 4: 23.1838

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), native S & C America.

Local/Common/Vernacular names: *Spider Lily, The beach spider lily* (English)

Pancratium L.

Pancratium parvum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 144.1850; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Pancratium parvum var. *malabaricum* Baker, Amaryllidaceae 119.1888

Pancratium parvum subsp. *malabaricum* (Baker) Traub, Pl. Life 23: 65.1967

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Dharwad, Uttara Kannada), Maharashtra; endemic.

Pancratium triflorum Roxb., Fl. Ind. ed. 1832 2: 126.1832; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Pancratium malabathricum Herb., Amaryllidaceae 202.1837

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu, West Bengal; endemic.

Pancratium zeylanicum L., Sp. Pl. 290.1753; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Distribution: Karnataka (Bengaluru), Kerala, Lakshadweep Islands, Tamil Nadu; Indonesia, Maldives Islands, Sri Lanka, Philippines.

Scadoxus Raf.

Scadoxus multiflorus (Martyn) Raf., Fl. Tellur. 4: 19.1838

Haemanthus multiflorus Martyn, Monogr. s.p., t. 1795

Amaryllis multiflora (Martyn) Tratt., Ausw. Merkw. Gartenpfl. 37.1821

Nerissa multiflorus (Martyn) Salisb., Gen. Pl. 131.1866

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical Africa.

Local/Common/Vernacular names: *Fire ball lily, Football lily, Mayflower* (English)

Zephyranthes Herb.

Zephyranthes carinata Herb., Bot. Mag. 52: t. 2594.1825

Amaryllis carinata (Herb.) Spreng., Syst. Veg. 4(2): 132.1827

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Dharwad, Mandya, Mysuru, Ramanagara, Tumakuru, Udupi), native to C America; cultivated throughout tropics.

Local/Common/Vernacular name: *Zephyr lily* (English)

Anacardiaceae

Anacardium L.

Anacardium occidentale L., Sp. Pl. 583. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 58. 1984. **Plate 3B**

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; native to America

Local/Common/Vernacular name: *Cashew nut* (English); *Geru* (Kannada)

Buchanania Spreng.

Buchanania axillaris (Desr.) Ramamoorthy in Saldanha & Nicolson, Fl. Hassan Dist. 374. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 58. 1984.

Mangifera axillaris Desr. in Lam., Encycl. Meth. Bot. 3: 687. 1789

Buchanania angustifolia Roxb., Pl. Coromandel 3: 79. t. 262. 1819

Buchanania intermedia Wight, Icon. Pl. Ind. Orient. 1: t. 81. 1838

Distribution: Karnataka (Ballari, Chikkaballapur, Chitradurga, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Tumakur), Andhra Pradesh, Assam, Bihar, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; Sri Lanka

Local/Common/Vernacular name: *Cuddaph Almond* (English); *Yemme murukalgida* (Kannada)

Buchanania cochinchinensis (Lour.) M.R.Almeida, Fl. Maharashtra 1: 287. 1996

Buchanania lanzan Spreng. in J. Bot. (Schr.) 2: 234. 1800; B.D.Sharma et al., Fl. Karnataka Anal. 58. 1984.

Buchanania latifolia Roxb., Fl. Ind. 2: 385. 1832

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkaballapur, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Maharashtra; China, Laos, Malaysia, Myanmar, Thailand, Vietnam.

Local/Common/Vernacular names: *Chironi, Murkali* (Kannada)

Buchanania lanceolata Wight, Icon. Pl. Ind. Orient. 1: t. 237 1839; B.D.Sharma et al., Fl. Karnataka Anal. 58. 1984.

Distribution: Karnataka (Chikkamagaluru, Kodagu, Shivamogga), Kerala, Tamil Nadu; Myanmar.

Holigarna Buch.-Ham. ex Roxb.

Holigarna arnottiana Hook. f., Fl. Brit. India 2: 36. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 59.

1984. **Plate 14F**

Holigarna longifolia Wight & Arn., Prodr. Fl. Ind. Orient. 1: 169. 1834, non Roxb.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Jungle Marking Nut* (English); *Sanna holegara* (Kannada)

Holigarna beddomei Hook.f., Fl. Brit. India 2: 36 1876; Kesh.Murthy & Yogan, Fl. Kodagu 123. 1990

Distribution: Karnataka (Kodagu, Uttara kannada), Kerala, Tamil Nadu; endemic.

Holigarna ferruginea Marchand, Rev. Anacard. 171. 1868; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.
Distribution: Karnataka (Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Tamil Nadu; endemic.

Holigarna grahamii (Wight) Kurz, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 42: 305. 1872; B.D.Sharma et al.,
Fl. Karnataka Anal. 59. 1984. **Plate 14G**

Semecarpus grahamii Wight, Icon. Pl. Ind. Orient. 1: t. 235. 1839

Holigarna wightii Balakr. in J. Bombay Nat. Hist. Soc. 63: 327. 1966

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Holigarna nigra Bourd., Indian J. For. 30: 95. 1904; C.J.Saldanha, Fl. Karnataka 2: 205. 1996

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Cherugeru* (Kannada)

Lannea A.Rich.

Lannea coromandelica (Houtt.) Merr., J. Arnold Arbor. 19: 353. 1939; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Dialium coromandelicum Houtt., Nat. Hist. ser. 2. 2: 39. t. 5. f. 2. 1774

Odina wodier Roxb., Fl. Ind., ed. 1832, 2: 293. 1832

Distribution: Karnataka (Ballari, Belagavi, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada), throughout India; Sri Lanka

Local/Common/Vernacular names: *Ajashringi, Udimara* (Kannada)

Mangifera L.

Mangifera indica L., Sp. Pl. 200. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout tropical regions of India; wild in E & W Ghats, NE India; Indomalaya.

Local/Common/Vernacular name: *Mango* (English); *Mavina mara* (Kannada)

Nothopegia Blume

Nothopegia beddomei Gamble, Bull. Misc. Inform. Kew 1918: 227. 1918; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Oolagera, Vyana* (Kannada)

Nothopegia colebrookiana (Wight) Blume in Mus. Bot. Lugd.-Bat 1: 203. 1850; C.J.Saldanha, Fl. Karnataka 2: 207. 1996

Pegia colebrookiana Wight, Icon. Pl. Ind. Orient. 1: t. 230. 1839

Distribution: Karnataka (Hassan), Kerala, Maharashtra, Tamil Nadu; endemic.

Nothopegia heyneana Gamble, Fl. Madras 265 1918; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Distribution: Karnataka (Hassan), Andhra Pradesh, Kerala, Maharashtra, Odisha, Tamil Nadu

Nothopegia racemosa (Dalzell) Ramamoorthy in Saldanha & Nicolson, Fl. Hassan Dist. 377. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Glycicarpus edulis Dalzell, J. Bombay Branch Roy. Asiat. Soc. 3: 70.1849

Glycicarpus racemosa Dalzell in J. Asiat. Soc. Bombay 3: 69. 1849

Nothopegia dalzellii Gamble, Fl. Madras 265 1918

Nothopegia dalzellii var. *angustifolia* Gamble, Fl. Madras 265.1918

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Nothopegia travancorica Bedd. ex Hook. f., Fl. Brit. India 2: 40. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan), Kerala, Tamil Nadu; endemic.

Rhus L.

Rhus mysorensis G.Don, Gen. Hist. 2:74.1832; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Distribution: Karnataka (Ballari, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Gujarat, Haryana, Punjab, Rajasthan; endemic.

Local/Common/Vernacular names: *Hulumari, Sabale, Sappl* (Kannada)

Schinus L.

Schinus molle L., Sp. Pl. 388. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 60. 1984.

Distribution: Karnataka (Bengaluru, Mysuru), Punjab, Tamil Nadu; native to tropical S America.cultivated elsewhere in tropics.

Local/Common/Vernacular names: *American pepper, Californian Pepper tree, False pepper, pepper tree, peppercorn tree, Peruvian masti, Peruvian pepper, Peruvian pepper tree, pirul* (English)

Semecarpus L. f.

Semecarpus anacardium L. f., Suppl. Pl. 182. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 60. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; Sri Lanka, Eastern Archipelago, North Australia

Local/Common/Vernacular names: *Marking nut* (English); *Gudde geru, Kari geru* (Kannada)

Semecarpus auriculata Bedd., Fl. Sylv. S. India 2: t. 232 1872; Kesh.Murthy & Yogan, Fl. Kodagu 126. 1990

Distribution: Karnataka (Kodagu, Shivamogga), Kerala, Tamil Nadu; endemic.

Semecarpus kathalekanensis Dasappa & Swam., Indian Forester 126: 78. 2000.

Distribution: Karnataka (Uttara Kannada), Goa; endemic

Spondias L.

Spondias indica (Wight & Arn.) Airy Shaw & Forman, Kew Bull. 21: 16. 1967; C.J.Saldanha, Fl. Karnataka 2: 209. 1996

Solenocarpus indicus Wight & Arn., Prodr. Fl. Ind. Orient. 1: 172. 1834

Distribution: Karnataka (Belagavi, Kodagu, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; Paleotropical

Spondias pinnata (L.f.) Kurz, Prelim. Rep. For. & Veg. Pegu Append. A.44. & B.42. 1875; B.D.Sharma et al., Fl. Karnataka Anal. 60. 1984.

Mangifera pinnata L. f., Suppl. Pl. 156. 1781

Spondias mangifera Willd., Sp. Pl. 2: 751. 1799

Spondias acuminata Roxb., Fl. Ind. 2: 453. 1832

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India; Paleotropical

Local/Common/Vernacular name: *Indian Hog Plum* (English); *Amate amara* (Kannada)

Terminthia Bernh.

Terminthia paniculata (Wall. ex G. Don) C.Y. Wu & T.L. Ming, Fl. Yunnan. 2: 408 1979

Rhus paniculata Wall. ex G. Don, Gen. Hist. 2: 73 1832; B.D.Sharma et al., Fl. Karnataka Anal. 59. 1984.

Toxicodendron paniculatum (Wall. ex G. Don) Kuntze, Revis. Gen. Pl. 1: 154. 1891

Distribution: Karnataka (Bengaluru), Andhra Pradesh; Bhutan, China, Myanmar.

Ancistrocladaceae

Ancistrocladus Wall.

Ancistrocladus heyneanus Wall. ex J. Graham, Cat. Pl. Bombay 28. 1839; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.

Plate 3C

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Annonaceae

Alphonsea Hook. f. & Thomson

Alphonsea lutea (Roxb.) Hook. f. & Thomson, Fl. Ind. 153. 1855; I. M. Turner, Nordic J. Bot. 33: 258. 2015.

Uvaria lutea Roxb., Pl. Coromandel 1: 32, t. 36. 1795

Alphonsea sclerocarpa Thwaites, Enum. Pl. Zeyl. 11. 1858

Bocagea lutea (Roxb.) Baill., Hist. Pl. 1: 215. 1868

Alphonsea madraspatana Bedd., Icon. Pl. Ind. Or. 19, t. 92. 1870

Distribution: Karnataka (), Arunachal Pradesh, Assam, Kerala, Maharashtra, Odisha, Tamil Nadu; Myanmar, Sri Lanka.

Annona L.

Annona glabra L., Sp. Pl. 537 1753

Annona palustris L., Sp. Pl. ed. 2 757 1762

Distribution: Karnataka (Chitradurga), Kerala; native to tropical America

Local/Common/Vernacular names: *Alligator apple, monkey apple pond apple, swamp apple* (English); *Kaadu seethapala* (Kannada)

Annona muricata L., Sp. Pl. 536 1753; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Distribution: Karnataka (Bengaluru, Davanagere), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Kerala, Maharashtra, Tamil Nadu, Uttar Pradesh; native to tropical America; introduced elsewhere

Annona reticulata L., Sp. Pl. 537 1753; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Davanagere, Kalburgi, Kodagu, Mysuru), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Delhi, Kerala, Madhya Pradesh, Maharashtra, Manipur, Odisha, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; native to tropical America.

Local/Common/Vernacular name: *Ramphal* (Kannada)

Annona squamosa L., Sp. Pl. 537 1753; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chitradurga, Davanagere, Dharwad, Kalburgi, Kodagu, Kolar, Mysuru, Tumakuru, Uttara Kannada), throughout India; native to tropical America

Local/Common/Vernacular name: *Sithaphal* (Kannada)

Artobotrys R. Br.

Artobotrys hexapetalus (L.f.) Bhandari, Baileya 12: 149 1965; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Annona hexapetala L.f., Suppl. Pl. 270 1782

Uvaria esculenta Roxb. ex Rottl. & Willd. in Rottler, Neue Schriften Ges. Naturf. Freunde Berlin 4: 201. 1803

Artobotrys odoratissimus R.Br., Bot. Reg. 5: t. 423 1820

Distribution: Karnataka (Belagavi, Bengaluru, Kalburgi, Kodagu, Mysuru), Andhra Pradesh, Assam, Bihar, Kerala, Madhya Pradesh, Maharashtra, Manipur, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Indomalaya.

Local/Common/Vernacular names: *Climbing ylang-ylang*, *The ylang-ylang vine* (English); *Manoranjini* (Kannada)

Artobotrys zeylanicus Hook.f. & Thomson, Fl. Ind. 1: 128 1855; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Cananga (Dunal) Hook.f. & Thomson

Cananga odorata (Lam.) Hook.f. & Thomson, Fl. Ind. 1: 130.1855

Uvaria odorata Lam., Encycl. 1: 595.1785

Unona odorata (Lam.) Dunal, Monogr. Anonac. 108.1817

Canangium odoratum (Lam.) Baill. ex King, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 61(2): 41.1892

Distribution: Karnataka (cultivated in gardens); native to NE Australia, cultivated throughout humid areas.

Local/Common/Vernacular names: *Cananga tree*, *Fragrant cananga*, *Macassar-oil plant*, *Perfume tree*, *Ylang-ylang* (English)

Cyathocalyx Champ. ex Hook. f. & Thomson

Cyathocalyx zeylanicus Champ. ex Hook.f. & Thomson, Fl. Ind. 1: 127 1855; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Distribution: Karnataka (Dakshina Kannada, Davanagere), Kerala, Tamil Nadu; Sri Lanka.

Desmos Lour.

Desmos lawii (Hook.f. & Thomson) Saff., Bull. Torrey Bot. Club 39: 506 1912

Plate 15A

Desmos chinensis auct.non Lour., 1790; C.J.Saldanha, Fl. Karnataka 1: 43. 1984

Unona discolor Vahl, Symb. Bot. 2: 63, t. 36 63 1791

Unona dunali Hook. f. & Thomson, Fl. Ind. 1: 131 131 1855

Unona lawii Hook.f. & Thomson, Fl. Ind. 1: 132 1855

Polyalthia lawii (Hook.f. & Thomson) Finet & Gagnep., Bull. Soc. Bot. France 53, M e moire4: 89. 1906

Desmos chinensis var. *lawii* (Hook.f. & Thomson) Barn, Bot. Zhurn. 59: 1774. 1974

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Friesodielsia Steenis

Friesodielsia sahyadrica N.V.Page & Survesw., Phytotaxa 158(3): 278. 2014

Distribution: Karnataka (Uttara Kannada); endemic.

Goniothalamus (Blume) Hook. f. & Thomson

Goniothalamus cardiopetalus (Dalzell) Hook.f. & Thomson, Fl. Ind. 1: 107 1855; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984.

Polyalthia cardiopetala Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 39 1850

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Goniothalamus wynadensis (Bedd.) Bedd., Icon. Pl. Ind. Or. 13. 1870; Kesh.Murthy & Yogan, Fl. Kodagu 31. 1990

Atrategia wynadensis Bedd. in Madras J. Lit. Sci. III, 1: 37. 1864

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Hubera Chaowasku

Hubera cerasoides (Roxb.) Chaowasku in Chaowasku et al., Phytotaxa 69: 47. 2012

Uvaria cerasoides Roxb., Pl. Coromandel 1: 30 1795

Guatteria cerasoides (Roxb.) Dunal, Monogr. Anon. 127. 1817

Unona cerasoides(Roxb.) Baill., Hist. Pl. 1: 272. 1868

Polyalthia cerasoides (Roxb.) Benth. & Hook. f. ex Bedd., Fl. Sylv. S. India , pl. 1 1869; C.J.Saldanha, Fl. Karnataka 1: 48. 1984

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Davanagere, Mandya, Mysuru), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Kerala, Maharashtra, Meghalaya, Odisha, Tamil Nadu, West Bengal; Bangladesh, Myanmar, Sri Lanka.

Meiogyne Miq.

Meiogyne pannosa (Dalzell) J.Sinclair, Sarawak Mus. J. 5: 604.1951; B.D.Sharma et al., Fl. Karnataka Anal. 3. 1984. Plate 15B

Unona pannosa Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 207.1851

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Meiogyne ramarowii (Dunn) Gandhi in Saldanha & Nicolson, Fl. Hassan 38. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Unona ramarowii Dunn, Bull. Misc. Inform. Kew 1914: 183 183 1914

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Miliusa Lesch. ex A. DC.

Miliusa eriocarpa Dunn. ex Gamble, Bull. Misc. Inform. Kew 1916: 58 1916; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Distribution: Karnataka (Dakshina Kannada, Mysuru, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Miliusa gokhalei Ratheesh, Sujanpal, Anil Kumar & Sivad., Phytotaxa 42: 27. 2012

Distribution: Karnataka (Kodagu), Kerala; endemic.

Miliusa horsfieldii (Bennett) Baill. ex Pierre, Fl. Forest. Cochinch., pl. 38 1881
Saccopetalum horsfieldii Bennett, Pl. Jav. Rar. 165, pl. 35. 1840
Saccopetalum arboreum Elmer, Leafl. Philipp. Bot. 5: 1739. 1913
Miliusa tectona C.E.Parkinson, Forest Fl. Andaman Isl. 75. 1923; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Saccopetalum lineatum Craib, Bull. Misc. Inform. Kew 1924: 82. 1924
Saccopetalum unguiculatum C.E.C.Fisch., Bull. Misc. Inform. Kew 1926: 454. 1926
Saccopetalum tectorum (Hutch. ex C.E.Parkinson) Chatterjee, Kew Bull. 3: 59 1948
 Distribution: Karnataka (Belagavi, Uttara Kannada)?, Australia, Andaman & Nicobar Islands; China, Indonesia, Laos, Malaysia, Myanmar, Philippines, Thailand.

Miliusa indica Lesch. ex A. DC., Mém. Soc. Phys. Genève 5: 212. 1832. Mem. Anonac. 37. 1832
Miliusa leschenaultii A. DC., Mém. Soc. Phys. Genève 5: 220, Pl. III. 1832. Mem. Anonac. 44, Pl. III. 1832
Miliusa montana var. *tomentosa* Thwaites, Enum. Pl. Zeyl. 10. 1858
Miliusa indica var. *tomentosa* (Thwaites) Hook. f. & Thomson, Fl. Brit. Ind. 1: 86. 1872
 Distribution: Karnataka (Chikkamagaluru, Kodagu, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Miliusa malnadensis N.V.Page & Nerlekar, Phytotaxa 245(1): 79. 2016. ('malnadense').
 Distribution: Karnataka (Chikkamagaluru: Kudremukh National Park); endemic.

Miliusa nilagirica Bedd., Icon. Pl. Ind. Or. 18. 1874
 Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Miliusa sclerocarpa (A. DC.) Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 41: 291. 1872
Uvaria sclerocarpa A. DC., Mém. Soc. Phys. Genève 5: 203. 1832. Mem. Anonac. 27. 1832
Saccopetalum sclerocarpum (A. DC.) Hook. f. & Thomson, Fl. Brit. Ind. 1: 88. 1872
Miliusa longiflora (Hook. f. f. & Thomson) Baill. ex Finet & Gagnep., Bull. Soc. Bot. France 53, Mémoire 4: 126. 1906
 Distribution: Karnataka(), Bihar, Sikkim, West Bengal; Bangladesh, Bhutan, Myanmar.

Miliusa tomentosa (Roxb.) J. Sinclair, Gard. Bull. Singapore 14: 378 1955; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Uvaria tomentosa Roxb., Pl. Coromandel 1: 31 1795
Saccopetalum tomentosum Hook. f. & Thomson, Fl. Ind. 1: 152 1855
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Uttara Kannada), Andhra Pradesh, Bihar, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh; Bangladesh, Nepal, Sri Lanka.

Miliusa velutina (DC.) Hook. f. & Thomson, Fl. Ind. 1: 151. 1855; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Uvaria velutina DC. in Dunal, Monogr. Fam. Anonac. 91. 1817
Guatteria villosa G. Don, Gen. Hist. 1: 100. 1831
Guatteria velutina A. DC., Mem. Soc. Phys. Genève 5: 218. 1832 & Mem. Anonac. 42. 1832
Uvaria villosa (G. Don) Roxb., Fl. Ind. 2: 664. 1832
Miliusa villosa (G. Don) W. Theob., Burmah, ed. 3. 2: 667. 1883
 Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Davanagere, Hassan, Kodagu, Tumakuru), Andhra Pradesh, Assam, Bihar, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Odisha, Uttar Pradesh, West Bengal; Bangladesh, Myanmar, Nepal.

Miliusa wightiana Hook. f. & Thomson, Fl. Ind. 1: 149 1855; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.
 Distribution: Karnataka (Hassan, Shivamogga), Kerala, Tamil Nadu; endemic.

Mitrephora (Blume) Hook. f. & Thomson

Mitrephora grandiflora Bedd., Icon. Pl. Ind. Or. 21,t. 101. 1870 & Fl. Sylv. S. India 1: t. 75 1871; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Kinginda grandiflora (Bedd.) Kuntze, Revis. Gen. Pl. 1: 7. 1891

Distribution: Karnataka (Dakshina Kannada, Mysuru, Shivamogga), Kerala; endemic.

Monoon Miq.

Monoon coffeoides (Thwaites ex Hook f. & Thomson) B. Xue & R. M. K. Saunders, Taxon 61: 1030. 2012

Guatteria coffeoides Thwaites ex Hook.f. & Thomson, Fl. Ind. 1: 141. 1855

Polyalthia coffeoides (Thwaites ex Hook. f. & Thomson) Bedd., Icon. Pl. Ind. Or. 11. 1869; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Distribution: Karnataka (Dakshina Kannada, Kodagu, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Monoon fragrans (Dalzell) B.Xue & R.M.K.Saunders, Taxon 61: 1031. 2012

Plate 15C

Guatteria fragrans Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 206. 1851

Polyalthia fragrans (Dalzell) Bedd., Fl. Brit. India 1: 63. 1872; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Monoon longifolium (Sonn.) B. Xue & R. M. K. Saunders, Taxon 61: 1032. 2012

Uvaria longifolia Sonn., Voy. Ind. Orient. 3: 260. 1782

Guatteria longifolia (Sonn.) Wall., Numer. List 644. 1832

Polyalthia longifolia (Sonn.) Thwaites, Enum. Pl. Zeyl. 398. 1864; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Sri Lanka, introduced to many tropical countries.

Local/Common/Vernacular name: *Ashoka mara* (Kannada)

Orophea Blume

Orophea malabarica Sasidh. & Sivar., Blumea 35(1): 269. 1990; Sankararao et al., J. Threatened Taxa 6996. 2015.

Distribution: Karnataka (Kodagu), Kerala; endemic.

Orophea sivarajanii Sasidh., Nordic J. Bot. 19(3): 301. 1999; Sankararao et al., J. Threatened Taxa 6996. 2015.

Distribution: Karnataka (Dakshina Kannad, Kodagu), Kerala; endemic.

Orophea uniflora Hook.f. & Thomson, Fl. Ind. 1: 110. 1855; C.J.Saldanha, Fl. Karnataka 1: 47. 1984

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Orophea zeylanica Hook.f. & Thomson, Fl. Ind. 1: 111. 1855; B.D.Sharma et al., Fl. Karnataka Anal. 4. 1984.

Distribution: Karnataka (Chikkamagaluru, Kodagu, Mysuru, Uttara Kannada), Kerala, Maharashtra; Sri Lanka.

Polyalthia Blume

Polyalthia suberosa (Roxb.) Benth. & Hook.f., Enum. Pl. Zeyl. 398.1864; B.D.Sharma et al., Fl. Karnataka Anal. 5. 1984.

Uvaria suberosa Roxb., Pl. Coromandel 1: 31, t. 34.1795

Distribution: Karnataka (Mysuru), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Odisha, Tamil Nadu, Uttar Pradesh, West Bengal; Bangladesh, China, Laos, Malaysia, Myanmar, Philippines, Sri Lanka, Thailand, Vietnam.

Sageraea Dalzell

Sageraea laurina Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 207. 1851

Plate 15D

Guatteria laurifolia J.Graham, Cat. Pl. Bombay 4 1839 non *Guatteria laurifolia* (Sw.) Dunal, 1817

Bocagea dalzellii Hook.f. & Thomson, Fl. Brit. India 1: 92.1872

Sageraea laurifolia (J.Graham) Blatt., J. Bombay Nat. Hist. Soc. 34: 294 1930; B.D.Sharma et al., Fl. Karnataka Anal. 5. 1984.

Distribution: Karnataka (Belagavi, Kodagu, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Uvaria L.

Uvaria concava Teijsm. & Binnend., Natuurk. Tijdschr. Ned.-Ind. 3: 331.1852

Uvaria lurida Hook. f. & Thomson, Fl. Ind. 101. 1855

Uvaria lurida var. *macrophylla* Hook. f. & Thomson, Fl. Ind. 101.1855

Uvaria narum var. *macrophylla* (Hook.f. & Thomson) Hook.f. & Thomson, Fl. Brit. Ind. 1: 50.1872

Uva concava (Teijsm. & Binn.) Kuntze, Revis. Gen. Pl. 1: 7.1891

Uva lurida (Hook.f. & Thomson) Kuntze, Revis. Gen. Pl. 1: 8.1891

Uvaria hookeri King, Ann. R. Bot. Gard.(Calcutta) 4: 28, t. 22. 1893; B.D.Sharma et al., Fl. Karnataka Anal. 5. 1984.

Uvaria lurida var. *sikkimensis* King, Ann. R. Bot. Gard.Calcutta 4: 29. t. 23.1893

Uvaria stellata Merr., Publ. Bur. Sci. Gov. Lab. 29: 14.1905

Uvaria eucineta Bedd. ex Dunn, Bull. Misc. Inform. Kew 1914: 182.1914; C.J.Saldanha, Fl. Karnataka 1: 50.1984.

Uvaria sympetala Merr., Univ. Calif. Publ. Bot. 15: 63.1929

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Assam, Bihar, Kerala, Maharashtra, Meghalaya, Odisha, Sikkim, Tamil Nadu, West Bengal; Australia, New Guinea, S & SE Asia.

Uvaria narum (Dunal) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 9. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 5. 1984.

Unona narum Dunal, Monogr. Anonac. 99.1817

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Apiaceae

Anethum L.

Anethum graveolens L., Sp. Pl. 263. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.

Anethum sowa Roxb. ex DC., Prodr. 4: 186. 1830

Peucedanum graveolens (L.) Hiern in Oliv., Fl. Trop. Africa 3: 19. 1871

Distribution: Karnataka (Belagavi, Hassan, Kalburgi, Yadgir), Bihar, Delhi, Himachal Pradesh, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh; native to Eurasia; cultivated elsewhere.

Local/Common/Vernacular names: *Dill*, *Indian dill* (English); *Sabsige* (Kannada)

Bunium L.

Bunium nothum (C.B.Clarke) P.K.Mukh., Bull. Bot. Surv. India 24: 43. 1983; C.J.Saldanha, Fl. Karnataka 2: 278. 1996
Carum nothum C.B.Clarke in Hook.f., Fl. Brit. India 2: 681. 1879
Bunium nilghirensis H.Wolff, Feddes Repert. 27:334. 1930
 Distribution: Karnataka (Mysuru), Kerala, Sri Lanka.

Bupleurum L.

Bupleurum ramosissimum Wight & Arn., Prodr. Fl. Ind. Orient. 1: 370. 1834; C.J.Saldanha, Fl. Karnataka 2: 278. 1996
Bupleurum mucronatum Wight & Arn., Prodr. Fl. Ind. Orient. 1: 370. 1834
Bupleurum virgatum Wight & Arn., Prodr. Fl. Ind. Orient. 1: 370. 1834, non Cav. 1827
Bupleurum falcatum var. *ramosissimum* (Wight & Arn.) Dalzell & A.Gibson, Bombay Fl. 108.1861
Bupleurum wightii P.K.Mukh., Indian Forester 95: 567. 1969.
Bupleurum wightii var. *ramosissimum* (Wight & Arn.) Chandrab. in Nair & & A.N.Henry, Fl. Tamil Nadu 1: 178.1983
Bupleurum ramosissimum var. *wightii* Bennet, J. Econ. Taxon. Bot. 5: 972.1984
Bupleurum ramosissimum var. *mucronatum* Panigrahi & S.C.Mishra, Bangladesh J. Bot. 14: 85.1985
Bupleurum ramosissimum var. *virgatum* (C.B.Clarke) Panigrahi & S.C.Mishra, Bangladesh J. Bot. 14: 86.1985
 Distribution: Karnataka (Chikkamagaluru, Dharwad, Mysuru), peninsular India; Sri Lanka.

Centella L.

Centella asiatica (L.) Urb., Fl. Bras. 11:287. t.78. f. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.
Hydrocotyle asiatica L., Sp. Pl. 234. 1753
 Distribution: Karnataka (throughout), throughout India; subtropical and tropical regions of both hemispherers
 Local/Common/Vernacular names: *Indian penny wort* (English); *Brahmi, Ilikivi gida, Ondelaga, Saraswati soppu* (Kannada)

Coriandrum L.

Coriandrum sativum L., Sp. Pl. 256. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.
 Distribution: Karnataka (cultivated throughout), cultivated throughout; native to Mediterranean region
 Local/Common/Vernacular name: *Coriander* (English); *Kothambari* (Kannada)

Cuminum L.

Cuminum cyminum L., Sp. Pl. 254 1753; B.D.Sharma et al., Fl. Karnataka Anal. 118. 1984.
 Distribution: Karnataka (Belagavi, Chitradurga, Hassan), Assam, Bihar, Delhi, Gujarat, Madhya Pradesh, Maharashtra, Manipur, Punjab, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh; native to Egypt; cultivated elsewhere.
 Local/Common/Vernacular name: *Cumin seed* (English); *Jeerige* (Kannada)

Daucus L.

Daucus carota L., Sp. Pl. 242. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.
 Distribution: Karnataka (Belagavi, Chitradurga, Kodagu), Assam, Goa, Himachal Pradesh, Jammu & Kashmir, Madhya Pradesh, Maharashtra, Manipur, Mizoram, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand; native to temperate regions of Europe, SW Asia; cultivated throughout world.
 Local/Common/Vernacular name: *Carrot* (English); *Kempu moolangi* (Kannada)

Eryngium L.

Eryngium foetidum L., Sp. Pl. 232. 1753; C.J.Saldanha, Fl. Karnataka 2: 279. 1996

Distribution: Karnataka (Belagavi, Dharwad), Arunachal Pradesh, Assam, Bihar, Kerala, Manipur, Meghalaya, Mizoram, Punjab, Tripura, Uttar Pradesh; native to C America.

Foeniculum Mill.

Foeniculum vulgare Mill., Gard. Dict. ed. 8. n. 1, 1768; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Distribution: Karnataka (Cultivated, Belagavi, Chitradurga), Assam, Bihar, Delhi, Goa, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh; native to Mediterranean region.

Local/Common/Vernacular name: *Fennel* (English); *Dodda sompu* (Kannada)

Heracleum L.

Heracleum candolleanum Gamble, Fl. Madras 565(399). 1919; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Heracleum rigens var. *candolleana* (Wight & Arn.) C.B.Clarke in Hook.f., Fl. Brit. India 2:715. 1879

Distribution: Karnataka (Chikkamagaluru, Mysuru), Kerala, Tamil Nadu; endemic.

Heracleum grande (Dalzell & A.Gibson) Mukhop., Sci. & Cult. 39(12): 542 1973; C.J.Saldanha, Fl. Karnataka 2: 281. 1996

Pastinaca grandis Dalzell & A.Gibson, Bombay Fl. 107. 1861

Peucedanum grande C.B.Clarke in Hook.f., Fl. Brit. India 2: 710 1879

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Heracleum hookerianum Wight & Arn., Prodr. Fl. Ind. Orient. 373 1834; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Distribution: Karnataka (Chikkaballapur), Tamil Nadu; endemic.

Heracleum rigens Wall. ex DC., Prodr. 4: 191. 1830; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Pastinaca candolleana Wight & Arn., Prodr. Fl. Ind. Orient. 1: 372. 1834

Pastinaca rigens Wight, Icon. Pl. Ind. Orient. 3: t. 1009. 1845

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Hassan, Kodagu, Mysuru, Shivamogga), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Heracleum spengelianum Wight & Arn., Prodr. Fl. Ind. Orient. 1: 372. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Pastinaca spengeliana Wight, Icon. Pl. Ind. Orient. 3: t. 1008. 1845

Distribution: Karnataka (Belagavi), Maharashtra, Tamil Nadu; endemic.

Karnataka P. K. Mukh. & Constance

Karnataka benthamii (C.B.Clarke) P.K.Mukh. & Constance, Brittonia 38: 145. 1986; C.J.Saldanha, Fl. Karnataka 2: 282. 1996

Schultzia ? benthamii C.B.Clarke in Hook.f., Fl. Brit. India 2: 696. 1879

Distribution: Karnataka (Dakshina Kannada); endemic.

Peucedanum L.

Peucedanum dhana Buch.-Ham. ex C.B.Clarke in Hook.f., Fl. Brit. India 2: 709. 1879; C.J.Saldanha, Fl. Karnataka 2: 282. 1996

Pastinaca glauca Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4: 293. 1859 non Peucedanum glaucum DC. 1830

var. *dhana*

Distribution: Karnataka (Belagavi), Andhra Pradesh, Bihar, Himachal Pradesh, Maharashtra, Rajasthan, Uttarakhand, West Bengal; Nepal.

var. ***dalzellii*** C.B.Clarke, Hook.f., Fl. Brit. India 2: 709. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Distribution: Karnataka (Belagavi, Dharwad), Andhra Pradesh, Madhya Pradesh, Maharashtra; endemic.

Pimpinella L.

Pimpinella adscendens Dalzell in J. Bot. 2: 261. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Distribution: Karnataka (Belagavi), Goa, Maharashtra; endemic.

Pimpinella candolleana Wight & Arn., Prodr. Fl. Ind. Orient. 1: 369. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Distribution: Karnataka (Chikkamagaluru, Mysuru), Kerala, Maharashtra, Tamil Nadu; endemic.

Pimpinella heyneana (Wall. ex DC.) Kurz, Gen. Pl. 1: 894 1867; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Helosciadium heyneanum Wall. ex DC., Prodr. 4: 106. 1830

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Assam, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu; Sri Lanka.

Pimpinella wallichiana (Miq.) Gandhi in Saldanha & Nicolson, Fl. Hassan Dist. 417. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Helosciadium wallichianum Miq. ex Hohen., Bot. Zeit. 7: 775. 1849

Pimpinella monoica Dalzell in Hook.'s J. Bot. Kew Gard. Misc. 3: 212. 1851

Pimpinella katrajensis Rao & Hemadri, Indian For. 102: 232. 1976

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Madhya Pradesh, Maharashtra, Sikkim, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Kaadu oma* (Kannada)

Pinda P. K. Mukh. & Constance

Pinda concanensis (Dalzell) P.K.Mukh. & Constance, Kew Bull. 41: 226. t. 1. 1986; C.J.Saldanha, Fl. Karnataka 2: 284. 1996

Heracleum concanense Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 260. 1850

Heracleum pinda Dalzell & A.Gibson, Bombay Fl. 107. 1861

Distribution: Karnataka (Chikkamagaluru, Shivamogga), Maharashtra; endemic.

Polyzygus Dalzell

Polyzygus tuberosus Dalzell ex Walp., Ann. Bot. Syst. (Walpers) 2(4): 715. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Plate 15E

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra

Trachyspermum Link

Trachyspermum ammi (L.) Sprague, Bull. Misc. Inform. Kew 1929: 228 1929; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Sison ammi L., Sp. Pl. 252 1753

Carum copticum (L.) Benth. & Hook.f. ex C.B.Clarke in Hook.f., Fl. Brit. India 2: 683 1879

Distribution: Karnataka (Belagavi, Bengaluru), Andhra Pradesh, Arunachal Pradesh, Bihar, Delhi, Gujarat, Himachal Pradesh, Madhya Pradesh, Maharashtra, Manipur, Punjab, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand, West Bengal

Local/Common/Vernacular name: *Carum* (English); *Oma* (Kannada)

Trachyspermum roxburghianum (DC.) H.Wolff, Fl. Siam. Enum. 1: 788. 1931; C.J.Saldanha, Fl. Karnataka 2: 285. 1996

Apium involucratum Roxb. ex Fleming, Asiat. Res. 11: 157. 1810

Ptychotis roxburghiana DC., Prodr. 4: 109. 1830

Pimpinella involucrata (Roxb.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 369. 1834

Carum stictocarpum C.B.Clarke in Hook.f., Fl. Brit. India 2: 681 1879

Trachyspermum involucratum (Roxb.) H. Wolff, Engler, Pflanzenr. 4: 228. 1927 non Maire 1922

Trachyspermum stictocarpum (C.B.Clarke) H. Wolff, Pflanzenr. (Engler) Umbellif.-Apioid.-Ammin. 89. 1927

Distribution: Karnataka (Ballari, Chikkamagaluru, Dharwad), Bihar, Maharashtra, Tamil Nadu, Uttarakhand

Local/Common/Vernacular names: *wild celery* (English); *Ajwain, Ajamoda* (Sanskrit)

Vanasushava P. K. Mukh. & Constance

Vanasushava pedata (Wight) P.K. Mukh. & Constance, Kew Bull. 29: 595 1974; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Heracleum pedatum Wight, Icon. Pl. Ind. Orient. t. 342 1840

Distribution: Karnataka (Belagavi), Kerala, Tamil Nadu; endemic.

Apocynaceae

Aganosma (Blume) G. Don

Aganosma cymosa (Roxb.) G. Don, Gen. Hist. 4: 77. 1837; B.D.Sharma et al., Fl Karnataka Anal. 161. 1984.

Echites cymosus Roxb., Fl. Ind. 2: 16. 1832

Aganosma cymosa var. *glabra* A.DC., Prodr. 8: 433. 1844

Aganosma doniana Wight, Icon. Pl. Ind. Orient. t. 1306. 1848

Aganosma cymosa var. *lanceolata* Hook.f., Fl. Brit. India 3: 665. 1882

Toxocarpus eriocarpus Hook.f., Fl. Brit. India 4: 15. 1883

Aganosma harmandiana Pierre in Spire & A. Spire, Caoutchouc Indo-Chine. 109. 1906

Aganosma cymosa var. *fulva* Craib, Repert. Spec. Nov. Regni Veg. 12: 393. 1913

var. **cymosa**

Distribution: Karnataka (Belagavi, Hassan, Mysuru, Tumakuru, Uttara Kannada), Peninsular to NE India; Bangladesh, China, Sri Lanka.

Local/Common/Vernacular names: *Chidaranabili, Chithra leena balli, Citraleena balli* (Kannada)

var. **elegans** (G.Don) Hook.f., Fl. Brit. India 3: 665. 1882; D.J. Middleton, Kew Bull. 51: 462. 1996.

Echites conferta Wall., Numer. List 1655. 1829

Aganosma elegans G.Don, Gen. Hist. 4: 77. 1837

Aganosma cymosa var. *conferta* Hook.f., Fl. Brit. India 3: 665. 1882

Distribution: Karnataka (), Peninsular India; Sri Lanka.

Aganosma dichotoma (Roth) K. Schum. in Engl. & Prantl, Nat. Pflanzenfam. IV. 2: 173. 1895

Echites dichotoma Roth, Nov. pl. sp. 136. 1821

Echites heynei Spreng., Syst. Veg. 1: 632. 1824

Aganosma caryophyllata auct. non G.Don, 1837; B.D.Sharma et al., Fl. Karnataka Anal. 160. 1984.

Distribution: Karnataka (Bengaluru, Shivamogga), Odisha, Tamil Nadu; cultivated elsewhere.

Local/Common/Vernacular names: *Malatilata* (Kannada); *Asanamallika, Asphota, Bhadramunja, Bhadravalli, Malati, Sumana, Vasanti, Visalyakrit, yuvati* (Sanskrit)

Allamanda L. ('Allemanda')

Allamanda blanchetii A.DC., Prodr. 8: 319. 1844

Allamanada violacea Gardner, Sert. Pl. 2: t. 41. 1844

Distribution: Karnataka (cultivated throughout), cultivated throughout; native to Brazil

Allamanda nobilis T.Moore, Gard. Chron. 1868: 180. 1868.

Allamanda cathartica var. *nobilis* (T.Moore) L.H.Bailey & Raffill in L.H.Bailey, Stand. Cycl. Hort. 1: 247. 1914.; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout except in Himalayas, occasionally naturalized; native to tropical America.

Local/Common/Vernacular names: *Allamanda gida, Haladi hoovu, Mithayi hoovu gida* (Kannada)

Allamanda schottii Pohl, Pl. Bras. Icon. Descr. 1: 73. 1827

Allamanda neriifolia Hook. in Curtis, Bot. Mag. 77: t. 4594. 1851; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.

Allamanda magnifica B.S.Williams, Cat. 1888: 19. 1888

Allamanda cathartica var. *schottii* (Pohl) L.H.Bailey & Raffill, Stand. Cycl. Hort. 1: 247. 1914

Distribution: Karnataka (cultivated Chikkamagaluru, Dakshina Kannada, Kodagu, Udupi), cultivated throughout except in Himalayas; native to Brazil.

Alstonia R. Br.

Alstonia macrophylla Wall. ex G. Don, Gen. hist. 4: 87. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.

Alstonia pangkorensis King & Gamble, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 74: 442. 1908

Distribution: Karnataka (cultivated in gardens & avenues), Andaman & Nicobar Islands; cultivated elsewhere.

Local/Common/Vernacular names: *Jantaala mara, Saptaparana* (Kannada)

Alstonia scholaris (L.) R. Br., Asclepiadaceae 65. 1810 & Mem. Wern. Nat. Hist. Soc. 1: 76. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.

Echites scholaris L., Mant. Pl. 1: 53. 1767

Alstonia kurzii Hook.f., Fl. Brit. India 3: 643. 1882

Distribution: Karnataka (cultivated in gardens & avenues throughout), throughout drier parts of India including Andaman & Nicobar Islands; SE Asia to Australia.

Local/Common/Vernacular names: *Devil's tree, Scholar tree* (English); *Haale, Haale mara, Kadu hale, Maddhaale* (Kannada); *Satvin* (Kunhi); *Saptaparni* (Sanskrit); *Paale* (Tulu)

Alstonia venenata R. Br., Asclepiadaceae 66. 1810 & Mem. Wern. Nat. Hist. Soc. 1: 77. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.

Alstonia venenata var. *pubescens* Monach., Pacific. Sci. 3: 156. 1949

Distribution: Karnataka (cultivated in gardens & avenues throughout), E to S India.

Local/Common/Vernacular name: *Poison Devil Tree* (English); *Adda sarpa* (Kannada)

Anodendron A.DC.

Anodendron paniculatum A.DC. in DC., Prodr. 8: 444. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984. **Plate 3E**

Echites paniculata auct. non Poir. 1812; Roxb., Fl. Ind. 2: 17. 1832

Anodendron rhinosporum Thwaites, Enum.P. Zeyl. 194. 1860

Anodendron manubriatum Merr., Philipp. J. Sci. 7: 333. 1912

Distribution: Karnataka (Dakshina Kannada, Kodagu, Mysuru, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Peninsular to NE India.

Asclepias L.

Asclepias curassavica L., Sp. pl. 215. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984 .**Plate 3D**

Distribution: Karnataka (cultivated as ornamental, also run wild throughout), cultivated as ornamental & also run wild throughout India; native to West Indies.

Local/Common/Vernacular names: *Kakathundi, Kurki* (Kannada)

Asclepias fruticosa L., Sp. pl. 216. 1753

Gomphocarpus fruticosus (L.) Aiton, Hort. Kew. ed. 2. 2: 80. 1811

Distribution: Karnataka (cultivated as an ornamental in Bengaluru), cultivated in various parts of India; native to West Indies.

Asclepias physocarpa (E.Mey.) Schltr., Bot. Jahrb. Syst. 21, Beibl. 54: 8. 1895; Gamble, Fl. Madras 2: 585. 1967.

Gomphocarpus physocarpus E.Mey. in E. Mey & Drège, Comm. Pl. Afr. Austr. 202. (1837) 1838

Distribution: Karnataka (cultivated as an ornamental in Bengaluru, Dakshina Kannada, Mysuru), cultivated in various parts of India; native to S & tropical Africa.

Beaumontia Wall.

Beaumontia grandiflora Wall., Tent. Fl. Napal. 15, t. 7. 1824

Plate 3F&15F

Distribution: Karnataka (Bengaluru), throughout India; Bangladesh, Bhutan, Laos, Myanmar, Nepal, Thailand, Vietnam.

Beaumontia jerdoniana Wight, Icon. Pl. Ind. Orient. 4: tt. 1314-15.1848; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.

Distribution: Karnataka (Belagavi, Hassan, Kodagu, Mysuru, Uttara Kannada), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Boucerosia Wight & Arn.

Boucerosia crenulata (Wall.) Wight & Arn., Contr. Bot. India: 34. 1834.

Boucerosia truncatocoronata Sedgw., J. Indian Bot. Soc. 2: 123. 1921

Caralluma truncatocoronata (Sedgw.) Gravely & Mayur., Bull. Madras Gov. Mus. 4: 21. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Caralluma nilagiriana Kumari & Subba Rao, J. Bombay Nat. Hist. Soc. 73: 194. 1976

Caralluma crenulata Wall., Pl. Asiat. Rar. 1: 6. 1829.

Distribution: Karnataka (Chitradurga, Dharwad, Uttara Kannada), Maharashtra, Tamil Nadu; endemic.

Boucerosia diffusa Wight, Icon. Pl. Ind. Orient. 4: t.1599. 1850

Desmidorchis diffusa (Wight) Kuntze, Revis. Gen. Pl. 2: 418. 1891

Caralluma diffusa (Wight) N.E.Br., Gard. Chron. 2: 369. 1892; Jagtap & N.P.Singh, Fasc. Fl. India 24: 200. 1999.

Distribution: Karnataka (Chitradurga), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Boucerosia indica (Wight & Arn.) Plowes, Haseltonia 3: 59. 1995.

Hutchinia indica Wight & Arn. in Wight, Contrib. Bot. India 35.1834

Boucerosia hutchinia Decne. in DC., Prodr. 8: 649. 1839 ('hutchinia')

Desmidorchis indica (Wight & Arn.) Kuntze, Revis. Gen. Pl. 2: 418. 1891

Caralluma indica (Wight & Arn.) N.E.Br., Gard. Chron. 2: 369. 1892; Jagtap & N.P.Singh, Fasc. Fl. India 24: 200. 1999.

Distribution: Karnataka (), Andhra Pradesh, Tamil Nadu; endemic.

Boucerosia umbellata (Haw.) Wight & Arn., Contr. Bot. India 34.1834

Caralluma umbellata Haw., Syn. Pl. Succ. 47.1812; Bhaskar & Kushalappa, Fl. Tumkur Dist. 61.

2013.

Plate 3G

Stapelia umbellata Donn ex Roxb., Pl. Coromandel 3: 36.1815

Desmidorchis umbellata (Haw.) Decne., Ann. Sci. Nat., Bot. II, 9: 266.1838

Boucerosia campanulata Wight, Icon. Pl. Ind. Orient. 4: t. 1287. 1848

Boucerosia lasiantha Wight, Icon. pl. Ind. orient. t. 1286. 1848

Boucerosia umbellata var. *campanulata* (Wight) Hook.f., Fl. Brit. India 4: 77. 1883

Stapelia callamulia Buch.-Ham. ex Hook.f., Fl. Brit. India 4: 77.1883

Caralluma campanulata (Wight) N. E. Br., Gard. Chron. 2: 369. 1892; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Caralluma lasiantha (Wight) N. E. Br., Gard. Chron. 2: 369. 1892

Desmidorchis umbellata (Haw.) M.R.Almeida, Fl. Maharashtra 3A: 242. 2001.

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkaballapur, Chitradurga, Dakshina Kannada, Davanegere, Dharwad, Hassan, Kalaburgi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada) Throughout drier parts of India; Myanmar.

Local/Common/Vernacular names: *Chikka kavali, Kar manji, Karande, Kwali balli* (Kannada); *Karvanda* (Kunhi)

Brachystelma R. Br.

Brachystelma ciliatum Arekal & T.M. Ramakrishna, Curr. Sci. 50: 145. 1981; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Distribution: Karnataka (Kolar); endemic.

Brachystelma edulis Collett & Hemsl., J. Linn. Soc., Bot. 28: 89, t. 14. 1890; S.R. Yadav, Salunkhe & G.B. Dixit, J. Bombay Nat. Hist. Soc. 86: 480. 1990.

Distribution: Karnataka (Dakshina Kannada, Udupi), Maharashtra; endemic.

Local/Common/Vernacular name: *Devara gedde* (Kannada)

Brachystelma elenaduensis M.B.S. Char, Curr. Sci. 47: 965-966, ff. A-B, 1-6. 1978; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Distribution: Karnataka (Tumakuru), Tamil Nadu; endemic.

Brachystelma kolarensis Arekal & T.M. Ramakrishna, Proc. Indian Acad. Sci. 90: 203. 1981; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Distribution: Karnataka (Kolar); endemic.

Brachystelma maculatum Hook.f., Fl. Brit. India 4: 65. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Brachystelma shrirangii Kambale, Gholve & Sardesai, Rheedia 26(2): 145. 2016

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Brachystelma volubile Hook.f., Fl. Brit. India 4: 65. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.
 Distribution: Karnataka (Hassan), Andhra Pradesh, Tamil Nadu; endemic.

Calotropis R. Br.

Calotropis gigantea (L.) R. Br. in W.T. Aiton, Hortus Kew. ed. 2. 2: 78. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Asclepias gigantea L., Sp. Pl. 214. 1753

Distribution: Karnataka (throughout), throughout India; Africa, Iran, Nepal, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Ekka*, *Ekkada gida*, *Kar yekkada gida*, *Yekkada gida* (Kannada); *Arka* (Sanskrit)

Calotropis procera (Aiton) R. Br. in W.T. Aiton, Hortus Kew. ed. 2. 2: 78. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Asclepias procera Aiton, Hort. kew. 1: 305. 1789

Calotropis hamiltonii Wight, Contr. Bot. India 53. 1834

Distribution: Karnataka (Belagavi, Bengaluru, Kalaburgi, Tumakuru, Yadgir), throughout tropical India; Africa, Iran, Iraq, Nepal.

Local/Common/Vernacular name: *Bili ekkada gida* (Kannada)

Caralluma R. Br.

Caralluma adscendens (Roxb.) Haw., Syn. pl. succ. 47. 1812; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Stapelia adscendens Roxb., Pl. Coromandell 1: 28, t. 30. 1795

Boucerosia adscendens (Roxb.) Wall., Numer. List 8146. 1847.

var. adscendens

Distribution: Karnataka (Chitradurga, Hassan, Tumakuru), Peninsular & E India; Myanmar, Sri Lanka.

var. gracilis Gravely & Mayur., Bull. Madras Gov. Mus. n.s., 4(1): 14. 1931; N. M. G. Babu, Fl. Ethnob. Chitradurga Dist. 348. 2011

Distribution: Karnataka (Chitradurga), Maharashtra, Tamil Nadu; endemic.

Caralluma attenuata Wight, Icon. Pl. Ind. Orient. t. 1268. 1848 & Ill. Ind. Bot. t. 155 b, f. B. 1850; Hook.f., Fl. Brit. India 4: 76. 1883.

Caralluma adscendens var. *attenuata* (Wight) Gravely & Mayur., Bull. Madras Gov. Mus. 4: 13. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Distribution: Karnataka (Ballari, Chikkamagaluru, Chitradurga, Hassan, Mysuru, Tumakuru, Vijayapura), Andhra Pradesh, Kerala, Maharashtra, Odisha, Tamil Nadu; endemic.

Caralluma fimbriata Wall., Pl. Asiat. Rar. 1: 7, t. 8. 1829

Caralluma adscendens var. *fimbriata* (Wall.) Gravely & Mayur., Bull. Madras Gov. Mus. 4: 12. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Distribution: Karnataka (Ballari, Chitradurga, Tumakuru), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Caralluma stalagmifera C.E.C.Fisch., Bull. Misc. Inform. 1925: 430. 1925; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Caralluma stalagmifera var. *longipetala* Karupp. & Pull., Rheedea 17: 42. 2007; Naik et al., J. Threatened Taxa 4(6): 2656. 2012.

Distribution: Karnataka (Chitradurga, Hassan, Kalaburgi, Yadgir), Tamil Nadu; endemic.

Local/Common/Vernacular name: *Mangana kodu* (Kannada)

Carissa L.

Carissa carandas L., Mant. Pl. 1: 52. 1767; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.
Carissa salicina Lam., Encycl. 1: 554. 1785; N.M.G.Babu, Fl. Ethnob. Chitradurga Dist. 338. 2011
 Distribution: Karnataka (wild/cultivated in most districts), throughout India; S & SE Asia.
 Local/Common/Vernacular names: *Dodda kavale*, *Kaare kayi*, *Kavale kaayi gida* (Kannada)

Carissa macrocarpa (Eckl.) A.DC., Prodr. 8: 336. 1844
Arduina macrocarpa Eckl., S. African Quart. J. 1: 372. 1830
Arduina grandiflora E.Mey., Comm. Pl. Afr. Austr. 190. 1838
Carissa grandiflora (E.Mey.) A.DC. in DC., Prodr. 8: 335. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.
Jasminonarium grandiflorum (E.Mey.) Kuntze, Revis. Gen. Pl. 2: 414. 1891
Jasminonarium macrocarpum (Eckl.) Kuntze, Revis. Gen. Pl. 2: 414. 1891
 Distribution: Karnataka (Uttara Kannada), cultivated throughout drier parts of India.

Carissa spinarum L., Mant. Pl. 2: 559. 1771.
Carissa inermis Vahl, Symb. bot. 3: 43. 1794; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.
Carissa axillaris Roxb., Fl. Ind. 1, 1: 691. 1820
Carissa hirsuta Roth, Nov. Pl. Sp. 128. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.
Carissa diffusa Roxb., Fl. Ind. 1, 2: 524. 1824
Carissa villosa Roxb., Fl. Ind. 2: 525. 1824
Carissa macrophylla A.DC. in DC., Prodr. 8: 332. 1844; Hook.f., Fl. Brit. India 3: 631. 1882.
Carissa paucinervia A.DC. in Dc., Prodr. 8: 333. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.
Carissa congesta Wight, Icon. Pl. Ind. Orient. 4: t. 1289. 1848; B.D.Sharma et al., Fl. Karnataka Anal. 161. 1984.
Carissa carandas var. *congesta* (Wight) Bedd., Fl. Sylv. S. India 156. 1871
Carissa dalzellii Bedd., Fl. Sylv. S. India 156. 1871
Carissa suavissima Bedd. ex Hook.f., Fl. Brit. India 3: 632. 1882
Carissa inermis var. *macrophylla* Haines, Bot. Bihar Orissa 535. 1922
Carissa gangetica Stapf ex Gamble, Fl. Madras 805. 1923
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Odisha, Tamil Nadu; endemic.
 Local/Common/Vernacular name: *Gundala mara* (Kannada)

Catharanthus G.Don

Catharanthus pusillus (Murray) G. Don, Gen. Hist. 4: 95. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.
Vinca pusilla Murray, Comm. Act. Goett. 3: 66, t. 2, f. 1. 1773
Vinca parviflora Retz., Observ. Bot. 2: 14. 1781
Lochnera pusilla (Murray) K. Schum. in Engl. & Prantl, Nat. Pflanzenfam. IV, 2: 145. 1895
 Distribution: Karnataka (throughout as a weed in cultivated fields), throughout India; Sri Lanka.
 Local/Common/Vernacular names: *Kasi kanagale*, *Visha Kanigale gida*, *Visha Kanigale soppu* (Kannada)

Catharanthus roseus (L.) G. Don, Gen. Hist. 4: 95. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.
Vinca rosea L., Syst. nat. ed. 10. 2: 944. 1759; Hook.f., Fl. Brit. India 3: 640. 1882.
Pervinca rosea (L.) Gaterau, Descr. Pl. Montauban 52. 1789
Lochnera rosea (L.) Rchb., Consp. regn. veg. 134. 1828
Ammocallis rosea (L.) Small, Fl. S.E. U.S. 936. 1903
 Distribution: Karnataka (cultivated as an ornamental throughout/naturalised), cultivated & also runs wild in India; native to Madagascar.
 Local/Common/Vernacular names: *Madagascar periwinkle* (English); *Ganesha huvi*, *Kanagelu*, *Kempu kasi kanigale*, *Nithya pushpa*, *Sadaa pushpa* (Kannada)

Cerbera L.

Cerbera odollam Gaertn., Fruct. sem. pl. 2: 193, t. 124. 1791; Hook.f., Fl. Brit. India 3: 638. 1882.

Cerbera manghas auct. non L., 1753; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.

Cerbera lactaria Buch.-Ham. ex Spreng., Syst. Veg. 1: 642. 1825

Tanghinia odollam (Gaertn.) G.Don, Gen. Hist. 4: 98. 1837

Odollamia malabarica Raf., Sylva Tellur. 162. 1838

Tanghinia lactaria (Buch.-Ham. ex Spreng.) G.Don, Hort. Brit. ed. 3: 461. 1839

Distribution: Karnataka (Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), Salt swamps of Andaman & Nicobar Islands, E & W coasts of India; Australia, China, Malaysia, Pacific Islands, Sri Lanka.

Local/Common/Vernacular names: *Honde, Kaande* (Kannada); *Tende mara* (Tulu)

Ceropegia L.

Ceropegia attenuata Hook., Icon. Pl. ser. 2, 5: t. 867. 1852; Hook.f., Fl. Brit. India 4: 67. 1883. **Plate 15G**

Distribution: Karnataka (Dakkisna Kannada, Udupi, Uttara Kannada), Goa, Maharashtra, Rajasthan; endemic.

Ceropegia barnesii E.A.Bruce & Chatterjee, Kew Bull. 1948: 62. 1948

Distribution: Karnataka (Chamarajanagar), Tamil Nadu; endemic.

Ceropegia bhatii S.R.Yadav & Shendage, Kew Bull. 65(1): 107. 2010

Distribution: Karnataka (Davangere-Malebennur Ghat); endemic.

Ceropegia bulbosa Roxb., Pl. Coromandel 1: 11, t. 7. 1795; B.D.Sharma et al., Fl. Karnataka Anal. 165. 1984.

Ceropegia acuminata Roxb., Pl. Coromandel 1: 12, t. 8. 1795

Ceropegia lushii J. Graham in Curtis, Bot. Mag. 61: t. 3300. 1834

Ceropegia esculenta Edgew. in J. Linn. Soc., Bot. 6: 204. 1862

Ceropegia bulbosa var. *esculenta* (Edgew.) Hook.f., Fl. Brit. India 4: 68. 1883

Ceropegia bulbosa var. *lushii* (J.Graham) Hook.f., Fl. Brit. India 4: 68. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Ceropegia discreta N. E. Br. in Bull. Misc. Inform. 1909: 262. 1909

Distribution: Karnataka (throughout), throughout India; Pakistan.

Ceropegia candelabrum L., Sp. Pl. 211. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Ceropegia biflora L., Sp. Pl. 211. 1753

Ceropegia tuberosa Roxb., Pl. Coromandel 1: 12, t. 9. 1795; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Ceropegia elliottii Hook.f., Fl. Brit. India 4: 70. 1883

Ceropegia candelabrum var. *biflora* (L.) Ansari, Fasc. Fl. India 16: 12. 1984

Ceropegia candelabrum var. *tuberosa* (Roxb.) N.P.Singh, Fl. E. Karnataka 1: 410. 1988

Distribution: Karnataka (Chikkmagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga Tumakuru, Udupi), Peninsular to NE India; Sri Lanka.

Ceropegia decaisneana Wight, Icon. Pl. Ind. Orient. t. 1259. 1848; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Ceropegia brevicollis Hook.f., Fl. Brit. India 4: 74. 1883, p.p.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Ceropegia elegans Wall. in Curtis, Bot. Mag. 57: t. 3015. 1830; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Ceropegia sphenantha Wight & Arn. in Wight, Contr. Bot. India 31. 1834

Ceropegia mysurensis Wight, Icon. Pl. Ind. Orient. t. 846. 1845; Hook.f., Fl. Brit. India 4: 69. 1883.

Ceropegia walkerae Wight, Icon. Pl. Ind. Orient. t. 1266. 1848

Distribution: Karnataka (Chikkamagaluru, Hassan, Mysuru, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Ceropegia fantastica Sedgw., J. Indian Bot. Soc. 2: 124. 1921; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Plate 16A

Distribution: Karnataka (Uttara Kannada), Goa, Maharashtra; endemic.

Ceropegia fimbriifera Bedd., Madras J. Lit. Sci. ser. 3, 1: 53. 1864 & Icon. pl. Ind. or. t. 172. 1874; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Distribution: Karnataka (Hassan, Kolar, Mysuru), Kerala, Tamil Nadu; endemic.

Ceropegia hirsuta Wight & Arn. in Wight, Contr. Bot. India 30. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Ceropegia jacquemontiana Decne. in DC., Prodr. 8: 641. 1844

Ceropegia ophiocephala Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 259. 1850

Ceropegia hirsuta var. *jacquemontiana* (Decne.) Hook.f., Fl. Brit. India 4: 72. 1883

Ceropegia hirsuta var. *ophiocephala* (Dalzell) Hook.f., Fl. Brit. India 4: 72. 1883

Ceropegia hirsuta var. *stenophylla* Hook.f., Fl. Brit. India 4: 72. 1883

Ceropegia hispida Blatt. & McCann, J. Bombay Nat. Hist. Soc. 35: 409. 1931

Distribution: Karnataka (Chikmagluru, Chitradurga, Davanagere, Hassan, Mysuru, Tumakuru), throughout India except Himalaya.

Ceropegia intermedia Wight, Icon. Pl. Ind. Orient. t. 1263. 1848 emend. H. Huber in Mem. Soc. Brot. 12: 61, t. 2, f. 26. 1957; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Distribution: Karnataka (Kodagu, Mysuru, Shivamogga), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Ceropegia jainii Ansari & B.G. Kulk., Bull. Bot. Surv. India 22(1–4): 221. 1982

Distribution: Karnataka, Maharashtra; endemic.

Ceropegia juncea Roxb., Pl. Coromandell: 12, t. 10. 1795; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Hassan, Kodagu, Mysuru, Tumakuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Ceropegia metziana Miq., Anal. bot. ind. 3: 11. 1852; Ansari in Fasc. Fl. India 16: 25. 1984.

Distribution: Karnataka (), Kerala, Tamil Nadu; endemic.

Ceropegia pusilla Wight & Arn. in Wight, Contr. Bot. India 31. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Distribution: Karnataka (Mysuru), Kerala, Maharashtra, Tamil Nadu; endemic.

Ceropegia spiralis Wight, Icon. pl. Ind. orient. t. 1267. 1848; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Plate 16B

Ceropegia munroii Wight, Icon. Pl. Ind. Orient. t. 1264. 1848

Distribution: Karnataka (Bengaluru, Chitradurga, Hassan, Tumakuru), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Chilocarpus Blume

Chilocarpus denudatus Blume, Bijdr., 1025. 1826; Bakh.f., Blumea 6: 385. 1950.

Hunteria atrovirens G.Don, Gen. Hist. 4: 105. 1837

Chilocarpus atrovirens (G.Don) Blume, Mus. Bot. 1: 153. 1850 ('atroviridis') ; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.

Chilocarpus malabaricus Bedd., Icon. Pl. Ind. Or. t. 175. 1874; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.

Winchia atroviridis (Blume) Kurz, Forest Fl. Burma 2: 170. 1877

Chilocarpus atroviridis Hook.f., Fl. Brit. India 3: 626. 1882 p.p.

Chilocarpus sunainaianus Yogan., Curr. Sci. 51: 902. 1982

Chilocarpus denudatus var. *nicobaricus* M.Gangop. & Chakrab., J. Econ. Taxon. Bot. 16: 37. 1992

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga, Uttara kannada), Andaman & Nicobar Islands, Kerala; S & SE Asia.

Chonemorpha G.Don

Chonemorpha fragrans (Moon) Alston in Ann. Roy. Bot. Gard. (Peradeniya) 11: 203. 1929; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.

Echites macrophylla auct. non Kunth, 1819; Roxb., Fl. Ind. 2: 13. 1832.

Echites fragrans Moon, Cat. Pl. Ceylon 20. 1824

Chonemorpha macrophylla G.Don, Gen. Hist. 4: 76. 1837

Epichysianthus macrophyllus (G.Don) Voigt, Hort. Suburb. Calcutt. 523. 1845

Chonemorpha griffithii Hook.f., Fl. Brit. India 3: 662. 1882

Beluttakaka griffithii (Hook.f.) Kuntze, Revis. Gen. Pl. 2: 413. 1891

Chonemorpha rheedei Ridl., Agric. Bull. Straits Fed. Malay States 10: 146. 1911

Chonemorpha valvata Chatterjee, Kew Bull. 2: 51. 1947

Chonemorpha grandiflora (Roth) M.R. & S.M. Almeida, J. Bombay Nat. Hist. Soc. 90: 427. 1994 based on *Echites grandiflora* Roth, 1821 non G. Mey., 1818

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Kodagu, Shivamogga, Udupi, Uttara Kannada), throughout India, Andaman & Nicobar Islands; S & SE Asia.

Cosmostigma Wight

Cosmostigma cordatum (Poir.) M.R.Almeida, Fl. Maharashtra 3A: 239. 2001

Asclepias racemosa Roxb., Fl. Ind. ed.2, 2: 32. 1832

Cosmostigma racemosum (Roxb.) Wight, Contr. Bot. India 42. 1834 & Icon. Pl. Ind. Orient. t. 591. 1842; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Cosmostigma acuminatum Wight, Icon. Pl. Ind. Orient. 4: t. 1270. 1848

Distribution: Karnataka (Hassan, Shivamogga, Tumakuru, Uttara Kannada, Udupi), Peninsular to E India; Indonesia, Myanmar, Sri Lanka.

Local/Common/Vernacular name: *Peru Kujumbe* (Tulu)

Cryptolepis R. Br.

Cryptolepis dubia (Burm.f.) M.R.Almeida, Fl. Maharashtra 3A: 221. 2001

Periploca dubia Burm.f., Fl. Indica 70. 1768

Cryptolepis buchananii Roem. & Schult., Syst. veg. 4: 409. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 170. 1984.

Nerium reticulatum Roxb., Fl. Ind. ed.2, 2: 8. 1832

Distribution: Karnataka (throughout), throughout India; China, Myanmar, Nepal, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Halu balli*, *Kare balli*, *Kari naamada balli*, *Kurubantana balli*, *Matle hambu*, *Metaguli hambu* (Kannada); *Karantha* (Kunhi); *Jambu patra sariva* (Sanskrit)

Cryptolepis grandiflora Wight, Icon. Pl. Ind. Orient. t. 831. 1845 & Ill. Ind. Bot. 2: 169, t. 155 b, f. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 170. 1984.

Distribution: Karnataka (Chikkaballapur, Kolar, Tumakuru), Andaman & Nicobar Islands, S & E India.

Local/Common/Vernacular name: *Hambu rubber gida* (Kannada)

Cryptolepis sinensis (Lour.) Merr., Philipp. J. Sci. 15: 254. 1919; Bullock, Kew Bull. 1955: 282. 1955.

Pergularia sinensis Lour., Fl. Cochinch. 169. 1790

Cryptolepis elegans Wall. ex G. Don, Gen. Hist. 4: 82. 1837; Hook.f., Fl. Brit. India 4: 6. 1883.

Distribution: Karnataka (Kolar), Andaman & Nicobar Islands, S to NE India including tropical E Himalaya; Cambodia, China, Indonesia, Malaysia, Vietnam.

Cryptostegia R. Br.

Cryptostegia grandiflora R. Br. in Bot. Reg. 5: t. 435. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 170. 1984.

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Dakshina Kannada, Kalaburgi, Mysuru, Raichur, Udupi, Uttara Kannada, Vijayapura, Yadgir), naturalised throughout India; native to tropical Africa & Madagascar.

Local/Common/Vernacular names: *Hambu rubber gida, Rubber hambu* (Kannada)

Cryptostegia madagascariensis Bojer ex Decne in DC., Prodr. 8: 492. 1844

Distribution: Karnataka (cultivated in Dakshina Kannada, Udupi), naturalised throughout India; native to Madagascar.

Cynanchum L.

Cynanchum alatum Wight & Arn. in Wight, Contr. Bot. India 57. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 166. 1984.

Distribution: Karnataka (Chikkamagaluru, Kodagu, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Cynanchum callialatum Buch.-Ham. ex Wight, Contr. Bot. India 56. 1834 & Icon. Pl. Ind. Orient. t. 1279. 1848; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Peninsular to NE India; Myanmar.

Cynanchum parviflorum (R.Br.) Alain, Bull. Torrey Bot. Club 90: 191. 1963

Distribution: Karnataka (Tumakuru) ?

Cynanchum tunicatum (Retz.) Alston in Trimen, Handb. Fl. Ceylon 6 (Suppl.): 194. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Periploca tunicata Retz., Observ. Bot. 2: 15. 1781

Cynanchum pauciflorum R. Br., Mem. Wern. Nat. Hist. Soc. 1: 45. 1811

Cynoctonum pauciflorum (R.Br.) Decne., Prodr. 8: 528. 1844

Cynanchum inconspicuum Griff., Not. Pl. Asiat. 4: 66. 1854

Vincetoxicum tunicatum (Retz.) Kuntze, Revis. Gen. Pl. 2: 423. 1891

Distribution: Karnataka (Belagavi, Bengaluru, Dharwad, Hassan, Mysuru, Tumakuru, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Kodu muruka balli* (Kannada)

Decalepis Wight & Arn.

Decalepis hamiltonii Wight & Arn. in Wight, Contr. bot. India 64. 1834; Wight, Icon. pl. Ind. orient. t. 1285. 1848; Hook.f., Fl. Brit. India 4: 11. 1883.

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Davanagere, Hassan, Kolar, Mysuru, Tumakuru, also cultivated for roots), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Magali beru, Makli beru* (Kannada)

Dregea E.Mey.
Wattakaka Hassk.

Dregea volubilis (L.f.) Benth. ex Hook. f., Fl. Brit. India 4: 46. 1883

Asclepias volubilis L. f., Suppl. pl. 170. 1782

Hoya viridiflora R. Br. in Mem. Wern. Nat. Hist. Soc. 1:26. 1809

Marsdenia volubilis (L.f.) T. Cooke, Fl. Bombay 2: 166. 1904

Wattakaka volubilis (L.f.) Stapf in Curtis, Bot. Mag. 148:t. 8976. 1923; B.D.Sharma et al., Fl. Karnataka Anal. 170. 1984.

var. *volubilis*

Distribution: Karnataka (throughout), throughout tropical India; Pakistan, Sri Lanka, SE Asia.

Local/Common/Vernacular names: *Dugdhiga*, *Hegal balli*, *Kadu hale balli* (Kannada); *Erme tajank*, *Petta tajank* (Tulu)

var. *angustifolia* Hook. f., Fl. Brit. India 4: 47. 1883

Marsdenia lanceolata T. Cooke, Fl. Bombay 2: 167. 1904

Marsdenia volubilis var. *angustifolia* (Hook.f.) Blatt. & McCann, J. Bombay Nat. Hist. Soc. 36: 532. 1933

Dregea angustifolia (Hook.f.) Santapau & Irani, Bot. Mem. Univ. Bombay 4: 41. 1962

Dregea lanceolata (T.Cooke) Santapau & Wagh, Bull. Bot. Surv. India 5: 106. 1964

Wattakaka lanceolata (T.Cooke) A.P. Jagtap & N.P. Singh, Kew Bull. 46: 359. 1991

Wattakaka angustifolia (Hook.f.) S.D. Deshp. in S.D.Deshp., B.D. Sharma & M.P. Nayar, Fl. Mahabaleshwar 1: 369. 1993

Distribution: Karnataka (), E. to Peninsular India.

var. *lacuna* (Buch.-Ham) ex Wight Hook. f., Fl. Brit. India 4: 47. 1883

Hoya lacuna Buch.-Ham. ex Wight, Contr. Bot. India 39. 1834

Wattakaka volubilis var. *lacuna* (Buch.-Ham) ex Wight M.A. Rahman & Wilcock, J. Econ. Taxon. Bot. 13: 183. 1989

Distribution: Karnataka (throughout), throughout India; Myanmar.

Genianthus Hook.f.

Genianthus laurifolius (Roxb.) Hook.f., Fl. Brit. India 4: 16. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Asclepias laurifolia Roxb., Fl. Ind. ed.2, 2: 49. 1832

Toxocarpus laurifolius (Roxb.) Wight, Contr. Bot. India 61. 1834 & Icon. Pl. Ind. Orient. 2:t. 598. 1843

Secamone laurifolia (Roxb.) K. Schum. in Engl. & Prantl, Nat. Pflanzenfam. 4(2): 263. 1893

Distribution: Karnataka (Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttar Kannada), Andaman & Nicobar Islands, Assam, Goa, Kerala, Maharashtra, Sikkim, West Bengal; Bangladesh, China? Myanmar

Gymnema R. Br.

Gymnema cuspidatum (Thunb.) Kuntze, Revis. Gen. Pl. 2: 420. 1891; Santapau Bot. Mem. Univ. Bombay 4: 50. 1962

Cynanchum cuspidatum Thunb., Observ. Cynanch. 50. 1821

Gymnema pergularioides Wight & Gardner in Thwaites, Enum. pl. zeyl. 198. 1860; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Bidaria cuspidata (Thunb.) H. Huber in Abeyw., Rev. Handb. Fl. Ceylon 1(1): 47. 1973

Distribution: Karnataka (Dharwad, Haveri), Maharashtra; Sri Lanka.

Gymnema decaisneanum Wight, Icon. Pl. Ind. Orient. 4: t. 1271. 1848

Gymnema hirsutum auct. non Wall., 1826; Wight & Arn. in Wight, Contr. Bot. India 44. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Gymnema sylvestre var. *decaisneana* (Wight) Thwaites, Enum. Pl. Zeyl. 3: 197. 1860

Gymnema mohanramii Karthik. & Moorthy, Fl. Pl. India 171. 2009

Distribution: Karnataka (Shivamogga), Peninsular to E & N India; Sri Lanka.

Gymnema inodorum (Lour.) Decne. in DC., Prodr. 8: 551. 1844

Cynanchum inodorum Lour., Fl. Cochinch. 1: 166. 1790

Asclepias tingens Roxb., Pl. Coromandel 3: 34, t. 239. 1815

Gymnema tingens (Roxb.) Spreng., Syst. Veg. ed. 16. 1: 844. 1825

Gymnema tingens var. *cordifolia* Wight, Icon. Pl. Ind. Orient. t. 593. 1843

Bidaria inodora Decne. in DC., Prodr. 8: 624. 1844

Bidaria tingens (Roxb.) Decne. in DC., Prodr. 8: 623. 1844

Distribution: Karnataka (Shivamogga), S to NE India, tropical Himalayas.

Gymnema montanum (Roxb.) Hook.f., Fl. Brit. India 4: 31. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Asclepias montana Roxb., Fl. Ind. ed. 2, 2: 45. 1832

Bidaria montana (Roxb.) M.A. Rahman & Wilcock, Blumea 34: 99. 1989

Distribution: Karnataka (Kodagu, Shivamogga), Peninsular to N India; endemic.

Gymnema sylvestre (Retz.) R. Br. ex Schult., Syst. veg. 6: 57. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Periploca sylvestris Retz., Observ. Bot. 2: 15. 1781

Marsdenia sylvestris (Retz.) P.I.Forst., Austral. Syst. Bot. 8: 694. 1995

Distribution: Karnataka (throughout), Peninsular to N & E India; Africa, S & SE Asia.

Local/Common/Vernacular names: *Small Indian Ipecacunha* (English); *Dhudkondi, Kadasige, Koda patter, Madhu nasha balli, Madhu nashini, Sanna aggera, Sanna gerese hambu, Sappe soppu, Uparsal, Urigamahari, Uriyamari* (Kannada); *Aja balli, Aja gandhini, Aja shrengi, Aja shringi, Akshi beshaja, Anyada, Aavarthini, Balachakshu, Bishani, Bhishaarni, Chakrashreni, Chakshubahla, Chakshushya, Grihadruma, Karnika, Kasheena vartha, Maurvi, Madhu nashini, Meda shringi, Meda singi, Medha shingi, Megha balli, Menda singi, Mesharingi, Mesha shringi* (Sanskrit)

Hemidesmus R. Br.

Hemidesmus indicus (L.) W. T. Aiton, Hortus kew. ed. 2. 2: 75. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 170. 1984.

Periploca indica L., Sp. pl. 211. 1753

var. **indicus**

Distribution: Karnataka (throughout), throughout India; Iran, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Bastarda sarsaparilla, False sarsaparilla, Fragrant root, Indian sarsaparilla* (English); *Anantha moola, Dhoothaphrynium, Haalu baeru, Haalu balli, Kanya baeru, Kari bandha, Kempu gandha, Makali beru, Namasoghade, Nammada beru, Nannari beru, Sogade beru, Sugandhi balli, Sugandhi beru* (Kannada); *Anantha, Anantha moola, Anantha sariva, Sariva* (Sanskrit)

var. **pubescens** (Wight & Arn.) Hook.f., Fl. Brit. India 4: 5. 1883

Hemidesmus pubescens Wight & Arn. in Wight, Contr. Bot. India 63. 1834

Distribution: Karnataka (throughout), throughout India; endemic.

Heterostemma Wight & Arn.
Oianthus Benth.

Heterostemma beddomei (Hook.f.) Swarupan. & Mangaly, Bot. J. Linn. Soc. 101: 254. 1989
Oianthus beddomei Hook. f., Fl. Brit. India 4: 49. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.
 Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Heterostemma dalzellii Hook.f., Fl. Brit. India 4: 48.1883
 Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.

Heterostemma disciflorum (Hook.f.) Swarupan. & Mangaly, Bot. J. Linn. Soc. 101: 256. 1989
Oianthus disciflorus Hook.f., Fl. Brit. India 4: 49. 1883
 Distribution: Karnataka (Uttara Kannada), Andhra Pradesh; endemic.

Heterostemma urceolatum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 4: 295. 1852
Oianthus urceolatus (Dalzell) Benth. in Hooker's Icon. Pl. ser. 3, 12: t. 1191. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.
 Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Holarrhena R. Br.

Holarrhena pubescens (Buch.-Ham.) Wall. ex G. Don, Gen. Hist. 4: 78. 1837; Panigrahi, Taxon 36: 466. 1987.

Nerium antidysentericum L., Sp. pl. 209. 1753 nom. ambig.

Echites antidysenterica Roth in Roem. & Schult., Syst. veg. 4: 394. 1819 non (L.) Roxb. ex Fleming 1810 nec (L.) Roxb. 1814

Echites pubescens Buch.-Ham., Trans. Linn. Soc. London 13: 524. 1822

Chonemorpha antidysenterica (Roth) G.Don, Gen. Hist. 4: 76.1837

Holarrhena codaga G.Don, Gen. Hist. 4: 78. 1837

Holarrhena antidysenterica Wall. ex A.DC. in DC., Prodr. 8: 413. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.

Holarrhena malaccensis Wight, Icon. Pl. Ind. Orient. t. 1298. 1848

Elytropus pubescens (Wall.) Miers, Apocyn. S. Amer. 114.1878

Distribution: Karnataka (throughout), throughout India including Himalayas; Africa, Bangladesh, Cambodia, China, Laos, Myanmar, Nepal, Thailand, Vietnam.

Local/Common/Vernacular names: *Bitter Oleander*, *Coorg milk plant*, *Dysentry rosebay*, *Tellichery bark* (English); *Haale*, *Halagatti*, *Hire kanigu*, *Kadu soge*, *Kodacaga*, *Kodasigae*, *Kodu muruke*, *Korasigina gida*, *Korchu mara*, *Kudasolu*, *Kuichi*, *Kurchi mara*, *Sanna haale* (Kannada); *Kuda* (Kunhi); *Kutaja* (Sanskrit); *Kudenchipoo* (Tulu)

Holostemma R. Br.

Holostemma ada-kodien Schult., Syst. Veg. 6: 95. 1820; K.M. Matthew, Fl. Tamilnadu Carnatic 3: 949. 1983.

Sarcostemma annulare Roth, Nov. pl. sp. 178. 1821

Holostemma rheedianum Spreng., Syst. Veg. ed. 16. 1: 851.1825

Holostemma rheedei Wall., Pl. Asiat. Rar. 2: 51. 1831

Asclepias annularia Roxb., Fl. Ind. ed.2, 2: 37. 1832

Holostemma brunonianum Royle, Ill. Bot. Himal. Mts. 276, t. 66, f.2. 1835

Holostemma annulare (Roth) K. Schum. in Engl. & Prantl, Nat. Pflanzenfam. IV, 2: 250. 1895; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Distribution: Karnataka (Ballari, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; China, Myanmar, Nepal, Pakistan, Sri Lanka, Thailand.

Local/Common/Vernacular names: *Arne beelu, Jeeva haale, Jeeva hale balli, Jeevanthi, Marulingana kasa, Pala kire soppu, Shihale* (Kannada); *Araka pushpi, Durdharsha, Jalamuka, Jaleruha, Jeevanthi, Kroorakarma, Ksheeradodi, Ksheerini, Kutumbini, Praharkutavi, Sheetha, Sheethala, Sheethaparani, Shirinthica, Sithaparini, Suryavalli, Vakrashalya* (Sanskrit); *Arane booru* (Tulu)

Hoya R. Br.

Hoya carnosa (L.f.) R.Br., Asclepiadeae 16.1810

Asclepias carnosa L.f., Suppl. Pl. 170.1782

Schollia carnosa (L.f.) Schrank ex Steud., Nomencl. Bot. 414, 746.1821

Distribution: Karnataka (cultivated in gardens, Bengaluru), Australia, China, Japan, Malaysia, Vietnam.

Hoya ovalifolia Wight & Arn. in Wight, Contr. Bot. India 37. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Hoya parviflora Wight, Contr. Bot. India 37.1834

Hoya pauciflora Wight, Icon. Pl. Ind. Orient. t. 1269. 1848 ('parviflora'); B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Hoya wightiana Thwaites, Enum. Pl. Zeyl. 199.1860

Distribution: Karnataka (Hassan, Mysuru, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Hoya pendula Wight & Arn., Contr. Bot. India 36.1834

Asclepias pendula Roxb., Fl. Ind. ed. 1832 2: 36.1832

Hoya pallida Dalzell & A.Gibson, Bombay Fl. 152.1861

Hoya wightii Hook.f., Fl. Brit. India 4: 59. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kalaburgi, Kodagu, Mysuru, Shivamogga, Uttara Kannada, Yadgir), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Ambri, Pipi balli* (Kannada); *Benduli* (Kunhi)

Hoya retusa Dalzell in Hooker's J. Bot. Kew Gard. Misc. 4: 294. 1854; B.D.Sharma et al., Fl. Karnataka Anal. 167. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Local/Common/Vernacular name: *Bendurki* (Kannada)

Ichnocarpus R. Br.

Ichnocarpus frutescens (L.) R. Br. in W. T. Aiton, Hortus kew. ed 2. 2: 69. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Apocynum frutescens L., Sp. pl. 213. 1753

Echites malabarica Lam., Dict. encycl. 2: 342. 1790

Quirivelia frutescens (L.) R. Br., Mem. Wern. Nat. Hist. Soc. 1: 61. 1811

Echites frutescens (L.) Roxb., Hort. Bengal. 20.1814

Chonemorpha malabarica (Lam.) G. Don, Gen. Hist. 4: 76. 1837

Beluttakaka malabarica (Lam.) Kuntze, Revis. Gen. Pl. 2: 413.1891

Distribution: Karnataka (throughout), throughout hotter regions of India; Australia, S & S E Asia.

Local/Common/Vernacular names: *Black creeper, Dark blue creeper* (English); *Gauri balli, Gopi, Haalu genasu, Kare anantha moola, Kare balli, Kare bantana balli, Kari hambu, Koogal balli, Malvi balli* (Kannada); *Anntha, Bhadra, Chnandana, Chandana gopa, Chandan saariva, Chinha dharani, Dheerga moola, Dridha bandhini, Gopa, Gopalli, Gopa vadhu, Gopini, Kala ghnteika, Kala peshi, Krisna, Krishna mooli, Krishna saariva Krishna valli, Maha shyama, Masura vindala, Palindi saariva, Shariva, Shyama, Shyamalata, Subhadra, Utpla saariva* (Sanskrit); *Illa booru* (Tulu)

Kamettia Kostel.

Kamettia caryophyllata (Roxb.) Nicolson & Suresh, Taxon 35: 354. 1986

Echites caryophyllata Roxb., Hort. Bengal. 20. 1814 ('coryophyllata').

Kamettia malabarica Kostel., Allg. Med.-Pharm. Fl. 3: 1062. 1834

Aganosma roxburghii G.Don, Gen. Hist. 4: 77. 1837

Ellertonia rheedei Wight, Icon. Pl. Ind. Orient. t. 1295. 1848 ('rheedii') nom. illeg.; B.D.Sharma et al., Fl. Karnataka Anal. 162. 1984.

Distribution: Karnataka (Chikkamagaluru, Uttara kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Odisha, Tamil Nadu; endemic.

Kopsia Blume

Kopsia fruticosa (Roxb.) A. DC., Prodr. 8: 352. 1844

Cerbera fruticosa Roxb., Bot. Reg. 5: t. 391. 1819

Kopsia vinciflora Blume, Bijdr. 1030. 1826

Calpicarpum roxburghii G.Don, Gen. Hist. 4: 100. 1837

Kopsia roxburghii (G.Don) Pharm. ex Wehmer, Pfl.-Stoffe 625. 1911

Distribution: Karnataka (cultivated as an ornamental in Bengaluru, Dakshina Kannada, Mysuru, Udupi), S & SE Asia; native to Myanmar.

Local/Common/Vernacular names: *Kopsia Merah*, *Pink Gardenia*, *Pink Kopsia*, *Shrub Vinca* (English)

Leptadenia R. Br.

Leptadenia pyrotechnica (Forssk.) Decne., Ann. Sci. Nat. Bot. ser. 2, 9: 270. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Cynanchum pyrotechnicum Forssk., Fl. Aegypt.-Arab. 53. 1775

Leptadenia spartium Wight, Contr. Bot. India 48. 1834

Distribution: Karnataka (Ballari), throughout India; Iran, Pakistan.

Leptadenia reticulata (Retz.) Wight & Arn. in Wight, Cat. Indian Pl. 89. 1834 & Contr. Bot. India 47. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Cynanchum reticulatum Retz., Observ. Bot. 2:15. 1781

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalaburgi, Mysuru, Raichur, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout India; Madagascar, Mauritius, Myanmar, Nepal, Singapore, Sri Lanka.

Local/Common/Vernacular names: *Hali balli*, *Pla thige balli* (Kannada)

Marsdenia R. Br.

Marsdenia floribunda (Brongn.) Schltr. in Urb., Symb. Antill. 1: 275. 1899

Stephanotis floribunda Brongn., Ann. Sci. Nat. Bot. ser. 2, 7: 30. 1837

Distribution: Karnataka (Cultivated & naturaliz), Cultivated & naturalized; native to Madagascar.

Marsdenia raziana Yogan. & Subr., Proc. Indian Acad. Sci. 83: 147. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannad, Udupi), Kerala; endemic.

Marsdenia tenacissima (Roxb.) Moon, Cat. Pl. Ceylon 21. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Asclepias tenacissima Roxb., Pl. Coromandel 3: 35, t. 240. 1815

Distribution: Karnataka (Kalaburgi), Peninsular to N & NE India, W Himalaya; Cambodia, China, Laos, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam].

Nerium L.

Nerium oleander L., Sp. pl. 209. 1753; Nicolson et al., Regnum Veg. 119 (Interpr. Rheede's Hort. malab.): 56. 1988.

Nerium indicum Mill., Gard. dict. ed. 8, no.2. 1768; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Nerium odoratum Sol. ex Aiton, Hort. Kew. ed. 1. 1: 297. 1789

Distribution: Karnataka (cultivated throughout) cultivated throughout India including Islands; native to Mediterranean region.

Local/Common/Vernacular names: *Kanagalu, Kanagile* (Kannada)

Oxystelma R. Br.

Oxystelma esculentum (L.f.) Sm. in Rees, Cycl. 1: 25. 1813

Periploca esculenta L.f., Suppl. Pl. 168. 1782

Oxystelma secamone (L.) H. Karst., Deut. Fl. 1031. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada), throughout plains & lower hills of India; Indoneia, Iran, Iraq, Nepal, Pakistan, Sri Lanka.

Parsonia R. Br.

Parsonia alboflavescens (Dennst.) Mabb., Taxon 26: 532. 1977; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Periploca alboflavescens Dennst., Schlüssel Hortus Malab. 12, 23. 1818

Echites laevigatus Moon, Cat. Pl. Ceylon 20. 1824

Parsonia helicandra Hook. & Arn., Bot. Beechey Voy. 197. 1837

Parsonia spiralis Wall. ex G. Don, Gen. Hist. 4: 80. 1837

Periploca spiralis Wall. ex G. Don, Gen. hist. 4: 80. 1837

Aganosma laevigata (Moon) J. Graham, Cat. Pl. Bombay 113. 1839

Heligme rheedei Wight, Icon. Pl. Ind. Orient. t. 1303. 1848

Aganosma concanensis Hook., Hooker's Icon. Pl. 9: t. 841. 1852

Spirostemon spiralis Griff., Not. Pl. Asiat. 4: 80. 1854

Parsonia laevigata (Moon) Alston, Ann. Roy. Bot. Gard. (Peradeniya) 11: 203. 1929

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, E, NE & Peninsular India; tropical Asia.

Pentatropis Wight & Arn.

Pentatropis capensis (L.f.) Bullock, Kew Bull. 1955: 284. 1955; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Cynanchum capense L.f., Suppl. pl. 168. 1782

Asclepias microphylla Roxb., Fl. Ind. (Ed. Carey) 2: 35. 1832

Pentatropis microphylla (Roxb.) Wight & Arn. in Wight, Contr. Bot. India 52. 1834; Wight, Icon. pl. Ind. orient. t. 352. 1840; Hook.f., Fl. Brit. India 4: 20. 1883.

Distribution: Karnataka (Ballari, Chitradurga, Dharwad, Kalaburgi, Mysuru, Raichur, Tumakuru, Vijayapura), throughout India, Iran, Pakistan, Sri Lanka.

Pergularia L.

Pergularia brunonianiana (Wight & Arn.) D. Dietr., Syn. Pl. 2: 894. 1840

Marsdenia brunonianiana Wight & Arn. in Wight. Contr. Bot. India 40. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Distribution: Karnataka (Dakshina Kannada, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Pergularia daemia (Forssk.) Chiov., Res. Sci. Somalia Ital. 1:115. 1916; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Asclepias daemia Forssk., Fl. Aegypt.-Arab. 51. 1775

Cynanchum extensum Jacq., Misc. Austriac. 2: 353. 1781

Daemia extensa (Jacq.) R. Br. ex Schult., Syst. Veg. 6: 112. 1820

Pergularia extensa (Jacq.) N.E. Br. in Harv., Fl. Cap. 4(1): 758. 1908

Distribution: Karnataka (throughout), throughout India including tropical Himalayas; Afghanistan, Iran, Middle East, S Asia, tropical Africa.

Local/Common/Vernacular names: *Halu kortiga, Kuntaga, Kurigana balli, Thalavarana balli, Uguru suttu balli* (Kannada)

Plumeria L.

Plumeria alba L., Sp. Pl. 210. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Distribution: Karnataka (cultivated as an ornamental throughout), cultivated in drier parts of India; native to C America & the Caribbean.

Local/Common/Vernacular name: *Deva Kanigalu* (Kannada)

Plumeria obtusa L., Sp. Pl. 210. 1753

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout India; native to C America & the Caribbean.

Plumeria rubra L., Sp. Pl. 209. 1753; Maheshw., Fl. Delhi 214. 1963.

Plumeria acuminata W.T. Aiton, Hortus kew. ed. 2. 2: 70. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Plumeria acutifolia Poir. in Lam., Encycl. suppl. 2: 667. 1812

Plumeria rubra var. *acuminata* (W.T.Aiton) R.S. Rao & Balamani, J. Indian Bot. Soc. 63: 227. 1984; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Distribution: Karnataka (cultivated as an ornamental throughout), cultivated in drier parts of India; native to tropical America.

Local/Common/Vernacular names: *Pagoda tree* (English); *Devakanigilu, Donne huvu, Gosampige, Jeeva kanegilu* (Kannada)

Rauvolfia L.

Rauvolfia serpentina (L.) Benth. ex Kurz, Forest Fl. Burma 2: 171. 1877; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Ophioxylon serpentinum L., Sp. Pl. 1043. 1753

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Shivamogga, udupi, Uttara Kannada, elsewhere cultivated), throughout tropical India; S & SE Asia.

Local/Common/Vernacular names: *Garuda paathaala, Sarpagandha* (Kannada); *Adki* (Kunhi)

Rauvolfia tetraphylla L., Sp. Pl. 208. 1753; A.S.Rao, Ann. Missouri Bot. Gard. 43: 285. 1956.

Rauvolfia hirsuta Jacq., Enum. Syst. Pl. 14. 1760

Rauvolfia tomentosa Jacq., Enum. Syst. Pl. 14. 1760

Rauvolfia canescens L., Sp. Pl. ed. 2. 303. 1762

Rauvolfia subpubescens L., Mant. Pl. 2: 345. 1771

Distribution: Karnataka (cultivated & naturalized throughout), cultivated & naturalized throughout India; native to West Indies.

Local/Common/Vernacular name: *Dodda Sarpag & ha* (Kannada)

Rauvolfia verticillata (Lour.) Baill., Mens. Soc. Linn. Paris 1: 768. 1888 & Hist. Pl. 13: 170 1895; Bull; Markgr., Blumea 30: 160. 1984.

Dissolena verticillata Lour., Fl. Cochinch. 1: 137. 1790

Tabernaemontana densiflora Wall. in Edwards' Bot. Reg. (Ed. Lindl.) 15: t. 1273, p.p. 1829

Ophioxylon Belagaviense Wight, Icon. Pl. Ind. Orient. 4(2): 2. 1848

Ophioxylon macrocarpum Wight, Icon. Pl. Ind. Orient. 4(2): 1. 1848

Ophioxylon neilgherrense Wight, Icon. pl. Ind. orient. 4(2): t. 1292. 1848

Ophioxylon zeylanicum Wight, Icon. pl. Ind. orient. 4(2): t. 1291. 1848

Rauvolfia decurva Hook.f., Fl. Brit. India 3: 633. 1882; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Rauvolfia densiflora (Wall.) Benth. ex Hook.f., Fl. Brit. India 3: 633. 1882; B.D.Sharma et al., l. Karnataka Anal. 163. 1984.

Rauvolfia densiflora var. *macrophylla* Hook.f., Fl. Brit. India 3: 633. 1882

Tabernaemontana subcapitata Hook.f. & Thomson, Fl. Brit. India 3: 633. 1882

Rauvolfia verticillata var. *macrophylla* (Hook.f.) Karthik. & Moorthy, Fl. Pl. India 137. 2009

Distribution: Karnataka (Belagavi, Davanagere, Kodagu, Mysuru, Shivamogga), Peninsular to NE India; S & SE Asia.

Local/Common/Vernacular name: *Pathala garudi* (Kannada); *Kudgi* (Kunhi)

Sarcostemma R. Br.

Sarcostemma brunonianum Wight & Arn. in Wight, Contr. Bot. India 59. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 169. 1984.

Sarcostemma viminalis subsp. *brunonianum* (Wight & Arn.) P.I. Forst., Austral. Syst. Bot. 5: 63. 1992

Distribution: Karnataka (Chitradurga, Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Sarcostemma intermedium Decne. in DC., Prodr. 8: 538. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 169. 1984.

Distribution: Karnataka (Chikkamagaluru, Dharwad, Hassan), Peninsular to N India; endemic.

Sarcostemma secamone (L.) Bennett, Indian Forester 95: 692. 1969

Periploca secamone L., Mant. Pl. 2: 216. 1771

Distribution: Karnataka (Chamarajanagar, Chitradurga, Mysuru), Tamil Nadu; endemic.

Local/Common/Vernacular names: *caustic bushes, climbing milkweeds* (English)

Sarcostemma stocksii Hook. f., Fl. Brit. India 4: 27. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 169. 1984.

Distribution: Karnataka (Dharwad, Gadag, Vijayapura), Gujarat, Maharashtra, Rajasthan; Iran, Pakistan.

Sarcostemma viminalis non R. Br., 1810; Wight & Arn. in Wight, Contr. Bot. India 59. 1834.

Euphorbia viminalis L., Sp. Pl. 452. 1753

Asclepias acida Roxb., Fl. Ind. ed.2, 2: 31. 1832

Sarcostemma brevistigma Wight & Arn. in Wight, Contr. Bot. India 59. 1834

Sarcostemma acidum (Roxb.) Voigt, Hort. Suburb. Calcutt. 542. 1845; B.D.Sharma et al., Fl. Karnataka Anal. 168. 1984.

Distribution: Karnataka (Belagavi, Ballari, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburgi, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Vijayapura, W Bengal, Yadgir), Peninsular to N & E India; Myanmar, Nepal, Pakistan, tropical Africa.

Local/Common/Vernacular names: *Hambu kalli, Konana karalu, Soma balli, Somalate, Vasukanthi* (Kannada)

Seshagiria Ansari & Hemadri

Seshagiria sahyadrica Ansari & Hemadri, Indian Forester 97: 126. 1971; M. Radh. Rao, Yogan., S.N. Ramaswamy & Kesh.Murthy in High. Pl. Indian Subcont. 1: 19. 1990. Plate 16C
 Distribution: Karnataka (Shivamogga), Maharashtra, Tamil Nadu; endemic.

Stapelia L.

Stapelia gigantea N. E. Br. in Gard. Chron. 1: 684, f. 112. 1877; S. R. Sriniv. in A. N. Henry et al., Fl. Tamil Nadu Anal. 2: 88. 1987.

Distribution: Karnataka (cultivated in Bengaluru), cultivated throughout India; native to Africa.

Stapelia grandiflora Masson, Stapel. nov. t. 11. 1797; Chandrab. & N.C. Nair, Fl. Coimbatore 179. 1988.

Distribution: Karnataka (cultivated in Bengaluru), cultivated throughout India; native to Africa.

Strophanthus DC.

Strophanthus wallichii A.DC. in DC., Prodr. 8: 418. 1844; Hook.f., Fl. Brit. India 3: 655. 1882.

Nerium caudatum Roxb., Fl. Ind. ed. 2, 2: 9. 1832

Distribution: Karnataka (cultivated in Bengaluru), Andaman & Nicobar Islands, C, E, NE & S India; Bangladesh, China, Laos, Malaysia, Thailand, Vietnam.

Strophanthus wightianus Wight, Icon. Pl. Ind. Orient. 4: t. 1301. 1848; Hook.f., Fl. Brit. India 3: 656. 1882.

Distribution: Karnataka (), Kerala, Tamil Nadu; endemic.

Tabernaemontana L.

Tabernaemontana alternifolia L., Sp. Pl. 211.1753

Plate 16D

Tabernaemontana crispa Roxb., Fl. Ind. 2: 24. 1824

Tabernaemontana heyneana Wall. in Edwards' Bot. Reg. 15:t.1273. 1829

Tabernaemontana oblonga Wall., Edwards's Bot. Reg. 15:t. 1273.1829

Ervatamia heyneana (Wall.) T. Cooke, Fl. Bombay 2: 134.1904

Pagiantha heyneana (Wall.) Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 546 1935

Pagiantha crispa (Roxb.) Markgr., Mitt. Bot. Staatssamml. München 1: 29.1950

Ervatamia alternifolia (L.) S.M. Almeida, J. Econ. Taxon. Bot. addl. ser. 8(1): 252. 1990

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Kalaburgi, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Maddarasa gida*, *Maddarasara*, *Maddle mara*, *Nagar kuda*, *Nanja battlu hoovu* (Kannada); *Nagkuda* (Kunhi); *Kokke kayi* (Tulu)

Tabernaemontana citrifolia L., Sp. Pl. 210.1753; Nilpure, Fl. Belagavi Dist.

Distribution: Karnataka (cultivated in Belagavi), cultivated in S India; native to West Indies.

Tabernaemontana dichotoma Roxb. ex Wall., Edwards's Bot. Reg. 15: t. 1273.1829

Tanghinia dichotoma (Roxb. ex Wall.) G.Don, Hort. Brit. ed. 3: 461.1839

Rejoua dichotoma (Roxb.) Gamble, Fl. Madras 812. 1923

Ervatamia dichotoma (Roxb. ex Wall.) Burkil, Bull. Misc. Inform. Kew 1935: 317. 1935

Pagiantha dichotoma (Roxb.) Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 546. 1935

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Shivamogga, Uttara Kannada), Kerala, Tamil nadu; Malaysia, Myanmar, Sri Lanka, Taiwan.

Tabernaemontana divaricata (L.) R. Br. ex Roem. & Schult., Syst. veg. 4: 427. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.
Nerium divaricatum L., Sp. Pl. 209. 1753
Nerium coronarium Jacq., Collectanea 1: 138. 1786
Tabernaemontana coronaria (Jacq.) Willd., Enum. Pl. 275. 1809
Tabernaemontana recurva Roxb., Hort. Bengal. 20. 1814
Tabernaemontana coronaria var. *plena* Roxb. ex Voigt, Hort. Suburb. Calcutt. 527. 1845
Ervatamia coronaria (Jacq.) Stapf in Oliv., Fl. Trop. Afr. 4: 127. 1902
Ervatamia recurva (Roxb. ex Lindl.) Lace, List Trees Shrubs Cl. Burma 91. 1913
Ervatamia divaricata (L.) Burkill, Rec. Bot. Surv. India 10: 320. 1925
Testudipes recurva (Roxb. ex Lindl.) Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 547. 1935
Ervatamia divaricata var. *plena* (Roxb. ex Voigt) M.R.Almeida, Fl. Maharashtra 3A: 207. 2001
 Distribution: Karnataka (cultivated throughout), cultivated throughout drier parts of India; Bangladesh, Bhutan, China, Myanmar, Nepal, Taiwan, Thailand.
 Local/Common/Vernacular name: *Nandibattlu* (Kannada); *Nandyavarath* (Sanskrit)

Tabernaemontana gamblei Subr. & A.N. Henry, Bull. Bot. Surv. India 12: 1. 1970; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Ervatamia caudata Gamble, Bull. Misc. Inform. 1921: 310. 1921 & Fl. Madras 813. 1923 non *Tabernaemontana caudata* Merr., 1909
 Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Telosma Coville

Telosma cordata (Burm.f.) Merr., Philipp. J. Sci. 19:372. 1921; Deb, Fl. Tripura 2: 36. 1983.

Asclepias cordata Burm. f., Fl. indica 72, t. 27. 1768

Pergularia minor Andrews, Bot. repos. t. 184. 1801; Hook. f., Fl. Brit. India 4:38.

Asclepias odoratissima Roxb., Fl. Ind. (Ed. Carey) 2: 46. 1832

Pergularia odoratissima (Roxb.) Wight, Contr. bot. India 43. 1834 & Icon. pl. Ind. orient. t. 414. 1840

Telosma minor (Andrews) Craib in Bull. Misc. Inform. 1911: 418. 1911

Distribution: Karnataka (cultivated or wild), cultivated or wild throughout India; China, Europe, Myanmar, N & S America, Pakistan, Vietnam.

Telosma pallida (Roxb.) Craib, Bull. Misc. Inform. 1911: 418. 1911; B.D.Sharma et al., Fl. Karnataka Anal. 169. 1984.

Asclepias pallida Roxb., Fl. Ind. ed. 2, 2: 48. 1832

Pergularia pallida (Roxb.) Wight & Arn. in Wight, Contr. Bot. India 42. 1834; Wight, Icon. Pl. Ind. Orient. t. 585. 1842; Hook.f., Fl. Brit. India 4: 38. 1883.

Distribution: Karnataka (Ballari, Dharwad, Shivamogga), throughout India including Himalayas; China, Myanmar, Nepal, Pakistan, Taiwan, Thailand, Vietnam.

Thevetia L. Nom. Cons.

Thevetia peruviana (Pers.) Merr., Philipp. J. Sci. 9: 130. 1914; B.D.Sharma et al., Fl. Karnataka Anal. 163. 1984.

Cerbera thevetia L., Sp. Pl. 209. 1753

Cerbera peruviana Pers., Syn. Pl. 1: 267. 1805

Thevetia neriifolia Juss. ex Steud., Nomencl. Bot. ed. 2, 2: 680. 1841 ('nereifolia').

Cascabela thevetia (L.) Lippold, Feddes Repert. 91: 52. 1980

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to C America & Mexico .

Local/Common/Vernacular names: *Haladi kanigilu*, *Kaadu kasi kanigalu*, *Kaaraveera* (Kannada)

Toxocarpus Wight & Arn.

Toxocarpus concanensis Hook. f., Fl. Brit. India 4: 14. 1883

Distribution: Karnataka (Chikkamagaluru), Goa, Maharashtra; endemic.

Toxocarpus kleinii Wight & Arn. in Wight, Contr. Bot. India 61. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 169. 1984.

Distribution: Karnataka (Hassan, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Peninsular to E India; Sri Lanka.

Tylophora R. Br.

Tylophora capparidifolia Wight & Arn. in Wight, Contr. Bot. India 51. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 169. 1984.

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Tylophora dalzellii Hook. f., Fl. Brit. India 4: 43. 1883; B.D.Sharma, Fl. Karnataka Anal. 169. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; endemic.

Tylophora fasciculata Buch.-Ham. ex Wight, Contr. Bot. India 50. 1834; Wight, Icon. Pl. Ind. Orient. t. 848. 1844 - 45 B.D.Sharma, Fl. Karnataka Anal. 169. 1984.

Pentasachme fasciculata (Buch.-Ham. ex Wight) M.R. Almeida, Fl. Maharashtra 3 (A & B): 253. 2001 ('Pentasacme').

Distribution: Karnataka (Belagavi, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; Nepal, Sri Lanka.

Tylophora flexuosa R. Br., Prodr. 460. 1810

Asclepias tetrapetala Dennst., Schlüssel Hortus Malab. 14, 18, 35. 1818

Asclepias tenuissima Roxb. ex Willd. in Schult., Syst. Veg. 6: 86. 1820

Tylophora tenuis Blume, Bijdr. 1062. 1827; B.D.Sharma, Fl. Karnataka Anal. 170. 1984

Tylophora tenuissima (Roxb. ex Willd.) Wight & Arn. in Wight, Contr. Bot. India 49. 1834

Tylophora carnosa auct. non Wall. ex Wight 1834. Dalzell & A.Gibson, Bombay Fl. 150. 1861

Vincetoxicum flexuosum (R.Br.) Kuntze, Revis. Gen. Pl. 2: 424. 1891

Tylophora tetrapetala (Dennst.) Suresh in Nicolson et al., Regnum Veg. 119 (Interpr. Rheede's Hort. malab.): 64. 1988

Tylophora tetrapetala var. *tenuissima* (Roxb. ex Willd.) Swarupan., Blumea 38: 234. 1993

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Udupi), Andaman & Nicobar Islands, Peninsular to NE India; SE Asia, Sri Lanka.

Tylophora indica (Burm.f.) Merr., Philipp. J. Sci. 19: 373. 1921; B.D.Sharma, Fl. Karnataka Anal. 169. 1984

Cynanchum indicum Burm. f., Fl. indica 70. 1768

Asclepias asthmatica L.f., Suppl. pl. 171. 1782

Asclepias prolifera Rottler ex Ainslie, Materia Ind. 2: 225. 1826

Tylophora asthmatica (L.f.) Wight & Arn. in Wight, Contr. Bot. India 51. 1834

var. indica

Distribution: Karnataka (throughout), throughout India including Andaman & Nicobar Islands; S & SE Asia.

Local/Common/Vernacular names: *Aadu muttada balli*, *Aadu muttada gida*, *Aadu muttada soppu*, *Antamula*, *Kiru manji balli* (Kannada)

var. glabra (Decne.) H. Huber in Abeyw., Rev. Handb. Fl. Ceylon 1: 43. 1973; A.P.Jagtap & N.P.Singh, Fasc. Fl. India 24: 161. 1999.

Tylophora asthmatica var. *glabra* Decne. in DC., Prodr. 8: 611. 1844 p.p. emend. Thwaites, Enum. Pl. Zeyl. 197. 1860

Distribution: Karnataka (Kolar), Kerala; Mauritius, Sri Lanka.

Tylophora iphisia Decne., Prodr. 8: 610.1844; Bhaskar & Kushalappa, Fl. Tumkur Dist. 65. 2013.
Distribution: Karnataka (Tumakuru), Tamil Nadu; endemic.

Tylophora multiflora (Wight & Arn.) Alston, Handb. Fl. Ceylon 6: 195.1931
Iphisia multiflora Wight & Arn., Contr. Bot. India 52.1834
Distribution: Karnataka (Dakshina Kannada), Tamil Nadu; Sri Lanka.

Tylophora pauciflora Wight & Arn. in Wight, Contr. Bot. India 49. 1834; B.D.Sharma, Fl. Karnataka Anal. 169. 1984
Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi), S to E India; Sri Lanka.

Tylophora rotundifolia Buch.-Ham. ex Wight, Contr. Bot. India 50. 1834; B.D.Sharma, Fl. Karnataka Anal. 169. 1984
Distribution: Karnataka (Davanagere, Hassan, Mysuru, Uttara Kannada), Peninsular to N & E India; Nepal.

Vallaris Burm. f.

Vallaris solanacea (Roth) Kuntze, Revis. Gen. Pl. 1: 147. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Peltanthera solanacea Roth, Nov. Pl. Sp. 132. 1821

Vallaris heynei Spreng., Syst. Veg. ed. 16. 1: 635.1825

Echites dichotomus Roxb., Fl. Ind. ed.2,2: 19.1832

Vallaris dichotoma (Roxb.) Wall. ex G.Don, Gen. Hist. 4: 79.1837

Vallaris assamensis Griff., Not. Pl. Asiat. 4: 77.1854

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Mysuru, Udupi, Uttara Kannada), cultivated throughout India; Cambodia, China, Indonesia, Laos, Myanmar, Pakistan, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular name: *Bugudi hambu* (Kannada)

Vinca L.

Vinca major L., Sp. pl. 209. 1753

Distribution: Karnataka (Cultivated in gardens), cultivated throughout tropical areas and also runs wild; native to Mediterranean region.

Local/Common/Vernacular names: *Bigleaf periwinkle, Blue periwinkle, greater periwinkle, Large periwinkle* (English)

Wrightia R. Br.

Wrightia arborea (Dennst.) Mabb., Taxon 26: 533. 1977

Periploca arborea Dennst., Schlüssel Hortus malab. 13, 23 & 25. 1818 & Fortsetz. Allg. Teutsch. Gart.-Mag. 3: 32, 41 & 83. 1818

Wrightia tomentosa Roem. & Schult., Syst. Veg. 4: 414. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Echites tomentosus Roth, Nov. Pl. Sp. 137.1821

Wrightia pubescens Roth, Nov. Pl. Sp. 120.1821

Wrightia mollissima Wall., Pl. Asiat. Rar.2: 39, t. 146. 1831

Nerium tomentosum (Roem. & Schult.) Roxb., Fl. Ind. 2, 2: 6 1832

Wrightia rheedei Kostel., Allg. Med.-Pharm. Fl. 3: 1060.1834

Wrightia coalita Buch.-Ham. ex Dillwyn, Rev. Hortus Malab. 40. 1839, nom. illeg. incl. type of *Periploca arborea* Dennst., 1818

Nerium coraea Buch.-Ham. ex A.DC., Prodr. 8: 407.1844

Wrightia wallichii A.DC. in DC., Prodr. 8: 405. 1844

Beaumontia wallichii (A.DC.) Walp., Ann. Bot. Syst. 3: 39.1852

Distribution: Karnataka (Belagavi Dakshina Kannada, Davanagere, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout tropical areas; China, Laos, Malaysia, Myanmar, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Benki gida, Biliganagal, Bilikada, Kaadu Kanagilu* (Kannada)

Wrightia tinctoria (Roxb.) R. Br., Asclepiadaceae 59.1810 & Mem. Wern. Nat. Hist. Soc. 1: 74. 1811;
B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.
Nerium tinctorium Roxb., Orient. Repert. 1: 39. 1791

subsp. *tinctoria*

Distribution: Karnataka (throughout), throughout tropical India; Myanmar, Sri Lanka, Timor.
Local/Common/Vernacular names: *Beppale mara*, *Hale*, *Hale mara*, *Kare kodasiga*, *Kodamurki*, *kodasige*,
Kodurka, *Nili*, *Pale* (Kannada); *Kala kuda* (Kunhi); *Krishna Kutaja* (Sanskrit); *Kooli bele thappu* (Tulu)

subsp. *rothii* (G.Don) Ngan, Ann. Missouri Bot. Gard. 52: 144.1965

Wrightia rothii G.Don, Gen. Hist. 4: 86. 1837

Wrightia tinctoria var. *rothii* (G.Don) Hook.f., Fl. Brit. India 3: 653. 1882; B.D.Sharma et al., Fl. Karnataka Anal. 164. 1984.

Distribution: Karnataka (Ballari, Bidar, Kalaburgi, Vijayapura), throughout India from Himalayas downwards excluding NE & NW India.

Aponogetonaceae

Aponogeton L. f.

Aponogeton crispus Thunb., Nov. Gen. Pl. 73.1784

Aponogeton echinatus Roxb., Fl. Ind. ed.2, 2: 210.1832; B.D.Sharma et al., Fl. Karnataka Anal. 299. 1984.

Spathium crispum (Thunb.) Voigt, Hort. Suburb. Calcutt. 694.1845

Spathium echinatum (Roxb.) Voigt, Hort. Suburb. Calcutt. 694.1845

Distribution: Karnataka (Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru), Maharashtra, Tamil Nadu; Sri Lanka.

Aponogeton natans (L.) Engl. & K.Krause, Pflanzenr. IV, 13: 11.1906; B.D.Sharma et al., Fl. Karnataka Anal. 299. 1984.

Saururus natans L., Mant. Pl. 227.1771

Aponogeton monostachyon L.f., Suppl. Pl. 214.1782

Potamogeton indicus Roth ex Roem. & Schult., Syst. Veg. 3: 516.1818

Spathium monostachyum (L.f.) Edgew., J. Asiat. Soc. Bengal part 2, Nat. Hist. 11: 148.1842

Aponogeton flavidum Ham. ex Hook.f., Fl. Brit. India 6: 564.1893

Distribution: Karnataka (throughout in tanks), throughout plains in India; Kerala; Australia, tropical Asia.

Local/Common/Vernacular name: *Kothi gedde* (Kannada)

Aponogeton nateshii S.R.Yadav, Rheedia 25(1): 9. 2015.

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Aponogeton undulatus Roxb., Fl. Ind. ed.2, 2: 211.1832 ('undulatum').

Aponogeton microphyllus Roxb., Fl. Ind. ed. 2, 2: 211.1832('microphyllum').

Spathium undulatum (Roxb.) Edgew., J. Asiat. Soc. Bengal 11: 145.1842

Ouvirandra undulata (Roxb.) Edgew., London J. Bot. 3: 405.1844

Spathium microphyllum (Roxb.) Voigt, Hort. Suburb. Calcutt. 694.1845

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi), throughout India; Bangladesh, Myanmar, Sri Lanka; cultivated elsewhere as aquarium plant.

Aquifoliaceae

Ilex L.

Ilex denticulata Wall. ex Wight, Illustr. 2: t. 142. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Distribution: Karnataka (Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; China, Sri Lanka.

Ilex malabarica Bedd., Fl. Sylv. t. 143. 1871; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Kodagu, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Ilex tadiandamolensis Kesh.Murthy, Yogan. & Vasud.Nair, Curr. Sci. 56: 270. 1987; C.J.Saldanha, Fl. Karnataka 2: 103. 1996

Distribution: Karnataka (Kodagu); endemic.

Araceae

Aglaonema Schott

Aglaonema commutatum Schott, Syn. Aroid. 123.1856

Aglaonema marantifolium var. *commutatum* (Schott) Engl., Monogr. Phan. 2: 441–442. 1879

Aglaonema robustum Alderw., Bull. Jard. Bot. Buitenzorg III, 4: 328.1922

Aglaonema commutatum var. *robustum* (Alderw.) Nicolson, Baileya 15: 125.1968

Distribution: Karnataka (cultivated as an ornamental)cultivated as an ornamental throughout India; native to Celebes & Philippines.

Local/Common/Vernacular names: *Aglaonema, Chinese evergreen, Poison dart plant* (English)

Alocasia Schott

Alocasia cucullata (Lour.) G.Don, Hort. Brit. ed. 3: 631.1839

Arum cucullatum Lour., Fl. Cochinch. 536.1790

Caladium cucullatum (Lour.) Pers., Syn. Pl. 2: 575.1807

Colocasia cucullata (Lour.) Schott, Melet. Bot. 18.1832

Caladium colocasia Schott ex Wight, Icon. Pl. Ind. Orient. 3: t. 787.1844

Distribution: Karnataka (cultivated as an ornamental)cultivated as an ornamental throughout India; Bangladesh, China, Laos, Myanmar, Nepal, Sri Lanka, Taiwan, Thailand, Vietnam.

Local/Common/Vernacular names: *Buddha's hand, Chinese ape, Chinese taro, elephant ear, Hooded dwarf* (English)

Alocasia fornicata (Roxb.) Schott, Oesterr. Bot. Wochensbl. 4: 410.1854; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Arum fornicatum Roxb., Fl. Ind. ed. 2, 3: 501.1832

Colocasia fornicata (Roxb.) Kunth, Enum. Pl. 3: 41.1841

Distribution: Karnataka (Hassan, Kalburgi, Yadgir), S to NE India; Bangladesh, Sri Lanka.

Alocasia macrorrhizos (L.) G.Don, Hort. Brit. ed. 3: 631.1839

Arum macrorrhizon L., Sp. Pl. 965.1753

Arum macrorrhizum L., Sp. Pl. 2: 965.1753

Arum peregrinum L., Sp. Pl. 966.1753

Arum indicum Lour., Fl. Cochinch. 536.1790

Caladium macrorrhizon (L.) R.Br., Prodr. Fl. Nov. Holl. 336.1810

Arum montanum Roxb., Fl. Ind. ed.2, 3: 497.1832

Arum rapiforme Roxb., Fl. Ind. ed. 2, 3: 497.1832

Colocasia macrorrhizos (L.) Schott, Melet. Bot. 18.1832

Colocasia indica (Lour.) Kunth, Enum. Pl. 3: 39.1841

Colocasia montana (Roxb.) Kunth, Enum. Pl. 3: 40.1841

Colocasia rapiformis (Roxb.) Kunth, Enum. Pl. 3: 40.1841

Alocasia indica (Lour.) Schott, Oesterr. Bot. Wochensbl. 4: 410.1854

Alocasia rapiformis (Roxb.) Schott, Prodr. Syst. Aroid. 157.1860

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), Peninsular India; Indonesia, Malaysia, Sri Lanka.

Local/Common/Vernacular names: *Marsanege, Mundi gedde* (Kannada)

Alocasia sanderiana W.Bull, Cat. 1894: 8.1894

Schizocasia sanderiana (W.Bull) Engl., Bot. Jahrb. Syst. 25: 26.1898

Distribution: Karnataka (cultivated as an ornamental) cultivated as an ornamental throughout India; native to Philippines.

Local/Common/Vernacular name: *Kris plant* (English)

Amorphophallus Blume ex Decne.

Amorphophallus bulbifer (Roxb.) Blume, Rumphia 1: 148.1837; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Arum bulbiferum Roxb., Fl. Ind. ed. 2, 3: 510.1832

Pythonium bulbiferum (Roxb.) Schott, Melet. Bot. 18.1832

Conophallus bulbifer (Roxb.) Schott., Syn. Aroid. 34.1856

Amorphophallus aculatum Hook.f., Fl. Brit. India 6: 515.1893

Amorphophallus taccoides Hook.f., Fl. Brit. India 6: 515.1893

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Hassan, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Peninsular to NE India; Bangladesh, Myanmar.

Local/Common/Vernacular name: *Kadu suvarna gedde* (Kannada); *Lutiche* (Kunabi)

Amorphophallus commutatus (Schott) Engl., Monogr. Phan. 2: 319.1879

Plate 16E

Conophallus commutatus Schott, Bonplandia (Hannover) 7: 28.1859

Amorphophallus commutatus var. *anmodensis* Sivad. & Jaleel, Rheedia 12: 159. 2002

Amorphophallus commutatus var. *wayanadensis* Sivad. & Jaleel, Rheedia 12: 163 2002

Amorphophallus commutatus var. *anshiensis* Punekar & Lakshmin., Fl. Anshi Natl. Park 514. 2011

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Local/Common/Vernacular name: *Kempu lutiche* (Kunabi)

Amorphophallus hohenackeri (Schott) Engl. & Gehrm., Pflanzenr. IV, 23C: 103 .1911; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Plate 16F

Rhaphiophallus hohenackeri Schott, Gen. Aroid. t. 27.1858

Corynophallus hohenackeri (Schott) Kuntze, Revis. Gen. Pl. 2: 741.1891

Hydrosme hohenackeri (Schott) Engl., Nat. Pflanzenfam. Nachtr. 1: 59.1897

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Kerala; endemic.

Amorphophallus konkanensis Hett., S.R.Yadav & K.S.Patil, Blumea 39: 289.1994

Distribution: Karnataka (Belagavi, Uttara kannada), Goa, Maharashtra; endemic.

Amorphophallus mysorensis E.Barnes & C.E.C.Fisch., Bull. Misc. Inform. Kew 1939: 661.1940; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Distribution: Karnataka (Mysuru); endemic.

Amorphophallus paeoniifolius (Dennst.) Nicolson, Taxon 26: 337.1977

Dracontium paeoniifolium Dennst., Schlüssel Hortus Malab. 38.1818

Arum campanulatum Roxb., Pl. Coromandel 3: 68.1820

Candarum roxburghii Schott., Melet. Bot. 17.1832

Amorphophallus campanulatus Decne., Nouv. Ann. Mus. Hist. Nat. 3: 366.1834

Amorphophallus dubius Blume, Rumphia 1: 142.1837; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Amorphophallus rex Prain ex Hook. f., Fl. Brit. India 6: 514.1896

Amorphophallus paeoniifolius var. *campanulatus* (Decne.) Sivad., Taxon 32: 130 1983

Distribution: Karnataka (Dakshina Kannada, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada, Yadgir), Peninsular to NE India; Pacefic Islands, S & SE Asia.

Local/Common/Vernacular name: *Kadu suvarna gedde* (Kannada)

Amorphophallus sylvaticus (Roxb.) Kunth, Enum. Pl. 3: 34.1841

Arum sylvaticum Roxb., Fl. Ind. ed. 2, 3: 511.1832

Pythonium sylvaticum (Roxb.) Wight, Icon. Pl. Ind. Orient. t. 802.1844

Brachyspatha sylvatica (Roxb.) Schott, Syn. Aroid. 35.1856

Syntherias sylvatica (Roxb.) Schott, Gen. Aroid. t. 28.1858; B.D.Sharma et al., Fl. Karnataka Anal. 298. 1984.

Distribution: Karnataka (Mysuru, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Anaphyllum Schott

Anaphyllum wightii Schott, Gen. Aroid. 83.1858

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Anthurium Schott

Anthurium acaule (Jacq.) Schott, Wiener Z. Kunst 1829(3): 828.1829

Pothos acaulis Jacq., Enum. Syst. Pl. 31. 1760

Pothos lanceolatus L., Sp. Pl. ed. 2.1373 1763

Anthurium lanceolatum (L.) Schott, Wiener Z. Kunst 3: 828.1829

Distribution: Karnataka (cultivated as ornamental pot plant in Belagavi, Bengaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Kolar, Mysuru), Kerala, Tamil Nadu, Sikkim, West Bengal; native to Americas.

Anthurium andraeanum Linden ex André, Ill. Hort. 24: 43.1877

Distribution: Karnataka (cultivated as ornamental and for cut flowers in Bengaluru, Dakshina Kannada, Hassan, Kodagu), Kerala, Tamil Nadu, Sikkim, West Bengal; native to Americas.

Anthurium schlechtendalii Kunth, Enum. Pl. 3: 75.1841

Anthurium tetragonum Hook. ex Schott, Prodr. Syst. Aroid. 475.1860

Distribution: Karnataka (cultivated as ornamental pot plant in Bengaluru), Kerala, Tamil Nadu, Sikkim, West Bengal; native to Americas.

Ariopsis Nimmo

Ariopsis peltata Nimmo, Cat. Pl. Bombay 252.1839; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, NE India, subtropical Himalays; Myanmar, SE Asia.

Arisaema C.Martius

Arisaema barnesii C.E.C.Fisch., Bull. Misc. Inform. Kew 1933: 342.1933; B.D.Sharma et al., Fl. Karnataka Anal. 296. 1984.

Arisaema tylophorum C.E.C.Fisch., Bull. Misc. Inform. Kew 1933: 346.1933

Distribution: Karnataka (Chamarajanagar), Kerala, Tamil Nadu; endemic.

Arisaema caudatum Engl., Monogr. Phan. 2: 559.1879

Distribution: Karnataka (Uttara Kannada), Goa, Maharashtra; endemic.

Arisaema jacquemontii Blume, Rumphia 1: 95.1836

Arisaema wightii Schott, Bonplandia (Hannover) 7: 26.1859

Distribution: Karnataka (?), Kerala, Tamil Nadu; Afghanistan, Bangladesh, Bhutan, China, Nepal, Pakistan.

Arisaema leschenaultii Blume, Rumphia 1: 93.1836; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Arisaema neglectum Schott, Bonplandia (Hannover) 7: 26.1859

Arisaema longicaudatum Blatt., J. Proc. Asiat. Soc. Bengal n.s., 26: 362.1931

Arisaema attenuatum E.Barnes & C.E.C.Fisch., Bull. Misc. Inform. Kew 1936: 275.1936

Arisaema peltatum C.E.C.Fisch., Bull. Misc. Inform. Kew 1936: 277.1936

Distribution: Karnataka (Belagavi, Chamarajanagar, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Arisaema murrayi (J.Graham) Hook., Bot. Mag. 74: t. 4388.1848

Arum murrayi J.Graham, Cat. Pl. Bombay 229.1839

var. **murrayi**

Distribution: Karnataka (Dakshina Kannada, Udupi), Kerala, Maharashtra, Nagar Haveli, Tamil Nadu; endemic.

var. **sahyadricum** (S.R.Yadav, K.S.Patil & Bachulkar) M.R.Almeida, Fl. Maharashtra 5B: 255.2009

Arisaema sahyadricum S.R.Yadav, K.S.Patil & Bachulkar, Willdenowia 23: 177.1993

Distribution: Karnataka (Belagavi), Kerala, Maharashtra, Tamil Nadu; endemic.

Arisaema tortuosum (Wall.) Schott, Melet. Bot. 17.1832; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Arum tortuosum Wall., Pl. Asiat. Rar. 2: 10.1830

Arum curvatum Roxb., Fl. Ind. ed. 2,3: 506 1832

Arisaema curvatum (Roxb.) Kunth, Enum. Pl. 3: 20.1841

Arisaema commutatum Schott, Bonplandia (Hannover) 7: 26.1859

Arisaema wightii Hook.f., Bot. Mag. 91: t. 5507. 1865.

Arisaema tortuosum var. *curvatum* (Roxb.) Engl., Pflanzenr. IV, 23F: 191.1920

Arisaema tortuosum var. *neglectum* (Schott) C. E. C. Fisch., Fl. Madras 1585 1931

subsp. **tortuosum**

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; Bhutan, China, Nepal, Sri Lanka.

subsp. **sivadasanii** (S.R.Yadav, K.S.Patil & Janarth.) Punekar & Kumaran, Novon 19: 394.2009

Arisaema sivadasanii S.R.Yadav, K.S.Patil & Janarth, Aroideana 20: 53.1997

Arisaema murrayi var. *sivadasanii* (S.R.Yadav, K.S.Patil & Janarth.) M.R.Almeida, Fl. Maharashtra 5B: 256.2009

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; Bhutan, China, Nepal, Sri Lanka.

Caladium Vent.

Caladium bicolor (Aiton) Vent., Mag. Encycl. 4: 464.1800

Arum bicolor Aiton, Hort. Kew. 3: 316.1789

Cyrtospadix bicolor (Aiton) Britton & P.Wilson, Bot. Porto Rico 5: 126.1924

Distribution: Karnataka (cultivated and sometimes naturalised in Belagavi, Bengaluru, Dakshina Kannada, Mysuru, Udupi), cultivated throughout India; native to S America.

Colocasia Schott

Colocasia affinis Schott, Bonplandia (Hannover) 7: 28.1859

Distribution: Karnataka (cultivated in Dakshina Kannada, Udupi), NE India; Bangladesh, China, Myanmar, Nepal

Colocasia esculenta (L.) Schott, Melet. Bot. 18.1832; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Arum colocasia L., Sp. Pl. 985.1753

Arum esculentum L., Sp. Pl. 965.1753

Arum chinense L., Amoen. Acad. 4: 234.1754

Caladium esculentum (L.) Vent., Descr. Pl. Nov. t. 30.1801

Caladium nymphaeifolium Vent., Descr. Pl. Nov. t. 30.1801

Arum nymphaeifolium (Vent.) Roxb., Hort. Bengal. 65.1814

Steudnera virosa (Roxb.) Prain, Bengal Pl. 1113.1903

Calla virosa Roxb., Fl. Ind. ed. 1832 3: 517.1832

Colocasia antiquorum Schott, Melet. Bot. 18.1832

Zantedeschia virosa (Roxb.) K.Koch, Index Seminum (B) 1854(App.): 9.1855

Leucocasia esculenta (L.) Nakai, Bull. Natl. Sci. Mus. Tokyo 31: 127.1952

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Yadgir) Peninsular to NE India; native to tropical Asia.

Local/Common/Vernacular names: *Kesave*, *Kesave gedde*, *Kesu*, *Kesuva* (Kannada); *Alu*, *Tero* (*Kunabi*)

Cryptocoryne Fischer ex Wydler

Cryptocoryne ciliata (Roxb.) Fisch. ex Wydler, Linnaea 5: 428.1830

Ambrosina ciliata Roxb., Pl. Coromandel 3: 90.1820

Ambrosina ciliaris Spreng., Syst. Veg. 3: 771.1826

Arum ciliatum (Roxb.) Loudon, Encycl. Pl. Suppl. 2: 150.1841; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Cryptocoryne alata Griff., Iconogr. Pl. Asiae Orient. 3: t. 170.1851

Cryptocoryne elata Griff., Not. Pl. Asiat. 3: 134.1851

Distribution: Karnataka (Mysuru), Maharashtra, Tamil Nadu; New Guinea, SE Asia, Vietnam.

Local/Common/Vernacular name: *Kachchi* (Kannada)

Cryptocoryne consobrina Schott, Bonplandia (Hannover) 5: 222.1857

Cryptocoryne wightii Schott, Prodr. Syst. Aroid. 17.1860; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Distribution: Karnataka (Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Cryptocoryne retrospiralis (Roxb.) Kunth, Enum. Pl. 3: 12.1841; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Ambrosina retrospiralis Roxb., Hort. Bengal. 65.1814

Ambrosina unilocularis Roxb., Fl. Ind. ed.2, 3: 493.1832

Ambrosina roxburghiana Voigt, Hort. Suburb. Calcutt. 685.1845

Cryptocoryne dalzellii Schott, Bonplandia (Hannover) 5: 221.1857

Cryptocoryne unilocularis (Roxb.) Schott, Bonplandia (Hannover) 5: 223.1857

Cryptocoryne roxburghiana (Voigt) Schott, Prodr. Syst. Aroid. 18.1860

Lagenandra dalzellii (Schott) Rataj, Stud. Ceskoslov. Akad. Ved 3: 101.1975

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, NE India, Tamil Nadu; Bangladesh, Myanmar.

Cryptocoryne sivadasanii Bogner, Willdenowia 34: 195.2004

Distribution: Karnataka (Dakshina Kannada, Udupi), Kerala; endemic.

Cryptocoryne spiralis (Retz.) Fisch. ex Wydler, Linnaea 5: 428.1830; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Arum spirale Retz., Observ. Bot. 1: 30.1779

Ambrosina spiralis (Retz.) Roxb., Hort. Bengal. 65.1814

Cryptocoryne huegelii Schott, Aroideae 8, t. 12.1853

Cryptocoryne tortuosa Blatt. & McCann, J. Bombay Nat. Hist. Soc. 35: 16.1931

var. *spiralis*

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Mysuru, Udupi, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; Bangladesh, Sri Lanka.

Local/Common/Vernacular name: *Neerina peepi* (Kannada)

var. *cognatoides* (Blatt. & McCann) S.R.Yadav, K.S.Patil & Bogner, Aqua Pl. 1993(2): 65.1993

Cryptocoryne cognatoides Blatt. & McCann, J. Bombay Nat. Hist. Soc. 35: 17.1931; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Distribution: Karnataka (Belagavi, Uttara Kannada), Goa, Maharashtra; endemic.

Dieffenbachia Schott**Dieffenbachia seguine (Jacq.) Schott, Melet. Bot. 20.1832**

Arum seguine Jacq., Enum. Syst. Pl. 31.1760

Caladium seguine (Jacq.) Vent., Descr. Pl. Nov. t. 30.1801

Dieffenbachia picta Schott, Oesterr. Bot. Wochensbl. 2: 68.1852

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical America.

Local/Common/Vernacular name: *Dumbcane* (English)

Epipremnum Schott**Epipremnum aureum (Linden & André) G.S.Bunting, Ann. Missouri Bot. Gard. 50: 28.1964**

Pothos aureus Linden & André, Ill. Hort. 27: 69.1880

Scindapsus aureus (Linden & André) Engl., Pflanzenr. IV, 23B: 80.1908

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to French Polynesia.

Epipremnum pinnatum (L.) Engl., Pflanzenr. IV, 23B: 60.1908; B.D.Sharma et al., Fl. Karnataka Anal. 298. 1984.

Pothos pinnatus L., Sp. Pl. ed. 2: 1324.1763

Polypodium laciniatum Burm.f., Fl. Indica 231.1768

Pothos caudatus Roxb., Fl. Ind. 1: 476.1820

Monstera caudata (Roxb.) Schott, Wiener Z. Kunst 4: 1028.1830

Scindapsus caudatus (Roxb.) Schott, Melet. Bot. 21.1832

Scindapsus pinnatus (L.) Schott, Melet. Bot. 21.1832

Rhaphidophora caudata (Roxb.) Schott, Prodr. Syst. Aroid. 382.1860

Rhaphidophora laciniata (Burm.f.) Merr., Philipp. J. Sci. 19: 342.1921

Distribution: Karnataka (cultivated throughout as a indoor and garden plant), Australia, SE Asia, S China, New Guinea, Melanesia; naturalized to West Indies

Local/Common/Vernacular names: *Australian native monstera, centipede tongavine, devil's ivy, golden pothos, hunter's robe, ivy arum, money plant, silver vine, Solomon Islands ivy, taro vine* (English)

Lagenandra Dalzell**Lagenandra meeboldii (Engl.) C.E.C.Fisch., Fl. Madras 1576.1931**

Cryptocoryne meeboldii Engl., Pflanzenr. IV, 23F: 234.1920

Distribution: Karnataka (Dakshina Kannada, Hassan, Mysuru, Shivamogga, Udupi), Kerala, Tamil Nadu; endemic.

Lagenandra ovata (L.) Thwaites, Enum. Pl. Zeyl. 334.1864; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Arum ovatum L., Sp. Pl. 967.1753

Caladium ovatum (L.) Vent., Descr. Pl. Nov. t. 30.1801

Cryptocoryne ovata (L.) Schott, Melet. Bot. 16.1832

Arisarum ovatum (L.) Raf., Fl. Tellur. 3: 63.1837

Lagenandra insignis Trimen, J. Bot. 23: 269.1885

Distribution: Karnataka (Belagavi, Chamarajanagar, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Lagenandra toxicaria Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4: 289.1852

Distribution: Karnataka (Dakshina Kannada, Udupi), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Monstera Adans.

Monstera deliciosa Liebm., Vidensk. Meddel. Naturhist. Foren. Kjøbenhavn 1: 19.1849

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Mexico to Panama.

Local/Common/Vernacular names: *cheese plant, Mexican breadfruit, Monster fruit, Swiss cheese plant, The Fruit Salad Plant* (English); *Monstera* (Kannada)

Philodendron Schott

Philodendron cordatum Kunth ex Schott, Syn. Aroid. 95.1856

Distribution: Karnataka (cultivated as an ornamental throughout), cultivated as an ornamental throughout India; native to Brazil.

Local/Common/Vernacular name: *Heart Leaf Philodendron* (English)

Philodendron giganteum Schott, Syn. Aroid. 89.1856

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Udupi, Uttara Kannada), cultivated as an ornamental throughout India; native to the West Indies.

Local/Common/Vernacular name: *Giant Philodendron* (English)

Philodendron lacerum (Jacq.) Schott, Wiener Z. Kunst 3: 780.1829

Arum lacerum Jacq., Pl. Hort. Schoenbr. 4: 35.1804

Caladium lacerum (Jacq.) Willd., Sp. Pl. 4: 491.1805

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Udupi, Uttara Kannada), cultivated as an ornamental throughout India; native to West Indies

Local/Common/Vernacular name: *Toothed philodendron* (English)

Pistia L.

Pistia stratiotes L., Sp. Pl. 963.1753; B.D.Sharma et al., Fl. Karnataka Anal. 297. 1984.

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular name: *Antaragange* (Kannada)

Pothos L.

Pothos scandens L., Sp. Pl. 968.1753; B.D.Sharma et al., Fl. Karnataka Anal. 298. 1984.

Distribution: Karnataka (Belagavi, Chickmagluru, Dakshina Kannada, Davanagere, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada, Yadgir), throughout India; China, Madagascar, S & SE India.

Local/Common/Vernacular names: *Adikabilu balli, Adike balli, Akki balli, Bhandarli, Murugina beelu* (Kannada)

Remusatia Schott

Remusatia vivipara (Roxb.) Schott, Melet. Bot. 18.1832; B.D.Sharma et al., Fl. Karnataka Anal. 298. 1984.

Arum viviparum Roxb., Hort. Bengal. 65.1814

Caladium viviparum (Roxb.) Nees, Melet. Bot. 18.1832

Colocasia vivipara (Roxb.) Thwaites, Enum. Pl. Zeyl. 336.1864

Distribution: Karnataka (Belagavi, Chamarajanagar, Chickmagluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Peninsular to NE India; Africa, Australia, Madagascar, S & SE Asia.

Local/Common/Vernacular name: *Mara kesu* (Kannada); *Kasalu* (*Kunabi*)

Rhaphidophora Hassk.

Rhaphidophora decursiva (Roxb.) Schott, Bonplandia (Hannover) 5: 45.1857

Pothos decursivus Roxb., Fl. Ind. 1: 456.1820

Monstera decursiva (Roxb.) Schott, Wiener Z. Kunst 4: 1028.1830

Scindapsus decursivus (Roxb.) Schott, Melet. Bot. 21.1832

Distribution: Karnataka (Belagavi, Kodagu), NE India; Bangladesh, Bhutan, Cambodia, China, Laos, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Rhaphidophora pertusa (Roxb.) Schott, Bonplandia (Hannover) 5: 45.1857

Pothos pertusus Roxb., Fl. Ind. 1: 455.1820

Monstera pertusa (Roxb.) Schott, Wiener Z. Kunst 4: 1028.1830

Scindapsus pertusus (Roxb.) Schott, Melet. Bot. 21.1832

Distribution: Karnataka (Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), Peninsular to NE India; S & SE Asia.

Local/Common/Vernacular name: *Creeping Philodendron* (English)

Sauromatum Schott

Sauromatum venosum (Dryand. ex Aiton) Kunth, Enum. Pl. 3: 28.1841

Arum venosum Dryand. ex Aiton, Hort. Kew. 3: 315.1789

Arum pedatum Willd., Enum. Pl. 54.1814

Arum guttatum Wall., Pl. Asiat. Rar. 2: 10.1830

Arum sessiliflorum Roxb., Fl. Ind. ed. 2, 3: 507 1832

Sauromatum guttatum Schott, Melet. Bot. 17.1832

Sauromatum pedatum (Link & Otto) Schott, Melet. Bot. 17.1832; B.D.Sharma et al., Fl. Karnataka Anal. 298. 1984.

Arisaema venosum (Dryand. ex Aiton) Blume, Rumphia 1: 109.1836

Alocasia pedata (Link & Otto) Raf., Fl. Tellur. 3: 64.1837

Desmesia venosum (Dryand. ex Aiton) Raf., Fl. Tellur. 3: 63.1837

Sauromatum sessiliflorum (Roxb.) Kunth, Enum. Pl. 3: 28.1841

Sauromatum simlense Schott, Oesterr. Bot. Z. 8: 349.1858

Sauromatum guttatum var. *pedatum* (Link & Otto) Engl., Pflanzenr. IV, 23F: 125.1920

Sauromatum guttatum var. *sessiliflorum* (Roxb.) Engl., Pflanzenr. IV, 23F: 125.1920

Sauromatum guttatum var. *simlense* (Schott) Engl., Pflanzenr. IV, 23F: 125.1920

Sauromatum guttatum var. *venosum* (Dryand. ex Aiton) Engl., Pflanzenr. IV, 23F: 125.1920

Typhonium venosum (Dryand. ex Aiton) Hett. & P.C.Boyce, Aroideana 23: 51.2000

Distribution: Karnataka (Belagavi, Dharwad, Hassan), Peninsular to NE India; Africa, Bhutan, China, Myanmar, Nepal.

Scindapsus Schott

Scindapsus officinalis (Roxb.) Schott, Melet. Bot. 21.1832

Pothos officinalis Roxb., Fl. Ind. ed. 1, 1: 452.1820

Monstera officinalis (Roxb.) Schott, Wiener Z. Kunst 4: 1028.1830

Distribution: Karnataka (cultivated as an ornamental), cultivated (and Wild) throughout India; Bangladesh, Bhutan, Nepal, Myanmar, Thailand, Laos, Cambodia, Vietnam.

Syngonium Schott

Syngonium auritum (L.) Schott, Wiener Z. Kunst 3: 780.1829

Arum auritum L., Syst. Nat. ed. 10 2: 1251. 1759

Caladium auritum (L.) Vent., Descr. Pl. Nov. t. 30.1801

Syngonium plumieri Schott, Syn. Aroid. 66.1856

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Udupi, Uttara Kannada), cultivated as an ornamental throughout India; native to tropical America.

Theriophonum Blume

Theriophonum dalzellii Schott, Aroideae 1: 15.1853; B.D.Sharma et al., Fl. Karnataka Anal. 298. 1984. **Plate 16G**

Tapinocarpus dalzellii Schott, Gen. Aroid. t. 15.1858

Theriophonum uniseriatum Blatt. & McCann, J. Bombay Nat. Hist. Soc. 35: 25.1931

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Theriophonum fischeri Sivad., Aroideana 4: 64.1981

Theriophonum indicum Engl., Pflanzenr. IV, 23F: 107.1920

Distribution: Karnataka (Belagavi), Kerala, Tamil Nadu; endemic.

Typhonium Schott

Typhonium bulbiferum Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4: 113.1852; Manikand.& Lakshminars., Fl. Rajiv Gandhi Nation. Park 400.2013.

Distribution: Karnataka (Dharwad, Mysuru, Uttara Kannada), Gujarat, Kerala, Maharashtra, Tamil Nadu; endemic.

Typhonium flagelliforme (Lodd.) Blume, Rumphia 1: 134.1837

Arum divaricatum L., Sp. Pl. 966.1753

Arum flagelliforme Lodd., Bot. Cab. 4: t. 396.1819

Typhonium cuspidatum Decne, Nouv. Ann. Mus. Hist. Nat. 3: 367.1834

Typhonium sylvaticum Voigt, Hort. Suburb. Calcutt. 686.1845

Arum angulatum Griff., Not. Pl. Asiat. 3: 143.1851

Typhonium flagelliferum Griff., Not. Pl. Asiat. 3: 144.1851

Heterostalis flagelliformis (Lodd.) Schott, Oesterr. Bot. Wochenbl. 7: 261.1857

Typhonium incurvatum Blatt. & McCann, J. Bombay Nat. Hist. Soc. 35: 22.1931

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi), Peninsular to NE India; Australia, Indo-China, S & SE Asia.

Typhonium roxburghii Schott, Aroideae 1: 12.1853

Typhonium divaricatum var. *roxburghii* (Schott) Engl., Monogr. Phan. 2: 612.1879

Typhonium schottii Prain, J. Asiatic Soc. Bengal, Pt. 2, Nat. Hist. 67: 303.1898

Typhonium amboinense Blatt. & McCann, J. Bombay Nat. Hist. Soc. 3: 23.1931

Distribution: Karnataka (Dakshina Kannada, Udupi), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka, SE Asia.

Xanthosoma Schott

Xanthosoma sagittifolium (L.) Schott, Melet. Bot. 19.1832; Parimala, Fl. Dharwad Dist. 490. 2013.

Arum sagittifolium L., Sp. Pl. 966.1753

Caladium sagittifolium (L.) Vent., Descr. Pl. Nov. t. 30.1801

Distribution: Karnataka (cultivated in Bengaluru, Dharwad, Udupi, Uttara Kannada), cultivated as an ornamental in India; native to West Indies.

Zantedeschia Spreng.

Zantedeschia aethiopica (L.) Spreng., Syst. Veg. 3: 765.1826

Calla aethiopica L., Sp. Pl. 968.1753

Colocasia aethiopica (L.) Link, Diss. Bot. 77.1795

Richardia aethiopica (L.) Spreng., Syst. Veg. 3: 765.1826

Otosma aethiopica (L.) Raf., New Fl. 2: 90.1837

Arodes aethiopicum (L.) Kuntze, Revis. Gen. Pl. 2: 740.1891

Distribution: Karnataka (cultivated as an ornamental), cultivated as an ornamental; native to S Africa.

Local/Common/Vernacular names: *Arum lily*, *Calla lily* (English)

Araliaceae

Aralia L.

Aralia leschenaultii (DC.) J.Wen, Brittonia 45: 53 1993

Panax leschenaultii DC., Prodr. 4: 254 1830

Pentapanax leschenaultii (DC.) Seem. in J. Bot. 2: 296 1864; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Distribution: Karnataka (Cultivated, Bengaluru), Kerala; China, Myanmar, Sri Lanka.

Hydrocotyle L.

Hydrocotyle javanica Thunb., Diss. Hydroc. n. 17. t. 2. 1798; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Hydrocotyle nepalensis Hook., Exot. Fl. 1:t. 30 1822

Hydrocotyle polyccephala Wight & Arn., Prodr. Fl. Ind. Orient. 1: 366. 1834

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga), throughout India; Tropical Africa to Indomalesia

Hydrocotyle sibthorpioides Lam., Encycl. 3: 153. 1789; B.D.Sharma et al., Fl. Karnataka Anal. 119. 1984.

Hydrocotyle rotundifolia Roxb. ex DC., Prodr. 4: 64. 1830

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Hassan, Mysuru, Uttara Kannada), Arunachal Pradesh, Bihar, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand, West Bengal; Paleotropical

Schefflera J. R. Forst. & G. Forst.

Schefflera actinophylla (Endl.) Harms in Nat. Pflanzenfam. 3(8): 36 1894

Brassaia actinophylla Endl., Nov. Stirp. Dec. 1: 89. 1839; C.J.Saldanha, Fl. Karnataka 2: 277. 1996

Distribution: Karnataka (Cultivated); Tropical Asia to Australia

Local/Common/Vernacular name: *Queensland Umbealla tree* (English)

Schefflera capitata (Wight & Arn.) Harms in Engl. & Prantl, Naturl. Pflanzenfam. 3(8): 36. 1894; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Paratropia capitata Wight & Arn., Prodr. Fl. Ind. Orient. 1: 378. 1834

Brassaia capitata (Wight & Arn.) C.B.Clarke in Hook. f., Fl. Brit. India 2: 732. 1879

Distribution: Karnataka (Mysuru, Shivamogga), Kerala, Tamil Nadu; India, Myanmar

Schefflera micrantha (C.B.Clarke) Gamble, Fl. Madras 569(402). 1919; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Heptapleurum rostratum var. *micranthum* C.B.Clarke, Fl. Brit. India 2: 729. 1879

Schefflera rostrata var. *micrantha* (C.B.Clarke) Maheshw., Bull. Bot. Surv. India 2: 376. 1960

Distribution: Karnataka (Chikkamagaluru, Shivamogga), Kerala, Tamil Nadu; endemic.

Schefflera rostrata (Wight) Harms in Engl. & Prantl, Naturl. Pflanzenfam. 3(8): 38. 1894; C.J.Saldanha, Fl. Karnataka 2: 274. 1996

Hedera rostrata Wight, Icon. Pl. Ind. Orient. 3: t. 1013. 1845

Distribution: Karnataka (Chikkamagaluru, Kodagu, Shivamogga), Kerala, Tamil Nadu; endemic.

Schefflera roxburghii Gamble, Fl. Madras 570 1919; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Distribution: Karnataka (Ballari, Kolar, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; Bangladesh? Sri Lanka?

Schefflera stellata (Gaertn.) Baill., Hist. Pl. 7: 161. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 120. 1984.

Heptapleurum stellatum Gaertn, Fruct. 2:472. t.178. 1791

Hedera obovata Wight, Icon. Pl. Ind. Orient. 3: t. 1011 1845

Distribution: Karnataka (Ballari, Chitradurga, Mysuru, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; Peninsular India, Sri Lanka.

Schefflera venulosa (Wight & Arn.) Harms in Engl. & Prantl, Naturl. Pflanzenfam. 3(8): 39. 1894; B.D.Sharma et al., Fl. Karnataka Anal. 121. 1984.

Aralia digitata Roxb., Fl. Ind. 2: 107. 1832

Paratropia venulosa Wight & Arn., Prodr. Fl. Ind. Orient. 1: 377. 1834

Heptapleurum venulosum (Wight & Arn.) Seem., J. Bot. 3: 80. 1865

Schefflera bengalensis Gamble, Bull. Misc. Inform. Kew 229. 1919

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout tropical and Subtropical India; China, Indochina, Malaysia, Myanmar.

Schefflera wallichiana (Wight & Arn.) Harms, Engl. & Prantl, Naturl. Pflanzenfam. 3(8): 38. 1894; B.D.Sharma et al., Fl. Karnataka Anal. 121. 1984.

Paratropia wallichiana Wight & Arn., Prodr. Fl. Ind. Orient. 1: 77. 1834

Heptapleurum wallichianum (Wight & Arn.) Seem., J. Bot. 3: 80 1865

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Arunachal Pradesh, Kerala, Meghalaya, Sikkim, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Betthu mara* (Kannada)

Arecaceae**Actinorhytis H.Wendl. & Drude**

Actinorhytis calapparia (Blume) H.Wendl. & Drude ex Scheff., Ann. Jard. Bot. Buitenzorg 1: 156.1876;
B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Areca calapparia Blume, Rumphia 2: t. 100.1843

Areca cocoides Griff., Calcutta J. Nat. Hist. 4: 454.1845

Seaforthia calapparia (Blume) Mart., Hist. Nat. Palm. 3: 313.1849

Ptychosperma calapparia (Blume) Miq., Fl. Ned. Ind. 3: 20.1855

Pinanga calapparia (Blume) H.Wendl., Palmiers 253.1878

Distribution: Karnataka (cultivated in Dakshina Kannada, Uttara Kannada), cultivated in gardens; native to New Guinea & The Solomon Islands.

Local/Common/Vernacular name: *Calappa Palm* (English)

Adonidia Becc.

Adonidia merrillii (Becc.) Becc., Philipp. J. Sci. 14: 329.1919; K. G. Bhat, Palms Karnataka 71.2011.

Normanbya merrillii Becc., Philipp. J. Sci., C 4: 606.1909

Veitchia merrillii (Becc.) H.E.Moore, Gentes Herb. 8: 501.1957

Distribution: Karnataka (cultivated in gardens), native to Indonesia & Philippines.

Local/Common/Vernacular names: *Christmas Palm, Kerpis Palm, Manila Palm* (English)

Aiphanes Willd.

Aiphanes horrida (Jacq.) Burret, Notizbl. Bot. Gart. Berlin-Dahlem 11: 575.1932; K. G. Bhat, Palms Karnataka 57.2011.

Caryota horrida Jacq., Fragm. Bot. 20.1801

Aiphanes aculeata Willd., Samml. Deutsch. Abh. Koningl. Akad. Wiss. Berlin 1803: 251.1806

Martinezia caryotifolia Kunth, Nov. Gen. Sp. 1: 305.1816

Aiphanes caryotifolia (Kunth) H.Wendl., Palmiers 230.1878

Tilmia caryotifolia (Kunth) O.F.Cook, Bull. Torrey Bot. Club 28: 565.1901

Distribution: Karnataka (cultivated in gardens), native to Northern S America & The Caribbean Islands.

Local/Common/Vernacular names: *Aculeata palm, Coyure palm, Ruffle palm, Spine palm* (English)

Archontophoenix H.Wendl. ex Drude

Archontophoenix alexandrae (F.Muell.) H.Wendl. & Drude, Linnaea 39: 212.1875; K. G. Bhat, Palms Karnataka 61.2011.

Ptychosperma alexandrae F.Muell., Fragm. 5: 47.1865

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Australia.

Local/Common/Vernacular names: *Alexander palm, Alexandra palm, King Alexander palm, King palm, Northern bangalow palm* (English)

Areca L.

Areca catechu L., Sp. Pl. 1189.1753; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Areca cathechu Burm.f., Fl. Indica 241.1768

Areca himalayana Griff. ex H.Wendl., Palmiers 231.1878

Areca catechu var. *deliciosa* Rau, J. Bombay Nat. Hist. Soc. 23:793. 1915

Distribution: Karnataka (Extensively cultivated except in dry eastern districts).widely cultivated in tropical Asia & Western Pacific region; presumably wild in Andaman & Nicobar Islands.

Local/Common/Vernacular names: *Areca nut, Beetle nut* (English); *Adike, Supaari* (Kannada); *Kangu* (Tulu)

Areca macrocalyx Zipp. ex Blume, Rumphia 2: 75.1839; K. G. Bhat, Palms Karnataka 63.2011.

Distribution: Karnataka (cultivated Dakshina Kannada), cultivated in gardens; native to Indonesia & Papua New Guinea.

Local/Common/Vernacular names: *Dwarf Betel Nut Palm, Highland Betel Nut Palm* (English)

Areca triandra Roxb. ex Buch.-Ham., Mem. Wern. Nat. Hist. Soc. 5: 310.1826; K. G. Bhat, Palms Karnataka 64.2011.

Distribution: Karnataka (cultivated in gardens of Belagavi, Bengaluru, Dakshina Kannada, Mysuru, Udupi), NE India, Andaman & Nicobar Islands; Bangladesh, Myanmar, Thailand, Vietnam.

Arenga Labill. Nom. Cons.

Arenga caudata (Lour.) H.E.Moore, Principes 4: 114.1960; K. G. Bhat, Palms Karnataka 43.2011.

Borassus caudatus Lour., Fl. Cochinch. 760.1790

Wallichia caudata (Lour.) Mart., Hist. Nat. Palm. 3: 315.1853

Blancoa caudata (Lour.) Kuntze, Gen. Pl. 2: 727.1891

Didymosperma caudatum (Lour.) H.Wendl. & Drude ex B.D.Jacks., Index Kew. 1: 756.1893

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Myanmar, Indo-China, SE Asia.

Arenga engleri Becc., Malesia 3: 184.1889; K. G. Bhat, Palms Karnataka 43.2011.

Didymosperma engleri (Becc.) Warb., Monsunia 1: t. 2, f. 1.1900

Distribution: Karnataka (cultivated), cultivated in gardens; native to Japan & Taiwan.

Arenga wightii Griff., Calcutta J. Nat. Hist. 5: 475.1845; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Plate 17A

Saguerus wightii (Griff.) H.Wendl. & Drude, Palmiers 256.1878

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Arenga, The Wild Coconut, Wight's Sago Palm, Wild Coconut* (English); *Dhadashi, Dodda salu* (Kannada)

Bactris Jacq. ex Scop.

Bactris major Jacq., Select. Stirp. Amer. Hist. ed. 2: 134.1781; K. G. Bhat, Palms Karnataka 59. 2011.

Distribution: Karnataka (cultivated in Dakshina Kannada), cultivated in gardens; native to S & C America.

Bismarckia Hildebr. & H.Wendl.

Bismarckia nobilis Hildebr. & H.Wendl., Bot. Zeitung (Berlin) 39: 94.1881; K. G. Bhat, Palms Karnataka 46. 2011.

Distribution: Karnataka (cultivated in gardens), native to Madagascar.

Local/Common/Vernacular names: *Bismarckia Palm, Nobilis Palm, The Bismarck Palm* (English)

Borassus L.

Borassus flabellifer L., Sp. Pl. 1187.1753; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Distribution: Karnataka (wild in coastal areas, elsewhere cultivated), native to coastal states, Bihar, Uttar Pradesh; Africa, Australia, tropical Asia.

Local/Common/Vernacular names: *Toddy palm* (English); *Tahatinunga, Thaale* (Kannada)

Calamus L.

Calamus brandisii Becc., Fl. Brit. India 6: 448.1892; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.
 Distribution: Karnataka (Uttara Kannada), Kerala, Tamil Nadu; endemic.

Calamus dransfieldii Renuka, Kew Bull. 42: 433.1987; K. G. Bhat, Palms Karnataka 13. 2011.
 Distribution: Karnataka (Dakshina Kannada, Mysuru, Udupi), Kerala; endemic.
 Local/Common/Vernacular name: *Meese betha* (Kannada)

Calamus gamblei Becc. In Hook. f., Fl. Brit. India 6: 453.1893; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.
 Distribution: Karnataka (Chamrajanagara, Chikkamagaluru, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Calamus hookerianus Becc., Ann. Roy. Bot. Gard. (Calcutta) 11(1): 226.1908; Renuka & Sreek. Field Guide Palms India 72. 2012
 Distribution: Karnataka (Uttara Kannada), Kerala, Tamil Nadu; endemic.

Calamus karnatakensis Renuka & Lakshmana, R. I. C. Bull. 9: 10.1990; K. G. Bhat, Palms Karnataka 14. 2011.
 Distribution: Karnataka (Kodagu, Shivamogga), Kerala; endemic.

Calamus lacciferus Lakshmana & Renuka, J. Econ. Taxon. Bot. 14: 707.1990; K. G. Bhat, Palms Karnataka 15. 2011.
 Distribution: Karnataka (Shivamogga), Kerala; endemic.

Calamus lakshmanae Renuka, J. Econ. Taxon. Bot. 14: 703.1990; K. G. Bhat, Palms Karnataka 16. 2011.
 Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Calamus metzianus Schldl., Linnaea 26: 727.1855; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.
Calamus rudentum Mart., Hist. Nat. Palm. 3: 340.1853
Calamus rivalis Thwaites ex Trimen, J. Bot. 23: 386 1885
Palmijuncus rivalis (Thwaites ex Trimen) Kuntze, Revis. Gen. Pl. 2: 733.1891
 Distribution: Karnataka (Dakshina Kannada, Udupi,Uttra Kannada), Kerala, Tamil Nadu; Sri Lanka.

Calamus nagbettai R.R. Fernandez & Dey, Indian Forester 96: 223.1970; B.D.Sharma et al., Fl. Karnataka Anal. 278. 1984.
 Distribution: Karnataka (Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), Kerala; endemic.
 Local/Common/Vernacular name: *Naga betha* (Kannada)

Calamus prasinus Lakshmana & Renuka, J. Econ. Taxon. Bot. 14: 705.1991; K. G. Bhat, Palms Karnataka 18. 2011.
 Distribution: Karnataka (Dakshina Kannada, Kodagu, Udupi), Kerala; endemic.
 Local/Common/Vernacular name: *Ontibetha* (Kannada)

Calamus pseudotenuis Becc. in Hook. f., Fl. Brit. India 6: 445.1892; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Tamil Nadu; Sri Lanka.
 Local/Common/Vernacular names: *Halu betha*, *Sanna betha* (Kannada)

Calamus rotang L., Sp. Pl. 325.1753; K. G. Bhat, Palms Karnataka 20. 2011.

Calamus monoecus Roxb., Fl. Ind. ed.2, 3: 783.1832

Calamus roxburghii Griff., Calcutta J. Nat. Hist. 5: 43.1845

Palmijuncus monoecus (Roxb.) Kuntze, Revis. Gen. Pl. 2: 733.1891

Distribution: Karnataka (cultivated), Andra Pradesh, Tamil Nadu; Sri Lanka

Calamus stoloniferus Renuka, J. Econ. Taxon. Bot. 14: 701.1990; K. G. Bhat, Palms Karnataka 21. 2011.

Distribution: Karnataka (Kodagu, Shivamogga); endemic.

Local/Common/Vernacular name: *Jedubetha* (Kannada)

Calamus thwaitesii Becc., Fl. Brit. India 6: 441.1892; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Plate 4A

Calamus thwaitesii var. *canaranus* Becc., Ann. Roy. Bot. Gard. (Calcutta) 11: 71.1908

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Handibetha* (Kannada)

Calamus travancoricus Bedd. ex Becc. in Hook. f., Fl. Brit. India 6: 452.1893; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Kari betha* (Kannada)

Calamus vattayila Renuka, Curr. Sci. 56: 1012.1987; K. G. Bhat, Palms Karnataka 23. 2011.

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Calamus viminalis Willd., Sp. Pl. 2: 203.1799; K. G. Bhat, Palms Karnataka 24. 2011.

Distribution: Karnataka (cultivated), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Kerala, Odisha, Sikkim, Tripura, West Bengal; Bangladesh, Indonesia, Malaysia, Myanmar.

Local/Common/Vernacular name: *Jungli beth* (Hindi)

Calamus wightii Griff., Palms Brit. E. Ind. t. 216C.1850; Renuka & Sreekum., Field Guide Palms India 132, 2012.

Calamus huegelianus Mart., Hist. Nat. Palm. 3: 338.1853; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Daemonorops melanolepis Mart., Hist. Nat. Palm. 3: 331.1853

Calamus melanolepis (Mart.) H.Wendl., Palmiers 237.1878

Palmijuncus huegelianus (Mart.) Kuntze, Revis. Gen. Pl. 2: 733.1891

Palmijuncus wightii (Griff.) Kuntze, Revis. Gen. Pl. 2: 733.1891

Distribution: Karnataka (Chikkamagaluruu, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Carpentaria Becc.

Carpentaria acuminata (H.Wendl. & Drude) Becc., Ann. Jard. Bot. Buitenzorg 2: 128.1885; K. G. Bhat, Palms Karnataka 72. 2011.

Kentia acuminata H.Wendl. & Drude, Linnaea 39: 207.1875

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Australia.

Local/Common/Vernacular names: *Carpentaria Palm*, *The Carpentaria Palm tree* (English)

Caryota L.

Caryota rumphiana Mart., Hist. Nat. Palm. 3: 195.1838; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Distribution: Karnataka (cultivated in Uttara Kannada), cultivated in gardens; native to Indonesia, Papua New Guinea & Philippines.

Local/Common/Vernacular names: *Albert palm*, *Fishtail palm* (English)

Caryota urens L., Sp. Pl. 1189.1753; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada, also cultivated as an ornamental), Peninsular to NE India, also cultivated as an ornamental; Malaysia, Myanmar, Nepal, Sri Lanka.

Local/Common/Vernacular names: *Fish tail Palm* (English); *Bagani mara, Baini mara* (Kannada); *Berlyamad (Kunabi); Endu* (Tulu)

Chamaedorea Willd.

Chamaedorea metallica O.F.Cook ex H.E.Moore, Principes 10: 45.1966; K. G. Bhat, Palms Karnataka 52. 2011.

Distribution: Karnataka (cultivated), cultivated in gardens; native to tropical & subtropical America.

Local/Common/Vernacular names: *Metallic Palm, Miniature Fishtail Palm* (English)

Chamaedorea seifrizii Burret, Notizbl. Bot. Gart. Berlin-Dahlem 14: 268.1938; K. G. Bhat, Palms Karnataka 52. 2011.

Chamaedorea erumpens H.E.Moore, Gentes Herb. 8: 232.1951

Distribution: Karnataka (cultivated), cultivated in gardens; native to tropical & subtropical America.

Local/Common/Vernacular names: *The Bamboo palm, The Lady Palm* (English)

Chamaerops L.

Chamaerops humilis L., Sp. Pl. 1187 1753

Phoenix humilis (L.) Cav., Icon. 2: 12.1793

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), cultivated in gardens; native to Mediterranean region.

Local/Common/Vernacular names: *European Fan Palm, Mediterranean Fan Palm, Moroccan Fan Palm* (English)

Coccothrinax Sarg.

Coccothrinax barbadensis (Lodd. ex Mart.) Becc., Webbia 2: 328.1908

Thrinax barbadensis Lodd. ex Mart., Hist. Nat. Palm. 3: 320.1853

Copernicia barbadensis (Lodd. ex Mart.) H.Wendl., Palmiers 241.1878

Distribution: Karnataka (Belagavi, Bengaluru?), cultivated in gardens; native to Caribbean Islands

Local/Common/Vernacular name: *Silver Thatch Palm* (English)

Cocos L.

Cocos nucifera L., Sp. Pl. 1188.1753; B.D.Sharma et al., Fl. Karnataka Anal. 294. 1984.

Cocos indica Royle, Ill. Bot. Himal. Mts. 395.1840

Cocos nana Griff., Not. Pl. Asiat. 3: 166.1851

Calappa nucifera (L.) Kuntze, Revis. Gen. Pl. 2: 982.1891

Distribution: Karnataka (cultivated throughout), cultivated, naturalized in Andaman & Nicobar Islands; throughout tropical coasts

Local/Common/Vernacular names: *Coconut palm* (English); *Tengina mara, Tengu* (Kannada); *Taare* (Tulu)

Corypha L.

Corypha umbraculifera L., Sp. Pl. 1178.1753; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada, cultivated elsewhere), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Talipot palm* (English); *Sritaale, Tale, Tali* (Kannada); *Pane* (Tulu)

Corypha utan Lam., Encycl. 2: 131.1786; K. G. Bhat, Palms Karnataka 24. 2011.

Distribution: Karnataka (cultivated), native to S & SE Asia & Australasia.

Local/Common/Vernacular names: *Cabbage Palm, Gebang Palm* (English)

Cyrtostachys Blume

Cyrtostachys renda Blume, Bull. Sci. Phys. Nat. Néerl. 1: 66 1838; K. G. Bhat, Palms Karnataka 75. 2011.

Bentinckia renda (Blume) Mart., Hist. Nat. Palm. 3: 316.1853

Cyrtostachys lakka Becc., Jard. Bot. Buitenzorg 2: 141.1885

Distribution: Karnataka (cultivated), cultivated in gardens; native to Indonesia, Malaysia & Singapore.

Local/Common/Vernacular names: *Lipstick palm, Red sealing wax palm* (English)

Daemonorops Blume

Daemonorops kurziana Hook.f. ex Becc.in Hook. f., Fl. Brit. India 6: 463.1893; K. G. Bhat, Palms Karnataka 23. 2011.

Distribution: Karnataka (cultivated in Lalbagh, Bengaluru), Andaman & Nicobar Islands; Myanmar.

Dictyosperma H.Wendl. & Drude

Dictyosperma album (Bory) Scheff., Ann. Jard. Bot. Buitenzorg 1: 157.1876

Areca alba Bory, Voy. îles Afrique 1: 306.1804

Linoma alba (Bory) O.F.Cook, J. Wash. Acad. Sci. 7: 123.1917

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Mascarene Islands.

Local/Common/Vernacular names: *Hurricane Palm, Princess Palm, Red Palm* (English)

Dypsis Noronha ex Mart.

Dypsis decaryi (Jum.) Beentje & J.Dransf., Palms Madagascar 187.1995; K. G. Bhat, Palms Karnataka 66. 2011.

Neodypsis decaryi Jum., Ann. Inst. Bot.-Géol. Colon. Marseille V, 1(1): 15.1933

Distribution: Karnataka (cultivated in gardens), throughout India; native to Madagascar.

Local/Common/Vernacular name: *The Triangle palm* (English)

Dypsis leptocheilos (Hodel) Beentje & J.Dransf., Palms Madagascar 176.1995; K. G. Bhat, Palms Karnataka 66. 2011.

Neodypsis leptocheilos Hodel, Palm J. 139: 9.1993

Distribution: Karnataka (cultivated in gardens), cultivated throughout India; native to Madagascar.

Local/Common/Vernacular names: *Redneck Palm, Teddy Bear Palm* (English)

Dypsis lutescens (H.Wendl.) Beentje & J.Dransf., Palms Madagascar 212.1995; K. G. Bhat, Palms Karnataka 67. 2011..

Chrysalidocarpus lutescens H.Wendl., Bot. Zeitung (Berlin) 36: 171.1878

Distribution: Karnataka (cultivated in gardens), cultivated throughout India; native to Madagascar.

Local/Common/Vernacular names: *Bamboo palm, Cane palm* (English)

Elaeis Jacq.

Elaeis guineensis Jacq., Select. Stirp. Amer. Hist. 280.1763; K. G. Bhat, Palms Karnataka 59. 2011.

Elaeis melanococca Gaertn., Fruct. Sem. Pl. 1: 18.1788

Distribution: Karnataka (cultivated in Bengaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Udupi, uttara Kannada), cultivated humid areas in India; native to West Africa.

Local/Common/Vernacular name: *Oil palm* (English)

Heterospathe Scheff.

Heterospathe elata Scheff., Ann. Jard. Bot. Buitenzorg 1: 162.1876; K. G. Bhat, Palms Karnataka 77. 2011.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Philippines.

Hyophorbe Gaertn.

Hyophorbe lagenicaulis (L.H.Bailey) H.E.Moore, Principes 20: 119.1976

Mascarena lagenicaulis L.H.Bailey, Gentes Herb. 6: 74.1942

Distribution: Karnataka (cultivated in gardens, Bengaluru), cultivated in gardens; native to Round Island & Mauritius.

Local/Common/Vernacular name: *The Bottle Palm* (English)

Hyophorbe verschaffeltii H.Wendl., Ill. Hort. 13(1): t. 462.1866

Areca verschaffeltii Lem., Ill. Hort. 13: t. 462.1866

Mascarena verschaffeltii (H.Wendl.) L.H.Bailey, Gentes Herb. 6: 76.1942

Distribution: Karnataka (cultivated in gardens, Bengaluru), cultivated in gardens; native to Mascarene Islands.

Local/Common/Vernacular names: *Spindle Pal, The Palmiste Marron* (English)

Latania Comm. ex Juss.

Latania loddigesii Mart., Hist. Nat. Palm. 3: 226.1838; K. G. Bhat, Palms Karnataka 47. 2011.

Cleophora loddigesii (Mart.) O.F.Cook, Natl. Hort. Mag. 20: 52.1941

Distribution: Karnataka (cultivated in gardens, Bengaluru), cultivated in gardens; native to Mauritius.

Local/Common/Vernacular name: *The blue latan palm* (English)

Latania lontaroides (Gaertn.) H.E.Moore, Principes 7: 85.1963; K. G. Bhat, Palms Karnataka 48. 2011.

Cleophora lontaroides Gaertn., Fruct. Sem. Pl. 2: 185.1791

Latania commersonii J.F.Gmel., Syst. Nat. 1035.1792

Latania rubra Jacq., Fragm. Bot. 13.1800

Cleophora commersonii (J.F.Gmel.) O.F.Cook, Natl. Hort. Mag. 20: 52.1941

Distribution: Karnataka (cultivated in gardens, Bengaluru), cultivated in gardens; native to Mascarene Islands

Local/Common/Vernacular name: *The Red Latan Palm* (English)

Licuala Thunb.

Licuala grandis H.Wendl., Ill. Hort. 27: t. 412.1880; K. G. Bhat, Palms Karnataka 37. 2011.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to the Solomon Islands & Vanuatu.

Local/Common/Vernacular names: *Palas Palm, The Ruffled Fan Palm, Vanuatu Fan Palm* (English)

Licuala paludosa Griff., Calcutta J. Nat. Hist. 5: 323.1844

Licuala paludosa var. *winkleriana* Becc., Webbia 5: 43.1921

Distribution: Karnataka (cultivated in gardens, Belagavi, Bengaluru), cultivated in gardens; native to SE Asia.

Licuala spinosa Wurmb, Verh. Batav. Genootsch. Kunsten 2: 474.1780; K. G. Bhat, Palms Karnataka 37. 2011.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens, wild in Andaman & Nicobar Islands; Cambodia, Malaysia, Myanmar, Philippines, Thailand, Vietnam.

Local/Common/Vernacular name: *The Mangrove Fan Palm* (English)

Livistona R.Br.

Livistona chinensis (Jacq.) R.Br. ex Mart., Hist. Nat. Palm. 3: 240.1838; B.D.Sharma et al., Fl. Karnataka Anal. 278. 1984.

Latania chinensis Jacq., Fragm. Bot. 16.1801

Saribus chinensis (Jacq.) Blume, Rumphia 2: 48.1838

Distribution: Karnataka (cultivated as an ornamental throughout), cultivated in gardens, probably wild in NE India; native to E Asia.

Local/Common/Vernacular names: *Chinese fan palm, Fountain palm* (English)

Livistona rotundifolia (Lam.) Mart., Hist. Nat. Palm. 3: 241.1838; K. G. Bhat, Palms Karnataka 35. 2011.

Corypha rotundifolia Lam., Encycl. 2: 131.1786

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated in gardens; native to Indonesia & Philippines.

Local/Common/Vernacular name: *Fountain palm* (English)

Normanbya F.Muell ex Becc.

Normanbya normanbyi (W.Hill) L.H.Bailey, Gentes Herb. 2: 188.1930; K. G. Bhat, Palms Karnataka 74. 2011.

Areca normanbyi (W.Hill) F.Muell., Fragm. 8: 235.1874

Cocos normanbyi W.Hill, Rep. Brisbane Bot. Gard. 1874: 6.1874

Ptychosperma normanbyi (W.Hill) F.Muell., Fragm. 11: 56.1880

Drymophloeus normanbyi (W.Hill) Benth. & Hook.f. ex Becc., Ann. Jard. Bot. Buitenzorg 2: 168.1885

Saguaster normanbyi (W.Hill) Kuntze, Revis. Gen. Pl. 2: 735.1891

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Australia.

Local/Common/Vernacular name: *Black palm* (English)

Phoenicophorium H.Wendl.

Phoenicophorium borsigianum (K.Koch) Stuntz, U.S.D.A. Bur. Pl. Industr. Invent. Seeds 31: 88.1914; K. G. Bhat, Palms Karnataka 74. 2011.

Astrocaryum borsigianum K.Koch, Wochenschr. Gärtnerie Pflanzenk. 2:401.1859

Phoenicophorium sechellarum H.Wendl., Ill. Hort. 12(Misc.): 5.1865

Areca sechellarum (H.Wendl.) Baill., Hist. Pl. 13: 348.1895

Stevensonia borsigiana (K.Koch) L.H.Bailey, Gentes Herb. 2:192.1930

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Seychelles Islands.

Local/Common/Vernacular names: *Latanier palm, Thief palm* (English)

Phoenix L.

Phoenix acaulis Roxb., Pl. Coromandel 3: 69.1820; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Phoenix acaulis var. *melanocarpa* Griff., Calcutta J. Nat. Hist. 5: 346.1845

Distribution: Karnataka (Mysuru, Uttara Kannada), Assam, Punjab, Uttar Pradesh, West Bengal; Myanmar, Nepal.

Phoenix canariensis Chabaud, Prov. Agric. Hort. Ill. 19: 293.1882; K. G. Bhat, Palms Karnataka 29. 2011.

Phoenix canariensis var. *porphyrococca* Vasc. & Franco, Portugaliae Act. Biol., Sér. B, Sist. 2: 313.1948

Distribution: Karnataka (cultivated in Soan's Form, Moodbidri, Dakshina Kannada), cultivated in gardens; native to Canary Islands.

Local/Common/Vernacular name: *Canary Island Date Palm* (English)

Phoenix dactylifera L., Sp. Pl. 1188.1753; K. G. Bhat, Palms Karnataka 30. 2011.

Palma dactylifera (L.) Mill., Gard. Dict. ed. 8 1.1768

Distribution: Karnataka (cultivated in gardens and farms), cultivated in Gujarat, Rajasthan and elsewhere in India; native to Middle East, Africa.

Local/Common/Vernacular names: *Date palm, Dates* (English); *Karjura* (Kannada)

Phoenix loureiroi Kunth, Enum. Pl. 3: 257.1841

Phoenix pusilla Lour., Fl. Cochinch. 614.1790

Phoenix ouseleyana Griff., Calcutta J. Nat. Hist. 5: 347.1845

Phoenix pedunculata Griff., Palms Brit. E. Ind. 139.1850

Phoenix humilis Royle ex Becc., Malesia 3: 373. 1890.

Phoenix humilis var. *loureiroi* (Kunth) Becc., Malesia 3: 382.1890

Phoenix humilis var. *pedunculata* (Griff.) Becc., Malesia 3: 384.1890

Phoenix loureiroi var. *pedunculata* (Griff.) Govaerts, World Checklist Palms 171.2005; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluruu, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Tumakur, Udupi, Uttara Kannada), Andhra Pradesh, Assam, Bihar, Himachal Pradesh, Kerala, Nagaland, Odisha, Tamil Nadu; Bangladesh, Bhutan, Cambodia, Laos, Myanmar, Nepal, Pakistan, Philippines, Thailand, Vietnam.

Phoenix paludosa Roxb., Fl. Ind. ed.2, 3: 789.1832; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Phoenix andamanensis W. Mill., J.G.Sm. & N.Taylor, Stand. Cycl. Hort. 5: 2594.1916

Distribution: Karnataka (Belagavi?), Andaman & Nicobar Islands, Odisha, West Bengal; Bangladesh, Indonesia, Malaysia, Myanmar.

Phoenix reclinata Jacq., Fragm. 1: 27.1801; K. G. Bhat, Palms Karnataka 30. 2011.

Phoenix spinosa Schumach. & Thonn., Beskr. Guin. Pl. 437.1827

Distribution: Karnataka (cultivated in Lalbagh, Bengaluru), cultivated in gardens; native to tropical Africa.

Local/Common/Vernacular name: *Senegal date palm* (English)

Phoenix roebelenii O'Brien, Gard. Chron. III, 6: 475.1889; K. G. Bhat, Palms Karnataka 31. 2011.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to S E Asia.

Local/Common/Vernacular names: *Miniature date palm, Pygmy date palm, Robellini* (English)

Phoenix sylvestris (L.) Roxb., Fl. Ind. ed. 1832 3: 787.1832; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Elate sylvestris L., Sp. Pl. 1189.1753

Distribution: Karnataka (throughout, more common in dry districts, also widely cultivated), throughout India except Himalayas.

Local/Common/Vernacular name: *Wild date palm* (English); *Eechalu mara* (Kannada)

Pinanga Blume

Pinanga dicksonii (Roxb.) Blume, Rumphia 2: 77.1839; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Plate 17B

Areca dicksonii Roxb., Fl. Ind. ed.2, 3: 616 1832

Seaforthia dicksonii (Roxb.) Mart., Hist. Nat. Palm. 3: 184.1838

Ptychosperma dicksonii (Roxb.) Miq., Fl. Ned. Ind. 3: 23.1855

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Kadu adike* (Kannada)

Pritchardia Seem. & H.Wendl.

Pritchardia pacifica Seem. & H.Wendl., Bonplandia (Hannover) 10: 197.1862; K. G. Bhat, Palms Karnataka 38 2011.

Distribution: Karnataka (cultivated in in gardens),widely cultivated in India; native to Tonga, Fiji & Samoa Islands.

Local/Common/Vernacular name: *Fiji Fan Palm* (English)

Ptychosperma Labill.

Ptychosperma elegans (R.Br.) Blume, Rumphia 2: 118.1843

Seaforthia elegans R.Br., Prodr. Fl. Nov. Holl. 267.1810

Archontophoenix elegans (R.Br.) F.M.Bailey, Cat. Pl. Queensland 50.1890

Saguaster elegans (R.Br.) Kuntze, Revis. Gen. Pl. 2: 735.1891

Distribution: Karnataka (Dakshina Kannada, Udupi), cultivated in gardens;native to Australia.

Local/Common/Vernacular name: *Solitaire palm* (English)

Ptychosperma macarthurii (H.Wendl. ex H.J.Veitch) H.Wendl. ex Hook.f., Rep. Progr. Condition Roy. Bot. Gard. Kew 1882: 55.1884

Kentia macarthurii H.Wendl. ex H.J.Veitch, Cat. Pl. 1879: 26.1879

Saguaster macarthurii (H.Wendl. ex H.J.Veitch) Kuntze, Revis. Gen. Pl. 2: 735 1891

Actinophloeus macarthurii (H.Wendl. ex H.J.Veitch) Becc. ex Raderm., Ann. Jard. Bot. Buitenzorg 35: 12.1925

Distribution: Karnataka (Dakshina Kannada, Udupi), cultivated in gardens;native to New Guinea.

Local/Common/Vernacular name: *Mac Arthur palm* (English)

Rhapis L. f. ex Aiton

Rhapis excelsa (Thunb.) Henry, J. Arnold Arbor. 11: 153.1930

Chamaerops excelsa Thunb., Syst. Veg. ed. 14 984.1784

Trachycarpus excelsus (Thunb.) H.Wendl., Bull. Soc. Bot. France 8: 429.1861

Distribution: Karnataka (Belagavi, Bengaluru, Mysuru), cultivated in gardens ; native to China & Taiwan.

Local/Common/Vernacular name: *Broadleaf Lady Palm* (English)

Rhapis humilis Blume, Rumphia 2: 54. 1839

Distribution: Karnataka (Belagavi), cultivated in gardens ; native to E Asia.

Local/Common/Vernacular name: *Slender Lady palm* (English)

Roystonea O.F.Cook

Roystonea regia (Kunth) O.F.Cook, Science II, 12: 479.1900; B.D.Sharma et al., Fl. Karnataka Anal. 295. 1984.

Oreodoxa regia Kunth, Nov. Gen. Sp. 1: 305.1816

Oenocarpus regius (Kunth) Spreng., Syst. Veg. 2: 140.1825

Distribution: Karnataka (cultivated throughout in gardens), cultivated in gardens;native to USA (Florida), Mexico, Central America & The Caribbean.

Local/Common/Vernacular names: *Bottle palm, Royal palm, Royal palm of Cuba* (English)

Sabal Adans.

Sabal mexicana Mart., Hist. Nat. Palm. 3: 246.1838

Inodes mexicana (Mart.) Standl., Contr. U. S. Natl. Herb. 13: 71.1920

Distribution: Karnataka (cultivated in in gardens, Belagavi), cultivated in gardens; native to N America.

Local/Common/Vernacular names: *Mexican Palmetto, Texas Palmetto, Texas Sabal Palm* (English)

Sabal minor (Jacq.) Pers., Syn. Pl. 1: 399.1805

Corypha minor Jacq., Hort. Bot. Vindob. 3: 8.1776

Distribution: Karnataka (cultivated in gardens Belagavi), cultivated in gardens; native to SE United States.

Local/Common/Vernacular names: *Bush palmetto, Dwarf Palmetto* (English)

Sabal palmetto (Walter) Lodd. ex Schult. & Schult.f., Syst. Veg. 7: 1487.1830; K. G. Bhat, Palms Karnataka 26. 2011.

Corypha palmetto Walter, Fl. Carol. 119. 1788

Chamaerops palmetto (Walter) Michx., Fl. Bor.-Amer. 1: 206.1803

Sabal blackburniana Schult. & Schult.f., Syst. Veg. 7: 1488.1830; K. G. Bhat, Palms Karnataka 31. 2011.

Inodes palmetto (Walter) O.F.Cook, Bull. Torrey Bot. Club 28: 532.1901

Distribution: Karnataka (cultivated in gardens Dharwad, Mysuru), cultivated in gardens; native to USA and tropical America.

Local/Common/Vernacular names: *Blue palmetto, Cabbage Palm, Cabbage palmetto, Carolina palmetto, Common palmetto Palmetto, Palmetto palm, Sabal palm, Swamp cabbage* (English)

Salacca Reinw.

Salacca zalacca (Gaertn.) Voss, Vilm. Blumengärtn. ed. 3 1: 1152.1895; K. G. Bhat, Palms Karnataka 11. 2011.

Calamus zalacca Gaertn., Fruct. Sem. Pl. 2: 267.1791

Salacca edulis Reinw., Syll. Pl. Nov. 2: 3.1825

Salacca rumphii Wall., Pl. Asiat. Rar. 3: t. 223.1831

Distribution: Karnataka (cultivated in Bengaluru, Dakshina Kannada), cultivated for fruits; native to Indonesia & Malaysia.

Local/Common/Vernacular name: *Salak palm* (English)

Syagrus Mart.

Syagrus romanzoffiana (Cham.) Glassman, Fieldiana, Bot. 31: 382.1968; K. G. Bhat, Palms Karnataka 56. 2011.

Cocos romanzoffiana Cham., Choris Voy. Pittor. (Chili) 5, t. 6.1822

Calappa romanzoffiana (Cham.) Kuntze, Revis. Gen. Pl. 2: 982.1891

Arecastrum romanzoffianum (Cham.) Becc., Agric. Colon. 10: 455.1916

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to South America.

Local/Common/Vernacular names: *Cocos Palm, Queen Palm* (English)

Syagrus schizophylla (Mart.) Glassman, Fieldiana, Bot. 31: 386.1968; K. G. Bhat, Palms Karnataka 57. 2011.

Cocos schizophylla Mart., Hist. Nat. Palm. 2: 119.1826

Calappa schizophylla (Mart.) Kuntze, Revis. Gen. Pl. 2: 982.1891

Arikury schizophylla (Mart.) Becc., Agric. Colon. 10: 445.1916

Arikuryroba schizophylla (Mart.) L.H.Bailey, Gentes Herb. 2: 196.1930

Distribution: Karnataka (cultivated in gardens of Bengaluru, Dharwad, Mysuru), cultivated in gardens; native to South America.

Local/Common/Vernacular name: *Arikury Palm* (English)

Thrinax L. f. ex Sw.

Thrinax parviflora Sw., Prodr. Veg. Ind. Occ. 57.1788; K. G. Bhat, Palms Karnataka 27. 2011.

Distribution: Karnataka (cultivated in gardens of Belagavi, Bengaluru), cultivated in gardens; native to Caribbean Islands.

Washingtonia H.Wendl.

Washingtonia filifera (Linden ex André) H.Wendl. ex de Bary, Bot. Zeitung (Berlin) 37: 61.1879; K. G. Bhat, Palms Karnataka 39. 2011.

Pritchardia filifera Linden ex André, Ill. Hort. 21: 28.1874

Brahea filifera (Linden ex André) W.Watson, Bull. Misc. Inform. Kew 1889: 296.1889

Neowashingtonia filifera (Linden ex André) Sudw., For. Trees Pacif. Slope 199.1908

Distribution: Karnataka (cultivated in gardens of Belagavi, Mysuru), cultivated in gardens; native to Western USA. Local/Common/Vernacular name: *Desert fan palm* (English)

Washingtonia robusta H.Wendl., Berliner Allg. Gartenzeitung 2: 198.1883

Neowashingtonia robusta (H.Wendl.) A.Heller, Cat. N. Amer. Pl. 3.1898

Washingtonia filifera var. *robusta* (H.Wendl.) Parish, Bot. Gaz. 44: 420.1907

Distribution: Karnataka (cultivated in gardens of Belagavi), cultivated in gardens; native to Mexico & USA.

Local/Common/Vernacular names: *Mexican Fan Palm, Mexican Washingtonia* (English)

Wodyetia A.K.Irvine

Wodyetia bifurcata A.K.Irvine, Principes 27: 163.1983; K. G. Bhat, Palms Karnataka 73. 2011.

Distribution: Karnataka (cultivated in gardens), cultivated as ornamental; native to Australia.

Local/Common/Vernacular name: *Foxtail palm* (English)

Aristolochiaceae

Aristolochia L.

Aristolochia acuminata Lam., Encycl. 1: 254.1783

Plate 4B

Aristolochia tagala Cham., Linnaea 7: 207.1832; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.

Aristolochia roxburghiana Klotzsch, Monatsber. Konigl. Preuss. Akad. Wiss. Berlin 1859: 596.1859

Aristolochia angulosa Wall. ex Duch., Prodr. 15(1): 480.1864

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Kolar, Raichur, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Peninsular to NE India; Australia, China, Indo-China, Pacific Islands, S & SE Asia.

Local/Common/Vernacular name: *Dodda eshwari balli* (Kannada)

Aristolochia bracteolata Lam., Encycl. 1: 258.1783; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.

Aristolochia bracteata Retz., Observ. Bot. 5: 29.1788

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chitradurga, Davanagere, Dharwad, Kalaburgi, Kodagu, Kolar, Raichur, Tumakuru, Uttara Kannada, Vijayapura, Yadgir), Andhra Pradesh, Delhi, Gujarat, Kerala, Madhya Pradesh, Odisha, Rajasthan Tamil Nadu; Pakistan, Bangladesh, Sri Lanka, Saudi Arabia & tropical Africa.

Local/Common/Vernacular names: *Kathe kiribina gida, Kuri gida, Sanjali hallu* (Kannada); *Kitamari* (Sanskrit)

Aristolochia grandis Craib, Bull. Misc. Inform. Kew 1914: 10 1914

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated as ornamental; native to tropical America.

Aristolochia indica L., Sp. Pl. 960.1753; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.

Plate 4C

Aristolochia lanceolata Wight, Icon. Pl. Ind. Orient. 5: t. 1858.1852

Aristolochia indica var. *lanceolata* (Wight) Duch., Prodr. 15(1): 479.1864

Aristolochia maysorensis Fisch. ex Duch., Prodr. 15(1): 479.1864

Aristolochia pandurata Wall. ex Duch., Prodr. 15(1): 479.1864

Distribution: Karnataka (throughout) throughout plains of India; Bangladesh, Bhutan, China, Pakistan, Sri Lanka. Local/Common/Vernacular names: *Eshwari balli, Eshwari beru, Eshwari beru balli* (Kannada); *Arkamula, Eswari* (Sanskrit)

Aristolochia labiata Willd., Mém. Soc. Imp. Naturalistes Moscou 2: 101.1809
Aristolochia ringens Link & Otto, Icon. Pl. Select. 33.1821
Aristolochia brasiliensis Mart., Nov. Gen. Sp. Pl. 1: 77.1824
Aristolochia ornithocephala Hook., Bot. Mag. 70: t. 4120 1844; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.
Howardia brasiliensis (Mart. & Zucc.) Klotzsch, Monatsber. Konigl. Preuss. Akad. Wiss. Berlin 1859: 608, t. 2, f. 14.1859
 Distribution: Karnataka (cultivated in gardens of Bengaluru, Kalaburgi, Kodagu, Mysuru, Yadgir) also cultivated in many states, native to tropical America.
 Local/Common/Vernacular names: *Mottled Dutchman's Pipe, Pelicanflower, Rooster Flower* (English)

Aristolochia littoralis Parodi, Anales Soc. Ci. Argent. 5: 155.1878
Aristolochia elegans Mast., Gard. Chron. n.s., 1885(2): 301.1885; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.
 Distribution: Karnataka (cultivated Bengaluru, Mysuru), cultivated as ornamental; native to Brazil.
 Local/Common/Vernacular names: *Calico flower, Elegant Dutchman's pipe* (English)

Thottea Rottb.

Thottea dalzellii (Hook.f.) Karthik. & Moorthy, Fl. Pl. India 156.2009
Bragantia dalzellii Hook.f., Fl. Brit. India 5: 73.1886
Apama dalzellii (Hook.f.) Kuntze, Revis. Gen. Pl. 1: 63.1891
 Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.
 Local/Common/Vernacular name: *Chakrani* (Kannada)

Thottea siliquosa (Lam.) Ding Hou, Blumea 27: 327.1981
Apama siliquosa Lam., Encycl. 1: 91 1783; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.
Bragantia wallichii R. Br. ex Wight & Arn., Cat. Ind. Pl. 57.1833
 Distribution: Karnataka (Chikmagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka
 Local/Common/Vernacular names: *Chakrapani, Charnike gida, Neeratte beru* (Kannada)

Thottea sivarajanii E.S.S.Kumar, A.E.S.Khan & Binu, Rheedia 10(2): 117.2000
 Distribution: Karnataka (Kodagu), Kerala; endemic.

Asparagaceae

Agave L.

Agave americana L., Sp. Pl. 323.1753; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.
Aloe americana (L.) Crantz, Inst. Rei Herb. 1: 466.1766
Agave ingens A.Berger, Hortus Mortolensis 12.1912; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.
Agave americana var. *marginata* Trel., Stand. Cycl. Hort. 1: 235.1914
 Distribution: Karnataka (Widely cultivated and naturalised throughout), throughout hotter parts of India; native to Mexico.
 Local/Common/Vernacular names: *Century plant* (English); *Ane kathale, Bhutale, Boodu kattale, Kannaru, Rakspatti* (Kannada)

Agave cantala Roxb. ex Salm-Dyck, Index Pl. Succ. Hort. Dyck. 1.1822; Roxb., Fl. Ind. ed.2, 2: 167.1832 ('cantula'); B.D.Sharma et al., Fl. Karnataka Anal. 283.1984.
 Distribution: Karnataka (Belagavi), throughout plains of India; native range obscure, probably native to C & Southern N America.
 Local/Common/Vernacular names: *Cantula, Cantula agave* (English)

Agave sisalana Perrine, Cogr. Doc. 564: 87.1838; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Agave rigida var. *sisalana* (Perrine) Engelm., Trans. Acad. Sci. St. Louis 3: 316.1875

Agave rigida var. *sisalana* Baker, Bull. Misc. Inform. Kew 1889: 254. 1889

Distribution: Karnataka (cultivated throughout), throughout plains of India; native to Mexico.

Local/Common/Vernacular names: *Aane kattaale*, *Kattaale* (Kannada)

Agave vera-cruz Mill., Gard. Dict. ed. 8,7 1768; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Distribution: Karnataka (Belagavi, Ballari), throughout plains of India; native to Mexico.

Agave vivipara L., Sp. Pl. 323.1753

Aloe vivipara (L.) Crantz, Inst. Rei Herb. 1: 466. 1766

Agave cuspidata Baker, Gard. Chron. n.s., 1877: 137.1877

Agave wightii J.R.Drumm. & Prain, Bengal Agric. Bull. 8: 15.1906

Distribution: Karnataka (cultivated and naturalised throughout), cultivated and naturalised throughout India; native to C America.

Asparagus L.

Asparagus aethiopicus L., Mant. Pl. 1.1770

Asparagopsis aethiopica (L.) Kunth, Enum. Pl. 5: 95.1850

Asparagus sprengeri Regel, Trudy Imp. S.-Peterburgsk. Bot. Sada 11: 302.1890

Protasparagus aethiopicus (L.) Oberm., S. African J. Bot. 2: 243.1983

Distribution: Karnataka (cultivated ornamental), cultivated ornamental; native to S Africa.

Asparagus africanus Lam., Encycl. 1: 295.1783

Protasparagus africanus (Lam.) Oberm., S. African J. Bot. 2: 243.1983

Distribution: Karnataka (Belagavi, Chitradurga), cultivated throughout; native tropical Africa & Middle East.

Asparagus asiaticus L., Sp. Pl. 313.1753; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Protasparagus asiaticus (L.) Kamble, J. Econ. Taxon. Bot. 15: 708 1992

Distribution: Karnataka (Ballari, Chitradurga, Dharwad, Vijayapura), throughout India; Africa, Australia, Madagascar.

Asparagus gonoclados Baker, J. Linn. Soc., Bot. 14: 627.1875; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.

Protasparagus gonoclados (Baker) Kamble, J. Econ. Taxon. Bot. 15: 709.1992

Distribution: Karnataka (throughout, sometimes cultivated as an ornamental), peninsular India; endemic.

Local/Common/Vernacular names: *Mahashatavari*, *Shathamooli* (Sanskrit)

Asparagus laevissimus Steud. ex Baker, J. Linn. Soc., Bot. 14: 623 1875; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.

Protasparagus laevissimus (Steud. ex Baker) Kamble, J. Econ. Taxon. Bot. 15: 709.1992

Distribution: Karnataka (Belagavi, Ballari, Chitradurga, Dharwad, Kalburgi, Mysuru), Andhra Pradesh, Kerala; endemic.

Asparagus officinalis L., Sp. Pl. 313.1753

Asparagus polyphyllus Steven, Bull. Soc. Imp. Naturalistes Moscou 30: 343.1857

Distribution: Karnataka (Belagavi), cultivated elsewhere; native to NW Africa, C & SW Asia, Europe.

Local/Common/Vernacular names: *Asparagus*, *Garden asparagus*, *Sparrow grass* (English)

Asparagus racemosus Willd., Sp. Pl. 2: 152.1799; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.
Asparagus acerosus Roxb., Fl. Ind. ed.2, 2: 150.1832
Asparagopsis hohenackeri Kunth, Enum. Pl. 5: 100 1850
Asparagopsis sarmentosa Dalzell & A.Gibson, Bombay Fl. 246.1861
Asparagus jacquemontii Baker, J. Linn. Soc., Bot. 14: 615.1875
Asparagus racemosus var. *javanicus* (Kunth) Baker, J. Linn. Soc., Bot. 14: 624.1875
Asparagus racemosus var. *zeylanicus* Baker, J. Linn. Soc., Bot. 14: 623 1875
Asparagus zeylanicus (Baker) Hook.f., Fl. Brit. India 6: 317.1892
Protasparagus jacquemontii (Baker) Kamble, J. Econ. Taxon. Bot. 15: 709.1992
Protasparagus racemosus var. *javanicus* (Kunth) Kamble, J. Econ. Taxon. Bot. 15: 709.1992
Protasparagus racemosus var. *subacerosus* (Baker) Kamble, J. Econ. Taxon. Bot. 15: 709.1992
Protasparagus zeylanicus (Hook.f.) Kamble, J. Econ. Taxon. Bot. 20: 269.1996
 Distribution: Karnataka (throughout), throughout India; pantropical.
 Local/Common/Vernacular names: *Halavu makkala balli*, *Halavu makkala tayi beru*, *Halavu makkala tayi gida*, *Satavari*, *Shatavari* (Kannada); *Shatavari* (Sanskrit)

Asparagus setaceus (Kunth) Jessop, Bothalia 9: 51.1966
Asparagus plumosus Baker, J. Linn. Soc., Bot. 14: 613.1875; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.
Protasparagus plumosus (Baker) Oberm., S. African J. Bot. 2: 244.1983
 Distribution: Karnataka (cultivated in gardens Belagavi, Bengaluru, Mysuru, Tumakuru); native to S Africa, cultivated elsewhere.
 Local/Common/Vernacular names: *Climbing asparagus*, *Common asparagus fern*, *Ferny asparagus*, *Lace fern* (English)

Beaucarnea Lem.
Nolina Michaux

Beaucarnea recurvata Lem., Ill. Hort. Misc.: 59.1861
Nolina recurvata (Lem.) Hemsl., Biol. Cent.-Amer., Bot. 3: 372. 1884
Dasyllirion recurvatum (Lem.) J.F.Macbr., Contr. Gray Herb. 56: 17.1918
 Distribution: Karnataka (cultivated as an ornamental), cultivated as an ornamental throughout India; native to Mexico.
 Local/Common/Vernacular names: *Elephant's foot*, *Ponytail palm* (English)

Chlorophytum Ker-Gawler

Chlorophytum arundinaceum Baker, J. Linn. Soc., Bot. 15: 323.1876; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.
Anthericum tuberosum Hook.f., Fl. Brit. India 6: 334.1892
 Distribution: Karnataka (Kalburgi, Hassan, Kodagu, Mysuru, Tumakuru, Yadgir), tropical parts of India; Myanmar.

Chlorophytum belagaumense Chandore., Malpure, Adsul & S.R.Yadav, Kew Bull. 67(3): 527. 2012
 Distribution: Karnataka (Belagavi); endemic.

Chlorophytum bharuchae Ansari, Sundararagh. & Hemadri, Indian Forester 96: 304. 1970; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984. Plate 17D
Chlorophytum borivilianum Santapau & R.R.Fern., J. Bombay Nat. Hist. Soc. 3: 898.1955
 Distribution: Karnataka (Belagavi, Shivamogga); endemic.
 Distribution: Karnataka (cultivated in Belagavi), Maharashtra; endemic.

Chlorophytum breviscapum Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 141. 1850 non Dammer 1912; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi); endemic.

Chlorophytum capense (L.) Voss, Vilm. Blumengärtn. Ed. 3 1: 1086.1895

Asphodelus capensis L., Syst. Nat. Ed. 10 2: 982.1759

Distribution: Karnataka (cultivated in most districts), cultivated throughout; native to Africa.

Chlorophytum comosum (Thunb.) Jacques, J. Soc. Imp. Centr. Hort. 8: 345.1862

Anthericum comosum Thunb., Prodr. Pl. Cap. 63.1794

Phalangium comosum (Thunb.) Poir., Encycl. 5: 252.1804

Caesia comosa (Thunb.) Spreng., Syst. Veg. 2: 88.1825

Hartwegia comosa (Thunb.) Nees, Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 15(2): 378.1831

Hollia comosa (Thunb.) Heynh., Alph. Aufz. Gew. 303.1846

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Africa.

Local/Common/Vernacular names: *Airplane plant, Ribbon plant, spider ivy, Spider plant, St. Bernard's lily* (English)

Chlorophytum filipendulum Baker, Trans. Linn. Soc. London, Bot. 1: 260. 1878.

subsp. **amanicense** (Engl.) Nordal & A.D.Poulsen in Fl. Trop. E. Afr. Antheric.: 59.1998

Chlorophytum amaniense Engl., Bot. Jahrb. Syst. 34: 157.1904

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Africa.

Chlorophytum glaucum Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 142.1850; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984. Plate 17E

Anthericum nimmonii J.Graham, Cat. Pl. Bombay 220.1839

Hartwegia nimmonii Dalzell ex Baker, J. Linn. Soc., Bot. 15: 324.1876

Distribution: Karnataka (Belagavi, Chikkamagaluru, Mysuru, Tumakuru, Uttara Kannada), Maharashtra; endemic.

Chlorophytum heynei Baker, J. Linn. Soc., Bot. 15: 322.1876; Bhaskar & Kushalappa, Fl. Tumkur Dist. 105. 2013.

Anthericum heynei Rottler ex Baker, J. Linn. Soc., Bot. 15: 323.1876

Chlorophytum heyneanum Wall. ex Hook.f., Fl. Brit. India 6: 333.1892

Distribution: Karnataka (Tumakuru, Uttara Kannada), Goa, Maharashtra; endemic.

Chlorophytum indicum (Willd. ex Schult. & Schult.f.) Dress, Baileya 9: 43.1961

Plate 17F

Ornithogalum indicum Willd. ex Schult. & Schult.f., Syst. Veg. 7: 535.1829

Phalangium indicum (Willd. ex Schult. & Schult.f.) Kunth, Enum. Pl. 4: 598.1843

Liliago indica (Willd. ex Schult. & Schult.f.) C.Presl, Abh. Königl. Böhm. Ges. Wiss. IV, 3: 534.1845

Phalangium attenuatum Wight, Icon. Pl. Ind. Orient. T. 2037.1853

Chlorophytum attenuatum (Wight) Baker, J. Linn. Soc., Bot. 15: 332.1876; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Kolar, Mysuru, Shivamogga, Tumakuru), Kerala; endemic.

Chlorophytum kolhapurense Sardesai, S.P.Gaikwad & S.R.Yadav, Kew Bull. 61: 269.2006 Plate 17G

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Chlorophytum laxum R.Br., Prodr. Fl. Nov. Holl. 277.1810; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.

Phalangium parviflorum Wight, Icon. Pl. Ind. Orient. T. 2039.1853

Anthericum parviflorum (Wight) Benth., Fl. Hongk. 373.1861

Phalangium laxum (R.Br.) F.Muell., Fragm. 7: 71.1869

Chlorophytum acaule Baker, J. Linn. Soc., Bot. 15: 327.1876; B.D.Sharma et al., Fl. Karnataka Anal. 285. 1984.

Distribution: Karnataka (throughout), throughout India; Paleotropical.

Local/Common/Vernacular name: *Sanna dravanti* (Kannada)

Chlorophytum malabaricum Baker, J. Linn. Soc., Bot. 15: 331.1876; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Plate 18A

Phalangium tuberosum Dalzell & A.Gibson, Bombay Fl. 251.1861

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Chitradurga, Kodagu, Shivamogga, Tumakuru), peninsular India; endemic.

Chlorophytum nimmonii Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 142 1850; B.D.Sharma et al., Fl. Karnataka Anal. 2786. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Dharwad, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Chlorophytum orchidastrum Lindl., Edwards's Bot. Reg. 10: t. 813.1824; Ramaswamy, M.R. Rao & Govindappa, Fl. Shivamogga Dist. 616. 2001.

Phalangium oligospermum Wight, Icon. Pl. Ind. Orient. t. 2038.1853

Chlorophytum petiolatum Baker, J. Linn. Soc., Bot. 15: 326.1876

Chlorophytum talbotii Rendle, Cat. Pl. Oban 113.1913

Distribution: Karnataka (cultivated Bengaluru, Shivamogga), cultivated in gardens of India; native W Africa.

Chlorophytum tuberosum (Roxb.) Baker, J. Linn. Soc., Bot. 15: 332.1876; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Anthericum tuberosum Roxb., Pl. Coromandel 2: 20. 1800.

Phalangium tuberosum (Roxb.) Kunth, Enum. Pl. 4: 598.1843

Liliago tuberosa (Roxb.) C.Presl, Abh. Königl. Böhm. Ges. Wiss. IV, 3: 534.1845

Chlorophytum anthericoideum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 141.1850

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Vijayapura, Chikkamagaluru, Kalburgi, Shivamogga, Uttara Kannada, Yadgir), Peninsular India; Africa, S Asia.

Local/Common/Vernacular name: *Dravanti* (Kannada); *Musli* (Sanskrit)

Cordyline Comm. ex R. Br.

Cordyline australis (G.Forst.) Endl., Prodr. Fl. Norfolk. 29 1833

Dracaena australis G.Forst., Fl. Ins. Austr. 24 1786

Distribution: Karnataka (cultivated in gardens), cultivated in gardens throughout; native to New Zealand.

Cordyline stricta (Sims) Endl., Ann. Wiener Mus. Naturgesch. 1: 162.1836

Dracaena stricta Sims, Bot. Mag. 52: t. 2575.1825

Charlwoodia stricta (Sims) Sweet, Fl. Australas. t. 18.1827

Terminalis stricta (Sims) Kuntze, Revis. Gen. Pl. 2: 716.1891

Taetsia stricta (Sims) Standl., Lista Pl. Salvador 50.1925

Distribution: Karnataka (Belagavi), cultivated in gardens throughout; native to Australia.

Cordyline terminalis (L.) Kunth, Abh. Königl. Akad. Wiss. Berlin 1842: 30.1844

Convallaria fruticosa L., Herb. Amb. 16 1754

Dracaena terminalis L., Syst. Nat. ed. 12 2: 246.1767

Taetsia terminalis (L.) W.Wight, Contr. U. S. Natl. Herb. 9: 382.1905

Distribution: Karnataka (Belagavi), cultivated in gardens throughout; native to South Sea Islands.

Dipcadi Medik.

Dipcadi montanum (Dalzell) Baker, J. Linn. Soc., Bot. 11: 398.1870; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Uropetalum montanum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 142 142.1850

Distribution: Karnataka (Belagavi, Kalburgi, Yadgir), Kerala, Maharashtra, Tamil Nadu

Dracaena Vand. ex L.

Dracaena aletriformis (Haw.) Bos in Fl. S. Afr. 5(3): 3.1992

Yucca aletriformis Haw., Philos. Mag. Ann. Chem. 5: 415 1831

Dracaena hookeriana K.Koch, Wochenschr. Vereines Beförd. Gartenbaues Königl. Preuss. Staaten 4: 394.1861; Seetharam, Kotresh & Uplaconkar, Fl. Dist 90. 2000.

Pleomele hookeriana (Kuntze) N.E.Br., Bull. Misc. Inform. Kew 1914: 278.1914

Distribution: Karnataka (cultivated in Kalburgi, Yadgir)

Dracaena braunii Engl., Bot. Jahrb. Syst. 15: 479.1892

Pleomele braunii (Engl.) N.E.Br., Bull. Misc. Inform. Kew 1914: 277.1914

Pleomele sanderiana (Sander) N.E.Br., Bull. Misc. Inform. Kew 1914: 279.1914

Distribution: Karnataka (cultivated in gardens), native to tropical E & W Africa.

Local/Common/Vernacular names: *Lucky bamboo, Ribbon plant* (English)

Dracaena draco (L.) L., Syst. Nat. ed. 12, 2: 246.1767

Asparagus draco L., Sp. Pl. ed. 2, 451.1762

Palma draco (L.) Mill., Gard. Dict. ed. 8, 11.1768

Stoerkia draco (L.) Crantz, Duab. Drac. Arbor. 25.1768

Drakaina draco (L.) Raf., Fl. Tellur. 4: 17.1838

Yucca draco (L.) Carrière, Rev. Hort. IV, 8: 389.1859

Distribution: Karnataka (Belagavi), cultivated in gardens throughout; native to Canary Islands.

Dracaena fragrans (L.) Ker Gawl., Bot. Mag. 27: t. 1081.1808

Aletris fragrans L., Sp. Pl. ed. 2, 456.1762

Pleomele fragrans (L.) Salisb., Prodr. Stirp. Chap. Allerton 245.1796

Sansevieria fragrans (L.) Jacq., Fragm. Bot. 29.1801

Dracaena smithii Baker ex Hook.f., Bot. Mag. 101: t. 6169.1875

Draco fragrans (L.) Kuntze, Revis. Gen. Pl. 2: 710.1891

Dracaena deremensis Engl., Bot. Jahrb. Syst. 32: 95.1902

Distribution: Karnataka (cultivated in gardens throughout), cultivated in gardens throughout; native to tropical E & W Africa.

Dracaena reflexa Lam., Encycl. 2: 324.1786; Seetharam, Kotresh & Uplaconkar, Fl. Dist 90. 2000.

Lomatophyllum reflexum (Lam.) Bojer, Hortus Maurit. 349.1837

Cordyline reflexa (Lam.) Endl., Cat. Horti Vindob. 1: 148.1842

Draco reflexa (Lam.) Kuntze, Revis. Gen. Pl. 2: 710.1891

Pleomele reflexa (Lam.) N.E.Br., Bull. Misc. Inform. Kew 1914: 279.1914

var. **angustifolia** Baker, J. Linn. Soc., Bot. 14: 531.1875

Cordyline marginata (Lam.) Endl., Cat. Horti Vindob. 1: 148.1842; Seetharam, Kotresh & Uplaonkar, Fl. Dist 90. 2000.

Draco marginata (Lam.) Kuntze, Revis. Gen. Pl. 2: 710.1891

Pleomele marginata (Lam.) N.E.Br., Bull. Misc. Inform. Kew 1914: 278.1914

Distribution: Karnataka (cultivated in Bengaluru, Kalburgi, Yadgir), cultivated in gardens; native to Madagascar and other Indian ocean Islands.

Dracaena spicata Roxb., Fl. Ind. ed.2, 2: 157.1832; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Dracaena wallichii Kunth, Abh. Königl. Akad. Wiss. Berlin 1842: 26.1844

Draco spicata (Roxb.) Kuntze, Revis. Gen. Pl. 2: 710.1891

Pleomele spicata (Roxb.) N.E.Br., Bull. Misc. Inform. Kew 1914: 279.1914

Distribution: Karnataka (Kodagu, Mysuru, Uttara Kannada), Andaman & Nicobar Islands. Assam Bangladesh, Myanmar.

Dracaena terniflora Roxb., Fl. Ind. ed. 1832 2: 159.1832; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Dracaena terniflora var. *heyneana* Hook.f., Fl. Brit. India 6: 329.1892

Pleomele terniflora (Roxb.) Merr., J. Arnold Arbor. 23: 157.1942

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Chitradurga, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Local/Common/Vernacular name: *Datuche zad* (*Kunabi*)

Drimia Jacq. ex Willd.

Drimia congesta Bullock, Bull. Misc. Inform. Kew 1932: 504.1932; Bhaskar & Kushalappa, Fl. Tumkur Dist. 171. 2013.

Distribution: Karnataka (Tumakuru)?

Drimia indica (Roxb.) Jessop, J. S. African Bot. 43: 272.1977; B.D.Sharma et al., Fl. Karnataka Anal. 278. 1984.

Erythronium indicum Rottler ex Spreng., Syst. Veg. 2: 97.1825

Scilla coromandeliana Roxb., Fl. Ind. ed. 2, 2: 147.1832

Scilla indica Roxb., Fl. Ind. ed. 2,2: 147 1832

Melanthium hyacinthoides Royle, Ill. Bot. Himal. Mts. 384.1839

Erythronium hyacinthoides Royle, Ill. Bot. Himal. Mts. 453.1840

Urginea indica (Roxb.) Kunth, Enum. Pl. 4: 333.1843

Urginea coromandeliana Wight, Icon. Pl. Ind. Orient. t. 2064.1853

Urginea wightiana Hook.f., Fl. Brit. India 6: 347.1892

Urginea govindappae A.Boraiah & Fathima, Bull. Bot. Surv. India 12: 128.1972

Urginea nagarjunae Hemadri & Swahari, Ancient Sci. Life 2: 105.1982

Drimia nagarjunae (Hemadri & Swahari) Anand Kumar, J. Econ. Taxon. Bot. 5: 962.1984

Thuranthos coromandelianus (Roxb.) Speta, Phyton (Horn) 38: 84.1998

Thuranthos govindappae (A.Boraiah & Fathima) Speta, Phyton (Horn) 38: 84. 1998

Thuranthos indicus (Roxb.) Speta, Phyton (Horn) 38: 84.1998

Thuranthos nagarjunae (Hemadri & Swahari) Speta, Phyton (Horn) 38: 84.1998

Thuranthos wightianus (Hook.f.) Speta, Phyton (Horn) 38: 85.1998

Indurgia coromandeliana (Roxb.) Speta, Staphia 75: 170.2001

Indurgia govindappae (A.Boraiah & Fathima) Speta, Staphia 75: 170.2001

Indurgia indica (Roxb.) Speta, Staphia 75: 170.2001

Indurgia nagarjunae (Hemadri & Swahari) Speta, Staphia 75: 170.2001

Indurgia wightiana (Hook.f.) Speta, Staphia 75: 170.2001

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Tumakuru, Udupi, Uttara Kannada), throughout tropical parts of India.

Local/Common/Vernacular names: *Adavi eerulli*, *Kaadu eerulli*, *Kaadu neerulli* (Kannada)

Drimia polyantha (Blatt. & McCann) Stearn, Ann. Mus. Goulandris 4: 209.1978; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Plate 18B

Urginea polyantha Blatt. & McCann, J. Bombay Nat. Hist. Soc. 32: 735.1928

Thuranthos polyanthus (Blatt. & McCann) Speta, Phyton (Horn) 38: 84.1998

Indurgia polyantha (Blatt. & McCann) Speta, Staphia 75: 170.2001

Distribution: Karnataka (Belagavi, Mysuru, Uttara Kannada), Maharashtra, Tamil Nadu; endemic.

Drimia wightii Lakshmin., Kew Bull. 58: 507.2003

Urginea congesta Wight, Icon. Pl. Ind. Orient. t. 2064.1853

Drimia congesta (Wight) Stearn, Ann. Mus. Goulandris 4: 208. 1978.

Thuranthos congestus (Wight) Speta, Phyton (Horn) 38: 81 1998

Indurgia congesta (Wight) Speta, Staphia 75: 170.2001

Distribution: Karnataka (Belagavi, Tumakuru), Andhra Pradesh, Maharashtra, Tamil Nadu; endemic.

Furcraea Vent.

Furcraea foetida (L.) Haw., Syn. Pl. Succ. 78.1812; B.D.Sharma et al., Fl. Karnataka Anal. 278. 1984.

Agave foetida L., Sp. Pl. 323.1753

Aloe foetida (L.) Crantz, Inst. Rei Herb. 1: 466.1766

Furcraea gigantea Vent., Bull. Soc. Philom. Paris 1: 65.1793

Agave gigantea (Vent.) D.Dietr., Syn. Pl. 2: 1192.1840

Fourcroya gigantea (Vent.) Hook., Bot. Mag. 86: t. 5163.1860

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), cultivated throughout; native to C America.

Local/Common/Vernacular name: *Seeme kattale* (Kannada)

Furcraea selloa K.Koch., Wochenschr. Vereines Beford. Gartenbaues Konigl. Preuss. Staaten 3: 22. 1860

Distribution: Karnataka (cultivated as an ornamental), cultivated as an ornamental throughout India; native to America.

Leedebouria Roth

Leedebouria karnatakensis Punekar & Lakshmin., Fl. Anshi Natatl. Park 500. 2011

Distribution: Karnataka (Uttara Kannada); endemic.

Leedebouria revoluta (L.f.) Jessop, J. S. African Bot. 36: 255.1970

Leedebouria hyacinthina Roth, Nov. Pl. Sp. 195.1821

Barnardia indica Wight, Icon. Pl. Ind. Orient. 6: t. 2041.1853

Scilla indica (Wight) Baker, Refug. Bot. 3(App.): 12. 1870.

Scilla hyacinthina (Roth) J.F.Macbr., Contr. Gray Herb. 56: 14.1918; B.D.Sharma et al., Fl. Karnataka Anal. 287. 1984.

Distribution: Karnataka (Belagavi, Bidar, Chitradurga, Dharwad, Kalburgi, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), Peninsular India; tropical E Africa.

Local/Common/Vernacular names: *Adavi ulla gaddi, Kadu bellulli* (Kannada)

Ophiopogon Ker-Gawl.

Ophiopogon intermedius D.Don, Prodr. Fl. Nepal. 48.1825; B.D.Sharma et al., Fl. Karnataka Anal. 281. 1984.

Flueggea japonica var. *intermedia* (D.Don) Schult.f., Syst. Veg. 7: 310.1829

Flueggea intermedia (D.Don) Kunth, Enum. Pl. 5: 306.1850

Flueggea wallichiana Kunth, Enum. Pl. 5: 303.1850

Ophiopogon indicus Wight, Icon. Pl. Ind. Orient. t. 2050.1853

Ophiopogon japonicus var. *intermedius* (D.Don) Maxim., Bull. Acad. Imp. Sci. Saint-Pétersbourg 15: 89.1871

Ophiopogon japonicus var. *wallichianus* (Kunth) Maxim., Bull. Acad. Imp. Sci. Saint-Petersbourg 7: 321.1872

Flueggea griffithii Baker, J. Linn. Soc., Bot. 17: 502.1879

Ophiopogon griffithii (Baker) Hook.f., Fl. Brit. India 6: 270.1892

Ophiopogon intermedius var. *gracilipes* Hook.f., Fl. Brit. India 6: 269.1892

Ophiopogon intermedius var. *macranthus* Hook.f., Fl. Brit. India 6: 269.1892

Ophiopogon intermedius var. *occidentalis* Hook.f., Fl. Brit. India 6: 269.1892

Ophiopogon intermedius var. *parviflorus* Hook.f., Fl. Brit. India 6: 269.1892

Ophiopogon intermedius var. *pauciflorus* Hook.f., Fl. Brit. India 6: 269.1892

Ophiopogon wallichianus (Kunth) Hook.f., Fl. Brit. India 6: 268.1892

Ophiopogon wallichianus var. *pauciflorus* Hook.f., Fl. Brit. India 6: 269.1892

Mondo japonicum var. *griffithii* (Baker) Farw., Amer. Midl. Naturalist 7: 42.1921

Mondo japonicum var. *intermedium* (D.Don) Farw., Amer. Midl. Naturalist 7: 42.1921

Mondo japonicum var. *wallichianum* (Kunth) Farw., Amer. Midl. Naturalist 7: 42.1921

Mondo intermedium (D.Don) L.H.Bailey, Gentes Herb. 2: 25.1929

Mondo wallichianum (Kunth) L.H.Bailey, Gentes Herb. 2: 21.1929

Ophiopogon parviflorus (Hook.f.) H.Hara, J. Jap. Bot. 40: 21.1965

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kalburgi, Mysuru, Shivamogga, Udupi, Uttara Kannada, Yadgir), throughout tropical parts of India; native to C America .

Peliosanthes Andrews

Peliosanthes teta Andrews, Bot. Repos. 10: t. 605. 1810.

subsp. **humilis** (Andrews) Jessop ex Gandhi, Fl. Hassan Distr. 799.1976; B.D.Sharma et al., Fl. Karnataka Anal. 281. 1984.

Peliosanthes humilis Andrews, Bot. Repos. 10: t. 634.1811

Peliosanthes courtallensis Wight, Icon. Pl. Ind. Orient. t. 2051.1853

Peliosanthes campanulata Wall. ex Baker, J. Linn. Soc., Bot. 17: 504.1879

Lourya campanulata Baill., Bull. Mens. Soc. Linn. Paris 1: 743.1888

Distribution: Karnataka (Chikkamagaluruu, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi), Kerala, Tamil Nadu; S & SE Asia.

Polianthes L.

Polianthes tuberosa L., Sp. Pl. 316.1753; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Agave tuberosa (L.) Thiede & Eggli, Kakteen And. Sukk. 50: 112.1999

Distribution: Karnataka (Widely cultivated), cultivated throughout plains of India; native to Mexico.

Local/Common/Vernacular name: *Sugandharaja* (Kannada)

Sansevieria Thunb.

Sansevieria cylindrica Bojer ex Hook., Bot. Mag. 85: t. 5093.1859; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Acyntha cylindrica (Bojer ex Hook.) Kuntze, Revis. Gen. Pl. 2: 699.1891

Cordyline cylindrica (Bojer ex Hook.) Britton, Bot. Porto Rico 5: 150.1923

Distribution: Karnataka (Belagavi, Bengaluru, Kalburgi, Vijayapura, Yadgir), cultivated in drier parts of India; native to tropical Africa.

Sansevieria hyacinthoides (L.) Druce, Rep. Bot. Exch. Club Soc. Brit. Isles 3: 423. 1914

Aloe hyacinthoides L., Sp. Pl. 321.1753

Aletris hyacinthoides (L.) L., Sp. Pl. ed. 2, 456.1762

Sanseverinia thyrsiflora Petagna, Inst. Bot. 3: 643.1787

Sansevieria thyrsiflora (Petagna) Thunb., Prodr. Pl. Cap. 65.1794

Acyntha thyrsiflora (Thunb.) Kuntze, Revis. Gen. Pl. 2: 699.1891

Cordyline hyacinthoides (L.) W.Wight, Contr. U. S. Natl. Herb. 9: 249.1904

Distribution: Karnataka (Belagavi), cultivated in drier parts of India; native to tropical Africa.

Sansevieria roxburghiana Schult. & Schult.f., Syst. Veg. 7: 357.1829; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Sansevieria zeylanica Roxb., Pl. Coromandel 2: 43, t.184.1805

Acyntha roxburghiana (Schult. & Schult.f.) Kuntze, Revis. Gen. Pl. 2: 699.1891

Cordyline roxburghiana (Schult. & Schult.f.) Merr., Enum. Philipp. Fl. Pl. 1: 206. 1923

Distribution: Karnataka (Belagavi, Bengaluru, Chikkaballapur, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kalburgi, Kolar, Mysuru, Tumakuru, Udupi, Uttara Kannada, Yadgir also cultivated as an ornamental), Goa, Kerala, Maharashtra, Tamil Nadu; Africa, China, Indonesia, Sri Lanka.

Local/Common/Vernacular name: *Manjina naaru* (Kannada)

Sansevieria stuckyi God.-Leb., Sansev. Gigant. Afr. Or. 13.1903

Acyntha stuckyi (God.-Leb.) Chiov., Fl. Somala 2: 422.1932

Distribution: Karnataka (cultivated in gardens), cultivated in gardens and parks in India; native to Africa..

Sansevieria trifasciata Prain, Bengal Pl. 2: 1054.1903

Aletris hyacinthoides var. *zeylanica* L.

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout plains in India; native to tropical Africa.

Local/Common/Vernacular names: *Mother-in-law tongue, snake plant* (English)

Yucca L.

Yucca aloifolia L., Sp. Pl. 319.1753

Yucca draconis L., Sp. Pl. 319.1753

Sarcococcus aloifolia (L.) Linding., Beih. Bot. Centralbl. 50(1): 446.1933

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Mysuru), cultivated in gardens; native to Bermuda, Mexico, USA.

Local/Common/Vernacular names: *Dagger plant, Spanish bayonet* (English)

Yucca gloriosa L., Sp. Pl. 319.1753

Yucca ensifolia Groenl., Rev. Hort. IV, 7: 433.1858

Yucca flexilis var. *ensifolia* (Groenl.) Baker, J. Linn. Soc., Bot. 18: 224.1880

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Kalburgi, Mysuru, Yadgir), cultivated in gardens; native to USA.

Local/Common/Vernacular name: *Spanish dagger* (English)

Asteraceae (Compositae)
Acanthospermum Schrank

Acanthospermum australe (Loeff.) Kuntze, Revis. Gen. Pl. 1: 303. 1891; Haleshi, Srungeswara, S.Vishwanath, Rajanna & B.Gowda, Rheedea 22: 33.2012.

Melampodium australe Loefl., Iter Hispan.: 268. 1758

Distribution: Karnataka (Bengaluru), weed in plains; native to S America.

Acanthospermum consobrinum S.F. Blake, Cont. U.S. Natl. Herb. 20: 383.1921; K. Ravikumar, N.M. Ganesh Babu, N. Balachandran & Vijayasankar, Rheedea 24: 50. 2014.

Distribution: Karnataka (Bengaluru), weed in plains; native to S America.

Acanthospermum hispidum DC., Prodr. 5: 522. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 134. 1984.

Acanthospermum humile var. *hispidum* (DC.) Kuntze, Revis. Gen. Pl. 1:303.1891

Acanthospermum australe sensu Raizada & V.S. Sharma in Indian Forester 88: 364. 1962. non (Loefl.) Kuntze, 1891.

Distribution: Karnataka (throughout), weed throughout India; native to S America.

Local/Common/Vernacular name: *Kadale mullu gida* (Kannada)

Acilepis H.Rob.

Acilepis aspera (Buch.-Ham.) H.Rob., Proc. Biol. Soc. Washington 112(1): 226. 1999

Vernonia aspera Buch.-Ham., Trans. Linn. Soc. London 14: 219. 1824; R.R. Rao et al., Fl. Ind. Enum.-Asterac. 86. 1988.

Eupatorium pyramidale D.Don, Prodr. Fl. Nepal. 170. 1825

Vernonia roxburghii Less., Linnaea 6: 674. 1831; Hook. f., Fl. Brit. India 3: 232. 1881.

Xipholepis aspera (Buch.-Ham.) Steetz in Peters, Reise Mossamb. Bot. 344.1863

Vernonia pyramidale (D.Don) S.N.Mitra, Indian Forester 99: 100. 1973

Distribution: Karnataka (throughout), throughout India.

Acilepis dendigulensis (DC.) H.Rob., Proc. Biol. Soc. Washington 112(1): 226.1999

Decaneurum dendigulense DC., Contr. Bot. India 7.1834

Gymnanthemum dendigulense (DC.) Walp., Repert. Bot. Syst. 2: 948. 1843

Lysistema dendigulense (DC.) Steetz in Peters, Reise Mossamb. Bot. 341.1863

Vernonia indica Wall. ex C.B. Clarke, Compos. Ind. 16. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Distribution: Karnataka (Belagavi, Ballari, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Arapu kanne* (Kannada)

Acilepis divergens (Roxb.) H.Rob. & Skvarla, Proc. Biol. Soc. Wash. 122(2): 140. 2009 140.2009

Eupatorium divergens Roxb., Fl. Ind. ed.2, 3: 414. 1832

Decaneurum divergens (Roxb.) DC., Prodr. 5: 68. 1834

Vernonia nilgheryensis DC., Prodr. 5: 32. 1836

Gymnanthemum divergens (Roxb.) Sch.Bip., Repert. Bot. Syst. 2: 948. 1843

Vernonia divergens (Roxb.) Edgew., J. Asiat. Soc. Bengal Part 2, Nat. Hist. 21: 172. 1853; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Cacalia divergens (Roxb.) Kuntze, Revis. Gen. Pl. 2:970.1891

Distribution: Karnataka (Belagavi, Ballari, Chamarajanagar, Chikkamagaluru, Chitradurga, Davanagere Hassan, Mysuru, Kodagu, Shivamogga, Uttara Kannada), throughout India;Myanmar.

Local/Common/Vernacular name: *Arapu kanne* (Kannada)

Acilepis gardneri (Thwaites) H.Rob. & Skvarla, Proc. Biol. Soc. Wash. 122(2): 141. 2009

Vernonia gardneri Thwaites, Enum. Pl. Zeyl. 161. 1860; Hook. f., Fl. Brit. India 3: 230. 1881; H.B.Naithani, Fl. Pl. India, Nepal & Bhutan 250. 1990.

Distribution: Karnataka(), S India ?

Acilepis ornata (Talbot) H.Rob. & Skvarla, Proc. Biol. Soc. Wash. 122(2): 141. 2009

Vernonia dalzelliana J.R. Drumm. & Hutch., Bull. Misc. Inform. 1909: 261. 1909; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Vernonia ornata Talbot, J. Bombay Nat. Hist. Soc. 11: 691. 1898 & For. Fl. Bombay 2: 137. 1911; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Acilepis dalzelliana (J.R.Drumm. & Hutch.) H.Rob., Proc. Biol. Soc. Washington 112(1): 226. 1999

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), N & S India; endemic.

Acilepis saligna (DC.) H.Rob., Proc. Biol. Soc. Washington 112(1): 226. 1999

Vernonia saligna DC., Prod 5: 33. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Conyza laeta C.B.Clarke, Compos. Ind. 14. 1876

Vernonia saligna var. *nilghirensis* Hook.f., Fl. Brit. India 3: 235. 1881

Cacalia saligna (DC.) Kuntze, Revis. Gen. Pl. 1:324.1891

Distribution: Karnataka (Uttara Kannada)

Acmella Rich. ex Pers.

Acmella calva (DC.) R.K.Jansen, Syst. Bot. Monogr. 8: 41. 1985.

Spilanthes acmella var. *calva* (DC.) C.B.Clarke ex C.B.Clarke, Comp. Ind. 138.1876

Distribution: Karnataka (Chitradurga, Hassan), throughout India; S & SE Asia.

Acmella ciliata (Kunth) Cass. in F.Cuvier, Dict. Sci. Nat., ed. 2. 24: 331. 1822

Spilanthes ciliata Kunth in Humb., Bonpl. & Kunth, Nov. Gen. Sp. 4: 208. 1820; Sivar. & Remesan, J. Econ. Taxon. Bot. 10: 145. 1987.

Distribution: Karnataka (throughout), throughout India except arid areas.

Acmella oleracea (L.) R.K.Jansen, Syst. Bot. Monogr. 8 65. 1985

Spilanthes oleracea L., Syst. Nat. ed. 12. 2: 534. 1767; R.R. Rao et al., Fl. Ind. Enum.-Asterac. 73. 1988.

Spilanthes acmella var. *oleracea* (L.) C.B.Clarke in Hook. f., Fl. Brit. Ind. 3: 307. 1881

Distribution: Karnataka (throughout), throughout tropical India; native to tropical America.

Local/Common/Vernacular name: *Toothache Plant* (English)

Acmella paniculata (Wall. ex DC.) R.K.Jansen, Syst. Bot. Monogr. 8: 67. 1985

Spilanthes calva DC. in Wight, Contrib. 19. 1834

Spilanthes paniculata Wall. ex DC., Prodr. 5: 625. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Spilanthes acmella var. *paniculata* (Wall. ex DC.) C.B.Clarke, Compos. Ind. 139. 1876

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; native to S America.

Local/Common/Vernacular name: *Sanna vana muguli* (Kannada)

Acmella radicans (Jacq.) R.K.Jansen, Syst. Bot. Monogr. 8: 69. 1985

Spilanthes radicans Jacq., Collectanea 3: 229. 1791; Sivar. & Remesan, J. Econ. Taxon. Bot. 10: 144. 1987.

Distribution: Karnataka (Chitradurga, Dharwad), N & S India; native to C America.

Acmella uliginosa (Sw.) Cass. in F.Cuvier's Dict. sci. nat. ed. 2. 24: 331. 1822

Spilanthes uliginosa Sw., Prodr. 110. 1788

Spilanthes iabadicensis A.H.Moore in Proc. Amer. Acad. Arts 42 (20): 524. 1907

Distribution: Karnataka (Dakshina Kannada, Udupi), throughout India except arid areas; native to S America.

Local/Common/Vernacular name: *Marsh Para Cress* (English)

Adenoon Dalzell

Adenoon indicum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 344. 1850; Hook.f., Fl. Brit. India 3: 229. 1881. Plate 18C

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Adenostemma J.R. & G. Forst.

Adenostemma lavenia (L.) Kuntze, Revis. Gen. Pl. 1: 304. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 134. 1984. Plate 4D

Verbesina lavenia L., Sp. Pl. 902. 1753

Adenostemma lavenia var. *viscosum* (J.R.Forst. & G.Forst.)M.R.Almeida, Fl. Maharashtra 3A:67.2001

var. **lavenia**

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; Paleotropical.

var. **angustifolium** (Arn.) J. Kost., Blumea 1: 475. 1935

Adenostemma angustifolium Arn., Nova Acta Phys.-Med. Acad. Caes. Leop.- Carol. Nat. Cur. 18: 347. 1836

Distribution: Karnataka (Shivamogga), throughout India; S & SE Asia.

var. **madurensse** (DC.) Panigrahi, Kew Bull. 30: 654. 1975; B.D.Sharma et al., Fl. Karnataka Anal. 134. 1984.

Adenostemma madurensse DC. in Wight, Contr. Bot. India 9. 1834

Adenostemma viscosum var. *reticulatum* (DC.) Hook.f., Fl. Brit. India 3: 243. 1881

Distribution: Karnataka (Hassan), S to NE India; endemic.

var. **reticulatum** (DC.) Panigrahi, Kew Bull. 30: 654. 1975

Adenostemma reticulatum DC. in Wight, Contr. Bot. India 8. 1834

Distribution: Karnataka (), Andaman & Nicobar Islands, S & E India.

var. **rugosum** (DC.) C.B.Clarke ex Kuntze, Revis. Gen. Pl. 1: 304. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 134. 1984.

Adenostemma rugosum DC. in Wight, Contr. Bot. India 8. 1834

Adenostemma viscosum var. *rugosum* (DC.) C.B.Clarke ex Hook.f., Fl. Brit. India 3: 243. 1881

Distribution: Karnataka (Kodagu, Uttara Kannada), Andaman & Nicobar Islands, Peninsular India.

Ageratina Spach

Ageratina adenophora (Spreng.) R.M. King & H. Rob., Phytologia 19: 211. 1970

Eupatorium glandulosum auct. non Michx., 1803; Kunth, Nov. Gen. Sp. 4: 96, t. 346. .

Eupatorium adenophorum Spreng., Syst. veg. ed. 16. 3: 420. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Kodagu, Kolar, Shivamogga), Weed throughout tropical India; native to Mexico & C America.

Local/Common/Vernacular names: *Crofton weed, Mexican devil, Sticky snakeroot* (English); *Indira Gandhi (Kunabi)*

Ageratina riparia (Regel) R.M. King & H. Rob., Phytologia 19: 216. 1970

Eupatorium riparium Regel, Gartenflora 15: 324, t. 525. 1866

Distribution: Karnataka (cultivated & naturalised), cultivated & naturalised in India; native to Mexico, The Caribbean.

Local/Common/Vernacular names: *Creeping Croftonweed, Mistflower* (English)

Ageratum L.

Ageratum conyzoides L., Sp. Pl. 839. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Distribution: Karnataka (throughout), throughout India; Pantropical.; native to tropical America.

Local/Common/Vernacular names: *Billygoat weed, Chik weed, Goat weed, White weed* (English); *Kempunola, Muguthi gida, Naayi tulasi, Shwetha shadevi* (Kannada)

Ageratum houstonianum Mill., Gard. Dict. ed. 8. no. 2. 1768

Carelia houstoniana (Mill.) Kuntze, Revis. Gen. Pl. 1: 325. 1891

Distribution: Karnataka (throughout), throughout India; native to C America.

Local/Common/Vernacular names: *Blue mink, Blue weed, Floss flower, Mexican paint brush, Pussy foot* (English)

Ambrosia L.

Ambrosia artemisifolia L., Sp. Pl. 988. 1753; Chatterji & Raizada, Indian Forester 74: 389. 1948.

Distribution: Karnataka (Bengaluru, Mysuru, Tumakuru), N India; native to N America.

Local/Common/Vernacular names: *American wormwood, Annual ragweed, Bitterweed, Blackweed, Carrot weed, Hayfever weed, Roman wormwood, stammerwort, stickweed, tassel weed* (English)

Ambrosia psilostachya DC., Prodr. 5: 526. 1836; Prasad et al., Curr. Sci. 104 (3): 294. 2013.

Distribution: Karnataka (Tumukuru); Native of temperate North America.

Local/Common/Vernacular names: *Perennial ragweed, Western ragweed* (English); *Sudugadu davana, Ketta davana* (Kannada).

Anaphalis DC.

Anaphalis lawii (Hook.f.) Gamble, Fl. Madras 695. 1921; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Anaphalis oblonga var. *lawii* Hook. f., Fl. Brit. India 3: 283. 1881

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Hassan, Kalaburgi, Kodagu, Mysuru, Yadgir), S to E India; endemic.

Anaphalis marcescens (Wight) C.B.Clarke in Hook.f., Fl. Brit. India 3: 286. 1881

Gnaphalium marcescens Wight, Icon. Pl. Ind. Orient. t. 1115. 1846

Distribution: Karnataka (Dakshina Kannada, Kodagu, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Anaphalis subdecurrens (DC.) Gamble, Fl. Madras 695. 1921; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Gnaphalium subdecurrens DC. in Wight, Contr. Bot. Ind. 21. 1834

Anaphalis oblonga DC., Prodr. 6: 274. 1838

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chikkamagaluru, Kodagu, Mysuru, Shivamogga), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Anaphalis wightiana (DC.) DC., Prodr. 6: 273. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Gnaphalium wightianum Wall ex DC. in Wight, Contr. Bot. India 21. 1834

Distribution: Karnataka (Chikkamagaluru, Coorg), Kerala, Tamil Nadu; endemic.

Argyranthemum Webb ex Sch.Bip.

Argyranthemum frutescens (L.) Sch.Bip. in Webb & Berth., Hist. Nat. Iles Canaries 2, 2:264 1844

Chrysanthemum frutescens L., Sp. Pl. 887. 1753

Distribution: Karnataka (cultivated), also elsewhere in India; native to Canary Islands.

Local/Common/Vernacular names: *Marguerite daisy, The Marguerite* (English)

Artemisia L.

Artemisia abrotanum L., Sp. Pl. 845. 1753; J. Graham, Cat. Pl. Bombay 102. 1839.

Distribution: Karnataka (cultivated in Bengaluru); native to Europe.

Local/Common/Vernacular names: *Lad's love, Southern wormwood, Southernwood* (English)

Artemisia japonica Thunb., Fl. Jap. 310. 1784

Artemisia parviflora Buch.-Ham. ex Roxb., Hort. Bengal. 61. 1814.

Artemisia japonica var. *parviflora* (Roxb. ex D. Don) Pamp., Nuovo Giorn. Bot. Ital. n.s. 34: 665. 1927

Artemisia japonica var. *purandharensis* M.R. Almeida, Fl. Maharashtra 3A: 70. 2001

Distribution: Karnataka (Belagavi); hilly areas throughout India; E & S Asia.

Local/Common/Vernacular name: *Japanese wormwood* (English)

Artemisia lactiflora Wall. ex DC., Prodr. 6: 115. 1838; Y.R. Ling, Bull. Bot. Surv. India 33: 303. 1994.

Distribution: Karnataka (cultivated in gardens), native to China.

Local/Common/Vernacular name: *The white mugwort* (English)

Artemisia nilagirica (C.B.Clarke) Pamp., Nuovo Giorn. Bot. Ital. n.s. 33: 452. 1926; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Artemisia vulgaris auct. non L., 1753; Hook. f., Fl. Brit. India 3: 325. 1881

Artemisia indica auct. non Willd., 1804; Wight, Icon. Pl. Ind. Orient. t. 1112. 1846

Artemisia vulgaris var. *nilagirica* C.B.Clarke, Compos. Ind. 162. 1876

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Chitradurga, Hassan, Kodagu, Mysuru, Tumakuru), throughout hilly regions of India; endemic.

Local/Common/Vernacular names: *Fleabane, Indian Wormwood, Mugwort* (English); *Mancha patre, Manji patre, Urigattige* (Kannada)

Artemisia pallens Wall. ex Besser, Nouv. Mém. Soc. Imp. Naturalistes Moscou 3: 85. 1834

Artemisia paniculata Roxb., Fl. Ind. ed.2, 3:418. 1832

Artemisia orientalis DC., Prodr., 6: 120. 1838

Artemisia absinthii Hook.f., Fl. Brit. India 3: 329. 1881

Distribution: Karnataka (cultivated in Bengaluru, Kolar, Mysuru, Tumakuru), cultivated in Andhra Pradesh, Maharashtra, Tamil Nadu; native to W Himalaya.

Local/Common/Vernacular name: *Dhavana* (Kannada); *Damanaka* (Sanskrit)

Artemisia parviflora Roxb. ex D. Don, Prodr. Fl. Nepal. 181. 1825

Distribution: Karnataka (Chamarajanagar, Shivamogga), Afghanistan, Bhutan, China, Myanmar, Nepal, Sri Lanka.

Artemisia princeps Pamp., Nuovo Giorn. Bot. Ital. n.s. 36: 444. 1930; S.M. Almeida, J. Econ. Taxon. Bot. addl. ser. 8(1): 236. 1990.

Distribution: Karnataka (cultivated in Gardens).

Local/Common/Vernacular names: *Japanese mugwort, Yellow-flower mugwort* (English)

Aster L.

Aster amellus L., Sp. Pl. 873. 1753; Hook. f., Fl. Brit. India 3: 249. 1881.

Distribution: Karnataka (cultivated), cultivated in many parts of India; native to Europe & C Asia.

Local/Common/Vernacular names: *Aster, The European Michaelmas Daisy* (English)

Bartlettina R.M.King & H.Rob.

Bartlettina sordida (Less.) R.M.King & H.Rob., Phytologia 22: 161. 1971

Eupatorium sordidum Less., Linnaea 6:403.1831

Conoclinium ianthinum C.Morren, Ann. Soc. Roy. Agric. Gand 5: 173. 1849

Eupatorium ianthinum (Morren) Hemsl., Biol. Cent.-amer., Bot. 2(7): 96. 1881; V. Chandras. in A.N. Henry et al., Fl. Tamil Nadu Anal. 2: 39. 1987.

Hebeclinium sordidum (Less.) Sch.Bip. ex J.Kost., Blumea 1: 494. 1935

Neobartlettia sordida (Less.) R.M.King & H.Rob., Phytologia 21: 296. 1971

Distribution: Karnataka (cultivated), cultivated in India; native to Mexico

Local/Common/Vernacular names: *Blue mist flower, Purple torch* (English)

Bellis L.

Bellis perennis L., Sp. Pl. 886. 1753

Erigeron perennis (L.) Sessé & Moc., Fl. Mexic., ed. 2, 184. 1894

Distribution: Karnataka (cultivated ornamental in Bengaluru), cultivated ornamental ;native to Europe.

Local/Common/Vernacular names: *Common daisy, English daisy, Lawn daisy* (English)

Bidens L.

Bidens bipinnata L., Sp. Pl. 832. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Bidens decomposita Wall. ex DC., Prodr. 5: 602. 1836

Bidens wallichii DC., Prodr. 5 598. 1836

Kerneria bipinnata (L.) Gren. & Godr., Fl. France 2: 169. 1850

Bidens pilosa var. *wallichii* (DC.) C.B.Clarke, Compos. Ind. 141. 1876

Bidens pilosa var. *bipinnata* (L.) Hook. f., Fl. Brit. India 3: 309. 1881

Bidens pilosa var. *decomposita* (Wall.) Hook. f., Fl. Brit. India 3: 310. 1881

Distribution: Karnataka (Belagavi, Bidar, Chikkamagaluru, Kolar, Mysuru, Tumakuru, Uttara Kannada), throughout India including Himalayas.

Local/Common/Vernacular name: *Para cress flower* (English); *Bende* (Kannada)

Bidens biternata (Lour.) Merr. & Sherff, Bot. Gaz. 88: 293. 1929; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Bidens pilosa non L., 1753; Hook. f., Fl. Brit. India 3: 309. 1881 p.p.

Coreopsis biternata Lour., Fl. Cochinch. 508. 1790

Actinea biternata (Lour.) Spreng., Syst. Veg. 3:574. 1826

Distribution: Karnataka (throughout), throughout India; Pantropical.

Bidens tripartita L., Sp. Pl.: 831. 1753

Bidens connata var. *comosa* A.Gray, Manual (Gray), ed. 5. 261. 1867

Bidens comosa (A.Gray) Wiegand, Bull. Torrey Bot. Club 24: 436. 1897; Babu, N.M.G., Fl. Ethnob. Chitradurga Dist. 299. 2011.

Distribution: Karnataka (Chitradurga), native to N America.

Blainvillea Cass.

Blainvillea acmella (L.) Philipson, Blumea 6: 350. 1950; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Plate 4E

Verbesina acmella L., Sp. Pl. 901. 1753

Spilanthes acmella auct. non Murray, 1774; Hook. f., Fl. Brit. India 3: 307. 1881

Pyrethrum acmella (L.) Medik., Act. Acad. Theod. Palat. 3: 243. t. 19. 1775

Eclipta latifolia L.f., Suppl. Pl. 378. 1782

Bidens acmella (L.) Lam., Encyc. 1(2): 415. 1785

Blainvillea rhomboidea Cass. in F. Cuvier's Dict. Sci. Nat. ed. 2. 29: 494. 1823

Blainvillea latifolia (L.f.) DC. in Wight, Contr. Bot. India 17. 1834

Coreopsis acmella (L.) E.H.L.Krause, Beih. Bot. Centralbl. 32(2): 340. 1914

Distribution: Karnataka (throughout), throughout India; Pantropical.

Blepharispermum DC.

Blepharispermum subsessile DC. in Wight, Contr. Bot. India 12. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Distribution: Karnataka (Belagavi, Ballari, Chamarajanagar, Chikkamagaluru, Davanagere, Dharwad, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Peninsular India.

Blumea DC.

Blumea axillaris (Lam.) DC., Prodr. 5: 434. 1836; Grierson & Spring. in Grierson & D.G. Long, Fl. Bhutan 2 (3): 1504. 2001.

Conyza axillaris Lam., Encycl. 2: 84. 1786

Gnaphalium solidaginoides Poir. in Lam., Encycl. Suppl. 2: 805. 1812

Erigeron molle D.Don, Prodr. Fl. Nepal. 172. 1825

Blumea wightiana DC. in Wight, Contr. Bot. India 14. 1834

Blumea solidaginoides (Poir.) DC., Prodr. 5: 443. 1836; M.R. Almeida, Fl. Maharashtra 3A: 85. 2001.

Blumea neilgherrensis Hook. f., Fl. Brit. India 3: 261. 1881

Blumea mollis (D.Don) Merr., Philipp. J. Sci. 5: 395. 1910

Distribution: Karnataka (Belagavi, Chamarajanagar, Chitradurga, Dakshina Kannada, Davanagere, Kalaburgi, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout India ; Paleotropical.

Local/Common/Vernacular name: *Gabbu soppu* (Kannada)

Blumea balsamifera (L.) DC., Prodr. 5:447.1836; Hook. f., Fl. Brit. India 3: 270. 1881.

Conyza balsamifera L., Sp. Pl. ed.2, 2:1208.1763

Pluchea balsamifera (L.) Less., Linnaea 6:150.1831

Distribution: Karnataka (), throughout India; S & SE Asia

Blumea barbata DC., Contr. Bot. India 14. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Blumea belangeriana DC., Prodr. 5:444.1836; B.D.Sharma et al., Fl. Karnataka Anal. 135. 1984.

Blumea metziana Sch.Bip. ex Hook.f., Fl. Brit. India 3: 266. 1881

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Peninsular to NE India; Sri Lanka.

Blumea bifoliata (L.) DC. in Wight, Contr. Bot. India 14. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

Conyza bifoliata L., Sp. Pl. ed. 2. 1207. 1763

Distribution: Karnataka (Belagavi, Bengaluru, Uttara Kannada), throughout India; S & SE Asia.

Blumea clarkei Hook.f., Fl. Brit. India 3 267.1881

Blumea malabarica Hook.f., Fl. Brit. India 3 267.1881

Distribution: Karnataka (Kodagu?, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; China, SE Asia.

Blumea eriantha DC., Contr. Bot. India 15. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Raichur, Shivamogga, Uttara Kannada), throughout except NE India; endemic.

Blumea fistulosa (Roxb.) Kurz, J. Asiat. Soc. Bengal Pt. 2, Nat. Hist. 46: 187. 1877; Randeria, Blumea 10: 256. 1960.

Conyzafistulosa Roxb., Fl. Ind. ed.2, 3: 422. 1832

Blumea glomerata DC., Contr. Bot. India 15. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

Blumeafistulosa var. *glomerata* (DC.) C.B.Clarke, Comp. Ind. 82. 1876

Blumea nodiflora Hook. f., Fl. Brit. India 3: 262. 1881

Distribution: Karnataka (Uttara Kannada), throughout India including tropical Himalayas.

Blumea hamiltonii DC., Prodr. 5: 439. 1836

Gnaphalium hamiltonii Wall., Numer. List n. 2938.1831

Blumea hieracifolia var. *hamiltonii* (DC.) C.B.Clarke, Compos. Ind. 82. 1876

Distribution: Karnataka (throughout), Almost throughout India.

Blumea hieracifolia (D.Don) DC., Contr. Bot. India 15, 1834; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

Conyzahirsuta L., Sp. Pl. 863.1753

Erigeron hieracifolium D.Don, Prodr. Fl. Nepal. 272. 1825

Conyzahieraciifolia Spreng., Syst. Veg. 3:514.1826

Blumeaflexuosa C.B.Clarke, Compos. Ind. 86. 1876

Blumeaflexuosa var. *peninsularis* Hook. f., Fl. Brit. India 3: 267. 1881

Blumeaflexuosa var. *zeylanica* Hook. f., Fl. Brit. India 3: 267. 1881

Blumea hieracifolia var. *flexuosa* (C.B.Clarke) Randeria, Blumea 10: 249. 1960; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

var. **hieracifolia**

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Hassan), throughout India; Australia, S & SE Asia.

var. **evolutior** C.B.Clarke, Comp. India 83.1876

Blumea macrostachya DC., Prodr. 5: 442. 1836; Burkill in Rec. Bot. Surv. India 10 (2): 306. 1925.

Blumea hieracifolia var. *macrostachya* (DC.) Hook.f., Fl. Brit. India 3: 263. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

Distribution: Karnataka (Hassan), throughout India; S & SE Asia.

Blumea hirsuta (Less.) M.R.Almeida, Fl. Maharashtra 3A:79.2001

Plucheahirsuta (L.) Less., Linnaea 6:150.1831

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Mysuru, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Peninsular & NE India.

Blumea lacera (Burm.f.) DC., Contr. Bot. India 14. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 136. 1984.

Conyza lacera Burm. f., Fl. indica 180, t. 59, f.1. 1768

Blumea glandulosa DC., Contr. Bot. India 14. 1834

Blumea heyneana DC., Contr. Bot. India 15. 1834

Baccharis auriculata Wall. ex DC., Prodr. 5: 442. 1836

Blumea cinerascens DC., Prodr. 5: 438. 1836

Blumea subcapitata DC., Prodr. 5: 439. 1836

Conyza necessaria Buch.-Ham. ex DC., Prodr. 5: 439. 1836

Blumea commersonii Edgew., J. Asiat. Soc. Bengal Part 2, Nat. Hist. 21(1): 38. 1852

Blumea fontinalis Edgew., J. Asiat. Soc. Bengal Part 2, Nat. Hist. 21(1): 173. 1852

Conyza lacera var. *wallichii* (DC.) DC. ex Miq., Fl. Ned. Ind. 2:42 1856

Conyza lacera var. *wightiana* (DC.) DC. ex Miq., Fl. Ned. Ind. 2:43 1856

Blumea lacera var. *hymenophylla* (DC.) C.B.Clarke, Compos. Ind. 77.1876

Blumea hieraciifolia Hook.f. & Thomson, Fl. Brit. India 3: 267.1881

Blumea lacera var. *cinerascens* (DC.) Hook.f., Fl. Brit. India 3: 263.1881

Blumea lacera var. *glandulosa* (DC.) Hook.f., Fl. Brit. India 3: 263.1881

Blumea villosa Sch.Bip. ex Hook. f., Fl. Brit. India 3: 263.1881

Conyza natans Buch.-Ham. ex Hook. f., Fl. Brit. India 3: 268.1881

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Raichur, Shivamogga, Udupi, Uttara Kannada), throughout plains of India; Paleotropical.

Blumea laciniata (Wall. ex Roxb.) DC., Prodr. 5: 436.1836; K. G. Bhat, Fl. S. Kanara 824. 2014.

Blumea acutata DC., Nouv. Ann. Mus. Paris 3: 409. 1834

Blumea glandulosa Benth., Fl. Hongk. 177. 1861.

Distribution: Karnataka (Dakshina Kannada, Udupi).

Blumea laevis (Lour.) Merr., Trans. Amer. Philos. Soc. n.s. 24: 386. 1935

Placus laevis Lour., Fl. Cochinch. 497. 1790

Distribution: Karnataka (Belagavi, Uttara Kannada), throughout India.

Blumea lanceolaria (Roxb.) Druce in Bot. Soc. Exch. Club Brit. Isles 4:609. 1917; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Conyza lanceolaria Roxb., Fl. Ind. ed.2, 3: 432. 1832

Blumea myriocephala DC., Prodr. 5: 445. 1836; Hook. f., Fl. Brit. India 3: 269. 1881.

Blumea sessilifolia auct. non DC., 1836; Kurz, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 46(2): 189. 1877

Blumea spectabilis DC., Prodr. 5: 445. 1836

Blumea squarrosa C.B.Clarke, Compos. Ind. 87.1876

Blumea wallichii C.B.Clarke, Compos. Ind. 87.1876

Blumea sessifolia var. *lanceolaria* (Roxb.) Kurz, J. Asiat. Soc. Bengal Part2, Nat. Hist. 46(2):189 1877

Blumea spectabilis var. *longifolia* Hook. f., Fl. Brit. India 3: 269. 1881

Blumea lanceolaria var. *spectabilis* (DC.) Randeria, Blumea 10: 220. 1960

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada),throughout India including Andaman & Nicobar Islands; China, S & SE Asia, Taiwan.

Blumea malcolmii (C.B.Clarke) Hook.f., Fl. Brit. India 3: 266. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Pluchea malcolmii C.B.Clarke, Compos. Ind. 95. 1876

Distribution: Karnataka (Belagavi, Dakshina Kannada), Kerala, Maharashtra, Nagar Haveli, Tamil Nadu; endemic.

Blumea membranacea DC., Prodr. 5:440.1836; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Blumea paucifolia DC., Prodr. 5: 446. 1836

Distribution: Karnataka (Belagavi, Chamarajanagar, Dakshina Kannada, Kalaburgi, Kodagu, Mysuru, Udupi, Uttara Kannada, Yadgir), Goa, Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Blumea obliqua (L.) Druce, Bot. Soc. Exch. Club Brit. Isles 4: 609. 1917; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Erigeron obliquum L., Mant. Pl. 2: 573. 1771

Blumea amplexens DC., Contr. Bot. India 13. 1834

Blumea arenaria DC., Contr. Bot. India 13. 1834

Blumea pubiflora DC., Prodr. 5: 434. 1836

Blumea amplexens var. *arenaria* (DC.) Hook.f., Fl. Brit. India 3: 260. 1881

Blumea amplexens var. *maritima* C.B.Clarke ex Hook. f., Fl. Brit. India 3: 260. 1881

Blumea amplexens var. *pubiflora* (DC.) Hook.f., Fl. Brit. India 3: 260. 1881

Blumea amplexens var. *tenella* Hook. f., Fl. Brit. India 3: 260. 1881

Blumea obliqua var. *arenaria* (DC.) Maheshw., J. Bombay Nat. Hist. Soc. 54: 805. 1957

Blumea obliqua var. *maritima* (C.B.Clarke) Maheshw., J. Bombay Nat. Hist. Soc. 54: 805. 1957

Blumea obliqua var. *pubiflora* (DC.) Maheshw., J. Bombay Nat. Hist. Soc. 54: 805. 1957

Blumea obliqua var. *tenella* (Hook.f.) Maheshw., J. Bombay Nat. Hist. Soc. 54: 805. 1957

Blumea obliqua var. *arenaria* (DC.) G.S.Giri & B.Mitra, Indian J. Forest. 7(2): 143. 1984

Blumea obliqua var. *maritima* (Hook.f.) G.S.Giri & B.Mitra, Indian J. Forest. 7(2): 143. 1984

Blumea obliqua var. *aurita* (L.f.) Naik & Bhog., Acta Bot. Indica 20: 49. 1992

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kalaburgi, Raichur, Uttara Kannada, Vijayapura), throughout India including Andaman & Nicobar Islands; S Asia.

Blumea oxyodonta DC., Contr. Bot. India 15. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; China, S & SE Asia.

Blumea procera DC., Prodr. 5: 445. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Distribution: Karnataka (Udupi? Uttara Kannada), Andaman & Nicobar Islands, NE India, tropical Himalayas, Upper Gangetic plains.

Blumea sessiliflora Decne., Nouv. Ann. Mus. Hist. Nat. 3: 140. 1834; N.P. Balakr., Fl. Jowai 1: 275. 1981.

Blumea fasciculata DC., Prodr. 5, 442. 1836

Distribution: Karnataka (throughout), Andaman & Nicobar Islands, Peninsular to NE India.

Blumea sinuata (Lour.) Merr., Trans. Amer. Philos. Soc. n.s. 24 : 388. 1935; K.K. Sharma & S. Sharma, Fl. Rajasthan 104. 1989.

Gnaphalium sinuatum Lour., Fl. cochinch. 2: 497. 1790

Conyzia laciniata Wall. ex Roxb., Fl. Ind. 3:427.1832

Blumea lacera var. *blumei* DC., Prodr. 5: 436. 1836

Distribution: Karnataka (throughout), throughout India including Andaman & Nicobar Islands.

Blumea virens DC. in Wight, Contr. Bot. India 14. 1834

Erigeron virens Russ. ex DC., Prodr. 5: 439. 1836

Blumea diffusa Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 46(2): 187. 1877

Blumea virens var. *minor* Hook. f., Fl. Brit. India 3: 264. 1881

Blumea laevis var. *minor* (Hook.f.) Karthik. & Moorthy, Fl. Pl. India 205. 2009

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi), Andaman & Nicobar Islands, Peninsular to NE India; S & SE Asia.

Blumeopsis Gagnep.

Blumeopsis flava (DC.) Gagnep., Bull. Mus. Hist. Nat. (Paris) 26: 76. 1920; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Blumea flava DC., Prodr. 5: 439. 1836

Laggera flava (DC.) Benth. in Benth. & Hook. f., Gen. Pl. 2: 290. 1873; Hook. f., Fl. Brit. India 3: 270. 1881.

Distribution: Karnataka (Hassan, Kodagu), throughout India; S Asia.

Caesulia Roxb.

Caesulia axillaris Roxb., Pl. Coromandel 1: 64, t. 93. 1798; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.

Distribution: Karnataka (Belagavi, Bidar, Dharwad, Kalaburgi, Uttara Kannada, Vijayapura, Yadgir), throughout India.

Calendula L.

Calendula officinalis L., Sp. Pl. 921. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Distribution: Karnataka (cultivated throughout, sometimes found as an escape), cultivated throughout India; native to Europe.

Local/Common/Vernacular names: *Common marigold, English marigold, Garden marigold, Pot marigold, Scottish marigold* (English)

Callistephus Cass., nom. cons.

Callistephus chinensis (L.) Nees, Gen. Sp. Aster. 222. 1832

Aster chinensis L., Sp. Pl. 872. 1753

Callistephus hortensis Cass. in F. Cuvier's Dict. Sci. Nat. ed. 2. 37: 491. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Distribution: Karnataka (cultivated throughout, sometimes as an escape), cultivated throughout India; native to E Asia.

Local/Common/Vernacular names: *Annual aster, China aster* (English)

Calyptocarpus Less.

Calyptocarpus vialis Less., Syn. Gen. Compos. 221. 1832

Synedrella vialis (Less.) A.Gray, Proc. Amer. Acad. Arts 17: 217. 1882

Distribution: Karnataka (weed throughout), cultivated throughout India; native to E Asia.

Local/Common/Vernacular names: *Creeping Cinderella-weed, Horseherb, Lawnflower, Straggler daisy* (English)

Carthamus L.

Carthamus lanatus L., Sp. Pl. 830. 1753

Centaurea lanata (L.) Lam. & DC., Fl. Franc. ed. 3. 4: 102. 1805

Kentrophyllum lanatum (L.) DC., Bot. Gallicum 1: 193. 1828

Distribution: Karnataka (Belagavi), cultivated throughout India; native to Mediterranean region.

Local/Common/Vernacular names: *Saffron thistle, Woolly distaff thistle* (English)

Carthamus tinctorius L., Sp. Pl. 830. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Distribution: Karnataka (cultivated throughout, sometimes as an escape), cultivated throughout India; Probably native to W Asia.

Local/Common/Vernacular name: *Safflower* (English); *Kusume* (Kannada)

Centaurea L.

Centaurea calcitrapa L., Sp. Pl. 917. 1753; Hook. f., Fl. Brit. India 3: 385. 1881.

Distribution: Karnataka (), NW Himalaya.

Local/Common/Vernacular names: *Caltrops*, *Centauree chausse-trappe*, *Chausse-trappe*, *Purple star-thistle* (English)

Centaurea cyanus L., Sp. Pl. 911. 1753; Hook. f., Fl. Brit. India 3: 384. 1881.

Distribution: Karnataka (Weed in cultivated fields & gardens).

Local/Common/Vernacular names: *Bachelor's button*, *Blue bottle*, *Cornflower* (English)

Centaurea melitensis L., Sp. Pl. 912. 1753; Hook. f., Fl. Brit. India 3: 385. 1881.

Distribution: Karnataka (Cultivated).

Local/Common/Vernacular names: *Centaury*, *Maltese star thistle*, *Napa thistle*, *Tocalote* (English)

Centipeda Lour.

Centipeda minima (L.) A. Braun & Asch., Index Sem. (Berlin) 1867: App. 6. 1868; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Artemisia minima L., Sp. Pl. 849. 1753

Centipeda orbicularis Lour., Fl. Cochinch. 493. 1790

Dichrocephala schmidii Wight, Icon. Pl. Ind. Orient. t. 1610. 1850

Distribution: Karnataka (Belagavi, Bengaluru, Davanagere, Hassan, Kodagu, Uttara Kannada), throughout India including Andaman & Nicobar Islands; Africa, Asia, Australia.

Local/Common/Vernacular name: *Chikoni* (Kannada)

Centratherum Cass.

Centratherum punctatum Cass., Bull. Sci. Soc. Philom. Paris 1817: 31. 1817; R.R. Rao et al., Fl. Ind. Enum. -Asterac. 23. 1988.

Distribution: Karnataka (cultivated as an ornamental, Bengaluru, Chitradurga, Dakshina Kannada, Mysuru, Udupi), native to tropical America.

Chaetoseris C.Shih

Chaetoseris cyanea (D.Don) C.Shih, Acta Phytotax. Sin. 29(5): 404. 1991

Sonchus cyaneus D.Don, Prodr. Fl. Nepal. 164. 1825

Lactuca hastata DC., Prodr. 7: 139. 1838; Hook. f., Fl. Brit. India 3: 407. 1881.

Cicerbita cyanea (D.Don) Beauverd in Bull. Soc. Bot. Genève 2: 132. 1910

Distribution: Karnataka (), Himalayas, E, N, NE & S India.

Chromolaena DC.

Chromolaena odorata (L.) R.M. King & H. Rob., Phytologia 20: 204. 1970; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Eupatorium odoratum L., Syst. Nat. ed. 10. 2: 1205. 1760

Distribution: Karnataka (throughout), throughout India; native to tropical America.

Local/Common/Vernacular names: *Communist grass*, *Communist weed* (English)

Chrysanthemum Rich.

Chrysanthemum carinatum Schousb., Iagttag. Vextrig. Marokko 198, t. 6. (1800) 1801; B.D.Naithani in Hajra et al., Fl. India 12: 74. 1995.

Distribution: Karnataka (cultivated), also elsewhere in India; probably native to N Africa.

Local/Common/Vernacular names: *Painted Daisy, Summer Chrysanthemum, Tricolor Chrysanthemum* (English)

Chrysanthemum coccineum Willd., Sp. Pl. 3: 2144. 1803; B.D.Naithani in Hajra et al., Fl. India 12: 74. 1995.

Distribution: Karnataka (cultivated), also elsewhere in India; native to Caucasus.

Local/Common/Vernacular names: *Painted Daisy, Persian chrysanthemum, Pyrethrum* (English)

Chrysanthemum foeniculaceum Steud., Nomencl. Bot. ed. 1. 1: 192. 1821; B.D.Naithani in Hajra et al., Fl. India 12: 75. 1995.

Distribution: Karnataka (cultivated), also elsewhere in India.

Chrysanthemum indicum L., Sp. Pl. 889. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Dendranthema indicum (L.) Des Moul., Actes Soc. Linn. Bordeaux 20: 562. 1855; Lauener, Notes Roy. Bot. Gard. Edinburgh 34: 352. 1975.

Distribution: Karnataka (cultivated), also elsewhere in India; native to China, Japan.

Local/Common/Vernacular name: *Indian chrysanthemum* (English)

Chrysanthemum morifolium Ramat., J. Hist. Nat. 2: 240. 1792; H. Hara, Fl. E. Himal. 335. 1966.

Matricaria morifolia (Ramat.) Ramat., J. Hist. Nat. 2: 240. 1792

Chrysanthemum sinense Sabine, Trans. Linn. Soc. London 14: 142. 1823; M.R. Almeida, Fl. Maharashtra 3A: 89. 2001.

Pyrethrum sinense (Sabine) DC., Prodr. 6: 62. 1838

Tanacetum sinense (Sabine) Sch. Bip., Tanacetum 50. 1844

Dendranthema sinensis (Sabine) Des Moul., Actes Soc. Linn. Bordeaux 20: 562. 1855

Distribution: Karnataka (cultivated), also elsewhere in India.

Distribution: Karnataka (cultivated), also elsewhere in India; native to China.

Local/Common/Vernacular names: *Florist's daisy, Hardy garden mum* (English)

Chrysanthemum myconis L., Sp. Pl. ed. 2. 1254. 1763

Distribution: Karnataka (cultivated), also elsewhere in India; native to Europe, Mediterranean region.

Local/Common/Vernacular name: *Corn marigold* (English)

Chrysanthemum parthenium (L.) Bernh., Syst. Verz. 145. 1800

Matricaria parthenium L., Sp. Pl. 890. 1753

Pyrethrum parthenium (L.) Sm., Flora Britannica 2. 1800

Distribution: Karnataka (cultivated), also elsewhere in India; native to Eurasia.

Cissampelopsis (DC.) Miq.

Cissampelopsis corymbosa (Wall. ex DC.) C.Jeffrey & Y.L.Chen, Kew Bull. 39(2): 341. 1984

Senecio corymbosus Wall. ex DC. in Wight, Contr. Bot. India 22. 1834

Distribution: Karnataka (Chamarajanagar, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Conyza Less., nom. cons.

Conyza bonariensis (L.) Cronquist, Bull. Torrey Bot. Club 70: 623. 1943; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Erigeron bonariensis L., Sp. Pl. 863. 1753

Erigeron crispus Pourr., Hist. & Mém. Acad. Roy. Sci. Toulouse 3: 318. 1788 p.p.

Erigeron linifolius Willd., Sp. Pl. 3: 1955. 1803

Coryza ambigua DC., Fl. franç. 6: 468. 1815

Conyza angustifolia Buch.-Ham. ex Roxb., Fl. Ind. ed. 2, 3: 429. 1832

Erigeron ambiguus auct. non Nutt. 1818. (DC.) Sch.Bip. in Webb & Berthel., Hist. nat. Iles Canaries sect. 3(2): 208. 1845

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; native to Europe.

Conyza japonica (Thunb.) Less. ex DC., Prodr. 5: 382. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Erigeron japonicum Thunb., Fl. Jap. 312. 1784

Eschenbachia japonica (Thunb.) J. Kost., Blumea 7: 290. 1952

Distribution: Karnataka (Hassan, Kalaburgi, Kodagu, Yadgir), throughout India; E, S & SE Asia.

Conyza leucantha (D.Don) Ludlow & P.H.Raven, Kew Bull. 17: 71. 1963; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Erigeron leucanthum D.Don, Prodr. Fl. Nepal. 171. 1825

Blumea leucanthema DC., Prodr. 5: 436. 1836; Hook. f., Fl. Brit. India 3: 270. 1881.

Conyza viscidula Wall. ex DC., Prodr. 5: 383. 1836; Hook. f., Fl. Brit. India 3: 258. 1881.

Blumeopsis falcata (D.Don) Merr., J. Arnold Arbor. 19: 70. 1938

Eschenbachia viscidula (Wall. ex DC.) J. Kost., Blumea 7: 290. 1952

Distribution: Karnataka (Davanagere, Hassan, Mysuru, Tumakuru, Uttara Kannada), throughout India; S & SE Asia.

Conyza stricta Willd., Sp. Pl. 3: 1922. 1803; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Conyza adenocarpa Dalzell & A.Gibson, Bombay Fl. 125. 1861

Eschenbachia stricta (Willd.) Raizada, Indian Forester 84: 467. 1959

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; Africa, S & SE Asia.

Conyza sumatrensis (Retz.) E. Walker, J. Jap. Bot. 46: 72. 1971; R.R. Rao et al., Fl. Ind. Enum. -Asterac. 28. 1988.

Erigeron sumatrensis Retz., Observ. Bot. 5: 28. 1787

Coreopsis L.

Coreopsis aristosa Michx., Fl. Bor.-Amer. 2: 140. 1803; M.R. Almeida, Fl. Maharashtra 3A: 92. 2001.

Bidens aristosa (Michx.) Britton, Bull. Torrey Bot. Club 20: 281. 1893

Distribution: Karnataka (cultivated in gardens)

Coreopsis tinctoria Nutt., J. Acad. Nat. Sci. Philadelphia 2:114.1821; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Diplosastera tinctoria (Nutt.) Tausch, Hort. Canal. t. 4.1823

Coreopsis atkinsoniana Douglas ex Lindl. in Edwards' Bot. Reg. 16: t. 1376. 1830; Deb, Fl. Tripura 2: 211. 1983.

Calliopsis atkinsoniana (Douglas) Hook., Fl. Bor.-Amer. 1: 310. 1833

Calliopsis tinctoria (Nutt.) DC., Prodr. 5:568.1836

Bidens tinctoria (Nutt.) Baill., Hist. Pl. 8: 305. 1882

Coreopsis tinctoria var. *atkinsoniana* (Douglas ex Lindl.) H.M.Parker ex E.B.Sm., Bot. Gaz. 136: 83. 1975

var. *tinctoria*

Distribution: Karnataka (cultivated in gardens), cultivated as an ornamental throughout; native to N America
 Local/Common/Vernacular name: *Plains coreopsis* (English)

var. *atropurpurea* Hook. in Curtis, Bot. Mag. 63: t. 3511. 1836; M. Sharma & H. Kaur, J. Econ. Taxon. Bot. 18: 726. 1994.

Distribution: Karnataka (cultivated in gardens), cultivated as an ornamental throughout; native to N America

Cosmos Cav.***Cosmos bipinnatus* Cav., Icon. 1: 10, t. 14. 1791; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.**

Distribution: Karnataka (cultivated throughout and also run wild), cultivated and also run wild throughout India; native to Mexico.

Local/Common/Vernacular names: *Garden cosmos, Mexican aster* (English)

***Cosmos caudatus* Kunth, Nov. Gen. Sp. 4(17): 188 (ed. fol.). 1818 & 4(17): 240 (ed. qto.). 1820**

Cosmea caudata (Kunth) Spreng., Syst. Veg. 3: 615. 1826

Bidens caudata (Kunth) Sch.Bip., Bot. Voy. Herald 7-8: 308. 1856; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Bidens artemisiifolia subsp. *caudata* (Kunth) Kuntze, Revis. Gen. Pl. 1: 321. 1891

Distribution: Karnataka (cultivated in gardens throughout, as an escape in Shivamogga), cultivated throughout India; native to C & S America.

Local/Common/Vernacular name: *Ulam raja* (English)

***Cosmos steenisiae* Veldkamp, Blumea 35: 479. 1991**

Cosmos calvus non Sherff, Publ. Field Columbian Mus., Bot. Ser. 8: 405. 1932; R.R. Rao et al., Fl. Ind. Enum. -Asterac. 28. 1988.

Distribution: Karnataka (Garden escape).

***Cosmos sulphureus* Cav., Icon. 1.1791**

Coreopsis artemisiaefolia Jacq., Icon. Pl. Rar. 3: 16, t. 595. 1786-1793

Cosmea sulphurea (Cav.) Willd., Sp. Pl. 3 2250. 1804

Bidens sulfurea (Cav.) Sch.Bip. in Seem., Narr. Voy. Herald 308. 1853

Bidens artemisiifolia (Jacq.) Kuntze, Revis. Gen. Pl. 1 321. 1891

Cosmos artemisiifolius (Jacq.) M.R.Almeida, Fl. Maharashtra 3A 93. 2001

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to C America.

Local/Common/Vernacular name: *Cosmos* (English) (Kannada)

Cotula L.***Cotula australis* (Sieber ex Spreng.) Hook. f., Fl. Nov.-Zel. 1: 128. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.**

Anacyclus australis Sieber ex Spreng., Syst. Veg. ed. 16, 3: 497. 1826

Distribution: Karnataka (Bengaluru, Chamarajanagar), native to Australia & New Zealand.

Local/Common/Vernacular names: *Australian waterbutton, Southern waterbuttons* (English)

***Cotula coronopifolia* L., Sp. Pl. 892. 1753; R.R. Rao et al., Fl. Ind. Enum. -Asterac. 29. 1988.**

Distribution: Karnataka (cultivated in gardens), cultivated as ornamental; native to S Africa.

Local/Common/Vernacular names: *Brass buttons, Button weed, Golden buttons* (English)

***Cotula turbinata* L., Sp. Pl. 892. 1753**

Cenia turbinata (L.) Pers., Syn. Pl. 2: 463. 1807; Hook. f., Fl. Brit. India 3: 316. 1881.

Distribution: Karnataka (cultivated in gardens) cultivated as ornamental; native to Australia & New Zealand.

Local/Common/Vernacular name: *Funnel Weed* (English)

Crassocephalum Moench

Crassocephalum crepidioides (Benth.) S. Moore, J. Bot. 50: 211. 1912; B.D.Sharma et al., Fl. Karnataka Anal. 139. 1984.

Erechtites valerianaefolia auct. non (Wolf) DC., 1838; C.E.C.Fisch. in Gamble, Fl. Madras 1882. 1936.

Gynura crepidioides Benth. in Hook., Niger Fl. 437. 1849

Distribution: Karnataka (Ballari, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi), naturalised almost throughout warmer parts of India; Pantropical.; native to tropical America.

Local/Common/Vernacular names: *Ebolo, Fireweed, Redflower ragleaf, Thickhead* (English)

Cyanthillium Blume

Cyanthillium cinereum (L.) H. Rob., Proc. Biol. Soc. Wash. 103: 252. 1990

Conyzia cinerea L., Sp. Pl. 862. 1753

Vernonia parviflora Reinw., Blume, Bijdr. Fl. Ned. Ind. 15: 893. 1826

Vernonia cinerea (L.) Less., Linnaea 4: 291. 1829; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Vernonia albicans DC., Contr. Bot. India 6. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Vernonia cinerea var. *parviflora* (Reinw.) DC., Prodr. 5: 24. 1836

Cyanthillium cinereum var. *parviflorum* (Reinw.) Karthik. & Moorthy, Fl. Pl. India 223. 2009

Distribution: Karnataka (throughout), throughout India; Paleotropical.

Local/Common/Vernacular names: *Kare hindi, Sahadevi* (Kannada)

Cyathocline Cass.

Cyathocline lutea Law ex Wight, Icon. Pl. Ind. Orient. t. 1150. 1846 & Calcutta J. Nat. Hist. 7: 158, t. 2, f. 1. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (Shivamogga, Uttara Kannada), peninsular India; endemic.

Cyathocline purpurea (D.Don) Kuntze, Revis. Gen. Pl. 1: 333. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Tanacetum purpureum D.Don, Prodr. Fl. Nepal. 181. 1825

Cyathocline lyrata Cass., Ann. Sci. Nat. (Paris) 17: 420. 1829

Dichrocephala minutifolia Vaniot, Bull. Acad. Int. Géogr. Bot. 12: 243. 1903

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Udupi, Uttara Kannada, Vijayapura), throughout India; SE Asia.

Cynara L.

Cynara cardunculus L., Sp. Pl. 827. 1753

Distribution: Karnataka (cultivated in gardens), cultivated as ornamental; native to Mediterranean region

Local/Common/Vernacular names: *Artichokethistle, Cardi, Cardone, Cardoni, Carduni* (English)

Cynara scolymus L., Sp. Pl. 827. 1753; H.J. Chowdhery in H.J. Chowdhery & Wadhwa, Fl. Himachal Pradesh 2: 382. 1984.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Mediterranean region.

Local/Common/Vernacular name: *Artichoke thistle* (English)

Dahlia Cav.

Dahlia × pinnata Cav., Icon. 1: 57, t. 80. 1791; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (cultivated & naturalized Belagavi), cultivated & naturalized in many states of India; native to Mexico

Local/Common/Vernacular name: *Garden dahlia* (English)

Dahlia imperialis Roezl ex Ortgies in Gartenflora 12: 243, tt. 407 & 408. 1863; H. Hara, Fl. E. Himal. 337. 1966.

Distribution: Karnataka (cultivated in gardens, also grows as an escape), cultivated in gardens; native to C America.

Local/Common/Vernacular name: *Bell tree dahlia* (English)

Dahlia rosea Cav., Icon. 3: 33, t. 265. 1794; Raizada & H.O. Saxena, Fl. Mussoorie 1: 355. 1978.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Mexico, Southern N America.

Local/Common/Vernacular names: *Dahlia, Pinnate dahlia* (English)

Dichrocephala L'Hér. ex DC.

Dichrocephala chrysanthemifolia (Blume) DC., Archives de Botanique 2: 518. 1833; Udayan et al., Indian For. 130: 558. 2004.

Cotula chrysanthemifolia Blume, Bijdr. 9: 18. 1826.

Distribution: Karnataka (Shivamogga), Kerala, Hiamalayas; Bhutan, Indonesia, Japan, Malaysia, Myanmar, Nepal, New Guinea, Philippines, Africa, Australia.

Dichrocephala integrifolia (L.f.) Kuntze, Revis. Gen. Pl. 1: 333. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Hippia integrifolia L.f., Suppl. Pl. 389. 1782

Grangea latifolia Poir. in Lam., Encycl. Suppl. 2: 826. 1812

Dichrocephala latifolia (Poir.) DC. in Wight, Contr. Bot. India 11. 1834

Dichrocephala bicolor (Roth) Schltdl., Linnaea 25: 209. 1852

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Davanagere, Hassan, Kalaburgi, Kodagu, Shivamogga, Uttara Kannada, Yadgir), throughout warmer parts of India; tropical Africa & Asia.

Dicoma Cass.

Dicoma tomentosa Cass. in Bull. Sci. Soc. Philom. Paris 1817: 12. 1817; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Dicoma lanuginosa DC. in Wight, Contr. Bot. India 26. 1834

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Tumakuru, Vijayapura), throughout India; Tropical Africa.

Local/Common/Vernacular names: *Navanangi, Sanne/Sanni gida* (Kannada)

Doellia Sch.Bip. ex Walp.

Doellia bovei (DC.) Anderb., Willdenowia 25(1): 21. 1995

Conyzia bovei DC., Ann. Sci. Nat., Bot. sér. 2, 2: 261. 1834

Blumea bovei (DC.) Vatke, Linnaea 39: 485. 1875; Babu, N.M.G., Fl. Ethnob. Chitradurga Dist. 301. 2011..

Erigeron bovei (DC.) Boiss., Fl. Orient. 3: 168. 1875

Distribution: Karnataka (Chitradurga, Gujarat), tropical Africa

Echinops L.

Echinops echinatus Roxb., Fl. Ind. ed. 2, 3: 447. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (Bengaluru, Ballari, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kalaburgi, Kolar, Mysuru, Tumakuru, Uttara Kannada, Vijayapura, Yadgir), throughout India; endemic.

Local/Common/Vernacular names: *Brhamadande, Kantulu, Nari mullu, Nayi mullu* (Kannada)

Eclipta L

Eclipta prostrata (L.) L., Mant. Pl. 2: 286. 1771; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Spilanthes pseudoacmella (L.) Murray in L., Syst. Veg. ed. 13. 610. 1774; H. Hara, Fl. E. Himal. 345. 1966.

Verbesina alba L., Sp. Pl. 902. 1753

Verbesina prostrata L., Sp. Pl. 902. 1753

Verbesina pseudoacmella L., Sp. Pl. 901. 1753

Eclipta erecta L., Mant. Pl. 2: 286. 1771

Eclipta alba (L.) Hassk., Pl. Jav. Rar. 528. 1848

Spilanthes acmella var. *calva* sensu Hook.f., Fl. Brit. India 3: 307. 1881 p.p. non C.B. Clarke, 1876

Eclipta prostrata var. *dixitii* AnandKumar & K.K. Khanna, J. Econ. Taxon. Bot. 23: 713. 1999

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular names: *Ajagara*, *Bhringraja*, *Bili bhringaraja*, *Garaga*, *Garagada soppu*, *Garagalu*, *Keshavadhana* (Kannada); *Maka* (*Kunabi*); *Bhringraja* (Sanskrit)

Elephantopus L.

Elephantopus scaber L., Sp. Pl. 814. 1753; Wight, Icon. Pl. Ind. Orient. t. 1086. 1846; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular names: *Ane kalu gida*, *Naayi nalige gida*, *Nela muchilu* (Kannada); *Marchand* (*Kunabi*)

Eleutheranthera Poit.

Eleutheranthera ruderalis (Sw.) Sch.Bip., Bot. Zeitung (Berlin) 24 165 1866

Melampodium ruderalis Sw., Fl. Ind. Occid. 3: 1372. 1806

Ogiera ruderalis (Sw.) Griseb., Pl. Wright. (Grisebach) 2: 513. 1862

Distribution: Karnataka (Dakshina Kannada, Udupi), native to tropical America.

Emilia Cass.

Emilia alstonii Fosberg, Ceylon J. Sci., Biol. Sci. 10: 66. 1972; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (Hassan), Sri Lanka.

Emilia exserta Fosberg, Ceylon J. Sci., Biol. Sci. 10: 65. 1972; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (Hassan, Kodagu, Shivamogga); Sr Lanka.

Emilia ramulosa Gamble, Bull. Misc. Inform. 1921: 120. 1921; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Distribution: Karnataka (Chamarajanagar, Shivamogga), SW Ghats.

Emilia scabra DC., Prodr. 6:303.1838; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Emilia sonchifolia var. *scabra* (DC.) Hook.f., Fl. Brit. India 3: 336. 1881

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru), Peninsular to NE India.

Emilia sonchifolia (L.) DC., Contr. Bot. India 24. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.
Cacalia sonchifolia L., Sp. Pl. 835. 1753
Hieracium javanicum Burm. f., Fl. Indica 174, t. 57, f. 1. 1768
Cacalia sagittata Vahl, Symb. Bot. 3: 91. 1794
Cacalia coccinea Sims in Curtis, Bot. Mag. 16: t. 564. 1802
Senecio sonchifolius (L.) Moench, Suppl. Meth. 231. 1802
Prenanthes javanica Willd., Sp. Pl. 3: 1534. 1803
Emilia flammea Cass. in F. Cuvier's Dict. Sci. Nat. ed. 2. 14: 406, t. 5. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.
Crassocephalum sonchifolium (L.) Less., Syn. Gen. Compos. 395. 1832
Emilia rigidula DC., Prodr. 6: 302. 1838
Emilia sagittata (Vahl) DC., Prodr. 6: 302. 1838
Emilia coccinea (Sims) G. Don in Sweet, Hort. Brit. ed. 3, 382. 1839
Emilia javanica (Burm.f.) C.B.Rob., Philipp. J. Sci. 3: 217. 1908
 Distribution: Karnataka (throughout), throughout India; Pantropical.
 Local/Common/Vernacular name: *Ili kivi* (Kannada)

Emilia zeylanica C.B.Clarke, Compos. Ind. 175. 1876; Hook. f., Fl. Brit. India 3: 336. 1881.
Emilia zeylanica var. *paludosa* Gamble, Fl. Madras 716. 1921
 Distribution: Karnataka (throughout), Peninsular to NE India; Sri Lanka.

Enydra Lour.

Enydra fluctuans Lour., Fl. cochinch. 511. 1790; Hook. f., Fl. Brit. India 3: 304. 1881.
Meyera fluctuans (Lour.) Spreng., Syst. Veg. ed. 16, 3: 602. 1826
 Distribution: Karnataka (throughout), throughout India; native to Asia.
 Local/Common/Vernacular names: *Harkuch, Helencha* (English)

Epaltes Cass.

Epaltes australis Less., Linnaea 5: 148. 1831
Sphaeromorphaea russeliana DC., Prodr. 6: 140. 1838
Erigerodes australe (Less.) Kuntze, Revis. Gen. Pl. 1:335.1891
Sphaeromorphaea australis (Less.) Kitam., Acta Phytotax. Geobot. 6: 80. 1937
 Distribution: Karnataka (Dakshina Kannada, Udupi), E & S India.

Epaltes divaricata (L.) Cass., Bull. Sci. Soc. Philom. Paris 1818: 139. 1818; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.
Ethulia divaricata L., Mant. Pl. 1: 110. 1767
Erigerodes divaricatum (L.) Kuntze, Revis. Gen. Pl. 1:335.1891
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kolar, Mysuru, Raichur, Tumakuru, Uttara Kannada), throughout except NW India; China, S & SE Asia.

Epaltes pygmaea DC., Prodr. 5: 461. 1836; Hook. f., Fl. Brit. India 3: 274. 1881.
Erigerodes pygmaeum (DC.) Kuntze, Revis. Gen. Pl. 1:335.1891
 Distribution: Karnataka (Bengaluru), S India.

Erechtites Raf.

Erechtites hieracifolia (L.) Raf. ex DC., Prodr. 6: 294. 1838; R.R. Rao et al., Fl. Ind. Enum. -Asterac. 36. 1988.
Senecio hieracifolius L., Sp. Pl. 866. 1753
 Distribution: Karnataka (Weed in Bengaluru, Mysuru), C & S India; native to Americas.
 Local/Common/Vernacular names: *American burnweed, Fireweed, Pilewort* (English)

Erechtites valerianifolius (Wolf ex Rchb.) DC., Prodr. 6: 295. 1838; Deb, Fl. Tripura 2: 216. 1983.

Senecio valerianifolius Wolf ex Rchb., Iconogr. Bot. Exot. 1: 59, t. 85. 1827

Distribution: Karnataka (throughout), throughout India; native to Americas.

Erigeron L.

Erigeron annuus (L.) Pers., Syn. Pl. 2: 431. 1807; Raizada & H.O. Saxena, Fl. Mussoorie 1: 375. 1978.

Aster annuus L., Sp. Pl. 875. 1753

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated; native to N America.

Local/Common/Vernacular names: *Annual fleabane, Daisy fleabane, Eastern daisy fleabane* (English)

Erigeron canadensis L., Sp. Pl. 863. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Conyza canadensis (L.) Cronquist, Bull. Torrey Bot. Club 70:632. 1943; T.A. Rao, Rec. Bot. Surv. India 18 (2): 36. 1961.

Erigeron canadensis L., Sp. Pl. 863. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Caenotus canadensis (L.) Raf., Fl. Tellur. 2: 50. 1837

Leptilon canadense (L.) Britton & Brown, Ill. Fl. N. U.S. 3: 391. 1898

Aster canadensis (L.) E.H.L.Krause, Deutschl. Fl. ed. 2. 13: 59. 1905

Trimorpha canadensis (L.) Lindm., Svensk Fanerogamfl. 529. 1918

Conyza canadensis (L.) Cronquist, Bull. Torrey Bot. Club 70:632. 1943; T.A. Rao, Rec. Bot. Surv. India 18 (2): 36. 1961.

Marsea canadensis (L.) V.M.Badillo, Bol. Soc. Venez. Ci. Nat. 10:256. 1946

Distribution: Karnataka (Chamarajanagar, Mysuru), throughout India. native to C & S America.

Local/Common/Vernacular names: *Canadian Fleabane, Canadian Horseweed, Coltstail, Horseweed, Marestail butter weed* (English)

Erigeron floribundus (Kunth) Sch.Bip. in Bull. Soc. Bot. France 12: 81. 1865

Conyza floribunda Kunth in Humb., Bonpl. & Kunth, Nov. Gen. Sp. 4 (ed folio): 57 & (ed. quarto) 73. 1820

Dimorphanthes floribunda (Kunth) Cass., Dict. Sci. Nat., ed. 2, 25: 96. 1822

Erigeron bonariensis var. *floribundus* (Kunth) Cuatrec., Trab. Mus. Nac. Ci. Nat., Ser. Bot. 33: 132. 1936

Conyza sumatrensis var. *floribunda* (Kunth) J.B.Marshall, Watsonia 10(2): 167. 1974

Distribution: Karnataka (Kodagu, Shivamogga, Uttara Kannada)

Erigeron karvinskianus DC., Prodr. 5: 285. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Erigeron mucronatus DC., Prodr. 5: 285. 1836

Felicia erigeroides DC., Prodr. 5: 219. 1836

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chikkamagaluru, Davanagere, Kodagu, Shivamogga), cultivated throughout; native to S America & Mexico.

Local/Common/Vernacular names: *Latin American fleabane, Mexican fleabane, Santa Barbara daisy, Spanish daisy* (English)

Erigeron speciosus (Lindl.) DC., Prodr. 5: 284. 1836; Mahab. in Anonymous, State Gazette. Maharashtra 4: 364. 1987.

Stenactis speciosus Lindl. in Edwards' Bot. Reg. 19: t. 1577. 1833

Distribution: Karnataka (Cultivated).

Local/Common/Vernacular name: *Aspen leabane* (English)

Erigeron sublyratus DC. in Wight, Contr. Bot. India 9. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Erigeron asteroides Roxb., Fl. Ind. ed.2, 3: 432. 1832

Erigeron hispidus DC. in Wight, Contrib. 9. 1834

Aster benghalensis B.Heyne ex DC., Prodr. 5: 292. 1836

Erigeron benghalensis (B.Heyne ex DC.) M.R.Almeida, Fl. Maharashtra 3A:103.2001

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), throughout India; S Asia.

Erigeron wightii DC., Prodr. 5: 286. 1836

Erigeron alpinus var. *wightii* (DC.) Hook.f., Fl. Brit. India 3: 255. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 140. 1984.

Distribution: Karnataka (Kodagu, Vijayapura), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Ethulia L.f.

Ethulia gracilis Del., in Caill., Voy. Meroe. 4: 334, t.64. 1827.

Distribution: Karnataka (Belagavi); North Cameroon, Central African Republic, Chad, Sudan, Ethiopia, Uganda, China.

Eupatorium L.

Eupatorium capillifolium (Lam.) Small ex Porter & Britton, Mem. Torrey Bot. Club 5: 311. 1894; B.D.Sharma et al., Fl. Karnataka Anal. 141. 1984.

Artemisia capillifolia Lam., Encycl. 1: 266. 1783

Artemisia tenuifolia Willd., Sp. Pl. 3: 1819. 1803

Eupatorium foeniculaceum Willd., Sp. Pl. 3: 1750. 1803

Traganthes tenuifolia Wallr., Sched. Crit. 1: 456. 1822

Mikania artemisioides Cass., Dict. Sci. Nat. 55: 130. 1828

Distribution: Karnataka (cultivated in Shivamogga), cultivated in India; native to South Eastern N America.

Local/Common/Vernacular name: *Dogfennel* (English)

Eupatorium repandum Willd., Sp. Pl. 3: 1767. 1803; R.R. Rao et al., Fl. Ind. Enum. -Asterac. 38. 1988.

Distribution: Karnataka (Mysuru), throughout except NW India; native to C & C America.

Eupatorium triplinerve Vahl, Symb. Bot. 3: 97. 1794; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Eupatorium ayapana Vent., Jard. Malmaison 3, t.3. 1803

Ayapana triplinervius (Vahl) R.M. King & H. Rob., Phytologia 20: 212. 1970

Distribution: Karnataka (cultivated in Bengaluru, Chikkamagaluru, Kodagu), cutivated in most gardens; native to tropical America.

Local/Common/Vernacular names: *Ayapana, Water Hemp* (English)

Farfugium Lindl.

Farfugium japonicum (L.) Kitam., Acta Phytotax. Geobot. 8: 268. 1939

Tussilago japonica L., Mant. Pl. 113. 1767

Arnica tussilaginea Burm. f., Fl. indica 182. 1768

Senecio tussilaginea (Burm.f.) Kuntze, Revis. Gen. Pl. 1: 364. 1891

Ligularia tussilaginea (Burm.f.) Makino, Bot. Mag. (Tokyo) 18: 52. 1904; R. Mathur in Hajra et al., Fl. India 13: 323. 1995.

Farfugium tussilagineum (Burm.f.) Kitam., Acta Phytotax. Geobot. 8: 73, 78. 1939

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout India;native to Europe & temperate Asia.

Local/Common/Vernacular names: *Giant leopard plant, Green leopard plant, Leopard plant* (English)

Felicia Cass., nom. cons.

Felicia heterophylla (Cass.) Grau in Mitt. Bot. Staatssamml. München 9: 558. 1973; M.R. Almeida, Fl. Maharashtra 3A: 105. 2001.

Charieis heterophylla Cass., Bull. Sci. Soc. Philom. Paris 1817: 68. 1817

Distribution: Karnataka (cultivated in Bengaluru), cutivated in most gardens; native to S Africa.

Local/Common/Vernacular names: *Kingfisher Daisy, The Blues Blue Daisy* (English)

Flaveria Juss.

Flaveria australasica Hook., J. Exped. Trop. Australia 118. 1848

Distribution: Karnataka (Mysuru), native to Australia

Flaveria trinervia (Spreng.) C. Mohr, Contr. U.S. Natl. Herb. 6: 810. 1901; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Oedera trinervia Spreng., Bot. Gart. Halle 63. 1800

Brotera trinervata (Willd.) Pers., Syn. Pl. 2: 498. 1807

Flaveria contrayerba Pers., Syn. Pl. 2: 489. 1807

Flaveria repanda Lag., Gen. Sp. Pl. 33. 1816

Flaveria trinervata (Willd.) Baill., Hist. Pl. 855. 1882

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kalaburgi, Mysuru, Raichur, Tumakuru, Vijayapura, Yadgi), introduced throughout India, native to Americas.

Local/Common/Vernacular name: *Clustered yellowtops* (English)

Gaillardia Foug. ('Gaillarda').

Gaillardia amblyodon J. Gay, Ann. Sci. Nat. Bot. ser. 2, 12: 62, 1839; R.R. Rao et al., Fl. Ind. Enum. - Asterac. 39. 1988.

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated elsewhere in India; native to USA.

Local/Common/Vernacular names: *Blunt-toothed blanketflower*, *Maroon blanke flower*, *Red gaillardia*, *Red Indian blanket* (English)

Gaillardia aristata Pursh, Fl. Amer. sept. 2: 573. 1814; Deb, Fl. Tripura 2: 219. 1983.

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated elsewhere in India; native to N America.

Local/Common/Vernacular names: *Common blanketflower*, *Common gaillardia* (English)

Gaillardia pulchella Foug., Mém. Acad. Sci. Paris 1786: 5. 1788; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Gaillardia bicolor Lam., Encycl., 2(2): 590. 1788

Distribution: Karnataka (throughout), naturalised throughout India; native to C United States of America.

Local/Common/Vernacular names: *Firewheel*, *Indian blanket*, *Indian blanketflower* (English)

Galinsoga Ruiz & Pav.

Galinsoga parviflora Cav., Icon. 3: 41, t. 281. 1794; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Wiborgia acmella Roth, Catal. Bot fasc. 2. 112. 1800

Galinsoga laciniata Retz. in Hoffm. Phytog. Blaetter i. 46. 1803

Vigolina acmella (Roth) Poir., Encycl. 8: 613. 1808

Wiborgia parviflora (Cav.) Kunth, Nov. Gen. Sp. 4(17): 201 (ed. fol.). 1820

Adventina parviflora (Cav.) Raf., New Fl. 1: 67. 1836

Sabazia microglossa DC., Prodr. 5: 497. 1836

Galinsoga hirsuta Baker, Thirsk Nat. Hist. Soc., Rep. Bot. Exch. Cl. 1861, 13. 1862

Distribution: Karnataka (throughout), naturalised throughout India; Pantropical. weed, native to S America

Gamochaeta Wedd.

Gamochaeta pensylvanica (Willd.) Cabrera, Bol. Soc. Argent. Bot. 9: 375. 1961; Grierson & Spring. in Grierson & D.G. Long, Fl. Bhutan 2(3): 1523. 2001.

Gnaphalium pensylvanicum Willd., Enum. Pl. 2: 867. 1809

Gnaphalium purpureum sensu Hook. f., Fl. Brit. India 3: 289. 1881 non L., 1753

Gnaphalium peregrinum Fernald in Rhodora 45: 479, t. 795. 1943

Distribution: Karnataka (Bengaluru, Mysuru), almost throughout India; native to N America.

Local/Common/Vernacular name: *Pennsylvania cudweed* (English)

Gerbera L., nom. cons.

Gerbera aurantiaca Sch.Bip., Flora 27(2): 781. 1844

Distribution: Karnataka (cultivated throughout except in dry districts), cultivated throughout India; native SE Africa.

Local/Common/Vernacular name: *The Hilton Daisy* (English)

Gerbera jamesonii Bolus, Gard. Chron. 1: 772, f. 122. 1889; M.R. Almeida, Fl. Maharashtra 3A: 108. 2001.

Distribution: Karnataka (cultivated throughout except in dry districts), cultivated throughout India; native SE Africa.

Local/Common/Vernacular names: *Barberton daisy, The Transvaal daisy* (English)

Glebionis Cass.

Glebionis coronaria (L.) Cass. ex Spach, Hist. Nat. Veg. 10:181. 1841

Chrysanthemum coronarium L., Sp. Pl. 890. 1753; Prain, Bengal Pl. 1: 455. 1963 (Repr.).

Matricaria coronaria (L.) Desr., Encycl. 3: 737. 1791

Pyrethrum indicum Roxb., Hort. Bengal. 62. 1814

Chrysanthemum roxburghii Desf. ex Cass., Tabl. École Bot., ed. 3 (Cat. Pl. Horti Paris.) 170. 1829

Pinardia coronaria (L.) Less., Syn. Gen. Compos. 255. 1832

Pinardia roxburghii (Desf. ex Cass.) Less., Fisch. & Mey., Index Sem. Hort. Petrop. 44 1837

Chamaemelum coronarium (L.) E.H.L.Krause, Deutschl. Fl. ed. 2. 13: 203. 1905

Glebionis roxburghii (Desf. ex Cass.) Tzvelev, Bot. Zhurn. (Moscow & Leningrad) 84(7): 117. 1999

Dendranthema coronaria (L.) M.R. Almeida, Fl. Maharashtra 3A: 97. 2001

Xanthophthalmum coronarium (L.) P.D. Sell, Fl. Great Britain & Ireland 4: 556. 2006; Grierson & Spring. in Grierson & D.G. Long, Fl. Bhutan 2(3): 1566. 2001.

Distribution: Karnataka (cultivated in gardens); native to Mediterranean & E Asia.

Local/Common/Vernacular names: *Chop suey green, Chrysanthemum greens, Crown daisy, Edible chrysanthemum, Garland chrysanthemum, Japanese-gree* (English)

Glossocardia Cass.

Glossocardia bidens (Retz.) Veldkamp, Blumea 35: 468. 1991

Zinnia bidens Retz., Observ. Bot. 5: 28. 1788

Glossogyne pinnatifida DC., Contr. Bot. India 19. 1834

Bidens pinnatifida DC., Prodr. 5: 593. 1836

Bidens rigida C.B.Clarke, Compos. Ind. 141. 1876

Glossogyne bidens (Retz.) Alston in Trimen, Handb. Fl. Ceylon 6 (suppl.): 168. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Neuractis bidens (Retz.) Veldkamp ex Mesfin, Kew Bull. 45: 144. 1990

Distribution: Karnataka (Bengaluru, Dharwad, Tumakuru), throughout India.

Local/Common/Vernacular name: *Kalla sabsige soppu* (Kannada)

Glossocardia bosvallia (L.f.) DC., Contr. Bot. India 19. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984 ('bosvallea').

Verbesina bosvallia L. f., Suppl. Pl. 379. 1782

Glossocardia linearifolia Cass., Dict. Sci. Nat. ed. 2, 9: 139. 1817

Glossocardia setosa Blatt. & Hallb., J. Bombay Nat. Hist. Soc. 26: 536. 1919

Distribution: Karnataka (throughout), throughout drier parts of India; endemic.

Local/Common/Vernacular names: *Adavi jirige, Ghatta rasava, Kaadu sabsige, Kallu sabsige, Parapataka, Pita padha, Pitari* (Kannada); *Charak, Parpata, Pithari, Raenu* (Sanskrit)

Gnaphalium L.

Gnaphalium polycaulon Pers., Syn. Pl. 2: 421. 1807; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Gnaphalium indicum auct. non L., 1753; Hook. f., Fl. Brit. India 3: 289. 1881

Gnaphalium multicaule Willd., Sp. Pl., ed. 4, 3: 388. 1803

Gnaphalium niliacum Raddi ex Spreng., Syst. Veg. ed. 16. 480. 1826

Gnaphalium floccosum DC., Prodr. 6: 234. 1838

Gnaphalium gracillimum Perr. ex DC., Prodr. 6: 232. 1838

Gnaphalium schomburgkii Sch.Bip., Reis. Br. 3: 1137. 1849

Distribution: Karnataka (throughout), throughout India; Pantropical.

Gnomophalium Greuter

Gnomophalium pulvinatum (Delile) Greuter, Willdenowia 33(2): 242. 2003

Gnaphalium pulvinatum Delile, Fl. Egypte 266. t. 44. 1812; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Gnaphalium depressum Roxb., Fl. Ind. ed. 2, 3: 425. 1832

Filago prostrata DC., Contr. Bot. India 22. 1834

Gnaphalium crispatum C.B.Clarke, Compos. Ind. 115. 1876

Homognaphalium pulvinatum (Delile) Fayed & Zareh, Willdenowia 18(2): 451. 1989

Distribution: Karnataka (Ballari, Bidar, Kalaburgi, Raichur, Yadgiri), throughout drier parts of India; Africa, China, S & W Asia.

Goniocaulon Cass.

Goniocaulon indicum (Klein ex Willd.) C.B. Clarke, Compos. Ind. 236. 1876

Serratula indica Klein ex Willd., Sp. Pl. 3: 1642. 1803

Goniocaulon glabrum Cass., Bull. Sci. Soc. Philom. Paris 1817:34. 1817; B.D.Sharma et al., Fl. Karnataka Anal. 142. 1984.

Athanasia indica Roxb., Fl. Ind. ed. 2, 3: 417. 1832

Amberboa indica (Klein ex Willd.) DC., Prodr. 6: 558. 1838

Distribution: Karnataka (Ballari, Kalaburgi, Tumakuru), throughout India; endemic.

Grangea Adans.

Grangea maderaspatana (L.) Poir. in Lam., Encycl. Suppl. 2: 825. 1812; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Artemisia maderaspatana L., Sp. Pl. 849. 1753

Cotula maderaspatana (L.) Willd., Sp. Pl., ed. 4, 3(3): 2170. 1803

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular names: *Dodda ghaadhaari, Dovana, Maachi pathre, Masi patre, Muttu karabe, Muttu karasi gida, Pancha pathre* (Kannada)

Guizotia Cass., nom. cons.

Guizotia abyssinica (L.f.) Cass. in F. Cuvier's Dict. Sci. Nat. ed. 2, 59: 248. 1829; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984; Hook.f., Fl. Brit. India 3: 308 1881.

Polymnia abyssinica L.f., Suppl. Pl. 383. 1782

Verbesina sativa Roxb. ex Sims, Bot. Mag. 26: t. 1017. 1807

Anthemis Mysurunsis DC., Prodr. 5: 551. 1836

Bidens ramtilla Wall. ex DC., Prodr. 5: 551. 1836

Distribution: Karnataka (cultivated throughout drier parts), cultivated throughout drier parts of India; native to tropical Africa.

Local/Common/Vernacular names: *Niger plant*, *Niger seeds* (English); *Goorellu*, *Gurellu*, *Hucchellu*, *Kaadellu*, *Raam thil* (Kannada)

Gynura Cass., nom. cons.

Gynura aurantiaca (Blume) Sch.Bip. ex DC., Prodr. 6: 300. 1838; N. Rani & K. M. Matthew in K.M. Matthew, Fl. Tamilnadu Carnatic 3: 798. 1983.

Cacalia aurantiaca Blume, Bijdr. 908. 1826

Crassocephalum aurantiacum (Blume) Kuntze, Revis. Gen. Pl. 1:331. 1891

rassocephalum crepidioides auct. non (Benth.) S. Moore, 1912; Ramam. in Saldanha & Nicolson, Fl. Hassan Dist. 610. 1976.

Distribution: Karnataka (Bengaluru, Chamarajanagar, Hassan), Andaman & Nicobar Islands, Kerala, Tamil Nadu; native to Indonesia..

Local/Common/Vernacular names: *Purple passion*, *Purple passion plant*, *Purple velvet plant*, *Velvet plant* (English)

Gynura bicolor (Roxb. ex Willd.) DC., Prodr. 6: 299. (1837) 1838

Cacalia bicolor Roxb. ex Willd., Sp. Pl. 3: 1731. 1803

Gynura angulosa DC., Prodr. 6: 298. (1837) 1838; Hook. f., Fl. Brit. India 3: 334. 1881.

Gynura angulosa var. *petiolata* Hook. f., Fl. Brit. India 3: 334. 1881

Distribution: Karnataka (Belagavi, Uttara Kannada), almost throughout India; SE Asia.

Gynura cusimbuia (D.Don) S.Moore, J. Bot. 50: 212. 1912; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Cacalia cusimbuia D.Don, Prodr. Fl. Nepal. 179. 1825

Cacalia angulosa Wall., Numer. List no. 3152. 1831

Gynura angulosa Hance, J. Bot. 21: 322. 1883.

Crassocephalum cusimbuia (D.Don) Kuntze, Revis. Gen. Pl. 1 331. 1891

Gynura cusimbuia var. *petiolata* (Hook.f.) R.Mathur in R.R. Rao et al., Fl. Ind. Enum.-Asterac. 42. 1988

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Uttara Kannada), throughout India; Bangladesh, Bhutan, China, Myanmar, Nepal, Thailand

Gynura divaricata (L.) DC., Prodr. 6:301.1838

Senecio divaricatus L., Sp. Pl. 866.1753

Gynura auriculata Cass., Opusc. phytol. 3: 100. 1834; Hook. f., Fl. Brit. India 3: 335. 1881.

Distribution: Karnataka (cultivated in gardens); cultivated in gardens in India; native to China & Vietnam.

Gynura lycopersicifolia DC., Prodr. 6: 300. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Cacalia lycopersicum DC., Prodr. 6: 301. 1838

subsp. *lycopersicifolia*

Distribution: Karnataka (Chikkamagaluru, Davanagere, Hassan, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

subsp. **taprobanensis** Grierson, Ceylon J. Sci., Biol. Sci. 11: 20. 1974

Distribution: Karnataka (Hassan), S to E India; Sri Lanka

Gynura nitida DC., Contr. Bot. India 24. 1834 & Prodr. 6: 299. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984; Hook.f., Fl. Brit. India 3: 333 1881.

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Hassan, Kalaburgi, Kodagu, Shivamogga, Tumakuru, Uttara kannada, Yadgir), Peninsular to E India; endemic.

Gynura travancorica W.W.Sm., Rec. Bot. Surv. India 4: 29. 1914

Distribution: Karnataka (Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Helianthus L.

Helianthus annuus L., Sp. Pl. 904. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Americas.

Local/Common/Vernacular names: *Sun flower* (English); *Arka pushpa, Hothu thirigu hoovu, Soorya kaanthi, Soorya mukhi, Sooryavarthi* (Kannada); *Adhityabhakata, Aaditya valli, Soovarchala* (Sanskrit)

Helianthus argophyllus Torr. & A. Gray, Fl. N. Amer. 2(2): 318. 1842

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to N America.

Local/Common/Vernacular name: *Silver leaf Sunflower* (English)

Helianthus debilis Nutt. in Trans. Amer. Philos. Soc. n.s. 7: 367. 1841; Chandrab. & N.C. Nair, Fl. Coimbatore 164. 1988.

subsp. **debilis**

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to N America.

Local/Common/Vernacular names: *Beach sunflower, Cucumberleaf sunflower, East Coast dune sunflower* (English)

subsp. **cucumerifolius** (Torr. & A.Gray) Heiser, Madroño. 13: 160. 1956

Helianthus cucumerifolius Torr. & A. Gray, Fl. N. Amer. 2(2): 319. 1842; Babu, Herb. Fl. Dehra Dun 270. 1977.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to N America.

Local/Common/Vernacular name: *Cucumberleaf sunflowe* (English)

Helianthus grosseserratus M. Martens, Index Seminum 4, adnot. 1839; Bull. Acad. Sci. Bruxelles 8(1): 67. 1841; Grierson & Spring. in Grierson & D.G. Long, Fl. Bhutan 2(3): 1610. 2001.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to Canada & United States of America.

Local/Common/Vernacular names: *Sawtooth sunflower, Thick-tooth sunflower* (English)

Helianthus lateiflorus Pers., Syn. Pl. 2: 476. 1807; M.R. Almeida, Fl. Maharashtra 3A: 114. 2001.

Distribution: Karnataka (cultivated), cultivated in India; native to N America.

Local/Common/Vernacular names: *Perennial Sunflower, Showy Sunflower* (English)

Helianthus mollis Lam., Encycl. 3: 85. 1789; M.R. Almeida, Fl. Maharashtra 3A: 114. 2001.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental native to Canada & USA.

Local/Common/Vernacular names: *Ashy sunflower, Downy sunflower* (English)

Helianthus pauciflorus Nutt., Gen. N. Amer. Pl. 2: 177. 1818

Harpalium rigidum Cass. in Bull. Sci. Soc. Philom. Paris 1818: 141. 1818

Helianthus rigidus (Cass.) Desf., Tabl. Ecole Bot. ed. 3. 184. 1829; M.R. Almeida, Fl. Maharashtra 3A: 114. 2001.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to Canada & USA.

Local/Common/Vernacular names: *Prairie sunflower, Stiff sunflower* (English)

Helianthus scaberrimus Elliott, Sketch Bot. S. Carolina 2: 423. 1823; Sundararagh. et al., Rec. Bot. Surv. India 21 (2): 45. 1981.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to Canada & USA.

Local/Common/Vernacular name: *cheerful sunflower* (English)

Helianthus tuberosus L., Sp. Pl. 905. 1753; Prain, Bengal Pl. 1: 451. 1963

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental; native to Canada & USA.

Local/Common/Vernacular names: *Jerusalem artichoke, Sunchoke, Sunroot* (English); *Artichoke gedde Sooryakanathi gedde* (Kannada)

Helichrysum Mill. ('Elichrysum')

Helichrysum buddleoides DC., Contr. Bot. India 20. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Gnaphalium buddleoides Sch.Bip., Bot. Zeitung (Berlin) 3:171.1845

Gnaphalium cynoglossoides Sch.Bip. ex Hook.f., Fl. Brit. India 3: 291. 1881

var. **buddleoides**

Distribution: Karnataka (Chikkamagaluru), S India; endemic.

var. **hookeriana** (Wight & Arn.) Hook. f., Fl. Brit. India 3: 291. 1881

Helichrysum hookerianum Wight & Arn. in DC., Prodr. 6: 201. (1837) 1838

Distribution: Karnataka (), Kerala, Tamil Nadu; Sri Lanka.

Helipterum DC.

Helipterum manglesii (Lindl.) F. Muell. ex Benth., Fl. Austral. 3: 640. 1867; M.R. Almeida, Fl. Maharashtra 3A: 115. 2001

Rhodanthe manglesii Lindl. in Edwards' Bot. Reg. 20: t. 1703. 1834

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Australia.

Local/Common/Vernacular names: *Australian Strawflower, Mangles Everlasting, Pink Sunray, Silver Bells, Timeless Rose* (English)

Helipterum roseum Benth., Fl. Austral. 3: 640. 1867; M.R. Almeida, Fl. Maharashtra 3A: 116. 2001.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Australia.

Local/Common/Vernacular name: *Pink paper Daisy* (English)

Kleinia Mill. *Notonia* DC.

Kleinia articulata (L.f.) Haw., Syn. Pl. Succ. 365. 1812; M.R. Almeida, Fl. Maharashtra 3A: 117. 2001.

Cacalia articulata L.f., Suppl. Pl. 364. 1782

Curio articulatus (L.f.) P.V.Heath, Calyx 5(4): 136, 1997

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to S Africa.

Local/Common/Vernacular names: *Candle plant, Hot Dog Cactus* (English)

Kleinia balsamica (Dalzell & A.Gibson) P. Halliday, Kew Bull. 42: 442. 1987

Notonia balsamica Dalzell & A. Gibson, Bombay Fl. 133. 1861

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Kleinia grandiflora (DC.) N.Rani in K.M. Matthew, Fl. Tamilnadu Carnatic 3(1): 801. 1983

Notonia corymbosa DC., Prodr. 6: 442.1838

Notonia grandiflora DC., Contr. Bot. India 24. 1834 & Prodr. 6: 442.1838; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Senecio indicus Backer in K. Heyne, Nutt. Pl. Ned. Ind. 4 :250.1917

Distribution: Karnataka (Chamarajanagar, Chitradurga), Peninsular India; Nepal, Sri Lanka.

Lactuca L.

Lactuca sativa L., Sp. Pl. 795. 1753; Deb, Fl. Tripura 2: 223. 1983.

Lactuca scariola var. *sativa* (L.) Hook. f., Fl. Brit. India 3: 404. 1881

Distribution: Karnataka (cultivated for leafy vegetable), cultivated throughout India; native to Mediterranean to Russia.

Local/Common/Vernacular name: *Lettuce* (English)

Lagascea Cav. ('Lagasca') nom. cons.

Lagascea mollis Cav., Anales Ci. Nat. 6: 332, t. 44. 1803; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Nocca mollis (Cav.) Jacq., Fragm. Bot. 58. 1806

Distribution: Karnataka (throughout), throughout plains of India; native to tropical America.

Laggera Sch.Bip. ex K. Koch

Laggera alata (D.Don) Sch.Bip. ex Oliv., Trans. Linn. Soc. London 39: 94. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Erigeron alatum D.Don, Prodr. Fl. Nepal. 171. 1825

Conyzza alata (D.Don) Roxb., Fl. Ind. ed. 2, 3:430 1832

Blumea alata (D.Don) DC., Prodr. 5: 448. 1836

Vernonia alata (D.Don) Heyne ex DC., Prodr., 5: 448. 1836

Distribution: Karnataka (Belagavi, Ballari, Chamarajanagar, Chitradurga, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; paleotropical.

Local/Common/Vernacular name: *Bettada kuppi gida* (Kannada)

Laggera aurita (DC.) Sch.Bip. ex Schweinf., Beitr. Fl. Aethiop. 151. 1867; B.D.Sharma et al., Fl.

Karnataka Anal. 143. 1984.

Distribution: Karnataka (Belagavi, Mysuru), Pakistan

Laggera crispata (Vahl) Hepper & J.R.I. Wood, Kew Bull. 38: 83. 1983

Conyzza crispata Vahl, Symb. Bot. 1: 71. 1790

Blumea crispata (Vahl) Merxm., Mitt. Bot. Staatssamml. München 20:7.1984

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Kolar, Mysuru, Uttara Kannada), throughout India; Africa.

Laggera pterodonta (DC.) Sch.Bip. ex Oliv. & Hiern in Oliv., Fl. Trop. Afr. 3: 324. 1877; Hook. f., Fl. Brit. India 3: 271. 1881.

Blumea pterodonta DC., Prodr. 5: 448. 1836

Laggera pterodonta (DC.) Benth., Genera Plantarum 2: 290. 1873.

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Hassan, Kodagu, Kolar, Uttara Kannada), throughout India; Myanmar; tropical Africa.

Lamprachaenium Benth.

Lamprachaenium microcephalum (Dalzell) Benth. in Benth. & Hook. f., Gen. Pl. 2: 226. 1873;
B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Plate 18D

Decaneurum microcephalum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 231. 1851

Baccharoides microcephalum (Dalzell) M.R.Almeida, J. Bombay Nat. Hist. Soc. 100: 570. 2003

Distribution: Karnataka (Belagavi, Chikkamagaluru, Shivamogga), Maharashtra; endemic.

Laphangium (Hillard & B.L.Burtt) Tzvelev

Laphangium luteoalbum (L.) Tzvelev, Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. 98(6): 105. 1994

Gnaphalium luteoalbum L., Sp. Pl. 851.1753; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Gnaphalium pallidum Lam., Encycl. 2: 750. 1790

Filaginella luteoalba (L.) Opiz, Seznam 44.1852

Gnaphalium luteoalbum var. *multiceps* (Wall. ex DC.) Hook.f., Fl. Brit. India 3:288.1881

Gnaphalium luteoalbum var. *pallidum* Hook.f., Fl. Brit. India 3:288.1881

Gnaphalium luteoalbum subsp. *pallidum* (Lam.) Maheshw., J. Bombay Nat. Hist. Soc. 57: 377. 1960

Pseudognaphalium luteoalbum (L.) Hilliard & B.L.Burtt, Bot. J. Linn. Soc. 82(3): 206.1981

Pseudognaphalium luteoalbum var. *pallidum* (Lam.) Karthik. & Moorthy, Fl. Pl. India 261. 2009

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), through tropical parts; pantropical.

Local/Common/Vernacular name: *Jersey cudweed* (English)

Launaea Cass.

Launaea acaulis (Roxb.) Babc. ex Kerr in Craib, Fl. Siam. 2: 299. 1936; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Prenanthes acaulis Roxb., Fl. Ind. ed.2, 3: 403. 1832

Lactuca glabra DC., Contr. Bot. India 26. 1834

Youngia acaulis (Roxb.) DC., Prodr. 7: 193. 1838

Microrhynchus glabra (DC.) Wight, Icon. Pl. Ind. Orient. t. 1145. 1846

Microrhynchus acaulis (Roxb.) Kurz, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 56(2): 208. 1877

Crepis acaulis (Roxb.) Hook. f., Fl. Brit. India 3: 396. 1881

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), throughout India including Himalayas ; China, S & SE Asia.

Launaea asplenifolia (Willd.) Hook. f., Fl. Brit. India 3: 415. 1881

Prenanthes asplenifolia Willd., Sp. Pl. 3: 1540. 1803

Distribution: Karnataka (throughout), almost throughout India including Himalayas.

Launaea intybacea (Jacq.) Beauverd, Bull. Soc. Bot. Geneve 2, 2: 114. 1910; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Lactuca intybacea Jacq., Icon. Pl. Rar. 1: 162. 1784

Cicerbita intybacea (Jacq.) Wallr., Sched. Crit. 434. 1822

Phoenixopus intybaceus (Jacq.) Less., Syn. Gen. Compos. 137. 1832

Lactuca heyneana DC., Contr. Bot. India 26. 1834

Lactuca remotiflora DC., Contr. Bot. India 26. 1834

Lactuca runcinata DC. in Wight, Contr. Bot. Ind. 26. 1834; Mamgain & R.R. Rao in Hajra et al., Fl. India 12: 284. 1995.

Launaea runcinata DC., Contr. Bot. India 26. 1834

Brachyramphus intybaceus (Jacq.) DC., Prodr. 7(1): 177. 1838

Sonchus calyculatus Roxb. ex DC., Prodr. (DC.) 7(1): 140.1838

Brachyramphus heyneanus (DC.) Wight, Icon. Pl. Ind. Orient. t. 1146. 1846

Launaea remotiflora (DC.) Stebbins, Indian Forest Rec. 1(6): 240. 1939

Brachyramphus remotiflorus (DC.) Kamelin, Opred. Rast. Sred. Azii 10: 628. 1993

Distribution: Karnataka (Belagavi, Chitradurga, Raichur), throughout India; native to tropical America.

Distribution: Karnataka (weedy in most parts), Peninsula to Himalayas; native to Americas.

Local/Common/Vernacular name: *Prickly Lettuce* (English)

Local/Common/Vernacular name: *Wild-lettuce* (English)

Launaea procumbens (Roxb.) Ramayya & Rajagopal, Kew Bull. 23:465.1969; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Prenanthes procumbens Roxb., Fl. Ind. ed. 2, 3: 404. 1832

Brachyramphus obtusus DC., Prodr. 7(1): 177. 1838

Launaea nudicaulis Hook. f., Fl. Brit. India 3: 416. 1881 non Less., 1832

Launaea fallax (Jaub. & Spach) Kuntze, Revis. Gen. Pl. 351. 1891

Paramicrorhynchus procumbens (Roxb.) Kirp., Fl. URSS 29: 237. 1964

Launaea procumbens (Roxb.) Amin, Student Fl. Egypt 84.. 1956.

Launaea obtusus (DC.) Mundhe & Almeida, J. Bombay Nat. Hist. Soc. 85(3): 523. 1989

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Kalaburgi, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru), throughout drier parts of India; China, C, S & SW Asia.

Launaea sarmentosa (Willd.) Sch.Bip. ex Kuntze, Revis. Gen. Pl. 1: 350. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Prenanthes sarmentosa Willd., Phytographia 10, t. 6, f. 2. 1794

Launaea pinnatifida Cass., Ann. Sci. Nat. (Paris) 23: 85. 1831

Microrhynchus sarmentosus (Willd.) DC., Prodr. 7: 181. 1838

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kalaburgi, Udupi, Uttara Kannada, Vijayapura, Yadgir), Islands & coastal areas, Rajasthan; Africa, Australia, China, SE Asia.

Launaea secunda (C.B.Clarke) Hook.f., Fl. Brit. India 3(8): 416. 1881

Chondrilla secunda Royle ex C.B.Clarke, Compos. Ind. 276. 1876

Microrhynchus secundus C.B.Clarke, Compos. Ind. 276. 1876

Zollikoferia secunda (C.B.Clarke) Boiss, Fl. Orient. Suppl. 323. 1888

Distribution: Karnataka (Tumakuru), Himachal Pradesh, Jammu & Kashmir, Uttaranchal; Afghanistan, Nepal, Pakistan.

Leucanthemum Mill.

Leucanthemum vulgare Lam., Fl. Franç. 2: 137. 1779; B.D.Naithani in Hajra et al., Fl. India 12: 55. 1995.

Chrysanthemum leucanthemum L., Sp. Pl. 888. 1753

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout India; native to Europe & temperate Asia.

Local/Common/Vernacular name: *Ox-eye daisy* (English)

Melampodium L.

Melampodium paludosum Kunth in Humb., Bonpl. & Kunth, Nov. Gen. Sp. 4 (ed. folio): 215 & (ed. quarto) 273. 1820

Distribution: Karnataka (cultivated as an ornamental), cultivated throughout India; native to Europe & temperate Asia.

Local/Common/Vernacular names: *Butter daisy*, *Gold Medallion Flower*, *Melampodium*, *Star Daisy* (English)

Melanthera J. P. Rohr.

Melanthera biflora (L.) Wild, Kirkia 5:4.1965

Verbesina biflora L., Sp. Pl. ed 2. 1272. 1763

Acmella biflora (L.) Spreng., Syst. Veg. 3 591 1826

Adenostemma biflorum (L.) Less., Syn. Gen. Compos. 156. 1832

Wedelia biflora (L.) DC., Contr. Bot. India 18. 1834

Wollastonia biflora (L.) DC., Prodr. 5: 546. 1836; Nicolson et al., Regnum Veg. 119 (Interpr. Rheede Hort. Malab.): 70. 1988.

Wedelia scandens C.B.Clarke, Compos. Ind. 136. 1876

Niebuhria biflora (L.) Britten, J. Bot. 39: 69. 1901

Stemmodontia biflora (L.) W.Wight, Contr. U.S. Natl. Herb. 9: 377. 1905

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Andaman & Nicobar Islands, E & W Coastal regions, E Himalaya

Local/Common/Vernacular name: *Melampodium* (English)

Mikania Willd., nom. cons.

Mikania cordata (Burm.f.) B.L.Rob., Contr. Gray Herb. 104: 65. 1934

Eupatorium cordatum Burm. f., Fl. Indica 176, t. 58, f. 2. 1768

Eupatorium volubile Norona, Verh. Batav. Genootsch. Kunst. Wet. 5, 4.1790

Eupatorium volubile (Poepp.) Vahl, Symb. Bot. 3:93.1794

Mikania volubilis (Vahl) Willd., Sp. Pl., ed. 4, 3(3): 1743. 1803

*Mikania scandens*auct. non (L.) Willd. 1803; Hook. f., Fl. Brit. India 3: 244. 1881

Distribution: Karnataka (throughout), Andaman & Nicobar Islands, NE India; native to tropical Americas.

Mikania micrantha Kunth, Nov. Gen. Sp. 4 (ed. folio): 105 & (ed. quarto)134. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Mikania cordata var. *indica* Kitam. in H. Hara, Fl. E. Himal. 341. 1966

Distribution: Karnataka (throughout), throughout India; native to tropical Americas

Montanoa La Llave & Lex.

Montanoa bipinnatifida (Kunth) K. Koch, Wochenschr. Vereines Beförd. Gartenbaues Königl. Preuss. Staaten 7: 406. 1864; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Polymnia grandis Hort.Berol. ex Kunth, Ind. Sem. Hort. Berol. 13. 1847

Uhdea bipinnatifida Kunth, Index Sem. Berlin 13. 1848

Montanoa heracleifolia Brongn. ex Groenl., Rev. Hort. 5: 544. 544 1857

Montanoa elegans K.Koch, Wochenschr. Vereines Beford. Gartenbaues Konigl. Preuss. Staaten 7: 408. 1864

Montanoa pyramidata Sch.Bip. ex Sch.Bip., Wochenschr. Vereines Beford. Gartenbaues Konigl. Preuss. Staaten 7: 408. 1864

Eriocoma elegans (C.Koch) Kuntze, Revis. Gen. Pl. 1 336.1891

Eriocoma pyramidata (Sch.Bip. ex C.Koch) Kuntze, Revis. Gen. Pl. 1 336.1891

Distribution: Karnataka (Kolar), naturalised throughout India; native to Mexico.

Local/Common/Vernacular names: *Daisy Tree, Pom Pom Tree, Tree Chrysanthemum* (English)

Montanoa hibiscifolia Benth., Vidensk. Meddel. Naturhist. Foren. Kjøbenhavn 1852(5-7): 89. 1853

Eriocoma hibiscifolia (Benth.) Kuntze, Revis. Gen. Pl. 1:336.1891

Distribution: Karnataka (Hassan, Kodagu), Kerala, Tamil Nadu;native to Mexico & C America.

Local/Common/Vernacular names: *Anzac flower, Bush daisy, montanoa, tree daisy* (English)

Moonia Arn.

Moonia arnottiana Wight, Icon. Pl. Ind. Orient. 3(4): 9, t. 1105. 1845; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Chrysogonum arnottianum C.B.Clarke, Compos. Ind. 132. 1876
Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; Sri Lanka.

Nanothamnus Thomson

Nanothamnus sericeus Thomson, J. Linn. Soc., Bot. 9: 342, t. 3. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Plate 18E

Distribution: Karnataka (Shivamogga), Maharashtra; endemic.

Oligochaeta (DC.) K.Koch

Oligochaeta ramosa (Roxb.) Wagenitz, Veröff. Geobot. Inst. ETH Stiftung Rübel. Zürich 37: 323. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 144. 1984.

Carduus ramosus Roxb., Fl. Ind. ed.2, 3: 407. 1832

Microlonchus divaricatus DC., Prodr. 6: 562. 1838

Tricholepis procumbens Wight, Icon. Pl. Ind. Orient. 3(4):12, t. 1139. 1845

Volutarella divaricata (DC.) Benth. in Benth. & Hook . f., Gen. Pl. 2. 476. 1873 excl. syns. *Centaurea patula* DC. & *Serratula divaricata* Fisch. & C.A. Mey

Volutarella ramosa (Roxb.) Santapau, Pl. Saurashtra 22. 1953

Amberboa ramosa (Roxb.) Jafri, Scientist 3: 29. 1959

Distribution: Karnataka (Belagavi, Ballari, Chitradurga, Kalaburgi, Kolar, Raichur, Tumakuru), throughout plains of India.

Parthenium L.

Parthenium hysterophorus L., Sp. Pl. 988. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Distribution: Karnataka (throughout), Weed throughout India; Pantropical., native to tropical America.

Local/Common/Vernacular name: *Congress gida* (Kannada)

Pentacalia Cass.

Pentacalia corymbosa (Benth.) Cuatrec., Phytologia 49: 253. 1981

Microchaete corymbosa Benth., Pl. Hartw. 196. 1845

Monticalia corymbosa (Benth.) C.Jeffrey, Kew Bull. 47: 70. 1992

Distribution: Karnataka (Chamarajanagar)

Pentanema Cass.

Pentanema cernuum (Dalzell) Ling, Acta Phytotax. Sin. 10: 180. 1965

Vicoa cernua Dalzell in Dalzell & A. Gibson, Bombay Fl. 126 & 314. 1861

Jacobaea cernua (Dalzell) Kuntze, Revis. Gen. Pl. 1: 347. 1891

Distribution: Karnataka (Belagavi, Mysuru, Shivamogga, Uttara Kannada), NE & NW India, throughout except; endemic.

Pentanema indicum (L.) Ling, Acta Phytotax. Sin. 10: 179. 1965

Inula indica L., Sp. Pl. ed. 2, 1236. 1763

Vicoa auriculata Cass., Ann. Sci. Nat. (Paris) 17: 418. 1829

Vicoa indica (L.) DC., Contr. Bot. India 10. 1834

Distribution: Karnataka (throughout), throughout India; S Asia, tropical Africa & Asia.

Local/Common/Vernacular name: *Mooguti soppu* (Kannada)

Phyllocephalum Blume

Phyllocephalum anthelminticum (L.) S.R. Paul & S.L. Kapoor, J. Econ. Taxon. Bot. 6: 728. 1985

Conyzanthelmintica L., Sp. Pl. ed. 2. 1207. 1763

Baccharoides anthelmintica (L.) Moench, Metod. 578. 1794

Vernonia anthelmintica (L.) Willd., Sp. Pl., ed. 4, 3(3): 1634. 1803; K. G. Bhat, Fl. S. Kanara 840. 2014.

Serratula anthelmintica (L.) Roxb., Hort. Bengal. 60. 1814

Ascaricida anthelmintica (L.) Sweet, Hort. Brit. 219. 1826

Centratherum anthelminticum (L.) Kuntze, Revis. Gen. Pl. 1: 320. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; S & SE Asia.

Local/Common/Vernacular names: *Kalajeerige, Vakushi* (Kannada)

Phyllocephalum lilacinum (Dalzell & A.Gibson) S.M.Almeida & M.R.Almeida in S.M. Almeida, J. Econ. Taxon. Bot. addl. ser. 8(1): 224. 1990

Decaneurum lilacinum Dalzell & A. Gibson, Bombay Fl. 314. 1861

Baccharoides lilacinum (Dalzell & A.Gibson) M.R.Almeida, Fl. Maharashtra 3A: 72. 2001

Distribution: Karnataka (Uttara Kannada), Goa, Maharashtra; endemic.

Phyllocephalum mayurii (C.E.C.Fisch.) Narayana, Curr. Sci. 51: 438. 1982

Centratherum mayurii C.E.C.Fisch., Bull. Misc. Inform. 1940: 45. 1940

Baccharoides mayurii (C.E.C.Fisch.) Punekar & Vasudeva Rao, J. Econ. Taxon. Bot. 29: 516. 2006

Distribution: Karnataka (Chikkamagaluru, Kodagu, Mysuru, Shivamogga), Tamil Nadu; endemic.

Phyllocephalum phyllolaenum (DC.) Narayana, Curr. Sci. 51(8): 438. 1982

Decaneurum phyllolaenum DC., Prodr. 7: 264. 1838

Gymnanthemum phyllolaenum (DC.) Sch.Bip. ex Walp., Repert. Bot. Syst. 2: 949. 1843

Centratherum phyllolaenum (DC.) Benth. ex C.B.Clarke, Compos. Ind. 4. 1876

Distribution: Karnataka (Hassan), Peninsular India to NE India; endemic.

Phyllocephalum scabridum (DC.) K.Kirkman, Rhodora 83: 10. 1981

Decaneurum scabridum DC., Contr. Bot. India 7. 1834

Decaneurum molle DC., Prodr. 5: 67. 1836

Gymnanthemum molle (DC.) Sch.Bip. ex Walp., Repert. Bot. Syst. 2: 948. 1843

Centratherum courtallense (Wight) Benth. & Hook.f. in Benth. & Hook. f., Gen. Pl. 2(1): 225. 1873

Centratherum molle (DC.) Benth. in Benth. & Hook. f., Gen. Pl. 2: 225. 1873

Centratherum hookeri C.B.Clarke, Compos. Ind. 3. 1876

Centratherum tenue C.B.Clarke, Compos. Ind. 4. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Decaneurum angustifolium Hook.f. & Thomson ex C.B.Clarke, Compos. Ind. 3. 1876

Centratherum metzianum Sch.Bip. ex Hook. f., Fl. Brit. India 3: 228. 1881

Centratherum ritchiei Hook. f., Fl. Brit. India 3: 228. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 138. 1984.

Cacalia clarkeana Kuntze, Revis. Gen. Pl. 1:323. 1891

Cacalia hookeri (C.B.Clarke) Kuntze, Revis. Gen. Pl. 2:970. 1891

Centratherum rangacharii Gamble, Bull. Misc. Inform. Kew 1920, 338. 1920

Phyllocephalum ritchiei (Hook.f. f.) Narayana, Curr. Sci. 51: 438. 1982

Phyllocephalum tenue (C.B.Clarke) Narayana, Curr. Sci. 51: 439. 1982

Phyllocephalum hookeri (C.B.Clarke) Uniyal in Hajra et al. (Eds.), Fl. India 13: 339. 1995

Baccharoides hookeri (C.B.Clarke) M.R. Almeida, Fl. Maharashtra 3A: 72. 2001

Baccharoides ritchiei (Hook.f. f.) M.R. Almeida, Fl. Maharashtra 3A: 73. 2001

Baccharoides scabridum (DC.) M.R.Almeida, Fl. Maharashtra 3A: 73. 2001

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), N & Peninsular India.

Pluchea Cass.

Pluchea ovalis (Pers.) DC., Prodr. 5: 450. 1836; Bhaskar & Kushalappa, Fl. Tumkur Dist. 73. 2013..
Baccharis ovalis Pers., Syn. Pl. 2(2): 424. 1807
Baccharis ovata Sieber ex DC., Prodr. 5: 450. 1836
 Distribution: Karnataka (Tumakuru), Maharashtra

Pluchea paniculata gardneri (Hook.f.) Karthik. & Moorthy, Fl. Pl. India 1: 259. 2009
Conyzapaniculata Willd., Sp. Pl., ed. 4, 3(3): 1920. 1803
Conyzadiffusa Roxb., Hort. Bengal. 61. 1814
Blumeamembranacea DC., Prodr. 5:440.1836; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.
Blumeasenecioides DC., Prodr. 5: 440. 1836
Conyzamembranacea Wall. ex DC., Prodr. 5:441. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 137. 1984.
Blumeahispidula DC. ex C.B.Clarke in Hook.f., Fl. Brit. India 3: 672. 1881
Blumeakingii C.B.Clarke ex Hook.f., Fl. Brit. India 3: 265. 1881
Pluchea senecioides (DC.) W. Theob. in Mason, Burmah ed 4, 2: 380. 1883
Blumeapaniculata (Willd.) M.R.Almeida, Fl. Maharashtra 3A:82.2001
 Distribution: Karnataka (Ballari, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Shivamogga, Uttara Kannada), throughout India; endemic.

var. paniculata

Distribution: Karnataka (throughout), throughout India.

var. jacquemontii (Hook.f.) Karthik. & Moorthy, Fl. Pl. India 1: 259. 2009

Blumeajacquemontii Hook.f., Fl. Brit. India 3: 265. 1881
Blumeakingii C.B.Clarke ex Hook.f., Fl. Brit. India 3: 265. 1881
Blumeamembranacea var. *jacquemontii* (Hook.f.) Randeria, Blumea 10: 271. 1960
 Distribution: Karnataka (throughout), throughout India.

var. subsimplex (Hook.f.) Karthik. & Moorthy, Fl. Pl. India 1: 259. 2009

Blumeasubsimplex DC., Prodr. 5: 441. 1836
Blumeamembranacea var. *subsimplex* (DC.) Hook.f., Fl. Brit. India 3: 265. 1881
 Distribution: Karnataka (throughout), throughout India.

Pluchea tomentosa DC., Contr. Bot. India 16. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Dharwad, Raichur, Vijayapura), throughout India.

Pseudoconyza Cuatrec

Pseudoconyza viscosa (Mill.) d'Arcy, Phytologia 25: 281. 1973

Conyzaviscosa Mill., Gard. Dict. ed. 8, no. 8. 1768

Conyzaurita L.f., Suppl. Pl. 367. 1782

Blumeaurita (L.f.) DC. in Wight, Contr. Bot. India 16. 1834; K.K. Sharma & S. Sharma, Fl. Rajasthan 98. 1989.

Laggeraurita (L.f.) Benth. ex C.B.Clarke, Compos. Ind. 92. 1876.

Distribution: Karnataka (), Plains of India.

Pulicaria Gaertn.

Pulicaria angustifolia DC., Prodr. 5: 479. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Callistephus concolor Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2:344.1850

Pulicaria indica Jaub. & Spach, Illustr. 4: 67. t. 343. 1852

Pulicaria stocksii Hook.f., Fl. Brit. India 3: 300.1881

Distribution: Karnataka (Ballari, Bidar, Chamarajanagar, Chikkamagaluru, Chitradurga, Shivamogga, Tumakuru), throughout India.

Rudbeckia L.

Rudbeckia amplexicaulis Vahl in Skr. Naturhist.- Selsk. 2(2): 29, t.4. 1793; M.R. Almeida, Fl. Maharashtra 3A: 129. 2001.

Dracopis amplexicaulis (Vahl) Cass. ex Less., Syn. Gen. Compos. 226 1832

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental in India; native to N America.

Local/Common/Vernacular names: *Clasping coneflower, clasping-leaf coneflower* (English)

Sclerocarpus Jacq.

Sclerocarpus africanus Jacq. Icon. Pl. Rar. 1: 17, t. 176. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Distribution: Karnataka (Belagavi, Uttara Kannada), throughout India including W Himalaya.

Scorzonera L.

Scorzonera hispanica L., Sp. Pl. 791. 1753; M.R. Almeida, Fl. Maharashtra 3A: 130. 2001.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental in India; native to N America.

Local/Common/Vernacular names: *Black oyster plant, Black salsify, Scorzonera, Serpent root, Spanish salsify, Viper's herb, Viper's-grass* (English)

Senecio L.

Senecio belagaviensis (Wight) C.B. Clarke, Compos. Ind. 200. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Madacarpus belagaviensis Wight, Calcutta J. Nat. Hist. 7: 157. 1847 & Icon. Pl. Ind. Orient. t. 1152. 1846

Distribution: Karnataka (Belagavi, Kodagu, Shivamogga, Uttara Kannada), Goa, Maharashtra; endemic.

Senecio bicolor Ces., Pass. & Gibelli, Comp. Fl. Ital. 2 (22): 517. 1878.

subsp. **cineraria** (DC.) Chater, Bot. J. Linn. Soc. 68: 273. 1974

Senecio cineraria DC., Prodr. 6: 374. (1837) 1838

Distribution: Karnataka (Cultivated).

Senecio bombayensis N.P. Balakr., J. Bombay Nat. Hist. Soc. 67: 61. 1970

Senecio reticulatus (Wight) C.B. Clarke, Compos. Ind. 199. 1876, non DC., 1837

Doronicum reticulatum Wight in Calcutta J. Nat. Hist. 7: 156. 1847 & Icon. Pl. Ind. Orient. t. 1151. 1846

Senecio grahami Hook. f., Fl. Brit. India 3: 347. 1881 non Benth., 1857

Distribution: Karnataka (Belagavi), C, N & Peninsular India.

Senecio candicans DC., Contr. Bot. India 22, 1834 & Prodr. 6: 369. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Cineraria sarmentosa Lesch. ex DC., Prodr. 6: 369. 1838

Distribution: Karnataka (Kolar), Tamil Nadu; endemic.

Senecio dalzellii C.B.Clarke, Compos. Ind. 201. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Senecio lawii C.B.Clarke, Compos. Ind. 201. 1876

Distribution: Karnataka (Belagavi, Chikkamagaluru, Uttara Kannada), Goa, Gujarat, Maharashtra; endemic.

Senecio edgeworthii Hook. f., Fl. Brit. India 3: 346. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Senecio tomentosus (Wight) C.B. Clarke, Compos. Ind. 200. 1876 non Michx., 1803

Doronicum tomentosum Wight, Icon Pl. Ind. Orient. t. 1151. 1846

Senecio lanuginosa var. *tomentosa* (Wight) M.R. Almeida, Fl. Maharashtra 3A: 132. 2001

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Senecio grahamii Hook.f., Fl. Brit. India 3: 347. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Distribution: Karnataka (Chikkamagaluru, Shivamogga, Uttara Kannada), peninsular India; endemic.

Senecio ludens C.B.Clarke, Compos. Ind. 199. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 145. 1984.

Doronicum walkeri Arn., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 18(1): 350. 1836

Distribution: Karnataka (Chikkamagaluru)

Senecio mayurii C.E.C.Fisch., Bull. Misc. Inform. 1939: 250. 1939; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Distribution: Karnataka (Mysuru), S India.

Senecio nudicaulis Buch.-Ham. ex D. Don, Prodr. Fl. Nepal. 179. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Distribution: Karnataka (Shivamogga), throughout India.

Senecio scandens Buch.-Ham. ex D. Don, Prodr. Fl. Nepal. 178. 1825

Senecio wightianus DC. in Wight, Contrib. Bot. India 22. 1834

Senecio intermedius Wight, Icon. Pl. Ind. Orient. t. 1135. 1846

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru), NE & S India, temperate Himalayas.

Senecio scopolii Hoppe & Hornsch. ex Bluff & Fingerh., Companion Fl. German. 2: 380. 1825

Senecio arachnoideus Sieber ex DC., Prodr. 6: 357. 1838

Madaractis arachnoidea C.B.Clarke, Comp. Ind. 199. 1876

Distribution: Karnataka (Uttara Kannada), Goa, Maharashtra; native to S Europe.

Senecio sempervivus Sch.Bip., Flora 28: 500. 1845; T. Cooke, Fl. Bombay 2: 54 1904.

Distribution: Karnataka (cultivated as an ornamental), cultivated in throughout; native to Africa & W Asia.

Senecio tenuifolius Burm.f., Fl. Ind. 181.1768; Ramamoorthy in Saldanha & Nicolson, Fl. Hassan Dist. 622.1976.

Senecio multifidus Willd., Sp. Pl., ed. 4, 3(3): 1986. 1803

Doronicum tenuifolium (Burm.f.) Wight, Icon. Pl. Ind. Orient. t. 1129. 1846

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kalaburgi, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Vijayapura Yadgir), S India; SE Asia.

Distribution: Karnataka (Hassan), peninsular India; endemic.

Local/Common/Vernacular names: *Slender-Leaf Senecio*, *Slender-leaved groundsel* (English)

Senecio vulgaris L., Sp. Pl. 867. 1753; Gamble, Fl. Madras 2: 509. 1967.

Distribution: Karnataka (weed in most districts), throughout India; Cosmopolitan.

Local/Common/Vernacular names: *Birdseed*, *Chikenweed*, *Common Butterweed*, *common groundsel*, *Grimsel*, *Grinsel*, *Ground Glutton*, *Groundsel*, *Grundsel*, *Grundy Swallow*, *Old-man-in-the-Spring*, *Old-man-of-the-spring*, *Ragwort*, *Simson*, *Squaw Weed* (English)

Senecio wightii (DC.) Benth. in Benth. & Hook. f., Gen. Pl. 2: 248. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Doronicum wightii DC., Contr. Bot. India 23. 1834

Senecio saxatilis Wall. ex DC., Prodr. 6: 367. 1838

Senecio pilosiusculus C.B.Clarke, Comp. Ind. 203. 1876

Senecio rabanii C.B.Clarke, Compos. Ind. 202. 1876

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), NE & S India; endemic.

Senecio zeylanicus DC., Prodr. 6: 365. 1838; Manjunatha, Krishna & Pullaiah, Fl. Davanagere Dist. 229. 2004.

Senecio gracilis Arn., Nov. Act. Nat. Cur. 18(1): 349. 1836

Distribution: Karnataka (Davanagere), Peninsular India; Sri Lanka.

Sigesbeckia L.

Sigesbeckia orientalis L., Sp. Pl. 900. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular name: *Antuvina gida* (Kannada)

Solidago L.

Solidago canadensis L., Sp. Pl. 878. 1753; Raizada & H.O. Saxena, Fl. Mussoorie 1: 372. 1978.

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Bengaluru Rurala, Chikkaballapur, Chitradurga, Kolar), cultivated for cut flowers in India, also an escape; native to north-eastern N America.

Local/Common/Vernacular names: *Canadian goldenrod, common goldenrod, giant goldenrod, meadow goldenrod, shorthair goldenrod, tall goldenrod* (English)

Solidago chilensis Meyen, Reise Erde 1: 311. 1834

Solidago microglossa DC., Prodr. 5: 332. 1836; Hajra in Hajra et al., Fl. India 12: 143. 1995.

Distribution: Karnataka (cultivated as an ornamental), cultivated in India, often found as an escape.; native to N America.

Solidago nemoralis Aiton, Hort. kew. ed. 1. 3: 213. 1789; Chandrab. & N.C. Nair, Fl. Coimbatore 163. 1988.

Distribution: Karnataka (cultivated as an ornamental), cultivated in India; native to N America

Local/Common/Vernacular names: *Dwarf goldenrod, Dyersweed goldenrod, Field goldenrod, Gray goldenrod, gray-stem Goldenrod, old-field Goldenrod, Prairie goldenrod* (English)

Sonchus L.

Sonchus arvensis L., Sp. Pl. 793. 1753; Hook. f., Fl. Brit. India 3: 414. 1881 p.p.

Distribution: Karnataka (throughout), throughout India; native to Europe.

Local/Common/Vernacular names: *Corn sow thistle, Dindle gutweed, Field milk thistle, Field sowthistle, Perennial sow-thistle, Swine thistle, Tree sow thistle* (English)

Sonchus asper (L.) Garsault, Fig. Pl. Méd. 4: t. 565. 1764 & Descr. Vertus Pl. 4: 332. 1767; Hook. f., Fl. Brit. India 3: 414. 1881.

Sonchus oleraceus var. *asper* L., Sp. Pl. 794. 1753

Distribution: Karnataka (Belagavi Kalaburgi, Yadgir), throughout India; N America; native to Europe

Local/Common/Vernacular names: *Prickly sow thistle, Sharp-fringed sow thistle, Spiny sow thistle, Spiny-leaved sow thistle* (English)

Sonchus brachyotus DC., Prodr. 7(1): 186. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Sonchus arvensis subsp. *brachyotus* (DC.) Kitam., Mem. Coll. Sci. Kyoto Imp. Univ. Ser. B Biol. 23: 148. 1956
Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Hassan, Mysuru), throughout India; Caucasus, China, Japan, Kazakhstan, Korea, Kyrgyzstan, Mongolia, Russia, Thailand.

Sonchus oleraceus L., Sp. Pl. 794. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dharwad, Hassan, Kalaburgi, Kodagu, Mysuru, Uttara Kannada, Vijayapura), throughout India

Distribution: Karnataka (throughout), throughout India; pantropical.

Local/Common/Vernacular names: *Annual sow thistle*, *Common sowthistle*, *Hare's colwort*, *Hare's thistle*, *Milky tassel*, *Smooth sow thistle*, *Sow thistle* (English)

Sonchus wightianus DC., Prodr. 7: 187. 1838; Bir & M. Sharma, Bull. Bot. Surv. India 26: 219. 1986.

Sonchus lachnocephalus Rech. f. in K. Danske Vid. Selsk., Biol. Skrift., 8(2) (Symb. afghan. 2): 202. 1955

Distribution: Karnataka (Chikkamagaluru, Davanagere, Tumakuru), throughout India; tropical to temperate areas.

Local/Common/Vernacular name: *Tambaku* (Kannada)

Sphaeranthus L.

Sphaeranthus africanus L., Sp. Pl. ed. 2. 1314. 1763; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Distribution: Karnataka (Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, plains of Peninsular to E India; Australia, Africa & Asia.

Sphaeranthus indicus L., Sp. Pl. 927. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Sphaeranthus hirtus Willd., Sp. Pl. 3: 2395. 1803

Distribution: Karnataka (throughout), throughout India; Paleotropical.

Local/Common/Vernacular names: *Karand*, *Karanda gida*, *Mudu gattana gida*, *Mundi kasa* (Kannada)

Sphaeranthus senegalensis DC., Prodr. 5: 370. 1836; Babu, Herb. Fl. Dehra Dun 283. 1977.

Distribution: Karnataka (throughout), throughout India including Andaman & Nicobar Islands; tropical Africa, Asia.

Spilanthes Jacq.

Spilanthes acmella (L.) Murray, Syst. Veg., ed. 13. 610. 1774.

Ceratocephalus acmella (L.) Kuntze, Revis. Gen. Pl. 1:326. 1891

Distribution: Karnataka (Tumakuru), throughout tropical areas.

Local/Common/Vernacular name: *Toothache Plant* (English)

Steirodiscus Less.

Steirodiscus tagetes (L.) Schltr., Bot. Jahrb. Syst. 27: 216. 1899

Othonna tagetes L., Syst. Nat. ed. 10. 1234. 1759

Gamolepis annua Less., Syn. gen. Compos. 251. 1832; R.K. Gupta, J. Bombay Nat. Hist. Soc. 58: 762. 1962.

Distribution: Karnataka (cultivated), cultivated in India; native to S Africa.

Struchium P. Browne

Sparganophorus Boehm ('Sphaganophorus')

Struchium sparganophora (L.) Kuntze, Revis. Gen. Pl. 1: 366. 1891; R.V. Nair, Bull. Bot. Surv. India 8: 202, f. 18. 1966.

Ethulia sparganophora L., Sp. Pl. ed. 2. 1171. 1763

Sparganophorus sparganophora (L.) C. Jeffrey, Kew Bull. 43: 272. 1988

Distribution: Karnataka (), throughout India; native to tropical America.

Symphyotrichum Nees

Symphyotrichum ericoides (L.) G.L.Nesom, Phytologia 77:280.1995

Aster ericoides L., Sp. Pl. 875. 1753; Raizada & H.O. Saxena, Fl. Mussoorie 1: 378. 1978.

Distribution: Karnataka (cultivated), cultivated throughout India; native to Europe & C Asia.

Local/Common/Vernacular names: *Heath aster, White aster* (English)

Synedrella Gaertn., nom. cons.

Synedrella nodiflora (L.) Gaertn., Fruct. Sem. Pl. 2: 456, t. 171, f. 7.1791; B.D.Sharma et al., Fl. Karnataka Anal. 146. 1984.

Verbesina nodiflora L., Cent. Pl. I. 28. 1755

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India including Himalayas; native to tropical America.

Tagetes L.

Tagetes erecta L., Sp. Pl. 887. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Mexico.

Local/Common/Vernacular name: *Marigold* (English); *Chedu hoo* (Kannada)

Tagetes minuta L., Sp. Pl. 887. 1753; Maheshw., J. Bombay Nat. Hist. Soc. 69: 451. 1972.

Distribution: Karnataka (cultivated in Bengaluru), cultivated in India; native to Americas.

Local/Common/Vernacular names: *Black mint, Southern cone marigold, Stinking roger* (English)

Tagetes patula L., Sp. Pl. 887. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Americas.

Local/Common/Vernacular name: *The French marigold* (English)

Tagetes tenuifolia Cav., Icon. 2: 54, t. 169. 1793; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Americas.

Local/Common/Vernacular names: *Lemon marigold, Signet marigold* (English)

Tanacetum L.

Tanacetum cinerariifolium (Trevir.) Sch.Bip., Tanacetum 58. 1844; H.J. Chowdhery in H.J. Chowdhery & Wadhwa, Fl. Himachal Pradesh 2: 421. 1984.

Pyrethrum cinerariifolium Trevir., Index Sem. (Bratislava) App. 2: 2. 1820

Chrysanthemum cinerariifolium (Trevir.) Vis., Fl. dalmat. 2: 88. 1847

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Balkans.

Local/Common/Vernacular name: *Dalmatian chrysanthemum* (English)

Taraxacum Weber ex F.H. Wigg., nom. cons.

Taraxacum insigne Ekman in Dahlst. Tarax. Scandinav. Exsicc. 3: 14. 1913

Distribution: Karnataka (cultivated).

Local/Common/Vernacular name: *Remarkable Dandelion* (English)

Taraxacum javanicum Soest, Wentia 10: 56, f. 27. 1963; M. Sharma & H. Kaur, J. Econ. Taxon. Bot. 18: 732. 1994.

Distribution: Karnataka (Bengaluru, Kodagu), throughout India; S & SE Asia.

Taraxacum officinale F.H.Wigg., Prim. Fl. Holsat. 56. 1780; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Distribution: Karnataka (Bengaluru), throughout India including Himalayas; cosmopolitan in temperate areas; native to Eurasia.

Tarlmounia H.Rob., S.C.Keeley, Skvarla & R.Chan

Tarlmounia elliptica (DC.) H.Rob., S.C.Keeley, Skvarla & R.Chan, Proc. Biol. Soc. Washington 121(1): 32. 2008

Vernonia elliptica DC. in Wight, Contr. bot. India 5. 1834; Uniyal in Hajra et al. (Eds.), Fl. India 13: 370. 1995.

Vernonia elaeagnifolia DC., Prodr., 5: 22. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Strobocalyx elaeagnifolia (DC.) Sch.Bip., Pollichia 18/19: 171. 1861

Cacalia elaeagnifolia (DC.) Kuntze, Revis. Gen. Pl. 2:970.1891

Distribution: Karnataka (cultivated), India, Myanmar, Thailand; introduced elsewhere.

Thymophylla Lag.

Thymophylla tenuifolia (Cass.) Rydb., N. Amer. Fl. 34:173.1915

Hymenatherum tenuifolium Cass., Dict. Sci. Nat., ed. 2, 22: 314. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 143. 1984.

Dyssodia tenuifolia (Cass.) Loes., Bull. Herb. Boissier ser. 2, 6: 866. 1906

Distribution: Karnataka (Bengaluru, Chitradurga, Dharwad), native to C America.

Tithonia Desf. ex Juss.

Tithonia diversifolia (Hemsl.) A.Gray, Proc. Amer. Acad. Arts 19: 5. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Mirasolia diversifolia Hemsl., Biol. Cent.-Amer., Bot. 2: 168, t. 47. 1881

Urbanisol tagetiflora var. *diversifolius* (Hemsl.) Kuntze, Rev. Gen. Pl. 1: 371.1891

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Introduced & naturalised; native to tropical America.

Local/Common/Vernacular names: *Japanese sunflower*, *Mexican sunflower*, *Mexican tournesol*, *Nitobe chrysanthemum*, *Tree marigold* (English)

Tithonia rotundifolia (Mill.) S.F. Blake, Contr. Gray Herb. 52: 41. 1917; Raizada, Indian Forester 92: 325. 1966.

Tagetes rotundifolia Mill., Gard. Dict. ed. 8. no. 4. 1768

Distribution: Karnataka (cultivated & naturalized in Belagavi, Uttara Kannada), cultivated & naturalized weed in Goa, Kerala, Maharashtra, Tamil Nadu; native to C & N America & The Caribbean Islands.

Tragopogon L.

Tragopogon dubius Scop., Fl. Carniol. ed. 2. 2: 95. 1772; Kachroo et al., Fl. Ladakh 106. 1977.

Tragopogon pratense auct. non L., 1753; Hook. f., Fl. Brit. India 3: 418. 1881

Distribution: Karnataka (naturalized in Kodagu ?), cultivated & naturalized cooler areas; native to S & C Europe & W Asia.

Local/Common/Vernacular names: *Common salsify*, *Goat's beard*, *Goatsbeard*, *Salsify*, *Western goat's-beard*, *Western salsify*, *Wild oysterplant*, *Yellow goat's beard*, *Yellow salsify* (English)

Tricholepis DC.

Tricholepis amplexicaulis C.B.Clarke, Compos. Ind. 240. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), peninsular India; endemic.

Tricholepis angustifolia DC., Prodr. 6: 564. (1837) 1838; Hook. f., Fl. Brit. India 3: 382. 1881.

Tricholepis glaberrima var. *angustifolia* (DC.) L.B.Chaudhary & A.K.Pandey, Rheedia 4: 45. 1994

Distribution: Karnataka (Dakshina Kannada, Udupi), S India.

Tricholepis glaberrima DC., Prodr. 6: 564. (1837) 1838; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), C, N & Peninsular India.
 Local/Common/Vernacular name: *Brahma dande* (Sanskrit)

Tricholepis radicans (Roxb.) DC. In Wight, Contr. Bot. India 25. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Carduus radicans Roxb., Fl. Ind. ed. 2, 3: 408. 1832

Distribution: Karnataka (Belagavi, Chitradurga, Davanagere, Hassan, Kalaburgi, Mysuru, Yadgir), Peninsular India, Rajasthan.

Local/Common/Vernacular names: *Dhana, Mandi kusubi gida* (Kannada)

Tridax L.

Tridax procumbens L., Sp. Pl. 900. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Distribution: Karnataka (throughout), throughout India; Pantropical.; native to S America.

Local/Common/Vernacular names: *Coat buttons, Tridax daisy* (English); *Adikke soppu, Kari balli, Sanna gabbushyavantige, Tike soppu* (Kannada)

Verbesina L.

Verbesina encelioides (Cav.) Benth. & Hook.f. ex A.Gray, Bot. California 1: 350. 1876 ('enceliodes'); B.D.Sharma et al., Fl. Karnataka Anal. 147. 1984.

Ximenesia encelioides Cav., Icon. 2: 60, t. 178. 1793

Distribution: Karnataka (Bengaluru, Kolar), throughout except N W India, Himalayas & Islands India; native to Mexico & USA.

Local/Common/Vernacular names: *American dogweed, Butter daisy, Cowpen daisy, Crown-beard, Golden crownbeard, South African daisy* (English)

Vernonia Schreb., nom. cons.

Vernonia altissima Nutt., Gen. N. Amer. Pl. 2: 134. 1818; Uniyal in Hajra et al., Fl. India 13: 394. 1995.

Distribution: Karnataka (cultivated), cultivated in S India; native to N America.

Local/Common/Vernacular name: *Tall ironweed* (English)

Vernonia arborea Buch.-Ham., Trans. Linn. Soc., London 14: 218. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Plate 18F

Monosis wightiana DC., Contr. Bot. India 1. 1834

Strobocalyx arborea (Buch.-Ham.) Sch.Bip., Jahresber. Pollichia 18/19: 171. 1861

Vernonia arborea var. *wightiana* Hook. f., Fl. Brit. India 3: 239. 1881

Gymnanthemum arboreum (Buch.-Ham.) H.Rob., Proc. Biol. Soc. Wash. 112:240.1999

Distribution: Karnataka (Hassan, Kodagu, Mysuru, Uttara Kannada), Andaman & Nicobar Islands, NE & Peninsular India; endemic.

Vernonia cinerascens Sch.Bip. in Schweinf., Beitr. Fl. Aethiop. 162.1867; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Orbivestus cinerascens (Sch.Bip.) H.Rob., Proc. Biol. Soc. Wash. 112(1): 231.1999

Distribution: Karnataka (Chikmagalur, Mysuru), Kerala, Maharashtra, Punjab, Tamil Nadu; Africa, Arabia, Pakistan?

Vernonia conyzoides DC., Contr. Bot. India 6. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Distribution: Karnataka (Chitradurga, Davanagere, Hassan), Gujarat, Kerala, Maharashtra, Rajasthan, Tamil Nadu; endemic.

Distribution: Karnataka (cultivated in Bengaluru, Dakshina Kannad, Kalaburgi, Mysuru, Udupi), throughout India;

Local/Common/Vernacular names: *Curtain creeper, Vernonia creeper* (English), *Kare hindi, Sahadevi* (Kannada)

Vernonia malabarica Hook. f., Fl. Brit. India 3: 230. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Cacalia malabarica (Hook.f.) Kuntze, Revis. Gen. Pl. 2:970.1891

Vernonia bababudensis Talbot, J. Bombay Nat. Hist. Soc. 11: 237. 1897

Distribution: Karnataka (Chikkamagaluru), SW Ghats; endemic.

Vernonia pectiniformis DC., Contr. Bot. India 2. 1834 & Prodr. 5: 31. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 148. 1984.

Cacalia pectiniformis Kuntze, Revis. Gen. Pl. 2:970.1891

Gymnanthemum pectiniforme (DC.) H.Rob., Proc. Biol. Soc. Washington 112(1): 242. 1999

Distribution: Karnataka (Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka..

Vernonia pulneyensis Gamble, Bull. Misc. Inform. 1920: 341. 1920

Cacalia travancorica Kuntze, Revis. Gen. Pl. 2:971.1891

Distribution: Karnataka (Chikmaguluru, Dakshina Kannada), Kerala, Tamil Nadu; endemic. Tamil Nadu; endemic.

Vittadinia A.Rich.

Vittadinia australis A.Rich., Ess. Fl. N. Zel. 251.1832; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Distribution: Karnataka (Kodagu), native to Australia.

Local/Common/Vernacular name: *White fuzzweed* (English)

Wedelia Jacq., nom. cons.

Wedelia chinensis (Osbeck) Merr., Philipp. J. Sci., C 12: 111. 1917; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Verbesina calendulacea L., Sp. Pl. 902. 1753

Solidago chinensis Osbeck, Dagb. Ostind. Resa 241.1757

Wedelia calendulacea auct. non Pers. 1807. (L.) Less., Syn. Gen. Compos. 222. 1832

Distribution: Karnataka (throughout), Andaman & Nicobar Islands, E & W Coasts, E Himalaya, NE India; native to China.

Local/Common/Vernacular name: *Kesa raji* (Kannada)

Wedelia glauca (Ortega) O. Hoffm. ex Hicken in Apuntes Hist. Nat. 2: 254. 1910; M.R. Almeida, Fl. Maharashtra 3A: 147. 2001.

Pascalia glauca Ortega, Nov. Pl. Descr. Dec. 39, t. 4. 1797

Distribution: Karnataka (Cultivated).

Wedelia trilobata (L.) Hitchc., Annual Rep. Missouri Bot. Gard. 4: 99. 1893; Sivar. & Pradeep, Indian J. Forest. 11: 161, f. 7. 1988.

Silphium trilobatum L., Syst. Nat. ed. 10. 232. 1759

Distribution: Karnataka (cultivated in gardens & naturalized throughout especially in W Ghats), cultivated; native to tropical America.

Xanthium L.

Xanthium strumarium L., Sp. Pl. 987. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Xanthium indicum J. König ex Roxb., Fl. Ind. ed. 2, 3: 601. 1832

Xanthium pungens Wallr. in Beitr. Bot. 1 : 231. 1844 emend. Widder, Repert. Spec. Nov. Regni. Veg. Beih. 20: 64. 1923 & Phyton (Horn) 12: 186. 1967; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Distribution: Karnataka (throughout), weedy throughout India; native to tropical America..

Local/Common/Vernacular name: *Cocklebur* (English); *Maralu ummatti* (Kannada)

Younghia Cass.

Younghia japonica (L.) DC., Prodr. 7: 194. 1838; R.K. Gupta, Fl. Nainital. 203. 1968.

Prenanthes japonica L., Mant. Pl. 1: 107. 1767

Younghia napifolia Wight, Icon. Pl. Ind. Orient. t. 1147. 1846

Crepis japonica (L.) Benth., Fl. Hongk. 194. 1861; Hook. f., Fl. Brit. India 3: 395. 1881.

Distribution: Karnataka (weed in Bengaluru), throughout India; native to E Asia.

Local/Common/Vernacular name: *Oriental False hawksbeard* (English)

Zinnia L., nom. cons.

Zinnia angustifolia Kunth, Nov. Gen. Sp. 4(17): 197 (ed. fol.). 1818

Zinnia linearis Benth., Pl. Hartw. 17. 1839; N.M.G. Babu, Fl. Ethnob. Chitradurga Dist. 321. 2011.

Crassina angustifolia (Kunth) Kuntze, Revis. Gen. Pl. 1:331 1891

Crassina linearis (Benth.) Kuntze, Revis. Gen. Pl. 1:331.1891

Distribution: Karnataka (cultivated in most districts), cultivated throughout India; native to Mexico.

Zinnia elegans L., Jacq., Collectanea 3: 152. 1789; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Distribution: Karnataka (cultivated), cultivated throughout; native to Mexico.

Zinnia linearifolia Moench, Suppl. Meth. 261. 1802

Distribution: Karnataka (cultivated in Belgaum), cultivated throughout in India; native to SW USA & S America.

Zinnia peruviana (L.) L., Syst. Nat. ed. 10. 190, 1221. 1759; B.D.Sharma et al., Fl. Karnataka Anal. 149. 1984.

Zinnia multiflora L., Sp. Pl. ed. 2. 1269. 1763

Zinnia pauciflora L., Sp. Pl. ed. 2. 1269. 1763

Zinnia tenuiflora Jacq., Icon. Pl. Rar. 3: t. 590. 1786; M.R. Almeida, Fl. Maharashtra 3A: 149. 2001.

Distribution: Karnataka (cultivated in Kolar, Tumakuru), cultivated throughout in India; native to Peru.

Local/Common/Vernacular name: *Peruvian Zinnia* (English)

Balanophoraceae

Balanophora J. R. Forst. & G. Forst.

Balanophora abbreviate Blume, Enum. Pl. Javae 1: 87.1827; B.D.Sharma et al., Fl. Karnataka Anal. 243. 1984.

Balaniella abbreviata (Blume) Tiegh., Ann. Sci. Nat., Bot. IX, 6: 181.1907

Distribution: Karnataka (Chitradurga, Mysuru), Andaman & Nicobar Islands, Peninsular, E & NE India; Africa, Madagascar, Pacific Islands, SE Asia.

Balanophora dioica R.Br. ex Royle, Ill. Bot. Himal. Mts. 330.1836; Hook.f., Fl.Brit.India 5:237.1856; Triveni et al., Nelumbo 58: 108. 2016.

Distribution: Karnataka (Chikkamagaluru, Kodagu), Arunachal Pradesh, Manipur, Meghalaya, West Bengal; Nepal.

Balanophora fungosa J. R. & G. Forst., Char. Gen. Pl. 50.1775; B.D.Sharma et al., Fl. Karnataka Anal. 243. 1984.

Langsdorffia indica Arn., Nat. Hist. 2: 37.1838

Balanophora indica (Arn.) Griff., Trans. Linn. Soc. London 20: 95.1846

Balanophora burmannica Griff., Proc. Linn. Soc. London 1: 219.1849

Balanophora picta Griff., Proc. Linn. Soc. London 1: 220.1849

Balanophora indica var. *minor* Eichler, Prodr. 17: 145.1873

Balanophora elkinsii Blatt., J. Bombay Nat. Hist. Soc. 33: 309.1929

Balanophora fungosa var. *indica* (Arn.) B.Hansen, Dansk Bot. Ark. 28: 100.1972

Balanophora indica var. *globosa* (Jungh.) Ban, Konsp. Sosud. Rast. Fl. V'etnama 2: 45.1996

Balanophora indica var. *agastyamalayana* M.B.Viswan. & Prem Kumar & N. Ramesh, Rheeda 10: 121.2000

Balanophora indica var. *tirunelveliensis* M.B.Viswan. & Prem Kumar & N.Ramesh, Rheeda 10: 123.2000

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Hassan, Kodagu, Mysuru), Peninsular to NE India; Australia, E, S & SE Asia, New Guinea, Pacefic Islands.

Balanophora polyandra Griff., Trans. Linn. Soc. London 20: 94.1851; B.D.Sharma et al., Fl. Karnataka Anal. 243. 1984.

Polyplethia polyandra (Griff.) Tiegh., Bull. Soc. Bot. France 43: 298.1896

Acroblastum polyandrum (Griff.) Setch., Univ. Calif. Publ. Bot. 19: 147.1935

Distribution: Karnataka (Mysuru), Arunachal Pradesh, Assam, Manipur, Maharashtra, Nagaland, Sikkim, West Bengal; Bhutan, China, Myanmar, Nepal.

Balsaminaceae

Impatiens L.

Impatiens aadishankarii Bhaskar & Sringsh, Taxon. Monogr. Impatiens W Ghats 118. 2012

Plate 19A

Distribution: Karnataka (Shivamogga); endemic.

Impatiens acaulis Arn., J. Hooker, Comp. Bot. Mag. 1: 325. 1835; B.D.Sharma et al., Fl. Karnataka Anal. 36. 1984.

var. **acaulis**

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Rajasthan, Tamil Nadu; Sri Lanka.

var. **granulata** Bhaskar, Razi & Yogan., Curr. Sci. 44 (5). 622. 1975; B.D.Sharma et al., Fl. Karnataka Anal. 36. 1984.

Distribution: Karnataka (Chikkamagaluru, Shivamogga, Uttar Kannada); endemic.

Impatiens agumbeana Bhaskar & Razi, J. Bombay Nat. Hist. Soc. 79: 382. 1983; B.D.Sharma et al., Fl. Karnataka Anal. 36. 1984.

Plate 19B

Distribution: Karnataka (Hassan, Mysuru, Shivamogga); endemic.

Impatiens aliciae C.E.C.Fisch., Bull. Misc. Inform. Kew 1934: 389. 1934; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala; endemic.

Impatiens bababudenensis Hook.f., Bull. Misc. Inform. Kew 295. 1910; Bhaskar, Taxon. Monogr. Impatiens W. Ghats 295. 2012

Distribution: Karnataka (Chikkamagaluru, Mysuru); endemic.

Impatiens balsamina L., Sp. Pl. 938. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

var. **balsamina**; C.J.Saldanha, Fl. Karnataka 2: 249. 1996

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada and also cultivated throughout); cultivated throughout India; native to SE Asia and also cultivated, introduced elsewhere.

Local/Common/Vernacular name: *Karnakundala* (Kannada)

var. **macrantha** Hook.f., Fl. Brit. India 454. 1874; Bhaskar, Taxon. Monogr. Impatiens W. Ghats 188. 2012

Distribution: Karnataka (Belagavi), W Peninsular India; endemic.

var. **micrantha** Hook.f., Fl. Brit. India 1: 454. 1874; C.J.Saldanha, Fl. Karnataka 2: 250. 1996

Distribution: Karnataka (Hassan, Kodagu); endemic.

var. **parusnathica** Hook.f., Rec. Bot. Surv. India 4 (3): 50. 1904; Bhaskar, Taxon. Monogr. Impatiens W. Ghats 188. 2012

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chikkaballapura), Jharkhand, Kerala, Tamil Nadu; endemic.

var. **rosea** (Lindley) Hook.f., Fl. Brit. India 1: 454. 1874

Impatiens rosea Lindl., Edward's Bot. Reg. Misc. 6. t. 27. 1841; C.J.Saldanha, Fl. Karnataka 2: 250. 1996

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Hassan, Kodagu, Shivamogga), Goa, Gujarat, Himalayas, Maharashtra, Tamil Nadu; endemic.

Impatiens barberi Hook.f., Rec. Bot. Surv. India 4: 45. 1906; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Hassan, Mysuru, Shivamogga); endemic.

Impatiens bhaskarii Desai et al. in Taiwania 54: 149. 2009; Dessai and Janarthanam, Rheedea 30. 2011

Distribution: Karnataka (Chikkamagaluru); endemic.

Impatiens bhimgadensis Bhaskar & Sringeswara, Curr. Sci. 114: 443. 2018

Distribution: Karnataka (Belagavi); endemic.

Impatiens brevicornis (E.Barnes) Bhaskar, Taxon. Monogr. Impatiens W. Ghats 124. 2012

Impatiens chinensis var. *brevicornis* Branes, J. Indian Bot. Soc. 18: 99. 1939; C.J.Saldanha, Fl. Karnataka 2: 251. 1996

Distribution: Karnataka (Chikkamagaluru, Hassan), Kerala, Tamil Nadu; endemic.

Impatiens chinensis L., Sp. Pl. 937. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Davanagere, Hassan, Mysuru), Andhra Pradesh, Arunachal Pradesh, Assam, Kerala, Maharashtra, Manipur, Meghalaya, Mizoram, Tamil Nadu, Tripura, West Bengal; China.

Impatiens clavata Bhaskar, Curr. Sci. 9: 1139. 2006

Distribution: Karnataka (Hassan); endemic.

Plate 19C

Impatiens crenata Bedd., Madras J. Lit. Sci. Ser. 2, 20: 69. t. 10. 1859

Distribution: Karnataka (Kodagu?), Tamil Nadu; endemic.

Impatiens cuspidata Wight & Arn., Comp. Bot. Mag. 1: 321. 1835; R.R.Rao & Razi, Fl. Mysore 485. 1981

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Impatiens dasysperma Wight in Madras J. Lit. Sci. ser. 1, 5: 7, t. 2. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Impatiens lucida B.Heyne ex Hook.f., Fl. Brit. India 1: 451. 1874

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Impatiens dendricola C.E.C.Fisch., Bull. Misc. Inform. Kew 157. 1935; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Kodagu), Kerala; endemic.

Impatiens denisonii Bedd., Madras J. Lit. Sci. Ser. 3, 1: 41. 1864; Bhaskar, Taxon. Monogr. Impatiens W. Ghats 65. 2012

Distribution: Karnataka (Mysuru?), Tamil Nadu; endemic.

Impatiens diversifolia B.Heyne ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 139. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Impatiens fasciculata Lam., Encycl. 1 (2): 363. 1785

Impatiens vivekananthanii J. Dessai & Janarth., Rheedea 21(1): 58. 2011

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Tamil Nadu; Sri Lanka.

Impatiens flaccida Arn., Companion Bot. Mag. 1: 322. 1835; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Chikkamagaluru, Mysuru), Kerala, Tamil Nadu; endemic.

Impatiens fruticosa Lesch. ex DC., Prodr. 1: 687. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 37. 1984.

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Impatiens gardneriana Wight, Icon. Pl. Ind. Orient. 3: t. 1050. 1846; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Plate 19D

Impatiens setosa Hook.f. & Thomson, Proc. Linn. Soc., Bot. 4: 123. 1860

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga), Kerala, Tamil Nadu; endemic.

Impatiens goughii Wight, Illustr. 1: 160. 1840; C.J.Saldanha, Fl. Karnataka 2: 253. 1996

Impatiens anamallayensis Bedd. in Madras J. Lit. Sci. n.s., 20: 68. 1859

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Impatiens henslowiana Arn., Companion Bot. Mag. 1: 322. 1836; C.J.Saldanha, Fl. Karnataka 2: 253. 1996

Impatiens albida Wight in Madras J. Lit. Sci. ser. 1, 5: 7. 1837

Distribution: Karnataka (Kodagu), Kerala; endemic.

Impatiens herbicola Hook. f., Bull. Misc. Inform. Kew 1911: 354. 1911; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Distribution: Karnataka (Hassan, Shivamogga), Kerala, Tamil Nadu; endemic.

Impatiens inconspicua B.Heyne ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 139. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Impatiens pusilla B.Heyne, Numer. List 4745 1831

Impatiens filiformis Wight & Arn., Prod. 140. 1834

var. **inconspicua**

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Uttara Kannada), Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

var. **jogensis** Bhaskar, Taxon. Monogr. Impatiens W. Ghats 145. 2012

Distribution: Karnataka (Shivamogga); endemic.

var. **nematostachys** (Hook.f.) Bhaskar, Taxon. Monogr. Impatiens W. Ghats 145. 2012

Distribution: Karnataka (Chikkamagaluru), Tamil Nadu; endemic.

var. **pushpagiriensis** L.Joseph & Bhaskar, Taxon. Monogr. Impatiens W. Ghats 147. 2012

Distribution: Karnataka (Kodagu); endemic.

Impatiens jerdoniae Wight, Icon. Pl. Ind. Orient. 4 (4): 15, t. 1602. 1850; Bhaskar & Razi, Indian J. Forest.

1: 191 – 198. 1978

Distribution: Karnataka (Kodagu, Mysuru), Tamil Nadu; endemic.

Impatiens kleiniformis Sedgw., Rec. Bot. Surv. India 6: 351. 1919; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Distribution: Karnataka (Belagavi, Uttara Kannada), Goa, Maharashtra; endemic.

Impatiens kodachadriensis Bhaskar & Sringesh, Taxon. Monogr. Impatiens W. Ghats 150. 2012

Plate 19E

Distribution: Karnataka (Shivamogga); endemic.

Impatiens kotebettii Bhaskar & Sringeswara, Webbia 72 (2) 165. 2017

Distribution: Karnataka (Kodagu); endemic.

Impatiens latifolia L., Sp. Pl. 937. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Uttara Kannada), Maharashtra; endemic.

Impatiens lawii Hook.f. & Thomson, J. Linn. Soc. Bot. 4: 122. 1860; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Uttara Kannada), Kerala, Maharashtra

Impatiens lawsonii Hook.f., Rec. Bot. Surv. India 4: 39 1906; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala; endemic.

Impatiens madapurae Bhaskar & Sringeswara, Webbia 72 (2) 165. 2017

Distribution: Karnataka (Kodagu); endemic.

Impatiens minor (DC.) Bennet, Indian J. For. 2: 283. 1979; C.J.Saldanha, Fl. Karnataka 2: 255. 1996

Balsamina minor DC., Prodr. 1: 686. 1824

Impatiens kleinii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 140.1834

var. **minor**

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra; endemic.

var. **hirsuta** Bhaskar, Taxon. Monogr. Impatiens W. Ghats 158. 2012

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Impatiens mullaingiriensis Bhaskar, Taxon. Monogr. Impatiens W. Ghats 159. 2012
Distribution: Karnataka (Chikkamagaluru); endemic.

Impatiens mysorensis Roth, Nov. Pl. Sp. 164. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Plate 19F

Balsamina mysorensis DC., Prodr. 1: 686. 1824

Distribution: Karnataka (Chikkaballapur, Chitradurga, Kolar, Mysuru); endemic.

Local/Common/Vernacular name: *Chikka basavanapada* (Kannada)

Impatiens oppositifolia L., Sp. Pl. 937. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 38. 1984.

Impatiens rosmarinifolia Retz., Observ. Bot. 5: 29 1788

Balsamina oppositifolia (L.) DC., Prodr. 1: 686. 1824

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Myanmar, Sri Lanka.

Impatiens pendula B.Heyne ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 136. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

Plate 19G

Distribution: Karnataka (Chikkamagaluru); endemic.

Impatiens pulcherrima Dalzell in Hook.'s J. Bot. Kew Gard. Misc. 2: 37. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

Distribution: Karnataka (Belagavi, Uttara Kannada), Goa, Maharashtra, Tamil Nadu; endemic.

Impatiens raziana Bhaskar & Razi, J. Bombay Nat. Hist. Soc. 79: 383. 1983; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Shivamogga); endemic.

Impatiens rupicola Hook.f., Bull. Misc. Inform. Kew 292. 1910; Bhaskar, Taxon. Monogr. Impatiens W. Ghats 172. 2012

Impatiens chinensis var. *rupicola* (Hook.f.) Bhaskar, Taxon. Monogr. Impatiens W. Ghats 172. 2012

var. **rupicola**

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

var. **nataliae** (Hook.f.) Bhaskar, Taxon. Monogr. Impatiens W. Ghats 174. 2012

Impatiens nataliae Hook. f., Bull. Misc. Inform. Kew 294. 1910

Distribution: Karnataka (Udupi); endemic.

Impatiens scabriuscula B.Heyne ex Wall., Fl. Ind. 2:464.1824; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

var. **scabriuscula**

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu), Kerala, Maharashtra, Tamil Nadu; endemic.

var. **alba** Bhaskar, Taxon. Monogr. Impatiens W. Ghats 212. 2012

Distribution: Karnataka (Chikkamagaluru); endemic.

var. **rosea** Bhaskar, Taxon. Monogr. Impatiens W. Ghats 213. 2012

Distribution: Karnataka (Chikkamagaluru); endemic.

Impatiens stocksii Hook.f. & Thomson, J. Linn. Soc. Bot. 4: 156. 1860; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu), Kerala; endemic.

Impatiens talbotii Hook.f., Rec. Bot. Surv. India 4: 47. 1906; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

Distribution: Karnataka (Belagavi, Shivamogga, Uttara Kannada); endemic.

Impatiens tenella B.Heyne ex Wight & Arn., Prodr. Fl. Ind. Orient. 140 1834; B.D.Sharma et al., Fl. Karnataka Anal. 39. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Impatiens trichocarpa Hook.f., Hooker's Icon. Pl. 30: t. 2914 1910; Bhaskar, Taxon. Monogr. Impatiens W. Ghats 218. 2012

Distribution: Karnataka (Kodagu), Tamil Nadu; endemic.

Impatiens wolfgangii Lali J. & V. Bhaskar, Phytotaxa 313 (3): 281.2017

Distribution: Karnataka (Chikkamagaluru: Bababudangiri); endemic.

Basellaceae

Basella L.

Basella alba L., Sp. Pl. 272.1753

Basella rubra L., Sp. Pl. 272.1753; B.D.Sharma et al., Fl. Karnataka Anal. 233. 1984.

Distribution: Karnataka (Cultivated throughout), throughout India; pantropical.; native to tropical Asia.

Local/Common/Vernacular names: *Buffalo spinach, Ceylon spinach Creeping spinach, Climbing spinach, Malabar spinach, Pui, Red vine spinach, Vine spinach* (English); *Baai basle, Basale soppu* (Kannada)

Begoniaceae

Begonia L.

Begonia albococcinea Hook., Bot. Mag. t. 4172. 1845; C.J.Saldanha, Fl. Karnataka 1: 288. 1984

Begonia grahamiana Wight, Icon. Pl. Ind. Orient. 4: t. 1811. 1848

Distribution: Karnataka (Dakshina Kannada), Kerala; endemic.

Begonia concanensis A.DC., Ann. Sci. Nat. ser. 4, 11: 126. 1859; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Maharashtra; endemic.

Begonia crenata Dryand., Trans. Linn. Soc. London 1: 164, t.14. 1791; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984. **Plate 20A**

Begonia canarana Miq., Ann. Mus. Bot. Lugduno-Batavi 3: 18 1853; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Begonia minima Bedd., Madras J. Lit. Sci. III, 1: 48. 1864

Begonia aliciae C.E.C.Fisch., Bull. Misc. Inform. 1939: 247. 1939

Distribution: Karnataka (Belagavi, Dakshina Kannada, Mysuru, Shivamogga, Udupi Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Begonia hirtella Link, Enum. Hort. Berol. Alt. 2: 296.1822

Distribution: Karnataka (cultivated), Kerala, also cultivated elsewhere; native to Brazil.

Begonia integrifolia Dalzell in Hook.'s J. Bot. Kew Gard. Misc. 3: 230. 1851; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Distribution: Karnataka (Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Begonia malabarica Lam., Encycl. 1: 393. 1785; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Diploclinium lindleyanum Wight, Icon. Pl. Ind. Orient. 5: t. 1817. 1852

Plate 4F

Begonia fallax A.DC., Prodr. 15(1): 329.1864

Begonia malabarica var. *rheedii* A.DC., Prodr. 15(1): 392.1864

Begonia malabarica var. *hydropila* C.B.Clarke, Fl. Brit. India 2: 655. 1879

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Anekivi hvu* (Kannada)

Begonia picta Sm., Exot. Bot. 2: 81. t. 101. 1804; C.J.Saldanha, Fl. Karnataka 1: 289. 1984

Distribution: Karnataka (Chikkamagaluru), Andhra Pradesh, Arunachal Pradesh, Bihar, Himachal Pradesh, Jammu & Kashmir, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Mizoram, Nagaland, Uttar Pradesh, Uttaranchal; China, Myanmar, Nepal.

Begonia subpeltata Wight, Icon. Pl. Ind. Orient. 5: t. 1812 1852; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Diploclinium arnottianum Wight, Icon. Pl. Ind. Orient. 5: t. 1815.1852

Diploclinium cordifolium Wight, Icon. Pl. Ind. Orient. 5: t. 1816.1852

Reichenheimia subpeltata (Wight) Klotzsch, Abh. Königl. Akad. Wiss. Berlin 1855: 175.1855

Begonia cordifolia (Wight) Thwaites, Enum. Pl. Zeyl. 129.1859

Begonia arnottiana (Wight) A.DC., Prodr. 15(1): 322.1864

Distribution: Karnataka (Chikkamagaluru), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Begonia trichocarpa Dalzell in Hook.'s J. Bot. Kew Gard. Misc. 3: 230. 1851; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Plate 20B

Distribution: Karnataka (Belagavi, Chikkamagaluru), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Bignoniaceae

Adenocalymma Mart. ex Meissin.

Adenocalymma comosum (Cham.) DC., Prodr. 9: 201.1845

Bignonia comosa Cham., Linnaea 7: 693. 1832

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to S America

Local/Common/Vernacular name: *Yellow trumpet vine* (English)

Amphilophium Kunth

Amphilophium buccinatorium (DC.) L.G.Lohmann, Ann. Missouri Bot. Gard. 2010

Bignonia cherere Lindl., Edwards's Bot. Reg. 15: t. 1301.1830

Pithecoctenium buccinatorium DC., Prodr. 9: 195.1845

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Mexico.

Local/Common/Vernacular name: *Mexican blood trumpet* (English)

Anemopaegma C.Mart. ex Meiss.**Anemopaegma chamberlainii** (Sims) Bureau & K.Schum., Fl. Bras. 8(2): 128.1896*Bignonia chamberlainii* Sims, Bot. Mag. 47: t. 2148 1820; Haines, Bot. Bihar Orissa 2: 653. 1922.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Brazil.

Local/Common/Vernacular name: *Yellow trumpet vine* (English)**Bignonia** L.**Bignonia aequinoctialis** L., Sp. Pl. 623. 1753*Bignonia incarnata* Aubl., Hist. Pl. Guiane 2: 645. 1775; T. Cooke, Fl. Bombay 1: 335.1902.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to New Guinea.

Bignonia binata Thunb., Pl. Bras. 3: 35. 1821*Bignonia noterophila* Mart. ex DC., Prodr. 9: 148.1845; Haines, Bot. Bihar Orissa 2: 652. 1922.*Bignonia purpurea* Lodd. ex Sweet, Bot. Mag. 95: t. 5800. 1869*Clytostoma noterophilum* (Mart. ex DC.) Bureau & K.Schum., Fl. Bras. 8(2): 153. 1896

Distribution: Karnataka (cultivated), cultivated in gardens India; native to Brazil.

Bignonia callistegioides Cham., Linnaea 7: 712. 1832; Haines, Bot. Bihar Orissa 2: 652. 1922.*Bignonia speciosa* Graham, Bot. Mag. 68: t. 3888 .1841*Cuspidaria callistegioides* (Cham.) DC., Prodr. 9:178. 1845*Clytostoma callistegioides* (Cham.) Bureau, Adansonia 8: 353.1868

Distribution: Karnataka (cultivated), cultivated in gardens in India; native to tropical America.

Bignonia diversifolia Kunth, Nov. Gen. Sp. Pl. 3: 133.1819*Cydistia diversifolia* (Kunth) Miers, Proc. Roy. Hort. Soc. London 3: 192.1863*Pleonotoma diversifolia* (Kunth) Bureau & K.Schum., Fl. Bras. 8(2): 274.1897

Distribution: Karnataka (cultivated), cultivated in gardens in India; native to tropical America.

Bignonia magnifica W. Bull, Gard. Chron. 2: 73. 1879; Haines, Bot. Bihar Orissa 2: 652. 1922.*Arrabidaea magnifica* (W.Bull) Sprague ex Steenis, Recueil Trav. Bot. Néerl. 24: 830. 1927*Saritaea magnifica* (W.Bull) Dugand, Caldasia 3: 263. 1945

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to United States.

Local/Common/Vernacular names: *funnel vine, Glow vine, Purple* (English)**Campsis** Lour.**Campsis grandiflora** (Thunb.) K. Schum. in Engl. & Prantl, Nat. Pflanzenfam. 4(3b): 230. 1894*Bignonia grandiflora* Thunb., Nova Acta Soc. Sci. Upsal. 4: 39 1783 & Fl. Jap. 253. 1784*Bignonia chinensis* Lam., Encycl. 1: 423. 1785*Campsis adrepens* Lour., Fl. Cochinch. 377. 1790*Tecoma grandiflora* Loisel., Herb. Gén. Amateur 5: t. 286. 1821*Tecoma chinensis* (Lam.) K. Koch, Dendrologie 2(1): 307. 1872*Gelsemium grandiflorum* (Thunb.) Kuntze, Revis. Gen. Pl. 2: 480.1891*Campsis chinensis* (Lam.) Voss, Vilm. Blumengärtn. ed. 3 1: 80.1 1894

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; China, Japan, Pakistan, Vietnam.

Campsis radicans (L.) Seem., J. Bot. 5: 372.1867; T. Cooke, Fl. Bombay 1: 335.1902.*Bignonia radicans* L., Sp. Pl. 624. 1753*Tecoma radicans* (L.) Juss., Gen. Pl. 139. 1789

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to N America.

Colea Bojer ex Meiss.

Colea colei (Bojer ex Hook.) M.L.Green, Stand.-Sp. Nom. Conserv. 55. 1926

Bignonia colei Bojer ex Hook., Bot. Mag. 55: t. 2817. 1828

Colea mauritiana Bojer ex DC., Prodr. 9: 241. 1845; T. Cooke, Fl. Bombay 1: 336. 1902.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Madagascar.

Crescentia L.

Crescentia cujete L., Sp. Pl. 2: 626. 1753; T. Cooke, Fl. Bombay 1: 336. 1902.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to tropical America.

Local/Common/Vernacular name: *The Calbash tree* (English)

Dolichandra Cham.

Dolichandra unguis-cati (L.) L.G.Lohmann, Nuevo Cat. Fl. Vasc. Venezuela 273. 2008

Bignonia unguis-cati L., Sp. Pl. 623. 1753

Bignonia gracilis Lodd., Bot. Cab. 18: t. 1705. 1831

Doxantha unguis-cati (L.) Miers, Proc. Roy. Hort. Soc. London 3: 189. 1863

Distribution: Karnataka (cultivated in gardens), cultivated throughout India; native to C & S America and the Caribbean

Local/Common/Vernacular names: *Cats claw creeper, Cat's claw trumpet, Funnel creeper* (English)

Dolichandrone Seem. Nom. cons.

Dolichandrone arcuata (Wight) C.B.Clarke in Hook., Fl. Brit. India 4: 380. 1884

Spathodea arcuata Wight, Icon. Pl. Ind. Orient. t. 1340. 1848

Spathodea crispaa Dalzell & A.Gibson, Bombay Fl. 160. 1868

Distribution: Karnataka (Belagavi), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Dolichandrone atrovirens (Roth) K.Schum. in Engl. & Prantl, Nat. Pflanzenfam. 4(3a): 240. 1894

Bignonia crispa Buch.-Ham. in Roxb., Fl. Ind. 3: 103. 1832

Spathodea crispa G.Don, Gen. Hist. 4: 222. 1838

Dolichandron crispa Seem., J. Bot. 8: 381. 1870; C.B.Clarke in Hook.f., Fl. Brit. India 4: 379. 1884

Distribution: Karnataka (Belagavi, Chitradurga, Davanagere, Kodagu, Mysuru, Tumakuru), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Oodi mara, Seerudure, Udi* (Kannada)

Dolichandrone falcata (Wall ex DC.) Seem., J. Bot. 8: 380. 1870; C.B.Clarke in Hook.f., Fl. Brit. India 4: 380. 1884

Bignonia spathacea Roxb., Pl. Coromandel 2: 24, t. 144. 1799

Spathodea falcata Wall. ex DC., Prodr. 9: 206. 1845

Dolicandrone lawii Seem., J. Bot. 8: 380. 1870; C.B.Clarke in Hook.f., Fl. Brit. India 4: 380. 1884

Dolichandrone falcata var. *lawii* Sprague, Kew Bull. 1919: 309. 1919

Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Kalaburgi, Uttar Kannada, Yadgir), Bihar, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Uttar Pradesh; endemic.

Local/Common/Vernacular names: *Godmurki, Oodi mara* (Kannada)

Dolichandrone serrulata (Wall. ex DC.) Seem., J. Bot. 8: 383. 1870

Bignonia serrulata Wall. ex DC., Biblioth. Universelle Genève 17: 124. 1838

Stereospermum serratum DC., Biblioth. Universelle Genève n.s., 17: 124. 1838

Spathodea serrulata (Wall. ex DC.) DC., Prodr. 9: 206. 1845

Distribution: Karnataka (), Tamil Nadu: Myanmar, Thailand, Vietnam.

Dolichandrone spathacea (L.f.) Seem., J. Bot. 1: 226. 1863

Bignonia spathacea L.f., Suppl. Pl. 283. 1782

Bignonia longissima Lour., Fl. Cochinch. 380. 1790

Spathodea longiflora P. Beau., Fl. Oware 1: 47 1805

Spathodea rheedei Spreng., Syst. Veg. 2: 835. 1825

Spathodea loureiriana DC., Prodr. 9: 209. 1845

Spathodea diepenhorstei Miq., Fl. Ned. Ind., Eerste Bijv. 754. 1861

Dolichandrone rheedei (Spreng.) Seem., J. Bot. 8: 380. 1870; Clarke in Hook. f., Fl. Brit. India 4: 379. 1884.

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Odisha, Tamil Nadu, West Bengal; S & SE Asia.

Fernandoa Welw. ex Seem.

Fernandia Baill.

Haplophragma Dop

Fernandoa adenophylla (Wall. ex G.Don) Steenis, Blumea 23: 135 1976

Bignonia adenophylla Wall. ex G.Don, Gen. Hist. 4: 22. 1837

Spathodea adenophylla A.DC., Prodr. 9: 206. 1845

Heterophragma adenophyllum (Wall. ex G.Don) Seem. ex Benth. & Hook.f., Gen. Pl. 2: 1047 1876; C.B.Clarke in Hook.f., Fl. Brit. India 4: 381. 1884.

Distribution: Karnataka (cultivated in gardens), Andaman & Nicobar Islands, Assam, Mizoram, planted elsewhere, Bangladesh, Myanmar.

Handroanthus Mattos

Handroanthus impetiginosus (Mart. ex DC.) Mattos, Loefgrenia 50: 2. 1970

Tecoma impetiginosa Mart. ex DC., Prodr. 9: 218. 1845

Tabebuia avellaneda Lorentz ex Griseb., Abh. Königl. Ges. Wiss. Göttingen 24: 258. 1879

Gelsemium avellaneda (Lorentz ex Griseb.) Kuntze, Revis. Gen. Pl. 3: 245. 1898

Tecoma avellaneda (Lorentz ex Griseb.) Speg., Cat. Descr. Maderas 379. 1910

Tabebuia impetiginosa (Mart. ex DC.) Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 11: 176. 1936

Handroanthus avellaneda (Lorentz ex Griseb.) Mattos, Loefgrenia 50: 4. 1970

Distribution: Karnataka (cultivated in Bengaluru, Belagavi, Mysuru), cultivated in gardens in India; native to C & S America.

Local/Common/Vernacular names: *Pink Ipe, Pink Lapacho* (English)

Heterophragma DC.

Heterophragma quadriloculare (Roxb.) K.Schum., Nat. Pflanzenfam. 4(3b): 243 1895

Bignonia quadriocularis Roxb., Pl. Coromandel 2: 24. t. 145. 1799

Spathodea roxburghii Spreng., Syst. Veg. 2: 835. 1825

Heterophragma roxburghii DC., Prod. 9: 210. 1845; C.B.Clarke in Hook.F., Fl. Brit. India 4: 381. 1884

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Adavi nugge, Baara nugge, Baechaadi, Genasang, Genasangu* (Kannada)

Jacaranda Juss.

Jacaranda acutifolia Bonpl., Pl. Aequinoct. 1: 59. 1808

Distribution: Karnataka (cultivated in Belagavi), native to Brazil.

Jacaranda mimosifolia D.Don, Bot. Reg. 8: t. 631. 1822; Haines, Bot. Bihar Orissa 2: 652. 1922.

Distribution: Karnataka (cultivated throughout), cultivated throughout; native to Brazil, cultivated in tropics.

Kigelia DC.

Kigelia africana (Lam.) Benth., Niger Fl. 463. 1849

Bignonia africana Lam., Encycl. 1: 424. 1785

Crescentia pinnata Jacq., Collectanea 3: 203. 1789

Kigelia pinnata (Jacq.) DC., Prodr. 9: 247. 1845; T. Cooke, Fl. Bombay 1: 336. 1902.

Tripinna africana (Spreng.) Voigt, Hort. Suburb. Calcutt. 479. 1845

Distribution: Karnataka (cultivated throughout plains), cultivated throughout plains of India; native to tropical Africa.

Local/Common/Vernacular names: *Cucumber tree, Sausage tree* (English); *Aane thordu, Mara sauthe* (Kannada)

Macfadyena A.DC.

Macfadyena unguis-cati (L.) A.H.Gentry, Brittonia 25: 236. 1973

Distribution: Karnataka (cultivated in gardens), cultivated in India; native to C & S America, West Indies.

Markhamia Seem. ex Baillon

Markhamia lutea (Benth.) K.Schum., Nat. Pflanzenfam. 4(3b): 242. 1895

Spathodea lutea Benth., Niger Fl. 461. 1849

Muenteria lutea (Benth.) Seem., J. Bot. 8: 211. 1870

Dolichandrone lutea (Benth.) Benth. ex B.D.Jacks., Index Kew. 1: 785. 1893

Dolichandrone hildebrandtii Baker, Bull. Misc. Inform. Kew 1894: 31. 1894

Dolichandrone platycalyx Baker, Bull. Misc. Inform. Kew 1894: 30. 1894

Markhamia hildebrandtii (Baker) Sprague in Hooker's Icon. Pl. 28: t. 2800. 1905

Markhamia platycalyx (Baker) Sprague, Hooker's Icon. Pl. 28: t. 2800. 1905

Distribution: Karnataka (cultivated in gardens of Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), cultivated in gardens; native to Africa.

Local/Common/Vernacular names: *Nile trumpet, Nile tulip, Siala tree* (English)

Markhamia stipulata (Wall.) Seem., J. Bot. 1: 226. 1863

Bignonia stipulata (Wall.) Roxb., Fl. Ind. ed.2, 3: 108. 1832

Spathodea stipulata Wall., Pl. Asiat. Rar. 3: 20. 1832

Dolichandrone stipulata (Wall.) Benth. & Hook.f., Gen. Pl. 2: 1046. 1876; C.B.Clarke in Hook.f., Fl. Brit. India 4: 379. 1884.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to Cambodia, China, Laos, Myanmar, Thailand, Vietnam.

Millingtonia L. f.

Millingtonia hortensis L. f., Suppl. Pl. 291. 1782; C.B.Clarke in Hook.f., Fl. Brit. India 4: 377. 1884.

Bignonia azedarachta König & Sims, Ann. Bot. 1: 578. 1805

Bignonia suberosa Roxb., Pl. Coromande 13: 11. t. 214. 1811 & Fl. Ind. 3: 111. 1832

Bignonia hortensis (L.f.) Oken, Allg. Naturgesch. 3(2): 1009. 1841

Distribution: Karnataka (cultivated throughout), cultivated throughout drier parts of India; Bangladesh, Cambodia, China, Laos, Myanmar, Thailand, Vietnam; probably native to Indonesia & Malaysia.

Local/Common/Vernacular names: *Indian Cork tree* (English); *Akasha mallige, Biratu, Birte mara* (Kannada)

Pajanelia DC.

Pajanelia longifolia (Willd.) K.Schum., Nat. Pflanzenfam. 4(3b): 244. 1895

Bignonia indica Lour., Fl. Cochinch.: 460. 1790.

Bignonia longifolia Willd., Sp. Pl. 3: 306. 1800

Bignonia pajanelia Buch.-Ham., Trans. Linn. Soc. London 13: 515. 1822

Bignonia multijuga Wall., Pl. Asiat. Rar. 1: 81. 1830; T. Cooke, Fl. Bombay 1: 335. 1902.

Bignonia macrostachya Wall. ex G.Don, Gen. Hist. 4: 22. 1837

Pajanelia multijuga (Wall.) DC., Prodr. 9: 227. 1845

Pajanelia rheedii C.B.Clarke in Hook. f., Fl. Brit. India 4: 384. 1884

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Assam, Goa, Kerala, Meghalaya; Bangladesh, Myanmar.

Local/Common/Vernacular names: *Alangi, Doundi, Mokkudu* (Kannada)

Pandorea (Endl.) Spach.

Pandorea jasminoides (Lindl.) K.Schum., Nat. Pflanzenfam. 4(3b): 230. 1894; T.Cooke, Fl. Bombay 2: 335. 1902.

Tecoma jasminoides Lindl., Bot. Reg. 23: t. 2002. 1837

Gelsemium jasminoides (Lindl.) Kuntze, Revis. Gen. Pl. 2: 479. 1891

Distribution: Karnataka (Cultivated in gardens), cultivated in gardens in India; native to Australia.

Local/Common/Vernacular name: *Bower of Beauty* (English)

Pandorea pandorana (Andrews) Steenis, Bull. Jard. Bot. Buitenzorg III, 10: 198. 1928

Bignonia pandorana Andrews, Bot. Repos. 2: t. 86. 1799

Tecoma australis R.Br., Prodr. Fl. Nov. Holl. 471. 1810; Haines, Bot. Bihar Orissa 2: 652. 1922.

Tecoma pandorana (Andrews) Skeels, Bull. Bur. Pl. Industr. U.S.D.A. 282: 62. 1913

Campsis pandorana (Andrews) Steenis in Fl. Males. 4: xxi. 1948

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Australia.

Local/Common/Vernacular name: *Wonga Wonga Vine* (English)

Parmentiera DC.

Parmentiera cereifera Seem., Bot. Voy. Herald 182. 1854; T. Cooke, Fl. Bombay 1: 336. 1902.

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Panama.

Local/Common/Vernacular name: *The candle tree* (English)

Pyrostegia C.Persl.

Pyrostegia venusta (KerGawler) Miers, Proc. Roy. Hort. Soc. London 3: 188. 1863

Bignonia venusta Ker Gawl., Bot. Reg. 3: t. 249. 1801

Bignonia ignea Vell., Fl. Flumin. 244. 1829

Tecoma venusta (KerGawler) Lem., Hort. Universel 1. 1843

Pyrostegia ignea (Vell.) C.Presl, Bot. Bemerk. 93. 1844; T. Cooke, Fl. Bombay 1: 335. 1902.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Brazil.

Radermachera Zoll. & Moritzi

Radermachera xylocarpa (Roxb.) Roxb. ex K.Schum., Nat. Pflanzenfam. 4(3b): 243. 1895

Bignonia xylocarpa Roxb., Fl. Ind. 3: 108. 1832

Tecoma xylocarpa (Roxb.) G. Don, Gen. Hist. 4: 225. 1838

Spathodea xylocarpa (Roxb.) T. Anderson ex Brandis., Forest Fl. N.W. India 349. 1874

Stereospermum xylocarpum (Roxb.) Benth. & Hook.f., Gen. Pl. 2: 1047. 1876; C.B.Clarke in Hook.f., Fl. Brit. India 4: 383. 1884.

Distribution: Karnataka (throughout), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Genasing, Konana kombu mara* (Kannada); *Kharsingi* (Kunhi)

Spathodea Pal.

Spathodea campanulata P.Beauv., Fl. Oware 1: 47 1805; T. Cooke, Fl. Bombay 1: 335.1902.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical Africa.

Local/Common/Vernacular names: *Tulip tree* (English); *Neeru kayi mara, Uchche kayi mara* (Kannada)

Stereospermum Cham.

Stereospermum chelonoides (L.f.) DC., Biblioth. Universelle Genève n.s., 17: 124. 1838

Bignonia chelonoides L. f., Suppl. Pl. 282.1782

Bignonia suaveolens Roxb., Fl. Ind. 3: 104.1832

Stereospermum suaveolens DC., Biblioth. Universelle Genève n.s., 17: 124 1838

Tecoma suaveolens (Roxb.) G.Don, Gen. Hist. 4: 224. 1838

Bignonia gratissima K.D.Koenig ex DC., Prodr. 9: 211.1845

Heterophragma chelonoides (L.f.) Dalzell & A.Gibson, Bombay Fl. 160. 1861

Heterophragma suaveolens (Roxb.) Dalzell & A. Gibson, Bombay Fl. 161. 1861

Spathodea suaveolens (Roxb.) Benth. & Hook.f., Gen. Pl. 2: 1047. 1876

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Davanagere), throughout India up to foot hills of Himalaya; Bangladesh, Myanmar, Sri Lanka.

Local/Common/Vernacular name: *Paadri* (Kannada)

Stereospermum colais (Buch.-Ham. ex Dillwyn) Mabb., Taxon 27: 553. 1978

Bignonia colais Buch.-Ham. ex Dillwyn, Rev. Hort. 6(26): 28. 1839

Stereospermum chelonoides var. *angustifolia* Haines, Descrip. List Trees S. Circle Cent. Prov. 169. 1916

Stereospermum angustifolia (Haines) Kew Bull.1922: 121. 1921

Stereospermum personatum var. *angustifolia* (Haines) Santisuk, Kew Bull. 28: 181. 1973

Stereospermum colais var. *angustifolium* (Haines) Bennet & Raizada, Indian J. Forest. 4(1): 67. 1981

Stereospermum colais var. *shendurunii* Sasidh., Sujanapal & Binoy, Rheedia 16: 37. 2006

Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Tumakuru, Uttara Kannada), S to E India; S & SE Asia.

Local/Common/Vernacular names: *Kaadu nugge, Kempudale, Paadari Mukarti* (Kannada); *Kudal* (Kunhi)

Stereospermum neuranthum Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 42(2): 91. 1873; C.B.Clarke in Hook.f., Fl. Brit. India 4: 383. 1884.

Stereospermum wallichii C.B.Clarke in Hook. f., Fl. Brit. India 4: 383.1884

Stereospermum grandiflorum Cubitt & W.W.Sm., Rec. Bot. Surv. India 4: 280. 1911

Radermachera wallichii (C.B.Clarke) Chatterjee, Bull. Bot. Soc. Bengal 2: 72. 1948

Distribution: Karnataka (Manipur), Cambodia, China, Laos, Myanmar, Thailand, Vietnam.

Stereospermum tetragonum DC., Biblioth. Universelle Genève n.s., 17: 124 1838

Dipterosperma personatum Hassk., Flora 25(2 Beibl.): 28.1842

Bignonia caudata DC., Prodr. 9: 166. 1845

Bignonia tetragona Wall. ex DC., Prodr. 9: 211. 1845

Stereospermum caudatum (DC.) Miq., Ann. Mus. Bot. Lugduno-Batavum 1: 200. 1863

Stereospermum personatum (Hassk.) Chatterjee, Bull. Bot. Soc. Bengal 2: 70 1948

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Arunachal Pradesh ? Bihar, Jharkhand, Odisha;Myanmar, Sri Lanka.

Local/Common/Vernacular names: *Lingadare mara, Malali* (Kannada)

Tabebuia Gomes ex DC.

Tabebuia aurea (Silva Manso) Benth. & Hook.f. ex S.Moore, Trans. Linn. Soc. London, Bot. 4: 423 1895

Bignonia aurea Silva Manso, Enum. Subst. Braz. 40 1836

Tecoma aurea (Silva Manso) DC., Prodr. 9: 222.1845

Tecoma argentea Bureau & K.Schum., Fl. Bras. 8(2): 332.1897

Tabebuia argentea (Bureau & K.Schum.) Britton, Sc. Surv. Porto Rico & Virgin Islands 6: 197.1925

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical S America.

Local/Common/Vernacular name: *Caribbean Trumpet Tree* (English)

Tabebuia heterophylla (DC.) Britton, Ann. Missouri Bot. Gard. 2: 48.1915

Raputia heterophylla DC., Mém. Mus. Hist. Nat. 9: 153.1822

Pilocarpus heterophyllum (DC.) A.Gray, Pl. Wright. 1: 170.1860

Tabebuia pentaphylla Hemsl., Biol. Cent.-Amer., Bot. 2: 495.1882

Tabebuia pallida subsp. *heterophylla* (DC.) Stehlé, Caribbean Forest. 6(Suppl.): 330.1945

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical America.

Local/Common/Vernacular name: *Pink trumpet tree* (English)

Tabebuia pallida (Lindl.) Miers, Proc. Roy. Hort. Soc. London 3: 199.1863

Bignonia pallida Lindl., Bot. Reg. 21: t. 965.1827

Tabebuia heterophylla subsp. *pallida* (Lindl.) Stehlé, Bull. Soc. Bot. France 93: 32 .1946

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical America.

Local/Common/Vernacular name: *White trumpet tree* (English)

Tabebuia rosea (Bertol.) Bertero ex A.DC., Prodr. 9: 215.1845

Tecoma rosea Bertol., Novi Comment. Acad. Sci. Inst. Bononiensis 4: 425.1840

Sparattosperma roseum (Bertol.) Miers, Proc. Roy. Hort. Soc. London 3: 199.1863

Couralia rosea (Bertol.) Donn.Sm., Bot. Gaz. 20: 9 1895

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical America.

Local/Common/Vernacular names: *Pink trumpet tree, Rosy trumpet tree* (English)

Tecoma Juss.

Tecoma capensis (Thunb.) Lindl., Bot. Reg. 13: t. 1117. 1828

Bignonia capensis Thunb., Prodr. Pl. Cap. 105. 1794

Tecomaria capensis (Thunb.) Spach, Hist. Nat. Veg. 9: 137. 1840; T. Cooke, Fl. Bombay 1: 335.1902.

Gelsemium capense (Lindl.) Kuntze, Revis. Gen. Pl. 2: 479. 1891

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to S Africa.

Local/Common/Vernacular name: *Cape honeysuckle* (English)

Tecoma castanifolia (D.Don) Melch., Ber. Deutsch. Bot. Ges. 59: 26.1941; K.G.Bhat, Fl. S. Kanara 807. 2014.

Stenolobium castanifolium D.Don, Edinburgh Philos. J. 9: 264.1823

Bignonia castaneifolia (D.Don) DC., Prodr. 9: 145.1845

Tecoma gaudichaudii DC., Prodr. 9: 223.1845

Stenolobium stans var. *castanifolium* (D.Don) Seem., J. Bot. 1: 88.1863

Gelsemium gaudichaudii (DC.) Kuntze, Revis. Gen. Pl. 2: 479.1891

Distribution: Karnataka (cultivated in gardens), cultivated in gardens; native to tropical Africa.

Local/Common/Vernacular name: *Chestnut leaf trumpet bush* (English)

Tecoma stans (L.) Juss. ex Kunth, Nov. Gen. Sp. Pl. 3: 144.1819

Stenolobium stans (L.) Seem., J. Bot. 1: 88. 1863

Gelsemium stans (L.) Kuntze, Revis. Gen. Pl. 2: 479.1891

Distribution: Karnataka (cultivated in gardens & road sides), cultivated throughout India; native to Americas.

Local/Common/Vernacular names: *Yellow bells, Yellow trumpet bush* (English); *Gante huvu* (Kannada)

Bixaceae

Bixa L.

Bixa orellana L., Sp. Pl. 512. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 12. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kodagu, Shivamogga, Uttara Kannada), throughout India; native to America

Local/Common/Vernacular name: *Butter-seed Tree* (English); *Rangumale* (Kannada)

Cochlospermum Kunth

Cochlospermum religiosum (L.) Alston in Trimen, Handb. Fl. Ceylon 6 (Suppl.): 14. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 12. 1984.

Bombax religiosum L., Sp. Pl. 512. 1753

Cochlospermum gossypium DC., Prodr. 1: 527. 1824

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Tumakuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Bihar, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; SE Asia.

Local/Common/Vernacular name: *Golden Silk Cotton Tree* (English); *Arishina booruga* (Kannada)

Boraginaceae

Adelocaryum Brand

Adelocaryum coelestinum (Lindl.) Brand, Repert. Spec. Nov. Regni Veg. 13: 549. 1915; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Cynoglossum coelestinum Lindl., Edwards's Bot. Reg. 25: t. 36. 1839

Echinospermum coelestinum (Lindl.) Wight, Icones Pl. Ind. Orient. 4: t. 1394. 1848

Paracaryum coelestinum (Lindl.) C.B.Clarke in Hook. f., Fl. Brit. India 4: 160. 1883

Paracaryopsis coelestina (Lindl.) R.R.Mill, Edinburgh J. Bot. 48: 57. 1991

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Maharashtra; endemic.

Adelocaryum malabaricum (C.B.Clarke) Brand, Repert. Spec. Nov. Regni Veg. 13: 549. 1915 **Plate 20C**

Paracaryum malabaricum C.B.Clarke in Hook. f., Fl. Brit. India 4: 160. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Cynoglossum malabaricum (C.B.Clarke) Riedl, Oesterr. Bot. Z. 119: 71. 1971

Paracaryopsis malabarica (C.B.Clarke) R. R. Mill, Edinburgh J. Bot. 48: 59. 1991

Distribution: Karnataka (Chamarajanagar, Chikkamagalur, Dakshina Kannada, Udupi, Uttara Kannada), Kerala, Maharashtra; endemic.

Coldenia L.

Lobophyllum F.Muell

Coldenia procumbens L., Sp. Pl. 1: 125. 1753; C.B.Clarke in Hook. f., Fl. Brit. India 4: 144. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Cordia furcans I. M. Johnst., J. Arnold Arbor. 32: 5. 1951

Distribution: Karnataka (throughout), throughout India; pantropical.

Local/Common/Vernacular names: *Hamsa padi*, *Thripakshi gida* (Kannada)

Cordia L.

Cordia alba (Jacq.) Roem. & Schult., Syst. Veg. 4: 466. 1819; Ramaswamy & Razi, Fl. Bangalore Dist. 481. 1973..

Varronia alba Jacq., Enum. Syst. Pl. 14. 1760

Calyptacordia alba (Jacq.) Britton, Sci. Surv. Porto Rico & Virgin Islands 6: 123. 1925

Gerascanthus albus (Jacq.) Borhidi, Acta Bot. Hung. 34: 403. 1988

Distribution: Karnataka (cultivated in Bengaluru), native to S America.

Cordia dichotoma G.Forst., Fl. Ins. Austr. 18. 1786; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Cordia obliqua Willd., Phytographia 4. 1794

Cordia indica Lam., Encycl. 7: 49. 1806

Cordia latifolia Roxb., Fl. Ind. 2: 330. 1824

Cordia wallichii G.Don, Gen. Hist. 4: 379. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Cordia griffithii C.B.Clarke in Hook. f., Fl. Brit. India 4: 139. 1883

Cordia obliqua var. *wallichii* (G.Don) C.B.Clarke in Hook. f., Fl. Brit. India 4: 137. 1883

Lithocardium griffithii Kuntze, Revis. Gen. Pl. 2: 977. 1891

Lithocardium obliquum Kuntze, Revis. Gen. Pl. 2: 977. 1891

Cordia lowiana Brandis, Indian Trees 479. 1906

Gerascanthus dichotomus (G.Forst.) Borhidi, Acta Bot. Hung. 34: 404. 1988

Gerascanthus griffithii (C.B.Clarke) Borhidi, Acta Bot. Hung. 34: 405. 1988

Distribution: Karnataka (throughout), Andhra Pradesh, Assam, Goa, Gujarat, Jharkhand, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Punjab & Tamil Nadu; Africa, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Challe hannu*, *Challe mara*, *Chelle kaayi*, *Gonne hannu mara*, *Mannadke*, *Solle mara* (Kannada); *Bhokar* (Kunhi); *Challangayi* (Tulu)

Cordia domestica Roth, Nov. Pl. Sp. 123. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Cordia fulvosa Wight, Icon. Pl. Ind. Orient. 4: t. 1380. 1848; C.B.Clarke in Hook.f., Fl. Brit. India 4: 140. 1883.

Cordia fulvosa var. *evolutior* C.B.Clarke in Hook. f., Fl. Brit. India 4: 140. 1883

Cordia myxa var. *domestica* (Roth) C.B.Clarke in Hook. f., Fl. Brit. India 4: 137. 1883

Lithocardium fulvosum Kuntze, Revis. Gen. Pl. 2: 977. 1891

Cordia evolutior (C.B.Clarke) Gamble, Fl. Madras 888. 1923; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Gerascanthus fulvosus (Wight) Borhidi, Acta Bot. Hung. 34: 404. 1988

Cordia domestica var. *evolutior* (C.B.Clarke) M.R.Almeida, Fl. Maharashtra 3A: 295. 2001

Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Hassan, Mysuru, Uttara Kannada), peninsular India; endemic.

Cordia macleodii Hook.f. & Thomson, J. Proc. Linn. Soc., Bot. 2: 128. 1858; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Hemigymnia macleodii Griff., Calcutta J. Nat. Hist. 3: 363. 1843

Lithocardium macleodii Kuntze, Revis. Gen. Pl. 2: 977. 1891

Gerascanthus macleodii (Hook.f. & Thomson) Borhidi, Acta Bot. Hung. 34: 405. 1988

Distribution: Karnataka (Belagavi, Chikkamagaluru, Mysuru, Uttara Kannada), throughout except NE India; endemic.

Local/Common/Vernacular name: *Bili celle* (Kannada)

Cordia monoica Roxb., Pl. Coromande 11: 43. 1796; C.B.Clarke in Hook. f., Fl. Brit. India 4: 137. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Lithocardium monoicum Kuntze, Revis. Gen. Pl. 2: 977. 1891

Gerascanthus monoicus (Roxb.) Borhidi, Acta Bot. Hung. 34: 406. 1988

Distribution: Karnataka (Chitradurga, Vijayapura), Peninsular India to Jharkhand; Arabia, Sri Lanka, tropical E Africa.

Cordia myxa L., Sp. Pl. 199. 1753; C.B.Clarke in Hook.f., Fl. Brit. India 4: 136. 1883.

Cordia sebestena Forssk., Fl. Aegypt.-Arab. 63. 1775

Sebestena officinalis Gaertn., Fruct. Sem. Pl. 1: 364. 1788

Cordia officinalis Lam., Tabl. Encycl. 2: 420. 1794

Ehretia glabra Roth ex Roem. & Schult., Syst. Veg. 4: 532. 1819

Cordia paniculata Roth, Nov. Pl. Sp. 125. 1821

Bourreria glabra G.Don, Gen. Hist. 4: 390. 1837

Sebestena myxa (L.) J. Presl, Wsobecný Rostl. 2: 1102. 1846

Lithocardium myxa Kuntze, Revis. Gen. Pl. 2: 977. 1891

Gerascanthus myxus (L.) Borhidi, Acta Bot. Hung. 34: 406. 1988

Distribution: Karnataka (throughout, both wild & cultivated), Sri Lanka, SE & W Asia.

Cordia nevillii Alston in Trimen, Handb. Fl. Ceylon 6(Suppl.): 199. 1931

Distribution: Karnataka (Chamrajanagara), Arabia, Sri Lanka, tropical Africa,

Cordia perrottetii Wight, Icon. Pl. Ind. Orient. 4: t. 1381. 1848; C.B.Clarke in Hook.f., Fl. Brit. India 4: 138. 1883.

Distribution: Karnataka (Ballari), Andhra Pradesh, Tamil Nadu; endemic.

Cordia sebestena L., Sp. Pl. 190. 1753.

Sebestena sebestena (L.) Britton, Fl. Miami 158. 1913

Distribution: Karnataka (cultivated in gardens), cultivated in gardens, common in Islands; native to West Indies.

Cordia sinensis Lam., Tabl. Encycl. 1: 423. 1792

Cornus gharaf Forssk., Fl. Aegypt.-Arab. 95. 1775

Cordia rothii Roem. & Schult., Syst. Veg. 4: 798. 1819; C.B.Clarke in Hook.f., Fl. Brit. India 4: 138. 1883.

Cordia reticulata Roth, Nov. Pl. Sp. 124. 1821

Cordia angustifolia Roxb., Fl. Ind. 2: 338. 1824

Cordia cuneata B.Heyne ex A.DC., Prodr. 9: 480. 1845

Cordia subopposita A.DC., Prodr. 9: 480. 1845

Cordia gharaf Ehrenb. ex Asch., Sitzungsber. Ges. Naturf. Freunde Berlin 46. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Lithocardium reticulatum Kuntze, Revis. Gen. Pl. 2: 977. 1891

Lithocardium rothii Kuntze, Revis. Gen. Pl. 2: 977. 1891

Gerascanthus sinensis (Lam.) Borhidi, Acta Bot. Hung. 34: 406. 1988

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Dharwad, Uttara Kannada), Peninsula to NW India; Africa, Middle East, Pakistan.

Cynoglossum L.

Cynoglossum ritchiei C.B.Clarke in Hook. f., Fl. Brit. India 4: 158. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Distribution: Karnataka (Belagavi); endemic.

Cynoglossum wallichii G.Don, Gen. Hist. 4: 354. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Cynoglossum glochidiatum Wall. ex Benth. In Royle, Ill. Bot. Himal. Mts. 306. 1836

Cynoglossum denticulatum A.DC., Prodr. 10: 150. 1846

Cynoglossum edgeworthii A.DC., Prodr. 10: 150. 1846

Cynoglossum denticulatum var. *zeylanicum* C.B.Clarke in Hook. f., Fl. Brit. India 4: 157. 1883

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Uttara Kannada), throughout India, tropical Africa & Asia.

Cynoglossum zeylanicum (Sw. ex Lehm.) Thunb. ex Brand, Repert. Spec. Nov. Regni Veg. 13(no. 381-382): 546. 1915 & Pflanzenr. IV, 252: 134 1921; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.
Echinospermum zeylanicum (Sw. ex Lehm.) Lehm., Pl. Asperif. Nucif. 1: 116. 1818
Cynoglossum furcatum Wall. ex Roxb., Fl. Ind.ed.1, 2: 6. 1824; C.B.Clarke in Hook. f., Fl. Brit. India 4: 155. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.
Cynoglossum heynei G.Don, Gen. Hist. 4: 354. 1837
 Distribution: Karnataka (throughout), throughout India; E, S & SE Asia.
 Local/Common/Vernacular names: *Amrada soppu, Jigti* (Kunhi)

Ehretia P. Brown

Ehretia acuminata R. Br., Prodr. Fl. Nov. Holl. 497. 1810; C.B.Clarke I

n Hook.f., Fl. Brit. India 4: 141. 1883.

Ehretia serrata Roxb., Fl. Ind. 2: 340. 1824

Ehretia pyrifolia D.Don, Prodr. Fl. Nepal. 102. 1825

Ehretia ovalifolia Hassk., Cat. Hort. Bot. Bogor. 137. 1844

Ehretia serrata var. *pyrifolia* (D.Don) DC., Prodr. 9: 503. 1845

Cordia thrysiflora Siebold & Zucc., Fl. Jap. Fam. Nat. 2: 26. 1846

Ehretia thrysiflora (Siebold & Zucc.) Nakai, Trees Shrubs Japan 327. 1922

Ehretia acuminata var. *pyrifolia* (D.Don) I.M.Johnst., J. Arnold Arbor. 32: 23. 1951

Ehretia acuminata var. *serrata* (Roxb.) I.M.Johnst., J. Arnold Arbor. 32: 23. 1951

Distribution: Karnataka (cultivated?), cultivated in S India, Arunachal Pradesh, Assam, Himachal Pradesh, Jammu & Kashmir, Meghalaya, Punjab, Sikkim, Uttarakhand, Uttar Pradesh, West Bengal; Australia, Bhutan, China, Indonesia, Japan, Laos, Nepal, New Guinea, Vietnam.

Local/Common/Vernacular name: *Koda tree* (English)

Ehretia laevis Roxb., Pl. Coromandel. 1: 42, t.56. 1796; C.B.Clarke in Hook.f., Fl. Brit. India 4: 141. 1883.

Ehretia aspera Willd., Phytographia 4. 1794

Ehretia heynei Roem. & Schult., Syst. Veg. 4: 532. 1819

Ehretia punctata Roth ex Roem. & Schult., Syst. Veg. 4: 532. 1819

Ehretia floribunda Royle, Ill. Bot. Himal. Mts. 306. 1836

Ehretia pubescens Benth. in Royle, Ill. Bot. Himal. Mts. 306. 1836

Bourreria aspera G.Don, Gen. Hist. 4: 390. 1837

Bourreria dichotoma Rottler ex G.Don, Gen. Hist. 4: 390. 1837

Bourreria punctata G.Don, Gen. Hist. 4: 390. 1837

Ehretia ovalifolia Wight, Icon. Pl. Ind. Orient. 4: t. 1383. 1848.

Ehretia laevis var. *floribunda* (Benth.) Brandis, Forest Fl. N.W. India 340. 1874

Ehretia canarensis Miq. ex C.B.Clarke, Fl. Brit. India 8: 142. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Ehretia championii Wight & Gardner ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 142. 1883

Ehretia cutranga C.B.Clarke in Hook.f., Fl. Brit. India 4: 142. 1883

Ehretia laevis var. *aspera* (Willd.) C.B.Clarke in Hook.f., Fl. Brit. India 4: 142. 1883

Ehretia laevis var. *canarensis* Miq. ex C.B.Clarke in Hook. f., Fl. Brit. India 4: 142. 1883

Ehretia laevis var. *pubescens* (Benth.) C.B.Clarke in Hook. f., Fl. Brit. India 4: 142. 1883

Ehretia aspera var. *obtusifolia* (Hochst. ex DC.) Parmar, Fl. Rajasthan 2: 523. 1991

Ehretia indica (Dennst. ex Kostel.) M.R.Almeida & S.M.Almeida, J. Bombay Nat. Hist. Soc. 90: 424. 1994

Distribution: Karnataka (throughout), throughout India; Africa, Australia, tropical Asia.

Local/Common/Vernacular name: *Bagari* (Kannada); *Datrang* (Kunhi)

Ehretia microphylla Lam., Tabl. Encycl. 1: 425. 1792

Cordia retusa Vahl, Symb. Bot. 2: 42. 1791

Ehretia buxifolia Roxb., Pl. Coromandel 1: 42, t.57. 1796

Carmona heterophylla Cav., Icon. 5: 22. 1799

Carmona microphylla (Lam.) G.Don, Gen. Hist. 4: 391. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Cordia coromandeliana Retz. ex A.DC., Prodr. 9: 509. 1845

Ehretia buxifolia var. *microphylla* (Lam.) DC., Prodr. 9: 509. 1845

Ehretia coromandeliana Retz. ex A.DC., Prodr. 9: 509. 1845

Carmona retusa (Vahl) Masam., Trans. Nat. Hist. Soc. Taiwan 30: 61. 1940

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kolar, Mysuru), Peninsula India, Nagaland; SE Asia.

Local/Common/Vernacular names: *Enne budugi, Kuruvinge* (Kannada)

Ehretia wightiana Wall. ex G.Don, Gen. Hist. 4: 388. 1837

Ehretia lanceolata B.Heyne ex C.B.Clarke in Hook. f., Fl. Brit. India 4: 143. 1883

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Heliotropium L.

Heliotropium amplexicaule Vahl, Symb. Bot. 3:21.1794; R.K.Swamy et al., Rheedia 25(2):148. 2015.

Heliotropium anchusifolium Poir. In Lam., Encycl. Suppl. 3:23.1813

Cochranea anchusifolia (Poir.) Gurke in Engl. & Prantl, Nat. Planzenfam. 4 (3a):97.1894

Distribution: Karnataka (Bengaluru, Dharwad), Africa, Australia, S & SE Asia, tropical America.

Heliotropium arborescens L., Syst. Nat. ed. 10 2: 913 1759

Heliotropium peruvianum L., Sp. Pl. ed. 2 187 1762; Duthie, Fl Upp. Gang. Pl. 2: 94. 1911.

Distribution: Karnataka (cultivated in gardens), cultivated in Indian gardens; Native to Peru.

Heliotropium bracteatum R.Br., Prodr. Fl. Nov. Holl. 493.1810; Bhaskar & Kushlappa, Fl. Tumkur Dist. 79. 2013.

Heliotropium laxiflorum Roth, Nov. Pl. Sp. 102.1821

Heliotropium bracteatum var. *leptostachyum* Benth., Fl. Austral. 4: 399.1868

Distribution: Karnataka (Chitradurga, Hassan, Tumakuru), Kerala, Tamil Nadu; Australia, SE Asia.

Heliotropium cornutum I.M.Johnst., Contr. Gray Herb. Harward Univ. 92: 90. 1930; C.E.C.Fisch. in Gamble, Fl. Madras 3: 1330 1957.

Heliotropium cornutum var. *pusilla* C.B.Clarke in Hook. f., Fl. Brit. India 4: 369. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Distribution: Karnataka (Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), Maharashtra; endemic.

Heliotropium indicum L., Sp. Pl. 1: 130. 1753; C.B.Clarke in Hook. f., Fl. Brit. India 4: 152. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Tiaridium indicum (L.) Lehm., Pl. Asperif. Nucif. 1: 14.1818

Heliophytum indicum (L.) DC., Prodr. 9: 556.1845

Distribution: Karnataka (throughout), throughout hotter parts of India; pantropical.

Local/Common/Vernacular names: *Bhoorundi, Chelu balad gida, Chelu kondi gida, Chelu mani* (Kannada); *Amardaaru, Bhoorundi, Bhoorischivatali, Duraalabha* (Sanskrit)

Heliotropium keralense Sivarajan & Manilal, J. Indian Bot. Soc. 51: 348. 1973

Distribution: Karnataka (Dakshina Kannada), Andaman & Nicobar Islands, Kerala, Tamil Nadu; endemic.

Heliotropium marifolium J.König ex Retz., Observ. Bot. 2: 8 1781; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Heliotropium scabrum Lehm., Pl. Asperif. Nucif. 70.1818

Cynoglossum marifolium Roxb., Fl. Ind. ed.1, 2: 8 1824

Bothriospermum marifolium A.DC., Prodr. 10: 116. 1846

Heliotropium marifolium var. *wallichii* C.B.Clarke in Hook. f., Fl. Brit. India 4: 152. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Distribution: Karnataka (throughout), throughout hotter parts India; S & SE Asia.

Heliotropium ovalifolium Forssk., Fl. Aegypt.-Arab. 38. 1775; C.B.Clarke in Hook. f., Fl. Brit. India 4: 150. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Heliotropium coromandelianum J.König ex Retz., Observ. Bot. 2: 9. 1781

Heliotropium gracile R.Br., Prodr. Fl. Nov. Holl. 493. 1810

Heliotropium polystachyum Poir., Encycl. Suppl. 3: 23. 1813

Heliotropium tomentosum Poir., Encycl. Suppl. 3: 23. 1813

Heliotropium coromandelianum var. *ovalifolium* Lehm., Pl. Asperif. Nucif. 47 1818

Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Hassan, Kalaburgi, Kodagu, Raichur, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), throughout plains India; Paleotropical.

Local/Common/Vernacular name: *Hasthishundi* (Sanskrit)

Heliotropium pycnophyllum Phil., Fl. Atacam. 38.1860

Heliotropium brevifolium Wall., Fl. Ind., ed. 1820, 2: 2.1824

Distribution: Karnataka (cultivated in Mysuru), native to Chile.

Heliotropium ramosissimum (Lehm.) Sieber ex DC., Prodr. 9:536. 1845; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Heliotropium crispum Desf., Fl. Atlant. 1:151.1798; M. Sharma & N. Vaid. J. Econ. Tax. Bot. 19 : 258. 1995

Distribution: Karnataka (Raichur), Afghanistan, Arabia, Iraq, Jordan, Iran, Pakistan, tropical Africa.

Heliotropium rariflorum Stocks, Hooker's J. Bot. Kew Gard. Misc. 4: 174.1852; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Distribution: Karnataka (Chitradurga, Dharwad), W India; Afghanistan, Pakistan, E & S E Africa.

Heliotropium scabrum Retz., Observ. Bot. 2: 8. 1781.

Distribution: Karnataka (Bengaluru, Dharwad, Mysuru, Tumakuru), peninsular & NW India; Cambodia, China (Hainan), Indonesia, Pakistan, Sri Lanka.

Heliotropium strigosum Willd., Sp. Pl. 1(2): 743. 1798; C. B. Clarke in Hook. f., Fl. Brit. India 4: 151. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Heliotropium compactum D.Don, Prodr. Fl. Nepal. 101 1825

Lithospermum chinense Hook. & Arn., Bot. Beechey Voy. 202.1837

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Uttara Kannada), throughout plains of India extending to W Himalaya; Afghanistan, Africa, Australia, Bhutan, Cambodia, China, Laos, Myanmar, Nepal, Pakistan, Thailand, Vietnam.

Local/Common/Vernacular names: *Chaelu baalada gida, Shvetha rangi Thaudatti* (Kannada)

Heliotropium subulatum (Hochst. ex A.DC.) Vatke, Linnaea 43(5-6): 316.1882; M. Sharma & N. Vaid, J. Econ. Tax. Bot. 19 : 258. 1995.

Distribution: Karnataka (Chitradurga, Vijayapura), Punjab; Africa, Pakistan

Local/Common/Vernacular name: *Hasthi shundi* (Sanskrit)

Heliotropium supinum L., Sp. Pl. 130. 1753; C.B.Clarke in Hook. f., Fl. Brit. India 4: 149. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Lithospermum heliotropoides Forssk., Fl. Aegypt.-Arab. 39. 1775

Heliotropium malabaricum Retz., Observ. Bot. 4: 24. 1786

Piptoclaina malabarica G.Don, Gen. Hist. 4: 364. 1837

Piptoclaina supina G.Don, Gen. Hist. 4: 364. 1837

Heliotropium supinum var. *malabaricum* C.B.Clarke in Hook. f., Fl. Brit. India 4: 149. 1883 ('malabarica').

Distribution: Karnataka (throughout), throughout except Himalaya & NE India; W Asia, C Africa, S Europe.

Heliotropium zeylanicum (Burm.f.) Lam., Encycl. 3: 94. 1789; C.B.Clarke in Hook. f., Fl. Brit. India 4: 148. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 177. 1984.

Messerschmidia hispida Benth. in Royle, Ill. Bot. Himal. Mts. 306. 1836

Tournefortia edgeworthii A.DC., Prodr. 9: 529. 1845

Tournefortia royleana A.DC., Prodr. 9: 527. 1845

Tournefortia subulata Hochst. ex A.DC., Prodr. 9: 528. 1845

Tournefortia zeylanica (Burm.f.) Wight, Ill. Ind. Bot. 2: t. 170. 1850

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Tumakuru, Vijayapura), throughout plains India; SE Asia, tropical Africa & Australia.

Local/Common/Vernacular name: *Hasthi shundi* (Sanskrit)

Rotula Lour.

Rotula aquatica Lour., Fl. Cochinch. 1: 121. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 178. 1984.

Rhabdia lycioides auct. non Mart., 1827; C.B.Clarke in Hook.f., Fl. Brit. India 4: 145. 1883, p.p.

Ehretia viminea Wall., Numer. List 906. 1829

Carmona lycioides G.Don, Gen. Hist. 4: 391. 1837

Ehretia cuneata Wight, Icon. Pl. Ind. Orient. 4: t. 1385. 1848

Rhabdia viminea (Wall.) Dalziel & Hook., Icon. Pl. 9: t. 823. 1851

Carmona viminea (Wall.) G. Don; C.B.Clarke in Hook. f., Fl. Brit. India 4: 145. 1883

Ehretia aquatica (Lour.) Gottschling & Hilger, Bot. Jahrb. Syst. 123: 263. 2001

Ehretia lycioides (G.Don) Wall. ex B.D.Jacks., Bot. Jahrb. Syst. 123(2): 263. 2001

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara kannada), Peninsula to N India; Bangladesh, China & SE Asia.

Tournefortia L.

Tournefortia reticosa Wight, Icon. Pl. Ind. Orient. 4: t. 1386. 1848; C.B.Clarke in Hook. f., Fl. Brit. India 4: 146. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Trichodesma R. Br.

Trichodesma inaequale Edgew., J. Asiat. Soc. Bengal, part 2, Nat. Hist. 21: 175. 1852

Trichodesma amplexicaul auct. non Roth, 1821; C.B.Clarke in Hook.f., Fl. Brit. India 4: 153. 1883, p.p.

Trichodesma sedgewickianum S. P. Banerji, Bull. Bot. Soc. Bengal 16:10. 1962

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Mysuru, Raichur), Gujarat, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Trichodesma indicum (L.) Lehm., Pl. Asperif. Nucif. 193. 1818; C. B. Clarke in Hook. f., Fl. Brit. India 4: 153. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Borago indica L., Sp. Pl. 137 1753

Borago spinulosa Roxb., Fl. Ind.ed. 1, 1: 459 1820

Trichodesma hirsutum Edgew., J. Asiat. Soc. Bengal Part 2, Nat. Hist. 21: 175 1853

Boraginella indica (L.) Kuntze, Revis. Gen. Pl. 2: 435 1891

var. *indicum*

Distribution: Karnataka (throughout), throughout India; Sri Lanka, Middle East, Mauritius.

Local/Common/Vernacular names: *Adhomuki, Katte tumbe soppu* (Kannada)

var. *amplexicaule* (Roth) T. Cooke, Fl. Bombay 2: 281.1967

Trichodesma amplexicaule Roth, Nov. Sp. Pl. 104.1821; C.B.Clarke in Hook.f., Fl. Brit. India 4: 153. 1883.

Distribution: Karnataka (Chitradurga), Maharashtra; endemic.

Trichodesma zeylanicum (Burm.f.) R.Br., Prodr. Fl. Nov. Holl. 496. 1810; C.B.Clarke in Hook. f., Fl. Brit. India 4: 154. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 176. 1984.

Borago zeylanica Burm.f., Fl. Indica 41. 1768

Pollichia zeylanica (Burm.f.) F.Muell., Syst. Census Austral. Pl. 100. 1882

Borraginoides zeylanica (Burm.f.) Hiern, Cat. Afr. Pl. 1: 720. 1898

Distribution: Karnataka (throughout), throughout plains India; Australia, Pacific Islands, SE Asia, tropical E Africa.

Local/Common/Vernacular name: *Ettina nalige gida* (Kannada)

Brassicaceae

Brassica L.

Brassica cretica Lam., Encycl. 1: 747 1785

Brassica oleracea var. *botrytis* L., Sp. Pl. 667. 1753; C.J.Saldanha, Fl. Karnataka 1: 321. 1984

Distribution: Karnataka (Belagavi, Davanagere); native to Europe.

Local/Common/Vernacular name: *Cauliflower* (English); *Hookosu* (Kannada)

Brassica juncea (L.) Czern. in Czern., Consp. Pl. Chark. 8. n. 5. 1859; B.D.Sharma et al., Fl. Karnataka Anal. 7. 1984.

Sinapis juncea L., Sp. Pl. 668. 1753

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Hassan, Kalburgi, Kodagu, Kolar, Shivamogga), throughout India; pantropical.

Local/Common/Vernacular name: *Indian Red Mustard* (English); *Kempusasive* (Kannada)

Brassica nigra (L.) K.Koch, Roehl., Deustchl. Fl. (ed.3) 4: 713. 1833; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Sinapis nigra L., Sp. Pl. 668. 1753

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Kalburgi, Kodagu, Mysuru), Arunachal Pradesh, Assam, Bihar, Delhi, Goa, Himachal Pradesh, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh; Europe, North Africa, South West Asia and temperate parts of India

Local/Common/Vernacular name: *Karisasive* (Kannada)

Brassica oleracea L., Sp. Pl. 667 1753

Brassica oleracea var. *capitata* L., Sp. Pl. 667. 1753; C.J.Saldanha, Fl. Karnataka 1: 321. 1984

Distribution: Karnataka (Belagavi, Davanagere, Kalburgi), Assam, Bihar, Delhi, Goa, Madhya Pradesh, Maharashtra, Manipur, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh; native to Europe

Local/Common/Vernacular name: *Cabbage* (English); *Kosu* (Kannada)

Brassica rapa L., Sp. Pl. 666 1753

Brassica campestris L., Sp. Pl. 666 1753; B.D.Sharma et al., Fl. Karnataka Anal. 7. 1984.

Distribution: Karnataka (Uttara Kannada), Arunachal Pradesh, Bihar, Himachal Pradesh, Madhya Pradesh, Maharashtra, Mizoram, Tamil Nadu, Uttar Pradesh

Cardamine L.

Cardamine africana L., Sp. Pl. 655. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Distribution: Karnataka (Chikkamagaluru, Mysuru), Arunachal Pradesh, Kerala, Tamil Nadu; Tropics of Asia

Cardamine hirsuta L., Sp. Pl. 655. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dharwad, Kodagu), Arunachal Pradesh, Assam, Himachal Pradesh, Kerala, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal; pantropical.

Local/Common/Vernacular name: *Kaadu saasuve* (Kannada)

Cardamine trichocarpa Hochst. ex A.Rich., Tent. Fl. Abyss. 1: 18. 1847; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Cardamine hirsuta var. *subumbellata* Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4:294. 1852

Cardamine subumbellata Hook. ex Hook.f. & T.Anderson, Hook.f., Fl. Brit. India 1: 138. 1872

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dharwad, Mysuru), Madhya Pradesh, Maharashtra, Manipur, Tamil Nadu; Tropical mountains of Africa & Asia

Eruca Mill.

Eruca vesicaria (L.) Cav., Gard. Dict. ed. 8 1 1768

Brassica eruca L., Sp. Pl. 667. 1753; C.J.Saldanha, Fl. Karnataka 1: 323. 1984

Eruca sativa Garsault, Traite Pl. Anim. 2: 166. t. 259. 1767. nom. inval.

Distribution: Karnataka (Dharwad); Mediterranean region

Iberis L.

Iberis amara L., Sp. Pl. 649 1753

Distribution: Karnataka (cultivated in Bengaluru, Dharwad, Mysuru), cultivated throughout India; native to Europe and N Africa.

Local/Common/Vernacular names: *Bitter candytuft*, *Rocket candytuft*, *Wild candytuft* (English)

Lepidium L.

Lepidium didymum L., Mant. Pl. 1: 92. 1767; C.J.Saldanha, Fl. Karnataka 1: 322. 1984

Senebiera pinnatifida DC. in Mem. Soc. Hist. Nat. Paris 144. t. 9. 1799

Coronopus didymus (L.) Sm., Fl. Brit. 2: 691. 1800

Senebiera didyma (L.) Pers., Syn. 2: 185. 1800

Distribution: Karnataka (Belagavi, Bengaluru, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru)

Lepidium sativum L., Sp. Pl. 644. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Davanagere, Dharwad, Kolar), throughout India; Native to Western Asia

Local/Common/Vernacular name: *Garden cress* (English); *Kurthike* (Kannada)

Raphanus L.

Raphanus raphanistrum L., Sp. Pl. 669 1753

subsp. *sativus* (L.) Domin, Beih. Bot. Centralbl. 26(2): 255 1910

Raphanus sativus L., Sp. Pl. 669 1753; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Distribution: Karnataka (Bengaluru, Chitradurga, Davanagere); Cultivated for its edible root

Rorippa Scop.

Rorippa indica (L.) Hiern, Cat. Afr. Pl. Welw. 26. 1896; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Sisymbrium indicum L., Mant. Pl. 1: 93. 1767

Nasturtium indicum (L.) DC., Prodr. 1: 139. 1824

Distribution: Karnataka (Belagavi, Bengaluru, Davanagere, Dharwad, Kodagu, Kolar, Mysuru, Tumakuru, Uttara Kannada), throughout India; Indomalesia

Local/Common/Vernacular name: *Kadu sasive* (Kannada)

Schouwia DC.

Schouwia purpurea (Forssk.) Schweinf., Bull. Herb. Boissier 4(App. 2): 183 1896

Thlaspi arabicum Vahl, Symb. 2: 76. 1790

Schouwia arabica (Vahl) DC., Syst. Nat. 2: 644. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 8. 1984.

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Kalburgi), Andhra Pradesh, Maharashtra; North East Africa to India.

Bromeliaceae**Ananas** Miller

Ananas comosus (L.) Merr., Interpr. Herb. Amboin. 133.1917; B.D.Sharma et al., Fl. Karnataka Anal. 281. 1984.

Distribution: Karnataka (cultivated in Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), cultivated in S & NE India; native to S America.

Local/Common/Vernacular name: *Pineapple* (English); *Ananas* (*Kunabi*)

Buddlejaceae**Buddleja** L.

Buddleja asiatica Lour., Fl. Cochinch. 72.1790; B.D.Sharma et al., Fl. Karnataka Anal. 171. 1984.

Buddleja neemda Buch.-Ham. ex Roxb., Fl. Ind.ed.1, 1: 411.1820

Buddleja discolor Roth, Nov. Pl. Sp. 8. 1821

Buddleja serrulata Roth, Nov. Pl. Sp. 82.1821

Buddleja subserrata Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 92.1825

Buddleja densiflora Blume, Bijdr. 7.1826

Buddleja sundaica Blume, Bijdr. 743.1826

Distribution: Karnataka (cultivated in Bengaluru, Chikkamagaluru, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Uttara kannada), E, S & SE Asia.

Local/Common/Vernacular names: *Hole lakki*, *Karakani* (Kannada)

Buddleja davidii Franch., Nouv Arch. Mus. Hist. Nat. II, 10: 65.1887; B.D.Sharma et al., Fl. Karnataka Anal. 171. 1984.

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), native to China.

Local/Common/Vernacular names: *Butterfly-bush*, *Orange eye*, *Summer lilac* (English)

Buddleja madagascariensis Lam., Encycl. 1: 513.1785; Ramaswamy & Razi, Fl. Bangalore Dist. 437. 1973.

Buddleja heterophylla Lindl., Edwards's Bot. Reg. 15: t. 1259.1829

Nicodemia madagascariensis (Lam.) R. Parker, Forest Fl. Punjab ed. 2, 357.1924. *Adenoplea madagascariensis* (Lam.) Eastw., Leafl. W. Bot. 1: 197.1936

Distribution: Karnataka (cultivated in Bengaluru), native to Madagascar.

Local/Common/Vernacular name: *Smokebush* (English)

Burmanniaceae**Burmannia L.**

Burmannia candelabrum Gagnep., Bull. Soc. Bot. France 54: 462.1907; Hajra, Fasc. Fl. India 19: 7. 1988.
 Distribution: Karnataka (Chikmagluru, Dakshina Kannad, Kodagu, Mysuru, Uttara Kannada), Kerala, Maharashtra, Meghalaya, Tamil Nadu; Bangladeah.

Burmannia candida Griff. ex Hook.f., Fl. Brit. India 5: 665.1888; Hajra, Fasc. Fl. India 19: 7. 1988.
 Distribution: Karnataka (Chikmagluru, Dakshina Kannad, Kodagu, Mysuru, Uttara Kannada), Kerala, Maharashtra, Meghalaya, Tamil Nadu; Bangladeah

Burmannia championii Thwaites, Enum. Pl. Zeyl. 325. 1864; Mesta et al., J. Threatened Taxa 3(1): 1465. 2011.

Distribution: Karnataka (Uttara Kannada), Andaman Islands, Kerala; Sri Lanka, Malayan peninsula, southern China and Japan, Taiwan.

Burmannia coelestis D.Don, Prodr. Fl. Nepal. 44.1825; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Burmannia uniflora Rottler ex Spreng., Syst. Veg. 4(2): 23.1827

Burmannia triflora Roxb., Fl. Ind. ed. 1832 2: 117.1832

Burmannia bifurca Ham. ex Hook.f., Fl. Brit. India 5: 665.1888

Distribution: Karnataka (Chikmagluru, Dakshina Kannad, Kodagu, Mysuru, Uttara Kannada), Peninsula to E & NE India;; Australia, China, Mauritius, S & SE Asia.

Burmannia disticha L., Sp. Pl. 287.1753; Hajra, Fasc. Fl. India 19:11.1988.

Distribution: Karnataka (Chikkamagaluru), E & NE India; Australasia, China, S & SE Asia

Burmannia indica Jonker, Monogr. Burmann. 161.1938; Hajra, Fasc. Fl. India 19: 7. 1988.

Distribution: Karnataka (Udupi, Uttara Kannada), Kerala, Maharashtra; endemic.

Burmannia pusilla (Miers) Thwaites, Enum. Pl. Zeyl. 325.1864

Gonianthes pusilla Miers, Trans. Linn. Soc. London 18: 537.1841

Cyananthus pusillus (Miers) Wall., Numer. List 9008.1849

Burmannia coelestis var. *pusilla* (Miers) Thwaites, Enum. Pl. Zeyl. 325.1864

Distribution: Karnataka (Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka, Cambodia, Laos, Vietnam.

Burseraceae**Boswellia** Roxb. ex Colebr.

Boswellia serrata Roxb. ex Colebr., Asiat. Res. 9: 379. t. 5. 1807; B.D.Sharma et al., Fl. Karnataka Anal. 43. 1984. Plate 5A

Boswellia glabra Roxb., Pl. Coromandel 3: t. 207. 1811

Distribution: Karnataka (Ballari, Belagavi, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Mysuru, Tumakuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Bihar, Delhi, Gujarat, Himachal Pradesh, Kerala, Madhya Pradesh, Maharashtra, Manipur, Punjab, Rajasthan, Tamil Nadu West Bengal; India

Local/Common/Vernacular names: *Indian olibanum tree* (English); *Dhoopada mara, Madi* (Kannada)

Bursera Jacq. ex L.

Bursera linanoe (La Llave) Rzed., Calderón & Medina, Acta Bot. Mex. 66: 103 2004

Amyris linanoe La Llave in Reg. Trimestre 1: 356 1832

Bursera delpechiana Poiss. ex Engl. in Monogr. Phan. 4: 53. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 43. 1984.

Distribution: Karnataka (Cultivated, Chitradurga, Tumakuru); Introduced, Native to Mexico

Local/Common/Vernacular name: *Mexican Linaloe tree* (English)

Canarium L.

Canarium strictum Roxb., Fl. Ind. 3: 138. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 44. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Arunachal Pradesh, Assam, Kerala, Maharashtra, Manipur, Meghalaya, Mizoram, Tamil Nadu, Tripura, West Bengal

Local/Common/Vernacular names: *The Black Dammer tree* (English); *Dhoopada kayi, Halemaddu* (Kannada)

Commiphora Jacq.

Commiphora berryi (Arn.) Engl. in A. & C. DC., Monogr. Phan. 4: 17. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 44. 1984.

Protium gileadense Wight & Arn., Prodr. Fl. Ind. Orient. 1: 177. 1834

Balsamodendron berryi Arn., Ann. Nat. Hist. 3: 86. 1839

Distribution: Karnataka (Kolar, Mandya, Mysuru, Tumakuru), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Indian Balm of Gilead* (English)

Commiphora caudata (Wight & Arn.) Engl. in A. & C. DC., Monogr. Phan. 4: 17. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 44. 1984.

Protium caudatum Wight & Arn., Prodr. Fl. Ind. Orient. 1: 176. 1834

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Hassan, Kodagu, Mysuru, Tumakuru), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka, India

Local/Common/Vernacular name: *Hill Mando* (English); *Konda mavu* (Kannada)

Commiphora wightii (Arn.) Bhandari, Bull. Bot. Surv. India 6: 327. 1965; B.D.Sharma et al., Fl. Karnataka Anal. 44. 1984.

Balsamodendron wightii Arn., Ann. Nat. Hist. 3: 86. 1839

Balsamodendrum mukul Hook. ex Stocks in Hooker's Kew J. Bot. 1: 259. 1848

Commiphora mukul (Hook. ex Stocks) Engl. in A. & DC., Monogr. Phan. 4: 26. 1883

Distribution: Karnataka (Ballari, Kalburgi, Mysuru, Raichuru, Koppala), Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Rajasthan, Tamil Nadu; Afghanistan, Central Asia, N Africa, Pakistan.

Local/Common/Vernacular names: *Gugal, Guggul, Gugul, Indian bdellium-tree, Mukul myrrh tree* (English); *Guggulu* (Kannada)

Garuga Roxb.

Garuga pinnata Roxb., Pl. Coromandel 3: t. 208. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 44. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; Indomalesia.

Buxaceae

Sarcococca Lindl.

Sarcococca coriacea (Hook.) cSweet, Hort. Brit. 491.1826

Buxus saligna D.Don, Prodr. Fl. Nepal. 63.1825

Pachysandra coriacea Hook., Exot. Fl. 2: t. 148.1825

Sarcococca pruniformis Lindl., Bot. Reg. 12: t. 1021.1826; B.D.Sharma et al., Fl. Karnataka Anal. 243. 1984.

Sarcococca nepalensis Royle, Ill. Bot. Himal. Mts. 327.1836

Sarcococca trinervia Wight, Icon. Pl. Ind. Orient. 5: t. 1877.1852

Sarcococca zeylanica Baill., Monogr. Buxac. 52.1859

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala, Tamil Nadu; Indonesia, Myanmar, Nepal, Thailand, Sri Lanka, Vietnam.

Cactaceae

Cereus Mill.

Cereus repandus (L.) Mill., Gard. Dict. ed. 8 5 1768

Cactus peruvianus L., Sp. Pl. 467. 1753; C.J.Saldanha, Fl. Karnataka 1: 140. 1984

Cereus peruvianus (L.) Mill., Gard. Dict. 4. 1768

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Vijayapura); Native to America

Harrisia Britton

Harrisia bonplandii (Parm.) Britton & Rose, Cact. 2: 157 1920

Cereus bonplandii Parm ex Pfeiffer, Enum. Diagn. Cact. 108. 1837; C.J.Saldanha, Fl. Karnataka 1: 140. 1984

Eriocereus bonplandii (Parm. ex Pfeiff.) Riccob., Bull. Reale Orto Bot. Girardino Colon Palermo 8: 238. 1909

Distribution: Karnataka (Bengaluru, Davanagere, Hassan); Native to South America

Nopalea Salm-Dyck

Nopalea cochenillifera (L.) Salm-Dyck, Cact. Hort. Dyck. 64 1850

Cactus cochenillifer L., Sp. Pl. 468. 1753; C.J.Saldanha, Fl. Karnataka 1: 141. 1984

Opuntia cochinellifera (L.) Mill., Gard. Dict. 4. 1768

Distribution: Karnataka (Belagavi, Bengaluru, Davanagere, Dharwad, Mysuru), Bihar, Maharashtra, Tamil Nadu, Tripura, Uttar Pradesh; Native to Mexico

Opuntia Mill.

Opuntia dillenii (KerGawler) Haw., Suppl. Pl. Succ. 79 1819; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Cactus dillenii Ker Gawl., Edward's Bot. Reg. 3: t.255. 1818

Cactus indicus Roxb., Fl. Ind. 2: 475. 1832

Opuntia stricta var. *dillenii* (KerGawler) L.D.Benson, Cact. Succ. J. (Los Angeles) 41: 126 1969

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Raichur), Bihar, Delhi, Madhya Pradesh, Meghalaya, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; native to America

Local/Common/Vernacular name: *Papas kalli* (Kannada)

Opuntia elatior (Willd.) Mill., Gard. Dict. 4. 1768; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Opuntia nigricans (Haw.) Haw., Syn. Pl. Succ. 189. 1812

Cactus elatior (Mill.) Willd., Enum. Pl. Suppl. 34. 1814

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Davanagere, Dharwad, Kalburgi, Mysuru), Andhra Pradesh, Assam, Bihar, Goa, Himachal Pradesh, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh; pantropical.

Opuntia ficus-indica (L.) Mill., Gard. Dict. ed. 8 2 1768

Cactus ficus-indica L., Sp. Pl. 468 1753

Opuntia vulgaris Mill., Gard. Dict. 1. 1768

Cactus monacanthos Willd., Enum. Pl. Suppl. 33. 1813

Opuntia monacantha (Willd.) Haw., Suppl. Pl. Succ. 81. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 116. 1984.

Distribution: Karnataka (Bengaluru, Davanagere, Hassan, Kalburgi), cultivated in drier parts of India; native to America

Opuntia stricta (Haw.) Haworth, Syn. Pl. Succ. 191. 1812; C.J.Saldanha, Fl. Karnataka 1: 141. 1984

Cactus strictus Haw. in Misc. 188. 1803

Distribution: Karnataka (Davanagere, Kodagu), cultivated /run wild in drier parts of India; native to Americas.

Calceolariaceae

Calceolaria L.

Calceolaria mexicana Benth., Pl. Hartw. 47. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 189. 1984.

Fagelia mexicana (Benth.) Kuntze, Revis. Gen. Pl. 2: 459. 1891

Distribution: Karnataka (Cahmarajanagar, Chikkamagaluru, Kodagu, Shivamogga), Peninsular India; native to Mexico.

Local/Common/Vernacular names: *Lady's purse, Pocketbookflower, Slipperflower, Slipperwort* (English)

Calophyllaceae

Calophyllum L.

Calophyllum apetalum Willd., Ges. Naturf. Freunde Berlin 5: 79 1811; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984. Plate 20D

Calophyllum decipiens Wight, Ill. Ind. Bot. 1: 128 1831

Calophyllum wightianum Wall. ex Planch. & Triana, Ann. Sci. Nat., Bot. IV, 15: 256 1861

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Hole hone, Irai, Kallu hone, Kalpoone, Shri hone* (Kannada)

Calophyllum austroindicum Kosterm. ex P.F.Stevens in J. Arnold Arbor. 61: 250 1980; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Calophyllum trapezifolium auct. non Thwaites 1864

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala, Tamil Nadu; endemic.

Calophyllum inophyllum L., Sp. Pl. 513 1753; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Mysuru, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Kerala, Maharashtra, Odisha, Tamil Nadu West Bengal; paleotropical

Calophyllum polyanthum Wall. ex Choisy, Descr. Guttif. Inde. 43 1849; C.J.Saldanha, Fl. Karnataka 1: 203. 1984

Calophyllum tomentosum Wight, Ill. Ind. Bot. 1: 128 1840

Calophyllum elatum Bedd., Fl. Sylv. S. India 1: t. 2 1869

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Arunachal Pradesh, Assam, Kerala, Maharashtra, Manipur, Meghalaya, Sikkim, Tamil Nadu, West Bengal.

Mammea L.

Mammea americana L., Sp. Pl. 512 1753; C.J.Saldanha, Fl. Karnataka 1: 209. 1984

Distribution: Karnataka (cultivated in Bengaluru), cultivated in gardens; native to tropical S America.

Local/Common/Vernacular names: *Mamey*, *Mamey apple*, *Mammee*, *Mammee apple*, *Santo Domingo apricot*, *South American apricot*, *Tropical apricot* (English)

Mammea suriga (Buch.-Ham. ex Roxb.) Kosterm., Pengum. Lemb. Pusat Penjel. Kehut. 72: 23 1961; B.D.Sharma et al., Fl. Karnataka Anal. 19. 1984.

Calophyllum suriga Buch.-Ham. ex Roxb., Fl. Ind. 2: 608 1824

Ochrocarpos longifolius Benth. & Hook.f. ex T.Anderson, Fl. Brit. India 1: 270 1874

Distribution: Karnataka (Belagavi, Dakshina Kannada, Shivamogga, Uttara Kannada, sometimes cultivated), Assam, Goa, Kerala, Maharashtra, Tamil Nadu, Uttar Pradesh, West Bengal; Sri Lanka.

Local/Common/Vernacular names: *Ceylon ironwood*, *Indian rose chestnut* (English); *Punya*, *Surige mara*, *Wundi* (Kannada)

Mesua L.

Mesua ferrea L., Sp. Pl. (ed. 2) 1: 734 1762; C.J.Saldanha, Fl. Karnataka 1: 210. 1984

Calophyllum nagassarium Burm.f., Fl. Indica 121. 1768

Mesua nagana Gardner, Calcutta J. Nat. Hist. 8: 4. 1847

Mesua nagassarium (Burm.f.) Kosterm., Ceylon J. Sci., Biol. Sci. 12: 71 1976

Mesua thwaitesii sensu, Kesh. Murthy & Yogan, 60. 1990, non Planch. & Triana 1861

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India; Indomalesia

Local/Common/Vernacular name: *Naagasampige* (Kannada)

Poeciloneuron Bedd.

Poeciloneuron indicum Bedd. in J. Linn. Soc., Bot. 8: 267 1865; B.D.Sharma et al., Fl. Karnataka Anal. 19. 1984. Plate 20E

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Mysuru, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Balige mara* (Kannada)

Campanulaceae

Asyneuma Griseb & Schenk.

Asyneuma fulgens (Wall.) Briquet, Candollea 4: 334. 1931

Campanula fulgens Wall. in Roxb., Fl. Ind. 2: 99. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984.

Distribution: Karnataka (Chikkamagaluru, Mysuru, Shivamogga), Kerala, Manipur, Meghalaya, Sikkim, Tamil Nadu, West Bengal; Bhutan, China, Myanmar, Nepal, Sri Lanka.

Campanula L.

Campanula alphonsei Wall. ex A. DC., Monogr. Camp. 294. 1830 & DC., Prodr. 7: 473.1839;

B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984.

Plate 20F

Distribution: Karnataka (Chikkamagaluru, Mysuru), Kerala, Maharashtra, Tamil Nadu; endemic.

Campanula dimorphantha Schweinf., Beitr. Fl. Aethiop. 140. 1867

Campanula benthamii Wall., Numer. List n. 1290.1829

Campanula canescens Wall. ex A. DC., Monogr. Camp. 292. 1830 & DC., Prodr. 7: 473. 1839 non Roth 1827. Hook. f. & Thomson, J. Linn. Soc. 2: 23. 1858; C.B.Clarke in Hook.f., Fl. Brit. India 3: 439. 1881.

Campanula wallichii Babu, J. Bombay Nat. Hist. Soc. 65: 808 1969. nom. Supefl.; B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984

Campanula colorata Collett, Fl. Simlensis 291. 1902 p.p.

Campanula benthamii Wall. ex Kitam., Fl. Afghanistan 377.1960

Distribution: Karnataka (Chikkamagaluru), throughout India; Africa, Afghanistan, China, Laos, Myanmar, Nepal, Pakistan, Sri Lanka, Vietnam.

Hippobroma G.Don

Hippobroma longiflora (L.) G.Don, Gen. Hist. 3: 717.1834

Lobelia longiflora L., Sp. Pl. 930.1753

Laurentia longiflora (L.) Peterm., Pflanzenreich 444 1845; B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984.

Isotoma runcinata Hassk., Bonplandia 7: 181 1859

Isotoma longiflora var. *runcinata* (Hassk.) Panigrahi, P.Daniel & M.V.Viswan., Indian J. Forest. 4: 151 1981

Solenopsis longiflora (L.) M.R.Almeida, Fl. Maharashtra 3A: 155. 2001

Distribution: Karnataka (Uttara Kannada), Andaman & Nicobar Islands, Maharashtra; native to carribean Islands, introduced and naturalized in tropics and subtropics.

Lobelia L.

Lobelia alsinoides Lam., Encycl. 3: 588. 1792

Lobelia triangulata Roxb., Hort. Bengal. 16. 1814

Lobelia trigona Roxb., Fl. Ind. 2: 111.1824; C.B.Clarke in Hook.f., Fl. Brit. India 3: 423. 1881.

Lobelia gratiolooides Roxb. ex C. Presl, Prodr. Monogr. Lobel. 50.1836

Dortmannia alsinodes (Lam.) Kuntze, Revis. Gen. Pl. 2: 972. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 151. 1984

Dortmannia trigona (Roxb.) Kuntze, Revis. Gen. Pl. 2: 380. 1891

Distribution: Karnataka (Belagavi, Bengaluru, Chikmagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Peninsula India extending eastwards to Assam; S & SE Asia, New Guinea.

Lobelia heyneana Schult., Syst. Veg. 5: 50.1819; B.D.Sharma et al., Fl. Karnataka Anal. 151. 1984.

Lobelia decurrens Roth, Nov. Pl. Sp. 145. 1821

Lobelia trialata Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 157. 1825; C.B.Clarke in Hook.f., Fl. Brit. India 3: 425. 1881.

Rapuntium arenariooides C.Presl, Prodr. Monogr. Lobel. 13. 1836

Rapuntium trialatum (Buch.-Ham. ex D.Don) C.Presl, Prodr. Monogr. Lobel. 13. 1836

Lobelia arenariooides (C.Presl.) A.DC., Prodr. 7: 367.1839

- Lobelia subincisa* Wall. ex A.DC., Prodr. 7: 367. 1839
Lobelia dichotoma Miq., Fl. Ned. Ind. 2: 576. 1857
Lobelia subracemosa Miq., Fl. Ned. Ind. 2: 576. 1857
Lobelia umbrosa Hochst. ex Hemsl. in Oliv., Fl. Trop. Afr. 3: 468. 1877
Lobelia trialata var. *lamiifolia* C.B.Clarke in Hook. f., Fl. Brit. India 3: 425. 1881
Lobelia zeylanica var. *walkeri* C.B.Clarke in Hook. f., Fl. Brit. India 3: 425. 1881
Dortmannia trialata (Buch.-Ham. ex D.Don) Kuntze, Revis. Gen. Pl. 2: 973. 1891
Lobelia trialata var. *umbrosa* (Hochst. ex Hemsl.) Chiov., Result. Sci. Miss. Stefan.-Paoli Somal. Ital. 1: 109. 1916
Lobelia aligera Haines, J. Proc. Asiat. Soc. Bengal 15: 316. 1920
Lobelia zeylanica var. *aligera* (Haines) Haines, Bot. Bihar Orissa 4: 501. 1922
Lobelia dichotoma var. *aligera* (Haines) E.Wimm., Ann. Naturhist. Mus. Wien 56: 345. 1948
Lobelia heyneana var. *lamiifolia* (C.B.Clarke) E.Wimm., Ann. Naturhist. Mus. Wien 56: 345. 1948
 Distribution: Karnataka (throughout), throughout India, Bhutan; Africa, China, New Guinea, S & SE Asia.

- Lobelia leschenaultiana** (C.Presl) Skottsb., Acta Horti Gothob. 4: 4. 1928
Lobelia excelsa Lesch. ex Roxb., Fl. Ind. ed.1, 2: 114. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 151. 1984.
Rapuntium leschenaultianum C.Presl, Prodr. Monogr. Lobel. 24. 1836
Lobelia aromatica Moon ex Wight, Icon. Pl. Ind. Orient. 4: t. 1172. 1848
Dortmannia leschenaultiana (C.Presl) Kuntze, Revis. Gen. Pl. 2: 972. 1891
 Distribution: Karnataka (Chamarajanagar), Kerala, Madhya Pradesh, Tamil Nadu; Sri Lanka.

- Lobelia nicotianifolia** Roth ex Schult., Syst. Veg. 5: 47. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 151. 1984.
Rapuntium nicotianifolium (Roth ex Schult.) C.Presl, Prodr. Monogr. Lobel. 24. 1836
Lobelia trichandra Wight, Icon. Pl. Ind. Orient. 4: t. 1171. 1848
Lobelia nicotianifolia var. *trichandra* (Wight) C.B.Clarke in Hook. f., Fl. Brit. India 3: 427. 1881
Dortmannia nicotianifolia (Roth ex Schult.) Kuntze, Revis. Gen. Pl. 2: 973. 1891
Lobelia courtallensis K.K.N.Nair, Proc. Indian Acad. Sci., B 87: 105. 1978
 Distribution: Karnataka (Chikkamagaluru, Chamarajanagar, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara kannada), Goa, Kerala, Maharashtra, Tamil Nadu; SE Asia, Sri Lanka, Taiwan.
 Local/Common/Vernacular names: *Kadu hoge soppu*, *Kadu thambaku* (Kannada)

- Lobelia zeylanica** L., Sp. Pl. 932. 1753
Lobelia hirta Wall., Fl. Ind., ed. 1, 2: 113. 1824
Lobelia gulparae Ham. ex Wall., Numer. List n. 1311. 1829
Lobelia affinis Wall. & G.Don, Gen. Hist. 3: 709. 1834
Rapuntium zeylanicum (L.) C.Presl, Prodr. Monogr. Lobel. 13. 1836
Lobelia subcuneata Miq., Fl. Ned. Ind. 2: 574. 1857
Lobelia lobbiana Hook.f. & Thomson, J. Proc. Linn. Soc., Bot. 2: 28. 1858
Lobelia affinis var. *lobbiana* (Hook.f. & Thomson) C.B.Clarke in Hook.f., Fl. Brit. India 3: 424. 1881
Dortmannia trigona var. *affinis* (C.Presl) Kuntze, Revis. Gen. Pl. 2: 380. 1891
Dortmannia zeylanica (L.) Kuntze, Revis. Gen. Pl. 2: 380. 1891
Lobelia succulenta var. *lobbiana* (Hook.f. & Thomson) E.Wimm., Ann. Naturhist. Mus. Wien 56: 361. 1948
Lobelia zeylanica var. *lobbiana* (Hook.f. & Thomson) Y.S.Lian in Fl. Reipubl. Popul. Sin. 73(2): 149. 1983
 Distribution: Karnataka (Dakshina Kannada, Uttara Kannada), Assam, Goa, Kerala, Sikkim, Maharashtra, Meghalaya, Tamil Nadu, West Bengal; China, Fiji Islands, New Guinea, S & SE Asia, Taiwan.

Wahlenbergia Schrad. ex Roth

Wahlenbergia erecta (Roth ex Roem. & Schult.) Tuynin Steenis, Fl. Males. 1, 6: 113. 1960; B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984.

Dentella erecta Roth ex Roem. & Schult., Syst. Veg. 5: 25. 1819

Wahlenbergia perotifolia Wight & Arn., Prod. Fl. Penin. Ind. Orient. 405. 1834

Cephalostigma schimperi Hochst. ex A. Richard, Tent. Fl. Abyss. 2: 2. 1851; C.B.Clarke in Hook.f., Fl. Brit. India 3: 428. 1881.

Cephalostigma erecta (Roth ex Roem. & Schult.) Vatke, Linnaea 38: 699. 1874 p. p.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; S & SE Asia, tropical Africa.

Wahlenbergia flexuosa (Hook.f. f. & Thomson) Thulin, Symb. Bot. Upsal. 21(1): 158. 1975

Cephalostigma flexuosum Hook. f. & Thomson, J. Linn. Soc. 2: 26. 1858; B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984.

Distribution: Karnataka (Chikkamagaluru, Uttara Kannada), Maharashtra; tropical Africa.

Wahlenbergia gracilis (G.Forst.) A.DC., Monogr. Campan. 142. 1830

Campanula gracilis G.Forst., Fl. Ins. Austr. 15. 1786

Lightfootia gracilis (G.Forst.) Miq., Fl. Ned. Ind. 2: 567. 1857

Cervicina gracilis (G.Forst.) Britten, Ill. Austral. Pl. Cook's Voy. 56. 1905

Distribution: Karnataka (cultivated and also as an escape), throughout India; Australia, E Asia, New Zealand, S Africa

Local/Common/Vernacular name: *Australian Bluebell* (English)

Wahlenbergia marginata (Thunb.) A. DC., Monogr. Camp. 143. 1830 & in DC., Prod. 7: 434. 1839; B.D.Sharma et al., Fl. Karnataka Anal. 150. 1984.

Campanula marginata Thunb., Fl. Japan 89. 1784

Campanula dehiscens Roxb ex Wall., Asiat. Res. 12: 571, t.6. 1816

Campanula agrestis Wall. in Roxb., Fl. Ind. 2: 97. 1824

Wahlenbergia indica A.DC., Monogr. Camp. 145. 1830

Wahlenbergia agrestis (Wall.) A. DC., Monogr. Camp. 145. 1830 & in DC., Prod. 7: 434. 1839

Wahlenbergia dehiscens (Roxb.) A. DC., Monogr. Camp. 145. 1830 & in DC., Prod. 7: 434. 1839

Distribution: Karnataka (Ballari, Chitradurga, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), throughout India; Australia, E, S & SE Asia, New Zealand, S Africa; naturalized in the Pacific Islands, N America.

Cannabaceae

Aphananthe Planch.

Aphananthe cuspidata (Blume) Planch. in DC., Prodr. 17: 209. 1873; C.J.Saldanha, Fl. Karnataka 1: 105. 1984

Cyclostemon cuspidatus Blume, Bijdr. 599. 1826

Gironniera reticulata Thwaites, Enum. Pl. Zeyl. 268. 1861

Gironniera cuspidata (Blume) Kurz, For. Fl. 2: 470. 1877

Distribution: Karnataka (Hassan, Kodagu, Mysuru, Uttara Kannada), Kerala, Tamil Nadu; India, Sri Lanka, Burma, Malesia, China

Cannabis L.

Cannabis sativa L., Sp. Pl. 1027 1753; Seeth. et al., Fl. 99. 2000

Distribution: Karnataka (Kalburgi), throughout India

Local/Common/Vernacular names: *Soft hemp, True hemp* (English); *Bhangi* (Kannada)

Celtis L.

Celtis philippensis Blanco, Fl. Filip. 197. 1837; C.J.Saldanha, Fl. Karnataka 1: 107. 1984

Celtis wightii Planch., Ann. Sci. Nat. Bot. ser. 3, 10:307.1848

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Andhra Pradesh, Kerala, Tamil Nadu; India, Malesia, Tropical Africa

Celtis tetrandra Roxb., Fl. Ind. 2: 63. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 257. 1984.

Celtis serotina Planch., Ann. Sci. Nat. Bot. ser. 3, 10: 301. 1848

Distribution: Karnataka (Chikkaballapur, Chikkamagaluru, Hassan, Kodagu, Kolar, Mysuru, Uttara Kannada), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Himachal Pradesh, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Rajasthan, Tamil Nadu, Uttaranchal; India, Malesia, Burma, Bangladesh.

Celtis timorensis Span. in Linnaea 15: 343. 1841; C.J.Saldanha, Fl. Karnataka 1: 107. 1984

Celtis cinnamomea Lindl. ex Planch., Ann. Sci. Nat. Bot. ser. 3, 10: 303. 1848

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Tamil Nadu; India, Burma, Malesia

Trema Lour.

Trema orientalis (L.) Blume in Mus. Bot. Lugd.-Bat. 2: 62. 1856; B.D.Sharma et al., Fl. Karnataka Anal. 257. 1984.

Celtis orientalis L., Sp. Pl. 1044. 1753

Sponia wightii Planch., Ann. Sci. Nat. Bot. ser. 3, 10: 322. 1848

Distribution: Karnataka (throughout), throughout India; Australia, China, Indonesia, Japan, Malaysia, Myanmar, Nepal, Pacific Islands Sri Lanka, Taiwan, Thailand, Vietnam.

Cannaceae

Canna L.

Canna × orchoides L.H.Bailey, Gentes Herb. 1: 120.1923

Distribution: Karnataka (cultivated throughout), origin not known, cultivated throughout tropica.

Canna flaccida Salisb., Icon. Stirp. Rar. 3.1791

Canna glauca var. *flaccida* (Salisb.) Willd., Sp. Pl. 1: 4.1797

Eurystylus flaccida (Salisb.) Bouché, Linnaea 18:485.1844

Distribution: Karnataka (Cultivated throughout), native to N America, cultivated elsewhere.

Canna indica L., Sp. Pl. 1.1753; B.D.Sharma et al., Fl. Karnataka Anal. 281. 1984.

Cannacorus indicus (L.) Medik., Hist. & Commentat. Acad. Elect. Sci. Theod.-Palat. 6: 379.1790

Canna orientalis Roscoe, Monandr. Pl. Scitam. t. 12.1826

Distribution: Karnataka (Cultivated throughout), cultivated throughout India; native to tropical America, pantropical.

Local/Common/Vernacular names: *Baale sugandhi, Kaadu baale, Samudra baale* (Kannada)

Capparaceae

Cadaba Forssk.

Cadaba fruticosa (L.) Druce, Bot. Exch. Club. Soc. Brit. Isles 3: 415. 1914; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Cleome fruticosa L., Sp. Pl. 671 1753

Cadaba indica Lam., Encycl. 1: 544. 1785

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Uttara Kannada, Vijayapura), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Gujarat, Himachal Pradesh, Jammu & Kashmir, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh; Bangladesh, Indo-China: Myanmar, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Kegnike, Maragacha* (Kannada)

Capparis L.

Capparis baducca L., Sp. Pl. 504 1753

Capparis rheedei DC., Prodr. 1: 246. 1824

Capparis heyneana Wall. ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 25. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Distribution: Karnataka (Belagavi, Dharwad, Shivamogga, Uttara Kannada), Goa, Kerala, Tamil Nadu; endemic.

Capparis cleghornii Dunn, Fl. Madras 46(33). 1915; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Capparis roxburghii auct non DC. 1824

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Tottilu darji* (Kannada)

Capparis decidua (Forssk.) Edgew. in J. Linn. Soc. Bot. 6: 184. 1862; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Sodada decidua Forssk., Fl. Aegypt.-Arab. 81. 1775

Capparis aphylla Roth, Nov. Pl. Sp. 238. 1821

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dharwad, Kalburgi, Vijayapura), Andhra Pradesh, Bihar, Delhi, Gujarat, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Tropical North East Africa to India

Local/Common/Vernacular name: *Kariuppigida* (Kannada)

Capparis divaricata Lam., Encycl. 1:606.1785; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Plate 5B

Capparis stylosa DC., Prodr. 1: 246. 1824

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Mandya, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Bihar, Kerala, Maharashtra, Tamil Nadu; Sri Lanka

Local/Common/Vernacular name: *Thottala* (Kannada)

Capparis grandiflora Wall. ex Hook.f. & Thomson, Fl. Brit. India 1: 174. 1872; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Capparis pyrifolia Wight & Arn., Icon. Pl. Ind. Orient. t. 1047. 1846, non Lam. 1785

Distribution: Karnataka (Bengaluru, Hassan, Kalburgi, Mysuru, Tumakuru), Kerala, Maharashtra, Tamil Nadu; endemic.

Capparis grandis L.f., Suppl. Pl. 263. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bidar, Chitradurga, Dharwad, Kalburgi, Kolar, Mysuru, Uttara Kannada), Andhra Pradesh, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu; India, Myanmar, Sri Lanka, Thailand.

Local/Common/Vernacular name: *Thotlu mullina gida* (Kannada)

Capparis moonii Wight, Illustr. 1: 35. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Mullukathari* (Kannada)

Capparis rotundifolia Rottler, Ges. Naturf. Freunde Berlin Neue Schriften 4: 185. 1803; B.D.Sharma et al., Fl. Karnataka Anal. 10. 1984.

Capparis pedunculosa Wall ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 27. 1834

Capparis longispina Hook.f. & Thomson, Hook.f., Fl. Brit. India 1: 176. 1872

Capparis pedunculosa var. *longispina* (Hook.f. & Thomson) T. Cooke, Fl. Bombay 1: 48. 1901

Distribution: Karnataka (Uttara Kannada), Andhra Pradesh, Goa, Maharashtra, Tamil Nadu; Myanmar, Sri Lanka.

Capparis sepiaria L., Syst. Nat. (ed.10) 2: 1071. 1759; B.D.Sharma et al., Fl. Karnataka Anal. 11. 1984.

Plate 5C

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Delhi, Goa, Gujarat, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Tropical Africa to Australia

Local/Common/Vernacular names: *Ippi, Kadukathari* (Kannada)

Capparis spinosa L., Sp. Pl. 1: 503. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 11. 1984.

Distribution: Karnataka (Kodagu), Bihar, Goa, Gujarat, Himachal Pradesh, Jammu & Kashmir, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal

Capparis tenera Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 41. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 11. 1984.

Distribution: Karnataka (Belagavi, Uttara Kannada), Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Kerala, Maharashtra, Manipur, Mizoram, Nagaland, West Bengal; Tropical Africa to Burma

Capparis tomentosa Lam., Encycl. 1: 606. 1785

Capparis floribunda Wight, Ill. Ind. Bot. 1: 35. 1840; K.G.Bhat, J. Bombay Nat. Hist. Soc., 90: 138. 1993

Distribution: Karnataka (Udupi), Kerala; Sri Lanka, Myanmar, extending to SE Asia

Capparis zeylanica L., Sp. Pl. (ed.2) 720. 1762; B.D.Sharma et al., Fl. Karnataka Anal. 11. 1984.

Capparis horrida L.f., Suppl. Pl. 264. 1781

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal; China, Indonesia, Indian Ocean islands, Myanmar, Nepal, Philippines, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Anthudikai, Govinda phala, Tottilu balli* (Kannada)

Cleoserrata H.H.Iltis

Cleoserrata speciosa (Raf.) Iltis, Novon 17: 448 2007

Cleome speciosa Kunth, Nov. Gen. Sp. 5: 84, t. 436. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Distribution: Karnataka (Davanagere, Shivamogga, Uttara Kannada); native to tropical America & West Indies

Crateva L.

Crateva nurvala Buch.-Ham. in Trans. Linn. Soc. London 15: 121. 1827; C.J.Saldanha, Fl. Karnataka 1: 319. 1984

Capparis magna Lour., Fl. Cochinch. 330. 1790

Crateva magna (Lour.) DC., Prodr. 1: 243. 1824

Crateva religiosa G.Forst. var. *nurvula* (Buch.-Ham.) Hook.f. & Thomson in Hook.f., Fl. Brit. India 1: 172. 1872

Crateva religiosa var. *nurvula* (Buch.-Ham.) Hook.f. & Thomson in Hook.f., Fl. Brit. India 1: 172. 1872

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kodagu, Mandya, Mysuru, Shivamogga, Uttara Kannada), Bihar, Maharashtra, Meghalaya, Rajasthan, Uttar Pradesh; South Asia, Indomalaya

Local/Common/Vernacular names: *Three leaved caper* (English); *Bilipatri, Nervala* (Kannada)

Maerua Forssk.

Maerua oblongifolia (Forssk.) A.Rich. in Guill. & Perr., Fl. Seneg. Tent. 1:32. t. 5. 1831; B.D.Sharma et al., Fl. Karnataka Anal. 11. 1984.

Capparis oblongifolia Forsk., Fl. Aegypt.-Arab. 99. 1775

Niebuhria oblongifolia (Forssk.) DC., Prodr. 1: 244. 1824

Maerua ovalifolia Combess. in Jacq., Voy.Inde. 4:23. t. 24. 1844

Maerua arenaria Hook. f. & Thomson, Fl. Brit. India 1: 171. 1872

Distribution: Karnataka (Bagalkot, Ballari, Belagavi, Chitradurga, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mandya, Mysuru, Raichur, Shivamogga, Tumakuru, Vijayapura), Andhra Pradesh, Bihar, Gujarat, Jammu & Kashmir, Maharashtra, Tamil Nadu, Uttar Pradesh, West Bengal; Arabia, tropical East Africa.

Local/Common/Vernacular name: *Necklace-berried caper* (English); *Nelasakregedde* (Kannada)

Caprifoliaceae**Lonicera L.**

Lonicera japonica Thunb., Syst. Veg. ed. 14, 216. 1784

Caprifolium japonicum (Thunb.) Dum.Cours., Bot. Cult. ed. 2, 7: 209. 1814

Distribution: Karnataka (cultivated in Belagavi, Bengaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru), cultivated in gardens; native to China & Japan.

Local/Common/Vernacular name: *Japanese honeysuckle* (English)

Lonicera leschenaultii Wall., Fl. Ind. 2: 178. 1824; K. Gandhi in Saldanha & Nicolson, Fl Hassan Dist. 594. 1976.

Caprifolium leschenaultii Kuntze, Revis. Gen. Pl. 1: 274. 1891

Distribution: Karnataka (Chamarajanagar, Hassan, Mysuru), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Handi pode kode, Nolle* (Kannada)

Caricaceae

Carica L.

Carica papaya L., Sp. Pl. 1036 1753; N.P.Singh, Fl. East. Karnataka 1: 321. 1988

Distribution: Karnataka (Bengaluru, Chitradurga, Kalburgi, Kodagu), throughout India; Cultivated, native to tropical America

Local/Common/Vernacular name: *Papayi hanna gida* (Kannada)

Caryophyllaceae

Arenaria L.

Arenaria neelgherrensis Wight & Arn., Prodr. Fl. Ind. Orient. 1: 43. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 15. 1984.

Distribution: Karnataka (Belagavi, Dharwad, Uttara Kannada), Himachal Pradesh, Jammu & Kashmir, Maharashtra, Punjab, Sikkim, Tamil Nadu, Uttar Pradesh; China, Nepal, Pakistan.

Cerastium L.

Cerastium glomeratum Thuill., Fl. Env. Paris ed. 2: 226. 1799; B.D.Sharma et al., Fl. Karnataka Anal. 15. 1984.

Cerastium vulgatum var. *glomeratum* (Thuill.) Edgew. & Hook. f., Fl. Brit. India 1: 228. 1874

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru), Bihar, Himachal Pradesh, Jammu & Kashmir, Maharashtra, Tamil Nadu, Uttar Pradesh, West Bengal; pantropical.

Drymaria Willd.

Drymaria cordata (L.) Willd. ex Schult., Syst. Veg. 5: 406. 1820; C.J.Saldanha, Fl. Karnataka 1: 152. 1984

Holosteum cordatum L., Sp. Pl. 88. 1753

subsp. **cordata**

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru), Andhra Pradesh, Arunachal Pradesh, Assam, Kerala, Madhya Pradesh, Manipur, Meghalaya, Nagaland, Punjab, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand, West Bengal; native to C & S America.

subsp. **diandra** (Blume) J.A.Duke, Ann. Miss. Bot. Gard. 48: 253. 1961; B.D.Sharma et al., Fl. Karnataka Anal. 15. 1984.

Drymaria diandra Blume, Bijdr. 62. 1825

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru); paleotropical

Polycarpea Lam.

Polycarpea aurea Wight & Arn., Ann. Nat. Hist. ser. 1: 91. 1839; B.D.Sharma et al., Fl. Karnataka Anal. 16. 1984.

Polycarpea corymbosa var. *aurea* (Wight & Arn.) Wight, Illustr. 2: 44, t.110. 1850

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dakshina Kannada, Kodagu, Vijayapura), Andhra Pradesh, Chattisgarh, Jharkhand, Madhya Pradesh, Odisha, Tamil Nadu; endemic.

Polycarpaea corymbosa (L.) Lam., Tabl. Encycl. 2: 129. 1792; B.D.Sharma et al., Fl. Karnataka Anal. 16. 1984.

Achyranthes corymbosa L., Sp. Pl. 205. 1753

Distribution: Karnataka (throughout), Andhra Pradesh, Bihar, Delhi, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; pantropical.

Polycarpon Loefl.

Polycarpon prostratum (Forssk.) Asch. & Schweinf., Oesterr. Bot. Z. 39: 128 1889; B.D.Sharma et al., Fl. Karnataka Anal. 16. 1984.

Pharnaceum depressum L., Mant. Pl. 2: 564 1771

Alsine prostrata Forssk., Fl. Aegypt.-Arab. 207 1775

Polycarpon loeflingii Wight & Arn. ex Benth., Gen. Pl. 1: 153 1862

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dakshina Kannada, Hassan, Kalburgi, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Uttara Kannada), throughout India; pantropical.

Stellaria L.

Stellaria media (L.) Vill., Hist. Pl. Dauphiné 3: 615 1789; B.D.Sharma et al., Fl. Karnataka Anal. 16. 1984.

Alsine media L., Sp. Pl. 272 1753

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Kodagu, Mysuru, Uttara Kannada), throughout India; pantropical.

Vaccaria Wolf

Vaccaria hispanica (Mill.) Rauschert, Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg, Math.-Naturwiss. Reihe 14: 496 1965

Saponaria vaccaria L., Sp. Pl. 409 1753; C.J.Saldanha, Fl. Karnataka 1: 155. 1984

Saponaria hispanica Mill., Gard. Dict. ed. 8 Errat. 1768

Distribution: Karnataka (cultivated in gardens), cultivated in gardens in India; native to Eurasia.

Local/Common/Vernacular names: *Cow basil, cow soapwort, Cowcockle, Cowherb, prairie carnation* (English)

Casurinaceae

Casuarina L.

Casuarina equisetifolia L., Amoen. Acad. 4: 143.1759; B.D.Sharma et al., Fl. Karnataka Anal. 263. 1984.

Casuarina indica Pers., Syn. Pl. 2: 531.1807

Casuarina litorea L. ex Fosberg & Sachet, Smithsonian Contr. Bot. 24: 4, 1975

Distribution: Karnataka (cultivated throughout), Andaman & Nicobar Islands, throughout coastal areas; Australia, Malaysia, Pacific Islands, S & SE Asia.

Local/Common/Vernacular name: *She oak* (English); *Sarve mara* (Kannada)

Celastraceae

Cassine L.

Cassine glauca (Rottb.) Kuntze, Rev. Gen. Pl. 1: 114. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Mangifera glauca Rottb., Nye Kongel. Norske Vidensk. Selsk. Skr. 2: 534. 1783

Elaeodendron glaucum (Rottb.) Pers., Syn. Pl. 1: 241 1805

Elaeodendron roxburghii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 157. 1834

Cassine albens (Retz.) Kosterm., Gard. Bull. Singapore 39: 178 1986

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Bihar, Himachal Pradesh, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal; Pakistan, Malaysia, Sri Lanka.

Local/Common/Vernacular names: *Mooguthi mara, Mookarike* (Kannada)

Cassine paniculata (Wight & Arn.) Lobb.-Callen, Adansonia ser. 2. 15: 220. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Elaeodendron paniculatum Wight & Arn., Prodr. Fl. Ind. Orient. 1: 157. 1834

Distribution: Karnataka (Belagavi, Davanagere, Hassan, Kodagu, Mysuru, Tumakuru), Kerala, Maharashtra, Tamil Nadu; Indo-Malaysia.

Catha Forssk. ex Scop.

Catha edulis (Vahl) Endl., Ench. Bot. 575 1841; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Distribution: Karnataka (cultivated in Bengaluru), cultivated in Maharashtra; native to Africa.

Celastrus L.

Celastrus paniculatus Willd., Sp. Pl. 1: 1125. 1797; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Celastrus rothianus Schult. in Roemer & Schultes, Syst. 5: 423. 1820 non Wight & Arn.

Ceanothus paniculatus Roth, Nov. Pl. Sp. 154 1821

Celastrus multiflorus Roxb., Fl. Ind. 2: 389. 1824

subsp. paniculatus

Distribution: Karnataka (throughout), throughout India; Australia, Bhutan, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Nepal, Pacific islands, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Black oil plant, Climbing staff tree, Intellect tree* (English); *Gangunde kayi, Jothishmati, Karigonne* (Kannada)

subsp. aggregatus K.T.Mathew ex K.M.Matthew, Kew Bull. 46: 540. 1991; Hegde & Hegde, J. Threat. Taxa 3: 1731. 2011

Distribution: Karnataka (Belagavi, Chamarajanagar, Chitradurga, Hassan, Kodagu, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Goa, Kerala, Maharashtra; endemic.

Euonymus L.

Euonymus angulatus Wight, Icon. Pl. Ind. Orient. 3: t. 1053. 1846; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Euonymus acutangulus Wight, Ill. 1: 178. 1840

Distribution: Karnataka (Chikkamagaluru, Kodagu), Kerala, Tamil Nadu; endemic.

Euonymus dichotomus B. Heyne ex Wall., Fl. Ind. 2: 410. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Euonymus indicus Heyne ex Wall., Fl. Ind. 2: 409. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 49. 1984.

Euonymus goughii Wight, Icon. Pl. Ind. Orient. 1: t. 215. 1839

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Maharashtra, Tamil Nadu; endemic.

Glyptopetalum Thwaites

Glyptopetalum grandiflorum Bedd., Icon. Pl. Ind. Or. 1: 21. 1871

Distribution: Karnataka (Kodagu), Kerala; endemic.

Gymnosporia (Wight & Arn.) Benth. & Hook. f.

Gymnosporia buchananii Loes., Bot. Jahrb. Syst. 28: 153 1900

Gymnosporia ovata (Wall. ex Wight & Arn.) M.A.Lawson, Fl. Brit. India 1: 619 1875

Maytenus ovata (Wall. ex Wight & Arn.) Loes., Nat. Pflanzenfam. 20B: 140 1942

Distribution: Karnataka (Chitradurga), peninsular India; tropical Africa.

Gymnosporia emarginata (Willd.) Thwaites, Enum. 409. 1864; C.J.Saldanha, Fl. Karnataka 2: 96. 1996.
Celastrus emarginatus Willd., Sp. Pl. 1: 1128. 1798
Celastrus montanus Roth, Roemer & Schultes, Syst. 5: 427. 1819
Gymnosporia montana (Roth) Benth., Fl. Austral. 1: 400. 1863
Maytenus emarginata (Willd.) Ding Hou, Steenis, Fl. Malesiana ser. 1, 6: 241. 1962
 Distribution: Karnataka (Ballari, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Tamil Nadu; Australia, SE Asia, Sri Lanka, Taiwan.
 Local/Common/Vernacular names: *Thorny staff tree* (English); *Tanasimale, Tandrasa* (Kannada)

Gymnosporia heyneana Wight & Arn., Prodr. Fl. Ind. Orient. 1: 159. 1834; C.J.Saldanha, Fl. Karnataka 2: 97. 1996
Celastrus heyneana Roth in Roem. & Schult., Syst. Veg. 5: 421. 1819, p.p.
Maytenus heyneana (Roth) D.C.S. Raju & Babu, Bull. Bot. Surv. India 10: 348. 1968
 Distribution: Karnataka (Chikkamagaluru, Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Gymnosporia senegalensis (Lam.) Loes., Bot. Jahrb. Syst. 17: 541 1893
Maytenus senegalensis (Lam.) Exell, Bol. Soc. Brot. II, 26: 223 1952; Seeth. et al., Fl. 49. 2000
 Distribution: Karnataka (Chitradurga, Dharwad, Kalburgi); tropical and subtropical Africa including Madagascar, Aldabra and Asia.
 Local/Common/Vernacular name: *Confetti tree* (English); *Tondarsi* (Kannada)

Gymnosporia wallichiana M.A.Lawson, Fl. Brit. India 1: 621 1875; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.
Maytenus wallichiana (Spreng.) D.C.S.Raju & Babu, Bull. Bot. Surv. India 10: 349 1969
 Distribution: Karnataka (Mysuru), Punjab, Tamil Nadu, West Bengal; Bangladesh, Pakistan.

Gymnosporia wightiana (Babu) R.S. Rao, Fl. Goa 1: 76 1985
Celastrus rothianus Wight & Arn., Prodr. Fl. Ind. Orient. 1: 159. 1834.
Celastrus serrulatus Roth, Nov. Pl. Sp. 156. 1821
Catha rothiana Walp., Rep. 1: 532. 1842
Gymnosporia rothiana (Walp.) M. A. Lawson, Hook.f., Fl. Brit India 1: 620. 1875
Maytenus wightiana Babu, Bull. Bot. Surv. India 10: 349. 1969
Maytenus rothiana (Walp.) Lobreau-Callen, Adansonia ser 2: 15: 233. 1975; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh; endemic.

Loeseneriella A.C. Sm.

Loeseneriella africana (Willd.) R.Wilczek, Fl. Congo Belge & Ruanda-Burundi 9: 154. 1960; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.
Tonsella africana Willd., Sp. Pl. 1: 194 1797
Hippocratea bourdillonii Gamble, Bull. Misc. Inform. Kew 1916: 132 1916
Hippocratea africana (Willd.) Loes. ex Engl., Veg. Erde 9(III 2): 240 1921
Loeseneriella bourdillonii (Gamble) Ramamoorthy in Saldanha & Nicolson, Fl. Hassan 315. 1976
 Distribution: Karnataka (Belagavi, Dharwad, Hassan, Mysuru, Shivamogga, Uttara Kannada); E Africa to S India

Loeseneriella arnottiana (Wight) A.C. Sm. in J. Arnold Arbor. 16: 174. 1945; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.
Hippocratea arnottiana Wight, Illustr. 1: 133, t. 46. f. 47. 1840
 Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; endemic.
 Local/Common/Vernacular name: *Matre balli* (Kannada)

Lophopetalum Wight ex Arn.

Lophopetalum wightianum Arn., Ann. Nat. Hist. 3: 151. 1839; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.

Lophopetalum fimbriatum Wight, Ill. 1: 178. 1840

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Arunachal Pradesh, Assam, Kerala, Maharashtra, Meghalaya, Tamil Nadu, Uttar Pradesh; Indo-Malaya.

Local/Common/Vernacular names: *Spindle tree* (English); *Banate, Boralu paale* (Kannada)

Maytenus Molina

Maytenus puberula (M.A.Lawson) Loesener, Engl. & Prantl., Pflanzenfam. 20: 136. 1942; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.

Gymnosporia puberula M. A. Lawson, Hook.f., Fl. Brit. India 1: 619. 1875

Distribution: Karnataka (Uttara Kannada), Maharashtra, Uttar Pradesh; endemic.

Microtropis Wall. ex Meisn.

Microtropis latifolia Wight ex M.A.Lawson, Fl. Brit. India 1: 613 1875; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Shivamogga), Kerala, Maharashtra, Tamil Nadu; endemic.

Microtropis stocksii Gamble, Bull. Misc. Inform. Kew 1916: 132 1916; B.D.Sharma et al., Fl. Karnataka Anal. 50. 1984.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Microtropis wallichiana Wight ex Thwaites, Enum. Pl. Zeyl. 71. 1858; C.J.Saldanha, Fl. Karnataka 2: 98. 1996

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Parnassia L.

Parnassia mysorensis H.Heyne ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 35. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 98. 1984.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Mysuru), Kerala, Sikkim, Tamil Nadu; Sikkim; China.

Pleurostyla Wight & Arn.

Pleurostyla opposita (Wall.) Alston in Trimen, Handb. Fl. Ceylon 6: (Suppl.) 48. 1931; C.J.Saldanha, Fl. Karnataka 2: 98. 1996

Celastrus opposita Wall., Roxb., Fl. Ind. 2: 398. 1824

Pleurostyla heynei Wight & Arn., Prodr. Fl. Ind. Orient. 1: 157. 1834

Pleurostyla wightii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 157. 1834

Distribution: Karnataka (Dakshina Kannada), Andhra Pradesh, Kerala, Tamil Nadu; India to Malaysia, N Australia, Sri Lanka.

Reissantia N. Halle

Reissantia grahamii (Wight) Ding Hou, Blumea 12: 33. 1963; B.D.Sharma et al., Fl. Karnataka Anal. 51. 1984.

Hippocratea grahamii Wight, Illustr. 1: 134. 1840

Pristimera grahamii (Wight) A.C. Sm. in J. Arnold Arbor. 26: 178. 1945

Distribution: Karnataka (Dharwad, Shivamogga, Uttara Kannada), Kerala; SE Asia.

Reissantia indica (Willd.) N.Hallé in Mem. Inst. Franc. Afrique Noire 64: 85. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 51. 1984.

Hippocratea indica Willd., Sp. Pl. 1: 193. 1797

Pristimera indica (Willd.) A.C.Sm., Amer. J. Bot. 28: 440. 1941

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Kerala, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Tripura, West Bengal; paleotropical.

Reissantia sessiliflora N.V.Page, Srivastav & R.Rao, Phytotaxa 297(2): 209. 2017

Distribution: Karnataka (Uttara Kannada); endemic.

Salacia L.

Salacia beddomei Gamble, Bull. Misc. Inform. Kew 1916: 133 1916

Distribution: Karnataka (Chikkamagaluru, Shivamogga), Kerala, Tamil Nadu; endemic.

Salacia chinensis L., Mant. Pl. 2: 293. 1771; C.J.Saldanha, Fl. Karnataka 2: 91. 1996

Salacia prinoides (Willd.) DC., Prodr. 1: 579. 1824

Salacia latifolia Wall. ex Lawson, Hook.f., Fl. Brit. India 1: 629. 1875

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Goa, Kerala, Maharashtra, Meghalaya, Odisha, Tamil Nadu, West Bengal; Cambodia, China, Laos, Myanmar, Sri Lanka, Thailand, Vietnam.

Salacia fruticosa Heyne ex Lawson, Hook.f., Fl. Brit. India 1: 628. 1875; C.J.Saldanha, Fl. Karnataka 2: 91. 1996

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala, Tamil Nadu; endemic.

Salacia gambleana Whiting & Kaul, Bull. Misc. Inform. Kew 1940: 300 1941

Salacia talbotii Gamble, Bull. Misc. Inf. Kew 133. 1916; B.D.Sharma et al., Fl. Karnataka Anal. 51. 1984.

Distribution: Karnataka (Shivamogga, Uttara Kannada), Maharashtra; endemic.

Salacia macroisperma Wight, Icon. Pl. Ind. Orient. 3: t. 962. 1845; C.J.Saldanha, Fl. Karnataka 2: 91. 1996

Distribution: Karnataka (Belagavi, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Balige balli* (Kannada)

Salacia malabarica Gamble, Bull. Misc. Inform. Kew 1916: 133. 1916; C.J.Saldanha, Fl. Karnataka 2: 91. 1996

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala, Tamil Nadu; endemic.

Salacia oblonga Wall. ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 106. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 51. 1984.

Distribution: Karnataka (Belagavi, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Salacia reticulata Wight, Ill. Ind. Bot. 1: 134. 1840; C.J.Saldanha, Fl. Karnataka 2: 91. 1996.

Distribution: Karnataka (Kodagu), Andaman & Nicobar Islands, Kerala, Tamil Nadu; Sri Lanka.

Ceratophyllaceae

Ceratophyllum L.

Ceratophyllum demersum L., Sp. Pl. 992. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 263. 1984.

Dichotophyllum demersum (L.) Moench, Methodus 345. 1794

Ceratophyllum verticillatum Roxb., Fl. Ind. ed. 1832, 3: 624. 1832

Distribution: Karnataka (Chitradurga, Davanagere, Dharwad, Hassan, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; cosmopolitan.

Local/Common/Vernacular name: *Honal* (Kannada)

Cleomaceae

Cleome L.

Cleome angustifolia Forssk., Fl. Aegypt.-Arab. 120 1775; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Cleome tenella L.f., Suppl. Pl. 300. 1782

Distribution: Karnataka (Chitradurga, Vijayapura), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Africa, Arabia.

Cleome aspera J.Koenig ex DC., Prodr. 1: 241. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dharwad, Mysuru, Vijayapura), Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Odisha, Tamil Nadu; India to Malesia

Cleome gynandra L., Sp. Pl. 671. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Gynandropsis pentaphylla (L.) DC., Prodr. 1: 238. 1824

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Kalburgi, Kodagu, Kolar, Mysuru, Uttara Kannada), throughout India; pantropical.

Local/Common/Vernacular names: *Ajaganda, Sirikalagida* (Kannada)

Cleome monophylla L., Sp. Pl. 672. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Uttara Kannada, Vijayapura), Andhra Pradesh, Bihar, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Sikkim, Tamil Nadu, Uttar Pradesh, West Bengal; Tropical Africa, India

Local/Common/Vernacular name: *Menasinakase* (Kannada)

Cleome rutidosperma DC., Prodr. 1: 241. 1824; K.G.Bhat, J. Bombay Nat. Hist. Soc., 90: 138. 1993

Distribution: Karnataka (Udupi), Kerala; Asia, Australia.

Cleome simplicifolia Hook.f. & Thomson, Fl. Brit. India 1: 169. 1872; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Distribution: Karnataka (Belagavi, Bidar, Kalburgi, Uttara Kannada), Andhra Pradesh, Goa, Gujarat, Madhya Pradesh, Maharashtra, Rajasthan, Uttar Pradesh; India

Corynandra Schrad. ex Spreng.

Corynandra chelidonii (L.f.) Cochrane & Iltis, Novon 23(1): 22. 2014

Cleome chelidonii L.f., Suppl. Pl. 300. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Polanisia chelidonii (L.f.) DC., Prodr. 1: 242. 1824

Cleome chelidonii var. *pallae* C.S.Reddy & V.S.Raju, J. Econ. Taxon. Bot. 25: 217. 2001

Distribution: Karnataka (Belagavi, Dharwad, Kalburgi, Mysuru, Uttara Kannada), Andhra Pradesh, Bihar, Goa, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu; Sri Lanka.

Corynandra felina (L.f.) Cochrane & Iltis, Novon 23(1): 23. 2014

Cleome felina L.f., Suppl. Pl. 300. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Dharwad, Raichur, Vijayapura), Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Puducherry, Tamil Nadu; Sri Lanka.

Corynandra viscosa (L.) Cochrane & Iltis, Novon 23(1): 24. 2014

Cleome viscosa L., Sp. Pl. 672. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 9. 1984.

Distribution: Karnataka (throughout), throughout India; pantropical.

Local/Common/Vernacular names: *Kadusasive, Nayisasuve* (Kannada)

Clusiaceae

Clusia L.

Clusia rosea Jacq., Enum. Syst. Pl. 34 1760; C.J.Saldanha, Fl. Karnataka 1: 211. 1984

Distribution: Karnataka (cultivated in gardens), cultivated in India; native to the Caribbean, Islands, Puerto Rico, USA(Florida).

Local/Common/Vernacular names: *Balsam apple, Copey, Pitch-apple, Scotch attorney, The autograph tree* (English)

Garcinia L.

Garcinia gummi-gutta (L.) N.Robson, Brittonia xx. 103. 1968; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Cambogia gummi-gutta L., Gen. Pl. ed. 5 522 1754

Mangostana cambogia Gaertn., Fruct. Sem. Pl. 2: 106 1790

Garcinia cambogia (Gaertn.) Desr., Encycl. 3: 701 1792

var. **gummi-gutta**

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Indonesia, Sri Lanka.

Local/Common/Vernacular names: *Brindleberry, Malabar tamarind, Pot tamarind* (English); *Muruganahuli* (Kannada); *Kodampuli* (Kodava)

var. **conicarpa** (Wight) N.P.Singh in Fl. India 3: 110. 1993

Garcinia conicarpa Wight, Icon. Pl. Ind. Orient. 1: t. 121. 1839; Kesh.Murthy & Yogan, Fl. Kodagu 59. 1990

Garcinia cambogia var. *conicarpa* (Wight) Anderson in Hook.f., Fl. Brit. India 1: 262 . 1874

Garcinia darwiniana Kesh.Murthy, Yogan. & K.Vasudev, Curr. Sci. 56: 425. 1987

Distribution: Karnataka (Kodagu, Shivamogga), Kerala; endemic.

Garcinia indica (Thouars) Choisy, Prodr. 1: 561 1824; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Plate 20G

Brindonia indica Thouars, Dict. Sci. Nat. 5: 340 1817

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Brindonia Tallow Tree, Goa-Butter, Indian Berry, Indian Gamboge, Indian Tallow Tree, Kokam Butter, Kokam butter tree, Mangosteen oil tree, Wild mangosteen* (English); *Dhupada mara, Kokam mara, Murgala, Murugina huli, Punarpuli, Tittidika* (Kannada); *kodam puli* (Kodava); *Bhirand* (Konkani)

Garcinia livingstonei T.Anderson, J. Linn. Soc., Bot. 9: 263 1867; C.J.Saldanha, Fl. Karnataka 1: 207. 1984

Distribution: Karnataka (introduced in Bengaluru), native to Africa.

Garcinia madruno (Kunth) Hammel., Ann. Missouri Bot. Gard. 76: 928 1989

Calophyllum madruno Kunth, Nov. Gen. Sp. 5: 202 1822

Rheedia madruno (Kunth) Planch. & Triana, Ann. Sci. Nat., Bot. IV, 14: 315 1860; C.J.Saldanha, Fl. Karnataka 1: 211. 1984

Distribution: Karnataka (cultivated in Bengaluru); native to C & S America.

Garcinia morella (Gaertn.) Desr., Encycl. 3: 701 1792; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Mangostana morella Gaertn., Fruct. Sem. Pl. 2: 106 1790

Garcinia pictoria Roxb., Fl. Ind. 2: 627. 1832

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Assam, Kerala, Maharashtra, Meghalaya, Tamil Nadu, West Bengal; Sri Lanka, Malesia

Local/Common/Vernacular names: *Ardala*, *Arisina guruge*, *Devana huli*, *Jirige huli*, *Murina huli*, *Ponpuli* (Kannada)

Garcinia spicata Hook.f. in J. Linn. Soc., Bot. 14: 486 1875; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Xanthochymus spicatus Wight & Arn., Prodr. Fl. Ind. Orient. 102 1834

Xanthochymus ovalifolius Bedd., Fl. Sylv. S. India xxi 1871

Garcinia ovalifolia (Roxb.) Hook.f. ex T.Anderson, Fl. Brit. India 1: 269 1874

Distribution: Karnataka (Belagavi, Dakshina Kannada, Shivamogga, Uttara Kannada), Andhra Pradesh, Assam, Kerala, Maharashtra, Meghalaya, Odisha, Tamil Nadu; Sri Lanka

Garcinia talbotii Raizada ex Santapau, Rec. Bot. Surv. India 16: 14 1960; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Plate 20H

Garcinia ovalifolius var. *macrantha* Hook.f. ex T.Anderson, Fl. Brit. India 1: 269.1874

Garcinia malabarica Talbot, J. Bombay Nat. Hist. Soc. 11: 234 1897

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Kanjeerakayi* (Kannada)

Garcinia xanthochymus Hook.f. ex T.Anderson, Fl. Brit. India 1: 269 1874; B.D.Sharma et al., Fl. Karnataka Anal. 18. 1984.

Garcinia tinctoria (DC.) Dunn, Fl. Madras 74 1915

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India; Bangladesh, Bhutan, Cambodia, China, Laos, Myanmar, Nepal, Thailand, Vietnam; cultivated elsewhere.

Colchicaceae

Disporum Salisb. ex D.Don

Disporum leschenaultianum D.Don, Proc. Linn. Soc. London 1: 45.1839; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Disporum leschenaultianum var. *angustifolium* C.E.C.Fisch., Fl. Madras 1522.1928

Distribution: Karnataka (Uttara Kannada)

Gloriosa L.

Gloriosa superba L., Sp. Pl. 305.1753; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Distribution: Karnataka (throughout), throughout tropical areas of India; paleotropical.

Local/Common/Vernacular names: *Cuntry aconite*, *Flame flower*, *Hary's plant*, *Malabar glory lily*, *Serpent's tounge*, *Superb lily* (English); *Agnishika*, *Akka thangi balli*, *Bali papu*, *Benne bidiru*, *Chauati hoovu*, *Gowri huvu*, *Huli uguru*, *Indrana gedde*, *Jokumarana balli*, *Kardi kannina gedde*, *Kenchan daapu*, *Koli kutumana gedde*, *Langulika*, *Naga leekana gedde*, *Nagulika*, *Onapou*, *Shivaraktha balli*, *Slvashakthi balli*, *Umbalagji gedde*, *Vana hoovu* (Kannada); *Agnimukhi*, *Agnishiksha*, *Ahijiva*, *Amrita*, *Annata*, *Dipta*, *Garbhanuta*, *Garbha pathani*, *Garbha gathini*, *hali*, *Halini*, *Haripriya*, *Indrpuspypika*, *Kalihari*, *Kandali*, *Longalika*, *Longuli*, *Nakta*, *Pushpa saurabha*, *Shakara pushpi*, *Sikhaajihva*, *Suvarna pushpa*, *Vanhivatra*, *Vidutajwala*, *Vishalya*, *Vranahrita* (Sanskrit); *Bali papu*, *Cauti puta gedde*, *Gonal thappu*, *Ken kannada pu* (Tulu)

Iphigenia Kunth

Iphigenia indica (L.) A.Gray ex Kunth, Enum. Pl. 1: 213.1833; B.D.Sharma et al., Fl. Karnataka Anal. 286. 1984.

Melanthium indicum L., Mant. Pl. 226.1771

Anguillaria indica (L.) R.Br., Prodr. Fl. Nov. Holl. 273.1810

Notocles indica (L.) Salisb., Gen. Pl. 54.1866

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dakshina Kannada, Hassan, Kalburgi, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Vijayapura), Peninsular India; S & SE Asia.

Local/Common/Vernacular names: *Kadu belluli, Neeru bale* (Kannada)

Iphigenia magnifica Ansari & R.S.Rao, Bull. Bot. Surv. India 20: 162.1979

Plate 21A

Distribution: Karnataka (Belagavi, Uttara Kannada), Maharashtra, Tamil Nadu; endemic.

Iphigenia mysorensis Arekal & S.N.Ramaswamy, Bot. Not. 125: 220.1972; B.D.Sharma et al., Fl. Karnataka Anal. 287. 1984.

Plate 21C

Distribution: Karnataka (Hassan, Mysuru, Tumakuru); endemic.

Iphigenia pallida Baker, J. Linn. Soc., Bot. 17: 451.1879; B.D.Sharma et al., Fl. Karnataka Anal. 287. 1984.

Plate 21B

Distribution: Karnataka (Belagavi, Dharwad, Kalburgi, Hassan, Shivamogga, Udupi, Uttara Kannada, Yadgir), SW India; endemic.

Iphigenia sahyadrica Ansari & R.S.Rao, Bull. Bot. Surv. India 20: 163.1979; B.D.Sharma et al., Fl. Karnataka Anal. 278. 1984.

Distribution: Karnataka (Shivamogga, Uttara Kannada)

Iphigenia stellata Blatt., J. Bombay Nat. Hist. Soc. 32: 734.1928

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Combretaceae

Anogeissus (DC.) Wall. ex Guill. & Perr.

Anogeissus latifolia (Roxb. ex DC.) Wall. ex Guillem. & Perr., Fl. Seneg. Tent. 7: 280. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 100. 1984.

Plate 5E

Conocarpus latifolius Roxb. ex DC., Prodr. 3: 16. 1828

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Bihar, Goa, Gujarat, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, Uttaranchal; Myanmar, Sri Lanka

Local/Common/Vernacular names: *Dindalu, Dindiga mara* (Kannada)

Bruguiera Savigny

Bruguiera cylindrica (L.) Blume, Enum. Pl. Jav. 1: 93. 1827; C.J.Saldanha, Fl. Karnataka 2: 58. 1996

Rhizophora cylindrica L., Sp. Pl. 443. 1753

Rhizophora caryophylloides Burm.f., Fl. Indica 109. 1768

Bruguiera caryophylloides (Burm.f.) Blume, Enum. Pl. Jav. 1: 93. 1827

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Maharashtra, Odisha, Tamil Nadu; Australia, China, Indonesia, Malaysia, Myanmar, New Guinea, Pacific islands Philippines, Sri Lanka, Thailand, Vietnam.

Bruguiera gymnorhiza (L.) Lam., Encycl. 4: 696. 1797; B.D.Sharma et al., Fl. Karnataka Anal. 100. 1984.

Rhizophora gymnorhiza L., Sp. Pl. 447. 1753

Bruguiera rheedei Blume, Enum. Pl. Jav. 1: 92. 1827

Bruguiera conjugata (L.) Merr. in Philipp. J. Sci. 9: 118. 1914

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada); paleotropical

Bruguiera parviflora (Roxb.) Wight & Arn. ex Griff., Trans. Med. Phys. Soc. Calc. 8: 10. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 100. 1984.

Rhizophora parviflora Roxb., Fl. Ind. 2: 461. 1832

Distribution: Karnataka (Uttara Kannada), Andaman & Nicobar Islands, Maharashtra, Odisha, West Bengal; Australia, Melanesia, SE Asia.

Combretum Loefl.

Combretum coccineum (Sonn.) Lam., Encycl. 1: 734. 1785; C.J.Saldanha, Fl. Karnataka 2: 48. 1996.

Cristaria coccinea Sonn., Voy. Indes Orient. 3: 284. 1782

Poivrea coccinea (Sonn.) Thouars, Mélang. Bot. 28. 1811

Grislea coccinea (Sonn.) Britton., Sci. Surv. Porto Rico & Virgin Islands 6: 25. 1925

Distribution: Karnataka (cultivated), cultivated throughout plains; native to Madagascar and Mauritius.

Local/Common/Vernacular names: *Flame wine, Scarlet comb* (English)

Combretum indicum (L.) DeFilipps, Useful Pl. Dominica 277 1998

Quisqualis indica L., Sp. Pl. ed. 2:556. 1762; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Distribution: Karnataka (cultivated Belagavi, Bengaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kodagu, Mysuru); native to Myanmar, cultivated and naturalised throughout paleotropics.

Local/Common/Vernacular names: *Chinese honeysuckle, Rangoon Creeper* (English); *Bombya mallige* (Kannada)

Combretum latifolium Blume, Bijdr. 641. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Combretum extensum Roxb. ex G. Don, Trans. Linn. Soc. London 15: 414. 1827

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Goa, Kerala, Maharashtra, Meghalaya, Mizoram, Tamil Nadu, West Bengal; Indo-China, Indonesia, Malaysia, Myanmar, New Guinea, Sri Lanka, Thailand.

Combretum malabaricum (Bedd.) Sujana, Ratheesh & Anil Kumar, J. Bot. Res. Inst. Texas 6(2): 453. 2012

Quisqualis malabarica Bedd., Icon. Pl. Ind. Or., 33, t. 155. 1874; K.G.Bhat, J. Bombay Nat. Hist. Soc., 99: 566. 2002

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala; endemic.

Combretum ovalifolium Roxb., Fl. Ind. 2: 226. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Combretum albidum G.Don in Trans. Linn. Soc. London 15: 429 1827

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkaballapur, Chitradurga, Dakshina Kannada, Dharwad, Kalburgi, Kolar, Mandya, Mysuru, Raichur, Shivamogga), Andhra Pradesh, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Manipur, Rajasthan, Tamil Nadu; India, Sri Lanka.

Local/Common/Vernacular name: *Edatiga* (Kannada)

Combretum razianum K. G. Bhat, Indian J. Forest. 28(2):210.2005; Sujana & Pramanik, Ann. Pl. Sci. 6(3): 1594.2017.

Combretum pilosum subsp. *razianum* (K.G.Bhat) Chakrab. & P. G. Diwakar, Indian J. Forest., 34 (4):483.2011

Distribution: Karnataka (Dakshina Kannada, Kerala); endemic.

Combretum shivannae Gholave, Kambale, Lekhak & S.R.Yadav, Kew Bull. 70:33.2015

Distribution: Karnataka (Uttara Kannada); endemic.

Getonia Roxb.**Getonia floribunda** Roxb., Fl. Corom. t.87. 1798; C.J.Saldanha, Fl. Karnataka 2: 47. 1996.**Plate 5D***Calycopteris floribunda* Lam., Tabl. Encycl. 2: 485. t. 357. 1793

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; SE Asia.

Local/Common/Vernacular names: *Baguli, Marasuthuballi* (Kannada)**Lumnitzera Willd.****Lumnitzera racemosa** Willd. in Neue Schriften Ges. Naturf. Freunde Westphalena 4: 187. 1803; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Goa, Maharashtra, Odisha, Tamil Nadu; paleotropical

Terminalia L.**Terminalia alata** Roth, Nov. Pl. Sp. 379. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.*Terminalia elliptica* Willd., Sp. Pl. 4: 969 1806*Pentaptera tomentosa* Roxb. ex DC., Prodr. 3: 14. 1828*Terminalia tomentosa* (Roxb. ex DC.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 314. 1834

Distribution: Karnataka (Ballari, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Tumakuru), throughout India; Myanmar, Indo-China, Sri Lanka, Thailand.

Local/Common/Vernacular name: *Karimatti* (Kannada)**Terminalia arjuna** (Roxb. ex DC.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 314. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.*Terminalia cuneata* Roth, Nov. Pl. Sp. 380 1821*Pentaptera arjuna* Roxb. ex DC., Prodr. 3: 14. 1828

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Kalburgi, Mysuru, Tumakuru, Uttara Kannada), throughout India; Sri Lanka

Local/Common/Vernacular names: *Bilimatti, Holematti* (Kannada)**Terminalia bellirica** (Gaertn.) Roxb., Pl. Coromandel 2:55, t. 198. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.*Myrobalanus bellirica* Gaertn., Fruct. 2: 90. t. 97. ff. a-d. 1791*Terminalia gella* Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 227. 1851; B.D.Sharma et al., Fl. Karnataka Anal. 102. 1984.

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Gujarat, Kerala, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh; Bangladesh, Indonesia, Indo-China, Malaysia, Myanmar, Philippines, Thailand.

Local/Common/Vernacular names: *Bahera, Bastard myrobalan, Beleric* (English); *Shantimara, Taremara* (Kannada)**Terminalia catappa** L., Syst. Nat. (ed.12) 2: 674. 1767; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Kalburgi, Kodagu, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Mizoram, Rajasthan, Tamil Nadu, Uttar Pradesh, also cultivated; throughout coastal Australia, Asia & Polynesia.

Local/Common/Vernacular names: *Bengal almond, Country almond, Indian almond, Malabar almond, Sea almond, tropical almond* (English)

Terminalia chebula Retz., Obs. Bot. 5: 31. 1789; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Myrobalanus chebula (Retz.) Gaertn., Fruct. 2: 91. t. 97. 1791

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; Bangladesh, Bhutan, Cambodia, India, Laos, Malaysia, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Chebulic myrobalan*, *yellow myrobalan* (English); *Alale kayi mara* (Kannada)

Terminalia crenulata Roth, Nov. Pl. Sp. 380. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 101. 1984.

Terminalia tomentosa var. *crenulata* (Roth) C.B.Clarke in Hook.f., Fl. Brit. India 2: 448. 1878

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Assam, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, West Bengal; Bangladesh, Cambodia, Laos, Nepal, Myanmar, Thailand, Vietnam.

Local/Common/Vernacular names: *Karimatti*, *Kirajul mathi* (Kannada)

Terminalia paniculata Roth, Nov. Pl. Sp. 383. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 102. 1984.

Pentaptera paniculata Roxb., Fl. Ind. 2: 442 1832

Distribution: Karnataka (throughout), Andhra Pradesh, Goa, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Flowering Murdah* (English); *Hanalu*, *Honagal*, *Honal*, *Honalu*, *Hongal*, *Hulive*, *Hulube*, *Huluvemara*, *Hunagalu*, *Kindal*, *Maruva*, *Mathi*, *Matti*, *Ulvi* (Kannada)

Commelinaceae

Belosynapsis Hassk.

Belosynapsis vivipara (Dalzell) C.E.C.Fisch., Bull. Misc. Inform. Kew 1928: 254. 1928; B.D.Sharma et al., Fl. Karnataka Anal. 288. 1984.

Cyanotis vivipara Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 226. 1851

Dalzellia vivipara (Dalzell) Hassk., Flora 48: 594. 1865

Tonningia vivipara (Dalzell) Kuntze, Revis. Gen. Pl. 2: 722. 1891

Distribution: Karnataka (Chikkamagaluru, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga), Kerala, Maharashtra, Tamil Nadu; endemic.

Commelina L.

Commelina attenuata K.D.Koenig ex Vahl, Enum. Pl. 2: 168. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 288. 1984.

Commelina rajmahlensis C.B.Clarke, J. Linn. Soc., Bot. 11: 444. 1871

Commelina repens Roxb. & H.Heyne ex C.B.Clarke in DC. & C.DC. Monogr. Phan. 3: 172 1881

Commelina clavata var. *attenuata* (K.D.Koenig ex Vahl) Trimen, Syst. Cat. Fl. Pl. Ceylon 95. 1885

Distribution: Karnataka (Bagalkot, Belagavi, Chitradurga, Davanagere, Mysuru, Raichur, Tumakuru, Vijayapura), Peninsular, N and NE India; Sri Lanka.

Commelina badamica Nandikar & Gurav, Telopea 18: 514. 2015

Distribution: Karnataka (Bagalkot); endemic.

Commelina benghalensis L., Sp. Pl. 41. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 288. 1984.

Distribution: Karnataka (throughout), throughout India; Paleotropical

Local/Common/Vernacular names: *Gubbachhi bale*, *Gubbachhi basale*, *Kanne soppu* (Kannada)

Commelina caroliniana Walter, Fl. Carol. 68.1788

Commelina hasskarlii C.B.Clarke, Commelyn. Cyrtandr. Bengal. t. 5.1874; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Distribution: Karnataka (Kalburgi, Mysuru, Shivamogga), throughout India; Bangladesh, Vietnam.

Commelina clavata C.B.Clarke, Commelyn. Cyrtandr. Bengal. t. 5.1874; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Commelina clavata var. *hohenackeri* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 171.1881

Commelina diffusa Zoll. ex C.B.Clarke, Monogr. Phan. 3: 171.1881., auct. non Burm. f., 1768.

var. **clavata**

Distribution: Karnataka (Belagavi, Bagalkot, Chamarajanagar, Chamarajanagar, Chikkamagaluru), Keral, Tamil Nadu; Indonesia, Sri Lanka

var. **clavatoides** Nandikar & Gurav, ined.

Commelina clavatoides Nampy & S. Joseph, J. Bot. Inst. Texas 6: 119. 2012

Distribution: Karnataka (Dakshina Kannada), Kerala; endemic.

Commelina coelestis Willd., Enum. Pl. 69.1809

Distribution: Karnataka (Belagavi), Tamil Nadu, Uttarakhand, West Bengal; native to Mexico.

Commelina diffusa Burm.f., Fl. Indica 18. 1768.

Commelina communis auct. non L., 1753; Dalzell & A.Gibson, Bombay Fl. 252.1861

Commelina nudiflora auct. non L. 1753; C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 144.1881

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular name: *Paradene* (Tulu)

Commelina ensifolia R.Br., Prodr. Fl. Nov. Holl. 269.1810; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Commelina ensifolia var. *pubescens* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 189.1881

Commelina lunata B.Heyne ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 188.1881

Commelina undulata var. *setosa* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 179.1881

Distribution: Karnataka (Bagalkot, Belagavi, Bengaluru, Ballari, Chamarajanagar, Davanagere, Hassan, Kodagu, Kolar, Tumakuru, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; Australia, New Guinea, SE Asia, Sri Lanka.

Commelina erecta L., Sp. Pl. 41.1753; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Davanagere, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), native to Mexico, Africa, Australia.

Commelina forsskalii Vahl, Enum. Pl. 2: 172.1805; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Distribution: Karnataka (Belagavi, Bagalkot, Chitradurga, Davanagere, Dharwad, Kalburgi, Mysuru, Yadgir), Goa, Gujarat, Maharashtra, Rajasthan, Tamil Nadu; Tropical Africa, Middle East.

Commelina hirsuta (Wight) C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 163.1881; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Heterocarpus hirsutus Wight, Icon. Pl. Ind. Orient. 6: 29.1853

Commelina nilagirica Steud. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 163.1881

Distribution: Karnataka (Belagavi), Tamil Nadu; endemic.

Commelina indehisca E.Barnes, J. Bombay Nat. Hist. Soc. 46: 74.1946; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Distribution: Karnataka (Chamarajanagar, Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Commelina longifolia Lam., Tabl. Encycl. 1: 129.1791; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Commelina salicifolia Roxb., Fl. Ind. 1: 176.1820

Distribution: Karnataka (Belagavi, Mysuru, Tumakuru), Andhra Pradesh, Kerala, Madya Pradesh, Maharashtra, Tamil Nadu; SE Asia.

Commelina maculata Edgew., Trans. Linn. Soc. London 20: 89.1851

Commelina obliqua var. *viscida* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 178.1881

Commelina paludosa var. *viscida* (C.B.Clarke) R.S.Rao & Kamathy, Bull. Bot. Surv. India 3: 168.1962

Distribution: Karnataka (Dakshina Kannada, Chitradurga, Dharwad, Mysuru, Udupi), throughout, China.

Commelina paleata Hassk., Pl. Jungh. 139.1852; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Trithyrocarpus paleatus (Hassk.) Hassk., Flora 49: 211.1866

Distribution: Karnataka (Belagavi, Chikkamagaluru, Kodagu, Mysuru, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu, Uttarakhand? ;Indonesia ?

Commelina paludosa Blume, Enum. Pl. Javae 1: 2.1827; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Commelina obliqua Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 45.1825

Distribution: Karnataka (Belagavi, Bidar, Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Raichur, Uttara Kannada), throughout India ; S E Asia , Sri Lanka.

Commelina subulata Roth, Nov. Pl. Sp. 23.1821; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Tradescantia biflora Willd. ex Schult. & Schult.f., Syst. Veg. 7: 1176.1830

Commelina violacea C.B.Clarke in Oliv., Fl. Trop. Afr. 8: 39.1901

Distribution: Karnataka (Bagalkote, Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Mysuru, Tumakuru), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Africa.

Commelina suffruticosa Blume, Catalogus 35.1823; B.D.Sharma et al., Fl. Karnataka Anal. 289. 1984.

Spathodithyros suffruticosus (Blume) Hassk., Commelin. Ind. 11.1870

Commelina rugulosa C.B.Clarke, J. Linn. Soc., Bot. 11: 446.1871

Commelina simsonii C.B.Clarke, J. Linn. Soc., Bot. 11: 446.1871

Distribution: Karnataka (Belagavi, Kodagu, Shivamogga, Uttara Kannada), throughout India; E & SE Asia, Nepal.

Commelina undulata R.Br., Prodr. Fl. Nov. Holl. 270.1810

Ananthopus undulatus (R.Br.) Raf., Fl. Tellur. 3: 70.1837

Commelina striata Edgew., Trans. Linn. Soc. London 20: 89.1851

Commelina longifolia Thwaites, Enum. Pl. Zeyl. 322. 1864.

Commelina kurzii C.B.Clarke, J. Linn. Soc., Bot. 11: 444.1871

Commelina obliqua var. *mathewii* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 878.1881

Distribution: Karnataka (Belagavi, Chamarajanagar, Uttara Kannada), throughout India;Bangladesh, Pakistan, Sri Lanka.

Cyanotis D.Don

Cyanotis adscendens Dalzell in Hooker's J. Bot. Kew Gard. Misc. 4: 343.1852; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Cyanotis sarmatica Wight, Icon. Pl. Ind. Orient. 6: 33, t. 2087.1863

Cyanotis sarmentosa Wight, Icon. Pl. Ind. Orient. 6: 33, t. 2087.1863

Cyanotis tuberosa var. *adscendens* (Dalzell) C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 249.1881

Distribution: Karnataka (Bagalkot, Belagavi, Bengaluru, Chikkamagaluru, Chamarajanagar, Dharwad, Mysuru, Shivamogga, Tumakuru), Andhra Pradesh, Assam, Bihar, Kerala, Maharashtra, West Bengal; Sri Lanka.

Cyanotis arachnoidea C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 250.1881; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Cyanotis pilosa Wight, Icon. Pl. Ind. Orient. t. 2083.1853

Cyanotis thwaitesii Hassk., Commelin. Ind. 136.1870

Cyanotis fasciculata var. *thwaitesii* (Hassk.) C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 254.1881

Tradescantia incana B.Heyne ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 250 1881

Cyanotis arachnoidea var. *obtusa* Trimen, J. Bot. 23: 266.1885

Tonningia arachnodea (C.B.Clarke) Kuntze, Revis. Gen. Pl. 2: 722.1891

Cyanotis obtusa (Trimen) Trimen, Handb. Fl. Ceylon 4: 812.1898

Cyanotis arachnoidea var. *thwaitesii* (Hassk.) R.S.Rao & Kammathy, Bull. Bot. Surv. India 6: 2.1964

Distribution: Karnataka (Belagavi Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Hassan, Kodagu, Mysuru, Tumakuru), Andhra Pradesh, Bihar, Maharashtra, Kerala, Odisha, Tamil Nadu; Sri Lanka. S & SE Asia.

Cyanotis arcotensis R.S.Rao, Blumea 14: 345.1967; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Cyanotis papilionacea auct. C.B.Clarke in DC., Monogr. Phan. 3: 246. 1881

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada?), Tamil Nadu; endemic.

Cyanotis axillaris (L.) D.Don ex Sweet, Hort. Brit. 430.1826

Commelina axillaris L., Sp. Pl. 42.1753

Tradescantia axillaris (L.) L., Mant. Pl. 2: 321.1771

Zygomenes axillaris (L.) Salisb., Trans. Hort. Soc. London 1: 271.1812

Tonningia axillaris (L.) Raf., Fl. Tellur. 2: 16.1837

Amischophacelus axillaris (L.) R.S.Rao & Kammathy, J. Linn. Soc., Bot. 59: 306.1966; B.D.Sharma et al., Fl. Karnataka Anal. 288. 1984.

Distribution: Karnataka (throughout), throughout India; Australia, S E Asia, West Indies.

Local/Common/Vernacular names: *Jgale, Nagalu kanne soppu* (Kannada)

Cyanotis burmanniana Wight, Icon. Pl. Ind. Orient. t. 2089.1853; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Cyanotis vaginata Wight, Icon. Pl. Ind. Orient. 6: 34, t. 2088.1853; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Cyanotis papilionacea var. *burmanniana* (Wight) C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 246. 1881

Tonningia vaginata (Wight) Kuntze, Revis. Gen. Pl. 2: 722.1891

Cyanotis papilionacea var. *vaginata* (Wight) C.E.C.Fisch., Fl. Madras 1549.1931

Distribution: Karnataka (Dakshina Kannada, Kodagu, Shivamogga, Udupi), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Cyanotis cristata (L.) D.Don, Prodr. Fl. Nepal. 46.1825; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Commelina cristata L., Sp. Pl. 42.1753

Tradescantia cristata (L.) L., Syst. Nat. ed. 12 2: 233.1767

Ephemerum cristatum (L.) Moench, Methodus 238.1794

Tradescantia imbricata Roxb., Fl. Ind. 2: 120.1824

Siphostigma cristata (L.) Raf., Fl. Tellur. 2: 16.1837

Cyanotis imbricata (Roxb.) Kunth, Enum. Pl. 4: 103.1843

Tonningia cristata (L.) Kuntze, Revis. Gen. Pl. 2: 722.1891

Distribution: Karnataka (Bagalkote, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; Bhutan, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Philippines, Sikkim, Sri Lanka, Thailand, Vietnam.

Cyanotis cucullata (Roth) Kunth, Enum. Pl. 4: 107.1843

Tradescantia cucullata Roth, Nov. Pl. Sp. 189.1821

Tradescantia glomerata B.Heyne ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 245.1881

Tonningia cucullata (Roth) Kuntze, Revis. Gen. Pl. 2: 722.1891

Zygomenes cucullata (Roth) R.S.Rao & Kammathy, Notes Roy. Bot. Gard. Edinburgh 25: 187.1964

Amischophacelus cucullata (Roth) R.S.Rao & Kammathy, J. Linn. Soc., Bot. 59: 306.1966; B.D.Sharma et al., Fl. Karnataka Anal. 288. 1984.

Cyanotis axillaris var. *cucullata* (Roth) Nandikar & Gurav, Taiwania 59: 297. 2014.

Distribution: Karnataka (Bagalkot, Ballari, Chitradurga, Hassan, Kalburgi, Kodagu, Mysuru, Tumakuru, Vijayapura), peninsular India, Rajasthan; endemic.

Cyanotis fasciculata (B.Heyne ex Roth) Schult. & Schult.f., Syst. Veg. 7: 1152 1830; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Tradescantia fasciculata B.Heyne ex Roth, Nov. Pl. Sp. 189.1821

Cyanotis decumbens Wight, Icon. Pl. Ind. Orient. 6: t. 2088.1853

Cyanotis lawiana Wight, Icon. Pl. Ind. Orient. 6: 33, t. 2086.1853

Cyanotis rosea Wight, Icon. Pl. Ind. Orient. 6: 33, t. 2086.1853

Cyanotis karliana Hassk., Commelin. Ind. 146: 1870; Karthik. et al., Fl. India. Enum. Monocot. 26.1989.

Cyanotis fasciculata var. *glabrescens* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 253.1881

Tonningia fasciculata (B.Heyne ex Roth) Kuntze, Revis. Gen. Pl. 2: 722.1891

Distribution: Karnataka (Bagalkot, Belagavi, Bengaluru, Chitradurga, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Tumakuru, Uttara kannada, Vijayapura, Yadgir), peninsular to N E India; endemic.

Cyanotis pilosa Schult. & Schult.f., Syst. Veg. 7: 1154. 1830.

Tradescantia pilosa Willd. ex Schult. & Schult.f., Syst. Veg. 7: 1155.1830

Cyanotis longifolia Wight, Icon. Pl. Ind. Orient. 6: 33, t. 2084.1853

Cyanotis wightii C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 250.1881

Tonningia pilosa (Schult. & Schult.f.) Kuntze, Revis. Gen. Pl. 2: 722.1891

Tonningia wightii (C.B.Clarke) Kuntze, Revis. Gen. Pl. 2: 722.1891

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Chitradurga, Kodagu, Tumakuru), Kerala, Maharashtra, Tamil Nadu, West Bengal; Sri Lanka.

Cyanotis tuberosa (Roxb.) Schult. & Schult.f., Syst. Veg. 7: 1153.1830; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Tradescantia tuberosa Roxb., Pl. Coromandel 1: 5.1799

Cyanotis concanensis Hassk., Commelin. Ind. 114.1870

Cyanotis stocksii Hassk., Commelin. Ind. 118.1870

Cyanotis thomsonii Hassk., Commelin. Ind. 133.1870

Tonningia tuberosa (Roxb.) Kuntze, Revis. Gen. Pl. 2: 722.1891

Cyanotis sahyadrica Blatt., J. Bombay Nat. Hist. Soc. 33: 77.1928

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Bidar, Chamarajanagar, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Raichur, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Odisha, Tamil Nadu; endemic.

Cyanotis villosa (Spreng.) Schult. & Schult.f., Syst. Veg. 7: 1155.1830; B.D.Sharma et al., Fl. Karnataka Anal. 290. 1984.

Tradescantia villosa Spreng., Syst. Veg. 2: 116.1825

Cyanotis lanceolata Wight, Icon. Pl. Ind. Orient. t. 2085.1853

Tradescantia cyanotidea Hassk., Commelin. Ind. 65.1870

Tonningia villosa (Spreng.) Kuntze, Revis. Gen. Pl. 2: 722.1891

Cyanotis cerifolia R.S.Rao & Kammathy, J. Linn. Soc., Bot. 59: 306.1966

Distribution: Karnataka (Chamarajanagar, Chitradurga, Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Dictyospermum Wight

Dictyospermum montanum Wight, Icon. Pl. Ind. Orient. 6: t. 2069.1853; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Tradescantia montana B.Heyne ex Wall., Numer. List 5203 B.1831

Commelina montana Steud., Nomencl. Bot. ed. 2, 1: 491.1840

Aneilema montanum (Wight) C.B. Clarke, Monogr. Phan. 3: 217 217.1881

Distribution: Karnataka (Chamarajanagar, Chikkamagalur, Dakshina Kannada, Hassan, Kodagu, Raichur, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Dictyospermum ovalifolium Wight, Icon. Pl. Ind. Orient. 6: t. 2070.1853; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Dictyospermum ovatum Hassk., Commelin. Ind. 24.1870

Dictyospermum wightii Hassk., Commelin. Ind. 19.1870

Aneilema ovalifolium (Wight) Hook.f. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 219.1881

Aneilema hallbergii Blatt., J. Bombay Nat. Hist. Soc. 33: 74.1928

Phaeneilema hallbergii (Blatt.) Raizada, Indian Forester 74: 335.1948

Distribution: Karnataka (Chamarajanagar, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Chatisgarh, Kerala, Tamil Nadu, Tripura; SE Asia, Sri Lanka.

Floscopia Lour.

Floscopia scandens Lour., Fl. Cochinch. 193.1790; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Tradescantia paniculata Roxb., Pl. Coromandel 1: 6.1799

Floscopia paniculata (Roxb.) Hassk., Pl. Jungh. 151.1852

Dithyrocarpus petiolatus Wight, Icon. Pl. Ind. Orient. 6: t. 2079.1853

Dithyrocarpus rothii Wight, Icon. Pl. Ind. Orient. 6: t. 2079.1853

Dithyrocarpus undulatus Wight, Icon. Pl. Ind. Orient. 6: t. 2079.1853

Lamprodithyros paniculatus (Roxb.) Hassk., Flora 46: 389.1863

Floscopia petiolata (Wight) Hassk., Commelin. Ind. 167.1870

Floscopia undulata (Wight) Hassk., Commelin. Ind. 166.1870

Distribution: Karnataka (Belagavi, Chikkamagalur, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Peninsular to NE India; Australia, SE Asia, Sri Lanka.

Murdannia Royle

Murdannia brownii Nandikar & Gurav, Taiwania 56: 227.2011

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Murdannia crocea (Griff.) Faden, Kew Bull. 32: 188.1977; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Aneilema crocea Griff., Not. Pl. Asiat. 3: 235.1851

subsp. **crocea**

Distribution: Karnataka (Dakshina Kannada, Udupi), Andaman & Nicobar Islands, peninsular India; Indo-china, Myanmar.

subsp. **ochracea** (Dalzell) Faden, Kew Bull. 32: 188.1977

Aneilema ochraceum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 135.1851

Dichaespermum repens Wight, Icon. Pl. Ind. Orient. 6: t. 2078.1853

Dichaespermum ochraceum (Dalzell) Hassk., Commelin. Ind. 41.1870

Phaeneilema ochraceum (Dalzell) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Murdannia ochracea (Dalzell) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Tumakuru), Kerala, Maharashtra, Tamil Nadu; endemic.

Murdannia dimorpha (Dalzell) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Aneilema dimorphum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 138.1851

Aneilema paniculatum Wight, Icon. Pl. Ind. Orient. t. 2075.1853

Phaeneilema dimorphum (Dalzell) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Murdannia edulis (Stokes) Faden, Taxon 29: 77.1980; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Commelina edulis Stokes, Bot. Mat. Med. 1: 184.1812

Commelina scapiflora Roxb., Fl. Ind. 1: 178.1820

Aneilema scapiflorum (Roxb.) Kostel., Allg. Med.-Pharm. Fl. 1: 127.1831

Murdannia scapiflora (Roxb.) Royle, Ill. Bot. Himal. Mts. 403.1840

Aneilema serotinum D.Don ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 200.1881

Phaeneilema scapiflorum (Roxb.) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Murdannia loureiroi (Hance) R.S.Rao & Kammathy, Notes Roy. Bot. Gard. Edinburgh 25: 184.1964

Distribution: Karnataka (Dharwad, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Sikkim, Tamil Nadu; China, Indonesia, Malaysia, Myanmar, Nepal, New Guinea, Philippines, Taiwan Thailand, Vietnam.

Murdannia esculenta (Wall. ex C.B.Clarke) R.S.Rao & Kammathy, Bull. Bot. Surv. India 3: 394.1962; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Aneilema esculentum Wall. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 206.1881

Commelina esculenta B.Heyne ex Hook.f., Fl. Brit. India 6: 377.1892

Aneilema pulneyense Fyson, Bull. Misc. Inform. Kew 1914: 332.1914

Phaeneilema pulneyense (Fyson) Raizada, Indian Forester 74: 335.1948

Murdannia gigantea (Vahl) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Commelina gigantea Vahl, Enum. Pl. 2: 177.1805

Aneilema giganteum (Vahl) R.Br., Prodr. Fl. Nov. Holl. 271.1810

Aneilema ensifolium Wight, Icon. Pl. Ind. Orient. t. 2072.1853

Prionostachys ensifolia Hassk. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 213.1881

Phaeneilema giganteum (Vahl) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Murdannia japonica (Thunb.) Faden, Taxon 26: 142 1977; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Commelina japonica Thunb., Trans. Linn. Soc. London 2: 332.1794

Commelina elata Vahl, Enum. Pl. Obs. 2: 178 1805

Commelina herbacea Roxb., Fl. Ind. ed.1,1: 179.1820

Dirtea japonica (Thunb.) Raf., Fl. Tellur. 3: 69.1837

Aneilema japonicum (Thunb.) Kunth, Enum. Pl. 4: 70.1843

Aneilema lineolatum (Blume) Kunth, Enum. Pl. 4: 69.1843

Aneilema latifolium Wight, Icon. Pl. Ind. Orient. t. 2072.1853

Aneilema herbaceum (Roxb.) Wall. ex C.B.Clarke, J. Linn. Soc., Bot. 11: 448.1870

Phaeneilema herbaceum (Roxb.) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Peninsular to NE India; Bhutan, China, Indonesia, Japan, Laos, Malaysia, Myanmar, Sikkim, Thailand.

Murdannia lanceolata (Wight) Kammathy, Bull. Bot. Surv. India 24: 206.1983

Dichaespermum lanceolatum Wight, Icon. Pl. Ind. Orient. 6: t. 2078.1853

Aneilema koenigii Wall. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 215 1881

Murdannia koenigii (Wall. ex C.B.Clarke) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930. *Aneilema koenigii* Wall. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 215 1881; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Udupi, Uttara Kannada), Tamil Nadu; Sri Lanka.

Murdannia lanuginosa (Wall. ex C.B.Clarke) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; B.D.Sharma et al., Fl. Karnataka Anal. 291. 1984.

Aneilema lanuginosum Wall. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 214.1881

Commelina lanuginosa B.Heyne ex Hook.f., Fl. Brit. India 6: 380.1892

Phaeneilema lanuginosum (Wall. ex C.B.Clarke) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Aneilema sienneum Blatt., J. Bombay Nat. Hist. Soc. 33: 75.1928

Phaeneilema sienneum (Blatt.) Raizada, Indian Forester 74: 336.1948

Murdannia sienna (Blatt.) Raizada, Indian Forester 84: 499.1958

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Murdannia loriformis (Hassk.) R.S.Rao & Kammathy, Bull. Bot. Surv. India 3: 393.1962; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Aneilema loriforme Hassk., Pl. Jungh. 143.1852

Aneilema terminale Wight, Icon. Pl. Ind. Orient. t. 2076.1853

Aneilema nudiflorum var. *terminale* (Wight) C.B.Clarke, Commelyn. Cyrtandr. Bengal. 211.1874

Murdannia malabarica var. *terminalis* (Wight) Santapau & S.K.Jain, Indian Forester 92: 643.1966

Distribution: Karnataka (Belagavi, Kodagu, Mysuru, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Murdannia nudiflora (L.) Brenan, Kew Bull. 7: 189.1952; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Commelina nudiflora L., Sp. Pl. 41. 1753.

Tradescantia malabarica L., Sp. Pl. ed. 2 412.1762

Ditelesia nudiflora (L.) Raf., Fl. Tellur. 3: 69.1837

Aneilema compressum Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 138.1851

Aneilema lancifolium Griff., Not. Pl. Asiat. 3: 236,1851

Aneilema malabaricum (L.) Merr., Philipp. J. Sci. 7: 232.1912

Murdannia malabarica (L.) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930

Phaeneilema malabaricum (L.) V.Naray., Indian Forest Rec., Bot. n.s., 3: 55.1941

Murdannia malabarica var. *compressa* (Dalzell) Santapau & S.K.Jain, Indian Forester 92: 643.1966

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; tropical Asia, naturalised elsewhere in tropics.

Murdannia pauciflora (G.Brückn.) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930

Aneilema pauciflorum Wight, Icon. Pl. Ind. Orient. t. 2077.1853

Dichaespermum aurantiacum Hassk. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 207.1881

Phaeneilema pauciflorum G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Murdannia wightii R.S.Rao & Kammathy, Notes Roy. Bot. Gard. Edinburgh 25: 184.1964; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; SE Asia.

Murdannia semiteres (Dalzell) Santapau, Poona Agric. Coll. Mag. 41(4): 15 1951; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Aneilema semiteres Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 138.1851

Dichaespermum juncoides Wight, Icon. Pl. Ind. Orient. 6: t. 2078.1853

Dichaespermum semiteres (Dalzell) Hassk., Commelin. Ind. 41.1870

Aneilema paniculatum Wall. ex C.B.Clarke, Monogr. Phan. 3: 215. 1881.

Commelina nimmoniana J.Graham ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 215.1881

Dichaespermum paniculatum Hassk. ex C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 215.1881

Phaeneilema paniculatum (Wall. ex C.B.Clarke) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Murdannia paniculata (Wall. ex C.B.Clarke) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930

Murdannia juncoidea (Wight) R.S.Rao & Kammathy, Bull. Bot. Surv. India 6: 3 1965

Murdannia nimmoniana (J.Graham ex C.B.Clarke) Bole & M.R.Almeida, J. Bombay Nat. Hist. Soc. 83: 593.1987

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Vijayapura), Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; tropical E Africa.

Murdannia simplex (Vahl) Brenan, Kew Bull. 7: 186.1952; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Commelina simplex Vahl, Enum. Pl. 2: 177. 1805

Aneilema simplex (Vahl) Kunth, Enum. Pl. 4: 71.1843

Aneilema secundum Wight, Icon. Pl. Ind. Orient. t. 2074.1853

Aneilema simplex (Vahl) C.B.Clarke, Monogr. Phan. 3: 212. 1881.

Aneilema rigidum Blatt., J. Bombay Nat. Hist. Soc. 33: 73.1928

Phaeneilema rigidum (Blatt.) Raizada, Indian Forester 74: 336.1948

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Udupi, Uttara Kannada), Peninsular to NE India; tropical Africa & Asia.

Murdannia spirata (L.) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Commelina spirata L., Mant. Pl. 2: 176.1771

Aneilema spiratum (L.) Sweet, Hort. Suburb. Lond. 12.1808

Aneilema spiratum (L.) R.Br., Prodr. 271. 1810.

Aphylax spiralis Salisb., Trans. Hort. Soc. London 1: 271.1812

Phaeneilema spiratum (L.) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; endemic.

Murdannia vaginata (L.) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Commelina vaginata L., Mant. Pl. 2: 177.1771

Aneilema vaginatum (L.) R.Br., Prodr. 271 271.1810

Aneilema vaginatum (L.) Wall., Numer. List 5212.1831

Dilasia vaginata (L.) Raf., Fl. Tellur. 4: 123. 1838

Phaeneilema vaginatum (L.) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Dictyospermum vaginatum (L.) D.Y.Hong, Acta Phytotax. Sin. 12: 477. 1974

Distribution: Karnataka (Belagavi, Uttara Kannada), throughout India; Australia, Myanmar, Sri Lanka.

Murdannia versicolor (Dalzell) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Aneilema versicolor Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 136.1851

Phaeneilema versicolor (Dalzell) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56. 1927

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Dharwad, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; SE Asia.

Murdannia zeylanica (C.B.Clarke) G.Brückn., Nat. Pflanzenfam. ed. 2 15a: 173.1930; .B.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Aneilema zeylanicum C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 204 1881

Aneilema zeylanicum var. *longiscapum* C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 204.1881

Phaeneilema zeylanicum (C.B.Clarke) G.Brückn., Notizbl. Bot. Gart. Berlin-Dahlem 10: 56.1927

Murdannia zeylanica var. *longiscpa* (C.B.Clarke) Raizada, Indian Forester 84: 499. 1958

Distribution: Karnataka (Shivamogga), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka

Pollia Thunb.

Pollia secundiflora (Blume) Bakh.f., Bkbn. Fl. Java 211: 10.1949; B.D.Sharma et al., Fl. Karnataka Anal. 292. 1984.

Commelina secundiflora Blume, Enum. Pl. Javae 1: 3.1827

Aneilema secundiflora (Blume) Kunth, Enum. Pl. 4: 69.1843

Aclisia indica Wight, Icon. Pl. Ind. Orient. t. 2068.1853

Pollia indica (Wight) Thwaites, Enum. Pl. Zeyl. 323.1864

Pollia didyma C.B.Clarke, J. Linn. Soc., Bot. 11: 451.1871

Pollia sorzogonensis var. *gigantea* (Hassk.) C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 127.1881

Pollia sorzogonensis var. *indica* (Wight) C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 127.1881

Aclisia secundiflora (Blume) Bakh.f., Fl. Java 3: 658 1968

Pollia secundiflora var. *gigantea* (Hassk.) Sanjappa, Fl. Ind. Enumerat. - Monocot. 30. 1989

Pollia secundiflora var. *indica* (Wight) Sanjappa, Fl. Ind. Enumerat. - Monocot. 30.1989

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi), Andaman & Nicobar Islands, S & NE India; Sri Lanka.

Rhopalephora Hassk.

Rhopalephora scaberrima (Blume) Faden, Phytologia 37: 480.1977; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Commelina scaberrima Blume, Enum. Pl. Javae 1: 4.1827

Aneilema scaberrimum (Blume) Kunth, Enum. Pl. 4: 69.1843

Tradescantia scaberrima (Blume) Hassk., Tijdschr. Natuurl. Gesch. Physiol. 10: 120.1843

Dictyospermum protensum Wight, Icon. Pl. Ind. Orient. 6: t.: 2071.1853

Aneilema protensum (Wight) Wall. ex Thwaites, Enum. Pl. Zeyl. 322.1864

Lamprodithyros protensus (Wight) Hassk., Naturw. Reise Mossambique 2: 529.1864

Piletocarpus protensus (Wight) Hassk., Commelin. Ind. 15.1870

Aneilema scaberrimum var. *fruticosum* C.B.Clarke in Hook.f., Fl. Brit. India 6: 383.1892

Phaeneilema scaberrimum (Blume) Raizada, J. Bombay Nat. Hist. Soc. 48: 677.1950

Dictyospermum scaberrimum (Blume) C.V.Morton ex Panigrahi, Phytologia 29: 338 1975

Rhopalephora scaberrima var. *fruticosa* (C.B.Clarke) Vrinda & Panikkar, J. Econ. Taxon. Bot. 23: 670.1999

Distribution: Karnataka (Hassan, Kodagu, Shivamogga, Uttara Kannada), Peninsular to NE India; Bhutan, China, Nepal, SE Asia, Taiwan.

Tradescantia L.

Tradescantia pallida (Rose) D.R.Hunt, Kew Bull. 30: 452.1975

Setcreasea pallida Rose, Acta Bot. Neerl. 4: 167.1955

Setcreasea purpurea Boom, Acta Bot. Neerl. 4: 167.1955

Setcreasea lanceolata Faruqi, Mehra & Celarier, Rhodora 64: 330.1962

Distribution: Karnataka (Cultivated in gardens), cultivated throughout drier parts; native to Mexico

Tradescantia sillamontana Matuda, Bol. Soc. Bot. México 18: 1.1955

Tradescantia pexata H.E.Moore, Baileya 8: 100.1960

Distribution: Karnataka (Cultivated in gardens), cultivated throughout drier parts; native to Mexico

Tradescantia spathacea Sw., Prodr. Veg. Ind. Occ. 57.1788

Tradescantia discolor L'Hér., Sert. Angl. 8.1789

Rhoeo discolor (L'Hér.) Hance, Ann. Bot. Syst. 3: 660.1852

Rhoeo spathacea (Sw.) Stearn, Baileya 5: 198.1957

Distribution: Karnataka (cultivated throughout), cultivated throughout plains; native to Mexico.

Local/Common/Vernacular name: *Moses-in-the-Cradle* (English); *Rhoeo gida* (Kannada)

Tradescantia zebrina Bosse, Vollst. Handb. Bl.-gärtn., ed. 2 4: 655. 1849

Zebrina pendula Schnizl., Bot. Zeitung (Berlin) 7: 870.1849

Commelina zebrina C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 318.1881

Tradescantia tricolor C.B.Clarke in DC. & C.DC., Monogr. Phan. 3: 318.1881

Tradescantia pendula (Schnizl.) D.R.Hunt, Kew Bull. 36: 197.1981

Distribution: Karnataka (throughout), native to Mexico.

Local/Common/Vernacular names: *Inch plant, Spiderwort, Wandering jew* (English)

Connaraceae

Connarus L.

Connarus monocarpus L., Sp. Pl. 675. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 60. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Uttara Kannada), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Indian Zebrawood* (English)

Connarus sclerocarpus (Wight & Arn.) Schellenb., Candollea 2: 98 1925; C.J.Saldanha, Fl. Karnataka 2: 188. 1996

Distribution: Karnataka (Dakshina Kannada), Kerala, Tamil Nadu; endemic.

Connarus wightii Hook. f., Fl. Brit. India 2: 51. 1876; B.D.Sharma et al., Fl. Karnataka Anal. 60. 1984.

Connarus ritchiei Hook. f., Fl. Brit. India 2: 51. 1876

Connarus paniculatus sensu Kesh. Murthy & Yogan, Fl. Coorg. 127 1990, non Roxb.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Rourea Aubl.

Rourea minor (Gaertn.) Alston, Trimen, Handb. Fl. Ceylon 6: 67. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 61. 1984.

Aegiceras minus Gaertn., Fruct. 1: 216. t.46. 1788

Connarus santalooides Vahl, Symb. Bot. 4: 87. 1794

Connarus roxburghii Hook. & Arn., Bot. Beechey Voy. 179.1833

Rourea santalooides (Vahl) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 144. 1834

Santaloides roxburghii (Hook. & Arn.) Kuntze, Revis. Gen. Pl. 1: 155. 1891

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Kerala, Maharashtra, Manipur, Meghalaya, Mizoram, Odisha, Tamil Nadu, West Bengal; Australia, China, Cambodia, Laos, Sri Lanka, Taiwan, Thailand, Vietnam.

Local/Common/Vernacular name: *Huleechala balli* (Kannada)

Rourea prainiana Talbot, For. Fl. Bombay 1: 368.1909

Tali prainiana (Talbot) M.R.Almeida, Fl. Maharashtra 1: 284.1996

Distribution: Karnataka (Belagavi, Chikkamagaluru, Shivamogga, Uttara Kannada), Kerala, Maharashtra; endemic.

Convolvulaceae

Aniseia Choisy

Aniseia martinicensis (Jacq.) Choisy, Convolv. Diss. Sec. 144.1837; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Convolvulus martinicensis Jacq., Stirp. Amer. Hist. 26, t. 17.1763

Convolvulus uniflorus Burm. f., Fl. Indica 47, t. 247. 1768

Ipomoea martinicensis (Jacq.) G. Mey., Prim. Fl. Esseq. 98.1818

Ipomoea uniflora (Burm.f.) Roem. & Schult., Syst. Veg. 4: 247.1819

Aniseia uniflora (Burm.f.) Choisy, Convolv. Orient. 101. 1833; T.Cooke, Fl. Bombay 2:232. 1903

Distribution: Karnataka (Dakshina Kannada), throughout India; pantropical.

Argyreia Lour.

Argyreia bracteata Choisy, Convolv. Orient. 39.1834

Ipomoea kleiniana Roem. & Schult., Syst. Veg., ed. 15, 4: 789.1819

Argyreia kleiniana Raizada, Indian Forester 92:302 .1966; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Distribution: Karnataka (Bengaluru, Davanagere, Tumakuru), Kerala, Tamil Nadu; endemic.

Argyreia cuneata (Willd.) Ker Gawl., Bot. Reg. 8: t. 661. 1822; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984. Plate 5F

Convolvulus cuneatus Willd., Sp. Pl. 1(2): 873. 1798

Ipomoea atrosanguinea Hook., Bot. Mag. 47: t. 2170. 1820

Letsomia cuneata (Willd.) Roxb., Fl. Ind. 1: 81.1824; Fl. Ind. 1: 491. 1832.

Rivea cuneata (KerGawler) Wight, Icon. Pl. Ind. Orient. 3(2): 8, t. 890. 1845

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Achari gida*, *Halle balli*, *Kallana gida*, *Kallana hambu* (Kannada)

Argyreia cymosa Sweet, Hort. Brit. 289. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Letsomia cymosa Roxb., Fl. Ind. 2: 82.1824; Fl. Ind. 1: 492.1832.

Distribution: Karnataka (Ballari, Bidar, Chitradurga, Kolar), Peninsula India; Philippines.

Argyreia daltonii C.B.Clarke in Hook.f., Fl. Brit. India 4: 190.1883; Bhaskar & Kushalappa, Fl. Tumkur Dist. 93.2013.

Distribution: Karnataka (Tumakuru), Kerala; endemic.

Argyreia elliptica Arn. ex Choisy, Convolv. Orient. 35.1834; Bhaskar & Kushalappa, Fl. Tumkur Dist. 93.2013.

Distribution: Karnataka (Chitradurga, Davanagere, Hassan, Mysuru, Shivamogga, Tumakuru), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Naayi ugani hambu* (Kannada)

Argyreia hirsuta Wight & Arn., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 18(1): 356c. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Rivea hirsuta Wight, Icon. Pl. Ind. Orient.3(2): t. 891.1845

Distribution: Karnataka (Chikkamagaluru, Mysuru, Shivamogga), Chattisgarh, Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Argyreia hookeri C.B.Clarke in Hook f., Fl. Brit. India 4:185. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Kodagu, Shivamogga, Uttara Kannada), W Peninsula India, Assam, Sikkim; Bhutan.

Argyreia involucrata C.B.Clarke in Hook f., Fl. Brit. India 4:187. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Argyreia involucrata var. *inequivalvis* C.B.Clarke in Hook f., Fl. Brit. India 4:187. 1883

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Argyreia lawii C.B.Clarke in Hook f., Fl. Brit. India 4:190. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala, Maharashtra; endemic.

Argyreia nellygherya Choisy, Convolv. Orient. 32. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Argyreia leschenaultii Choisy, Convolv. Orient. 31. 1834; C.B.Clarke in Hook.f., Fl. Brit. India 4: 188. 1883.

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Argyreia nervosa (Burm.f.) Bojer, Hortus Maurit. 224.1837; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Convolvulus nervosus Burm. f., Fl. Indica 48, pl. 20, f. 1. 1768

Convolvulus speciosus L. f., Suppl. Pl. 137.1781

Ipomoea speciosa (L.f.) Pers., Syn. Pl. 1: 183. 1805

Letsomia nervosa (Burm.f.) Roxb., Fl. Ind. 2: 78.1824

Argyria speciosa (L.f.) Sweet, Hort. Brit. Ed.2, 373.1830; C.B.Clarke in Hook.f., Fl. Brit. India 4: 185. 1883.

Rivea nervosa (Burm.f.) Hallier f., Bull. Herb. Boissier 5: 381.1897

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Mysuru, Raichur, Tumakuru, Udupi, Uttara Kannada),throughout India; China, Indonesia, Mauritius

Local/Common/Vernacular name: *Chandra pada* (Kannada)

Argyreia osyrensis (Roth) Choisy in DC., Prodr. 9: 334. 1845

Ipomoea imbricata Roth, Nov. Pl. Sp. 112. 1821

Ipomoea osyrensis Roth, Nov. Pl. Sp. 117. 1821

Letsomia aggregata Roxb., Fl. Ind. ed.2, 1: 488. 1832; C.B.Clarke in Hook.f., Fl. Brit. India 4: 191. 1883

Argyreia aggregata (Roxb.) Choisy, Convolv. Orient. 45.1834

Letsomia aggregata var. *osyrensis* (Roth) C. B. Clarke in Hook f., Fl. Brit. India 4:192. 1883

Argyreia imbricata (Roth) Santapau & V.Patel, Trans. Bose Res. Inst. Calcutta 22 (Convolv. Bomb.: Addit. & Corr.) 40.1958; B.D.Sharma et al., Fl. Karnataka Anal. 179. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Davanagere, Dharwad, Hassan, Mysuru, Tumakuru), Kerala, Maharashtra, Tamil Nadu; China, S & SE Asia.

Local/Common/Vernacular name: *Juju garanga balli* (Kannada)

Argyreia pilosa Wight & Arn., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 18(1): 356c. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Argyreia pomacea Choisy, Convolv. Orient. 32. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Letsomia pomacea Roxb., Fl. Ind. 1: 493. 1832

Rivea pomacea Wight, Icon. Pl. Ind. Orient.3(1): t. 888.1845

Distribution: Karnataka (Bengaluru, Davanagere, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Argyreia populifolia Choisy, Convolv. Orient. 32. 1834 & in DC., Prodr. 9: 329. 1845; C.B.Clarke in Hook.f., Fl. Brit. India 4: 187. 1883.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Tumakuru), Kerala, Tamil Nadu; Sri Lanka.

Argyreia serecia Dalzell & A.Gibson, Bombay Fl. 169.1861; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984. Plate 21F

Ipomaea bracteata auct. non Cav. 1799: J.Graham, Cat. Bombay Pl. 131.

Distribution: Karnataka (Belagavi, Ballari, Chikkamagaluru, Chitradurga, Dharwad, Kolar, Mysuru, Tumakuru, Uttara Kannada), Dadra, Kerala, Maharashtra, Nagar Haveli, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Samudr balli*, *Yemme balli* (Kannada)

Bonamia Thouars

Bonamia semidigyna (Roxb.) Hallier f., Bot. Jahrb. Syst. 16: 528.1893

Convolvulus semidigynus Roxb., Fl. Ind., ed.1, 2: 47. 1824; Fl.Ind. ed.2, 1: 468. 1832.

Breweria cordata Blume, Bijdr. Fl. Ned. Ind. 13: 722. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Breweria roxburghii Choisy, Mém. Soc. Phys. Genève 6: 493.1833

Prevostea cordata Hallier f., Bot. Jahrb. Syst. 18: 93.1894

Bonamia cordata (Hallier f.) Hallier f., Bull. Herb. Boissier 7: 43.1899

Distribution: Karnataka (Uttara Kannada, Udupi), Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Breweria R. Br.

Breweria evolvuloides Choisy, Mém. Soc. Phys. Genève 6: 494.1833; C.B.Clarke in Hook f., Fl. Brit. India 4: 224. 1883.

Seddera evolvuloides Wight, Icon. Pl. Ind. Orient. t. 1369. 1848

Distribution: Karnataka (), Kerala, Tamil Nadu; endemic.

Camonea Raf.

Camonea umbellata (L.) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 583. 2017.

Convolvulus umbellatus L., Sp. Pl. 155.1753

Convolvulus cymosus Desr. in Lam., Encycl. 3: 556. 1792

Ipomoea bifida Roem. & Schult., Syst. Veg., ed. 15 bis 4: 241. 1819

Ipomoea heynei Roem. & Schult., Syst. Veg., ed. 15 bis 4: 237. 1819

Ipomoea rothii Roem. & Schult., Syst. Veg., ed. 15 bis 4: 241. 1819

Ipomoea cymosa (Desr.) Roem. & Schult., Syst. Veg. 6: 241.1820; C.B.Clarke in Hook.f., Fl. Brit. India 4: 211.1883.

Convolvulus blandus Roxb., Fl. Ind. 2: 50. 1824; Fl.Ind. 2: 470. 1832.

Convolvulus pentagonus Roxb., Fl. Ind. Ed. 1, 2: 72. 1824; Fl.Ind. ed. 2, 2: 485. 1832.

Convolvulus rothii Spreng., Syst. Veg. ed. 16, 1: 600. 1824

Ipomoea caliginosa Choisy, Mém. Soc. Phys. Genève 4: 461. 1833

Merremia umbellata (L.) Hallier f., Bot. Jahrb. Syst. 16(4-5): 552.1893; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), throughout India; Pantropical.

Camonea vitifolia (Burm.f.) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 583. 2017.

Convolvulus angularis Burm.f., Fl. Indica 46.1768

Ipomoea vitifolia (Burm.f.) Blume, Bijdr. Fl. Ned. Ind. 13: 709. 1825; C.B.Clarke in Hook.f., Fl. Brit. India 4: 213. 1883.

Ipomoea vitifolia var. *angularis* (Burm.f.) Choisy, Mém. Soc. Phys. Genève 4: 461. 1833

Merremia vitifolia (Burm.f.) Hallier.f., Bot. Jahrb. Syst. 16: 552. 1893; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), throughout India except NW desert areas; S & SE Asia.

Convolvulus L.

Convolvulus arvensis L., Sp. Pl. 1: 153. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Convolvulus malcolmii Roxb., Fl. Ind 2:55. 1824

Convolvulus chinensis Ker Gawler in Edwards's Bot. Reg. t. 322. 1878

Distribution: Karnataka (Belagavi, Ballari, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), throughout India; Americas, Asia, Europe.

Local/Common/Vernacular name: *Narangi balli* (Kannada)

Convolvulus rotundifolius Choisy, Convolv. Orient. 95. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Retzia pilosa Rottler ex Choisy in DC., Prodr. 9: 403. 1845

Convolvulus stocksii auct. non Boiss., 1875; Seetharam, Kotresh & Uploankar, Fl. Dist. 53. 2000.

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Dharwad, Kalaburgi, Tumakuru, Vijayapura, Yadgir); Peninsula India; Afghanistan, Pakistan, Turkestan,

Cressa L.

Cressa cretica L., Sp. Pl. 223. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Distribution: Karnataka (Ballari, Dakshina Kannada, Udupi, Uttara Kannada), throughout plains excluding NE India; pantropical.

Cuscuta L.

Cuscuta campestris Yunck., Mem. Torrey Bot. Club 18(2): 138. 1932; B.D.Sharma et al., Fl. Karnataka Anal. 185. 1984.

Distribution: Karnataka (Ballari), throughout; Africa, Americas, Asia, Australia, Europe, Pacific Islands.

Cuscuta chinensis Lam., Encycl. 2(1): 229. 1786; C.B.Clarke in Hook.f., Fl. Brit. India 4: 226. 1883.

Grammica aphylla Lour., Fl. Cochinch. 171. 1790

Cuscuta carinata R. Br., Prodr. 491. 1810

Cuscuta sulcata Roxb., Fl. Ind. 1: 447. 1832

Cuscuta chinensis var. *carinata* (R.Br.) Engelm., Trans. Acad. Sci. St. Louis 1(3): 480. 1859

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Dharwad, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; Australia, Middle East, SE Asia

Cuscuta arabica Roth, Nov. Pl. sp. 100. 1821

Cuscuta arabica Wight, Icon. Pl. Ind. Orient. , t. 1371 1850

Cuscuta boissieri Stocks, Hooker's J. Bot. Kew Gard. Misc. 4: 173. 1852

Cuscuta oxypetala Boiss., Diagn. Pl. Orient., ser. 2 3: 130. 1856

Distribution: Karnataka (Mysuru), S to W India; Pakistan, Middle East, N Africa.

Local/Common/Vernacular name: *Badanike* (Kannada)

Cuscuta reflexa Roxb., Pl. Coromandel 2: 3, t. 104. 1798; B.D.Sharma et al., Fl. Karnataka Anal. 185. 1984.

Cuscuta verrucosa Sweet, Brit. Fl. Gard. 1: t. 6. 1823

Cuscuta hookeri Sweet, Hort. Brit. 290. 1826

Cuscuta macrantha G.Don, Gen. Hist. 4: 305. 1838

Cuscuta elatior Choisy, Mém. Soc. Phys. Genève 9: 273. 1841

Monogynella reflexa (Roxb.) Holub, Folia Geobot. Phytotax. 12(4): 429. 1977

Distribution: Karnataka (throughout), throughout India; Afghanistan, S & SE Asia.

Local/Common/Vernacular names: *Aakash balli*, *Badanike* (Kannada); *Aakashavalle* (Sanskrit)

Distimake Raf.

Distimake aegyptius (L.) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 573. 2017.

Ipomoea aegyptia L., Sp. Pl. 1: 162. 1753

Convolvulus pentaphyllus L., Sp. Pl. ed. 2: 223. 1762

Convolvulus vitifolius Burm. f., Fl. Indica 45, t. 18, f. 1. 1768

Ipomoea pentaphylla (L.) Jacq., Collectanea 2: 297. 1788; C.B.Clarke in Hook.f., Fl. Brit. India 4: 193. 1883.

Convolvulus munitus Wall., Numer. List no. 1354 1829

Batatas pentaphylla (L.) Choisy, Convolv. Orient. 54. 1834

Spiranthera pentaphylla (L.) Bojer, Hortus Maurit. 226. 1837

Merremia pentaphylla (L.) Hallier f., Bot. Jahrb. Syst. 16: 552. 1893

Operculina aegyptia (L.) House, Bull. Torrey Bot. Club 33: 502. 1906

Operculina pentaphylla (L.) House, Bull. Torrey Bot. Club 33: 502. 1906

Merremia aegyptia (L.) Urb., Symb. Antill. 4: 505. 1910; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Distribution: Karnataka (Bengaluru, Chitradurga, Dharwad, Kalaburgi, Raichur), Peninsula India extending to Punjab; Pantropical.

Local/Common/Vernacular names: *Hairy Merremia*, *Hairy Woodrose* (English)

Distimake dissectus (Jacq.) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 574. 2017.

Convolvulus dissectus Jacq., Obs. Bot. 2: 4. 1767

Ipomoea sinuata Ortega, Nov. Pl. Descr. Dec. 7: 84. 1798; C.B.Clarke in Hook.f., Fl. Brit. India 4: 214. 1883.

Ipomoea dissecta (Jacq.) Pursh, Fl. Amer. Sept. (Pursh) 1: 145. 1813.

Merremia dissecta (Jacq.) Hallier f., Bot. Jahrb. Syst. 16: 552. 1893; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Operculina dissecta (Jacq.) House, Bull. Torrey Bot. Club 33: 500. 1906

Distribution: Karnataka (Belagavi), throughout plains of India; Australia, China, S & SE Asia; native to tropical America.

Distimake kentrocaulos (C.B.Clarke) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 574. 2017.

Ipomoea tuberosa A.Rich., Tent. Fl. Abyss. 2: 67. 1850

Ipomoea kentrocaulos C.B.Clarke, Hook.f., Fl. Brit. India 4: 213. 1883

Operculina kentrocaulos Hallier f., Bot. Jahrb. 18119. 1894

Merremia kentrocaulos (C.B.Clarke) Hallier f., Bull. Herb. Boissier 7: 410. 1899.

Distribution: Karnataka (Kalaburgi, Yadgir), Peninsula extending to NW India), Sri Lanka, Malaysia, tropical Africa.

Distimake quinquefolius (L.) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 575. 2017.

Ipomoea quinquefolia L., Sp. Pl. 162. 1753

Convolvulus quinquefolius (L.) L., Syst. Nat. (ed. 10) 923. 1759

Batatas quinquefolia (L.) Choisy, Convolv. Orient. 127. 1834

Merremia quinquefolia (L.) Hallier f., Bot. Jahrb. Syst. 16(4–5): 552. 1893; Parimala, Fl. Dharwad Dist. 317. 2011.

Distribution: Karnataka (Dharwad), Uttar Pradesh; Pantropical, native to tropical America.

Distimake rhyncorhiza (Dalzell) A.R.Simoes & Staples, Bot. J. Linn. Soc. 183: 575. 2017.

Ipomoea rhyncorhiza Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 179. 1851

Merremia rhyncorhiza (Dalzell) Hallier f., Bot. Jahrb. Syst. 16 (4–5): 552. 1893. (as "rhynchorrhiza"); B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984. Plate 21G

Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.

Erycibe Roxb.

Erycibe paniculata Roxb., Pl. Coromandel 2: 31, t. 154. 1802; Fl. Ind. 1:585. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Catonia glauca Vahl, Skr. Nat. Hist. Selsk. Kiobenb. 6:98. 1810

Erimatalia rheedei Roem. & Schult., Syst. Veg., ed. 15 bis 5: 331. 1819

var. *paniculata*

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India including Andaman & Nicobar Islands; Australia, Malaysia, Myanmar, Sri Lanka.

Local/Common/Vernacular names: *Singara balli*, *Sundra balli* (Kannada)

var. *wightiana* C.B.Clarke in Hook f., Fl. Brit. India 4:180. 1883

Catonia elliptica Vahl in Skrивt. Nat. Hist. Selsk. Kiobenb. 6: 101. 1810

Erycibe wightiana Graham, Cat. Pl. Bomb. 137. 1839 ex Dalzell

Distribution: Karnataka (), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Evolvulus L.**Evolvulus alsinoides** (L.) L., Sp. Pl. ed. 2. 392. 1762; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus alsinoides L., Sp. Pl. 1: 157. 1753

Evolvulus hirsutulus Choisy, Mém. Soc. Phys. Genève 8: 76.1837

Distribution: Karnataka (throughout), throughout India; paleotropical.

Local/Common/Vernacular names: *Vishnu kranthi*, *Vishnu kranthi soppu* (Kannada)

Evolvulus nummularius (L.) L., Sp. Pl. ed. 2. 391. 1762

Convolvulus nummularius L., Sp. Pl. 1: 157. 1753

Volvulopsis nummularium (L.) Roberty, Candollea 14: 28.1952

Distribution: Karnataka (throughout), naturalized throughout plains of India, Asia, Africa, native to Americas.

Hewittia Wight & Arn.**Hewittia malabarica** (L.) Suresh, Interpret. Van Rheede's Hort. Malab. 88.1988

Convolvulus malabaricus L., Sp. Pl. 1: 155. 1753

Convolvulus sublobatus L. f., Suppl. Pl. 135.1781

Convolvulus bicolor Vahl, Symb. Bot. 3: 25.1794

Convolvulus bracteatus Vahl, Symb. Bot. 3: 25.1794

Convolvulus involucratus Ker Gawl., Bot. Reg. 4: t. 318. 1818

Calystegia keriana Sweet, Hort. Brit. 287. 1826

Shutereia bicolor (Vahl) Choisy, Mém. Soc. Phys. Genève 6: 486.1833

Argyreia malabarica Choisy, Convolv. Orient. 38. 1834; C.B.Clarke in Hook.f., Fl. Brit. India 4: 189. 1883.

Hewittia bicolor (Vahl) Wight & Arn., Madras J. Lit. Sci. 1(5): 22. 1837; C.B.Clarke in Hook.f., Fl. Brit. India 4: 216. 1883.

Palmia bicolor Endl. ex Dalzell & A.Gibson, Bombay Fl. 163. 1861

Hewittia sublobata (L.f.) Kuntze, Revis. Gen. Pl. 2: 441.1891; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Shutereia sublobata (L.f.) House, Bull. Torrey Bot. Club 33: 318.1906

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu,Udupi, Uttara kannada), Peninsula India, Assam; pantropical.

Ipomoea L.**Ipomoea alba** L., Sp. Pl. 1: 161.1753; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Ipomoea bona-nox L., Sp. Pl. ed. 2, 1: 228. 1762

Convolvulus bona-nox (L.) Spreng., Syst. Veg. 1: 600.1825; C.B.Clarke in Hook.f., Fl. Brit. India 4: 197. 1883.

Calonyction speciosum Choisy, Convolv. Orient. 59.1833

Calonyction bona-nox (L.) Bojer, Hortus Maurit. 227.1837

Convolvulus muricatum G.Don, Gen. Hist. 4: 264. 1837; C.B.Clarke in Hook.f., Fl. Brit. India 4: 197. 1883.

Convolvulus aculeatus var. *bona-nox* (L.) Kuntze, Revis. Gen. Pl. 3(3): 212.1898

Calonyction album (L.) House, Bull. Torrey Bot. Club 31(11): 591.1904

var. alba

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical America.
 Local/Common/Vernacular names: *American moon flower*, *Good night flower* (English); *Chandra kanti*, *Chandra pushpa*, *Kaadu katti kayi* (Kannada); *Chandra kanthi*, *Mandavalli* (Sanskrit)

var. *grandiflora* C.B.Clarke in Hook f., Fl. Brit. India 4:197. 1883

Ipomoea noctiluca Herb., Bot. Reg. 11:t. 917, in nota. 1825

Ipomoea aculeata Blume, Bijdr. Fl. Ned. Ind. 13: 715. 1826

Ipomoea noctiflora Griff., Not. Pl. Asiat. 4: 286. 1854

Distribution: Karnataka (cultivated throughout), cultivated throughout India; Bangladesh, E Asia, tropical Africa.

Ipomoea aquatica Forssk., Fl. Aegypt.-Arab. 44.1775; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus repens Vahl, Symb. Bot. 2: 17. 1790

Convolvulus adansonii Desr. in Lam. Encyc. 3: 560. 1792

Ipomoea reptans Poir., Encycl., Suppl. 3(2): 460.1814; C.B.Clarke in Hook.f., Fl. Brit. India 4: 210.1883.

Ipomoea repens Roth, Nov. Pl. Sp. 110.1821

Ipomoea subdentata Miq., Fl. Ned. Ind. 2: 614. 1856

Distribution: Karnataka (throughout in ponds, tanks), throughout India; paleotropical.

Local/Common/Vernacular names: *Bale soppu*, *Bili hambu*, *Neeru bili gedde hambu*, *Neeru bili gedde hoovu*, *Neeru bili hambu*, *Neeru hambu*, *Rakshan karalu* (Kannada); *Kadambi*, *Kalaka*, *Kalambi*, *Kalambika*, *Kalashaka*, *Kechuka*, *Naadi*, *Naadika*, *Naalika*, *Pattashaka*, *Pechu*, *Pechuli*, *Shatha parana*, *Shraddha shaka*, *Vishvarochanna* (Sanskrit)

Ipomoea arachnosperma Welw., Apont. 588.1859

Convolvulus dichrous Roem. & Schult., Syst. Veg., ed. 15 bis 4: 268. 1819

Ipomoea pilosa Sweet, Hort. Brit. 289.1827; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Ipomoea dichroa Choisy in DC., Prodr. 9: 364. 1845

Distribution: Karnataka (Ballari), throughout except NE India; Pakistan, Africa.

Ipomoea asarifolia (Desr.) Roem. & Schult., Syst. Veg. 4: 251.1819; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus asarifolius Desr., Encycl. 3: 562.1789

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Davanagere, Tumakuru), Maharashtra; native to Americas.

Local/Common/Vernacular name: *Ginger-leaf morning-glory* (English)

Ipomoea batatas (Choisy) Benth. ex C.B.Clarke in Hook. f., Fl. Brit. India 4: 201.1883; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Plate 5G

Aniseia barlerioides Choisy, Mém. Soc. Phys. Genève 6: 484.1834

Distribution: Karnataka (Belagavi, Chitradurga, Davanagere, Hassan, Mysuru, Tumakuru, Uttara Kannada), throughout India except Himalayas; Myanmar.

Ipomoea batatas (L.) Lam., Encycl. 2: 465. 1793; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus batatas L., Sp. Pl. 1: 154. 1753

Convolvulus edulis Thunb., Syst. Veg. (ed. 14) 203.1784

Convolvulus candidans Soland. ex Sims, Bot. Mag. 39:t. 1603. 1813

Batatas edulis (Thunb.) Choisy, Convolv. Orient. 53. 1833

Ipomoea batatas var. *edulis* (Thunb.) Makino, Somoku-Dzusetsu 4:t. 25 1903

Distribution: Karnataka (cultivated throughout), cultivated throughout India; Indonesia, Japan, Laos, Malaysia, Nepal, New Guinea, Pakistan, Philippines, Sri Lanka, Thailand, Vietnam; Africa, Americas, Australia, Pacific Islands; native to tropical America..

Local/Common/Vernacular names: *Sweet potato* (English); *Bella genasu*, *Genasu*, *Genasu gedde*, *Sihi genasu* (Kannada); *Kanda granthi*, *Pichhila*, *Tindalu*, *Tindithaka*, *Rakthalu*, *Roma shakandaka*, *Swadu kandaka* (Sanskrit); *Kireng* (Tulu)

Ipomoea biflora (L.) Pers., Syn. Pl. 1: 183. 1805

Convolvulus biflorus L., Sp. Pl. ed. 2., App. 1668. 1763

Convolvulus sinensis Desr., Encycl. 3(2): 557. 1789

Ipomoea timorensis Blume, Bijdr. Fl. Ned. Ind. 711. 1825

Convolvulus hardwickii Spreng., Syst. Veg. 4: 60. 1827

Convolvulus calycinus Roxb., Fl. Ind. 1: 471. 1832

Aniseia biflora (L.) Choisy, Mém. Soc. Phys. Genève 6: 483. 1833

Aniseia calycina (Roxb.) Choisy, Mém. Soc. Phys. Genève 6: 481. 1834

Ipomoea sinensis (Desv.) Choisy, Mém. Soc. Phys. Genève 6: 459. 1834

Ipomoea calycina auct. non Messin. 1869. (Roxb.) Benth. ex C. B. Clarke in Hook. f., Fl. Brit. India 4: 201. 1883

Ipomoea cynanchifolia C.B. Clarke in Hook. f., Fl. Brit. India 4: 208. 1883

Ipomoea hardwickii (Spreng.) Hemsl., J. Linn. Soc., Bot. 26(174): 160. 1890

Distribution: Karnataka (throughout Eastern districts), throughout except drier W parts; Africa, Australia, China, Indonesia, Japan, Myanmar, Vietnam.

Ipomoea cairica (L.) Sweet, Hort. Brit., ed. 1, 287. 1827; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus cairicus L., Syst. Nat. (ed. 10) 2: 922. 1759

Ipomoea palmata Forssk., Fl. Aegypt.-Arab. 43. 1775; C.B.Clarke in Hook.f., Fl. Brit. India 4: 214. 1883.

Convolvulus tuberculatus Desr. in Lam., Encycl. 3(2): 545. 1792

Ipomoea stipulacea Jacq., Pl. Hort. Schoenbr. 2: 39, t. 199. 1797

Convolvulus heptaphyllus Rottler, Neue Schriften Ges. Naturf. Freunde Berlin 4: 196. 1803

Ipomoea pendula R.Br., Prodr. Fl. Nov. Holland. 486. 1810

Ipomoea tuberculata (Desr.) Roem. & Schult., Syst. Veg. 4: 208. 1819

Ipomoea pulchella Roth, Nov. Pl. Sp. 115. 1821

Convolvulus bellus Spreng., Syst. Veg. ed. 16, 1: 590. 1824

Convolvulus digitatus Roxb., Fl. Ind. 1: 479. 1832

Ipomoea sepiarea var. *stipulacea* (Koenig ex Roxb.) C. B. Clarke in Hook. f., Fl. Brit. India 4: 209. 1883

Distribution: Karnataka (throughout), throughout India; pantropical.; native to N Africa, Arabian Peninsula.

Local/Common/Vernacular names: *Cairo Morning Glory, Coast Morning Glory creeper, Messina Creeper, Mile-a-minute Vine, Railway creeper* (English); *Bekkina hejje balli, Kothi hejje balli* (Kannada)

Ipomoea carnea Jacq., Enum. Syst. Pl. 13. 1760; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Distribution: Karnataka (throughout plains in water bodies), throughout plains of India in moist area), native to tropical America.

Local/Common/Vernacular name: *The pink morning glory* (English)

Ipomoea cheirophylla O'Donell, Lilloa 29: 141. 1959

Ipomoea digitata L., Syst. Nat. ed. 10: 924. 1759; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Davanagere, Shivamogga, Uttara Kannada), throughout India; pantropical.

Local/Common/Vernacular names: *Bhuja gumbala, Bhumi chakra gedde, Gudde genasu* (Kannada)

Ipomoea cholulensis Kunth, Nov. Gen. Sp. ed. 4, 3: 112. 1818

Quamoclit coccinea (L.) Moench, Methodus 453. 1794

Distribution: Karnataka (cultivated & naturalised in plains), throughout hotter parts of India; pantropical; native to tropical America.

Ipomoea coptica (L.) Roth ex Roem. & Schult., Syst. Veg. 4: 208. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus copticus L., Mant. Pl. App. 559. 1771

Convolvulus stipulatus Desr. in Lam., Encycl. 3: 546. 1792

Ipomoea dissecta Willd., Phytographia 5: t. 2, f. 3. 1794; C.B.Clarke in Hook.f., Fl. Brit. India 4: 200. 1883.

Convolvulus thonningii Schumach. & Thonn., Beskr. Guin. Pl. 98. 1827

Distribution: Karnataka (Davanagere, Mysora, Uttara Kannada), Peninsula India; Australia, Africa, Sri Lanka.

Ipomoea eriocarpa R. Br., Prodr. 484. 1810; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Convolvulus hispidus Vahl, Symb. Bot. 3: 29.1794

Ipomoea sessiliflora Roth, Nov. Pl. Sp. 116.1821

Convolvulus eriocarpus (R.Br.) Spreng., Syst. Veg. 1: 598.1825

Convolvulus sessiliflorus (Roth) Spreng., Syst. Veg. 1: 599.1825

Ipomoea horsefieldiana Miq., Fl. Ned. Ind. 2: 611.1857

Ipomoea sindica Stapf, Bull. Misc. Inform. Kew 346. 1894

Distribution: Karnataka (throughout), throughout India; paleotropical.

Local/Common/Vernacular names: *Mulli balli* (Kannada); *Nakari, Seetha valli* (Sanskrit)

Ipomoea fistulosa Mart. ex Choisy, DC., Prodr. 9: 349.1845

Batatas crassicaulis Benth., Bot. Voy. Sulphur 134.1845

Ipomoea crassicaulis (Benth.) B. L. Rob., Proc. Amer. Acad. Arts 51(10): 530.1916

Ipomoea carnea subsp. *fistulosa* (Martius ex Choisy) D. F. Austin, Taxon 26: 237. 1977; B.D.Sharma et al., Fl. Karnataka Anal. 181. 1984.

Distribution: Karnataka (throughout), throughout India; pantropical.

Ipomoea grandifolia (Dammer) O'Donell, Arq. Mus. Paranaense 9: 222, f. 6. 1952

Jacquemontia grandifolia Dammer, Bot. Jahrb. Syst. 23(Beibl. 57): 41.1897

Distribution: Karnataka (Dakshina Kannada, Udupi), throughout drier parts of India; native to tropical America

Ipomoea hederifolia L., Syst. Nat. ed. 10:925.1759; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Ipomoea luteola Jacq., Collectanea 2: 266.1788

Ipomoea phoenicea Roxb., Fl. Ind., ed. 1,2: 92. 1824; Fl.Ind., ed.1: 502.1832.

Convolvulus luteolus (Jacq.) Spreng., Syst. Veg. 1: 599.1825

Convolvulus phoeniceus (Roxb.) Spreng., Syst. Veg. 1: 596.1825

Quamoclit phoenicea Choisy, Mém. Soc. Phys. Genève 6: 433. 1833

Quamoclit hederifolia (L.) G. Don, Gen. Hist. 4: 259.1838

Distribution: Karnataka (throughout), cultivated & naturalised throughout India; pantropical.; native to tropical America.

Local/Common/Vernacular names: *Ivy-leaved Morning Glory, Scarlet Creeper, Scarlet Morning Glory, Star Ipomoea* (English); *Halle balli, Kama lata, Nakshatra hoovu, Nalle balli* (Kannada)

Ipomoea horsfalliae Hook., Bot. Mag., t. 3315.1834; Anon., Beautiful Climb. Creep. Lalbagh 56. 2004.

Distribution: Karnataka (cultivated in Bengaluru), native to the Caribbean & Brazil.

Local/Common/Vernacular names: *Cardinal Creeper, Lady Doorly's Morning Glory* (English)

Ipomoea illustris Prain, Bengal Pl. 2: 735. 1903; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dharwad, Hassan, Mysuru, Shivamogga, Uttara Kannada), Maharashtra, West Bengal; SE Asia.

Local/Common/Vernacular names: *Karugani hambu, Kugini balli* (Kannada)

Ipomoea imperati (Vahl) Griseb., Cat. Pl. Cub. 203. 1866

Convolvulus littoralis L., Syst. Nat. ed. 10: 924.1759

Convolvulus imperati Vahl, Symb. Bot. 1: 17. 1790

Ipomoea stolonifera (Cirillo) J.F.Gmelin, Syst. Veg. 1: 345.1796

Ipomoea carnosa R. Br., Prodr. 485. 1810; C.B.Clarke in Hook.f., Fl. Brit. India 4: 213.1883.

Ipomoea acetosifolia (Vahl) Roem. & Schult., Syst. Veg. 4: 246.1819

Batatas littoralis (L.) Choisy, Mém. Soc. Phys. Genève 6: 506.1833

Batatas acetosifolia (Vahl) Choisy, Convolv. Orient. 124.1834

Convolvulus acetosifolius Vahl, Bull. Soc. Imp. Naturalistes Moscou 73.1840

Distribution: Karnataka (throughout coastal areas), throughout India along coastal areas; pantropical. & warm temperate littoral species.

Local/Common/Vernacular names: *Beach morning glory, Fiddle-leaf morning-glory, White-flower beach* (English)

- Ipomoea indica** (Burm.) Merr., Interpr. Herb. Amboin. 445. 1917
Convolvulus indicus Burm., Index Univ. Herb. Amb. 7: 6. 1755
Convolvulus acuminatus Vahl, Symb. Bot. 3: 26.1794
Ipomoea congesta R. Br., Prodr. 485.1810
Ipomoea mutabilis Lindl., Bot. Reg. 1: 39.1813
Ipomoea cathartica Poir. in Lam., Encycl., Suppl. 4(2): 633.1816
Ipomoea acuminata (Vahl) Roem. & Schult., Syst. Veg. 4: 228.1819
Convolvulus congestus (R.Br.) Spreng., Syst. Veg. 1: 601.1825
Ipomoea cataractae Endl., Prodr. Fl. Norfolk. 53. 1833
Pharbitis insularis Choisy, Mém. Soc. Phys. Genève 6: 439.1833
Ipomoea learii Knight ex Paxton in Paxton's Mag. Bot. 6: 267.1839
Ipomoea insularis (Choisy) Steud., Nomencl. Bot. (ed. 2) 1: 817. 1840
Pharbitis learii (Knight ex Paxton) Lindl., Edwards's Bot. Reg. 27:pl. 56 1841; Dalzell & A.Gibson, Suppl. 58.1861
Pharbitis acuminata (Vahl) Choisy in DC., Prodr. 9: 342.1845
Pharbitis acuminata var. *congesta* (R.Br.) Choisy, Prodr. 9: 343 1845
Pharbitis cathartica (Poiret) Choisy in DC., Prodr. 9: 342.1845
Ipomoea indica var. *acuminata* (Vahl) Fosberg, Bot. Not., Suppl. 129: 38.1976
Pharbitis indica (Burm.f.) R.C.Fang, Fl. Reipubl. Popularis Sin. 64(1): 105.1979
Ipomoea amoena Blume, Bijdr. Fl. Ned. Ind. 718.1985
 Distribution: Karnataka (cultivated), cultivated ? throughout India; circumtropical.
 Local/Common/Vernacular names: *Blue dawn flower*, *Blue morning glory*, *Island morning glory*, *Morning Glory*, *Ocean blue morning glory* (English)

Ipomoea laciniata C.B.Clarke in Hook f., Fl. Brit. India 4:200. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Pharbitis laciniata Dalzell in Hook., Kew J. Bot. 3:167.1851

Distribution: Karnataka (Belagavi, Chikmagaluru, Shivamogga), Maharashtra, Tamil Nadu; endemic.

Ipomoea littoralis (L.) Boiss., Fl Orient. 4: 112. 1879; K.G.Bhat & Vishwanatha, Indian J. For. 32: 175. 2009.

Convolvulus denticulatus Desr. in Lam., Encycl. 3(2): 540. 1789

Ipomoea denticulata (Desr.) Choisy, Mém. Soc. Phys. Genève 6: 447.1833; C.B.Clarke in Hook.f., Fl. Brit. India 4: 208. 1883.

Ipomoea choisiana Wight ex Safford, Contr. U.S. Natl. Herb. 9: 298.1905

Distribution: Karnataka (Dakshina Kannada, Udupi), Andaman & Nicobar Islands; Australia, Indonesia, Malaysia, Polynesia, Sri Lanka, Seychelles

Ipomoea marginata (Desr.) Verdc., Kew Bull. 42: 658. 1987

Convolvulus marginatus Desr. in Lam., Encycl. 3: 558. 1792

Convolvulus stipulaceus Roxb., Fl. Ind. 2:71.1824; Fl.Ind. 1: 484.1832.

Ipomoea sepiaria Koenig ex Roxb., Fl. Ind. 2: 90.1824; C.B.Clarke in Hook.f., Fl. Brit. India 4: 208. 1883.

Ipomoea verrucosa Blume, Bijdr. Fl. Ned. Ind. 13: 718.1825

Convolvulus verrucosus (Blume) D. Dietr., Syn. Pl. 1: 670.1839

Ipomoea subtrilobans Miq., Fl. Ned. Ind. 2: 615. 1856

Distribution: Karnataka (Chitradurga, Dharwad), throughout India; Paleotropical.

Local/Common/Vernacular names: *Hedge bind-weed*, *Purple heart glory* (English); *Nagini*, *Putra kanda*, *Thali kiru soppu* (Kannada); *Aaku karni*, *Ashra bindhu chhada*, *Laksmana*, *Manjika*, *Naga putri*, *Naga vha*, *Nagini*, *Puchhada*, *Putrada*, *Putrajanani*, *Putra kanda*, *Tulini* (Sanskrit)

Ipomoea mauritiana Jacq., Collectanea 4: 216.1790*Convolvulus paniculatus* L., Sp. Pl. 156. 1753

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kalaburgi, Udupi, Uttara kannad, Yadgir), Goa, Kerala, Maharashtra, Tamil Nadu; pantropical.

Local/Common/Vernacular names: *Ajigator* *yam*, *Giant potato*, *Milky yam*, *White yam* (English); *Bhoochakra gedde*, *Bhuja gumbala*, *Bhumi chakra gedde*, *Gudde genasu*, *Nel gumbala*, *Nela gudda genasu*, *Vidari* (Kannada); *Bhuksa maandi*, *Bhumi kush manda*, *Gandha phala*, *Gajavaji priya*, *Gajhist*, *Ksheera vidaari*, *Mahashweta*, *Payasvini*, *Shresta kan a*, *Shri galika*, *Vidari* (Sanskrit)

Ipomoea maxima Don ex Sweet, Hort. Brit. ed. 2:372.1830; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Distribution: Karnataka (Bengaluru, Chitradurga, Davanagere, Kalaburgi, Mysuru, Raichur, Uttara Kannada), Peninsular India.

Local/Common/Vernacular name: *Thali kire soppu* (Kannada)

Ipomoea muricata (L.) Jacq., Pl. Rar. Hort. Schoenbr. 3: 40.1798; Kesh. Murthy & Yogan., Fl. Coorg 300.1990.*Convolvulus muricatus* L., Mant. Pl. 1: 44.1767*Ipomoea bona-nox* var. *purpurascens* Ker Gawl., Bot. Reg. 4:t 290.1818*Calonyction muricatum* (L.) G. Don, Gen. Hist. 4: 264.1838

Distribution: Karnataka (Hassan, Kodagu, Mysuru), throughout India; pantropical.; native to S America.

Ipomoea nil (L.) Roth, Catal. Bot. 1: 36. 1797; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.*Convolvulus hederaceus* L., Sp. Pl. 1: 154.1753*Convolvulus nil* L., Sp. Pl. ed. 2. 1: 219. 1762*Ipomoea scabra* Forssk., Fl. Aegypt.-Arab. 44.1775

Ipomoea hederacea (L.) Jacq., Collectanea 1: 124, t. 36.1787; C.B.Clarke in Hook.f., Fl. Brit. India 4: 199. 1883.

Ipomoea coerulasca Roxb., Fl. Ind. 2: 90. 1824*Ipomoea setosa* Blume, Bijdr. Fl. Ned. Ind. 714.1825*Pharbitis hederacea* (L.) Choisy, Mém. Soc. Phys. Genève 6: 440.1833*Pharbitis nil* (L.) Choisy, Mém. Soc. Phys. Genève 6: 439.1833*Pharbitis variifolia* Decne., Nouv. Ann. Mus. 3: 390. 1834*Pharbitis diversifolia* Lindl. in Edwards's Bot. Reg. 23:t. 1988. 1837*Pharbitis barbata* G.Don, Gen. Hist. 4: 263. 1838*Pharbitis punctata* G.Don, Gen. Hist. 4: 263. 1838*Pharbitis purshii* G.Don, Gen. Hist. 4: 263. 1838*Ipomoea trichocalyx* Steud., Nomencl. Bot. ed. 2, 2(1): 819. 1840*Ipomoea hederacea* var. *himalaica* C.B.Clarke in Hook f., Fl. Brit. India 4:199. 1883*Ipomoea hederacea* var. *integrifolia* Hallier f., Jahrb. Hamburg. Wiss. Anst. Beih. 16: 3—42. 1899*Ipomoea nil* var. *setosa* (Blume) Boerl., Handl. Fl. Ned. Ind. 2(2): 511.1899

Distribution: Karnataka (cultivated & run wild throughout) cultivated & run wild throughout India; pantropical.; native to S America.

Local/Common/Vernacular names: *Morning Glory* (English); *Chita bogari*, *Gauri bheea*, *Kolli bheea*, *Musukina bogari* (Kannada); *Krishan bheea*, *Nila pushpin*, *Nikalni* (Sanskrit)

Ipomoea obscura (L.) Ker Gawl., Bot. Reg. 3: t. 239. 1817; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.*Convolvulus obscurus* L., Sp. Pl. ed. 2., 2: 220. 1762*Ipomoea insuavis* Blume, Bijdr. Fl. Ned. Ind. 716.1825*Ipomoea ocularis* Bartl., Linnaea 14(Lit.): 126. 1840*Ipomoea ochroleuca* Span., Linnaea 15: 340.1841

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular names: *Bili chita bogari*, *Bokadi*, *Godla jadi*, *Muguthi balli*, *Vaaji gandha* (Kannada); *Ajanthri*, *Lakshmana*, *Vacha gandha* (Sanskrit)

Ipomoea parasitica (Kunth) G. Don, Gen. Hist. 4: 275. 1838; Biju, Rheedia 12: 77. 2002.

Convolvulus parasiticus Kunth, Nov. Gen. Sp. ed 4, 3: 103. 1818

Ipomoea perlonga Robinson, Proc. Amer. Acad. Arts 22: 312. 1894; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Distribution: Karnataka (Dharwad, Uttara Kannada), Andhra Pradesh, Chattisgarh, Kerala, Maharashtra, Tamil Nadu; native to Americas.

Ipomoea pes-caprae (L.) R. Br. in Tuckey, Narr. Exped. Zaire 477. 1818; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Convolvulus pes-caprae L., Sp. Pl. 1: 159. 1753

Ipomoea biloba Forssk., Fl. Aegypt.-Arab. 44. 1775; C.B.Clarke in Hook.f., Fl. Brit. India 4: 212. 1883.

Convolvulus maritimus Desr. In Lam., Encycl. 3(2):550. 1792

Convolvulus bilobatus Roxb., Fl. Ind. 2: 73. 1824

Batatas maritimus(Desr.) Bojer, Hortus Maurit. 225. 1837

Distribution: Karnataka (common in coastal areas of Dakshina Kannada, Udupi, Uttara Kannada, introduced in Bengaluru, Kalaburgi, Kodagu, Mysuru, Raichur, Yadgir), throughout coastal areas & Islands of India, elsewhere introduced; pantropical. littoral species.

Local/Common/Vernacular names: *Goats foot creeper, Hare leaf, Rabbit weed, Red adambo, Sea side potato* (English); *Aadu balli, Adambu balli, Bangada balli, Kempu bangada balli* (Kannada); *Chagalantri, Gava patta, Manmatha, Mar valli, Maryada, Raktha pushpa, Sagara, Sagara mekala, Yugma patra* (Sanskrit); *Adumbu booru* (Tulu)

Ipomoea pes-tigridis L., Sp. Pl. 1: 162. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Ipomoea hepaticifolia L., Sp. Pl. 1: 161. 1753

Convolvulus pes-tigridis (L.) Spreng., Syst. Veg. 1: 592. 1825

Ipomoea capitellata Choisy, Mém. Soc. Phys. Genève 6: 457. 1833

Ipomoea pes-tigridis var. *hepaticifolia* (L.) C.B.Clarke in Hook.f., Fl. Brit. India 4: 204. 1883

Distribution: Karnataka (throughout), throughout India; Paleotropical.

Local/Common/Vernacular name: *Tiger's foot bind weed* (English); *Kama lathe* (Kannada); *Vyagra naki* (Sanskrit)

Ipomoea pileata Roxb., Fl. Ind. 2: 94. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Ipomoea involucrata P. Beauv., Fl. Oware 52, t. 89. 1817

Convolvulus pileatus (Roxb.) Spreng., Syst. Veg. 4(Cur. Post): 61. 1827

Distribution: Karnataka (Dakshina Kannada), Peninsula India eastwards to Assam; SE Asia, tropical Africa.

Ipomoea purpurea (L.) Roth, Bot. Abh. 27. 1787; Anon., Beautiful Climb. Creep. Lalbagh 53. 2004.

Convolvulus purpureus L., Sp. Pl. ed. 2. 1: 219. 1762

Pharbitis hispida Choisy, Convolv. Orient. 56. 1833

Pharbitis diversifolius Lindl. in Edwards's Bot. Reg. 23: t. 1988. 1837

Pharbitis purpurea (L.) Voigt, Hort. Suburb. Calcutt. 354. 1845

Ipomoea purpurea (L.) Lam., Tabl. Encycl. 1: 466. 1791; C. B. Clarke in Hook f., Fl. Brit. India 4: 199. 1883; Anon., Beautiful Climb. Creep. Lalbagh 53. 2004.

Distribution: Karnataka (cultivated & naturalised throughout), cultivated & naturalised throughout India; Pantropical.; native to Mexico & C America.

Local/Common/Vernacular names: *or Common Morning Glory, Purple* (English)

Ipomoea quamoclit L., Sp. Pl. 1: 159. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Convolvulus pinnatus Desr., Encycl. 3: 567. 1789

Convolvulus quamoclit (L.) Spreng., Syst. Veg. 1: 591. 1825

Quamoclit vulgaris Choisy, Mém. Soc. Phys. Genève 6: 434. 1834

Quamoclit pinnata (Desr.) Bojer, Hortus Maurit. 224. 1837

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical America.

Local/Common/Vernacular names: *American bell flower, China creeper, Cyprus vine, Indian forget-me-not, Indian pink, Red Jasmine* (English); *Handara hoovu, Kama lathe, Kasi ratna, Kempu malle, Kempu mallige* (Kannada); *Kama latha, Shwetha tahru latha* (Sanskrit)

Ipomoea rumicifolia Choisy, Mem. Soc. Phys. Geneve 4:447. 1833

Convolvulus lasiospermus Vis., Pl. Aeg. Nub. 13. t. 1. f. 2. 1836

Distribution: Karnataka (), Kerala, Tamil Nadu; Arabia, tropical Africa.

Ipomoea soluta Kerr, Bull. Misc. Inform. Kew 1941(1): 18. 1941; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Distribution: Karnataka (Shivamogga), Maharashtra; Laos, Myanmar, Thailand.

Ipomoea staphylina Roem. & Schult., Syst. Veg. 4: 249. 1819; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Ipomoea racemosa Roth, Nov. Pl. Sp. 115. 1821

Distribution: Karnataka (throughout), throughout India; Malaysia, Sri Lanka.

Local/Common/Vernacular name: *Jigune balli* (Kannada)

Ipomoea tricolor Cav., Icon. 3: 5, t. 208. 1795; Anon., Beautiful Climb. Creep. Lalbagh 55. 2004.

Ipomoea rubrocaerulea Hook., Bot. Mag. 8: t. 3297. 1834; T. Cooke, Fl. Bombay 2:252. 1903.

Convolvulus rubrocaeruleus (Hook.) Dietrich, Syn. Pl. 1: 670. 1839

Quamoclit mutica Choisy in DC., Prodr. 9: 335?1845

Pharbitis rubrocaeruleus (Hook.) Planch., Fl. Serres Jard. Eur. 9: 281, t. 966. 1853-1854

Distribution: Karnataka (cultivated in Bengaluru), cultivated in Gardens; native to Mexico.

Local/Common/Vernacular names: *Granny vine, Morning Glory* (English)

Ipomoea triloba L., Sp. Pl. 161. 1753; Pramila, Fl. Dharwad Dist. 314. 2011.

Convolvulus trilobus (L.) Desr., Encycl. 3(2): 564. 1792

Batatas triloba (L.) Choisy, Mém. Soc. Phys. Genève 8(1): 49. 1838

Ipomoea triloba var. *quinquefolia* Kuntze, Revis. Gen. Pl. 2: 466. 1891

Distribution: Karnataka (Dharwad), throughout tropical areas of India; Pantropical.; Native to tropical America.

Local/Common/Vernacular names: *Aiea morning glory, Little bell morning glory* (English)

Ipomoea tropica Santapau & V. Patel, J. Bombay Nat. Hist. Soc. 54: 799. 1957; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Distribution: Karnataka (Chikkamagaluru), Maharashtra; endemic.

Ipomoea tuberculata Ker Gawl., Bot. Reg. 1, t. 86. 1816.

Ipomoea dasysperma Jacq., Ecl. Pl. Rar. 1(9-10): 32, t. 89. 1816; C.B.Clarke in Hook.f., Fl. Brit. India 4: 210. 1883.

Convolvulus dasyspermus Spreng., Syst. Veg. ed. 16, 1: 591. 1824

Convolvulus pedatus Roxb., Fl. Ind. 2: 63. 1824; Fl. Ind. 1: 478. 1832.

Distribution: Karnataka (throughout), Deccan Peninsula extending up to Himachal Pradesh, also cultivated), tropical Africa & Asia.

Ipomoea turbinata Lag., Gen. Sp. Pl. 10. 1816; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Calonyction speciosum var. *muricatum* (L.) Choisy, Prodr. 9: 345. 1845

Distribution: Karnataka (Belagavi, Ballari, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Mysuru, Udupi), Andhra Pradesh, Chattisgarh, Himachal Pradesh, Kerala, Punjab, Sikkim, Tamil Nadu, Uttarakhand, also cultivated; America, China, S & SE Asia, Africa.

Local/Common/Vernacular name: *Purple moonflower* (English); *Tigane maari balli* (Kannada)

***Ipomoea violacea* L., Sp. Pl. 161. 1753**

Convolvulus grandiflorus Jacq., Hort. Bot. Vindob. 3: 39, t. 69. 1776

Convolvulus macranthus Kunth, Nov. Gen. Sp. ed.4, 3: 95. 1818

Ipomoea macrantha Roem. & Schult., Syst. Veg. 4: 251. 1819

Convolvulus tuba Schltdl., Linnaea 6(4): 735. 1831

Calonyction asperum Choisy, Convolv. Orient. 60. 1834

Calonyction grandiflorum (Jacq.) Choisy, Convolv. Orient. 60. 1834

Calonyction muticum Decne., Nouv. Ann. Mus. 3: 390. 1834

Ipomoea glaberrima Bojer ex Hook., J. Bot. 1: 357. 1834

Calonyction comespermum Bojer, Hortus Maurit. 228. 1837

Calonyction comorensis Bojer, Hortus Maurit. 228. 1837

Ipomoea tuba (Schltdl.) G. Don, Gen. Hist. 4: 271. 1837; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Calonyction tuba (Schltdl.) Colla, Mem. Nov. Sp. Calon. 15. 1840

Ipomoea grandiflora C.B.Clarke in Hook. f. Fl. Brit. Ind. 4: 198. 1883

Ipomoea grandiflora (Jacq.) Hallier f., Bot. Jahrb. Syst. 18(1-2): 153. 1893

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Peninsula India extending to Meghalaya, Uttar Pradesh; Pantropical.

***Ipomoea wightii* (Wall.) Choisy, Mém. Soc. Phys. Genève 6: 470. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.**

Convolvulus wightii Wall., Pl. Asiat. Rar. 2: 55, t. 171. 1831

Distribution: Karnataka (Chikkamagaluru, Mysuru), Andhra Pradesh, Kerala, Odisha, Tamil Nadu; endemic.

Jacquemontia* Choisy**Jacquemontia ovalifolia* (Choisy) Hallier f., Bot. Jahrb. Syst. 16: 543. 1892**

Convolvulus coeruleus Schum. & Thonn., Beskr. Guin. Pl. 101. 1827

Ipomoea ovalifolia Choisy, Mém. Soc. Phys. Genève 6: 449. 1834

Jacquemontia coerulea (Schum. & Thonn.) Choisy ex G. Don, Gen. Hist. 4: 283. 1838; C.B.Clarke in Hook.f., Fl. Brit. India 4: 220. 1883.

Distribution: Karnataka (cultivated throughout), cultivated in gardens; native to Africa, Mexico, USA.

***Jacquemontia paniculata* (Burm.) Hallier.f., Bot. Jahrb. Syst. 16: 541. 1893; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.**

Convolvulus parviflorus Vahl, Symb. Bot. 3: 29. 1793

Jacquemontia violacea (Vahl) Choisy, Mém. Soc. Phys. Genève 8(1): 61. 1838

Jacquemontia parviflora (Vahl) Roberty, Candollea 14: 32. 1952

Distribution: Karnataka (cultivated throughout), cultivated throughout India; Africa, Australia, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, New Guinea, Pacific Islands. Philippines, Sri Lanka, Thailand, Vietnam.

***Jacquemontia pentantha* (Jacq.) G. Don, Gen. Hist. 4: 283. 1838**

Convolvulus pentanthus Jacq., Collectanea 4: 210. 1791

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical S America.

Lettsomia* Roxb.**Lettsomia elliptica* (Roth) Wight ex C.B.Clarke in Hook. f., Fl. Brit. India 4: 192. 1883**

Ipomoea elliptica Roth, Nov. Pl. Sp. 113. 1821

Ipomoea laurifolia Sweet, Hort. Brit. 288. 1826

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Peninsula India; Sri Lanka.

Merremia Dennst. ex Endl.

Merremia emarginata (Burm.f.) Hallier f., Bot. Jahrb. Syst. 16(4-5): 552.1893

Evolvulus emarginatus Burm. f., Fl. Indica 77, t. 30, f.1. 1768

Convolvulus reniformis Roxb., Fl. Ind. 1: 481.1832

Ipomoea reniformis (Roxb.) Choisy, Mém. Soc. Phys. Genève 6: 446.1833; C.B.Clarke in Hook.f., Fl. Brit. India 4: 206.1883.

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chtradurga, Dakshina Kannada, Dharwad, Hassan, Udupi, Uttara Kannda, Vijayapura), extending from Deccan Peninsula to Punjab, West Bengal; SE Asia, tropical Africa.

Merremia gangetica Cufod., Bull. Jard. Bot. État Bruxelles 31(Suppl.): 743. 1961; Seetharam, Kotresh & Uploankar, Fl. Dist. 53. 2000.

Distribution: Karnataka (Bengaluru, Chitradurga, Davanagere, Dharwad, Kalaburgi, Tumakuru, Yadgir), throughout India; Paleotropical.

Local/Common/Vernacular name: *Ili kivi* (Kannada)

Merremia hederacea (Burm.) Hallier.f., Bot. Jahrb. Syst. 18: 118. 1894

Evolvulus hederaceus Burm., Fl. Ind. 77. t. 30, f. 2. 1768

Convolvulus acetosellifolius Desr., Encycl. 3: 564.1792

Convolvulus dentatus Vahl, Symb. Bot. 3: 25.1794

Convolvulus flavus Willd., Sp. Pl. 1(2):852. 1797; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.

Ipomoea chryseides Ker Gawler, Bot. Reg. 4: t. 270. 1818; C.B.Clarke in Hook.f., Fl. Brit. India 4: 206.1883.

Ipomoea dentata (Vahl) Roem. & Schult., Syst. Veg. 4: 789.1819

Convolvulus chryseides (KerGawler) Spreng., Syst. Veg. 1: 598.1825; C.B.Clarke in Hook.f., Fl. Brit. India 4: 206.1883.

Convolvulus rufescens Choisy, Convolv. Orient. 97. 1834

Lepistemon muricatum Span., Linnaea 15: 339.1841

Ipomoea acetosellifolia (Desr.) Choisy in DC., Prodr. 9: 383. 1845

Ipomoea subtriflora Zoll. & Moritzi, Syst. Verz. 51.1845

Merremia chryseides (KerGawler) Hallier f., Bot. Jahrb. Syst. 16(4-5): 552.1893; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984 ('chrysoides').

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Dharwad, Mysuru, Udupi, Uttara Kannada), throughout India; Paleotropical.

Neopeltis Wall.

Neopeltis malabarica Van Ooster, Blumea 5: 268. 1942; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Distribution: Karnataka (Uttara Kannada), Kerala, Maharashtra; endemic.

Neopeltis racemosa Wall. in Roxb., Fl. Ind. 2: 44. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Neopeltis ovata Wall., Numer. List No. 1323. 1829

Neopeltis bracteata Griff., Not. Pl. Asiat. 4: 285. 1854

Neopeltis intermedia Griff., Not. Pl. Asiat. 4: 285. 1854

Distribution: Karnataka (Uttara Kannada), Kerala, Indonesia, Malaysia, Myanmar.

Operculina S. Manso

Operculina turpethum (L.) S. Manso, Enum. Subst. Braz. 16. 1836

Convolvulus turpethum L., Sp. Pl. 1: 155. 1753

Convolvulus anceps L., Mant. Pl. 1: 43. 1767

Convolvulus triqueter Vahl, Symb. Bot. 3: 30. 1794

Ipomoea turpethum (L.) R. Br., Prodr. 485. 1810; B.D.Sharma et al., Fl. Karnataka Anal. 183. 1984.

Ipomoea anceps (L.) Roem. & Schult., Syst. Veg. 4: 231. 1819

Ipomoea triquetra Roem. & Schult., Syst. Veg., ed. 15 bis 4: 231. 1819

Spiranthera turpethum (L.) Bojer, Hortus Maurit. 226. 1837

Argyreia alulata Miq., Fl. Ned. Ind. 2: 587. 1856

Merremia turpethum (L.) Rendle in Oliv., Fl. Trop. Afr. 4: 102. 1905

Ipomoea turpethum var. *humilior* Haines, Bot. Bihar Orissa 2: 600. 1922

Operculina turpethum var. *humilior* (Haines) K.K.Khanna, Fl. Bihar, Analysis 340. 2001

Distribution: Karnataka (Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), throughout India; Asia, E Africa, Australia, introduced in N & S America, Pacific Islands, West Indies.

Porana Burm. f.

Porana malabarica C.B.Clarke in Hook. f., Fl. Brit. India 4: 223. 1883

Distribution: Karnataka (Dakshina Kannada, Mysuru, Uttara Kannada), Kerala, Maharashtra; endemic.

Poranopsis Roberty

Poranopsis paniculata (Roxb.) Roberty, Candollea 14: 26. 1952

Porana paniculata Roxb., Pl. Coromandel 3: 31. t. 235. 1815; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Dinetus paniculatus (Roxb.) Sweet, Hort. Brit. 289. 1826

Distribution: Karnataka (Bengaluru, Mysuru, Vijayapura), throughout India; Bhutan, China, Indonesia, Myanmar, Nepal, Pakistan.

Quamoclit Mill.

Quamoclit longiflora (Willd.) G. Don, Gen. Hist. 4: 259. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 182. 1984.

Ipomoea longiflora R. Br., Prodr. 484. 1810

Distribution: Karnataka (Uttara Kannada), throughout drier parts of India; native to tropical America.

Rivea Choisy

Rivea hypocarteriformis Choisy, Mém. Soc. Phys. Genève 6: 408. 1833; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Convolvulus candidans auct. non Roth, Roem. & Schult., Syst. Veg. 4: 302. 1819

Argyreia bona-nox Sweet, Hort. Brit. 289. 1827

Argyreia uniflor Sweet, Hort. Brit. 289. 1827

Letsomia bona-nox Roxb., Fl. Ind. 1: 494. 1832

Letsomia uniflora Roxb., Fl. Ind. 1: 495. 1832

Rivea bona-nox Choisy, Mém. Soc. Phys. Genève 4: 409. 1833

Distribution: Karnataka (throughout), throughout India in dry plains; Sri Lanka.

Local/Common/Vernacular names: *Common night glory, Midnapur creeper* (English)

Rivea ornata Choisy, Mém. Soc. Phys. Genève 4: 409. 1833; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.
Convolvulus candicans Roth, Nov. Pl. Sp. 106. 1821
Letsomia ornata Roxb., Fl. Ind. Ed. 1, 2: 86. 1824; Fl. Ind. ed. 2, 1: 496. 1832.
Argyreia ornata Sweet, Hort. Brit. 289. 1826
Convolvulus tarita Buch.-Ham. ex Wall., Numer. List No. 2253. 1830
 Distribution: Karnataka (Kolar, Vijayapura), Andhra Pradesh, Tamil Nadu; Sri Lanka.

Seddera Hochst.

Seddera latifolia Hochst. & Steud., Flora 27, Beil.: 8, t. 5. 1844
Breweria latifolia (Hochst.) Benth., Gen. Pl. 2: 877. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 180. 1984.
 Distribution: Karnataka (E Karnataka), W India; Arabia, E Africa, Socotra, Pakistan.

Stictocardia Hallier f.

Stictocardia beraviensis (Vatke) Hallier f., Bot. Jahrb. Syst. 18(1–2): 159. 1894
Ipomoea beraviensis Vatke, Linnaea 43: 514. 1882; Anon., Beautiful Climb. Creep. Lalbagh 50. 2004.
 Distribution: Karnataka (cultivated in Bengaluru, Dakshina Kannada, Udupi), native to tropical Africa.

Stictocardia tiliifolia (Desr.) Hallier f., Bot. Jahrb. Syst. 18: 159. 1894
Ipomoea campanulata L., Sp. Pl. 1: 160. 1753; C.B.Clarke in Hook.f., Fl. Brit. India 4: 211. 1883.
Ipomoea grandiflora (L.f.) Lam., Tabl. Encycl. 1: 467. 1791
Convolvulus tiliifolius Desr. in Lam., Encycl. 3: 544. 1792
Convolvulus gangeticus Roxb., Hort. Bengal. 13. 1814
Ipomoea benghalensis Roem. & Schult., Syst. Veg. 4: 229. 1819
Ipomoea tiliifolia (Desr.) Roem. & Schult., Syst. Veg. 4: 229. 1819
Convolvulus campanulatus (L.) Spreng., Syst. Veg. 1: 608. 1825
Ipomoea gangetica Sweet, Hort. Brit. 288. 1830
Rivea tiliifolia (Desr.) Choisy, Mém. Soc. Phys. Genève 6: 407. 1833
Convolvulus platypeltis Zipp. ex Span., Linnaea 15: 338. 1841
Argyreia tiliifolia (Desr.) Wight, Icon. Pl. Ind. Orient. 4: 12, t. 1358. 1848; C.B.Clarke in Hook f., Fl. Brit. India 4: 184. 1883.
 Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Dharwad, Shivamogga, Uttara Kannada), throughout plains of India except dry Western areas; Australia, N & C America, S & SE Asia, West Indies, Pacific Islands.
 Local/Common/Vernacular name: *Kari ugunia hambu* (Kannada)

Turbina Raf.

Turbina corymbosa (L.) Raf., Fl. Tellur. 4: 81. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.
Convolvulus corymbosus L., Syst. Nat. (ed. 10) 923. 1759
Legendrea mollissima Webb & Berthel., Hist. Nat. Îles Canaries 3: 27, t. 137. 1844; T. Cooke, Fl. Bombay 2: 261. 1903.
 Distribution: Karnataka (Uttara Kannada, Vijayapura) cultivated India; native to Canary Islands.

Xenostegia D.F.Austin & Staples

Xenostegia tridentata (L.) D.F.Austin & Staples, Brittonia 32: 533. 1980.
Convolvulus tridentatus L., Sp. Pl. 1: 157. 1753
Evolvulus tridentatus (L.) L., Sp. Pl. ed. 2: 392. 1762
Ipomoea tridentata (L.) Roth, Arch. Bot. (Leipzig) 1(2): 38. 1798; C.B.Clarke in Hook.f., Fl. Brit. India 4: 205. 1883.
Merremia tridentata (L.) Hallier f., Bot. Jahrb. Syst. 16(4–5): 552, in obs. 1893

subsp. *tridentata*

Distribution: Karnataka (Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kalaburgi, Kolar, Mysuru, Raichur, Tumakuru, Udupi, Uttara Kannada, Vijayapura), throughout except NW India; Paleotropical.

subsp. *hastata* (Ooststr.) Parmar, J. Econ. Taxon. Bot. 18: 251. 1994.

Merremia hastata Hallier f., Bot. Jahrb. Syst. 16: 552. 1893

Merremia tridentata subsp. *hastata* (Hallier f.) Ooststr., Blumea 3(2): 317. 1939; B.D.Sharma et al., Fl. Karnataka Anal. 184. 1984.

Distribution: Karnataka (Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kalaburgi, Kolar, Mysuru, Raichur, Tumakuru, Udupi, Uttara Kannada, Vijayapura), throughout except NW India; Borneo, Caroline Is., China Southeast, Jawa, Lesser Sunda Is., Malaya, Maluku, Philippines, Sri Lanka, Sulawesi, Sumatera, Taiwan.

Cornaceae

Alangium Lam.

Alangium salviifolium (L.f.) Wangerin in Engl., Pflanzenreich Alangiac. 9. 1910; B.D.Sharma et al., Fl. Karnataka Anal. 121. 1984; Gamble, Fl. Pres. Madras 572 (404). 1919; C.J.Saldanha, Fl. Karnataka 2: 64. 1996.

Grewia salviifolia L. f., Suppl. Pl. 409. 1781

Alangium lamarckii Thwaites, Enum. Pl. Zeyl. 133. 1859

subsp. *salviifolium*; C.J.Saldanha, Fl. Karnataka 2: 64. 1996

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Dharwad, Kalburgi, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura); Cambodia, China, Indonesia, Laos, Malaysia, Nepal, Philippines, Sri Lanka, SE Africa, Thailand, Vietnam.

Local/Common/Vernacular names: *Sage-leaved alangium* (English); *Ankola, Ankule mara* (Kannada)

subsp. *hexapetalum* (Lam.) Wangerin in Engl., Pflanzenreich Alangiac. 9. 1910; C.J.Saldanha, Fl. Karnataka 2: 64. 1996

Plate 6A

Alangium hexapetalum Lam., Encycl. 1: 175. 1783

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga), Kerala, Tamil Nadu; Indomalasia.

Local/Common/Vernacular name: *Guddada goni* (Kannada)

Mastixia Blume

Mastixia arborea (Wight) C.B.Clarke in Hook.f., Fl. Brit. India 2(6): 745. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 121. 1984.

Mastixia pentandra auct. non Blume, 1825; Ramaswamy et al., Fl. Shimoga 266. 2001.

Bursinopetalum arboreum Wight, Icon. Pl. Ind. Orient. 3: t. 956. 1845

subsp. arborea

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Gulla, Gulle mara* (Kannada)

subsp. **meziana** (Wangerin) K.M.Matthew ex Ramamoorthy, Blumea 23: 89. 1976; C.J.Saldanha, Fl. Karnataka 2: 66. 1996.

Mastixia meziana Wangerin, Feddes Repert. 4: 336. 1907

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Costaceae

Chamaecostus C.D.Specht & D.W.Stev.

Chamaecostus cuspidatus (Nees & Mart.) C.D.Specht & D.W.Stev., Taxon 55: 158.2006

Globba cuspidata Nees & Mart., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 11: 28.1823

Costus igneus N.E.Br., Ill. Hort. 31: 25.1884

Costus cuspidatus (Nees & Mart.) Maas, Acta Bot. Neerl. 24: 469.1976

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Brazil.

Local/Common/Vernacular names: *Fiery Costus, Insulin plant, Spiral Flag* (English)

Cheilocostus C.Specht

Cheilocostus speciosus (J.Koenig) C.Specht, Taxon 55: 159.2006

Banksea speciosa J.König, Observ. Bot. 3: 75.1783

Costus speciosus (J.Koenig) Sm., Trans. Linn. Soc. London 1: 249.1791; B.D.Sharma et al., Fl. Karnataka Anal. 280. 1984.

Tsiana speciosa (J.Koenig) J.F.Gmel., Syst. Nat. 2: 9.1791

Planera speciosa (J.Koenig) Giseke, Prael. Ord. Nat. Pl. 244.1792

Kaempferia speciosa (J.Koenig) Thunb., Fl. Jav. 8 1825

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakhsina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Peninsular to NE India & Andaman & Nicobar Islands; E, S & SE Asia.

Local/Common/Vernacular names: *Chamlasulli, Changaala koshtha, Nari kabbu* (Kannada); *Kosta, Piyo kando (Kunabi); Kemuka* (Sanskrit)

Costus L.

Costus pictus D.Don, Edwards's Bot. Reg. 19: t. 1594.1833; K. G. Bhat, Fl. S. Kanara 145. 2014.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to C & S America.

Local/Common/Vernacular name: *Insulin plant* (English)

Crassulaceae

Bryophyllum Salisb.

Bryophyllum pinnatum (Lam.) Oken, Allg. Naturgesch. 3: 1966. 1841; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Cotyledon pinnata Lam., Dict. 2: 141.1786

Bryophyllum calycinum Salisb., Parad. Lond. t.3. 1805

Kalanchoe pinnata (Lam.) Pers., Syn. 1: 446. 1805

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Himachal Pradesh, Maharashtra, Odisha, Tamil Nadu, West Bengal; pantropical.

Local/Common/Vernacular names: *Air plant, Cathedral bells, Goethe plant Life plant, Miracle leaf* (English); *Kadubasale* (Kannada)

Crassula L.

Crassula schimperi Fisch. & C.A.Mey., Index Seminum (LE) 8: 56.1842

Tillaea pentandra Royle ex Edgew., Trans. Linn. Soc. 20: 50. 1846; C.J.Saldanha, Fl. Karnataka 1: 363. 1984.

Tillaea schimperi (Fisch. & C.A. Mey.) M.G.Gilbert, H. Ohba & K.T. Fu, Novon 10(4): 366. 2000

Distribution: Karnataka (Mysuru); Bhutan, China, Jammu & Kashmir, Nepal, Pakistan, SW Asia, tropical Africa,

Kalanchoe Adans.

Kalanchoe bhidei T. Cooke., Fl. Bombay 1: 467. 1903; B.D.Sharma et al., Fl. Karnataka Anal. 98. 1984.

Distribution: Karnataka (Bengaluru, Chikkaballapur, Chitradurga, Hassan, Kolar, Mandya, Mysuru), Maharashtra, Tamil Nadu; endemic.

Kalanchoe blossfeldiana Poelln., Repert. Spec. Nov. Regni Veg. 35(918/927): 159 1934

Distribution: Karnataka (Chitradurga), cultivated throughout India; native to Madagascar; cultivated throughout tropics.

Local/Common/Vernacular names: *Christmas kalanchoe, Flaming Katy, Florist kalanchoe, Madagascar widow's-thrill* (English)

Kalanchoe integra (Medik.) Kuntze, Rev. Gen. 1: 229. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Cotyledon integra Medik., Acta Acad. Theod. Palat. 3: 200. t. 9. 1775

Kalanchoe spatulata DC., Pl. Hist. Succ. T. 65. 1801

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Mysuru, Vijayapura), Jammu & Kashmir, Maharashtra, Mizoram, Tripura, Uttar Pradesh, Uttarakhand; Bhutan, Cambodia, China, Indonesia, Laos, Malaysia, Nepal, Philippines, Sikkim, Thailand, Vietnam, W Asia; introduced elsewhere.

Kalanchoe laciniata (L.) DC., Pl. Hist. Succ. t. 100 1802; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Cotyledon laciniata L., Sp. Pl. 430. 1753

Kalanchoe floribunda Wight & Arn., Prodr. Fl. Ind. Orient. 1: 359. 1834

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Vijayapura), Andhra Pradesh, Arunachal Pradesh, Bihar, Kerala, Maharashtra, Tamil Nadu, Tripura, West Bengal; Africa, S America, West Asia.

Local/Common/Vernacular names: *Christmas tree plant, Fir-tree kalanchoe* (English); *Gandukaalinga, Nununni gida* (Kannada)

Kalanchoe lanceolata (Forssk.) Pers., Syn. Pl. 1: 446 1805; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Cotyledon lanceolata Forssk., Fl. Aegypt.-Arab. 89 1775

Kalanchoe glandulosa Hochst. ex A.Rich., Fl. Abyss. 1: 312. 1847

Distribution: Karnataka (Belagavi, Dharwad, Kolar, Mysuru, Tumakuru, Vijayapura), Andhra Pradesh, Bihar, Madhya Pradesh, Maharashtra, Tamil Nadu; Pantropical.

Local/Common/Vernacular name: *Kumudagedde* (Kannada)

Kalanchoe olivacea Dalzell, Bombay Fl. 313. 1861; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Distribution: Karnataka (Belagavi, Chikkaballapur, Kalburgi, Kolar, Mysuru), Maharashtra, Tamil Nadu; endemic.

Kalanchoe tubiflora (Harv.) Raym.-Hamet, Beih. Bot. Centralbl. 29(2): 41 1912; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Bryophyllum tubiflorum Harv., Fl. Cap. 2: 380. 1862

Distribution: Karnataka (Cultivated, Belagavi), cultivated throughout plains; native to Madagascar.

Umbilicus DC.

Umbilicus rupestris (Salisb.) Dandy in Riddelsdell, Hedley & Price, Fl. Gloucestershire 611. 1948;
C.J.Saldanha, Fl. Karnataka 1: 363. 1984

Cotyledon rupestris Salisb., Prodr. 307. 1796

Distribution: Karnataka (cultivated in Bengaluru, Chikkamagaluru), cultivated in gardens.' Native to Europe.

Local/Common/Vernacular names: *Navelwort, Penny-pies, Wall Pennywort* (English)

Cucurbitaceae**Benincasa Savi**

Praecitrullus Pangalo

Benincasa fistulosa (Stocks) H.Schaef. & S.S.Renner, Taxon 60: 133. 2011

Citrullus fistulosus Stocks, Hooker's J. Bot. Kew Gard. Misc. 3: 74, t. 3. 1851

Citrullus vulgaris var. *fistulosus* (Stocks) J.L.Stewart, Punjab Pl. 96. 1869

Praecitrullus fistulosus (Stocks) Pangalo, Bot. Zhurn. S.S.S.R. 29: 203. 1944

Colocynthis citrullus var. *fistulosus* (Stocks) Chakrav., Rec. Bot. Surv. India 17(1): 116. 1959

Citrullus lanatus var. *fistulosus* (Stocks) Chakrav., Fasc. Fl. India 11: 23. 1982

Distribution: Karnataka (cultivated throughout), cultivated throughout plains of India; Pakistan; cultivated in the entire range of distribution and outside.

Local/Common/Vernacular name: *White or Ash gourd* (English); *Boodu Kumbala* (Kannada); *Kooshmanda* (Sanskrit)

Benincasa hispida (Thunb.) Cogn. in A. & C. DC., Monogr. Phan. 3: 513. 1881; C.J.Saldanha, Fl. Karnataka 1: 307. 1984

Cucurbita pruriens Parkinson, J. Voy. South Seas 44. 1773

Cucurbita hispida Thunb., Nov. Acta Regiae Soc. Sci. Upsal. 4: 38. 1783

Benincasa cerifera Savi, Biblioth. Ital. 9: 158. 1818

Cucurbita alba Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 344. 1834

Benincasa pruriens f. *hispida* (Thunb.) W.J.de Wilde & Duyfjes, Sandakania 17: 47. 2008

Distribution: Karnataka (cultivated throughout), throughout India; throughout tropical Asia; presumably wild in Indonesia.

Local/Common/Vernacular names: *Ash gourd, Ash or White pumpkin, Chinese preserving melon, Tallow gourd, The winter melon, Wax gourd, White gourd, Winter gourd* (English); *Bhoodu kumbala kayi* (Kannada)

Blastania Kotschy & Peyr. 3

Blastania garcinii (Burm.f.) Cogn. in A. & C. DC., Monogr. Phan. 3: 629. 1881

Sicyos garacini Burm.f., Fl. Ind. 211. 1768

Bryonia garcinii (Burm.f.) Willd., Sp. Pl. 4(1): 623. 1805 ('garcini').

Pilogyne garcini (Burm.f.) Wight, Ann. Mag. Nat. Hist. 8: 268 1843

Ctenolepis garcinii (Burm.f.) Benth. & Hook.f., Gen. Pl. 1(3): 832. 1867; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.

Distribution: Karnataka (kalburgi, Raichur, Vijayapura), Andhra Pradesh, Delhi, Gujarat, Haryana, Kerala, MadhyaPradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Sri Lanka.

Cayaponia Silva Manso

Cayaponia laciniosa (L.) C.Jeffrey, Kew Bull. 15: 346 1962

Bryonia laciniosa L., Sp. Pl. 1013 1753

Aechmandra zeylanica Thwaites, Enum. Pl. Zeyl. 2: 125. 1859

Cerasiocarpum zeylanicum Hook.f., Gen. Pl. 1: 832. 1867.

Distribution: Karnataka (Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Odisha, Tamil Nadu, West Bengal; Bangladesh, Myanmar, Sri Lanka

Citrullus Schrad.

Citrullus colocynthis (L.) Schrad., Linnaea 12: 414 1838; B.D.Sharma et al., Fl. Karnataka Anal. 111. 1984.

Cucumis colocynthis L., Sp. Pl. 1011.1753

Colocynthis vulgaris Schrad., Index Seminum (GOET) 1833: 2 1833

Distribution: Karnataka (Ballari, Chitradurga, Davanagere, Kalaburagi, Raichur, Shivamogga, Uttara Kannada, Yadgiri), Andhra Pradesh, Assam, Bihar, Delhi, Goa, Gujarat, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Africa, Afghanistan, Myanmar, Middle East, Pakistan, Sri Lanka, W Asia.

Local/Common/Vernacular name: *Havu mekke kayi* (Kannada)

Citrullus lanatus (Thunb.) Matsum. & Nakai, Index Seminum (TI) 1916: 30.1916; B.D.Sharma et al., Fl. Karnataka Anal. 111. 1984.

Cucurbita citrullus L., Sp. Pl. 1010. 1753

Momordica lanata Thunb., Prodr. Pl. Cap. 13. 1794

Citrullus vulgaris Schrad. ex Eckl. & Zey., Enum. Pl. Afr. Austral. 279. 1836

Citrulus fistulosus Stocks in Hook., J. Bot. 3: 74. 1851

Citrullus colocynthis var. *lanatus* (Thunb.) Matsum. & Nakai, Index Seminum (TI) 1916: 30 1916

Citrullus lanatus var. *fistulosus* (J.L.Steward) Babu, Herb. Fl. Dehra Dun 194. 1977

Citrullus lanatus var. *albidus* (Chakrav.) Maheshw., Indian J. Forest. 1: 180.1978

Citrullus lanatus var. *minor* (Chakrav.) Maheshw., Indian J. Forest. 1: 180.1978

Citrullus lanatus var. *oblongus* (Chakrav.) Maheshw., Indian J. Forest. 1: 181. 1978

Citrullus lanatus var. *pulcherrimus* (Chakrav.) Maheshw., Indian J. Forest. 1: 180.1978

Citrullus lanatus var. *pumilus* (Chakrav.) Maheshw., Indian J. Forest. 1: 181. 1978

Citrullus lanatus var. *rotundus* (Chakrav.) Maheshw., Indian J. Forest. 1: 180. 1978

Citrullus lanatus var. *shami* (Chakrav.) Maheshw., Indian J. Forest. 1: 181. 1978

Citrullus lanatus var. *variegatus* (Chakrav.) Maheshw., Indian J. Forest. 1: 180. 1978

Citrullus lanatus var. *virgatus* (Chakrav.) Maheshw., Indian J. Forest. 1: 181. 1978

Citrullus lanatus var. *viridis* (Chakrav.) Maheshw., Indian J. Forest. 1: 180. 1978

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Davanagere, Hassan, Mysuru, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Assam, Bihar, Jharkhand, Delhi, Gujarat, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand, West Bengal; native to tropical Africa; cultivated all over the world.

Local/Common/Vernacular names: *Water melon* (English); *Dodda hale, Havu makke kayi, Indravaruni, Kallangadi hannu* (Kannada); *Bacchangaayi* (Tulu)

Coccinia Wight & Arn.

Coccinia grandis (L.) Voight, Hort. Suburb. Calcutt. 59. 1845; B.D.Sharma et al., Fl. Karnataka Anal. 111. 1984.

Bryonia grandis L., Mant. 1: 126. 1767

Momordica monadelpha Roxb., Fl. Ind. ed.2, 3: 708. 1832

Coccinia indica Wight & Arn., Prodr. Fl. Ind. Orient. 1: 347. 1834

Cucurbita dioica Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 347. 1834

Coccinia wightiana M.Roem., Fam. Nat. Syn. Monogr. 2: 93. 1846

Cephalandra indica Naudin, Ann. Sci. Nat. Bot. ser. 5, 5 : 16.1866

Cephalandra grandis Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 46(2): 103. 1877

Cephalandra indica var. *palmata* C.B.Clarke in Hook. f., Fl. Brit. India 2: 621. 1879

Coccinia cordifolia var. *wightiana* (M.Roem.) Cogn. in A. & C. DC., Monogr. Phan. 3: 531. 1881

Coccinia grandis var. *wightiana* (M.Roem.) Greb. in R. Mansfeld & J.Schultze-Motel, Verz. Landwirtsch. u. Gaertn. Kulturpfl. 2: 929. 1986

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Kalaburagi, Mysuru, Vijayapura), throughout India including Islands; Africa, E, S & SE Asia.

Local/Common/Vernacular names: *Baby watermelon*, *Gentleman's toes*, *Ivy gourd*, *Little gourd*, *Tindora* (English); *Kage tonde*, *Sihithonde*, *Thonde balli*, *Thonde kaayi* (Kannada); *Manoli* (Tulu)

Corallocarpus Welw. ex Benth. & Hook. f.

Corallocarpus conocarpus (Dalzell & A.Gibson) Hook.f. in Hook. f., Fl. Brit. India 2: 628. 1879; C.J.Saldanha, Fl. Karnataka 1: 294. 1984

Aechmandra conocarpa Dalzell & A.Gibson, Bombay. Fl. 100.1861

Distribution: Karnataka (Dharwad, Tumakuru), Gujarat, Maharashtra, Rajasthan, Tamil Nadu; Pakistan ?

Corallocarpus epigaeus (Rottler) Hook.f. in Hook. f., Fl. Brit. India 2: 628. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.

Bryonia epigaea Rottler Ges., Naturf. Freunde Berlin Neue Schriften ser. 4. 212. 1803

Bryonia sinuata Wall., Numer. List 6711D 1832

Aechmandra epigaea (Rottler) Arn., J. Bot. 3: 274. 1841

Rhynchosarpa epigaea (Rottler) Nandin, Ann. Sci. Nat. Bot. ser. 4, 10 : 179. 1862

Rhynchosarpa epigaea var. *gracilipes* Naudin, Ann. Sci. Nat. Bot. ser. 4, 16: 179. 1862

Corallocarpus gracilipes (Naudin) Cogn. in A. & C. DC., Monogr. Phan. 3: 656. 1881

Corallocarpus palmatus Cogn. in A. & C. DC., Monogr. Phan. 3: 648. 1881; Chakrav., Rec. Bot. Surv. Ind. 17(1): 170. 1959.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Kalaburagi, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Andhra Pradesh, Assam, Bihar, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Puducherry, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Pakistan, Sri Lanka, tropical East Africa.

Local/Common/Vernacular name: *Akashagaruda balli* (Kannada)

Ctenolepis Hook.f.

Ctenolepis garcini (L.) C.B.Clarke in Hook.f., Hook.f., Fl. Brit. India 2: 629. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.

Sicyos garcini Burm. f., Fl. Indica 211. 1768

Blastania garcini (L.) Cogn. in A. & C. DC., Monogr. Phan. 3: 629. 1881

Distribution: Karnataka (Belagavi, Bengaluru, Kalaburagi, Kolar, Mandya, Raichur, Vijayapura), Andhra Pradesh, Delhi, Gujarat, Haryana, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Sri Lanka.

Cucumis L.

Cucumis hystrix Chakrav., J. Bombay Nat. Hist. Soc. 50: 896. 1952

Cucumis muriculatua Chakrav., J. Bombay Nat. Hist. Soc. 50: 896. 1952

Distribution: Karnataka (Chitradurga), Arunachal Pradesh, Assam, Meghalaya, Mizoram, Nagaland; Bangladesh, China, Myanmar, Thailand.

Cucumis indicus Ghebret. & Thulin, Novon 17: 177 2007

Melothria ritchiei Chakrav., J. Bombay Nat. Hist. Soc. 50: 898. 1952

Cucumella ritchiei (Chakrav.) C.Jeffrey, Kew Bull. 19: 215. 1965; C.J.Saldanha, Fl. Karnataka 1: 295. 1984

Distribution: Karnataka (Belagavi, Shivamogga), Kerala, Maharashtra; endemic.

Cucumis leiospermus (Wight & Arn.) Ghebret. & Thulin, Novon 17: 177 2007

Bryonia leiosperma Wight & Arn., Prodr. Fl. Ind. Orient. 1: 1: 345. 1834; C.J.Saldanha, Fl. Karnataka 1: 301. 1984

Mukia leiosperma (Wight & Arn.) Wight, Ann. Mag. Nat. Hist. ser. 8: 208. 1842

Melothria leiosperma (Wight & Arn.) Cogn. in A. & C. DC., Monogr. Phan. 3: 622. 1881

var. **leiospermus**

Distribution: Karnataka (Mysuru), Andhra Pradesh, Arunachal Pradesh, Assam, Himachal Pradesh, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Nagaland, Punjab, Rajasthan, Sikkim, Tamil Nadu, Uttar Pradesh; China, Malaysia, Myanmar, Sri Lanka, Taiwan.

var. **triloba** (C.B.Clarke) Sanjappa *Comb. nov.*

Mukia leiosperma var. *triloba* (C.B.Clarke) in Hook. f., Fl. Brit. India. 2: 1879; Prain, Beng. Pl. 1: 527. 1903

Distribution: India (throughout), Malaysia, Myanmar.

Cucumis maderaspatanus L., Sp. Pl. 1012. 1753

Bryonia cordifolia L., Sp. Pl. 2: 1012. 1753

Cucumis maderaspatanus L., Sp. Pl. 1012. 1753

Bryonia scabrella L.f., Suppl. Pl. 424. 1782

Bryonia maderaspatana (L.) Lam., Encycl. 1: 496 1785

Bryonia callosa Rottler, Neue Schrift. Ges. Nat. Freunde Zu Berlin 4: 210. 1803 ('colosa').

Bryonia gracilis Wall., Numer. List 6714. 1832

Bryonia scabra Rottler ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 345 1834

Mukia scabrella (L.f.) Arn., J. Bot. 3: 276. 1841; C.B.Clarke in Hook.f., Fl. Brit. India 2: 623. 1879 excl.syn.

Mukia maderaspatana (L.) M.Roem., Fam. Nat. Syn. Monogr. 2: 47. 1846; B.D.Sharma et al., Fl. Karnataka Anal. 114. 1984.

Mukia rottleri M.Roem., Fam. Nat. Syn. Monogr. 2: 47. 1846

Coccinia cordifolia (L.) Cogn. in A. & C. DC., Monogr. Phan. 3: 623. 1881

Melothria maderaspatana (L.) Cogn. in A. & C. DC., Monogr. Phan. 3: 623. 1881; Cooke, Fl. Pres. Bombay 1: 539. 1903.

Distribution: Karnataka (throughout), throughout India; Andhra Pradesh, Arunachal Pradesh, Bihar, Delhi, Gujarat, Himachal Pradesh, Madhya Pradesh, Maharashtra, Mizoram, Rajasthan, Tamil Nadu, Tripura; Bhutan, China, Myanmar, Nepal, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Gubbi Sauthé kayi*, *Kaduvate balli*, *Mani tonde* (Kannada)

- Cucumis melo** L., Sp. Pl. 1011. 1753; C.J.Saldanha, Fl. Karnataka 1: 296. 1984
Cucumis pubescens Willd., Sp. Pl. 4 : 614. 1805; Gamble, Fl. Madras 3:535. 1919.
Cucumis momordica Roxb., Fl. Ind.3: 720 1832
Cucumis trigonus Roxb., Fl. Ind. 2 :722. 1832; C.B.Clarke in Hook.f., Fl. Brit. India 2: 619. 1879.
Cucumis turbinatus Roxb., Fl. Ind. 3: 723.1832
Cucumis utilissimus Roxb., Fl. Ind. 3: 721.1832
Cucumis pyriformis Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 342 1834
Momordica sativa Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 341. 1834
Cucumis aromaticus Royle, Illust. Bot. Himal. Mts. 220 1835
Cucumis pseudocolocynthis Royle, Illust. Bot. Himal. Mts. 220 1835
Cucumis melo var. *agrestis* Naudin, Ann. Sci. Nat. Bot. ser.4, 11:73.1859 & 12: 110. 1859; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
Cucumis melo var. *saharunporensis* Naudin, Ann. Sci. Nat., Bot., sér. 5 5: 11.1866; T.Cooke, Fl. Bombay 1: 535, 1903.
Cucumis callosus (Rottler) Cogn. in Engl., Pflanzenr. 4. 275, 2: 129. 1924; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
 Distribution: Karnataka (cultivated in throughout), cultivated and wild throughout India; cultivated throughout the world.
 Local/Common/Vernacular names: *Muskmelon* (English); *Budame kayi Halmekke balli, Ibbudala, Karbbuja, Kekkerike, Mekke kayi, Mekke kayi balli, Sauthé* (Kannada)
- Cucumis prophetarum** L., Cent. Pl. I 32 1755; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
 Distribution: Karnataka (Ballari, Chikkamagaluru, Hassan, Kalaburagi, Kodagu, Mysuru, Uttara Kannada), Andhra Pradesh, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu; Africa, Middle East, Pakistan, W Asia.
- Cucumis ritchiei** (C.B.Clarke) Ghebret. & Thulin, Novon 17: 178 2007
Dicoelospermum ritchiei C.B.Clarke in Hook.f., Fl. Brit. India 2:630. 1879; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Shivamogga), Gujarat, Kerala, Maharashtra, Punjab, Tamil Nadu; endemic.
- Cucumis sativus** L., Sp. Pl. 1012. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
Cucumis hardwickii Royle, Ill. Bot. Himal. Mts. 220.1835
Cucumis setosus Cogn. in A. & C. DC., Monogr. Phan. 3: 491. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
 Distribution: Karnataka (cultivated throughout), throughout India; cultivated throughout tropics, native to N India.
 Local/Common/Vernacular names: *Cucumber* (English); *Mullu sauthe, Sauthe balli, Sauthe kaayi* (Kannada)

Cucurbita L.

- Cucurbita argyrosperma** C.Huber, Cat. Graines 8. 1867
 Distribution: Karnataka (cultivated), cultivated throughout India; native to C America.
 Local/Common/Vernacular names: *Cilver-seed gourd* *Cushaw pumpkin, Japanese pie pumpkin, Pipian* (English)
- Cucurbita maxima** Duchesne in Lam., Encycl. 2: 151.1786; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.
Cucurbita maxima var. *badagarensis* Mudaliar, J. Bombay Nat. Hist. Soc. 49: 242.1950
 Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to C America; cultivated throughout the World.
 Local/Common/Vernacular names: *Squash gourd* (English); *Kumbala Sigumbala, Sihi kumbala kaayi* (Kannada)

Cucurbita moschata Duchesne, Ess. Hist. Nat. Courges 7 1786; Datar et al., Rheedia 134. 2005

Cucurbita pepo var. *moschata* (Duchesne) Duchesne, Encycl. 2: 152 1786

Distribution: Karnataka (throughout), throughout India; cultivated throughout tropics & subtropics, native to C & S America.

Local/Common/Vernacular names: *Butternut Squash*, *Canada crookneck Canada pumpkin*, *China Crook-neck Squash*, *Crook-neck Squash*, *musky gourd*, *musky pumpkin*, *musky squash*, *West Indian pumpkin*, *winter crookneck squash*, *winter squash* (English); *Kumbala* (Kannada)

Cucurbita pepo L., Sp. Pl. 1010. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.

subsp. **pepo**

Distribution: Karnataka (cultivated throughout); cultivated throughout tropics & subtropics, native to C & S America.

Local/Common/Vernacular names: *Butternut Squash*, *Canada crookneck Canada pumpkin*, *China Crook-neck Squash*, *Crook-neck Squash*, *musky gourd*, *musky pumpkin*, *musky squash*, *West Indian pumpkin*, *winter crookneck squash*, *winter squash* (English); *Kumbala* (Kannada)

subsp. **ovifera** (L.) D.S.Decker, Econ. Bot. 42: 11. 1988

Cucurbita ovifera L., Mant. Pl. 1: 126. 1767

Cucurbita pepo var. *ovifera* (L.) Alef, Landw. Fl. 224. 1866

Distribution: Karnataka (cultivated throughout), cultivated throughout India; cultivated throughout tropical and subtropical countries.

Diplocyclos (Endl.) Post & Kuntze

Diplocyclos palmatus (L.) C.Jeffrey, Kew Bull. 15: 352. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 112. 1984.

Bryonia palmata L., Sp. Pl. 1012. 1753 excl. syn.

Coccinia palmata M.Roem., Syn. Peponif. 93. 1846

Bryonopsis laciniosa var. *walkeri* Chakrav., Rec. Bot. Surv. India 17(1):138. 1959

Diplocyclos palmatus var. *walkeri* (Chakrav.) Babu, Herb. Fl. Dehra Dun 198. 1977

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalaburagi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; tropical Africa, tropical Asia and Australia.

Local/Common/Vernacular names: *Anga tonde balli*, *Haagala beelu*, *Malingana balli*, *Shivalinga balli* (Kannada)

Gymnopetalum Arn.

Gymnopetalum chinense (Lour.) Merr., Philipp. J. Sci. 15: 256.1919

Bryonia cochinchinensis Lour., Fl. Cochinch., 595. 1790

Gymnopetalum cochinchinense (Lour.) Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 40(2):57.1871; K.G. Bhat, J. Bombay Nat. Hist. Soc. 90:138.1993.

Distribution: Karnataka (Dakshina Kannada), Andhra Pradesh, Bihar, Manipur, Tripura, Uttaranchal

Gymnopetalum tubiflorum (Wight & Arn.) Cogn. in A. & C. DC., Monogr. Phan. 3: 388.1881

Bryonia tubiflora Wight & Arn., Prodr. Fl. Ind. Orient. 1:347.1834

Gymnopetalum wightii Arn., Madras J. Lit. Sci. 8: 52. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Gynostemma Blume

Gynostemma pentaphyllum (Thunb.) Makino, Bot. Mag. (Tokyo) 16: 179. 1902; Jayanthi et al., Indian J. For. 39 (1): 79. 2016.

Vitis pentaphylla Thunb., Syst. Veg. ed. 14 244. 1784.

Distribution: Karnataka (Chamarajanagar), Andhra Pradesh, Assam, Himachal Pradesh, Manipur, Meghalaya, Mizoram, Sikkim, Tamil Nadu, Tripura, Uttarkhand, Uttar Pradesh, West Bengal; Bangladesh, Bhutan, China, Japan, Korea, Laos, Malaysia, Myanmar, Nepal, New Guinea, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam.

Kedrostis Medik.

Kedrostis courtallensis (Arn.) C. Jeffrey, Kew Bull. 15: 353. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Bryonopsis courtallensis Arn., J. Bot. 3: 274. 1841

Bryonopsis bennettii Miq., Fl. Ned. Ind. 1: 657. 1855

Cerasiocarpum bennettii (Miq.) Cogn. in A. & C. DC., Monogr. Phan. 3: 729. 1881

Kedrostis bennettii (Miq.) W.J.de Wilde & Duyfjes, Reinwardtia 12(2): 130. 2004

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Odisha, Tamil Nadu and West Bengal; Bangladesh, Myanmar and Sri Lanka.

Kedrostis foetidissima (Jacq.) Cogn. in A. & C. DC., Monogr. Phan. 3: 634. 1881; C.J.Saldanha, Fl. Karnataka 1: 298. 1984.

Trichosanthes foetidissima Jacq., Collectanea 2: 841. 1788

Bryonia rostrata Rottler, Neue Schriften der Ges. Naturf. Freunde Berlin 4: 212. 1803

Rhynchosarpa foetida auct. non Schrad., 1838; C.B.Clarke in Hook.f., Fl. Brit. India 2: 627. 1879.

Aechmandra rostrata (Rottler) Arn., J. Bot. 3: 274. 1841

Rynchosarpa rostrata (Rottler) Naudin, Ann. Sci. Nat., Bot. sér. 4, 16: 177. 1862

Kedrostis rostrata (Rottler) Cogn. in A. & C. DC., Monogr. Phan. 3: 636. 1881; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Distribution: Karnataka (cultivated in Belagavi, Chikmagaluru, Chitradurga, Uttara Kannada), Andhra Pradesh, Gujarat, Madhya Pradesh, Maharashtra, Tamil Nadu; native to W Africa; cultivated elsewhere.

Lagenaria Ser.

Lagenaria siceraria (Molina) Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 3: 435 1930; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Cucurbita lagenaria L., Sp. Pl. 1010. 1753

Cucurbita siceraria Molina, Sag. Stor. Nat. Chili 133. 1782

Lagenaria vulgaris Ser., Mém. Soc. Phys. Genève 3(1): 25. 1825; C.B.Clarke in Hook.f., Fl. Brit. India 2: 613. 1879.

Lagenaria hispida Ser., Prodr. 3: 299. 1828

Lagenaria idolatrica (Willd.) Ser. ex Cogn., Prodr. 3: 299. 1828

Lagenaria vittata Ser., Prodr. 3: 299. 1828

Lagenaria leucantha (Duchesne) Rusby, Mem. Torrey Bot. Club 6: 43. 1896

Distribution: Karnataka (cultivated throughout), cultivated throughout; native to Africa, cultivated elsewhere.

Local/Common/Vernacular names: *Bottle gourd* (English); *Hala gumbala, Kahi sore balli, Sore kaayi* (Kannada)

Luffa Mill.

Luffa acutangula (L.) Roxb., Hort. Bengal. 70. 1841; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Cucumis acutangulus L., Sp. Pl. 1011. 1753

Cucurbita umbellata Klein ex Willd., Sp. Pl., ed. 4(1): 608. 1805

Cucurbita acutangula (L.) Blume, Bijdr. 932. 1826

Luffa amara Roxb., Fl. Ind. ed. 1832 3: 715. 1832

Cucumis operculatus Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 343. 1834

Luffa kleinii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 344. 1834

Luffa umbellata (Klein ex Willd.) M.Roem., Fam. Nat. Syn. Monogr. 2: 63. 1846

Luffa acutangula var. *amara* (Roxb.) C.B.Clarke in Hook.f., Fl. Brit. India 2: 615. 1879

Luffa hermaphrodita Singh & Bhandari, Baileya 11(4): 136, Fig. 13. 1964

Distribution: Karnataka (throughout), native to & cultivated throughout India; cultivated throughout tropics.

Local/Common/Vernacular name: *Ridge or Ribbed gourd* (English); *Heere kayi* (Kannada)

Luffa cylindrica (L.) M.Roem., Syn. Monogr. 2: 63. 1846; C.J.Saldanha, Fl. Karnataka 1: 308. 1984

Momordica cylindrica L., Sp. Pl. 1009.1753

Luffa aegyptica Mill., Gard. Dict. ed. 8: sp.1768; C.B.Clarke in Hook.f., Fl. Brit. India 2: 614.1879.

Bryonia cheiophylla Wall., Numer. List 6715A. 1832

Luffa pentandra Roxb., Fl. Ind. ed.2, 3: 712.1832

Cucumis pentandrus Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 343. 1834

Distribution: Karnataka (cultivated throughout), cultivated and wild throughout; cultivated in tropical countries.

Local/Common/Vernacular name: *Bath sponge gourd* (English); *Boluheere kaayi Tuppada hirekayi* (Kannada)

Melothria L.

Melothria angulata Chakrav., J. Bombay Nat. Hist. Soc. 50: 899.1952; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Zehneria angulata (Chakrav.) J.L.Ellis, Bull. Bot. Surv. India 9: 8.1968

Solena angulata (Chakrav.) Babu, Herb. Fl. Dehra Dun 203.1977

Distribution: Karnataka (Kodagu, Mysuru, Shimoga), S India; endemic.

Momordica L.

Momordica charantia L., Sp. Pl. 1009. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Momordica charantia var. *B* L., Sp. Pl.1009.1753

Momordica indica L., Herb. Amb. 24.1754

Momordica muricata Willd., Sp. Pl. 4(1): 602. 1805

Momordica charantia var. *muricata* (Willd.) Chakrav., Fasc. Fl. India 11: 92. 1982

Distribution: Karnataka (cultivated throughout), native to cultivated throughout India; cultivated throughout tropics.

Local/Common/Vernacular name: *Bitter gourd* (English); *Hagala kayi* (Kannada)

Momordica cochinchinensis (Lour.) Spreng., Syst. Veg. 3: 14. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Muricia cochinchinensis Lour., Fl. Cochinch. 2: 596. 1790

Momordica mixta Roxb., Fl. Ind. 2: 709.1832

Momordica macrophylla Gage, Rec. Bot. Surv. India 3(1):61. 1904; Chakrav. Rec. Bot. Surv. India 17(1):94.1959.

Distribution: Karnataka (cultivated throughout), throughout India; Australia, Bangladesh, New Guinea, SE Asia.

Momordica cymbalaria Fenzl ex Naudin, Ann. Sc. Nat. ser 4, 12: 134. 1859; C.J.Saldanha, Fl. Karnataka 1: 299. 1984.

Luffa tuberosa Roxb., Fl. Ind. ed. 2. 3: 717. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Cucumis tuberosus Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 344. 1834

Momordica tuberosa (Roxb.) Cogn. in A. & C. DC., Monogr. Phan. 3: 454. 1881 non Dennst. 1818

Distribution: Karnataka (Ballari, Chitradurga, Davanagere, Dharwad, Kalaburagi, Raichur, Shivamogga, Vijayapura, Yadgir), Andhra Pradesh, Madhya Pradesh, Maharashtra, Tamil Nadu; tropical Africa.

Local/Common/Vernacular names: *Kadavanchi, Karchi kayi* (Kannada)

Momordica denudata (Thwaites) C.B. Clarke in hook.f., Fl. Brit. India 2: 618. 1879

Momordica dioica var. *denudata* Thwaites, Enum. Pl. Zeyl. 2: 126. 1859

Distribution: Karnataka (), Gujarat, Kerala, Maharashtra; Sri Lanka.

Momordica dioica Roxb. ex Willd., Sp. Pl. 4: 605. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 113. 1984.

Momordica roxburghii G.Don, Syst. Gard. 3: 35. 1834

Momordica triloba Wight, Herb. ex Steud. Nomencl. ed. 2 11: 155. 1841

Momordica wallichii Roem., Syn. Pepon. 2: 58. 1846

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada, Vijayapura), throughout India; Bangladesh, China, Myanmar, Nepal, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Gidda hagala*, *Kadu haagala kaayi*, *Karichi balli*, *Midi hagala* (Kannada)

Momordica subangulata Blume, Bijdr. 928. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 114. 1984.

Momordica renigera Wall. ex G. Don, Gen. Hist. 3: 36. 1834

Momordica subangulata subsp. *renigera* (Wall. ex G. Don) W.J.de Wilde, Bot. Zhurn. (Moscow & Leningrad) 87(3): 147. 2002

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi), Kerala, Maharashtra, Meghalaya, Sikkim; Bangladesh, China, Indonesia, Laos, Malaysia, Myanmar, Thailand, Vietnam.

Sicyos L.

Sicyos edulis Jacq., Enum. Syst. Pl. 32. 1760

Sechium edule (Jacq.) Sw., Fl. Ind. Occid. 2: 1150. 1800; B.D.Sharma et al., Fl. Karnataka Anal. 114. 1984.

Sechium americanum Poir. in Lam., Encycl. 7: 50. 1806

Sechium chayota Jacq. ex Hemsl., Biol. Cent.-Amer., Bot. 1(6): 491. 1880

Distribution: Karnataka (cultivated throughout) widely cultivated in Andhra Pradesh, Himachal Pradesh, Karnataka, Manipur, Maharashtra, Meghalaya, Mizoram, Nagaland, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh and West Bengal; runs wild in NE India), native of tropical America.

Local/Common/Vernacular name: *Chow-chow* (English); *Seeme badane kayi* (Kannada)

Solena Willd.

Solena amplexicaulis (Lam.) Gandhi in Saldanha & Nicolson, Fl. Hassan Dist. 179. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 114. 1984.

Bryonia amplexicaulis Lam., Encycl. 1: 496. 1785

Solena heterophylla Lour., Fl. Cochinch. 514. 1790

Bryonia umbellata Klein ex Willd., Sp. Pl. 4: 618. 1805

Zehneria umbellata (Klein ex Willd.) Thwaites, Enum. Pl. Zeyl. 125. 1859

Melothria amplexicaulis (Lam.) Cogn. in A. & C. DC., Monogr. Phan. 3: 621. 1881

Melothria heterophylla (Lour.) Cogn. in A. & C. DC., Monogr. Phan. 3: 618. 1881

Melothria ovata Cogn. in Engl. Pflanzenr. IV. 275, 1 (Heft 66): 114. 1916

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Manipur, Tamil Nadu; Afghanistan, Bangladesh, China, Indonesia, Malaysia, Nepal, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular name: *Gouti gida* (Kannada); *Karolu* (Tulu)

Trichosanthes L.

Trichosanthes anaimalaiensis Bedd., Madras J. Lit. Sci. ser. 3, 1: 47. 1864; C.J.Saldanha, Fl. Karnataka 1: 304. 1984

Trichosanthes anamalayana Bedd. in Trans. Linn. Soc. 25: 217. 1865

Trichosanthes bracteata var. *tomentosa* (Heyne ex C.B. Clarke) Chakrav., Rec. Bot. Surv. India 17(1): 47. 1959, nom. illeg.

Distribution: Karnataka (Bengaluru, Hassan), Andaman & Nicobar Islands, Andhra Pradesh, Arunachal Pradesh, Kerala, Maharashtra, Tamil Nadu, Tripura; endemic.

Trichosanthes bracteata (Lam.) Voigt, Hort. Suburb. Calcutt. 58. 1845

Trichosanthes tricuspidata auct. non Lour. 1790; C.J. Saldanha & M.S.E. Rao in C.J. Saldanha, Fl. Karnataka 1: 305. 1984.

Trichosanthes tricuspidata Lour., Fl. Cochinch. 589. 1790; C.J. Saldanha, Fl. Karnataka 1: 305. 1984

Modecca bracteata Lam., Encycl. 4: 210. 1797

Trichosanthes palmata Roxb., Fl. Ind. 3: 704. 1832

Trichosanthes palmata var. *tomentosa* Heyne ex C.B. Clarke, Fl. Brit. India 2(6): 607. 1879

Trichosanthes burmensis var. *alba* Kundu, J. Bombay Nat. Hist. Soc. 43: 381. 1943

Distribution: Karnataka (Bengaluru), Andaman & Nicobar Islands, Arunachal Pradesh, Kerala, Maharashtra, Tamil Nadu; Tripura; Indonesia, Myanmar.

Local/Common/Vernacular name: *Kagemari balli* (Kannada)

Trichosanthes cucumerina L., Sp. Pl. 1008. 1753; B.D. Sharma et al., Fl. Karnataka Anal. 114. 1984.

Trichosanthes anguina L., Sp. Pl. 1008. 1753; C.B. Clarke in Hook.f., Fl. Brit. India 2: 610. 1879.

Cucumis anguinus (L.) L., Sp. Pl. ed. 2 1437 1763.

Trichosanthes cucumerina var. *anguina* (L.) Haines, Bot. Bihar Orissa 388. 1922

Trichosanthes brevibracteata Kundu, J. Bot. 77: 10. 1939

Trichosanthes brevibracteata var. *longirostrata* Kundu, J. Bot. 77: 11. 1939

Trichosanthes brevibracteata var. *sublobata* Kundu, J. Bot. 77: 11. 1939

Trichosanthes pachyrrhachis Kundu, J. Bot. 77: 9. 1939

Distribution: Karnataka (cultivated throughout), cultivated throughout India, E Africa and Malaysia; widely cultivated for fruits as vegetable.

Local/Common/Vernacular names: *Snake gourd* (English); *Kahi padavala, Padavala kaayi* (Kannada)

Trichosanthes lepiniana (Naudin) Cogn. in A. & C. DC., Monogr. Phan. 3: 377. 1881; C.J. Saldanha, Fl. Karnataka 1: 304. 1984.

Involucaria lepiniana Naudin in Huber, Cat. 11. 1868

Distribution: Karnataka (Hassan, Shivamogga), Nagaland, Sikkim, Tamil Nadu, West Bengal; Indomalesia

Trichosanthes lobata Roxb., Fl. Ind. 3: 703. 1832; C.J. Saldanha, Fl. Karnataka 1: 305. 1984.

Trichosanthes perrottetiana Cogn. in A. & C. DC., Monogr. Phan. 3: 362. 1881

Anguina perrottetiana (Cogn.) Kuntze, Revis. Gen. Pl. 1: 254. 1891

Trichosanthes villosula var. *nilgirrensis* Kundu, J. Bombay Nat. Hist. Soc. 43(3): 375. 1942

Distribution: Karnataka (Hassan, Mysuru), Andhra Pradesh, Kerala, Puducherry, Tamil Nadu, Uttar Pradesh, West Bengal; China.

Trichosanthes nervifolia L., Sp. Pl. 1008. 1753; B.D. Sharma et al., Fl. Karnataka Anal. 114. 1984.

Trichosanthes cuspidata Lam., Encycl. 1: 188. 1783

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Goa, Kerala, Madhya Pradesh, Manipur, Meghalaya, Mizoram, Nagaland, Rajasthan, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; Sri Lanka.

Local/Common/Vernacular name: *Kaadubimbe* (Kannada)

Trichosanthes villosula Cogn. in A. & C. DC., Monogr. Phan. 3: 362. 1881

Anguina villosula (Cogn.) Kuntze, Revis. Gen. Pl. 1: 254. 1891

Distribution: Karnataka (Mysuru), Andhra Pradesh, Assam, Kerala, Nagaland, Tamil Nadu, West Bengal; Bangladesh, China.

Trichosanthes wallichiana (Ser.) Wight, Ann. Mag. Nat. Hist. 8: 270. 1842; B.D.Sharma et al., Fl. Karnataka Anal. 115. 1984.

Involucraria wallichiana Ser., Mem. Soc. Phy. Hist. Nat. Geneve 3(1): 25, t. 5. 1825

Trichosanthes grandibracteata Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 98. 1877

Trichosanthes multiloba C.B.Clarke in Hook. f., Fl. Brit. India 2: 608. 1879

Trichosanthes palmata var. *scotanthus* C.B.Clarke in Hook. f., Fl. Brit. India 2: 607. 1879

Trichosanthes bracteata var. *scotanthus* (C.B.Clarke) Handel-Mazzetti, Symb. Sin. 7(4): 1065. 1936

Distribution: Karnataka (Uttara Kannada), Arunachal Pradesh, Assam, Bihar, Himachal Pradesh, Maharashtra, Manipur, Meghalaya, Nagaland, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand, West Bengal; China, Nepal.

Zanonia L.

Zanonia indica L., Sp. Pl. ed. 2, 1457. 1763; B.D.Sharma et al., Fl. Karnataka Anal. 115. 1984.

Zanonia indica var. *pubescens* Cogn. in A. & C. DC., Monogr. Phan. 3: 927. 1881

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Assam, Goa, Kerala, Maharashtra, Meghalaya, Sikkim, Tamil Nadu, West Bengal; China, Indo-China, New Guinea, SE Asia, Sri Lanka.

Zehneria Endl.

Zehneria maysorensis (Wight & Arn.) Arn., J. Bot. (Hooker) 3: 275. 1841 ('maysorensis'); B.D.Sharma et al., Fl. Karnataka Anal. 115. 1984.

Bryonia mysorensis Wight & Arn., Prodr. Fl. Ind. Orient. 1: 345. 1834

Bryonia maysorensis Arn. in Hooker's J. Bot. 3 : 275. 1841

Melothria perpusilla var. *subtruncata* Cogn. in A. & C. DC., Phan 3: 606. 1881

Melothria mucronata var. *umbellata* Chakrav., Rec. Bot. Surv. India 17(1): 150. 1959

Melothria maysorensis (Wight & Arn.) C.C. Chang, Acta Phytotax. Sin. 8(4): 344. 1963

Zehneria maysorensis var. *umbellata* (Chakrav.) Kumari in A.N.Henry & N.C.Nair, Fl. Tamil Nadu Ind., Ser. 1: 175. 1983

Zehneria maysorensis var. *oblonga* V.P.Prasad & M.Prasad, J. Econ. Taxon. Bot. 17(2): 471. 1993

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Udupi), Andhra Pradesh, Assam, Kerala, Maharashtra, Meghalaya, Tamil Nadu; China, Malaysia.

Zehneria mucronata (Blume) Miq., Fl. Ned. Ind. 1(1): 656. 1855; Ramaswamy & Razi, Fl. Bangalore Dist. 597. 1973.

Bryonia mucronata Blume, Bijdr. 923. 1826

Melothria mucronata (Blume) Cogn. in A. & C. DC., Monogr. Phan. 3: 608. 1881

Pilogyne mucronata (Blume) W.J.de Wilde & Duyfjes, Reinwardtia 12: 410. 2009

Distribution: Karnataka (Bengaluru), cultivated elsewhere in India; SE Asia.

Distribution: Karnataka (Cultivated in Bengaluru),

Zehneria perpusilla (Blume) Bole & M.R. Almeida, J. Bombay Nat. Hist. Soc. 79(2): 315. 1983

Cucurbita perpusilla Blume, Cat. Hort. Buitenz. 105. 1823

Melothria perpusilla (Blume) Cogn. in A. & C. DC., Monogr. Phan. 3:607. 1881; Cooke, Fl. Pres. Bombay 1: 540. 1903.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Kodagu, Shivamogga, Uttara kannada), Peninsular to NE India; China, Japan, Malaysia, Myanmar.

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Kodagu, Shimoga, Uttara Kannada), Peninsula extending to NE India; China, Japan, Malaysia, Myanmar.

Zehneria scabra Sond. in Harv.& Sond., Fl. Cap. 2: 486. 1862; B.D.Sharma et al., Fl. Karnataka Anal. 115. 1984.

Bryonia scabra L.f., Suppl. Pl. 424. 1782.

Bryonia scabra Thunb., Prodr. Pl. Cap. 13. 1794.

Pilogyne scabra (L.f.) W.J. de Wilde & Duyfjes, Blumea 55(3): 294.2010

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Manipur, Tamil Nadu, Uttar Pradesh; Paleotropical

Cyperaceae

Actinoscirpus (Ohwi) R.W.Haines & Lye

Actinoscirpus grossus (L.f.) Goetgh. & D.A.Simpson, Kew Bull. 46: 171 1991.

Schoenoplectus grossus (L.f.) Palla, Allg. Bot. Z. Syst. 17 (Beih.): 3. 1911; Rao et al., J. Econ. Taxon. Bot. 35: 155. 2011.

Distribution: Karnataka (Uttara Kannada), Assam, Kerala; Trop. & Subtrop. Asia to N. Australia.

Bolboschoenus Asch. ex Palla

Bolboschoenus maritimus (L.) Palla, Syn. Deut. Schweiz. Fl. ed. 3: 2532.1905; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Scirpus maritimus L., Sp. Pl. 51.1753

Reigera maritima (L.) Opiz, Seznam 83.1852

Schoenoplectus maritimus (L.) Lye, Blyttia 29: 145.1971

subsp. **maritimus**

Distribution: Karnataka (Belagavi, Bengaluru, Dharwad, Hassan, Mysuru, Raichur), throughout India except NE region.

subsp. **affinis** (Roth) T.Koyama, Brittonia 31: 284.1979

Scirpus affinis Roth, Syst. Veg. 2: 140. 1817

Scirpus strobilinus Roxb., Fl. Ind. 1, 1: 219. 1820

Scirpus maritimus var. *affinis* (Roth) C.B. Clarke, Fl. Brit. India 6(19): 659. 1893

Bolboschoenus affinis (Roth) Drobow, Trudy Bot. Muz. Imp. Akad. Nauk 16: 139.1916

Bolboschoenus strobilinus (Roxb.) V.I.Krecz., Fl. Tadzhikskoi SSR 2: 47.1963

Scirpus maritimus subsp. *affinis* (Roth) Norl., Bot. Not. 125: 404.1972

Bolboschoenus maritimus var. *affinis* (Roth) Parmar, Fl. Rajasthan 3: 888.1993

Distribution: Karnataka (Belagavi, Vijayapura), throughout except E Peninsulaar India; Asia

Bulbostylis Kunth

Bulbostylis barbata (Rottb.) C.B.Clarke in Hook.f.,Fl. Brit. India 6: 651 1893; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Scirpus barbatus Rottb., Descr. Icon. Rar. Pl. 52.1773

Cyperus barbata (Rottb.) Poir., Encycl. Suppl. 5: 186.1817

Scirpus monander Roxb., Fl. Ind. 1: 222 1820

Isolepis wallichiana Schult., Mant. 2: 533.1824

Fimbristylis barbata (Rottb.) Benth., Fl. Austral. 7: 321.1878

Iria barbata (Rottb.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Stenophyllum barbatus (Rottb.) T.Cooke, Fl. Bombay 2: 887.1908

Abildgaardia wallichiana (Schult.) Lye, Nordic J. Bot. 3: 239.1983

subsp. *barbata*

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Chamarajanagar, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Vijayapura), throughout India: warmer regions of Old world & Southern USA.

Local/Common/Vernacular name: *Chavuri hullu* (Kannada)

subsp. *pulchella* (Thwaites) T.Koyama, Bot. Mag. (Tokyo) 93: 341.1980

Isolepis pulchella Thwaites, Enum. Pl. Zeyl. 350.1864

Fimbristylis pulchella (Thwaites) Trimen, Syst. Cat. Fl. Pl. Ceylon 102.1885

Bulbostylis barbata var. *pulchella* (Thwaites) C.B.Clarke in Hook.f., Fl. Brit. India 6: 652.1893

Distribution: Karnataka (Belagavi, Dakshina Kannada, Uttara Kannada), Peninsular India; Sri Lanka.

***Bulbostylis densa* (Wall.) Hand.-Mazz., Vegetationsbilder 20(7): 16.1930; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.**

Scirpus densus Wall. in Roxb., Fl. Ind. ed.1, 1: 231.1820

Isolepis densa (Wall.) Schult., Mant. 2: 70.1824

Bulbostylis trifida Nees, Fl. Bras. 2(1): 84.1842

Bulbostylis capillaris f. *densus* (Wall.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 652.1894

Stenophyllum densum f. *trifidus* (Wall.) H.Pfeiff., Bot. Arch. 6: 188.1924

Fimbristylis densa (Wall.) T.Koyama & T.I.Chiang, Quart. J. Taiwan Mus. 13: 229.1960

Abildgaardia densa (Wall.) Lye., Bot. Not. 127: 496.1974

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India: throughout old world tropics.

***Bulbostylis puberula* C.B.Clarke in Hook.f., Fl. Brit. India 6: 652.1893; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.**

Scirpus puberulus Poir., Encycl. 6: 767.1805

Isolepis puberula (Poir.) Kunth, Enum. Pl. 2: 213.1837

Stenophyllum puberulus (Poir.) Killip, J. Siam Soc., Nat. Hist. Suppl. 7: 57.1927

Bulbostylis puberula var. *gracilis* C.E.C.Fisch., Fl. Madras 1662.1931

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Udupi, Uttara Kannada), Peninsular India; tropical Africa & Asia, Southern USA.

***Bulbostylis subspinescens* C.B.Clarke in Hook. F., Fl. Brit. India 6: 652.1893; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.**

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dakshina Kannada, Udupi), peninsular India; endemic.

Carex L.

***Carex baccans* Nees, Contr. Bot. India 122.1834; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.**

Carex baccans var. *siccifructus* C.B.Clarke in Hook.f., Fl. Brit. India 6: 723.1894

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Kodagu), Peninsular, Central & N E India; tropical Asia.

***Carex filicina* Nees, Contr. Bot. India 123.1834; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.**

Carex meiogyna Nees, Contr. Bot. India 123.1834

Carex filicina var. *meiogyna* (Nees) Strachey, Cat. Pl. Kumaon 73.1854

Carex nilagirica Hochst. ex Steud., Syn. Pl. Glumac. 2: 207.1855

Carex glauca Boeckeler, Linnaea 40: 353.1876; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Carex filicina var. *microgyna* C.B.Clarke in Hook.f., Fl. Brit. India 6: 718.1894

Carex leptocarpa C.B.Clarke in Hook.f., Fl. Brit. India 6: 719.1894

Carex saturata C.B.Clarke, J. Linn. Soc., Bot. 37: 12.1904

Carex filicina var. *leptocarpa* (C.B.Clarke) Kük., Pflanzenr. IV, 20(38): 274.1909

Carex filicina var. *saturata* (C.B.Clarke) Kük., Pflanzenr. IV, 20(38): 275.1909

Carex caricina (D.Don) Ghildyal & U.C.Battach., J. Econ. Taxon. Bot. 7: 602.1986

Carex caricina var. *glaucina* (Boeckeler) Ghildyal & U.C.Battach., J. Econ. Taxon. Bot. 7: 602.1986

Carex caricina var. *minor* Ghildyal & U.C.Battach., J. Econ. Taxon. Bot. 7: 602.1986

Carex caricina var. *microgyna* (C.B.Clarke) Karth., Fl. Ind. Enum. Monocot.: 35.1989

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluruu, Dakshina Kannada, Hassan, Kodagu, Uttara Kannada), Peninsular to N E India; E, S & SE Asia

Carex foliosa D.Don, Trans. Linn. Soc. London 14(2): 327.1824; Seetharam, Kotresh & Uplaonkar, Fl. Dist 57. 2000.

Distribution: Karnataka (Hassan, Mysuru, Uttara Kannada, Kalburgi? Yadgir?), N E India; Pakistan.

Carex indica L., Mant. Pl. 2: 574.1771; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Carex bengalensis Thwaites, Enum. Pl. Zeyl. 355.1864

Carex indica var. *laetebrunnea* C.B.Clarke in Hook.f., Fl. Brit. India 6: 715.1894

Carex indica var. *milnei* C.B.Clarke in Hook.f., Fl. Brit. India 6: 715.1894

Carex milnei Boott ex C.B.Clarke in Hook.f., Fl. Brit. India 6: 715.1894

Carex indica subsp. *laetebrunnea* (C.B.Clarke) T.Koyama in Dasanayake & Fosberg, Revised Handb. Fl. Ceylon 5: 374.1985

Distribution: Karnataka (Chikkamagaluru, Hassan, Uttara Kannada), S India; Australia, SE Asia,

Carex lindleyana Nees, Contr. Bot. India 121.1834; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Carex thyrsiflora Boott, Ill. Carex 1: 12.1858

Distribution: Karnataka (Chikkamagaluru, Shivamogga). S India; Sri Lanka.

Carex maculata Boott, Trans. Linn. Soc. London 20: 128.1846; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Distribution: Karnataka (Hassan), S & N E India; Asia.

Carex raphidocarpa Nees, Contr. Bot. India 122.1834

Carex mercarensis Hochst. ex Steud., Syn. Pl. Glumac. 2: 194.1855

Carex mercarensis var. *major* Steud., Syn. Pl. Glumac. 2: 194.1855

Carex raphidocarpa var. *major* (Steud.) Kük., Pflanzenr. IV, 20(38): 277.1909

Carex raphidocarpa var. *mercarenensis* (Hochst. ex Steud.) Kük., Pflanzenr. IV, 20(38): 277.1909

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), peninsular India; endemic.

Carex speciosa Kunth, Enum. Pl. 2: 504.1837; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Distribution: Karnataka (Mysuru, Uttara Kannada), throughout except W & NW India; S E Asia.

***Courtoisina* J. Sojak**

Courtoisina cyperoides (Roxb.) Soják, Cas. Nár. Mus., Odd. Prír. 148: 193.1980; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Kyllinga cyperoides Roxb., Fl. Ind. ed.1, 1: 182.1820

Mariscus cyperoides (Roxb.) A.Dietr., Sp. Pl. 2: 348.1832

Courtoisia cyperoides (Roxb.) Nees, Linnaea 9: 286.1834

Indocourtoisia cyperoides (Roxb.) Bennet & Raizada, Indian Forester 107: 432.1981

Pseudomariscus cyperoides (Roxb.) Rauschert, Taxon 31: 559.1982

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Dharwad, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Uttara Kannada), throughout except N W India; Tropical Africa, Madagascar.

Local/Common/Vernacular name: *Bili sanna Jambu hullu* (Kannada)

Cyperus L.

Cyperus alopecuroides Rottb., Descr. Icon. Rar. Pl. 38.1773; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Juncellus alopecuroides (Rottb.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 595.1893

Chlorocyperus alopecuroides (Rottb.) Grossh., Fl. Kavkaza 1: 150.1928

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Vijayapura, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Kalburgi, Hassan, Kodagu, Mandya, Mysuru, Uttara Kannada), throughout India; Africa, Australia, S & SE Asia, Madagascar, West Asia.

Local/Common/Vernacular name: *Bili jambu hullu* (Kannada)

Cyperus alternifolius L., Mant. Pl. 1: 28.1767; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Eucyperus alternifolius (L.) Rikli, Jahrb. Wiss. Bot. 27: 568.1895

subsp. *alternifolius*

Distribution: Karnataka (cultivated in gardens of Belagavi, Bengaluru, Dharwad, Mysuru), native to Arabia, Madagascar, tropical Africa.

subsp. *flabelliformis* Kük. in H.G.A.Engler, Pflanzenr., IV, 20(101): 193. 1936.

Cyperus involucratus Rottb., Descr. Pl. Rar. 22.1772; V.P.Prasad & N.P.Singh, Sedges Karnataka 101. 2002.

Cyperus flabelliformis Rottb., Descr. Icon. Rar. Pl. 42.1773; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Cyperus alternifolius var. *flabelliformis* (Rottb.) M.R.Almeida, Fl. Maharashtra 5B: 304.2009

Distribution: Karnataka (cultivated in gardens of Belagavi, Bengaluru, Dakshina Kannada, Dharwad, Mysuru Udupi), native to Arabia & tropical Africa.

Cyperus alulatus J.Kern, Reinwardtia 1: 463.1952; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Distribution: Karnataka (Belagavi, Vijayapura, Chikkamagaluru, Chitradurga, Mysuru, Shivamogga), throughout India; Pakistan

Cyperus amabilis Vahl, Enum. Pl. 2: 318.1805; V.P. Prasad & N. P. Singh, Sedges Karnataka 77.2002.

Distribution: Karnataka (Raichur), throughout except NW India; tropical Africa, Asia, America.

Cyperus arenarius Retz., Observ. Bot. 4: 9.1786; B.D.Sharma et al., Fl. Karnataka Anal. 304. 1984.

Scirpus glomeratus B.Heyne ex Wall., Numer. List 3460.1831

Distribution: Karnataka (Ballari, Chitradurga, Dakshina Kannada, Udupi, Uttara Kannada), throughout India; Iran, Pakistan, Sri Lanka, Vietnam.

Cyperus articulatus L., Sp. Pl. 44.1753; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Chlorocyperus articulatus (L.) Rikli, Jahrb. Wiss. Bot. 27: 563 1895

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Vijayapura, Chitradurga, Davanagere, Dharwad, Hassan, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada), Peninsular to Eastern India; Indo-China, Mediterranean, tropical Africa, Americas.

Local/Common/Vernacular name: *Yele tollu Jambu hullu* (Kannada)

Cyperus bulbosus Vahl, Enum. Pl. 2: 342.1805; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus polyphyllus Vahl, Enum. Pl. 2: 317.1805

Cyperus oleraceus Roxb. ex Nees, Contr. Bot. India 81.1834

Distribution: Karnataka (Bengaluru, Mysuru, Uttara Kannada), throughout, Asia, Australia, tropical Africa.

Cyperus castaneus Willd., Sp. Pl. 1: 278.1797; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus flavescens var. *castaneus* (Willd.) Pursh, Fl. Amer. Sept. 1: 52.1813

Cyperus castaneus var. *brevimucronatus* Kük., Pflanzenr. IV, 20(101): 264.1936

Distribution: Karnataka (Bengaluru, Chitradurga, Dakshina Kannada, Dharwad, Mandya, Mysuru, Raichur, Udupi), Peninsular India and Himalayas; Asia, tropical Australia.

Local/Common/Vernacular name: *Chandu cheni hullu* (Kannada)

Cyperus cephalotes Vahl, Enum. Pl. 2: 311.1805; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus monocephalus Roxb., Fl. Ind. 1: 193.1820

Anosporum monocephalum (Roxb.) Nees, Linnaea 9: 287.1834

Cyperus monocephalooides Roxb. ex Nees, Contr. Bot. India 92.1834

Anosporum cephalotes (Vahl) Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 39(2): 84. 1870

Cyperus natans Buch.-Ham. ex C.B.Clarke, J. Linn. Soc., Bot. 21: 34.1884

Ungeria monocephala (Roxb.) Nees ex C.B.Clarke, J. Linn. Soc., Bot. 21: 34.1884

Distribution: Karnataka (Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), throughout except NW India & Himalayas ;E & SE Asia, tropical Australia

Cyperus clarkei T.Cooke, Fl. Bombay 2: 873.1908

Mariscus bulbosus C.B.Clarke in Hook.f.,Fl. Brit. India 6: 620.1893; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Mariscus clarkei (T.Cooke) T.Koyama, J. Jap. Bot. 51: 316.1976

Distribution: Karnataka (Belagavi, Vijayapura, Dharwad, Mysuru, Uttara Kannada), Western Peninsular & N W India;Sri Lanka.

Cyperus compactus Retz., Observ. Bot. 5: 10.1788

Cyperus dilutus Vahl, Enum. Pl. 2: 357. 1805

Cyperus spinulosus Roxb., Fl. Ind. 1: 207. 1820

Cyperus dilutus var. *macrostachys* Boeckeler, Linnaea 36: 355.1870

Mariscus compactus (Retz.) Bold., Zakfl. Java 77.1916; V.P.Prasad & N. P. Singh, Sedges Karnataka 253. 2002..

Cyperus compactus var. *macrostachys* (Boeckeler) Kük., Pflanzenr. IV, 20(101): 424.1936

Mariscus compactus var. *macrostachys* (Boeckeler) F.C. How, Flora of Kwangchow 747.1956

Mariscus compactus var. *macrostachys* (Boeckeler) H.B.Naithani & S.Biswas, Flow. Pl. India, Nepal & Bhutan 471. 1990

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Dharwad, Mysuru, Tumakuru, Uttara Kannada), throughout India; China, S EAsia, Sri Lanka, Taiwan.

Cyperus compressus L., Sp. Pl. 46.1753; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus pectinatus Roxb., Fl. Ind. 1: 195.1820

Cyperus compressus var. *capillaceus* C.B.Clarke in Urb., Symb. Antill. 2: 32.1900

Chlorocyperus compressus (L.) Palla, Denkschr. Kaiserl. Akad. Wiss., Math.-Naturwiss. Kl. 84: 451.1909

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Vijayapura, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India, Cosmopolitan.

Local/Common/Vernacular names: *Usumani hullu*, *Vusumani hullu* (Kannada)

Cyperus corymbosus Rottb., Descr. Icon. Rar. Pl. 42.1773; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus koenigii Vahl, Enum. Pl. 2: 302.1805

Cyperus bengalensis Spreng., Neue Entdeck. Pflanzenk. 3: 101.1822

Cyperus nudus Roxb., Fl. Ind. ed. 2, 1: 209.1832

Cyperus tegetiformis Roxb. ex Arn., Contr. Bot. India 89.1834

Cyperus corymbosus var. *pangorei* C.B.Clarke, J. Linn. Soc., Bot. 21: 159.1884

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dharwad, Hassan. Kodagu, Mandya, Mysuru, Shivamogga, Uttara Kannada), throughout; Myanmar, Nepal, Sri Lanka, tropical W Africa & S America.

Local/Common/Vernacular names: *Bhandre hullu*, *Jake* (Kannada)

Cyperus cuspidatus Kunth, Nov. Gen. Sp. 1: 204.1817; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.
Cyperus angustifolius Nees, Contr. Bot. India 79.1834
Cyperus cuspidatus var. *burchellii* C.B.Clarke, J. Linn. Soc., Bot. 21: 90.1884
Dichostylis cuspidata (Kunth) Palla, Bot. Jahrb. Syst. 10: 296.1888
Cyperus uncinatus C.B.Clarke in Oliv., Fl. Trop. Afr. 8: 328. 1901
Cyperus gratus C.B.Clarke, Bot. Jahrb. Syst. 38: 133.1906
Cyperus uncinatus var. *gratus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 261.1936
 Distribution: Karnataka (throughout), throughout India; pantropical.

Cyperus cyperinus (Retz.) Suringar, Cyperus Mal. Arch. 154.1898
Kyllinga cyperina Retz., Observ. Bot. 6: 21. 1791
Kyllinga bulbosa J.Koenig ex Vahl, Enum. Pl. 2: 376. 1805.
Mariscus cyperinus (Retz.) Vahl, Enum. Pl. 2: 377.1805; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984.
Mariscus pictus Nees, Contr. Bot. India 90.1834; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984.
Cyperus umbellatus var. *laxatus* C.B.Clarke, J. Linn. Soc., Bot. 21: 201. 1884
Mariscus cyperinus var. *bengalensis* C.B.Clarke in Hook.f., Fl. Brit. India 6: 621.1893
Mariscus cyperinus var. *maximus* C.B.Clarke, J. Linn. Soc., Bot. 34: 43. 1898
Cyperus cyperoides subsp. *cyperinus* (Retz.) Kük., Bot. Jahrb. Syst. 59: 46.1925
Cyperus cyperinus var. *bengalensis* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 520.1936
Cyperus cyperinus var. *laxatus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 519.1936
Cyperus cyperinus var. *maximus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 520.1936
Mariscus cyperinus subsp. *laxatus* (C.B.Clarke) T.Koyama, Gard. Bull. Singapore 30: 157.1977
Mariscus cyperinus var. *pictus* (Nees) Karthik., Fl. Ind. Enumerat. - Monocot. 63 1989
 Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Vijayapura, Chikkamagalurru, Dakshina Kannada, Dharwad, Kalburgi, Hassan, Kodagu, Mandya, Raichur, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout except NE India; tropical Africa, Australia, E & SE Asia. Asia
 Local/Common/Vernacular name: *Anantha gonde hullu* (Kannada)

Cyperus cyperoides (L.) Kuntze, Revis. Gen. Pl. 3(2): 333.1898
Scirpus cyperoides L., Mant. Pl. 2: 181 1771
Kyllinga sumatrensis Retz., Observ. Bot. 4: 13.1786
Kyllinga umbellata var. *sumatrensis* (Retz.) Willd., Sp. Pl. 1: 258.1797
Cyperus biglumis C.B.Clarke, J. Linn. Soc., Bot. 21: 199.1884
Mariscus sieberianus Nees ex C.B.Clarke in Hook.f., Fl. Brit. India 6: 622.1893
Mariscus sieberianus var. *evolutior* C.B.Clarke in Hook.f., Fl. Brit. India 6: 622.1893
Mariscus sieberianus var. *khasianus* C.B.Clarke in Hook.f., Fl. Brit. India 6: 623.1893
Mariscus sublimis C.B.Clarke in T.A. Durand & H. Schinz, Consp. Fl. Afric. 5: 594.1894
Cyperus sieberianus (Nees ex C.B.Clarke) K.Schum., Notizbl. Königl. Bot. Gart. Berlin 2: 96.1898
Cyperus konkanensis T.Cooke, Fl. Bombay 2: 874.1908
Mariscus umbellatus var. *evolutior* (C.B.Clarke) E.G. Camus, Fl. Indo-Chine 7: 78 78. 1912
Cyperus cyperoides var. *khasianus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 518.1936
Cyperus cyperoides var. *subcompositus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 516.1936
Cyperus subumbellatus var. *sublimis* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 525.1936
Cyperus cylindrostachyus var. *subcompositus* (C.B.Clarke) Skottsb., Acta Horti Gothob. 15: 300.1944
Cyperus sublimis (C.B.Clarke) Dandy, Cat. Vasc. Pl. S. Tome 362.1944
Mariscus umbellatus var. *subcompositus* (C.B.Clarke) T.Tang & F.T.Wang, Fl. Reipubl. Popularis Sin. 11: 181.1961
Mariscus sumatrensis (Retz.) J.Raynal, Adansonia n.s., 15: 110.1975; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984 (*Mariscus sumatrensis* Retz.).
Mariscus sumatrensis var. *evolutior* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 64.1989
Mariscus sumatrensis var. *khasianus* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 64.1989
Mariscus sumatrensis var. *subcompositus* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 64.1989
Mariscus concinnus var. *evolutior* (C.B.Clarke) Panigrahi, Fl. Bilaspur Distr. 2: 699.1999
Mariscus concinnus var. *khasianus* (C.B.Clarke) Panigrahi, Fl. Bilaspur Distr. 2: 699.1999
Mariscus concinnus var. *subcompositus* (C.B.Clarke) Panigrahi, Fl. Bilaspur Distr. 2: 699.1999
 Distribution: Karnataka (Bengaluru, Bidar, Chamarajanagar, Chikkamagaluru, Dharwad, Hassan, Kodagu, Mandya, Shivamogga, Uttara Kannada), throughout India: subtropical regions of Africa, Asia, Australia

Cyperus difformis L., Cent. Pl. II 6.1756

Distribution: Karnataka (throughout), throughout India; tropical & subtropical regions of old world.

Local/Common/Vernacular name: *Kari sanna jambu hullu* (Kannada)

Cyperus diffusus Vahl, Enum. Pl. 2: 321.1805

Cyperus pubisquama Steud., Syn. Pl. Glumac. 2: 20.1854; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus diffusus var. *macrostachyus* Boeckeler, Linnaea 35: 534.1868

Eucyperus diffusus (Vahl) Rikli, Jahrb. Wiss. Bot. 27: 568.1895

Cyperus diffusus subsp. *macrostachyus* (Boeckeler) T.Koyama, Gard. Bull. Singapore 30: 139.1977

Cyperus laxus var. *macrostachyus* (Boeckeler) Karth., Fl. Ind. Enum. Monocot.: 46 1989

Cyperus laxus subsp. *macrostachyus* (Boeckeler) V.P.Prasad & N.P.Singh, Phytotaxonomy 1: 64.2001

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Udupi, Uttara Kannada) W Peninsular to N E India; China, Taiwan, S E Asia to New Guinea.

Cyperus digitatus Roxb., Fl. Ind. 1: 209.1820; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus glomeratus Roxb. ex Arn., Contr. Bot. India 85.1834

Cyperus digitatus var. *hookeri* (Boeckeler) C.B.Clarke in Hook.f., Fl. Brit. India 6: 618.1893; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus oatesii C.B.Clarke in Hook.f., Fl. Brit. India 6: 618.1893

Cyperus digitatus var. *oatesii* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 56.1935

subsp. **digitatus**

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Hassan), throughout India; Pantropical.

subsp. **auricomus** (Sieber ex Spreng.) Kük., Bot. Not. 1934: 65.1934

Cyperus auricomus Sieber ex Spreng., Syst. Veg. 1: 230.1824

Papyrus elatus Nees, Contr. Bot. India 88.1834

Cyperus auricomus var. *khassiana* C.B.Clarke, J. Linn. Soc., Bot. 21: 189.1884

Cyperus auricomus var. *minor* C.B.Clarke in Oliv., Fl. Trop. Afr. 8: 373.1901

Papyrus auricomus (Sieber ex Spreng.) Schrad., Index Seminum (GOET) 1835: 5.1935

Distribution: Karnataka (Hassan); Soutern Peninsular to N E India; Myanmar, Sri Lanka.

Cyperus distans L.f., Suppl. Pl. 103.1782; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus nutans Sieber ex C.Presl, Isis (Oken) 21: 271.1828.

Cyperus distans var. *pseudonutans* Kük., Pflanzenr. IV, 20(101): 140.1935; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Distribution: Karnataka (throughout), througout India;Pantropical.

Cyperus dives Delile, Descr. Egypte, Hist. Nat. 149.1813

Cyperus immensus C.B.Clarke, J. Linn. Soc., Bot. 20: 294.1883

Cyperus exaltatus var. *dives* (Delile) C.B.Clarke, J. Linn. Soc., Bot. 21: 187.1884; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Cyperus immensus var. *taylorii* C.B.Clarke in Oliv., Fl. Trop. Afr. 8: 372.1901

Cyperus petherickii C.B.Clarke in Oliv., Fl. Trop. Afr. 8: 371.1901

Cyperus immensus var. *petherickii* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 67.1935

Distribution: Karnataka (Ballari, Chikkamagaluru, Mysuru, Shivamogga, Uttara Kannada), Peninsular & E India; Africa, Pakistan.

Cyperus dubius Rottb., Descr. Icon. Rar. Pl. 20.1773

Mariscus dregeanus Kunth, Enum. Pl. 2: 120.1837

Cyperus triceps Endl., Cat. Horti Vindob. 1: 94.1842

Mariscus dubius (Rottb.) Kük. ex G.E.C.Fischer, Fl. Madras 1644.1931; B.D.Sharma et al., Fl. Karnataka Anal. 305. 1984.

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Uttara Kannada), S to E India; China, Sri Lanka, S E Asia, tropical Africa.

Local/Common/Vernacular name: *Ananata gonde hullu* (Kannada)

Cyperus esculentus L., Sp. Pl. 45.1753; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Pterocyperus esculentus (L.) Opiz, Seznam 80.1852

Cyperus esculentus var. *helodes* (Schrad. ex Nees) C.B.Clarke, J. Linn. Soc., Bot. 21: 181.1884

Cyperus esculentus var. *sprucei* C.B.Clarke, J. Linn. Soc., Bot. 21: 181.1884

Pycreus esculentus (L.) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(3): 147.1932

Distribution: Karnataka (Belagavi, Bengaluru, Uttara Kannada), throughout except E & N E India; Africa, Australia, Meditrranian region, S E Asia.

Local/Common/Vernacular name: *Nela badami hullu* (Kannada)

Cyperus exaltatus Retz., Observ. Bot. 5: 11.1788; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Cyperus odoratus Burm.f., Fl. Indica 20.1768

Cyperus umbellatus Roxb., Fl. Ind. 1: 205.1820

Cyperus tokiensis C.B.Clarke, Kew Bull., Addit. Ser. 8: 9.1908

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Uttara Kannada); throughout India;tropical Africa, S Australia, E & S E Asia.

Local/Common/Vernacular names: *Kempu jambu*, *Kempu jambu hullu* (Kannada)

Cyperus fuscus L., Sp. Pl. 46 1753; N.M.G.Babu, Fl. Ethnobot. Chitradurga Dist. 594. 2011.

Distribution: Karnataka (Chitrdurga)

Cyperus haspan L., Sp. Pl. 45.1753; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Cyperus juncoides Lam., Tabl. Encycl. 1: 147.1791

Cyperus hyemalis Pursh ex C.B.Clarke, J. Linn. Soc., Bot. 21: 120.1884

Cyperus haspan subsp. *juncoides* (Lam.) Kük., Pflanzenr. IV, 20(101): 249.1936; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Distribution: Karnataka (throughout), throughout India; throughout tropical & subtropical regions.

Local/Common/Vernacular name: *Chikka tunge hullu* (Kannada)

Cyperus iria L., Sp. Pl. 45.1753; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Cyperus paniciformis Franch. & Sav., Enum. Pl. Jap. 2: 537.1878

Cyperus iria var. *paniciformis* C.B.Clarke in Hook.f., Fl. Brit. India 6: 607.1893; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Chlorocyperus iria (L.) Rikli, Jahrb. Wiss. Bot. 27: 564.1895

Distribution: Karnataka (throughout), throughout India; Tropical E Africa, Asia, Australia; introduced elsewhere.

Local/Common/Vernacular names: *Dabbe jambu hullu*, *Dodda jambu hullu* (Kannada)

Cyperus javanicus Houtt., Nat. Hist. t. 88, f. 1.1782

Cyperus pennatus Lam., Tabl. Encycl. 1: 144.1791

Mariscus albescens Gaudich., Voy. Uranie 415.1829

Mariscus pennatus (Lam.) Domin, Biblioth. Bot. 85: 440.1915

Mariscus javanicus (Houtt.) Merr. & F.P.Metcalf, Lingnan Sci. J. 21: 4.1945; V.P. Prasad & N.P.Singh,Sedges Karnataka 258. 2002.

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), throughout coastal areas of India both main land and Islands;tropical coasts of Africa, Australia, E & SE Asia, Pacefic Islands

Cyperus laevigatus L., Mant. Pl. 2: 179.1771; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Juncellus laevigatus (L.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 596.1893

Juncellus laevigatus var. *junciformis* C.B.Clarke, Fl. Brit. India 6: 596.1893

Chlorocyperus laevigatus (L.) Palla, Allg. Bot. Z. Syst. 6: 221.1900

Acorellus laevigatus (L.) Palla, Syn. Deut. Schweiz. Fl. ed. 3, 2: 2558.1905

Distribution: Karnataka (Belagavi, Ballari, Chitradurga, Chitradurga, Hassan, Mysuru, Raichur, Tumakuru), throughout except N E India & Himalayas; Africa, Australia, Europe, S America, W Asia.

Cyperus laxus Lam., Tabl. Encycl. 1: 146.1791; V.P.Prasad & N.P.Singh Sedges Karnataka 104. 2002 (as subsp. *laxus*).

Cyperus homoliria Steud., Syn. Pl. Glumac. 2: 23. 1854

Distribution: Karnataka (Dakshina Kannada), throughout India; China, S E Asia, Taiwan.

Cyperus macer C.B.Clarke, J. Linn. Soc., Bot. 21: 160.1884; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Distribution: Karnataka (Uttara Kannada), Peninsular to N E India; Bangladesh, Myanmar.

Cyperus maderaspatanus Willd., Sp. Pl. 1: 278.1797

Mariscus maderaspatanus (Willd.) Napper, J. E. Africa Nat. Hist. Soc. Natl. Mus. 28: 10.1971; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984.

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Peninsular India; S E Asia, tropical E Africa.

Cyperus malaccensis Lam., Tabl. Encycl. 1: 146.1791; V.P.Prasad & N.P.Singh Sedges Karnataka 107. 2002.

Cyperus incurvatus Roxb., Fl. Ind. 1,1: 200 1820

Cyperus wallichii Nees, Contr. Bot. India 83.1834

Chlorocyperus malaccensis (Lam.) Palla, Allg. Bot. Z. Syst. 17(Beil.): 6.1911

Distribution: Karnataka (Dakshina Kannada, Kalburgi, Udupi, Uttara Kannada, Yadgir), W Peninsular to E India; Australia, Asia.

Cyperus meboldii Kük., Repert. Spec. Nov. Regni Veg. 18: 345 1922; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Distribution: Karnataka (Belagavi, Vijayapura, Shivamogga, Uttara Kannada), W Peninsular to N W India; E Africa.

Cyperus michelianus (L.) Delile, Descr. Egypte, Hist. Nat. 3: 50.1813

Scirpus michelianus L., Sp. Pl. 52.1753; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Isolepis micheliana (L.) Roem. & Schult., Syst. Veg. 2: 114.1817

Fimbristylis micheliana (L.) Rchb., Fl. Germ. Excurs. 75.1830

Dichostylis micheliana (L.) Nees, Contr. Bot. India 94.1834

Cyperus pygmaeus var. *micheliana* (L.) Boeckeler, Linnaea 35: 494.1868

Juncellus michelianus (L.) Blatt. & McCann, J. Bombay Nat. Hist. Soc. 37: 33.1934

subsp. **micheliana**

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Chitradurga, Dakshina Kannada, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; Africa, Australia, Meditrranea, S & SE Asia.

subsp. *pygmaeus* (Rottb.) Asch. & Graebn., Syn. Mitteleur. Fl. 2(2): 273. 1904

Cyperus pygmaeus Rottb., Descr. Icon. Rar. Pl. 20. 1773

Dichostylis pygmaea (Rottb.) Nees, Linnaea 9: 289. 1834

Pycreus pygmaeus (Rottb.) Nees, Linnaea 9: 283. 1834

Juncellus pygmaeus (Rottb.) C.B.Clarke in Fl. Brit. India 6: 596. 1893

Distribution: Karnataka (Bengaluru, Chitradurga, Hassan), throughout India; Paleotropical.

Cyperus mitis Steud., Syn. Pl. Glumac. 2: 316. 1855; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus tenuiflorus Roxb., Fl. Ind. 1: 199. 1820

Cyperus stenostachyus var. *indica* C.B.Clarke in Hook.f., Fl. Brit. India 6: 614. 1893

Distribution: Karnataka (Chitradurga, Hassan), Peninsular India; Australia, Madagascar, S E Asia.

Cyperus mollipes (C.B.Clarke) K.Schum., Pflanzenw. Ost-Afrikas C: 122. 1895

Mariscus mollipes C.B.Clarke in T.A. Durand & H. Schinz, Consp. Fl. Afric. 5: 590. 1894

Ascopholis gamblei C.E.C.Fisch., Bull. Misc. Inform. Kew 1931: 105. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Hassan), Kerala, Tamil Nadu; endemic.

Cyperus niveus Retz., Observ. Bot. 5: 12. 1788; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Distribution: Karnataka (Bengaluru), throughout India; Afghanistan, China, Myanmar, Thailand.

Cyperus nutans Vahl, Enum. Pl. Obs. 2: 363. 1805.

Cyperus myurosensis B.Heyne ex Wall., Numer. List 3346. 1831

Cyperus eleusinoides Kunth, Enum. Pl. 2: 39. 1837

Cyperus indicus Boeckeler, Beitr. Cyper. 1: 6. 1888

Cyperus nutans var. *eleusinoides* (Kunth) Haines, Bot. Bihar Orissa 5: 898. 1924; V.P.Prasad & N.P.Singh, Sedges Karnataka 112. 2002.

Cyperus nutans subsp. *eleusinoides* (Kunth) T.Koyama, Gard. Bull. Singapore 30: 136. 1977; B.D.Sharma et al., Fl. Karnataka Anal. 306. 1984.

Distribution: Karnataka (throughout), throughout India; Australia, E & SE Asia, tropical Africa.

Cyperus odoratus L., Sp. Pl. 46. 1753.

Diclidium odoratum (L.) Schrad. ex Nees, Fl. Bras. 2(1): 54. 1842

Torulinium odoratum (L.) S.S.Hooper, Kew Bull. 26: 579. 1972

Distribution: Karnataka (Mysuru)

Cyperus pangorei Rottb., Descr. Icon. Rar. Pl. 31. 1773; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus tegetum Roxb., Fl. Ind. 1: 211. 1820

Papyrus corymbosus Nees, Contr. Bot. India 89. 1834

Papyrus dehiscens Nees, Contr. Bot. India 89. 1834

Cyperus pangorei var. *ambiguus* C.B.Clarke, J. Linn. Soc., Bot. 21: 162. 1884

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; S Asia.

Local/Common/Vernacular name: *Murumule dodda jambu hullu* (Kannada)

Cyperus paniceus (Rottb.) Boeckeler, Linnaea 36: 381. 1870

Kyllinga panicea Rottb., Descr. Icon. Rar. Pl. 15. 1773

Mariscus paniceus (Rottb.) Vahl, Enum. Pl. 2: 373. 1805; V.P.Prasad & N.P.Singh, Sedges Karnataka 261. 2002.

Mariscus paniceus var. *roxburghianus* C.B.Clarke in Hook.f., Fl. Brit. India 6: 621. 1893

Mariscus roxburghianus (C.B.Clarke) Alston in Trimen, Handb. Fl. Ceylon 6(Suppl.): 309. 1931

Cyperus paniceus var. *roxburghianus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 526. 1936

Distribution: Karnataka (Bengaluru, Vijayapura, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout except N W India & Himalayas; S E Asia & Sri Lanka.

Cyperus pilosus Vahl, Enum. Pl. 2: 354.1805; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus fimbriatus Nees, Contr. Bot. India 86.1834

Cyperus marginellus Nees, Contr. Bot. India 83.1834

Cyperus obliquus Nees, Contr. Bot. India 86.1834

Cyperus heyneanus Boeckeler, Flora 42: 440.1859

Cyperus pilosus var. *obliquus* (Nees) C.B.Clarke, J. Linn. Soc., Bot. 21: 151.1884

Cyperus pilosus var. *polyanthus* C.B.Clarke, J. Linn. Soc., Bot. 21: 151.1884

Duval-jouvea pilosa (Vahl) Palla, Syn. Deut. Schweiz. Fl. ed. 3, 3: 2555.1905

Cyperus pilosus var. *fimbriatus* (Nees) Kük., Pflanzenr. IV, 20(101): 94.1935

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mandy, Mysuru, Shivamogga, Udupi, Uttara Kannada)

Cyperus platystylis R.Br., Prodr. Fl. Nov. Holl. 214.1810; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus pallidus Nees, Contr. Bot. India 79.1834

Anosporum pallidum Boeckeler, Linnaea 36: 412.1870

Cyperus fluitans Buch.-Ham. ex C.B.Clarke, J. Linn. Soc., Bot. 21: 118.1884

Distribution: Karnataka (Shivamogga, Uttara Kannada), throughout except N E India; Sri Lanka, Taiwan, S E Asia, Australia.

Cyperus procerus Rottb., Descr. Icon. Rar. Pl. 29.1773; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus carnosus B.Heyne ex Wall., Numer. List 3355 A.1831

Cyperus griffithianus Boeckeler, Linnaea 35: 601.1868

Cyperus heynei Boeckeler, Linnaea 35: 600.1868

Cyperus procerus var. *lasiorrhachis* C.B.Clarke in Hook.f., Fl. Brit. India 6: 610.1893

Duval-jouvea procera (Rottb.) H.Pfeiff., Mitt. Inst. Allg. Bot. Hamburg 7: 167.1928

Cyperus procerus var. *griffithianus* (Boeckeler) Kük., Pflanzenr. IV, 20(101): 91.1935

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Dharwad, Mysuru, Udupi, Uttara Kannada), Peninsular to E India; Australia. E & S E Asia.

Local/Common/Vernacular name: *Suvata bandhu* (Kannada)

Cyperus prolifer Lam., Tabl. Encycl. 1: 147.1791

Cyperus isocladus Kunth, Enum. Pl. 2: 37.1837

Cyperus prolifer var. *isocladus* (Kunth) Kük., Pflanzenr. IV, 20(101): 257.1936

Distribution: Karnataka (Belagavi)

Cyperus pulchellus R.Br., Prodr. Fl. Nov. Holl. 213 1810; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus sorostachyus Boeckeler, Linnaea 35: 588 1868

Cyperus zambesiensis C.B.Clarke, Trans. Linn. Soc. London, Bot. 4: 53 1894

Cyperus glaucophyllum var. *zambesiensis* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 203.1936

Sorostachys pulchellus (R.Br.) Lye, Nordic J. Bot. 1: 189.1981

Distribution: Karnataka (throughout), S, C & E India; tropical Africa, Asia, Australia.

Cyperus rotundus L., Sp. Pl. 45.1753; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Cyperus maritimus Bojer, Hortus Maurit. 378.1837

Chlorocyperus rotundus (L.) Palla, Allg. Bot. Z. Syst. 6: 61.1900

Pycnus rotundus (L.) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(3): 147.1932

Distribution: Karnataka (throughout), throughout India; pantropical.

Local/Common/Vernacular names: *Bhadra hullu*, *Bhadra mushti*, *Jake*, *Jekina gedde*, *Kanhari hullu*, *Kannari hullu*, *Tunge hullu* (Kannada); *Mustaka* (Sanskrit)

Cyperus rubicundus Vahl, Enum. Pl. 2: 308. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Cyperus teneriffae Poir., Encycl. 7: 245. 1806

Cyperus coromandelinus Spreng., Syst. Veg. 1: 217. 1824

Cyperus wightii Nees, Contr. Bot. India 78. 1834

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Davanagere, Dharwad, Hassan, Kodagu, Mandya, Mysuru, Tumakuru), Peninsular India; Africa, Arabia, Australia, Madagascar.

Cyperus squarrosus L., Cent. Pl. II 6. 1756

Cyperus aristatus Rottb., Descr. Icon. Rar. Pl. 22. 1773

Cyperus inflexus Muhl., Descr. Gram. 16. 1817

Cyperus versicolor Nees, Contr. Bot. India 78. 1834

Pycneurus squarrosus (L.) Nees, Linnaea 9: 283. 1834

Cyperus aristatus var. *versicolor* (Nees) Boeckeler, Linnaea 35: 500. 1868

Dichostylis aristata (Rottb.) Palla, Bot. Jahrb. Syst. 10: 299. 1888

Dichostylis squarrosa (L.) Palla, Bot. Jahrb. Syst. 10: 296. 1888

Mariscus squarrosus (L.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 623. 1893; V.P.Prasad & N.P.Singh, Sedges Karnataka 261. 2002.

Distribution: Karnataka (throughout), throughout India; throughout tropical & subtropical regions of Africa, America, Asia, Australia, sometimes extending to temperate areas.

Cyperus stoloniferus Retz., Observ. Bot. 4: 10. 1786; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Cyperus arenarius Hance ex C.B.Clarke, J. Linn. Soc., Bot. 21: 173. 1884.

Distribution: Karnataka (Belagavi, Ballari, Vijayapura, Chitradurga, Dakshina Kannada, Mandya, Mysuru, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Peninsular India; Australia, Madagascar, Mauritius, S E Asia, Sr Lanka.

Cyperus surinamensis Rottb., Descr. Icon. Rar. Pl. 35. 1773; K.G.Bhat, Indian J. For. 36(3): 365. 2013.

Distribution: Karnataka (Dakshina Kannada), native to tropical America

Cyperus tenuiculmis Boeckeler, Linnaea 36: 286. 1870; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Distribution: Karnataka (Davanagere, Hassan, Kodagu, Mysuru), throughout India in moist areas; S & S E Asia, tropical Africa & Australia.

Cyperus tenuispica Steud., Syn. Pl. Glumac. 2: 11. 1854; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Cyperus leptostachys Nees, Linnaea 9: 285. 1834

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; tropical Africa, Asia & Australia, Mauritius, Sri Lanka.

Local/Common/Vernacular name: *Honnehu hullu* (Kannada)

Cyperus tuberosus Rottb., Descr. Icon. Rar. Pl. 28. 1773

Chlorocyperus salaamensis Palla, Allg. Bot. Z. Syst. 9: 70. 1903

Cyperus rotundus subsp. *tuberosus* (Rottb.) Kük., Pflanzenr. IV, 20(101): 113. 1935

Distribution: Karnataka (Mysuru ?), Peninsular India; Africa, Australia, Mauritius, Sri Lanka.

Diplacrum R. Br.

Diplacrum africanum (Benth.) C.B.Clarke in T.A. Durand & H. Schinz, Consp. Fl. Afric. 5: 668. 1894

Scleria africana Benth., Gen. Pl. 3: 1071. 1883; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Scleria reticulata auct. non (Holttum) J.Kern, 1961; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Distribution: Karnataka (Belagavi, Shivamogga, Uttara Kannada), Peninsular India; W tropical Africa.

Diplacrum caricinum R.Br., Prodr. Fl. Nov. Holl. 241.1810

Scleria caricina (R.Br.) Benth., Fl. Austral. 7: 426.1878; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Mysuru, Udupi, Uttara Kannada), Peninsular to N E India including Andaman & Nicobar Islands; Australia, E & SE Asia, Micronesia.

Eleocharis R. Br.

Eleocharis acutangula (Roxb.) Schult., Mant. 2: 91.1824; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Scirpus acutangulus Roxb., Fl. Ind. 1: 216.1820

Scirpus medius Roxb., Fl. Ind. 1,1: 216.1820

Eleocharis fistulosa Schult., Mant. 2: 89.1824

Baeothryon fistulosum (Schult.) A.Dietr., Sp. Pl. 2: 94.1832

Limnochloa acutangula (Roxb.) Nees, Contr. Bot. India 114.1834

Limnochloa fistulosa (Schult.) Nees, Linnaea 9: 294.1834

Limnochloa media (Roxb.) Nees, Contr.Bot. Ind.114.1834

Scirpus fistulosus (Schult.) Kuntze, Revis. Gen. Pl. 2: 757.1891

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Dakshina Kannada, Davanagere, Hassan, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), W Peninsular to N E India; E Asia, Madagascar, S America, Tropical Africa, West Indies.

Eleocharis atropurpurea (Retz.) J.Presl & C.Presl, Reliq. Haenk. 1: 196 1828; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Scirpus atropurpureus Retz., Observ. Bot. 5: 14.1788

Isolepis atropurpurea (Retz.) Roem. & Schult., Syst. Veg. 2: 106.1817

Apostemon atropurpureum (Retz.) Raf., J. Phys. Chim. Hist. Nat. Arts 89: 105.1819

Heleocharis atropurpurea (Retz.) C. Presl, Reliq. Haenk. 1: 196.1828

Eleogiton atropurpurea (Retz.) A.Dietr., Sp. Pl. 2: 97.1832

Eleogenus atropurpureus (Retz.) Nees, Contr. Bot. India 113.1834

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Tumakuru, Udupi, Uttara Kannada), throughout India; Asia, America, Australia, S Europe.

Eleocharis congesta D.Don, Prodr. Fl. Nepal. 41.1825

Scirpus congestus (D.Don) Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 41.1825

Eleocharis purpurascens Boeckeler, Linnaea 36: 455.1870

Eleocharis subvivipara Boeckeler, Linnaea 36: 424.1870

Scirpus purpurascens (Boeckeler) Kuntze, Revis. Gen. Pl. 2: 758.1891

Scirpus subviviparus (Boeckeler) Kuntze, Revis. Gen. Pl. 2: 758.1891

Eleocharis congesta subsp. *subvivipara* (Boeckeler) T.Koyama, Micronesica 1: 78.1964

Distribution: Karnataka (Belagavi, Davanagere, Hassan, Kodagu, Mysuru, Uttara Kannada), throughout India; E, S & SE Asia, Madagascar.

Eleocharis dulcis Trin. ex Hensch., Vita Rumphii 186.1833; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Scirpus plantaginoides Rottb., Descr. Icon. Rar. Pl. 45.1773

Scirpus plantagineus Retz., Observ. Bot. 5: 14.1788

Hippuris indica Lour., Fl. Cochinch. 16.1790

Scirpus tuberosus Roxb., Pl. Coromandel 3: 25, t.231.1815

Eleocharis plantaginea (Retz.) Roem. & Schult., Syst. Veg. 2: 150.1817

Scirpus dubius Roxb., Fl. Ind. 1: 215.1820

Scirpus tumidus Roxb., Fl. Ind. 1: 212.1820

Eleocharis tumida (Roxb.) Schult., Mant. 2: 86.1824

Eleocharis equisetina J.Presl & C.Presl, Reliq. Haenk. 1: 195.1828

Limnochloa tumida (Roxb.) Nees, Contr. Bot. India114.1834

Eleocharis plantaginoides (Rottb.) Domin, Biblioth. Bot. 85: 445.1915

Eleocharis indica (Lour.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 621.1917

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India except N India & Himalayas; Australia, Madagascar, E, S & SE Asia, W Africa.

Eleocharis geniculata (L.) Roem. & Schult., Syst. Veg. 2: 150.1817; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Scirpus capitatus L., Sp. Pl. 48.1753

Scirpus geniculatus L., Sp. Pl. 48.1753

Eleocharis capitata (L.) R.Br., Prodr. Fl. Nov. Holl. 225.1810

Bulbostylis geniculatus (L.) Steven, Mém. Soc. Imp. Naturalistes Moscou 5: 355. 1814

Megadenus capitatus (L.) Raf., Neogenyton 4.1825

Eleogenus capitatus (L.) Nees, Contr. Bot. India 112.1834

Limnochloa geniculata (L.) Nees, Fl. Bras. 2(1): 99.1842

Chlorocharis geniculata (L.) Rikli, Jahrb. Wiss. Bot. 27: 564.1895

Trichophyllum capitatum (L.) House, Amer. Midl. Naturalist 6: 204.1920

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Chitradurga, Dakshina Kannada, Dharwad, Kalburgi, Hassan, Kodagu, Mandya, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Peninsula to NE India; Pantropical.

Eleocharis lankana T.Koyama, Bull. Natl. Sci. Mus. Tokyo 17(1): 67. 1974.

subsp. **mohamadii** Wad.Khan, J. Econ. Taxon. Bot. 22(3): 560. 1999

Distribution: Karnataka (Belagavi), Madhya Pradesh; endemic.

Eleocharis retroflexa (Poir.) Urb., Symb. Antill. 2: 165. 1900.

subsp. **chaetaria** (Roem. & Schult.) T.Koyama, Bull. Natl. Sci. Mus. Tokyo n.s., 17: 68.1974; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Eleocharis chaetaria Roem. & Schult., Syst. Veg. 2: 154.1817

Scirpus chaetarius (Roem. & Schult.) Spreng., Syst. Veg. 1: 203.1824

Trichophyllum chaetaria (Roem. & Schult.) House, Amer. Midl. Naturalist 6: 204.1920

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; Old World tropics.

Eleocharis spiralis (Rottb.) Roem. & Schult., Syst. Veg. 2: 155.1817; V.P.Prasad & N.P.Singh, Sedges Karnataka 148. 2002.

Scirpus spiralis Rottb., Descr. Icon.Rar. Pl. 42.1773

Heleocharis spiralis (Rottb.) R. Br., Prodr. 312.1810

Limnochloa spiralis (Rottb.) Nees, Contr. Bot. India 114.1834

Scirpus mutatus Roxb. ex C.B.Clarke in Hook.f., Fl. Brit. India.6: 627 1893

Distribution: Karnataka (Dakshina Kannada,Udupi, Uttara Kannada), W Peninsular India; Pantropical.

Eriophorum L.

Eriophorum comosum (Wall.) Nees, Contr. Bot. India 110.1834

Scirpus comosus Wall., Fl. Ind. 1: 234.1820

Scirpus elongatus Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 40.1825

Trichophorum comosum (Wall.) A.Dietr., Sp. Pl. 2: 202.1833

Eriophorum arundinaceum Nees, Contr. Bot. India 110.1834

Eriophorum cannabina Royle, Ill. Bot. Himal. Mts. 415.1840

Erioscirpus comosus (Wall.) Palla, Bot. Zeitung (Berlin) 54: 151.1896

Distribution: Karnataka (Mysuru),

Fimbristylis Vahl

Fimbristylis acuminata Vahl, Enum. Pl. 2: 285.1805; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Scirpus scaber Roxb., Fl. Ind. 1: 220.1820

Fimbristylis scabra (Roxb.) Schult., Mant. 2: 59.1824

Eleogiton scabra (Roxb.) A.Dietr., Sp. Pl. 2: 99.1832

Iria acuminata (Vahl) Kuntze, Revis. Gen. Pl. 2: 751.1891

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mandya, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; S & S E Asia, tropical Australia.

Fimbristylis aestivalis Vahl, Enum. Pl. 2: 288.1805; B.D.Sharma et al., Fl. Karnataka Anal. 308. 1984.

Scirpus aestivalis (Vahl) Retz., Observ. Bot. 4: 12.1786

Iria aestivalis (Vahl) Kuntze, Revis. Gen. Pl. 2: 751.1891

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mandya, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; S & S E Asia, tropical Australia.

Fimbristylis alboviridis C.B.Clarke in Hook.f., Fl. Brit. India 6: 638.1893; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Cyperus alboviridis C.B.Clarke ex Scott-Elliott, J. Linn. Soc., Bot. 29: 62. 1891

Distribution: Karnataka (Belagavi, Hassan), Andaman & Nicobar Islands, Peninsular & N E India; S & S E Asia.

Fimbristylis aphylla Steud., Syn. Pl. Glumac. 2: 114.1855; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Fimbristylis quadrangularis var. *crassa* C.B.Clarke in Hook.f., Fl. Brit. India 6: 644.1893

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Udupi, Uttara Kannada), throughout except N E India & Himalayas; S & SE Asia, tropical S Africa.

Fimbristylis argentea (Rottb.) Vahl, Enum. Pl. 2: 294.1805; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Scirpus argenteus Rottb., Descr. Icon. Rar. Pl. 51.1773

Scirpus monander Rottb., Descr. Icon. Rar. Pl. 50. 1773.

Fimbristylis monandra (Rottb.) F.Muell., Fragm. 1: 195.1859

Iria monandra (Rottb.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Distribution: Karnataka (Bengaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Tumakuru, Udupi, Uttara Kannada), Peninsular to E India; Mauritius, SE Asia, Sri Lanka.

Local/Common/Vernacular name: *Bettada usumani hullu* (Kannada)

Fimbristylis bispicula Govind., Rheedia 6: 59.1996; V.P.Prasad & N.P.Singh, Sedges Karnataka 161. 2002.

Distribution: Karnataka (Shivamogga), Andhra Pradesh, Maharashtra; endemic.

Fimbristylis bisumbellata (Forssk.) Bubani, Dodecanthea 30.1850; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Scirpus bisumbellatus Forssk., Fl. Aegypt.-Arab. 15.1775

Scirpus pallescens Roxb., Fl. Ind. 1: 229.1820

Iria bisumbellata (Forssk.) Kuntze, Revis. Gen. Pl. 2: 752.1891

Fimbristylis dichotoma subsp. *bisumbellata* (Forssk.) Luceño, Anales Jard. Bot. Madrid 57: 176.1999

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Uttara Kannada, Vijayapura), throughout India; Paleotropical extending to Mediterranean region.

Local/Common/Vernacular names: *Kadu s absige hullu*, *Niru sabsige hullu* (Kannada)

Fimbristylis cinnamometorum (Vahl) Kunth, Enum. Pl. 2: 229.1837; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Scirpus cinnamometorum Vahl, Enum. Pl. 2: 278.1805

Isolepis cinnamometorum (Vahl) Roem. & Schult., Syst. Veg. 2: 119.1817

Trichelostylis cinnamometorum (Vahl) Nees, Linnaea 9: 290.1834

Abildgaardia cinnamometorum (Vahl) Thwaites, Enum. Pl. Zeyl. 347.1864

Iria cinnamometorum (Vahl) Kuntze, Revis. Gen. Pl. 2: 753.1891

Distribution: Karnataka (Dakshina Kannada, Hassan, Udupi, Uttara Kannada), Peninsular to N E India; S & SE Asia, tropical Australia

Fimbristylis complanata (Retz.) Link, Hort. Berol. 1: 292.1827; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Scirpus complanatus Retz., Observ. Bot. 5: 14.1788

Cyperus complanatus (Retz.) Willd., Sp. Pl. 1: 270.1797

Isolepis complanata (Retz.) Roem. & Schult., Syst. Veg. 2: 119.1817

Trichelostylis complanata (Retz.) Nees, Contr. Bot. India 103.1834

Fimbristylis autumnalis var. *complanata* (Retz.) Kük., Bot. Jahrb. Syst. 59: 50.1924

Fimbristylis autumnalis var. *complanata* (Retz.) Barros, Anales Mus. Argent. Ci. Nat. Bernardino Rivadavia 41: 334.1945

Distribution: Karnataka (Bengaluru, Ballari, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru), (Andaman & Nicobar Islands, Andhra Pradesh, Assam, Chattisgarh, Himachal Pradesh, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Nagaland, Punjab, Sikkim, Tamil Nadu, West Bengal; tropical Africa, S America and Asia.

Fimbristylis consanguinea Kunth, Enum. Pl. 2: 228.1837

Iria consanguinea (Kunth) Kuntze, Revis. Gen. Pl. 2: 753.1891

Fimbristylis complanata var. *kraussiana* C.B.Clarke in Hook.f., Fl. Brit. India 6: 646.1893

Fimbristylis kraussiana Hochst. ex Hook.f., Handb. Fl. Ceylon 5: 63.1900

Fimbristylis horsfieldii C.B.Clarke, Kew Bull., Addit. Ser. 8: 25.1908

Distribution: Karnataka (Shivamogga), S Peninsular India; China, Madagascar, Sri Lanka, SAfrica, S E Asia.

Fimbristylis contorta C.E.C.Fisch., Bull. Misc. Inform. Kew 1931: 45.1931; Bhaskar & Kushalappa, Fl. Tumkur Dist. 104. 2013.

Distribution: Karnataka (Tumakuru)

Fimbristylis cymosa R.Br., Prodr. Fl. Nov. Holl. 228.1810; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Scirpus cymosus (R.Br.) Poir., Encycl. Suppl. 5: 99.1816

Scirpus strictus Roxb., Fl. Ind. 1,1: 226 1820

Fimbristylis spathacea Roth, Nov. Pl. Sp. 24 1821

Isolepis stricta (Roxb.) A.Dietr., Sp. Pl. 2: 121.1832

Iria cymosa (R.Br.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Fimbristylis cymosa subsp. *spathacea* (Roth) T.Koyama, Micronesica 1: 83.1964

Fimbristylis cymosa var. *spathacea* (Roth) T.Koyama, J. Jap. Bot. 46: 66.1971

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Vijayapura, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Mandya, Mysuru, Raichur, Udupi, Uttara Kannada), Peninsular & E India including Andaman & Nicobar Islands; tropical Asia & EAfrica..

Fimbristylis dichotoma (L.) Vahl, Enum. Pl. 2: 287.1805

Scirpus dichotomus L., Sp. Pl. 50.1753

Scirpus diphylloides Retz., Observ. Bot. 5: 15.1788

Fimbristylis diphylloides (Retz.) Vahl, Enum. Pl. 2: 289.1805

Scirpus arvensis Roxb., Fl. Ind. 1: 224.1820

Scirpus brevifolius Roxb., Fl. Ind. 1: 229.1820

Scirpus glomeratus Roxb., Fl. Ind. 1: 244. 1820.

Scirpus miliaceus Roxb., Fl. Ind. 1: 231.1820

Fimbristylis brevifolia (Roxb.) Wall. ex Schult., Mant. 2: 54.1824

Fimbristylis royeriana Nees, Contr. Bot. India 99.1834

Fimbristylis diphylloides var. *annua* (All.) C.B. Clarke, Fl. Brit. India 6(19): 637.1893

Fimbristylis diphylloides var. *nilagirica* C.B. Clarke in Hook.f., Fl. Brit. India 6: 637.1893

Fimbristylis annua var. *diphylloides* (Retz.) Kük., Wiss. Erg. Schwed. Rhod.-Kongo Exped. Erg.: 5 1921

Fimbristylis annua var. *paucispiculata* Blatt. & McCann, J. Bombay Nat. Hist. Soc. 37: 544.1934; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Fimbristylis dichotoma var. *paucispiculata* (Blatt. & McCann) Panigrahi, Bull. Bot. Surv. India 11: 111.1971

Fimbristylis dichotoma var. *nilagirica* (C.B.Clarke) Karthik., Fl. Ind. Numerat. - Monocot. 51.1984

Fimbristylis dichotoma var. *poladpurensis* D.P.Chavan, J. Econ. Taxon. Bot. 30: 721. 2006

subsp. **dichotoma**

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu.

Local/Common/Vernacular names: *Kadu sabsige hullu, Neeru sabsige hullu* (Kannada)

subsp. **podocarpa** (Nees) T.Koyama, Micronesica 1: 87.1964

Fimbristylis tomentosa Vahl, Enum. Pl. 2: 290.1805; V.P.Prasad & N.P.Singh, Sedges Karnataka 214. 2002.

Fimbristylis cincta Nees, Contr. Bot. India 98.1834

Fimbristylis podocarpa Nees, Contr. Bot. India 98.1834; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Fimbristylis diphylloides var. *podocarpa* (Nees) Kük., Bot. Jahrb. Syst. 69: 257.1938

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan Shivamogga, Udupi, Uttara Kannada), Peninsular India including Bihar & W Himalaya; Madagascar, S & SE Asia, Tropical Africa.

Fimbristylis dimorphonucifera Govind., Rheedea 7: 122.1997; V.P.Prasad & N.P.Singh, Sedges Karnataka 174. 2002.

Distribution: Karnataka (Dakshina Kannada, Udupi); endemic.

Fimbristylis dipsacea (Rottb.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 635.1893; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Scirpus dipsaceus Rottb., Descr. Icon. Rar. Pl. 56.1773

Echinolytrum dipsaceum (Rottb.) Desf., J. Bot. (Desvaux) 1: 20.1808

Isolepis dipsacea (Rottb.) Roem. & Schult., Syst. Veg. 2: 119.1817

Distribution: Karnataka (Dakshina Kannada, Hassan, Mysuru, Shivamogga, Udupi), throughout except NW India, Himalayas & Islands; S SE Asia & tropical Africa.

Fimbristylis eragrostis (Nees) Hance, J. Linn. Soc., Bot. 13: 132.1873

Abildgaardia eragrostis Nees, Contr. Bot. India 95.1834

Fimbristylis nigrobrunnea Thwaites, Enum. Pl. Zeyl. 434.1864

Iria eragrostis (Nees) Kuntze, Revis. Gen. Pl. 2: 753,1891

Fimbristylis eragrostis var. *nigrobrunnea* (Thwaites) S.S.Hooper, Fl. Hassan Distr. 678.1976; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), throughout India; Australia, China, Indonesia, Laos, Malaysia, Papua New Guinea, Sri Lanka, Taiwan, Thailand, Vietnam.

Fimbristylis falcata (Vahl) Kunth, Enum. Pl. 2: 239.1837; B.D.Sharma et al., Fl. Karnataka Anal. 309. 1984.
Scirpus falcatus Vahl, Enum. Pl. 2: 275.1805
Isolepis falcata (Vahl) Roem. & Schult., Syst. Veg. 2: 118.1817
Fimbristylis junciformis (Nees) Kunth, Enum. Pl. 2: 239.1837
Fimbristylis junciformis var. *conostachya* C.B.Clarke, J. Bot. (Morot) 3: 197.1889
Iria falcata (Vahl) Kuntze, Revis. Gen. Pl. 753.1891
Fimbristylis falcata var. *latifolia* (Nees) H.B.Naithani & M.B.Raizada, Indian Forester 103: 414.1977
 Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Chikkamagaluru, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; S & SE Asia.

Fimbristylis ferruginea (L.) Vahl, Enum. Pl. 2: 291.1805; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.
Scirpus ferrugineus L., Sp. Pl. 50.1753
Scirpus trannebbariensis Roth, Catal. Bot. 3: 6.1806
Scirpus globulosus Roxb., Fl. Ind. 1: 217.1820
Scirpus tristachyos Roxb., Fl. Ind. 1: 221.1820
Isolepis ferruginea (L.) Schltdl., Linnaea 6: 27.1831
Fimbristylis roxburghii A.Dietr., Sp. Pl. 2: 162.1832
Fimbristylis sieberiana Kunth, Enum. Pl. 2: 237.1837
Fuirena pubescens auct. non (Poir.) Kunth, 1837; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.
Fimbristylis andamanica Kurz, Rep. Veg. Andaman Isl. App. B: 21.1870
Fimbristylis ferruginea var. *sieberiana* (Kunth) Boeckeler, Linnaea 37: 17.1871
Iria ferruginea (L.) Kuntze, Revis. Gen. Pl. 2: 752.1891
Fimbristylis ferruginea subsp. *sieberiana* (Kunth) Lye, Nordic J. Bot. 2: 335.1982
 Distribution: Karnataka (throughout), throughout except NE India; Pantropical.

Fimbristylis insignis Thwites, Enum. Pl. Zeyl. 349.1864
Fimbristylis thwaitesii Boeckeler, Linnaea 37: 34 1871
Iria insignis (Thwaites) Kuntze, Revis. Gen. Pl. 2: 753.1891
 Distribution: Karnataka (Chitradurga), S India; China, SE Asia, Sri Lanka.

Fimbristylis karthikeyanii Kottaim. & Gnanasek., Telopea 18: 433. 2015
Fimbristylis breviculma Govind., Rheedea 6: 61.1996 non Boeck. 1858; V.P.Prasad & N.P.Singh, Sedges Karnataka 161. 2002.
 Distribution: Karnataka (Belagavi, Shivamogga); endemic.

Fimbristylis kingii Gamble ex Boeckeler, Beitr. Cyper. 2: 40.1890; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984 ('Fimbristylis kingii Clarke').
 Distribution: Karnataka (Bengaluru, Chitradurga, Shivamogga); endemic.

Fimbristylis lawiana (Boeckeler) J.Kern, Reinwardtia 4: 96.1956; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.
Scirpus lawianus Boeckeler, Linnaea 36: 497.1870
Trichelostylis digitata Dalzell ex Boeckeler, Flora 61: 35.1878
 Distribution: Karnataka (Belagavi, Vijayapura, Uttara Kannada), Goa, Gujarat, Maharashtra; endemic.

Fimbristylis littoralis Gaudich., Voy. Uranie 413.1829
Fimbristylis emarginata Wight ex Wall., Numer. List 3500.1831
 Distribution: Karnataka (throughout), throughout India; Pantropical.

Fimbristylis merrillii J.Kern, Blumea 8: 135.1955; V.P.Prasad & N.P.Singh, Sedges Karnataka 189. 2002.
 Distribution: Karnataka (Belagavi, Bengaluru, Dharwad, Hassan, Mysuru, Uttara Kannada), W India; China, S E Asia, Australia..

Fimbristylis microcarya F.Muell., Fragm. 1: 200.1859; V.P.Prasad & N.P.Singh, Sedges Karnataka 189. 2002.

Iria microcarya (F.Muell.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Fimbristylis complanata var. *microcarya* (F.Muell.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 646.1893

Fimbristylis autumnalis var. *microcarya* (F.Muell.) Kük., Bot. Jahrb. Syst. 69: 258. 1938

Distribution: Karnataka (Belagavi Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), Peninsular India & Himalayas; E Asia, Australia.

Fimbristylis monospicula Govind., Proc. Indian Acad. Sci., B 79: 169.1974; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Distribution: Karnataka (Mysuru); endemic.

Fimbristylis monticola Hochst. ex Steud., Syn. Pl. Glumac. 2: 111.1855

Fimbristylis tenuifolia Thwaites, Enum. Pl. Zeyl. 434.1864

Distribution: Karnataka (Dakshina Kannada, Udupi)

Fimbristylis narayanii C.E.C.Fisch., Bull. Misc. Inform. Kew 1931: 46.1931; V.P.Prasad & N.P.Singh, Sedges Karnataka 195. 2002.

Distribution: Karnataka (Dakshina Kannada, Udupi), S India & NW Himalaya.

Fimbristylis ovata (Burm.f.) J.Kern, Blumea 15: 126.1967; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Carex ovata Burm.f., Fl. Indica 194.1768

Cyperus monostachyos L., Mant. Pl. 2: 180.1771

Abildgaardia monostachya (L.) Vahl, Enum. Pl. 2: 296.1805

Cyperus indicus Rich. ex Pers., Syn. Pl. 1: 65. 1805.

Scirpus schoenoides Roxb., Fl. Ind. 1: 221.1820

Iria indica (Rich. ex Pers.) Schult., Mant. 2: 94.1824

Abildgaardia indica (Rich. ex Pers.) Nees, Linnaea 9: 289.1834

Abildgaardia rottboelliana Nees ex Hook. & Arn., Bot. Beechey Voy. 272.1838

Fimbristylis monostachya (L.) Hassk., Pl. Jav. Rar. 61.1848

Iria monostachya (L.) Kuntze, Revis. Gen. Pl. 2: 752.1891

Scirpus monostachyus (L.) Kuntze, Revis. Gen. Pl. 3(2): 337.1898

Abildgaardia ovata (Burm.f.) Kral, Sida 4: 72.1971

Distribution: Karnataka (throughout), throughout India; Pantropical.

Fimbristylis polytrichoides (Retz.) R.Br., Prodr. Fl. Nov. Holl. 226.1810; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Scirpus polytrichoides Retz., Observ. Bot. 4: 11.1786

Apostemon polytrichoides (Retz.) Raf., J. Phys. Chim. Hist. Nat. Arts 89: 105.1819

Iria polytrichoides (Retz.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Distribution: Karnataka (Dakshina Kannada, Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Peninsular & E India; tropical Africa, Asia & Australia

Fimbristylis pseudomicrocarya Govind., Proc. Indian Acad. Sci., Pl. Sci. 100: 77.1990; V.P.Prasad & N.P.Singh, Sedges Karnataka 201. 2002.

Distribution: Karnataka (Chikkamagalur, Dakshina Kannada, Udupi), Kerala; endemic.

Fimbristylis pubisquama J.Kern, Blumea 8: 131 1955; Rao et al., J. Econ. Taxon. Bot. 35: 154. 2011.

Fimbristylis compressa Boeckeler, Linnaea 38: 387 1874.

Distribution: Karnataka (Uttara Kannada), Kerala; Sri Lanka, Cambodia, Myanmar, Thailand, Vietnam, North Australia.

Fimbristylis pustulosa Govind., Proc. Indian Acad. Sci., Pl. Sci. 90: 444.1981; Kesh. Murthy & Yogan., Fl. Coorg 508. 1990.
 Distribution: Karnataka (Kodagu), Tamil Nadu; endemic.

Fimbristylis quinquangularis (Vahl) Kunth, Enum. Pl. 2: 229.1837
Scirpus miliaceus L., Syst. Nat. ed. 10 2: 868. 1759.
Fimbristylis miliacea (L.) Vahl, Enum. Pl. 2: 229.1805; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.
Scirpus benghalensis Pers., Syn. Pl. 1: 68.1805
Scirpus quinquangularis Vahl, Enum. Pl. 2: 279.1805
Fimbristylis benghalensis (Pers.) Roem. & Schult., Syst. Veg. 2: 94.1817
Scirpus pentagonus Roxb., Fl. Ind. 1: 218.1820
Scirpus plantagineus Roxb., Fl. Ind. 1: 212. 1820.
Isolepis pentagona (Roxb.) Schult., Mant. 2: 69.1824

Scirpus salbundius Buch.-Ham. ex Wall., Numer. List 3499 1831
Scirpus tetragonus Roxb., Fl. Ind. ed. 2 ,1: 228.1832
Trichelostylis miliacea (L.) Nees, Contr. Bot. India 103 1834
Trichelostylis quinquangularis (Vahl) Nees, Contr. Bot. India 104 1834
 Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hasan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada),throughout; Asia except West Asia, tropical Africa & Australia.

Fimbristylis schoenoides (Retz.) Vahl, Enum. Pl. 2: 286.1805; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Scirpus schoenoides Retz., Observ. Bot. 5: 14. 1788.
Scirpus bispicatus J.König ex Roxb., Fl. Ind. 1: 220.1820
Scirpus monostachyos J.König ex Roxb., Fl. Ind. 1: 219.1820
Isolepis monostachya (J.Koenig) ex Roxb.) Spreng., Neue Entdeck. Pflanzenk. 3: 11.1822
Isolepis bispicata (J.Koenig) ex Roxb.) Schult., Mant. 2: 61.1824
Iria bispicata (J.Koenig) ex Roxb.) Kuntze, Revis. Gen. Pl. 2: 753.1891
Iriha schoenoides (Retz.) Kuntze, Revis. Gen. Pl. 2: 752.1891

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamogluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru, Raichur, Shivamogga, Udupi,Uttra Kannada)

Local/Common/Vernacular names: *Gedde gundu hullu, Kadu gandu hullu* (Kannada)

Fimbristylis semidisticha Govind., Proc. Indian Acad. Sci., B 78: 51.1973; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru), Tamil Nadu; endemic.

Fimbristylis sericea (Poir.) R.Br., Prodr. Fl. Nov. Holl. 228.1810; Bhaskar & Kushalappa, Fl. Tumkur Dist.105. 2013.

Scirpus sericeus Poir., Encycl. Suppl. 5: 99.1817
Iria sericea (Poir.) Kuntze, Revis. Gen. Pl. 2: 753.1891
 Distribution: Karnataka (Tumakuru),

Fimbristylis squarrosa Vahl, Enum. Pl. Obs. 2: 289. 1805.

var. **esquarrosa** Makino, Bot. Mag. (Tokyo) 17: 47.1903; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.
Fimbristylis velata R.Br., Prodr. Fl. Nov. Holl. 227.1810
Scirpus velatus (R.Br.) Poir., Encycl. Suppl. 5: 99.1817
Iria velata (R.Br.) Kuntze, Revis. Gen. Pl. 2: 753.1891
Fimbristylis squarrosa var. *velata* (R.Br.) C.B.Clarke ex Cheeseman, Man. New Zealand Fl. 770.1906
Fimbristylis squarrosa subsp. *esquarrosa* (Makino) T.Koyama, Fl. Taiwan 5: 247.1978
 Distribution: Karnataka (Hassan, Mysuru, Uttara Kannada), S India hill stations; Australia, New Zealand,, E & SE Asia, Polynesia.

Fimbristylis tenera Schult., Mant. 2: 57.1824; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Scirpus tenellus Roxb., Fl. Ind. 1: 224.1820

Trichelostylis tenella Nees, Contr. Bot. India 103.1834

Fimbristylis glabra Hochst. ex Steud., Syn. Pl. Glumac. 2: 111.1855; B.D.Sharma et al., Fl. Karnataka Anal. 310. 1984.

Fimbristylis oxylepis Steud., Syn. Pl. Glumac. 2: 110.1855

Iria tenera (Schult.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Fimbristylis tenera var. *oxylepis* (Steud.) C.B. Clarke, Fl. Brit. India 6: 642.1893; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Vijayapura, Chitradurga, Dakshina Kannada, Hassan, Mandya, Mysuru, Raichur, Shivamogga, Udupi, Uttara Kannada), Peninsular to N India; Sri Lanka, tropical Africa.

Fimbristylis tetragona R.Br., Prodr. Fl. Nov. Holl. 226.1810; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Scirpus tetragonus (R.Br.) Poir., Encycl. Suppl. 5: 98. 1817.

Iria tetragona (R.Br.) Kuntze, Revis. Gen. Pl. 2: 753.1891

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mandya, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India ; Tropical E, S & SE Asia & Australia.

Fimbristylis tuckeri Viji , Pandur. & Deepu, Kew Bull. 71(3)-38: 1. 2016.

Distribution: Karnataka (Kodagu), Kerala; endemic.

Fimbristylis virella Govind., Rheedea 8: 93.1998; V.P.Prasad & N.P.Singh, Sedges Karnataka 216. 2002.

Distribution: Karnataka (Udupi), Kerala; endemic.

Fimbristylis woodrowii C.B.Clarke, Bull. Misc. Inform. Kew 1898: 227.1898; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Udupi,Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Fuirena Rottb.

Fuirena ciliaris (L.) Roxb., Hort. Bengal. 81.1814; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Scirpus ciliaris L., Mant. Pl. 2: 182.1771

Fuirena glomerata Lam., Tabl. Encycl. 1: 150.1791

Scirpus pilosus Retz., Observ. Bot. 6: 19. 1791

Scirpus aristatus Willd., Sp. Pl. 1: 300. 1797

Fuirena rottboellii Nees, Contr. Bot. India 94.1834

Fuirena tuvensis auct. non M.B.Deshp. & Shah, 1969; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mandya, Mysuru, Shivamogga, Udupi, Uttara Kannada),throughout warmer parts; Paleotropical.

Local/Common/Vernacular name: *Petlu gori hullu* (Kannada)

Fuirena cuspidata (Roth) Kunth, Enum. Pl. 2: 187.1837

Scirpus cuspidatus Roth, Nov. Pl. Sp. 31.1821

Isolepis cuspidata (Roth) A.Dietr., Sp. Pl. 2: 120.1832

Fuirena wallichiana Kunth, Enum. Pl. 2: 182.1837; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Fuirena wallichiana var. *evoluta* C.B.Clarke in Hook.f.,Fl. Brit. India 6: 666.1893

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Chikkamagaluru, Dharwad, Hassan, Mysuru)

Fuirena trilobites C.B.Clarke in Hook.f., Fl. Brit. India 6: 666.1893; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Hassan, Kolar, Raichur, Tumakuru, Uttara Kannada)

Fuirena umbellata Rottb., Descr. Icon. Rar. Pl. 70.1773; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Fuirena pentagona Wight & Arn. ex Nees, Contr. Bot. India 93.1834

Scirpus umbellatus (Rottb.) Kuntze, Revis. Gen. Pl. 3(2): 337.1898

Fuirena seriata C.B.Clarke, Bull. Soc. Bot. France 54(8): 28.1907

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada)

Fuirena uncinata (Willd.) Kunth, Enum. Pl. 2: 184.1837; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Scirpus uncinatus Willd., Sp. Pl. 1: 300.1797

Isolepis uncinata (Willd.) Roem. & Schult., Syst. Veg. 2: 111.1817

Fuirena capitata T.Koyama in Dasanayake & Fosberg, Revised Handb. Fl. Ceylon 5: 151.1985

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Hassan, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Peninsular India; Sri Lanka.

Hypolytrum L. C. Rich.

Hypolytrum nemorum (Vahl) Spreng., Syst. Veg. 1: 233.1825; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Schoenus nemorum Vahl, Symb. Bot. 3: 8.1794

Hypolytrum latifolium Rich. ex Pers., Syn. Pl. 1: 70.1805

Hypaelyptum nemorum (Vahl) P.Beauv., Fl. Oware 2: 13.1810

Tunga diandra Roxb., Fl. Ind. 1: 188.1820

Hypolytrum diandrum (Roxb.) A.Dietr., Sp. Pl. 2: 365.1833

Hypolytrum giganteum Wall. ex Nees, Contr. Bot. India 93.1834

Hypolytrum wightianum Boeckeler, Linnaea 37: 130.1871

Hypolytrum penangense C.B.Clarke in Hook.f., Fl. Brit. India 6: 679.1894

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Peninsular to NE India including Andaman & Nicobar Islands; Australia, Fiji, Polynesia, tropical E & SE Asia.

Kyllinga Rottb.

Kyllinga brevifolia Rottb., Descr. Icon. Rar. Pl. 13.1773; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Cyperus brevifolius (Rottb.) Hassk., Cat. Hort. Bot. Bogor. 24.1844

Kyllinga intermegia var. *oligostachya* C.B.Clarke, J. Linn. Soc., Bot. 36: 224.1903

var. brevifolia

Distribution: Karnataka (Belagavi, Bengaluru, Vijayapura, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mandya, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; pantropical.

var. stellulata (J.V.Suringar) Ohwi, Bot. Mag. (Tokyo) 56: 199.1942

Cyperus brevifolius var. *stellulatus* (J.V.Suringar) Kük., Pflanzren. IV, 20(101): 603.1936

Kyllinga melanosperma var. *stellulata* (J.V.Suringar) Tang & F.T.Wang, Fl. Reipubl. Popul. Sin. 11: 189.1961

Cyperus brevifolius var. *stellulatus* (J.V.Suringar) Hooper, Fl. Hassan 687. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Distribution: Karnataka (Hassan), S India, China, Indonesia, Papua New Guinea, Philippines.

Kyllinga eglandulosa Govind. & Ramani, J. Econ. Taxon. Bot. 18: 335.1994; V.P. Prasad & N.P.Singh, Sedges Karnataka 216. 2002.

Distribution: Karnataka (Chikkamagaluru); endemic.

Kyllinga melanosperma Nees, Contr. Bot. India 91.1834; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Kyllinga brevifolia Nees, Contr. Bot. India 91. 1834.

Cyperus melanosperma (Nees) Suringar, Het Gesl. Cyp. Mal. Archip. 50, pl. 2, f. 8 50.1898

Cyperus melanospermus (Nees) Suringar, Cyperus Mal. Arch. 50.1898

subsp. **melanosperma**

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Davanagere, Hassan, Kodagu, Mysuru, Uttara Kannada), Peninsular to E India; S & SE Asia, tropical Africa.

subsp. **bifolia** (Miq.) Karthik., Fl. Ind. Enumerat. - Monocot. 60.1989

Kyllinga bifolia Miq., Fl. Ned. Ind. 3: 293.1856; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Cyperus melanospermus subsp. *bifolius* (Miq.) J.Kern, Reinwardtia 3: 62.1954

Distribution: Karnataka (Belagavi, Kodagu, Uttara Kannada), S to N India; SE Asia,

Kyllinga nemoralis (J.R.Forst. & G.Forst.) Dandy ex Hutch. & Dalziel, Fl. W. Trop. Afr. 2: 487.1936; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Thryocephalon nemorale J.R.Forst. & G.Forst., Char. Gen. Pl. 65.1775

Kyllinga monocephala Stokes, Bot. Mat. Med. 1: 120. 1812.

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Chitradurga, Davanagere, Dharwad, Kalburgi, Hassan, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout India; pantropical.

Local/Common/Vernacular name: *Choodle* (Kannada)

Kyllinga odorata Vahl, Enum. Pl. Obs. 2: 382. 1805.

subsp. **cylindrica** (Nees) T.Koyama, Gard. Bull. Singapore 30: 161.1977

Kyllinga odorata auct. Non Vahl, 1805; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Kyllinga cylindrica Nees, Contr. Bot. India 91.1834

Kyllinga odorata var. *cylindrica* (Nees) Kük., J. Straits Branch Roy. Asiat. Soc. 76: 80.1917

Cyperus sesquiflorus var. *cylindrica* (Nees) Kük., Pflanzenr. 4(20, Heft. 101): 593.1935

Cyperus sesquiflorus var. *cylindricus* (Nees) Kük., Pflanzenr. IV, 20(101): 593.1936

Distribution: Karnataka (Bengaluru, Hassan, Kodagu, Mysuru), throughout India; S E asia, tropical Africa.

Kyllinga squamulata Vahl, Enum. Pl. 2: 381.1805; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Kyllinga squamulosa Kunth, Enum. Pl. 2: 135.1837

Kyllinga metzii Hochst. ex Steud., Syn. Pl. Glumac. 2: 70.1854

Cyperus metzii (Hochst. ex Steud.) Mattf. & Kük., Pflanzenr. IV, 20(101): 25.1935

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Udupi, Uttara Kannada), throughout India; tropical Africa & America.

Kyllinga tenuifolia Steud., Syn. Pl. Glumac. 2: 69.1854; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Cyperus tenuifolius (Steud.) Dandy, Cat. Vasc. Pl. S. Tome 363.1944

Distribution: Karnataka (Bengaluru, Bidar, Vijayapura, Chikkamagaluru, Dakshina Kannada, Kalburgi, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi), S to NE India; Africa, Australia, China, SE Asia

Lipocarpha R.Br.

Lipocarpha chinensis (Osbeck) J.Kern, Blumea, Suppl. 4: 167.1958; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Scirpus chinensis Osbeck, Dagb. Ostind. Resa 220.1757

Hypolytrum argenteum Kunth, Nov. Gen. Sp. 1: 218.1816

Lipocarpha argentea (Kunth) R.Br., Narr. Exped. Zaire 477.1818

Tunga laevigata Roxb., Fl. Ind. 1: 188.1820

Hypolytrum laevigatum (Roxb.) Spreng., Syst. Veg. 1: 233 1824

Lipocarpha humboldtiana Nees, Linnaea 9: 287.1834

Lipocarpha laevigata (Roxb.) Nees, Contr. Bot. India 92 1834

Schoenus laevigatus Roxb. ex Nees, Contr. Bot. India 92.1834

Rikliella chinensis (Osbeck) M.R.Almeida, Fl. Maharashtra 5B: 381.2009

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mandya, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India ; S & SE Asia, tropical & S Africa, tropical Australia.

Lipocarpha gracilis (Rich. ex Pers.) Nees, Linnaea 9: 287.1834

Hypaelyptum sphacelatum Vahl, Enum. Pl. Obs. 2: 283.1805

Hypolytrum gracile Rich. ex Pers., Syn. Pl. 1: 70.1805

Tunga triceps Roxb., Fl. Ind. 1: 187.1820

Hypaelyptum gracile (Rich. ex Pers.) Steud., Nomencl. Bot. 1: 419.1821

Hypolytrum triceps (Roxb.) A.Dietr., Sp. Pl. 2: 363.1832

Hypolytrum ceylanicum B.Heyne ex Nees, Linnaea 9: 288.1834

Lipocarpha triceps (Roxb.) Nees, Contr. Bot. India 92.1834

Lipocarpha sphacelata (Vahl) Kunth, Enum. Pl. 2: 267 1837; B.D.Sharma et al., Fl. Karnataka Anal. 312. 1984.

Lipocarpha sphacelata var. *gracilis* (Rich. ex Pers.) Boeckeler, Linnaea 37: 116.1871

Cyperus ceylanicus (Vahl) T.Koyama, Bot. Mag. (Tokyo) 73: 438.1960

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Mandya, Mysuru, Shivamogga, Tumakuru, Udupi,Uttara Kannada), throughout India; Nepal, Thailand, tropical Africa & America.

Lipocarpha hemisphaerica (Roth) Goetgh., Wageningen Agric. Univ. Pap. 89(1): 37.1989

Isolepis hemisphaerica (Roth) A.Dietr., Sp. Pl. 2: 109.1832

Hemicarpha isolepis Nees, Edinburgh New Philos. J. 17: 263.1834; B.D.Sharma et al., Fl. Karnataka Anal. 311. 1984.

Scirpus isolepis (Nees) Boeckeler, Linnaea 36: 498.1870

Distribution: Karnataka (Hassan, Mysuru), Peninsular to E India,tropical Africa, Thailand

Local/Common/Vernacular name: *Sanna gundu hullu* (Kannada)

Lipocarpha kernii (Raymond) Goetgh., Wageningen Agric. Univ. Pap. 89(1): 42.1989

Scirpus kernii Raymond, Naturaliste Canad. 86: 230.1959

Isolepis kernii (Raymond) Lye, Bot. Not. 124: 479.1971

Rikliella kernii (Raymond) J.Raynal, Adansonia n.s., 13: 155.1973; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Distribution: Karnataka (Hassan), Peninsular India; tropical Africa.

Lipocarpha squarrosa (L.) Goetgh., Wageningen Agric. Univ. Pap. 89(1): 71.1989

Scirpus squarrosus L., Mant. Pl. 2: 181.1771

Isolepis squarrosa (L.) Roem. & Schult., Syst. Veg. 2: 111.1817

Scirpus squamosus Link, Jahrb. Gewächsk. 1(3): 79.1820

Rikliella squarrosa (L.) J.Raynal, Adansonia n.s., 13: 154.1973; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Distribution: Karnataka (throughout), throughout India ; S & SE Asia.

Local/Common/Vernacular name: *Kadu usumani hullu* (Kannada)

Pycreus P. Beauv.

Pycreus diaphanus (Schrad. ex Roem. & Schult.) S.S.Hooper & T.Koyama, J. Jap. Bot. 51: 316.1976; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984.

Cyperus diaphanus Schrad. ex Roem. & Schult., Mant. 2: 477.1824

Cyperus latispicatus var. *acaulis* C.B.Clarke, J. Linn. Soc., Bot. 21: 41.1884

Pycreus latispicatus (Boeckeler) C.B.Clarke in Hook.f., Fl. Brit. India 6: 696.1894

Cyperus latispicatus Boeckeler ex C.B.Clarke, J. Linn. Soc., Bot. 36: 202. 1903

Cyperus latispicatus var. *diaphanus* (Schrad. ex Roem. & Schult.) Kük., Pflanzenr. IV, 20(101): 393.1936

Cyperus latispicatus var. *gracilescens* Kük., Pflanzenr. IV, 20(101): 393.1936

Cyperus latispicatus var. *laxus* Kük., Pflanzenr. IV, 20(101): 393.1936

Cyperus diaphanus var. *gracilescens* (Kük.) H.O.Saxena, Indian Forester 99: 505.1973

Pycreus diaphanus var. *gracilescens* (Kük.) S.S.Hooper, Fl. Hassan Distr. 692.1976; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984.

Pycreus diaphanus var. *acaulis* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 64.1989

Pycreus diaphanus var. *laxus* (Kük.) H.B.Naithani & S.Biswas, Flow. Pl. India, Nepal & Bhutan 471.1990

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Peninsular India; C and SE Asia

Pycreus flavidus (Retz.) T.Koyama, J. Jap. Bot. 51: 316.1976

Cyperus globosus auct non Forsskal, All., Fl. Pedem. 1: 49.1785

Cyperus flavidus Retz., Observ. Bot. 5: 13. 1789.

Cyperus strictus Lam., Tabl. Encycl. 1: 146.1791

Cyperus capillaris J.König ex Roxb., Fl. Ind. 1: 194.1820

Cyperus lamarckianus Schult., Mant. 2: 108.1824

Pycreus globosus Rchb., Fl. Germ. Excurs. 140. 1831; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Cyperus nilagiricus Hochst. ex Steud., Syn. Pl. Glumac. 2: 2.1854

Cyperus globosus var. *strictus* (Lam.) C.B.Clarke, J. Linn. Soc., Bot. 20:280.1883; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Cyperus globosus var. *tortuosa* C.B.Clarke, J. Linn. Soc., Bot. 20: 280. 1883

Cyperus globosus var. *khasianus* C.B.Clarke, J. Linn. Soc., Bot. 21: 48.1884

Cyperus globosus var. *nilagiricus* (Hochst. ex Steud.) C.B.Clarke, J. Linn. Soc., Bot. 21: 49.1884; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Cyperus semiteres B.Heyne ex C.B.Clarke, J. Linn. Soc., Bot. 21: 49.1884

Pycreus capillaris (J.Koenig ex Roxb.) Nees ex C.B.Clarke in Hook.f., Fl. Brit. India 6: 591.1893

Chlorocyperus nilagiricus (Hochst. ex Steud.) Rikli, Jahrb. Wiss. Bot. 27: 564.1895

Pycreus globosus var. *erectus* C.B.Clarke, J. Linn. Soc., Bot. 34: 15.1898

Chlorocyperus globosus (Rchb.) Palla, Allg. Bot. Z. Syst. 6: 61.1900

Pycreus globosus var. *nilagiricus* (Hochst. ex Steud.) C.B.Clarke, J. Linn. Soc., Bot. 36(251): 204.1903

Pycreus globosus var. *strictus* (Lam.) Domin, Biblioth. Bot. 85: 419.1915; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Pycreus strictus (Lam.) Alston in Trimen, Handb. Fl. Ceylon 4(Suppl.): 307.1931

Cyperus globosus var. *erectus* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 356.1936

Cyperus flavidus var. *erectus* (C.B.Clarke) Korlah., Bull. Bot. Surv. India 9: 237.1968

Cyperus flavidus var. *khasianus* (C.B.Clarke) Korlah., Bull. Bot. Surv. India 9: 237.1968

Pycreus flavidus var. *erectus* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 65.1989

Pycreus flavidus var. *khasianus* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 65.1989

Pycreus lamarckianus (Schult.) M.R.Almeida, Fl. Maharashtra 5B: 372.2009

Pycreus lamarckianus var. *strictus* (Lam.) M.R.Almeida, Fl. Maharashtra 5B: 377.2009

Distribution: Karnataka (throughout), throughout India; Africa, E & SE Asia, S Europe.

Pycreus kanarensis V.P.Prasad & N.P.Singh, J. Econ. Taxon. Bot. 21: 667 .1998; Sedges Karnataka 272. 2002.

Pycreus flavescens auct. non (L.) P.Beauv. ex Rchb., 1830; B.D.Sharma et al., Fl. Karnataka Anal. 313. 1984. Distribution: Karnataka (Belagavi, Uttara Kannada); endemic.

Pycreus macrostachyos (Lam.) J.Raynal, Kew Bull. 23: 314.1969; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Cyperus macrostachyos Lam., Tabl. Encycl. 1: 147.1791

Cyperus albomarginatus Mart. & Schrad. ex Nees, Fl. Bras. 2(1): 9.1842

Cyperus retusus A.Rich., Tent. Fl. Abyss. 2: 485.1850

Cyperus hochstetteri var. *russa* C.B.Clarke, J. Linn. Soc., Bot. 20: 282 1883

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Mandya, Mysuru, Udupi, Uttara Kannada), Peninsular India; Pantropical.

Pycreus mahadevanii Govind., J. Econ. Taxon. Bot. 20: 299.1996; V.P.Prasad & N.P.Singh, Sedges Karnataka 275. 2002.

Distribution: Karnataka (Shivamogga); endemic.

Pycreus malabaricus C.B.Clarke, J. Linn. Soc., Bot. 34: 12.1898

Cyperus malabaricus (C.B.Clarke) T.Cooke, Fl. Bombay 2: 856.1908

Cyperus decumbens Govind., J. Indian Bot. Soc. 52: 72.1974

Cyperus atroglumosa Govind., Proc. Indian Acad. Sci., B 81: 187.1975

Cyperus lurida Govind., Proc. Indian Acad. Sci., B 81: 189.1975

Cyperus plurinodosa Govind., Proc. Indian Acad. Sci., B 81: 192.1975

Cyperus stricticulmis Govind., Proc. Indian Acad. Sci., B 81: 194.1975

Pycreus atroglumosus (Govind.) P.Singh & V.Singh, J. Econ. Taxon. Bot. 5: 467.1984

Pycreus decumbens (Govind.) P.Singh & V.Singh, J. Econ. Taxon. Bot. 5: 467.1984

Pycreus lurida (Govind.) P.Singh & V.Singh, J. Econ. Taxon. Bot. 5: 467.1984

Pycreus plurinodosus Pycreus plurinodosus (Govind.) P.Singh & V.Singh, J. Econ. Taxon. Bot. 5: 467.1984

Pycreus stricticulmis (Govind.) P.Singh & V.Singh, J. Econ. Taxon. Bot. 5: 467.1984

Pycreus govindarajalui V.S.Raju, Taxon 40: 103.1991

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), W Peninsular India; Myanmar.

Pycreus polystachyos (Rottb.) P.Beauv., Fl. Oware 2: 48.1816; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Cyperus paniculatus Rottb., Descr. Icon. Rar. Pl. 40.1773

Cyperus polystachyos Rottb., Descr. Icon. Rar. Pl. 39.1773

Cyperus caespitosus Poir., Encycl. 7: 250.1806

Pycreus paniculatus (Rottb.) Nees, Linnaea 9: 283.1834

Cyperus thouarsii Kunth, Enum. Pl. 2: 12.1837

Cyperus hahnianus Boeckeler, Flora 61: 138.1878

Cyperus hookerianus Arn. ex C.B.Clarke, J. Linn. Soc., Bot. 20: 280.1883

Cyperus polystachyos var. *hookerianus* C.B.Clarke, J. Linn. Soc., Bot. 20: 280.1883

Cyperus polystachyus var. *thouarsii* (Kunth) C.B.Clarke, J. Linn. Soc., Bot. 20: 280.1883

Cyperus polystachyos var. *paniculatus* (Rottb.) C.B.Clarke, J. Linn. Soc., Bot. 21: 55.1884

Cyperus circinatus Ridl., J. Linn. Soc., Bot. 27: 64.1890

Pycreus polystachyos var. *paniculatus* (Rottb.) Donn.Sm., Enum. Pl. Guatemp. 3: 86.1893

Pycreus caespitosus (Poir.) C.B.Clarke in T.A. Durand & H. Schinz, Conspl. Fl. Afric. 5: 535.1894

Chlorocyperus polystachyus (Rottb.) Rikli, Jahrb. Wiss. Bot. 27: 563.1895

Pycreus polystachyos var. *hahnianus* (Boeckeler) C.B.Clarke in Urb., Symb. Antill. 2: 18.1900

Cyperus polystachyos var. *hahnianus* (Boeckeler) Kük., Pflanzenr. IV, 20(101): 370.1936

Cyperus polystachyos var. *thouarsii* (Kunth) Kük., Pflanzenr. IV, 20(101): 373.1936

Pycreus odoratus var. *hahnianus* (Boeckeler) Stehlé, Caribbean Forester 5: 184.1944

Pycreus polystachyos var. *circinatus* (Ridl.) H.B.Naithani & S.Biswas, Flora. Pl. India, Nepal & Bhutan 472.1990

var. *polystachyos*

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada)

var. *microdontus* (Torr.) Govaerts, World Checklist Cyperaceae 610.2007

Cyperus polystachyos var. *holosericeus* (Link) C.B.Clarke, J. Linn. Soc., Bot. 21: 55 1884

Pycreus polystachyos var. *laxiflorus* (Benth.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 592 1893; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Distribution: Karnataka (Bengaluru, Chitradurga, Dakshina Kannada, Hassan, Tumakuru)

Distribution: Karnataka (Hassan)

***Pycreus pumilus* (L.) Nees, Linnaea 9: 283.1834; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.**

Cyperus pumilus L., Cent. Pl. 2: 6.1756

Cyperus punctatus Roxb., Fl. Ind. 1: 197.1820

Cyperus leucolepis J.Carey ex C.B.Clarke, J. Linn. Soc., Bot. 21: 61.1884

Cyperus pumilus var. *punctatus* C.B.Clarke, J. Linn. Soc., Bot. 21: 46.1884

Cyperus pumilus var. *leucolepis* (C.B.Clarke) Kük., Pflanzenr. IV, 20(101): 378.1936

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluruu, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Dharwad, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Mahsrashtra, Tamil Nadu; tropical Africa, E & SE Asia, Australia.

***Pycreus puncticulatus* (Vahl) Nees, Fl. Bras. 2(1): 10.1842; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.**

Cyperus puncticulatus Vahl, Enum. Pl. 2: 348.1805

Cyperus puncticulatus var. *quinquagintiflorus* C.B.Clarke, J. Linn. Soc., Bot. 21: 69.1884

Pycreus puncticulatus var. *quinquagintiflorus* (C.B.Clarke) C.B.Clarke in Hook.f., Fl. Brit. India 6: 594.1893

Distribution: Karnataka (Chikkamagaluru, Mysuru, Uttara Kannada), S India, Indonesia, Sri Lanka.

***Pycreus sanguinolentus* (Vahl) Nees, Linnaea 9: 283.1834; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.**

Cyperus sanguinolentus Vahl, Enum. Pl. 2: 351.1805

Cyperus eragrostis var. *micronyx* C.B.Clarke, J. Linn. Soc., Bot. 21: 59.1884

Cyperus sanguinolentus subsp. *sacer* Korlahalli, Bull. Bot. Surv. India 9: 235.1968

Pycreus sanguinolentus subsp. *sacer* (Korlahalli) Karthik., Fl. Ind. Enumerat. - Monocot. 66.1989

Pycreus sanguinolentus var. *micronyx* (C.B.Clarke) Karthik., Fl. Ind. Enumerat. - Monocot. 66.1989

Pycreus sanguinolentus subsp. *micronyx* (C.B.Clarke) H.B.Naithani & S.Biswas, Flow. Pl. India, Nepal & Bhutan 472.1990

Pycreus sanguinolentus subsp. *sacer* (Korlahalli) H.B.Naithani & S.Biswas, Flow. Pl. India, Nepal & Bhutan 473. 1990

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluruu, Chitradurga, Davanagere, Hassan, Kodagu, Mysuru, Uttara Kannada), Goa, Kerala, Mahsrashtra, Tamil Nadu; Sri Lanka, Timor

***Pycreus stramineus* C.B.Clarke in Hook.f., Fl. Brit. India 6: 589.1893; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.**

Cyperus stramineus Nees, Contr. Bot. India 74.1834

Cyperus coromandelinus Boeckeler, Linnaea 35: 480. 1868.

Distribution: Karnataka (Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi)

Queenslandiella Dom.

Queenslandiella hyalina (Vahl) Ballard, Hooker's Icon. Pl. 33: t. 3208.1933; B.D.Sharma et al., Fl. Karnataka Anal. 314. 1984.

Cyperus hyalinus Vahl, Enum. Pl. 2: 320.1805

Pycrus hyalinus (Vahl) Domin, Biblioth. Bot. 85: 417.1916

Mariscopsis hyalinus (Vahl) Ballard, Bull. Misc. Inform. Kew 1932: 458.1932

Kyllinga hyalina (Vahl) T.Koyama, J. Jap. Bot. 51: 317.1976

Distribution: Karnataka (Bengaluru, Davanagere, Hassan, Mandya, Mysuru), Peninsular to E India; E Africa, Sri Lanka, SE Asia, tropical Australia.

Remirea Aubl.

Remirea maritima Aubl., Hist. Pl. Guiane 1: 45.1775; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Miegia maritima (Aubl.) Willd., Sp. Pl. 1(1): 311.1797

Remirea pedunculata R.Br., Prodr. Fl. Nov. Holl. 236.1810

Remirea wightiana Nees, Contr. Bot. India 92.1834

Remirea maritima var. *pedunculata* (R.Br.) Benth., Fl. Austral. 7: 347.1878

Duval-jouvea maritima (Aubl.) Palla, Denkschr. Kaiserl. Akad. Wiss., Wien. Math.-Naturwiss. Kl. 89: 500.1913

Cyperus pedunculatus (R.Br.) J.Kern, Acta Bot. Neerl. 7: 798.1958; B.D.Sharma et al., Fl. Karnataka Anal. 307. 1984.

Mariscus pedunculatus (R.Br.) T.Koyama, Gard. Bull. Singapore 30: 159.1977

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), throughout Coastal areas in India; Pantropical. coasts.

Rhynchospora Vahl

Rhynchospora colorata (L.) H.Pfeiff., Repert. Spec. Nov. Regni Veg. 38: 89. 1935

Schoenus coloratus L., Sp. Pl. 43.1753

Kyllinga monocephala Rottb., Descr. Icon. Rar. Pl. 13. 1773.

Cyperus kyllingia Endl., Cat. Horti Vindob. 1: 94.1842

Dichromena colorata (L.) C.L.Hitchc., Rep. (Annual) Missouri Bot. Gard. 4: 141.1893

Kyllinga colorata (L.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 630.1917

Distribution: Karnataka (Bengaluru, Mysuru),

Rhynchospora corymbosa (L.) Britton, Trans. New York Acad. Sci. 11: 84.1892; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Scirpus corymbosus L., Cent. Pl. 2: 7.1756

Rhynchospora aurea Vahl, Enum. Pl. Obs. 2: 229.1805

Schoenus corymbosus (L.) Pers., Syn. Pl. 1: 59.1805

Schoenus articulatus Roxb., Fl. Ind. 1: 184.1820

Rhynchospora articulata (Roxb.) Schult., Mant. 2: 49.1824

Scirpus bangalorensis B.Heyne ex Wall., Numer. List 3464.1831

Cephaloschoenus articulatus (Roxb.) Nees, Edinburgh New Philos. J. 17: 266.1834

Scirpus umbellatus Roxb. ex Kunth, Enum. Pl. 2: 294. 1837.

Calyptrostylis articulata (Roxb.) Nees, J. Bot. (Hooker) 2: 394.1840

Dichromena corymbosa (L.) J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 11: 6.1931

Distribution: Karnataka (Belagavi, Bengaluru, Hassan, Mandya, Mysuru, Shivamogga, Uttara Kannada), throughout India; Pantropical.

Rhynchospora rugosa (Vahl) Gale, Rhodora 46: 275.1944; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Schoenus rugosus Vahl, Eclog. Amer. 2: 5. 1798

Rhynchospora glauca Vahl, Enum. Pl. Obs. 2: 233.1805

Distribution: Karnataka (?), Peninsula to NE India;tropical and subtropical areas

Rhynchospora wightiana (Nees) Steud., Syn. Pl. Glumac. 2: 148.1855; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Haplostylis wightianus Nees, Nov. Actorum Acad. Caes. Leop.-Carol. Nat. Cur. 19(Suppl. 1): 101.1843

Rhynchospora discolor Hochst. ex Steud., Syn. Pl. Glumac. 2: 148.1855

Distribution: Karnataka (Belagavi, Ballari, Chamarajanagar, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), W Peninsular India. S E Asia.

Schoenoplectiella Lye

Schoenoplectiella articulata (L.) Lye, Lidia 6: 20.2003

Scirpus articulatus L., Sp. Pl. 47.1753

Scirpus incurvatus Roxb., Fl. Ind. 1: 216.1820

Scirpus subarticulatus Roxb., Fl. Ind. 1: 215.1820

Eleocharis incurvata (Roxb.) Schult., Mant. 2: 92.1824

Holoschoenus incurvatus (Roxb.) A.Dietr., Sp. Pl. 2: 165.1832

Holoschoenus subarticulatus (Roxb.) A.Dietr., Sp. Pl. 2: 166.1832

Isolepis incurvata (Roxb.) Nees, Contr. Bot. India 108.1834

Schoenoplectus articulatus (L.) Palla, Bot. Jahrb. Syst. 10: 299.1888; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Distribution: Karnataka (throughout), throughout india; Paleotropical & Mediterranean.

Schoenoplectiella juncoides (Roxb.) Lye, Lidia 6: 25.2003

Scirpus juncoides Roxb., Fl. Ind. 1: 218.1820

Schoenoplectus juncoides (Roxb.) Palla, Bot. Jahrb. Syst. 10: 299.1888; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Scirpus erectus C.B.Clarke, Fl. Brit. India 6: 656.1893

Hymenocheata juncoides (Roxb.) Nakai, Bull. Natl. Sci. Mus. Tokyo 31: 133.1952

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; Australia, madagascar, E & SE Asia,

Schoenoplectiella lateriflora (J.F.Gmel.) Lye, Lidia 6: 25.2003

Scirpus lateriflorus J.F.Gmel., Syst. Nat. 2: 127.1791

Scirpus supinus var. *lateriflorus* (J.F.Gmel.) T.Koyama, J. Fac. Sci. Univ. Tokyo, Sect. 3, Bot. 7: 302.1958

Schoenoplectus lateriflorus (J.F.Gmel.) Lye, Bot. Not. 124: 290.1971; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Schoenoplectus supinus subsp. *lateriflorus* (J.F.Gmel.) Soják, Cas. Nár. Mus., Odd. Prír. 141: 62.1972

Distribution: Karnataka (throughout), throughout except NW India; China, S & SE Asia, Australia.

Schoenoplectiella senegalensis (Steud.) Lye, Lidia 6: 27.2003

Isolepis senegalensis Steud., Syn. Pl. Glumac. 2: 96.1855

Schoenoplectus senegalensis (Steud.) Palla, Bot. Jahrb. Syst. 10: 299.1888

Scirpus jacobi C.E.C.Fisch., Bull. Misc. Inform. Kew 1931: 103.1931

Schoenoplectus jacobi (C.E.C.Fisch.) Lye, Bot. Not. 124: 290.1971

Schoenoplectus senegalensis (Hochst. ex Steud.) Palla ex J.Raynal, Cat. Pl. Vasc. Niger 344. 1976; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Distribution: Karnataka (Bengaluru, Hassan, Mysuru, Tumakuru), Peninsular India; Africa.

Schoenoplectiella supina (L.) Lye, Lidia 6: 27.2003

Scirpus supinus L., Sp. Pl. 49.1753

Distribution: Karnataka (), throughout India; Africa, Afghanistan, Australia, Europe, China, Indonesia, Kazakhstan, Madagascar, Malaysia, Myanmar, Nepal, Pakistan, Philippines, Russia, Sri Lanka, S. America, SW Asia, Thailand, Uzbekistan, Vietnam.

Schoenoplectus (Rchb.) Palla,

Schoenoplectus corymbosus (Roth ex Roem. & Schult.) J.Raynal, Cat. Pl. Vasc. Niger 343.1976; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Isolepis corymbosa Roth ex Roem. & Schult., Syst. Veg. 2: 110.1817

Scirpus corymbosus (Roth ex Roem. & Schult.) B.Heyne, Nov. Pl. Sp. 28. 1821.

Malacochaete corymbosa (Roth ex Roem. & Schult.) Arn., Cat. Ind. Pl. 143.1837

Distribution: Karnataka (Bengaluru, Hassan, Mysuru), throughout Indian plains; Paleotropical.

Schoenoplectus litoralis (Schrad.) Palla, Bot. Jahrb. Syst. 10: 299. 1888.

subsp. **thermalis** (Trab.) S.S.Hooper, Fl. Hassan Distr. 698.1976; B.D.Sharma et al., Fl. Karnataka Anal. 315. 1984.

Scirpus subulatus Vahl, Enum. Pl. 2: 268.1805

Scirpus pectinatus Roxb., Fl. Ind. 1: 220.1820

Malacochaete pectinata (Roxb.) Nees, Linnaea 9: 292.1834

Scirpus thermalis Trab., Fl. Algérie Monocot.: 99.1895

Scirpus wardianus J.R.Drumm., J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 73: 147.1904

Scirpus litoralis subsp. *thermalis* (Trab.) Murb., Acta Univ. Lund. n.s., 2(18) 3: 18.1922

Scirpus litoralis var. *subulatus* (Vahl) Chiov., Pl. Nov. Aethiop. 16.1928

Schoenoplectus litoralis subsp. *subulatus* (Vahl) Soják, Cas. Nár. Mus., Odd. Prír. 141: 62.1972

Scirpus litoralis subsp. *subulatus* (Vahl) M.A.Wadood Khan, Rheedia 10: 28.2000

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina kannada, Dharwad, Kalburgi, Hassan, Mysuru, Raichur, Uttara Kannada, Vijayapura), throughout India; Madagascar, Micronesia, S & SE Asia, tropical & S Africa.

Schoenoplectus mucronatus (L.) Palla, Bot. Jahrb. Syst. 10: 299.1888; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Scirpus mucronatus L., Sp. Pl. 50.1753

Isolepis mucronata (L.) Fourr., Ann. Soc. Linn. Lyon n.s., 17: 173.1869

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; paleotropical.

Local/Common/Vernacular name: *Hommagali hullu* (Kannada)

Scleria Berg.**Scleria annularis** Steud., Syn. Pl. Glumac. 2: 176.1855; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Hypoporum annulare Nees, Linnaea 9: 303.1834

Distribution: Karnataka (Uttara Kannada), throughout except N India & Himalayas; China, SE Asia.

Scleria biflora Roxb., Fl. Ind. ed.2, 3: 573.1832

Scleria ferruginea Ohwi, Acta Phytotax. Geobot. 7: 37.1938

Scleria biflora subsp. *ferruginea* J.Kern, Reinwardtia 6: 76.1961

Distribution: Karnataka (Dakshina Kannada, Hassan, Udupi, Uttara Kannada), W Peninsular to E India; China, Sri Lanka, S E Asia.

Scleria corymbosa Roxb., Fl. Ind. ed. 2, 3: 574.1832; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Scleria major Moon, Cat. Pl. Ceylon 62.1824

Scleria androgyna Nees, Contr. Bot. India 117.1834

Scleria corymbifera Boeckeler, Linnaea 38: 537.1874

Scleria ridleyi C.B.Clarke in Hook.f., Fl. Brit. India 6: 686.1894

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), throughout except Nw India; China, Sri Lanka & SE Asia.

Scleria foliosa Hochst. ex A.Rich., Tent. Fl. Abyss. 2: 509.1850; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Distribution: Karnataka (Hassan), S India, Madagascar, tropical & S Africa.

Scleria levis Retz., Observ. Bot. 4: 13.1786; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Carex laevis (Retz.) J.F.Gmel., Syst. Nat. 2: 138.1791

Scleria hebecarpa Nees, Contr. Bot. India 117.1834

Scleria pubescens Steud., Syn. Pl. Glumac. 2: 168.1855

Scleria hebecarpa var. *pubescens* (Steud.) C.B.Clarke in Hook.f., Fl. Brit. India 6: 689.1894

Scleria levis var. *pubescens* (Steud.) C.Z.Zheng, Fl. Zhejiang 7: 286.1993

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India except ; E & SE Asia, tropical Australia

Scleria lithosperma (L.) Sw., Prodr. Veg. Ind. Occ. 18.1788; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Scirpus lithospermus L., Sp. Pl. 51.1753

Hypoporum lithospermum (L.) Nees, Contr. Bot. India 117.1834

Hypoporum purpurascens Nees, Linnaea 9: 303.1834

Scleria wightiana Steud., Syn. Pl. Glumac. 2: 176.1855

var. **lithosperma**

Distribution: Karnataka (throughout), throughout India except desert areas; Pantropical.

var. **linearis** Benth., Fl. Austral. 7: 430. 1878; Dey & Prasanna, Indian J. For. 32 (3): 511. 2009.

Distribution: Karnataka (Bengaluru), Andhra Pradesh, Bihar, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; Malaya, New Guinea, Phillipines, Queensland, Sri Lanka, Thailand, Vietnam.

Scleria multilacunosa T.Koyama, Bull. Natl. Sci. Mus. Tokyo n.s., 17: 71.1974; Manikand.& Lakshminars., Fl. Rajiv Gandhi Nation. Park 446.2013.

Distribution: Karnataka (Belagavi, Mysuru)

Scleria pergracilis (Nees) Kunth, Enum. Pl. 2: 354.1837; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Hypoporum pergracile Nees, Edinburgh New Philos. J. 1834: 267.1834

Scleria salebrosa Boeckeler ex C.B.Clarke in T.A. Durand & H. Schinz, Consp. Fl. Afric. 5: 673.1894

Distribution: Karnataka (Hassan), throughout except N W india; Sri Lanka, SE Asia & Australia.

Scleria poiformis Retz., Observ. Bot. 4: 13.1786; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984 ('poaeformis').

Carex poiformis (Retz.) J.F.Gmel., Syst. Nat. 2: 138.1791

Distribution: Karnataka (Uttara Kannada), S to NE India; Sri Lanks, E & SE Asia, tropical Africa & Australia.

Scleria psilorrhiza C.B.Clarke, Fl. Brit. India 6: 691. 1894; Dey & Prasanna, Indian J. For. 32 (3): 511. 2009.

Distribution: Karnataka (Uttara Kannada), Assam, Himachal Pradesh, Madhya Pradesh, Maharastra, Odisha; Bangladesh, Cambodia, China South-Central, India, Jawa, New Guinea, Northern Territory, Philippines, Queensland, Thailand, Western Australia.

Scleria stocksiana Boeckeler, Linnaea 38: 474.1874; B.D.Sharma et al., Fl. Karnataka Anal. 316. 1984.

Distribution: Karnataka (Belagavi, Dharwad, Uttara Kannada), Peninsular India.

- Scleria terrestris** (L.) Fassett, Rhodora 26: 159.1924
Zizania terrestris L., Sp. Pl. 2: 991.1753
Schoenus paniculatus Burm.f., Fl. Indica 19.1768
Diaphora cochinchinensis Lour., Fl. Cochinch. 578.1790
Scleria elata Thwaites, Enum. Pl. Zeyl. 353.1864
Scleria melanostoma Boeckeler, Linnaea 38: 514.1874
Scleria chinensis var. *biauriculata* C.B.Clarke in Hook.f., Fl. Brit. India 6: 690.1894
Scleria elata var. *decolorans* C.B.Clarke in Hook.f., Fl. Brit. India 6: 690.1894
Scleria elata var. *latrix* C.B.Clarke in Hook.f., Fl. Brit. India 6: 690.1894
Scleria scrobiculata C.B.Clarke, J. Linn. Soc., Bot. 36: 266. 1903
Scleria cochinchinensis (Lour.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 646.1917
Scleria terrestris var. *decolorans* (C.B.Clarke) Fassett, Rhodora 26: 160.1924
Scleria terrestris var. *latrix* (C.B.Clarke) Fassett, Rhodora 26: 160.1924
 Distribution: Karnataka (Belagavi, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India; Australia, E & SE Asia.

- Scleria tessellata** Willd., Sp. Pl. 4: 315.1805; V.P.Prasad & N.P.Singh, Sedges Karnataka 327. 2002..
Carex indica J.Koenig ex Willd., Sp. Pl. 4: 315. 1805.
Scleria glandiformis Boeckeler, Flora 41: 647.1858
 Distribution: Karnataka (Belagavi, Mysuru), throughout India except arid deserts; madagascar, tropical Africa.

Daphniphyllaceae
Daphniphyllum Blume

- Daphniphyllum neilgherrense** (Wight) K.Rosenthal, Pflanzenr. IV, 147a: 7.1919
Goughia neilgherrensis Wight, Icon. Pl. Ind. Orient. 5: t. 1878.1852
Daphniphyllum glaucescens var. *concolor* Müll.Arg., Prodr. 16(1): 4.1869
Daphniphyllum glaucescens subsp. *neilgherrense* (Wight) T.C.Huang, Taiwania 12: 195.1966
 Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Dichapetalaceae
Dichapetalum Thouars

- Dichapetalum gelonioides** (Roxb.) Engl. in Engl. & Prantl, Naturl. Pflanzenfam. 3(4): 348. 1896;
 B.D.Sharma et al., Fl. Karnataka Anal. 47. 1984.
Moacurra gelonioides Roxb., Fl. Ind. 2: 69. 1832
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga), Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Kerala, Maharashtra, Tamil Nadu; Indonesia, Sri Lanka.
 Local/Common/Vernacular name: *Kaadu tengu* (Kannada)

Dilleniaceae
Dillenia L.

- Dillenia bracteata** Wight, Icon. Pl. Ind. Orient. 2: t. 358. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 2. 1984.
Wormia bracteata (Wight) Hook.f. & Thomson, Fl. Ind. 68. 1855
 Distribution: Karnataka (Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.
 Local/Common/Vernacular names: *Bettada kanigilu, kanigulu* (Kannada)

- Dillenia indica** L., Sp. Pl. 535. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 2. 1984.
Dillenia speciosa Thunb., Trans. Linn. Soc. London 1: 200. 1791
 Distribution: Karnataka (Belagavi, Bengaluru, Kodagu, Shivamogga), throughout India; Bhutan, China, Indonesia, Laos, Malaysia, Myanmar, Nepal, Philippines, Sri Lanka, Thailand, Vietnam.

Dillenia pentagyna Roxb., Pl. Coromandel 1: t. 20. 1795; B.D.Sharma et al., Fl. Karnataka Anal. 2. 1984.

Dillenia pilosa Roxb., Fl. Ind. 2: 652. 1832

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Shivamogga, Uttara Kannada), throughout India; Bhutan, China, Indonesia, Malaysia, Myanmar, Nepal, Thailand, Vietnam.

Local/Common/Vernacular name: *Maratega* (Kannada)

Dillenia retusa Thunb., Trans. Linn. Soc. London 1: 200. 1791; B.D.Sharma et al., Fl. Karnataka Anal. 2. 1984.

Wormia retusa (Thunb.) Hook.f. & Thomson, Fl. Ind. 68. 1855

Distribution: Karnataka (Mysuru ?), Kerala, Tamil Nadu; Sri Lanka.

Dioscoreaceae

Dioscorea L.

Dioscorea alata L., Sp. Pl. 1033.1753; B.D.Sharma et al., Fl. Karnataka Anal. 283. 1984.

Dioscorea globosa Roxb., Fl. Ind. ed.2, 3: 797.1832

Dioscorea purpurea Roxb., Fl. Ind. ed.2, 3: 799.1832

Dioscorea rubella Roxb., Fl. Ind. ed.2. 3: 798.1832

Polynome alata (L.) Salisb., Gen. Pl. 12. 1866

Dioscorea alata var. *globosa* (Roxb.) Prain, Bengal Pl. 2: 1067.1903

Distribution: Karnataka (Widely cultivated), native to SE Asia.

Local/Common/Vernacular names: *Ivana genasu*, *Mundi genasu*, *Tuna genasu* (Kannada)

Dioscorea belophylla (Prain) Voigt ex Haines, Forest Fl. Chota Nagpur 530.1910; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Dioscorea sagittata Royle, Ill. Bot. Himal. Mts. 378.1839

Dioscorea nummularia var. *belophylla* Prain, Bengal Pl. 2: 1065.1903

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Mysuru, Udupi, Uttara Kannada), throughout except dry NW India; endemic.

Dioscorea bulbifera L., Sp. Pl. 1033.1753; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Dioscorea crispata Roxb., Fl. Ind. ed.2, 3: 802.1832

Dioscorea heterophylla Roxb., Fl. Ind. ed.2, 3: 804.1832

Dioscorea pulchella Roxb., Fl. Ind. ed.2, 3: 801.1832

Helmia bulbifera (L.) Kunth, Enum. Pl. 5: 435.1850

Polynome bulbifera (L.) Salisb., Gen. Pl. 12.1866

Dioscorea bulbifera var. *crispata* (Roxb.) Prain, Bengal Pl. 2: 1066.1903

Dioscorea bulbifera var. *pulchella* (Roxb.) Prain, Bengal Pl. 2: 1066.1903

Dioscorea bulbifera var. *sativa* Prain, Bengal Pl. 2: 1066.1903

Distribution: Karnataka (throughout), throughot India except in deserts & arid areas; Sri Lanka.

Local/Common/Vernacular names: *Handi gedde*, *Heggenasu*, *Kunta genasu* (Kannada); *Karande* (*Kunabi*); *Varahi kanda* (Sanskrit)

Dioscorea esculenta (Lour.) Burkill, Gard. Bull. Straits Settlem. 1: 396.1917

Oncus esculentus Lour., Fl. Cochinch. 194.1790

Oncorhiza esculentus (Lour.) Pers., Syn. Pl. 1: 374.1805

Dioscorea aculeata Roxb., Fl. Ind. ed. 2, 3: 800.1832

Dioscorea fasciculata Roxb., Fl. Ind. ed. 2, 3: 801.1832

Dioscorea spinosa Roxb. ex Hook.f., Fl. Brit. India 6: 291.1892

Dioscorea fasciculata var. *spinosa* Prain, Bengal Pl. 2: 1066.1903

Dioscorea aculeata var. *spinosa* (Prain) Roxb. ex Prain & Burkill, J. Proc. Asiat. Soc. Bengal 10: 20.1914

Dioscorea esculenta var. *fasciculata* (Roxb.) Prain & Burkill, J. Proc. Asiat. Soc. Bengal 10: 19.1914

Distribution: Karnataka (Dakshina Kannada, Udupi)

Dioscorea glabra Roxb., Fl. Ind. ed. 1832 3: 804.1832; Manikand.& Lakshminars., Fl. Rajiv Gandhi Nation. Park 400.2013.

Dioscorea nummularia Roxb., Fl. Ind. ed. 1832 3: 804.1832

Dioscorea glabra var. *longifolia* Prain & Burkill, J. Proc. Asiat. Soc. Bengal 10: 37.1914

Dioscorea oryzetorum var. *latifolia* Prain & Burkill, Bull. Misc. Inform. Kew 1927: 243.1927

Dioscorea oryzetorum var. *mediifolia* Prain & Burkill, Bull. Misc. Inform. Kew 1927: 243 1927

Dioscorea glabra var. *vera* Prain & Burkill, Ann. Roy. Bot. Gard. (Calcutta) 14: 357.1936

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Davanagere) Peninsular India; S & SE Asia.

Dioscorea hamiltonii Hook.f., Fl. Brit. India 6: 294.1892; Manikand.& Lakshminars., Fl. Rajiv Gandhi Nation. Park 400.2013.

Dioscorea persimilis Prain & Burkill, J. Proc. Asiat. Soc. Bengal 4: 454.1908

Distribution: Karnataka (Kodagu, Mysuru), Assam, Sikkim, Bhutan, China, Myanmar, Nepal, Taiwan,, Thailand, Vietnam.

Dioscorea hispida Dennst., Schlüssel Hortus Malab. 33.1818; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Dioscorea mollissima Blume, Enum. Pl. Javae 1: 21.1827

Dioscorea daemonia Roxb., Fl. Ind. ed.2, 3: 805.1832

Helmia daemonia (Roxb.) Kunth, Enum. Pl. 5: 439.1850

Helmia hirsuta (Blume) Kunth, Enum. Pl. 5: 438.1850

Dioscorea daemonia var. *reticulata* Hook.f., Fl. Brit. India 6: 289.1892

Dioscorea triphylla var. *daemonia* (Roxb.) Prain & Burkill, J. Proc. Asiat. Soc. Bengal 10(1): 26.1914

Dioscorea triphylla var. *mollissima* (Blume) Prain & Burkill, J. Proc. Asiat. Soc. Bengal 10(1): 26.1914

Dioscorea triphylla var. *reticulata* (Hook.f.) Prain & Burkill, J. Proc. Asiat. Soc. Bengal 10: 26.1914

Dioscorea hispida var. *daemonia* (Roxb.) Prain & Burkill, Ann. Roy. Bot. Gard. (Calcutta) 14: 192.1936

Dioscorea hispida var. *reticulata* (Hook.f.) Sanjappa, Fl. Ind. Enumerat. - Monocot. 74 1989

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), from Peninsular to NE India, S & SE Asia.

Dioscorea intermedia Thwaites, Enum. Pl. Zeyl. 326.1864; Seetharam, Kotresh & Uplaonkar, Fl. Dist 57. 2000.

Distribution: Karnataka (Kalburgi, Yadgir), Sri Lanka.

Dioscorea oppositifolia L., Sp. Pl. 1033.1753; B.D.Sharma et al., Fl. Karnataka Anal. 278. 1984.

Distribution: Karnataka (throughout), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Bellarayi*, *Hakki genasu*, *Kadu genasu* (Kannada); *Sarpkhyā* (Sanskrit)

Dioscorea pentaphylla L., Sp. Pl. 1032.1753; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Dioscorea triphylla L., Sp. Pl. 1032.1753

Botryosicyos pentaphyllum (L.) Hochst., Flora 27(Bes. Beil.): 3.1844

Dioscorea pentaphylla var. *communis* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10(1): 23.1914

Dioscorea pentaphylla var. *linnaei* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10(1): 23.1914

Dioscorea pentaphylla var. *rheedei* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10(1): 23.1914

Distribution: Karnataka (throughout), from Peninsular to NE India; S & SE Asia, tropical Africa.

Local/Common/Vernacular name: *Kaadu gumbala* (Kannada)

Dioscorea pubera Blume, Enum. Pl. Javae 1: 21.1827

Dioscorea anguina Roxb., Fl. Ind. ed.2, 3: 803.1832

Distribution: Karnataka (Dakshina Kannada, Udupi)

Dioscorea spicata Roth, Nov. Pl. Sp. 371. 1821; Bhaskar & Kushalappa, Fl. Tumkur Dist. 105. 2013.

Dioscorea nummularia var. *glaucia* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10: 35. 1914

Dioscorea spicata var. *anamallayana* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10: 29. 1914

Dioscorea spicata var. *parvifolia* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10: 29. 1914

Distribution: Karnataka (Tumakuru)

Dioscorea tomentosa J.König ex Spreng., Pl. Min. Cogn. Pug. 2: 92.1815; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Helmia tomentosa (J.Koenig) ex Spreng.) Kunth, Enum. Pl. 5: 439.1850

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chitradurga, Dakshina Kannada, Davanagere, Kalburgi, Kodagu, Shivamogga, Tumakuru, Udupi, Yadgir)

Local/Common/Vernacular names: *Bili genasu, Mullu genasu* (Kannada)

Dioscorea villosa L., Sp. Pl. 1033.1753; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Dioscorea sativa L., Sp. Pl. 1033.1753

Merione villosa (L.) Salisb., Gen. Pl. 12.1866

Distribution: Karnataka (Dharwad)

Dioscorea wallichii Hook.f., Fl. Brit. India 6: 295.1892; B.D.Sharma et al., Fl. Karnataka Anal. 284. 1984.

Dioscorea wallichii var. *christiei* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10: 31.1914

Dioscorea wallichii var. *vera* Prain & Burkhill, J. Proc. Asiat. Soc. Bengal 10: 31.1914

Distribution: Karnataka (Chikkamagalur, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), S to NE India; Myanmar.

Tacca J. R. Forst & G. Forst.

Tacca leontopetaloides (L.) Kuntze, Revis. Gen. Pl. 2: 704.1891; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Leontice leontopetaloides L., Sp. Pl. 313. 1753

Distribution: Karnataka (Belagavi, Kodagu, Mysuru, Uttara Kannada), Peninsular to NE India & Andaman & Nicobar Islands; S & SE Asia.

Dipterocarpaceae

Dipterocarpus C.F.Gaertn .

Dipterocarpus bourdillonii Brandis in Hook., Ic. Pl. t. 25. 1895; Ramesh & Pascal, J. Bombay Nat. Hist. Soc., 90: 324. 1993

Distribution: Karnataka (Kodagu), Kerala; endemic.

Dipterocarpus indicus Bedd., Conserv. For. Rep. 17 1864; B.D.Sharma et al., Fl. Karnataka Anal. 19. 1984.

Distribution: Karnataka (Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Hopea Roxb.

Hopea canarensis Hole, Indian Forester 44: 575 1918; B.D.Sharma et al., Fl. Karnataka Anal. 19. 1984.
 Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga); endemic.

Hopea erosa (Bedd.) Slooten, Reinwardtia 3: 318.1956
Balanocarpus erosus Bedd., Fl. Sylv. S. India Manual: 237. 1872
 Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Hopea glabra Wight & Arn., Prodr. Fl. Ind. Orient. 85 1834; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.
 Distribution: Karnataka (Dakshina Kannada), Kerala, Tamil Nadu; endemic.

Hopea jacobi C.E.C.Fisch., Bull. Misc. Inform. Kew 1932: 245 1932; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.
 Distribution: Karnataka (Kodagu), Kerala; endemic.

Hopea odorata Roxb., Pl. Coromandel 3: 7.1811
Hopea wightiana Wall. ex Wight & Arn., Prodr. Fl. Ind. Orient. 1: 85. 1834
Hopea decandra Buch.-Ham. ex Wight, Ill. Ind. Bot. 1: 88.1840
 Distribution: Karnataka (cultivated?), Andaman & Nicobar Islands; Bangladesh, Cambodia, Laos, Malaysia, Myanmar, Thailand, Vietnam.
 Local/Common/Vernacular name: *white thingan* (English)

Hopea parviflora Bedd., Fl. Sylv. S. India t. 7 1869; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984 **Plate 22A**

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.
 Local/Common/Vernacular names: *Hopea*, *Iron wood of Malabar*; *White Kongu* (English); *Bovu mara*, *Kiralbhogi*, *Sannele bogi* (Kannada)

Hopea ponga (Dennst.) Mabb., Taxon 28: 587 1979; C.J.Saldanha, Fl. Karnataka 1: 194. 1984 **Plate 22B**
Artocarpus ponga Dennst., Schlüssel Hortus Malab. 30 1818

var. **ponga**
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.
 Local/Common/Vernacular name: *Haiga* (Kannada)

var. **cauveriana** Kesh. Murthy & Yogan., Curr. Sci. 56 (11): 544. 1987; Kesh.Murthy & Yogan, Fl. Kodagu 62. 1990
 Distribution: Karnataka (Kodagu); endemic.

Hopea racophloea Dyer, Fl. Brit. India 1: 310 1874; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.
Hopea malabarica Bedd., Icon. Pl. Ind. Or. 1: 42 1874
 Distribution: Karnataka (Dakshina Kannada, Shivamogga), Kerala, Tamil Nadu; endemic.

Shorea Roxb. ex C. F. Gaertn.

Shorea roxburghii G.Don, Gen. Hist. 1: 813 1831; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.
Shorea laccifera B.Heyne ex Wall., Numer. List 967 1829
Shorea talura Roxb., Fl. Ind. ed. 1832 2: 618 1832
Vatica laccifera Wight & Arn., Prodr. Fl. Ind. Orient. 84 1834
 Distribution: Karnataka (Bengaluru, Davanagere, Hassan, Kodagu, Mandya, Mysuru, Tumakuru, Uttara Kannada), Andhra Pradesh, Chattisgarh, Kerala, Tamil Nadu; Cambodia, Laos, Malaysia, Myanmar, Thailand, Vietnam.
 Local/Common/Vernacular name: *Jalari* (Kannada)

Shorea tumbuggaia Roxb., Fl. Ind. ed. 1832 2: 617 1832; B.D.Sharma et al., Fl. Karnataka Anal. 2. 1984.
 Distribution: Karnataka (Kolar? Mysuru), Andhra Pradesh, Tamil Nadu; endemic.

Vateria L.

Vateria indica L., Sp. Pl. 515 1753; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.

Plate 22C

Vateria malabarica Blume in Mus. Bot. 2: 29 1856

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Dhoopada mara* (Kannada)

Vatica L.

Vatica chinensis L., Mant. Pl. 2: 242 1771; B.D.Sharma et al., Fl. Karnataka Anal. 20. 1984.

Vatica roxburghiana Blume in Mus. Bot. 2: 31 1856

Distribution: Karnataka (Dakshina Kannada, Mysuru), Kerala, Tamil Nadu; endemic.

Droseraceae

Drosera L.

Drosera burmanni Vahl, Symb. Bot. 3: 50. 1794; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Distribution: Karnataka (throughout), Andhra Pradesh, Assam, Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Mizoram, Odisha, Tamil Nadu; West Africa to North East Australia

Local/Common/Vernacular name: *Hula hiduka gida* (Kannada)

Drosera indica L., Sp. Pl. 282. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 99. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Vijayapura), Andhra Pradesh, Bihar, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; tropical Africa to Australia

Drosera peltata Thunb., Drosera 7 1797; Yogan et al., Fl. Chikmagalur 136. 1981

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Mysuru); India to Australia.

Ebenaceae

Diospyros L.

Diospyros affinis Thwaites, Enum. Pl. Zeyl. 179. 1860; Ramesh & Pascal, J. Bombay Nat. Hist. Soc., 90: 324. 1993

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Kunchigana mara* (Kannada)

Diospyros bourdillonii Brandis, Indian Trees 435. 1906; Ramesh & Pascal, J. Bombay Nat. Hist. Soc., 90: 324. 1993

Distribution: Karnataka (Dakshina Kannada, Kodagu), Kerala, Tamil Nadu; endemic.

Diospyros buxifolia (Blume) Hiern in Trans. Cambridge Philos. Soc. 12: 218. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 155. 1984.

Leucoxylum buxifolium Blume, Bijdr. 1169. 1826

Diospyros microphylla Bedd., Fl. Sylv. S. India 145 1872

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Indo-China, New Guinea, Madagascar, SE Asia.

Local/Common/Vernacular name: *Kunchigana mara* (Kannada)

Diospyros candolleana Wight, Icon. Pl. Ind. Orient. 4: t. 1221. 1848; B.D.Sharma et al., Fl. Karnataka Anal. 155. 1984.

Diospyros canarica Bedd., Icon. Pl. Ind. Or. t.134. 1874

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Kare mara, Kari* (Kannada)

Diospyros chloroxylon Roxb., Pl. Coromandel 1: 38. t. 49. 1795; B.D.Sharma et al., Fl. Karnataka Anal. 155. 1984.

Diospyros capitulata Wight, Icon. Pl. Ind. Orient. 4: T. 1224. 1848

Diospyros chloroxylon var. *cupulosa* V.Singh, Monogr. Ind. *Diospyros* 73. 2005

Distribution: Karnataka (Ballari, Bidar, Chitradurga, Kalaburagi, Kolar, Uttara Kannada, Yadgir), Andhra Pradesh, Chattisgarh, Gujarat, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, West Bengal; Bangladesh, Myanmar, Sri Lanka.

Local/Common/Vernacular names: *Coromandel ebony, Green ebony perismon Ninai* (English); *Handjee, Kadu gariathu mara, Nensi* (Kannada)

Diospyros crumenata Thwaites, Enum. Pl. Zeyl. 179. 1860; B.D.Sharma et al., Fl. Karnataka Anal. 155. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Uttara Kannada), Andaman & Nicobar Islands, Goa, Kerala, Maharashtra, Tamil Nadu; Myanmar, Sri Lanka.

Local/Common/Vernacular names: *Kantumri, Karitumari, Thumri* (Kannada)

Diospyros ebenum J.Koenig ex Retz., Physiogr. Salsk. Mag. 1: 176. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 155. 1984.

Diospyros glaberrima (L.) Rottler, Nye Saml. Kongel. Vidensk. Danske Selsk. Skr. 2: 540.1783

Diospyros ebenaster Retz., Observ. Bot. 5: 31.1788

Diospyros melanoxylon Willd., Sp. Pl. 4: 1109.1806 non Roxb. 1795

Diospyros membranacea A.DC., Prodr. 8: 227.1844

Diospyros assimilis Bedd., Rep. Forest. Madr. 20 1866

Diospyros ebenum var. *glaberrima* (L.f.) Bakh.f., Bull. Jard. Bot.Buitz. Ser.3, 15(3):216.1938

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Goa, Kerala, Maharashtra, Odisha, Tamil Nadu; S & SE Asia, New Caledonia, New Guinea, Australia C America.

Local/Common/Vernacular names: *Bale, Balemara, Kari, Karikhanamara, Karimara, Mallai* (Kannada)

Diospyros exsculpta Buch.-Ham., Trans. Linn. Soc. London 15: 110 1826 non Dalzell 1861

Diospyros tomentosa Roxb., Fl. Ind. ed. 1832, 2: 532 1832 non Poir. 1804; B.D.Sharma et al., Fl. Karnataka Anal. 157. 1984.

Distribution: Karnataka (Uttara Kannada), Andhra Pradesh, Maharashtra, Tamil Nadu, West Bengal.

Diospyros foliosa (Rich ex A.Gray) Bakh., Bull. Jard. Bot. Buitenzorg 3, 15: 447 1941; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Maba foliosa Rich ex A.Gray, Proc. Amer. Acad. Arts 5: 326.1862

Ebenus foliosa (Rich ex A.Gray) Kuntze, Revis. Gen. Pl. 2: 408.1891

Diospyros elliptica var. *foliosa* (Rich ex A.Gray) A.C.Sm., J. Arnold Arbor. 52: 377.1971

Distribution: Karnataka (Shivamogga), Kerala, Tamil Nadu; Pacific Islands?

Diospyros ghatensis B.R.Ramesh & D.De Franceschi, Blumea 38: 131. 1993

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Diospyros insignis Thwaites, Enum. Pl. Zeyl. 180.1860

Diospyros insignis var. *parvifolia* Kosterm., Revised Handb. Fl. Ceylon 3: 45.1981

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Diospyros lanceifolia Roxb., Fl. Ind. ed. 1832 2: 537 1832

Diospyros amoena Wall. ex G.Don, Gen. Hist. 4: 40.1837

Diospyros grata Wall. ex A.DC., Prodr. 8: 232.1844

Diospyros lucida Wall. ex A.DC., Prodr. 8: 233.1844

Diospyros multiflora Wall. ex A.DC., Prodr. 8: 231.1844

Diospyros hirsuta Hiern, Trans. Cambridge Philos. Soc. 12: 164. 1873; C.J.Saldanha, Fl. Karnataka 1: 336. 1984.

Diospyros pachyphylla C.B.Clarke in Hook. f., Fl. Brit. India 3: 568.1882

Diospyros clavigera var. *pachyphylla* (C.B.Clarke) Ridl., Fl. Malay Penins. 2: 288.1923

Diospyros lonchophylla Hiern, J. Bot. 63(Suppl.): 63.1925

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga, Uttara Kannada), E & NE India; S & SE Asia.

Diospyros malabarica (Desr.) Kostel., Allg. Med.-Pharm. Fl. 3: 109. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Embryopteris peregrina Gaertn., Fruct. Sem. Pl. 1: 145.1788

Embryopteris glutinifera Roxb., Pl. Coromandel 1: 49.1796

Diospyros embryopteris Pers., Syn. Pl. 2: 624. 1807

Diospyros glutinifera (Roxb.) Wall., Numer. List 4123 B.1831

Diospyros glutinosa J.Koenig ex Roxb., Fl. Ind. ed. 1832 2: 533.1832

Diospyros citrifolia Wall. ex A.DC., Prodr. 8: 285.1844

Diospyros peregrina (Gaertn.) Gurke in Nat. Pflanzenfam. 4(1): 164 1891

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Mandya, Mysuru, Shivamogga, Uttara Kannada), Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; India, Sri Lanka.

Local/Common/Vernacular names: *Gaub tree, Indian Perismmon, River ebony, Wild mangosteen* (English); *Holeupare* (Kannada)

Diospyros melanoxylon Roxb., Pl.Coromandel 1:t.46.1795; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Diospyros dubia Wall. ex A.DC., Prodr. 8: 223 1844

Diospyros roylei Wall. ex A.DC., Prodr. 8: 239.1844

Diospyros exsculpta Bedd., Fl. Sylv. S. India t. 66. 1870.

Diospyros wightiana Wall. ex Bedd, Fl. Syl. T. 67. 1870

var. **melanoxyton**

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburagi, Kodagu, Kolar, Mandya, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Bihar, Chattisgarh, Goa, Gujarat, Jharkhand, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttaranchal, Uttar Pradesh, West Bengal; Sri Lanka.

Local/Common/Vernacular names: *Abanasi tumari, Balai, Beedi yele, Kendu, Makartendu, Temru, Thumbri, Toomri, Tupra* (Kannada)

var. **tupru** (Buch.-Ham.) V.Singh in Monogr. Ind. Diospyros 159 2005

Diospyros rubiginosa Roth, Nov. Pl. Sp. 385.1821

Diospyros tupru Buch.-Ham. in Trans. Linn. Soc. London 5:111. 1827; C.J.Saldanha, Fl. Karnataka 1: 339. 1984.

Diospyros montana B.Heyne ex A.DC., Prodr. 8: 239.1844

Diospyros exsculpta Dalzell & A.Gibson, Bombay Fl. 142. 1861.

Distribution: Karnataka (Chitradurga, Davanagere, Kodagu); throughout except NW & NE India; Afghanistan, Sri Lanka.

Local/Common/Vernacular names: *Bandada mara, Kakatinduka, Timburni, Tindura, Tuparanda, Tupru* (Kannada)

Diospyros montana Roxb., Pl. Coromandel 1: 37. 1795.

Diospyros cordifolia Roxb., Pl. Coromandel 1:38, t. 50. 1795

Diospyros glauca Rottler, Neue Schriften Ges. Naturf. Freunde Berlin 4: 221.1803

Diospyros orixensis Klein ex Willd., Sp. Pl. 4: 407.1805

Diospyros bracteata Roxb., Fl. Ind. ed. 1832 2: 539 1832

Diospyros goindu Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4: 111.1852

Diospyros auriculata Wight ex Hiern, Trans. Cambridge Philos. Soc. 12: 188.1873

Diospyros montana var. *cordifolia* (Roxb.) Clarke in Hook.f., Fl. Brit. India 3: 555. 1882

Diospyros humilis Bourd., J. Bombay Nat. Hist. Soc. 12: 352.1899

Diospyros kanjilalii Duthie, Indian Forester 31: 307.1905

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkaballapur, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India except Jammu & Kashmir and Andaman & Nicobar Islands; Sri Lanka, SE Asia to N Australia.

Local/Common/Vernacular names: *Balagunte, Jagalaganti* (Kannada)

Diospyros neilgerrensis (Wight) Kosterm., Ceylon J. Sci., Biol. Sci. 12: 106.1977

Plate 22D

Maba neilgerrensis Wight, Icon. Pl. Ind. Orient. 4: 10.1848

Diospyros ferrea var. *neilgerrensis* (Wight) Bakh., Bull. Jard. Bot. Buitenzorg III, 15: 57.1937

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Daman & Diu, Goa, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Karimaru, Ruktroora, Sunnala* (Kannada)

Diospyros nilagirica Bedd., Ic. t. 136. 1868-1874; Ramesh & Pascal, J. Bombay Nat. Hist. Soc., 90: 324. 1993.

Diospyros hirsuta f. *nilagirica* (Bedd.) Hiern, Trans. Cambridge Philos. Soc. 12: 164.1873

Distribution: Karnataka (Kodagu), introduced elsewhere, Kerala, Tamil Nadu; endemic.

Diospyros oocarpa Thwaites, Enum. 180. 1860; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Assam, Goa, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; Sri Lanka, SE Asia.

Local/Common/Vernacular name: *Kalu kadumberiya* (Kannada)

Diospyros ovalifolia Wight, Icon. Pl. Ind. Orient. 4: T. 1227. 1848; C.J.Saldanha, Fl. Karnataka 1: 340. 1984.

Distribution: Karnataka (Davanagere, Mysuru), Andhra Pradesh, Assam, Kerala, Madhya Pradesh, Tamil Nadu; Malaysia, Sri Lanka.

Diospyros paniculata Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4:109. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Shivamogga, Uttara Kannada), Chattisgarh, Goa, Kerala, Tamil Nadu, West Bengal; endemic.

Local/Common/Vernacular name: *Ebony* (English); *Kur-koomar* (Kannada)

Diospyros pruriens Dalzell, Hooker's J. Bot. Kew Gard. Misc. 4:110. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Karimaru* (Kannada)

Diospyros ridleyi Bakh., Gard. Bull. Straits Settlem. 7: 183.1933

Diospyros pyrrhocarpa var. *andamanica* Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 46(2): 236.1877

Diospyros pyrrhocarpoides B.R.Ramesh & D.De Franceschi, Blumea 38: 133.1993

Distribution: Karnataka (Hassan), Andaman & Nicobar Islands, Kerala;Malaysia.

Diospyros sahayadryensis P.Daniel & Vajr., J. Econ. Taxon. Bot. 3: 675 1982; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Holochilus micranthus Dalzell in Hooker's J. Bot. Kew Gard. Misc. 4: 291 1852 non *Diospyros micrantha* Sandwith 1950

Maba micrantha (Dalzell) Hiern, Trans. Cambridge Philos. Soc. 12: 133 1873

Ebenus micrantha (Dalzell) Kuntze, Revis. Gen. Pl. 2: 408.1891

Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.

Diospyros saldanhae Kosterm. in J. Bombay Nat. Hist. 74: 326. 1978; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Plate 22E

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Goa, Kerala; endemic.

Diospyros sylvatica Roxb., Pl. Coromandel 1:37, t. 47. 1795; B.D.Sharma et al., Fl. Karnataka Anal. 157. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andhra Pradesh, Bihar, Jharkhand, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, West Bengal; Sri Lanka.

Local/Common/Vernacular names: *Akkasarali, Bilisarali, Kakisarali* (Kannada)

Diospyros trichophylla Alston in Trimen, Handb. Fl. Ceylon 6(Suppl.): 181. 1931

Distribution: Karnataka (Kodagu?), Andaman & Nicobar Islands, Kerala; Tamil Nadu; Sri Lanka.

Diospyros vera (Lour.) A.Chev., Cat. Pl. Jard. Bot. Saigon 31 1919

Ebenoxylum verum Lour., Fl. Cochinch. 613 1790

Ferreola buxifolia (Rottb.) Roxb., Pl. Coromandel 1: 35.1795

Maba buxifolia (Rottb.) Pers., Syn. Pl. 2: 606 1807

Maba angustifolia Miq., Anal. Bot. Ind. 3: 13. 1852

Maba nigrescens Dalzell in Dalzell & A.Gibson, Bombay Fl. 142. 1861

Ebenus nigrescens (Dalzell) Kuntze, Revis. Gen. Pl. 2: 408.1891

Diospyros ferrea (Willd.) Bakh., Gard. Bull. Straits Settlem. 7: 162 1933; B.D.Sharma et al., Fl. Karnataka Anal. 156. 1984.

Diospyros nigrescens (Dalzell) Saldanha in Saldanha & Nicolson, Fl. Hassan 197. 1976

Diospyros angustifolia (Miq.) Kosterm., Ceylon J. Sci. 12: 106. 1977

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Shivamogga, Uttara Kannada), Peninsular India; tropics of the old world.

Local/Common/Vernacular names: *Persimmon, Philippines ebony* (English); *Karugana, Piccane* (Kannada)

Elaeagnaceae

Elaeagnus L.

Elaeagnus conferta Roxb., Fl. Ind. 1: 460. 1820; B.D.Sharma et al., Fl. Karnataka Anal. 240. 1984.

Elaeagnus arborea Roxb., Fl. Ind. 1: 461. 1820

Elaeagnus kologa Schltdl., Prodr. 14: 611.1857; B.D.Sharma et al., Fl. Karnataka Anal. 240. 1984.

Elaeagnus wallichiana Schltdl., Prodr. 14: 612. 1857

Elaeagnus arborea var. *maculata* Servett., Beih. Bot. Centralbl. 25(2): 89. 1909

Elaeagnus conferta subsp. *nigricans* var. *maculata* Serverttaz, Beihe. Bot. Centralbl. 25(2): 72. 1909

Elaeagnus conferta var. *calcuttensis* Servett., Beih. Bot. Centralbl. 25(2): 89. 1909

Elaeagnus conferta subsp. *nigricans* var. *maculata* (Serverttaz) D.Basu, J. Econ. Taxon. Bot. 7: 658. 1986

Elaeagnus conferta var. *maculata* (Servett.) D. Basu, J. Econ. Taxon. Bot. 7: 427. 1986

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout including Andaman & Nicobar Islands except NW India & Deserts; China, Myanmar, Laos, Thailand, S & SE Asia.

Local/Common/Vernacular names: *Gunna madapatte balli, Halige balli, Hulige, Hundase beelu* (Kannada); *Ambgool (Kunabi)*

Elaeagnus indica Servett., Bull. Herb. Boissier II, 8: 393.1908; B.D.Sharma et al., Fl. Karnataka Anal. 240. 1984.

Distribution: Karnataka (Bengaluru), Kerala, Tamil Nadu; SE Asia.

Elaeagnus latifolia L., Sp. Pl. 121. 1753.

Elaeagnus thwaitesii Schltdl., Prodr. 14: 611. 1857

Elaeagnus kologa subsp. *wightii* Servett., Bull. Herb. Boissier II, 8: 394. 1908

Distribution: Karnataka (Belagavi), Peninsula, tropical Himalayas, NE India; Bangladesh, China, Myanmar, SE Asia, Sri Lanka.

Elaeagnus triflora Roxb., Fl. Ind. ed.1, 1: 459. 1820

Elaeagnus ferruginea Rich., Mém. Soc. Hist. Nat. Paris 1: 404. 1792

Elaeagnus latifolia var. *triflora* (Roxb.) Schltdl., Prodr. 14: 610. 1857

Distribution: Karnataka (cultivated), Peninsula, tropical Himalayas, NE India; Australia, Myanmar, S & SE Asia, Taiwan.

Elaeocarpaceae

Elaeocarpus L.

Elaeocarpus hygrophilus Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 43(2): 133.1874

Elaeocarpus madopetalus Pierre, Fl. Forest. Cochinch. t. 150.1888

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Arunachal Pradesh, Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Elaeocarpus lanceifolius Roxb., Fl. Ind. ed. 1832 2: 598.1832 ('lanceaefolius'); B.D.Sharma et al., Fl. Karnataka Anal. 32. 1984.

Elaeocarpus serrulatus Benth. in Hooker's J. Bot. Kew Gard. Misc. 3: 263.1851non Roxb. 1832

Distribution: Karnataka (Shivamogga, Uttara Kannada), Andhra Pradesh, Assam, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim, West Bengal; Bhutan, Cambodia, China, Laos, Malaysia, Myanmar, Nepal, Thailand, Vietnam.

Elaeocarpus munroii Mast., Fl. Brit. India 1: 407 1874('munronii'); B.D.Sharma et al., Fl. Karnataka Anal. 32. 1984.

Plate 22F

Monocera munroii Wight, Ill. Ind. Bot. 1: 84 1838 ('munronii').

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Kerala, Maharashtra, Tamil Nadu; endemic.

Elaeocarpus serratus L., Sp. Pl. 515 1753; B.D.Sharma et al., Fl. Karnataka Anal. 32. 1984.

Elaeocarpus perim-kara DC., Prodr. 1: 519.1824

Elaeocarpus ganitrus Roxb. ex G.Don, Gen. Hist. 1: 559.1831

Elaeocarpus perincara Buch.-Ham., Trans. Linn. Soc. London 17: 181. 1835

Elaeocarpus cuniatus Wight, Ill. Ind. Bot. 1: 83.1840

Ganitrus roxburghii Wight, Ill. Ind. Bot. 1: 83.1840

Elaeocarpus malabaricus Oken, Allg. Naturgesch. 3(2): 1194.1841

Elaeocarpus sphaericus (Gaertn.) K.Schum., Nat. Pflanzenfam. 3(6): 5. 1890

var. *serratus*

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Beegada mara* (Kannada)

var. *weibelii* Zmartzy, Kew Bull. 56: 437.2001

Distribution: Karnataka (Kodagu), Kerala; endemic.

Elaeocarpus tuberculatus Roxb., Fl. Ind. ed. 1832, 2: 594 1832; B.D.Sharma et al., Fl. Karnataka Anal. 33. 1984.

Elaeocarpus bilocularis Roxb. ex Wight & Arn., Prodr. Fl. Ind. Orient. 83. 1834

Monocera tuberculata (Roxb.) Wight & Arn., Prodr. Fl. Ind. Orient. 83 1834

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Maharashtra, Tamil Nadu; Indomalesia

Local/Common/Vernacular names: *Bhtali*, *Dande mara*, *Dandele mara*, *Kunge mara*, *Kyasatte mara* (Kannada)

Elaeocarpus variabilis Zmarzty, Kew Bull. 56: 429.2001

Elaeocarpus tectorius auct. non (Lour.) Poir., 1812; Ramamoorthy in Saldanha & Nicolson

Elaeocarpus oblongus Wight & Arn., Prodr. Fl. Ind. Orient. 1: 82.1834 descr. only, non Gaert. 1791

Elaeocarpus glandulosus Wall. ex Merr., J. Arnold Arbor. 32: 194 1951; C.J.Saldanha, Fl. Karnataka 1: 211. 1984; Matthew, Illust. Fl. Palni Hills 79.1996.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Kerala, Maharashtra, Tamil Nadu; endemic.

Elatinaceae

Bergia L.

Bergia ammannioides Roxb. ex Roth, Nov. Pl. Sp. 219 1821; B.D.Sharma et al., Fl. Karnataka Anal. 17. 1984.

Lechea verticillata Willd., Sp. Pl. 1: 495 1797

Distribution: Karnataka (throughout), Andhra Pradesh, Bihar, Goa, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; Pantropical.

Local/Common/Vernacular name: *Nandi kallu soppu* (Kannada)

Bergia capensis L., Mant. Pl. 2: 241 1771; B.D.Sharma et al., Fl. Karnataka Anal. 17. 1984.

Bergia verticillata Willd., Sp. Pl. 2: 770 1799

Distribution: Karnataka (Belagavi, Dakshina Kannada, Uttara Kannada), Andhra Pradesh, Bihar, Delhi, Goa, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; Pantropical.

Elatine L.

Elatine ambigua Wight, Bot. Misc. 2: 103 1831; B.D.Sharma et al., Fl. Karnataka Anal. 17. 1984.

Distribution: Karnataka (Mysuru), Andhra Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Rajasthan, Tamil Nadu; China, Indonesia, Japan, Malaysia, Turkey, Vietnam, introduced elsewhere.

Ericaceae

Gaultheria Kalm ex L.

Gaultheria fragrantissima Wall., Asiat. Res. 13: 397.1820; B.D.Sharma et al., Fl. Karnataka Anal. 151. 1984.

Andromeda katagherensis Hook., Hooker's Icon. Pl. 3: t. 246.1839

Leucothoe katagherensis DC., Prodr. 7: 606.1839

Gaultheria hirsuta Gardner ex C.B.Clarke in Hook.f., Fl. Brit. India 3: 458.1882

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu, West Bengal, N E India; Bhutan, China, Myanmar, Nepal, Sri Lanka.

Vaccinium L.

Vaccinium neilgherrense Wight, Calcutta J. Nat. Hist. 8: 173.1847; B.D.Sharma et al., Fl. Karnataka Anal. 151. 1984.

Epigynium neilgherense (Wight) Klotzsch, Linnaea 24: 50.1852

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Kerala, Tamil Nadu; endemic.

Eriocaulaceae**Eriocaulon L.**

Eriocaulon achiton Körn., Linnaea 27: 630.1856; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Uttara Kannada), Goa, Maharashtra, Meghalaya; Bangladesh, Pakistan, Thailand, Vietnam

Eriocaulon albotetrandra P.Biju, Josekutty & Augustine, Webbia 73 (1): 1. 2017.

Distribution: Karnataka (Uttara Kannada); endemic.

Eriocaulon anshiense Punekar, Malpure & Lakshmin., Sida 21: 626.2004; Punekar & Lakshmin. Fl. Anshi Natl. Park 525. 2011.

Distribution: Karnataka (Uttara Kannada-Anshi National Park); endemic.

Eriocaulon balakrishnanii Punekar, Lakshmin. & Vasudeva Rao, Nord. J. Bot. 22: 609. 2003; Punekar & Lakshmin. Fl. Anshi Natl. Park 526. 2011.

Distribution: Karnataka (Uttara Kannada-Anshi National Park); endemic.

Eriocaulon barba-caprae Fyson, J. Indian Bot. 2: 197.1921; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Distribution: Karnataka (Chikkamagaluru); endemic.

Eriocaulon belgaumensis Shimpale & S. R. Yadav, Kew Bull. 65: 337. 2010

Distribution: Karnataka (Belagavi); endemic.

Eriocaulon breviscapum Körn., Linnaea 27: 676.1856; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Distribution: Karnataka (Belagavi, Shivamogga, Uttara Kannada); endemic.

Eriocaulon brownianum Mart., Pl. Asiat. Rar. 3: 25.1832

Eriocaulon nilagirense Steud., Syn. Pl. Glumac. 2: 271.1855; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Eriocaulon brownianum var. *nilagirense* (Steud.) Fyson, J. Indian Bot. 2: 262.1921

Distribution: Karnataka (Chikkamagaluru, Hassan), Assam, Karnataka, Kerala, Maharashtra, Manipur, Meghalaya, Tamil Nadu; Bangladesh, Cambodia, China, Indonesia, Lao, Myanmar, Sri Lanka, Thailand; Vietnam

Eriocaulon cinereum R.Br., Prodr. Fl. Nov. Holl. 254.1810; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Leucocephala spathacea Roxb., Fl. Ind. ed.2, 3: 613.1832

Eriocaulon sieboldianum Siebold & Zucc. ex Steud., Syn. Pl. Glumac. 2: 272.1855

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Dharwad, Hassan, , Kolar, Mysuru, Tumakuru, Udupi, Uttara Kannada), peninsular to NE India;paleotropical.

Eriocaulon conicum (Fyson) C.E.C.Fisch., Fl. Madras 1620.1931; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Eriocaulon dianae var. *conicum* Fyson, J. Indian Bot. 2: 260.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Chamarajanagar, Dakshina Kannada), Kerala, Maharashtra, Odisha, Tamil Nadu; endemic.

Eriocaulon cuspidatum Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 281.1851; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Distribution: Karnataka (Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Eriocaulon dalzellii Körn., Linnaea 27: 605.1856; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Eriocaulon rivulare Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 280.1851; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Eriocaulon dalzellii var. *glabratum* Moldenke, Phytologia 28: 192.1974

Distribution: Karnataka (Belagavi, Mysuru, Shivamogga, Uttara Kannada), Goa, Maharashtra; endemic.

Eriocaulon duthiei Hook.f., Fl. Brit. India 7: 578.1893

Distribution: Karnataka (Belagavi), Chattisgarh, Madhya Pradesh, Maharashtra; endemic.

Eriocaulon elenorae Fyson, J. Indian Bot. 2: 316.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Hassan, Kodagu, Mysuru, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Eriocaulon eurypeplon Körn., Linnaea 27: 685.1856; Punekar & Lakshmin. Fl. Anshi Natl. Park 527. 2011.

Eriocaulon vanheurckii Müll.Arg., Observ. Bot. Descript. Pl. Nov. Herb. Van Heurckiani 1: 98.1870; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Eriocaulon bombayanum Ruhland, Pflanzenr. IV, 30: 104.1903

Eriocaulon thomasii Fyson, J. Indian Bot. 2: 318.1921

Eriocaulon indicum Moldenke, Phytologia 3: 163 1949

Eriocaulon keralense R.Ansari, V.J.Nair & N.C.Nair, J. Econ. Taxon. Bot. 11: 235.1987

Distribution: Karnataka (Belagavi, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Eriocaulon fluviatile Trimen, J. Bot. 23: 270.1885; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Shivamogga, Uttara Kannada), Kerala, Madhya Pradesh, Maharashtra; Sri Lanka.

Eriocaulon hamiltonianum Mart., Pl. Asiat. Rar. 3: 29.1832; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Eriocaulon oinereum Buch.-Ham. ex Mart., Pl. Asiat. Rar. 3: 29.1832

Eriocaulon hamiltonianum var. *minor* Fyson, J. Indian Bot. 2: 313.1921

Distribution: Karnataka (?), N & NE India; endemic.

Eriocaulon heterolepis Steud., Syn. Pl. Glumac. 2: 271.1855; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Eriocaulon dianae Fyson, J. Indian Bot. 2: 259.1921; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Eriocaulon dianae var. *longibracteatum* Fyson, J. Indian Bot. 2: 259.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Udupi, Uttara Kannada), Goa, Gujarat, Kerala, Maharashtra, Rajasthan; endemic.

Eriocaulon hookerianum Stapf, Trans. Linn. Soc. London, Bot. 4: 243.1894; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Hassan); Indonesia, Malaysia, Papua New Guinea, Vietnam.

Eriocaulon infirmum Steud., Syn. Pl. Glumac. 2: 271.1855; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Hassan); Myanmar, Thailand.

Eriocaulon kanarensse Punekar, Watve & Lakshmin., Sida 21: 628.2004

Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.

Eriocaulon lanceolatum Miq. ex Körn., Linnaea 27: 656.1856; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Eriocaulon lanceolatum var. *pilosum* Moldenke, Phytologia 3: 164.1949

Eriocaulon kasaragodense R.Ansari, V.J.Nair & N.C.Nair, J. Econ. Taxon. Bot. 11: 235.1988

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Eriocaulon longicuspe Hook.f., Fl. Brit. India 6: 573.1893

Eriocaulon polycephalum Hook.f., Fl. Brit. India 6: 573.1893; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Eriocaulon longicuspe var. *polycephalum* (Hook.f.) Fyson, J. Indian Bot. Soc. 2: 308. 1921

Distribution: Karnataka (Chikkamagalurru, Hassan), Kerala, Madhya Pradesh, Maharashtra Tamil Nadu; Sri Lanka.

Eriocaulon margaretae Fyson, J. Indian Bot. 2: 316.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Chamarajanagar, Shivamogga), Maharashtra; endemic.

Eriocaulon meeboldii R.Ansari & N.P.Balakr., Eriocaul. India 139.1994

Eriocaulon hamiltonianum var. *minimum* Fyson, J. Indian Bot. 2: 313.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984 ('minima').

Distribution: Karnataka (Belagavi, Uttara Kannada); endemic.

Eriocaulon minutum Hook.f., Fl. Brit. India 6: 579.1893; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Eriocaulon humile Moldenke, Phytologia 3: 162.1949

Distribution: Karnataka (Belagavi, Kodagu, Uttara Kannada), Maharashtra, Rajasthan, Tamil Nadu; endemic.

Eriocaulon odoratum Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 280.1851

Eriocaulon collinum Hook.f., Fl. Brit. India 6: 584.1893; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Eriocaulon christopheri Fyson, Bull. Misc. Inform. Kew 1914: 330.1914

Eriocaulon oliveri Fyson, Bull. Misc. Inform. Kew 1914: 331.1914

Eriocaulon collinum var. *nanum* Moldenke, Phytologia 28: 101.1974

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Davanagere, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra and Tamil Nadu; Sri Lanka.

Eriocaulon palghatense R. Ansari & N. P. Balakr., Eriocaul. India 111.1994

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttra Kannada), Goa, Kerala; endemic.

Eriocaulon parviflorum (Fyson) R.Ansari & N.P.Balakr., Eriocaul. India 53.1994

Eriocaulon dianae var. *parviflorum* Fyson, J. Indian Bot. 2: 260.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Dharwad, Kodagu, Mysuru, Uttara Kannada), Goa, Kerala, Madhya Pradesh, Maharashtra; endemic.

Eriocaulon peninsulare Punekar & Lakshmin., Nordic J. Bot. 23: 173.2004; Punekar & Lakshmin. Fl. Anshi Natl. Park 530. 2011.

Distribution: Karnataka (Uttara Kannada-Anshi National Park); endemic.

Eriocaulon quinquangulare L., Sp. Pl. 87.1753; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Sphaerochloa quinquangularis (L.) P.Beauv. ex Desv., Ann. Sci. Nat. (Paris) 13: 47.1828

Eriocaulon melaleucum Bong., Mém. Acad. Imp. Sci. St.-Pétersbourg, Sér. 6, Sci. Math. VI, 1: 629 1831

Leucocephala graminifolia Roxb., Fl. Ind. ed. 1832 3: 612.1832

Eriocaulon martianum Wall. ex Körn., Linnaea 27: 642.1854

Eriocaulon quinquangulare var. *martianum* Fyson, J. Indian Bot. 2: 204.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Ballari, Chamrajanagar, Chitradurga, Dakshina Kannada, Davanagere, Kalburgi, Hassan, Kodagu, Kolar, Mysuru, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout India; Sri Lanka.

Local/Common/Vernacular names: *Gomajjige gida, Gomti* (Kannada)

Eriocaulon richardianum (Fyson) R.Anvari & N.P.Balakr., Eriocaul. India 51.1994

Eriocaulon dianae var. *richardianum* Fyson, J. Indian Bot. 2: 260.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Mysuru, Udupi, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Eriocaulon ritchieanum Ruhland, Pflanzenr. IV, 30: 73.1903; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Eriocaulon horsley-kundae Fyson, J. Indian Bot. 3: 13.1922

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Goa, Maharashtra; endemic.

Eriocaulon robustobrownianum Ruhland, Pflanzenr. IV, 30: 77.1903; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Eriocaulon mysorense Fyson, Bull. Misc. Inform. Kew 1914: 331.1914; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Eriocaulon rhodae Fyson, J. Indian Bot. 2: 264.1921; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Distribution: Karnataka (Belagavi, Ballari, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Eriocaulon sedgwickii Fyson, J. Indian Bot. 2: 260.1921; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Distribution: Karnataka (Chikkamagaluru, Kalburgi), Maharashtra; endemic.

Eriocaulon setaceum L., Sp. Pl. 87.1753; B.D.Sharma et al., Fl. Karnataka Anal. 302. 1984.

Eriocaulon intermedium Körn., Linnaea 27: 601.1856

Eriocaulon fluitans Griff. ex Moldenke, Known Geogr. Distrib. Mem. Erioc. 34. 1946

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala; Sri Lanka.

Eriocaulon sexangulare L., Sp. Pl. 87.1753; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Eriocaulon setaceum B.Heyne ex Wall., Numer. List 6073. 1831.

Eriocaulon wallichianum Mart., Pl. Asiat. Rar. 3: 26 1832

Eriocaulon wallichianum Wight ex Kunth, Enum. Pl. 3: 566. 1841.

Eriocaulon wallichianum Thwaites, Enum. Pl. Zeyl. 341. 1864.

Eriocaulon wallichianum var. *tenellum* Wight ex Moldenke, Phytologia 3: 397.1950

Eriocaulon sexangulare var. *micronesicum* Moldenke, Phytologia 43: 355 1979

Distribution: Karnataka (Dakshina Kannada, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Africa, Cambodia, China, Indonesia, Japan, Laos, Madagascar, Malaysia, Myanmar, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam.

Eriocaulon sharmae R.Ansari & N.P.Balakr., Eriocaul. India 36.1994

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Eriocaulon sollyanum Royle, Ill. Bot. Himal. Mts. 409, t. 97, f. 1.1840

Eriocaulon quinquangular Mart. ex Korn, Linnaea 645.1854

Eriocaulon subalatum Bojer ex Körn., Linnaea 27: 645.1856

Eriocaulon trilobum Buch.-Ham. ex Körn., Linnaea 27: 645.1856

Eriocaulon trilobum Korn. apud Sprague, Kew Bull. 385.1933

Distribution: Karnataka (Belagavi Dakshina Kannada, Udupi), kerala, Bangladesh, China, Indonesia, Nepal, Sri Lanka, Thailand.

Eriocaulon stellulatum Korn., Linnaea 27: 620.1856; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Eriocaulon stellulatum var. *laosense* Moldenke, Phytologia 7: 119.1960

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Eriocaulon thwaitesii Korn., Linnaea 27: 627.1856; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Eriocaulon neesianum Korn., Linnaea 27: 628.1856

Eriocaulon mariae Fyson, Bull. Misc. Inform. Kew 1914: 331.1914

Eriocaulon gamblei C.E.C.Fisch., Bull. Misc. Inform. Kew 1930: 160.1930

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Eriocaulon truncatum Buch.-Ham. ex Mart., Pl. Asiat. Rar. 3: 29.1832; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Tumakuru, Udupi, Uttara Kannada), Andaman and Nicobar Islands, Andhra Pradesh, Assam, Bihar, Goa, Kerala, Meghalaya, Maharashtra, Manipur, Madhya Pradesh, Tamil Nadu, Orissa, Sikkim, West Bengal; Africa (Tanzania) Bangladesh, Bhutan, China, Indonesia, Japan, Malaysia, Myanmar, Papua New Guinea, Philippines, Sri Lanka, Taiwan, Thailand.

Eriocaulon tuberiferum A.R.Kulk. & Desai, J. Bombay Nat. Hist. Soc. 71: 81.1974

Distribution: Karnataka (Belagavi), Maharashtra; endemic.

Eriocaulon walkeri Hook.f., Fl. Brit. India 6: 583.1893

Eriocaulon quadrangulare var. *argenteum* Thwaites, Enum. Pl. Zeyl. 341.1864

Eriocaulon quadrangulare var. *walkeri* Fyson, J. Indian Bot. 2: 204.1921; B.D.Sharma et al., Fl. Karnataka Anal. 301. 1984.

Distribution: Karnataka (Uttara Kannada), Kerala? Tamil Nadu?; Sri Lanka.

Eriocaulon wightianum Mart., Pl. Asiat. Rar. 3: 28.1832; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Eriocaulon sericans Mart., Pl. Asiat. Rar. 3: 29.1832

Eriocaulon helferi Hook.f., Fl. Brit. India 6: 583.1893

Eriocaulon wightianum var. *helferi* (Hook.f.) Fyson, J. Indian Bot. 2: 266 1921

Distribution: Karnataka (Kodagu, Uttara Kannada), Andaman & Nicobar Islands, Maharashtra, Myanmar, Thailand.

Eriocaulon xeranthemum Mart., Pl. Asiat. Rar. 3: 29.1832; B.D.Sharma et al., Fl. Karnataka Anal. 303. 1984.

Eriocaulon pygmaeum Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 281.1851

Distribution: Karnataka (Belagavi, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands Assam, Bihar, Goa, Karnataka, Kerala, Maharashtra, Meghalaya, West Bengal; Myanmar, Nepal.

Erythroxylaceae

Erythroxylum P. Browne

Erythroxylum coca Lam., Encycl. 2: 393 1786; B.D.Sharma et al., Fl. Karnataka Anal. 33. 1984.

Distribution: Karnataka (Uttara Kannada), Bihar, Maharashtra, Manipur, Tamil Nadu; native to tropical America.

Local/Common/Vernacular names: *Coca leaf, Cocaine plant, English coca* (English)

Erythroxylum monogynum Roxb., Pl. Coromandel 1: 61, t.88. 1798; B.D.Sharma et al., Fl. Karnataka Anal. 33. 1984.

Sethia indica DC. nom. superfl., Prodr. 1: 576. 1824

Erythroxylum indicum (DC.) Bedd., Fl. Sylv. t. 81. 1871

Distribution: Karnataka (Ballari, Bengaluru, Chikkaballapur, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Tumakuru), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Bastard sandal* (English); *Devadaru* (Kannada)

Erythroxylum obtusifolium Hook.f., Fl. Brit. India 1: 415 1874

Distribution: Karnataka (Dakshina Kannada), Tamil Nadu; Sri Lanka.

Euphorbiaceae

Acalypha L.

Acalypha alnifolia Klein ex Willd., Sp. Pl. 4: 525. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 243. 1984.

Acalypha cylindrica Roxb., Fl. Ind. ed. 1832 3: 678.1832

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Davanagere, Kolar, Mysuru, Raichur), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Acalypha brachystachya Hornem., Enum. Pl. Hort. Hafn. rev. ed.: 1.1807 & Cat. Hort. Hafn. Willd. Enum. Hort. Berol 992. 1809; Govaerts et al., World Checkl.& Bibl. Euphorbiaceae 1: 40. 2000.; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Acalypha supera Forssk., Fl. Aegypt.-Arab. 162. 1775.

Acalypha conferta Roxb., Fl. Ind. ed. 1832 3: 677.1832

Ricinocarpus brachystachyus (Hornem.) Kuntze, Revis. Gen. Pl. 2: 617.1891

Distribution: Karnataka (Chikkamagaluru, Davanagere, Mysuru, Shivamogga), Andhra Pradesh, Himachal Pradesh, Jammu & Kashmir, Kerala, Maharashtra, Meghalaya, Odisha, Uttarakhand, Uttar Pradesh; S to SE Asia, tropical Africa.

Acalypha ciliata Forssk., Fl. Aeg.-Arab. 162. 1775; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Acalypha rubra Wight ex Wall., Numer. List 7781.1847

Ricinocarpus ciliatus (Forssk.) Kuntze, Revis. Gen. Pl. 2: 617.1891

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; tropical Africa, Arabia, Indonesia, Nepal, Pakistan, Sri Lanka.

Acalypha filiformis Poir., Encycl. 6: 205. 1804.

var. **ovalifolia** (Baill.) Govaerts, World Checklist Bibliogr. Euphorbiaceae 62 2000

Acalypha ovalifolia Baill., Recueil Observ. Bot. 1: 269 1861; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Acalypha reticulata var. *ovalifolia* (Baill.) Müll.Arg., Linnaea 34: 32.1865

Distribution: Karnataka (cultivated in Bengaluru)

Acalypha fruticosa Forssk., Fl. Aeg.-Arab. 161. 1775; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Acalypha betulina Retz., Observ. Bot. 5: 30. 1789

Acalypha capitata Wall., Numer. List 7783A, B. 1847

Ricinocarpus fruticosus (Forssk.) Kuntze, Revis. Gen. Pl. 2: 618. 1891

Distribution: Karnataka (Bengaluru, Mandya, Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; tropical Africa, Arabia, Indonesia, Middle East, Myanmar, Sri Lanka.

Acalypha hispida Burm.f., Fl. Indica 303. 1768.

Caturus spiciflorus auct. non L. 1767; Roxb., Fl. Ind. ed. 2, 3: 760. 1832

Distribution: Karnataka (cultivated as ornamental Belagavi, Bengaluru, Kalaburagi, Mysuru), cultivated throughout; native to Oceania.

Local/Common/Vernacular names: *Fox tail, Philippines Medusa, Red hot cat's tail* (English)

Acalypha indica L., Sp. Pl. 1003. 1753.

Plate 6C

Acalypha canascens Wall., Numer. List 7785. 1847

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Kalaburagi, Kodagu, Mysuru, Raichur, Tumakuru, Uttara Kannada, Vijayapura), throughout tropical areas of India; Cambodia, China, Indonesia, Japan, Malaysia, Philippines, Sri Lanka, Taiwan Thailand, Vietnam; tropical Africa; naturalized in tropical America.

Local/Common/Vernacular names: *Kuppe gida, Kuppi gida, Okkalugodi gida, Tappa kire* (Kannada)

Acalypha integrifolia Willd., Sp. Pl. 4: 530. 1805.

subsp. **marginata** (Poir.) Couda in Kew Bull. 34: 42. 1979

Tragia marginata Poir. in Lam., Encycl. 7: 725. 1806

Acalypha marginata (Poir.) Spreng., Syst. 3: 879. 1826; Mull.Arg. in DC., Prodr. 15(2): 850. 1866.

Distribution: Karnataka (cultivated), cultivated as ornamental throughout India.

Acalypha lanceolata Willd., Sp. Pl. 4: 524. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Acalypha hispida Wall., Numer. List 7780. 1847.

Acalypha virginica Wall., Numer. List 7779. 1847

Acalypha fallax Mull.Arg., Linnaea 34: 43. 1865

Acalypha wightiana Müll.Arg., Linnaea 34: 43. 1865

Acalypha albicans B.Heyne ex Hook.f., Fl. Brit. India 5: 417. 1887.nom. invalid.

Acalypha collina Hayne ex Hook.f., Fl. Brit. India 5: 417. 1887.nom. invalid.

Acalypha floribunda B.Heyne ex Hook.f., Fl. Brit. India 5: 417. 1887.nom. invalid.

Acalypha indica K.Schum. & Hollr., Fl. Kais. Wilh. Land 75. 1889.

Ricinocarpus lanceolatus (Willd.) Kuntze, Revis. Gen. Pl. 2: 617. 1891

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dharwad, Mysuru, Tumakuru, Uttara Kannada, Vijayapura), Andaman and Nicobar Islands, Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu Uttar Pradesh, West Bengal; Australia, Pacific islands, S & SE Asia.

Acalypha malabarica Müll.Arg., Linnaea 34: 42. 1865 & in DC., Prodr. 15(2): 871. 1866; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Ricinocarpus malabaricus (Müll.Arg.) Kuntze, Revis. Gen. Pl. 2: 618. 1891

Distribution: Karnataka (Bidar, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Mysuru, Tumakuru, Uttara Kannada), Andhra Pradesh, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh; endemic.

Acalypha stachyura Pax, Repert. Spec. Nov. 7: 110. 1909

Acalypha macrophylla Ule, Verh. Bot. Ver. Brandenb. 1: 79. 1908, nom. illeg.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental throughout India.

Acalypha wilkesiana Müll.Arg. in DC., Prodr. 15: 817. 1866; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Acalypha amentacea subsp. *wilkesiana* (Müll.Arg.) Fosberg, Smithsonian Contr. Bot. 45: 10 1980

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental throughout India; native to Oceania(Vanuatu).

Local/Common/Vernacular names: *Copper leaf, Jacob's coat* (Kannada)

Agrostistachys Dalzell

Agrostistachys borneensis Becc. in Nelle For. Borneo 331. 1902; C.J.Saldanha, Fl. Karnataka 2: 115. 1996

Sarcoclinium longifolium Wight, Icon. Pl. Ind. Orient. 5: 24, t. 1887-88. 1852, non *Agrostistachys longifolia* Kurz, 1875

Agrostistachys longifolia (Wight) Trimen, Syst. Cat. Fl. Pl. Ceylon 81. 1885

Agrostistachys meeboldii Pax & K. Hoffm. in Engl., Pflanzenr., Euphorb.-Chrozoph. 100. 1912

Distribution: Karnataka (Chikkamagaluru, Kodagu, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Agrostistachys indica Dalzell in Hook.'s J. Bot. Kew Gard. Misc. 2: 41. 1850; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Plate 6B

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andaman & Nicobar, Goa, Kerala, Maharashtra, Tamil Nadu; Laos, Myanmar, New Guinea, Philippines, Sri Lanka, Thailand, Vietnam

Aleurites J. R. Forst. & G. Forst.

Aleurites moluccanus (L.) Willd., Sp. Pl. ed. 4, 4: 590. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 245. 1984.

Manihot moluccana (L.) Crantz, Inst. Rei Herb. 1: 167. 1766

Aleurites triloba J.R.Forst. & G.Forst., Char. Gen. Pl. 56. 1775

Camirium cordifolium Gaertn., Fruct. Sem. Pl. 2: 194. 1791

Ricinus dicoccus Roxb., Fl. Ind. ed.2, 3: 690. 1832

Mallotus moluccanus (L.) Müll.Arg., Linnaea 34: 185. 1865

Rottlera moluccana (L.) Scheff., Ann. Mus. Bot. Lugduno-Batavi 4: 122. 1869

Camerium moluccanum (L.) Kuntze, Revis. Gen. Pl. 2: 595. 1891

Distribution: Karnataka (cultivated throughout), cultivated in Assam, Kerala, Maharashtra, Odisha and Tamil Nadu and West Bengal), native of S. E. Asia and Pacific Islands. Widely cultivated in tropics.

Local/Common/Vernacular names: *buah keras or kukui nut tree, Candle nut tree, Indian walnut, varnish tree* (English)

Baliospermum Blume

Baliospermum solanifolium (Burm.) Suresh in Nicolson et al., Interpr. Rheede's Hort. Malab. (Regnum Veg. 119) 106. 1988

Plate 6D

Croton solanifolius Burm., Fl. Malab. 6. 1769

Jatropha montana Willd., Sp. Pl. 4: 563. 1805

Croton solanifolius Geiseler, Croton. Monogr. 74. 1807.

Baliospermum axillare Blume, Bijdr. 604. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 245. 1984.

Croton polyandrus Roxb., Fl. Ind. ed.2, 3: 682. 1832

Croton roxburghii Wall., Rep. Calcutta Bot. Gard. to G.A. Bushby 20. 1840

Baliospermum indicum Decne. in Jacquem., Voy. Inde 4: 154. 1841

Ricinus montanus (Willd.) Wall., Numer. List 7727. 1847

Baliospermum polyandrum (Roxb.) Wight, Icon. Pl. Ind. Orient. 5: t. 1885. 1852

Baliospermum angulare Decne. ex Baill., Étude Euphorb. 395. 1858

Baliospermum moritzianum Baill., Étude Euphorb. 395. 1858

Baliospermum montanum (Willd.) Mull.Arg. in DC., Prodr. 15(2): 1125. 1866

Baliospermum axillare var. *dioica* Haines, For. Fl. Chotta Nagpur 583. 1919 & Bot. Bihar Orissa 115. 1921

Baliospermum raziana Keshav, Murthy & Yogan. Curr. Sci. 56(10): 486. 1987

Distribution: Karnataka (throughout), throughout India; Bangladesh, Bhutan, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Nepal, Pakistan, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular name: *Danti* (Kannada); *Vayuche* (Kunabi); *Naga danti* (Sanskrit)

Blachia Baill.

Blachia andamanica (Kurz) Hook.f., Fl. Brit. India 5: 403. 1887.

subsp. **Denudata** (Benth.) N.P.Balakr. & Chakrab., Proc. Indian Acad. Sci. (Pl. Sci.) 99: 571, f. 2. 1989
Plate 22G

Croton umbellatus Dalzell & A.Gibson, Bombay Fl. 231. 1861, excl. quoad Wight, Icon. Pl. Ind. Orient. t. 1874, non *Blachia umbellata* (Willd.) Baill. 1858

Blachia denudata Benth., J. Linn. Soc. Bot. 17: 226. 1880; B.D.Sharma et al., Fl. Karnataka Anal. 245. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Blachia calycina Benth., J. Linn. Soc. Bot. 17: 226. 1880; B.D.Sharma et al., Fl. Karnataka Anal. 245. 1984.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Blachia umbellata Gamble, Fl. Madras 2(7): 1338. 1925, p.p. non (Willd.) Baill., 1858

Croton umbellatus Willd., Sp. Pl. 4: 545. 1805.

Codiaeum umbellatus (Willd.) Muell.-Arg., DC., Prodr. 15: 1118. 1866

Blachia reflexa Benth., J. Linn. Soc. Bot. 17: 226. 1880

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Shivamogga, Uttara Kannada), Assam, Kerala, Tamil Nadu; Sri Lanka.

Chrozophora A.Juss.

Chrozophora plicata (Vahl) A.Juss. ex Spreng., Syst. Veg. 3: 850 1826; N.P.Singh, Fl. East. Karnataka 2: 561. 1988

Croton hastatus Burm.f., Fl. Indica 305. 1768

Croton tinctorius Burm.f., Fl. Indica 304. 1768

Croton plicatus Vahl, Symb. Bot. 1: 78 1790

Chrozophora burmanni Spreng., Syst. Veg. 3: 851. 1826

Croton asper J.König ex Roxb., Fl. Ind. ed.2, 3: 681. 1832

Chrozophora prostrata Dalzell & Gibson, Bombay Fl. 233. 1861

Chrozophora parvifolia Klotsch ex Schweinf., Pl. Quaed. Nilot. 11. 1862

Chrozophora plicata var. 2 & 3 Hook.f., Fl. Brit. India 5: 410. 1887

Tournesol plicata (Vahl) Kuntze, Revis. Gen. Pl. 2: 621. 1891

Chrozophora prostrata var. *parvifolia* (Klotzsch ex Schweinf.) N.P.Balakr., Bull. Bot. Surv. India 15: 7 1976

Distribution: Karnataka (throughout), throughout plains, Africa, Middle East, Meditrannian areas, S and SE Asia.

Local/Common/Vernacular name: *Linga menasu* (Kannada)

Chrozophora rattleri (Geiseler) A.Juss. ex Spreng., Euphorb. Gen. 28. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 246. 1984.

Croton rattleri Geiseler, Croton Monogr. 54. 1807

Chrozophora tinctoria auct. non (L.) A.Juss., 1824; Hook.f., Fl. Brit. India 5: 408. 1887, p. p.

Chrozophora plicata var. *rattleri* (Geiseler) Mull.Arg. in DC., Prodr. 15(2): 747. 1866

Distribution: Karnataka (throughout), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Bihar, Chattisgarh, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh West Bengal; S & SE Asia, tropical Africa.

Local/Common/Vernacular names: *Anga menasu*, *Gurugu*, *Suttu mare* (Kannada)

Cleidion Blume

Cleidion javanicum Blume, Bijdr. 613. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 246. 1984. **Plate 6E**

Acalypha spiciflora Burm.f., Fl. Ind.ica 203, t. 61, f. 2. 1768, nom. confusum.

Cleidion spiciflorum auct. non Burm.f., 1768; Govaerts et al., World Checkl. & Bibl. Euphorbiaceae 1: 358. 2000.

Lasiostyles salicifolia C.Presl, Abh. Königl. Böhm. Ges. Wiss. V, 3: 579. 1845

Rottlera urandra Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 229. 1851

Tetraglossa indica Bedd., Madras J. Lit. Sci. II, 22: 70. 1861

Cleidion javanicum var. *longipedicellatum* Chakrab. & M.Gangop., J. Econ. Taxon. Bot. 12: 491. 1989

Cleidion javanicum var. *moniliflorum* (Airy Shaw) Chakrab. & M.Gangop., J. Econ. Taxon. Bot. 12: 493. 1989 non *Cleidion moniliflorum* Airy Shaw, 1980

Cleidion spiciflorum var. *moniliflorum* (Airy Shaw) Radcl.-Sm. & Govaerts, Kew Bull. 52: 478. 1997 non *Cleidion moniliflorum* Airy Shaw, 1980

Cleidion spiciflorum var. *longipedicellatum* (Chakrab. & M.Gangop.) Susila & N.P.Balakr., J. Econ. Taxon. Bot. 28: 520. 2005

Distribution: Karnataka (throughout), Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Bihar, Kerala, Maharashtra, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim, Tamil Nadu, Tripura West Bengal; Australia, Brunei, China, Indonesia, Malaysia, Myanmar, Philippines, Sri Lanka.

Local/Common/Vernacular name: *Ankla* (Kannada)

Codiaeum A.Juss.

Codiaeum variegatum (L.) Rumph. ex A.Juss., Euphorb. Gen. 80, 111, t. 9, f. 30. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Croton variegatus L., Sp. Pl. 2: 1199. 1753 ('variegatum') & ed. 3, 1424. 1764

Crozophyla variegata (L.) Raf., Sylva Tellur. 64. 1838

Oxydectes variegata (L.) Kuntze, Revis. Gen. Pl. 2: 613. 1891

Phyllaurea variegata (L.) W.Wight, Contr. U. S. Natl. Herb. 9: 352. 1905

Distribution: Karnataka (cultivated for the ornamental), cultivated for the ornamental foliage in gardens, throughout in tropical and subtropical regions), native of Indonesia, Malaysia, Oceania.

Croton L.

Croton aromaticus L., Sp. Pl. 1005. 1753; C.J.Saldanha, Fl. Karnataka 2: 126. 1996

Croton tiliifolius var. *aromaticus* (L.) Lam., Encycl. 2: 206. 1786

Cascarilla aromatica (L.) Raf., Sylva Tellur. 62. 1838

Oxydectes aromatica (L.) Kuntze, Revis. Gen. Pl. 2: 611. 1891

Distribution: Karnataka (Chikkamagaluru, Kolar, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Croton caudatus Geiseler, Crot. Monogr. 73. 1807; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Croton denticulatus Blume, Bijdr. 603.1826

Croton drupaceus Roxb., Fl. Ind. ed.2, 3: 684.1832

Croton malvifolius Griff., Itin. Pl. Khasyah Mts. 200.1848

Croton caudatus var. *denticulatus* Müll.Arg., Prodr. 15(2): 599.1866

Oxydectes caudata (Geiseler) Kuntze, Revis. Gen. Pl. 2: 611.1891

Croton caudatus var. *obovoideus* N.P.Balakr. & Chakrab., Bull. Bot. Surv. India 25: 190. 1985

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Kolar, Shivamogga), Arunachal Pradesh, Assam, Kerala, Meghalaya, Nagaland, Odisha, Sikkim, Tamil Nadu, West Bengal; S & SE Asia.

Croton gibsonianus Nimmo in J. Graham, Cat. Pl. Bombay 251. 1839; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Oxydectes gibsoniana (Nimmo) Kuntze, Revis. Gen. Pl. 2: 611.1891

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Shivamogga, Uttara Kannada), Kerala, Maharashtra; endemic.

Croton hirtus L'Hér., Stirp. Nov. 17.1785; K.G. Bhat, J. Bombay Nat Hist. Soc. 111(3):248.2014.

Croton glandulosus var. *hirtus* (L'Her.) Mull.Arg. in DC., Prodr. 15(2): 684. 1866

Distribution: Karnataka (Bengaluru, Udupi), Kerala, Tamil Nadu; native to C & S America; introduced weed elsewhere.

Croton klotzschianus (Wight) Thwaites, Enum. Pl. Zeyl. 276 1861; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Tiglium klotzschianum Wight, Icon. Pl. Ind. Orient. 5: t. 1914.1852

Croton caudatus var. *klotzschianus* Müll.Arg., Prodr. 15(2): 599.1866

Distribution: Karnataka (Bengaluru), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Croton laccifer L., Sp. Pl. 1005. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Aleurites laccifera (L.) Willd., Sp. Pl. 4: 590.1805

Oxydectes laccifera (L.) Kuntze, Revis. Gen. Pl. 2: 610.1891

Croton aromaticus var. *lacciferus* (L.) Trimen, Handb. Fl. Ceylon 4: 48.1898

Distribution: Karnataka (Chikkamagaluru, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Croton lawianus Nimmo ex Dalzell & A.Gibson in J.Graham, Cat. Pl. Bombay 251. 1839; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Trigonostemon lawianus (Nimmo ex Dalzell & A.Gibson) Müll.Arg., Linnaea 34: 212.1865

Distribution: Karnataka (Chikkamagaluru, Mysuru), Maharashtra; endemic.

Croton malabaricus Bedd., For. Man. Bot. 204. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Distribution: Karnataka (Hassan), Kerala, Tamil Nadu; endemic.

Croton persimilis Mull.Arg., Linnaea 34: 116. 1865 & in DC., Prodr. 15(2): 618. 1866 excl. var. *glabrata* emend. Philcox in Dassan. & Clayton, Rev. Handb. Fl. Ceylon 11: 92. 1997; Chakrab. & N.P. Balakr., J. Econ. Taxon. Bot. 30: 298, f. 1. 2006.

Croton oblongifolius Roxb., Fl. Ind.ed.2, 3: 685. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Croton boragatch Roxb. ex Wall., Numer. List 7735.1847

Croton elaeocarpifolius Wall., Numer. List 7734.1847

Oxydectes oblongifolia Kuntze, Revis. Gen. Pl. 2: 612.1891

Croton roxburghii N.P.Balakr., Bull. Bot. Surv. India 3: 39. 1962.

Croton virbalae M.R. Almeida, J. Bombay Nat. Hist. Soc. 100: 580. 2003

Distribution: Karnataka (Belagavi, Dakshina Kannada, Dharwad, Udupi, Uttara Kannada), extending from Peninsula to NE India; Bangladesh, Bhutan, Cambodia, China, Laos, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Croton rheedei J.Graham, Cat. Pl. Bombay 182.1839; Hook.f., Fl. Brit. India 5: 395. 1887.

Croton gibsonii J.Graham, Cat. Pl. Bombay 182.1839

Oxydectes rheedei (J.Graham) Kuntze, Revis. Gen. Pl. 2: 612.1891

Distribution: Karnataka (), Kerala, Maharashtra; endemic.

Croton tiglum L., Sp. Pl. 1004. 1753; C.J.Saldanha, Fl. Karnataka 2: 128. 1996

Croton jamalgota Buch.-Ham., Trans. Linn. Soc. London 14: 258.1824

Croton pavana Buch.-Ham., Trans. Linn. Soc. London 14: 259.1825('parana', 'parona' & 'parane').

Kurkas tiglum (L.) Raf., Sylva Tellur. 62.1838

Tiglum officinale Klotzsch, Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur. 19(Suppl. 1): 418.1843

Oxydectes pavana (Buch.-Ham.) Kuntze, Revis. Gen. Pl. 2: 610.1891

Oxydectes tiglum (L.) Kuntze, Revis. Gen. Pl. 2: 614. 1891

Croton officinalis (Klotzsch) Alston, Handb. Fl. Ceylon 4(Suppl.): 264.1931

Croton himalaicus D.G.Long, Notes Roy. Bot. Gard. Edinburgh 44: 170. 1986

Distribution: Karnataka (cultivated), Peninsula extending to Sikkim and NE India; Bangladesh, Bhutan, Cambodia, China, Indonesia, Japan, Malaysia, Myanmar, Nepal, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam.

Local/Common/Vernacular names: *Danti, Japalabeeja, Nepala, Nervalam* (Kannada)

Croton zeylanicus Mull.Arg., Linnaea 34: 107. 1865 & in DC., Prodr. 15(2): 581. 1866; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Croton hypoleucus Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 123. 1851 non Schlecht. 1847

Oxydectes zeylanica (Müll.Arg.) Kuntze, Revis. Gen. Pl. 2: 613.1891

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada), Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Dalechampia L.

Dalechampia stenoloba Raghavan & B.G.Kulk., Kew Bull. 35: 325. 1980; B.D.Sharma et al., Fl. Karnataka Anal. 247. 1984.

Distribution: Karnataka (Chikkamagaluru); endemic.

Dalechampia velutina Wight, Icon. Pl. Ind. Orient.t. 1881. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 248. 1984.

Dalechampia scandens var. *velutina* (Wight) Mull.Arg. in DC., Prodr. 15(2): 1245. 1866; Chakrab. & N.P.Balakr., J. Econ. Taxon. Bot. 32: 603. 2008.

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Dimorphocalyx Thwaites

Dimorphocalyx glabellus Thwaites, Enum. Pl. Zeyl. 278. 1861; Hook.f., Fl. Brit. India 5: 403. 1887 excl. syn. *Trigonostemon lawianus* (Nimmo ex Dalzell & A.Gibson) Müll.Arg.

var. **glabellus**

Croton ramiflorus J.Graham, Cat. Pl. Bombay 182.1839; B.D.Sharma et al., Fl. Karnataka Anal. 248. 1984.

Oxydectes ramiflora (J.Graham) Kuntze, Revis. Gen. Pl. 2: 612.1891

Dimorphocalyx ramiflorus (J.Graham) M.R.Almeida, Fl. Maharashtra 4B: 301.2003

Distribution: Karnataka (Belagavi, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Odisha, Tamil Nadu; Sri Lanka.

var. **lawianus** (Hook.f.) Chakrab. & N.P.Balakr., Proc. Indian Acad. Sci. (Pl. Sci.) 100: 296. 1990; C.J.Saldanha, Fl. Karnataka 2: 130. 1996.

Plate 6F&23A

Dimorphocalyx lawianus Hook.f., Fl. Brit. India 5: 404. 1887 excl. syn. *Trigonostemon lawianus* Muell.Arg.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Epiprinus Griff.

Epiprinus mallotiformis (Müll.Arg.) Croizat, J. Arnold Arbor. 23: 53. 1942; B.D.Sharma et al., Fl. Karnataka Anal. 248. 1984.

Sympphyllia mallotiformis Muell.Arg., Linnaea 34: 156.1865

Cephalocroton indicus Bedd., Fl. Sylv. t. 261.1872

Adenochlaena indica Bedd. ex Hook.f., Fl. Brit. India 5: 418.1887

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Euphorbia L.

Euphorbia antiquorum L., Sp. Pl. 450. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 248. 1984.

Euphorbia arborescens Roxb., Fl. Ind. ed.2, 22: 468.1832

Euphorbia antiquorum var. *polygona* Wight, Icon. Pl. Ind. Orient. 2: t. 897. 1843

Euphorbia mayuranathanii Croizat in Hooker's Icon. Pl. 35 t. 3404. 1940; Binojk. & N.P. Balakr., Euphorbia J. 10: 27. 1996.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Kalaburagi, Kodagu, Mysuru, Raichur, Uttara Kannada, Vijayapura), throughout India; Bangladesh, China, Indonesia, Malaysia, Myanmar, Pakistan, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Indian spurge, Malayan spurge, Square milk hedge, Square spurge, Triangular milkwort, Triangular splurge* (English); *Cedara kalli, Chandara galli, Jadegalli, Manda kalli, Mondu kalli, Mulajemudu, Mundu kalli* (Kannada)

Euphorbia atoto G.Forst., Fl. Ins. Austr. 36 1786; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Anisophyllum atoto (G.Forst.) Klotsch & Garcke, Abh. Königl. Akad. Wiss. Berlin 1859: 37.1860

Chamaesyce atoto (G.Forst.) Croizat, Fl. Hawaïensis 190: Chamaesyce leafl. 4.1936

Distribution: Karnataka (Dakshina Kannada), Cambodia, China, Indonesia, Japan, Laos, Malaysia, Myanmar, Philippines, Sri Lanka, Thailand, Vietnam; Australia, Pacific islands

Euphorbia bonplandii Sweet, Hort. Brit. ed. 2: 456. 1830; Manikand. & Lakshmin., Fl. Rajiv Gandhi Nation Park 361. 2013

Euphorbia marginata Kunth, Nov. Gen. Sp. 2: 61. 1817

Distribution: Karnataka (Kodagu, Mysuru), cultivated; native to Mexico

Euphorbia caducifolia Haines, Indian Forester 40: 154 1914 & Bot. Bihar Orissa 1: 143. 1925; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Distribution: Karnataka (Ballari, Kolar), Andhra Pradesh, Bihar, Chattisgarh, Gujarat, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Uttar Pradesh; endemic.

Local/Common/Vernacular name: *Balagajji* (Kannada)

Euphorbia clarkeana Hook.f., Fl. Brit. India 5: 253. 1887; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Euphorbia heyneana auct. non Spreng. 1826: Boiss. in DC., Prodr.15(2): 35. 1862

Euphorbia clarkeana var. *erecta* Hook. f., Fl. Brit. India 5: 254. 1887

Chamaesyce clarkeana (Hook.f.) Sojak in Cas. Nar. Mus., Odd. Prir.140: 169. 1972

Distribution: Karnataka (Chitradurga, Mysuru, Tumakuru), Andhra Pradesh, Delhi, Gujarat, Haryana, Himachal Pradesh, Jammu & Kashmir, Maharashtra, Punjab, Rajasthan, Uttar Pradesh, Uttarakhand; Afghanistan, Pakistan.

Euphorbia coccinea B.Heyne ex Roth, Nov. Pl. Sp. 228. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Chamaesyce coccinea (B.Heyne ex Roth) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 169. 1972

Distribution: Karnataka (Belagavi, Dharwad, Uttara Kannada, Vijayapura), Goa, Gujarat, Maharashtra, Rajasthan; endemic.

Euphorbia concanensis Janarth. & S.R.Yadav, Rheedea 5: 148. 1995

Chamaesyce concanensis (Janarth. & S.R.Yadav) M.R.Almeida, S.Dutta & S.M.Almeida, J. Bombay Nat. Hist. Soc. 100: 580. 2003

Chamaesyce concanensis (Janarth. & S.R.Yadav) V.S.Raju, J. Econ. Taxon. Bot. 28: 91. 2004

Distribution: Karnataka (Uttara Kannada), Goa, Maharashtra; endemic.

Euphorbia corrigioloides Boiss., Cent. Euphorb. 11. 1860& in DC., Prodr. 15(2): 32.1862; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Euphorbia disticha Engelm. ex Boiss., Prodr. 15(2): 32.1862

Chamaesyce corrigioloides (Boiss.) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 169. 1972

Distribution: Karnataka (Ballari, Bengaluru, Dharwad Ramanagaram), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

Euphorbia cotinifolia L., Sp. Pl. 453. 1753; C.J.Saldanha, Fl. Karnataka 2: 141. 1996

Distribution: Karnataka (cultivated), cultivated throughout India; native to Mexico and South America.

Local/Common/Vernacular names: *Caribbean copper plant, Mexican shrubby spurge Smoketree spurge, tropical smoke bush* (English)

Euphorbia cristata Heyne ex Roth, Nov. Pl. Sp. 226. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Chamaesyce cristata (Heyne ex Roth) G.L. Webster, J. Arnold Arbor. 48: 424. 1967

Distribution: Karnataka (Ballari, Bengaluru, Bidar, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburagi, Kolar, Mandya, Mysuru, Tumakuru, Vijayapura); Andhra Pradesh, Kerala, Jammu & Kashmir, Madya Pradesh, Maharashtra, Tamil Nadu; Bangladesh, Myanmar, Sri Lanka, Thailand.

Euphorbia deccanensis V.S.Raju, Taxon 34: 519 1985

Euphorbia linearifolia Roth, Nov. Pl. Sp. 224. 1821 non Willd. 1799; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Chamaesyce linearifolia (Roth) Soják, Cas. Nár. Mus., Odd. Prír. 140: 169.1972

Euphorbia katrajensis var. *kasargodensis* V.J.Nair, Binojk. & R.Ansari, J. Econ. Tax. Bot. 14:471.1990

Chamaesyce katrajensis var. *kasargodensis* (V.J.Nair et al.) V.S.Raju, J. Econ. Tax. Bot.28:92.2004

Distribution: Karnataka (Dakshina Kannada, Davanagere, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Euphorbia dracunculoides Lam., Encycl. 2: 428. 1788; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Euphorbia lanceolata Spreng., Mant. Prim. Fl. Hal. 41.1807

Euphorbia angustifolia Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 62. 1825, non (Haw.) Steud.,1821

Tithymalus dracunculoides (Lam.) Klotzsch & Garcke, Abh. Akad. Berlin 1859: 84. 1860

Euphorbia hamiltonii Oudejans, Phytologia 67: 46. 1989

Distribution: Karnataka (throughout),throughout plains; Afghanistan, Africa, China, Egypt, Europe,Nepal, Pakistan, West Asia.

Local/Common/Vernacular name: *Bili kaddi* (Kannada); *Sankini* (Sanskrit)

Euphorbia elegans Spreng., Syst. Veg. 3: 794. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Euphorbia variegata auct. non Sims 1815; Roth, Nov. Pl. Sp. 225. 1821

Euphorbia dichotoma Roxb., Fl. Ind. ed.2, 2: 471. 1832

Euphorbia platylepis Decne. in Jaquem., Voy. Inde 4: 156. 1844

Euphorbia strobilifera Dalz., Hooker's J. Bot. Kew Gard. Misc. 3: 229. 1851

Chamaesyce elegans (Spreng.) Sojak in Cas. Nar. Mus., Odd. Prir. 140: 169. 1972

Distribution: Karnataka (Belagavi, Chikkamagaluru, Kalaburagi, Vijayapura), Andhra Pradesh, Chattisgarh, Gujarat, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.

Euphorbia erythroclada Boiss., DC., Prodr. 15: 25. 1862; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Chamaesyce erythroclada (Boiss.) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 169. 1972

Distribution: Karnataka (Chikkamagaluru, Kodagu, Mysuru, Uttara Kannada), Goa, Madhya Pradesh, Maharashtra; endemic.

Euphorbia fusiformis Buch.-Ham. ex D.Don, Prodr. Fl. Nepal 62. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Euphorbia acaulis Roxb., Fl. Ind. 2: 472. 1832; Duthie, Fl. Gangetic Plain 3: 74. 1906.

Distribution: Karnataka (Ballari, Chikkamagaluru, Davanagere, Mysuru, Tumakuru, Uttara Kannada), Andhra Pradesh, Bihar, Goa, Gujarat, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttarakhand, West Bengal; Nepal.

Local/Common/Vernacular name: *Kundali kivi gedde* (Kannada)

Euphorbia gokakensis Malpure, Chandore & S.R.Yadav, Nordic J. Bot.: 2016

Distribution: Karnataka (Belgavi:Gokak); endemic.

Plate 23B

Euphorbia heterophylla L., Sp. Pl. 453. 1753.

Poinsettia heterophylla (L.) Klotzsch & Garcke, Monatsb. Preuss. Akad. Berlin 1859: 253. 1859

var. **heterophylla**

Euphorbia geniculata Ortega, Nov. Pl. Rar. Descr. Dec. 2: 18. 1797

Euphorbia prunifolia Jacq., Pl. Rar. Hort. Schoenbr. 3: 15, t. 277. 1798

Distribution: Karnataka (weed throughout drier areas), weed throughout in tropical areas; native of tropical and subtropical America; pantropical weed.

Local/Common/Vernacular names: *Desert Poinsettia, Fire on the Mountain, Japanese poinsettia, Kaliko Plant, Mexican Fireplant, Milkweed, Painted Euphorbia, Painted leaf, Painted Spurge, Wild Poinsettia* (English); *Beedi soppu, Halalbu* (Kannada)

var. **cyathophora** (J.Murr.) Griseb., Fl. Brit. W. Indies 45, 54. 1859; Boiss. in DC., Prodr. 15 (2): 72. 1862.

Euphorbia cyathophora Murray, Commentat. Soc. Regiae Sci. Gott. 7: 81. 1786; C.J.Saldanha, Fl. Karnataka 2: 135. 1996

Poinsettia cyathophora (Murray) Klotzsch & Garcke, Monatsber. Königl. Preuss. Akad. Wiss. Berlin 1859: 253. 1859

Distribution: Karnataka (cultivated and naturalized throughout), cultivated and naturalized throughout India; native to N & S America.

Local/Common/Vernacular names: *Dwarf poinsettia, Fire-on-the-mountain, Painted leaf* (English)

Euphorbia heyneana Spreng., Syst. Veg. 3: 791 1826

Euphorbia microphylla B. Heyne ex Roth, Nov. Pl. Sp. 229. 1821 non Lam., 1788

Euphorbia orbiculata Miq., Fl. Ned. Ind. 1(2): 421. 1859 non Kunth, 1817

Euphorbia wallichiana Boiss., Prodr. 15(2): 30. 1862

Euphorbia bombayensis Santapau, Bull. Bot. Soc. Bengal 8: 17. 1955

Chamaesyce heyneana (Spreng.) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 169. 1972

Distribution: Karnataka (Belagavi, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburagi, Mysuru, Raichur Tumakuru), Andhra Pradesh, Bihar, Chhattisgarh, Goa, Gujarat, Himachal Pradesh, Jammu & Kashmir, Kerala, Lakshadweep, Madhya Pradesh, Maharashtra, Odisha, Punjab, Tamil Nadu, West Bengal; Africa, China, S & SE Asia; Indian Ocean Islands (Mauritius, Seychelles).

Local/Common/Vernacular name: *Sannele gida* (Kannada)

Euphorbia hirta L., Sp. Pl. 454. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 249. 1984.

Euphorbia pilulifera L., Sp. Pl. 454. 1753, p. p.; Hook.f., Fl. Brit. India 5: 250. 1887.

Chamaesyce pilulifera (L.) Small, Fl. S.E.U.S. 714. 1903

Chamaesyce hirta (L.) Millsp., Publ. Field Columbian Mus., Bot. Ser. 2: 303. 1909

Distribution: Karnataka (Throughout), throughout India; pantropical.

Local/Common/Vernacular names: *Ace gida*, *Halukudi Nenekki soppu*, *Karihalsoppu* (Kannada); *Dugudhika* (Sanskrit)

Euphorbia hypericifolia L., Sp. Pl. 454. 1753; Binojk. & N.P.Balakr., Bull. Bot. Surv. India 35: 74. 1993 (1997) & Genus Euphorbia in India 163, f. 17. 2010.

Chamaesyce hypericifolia (L.) Millsp., Publ. Field Columbian Mus., Bot. Ser. 2: 302. 1909

Distribution: Karnataka (throughout dry areas), throughout hotter parts of India; Pantropical.

Euphorbia hyssopifolia L., Syst. Nat. ed. 10, 2: 1048. 1759; Binojk. & N.P. Balakr., Genus Euphorbia in India 166, t. 31. 2010.

Chamaesyce hyssopifolia (L.) Small, Bull. New York Bot. Gard. 3: 429. 1905

Distribution: Karnataka (throughout), Andhra Pradesh, Maharashtra, Kerala, Tamil Nadu; native to Neotropics.

Euphorbia indica Lam., Encycl. 2: 423 1788; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Euphorbia bracteolaris Boiss., Cent. Euphorb. 8. 1860 & in DC., Prodr. 15(2): 22. 1862

Euphorbia indica var. *angustifolia* Boiss., Prodr. 15(2): 22. 1862

Chamaesyce indica (Lam.) Croizat, Lilloa 8: 406. 1942

Euphorbia hypericifolia var. *coimbatorensis* Chandrab., Fl. Coimbatore 266. 1988

Distribution: Karnataka (throughout), throughout India Africa and Asia; tropical weed.

Euphorbia lacei Craib, Bull. Misc. Inform. Kew 1911: 456. 1911; Binojk. & N.P.Balakr., Genus Euphorbia in India 304, f. 60, t. 13. 2010.

Euphorbia trigona Roxb., Fl. Ind. 2: 468. 1832, non Mill. 1768

Euphorbia barnhartii Croizat, Euphorb. Antiq. Offic. 25, 54. 1934; Santapau in Bull. Bot. Soc. Bengal 8: 8. 1954.

Distribution: Karnataka (cultivated and naturalized), Andhra Pradesh, Bihar, Kerala, Maharashtra, Meghalaya, Nagaland, Tamil Nadu, Uttar Pradesh, West Bengal; native to Molukas.

Euphorbia laciniata Panigrahi, Kew Bull. 30: 531. 1975; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Euphorbia fimbriata B.Heyne ex Roth, Nov. Pl. Sp. 227. 1821, non Scopoli 1788; Gamble, Fl. Madras 2(7): 1274. 1925.

Euphorbia cristata var. *major* Roth., Nov. Pl. Sp. 226. 1821

Euphorbia fimbriata var. *laxa* Boiss. in DC., Prodr. 15(2): 19. 1862, p. p.

Chamaesyce fimbriata (B.Heyne ex Roth) R.R.Rao & Razi, J. Mysuru Univ. B. 23: 53. 1970

Chamaesyce laciniata (Panigrahi) V.S. Raju & P.N.Rao, Phytologia 37: 454. 1977

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Mysuru, Tumakuru, Vijayapura), Andhra Pradesh, Maharashtra, Tamil Nadu; endemic.

Euphorbia lactea Haw., Syn. Pl. Succ. 127. 1812; Binojk. & N.P.Balakr., Genus Euphorbia in India 307, f. 61, t. 14. 2010.

Distribution: Karnataka (Dharawad, Kodagu), Andhra Pradesh, Maharashtra, Kerala, Tamil Nadu Uttar Pradesh, West Bengal; Africa, Europe, S & SE Asia, cultivated in tropical Africa & Asia.

Euphorbia lawsonii Binojk. & Dwarakan, Rheedia 4: 60, f. 1. 1994; Binojk. & N.P.Balakr., Genus Euphorbia in India 120. 2010.

Chamaesyce lawsonii (Binojk. & Dwarakan) V.S.Raju, J. Econ. Taxon. Bot. 28: 92. 2004

Distribution: Karnataka (Ballari), Tamil Nadu; endemic.

Euphorbia leucocephala Lotsy, Bot. Gaz. 20: 350. 1895; Binojk. & N.P. Balakr., Indian J. Forest. 15: 181. 1992 & Genus Euphorbia in India 100, f. 2. 2010.

Distribution: Karnataka (cultivated as ornamental), cultivated as ornamental throughout India; native to Southern Mexico to El Salvador.

Local/Common/Vernacular names: *Little Christmas Flower, Pascuita, Snow bush, Snow flake euphorbia, Snows of Kilimanjaro, White laced euphorbia, White small leaf Poinsettia* (English)

Euphorbia longistyla Boiss., Cent. Euphorb. 9. 1860; C.J.Saldanha, Fl. Karnataka 2: 138. 1996.

Chamaesyce longistyla (Boiss.) Croizat in O.Deg., Fl. Hawaïensis Fam. 190, Chamaesyce, leafl. 2. 1937, in obs. et in adnot.; V.S.Raju & P.N.Rao, Phytologia 37(5): 454. 1977.

Distribution: Karnataka (Ballari), Andhra Pradesh; endemic.

Euphorbia meenae S.Carter, Cact. Succ. J. (Los Angeles) 72: 210, ff. 5 & 6. 2000; Binojk. & N.P.Balakr., Genus Euphorbia in India 370. 2010.

Distribution: Karnataka (Chikkamagaluru), Tamil Nadu; endemic.

Euphorbia milii Des Moul., Bull. Hist. Nat. Soc. Linn. Bordeaux. 1: 27. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Euphorbia splendens Bojer ex Hook. in Bot. Mag. 56: t. 2902. 1829

Euphorbia bojeri Hook., Bot. Mag. 63: t. 3527. 1836; R.Parker, Forest Fl. Punjab 447. 1918.

Distribution: Karnataka (cultivate throughout), cultivate throughout; native of Madagascar.

Local/Common/Vernacular names: *Christ plant, Christ thorn, Crown of thorns* (English)

Euphorbia nerifolia L., Sp. Pl. 451. 1753, p. p.; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984; Binojk. & N.P.Balakr., Genus Euphorbia in India 310, f. 62, tt. 16 & 17 2010.

Euphorbia ligularia Roxb. ex Buch.-Ham., Trans. Linn. Soc. London 14: 285. 1825, nom. Illeg.

Distribution: Karnataka (throughout), throughout plains; tropical Asia.

Local/Common/Vernacular name: *Indian spurge tree* (English); *Yele kalli* (Kannada)

Euphorbia nivulia Buch.-Ham., Trans. Linn. Soc. London 14: 286. 1824; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Euphorbia nerifolia Roxb., Fl. Ind. ed. 1832 2: 467. 1832.

Distribution: Karnataka (throughout), throughout drier parts of India; Bangladesh, Myanmar, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Dabha galli, Dubba kalli Dundu kalli, Yele kalli* (Kannada)

Euphorbia notoptera Boiss. in DC., Prodr. 15: 26. 1862; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Chamaesyce notoptera (Boiss.) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 169. 1972

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Hassan, Mysuru, Uttara Kannada), Goa, Maharashtra; endemic.

Euphorbia orbiculata Kunth in H.B.K., Nov. Gen. Sp. 2: 52. 1817

Anisophyllum orbiculatum (Kunth) Klotzsch & Garcke, Abh. Konigl. Akad. Wiss. Berlin 1859: 24. 1860

Chamaesyce orbiculata (Kunth) Sojak, Cas. Nar. Mus., Odd. Prir. 148: 199. 1980

Distribution: Karnataka (Mysuru), nqutive to tropical America.

Euphorbia pallens Dillwyn, Rev. Hortus Malab. 54. 1839; V.S. Raju, Rheedea 11: 45. 2001. Ven-pala Rheede, Hortus Malab. 10: 115, t. 58. 1690.

Euphorbia articulata Dennst., Schluss. Hort. Malab. 37. 1818; C.J.Saldanha, Fl. Karnataka 2: 134. 1996

Euphorbia halophila Miq., Anal. Bot. Ind. 3: 16. 1852; Boiss. in DC., Prodr. 15 (2): 13. 1862.

Chamaesyce pallens (Dillwyn) V.S.Raju, J. Econ. Taxon. Bot. 28: 92. 2004

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Odisha; Australia, China, Indonesia, Malaysia, Pacefic Islands, Sri Lanka.

Euphorbia parviflora L., Syst. Nat. ed. 10 2: 1047. 1759.

Euphorbia hypericifolia var. *parviflora* auct. non Prain, 1903; Saldanha, Fl. Karnataka 2, 139. 1996,

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kalaburagi, Mysuru, Raichur), Andhra Pradesh, Bihar, Goa, Gujarat, Himachal Pradesh, Madhya Pradesh, Maharashtra, Rajasthan, Uttar Pradesh; Indomalesia

Euphorbia perbracteata Gage, Bull. Misc. Inform. Kew 238. 1914; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Tithymalus perbracteatus (Gage) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 175. 1972. *Euphorbia laeta* auct. non B.Heyne ex Roth, 1821: Sedgw., J. Bombay Nat. Hist. Soc. 26: 599. 1919

Distribution: Karnataka (Bidar, Dharwad, Kalaburagi), Andhra Pradesh, Bihar, Chattisgarh, Daman, Goa, Madhya Pradesh, Maharashtra, Odisha, Uttar Pradesh; endemic.

Euphorbia prostrata Aiton, Hort. Kew (ed. 1) 2: 139. 1789; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Chamaesyce prostrata (Aiton) Small, Fl. SE. U.S. 713. 1903

Euphorbia chamaesyce auct. non L. 1753: Raizada, Suppl. Fl. Gangetic Plain 247. 1976

Distribution: Karnataka (throughout), throughout India including Andaman & Nicobar Islands; naturalized throughout Old World tropics; native to tropical and subtropical America;

Euphorbia pulcherrima Willd. ex Klotzsch. in Otto & Dietr., Allg. Gartenz. 2: 27. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.

Poinsettia pulcherrima (Willd. ex Klotzsch) R.Graham, Edinburgh New Philos. J. 20: 412. 1836

Distribution: Karnataka (cultivated as an ornamental throughout), cultivated as an ornamental throughout India; native of Mexico & C America.

Local/Common/Vernacular name: *The poinsettia* (English)

Euphorbia pycnostegia Boiss., Cent. Euphorb. 9. 1860& in DC., Prodr. 15(2): 18. 1862; C.J.Saldanha, Fl. Karnataka 2: 140. 1996.

Euphorbia pycnostegia var. *laxa* Boiss. in DC., Prodr. 15(2): 19. 1862

Euphorbia zornioides Boiss. in DC., Prodr. 15(2): 19. 1862; Hook.f., Fl. Brit. India 5: 246. 1887.

Euphorbia pycnostegia var. *zornioides* (Boiss.) Santapau, Bull. Bot. Soc. Bengal 8: 11. 1954

Chamaesyce pycnostegia (Boiss.) Sojak, Cas. Nar. Mus., Odd. Prir. 140: 170. 1972

Chamaesyce zornioides (Boiss.) Sojak in Cas. Nar. Mus., Odd. Prir. 140: 170. 1972

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Shivamogga, Uttara Kannada), Andhra Pradesh, Daman, Diu, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.

Euphorbia ridleyi Croizat, Gard. Bull. Straits Settlem. 9: 147 1937

Euphorbia synadenium Ridl., J. Straits Branch Roy. Asiat. Soc. 61: 36 1912; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.

Distribution: Karnataka (Tumakuru), SE Asia.

Euphorbia rosea Retz., Obs. Bot. 4: 26. 1786; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.

Euphorbia satureioides Lam., Encycl. 2: 424. 1788

Euphorbia auricularia Boiss., Cent. Euphorb. 17. 1860 & in DC., Prodr. 15(2): 50. 1862

Chamaesyce rosea (Retz.) G.L. Webster, J. Arnold Arb. 48: 423. 1967, in adnot.

Chamaesyce auricularia (Boiss.) V.S.Raju & P.N.Rao, Phytologia 37: 453. 1977

Distribution: Karnataka (Ballari, Bengaluru, Dakshina Kannada, Dharawad, Hassan, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, Uttarakhand, Uttar Pradesh; Afghanistan, Iran, Nepal, Pakistan, Sri Lanka.

Euphorbia rothiana Spreng., Syst. Veg. 3: 796. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 250. 1984.
Euphorbia laeta B.Heyne ex Roth, Nov. Pl. Sp. 230. 1821
Euphorbia glauca Roxb., Fl. Ind. ed. 22: 473. 1832
Euphorbia oreophila Miq., Annal. Bot. Ind. 3: 17. 1852; Sedgw., J. Bombay Nat. Hist. Soc. 26: 599. 1919.
Euphorbia wightiana Boiss., Cent. Euphorb. 37.1860
Tithymalus rothianus (Spreng.) Klotzsch & Garske, Abh. Konigl. Akad. Wiss. Berlin 1859: 89. 1860
Euphorbia divergens Klotzsch, Bot. Ergebni. Reise Waldemar 114.1862
Euphorbia oreophila var. *pubescens* Boiss. in DC., Prodr. 15(2): 156. 1862
Euphorbia oreophila var. *wightiana* Boiss. in DC., Prodr. 15(2): 156. 1862
Tithymalus divergens Klotzsch, Bot. Ergebni. Reise Waldemar 114.1862
Euphorbia rothiana var. *pubescens* (Boiss.) Gamble, Fl. Madras 2(7): 1278. 1925
 Distribution: Karnataka (throughout), S India to Gujarat, Uttar Pradesh; Sri Lanka.

Euphorbia senguptae N.P.Balakr. & Subr., Bull. Bot. Surv. India 2: 175 1960
Chamaesyce senguptae (N.P.Balakr. & Subr.) V.S.Raju & P.N.Rao, Phytologia 37: 454 1977
 Distribution: Karnataka (Chitradurga), Andhra Pradesh; endemic.

Euphorbia serpens Kunth in H.B.K., Nov., Gen. Sp. (quarto ed.) 2: 52. 1817
Euphorbia microphylla auct. non B.Heyne ex Roth 1821; Hook.f., Fl. Brit. India 5: 252. 1887, p.p.
Chamaesyce serpens (Kunth) Small, Fl. S.E. US. 709. 1903
Euphorbia orbiculata var. *jawaharii* Rajagopal & Panigrahi, Taxon 17: 547. 1968
 Distribution: Karnataka (throughout), throughout plains of India; native of tropical America; pantropical weed.

Euphorbia thymifolia L., Sp. Pl. 454. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.
Anisophyllum thymifolium (L.) Haw., Syn. Pl. Succ. 160.1812
Euphorbia rubicunda Blume, Catalogus 73.1823
Euphorbia foliata Buch.-Ham. ex Dillwyn, Rev. Hortus Malab. 52.1839
Chamaesyce thymifolia (L.) Millsp., Publ. Field Columbian Mus., Bot. Ser. 2(11): 412. 1916
 Distribution: Karnataka (throughout), throughout India; Pantropical weed.
 Local/Common/Vernacular names: *Bili chitra pala*, *Citra ballare* (Kannada); *Laghududhica* (Sanskrit)

Euphorbia tirucalli L., Sp. Pl. 452. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.
 Distribution: Karnataka (cultivated and naturalized throughout), cultivated and naturalized throughout India excluding Andaman & Nicobar Islands; native to Africa, widely cultivated in Asia.
 Local/Common/Vernacular names: *Kalli gida*, *Kollu galli*, *tirukalli* (Kannada); *Trikantaka* (Sanskrit)

Euphorbia tithymaloides L., Sp. Pl. 453 1753
Pedilanthus tithymaloides (L.) Poit., Ann. Mus. Natl. Hist. Nat. Paris 19: 390, t. 19. 1812; B.D.Sharma et al., Fl. Karnataka Anal. 254. 1984.
 Distribution: Karnataka (cultivated throughout as hedge plant) cultivated throughout tropical areas; native to West Indies.
 Local/Common/Vernacular names: *Buck-thorn*, *Christmas candle*, *Cimora misha*, *Devil's backbone*, *Fiddle flower*, *Ipecacuanha*, *Jacob's ladder*, *Japanese poinsettia*, *Jewbush*, *Jew's slipper*, *Milk-hedge*, *Myrtle-leaved spurge*, *Padus-leaved clipper plant*, *Red slipper spurge*, *Redbird cactus*, *Redbird flower*, *Slipper flower*, *Slipper plant*, *Slipper spurge*, *Timora misha*, *Zig-zag plant* (English)

Euphorbia tortilis Rottler ex Ainslie, Materia indica 2: 424 1826; Binojk. & N.P.Balakr., Genus Euphorbia in India 328, f. 67. 2010.
 Distribution: Karnataka (Chitradurga), Andhra Pradesh, Kerala and Tamil Nadu; Sri Lanka.
 Local/Common/Vernacular name: *Mandagalli* (Kannada); *Vajra vriksa* (Sanskrit)

Euphorbia umbellata (Pax) Bruyns, Euphorbia World 3: 5 2007

Synadenium grantii Hook. f., Bot. Mag. t. 5633. 1867; Binojk. & N.P. Balakr., Rheedia 3: 136, f. 1. 1993;
B.D.Sharma et al., Fl. Karnataka Anal. 256. 1984.

Synadenium umbellatum Pax, Bot. Jahrb. Syst. 19: 125 1894

Synadenium umbellatum var. *puberulum* N.E.Br., Fl. Trop. Afr. 6(1): 464 1911

Distribution: Karnataka (cultivated as a hedge plant throughout), cultivated as a hedge plant throughout India; native of tropical Africa.

Local/Common/Vernacular name: *African milk bush* (English)

Excoecaria L.**Excoecaria agallocha** L., Syst. Nat. 2: 1288. 1759; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.

Excoecaria camettia Willd., Sp. Pl. 4: 864 1806

Stillingia agallocha (L.) Baill., Étude Euphorb. 518. 1858

Excoecaria agallocha var. *camettia* (Willd.) Müll.Arg. in DC., Prodr. 15(2): 1221. 1866

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Odisha, Puducherry, Tamil Nadu, West Bengal; Australasia, Pacific Islands, S & SE Asia.

Local/Common/Vernacular names: *Blinding tree* (English); *Harogida, Thilla* (Kannada)

Excoecaria oppositifolia Griff., Calcutta J. Nat. Hist. 4: 386. 1844.

var. *crenulata* (Wight) Chakrsab. & M.Gangop., J. Econ. Taxon. Bot. 18: 208 1994

Excoecaria crenulata Wight, Icon. Pl. Ind. Orient. 5: t. 1865. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.

Excoecaria robusta Hook. f., Fl. Brit. India 5: 474. 1888

Distribution: Karnataka (Hassan, Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Falconeria Royle**Falconeria insignis** Royle, Ill. Bot. Himal. Mts. 354. t.98. f.2. 1839; C.J.Saldanha, Fl. Karnataka 2: 160. 1996.

Falconeria wallichiana Royle, Ill. Bot. Himal. Mts. 354, t. 84a or 98, f. 3. 1839

Falconeria malabarica Wight, Icon. Pl. Ind. Orient. 5: t. 1866. 1852

Gymnobotrys lucida Wall. ex Baill., Étude Euphorb. 527. 1858

Excoecaria insignis (Royle) Mull.Arg. in DC., Prod. 15(2): 1212. 1866

Carumbium insigne (Royle) Kurz, Rep. For. Veg. Pegu, App. A, 114. 1875

Sapium insigne (Royle) Benth. & Hook. f., Gen. Pl. 3: 335. 1880

Sapium insigne var. *malabaricum* (Wight) Hook.f., Fl. Brit. India 5: 472. 1888

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada); Arunachal Pradesh, Andhra Pradesh Assam, Bihar, Goa, Himachal Pradesh, Kerala Maharashtra, Meghalaya, Odisha, Sikkim, Tamil Nadu Uttarakhand, Uttar Pradesh, West Bengal; Bangladesh, Bhutan, Cambodia, China, Laos, Malaysia, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular name: *Kannu pade* (Kannada); *Hura (Kunabi)*

Givotia Griff.**Givotia moluccana** (L.) Sreem., Taxon 24: 696 1975

Croton moluccanus L., Sp. Pl. 1005. 1753, p. p. quoad spec. Hermann.

Givotia rottleriformis Griff. ex Wight, Calcutta J. Nat. Hist. 4: 388. 1844 & Icon. Pl. Ind. Orient. 5: 24, t. 1889. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.

Distribution: Karnataka (throughout dry districts), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *White catamaran tree* (English); *Bili polik, Bili thaale, Boodhi taale, Burle, Kaributti* (Kannada)

Hevea Aubl.
Caoutchoua Gmelin
Siphonia Richard.

Hevea brasiliensis (Willd. ex A.Juss.) Müll.Arg., Linnaea 34: 204. 1865; B.D.Sharma et al., Fl. Karnataka Anal. 252. 1984.

Siphonia brasiliensis Willd. ex A.Juss., Euphorb. Gen. 113, t. 12 1824

Distribution: Karnataka (cultivated in Chikkamagaluru, Dakshina Kannada, Kodagu), cultivated in Assam, Andaman & Nicobar Islands, Assam, Goa Karnataka, Kerala, Manipur, Tamil Nadu; native to Brazil; widely cultivated throughout the tropics.

Local/Common/Vernacular names: *Para Rubber tree, Rubber tree* (English); *Rubber gida, Rubber mara* (Kannada)

Homonoia Lour.

Homonoia intermedia Haines, Bot. Bihar Orissa 2: 111. 1921

Distribution: Karnataka (Belagavi), Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu; endemic.

Homonoia retusa (J.Graham ex Wight) Müll.Arg., Linnaea 34: 200. 1865& in DC., Prodr. 15(2): 1022. 1866; B.D.Sharma et al., Fl. Karnataka Anal. 252. 1984.

Adelia retusa J.Graham ex Wight, Icon. Pl. Ind. Orient. 5: t. 1869. 1852

Distribution: Karnataka (Ballari, Dakshina Kannada, Kodagu, Kolar, Mandya, Raichur, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Homonoia riparia Lour., Fl. Cochinch. 637. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 252. 1984.

Croton salicifolius Geiseler, Croton. Monogr. 6. 1807

Adelia neriifolia B.Heyne ex Roth, Nov. Pl. Sp. 375. 1821

Ricinus salicinus Hassk., Tijdschr. Ned.-Indië 10: 142. 1843

Haematospermum neriifolium (B.Heyne ex Roth) Wall. ex Voigt, Hort. Suburb. Calcutt. 157. 1845

Haematospermum salicinum (Hassk.) Baill., Étude Euphorb. 293. 1858

Spathiostemon salicinus (Hassk.) Hassk., Hort. Bogor. Desc. 41. 1858

Distribution: Karnataka (throughout), throughout India; Brunei, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam.

Local/Common/Vernacular names: *Water croton* (English); *Abbe koralu, Halla nokki, Hole baysa, Hole mekki, Hole nage, Hole nekki, Kiral bogi, Kshoodra pashana bhedi, Neer nekki, Neerganigile, Niganagile* (Kannada); *Jalajabmu, Kshoodra pashana bheda, Pashana beda* (Sanskrit)

Hura L.

Hura crepitans L., Sp. Pl. 1008. 1753

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated in gardens throughout plains; native to S America.

Local/Common/Vernacular name: *Pirangi gundina mara* (Kannada)

Jatropha L.

Jatropha curcas L., Sp. Pl. 1006. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 252. 1984.

Distribution: Karnataka (cultivated and naturalized throughout), cultivated and naturalized throughout India; native to tropical America.

Local/Common/Vernacular names: *Barbados nut, physic nut, Poision nut, purging nut* (English); *Adalu haralu, Bettada haralu Kadandla, Karnochhi, Mara haralu, Turuku haralu* (Kannada)

Jatropha glandulifera Roxb., Fl. Ind. 3: 688. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Adenoropium roxburghii Kostel., Allg. Med.-Pharm. Fl. 5: 1750.1836

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Kalaburagi, Raichur, Vijayapura), Andhra Pradesh,

Madhya Pradesh, Maharashtra, Tamil Nadu, West Bengal; tropical Africa, Deccan plateau, Sri Lanka

Local/Common/Vernacular names: *Jangli Airand, Totlagida* (Kannada)

Jatropha gossypiifolia L., Sp. Pl. 1006. 1753('gossipifolia'); B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Manihot gossypiifolia (L.) Crantz, Inst. Rei Herb. 1: 167.1766

Adenoropium gossypiifolium (L.) Pohl, Pl. Bras. Icon. Descr. 1: 16.1826

Distribution: Karnataka (naturalized throughout), naturalized throughout India; native to Brazil.

Local/Common/Vernacular names: *Chikka kadu haralu, Hathi yele haralu, Vilayati haralu* (Kannada)

Jatropha heynei N.P.Balakr., Bull. Bot. Surv. India 3: 40. 1961; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Jatropha heterophylla B.Heyne ex Hook.f., Fl. Brit. India 5: 382. 1887 non Steud. 1840

Distribution: Karnataka (Ballari, Chitradurga, Mysuru), Andhra Pradesh, Madhya Pradesh, Tamil Nadu; endemic.

Jatropha integerrima Jacq., Enum. Syst. Pl. 32. 1763

Jatropha integerrima var. *coccinea* (Link) N.P.Balakr., Bull. Bot. Surv. India 22: 176 1980

Jatropha integerrima var. *latifolia* (Pax) N.P.Balakr., Bull. Bot. Surv. India 22: 176. 1980

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Cuba.

Jatropha multifida L., Sp. Pl. 1006. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Distribution: Karnataka (cultivated in gardens throughout), cultivated throughout plains in India; native to tropical & subtropical America.

Jatropha podagraria Hook., Bot. Mag. 74: t. 4376. 1848; C.J.Saldanha, Fl. Karnataka 2: 149. 1996.

Distribution: Karnataka (cultivated throughout), cultivated ornamental throughout plains; native to C America.

Jatropha villosa Wight, Icon. Pl. Ind. Orient. 4: 1. 1848 non Baillon 1863; C.J.Saldanha, Fl. Karnataka 2: 149. 1996.

Jatropha peltata Wight, Icon. Pl. Ind. Orient. 4: t. 1169. 1848 non Cerv. 1794, nec Steud. 1840

Jatropha wightiana Muell.Arg. in DC., Prodr. 15: 1080. 1866

Jatropha villosa var. *ramnadensis* Ramam., Bull. Bot. Surv. India 9: 278, ff. 1 - 11. 1967

Distribution: Karnataka (Mysuru), Andhra Pradesh, Tamil Nadu; endemic.

Macaranga Thouars

Macaranga indica Wight, Icon. Pl. Ind. Orient. 5: t. 1883. 1852; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Trewia hernandifolia Roth, Nov. Pl. Sp. 374. 1821

Tanarius indicus (Wight) Kuntze, Revis. Gen. Pl. 2: 619.1891

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Bihar, Kerala, Madhya Pradesh, Meghalaya, Odisha, Sikkim, Tamil Nadu, Uttaranchal, West Bengal; China, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Bettadavare, Makaranda mara* (Kannada)

Macaranga peltata (Roxb.) Muell.Arg. in DC., Prodr. 15: 1010. 1866; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Osyris peltata Roxb., Fl. Ind. 3: 755. 1832

Mappa peltata (Roxb.) Wight, Icon. Pl. Ind. Orient. 4: t. 817. 1844

Macaranga flexuosa Wight, Icon. Pl. Ind. Orient. 5(2): 23. 1852

Macaranga roxburghii Wight, Icon. Pl. Ind. Orient. 6: t. 1949. 1853

Macaranga tomentosa Wight, Icon. Pl. Ind. Orient. 6: t. 1949. 1853

Macaranga wightiana Baill., Étude Euphorb. 432. 1858

Tanarius flexuosus (Wight) Kuntze, Revis. Gen. Pl. 2: 620. 1891

Tanarius peltatus (Roxb.) Kuntze, Revis. Gen. Pl. 2: 620. 1891

Tanarius roxburghii (Wight) Kuntze, Revis. Gen. Pl. 2: 619. 1891

Tanarius tomentosus (Wight) Kuntze, Revis. Gen. Pl. 2: 620. 1891

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Odisha, Sikkim, Tamil Nadu, West Bengal; Myanmar, Sri Lanka.

Local/Common/Vernacular names: *Chanda kanne*, *Chandrakala*, *Chenthakanni*, *Chinthakanni*, *Kanchupranti*, *Uppaligi* (Kannada); *Chanda* (*Kunabi*)

Mallotus Lour.

Mallotus aureopunctatus (Dalzell) Müll.Arg. in DC., Prodr. 15: 973. 1866; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Rottlera aureopunctata Dalzell in Hook.'s J.Bot. Kew Gard. Misc. 3:122. 1851

Mallotus lawii Muell.Arg. in Linnaea 34: 192. 1865

Distribution: Karnataka (Belagavi, Dakshina Kannada, Kodagu, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Mallotus beddomei Hook. f., Fl. Brit. India 5: 438. 1887; B.D.Sharma et al., Fl. Karnataka Anal. 253. 1984.

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Mallotus nudiflorus (L.) Kulju & Welzen, Blumea 52(1): 124. 2007

Trewia nudiflora L., Sp. Pl. 1193. 1753; C.J.Saldanha, Fl. Karnataka 2: 165. 1996

Tetragastris ossea Gaertn., Fruct. Sem. Pl. 2: 130. 1790

Rottlera indica Willd., Gött. J. Naturwiss. 1: 8. 1797

Trewia macrophylla Roth, Nov. Pl. Sp. 373. 1821

Trewia macrostachya Klotzsch, Bot. Ergebn. Reise Waldemar 117. 1862

Trewia nudiflora var. *tomentosa* Susila & N.P.Balakr., J. Econ. Taxon. Bot. 22: 351. 1998

Distribution: Karnataka (Hassan, Kodagu, Mysuru, Raichur, Shivamogga, Uttara Kannada); throughout hotter parts of India; Bangladesh, Indonesia, Malaysia, Myanmar, Sri Lanka.

Mallotus philippensis (Lam.) Muell.Arg. in Linnaea 34: 196. 1865; C.J.Saldanha, Fl. Karnataka 2: 152. 1996

Croton philippensis Lam., Encycl. 2: 206. 1786

Rottlera tinctoria Roxb., Pl. Coromandel 2: 36. 1802

Mallotus philippensis var. *tomentosus* Gamble, Fl. Madras 1322. 1925

Distribution: Karnataka (Throughout), throughout India; Australia, China, Indo-China, S & SE Asia.

Local/Common/Vernacular name: *Kumkumadamara* (Kannada)

Mallotus polycarpus (Benth.) Kulju & Welzen, Blumea 52: 130. 2007

Trewia polycarpa Benth., Gen. Pl. 3: 318. 1880

Trewia nudiflora var. *polycarpa* (Benth.) Susila & N.P.Balakr., J. Econ. Taxon. Bot. 22: 351. 1998

Distribution: Karnataka (Mysuru, Shivamogga), Goa, Kerala, Madhya Pradesh, Maharashtra; endemic.

Mallotus repandus (Rottler ex Willd.) Mull.Arg., Linnaea 34: 197. 1865 & in DC., Prodr. 981. 1866;
 B.D.Sharma et al., Fl. Karnataka Anal. 254. 1984.
Croton repandus Rottler ex Willd., Ges. Naturf. Freunde Berlin Neue Schriften 4: 206. 1803
Croton rhombifolius Willd., Sp. Pl. 4: 555. 1805
Rottlera scabrifolia A.Juss., Euphorb. Gen. 111. 1824
Rottlera viscosa Blume, Bijdr. 608. 1826
Rottlera trinervis Zipp. ex Span., Linnaea 15: 348. 1841
Rottlera rhombifolia (Willd.) Thwaites, Enum. Pl. Zeyl. 272. 1861
Mallotus repandus var. *scabrifolius* (A.Juss.) Müll.Arg., Prodr. 15(2): 982. 1866
Trewia nudiflora var. *dentata* Susila & N.P.Balakr., J. Econ. Taxon. Bot. 22: 352. 1998
 Distribution: Karnataka (throughout), throughout drier parts of India including Andaman & Nicobar Islands; Australia, Bangladesh, Bhutan, Brunei, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Nepal, New Caledonia, New Guinea, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam.

Mallotus resinosus (Blanco) Merr., Sp. Blancoan. 222. 1918; B.D.Sharma et al., Fl. Karnataka Anal. 254. 1984.

Adelia resinosa Blanco, Fl. Philip., ed. 2: 562. 1845
Claoxylon muricatum Wight, Icon. Pl. Ind. Orient. 5: 24, t. 1886. 1852
Axenfeldia intermedia Baill., Etude Euphorb. 419. 1858
Mallotus muricatus (Wight) Mull.Arg., Linnaea 34: 191. 1865
Mallotus stenanthus Mull.Arg., Linnaea 34: 191. 1865
Mallotus andamanicus Hook.f., Fl. Brit. India 5: 439. 1887
Mallotus walkerae Hook.f., Fl. Brit. India 5: 437. 1887
Mallotus walkerae var. *laxiflorus* Hook.f., Fl. Brit. India 5: 437. 1887

Mallotus muricatus var. *walkerae* (Hook.f.) Pax & K.Hoffm. in Engl., Pflanzenr. IV. 147. vii: 190. 1914
Mallotus intermedius (Baill.) N.P.Balakr., Bull. Bot. Surv. India 10: 245. 1968

Mallotus resinosus var. *subramanyamii* (J.L.Ellis) Chakrab., J. Econ. Taxon. Bot. 6: 704. 1985
Mallotus subramanyamii J.L.Ellis, Bull. Bot. Surv. India 25: 199. 1985 ('subramanyamili').
Mallotus resinosus var. *muricatus* (Wight) N.P.Balakr. & Chakrab., Rheedea 1: 39. 1991

Mallotus resinosus var. *stenanthus* (Mull.Arg.) Susila & N.P.Balakr. in N.P. Balakr. & Chakrab., Fam. Euphorb. India 151. 2007

Distribution: Karnataka (Belagavi, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; Australia, Brunei, China, Indonesia, Malaysia, Myanmar, New Guinea, Philippines, Sri Lanka, Thailand.

Mallotus rhamnifolius (Rottler ex Willd.) Mull.Arg., Linnaea 34: 196. 1865

Croton nervosus Rottler, Neue Schriften Ges. Naturf. Freunde Berlin 4: 190. 1803
Croton rhamnifolius Rottler ex Willd., Neue Schriften Ges. Naturf. Freunde Berlin 4: 190. 1803

Croton reticulatus Willd., Sp. Pl. 4: 545. 1805.

Mallotus micranthus Mull.Arg., Linnaea 34: 191. 1865

Mallotus zeylanicus Mull.Arg., Linnaea 34: 195. 1865

Oxydectes reticulata (Willd.) Kuntze, Revis. Gen. Pl. 2: 612. 1891

Distribution: Karnataka (Chamarajanagar?), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Mallotus tetracoccus (Roxb.) Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 41(2): 245. 1873

Rottlera ferruginea Roxb., Fl. Ind. 3: 828. 1832

Rottlera tetracocca Roxb., Fl. Ind. 3: 826. 1832

Rottlera peltata Wight, Icon. Pl. Ind. Orient. 5: t. 1873. 1852

Mallotus ferrugineus (Roxb.) Mull.Arg. in DC., Prodr. 15(2): 982. 1866

Mallotus albus var. *occidentalis* Hook. f., Fl. Brit. India 5: 429. 1887

Distribution: Karnataka (Belagavi, Kodagu, Shimoga, Uttara Kannada), Arunachal Pradesh, Assam, Goa, Kerala, Maharashtra, Meghalaya, Tamil Nadu, West Bengal; Bangladesh, China, Myanmar, Nepal, Sri Lanka, Thailand.

Local/Common/Vernacular names: *Kukbuta, Uppali mara* (Kannada)

Manihot Mill.

Manihot carthagenensis (Jacq.) Müll.Arg., Prodr. 15(2): 1073. 1866.

subsp. **glaziovii** (Müll.Arg.) Allem, Novon 11: 160 2001

Manihot glaziovii Muell.Arg. in Mart., Fl. Bras. 11: 471. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 254. 1984.

Distribution: Karnataka (cultivated in Ballari, Bengaluru, Hassan, Mysuru, Tumakuru), cultivated throughout in India; native to Brazil; cultivated throughout tropics.

Local/Common/Vernacular name: *Ceara rubber* (English)

Manihot esculenta Crantz., Inst. Rei. Herb. 1: 167. 1766; B.D.Sharma et al., Fl. Karnataka Anal. 254. 1984.

Jatropha manihot L., Sp. Pl. 1007. 1753

Janipha manihot (L.) Kunth, Nov. Gen. Sp. 2: 108. 1817

Manihot utilissima Pohl, Pl. Bras. Icon. Descr. 1: 55 1827, nom. superfl.

Manihot edulis A.Richard, Hist. Fis. Cuba, Bot. 2: 208. 1850

Distribution: Karnataka (cultivated for edible tuberous roots), cultivated Andaman & Nicobar Islands, Andhra Pradesh, Assam, Bihar, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Odisha, Sikkim, Tamil Nadu, Uttar Pradesh, West Bengal; native to Brazil, now widely cultivated in tropics.

Local/Common/Vernacular names: *Cassava*, *Tapioca* (English); *Kolgenasu* (Kannada)

Micrococca Benth.

Micrococca mercurialis (L.) Benth. in Hook., Niger Fl. 503. 1849; Haines, Bot. Bihar Orissa 2: 116. 1921.

Tragia mercurialis L., Sp. Pl. 980. 1753

Mercurialis alternifolia Lam., Encycl. 4: 120. 1797

Claoxylon mercurialis (L.) Thwaites, Enum. Pl. Zeyl. 271. 1861; Susila & N.P.Balakr., Rheedia 5(2): 131. 1995.

Microstachys mercurialis (L.) Dalzell & A.Gibson, Bombay Fl. 227. 1861

Distribution: Karnataka (Mysuru), Andhra Pradesh, Bihar, Goa, Himachal Pradesh, Haryana, Kerala, Lakshadweep, Madhya Pradesh, Maharashtra, Odisha, Puducherry, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Middle East, Myanmar, Sri Lanka, Thailand, tropical Africa.

Microstachys A.Juss.

Sebastiania Spreng.

Microstachys chamaelea (L.) Mull.Arg., Linnaea 32: 95. 1893

Tragia chamaelea L., Sp. Pl. 981. 1753; C.J.Saldanha, Fl. Karnataka 2: 162. 1996.

Cnemidostachys chamaelaea (L.) Spreng., Syst. Veg. 3: 835. 1826

Elachocroton asperococcum F.Muell., Hooker's J. Bot. Kew Gard. Misc. 9: 17. 1857

Cnemidostachys linearifolia Miq., Fl. Ned. Ind., Eerste Bijv. 460. 1861

Excoecaria chamaelea (L.) Baill., Adansonia 4: 323. 1864

Sebastiania chamaelea (L.) Mull.Arg. in DC., Prodr. 15(2): 1175. 1866

Stillingia chamaelea (L.) Müll.Arg. in Prodr. 15(2): 1175. 1866

Distribution: Karnataka (Ballari, Bengaluru, Chikkaballapur, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Andaman & Nicobar Islands, Peninsula to Bihar and W Bengal; paleotropical.

Paracroton Miq.

Paracroton integrifolius (Airy Shaw) N.P.Balakr. & Chakrab., Kew Bull. 48: 718. 1993

Plate 6G

Ostodes integrifolia Airy Shaw, Kew Bull. 16: 362. 1963

Fahrenheitia integrifolia (Airy Shaw) Airy Shaw, Kew Bull. 20: 410. 1966

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Paracroton pendulus (Hassk.) Miq., Fl. Ned. Ind. 1(2): 382 1859

Croton pendulus Hassk., Pl. Jav. Rar. 266 1848

subsp. **zeylanicus** (Thwaites) N.P.Balakr. & Chakrab., Kew Bull. 48: 719. 1993

Desmostemon zeylanica Thwaites, Enum. Pl. Zeyl. 278. 1861; C.J.Saldanha, Fl. Karnataka 2: 142. 1996

Ostodes zeylanica (Thwaites) Müll.Arg. in Linnaea 34: 214. 1865

Trigonostemon zeylanicus (Thwaites) Müll.Arg., Linnaea 34: 213. 1865

Fahrenheitia zeylanica (Thw.) Airy Shaw, Kew Bull. 20: 410. 1966

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Kolar, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Bale, Sotige* (Kannada)

Reutealis Airy Shaw

Reutealis trisperma (Blanco) Airy Shaw, Kew Bull. 20: 395. 1966

Aleurites trispermus Blanco, Fl. Filip. 755. 1837

Distribution: Karnataka (cultivated as an avenue tree in Bengaluru); native to the Philippines

Local/Common/Vernacular name: *Philippine tung* (English)

Ricinus L.

Ricinus communis L., Sp. Pl. 1007. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 256. 1984.

Croton spinosus L., Sp. Pl. 1005. 1753

Ricinus speciosus Burm.f., Fl. Indica 207. 1768

Ricinus spectabilis Blume, Bijdr. 623. 1826

Distribution: Karnataka (cultivated and naturalized throughout) cultivated in S India, naturalized throughout except temperate areas, Pantropical.; native of Africa.

Local/Common/Vernacular names: *Castor, Castor oil plant* (English); *Haralau, Oudala gida* (Kannada)

Suregada Roxb. ex Rottler

Suregada lanceolata (Willd.) Kuntze, Revis. Gen. Pl. 2: 619 1891

Gelonium lanceolatum Willd., Sp. Pl. 4: 832 1806; B.D.Sharma et al., Fl. Karnataka Anal. 251. 1984.

Suregada angustifolia Baill., Étude Euphorb. 396. 1858

Gelonium angustifolium Müll.Arg. in DC., Prodr. 15: 1128. 1866

Distribution: Karnataka (Bengaluru), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Tragia L.

Tragia bicolor Miq., Linnaea 26: 222 1854; B.D.Sharma et al., Fl. Karnataka Anal. 256. 1984.

Tragia miquelianiana Müll.Arg. in DC., Prodr. 15(2): 942. 1866

Tragia miquelianiana var. *bicolor* (Miq.) Mull.Arg. in DC., Prodr. 15(2): 942. 1866

Distribution: Karnataka (Chikkamagaluru, Shivamogga), Kerala, Tamil Nadu; endemic.

Tragia hildebrandtii Müll.Arg., Breman, Abh. 7: 26. 1880; C.J.Saldanha, Fl. Karnataka 2: 165. 1996
Tragia involucrata var. *cannabina* auct. non Müll.Arg., 1866; Saldanha in Saldanha, Fl. Karnataka 2: 165. 1996
 Distribution: Karnataka (Belagavi, Chitradurga, Dharwad, Kalaburagi, Mysuru, Raichur), Maharashtra; tropical Africa.
 Local/Common/Vernacular name: *Thindu toppalu gida* (Kannada)

Tragia hispida Willd., Sp. Pl. 4: 323. 1805; B.D.Sharma et al., Fl. Karnataka Anal. 256. 1984.
Tragia involucrata auct. non L., 1753; Duthie, Fl. Gangetic Plain 2: 114. 1905, p. p., non L. 1753.
Tragia involucrata var. *hispida* (Willd.) Mull.Arg. in DC., Prodr. 15(2): 943. 1866
 Distribution: Karnataka (throughout), Assam, Bihar, Kerala, Maharashtra, Uttarakhand, West Bengal; Bangladesh.
 Local/Common/Vernacular names: *Churuchangee soppu, Curca, Turike balli, Turike soppu* (Kannada)

Tragia involucrata L., Sp. Pl. 980. 1753; Hook.f., Fl. Brit. India 5: 465. 1888, p. p. excl. *Tragia hispida* Willd. & vars; Prain, Bengal Pl. 2: 952. 1903.
Croton urens L., Sp. Pl. 1005. 1753
Tragia involucrata var. *rheediana* Müll.Arg., Prodr. 15(2): 943. 1866
 Distribution: Karnataka (throughout), throughout India; Bangladesh, China, Myanmar, Sri Lanka.
 Local/Common/Vernacular names: *Indian stinging nettle* (English); *Antu sura surige, Dulagondi, Haligilu, Kiriberalu* (Kannada)

Tragia montana (Thwaites) Mull.Arg., Linnaea 34: 183. 1865 & in DC., Prodr. 15(2): 944. 1866; Chakrab. & N.P.Balakr., Rheedia 16: 20. 2006.
Tragia involucrata var. *montana* Thwaites, Enum. Pl. Zeyl. 270. 1861
Tragia involucrata var. *cordata* Mull.Arg., l. c. 943. 1866
Tragia muelleriana Pax & K.Hoffm., l. c. 80. 1919, excl. var. *unicolor*
Tragia muelleriana var. *unicolor* auct. non (Mull.Arg.) Pax & K.Hoffm., 1919; A.N.Londhe in N.P. Singh et al., Fl. Maharashtra State: Dicot 2: 904. 2001, p. p., excl. syn.
 Distribution: Karnataka (), Kerala, Maharashtra Tamil Nadu; Sri Lanka.
 Local/Common/Vernacular name: *Dulagondi* (Kannada)

Tragia pluknetii Radcl.-Sm., Kew Bull. 37: 688. 1983
Croton hastatus L., Sp. Pl. 1005. 1753.
Tragia cannabina L.f., Suppl. Pl. 415 1782; B.D.Sharma et al., Fl. Karnataka Anal. 256. 1984.
Tragia involucrata var. *cannabina* Müll.Arg., Prodr. 15(2): 944. 1866
Tragia cannabina var. *hastata* (L.) Pax & K.Hoffm., Pflanzenr. IV, 147, IX: 85. 1919
 Distribution: Karnataka (Ballari, Chitradurga, Dharwad, Kalaburagi, Kodagu, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Goa, Gujarat, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, West Bengal; Africa, Sri Lanka.
 Local/Common/Vernacular names: *Churachrike gida, Kiri turachi, Sanna turachi* (Kannada)

Tragia praetervisa Chakrab. & N.P.Balakr., Rheedia 16: 23 2006
Tragia miquelianoides var. *unicolor* Müll.Arg., Prodr. 15(2): 943. 1866
Tragia muelleriana var. *unicolor* (Müll.Arg.) Pax & K.Hoffm. in Engl., Pflanzenr. IV, 147, 9: 81. 1919; B.D.Sharma et al., Fl. Karnataka Anal. 257. 1984
Tragia praetervisa var. *unicolor* (Mull.Arg.) Chakrab. & N.P.Balakr., Rheedia 16: 25. 2006; N.P.Balakr. & Chakrab., Fam. Euphorb. India 184. 2007.
 Distribution: Karnataka (Ballari, Chikkamagaluru, Chitradurga, Kalaburagi, Kodagu, Tumakuru, Uttara Kannada), Goa, Kerala, Maharashtra, Mizoram, Tamil Nadu, West Bengal; Sri Lanka.

Vernicia Lour.

Vernicia cordata (Thunb.) Airy Shaw, Kew Bull. 20: 394.1966

Dryandra cordata Thunb., Nova Acta Regiae Soc. Sci. Upsal. 4: 38 1783

Elaeococca cordata (Thunb.) A.Juss, Euphorb. Gen. 38.1824

Dryandra verrucosa Royle, Ill. Bot. Himal. Mts. 328, 452.1836

Aleurites cordata R. Br. ex Steud., Nomencl. Bot. 1: 49. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 244. 1984.

Camerium cordatum (Thunb.) Kuntze, Revis. Gen. Pl. 2: 595.1891

Distribution: Karnataka (Chikkamagaluru), Sikkim, cultivated in S & SE Asia; native to Japan & China.

Local/Common/Vernacular names: *Japanese tung oil tree, Japanese wood-oil-tree* (English)

Vernicia fordii (Hemsl.) Airy Shaw, Kew Bull. 20: 394 1966

Aleurites fordii Hemsl., Hooker's Icon. Pl. 29: t. 2801 1906; C.J.Saldanha, Fl. Karnataka 2: 116. 1996.

Distribution: Karnataka (cultivated), Assam, Bihar, Odisha, Sikkim, West Bengal; native to China, Myanmar & Vietnam.

Local/Common/Vernacular names: *Candle nut tree, Chinese tung oil tree, Chinese wood-oil tree, Kalo nut, Kalo nut tree, Tung nut, Tung oil tree* (English)

Vernicia montana Lour., Fl. Cochinch. 586. 1790; Stuppy et al. Blumea 44: 92, f. 3.1999.

Elaeococca montana (Lour.) Oken, Allg. Naturgesch. 3(3): 1599.1841

Aleurites montana (Lour.) E.H. Wilson, Bull. Imp. Inst. Gr. Brit. 11: 460. 1913; Mooney, Suppl. Bot. Bihar Orissa 33. 1950.

Distribution: Karnataka (cultivated), cultivated in Arunachal Pradesh, Assam, Bihar, Meghalaya, Odisha, Sikkim, Tamil Nadu, West Bengal; native to China; cultivated in tropics.

Local/Common/Vernacular names: *Cantonese wood-oil tree, China wood oil, Hong Kong wood oil tree, Mu oil tree, Tung-oil tree* (English)

Flagellariaceae

Flagellaria L.

Flagellaria indica L., Sp. Pl. 333.1753; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Flagellaria indica var. *minor* Hook.f., Fl. Brit. India 6: 391.1892

Distribution: Karnataka (Dakshina Kannada, Kodagu, Udupi, Uttara Kannada), throughout coastal areas including Andaman & Nicobar Islands; paleotropical coasts.

Gentianaceae

Canscora Lam.

Canscora alata (Roth) Wall., Numer. List 4363.1831

Exacum alatum Roth, Syst. Veg. 3: 159.1818

Pladeria decussata Roxb., Fl. Ind.ed.1,1: 418.1820

Canscora decussata (Roxb.) Schult. & Schult.f., Mant. 3: 229.1827; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984.

Distribution: Karnataka (Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru); Peninsular India, Madagascar, tropical Africa.

Canscora diffusa (Vahl) R.Br. ex Roem. & Schult., Syst. Veg. 3: 301.1818; B.D.Sharma et al., Fl.

Karnataka Anal. 172. 1984.

Plate 8A

Gentiana diffusa Vahl, Symb. Bot. 3: 47.1794

Exacum diffusum (Vahl) Willd., Sp. Pl. 1: 637.1798

Canscora lawii Wight, Icon. Pl. Ind. Orient. 4: t. 1327.1848

Canscora tenella Wall. ex Wight, Icon. Pl. Ind. Orient. 4: t. 1327.1848

Canscora decurrens Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 136.1850; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984.

Canscora pauciflora Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 136.1850; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984.

Canscora stricta Sedgw., J. Indian Bot. 2: 126.1921

Canscora khandalensis Santapau, Kew Bull. 3: 485.1949

Canscora diffusa var. *tetraptera* Naik & Pokle, J. Econ. Taxon. Bot. 7: 673.1986

Canscora sanjappae Diwakar & R.Kr.Singh, Indian J. Forest. 32: 337.2009

Canscora devendrae Diwakar & R.Kr.Singh, Indian J. Forest. 34: 249.2011

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Kalaburgi, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Udupi, Uttara Kannada, Vijayapura, Yadgir), S & S E Asia, tropical Africa.

Local/Common/Vernacular name: *Shanka huli* (Kannada)

Canscora heteroclita (L.) Gilg, Nat. Pflanzenfam. 4(2): 76.1895; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984.

Gentiana heteroclita L., Mant. Pl. 2: 560.1771

Centaurium malabaricum Borkh., Arch. Bot. (Leipzig) 1(1): 29.1796

Exacum heteroclitum (L.) Willd., Sp. Pl. 1: 639.1798

Pladera sessiliflora Roxb., Fl. Ind. 1: 416.1820

Canscora sessiliflora (Roxb.) Schult., Mant. 3: 230.1827

Distribution: Karnataka (Bengaluru, Ballari, Kolar, Raichur, Tumakuru), Kerala, Tamil Nadu; Sri Lanka.

Canscora perfoliata Lam., Encycl. 1: 601.1785; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984

Plate 23C

Canscora ventricosa J.F.Gmel., Syst. Nat. 262.1791

Pootia triflora Dennst., Reis. Br.-Guiana 37.1818

Pladera alata Buch.-Ham. ex Wall., Numer. List 4364 B.1831

Canscora grandiflora Wight, Icon. Pl. Ind. Orient. 4: t. 1326.1848

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Centaurium Hill.

Centaurium centaurioides (Roxb.) R.S. Rao & Hemadri, J. Bombay Nat. Hist. Soc. 67: 357. 1970; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984.

Chironia centaurioides Roxb., Fl. Ind. 2: 283.1824

Chironia carinata Banks ex Roxb., Fl. Ind. ed.2, 1: 584.1832

Erythraea roxburghii G.Don, Gen. Hist. 4: 206.1837

Erythraea centaurioides (Roxb.) Voigt, Hort. Suburb. Calcutt. 520.1845

Centaurium meyeri auct. non (Bunge) Druce, 1917; N.M.G. Babu, Fl. Ethnobot. Chitradurga Dist. 366.2011.

Centaurium roxburghii (D.Don) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 614.1917

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Hassan, Kalaburgi, Mysuru, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout planis of India; endemic.

Enicostema Blume

Enicostema axillare (Poir. ex Lam.) A.Raynal, Adansonia n.s., 9: 75.1969

Gentiana verticillaris Retz., Observ. Bot. 2: 15.1781

Gentiana axillaris Poir. ex Lam., Tabl. Encycl. i. 487. 1793

Exacum hyssopifolium Willd., Sp. Pl. 1: 640.1798

Hippion hyssopifolium (Willd.) Spreng., Syst. Veg. 1: 589.1825

Slevogtia orientalis Griseb., Prodr. 9: 65.1845

Slevogtia maritima Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 137.1850

Hippion orientale (Griseb.) Dalzell & A.Gibson, Bombay Fl. 157.1861

Enicostema verticillare (Retz.) Baill., Hist. Pl. 10: 131.1888

Hippion verticillatum var. *hyssopifolium* (Willd.) Kuntze, Revis. Gen. Pl. 2: 428.1891

Hippion verticillatum var. *maritimum* (Dalzell) Kuntze, Revis. Gen. Pl. 2: 428.1891

Enicostema hyssopifolium (Willd.) Verd., Bothalia 7: 462.1961; B.D.Sharma et al., Fl. Karnataka Anal. 172. 1984.

Distribution: Karnataka (Belagavi, Ballari, Chitradurga, Dharwad, Kalaburgi, Mysuru, Raichur, Tumakuru, Uttara Kannada, Vijayapura, Yadgir), Andhra Pradesh, Maharashtra, Tamil Nadu; SE Asia, tropical Africa.

Local/Common/Vernacular name: *Bili ranjaka* (Kannada)

Exacum L.

Exacum atropurpureum Bedd., Icon. Pl. Ind. Or. 1: 24.1869; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Exacum atropurpureum var. *palghatense* Gamble, Fl. Madras 874.1935

Distribution: Karnataka (Uttara Kannada), Kerala, Tamil Nadu; endemic.

Exacum lawii C.B.Clarke in Hook.f., Fl. Brit. India 4: 98.1883; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984. Plate 23D

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Uttara Kannada), Kerala, Maharashtra; endemic.

Exacum pedunculatum L., Sp. Pl. 112.1753; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Exacum carinatum Roxb., Fl. Ind. 1: 415.1820; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Exacum sulcatum Roxb., Fl. Ind. 1: 415.1820

Sebaea carinata (Roxb.) Spreng., Syst. Veg. 1: 426.1825

Sebaea sulcatum (Roxb.) Spreng., Syst. Veg. 1: 426.1825

Exacum rivulare Edgew., J. Asiat. Soc. Bengal Pt. 2, Nat. Hist. 21 (2): 175.1853

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kalaburgi, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), throughout India except in Jammu & Kashmir; Malaysia, Myanmar, Sri Lanka.

Exacum pumilum Griseb., Prodr. 9: 46 1845; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Distribution: Karnataka (Belary, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kalaburgi, Kolar, Raichur, Uttara Kannada, Vijayapura), Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; endemic.

Exacum sessile C.B.Clarke, J. Linn. Soc., Bot. 14: 427.1875

Distribution: Karnataka (Dakshina Kannada, Hassan, Shivamogga, Udupi), Kerala, Tamil Nadu; Sri Lanka.

Exacum tetragonum Roxb., Fl. Ind. ed. 1, 1: 413.1820; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Exacum bicolor Roxb., Fl. Ind. ed. 1, 1: 413.1820; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Exacum grandiflorum Wall. ex G.Don, Gen. Hist. 4: 212.1837

Exacum stylosum Wall. ex G.Don, Gen. Hist. 4: 212.1837

Exacum tetragonum var. *roseum* Royle ex D.Don, Trans. Linn. Soc. London 17: 532.1837

Canscora justicioides Griff. ex Voigt, Hort. Suburb. Calcutt. 520.1845

Exacum perrottetii Griseb., Prodr. 9: 45.1845; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Exacum tetragonum var. *bicolor* (Roxb.) Hook., Bot. Mag. 73: t. 4340.1847

Exacum tetragonum var. *khasianum* C.B.Clarke, J. Linn. Soc., Bot. 14: 424.1875

Exacum metzianum Hohen. ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 96.1883

Exacum tetragonum var. *stylosum* (Wall. ex G.Don) C.B.Clarke in Hook.f., Fl. Brit. India 4: 95.1885

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout tropical and subtropical areas of India; Australia, Cambodia, China, Laos, Malaysia, Myanmar, Nepal, New Guinea, Philippines, Sikkim, Vietnam.

Exacum wightianum Arn., Ann. Sci. Nat., Bot. 2, 11: 176.1839; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Lisanthus zeylanicus Spreng., Syst. Veg. 1: 586.1824

Exacum foliosum Griseb., Prodr. 9: 45.1845

Distribution: Karnataka (Uttara Kannada), Kerala, Tamil Nadu; endemic.

Fagraea Thunb.

Fagraea ceilanica Thunb., Kongl. Vetensk. Acad. Nya Handl. 3: 132. 1782

Fagraea malabarica Blume, Rumphia 2: 29.1838

Fagraea coromandelina Wight, Icon. Pl. Ind. Orient. 4: t. 1316.1848

Gardenia heteroclyta J.Koenig ex Blume, Mus. Bot. 1: 164.1850

Fagraea khasiana Benth., J. Proc. Linn. Soc., Bot. 1: 99.1857

Fagraea prainii Gand., Bull. Soc. Bot. France 65: 58.1918

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Ginnannu* (Kannada)

Gentiana L.

Gentiana pedicellata (D.Don) Wall., Numer. List 4394.1831; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Gentiana orbiculata B.Heyne ex Wall., Numer. List 4395.1831

Gentiana pumila Griff., Ic. Pl. Asiat. t. 384.1854

Gentiana quadrifaria var. *pilosula* C.B.Clarke in Hook.f., Fl. Brit. India 4: 112.1883

Gentiana pedicellata var. *wallichii* Kusn., Trudy Imp. St.-Peterburgsk. Bot. Sada 15: 402.1898

Gentiana quadrifaria var. *zeylanica* (Griseb.) Kusn., Trudy Imp. S.-Peterburgsk. Bot. Sada 15: 403.1898

Gentianodes pedicellata (D.Don) Omer, Ali & Qaiser, Pakistan J. Bot. 20: 14.1988

Distribution: Karnataka (Mysuru), Kerala, Himalayas of India; Bhutan, China, Myanmar, Nepal, Pakistan.

Gentiana quadrifaria Blume, Bijdr. 847.1826; B.D.Sharma et al., Fl. Karnataka Anal. 173. 1984.

Gentiana pedicellata var. *wightii* Kusn., Trudy Imp. S.-Peterburgsk. Bot. Sada 15: 401.1904

Varasia quadrifaria (Blume) Sojak, Nar. Mus., Odd. P?ír. 148: 202.1980

Gentiana quadrifaria var. *wightii* (Kusn.) T.N.Ho, Acta Phytotax. Sin. 38: 11.2000.

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; China Myanmar, Sri Lanka.

Hoppea Willd.

Hoppea dichotoma Willd., Neue Schriften Ges. Naturf. Freunde Berlin 3: 435.1801; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Pladera pusilla Roxb., Fl. Ind. ed. 1832 1: 403.1832

Cicendia roxburghii Griseb., Gen. Sp. Gent. 160.1838

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Kalaburgi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), Andhra Pradesh, Bihar, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, West Bengal; endemic.

Hoppea fastigiata (Griseb.) C.B.Clarke in Hook.f., Fl. Brit. India 4: 100.1883; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Cicendia fastigiata Griseb., Gen. Sp. Gent. 158.1838

Exacum fastigiatum Arn. ex Griseb., Gen. Sp. Gent. 158.1838

Pladera pusilla Thwaites, Enum. Pl. Zeyl. 204. 1860.

Pladera fastigiata C.B.Clarke, J. Linn. Soc., Bot. 14: 429.1875

Exacum concinnum Miq. ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 100.1883

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu; Sri Lanka.

Swertia L.

Swertia angustifolia Buch.-Ham. ex D. Don, Prodr. Fl. Nepal. 127.1825

Ophelia pulchella D.Don, London Edinburgh Philos. Mag. & J. Sci. 8: 77.1836

Ophelia elegans Wight, Icon. Pl. Ind. Orient. 4: t. 1331.1848

Swertia affinis C.B.Clarke in Hook.f., Fl. Brit. India 4(10): 126.1883

Swertia pulchella var. *elegans* (Wight) C. B. Clarke in Hook. f., Fl. Brit. India 4:125.1883

Swertia angustifolia var. *hamiltoniana* Burkhill, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 2: 374.1906

Swertia angustifolia var. *pulchella* (D.Don) Burkhill, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 2: 375.1906; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Distribution: Karnataka (Ballari, Chamarajanagar, Chikkamagaluru), Andhra Pradesh, Kerala, Jammu & Kashmir, Sikkim, Tamil Nadu; Bhutan, China, Myanmar, Nepal.

Swertia beddomei C.B.Clarke in Hook.f., Fl. Brit. India 4: 127.1883; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada), Kerala, Tamil Nadu; endemic.

Swertia corymbosa Wight ex Griseb., Gen. Sp. Gent. 317.1839; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Ophelia corymbosa Griseb., Gen. Sp. Gent. 317.1839

Ophelia grisebachiana Wight, Icon. Pl. Ind. Orient. 4: t. 1330.1848

Ophelia pauciflora Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 211.1851

Ophelia umbellata C.B.Clarke, J. Linn. Soc., Bot. 14: 446.1875

Distribution: Karnataka (Chikkamagaluru, Hassan, Kalaburgi, Kodagu, Mysuru, Shivamogga, Uttara Kannada, Yadgir), Kerala, Tamil Nadu; endemic.

Swertia decussata Nimmo ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 127.1883

Ophelia densifolia Griseb., Prodr. 9: 125.1845

Ophelia alba Wight & Arn., Ill. Ind. Bot. 2: t. 157.1850

Ophelia multiflora Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 135.1850

Swertia densifolia (Griseb.) Kashyapa, Kew Bull. 15: 42.1961; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala, Maharashtra, Tamil Nadu; endemic.

Swertia lawii Burkil, J. Proc. Asiat. Soc. Bengal 2: 379. 1906

Plate 23E

Ophelia lawii Wight & Arn., Ill. Ind. Bot. 2: 175.1850

Distribution: Karnataka (Belgaum, Shivamogga, Udupi), Kerala; endemic.

Swertia minor Knobl., Abh. Königl. Akad. Wiss. Berlin 1891: 338 1892; B.D.Sharma et al., Fl. Karnataka Anal. 174. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Mysuru, Shivamogga), Kerala, Tamil Nadu; endemic.

Swertia raveendrae Shahina & Nampy, Phytotaxa 195 (1):44.2015

Distribution: Karnataka (Chikkamagaluru); endemic.

Swertia trichotoma Wall. ex C. B. Clarke in Hook. f., Fl. Brit. India 4: 126. 1883

Ophelia trichotoma Wight & Arn., Ill. Ind. Bot. 2: t. 157b.1850

Distribution: Karnataka (Chamarajanagar), Kerala, Tamil Nadu; endemic.

Gesneriaceae

Aeschynanthus Jack.

Aeschynanthus perrottetii A.DC. in DC., Prodr. 9: 261.1845; B.D.Sharma et al., Fl. Karnataka Anal.

196. 1984

Plate 23F

Aeschynanthus grandiflorus J.Graham, Cat. Pl. Bombay 146.1839

Aeschynanthus ceylanicus Wight, Icon. Pl. Ind. Orient. 4: t. 1347.1848

Trichosporum perrottetii Kuntze, Revis. Gen. Pl. 2: 478.1891

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Didymocarpus Wall. ex Buch.-Ham.

Didymocarpus subalternans Wall. ex R.Br., Pl. Jav. Rar. 118.1840; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Distribution: Karnataka (Shivamogga), W Himalaya; Nepal.

Epithema Blume

Epithema carnosum Benth., Scroph. Ind. 57.1835; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Aikinia carnosa G.Don, Gen. Hist. 4: 665.1838

Distribution: Karnataka (Shivamogga, Uttara Kannada), Andhra Pradesh, Maharashtra, Meghalaya, Odisha, Tamil Nadu, Uttaranchal, W Bengal; Bhutan, China, Laos, Myanmar, Nepal, Thailand.

Epithema ceylanicum Gardner, Calcutta J. Nat. Hist 6: 492.1846

Epithema carnosum var. *ceylanicum* (Gardner) C.B.Clarke in A.DC. & C.DC. Monogr. Phan.5(1): 178.1883

Epithema carnosum var. *dentatum* C.B.Clarke in A.DC. & C.DC. in A.DC. & C.DC., Monogr. Phan. 5(1): 177.1883

Epithema carnosum var. *hispidum* C.B.Clarke in A.DC., Monogr. Phan. 5: 178.1883; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Epithema dentatum (C.B.Clarke) Hilliard & B.L.Burtt, Edinburgh J.Bot. 54: 111.1997

Epithema dentatum subsp. *hispidum* (C.B.Clarke) Hilliard & B.L.Burtt, Edinburgh J. Bot. 54: 112 1997

Distribution: Karnataka (Belgavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, Goa, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil nadu, W Bengal; Cambodia, Myanmar, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam.

Epithema pusillum (C.B.Clarke) Bransgrove, Gard. Bull. Singapore 67(1): 202. 2015

Epithema carnosum var. *pusillum* C.B.Clarke in A.DC., Monogr. Phan. 5: 178.1883; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Epithema dentatum var. *pusillum* (C.B.Clarke) Hilliard & B.L.Burtt, Edinburgh J. Bot. 54: 112.1997

Distribution: Karnataka (Shivamogga), Maharashtra-Ratnagiri; endemic.

Gloxinia L.Herit.

Gloxinia perennis (L.) Druce, Rep. Bot. Exch. Club Soc. Brit. Isles 3: 418.1914; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Martynia perennis L., Sp. Pl. 618 1753

Gloxinia maculata L'Hér., Hortus Kew. 2: 331.1789

Distribution: Karnataka (Kodagu), cultivated in cooler parts of India; native to C & S America.

Henckelia Spreng.

Henckelia incana (Vahl) Spreng., Syst. Veg. 1: 38.1824

Rottlera incana Vahl, Enum. Pl. 1: 88.1804

Didymocarpusrottlerianus Wall. ex R.Br., Cyrtandreae 119.1839

Gratiola montana Rottler ex DC., Prodr. 9: 268.1845

Didymocarpus tomentosus Wight, Icon. Pl. Ind. Orient. 4: t. 1349.1848; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chitradurga, Tumakuru), Kerala, Tamil Nadu; S & SE Asia.

Henckelia innominata (B.L.Burtt) A.Weber & B.L.Burtt, Beitr. Biol. Pflanzen 70: 347.1998

Didymocarpus innominatus B.L.Burtt, Notes Roy. Bot. Gard. Edinburgh 21: 201. 1954; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Distribution: Karnataka (Chitradurga), Kerala, Tamil Nadu; endemic.

Jerdonia Wight

Jerdonia indica Wight, Icon. Pl. Ind. Orient. 4: t. 1352.1848; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Plate 23G

Distribution: Karnataka (Hassan, Kodagu, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Microchirita (C.B.Clarke) Yin Z.Wang

Microchirita hamosa (R.Br.) Yin Z.Wang, J. Syst. Evol. 49: 60.2011

Chirita hamosa R.Br., Cyrtandreae 117.1839; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Didymocarpus cristatus Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 225.1851

Chirita hamosa var. *finlaysonii* C.B.Clarke in A.DC., Monogr. Phan. 5: 128.1883

Chirita hamosa var. *unifolia* C.B.Clarke in A.DC., Monogr. Phan. 5: 129.1883

Roettlera hamosa (R.Br.) Kuntze, Revis. Gen. Pl. 2: 475.1891

Chirita cristata (Dalzell) B.L.Burtt, Notes Roy. Bot. Gard. Edinburgh 23: 96.1960

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Mysuru, Shivamogga, Uttara Kannada), peninsular India, Bihar; China, Laos, Malaysia, Thailand, Vietnam.

Microchirita sahyadriensis (Punekar & Lakshmin.) A.Weber & D. J. Middleton, Taxon 60: 779. 2011; Punekar & Lakshmin., Fl. Anshi Natl. Park 338. 2011.

Chirita sahyadriensis Punekar & Lakshmin., Folia Malaysiana 10: 18.2009

Distribution: Karnataka (Uttara Kannada); endemic.

Rhynchoglossum Blume

Rhynchoglossum notonianum (Wall.) B.L. Burtt, Notes Roy. Bot. Gard. Edinburgh 24(2): 170 .1962; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Wulfenia notoniana Wall., Tent. Fl. Napal. pt. 2: 46. 1826

Klugia notoniana A.DC. in DC., Prodr. 9: 276.1845

Distribution: Karnataka (Chamarajanagar, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Maharashtra; S Asia.

Rhynchoglossum obliquum Blume, Bijdr. 741.1826; B.D.Sharma et al., Fl. Karnataka Anal. 197. 1984.

Wulfenia obliqua Wall., Tent. Fl. Napal. 45, t. 35.1826

Antonia obliqua (Blume) R.Br., Pl. Asiat. Rar. 3: 65.1832

Loxotis obliqua (Blume) R.Br., Pl. Jav. Rar. 102.1838

Rhynchoglossum obliquum var. *parviflorum* C.B.Clarke in A. DC., Monogr. Phan. 5: 162.1883

Distribution: Karnataka (Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Goa, Maharashtra; S & SE Asia.

Rhynchotechum Blume

Rhynchotechum permolle (Nees) B.L. Burtt, Notes Roy. Bot. Gard. Edinburgh 24: 39.1962

Isanthera permollis Nees, Trans. Linn. Soc. London 17(1): 82.1834

Distribution: Karnataka (Chikkamagaluru, Hassan, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Gisekiaceae

Gisekia L.

Gisekia pharnaceoides L., Mant. Pl. 2: 562. 1771; B.D.Sharma et al., Fl. Karnataka Anal. 117. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannada, Mysuru, Tumakuru, Vijayapura), Andhra Pradesh, Delhi, Gujarat, Maharashtra, Odisha, Punjab, Rajasthan, Tamil NaduUttar Pradesh; paleotropical

Goodeniaceae

Scaevola L.

Scaevola plumieri (L.) Vahl, Symb. Bot. 2: 36.1791

Lobelia plumieri L., Sp. Pl. 929.1753

Scaevola lobelia L., Syst. Veg. ed. 13 2: 178.1774

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada)), throughout coastal areas; pantropical coasts.

Scaevola taccada (Gaertn.) Roxb., Hort. Bengal. 15.1814

Scaevola sericea Vahl, Symb. Bot. 2: 37.1791

Scaevola koenigii Vahl, Symb. Bot. 3: 36.1794

Scaevola lobelia var. *sericea* (Vahl) Benth., Hooker's J. Bot. Kew Gard. Misc. 4: 297.1852

Scaevola frutescens var. *sericea* (Vahl) Merr., Philipp. J. Sci. 7(5): 354.1912

Distribution: Karnataka (Dakshina Kannada, Udupi, Uttara Kannada), throughout coastal areas; pantropical coasts

Haloragaceae

Haloragis J. R. Forst. & G. Forst.

Haloragis oligantha Wight & Arn., Prodr. Fl. Ind. Orient. 1: 338. 1834; C.J.Saldanha, Fl. Karnataka 2: 8. 1996

Myriophyllum intermedium auct non DC. 1828

Myriophyllum oliganthum (Wight & Arn.) F.Muell., Fragm. Phyt. Austr. 10: 54. 1876

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkaballapura, Chikkamagaluru, Dharwad, Hassan, Kalaburgi, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada), Kerala, Meghalaya, Tamil Nadu; Australia, Malaysia, Myanmar.

Myriophyllum L.

Myriophyllum tuberculatum Roxb., Hort. Bengal. 12. 1814; B.D.Sharma et al., Fl. Karnataka Anal. 98. 1984.

Myriophyllum spathulatum Blatter & Hallb., J. Ind. Bot. Soc. 2: 44. 1921

Distribution: Karnataka (Mysuru, Uttara Kannada), Assam, Bihar, Kerala, Madhya Pradesh, Maharashtra, Manipur, Odisha, Tripura, West Bengal; Australia, Indonesia, Malaysia,

Heliconiaceae

Heliconia L.

Heliconia angusta Vell., Fl. Flumin. 3: 106, t. 20.1829

Heliconia bicolor Benth., Botanist 3: t. 101.1839

Heliconia angustifolia Hook., Bot. Mag. 75:t. 4475.1849

Bihai angustifolia (Hook.) Kuntze, Revis. Gen. Pl. 2: 685.1891

Bihai angusta (Vell.) Griggs, Bull. Torrey Bot. Club 42: 327.1915

Distribution: Karnataka (Belagavi, Bengaluru, Kodagu), cultivated as ornamental throughout India; native to Brazil.

Heliconia bihai (L.) L., Mant. Pl. 2: 211 1771

Musa bihai L., Sp. Pl. 1043.1753

Bihai bihai (L.) Griggs, Bull. Torrey Bot. Club 31: 445.1904

Distribution: Karnataka (Belagavi, Bengaluru, Kodagu), cultivated as ornamental throughout India; native to Northern S America & West Indies.

Heliconia psittacorum L.f., Suppl. Pl. 158.1782; K. G. Bhat, Fl. S. Kanara 140.2014.

Distribution: Karnataka (cultivated as ornamental throughout), cultivated as ornamental throughout India; native to Brazil

Heliconia rostrata Ruiz & Pavon, Fl. Peruv. 3: 71 1802; K. G. Bhat, Fl. S. Kanara 140.2014.

Bihai rostrata (Ruiz & Pav.) Griggs, Bull. Torrey Bot. Club 31: 445.1904

Distribution: Karnataka (cultivated as ornamental throughout), cultivated as ornamental throughout India; native to Peru, Bolivia, Columbia & Ecuador.

Hernandiaceae**Gyrocarpus** Jacq.

Gyrocarpus americanus Jacq., Select. Stirp. Amer. Hist. 282.1763; B.D.Sharma et al., Fl. Karnataka Anal. 235. 1984.

Gyrocarpus jacquinii Gaertn., Fruct. Sem. Pl. 2: 92.1790

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Davenegere, Dharwad, Kalaburgi, Mysuru, Raichur, Tumakuru, Vijayapura), throughout drier parts of India; pantropical.

Local/Common/Vernacular names: *Helicopter Tree, Propeller Tree, Shitwood, Stinkwood, Whirly Whirly Tree* (English); *Bendoli, Gali mara, Guthi mara* (Kannada)

Hydatellaceae**Trithuria** Hook. f.

Trithuria konkanensis S.R.Yadav & Janarth., Rheedea 4: 18.1994

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Goa, Maharashtra; endemic.

Hydrocharitaceae**Blyxa** Nronha ex Thouars

Blyxa aubertii Rich., Mem. Cl. Sci. Math. Inst. Natl. France 1811(2): 19.1814

Blyxa ceylanica Hook.f. in Hook.f., Fl. Brit. India 5: 661.1888

Distribution: Karnataka (Belagavi, Hassan, Mysuru), throughout India; pantropical.

Blyxa echinosperma (C.B.Clarke) Hook.f., Fl. Brit. India 5: 661.1888; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Hydrotrophus echinospermus C.B.Clarke, J. Linn. Soc., Bot. 14: 8.1873

Blyxa talbotii Hook.f., Fl. Brit. India 5: 661.1888

Blyxa echinospermoides Blatt., J. & Proc. Asiat. Soc. Bengal 26: 354 1931

Blyxa aubertii var. *echinosperma* (C.B.Clarke) C.D.K.Cook & Luond, Aquatic Bot. 15: 14. 1983

Distribution: Karnataka (Belagavi, Bengaluru, Dakshina Kannad, Davanagere, Hassan, Shivamogga, Uttara Kannada), Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu; Australasia, Indo-China, E, S & SE Asia.

Local/Common/Vernacular name: *Jala shulaka* (Kannada)

Blyxa mangalensis K.Rashmi & G.Krishnak. Kew Bull. 72(1)-2: 1. 2016

Distribution: Karnataka (Dakshina Kannada), Kerala; endemic

Blyxa octandra (Roxb.) Planch. ex Thwaites, Enum. Pl. Zeyl. 332. 1864; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Vallisneria octandra Roxb., Pl. Coromandel 2: 34.1802

Blyxa roxburghii Rich., Mem. Cl. Sci. Math. Inst. Natl. France 1811(2): 77.1814

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Shivamogga, Udupi, Uttara Kannada), throughout plains; Australia, Bangladesh, China, Myanmar, Papua New Guinea, Sri Lanka, Vietnam.

Halophila Thours

Halophila beccarii Asch., Nuovo Giorn. Bot. Ital. 3: 302.1871; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Distribution: Karnataka (Dakshina Kannad), S India; China, Indonesia, Malaysia, Myanmar, Philippines, Sri Lanka, Taiwan, Vietnam.

Hydrilla Rich.

Hydrilla verticillata (L.f.) Royle, Ill. Bot. Himal. Mts. t. 376.1839; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Serpicula verticillata L.f., Suppl. Pl. 416.1782

Vallisneria verticillata (L.f.) Roxb., Fl. Ind. 2, 3: 751 1832

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Kalburgi, Hassan, Kodagu, Mysuru, Tumakuru, Udupi, Uttara Kannada, Vijayapura, Yadgir), throughout plains; paleotropical.

Local/Common/Vernacular names: *Pachi, Pachi gida* (Kannada)

Najas L.

Najas graminea Delile, Descr. Egypte, Hist. Nat. 282.1813; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Najas seminuda Griff. ex Voigt, Hort. Suburb. Calcutt. 694.1845

Caulinia graminea (Delile) Tzvelev, Novosti Sist. Vyssh. Rast. 13: 20.1976

Distribution: Karnataka (Bengaluru, Chitradurga, Dakshina Kannada, Dharwad, Mandya, Mysuru, Udupi, Uttara Kannada), throughout plains of India; Europe, Eurasia, pantropical.

Local/Common/Vernacular name: *Rice field nymph* (English)

Najas indica (Willd.) Cham., Linnaea 4: 501.1829; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Caulinia indica Willd., Mém. Acad. Roy. Sci. Hist. (Berlin) 1798: 89.1801

Najas minor var. *indica* (Cham.) A.Braun, J. Bot. 2: 278.1864

Najas lacerata Rendle, Trans. Linn. Soc. London, Bot. 5: 416.1899

Distribution: Karnataka (Hassan, Kodagu, Mysuru), throughout India; throughout tropical and subtropical Asia.

Najas minor All., Auct. Syn. Stirp. Horti Taur. 3.1773; B.D.Sharma et al., Fl. Karnataka Anal. 300. 1984.

Fluvialis minor (All.) Pers., Syn. Pl. 2: 530.1807

Ittnera minor (All.) C.C.Gmel., Fl. Bad. 3: 592.1808

Najas dichotoma Roxb., Fl. Ind. ed. 1832 3: 749.1832

Caulinia minor (All.) Coss. & Germ., Fl. Descr. Anal. Paris 575.1845

Najas ternata Roxb. ex Griff., Not. Pl. Asiat. 3: 183.1851

Distribution: Karnataka (Bengaluru), throughout plains of India; Europe, Eurasia, pantropical.

Nechamandra Planch.

Nechamandra alternifolia (Roxb. ex Wight) Thwaites, Enum. Pl. Zeyl. 332 1864; B.D.Sharma et al., Fl. Karnataka Anal. 264. 1984.

Vallisneria alternifolia Roxb. ex Wight, Bot. Misc. 2: 344. 1831

Nechamandra roxburghii Planch, Ann. Sci. Nat., Bot. III, 11: 78.1849

Lagarosiphon roxburghii (Planch.) Benth., Gen. Pl. 3: 451.1883

Lagarosiphon alternifolia (Roxb. ex Wight) Druce, Bot. Soc. Exch. Club Brit. Isles 4: 630.1917

Distribution: Karnataka (Mandya, Mysuru, Tumakuru) peninsula to NE India; Bangladesh, China, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Ottelia Pers.

Ottelia alismoides (L.) Pers., Syn. Pl. 1: 400.1805; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Stratiotes alismoides L., Sp. Pl. 535.1753

Damasonium indicum Willd., Sp. Pl. 2: 276.1799

Damasonium alismoides (L.) R.Br., Prodr. Fl. Nov. Holl. 344.1810

Ottelia indica (Willd.) Planch. ex Dalzell & A.Gibson, Bombay Fl. 278.1861

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Kalburgi, Dakshina Kannad, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Vijayapura, Yadgir), throughout plains of India; pantropical.

Local/Common/Vernacular names: *Hasiru niru patre*, *Kotta genasu* (Kannada)

Vallisneria L.

Vallisneria spiralis L., Sp. Pl. 1015. 1753; B.D.Sharma et al., Fl Karnataka Anal. 264. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chitradurga, Davanagere, Dharwad, Kalburgi, Hassan, Mandya, Mysuru, Shivamogga, Tumakuru, Uttara Kannada. Vijayapura), throughout India; paleotropical.

Local/Common/Vernacular names: *Halasarpi*, *Hasiru ribbon gida*, *Kudure balada gida* (Kannada)

Hydroleaceae**Hydrolea L.**

Hydrolea zeylanica (L.) Vahl, Symb. Bot. 2: 46.1791; B.D.Sharma et al., Fl. Karnataka Anal. 175. 1984.

Nama zeylanica L., Sp. Pl. 226.1753

Steris aquatica Burm. f., Fl. Indica 73, t. 39, f. 3.1768

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannad, Kodagu, Mysuru, Udupi, Uttara Kannada), throughout India; tropical Asia.

Hypericaceae**Hypericum L.**

Hypericum humifusum L., Sp. Pl. 785.1753

Holosepalum humifusum (L.) Fourr., Ann. Soc. Linn. Lyon n.s., 16: 352. 1868

Hypericum rubrum Wight ex Dyer in Hook.f., Fl. Brit. India 1: 255.1874

Distribution: Karnataka (), Kerala, Tamil Nadu; Europe, tropical Africa

Local/Common/Vernacular name: *Trailing St John's wort* (English)

Hypericum japonicum Thunb., Syst. Veg. ed. 14 702 1784; B.D.Sharma et al., Fl. Karnataka Anal. 17. 1984.

Hypericum pusillum Choisy, Prodr. Monogr. Hypéric. 50.1821

Brathys japonica (Thunb.) Wight, Icon. Pl. Ind. Orient. 1: 113.1839

Hypericum calycatum Jacquem. ex Dyer in Hook.f., Fl. Brit. India 1: 256.1874

Hypericum japonicum var. *major* Fyson, Fl. South Ind. Hill Stations 1: 49. 1932

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India; Australia, Bhutan, Cambodia, China, Indonesia, Japan, S Korea, Laos, Malaysia, Myanmar, Nepal, New Zealand, Philippines, Sri Lanka, Thailand, Vietnam.

Hypericum mysurense Wall. ex Wight & Arn., Prodr. Fl. Ind. Orient. 99 1934; B.D.Sharma et al., Fl.

Karnataka Anal. 17. 1984.

Norysca mysurensis (Wall. ex Wight & Arn.) Wight, Icon. Pl. Ind. Orient. 1: t. 56 1838

Distribution: Karnataka (Bengaluru, Chikkaballapura, Chikkamagaluru, Hassan, Kodagu, Mysuru), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Chinnadaavare gida* (Kannada)

Hypericum wightianum Wall. ex Wight & Arn., Prodr. Fl. Ind. Orient. 1:99.1834; C.J.Saldanha, Fl. Karnataka 1: 209. 1984

Distribution: Karnataka (Chikkamagaluru), Arunachal Pradesh, Assam, Jammu & Kashmir, Kerala, Manipur, Meghalaya, Nagaland, Sikkim, Tamil Nadu, Uttar Pradesh, West Bengal; Bhutan, China, Laos, Myanmar, Sri Lanka, Thailand.

Hypoxidaceae

Curculigo Gaertn.

Curculigo orchoides Gaertn., Fruct. Sem. Pl. 1: 63.1788; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Curculigo malabarica Wight, Icon. Pl. Ind. Orient. t. 2043.1853

Hypoxis orchoides (Gaertn.) Kurz, Ann. Mus. Bot. Lugduno-Batavi 4: 177.1869

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Kalburgi, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Yadgir), from Peninsular to NE India; SE Asia.

Local/Common/Vernacular names: *Nela tale*, *Nela tengu*, *Neladali*, *nelatati gadde* (Kannada); *Nelamanda* (Tulu)

Hypoxis L.

Hypoxis aurea Lour., Fl. Cochinch. 200. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Curculigo graminifolia Nimmo, Cat. Pl. Bombay 215.1839

Distribution: Karnataka (Belagavi, Bengaluru, Chamarajanagar, Chikkamagaluru, Hassan, Shivamogga, Uttara Kannada), peninsular to NE india; S & SE Asia.

Molineria Colla

Molineria capitulata (Lour.) Herb., Amaryllidaceae 84.1837

Leucojum capitatum Lour., Fl. Cochinch. 199.1790

Curculigo capitulata (Lour.) Kuntze, Revis. Gen. Pl. 2: 703.1891

Distribution: Karnataka (cultivated in Belagavi Dakshina Kannada, Udupi), native to E, S & SE Asia & Australasia.

Molineria trichocarpa (Wight) N.P.Balakr., J. Bombay Nat. Hist. Soc. 63: 330.1967; B.D.Sharma et al., Fl. Karnataka Anal. 282. 1984.

Hypoxis brachystachya Wight, Icon. Pl. Ind. Orient. 4:t. 2046.1853

Hypoxis latifolia Wight, Icon. Pl. Ind. Orient. 4:t. 2044.1853

Hypoxis leptostachya Wight, Icon. Pl. Ind. Orient. 4:t. 2045.1853

Hypoxis pauciflora Wight, Icon. Pl. Ind. Orient. 4:t. 2046.1853

Hypoxis trichocarpa Wight, Icon. Pl. Ind. Orient. 4:t. 2045.1853

Molineria finlaysoniana Baker, J. Linn. Soc., Bot. 17: 121.1878

Curculigo finlaysoniana (Baker) Wall. ex Hook.f., Fl. Brit. India 6: 279.1892

Curculigo leptostachya (Wight) Alston, Kandy Fl. 72.1938

Molineria leptostachya (Wight) Abeyw., Ceylon J. Sci., Biol. Sci. 2: 143.1959

Curculigo trichocarpa (Wight) Bennet & Raizada, Indian J. Forest. 4: 68.1981

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Shivamogga, Tumakuru), Kerala, Tamil Nadu; Sri Lanka.

Icacinaceae**Apodytes E.Mey. ex Arn.**

Apodytes dimidiata E.Mey. ex Arn. in Hook.'s J. Bot. 3: 155. 1840; C.J.Saldanha, Fl. Karnataka 2: 104. 1996.

Plate 8B

Apodytes acutifolia ARich, Tent. Fl. Abyss. 1: 92. 1847

Apodytes beddomei Mast. in Hook. f., Fl. Brit. India 1: 588. 1875

Apodytes dimidiata var. *acutifolia* (A.Rich.) Boutique, SFl. Congo Belge 9: 274. 1960

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Assam, Kerala, Meghalaya, Mizoram, Tamil Nadu; China, Indonesia, Malaysia, Myanmar, Philippines, Sri Lanka, Thailand; tropical and subtropical Africa.

Local/Common/Vernacular name: *White Pear* (English); *Kainka* (Kannada)

Miquelia Meisn.

Miquelia dentata Bedd., Madras J. Lit. Sci. ser. 3, 1: 38. 1864; B.D.Sharma et al., Fl. Karnataka Anal. 48. 1984.

Distribution: Karnataka (Dakshina Kannada, Hassan, Kodagu, Shivamogga), Kerala, Tamil Nadu; endemic.

Nothapodytes Blume

Nothapodytes nimmoniana (J.Graham) Mabb. in Manilal, Bot. Hist. Hort. Malab. 88. 1980; C.J.Saldanha, Fl. Karnataka 2: 105. 1996

Premna nimmoniana J.Graham, Cat. Pl. Bombay 155. 1839

Stemonurus foetidus Wight, Icon. Pl. Ind. Orient. 3: t. 955. 1845

Mappia foetida (Wight) Miers, Ann. Mag. Nat. Hist. ser. 2, 9: 395. 1852

Mappia tomentosa Miers, Ann. Mag. Nat. Hist. II, 9: 397 1852

Mappia wightiana Miers, Ann. Mag. Nat. Hist. II, 9: 397 1852

Nothapodytes foetida (Wight) Sleumer in Notizbl. Bot. Gart. Berlin-Dahlem. 15: 247. 1940

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Assam, Goa, Kerala, Maharashtra, Sikkim, Tamil Nadu, West Bengal; E & SE Asia.

Local/Common/Vernacular name: *Durvasane mara* (Kannada)

Sarcostigma Wight & Arn.

Sarcostigma kleinii Wight & Arn., Edinb. New Phil. J. 14: 299. 1833; B.D.Sharma et al., Fl. Karnataka Anal. 48. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Maharashtra, Tamil Nadu; endemic.

Iridaceae**Iris L.**

Iris domestica (L.) Goldblatt & Mabb., Novon 15: 129.2005

Epidendrum domesticum L., Sp. Pl. 952.1753

Moraea chinensis (L.) Thunb., Nova Acta Regiae Soc. Sci. Upsal. 4: 39.1783

Pardanthus chinensis (L.) Ker Gawl., Ann. Bot. 1: 247.1804

Belamcanda chinensis (L.) DC., Liliac. 3: t. 121.1805; B.D.Sharma et al., Fl. Karnataka Anal. 281. 1984.

Gemmungia chinensis (L.) Kuntze, Revis. Gen. Pl. 2: 701.1891

Vanilla domestica (L.) Druce, Rep. Bot. Exch. Club Soc. Brit. Isles 3: 425.1914

Distribution: Karnataka (cultivated as an ornamental in Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Tumakuru, Uttara Kannada), cultivated throughout India; native to China.

Local/Common/Vernacular names: *Blackberry lily*, *Leopard flower*, *Leopard lily* (English)

Juncaceae**Juncus L.**

Juncus maritimus Lam., Encycl. 3: 264.1789; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Distribution: Karnataka (Vijayapura); Africa, Asia, Europe.

Local/Common/Vernacular name: *Sea rush* (English)

Juncus prismatocarpus R.Br., Prodr. Fl. Nov. Holl. 1: 259.1810; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Distribution: Karnataka (Belagavi, Chamarajanagar, Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Australia, E, S & SE Asia, New Zealand.

Luzula DC.

Luzula campestris (L.) DC., Fl. Franç. ed. 3, 3: 161.1805; B.D.Sharma et al., Fl. Karnataka Anal. 293. 1984.

Distribution: Karnataka (Kodagu), throughout cooler parts of India; cosmopolitan.

Local/Common/Vernacular names: *Field wood-rush, Good Friday grass, Sweep's brush* (English)

Lamiaceae**Anisochilus Wall.ex Benth.**

Anisochilus adenanthus Dalzell in Dalzell & A.Gibson, Bombay Fl. 206.1861

Anisochilus verticillatus Hook.f., Fl. Brit. India 4: 629.1885; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Distribution: Karnataka (Belgavi, Dharwad), Andhra Pradesh, Kerala; endemic.

Anisochilus carnosus (L.f.) Wall., Pl. Asiat. Rar. 2: 18 1830

Plate 8C

Lavandula carnosa L.f., Suppl. Pl. 273.1782

Plectranthus carnosus (L.f.) Sm., Cycl. 27: 7.1814

Plectranthus strobiliferus Roxb., Hort. Bengal. 45.1814

Plectranthus dubius Spreng., Syst. Veg., ed. 16, 2: 691. 1825, nom. illeg.

Anisochilus carnosus var. *purpurascens* Benth. in Wall., Pl. Asiat. Rar. 2: 18.1830

Anisochilus carnosus var. *glabrus* (Schrad.) Benth., Labiatarum Genera & Species: 711. 1835. ('*glabrum*').

Anisochilus glaber Schrad., Linnaea 10(Lit.): 69. 1836

Anisochilus carnosus var. *glabrior* (Schrad.) Benth. in DC., Prodr. 12: 81.1848

Anisochilus carnosus var. *villosior* Benth. in DC., Prodr. 12: 81.1848

Anisochilus carnosus var. *viridis* Benth. in DC., Prodr. 12: 81.1848

Anisochilus crassus Benth. in DC., Prodr. 12: 81.1848

Anisochilus eriocephalus Benth. in Dc., Prodr. 12: 81.1848; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Anisochilus decussatus Dalzell & A.Gibson, Bombay Fl. 206.1861

Anisochilus rupestris Wight ex Hook.f., Fl. Brit. India 4: 627.1885

Anisochilus carnosus var. *eriocephalus* (Benth.) T.Cooke, Fl. Bombay 2:450.1906

Anisochilus carnosus var. *eriocephalus* (Benth.) S. R. Paul, New Bot. 6: 45. 1979

Distribution: Karnataka (Belgavi, Ballari, Bengaluru, Chamarajanagar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada, Vijayapura), Assam, peninsular India, Sikkim,; China, Myanmar, Nepal, Sri Lanka, Thailand.

Local/Common/Vernacular name: *Karpoorada gida* (Kannada)

Anisochilus dysophylloides Benth., Pl. Asiat. Rar. 2: 19.1830; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Anisochilus sericeus Benth. in Dc., Prodr. 12: 82.1848

Anisochilus albidus Wight, Icon. Pl. Ind. Orient. 4: t. 1435.1849

Anisochilus purpureus Wight, Icon. Pl. Ind. Orient. 4: t. 1435.1849

Anisochilus dysophylloides var. *purpureus* (Wight) Gamble, Fl. Madras 2: 1128.1924

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Tamil Nadu; endemic.

Anisochilus paniculatus Benth. in Dc., Prodr. 12: 82.1848; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Anisochilus suffruticosus Thwaites, Enum. Pl. Zeyl. 238.1860

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Anisochilus plantagineus Hook. f., Fl. Brit. India 4: 628.1885

Distribution: Karnataka (Uttara Kannada); endemic.

Anisomeles R.Br.

Anisomeles heyneana Benth., Pl. Asiat. Rar. 1: 59.1830; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Distribution: Karnataka (Davanagere, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Oshthapala* (Sanskrit)

Anisomeles indica (L.) Kuntze, Revis. Gen. Pl. 2: 512.1891; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984. **Plate 8D**

Nepeta indica L., Sp. Pl. 571.1753

Ballota disticha L., Mant. Pl. 1: 83.1767

Marrubium indicum (L.) Burm.f., Fl. Indica 127.1768

Monarda zeylanica Burm.f., Fl. Indica 12.1768

Nepeta amboinica L.f., Suppl. Pl. 273.1782

Ajuga disticha (L.) Roxb., Hort. Bengal. 44.1814

Anisomeles disticha (L.) B.Heyne ex Roth, Nov. Pl. Sp. 254.1821

Nepeta disticha (L.) Blume, Bijdr. 823.1826

Epimeredi indicus (L.) Rothm., Feddes Repert. Spec. Nov. Regni Veg. 53: 12.1944

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere Dharwad, Hassan, Kodagu, Kolar, Mandya, Mysuru, Raichur, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout plains of India; Cambodia, China, Laos, Malaysia, Myanmar, Philippines, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *cat mint, Indian cat mint* (English); *Dodda thumbe, Haccu matti gida, Hennu karee tumbe, Hennu tumbe, Huchhu moothi gida, Manga mari soppu* (Kannada)

Anisomeles malabarica (L.) R.Br. ex Sims, Bot. Mag. 46: t. 2071 1819; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984. **Plate 8E**

Nepeta malabarica L., Mant. Pl. 2: 566.1771

Ajuga fruticosa Roxb., Hort. Bengal. 44.1814

Anisomeles intermedia Wight ex Benth., Labiat. Gen. Spec. 703 1835

Epimeredi malabaricus (L.) Rothm., Feddes Repert. Spec. Nov. Regni Veg. 53: 12.1944

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Hassan, Kalaburgi, Kodagu, Kolar, Mandya, Mysuru, Raichur, Tumakuru, Uttara Kannada), throughout India; Australia, Mauritius, New Guinea, Reunion, S & SE Asia.

Local/Common/Vernacular names: *Malabar cat mint* (English); *Gandu birana gida, Gandu kari tumbe, Hetthunbe, Kaadu thumbe* (Kannada)

Basilicum Moench.

Basilicum polystachyon (L.) Moench, Suppl. Meth. 143.1802; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Ocimum polystachyon L., Mant. Pl. 2: 567. 1771

Moschosma polystachyon (L.) Benth., Pl. Asiat. Rar. 2: 13.1830

Perxo polystachyon (L.) Raf., Autik. Bot. 121.1840

Distribution: Karnataka (Dakshina Kannada, Shivamogga, Udupi) throughout India; Africa, Australia, E, S & SE Asia.

Local/Common/Vernacular name: *Musk Basil* (English)

Callicarpa L.

Callicarpa tomentosa (L.) L., Syst. Veg. ed. 13 153.1774; B.D.Sharma et al., Fl. Karnataka Anal. 214. 1984.

Tomex tomentosa L., Sp. Pl. 118.1753

Callicarpa lanata L., Mant. Pl. 2: 331.1771

Callicarpa wallichiana Walp., Repert. Bot. Syst. 4: 125 1845

Callicarpa arborea Miq. ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 567 1885

Callicarpa farinosa Roxb. ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 567 1885

Callicarpa lobata C.B.Clarke in Hook.f., Fl. Brit. India 4: 566 1885

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Local/Common/Vernacular names: *Arthigida, Dodda natada gida, Rishipathiri* (Kannada)

Clerodendrum L.

Clerodendrum calamitosum L., Mant. Pl. 1: 90.1767; K. G. Bhat, Fl. S. Kanara 746. 2014.

Volkameria alternifolia Burm.f., Fl. Indica 137.1768

Distribution: Karnataka (cultivated in gardens of Dakshina Kannada, Udupi), Andaman & Nicobar Islands, Kerala, Maharashtra, Tamil Nadu, Uttar Pradesh; Malaysia, Indonesia, Philippines, Thailand.

Local/Common/Vernacular name: *White Butterfly* (English)

Clerodendrum chinense (Osbeck) Mabb., Plant-book Repr.: 707.1989

Cryptanthus chinensis Osbeck, Dagr. Ostind. Resa 215.1757

Volkameria fragrans Vent., Jard. Malmaison 70.1804

Clerodendrum fragrans var. *pleniflorum* Schauer in DC., Prodr. 11: 666.1847

Clerodendrum philippinum Schauer in DC., Prodr. 11: 667.1847; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Clerodendrum lasiocephalum C.B.Clarke in Hook.f., Fl. Brit. India 4: 594.1885

Ovidea fragrans (Vent.) Hitchc., Mem. Torrey Bot. Club 12: 63.1902

Clerodendrum fragrans f. *pleniflorum* (Schauer) Standl. & Steyermark., Publ. Field Mus. Nat. Hist., Bot. Ser. 22: 272.1940

Clerodendrum japonicum var. *pleniflorum* (Schauer) Maheshw., Taxon 15: 43.1966

Distribution: Karnataka (cultivated in Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), throughout India; cultivated in tropical and subtropical Asia; native to China & Japan.

Local/Common/Vernacular name: *Mysore mallige* (Kannada)

Clerodendrum glandulosum Lindl., Edwards's Bot. Reg. 30: t. 19.1844

Clerodendrum colebrookianum Walp., Repert. Bot. Syst. 4: 114.1845; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Clerodendrum colebrookianum var. *denticulatum* C.B.Clarke in Hook. f., Fl. Brit. India 4: 594. 1885

Clerodendrum colebrookianum var. *henryanum* Moldenke, Phytologia 52: 330.1983

Distribution: Karnataka (probably introduced Kolar), NE India; Bangladesh, Bhutan, China, Indonesia, Laos, Malaysia, Myanmar, Nepal, Thailand, Vietnam.

Clerodendrum indicum (L.) Kuntze, Revis. Gen. Pl. 2: 586.1891; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Siphonanthus indicus L., Sp. Pl. 109.1753

Ovieda mitis L., Sp. Pl. ed. 2, 889.1763

Clerodendrum siphonanthus R.Br., Hortus Kew. 4: 65.1812; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Clerodendrum verticillatum Roxb. ex D.Don, Prodr. Fl. Nepal. 102.1825

Ovieda verticillatum Roxb. ex D.Don, Prodr. Fl. Nepal. 102.1825

Distribution: Karnataka (cultivated in gardens Belagavi, Chikkamagaluru, Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), throughout India; Bhutan, Cambodia, China, Indonesia, Laos, Myanmar, Malaysia, Nepal, Sri Lanka, Thailand, Vietnam; introduced elsewhere.

Local/Common/Vernacular names: *Bowing lady, Glory bowee, Indian glory bower, Sky rocket, Tube flower, Turk's turban* (English); *Bharangi, Hunjika* (Kannada); *Bharagi* (Sanskrit)

Clerodendrum infortunatum L., Sp. Pl. 637.1753

Clerodendrum viscosum Vent., Jard. Malmaison t. 25.1803; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Ovieda infortunata (L.) Baill., Hist. Pl. 11: 95.1891

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Indonesia, China, Laos, Malaysia, Myanmar, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *The Hill glory bower* (English); *Ittai mara, Madras mallige, Thaggi gida* (Kannada); *Sthouneyeka* (Sanskrit); *Ittevu tappu* (Tulu)

Clerodendrum laevifolium Blume, Bijdr. 808.1826

Clerodendrum nutans Wall. ex D.Don, Prodr. Fl. Nepal. 103.1825

Clerodendrum wallichii Merr., J. Arnold Arbor. 33: 220.1952

Distribution: Karnataka (cultivated in gardens Belagavi, Chikkamagaluru, Dakshina Kannada, Mysuru, Udupi, Uttara Kannada), cultivated throughout India; Bangladesh, China, Myanmar, Vietnam.

Local/Common/Vernacular names: *Bridal Veil, Nodding Clerodendron, Nutan bleeding heart, Wallich's glorybower* (English)

Clerodendrum paniculatum L., Mant. Pl. 1: 90.1767

Caprifolium paniculatum Noronha, Verh. Batav. Genootsch. Kunsten 5: 9.1790

Clerodendrum diversifolium Vahl, Symb. Bot. 2: 75.1791

Clerodendrum pyramidale Andrews, Bot. Repos. 10: t. 628.1811

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), cultivated & naturalised in humid areas; Australia, Bangladesh, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Sri Lanka, Taiwan, Vietnam.

Local/Common/Vernacular names: *The Pagoda flower* (English); *Ratha pushpa, Tatha hoovu, Theru hoovu* (Kannada)

Clerodendrum phlomidis L.f., Suppl. Pl. 292 1782; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Volkameria multiflora Burm.f., Fl. Indica 137.1768

Clerodendrum phlomidis var. *ruberum* Roxb. ex Voigt, Hort. Suburb. Calcutt. 465.1845

Clerodendrum multiflorum (Burm.f.) Kuntze, Revis. Gen. Pl. 2: 526.1891

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Bidar, Chitradurga, Dharwad, Davanagere, Hassan, Kalaburgi, Kolar, Mysuru, Tumakuru, Uttara Kannada, Yadgir), throughout drier parts of India; Myanmar, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Husu lanagi*, *Ibbane gida*, *Madras mallige*, *Taggi gida*, *Thantu gida* (Kannada)

Clerodendrum splendens G.Don, Edinburgh Philos. J. 11: 349.1824; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Clerodendrum aurantium G.Don, Edinburgh Philos. J. 11: 349.1824

Siphonanthus splendens (G.Don) Hiern, Cat. Afr. Pl. 1: 841.1900

Distribution: Karnataka (cultivated in Bengaluru, Dakshina Kannada, Udupi, Tumakuru), cultivated throughout India; native to C & W Africa, widely cultivated in temperate parts of Africa, Asia, Australia, Europe, N America, West Indies.

Local/Common/Vernacular names: *Glaming glory bower*, *Glory tree* (English)

Clerodendrum thomsoniae Balf.f., Gard. Chron. 1862: 70.1862; G.K.Bhat, Fl. S. Kanara 748.2014.

Clerodendrum thomsoniae var. *balfourii* B.D.Jacks. ex Dombrain, Fl. Mag. (London) 4: t. 255.1865

Clerodendrum balfourii (B.D.Jacks. ex Dombrain) Dombrain, Fl. Mag. (London) 8: t. 432.1869

Distribution: Karnataka (cultivated in gardens of Bengaluru, Dakshina Kannada, Udupi, Tumakuru), cultivated throughout India; native to W tropical Africa.

Local/Common/Vernacular names: *Bagflower*, *Bleeding-heart vine* *Bleeding glory-bower*, *Glory-bowe* (English)

Colebrookea Sm.

Colebrookea oppositifolia Sm., Exot. Bot. 2: 111.1806; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Buchanania oppositifolia Sm., Exot. Bot. 2: 115.1806

Colebrookea ternifolia Roxb., Pl. Coromandel 3: 40, t.245.1815

Elsholtzia oppositifolia (Sm.) Poir., Dict. Sci. Nat. ed. 2, 14: 366.1819

Sussodia oppositifolia (Sm.) Buch.-Ham., Prodr. Fl. Nepal. 104.1825

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; China, Myanmar, Nepal, Thailand.

Local/Common/Vernacular name: *Thuggi gida* (Kannada); *Bhamni* (*Kunabi*)

Congea Roxb.

Congea tomentosa Roxb., Pl. Coromandel 3: 90, t.293.1820; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Roscoea villosa Roxb., Fl. Ind. ed. 1832 3: 55.1832

Congea villosa (Roxb.) Voigt, Hort. Suburb. Calcutt. 473.1845

Distribution: Karnataka (Bengaluru, Davanagere), Andhra Pradesh, Assam, Manipur, Odisha, Tamil Nadu, West Bengal; Bangladesh, China, Laos, Myanmar, Thailand, Vietnam; introduced elsewhere and naturalised.

Local/Common/Vernacular names: *Shower of Orchid*, *Shower Orchid*, *Wooly Congea* (English)

Endostemon N. E. Br.

Endostemon viscosus (Roth) M.R.Ashby, J. Bot. 74: 126.1936; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Ocimum menthooides B.Heyne ex Roth, Nov. Pl. Sp. 274.1821

Ocimum viscosum Roth, Nov. Pl. Sp. 274.1821

Plectranthus viscosus (Roth) Spreng., Syst. Veg. 2: 691.1825

Ocimum diffusum Benth., Pl. Asiat. Rar. 2: 14.1830

Orthosiphon diffusus (Benth.) Benth., Prodr. 12: 50.1848

Orthosiphon hispidus Benth., Prodr. 12: 50.1848

Ocimum adscendens Wight ex Hook.f., Fl. Brit. India 4: 614.1885

Ocimum glaucum B.Heyne ex Hook.f., Fl. Brit. India 4: 614.1885

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Hassan, Kodagu, Mysuru, Tumakuru), Kerala, Tamil Nadu; Sri Lanka.

Gmelina L.

Gmelina arborea Roxb., Hort. Bengal. 46.1814 & Pl. Corom. 3:42, t.246.1815; B.D.Sharma et al., Fl. Karnataka Anal. 216. 1984.

Gmelina rheedei Hook., Bot. Mag. 74: t. 4395.1848

Gmelina arborea var. *glaucescens* C.B.Clarke in Hook. f., Fl. Brit. India 4: 582.1885

Gmelina arborea var. *canescens* Haines, Forest Fl. Chota Nagpur 82.1910

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalaburgi, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Maharashtra, Tamil Nadu, E & NE India; China, Indonesia, Laos, Malaysia, Myanmar, Nepal, Philippines, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Candahar tree*, *Coomb teak*, *Kashmere tree* (English); *Gamhari*, *Gandhari gida*, *Guli*, *Gumti*, *Gupsi*, *Kashmiri mara*, *Kkole*, *Kooli mara*, *Kuli*, *Kumada mara*, *Kumathi kadi*, *Kumbadi*, *Kumbal*, *Kumulu*, *Kusa*, *Shivane mara*, *Shivanni*, *Sriparni*, *Triparni* (Kannada); *Shivan* (*Kunabi*); *Gamhari*, *Ashvetha*, *Bhadraparni*, *Gaandharya*, *Gandhaari*, *Gopabhadrika*, *Grishti*, *Hira*, *Kaakodumbari*, *Kaashmeeri*, *Kambharika*, *Kataphala*, *Krishna*, *Krishnaphala*, *Krishnavrintika*, *Ksheerini*, *Kumuda*, *Madhubhadra*, *Madhumati*, *Madhuparnika*, *Madhurasa*, *Mahabhadra*, *Mahakumda*, *Modini*, *Peetaphala*, *Peetarohini*, *Rohini*, *Sahadabhadra*, *Sarvathobhadra*, *Sarvathobhadrika*, *Shreeparni*, *Snigdhaparni*, *Sthoolathvacha*, *Subhadra*, *Sudridhavacha*, *Suphala*, *Svarubhadra*, *Vataha*, *Vidaari*, *Vidaarini* (Sanskrit)

Gmelina asiatica L., Sp. Pl. 626.1753; B.D.Sharma et al., Fl. Karnataka Anal. 216. 1984.

Gmelina lobata Gaertn., Fruct. Sem. Pl. 1: 268.1788

Gmelina parvifolia Roxb., Pl. Coromandel 2: 32.1802

Premna parvifolia Roth, Nov. Pl. Sp. 388.1821

Distribution: Karnataka (Bengaluru, Chitradurga, Kodagu, Mysuru, Tumakuru), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Gujarat, Kerala, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Bangladesh, Cambodia, China, Indonesia, Malaysia, Myanmar, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Asian bush beech*, *Asiatic beech berry*, *Small Kashmiri tree* (English); *Guldu*, *Guludu mara*, *Kal shivaani*, *Kiru shivaani*, *Kumathi gida*, *Shivaani mara* (Kannada)

Gmelina philippensis Cham., Linnaea 7: 109.1832; B.D.Sharma et al., Fl. Karnataka Anal. 216. 1984.

Distribution: Karnataka (cultivated in Bengaluru), cultivated in gardens throughout India; native to Philippines; cultivated and naturalised elsewhere.

Local/Common/Vernacular names: *Hedgehog*, *Yellow Parrot's beek* (English); *Guldu mara* (Kannada)

Gomphostemma Wall. ex Benth.

Gomphostemma heyneanum Wall. ex Benth., Pl. Asiat. Rar. 2: 12.1830; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Distribution: Karnataka (cultivated in Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Gomphostemma strobilinum Wall. ex Benth., Pl. Asiat. Rar. 2: 12.1830; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Distribution: Karnataka (Chikkamagaluru, Uttara Kannada), Kerala, Tamil Nadu; Bangladesh, Myanmar, Thailand, Vietnam.

Holmskioldia Retz.

Holmskioldia sanguinea Retz., Observ. Bot. 6: 31.1791; B.D.Sharma et al., Fl. Karnataka Anal. 216. 1984.

Hastingia scandens Roxb., Fl. Ind. ed. 1832 3: 66.1832

Distribution: Karnataka (cultivated throughout in gardens), indigenous in NE India, cultivated throughout India and tropical Asia.

Hyptis Jacq.

Hyptis capitata Jacq., Collectanea 1: 102. 1787; K.G.Bhat, Indian J. for. 39 (1): 67. 2016.

Distribution: Karnataka (Udupi), Andaman Islands, Goa, Kerala, Tripura, West Bengal; native to Tropical America.

Hyptis suaveolens (L.) Poit., Ann. Mus. Hist. Nat. 7: 472.1806; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Ballota suaveolens L., Syst. Nat. ed. 10 2: 1100.1759

Marrubium indicum Blanco, Fl. Filip. 477. 1837.

Bystropogon suaveolens (L.) L'Hér., Sert. Angl. 19.1789

Mesosphaerum suaveolens (L.) Kuntze, Revis. Gen. Pl. 2: 525.1891

Distribution: Karnataka (throughout), throughout tropicl ares of India; Pantropical.; native to tropical America.

Local/Common/Vernacular names: *American mint* (English); *Bhustroona gida, Ganga tulasi, Nayi tulsi, Vilayati tulsi* (Kannada); *Bhustrana* (Sanskrit)

Isodon (Benth.) Spach.

Isodon coetsa (Buch.-Ham. ex D.Don) Kudô, Mem. Fac. Sci. Taihoku Imp. Univ. 2: 131.1929

Plectranthus coetsa Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 117.1825; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Ocimum coetsa (Buch.-Ham. ex D.Don) Spreng., Syst. Veg. 4(2): 223.1827

Plectranthus macraei Benth., Labiat. Gen. Spec. 42.1832

Plectranthus rufescens Benth., Prodr. 12: 59.1848

Plectranthus coetsa var. *hookeri* Hook.f., Fl. Brit. India 4: 620.1885

Rabdosia coetsa (Buch.-Ham. ex D.Don) H.Hara, J. Jap. Bot. 47: 194.1972

Rabdosia coetsa var. *hookeri* (Hook.f.) V.Singh & P.Singh, J. Econ. Taxon. Bot. 15: 705.1992

Distribution: Karnataka (Chikkamagaluru),Kerala, Maharashtra, Tamil Nadu; Bangladesh, China, Laos, Myanmar, Nepal, Sri Lanka, Thailand, Vietnam.

Isodon japonicus (Burm.f.) H.Hara, Enum. Sperm. Jap. 1: 206.1948

Scutellaria japonica Burm.f., Fl. Indica 130.1768

Isodon glaucocalyx var. *japonicus* (Burm.f.) Kudô, Mem. Fac. Sci. Taihoku Imp. Univ. 2: 127.1929

Plectranthus japonicus (Burm.f.) Koidz., Bot. Mag. (Tokyo) 43: 386.1929; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Amethystanthus japonicus (Burm.f.) Nakai, Bot. Mag. (Tokyo) 48: 788.1934

Distribution: Karnataka (Mysuru); China, Japan, Korea, Russia.

Isodon lophanthoides (Buch.-Ham. ex D.Don) H.Hara, J. Jap. Bot. 60: 235.1985

Hyssopus lophanthoides Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 110.1825

Plectranthus gerardianus Benth., Pl. Asiat. Rar. 2: 17.1830

Plectranthus wightii J.Graham, Cat. Pl. Bombay 148.1839

Plectranthus stocksii Hook.f., Fl. Brit. India 4: 618.1885; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Plectranthus striatus var. *gerardianus* (Benth.) Hand.-Mazz., Acta Horti Gothob. 13: 379.1939

Rabdosia lophanthoides (Buch.-Ham. ex D.Don) H.Hara, J. Jap. Bot. 47: 197.1972

Rabdosia lophanthoides var. *gerardiana* (Benth.) H.Hara, J. Jap. Bot. 47: 197.1972

Plectranthus lophanthoides (Buch.-Ham. ex D.Don) Grierson & D.G.Long, Notes Roy. Bot. Gard. Edinburgh 40: 177.1982

Plectranthus lophanthoides var. *gerardianus* (Benth.) Bennet, J. Econ. Taxon. Bot. 5: 451.1984

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; China, Laos, Myanmar, Nepal, Thailand, Vietnam.

Isodon nigrescens (Benth.) H.Hara, J. Jap. Bot. 60: 236.1985

Plectranthus nigrescens Benth., Labiat. Gen. Spec. 710.1835; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Rabdosia nigrescens (Benth.) H.Hara, J. Jap. Bot. 47: 198.1972

Distribution: Karnataka (Davanagere, Mysuru, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Isodon nilgherricus (Benth.) H.Hara, J. Jap. Bot. 60: 236.1985

Plectranthus nilgherricus Benth., Prodr. 12: 57.1848; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Plectranthus scabrellus Benth., Prodr. 12: 58.1848

Rabdosia nilgherrica (Benth.) H.Hara, J. Jap. Bot. 47: 198.1972.

Distribution: Karnataka (Chamarajanagar), Kerala, Tamil Nadu; endemic.

Lavandula L.

Lavandula bipinnata (Roth) Kuntze, Revis. Gen. Pl. 2: 521.1891; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Bystropogon bipinnatus Roth, Nov. Pl. Sp. 225.1821

Lavandula bipinnata var. *burmanniana* Kuntze, Revis. Gen. Pl. 2: 521.1891

Lavandula bipinnata var. *intermedia* Kuntze, Revis. Gen. Pl. 2: 521.1891

Lavandula bipinnata var. *rothiana* Kuntze, Revis. Gen. Pl. 2: 521.1891

Distribution: Karnataka (Belagavi, Ballari, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kalaburgi, Kodagu, Mysuru, Raichur, Tumakuru, Uttara Kannada, Vijayapura), Maharashtra; endemic.

Local/Common/Vernacular name: *Jungle lavender* (English)

Lavandula canariensis (L.) Mill., Gard. Dict. ed. 8:4.1768; Seetharam, Kotresh & Uplaonkar, Fl. Dist 57. 2000.

Lavandula multifida subsp. *canariensis* (Mill.) Pit. & Proust, Iles Canaries 299.1909

Distribution: Karnataka (cultivated in Kalaburgi, Yadgir), cultivated throughout plains; native to Canary Islands

Local/Common/Vernacular name: *Canary island lavender* (English)

Leonotis (Pers.) R. Br.

Leonotis nepetifolia (L.) R.Br., Hortus Kew. 3: 409.1811; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Phlomis nepetifolia L., Sp. Pl. 586.1753

Leonurus nepetifolius (L.) Mill., Gard. Dict. ed. 8.2.1768

Distribution: Karnataka (throughout), throughout India; pantropical.

Local/Common/Vernacular names: *Christmas candlestick*, *Klip dagga*, *Lion's ear* (English); *Aane thumbe*, *Gova gadde*, *Kaadu hunne gida*, *Kaadu tumbe*, *Mullu thumbe*, *Pruthu beeja*, *Ranaberi* (Kannada)

Leonurus L.

Leonurus sibiricus L., Sp. Pl. 584.1753; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Phlomis sibirica (L.) Medik., Bot. Beob. 1783 124.1784

Panzeria sibirica Steud., Nomencl. Bot. ed. 2, 2: 265.1841

Lamium sibiricum (L.) Cordem., Fl. Réunion 488.1895

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Hassan, Kodagu, Mysuru), throughout India; native to China, Mongolia and Russia; pantropical.

Local/Common/Vernacular names: *Honeyweed*, *Siberian Motherwort* (English)

Leucas R. Br.

Leucas angustissima Sedgw., J. Indian Bot. Soc. 2: 123.1921; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.

Leucas aspera (Willd.) Link, Enum. Hort. Berol. Alt. 2: 113.1822; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Phlomis aspera Willd., Enum. Pl. 621.1809

Phlomis dimidiata Roth, Nov. Pl. Sp. 264.1821

Phlomis plukenetii Roth, Nov. Pl. Sp. 261 1821

Leucas dimidiata (Roth) Spreng., Syst. Veg. 2: 743.1825

Leucas plukenetii (Roth) Spreng., Syst. Veg. 2: 743.1825

Phlomis esculenta Roxb., Fl. Ind. ed. 1832 3: 10.1832

Leucas obliqua Buch.-Ham. ex Dillwyn, Rev. Hortus Malab. 57.1839

Phlomis obliqua Buch.-Ham. ex Hook.f., Fl. Brit. India 4: 690.1885

Distribution: Karnataka (Ballari, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Uttara Kannada), throughout plains, Andaman & Nicobar Islands; Mauritius, S & SE Asia to Papua New Guinea.

Local/Common/Vernacular names: *White dead nettle* (English); *Olle thumbe*, *Thumbe*, *Thumbe gida* (Kannada)

Leucas beddomei (Hook.f.) Sunojk. & P.Mathew, Rheedia 12: 170 2002

Leucas hirta var. *beddomei* Hook.f., Fl. Brit. India 4: 687.1885

Distribution: Karnataka (Kodagu ?), Kerala, Tamil Nadu; endemic.

Leucas biflora (Vahl) Sm., Cycl. 20(2): 7.1812; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Nepeta indica Burm.f., Fl. Indica 126. 1768.

Phlomis biflora Vahl, Symb. Bot. 3: 77.1794

Leucas procumbens Desf., Mém. Mus. Hist. Nat. 11: 7.1824

Blandinia biflora (Vahl) Raf., Fl. Tellur. 3: 88.1837

Leucas biflora var. *procumbens* (Desf.) Gamble, Fl. Madras 2:1151.1924

Distribution: Karnataka (Belagavi, Dakshina Kannada, Davanagere, Dharwad, Kalaburgi, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Island, Peninsular India, West Bengal; Bangladesh, Sri Lanka.

Leucas cephalotes (Roth) Spreng., Syst. Veg. 2: 743.1825; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Phlomis cephalotes Roth, Nov. Pl. Sp. 262.1821

Leucas capitata Desf., Mém. Mus. Hist. Nat. 11: 8.1824

Distribution: Karnataka (Bidar, Chikkamagaluru, Dharwad, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; Afghanistan, China, Fizi Islands, Iran, S Asia.

Local/Common/Vernacular names: *Dhrona pushpi*, *Tumbe* (Kannada)

Leucas chinensis (Retz.) Sm., Cycl. 20(2): 8.1812; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Phlomis chinensis Retz., Observ. Bot. 2: 19.1781

Phlomis montana J.Koenig ex Roxb., Fl. Ind. ed. 1832 3: 11.1832

Blandinia chinensis (Retz.) Raf., Fl. Tellur. 3: 88.1837

Distribution: Karnataka (Bengaluru, Tumakuru), Andhra Pradesh, Kerala, Tamil Nadu; China, Taiwan.

Leucas ciliata Benth., Pl. Asiat. Rar. 1: 61.1830

Leucas ciliata var. *hirsuta* Benth., Labiat. Gen. Spec. 614.1834

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), throughout India; China, S & SE Asia.

Leucas clarkei Hook.f., Fl. Brit. India 4: 688.1885; V. Singh, Monogr. Indian Leucas 65.2001.

Leucas stricta var. *clarkei* (Hook.f.) S.R.Paul, Bangladesh J. Bot. 6: 42.1977

Distribution: Karnataka (Chitradurga), Peninsular to N & NE India; endemic.

Leucas decemdentata (Willd.) Sm., Cycl. 20(2): 6.1812

Stachys decemdentata G.Forst., Fl. Ins. Austr. 91.1786

Phlomis decemdentata Willd., Sp. Pl. 3: 124.1800

Leucas mollissima Wall. ex Benth., Pl. Asiat. Rar. 1: 62.1830; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Phlomis moluccana Roxb., Fl. Ind. ed. 1832 3: 11.1832

Leucas mollissima var. *scaberula* Hook.f., Fl. Brit. India 4: 682.1885

Leucas montana var. *pilosa* Haines, Bot. Bihar Orissa 4: 749.1922

Leucas montana var. *strigosa* (Benth.) S.R.Paul, Bangladesh J. Bot. 6: 42.1977

Leucas flaccida var. *scaberula* (Hook.f.) S.R.Sriniv. in Fl. Tamil Nadu, India 2: 176.1987

Leucas mollissima subsp. *scaberula* (Hook.f.) Murti & Panigrahi, Fl. Bilaspur Distr. 2: 487. 1999

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Mysuru, Uttara Kannada), Peninsular to NE India; Australia, China, Europe, SE Asia, Taiwan.

Leucas diffusa Benth., Labiat. Gen. Spec. 615.1834; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Phlomis diffusa Rottler ex Hook.f., Fl. Brit. India 4: 689.1885

Distribution: Karnataka (Chitradurga, Vijayapura), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Leucas eriostoma Hook.f., Fl. Brit. India 4: 686.1885; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Leucas lepidota Wight ex Hook.f., Fl. Brit. India 4: 686.1885

Phlomis eriostoma B.Heyne ex Hook.f., Fl. Brit. India 4: 686.1885

var. **eriostoma**

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Udupi), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; endemic.

var. **lanata** Hook.f., Fl. Brit. India 4: 686.1885

Leucas eriostoma var. *stocksii* Hook.f., Fl. Brit. India 4: 686.1885

Distribution: Karnataka (Chikkamagaluru), Tamil Nadu; endemic.

Leucas hirta (B.Heyne ex Roth) Spreng., Syst. Veg. 2: 743.1825; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Phlomis hirta B.Heyne ex Roth, Nov. Pl. Sp. 264.1821

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Leucas laenceaefolia Desf., Mém. Mus. Hist. Nat. Paris 11:5.1824; Hook. f., Fl. Brit. India 4: 685. 1885.

Distribution: Karnataka (), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Leucas lamiifolia Desf., Mém. Mus. Hist. Nat. 11: 4.1824; Ramaswamy, M. R. Rao & Govindappa, Fl. Shivamogga Dist. 477. 2001.

Distribution: Karnataka (Shivamogga), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Leucas lanata Benth., Pl. Asiat. Rar. 1: 61.1830; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Leucas collina Dalzell in Hooker's J. Bot. Kew Gard. Misc. 2: 338.1850

Leucas lanata var. *nagpurensis* C.B.Clarke ex Haines, Bot. Bihar Orissa 4: 747.1922

Distribution: Karnataka (Belagavi, Ballari, Chikkamagaluru, Mysuru, Shimoga, Tumakuru, Uttara Kannada), throughout India; China, S & SE Asia.

Leucas lavandulifolia Sm., Cycl. 20(2): 2.1812; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Leonurus indicus L., Syst. Nat. ed. 10 2: 1101.1759

Leonurus malebaricus J.Koenig ex Rottb., Pl. Hort. Univ. Rar. Prog. 20.1773

Phlomis linifolia Roth, Nov. Pl. Sp. 260 1821

Leucas linifolia (Roth) Spreng., Syst. Veg. 2: 743.1825

Heterpta lavandulifolia (Sm.) Raf., Fl. Tellur. 3: 88.1837

Leucas indica (L.) Vatke, Oesterr. Bot. Z. 25: 95.1875

Leucas malabarica W.Theob., Burmah ed. 3, 2: 296.1883

Leucas linifolia var. *decipiens* Hook.f., Fl. Brit. India 4: 691.1885

Leucas lavandulifolia var. *decipiens* (Hook.f.) Chandrab. & S.R.Sriniv., Bull. Bot. Surv. India 17: 165.1978

Leucas lavanulifolia var. *nagalapuramiana* Chandrab. & S.R.Sriniv., Bull. Bot. Surv. India 17: 165.1978

Leucas indica var. *decipiens* (Hook.f.) Bennet, J. Econ. Taxon. Bot. 5: 452.1984

Distribution: Karnataka (throughout), Andaman & Nicobar Islands, throughout plains of India; pantropical.

Local/Common/Vernacular names: *Gantu thumbe*, *Kanijali gida*, *Thumbe* (Kannada); *Drona pushpi* (Sanskrit)

Leucas longifolia Benth., Labiat. Gen. Spec. 744.1835; B.D.Sharma et al., Fl. Karnataka Anal. 221.

1984.

Distribution: Karnataka (Vijayapura), Gujarat, Maharashtra; Sri Lanka, West Indies.

Leucas marrubiooides Desf., Mém. Mus. Hist. Nat. 11: 6.1824; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Leucas martinicensis (Jacq.) R.Br., Hortus Kew. 3: 409.1811; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Clinopodium martinicense Jacq., Enum. Syst. Pl. 25.1760

Phlomis martinicensis (Jacq.) Sw., Prodr. Veg. Ind. Occ. 88.1788

Distribution: Karnataka (Belagavi, Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Vijayapura), throughout tropical plains of India; pantropical.

Leucas montana (Roth) Spreng., Syst. Veg. 2: 742.1825

Phlomis montana Roth, Nov. Pl. Sp. 263. 1821.

Leucas hamiltoniana Benth., Pl. Asiat. Rar. 1: 62.1830

Leucas montana var. *wightii* Hook.f., Fl. Brit. India 4: 682.1885

Leucas montana var. *parviflora* Haines, Bot. Bihar Orissa 4: 749.1922

Distribution: Karnataka (Ballari, Chikkamagaluru, Mysuru, Tumakuru), Peninsular to E India; endemic.

Leucas nepetifolia Benth., Pl. Asiat. Rar. 1: 62.1830; B.D.Sharma et al., Fl. Karnataka Anal. 221. 1984.

Distribution: Karnataka (Kolar, Mysuru, Shivamogga), Andhra Pradesh, Tamil Nadu; endemic.

Leucas nutans (Roth) Spreng., Syst. Veg. 2: 743.1825

Phlomis nutans Roth, Nov. Pl. Sp. 263.1821

Leucas decurva Benth., Pl. Asiat. Rar. 1: 60.1830

Distribution: Karnataka (Raichur, Tumakuru, Vijayapura), throughout except NE India & Islands; Myanmar, Pakistan.

Leucas prostrata (Hook.f.) Gamble, Fl. Madras 1154.1924; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Leucas hirta var. *prostrata* Hook.f., Fl. Brit. India 4: 687 1885

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Leucas pubescens Benth., Labiat. Gen. Spec. 610.1834; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Distribution: Karnataka (Chamarajanagara, Tumakuru), Kerala, Tamil Nadu; endemic.

Leucas rosmarinifolia Benth., Pl. Asiat. Rar. 1: 61.1830

Distribution: Karnataka (Chamarajanagar?), Kerala, Tamil Nadu; endemic.

Leucas sivadasaniana Sunojk., Novon 18:247.2008

Plate 24A

Distribution: Karnataka (Udupi); endemic.

Leucas stelligera Wall. ex Benth., Pl. Asiat. Rar. 1: 61.1830; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Leucas eriostoma var. *longifolia* Hook.f., Fl. Brit. India 4: 686.1885

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India except NE India & Islands; Sri Lanka

Leucas stricta Benth. in Wall., Pl. Asiat. Rar. 1: 61.1830; Bhaskar & Kushalappa, Fl. Tumkur Dist. 163. 2013.

Phlomis stricta B.Heyne ex Hook.f., Fl. Brit. India 4: 688.1885

Distribution: Karnataka (Chitradurga, Mysuru, Kodagu, Tumakuru), throughout India; Myanmar.

Leucas suffruticosa Benth., Labiat. Gen. Spec. 611.1834; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Distribution: Karnataka (Chamrajanagara, Tumakuru), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Leucas urticifolia (Vahl) Sm., Cycl. 20(2): 4.1812; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Phlomis urticifolia Vahl, Symb. Bot. 3: 76.1794

Eneodon urticifolia (Vahl) Raf., Fl. Tellur. 3: 88.1837

Distribution: Karnataka (Dharwad, Kodagu, Mysuru, Raichur, Vijayapura), E Africa, Middle East, Pakistan.

Leucas vestita Benth., Pl. Asiat. Rar. 1: 61.1830; G. K. Bhat, Fl. S. Kanara 753. 2014.

Leucas ciliata var. *vestita* (Benth.) Sunojk., Candollea 63: 83.2008

Distribution: Karnataka (Dakshina Kannada), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Leucas wightiana Wall. ex Benth., Pl. Asiat. Rar. 1: 60.1830

Leucas aspera var. *wightiana* (Wall. ex Benth.) Benth., Labiat. Gen. Spec. 616.1834

Distribution: Karnataka (), Andhra Pradesh, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Leucas zeylanica (L.) W.T.Aiton, Hortus Kew. 3: 409.1811; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Phlomis indica L., Sp. Pl. 586.1753

Phlomis zeylanica L., Sp. Pl. 586.1753

Leonurus marrubiastrum Burm.f., Fl. Indica 127.1768

Leucas indica (L.) Sm., Cycl. 20(2): 5. 1812.

Leucasia zeylanica (L.) Raf., Fl. Tellur. 3: 88.1837

Distribution: Karnataka (Chikkamagaluru, Kalaburgi, Mysuru, Shivamogga, Vijayapura), throughout except Islands; E, S & SE Asia.

Local/Common/Vernacular name: *Huccchu tumbe* (Kannada)

Mentha L.

Mentha × piperita L., Sp. Pl. 576.1753; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Mentha aquatica f. *piperita* (L.) G.Mey., Chloris Han. 290,1836

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Europe & Middle East.

Local/Common/Vernacular name: *Peppermint* (English)

Ocimum L.

Ocimum americanum L., Cent. Pl. I:15.1755; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Ocimum canum Sims, Bot. Mag. 51:t. 2452.1823

Ocimum album Roxb., Fl. Ind. ed. 1832 3: 15.1832

Ocimum dichotomum Hochst. ex Benth., Prodr. 12: 39.1848

Distribution: Karnataka (throughout), throughout plains of India; pantropical.

Local/Common/Vernacular name: *Nayi tulasi* (Kannada)

Ocimum basilicum L., Sp. Pl. 597.1753; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Ocimum thyrsiflorum L., Mant. Pl. 1: 84.1767

Ocimum barrelieri Roth, Nov. Pl. Sp. 278.1821

Plectranthus barrelieri (Roth) Spreng., Syst. Veg. 2: 691.1825

Ocimum anisatum Benth., Labiat. Gen. Spec. 4.1832

Ocimum caryophyllum Roxb., Fl. Ind. ed. 1832 3: 16.1832

Ocimum ciliare B.Heyne ex Hook. f., Fl. Brit. India 4: 608.1885

Distribution: Karnataka (often cultivated in gardens Bengaluru, Chitradurga, Dakshina Kannada, Davanagere, Kalaburgi, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Yadgir), Peninsular India; paleotropical.

Local/Common/Vernacular names: *Bu tulsi, Kama kastoori* (Kannada)

Ocimum filamentosum Forssk., Fl. Aegypt.-Arab. 108.1775

Ocimum indicum Roth, Nov. Pl. Sp. 273.1821

Plectranthus indicus (Roth) Spreng., Syst. Veg. 2: 691.1825

Ocimum cristatum Buch.-Ham. & J.Koenig ex Roxb., Fl. Ind. ed. 2, 3: 19.1832

Ocimum pubescens Bojer ex Benth., Prodr. 12: 35.1848

Ocimum adscendens Willd., Sp. Pl. 3: 166. 1800.

Ocimum verticillatum B.Heyne ex Hook.f., Fl. Brit. India 4: 609.1885

Becium filamentosum (Forssk.) Chiov., Nuovo Giorn. Bot. Ital. n.s., 26: 162.1919

Distribution: Karnataka (Bengaluru, Chikkballapur, Chitradurga, Hassan, Kodagu, Kolar, Mysuru, Ramanagara, Tumakuru, Uttara Kannada), peninsular India; native to tropical America.

Local/Common/Vernacular names: *Heddumbe gida, Mara tulasi* (Kannada)

Ocimum gratissimum L., Sp. Pl. 1197.1753; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Ocimum zeylanicum Medik., Hist. & Commentat. Acad. Elect. Sci. Theod.-Palat. 4: 197.1780

Ocimum urticifolium Roth, Catal. Bot. 2: 52.1800

Ocimum viridiflorum Roth, Catal. Bot. 2: 54.1800

Ocimum gratissimum var. *suave* (Willd.) Hook.f., Fl. Brit. India 4: 609.1885

Ocimum robustum B.Heyne ex Hook.f., Fl. Brit. India 4: 608.1885

Distribution: Karnataka (Belagavi, Chitradurga, Dakshina Kannada, Davanagere, Mysuru, Tumakuru, Udupi), Kerala, Maharashtra, Tamil Nadu; pantropical.

Local/Common/Vernacular names: *Nimbe tulasi*, *Rama tulasi* (Kannada)

Ocimum kilimandscharicum Gürke, Pflanzenw. Ost-Afrikas C: 349.1895; B.D.Sharma et al., Fl. Karnataka Anal. 222. 1984.

Distribution: Karnataka (Shivamogga), Jammu & Kashmir, Maharashtra, Karnataka, Kerala, Uttarakhand, Uttar Pradesh, West Bengal; tropical E Africa.

Local/Common/Vernacular names: *African Basil*, *Camphor Basil*, *Camphor-Scented Basil*, *Fever plant*, *Hoary Basil*, *Kilimanjaro Basil*, *Perennial Basil* (English); *Karpura tulasi* (Kannada)

Ocimum tenuiflorum L., Sp. Pl. 597.1753

Ocimum monachorum L., Mant. Pl. 1: 85.1767

Ocimum sanctum L., Mant. Pl. 1: 85.1767; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Ocimum inodorum Burm.f., Fl. Indica 130.1768

Lumnitzera tenuiflora (L.) Spreng., Syst. Veg. 2: 687.1825

Plectranthus monachorum (L.) Spreng., Syst. Veg. 2: 690.1825

Ocimum hirsutum Benth., Pl. Asiat. Rar. 2: 14.1830

Ocimum villosum Roxb., Fl. Ind. ed. 1832 3: 13.1832

Moschosma tenuiflorum (L.) Heynh., Nom. Bot. Hort. 532.1841

Ocimum sanctum var. *hirsutum* (Benth.) Hook.f., Fl. Brit. India 4: 609.1885

Ocimum subserratum B.Heyne ex Hook.f., Fl. Brit. India 4: 609.1885

Geniosporum tenuiflorum (L.) Merr., Philipp. J. Sci. 19: 379.1921

Distribution: Karnataka (throughout wild & cultivated), native to Indian subcontinent.

Local/Common/Vernacular names: *Sacred basil* (English); *Kari tulasi*, *Krishna tulasi*, *Sri tulasi*, *Tulasi*,

Vishnu tulasi (Kannada)

Origanum L.

Origanum vulgare L., Sp. Pl. 590.1753; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Origanum creticum L., Sp. Pl. 589.1753

Origanum laxiflorum Royle ex Benth., Hooker's J. Bot. Kew Gard. Misc. 3: 376.1833

Origanum venosum Willd. ex Benth., Linnaea 11: 239.1837

Thymus origanum Kuntze, Taschen-Fl. Leipzig 106.1867

Distribution: Karnataka (Bengaluru, Chikkaballapura, Kolar, Mandya, Mysuru, Tumakuru), cultivated as pot plant in plains of India; native to Eurasia & Mediterranean region

Local/Common/Vernacular names: *Oregano*, *Wild marjoram* (English); *Maruga* (Kannada)

Orthosiphon Benth.

Orthosiphon pallidus Royle ex Benth., Hooker's J. Bot. Kew Gard. Misc. 3: 370.1833; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Orthosiphon inodorus K.D.Koenig ex Hook.f., Fl. Brit. India 4: 613.1885

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dharwad, Kalaburgi, Kolar, Mysuru, Raichur, Tumakuru, Uttara Kannada, Yadgir), Kerala, Maharashtra, Tamil Nadu; Arabia, Pakistan.

Orthosiphon rubicundus (D.Don) Benth., Pl. Asiat. Rar. 2: 14.1830; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Ocimum rubicundum Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 116.1825

Plectranthus rubicundus D.Don, Prodr. Fl. Nepal. 116.1825

Lumnitzera rubicunda (D.Don) Spreng., Syst. Veg. 4(2): 223.1827

Orthosiphon rubicundus var. *canescens* Benth., Pl. Asiat. Rar. 2: 14.1830

Orthosiphon rubicundus var. *mollissimus* Benth., Pl. Asiat. Rar. 2: 14.1830

Ocimum tuberosum Roxb., Fl. Ind. ed. 1832 3: 18.1832

Plectranthus mollissimus Wall. ex Benth., Labiat. Gen. Spec. 26.1832

Plectranthus tuberosus Roxb. ex Benth., Labiat. Gen. Spec. 26.1832

Ocimum mollissimum Wall. ex Hook.f., Fl. Brit. India 4: 615.1885

Orthosiphon rubicundus var. *hohenackeri* Hook.f., Fl. Brit. India 4: 615.1885

Orthosiphon rubicundus var. *macrocarpus* Prain, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 66: 520.1897

Distribution: Karnataka (Chikkamagaluru, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga), peninsular India to W Himalaya; Myanmar, Nepal, Pakistan.

Local/Common/Vernacular name: *Redjava tea* (English)

Orthosiphon thymiflorus (Roth) Sleesen, Reinwardtia 5: 42.1959; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Ocimum thymiflorum Roth, Nov. Pl. Sp. 269.1821

Plectranthus thymiflorus (Roth) Spreng., Syst. Veg. 2: 690.1825

Orthosiphon glabratus Benth., Pl. Asiat. Rar. 2: 14.1830

Orthosiphon viscosus Benth., Pl. Asiat. Rar. 2: 14.1830

Orthosiphon tomentosus var. *parviflorus* Benth., Prodr. 12: 51.1848

Orthosiphon tomentosus var. *glabratus* (Benth.) Hook.f., Fl. Brit. India 4: 614.1885

Orthosiphon tomentosus var. *rubiginosus* C.B.Clarke ex Hook.f., Fl. Brit. India 4: 614.1885

Orthosiphon tomentosus var. *viscosus* (Benth.) Hook.f., Fl. Brit. India 4: 614.1885

Orthosiphon glabratus var. *parviflorus* (Benth.) Gamble, Fl. Pres. Madr. 2: 1114.1924

Distribution: Karnataka (Ballari, Bidar, Chikkamagaluru, Chitradurga, Dharwad, Kodagu, Mysuru, Tumakuru, Uttara Kannada), peninsular to E India; SE Asia.

Local/Common/Vernacular name: *Kadu sanna patri gida* (Kannada)

Platostoma P.Beauv.

Platostoma africanum P.Beauv., Fl. Oware 2: 61.1818

Ocimum flaccidum A.Rich., Tent. Fl. Abyss. 2: 179.1850

Platostoma flaccidum (A.Rich.) Benth.in Hook., Fl. Brit. India 4: 611.1885; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Distribution: Karnataka (Dharwad, Mysuru, Uttara Kannada, Udupi), Maharashtra; tropical Africa.

Platostoma hispidum (L.) A.J.Paton, Kew Bull. 52: 273.1997

Gomphrena hispida L., Sp. Pl. ed. 2 326.1762

Prunella indica Burm.f., Fl. Indica 130.1768

Ocimum capitellatum L.f., Suppl. Pl. 276.1782

Ocimum capitatum Roth, Nov. Pl. Sp. 276.1821

Acrocephalus indicus (Burm.f.) Kunze, Revis. Gen. Pl. 2: 511.1891; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Acrocephalus hispidus (L.) Nicolson & Sivad., Taxon 29: 324.1980

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; SE Asia.

Plectranthus L.Herit.

Plectranthus amboinicus (Lour.) Spreng., Syst. Veg. 2: 690.1825

Coleus amboinicus Lour., Fl. Cochinch. 372.1790

Coleus aromaticus Benth., Pl. Asiat. Rar. 2: 16.1830

Coleus crassifolius Benth., Pl. Asiat. Rar. 2: 15.1830

Majana amboinica (Lour.) Kuntze, Revis. Gen. Pl. 2: 524.1891

Distribution: Karnataka (cultivated as pot herb throughout, some places naturalised), cultivated throughout drier parts of India; native to E Africa.

Local/Common/Vernacular names: *Allherb*, *Big thyme*, *Broad leaf thyme*, *Broadleaf thyme*, *Cuban oregano*, *Country borage*, *French thyme*, *Indian borage*, *Indian mint*, *Mexican mint*, *Mexican thyme*, *Mother of herbs*, *Poor man pork*, *Queen of herbs*, *Soup mint*, *Spanish thyme*, *Thick leaf thyme*, *Three-in-one herb* (English); *Doddapatri*, *Sambara balli* (Kannada)

Plectranthus barbatus Andrews, Bot. Repos. 10: t. 594.1810; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Ocimum asperum Roth, Nov. Pl. Sp. 268.1821

Plectranthus asper (Roth) Spreng., Syst. Veg. 2: 690.1825

Coleus barbatus (Andrews) Benth. ex G.Don, Hort. Brit. 483.1830

Orthosiphon asperus (Roth) Benth. ex Sweet, Hort. Brit. ed. 2: 413.1830

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Kodagu, Mysuru, Tumakuru, Uttara Kannada), throughout India; Madagascar, S & W Asia, tropical Africa.

Plectranthus beddomei Raizada, Indian Forester 84: 503.1958

Coleus urticifolius Benth., Prodr. 12: 78.1848

Plectranthus urticifolius (Benth.) Hook.f., Fl. Brit. India 4: 622.1885; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Plectranthus bishopianus Gamble, Fl. Madras 1122.1924; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Distribution: Karnataka (Chikkamagaluru), Tamil Nadu; endemic.

Plectranthus caninus Roth, Nov. Pl. Sp. 279.1821; B.D.Sharma et al., Fl. Karnataka Anal. 223. 1984.

Coleus spicatus Benth., Pl. Asiat. Rar. 2: 15.1830

Plectranthus monadelphus (R.Br. ex Roth) Buch.-Ham. ex Wall., Numer. List 2729.1831

Coleus heynei Benth., Labiat. Gen. Spec. 50.1832

Coleus caninus (Roth) Vatke, Linnaea 37: 318.1872

Majana canina (Roth) Kuntze, Revis. Gen. Pl. 2: 524.1891

Majana spicata (Benth.) Kuntze, Revis. Gen. Pl. 2: 524.1891

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Chitradurga, Hassan, Kolar, Mysuru), S India; Australia, tropical & S Africa, S Asia.

Local/Common/Vernacular names: *Dogbane*, *Piss-off plan*, *Scaredy Cat Plant* (English)

Plectranthus crameri R.H.Willemse, Blumea 25: 510.1979; Manjunatha, Krishna & Pullaiah 322. 2004.

Coleus mollis Benth., Prodr. 12: 77.1848

Distribution: Karnataka (Davanagere, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Plectranthus hadiensis (Forssk.) Schweinf. ex Sprenger, Wiener Ill. Gart.-Zeitung 19: 2.1894; R. R. Rao & Razi, Synop. Fl. Mysore Dist.

Ocimum hadiense Forssk., Fl. Aegypt.-Arab. 109.1775

Plectranthus forsskaolii Vahl, Symb. Bot. 1: 44.1790

Distribution: Karnataka (Mysuru), tropical & S Africa, Middle East.

Plectranthus malabaricus (Benth.) R.H.Willemse, Blumea 25: 509.1979; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Coleus malabaricus Benth., Pl. Asiat. Rar. 2: 16.1830

Coleus macraei Benth., Labiat. Gen. Spec. 58.1832

Coleus ovatus Benth., Labiat. Gen. Spec. 57.1832

Coleus leptostachys Benth., Prodr. 12: 77.1848

Coleus walkeri Benth., Prodr. 12: 77.1848

Plectranthus malabaricus var. *leptostachys* (Benth.) R.H.Willemse, Blumea 25: 510.1979

Distribution: Karnataka (Hassan, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Plectranthus mollis (Aiton) Spreng., Syst. Veg. 2: 690.1825; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Ocimum maypurensse Roth, Nov. Pl. Sp. 271.1821

Ocimum cordifolium Buch.-Ham. ex D.Don, Prodr. Fl. Nepal. 116.1825

Plectranthus cordifolius D.Don, Prodr. Fl. Nepal. 116.1825

Plectranthus maypurensis (Roth) Spreng., Syst. Veg. 2: 691.1825

Plectranthus secundus Roxb., Fl. Ind. ed. 1832 3: 20.1832

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Kerala, Maharashtra, Sikkim, Tamil Nadu, Uttaranchal; Nepal.

Local/Common/Vernacular name: *Nona kasa* (Kannada)

Plectranthus parvifolius Talbot, J. Bombay Nat. Hist. Soc. 11: 238.1897; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Distribution: Karnataka (Uttara Kannada); endemic.

Plectranthus rotundifolius (Poir.) Spreng., Syst. Veg. 2: 690.1825

Germanea rotundifolia Poir., Encycl. 2: 763.1788

Plectranthus tuberosus Blume, Bijdr. 838. 1826.

Coleus rugosus Benth., Pl. Asiat. Rar. 2: 15.1830

Coleus tuberosus (Blume) Benth., Labiat. Gen. Spec. 59.1832

Coleus parviflorus Benth., Prodr. 12: 72.1848

Majana tuberosa (Blume) Kuntze, Revis. Gen. Pl. 2: 524.1891

Solenostemon rotundifolius (Poir.) J.K.Morton, J. Linn. Soc., Bot. 58: 272.1962

Calchas parviflorus (Benth.) P.V.Heath, Calyx 6: 51.1999

Distribution: Karnataka (Dakshina Kannada, Udupi), native to tropical Africa.

Local/Common/Vernacular names: *Country potato Chinese potato* (English); *Sambraani gida, Sambrali* (Kannada); *Kuka* (Konkani)

Plectranthus scutellarioides (L.) R.Br., Prodr. Fl. Nov. Holl. 506.1810; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Ocimum scutellarioides L., Sp. Pl. ed. 2 833.1763

Plectranthus aromaticus Roxb., Hort. Bengal. 45.1814

Coleus scutellarioides (L.) Benth., Pl. Asiat. Rar. 2: 16.1830

Coleus blumei Benth., Labiat. Gen. Spec. 56.1832

Coleus scutellarioides var. *blumei* (Benth.) Miq., Fl. Ned. Ind. 2: 950.1858

Majana scutellarioides var. *blumei* (Benth.) Kuntze, Revis. Gen. Pl. 2: 524.1891

Plectranthus blumei (Benth.) Launert, Mitt. Bot. Staatssamml. München 7: 301.1968

Solenostemon scutellarioides (L.) Codd, Bothalia 11: 439.1975

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to SE Asia.

Local/Common/Vernacular name: *Painted nettle* (English)

Pogostemon Desf.

Pogostemon auricularius (L.) Hassk., Tijdschr. Natuurl. Gesch. Physiol. 10: 127.1843; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Mentha auricularia L., Mant. Pl. 1: 81.1767

Mentha foetida Burm.f., Fl. Indica 126.1768

Dysophylla auricularia (L.) Blume, Bijdr. 826.1826

Eusteralis auricularia (L.) M.R.Almeida, Fl. Maharashtra 4A: 146.2003

Distribution: Karnataka (Ballari, Chikkamagaluru, Kodagu, Mysuru), Kerala, Tamil Nadu; Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Philippines, Sri Lanka, Thailand, Vietnam.

Pogostemon benghalensis (Burm.f.) Kuntze, Revis. Gen. Pl. 2: 529.1891; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.

Origanum benghalense Burm.f., Fl. Indica 128.1768

Origanum indicum Roth, Nov. Pl. Sp. 265.1821

Mentha integra Buch.-Ham. ex Benth., Pl. Asiat. Rar. 1: 31.1830

Pogostemon frutescens J.Graham, Cat. Pl. Bombay 149.1839

Pogostemon purpuricaulis Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 336.1850

Pogostemon indicus (Roth) Kuntze, Um die Erde 500.1881

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Shivamogga, Tumakuru, Uttara Kannada), Peninsular to E India; endemic.

Local/Common/Vernacular name: *Naati pachi thene* (Kannada)

Pogostemon cruciatus (Benth.) Kuntze, Revis. Gen. Pl. 2: 530.1891 ('*cruciatum*'); B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Mentha quadrifolia D.Don, Prodr. Fl. Nepal. 113.1825

Dysophylla cruciata Benth., Pl. Asiat. Rar. 1: 30.1830

Dysophylla tetraphylla Wight, Icon. Pl. Ind. Orient. t. 1444.1849

Pogostemon tetraphylla (Wight) F.Muell., Fragm. 4: 47.1863

Anuragia cruciata (Benth.) Raizada, Suppl. Duthie's Fl. Upper Gangetic Plain 218.1976

Eusteralis cruciata (Benth.) Panigrahi, Phytologia 32: 478.1976

Distribution: Karnataka (Hassan, Kodagu, Shivamogga), Peninsular India to Himalayas & NE India; Nepal.

Pogostemon deccanensis (Panigrahi) Press, Bull. Brit. Mus. (Nat. Hist.), Bot. 10: 73.1982 **Plate 24B**

Eusteralis deccanensis Panigrahi, Phytologia 32: 465.1976

Distribution: Karnataka (Belagavi, Dakshina Kannada, Udupi, Uttara Kannada), Kerala, Maharashtra; endemic.

Pogostemon erectus (Dalzell) Kuntze, Revis. Gen. Pl. 2: 530.1891
Dysophylla erecta Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 337.1850
Dysophylla gracilis Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 337.1850
Pogostemon gracilis (Dalzell) F.Muell., Fragm. 4: 47.1863
Anuragia erecta (Dalzell) Raizada, Suppl. Duthie's Fl. Upper Gangetic Plain 219.1976
Eusteralis erecta (Dalzell) Panigrahi, Phytologia 32: 477.1976; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.
Eusteralis gracilis (Dalzell) Panigrahi, Phytologia 32: 476.1976
Eusteralis tomentosa var. *gracilis* (Dalzell) Bennet & Raizada, Indian Forester 108: 303.1982
Pogostemon erectus var. *gracilis* (Dalzell) Panigrahi, J. Orissa Bot. Soc. 6: 52.1984
 Distribution: Karnataka (Uttara Kannada), Maharashtra; endemic.

Pogostemon heyneanus Benth., Pl. Asiat. Rar. 1: 31. 1830; G. K. Bhat, Fl. S. Kanara 757.2014.
Pogostemon suavis Ten., Giorn. Bot. Ital. 2: 56.1847
Pogostemon patchouly var. *suavis* (Ten.) Hook.f., Fl. Brit. India 4: 634.1885
 Distribution: Karnataka (Dakshina Kannada, Kodagu, Shivamogga, Udupi, Uttara Kannada), peninsular India; SE Asia, Sri Lanka.
 Local/Common/Vernacular names: *False patchpouli*, *Indian patchouli*, *Java patchouli* (English); *Pacche koralu*, *Pachhe kadiru*, *Pachhe thene* (Kannada)

Pogostemon mollis Benth., Labiat. Gen. Spec. 155.1833; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.
 Distribution: Karnataka (Belagavi, Hassan, Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Pogostemon myosuroides (Roth) Kuntze, Revis. Gen. Pl. 2: 530.1891; B.D.Sharma et al., Fl. Karnataka Anal. 224. 1984.
Mentha myosuroides Roth, Nov. Pl. Sp. 257.1821
Dysophylla myosuroides (Roth) Benth. ex Wall., Numer. List 1547.1829
Eusteralis myosuroides (Roth) M.R.Almeida, Fl. Maharashtra 4A: 147.2003
 Distribution: Karnataka (Ballari, Chikkamagaluru), Maharashtra; endemic.

Pogostemon paniculatus (Willd.) Benth., Pl. Asiat. Rar. 1: 30.1830; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.
Elsholtzia paniculata Willd., Sp. Pl. 3: 59.1800
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.
 Local/Common/Vernacular name: *Pachhe thene* (Kannada)

Pogostemon peethapushpam Pradeep, Candollea 523: 419. 1998
 Distribution: Karnataka (Chikkamagaluru), Kerala; endemic.

Pogostemon plectrantoides Desf., Mém. Mus. Hist. Nat. 2: 155.1815; G. K. Bhat, Fl. S. Kanara 757.2014.
Ocimum bengalense Poir., Encycl. 4: 608.1798
Mentha secunda Roxb., Hort. Bengal. 44.1814
Wensea pyramidata J.C.Wendl., Coll. Pl. 3: 26.1819
Mentha fruticosa Roxb. ex D.Don, Prodr. Fl. Nepal. 115.1825
 Distribution: Karnataka (Dakshina Kannada, Udupi), Meghalaya, W Bengal

Pogostemon pubescens Benth., Prodr. 12: 152.1848; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.
 Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Pogostemon purpurascens Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 337.1850; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.
 Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Shivamogga, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Pogostemon quadrifolius (Benth.) F.Muell., Fragm. 5: 200.1866

Dysophylla quadrifolia Benth., Pl. Asiat. Rar. 1: 30.1830

Mentha quadrifolia Roxb., Fl. Ind. ed. 1832 3: 4. 1832.

Mentha stellata Buch.-Ham. ex Roxb., Fl. Ind. ed. 1832 3: 5.1832

Dysophylla rupestris Dalzell, Hooker's J. Bot. Kew Gard. Misc. 3: 120.1851

Eusteralis quadrifolia (Benth.) Panigrahi, Phytologia 32: 478.1976; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Eusteralis malabarica Majumdar, J. Bombay Nat. Hist. Soc. 74: 385.1977; B.D.Sharma et al., Fl. Karnataka Anal. 219. 1984.

Eusteralis stellata var. *roxburghiana* (H.Keng) Bennet & Raizada, Indian Forester 108: 303.1982

Distribution: Karnataka (Belagavi, Ballari, Chikkamagaluru, Dakshina Kannada, Shivamogga, Udupi, Uttara Kannada), Andhra Pradesh, Assam, Kerala, Maharashtra, Meghalaya, Odisha, W Bengal; Bangladesh, Myanmar.

Pogostemon speciosus Benth., Pl. Asiat. Rar. 1: 31.1830; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; endemic.

Pogostemon stellatus (Lour.) Kuntze, Revis. Gen. Pl. 2: 429.1891

Mentha stellata Lour., Fl. Cochinch. 361. 1790.

Mentha quaternifolia B.Heyne ex Roth, Nov. Pl. Sp. 256.1821

Mentha verticillata (Benth.) Roxb., Fl. Ind. ed. 1832 3: 5.1832

Dysophylla tomentosa Dalzell, Hooker's J. Bot. Kew Gard. Misc. 2: 337.1850

Pogostemon verticillatus (Benth.) Miq., Fl. Ned. Ind. 2: 965.1859

Anuragia tomentosa (Dalzell) Raizada, Suppl. Duthie's Fl. Upper Gangetic Plain vii.1976

Anuragia verticillata (Benth.) Raizada, Suppl. Duthie's Fl. Upper Gangetic Plain vii.1976

Eusteralis tomentosa (Dalzell) Panigrahi, Phytologia 32: 477.1976; B.D.Sharma et al., Fl. Karnataka Anal. 220. 1984.

Pogostemon verticillatus (Roxb.) Bhatti & Ingr., Bull. Nat. Hist. Mus. London, Bot. 27(2): 115. 1997.

Distribution: Karnataka (Dakshina Kannada, Davanagere, Kalaburgi, Hassan, Kodagu, Shivamogga, Uttara Kannada, Yadgir), Peninsular India; Australia, S E Asia.Sri Lanka.

Pogostemon wightii Benth., Labiat. Gen. Spec. 156.1833; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Pogostemon hirsutus Wight, Icon. Pl. Ind. Orient. t. 1442.1849

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Premna L.

Premna coriacea C.B.Clarke in Hook. f., Fl. Brit. India 5: 573.1888; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.

var. **coriacea**

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Udupi, Uttara Kannada), Andaman & Nicobar Islands, S India to NE India; Bangladesh, Bhutan, Cambodia, Laos, Myanmar, Thailand, Vietnam.

Local/Common/Vernacular names: *Chaankhaari*, *Ishe*, *Javangi balli* (Kannada)

var. **villosa** (C.B.Clarke) A.Rajendran & P.Daniel, Bull. Bot. Surv. India 38: 43.2001

Premna villosa C.B.Clarke in Hook.f.,Fl. Brit. India 4: 573.1885; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.

Gumira villosa (C.B.Clarke) Kuntze, Revis. Gen. Pl. 2: 508.1891

Premna coriacea (C.B.Clarke) A. Rajendran & P. Daniel, Bull. Bot. Surv. India 38(1–4): 43. 1996.

Distribution: Karnataka (Ballari, Kodagu) , Kerala, Tamil Nadu; Bangladesh, China, Laos, Thailand, Vietnam.

Premna mollissima Roth, Nov. Pl. Sp. 286.1821*Premna latifolia* Roxb., Fl. Ind. ed. 1832 3: 76.1832; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.*Premna viburnoides* Wall. ex Schauer, Prodr. 11: 635.1847*Premna latifolia* var. *cuneata* C.B.Clarke in Hook.f., Fl. Brit. India 4: 578.1885*Premna latifolia* var. *mucronata* (Roxb.) C.B.Clarke in Hook.f., Fl. Brit. India 4: 578.1885*Premna latifolia* var. *viburnoides* C.B.Clarke in Hook.f., Fl. Brit. India 4: 578.1885*Gumira mollissima* (Roth) Kuntze, Revis. Gen. Pl. 2: 508.1891*Premna latifolia* var. *gamblei* Haines, Bot. Bihar Orissa 717.1925*Premna latifolia* var. *henryi* D.Naras. ex A.Rajendran & P.Daniel, Indian Verbenac. 253.2002

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Mysuru, Tumakuru, Udupi, Uttara Kannada), throughout India; Bangladesh, Bhutan, China, Myanmar, Cambodia, Laos, Nepal, Pakistan, Philippines, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *Adrae soppu, Nergula* (Kannada)**Premna serratifolia** L., Mant. Pl. 2: 253.1771*Cornutia corymbosa* Burm.f., Fl. Indica 132.1768*Premna obtusifolia* R.Br., Prodr. Fl. Nov. Holl. 512.1810; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.*Premna spinosa* Roxb., Fl. Ind. ed. 1832 3: 77.1832*Premna corymbosa* (Burm.f.) Schauer, Prodr. 11: 632.1847*Premna littoralis* King & Gamble, Bull. Misc. Inform. Kew 1908: 107.1908*Premna angustior* (C.B.Clarke) Ridl., Fl. Malay Penins. 2: 619.1923*Premna integrifolia* var. *obtusifolia* (R.Br.) C.Pei, Mem. Sci. Soc. China 3: 75.1932*Premna obtusifolia* var. *angustior* (C.B.Clarke) Moldenke, Phytologia 5: 87.1954*Premna obtusifolia* var. *minor* (Ridl.) Moldenke, Phytologia 5: 88.1954*Premna serratifolia* var. *minor* (Ridl.) A.Rajendran & P.Daniel, Bull. Bot. Surv. India 38: 44. 2001

Distribution: Karnataka (Chitradurga, Dakshina Kannada, Udupi, Uttara Kannada), China, S & SE Asia, Australia, Taiwan.

Premna tomentosa Willd., Sp. Pl. 3: 314.1800; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.*Premna latifolia* Thwaites, Enum. Pl. Zeyl. 242. 1861.

Distribution: Karnataka (Bengaluru, Chamarajanagara, Chitradurga, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Tumakuru), Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Bachanige mara, Eeji gida, Narale, Naruvale, Narve, Navaraalu, Thavadike* (Kannada)**Rothea Raf.****Rothea incisa** (Klotzsch) Steane & Mabb., Novon 8: 205.1998; K. G. Bhat, Fl. S. Kanara 747. 2014*Clerodendrum incisum* Klotzsch, Naturw. Reise Mossambique 257. 1861*Clerodendrum macrosiphon* Hook.f., Bot. Mag. 109: t. 6695 1883*Clerodendrum incisum* var. *macrosiphon* (Hook.f.) Baker in Oliv., Fl. Trop. Afr. 5: 308.1900

Distribution: Karnataka (cultivated as hedge plant in Dakshina Kannada, Udupi), cultivated in S India; native to Tanzania.

Rothea serrata (L.) Steane & Mabb., Novon 8: 206 1998*Volkameria serrata* L., Mant. Pl. 1: 90.1767*Volkameria herbacea* Roxb., Hort. Bengal. 46.1814*Clerodendrum serratum* (L.) Moon, Cat. Pl. Ceylon 46.1824; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.*Cyclonema serratum* (L.) Hochst., Flora 25: 227.1842*Clerodendrum herbaceum* Roxb. ex Schauer, Prodr. 11: 675.1847*Clerodendrum serratum* var. *wallichii* C.B.Clarke in Hook.f., Fl. Brit. India 4: 592.1885*Clerodendrum serratum* var. *herbaceum* (Roxb. ex Schauer) C.Y.Wu in Fl. Yunnanica 1: 468. 1977

Distribution: Karnataka (throughout), Peninsular India; China, S & SE Asia.

Local/Common/Vernacular names: *Bhaarangi, Bharangi gida, Gantu bharangi, Gungurada hoovu, Kariteggi, Kir taggi gida, Nirvisha* (Kannada); *Bharang* (Kunabi)

Salvia L.

Salvia aegyptiaca L., Sp. Pl. 23.1753; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Thymus syrticus Spreng., Syst. Veg. 2: 697.1825

Salvia pumila Benth., Labiat. Gen. Spec. 726.1835

Pleudia aegyptiaca (L.) Raf., Fl. Tellur. 3: 94.1837

Distribution: Karnataka (Vijayapura), NW India; Afghanistan, Arabian peninsula, Cape Verde Islands, Canary Islands, Iran, Pakistan, tropical Africa.

Local/Common/Vernacular name: *Egyptian sag* (English)

Salvia coccinea Buc'hoz ex Etl., Salv. 23.1777; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Horminum coccineum (Buc'hoz ex Etl.) Moench, Methodus 377.1794

Salvia ciliata Benth., Labiat. Gen. Spec. 286.1833

Distribution: Karnataka (cultivated in gardens almost throughout), cultivated throughout plains; native to Mexico.

Salvia leucantha Cav., Icon. 1: 16.1791; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Mexico.

Local/Common/Vernacular name: *Mexican bush sage* (English)

Scutellaria L.

Scutellaria barbata D.Don, Prodr. Fl. Nepal. 109.1825

Scutellaria rivularis Wall. ex Benth., Pl. Asiat. Rar. 2: 66.1831; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Scutellaria squamulosa A.Ham., Esq. Monogr. Scutellaria 35.1832

Scutellaria minor var. *indica* Benth., Prodr. 12: 427.1848

Distribution: Karnataka (Mysuru), Peninsular to E India; China, Japan, Korea, Laos, Myanmar, Nepal, Thailand, Vietnam.

Local/Common/Vernacular name: *Barbed skullcap* (English)

Scutellaria colebrookiana Wall., Pl. Asiat. Rar. 1: 67.1830; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Scutellaria violacea var. *colebrookiana* Hook.f., Fl. Brit. India 4:668.1885

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chikkamagaluru, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Scutellaria discolor Colebr., Pl. Asiat. Rar. 1: 66.1830; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; S & SE Asia.

Scutellaria scandens D.Don, Prodr. Fl. Nepal. 110.1825; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Scutellaria angulosa Benth., Pl. Asiat. Rar. 1: 67.1830

Scutellaria celtidifolia A.Ham., Esq. Monogr. Scutellaria 27.1832

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Scutellaria violacea B.Heyne ex Benth., Pl. Asiat. Rar. 1: 66.1830; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Scutellaria indica Roxb., Fl. Ind. ed. 1832 3: 24.1832

Scutellaria floribunda Benth., Prodr. 12: 418.1848

Distribution: Karnataka (Mysuru), Andhra Pradesh, Kerala, Tamil Nadu; Indonesia, Nepal, Sri Lanka.

Scutellaria wightiana Benth., Pl. Asiat. Rar. 1: 67.1830; B.D.Sharma et al., Fl. Karnataka Anal. 225. 1984.

Scutellaria violacea var. *hispidior* Hook.f., Fl. Brit. India 4:668.1885

Distribution: Karnataka (Mysuru), Tamil Nadu; endemic.

Sphenodesme Jack.

Sphenodesme involucrata (C.Presl) B.L.Rob., Proc. Amer. Acad. Arts 51: 531. 1916.

var. **paniculata** (C.B.Clarke) Munir, Gard. Bull. Singapore 21: 338.1966

Sphenodesme paniculata C.B.Clarke in Hook.f., Fl. Brit. India 4: 600.1885; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Sympcorema Roxb.

Sympcorema involucratum Roxb., Pl. Coromandel 2: 46, t.186.1805; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.

Analectis speciosa Vahl, Kongel. Danske Vidensk. Selsk. Skr. 6: 94.1810

Congea involucratum (Roxb.) Wall., Numer. List 1740.1829

Distribution: Karnataka (Dakshina Kannada, Hassan, Udupi, Uttara Kannada), peninsular to NE India; Myanmar, Sri Lanka, Thailand.

Sympcorema polyandrum Wight, Icon. Pl. Ind. Orient. t. 363.1840; B.D.Sharma et al., Fl. Karnataka Anal. 217. 1984.

Tectona theca Lour., Fl. Cochinch. 137.1790

Jatus grandis (L.f.) Kuntze, Revis. Gen. Pl. 2: 508.1891

Distribution: Karnataka (Belagavi, Dharwad, Raichur, Uttara Kannada), Andhra Pradesh, Odisha, Tamil Nadu; endemic.

Distribution: Karnataka (cultivated throughout), throughout India up to terai Himalayas, also cultivatrd in Andaman & Nicobar Islands; S & SEAsia.

Local/Common/Vernacular names: *Teak tree* (English); *Saguvani, Saguvani mara, Tega, Thega* (Kannada); *Sag (Kunabi)*

Tectona L.f.

Tectona grandis L.f., Suppl. Pl. 151.1782;B. D. Sharma et al., Fl. Karnataka Anal. 217.1984.

Tectona theca Lour., Fl. Cochinch. 137.1790.

Theka grandis (L.f.) Lam., Tabl. Encycl. 2: 111.1797.

Jatus grandis (L.f.) Kuntze, Revis. Gen. Pl. 2: 508.1891.

Karnataka (cultivated throughout), throughout India up to terai Himalayas.

Local names: Teak tree (English), Saaguvani mara, Thega (Kannada)

Teucrium L.

Teucrium heynei V.S.Kumar & Chakrab., Phytotaxa 243: 197. 2016.

Teucrium tomentosum B.Heyne ex Benth., Pl. Asiat. Rar. 1: 58. 1830. non Lam., 1779.; Kesh. Murthy & Yogan., Fl. Coorg. 1990.

Distribution: Karnataka (Belagavi, Hassan, Kodagu, Mysuru, Tumakuru, Uttara Kannada), Kerala, Tamil Nadu; Sri Lanka.

Teucrium polium L., Sp. Pl. 566.1753

Teucrium tomentosum Lam., Fl. Franç. 2: 407.1779

Chamaedrys polium (L.) Raf., Fl. Tellur. 3: 85.1837

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Udupi), native to the western Mediterranean region,

Local/Common/Vernacular name: *Felty germander* (English)

Vitex L.f.

Vitex altissima L.f., Suppl. Pl. 294.1782; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Vitex alata Willd., Neue Schriften Ges. Naturf. Freunde Berlin 4: 203.1803

Vitex latifolia Wight ex Steud., Nomencl. Bot. ed. 2, 2: 777.1841

Vitex zeylanica Turcz., Bull. Soc. Imp. Naturalistes Moscou 36(2): 223.1863

Vitex appendiculata Rottler ex C.B.Clarke in Hook. F., Fl. Brit. India 4: 584.1885

Vitex altissima var. *alata* (Willd.) Trimen, Handb. Fl. Ceylon 3 : 358. 1895

Distribution: Karnataka (Bengaluru, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Peninsular India; Bangladesh, Indo-China, SE Asia, Sri Lanka.

Local/Common/Vernacular names: *Baingani*, *Bharanege*, *kaadu Navile*, *Myrole*, *Navatari*, *Navladi*, *Tore nekki mara* (Kannada); *Bailada* (*Kunabi*); *Myrolu* (Tulu)

Vitex leucoxylon L.f., Suppl. Pl. 293.1782; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Wallrothia leucoxylon (L.f.) Roth, Nov. Pl. Sp. 319.1821

Vitex saligna Roxb., Fl. Ind. ed. 1832 3: 75.1832

Vitex rheedei Kostel., Allg. Med.-Pharm. Fl. 3: 826.1834

Vitex leucoxylon var. *zeylanica* Moldenke, Phytologia 21: 419.1971

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Indonesia, Malaysia, Myanmar, Pakistan, Sri Lanka.

Local/Common/Vernacular names: *Hole lakki*, *Hole nekki*, *Kariyale*, *Senkeni* (Kannada)

Vitex negundo L., Sp. Pl. 638.1753; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Vitex incisa Lam., Encycl. 2: 612.1788

Agnus-castus incisa (Lam.) Carrière, Rev. Hort. 42: 415.1871

Agnus-castus negundo (L.) Carrière, Rev. Hort. 42: 415.1871

Vitex negundo var. *incisa* (Lam.) C.B.Clarke in Hook.f., Fl. Brit. India 4: 584.1885

Vitex agnus-castus var. *negundo* (L.) Kuntze, Revis. Gen. Pl. 2: 511.1891

Vitex negundo var. *intermedia* (C.P'ei) Moldenke, Revista Sudamer. Bot. 5: 2.1937; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Vitex negundo var. *purpurascens* Sivar. & Moldenke, Phytologia 28: 404.1974

Distribution: Karnataka (throughout), throughout India; S & SE Asia.

Local/Common/Vernacular names: *Kari lakki gida*, *Kari nekke*, *Lakki gida*, *Lakki soppu*, *Neergundi*, *Nekki* (Kannada); *Nirgundi* (*Kunabi*); *Nekki tappu* (Tulu)

Vitex peduncularis Wall. ex Schauer in DC., Prodr. 11: 687 1847; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Vitex morava Buch.-Ham. ex Wall., Numer. List 1752.1829

Vitex peduncularis var. *roxburghiana* C.B.Clarke in Hook.f., Fl. Brit. India 4: 587.1885

Distribution: Karnataka (Tumakuru, Uttara Kannada), Kerala, Tamil Nadu; Indo-China, S & SE Asia.

Vitex pinnata L., Sp. Pl. 638.1753; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Vitex arborea Roxb., Fl. Ind. ed. 1832 3: 73.1832

Vitex digitata Wight ex Steud., Nomencl. Bot. ed. 2, 2: 777.1841

Pistaciovitex pinnata (L.) Kuntze, Lex. Gen. Phan. 442.1903

Distribution: Karnataka (Mysuru), Kerala, Tamil Nadu; Indo-China, S & SE Asia.

Vitex trifolia L., Sp. Pl. 638.1753; B.D.Sharma et al., Fl. Karnataka Anal. 218. 1984.

Vitex indica Mill., Gard. Dict. ed. 8 in errata. 1768

Vitex triphylla Royle, Ill. Bot. Himal. Mts. 299.1836

Vitex agnus-castus var. *trifolia* (L.) Kurz, Forest Fl. Burma 270.1877

Distribution: Karnataka (Dakshina Kannada, Davanagere, Mysuru, Tumakuru, Udupi, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Afghanistan, Australia, Iran, Madagascar, Pacific Islands, E, S & SE Asia.

Local/Common/Vernacular name: *Neer Lakki* (Kannada)

Volkameria L.

Volkameria aculeata L., Sp. Pl. 637. 1753

Clerodendrum aculeatum (L.) Schltdl., Linnaea 6: 750.1831

Ovieda aculeata (L.) Baill., Hist. Pl. 11: 95.1891

Distribution: Karnataka (cultivated as hedge plant in Dakshina Kannada, Udupi), native to West Indies.

Local/Common/Vernacular names: *Amourette, Bitter fence, Coffee fence, Cough bush, Hague bush, Miracle fence, Prayer berry, Prickly wild coffee, Privy hedge* (Kannada)

Volkameria heterophylla Poir., Encycl. 8: 687.1808

Clerodendrum heterophyllum (Poir.) R.Br., Hortus Kew. 4: 64.1812

Distribution: Karnataka (cultivated as hedge plant in Dakshina Kannada, Udupi), cultivated in gardens; native to Mauritius & Reunion.

Volkameria inermis L., Sp. Pl. 637.1753

Ovieda inermis (L.) Burm.f., Fl. Indica 136.1768

Clerodendrum inerme (L.) Gaertn., Fruct. Sem. Pl. 1: 271.1788; B.D.Sharma et al., Fl. Karnataka Anal. 215. 1984.

Clerodendrum coromandelianum Spreng., Syst. Veg. 2: 758.1825

Volkameria neriifolia Roxb., Fl. Ind. ed. 1832 3: 64.1832

Clerodendrum neriifolium (Roxb.) Wall. ex Steud., Nomencl. Bot. ed. 2, 1: 383.1840

Clerodendrum neriifolium var. *macrocarpum* Wall. ex C.B.Clarke in Hook.f., Fl. Brit. India 4: 589.1885

Clerodendrum inerme var. *macrocarpum* (Wall. ex C.B.Clarke) Moldenke, Phytologia 22: 6.1971

Distribution: Karnataka (throughout, often cultivated), throughout India except Himalayas; Australia, Brazil, Pacific Islands, tropical Asia, Hawaii Islands.

Local/Common/Vernacular names: *Glory bower, Indian Pivot* (English); *Kundali gida, Vishamdhari, Vishampuri* (Kannada)

Lauraceae

Actinodaphne Nees

Actinodaphne bourdillonii Gamble, Bull. Misc. Inform. Kew 1925: 129. 1925; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Mysuru, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Actinodaphne gullavarai (Buch.-Ham. ex Nees) M.R.Almeida, Fl. Maharashtra 4A: 251 2003

Litsea angustifolia Blume, Bijdr. Fl. Ned. Ind. 566 1825

Laurus gullavarai Buch.-Ham. ex Nees in Pl. Asiat. Rar. 2: 65 1831

Actinodaphne angustifolia Nees in Pl. Asiat. Rar. 3: 34 1832; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Actinodaphne areolata Blume in Mus. Bot. 1: 34 1849

Distribution: Karnataka (Belagavi, Davanagere, Kodagu, Shivamogga, Uttara Kannada), Goa, Maharashtra;

Local/Common/Vernacular name: *Tud genasu* (Kannada); *Amberi* (*Kunabi*)

Actinodaphne hookeri Meisn., Prodr. 15(1): 218 1864

Actinodaphne angustifolia Hook.f. & Thomson ex Meisn., Prodr. 15(1): 218. 1864.

Jozoste hookeri Kuntze, Revis. Gen. Pl. 2: 569.1891

Distribution: Karnataka (Davangere, Uttara Kannada), Peninsular to NE India; endemic.

Local/Common/Vernacular names: *Galavara*, *Heggodi mara*, *Tudgenasu* (Kannada)

Actinodaphne madraspatana Bedd. ex Hook.f., Fl. Brit. India 5: 149 1886; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Distribution: Karnataka (Dakshina Kannada), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Actinodaphne tadulingamii Gamble, Bull. Misc. Inform. Kew 1925; C.J.Saldanha, Fl. Karnataka 1: 58. 1984

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada), Kerala, Tamil Nadu; endemic.

Actinodaphne wightiana (Kuntze) Noltie, Regnum Veg. 145: 495 2005

Actinodaphne hirsuta Hook. f., Fl. Brit. India 5: 152. 1886, non Blume 1851; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Jozoste wightiana Kuntze, Revis. Gen. Pl. 2: 570 1891

Actinodaphne malabarica N.P.Balakr., J. Bombay Nat. Hist. Soc. 63: 329. 1967

Distribution: Karnataka (Hassan, Kodagu, Mysuru, Shimoga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Alseodaphne Nees

Alseodaphne semecarpifolia Nees in Wall., Pl. Asiat. Rar. 2: 72. 1831. var. *semecarpifolia*; C.J.Saldanha, Fl. Karnataka 1: 59. 1984

Alseodaphne semecarpifolia var. *angustifolia* Meisn. in DC., Prodr. 15: 28. 1864

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Kachu* (Kannada); *Phadgus* (Kunabi)

Beilschmiedia Nees

Beilschmiedia dalzellii (Meisn.) Kostermans in Reinwardtia 6:282. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Beilschmiedia fagifolia var. *dalzellii* Meisn. in DC., Prodr. 15(1): 64. 1864

Distribution: Karnataka (Belagavi, Mysuru, Uttara Kannada), Goa, Maharashtra; endemic.

Beilschmiedia gemmiflora (Blume) Kosterm., J. Sci. Res. (Indonesia) 1:115. 1952; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Laurus gemmiflora Blume, Bijdr. Fl. Ned. Ind. 11: 555. 1826

Polyadenia gemmiflora (Blume) Nees, Syst. Laur. 576. 1836

Daphnidium gemmiflorum (Blume) Blume, Mus. Bot. 1:352. 1851

Lindera gemmiflora (Blume) Boerl., Handl. Fl. Ned. Ind. (Boerlage) 3(1): 147. 1900

Beilschmiedia bourdillonii Brandis, Indian Trees 528. 1906

Distribution: Karnataka (Dakshina Kannada, Shivamogga), Goa, Kerala; Indonesia, New Guinea.

Beilschmiedia roxburghiana Nees, Pl. Asiat. Rar. 2: 69. 1831; Punekar & Lakshmin., Fl. Anshi Natl. Park 392. 2011

Distribution: Karnataka (Uttara Kannada), Andhra Pradesh, Goa, Maharashtra, Odisha, Tamil Nadu; endemic.

Beilschmiedia wightii (Nees) Benth. ex Hook. f., Fl. Brit. India 5: 124. 1886; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Haasia wightii Nees, Syst. Laurin. 676. 1836

Distribution: Karnataka (Chikkamagaluru, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Cassytha Osbeck

Cassytha capillaris Meisn., Prodr. 15(1): 252 1864; B.D.Sharma et al., Fl. Karnataka Anal. 239. 1984.

Distribution: Karnataka (Chikkamagaluru, Kalaburagi, Shivamogga), Kerala, Tamil Nadu; Sri Lanka.

Cassytha filiformis L., Sp. Pl. 35. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 239. 1984.

Distribution: Karnataka (Throughout), throughout India; pantropical.

Local/Common/Vernacular name: *Aakash balli* (Kannada)

Cinnamomum Schaeff.

Cinnamomum camphora (L.) J.Presl, Prir. Rostlin 2: 36 1825; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Laurus camphora L., Sp. Pl. 369.1753

Persea camphora (L.) Spreng., Syst. Veg. 2: 268.1825

Camphora camphora (L.) H.Karst., Ill. Repet. Pharm.-Med. Bot. 504.1886

Camphorina camphora (L.) Farw., Druggists' Circular 62: 535.1918

Distribution: Karnataka (Bengaluru, Chikkaballapur, Chikkamagaluru), Assam, Bihar, Himachal Pradesh, Kerala, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh; native to E Asia & Indo-China; cultivated throughout tropics.

Local/Common/Vernacular names: *Camphor laurel*, *Camphor tree*, *Camphor wood* (English)

Cinnamomum cassia (L.) J.Presl, Prir. Rostlin 2: 44. 1825

Laurus cassia L., Sp. Pl. 369. 1753

Cinnamomum nitidum Hook., Exot. Fl. 3: t. 176.1825

Persea cassia (L.) Spreng., Syst. Veg. 2: 267.1825

Camphorina cassia (Nees & T.Nees) Farw., Druggists' Circular 62: 535.1918

Neolitsea cassia (L.) Kosterm., J. Sci. Res. (Jakarta) 1: 85.1952

Distribution: Karnataka (cultivated in Uttara Kannada), cultivated in Peninsular India and S & SE Asia; native to China.

Local/Common/Vernacular names: *Chinese cassia*, *Chinese cinnamon Cinnamon* (English)

Cinnamomum goaense Kosterm., Bull. Bot. Surv. India 25: 94. 1985.

Distribution: Karnataka (Uttara Kannada), Kerala; endemic

Cinnamomum macrocarpum Hook. f., Fl. Brit. India 5: 133. 1886; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Chikkaballapur, Chikkamagaluru, Hassan, Kodagu, Kolar, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Lavange patre* (Kannada); *Teja patre* (Sanskrit)

Cinnamomum malabatum (Burm.f.) Blume, Bijdr. 568. 1826; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Laurus malabatum Burm. f., Fl. Indica 92. 1768

Cinnamomum malabathricum Lukman., Nomencl. Icon. Cannel. 8.1889

Cinnamomum malabatum var. *rheedei* Lukman., Nomencl. Icon. Cannel. 8.1889

Cinnamomum rheedii Lukman., Nomencl. Icon. Cannel. 8.1889

Distribution: Karnataka (Hassan, Kodagu, Mysuru, Shimoga, Uttara Kannad), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Adavulavangapatte*, *Lavanga patra*, *Lavangada yele* (Kannada)

Cinnamomum riparium Gamble, Bull. Misc. Inform. Kew 1925. 128. 1925; C.J.Saldanha, Fl. Karnataka 1: 62. 1984

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; endemic.

Cinnamomum sulphuratum Nees in Wall., Pl. Asiat. Rar. 2: 74. 1831; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Chikkamagaluru, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga Uttara Kannada), Goa, Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Pinga dalchinni* (Kannada)

Cinnamomum travancoricum Gamble, Bull. Misc. Inform. Kew 1925; C.J.Saldanha, Fl. Karnataka 1: 62. 1984

Distribution: Karnataka (Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Cinnamomum verum Presl, Prir. Rostl. 2: 36. t. 7. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Cinnamomum zeylanicum Blume, Bijdr. 11, 568 1825

Distribution: Karnataka (cultivated in Belagavi, Bengaluru Chikkamagaluru, Chitradurga, Dakshina Kannada, Dharwad, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu, Tripura; cultivated in Madagascar, Seychelles ; native to Sri Lanka.

Local/Common/Vernacular names: *Ceylon cinnamon tree*, *Cinnamo*, *True cinnamon tree* (English); *Chakke mara* (Kannada)

Cinnamomum wightii Meisn. in DC., Prodr. 15(1): 11. 1864; C.J.Saldanha, Fl. Karnataka 1: 63. 1984

Cinnamomum ovalifolium Gardner ex Meisn. in DC., Prodr. 15(1): 11.1864

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu; Sri Lanka.

Cryptocarya R. Br.

Cryptocarya beddomei Gamble, Bull. Misc. Inform. Kew 1925: 127.1925; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Dakshina Kannada, Mysuru, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Cryptocarya lawsonii Gamble, Bull. Misc. Inform. Kew 1925; C.J.Saldanha, Fl. Karnataka 1: 64. 1984

Distribution: Karnataka (Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Cryptocarya procera Talbot, Forest Fl. Bombay 2:385. 1911; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Uttara Kannada); endemic.

Cryptocarya stocksii Meisn. in DC., Prodr. 15:71.1864; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Cryptocarya neilgherrensis Meisn. in DC., Prodr. 15(1): 71.1864; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Chikkamagaluru, Mysuru, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Cryptocarya wightiana Thwaites, Enum. Pl. Zeyl. 254. 1861; C.J.Saldanha, Fl. Karnataka 1: 64. 1984

Cryptocarya floribunda Wight, Icon. Pl. Ind. Orient. 5: t. 1829.1852

Cryptocarya wightiana var. *lanceolata* Meisn., Prodr. 15(1): 70.1864

Cryptocarya wightiana var. *parvifolia* Meisn., Prodr. 15(1): 70.1864

Cryptocarya bourdillonii Gamble, Bull. Misc. Inform. Kew 1925: 127.1925; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Maharashtra, Tamil Nadu; Sri Lanka.

Litsea Lam.

Litsea bourdillonii Gamble, Bull. Misc. Inform. Kew 1925:131.1925; C.J.Saldanha, Fl. Karnataka 1: 65. 1984.

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Litsea coriacea Hook. f., Fl. Brit. India 5: 166. 1886; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Litsea deccanensis Gamble, Fl. Madras 1235(864). 1925; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular name: *Dodda kana lakki* (Kannada)

Litsea floribunda (Blume) Gamble, Fl. Madras 1238. 1925; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Cylcodaphne floribunda Blume, Mus. Bot. Lugd.-Bat. 1: 387. 1851

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular names: *Halu maddi, Neeralagi* (Kannada)

Litsea ghatica C.J.Saldanha, Fl. Karnataka 1: 67. 1984

Distribution: Karnataka (Belagavi, Hassan, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra; endemic.

Litsea glabrata Hook.f., Fl. Brit. India 5: 174.1886; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Malapoenna glabrata Kuntze, Revis. Gen. Pl. 2: 572.1891

Distribution: Karnataka (Chikkmagaluru), Kerala, Tamil Nadu; endemic.

Litsea glutinosa (Lour.) C.B.Rob., Philipp. J. Sci. 6: 321. 1911; C.J.Saldanha, Fl. Karnataka 1: 67. 1984

Sebifera glutinosa Lour., Fl. Cochinch. 638.1790

Laurus involucrata J.Koenig ex Retz., Observ. Bot. 6: 27.1791

Litsea chinensis Lam., Encycl. 3: 574.1792; B.D.Sharma et al., Fl. Karnataka Anal. 237. 1984.

Tetranthera apetala Roxb., Pl. Coromandel 2: 25.1800

Litsea apetala (Roxb.) Pers., Syn. Pl. 2: 4.1806

Lepidadenia wightiana Nees, Edinburgh New Philos. J. 15: 379.1833

Distribution: Karnataka (Belagavi, Dharwad, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Bihar, Himachal Pradesh, Kerala, Maharashtra, Odisha, Punjab, Tamil Nadu, Tripura, West Bengal; Australia, S & SE Asia.

Litsea laevigata (Nees) Gamble, Fl. Madras 123. 1925; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Tetranthera attenuata var. *laevigata* Nees, Syst. Laurin. 677. 1836

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Litsea lakshmanniana Kesh.Murthy & Yogan., Curr. Sci. 56: 371. 1987; Kesh.Murthy & Yogan, Fl. Kodagu 384. 1990

Distribution: Karnataka (Chikkamagaluru, Kodagu); endemic.

Litsea mysorensis Gamble, Bull. Misc. Inform. Kew 1925; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Mysuru), Kerala, Tamil Nadu; endemic.

Litsea oblonga Hook.f., Fl. Brit. India 5: 168 1886 non Nees 1836 nec Meisn. 1864; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Distribution: Karnataka (Dakshina Kannada, Uttara Kannada), S to NE India

Litsea oleoides Hook.f., Fl. Brit. India 5: 175. 1886; Udayan et al., Indian For., 130: 560. 2004

Distribution: Karnataka (Chikkamagaluru), Kerala, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Kaadubende* (Kannada)

Litsea stocksii (Meisn.) Hook. f., Fl. Brit. India 5: 176. 1886; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Tetranthera oblonga var. *stocksii* Meisn., DC., Prodr. 15: 205. 1864

Litsea stocksii var. *glabrescens* (Meisn.) Hook.f., Fl. Brit. India 5: 176. 1886; C.J.Saldanha, Fl. Karnataka 1: 69. 1984.

Litsea vartakii M.R.Almeida, J. Bombay Nat. Hist. Soc. 86: 180. 1989

Litsea josephii S.M. Almeida, Fl. Savantwadi 1: 364. 1990

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Kodagu, Shivamogga, Uttara Kannada), Goa, Kerala, Maharashtra, Tamil Nadu; endemic.

Litsea travancorica Gamble, Bull. Misc. Inform. Kew 1925: 132. 1925; Punekar & Lakshmin., Fl. Anshi Natl. Park 397. 2011

Distribution: Karnataka (Uttara Kannada), Kerala, Tamil Nadu; endemic.

Litsea wightiana (Nees) Hook. f., Fl. Brit. India 5: 177. 1886; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Clycodaphne wightiana Nees in Wall., Pl. Asiat. Rar. 2: 68. 1831

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Kolar, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Konda hippa* (Kannada)

Neocinnamomum H. Liu

Neocinnamomum caudatum (Nees) Merrill, Contr. Arnold Arbor. 8: 64. 1934

Cinnamomum caudatum Nees, Pl. Asiat. Rar. 2: 76 1831; B.D.Sharma et al., Fl. Karnataka Anal. 236. 1984.

Distribution: Karnataka (Kodagu), Andhra Pradesh, Assam, Sikkim; Bhutan, China, Myanmar, Nepal, Thailand, Vietnam.

Neolitsea (Benth. & Hook. f.) Merr.

Neolitsea scrobiculata (Meisn.) Gamble, Fl. Madras 1240.1925; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Litsea scrobiculata Meisn. in DC., Prodr. 15(1): 223. 1864

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Kerala, Tamil Nadu; endemic.

Neolitsea umbrosa (Nees) Gamble, Pl. Wilson. 2: 79. 1914

Tetradenia foliosa Nees, Pl. Asiat. Rar. 2: 64. 1831

Litsea foliosa (Nees) Nees, Syst. Laur. 622. 1836

Litsea foliosa var. *impunctata* Meisn. in DC., Prodr. 15(1): 222. 1865

Litsea umbrosa var. *khasiana* Meisn. in DC., Prodr. 15(1): 223. 1865

Litsea umbrosa var. *impunctata* (Meisn.) Hook.f., Fl. Brit. India 5: 180. 1886

Neolitsea foliosa (Nees) Gamble, Fl. Madras 1240.1925; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Neolitsea foliosa var. *caesia* (Meisn.) Gamble, Fl. Madras 1240.1925

Neolitsea umbrosa var. *impunctata* (Meisn.) V. Singh & P. Singh, J. Econ. Taxon. Bot. 15(3): 705. 1991

Distribution: Karnataka (Mysuru, Shivamogga, Uttara Kannada), Himachal Pradesh, Meghalaya; S & SE Asia.

Neolitsea zeylanica (Nees & T. Nees) Merr., Philipp. J. Sci. Suppl. 1: 57. 1906; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Litsea zeylanica Nees & T.Nees, Amoen. Bot. Benn. Fasc. 1: 58. t.5. 1823

Tetradenia zeylanica (Nees & T. Nees) Nees, Pl. Asiat. Rar. 2: 64.1831

Distribution: Karnataka (Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Assam, Goa, Maharashtra, Manipur, Odisha, Tamil Nadu; S & SE Asia.

Ocotea Aubl.

Ocotea lancifolia (Schott) Mez, Jahrb. Königl. Bot. Gart. Berlin 5: 254. 1889

Persea lancifolia Schott, Syst. Veg. 4(2): 405.1827

Ocotea ligustrina Nees, Pl. Asiat. Rar. 2: 71 .1831

Laurus lanceolaria Roxb., Fl. Ind., ed. 1832, 2: 309.1832

Strychnodaphne lanceolata Nees, Linnaea 8: 39.1833

Ocotea lanceolata (Nees) Nees in Wall., Pl. Asiat. Rar. 2: 71. 1831 & Syst. Laur. 474.1836; C.J.Saldanha, Fl. Karnataka 1: 72. 1984

Oreodaphne lancifolia (Schott) Nees, Syst. Laur. 410.1836

Phoebe lanceolata (Nees) Nees, Syst. Laur. 109.1836

Distribution: Karnataka (Kodagu), Kerala, Tamil Nadu.

Persea Mill.

Persea americana Mill., Gard. Dict. ed. 8 1768; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Laurus persea L., Sp. Pl. 370 1753

Distribution: Karnataka (cultivated in Bengaluru, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shimoga, Uttara Kannada), cultivated in Peninsular India ; native to tropical America.

Local/Common/Vernacular names: *Alligator pear, Avocado, Butter fruit* (English); *Avocado, Butter fruit* (Kannada)

Persea macrantha (Nees) Kosterm., Reinwardtia 6: 193. 1962; B.D.Sharma et al., Fl. Karnataka Anal. 238. 1984.

Machilus macrantha Nees, Wall., Pl. Asiat. Rar. 2: 70. 1831

Machilus glaucescens Wight, Icon. Pl. Ind. Orient. 5: 12, t.1825. 1852 non. (Nees) H.W.Li 1988

Persea glaucescens (Wight) N.P. Singh, Fl. East. Karnataka, 2: 548. 1988

Distribution: Karnataka (Belagavi, Chikkaballapur, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Goa, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Gula mavu, Koluru mauvu, Kulur mara, Kurma mara* (Kannada)

Phoebe Nees

Phoebe cathia (D.Don) Kosterm., Nat. Hist. Bull. Siam Soc. 25(3–4): 44.1974; B.D.Sharma et al., Fl. Karnataka Anal. 239. 1984.

Cinnamomum cathia D.Don, Prodr. Fl. Nepal. 66.1825

Phoebe paniculata (Nees) Nees, Syst. Laurin. 105. 1836

Distribution: Karnataka (Chikkamagaluru, Kodagu, Kolar), Andhra Pradesh, Assam, Kerala, Tamil Nadu; Myanmar, Nepal, Pakistan.

Local/Common/Vernacular name: *Kara mara* (Kannada)

Phoebe wightii Meisn. in DC., Prodr. 15: 38. 1864; B.D.Sharma et al., Fl. Karnataka Anal. 239. 1984.

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Lecythidaceae**Barringtonia** J. R. Forst. & G. Forst.

Barringtonia acutangula (L.) Gaertn., Fruct. 2: 97, t.101. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 105. 1984. **Plate 8F**

Eugenia acutangula L., Sp. Pl. 471. 1753

Distribution: Karnataka (Belagavi, Mysuru, Uttara Kannada), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Mizoram, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; Australia, tropical Asia

Local/Common/Vernacular name: *Indian oak* (English); *Neeru kanigalu* (Kannada)

Barringtonia racemosa (L.) Spreng., Syst. Veg. 3:127.1826; B.D.Sharma et al., Fl. Karnataka Anal. 105. 1984.

Eugenia racemosa L., Sp. Pl. 471. 1753

Distribution: Karnataka (Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Kerala, Maharashtra, Tamil Nadu; E Africa to W Pacific

Local/Common/Vernacular name: *Seaside Indian oak* (English); *Samudra phala* (Kannada)

Careya Roxb.

Careya arborea Roxb., Pl. Coromandel 3: 14, t. 218. 1811; B.D.Sharma et al., Fl. Karnataka Anal. 105. 1984. **Plate 8G**

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), throughout India; Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Ceylon oak, Slow Match Tree* (English); *Gobbala kayi, Kavulu mara* (Kannada)

Couroupita Aubl.

Couroupita guianensis Aubl., Hist. Pl. Guiane 708. t. 282. 1775; B.D.Sharma et al., Fl. Karnataka Anal. 105. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Davanagere, Mysuru), cultivated throughout India; native to S America

Local/Common/Vernacular name: *Cannon-ball tree* (English); *Shivalinga pushpa* (Kannada)

Leguminosae-Caesalpinoideae**Acacia** Mill.

Acacia auriculiformis Benth., London J. Bot. 1: 377. 1842; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Racosperma auriculiforme (Benth.) Pedley, Bot. J. Linn. Soc. 92(3): 247.1986

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Australia; cultivated in tropics.

Local/Common/Vernacular name: *Bengalijali* (Kannada)

Acacia decurrens Willd., Sp. Pl. 4(2): 1072. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Racosperma decurrens (Willd.) Pedley, Austrobaileya 2(4): 358.1987

Distribution: Karnataka (Mysuru), Himachal Pradesh, Manipur, Sikkim, Tamil Nadu, Uttar Pradesh, West Bengal; native to Australia.

Acacia mangium Willd., Sp. Pl. 4(2): 1053. 1806

Racosperma mangium (Willd.) Pedley, Austrobaileya 2(4): 352. 1987

Distribution: Karnataka (cultivated in Chitradurga), cultivated in Delhi, Pondicherry; native to Australia, Indonesia, Papua New Guinea.

Local/Common/Vernacular name: *Pencil mara* (Kannada)

Acrocarpus Wight ex Arn.

Acrocarpus fraxinifolius Arn. in Mag. Zool. Bot. 2: 547. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 86. 1984.

Distribution: Karnataka (cultivated in Belagavi, Dakshina Kannada, Kodagu, Shivamogga), Assam, Kerala, Maharashtra, Meghalaya, Mizoram, Sikkim, Tamil Nadu, West Bengal; S & SE Asia.

Local/Common/Vernacular name: *Houtige mara* (Kannada)

Adenanthera L.

Adenanthera pavonina L., Sp. Pl. 384. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Distribution: Karnataka (Belagavi, Chitradurga, Kalburgi, Mysuru, Uttara Kannada), Andaman & Nicobar Islands, Assam, Bihar, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; China, S & SE Asia; introduced and cultivated in other tropical countries

Local/Common/Vernacular names: *Red Bead Tree, Red Sandalwood* (English); *Aane gulaganji* (Kannada)

Albizia Durazz.

Albizia amara (Roxb.) B.Boivin, Encycl. 19: 34. 1838; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Plate 7B

Mimosa amara Roxb., Pl. Coromandel 2:13, t.122. 1799

Mimosa pulchella Roxb., Fl. Ind.2: 548. 1832

Acacia wightii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 274. 1834

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Tumakuru), Andhra Pradesh, Delhi, Goa, Gujarat, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Tamil Nadu, Uttar Pradesh; E Africa, Sri Lanka.

Local/Common/Vernacular name: *Oil cake tree* (English); *Chigare* (Kannada)

Albizia chinensis (Osbeck) Merr., Amer. J. Bot. 3: 575. 1916; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Plate 7C

Mimosa chinensis Osbeck, Dabgok. Ostind. Resa 233. 1757

Mimosa marginata Lam., Encycl.1(1): 12. 1783

Acacia stipulata DC., Prodr. 2: 469. 1825

Mimosa stipulacea Roxb., Fl. Ind. 2: 549. 1832

Albizia stipulata (DC.) Boivin, Encycl. 19(2): 33. 1838

Albizia marginata (Lam.) Merr., Philipp. J. Sci. 5: 23. 1910

Distribution: Karnataka (Belagavi, Chikkamagalur, Dakshina kannada, Davanagere, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), throughout India; China, Indo-China, Papua New Guinea, S & SE Asia.

Local/Common/Vernacular name: *Stipulate siris* (English); *Hottebage* (Kannada)

Albizia julibrissin Durazz. in Mag. Tosc. 3. 4. 11. 1772; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

var. **julibrissin**

Distribution: Karnataka (Bidar, Kalburgi), Arunachal Pradesh, Assam, Himachal Pradesh, Jammu & Kashmir, Manipur, Meghalaya, Sikkim, Tripura, Uttar Pradesh, Uttarakhand; Indonesia, Iran, Iraq, E & S Asia; cultivated elsewhere.

Local/Common/Vernacular names: *Silktree, Silktree Albizia* (English)

var. **mollis** (Wall.) Benth., London J. Bot. 3: 91. 1844; N.P.Singh, Fl. East. Karnataka 1: 295. 1988.

Albizia mollis (Wall.) Boivin, Encycl. 19(2): 33. 1838

Albizia julibrissin subsp. *mollis* (Wall.) Ali & Quraishi, Pakistan J. Forest. 17(3): 368. 1967

Distribution: Karnataka (Ballari, Bidar), Himachal Pradesh, Jammu & Kasmir, Uttarakand, Sikkim; Bhutan, Nepal.

Albizia lathamii Hole, Indian Forest Rec. 4. 153. 1913; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Distribution: Karnataka (Ballari, Kalburgi, Tumakuru), Kerala, Madhya Pradesh, Tamil Nadu; endemic.

Albizia lebbeck (L.) Benth., London J. Bot. 3: 87. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Plate 7D

Mimosa lebbeck L., Sp.Pl.516. 1753

Acacia lebbeck(L.) Willd., Sp. Pl. 4: 1066. 1806

Mimosa sirissa Roxb., Fl. Ind. 2: 544.1832

Feuilleea lebbeck(L.) Kuntze, Revis.Gen.Pl. 1: 188.1891

Albizia lebbeck var. *pubescens* Haines, Bot. Bihar & Orissa 3:332.1922

Albizia lebbeck var. *rostrata* Haines, Bot. Bihar & Orissa 3:332.1922

Distribution: Karnataka (throughout), throughout India; paleotropical,
Local/Common/Vernacular name: *Bage mara* (Kannada)

Albizia odoratissima (L.f.) Benth., London J. Bot. 3: 88. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Mimosa odoratissima L. f., Suppl. Pl. 437. 1781

Acacia odoratissima (L.f.) Willd., Sp. Pl., ed. 4, 4(2): 1063. 1806

Albizia odoratissima var. *mollis* Baker in Hook.f., Fl.Brit.India 2:299.1878

Feuilleea odoratissima (L.f.) Kuntze, Revis. Gen. Pl. 1: 188.1891

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Uttara Kannada), throughout India; China, Indo-China, S & SE Asia.

Albizia procera (Roxb.) Benth., London J. Bot. 3: 89. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Mimosa procera Roxb., Pl. Coromandel 1:12, t. 121. 1799

Acacia procera (Roxb.) Willd., Sp. Pl., ed. 4, 4(2): 1063. 1806

Mimosa elata Roxb., Fl. Ind. 2: 546.1832

Acacia elata (Roxb.) Voigt, Hort. Suburb. Calcutt. 261. 1845

Acacia procera var. *elata* (Roxb.) Baker in Hook.f., Fl.Brit.India 2:299.1878

Feuilleea procera (Roxb.) Kuntze, Revis. Gen. Pl. 1: 188.1891

Distribution: Karnataka (Belagavi, Bidar, Chikkamagaluru, Dharwad, Kalburgi, Uttara Kannada), throughout India; Australia, China, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Hill Siris*, *Silver Bark Rain Tree*, *Tall Albizia*, *White Siri* (English); *Belati* (Kannada)

Albizia saman (Jacq.) Merr., Select Pl. 12. 1876

Mimosa saman Jacq., Frigm. Bot. 15, pl. 9. 1800; N.P.Singh, Fl. East. Karnataka 1: 300. 1988

Inga saman (Jacq.) Willd., Sp. Pl., ed. 4, 4(2): 1024. 1806

Pithecellobium saman (Jacq.) Benth., London J. Bot. 3: 216. 1844

Feuilleea saman (Jacq.) Kuntze, Revis. Gen. Pl. 1: 189. 1891

Enterolobium saman (Jacq.) Prain, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 66: 252. 1897

Zygia saman (Jacq.) Lyons, Pl. Names Sci. & Pop ed. 2, 503.1907

Samanea saman (Jacq.) Merr., J. Wash. Acad. Sc. 6: 47. 1916

Distribution: Karnataka (cultivated throughout), cultivated throughout; native to C & S America.

Local/Common/Vernacular name: *Rain tree* (English)

Archidendron F.Muell

Archidendron bigeminum (L.) I.C.Nielsen, Opera Bot. 76: 73. 1985

Mimosa bigemina L., Sp. Pl. 1: 517. 1753; Yogan et al., Fl. Chikmagalur 132. 1981

Mimosa monadelpha Roxb., Fl. Ind. 2: 544. 1832

Pithecellobium bigeminum (L.) Mart., Flora 20(2, Beibl.): 115. 1837

Pithecellobium gracile Bedd., Madras J. Lit. Sci. Ser. 3, 44. 1864

Pithecellobium nicobaricum Prain, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 66(1): 267. 1897

Abarema bigemina (L.) Kosterm., Bull. Org. Sci. Res. Indonesia No. 20, 51. 1954

Abarema monadelpha (Roxb.) Kosterm., Reinwardtia 3: 11. 1954

Abarema monadelpha var. *gracile* (Bedd.) Kosterm., Bull. Org. Sci. Res. Indonesia No. 20, 51. 1954

Archidendron monadelphum (Roxb.) I.C.Nielsen, Adansonia 19: 21. 1979

Archidendron monadelphum var. *gracile* (Bedd.) Sanjappa, Legumes of India 62. 1992

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Goa, Karnataka, Kerala, Maharashtra, Meghalaya, Mizoram, Sikkim, Tamil Nadu, Tripura, West Bengal; S Asia.

Local/Common/Vernacular names: *Doldatte mara*, *Kadu kondemara* (Kannada)

Biancaea Tod.

Biancaea decapetala (Roth) O.Deg., Fl. Hawaiiensis K7. 1936; E.Gagnon & G.P.Lewis, PhytoKeys 71: 69. 2016

Reichardia decapetala Roth, Nov. Pl. Sp. 212. 1821

Caesalpinia sepiaria Roxb., Fl. Ind. 2: 360. 1832

Caesalpinia decapetala (Roth) Alston in Trimen, Handb. Fl. Ceylon 6(Suppl.) 89. 1931; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Plate 7F

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kolar, Mysuru, Shivamogga), throughout tropical parts; tropical Asia; introduced in Africa, Australia, Caribbean Islands.

Local/Common/Vernacular name: *Mysore thorn* (English); *Kurudu gejjuga* (Kannada)

Biancaea sappan (L.) Tod., Hort. Bot. Panorm. 1(1): 3. 1875

Caesalpinia sappan L., Sp. Pl. 381. 1753; C.J.Saldanha, Fl. Karnataka 1: 380. 1984.

Distribution: Karnataka (cultivated), Arunachal Pradesh, Assam, Bihar, Goa, Himachal Pradesh, Kerala, Maharashtra, Punjab, Sikkim, Tamil Nadu, Uttar Pradesh, West Bengal; China, Indo-China, S & SE Asia.

Local/Common/Vernacular name: *Sappan wood* (English); *Patanga-Chekke Sappanga* (Kannada)

Caesalpinia L.

Caesalpinia crista L., Sp. Pl. 380. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Guilandina nuga L., Sp. Pl. (ed. 2) 546. 1762

Caesalpinia nuga (L.) Aiton, Hort. Kew (ed.2) 73: 32. 1811

Caesalpinia paniculata (Lam.) Roxb., Prodr. 2: 482. 1825

Distribution: Karnataka (Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Hassan, Mysuru, Raichur, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Bihar, Delhi, Goa, Maharashtra, Odisha, Tamil Nadu, Uttar Pradesh; SE Asia.

Local/Common/Vernacular name: *Sanna gejjuga* (Kannada)

Caesalpinia globulorum Bakh.f. & P.Royen, Blumea 12: 62. 1963

Caesalpinia major (Medik.) Dandy & Exell, J. Bot. 76: 180. 1938; Udayan & Ravikumar, Indian J. For., 26: 385. 2003.

Distribution: Karnataka (Dakshina Kannada), Madhya Pradesh; Indo-China, Madagascar, Maldives, Pacific Islands, S & SE Asia.

Caesalpinia pulcherrima (L.) Sw., Obs. 166. 1791; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Poinciana pulcherrima L., Sp. Pl. 1: 380. 1753

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Mysuru, Tumakuru), Andaman & Nicobar Islands, Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Goa, Gujarat, Madhya Pradesh, Maharashtra, Manipur, Mizoram, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; native to West Indies.

Local/Common/Vernacular names: *Dwarf Poinciana, Mexican Bird of Paradise, Peacock Flower, Poinciana, Pride of Barbados, Red Bird of Paradise* (English); *Ratnagandhi* (Kannada)

Calliandra Benth.

Calliandra brevipes Benth. in Hooker's J. Bot. 2: 140. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 94. 1984.

Distribution: Karnataka (Kodagu, Mysuru), cultivated throughout India; native to SE Brazil, Uruguay and N Argentina.

Local/Common/Vernacular name: *Pink Powderpuff* (English)

Calliandra haematocephala Hassk., Retzia 1: 216. 1855; Seeth. et al., Fl. 75. 2000.

Distribution: Karnataka (cultivated throughout), cultivated throughout; native to tropical America.

Local/Common/Vernacular names: *Blood red tassel flower, Red powder puff, Rose Cascade* (English)

Calliandra tweedii Benth. in Hook. London J. Bot. 2. 140. 1840; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated throughout India; native to

Local/Common/Vernacular names: *Mexican Flame Bush, Red Flash, Red Tassel Flower* (English)

Cassia L.

Cassia fistula L., Sp. Pl. 377. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Cassia rhombifolia Roxb., Fl. Ind. 3: 334. 1832

Distribution: Karnataka (throughout), throughout India; S & SE Asia; cultivated throughout tropics.

Local/Common/Vernacular names: *Indian Laburnum, Pudding-pipe Tree, Purging Cassia, Purging Fistula* (English); *Kakke gida* (Kannada)

Cassia javanica L., Sp. Pl. 1: 379. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Distribution: Karnataka (Kalburgi, Kodagu, Mysuru), throughout India; Australia, China, Indo-China, S & SE Asia; elsewhere cultivated.

Local/Common/Vernacular names: *Pink and White Shower, Pink Cassia* (English)

Cassia renigera Wall. ex Benth., Trans. Linn. Soc. 27: 518. 1871; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Cassia javanica subsp. *renigera* (Wall. ex Benth.) K.Larsen, Nordic J. Bot. 13(4): 404. 1993

Distribution: Karnataka (cultivated in Bengaluru, Mysuru), cultivated in Bihar, Delhi, Madhya Pradesh, Maharashtra, Manipur, Mizoram, Nagaland, Punjab, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; native to Myanmar, cultivated elsewhere.

Local/Common/Vernacular names: *Burmese Cassia, Burmese Pink Cassia* (English)

Cassia roxburghii DC., Prodr. 2: 489. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Cassia marginata Roxb., Fl. Ind. 2: 338. 1832

Distribution: Karnataka (Belagavi, Bengaluru, Kalburgi, Kodagu, Mysuru), Andhra Pradesh, Bihar, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Pondicherry, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Sri Lanka; introduced elsewhere.

Local/Common/Vernacular name: *Red Cassia* (English)

Chamaecrista Moench

Chamaecrista absus (L.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 664. 1982

Cassia absus L., Sp. Pl. 376. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Senna absus (L.) Roxb., Fl. Ind. 2: 340.1832

Grimaldia absus (L.) Britton & Rose, N. Amer. Fl. 23(5): 299. 1930

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Tamil Nadu; pantropical.

Local/Common/Vernacular name: *Kadu huliga* (Kannada)

Chamaecrista kleinii (Wight & Arn.) V.Singh, J. Econ. Taxon. Bot. 16(3): 600.1992

Cassia kleinii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 293. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Distribution: Karnataka (Bengaluru, Chitradurga, Dakshina Kannada, Kalburgi, Shivamogga), Andhra Pradesh, Bihar, Gujarat, Himachal Pradesh, Kerala, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Sri Lanka.

Local/Common/Vernacular name: *Klein's dwarf senna* (English)

Chamaecrista lechenaultiana (DC.) Degener, Fl. Haw., Fam. 169b. 1934

Cassia lechenaultiana DC., Mem. Soc. Phys. Geneve 2: 132. 1824 ('lechenaultiana').

Cassia wallichiana DC., Mém. Soc. Phys. Genève 2: 132.1824

Cassia mimosoides subsp. *lechenaultiana* (DC.) Ohashi, J. Jap. Bot. 50(10): 308. 1975

Chamaecrista wallichiana (DC.) V.Singh, J. Econ. Taxon. Bot. 16(3): 600.1992

Distribution: Karnataka (Uttara kannada, Udupi), Cambodia, Indonesia, Laos, Malaysia, Myanmar, Papua New Guinea, Thailand, Vietnam.

Chamaecrista mimosoides (L.) Greene, Pittonia 4:27.1899

Cassia mimosoides L., Sp. Pl. 379. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Distribution: Karnataka (throughout), throughout India; paleotropical.

Local/Common/Vernacular name: *Mimosa like senna* (English)

Chamaecrista nigricans (Vahl) Greene, Pittonia 4: 30. 1899

Cassia nigricans Vahl, Symb. Bot. 1: 30. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Distribution: Karnataka (Dakshina Kannada, Shivamogga, Uttara Kannada), Andhra Pradesh, Kerala, Maharashtra, Tamil Nadu; Middle East, tropical Africa.

Chamaecrista pumila (Lam.) K.Larsen, Nordic J. Bot. 13. 1993

Cassia pumila Lam., Encycl. 1: 651. 1785; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Distribution: Karnataka (Belagavi, Bidar, Chitradurga, Dharwad, Kalburgi, Mysuru, Raichur, Tumakuru, Uttara Kannada, Vijayapura), throughout India; Australia, China, Indo-China, S & SE Asia.

Local/Common/Vernacular name: *Dwarf senna* (English); *Nela tagache* (Kannada)

Chamaecrista rotundifolia (Pers.) Greene, Pittonia 4: 31. 1899; Rao et al., Rheedia 22: 66. 2012

Cassia rotundifolia Pers., Syn. Pl. 1: 456. 1805

Cassia bifoliolata DC. in Collad., Hist. Nat. Mid.Casses 120. t. 9.1816

Chamaecrista bifoliolata (Collad.) Greene, Pittonia 4: 31. 1899

Distribution: Karnataka (Bengaluru); native to tropical America; introduced elsewhere.

Colvillea Bojer

Colvillea racemosa Bojer, Bot. Mag. tt. 3325, 3326. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 90. 1984.

Distribution: Karnataka (Bengaluru, Kolar, Mysuru), Bihar, Maharashtra, Tamil Nadu, Uttar Pradesh, West Bengal; widely cultivated in tropics; native to Madagascar.

Local/Common/Vernacular name: *Colville's Glory* (English)

Delonix Raf.

Delonix elata (L.) Gamble, Fl. Madras 396(280). 1919; B.D.Sharma et al., Fl. Karnataka Anal. 90. 1984.

Poinciana elata L., Cent. Pl. 2: 16. 1756

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Davanagere, Kalburgi), Andhra Pradesh, Bihar, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; native to Africa

Local/Common/Vernacular names: *White Gulmohar* (English); *Bili gulmohr, Kempukkenjiga, Nirangi, Vatanarayana, Vayunarayani gida* (Kannada)

Delonix regia (Hook.) Raf., Fl. Tell. 2: 92. 1836; B.D.Sharma et al., Fl. Karnataka Anal. 90. 1984.

Poinciana regia Hook., Bot. Mag. 56: t. 2884. 1829

Distribution: Karnataka (cultivated throughout), throughout India; cultivated throughout tropics; native to Madagascar.

Local/Common/Vernacular names: *Flamboyant tree, Flame tre, Royal Poinciana* (English); *Kempu torai* (Kannada)

Desmanthus Willd.

Desmanthus pernambucanus (L.) Thell., Mém. Soc. Sci. Nat. Math. Cherbourg, sér. 4. 38: 256. 1912

Mimosa pernambucana L., Sp. Pl. 1: 519. 1753

Desmanthus virgatus (L.) Willd., Sp. Pl. 4: 1047. 1806; C.J.Saldanha, Fl. Karnataka 1: 405. 1984

Distribution: Karnataka (Bengaluru), introduced and naturalised throughout India; cultivated and also naturalised throughout tropics; native to tropical America, pantropical.

Local/Common/Vernacular names: *Acacia savane Desmanto, Acacia courant, Dwarf koa, Prostrate bundleflower, Wild tantan* (English)

Dichrostachys (DC.) Wight & Arn.

Dichrostachys cinerea (L.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 271. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Mimosa cinerea L., Sp. Pl. 520. 1753

Cailliea cinerea (L.) Roberty, Bull. Inst. Franc. Afr. Noire, ser. A, 16: 345, 1954

Distribution: Karnataka (Ballari, Bengaluru, Bidar, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kolar, Mysuru, Raichur, Tumakuru), Andhra Pradesh, Delhi, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Australia, Africa, Indonesia, Middle East, Sri Lanka, Thailand.

Local/Common/Vernacular name: *Vaduvarada gida* (Kannada)

Entada Adans.

Entada rheedii Spreng., Syst. Veg. ed. 16, 2: 325. 1825

Mimosa entada L., Sp. Pl. 1: 518. 1753

Entada monostachya DC., Prodr. 2: 425. 1825

Entada pursaetha DC., Prodr. 2: 425. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Entada scandens auct. non Benth., 1841; Hook.f., Fl. Brit. India 2: 86. 1878.

Pusaetha entada Kuntze, Revis. Gen. Pl. 1: 204. 1891

Distribution: Karnataka (Belagavi, Dakshina Kannada, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Arunachal Pradesh, Bihar, Goa, Kerala, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Mizoram, Nagaland, Odisha, Sikkim, Tamil Nadu, Tripura, West Bengal; Australasia, China, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Bell mimosa, Chinese lantern tree, Kalahari Christmas tree Sicklebush* (English); *Ganape kai* (Kannada)

Guillandina L.

Guillandina bonduc L., Sp. Pl. 381. 1753

Caesalpinia crista Thunb., Fl. Jap. 179. 1784.

Caesalpinia bonducella (L.) Fleming, Asiat. Res. 11: 159. 1810

Caesalpinia bonduc (L.) Roxb., Fl. Ind. 2: 362. 1832; C.J.Saldanha, Fl. Karnataka 1: 378. 1984

Distribution: Karnataka (throughout), throughout India; Pantropical.

Local/Common/Vernacular names: *Bonduc Nut* (English); *Gajjaga, Heggejuga* (Kannada)

Haematoxylum Gronov.

Haematoxylum campechianum L., Sp. Pl. 1: 384. 1753; C.J.Saldanha, Fl. Karnataka 1: 394. 1984.

Distribution: Karnataka (Bengaluru), Bihar, Delhi, West Bengal; native to S Mexico & northern C America.

Local/Common/Vernacular names: *Bloodwoodtree, Llogwood* (English)

Hultholia E.Gagnon & G.P.Lewis

Hultholia mimosoides (Lam.) E.Gagnon & G.P.Lewis, PhytoKeys 71: 61. 2016.

Caesalpinia mimosoides Lam., Encycl. 1: 452.1785; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Caesalpinia resupinata Roxb., Fl. Ind. 2: 362.1832

Caesalpinia simora Roxb., Fl. Ind. 2: 359.1832

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Goa, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Tripura; Bangladesh, China, Laos, Myanmar, Thailand, Vietnam.

Local/Common/Vernacular name: *Kenjige mullu* (Kannada)

Leucaena Benth.

Leucaena leucocephala (Lam.) de Wit, Taxon 10: 54. 1961 & 24: 352. 1975; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Mimosa leucocephala Lam., Encycl. 1(1): 12. 1783

Acacia leucocephala (Lam.) Link, Enum. Hort. Berol. Alt. 2: 444. 1822

Leucaena glauca Benth. in Hooker's J. Bot. 4: 416. 1842

Distribution: Karnataka (cultivated throughout),cultivated and also as an escape throughout India; native to tropical America; pantropical.

Local/Common/Vernacular names: *Jumbay, White leadtree, White popinac* (English); *Psubabul* (Kannada)

Libidibia (DC.) Schltdl

Libidibia coriaria (Jacq.) Schltdl., Linnaea 5: 193. 1830; E.Gagnon & G. P. Lewis, Phytokeys 71: 95.2016

Poinciana coriaria Jacq., Select. Am. 123. t. 175. f. 36. 1781

Caesalpinia coriaria (Jacq.) Willd., Sp. Pl. 2: 532. 1799; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Davanagere, Dharwad, Kalburgi, Kodagu, Mysuru, Uttara Kannada, Vijayapura), Andaman & Nicobar Islands, Andhra Pradesh, Bihar, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu,Uttar Pradesh, West Bengal; native to tropical America

Local/Common/Vernacular name: *Divi Divi* (English); *Seeme Alde Mara* (Kannada)

Libidibia ferrea (Mart. ex Tul.) L.P.Queiroz, Leguminosas da Caatinga: 130. 2009; E.Gagnon & G.P.Lewis, Phytokeys 71: 98.2016.

Caesalpinia ferrea C. Mart, Reise Bras. 2: 611. 1828; Sen & Naskar, Bull. Bot. Surv. India 7: 36. 1965.

Distribution: Karnataka (Bengaluru), Gujarat, Kerala, Maharashtra, Tamil Nadu, W Bengal; native to Brazil, cultivated elsewhere in tropics.

Lysiloma Benth.

Lysiloma latisiliquum (L.) Benth., Trans. Linn. Soc. London 30: 534. 1875

Mimosa latisiliqua L., Sp. Pl. 1: 519. 1753

Acacia latisiliqua (L.) Willd., Sp. Pl., ed. 4, 4(2): 1067. 1806

Leucaena latisiliqua (L.) Gillis & Stearn, Taxon 23: 190. 1974

Distribution: Karnataka (Chitradurga, Dharwad); native to Caribbean, C America, Mexico, USA.

Local/Common/Vernacular names: *False tamarind*, *Wild tamarind* (English)

Mezoneuron Wight & Arn.

Mezoneuron cucullatum (Roxb.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 283. 1834

Caesalpinia cucullata Roxb., Fl. Ind. 2: 358. 1832; C.J.Saldanha, Fl. Karnataka 1: 391. 1984.

Distribution: Karnataka (Belagavi, Chikkamagaluru, Hassan, Kodagu, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Bihar, Kerala, Maharashtra, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim, Tamil Nadu, West Bengal; China, Indo-China, S & SE Asia.

Local/Common/Vernacular name: *Brasiletto climber* (English); *Mullodu balli* (Kannada)

Mimosa L.

Mimosa diplosticha Sauvalle, Anal. Acad. Ci. Habana v. 405. 1868

Mimosa invisa C.Mart., Flora 20: Biebl. 8: 121. 1837; C.J.Saldanha, Fl. Karnataka 1: 408. 1984.

Distribution: Karnataka (Cultivated), Kerala, Tamil Nadu; native to tropical America.

Mimosa hamata Willd., Sp. Pl. 4: 1033. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bidar, Chitradurga, Dharwad, Kalburgi, Mysuru, Raichur, Shivamogga), Andhra Pradesh, Delhi, Gujarat, Haryana, Kerala, Madhya Pradesh, Maharashtra, Pondicherry, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Pakistan.

Local/Common/Vernacular name: *Sagari mullu* (Kannada)

Mimosa prainiana Gamble, Fl. Madras 421. 1919; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Distribution: Karnataka (Ballari, Bidar, Kalburgi, Raichur, Vijayapura), Andhra Pradesh, Bihar, Gujarat, Madhya Pradesh, Maharashtra, Tamil Nadu; endemic.

Mimosa pudica L., Sp. Pl. 518. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Distribution: Karnataka (throughout), throughout India; native to tropical America; pantropical.

Local/Common/Vernacular name: *Touch me not* (English); *Muttidare Muni* (Kannada)

Mimosa rubicaulis Lam., Encycl. 1: 20. 1783; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Mimosa octandra Roxb., Pl. Coromandel 2:55, t. 200. 1805

Mimosa mutabilis Roxb., Fl. Ind. 2:564.1832

Distribution: Karnataka (Bengaluru, Mysuru), throughout India; Afghanistan.

Local/Common/Vernacular names: *Rasne*, *Uri seege* (Kannada)

Moullava Adans.

Moullava spicata (Dalzell) Nicolson in Manilal, Bot. Hist. Hort. Malab. 181. 1981; C.J.Saldanha, Fl. Karnataka 1: 392. 1984.

Plate 24G

Caesalpinia spicata Dalzell in Hooker's J. Bot. Kew Gard. Misc. 3: 90. 1851

Wagatea spicata (Dalzell) Wight, Icon. Pl. Ind. Orient. 6: t. 1995. 1853

Distribution: Karnataka (Chikkamagaluru, Dakshina Kannada, Udupi, Uttara Kannada), Goa, Gujarat, Kerala, Maharashtra; endemic.

Local/Common/Vernacular name: *Orange brasiletto climber* (English); *Gajjiga balli* (Kannada)

Neptunia Lour.

Neptunia oleracea Lour., Fl. Cochinch. 654. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Mimosa natans L.f., Suppl. Pl. 439. 1782

Mimosa prostrata Lam., Encycl. 1(1): 10. 1783

Desmanthus natans Willd., Sp. Pl. 4: 1044. 1806

Neptunia plena Lindl. in Edwards's Bot. Reg. t 3.1846

Neptunia prostrata (Lam.) Baill., Bull. Mens. Soc. Linn. Paris 1: 356. 1883

Neptunia natans (L.f.) Druce, Rep. Bot. Exch. Cl. Brit. Isles 1916, 637.1917

Distribution: Karnataka (Belagavi, Chitradurga, Uttara Kannada), throughout India; pantropical.

Neptunia triquetra (Vahl) Benth. in Hooker's J. Bot. Kew Gard. Misc. 4: 355. 1842; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Mimosa triquetra Vahl, Symb. Bot. 3: 102. 1794

Desmanthus triquetrus Willd., Sp. Pl. 4: 1045. 1806

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Dharwad, Kalburgi, Vijayapura), Andhra Pradesh, Dadra-Nagar-Haveli, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Pondicherry, Punjab, Rajasthan, Tamil Nadu; Myanmar.

Parkia R. Br.

Parkia biglandulosa Wight & Arn., Prodr. Fl. Ind. Orient. 1: 279. 1834; N.P.Singh, Fl. East. Karnataka 1: 300. 1988.

Mimosa pedunculata Roxb., Fl. Ind. 2:551.1832

Parkia pedunculata (Roxb.) J.F.Macbr., Contr. Gray Herb. 59: 19. 1919

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to Myanmar.

Local/Common/Vernacular name: *Tennis ball tree* (English)

Parkia timoriana (DC.) Merr., Philipp. J. Sci., C 5: 33. 1910

Inga timoriana DC., Prodr. 2: 442. 1825

Parkia roxburghii G.Don, Gen. Hist. 2: 397. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 95. 1984.

Distribution: Karnataka (Mysuru), Arunachal Pradesh, Assam, Meghalaya, Mizoram, Tamil Nadu, Tripura, West Bengal; Bangladesh, Myanmar, Thailand, SE Asia.

Parkinsonia L.

Parkinsonia aculeata L., Sp. Pl. 375. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Mysuru), cultivated throughout drier parts of India; native to tropical N America.

Local/Common/Vernacular names: *jelly bean tree, Jerusalem thorn* (English)

Peltophorum (Vogel) Benth.

Peltophorum africanum Sond., Linnaea 23: 35. 1850; Basu & S.N. Chakravorty, J. Econ. Taxon. Bot. 10: 173. 1987.

Distribution: Karnataka (cultivated in gardens), Delhi, Tamil Nadu, Uttarakhand, West Bengal); native to Southern tropical and S Africa.

Peltophorum dasyrachis (Miq.) Baker in Hook.f., Fl. Brit. India2: 257. 1878; Gowda & Krishnasw., Pl. Wealth Lalbagh 112. 1968.

Caesalpinia dasyrachis Miq., Fl. Ind. Bat. suppl. 29. 1861.

Distribution: Karnataka (cultivated in Lalbagh, Bengaluru); native to Indonesia.

Peltophorum dubium (Spreng.) Taub. in Engl. & Prantl, Nat. Pflanzenfam. 3, 3: 176. 1892; Gowda & Krishnasw., Pl. Wealth Lalbagh 112. 1968.

Caesalpinia dubia Spreng., Syst. Veg. 2: 343: 1825.

Distribution: Karnataka (cultivated in Lalbagh, Bengaluru); native to Brazil.

Peltophorum pterocarpum (DC.) K.Heyne, Nutt. Pl. Ned. Ind. (ed.2) 2:755.1927; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Inga pterocarpa DC., Prodr. 2: 441. 1825

Peltophorum ferrugineum (Decne.) Benth., Fl. Australiensis 2: 279. 1864

Distribution: Karnataka (throughout), cultivated throughout India, wild in Andaman & Nicobar Islands; Australasia, SE Asia.

Local/Common/Vernacular names: *Copper pod, Flamboyant, Golden, Yellow Flamboyant, Yellow Flame Tree, Yellow Poinciana* (English)

Pithecellobium Mart.

Pithecellobium dulce (Roxb.) Benth. in London J. Bot. 3: 199. 1844; B.D.Sharma et al., Fl. Karnataka Anal. 96. 1984.

Mimosa dulcis Roxb., Pl. Coromandel 1: 67. t.99. 1798

Inga dulcis (Roxb.) Willd., Sp. Pl. 4: 1005. 1806

Albizia dulcis (Roxb.) F.Muell., Sel. Pl. Indust. Cult. 12. 1872

Feuilleea dulcis (Roxb.) Kuntze, Revis. Gen. Pl. 1: 184. 1891

Zygia dulcis (Roxb.) Lyons, Pl. Nam. Ed. 2: 503. 1907

Distribution: Karnataka (throughout), throughout India; native tropical America.

Local/Common/Vernacular names: *Madras thorn, Manila tamarind* (English); *Seeme hunase* (Kannada)

Prosopis L.

Prosopis cineraria (L.) Druce, Bot. Exch. Club. Soc. Isles 3: 422. 1914; B.D.Sharma et al., Fl. Karnataka Anal. 96. 1984.

Mimosa cineraria L., Sp. Pl. 2: 1500. 1753

Prosopis spicigera L., Mant. Pl. 68. 1767

Prosopis spicata Burm.f., Fl. Indica 102. 1768

Adenanthera aculeata Roxb., Fl. Ind. 2: 37.1832

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Mysuru, Tumakuru), Andhra Pradesh, Bihar, Delhi, Gujarat, Haryana, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal; Afghanistan, Middle East, Pakistan.

Local/Common/Vernacular name: *Banni* (Kannada)

Prosopis glandulosa Torr., Ann. Lyceum Nat. Hist. New York 2. 192. 1828; B.D.Sharma et al., Fl. Karnataka Anal. 96. 1984.

Algarobia glandulosa (Torr.) Torr. & A. Gray, Fl. N. Amer. 1(3): 399 399 1840

Neltuma glandulosa (Torr.) Britton & Rose, N. Amer. Fl. 23(3): 186. 1928

Distribution: Karnataka (Bengaluru), Delhi, Haryana, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; introduced in tropics; native to USA.

Prosopis juliflora (Sw.) DC., Prodr. 2: 447. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 96. 1984.

Mimosajuliflora Sw., Prodr. 85. 1788

Acacia juliflora (Sw.) Willd., Sp. Pl., ed. 4, 4(2): 1076. 1806

Algarobia juliflora (Sw.) Heynh., Alph. Aufz. Gew. 18. 1846

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Dharwad, Kalburgi, Raichur, Tumakuru, Vijayapura), throughout except NE India.; pantropical; native to tropical S America.

Local/Common/Vernacular name: *Mesquite* (English); *Ballari jali* (Kannada)

Pterolobium R. Br. ex Wight & Arn.

Pterolobium hexapetalum (Roth) Santapau & Wagh, Bull. Bot. Surv. India 5: 108. 1964; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Reichardia hexapetala Roth, Nov. Pl. Sp. 210. 1821

Caesalpinia lacerans Roxb., Fl. Ind. 2: 367. 1832

Pterolobium indicum A.Rich., Fl. Abyss. 1: 247. 1847

Distribution: Karnataka (Ballari, Bengaluru, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Tumakuru, Yadgiri), Andhra Pradesh, Kerala, Pondicherry, Sikkim, Tamil Nadu, Uttar Pradesh, West Bengal; Bangladesh, Bhutan, Myanmar.

Local/Common/Vernacular names: *The Indian Redwing, White brasiletto climber* (English); *Bada bakka* (Kannada)

Senegalia Raf.

Senegalia caesia (L.) Maslin, Seigler & Ebinger, Blumea 58(1): 40. 2013

Mimosa caesia L., Sp. Pl. 522. 1753

Acacia caesia (L.) Willd., Sp. Pl. 4: 1090. 1806; C.J.Saldanha, Fl. Karnataka 1: 396. 1984.

Acacia intsia auct. non (L.) Willd., Prodr. Fl. Ind. Orient. 1: 278. 1834

Acacia intsia var. *caesia* (L.) Baker in Hook.f. Fl. Brit. India 2(5): 297. 1878

Acacia columnaris Craib., Bull. Misc. Inform. Kew 1915(10): 410. 1915

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Bihar, Jharkhand, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, W Bengal; Bangladesh, Bhutan, Cambodia, China, Laos, Myanmar, Sri Lanka, Taiwan, Thailand, Vietnam.

Local/Common/Vernacular name: *Black catechu* (English); *Kadu seege* (Kannada)

Senegalia caffra (Thunb.) P.J.H.Hurter & Mabb., Mabberley's Pl.-Book 1021. 2008

Acacia caffra Willd., Sp. Pl. 4: 1078. 1806; Gowda & Krishnasw., Pl. Wealth Lalbagh 1. 1968.

Distribution: Karnataka (cultivated in Lalbagh, Bengaluru), native to tropical Africa.

Senegalia catechu (L.f.) P.J.H.Hurter & Mabb., Mabberley's Pl.-Book 1021. 2008

Mimom catechu L.f., Suppl. Pl. 439. 1781; Roxb., Pl. Coromandel 2: t 175. 1802.

Acacia catechu (L.f.) Willd., Sp. Pl. 4: 1079. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Acacia wallichiana DC., Prodr. 2: 458. 1825

Mimosa catechoides Roxb., Fl. Ind. 2: 562. 1832

Acacia catechoides (Roxb.) Benth., London J. Bot. 1: 510. 1842

Acacia catechu var. *catechoides* (Roxb.) Prain, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 66: 508. 1898

Acacia catechu var. *wallichiana* P.C. Huang, Sylva Sinica? 2: 1259, f. 579, 5-6. 1985

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chitradurga, Dharwad, Kalburgi, Mysuru, Uttara Kannada), throughout India; Bhutan, China, Myanmar, Nepal, Pakistan, Thailand.

Local/Common/Vernacular name: *Kaggali* (Kannada)

Senegalia chundra (Roxb. & Rottler) Maslin, Nuytsia 22(6): 466. 2012

Mimosa chundra Rottler, Ges. Naturf. Freunde Berlin Neue Schriften 4: 207. 1803

Acacia chundra (Rottler) Willd., Sp. Pl. 4: 1078. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Mimosa sundra Roxb., Pl. Coromandel 3: 19. 1811; Hort. Bengal. 41. 1814; Fl. Ind. ed. 1832, 2: 562. 1832.

Acacia sundra (Roxb.) DC., Prodr. 2: 458. 1825

Acacia catechu var. *sundra* (Roxb.) Prain, J. Asiat. Soc. Bengal Part 2, Nat. Hist. 66: 508. 1898

Distribution: Karnataka (throughout), Andhra Pradesh, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu; China, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Kach, Kempujali* (Kannada)

Senegalia ferruginea (DC.) Pedley, Bot. J. Linn. Soc. 92(3): 250. 1986

Acacia ferruginea DC., Prodr. 2: 458. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Mimosa ferruginea Roxb., Fl. Ind. 2: 561. 1832 non Rottl. 1803

Distribution: Karnataka (Ballari, Bengaluru, Chikkamagaluru, Davanagere, Hassan, Kalburgi, Kodagu, Kolar, Mysuru), Andhra Pradesh, Bihar, Dadra-Nagar-Haveli Delhi, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Myanmar, Sri Lanka.

Local/Common/Vernacular name: *Banni mara* (Kannada)

Senegalia hohenackeri (Craib) Ragupathy, Seigler, Ebinger & Maslin, Phytotaxa 162(3): 175. 2014

Acacia hohenackeri Craib, Bull. Misc. Inform. 1915(10):408. 1915; C.J.Saldanha, Fl. Karnataka 1: 398. 1984

Distribution: Karnataka (Kodagu, Uttara Kannada), Andhra Pradesh, Kerala, Tamil Nadu; endemic.

Senegalia intsia (L.) Maslin, Seigler & Ebinger, Blumea 58(1): 40. 2013

Mimosa cassia L., Sp. Pl. 522. 1753 pro parte.

Mimosa intsia L., Sp. Pl. 522. 1753

Acacia intsia (L.) Willd. in Hook.f., Fl. Brit. India 2: 297. 1878

Distribution: Karnataka (), throughout up to 1700 m in Himalayas; Sri Lanka, Myanmar, Malay Peninsula.

Senegalia megaladena (Desv.) Maslin, Seigler & Ebinger, Blumea 58:41. 2013

Acacia megaladena Desv., J. Bot. 3: 69. 1814; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Acacia arrophula D.Don, Prodr. Fl. Nepal. 247. 1825

Acacia pennata var. *arrophula* (D.Don) Baker, Hook.f., Fl. Brit. India 2: 298. 1878

Distribution: Karnataka (Belagavi, Hassan), Andaman & Nicobar Islands, Arunachal Pradesh, Jharkhand, Mizoram, Rajasthan, Sikkim, West Bengal; Bangladesh, Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Nepal, Philippines, Thailand, Vietnam.

Senegalia pennata (L.) Maslin, Nuytsia 22(6): 466. 2012

Mimosa pennata L., Sp. Pl. 522. 1753

Acacia pennata (L.) Willd., Sp. Pl. 4: 1090. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Acacia hainanensis Hayata, Icon. Pl. Formosan. 3: 86. 1913

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout India; China, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Rusty mimosa* (English); *Bili seege*, *Kaadu seege*, *Mullu seege* (Kannada)

Senegalia polyacantha (Willd.) Seigler & Ebinger, Phytologia 91(1): 28. 2009

Acacia polyacantha Willd., Sp. Pl. 4: 1079. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Mimosa suma Roxb., Fl. Ind. 2: 563. 1832

Acacia suma (Roxb.) Voigt, Hort. Suburb. Calcutt. 260. 1845

Senegalia suma (Roxb.) Britton & Rose, N. Amer. Fl. 23(2): 113. 1928

Distribution: Karnataka (Ballari, Belagavi, Bidar, Chitradurga, Dharwad, Hassan, Kolar, Mysuru, Tumakuru, Uttara Kannada), Andhra Pradesh, Bihar, Chattisgarh, Delhi, Goa, Gujarat, Jharkhand, Maharashtra, Tamil Nadu, West Bengal; Africa, Madagascar, Mauritius, Sri Lanka.

Local/Common/Vernacular names: *White cutch* (English); *Mugli*, *Sike gida*, *Sujalu* (Kannada)

Plate 7A

Senegalia rugata (Lam.) Britton & Rose, N. Amer. Fl. 23(2): 120. 1928

Mimosa rugata Lam., Encycl. 1:20. 1783

Mimosa sinuata Lour., Fl. Cochinch. 653. 1790

Mimosa concinna Willd., Sp. Pl. 4: 1039. 1806

Acacia concinna (Willd.) DC., Prodr. 2: 464. 1825

Acacia rugata Buch.-Ham. ex Benth., London J. Bot. 1: 514. 1842

Acacia concinna var. *rugata* (Benth.) Baker in Hook. f., Fl. Brit. India 2: 297. 1878

Acacia sinuata (Lour.) Merr., Trans. Amer. Philos. Soc. n.s.24: 186. 1935; B.D.Sharma et al., Fl.

Karnataka Anal. 93. 1984.

Acacia gamblei Bahadur & R.C.Gaur, Acta Bot. Indica 4: 67. 1976

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), throughout; Bhutan, Cambodia, China, Bangladesh, Indonesia, Laos, Myanmar, Nepal, Philippines, Papua New Guinea, Singapore, Thailand, Vietnam.

Local/Common/Vernacular names: *Soap pod*, *Washing pod* (English); *Seege*, *Shikekai* (Kannada)

Senegalia senegal (L.) Britton, Sci. Surv. Porto Rico & Virgin Islands 6: 538. 1930

Acacia senegal (L.) Willd., Sp. Pl., ed. 4(2): 1077. 1806

Acacia rupestris Stock ex Boiss., Fl. Orient. 2:638. 1853; Brandis, For. Fl. 184. 1874.

Distribution: Karnataka (Chitradurga), Delhi, Gujarat, Haryana, Madhya Pradesh, Punjab, Rajasthan, Uttar Pradesh; Arabia, Pakistan, tropical Africa.

Local/Common/Vernacular names: *Gum Acacia*, *Gum Arabic Tree*, *Gum Senegal Tree* (English); *Mugli* (Kannada)

Senegalia torta (Roxb.) Maslin, Seigler & Ebinger, Blumea 58:42.2013

Mimosa torta Roxb., Fl. Ind. 2: 566. 1832

Acacia torta (Roxb.) Craib, Bull. Misc. Inform. Kew 1915: 410. 1915; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Dakshina Kannada, Davanagere, Hassan, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), throughout except NE India; Pakistan, Thailand.

Local/Common/Vernacular name: *Kempu seege* (Kannada)

Senna Mill.

Senna alata (L.) Roxb., Fl. Ind. 2:2.1824

Cassia alata L., Sp. Pl. 378. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Herpetica alata (L.) Raf., Sylva Tellur. 123. 1838

Distribution: Karnataka (Belagavi, Chikkamagaluru, Chitradurga, Dakshina Kannada, Kodagu), throughout India, often naturalised; pantropical; native to S America.

Local/Common/Vernacular names: *Ringworm Cassia*, *Ringworm Plant Ringworm Shrub*, *Roman Candle Tree*, *Seven Golden Candlesticks*, *Winged Senna* (English)

Senna alexandrina Mill., Gard. Dict., ed. 8. n. 1. 1768

Cassia senna L., Sp. Pl. 377. 1753

Cassia angustifolia Vahl, Symb. Bot. 1: 20. 1790; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Cassia acutifolia Delile, Fl. Egypte 219. t. 27. 1813

Senna acutifolia (Delile) Batka, Proc. Linn. Soc. 2: 282.1854

Senna angustifolia (Vahl) Batka, Proc. Linn. Soc. 2: 282.1854

Distribution: Karnataka (Ballari, Davanagere, Kodagu) Andhra Pradesh, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Tamil Nadu, Uttar Pradesh; native to Africa and Arabia; introduced elsewhere.

Local/Common/Vernacular name: *Tinnevelly senna* (English); *Nelavarike* (Kannada)

Senna auriculata (L.) Roxb., Fl. Ind. 2: 349. 1832*Cassia auriculata* L., Sp. Pl. 379. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Raichur, Shivamogga, Tumakuru, Vijayapura), Andhra Pradesh, Bihar, Delhi, Goa, Gujarat, Haryana, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Odisha, Pondicherry, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal, Yanam; Myanmar, Sri Lanka; introduced elsewhere.

Local/Common/Vernacular names: *Eared senna*, *Tanner's cassia* (English); *Honnambri gida*, *Olle thangadi* (Kannada)

Senna floribunda (Cav.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 360. 1982*Cassia floribunda* Cav., Desc. Pl. 132. 1802; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.*Cassia leavigata* Willd., Enum. Hort. Berol. 441. 1809*Chamaefistula floribunda* (Cav.) G.Don, Gen. Hist. 2: 452. 1832

Distribution: Karnataka (Chikkamagaluru, Hassan, Kodagu, Shivamogga); native to tropical America; introduced elsewhere.

Senna hirsuta (L.) H.S.Irwin & Barneby, Phytologia 44: 499. 1979*Cassia hirsuta* L., Sp. Pl. 378. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.*Ditremexa hirsuta* (L.) Britton & Wilson, Sci. Surv. Porto Rico & Virgin Islands 5:372. 1924

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Shivamogga, Uttara Kannada), throughout India; pantropical; native to tropical America.

Local/Common/Vernacular name: *American hairy senna* (English)

Senna italica Mill., Gard. Dict. 1768; C.J.Saldanha, Fl. Karnataka 1: 383. 1984*Cassia obovata* Collad., Hist. Nat. Méd. Casses 92. t. 15. 1816*Senna obovata* (Collad.) Link, Handbuch 2: 140. 1831*Cassia obtusa* (Roxb.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 288. 1832*Senna obtusa* Roxb., Fl. Ind. 2: 344. 1832*Cassia italicica* (Mill.) F.W.Andrews, Fl. Anglo-Egypt. Sudan 2: 117. 1952

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Kalburgi, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Daman, Delhi, Diu, Gujarat, Haryana, Himachal Pradesh, Jammu & Kashmir, Kerala, Madhya Pradesh, Maharashtra, Punjab, Tamil Nadu; Africa, Middle East, S Asia; introduced elsewhere.

Local/Common/Vernacular name: *Italian senna* (English); *Neladavare* (Kannada)

Senna montana (Roth) V.Singh, J. Econ. Taxon. Bot. 16(3): 600. 1992*Cassia montana* Roth, Nov. Pl. Sp. 214. 1821; B.D.Sharma et al., Fl. Karnataka Anal. 88. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Chitradurga, Hassan, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada), Andhra Pradesh, Gujarat, Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Deccan senna* (English); *Betta thangadi* (Kannada)

Senna multiglandulosa (Jacq.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 357. 1982*Cassia tomentosa* L.f., Suppl. Pl. 231. 1781; B.D.Sharma et al., Fl. Karnataka Anal. 90. 1984.*Cassia multiglandulosa* Jacq., Icon. Pl. Rar. 1(3): 8, t.72. 1783*Adipera tomentosa* (L.f.) Britton & Rose, N. Amer. Fl. 23(4): 241. 1930

Distribution: Karnataka (Chikkamagaluru, Mysuru), Tamil Nadu; native tropical S America, introduced elsewhere in tropics.

Local/Common/Vernacular name: *Butter cup bush* (English)

Senna multijuga (Rich.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 492. 1982

Cassia multijuga Rich., Act. Soc. Hist. Nat. 1: 108. 1792; C.J.Saldanha, Fl. Karnataka 1: 383. 1984.

Peiranisia multijuga (Rich.) Britton & Rose, Sci. Surv. Porto Rico & Virgin Islands 6: 350. 1926

Chamaesenna multijuga (Rich.) Pittier, Third Conf. Interamer. Agric. Caracas 376.1945

Distribution: Karnataka (Dakshina Kannada), Uttar Pradesh, West Bengal; native to tropical America .

Local/Common/Vernacular name: *Leafy Cassia* (English)

Senna obtusifolia (L.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 252. 1982

Cassia obtusifolia L., Sp. Pl. 1: 377. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Senna toroides Roxb., Fl. Ind. 2: 341.1832

Diallobus tora (L.) Raf., Sylva Tellur. 128. 1838

Cassia tora var. *obtusifolia* (L.) Haines, Bot. Bihar & Orissa 3:304.1922

Distribution: Karnataka (Ballari, Belagavi, Dharwad, Kalburgi, Shivamogga, Vijayapura), Andhra Pradesh, Bihar, Dadra-Nagar-Haveli, Delhi, Goa, Gujarat, Haryana, Himachal Pradesh, Jammu-Kashmir, Kerala, Madhya Pradesh, Maharashtra, Odisha, Pondicherry, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; native to tropical America;introduced elsewhere in tropics.

Local/Common/Vernacular names: *Coffee Weed, Low Senna, Sickle pod* (English)

Senna occidentalis (L.) Link, Handbuch 2: 140. 1831

Cassia occidentalis L., Sp. Pl. 377. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Ditremexa occidentalis (L.) Britton & Wilson, Sci. Surv. Porto Rico & Virgin Islands 5: 372. 1924

Distribution: Karnataka (throughout), throughout India; native to tropical America;pan-tropical weed.

Local/Common/Vernacular names: *Coffee Senna, Coffee Weed, Negro coffee, Stinking Weed* (English); *Kolthogache* (Kannada)

Senna siamea (Lam.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35(1): 98. 1982

Cassia siamea Lam., Encycl. 1: 648. 1785; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Cassia floridula Vahl, Symb. Bot. 3: 57. 1794

Cassia sumatrana Roxb. ex Hornem., Suppl. Hort. Bot. Hafn. 135. 1819

Senna sumatrana Roxb., Fl. Ind. 2: 347.1832

Sciacassia siamea (Lam.) Britton & Rose, N. Amer. Fl. 23(4): 252. 1930

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kodagu, Kolar, Mysuru, Shivamogga), throughout plains; SE Asia

Local/Common/Vernacular names: *Iron Wood Tree, Siamese Senn, Siamese tree senna* (English)

Senna sophera (L.) Roxb., Fl. Ind. 2: 347. 1832

Cassia sophera L., Sp. Pl. 379. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Cassia occidentalis var. *sophera* (L.) Kuntze, Revis. Gen. Pl. 1: 170.1891

Ditremexa sophera (L.) Britton & Wilson, Sci. Surv. Porto Rico & Virgin Islands 5: 372. 1924

Senna occidentalis var. *sophera* (L.) X.Y.Zhu., Legumes of China 29 29 2007

Distribution: Karnataka (Ballari, Belagavi, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Kalburgi, Kodagu, Mysuru, Raichur, Shivamogga, Uttara Kannada, Vijayapura), Tamil Nadu; native to tropical America;introduced elsewhere.

Local/Common/Vernacular name: *Foetid senna* (English); *Kasamarda* (Kannada)

Senna spectabilis (DC.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 600. 1982

Cassia spectabilis DC., Cat. Pl. Horti Monsp. 90. 1813; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Pseudocassia spectabilis (DC.) Britton & Rose, N. Amer. Fl. 23(4): 230. 1930

Distribution: Karnataka (Bengaluru, Chamarajanagar, Chikkaballapura, Kalburgi, Kolar, Mysuru, Tumakuru), cultivated throughout tropical parts; native to tropical S America; cultivated in tropics.

Local/Common/Vernacular name: *Golden Shower* (English)

Senna splendida (Vogel) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 190. 1982

Cassia splendida Vogel, Gen. Cass. Syn. 17. 1837; Seeth. et al., Fl. 71. 2000.

Distribution: Karnataka (Kalburgi); native to tropical S America; cultivated elsewhere.

Senna sulfurea (Collad.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 78. 1982

Wellia tagera Rheed, Hort. Ind. Malabar. 6: 17,t.9-10 17.1686

Cassia arborescens Vahl, Symb. Bot. 3: 56. 1794

Cassia sulfurea Collad., Hist. Nat. Med. Casses 84. 1816

Senna arborescens (Vahl) Roxb., Fl. Ind. 3: 345. 1832

Cassia enneaphylla Wight & Arn., Prodr. Fl. Ind. Orient. 1: 289. 1834

Distribution: Karnataka (Ballari, Chikkamagaluru, Chitradurga, Dharwad, Hassan, Kolar), throughout India; Bangladesh, China, Indonesia, Thailand; introduced elsewhere.

Local/Common/Vernacular name: *Sulphur-flowered senna* (English); *Adavi thangadi* (Kannada)

Senna surattensis (Burm.f.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35. 1982

Cassia surattensis Burm. f., Fl. Indica 97. 1768; C.J.Saldanha, Fl. Karnataka 1: 386. 1984.

Cassia glauca Lam., Encycl. 1: 647. 1785; B.D.Sharma et al., Fl. Karnataka Anal. 89. 1984.

Cassia suffruticosa Koenig ex Roth., Nov. Pl. Sp. 213. 1821

Senna speciosa Roxb., Fl. Ind. 2: 347.1832

Cassia glauca var. *suffruticosa* (Koenig ex Roth.) Baker, Hook.f., Fl. Brit. India 2: 265. 1878

Psilorhegma suffruticosa (Roth) Britton, N. Amer. Fl. 23(4): 255. 1930

Psilorhegma glauca (Lam.) O. Deg. & I. Deg., Fl. Hawaïensis Fam. 169b. 1959

Cassia surattensis var. *suffruticosa* (Roth) Ali, Fl. W. Pakistan 54.1973

Cassia surattensis subsp. *glauca* (Lam.) K.Larsen & S.S.Larsen, Fl. Cambodge, Laos & Vietnam 18: 102. 1980

Cassia surattensis subsp. *suffruticosa* (Roth) K.Larsen & S.S.Larsen, Fl. Cambodge, Laos & Vietnam 18: 102. 1980

Senna surattensis var. *glauca* (Lam.) X.Y. Zhu, Legumes of China 31. 2007

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kolar, Mysuru, Uttara Kannada, Vijayapura), throughout India, also cultivated; Australia, Bangladesh, Indonesia, Malaysia, Myanmar, Papua New Guinea; cultivated throughout tropics.

Senna timoriensis (DC.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35(1): 98. 1982

Cassia timoriensis DC., Prodr. 2: 499. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 90. 1984.

Senna glauca Roxb., Fl. Ind. 2: 351.1832

Cassia goensis Dalzell in Hooker's J. Bot. Kew Gard. Misc. 4: 112. 1852

Distribution: Karnataka (Bengaluru, Dakshina Kannada, Shivamogga, Uttara Kannada), Andhra Pradesh, Arunachal Pradesh, Assam, Daman, Diu, Goa, Kerala, Maharashtra, Manipur, Meghalaya, Mizoram, Nagaland, Tamil Nadu, Tripura, West Bengal; Australia, Indonesia, Malaysia.

Local/Common/Vernacular names: *Limestone Cassia*, *Timor senna* (English)

Senna tora (L.) Roxb., Fl. Ind. 2: 340. 1832

Cassia tora L., Sp. Pl. 376. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 90. 1984.

Emelista tora (L.) Britton & Rose ex Britton & P. Wilson, Sci. Surv. Porto Rico & Virgin Islands 5: 371.1924

Distribution: Karnataka (throughout), Tamil Nadu; pantropical.

Local/Common/Vernacular name: *Foetid cassia* (English); *Chagche* (Kannada)

Senna uniflora (Mill.) H.S.Irwin & Barneby, Mem. New York Bot. Gard. 35: 258. 1982

Cassia uniflora Mill., Gard. Dict., ed. 8. n. 5. 1768; N.P.Singh, Fl. East. Karnataka 1: 278. 1988.

Cassia sericea Sw., Fl. Ind. Occ. 2: 724. 1798

Serceocassia uniflora (Mill.) Britton, N. Amer. Fl. 23(4): 246.1930

Distribution: Karnataka (cultivated throughout), cultivated throughout India; native to tropical S America.

Vachellia Wight & Arn.

Vachellia eburnea (L.f.) P.J.H.Hurter & Mabb., Mabberley's Pl.-Book 1021. 2008

Mimosa eburnea L.f., Suppl. Pl. 437. 1781

Acacia eburnea (L.f.) Willd., Sp. Pl. 4: 1081. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Acacia roxburghii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 276. 1834

Acacia campbellii Arn., Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 18: 333. 1836

Acacia minutifolia Ragup., A.Mahad. & Thoth., Indian J. Forest. 14: 65 65 1991non F.muell.1874

Distribution: Karnataka (Ballari, Belagavi, Chitradurga, Kalburgi, Mysuru, Raichur, Tumakuru, Vijayapura, Yadgir), Andhra Pradesh, Arunachal Pradesh, Bihar, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh; Afghanistan, Arabia, Pakistan, Sri Lanka.

Local/Common/Vernacular name: *Black babool* (English); *Pikajali* (Kannada)

Vachellia farnesiana (L.) Wight & Arn., Prodr. Fl. Ind. Orient. 1: 272. 1834

Mimosa farnesiana L., Sp. Pl. 521. 1753

Acacia farnesiana (L.) Willd., Sp. Pl. 4: 1083. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 92. 1984.

Distribution: Karnataka (Belagavi, Bengaluru, Chitradurga, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Raichur), throughout India; probably native to Mexico & C America; native to C & S America, elsewhere naturalised in tropics.

Local/Common/Vernacular names: *Cassie flower*, *Fragrant acacia*, *Needle bush*, *Sweet acaia*, *Sweet wattle* (English); *Kastoorijali* (Kannada)

Vachellia horrida (L.) Kyal. & Boatwr., Bot. J. Linn. Soc. 172(4): 513. 2013

Mimosa horrida L., Sp. Pl. 1: 521. 1753

Mimosa cornigera L.f., Suppl. Pl. 438. 1781

Mimosa latronum L.f., Suppl. Pl. 438. 1781

Acacia horrida (L.) Willd., Sp. Pl. 4(2): 1082. 1806; Seetharam, et al., Fl. 73. 2000

Acacia latronum (L.f.) Willd., Sp. Pl. 4: 1077. 1806

Distribution: Karnataka (Bagalkot, Ballari, Belagavi, Chitradurga, Dharwad, Kalburgi, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Delhi, Goa, Kerala, Maharashtra, Odisha, Rajasthan, Tamil Nadu; Myanmar.

Local/Common/Vernacular names: *Anegobli*, *Donni*, *Vattajali* (Kannada)

Vachellia leucophloea (Roxb.) Maslin, Seigler & Ebinger, Blumea 58:42.2013

Mimosa leucophloea Roxb., Pl. Coromandel 2:27, t. 150. 1800

Acacia alba (Rottler) Willd., Enum. Pl. Berol. Suppl. 68. 1803

Mimosa alba Rottler, Nov. Act. Ber. 208. 1803

Acacia leucophloea (Roxb.) Willd., Sp. Pl. 4: 1083. 1806; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Bidar, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kolar, Mysuru, Raichur, Tumakuru, Vijayapura), Andhra Pradesh, Delhi, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal; Indonesia Mauritius, Malaysia, Myanmar, Pakistan, Sri Lanka, Thailand, Vietnam.

Local/Common/Vernacular names: *White babool* (English); *Biljali*, *Naibela* (Kannada)

Vachellia nilotica (L.) P.J.H.Hurter & Mabb., Mabberley's Pl.-Book 1021. 2008.

subsp. **cupressiformis** (J.L.Stewart) Ragupathy, Seigler, Ebinger & Maslin, Phytotaxa 162(3): 177.2014

Acacia arabica var. *cupressiformis* J.L.Stewart, Punjab. Pl. 51. 1869

Acacia nilotica subsp. *cupressiformis* (J.L.Stewart) Ali & Faruqui, Pak. J. Bot. 1: 4. 1969

Acacia nilotica subsp. *indica* var. *cupressiformis* (J.L.Stewart) Vajrav. & Kamble, J. Bombay Nat. Hist Soc. 79: 708.1983

Distribution: Karnataka (Bagalkot, Chamarajanagar, Dharwad, Mysuru, Raichur, Vijayapura, Yadgiri), Andhra Pradesh, Gujarat, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Pakistan.

subsp. **indica** (Benth.) Kyal. & Boatwr., Bot. J. Linn. Soc. 172(4): 515. 2013

Mimosa arabica Lam., Encycl. 1(1): 19. 1783

Acacia arabica var. *indica* Benth., London J. Bot. 1: 500. 1842

Acacia arabica auct. non (Lam.) Willd. 1805; Baker in Hook.f., Fl. Brit. India 2: 293. 1878

Acacia nilotica var. *indica* (Benth.) A.F.Hill, Bot. Mus. Leafl. 8: 99. 1940

Acacia nilotica subsp. *Indica* (Benth.) Brenan, Kew Bull. 12: 84. 1957; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Distribution: Karnataka (Bagalkot, Ballari, Belagavi, Bengaluru, Bidar, Chitradurga, Davanagere, Hassan, Kalburgi, Kolar, Koppala, Mysuru, Raichur, Shivamogga, Tumakuru, Vijayapura, Yadgiri), throughout India; paleotropical

Local/Common/Vernacular names: *Babool Gum Arabic Tree*, *Indian gum arabic* (English); *Gobli, Kari jali* (Kannada)

Vachellia planifrons (Koenig ex Wight & Arnott) Ragupathy, Seigler, Ebinger & Maslin, Phytotaxa 162(3): 177. 2014

Acacia planifrons Wight & Arn., Prodr. Fl. Ind. Orient. 1: 276. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 93. 1984.

Distribution: Karnataka (Ballari), Andhra Pradesh, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu; Sri Lanka.

Local/Common/Vernacular names: *Umbrella thorn*, *Umbrella thorn babool* (English); *Budde jali* (Kannada)

Vachellia tomentosa (Rottler) Maslin, Seigler & Ebinger, Blumea 58:43. 2013

Mimosa tomentosa Rottler, Neue Schriften Ges. Naturf. Freunde Berlin 4: 208. 1803

Acacia tomentosa Willd., Sp. Pl. 4: 1087. 1806; C.J.Saldanha, Fl. Karnataka 1: 402. 1984.

Distribution: Karnataka (Dharwad), Andhra Pradesh, Gujarat, Goa, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh, West Bengal), Bangladesh, Indonesia, Myanmar, Sri Lanka, Thailand, Vietnam.

Xylia Benth.

Xylia xylocarpa (Roxb.) Taub., Bot. Centralbl. 47: 395. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 96. 1984.

Mimosa xylocarpa Roxb., Pl. Coromandel 1: 68. t. 100. 1795

Acacia xylocarpa (Roxb.) Willd., Sp. Pl., ed. 4, 4(2): 1055. 1806

Inga xylocarpa (Roxb.) DC., Prodr. 2: 439. 1825

Xylia dolabriformis Benth. in Hooker's J. Bot. 4: 417. 1842

Distribution: Karnataka (Belagavi, Chikkamagaluru, Dakshina Kannada, Davanagere, Dharwad, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Bihar, Goa, Gujarat, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Odisha, Tamil Nadu, West Bengal; Bangladesh, Cambodia, Laos, Myanmar, Philippines, Singapore, Thailand, Vietnam.

Local/Common/Vernacular name: *Burma iron wood* (English); *Jambe* (Kannada)

Leguminosae-Cercidoideae

Bauhinia L.

Bauhinia acuminata L., Sp. Pl. 1: 375. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Bauhinia linnaei Ali, Portug. Acta Biol., Ser. B 8: 243. 1966

Pauletia acuminata (L.) A.Schimtz, Bull. Jard. Bot. Natl. Belg. 43:387. 1973

Distribution: Karnataka (Kalburgi, Kodagu, Uttara Kannada), Peninsular to NE India, Andaman & Nicobar Islands; SE Asia.

Local/Common/Vernacular name: *Dwarf white Bauhinia* (English)

Bauhinia × blakeana Dunn, J. Bot. 46: 325. 1908

Distribution: Karnataka (cultivated in Bengaluru), cultivated throughout in gardens.

Local/Common/Vernacular name: *Hong Kong Orchid Tree* (English)

Note: A hybrid between *Bauhinia variegata* and *Bauhinia purpurea*.

Bauhinia galpinii N.E.Br., Gard., Chron. 9. 728. 1891; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Distribution: Karnataka (cultivated in Bengaluru, Davanagere, Kodagu, Mysuru), cultivated as an ornamental in India; native to S Africa.

Local/Common/Vernacular name: *Red orchid bush* (English)

Bauhinia monandra Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 42(2): 73. 1873; Kesh.Murthy & Yogan, Fl. Kodagu 167. 1990.

Caspareaopsis monandra (Kurz) Britton & Rose, N. Amer. Fl. 23(4): 217. 1930

Distribution: Karnataka (cultivated in Kodagu), cultivated in Maharashtra, Tamil Nadu; native to Brazil.

Local/Common/Vernacular name: *Pink orchid tree* (English)

Bauhinia phoenicea Wight & Arn., Prodr. Fl. Ind. Orient. 1: 296. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Phanera phoenicea (Wight & Arn.) Benth. in Miq., Pl. Jungh. 262. 1852

Bauhinia benthamii Bedd., Fl. Sylv. S. India 93. 1872

Perlibia phoenicea (Wight & Arn.) A.Schimtz, Bull. Soc. Roy. Bot. Belgique 110:15. 1977

Distribution: Karnataka (Dakshina Kannada, Davanagere, Hassan, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Kerala, Maharashtra, Tamil Nadu; endemic.

Local/Common/Vernacular name: *Crimson mountain Ebony* (English); *Kempu yatha Balli* (Kannada)

Bauhinia purpurea L., Sp. Pl. 1: 375. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Bauhinia coromandeliana DC., Prod. 2:515. 1825

Bauhinia triandra Roxb., Fl. Ind. 2:320. 1832

Telisstra purpurea (L.) Raf., Sylva Tellur. 112. 1838

Phanera purpurea (L.) Benth. in Miq., Pl. Jungh. 262. 1852

Caspareaopsis purpurea (L.) Pittier in Pittier et al., Cat. Fl. Venez. 1 (3rd Conf. Interamer. Agric., No. 20): 363. 1945

Perlibia purpurea (L.) A.Schimtz, Bull. Jard. Bot. Natl. Belg. 43:382. 1973

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Hassan, Kalburgi, Kolar, Mysuru, Shivamogga, Tumakuru, Uttara Kannada, Vijayapura), throughout India; China, S & SE Asia; cultivated throughout tropics.

Local/Common/Vernacular names: *Geranium tree*, *Orchid tree*, *Purple Bauhinia* (English); *Basavanapada*, *Kempu mandara*, *Kempukajivala*, *Sarul*, *Ulipe* (Kannada)

Bauhinia racemosa Lam., Encycl. 1: 390. 1785; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Bauhinia parviflora Vahl, Symb. Bot. 3: 55. 1794

Piliostigma racemosum (Lam.) Benth. in Miq., Pl. Jungh. 262. 1852

Pauletia racemosum (L.) A.Schimtz, Bull. Soc. Roy. Bot. Belgique 110:15. 1977

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Dharwad, Hassan, Kalburgi, Kodagu, Mysuru, Raichur, Shivamogga, Tumakuru, Vijayapura), Andhra Pradesh, Bihar, Delhi, Goa, Gujarat, Himachal Pradesh, Kerala, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal; China, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Common Mountain Ebony* (English); *Aatalumara*, *Akalu pada*, *Doddabanni*, *Vana sampige* (Kannada)

Bauhinia tomentosa L., Sp. Pl. 1: 375. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Pauletia tomentosa (L.) A.Schimtz, Bull. Jard. Bot. Natl. Belg. 43:393. 1973

Distribution: Karnataka (Chitradurga, Kalburgi, Kolar, Mysuru, Shivamogga, Uttara Kannada), Andaman & Nicobar Islands, Andhra Pradesh, Assam, Bihar, Delhi, Goa, Gujarat, Maharashtra, Tamil Nadu, Uttar Pradesh; tropical Africa, Asia.

Local/Common/Vernacular names: *Bell Bauhinia*, *St. Thomas tree*, *Yellow Bauhinia* (English); *Kadatti*, *Karnanaspu* (Kannada); *Pitakanchana* (Sanskrit)

Bauhinia variegata L., Sp. Pl. 375. 1753; B.D.Sharma et al., Fl. Karnataka Anal. 87. 1984.

Bauhinia candida auct. non Aiton, 1789; DC., Prodr. 2: 513.1825.

Phanera variegata (L.) Benth. in Miq., Pl. Jungh. 261. 1852

Perlabia variegata (L.) A.Schimtz, Bull. Jard. Bot. Natl. Belg. 43:386.1973

Distribution: Karnataka (Belagavi, Bengaluru, Bidar, Chikkamagaluru, Chitradurga, Davanagere, Kalburgi, Kodagu, Kolar, Mysuru, Tumakuru), throughout India; Indo-China, S Asia, native to China

Local/Common/Vernacular names: *Buddhist Bauhinia*, *Variegated Bauhinia* (English); *Arisantige*, *Ayata*, *Bilikanjivala*, *Karalabhogi*, *Kempumandara*, *Ulipe* (Kannada)

Phanera Lour.

Phanera retusa (Roxb.) Benth. in Miq., Pl. Jungh. 263.1852

Bauhinia retusa Roxb., Fl. Ind. 2: 322. 1832 non Poir. 1811 nec Roxb. ex DC. 1825; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Bauhinia roxburghiana Voigt, Hort. Suburb. Calcutt. 254. 1845

Lasiobema retusum (Roxb.) de Wit, Reinwardtia 3: 538. 1856

Bauhinia semla Wunderlin, Taxon 25(2-3): 362.1976

Distribution: Karnataka (Mysuru), Andhra Pradesh, Bihar, Himachal Pradesh, Madhya Pradesh, Maharashtra, Odisha, Punjab, Uttar Pradesh

Phanera vahlii (Wight & Arn.) Benth. in Miq., Pl. Jungh. 263.1852

Bauhinia vahlii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 297. 1834; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Distribution: Karnataka (Bengaluru, Chikkamagaluru), throughout India; Nepal, Pakistan.

Local/Common/Vernacular names: *Aane pada*, *Hepparige*, *Kambihu* (Kannada)

Piliostigma Hochst.

Piliostigma foveolatum (Dalzell) Thoth., Bull. Bot. Soc. Bengal 19: 131. 1967

Bauhinia foveolata Dalzell, J. Linn. Soc. Bot. 13: 188. 1873; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Bauhinia lawii Baker, Hook.f., Fl. Brit. India 2: 277. 1878

Distribution: Karnataka (Belagavi, Uttara Kannada), Dadra, Gujarat, Maharashtra, Nagar Haveli; endemic.

Local/Common/Vernacular name: *Basavanapada* (Kannada)

Piliostigma malabaricum (Roxb.) Benth. in Miq., Pl. Jungh. 261. 1852

Bauhinia malabarica Roxb., Fl. Ind. 2: 321. 1832; B.D.Sharma et al., Fl. Karnataka Anal. 91. 1984.

Pauletia acida (Korth.) Hassk., Flora 31: 578. 1848

Piliostigma acidum (Reinw. ex Korth.) Benth. in Miq., Pl. Jungh. 261. 1852

Bauhinia malabarica var. *reniformis* Royle ex Baker in Hook.f., Fl. Brit. India 2:277.1878

Bauhinia acida Reinw. ex Korth., Ver. Nat. Gesch, Ned. Bezitt. Bot. 86.1941

Piliostigma malabaricum var. *acidum* (Reinw. ex Korth.) de Wit, Reinwardtia 3:351.1956

Piliostigma malabaricum var. *reniformis* Royle ex Baker Thoth., Bull. Bot. Soc. Bengal 19: 131. 1967

Distribution: Karnataka (Ballari, Belagavi, Bengaluru, Chikkamagaluru, Dakshina Kannada, Kalburgi, Kodagu, Mysuru, Shivamogga, Uttara Kannada), Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Himachal Pradesh, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Tamil Nadu, Tripura, Uttar Pradesh, Uttarakhand, West Bengal; Australia, Indo-China, S & SE Asia.

Local/Common/Vernacular names: *Malabar Mountain Ebony* (English); *Basavana pada*, *Cheppura*, *Kogele*, *Mandara*, *Nimbede*, *Pulikoelepukke*, *Sappara*, *Shadhu* (Kannada)