

REFERENCES

- Acín**, P., G. Rosell, A. Guerrero and C. Quero, 2010. Sex pheromone of the Spanish population of the beet armyworm *Spodoptera exigua*. *J. Chem. Ecol.*, **36**: 778-786.
- Adachi**, Y., N. D. Do, M. Kinjo, S. Makisako, R. Yamakawa, K. Mori and T. Ando, 2010. Positions and stereochemistry of methyl branches in the novel sex pheromone components produced by a lichen moth, *Lyclene dharma dharma*. *J. Chem. Ecol.*, **36**: 814-823.
- Adati**, T. and S. Tatsuki 1999. Identification of female sex pheromone of the legume pod borer, *Maruca vitrata* and antagonistic effects of geometrical isomers. *J. Chem. Ecol.*, **25**: 105-115.
- Adati, T., S. Kato, W. Toriumi, K. Kawazu, W. Susila, K. Sumiartha, G. Susatika, Dadang and I. M. Samudra, 2013. Field trials of synthetic sex pheromone lures for the large cabbage-heart caterpillar moth, *Crocidolomia pavonana* (Lepidoptera: Crambidae) in Indonesia. *Appl. Entomol. Zool.*, **48**: 113-118.
- Adler**, V. E., M. Jacobson, J. F. Edmiston and M. H. Fleming, 1976. (*Z,E*)-9,12-Tetradecadien-1-ol acetate: a potent attractant for male *Ephestiodes infimella*. *J. Econ. Entomol.*, **69**: 706-708.
- Adler, V. E., 1983. Adult Sesiidae of Maryland collected in traps baited with isomers of 3,13-octadecadien-1-ol acetate fruit orchards. *J. Environ. Sci. Health.*, **A 18**: 611-619.
- Albages**, R., R. Canela, M. Eizaguirre, M. Riba and A. Guerrero, 1988. (*Z*)-9-Tetradecen-1-ol, an attractant for *Athetis Hospes* Freyer. *Colloq. INRA (Mediateurs Chim.: Comportement Syst. Lepidopteres, Appl. Agron.)* **46**: 55-58. [*Chem. Abst.*, 110: 52909n]
- Albert**, von R., H. Bogenschütz and E. König, 1984. Investigation on the use of pheromone-baited traps for monitoring the population dynamics of *Operophtera brumata* L. (Lepid., Geometridae). *Z. Angew. Entomol.*, **98**: 286-298.
- Alford**, D. V., 1978. Observations on the specificity of pheromone-baited traps for *Cydia funebrana* (Treitschke) (Lepidoptera: Tortricidae). *Bull. Entomol. Res.*, **68**: 97-103.
- Alford, A. R., P. J. Silk, M. McClure, C. Gibson and J. Fitzpatrick, 1983. Behavioural effects of secondary components of the sex pheromone of the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **115**: 1053-1058.
- Alford, A. R. and P. J. Silk, 1984. Behavioral effects of secondary components of sex pheromone of western spruce budworm (*Choristoneura occidentalis*) Free. *J. Chem. Ecol.*, **10**: 265-270.
- Allison**, J. D. and R. T. Cardé, 2006. Heritable variation in the sex pheromone of the almond moth, *Cadra cautella*. *J. Chem. Ecol.*, **32**: 621-641.

Alm, S. R., R. N. Williams, D. M. Pavuk, J. W. Snow and M. A. Heinlein, 1989. Distribution and seasonal flight activity of male grape root borers (Lepidoptera: Sesiidae) in Ohio. *J. Econ. Entomol.*, **82**: 1604-1608.

Altamar-Varón, P., D. Pérez-Maldonado, D. Rodríguez-Caicedo, C. Guerrero-Perilla and E. Coy-Barrera, 2016. Chemical composition of the low-polar fraction of the *Copitarsia uncilata* Burgos & Leiva (Lepidoptera: Noctuidae) eversible pheromone gland. *Neotropical Entomol.*, **45**: 734-739.

Altesor, P., C. Rossini, P. H. G. Zarbin and A. Gonzalez, 2009. Sex pheromone of the bud borer *Epinotia aporema*: chemical identification and male behavioral response. *J. Chem. Ecol.*, **35**: 349-354.

Anagnostakis, S. L., K. M. Welch, J. W. Snow, K. Scarborough and T. D. Eichlin, 1994. The rediscovery of the clearwing chestnut moth, *Synanthedon castaneae* (Busck) (Lepidoptera: Sesiidae) in Connecticut. *J. N.Y. Entomol. Soc.*, **102**: 111-112.

Ando, T., S. Yoshida, S. Tatsuki and N. Takahashi, 1975. Lepidopterous sex attractants. *Agric. Biol. Chem.*, **39**: 1163-1165.

Ando, T., S. Yoshida, S. Tatsuki and N. Takahashi, 1977. Sex attractants for male Lepidoptera. *Agric. Biol. Chem.*, **41**: 1485-1492.

Ando, T., H. Kuroko, S. Nakagaki, O. Saito, T. Oku and N. Takahashi, 1978. Two-component sex attractants for male moths of the subfamily Tortricinae (Lepidoptera). *Agric. Biol. Chem.*, **42**: 1081-1083.

Ando, T., T. Koshihara, H. Yamada, M. H. Vu, N. Takahashi and Y. Tamaki, 1979. Electroantennogram activities of sex pheromone analogues and their synergistic effect on field attraction in the diamondback moth. *Appl. Entomol. Zool.*, **14**: 362-364.

Ando, T., K. Kishino, S. Tatsuki and N. Takahashi, 1980 (a). Sex pheromone of the rice green caterpillar: chemical identification of three components and field tests with synthetic pheromones. *Agric. Biol. Chem.*, **44**: 765-775.

Ando, T., O. Saito, K. Arai and N. Takahashi, 1980 (b). (*Z*)- and (*E*)-12-Tetradecenyl acetates: sex pheromone components of oriental corn borer (Lepidoptera: Pyralidae). *Agric. Biol. Chem.*, **44**: 2643-2649.

Ando, T., H. Kuroko, S. Nakagaki, O. Saito, T. Oku and N. Takahashi, 1981. Multi-component sex attractants in systematic field tests of male Lepidoptera. *Agric. Biol. Chem.*, **45**: 487-495.

Ando, T., M. H. Vu, S. Yoshida, N. Takahashi, S. Tatsuki, K. Katagiri, A. Yamane, T. Ikeda and S. Yamazaki, 1982. (*5Z,7E*)-5,7-Dodecadien-1-ol: female sex pheromone of the pine moth *Dendrolimus spectabilis* Butler. *Agric. Biol. Chem.*, **46**:

709-715.

- Ando, T., K. Taguchi, M. Uchiyama and T. Horikawa, 1985 (a). Female sex pheromone of the tea leafroller, *Caloptilia theivora* Walsingham (Lepidoptera: Gracillariidae). *Agric. Biol. Chem.*, **49**: 233-234.
- Ando, T., K. Taguchi, M. Uchiyama, T. Ujiye and H. Kuroko, 1985 (b). (7Z,11Z)-7,11-Hexadecadienol: sex attractant of the citrus leafminer moth, *Phyllocnistis citrella* Stainton (Lepidoptera, Phyllocnistidae). *Agric. Biol. Chem.*, **49**: 3633-3635.
- Ando, T., M. Koike, M. Uchiyama and H. Kuroko, 1987. Lepidopterous sex attractants with a conjugated diene system. *Agric. Biol. Chem.*, **51**: 2691-2694.
- Ando, T., H. Ohsawa, T. Ueno, H. Kishi, Y. Okamura and S. Hashimoto, 1993. Hydrocarbons with a homoconjugated polyene system and their monoepoxy derivatives: sex attractants of geometrid and noctuid moths distributed in Japan. *J. Chem. Ecol.*, **19**: 787-798.
- Ando, T., H. Kishi, N. Akashio, X.-R. Qin, N. Saito, H. Abe and S. Hashimoto, 1995 (a). Sex attractants of geometrid and noctuid moths: chemical characterization and field test of monoepoxides of 6,9-dienes and related compounds. *J. Chem. Ecol.*, **21**: 299-311.
- Ando, T., R. Ohno, T. Kokuryu, A. Funayoshi and M. Nomura, 1995 (b). Identification of female sex pheromone components of rice looper, *Plusia festucae* (L.), (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **21**: 1181-1190.
- Ando, T., K. Ohtani, M. Yamamoto, T. Miyamoto, X.-R. Qin and Witjaksono, 1997. Sex pheromone of Japanese giant looper, *Ascotis selenaria cretacea*: identification and field tests. *J. Chem. Ecol.*, **23**: 2413-2423.
- Ando, T., S. Inomata, R. Shimada, M. Nomura, S. Uehara and G.-Q. Pu, 1998. Sex pheromones of *Thysanoplusia intermixta* and *T. orichalcea*: identification and field tests. *J. Chem. Ecol.*, **24**: 1105-1116.
- Anglade**, P., J. Stockel and I.W.G.O. cooperators, 1984. Intraspecific sex-pheromone variability in the European corn borer, *Ostrinia nubilalis* Hbn. (Lepidoptera, Pyralidae). *Agronomie*, **4**: 183-187.
- Anshelevich**, L., M. Kehat, E. Dunkelblum and S. Greenberg, 1993. Sex pheromone traps for monitoring the honeydew moth, *Cryptoblabes gnidiella*: effect of pheromone components, pheromone dose, field aging of dispenser, and type of trap on male captures. *Phytoparasitica*, **21**: 189-198.
- Arai**, K., T. Ando, A. Sakurai, H. Yamada, T. Koshihara and N. Takahashi, 1982. Identification of the female sex pheromone of the cabbage webworm. *Agric. Biol. Chem.*, **46**: 2395-2397.
- Arakawa**, A., F. Mochizuki, Y. Yumoto and K. Fukumoto, 2009. Sex pheromone components of *Toleria romanovi* and insect attractant

containing octadecadienol and octadecadienyl acetate. *Jpn. Kokai Tokkyo Koho* JP 2009209104. [SciFinder. AN 2009:1131520]

- Arn, H., C. Schwarz, H. Limacher and E. Mani, 1974. Sex attractant inhibitors of the codling moth *Laspeyresia pomonella* L. *Experientia*, 30: 1142-1144.
- Arn, H., E. Städler and S. Rauscher, 1975. The electroantennographic detector - a selective and sensitive tool in the gas chromatographic analysis of insect pheromones. *Z. Naturforsch.*, 30c: 722-725.
- Arn, H., B. Delley, M. Baggiolini and P. J. Charmillot, 1976(a). Communication disruption with sex attractant for control of the plum fruit moth, *Grapholita funebrana*: a two-year field study. *Entomol. Exp. Appl.*, 19: 139-147.
- Arn, H., S. Rauscher, H.-R. Buser and W. L. Roelofs, 1976(b). Sex pheromone of *Eupoecilia ambiguella*: *cis*-9-dodecenyl acetate as a major component. *Z. Naturforsch.*, 31c: 499-503.
- Arn, H., E. Priesner, H. Bogenschütz, H. R. Buser, D. L. Struble, S. Rauscher and S. Voerman, 1979. Sex pheromone of *Tortrix viridana*: (Z)-11-tetradecenyl acetate as the main component. *Z. Naturforsch.*, 34c: 1281-1284.
- Arn, H., E. Städler, S. Rauscher, H. R. Buser, H. Mustaparta, P. Esbjerg, H. Philipsen, O. Zethner, D. L. Struble and R. Bues, 1980. Multicomponent sex pheromone in *Agrotis segetum*: preliminary analysis and field evaluation. *Z. Naturforsch.*, 35c: 986-989.
- Arn, H., P. Esbjerg, R. Bues, M. Tóth, G. Szöcs, P. Guerin and S. Rauscher, 1983. Field attraction of *Agrotis segetum* males in four European countries to mixtures containing three homologous acetates. *J. Chem. Ecol.*, 9: 267-276.
- Arn, H., P. M. Guerin, H. R. Buser, S. Rauscher and E. Mani, 1985. Sex pheromone blend of the codling moth, *Cydia pomonella*: evidence for a behavioral role of dodecan-1-ol. *Experientia*, 41: 1482-1484.
- Arn, H., M. Tóth and E. Priesner, 1986(a). "List of sex pheromones of Lepidoptera and related attractants" OILB-SROP / IOBC-WPRS, Paris, pp. 123.
- Arn, H., S. Rauscher, H.-R. Buser and P. M. Guerin, 1986(b). Sex pheromone of *Eupoecilia ambiguella* female: analysis and male response to ternary blend. *J. Chem. Ecol.*, 12: 1417-1429.
- Arn, H., S. Rauscher, P. Guerin and H.-R. Buser, 1988. Sex pheromone blends of three tortricid pests in European vineyards. *Agric. Ecosys. Environ.*, 21: 111-117.
- Arn, H., M. Tóth and E. Priesner, 1992. "List of sex pheromones of Lepidoptera and related attractants" OILB-SROP / IOBC-WPRS, Secrétariat Général, F-84143 Montfavet, France, ISBN 92-9067-044-4. 179 pp.

Arn, H., M. Tóth and E. Priesner, 1997. "List of sex pheromones of Lepidoptera and related attractants, supplement 1992-1996" *Techno. Trans. Mating Disrup.*, 20 : 257-293.

Asaro, C., M. J. Dalusky and C. W. Berisford, 2001. Quantity and ratio of pheromone components among multiple generations of the nantucket pine tip moth (Lepidoptera: Tortricidae) in Georgia and Virginia. *Environ. Entomol.*, 30: 1006-1011.

Arsura, E., A. Capizzi, P. Piccardi and P. Spinelli, 1977. (Z)-9-Tetradecen-1-ol and (Z)-9-tetradecenyl acetate: a potent attractant system for male *Sesamia cretica* Led. (Lep., Noctuidae). *Experientia*, 33: 1423-1424.

Attygalle, A. B., J. Schwarz, O. Vostrowsky and H. J. Bestmann, 1986. Individual variation in the sex pheromone components of the false codling moth, *Cryptophlebia leucotreta* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, 41c: 1077-1081.

Attygalle, A. B., M. Herrig, O. Vostrowsky and H. J. Bestmann, 1987. Technique for injecting intact glands for analysis of sex pheromones of Lepidoptera by capillary gas chromatography: reinvestigation of pheromone complex of *Mamestra brassicae*. *J. Chem. Ecol.*, 13: 1299-1311.

Attygalle, A. B., J. Schwarz and N. E. Gunawardena, 1988 (a). Sex pheromone of brinjal shoot and pod borer *Leucinodis orbonalis* Guenée (Lepidoptera: Pyralidae: Pyraustinae). *Z. Naturforsch.*, 43c: 790-792.

Attygalle, A. B., C.-H. Wu, J. Schwarz, O. Vostrowsky, I. Hasenfuss and H. J. Bestmann, 1988 (b). Sex pheromone of female *Myelois cribrella* Hübner (Lepidoptera: Pyralidae). Chemical identification, electrophysiological evaluation, and field attractancy tests. *J. Chem. Ecol.*, 14: 485-494.

Attygalle, A. B., G. N. Jham, A. Svatos, R. T. S. Frighetto and J. Meinwald, 1995. Microscale, random reduction to the characterization of $(3E,8Z,11Z)$ -3,8,11-tetradecatrienyl acetate, a new lepidopteran sex pheromone. *Tetrahedron Lett.*, 36: 5471-5474.

Attygalle, A. B., G. N. Jham, A. Svatos, R. T. S. Frighetto, F. A. Ferrara, E. F. Vilela, M. A. Uchoa-Fernandes and J. Meinwald, 1996. $(3E,8Z,11Z)$ -3,8,11-Tetradecatrienyl acetate, major sex pheromone component of the tomato pest *Scrobipalpuloides absoluta* (Lepidoptera: Gelechiidae). *Bioorg. Medic. Chem.*, 4: 305-314.

Avand-Faghah, A. and B. Frérot, 2008. Identification of the sex pheromone of *Sesamia cretica* Lederer. *J. Chem. Ecol.*, 34: 103-106.

Ayre, G. L., W. J. Turnock and D. L. Struble, 1982 (a). Spatial and temporal variability in sex attractant trap catches of *Scotogramma trifolii* (Lepidoptera: Noctuidae) in relation to its biology in

Manitoba. *Can. Entomol.*, **114**: 145-154.

Ayre, G. L., W. J. Turnock and D. L. Struble, 1982(b). Spatial and temporal variability in sex attractant trap catches of *Euxoa messoria* and *E. ochrogaster* (Lepidoptera: Noctuidae) in relation to their biology in Manitoba. *Can. Entomol.*, **114**: 993-1001.

Babu, V. R., S. Satpathy, B. V. S. Reddy, 2021. Identification and characterisation of female released sex pheromone components of jute semilooper, *Anomis sabulifera* Guenée (Lepidoptera: Noctuidae). *J. Environ. Biol.*, **42**: 254-264.

Bacon, O. G., J. N. Seiber and G. G. Kennedy, 1976. Evaluation of survey trapping techniques for potato tuberworm moths with chemical baited traps. *J. Econ. Entomol.*, **69**: 569-572.

Bailey, J. B., L. M. McDonough and M. P. Hoffmann, 1986. Western avocado leafroller, *Amorbia cuneana* (Walsingham), (Lepidoptera: Tortricidae). Discovery of populations utilizing different ratios of sex pheromone components. *J. Chem. Ecol.*, **12**: 1239-1245.

Baker, J. L., A. S. Hill, R. T. Cardé, A. Kurokawa and W. L. Roelofs, 1975. Sex pheromone field trapping of the omnivorous leafroller, *Platynota stultana*. *Environ. Entomol.*, **4**: 90-92.

Baker, R., J. W. S. Bradshaw and W. Speed, 1982. Methoxymercuration-demercuration and mass spectrometry in the identification of the sex pheromones of *Panolis flammea*, the pine beauty moth. *Experientia*, **38**: 233-234.

Baker, R., R. H. Herbert, N. N. Neequaye and K. N. Rao, 1985. Sex pheromone of tobacco stem borer, *Scrobipalpa heliopa* (Lower) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **11**: 989-998.

Baker, T. C., W. Francke, C. Löfstedt, B. S. Hansson, J.-W. Du, P. L. Phelan, R. S. Vetter and R. Youngman, 1989. Isolation, identification and synthesis of sex pheromone components of the carob moth, *Ectomyelois ceratoniae*. *Tetrahedron Lett.*, **30**: 2901-2902.

Baker, T. C., W. Francke, J. G. Millar, C. Löfstedt, B. Hansson, J.-W. Du, P. L. Phelan, R. S. Vetter, R. Youngman and J. L. Todd, 1991. Identification and bioassay of sex pheromone components of carob moth, *Ectomyelois ceratoniae* (Zeller). *J. Chem. Ecol.*, **17**: 1973-1988.

Baltensweiler, W., E. Priesner, H. Arn and V. Delucchi, 1978. Evidence for different sexual attractants used by the larch form and the cembran pine form of the grey larch bud moth (*Zeiraphera diniana* Gn.). *Mitt. Schweiz. Entomol. Ges.*, **51**: 133-142.

Baltensweiler, W. and E. Priesner, 1988. A study of pheromone polymorphism in *Zeiraphera diniana* Gn. (Lep., Tortricidae). 3. Specificity of attraction to synthetic pheromone source by different male response types from two host races. *J. Appl. Entomol.*, **106**: 217-231.

Barrer, P. M., M. J. Lacey and A. Shani, 1987. Variation in relative quantities of airborne sex pheromone components from individual female *Ephestia cautella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, 13: 639-653.

Barros-Parada, W., J. Bergmann, T. Curkovic, C. Espinosa, E. Fuentes-Contreras, J. Guajardo, H. Herrera, S. Morales, A. F. O. Queiroz and A. Vidal, 2020. 3,7-Dimethylpentadecane: a novel sex pheromone component from *Leucoptera sinuella* (Lepidoptera: Lyonetiidae). *J. Chem. Ecol.*, 46: 820-829.

Barry, M. W., D. G. Nielsen, F. F. Purrington and J. H. Tumlinson, 1978. Attractivity of pheromone blends to male peachtree borer, *Synanthedon exitiosa*. *Environ. Entomol.*, 7: 1-3.

Bartell, R. J., T. E. Bellas and C. P. Whittle, 1988. Evidence for biological activity of two further alcohols in the sex pheromone of female *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *J. Aust. Entomol. Soc.*, 27: 11-12.

Bathon, H. and M. Glas, 1983. The geographical distribution of the cereal leaf roller, *Cnephiasia pumicana* Zeller (Lepidoptera: Tortricidae), in the Federal Republic of Germany. Preliminary results of a pheromone trap survey. *Nachrichtenbl. Deut. Pflanzenschutzd.*, 35: 81-86.

Batista-Pereira, L. G., E. M. Santangelo, K. Stein, C. R. Unelius, A. E. Eiras and A. G. Correa, 2002. Electrophysiological studies and identification of possible sex pheromone components of Brazilian populations of the sugarcane borer, *Diatraea saccharalis*. *Z. Naturforsch.*, 57c: 753-758.

Batista-Pereira, L. G., K. Stein, A. F. de Paula, J. A. Moreira, I. Cruz, M. de L. C. Figueiredo, J. Perri Jr. and A. G. Correa, 2006. Isolation, identification, synthesis, and field evaluation of the sex pheromone of the Brazilian population of *Spodoptera frugiperda*. *J. Chem. Ecol.*, 32(5): 1085-1099.

Bavaresco, A., S. Nunez, M. S. Garcia, M. Botton and J. Sant'Ana, 2005. Attraction of males of the South American tortricid moth *Argyrotaenia sphaleropa* (Meyrick) (Lepidoptera: Tortricidae) to the components of the synthetic sexual pheromone in persimmon. *Neotrop. Entomol.*, 34: 619-625.

Beauvais, F., G. Biwer and P. J. Charmillot, 1977. Tests of sex trapping of *Grapholitha janthinana* Dup. with a mixture of Z-8 dodecenyl acetate containing 28 p. 100 of its isomer E. *Ann. Zool. Ecol. Anim.*, 9: 457-467.

Becker, D., T. Kimmel, R. Cyjon, I. Moore, M. Wysoki, H. J. Bestmann, H. Platz, K. Roth and O. Vostrowsky, 1983. (3Z,6Z,9Z)-3,6,9-Nonadecatriene -- a component of the sex pheromonal system of the giant looper, *Boarmia (Ascotis) selenaria* Schiffermüller (Lepidoptera: Geometridae). *Tetrahedron Lett.*,

24: 5505-5508.

Becker, D., R. Cyjon, A. Cossé, I. Moore, T. Kimmel and M. Wysoki, 1990. Identification and enantioselective synthesis of (*Z,Z*)-6,9-*cis*-3*S*,4*R*-epoxynonadecadiene, the major sex pheromone component of *Boarmia selenaria*. *Tetrahedron Lett.*, 31: 4923-4926.

Bédard, C., R. Gries, G. Gries and R. Bennett, 2002. Sex pheromone and diel periodicity of *Cydia strobilella* (Lepidoptera: Tortricidae) pheromonal communication. *Can. Entomol.*, 134: 847-850.

Beevor, P. S., D. R. Hall, R. Lester, R. G. Poppi, J. S. Read and B. F. Nesbitt, 1975. Sex pheromones of the armyworm moth, *Spodoptera exempta* (Wlk.). *Experientia*, 31: 22-23.

Beevor, P. S., D. R. Hall, B. F. Nesbitt, V. A. Dyck, G. Arida, P. C. Lippold and H. Oloumi-Sadeghi, 1977. Field trials of the synthetic sex pheromones of the stripe rice borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae), and of related compounds. *Bull. Entomol. Res.*, 67:439-447.

Beevor, P. S., A. Cork, D. R. Hall, B. F. Nesbitt, R. K. Day and J. D. Mumford, 1986. Components of female sex pheromone of cocoa pod borer moth, *Conopomorpha cramerella*. *J. Chem. Ecol.*, 12: 1-23.

Beevor, P. S., O. Youm, D. R. Hall and A. Cork, 1999. Identification and field evaluation of components of female sex pheromone of millet stem borer, *Coniesta ignefusalis*. *J. Chem. Ecol.*, 25(12): 2643-2663.

Bell, T. W. and J. Meinwald, 1986. Pheromones of two arctiid moths (*Creationotos transiens* and *C. gangis*): chiral components from both sexes and achiral female components. *J. Chem. Ecol.*, 12: 385-409.

Bellas, T. E., R. J. Bartell and A. Hill, 1983. Identification of two components of the sex pheromone of the moth, *Epiphyas postvittana* (Lepidoptera, Tortricidae). *J. Chem. Ecol.*, 9: 503-512.

Bengtsson, M., St. Rauscher, H. Arn, W.-C. Sun and G. D. Prestwich, 1990. Fluorine-substituted pheromone components affect the behavior of the grape berry moth. *Experientia*, 46: 1211- 1213.

Benn, M. H., R. A. Galbreath, V. A. Holt and H. Young, 1981. A sex attractant for *Sperchia intractana* Walker (Lepidoptera: Tortricidae) in New Zealand. *Z. Naturforsch.*, 36 c: 178-179.

Benn, M. H., R. A. Galbreath, V. A. Holt, H. Young, G. Down and E. Priesner, 1982. The sex pheromone of the silver Y moth *Chrysodeixis eriosoma* (Doubleday) in New Zealand. *Z. Naturforsch.*, 37 c: 1130-1135.

Benz, G. and G. von Salis, 1973. Use of synthetic sex attractant of larch bud moth *Zeiraphera diniana* (Gn.) in monitoring traps under different conditions, and antagonistic action of cis-isomer. *Experientia*, 29: 729-730.

Berger, R. S., 1966. Isolation, identification, and synthesis of the sex attractant of the cabbage looper, *Trichoplusia ni*. *Ann. Entomol. Soc. Am.*, 59: 767-771.

Berger, R. S. and T. D. Canerday, 1968. Specificity of the cabbage looper sex attractant. *J. Econ. Entomol.*, 61: 452-454.

Bergmann, J., L. Reyes-Garcia, C. Ballesteros, Y. Cuevae, M. F. Flores, and T. Curkovic, 2016. Identification of the female sex pheromone of the leafroller *Proeulia triquetra* Obraztsov (Lepidoptera: Tortricidae). *Neotrop. Entomol.*, 45: 351-356.

Beroza, M., K. Katagiri, Z. Iwata, H. Ishizuka, S. Suzuki and B. A. Bierl, 1973(a). Disparlure and analogues as attractants for two Japanese lymantriid moths. *Environ. Entomol.*, 2: 966.

Beroza, M., A. A. Punjabi and B. A. Bierl, 1973(b). Disparlure and analogues as attractants for *Lymantria obfuscata*. *J. Econ. Entomol.*, 66: 1215-1216.

Beroza, M., B. A. Bierl and H. R. Moffitt, 1974. Sex pheromones: (*E,E*)-8,10-dodecadien-1-ol in the codling moth. *Science*, 183: 89-90.

Bestmann, H. J., O. Vostrowsky, K.-H. Koschatzky, H. Platz, T. Brosche, I. Kantardjiew, M. Rheinwald and W. Knauf, 1978(a). (*Z*)-5-Decenyl acetate, a sex attractant for the male turnip moth *Agrotis segetum* (Lepidoptera). *Angew. Chem. Int. Ed. Engl.*, 17: 768.

Bestmann, H. J., O. Vostrowsky, K. H. Koschatzky, H. Platz, A. Szymanska and W. Knauf, 1978(b). Pheromone XVII. (*Z*)-11-Hexadecenylacetat, ein Sexuallockstoff des Pheromonsystems der Kohleule *Mamestra brassicae*. *Tetrahedron Lett.*, 605-608.

Bestmann, O. Vostrowsky, H. Platz, T. Brosche, K. H. Koschatzky, 1979. Pheromone XXIII. (*Z*)-7-Tetradecenylacetat, ein Sexuallockstoff für Männchen von *Amnathes c-nigrum* (Noctuidae, Lepidoptera). *Tetrahedron Lett.*, 497-500.

Bestmann, H. J., T. Brosche, K. H. Koschatzky, K. Michaelis, H. Platz, O. Vostrowsky and W. Knauf, 1980. Pheromone XXX. Identifizierung eines neuartigen Pheromonkomplexes aus der Graseule *Scotia exclamationis* (Lepidoptera). *Tetrahedron Lett.*, 21: 747-750.

Bestmann, H. J. and O. Vostrowsky, 1982(a). Insect pheromone. *Naturwissenschaften*, 69: 457-471.

Bestmann, H. J., K. H. Koschatzky, H. Platz, J. Süß O. Vostrowsky, W. Knauf, G. Burghrdt and I. Schneider, 1982(b). Pheromones, XL. Synthesis of the pheromone complex of Lasiocampidae species (Lepidoptera); a sex attractant for *Dendrolimus pini*. *Liebigs*

Ann. Chem., 1359-1365.

- Bestmann, H. J., T. Brosche, K. H. Koschatzky, K. Michaelis, H. Platz, K. Roth, J. Süß O. Vostrowsky and W. Knauf, 1982(c). Pheromone-XLII. 1,3,6,9-Nonadecatetraen, das Sexualpheromon des Frostspanners *Operophtera brumata* (Geometridae). *Tetrahedron Lett.*, 23: 4007-4010.
- Bestmann, H. J., K. H. Koschatzky and O. Vostrowsky, 1982(d). Stereoselective syntheses of 1,5- and 1,6-alkadienes as sex pheromones and pheromone analogues. *Liebigs Ann. Chem.*, 1478-1494.
- Bestmann, H.-J., A. Attygalle, H. Platz and O. Vostrowsky, 1985. (Z)-9-Dodecenyl acetate, a component of the sex pheromone of *Cnephiasia longana* Haworth (Lepidoptera: Tortricidae). *Z. Naturforsch.*, 40c: 667-669.
- Bestmann, H.-J., A. B. Attygalle, T. Brosche, J. Erler, H. Platz, J. Schwarz, O. Vostrowsky, C.-H. Wu, K. E. Kaissling and T.-M. Chen, 1987. Identification of three sex pheromone components of the female Saturniid moth *Antheraea pernyi* (Lepidoptera: Saturniidae). *Z. Naturforsch.*, 42c, 631-636.
- Bestmann, H. J., A. B. Attygalle, J. Schwarz, O. Vostrowsky and W. Knauf, 1988(a). Identification of sex pheromone components of *Spodoptera sunia* Guenée (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, 14: 683-690.
- Bestmann, H. J., J. Erler and O. Vostrowsky, 1988(b). Determination of diel periodicity of sex pheromone release in three species of Lepidoptera by 'closed-loop-stripping'. *Experientia*, 44: 797-799.
- Bestmann, H. J., A. B. Attygalle, J. Schwarz, W. Garbe, O. Vostrowsky and I. Tomida, 1989. Pheromones, 71. Identification and synthesis of female sex pheromone of eri-silkworm, *Samia cynthia ricini* (Lepidoptera: Saturniidae). *Tetrahedron Lett.*, 30: 2911-2914.
- Bestmann, H. J., F. Kern, D. Schäfer, O. Vostrowsky and I. Hasenfuss, 1991. (11Z,13Z)-11,13-Hexadecadienyl acetate, the sex pheromone of females of *Notodonta torva*. *Naturwissenschaften*, 78: 465-467.
- Bestmann, H. J., F. Kern, A. Mineif, H. Platz and O. Vostrowsky, 1992(a). Pheromone, 84. The sex pheromone complex of the arctiid moth *Arctia caja* (Lepidoptera: Arctiidae). *Z. Naturforsch.*, 47c: 132-135.
- Bestmann, H. J., J. Erler, W. Garbe, F. Kern, V. Martischonok, D. Schäfer, O. Vostrowsky and L. T. Wasserthal, 1992(b). Pheromone components of the female elephant hawk-moth, *Deilephila elpenor*, and the silver-striped hawk-moth, *Hippotion celerio*. *Experientia*, 48: 610-613.
- Bestmann, H. J., F. Kern, D. Schäfer and O. Vostrowsky, 1993(a). (11Z,13Z)-11,13-Hexadecadienal, the sex pheromone of females of *Notodonta dromedarius*. *Naturwissenschaften*, 80: 271-273.

Bestmann, H. J., F. Kern, G. G. Melikyan, D. Schäfer, O. Vostrowsky, E. V. Babayan and Sh. O. Badanyan, 1993(b). Pheromone, 88. Sex pheromone components of female *Euzophera punicaella* M. (Lepidoptera, Pyralidae). *Z. Naturforsch.*, 48c: 110-112.

Bestmann, H. J., F. Kern, E. Janssen and I. Hasenfust 1993(c). Pheromone, 93. The sex pheromone of the cosmopterigid moth *Limnaecia phragmitella* (Lepidoptera: Cosmopterigidae). *Z. Naturforsch.*, 48c: 515-518.

Bestmann, H. J., A. B. Attygalle, W. Garbe, F. Kern, V. Martichonok, D. Schäfer, O. Vostrowsky and I. Hasenfub 1993(d). Chemical structure and final steps of biosynthesis of the female sex pheromone of *Gastropacha quercifolia* (Lepidoptera: Lasiocampidae). *Insect Biochem. Molec. Biol.*, 23: 791-799.

Bestmann, H. J., E. Janssen, F. Kern, D. Schäfer and O. Vostrowsky, 1994. Pheromones, 97. The sex pheromone complex of the female arctiid moth *Thyria jacobaeae* (Lepidoptera, Arctiidae). *Z. Naturforsch.*, 49c: 276-279.

Bhanu, K. R. M., D. R. Hall, R. V. Awalekar, M. Chandrashekharaiyah, T. N. Divya, M. S. Prabhakara and K. P. Jayanth, 2018. Identification and field evaluation of female sex pheromone of leaf-eating caterpillar, *Opisina arenosella* (Lepidoptera: Oecophoridae). *Int. J. Tropic. Insect Sci.*, 38: 274-282.

Bhardwaj, S. P., 1987. Investigations on the response of lepidopteran sex pheromones of temperate fruit pests in Himachal Pradesh, India. *Agric. Ecosyst. Environ.*, 19: 87-91.

Biddinger, D. J., W. Roelofs and A. J. Howitt, 1994. The development of a sex pheromone lure for the American plum borer, *Euzophera semifuneralis* (Lepidoptera: Pyralidae), a major pest of cherry in Michigan. *Great Lakes Entomol.*, 26: 311-317.

Bierl, B. A., M. Beroza and C. W. Collier, 1970. Potent sex attractant of the gypsy moth: its isolation, identification, and synthesis. *Science*, 170: 87-89.

Bierl, B. A., M. Beroza, R. T. Staten, P. E. Sonnet and V. E. Adler, 1974. The pink bollworm sex attractant. *J. Econ. Entomol.*, 67: 211-216.

Bierl, B. A., M. Beroza, V. E. Adler, G. Kasang, H. Schröter and D. Schneider, 1975. The presence of disparlure, the sex pheromone of the gypsy moth, in the female nun moth. *Z. Naturforsch.*, 30 c: 672-675.

Birch, M. C., 1977. Response of both sexes of *Trichoplusia ni* (Lepidoptera: Noctuidae) to virgin females and to synthetic pheromone. *Ecol. Entomol.*, 2: 99-104.

Biwer, G., B. Lalanne-Cassou, C. Descoins and D. Samain, 1975. Sex trapping of *Sterrhia biselata* (Lepidoptera, Geometridae,

Sterrhinae) by (7E,9Z)-dodecadienyl acetate, a sex pheromone for *Lobesia botrana* (Lepidoptera, Tortricidae, Olethreutinae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **280**: 1469-1471.

Biwer, G., C. Descoins, M. Gallois, E. Priesner, J. P. Chambon, G. Genestier and M. Martinez, 1978 (a). Study of the sex pheromone blend produced by the virgin female of the cereal tortrix: *Cnephiasia pumicana* Zeller. *Ann. Zool. Ecol. Anim.*, **10**: 129-138.

Biwer, G. and C. Descoins, 1978 (b). Approach to a mechanism of sexual isolation between four species of tortricids of the *Grapholitha* genus. *C. R. Hebd. Seances Acad. Sci., Ser. D*, **286**: 875-877.

Biwer, G., C. Descoins and M. Gallois, 1979. Volatile components from the sex pheromone-producing gland of *Grapholitha molesta* Busk virgin females (Lepidoptera Tortricidae, Olethreutinae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **288**: 413-416.

Bjostad, L. B., E. F. Taschenberg and W. L. Roelofs, 1980 (a). Sex pheromone of the chokecherry leafroller moth, *Sparganothis directana*. *J. Chem. Ecol.*, **6**: 487-498.

Bjostad, L. B., L. K. Gaston, L. L. Noble, J. H. Moyer and H. H. Shorey, 1980 (b). Dodecyl acetate, a second pheromone component of the cabbage looper moth, *Trichoplusia ni*. *J. Chem. Ecol.*, **6**: 727-734.

Bjostad, L. B., E. F. Taschenberg and W. L. Roelofs, 1980 (c). Sex pheromone of the woodbine leafroller moth, *Sparganothis* sp. *J. Chem. Ecol.*, **6**: 797-804.

Bjostad, L. B., E. Gurevitz, S. Gothilf and W. L. Roelofs, 1981. Sex attractant for the honeydew moth, *Cryptoblabes gnidiella*. *Phytoparasitica*, **9**: 95-99.

Bjostad, L. B., C. E. Linn, J.-W. Du and W. L. Roelofs, 1984. Identification of new sex pheromone components in *Trichoplusia ni*, predicted from biosynthetic precursors. *J. Chem. Ecol.*, **10**: 1309-1323.

Blair, B. W. and J. Tannock, 1978. A further note on the possible pheromone for *Spodoptera tritubata* (Walker) (Lepidoptera, Noctuidae). *Rhod. J. Agric. Res.*, **16**: 221-223.

Blassioli-Moraes, M. C., M. Borges, A. R. Viana, R. A. Laumann, J. E. Miranda, D. M. Magalhães and M. A. Birkett, 2016. Identification and field evaluation of the sex pheromone of a Brazilian population of *Spodoptera cosmioides*. *Pesq. Agropec. Bras.*, **51**: 545-554.

Bogenschütz, von H., W. Knauf, E. J. Tröger, H. J. Bestmann and O. Vostrowsky, 1985. Pheromones 49: males of Geometridae caught by C₁₉-polyenes in pine stands. *Z. Angew. Entomol.*, **100**: 349-354.

Boness, M., 1978. Experiences with sex pheromone of Lepidoptera. *Anz. Schädlingskde.*, **51**: 161-166.

- Boo**, K. S. and C. H. Jung **1998** (a). Field tests of synthetic sex pheromone of the apple leafminermoth, *Phyllonorycterringoniella*. *J. Chem. Ecol.*, **24** (12): 1939-1947.
- Boo, K. S. and J. W. Park, **1998** (b). Sex pheromone composition of the Asian corn borer moth, *Ostrinia furnacalis* (Guenée) (Lepidoptera: Pyralidae) in South Korea. *J. Asia Pacific Entomol.*, **1**: 77-84.
- Booij**, C. J. H. and S. Voerman, **1984** (a). New sex attractants for 35 tortricid and 4 other lepidopterous species, found by systematic field screening in the Netherlands. *J. Chem. Ecol.*, **10**: 135-144.
- Booij, C. J. H. and S. Voerman, **1984** (b). Sex attractants for the fruit-attacking noctuids *Orthosia incerta* and *Orthosia cruda*. *Entomol. Exp. Appl.*, **36**: 15-16.
- Booij, C. J. H. and S. Voerman, **1984** (c). (*Z*)-11-Hexadecenyl compounds as attractants for male microlepidoptera of the subfamilies Argyresthiinae, Glyphipteryginae, and Crambinae. *Entomol. Exp. Appl.*, **36**: 47-53.
- Booij, C. J. H., **1984** (d). *Cydia indivisa* (Danilevski), een nieuwe bladrollersoort voor de Nederlandse fauna, gevangen met sekslokstoffen (Lepidoptera: Tortricidae). *Entomol. Ber.*, **44**: 133-135.
- Booij, C. J. H. and S. Voerman, **1984** (e). Sex attractant for the poplar shoot-borer, *Gypsonoma aceriana* (Duponchel). *Z. Angew. Entomol.*, **97**: 176-179.
- Booij, C. J. H. and S. Voerman, **1985**. New lepidopteran sex attractants found by systematic field screening of blends containing (*Z*)-11- and (*E*)-11-tetradecenal. *J. Chem. Ecol.*, **11**: 1333-1339.
- Booij, C. J. H., S. Voerman and L. P. M. Willemse, **1986**. Conifer-infesting tortricids in Western Europe: a search for more potent and selective sex attractants. *J. Appl. Entomol.*, **102**: 253-259.
- Borek**, V., B. Kalinová, I. Valterová, R. Hochmut and J. Vrkoc~, **1991**. Sex pheromone gland volatiles from *Hypsipyla grandella* females (Lepidoptera, Pyralidae, Phycitinae). *Acta. Entomol. Boemoslov.*, **88**: 181-186.
- Borg-Karlsson**, A.-K. and R. Mozūraitis, **1996**. Solid phase micro extraction technique used for collecting semiochemicals. Identification of volatiles released by individual signalling *Phyllonorycter sylvella* moths. *Z. Naturforsch.*, **51c**: 599-602. (See Ph-3-2-6)
- Bosa**, C. F., A. M. C. Prado, T. Fukumoto, M. Bengtsson and P. Witzgall, **2005**. Pheromone-mediated communication disruption in Guatemalan potato moth, *Tecia solanivora*. *Entomol. Exp. Appl.*, **114**: 137-142.
- Bosson**, G. A. and M. Gallois, **1982**. Analysis of pheromonal secretion in virgin females of *Hypsipyla robusta* (Moore) (Lepidoptera,

Pyralidae, Phycitinae) from young mahogany shoots. *C. R. Acad. Sci., Ser. 3*, **294**: 819-822.

Bourdouxhe, L. 1982. Results of two years of sex trapping of *Cryptophlebia leucotreta* Meyr. in Senegal. *FAO Plant Prot. Bull.*, **30**: 125-129.

Bouwer, M. C., B. Slippers, D. Degefu, M. J. Wingfield, S. Lawson and E. R. Rohwer, 2015. Identification of the sex pheromone of the tree infesting cossid moth *Coryphodema tristis* (Lepidoptera: Cossidae). *PloS ONE.*, **10**: e0118575.

Bouwer, M. C., B. Slippers, M. J. Wingfield, J. D. Allison and E. R. Rohwer, 2017. Optimization of pheromone traps for *Coryphodema tristis* (Lepidoptera: Cossidae). *J. Econ. Entomol.*, **110**: 1603-1610.

Bouwer, M. C., L.-M. Scheepers, B. Slippers, E. R. Rohwer and J. D. Allison, 2021. The sex pheromone of the pine brown-tail moth, *Euproctis terminalis* (Lepidoptera: Erebidae). *J. Chem. Ecol.*, **47**: 732-739.

Bradshaw, J. W. S., R. Baker, C. Longhurst, J. C. Edwards and J. C. Lisk, 1983. Optimization of a monitoring system for the pine beauty moth, *Panolis flammea* (Denis & Schiffermüller), using sex attractants. *Crop Prot.*, **2**: 63-73.

Brady, U. E., J. H. Tunlinson III, R. G. Brownlee and R. M. Silverstein, 1971(a). Sex stimulant and attractant in the Indian meal moth and in the almond moth. *Science*, **171**: 802-804.

Brady, U. E. and D. A. Nordlund, 1971(b). *Cis-9,trans-12 tetradecadien-1-yl acetate* in the female tobacco moth *Ephestia elutella* (Hübner) and evidence for an additional component of the sex pheromone. *Life Sci.*, **10**: 797-801.

Brady, U. E. and M. C. Ganyard, Jr., 1972(a). Identification of a sex pheromone of the female beet armyworm, *Spodoptera exigua*. *Ann. Entomol. Soc. Am.*, **65**: 898-899.

Brady, U. E. and R. C. Daley, 1972(b). Identification of a sex pheromone from the female raisin moth, *Cadra figulilella*. *Ann. Entomol. Soc. Am.*, **65**: 1356-1358.

Brady, U. E., 1973. Isolation, identification and stimulatory activity of a second component of the sex pheromone system (complex) of the female almond moth, *Cadra cautella* (Walker). *Life Sci.*, **13**: 227-235.

Brennan, B. M., 1977. Attraction of *Ereunetis simulans* (Lepidoptera: Tineidae) to the sex attractant of peachtree borer, *Sanninoidae exitiosa* (Lepidoptera: Sesiidae). *Proc. Hawaii. Entomol. Soc.*, **22**: 457-460.

Breuer, M., H.-G. Kontzog, A. Guerrero, F. Camps and A. de Loof, 2003.

Field trials with the synthetic sex pheromone of the oak processionary moth *Thaumetopoea processionea*. *J. Chem. Ecol.*, **29**(11): 2461-2468.

Brewer, J. W., F. Krampl and V. Skuhravy', 1985. Forest Lepidoptera attracted to six synthetic pheromones in Czechoslovakia. *Z. Angew. Entomol.*, 100: 372-381.

Brown, L. N. and J. W. Snow, 1985(a). The blackberry clearwing borer, *Pennisetia marginata* (Harris): a first report in Florida. *Florida Entomol.*, 68: 699-700.

Brown, L. N., 1985(b). New records of the rare clearwing moth, *Alcathoe carolinensis* Engelhardt, (Sesiidae) in Florida. *Florida Entomol.*, 68: 700-701.

Brown, L. N., T. D. Eichlin and J. W. Snow, 1985(c). A new species of clearwing moth, *Carmenta laurelae* (Sesiidae), from Florida. *J. Lepido. Soc.*, 39: 262-265.

Brown, L. N., T. D. Eichlin and J. W. Snow, 1985(d). Ecological notes on *Synanthedon dominicki* Duckworth and Eichlin (Sesiidae) in Florida and first description of the female. *J. Lepido. Soc.*, 39: 196-200.

Brown, L. N. and J. W. Snow, 1986(a). First record of the clearwing moth, *Carmenta odda* (Lepidoptera: Sesiidae) in Florida. *Florida Entomol.*, 69: 423-424.

Brown, L. N. 1986(b). First records of three clearwing moths (*Synanthedon tabaniformis*, *Synanthedon proxima*, and *Synanthedon castaneae*) in Florida. *Florida Scientist*, 49: 80-81.

Brückner, C., E. Beschmann, R. Becker and W. Seufert, 1988. A new highly effective synthetic pheromone mimic for *Lobesia botrana* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, 43 c: 315-318.

Bûda, V., V. Karalius and P. Ivinskis, 1987. Sex attractants for *Pyrausta ostrinalis* Hbn. (Lepidoptera, Pyralidae) and leafroller *Apotomis capreana* Hbn. (Lepidoptera, Tortricidae). *Khemoretspsiya Nasekomykh*, 9: 84-88.

Bûda, V., U. Mäeorg, V. Karalius, G. H. L. Rothschild, S. Kolonistova, P. Ivinskis and R. Mozūraitis, 1993. C_{18} Dienes as attractants for eighteen clearwing (Sesiidae), tineid (Tineidae), and choreutiid (Choreutidae) moth species. *J. Chem. Ecol.*, 19: 799-813.

Buechi, R., E. Priesner and R. Brunetti, 1982. Sympatric occurrence of two pheromonal strains of the European corn borer, *Ostrinia nubilalis* Hbn., in southern Switzerland. *Mitt. Schweiz. Entomol. Ges.*, 55: 33-53.

Buleza, V. V., 1991. Role of sex pheromone components in male reproduction behavior of black cutworm moth *Agrotis ipsilon*. *Dokl. Akad. Nauk SSSR*, 318: 1265-1268. [Chem. Abst., 116: 53588t]

Burghardt, G., W. Knauf, H. J. Bestmann, O. Vostrowsky and K. H. Koschatzky, 1980. Synthetic pheromones and their inhibitors for *Choristoneura murinana* Hbn. (silver-fir shoot tortricid. Lepidoptera: Tortricidae). *Anz. Schädlingskde.*, 53: 49-51.

Burns, E. L. and P. E. A. Teal, 1989. Response of male potato stem borer moths, *Hydraecia micacea* (Esper) to conspecific females and synthetic pheromone blends in the laboratory and field. *J. Chem. Ecol.*, 15: 1365-1378.

Buser, H.-R., S. Rauscher and H. Arn, 1974. Sex pheromone of *Lobesia botrana*: (E,Z)-7,9-dodecadienyl acetate in the female grape vine moth. *Z. Naturforsch.*, 29 c: 781-783.

Buser, H. R., P. M. Guerin, M. Tóth, G. Szöcs, A. Schmid, W. Francke and H. Arn, 1985. (Z,Z)-6,9-Nonadecadien-3-one and (Z,Z,Z)-3,6,9-nonadecatriene: identification and synthesis of sex pheromone components of *Peribatodes rhomboidaria*. *Tetrahedron, Lett.*, 26: 403-406.

Butenandt, von A., R. Beckmann, D. Stamm and E. Hecker, 1959. Über den Sexual-Lockstoff des Seidenspinners *Bombyx mori*. Reindarstellung und Konstitution. *Z. Naturforsch.*, 14 b: 283-284.

Butenandt, A., R. Beekmann and E. Hecker, 1961. Über den Sexuallockstoff des Seidenspinners, I. Der biologische Test und Isolierung des reinen Sexuallockstoffes Bombykol. *Hoppe-Seyler's Z. Physiol. Chem.*, 324: 71-83.

Butler, L. I., J. E. Halfhill, L. M. McDonough and B. A. Butt, 1977. Sex attractant of the alfalfa looper *Autographa californica* and the celery looper *Anagrapha falcifera* (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, 3: 65-70.

Byers, J. R., E. W. Underhill, W. F. Steck, M. D. Chisholm and P. E. A. Teal, 1981. Biosystematics of the genus *Euxoa* (Lepidoptera: Noctuidae). XV. Sex pheromone cross attractancy among the three closely related species of the *declarata* group. *Can. Entomol.*, 113: 235-243.

Byers, J. R. and D. L. Struble, 1987. Monitoring population levels of eight species of noctuids with sex-attractant traps in southern Alberta, 1978-1983: specificity of attractants and effect of target species abundance. *Can. Entomol.*, 119: 541-556.

Byers, J. R. and D. L. Struble, 1990. Identification of sex pheromones of two sibling species in dingy cutworm complex, *Feltia jaculifera* (Gn.) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, 16: 2981-2992.

Byers, J. R. and C. E. Herle, 1997. Identification of sex pheromones of *Leucania anteoclara* Sm. and *Leucania commoides* Gn. (Lepidoptera: Noctuidae; Hadeninae). *J. Chem. Ecol.*, 23 (2): 473-485.

Cabrera, A., A. Eiras, G. Gries, R. Gries, N. Urdaneta, B. Miras, C.

Badji and K. Jaffe, 2001. Sex pheromone of tomato fruit borer, *Neoleucinodes elegantalis*. *J. Chem. Ecol.*, 27(10): 2097-2107.

Campbell, C. A. M., E. L. Tregidga, D. R. Hall, T. Ando and M. Yamamoto, 2007. Components of the sex pheromone of the currant pug moth *Eupithecia assimilata*, a re-emergent hop pest in UK. *Entomol. Exp. Appl.*, 122: 265-269.

Campion, D. G., B. W. Bettany, B. F. Nesbitt, P. S. Beevor, R. Lester and R. G. Poppi, 1974. Field studies of the female sex pheromone of the cotton leafworm *Spodoptera littoralis* (Boisd.) in Cyprus. *Bull. Entomol. Res.*, 64: 89-96.

Campion, D. G., 1975. Sex pheromones and their uses for control of insects of the genus *Spodoptera*. *Meded. Fac. Landbouww. Rijksuniv. Gent.*, 40: 283-292.

Campion, D. G., L. J. McVeigh, J. Polyrakis, S. Michaelakis, G. N. Stravrakakis, P. S. Beevor, D. R. Hall and B. F. Nesbitt, 1979. Laboratory and field studies of the female sex pheromone of the olive moth, *Prays oleae*. *Experientia*, 35: 1146-1147.

Campion, D. G., P. Hunter-Jones, L. J. McVeigh, D. R. Hall, R. Lester and B. F. Nesbitt, 1980. Modification of the attractiveness of the primary pheromone component of the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae), by secondary pheromone components and related chemicals. *Bull. Entomol. Res.*, 70: 417-434.

Can, F., K. A. Efetov, J. Burman, K. Kaya, E. E. Kucherenko, B. Ulaşlı and G. M. Tarmann, 2019. A study of the Zygaenidae (Lepidoptera) fauna of Central Anatolia, Turkey. *Turk. J. Entomol.*, 43: 189-199.

Cao, W.-H. and H.-J. He, 2002. Identification of sex pheromone of the moth, *Tyta luctuosa*. *Act. Entomol. Sinica (Kunchong Xuebao)*, 45: 30-34.

Cao, W.-H., R. E. Charlton, J. R. Nechols and M. J. Horak, 2003. Sex pheromone of the noctuid moth, *Tyta luctuosa* (Lepidoptera: Noctuidae), a candidate biological control agent of field bindweed. *Environ. Entomol.*, 32(1): 17-22.

Capek, M., 1979. The influence of orientation of pheromone traps on their attractivity for gypsy moth (*Lymantria dispar* L.). *Lesnictivi*, 25: 301-306.

Capizzi, A., C. Tonini, E. Arsura, G. Guglielmetti, P. Massardo and P. Piccardi, 1983. Sex pheromone components of the European goat moth, *Cossus cossus*. *J. Chem. Ecol.*, 9: 191-200.

Cardé, R. T., J. Kochansky, J. F. Stimmel, A. G. Wheeler, Jr. and W. L. Roelofs, 1975. Sex pheromone of the European corn borer (*Ostrinia nubilalis*): cis- and trans-responding males in Pennsylvania. *Environ. Entomol.*, 4: 413-414.

Cardé, R. T., A. M. Cardé, A. S. Hill and W. L. Roelofs, 1977(a). Sex pheromone specificity as a reproductive isolating mechanism among the sibling species *Archips argyrospilus* and *A. mortuanus* and other sympatric tortricine moths (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, 3: 71-84.

Cardé, R. T., C. C. Doane, T. C. Baker, S. Iwaki and S. Marumo, 1977(b). Attractancy of optically active pheromone for male gypsy moths. *Environ. Entomol.*, 6: 768-772.

Cardé, A. M., T. C. Baker and R. T. Cardé, 1979. Identification of a four-component sex pheromone of the female Oriental fruit moth, *Grapholita molesta* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, 5: 423-427.

Carney, R. L. and A. S. T. Lui, 1982. Insect pheromone. *U. S. Patent* 4,357,474. [*Chem. Abst.*, 98: 125738b]

Castellari, P. L., 1985. Oblique-banded leaf roller *Choristoneura lafauryana* Rag. (Lepidoptera, Tortricidae): field tests with synthetic sex pheromone blends. *Bull. Ist. Entomol. Univ. Bologna*, 39: 243-260.

China, 1978. Isolation, identification, and synthesis of the sex pheromone of *Argyroploce (Eucosma) schistaceana* Snellen (Lepidoptera: Tortricidae). *K'o Hsueh T'ung Pao*, 23: 438-441, 435.

China, 1979. Isolation, identification and synthesis of EAG-active components of the pine-caterpillar moth sex pheromone. *K'o Hsueh T'ung Pao*, 21: 1004-1008.

Chamberlain, D. J., P. S. Beevor, A. Cork and D. R. Hall, 2003. (*E*)-8-Dodecenyl acetate: major component of the female sex pheromone of a Macadamia nut borer, *Ecdytolopha torticornis*. *Entomol. Exp. Appl.*, 107: 91-93.

Chambon, J. P. and J. d'Aguilar, 1974. Remarks on the specificity of some synthetic sexual pheromones: systematic study of tortricids in orchards. *Ann. Zool. Ecol. Anim.*, 6: 423-430.

Chang, V. C. S., 1995. Trapping *Cryptophlebia illepida* and *C. ombrodelta* (Lepidoptera: Tortricidae) in macadamia in Hawaii. *Int. J. Pest Manage.*, 41(2): 104-108.

Charlton, R. E., J. A. Wyman, J. R. McLaughlin, J.-W. Du and W. L. Roelofs, 1991. Identification of sex pheromone of tomatopinworm, *Keiferia lycopersicella* (Wals.). *J. Chem. Ecol.*, 17: 175-183.

Chen, X., K. Nakamura, T. Nakanishi, T. Nakashima, M. Tokoro, F. Mochizuki and T. Fukumoto, 2006. Female sex pheromone of a carpenter moth, *Cossus insularis* (Lepidoptera: Cossidae). *J. Chem. Ecol.*, 32(3): 669-679.

Chen, G.-F., M.-L. Sheng, T. Li, J. G. Millar and Q.-H. Zhang, 2010. Synergistic sex pheromone components of the grey-spotted tussock moth, *Orgyia ericae*. *Entomol. Exp. Appl.*, 136: 227-234.

- Cheng**, Z.-Q., J.-C. Xiao, X.-T. Huang, D.-L. Chen, J.-Q. Li, Y.-S. He, S.-R. Huang, Q.-C. Luo, C.-M. Yang and T.-H. Yang, 1981. Sex pheromone components isolated from Chaina corn borer, *Ostrinia furnacalis* Guenée (Lepidoptera: Pyralidae), (*E*)- and (*Z*)-12-tetradecenyl acetates. *J. Chem. Ecol.*, 7: 841-851.
- Cheng, Z.-Q. 1989. Sex pheromones of two species of corn borers living in China. *Acta Phytopathol. Entomol. Hung.*, 24: 237-240.
- Cheng, X., J. Wang and J. Zhang, 2007. Synthesis and field tests of sex attractant for *Holcocerus hippophaecolus* (Lepidoptera: Cossidae). *Linye Kexue*, 43(5): 74-77. [SciFinder, 2007:697769]
- Chisholm**, M. D., W. F. Steck, A. P. Arthur and E. W. Underhill, 1975. Evidence for *cis*-11-hexadecen-1-ol acetate as a major component of the sex pheromone of the bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae). *Can. Entomol.*, 107: 361-366.
- Chisholm, M. D., E. W. Underhill and W. F. Steck, 1979. Field trapping of the diamondback moth *Plutella xylostella* using synthetic sex attractants. *Environ. Entomol.*, 8: 516-518.
- Chisholm, M. D., E. W. Underhill, W. Steck, K. N. Slessor and G. G. Grant, 1980(a). (*Z*)-5, (*E*)-7-Dodecadienal and (*Z*)-5, (*E*)-7-dodecadien-1-ol, sex pheromone components of the forest tent caterpillar, *Malacosoma disstria*. *Can. Entomol.*, 9: 278-282.
- Chisholm, M. D., W. Steck and E. W. Underhill, 1980(b). Effects of additional double bonds on some olefinic moth sex attractants. *J. Chem. Ecol.*, 6: 203-212.
- Chisholm, M. D., P. Palaniswamy and E. W. Underhill, 1982. Orientation disruption of male forest tent caterpillar, *Malacosoma disstria* (Hübner) (Lepidoptera: Lasiocampidae), by air permeation with sex pheromone components. *Environ. Entomol.*, 11: 1248-1250.
- Chisholm, M. D., W. F. Steck, E. W. Underhill and P. Palaniswamy, 1983. Field trapping of diamondback moth *Plutella xylostella* using an improved four-component sex attractant blend. *J. Chem. Ecol.*, 9: 113-118.
- Chisholm, M. D., D. W. Reed, E. W. Underhill, P. Palaniswamy and J. W. Wong, 1985(a). Attraction of tortricid moths of subfamily Olethreutinae to field traps baited with dodecadienes. *J. Chem. Ecol.*, 11: 217-230.
- Chisholm, M. D. and D. W. Reed, 1985(b). 9,11-Dodecadienyl alcohols, acetates, or aldehydes as synthetic chemical sex attractants for four Lepidoptera: *Cosmopterix gemmiferella* (Clemens), *Dichrorampha simulana* (Clemens), *Tortricidiatestacea* (Packard), and an *Ancylis* sp.. *J. Chem. Ecol.*, 11: 1341-1347.
- Cho**, J. R., H. H. Park, B. Y. Seo, J. H. Kim, K. H. Yum, J. K. Jung and C. Y. Yang, 2014. Identification and application of the sex pheromones of the rice green caterpillar *Naranga aenescens* In

Korea. *J. Asia-Pac. Entomol.*, **17**: 191-197.

Choi, M.-Y., H. Lim, K. C. Park, R. Adlof, S. Wang, A. Zhang and R. Jurenka **2007**. Identification and biosynthetic studies of the hydrocarbon sex pheromone in *Utetheisa ornatrix*. *J. Chem. Ecol.*, **33** (7): 1336-1345.

Choi, K. S., J. M. Lee, J. H. Park, J. R. Cho, J. H. Song, D. S. Kim and K. S. Boo, **2009**(a). Sex pheromone composition of the cotton caterpillar, *Palpita indica* (Lepidoptera: Pyralidae), in Korea. *J. Asia-Pac. Entomol.*, **12**: 269-275.

Choi, K. S., J. R. Cho, J. H. Song, D.-S. Kim and K. S. Boo, **2009**(b). Sex pheromone composition of the variegated cutworm, *Peridroma saucia* (Lepidoptera: Noctuidae), in Korea. *J. Asia-Pac. Entomol.*, **12**: 71-77.

Choi, K. S., Y. M. Park, K. H. Choi, D. H. Kim and D. S. Kim, **2012**. Sex pheromone composition of *Ascotis selenaria* (Lepidoptera: Geometridae) and its regional variation in Korea. *J. Asia-Pac. Entomol.*, **15**: 413-418.

Chow, Y. S., Y. M. Lin and C. L. Hsu, **1977**. Sex pheromone of the diamondback moth (Lepidoptera: Plutellidae). *Bull. Inst. Zool., Academia Sinica*, **16**: 99-105.

Chow, Y. S., R. S. Tsai, G. Gries, R. Gries and G. Khaskin, **2001**. Preliminary identification of the sex pheromone of *Orgyia postica* (Walker) (Lepidoptera: Lymantriidae). *Entomol. Sinica*, **8**: 13-20.

Cichón, L. I., P. Trematerra, M. D. A. Coracini, D. Fernandes, M. Bengtsson and P. Witzgall, **2004**. Sex pheromone of *Argyrotaenia pomililiana* (Lepidoptera: Tortricidae), a leafroller pest of apples in Argentina. *J. Econ. Entomol.*, **97**(3): 946-949.

Clark, J. D. and K. F. Haynes, **1990**. Sex attractant for the bluegrass webworm (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **83**: 856-859.

Clarke, C. A., A. Cronin, W. Francke, P. Philipp, J. A. Pickett, L. J. Wadhams and C. M. Woodcock, **1996**. Mating attempts between the scarlet tiger moth, *Callimorpha dominula* L., and the cinnabar moth, *Tyria jacobaeae* L. (Lepidoptera: Arctiidae), involve a common sex pheromone composition. *Experientia*, **52**: 636-638.

Clearwater, J. R., R. A. Galbreath, M. H. Benn and H. Young, **1986**. Female-produced sexual pheromone of *Sceliodes cordalis* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **12**: 1943-1964.

Clearwater, J. R., S. P. Foster, S. J. Mugglestone, J. S. Dugdale and E. Priesner, **1991**. Intraspecific variation and interspecific differences in sex pheromones of sibling species in *Ctenopseustis obliquana* complex. *J. Chem. Ecol.*, **17**: 413-429.

Coffelt, J. A., K. W. Vick, P. E. Sonnet and R. E. Doolittle, **1979**.

Isolation, identification, and synthesis of a female sex pheromone of the navel orangeworm, *Amyelois transitella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **5**: 955-966.

Conner, W. E., T. Eisner, R. K. V. Meer, A. Guerrero, D. Ghiringelli and J. Meinwald, **1980**. Sex attractant of an arctiid moth (*Uteheisa ornatrix*): a pulsed chemical signal. *Behav. Ecol. Sociobiol.*, **7**: 55-63.

Coracini, M. D. A., M. Bengtsson, A. Reckziegel, J. Lofqvist, W. Francke, E. F. Vilela, A. E. Eiras, A. Kovaleski and P. Witzgall, **2001**. Identification of a four-component sex pheromone blend in *Bonagota cranaodes* (Lepidoptera: Tortricidae). *J. Econ. Entomol.*, **94**: 911-914.

Coracini, M. D. A., M. Bengtsson, A. Reckziegel, A. E. Eiras, E. F. Vilela, P. Anderson, W. Francke, J Lofqvist and P. Witzgall, **2003**. Behavioural effects of minor sex pheromone components in Brazilian apple leafroller *Bonagota cranaodes* (Lep., Tortricidae). *J. Appl. Entomol.*, **127**: 427-434.

Cork, A., P. S. Beevor, D. R. Hall, B. F. Nesbitt, G. S. Arida and O. Mochida, **1985** (a). Components of the female sex pheromone of the yellow stem borer, *Scirpophaga incertulas*. *Entomol. Exp. Appl.*, **37**: 149-153.

Cork, A., P. S. Beevor, D. R. Hall, Nesbitt B. F. and D. G. Campion, **1985** (b). A sex attractant for the spotted bollworm *Earias vittella*. *Trop. Pest Manag.*, **31**: 156-158.

Cork, A., D. J. Chamberlain, P. S. Beevor, D. R. Hall, B. F. Nesbitt, D. G. Campion and M. R. Attique, **1988**. Components of female sex pheromone of spotted bollworm, *Earias vittella* F. (Lepidoptera: Noctuidae): identification and field evaluation in Pakistan. *J. Chem. Ecol.*, **14**: 929-945.

Cork, A., J. Murlis and T. Megenasa, **1989**. Identification and field testing of additional components of female sex pheromone of African armyworm, *Spodoptera exempta* (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **15**: 1349-1364.

Cork, A., M. Agyen-Sampong, S. J. Fannah, P. S. Beevor and D. R. Hall, **1991**. Sex pheromone of female African white rice stem borer, *Maliarpha separatella* (Lepidoptera: Pyralidae) from Sierra Leone: identification and field testing. *J. Chem. Ecol.*, **17**: 1205-1219.

Cork, A., K. S. Boo, E. Dunkelblum, D. R. Hall, K. Jee-Rajunga, M. Kehat, E. Kong Jie, K. C. Park, P. Tepgidagarn and Liu Xun, **1992**. Female sex pheromone of Oriental tobacco budworm, *Helicoverpa assulta* (Guenée) (Lepidoptera: Noctuidae): identification and field testing. *J. Chem. Ecol.*, **18**: 403-418.

Cork, A., S. N. Alam, A. Das, C. S. Das, G. C. Ghosh, D. I. Farman, D. R. Hall, N. R. Maslen, K. Vedham, S. J. Phythian, F. M. A. Rouf and K. Srinivasan, **2001**. Female sex pheromone of brinjal fruit and shoot borer, *Leucinodes orbonalis* blend optimization. *J. Chem. Ecol.*, **27** (9): 1867-1877.

- Cork, A. and E. A. Lobos, 2003. Female sex pheromone components of *Helicoverpa gelotopoeon*: first heliothine pheromone without (Z)-11-hexadecenal. *Entomol. Exp. Appl.*, 107: 201-206.
- Coroiu**, I., GH. Stan, N. Tomescu, M. C. Roman, A. T. Dragotel, I. Oprean and L. Pop, 1986. Attractant and specificity of some pheromonal synthetic compounds in *Agrotis segetum* (Lepidoptera: Noctuidae). *Rev. Roum. Biol., Ser. Biol. Anim.*, 31: 109-118.
- Cortés**, A. M. P., P. H. G. Zarbin, D. M. Takiya, J. M. S. Bento, A. S. Guidolin and F. L. Consoli, 2010. Geographic variation of sex pheromone and mitochondrial DNA in *Diatraea saccharalis* (Feb., 1794) (Lepidoptera: Crambidae). *J. Insect Physiol.*, 56: 1624-1630.
- Cory**, H. T., G. E. Daterman, G. D. Daves, Jr., L. L. Sower, R. F. Shepherd and C. J. Sanders, 1982. Chemistry and field evaluation of the sex pheromone of western spruce budworm, *Choristoneura occidentalis*, Freeman. *J. Chem. Ecol.*, 8: 339-350.
- Cossé**, A. A., R. Cyjon, I. Moore, M. Wysoki and D. Becker, 1992. Sex pheromone components of the giant looper, *Boarmia selenaria* Schiff. (Lepidoptera: Geometridae): identification, synthesis, electrophysiological evaluation, and behavioral activity. *J. Chem. Ecol.*, 18: 165-181.
- Cowles**, R. S., J. G. Millar, E. J. Perry, J. S. McElfresh and N. Garrison, 1996. Identification of the sex pheromone of the western poplar clearwing moth (Lepidoptera: Sesiidae). *Environ. Entomol.*, 25; 109-113.
- Cuevas**, P., R. Montoya, X. Belles, F. Camps, J. Coll, A. Guerrero and M. Riba, 1983. Initial field trials with the synthetic sex pheromone of the processionary moth *Thaumetopoea pityocampa* (Denis and schiff.). *J. Chem. Ecol.*, 9: 85-93.
- Curtis**, C. E., P. J. Landolt, R. R. Heath and R. Murphy, 1989. Attraction of western grapeleaf skeletonizer males (Lepidoptera: Zygaenidae) to *S*- (+)-2-butyl-(Z)-7-tetradecenoate. *J. Econ. Entomol.*, 82: 454-457.
- Dahm**, K. H., I. Richter, D. Meyer and H. Röller, 1971. The sex attractant of the Indian-meal moth *Plodia interpunctella* (Hübner). *Life Sci.*, 10: 531-539.
- Dai**, X.-J., S.-F. Xu, M.-Z. Wang, Y.-X. Zhu, X.-H. Tang, J.-W. Zhu, J.-W. Du, T.-X. Dong and M.-Z. Du, 1988. E-10,Z-12-Hexadecadienyl acetate - sex pheromone of the mulberry white caterpillar *Rondotia menciana* Moore (Lepidoptera: Bombycidae). *Kexue Tongbao*, 33: 1575-1576.
- Daimon**, T., T. Fujii, M. Yago, Y.-F. Hsu, Y. Nakajima, T. Fujii, S. Katsuma, Y. Ishikawa and T. Shimada, 2012(a). Female sex pheromone and male behavioral responses of the bombycid moth *Triloche varians*: comparison with those of the domesticated silkworm *Bombyx mori*. *Naturwissenschaften*, 99: 207-215.

Daimon, T., T. Fujii, T. Fujii, T. Yokoyama, S. Katsuma, T. Shinoda, T. Shimada and Y. Ishikawa, 2012 (b). Reinvestigation of the sex pheromone of the wild silkmoth *Bombyx mandarina*: the effects of bombykal and bombykyl acetate. *J. Chem. Ecol.*, 38:1031-1035.

Daterman, G. E., L. J. Peterson, R. G. Robbins, L. L. Sower, G. D. Daves, Jr. and R. G. Smith, 1976. Laboratory and field bioassay of the Douglas-fir tussock moth pheromone, (Z)-6-heneicosene-11-one. *Environ. Entomol.*, 5: 1187-1190.

Daterman, G. E., R. G. Robbins, T. D. Eichlin and J. Pierce, 1977. Forest Lepidoptera attracted by known sex attractants of western spruce budworms, *Choristoneura* spp. (Lepidoptera: Tortricidae). *Can. Entomol.*, 109: 875-878.

Daterman, G. E., H. T. Cory, L. L. Sower and G. D. Daves Jr., 1984. Sex pheromone of a conifer-feeding budworm, *Choristoneura retiniana*, Walsingham. *J. Chem. Ecol.*, 10: 153-160.

David, H., B. F. Nesbitt, S. Easwaramoorthy and V. Nandagopal, 1985. Application of sex pheromones in sugarcane pest management. *Proc. Indian Acad. Sci.*, 94: 333-339.

Davis, F. M., T. G. Bird, A. E. Knutson and S.-S. Ng, 1986. Evaluation of trapping methods for southwestern corn borer (Lepidoptera: Pyralidae) using synthetic sex pheromone. *J. Econ. Entomol.*, 79: 1385-1389.

Davis, H. G. and L. M. McDonough, 1981. Sex attractant for the filbertworm, *Melisopus latiferranus* (Walsingham). *Environ. Entomol.*, 10: 390-391.

Davis, H. G., L. M. McDonough, A. K. Burditt, Jr. and B. A. Bierl-Leonhardt, 1984. Filbertworm sex pheromone. Identification and field tests of (E,E)- and (E,Z)-8,10-dodecadien-1-ol acetate. *J. Chem. Ecol.*, 10: 53-61.

Davis, H. G., L. M. McDonough, C. L. Smithhisler, D. F. Brown, B. A. Leonhardt, S. Voerman and P. S. Chapman, 1991. Mint root borer, *Fumibotys fumalis* (Guenee): identification and field tests of female sex pheromone gland components. *J. Chem. Ecol.*, 17: 249-258.

Davis, H. G., L. M. McDonough, P. S. Chapman, C. L. Smithhisler, D. D. Bowers and S. Voerman, 1993. Sugar beet crown borer, *Hulstia undulatella* (Clemens): identification and field tests of female sex pheromone gland components. *J. Chem. Ecol.*, 19: 433-440.

DeBarr, G. L., L. R. Barber, C. W. Berisford and J. C. Weatherby, 1982. Pheromone traps detect webbing coneworms in loblolly pine seed orchards. *Southern J. Appl. Forestry.*, 6: 122-127.

Decamps, C., P. du Merle, C. Gourio and G. Luquet, 1981. Attraction of species of the genus *Zygaena* by tortricid sex pheromone components (Lepidoptera, Zygaenidae and Tortricidae). *Ann. Soc. Entomol. Fr.*, 17: 441-448.

- Deland**, J.-P., R. Gries, G. Gries, G. J. R. Judd and B. D. Roitberg, 1993. Sex pheromone components of the fruit-tree leaf roller, *Archips argyrospilus* (Walker) (Lepidoptera: Tortricidae), in British Columbia. *J. Chem. Ecol.*, **19**: 2855-2864.
- Delisle**, J., 1992. Monitoring the seasonal male flight activity of *Choristoneura rosaceana* (Lepidoptera: Tortricidae) in eastern Canada using virgin females and several different pheromone blends. *Environ. Entomol.*, **21**(5): 1007-1012.
- Deng**, J., C. Lan, J. Zhou, Y. Yao, X. Yin, K. Fu, X. Ding, W. Guo, W. Liu, N. Wang and F. Wang, 2023. Analysis of sex pheromone production and field trapping of the Asian corn borer (*Ostrinia furnacalis* Guene) in Xinjiang, China. *J. Integrat. Agric.*, **22**: 1093-1103.
- Descoins**, C., E. Priesner, M. Gallois, H. Arn and G. Martin, 1978. Pheromonal secretion by virgin females of *Mamestra brassicae* L. and of *Mamestra oleracea* L. (Lepidopteras Noctuidae, Hadenniae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **286**: 77-80.
- Descoins, C., J. Einhorn, B. Lalanne-Cassou, M. Pralavorio, N. Deschamps and M. Gallois, 1985. Study of the pheromone secretion of the European carnation tortrix: *Cacoecimorpha pronubana* (Hb.), Lep. Tortricidae, Tortricinae. *Agronomie*, **5**: 19-26.
- Descoins, C., B. Lalanne-Cassou, C. Malosse and M.-L. Milat, 1986. Analysis of the sex pheromone produced by the virgin female of *Mocis latipes* (Guenée), Noctuidae, Catocalinae, from Guadeloupe (French Antilla). *C. R. Acad. Sci., Ser. 3*, **302**: 509-512.
- Descoins, C., J. F. Silvain, B. Lalanne-Cassou and H. Chéron, 1988. Monitoring of crop pests by sexual trapping of males in the French West Indies and Guyana. *Agric. Ecosystems Environ.*, **21**: 53-65.
- Descoins, C., B. Lalanne-Cassou, B. Frérot, C. Malosse and M. Renou, 1989. Comparative study of pheromonal secretions produced by arctiid and ctenuchid female moths (insects, Lepidoptera) from the neotropical area. *C. R. Acad. Sci., Ser. 3*, **309**: 577-581.
- Descoins, C., C. Malosse, M. Renou, B. Lalanne-Cassou and J. L. D. d'Aubigny, 1990. Chemical analysis of the pheromone blends produced by males and females of the neotropical moth, *Mocis megas* (Guénée) (Lepidoptera, Noctuidae, Catocalinae). *Experientia*, **46**: 536-539.
- Dix**, M. E., A. Tagesstad, J. D. Stein and M. Jacobson, 1984. Detecting tip mining Olethreutinae (Tortricidae) moths in the northern and central Great Plains with synthetic attractants. *USDA Forest Serv. Res. Note RM 445*: 5.
- Dix**, M. E. and E. W. Underhill, 1988. Sex pheromone of *Retinia metallica* (Busck) (Lepidoptera: Tortricidae): identification and field studies. *Can. Entomol.*, **120**: 721-726.

- Do**, N. D., M. Kinjo, T. Taguri, Y. Adachi, R. Yamakawa and T. Ando, 2009. Synthesis and field evaluation of methyl-branched ketones, sex pheromone components produced by Lithosiinae female moths in the family of Arctiidae. *Biosci. Biotechnol. Biochem.*, 73: 1618-1622.
- Do**, N. D., K. Ohbayashi, H. Naka, K. Nakada and T. Ando, 2011. Identification and field evaluation of sex pheromone components of the pear barkminer moth, *Spulerina astaurota*. *J. Chem. Ecol.*, 37: 1222-1230.
- Doolittle**, R. E., A. Tagestad and M. E. McKnight, 1976 (a). Trapping carpenterworms and aspen carpenterworms with sex attractants in North Dakota. *Environ. Entomol.*, 5: 267-269.
- Doolittle, R. E., W. L. Roelofs, J. D. Solomon, R. T. Cardé and M. Beroza, 1976 (b). (*Z,E*)-3,5-Tetradecadien-1-ol acetate sex attractant for the carpenterworm moth *Prionoxystus robiniae* (Peck) (Lepidoptera: Cossidae). *J. Chem. Ecol.*, 2: 399-410.
- Doolittle, R. E. and J. D. Solomon, 1986. Stereoselective synthesis of (*Z,E*)-3,5-tetradecadienyl acetate: sex attractant for carpenterworm moth, *Prionoxystus robiniae* (Peck) (Lepidoptera: Cossidae) and effect of isomers and monounsaturated acetates on its attractiveness. *J. Chem. Ecol.*, 12: 619-633.
- Downham**, M. C. A., D. R. Hall, D. J. Chamberlain, A. Cork, D. I. Farman, M. Tamo, D. Dahounto, B. Datinon and S. Adetonah, 2003. Minor components in the sex pheromone of legume podborer: *Maruca vitrata* development of an attractive blend. *J. Chem. Ecol.*, 29(4): 989-1011.
- Du**, J.-W., Y.-X. Zhu, T.-P. Zang, S.-F. Xu and X.-J. Dai, 1986. Studies on the precise blending of the pheromone components of the Asian corn borer, *Ostrinia furnacalis* Guenée (Lep. Pyralidae). *Contr. Shanghai Inst. Entomol.*, 6: 17-22.
- Du, J.-W., C. Löfstedt and J. Löfqvist, 1987 (a). Repeatability of pheromone emissions from individual female ermine moths *Yponomeuta padellus* and *Yponomeuta rorellus*. *J. Chem. Ecol.*, 13: 1431-1441.
- Du, J., X. Dai, S. Xu, X. Tang, T. Zhang, Y. Zhu and M. Wang, 1987 (b). Studies on the sex pheromone of the yellow rice borer, *Scirpophaga incertulas* (Walker), Pyralidae, Lepidoptera: identification of chemical structure and field trapping tests. *Sci. Sin., Ser. B*, 30: 967-973.
- Du**, T., J. Xiong, Z. Wang and F. Kong, 1989. (*7Z,11Z*)-7,11-Hexadecadienal: sex attractant of *Phyllocnistis wampella* Liu et Zeng. *Kunchong Zhishi*, 26: 147-149. [Chem. Abst., 112: 50570n]
- Du**, Y., P. Li, Z. Chen, Y. Lin, Y. Wang and Y. Qin, 2013. Field trapping of male *Phyllonorycter ringoniella* using variable ratios of pheromone components. *Entomol. Exp. Appl.*, 146: 357-363.

- Duckworth**, W. D. and T. D. Eichlin, 1977 (a). A new species of clearwing moth from southcentral Texas. *Pan-Pac. Entomol.*, **53**: 175-178.
- Duckworth, W. D. and T. D. Eichlin, 1977 (b). Two new species of clearwing moths (Sesiidae) from eastern North America clarified by sex pheromones. *J. Lepi. Soc.*, **31**: 191-196.
- Duff**, C. M., G. Gries, K. Mori, Y. Shirai, M. Seki, H. Takikawa, T. Sheng, K. N. Slessor, R. Gries, C. T. Maier and D. C. Ferguson, 2001. Dose pheromone biology of *Lambdina athasaria* and *L. pellucidaria* contribute to their reproductive isolation? *J. Chem. Ecol.*, **27** (3): 431-442.
- Dung**, T. T., D. K. Hanh, T. P. Linh, D. T. Chi, T. T. Thy, L. V. Vang and T. V. Phen, 2021. Preliminary identification of the sex pheromones of the citrus fruit borer (*Citripestis sagitiferella* Moore) inhabiting the Mekong Delta of Vietnam. *Int. J. Plant. Anim. Environ. Sci.*, **11**: 194-205.
- Dunkelblum**, E., S. Gothilf and M. Kehat, 1980. Identification of the sex pheromone of the cotton bollworm, *Heliothis armigera*, in Israel. *Phytoparasitica*, **8**: 209-211.
- Dunkelblum, E., S. Gothilf and M. Kehat, 1981. Sex pheromone of the tomato looper, *Plusia chalcites* (Esp.). *J. Chem. Ecol.*, **7**: 1081-1088.
- Dunkelblum, E., M. Kehat, S. Gothilf, S. Greenberg and B. Sklarsz, 1982. Optimized mixture of sex pheromonal components for trapping of male *Spodoptera littoralis* in Israel. *Phytoparasitica*, **10**: 21-26.
- Dunkelblum, E. and S. Gothilf, 1983. Sex pheromone components of the gamma moth, *Autographa gamma* (L.) (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **38c**: 1011-1014.
- Dunkelblum, E., R. Snir, S. Gothilf and I. Harpaz, 1987 (a). Identification of sex pheromone components from pheromone gland volatiles of the tomato looper, *Plusia chalcites* (Esp.). *J. Chem. Ecol.*, **13**: 991-1003.
- Dunkelblum, E., M. Kehat, M. Harel and D. Gordon, 1987 (b). Sexual behaviour and pheromone titer of the *Spodoptera littoralis* female moth. *Entomol. Exp. Appl.*, **44**: 241-247.
- Dunkelblum, E. and M. Kehat, 1989. Female sex pheromone components of *Heliothis peltigera* (Lepidoptera: Noctuidae). Chemical identification from gland extracts and male response. *J. Chem. Ecol.*, **15**: 2233-2245.
- Dunkelblum, E. and M. Mazor, 1993. Chemical characterization and species specificity of sex pheromones of Plusiinae moths in Israel. *Arch. Insect Biochem. Physiol.*, **22**: 413-424.
- Efetov**, K. A., F. Can, T. B. Toshova and M. Subchev, 2010. New sex attractant for *Jordanita anatolica* (Naufock) (Lepidoptera: Zygaenidae: Procridinae). *Acta Zool. Bulg.*, **62**: 315-319.

- Efetov, K. A., E. V. Parshkova, M.Y. Baevsky and A. I. Poddubov, 2014. Sec-butyl ester of dodecenoate: synthesis and attractive properties. *Ukr. Biochem. J.*, **86**: 175-182.
- Efetov, K. A., E. E. Kucherenko, E. V. Parshkova and G. M. Tarmann, 2016. 2-Butyl 2-dodecenoate, a new sex attractant for *Jordanita (Tremewania) notata* (Zeller, 1847) and some other Procridinae species (Lepidoptera: Zygaenidae). *SHILAP Rev. Lepid.*, **44**: 519-527.
- Efetov, K. A., E. E. Kucherenko and G. M. Tarmann, 2019. New synthetic sex attractants for the males of two endemic Iberian Procridinae species (Lepidoptera: Zygaenidae). *SHILAP Rev. Lepid.*, **47**: 307-315.
- Efetov, K. A. and E. E. Kucherenko, 2020. Structural analysis of sex pheromones and attractants in Zygaenidae (Insecta, Lepidoptera): biochemical and evolutionary aspects. *J. Evol. Biochem. Physiol.*, **56**: 381-395.
- Efetov, K. A. and E. E. Kucherenko, 2021. Enantiomers of 2-butyl dodec-2-enoate as sex attractants. *Rus. J. Bioorg. Chem.*, **47**: 1043-1050.
- Eichlin, T. D. and S. Passoa, 1983. A new clearwing moth (Sesiidae) from Central America: a stem borer in *Mimosa pigra*. *J. Lepid. Soc.*, **37**: 193-206.
- Einhorn, J., J.-Y. Lallemand, P. Zagatti, M. Gallois, H. Virelizier, J. Riom and P. Menassieu, 1982. Isolation and identification of the sex pheromone blend of *Hyphantria cunea* (Drury) (Lepidoptera, Arctiidae). *C. R. Acad. Sci., Ser. 2*, **294**: 41-44.
- Einhorn, J., P. Menassier, D. Michelot and J. Riom, 1983. Sex trapping of the processionary moth *Thaumetopoea pityocampa* Schiff. (Lep., Notodontidae) with synthetic attractants. First experiments in the south-west of France. *Agronomie*, **3**: 499-505.
- Einhorn, J., B. Boniface, M. Renou and M.-L. Milat, 1984 (a). Study on the sex pheromone of *Arctia villica* L. (Lepidoptera, Arctiidae). *C. R. Acad. Sci., Ser. 3*, **298**: 573-576.
- Einhorn, J., F. Beauvais, M. Gallois, C. Descoins and R. Causse, 1984 (b). Secondary components of the codling moth (*Cydia pomonella* L.) (Lepidoptera, Tortricidae) sex pheromone. *C. R. Acad. Sci., Ser. 3*, **299**: 773-778.
- Einhorn, J., P. Witzgall, H. audemard, B. Boniface and R. Causse, 1986. Secondary components of the codling moth (*Cydia pomonella* L.) (Lepidoptera, Tortricidae) sex pheromone. II. First results on the behavioural effects. *C. R. Acad. Sci., Ser. 3*, **302**: 263-266.

Eiras, A. E., A. Kovaleski, E. F. Vilela, J. P. Champon, C. R. Unelius, A.-K. Borg-Karlsson, I. Liblikas, R. Mozūraitis, M. Bengtsson and P. Witzgall, 1999. Sexpheromone of the Brazilian apple leafroller, *Bonagota cranaodes* Meyrick (Lepidoptera, Tortricidae). *Z. Naturforsch.*, **54c**: 595-601.

El-Sayed, A., J. Gödde, P. Witzgall and H. Arn, 1999. Characterization of pheromone blend for grapevine moth, *Lobesia botrana* by using flight track recording. *J. Chem. Ecol.*, **25**(2): 389-400.

El-Sayed, A. M., H. M. Fraser and R. M. Trimble, 2001. Chemical identification and behavioral activity of (*Z*)-11-tetradecenal in an eastern North American population of the obliquebanded leafroller. *Can. Entomol.*, **133**: 365-374.

El-Sayed, A. M., J. Delisle, N. De Lury, L. J. Gut, G. J. R. Judd, S. Legrand, W. H. Reissig, W. L. Roelofs, C. R. Unelius and R. M. Trimble, 2003. Geographic variation in pheromone chemistry, antennal electrophysiology, and pheromone-mediated trap catch of North American populations of the obliquebanded leafroller. *Environ. Entomol.*, **32**: 470-476.

El-Sayed, A. M., A. R. Gibb, D. M. Suckling, B. Bunn, S. Fielder, D. Comeskey, K. A. Moss, L. A. Manning, S. P. Foster, B. D. Morris, T. Ando and K. Mori, 2004(a). Identification and synthesis of the sex pheromone of painted apple moth. *HortResearch Client Report* No: 13463, Contact No: 18879.

El-Sayed, A. M., L. I. Wainman, E. M. Santangelo, C. R. Unelius and R. M. Trimble, 2004(b). Relative attractiveness of (10*E*)-dodecen-1-yl acetate and (4*E*,10*E*)-dodecadien-1-yl acetate to male spotted tentiform leafminers *Phyllonorycter blancardella* (F.). *J. Chem. Ecol.* **30**: 1827-1838.

El-Sayed, A. M., A. R. Gibb, D. M. Suckling, B. Bunn, S. Fielder, D. Comeskey, L. A. Manning, S. P. Foster, B. D. Morris, T. Ando and K. Mori, 2005(a). Identification of sex pheromone components of the painted apple moth: a tussock moth with a thermally labile pheromone component. *J. Chem. Ecol.*, **31**(3): 621-646.

El-Sayed, A. M., L. I. Wainman, E. M. Santangelo, C. R. Unelius and R. M. Trimble, 2005(b). Sex pheromone of a North American population of the spotted tentiform leafminer, *Phyllonorycter blancardella*. *Entomol. Exp. Appl.*, **116**: 143-148.

El-Sayed, A. M., A. R. Gibb and D. M. Suckling, 2005(c). Chemistry of the sex pheromone gland of the fall webworm, *Hyphantria cunea*, discovered in New Zealand. *New Zealand Plant Protection*, **58**: 31-36.

El-Sayed, A. M., V. J. Mitchell, L.-A. M. Manning and D. M. Suckling, 2011. New sex pheromone blend for the lightbrown apple moth, *Epiphyas postvittana*. *J. Chem. Ecol.*, **37**: 640-646.

El-Sayed, A. M., A. R. Gibb, V. J. Mitchell, L. M. Manning, J. Revell,

B. Thistleton and D. M. Suckling, 2013. Identification of the sex pheromone of *Conogethes pluto*: a pest of Alpinia. *Chemoecology*, 23: 93-101.

El-Sayed, A. M. and L.-A. M. Manning, 2022. Identification of the sex pheromone of the pink grass worm, *Tmetolophota atristriga*, reveals possible population differences in male response to sex pheromone. *J. Chem. Ecol.*, 48: 683-689.

Evenden, M. L., J. H. Borden, G. A. Van Sickle and G. Gries, 1995. Development of a pheromone-based monitoring system for western hemlock looper (Lepidoptera: Geometridae): effect of pheromone dose, lure age, and trap type. *Environ. Entomol.*, 24(4): 923-932.

Evenden, M. L. and R. Gries, 2006. Sex pheromone of the large aspen tortrix, *Choristoneura conflictana* (Lepidoptera: Tortricidae). *Chemoecology* 16: 115-122.

Evenden, M.L., B. A. Mori, R. Gries and J. Otani, 2010. Sex pheromone of the red clover casebearer moth, *Coleophora deauratella*, an invasive pest of clover in Canada. *Entomol. Exp. Appl.*, 137: 255-261.

Fang, Y.-L., J.-H. Sun, C.-H. Zhao and Z.-N. Zhang, 2005. Sex pheromone components of the sandthorn carpenterworm, *Holococerus hippophaecolus*. *J. Chem. Ecol.* 31: 39-48.

Farine, J.-P., B. Frérot and J. Isart, 1981. Chemical isolation factor in pheromonal secretion of two noctuids (Hadeninae): *Mamestra brassicae* (L.) and *Pseudaletia unipuncta* (Haw.). *C. R. Acad. Sci., Ser. 3*, 292: 101-104.

Félix, A.-E., G. Genestier, C. Malosse, P.-A. Calatayud, B. L. Rü, J.-F. Silvain and B. Frérot, 2009. Variability in pheromone communication among different haplotype populations of *Busseola fusca*. *J. Chem. Ecol.*, 35: 618-623.

Félix, A.-E., P.-A. Calatayud, B. Le Ru, J.-F. Silvain and B. Frérot, 2011. Sex pheromone composition and reproductive isolation in two *Busseola* species (Lepidoptera: Noctuidae) in Kenya. *Chemoecology*, 21: 107-111.

Ferrao, P., G. Gries, P. D. C. Wimalaratne, C. T. Maier, R. Gries, K.N. Slessor and J. Li 1998. Sex pheromone of apple blotch leafminer, *Phyllonorycter crataegella*, and its effect on *P. mespiliella* pheromone communication. *J. Chem. Ecol.*, 24(12): 2059-2078.

Ferrara, F. A. A., E. F. Vilela, G. N. Jham, A. E. Eiras, M. C. Picanco, A. B. Attygalle, A. Svatos, R. T. S. Frighetto and J. Meinwald, 2001. Evaluation of the synthetic major component of the sex hormone of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, 27(5): 907-917.

Filho, M. M., E. F. Vilela, A. B. Attygalle, J. Meinwald, A. Svatos

and G. N. Jham, 2000. Field trapping of tomato moth, *Tuda absoluta* with pheromone traps. *J. Chem. Ecol.*, 26(4): 875-881.

Fitzpatrick, S. M. and J. T. Troubridge, 1992. Relative efficacies of two commercial pheromone blends for monitoring the blackheaded fireworm (Lepidoptera: Tortricidae) of Cranberries. *J. Econ. Entomol.*, 85(3): 947-949.

Fleischer, S. J., C. L. Harding, P. E. Blom, J. White and J. Grehan, 2005. *Spodoptera frugiperda* pheromone lures to avoid nontarget captures of *Leucania phragmatidicola*. *J. Econ. Entomol.*, 98: 66-71.

Flint, H. M., M. Balasubramanian, J. Campero, G. R. Strickland, Z. Ahmad, J. Barral, S. Barbosa and A. F. Khail, 1979. Pink bollworm: response of native males to ratios of Z,Z-and Z,E-isomers of gossyplure in several cotton growing areas of the world. *J. Econ. Entomol.*, 72: 758-762.

Foster, S. P., J. R. Clearwater, S. J. Mugglestone, J. S. Dugdale and W. L. Roelofs, 1986. Probable sibling species complexes within two described New Zealand leafroller moths. *Naturwissenschaften*, 73: 156-158.

Foster, S. P. and W. L. Roelofs, 1987(a). Sex pheromone differences in populations of the brownheaded leafroller, *Ctenopseustis obliquana*. *J. Chem. Ecol.*, 13: 623-629.

Foster, S. P., J. R. Clearwater and W. L. Roelofs, 1987(b). Sex pheromone of *Planotortrix* species found on mangrove. *J. Chem. Ecol.*, 13: 631-637.

Foster, S. P. and J. S. Dugdale, 1988. A comparison of morphological and sex pheromone differences in some New Zealand Tortricinae moths. *Biochem. System. Ecol.*, 16: 227-232.

Foster, S. P., J. R. Clearwater and S. J. Mufflestone, 1989. Intraspecific variation of two components in sex pheromone gland of *Planotortrix excessana* sibling species. *J. Chem. Ecol.*, 15: 457-465.

Foster, S. P., J. R. Clearwater, S. J. Mufflestone and P. W. Shaw, 1990. Sex pheromone of a *Planotortrix excessana* sibling species and reinvestigation of related species. *J. Chem. Ecol.*, 16: 2461-2474.

Foster, S. P., J. S. Dugdale and C. S. White, 1991. Sex pheromones and the status of greenheaded and brownheaded leafroller moths in New Zealand. *New Zealand J. Zool.*, 18: 63-74.

Foster, S. P., D. Steven, C. McKenna and S. J. Mufflestone, 1993. Sex pheromones of two New Zealand tortricid moths, *Cnephacia jactatana* and *Merophyas leucaniana*. *New. Zeal. J. Zool.*, 20: 81-88.

Foster, S. P. and W. P. Thomas, 2000. Identification of a sex pheromone component of the raspberry budmoth, *Heterocrossa rubophaga*. *J. Chem. Ecol.*, 26(11): 2549-2555.

- Foster, S. P., B. D. Morris, L. D. Charlet, T. Gross and S. Grugel, 2003. Identification of the sex pheromone of *Cochylis arthuri* (Lepidoptera: Cochylidae). *Can. Entomol.*, **135**: 713-719.
- Fouda**, H. G., J. N. Seiber and O. G. Bacon, 1975. A potent sex attractant for the potato tuberworm moth. *J. Econ. Entomol.*, **68**: 423-427.
- Francke**, W., S. Franke, M. Toth, G. Szöcs, P. Guerin and H. Arn, 1987. Identification of 5,9-dimethylheptadecane as a sex pheromone of the moth *Leucoptera scitella*. *Naturwissenschaften*, **74**: 143-144.
- Francke, W., M. Toth, G. Szöcs, W. Krieg, H. Ernst and E. Buschmann, 1988. Identification and synthesis of dimethylalkanes as sex attractants of female leaf miner moths (Lyonetiidae). *Z. Naturforsch.*, **43c**: 787-789.
- Francke, W., E. Plass, N. Zimmermann, H. Tietgen, T. Tolasch, S. Franke, M. Subchev, T. Toshova, J. A. Pickett, L. J. Wadhams and C. M. Woodcock, 2000. Major sex pheromone component of female herald moth *Scoliopteryx libatrix* is the novel branched alkene (6Z,13)-methylheneicosene. *J. Chem. Ecol.*, **26**(5): 1135-1149.
- Francke, W., S. Franke, J. Bergmann, T. Tolasch, M. Subchev, A. Mircheva, T. Toshova, A. Svatos, B. Kalinová, Z. Karpati, G. Szöcs and M. Tóth, 2002. Female sex pheromone of *Cameraria ohridella* Desch. and Dim. (Lepidoptera: Gracillariidae): structure confirmation, synthesis and biological activity of (8E,10Z)-8,10-tetradecadienal and some analogues. *Z. Naturforsch.*, **57c**: 739-752.
- Francke, W., V. Karalius, E. Plass, L. Lehmann, A. D. Santos, V. Buda, A.-K. Borg-Karlson and R. Mozūraitis, 2004. New type of Sesiidae sex pheromone identified from the hornet moth *Sesia apiformis*. *J. Chem. Ecol.*, **30**(4): 805-817.
- Frerot**, B., E. Priesner and M. Gallois, 1979(a). A sex attractant for the green budworm moth, *Hedya nubiferana*. *Z. Naturforsch.*, **34c**: 1248-1252.
- Frerot, B., C. Descoins, B. Lalanne-Cassou, P. Saglio and F. Beauvais, 1979(b). Field trapping of tortricids from apple orchards with synthetic sex attractants. *Ann. Zool. Ecol. Anim.*, **11**: 617-636.
- Frerot, B., B. Boniface, J.-P. Chambon and Y. Meritan, 1982(a). Trapping with synthetic sex attractants to determine the distribution of three tortricid moths of apple orchards in the Paris area. *Agronomie*, **2**: 885-893.
- Frerot, B., M. Gallois, M. Lettere, J. Einhorn, D. Michelot and C. Descoins, 1982(b). Sex pheromone of *Pandemis heparana* (Den. and Schiff.) (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **8**: 663-670.
- Frerot, B., M. Renou, M. Gallois and C. Descoins, 1983. A sex attractant for *Archips xylosteana* L. (Lepid., Tortricidae, Tortricinae). *Agronomie*, **3**: 173-178.

- Frerot, B., M. Renou, C. Descoins and M. Gallois, 1984. A sex pheromone for *Argyrotaenia pulchellana* (Haw.) (Lepidoptera, Tortricinae). *Agronomie*, 4: 565-572.
- Frerot, B., J.-P. Chambon, B. Boniface and C. Descoins, 1985. Sexual trapping of various Lepidoptera males by synthetic attractants. *Ann. Soc. Entomol. Fr.*, 21: 297-305.
- Frérot, B., C. Malosse, M.-L. Milat, J.-Y. Lallemand, J. Soulié and A. Brunetiére, 1986. Isolation and identification of three constituents of the sexual pheromone of *Zeuzera pyrina* L. *C. R. Acad. Sci., Ser. 2*, 302: 413-415.
- Frérot, B., J.-P. Chambon and C. Cocquempot, 1987(a). Some results on sexual trapping of Lepidoptera Noctuidae and Geometridae. *Ann. Soc. Entomol. Fr.*, 23: 151-154.
- Frerot, B., M. Gallois, C. Malosse and R. Bues, 1987 (b). Identification of the sex pheromone of *Mamestra biren* (Goeze) ex *M. glauca* (Hb.). *C. R. Acad. Sci., Ser. 3*, 304: 139-142.
- Frérot, B., M. Renou, C. Malosse and C. Descoins, 1988 (a). Isolation and identification of pheromone compounds in female moths of *Tyria jacobaeae* (Lepidoptera, Arctiidae). Biological characterization of the absolute configuration of the main component. *Entomol. Exp. Appl.*, 46: 281-289.
- Frerot, B., J.-R. Pougny, M.-L. Milat, P. Rollin and C. Malosse, 1988 (b). Study of the pheromonal secretion of the tiger moth, *Cymbalophora pudica* (Esper) and enantioselectivity of its pheromonal perception. *C. R. Acad. Sci., Ser. 3*, 306: 157-160.
- Frerot, B., B. Lalanne-Cassou, M. Gallois, C. Malosse, A.-H. Cain and R. Bues, 1988 (c). Physico-chemical analysis of the sex pheromone blend produced by the Hadeninae noctuid moth: *Mamestra suasa* (D. et S.). *C. R. Acad. Sci., Ser. 3*, 307: 785-788.
- Frérot, B., C. Malosse, M. L. Milat, G. Demolin, J. C. Martin, M. Khemici, M. Zamoun and M. Gachi, 1990 (a). Chemical analysis of the sex pheromone glands of *Thaumetopoea bonjeani* (Powell) (Lep., Thaumetopoeidae). *J. Appl. Entomol.*, 109: 210-212.
- Frérot, B., J. P. Chambon, C. Malosse and G. Genestier, 1990 (b). A sex attractant for *Lobesia bicinctana* (Dup.) (Lep., Tortricidae). *J. Appl. Entomol.*, 110: 365-369.
- Frérot, B. and S. P. Foster, 1991. Sex pheromone evidence for two distinct taxa within *Graphania mutans* (Walker). *J. Chem. Ecol.*, 17: 2077-2093.
- Frérot, B., C. Malosse, R. D. Chenon, P. H. Ducrot and A.-H. Cain, 1993 (a). Identification of the sex pheromone components in *Pareuchaetes pseudoinsulata* Rego Barros (Lepidoptera, Arctiidae). *C. R. Acad. Sci., Ser. 2*, 317: 1045-1050.
- Frérot, B., J. S. Dugdale and S. P. Foster, 1993 (b). Chemotaxonomy of some species of moths in the New Zealand genus *Graphania*.

based on sex pheromones. *New Zealand J. Zool.*, **20**: 71-80.

Frerot, B. and G. Demolin, 1993 (c). Sex pheromone of the processionary moths and biosystematic considerations within the genus *Thaumetopoea* (*Thaumetopoeidae Thaumetopoeinae*). *Boll. Zool. Agric. Bachic. Ser. II*, **25**(1): 33-40.

Frérot, B., J.-P. Marro and C. Malosse, 1995. In vitro incubation of sex pheromone gland and identification of pheromone components in *Cydia splendana* (Hb). *C. R. Acad. Sci., Life Sciences*, **318**; 447-451.

Frilli, F., 1974. Observations on the use of *Cydia molesta* pheromones. *Entomologica*, **10**: 31-56.

Fu, X.-Y. X.-Q. Pan and X.-Z. Meng, 2002. Female sex pheromone of the Chinese tortrix *Cydia trasias*. *Entomol. Exp. Appl.*, **103**: 183-186.

Fu, X., J. Tabata, Y. Huang, T. Takanashi, S. Ohno, H. Honda, S. Tatsuki and Y. Ishikawa, 2004. Female sex pheromone of *Ostrinia orientalis* - throwing a light on the relationship between *O. orientalis* and the European corn borer, *O. nubilalis*. *Chemoecology*, **14**: 175-180.

Fu, N., F. H. Magsi, Y. Zhao, X. Cai, Z. Li, L. Bian, C. Xiu, Z. Chen and Z. Luo, 2022. Identification and field evaluation of sex pheromone components and its antagonist produced by a major tea pest, *Archips strojny* (Lepidoptera: Tortricidae). *Insects*, **13**: 1056.

Fujii, T., D. Iwai, F. Mochizuki and K. Fukumoto, 2010 (a). Sex pheromones and sex attractants from *Platyptilia ignifera*. *Jpn. Kokai Tokkyo Koho* JP 2010018543. [SciFinder. AN 2010:110975]

Fujii, T., R. Nakano, Y. Takubo, S. Qian, R. Yamakawa, T. Ando and Y. Ishikawa, 2010 (b). Female sex pheromone of a lichen moth *Eilema japonica* (Arctiidae, Lithosiinae). *J. Insect Physiol.*, **56**: 1986-1991.

Fujii, T., H. Miyake and Y. Nasu, 2013 (a). Sex attractants of *Carposina coreana*. *Jpn. Kokai Tokkyo Koho* JP 2013241377.

Fujii, T., R. Yamakawa, Y. Terashima, S. Imura, K. Ishigaki, M. Kinjo and T. Ando, 2013 (b). Propionates and acetates of chiral secondary alcohols, novel sex pheromone components produced by a lichen moth *Barsine expressa* (Arctiidae: Lithosiinae). *J. Chem. Ecol.*, **39**: 28-36.

Gago, R., J. D. Allison, J. S. McElfresh, K. F. Haynes, J. McKenney, A. Guerrero and J. G. Millar, 2013. A tetraene aldehyde as the major sex pheromone component of the promethea moth (*Callosamia promethea* (Drury)). *J. Chem. Ecol.*, **39**: 1263-1272.

Galbreath, R. A., M. H. Benn, H. Young, V. A. Holt, 1985. Sex pheromone components in the New Zealand greenheaded leafroller *Planotortrix excessana* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **40 c**: 266-271.

Gemeno, C. and K. F. Haynes 1998. Chemical and behavioral evidence for a third pheromone component in a North American population of the black cutworm moth, *Agrotis ipsilon*. *J. Chem. Ecol.*, **24**(6): 999-1011.

Gemeno, C., A. F. Lutfallah and K. F. Haynes, 2000. Pheromone blend variation and cross-attraction among populations of the black cutworm moth (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **93**(6): 1322-1328.

Gentry, C. R., M. Beroza and J. L. Blythe, 1975. Pecan bud moth: captures in Georgia in traps baited with the pheromone of the oriental fruit moth. *Environ. Entomol.*, **4**: 227-228.

Germinara, G. S., A. Elgargoti, A. D. Cristofaro and G. Rotundo, 2007. Female sex pheromone of *Sesamia cretica*: chemical and behavioural evidence for a three-component blend. *Entomol. Exp. Appl.*, **124** (2): 213-219.

Ghizdavu, I., F. Hodosan, N. Popovici, A. Botar and A. Barabas, 1979. A sex attractant for *Autographa (Plusia) gamma* L. moth. *Rev. Roum. Biol., Ser. Biol. Anim.*, **24**: 87-90.

Ghizdavu, I., F. P. Hodosan, I. Oprean, A. Gocan, H. Ciupe and S. Matic, 1985. Sex attractants for five species of Noctuidae determined using traps. *Rev. Roum. Biol., Ser. Biol. Anim.*, **30**: 25-29.

Ghizdavu, I., F. P. Hodosan and I. Oprean, 1987. Attractifs sexuels spécifiques pour *Adoxophyes orana* F. v. R. et *Archips crataegana* Hb. *Rev. Roum. Biol., Ser. Biol. Anim.*, **32**: 23-27.

Gibb, A. R., L. E. Jamieson, D. M. Suckling, P. Ramankutty and P. S. Stevens, 2005. Sex pheromone of the citrus flower moth *Prays nephelomina*: pheromone identification, field trapping trials, and phenology. *J. Chem. Ecol.*, **31**(7): 1633-1644.

Gibb, A. R., D. M. Suckling, B. D. Morris, T. E. Dawson, B. Bunn, D. Comeskey and J. J. Dymock, 2006(a). (Z)-7-Tricosene and monounsaturated ketones as sex pheromone components of the Australian guava moth, *Coscinoptyla improbana*: identification, field trapping, and phenology. *J. Chem. Ecol.*, **32**(1): 221-237.

Gibb, A. R., D. Comeskey, L. Berndt, E. G. Brockerhoff, A. M. El-Sayed, H. Jactel and D. M. Suckling, 2006(b). Identification of sex pheromone components of a New Zealand geometrid moth, the common forest looper *Pseudocoremia suavis*, reveals a possible species complex. *J. Chem. Ecol.*, **32**(4): 865-879.

Gibb, A. R., B. Pineese, D. Tenakanai, A. P. Kawi, B. Bunn, P. Ramankutty and D. M. Suckling, 2007(a). (Z)-11-Hexadecenal and (3Z,6Z,9Z)-tricosatriene: sex pheromone components of the red banded mango caterpillar *Deanolis sublimbalis*. *J. Chem. Ecol.*, **33** (3): 579-589.

Gibb, A. R., D. M. Suckling, A. M. El-Sayed, B. Bohman, C. R. Unelius, J. J. Dymock, M. L. Larsen and B. E. Willoughby, 2007(b).

(11Z,13E)-Hexadecadien-1-yl acetate: sex pheromone of the grass webworm *Herpetogramma licarsalis* - Identification, synthesis, and field bioassays. *J. Chem. Ecol.*, **33** (4): 839-847.

Gibb, A. R., D. M. Suckling, S. Fielder, B. Bunn, L. E. Jamieson, M. L. Larsen, G. H. Walter and D. J. Kriticos, 2008. Major sex pheromone components of the Australian gum leaf skeletonizer *Uraba lugens*: (10E,12Z)-hexadecadien-1-yl acetate and (10E,12Z)-hexadecadien-1-ol. *J. Chem. Ecol.*, **34** (9): 1125-1133.

Gigon, F., 1980. Utilisation des piéges à phéromone sexuelle synthétique dans l'étude faunistique des tordeuses (Tortricidae s. str.) de vergers neuchatelois. *Bulletin de la Société Neuchâteloise des Sciences Naturelles*, **103**: 45-56.

Gillespie, D. R., G. G. S. King, V. A. Salas-Reyes and K. N. Slessor, 1984. Determination of the sex-pheromone components of *Cheimophila salicella* (Lepidoptera: Oecophoridae), a pest of blueberry in British Columbia. *Can. Entomol.*, **116**: 1397-1402.

Gothilf, S., M. Kehat, M. Jasobcon and R. Galun, 1978. Sex attractants for male *Heliothis armigera* (Hbn.). *Experientia*, **34**: 853-854.

Granges, J. and M. Baggolini, 1971. Une phéromone sexuelle synthétique attractive pour le carpocapse des prunes (*Grapholitha funebrana* Tr. [Lép. Tortricidae]). *Rev. Suisse Vitic. Arboric.*, **3**: 93-94.

Grant, G. G., 1977. Interspecific pheromone responses of tussock moths and some isolating mechanisms of eastern species. *Environ. Entomol.*, **6**: 739-742.

Grant, G. G. and K. N. Slessor, 1983. Sex pheromone of the maple leaf roller *Cenopis acerivora* (Lepidoptera: Tortricidae). *Can. Entomol.*, **115**: 189-194.

Grant, G. G., L. MacDonald, D. Frech, K. Hall and K. N. Slessor, 1985. Sex attractants for some eastern species of *Rhyacionia*, including a new species, and *Eucosma gloriola* (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 1489-1496.

Grant, G. G., Y. H. Prévost, K. N. Slessor, G. G. S. King and R. J. West, 1987(a). Identification of the sex pheromone of the spruce coneworm, *Dioryctria reniculelloides* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **16**: 905-909.

Grant, G. G., D. Frech, L. MacDonald, K. N. Slessor and G. G. S. King, 1987(b). Copulation releaser pheromone in body scales of female whitemarked tussock moth, *Orgyia leucostigma* (Lepidoptera: Lymantriidae): identification and behavioral role. *J. Chem. Ecol.*, **13**: 345-356.

Grant, G. G., W. H. Fogal, R. J. West, K. N. Slessor and G. E. Miller, 1989. A sex attractant for the spruce seedmoth, *Cydia strobilella* (L.), and the effect of lure dosage and trap height on capture of male moths. *Can. Entomol.*, **121**: 691-697.

- Grant G. G., B. Pendrel, K. M. Slessor, X. Z. Meng and W. E. Miller, 1991. Identification of sex pheromone components for two lepidopteran defoliators, the oak olethreutid leafroller, *Pseudexentera spoliana* (Clemens), and the aspen leafroller, *Pseudexentera oregonana* (Walsingham). *Can. Entomol.*, 123: 1209-1218.
- Grant, G. G., S. A. Katovich, D. J. Hall, D. A. Lombardo and K. N. S. A. Lessor, 1993. Sex pheromone identification and trapping of *Dioryctria resinosella* (Lepidoptera: Pyralidae). *Environ. Entomol.*, 22(1): 154-161.
- Grant, G. G., D. Langevin, J. Liska, P. Kapitola and J. M. Chong, 1996. Olefin inhibitor of gypsy moth, *Lymantria dispar*, is a synergistic pheromone component of nun moth, *L. monacha*. *Naturwissenschaften*, 83: 328-330.
- Grant, G. G., P. D. Groot, D. Langevin, S. A. Katovich, K. N. Slessor and W. E. Miller, 2002. Sex attractants and seasonal flight patterns for three *Eucosma* (Lepidoptera: Tortricidae) species sympatric in eastern pine seed orchards and plantations. *Can. Entomol.*, 134: 391-401.
- Grant, G. G., K. N. Slessor, Weiliu and M. M. Abou-Zaid, 2003. (Z,Z)-6,9-Heneicosadien-11-one, labile sex pheromone of the whitemarked tussock moth, *Orgyia leucostigma*. *J. Chem. Ecol.*, 29(3): 589-601.
- Grant, G. G., M. D. Coppens, L. K. Hartling, D. O'Shea, D. Winter, J. Gordon, J. Rudderham and W. Liu, 2008. Operational monitoring of the whitemarked tussock moth with Multipher traps generating the unstable pheromone component (Z,Z)-6,9-heneicosadien-11-one from a precursor. *Entomol. Exp. Appl.*, 126: 174-178.
- Gray**, T. G., K. N. Slessor, G. G. Grant, R. F. Shepherd, E. H. Holsten and A. S. Tracey, 1984(a). Identification and field testing of pheromone components of *Choristoneura orae* (Lepidoptera: Tortricidae). *Can. Entomol.*, 116: 51-56.
- Gray, T. G., K. N. Slessor, R. F. Shepherd, G. G. Grant and J. F. Manville, 1984(b). European pine shoot moth, *Rhyacionia buoliana* (Lepidoptera: Tortricidae): identification of additional pheromone components resulting in an improved lure. *Can. Entomol.*, 116: 1525-1532.
- Gray, T. G. and G. Gries, 1993. Sex pheromone components of an undescribed *Choristoneura* species (Lepidoptera: Tortricidae) on lodgepole pine in British Columbia. *J. Entomol. Soc. Brit. Columb.*, 90: 13-18.
- Gray, T. G., R. F. Shepherd, G. Gries and R. Gries, 1996. Sex pheromone component of the western blackheaded budworm, *Acleris gloverana* Walsingham (Lepidoptera: Tortricidae). *Can. Entomol.*, 128: 1135-1142.

Green, N., M. Jacobson and J. C. Keller, 1969. Hexalure, an insect sex attractant discovered by empirical screening. *Experientia*, 25: 682-683.

Greenfield, M. D. and M. G. Karandinos, 1979. Resource partitioning of the sex communication channel in clearwing moths (Lepidoptera: Sesiidae) of Wisconsin. *Ecol. Monogr.*, 49: 403-426.

Greenway, A. R., 1984. Sex pheromone of the pea moth, *Cydia nigricana* (F.) (Lepidoptera: Olethreutidae). *J. Chem. Ecol.*, 10: 973-982.

Griepink, F. C., T. A. Beek, J. H. Visser, S. Voerman and A. Groot, 1995. Isolation and identification of sex pheromone of *Symmetrischema tangolias* (Gyen) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, 21: 2003-2013.

Griepink, F. C., T. A. Beek, M. A. Posthumus, A. Groot, J. H. Visser and S. Voerman, 1996. Identification of the sex pheromone of *Scrobipalpula absoluta*; determination of double bond positions in triple unsaturated straight chain molecules by means of dimethyl disulphide derivatization. *Tetrahedron Lett.*, 37; 411-414.

Gries, G., R. Gries, J. H. Borden, J.-X. Li, K. N. Slessor, G. G. S. King, W. W. Bowers, R. J. West and E. W. Underhill, 1991. 5,11-Dimethyl-heptadecane and 2,5-dimethylheptadecane: sex pheromone components of the geometrid moth, *Lambdina fiscellaria fiscellaria*. *Naturwissenschaften*, 78: 315-317.

Gries, G., R. Gries, E. W. Underhill and L. Humble, 1993 (a). (6Z,9Z-3R,4S)-Epoxy-heptadecadiene: major sex pheromone component of the larch looper, *Semiothisa sexmaculata* (Packard) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 19: 843-850.

Gries, G., R. Gries, S. H. Krannitz, J. Li, G. G. S. King, K. N. Slessor, J. H. Borden, W. W. Bowers, R. J. West and E. W. Underhill, 1993 (b). Sex pheromone of the western hemlock looper, *Lambdina fiscellaria lugubrosa* (Hulst) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 19: 1009-1019.

Gries, G., G. G. S. King, R. Gries, P. D. C. Wimalaratne, T. G. Gray, R. F. Shepherd, J.-X. Li, K. N. Slessor and G. Khaskin, 1993 (c). 3,13-Dimethylheptadecane: major sex pheromone component of the western false hemlock looper, *Nepytiia freemani* Munroe (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 19: 1501-1510.

Gries, G., H. L. McBrien, R. Gries, J.-X. Li, G. J. R. Judd, K. N. Slessor, J. H. Borden, R. F. Smith, M. Christie, J. T. Troubridge, P. D. C. Wimalaratne and E. W. Underhill, 1993 (d). (E4, E10)-Dodecadienyl acetate: novel sex pheromone component of tentiform leafminer, *Phyllonorycter mespilella* (Hübner) (Lepidoptera: Gracillariidae). *J. Chem. Ecol.*, 19: 1789-1798.

Gries, G., J.-X. Li, R. Gries, W. W. Bowers, R. J. West, P. D. C. Wimalaratne, G. Khaskin, G. G. S. King and K. N. Slessor, 1994 (a). (E)-11,13-Tetradecadienal: major sex pheromone component of the eastern blackheaded budworm, *Acleris variana* (Fern.)

(Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **20**: 1-8.

Gries, R., G. Gries, J.-X. Li, C. T. Maier, C. R. Lemmon and K. N. Slessor, 1994 (b). Sex pheromone components of the spring hemlock looper, *Lambdina athasaria* (Walker) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **20**: 2501-2511.

Gries, G., R. Gries, G. Khaskin, K. N. Slessor, G. G. Grant, J. Liska and P. Kapitola, 1996. Specificity of nun and gypsy moth sexual communication through multiple-component pheromone blends. *Naturwissenschaften*, **83**: 382-385.

Gries, G., K. N. Slessor, R. Gries, G. Khaskin, P. D. C. Wimalaratne, T. G. Gray, G. G. Grant, A. S. Tracey and M. Hulme, 1997 (a). (*Z*)6, (*E*)8-Heneicosadien-11-one: synergistic sex pheromone component of Douglas-fir tussock moth, *Orgyia pseudotsugata* (McDunnough) (Lepidoptera: Lymantriidae). *J. Chem. Ecol.*, **23** (1): 19-34.

Gries, R., G. Gries, G. G. S. King and C. T. Maier, 1997 (b). Sex pheromone components of the apple leafminer, *Lyonetia prunifoliella*. *J. Chem. Ecol.*, **23** (4): 1119-1130.

Gries, R., D. Holden, G. Gries, P. D. C. Wimalaratne, K. N. Slessor and C. Saunders, 1997 (c). 3*Z*-*cis*-6,7-*cis*-9,10-Di-epoxy-heneicosene: Novel class of Lepidopteran pheromone. *Naturwissenschaften*, **84**: 219-221.

Gries, R., E. Dunkelblum, G. Gries, F. Baldilla, C. Hernandez, F. Alvarez, A. Perez, J. Velasco and A. C. Oehlschlager, 1998. Sex pheromone components of *Diatraea considerata* (Heinrich) (Lep., Pyralidae). *J. Appl. Entomol.*, **122**: 265-268.

Gries, G., J. Clearwater, R. Gries, G. Khaskin, S. King and P. Schaefer 1999 (a). Synthetic sex pheromone components of white-spotted tussock moth, *Orgyia thyellina*. *J. Chem. Ecol.*, **25** (5): 1091-1104.

Gries, G., R. Gries, P. W. Schaefer, T. Gotoh and Y. Higashiura, 1999 (b). Sex pheromone components of pink gypsy moth, *Lymantria mathura*. *Naturwissenschaften*, **86**: 235-238.

Gries, G., P. W. Schaefer, G. Khaskin, R. Hahn, R. Gries and J.-T. Chao, 1999 (c). Sex pheromone components of Casuarina moth, *Lymantria xyloina*. *J. Chem. Ecol.*, **25** (11): 2535-2545.

Gries, G., P. W. Schaefer, R. Gries, J. Liska and T. Gotoh, 2001. Reproductive character displacement in *Lymantria monacha* from northern Japan?. *J. Chem. Ecol.*, **27** (6): 1163-1176.

Gries, G., P. W. Schaefer, R. Gries, Y.-B. Fan, Y. Higashiura and B. Tanaka, 2002. 2-Methyl-(*Z*)-7-octadecene: sex pheromone of allopatric *Lymantria lucescens* and *L. serva*. *J. Chem. Ecol.*, **28** (3): 469-478.

Gries, R., G. Khaskin, E. Khaskin, J. L. Foltz, P. W. Schaefer and G. Gries, 2003. Enantiomers of (*Z,Z*)-6,9-heneicosadien-11-ol: sex pheromone components of *Orgyia detrita*. *J. Chem. Ecol.*,

29 (10) : 2201-2212.

- Gries, R., A. Reckziegel, H. Bogenschütz, H.-G. Kontzog, C. Schlegel, W. Francke, J. G. Millar and G. Gries, **2004**.
*(Z,Z)-11,13-Hexadecadienyl acetate and
(Z,E)-11,13,15-hexadecatrienyl acetate: synergistic sex
pheromone components of oak processionary moth, Thaumetopoea
processionea (Lepidoptera: Thaumetopoeidae).* *Chemoecology*, **14** (2) : 95-100.
- Gries, R., G. Khaskin, P. W. Schaefer, R. Hahn, T. Gotoh and G. Gries, **2005** (a). *(7R,8S)-cis-7,8-Epoxy-2-methyloctadec-17-ene: a novel
trace component from the sex pheromone gland of gypsy moth,
Lymantria dispar.* *J. Chem. Ecol.*, **31** : 49-62.
- Gries, R., G. Khaskin, J. Cleawater, D. Hasman, P. W. Schaefer, E. Khaskin, O. Miroshnychenko, G. Hosking and G. Gries, **2005** (b).
*(Z,Z)-6,9-Heneicosadien-11-one: major sex pheromone component
of painted apple moth, Teia anartoides.* *J. Chem. Ecol.*, **31** (3) : 603-620.
- Gries, R., G. Khaskin, T. Gotoh, P. W. Schaefer and G. Gries, **2005** (c).
*(7Z,9E)-2-Methyl-7,9-octadecadiene: a sex pheromone component
of Lymantria bantaizana.* *J. Chem. Ecol.*, **31** (4) : 879-891.
- Gries, R., G. Gries, P. W. Schäfer, H. J. S. Yoo and M. Greaves, **2005** (d).
*(Z,E)-6,8-Heneicosadien-11-one: major sex-pheromone component
of Orgyia vetusta (Lepidoptera: Lymantriidae).* *Can. Entomol.*, **137** : 471-475.
- Gries, R., G. Khaskin, Z.-X. Tan, B.-G. Zhao, G. G. S. King, A. Miroshnychenko, G.-Q. Lin, M. Rhainds and G. Gries, **2006** (a).
*(1S)-1-Ethyl-2-methylpropyl 3,13-dimethylpentadecanoate:
major sex pheromone component of Paulownia bagworm, Clania
variegata.* *J. Chem. Ecol.*, **32** (8) : 1673-1685.
- Gries, R., G. Khaskin, H. Daroogheh, C. Mart, S. Karadag, M. Kubilay Er, R. Britton and G. Gries, **2006** (b).
*(2S,12Z)-2-Acetoxy-12-heptadecene: Major sex pheromone
component of pistachio twigborer, Kermaniapistaciella.* *J. Chem. Ecol.*, **32** (12) : 2667-2677.
- Gries, R., P. W. Schaefer, R. Hahn, G. Khaskin, G. Ramaseshiah, B. Singh, G. K. Hehar and G. Gries, **2007**. Sex pheromone components
of Indian gypsy moth, *Lymantria obfuscata*. *J. Chem. Ecol.*, **33** (9) : 1774-1786.
- Grodner**, J., K. Świech, M. Jakubowska and J. Bocianowski, **2018**.
Attraction of moths of two Noctuidae species to field traps baited
with a mixture of two to three homologous acetates in Poland.
J. Econ. Entomol., **111** : 1664-1673.
- Groot**, A. T., S. Nojima, J. J. Heath, B. Ammagarahalli, M. van Wijk, A. Claßen, R. G. Santangelo, J. Lopez and C. Schal, **2018**. Alcohol contributes to attraction of *Heliothis* (= *Chloridea*) *virescens* males to females. *J. Chem. Ecol.*, **44** : 621-630.

- Gu**, P., C.-L. Peng, J. Li, C.-H. Feng, P. Li and Y.-J. Du, 2013. Optimization and field evaluation of the sex pheromone of *Trichophysetis cretacea* (Lepidoptera, Crambidae) for pest monitoring. *J. Appl. Entomol.*, **137**: 462-468.
- Guerin**, P. M., H. Arn, C. Blaser and M. Lettéré, 1983. Z,E-8,10-dodecadien-1-ol, attractant for male *Pammene rhediella*. *Entomol. Exp. Appl.*, **33**: 346-347.
- Guerin, P. M., W. Baltensweiler, H. Arn and H.-R. Buser, 1984. Host race pheromone polymorphism in the larch budmoth. *Can. Entomol.*, **40**: 892-894.
- Guerin, P. M., H. Arn, H. R. Buser and P. J. Charmillot, 1986(a). Sex pheromone of *Adoxophyes orana*: additional components and variability in ratio of (Z)-9- and (Z)-11-tetradecenyl acetate. *J. Chem. Ecol.*, **12**: 763-772.
- Guerin, P. M., H. Arn, H. R. Buser, P. Charmillot, M. Tóth and G. Szira'ki, 1986(b). Sex pheromone of *Grapholita funebrana*. Occurrence of Z-8- and Z-10-tetradecenyl acetate as secondary components. *J. Chem. Ecol.*, **12**: 1361-1368.
- Guerin, P. M., H. R. Buser, M. Tóth, E. Höbaus, A. Schmid and H. Arn, 1986(c). Sex pheromone of *Sparganothis pilleriana*: E- and Z-11-tetradecenyl acetates as essential components. *Entomol. Exp. Appl.*, **40**: 137-140.
- Guerrero**, A., F. Camps, J. Coll, M. Riba, J. Einhorn, C. Descoins and J. Y. Lallemand, 1981. Identification of a potential sex pheromone of the processionary moth, *Thaumetopoea pityocampa* (Lepidoptera, Notodontidae). *Tetrahedron Lett.*, **22**: 2013-2016.
- Guerrero, A., R. Canela, M. Riba, M. Eizaguirre, R. Albajes and V. Sarto, 1986. A sex attractant for males of the noctuid moth *Athetis hospes*. *Entomol. Exp. Appl.*, **42**: 97-100.
- Guleria**, N., S. M. Nebapure, P. D. K. Jayanthi, S. B. Suby and M. G. Deeksha, 2023. Electrophysiological and behavioral responses of spotted stemborer, *Chilo partellus*, to sex pheromone components and their blends. *J. Chem. Ecol.*, **49**: 155-163.
- Guo**, G., Z. Li, G. Lin, P. Wu, X. Zhang, H. Liu and X. Yang, 1990(a). Identification and synthesis of insect pheromone. XXVIII. Sex pheromone of poplar pale clearwing moth, *Sphecia sinensis* Hsu. *J. Chem. Ecol.*, **16**: 819-826.
- Guo, G.-Z., H.-Q. Liu, G.-Q. Lin, C.-H. Wu, L.-Z. Meng and D.-M. Chen, 1990(b). Insect sex pheromone of vine tree borer *Paranthrene regalis*. *Youji Huaxue*, **10**: 504-506.
- Guo**, Q., S. Yang, F. Zhuo, W. Xiang, C. Feng, S. Zhang and Y. Du, 2022. Effect of age, daily rhythm and mating on pheromone titres and trapping of *Glyphodes pyloalis* (Lepidoptera, Crambidae). *J. Appl. Entomol.*, **146**: 875-884.

Hai, T. V., L. V. Vang, P. K. Son, S. Inomata and T. Ando, 2002. Sex attractants for moths of Vietnam: field attraction by synthetic lures baited with known lepidopteran pheromones. *J. Chem. Ecol.*, **28**(7): 1473-1481.

Hall, D. R. and J. S. Read, 1979. Sex attractants for two zygaenid moths. *J. Entomol. Soc. South Afr.*, **42**: 115-119.

Hall, D. R., P. S. Beevor, R. Lester and B. F. Nesbitt, 1980. (*E,E*)-10,12-Hexadecadienal: a component of the female sex pheromone of the spiny bollworm, *Earias insulana* (Boisd.) (Lepidoptera: Noctuidae). *Experientia*, **36**: 152-154.

Hall, D. R., P. S. Beevor, A. Cork, R. Lester, B. F. Nesbitt, G. K. C. Nyirenda, D. D. N. Phiri, B. W. Blair and J. Tannock, 1981. The female sex pheromone of the maize stalk-borer *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae): identification and initial field trials. *Zimbabwe J. Agric. Res.*, **19**: 111-122.

Hall, D. R., P. S. Beevor, A. Cork, B. F. Nesbitt and E. A. S. La Croix, 1984. (*Z*)-8-Dodecenyl acetate: the major component of the female sex pheromone of *Cryptophlebia batrachopa*, a tortricid pest of macadamia in Malawi. *Entomol. Exp. Appl.*, **35**: 33-36.

Hall, D. R., A. Cork, R. Lester, B. F. Nesbitt and P. Zagatti, 1987. Sex pheromone of rice moth, *Corcyra cephalonica* Stainton. II. Identification and role of female pheromone. *J. Chem. Ecol.*, **13**: 1575-1589.

Hall, D. R., P. S. Beevor, D. G. Campon, D. J. Chamberlain, A. Cork, R. D. White, A. Almestral and T. J. Henneberry, 1992. Nitrate esters: Novel sex pheromone components of the cotton leafperforator, *Bucculatrix thurberiella* Busck. (Lepidoptera: Lyonetiidae). *Tetrahedron Lett.*, **33**: 4811-4814.

Hall, D. R., P. S. Beevor, D. G. Campon, D. J. Chamberlain, A. Cork, R. White, A. Almestral, T. J. Henneberry, Nandagopal, V. Wightman, J. A. and G. V. R. Rao, 1993. Identification and synthesis of new pheromones. *Bulletin OILB SROP*, **16**: 1-9.

Hall, D. R., D. Farman, J. C. Domínguez and J. A. Pajares, 2017. Female sex pheromone of the cone moth, *Dioryctria mendacella*: investigation of synergism between Type I and Type II pheromone components. *J. Chem. Ecol.*, **43**: 433-442.

Halperin, J. and Y. Golan, 1982. Attraction of males of *Thaumetopoea wilkinsoni* Tams to the sex pheromone of *Th. pityocampa* Schiff. *Phytoparasitica*, **10**: 140-141.

Han, G.-B., R.-Y. Ma, J.-W. Du and C. Hu, 1999. The sex pheromone communication system of adult jujube leafroller *Ancylis sativa* Liu. *Act. Entomol. Sinica (Kunchong Xuebao)*, **42**: 25-30.

Han, K.-S., J.-K. Jung,, K.-H. Choi,, S.-W. Lee,, and K.-S. Boo, 2000. Sex pheromone composition and male trapping of the peach fruit

moth, *Carposina sasakii* (Matsumura) (Lepidoptera: Carposinidae) in Korea. *J. Asia Pacific Entomol.*, 3: 83-88.

Han, K.-S., J.-K. Jung, K.-H. Choi, S.-W. Lee and K.-S. Boo, 2001. Sex pheromone composition and male trapping of the oriental fruit moth, *Grapholita molesta* (Lepidoptera: Tortricidae) in Korea. *J. Asia Pacific Entomol.*, 4: 31-35.

Hansen, K., 1984. Discrimination and production of disparlure enantiomers by the gypsy moth and the nun moth. *Physiol. Entomol.*, 9: 9-18.

Hansson, B. S., M. Tóth, C. Löfstedt, G. Szöcs, M. Subchev and J. Löfqvist, 1990(a). Pheromone variation among eastern European and a western Asian population of the turnip moth *Agrotis segetum*. *J. Chem. Ecol.*, 16: 1611-1622.

Hansson, B. S., G. Szöcs, F. Schmidt, W. Francke, C. Löfstedt and M. Tóth, 1990(b). Electrophysiological and chemical analysis of sex pheromone communication system of the mottled umber, *Erannis defoliaria* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 16: 1887-1897.

Hansson, B. S., A. Blackwell, E. Hallberg and J. Löfqvist, 1995. Physiological and morphological characteristics of the sex pheromone detecting system in male corn stemborers, *Chilo partellus* (Lepidoptera: Pyralidae). *J. Insect Physiol.*, 41(2): 171-178.

Hanula, J. L., C. W. Berisford and G. L. DeBarr, 1984. Pheromone cross-attraction and inhibition among four coneworms, *Dioryctria* spp. (Lepidoptera: Pyralidae) in a loblolly pine seed orchard. *Environ. Entomol.*, 13: 1298-1301.

Harris, M. K., A. A. A. Fu, H. Nunez, E. Aranda-Herera, J. A. Moreira, J. S. McElfresh and J. G. Millar, 2008. A new pheromone race of *Acrobasis nuxvorella* (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, 101: 769-776.

Hathaway, D. O. and G. Tamaki, 1981. Another olethreutine, *Phaneta latens* (Lepidoptera: Tortricidae), attracted to the sex pheromone of the codling moth. *J. Entomol. Soc. Brit. Columbia*, 78: 24-27.

Hattori, M., S. Wakamura, K. Igita, K. Yasuda and Tridjaka, 2001. Comparison of the characteristics and sex pheromone of *Etiella behrii* (Zeller), a newly identified pod borer of soybean in Indonesia, with *E. zinckenella* (Treit.). *Jpn. Int. Res. Quart.*, 35: 19-24.

Haynes, K. F., 1987. Identification of sex pheromone of calendula plume moth, *Platyptilia williamsii*. *J. Chem. Ecol.*, 13: 907-916.

Haynes, K. F., 1990. Identification of sex pheromone of bristly cutworm, *Lacinipolia renigera* (Stephens). *J. Chem. Ecol.*, 16: 2615-2621.

Haynes, K. F., K. V. Yeargan, J. G. Millar and B. B. Chastain, 1996.

Identification of sex pheromone of *Tetanolita mynesalis* (Lepidoptera: Noctuidae), a prey species of bolas spider, *Mastophora hutchinsoni*. *J. Chem. Ecol.*, **22**: 75-89.

He, Y., H. Xie and S. Chi, 2016. Identification of female main component sex pheromones of *Orthaga achatina* (Lepidoptera: Pyralidae) and their attractiveness to moths. *Agric. Sci. Technol.*, **17**: 920-922.

Heath, R. R., J. H. Tumlinson, N. C. Leppla, J. R. McLaughlin, B. Dueben, E. Dundulis and R. H. Guy, 1983. Identification of a sex pheromone produced by female velvetbean caterpillar moth. *J. Chem. Ecol.*, **9**: 645-656.

Heath, R. R. and P. J. Landolt, 1987. (Z)-5-Decen-1-ol acetate: the major sex pheromone component of *Argyrogramma verruca* (Fabricius) (Lepidoptera: Noctuidae: Plusiinae). *Environ. Entomol.*, **16**: 1030-1033.

Heath, R. R., P. E. A. Teal, N. D. Epsky, B. D. Dueben, S. D. Hight, S. Bloem, J. E. Carpenter, T. J. Weissling, P. E. Kendra, J. Cibrian-Tovar and K. A. Bloem, 2006. Pheromone-based attractant for males of *Cactoblastis cactorum* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **35**: 1469-1476.

Hedin, P. A., F. M. Davis, J. C. Dickens, M. L. Burks, T. G. Bird and A. E. Knutson, 1986. Identification of the sex attractant pheromone of the southwestern corn borer *Diatraea grandiosella* Dyar. *J. Chem. Ecol.*, **12**: 2051-2063.

Hedlin, A. F., J. Weatherston, D. S. Ruth and G. E. Miller, 1983. Chemical lure for male Douglas-fir cone moth, *Barbara colfaxiana* (Lepidoptera: Olethreutidae). *Environ. Entomol.*, **12**: 1751-1753.

Heikinheimo, O., 1978. Feromonfangst som metod för faunistiska undersökningar. *Norw. J. Entomol.*, **25**: 102.

Henderson, H. E., F. L. Warren, O. P. H. Augustyn, B. V. Burger, D. F. Schneider, P. R. Boshoff, H. S. C. Spies and H. Geertsema, 1973. Isolation and structure of the sex-pheromone of the moth, *Nudaurelia cytherea cytherea*. *J. Insect Physiol.*, **19**: 1257-1264.

Hendricks, D. E., B. A. Leonhardt and T. N. Shaver, 1989. Development of optimized blends of two sex pheromone components impregnated in PVC dispensers for tobacco budworm bait. *Southwestern Entomol.*, **14**: 17-25.

Hendrikse, A., W. M. Herrebout and C. J. Persoons, 1982. Sex pheromones, attractants and inhibitors in small ermine moths: a comparison of bioassays. *Meded. Fac. Landbouww. Rijksuniv. Gent.*, **47**: 493-502.

Hendry, L. B., J. Jugovich, L. Roman, M. E. Anderson and R. O. Mumma, 1974. *Cis*-10-tetradecenyl acetate, an attractant component in the sex pheromone of the oak leaf roller moth (*Archips semiferanus*

Walker). *Experientia*, **30**: 886-887.

Herrera, H., W. Barros-Parada, M. F. Flores, W. Francke, E. Fuentes-Contreras, M. Rodriguez, F. Santis, P. H. G. Zarbin and J. Bergmann, **2016**. Identification of a novel moth sex pheromone component from *Chilecomadia valdiviana*. *J. Chem. Ecol.*, **42**: 908-918.

Hill, A., R. Cardé, A. Comeau, W. Bode and W. Roelofs, **1974**. Sex pheromone of the tufted apple bud moth (*Platynota idaeusalis*). *Environ. Entomol.*, **3**: 249-252.

Hill, A. S. and W. L. Roelofs, **1975**(a). Sex pheromone components of the omnivorous leafroller moth, *Platynota stultana*. *J. Chem. Ecol.*, **1**: 91-99.

Hill, A. S., R. T. Cardé, H. Kido and W. L. Roelofs, **1975**(b). Sex pheromone of the orange tortrix moth, *Argyrotaenia citrana* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **1**: 215-224.

Hill, A. S., C. W. Berisford, U. E. Brady and W. L. Roelofs, **1976**. Sex pheromone of the pitch pine tip moth, *Rhyacionia rigidana*. *Environ. Entomol.*, **5**: 959-960.

Hill, A. S., R. T. Cardé, W. M. Bode and W. L. Roelofs, **1977**. Sex pheromone components of the variegated leafroller moth, *Platynota flavedana*. *J. Chem. Ecol.*, **3**: 369-376.

Hill, A. S. and W. L. Roelofs, **1978**. Two sex attractants for male speckled green fruitworm moths, *Orthosia hibisci* Guenée (Lepidoptera: Noctuidae). *J. New York Entomol. Soc.*, **86**: 296.

Hill, A. S. and W. L. Roelofs, **1979**(a). Sex pheromone components of the obliquebanded leafroller moth, *Choristoneura rosaceana*. *J. Chem. Ecol.*, **5**: 3-11.

Hill, A. S., R. W. Rings, S. R. Swier and W. L. Roelofs, **1979**(b). Sex pheromone of the black cutworm moth, *Agrotis ipsilon*. *J. Chem. Ecol.*, **5**: 439-457.

Hill, A. S. and W. L. Roelofs, **1980**. A female-produced sex pheromone component and attractant for males in the armyworm moth, *Pseudaletia unipuncta*. *Environ. Entomol.*, **9**: 408-411.

Hill, A. S., C. W. Berisford, U. E. Brady and W. L. Roelofs, **1981**(a). Nantucket pine tip moth, *Rhyacionia frustrana*: identification of two sex pheromone components. *J. Chem. Ecol.*, **7**: 517-528.

Hill, A. S. and W. L. Roelofs, **1981**(b). Sex pheromone of the saltmarsh caterpillar moth, *Estigmene acrea*. *J. Chem. Ecol.*, **7**: 655-668.

Hill, A. S., B. G. Kovalev, L. N. Nikolaeva and W. L. Roelofs, **1982**. Sex pheromone of the fall webworm moth, *Hyphantria cunea*. *J. Chem. Ecol.*, **8**: 383-396.

Hillier, N. K., P. L. Dixon, W. D. Seabrook and D. J. Larson, **2002**. Field testing of synthetic attractants for male *Grapholita*

libertina (Lepidoptera: Tortricidae). *Can. Entomol.*, **134**: 657-665.

Himeno, K. and H. Honda, 1992. (*E, Z*) - and (*E, E*) -10,12-Hexadecadienals, major components of female sex pheromone of the cotton leaf-roller, *Notarcha derogata* (Fabricius) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **27**: 507-515.

Hirai, Y., H. Kimura, K. Kawasaki and Y. Tamaki, 1978. (*Z*) -11-Hexadecenyl acetate: a sex-pheromone component of the cabbage armyworm moth, *Mamestra brassicae* Linné. *Appl. Entomol. Zool.*, **13**: 136-137.

Hirano, C., H. Muramoto and M. Horiike, 1976. Sex pheromone produced by female sweet potato leaf folder moth. *Naturwissenschaften*, **63**: 439.

Ho, H.-Y., Y.-T. Tao, R.-S. Tsai, Y.-L. Wu, H.-K. Tseng and Y.-S. Chow, 1996. Isolation, identification, and synthesis of sex pheromone components of female tea cluster caterpillar, *Andracia bipunctata* Walker (Lepidoptera: Bombycidae) in Taiwan. *J. Chem. Ecol.*, **22**: 271-285.

Ho, H. Y., R. S. Tsai, E. L. Hsu, Y. S. Chow and R. Kou, 2002. Investigation of possible sex pheromone components of female *loreyi* leafworm *Acantholeucania loreyi* (Duponchel) (Lepidoptera: Noctuidae) in Taiwan. *Zool. Studies*, **41**(2): 188-193.

Hoddle, M. S., J. G. Millar, C. D. Hoddle, Y. Zou and J. S. McElfresh, 2009. Synthesis and field evaluation of the sex pheromone of *Stenoma catenifer* (Lepidoptera: Elachistidae). *J. Econ. Entomol.*, **102**: 1460-1467.

Hoffmann, M. P., L. M. McDonough and J. B. Bailey, 1983. Field test of the sex pheromone of *Amorbia cuneana* (Walsingham) (Lepidoptera: Tortricidae). *Environ. Entomol.*, **12**(5): 1387-1390.

Holden, D. G., 2000. Sex pheromone components of satin moth, *Leucoma salicis* (L.) (Lepidoptera: Lymantriidae). Master Thesis, Simon Fraser University, Burnaby, B.C., Canada.

Honda, H., T. Ando, Y. Ogura, T. Teramine and R. Ueda, 1990. Identification of the sex pheromone of the mulberry pyralid, *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **25**: 265-272.

Honda, H., K. Himeno and Y. Yoshiyasu, 1994. Chemotaxonomy of the cotton leaf-roller (Lepidoptera: Pyralidae) in Japan with special reference to differences in sex pheromones. *Appl. Entomol. Zool.*, **29**(3): 323-330.

Honda, H. and T. Kimura, 2004. (*E*) -10-Hexadecenal as a sex attractant of *Pleuroptia chloropahanta* (Butler) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **48**: 57-60. (in Japanese)

Honda, H., R. Yamasaki, Y. Sumiuchi, T. Uehara, S. Matsuyama, T. Ando and H. Naka, 2015. Hybrid sex pheromones of the hibiscus

flower-bud borer, *Rehimena surusalis*. *J. Chem. Ecol.*, **41**: 1043-1049.

Honma, K., K. Kawasaki and Y. Tamaki, 1978. Sex-pheromone activities of synthetic 7-alken-11-ones for male peach fruit moths, *Carposina nipponensis* Walsingham (Lepidoptera: Carposinidae). *Jap. J. Appl. Entomol. Zool.*, **22**: 87-91.

Horák, A. and I. Hrdý, 1988. Attractiveness of (E)-10-dodecen-1-yl acetate for the pea moth, *Cydia nigricana*: modification by codlemone, codlemone acetate and dodecyl acetate. *Acta Entomol. Bohemoslov.*, **85**: 174-181.

Horak, M., C. P. Whittle, T. E. Bellas and E. R. Rumbo, 1988. Pheromone gland components of some Australian tortricids in relation to their taxonomy. *J. Chem. Ecol.*, **14**: 1163-1175.

Hrdý, I., J. Marek and F. Krampf, 1979. Sexual pheromone activity of 8-dodecenyl and 11-tetradecenyl acetates for males of several lepidopteran species in field trials. *Acta Entomol. Bohemoslov.*, **76**: 65-84.

Hrdý, I., J. Liska, J. Vrkoc and R. Hochmut, 1989. New records on sex attractants for males and faunistic comments on moth (Lepidoptera) from Czechoslovakia. *Acta Entomol. Bohemoslov.*, **86**: 252-268.

Hrudová, E, 2003. The presence of non-target lepidopteran species in pheromone traps for fruit tortricid moths. *Plant Protect. Sci.*, **39**: 126-131.

Hsiao, S. N. 1990. Bioassay of sex pheromone of the smaller tea tortrix, *Adoxophyes* sp. (Lepidoptera: Tortricidae). *Chin. J. Entomol.*, **10**: 443-450.

Hu, Wenli and J. Du, 2005. Sex pheromone secondary components of Indian meal moth *Plodia interpunctella* in China. *Yingyong Shengtai Xuebao*, **16**(9): 1751-1755. [SciFinder, 2005:1285000]

Hu, D.-H., J. He, Y.-W. Zhou, J.-T. Feng and X. Zhang, 2012. Synthesis and field evaluation of the sex pheromone analogues to soybean pod borer *Leguminivora glycinivorella*. *Molecules*, **17**: 12140-12150.

Huang, Y.-P., S. Tatsuki, C.-G. Kim, S. Hoshizaki, Y. Yoshiyasu, H. Honda and Y. Ishikawa 1997. Identification of sex pheromone of adzuki bean borer, *Ostrinia sacculalis*. *J. Chem. Ecol.*, **23**(12): 2791-2802.

Huang, Y., T. Takanashi, S. Hoshizaki, S. Tatsuki, H. Honda, Y. Yoshiyasu and Y. Ishikawa 1998(a). Geographic variation in sex pheromone of Asian corn borer, *Ostrinia furnacalis*, in Japan. *J. Chem. Ecol.*, **24**(12): 2079-2088.

Huang, Y., S. Tatsuki, C.-G. Kim, S. Hoshizaki and Y. Ishikawa, 1998(b). Identification of the sex pheromone of *Ostrinia*

palustralis. *Entomol. Exp. Appl.*, **86**: 313-318.

Huang, Y., H. Honda, Y. Yoshiyasu, S. Hoshizaki, S. Tatsuki and Y. Ishikawa, 1998(c). Sex pheromone of the butterbur borer, *Ostrinia zaguliaevi*. *Entomol. Exp. Appl.*, **89**: 281-287.

Huang, Y., T. Takanashi, S. Hoshizaki, S. Tatsuki and Y. Ishikawa, 2002. Female sex pheromone polymorphism in adzuki bean borer, *Ostrinia scapulalis*, is similar to that in European corn borer, *O. nubilalis*. *J. Chem. Ecol.*, **28**(3): 533-539.

Huang, J., L. J. Gut and M. J. Grieshop, 2022. Attraction of *Choristoneura rosaceana* to pheromone blends in ratios produced in glands or emitted by females. *J. Appl. Entomol.*, **146**: 734-742.

Hummel, H. E., L. K. Gaston, H. H. Shorey, R. S. Kaae, K. J. Byrne and R. M. Silverstein, 1973. Clarification of the chemical status of the pink bollworm sex pheromone. *Science*, **181**: 873-875.

Hung, C. C., J. S. Hwang, M. D. Hung, Y. P. Yen and R. F. Hou, 2001. Isolation, identification and field tests of the sex pheromone of the carambola fruit borer, *Eucosma notanthes*. *J. Chem. Ecol.*, **27** (9): 1855-1866.

Hung, C. C., M. D. Hung and W. L. Wang, 2006. Development of formulations of sex attractant of *Conopomorpha sinensis* on *Litchi chinesis*. *Zhiwu Baohu Xuehui Huikan*, **48**(3): 189-202. [SciFinder, 2007:3272]

Hwang, J. S., C. C. Hung and R. F. Hou, 1996(a). A bioassay method of the sex pheromone of the carambola fruit borer, *Eucosma notanthes* Meyrick. *Zhiwu Baohu Xuehui Huikan*, **38**(2): 119-127. [Chem. Abst., **126**: 2969s]

Hwang, J. S., C. C. Hung, Y. P. Yen and C. N. Chen, 1996(b). Sex attractant and trap designs for trapping litchi fruit borer, *Conopomorpha sinensis* Bradley. *Zhiwu Baohu Xuehui Huikan (Plant Protection Bulletin)*, **38**(2): 129-136.

Ichikawa, A., T. Yasuda and S. Wakamura, 1995. Absolute configuration of sex pheromone for tea tussock moth, *Euproctis pseudoconspersa* (Strand) via synthesis of (R)- and (S)-10,14-dimethyl-1-pentadecyl isobytyrates. *J. Chem. Ecol.*, **21**: 627-634.

Inomata, S., M. Komoda, H. Watanabe, M. Nomura and T. Ando, 2000. Identification of sex pheromones of *Anadevidia peponis* and *Macdunnoughia confusa*, and field tests of their role in reproductive isolation of closely related Plusiinae moths. *J. Chem. Ecol.*, **26**: 443-454.

Inomata, S., S. Tsuchiya, K. Ikeda, O. Saito and T. Ando, 2002. Identification of sex pheromone components secreted by female moths of *Peridroma saucia* (Noctuidae: Noctuinae). *Biosci. Biotechnol. Biochem.*, **66**(11): 2461-2464.

Inomata, S., A. Watanabe, M. Nomura and T. Ando, 2005. Mating

communication systems of four Plusiinae species distributed in Japan: identification of the sex pheromones and field evaluation. *J. Chem. Ecol.*, **31**(6): 1429-1442.

Ishikawa, Y., T. Takanashi and Y. Huang, 1999(a). Comparative studies on the sex pheromones of *Ostrinia* spp. In Japan: the burdock bore, *Ostrinia zealis*. *Chemoecology*, **9**: 25-32.

Ishikawa, Y., T. Takanashi, C.-G. Kim, S. Hoshizaki, S. Tatsuki and Y. Huang, 1999(b). *Ostrinia* spp. in Japan: their host plants and sex pheromones. *Entomol. Exp. Appl.*, **91**: 237-244.

Islam, MD. A., M. Yamamoto, M. Sugie, H. Naka, J. Tabata, Y. Arita and T. Ando, 2007. Synthesis and characterization of 2,13- and 3,13-octadecadienals for the identificatin of the sex pheromone secreted by a clearwing moth. *J. Chem. Ecol.*, **33**(9): 1763-1773.

Islam, MD. A., R. Yamakawa, N. D. Do, N. Numakura, T. Suzuki and T. Ando, 2009. Instrumental analysis of terminal-conjugated dienes for reexamination of the sex pheromone secreted by a nettle moth, *Parasa lepida lepida*. *Biosci. Biotechnol. Biochem.*, **73**: 1156-1162.

Iwaki, S., S. Marumo, T. Saito, M. Yamada and K. Katagiri, 1974. Synthesis and activity of optically active disparlure. *J. Amer. Chem. Soc.*, **96**: 7842-7844.

Jaastad, G., M. Bengtsson, P. Anderson, S. Kobro, G. Knudsen and P. Witzgall, 2002. Sex pheromone of apple fruit moth *Argyresthia conjugella* (Lepidoptera: Argyresthiidae). *Agric. Forest Entomol.*, **4**: 233-236.

Jacobson, M., R. E. Redfern, W. A. Jones, M. H. Aldridge, 1970. Sex pheromones of the southern armyworm moth: isolation, identification, and synthesis. *Science*, **170**: 542-544.

Jacobson, M. and D. T. Jennings, 1978. Attraction of *Rhyacionia neomexicana* (Dyar) to synthetic pheromones. *J. Environ. Sci. Health, A* **13**: 429-443.

Jain, S. C., D. E. Dussourd, W. E. Conner, T. Eisner, A. Guerrero and J. Meinwald, 1983. Polyene pheromone components from an arctiid moth (*Utetheisa ornatrix*): characterization and synthesis. *J. Org. Chem.*, **48**: 2266-2270.

James, D. G., A. Cosse, L. C. Wright and J. Perez, 2001. Pheromone trapping of *Synanthedon tipuliformis* (Lepidoptera: Sesiidae) in Washington red currants. *Environ. Entomol.*, **30**: 663-666.

Jennings, D. T. and M.W. Houseweart, 1983. Field attractiveness of (E)- and (Z)-11-tetradecenal pheromone blends to male spruce budworm moths, *Choristoneura fumiferana* (Clemens). *J. Chem. Ecol.*, **9**: 1327-1332.

Jham, G. N., A. A. da Silva, E. R. Lima and P. Viana, 2005. Identification (GC and GC-MS) of unsaturated acetates in *Elasmopalpus lignosellus*

and their biological activity (GC-EAD and EAG). *J. Sep. Sci.*, **28**: 281-285.

Jham, G. N., A. A. da Silva, E. R. Lima and P. A. Viana, **2007**. Identification of acetates in *Elasmopalpus lignosellus* pheromone glands using a newly created mass spectral database and Kóvats retention indices. *Quim. Nova*, **30** (4): 916-919.

Jiang, N.-J., B.-T. Mo, H. Guo, J. Yang, R. Tang and C.-Z. Wang, **2022**. Revisiting the sex pheromone of the fall armyworm *Spodoptera frugiperda*, a new invasive pest in South China. *Insect Sci.*, **29**: 865-878.

Jing, X.-Y., J.-T. Zhang, Y.-Q. Luo, S.-X. Zong, P.-H. Liu and J.-X. Jia, **2010**. Identification and field evaluation of the female sex pheromone of the sand *Salix* carpenterworm, *Holcocerus arenicola* Staudinger (Lepidoptera: Cossidae). *Z. Naturforsch.*, **65c**: 403-411.

Jones, R. L. and A. N. Sparks, **1979**. (Z)-9-Tetradecen-ol acetate. A secondary sex pheromone of the fall armyworm, *Spodoptera frugiperda* (J. E. Smith). *J. Chem. Ecol.*, **5**: 721-725.

Jordan, T. A., A. Zhang and D. G. Pfeiffer, **2013**. Blend chemistry and field attraction of commercial sex pheromone lures to grape berry moth (Lepidoptera: Tortricidae), and a nontarget tortricid in vineyards. *Environ. Entomol.*, **42**: 558-563.

Judd, G. J. R., R. Gries, V. M. Aurelian and G. Gries, **2011**. 3Z,13Z-octadecadienyl acetate: sex pheromone of the apple clearwing moth in British Columbia. *Can. Entomol.*, **143**: 236-244.

Judd, G. J. R., R. Gries, C. Teasdale and G. Gries, **2012**. Identification of a sex pheromone component for *Pennisetia marginata* (Lepidoptera: Sesiidae). *Can. Entomol.*, **144**: 798-807.

Jung, C. R., Y. J. Park and K. S. Boo, **2003**. Optimal sex pheromone composition for monitoring *Spodoptera exigua* (Lepidoptera: Noctuidae) in Korea. *J. Asia-Pac. Entomol.*, **6**: 175-182.

Kaae, R. S., H. H. Shorey, S. U. McFarland and L. K. Gaston, **1973** (a). Sex pheromones of Lepidoptera. XXXVII. Role of sex pheromones and other factors in reproductive isolation among ten species of Noctuidae. *Ann. Entomol. Soc. Am.*, **66**: 444-448.

Kaae, R. S., H. H. Shorey and L. K. Gaston, **1973** (b). Pheromone concentration as a mechanism for reproductive isolation between two lepidopterous species. *Science*, **179**: 487-489.

Kakizaki, M. and H. Sugie, **2003**. Sex pheromone of the flax budworm, *Heliothis maritima adaucta* Butler (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **38**(1): 73-78.

Kalinová, B., A. Minaif and L. Kotera, **1994**. Sex pheromone characterization and field trapping of the European corn borer,

Ostrinia nubilalis (Lepidoptera: Pyralidae), in South Moravia and Slovakia. *Eur. J. Entomol.*, **91**: 197-203.

Kalinová, B., A. Svatos, J. Kindl, O. Hovorka, I. Hrdý, J. Kuldova and M. Hoskovec, 2003. Sex pheromone of horse-chestnut leafminer *Cameraria ohridella* and its use in a pheromone-based monitoring system. *J. Chem. Ecol.*, **29**(2): 387-404.

Kalinová, B., J. Kindl, O. Hovorka, M. Hoskovec and A. Svatos, 2005. (11Z)-Hexadec-11-enal enhances the attractiveness of *Diatraea saccharalis* main pheromone component in wind tunnel experiments. *J Appl. Entomol.*, **129** (2): 70-74.

Kalinová, B., P. Jiros, J. Zd'arek, X. Wen and M. Hoskovec, 2006. GC x GC/TOF MS technique. A new tool in identification of insect pheromones: analysis of the persimmon bark borer sex pheromone gland. *Talanta* **69**: 542-547.

Kalinová, B., R. R. D. Nascimento, M. Hoskovec, A. L. Mendonça, E. L. Silva, M. R. T. D. Freitas, C. R. Cabral-Jr, C. E. Silva, A. E. G. Sant'Ana and A. Svatoš, 2012. Identification of two components of the female sex pheromone of the sugarcane-borer *Diatraea flavipennella* (Lepidoptera: Crambidae). *J. Appl. Entomol.*, **136**: 203-211.

Kamm, J. A. and L. M. McDonough, 1979. Sex attractant for a plume moth, *Oidaematophorus monodactylus* (L.) (Lepidoptera: Pterophoridae). *Pan-Pacific Entomol.*, **55**: 284-286.

Kamm, J. A. and L. M. McDonough, 1980. Synergism of the sex pheromone of the granberry girdler. *Environ. Entomol.*, **9**: 795-797.

Kamm, J. A., L. M. McDonough and C. L. Smithhisler, 1982. Sex attractant for *Protagrotis obscura*, a pest of grass grown for seed. *Environ. Entomol.*, **11**: 118-120.

Kamm, J. A. and L. M. McDonough, 1985. Sex attractant for *Diarsia pseudorosaria*, a defoliator of ryegrass (Lepidoptera: Noctuidae). *Pan-Pacific Entomol.*, **61**: 299-302.

Kamm, J. A., L. M. McDonough and K. E. Rowe, 1989. Effect of dosage and ratio of sex pheromone components on behavior of *Chrysoteuchia topiaria* (Zeller) (Lepidoptera: Pyralidae). *Environ. Entomol.*, **18**: 368-372.

Kang, S.-K., B.-H. Moon, J.-O. Lee and H.-G. Goh, 1985. A Convergent synthesis of (Z)-13-octadecen-1-yl acetate, the pheromone mimic of the rice leaf folder moth and its biological activity test. *Bull. Korean Chem. Soc.*, **6**; 228-230.

Kang, S.-K. and K.-Y. Seol, 1995. Sex pheromone of the perilla leaf pyralid moth: isolation and identification. *Bull. Korean Chem. Soc.*, **16**; 688-689.

Kanno, H., L. P. S. Kuenen, K. A. Klingler, J. G. Millar and R. T. Cardé, 2010. Attractiveness of a four-component pheromone blend to male navel orangeworm moths. *J. Chem. Ecol.*, **36**:

584-591.

Karalius, V., R. Mozūraitis, J. Miatleuski, V. Bûda and P. Ivinskis, 2001. Sex attractants for six clearwing and tineid species (Lepidoptera, Sesiidae and Tineidae) from Kazakhstan and Lithuania. *Z. Naturforsch.*, 56c: 1120-1125.

Karalius, V., R. Mozūraitis, V. Bûda, I. Libikas and A.-K. Borg-Karlson, 2005. Sex attractants for six moth species of the families Brachodidae, Choreutidae and Tineidae from Kazakhstan and Lithuania. *Z. Naturforsch.*, 60c: 625-631.

Karalius, V. and V. Bûda, 2006. Sex attractants for clearwing moths *Synanthedon mesiaeformis* (Herrich-Schaffer) and *S. flaviventris* (Staudinger) (Lepidoptera, Sesiidae). *Acta Zool. Lit.*, 16: 198-203.

Karandinos, M. G., J. H. Tunlinson and T. D. Eichlin, 1977. Field evidence of synergism and inhibition in the Sesiidae sex pheromone system. *J. Chem. Ecol.*, 3: 57-64.

Kasang, G., K. E. Kaissling, O. Vostrowsky and H. J. Bestmann, 1978 (a). Bombykal, a second pheromone component of the silkworm moth *Bombyx mori* L. *Angew. Chem. Int. Ed. Engl.*, 17: 60.

Kasang, G. and D. Schneider, 1978 (b). The silkworm moth *Bombyx mori*. Presence of the (E,E)-stereoisomer of bombykol in the female pheromone gland. *Naturwissenschaften*, 65: 337-338.

Katovich, S. A., P. D. Swedenborg, M. Giblin and E. W. Underhill, 1989. Evidence for (E,Z)-8,10-dodecadienyl acetate as the major component of the sex pheromone of the eastern pine seedworm, *Cydia toreuta* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, 15: 581-590.

Kawahara, A. Y., M. Jones, Q. Jia, S. L. LaPointe and P. A. Stansly, 2013. A synthetic pheromone for *Phyllocnistis citrella* (Lepidoptera: Gracillariidae) attracts multiple leafminer species. *Florida Entomol.*, 96: 1213-1216.

Kawazu, K., J. Hasegawa, H. Honda, Y. Ishikawa, S. Wakamura, H. Sugie, H. Kamiwada, T. Kamimuro, Y. Yoshiyasu and S. Tatsuki, 2000. Geographical variation in female sex pheromones of the rice leaffolder moth, *Cnaphalocrocis medinalis*: identification of pheromone components in Japan. *Entomol. Exp. Appl.*, 96: 103-109.

Kawazu, K., O. Setokuchi, K. Kohno, K. Takahashi, Y. Yoshiyasu and S. Tatsuki, 2001. Sex pheromone of the rice leaffolder moth, *Cnaphalocrocis medinalis* (Lepidoptera: Crambidae): Synthetic Indian and Philippine blends are not attractive to male *C. medinalis*, but are attractive to *C. pilosa* in the South-Western islands n Japan. *Appl. Entomol. Zool.*, 36: 471-474.

Kawazu, K., K. Nagata, Z. Zhang, H. Sugie and S. Tatsuki, 2002. Comparison of attractiveness in Japan and China of three synthetic pheromone blends based on geographic variations in the rice

leaffolder, *Cnaphalocrocis medinalis* (Lepidoptera: Pyralidae).
Bull. Entomol. Res., **92**: 295-299.

Kawazu, K., H. Honda, S. Nakamura and T. Adati, 2007. Identification of sex pheromone components of the box tree pyralid, *Glyphodes perspectalis*. *J. Chem. Ecol.*, **33**: 1978-1985.

Kawazu, K., T. Adati, Y. Yosiyasu, K. Sumiartha, W. Susila, P. Sudiarta, H. Purwanto, and S. Tatsuki, 2009. Sex pheromone components of the rice leaffolder, *Cnaphalocrocis medinalis* (Lepidoptera: Crambidae), in Indonesia. *J. Asia Pacific Entomol.*, **12**: 97-99.

Keaster, A. J., J. A. Grundler, W. S. Craig and M. A. Jackson, 1996. Noctuidmoths and other insects captured in wing-style traps baited with black cutworm (Lepidoptera: Noctuidae) pheromone on offshore oil platforms in the Gulf of Mexico, 1988-1991. *J. Kansas Entomol. Soc.*, **69**: 17-25.

Keathley, C. P., L. L. Stelinski and S. L. Lapointe, 2013. Attraction of a native Florida leafminer, *Phyllocnistis insignis* (Lepidoptera: Gracillariidae), to pheromone of an invasive citrus leafminer, *P. citrella*: Evidence for mating disruption of a native non-target species. *Florida Entomol.*, **96**: 877-886.

Kehat, M., S. Greenberg and Y. Tamaki, 1976. Field evaluation of the synthetic sex pheromone, as an attractant for males of the cotton leafworm, *Spodoptera littoralis* (Boisd.), in Israel. *Appl. Entomol. Zool.*, **11**: 45-52.

Kehat, M., S. Gothilf, E. Dunkelblum and S. Greenberg, 1980. Field evaluation of female sex pheromone components of the cotton bollworm, *Heliothis armigera*. *Entomol. Exp. Appl.*, **27**: 188-193.

Kehat, M. and E. Dunkelblum, 1990. Behavioral responses of male *Heliothis armigera* (Lepidoptera: Noctuidae) moths in a flight tunnel to combinations of components identified from female sex pheromone glands. *J. Insect Behavior*, **3**: 75-83.

Kehat, M., A. Eitam, D. Blumberg, E. Dunkelblum, L. Anshelevich and D. Gordon, 1992. Sex pheromone traps for detecting and monitoring the raisin moth, *Cadra figulilella*, in date palm plantations. *Phytoparasitica*, **20**: 99-106.

Khrimian, A., D. R. Lance, M. Schwarz, B. A. Leonhardt and V. Mastro, 2008. Sex pheromone of browntail moth, *Euproctis chrysorrhoea* (L.): synthesis and field development. *J. Agric. Food Chem.*, **56**: 2452-2456.

Kim, J. and I.-K. Park, 2013. Female sex pheromone components of the box tree pyralid, *Glyphodes perspectalis*, in Korea: Field test and development of film-type lure. *J. Asia-Pac. Entomol.*: **16**, 473-477.

Kim, J., K.-S. Cho, C. Y. Yang, C. G. Park, 2014 (a). Identification and field evaluation of the female sex pheromone of *Stathmopoda masinsissa* in Korea. *Chemoecology*, **24**: 253-259.

- Kim, J., H. S. Roh, B. K. Byun and C. G. Park, 2014 (b). A co-attractant mixture of (*Z*)-11-tetradecenyl acetate and (*Z*)-9-dodecenyl acetate for *Neocalyptis angustilineata* and *Homona magnanima*. *J. Asia-Pac. Entomol.*, **17**: 659-661.
- Kim, J., B.-K. Byun, H.-W. Oh, S. A. Jang and C. G. Park, 2016. *Athetis dissimilis* (Lepidoptera: Noctuidae) is attracted to the sex pheromone of *Euzophera batangensis* (Lepidoptera: Pyralidae). *J. Asia-Pac. Entomol.*, **19**: 841-845.
- Kindl**, J., P. Jiroš, B. Kalinová, P. Žáček and I. Valterová, 2012. Females of the bumblebee parasite, *Aphomia sociella*, excite males using a courtship pheromone. *J. Chem. Ecol.*, **38**: 400-407.
- King**, G. G. S., R. Gries, G. Gries and K. N. Slessor, 1995. Optical isomers of 3,13-dimethylheptadecane: sex pheromone components of the western false hemlock looper, *Nepytiā freemani* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **21**: 2027-2045.
- Kinjo**, M., M. Nagamine, H. Sugie and Y. Tamaki, 1996. Sex pheromone of the sugarcane shoot borer, *Tetramoera schistaceana* Snellen (Lepidoptera: Tortricidae: Olethreutinae). *Jpn. J. Appl. Entomol. Zool.*, **40**; 191-197.
- Klimetzek**, D., G. Loskant, J. P. Vité and K. Mori, 1976. Disparlure: differences in pheromone perception between gypsy moth and nun moth. *Naturwissenschaften*, **63**: 581-582
- Klun**, J. A. and T. A. Brindley, 1970. *cis*-11-Tetradecenyl acetate, a sex stimulant of the European corn borer. *J. Econ. Entomol.*, **63**: 779-780.
- Klun, J. A. and J. F. Robinson, 1972. Olfactory discrimination in the European corn borer and several pheromonally analogous moths. *Ann. Entomol. Soc. Am.*, **65**: 1337-1340.
- Klun, J. A., O. L. Chapman, K. C. Mattes, P. W. Wojtkowski, M. Beroza and P. E. Sonnet, 1973. Insect sex pheromones: minor amount of opposite geometrical isomer critical to attraction. *Science*, **181**: 661-663.
- Klun, J. A. and cooperators, 1975. Insect sex pheromones: intraspecific pheromonal variability of *Ostrinia nubilalis* in North America and Europe. *Environ. Entomol.*, **4**: 891-894.
- Klun, J. A. and G. A. Junk, 1977. Iowa European corn borer sex pheromone. Isolation and identification of four C₁₄ esters. *J. Chem. Ecol.*, **3**: 447-459.
- Klun, J. A. and S. Maini, 1979. Genetic basis of an insect chemical communication system: the European corn borer. *Environ. Entomol.*, **8**: 423-426.
- Klun, J. A., B. A. Bierl-Leonhardt, M. Schwarz, J. A. Litsinger, A. T. Barrion, H. C. Chiang and Z. Jiang, 1980 (a). Sex pheromone

of the Asian corn borer moth. *Life Sci.*, **27**: 1603-1606.

Klun, J. A., J. R. Plimmer, B. A. Bierl-Leonhardt, A. N. Sparks, M. Primiani, O. L. Chapman, G. H. Lee and G. Lepone, **1980**(b). Sex pheromone chemistry of female corn earworm moth, *Heliothis zea*. *J. Chem. Ecol.*, **6**: 165-175.

Klun, J. A., B. A. Bierl-Leonhardt, J. R. Plimmer, A. N. Sparks, M. Primiani, O. L. Chapman, G. Lepone and G. H. Lee, **1980**(c). Sex pheromone chemistry of the female tobacco budworm moth, *Heliothis virescens*. *J. Chem. Ecol.*, **6**: 177-183.

Klun, J. A., K. F. Haynes, B. A. Bierl-Leonhardt, M. C. Birch and J. R. Plimmer, **1981**. Sex pheromone of the female artichoke plume moth, *Platyptilia carduidactyla*. *Environ. Entomol.*, **10**: 763-765.

Klun, J. A., B. A. Leonhardt, J. D. Lopez, Jr. and L. E. Lachance, **1982**. Female *Heliothis subflexa* (Lepidoptera: Noctuidae) sex pheromone: chemistry and congeneric comparisons. *Environ. Entomol.*, **11**: 1084-1090.

Klun, J. A., C. C. Blickenstaff, M. Schwarz, B. A. Leonhardt and J. R. Plimmer, **1983**. Western bean cutworm, *Loxagrotis albicosta* (Lepidoptera: Noctuidae): female sex pheromone identification. *Environ. Entomol.*, **12**: 714-717.

Klun, J. A., B. A. Leonhardt, M. Schwarz, A. Day and A. K. Raina, **1986**. Female sex pheromone of the pickleworm, *Diaphania nitidalis* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **12**: 239-249.

Klun, J. A., J. W. Neal, Jr. and B. A. Leonhardt, **1989**. Female sex pheromone of iris borer, *Macronoctua onusta* (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **15**: 1559-1566.

Klun, J. A., M. Schwarz, B. A. Leonhardt and W. W. Cantelo, **1990**. Sex pheromone of the female squash vine borer (Lepidoptera: Sesiidae). *J. Entomol. Sci.*, **25**(1): 64-72.

Klun, J. A., Y. N. Baranchikov, V. C. Mastro, Y. Hijji, J. Nicholson, I. Ragenovich and T. A. Vshivkova, **2000**. A sex attractant for the Siberian moth *Dendrolimus superans sibiricus* (Lepidoptera: Lasiocampidae). *J. Entomol. Sci.*, **35**(2): 158-166.

Knauf, W., H. J. Bestmann, K. H. Koschatzky, J. Süss and O. Vostrowsky, **1979**. Studies on the attractiveness of sex attractants towards *Tortrix viridana* L. and *Panolis flammea* Schiff. *Z. Angew. Entomol.*, **88**: 307-312.

Knauf, W., H. J. Bestmann and O. Vostrowsky, **1984**. Responses of male winter moths (*Operophtera brumata*) to their sex attractant (3Z,6Z,9Z)-1,3,6,9-nona-decatetraene and to some structural analogues. *Entomol. Exp. Appl.*, **35**: 208-210.

Kochansky, J., R. T. Cardé, J. Liebherr and W. L. Roelofs, **1975**(a). Sex pheromone of the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae), in New York. *J. Chem. Ecol.*, **1**:

225-231.

Kochansky, J., J. Tette, E. F. Taschenberg, R. T. Cardé, K.-E. Kaissling and W. L. Roelofs, 1975(b). Sex pheromone of the moth, *Antheraea polyphemus*. *J. Insect Physiol.*, **21**: 1977-1983.

Kochansky, J. P., W. L. Roelofs and P. Sivapalan, 1978. Sex pheromone of the tea tortrix moth (*Homona coffearia* Neitner). *J. Chem. Ecol.*, **4**: 623-631.

Kochansky, J., A. Hill, J. W. Neal Jr., J.-A. Bentz and W. L. Roelofs, 1996. The pheromone of the eastern tent caterpillar, *Malacosoma americanum* (F.) (Lepidoptera: Lasiocampidae). *J. Chem. Ecol.*, **22**: 2251-2261.

Komai, F. 1999. A taxonomic review of the genus *Grapholita* and allied genera (Lepidoptera: Tortricidae) in the Palaearctic region. *Ent. Scand. Suppl.* **55**: 1-226.

Kohno, K., H. Iida, M. Takeda and K. Honda, 2014. Non-target lepidopteran insects specifically attracted to sex pheromone lures for *Helicoverpa armigera* and *H. assulta* (Lepidoptera: Noctuidae). *Jpn. J. Appl. Entomol. Zool.*, **58**: 343-350. (in Japanese)

Kong, F.-L., S.-D. Yu, Y.-Q. Yu, X.-H. Hu, J.-W. Xu and P.-C. Zhu, 1990. Sex pheromone of eggplant borer, *Leucinodes orbonalis* Guenée. *Youji Huaxue*, **10**: 258-260.

Kong, X., Z. Zhang, H. Wang and C. Zhao, 2005. Study on trace component in sex pheromone of *Dendrolimus* spp. *Sepu*, **23**(4): 370-373. [SciFinder, 2005:824663]

Kong, X. B., Z. Zhang, C. H. Zhao and H. B. Wang, 2007(a). Female sex pheromone of the Yunnan pine caterpillar moth *Dendrolimus houi*: first (E,Z)-isomers in pheromone components of *Dendrolimus* spp. *J. Chem. Ecol.*, **33** (7): 1316-1327.

Kong, X.-B., C.-H. Zhao and R. Wang, 2007(b). Sex pheromone of the larch caterpillar moth, *Dendrolimus superans*, from northeastern China. *Entomol. Exp. Appl.*, **124** (1): 37-44.

Kong, X.-B., X.-L. Sun, H.-B. Wang, Z. Zhang, C-H. Zhao and K. C. J. H. Booij, 2011. Identification of components of the female sex pheromone of the Simao pine caterpillar moth, *Dendrolimus kikuchii* Matsumura. *J. Chem. Ecol.*, **37**: 412-419.

Kong, X.-B., K.-W. Liu, H.-B. Wang, S.-F. Zhang and Z. Zhang. 2012 (a). Identification and behavioral evaluation of sex pheromone components of the Chinese pine caterpillar moth, *Dendrolimus tabulaeformis*. *PLOS ONE.*, **7**: e33381.

Kong, X.-B., H.-B. Wang and Z. Zhang, 2012 (b). Determination of the absolute configuration of an EAG active component in the sex pheromone gland of *Semiothisa cirraria* Bremer et Grey (Lepidoptera: Geometridae). *Acta Entomol. Sinica*, **55**: 162-167.

Konjuchov, V. P., B. G. Kovalev and N. R. Sattar-Zade, 1983. Isolation and identification of *Heliothis armigera* Hb. sex pheromone components. *Bioorganisk Khimiya*, 9: 1422-1428.

Konno, Y., K. Arai, K. Sekiguchi and Y. Matsumoto, 1982. (E)-10-Hexadecenal, a sex pheromone component of the yellow peach moth, *Dichocrocis punctiferalis* Guenée (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, 17: 207-217.

Konyukhov, V. P. and B. G. Kovalev, 1988. Sex pheromone of lackey moth (*Malacosoma neustria* L.) females. *Bioorg. Khim*, 14: 268-272. [Chem. Abst., 108: 164997e]

Kosaki, A., T. Uehara, H. Naka, S. Matsuyama, T. Ando and H. Honda, 2021. Identification and behavioral assays of sex pheromone components in *Smerinthus tokyonis* (Lepidoptera: Sphingidae). *Appl. Entomol. Zool.*, 56: 373-378.

Koshihara, T., H. Yamada, Y. Tamaki and T. Ando, 1978. Field attractiveness of the synthetic sex-pheromone of the diamondback moth, *Plutella xylostella* (L.). *Appl. Entomol. Zool.*, 13: 138-141.

Kou, R., Y.-S. Chow, S. Takahashi and R. Yamaoka, 1989. Evidence for the presence and nature of a female sex pheromone of *Brithys crini* Fabricius (Lepidoptera: Noctuidae). *Chung Yang Yen Chiu Yuan Chih Wu Yen Chiu So Chuan K'an* (Phytochem. Ecol.), 429-434. [Chem. Abst., 113: 56115k]

Kou, R., D. S. Tang, Y. S. Chow and H. K. Tseng, 1990. Sex pheromone components of female smaller tea tortrix moth, *Adoxophyes* sp. (Lepidoptera: Tortricidae) in Taiwan. *J. Chem. Ecol.*, 16: 1409-1415.

Kou, R., Y. S. Chow, S. Takahashi and R. Yamaoka, 1991. Evidence for presence and nature of female sex pheromone of *Brithys crini* Fabricius (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, 17: 2035-2042.

Kou, R., H. Y. Ho, H. T. Yang, Y. S. Chow and H. J. Wu, 1992 (a). Investigation of sex pheromone components of female Asian corn borer, *Ostrinia furnacalis* (Hübner) (Lepidoptera: Pyralidae) in Taiwan. *J. Chem. Ecol.*, 18: 833-840.

Kou, R., Y.-S. Chow and H.-Y. Ho, 1992 (b). Chemical composition of sex pheromone gland extract in female oriental armyworm *Pseudaletia separata* Walker (Lepidoptera: Noctuidae) in Taiwan. *Bull. Inst. Zool., Academia Sinica*, 31: 246-250.

Kou, R., 1992c. Calling behavior and pheromone titer in the smaller tea tortrix moth, *Adoxophyes* sp. (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, 18: 855-861.

Kovaleski, A., M. Botton, O. Nakano, E. F. Vilela and A. E. Eiras,

2003. Concentration and duration of the synthetic sex pheromone of *Bonagota cranaodes* (Meyrick) (Lepidoptera: Tortricidae) in an apple orchard. *Neotropical Entomol.*, **32**(1): 45-48.

Kovalev, B. G., D. V. Sokolova, T. S. Bolgar and N. P. Sekerskaya, 1985. Identification of the sex pheromone of *Cacocimorpha pronubana*. *Khim. Prir. Soedin.*, 110-112. [Chem. Abst., **102**: 182701j]

Kovalev, B. G. and L. A. Nikolaeva, 1986. Attractiveness of some epoxy compounds for males of a tiger moth, *Phragmatobia fuliginosa* (Arctiidae), and of a geometrid moth, *Caustoloma (Therapis) flaviguria* (Geometriidae). *Zool Zh.*, **56**: 802-804.

Kovalev, B. G. and S. Samoilov, 1987. Sex pheromone of female *Eurhodope advenella* Zck. (Lepidoptera, Pyralidae): identification by GLC and EAG methods. *Khemoretspsiya Nasekomykh*, **9**: 26-29.

Kovalev, B. G., J. H. Tamlinson, T. S. Bolgar, T. P. Kolomoets, V. M. Kovalenko and R. R. Heath, 1988. Isolation and identification of a component in the sex pheromone of *Aegeria apiformis*. *Khim. Prir. Soedin.*, 126-128. [Chem. Abst., **108**: 202002m]

Kovalev, B. G., K. Lofsteot, S. F. Nedopekina, A. M. Sorochinskaya, K. S. Artokhin, E. A. Nesterov and E. Viik, 1991. Isolation and identification of the female sex pheromone of *Heliothis maritima* (Lepidoptera, Noctuidae). *Khim. Prir. Soedin.*, 574-577. [Chem. Abst., **118**: 54297a]

Kovalev, B. G., T. S. Bolgar, P. A. Zubov, D. G. Zharkov, M. Golosova, E. A. Nesterov and M. S. Tvaradze, 1993. Identification of additional components of the sex pheromone of *Dendrolimus pini*. *Chem. Nat. Compd.*, **29**: 135-136.

Kovaleva, A. S., L. L. Ivanov, I. B. Piatnova, B. G. Kovalev, V. A. Minyailo and A. K. Minyailo, 1975. Results of testing synthetic compounds as to their attractiveness to insects in nature. *Khemoretspsiya Nasekomykh*, **2**: 159-164. [Chem. Abst., **85**: 105340]

Kozaki, J., S. Nakagaki, R. Sato, M. Yukinari, H. Sugie, H. Noguchi and Y. Tamaki, 1984. Sex pheromone of the common apple leafroller, *Hoshinoia longicellana* Walsingham (Lepidoptera: Tortricidae): isolation and identification. *Appl. Entomol. Zool.*, **19**: 382-388.

Kozlov, M. V., J. Zhu, P. Philipp, W. Francke, E. L. Zvereva, B. S. Hansson and C. Löfstedt, 1996. Pheromone specificity in *Eriocrania semipurpurella* (Stephens) and *E. sangii* (Wood) (Lepidoptera: Eriocraniidae) based on chirality of semiochemicals. *J. Chem. Ecol.*, **22**: 431-454.

Krasnoff, S. B., K. W. Vick and J. A. Coffelt, 1984. (*Z,E*)-9,12-Tetradecadien-1-ol: a component of the sex pheromone of *Ephestia elutella* (Hübner) (Lepidoptera: Pyralidae). *Environ. Entomol.*, **13**: 765-767.

Krasnoff, S. B. and W. L. Roelofs, 1988. Sex pheromone released as

an aerosol by the moth *Pyrrharctia isabella*. *Nature*, 333: 263-265.

Krishnakumari, B., A. L. Prasuna, K. N. Jyothi, M. Y. Valli, S. Sighamony, A. R. Prasad and J. S. Yadav 1998. Behavioural and electrophysiological responses of *Achaea janata* Linn. (Lepidoptera: Noctuidae) males to synthetic female-produced sex pheromone components. *J. Entomol. Res.*, 22(3): 197-202.

Krivokhizhin, V. I., V. V. Brovko, V. A. Minyailo and G. S. Zolotarenko, 1984. *Cis*-11-hexadecen-1-yl acetate as a sexual attractant for cutworm moths (Lepidoptera, Noctuidae) in Western Siberia. *Sib. Vestn. S-kh. Nauki*, 33-38. [*Chem. Abst.*, 101: 124847e]

Krokos, F. D., A. Ameline, J. Bau, A. Sans, M. Konstantopoulou, B. Frerot, A. Guerrero, M. Eizaguirre, C. Malosse, O. Etchepare, R. Albajes and B. E. Mazomenos, 2002. Comparative studies of female sex pheromone components and male response of the corn stalk borer *Sesamia nonagrioides* in three different populations. *J. Chem. Ecol.*, 28(7): 1463-1472.

Kuenen, L. P. S., J. W. Peacock, P. J. Silk and S. S. Wright, 1990. Identification of the primary female sex pheromone component of the mimosa webworm (Lepidoptera: Plutellidae). *Environ. Entomol.*, 19: 1-7.

Kuenen, L. P. S., J. S. McElfresh and J. G. Millar, 2010. Identification of critical secondary components of the sex pheromone of the navel orangeworm (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, 103: 314-330.

Kurihara, M., K. Usui, K. Uchiumi and S. Tatsuki, 1991. Sex pheromone of the lawn grass cutworm moth, *Spodoptera depravata* (Butler) (Lepidoptera: Noctuidae). *Jap. J. Appl. Entomol. Zool.*, 35: 323-324.

Kuwahara, Y., C. Kitamura, S. Takahashi, H. Hara, S. Ishii and H. Fukami, 1971(a). Sex pheromone of the almond moth and the Indian meal moth: *cis*-9, *trans*-12-tetradecadienyl acetate. *Science*, 171: 801-802.

Kuwahara, Y., H. Hara, S. Ishii and H. Fukami, 1971(b). The sex pheromone of the Mediterranean flour moth. *Agric. Biol. Chem.*, 35: 447-448.

Kuwahara, Y. and J. E. Casida, 1973. Quantitative analysis of the sex pheromone of several phycitid moths by electron-capture gas chromatography. *Agric. Biol. Chem.*, 37: 681-684.

Kuwahara, Y. and L. T. M. Yen, 1979. Identification of bombykol in *Bombyx mandarina* females (Lepidoptera: Bombycidae). *Appl. Entomol. Zool.*, 14: 114-117.

Kuwahara, Y., N. Mori, S. Yamada and T. Nemoto, 1984. Evaluation of bombykol as the sex pheromone of *Bombyx mandarina* (Lepidoptera: Bombycidae). *Appl. Entomol. Zool.*, 19: 265-267.

Lacey, M.J. and C.J. Sanders, 1992. Chemical composition of sex

pheromone of oriental fruit moth and rates of release by individual female moths. *J. Chem. Ecol.*, **18**: 1421-1435.

La Croix, E. A. S., H. Z. Thindwa and D. R. Hall, 1985. Field studies with the synthetic female sex pheromones of *Cryptophlebia batrachopa* and *C. leucotreta*, pests of macadamia in Malawi. *Trop. Pest Manag.*, **31**: 189-191.

Lalanne-Cassou, B. and B. Frerot, 1980. Analysis of pheromonal secretion of South African carnation tortrix, *Epichoristodes acerbella* (Walk.) (Lepidoptera: Tortricidae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **291**: 845-848.

Landolt, P. J. and C. E. Curtis, 1982. Interspecific sexual attraction between *Pyralis farinalis* L. and *Amyelois transitella* (Walker) (Lepidoptera: Pyralidae). *J. Kansas Entomol. Soc.*, **55**: 248-252.

Landolt, P. J., J. D. Solomon and R. E. Doolittle, 1985. Attraction of male *Prionoxystus piger* (Lepidoptera: Cossidae) to isomers of 3,5-tetradecadien-1-ol acetate. *Florida Entomol.*, **68**: 228-230.

Landolt, P. J., R. R. Heath, P. E. Sonnet and K. Matsumoto, 1986 (a). Attraction of *Harrisina americana* and *Acoloithus falsarius* males (Lepidoptera: Zygaenidae) to (*R*) -(-)-2-butyl-(*Z*)-7-tetradecenoate. *Environ. Entomol.*, **15**: 959-962.

Landolt, P. J., R. R. Heath and N. C. Leppla, 1986 (b). (*Z,Z,Z*)-3,6,9-Eicosatriene and (*Z,Z,Z*)-3,6,9-heneicosatriene as sex pheromone components of a grass looper, *Mocis disseverans* (Lepidoptera: Noctuidae). *Environ. Entomol.*, **15**: 1272-1274.

Landolt, P. J. and R. R. Heath, 1986 (c). A sex attractant synergist for *Trichoplusia oxygramma* (Lepidoptera: Noctuidae). *Florida Entomol.*, **69**: 425-426.

Landolt, P. J., J. H. Tumlinson and M. M. Brennan, 1989 (a). Attraction of *Amphion floridensis* (Lepidoptera: Sphingidae) to bombykal, (*E,Z*)-10,12-hexadecadienal. *Florida Entomol.*, **72**: 324-327.

Landolt, P. J. and R. R. Heath, 1989 (b). Lure composition, component ratio, and dose for trapping male *Mocis latipes* (Lepidoptera: Noctuidae) with synthetic sex pheromone. *J. Econ. Entomol.*, **82**: 307-309.

Landolt, P. J. and R. R. Heath, 1991. Zygaenidae trapped with enantiomers of 2-butyl (*Z*)-7-tetradecenoate. *J. Lepido. Soc.*, **45**: 63-65.

Landolt, P. J., M. Tóth, W. Francke and K. Mori, 1996. Attraction of moths (Lepidoptera: Arctiidae, Geometridae, Noctuidae) to enantiomers of several epoxydienes. *Florida Entomol.*, **79**(3): 392-397.

Landolt, P. J. and C. L. Smithhisler 1998. Isolation and identification

of female sex pheromone and development of a sex attractant for *Lacanobia subjuncta*. *J. Chem. Ecol.*, **24**(8): 1369-1379.

Landolt, P. J., C. Smithhisler, T. Adams and R. S. Zack, 2003. An improved multi-component sex attractant for trapping male western yellowstriped armyworm, *Spodoptera praefica* (Grote) (Lepidoptera: Noctuidae). *Agric. Forest Entomol.*, **5**: 333-339.

Landolt, P. J., C. Guédot and R. S. Zack, 2011. Spotted cutworm, *Xestia c-nigrum* (L.) (Lepidoptera: Noctuidae) responses to sex pheromone and blacklight. *J. Appl. Entomol.*, **135**: 593-600.

Landolt, P. J. and D. Roberts, 2012. A sex attractant for trapping *Crambus cypridalis* (Lepidoptera: Crambidae). *Florida Entomol.*, **95**: 194-195.

Lapointe, S. L., D. G. Hall, Y. Murata, A. L. Parra-pedrazzoli, J. M. S. Bento, E. F. Vilela and W. S. Leal, 2006. Field evaluation of a synthetic female sex pheromone for the leafmining moth *Phyllocnistis citrella* (Lepidoptera: Gracillariidae) in Florida citrus. *Florida Entomol.*, **89** (2), 274-276.

Lapointe, S. L., W. Barros-Parada, E. Fuentes-Contreras, H. Herrera, T. Kinsho, Y. Miyake, R. P. Niedz and J. Bergmann 2017. Use of mixture designs to investigate contribution of minor sex pheromone components to trap catch of the carpenterworm moth, *Chilecomadia valdiviana*. *J. Chem. Ecol.*, **43**: 1046-1655.

Lassance, J.-M., G. P. Svensson, M. V. Kozlov, W. Francke and C. Löfstedt, 2019. Pheromones and barcoding delimit boundaries between cryptic species in the primitive moth genus *Eriocrania* (Lepidoptera: Eriocraniidae). *J. Chem. Ecol.*, **45**: 429-439.

Leal, W. S., J. M. S. Bento, Y. Murata, M. Ono, J. R. P. Parra and E. F. Vilela, 2001. Identification, synthesis, and field evaluation of the sex pheromone of the citrus fruit borer *Ecdytolopha aurantiana*. *J. Chem. Ecol.*, **27**(10): 2041-2051.

Leal, W. S., A. L. Parra-Pedrazzoli, K.-E. Kaissling, T. I. Morgan, F. G. Zalom, D. J. Pesak, E. A. Dundulis, C. S. Burks and B. S. Higbee, 2005. Unusual pheromone chemistry in the navel orangeworm: novel sex attractants and a behavioral antagonist. *Naturwissenschaften*, **92**: 139-146.

Leal, W. S., A. L. Parra-Pedrazzoli, A. A. Cossé, Y. Murata, J. M. S. Bento and E. F. Vilela, 2006. Identification, synthesis, and field evaluation of the sex pheromone from the citrus leafminer, *Phyllocnistis citrella*. *J. Chem. Ecol.*, **32**(1): 155-168.

Lebedeva, K. V., V. V. Voronkova and N. I. Bocharova, 1991. Pheromone of the turnip moth (*Agrotis segetum*). *Agrokhimiya*, 99-114. [Chem. Abst., **116**: 209681f]

Lebedeva, K. V., N. V. Vendilo, S. A. Kurbatov, V. A. Pletnev, V. L. Ponomarev, Y. B. Pyatnova and N. I. Bocharova, 2000(a). Identification of the pheromone of eastern-meadow cutworm *Mythimna separata* (Lepidoptera: Noctuidae). *Agrokhimiya*, (5): 57-69. [Chem. Abst., 134: 278087]

Lebedeva, K. V., N. V. Vendilo, V. A. Pletnev, V. L. Ponomarev, S. A. Kurbatov, V. V. Voronkova and V. A. Shchennikov, 2000(b). Study of the pheromone of the heart-and-dart moth, *Agrotis exclamatio*nis (Lepidoptera: Noctuidae). *Agrokhimiya*, (8): 71-75. [Chem. Abst., 134: 262292]

Lecheva, I., 1999. Possibilities of implementation of synthetic sex pheromones for the geometrid *Erannis bajaria* (Lepidoptera: Geometridae). *Restenievud. Nauki.*, 36(7-8): 397-400. [Chem. Abs., 132: 318901]

Lee, S., D.-W. Lee and K. S. Boo, 2005. Sex pheromone composition of the diamondback moth, *Plutella xylostella* (L.) in Korea. *J. Asia-Pacific Entomol.*, 8: 243-248.

Lee, S.-C., E.-C. Koo, D.-H. Lee, S.-M. Seo, M.-J. Huh, I Nam & J.-H. Park and I.-K. Park, 2020. Identification of the female-produced sex pheromone of the dotted white geometrid *Naxa seriaria* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 46: 927-934.

Lee, S.-C., J.-W. Lee, D. H. Lee, M.-J. Huh, I. Nam, J.-H. Park, M. Jung and I.-K. Park, 2022. Identification of sex pheromone components of Korean *Dioryctria abietella* (Lepidoptera: Pyralidae) population and synergism of pheromone and pine cone volatile blends. *J. Econ. Entomol.*, 115: 178-186.

Leeuw, I., 1983. *Thyris maculata* (Thyrididae) and three species of clearwing moths (Sesiidae) attracted to an artificial chemical bait. *J. Lepi. Soc.*, 37: 87-88.

Legrand, S., M. Botton, M. Coracini, P. Witzgall and C. R. Unelius, 2004. Synthesis and field tests of sex pheromone components of the leafroller *Argyrotaenia sphaleropa*. *Z. Naturforsch.*, 59c: 708-712.

Lemmen, J. and M. Evenden, 2009. Peripheral and behavioral plasticity of pheromone response and its hormonal control in a long-lived moth. *J. Exp. Biol.*, 212: 2000-2006.

Leonhardt, B. A., J. W. Neal, Jr., J. A. Klun, M. Schwarz and J. R. Plimmer, 1983. An unusual lepidopteran sex pheromone system in the bagworm moth. *Science*, 219: 314-316.

Leonhardt, B. A., V. C. Mastro, M. Schwarz, J. D. Tang, R. E. Charlton, A. Pellegrini-Toole, J. D. Warthen, Jr., C. P. Schwalbe and R. T. Cardé, 1991. Identification of sex pheromone of browntail moth, *Euproctis chrysorrhoea* (L.) (Lepidoptera: Lymantriidae). *J. Chem. Ecol.*, 17: 897-910.

- Leskey**, T. C., J. C. Bergh, J. F. Walgenbach and A. J. Zhang, 2006. Attractiveness and specificity of pheromone-baited traps for male dogwood borer, *Synanthedon scitula* Harris (Lepidoptera: Sesiidae). *Environ. Entomol.*, 35: 268-275.
- Levi-Zada**, A., D. Fefer, L. Anshelevitch, A. Litovsky, M. Bengtsson, G. Gindin and V. Soroker, 2011(a). Identification of the sex pheromone of the lesser date moth, *Batrachedra amydraula*, using sequential SPME auto-sampling. *Tetrahedron Lett.*, 52: 4550-4553.
- Levi-Zada, A., S. Ben-Yehuda, E. Dunkelblum, G. Gindin, D. Fefer, A. Protasov, T. Kuznetsowa, S. Manulis-Sasson and Z. Mendel, 2011(b). Identification and field bioassays of the sex pheromone of the yellow-legged clearwing *Synanthedon vespiformis* (Lepidoptera: Sesiidae). *Chemoecology*, 21: 227-233.
- Levi-Zada, A., A. Sadowsky, S. Dobrinin, M. David, T. Ticuchinski, D. Fefer, A. Greenberg and D. Blumberg, 2013. Reevaluation of the sex pheromone of the lesser date moth, *Batrachedra amydraula*, using autosampling SPME-GC/MS and field bioassays. *Chemoecology*, 23: 13-20.
- Lewis**, T., C. Wall, E. D. M. Macaulay and A. R. Greenway, 1975. The behavioural basis of a pheromone monitoring system for pea moth, *Cydia nigricana*. *Ann. Appl. Biol.*, 80: 257-274.
- Li**, J., R. Gries, G. Gries, K. N. Slessor, G. G. S. King, W. W. Bowers and R. J. West, 1993(a). Chirality of 5,11-dimethylheptadecane, the major sex pheromone component of the hemlock looper, *Lambdina fiscellaria* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 19: 1057-1062.
- Li, J.-X., G. Gries, R. Gries, J. Bikic and K. N. Slessor, 1993(b). Chirality of synergistic sex pheromone components of the western hemlock looper *Lambdina fiscellaria lugubrosa* (Hulst) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 19: 2547-2561.
- Li, Z.-M., E.-Y. Yao, T.-L. Liu, Z.-P. Liu, S.-H. Wang, H.-Q. Zhu, G. Zhao and Z.-L. Ren, 1993(c). Structural elucidation of sex pheromone components of the Geometridae *Semiothisa cinerearia* (Bremer et Grey) in China. *Chinese J. Chem.*, 11: 251-256.
- Li, W.-W., X.-Z. Meng, Y.-H. Deng and X.-G. Hu, 1993(d). Experimental effects on field trapping of noctuids using several sex attractants. *Entomol. Knowledge*, 30: 38-40.
- Liebold**, A. M. and W. J. A. Volney, 1985. Effects of attractant composition and release rate on attraction of male *Choristoneura retiniana*, *C. occidentalis*, and *C. carana* (Lepidoptera: Tortricidae). *Can. Entomol.*, 117: 447-457.
- Liebold, A. M. and P. J. Silk, 1991. Capture of *Choristoneura pinus maritima* in traps baited with *C. pinus* pinus pheromone components (Tortricidae). *J. Lepidop. Soc.*, 45: 172-173.

- Lin**, Y. M., Y. S. Chow and H. C. Tzeng, 1982. Field trapping of the diamondback moth *Plutella xylostella* (Linnaeus) and *Pseudaletia separata* Walker using the synthetic sex pheromone of the diamondback moth pest of cruciferous vegetables, Taiwan. *Bull. Inst. Zool. Acad. Sin.*, 21: 121-127.
- Lin**, G.-Q., G.-Z. Guo, Y.-W. Wu, B.-Q. Wu, L.-G. Liu and W.-S. Zhou, 1984. Study on identification and syntheses of insect pheromone XV. Multiple sex pheromone of *Ancylis sativa*. *Kexue Tongbao*, 29: 1670-1674.
- Lin**, G.-Q., 1993. Structural elucidation and chiral syntheses of insect pheromones and extension of the synthetic approaches. *Pure Appl. Chem.*, 65: 1233-1240.
- Linn**, C. E. and W. L. Roelofs, 1983. Effect of varying proportions of the alcohol component on sex pheromone blend discrimination in male Oriental fruit moths. *Physiol. Entomol.*, 8: 291-306.
- Linn**, C. E. Jr., L. B. Bjostad, J. W. Du and W. L. Roelofs, 1984. Redundancy in a chemical signal: Behavioral responses of male *Trichoplusia ni* to a 6-component sex pheromone blend. *J. Chem. Ecol.*, 10: 1635-1658.
- Linn**, C. E., Jr., J. Du, A. Hammond and W. L. Roelofs, 1987. Identification of unique pheromone components for soybean looper moth *Pseudoplusia includens*. *J. Chem. Ecol.*, 13: 1351-1360.
- Linnaluoto**, E. T. and S. Koponen, 1983. Northern moths (Lepidoptera) attracted by synthetic pheromones. *Ann. Entomol. Fenn.*, 49: 64.
- Liska**, J., R. Hochmut and J. Jurik, 1990. Field trials with sexual attractants of *Zeiraphera griseana* (Lepidoptera, Tortricidae) in the High Tatras. *Acta Entomol. Bohemoslov.*, 87: 81-86.
- Liu**, Y.-X. and X.-Z. Meng, 2002(a). Identification of sex pheromone components for *Acleris fimbriana* (Lepidoptera: Tortricidae). *Can. Entomol.*, 134: 511-518.
- Liu**, Y.-X. and X.-Z. Meng, 2002(b). Behavioral response of male *Acleris fimbriana* Thunberg et Becklin (Lepidoptera: Tortricidae) to synthetic sex pheromones. *Acta Entomol. Sinica (Kunchong Xuebao)*, 45: 436-440.
- Liu**, Y.-X. and X.-Z. Meng, 2003. Trapping effect of synthetic sex pheromone of *Acleris fimbriana* (Lepidoptera: Tortricidae) in Chinese northern orchard. *Z. Naturforsch.*, 58c: 421-425.
- Liu**, F., L. Guo, Q.-C. Li, S.-F. Zhang, X.-B. Kong, Z. Zhang and F.-Q. Huang, 2019. Identification of the active components of sex pheromone of female moths of *Micromelalopha sieversi* (Lepidoptera: Notodontidae). *Acta Entomol. Sinica*, 62: 814-822.
- Liu**, F., L. Guo, S. Zhang, X. Kong and Z. Zhang, 2021. Synthesis and bioactivity of (13Z,15E)-octadecadienal: a sex pheromone

component from *Micromelalopha siversi* Staudinger (Lepidoptera: Notodontidae). *Pest Manag. Sci.*, **77**: 264-272.

Liu, H., R. Dou, J. Wang, Y. Zhu, J. Li, J. Liu, Y. Wang, J. Zhang and X. Jing, 2022. Evaluation of sex attractants for monitoring of *Illiberis ulmivora* Graeser (Lepidoptera: Zygaenidae: Procridinae): height, distance, placement, climatic conditions. *Entomol. Res.*, **52**: 99-107.

Löfstedt, C., J. N. C. VanDerPers, J. Lofqvist, B. S. Lanne, M. Appelgren, G. Bergström and B. Thelin, 1982. Sex pheromone components of the turnip moth, *Agrotis segetum*: chemical identification, electrophysiological evaluation and behavioral activity. *J. Chem. Ecol.*, **8**: 1305-1321.

Löfstedt, C., J. N. C. Van der Pers, J. Löfqvist and B. S. Lanne, 1983. Sex pheromone of the cone pyralid *Dioryctria abietella*. *Entomol. Exp. Appl.*, **34**: 20-26.

Löfstedt, C. and J. N. C. van der Pers, 1985(a). Sex pheromones and reproductive isolation in four European small ermine moths. *J. Chem. Ecol.*, **11**: 649-666.

Löfstedt, C., B. S. Lanne, J. Löfqvist, M. Appelgren and G. Bergström, 1985(b). Individual variation in the pheromone of the turnip moth, *Agrotis segetum*. *J. Chem. Ecol.*, **11**: 1181-1196.

Löfstedt, C., C. E. Linn, Jr. and J. Löfqvist, 1985(c). Behavioral responses of male turnip moths, *Agrotis segetum*, to sex pheromone in a flight tunnel and in the field. *J. Chem. Ecol.*, **11**: 1209-1221.

Löfstedt, C., W. M. Herrebout and J.-W. Du, 1986(a). Evolution of the ermine moth pheromone tetradecyl acetate. *Nature*, **323**: 621-623.

Löfstedt, C., J. Löfqvist, B. S. Lanne, J. N. C. Van Der Pers and B. S. Hansson, 1986(b). Pheromone dialects in European turnip moths *Agrotis segetum*. *Oikos*, **46**: 250-257.

Löfstedt, C. and W. M. Herrebout, 1988. Sex pheromones of three small ermine moths found on the European spindle tree. *Entomol. Exp. Appl.*, **46**: 29-38.

Löfstedt, C., B. S. Hansson, H. J. Dijkerman and W. M. Herrebout, 1990. Behavioural and electrophysiological activity of unsaturated analogues of the pheromone tetradecyl acetate in the small ermine moth *Yponomeuta rorellus*. *Physiol. Entomol.*, **15**: 47-54.

Löfstedt, C., B. S. Hansson, M. Tóth, G. Szöcs, V. Buda, M. Bengtsson, N. Ryholm, M. Svensson and E. Priesner, 1994. Pheromone differences between sibling taxa *Diachrysia chrysitis* (Linnaeus, 1758) and *D. tutti* (Kostrowicki, 1961) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **20**: 91-109.

Löfstedt, C., J. Zhu, M. V. Kozlov, V. Buda, E. V. Jirle, S. Hellqvist, J. Löfqvist, E. Plass, S. Franke and W. Francke, 2004.

Identification of the sex pheromone of the currant shoot borer *Lampronia capitella*. *J. Chem. Ecol.*, **30**(3): 643-658.

Löfstedt, C., G. P. Svensson, E. V. Jirle, O. Rosenberg, A. Roques and J. G. Millar, 2012. (3Z,6Z,9Z,12Z,15Z)-pentacosapentaene and (9Z,11E)-tetradecadienyl acetate: sex pheromone of the spruce coneworm *Dioryctria abietella* (Lepidoptera: Pyralidae). *J. Appl. Entomol.*, **136**: 70-78.

Lonergan, G. C., B. M. Ponder, W. D. Seabrook and L. R. Kipp, 1989. Sex pheromone components of the blueberry leaf tier moth *Croesia curvalana* Kearfott (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **15**: 2495-2506.

Lopez, J. D., Jr. and T. N. Shaver, 1990. Plastic laminate dispenser for components of the *Heliothis zea* sex pheromone. *Southwest. Entomol.*, **15**(1): 1-8.

Lu, P.-F., H.-L. Qiao and Y.-Q Luo, 2013. Female sex pheromone blends and male response of the legume pod borer, *Maruca vitrata* (Lepidoptera: Crambidae), in two populations of mainland China. *Z. Naturforsch.*, **68 c**: 416-427.

Luo, Z.-X., Z.-Q. Li, X.-M. Cai, L. Bian and Z. M. Chen, 2017. Evidence of pre-mating isolation between two sibling moths: *Ectropis griseascens* and *Ectropis obliqua* (Lepidoptera: Geometridae). *J. Econ. Entomol.*, **110**: 2364-2370.

Luo, Z., X. Cai, Z. Li, L. Bian, Z. Xin, F. Hussain and Z. Chen, 2019. Identification of sex pheromone of *Miltochrista striata* (Lepidoptera: Arctiidae). *J. Econ. Entomol.*, **112**: 665-670.

Lynch, R. E., J. A. Klun, B. A. Leonhardt, M. Schwarz and J. W. Garner, 1984. Female sex pheromone of the lesser cornstalk borer, *Elasmopalpus lignosellus* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **13**: 121-126.

Ma, T., Y. Li, Z. Sun and X. Wen, 2014. (Z,E)-9,12-Tetradecadien-1-ol: A major sex pheromone component of *Euzophera pyriella* (Lepidoptera: Pyralidae) in Xinjiang, China. *Florida Entomol.*, **97**: 496-503.

Ma, T., Z. Liu, J. Lu, Z. Sun, Y. Li, X. Wen and Y. Cui, 2015. A key compound: (Z)-9-tetradecen-1-ol as sex pheromone active component of *Hypsipyla robusta* (Lepidoptera: Pyralidae). *Chemoecology*, **25**: 325-330.

Ma, T., Q. Xiao, Y.-G. Yu, C. Wang, C.-Q. Zhu, Z.-H. Sun, X.-Y. Chen and X.-J. Wen, 2016. Analysis of tea geometrid (*Ectropis griseascens*) pheromone gland extracts using GC-EAD and GCxGC/TOFMS. *J. Agric. Food Chem.*, **64**: 3161-3166.

Ma, T., Z. Liu, C. Wang, S. Zhang, X. Shi, Z. Sun, X. Chen, C. Jia, C. Wang, Y. He and X. Wen, 2017. Production, identification,

and field evaluation of sex pheromone from calling females in *Diaphania angustalis* (Lepidoptera: Crambidae). *Environ. Sci. Pollut. Res. Int.*, **24**: 24485-24493.

Mafi, S. A., L. V. Vang, Y. Nakata, N. Ohbayashi, M. Yamamoto and T. Ando, 2005. Identification of the sex pheromone of the citrus leafminer (*Phyllocnistis citrella* Stainton, Lepidoptera: Gracillariidae) with a trial of control by the communication disruption method. *J. Pestic. Sci.*, **30**(4): 361-367.

Magsi, F. H., Z. Li, X. Cai, M. Yamamoto, L. Bian, Y. Zhao, L. Zhou, C. Xiu, N. Fu, T. Ando, Z. Luo and Z. Chen, 2022. Identification of a unique three-component sex pheromone produced by the tea black tussock moth, *Dasychira baibarana* (Lepidoptera: Erebidae: Lymantriinae). *Pest Manag. Sci.*, **78**: 2607-2617.

Maier, C. T., R. Gries and G. Gries 1998. Sex pheromone components of pitch pine looper, *Lambdina pellucidaria*. *J. Chem. Ecol.*, **24**(3): 491-500.

Maini, S., 1973. Preliminary tests with synthetic sex attractants of the European corn borer. *Inf. Tore Fitopatol.*, **9**: 11-14.

Malo, E. A., J. C. Rojas, G. Lopez-Guillen and J. F. Barrera, 2009. Chemical analysis of female volatiles and field response of the coffee leafminer moth (Lepidoptera: Lyonetiidae) to stereoisomers of its major sex pheromone component. *Florida Entomol.*, **92**: 548-553.

Malosse, C., C. Descoins and B. Frerot, 1993. Identification of minor component of the sex pheromone of the leopard moth (*Zeuzera pyrina* L.; Lepidoptera, Cossidae) by capillary gas chromatography. *J. High Resol. Chromat.*, **16**: 123-125.

Matsuoka, K., M. Yamamoto, R. Yamakawa, M. Muramatsu, H. Naka, Y. Kondo and T. Ando, 2008. Identification of novel C₂₀ and C₂₂ trienoic acids from arctiid and geometrid female moths that produce polyenyl Type II sex pheromone components. *J. Chem. Ecol.*, **34**: 1437-1445.

Masnyk, M., J. Fried and W. L. Roelofs, 1989. Synthesis and pheromone properties of (Z)-7,7-difluoro-8-docecetyl acetate, a difluoro derivative of the sex pheromone of the Oriental fruit moth. *Tetrahedron Lett.*, **30**: 3243-3246.

Mayer, M. S., J. R. McLaughlin and E. R. Mitchell, 1995. Use of paired and single traps to assess perception and discrimination of sex pheromone mixtures in the field by *Trichoplusia ni* (Hubner). *J. Chem. Ecol.*, **21**: 2057-2068.

Mazomenos, B. E., D. Bardas and A. Pantazimazomenou, 1989(a). Sesamianonagrioides sex pheromone: field attractiveness use of pheromone traps for monitoring and control. *Acta Phytopathol. Entomol. Hung.*, **24**: 153-156.

Mazomenos, B. E., 1989(b). Sex pheromone components of corn stalk borer *Sesamia nonagrioides* (Lef.). Isolation, identification, and

field tests. *J. Chem. Ecol.*, **15**: 1241-1247.

Mazomenos, B. E., D. Raptopoulos, I. Lefkidou and A. Mazomenos-Pantazi, 1994. Female sex pheromone components of jasmine moth *Palpita unionalis* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **20**: 745-751.

Mazor, M., M. Harel, S. Levsky and E. Dunkelblum, 1991. Sex pheromone components of female *Cornutiplusia circumflexa*. *Entomol. Exp. Appl.*, **60**: 167-172.

Mazor, M. and E. Dunkelblum 1992. Role of sex pheromone components in behavioral reproductive isolation between *Autographa gamma* (L.) and either *Trichoplusiani* (Hubner) or *Chrysodeixis chalcites* (Esp.) (Lepidoptera: Noctuidae: Plusinae). *J. Chem. Ecol.*, **18**(12): 2373-2384.

McBrien, H., G. Gries, R. Gries, J. H. Borden, G. J. R. Judd, G. G. S. King and K. N. Slessor, 1992. Sex pheromone components of the eyespotted bud moth, *Spilonota ocellana* (Denis and Schiffermüller) (Lepidoptera: Olethreutidae). *Can. Entomol.*, **124**: 1391-1394.

McBrien, H. L., G. J. R. Judd, A. K. Minks, J. W. Johnson, R. F. Smith, H. M. A. Thistlewood and J. T. Troubridge, 1994. Response of male eye-spotted bud moth, *Spilonota ocellana* (Lepidoptera: Tortricidae), to different pheromone blends in North America and the Netherlands. *J. Chem. Ecol.*, **20**: 625-630.

McDonough, L. M. and J. A. Kamm, 1979. Sex pheromone of the cranberry girdler, *Chrysoteuchia topiaria* (Zeller) (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **5**: 211-219.

McDonough, L. M., J. A. Kamm and B. A. Bierl-Leonhardt, 1980. Sex pheromone of the armyworm, *Pseudaletia unipuncta* (Haworth) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **6**: 565-572.

McDonough, L. M., M. P. Hoffmann, B. A. Bierl-Leonhardt, C. L. Smithhisler, J. B. Bailey and H. G. Davis, 1982(a). Sex pheromone of the avocado pest, *Amorbia cuneana* (Walsingham) (Lepidoptera: Tortricidae): structure and synthesis. *J. Chem. Ecol.*, **8**: 255-265.

McDonough, L. M., D. S. Moreno, B. A. Bierl-Leonhardt and B. A. Butt, 1982(b). Isolation and identification of a component of the female sex pheromone gland attractive to male *Xylomyges curialis*. *Environ. Entomol.*, **11**: 660-662.

McDonough, L. M., J. A. Kamm, D. A. George, C. L. Smithhisler and S. Voerman, 1982(c). Sex attractant for the western lawn moth, *Tehama bonifatella* Hulst. *Environ. Entomol.*, **11**: 711-714.

McDonough, L. M., J. B. Bailey, M. P. Hoffmann, B. A. Leonhardt, D. F. Brown, C. L. Smithhisler and K. Olsen, 1986. *Sabulodes caberata* Guenée (Lepidoptera: Geometridae). Components of its sex pheromone gland. *J. Chem. Ecol.*, **12**: 2107-2116.

McDonough, L. M., H. G. Davis and S. Voerman, 1987. Blackheaded

fireworm: laboratory and field studies of its sex pheromone. *J. Chem. Ecol.*, **13**: 1235-1242.

McDonough, L. M., R. D. Eikenbary, M. T. Smith, H. G. Davis, C. L. Smithhisler, G. A. Hedger, R. K. Campbell, J. A. Payne, W. Reid and J. R. McVay, 1990(a). Sex pheromone of hickory shuckworm *Cydia caryana*. Development of an effective field lure. *J. Chem. Ecol.*, **16**: 317-324.

McDonough, L. M., H. G. Davis, C. L. Smithhisler, S. Voerman and P. S. Chapman, 1990(b). Apple ermine moth, *Yponomeuta malinellus* Zeller. Two components of female sex pheromone gland highly effective in field trapping tests. *J. Chem. Ecol.*, **16**: 477-486.

McDonough, L. M., H. G. Davis, P. S. Chapman and C. L. Smithhisler, 1993. Response of male codling moths (*Cydia pomonella*) to components of conspecific female sex pheromone glands in flight tunnel tests. *J. Chem. Ecol.*, **19**: 1737-1748.

McDonough, L. M., A. L. Averill, H. G. Davis, C. L. Smithhisler, D. A. Murray, P. S. Chapman, S. Voerman, L. J. Dapsis and M. M. Averill, 1994. Sex pheromone of cranberry fruitworm, *Acrobasis vaccinii* Riley (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **20**: 3269-3279.

McDonough, L. M., H. G. Davis, P. S. Chapman and C. L. Smithhisler, 1995. Codling moth, *Cydia pomonella*, (Lepidoptera: Tortricidae): is its sex pheromone multicomponent? *J. Chem. Ecol.*, **21**: 1065-1071.

McElfresh, J. S. and J. G. Millar 1999(a). Sex pheromone of the common sheep moth, *Hemileuca eglanterina*, from the San Gabriel Mountains of California. *J. Chem. Ecol.*, **25**(4): 687-709.

McElfresh, J. S. and J. G. Millar 1999(b). Sex pheromone of Nuttall's sheep moth, *Hemileuca nuttalli*, from the Eastern Sierra Nevada Mountains of California. *J. Chem. Ecol.*, **25**(4): 711-726.

McElfresh, J. S. and J. G. Millar 1999(c). Sex attractant pheromone of saturniid moth, *Coloradia velda*. *J. Chem. Ecol.*, **25**(5): 1067-1078.

McElfresh, J. S. and J. G. Millar, 1999(d). Geographic variation in sex pheromone blend of *Hemileuca electra* from southern California. *J. Chem. Ecol.*, **25**(11): 2505-2525.

McElfresh, J. S., X. Chen, D. W. Ross and J. G. Millar, 2000. Sex pheromone blend of the pandora moth (Lepidoptera: Saturniidae), an outbreak pest in pine forests (Pinaceae). *Can. Entomol.*, **132**: 775-787.

McElfresh, J. S., J. G. Millar and D. Rubinoff, 2001(a). (E4, Z9)-Tetradecadienal, a sex pheromone for three north American moth species in the genus *Saturnia*. *J. Chem. Ecol.*, **27**(4): 791-806.

McElfresh, J. S., A. M. Hammond and J. G. Millar, 2001 (b). Sex pheromone components of the buck moth *Hemileuca maia*. *J. Chem. Ecol.*, 27 (7): 1409-1422.

McElfresh, J. S. and J. G. Millar, 2008. Sex pheromone of the saturniid moth, *Hemileuca burnsi*, from the western Mojave Desert of California. *J. Chem. Ecol.*, 34 (9): 1115-1124.

McElfresh, J. S., J. A. Moreira, E. E. Grafton-Cardwell, D. H. Headrick, J. M. Heraty, M. Guillen and J. G. Millar, 2009. Novel lepidopteran sex pheromone components from *Marmara gulosa* (Lepidoptera: Gracillariidae). *J. Econ. Entomol.*, 102: 574-584.

McLaren, G. F. and D. M. Suckling, 1993. Pheromone trapping of orchard lepidopterous pests in Central Otago, New Zealand. *New Zealand J. Crop Horticul. Sci.*, 21: 25-31.

McLaughlin, J. R. and R. R. Heath, 1989. Field trapping and observations of male velvetbean caterpillar moths and trapping of *Mocis* spp. (Lepidoptera: Noctuidae: Catacolinae) with calibrated formulations of sex pheromone. *Environ. Entomol.*, 18: 933-938.

McLeod, D. G. R., T. Nagai, A. N. Starratt, C. Bonenfant, E. W. Rud and S. Belloncik, 1982. Field and laboratory evaluation of a sex attractant for the white cutworm, *Euxoa scandens* (Lepidoptera: Noctuidae). *Can. Entomol.*, 114: 1101-1103.

McNair, C., G. Gries, and R. Gries 1999. Sex pheromone components of *Enarmonia formosana* (Lepidoptera: Tortricidae). *Can. Entomol.* 131: 85-92.

McQuate, G. T., A. Cossé, C. D. Sylva and J. A. MacKay, 2019. Field evaluation of a binary sex pheromone for sweetpotato vine borer (Lepidoptera: Crambidae) in Hawaii. *J. Insect Sci.*, 19: iez008.

Meijer, G. M., F. J. Ritter, C. J. Persoons, A. K. Minks and S. Voerman, 1972. Sex pheromone of summer fruit tortrix moth *Adoxophyes orana*: two synergistic isomers. *Science*, 175: 1469-1470.

Meng, X.-Z., 1983 (a). Synthesis of the three active components of the sex pheromone of the masson pine caterpillar moth - (*Z,E*)-5,7-dodeca-dien-1-ol, acetate and propionate. *Kexue Tongbao*, 28: 141.

Meng, X.-Z. and H.-M. Wang, 1983 (b). Synthesis and field experiment of sex attractant for pine caterpillar, *Dendrolimus tabulaeformis*. *Scientia Silvae Sinicae*, 19: 137-140.

Meyer, W. L., G. L. Debarr, C. W. Berisford, L. R. Barber and W. L. Roelofs, 1982. Identification of the sex pheromone of the webbing coneworm moth, *Dioryctria disclusa* (Lepidoptera: Pyralidae). *Environ. Entomol.*, 11: 986-988.

Meyer, W. L., R. S. Cameron, A. Tamhankar, G. DeBarr, C. W. Berisford and W. L. Roelofs, 1984. Sex pheromone of the blister coneworm moth, *Dioryctria clarioralis* (Lepidoptera: Pyralidae).

Environ. Entomol., 13: 854-858.

Meyer, W. L., G. L. Debarr, J. L. Hanula, B. Kovalev, R. S. Cameron, C. W. Berisford and W. L. Roelofs, 1986. Z-11-Hexadecenyl acetate, a sex pheromone component for the southern pine coneworm, *Dioryctria amatella* (Lepidoptera: Pyralidae). *Environ. Entomol.*, 15: 316-320.

Meyer, W. L., W. S. Cranshaw and T. D. Eichlin, 1988. Flight patterns of clearwing borers in Colorado based on pheromone trap captures. *Southwest. Entomol.*, 13: 39-46.

Meyer, W. L., and Cranshaw, W. S. 1994. Capture of clearwing borers (Lepidoptera: Sesiidae) with three synthetic attractants in Colorado. *Southwest Entomol.*, 19: 71-76.

Millar, J. G., E. W. Underhill, M. Giblin and D. Barton, 1987. Sex pheromone components of three species of *Semiothisa* (Geometridae), (*Z,Z,Z*)-3,6,9-heptadecatriene and two monoepoxydiene analogs. *J. Chem. Ecol.*, 13: 1371-1383.

Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1990 (a). (*3Z,6Z,9Z*)-Nonadecatriene and enantiomers of (*3Z,9Z*)-*cis*-6,7-epoxy-nonadecadiene as sex attractants for two geometrid and one noctuid moth species. *J. Chem. Ecol.*, 16: 2153-2166.

Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1990 (b). *3Z,6Z,9Z*-Trienes and unsaturated epoxides as sex attractants for geometrid moths. *J. Chem. Ecol.*, 16: 2307-2316.

Millar, J. G., M. Giblin, D. Barton, A. Morrison and E. W. Underhill, 1990 (c). Synthesis and field testing of enantiomers of *6Z,9Z-cis*-3,4-epoxydienes as sex attractants for geometrid moths. *J. Chem. Ecol.*, 16: 2317-2339.

Millar, J. G., M. Giblin, D. Barton, D. A. Reynard, G. B. Neill and E. W. Underhill, 1990 (d). Identification and field testing of female-produced sex pheromone components of the spring cankerworm, *Paleacrita vernata* Peck (Lepidoptera: Geometridae). *J. Chem. Ecol.*, 16: 3393-3409.

Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1991 (a). Synthesis and field screening of chiral monounsaturated epoxides as lepidopteran sex attractants and sex pheromone components. *J. Chem. Ecol.*, 17: 911-929.

Millar, J. G., M. Giblin, D. Barton, J. W. Wong and E. W. Underhill, 1991 (b). Sex attractants and sex pheromone components of noctuid moths *Euclidea cuspidea*, *Caenurgina distincta*, and geometrid moth *Eupithecia annulata*. *J. Chem. Ecol.*, 17: 2095-2111.

Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1991 (c). Chiral lepidopteran sex attractants: blends of optically active C_{20} and C_{21} diene epoxides as sex attractants for geometrid and noctuid moths (Lepidoptera). *Environ. Entomol.*, 20(2): 450-457.

- Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1992(a). Sex pheromone components of the geometrid moths *Lobophora nivigerata* and *Epirrhoë sperryi*. *J. Chem. Ecol.*, **18**: 1057-1068.
- Millar, J. and R. E. Rice, 1992(b). Reexamination of the female sex pheromone of the peach twig borer: field screening of minor constituents of pheromone gland extracts and of pheromone analogs. *J. Econ. Entomol.*, **85**: 1709-1716.
- Millar, J. G., A. E. Knudson, J. S. McElfresh, R. Gries, G. Gries and J. H. Davis, 1996. Sex attractant pheromone of the pecan nut casebearer (Lepidoptera: Pyralidae). *Bioorg. Medic. Chem.*, **4**: 331-339.
- Millar, J., J. S. McElfresh and F. de A. Marques, 2002. Unusual acetylenic sex pheromone of grape leaffolder (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **95**(4): 692-698.
- Millar, J. G., G. G. Grant, J. S. McElfresh, W. Strong, C. Rudolph, J. D. Stein and J. A. Moreira, 2005. (3Z,6Z,9Z,12Z,15Z)-Pentacosapentaene, a key pheromone component of the fir coneworm moth, *Dioryctria abietivorella*. *J. Chem. Ecol.*, **31**(5): 1229-1234.
- Millar, J. G., M. Hoddle, J. S. McElfresh, Y. Zou and C. Hoddle, 2008. (9Z)-9,13-Tetradecadien-11-ynal, the sex pheromone of the avocado seed moth, *Stenoma catenifer*. *Tetrahedron Lett.*, **49**: 4820-4823.
- Millar, J. G., J. S. McElfresh, C. Romero, M. Vila, N. Mari-Mena and C. Lopez-Vaamonde, 2010. Identification of the sex pheromone of a protected species, the Spanish moon moth *Graellsia isabellae*. *J. Chem. Ecol.*, **36**: 923-932.
- Millar, J. G., K. F. Haynes, A. T. Dossey, J. S. McElfresh and J. D. Allison, 2016. Sex attractant pheromone of the luna moth, *Actias luna* (Linnaeus). *J. Chem. Ecol.*, **42**: 869-876.
- Miller**, J. R., T. C. Baker, R. T. Cardé and W. L. Roelofs, 1976. Reinvestigation of oak leaf roller sex pheromone components and the hypothesis that they vary with diet. *Science*, **192**: 140-143.
- Miller, J. R. and W. L. Roelofs, 1978. Gypsy moth responses to pheromone enantiomers as evaluated in a sustained-flight tunnel. *Environ. Entomol.*, **7**: 42-44.
- Miller, D. R., J. G. Millar, A. Mangini, C. M. Crowe and G. G. Grant, 2010. (3Z,6Z,9Z,12Z,15Z)-Pentacosapentaene and (Z)-11-hexadecenyl acetate: sex attractant blend for *Dioryctria amatella* (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **103**: 1216-1221.
- Minks**, A. K. and S. Voerman, 1973(a). Sex pheromone of the summerfruit tortrix moth, *Adoxophyes orana*: trapping performance in the field. *Entomol. Exp. Appl.*, **16**: 541-549.
- Minks, A. K., W. L. Roelofs, F. J. Ritter and C. J. Persoons, 1973(b).

Reproductive isolation of two tortricid moth species by different ratios of a two-component sex attractant. *Science*, **180**: 1073-1074.

Minks, A. K., S. Voerman and M. Van de Vrie, 1976. A sex attractant for the cherry-bark tortrix moth, *Enarmonia formosana*. *Entomol. Exp. Appl.*, **19**: 301-302.

Minks, A. K., S. Voerman and J. Theunissen, 1994. Improved sex attractant for the leek moth, *Acrolepiopsis assectella* Zell. (Lep., Acrolepiidae). *J. Appl. Entomol.*, **117**: 243-247.

Minyailo, V. A., B. G. Kovalev, E. I. Kirov and A. K. Minyailo, 1977. On the attractiveness of disparlure, sex pheromone of the *Porthetria dispar* females, for the *Zanclognatha lunaris* males. *Zool. Zh.*, **56**: 309-310.

Mitchell, E. R., J. C. Webb, A. H. Baumhover, R. W. Hines, J. W. Stanley, R. G. Endris, D. A. Lindquist and S. Masuda, 1972. Evaluation of cylindrical electric grids as pheromone traps for loopers and tobacco hornworms. *Environ. Entomol.*, **1**: 365-368.

Mitchell, E. R. and J. H. Tumlinson, 1973 (a). An attractant for males of *Spodoptera dolichos* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **66**: 917-918.

Mitchell, 1973 (b). Nocturnal activity of adults of three species of loopers, based on collections in pheromone traps. *Environ. Entomol.*, **2**: 1078-1080.

Mitchell, E. R. and R. E. Doolittle, 1976. Sex pheromones of *Spodoptera exigua*, *S. eridania*, and *S. frugiperda*: bioassay for field activity. *J. Econ. Entomol.*, **69**: 324-326.

Mitchell, E. R., J. H. Tumlinson and A. H. Baumhover, 1978. *Heliothis virescens*: attraction of males to blends of (Z)-9-tetradecen-1-ol formate and (Z)-9-tetradecenal. *J. Chem. Ecol.*, **4**: 709-716.

Mitchell, E. R., H. Sugie and J. H. Tumlinson, 1983. *Spodoptera exigua*: capture of feral males in traps baited with blends of pheromone components. *J. Chem. Ecol.*, **9**: 95-104.

Mitchell, E. R., J. H. Tumlinson and J. N. McNeil, 1985. Field evaluation of commercial pheromone formulations and traps using a more effective sex pheromone blend for the fall armyworm (Lepidoptera: Noctuidae). *J. Econ. Entomol.*, **78**: 1364-1369.

Mitchell, E. R. and J. H. Tumlinson, 1994. Response of *Spodoptera exigua* and *S. eridania* (Lepidoptera: Noctuidae) males to synthetic pheromone and *S. exigua* females. *Florida Entomol.*, **77**(2): 237-247.

Mizukoshi, T., 2006. Non-target species caught by synthetic sex pheromone trap of lepidotriterous pests on apple trees in Hokkaido. *Jpn. J. Appl. Entomol. Zool.*, **50** (3): 231-239.

Mochida, O., G. S. Arida, S. Tatsuki and J. Fukami, 1984. A field

test on a third component of the female sex pheromone of the rice striped stem borer, *Chilo suppressalis*, in the Philippines. *Entomol. Exp. Appl.*, **36**: 295-296.

Mochizuki, F., T. Shibuya, T. Ihara and S. Wakamura, 1993. Electrophysiological responses of the male antenna to compounds found in the female sex pheromone gland of *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **28**: 489-496.

Mochizuki, F., M. Takai, T. Shibuya, and S. Wakamura, 1994. Field trap responses of male *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) to (Z,E)-9,12-tetradecadien-1-ol and (Z)-9-tetradecenyl acetate. *Appl. Entomol. Zool.*, **29**: 436-438.

Mochizuki, F. and K. Fukumoto, 2003. Insect attractants for *Grapholita dimorpha*. *Jpn. Kokai Tokkyo Koho*, JP 2003-238311 [2002-42519].

Mochizuki, F. and Y. Yumoto, 2009(a). Hexadecenal mixture-containing sex pheromones of *Pleuroptya ruralis*, and their use as attractants. *Jpn. Kokai Tokkyo Koho* JP 2009018994. [SciFinder. AN 2009:111797]

Mochizuki, F. Y. Aoki, H. Sugie, J. Tabata, A. Takahashi and H. Higuchi, 2009(b). Sex attractant of the bean webworm, *Pleuroptya ruralis* (Crambidae: Pyraustinae). *Jpn. J. Appl. Entomol. Zool.*, **53**: 53-56.

Molnár, B. P., A. Tröger, T. B. Toshova, M. Subchev, E. J. van Nieukerken, J. C. Koster, G. Szöcs, M. Tóth and W. Francke, 2012. Identification of the female-produced sex pheromone of *Tischeria ekebladella*, an oak leafmining moth. *J. Chem. Ecol.*, **38**: 1298-1305.

Molnár, P. B., C. Bognár, A. L. Erdei, T. Fujii, P. Vági, J. K. Jósvai and Z. Kárpáti 2018. Identification of the female-produced sex pheromone of an invasive greenhouse pest, the European pepper moth (*Duponchelia fovealis*) *J. Chem. Ecol.*, **44**: 257-267.

Monti, L., B. Lalanne-Cassou, P. Lucas, C. Malosse and J.-F. Silvain, 1995. Differences in sex pheromone communication systems of closely related species: *Spodoptera latifascia* (Walker) and *S. descoinsi* Lalanne-Cassou & Silvain (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **21**: 641-660.

Moreira, J. A., J. S. McElfresh and J. G. Millar, 2006. Identification, synthesis, and field testing of the sex pheromone of the citrus leafminer, *Phyllocnistis citrella*. *J. Chem. Ecol.*, **32**(1): 169-194.

Mori, K., H. Harada, P. Zagatti, A. Cork and D. R. Hall, 1991. Synthesis and biological activity of four stereoisomers of 6,10,14-trimethyl-2-pentadecanol, the female-produced sex pheromone of rice moth (*Coryza cephalonica*). *Liebigs Ann. Chem.*, 259-267.

Mori, K., T. Tashiro, B. Zhao, D. M. Suckling and A. M. El-Sayed, 2010. Pheromone synthesis. Part 243: Synthesis and biological

evaluation of (*3R,13R,1'S*)-1'-ethyl-2'-methylpropyl 3,13-dimethylpentadecanoate, the major component of the sex pheromone of Paulownia bagworm, *Clania variegate*, and its stereoisomers. *Tetrahedron*, **66**: 2642-2653.

Mõttus, E., Z. Klukowski, I. Liblikas, M. Stachowiak and J. Viidalepp, 1998. Sex attractant dispensers for monitoring antler moth *Cerapteryx graminis* (L.) (Lep., Noctuidae): Effect of component ratio and dispenser type. *J. Appl. Entomol.*, **122**: 193-200.

Mozūraitis, R., V. Buda, A.-K. Borg-Karlson, P. Ivinskis, V. Karalius, M. Laanmaa and D. Pleys 1998. New sex attractants and inhibitors for 17 moth species from the families Gracillariidae, Tortricidae, Yponomeutidae, Oecophoridae, Pyralidae and Gelechiidae. *J. Appl. Entomol.*, **122**: 441-452.

Mozūraitis, R., V. Buda and J. Metleuski 1999 (a). New sex attractants for five *Chamaesphecia* species (Lepidoptera, Sesiidae) from the Ukraine and Turkmenistan. *Z. Naturforsch.*, **54c**: 253-258.

Mozūraitis, R., A.-K. Borg-Karlson, V. Buda and P. Ivinskis, 1999 (b). Sex pheromone of the spotted tentiform leafminer moth *Phyllonorycter blancardella* (Fabr.) (Lep., Gracillariidae). *J. Appl. Entomol.*, **123**: 603-606.

Mozūraitis, R., V. Buda, V. Jonusaite, A.-K. Borg-Karlson and R. Noreika, 2000. Sex pheromones of *Phyllonorycter acerifoliella* and *Ph. heegerella* and communication peculiarities in three species of leafmining moths. *Entomol. Exp. Appl.*, **94**: 15-23.

Mozūraitis, R., V. Buda, I. Liblikas, C. R. Unelius and A.-K. Borg-Karlson, 2002. Parthenogenesis, calling behavior, and insect-released volatiles of leafminer moth *Phyllonorycter emberizaepenella*. *J. Chem. Ecol.*, **28**(6): 1191-1208.

Mozūraitis, R., V. Karalius, V. Buda and A.-K. Borg-Karlson, 2006. Inter- and intraspecific activities of compounds derived from sex pheromone glands of currant borer, *Synanthedon tipuliformis* (Clerck) (Lepidoptera: Sesiidae). *Z. Naturforsch.*, **61 c**, 278-284.

Mozūraitis, R. and V. Karalius, 2007. Identification of minor sex pheromone components of the poplar clearwing moth *Paranthrene tabaniformis* (Lepidoptera, Sesiidae). *Z. Naturforsch.*, **62c**: 138-142.

Mozūraitis, R., I. Liblikas and R. Noreika, 2008. Sex pheromone communication of tentiform leaf-miners *Phyllonorycter insignitella* and *Ph. nigrescentella* from two related species groups. *Chemoecol.*, **18**: 171-176.

Mozūraitis, R. and V. Buda, 2013. Intra- and interspecific activities of semiochemicals from the sex pheromone gland of the Welsh clearwing, *Synanthedon scoliaeformis*. *J. Chem. Ecol.*, **39**: 1066-1069.

- Muggleston**, S. J. and S. P. Foster, 1989. Sustained-flight tunnel responses of male lightbrown apple moth to synthetic sex pheromone. *Physiol. Entomol.*, 14: 443-449.
- Mukunthan**, N. and B. Singaravelu, 2005. Evaluation of synthetic sex pheromone of sugarcane top borer (*Scirpophaga excerptalis* Walk.). *Sugar Tech.*, 7: 168-170.
- Murakami**, Y., H. Sugie, T. Fukumoto and F. Mochizuki, 2005. Sex pheromone of *Grapholita dimorpha* Komai (Lepidoptera: Tortricidae), and its utilization for monitoring. *Appl. Entomol. Zool.*, 40(3): 521-527.
- Muraki**, Y. T. Taguri, R. Yamakawa and T. Ando, 2014. Synthesis and field evaluation of stereoisomers and analogues of 5-methylheptadecan-7-ol, an unusual sex pheromone component of the lichen moth, *Miltochrista calamine*. *J. Chem. Ecol.*, 40: 250-258.
- Muraki, Y., R. Yamakawa, M. Yamamoto, H. Naka, A. Honma, J. Mappes, K. Suisto and T. Ando, 2017. GC/FT-IR analysis of novel 4,6,9-triene and 2,4,6,9-tetraene occurring in a female pheromone gland of *Arctia plantaginis* (Erebidae: Arctiinae). *Am. J. Chem. Anal.*, 8: 645-656.
- Myerson**, J., W. F. Haddon and E. L. Soderstrom, 1982. sec-Butyl (Z)-7-tetradecenoate. A novel sex pheromone component from the western grapeleaf skeletonizer, *Harrisina brillians*. *Tetrahedron Lett.*, 23: 2757-2760.
- Nagaki**, M., K. Miyata, T. Musashi, J. Kawakami, N. Ohya and H. Sagami, 2009. Insect pheromone-like activity of several isoprenoids against *Phyllonorycterringoniella* (Matsumura). *Trans. Mater. Res. Soc. Jpn.*, 34: 575-578.
- Nagano**, M., Y. Sato, M. Sakai and H. Fujiwara, 1979. Cross-attractiveness of synthetic sex pheromones of lepidopterous species. *J. Takeda Res. Lab.*, 38: 209-215.
- Nagayama**, A., S. Wakamura, N. Taniai and N. Arakaki, 2006. Reinvestigation of sex pheromone components and attractiveness of synthetic sex pheromone of the pink borer, *Sesamia inferens* Walker (Lepidoptera: Noctuidae) in Okinawa. *Appl. Entomol. Zool.* 41 (3): 399-404.
- Naka**, H., L. V. Vang, S. Inomata, T. Ando, T. Kimura, H. Honda, K. Tsuchida and H. Sakurai, 2003. Sex pheromone of the persimmon fruit moth, *Stathmopoda masinissa*: identification and laboratory bioassay of (4E,6Z)-4,6-hexadecadien-1-ol derivatives. *J. Chem. Ecol.*, 29(11): 2447-2459.
- Naka, H., T. Nakazawa, M. Sugie, M. Yamamoto, Y. Horie, R. Wakasugi, Y. Arita, H. Sugie, K. Tsuchida and T. Ando, 2006. Synthesis and characterization of 3,13- and 2,13-octadecadienyl compounds for identification of the sex pheromone secreted by a clearwing

moth, *Nokona pernix*. *Biosci. Biotechnol. Biochem.*, **69**(2) : 508-516.

Naka, H., S. Inomata, K. Matsuoka, M. Yamamoto, H. Sugie, K. Tsuchida, Y. Arita and T. Ando, 2007. Sex pheromone of two Melittini species, *Macroscellesia japonica* and *M. longipes*: identification and field attraction. *J. Chem. Ecol.*, **33** (3) : 591-601.

Naka, H., Y. Horie, F. Mochizuki, L. V. Vang, M. Yamamoto, T. Saito, T. Watarai, K. Tsuchida, Y. Arita and T. Ando, 2008. Identification of the sex pheromone secreted by *Synanthedon hector* (Lepidoptera: Sesiidae). *Appl. Entomol. Zool.*, **43** : 467-474.

Naka, H., M. Mochizuki, K. Nakada, N. D. Do, T. Yamauchi, Y. Arita and T. Ando, 2010. Female sex pheromone of *Glossosphecia romanovi* (Lepidoptera: Sesiidae): Identification and field attraction. *Biosci. Biotechnol. Biochem.*, **74** : 1943-1946.

Naka, H., T. Suzuki, T. Watarai, Y. Horie, F. Mochizuki, A. Mochizuki, K. Tsuchida, Y. Arita and T. Ando, 2013. Identification of the sex pheromone secreted by *Synanthedon tenuis* (Lepidoptera: Sesiidae). *Appl. Entomol. Zool.*, **48** : 27-33.

Naka, H., S. Kubota, F. Kuchiki, H. Shibusaki, M. Yamamoto, Y. Soejima, Y. Arita and T. Ando, 2018. Identification and field attraction of the female sex pheromone of a kiwifruit pest, *Nokona feralis* (Lepidoptera: Sesiidae). *Biosci. Biotechnol. Biochem.*, **82** : 1468-1472.

Nam, I., J.-H. Park, J.-W. Lee, D. H. Lee, W. Jung, U. Lee and I.-K. Park, 2022. Field confirmation of (*Z*)-9-heptacosene and (*3Z,6Z,9Z*)-tricosatriene as key sex pheromone components of Korean *Conogethes punctiferalis* Guenée (Lepidoptera: Crambidae). *J. Econ. Entomol.*, **115** : 1976-1986.

Nässig, W. A. and W. Thomas, 1991. *Cnephasia ecullyana* Réal, 1951, a species native to Central Europe (Lepidoptera, Tortricidae). *Nota lepid.*, **14** : 41-51.

Nasu, Y., K. Tamashima, M. Shibao, S. Yoshimatsu and T. Naito, 2010. Discovery of *Carposina nipponensis* Walsingham and Carposinids caught by synthetic sex pheromone trap for *C. sasakii* Matsumura in Japan (Lepidoptera: Carposinidae). *Jpn. J. Appl. Entomol. Zool.*, **54** : 115-126.

Navarro, I., E. Mas, G. Fabriàs and F. Camps, 1997. Identification and biosynthesis of (*E,E*)-10,12-tetradecadienyl acetate in *Spodoptera littoralis* female sex pheromone gland. *Bioorg. Med. Chem.*, **5** : 1267-1274. (See Ph-4-1-102: pheromone biosynthesis)

Neal, J. W., Jr, 1981. Timing insecticide control of rhododendron borer with pheromone trap catches of males. *Environ. Entomol.*, **10** : 264-266.

Neal, J. W., Jr., J. A. Klun, B. A. Bierl-Leonhardt and M. Schwarz,

1982. Female sex pheromone of *Choristoneura parallela* (Lepidoptera: Tortricidae). *Environ. Entomol.*, **11**: 893-896.
- Neal, J. W., Jr. and T. D. Eichlin, 1983. Seasonal response of six male Sesiidae of woody ornamentals to clearwing borer (Lepidoptera: Sesiidae) lure. *Environ. Entomol.*, **12**: 206-209.
- Negishi, T., T. Ishiwatari, M. Uchida and S. Asano, 1979. A new synergist, 11-methyl-*cis*-9,12-tridecadienyl acetate, for the smaller tea tortrix sex attractant. *Appl. Entomol. Zool.*, **14**: 478-483.
- Nesbitt, B. F., P. S. Beevor, R. A. Cole, R. Lester and R. G. Poppi, 1973. Sex pheromones of two noctuid moths. *Nature*, **244**: 208-209.
- Nesbitt, B. F., P. S. Beevor, R. A. Cole, R. Lester and R. G. Poppi, 1975a. The isolation and identification of the female sex pheromones of the red bollworm moth, *Diparopsis castanea*. *J. Insect Physiol.*, **21**: 1091-1096.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester and V. A. Dyck, 1975b. Identification of the female sex pheromones of the moth, *Chilo suppressalis*. *J. Insect Physiol.*, **21**: 1883-1886.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester and V. A. Dyck, 1976. Identification of the female sex pheromone of the purple stem borer moth, *Sesamia inferens*. *Insect Biochem.*, **6**: 105-107.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester, M. Sternlicht and S. Goldenberg, 1977. Identification and synthesis of the female sex pheromone of the citrus flower moth, *Prays citri*. *Insect Biochem.*, **7**: 355-359.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester, J. C. Davies and K. V. Seshu Reddy, 1979(a). Components of the sex pheromone of the female spotted stalk borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae): identification and preliminary field trials. *J. Chem. Ecol.*, **5**: 153-163.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall and R. Lester, 1979(b). Female sex pheromone components of the cotton bollworm, *Heliothis armigera*. *J. Insect Physiol.*, **25**: 535-541.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester and J. R. Williams, 1980(a). Components of the sex pheromone of the female sugar cane borer, *Chilosacchariphagus* (Bojer) (Lepidoptera: Pyralidae). Identification and field trials. *J. Chem. Ecol.*, **6**: 385-394.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall and R. Lester, 1980(b). (*Z*)-9-Hexadecenal: a minor component of the female sex pheromone of *Heliothis armigera* (Hübner) (Lepidoptera, Noctuidae). *Entomol. Exp. Appl.*, **27**: 306-308.
- Nesbitt, B. F., P. S. Beevor, A. Cork, D. R. Hall, R. M. Murillo and H. R. Leal, 1985. Identification of components of the female sex pheromone of the potato tuber moth, *Scrobipalpopsis solanivora*. *Entomol. Exp. Appl.*, **38**: 81-85.

Nesbitt, B. F., P. S. Beevor, A. Cork, D. R. Hall, H. David and V. Vandagopal, 1986. The female sex pheromone of sugarcane stalk borer, *Chilo auricilius*. Identification of four components and field tests. *J. Chem. Ecol.*, **12**: 1377-1388.

Newton, P. J., C. D. Thomas, V. C. Mastro and C. P. Schwalbe, 1993. Improved two-component blend of the synthetic female sex pheromone of *Cryptophlebia leucotreta*, and identification of an attractant for *C. peltastica*. *Entomol. Exp. Appl.*, **66**: 75-82.

Nielsen, D. G., F. F. Purrington, J. H. Tumlinson, R. E. Doolittle and C. E. Yonce, 1975. Response of male clearwing moths to caged virgin females, female extracts, and synthetic sex attractants. *Environ. Entomol.*, **4**: 451-454.

Nielsen, D. G., F. F. Purrington, R. L. Campbell, T. R. Wilmot, J. Capizzi and J. H. Tumlinson, 1978(a). Sex attractants for sequoia pitch moth and strawberry crown moth. *Environ. Entomol.*, **7**: 544-546.

Nielsen, D. G. and F. F. Purrington, 1978(b). Field attraction of male *Podosesia syringae* and *P. aureocincta* to blends of synthetic compounds. *Environ. Entomol.*, **7**: 708-710.

Nielsen, D. G., F. F. Purrington and G. F. Shambaugh, 1979. EAG and field responses of sesiid males to sex pheromones and related compounds. "Pheromone of the Sesiidae," ed. by J. W. Neal, Jr., ARR-NE-6, U. S. Department of Agriculture, Beltsville, Maryland, pp. 11-26.

Niogret, J., A. Ekayanti and A. Zhang, 2022. Sex Pheromone of cocoa pod borer, *Conopomorpha cramerella*: field activity evaluation of pheromone formulations in an Indonesia plantation. *Insects*, **13**: 663.

Niwa, C. G., L. L. Sower and G. E. Daterman, 1987. Chemistry and field evaluation of the sex pheromone of ponderosa pine tip moth, *Rhyacionia zozana* (Lepidoptera: Tortricidae). *Environ. Entomol.*, **16**: 1287-1290.

Niwa, C. G. and L. L. Sower, 1992. Sex pheromone chemistry of the southwestern pine tip moth (Lepidoptera: Tortricidae). *Environ. Entomol.*, **21**: 1410-1413.

Noguchi, H., Y. Tamaki and T. Yushima, 1979. Sex pheromone of the tea tortrix moth: isolation and identification. *Appl. Entomol. Zool.*, **14**: 225-228.

Noguchi, H., Y. Tamaki, S. Arai, M. Shimoda and I. Ishikawa, 1981. Field evaluation of synthetic sex pheromone of the oriental tea tortrix moth, *Homona magnanima* Diakonoff (Lepidoptera: Tortricidae). *Jap. J. Appl. Entomol. Zool.*, **25**: 170-175.

Noguchi, H., H. Sugie, Y. Tamaki and Y. Oomasa, 1985. Sex-pheromone components and related compounds released by virgin females of *Adoxophyes* sp. and *Adoxophyes orana fasciata* (Lepidoptera: Tortricidae). *Jap. J. Appl. Entomol. Zool.*, **29**: 278-283.

- Nojima**, S., A. Classen, A. T. Groot and C. Scha, 2018. Qualitative and quantitative analysis of chemicals emitted from the pheromone gland of individual *Heliothis subflexa* females. *PLoS ONE*, **13**: e0202035.
- Novák**, L., M. Tóth, J. Balla and C. Szántay, 1979. Sex pheromone of the cabbage armyworm, *Mamestra brassicae*: isolation, identification and stereocontrolled synthesis. *Acta Chim. Acad. Sci. Hung.*, **102**: 135-140.
- Novotný**, Von J., J. Patočka, I. Hrdý and J. Vrkoč, 1990. Monitoring of *Tortrix viridana* L. by means of the synthetic sexual pheromone. *Anz. Schädl. Pflanz. Umwelt.*, **63**: 125-129.
- Nunez**, S., J. J. De Vlieger, J. J. Rodriguez, C. J. Persoons and I. Scatoni, 2002. Sex pheromone of South American tortricid moth *Argyrotaenia sphaleropa*. *J. Chem. Ecol.*, **28**(2): 425-432.
- Odell**, T. M., C.-H. Xu, P. W. Schaefer, B. A. Leonhardt, D.-F. Yao and X.-D. Wu, 1992. Capture of gypsy moth, *Lymantria dispar* (L.), and *Lymantria mathura* (L.) males in traps baited with disparlure enantiomers and olefin precursor in the People's Republic of China. *J. Chem. Ecol.*, **18**: 2153-2159.
- Oetting**, R. D., 1977. Immature stages and biology of *Chionodes psiloptera* a pest in bluegrass seed fields. *Pan-Pac. Entomol.*, **53**: 258-268.
- Ohmasa**, Y., S. Wakamura, S. Kozaki, H. Sugie, M. Horike, C. Hirano and S. Mori, 1991. Sex pheromone of the fruit-piercing moth, *Oraesia excavata* (Butler) (Lepidoptera: Noctuidae): isolation and identification. *Appl. Entomol. Zool.*, **26**: 55-62.
- Ohshima**, M., Y. Kuwahara, S. Matsuyama and T. Suzuki, 1993. Identification of the sex pheromone of the European grain moth *Nemapogon granellus*: (3Z,13Z)-3,13-octadecadienyl acetate. *J. Pesticide Sci.*, **18**: 163-167.
- Ohta**, K., S. Tatsuki, K. Uchiumi, M. Kurihara and J. Fukami, 1976. Structures of sex pheromones of rice stem borer. *Agric. Biol. Chem.*, **40**: 1897-1899.
- Oleander**, A., D. R. Hall, D. P. Bray and J. P. J. Burman, 2019. Identification of female sex pheromone for monitoring the barred tooth striped moth, *Trichopteryx polycommata*, a priority conservation species. *J. Chem. Ecol.*, **45**: 649-656.
- Oliver**, J. E., J. C. Dickens, M. Zlotina, V. C. Mastro and G. I. Yurchenko 1999. Sex attractant of the rosy Russian gypsy moth (*Lymantria mathura* Moore). *Z. Naturforsch.*, **54c**: 387-394.
- Ono**, T., R. E. Charlton and R. T. Cardé, 1990. Variability in pheromone composition and periodicity of pheromone titer in potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *J.*

Chem. Ecol., 16: 531-542.

- Ono, T., P. Chouvalitwongporn and T. Saito, 1997. Comparison of the sex pheromone system between Japanese and Thai populations of the potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *Appl. Entomol. Zool.*, 32: 514-517.
- Ono, M., Y. Murata, J. M. S. Vento, W. S. Leal, E. F. Vilela, J. R. P. Para and A. J. Aires, 2001. Dodecenyl acetate and dodecenol as sex attractants for citrus borer moth. *Jpn. Kokai Tokkyo Koho*, JP 2001-81002A.
- Ostrauskas, H., 2003. Moths caught in pheromone traps for southern armyworm (*Spodoptera eridania* Cr.), fall armyworm (*S. frugiperda* Sm.), and Egyptian cotton leafworm (*S. littoralis* Bsd.) (Noctuidae, Lepidoptera) during 1999-2001 in Lithuania. *Acta Zool. Lit.*, 13: 411-424.
- Ostrauskas, H., 2004. Moths caught in pheromone traps for American white moth (*Hyphantria cunea* Dr.) (Arctiidae, Lepidoptera) in Lithuania during 2001. *Acta Zool. Litua.*, 14: 66-74.
- Ostrauskas, H., P. Ivinskis and V. Bûda, 2010(a). Moths caught in traps during a survey of *Choristoneura fumiferana* (Clemens, 1865) - Lepidoptera, Tortricidae - in Lithuania. *Acta Zool. Lit.*, 20: 156-161.
- Ostrauskas, H., P. Ivinskis and V. Bûda, 2010(b). Moths caught in traps during the survey of *Choristoneura Rosaceana* (Lepidoptera, Tortricidae) in Lithuania. *Acta Zool. Lit.*, 20: 242-249.
- Palacio-Cortes, A. M., M. G. Fonseca and P. H. G. Zarbin, 2014. Toward the identification of the sex pheromone of *Diatraea indigenella* Dyar & Heinrich (Lepidoptera: Crambidae): calling behavior and chemical structure of a major component. *Neotrop. Entomol.*, 43: 526-531.
- Park, K. C., A. Cork, K. S. Boo and D. R. Hall, 1994. Biological activity of female sex pheromone of the oriental tobacco budworm, *Helicoverpa assulta* (Guenee) (Lepidoptera: Noctuidae): electroantennography, wind tunnel observation and field trapping. *Korean J. Appl. Entomol.*, 33(1): 26-32.
- Park, K. C., A. Cork and K. S. Boo, 1996. Intrapopulational changes in sex pheromone composition during scotophase in oriental tobacco budworm, *Helicoverpa assulta* (Guenée) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, 22(6): 1201-1210.
- Park, S.-C. and S. P. Foster, 1998. Comparative sex pheromone-associated behavior of *Planotortrix notophaea* and *P. octo* (Lepidoptera: Tortricidae). *J. Asia Pacific Entomol.*, 1: 91-98.

- Park, J. H., K. S. Han, K. Mori and K. S. Boo, 2002. Right stereoisomers for sex pheromone components of the apple leafminer, *Lyonetia prunifoliella*, in Korea. *J. Chem. Ecol.*, **28**(12): 2515-2525.
- Park, H.-K., H.-R. Lee, C.-S. Jung and S.-H. Koh, 2019. Synergic effect of sex pheromone (*7R,8S*)-*cis*-7,8-epoxy-2-methyloctadec-17-ene on (+)-disparlure of the Asian gypsy moth, *Lymantria dispar*, in Korea. *J. Forest. Res.*, **30**: 1119-1122.
- Pasek, J. E. and M. E. Dix, 1989. Life history of a ponderosa pine coneworm, *Dioryctria auranticella* (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **82**: 879-885.
- Peltotalo, P. and T. Tuovinen, 1986. Specificity of pheromone preparations for lepidopterous pests. *Ann. Agric. Fenniae*, **25**: 139-146.
- Pelozuelo, L., C. Malosse, G. Genestier, H. Guenego and B. Frerot, 2004. Host-plant specialization in pheromone strains of the European corn borer *Ostrinia nubilalis* in France. *J. Chem. Ecol.*, **30**(2): 335-352.
- Peña, A., H. Arn, H.-R. Buser, S. Rauscher, F. Bigler, R. Brunetti, S. Maini and M. Tóth, 1988. Sex pheromone of European corn borer, *Ostrinia nubilalis*: polymorphism in various laboratory and field strains. *J. Chem. Ecol.*, **14**: 1359-1366.
- Peng, C.-L., P. Gu, J. Li, Q.-Y. Chen, C.-H. Feng, H.-H. Luo and Y.-J. Du, 2012. Identification and field bioassay of the sex pheromone of *Trichophysetis cretacea* (Lepidoptera: Crambidae). *J. Econ. Entomol.*, **105**: 1566-1572.
- Persoons, C. J., A. K. Minks, S. Voerman, W. L. Roelofs and F. J. Ritter, 1974. Sex pheromones of the moth, *Archips podana*: isolation, identification and field evaluation of two synergistic geometrical isomers. *J. Insect. Physiol.*, **20**: 1181-1188.
- Persoons, C. J., S. Voerman, P. E. J. Verwiel, P. J. F. Nooijen, W. J. Nooijen, F. J. Ritter and A. K. Minks, 1976(a). Sex pheromone of the potato tuberworm moth, *Phthorimaea operculella*: isolation and identification. *Med. Fac. Landbouww. Rijksuniv. Gent.*, **41**: 945-948.
- Persoons, C. J., S. Voerman, P. E. J. Verwiel, F. J. Ritter, W. J. Nooyen and A. K. Minks, 1976(b). Sex pheromone of the potato tuberworm moth, *Phthorimaea operculella*: isolation, identification and field evaluation. *Entomol. Exp. Appl.*, **20**: 289-300.
- Persoons, C. J., F. J. Ritter, D. Hainaut and J. P. Demoute, 1976(c). Sex pheromone of the false codling moth *Cryptophlebia* (= *Argyroploce*) *leucotreta* (Lepidoptera: Tortricidae) *trans*-8-dodecenyl acetate, a corrected structure. *Med. Fac. Landbouww. Rijksuniv. Gent.*, **41**: 937-943.
- Persoons, C. J., F. J. Ritter and W. J. Nooyen, 1977. Sex pheromone of the false codling moth *Cryptophlebia leucotreta* (Lepidoptera:

Tortricidae). Evidence for a two-component system. *J. Chem. Ecol.*, **3**: 717-722.

Persoons, C. J., C. van der Kraan, W. J. Nooijen, F. J. Ritter, S. Voerman and T. C. Baker, **1981**. Sex pheromone of the beet armyworm, *Spodoptera exigua*: isolation, identification and preliminary field evaluation. *Entomol. Exp. Appl.*, **30**: 98-99.

Persoons, C. J., S. Voerman, C. van der Kraan and C. J. H. Booij, **1984**. Sex pheromone of the leafroller moth *Syndemis musculana* (Tortricidae): preliminary identification and field evaluation. *Entomol. Exp. Appl.*, **35**: 101-103.

Persoons, C. J., J. D. Vos, J. S. Yadav, A. R. Prasad, S. Sighomony, K. N. Jyothi and A. L. Prasuna, **1993**. Indo-Dutch cooperation on pheromones of Indian agricultural pest insects: sex pheromone components of *Diacrisia obliqua* (Arctiidae), *Achaea janata* (Noctuidae) and *Amsacta albistriga* (Arctiidae). *Bulletin OILB SROP*, **16**: 136-140.

Piccardi, P., A. Capizzi, G. Cassani, P. Spinelli, E. Arsura and P. Massardo, **1977**. A sex pheromone component of the old world bollworm *Heliothis armigera*. *J. Insect Physiol.*, **23**: 1443-1445.

Picimbon, J. F., C. Gadenne, J. M. Bécard, J. L. Clément and L. Sreng, **1997**. Sex pheromone of the French black cutworm moth, *Agrotis ipsilon* (Lepidoptera: Noctuidae): identification and regulation of a multicomponent blend. *J. Chem. Ecol.*, **23**(1): 221-230.

Pineda-Ríos, J. M., J. C. Tovar, J. M. Sámano, L. F. S. Abarca, R. M. L. Romero and E. de J. A. Suárez, **2016**. La composición y proporción de los componentes de la feromona sexual de *Hypsipyla grandella* Zeller (Lepidoptera: Pyralidae) varía dependiendo de la técnica utilizada para su aislamiento. *Entomotropica*, **31**: 172-185.

Pires, E. V., A. L. Mendonça, L. Vanícková, N. S. J. Serra, R. C. C. da Silva, D. C. dos Santos, R. S. Campos, A. E. G. Sant'Ana and R. R. do Nascimento, **2016**. Identification and field and laboratory tests of the sex pheromone of *Cerconota anonella* Sepp. (Lepidoptera: Oecophoridae). *J. Appl. Entomol.*, **140**: 72-80.

Pletnev, V. A., V. L. Ponomarev, N. V. Vendilo, S. A. Kurbatov and K. V. Lebedeva, **2000**. Search of the pheromone of Siberian silkworm *Dendrolimus superans sibiricus* (Lepidoptera: Lasiocampidae). *Agrokhimiya*, (6): 67-72. [*Chem. Abst.*, **134**: 263662]

Polavarapu, S. and G. Lonergan, **1998**. Sex pheromone of *Choristoneura parallela* (Lepidoptera: Tortricidae): components and development of a pheromone lure for population monitoring. *Environ. Entomol.*, **27**(5): 1242-1249.

Ponder, B. M. and W. D. Seabrook, **1991**. Sensitivity of blueberry

leafier moths (Lepidoptera: Tortricidae) (Kearfott) to their own sex pheromone: mating bioassay, electroantennogram, and trap attractancy studies. *Can. Entomol.*, **123**: 231-238.

Pope, M. M., L. K. Gaston and T. C. Baker, 1982. Composition, quantification, and periodicity of sex pheromone gland volatiles from individual *Heliothis virescens* females. *J. Chem. Ecol.*, **8**: 1043-1055.

Pope, M. M., L. K. Gaston and T. C. Baker, 1984. Composition, quantification, and periodicity of sex pheromone volatiles from individual *Heliothis zea* females. *J. Insect Physiol.*, **30**: 943-945.

Priesner, E., M. Jacobson, H. J. Bestmann, 1975. Structure-response relationships in noctuid sex pheromone reception. *Z. Naturforsch.*, **30c**: 283-293.

Priesner, E., H.-J. Bestmann, O. Vostrowsky and P. Rösel, 1977. Sensory efficacy of alkyl-branched pheromone analogues in noctuid and tortricid Lepidoptera. *Z. Naturforsch.*, **32c**: 979-991.

Priesner, E., H. Bogenschütz, W. Altenkirch and H. Arn, 1978. A sex attractant for the pine beauty moth, *Panolis flammea*. *Z. Naturforsch.*, **33c**: 1000-1002.

Priesner, E., H. Bogenschütz and H. Arn, 1980(a). A sex attractant for the European fir budworm moth, *Choristoneura murinana*. *Z. Naturforsch.*, **35c**: 390-398.

Priesner, E., 1980(b). Sex attractant system in *Polia pisi* L. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **35c**: 990-994.

Priesner, E. and H. Bogenschütz, 1982(a). Further field experiments on the specificity of synthetic sex-attractants in *Choristoneura murinana*. *Z. Angew. Entomol.*, **94**: 331-351.

Priesner, E., W. Altenkirch, W. Baltensweiler and H. Bogenschütz, 1982(b). Evaluation of (Z)-5-decen-1-ol as an attractant for male larch casebearer moths, *Coleophora laricella*. *Z. Naturforsch.*, **37c**: 953-966.

Priesner, E. and M. Schroth, 1983. Supplementary data on the sex attractant system *Panolis flammea*. *Z. Naturforsch.*, **38c**: 870-873.

Priesner, E., C.M. Naumann and J. Stertenbrink, 1984(a). Specificity of synthetic sex-attractants in Zygaena moths. *Z. Naturforsch.*, **39c**: 841-844.

Priesner, E., 1984(b). Analysis of a sex-attractant system in the noctuid moth *Rhyacia baja* Schiff. *Z. Naturforsch.*, **39c**: 845-848.

Priesner, E., 1984(c). The pheromone receptor system of male *Eulia ministrana* L., with notes on other Cnephasiini moths. *Z. Naturforsch.*, **39c**: 849-852.

- Priesner, E., H. Bogenschütz, R. Albert, D. W. Reed and M. D. Chisholm, 1984 (d). Identification and field evaluation of a sex pheromone of the European pine moth. *Z. Naturforsch.*, **39c**: 1192-1195.
- Priesner, E., 1985 (a). Specificity of sexual attractants in *Xestia triangulum* Hufn. and *X. ditrapezium* Schiff. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **40c**: 939-942.
- Priesner, E., 1985 (b). Inhibitors of sexual attraction in the moth *Agrotis exclamationis*. *Z. Naturforsch.*, **40c**: 943-945.
- Priesner, E., 1985 (c). Species-specific sex-attractants for male *Diachrysia chrysitis* (L.) and *D. tutti* (Koster.) (Lepidoptera, Noctuidae: Plusiinae). *Mitt. Schweiz. Entomol. Ges.*, **58**: 373-391.
- Priesner, E., G. Dobler and S. Voerman, 1986 (a). Synergism of positional isomers in sex-attractant systems of clearwing moths (Sesiidae). *Entomol. Exp. Appl.*, **41**: 311-313.
- Priesner, E., P. Witzgall and S. Voerman, 1986 (b). Field attraction response of raspberry clearwing moths, *Pennisetia hylaeiformis* Lasp. (Lepidoptera: Sesiidae), to candidate pheromone chemicals. *J. Appl. Entomol.*, **102**: 195-210.
- Priesner, E., 1987 (a). (*Z*)-5-Dodecen-1-ol, another inhibitor of pheromonal attraction in *Coleophora laricella*. *Z. Naturforsch.*, **42c**: 1349-1351.
- Priesner, E., 1987 (b). (*Z*)-3-Tetradecenyl acetate as a sex-attractant component in Gelechiinae and Anomologinae (Lepidoptera: Gelechiidae). *Z. Naturforsch.*, **42c**: 1352-1354.
- Priesner, E., H. Bogenschütz and C. Löfstedt, 1988 (a). Secondary sex pheromone components of *Choristoneura murinana* Hb. (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **43c**: 918-928.
- Priesner, E., 1988 (b). (*Z*)-3- and (*Z*)-5-Decenyl acetates, sex-attractant components for male *Eustrotia uncula* Cl. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **43c**: 961-962.
- Priesner, E., 1989 (a). Two-component sexual attractant for male *Batrachedra pinicolella* (Zell.) (Lepidoptera: Batrachedridae). *Z. Naturforsch.*, **44c**: 1061-1062.
- Priesner, E., D. W. Reed, E. W. Underhill and H. Bogenschütz, 1989 (b). (*Z,Z*)-7,9-Dodecadienyl acetate, sexpheromone of *Epinotia tatedella* Clerck (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **15**: 2457-2464.
- Priesner, E. and K. Spatenka, 1990. The distribution of *Pennisetia bohemica* Kralicek & Povolny, 1974 (Lepidoptera: Sesiidae) in central Europa as determined by pheromone trapping. *Mitt. Schweiz. Entomol. Ges.*, **63**: 87-98.
- Priesner, E. and Q.-H. Zhang, 1991. Field response of male eastern larch casebearer *Coleophora dahurica* Flkv. (Lep., Coleophoridae)

to synthetic sex-attractant and attraction-inhibitors. *J. Appl. Entomol.*, **112**: 321-326.

Priesner, E., 1993. Pheromontest an einer südbayerischen Population von *Synanthesdon soffneri* Spatenka, 1983 (Lepidoptera: Sesiidae). *NachrBl. Bayer. Entomol.*, **42**(4): 97-107.

Pu, G.-Q., M. Yamamoto, Y. Takeuchi, H. Yamazawa and T. Ando 1999. Resolution of epoxydienes by reversed-phase chiral HPLC and its application to stereochemistry assignment of mulberry looper sex pheromone. *J. Chem. Ecol.*, **25**(5): 1151-1162.

Qin, X.-R., T. Ando, M. Yamamoto, M. Yamashita, K. Kusano and H. Abe, 1997. Resolution of pheromonal epoxydienes by chiral HPLC, stereochemistry of separated enantiomers, and their field evaluation. *J. Chem. Ecol.*, **23**(5): 1403-1417.

Quartey, G. K. and T. H. Coaker, 1993. Role of sex pheromone components in the orientation behaviour of *Ephestia cautella*. *Entomol. Exp. Appl.*, **66**: 237-245.

Quero, C., F. Camps and A. Guerrero, 1995. Behavior of processionary males (*Thaumetopoea pityocampa*) induced by sex pheromone and analogs in a wind tunnel. *J. Chem. Ecol.*, **21**: 1957-1969.

Quero, C., D. A. Malo, G. Fabriàs, F. Camps, P. Lucas, M. Renou and A. Guerrero, 1997. Reinvestigation of female sex pheromone of processionary moth (*Thaumetopoea pityocampa*): No evidence for minor components. *J. Chem. Ecol.*, **23**(3): 713-726.

Quero, C., J. Bau, A. Guerrero, M. Breuer, A. D. Loof, H.-G. Kontzog and F. Camps, 2003. Sex pheromone of the oak processionary moth *Thaumetopoea processionea*. Identification and biological activity. *J. Agric. Food Chem.*, **51**: 2987-2991.

Rahmani, R., D. Carrasco, G. P. Svensson, H. Roweck, N. Ryholm, M. C. Larsson and E. Hedenstroem, 2020. Identification and Synthesis of Putative Pheromone Components of the Threatened Salt Marsh Bagworm Moth, *Whittleia retiella* (Lepidoptera: Psychidae) *J. Chem. Ecol.*, **46**: 115-127.

Raina, A. K., J. A. Klun, M. Schwarz, A. Day, B. A. Leonhardt and L. W. Douglass, 1986(a). Female sex pheromone of the melonworm, *Diaphania hyalinata* (Lepidoptera: Pyralidae), and analysis of male responses to pheromone in a flight tunnel. *J. Chem. Ecol.*, **12**: 229-237.

Raina, A. K., J. A. Klun, J. D. Lopez and B. A. Leonhardt, 1986(b). Female sex pheromone of *Heliothis phloxiphaga* (Lepidoptera: Noctuidae): chemical identification, male behavioral response in the flight tunnel, and field tests. *Environ. Entomol.*, **15**: 931-935.

Ramachandran, R., P. Caballero and Z. R. Khan, 1990. Pheromone components of rice leaffolders (LF) *Cnaphalocrocis medinalis* and

Marasmia patnalis. *Int. Rice Res. New.*, **15**: 25-26.

Raman, K. V., 1982. Field trials with the sex pheromone of the potato tuberworm. *Environ. Entomol.*, **11**: 367-370.

Ramaswamy, S. B., S. A. Randle and W. K. Ma, 1985. Field evaluation of the sex pheromone components of *Heliothis virescens* (Lepidoptera: Noctuidae) in cone traps. *Environ. Entomol.*, **14**: 293-296.

Rao, A. G., K. Krishnaiah, D. R. Hall, A. Cork and D. D. R. Reddy, 1993. Evaluation of sex pheromone components of rice leaf folder, *Cnaphalocrocis medinalis* Guenée. *Current Sci.*, **65**: 355-356.

Rao, A. G., D. D. R. Reddy, K. Krishnaiah, P. S. Beevor, A. Cork and D. R. Hall, 1995. Identification and field optimisation of the female sex pheromone of the rice leaf folder, *Cnaphalocrocis medinalis* in India. *Entomol. Exp. Appl.*, **74**: 195-200.

Rauscher, S., H. Arn and P. Guerin, 1984. Effects of dodecyl acetate and Z-10-tridecenyl acetate on attraction of *Eupoecilia ambiguella* males to the main sex pheromone component, Z-9-dodecenyl acetate. *J. Chem. Ecol.*, **10**: 253-264.

Razov, J., K. A. Efetov, K. Franin, T. B. Toshova and M. A. Subchev, 2017. The application of sex pheromone traps for recording the Procridinae fauna (Lepidoptera: Zygaenidae) in Croatia. *Entomol. Gaz.*, **68**: 49-53.

Read, J. S., P. H. Hewitt, F. L. Warren and A. C. Myberg, 1974. Isolation of the sex pheromone of the moth *Argyroploce leucotreta*. *J. Insect Physiol.*, **20**: 441-450.

Read, J. S. and P. S. Beevor, 1976. Analytical studies on the sex pheromone complex of *Ephestia cautella* (Walker) (Lepidoptera: Phycitidae). *J. Stored Prod. Res.*, **12**: 55-57.

Read, J. S. and C. P. Haines, 1979. Secondary pheromone components and synergism in stored-products Phycitinae. *J. Chem. Ecol.*, **5**: 251-257.

Reed, D. W., M. D. Chisholm and E. W. Underhill, 1984. Sex attractant for three species of the genus *Oncocnemis*: *O. chandleri* (Grt.), *O. cibalis* (Grt.), and *O. mackiei* (B. & Benj.) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **10**: 1701-1706.

Reed, D. W. and M. D. Chisholm, 1985 (a). Field trapping of three *Epinotia* species with (Z,Z)-7,9-dodecadienyl acetate (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **11**: 1389-1398.

Reed, D. W. and M. D. Chisholm, 1985 (b). Attraction of moth species of Tortricidae, Gelechiidae, Geometridae, Drepanidae, Pyralidae, and Gracillariidae families to field traps baited with conjugated dienes. *J. Chem. Ecol.*, **11**: 1645-1657.

Reed, D. W., E. W. Underhill and E. M. Giblin, 1987. Attraction of

sphingid moths (Lepidoptera: Sphingidae) to 10,12-hexadecadienyl aldehydes and acetates: evidence of pheromone components. *J. Chem. Ecol.*, **13**: 931-942.

Renou, M., C. Descoins, E. Priesner, M. Gallois and M. Lettere, **1979**. Z-7-Tetradecenal, the main component of the sex pheromone of the olive moth *Prays oleae*. *C. R. Hebd. Seances Acad. Sci., Ser. D*, **288**: 1559-1562.

Renou, P. M., Ch. Descoins, J. Y. Lallemand, E. Priesner, M. Lettere and M. Gallois, **1980**. (E)-3-dodecenyl acetate, main component of the sex pheromone of the sugarbeet moth, *Scrobipalpa ocellatella* Boyd. (Lepidoptera, Gelechiidae). *Z. Angew. Entomol.*, **90**: 275-289.

Renou, M., C. Descoins, E. Priesner, M. Gallois and M. Lettere, **1981** (a). A study of the sex pheromone of the leek moth, *Acrolepiopsis assectella* (Lepidoptera: Acrolepiidae). *Entomol. Exp. Appl.*, **29**: 198-208.

Renou, M., B. Lalanne-Cassou, B. Frérot, M. Gallois and C. Descoins, **1981** (b). Composition of the pheromonal secretion produced by virgin females of *Mamestra (Polia) pisi* L. (Lepidoptera, Noctuidae, Hadeninae). *C. R. Acad. Sci., Ser. 3*, **292**: 1117-1119.

Renou, M., B. Lalanne-Cassou, J.-C. Doré and M.-L. Milat, **1988**. Electroantennographic analysis of sex pheromone specificity in neotropical Catocalinae (Lepidoptera: Noctuidae): a multivariate approach. *J. Insect Physiol.*, **34**: 481-488.

Renou, M., **1991**. Sex pheromone reception in the moth, *Mamestra thalassina*. Characterization and distribution of two types of olfactory hairs. *J. Insect Physiol.*, **37**: 617-626.

Rhainds, M., G. Gries, J.-X. Li, R. Gries, K. N. Slessor, C. M. Chinchilla and A. C. Oehlschlager, **1994**. Chiral esters: sex pheromone of the bagworm, *Oiketicus kirbyi* (Lepidoptera: Psychidae). *J. Chem. Ecol.*, **20**: 3083-3096.

Riba, M., J. A. Rosell, M. Eizaguirre, R. Canela and A. Guerrero, **1990**. Identification of a minor component of the sex pheromone of *Leucoptera malifoliella* (Lepidoptera, Lyonetiidae). *J. Chem. Ecol.*, **16**: 1471-1483.

Ritter, F. J. and C. J. Persoons, **1975**. Recent development in insect pheromone research, in particular in the Netherlands. *Neth. J. Zool.*, **25**: 261-275.

Robacker, D. C. and K. J. R. Johnson, **1989**. Pheromone emission and blend percentages in *Eoreuma loftini* determined by two methods. *J. Chem. Ecol.*, **15**: 105-120.

Rocchini, L. A., B. S. Lindgren and R. G. Bennett, **2003**. Douglas-fir pitch moth, *Synanthedon novaroensis* (Lepidoptera: Sesiidae) in North-Central British Columbia: flight period and the effect of trap type and pheromone dosage on trap catches. *Environ. Entomol.*,

32: 208-213.

Rodríguez, S. A. and M. A. Nazareno 2018. Identification of the sex pheromone of female *Eurata patagiata*. *Southwestern Entomol.*, 43: 81-90.

Roelofs, W. L. and H. Arn. 1968 (a). Sex attractant of the red-banded leafroller moth. *Nature*, 219: 513.

Roelofs, W. L. and A. Comeau, 1968 (b). Sex pheromone perception. *Nature*, 220: 600-601.

Roelofs, W. L., A. Comeau and R. Selle, 1969. Sex pheromone of the oriental fruit moth. *Nature*, 224: 723.

Roelofs, W. L. and A. Comeau, 1970 (a). Lepidopterous sex attractants discovered by field screening tests. *J. Econ. Entomol.*, 63: 969-974.

Roelofs, W. L. and J. P. Tette, 1970 (b). Sex pheromone of the oblique-banded leaf roller moth. *Nature*, 226: 1172.

Roelofs, W. L. and A. Comeau, 1971 (a). Sex attractants in Lepidoptera. "Pesticide Chemistry, vol. III, Chemical Releasers in Insects," ed. by A. S. Tobotu, Cordon of Beach, New York, N. Y., pp. 91-114.

Roelofs, W. L. and R. T. Cardé, 1971 (b). Hydrocarbon sex pheromone in tiger moths (Arctiidae). *Science*, 171: 684-686.

Roelofs, W. L., J. P. Tette, E. F. Taschenberg and A. Comeau, 1971 (c). Sex pheromone of the grape berry moth: identification by classical and electroantennogram methods, and field tests. *J. Insect Physiol.*, 17: 2235-2243.

Roelofs, W. L., R. Cardé, G. Benz and G. von Salis, 1971 (d). Sex attractant of the larch bud moth found by electroantennogram method. *Experientia*, 27: 1438-1439.

Roelofs, W., A. Comeau and G. M. Hill, 1971 (e). Sex attractant of the codling moth: characterization with electroantennogram technique. *Science*, 174: 297-299.

Roelofs, W. L., R. T. Cardé, R. J. Bartell and P. G. Tierney, 1972. Sex attractant trapping of the European corn borer in New York. *Environ. Entomol.*, 1: 606-608.

Roelofs, W., J. Kochansky, R. Cardé, H. Arn and S. Rauscher, 1973. Sex attractant of the grape vine moth, *Lobesia botrana*. *Mitt. Schweiz. Entomol. Ges.*, 46: 71-73.

Roelofs, W. L. and R. T. Cardé, 1974 (a). Oriental fruit moth and lesser appleworm attractant mixtures refined. *Environ. Entomol.*, 3: 586-588.

Roelofs, W. L. and R. T. Cardé, 1974 (b). Sex pheromones in the reproductive isolation of lepidopterous species.

"North-Holland Research Monograph Frontiers of Biology, vol. 32, Pheromones," ed. by M. C. Birch, North-Holland Publishing Co., Amsterdam, pp. 96-114.

- Roelofs, W. L., A. S. Hill, R. T. Cardé and T. C. Baker, 1974(c). Two sex pheromone components of tobacco budworm moth, *Heliothis virescens*. *Life Sci.*, 14: 1555-1562.
- Roelofs, W. L., A. Hill, R. Cardé, J. Tette, H. Madsen and J. Vakenti, 1974(d). Sex pheromones of the fruittree leafroller moth, *Archips argyrospilus*. *Environ. Entomol.*, 3: 747-751.
- Roelofs, W., A. Hill and R. Cardé, 1975(a). Sex pheromone components of the redbanded leafroller, *Argyrotaenia velutinana* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, 1: 83-89.
- Roelofs, W., J. Kochansky, E. Anthon, R. Rice and R. Cardé, 1975(b). Sex pheromone of the peach twing borer moth (*Anarsia lineatella*). *Environ. Entomol.*, 4: 580-582.
- Roelofs, W. L., J. P. Kochansky, R. T. Carde, G. G. Kennedy, C. A. Henrick, J. N. Labovitz and V. L. Corbin, 1975(c). Sex pheromone of the potato tuberworm moth, *Phthorimaea operculella*. *Life Sci.*, 17: 699-706.
- Roelofs, W., A. Hill, A. Cardé, R. Cardé, H. Madsen and J. Vakenti, 1976(a). Sex pheromone of the European leafroller, *Archips rosanus*. *Environ. Entomol.*, 5: 362-364.
- Roelofs, W., A. Carde, A. Hill and R. Carde, 1976(b). Sex pheromone of the threelined leafroller, *Pandemis limitata*. *Environ. Entomol.*, 5: 649-652.
- Roelofs, W. L., R. F. Lagier and S. C. Hoyt, 1977(a). Sex pheromone of the moth, *Pandemis pyrusana*. *Environ. Entomol.*, 6: 353-354.
- Roelofs, W. L., W. H. Reissig and R. W. Weires, 1977(b). Sex attractant for the spotted tentiform leaf miner moth, *Lithoccolletis blancardella*. *Environ. Entomol.*, 6: 373-374.
- Roelofs, W. L., A. S. Hill, C. W. Berisford and J. F. Godbee, 1979. Sex pheromone of the subtropical pine tip moth, *Rhyacionia subtropica*. *Environ. Entomol.*, 8: 894-895.
- Roelofs, W. L., A. J. Tamhankar, A. Comeau, A. S. Hill and E. F. Taschenberg, 1980. Moth activity periods and identification of the sex pheromone of the uglynest caterpillar, *Archips cerasivoranus*. *Ann. Entomol. Soc. Am.*, 73: 631-634.
- Roelofs, W. L. and R. L. Brown, 1982(a). Pheromones and evolutionary relationships of Tortricidae. *Ann. Rev. Ecol. Syst.*, 13: 395-422.
- Roelofs, W. L., A. S. Hill, C. E. Linn, J. Meinwald, S. C. Jain, H. J. Herbert and R. F. Smith, 1982(b). Sex pheromone of the winter moth, a geometrid with unusually low temperature precopulatory responses. *Science*, 217: 657-659.

- Roelofs, W. L. and L. Bjostad, 1984. Biosynthesis of lepidopteran pheromones. *Bioorg. Chem.*, **12**: 279-298.
- Rojas, J. C., L. Cruz-López, E. A. Malo, O. Diaz-Gomez, G. Calyecac and J. C. Tovar, 2006. Identification of the sex pheromone of *Copitarsia decolora* (Lepidoptera: Noctuidae). *J. Econ. Entomol.*, **99**: 797-802.
- Rossi, R., A. Carpita, P. Piccardi, C. Tonini, G. Guglielmetti, A. Niccoli and G. A. Fenili, 1982. Identification of (Z)-11-tetradecen-1-yl acetate as a component of *Epichoristodes acerbella* sex pheromone. *Frust. Entomol. Nuova serie*, **2**: 113-120.
- Rothschild, G. H. L., 1975. Attractants for monitoring *Pectinophora scutigera* and related species in Australia. *Environ. Entomol.*, **4**: 983-985.
- Rothschild, G. H. L., 1978. Attractants for *Heliothis armigera* and *H. punctigera*. *J. Aust. Entomol. Soc.*, **17**: 389-390.
- Rothschild, G. H. L., B. F. Nesbitt, P. S. Beevor, A. Cork, D. R. Hall and R. A. Vickers, 1982. Studies of the female sex pheromone of the native budworm, *Heliothis punctiger*. *Entomol. Exp. Appl.*, **31**: 395-401.
- Rotundo, G. and E. Tremblay, 1982. Preliminary observations on the sexual pheromone of *Opogona sacchari* (Bojer) (Lepidoptera, Hieroxestidae). *Boll. Lab. Entomol. Agrar. F.S.*, **39**: 123-132.
- Rotundo, G., 1983. Identification of the primary component of the sex pheromone of *Blepharita solieri* (Bdv.) (Lepidoptera, Noctuidae). *Boll. Lab. Entomol. Agrar. F.S.*, **40**: 179-186.
- Rotundo, G., R. Giacometti and A. Rotundo, 1984. Identification of the sex attractants of chestnut tortricids by means of electroantennographic and field studies. *Boll. Lab. Entomol. Agrar. F.S.*, **41**: 211-223.
- Rotundo, G. and E. Tremblay, 1985(a). Field evaluation of female sex pheromone of the artichoke moth, *Gortyna xanthenes*. *Entomol. Exp. Appl.*, **38**: 201-204.
- Rotundo, G. and R. Giacometti, 1985(b). Tortricidae Olethreutini caught with attractant traps in a chestnut-growing area. *Boll. Lab. Entomol. Agrar. F.S.*, **42**: 167-189.
- Rotundo, G. and R. Giacometti, 1988. Individualization of an attractive mixture for males of *Cydia fagiglandana* Z. (Lep. Tortricidae) by means of a field study. *Boll. Lab. Entomol. Agrar. F.S.*, **45**: 81-97.
- Rotundo, G., R. Giacometti and A. de Cristofaro, 1991. Individual

mixtures attractive to males of *Pammene fasciana* L. (Lep., Tortricidae) using electroantennograms and field studies. *Boll. Lab. Entomol. Agrar. F.S.*, **48**: 89-104.

Rotundo, G., G. S. Germinara and A. de Cristofaro, 2004. Chemical electrophysiological and behavioral investigations on the sex pheromone of lackey moth, *Malacosoma neustria*. *J. Chem. Ecol.* **30**: 2057-2069.

Rudzi•ski, K. J., D. Staszek, M. Asztemborska, L. Sukovata, J. Raczko, M. Cie•lak, A. Kolk and R. Szmigielski, 2022. Newly discovered components of *Dendrolimus pini* sex pheromone. *Insects*, **13**: 1063.

Ruiz, K. P., A. Bruce, N. E. Chérémond, C. A. Stratton, E. G. Murrell, S. Gillette and W. R. Morrison III, 2022. Field trapping and flight capacity of *Eucosma giganteana* (Riley) (Lepidoptera: Tortricidae) in response to behaviorally active congeneric semiochemicals in novel silfower agroecosystems. *Insects*, **13**: 350.

Rule, G. S. and W. L. Roelofs, 1989. Biosynthesis of sex pheromone components from linolenic acid in arctiid moths. *Arch. Insect Biochem. Physiol.*, **12**: 89-97.

Ryall, K., P. J. Silk, J. Wu, P. Mayo, M. A. Lemay and D. MaGee, 2010. Sex pheromone chemistry and field trapping studies of the elm spanworm *Ennomos subsignaria* (Hübner) (Lepidoptera: Geometridae). *Naturwissenschaften*, **97**: 717-724.

Safonkin, A. F. and V. V. Buleza, 1988. Role of sexual feromones in interspecific isolation of leafrollers (Lepidoptera: Tortricidae). *Zhurnal Obschiy Biologiya (J. General Biology)*, **49**: 396-400.

Safonkin, A. F. and T. A. Triseleva, 2003. Identification of the sex pheromone components of the apple surface eating tortrix *Archips podana* Scop. (Lepidoptera: Tortricidae). *Agrokhimiya*, (3): 62-64.

Saglio, P., E. Priesner, C. Descoins and M. Gallois, 1977(a). A synthetic sex attractant for *Sparganothis pilleriana* (Schiffermüller). *C. R. Hebd. Séances Acad. Sci., Ser. D*, **284**: 2007-2010.

Saglio, P., C. Descoins, M. Gallois, M. Lettere, D. Jaouen and J. Mercier, 1977(b). Study of the sex pheromone of *Eupoecilia (Clyisia) ambigua* Hb. Lepidoptera Tortricoidea, Cochyliidae. *Ann. Zool. Entomol. Anim.*, **9**: 553-562.

Sanders, C. J., G. E. Daterman, R. F. Shepherd and H. Cerezke, 1974. Sex attractants for two species of western spruce budworm, *Choristoneura biennis* and *C. viridis* (Lepidoptera: Tortricidae). *Can. Entomol.*, **106**: 157-159.

Sanders, C. J. and J. Weatherston, 1976. Sex pheromone of the eastern

spruce budworm (Lepidoptera: Tortricidae): optimum blend of *trans*- and *cis*-tetradecenal. *Can. Entomol.*, **108**: 1285-1290.

Sanders, C.J. 1984. Sex pheromone of the spruce budworm (Lepidoptera: Tortricidae): evidence for a missing component. *Can. Entomol.*, **116**: 93-100.

Sanders, C. J., 1993. Non-target Lepidoptera captured in traps baited with spruce budworm pheromone in Ontario. *Can. Entomol.*, **125**: 1067-1076.

Sans, A., M. Riba, M. Eizaguirre and C. Lopez, 1997. Electroantennogram, wind tunnel and field responses of male Mediterranean corn borer, *Sesamia nonagrioides*, to several blends of its sex pheromone components. *Entomol. Exp. Appl.*, **82**(2): 121-127.

Santangelo, E. M., M. Coracini, P. Witzgall, A. G. Correa and C. R. Unelius, 2002. Identification, syntheses, and characterization of the geometric isomers of 9,11-hexadecadienol from female pheromone glands of the sugar cane borer *Diatraea saccharalis*. *J. Nat. Prod.*, **65**: 909-915.

Sartwell, C., G. E. Daterman and L. L. Sower, 1985. A synthetic attractant for male moths of a biotype in the *Cydia piperana* complex (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 1151-1152.

Sasaerila, Y., G. Gries, G. Khaskin, R. Gries and Hardi, 1997. Identification of sex pheromone components of nettle caterpillar, *Setothosea asigna*. *J. Chem. Ecol.*, **23**(9): 2187-2196.

Sasaerila, Y., R. Gries, G. Gries, G. Khaskin, S. King and T. C. Boo, 2000(a). Decadienoates: sex pheromone components of nettle caterpillars *Darna trima* and *D. bradleyi*. *J. Chem. Ecol.*, **26**(8): 1969-1981.

Sasaerila, Y., R. Gries, G. Gries, G. Khaskin and Hardi, 2000(b). Sex pheromone components of nettle caterpillar, *Setora nitens*. *J. Chem. Ecol.*, **26**(8): 1983-1990.

Sato, R., K. Yaginuma and M. Kumakura, 1979. Sex attractant of the stone leek miner, *Acrolepia alliella* Semenov et Kuznesov: (*Z*)-11-hexadecenyl acetate, (*Z*)-11-hexadecenal and (*Z*)-11-hexadecenol. *Jap. J. Appl. Entomol. Zool.*, **23**: 115-117.

Sato, R., N. Abe, P. Sonnet, H. Sugie and Y. Tamaki, 1985. Biological activity of (*R*)- and (*S*)-14-methyl-1-octadecene, as the chiral component of the sex pheromone of the peach leafminer moth, *Lyonetia clerkella* Linné (Lepidoptera: Lyonetiidae). *Appl. Entomol. Zool.*, **20**: 411-415.

Sato, R., N. Abe, H. Sugie, M. Kato, K. Mori and Y. Tamaki, 1986. Biological activity of the chiral sex pheromone of the peach leafminer moth, *Lyonetia clerkella* Linné (Lepidoptera: Lyonetiidae). *Appl. Entomol. Zool.*, **21**: 478-480.

Sato, Y., S. Takahashi, M. Sakai and T. Kodama, 1980. Attractiveness of the synthetic sex pheromone to the males of the armyworm,

Leucania separata Walker and the lorey leafworm, *Leucania loreyi* Duponchel (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **15**: 334-340.

Saveer, A. M., E. Hatano, A. Wada-Katsumata, R. L. Meagher and C. Schal, 2023. Nonanal, a new fall armyworm sex pheromone component, significantly increases the efficacy of pheromone lures. *Pest Manag. Sci.*, **79**: 2831-2839.

Schaefer, P. W., G. Gries, R. Gries and D. Holden 1999. Pheromone components and diel periodicity of pheromonal communication in *Lymantria fumida*. *J. Chem. Ecol.*, **25**(10): 2305-2312.

Schal, C., R. E. Charlton and R. T. Cardé, 1987. Temporal patterns of sex pheromone titers and release rates in *Holomelina lamae* (Lepidoptera: Arctiidae). *J. Chem. Ecol.*, **13**: 1115-1129.

Schläger, S., F. Beran, A. T. Groot, C. Ulrichs, D. Veit, C. Paetz, B. R. M. Karumuru, R. Srinivasan, M. Schreiner and I. Mewi, 2015. Pheromone blend analysis and cross-attraction among populations of *Maruca vitrata* from Asia and West Africa. *J. Chem. Ecol.*, **41**: 1155-1162.

Schlamp, K. K., R. Gries, G. Khaskin, K. Brown, E. Khaskin, G. J. R. Judd and G. Gries, 2005. Pheromone components from body scales of female *Anarsia lineatella* induce contacts by conspecific males. *J. Chem. Ecol.* **31**(12): 2897-2911.

Schmidt, B. C., J. Roland and D. Wakarchuk, 2003. Evaluation of synthetic pheromones for monitoring forest tent caterpillar (Lepidoptera: Lasiocampidae) populations. *Environ. Entomol.*, **32**(1): 214-219.

Schneider, D., S. Schulz, R. Kittmann and P. Kanagaratman, 1992. Pheromones and glandular structures of both sexes of the weed-defoliator moth *Pareuchaetes pseudoinsulata* Rego Barros (Lep., Arctiidae). *J. Appl. Entomol.*, **113**: 280-294.

Schneider, D., S. Schulz, E. Priesner, J. Ziesmann and W. Francke 1998. Autodetection and chemistry of female and male pheromone in both sexes of the tiger moth *Panaxia quadripunctaria*. *J. Comp. Physiol. A*, **182**: 153-161.

Schönherr, von J., 1972. Attraction of disparlure to the nun moth, *Lymantria monacha* L. *Z. Angew. Entomol.*, **71**: 260-263.

Schwarz, M., J. A. Klun, E. R. Hart, B. A. Leonhardt and J. C. Weatherby, 1983(a). Female sex pheromone of the yellowheaded fireworm, *Acleris minuta* (Lepidoptera: Tortricidae). *Environ. Entomol.*, **12**: 1253-1256.

Schwarz, M., J. A. Klun, B. A. Leonhardt and D. T. Johnson, 1983(b). (*E,Z*)-2,13-Octadecadien-1-ol acetate. A new pheromone structure for sesiid moths. *Tetrahedron Lett.*, **24**: 1007-1010.

Scott, C. D., M. L. Winston, K. N. Slessor, G. G. S. King and G. G.

Grant, 1984. The biology and pheromone-based monitoring of the driedfruit moth, *Vitula edmandsae serratilineella* (Lepidoptera: Pyralidae). *Can. Entomol.*, 116: 1007-1013.

Sekul, A. A. and A. N. Sparks, 1967. Sex pheromone of the fall armyworm moth: isolation, identification, and synthesis. *J. Econ. Entomol.*, 60: 1270-1272.

Seol, K. Y., H. Honda, K. Usui, T. Ando and Y. Matsumoto, 1987. 10,12,14-Hexadecatrienyl acetate: sex pheromone of the mulberry pyralid, *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae). *Agric. Biol. Chem.*, 51: 2285-2287.

Senda, S., T. Omata, Y. Ninomiya, M. Aburatani and T. Takeuchi, 1991. Attractant compositions for fall webworm. *European Patent Application*, EP 474,486.

Sharp, J. L., J. R. McLaughlin, J. James, T. D. Eichlin and J. H. Tunlinson, 1978. Seasonal occurrence of male Sesiidae in north central Florida determined with pheromone trapping methods. *Florida Entomol.*, 61: 245-250.

Shaver, T. N., H. E. Brown, H. J. Williams, L. T. Woods and J. Worley, 1988. Components of female sex pheromone of *Eoreuma loftini* Dyar. *J. Chem. Ecol.*, 14: 391-399.

Shaver, T. N., D. E. Hendricks and J. D. Lopez, Jr., 1989. Influence of (Z)-11-hexadecen-1-ol on field performance of *Heliothis virescens* pheromone in a PVC dispenser as evidenced by trap capture. *J. Chem. Ecol.*, 15: 1637-1644.

Shaver, T. N., H. E. Brown and D. E. Hendricks, 1990. Development of pheromone lure for monitoring field populations of *Eoreuma loftini* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, 16: 2393-2399.

Shearer, P. and H. Riedl, 1994. Comparison of pheromone trap bioassays for monitoring insecticide resistance of *Phyllonorycter elmaella* (Lepidoptera: Gracillariidae). *J. Econ. Entomol.*, 87(6): 1450-1454.

Shibasaki, H., M. Yamamoto, Q. Yan, H. Naka, T. Suzuki and T. Ando, 2013. Identification of the sex pheromone secreted by a nettle moth, *Monema flavescens*, using gas chromatography/Fourier transform infrared spectroscopy. *J. Chem. Ecol.*, 39: 350-357.

Shimuzu, N. and Y. Kuwahara, 2009. Female sex pheromone of a Japanese population of allium leafminer, *Acrolepiopsis sapporensis* (Lepidoptera: Acrolepiidae). *J. Pestic. Sci.*, 34: 181-183.

Shoji, A., Y. Adachi, H. Naka, T. Ando and Y. Arita, 2009. *Nokona regalis* (Lepidoptera, Sesiidae): a late collection record in autumn and a sex attractant. *Trans. Lipid. Soc. Jpn.*, 60: 193-195.

Shorey, H. H., L. K. Gaston and J. S. Roberts, 1965. Sex pheromones of noctuid moths. VI. Absence of behavioral specificity for the female sex pheromones of *Trichoplusia ni* versus *Autographa*

californica, and *Heliothis zea* versus *H. virescens* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **58**: 600-603.

Shry, Y. L. and Y. S. Chow, 1985. Field attraction experiment of the synthetic sex pheromone of the smaller tea tortrix and oriental tea tortrix in Taiwan. *J. Agric. Assoc. China*, **132**: 108.

Siderhurst, M. S., E. B. Jang, A. H. Hara and P. Conant, 2007. n-Butyl (E)-7,9-decadienoate: sex pheromone component of the nettle caterpillar, *Darna pallivitta*. *Entomol. Exp. Appl.*, **125** (1): 63-69.

Silk, P. J., S. H. Tan, C. J. Wiesner, R. J. Ross and G. C. Lonergan, 1980. Sex pheromone chemistry of the eastern spruce budworm, *Choristoneura fumiferana*. *Environ. Entomol.*, **9**: 640-644.

Silk, P. J., C. J. Wiesner, S. H. Tan, R. J. Ross and G. G. Grant, 1982. Sex pheromone chemistry of the western spruce budworm *Choristoneura occidentalis* Free. *J. Chem. Ecol.*, **8**: 351-362.

Silk, P. J., L. P. S. Kuenen and G. C. Lonergan, 1985 (a). A biologically active analogue of the primary sex-pheromone components of spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 257-260.

Silk, P. J., L. P. S. Kuenen, S. H. Tan, W. L. Roelofs, C. J. Sanders and A. R. Alford, 1985 (b). Identification of sex pheromone components of jack pine budworm, *Choristoneura pinus pinus* Freeman. *J. Chem. Ecol.*, **11**: 159-167.

Silk, P. J., E. W. Butterworth, L. P. S. Kuenen, C. J. Northcott and E. G. Kettela, 1988 (a). Sex pheromone of purplestriped shootworm, *Zeiraphera unfortunana* Powell. *J. Chem. Ecol.*, **14**: 1417-1425.

Silk, P. J. and L. P. S. Kuenen, 1988 (b). Sex pheromones and behavioral biology of the coniferophagous *Choristoneura*. *Ann. Rev. Entomol.*, **33**: 83-101.

Silk, P. J., E. W. Butterworth, L. P. S. Kuenen, C. J. Northcott, E. Dunkelblum and E. G. Kettela, 1989. Identification of sex pheromone component of spruce budmoth *Zeiraphera canadensis*. *J. Chem. Ecol.*, **15**: 2435-2444.

Silk, P. J., L. P. S. Kuenen, A. R. Alford, G. C. Lonergan and G. G. Grant, 1997. Sex pheromone components of the oak leaf shredder, *Croesia semipurpurana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **129**: 1001-1008.

Silk, P. J., E. Eveleigh, L. Roscoe, K. Burgess, S. Weatherby, G. Leclair, P. Mayo, M. Brophy, 2017. Unsaturated cuticular hydrocarbons enhance responses to sex pheromone in spruce budworm, *Choristoneura fumiferana*. *J. Chem. Ecol.*, **43**: 753-762.

Silva, E. C. A. D., P. J. Silk, P. Mayo, N. K. Hillier, D. Magee and G. C. Cutler, 2013. Identification of sex pheromone components

of blueberry spanworm *Itame argillacea* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **39**: 1169-1181.

Silva, M. R. da, A. M. P. Cortes, G. P. Svensson, C. Löfstedt, E. R. Limac and P. H. G. Zarbin, 2021. Identification of two additional behaviorally active gland constituents of female *Diatraea saccharalis* (Fabricius) (Lepidoptera: Crambidae). *J. Braz. Chem. Soc.*, **32**: 225-230.

Simmons, R. B., W. E. Conner and R. B. Davidson, 1998. Identification of major components of the female pheromone glands of *Euchaetes egle* Drury (Arctiidae). *J. Lepid. Soc.*, **52**: 356-363.

Sinclair, E. R. and P. Sinclair, 1980. Trapping adult macadamia nut borer, *Cryptophlebia ombrodelta* (Lower) (Lepidoptera: Tortricidae). *J. Aust. Entomol. Soc.*, **19**: 211-216.

Slessor, K. N., J. Raine, G. G. S. King, S. J. Clements and S. A. Allan, 1987. Sex pheromone of blackheaded fireworm, *Rhopobota naevana* (Lepidoptera: Tortricidae), a pest of cranberry. *J. Chem. Ecol.*, **13**: 1163-1170.

Smith, R. G., G. E. Daterman, G. D. Daves, Jr., K. D. McMurtrey and W. L. Roelofs, 1974. Sex pheromone of the European pine shoot moth: chemical identification and field tests. *J. Insect Physiol.*, **20**: 661-668.

Smith, R. G., G. E. Daterman and G. D. Daves, Jr., 1975. Douglas-fir tussock moth: sex pheromone identification and synthesis. *Science*, **188**: 63-64.

Smith, L. M., R. G. Smith, T. M. Loehr and G. D. Daves, Jr., 1978. Douglas fir tussock moth pheromone: identification of a diene analogue of the principal attractant and synthesis of stereochemically defined 1,6-, 2,6-, and 3,6-heneicosadien-11-ones. *J. Org. Chem.*, **43**: 2361-2366.

Smith, M. T., L. M. McDonough, S. Voerman, H. G. Davis, R. D. Eikenbary and C. L. Smithhisler, 1987. Hickory shuckworm *Cydia caryana*: electroantennogram and flight tunnel studies of potential sex pheromone components. *Entomol. Exp. Appl.*, **44**: 23-30.

Snow, J. W., T. D. Eichlin and J. H. Tunlinson, 1985. Seasonal captures of clearwing moths (Sesiidae) in traps baited with various formulations of 3,13-octadecadienyl acetate and alcohol. *J. Agric. Entomol.*, **2**: 73-84.

Snow, J. W., M. Schwarz and J. A. Klun, 1987. The attraction of the grape root borer, *Vitacea polistiformis* (Harris) (Lepidoptera: Sesiidae) to (*E,Z*)-2,13-octadecadienyl acetate and the effects of related isomers on attraction. *J. Entomol. Sci.*, **22**: 371-374.

Snow, J. W., M. Schwarz and T. D. Eichlin, 1989. Captures of clearwing moths (Lepidoptera: Sesiidae) with various octadecadienyl acetates and alcohols in central Georgia during 1983-1985. *Environ. Entomol.*, **18**: 216-222.

Snow, W. J., D. T. Johnson and J. R. Meyer, 1991. The seasonal occurrence of the grape root borer, (Lepidoptera: Sesiidae) in the eastern United States. *J. Entomol. Sci.*, 26: 157-168.

Soares, A. M. L., P. H. B. França, M. F. Triana, J. M. D. Santos, N. S. Dias-Pini, H. F. Goulart, J. X. Araújo-Júnior and A. E. G. Santana, 2020. Identification of Δ 6-unsaturated, monoenyl type I pheromone compounds from the cashew stem borer *Anthistarcha binocularis* (Lepidoptera: Gelechiidae). *Pest Manag. Sci.*, 76: 1435-1442.

Soderstrom, E. L., D. G. Brandl, J. Myerson, R. G. Buttery and B. E. Mackey, 1985. Sex pheromone for attracting western grapeleaf skeletonizer (Lepidoptera: Zygaenidae). *J. Econ. Entomol.*, 78: 799-801.

Soldán, T., and Spitzer, K. 1983. Some moths recorded at sex pheromone traps in Mitidja, Algeria (Lepidoptera: Tortricidae, Pyralidae, Noctuidae). *Acta Entomol. Bohemoslov.*, 80: 395-398.

Solomon, J. D., M. E. Dix and R. E. Doolittle, 1978. Attractiveness of the synthetic carpenterworm sex attractant increased by isomeric mixtures and prolonged by preservatives. *Environ. Entomol.*, 7: 39-41.

Solomon, J. D., F. L. Oliveria, J. H. Tunlinson and R. E. Doolittle, 1982. Occurrence of clearwing borers (Sesiidae) in west central Mississippi. *J. Georgia Entomol. Soc.*, 17: 4-12.

Song, Y. H., H. Y. Song, H. K. Kim, Y. D. Chang and P. C. Lippold, 1978. Studies on the synthetic pheromones of striped rice borer and tortricid insect pests. *Kor. J. Plant Prot.*, 17: 41-47.

Sower, L. L., G. E. Daterman, C. Sartwell and H. T. Cory, 1979. Attractants for the Western pine shoot borer, *Eucosma sonomana*, and *Rhyacionia zozana* determined by field screening. *Environ. Entomol.*, 8: 265-267.

Spitzer, K., M. Rejmánek, T. Soldán and J. Zelený, 1983. Flight activity of some moths (Lepidoptera) monitored by sex pheromones in Vietnam: dry season. *Acta Entomol. Bohemoslov.*, 80: 413-418.

Sreng, I., B. Maume and B. Frérot, 1985. Analysis of the pheromonal secretion of virgin females of *Sesamia nonagrioides* (Lef.) (Lepidoptera, Noctuidae). *C. R. Acad. Sci., Ser. 3*, 301: 439-442.

Stan, G. and L. Pop, 1992. Attractiveness of pheromone variants with (Z)-11-hexadecenyl acetate for males of some lepidopteran species. *Soc. Lepid. Rom. Bull. Inf.*, 10: 10-15.

Starratt, A. N., K. H. Dahm, N. Allen, J. G. Hildebrand, T. L. Payne and H. Röller, 1979. Bombykal, a sex pheromone of the sphinx moth *Manduca sexta*. *Z. Naturforsch.*, 34 c, 9-12.

- Steck, W. F., B. K. Bailey, E. W. Underhill and M. D. Chisholm, 1976. A sex attractant for the great dart, *Eurois occulta*: a mixture of (*Z*)-9-tetradecen-1-ol acetate and (*Z*)-11-hexadecen-1-ol acetate. *Environ. Entomol.*, 5: 523-526.
- Steck, W., E. W. Underhill, M. D. Chisholm, B. K. Bailey, J. Loeffler and C. G. Devlin, 1977(a). Sex attractants for males of 12 moth species found in Western Canada. *Can. Entomol.*, 109: 157-160.
- Steck, W., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1977(b). A sex attractant for male moths of the glassy cutworm *Crymodes devastator* (Brace): a mixture of *Z*-11-hexadecen-1-yl acetate, *Z*-11-hexadecenal and *Z*-7-dodecen-1-yl acetate. *Environ. Entomol.*, 6: 270-273.
- Steck, W., B. K. Bailey, M. D. Chisholm and E. W. Underhill, 1978. A sex attractant for males of the cutworm *Euxoa pleuritica* (Lepidoptera: Noctuidae). *Can. Entomol.*, 110: 775-777.
- Steck, W. F., D. L. Struble, C. E. Lilly, M. D. Chisholm, E. W. Underhill and G. E. Swailes, 1979(a). A sex attractant for males of the early cutworm, *Euxoa tristicula* (Lepidoptera: Noctuidae). *Can. Entomol.*, 111: 337-341.
- Steck, W. F., E. W. Underhill, M. D. Chisholm and H. S. Gerber, 1979(b). Sex attractant for male alfalfa looper moths, *Autographa californica*. *Environ. Entomol.*, 8: 373-375.
- Steck, W. F., M. D. Chisholm, B. K. Bailey and E. W. Underhill, 1979(c). Moth sex attractants found by systematic field testing of 3-component acetate-aldehyde candidate lures. *Can. Entomol.*, 111: 1263-1269.
- Steck, W. F., E. W. Underhill, M. D. Chisholm and J. R. Byers, 1979(d). Sex attractants for *Agrotis venerabilis* and *Euxoa albipennis* based on (*Z*)-5-decanyl acetate and (*Z*)-7-dodecanyl acetate. *Environ. Entomol.*, 8: 1126-1128.
- Steck, W., E. W. Underhill and M. D. Chisholm, 1980(a). Trace components in lepidopterous sex attractants. *Environ. Entomol.*, 9: 583-585.
- Steck, W., M. D. Chisholm, E. W. Underhill and C. C. Peters, 1980(b). Optimized conditions for sex attractant trapping male redbacked cutworm moths, *Euxoa ochrogaster* (Guenée). *J. Chem. Ecol.*, 6: 585-591.
- Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1980(c). Improved sex attractant blend for adult males of the glassy cutworm, *Crymodes devastator* (Lepidoptera: Noctuidae). *Can. Entomol.*, 112: 751-752.
- Steck, W., E. W. Underhill and M. D. Chisholm, 1982(a). Structure-activity relationships in sex attractants for North American noctuid moth. *J. Chem. Ecol.*, 8: 731-754.
- Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1982(b). Trace co-attractants in synthetic sex lures for 22 noctuid moths.

Experientia, 38: 94-96.

Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1982 (c). (Z)-7-Tetradecenal, a seasonally dependent sex pheromone of the w-marked cutworm, *Spaelotis clandestina* (Harris) (Lepidoptera: Noctuidae). *Environ. Entomol.*, 11: 1119-1122.

Steck, W., E. W. Underhill, M. D. Chisholm and B. K. Bailey, 1982 (d). (Z)-5-Tetradecenyl acetate, a trapping synergist for the major sex pheromone of the spotted cutworm, *Amathes c-nigrum*. *Entomol. Exp. Appl.*, 31: 328-330.

Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1982 (e). A 4-component sex attractant for male moths of the armyworm, *Pseudaletia unipuncta*. *Entomol. Exp. Appl.*, 32: 302-304.

Steck, W. F., E. W. Underhill and P. Palaniswamy, 1986. Sex pheromone of the cutworm *Feltia jaculifera* (Lepidoptera: Noctuidae). *Environ. Entomol.*, 15: 40-43.

Steinbauer, M. J., F. Östrand, T. E. Bellas, A. Nilsson, F. Andersson, E. Hedenström, M. J. Lacey and F. P. Schiestl, 2004. Identification, synthesis and activity of sex pheromone gland components of the autumn gum moth (Lepidoptera: Geometridae), a defoliator of *Eucalyptus*. *Chemoecology*, 14: 217-223.

Stevens, R. E., C. Sartwell, T. W. Koerber, G. E. Daterman, L. L. Sower and J. A. Powell, 1980. Western *Rhyacionia* (Lepidoptera: Tortricidae, Olethreutinae) pine tip moths trapped using synthetic sex attractants. *Can. Entomol.*, 112: 591-603.

Stevens, R. E., C. Sartwell, T. W. Koerber, J. A. Powell, G. E. Daterman and L. L. Sower, 1985. Forest tortricids trapped using *Eucosma* and *Rhyacionia* synthetic sex attractants. *J. Lepidop. Soc.*, 39: 26-32.

Stockel, J. and P. Anglade, 1977. Influence de la concentration en isomères cis et trans dans la phéromone sexuelle d'*Ostrinia nubilalis* Hb. sur le comportement d'orientation de cette espèce et sur celui de *Pyrausta aurata* Sc. *C. R. Hebd. Séances Acad. Sci., Ser. D*, 285: 61-64.

Strong, W. B., J. G. Millar, G. G. Grant, J. A. Moreira, J. M. Chong and C. Rudolph, 2008. Optimization of pheromone lure and trap design for monitoring the fir coneworm, *Dioryctria abietivorella*. *Entomol. Exp. Appl.*, 126: 67-77.

Struble, D. L. and G. E. Swailes, 1975 (a). A sex attractant for the clover cutworm, *Scotogramma trifolii* (Rottenberg), a mixture of Z-11-hexa-decen-1-ol acetate and Z-11-hexadecen-1-ol. *Environ. Entomol.*, 4: 632-636.

Struble, D. L., M. Jacobson, N. Green and J. D. Warthen, 1975 (b). Bertha armyworm (Lepidoptera: Noctuidae): detection of a sex pheromone and the stimulatory effect of some synthetic chemicals. *Can. Entomol.*, 107: 355-359.

- Struble, D. L., G. E. Swailes, W. F. Steck, E. W. Underhill and M. D. Chisholm, 1976. A sex attractant for the adult males of variegated cutworm, *Peridroma saucia*. *Environ. Entomol.*, 5: 988-990.
- Struble, D. L. and C. E. Lilly, 1977(a). An attractant for the beet webworm, *Loxostege sticticalis* (Lepidoptera: Pyralidae). *Can. Entomol.*, 109: 261-266.
- Struble, D. L. and G. E. Swailes, 1977(b). A sex attractant for the adult males of the army cutworm, *Euxoa auxiliaris*: a mixture of Z-5-tetradecen-1-yl acetate and E-7-tetradecen-1-yl acetate. *Environ. Entomol.*, 6: 719-724.
- Struble, D. L., G. E. Swailes and G. L. Ayre, 1977(c). A sex attractant for males of the darksided cutworm, *Euxoa messoria* (Lepidoptera: Noctuidae). *Can. Entomol.*, 109: 975-980.
- Struble, D. L., G. E. Swailes, W. F. Steck, E. W. Underhill and M. D. Chisholm, 1977(d). A sex attractant for *Leucania commoides*: a mixture of Z-9-tetradecen-1-yl acetate, Z-11-hexadecen-1-yl acetate, and Z-11-hexadecen-1-ol. *Can. Entomol.*, 109: 1393-1398.
- Struble, D. L. and G. E. Swailes, 1978(a). A sex attractant for adult males of the pale western cutworm, *Agrotis orthogonia* (Lepidoptera: Noctuidae). *Can. Entomol.*, 110: 769-773.
- Struble, D. L. and G. E. Swailes, 1978(b). Sex attractant for adult males of the redbacked cutworm, *Euxoa ochrogaster*. *Environ. Entomol.*, 7: 384-386.
- Struble, D. L., H. R. Buser, H. Arn and G. E. Swailes, 1980(a). Identification of sex pheromone components of redbacked cutworm, *Euxoa ochrogaster*, and modification of sex attractant blend for adult males. *J. Chem. Ecol.*, 6: 573-584.
- Struble, D. L., H. Arn, H. R. Buser, E. Städler and J. Freuler, 1980(b). Identification of 4 sex pheromone components isolated from calling females of *Mamestra brassicae*. *Z. Naturforsch.*, 35 c: 45-48.
- Struble, D. L., 1981(a). Modification of the attractant blend for adult males of the army cutworm, *Euxoa auxiliaris* (Grote), and the development of an alternate 3-component attractant blend for this species. *Environ. Entomol.*, 10: 167-170.
- Struble, D. L., 1981(b). A four-component pheromone blend for optimum attraction of redbacked cutworm males, *Euxoa ochrogaster* (Guenée). *J. Chem. Ecol.*, 7: 615-625.
- Struble, D. L., W. F. Steck, G. E. Swailes, M. D. Chisholm, E. W. Underhill and C. E. Lilly, 1982. Two two-component sex attractant blends for adult males of the striped cutworm, *Euxoa tessellata* (Lepidoptera: Noctuidae). *Can. Entomol.*, 114: 359-362.
- Struble, D. L., 1983(a). Sex pheromone components of *Euxoa drewseni*: chemical identification, electrophysiological evaluation, and

field attractancy tests. *J. Chem. Ecol.*, **9**: 335-346.

Struble, D. L. and K. W. Richards, 1983 (b). Identification of sex pheromone components of the female driedfruit moth, *Vitula edmandsae serratilineella*, and a blend for attraction of male moths. *J. Chem. Ecol.*, **9**: 785-801.

Struble, D. L., G. L. Ayre and J. R. Byers, 1984. Minor sex-pheromone components of *Mamestra configurata* (Lepidoptera: Noctuidae) and improved blends for attraction of male moths. *Can. Entomol.*, **116**: 103-105.

Struble D. L. and J. R. Byers, 1985. Identification of sex-pheromone components of the sibling species *Euxoa ridingsiana* (Grt.) and *Euxoa maimes* (Sm.) (Lepidoptera: Noctuidae), and blends for their specific attraction. *Can. Entomol.*, **117**: 495-504.

Struble, D. L., J. R. Byers, D. G. R. McLeod and G. L. Ayre, 1987 (a). Sex pheromone components of an Alberta population of European corn borer, *Ostrinia nubilalis* (Hbn.) (Lepidoptera: Pyralidae). *Can. Entomol.*, **119**: 291-299.

Struble, D. L. and J. R. Byers, 1987 (b). Identification of sex pheromone components of darksided cutworm, *Euxoa messoria*, and modification of sex attractant blend for adult males. *J. Chem. Ecol.*, **13**: 1187-1199.

Struble, D. L., G. L. Ayre and J. R. Byers, 1987 (c). Sex attractant blends for strawberry cutworm, *Amphipoea interoceanica* (Smith), and a closely related species, *Amphipoea americana* (Speyer) (Lepidoptera: Noctuidae). *Can. Entomol.*, **119**: 301-303.

Struble, D. L., J. R. Byers, R. F. Shepherd and T. G. Gray, 1989. Identification of sex pheromone components of the black army cutworm, *Actebia fennica* (Tauscher) (Lepidoptera: Noctuidae), and a sex attractant blend for adult males. *Can. Entomol.*, **121**: 557-563.

Su, M.-W., Y.-L. Fang, W.-Q. Tao, G.-Z. Yan, W.-E. Ma and Z.-N. Zhang, 2008. Identification and field evaluation of the sex pheromone of an invasive pest, the fall webworm *Hyphantria cunea* in China. *Chinese Sci. Bull.*, **53**: 555-560.

Subchev, M. A. and N. V. Kuznetzov, 1974. Preliminary identification of the sex pheromone of *Rhynchopacha* sp. (Lepidoptera: Gelechiidae) by field screening tests. *Dokl. Bolg. Akad. Nauk. (C. R. Acad. Blug. Sci.)*, **27**: 993-994.

Subchev, M. A., 1981. Z-11-Hexadecenyl acetate: sexual attractant for males of eight species of moths (Insecta, Lepidoptera). *Ekologiya*, **9**: 59-62.

Subchev, M. A. and I. L. Stoilov, 1984. Co-attractants for *Mamestra brassicae* L. and *Mamestra suasa* Denis & Schiff. (Lepidoptera: Noctuidae). *Dokl. Bolg. Akad. Nauk. (C. R. Acad. Blug. Sci.)*, **37**: 353-354.

- Subchev, M. A., J. A. Ganev, O. Vostrowsky and H. J. Bestmann, 1986 (a). Screening and use of sex attractants in monitoring of Geometrid moths in Bulgaria. *Z. Naturforsch.*, **41 c**: 1082-1086.
- Subchev, M. A., J. A. Ganev and I. L. Stoilov, 1986 (b). Noctuid male sex attractants - *cis*-11-hexadecenyl compounds alone and in mixtures. *Ekologiya*, **18**: 33.
- Subchev, M. A., L.S. Stanimirova and T. S. Milkova, 1987. The effect of compounds related to *cis*-11-hexadecenyl acetates on its attractiveness to the males of *Mamestra brassicae* L. (Lepidoptera: Noctuidae) and some other noctuid species. *Folia Biol.*, **35**: 143-150.
- Subchev, M. A., I. A. Krusteva, J. A. Ganev and T. S. Milkova, 1990. Some new lepidopteran sex attractants and attractant inhibitors in Bulgaria. *J. Appl. Entomol.*, **109**: 189-193.
- Subchev, M., 1992. Sex attractant for *Trachea atriplicis* L. (Lepidoptera: Noctuidae) males. *Dokl. Bulg. Akad. Nauk. (C. R. Acad. Blag. Sci.)*, **45**: 121-123.
- Subchev, M. and L. Pop, 1994. Aldehyde sex attractants for *Agrochola* (Lepidoptera: Noctuidae) species. *Comptes-rendus de l'Académie bulgare des sciences*, **47** (4): 101-104.
- Subchev, M., A. Botar and S. Voerman, 1995. Sex attractant and attractant inhibitors for *Atethmia ambusta* F. (Lep., Noctuidae). *J. Appl. Entomol.*, **119**: 207-209.
- Subchev, M. A., M. Tóth, G. Szöcs, G. Stan and A. Botar, 1996. Evidence for geographical differences in pheromonal responses of male *Amathes c-nigrum* L. (Lep., Noctuidae). *J. Appl. Entomol.*, **120**: 615-617.
- Subchev, M., A. Harizanov, W. Francke, S. Franke, E. Plass, A. Reckziegel, F. Schröder, J. A. Pickett, L. J. Wadhams and C. M. Woodcock 1998. Sex pheromone of female vine bud moth, *Theresimima ampelophaga* comprises (2S)-butyl (7Z)-tetradecenoate. *J. Chem. Ecol.*, **24** (7): 1141-1151. [Erratum: (2S)-butyl → (2R)-butyl, *J. Chem. Ecol.*, **25**: 1203]
- Subchev, M. A., M. Toth, D. Wu and Z. Imrei, 1999. (Z)-8-Tridecenyl acetate - A sex attractant for three microlepidopteran species (Lepidoptera: Cosmopterigidae, Symmocidae). *Dokl. Bulg. Akad. Nauk. (C. R. Acad. Blag. Sci.)*, **52** (7-8): 113-114.
- Subchev, M., T. Toshova, L. Stanimirova, G. Stan, G. Embacher, W. Francke, A. Reckziegel, J. T. Ferreira and E. Priesner, 2000 (a). 1-Methylhexyl octanoate, a new lepidopteran sex pheromone from the bagworm, *Megalophanes viciella*. *J. Chem. Ecol.*, **26** (2): 487-495.
- Subchev, M., M. Toth, D. Wu, L. Stanimirova, T. Toshova and Z. Karpati, 2000 (b). Sex attractant for *Diloba caeruleocephala* (L.), (Lep., Dilobidae) (Z)-8-tridecenyl acetate. *J. Appl. Entomol.*, **124**:

197-199.

Subchev, M., A. Mircheva, J. Pickett, L. Wadhams, C. Woodcock, A. D. Santos, S. Franke and W. Francke, 2003. Sex pheromone of the leaf-miner *Phyllonorycter platani*: (Z10)-tetradecenyl acetate. *J. Chem. Ecol.*, **29**(10): 2391-2396.

Subchev, M., T. Toshova, C. Koshio, S. Franke, A. Tröger, R. Twele, W. Francke, J. A. Pickett, L. J. Wadhams and C. M. Woodcock, 2009. Identification and biological activity of sex pheromone components from females of the plumm moth *Illiberis rotundata* Jordan (Lepidoptera: Zygaenidae: Procridinae). *Chemoecol.*, **19**: 47-54.

Subchev, M. A., K. A. Efetov, T. B. Toshova, E. V. Parshkova, W. Tóth and W. Francke, 2010. New sex attractants for species of the zygaenid subfamily Procridinae (Lepidoptera: Zygaenidae). *Entomol. Gen.*, **32**: 243-250.

Subchev, M., C. Koshio, T. Toshova, K. A. Efetov and W. Francke, 2013. (2R)-butyl (7Z)-dodecenoate, a main sex pheromone component of *Illiberis (Primilliberis) pruni* Dyar (Lepidoptera: Zygaenidae: Procridinae)? *Acta Zool. Bulg.*, **65**: 391-396.

Subchev, M., 2014. Sex pheromone communication in the family Zygaenidae (Insecta: Lepidoptera): a review. *Acta Zool. Bulg.*, **66**: 147-157.

Suckling, D. M., A. R. Gibb, H. Gourlay, P. Conant, C. Hirayama, R. Leen and G. Szocs, 2000. Sex attractant for the gorse biocontrol agent *Agonopterix ulicetella* (Oecophoridae). *Proc. N. Z. Plant Prot. Conf.*, **53**: 66-70. [*Chem. Abst.*, **135**: 57326]

Suckling, D. M., A. R. Gibb, J. M. Daly, D. J. Rogers and G. P. Walker, 2002. Improving the pheromone lure for diamondback moth. *NZ Plant Prot.*, **55**: 182-187.

Suckling, D. M., A. R. Gibb, G. M. Burnip, C. Snelling, J. De Rutter, G. Langford and A. M. El-Sayed, 2005(a). Optimization of pheromone lure and trap characteristics for currant clearwing, *Synanthedon tipuliformis*. *J. Chem. Ecol.*, **31**(2): 393-406.

Suckling, D. M., A. R. Gibb, P. R. Dentener, D. S. Seldon, G. K. Clare, L. Jamieson, D. Baird, D. J. Kriticos and A. M. El-Sayed, 2005(b). *Uraba lugens* (Lepidoptera: Nolidae) in New Zealand: pheromone trapping for delimitation and phenology. *J. Econ. Entomol.*, **98**: 1187-1192.

Sugie, H., K. Yaginuma, Y. Tamaki, 1977. Sex pheromone of the Asiatic leafroller, *Archippus breviplicanus* Walsingham (Lepidoptera: Tortricidae): isolation and identification. *Appl. Entomol. Zool.*, **12**: 69-74.

Sugie, H., Y. Tamaki, R. Sato and M. Kumakura, 1984(a). Sex pheromone of the peach leafminer moth, *Lyonetia clerkella* Linné: isolation and identification. *Appl. Entomol. Zool.*, **19**: 323-330.

- Sugie, H., Y. Tamaki, S. Shirasaki and T. Kitamura, 1984 (b). Further studies on the sex pheromone of the summer fruit tortrix moth, *Adoxophyes orana fasciata*. *Jap. J. Appl. Entomol. Zool.*, **28**: 156-160.
- Sugie, H., Y. Tamaki, K. Kawasaki, M. Wakou, T. Oku, C. Hirano and M. Horike, 1986. Sex pheromone of the apple leafminer moth, *Phyllonorycter ringoniella* (Matsumura) (Lepidoptera: Gracillariidae): activity of geometrical isomers of tetradecadienyl acetates. *Appl. Entomol. Zool.*, **21**: 578-581.
- Sugie, H., K. Kawasaki, H. Higuchi, C. Kitamura and Y. Tamaki, 1990. Identification of the sex pheromone of the three-spotted plusia, *Acanthoplusia agnata* Staudinger (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **25**: 467-473.
- Sugie, H., K. Kawasaki, S. Nakagaki and N. Iwata, 1991 (a). Identification of sex pheromone of the semi-looper, *Autographa nigrisigna* Walker (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **26**: 71-76.
- Sugie, H., S. Tatsuki, S. Nakagaki, C. B. J. Rao and A. Yamamoto, 1991 (b). Identification of the sex pheromone of the oriental tobacco budworm, *Heliothis assulta* (Guenée) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **26**: 151-153.
- Sugie, H., H. Noguchi, K. Shimizu, T. Kawana, H. Fukuda, A. Yamamoto, T. Fukumoto and T. Ihara, 1993. Attractants containing sex pheromones for *Mamestra brassicae*. *Jpn. Kokai Tokkyo Koho* JP 05,255,009. [*Chem. Abst.* vol.120: No.7 p71571u]
- Sugie, H., H. Noguchi, K. Shimizu, T. Kawana, H. Fukuda, A. Yamamoto, T. Fukumoto and T. Ihara, 1994. Insect attractants containing Z-11-hexadecenal, Z-9-hexadecenal, and Z-11-hexadecenol for *Pediasia teterrellus*. *Jpn. Kokai Tokkyo Koho* JP 06 56,608. [*Chem. Abst.* vol.121: No.3 p29291y]
- Sugie, H., J. Yase, K. Fujita and Y. Shirai, 2003. A sex attractant of the cabbage webworm, *Hellula undalis* Fabricius (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **38**(1): 45-48.
- Svatos**, A., A. B. Attygalle, G. N. Jham, R. T. S. Frighetto, E. F. Vilela, D. Saman and J. Meinwald, 1996. Sex pheromone of tomato pest *Scrobipalpuloides absoluta* (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **22**: 787-800
- Svatos, A., B. Kalinová, M. Oskovee, J. Kindl, O. Hovorka and I. Hrdý, 1999. Identification of a new Lepidopteran sex pheromone in picogram quantities using an antennal biodetector: (8E,10Z)-tetradeca-8,10-dienal from *Cameraria ohridella*. *Tetrahedron Lett.*, **40**: 7011-7014.
- Svensson**, M. G. E., M. Bengtsson and J. Löfqvist, 1997. Individual variation and repeatability of sex pheromone emission of female turnip moths *Agrotis segerum*. *J. Chem. Ecol.* **23**: 1833-1850.

Svensson, G. P., V. Tönnberg and L. Sigsgaard, 2019. Identification and field evaluation of (*E*)-11,13-tetradecadienal as sex pheromone of the strawberry tortrix (*Acleris comariana*). *J. Appl. Entomol.*, **143**: 535-541.

Svensson, G. P., O. Anderbrant, E. Öberg, E. V. Jirle, S. Hellqvist and C. Löfstedt, 2023. Identification of (*E*)- and (*Z*)-11-tetradecenyl acetate as sex pheromone components of the currant pest *Euhypomeutoides albithoracellus*. *J. Appl. Entomol.*, **147**: 313-319.

Sweeney, J. D., J. A. McLean and L. M. Friskie, 1990. Roles of minor components in pheromone-mediated behavior of western spruce budworm moths. *J. Chem. Ecol.*, **16**: 1517-1530.

Sziráki, G., 1978. Examinations on tortricid moths trapped by synthetic attractants (Lepidoptera). *Folia Entomol. Hung.*, **31**: 259-264.

Sziráki, G., 1980. Notes on Coleophora and Cnephasia species trapped by synthetic attractants (Lepidoptera: Coleophoridae and Tortricidae). *Folia Entomol. Hung.*, **33**: 161-166.

Szöcs, G., M. Tóth and L. Novák, 1981. Sex attractants for eight lepidopterous species. *Z. Angew. Entomol.*, **91**: 272-280.

Szöcs, G., M. Tóth and L. Novák, 1983. Sex attractants and inhibitors for lepidopterous species found by field screening of olefinic compounds in Hungary. *Z. Angew. Entomol.*, **96**: 56-67.

Szöcs, G., M. Tóth, H. J. Bestmann and O. Vostrowsky, 1984. A two-component sex attractant for males of the geometrid moth *Alsophila quadripunctata*. *Entomol. Exp. Appl.*, **36**: 287-291.

Szöcs, G., M. Schwarz, G. Sziraki, M. Toth, J. A. Klun and B. A. Leonhardt, 1985. Sex pheromone of the female currant borer, *Synanthedon tipuliformis*: identification and field evaluation. *Entomol. Exp. Appl.*, **39**: 131-133.

Szöcs, G., M. Tóth, H. J. Bestmann, O. Vostrowsky, R. R. Heath and J. H. Tunlinson, 1987. Polyenic hydrocarbons as sex attractants for geometrids and amatids (Lepidoptera) found by field screening in Hungary. *Z. Naturforsch.*, **42 c**: 165-168.

Szöcs, G., M. Tóth, Gy. Sziráki and M. Schwarz, 1989. 2,13- and 3,13-Octadecadienyl compounds composing sex attractants for tineid and sesiid moths (Lepidoptera). *Biochem. System. Ecol.*, **17**: 417-422.

Szöcs, G., L. A. Miller, W. Thomas, R. A. Vickers, G. H. L. Rothschild, M. Schwarz and M. Tóth, 1990. Compounds modifying male responsiveness to main female sex pheromone component of the currant borer, *Synanthedon tipuliformis* Clerk (Lepidoptera: Sesiidae) under field conditions. *J. Chem. Ecol.*, **16**: 1289-1305.

- Szöcs, G., V. Bu'da, P. Charmillot, P. Esbjerg, B. Freier, R. Gottwald, B. Kovalev, S. Maini, M. G. Solomon, O. Sorum, M. Subchev, M. Tóth and M. Van de Veire, 1991. Field tests of (*E,Z*)-3,13-octadecadien-1-ol acetate: a sex attractant synergist for male currant borer, *Synanthedon tipuliformis*. *Entomol. Exp. Appl.*, **60**: 283-288.
- Szöcs, G., A. Raina, M. Tóth and B. A. Leonhardt, 1993 (a). Sex pheromone components of *Heliothis maritima*: chemical identification, flight tunnel and field tests. *Entomol. Exp. Appl.*, **66**: 247-253.
- Szöcs, G., M. Tóth, W. Francke, F. Schmidt, P. Philipp, W. A. König, K. Mori, B. S. Hansson and C. Löfstedt, 1993 (b). Species discrimination in five species of winter-flying geometrids (Lepidoptera) based on chirality of semiochemicals and flight season. *J. Chem. Ecol.*, **19**: 2721-2735.
- Szöcs, G., M. Tóth, Z. Kárpáti, J. Zhu, C. Löfstedt, E. Plass and W. Francke, 2004. Identification of polyenic hydrocarbons from the northern winter moth, *Operophtera fagata*, and development of a species specific lure for pheromone traps. *Chemoecology*, **14**(1): 53-58.
- Szöcs, G., M. Tóth and K. Mori, 2005. Absolute configuration of the major sex pheromone component of the satin moth, *Leucoma salicis*, verified by field trapping test in Hungary. *Chemoecology*, **15**(2): 127-128.
- Tabata**, J., T. Takanashi and Y. Ishikawa, 2003. Pheromone analysis of wild female moths with a PBAN C-terminal peptide injection for an estimation of assortative mating in adzuki bean borer, *Ostrinia scapulalis*. *J. Chem. Ecol.*, **29**(12): 2749-2759.
- Tabata, J., S. Hoshizaki, S. Tatsuki and Y. Ishikawa, 2006. Heritable pheromone blend variation in a local population of the butterbur borer moth *Ostrinia zaguliaevi* (Lepidoptera: Crambidae). *Chemoecology* **16**: 123-128.
- Tabata, J., T. Yokosuka, M. Hattori, M. Ohashi, H. Noguchi and H. Sugie, 2008 (a). Sex attractant pheromone of the limabean pod borer, *Etiella zinckenella* (Treitschke) (Lepidoptera: Pyralidae), in Japan. *Appl. Entomol. Zool.*, **43**: 351-358.
- Tabata, J., Y. Huang, S. Ohno, Y. Yoshiyasu, H. Sugie, S. Tatsuki and Y. Ishikawa, 2008 (b). Sex pheromone of *Ostrinia* sp. newly found on the leopard plant *Farfugium japonicum*. *J. Appl. Entomol.*, **132**: 566-574.
- Tabata, J., M. Minamishima, H. Sugie, T. Fukumoto, F. Mochizuki and Y. Yoshiyasu, 2009. Sex pheromone components of the pear fruit moth, *Acrobasis pyrivorella* (Matsumura). *J. Chem. Ecol.*, **35**: 243-249.
- Tabata, J. and S. Kubota, 2017. Sex pheromone of the iris gelechiid moth *Monochroa divisella*. *Entomol. Exp. Appl.*, **164**: 113-119.

Tabata, J., R. Nakano, H. Yasui, K. Nakamura, K. Takehara, H. Matsuda, Y. Ikenoue, Y. Kusuhata, K. Kinjo, K. Nakama, Y. Tsuha and Y. Ishikawa, 2023. Sex pheromone of the fall armyworm, *Spodoptera frugiperda*: identification of a trace component that enhances attractiveness and specificity. *Entomol. Exp. Appl.*, **171**: 535-541.

Taft, W. H., 2005. *Synanthedon bolteri* (Sesiidae) in Michigan. *J. Lepid. Soc.*, **59**: 235-236.

Tagestad, A. D., 1975. Spruce budworm detection in North Dakota shelterbelts and nurseries with a synthetic sex attractant. *North Dakota Farm Res.*, **33**: 17-19.

Taguri, T., K. Yaginuma, M. Yamamoto, T. Fujii and T. Ando, 2014. Enantiospecific synthesis and field evaluation of four stereoisomers of 10,14-dimethyloctadec-1-ene, a sex pheromone component secreted by female moths of the apple leafminer. *Biosci. Biotechnol. Biochem.*, **78**: 761-765.

Takács, S., G. Gries and R. Gries, 2001. Communication ecology of webbing clothes moth: 4. Identification of male- and female-produced pheromones. *Chemoecology*, **11**: 153-159.

Takahashi, S., M. Kawaradani, Y. Sato and M. Sakai, 1979. Sex pheromone components of *Leucania separata* Walker and *Leucania loreyi* Duponchel. *Jap. J. Appl. Entomol. Zool.*, **23**: 78-81.

Takahashi, S., M. Kawaradani, Y. Sato and M. Sakai, 1980. The components of the female sex pheromone of the lorey leafworm, *Leucania loreyi* Duponchel. *Appl. Entomol. Zool.*, **15**: 499-501.

Takahashi, S., Y. Sato and T. Kodama, 1983. Attractiveness of synthetic sex pheromone to males of the lorey leafworm, *Acantholeucania loreyi* (Duponchel). *Appl. Entomol. Zool.*, **18**: 435-437.

Takanashi, T., S. Ohno, Y. Huang, S. Tatsuki, H. Honda and Y. Ishikawa 2000. A sex pheromone component novel to *Ostrinia* identified from *Ostrinia latipennis* (Lepidoptera: Crambidae). *Chemoecology*, **10**: 143-147.

Tamaki, Y., H. Noguchi, T. Yushima, C. Hirano, K. Honma and H. Sugawara, 1971(a). Sex pheromone of the summerfruit tortrix: isolation and identification. *Kontyu*, **39**: 338-340.

Tamaki, Y., H. Noguchi, T. Yushima and C. Hirano, 1971(b). Two sex pheromones of the smaller tea tortrix: isolation, identification, and synthesis. *Appl. Entomol. Zool.*, **6**: 139-141.

Tamaki, Y., H. Noguchi and T. Yushima, 1973. Sex pheromone of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae): isolation, identification, and synthesis. *Appl. Entomol. Zool.*, **8**: 200-203.

Tamaki, Y. and T. Yushima, 1974. Sex pheromone of the cotton leafworm, *Spodoptera littoralis*. *J. Insect Physiol.*, **20**: 1005-1014.

- Tamaki, Y., K. Honma and K. Kawasaki, 1977(a). Sex pheromone of the peach fruit moth, *Carposina nipponensis* Walsingham (Lepidoptera: Carposinidae): isolation, identification and synthesis. *Appl. Entomol. Zool.*, **12**: 60-68.
- Tamaki, Y., K. Kawasaki, H. Yamada, T. Koshihara, N. Osaki, T. Ando, S. Yoshida and H. Kakinohana, 1977(b). (*Z*)-11-Hexadecenal and (*Z*)-11-hexadecenyl acetates: sex-pheromone components of the diamondback moth (Lepidoptera: Plutellidae). *Appl. Entomol. Zool.*, **12**: 208-210.
- Tamaki, Y., T. Yushima, M. Oda, K. Kida, K. Kitamura, S. Yabuki and J. H. Tunlinson, 1977(c). Attractiveness of 3,13-octadecadienyl acetates for males of clearwing moths. *Jap. J. Appl. Entomol. Zool.*, **21**: 106-107.
- Tamaki, Y., H. Noguchi, H. Sugie, R. Sato and A. Kariya, 1979. Minor components of the female sex-attractant pheromone of the smaller tea tortrix moth (Lepidoptera: Tortricidae): isolation and identification. *Appl. Entomol. Zool.*, **14**: 101-113.
- Tamaki, Y., H. Noguchi, H. Sugie, A. Kariya, S. Arai, M. Ohba, T. Terada, T. Suguro and K. Mori, 1980. Four-component synthetic sex pheromone of the smaller tea tortrix moth: field evaluation of its potency as an attractant for male moth. *Jap. J. Appl. Entomol. Zool.*, **24**: 221-228.
- Tamaki, Y., H. Sugie, M. Osakabe and P. Sonnet, 1983. Biological activities of *R*- and *S*-10-methyldodecyl acetates, the chiral component of the sex pheromone of the smaller tea tortrix moth (*Adoxophyes* sp., Lepidoptera: Tortricidae). *Appl. Entomol. Zool.*, **18**: 292-294.
- Tamaki, Y., H. Sugie, Y. Ando, A. Yamashita, S. Ooya, H. Kamiwada, H. Suzuki and T. Fukumoto, 1996. Insect attractants for *Aedia leucomelas*. *Jpn. Kokai Tokkyo Koho*, JP 08,295 602 [96,295,602].
- Tamhankar**, A. J., T. P. Rajendran, M. W. Bhamburkar and M. R. Harwalkar, 1993. Optimum pheromone blend for pink bollworm males: evaluation with two experimental designs. *Int. J. Pest Manage.*, **39**(1): 111-112.
- Tan**, Z.-X., Y.-W. Wu, G.-Q. Lin, B.-C. Wu, H.-Q. Liu, X.-Y. Xu, W.-S. Zhou, G.-Q. Pu and M.-S. Zhang, 1984. Study on identification and synthesis of insect pheromone XVII. The sex pheromone of *Euproctis similis xanthocampa*. *Huaxue Xuebao (Acta Chimica Sinica)*, **42**: 1178-1182.
- Tan, Z.-X., G.-Q. Lin, H.-Q. Liu, G.-Q. Pu, H.-M. Du, J.-P. Mao, L.-Z. Meng and C.-H. Wu, 1992. Identification and synthesis of insect pheromone XXX. Sex pheromone of mulberry clearwing moth *Paradoxecia pieli* Lieu. *J. Chem. Ecol.*, **18**: 419-425.
- Tan, Z.-X., R. Gries, G. Gries, G.-Q. Lin, G.-Q. Pu, K. N. Slessor and J.-X. Li, 1996. Sex pheromone components of mulberry looper, *Hemerophila atrilineata* Butler (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **22**: 2263-2271.

- Tanaka**, A., K. Yasuda and S. Wakamura 2001. Sex pheromone of *Acrolepiopsis nagaimo* Yasuda (Lepidoptera: Acrolepiidae), a predominant leafminer on Chinese yam in Tottori Prefecture: field evaluation and activity of the third component. *Appl. Entomol. Zool.*, **36**(2): 203-207.
- Tanaka**, M., K. Abe, T. Ando and L. V. Vang, 2007. Identification of the sex pheromone secreted by females of *Grapholita inopinata* Heinrich (Lepidoptera: Tortricidae). *Res. Bull. Plant Prot. Japan*, **43**: 17-22.
- Taschenberg**, E. F., R. T. Cardé, A. Hill, J. P. Tette and W. L. Roelofs, 1974. Sex pheromone trapping of the grape berry moth. *Environ. Entomol.*, **3**: 192-194.
- Tatsuki**, S., M. Kurihara, K. Usui, Y. Ohguchi, K. Uchiumi and J. Fukami, 1983. Sex pheromone of the rice stem borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae): the third component, Z-9-hexadecenal. *Appl. Entomol. Zool.*, **18**: 443-446.
- Tatsuki, S., H. Sugie, K. Usui, J. Fukami, M. H. Sumartaputra and A. N. Kuswadi, 1985. Identification of possible sex pheromone of the yellow stem borer moth, *Scirpophaga incertulas* (Walker) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **20**: 357-359.
- Teal**, P. E. A., R. R. Heath, J. H. Tumlinson and J. R. McLaughlin, 1981. Identification of a sex pheromone of *Heliothis subflexa* (Gn.) (Lepidoptera: Noctuidae) and field trapping studies using different blends of components. *J. Chem. Ecol.*, **7**: 1011-1022.
- Teal**, P. E. A., J. R. McLaughlin, J. H. Tumlinson and R. R. Rush, 1982. A sex attractant for *Platysenta videns* (Gn.) similar to the sex pheromone of *Heliothis virescens* (F.). *Florida Entomol.*, **65**: 583-584.
- Teal**, P. E. A., R. J. West and J. E. Laing, 1983. Identification of a blend of sex pheromone components of the potato stem borer (Lepidoptera: Noctuidae) for monitoring adults. *Proc. Entomol. Soc. Ont.*, **114**: 15-19.
- Teal**, P. E. A., J. H. Tumlinson, J. R. McLaughlin, R. Heath and R. A. Rush, 1984. (Z)-11-Hexadecen-1-ol: a behavioral modifying chemical present in the pheromone gland of female *Heliothis zea* (Lepidoptera: Noctuidae). *Can. Entomol.*, **116**: 777-779.
- Teal**, P. E. A., E. R. Mitchell, J. H. Tumlinson, R. R. Heath, and H. Sugie, 1985. Identification of volatile sex pheromone components released by the southern armyworm, *Spodoptera eridania* (Cramer). *J. Chem. Ecol.*, **11**: 717-725.
- Teal**, P. E. A., J. H. Tumlinson and R. R. Heath, 1986. Chemical and behavioral analyses of volatile sex pheromone components released by calling *Heliothis virescens* (F.) females (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **12**: 107-126.
- Teal**, P. E. A., R. R. Heath, B. D. Dueben, J. A. Coffelt and K. W.

Vick, 1995. Production and release of (*Z,E*)-9,12-tetradecadienal by sex pheromone glands of females of *Plodia interpunctella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **21**: 787-799.

Teasdale, C., G. J. R. Judd, R. Gries and G. Gries, 2013. Evaluation of synthetic sex pheromone for monitoring and management of raspberry crown borer *Pennisetia marginata* (Lepidoptera: Sesiidae). *Agric. Forest Entomol.*, **15**: 285-293.

Teixeira, A. L., J. S. Oliveira, E. F. Vilela, E. R. Lima and H. O. Reis, 1989. Sex pheromone components of *Spodoptera latifascia* Walker (Lepidoptera: Noctuidae). *Revista Ceres*, **36**: 373-381.

Terytze, K., H. Adam and B. Kovaljev, 1987. Use of pheromone traps for monitoring harmful Lepidoptera species in cabbage. *Arch Phytopathol. Pflanzenschutz, Berlin*, **6**: 465-473.

Thomson, D. R., N. P. D. Angerilli, C. Vincent and A. P. Gaunce, 1991. Evidence for regional differences in the response of obliquebanded leafroller (Lepidoptera: Tortricidae) to sex pheromone blends. *Environ. Entomol.*, **20**: 935-938.

Tiberi, R. and A. Niccoli, 1984. Observations over several years on the use of traps with the sex pheromone of *Thaumetopoea pityocampa* (Den. et Schiff.) (Lepidoptera, Thaumetopoeidae). *Redia*, **67**: 129-144.

Tonini, C., G. Cassani, P. Piccardi, S. Maini, P. L. Castellari and E. Pasqualini, 1982. Sex pheromone components of the leafroller moth *Pandemis cerasana*. *J. Insect Physiol.*, **28**: 443-446.

Tonini, C., G. Cassani, P. Massardo, G. Guglielmetti and P. L. Castellari, 1986. Study of female sex pheromone of leopard moth, *Zeuzera pyrina* L. Isolation and identification of three components. *J. Chem. Ecol.*, **12**: 1545-1558.

Toshova, T., M. Subchev, E. Plass and W. Francke, 2003. *Scoliopteryx libatrix* (L.) (Lep., Noctidae) male reaction to the synthetic main sex pheromone component and its isomers - wind tunnel and field investigations. *J. Appl. Entomol.*, **127**: 195-199.

Tóth, M., G. Szöcs, L. Novák and Cs. Szántay, 1979. Post season field screening of lepidopterous sex attractant candidates in Hungary. *Acta Phytopathol. Acad. Sci. Hung.*, **14**: 195-199.

Tóth, M., J. Jakab and L. Novák, 1980. Identification of two components from the sex pheromone system of the white-line dart moth, *Scotia segetum* (Schiff.) (Lep., Noctuidae). *Z. Angew. Entomol.*, **90**: 505-510.

Tóth, M., G. Szöcs, B. Majoros, T. E. Bellas and L. Novák, 1983. Experiments with a two-component sex attractant of the silver Y moth (*Autographa gamma* L.), and some evidence for the presence of both components in natural female sex pheromone. *J. Chem. Ecol.*, **9**: 1317-1325.

- Tóth, M., T. E. Bellas and G. H. L. Rothschild, 1984. Role of pheromone components in evoking behavioral responses from male potato tuberworm moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **10**: 271-280.
- Tóth, M., P. M. Guerin, H.-R. Buser, H. Müller, G. Szöcs, G. Sziráki and H. Arn, 1985. Z-11-Tetradecenyl acetate: sex attractant of *Agapeta zoegana* (Lepidoptera: Tortricidae), a potential species for the biological control of knapweed. *Can. Entomol.*, **117**: 1163-1165.
- Tóth, M., G. Sziráki and G. Szöcs, 1986(a). Sex attractants for *Pexicopia malvella* and *Platyedra subcinerea*, pests of malvaceous ornamentals and cotton. *Entomol. Exp. Appl.*, **40**: 147-150.
- Tóth, M., G. Szöcs, C. Löfstedt, B. S. Hansson and M. Subchev, 1986(b). Sex pheromone components of *Mamestra suasa*: chemical analysis, electrophysiological activity, wind tunnel activity and field tests in two European countries. *Entomol. Exp. Appl.*, **42**: 291-299.
- Tóth, M., G. Szöcs, W. Francke, P. M. Guerin, H. Arn and A. Schmid, 1987. Field activity of sex pheromone components of *Peribatodes rhomboidaria*. *Entomol. Exp. Appl.*, **44**: 199-204.
- Tóth, M., G. Szöcs, J. Molnár and I. Szarukán, 1988. Field tests with sex attractants of *Diachrysia chrysitis* and *D. tutti* (Lepidoptera: Noctuidae) at several sites in Hungary. *Z. Naturforsch.*, **43 c**: 463-466.
- Tóth, M., C. Löfstedt, B. S. Hansson, G. Szöcs and A. I. Farag, 1989(a). Identification of four components from the female sex pheromone of the lima-bean pod borer, *Etiella zinckenella*. *Entomol. Exp. Appl.*, **51**: 107-112.
- Tóth, M., H. R. Buser, A. Peña, H. Arn, K. Mori, T. Takeuchi, L. N. Nikolaeva and B. G. Kovalev, 1989(b). Identification of (3Z,6Z)-1,3,6-9,10-epoxyheneicosatriene and (3Z,6Z)-1,3,6-9,10-epoxyeicosatriene in the sex pheromone of *Hyphantria cunea*. *Tetrahedron Lett.*, **30**: 3405-3408.
- Tóth, M., G. Helmchen, U. Leikauf, GY. Sziráki and G. Szöcs, 1989(c). Behavioral activity of optical isomers of 5,9-dimethylheptadecane, the sex pheromone of *Leucoptera scitella* L. (Lepidoptera: Lyonetidae). *J. Chem. Ecol.*, **15**: 1535-1543.
- Tóth, M., G. Szöcs, C. Löfstedt, B. S. Hansson, F. Schmidt and W. Francke, 1991(a). Epoxyheptadecadienes identified as sex pheromone components of *Tephritis arenacearia* Hbn. (Lepidoptera: Geometridae). *Z. Naturforsch.*, **46 c**: 257-263.
- Tóth, M. and G. Szöcs, 1991(b). A sex attractant for male *Swammerdamia pyrella*. *Entomol. Exp. Appl.*, **61**: 95-96.
- Tóth, M. and G. Szöcs, 1991(c). (Z)-7-Dodecenol attractive to male

Agrotis segetum Den. & Schiff. (Lep., Noctuidae) in the field.
J. Appl. Entomol., **112**: 202-206.

Tóth, M., H. R. Buser, P. M. Guerin, H. Arn, F. Schmidt, W. Francke and G. Szöcs, 1992(a). *Abraxas grossulariata* L. (Lepidoptera: Geometridae): identification of (3Z,6Z,9Z)-3,6,9-heptadecatriene and (6Z,9Z)-6,9-cis-3,4-epoxyheptadecadiene in the female sex pheromone. *J. Chem. Ecol.*, **18**: 13-25.

Tóth, M. and R. E. Doolittle, 1992(b). Sex attractants for the white-barred groundling moth (*Recurvaria leucatella* Clerck) and the lesser budmoth (*R. nanella* Hübner) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **18**: 1093-1105.

Tóth, M., C. Löfstedt, B. W. Blair, T. Cabello, A. I. Farag, B. S. Hansson, B. G. Kovalev, S. Maini, E. A. Nesterov, I. Pajor, A. P. Sazonov, I. V. Shamshev, M. Subchev and G. Szöcs, 1992(c). Attraction of male turnip moths *Agrotis segetum* (Lepidoptera: Noctuidae) to sex pheromone components and their mixtures at 11 sites in Europe, Asia, and Africa. *J. Chem. Ecol.*, **18**: 1337-1347.

Tóth, M., G. Szöcs, G. Sziraki and W. Sauter, 1992(d). Sex attractants for male Microlepidoptera found in field trapping tests in Hungary. *J. Appl. Entomol.*, **113**: 342-355.

Tóth, M., R. Gottwald, G. Szöcs and B. Freier, 1992(e). (Z)-11-Hexadecenyl acetate attracting male *Oriamusculosa* Hbn. (Lep., Noctuidae). *J. Appl. Entomol.*, **113**: 509-512.

Tóth, M., G. Szöcs and M. Bengtsson, 1992(f). A sex attractant for male *Orthosia gothica* L. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **47c**: 613-617.

Tóth, M., G. Szöcs and M. Bengtsson, 1993. Optimization of male sex attractants for *Orthosia cruda* Schiff., *O. munda* Schiff. and *O. stabilis* Schiff. (Lep., Noctuidae). *J. Appl. Entomol.*, **115**: 342-349.

Tóth, M., G. Szöcs, W. Francke, F. Schmidt, P. Philipp, C. Löfstedt, B. S. Hansson and A. I. Farag, 1994(a). Pheromonal production of and response to optically active epoxydienes in some geometrid moths (Lepidoptera: Geometridae). *Z. Naturforsch.*, **49 c**: 516-521.

Tóth, M., G. Szöcs and G. Sziráki, 1994(b). Male sex attractants for some Lepidopterous pests of the genera *Evergestis* and *Pyrausta* (Lepidoptera: Pyralidae). *Acta Phytopathol. Entomol. Hung.*, **29**: 155-159.

Tóth, M., G. Szöcs, E. J. van Nieukerken, P. Philipp, F. Schmidt and W. Francke, 1995. Novel type of sex pheromone structure identified from *Stigmella malella* (Stainton) (Lepidoptera: Nepticulidae). *J. Chem. Ecol.*, **21**: 13-27.

Tóth, M., N. S. Talekar and G. Szöcs, 1996. Optimization of blends of synthetic pheromone components for trapping male limabean

pod borers (*Etiella zinckenella* Tr.) (Lepidoptera: Phycitidae): Preliminary evidence on geographical differences. *Bioorg. Medic. Chem.*, **4**: 495-497.

Tsai, R.-S., E.-C. Yang, C.-Y. Wu, H.-K. Tseng and Y.-S. Chow **1999**. A potent sex attractant of the male tea tussock moth, *Euproctis pseudoconspersa* (Strand) (Lepidoptera: Lymantriidae) in Taiwan: Field and EAG responses. *Zool. Studies*, **38**(3): 301-306.

Tumlinson, J. H., E. R. Mitchell, S. M. Browner and D. A. Lindquist, **1972**. A sex pheromone for the soybean looper. *Environ. Entomol.*, **1**: 466-468.

Tumlinson, J. H., C. E. Yonce, R. E. Doolittle, R. R. Heath, C. R. Gentry and E. R. Mitchell, **1974**. Sex pheromones and reproductive isolation of the lesser peachtree borer and the peachtree borer. *Science*, **185**: 614-616.

Tumlinson, J. H., D. E. Hendricks, E. R. Mitchell, R. E. Doolittle and M. M. Brennan, **1975**. Isolation, identification, and synthesis of the sex pheromone of the tobacco budworm. *J. Chem. Ecol.*, **1**: 203-214.

Tumlinson, J. H., E. R. Mitchell and P. E. Sonnet, **1981**. Sex pheromone components of the beet armyworm, *Spodoptera exigua*. *J. Environ. Sci. Health*, **A 16**: 189-200.

Tumlinson, J. H., E. R. Mitchell, P. E. A. Teal, R. R. Heath and L. J. Mengelkoch, **1986**. Sex pheromone of fall armyworm, *Spodoptera frugiperda* (J. E. Smith). Identification of components critical to attraction in the field. *J. Chem. Ecol.*, **12**: 1909-1926

Tumlinson, J. H., M. M. Brennan, R. E. Doolittle, E. R. Mitchell, A. Brabham, B. E. Mazomenos, A. H. Baumhover and D. M. Jackson, **1989**. Identification of a pheromone blend attractive to *Manduca sexta* (L.) males in a wind tunnel. *Arch. Insect Biochem. Physiol.*, **10**: 255-271.

Tumlinson, J. H., E. R. Mitchell and H.-S. Yu, **1990**. Analysis and field evaluation of volatile blend emitted by calling virgin females of beet armyworm moth, *Spodoptera exigua* (Hübner). *J. Chem. Ecol.*, **16**: 3411-3423.

Tumlinson, J. H., E. R. Mitchell, R. E. Doolittle and D. M. Jackson, **1994**. Field tests of synthetic *Manduca sexta* sex pheromone. *J. Chem. Ecol.*, **20**: 579-591.

Turgeon, J. J. and G. G. Grant, **1988**. Response of *Zeiraphera canadensis* (Lepidoptera: Tortricidae: Olethreutinae) to candidate sex attractants and factors affecting trap catches. *Environ. Entomol.*, **17**: 442-447.

Twidle, A. M., D. Barker and D. M. Suckling, **2020**. (7Z)-Tricosene improves pheromone trap catch of raspberry bud moth, *Heterocrossa rubophaga*. *J. Chem. Ecol.*, **46**: 830-834.

Uehara, T., H. Naka, S. Matsuyama, T. Ando and H. Honda, **2012**.

Identification and field evaluation of sex pheromones in two hawk moths *Deilephila elpenor lewisii* and *Theretra oldenlandiae oldenlandiae* (Lepidoptera: Sphingidae). *Appl. Entomol. Zool.*, **47** (3): 227-232.

Uehara, T., H. Naka, S. Matsuyama, L. V. Vang, T. Ando and H. Honda, 2013. Identification of conjugated pentadecadienals as sex pheromone components of the sphingid moth, *Dolbina tancrei*. *J. Chem. Ecol.*, **39**: 1441-1447.

Uehara, T., H. Naka, S. Matsuyama, T. Ando and H. Honda, 2015. Identification of the sex pheromone of the diurnal hawk moth, *Hemaris affinis*. *J. Chem. Ecol.*, **41**: 9-14.

Uehara, T., H. Kitahara, H. Naka, S. Matsuyama, T. Ando and H. Honda, 2016. Single-component pheromone consisting of bombykal in a diurnal hawk moth, *Neogurelca himachala sangaica*. *J. Chem. Ecol.*, **42**: 517-522.

Ujiye, T., M. Wakou, T. Oku, K. Homma, K. Kawasaki, Y. Tamaki and H. Sugie, 1986. Sex pheromone of the apple leafminer moth (*Phyllonorycter ringoniella*): isolation and identification of the pheromone. *Jap. J. Appl. Entomol. Zool.*, **30**: 268-271.

Ujváry, I., J. C. Dickens, J. A. Kamm and L. M. McDonough, 1993. Natural product analogs: stable mimics of aldehyde pheromones. *Arch. Insect Biochem. Physiol.*, **22**: 393-411

Underhill, E. W., W. F. Steck and M. D. Chisholm, 1976. Sex pheromone of the clover cutworm moth, *Scotogramma trifolii*: isolation, identification and field studies. *Environ. Entomol.*, **5**: 307-310.

Underhill, E. W., M. D. Chisholm and W. Steck, 1977 (a). Olefinic aldehydes as constituents of sex attractants for noctuid moths. *Environ. Entomol.*, **6**: 333-337.

Underhill, E. W., W. F. Steck and M. D. Chisholm, 1977 (b). A sex pheromone mixture for the bertha armyworm moth, *Mamestra configurata*: (Z)-9-tetradecen-1-ol acetate and (Z)-11-hexadecen-1-ol acetate. *Can. Entomol.*, **109**: 1335-1340.

Underhill, E. W., W. Steck, M. D. Chisholm, H. A. Worden and J. A. G. Howe, 1978. A sex attractant for the cottonwood crown borer, *Aegeria tibialis* (Lepidoptera: Sesiidae). *Can. Entomol.*, **110**: 495-498.

Underhill, E. W., A. P. Arthur, M. D. Chisholm and W. F. Steck, 1979. Sex pheromone components of the sunflower moth, *Homoeosoma electellum*: Z-9, E-12-tetradecadienol and Z-9-tetradecenol. *Environ. Entomol.*, **8**: 740-743.

Underhill, E. W., M. D. Chisholm and W. Steck, 1980. (E)-5, (Z)-7-Dodecadienal, a sex pheromone component of the western tent caterpillar, *Malacosoma californicum* (Lepidoptera: Lasiocampidae). *Can. Entomol.*, **112**: 629-631.

Underhill, E. W., W. F. Steck, J. R. Byers and M. D. Chisholm, 1981.

Biosystematics of the genus *Euxoa* (Lepidoptera: Noctuidae). XVI. (Z)-5-Decenyl acetate a sex attractant for three closely related species, *Euxoa declarata*, *Euxoa campestris*, and *Euxoa rockburnei*. *Can. Entomol.*, **113**: 245-249.

Underhill, E. W., C. E. Rogers, M. D. Chisholm and W. F. Steck, 1982. Monitoring field populations of the sunflower moth, *Homoeosoma electellum* (Lepidoptera: Pyralidae), with its sex pheromone. *Environ. Entomol.*, **11**: 681-684.

Underhill, E. W., P. Palaniswamy, S. R. Abrams, B. K. Bailey, W. F. Steck and M. D. Chisholm, 1983. Triunsaturated hydrocarbons, sex pheromone components of *Caenurgina erechtea*. *J. Chem. Ecol.*, **9**: 1413-1423.

Underhill, E. W., A. P. Arthur and P. G. Mason, 1986. Sex pheromone of the banded sunflower moth, *Cochylis hospes* (Lepidoptera: Cochylidae): identification and field trapping. *Environ. Entomol.*, **15**: 1063-1066.

Underhill, E. W., J. G. Millar, R. A. Ring, J. W. Wong, D. Barton and M. Giblin, 1987(a). Use of a sex attractant and an inhibitor for monitoring winter moth and Bruce spanworm populations. *J. Chem. Ecol.*, **13**: 1319-1330.

Underhill, E. W., C. E. Rogers and L. R. Hogge, 1987(b). Sex attractants for two sunflower pests, *Eucosma womonana* (Lepidoptera: Tortricidae) and *Isophrictis similiella* (Lepidoptera: Gelechiidae). *Environ. Entomol.*, **16**: 463-466.

Unelius, C. R., A. Eiras, P. Witzgall, M. Bengtsson, A. Kovaleski, E. F. Vilela and A.-K. Borg-Karlson, 1996. Identification and synthesis of the sex pheromone of *Phteochoroa cranaodes* (Lepidoptera: Tortricidae). *Tetrahedron Lett.*, **37**: 1505-1508.

Usui, K., M. Kurihara, K. Uchiumi, J. Fukami and S. Tatsuki, 1987. Sex pheromone of the female large cabbage-heart caterpillar, *Crocidiolomia binotalis* Zeller (Lepidoptera: Pyralidae). *Agric. Biol. Chem.*, **51**: 2191-2195.

Vakenti, J. M., A. P. Gaunce, K. N. Slessor, G. G. S. King, S. A. Allan, H. F. Madsen and J. H. Borden, 1988. Sex pheromone components of the oblique-banded leafroller, *Choristoneura rosaceana* in the Okanagan Valley of British Columbia. *J. Chem. Ecol.*, **14**: 605-621.

Van de Veire, M. and P. Dirinck, 1986. Sex pheromone components of the cabbage armyworm, *Mamestra brassicae*: isolation, identification and field experiments. *Entomol. Exp. Appl.*, **41**: 153-155.

Vang, L. V., S. Inomata, M. Kinjo, F. Komai and T. Ando, 2005. Sex pheromones of five olethreutine species (Lepidoptera: Tortricidae) associated with the seedlings and fruits of mangrove plants in the Ryukyu Islands, Japan: identification and field evaluation. *J. Chem. Ecol.*, **31**(4): 859-878.

Vang, L. V., M. Ishitani, F. Komai, M. Yamamoto and T. Ando, 2006.

Sex pheromone of the soybean pod borer, *Leguminivora glycinvorella* (Lepidoptera: Tortricidae): Identification and field evaluation. *Appl. Entomol. Zool.*, **41** (3): 507-513.

Vang, L. V., Md. A. Islam, N. D. Do, T. V. Hai, S. Koyano, Y. Okahana, N. Ohbayashi, M. Yamamoto and T. Ando, 2008. 7,11,13-Hexadecatrienal identified from female moths of the citrus leafminer as a new sex pheromone component: synthesis and field evaluation in Vietnam and Japan. *J. Pestic. Sci.*, **33**: 152-158.

Vang, V. L., N. D. Do, L. K. An, P. K. Son and T. Ando, 2011. Sex pheromone components and control of the citrus pock caterpillar, *Prays endocarpa*, found in the Mekong Delta of Vietnam. *J. Chem. Ecol.*, **37**: 134-140.

Vang, L. V., C. N. Q. Khanh, H. Shibasaki and T. Ando, 2012. Female sex pheromone secreted by *Carmenta mimosa* (Lepidoptera: Sesiidae), a biological control agent for an invasive weed in Vietnam. *Biosci. Biotechnol. Biochem.*, **76**: 2153-2155.

Vang, L. V., H. N. Thuy, C. N. Q. Khanh, P. K. Son, Q. Yan, M. Yamamoto, U. Jinbo and T. Ando, 2013. Sex pheromones of three citrus leafrollers, *Archips atrolucens*, *Adoxophyes privatana*, and *Homona* sp., inhabiting the Mekong Delta of Vietnam. *J. Chem. Ecol.*, **39**: 783-789.

Vang, L. V., Q. Yan, N. T. N. Nghia, C. N. Q. Khanh and T. Ando, 2018. Unsaturated cuticular hydrocarbon components of the sex pheromone of eggplant fruit borer, *Leucinodes orbonalis* Guenée (Lepidoptera: Crambidae). *J. Chem. Ecol.*, **44**: 631-636.

Vang, L. V., T. T. Thy, D. K. Hanh, T. P. Linh, M. Yamamoto and T. Ando, 2020. Sex pheromone analysis and effective attraction of males of the cabbage webworm, *Hellula undalis*, inhabiting the Mekong Delta of Vietnam. *J. Asia-Pac. Entomol.*, **23**: 935-941.

Velcheva, N. and T. Milkova, 1990. Sex attractant for *Exapate congelatella* Cl (Lepidoptera, Tortricidae) - a new species for the Bulgarian fauna. *Ekologiya*, **23**: 75-78.

Vetter, R. S. and T. C. Baker, 1983. Behavioral responses of male *Heliothis virescens* in a sustained-flight tunnel to combinations of seven compounds identified from female sex pheromone glands. *J. Chem. Ecol.*, **9**: 747-759.

Vick, K. W., H. C. F. Su, L. L. Sower, P. G. Mahany and P. C. Drummond, 1974. (*Z,E*) -7,11-Hexadecadien-1-ol acetate: the sex pheromone of the Angoumois grain moth, *Sitotroga cerealella*. *Experientia*, **30**: 17-18.

Vickers, N. J., 2002. Defining a synthetic pheromone blend attractive to male *Heliothis subflexa* under wind tunnel conditions. *J. Chem. Ecol.*, **28** (6): 1255-1267.

- Voerman**, S. and W. M. Herrebout, 1978(a). A sex attractant for the leaf miner moth *Lithocolletis corylifoliella* and its influence on that of *L. blancardella* (Lep., Gracillariidae). *Entomol. Exp. Appl.*, **23**: 96-98.
- Voerman, S., A. K. Minks, G. Vanwetswinkel and J. H. Tumlinson, 1978(b). Attractivity of 3,13-octadecadien-1-ol acetates to the male clearwing moth *Synanthedon myopaeformis* (Borkhausen) (Lepidoptera, Sesiidae). *Entomol. Exp. Appl.*, **23**: 301-304.
- Voerman, S. and G. H. L. Rothschild, 1978(c). Synthesis of the two components of the sex pheromone system of the potato tuberworm moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) and field experience with them. *J. Chem. Ecol.*, **4**: 531-542.
- Voerman, S., H. Audemard and E. Priesner, 1983. Sex attractants for clearing moths: *Synanthedon vespiformis* and *Chamaesphecia tenthrediniformis* (and/or *C. empiformis*). *Entomol. Exp. Appl.*, **34**: 203-205.
- Voerman, S., C. J. Persoons and E. Priesner, 1984. Sex attractant for currant clearwing moth, *Synanthedon tipuliformis* (Clerck) (Lepidoptera: Sesiidae). *J. Chem. Ecol.*, **10**: 1371-1376.
- Voerman, S., 1988. The pheromone bank: a collection of unsaturated compounds indispensable for discovery of sex attractants for Lepidoptera. *Agric. Ecosyst. Environ.*, **21**: 31-41.
- Voerman, S., 1991. A mono-unsaturated acetate as sex attractant for the leaf miner moth *Lithocolletis corylifoliella* (Hübner) (Lep., Gracillariidae). *J. Appl. Entomol.*, **112**: 207-210.
- Vrkoc**, J., B. G. Kovalev and V. A. Starets, 1981. Isolation and identification of the main component of the female sexual pheromone of *Mamestra suasa* (Lepidoptera, Noctuidae). *Acta Entomol. Bohemoslov.*, **78**: 353-357.
- Vrkoc, J., V. P. Konyukhov and B. G. Kovalev, 1983. Isolation and identification of main components of the pheromone complex of the moth *Agrotis exclamacionis* (Lepidoptera, Noctuidae). *Acta Entomol. Bohemoslov.*, **80**: 184-189.
- Vu**, M. H., T. Ando, N. Takahashi, S. Tatsuki, A. Yamane, T. Ikeda and S. Yamazaki, 1980. Identification of the female sex pheromone of the pine moth. *Agric. Biol. Chem.*, **44**: 231-233.
- Wakamura**, S., 1978. Sex attractant pheromone of the common cutworm moth, *Agrotis fucosa* Butler (Lepidoptera: Noctuidae): isolation and identification. *Appl. Entomol. Zool.*, **13**: 290-295.
- Wakamura, S., 1980. Sex attractant pheromone of the common cutworm moth *Agrotis fucosa* Butler: field evaluation. *Appl. Entomol. Zool.*, **15**: 167-174.
- Wakamura, S., 1985. Identification of sex-pheromone components of the podborer, *Matsumuraeses falcana* (Walshingham) (Lepidoptera:

Tortricidae). *Appl. Entomol. Zool.*, **20**: 189-198.

Wakamura, S., D. L. Struble, H. Matsuura, M. Sato and K. Kegasawa, 1986(a). Sex pheromone of the black cutworm moth, *Agrotis ipsilon* Hufnagel (Lepidoptera: Noctuidae): attractant synergist and improved formulation. *Appl. Entomol. Zool.*, **21**: 299-304.

Wakamura, S. and K. Kegasawa, 1986(b). Sex pheromone of the podborer, *Matsumuraes falcana* (Walshingham) (Lepidoptera: Tortricidae): activity of the third component, (*E,Z*)-7,9-dodecadienyl acetate, and 3-component formulation. *Appl. Entomol. Zool.*, **21**: 334-339.

Wakamura, S., 1987. Sex pheromone of the beet armyworm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae): Field attraction of male moths in Japan to (*Z,E*)-9,12-tetradecadienyl acetate and (*Z*)-9-tetradecen-1-ol. *Appl. Entomol. Zool.*, **22**: 348-351.

Wakamura, S., T. Yasuda, A. Ichikawa, T. Fukumoto and F. Mochizuki, 1994. Sex attractant pheromone of the tea tussock moth, *Euproctis pseudoconspersa* (Strand) (Lepidoptera: Lymantriidae): identification and field attraction. *Appl. Entomol. Zool.*, **29**: 403-411

Wakamura, S., T. Yasuda, M. Watanabe, K. Kiguchi, M. Shimoda and T. Ando, 1996(a). Sex pheromone of the sweetpotato hornworm, *Agrius convolvuli* (L.) (Lepidoptera: Sphingidae): identification of a major component and its activity in a wind tunnel. *Appl. Entomol. Zool.*, **31**: 171-174.

Wakamura, S., A. Ichikawa, T. Yasuda, N. Arakaki and T. Fukumoto, 1996(b). EAG and field responses of the male tea tussock moth, *Euproctis pseudoconspersa* (Strand) (Lepidoptera: Lymantriidae) to (*R*)- and (*S*)-enantiomers and racemic mixture of 10,14-dimethylpentadecyl isobutyrate. *Appl. Entomol. Zool.*, **31**(4): 623-625.

Wakamura, S., N. Arakaki, T. Yasuda and K. Kawasaki, 1997. Sex pheromone of *Statherotis discana* (Felder et Rogenhofer) (Lepidoptera: Tortricidae), a pest of litchi leaves: identification and field attraction. *Appl. Entomol. Zool.*, **32**(1): 267-269.

Wakamura, S., N. Arakaki, K. Kinjo and T. Yasuda 1998. Sex pheromone of the cotton caterpillar, *Diaphania indica* (Saunders) (Lepidoptera: Pyralidae): Identification and field attraction. *Appl. Entomol. Zool.*, **33**(3): 429-434.

Wakamura, S., M. Hattori, K. Igita, K. Yasuda and Tridjaka, 1999. Sex pheromone of *Etiella behrii*, a pod borer of soybean in Indonesia: identification and field attraction. *Entomol. Exp. Appl.*, **91**: 413-420.

Wakamura, S., N. Arakaki, M. Yamamoto, S. Hiradate, H. Yasui, T. Yasuda, and T. Ando, 2001(a). Posticlure: a Novel *trans*-Epoxide as a sex Pheromone Component of the Tussock Moth, *Orgyia postica* (Walker). *Tetrahedron Lett.*, **42**(4): 687-689.

Wakamura, S., A. Tanaka, K. Yasuda and T. Yasuda, 2001(b). Sex pheromone of *Acrolepiopsis nagaimo* Yasuda (Lepidoptera: Acrolepiidae), a

predominant leafminer on Chinese yam in Tottori Prefecture: identification and field attraction. *Appl. Entomol. Zool.*, **36**(1): 71-76.

Wakamura, S., N. Arakaki, H. Ono and H. Yasui, 2001(c). Identification of novel sex pheromone components from a tussock moth, *Euproctis pulvrea*. *Entomol. Exp. Appl.*, **100**: 109-117.

Wakamura, S., N. Arakaki, H. Yamazawa, N. Nakajima, M. Yamamoto and T. Ando, 2002. Identification of epoxyhenicoxadiene and novel diepoxy derivatives as sex pheromone components of the clear-winged tussock moth *Perina nuda*. *J. Chem. Ecol.*, **28**(3): 449-467.

Wakamura, S. and N. Arakaki, 2004. Sex pheromone components of pyralid moths *Terastia subjectalis* and *Agathodes ostentalis* feeding on the coral tree, *Erithrina variegata*: two sympatric species share common components in different ratios. *Chemoecology*, **14**: 181-185.

Wakamura, S., N. Arakaki, M. Yamamoto, S. Hiradate, H. Yasui, K. Kinjo, T. Yasuda, H. Yamazawa and Tetsu Ando, 2005(a). Sex pheromone and related compounds in the Ishigaki and Okinawa strains of the tussock moth *Orgyia postica* (Walker) (Lepidoptera: Lymantriidae). *Biosci. Biotechnol. Biochem.*, **69** (5): 957-965.

Wakamura, S., N. Arakaki and K. Kinjo, 2005(b). Sex pheromone components of an olethreutid moth, *Strepsicrates semicanella* (Walker) (Lepidoptera: Tortricidae), a pest of guava and eucalyptus in Okinawa. *Appl. Entomol. Zool.*, **40**(4): 637-642.

Wakamura, S., H. Tanaka, Y. Masumoto, H. Sawada and N. Toyohara, 2007(a). Sex pheromone of the blue-striped nettle grub moth *Parasa lepida* (Cramer) (Lepidoptera: Limacodidae): identification and field attraction. *Appl. Entomol. Zool.*, **42** (3): 347-352.

Wakamura, S., T. Yasuda, Y. Hirai, H. Tanaka, T. Doki, Y. Nasu, M. Shibao, A. Yunotani and K. Kadono, 2007(b). Sex pheromone of the oriental tussock moth *Artaxa subflava* (Bremer) (Lepidoptera: Lymantriidae): identification and field attraction. *Appl. Entomol. Zool.*, **42** (3): 375-382.

Wakamura, S., S. Ohno, N. Arakaki, T. Kohama, D. Haraguchi and H. Yasui, 2010. Identification and field activity of the sex pheromone component of the sweetpotato vine borer moth *Omphisa anastomosalis* (Lepidoptera: Crambidae). *Appl. Entomol. Zool.*, **45** (4): 635-640.

Wakamura, S., H. Yasui, M. Shimatani, T. Tokashiki, T. Okuhira, N. Fujiwara-Tsujii, M. Ishikawa, T. Uesato, H. Zukeyama and H. Tobaru, 2011. Sex pheromone of the African armyworm *Spodoptera exempta* (Lepidoptera: Noctuidae): identification of components of the Okinawan population and formulation for population monitoring. *Appl. Entomol. Zool.*, **46**: 415-420.

Wakamura, S., N. Arakaki, S. Yoshimatsu, 2021. Sex pheromone of the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) of a "Far East" population from Okinawa, Japan. *Appl. Entomol. Zool.*, **56**: 19-25.

Walker, P. W., G. R. Allen, N. W. Davies, J. A. Smith, P. P. Molesworth, A. Nilsson, F. Andersson and E. Hedenström, 2009. Identification, synthesis and field testing of (3Z,6Z,9Z)-3,6,9-henicosatriene, a second bioactive component of the sex pheromone of the autumn gum moth, *Mnesampela privata*. *J. Chem. Ecol.*, **35**: 1411-1422.

Wall, C., A. G. Greenway and P. E. Burt, 1976. Electroantennographic and field responses of the pea moth, *Cydia nigricana*, to sex attractants and related compounds. *Physiol. Entomol.*, **1**: 151-157.

Wall, C., R. Bourneville, A. R. Greenway and J. Einhorn, 1987. Efficacy and stability of sex-attractant lures for *Cydia nigricana* (F.) and *Cydia medicaginis* (Kuzn.). *Crop Prot.*, **6**: 7-12.

Wallin, E. A., B. Kalinová, J. Kindl, E. Hedenström and I. Valterová, 2020. Stereochemistry of two pheromonal components of the bumblebee wax moth, *Aphomia sociella*. *Sci. Rep.*, **10**: 2094.

Wallner, W. E., R. T. Cardé, C. Xu, R. M. Weseloh, S. Xilin, J. Yan and P. W. Schaefer, 1984. Gypsy moth (*Lymantria dispar* L.) attraction to disparlure enantiomers and the olefin precursor in the People's Republic of China. *J. Chem. Ecol.*, **10**: 753-757.

Wang, X.-W. and M.-Y. Liu 1997. Extract components of single sex pheromone gland of oriental armyworm *Mythimna separata* Walker with trace analytical techniques. *Acta Entomol. Sinica. (Kunchong Xuebao)*, **40**: 158-165.

Wang, H.-L., G. P. Svensson, O. Rosenberg, M. Bengtsson, E. V. Jirle and C. Löfstedt, 2010. Identification of the sex pheromone of the spruce seed moth, *Cydia strobilella* L. *J. Chem. Ecol.*, **36**: 305-313.

Wang, H. L., G. P. Svensson, J. Jakobsson, E. V. Jirle, O. Rosenberg, W. Francke, O. Anderbrant, J. G. Millar and C. Löfstedt, 2015 (a). Sex pheromone of the cloaked pug moth, *Eupithecia abietaria* (Lepidoptera: Geometridae), a pest of spruce cones. *J. Appl. Entomol.*, **139**: 352-360.

Wang, H.-L., H. Geertsema, E. J. van Nieukerken and C. Löfstedt, 2015 (b). Identification of the female-produced sex pheromone of the leafminer *Holocacista capensis* infesting grapevine in South Africa. *J. Chem. Ecol.*, **41**: 724-731.

Wang, H.-L., M. Baldessari, G. Anfora, E. J. van Nieukerken and C. Löfstedt, 2019. Sex pheromones of two leafminer species,

Antispila oinophylla and *Holocacista rivillei* (Lepidoptera: Heliozelidae) infesting grapevine in Italy. *J. Chem. Ecol.*, **45**: 1-8.

Wang, C., S. Zhang, M.-B. Guo, X.-Q. Hou, B. Yang and G.-R. Wang, 2022. Optimization of a pheromone lure by analyzing the peripheral coding of sex pheromones of *Spodoptera frugiperda* in China. *Pest Manag. Sci.*, **78**: 2995-3004.

Wang, C., Y.-H. Li, L. Wang, B. Yang and G.-R. Wang, 2023. Development of a new sex attractant via the peripheral coding of pheromones in *Mythimna loreyi*. *J. Agric. Food Chem.*, **71**: 2795-2803.

Warthen, J. D., J. A. Klun and E. D. DeVilbiss, 1996. Structure-activity relationship observations for the bagworm moth pheromone. *J. Chem. Ecol.*, **22**: 1315-1324.

Weatherston, J., W. Roelofs, A. Comeau and C. J. Sanders, 1971. Studies of physiologically active arthropod secretions. X. Sex pheromone of the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **103**: 1741-1747.

Weatherston, J., L. M. Davidson and D. Simonini, 1974 (a). Attractants for several male forest Lepidoptera. *Can. Entomol.*, **106**: 781-782.

Weatherston, J. and W. Maclean, 1974 (b). The occurrence of (*E*)-11-tetradecen-1-ol, a known sex attractant inhibitor, in the abdominal tips of virgin female eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **106**: 281-284.

Weatherston, J., J. E. Percy and L. M. MacDonald, 1976. Field testing of *cis*-11-tetradecenal as attractant or synergist in Tortricinae. *Experientia*, **32**: 178-179.

Weatherston, J., A. F. Hedlin, D. S. Ruth, L. M. MacDonald, C. C. Leznoff and T. M. Fyles, 1977. Chemical and field studies on the sex pheromones of the cone and seed moths *Barbara colfaxiana* and *Laspeyresia youngana*. *Experientia*, **33**: 723-725.

Weatherston, J., G. G. Grant, L. M. MacDonald, D. Frech, R. A. Werner, C. C. Leznoff and T. M. Fyles, 1978. Attraction of various tortricine moths to blends containing *cis*-11-tetradecenal. *J. Chem. Ecol.*, **4**: 543-549.

Wei, H.-Y. and J.-W. Du, 2003. Identification of active components of sex pheromone for *Sidemia depravata* and field trapping. *Yingyong Shengtai Xuebao*, **14**: 730-732.

Wei, Z.-H. and F.-M. Pan, 1985. Preliminary report on the sex pheromone of the armyworm *Mythimna separata*. *Kunchong Xuebao (Acta Entomol. Sinica)*, **28**: 348-350.

Wen, X.-J., B. Kalinová, M.-G. Liu and M. Hoskovec, 2009. Candidate

sex pheromone components of persimmon bark borer, *Euzophera batangensis*. *Linye Kexue*, **45**: 83-89. [SciFinder. AN2010:149071]

Wen, X., T. Ma, Z. Liu, C. Sun and W. Cai, 2014. Sex attractant for controlling *Diaphania glauculalis*. CN 103875661. [CAN 161:180686]

Whittle, C. P., T. E. Bellas, M. Horak and B. Pineese, 1987. The sex pheromone and taxonomic status of *Homona spargotis* Meyrick sp. rev., an Australian pest species of the *Coffearia* group (Lepidoptera: Tortricidae: Tortricinae). *J. Aust. Entomol. Soc.*, **26**: 169-179.

Whittle, C. P., T. E. Bellas and A. L. Bishop, 1991. Sex pheromone of lucerne leafroller, *Merophyas divulsana* (Walker) (Lepidoptera: Tortricidae): evidence for two distinct populations. *J. Chem. Ecol.*, **17**: 1883-1895.

Whittle, C. P., R. A. Vickers, L. S. Kuniata, T. E. Bellas and E. R. Rumbo, 1995. Identification of an attractant for the caneborer *Sesamia grisescens* Walker (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **21**: 1409-1420.

Wiesner, C. J., P. J. Silk, S.-H. Tan, P. Palaniswamy and J. O. Schmidt, 1979. Components of the sex pheromone gland of the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **111**:1311.

Willemse, L. P. M., C. J. H. Booij and S. Voerman, 1987. New sex attractants for male Lepidoptera (Coleophoridae, Gelechiidae, Momphidae, Oecophoridae, and Yponomeutidae) found by field screening in the Netherlands. *J. Appl. Entomol.*, **103**: 508-515.

Williams, R. N., D. T. Johnson and E. Priesner 1999. *Synanthedon rileyana* (H. Edwards) response to selected clearwing pheromone blends. *J. Entomol. Sci.*, **34** (2): 219-224.

Witjaksono, K. Ohtani, M. Yamamoto, T. Miyamoto and T. Ando 1999. Responses of Japanese giant looper male moth to synthetic sex pheromone and related compounds. *J. Chem. Ecol.*, **25**(7): 1633-1642.

Witzgall, P. and B. Frérot, 1989(a). Pheromone emission by individual females of carnation tortrix, *Cacoecimorpha pronubana*. *J. Chem. Ecol.*, **15**: 707-717.

Witzgall, P., W. Sauter, H.-R. Buser, S. Rauscher, H. Arn, P. J. Charmillot and Th. Wildbolz, 1989(b). Use of pheromone chemistry to identify *Grapholita lobarzewskii* as an occasional pest of apple and plum. *Entomol. Exp. Appl.*, **53**: 133-136.

Witzgall, P., 1990. Attraction of *Cacoecimorpha pronubana* male moths to synthetic sex pheromone blends in the wind tunnel. *J. Chem. Ecol.*, **16**: 1507-1515.

Witzgall, P., M. Bengtsson, H.-R. Buser, P.-J. Chambon, E. Priesner, T. Wildbolz and H. Arn, 1991(a). Sex pheromone of *Spilonota*

ocellana and *Spilonota laricana*. *Entomol. Exp. Appl.*, **60**: 219-223.

Witzgall, P. P., B. Frérot and C. Malosse, 1991(b). Identification of the major component of the pheromone secretion of *Notocelia uddmanniana* L. (Lep., Tortricidae). *J. Appl. Entomol.*, **112**: 71-75.

Witzgall, P., M. Bengtsson, C. R. Unelius and J. Löfqvist, 1993. Attraction of pea moth *Cydia nigricana* F. (Lepidoptera: Tortricidae) to female sex pheromone (*E,E*)-8,10-dodecadien-1-yl acetate, is inhibited by geometric isomers *E,Z*, *Z,E*, and *Z,Z*. *J. Chem. Ecol.*, **19**: 1917-1928.

Witzgall, P., J.-P. Chambon, M. Bengtsson, C. R. Unelius, M. Appelgren, G. Makranczy, N. Muraleedharan, D. W. Reed, K. Hellrigl, H.-R. Buser, E. Hallberg, G. Bergström, M. Tóth, C. Löfstedt and J. Löfqvist, 1996. Sex pheromone and attractants in the Eucosmini and Grapholitini (Lepidoptera, Tortricidae). *Chemoecology*, **7**: 13-23.

Witzgall, P., M. Bengtsson and R. M. Trimble, 2000. Sex pheromone of grape berry moth (Lepidoptera: Tortricidae). *Environ. Entomol.*, **29**(3): 433-436.

Witzgall, P., M. Tasin, H.-R. Buser, G. Wegner-Kiß, V. S. M. Mancebón, C. Ioriatti, A.-C. Bäckman, M. Bengtsson, L. Lehmann and W. Francke, 2005. New pheromone components of the grapevine moth *Lobesia botrana*. *J. Chem. Ecol.* **31**(12): 2923-2932.

Wong, J. W., P. Palaniswamy, E. W. Underhill, W. F. Steck and M. D. Chisholm, 1984(a). Novel sex pheromone components from the fall cankerworm moth, *Alsophila pometaria*. *J. Chem. Ecol.*, **10**: 463-473.

Wong, J. W., P. Palaniswamy, E. W. Underhill, W. F. Steck and M. D. Chisholm, 1984(b). Sex pheromone components of fall cankerworm moth, *Alsophila pometaria*. Synthesis and field trapping. *J. Chem. Ecol.*, **10**: 1579-1596.

Wong, H. R., J. A. Drouin and C. L. Rentz, 1985(a). *Petrova albicapitana* (Busck) and *P. metallica* (Busck) (Lepidoptera: Tortricidae) in *Pinus contorta* Dougl. stands of Alberta. *Can. Entomol.*, **117**: 1463-1470.

Wong, J. W., E. W. Underhill, S. L. MacKenzie and M. D. Chisholm, 1985(b). Sex attractants for geometrid and noctuid moths. Field trapping and electroantennographic responses to their hydrocarbons and monoepoxydiene derivatives. *J. Chem. Ecol.*, **11**: 727-756.

Wu, D.-M. and G.-L. Li, 1982. EAG analysis and field attractant effect of sex pheromone of the striped sugarcane borer. *Kunchong Xuebao (Acta Entomol. Sinica)*, **25**: 358-362.

Wu, D.-M., Y.-H. Yan, B.-H. Wang and H.-L. Huang, 1984. The chemical structure and field attractant effect of the sex pheromone of

sugarcane stem borer *Chilo infuscatellus* (Snellen). *Kunchong Xuebao (Acta Entomol. Sinica)*, **27**: 368-374.

Wu, D.-M. and J.-R. Cui, 1986(a). Electrophysiological responses of the purple stem borer to sex pheromone and related compounds. *Kunchong Xuebao (Acta Entomol. Sinica)*, **29**: 239-245.

Wu, D.-M., Z.-R. Ding, J.-R. Cui and Y.-H. Yan, 1986(b). The study of sex attractant of the pine tip moth *Dioryctria rubella* Hampson. *Scientia Silvae Sinicae*, **22**: 368-372.

Wu, D., Y. Yan, F. Kong, W. Li, S. Yu, J. Li and W. Zeng, 1990(a). Sex pheromone of the sugarcane striped borer -- identification of three components. *Youji Huaxue, 10*: 47-49. [Chem. Abst., **112**: 174029n]

Wu, D.-M., Y.-H. Yan, J.-R. Cui, M.-Y. Liu, D.-F. Ren, A. Chen and Q. Zhang, 1990(b). Sex pheromone of sugarcane tip borer *Scirpophaga excerpalis* Walker. *Chin. Sci. Bull.*, **35**: 1469-1473.

Wu, D.-M., J.-R. Cui and Y.-H. Yan, 1993. Electrophysiological analysis and field trapping results of sex pheromone of the reed stalk borer. *Acta Phytophysiolica Sinica*, **20**(3): 229-232.

Wu, W.-Q., B. S. Hansson and C. Löfstedt, 1995. Electrophysiological and behavioural evidence for a fourth sex pheromone component in the turnip moth, *Agrotis segetum*. *Physiol. Entomol.*, **20**; 81-92.

Wu, W.-Q, C. B. Cottrell, B. S. Hansson and C. Löfstedt 1999. Comparative study of pheromone production and response in Swedish and Zimbabwean populations of turnip moth, *Agrotis segetum*. *J. Chem. Ecol.*, **25**(1): 177-196.

Wunderer, H., K. Hansen, T. W. Bell, D. Schneider and J. Meinwald, 1986. Sex pheromones of two Asian moths (*Creatonotos transiens*, *C. gangis*; Lepidoptera - Arctiidae): behavior, morphology, chemistry and electrophysiology. *Exp. Biol.*, **46**: 11-27.

Xiao, W. S. Matsuyama, T. Ando, J. G. Millar and H. Honda, 2012. Body wax unsaturated hydrocarbons as synergist of sex pheromone in the yellow peach moth, *Conogethes punctiferalis* (Crambidae: Lepidoptera). *J. Chem. Ecol.*, **38**: 1143-1150.

Yadav, J. S., M. Y. Valli and A. R. Prasad, 2001. Isolation, identification, synthesis, and bioefficacy of female *Diacrisia oblique* (Arctiidae) sex pheromone blend. An Indian agricultural pest. *Pure Appl. Chem.*, **73**: 1157-1162.

Yaginuma, K., M. Kumakura, Y. Tamaki, T. Yushima and J. H. Tunlinson, 1976. Sex attractant for the cherry tree borer, *Synanthedon hector* Butler (Lepidoptera: Sesiidae). *Appl. Entomol. Zool.*, **11**: 266-268.

Yamada, H. and T. Koshihara, 1980. Interspecific attractiveness of female sex pheromone components of the diamondback moth, *Plutella xylostella* (L.), and their analogue to the stone leek miner, *Acrolepiopsis sapporensis* Matsumura. *Kontyū*, 48: 104-110.

Yamakawa, R., N. D. Do, Y. Adachi, M. Kinjo and T. Ando, 2009. (6Z,9Z,12Z)-6,9,12-Octadecatriene and (3Z,6Z,9Z,12Z)-3,6,9,12-icosatetraene, the novel sex pheromones produced by emerald moths. *Tetrahedron Lett.*, 50: 4738-4740.

Yamakawa, R., N. D. Do, K. Kinjo, Y. Terashima and T. Ando, 2011(a). Novel components of the sex pheromones produced by emerald moths: Identification, synthesis, and field evaluation. *J. Chem. Ecol.*, 37: 105-113.

Yamakawa, R., R. Kiyota, T. Taguri and T. Ando, 2011(b). (5R,7R)-5-Methylheptadecan-7-ol: a novel sex pheromone component produced by a female lichen moth, *Miltachrista calamine*, in the family Arctiidae. *Tetrahedron Lett.*, 52: 5808-5811.

Yamakawa, R., Y. Takubo, K. Ohbayashi, H. Naka and T. Ando, 2012(a). Female sex pheromone of *Cystidia couaggaria couaggaria* (Lepidoptera: Geometridae): Identification and field attraction. *Biosci. Biotechnol. Biochem.*, 76: 1303-1307.

Yamakawa, R., Y. Takubo, H. Shibasaki, Y. Murakami, M. Yamamoto and T. Ando, 2012(b). Characterization of epoxytrienes derived from (3Z,6Z,9Z)-1,3,6,9-tetraenes, sex pheromone components of arctiid moths and related compounds. *J. Chem. Ecol.*, 38: 1042-1049.

Yamamoto, M., Y. Takeuchi, Y. Ohmasa, H. Yamazawa and T. Ando, 1999. Chiral HPLC resolution of monoepoxides derived from 6,9-dienes and its application to stereochemistry assignment of fruit-piercing noctuid pheromone. *Biomed. Chromatogr.*, 13: 410-417.

Yamamoto, M., M. Kiso, H. Yamazawa, J. Takeuchi and T. Ando, 2000. Identification of chiral sex pheromone secreted by giant geometrid moth, *Biston robustum* Butler. *J. Chem. Ecol.*, 26(11): 2579-2590.

Yamamoto, M., T. Kamata, N. D. Do, Y. Adachi, M. Kinjo and T. Ando, 2007. A novel lepidopteran sex pheromone produced by females of a Lithosiinae species, *Lyclene dharma dharma*, in the family of Arctiidae. *Biosci. Biotechnol. Biochem.*, 71 (11): 2860-2863.

Yamamoto, M., R. Yamakawa, T. Oga, Y. Takei, M. Kinjo and T. Ando, 2008. Synthesis and chemical characterization of hydrocarbons with a 6,9,11-, 3,6,9,11-, or 1,3,6,9-polyene system, pheromone candidates in Lepidoptera. *J. Chem. Ecol.*, 34 (8): 1057-1064.

Yamaoka, R., H. Fukami and S. Ishii, 1976. Isolation and identification of the female sex pheromone of the potato tuberworm moth, *Phtorimaea operculella* (Zeller). *Agric. Biol. Chem.*, 40:

1971-1977.

- Yamaoka, R., Y. Shiraishi, T. Ueno, Y. Kuwahara and H. Fukami, 1985. Structure elucidation of koiganal I and II, the sex pheromones of the webbing clothes moth, using capillary GC/MS. *Mass Spectro.*, **33**: 189-195.
- Yamaoka, R., Y. Kuwabara, M. Ogawa, S. Nakamura and R. Muneyuki, 1990. Preparation of *trans*-2-octadecenal as insect attractant, for *Tinea pellionella*. *Jpn. Kokai Tokkyo Koho* JP 02,243,605.
- Yamazawa, H., N. Nakajima, S. Wakamura, N. Arakaki, M. Yamamoto and T. Ando, 2001. Synthesis and characterization of diepoxyalkenes derived from (3Z,6Z,9Z)-trienes: lymantriid sex pheromones and their candidates. *J. Chem. Ecol.*, **27**(11): 2153-2167.
- Yan, Q., L. V. Vang, C. N. Q. Khanh, H. Naka and T. Ando, 2014. Reexamination of the female sex pheromone of the sweet potato vine borer moth: identification and field evaluation of a tricosatriene. *J. Chem. Ecol.*, **40**: 590-598.
- Yan, Q., A. Kanegae, T. Miyachi, H. Naka, H. Tatsuta and T. Ando, 2015(a). Female sex pheromones of two Japanese saturniid species, *Rhodinia fugax* and *Loepa sakaei*: identification, synthesis, and field evaluation. *J. Chem. Ecol.*, **41**: 1-8.
- Yan, Q., K. Kuriyama, K. Nishikawa, S. Tominaga, H. Tatsuta, T. Ando and H. Naka, 2015(b). (Z)-13-Hexadecenyl acetate: a novel moth sex pheromone component from *Herpetogramma submarginalis* (Lepidoptera: Crambidae). *J. Chem. Ecol.*, **41**: 441-445.
- Yan, Q., M.-Y. Zheng, J.-W. Xu, J.-F. Ma, Y. Chen, Z.-P. Dong, L. Liu, S.-L. Dong and Y.-N. Zhang, 2018(a). Female sex pheromone of *Athetis lepigone* (Lepidoptera: Noctuidae): Identification and field evaluation. *J. Appl. Entomol.*, **142**: 125-130.
- Yan, Q., H.-D. Li, Y. Chen, Z.-F. Ye, X.-Y. You, J. Zhou, L.-F. Mu, S.-J. Liu, X.-B. Kong, S. A. Khuhro and S.-L. Dong, 2018(b). Identification and field evaluation of the sex pheromone of *Orthaga achatina* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **44**: 886-893.
- Yan, Q., A. Fujino, H. Naka, S.-L. Dong and T. Ando, 2018(c). Chemical analysis of the female sex pheromone in *Palpita nigropunctalis* (Lepidoptera: Crambidae). *J. Asia-Pac. Entomol.*, **21**: 1283-1288.
- Yan, J., M. Zhang, A. Ali, X. Du, X. Mei and Y. Gao, 2022. Optimization and field evaluation of sex-pheromone of potato tuber moth, *Pthorimaea operculella* Zeller (Lepidoptera: Gelechiidae). *Pest Manag. Sci.*, **78**: 3903-3911.
- Yang, C.-Y., S. Lee, K.-S. Choi, H.-Y. Jeon and K.-S. Boo, 2007. Sex pheromone production and response in Korean populations of the diamondback moth, *Plutella xylostella*. *Entomol. Exp. Appl.*, **124**: 293-298.

- Yang, C. Y., J. R. Cho, T. J. Kang and H. Y. Jeon, 2008. Identification and field testing of sex pheromone components of a Korean population of the allium leafminer, *Acrolepiopsis sapporensis*. *Entomol. Exp. Appl.*, 129: 216-222.
- Yang, C. Y., K. S. Han and K. S. Boo, 2009(a). Sex pheromones and reproductive isolation of three species in genus *Adoxophyes*. *J. Chem. Ecol.*, 35: 342-348.
- Yang, C. Y., J. Kim, T. J. Kang and H. Y. Jeon, 2009(b). Identification and field bioassays of the sex pheromone of *Synanthedon haitangvora*. *J. Chem. Ecol.*, 35: 1197-1201.
- Yang, C. Y., S. J. Kim and S. G. Lee, 2011(a). Identification and field evaluation of the sex pheromone of *Synanthedon bicingulata* (Staudinger). *J. Chem. Ecol.*, 37: 398-402.
- Yang, C. Y., M. R. Cho and D. R. Choi, 2011(b). Identification and field evaluation of the sex pheromone components of a Korean population of *Glossosphecia romanovi*. *Entomol. Exp. Appl.*, 138: 244-248.
- Yang, C. Y., C. H. Paik, G. H. Lee and J. Y. Park, 2011(c). Female sex pheromone of the gelechiid moth *Scrobipalpa salinella* (Zeller). *J. Chem. Ecol.*, 37: 1216-1221.
- Yang, C. Y., H. S. Lee and C. G. Park, 2012. Sex pheromone of the smaller clearwing moth *Synanthedon tenuis* (Butler). *J. Chem. Ecol.*, 38: 1159-1162.
- Yang, C. Y., K. S. Choi and M. R. Cho, 2013. (*E*) -5-Hexadecenyl acetate: a novel moth sex pheromone component from *Stathmopoda auriferella*. *J. Chem. Ecol.*, 39: 555-558.
- Yang, M.-H., H.-X. Liu, J.-L. Liu, X.-Y. Jing, J.-T. Zhang, L.-H. Fan and S.-F. Wang, 2015. Sex pheromone components of the carpenterworm moth, *Holcocerus vicarious*. *Entomol. Exp. Appl.*, 154: 199-205.
- Yang, Y., L. Zhang, F. Guo, Y. Long, Y. Wan and X. Wan, 2016. Reidentification of sex pheromones of tea geometrid *Ectropis oblique* Prout (Lepidoptera: Geometriidae). *J. Econ. Entol.*, 109: 167-175.
- Yao**, E.-Y., Z.-M. Li, Z.-Q. Luo, Z.-Z. Shang, K.-S. Yin and B.-B. Hong, 1991. Report on structural elucidation of sex pheromone components of a tea pest (*Ectropis obliqua* Prout). *Prog. Nat. Sci.*, 1: 566-569.
- Yasuda**, T., S. Yoshii and S. Wakamura, 1994. Identification of the sex attractant pheromone of the browntail moth, *Euproctis similis* (Fuessly) (Lepidoptera: Lymantriidae). *Appl. Entomol. Zool.*, 29: 21-30.
- Yasuda, T., S. Wakamura and N. Arakaki, 1995. Identification of sex attractant pheromone components of the tussock moth, *Euproctis*

taiwana (Shiraki) (Lepidoptera: Lymantriidae). *J. Chem. Ecol.*, **21**: 1813-1822.

Yasui, H., S. Wakamura, N. Arakaki, H. Irei, C. Kiyuna, H. Ono, H. Yamazawa and T. Ando, 2005. Identification of a sex pheromone component of the geometrid moth *Milionia basalis pryeri*. *J. Chem. Ecol.*, **31**(3): 647-656.

Yeh, S.-F., K.-C. Lee, K.-T. Chang, F.-C. Yen and J.-S. Hwang, 1989. Sex pheromone components from Asian corn borer, *Ostrinia furnacalis* (Hubner) (Lepidoptera: Pyralidae) in Taiwan. *J. Chem. Ecol.*, **15**: 497-505.

Youm, O. and P. S. Beevor, 1995. Field evaluation of pheromone-baited traps for *Coniesta ignefusalis* (Lepidoptera: Pyralidae) in Niger. *J. Econ. Entomol.*, **88**(1): 65-69.

Young, H., R. A. Galbreath, M. H. Benn, V. A. Holt and D. L. Struble, 1985. Sex pheromone components in the New Zealand brownheaded leafroller *Ctenopseustis obliquana* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **40c**: 262-265.

Yushima, T., Y. Tamaki, S. Kamano and M. Oyama, 1974. Field evaluation of a synthetic sex pheromone, "litlure", as an attractant for males of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **9**: 147-152.

Zagatti, P., B. Lalanne-Cassou, C. Descoins and M. Gallois, 1983(a). New results on the sex pheromone of the false codling moth, *Cryptophlebia leucotreta* Meyr. (Lepidoptera, Tortricidae). *Agronomie*, **3**: 75-80.

Zagatti, P., G. A. Bosson, J. Etienne, J. Breniére, C. Descoins and M. Gallois, 1983(b). Sex pheromone of an African rice borer moth: *Chilo zacconius* Blesz. (Lepidoptera, Pyralidae). *C. R. Acad. Sci., Ser. 3*, **296**: 85-88.

Zagatti, P., A. Hamadoun, M. Lettere, D. Bordat, M. Gallois and C. Malosse, 1988. Sex pheromone of *Sesamia calamistis* (Hampson) (Lepidoptera: Noctuidae). *C. R. Acad. Sci., Ser. 3*, **307**: 837-840.

Zagatti, P., M. Renou, C. Malosse, B. Frérot, C. Pavis, M. Lettere, C. Descoins, A. Permana, Y. Pivot and F. Leclant, 1991. Sex pheromone of the European sunflower moth, *Homoeosoma nebulosum* (Den. & Schiff.) (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **17**: 1399-1414.

Zagatti, P., P. Lucas, P. Genty, S. Arango, C. Malosse and F. Tellier, 1996. Sex pheromone of *Stenoma cecropia* Meyrick (Lepidoptera: Elachistidae). *J. Chem. Ecol.*, **22**: 1103-1121

Zarbin, P. H. G., L. M. Lorini, B. G. Ambrogi, D. M. Vidal and E. R. Lima, 2007. Sex pheromone of *Lonomia obliqua*: daily rhythm of production, identification, and synthesis. *J. Chem. Ecol.*, **33**(3): 555-565.

- Zeleny, J., K. Spitzer and J. Ondracek, 1982. Tropical moths (Lepidoptera) attracted by synthetic sex pheromones in Vietnamese agro-ecosystems. *Acta Entomol. Bohemoslov.*, **80**: 413-418.
- Zhan, Y.-D., Y.-J. Liu, J.-H. Liu and Y. Liu, 2023. Pheromones emitted by both female and male moths regulate coordination between the sexes for *Agriphila aeneociliella* (Lepidoptera: Crambidae). *Insect Sci.*, **30**: XXX-XXX.
- Zhang, X.-H., G.-Z. Guo, G.-Q. Lin, Y.-W. Wu, P.-H. Wu, Z.-Y. Li and K.-N. Wei, 1986. Studies on identification and syntheses of insect pheromones. XXII Sex pheromone of poplar twig clearwing moth *Paranthrene tabaniformis* Rott -- structure and synthesis. *J. Chem. Ecol.*, **12**: 1263-1271.
- Zhang, Q.-H., E. Priesner, H.-X. Wei and Q.-C. Chen, 1993. Sex-attractant and attraction-inhibitors in Chinese larch casebearer, *Coleophora sinensis* Yang (Lep., Coleophoridae). *J. Appl. Entomol.*, **116**: 516-522.
- Zhang, Q.-H., F. Schlyter, D. Chu, X.-Y. Ma and Y. Ninomiya, 1996. Female calling behaviour and response of males to synthetic pheromone and virgin females in spring and summer generations of *Hyphantria cunea* (Lep., Arctiidae) (Drury) in Northeastern China. *J. Appl. Entomol.*, **120**: 467-476.
- Zhang, J.-T., Han Y. and Meng X.-Z., 2001. Sex pheromone of the carpenterworm, *Holcocerus insularis* (Lepidoptera, Cossidae). *Z. Naturforsch.*, **56c**: 423-429.
- Zhang, A.-B., Z.-J. Wang, S.-J. Tan and D.-M. Li, 2003 (a). Monitoring the masson pine moth, *Dendrolimus punctatus* (Walker) (Lepidoptera: Lasiocampidae) with synthetic sex pheromone-baited traps in Qianshan County, China. *Appl. Entomol. Zool.*, **38** (2): 177-186.
- Zhang, A. and S. Polavarapu, 2003 (b). Sex pheromone of the cranberry blossomworm, *Epiglaea apiata*. *J. Chem. Ecol.*, **29** (9): 2153-2164.
- Zhang, A. and S. Polavarapu, 2004. Identification of a sex pheromone component for the blueberry leafminer, *Caloptilia porphyretica*. *J. Chem. Ecol.*, **30**: 1531-1545.
- Zhang, A., T. C. Leskey, J. C. Bergh and J. F. Walgenbach, 2005. Sex pheromone of the dogwood borer, *Synanthedon scitula*. *J. Chem. Ecol.*, **31** (10): 2463-2479.
- Zhang, J.-T., X.-Y. Jing, Y.-Q. Luo, Z.-W. Li, S.-X. Zong and M.-H. Yang, 2009. The sex pheromone of the sand sagebrush carpenterworm, *Holcocerus artemisiae* (Lepidoptera, Cossidae). *Z. Naturforsch.*, **64c**: 590-596.
- Zhang, J., H. Liu, W. Zhao, J. Liu and S. Zong, 2011. Identification of the sex pheromone of *Isoceras sibirica* Alpheraky (Lepidoptera, Cossidae). *Z. Naturforsch.*, **66c**: 527-533.

- Zhao, C.-H., Q. Li, X.-Y. Guo and X.-Y. Wang, 1993. New components of sex pheromone in the pine caterpillar moth, *Dendrolimus punctatus*: identification of chemical structures and field tests. *Kunchong Xuebao (Acta Entomol. Sinica)*, 36(2): 247-250.
- Zhao, C.-H., J. G. Millar, K.-H. Pan and C.-S. Xu 1998. Responses of tea tussock moth, *Euproctis psudoconspersa*, to its pheromone, (*R*) -10,14-dimethylpentadecyl isobutyrate, and to the *S*-enantiomer of its pheromone. *J. Chem. Ecol.*, 24(8): 1347-1353.
- Zhou, Z. L., 1995. Preliminary study on the mass sex pheromone trapping against the grape clearwing moth. *Acta Phytophylacica Sinica*, 22: 17-21.
- Zhu, P.-C., F.-L. Kong, S.-D. Yu, Y.-Q. Yu, S.-P. Jin, X.-H. Hu and J.-W. Xu, 1987(a). Identification of the sex pheromone of eggplant borer *Leucinodes orbonalis* Guenée (Lepidoptera: Pyralidae). *Z. Naturforsch.*, 42 c: 1347-1348.
- Zhu, P.-C., F.-L. Kong and Z.-H. Wang, 1987(b). Sex pheromone components of purple stem borer *Sesamia inferens* (Walker). *J. Chem. Ecol.*, 13: 983-989.
- Zhu, P.-C., F.-L. Kong and Y.-Q. Yu, 1987(c). Sex pheromone of Oriental armyworm *Mythimna separata* Walker. *J. Chem. Ecol.*, 13: 977-981.
- Zhu, J.-W., M. V. Kozlov, P. Philipp, W. Francke and C. Löfstedt, 1995(a). Identification of a novel moth sex pheromone in *Eriocrania cicatricella* (Zett.) (Lepidoptera: Eriocraniidae) and its phylogenetic implications. *J. Chem. Ecol.*, 21: 29-43.
- Zhu, J.-W., C. Löfstedt, P. Philipp, W. Francke, T. Tammaru and E. Haukioja, 1995(b). A sex pheromone component to the Geometridae identified from *Epirrita autumnata*. *Entomol. Exp. Appl.*, 75; 159-164.
- Zhu, J., N. Ryrholm, H. Ljungberg, B. S. Hansson, D. Hall, D. Reed and C. Löfstedt, 1996. Olefinic acetates, •-9,11-14:OAc and •-7,9-12:OAc used as sex pheromone components in three geometrid moths, *Idaea aversata*, *I. straminata*, and *I. biselata* (Geometridae, Lepidoptera). *J. Chem. Ecol.*, 22: 1505-1526.
- Zhu, J., C. Ryne, C. R. Unelius, P. G. Valeur and C. Löfstedt, 1999. Reidentification of the female sex pheromone of the Indian meal moth, *Plodia interpunctella*: evidence for a four-component pheromone blend. *Entomol. Exp. Appl.*, 92: 137-146.
- Zhu, J. and K. F. Haynes, 2004. Sex pheromone components of the bronzed cutworm, *Nehelodes minians*, a prey species of a bolas spider, *Mastophora hutchinsoni*. *J. Chem. Ecol.* 30: 2047-2056.
- Zhu, J.-W., S. Polavarapu, K-C. Park, C. Garvey, D. Mahr, S. Nojima, W. Roelofs and T. Baker, 2009. Reidentification of pheromone composition of *Sparganothis sulfureana* (Clemens) and evidence of geographic variation in male responses from two US states.

J. Asia Pacific Entomol., **12**: 247-252.

Zilahi-Balogh, G. M. G., N. P. D. Angerilli, J. H. Borden, M. Meray, M. Tulung and D. Sembel, 1995. Regional differences in pheromone responses of diamondback moth in Indonesia. *Int. J. Pest Manage.*, **41**(4): 201-204.