

Pituitary adenoma

Lester D.R. Thompson, MD, FASCP




Figure 1. A: An ectopic pituitary adenoma within the sphenoid sinus has an intact respiratory epithelium (top) and a sheetlike growth of cells (below). B: A trabecular arrangement is noted in many tumors.

Pituitary tumors account for approximately 15% of all intracranial neoplasms. They can be either benign or malignant, primary or secondary, and epithelial or nonepithelial. Pituitary carcinoma can only be diagnosed when there is confirmed metastatic disease. The vast majority are primary epithelial pituitary adenomas, which generally arise from hormone-producing cells, regardless of whether a particular tumor is active or inactive (i.e., non-hormone-producing). Pituitary adenomas are further classified on the basis of size, function, cell type, hormone production, location, and biologic behavior. Most pituitary adenomas are benign neoplasms, either null-cell or prolactin-producing micro-

adenomas (<1 cm in their greatest dimension—usually seen radiographically) that are seen within the pituitary gland. Most pituitary adenomas occur in women; they can arise at any age, but they are more common in the third through sixth decades of life. Patients with a pituitary adenoma present with either an endocrinopathy or a mass effect. Nonfunctioning adenomas exhibit no biochemical or clinical evidence of excessive hormone production. Patients with functional tumors present with specific syndromes, such as Cushing's syndrome (adrenocorticotrophic hormone-producing).

Histologically, pituitary adenomas are classified as




Figure 2. A: Small- to medium-size cells with abundant pale and eosinophilic cytoplasm surround nuclei with coarse nuclear chromatin. B: The neoplastic cells are strongly immunoreactive with chromogranin.

basophilic, eosinophilic, and null cell; each has a host of different subtypes whose classification is based on specific ultrastructural and immunophenotypic features. Cells are usually arranged in sheets or in gland-like or follicular structures (figure 1). The size of the cells varies from small to medium. Some cells have eosinophilic to basophilic granular cytoplasm, and others have clear cytoplasm (figure 2). Immunophenotypically, these cells react with keratin, chromogranin, and synaptophysin; they also demonstrate specific peptides in many cases.

The transsphenoid approach to surgical excision is the most effective treatment, with a cure rate that ranges from 75 to 93%. Irradiation has also been used; among adults, its cure rate is 15% and its clinical improvement rate is 30%. Specific symptoms caused by functional tumors can be managed with pharmacotherapy, but with mixed results.


Suggested reading

Asa SL. Tumors of the pituitary gland. In: Atlas of Tumor Pathology. 3rd series, fascicle 22. Washington, D.C.: Armed Forces Institute of Pathology, 1998.

DeLellis RA, Lloyd RV, Heitz PU, Eng C, eds. Pathology and Genetics of Tumours of Endocrine Organs. World Health Organization Classification of Tumours. Lyon: IARC Press, 2004:10-47.

OTOVENT[®]

Treatment for Negative Pressure & Otitis



The Otovent Auto-Inflation Kit is designed to treat many of the complications associated with negative ear pressure caused by Eustachian tube dysfunction. The Otovent method can prevent or treat Otitis Media and can be an attractive alternative to surgery.

Clinical Study:
S-E Stangerup M.D., J. Sederberr-Olsen M.D.,
V. Balle M.D. Auto-Inflation as a treatment of
Secretory Otitis Media, Arch Otolaryngology
Head Surg. 1992; 118: 149-154



Benefits include:

- High compliance
- Visually verifiable compliance
- Frequent, regular treatments
- Effective without drugs
- May avoid / delay surgery
- Painless
- Low treatment costs
- No known side effects
- Early, satisfying results

INVOTEC INTERNATIONAL[™]

6833 Phillips Industrial Boulevard • Jacksonville, Florida 32256 • U.S.A.
Tel: 800-998-8580 / 904-880-1229 • Fax: 904-886-9517 • www.invotec.net

Circle 109 on Reader Service Card

Serenocem[™]

Otologic Cement

Serenocem[™] was designed specifically for non-weight bearing applications in otologic surgery including:

- Reconstruction of the ossicular chain where cement can be used to repair the bony ossicles in their normal position
- Acoustic meatal wall reconstruction
- Cementation of cochlear implant electrodes



Use Serenocem[™] to bridge the eroded incus with the stapes

In its mastoidal course, the electrode is fixated with Serenocem[™]



- Highly biocompatible
- Moldable on placement
- Easy to trim when set
- High compressive strength
- Non exothermic snap set
- Chemical adhesion to bone and metals
- Bioactive/osteoconductive ion release
- Minimal biodegeneration or absorption
- Minimal shrinkage upon setting

for a FREE Surgical CD contact:

INVOTEC INTERNATIONAL

6833 Phillips Industrial Blvd • Jacksonville, Florida 32256 • USA
Tel: 800-998-8580 / 904-880-1229 • Fax: 904-886-9517 • www.invotec.net

Circle 110 on Reader Service Card