

Bungartz, F., Ziemmeck, F., Yáñez Ayabaca, A., Nugra, F., Aptroot, A. (2013). CDF Checklist of Galapagos Lichenized Fungi - FCD Lista de especies de Hongos liquenizados Galápagos. In: Bungartz, F., Herrera, H., Jaramillo, P., Tirado, N., Jiménez-Uzcátegui, G., Ruiz, D., Guézou, A. & Ziemmeck, F. (eds.). Charles Darwin Foundation Galapagos Species Checklist - Lista de Especies de Galápagos de la Fundación Charles Darwin. Charles Darwin Foundation / Fundación Charles Darwin, Puerto Ayora, Galapagos:
<http://checklists.datazone.darwinfoundation.org/true-fungi/lichens/> Last updated: 19 Sep 2013

CDF Checklist of Galapagos Lichenized Fungi

Frank Bungartz, Frauke Ziemmeck, Alba Yáñez Ayabaca, Fredy Nugra, André Aptroot

Last updated: 19 Sep 2013

Abstract

This Checklist of Galapagos Lichenized Fungi includes a total of all 794 taxa reported from the Galapagos Islands.

For each name, detailed information is provided: its Galapagos distribution in islands groups or bioregions generated from the specimen records, comments about the taxonomy (especially synonyms), the origin (native and introduced), taxon status (accepted vs. rejected records) and relevant literature references.

Introduction

This publication lists all species of Galapagos Lichenized Fungi currently known.

This checklist includes all lichen fungi that are currently known to occur in the Galapagos Islands. This list also includes some fungi typically associated with lichens (lichenicolous fungi, i.e., fungi parasitic or parasymbiotic on lichens) or often treated by lichenologists even though they are not lichenized.

Lichens do not represent a taxonomic group, but are a classic example of a symbiosis, i.e., two very different organisms living together to form a new entity.

The outside shape and structure that we notice as a lichen is built by a fungus. Inside, this macroscopic structure hosts microscopic cells of green algae (Chlorophyta) or blue-green photosynthetic bacteria (Cyanobacteria).

The fungi that participate in this symbiosis belong to two taxonomic divisions (or phyla): Basidiomycota & Ascomycota.

Methods

This checklist of all known Galapagos Lichenized Fungi is automatically generated using the online database of the Charles Darwin Foundation Galapagos Species Checklist.

All CDF Galapagos Species Checklists represent the synthesis of many different records: literature citations, data from previously unpublished reports (grey literature), specimen records of natural history collections located in Galapagos and all over the world. To the best of their knowledge authors of the individual checklists revised all available data. When new information becomes available, the taxonomy of a group changes or new species are discovered, the CDF online database and thus this publication becomes updated.

For many poorly known species groups the higher taxonomic classification still regularly changes according to how our knowledge about species being related changes. In many well known groups the phylogeny is somewhat stable, but to avoid confusion, in particular for groups where taxonomic changes are frequent, all checklists presented here are sorted alphabetical according to genus name and specific epithet. Please refer to the website for the currently accepted taxonomic hierarchy of each group.

Please be aware that the distribution presented here is automatically generated from specimen records and does not always accurately reflect the known distribution for all species.

For marine species, the distribution generally refers to the five main bioregions of the archipelago (Far Northern, Northern, Western, South Eastern and the Elisabeth Bay Bioregion). For the terrestrial species more than 120 islands, islets and small rocks have been aggregated into Islands Groups, thus, for example, the island group "Santa Cruz" includes smaller islands like Santa Fé, Plaza Norte, Plaza Sur, Baltra, Daphne Mayor, Daphne Minor, and others.

IUCN red-list assessments presented here may deviate from the global IUCN list for the following reasons:

- for well known species groups like vascular plants or vertebrates updates proposed to the IUCN are shown instead of the outdated, but currently accepted status;
- for poorly known species groups (e.g., lichenized fungi) a general assessment is currently not possible and the list presented here is a regional red-list list for Galapagos archipelago.

Numbers of the species included in this list are auto-generated. Adding up the number of species in each category will not always equal the total number indicated. Some species have insufficient data to be categorized while others (e.g., category eradicated) will not be included in the total.

Results

Names of taxa included in this checklist: 794 (577 accepted, 5 unidentified taxon, 13 doubtful, 39 preliminary identification, 12 problematic, 8 new to science), 140 rejected.

Origin of the taxa included: 1 questionable native, 41 endemic, 556 indigenous, 3 questionable endemic.

1. *Acantholichen pannariooides* P.M. Jørg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Jørgensen, P.M. et al. (1998), Lawrey, J.D. et al. (2009), Lücking, R. et al. (2009), Yáñez-Ayabaca, A. et al. (2012).

2. *Acarospora americana* H. Magn.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

References: Knudsen, K. et al. (2012).

3. *Acarospora chrysops* (Tuck.) H. Magn.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide K. Knudsen (pers. comm.) this taxon was misidentified in Elix & McCarthy (1998) as *A. citrina* and in Weber (1986) as *A. schleicheri*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Weber, W.A. et al. (1986).

4. *Acarospora sparsiuscula* H. Magn.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

References: Knudsen, K. et al. (2012).

5. *Acarospora trachyticola* (Müll. Arg.) Hue

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

References: Knudsen, K. et al. (2012).

6. *Actinoplaca strigulacea* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

7. *Actinoplaca vulgaris* (Müll. Arg.) V■zda & Poelt

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Tricharia vulgaris, fide Elix & McCarthy (1998); specimen in COLO 193173 (L-40433, det. V■zda):

Santa Cruz, on leaves of Coffea, Horneman garden, p.p. (with Aspidothelium scutellicarpum Lücking;
originally identified by V■zda as Aspidothelium fugiens)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

8. *Aderkomyces papilliferus* (Lücking) Lücking, Sérus. & V■zda

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

9. *Agonimia opuntiella* (Buschardt & Poelt) V■zda

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela.

10. *Agonimia pacifica* (H. Harada) Diederich

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón, Santa Cruz.

11. *Agonimia tristicula* (Nyl.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

12. *Alyxoria ochrocheila* (Nyl.) Ertz & Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Opegrapha ochrocheila Nyl.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Unknown.

References: Ertz, D. et al. (2010).

13. *Alyxoria varia* (Pers.) Ertz & Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Opegrapha varia Pers.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Unknown.

References: Ertz, D. et al. (2010).

14. *Angiactis spinicola* Aptroot & Sparrius

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR. Galapagos: Santa Cruz Island, coastal cliffs E of Puerto Ayora near Charles Darwin Research Station, on spines of *Scutia spicata*, 0°44'45.0"S, 90°17'39.0"W, 20 m alt., 29 May 2005, Aptroot 63413 (CDS 30168, holotype; ABL, BM, isotypes).

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, San Cristóbal, Santa Cruz.

References: Aptroot, A. et al. (2007).

15. *Anisomeridium albisedum* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón, Santa Cruz.

16. *Anisomeridium biforme* (Borrer) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

17. *Anisomeridium leptospermum* (Zahlbr.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

18. *Anisomeridium polypori* (Ellis & Everh.) M.E. Barr

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz, Santiago.

19. *Anisomeridium subprostans* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

20. *Anisomeridium tamarindi* (Fée) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

21. *Anisomeridium tuckerae* R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Original spelling as *A. tuckeri* is incorrect; species named after S. C. Tucker, correct spelling therefore: *A. tuckerae*; Syn.: *Ditremis tuckerae*, fide Elix & McCarthy (1998).

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998).

22. *Arthonia antillarum* (Fée) Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

23. *Arthonia cinnabarina* (DC.) Wallr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Arthonia tumidula*, fide Elix & McCarthy (1998), *Arthonia gregaria*, fide Weber (1986): 488

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

24. *Arthonia follmanniana* Diederich

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

References: Diederich, P. et al. (1995), Elix, J.A. et al. (1998).

25. *Arthonia platygraphidea* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Fernandina.

26. *Arthonia sanguinea* (Willey) Zahlbr.

Taxon status: Identification not yet confirmed.

F. Bungartz: material needs verification

Origin: Native, Indigenous.

Galapagos Distribution: Isabela.

27. *Arthonia speciosa* (Müll. Arg.) Grube

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Santa Cruz.

References: Grube, M. et al. (2007).

28. *Arthopyrenia cerasi* (Schrad.) A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

29. *Arthopyrenia cinchonae* (Ach.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Specimen in COLO: Santa Cruz, on Scalesia, Bella Vista, Weber L-40225, det. Aptroot, 1991

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

30. *Arthothelium galapagoense* Huneck & Follmann

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

31. *Aspidothelium cinerascens* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

32. *Aspidothelium scutellicarpum* Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: in Weber (1986) erroneously cited as Aspidophyllum fugiens; material Weber 285 (L-40433).

The material was originally determined by Vezda, but with publication of Lücking (2008) the species concept has changed; the Galapagos specimens has perithecia with disk-like, dentate expansion and not setae or hairs and thus belongs to A. scutellicarpum.

Origin: No Data.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Lücking, R. et al. (2008), Weber, W.A. et al. (1986).

33. *Asterothyrium rotuliforme* (Müll. Arg.) Sérus. & J.R. De Sloover

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

34. *Astrothelium variolosum* (Ach.) Müll. Arg.

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz & R. Miranda: According to Harris (1995) Trypethelium ochroleucum and Astrothelium variosolum are identical in all characters but their stromata formation; Trypethelium nitidiusculum is identical to Trypethelium ochroleucum in all characters but the presence of lichexanthone. It is possible that all three taxa refer to one variable species only.

Origin: No Data.

Galapagos Distribution: Isabela, Pinta.

References: Harris, R.C. et al. (1995), Kirk, P. (ed.) et al. (2010).

35. *Aulaxina opegraphina* Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

36. ***Bacidia heterochroa*** (Müll. Arg.) Zahlbr.

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela, Santa Cruz.

37. ***Bacidia insularis*** Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

38. ***Bacidia millegrana*** (Taylor) Zahlbr.

Taxon status: Identification not yet confirmed.

Lich. Exs. COLO 121; A. Aptroot doubts this determination (pers. comm.)

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1981).

39. ***Bacidia polychroa*** (Th. Fr.) Körb.

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela, Santa Cruz.

40. ***Bacidia rubella*** (Hoffm.) A. Massal.

Taxon status: Identification not yet confirmed.

specimen in COLO (L-41125); A. Aptroot doubts this determination (pers. comm.)

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

41. ***Bacidina apiahica*** (Müll. Arg.) V■zda

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Bacidia apiahica*, fide A. Aptroot (pers. comm.), *Woessia apiahica*, fide Index Fungorum

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

42. ***Bacidina chlorotica*** (Nyl.) V■zda & Poelt

Taxon status: Identification not yet confirmed.

Syn.: *Bacidia chloroticula*

Origin: No Data.

Galapagos Distribution: Santa Cruz.

43. *Bacidina delicata* (Larbal. ex Leight.) V. Wirth & Vázquez

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela, Santa Cruz.

44. *Bacidina pallidocarnea* (Müll. Arg.) Vázquez

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

45. *Bactrospora acicularis* (C.W. Dodge) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, San Cristóbal, Santa Cruz, Santa Fé.

References: Aptroot, A. et al. (2008).

46. *Bactrospora denticulata* (Vain.) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008).

47. *Bactrospora myriadea* (Fée) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008).

48. *Bathelium degenerans* (Vain.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

49. *Bathelium feei* (C.F.W. Meissn.) Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Bathelium carolinianum*.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz.

50. *Biatoropsis usnearum* Räsänen

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santiago.

51. *Brigantiae a leucoxantha (sorediate form)* (Spreng.) R. Sant. & Hafellner

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: the Galapagos material is densely diffusely sorediate across the entire thallus and therefore not *B. leucoxantha* s.str.; the specimen collected by Pike (COLO L-55433) examined by J. Hafellner 1983 identified as *B. leucoxantha* is also densely sorediate.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

52. *Brigantiae a leucoxantha s.str.* (Spreng.) R. Sant. & Hafellner

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: most Galapagos material is densely sorediate and therefore does not belong to *B. leucoxantha* s.str. However, some material not forming soredia has been found, those esorediate specimens may not be identical with the sorediate forms because they are known from much drier habitats only.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz.

53. *Buellia coccinea s.l.* (Fée) Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

54. *Buellia erubescens* Arnold

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Unknown.

55. *Buellia galapagona* W.A. Weber

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Near Threatened.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Roth, K. et al. (1978), Weber, W.A. et al. (1971), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

56. ***Buellia halonia*** (Ach.) Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz, Santiago.

57. ***Buellia lauricassiae*** (Fée) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: No Data.

Galapagos Distribution: Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

58. ***Buellia mamillana s.l.*** (Tuck.) W.A. Weber

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Rinodina mamillana*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Darwin, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

59. ***Buellia oidalea*** (Nyl.) Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, San Cristóbal.

60. ***Buellia rufofuscescens*** Stizenb.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

61. ***Buellia spuria*** (Schaer.) Anzi

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Española, Isabela, Santa Cruz, Santa Fé.

62. ***Buellia stellulata*** (Taylor) Mudd

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Española, Floreana, Isabela, Santa Cruz, Santiago.

63. *Buellia straminea* Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Buellia xanthinula* auct. non (Müll. Arg.) Malme, fide Elix & McCarthy (1998); in Dodge (1936) as *Buellia xanthinula*

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

64. *Buellia subdisciformis* agg. (Leight.) Vain.

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela.

65. *Buellia sulphurica* Bungartz & Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR. Galápagos: Isabela Island, Volcán Alcedo, upper NNW-exposed slope inside the crater, 0°27'S, 91°7'W, 1055 m, open vegetation with *Adianthus concinnum* and scattered shrubs of *Tournefortia rufosericea* among basalt blocks in the vicinity of the sulfur vents, on basalt, March 2006, Aptroot 64881(holotype CDS-31458, isotype ABL).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

References: Lumbsch, H.T. et al. (2010).

66. *Buellia trachyspora* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

67. *Bulbothrix bulbillosa* Benatti, Spielmann & Bungartz ined.

Taxon status: Unpublished name (Nomen nudum).

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, San Cristóbal.

68. *Bulbothrix laevigatula* (Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia laevigatula*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

69. *Bulbothrix lyngei* Benatti & Marcelli

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

70. *Bulbothrix scortella* (Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: only one specimen (Aptroot 65313) has a brown lower side throughout.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

71. *Bulbothrix subdissecta* (Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

72. *Byssoloma discordans* (Vain.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

73. *Byssoloma leucoblepharum* (Nyl.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

74. *Byssoloma subdiscordans* (Nyl.) P. James

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Byssoloma rotuliforme*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

75. *Byssoloma tricholomum* (Mont.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

searched for specimen in COLO, but not found: COLO (L-63764), coll.: Lanier, in packet with *Tapellaria epiphylla*, det.: Vezda

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

76. *Calicium robustellum* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

77. *Calopadia bonitensis* Cáceres & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

78. *Calopadia cinereopruinosa* Bungartz & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Type:—ECUADOR. Galápagos: Isla Sán Cristóbal, area W of Cerro Pelado on the way to El Ríososo, 0°52'S, 89°28'W, 400 m, transition zone, open *Psidium guajava* shrubland with *Macraea loricifolia* and dominant annual herb *Malachra capitata*, on bark and wood, dead twigs of *Psidium guajava*, sunny, wind- and rain-exposed, August 2008, Bungartz 8489 (holotype CDS-41135).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz.

References: Lumbsch, H.T. et al. (2010).

79. *Calopadia editae* Vázquez ex Chaves & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Calopadia pruinosa* Lücking & Chavez nom. nud.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz, Santiago.

References: Lumbsch, H.T. et al. (2010).

80. *Calopadia foliicola* (Fée) Vázquez

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Lopadium foliicola*, fide Weber (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998).

81. *Calopadia fusca* (Müll. Arg.) Vitzada

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lopadium fuscum, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998).

82. *Calopadia perpallida* (Nyl.) Vitzada

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

83. *Calopadia puiggarii* (Müll. Arg.) Vitzada

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lopodium puiggarii, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: San Cristóbal, Santa Cruz.

References: Elix, J.A. et al. (1998).

84. *Calopadia subcoeruleascens* (Zahlbr.) Vitzada

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, Santa Cruz, Santiago.

85. *Caloplaca aphanotripta* (Nyl.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

86. *Caloplaca arenaria agg.* (Pers.) Müll. Arg.

Taxon status: Identification not yet confirmed.

F. Bungartz & U. Søchting: specimens belong to at least one undescribed, possibly several species

Origin: No Data.

Galapagos Distribution: Pinta, San Cristóbal, Santa Cruz, Santiago.

87. *Caloplaca brouardii* (de Lesd.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide A. Aptroot: Caloplaca brouardii

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

88. *Caloplaca camptidia* (Tuck.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

89. *Caloplaca cf. sonorae* Wetmore

Taxon status: Identification not yet confirmed.

F. Bungartz & U. Søchting: not the correct name; together with specimens preliminary identified as *C. arenaria* agg. this species belong to at least one undescribed, possibly several species

Origin: No Data.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz, Santiago.

90. *Caloplaca chlorina* (Flot.) Sandst.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón, San Cristóbal, Santa Cruz.

91. *Caloplaca chrysodeta* (Vain. ex Räsänen) Dombr.

Taxon status: Accepted name; taxon occurs in Galapagos.

U. Søchting: No molecular data are available, but the morphology and secondary chemistry conform with *C. chrysodeta*, which is recorded as the only leprose Caloplaca from the Sonoran Desert.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Unknown.

92. *Caloplaca cupulifera* (Vain.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

93. *Caloplaca diplacia* (Ach.) Riddle

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

94. *Caloplaca epiphora* (Taylor) C.W. Dodge

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

95. *Caloplaca flavovirescens s.l.* (Wulfen) Dalla Torre & Sarnth.

Taxon status: Identification not yet confirmed.

F. Bungartz & U. Søchting: the Galapagos specimens belong to an undescribed species in the Caloplaca flavovirescens-Group

Origin: No Data.

Galapagos Distribution: Santa Cruz, Santiago.

96. *Caloplaca floridana* (Tuck.) S.C. Tucker

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Placodium floridanum*; specimen in COLO (56167), coll.: Pike

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

97. *Caloplaca furfuracea* H. Magn.

Taxon status: Identification not yet confirmed.

F. Bungartz: Superficially similar to *C. wrightii*, but with different chemistry.

Origin: No Data.

Galapagos Distribution: Isabela.

98. *Caloplaca isidiosa* (Vain.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Darwin, Española, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

99. *Caloplaca phyllidizans* Wetmore

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santiago.

100. *Caloplaca sideritis* (Tuck.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz.

101. *Caloplaca squamosa* (B. de Lesd.) Zahlbr. s.l.

Taxon status: Identification not yet confirmed.

F. Bungartz & U. Søchting: taxonomically problematic group of species including, commonly called C. subsoluta or C. modesta, the Galapagos specimens probably several new species

Origin: No Data.

Galapagos Distribution: Isabela, San Cristóbal, Santiago.

102. *Caloplaca subsoluta* (Nyl.) Zahlbr.

Taxon status: Identification not yet confirmed.

Syn.: Caloplaca modesta, fide F. Bungartz annotations, 2008; F. Bungartz & U. Søchting: taxonomically problematic group of species including, commonly called C. subsoluta or C. modesta, the Galapagos specimens probably several new species

Origin: No Data.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

103. *Caloplaca wrightii* (Willey) Fink

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

104. *Candelaria pacifica* M. Westb. & Arup

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: first published as nomen nudum by M. Westberg in Nash et al. (2002), then validated in M. Westb. & Arup, Bibl. Lich. 106: 358 (2011).

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela.

References: Nash III, T.H. et al. (2002), Westberg, M. et al. (2011).

105. *Candelariella corallizoides* Westberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

106. *Canoparmelia caroliniana* (Nyl.) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Parmelia caroliniana, fide Elix (1989), Pseudoparmelia caroliniana

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998).

107. *Canoparmelia raunkiaeri* (Vain.) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

108. *Canoparmelia texana* (Tuck.) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

109. *Celothelium dominicanum* (Vain.) M.B. Aguirre

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz & R. Miranda: previously misidentified as *C. aciculiferum*.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

110. *Chaenotheca chloroxantha* Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal.

111. *Chaenothecopsis savonica s.l.* (Räsänen) Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

112. *Chiodecton malmei* G. Thor

Taxon status: Identification not yet confirmed.

F. Bungartz: very similar, but spores of different size

Origin: No Data.

Galapagos Distribution: Santiago.

113. *Chrysotrichia aff. occidentalis* Elix & Kantivalis

Taxon status: Taxonomic status unresolved or unrevised.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, San Cristóbal, Santiago.

114. *Chrysotrichia xanthina* (Vain.) Kalb

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Chrysotrichia candelaris*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Kalb, K. et al. (2001).

115. *Cladia aggregata* (Sw.) Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela, Santa Cruz.

References: Ahti, T. et al. (2000), Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986), Yáñez-Ayabaca, A. et al. (2013).

116. *Cladonia aff. ramulosa* (With.) J.R. Laundon

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Cladonia adspersa*, fide Ahti (2000), *Cladonia pityrea*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Yáñez-Ayabaca, A. et al. (2013).

117. *Cladonia aff. sphacelata* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Not In Galapagos, Not In Galapagos.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986), Yáñez-Ayabaca, A. et al. (2013).

118. *Cladonia arbuscula* ssp. *boliviiana* Ahti

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Cladina arbuscula* ssp. *boliviiana*; specimens in H and LSU: Isabela, Volcán Alcedo, 1970, Prichard s.n.

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Isabela, Santa Cruz.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

119. *Cladonia arcuata* Ahti

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Cladina arcuata*; in Weber (1986) as *Cladina sandstedei*

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

120. *Cladonia bungartzii* Stenroos & Ahti

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

Galapagos Distribution: Pinta.

References: Yáñez-Ayabaca, A. et al. (2013).

121. *Cladonia cartilaginea* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide T. Ahti annotations 2010 (new to Galapagos): CDS 33156

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010), Yáñez-Ayabaca, A. et al. (2013).

122. *Cladonia ceratophylla* (Sw.) Spreng.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Ahti, T. et al. (2000), Dodge, C.W. et al. (1935), Elix, J.A. et al. (1998), Stewart, A. et al. (1912), Svenson, H.K. et al. (1935), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986), Yáñez-Ayabaca, A. et al. (2013).

123. *Cladonia chlorophaea* (Flörke ex Sommerf.) Spreng.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Cladonia balfourii, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Weber, W.A. et al. (1981), Yáñez-Ayabaca, A. et al. (2013).

124. *Cladonia confusa f. bicolor* (Müll. Arg.) Ahti

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Cladonia polia, fide A. Aptroot (pers. comm.), Cladina confusa f. bicolor

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

125. *Cladonia confusa f. confusa* (R. Sant.) Follmann & Ahti

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Cladonia galapagensis Ahti, fide Ahti (2000), Cladina confusa f. confusa

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

126. *Cladonia corniculata* Ahti & Kashiwadani

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Ahti, T. et al. (2000), Elix, J.A. et al. (1998).

127. *Cladonia corymbites* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

128. *Cladonia corymbosula* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide annotations T. Ahti, 2010 (new to Galapagos, new records from the Venezuelan Andes): CDS 31848

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Yáñez-Ayabaca, A. et al. (2013).

129. *Cladonia dactylota* Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Cladonia subcariosa, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Yáñez-Ayabaca, A. et al. (2013).

130. *Cladonia didyma* (Fée) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Ahti, T. et al. (2000), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986), Yáñez-Ayabaca, A. et al. (2013).

131. *Cladonia grayi* G. Merr. ex Sandst.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Yáñez-Ayabaca, A. et al. (2013).

132. *Cladonia macilenta var. bacillaris* (Genth) Schaer.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide T. Ahti annotations 2010: CDS 27818; first reported by Weber (1986) as *C. macilenta* ssp. *theiophila*

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

133. *Cladonia nana* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

134. *Cladonia polyscypha* Ahti & L. Xavier

Taxon status: Accepted name; taxon occurs in Galapagos.

fide annotations T. Ahti, 2010 (new to Galapagos, range poorly known): CDS 31086

Origin: Not In Galapagos, Not In Galapagos.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, Santa Cruz, Santiago.

References: Yáñez-Ayabaca, A. et al. (2013).

135. *Cladonia pulverulenta* (L. Scriba) Ahti

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Yáñez-Ayabaca, A. et al. (2013).

136. *Cladonia pyxidata* (L.) Hoffm.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide annotations T. Ahti, 2010 (new to Galapagos, recently found in continental Ecuador): CDS 31422

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

References: Yáñez-Ayabaca, A. et al. (2013).

137. *Cladonia scholanderi* Abbayes

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Cladonia sphacelata*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, Santa Cruz.

References: Ahti, T. et al. (2000), Yáñez-Ayabaca, A. et al. (2013).

138. *Cladonia strepsilis* (Ach.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Yáñez-Ayabaca, A. et al. (2013).

139. *Cladonia subradiata* (Vain.) Sandst.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Cladonia subulata*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Ahti, T. et al. (2000), Elix, J.A. et al. (1998), Yáñez-Ayabaca, A. et al. (2013).

140. *Cladonia subsquamosa* Kremp.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz.

References: Weber, W.A. et al. (1986), Yáñez-Ayabaca, A. et al. (2013).

141. *Cladosporium arthoniae* M.S. Christ. & D. Hawksw.

Taxon status: Identification not yet confirmed.

Index Fungorum: anamorphic Davidiella

Origin: No Data.

Galapagos Distribution: Santiago.

References: Kirk, P. (ed.) et al. (2010).

142. *Coccocarpia delicatula* Bungartz, Ziemmeck & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR. Galápagos: Isla Sán Cristóbal, area W of Cerro Pelado on the way to El Ripioso, 0°52'S, 89°28'W, 400 m, transition zone, open *Psidium guajava* shrubland with *Macraea laricifolia* and dominant annual herb *Malachra capitata*, on bark and wood, dead twigs of *Psidium guajava*, sunny, wind- and rain-exposed, August 2008, Bungartz 8496 (holotype CDS-41142).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Lumbsch, H.T. et al. (2010).

143. *Coccocarpia domingensis* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Pinta, San Cristóbal, Santa Cruz.

144. *Coccocarpia erythroxyli* (Spreng.) Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Pannaria molybdaea*, fide Weber (1986) & Elix & McCarthy (1998), *Coccocarpia pellita* var. *parmeliooides*, fide Elix & McCarthy (1998); 2 specimens in COLO: 40290, Santa Cruz, Stewart, 1912, det. Dodge 1935 & Santiago, Pike 2718, det. Arvidsson

Origin: Native, Indigenous.

IUCN Red List: Regionally Extinct.

Galapagos Distribution: Floreana, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1935), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

145. *Coccocarpia filiformis* Arv.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz.

146. *Coccocarpia palmicola* (Spreng.) Arv. & D.J. Galloway

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Floreana, Isabela, Marchena, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

147. *Coccocarpia pellita* (Ach.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Svenson, H.K. et al. (1935).

148. *Coccocarpia prostrata* Lücking, Aptroot & Sipman

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

149. *Coenogonium aff. luteolum* (Kalb) Kalb & Lücking

Taxon status: Unpublished name (Nomen nudum).

F. Bungartz: most specimens of *C. strigosum* possibly belong here

Origin: No Data.

Galapagos Distribution: Santa Cruz.

150. *Coenogonium flavum* (Malcolm & Vizzda) Malcolm

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

151. *Coenogonium interplexum* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

152. *Coenogonium pineti* (Schrad. ex Ach.) Lücking & Lumbsch

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

153. *Coenogonium strigosum* Rivaz Plata, Lücking & Chaves

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) probably as *Dimerella lutea*, fide A. Aptroot (pers. comm.); most Galapagos specimens most likely belong to a yet unpublished species *C. aff. luteolum*, sensu (Lücking, pers. com.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz, Santiago.

154. *Collema furfuraceum* Du Rietz

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2008), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

155. *Collema leptaleum* Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2008).

156. *Collema pulcellum* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

157. *Collema texanum* Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2008).

158. *Collemopsidium sublitorale* (Leight.) Grube & B.D. Ryan

Taxon status: Accepted name; taxon occurs in Galapagos.

Syns.: Pyrenocollema sublitorale, Arthopyrenia sublitoralis.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

References: Santesson, R. et al. (1939).

159. *Compsocladium kalbii* Frisch

Taxon status: Identification not yet confirmed.

F. Bungartz: material identified originally by Aptroot as Compsocladium archboldianum, but this species does not occur in South America and if correct the identification likely refers to Compsocladium kalbii Frisch, unfortunately all material missing from CDS!

Origin: Native, Indigenous.

Galapagos Distribution: Santa Cruz.

References: Frisch, A. et al. (2007).

160. *Cora glabrata* (Spreng.) Fr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Dictyonema glabratum fide Bungartz 2010; In Dodge (1935) und Weber (1966) as Cora pavonia

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1935), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986),

Yáñez-Ayabaca, A. et al. (2012).

161. *Cresponea flava* (Vain.) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (2008).

162. *Cryptothecia darwiniana* Bungartz & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

Galapagos Distribution: Floreana, Santa Cruz, Santiago.

References: Bungartz, F. et al. () .

163. *Cryptothecia galapagoana* Bungartz & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

Galapagos Distribution: Pinzón, Santa Cruz.

References: Bungartz, F. et al. () .

164. *Cryptothecia striata* G. Thor

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz.

References: Bungartz, F. et al. () .

165. *Cyphellostereum imperfectum* Lücking, Barillas & Dal Forno

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santiago.

References: Yáñez-Ayabaca, A. et al. (2012).

166. *Cyphellostereum sp. 1*

Taxon status: Unpublished name (Nomen nudum).

F. Bungartz: Dictyonema phyllogenum f. defectum was described for a morphotype with a very poorly developed fungal sheath (Lücking 2008). Fig 258 E-F in Lücking (2008), however, depicts cyanobacterial trichomes that are not closely bundled as it is the case in the Galapagos material. Also, trichomes of Dictyonema phyllogenum f. defectum are appressed rather than erect, as seen in the Galapagos material. The Galapagos specimens grow on a variety of substrates, one also found on leaves of the endemic Miconia robertsoniana, but most were collected on bryophytes. The micromorphology of these specimens is quite similar, but not identical with free-living forms of Scytonema, since Cyphellostereum sp. 1 clearly presents fungal sheath cells surrounding the cyanobacteria.

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Yáñez-Ayabaca, A. et al. (2012).

167. *Dibaeis sorediata* Kalb & Gierl

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: material confirmed per TLC by K. Kalb, December 2011

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santiago.

168. *Dictyographa arabica* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Opegrapha arabica (Müll. Arg.) Vain.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Unknown.

References: Ertz, D. et al. (2010).

169. *Dictyonema galapagoense* Yáñez, Dal-Forno & Bungartz

Taxon status: Accepted name; taxon occurs in Galapagos.

Holotype Bungartz 8517 (CDS 41163)

Origin: Native, Endemic.

IUCN Red List: Endangered.

Galapagos Distribution: Floreana, San Cristóbal.

References: Yáñez-Ayabaca, A. et al. (2012).

170. *Dictyonema pectinatum* Dal Forno, Yáñez & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Yáñez-Ayabaca, A. et al. (2012).

171. *Dictyonema schenkianum* (Müll. Arg.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: the correct spelling is 'schenkianum' because the basionym is *Laudatea schenkiana* Müll. Arg., Parmastro erroneously spelled the epithet with 'ck' as '*Dictyonema sericeum* f. *schenckianum*'.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Kirk, P. (ed.) et al. (2010), Yáñez-Ayabaca, A. et al. (2012).

172. *Dictyonema sericeum* (Sw.) Berk.

Taxon status: Accepted name; taxon occurs in Galapagos.

reported by Weber (1986) as *Dictyonema guadelupense*; only the material forming bracket-like shelves belongs here

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Lawrey, J.D. et al. (2009), Weber, W.A. et al. (1986).

173. *Dimidiographa loandensis* (Nyl.) Ertz, Bungartz & Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz, Santa Fé, Santiago.

References: Ertz, D. et al. (2009), Ertz, D. et al. (2010).

174. *Diorygma poitaei* (Fée) Kalb, Staiger & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Graphina virginea, fide F. Bungartz 2008

Origin: Native, Indigenous.

IUCN Red List: Not Evaluated.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

175. *Diploschistes actinostomus* (Pers.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

176. *Diploschistes badius* Lumbsch & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: previously misidentified as *D. aeneus*, but material contains gyrophoric acid

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

177. *Diploschistes cinereocaesius* (Sw.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Diploschistes scruposus*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

178. *Diploschistes euganeus* (A. Massal.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

179. *Diploschistes muscorum* ssp. *bartlettii* Lumbsch

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Lumbsch, H.T. et al. (1987).

180. *Diploschistes rampoddensis* (Nyl.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, Santa Cruz.

181. *Dirina approximata* Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Dirina paradoxa* ssp. *approximata* (Fée) Tehler & *Dirina herrei* Zahlbr. fide Tehler et al. (2013); Type: Herre A.W.C.T. s.n. [W - lectotype selected by Tehler et al. (2013), B, BM, G, GBG, H, KASSEL, L, LD, M, NY, S-L6, UC, UPS - isolectotypes], material not seen by F. Bungartz (no specimens in COLO?); Tehler et al. (2013) also include the sorediate forms rarely found in Galapagos here.

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Aptroot, A. et al. (2008), Elix, J.A. et al. (1998), Tehler, A. et al. (2013), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

182. *Dirina pacifica* Tehler & Ertz

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: previously reported as *D. catalinariae* f. *catalinariae* and *D. catalinariae* f. *sorediata*; according to Aptroot & Sparrius (2008) reports of *Roccellina badia* by Weber (1986) and Elix & McCarthy (1998) belong to *D. catalinariae* and thus must be considered records of *D. pacifica*.

Origin: Native, Indigenous.

Galapagos Distribution: Unknown.

References: Aptroot, A. et al. (2008), Elix, J.A. et al. (1998), Tehler, A. et al. (2013), Weber, W.A. et al. (1986).

183. *Dirinaria aegialita* (Afzel. ex Ach.) B.J. Moore

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Physcia aegialita*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1936), Weber, W.A. et al. (1966).

184. *Dirinaria appianata* (Fée) D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

According to A. Aptroot (pers. comm.) in Weber (1986) probably also as *Dirinaria leopoldii*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Weber, W.A. et al. (1986).

185. *Dirinaria confluens* (Fr.) D.D. Awasthi

Taxon status: Taxonomic status unresolved or unrevised.

Syn.: *Physcia confluens*, fide Elix & McCarthy (1998); the only CDS specimen (Ap 63111) has a UV-medulla and otherwise looks like *D. confusa*, this record is thus probably a misidentification, but check COLO loan, and do chemistry!!!, fide F. Bungartz

Origin: No Data.

Galapagos Distribution: Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

186. *Dirinaria confusa* D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

A. Aptroot (pers. comm.) suspects that Weber's specimens are misidentification of *Dirinaria aegialita*. However, TLC by K.Kalb & F. Bungartz confirms that the specimens have indeed been correctly identified as *D. confusa*.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinzón, Santa Cruz, Santiago.

References: Weber, W.A. et al. (1986).

187. *Dirinaria consimilis* (Stirt.) D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

188. *Dirinaria leopoldii* (Stein) D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

One specimen of Weber cited in Awasthi (1975): South Plaza , Weber L-40110; Two specimens in COLO (L-40736, L-40762), coll. Itow

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Santa Cruz, Santiago.

References: Awasthi, D.D. et al. (1975), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

189. *Dirinaria neotropica* Kalb

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Genovesa, Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

190. *Dirinaria papillulifera* (Nyl.) D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santiago.

191. *Dirinaria picta* (Sw.) Schaer. ex Clem.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Physcia picta*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

192. *Dyplobelia afzelii* (Ach.) A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Graphis afzelii*

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Floreana, Santiago.

References: Bungartz, F. et al. (2009), Weber, W.A. et al. (1986).

193. *Echinoplaca leucotrichoides* (Vain.) R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

194. *Endocarpon nigromarginatum* H. Harada

Taxon status: Accepted name; taxon occurs in Galapagos.

material confirmed by O. Breuss

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón.

195. *Endocarpon pallidulum* (Nyl.) Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

O. Breuss: previously rejected, but one record now confirmed (Bungartz, F. 6527, CDS 34745); F. Bungartz: in previous versions of this checklist erroneously spelled as *Endocarpon pallidellum*.

Origin: Not In Galapagos, Not In Galapagos.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

196. *Endocarpon petrolepideum* (Nyl.) Hue

Taxon status: Accepted name; taxon occurs in Galapagos.

material confirmed by O. Breuss

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

197. *Endocarpon pusillum* Hedw.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: previously already confirmed by Weber (1986), material confirmed by Breuss, but some specimens are *Endocarpon pallidellum* Ach.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

198. *Enterographa anguinella* (Nyl.) Redinger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2008).

199. *Enterographa leucolyta* (Nyl.) Redinger

Taxon status: Identification not yet confirmed.

F. Bungartz: identification cannot be verified, specimen missing (Aptroot 63270, CDS 30006; unicate loan not returned)

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (2008).

200. *Enterographa pallidella* (Nyl.) Redinger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz, Santiago.

References: Aptroot, A. et al. (2008).

201. *Enterographa subserialis* (Nyl.) Redinger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2008).

202. *Erioderma mollissimum* (Samp.) Du Rietz

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

203. *Erioderma sorediatum* D.J. Galloway & P.M. Jørg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

204. *Everniastrum vexans* (Zahlbr. ex W.L. Culb. & C.F. Culb.) Hale ex Sipman

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Parmelia vexans

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

205. *Fellhanera encephalarti* (Vezda) Vezda

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

206. *Fellhanera parvula* (Vezda) Vezda

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

207. *Fellhanera rubida* (Müll. Arg.) Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz.

208. *Fellhanera stanhopiae* (Müll. Arg.) Lücking, Lumbsch & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

209. *Fellhanera sublecanorina* (Nyl.) Vezda

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Bacidia sublecanorina, fide A. Aptroot (pers. comm.); specimen in COLO (63764, in packet of Tapellaria epiphylla), det. Vezda

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

210. *Fissurina cf. comparilis* (Nyl.) Nyl.

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

F. Bungartz: preliminary identification based on a single specimen (Aptroot 63929) only.

Origin: No Data.

Galapagos Distribution: Floreana, Santa Cruz.

References: Bungartz, F. et al. (2009).

211. *Fissurina columbina* (Tuck.) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Bungartz, F. et al. (2009).

212. *Fissurina dumastioides* (Fink) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Bungartz, F. et al. (2009).

213. *Flakea papillata* O.E. Erikss.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Agonimia papillata, fide Index Fungorum and A. Aptroot; Syn.: Psoroglaena cubensis auct. non Müll.

Arg., fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Muggia, L. et al. (2009), Weber, W.A. et al. (1993).

214. *Flavoparmelia leucoxantha* (Müll. Arg.) Hale ex DePriest & B.W. Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pseudoparmelia leucoxantha.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, Santa Cruz.

215. *Glyphis cicatricosa* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

216. *Glyphis scyphulifera* (Ach.) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, San Cristóbal, Santiago.

References: Bungartz, F. et al. (2009).

217. *Gomphillus ophiosporus* Kalb & Vázquez

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

218. *Graphis acharii* Fé

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

219. *Graphis adpressa* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Bungartz, F. et al. (2009).

220. *Graphis anfractuosa* (Eschw.) Eschw.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Bungartz, F. et al. (2009).

221. *Graphis caesiella* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Bungartz, F. et al. (2009), Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

222. *Graphis cincta* (Pers.) Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

223. *Graphis cleistomma* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela, Santiago.

References: Bungartz, F. et al. (2009).

224. *Graphis conferta* Zenker

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

225. *Graphis crebra* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

226. *Graphis dichotoma* (Müll. Arg.) Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

References: Bungartz, F. et al. (2009).

227. *Graphis disserpens* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

228. *Graphis elongata* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

229. *Graphis flavominiata* Moncada & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

230. *Graphis glaucescens* Féé

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Graphis bulacana Vain.

Origin: Introduced, Questionable Native.

IUCN Red List: Not Evaluated.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Bungartz, F. et al. (2009).

231. *Graphis handelii* Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

232. *Graphis immersella* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

233. *Graphis intricata* Féé

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

234. *Graphis leptospora* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

235. *Graphis longula* Kremp.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

236. *Graphis modesta* Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

237. *Graphis oxyclada* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

References: Kirk, P. (ed.) et al. (2010).

238. *Graphis paradiisperpens* Sipman & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

239. *Graphis pedunculata* Bungartz & Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Holotype Bungartz 5701 (CDS 28799)

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

References: Bungartz, F. et al. (2009).

240. *Graphis phaeospora* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz.

References: Bungartz, F. et al. (2009).

241. *Graphis platycarpa* Eschw.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

242. *Graphis rimulosa* (Mont.) Trevis.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

243. *Graphis subchrysocarpa* Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1993) as Phaeographina chrysocarpa

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Pinta.

References: Bungartz, F. et al. (2009), Weber, W.A. et al. (1993).

244. *Graphis tenella* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

245. *Graphis vestitoides* (Fink) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

246. *Gyalectidium catenulatum* (Cavalc. & A.A. Silva) L.I. Ferraro, Lücking & Sérus.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

247. *Gyalectidium eskuchei* Sérus. & J.R. De Sloover

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

248. *Gyalectidium filicinum* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

249. *Gyalectidium imperfectum* Vitzada

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

250. *Gyalideopsis subaequatoriana* Lücking & W.R. Buck

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

251. *Gyalideopsis vulgaris* (Müll. Arg.) Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

252. *Haematomma persoonii* (Fée) A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1966, 1986) and Elix & McCarthy (1998) as *Haematomma puniceum*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

253. *Helminthocarpon leprevostii* Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. () .

254. *Heppia despreauxii* (Mont.) Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

255. *Herpothallon aff. confluenticum* Aptroot & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Bungartz, F. et al. () .

256. *Herpothallon echinatum* Aptroot, Lücking & Will-Wolf

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Santa Cruz.

References: Bungartz, F. et al. () .

257. *Herpothallon granulare* (Sipman) Aptroot & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. () .

258. *Herpothallon hyposticticum* Bungartz & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. () .

259. *Herpothallon rubrocinctum* (Ehrenb.: Fr.) Aptroot, Lücking & G. Thor

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Herpothallon sanguineum, Chiodescon sanguineum and C. rubrocinctum, fide Aptroot et al. (2009)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2009), Dodge, C.W. et al. (1935), Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

260. *Herpothallon rubroechinatum* Frisch & G. Thor

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Bungartz, F. et al. () .

261. *Herpothallon saxorum* Bungartz & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Questionable Endemic.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. () .

262. *Heterocyphelium leucampyx* (Tuck.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela, Pinta.

263. *Heterodermia albicans* (Pers.) Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Physcia albicans*, fide F. Bungartz, *Physcia crispa*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Wolf.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

264. *Heterodermia antillarum* (Vain.) Kurok.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

265. *Heterodermia appalachensis* Kurok.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz & K. Kalb: this species is reported endemic to the North American Appalachians, but very similar to *H. lutescens*, the only difference being the soralia (capitate to labriform in *H. appalachensis*; subcapitate in *H. lutescens*), both forms occur in Galapagos and quite possibly belong to one and the same species

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

266. *Heterodermia circinalis* (Zahlbr.) W.A. Weber

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Anaptychia leucomelaena* f. *circinalis*, *Anaptychia circinalis*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, Santiago.

References: Weber, W.A. et al. (1981).

267. *Heterodermia comosa* (Eschw.) Follmann & Redón

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Anaptychia comosa, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Floreana, Isabela, Pinta, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

268. *Heterodermia coralliphora* (Taylor) Skorepa

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Anaptychia coralliphora [corallophora] Taylor; F. Bungartz & K. Kalb: the spelling "corallophora" is an orthographical error that needs to be corrected to "corallifera".

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

269. *Heterodermia diademata* (Taylor) D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: wrongly keyed in Martins (2007); the species is fertile and has no vegetative propagules.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela.

References: Martins, M.F.N. et al. (2007).

270. *Heterodermia galactophylla* (Tuck.) W.L. Culb.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz.

271. *Heterodermia japonica* (M. Satô) Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz, Santiago.

272. *Heterodermia leucomela* ssp. *boryi* (Fée) Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz.

273. *Heterodermia leucomela* ssp. *leucomela* (Fée) Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

274. *Heterodermia lutescens* (Kurok.) Follmann

Taxon status: Taxonomic status unresolved or unrevised.

Syn.: Anaptychia lutescens; F. Bungartz: How does this differ from *H. appalachensis*?

Origin: No Data.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

275. *Heterodermia obscurata* (Nyl.) Trevis.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Anaptychia obscurata, fide Index Fungorum

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

276. *Heterodermia podocarpa* (Bél.) D.D. Awasthi

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Anaptychia podocarpa, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1936), Weber, W.A. et al. (1966).

277. *Heterodermia speciosa* (Wulfen) Trevis.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

278. *Heterodermia squamulosa* (Degel.) W.L. Culb.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) and Elix & McCarthy (1998) as *Heterodermia lepidota*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

279. *Heterodermia verrucifera* (Kurok.) W.A. Weber

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Anaptychia leucomelaena f. verrucifera

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Weber, W.A. et al. (1981).

280. *Hyperphyscia adglutinata* (Flörke) H. Mayrhofer & Poelt

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Physcia adglutinata*, fide Elix & McCarthy (1998); In Weber (1986) as *Phaeophyscia hispidula*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Dodge, C.W. et al. (1936), Weber, W.A. et al. (1966).

281. *Hyperphyscia pandani* (H. Magn.) Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, San Cristóbal.

282. *Hypotrachyna costaricensis* (Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

283. *Hypotrachyna exsplendens* (Hale) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Santa Cruz.

284. *Hypotrachyna isidiocera* (Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia isidiocera*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

285. *Hypotrachyna microblasta* (Vain.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia microblasta*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

286. *Hypotrachyna osseoalba* (Vain.) Y.S. Park & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

287. *Hypotrachyna sanjosensis* Elix, T.H. Nash & Sipman

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, Santa Cruz.

288. *Julella asema* R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, Santa Cruz.

289. *Julella avicenniae* (Borse) K.D. Hyde

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: non-lichenized phytopathogen

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

290. *Julella cf. vitrispora* (Cooke & Harkn.) M.E. Barr

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

F. Bungartz & R. Miranda: the material is similar to *J. vitrispora* but does not agree in all characters; possibly an undescribed species.

Origin: Native, Indigenous.

Galapagos Distribution: Española, Floreana, Pinzón, Santa Cruz.

291. *Julella geminella* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

292. *Lecanactis abietina* (Ehrh. ex Ach.) Körb.

Taxon status: Accepted name; taxon occurs in Galapagos.

fide annotations J.M. Egea & P. Torrente, 1990 and F. Bungartz, 2009 (FH-TUCK 197166)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

293. *Lecanactis epileuca* (Nyl.) Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

294. *Lecanographa brattiae* (Egea & Ertz) Ertz & Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Opegrapha brattiae Ertz & Tehler

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Unknown.

References: Ertz, D. et al. (2010).

295. *Lecanographa hypothallina* (Zahlbr.) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Pinzón, Santa Cruz.

References: Aptroot, A. et al. (2008).

296. *Lecanographa laingiana* Diederich, Egea & Sipman

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, Pinta, Santa Cruz, Santa Fé.

References: Aptroot, A. et al. (2008).

297. *Lecanographa lyncea* (Sm.) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008).

298. *Lecanographa microcarpella* (Müll. Arg.) Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

In Aptroot & Sparrius (2008) as Lecanographa illecebrosula, fide F. Bungartz annotations, 2008 and fide D. Ertz annotations, 2011.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2008).

299. *Lecanographa subcaesioides* Egea & Torrente

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, Santa Cruz, Santiago, Wolf.

References: Aptroot, A. et al. (2008).

300. *Lecanora avium* (Zahlbr.) Hertel

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lecidea chilena, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

301. *Lecanora caesiorubella* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lecanora caesiorubella ssp. glaucomodes, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

302. *Lecanora expallens* Ach.

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela, Santiago.

303. *Lecanora floridula* Lumbsch

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Guderley, R. et al. (1999).

304. *Lecanora galactiniza* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

305. *Lecanora legalloana* Elix & Øvstedal

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

306. *Lecanora leprosa* Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Guderley, R. et al. (1999), Weber, W.A. et al. (1986).

307. *Lecanora oreinoides* (Körb.) Hertel & Rambold

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lecidea oreinoides

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

308. *Lecanora pseudopinguis* W.A. Weber

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

309. *Lecanora schindleri* Guderley

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Guderley, R. et al. (1999).

310. *Lecanora strobilina* (Spreng.) Kieff.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santiago.

311. *Lecanora subcrenulata* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

312. *Lecanora subimmergens* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

313. *Lecanora subimmersa* ssp. *ramboldii* Lumbsch & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: Both chemotypes occur in Galapagos (previously only *L. subimmersa* ssp. *subimmersa* reported by Guderley 1999)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal.

References: Guderley, R. et al. (1999).

314. *Lecanora subimmersa* ssp. *subimmersa* (Fée) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, Pinzón, San Cristóbal, Santiago.

References: Guderley, R. et al. (1999).

315. *Lecanora substrobilina* Printzen

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

316. *Lecanora sulfurescens* Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Darwin, Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Guderley, R. et al. (1999), Weber, W.A. et al. (1986).

317. *Lecanora tropica* Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Guderley, R. et al. (1999).

318. *Lecidopyrenopsis corticola* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta.

319. *Leiorreuma sericeum* (Eschw.) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Fernandina, Isabela, Pinta, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

320. *Lempholemma dussii* (Vain.) Zahlbr.

Taxon status: Identification not yet confirmed.

In Weber (1993) and Elix & McCarthy (1998) erroneously as Lempholemma dussiana; Specimens in COLO: Weber & Lanier (L-62925), Weber (L-63035); material being revised by M. Schultz

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

321. *Lepraria lobificans* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

322. *Lepraria tenellum* (Tuck.) Lendemer & Hodkinson

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Stereocaulon tenellum Tuck., Leprocaulon tenellum (Tuck.) Nyl., Stereocaulon albicans, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Unknown.

References: Elix, J.A. et al. (1998), Lendemer, J.C. et al. (2013), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

323. *Lepraria vouauxii* (Hue) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

324. *Leprocollema novocaldonianum* A.L. Sm. [as 'nova-caledonianum']

Taxon status: Identification not yet confirmed.

F. Bungartz: previously listed as "Lemmopsis novae-caledoniarum fide Aptroot", but this combination is problematic, because this has never been published; M. Schultz: the specimen is currently being revised and it is probably not identical with Leprocollema novacaledoniarum, but possibly a new species.

Origin: No Data.

349. *Minutoexcipula sp. I*

Taxon status: Taxon not identified to species, subspecies, form or variety.

Index Fungorum: anamorphic Pezizomycotina

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

350. *Mycocalicium americanum* (R. Sant.) Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santa Fé, Santiago.

351. *Mycomicrothelia subfallens* (Müll. Arg.) D. Hawksw.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

352. *Mycomicrothelia thelena* (Ach.) D. Hawksw.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, San Cristóbal.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

353. *Mycoporum compositum* (A. Massal.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: Aptroot 63342, material needs verification

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinta, Santa Cruz, Santiago.

354. *Mycoporum eschweileri* (Müll. Arg.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Tomasellia eschweileri (Müll. Arg.) R.C. Harris

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

355. *Mycoporum sparsellum* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

356. *Myriotrema olivaceum* Féé

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungarz: Santiago, on Zanthoxylum, coll. Pike ID18-19, det. Hale (OSC 101980); specimen in COLO not seen (COLO 55414)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

357. *Nelsenium usnicum* (Sipman) Lendemer & Hodkinson

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lepraria usnica Sipman

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Unknown.

References: Lendemer, J.C. et al. (2013).

358. *Normandina pulchella* (Borrer) Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

359. *Ochrolechia africana* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1966, 1986) and Elix & McCarthy (1998) as *Ochrolechia pallescens*

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

360. *Ochrolechia pallescens s.l.* (L.) A. Massal.

Taxon status: Identification not yet confirmed.

Syn.: *Lecanora pallescens*, fide Elix & McCarthy (1998); chemical reactions agree with *O. pallescens*, but the Galapagos material grows in arid vegetation

Origin: No Data.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

361. *Ochrolechia subpallescens* Verseghy

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.
IUCN Red List: Data Deficient.
Galapagos Distribution: Unknown.

362. *Opegrapha agelaeotera* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Indigenous.
IUCN Red List: Least Concern.
Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

363. *Opegrapha astraea* Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Indigenous.
IUCN Red List: Data Deficient.
Galapagos Distribution: Floreana, Santa Cruz.

364. *Opegrapha cactacearum* Riedl

Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Indigenous.
IUCN Red List: Least Concern.
Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé.

365. *Opegrapha cf. foreaui* (Moreau) Hafellner & R. Sant.

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.
Origin: Native, Indigenous.
IUCN Red List: Data Deficient.
Galapagos Distribution: Santa Cruz.

366. *Opegrapha diagrapha* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Indigenous.
IUCN Red List: Data Deficient.
Galapagos Distribution: Santa Cruz.

367. *Opegrapha difficilior* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Indigenous.
IUCN Red List: Data Deficient.
Galapagos Distribution: Pinzón, Santa Cruz, Santiago.

368. *Opegrapha herbarum* Mont.

Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Indigenous.
IUCN Red List: Data Deficient.
Galapagos Distribution: Isabela, Pinta, Santa Cruz.

369. *Opegrapha melanospila* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

370. *Opegrapha sp. nov. 1 "norstictica"*

Taxon status: Unpublished name (Nomen nudum).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

371. *Opegrapha trilocularis* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Santa Cruz.

372. *Opegrapha trochodes* Coppins, F. Berger & Ertz

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinta, Santiago.

373. *Opegrapha vulgata* (Ach.) Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Santa Cruz.

374. *Opegrapha xerica* Torrente & Egea

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela.

375. *Pannaria lurida* (Mont.) Nyl.

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz: record based on a minute specimen fragment collectd by Snodgrass & Heller in Fenandina and isolated by B. Weber from the original collection of *Pseudocyphellaria aurata*; locality data are very sparse (Fernandina, 200ft.) and it is not possible to say what original collection the material was isolated from. The specimen was identified by first by Weber as *Erioderma cf. wrightii* Tuck., then annotated by P.M.

Joergensen 1978 as *Pannaria lurida* (Mont.) Nyl.

Origin: No Data.

Galapagos Distribution: Unknown.

376. *Parapyrenis "portoricensis"*

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz: nomen nudum; specimen identified by A. Aptroot (Aptroot 63057) agrees well with *P. aurora*, the only difference being that the Galapagos material has perithecia are broadly elongate, possibly an artifact caused by the perithecia being formed in between the wood fibers of the substrate.

Origin: No Data.

Galapagos Distribution: Santa Cruz.

377. *Parapyrenis aurora* (Zahlbr.) Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

378. *Parmeliella pannosa* (Sw.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: first reported as *Parmeliella pannosa*, which is the correct identification, the Galapagos material does not belong to *Parmiella mariana* as suggested by Elix & McCarthy (1998); Index Fungorum suggests that *P. pannosa* is a synonym of *P. mariana* based on the Australian checklist, this is erroneous, both taxa are good species, *P. pannosa*, however, does not exist in Australia

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

379. *Parmeliella stylophora* (Vain.) P.M. Jørg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

380. *Parmelinella wallichiana* (Taylor) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

381. *Parmelinopsis horrescens* (Taylor) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

382. *Parmelinopsis minarum* (Vain.) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

383. *Parmelinopsis spumosa* (Asahina) Elix & Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santiago.

384. *Parmotrema aberrans* Vain. & Abbeys

Taxon status: Accepted name; taxon occurs in Galapagos.

A. Spielmann: previously erroneously identified as *P. neotropicum*, *P. flavescens* or *P. xanthinum*

Origin: No Data.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santiago.

385. *Parmotrema aff. subisidiosum* (Müll. Arg.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

A. Spielmann: with unusually well developed ciliate isidia

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

386. *Parmotrema cf. bangii* (Vain.) Hale

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

A. Spielmann: only one songle specimen at CDS that still needs TLC for confirmation.

Origin: Not In Galapagos, Not In Galapagos.

Galapagos Distribution: Unknown.

387. *Parmotrema clavuliferum* (Räsänen) Streimann

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Rimelia clavulifera*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

388. *Parmotrema conformatum* (Vain.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

389. *Parmotrema crinitum* (Ach.) M. Choisy

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia crinita*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

390. *Parmotrema cristiferum* (Taylor) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia cristifera*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

391. *Parmotrema dilatatum* (Vain.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

392. *Parmotrema dominicanum* (Vain.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia dominicana*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

393. *Parmotrema eborinum* (Hale) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

no description found, keys out in Sipman's key sensu Hale, also annot. specimens from Hale in COLO loan(L-40237,L-40505);without apothecia,isidia,soredia,± lobulate(i.e., broader lobes with smaller marginal lobes),medulla P+ red,K+ yellow turning orange, C-

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, San Cristóbal.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

394. *Parmotrema endosulphureum* (Hillmann) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia endosulphurea*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

395. *Parmotrema grayanum* (Hue) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

396. *Parmotrema internexum* (Nyl.) Hale ex DePriest & B.W. Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Santiago.

397. *Parmotrema latissimum* (Fée) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, San Cristóbal.

398. *Parmotrema mellissii* (C.W. Dodge) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

399. *Parmotrema praesorediosum* (Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Parmelia praesorediosa*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

400. *Parmotrema reticulatum* (Taylor) M. Choisy

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Rimelia reticulata*, *Parmelia reticulata*, *Parmotrema leucosemethetum*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

401. *Parmotrema sancti-angeli* (Lynge) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

402. *Parmotrema subisidiosum* (Müll. Arg.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Rimelia subisidiosa

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

403. *Parmotrema tinctorum* (Despr. ex Nyl.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Parmelia tinctorum, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago, Wolf.

References: Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

404. *Parmotrema ultralucens* (Krog) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Parmelia subcrinita, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

405. *Parmotrema xanthinum* (Müll. Arg.) Hale

Taxon status: Taxonomic status unresolved or unrevised.

Specimen in COLO (40757) not seen: Cavagnarao s.n., Pinzón, caldera rim, 300 m; F. Bungartz: needs verification

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

406. *Peltigera dolichorrhiza* (Nyl.) Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

407. *Peltigera ulcerata* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

previously misidentified as *P. didactyla* (originally reported by Weber 1986 as *P. erumpens*, later re-identified as *P. spuria* and subsequently cited by Elix & McCarthy as *P. didactyla*), the specimens are not tomentose and thus misidentifications of *P. ulcerata* Müll. Arg.

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Isabela, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

408. *Peltula bolanderi* (Tuck.) Wetmore

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal.

409. *Peltula euploca* (Ach.) Poelt

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Heppia euploca*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Darwin, Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago, Wolf.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

410. *Peltula impressa* (Vain.) Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

material confirmed by M. Schultz

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

411. *Peltula omphaliza* (Nyl.) Wetmore

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española.

References: Kirk, P. (ed.) et al. (2010).

412. *Peltula placodizans* (Zahlbr.) Wetmore

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

413. *Pertusaria albinea* Tuck. [non Müll. Arg.]

Taxon status: Identification not yet confirmed.

A. Aptroot: Uncertain what this species is, depends on the chemistry; Type (FH): on bark, Galapagos Islands, coll. Rev. T. Hill, Hassler Expedition (F. Bungartz: specimen not found during visit to FH, possibly on loan to I. Messuti, one specimen collected and identified by Weber as *P. albinea*, L-40346, COLO 188868).

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

414. *Pertusaria hypothamnolica* Dibben

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz.

415. *Pertusaria pustulata* (Ach.) Duby

Taxon status: Identification not yet confirmed.

In Weber (1986) probably as *Pertusaria bispora*

Origin: No Data.

Galapagos Distribution: Isabela, Santa Cruz.

416. *Pertusaria tejocotensis* de Lesd.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Isabela, Pinta, Pinzón, Santa Cruz, Santa Fé, Santiago.

417. *Pertusaria texana* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Pertusaria bispora* Farl. ex Lindner; Type in Farlow Herbarium (FH): Genovesa, A. Stewart, 153, Isotype in CAS, No. 119734; FH 197376 is *Pertusaria texana*, fide F. Bungartz annotation, 2009

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

418. *Pertusaria xanthodes* Müll. Arg.

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Isabela, Santa Cruz.

419. *Phaeographis atromaculata* (A.W. Archer) A.W. Archer

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Phaeographis kalbii*, fide Bungartz et al. (2009)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Fernandina, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

420. *Phaeographis brasiliensis* (A. Massal.) Kalb & Matthes-Leicht

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

421. *Phaeographis decipiens* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela.

References: Bungartz, F. et al. (2009).

422. *Phaeographis dendritica* (Ach.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

423. *Phaeographis fusca* Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

424. *Phaeographis intricans* (Nyl.) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

425. *Phaeographis leiogrammodes* (Kremp.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

426. *Phaeographis lobata* (Eschw.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

427. *Phaeographis major* (Kremp.) Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

428. *Phaeographis punctiformis* (Eschw.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

429. *Phaeographis striata* Bungartz

Taxon status: Accepted name; taxon occurs in Galapagos.

Holotype Bungartz 6606 (CDS 34826)

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal.

References: Bungartz, F. et al. (2009).

430. *Phaeophyscia pusilloides* (Zahlbr.) Essl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

431. *Phyllopsora confusa* Swinscow & Krog

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

432. *Phyllopsora intermediella* (Nyl.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Santa Cruz, Santiago.

433. *Phyllopsora kalbii* Brako

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

434. *Phyllopsora parvifolia* (Pers.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

435. *Physcia atrostriata* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Moberg, R. et al. (1990).

436. *Physcia crispa* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santiago.

References: Elix, J.A. et al. (1998).

437. *Physcia decorticata* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

438. *Physcia dubia* (Hoffm.) Lettau

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

439. *Physcia erumpens* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz, Santiago.

440. *Physcia integrata* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

441. *Physcia kalbii* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

442. *Physcia lobulata* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

443. *Physcia lopezii* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

444. *Physcia mexicana* B. de Lesd.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: P. mexicana was not included in Moberg (1990), possible syntype material in UPS is in very poor condition and until now was thought to be synonymous with P. aipolia (Moberg, pers. com.); Galapagos material examined by K. Kalb & F. Bungartz does not contain zeorin but it clearly does have a K+ yellow medulla, it is thus chemically distinct from Physcia aipolia and most likely corresponds to what has been described as Physcia mexicana.

Origin: Not In Galapagos, Not In Galapagos.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Moberg, R. et al. (1990), Weber, W.A. et al. (1986).

445. *Physcia poncinsii* Hue

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz.

446. *Physcia rolffii* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz, Santiago.

447. *Physcia sinuosa* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santiago.

448. *Physcia sorediosa* (Vain.) Lyngé

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinzón, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

449. *Physcia undulata* Moberg

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón.

450. *Physma byrsinum* (Ach.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: only one single, historic specimen (Sipman L-70, COLO L-63545, from Cerro Azul, Isabela, H. Sipman L-70, 22-25 June 1976).

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Isabela.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

451. *Piccolia conspersa* (Fée) Hafellner

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinta, Santa Cruz.

452. *Placidium squamulosum* (Ach.) Breuss

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Unknown.

453. *Polychidium muscicola* (Sw.) Gray

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

454. *Polychidium stipitatum* V■zda & W.A. Weber

Taxon status: Identification not yet confirmed.

F. Bungartz: Aptroot 64694 material needs verification

Origin: No Data.

Galapagos Distribution: Santa Cruz.

455. *Porina atrocoerulea* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

456. *Porina cestrensis* (Tuck.) Müll. Arg. agg.

Taxon status: Identification not yet confirmed.

F. Bungartz: a group of very similar species; material needs further studies.

Origin: No Data.

Galapagos Distribution: Santa Cruz.

457. *Porina conspersa* Malme

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz & R. Miranda: only one single fertile specimen in CDS, all others sterile and thus referred to *P. distans* (according to Lücking 2008).

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010), Lücking, R. et al. (2008).

458. *Porina cubana* V■zda

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

459. *Porina distans* V■zda & Vivant

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz & R. Miranda: the sterile material with coralloid isidia most likely belongs to *P. conspersa*, but

since no perithecia could be found treated here according to Lücking (2008) as *P. distans*.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz.

References: Lücking, R. et al. (2008).

460. *Porina guentheri* (Flot.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón, Santa Cruz.

461. *Porina leptalea* (Durieu & Mont.) A.L. Sm.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

462. *Porina melanops* Malme

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz.

463. *Porina nitidula* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

464. *Porina nucula* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

465. *Porina tetramera* (Malme) R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

466. *Pseudocyphellaria argyracea* (Delise) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Floreana, Pinzón, San Cristóbal, Santa Cruz.

References: Elix, J.A. et al. (1998), Galloway, D.J. et al. (1990), Weber, W.A. et al. (1986).

467. *Pseudocyphellaria aurata* (Ach.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Sticta aurata, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Andersson, N. J. et al. (1855), Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Hooker, J.D. et al. (1847), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

468. *Pseudocyphellaria crocata* (L.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pseudocyphellaria mougeotiana var. aurigera, Pseudocyphellaria xantholoma, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Galloway, D.J. et al. (1990), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

469. *Pseudocyphellaria dozyana* (Mont. & Bosch) D.J. Galloway

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Pseudocyphellaria mougeotiana var. aurigera (=? Pseudocyphellaria crocata), fide Weber (1993)

Origin: Native, Questionable Endemic.

IUCN Red List: Vulnerable.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Galloway, D.J. et al. (1985), Galloway, D.J. et al. (1990), Weber, W.A. et al. (1993), Weber, W.A. et al. (1986).

470. *Pseudopyrenula diluta* (Fée) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, Santa Cruz, Santiago.

471. *Pseudopyrenula subnudata* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pseudopyrenula subgregaria, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

472. *Psilolechia lucida* (Ach.) M. Choisy

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela.

473. *Psora nipponica* (Zahlbr.) Gotth. Schneid.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Psora novomexicana, fide Elix & McCarthy (1998), Toninia novomexicana: specimen in COLO (L-44020), Cavagnar s.n., Pinzón, det. E. Timdal, 1990

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998).

474. *Psoroglaena cubensis* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: several specimens confirmed at CDS, but many other identifications erroneous, e.g., a specimen in FH is based on a misidentification of Peltula omphaliza (FH 197404); also some taxonomic confusion: Psoroglaena cubensis auct. non Müll. Arg., fide Elix & McCarthy (1998) is synonymous with Flakea papillata.

Origin: Not In Galapagos, Not In Galapagos.

Galapagos Distribution: Española, Pinta, Santa Cruz, Santiago.

References: Dodge, C.W. et al. (1936), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

475. *Psorotichia cf. hassei* Fink ex J. Hedrick

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: San Cristóbal.

476. *Psorotichia murorum* A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

First erroneously identified as Psorotichia schaeferi by M. Schultz in 2006.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Schultz, M. et al. (2008).

477. *Pterygiopsis guyanensis* Schultz

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Fé.

478. *Pyrenographa irregularis* (Wehm.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: ?*Pyrenographa xylographoides* Aptroot (with submuriform spores, see comments in Harris 1995); basionym: *Phaeopeltosphaeria irregularis* Wehmeyer; Type. ECUADOR. Galapagos: South Seymour Island, on dead, decorticated wood of *Bursera graveolens*, 6 Sep 1945, Martin 6251 (NY, isotype).

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (1991), Harris, R.C. et al. (1995), Martin, G.W. et al. (1948).

479. *Pyrenopsis portoricensis* Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinta, Santiago.

480. *Pyrenothrix nigra* Riddle

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

481. *Pyrenula adacta* Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Pyrenula marginatula* Müll. Arg., *Pyrenula caraibica* Aptroot & Etayo

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013).

482. *Pyrenula aggregata* (Fée) Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Pyrenula costaricensis* Müll. Arg.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Aptroot, A. et al. (2013).

483. *Pyrenula anomala* (Ach.) A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Aptroot, A. et al. (2013).

484. *Pyrenula aspistea* (Afzel.) Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Specimen in COLO: Itow (L-40634), det. Aptroot, 1991; specimens in CDS identified by Aptroot as *P. aspistea* were misidentifications of *Pyrenula costaricensis*, fide annot. R. Miranda, 2010

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (2013), Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

485. *Pyrenula astroidea* (Fée) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Parmentaria astroidea, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz.

References: Aptroot, A. et al. (2013), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

486. *Pyrenula bahiana* Malme

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pyrenula concatervans (Nyl.) R.C. Harris; Specimens in COLO: Santa Cruz: Herre 41 (L-41177), Weber (L-40220), Itow (L-40728), Fernandina: Cavagnaro (L-40469), Floreana: Weber & Lanier (L-62944)

Origin: Native, Indigenous.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013).

487. *Pyrenula breutelii* (Müll. Arg.) Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pyrenula macularis; in Weber (1986) probably as *Anthracothecium leucostomum*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013).

488. *Pyrenula cerina* Eschw.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pyrenula aurantiaca, fide Elix & McCarthy (1998); F. Bungartz: species not included in the key by Aptroot (2012), only listed as accepted in the appendix

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Genovesa, Isabela, Marchena, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2013), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

489. *Pyrenula cf. acutispora* Kalb & Hafellner

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

F. Bungartz & R. Miranda: spore measurements of material do not agree with *P. acutispora* s.str.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, Santa Cruz.

490. *Pyrenula cf. cuyabensis* (Malme) R.C. Harris

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

Basionym: *Parathelium cuyabense* Malme

Origin: Native, Indigenous.

Galapagos Distribution: San Cristóbal, Santa Cruz.

491. *Pyrenula cf. dermatodes* (Borrer) Schaer.

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

Origin: Native, Indigenous.

Galapagos Distribution: Santa Cruz.

492. *Pyrenula confinis* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz.

References: Aptroot, A. et al. (2013).

493. *Pyrenula cruenta* (Mont.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Melanotheca cruenta*, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

494. *Pyrenula dermatodes* (Borrer) Schaer.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Aptroot, A. et al. (2013), Kirk, P. (ed.) et al. (2010).

495. *Pyrenula erumpens* R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Specimen in COLO: Santa Cruz, on Pisonia, Itow (L-40623), det. Aptroot, 1991

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (2013), Elix, J.A. et al. (1998), Weber, W.A. et al. (1993).

496. *Pyrenula fetivica* (Kremp.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn. *Pyrenula subcongruens* Müll. Arg.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (2013), Kirk, P. (ed.) et al. (2010).

497. *Pyrenula massariospora* (Starbäck) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013), Kirk, P. (ed.) et al. (2010).

498. *Pyrenula microtheca* R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: non *Pyrenula microcarpa* Müll. Arg.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Aptroot, A. et al. (2013).

499. *Pyrenula neosandwicensis* Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: previously treated as *P. aff. duplicans*, Syn.: *Anthracothecium sandwicense* Zahlbr.; not identical with *Pyrenuma sandwicensis* Zahlbr., which is a synonym of *P. fetivica* (Kempelh.) Müll. Arg.

Origin: Native, Indigenous.

Galapagos Distribution: Floreana, Santa Cruz.

References: Aptroot, A. et al. (2013).

500. *Pyrenula nitidula* (Bres.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz.

References: Aptroot, A. et al. (2013), Kirk, P. (ed.) et al. (2010).

501. *Pyrenula ochraceoflava* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Anthracothecium ochraceoflavum*, fide Elix & McCarthy (1998), *Verrucaria ochraceoflava*, fide Index Fungorum

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal,

Santa Cruz, Santa Fé, Santiago, Wolf.

References: Aptroot, A. et al. (2013), Elix, J.A. et al. (1998), Stewart, A. et al. (1912), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

502. *Pyrenula ochraceoflavens* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz & R. Miranda: very few specimens deviate sufficiently in spore size from *P. ochraceoflava* and the two taxa might refer to the same species.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Aptroot, A. et al. (2013), Kirk, P. (ed.) et al. (2010).

503. *Pyrenula quassiaecola* Fée

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013).

504. *Pyrenula thelomorpha* Tuck.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2013), Kirk, P. (ed.) et al. (2010).

505. *Pyrgidium montellicum* (Beltr.) Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

506. *Pyrgillus javanicus* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

507. *Pyrrhospora quernea* (Dicks.) Körb.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: No Data.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal.

508. *Pyxine albovirens* (G. Mey.) Aptroot 1987

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

509. *Pyxine berteroana* (Fée) Imshaug

Taxon status: Accepted name; taxon occurs in Galapagos.

The spelling P. berteriana is an orthographical error, fide F. Bungartz

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

510. *Pyxine cocoës* (Sw.) Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

511. *Pyxine eschweileri* (Tuck.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Weber, W.A. et al. (1986).

512. *Pyxine petricola* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pyxine pringlei, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

513. *Pyxine subcinerea* Stirn.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Pyxine caesiopruinosa, fide Elix & McCarthy (1998), Pyxine connectens, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

514. *Ramalina anceps* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2007), Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

515. *Ramalina aspera* Räsänen

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Ramalina denticulata, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007), Brodo, I. M. et al. (2001), Weber, W.A. et al. (1986).

516. *Ramalina campylospora* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Pinzón, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2007).

517. *Ramalina complanata* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Ramalina indica auct. non Fr., Ramalina subasperata auct. non Nyl., Ramalina interponens auct. non Nyl., Ramalina subfraxinea auct. non Nyl., fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007), Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

518. *Ramalina darwiniana* var. *curvula* Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: Ecuador, Galapagos, Santa Cruz Island, Puerto Ayora, near the Charles Darwin Research Station, 0°44'32"S, 90°18'10"W, alt. 5 m, on twigs of coastal shrubs, 24 May 2005, A. Aptroot 63029 (CDS no. 29757—holotype; ABL—isotype).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Floreana, Isabela, Santa Cruz.

References: Aptroot, A. et al. (2007).

519. *Ramalina darwiniana* var. *darwiniana* Aptroot & Bungartz

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR, Galapagos, Plaza Sur Island, 0°34'59"S, 90°9'54"W, alt. 1 m, coastal zone; eastern part with scattered and low vegetation of Sesuvium portulacastrum & Tiquilia galapagoa with occasional Opuntia, western part also with scattered shrubs of Grabowskia boerhaaviaefolia, Maytenus octogona, and Castela galapageia, on wood, twig, 21-Feb-2006, A. Aptroot, 64433 (CDS no. 31001—holotype; ABL—isotype).

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007).

520. *Ramalina fragilis* Aptroot & Bungartz

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Niebla sp. nov. ined.; Typus: ECUADOR, Galapagos, San Cristóbal, near Tortugueria Cerro Colorado, on lava cliff, 130 m alt., 2 June 2005, A. Aptroot 63419 (CDS 30174—holotypus; ABL—isotypus).

Origin: Native, Endemic.

IUCN Red List: Vulnerable.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Fé.

References: Aptroot, A. et al. (2007), Weber, W.A. et al. (1986).

521. *Ramalina furcellangulida* Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Typus: Ecuador, Galapagos, Isabela, Volcán Alcedo, highest cinder cone along the trail going up the east slope, on bark of Bursera graveolens, 250 m alt., 10 March 2006, A. Aptroot 65029 (CDS 31611—holotypus; ABL—isotypus).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2007).

522. *Ramalina montagnei* De Not.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2007).

523. *Ramalina pacifica* Asah.

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz: some specimens of R. sorediantha with more flattened lobes may belong to this species.

Origin: No Data.

IUCN Red List: Not Evaluated.

Galapagos Distribution: Pinzón, San Cristóbal, Santa Cruz.

References: Aptroot, A. et al. (2007).

524. *Ramalina peruviana* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007), Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

525. *Ramalina polyformata* Aptroot

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: Ecuador, Galapagos, Santa Cruz Island, on coastal lava cliffs E of Puerto Ayora near Charles Darwin Research Station, 20 m alt., 29 May 2005, A. Aptroot 63412 (CDS 30176—holotype; ABL—isotype).

Origin: Native, Endemic.

IUCN Red List: Vulnerable.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007).

526. *Ramalina puiggarii* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Ramalina linearis, fide Aptroot & Bungartz (2007)

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Española, Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2007), Dodge, C.W. et al. (1935), Dodge, C.W. et al. (1936), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

527. *Ramalina sideriza* Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007).

528. *Ramalina sorediantha* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007), Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

529. *Ramalina sorediosa* (B. de Lesd.) Landrón

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Ramalina furcellata*, fide Aptroot & Bungartz (2007); In Stewart (1912) & Weber (1966, 1981) as *Ramalina farinacea*; In Dodge (1936) & Weber (1966) as *Ramalina dasypoga*

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2007), Dodge, C.W. et al. (1936), Kashiwadani, H. et al. (1993), Landrón C.I. et al. (1972), Stewart, A. et al. (1912), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

530. *Ramalina usnea* (L.) R. Howe

Taxon status: Accepted name; taxon occurs in Galapagos.

In Stewart (1912) and Linder (1934) as *Alectoria sarmentosa*, fide Weber (1966)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2007), Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Linder, D.H. et al. (1934), Stewart, A. et al. (1912), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

531. *Ramboldia heterocarpa* (Fée) Kalb, Lumbsch & Elix

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

532. *Ramonia valenzueliana* (Mont.) Stizenb.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

533. *Redonographa galapagoensis* Bungartz & Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR. Galapagos: Santiago Island, ca. 5 km inland from the E-coast; 0° 16' S, 90° 37' W; Bungartz 5208 (CDS 29421, holotype); previously reported as *Carbacanthographis saxiseda* (Bungartz et al., 2010) but was found to represent an undescribed taxon (Lücking et al. 2013).

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Unknown.

References: Bungartz, F. et al. (2009), Elix, J.A. et al. (1998), Lücking, R. et al. (2103), Weber, W.A. et al. (1993).

534. *Redonographa saxorum* (Egea & Torrente) Lücking & Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Graphis saxorum, Carbacanthographis saxorum.

Origin: Native, Indigenous.

Galapagos Distribution: Unknown.

References: Bungartz, F. et al. (2009), Lücking, R. et al. (2103).

535. *Rhizoplacea minarum* Kalb, Elix & Bungartz ined.

Taxon status: Unpublished name (Nomen nudum).

Origin: Native, Questionable Endemic.

Galapagos Distribution: Isabela, Santiago.

536. *Rinodina lepida* (Nyl.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela.

537. *Rinodina olea* Bagl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Specimen of 'Buellia myriocarpa' FH 1108 (00197447) identified by J.W. Sheard, 2009 as *R. olea*.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

538. *Roccella albida* Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Holotype Tehler 8653 (CDS 40614)

Origin: Native, Endemic.

IUCN Red List: Endangered.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Tehler, A. et al. (2009).

539. *Roccella galapagoensis* Follmann

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Roccella geniculata, R. glebaria, R. obscurissima Follmann & B. Werner, R. geniculata Follmann & B. Werner, R. fusca B. Werner nom. nud., R. glebaria B. Werner & Follmann nom. nud., R. colonii Follmann, R. capitata Follmann nom. nud., fide Tehler et al. (2009); Holotype: Galápagos, South Plaza Island, abundant on seaward vertical sides of boulders on highest points (S side), ca. 50 ft elev., 1964, W.A. Weber L-39130, Lichenes Exsiccati no. 112; Isotypes: COLO-184895, 186050, UPS, M-0024655, H, S-L21409

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Schofield, E.K. et al. (1984), Tehler, A. et al. (2009), Tehler, A. et al.

(2007), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

540. *Roccella gracilis* Bory

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Roccella babingtonii*, R. *peruensis*, R. *difficilis*, fide Tehler (2002) and Tehler et al. (2009)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Darbshire, O.V. et al. (1935), Dodge, C.W. et al. (1935), Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Tehler, A. et al. (2009), Tehler, A. et al. (2002), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

541. *Roccella margaritifera* B. Werner & Follmann

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Roccella octopodioides* nom. nud., *Roccella margaritifera* f. *octopodioides* fide Tehler et al. (2009);

Holotype: Galápagos, San Cristóbal Island, Lobería Pto. Baquerizo, 1991, Sánchez-Pinto 6616 (B-128629), TFMC

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Genovesa, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Tehler, A. et al. (2009), Tehler, A. et al. (2007).

542. *Roccella nigerrima* (Darb.) Follmann

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Roccellodea nigerrima*, *Roccella translucida* Follmann & B. Werner, *Roccella incurvata* B. Werner, *Roccella kappenniana* Follmann & B. Werner, *Roccella botrytis* B. Werner nom. nud., *Roccella floribrassica* B. Werner nom. nud., *Roccella stipitata* B. Werner & Follmann nom. nud., *Roccella floteana* Follmann nom. nud., *Roccella floreana* B. Werner nom. nud. fide Tehler et al. (2009); Lectotype selected by Tehler (2007): Galápagos Islands, 1872, Hill s.n. (FH)

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Tehler, A. et al. (2009).

543. *Roccellographa circumscripta* (Taylor) Ertz & Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Sclerophytomyces circumscripti* Cif. & Tomas.; *Sclerophyton cricumscriptum* (Taylor) Zahlbr., *Peterjamesia circumscripta* (Taylor) D. Hawksw.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Unknown.

References: Aptroot, A. et al. (2008), Ertz, D. et al. (2010).

544. *Sarcographa medusulina* (Nyl.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

545. *Sarcographa ramificans* (Kremp.) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Bungartz, F. et al. (2009).

546. *Sarcographa tricosa* (Ach.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

547. *Schismatomma spierii* Aptroot & Sparrius

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR. Galapagos: Isabela Island, Cerro Alcedo, highest cinder cone along the trail going up the E slope, 0° 23'37"S, 91°01'31"W, 250 m, arid lowlands with basalt outcrops, at the bottom scattered Bursera graveolens trees and shrubs of Castela galapageia, on bark of Bursera graveolens, 10 Mar 2006, Aptroot 65014 (holotype CDS 31595; isotype ABL).

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

548. *Schistophoron tenuie* Stirz.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Pinta, Santa Cruz.

References: Elix, J.A. et al. (1998), Tehler, A. et al. (2009), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

549. *Schistophoron variabile* Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Regionally Extinct.

Galapagos Distribution: Española, Floreana, Santa Cruz.

550. *Sclerococcum* sp. 1

Taxon status: Taxon not identified to species, subspecies, form or variety.

Index Fungorum: anamorphic Pezizomycotina

Origin: No Data.

Galapagos Distribution: San Cristóbal, Santa Cruz.

551. *Sclerophyton murex* Egea & Torrente ex Sparrius

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008).

552. *Sclerophyton vertex* Sparrius

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2008).

553. *Sigridea leptothallina* (Malme) Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

The spelling in Index Fungorum is incorrect, the correct name is Sigridea leptothallina [Tehler, A. 1993. The genus Sigridea [Roccellaceae, Arthoniales, Euascomycetes. Nova Hedwigia 57(3-4): 417-435]

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela, San Cristóbal, Santa Fé.

References: Aptroot, A. et al. (2008).

554. *Sphinctrina leucopoda* s.l. Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Sphinctrina podocarpa, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Fé.

555. *Sphinctrina tubiformis* A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Index Fungorum as Sphinctrina tubiformis

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, Santiago.

556. *Spilonema revertens* Nyl.

Taxon status: Identification not yet confirmed.

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

557. *Sporopodium aff. phyllocharis* (Mont.) A. Massal.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

558. *Sporopodium leprieurii* Mont.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

559. *Staurothele lecideoides* B. de Lesd.

Taxon status: Identification not yet confirmed.

material examined by Breuss, morphologically identical to *S. lecideoides*, but no mature perithecia found, perithecia contain hymenial algae

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

560. *Stereocaulon azulense* Yoshim. & W.A. Weber

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Marchena, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

561. *Stereocaulon microcarpum* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

562. *Stereocaulon weberi* I.M. Lamb

Taxon status: Accepted name; taxon occurs in Galapagos.

Holotype in FH

Origin: Native, Endemic.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Lamb, I.M. et al. (1977), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

563. *Sticta beauvoisii* Delise

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Kirk, P. (ed.) et al. (2010), McDonald, T. et al. (2003).

564. *Sticta carolinensis* T. McDonald

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz.

References: Kirk, P. (ed.) et al. (2010), McDonald, T. et al. (2003).

565. *Sticta damicornis* (Sw.) Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Index Fungorum: the species epithet was originally published a 'damaecornis'.

Origin: Native, Indigenous.

IUCN Red List: Critically Endangered.

Galapagos Distribution: Santa Cruz.

566. *Sticta dichotoma* s.l. Delise

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1966).

567. *Sticta fuliginosa* (Dicks.) Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), McDonald, T. et al. (2003), Weber, W.A. et al. (1986).

568. *Sticta weigelii* (Ach.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Stewart (1912) as *Sticta quercizans*, fide Weber (1966); In Weber (1966) as *Sticta weigelii* var. *peruviana*;

Syn.: *Sticta quercizans*, fide Elix & McCarthy (1998), *Stictina quercizans*, fide Index Fungorum

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Svenson, H.K. et al.

(1935), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

569. *Strigula cf. obtecta* (Vain.) R. C. Harris

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, Santa Cruz.

570. *Strigula microspora* Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

571. *Strigula nitidula* Mont.

Taxon status: Accepted name; taxon occurs in Galapagos.

Specimen in COLO: Santa Cruz: on leaves of Eugenia jambos, along trail to Horneman place just above Bellavista, 64131 p.p. (with Byssoloma subdiscordans), det. Vezda; F. Bungartz: specimen not seen!

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

572. *Strigula phaea* (Ach.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

573. *Strigula phyllogena* (Müll. Arg.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: No Data.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

574. *Strigula smaragdula* Fr.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz & R. Miranda: specimen material identified by R. Lücking very poor.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

575. *Strigula subtilissima* (Fée) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Specimen in COLO: Santa Cruz: on leaves of Eugenia jambos, along trail to Horneman place just above Bellavista, 64131 p.p. (with *Byssoloma subdiscordans*), det. Vezda, F. Bungartz: specimen not seen!

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

576. *Strigula viridiseda* (Nyl.) R.C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal.

References: Kirk, P. (ed.) et al. (2010).

577. *Synalissa mattogrossensis* (Malme) Henssen

Taxon status: Accepted name; taxon occurs in Galapagos.

material confirmed by M. Schultz

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Española, Santa Cruz, Santiago.

578. *Syncesia farinacea* (Fée) Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Chiodecton farinaceum*

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, Pinzón, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008).

579. *Syncesia graphica* (Fr.) Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Chiodecton myrticola*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008), Weber, W.A. et al. (1986).

580. *Syncesia leprobola* Nyl. ex Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Chiodecton farinaceum*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Aptroot, A. et al. (2008), Weber, W.A. et al. (1986).

581. *Syncesia psaroleuca* (Nyl.) Tehler

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as Chiodecton effusum, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Aptroot, A. et al. (2008), Weber, W.A. et al. (1986).

582. *Taeniolella sp. 1*

Taxon status: Taxon not identified to species, subspecies, form or variety.

Index Fungorum: anamorphic Glyphium

Origin: No Data.

Galapagos Distribution: Isabela, Santa Cruz.

583. *Tapellaria epiphylla* (Müll. Arg.) R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

584. *Tapellaria granulosa* Lücking & Rivas Plata

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

585. *Tapellaria malmei* R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

586. *Tapellaria nana* (Fée) R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

587. *Tapellaria nigrata* (Müll. Arg.) R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: San Cristóbal, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

588. *Tapellaria phyllophila* (Stirt.) R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

589. *Teloschistes chrysophthalmus* (L.) Th. Fr.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Vulnerable.

Galapagos Distribution: Isabela.

590. *Teloschistes flavicans* (Sw.) Norman

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Dodge, C.W. et al. (1936), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Weber, W.A. et al. (1981), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

591. *Tephromela rhizophorae* Kalb

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Kalb, K. et al. (2008).

592. *Tephromela weberi* Kalb, Aptroot & Bungartz ined.

Taxon status: Unpublished name (Nomen nudum).

Origin: No Data.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santiago.

593. *Thalloloma cinnabarinum* (Fée) Staiger

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Endangered.

Galapagos Distribution: Floreana, Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Bungartz, F. et al. (2009).

594. *Thamnolia vermicularis* (Sw.) Ach. ex Schaer.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

595. *Thelenella inductula* (Nyl.) H. Mayrhofer

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

596. *Thelenella muscorum* (Fr.) Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

597. *Thelenella sastreana* R. C. Harris

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

598. *Thelopsis isiaca* Stitzenb.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

599. *Thelopsis rubella* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

600. *Thelotrema monosporum* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela.

References: Kirk, P. (ed.) et al. (2010).

601. *Thelotrema pachysporum* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

602. *Trapelia coarctata* (Turner ex Sm.) M. Choisy

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

603. *Trapelia involuta* (Taylor) Hertel

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

604. *Trapelia placodioides* Coppins & P. James

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz.

605. *Trapeliopsis flexuosa* (Fr.) Coppins & P. James

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santiago.

606. *Trapeliopsis glaucopholis* (Nyl.) Printzen & McCune

Taxon status: Identification not yet confirmed.

F. Bungartz: material needs verification

Origin: No Data.

Galapagos Distribution: Santa Cruz.

607. *Trapeliopsis granulosa* (Hoffm.) Lumbsch

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Lecidea granulosa, fide Elix & McCarthy (1998); specimen in COLO (63337), Santa Cruz, on plant debris, saddle between El Puntudo and Cerro Crocker, 700 m

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santa Cruz.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

608. *Tricharia melanothrix* Fée

Taxon status: Identification not yet confirmed.

F. Bungartz: folicolous specimens in COLO det. as *T. melanothrix* need revision.

Origin: No Data.

Galapagos Distribution: Santa Cruz.

609. *Tricharia vainioi* R. Sant.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

610. *Trichothelium epiphyllum* Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

611. *Trichothelium montanum f. montanum* Lücking

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

612. *Trypethelium aeneum* (Eschw.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: material needs verification

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Isabela, Santiago.

613. *Trypethelium aff. marcidum* (Fée) Müll. Arg.

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz & R. Miranda: specimens with different chemistry and spore size from *T. marcidum* s.str.

Origin: Native, Indigenous.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

614. *Trypethelium eluteriae* Spreng.

Taxon status: Accepted name; taxon occurs in Galapagos.

F. Bungartz: specimens confirmed by R. Miranda & R. Lücking

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

615. *Trypethelium nitidiusculum* (Nyl.) R.C. Harris

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz & R. Miranda: According to Harris (1995) Trypethelium ochroleucum and Astrothelium variosolum are identical in all characters but their stromata formation; Trypethelium nitidiusculum is identical to Trypethelium ochroleucum in all characters but the presence of lichexanthone. It is possible that all three taxa refer to one variable species only.

Origin: No Data.

Galapagos Distribution: Isabela, Pinta, Pinzón, Santa Cruz.

References: Harris, R.C. et al. (1995).

616. *Trypethelium ochroleucum* (Eschw.) Nyl.

Taxon status: Taxonomic status unresolved or unrevised.

F. Bungartz & R. Miranda: According to Harris (1995) Trypethelium ochroleucum and Astrothelium variosolum are identical in all characters but their stromata formation; Trypethelium nitidiusculum is identical to Trypethelium ochroleucum in all characters but the presence of lichexanthone. It is possible that all three taxa refer to one variable species only.

Origin: No Data.

Galapagos Distribution: Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Harris, R.C. et al. (1995), Weber, W.A. et al. (1986).

617. *Trypethelium tropicum* (Ach.) Müll. Arg.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

618. *Tylophorella pyrenocarpoides* (Müll. Arg.) Egea

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz, Santa Fé.

619. *Tylophoron cf. protrudens* Nyl.

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

F. Bungartz: Identification problematic; specimens need TLC, spot tests: C+ red, but UV-

Origin: No Data.

Galapagos Distribution: Isabela, Pinta, Santa Cruz, Santiago.

620. *Tylophoron galapagoense* Bungartz, Ertz, Diederich & Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Holotype Ertz 11794 (CDS 37153)

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

References: Ertz, D. et al. (2011).

621. *Tylophoron hibernicum* (D. Hawksw., Coppins & P. James) Ertz, Diederich, Bungartz & Tibell

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Blarneya hibernica D. Hawksw., Coppins & P. James

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Fernandina, San Cristóbal, Santa Cruz, Santiago.

References: Ertz, D. et al. (2011).

622. *Tylophoron moderatum* Nyl.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

623. *Usnea angulata* Ach.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1981, 1986) and Elix & McCarthy (1998) as *Usnea paradoxa*; no recent specimens; all collections from before 1972, presumed extinct!

Origin: Native, Indigenous.

IUCN Red List: Regionally Extinct.

Galapagos Distribution: Santa Cruz.

References: Elix, J.A. et al. (1998), Truong, C. et al. (2013), Weber, W.A. et al. (1981), Weber, W.A. et al. (1986).

624. *Usnea baileyi* (Stirt.) Zahlbr.

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) as *Usnea antillarum*, fide A. Aptroot (pers. comm.)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Fernandina, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Truong, C. et al. (2013).

625. *Usnea brattiae* P. Clerc

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

626. *Usnea cf. bornmuelleri* J. Steiner

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

Origin: Native, Indigenous.

IUCN Red List: Not Evaluated.

Galapagos Distribution: Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz.

627. *Usnea cirrosa (classical)* Motyka

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz.

628. *Usnea cirrosa (hirto-cirrosa)* Motyka

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santiago.

629. *Usnea columbiana* Motyka

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

630. *Usnea cornuta* Körb.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz, Santiago.

631. *Usnea dodgei* Motyka

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

Galapagos Distribution: Isabela, San Cristóbal.

References: Truong, C. et al. (2013).

632. *Usnea dorogawensis* Asahina

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: JAPAN. Honshu: Prov. Yamato, Dorogawa, Amakawa-mura, Yoshino-gun, 1952, Togashi (lectotype TNS!); % C/M/A: 4/37.5/16; contains usnic, lobaric, norstictic, stictic and constictic acids, unknown US6 (Ohmura 2001).

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Ohmura, Y. et al. (2001), Truong, C. et al. (2011).

633. *Usnea erinacea* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Floreana, Pinzón.

References: Clerc, P. et al. (2008), Clerc, P. et al. (2011), Truong, C. et al. (2011).

634. *Usnea galapagona* Truong & P.Clerc

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: Ecuador. Galapagos: Isla Sán Cristóbal, Cerro Mundo, at the top of the rock cliffs on the S side close to the summit, 00°53'S, 89°34'W, 282 m, transition zone with Bursera graveolens, Croton scouleri and Jasminocereus thouarsii, on Jasminocereus thouarsii on the ridge, August 2008, Clerc & Truong 08-405(holotype CDS, isotypes G, CMA: 16/3/61.5; chemistry: usnic acid, unknown medullary metabolite reacting UV+ green after charring).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinta, San Cristóbal, Santa Cruz.

References: Herrera-Campos, M. et al. (1998), Lumbsch, H.T. et al. (2010).

635. *Usnea grandisora* Truong & P. Clerc

Taxon status: Accepted name; taxon occurs in Galapagos.

Type: ECUADOR. Galapagos: Santa Cruz Island, above Mina Granillo, 607 m, upper transition zone, on branches of Scalesia, Truong 1122 (holotype CDS 39433, isotypes G, UPS; % C/M/A: 11/20.5/38. Chemistry: usnic, salazinic, galbinic and norstictic acids.)

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Ohmura, Y. et al. (2010), Swinscow, T. et al. (1979), Truong, C. et al. (2011).

636. *Usnea hirta* ssp. *trachista* (Motyka) P. Clerc

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal.

637. *Usnea mexicana* Vain.

Taxon status: Accepted name; taxon occurs in Galapagos.

Erronously reported as Usnea longissima, U. amabilis or U. arthroclada by Farlow (1902), Stewart (1912), Weber (1966, 1986), Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Floreana, Isabela, Pinta, Pinzón, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Stewart, A. et al. (1912), Truong, C. et al. (2013), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

638. *Usnea parvula* Motyka

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela.

639. *Usnea poliotrix* Kremp.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

References: Motyka, J. et al. (1936–38), Truong, C. et al. (2011), Vareschi, V. et al. (1973).

640. *Usnea rubicunda* Stirt.

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Usnea rubiginea*, fide Elix & McCarthy (1998), *U. rubiginosa*, fide Index Fungorum; In Farlow (1902), Stewart (1912) and Weber (1966) as *Usnea ceratina*; Type: ENGLAND: 1879, Holmes (holotype BM!; contains usnic, stictic, constictic, menegazziaic and norstictic acids.).

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Fernandina, Floreana, Isabela, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Clerc, P. et al. (2008), Elix, J.A. et al. (1998), Farlow, W.G. et al. (1902), Ohmura, Y. et al. (2008), Ohmura, Y. et al. (2001), Stewart, A. et al. (1912), Truong, C. et al. (2011), Weber, W.A. et al. (1966), Weber, W.A. et al. (1986).

641. *Usnea subcornuta* Stirt.

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Pinzón, San Cristóbal, Santa Cruz.

642. *Usnea subdasaea* Truong & P. Clerc

Taxon status: Unpublished name (Nomen nudum).

Type: Ecuador. Galapagos, Isabela, road to Sierra Negra crater, close to la Esperanza, 306 m, farming areas in the humid zone, living fencepost (still with bark) Truong 1194 (holotype CDS 39505, isotypes G, UPS; % C/M/A: 5/30/30. contains usnic, salazinic, galbinic, norstictic acids.)

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz.

References: Clerc, P. et al. (2008), Herrera-Campos, M. et al. (2001), Truong, C. et al. (2011).

643. *Usnea subscabrosa* Nyl. ex Motyka

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, Santa Cruz.

644. *Vainionora sp. nov.*

Taxon status: Unpublished name (Nomen nudum).

Origin: Native, Endemic.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santa Cruz.

645. *Verrucaria lecideoides* (A. Massal.) Gueidan & C. Roux

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Santiago.

646. *Verrucaria xyloxena* Norman

Taxon status: Accepted name; taxon occurs in Galapagos.

material confirmed by O. Breuss (2011)

Origin: No Data.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal.

647. *Vouauxiomycetes sp. I*

Taxon status: Taxon not identified to species, subspecies, form or variety.

Index Fungorum: anamorphic Abrothallus (Pezizomycotina)

Origin: No Data.

Galapagos Distribution: Santa Cruz.

References: Kirk, P. (ed.) et al. (2010).

648. *Xanthomendoza weberi* (S.Y. Kondr. & Kärnefelt) L. Lindblom

Taxon status: Accepted name; taxon occurs in Galapagos.

In Weber (1986) and Elix & McCarthy (1998) as Xanthoria candelaria, fide F. Bungartz annotations, 2006

Origin: Native, Indigenous.

IUCN Red List: Near Threatened.

Galapagos Distribution: Isabela, Santa Cruz, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

649. *Xanthoparmelia congensis* (J. Steiner) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Indigenous.

IUCN Red List: Data Deficient.

Galapagos Distribution: Isabela, San Cristóbal, Santa Cruz, Santiago.

650. *Xanthoparmelia subramigera* (Gyeln.) Hale

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: Parmelia subramigera, fide Elix & McCarthy (1998)

Origin: Native, Indigenous.

IUCN Red List: Least Concern.

Galapagos Distribution: Española, Floreana, Isabela, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Elix, J.A. et al. (1998), Weber, W.A. et al. (1986).

651. *Xylographa parallela* (Ach.) Fr.

Taxon status: Identification not yet confirmed.

F. Bungartz: possibly misidentifications of *Pyrenographa irregularis* (Wehm.) R.C. Harris

Origin: No Data.

Galapagos Distribution: Santa Cruz.

652. *Xylographa vitiligo* (Ach.) J.R. Laundon

Taxon status: Identification not yet confirmed.

F. Bungartz: possibly misidentifications of *Pyrenographa irregularis* (Wehm.) R.C. Harris

Origin: No Data.

Galapagos Distribution: Santa Cruz.

653. *cf. Baflavia flavesrens* (Lücking) Lücking

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

F. Bungartz: material most likely belongs to an undescribed Eugeniella

Origin: No Data.

Galapagos Distribution: Floreana, Santa Cruz.

654. *cf. Nadvornikia sp.*

Taxon status: The identification of this taxon or its occurrence in Galapagos is doubtful.

Origin: No Data.

Galapagos Distribution: Pinta.

Rejected taxa

1. *Acarospora cf. fuscata* (Nyl.) Arnold

F. Bungartz: material previously identified as A. cf. fuscata has been annotated by K. Knudsen as A. sparsiuscula.

2. *Acarospora citrina* (Taylor) Zahlbr.

Misidentification for A. chrysops, fide K. Knudsen (pers. comm., 2007)

3. *Acarospora schleicheri* (Ach.) A. Massal.

Misidentification for A. chrysops, fide K. Knudsen (pers. comm., 2007)

4. *Acarospora thelococcoides* (Nyl.) Zahlbr.

F. Bungartz: the only specimen identified by Aptroot (Aptroot 64834) as A. thelococcoides is A. americana according to K. Knudsen.

5. *Alectoria sarmentosa* (Ach.) Ach.

Misidentifications of *Ramalina usnea*, fide Weber (1966) & fide F. Bungartz annotations, 2009

6. *Anisomeridium anisolobum* (Müll. Arg.) Aptroot

F. Bungartz & R. Miranda: probably all misidentifications of A. leptospermum; Syn.: *Ditremis anisoloba*, fide Elix & McCarthy (1998); one specimen in COLO has not yet been revised (L-40404) det. Aptroot, 1991

7. *Arthonia parantillarum* Aptroot
Misidentifications of *A. antillarum*, fide Bungartz (2010)
8. *Arthopyrenia punctiformis* A. Massal.
F. Bungartz & R. Miranda: specimens previously cited as *A. punctiformis* are based on misidentifications.
9. *Arthothelium spilomatooides* (Nyl.) Zahlbr.
Misidentification of *Arthothelium galapagoense*, fide Weber (1981)
10. *Aspidothelium fugiens* (Müll. Arg.) R. Sant.
F. Bungartz: in Weber (1986) erroneously cited as *Aspidophyllum fugiens*; material Weber 285 (L-40433).
The material was originally determined by Vezda, but with publication of Lücking (2008) the species concept has changed; the Galapagos specimens have perithecia with disk-like, dentate expansion and not setae or hairs and thus belongs to *A. scutellicarpum*
11. *Buellia xanthinula* (Müll. Arg.) Malme
Misidentification of *Buellia straminea*, fide Weber (1986): 488; misidentification of *Buellia flavoareolata*, fide F. Bungartz annotations, 2009
12. *Bulbothrix goebelii* (Zenker) Hale
M. Benatti, A. Spielmann & F. Bungartz: specimens previously identified as *B. goebelii* have isidia and thus don't belong to *B. goebelii*; most material belongs to *B. subdissecta*.
13. *Calicium abietinum* Pers.
Misidentifications of *Chaenothecopsis aff. savonica* fide Tibell (2010)
14. *Calicium hyperelloides* Nyl.
Possibly misidentifications of *C. robustellum* fide Tibell (2010)
15. *Caloplaca cirrochroa* (Ach.) Th. Fr.
Syn.: *Placodium cirrochroum*; misidentifications of *C. cf. cupulifera* fide U. Søchting
16. *Caloplaca elegans* (Link) Th. Fr.
Misidentifications of *Caloplaca isidiosa* fide U. Søchting, 2010
17. *Canoparmelia crozalsiana* (de Lesd.) Elix & Hale
F. Bungartz: misidentifications, all specimens belong to *C. cf. raunkierii*
18. *Carbacanthographis saxiseda* (Zahlbr.) Bungartz
F. Bungartz: all Galapagos specimens previously identified as *C. saxiseda* by Bungartz & Lücking (2009) belong to *Redonographa galapagoensis*; they were originally reported by Weber (1993) as *Graphina saxiseda* and Elix & McCarthy (1998) as *Graphis saxiseda*.
19. *Celothelium aciculiferum* (Nyl.) Vain.
F. Bungartz & R. Miranda: misidentifications of *Celothelium dominicanum*.
20. *Chaenotheca brunneola* (Ach.) Müll. Arg.
L. Tibell: *Chaenothecopsis savonica* s.l.
21. *Chiodecton myrticola* Feé
Acc. name: *Syncesia myrticola*; Misidentification of *Syncesia psaroleuca*, fide F. Bungartz annotations, 2008

22. *Chrysotrichia candelaris* (L.) J.R. Laundon
Misidentification of *Chrysotrichia xanthina*, fide A. Aptroot (pers. comm.)
23. *Cladina sandstedei* (Abbayes) Ahti
Misidentification of *Cladonia arcuata* (see Yáñez-Ayabaca et al. 2013)
24. *Cladonia coccifera* agg. (L.) Willd.
F. Bungartz: misidentifications of *C. corymbosula* (see Yáñez-Ayabaca et al. 2013)
25. *Cladonia furcata* f. *adspersa* (Flörke) Vain.
F. Bungartz: most likely misidentifications of *C. sphacelata*
26. *Cladonia granulosa* (Vain.) Ahti
F. Bungartz: one specimen [Bungartz 3272 (CDS 26911)] first identified by T. Ahti as *C. granulosa* (annotation 2010); later revised to *C. polycypha* (annotation 2011) see also Yáñez-Ayabaca et al. (2013)
27. *Cladonia macilentoidea* Ahti & Fleig
F. Bungartz & A. Yáñez: misidentifications of *C. macilenta*
28. *Cladonia polycarpoides* Nyl.
Syn.: *Cladonia subcariosa*, fide Elix & McCarthy (1998); Yáñez-Ayabaca: Reports of *C. polycarpoides* Nyl. and *C. subcariosa* (Weber 1986) based on misidentifications of *C. dactylota* (see Yáñez-Ayabaca et al. 2013)
29. *Cladonia rangiferina* (L.) Weber ex F.H. Wigg.
F. Bungartz: the report in Hooker (1847) is most certainly not the arctic alpine *C. rangiferina*, but a similar species of reindeer lichens, most likely *C. confusa* f. *bicolor*
30. *Cladonia rappii* A. Evans
F. Bungartz: only one minute, fragmentary specimen in CDS (Aptroot 65573 B); originally identified by A. Aptroot as *C. rappii*, annotated by T. Ahti as *C. sp.* (probably undescribed, *C. aff. rappii*); however, the fragmentary minute material is extremely similar to basal squamules of *C. pulverulenta*, which also contains fumarprotocetraric acid, no typical material of *C. rappi* has been reported from Galapagos
31. *Cladonia subulata* (L.) Weber ex F.H. Wigg.
F. Bungartz & A. Yáñez: specimen in COLO (63520): Isabela, Cerro Azul, det. Thomson is misidentification of *Cladonia subradiata* (see Yáñez-Ayabaca et al. 2013)
32. *Cladonia symphoriza* Nyl.
Misidentification of *Cladonia didyma*, fide F. Bungartz annotation, 2006
33. *Coccotrema colobinum* (Tuck.) Messuti
F. Bungartz: The type specimen of *Pertusaria colobina* Tuck., later transferred by Messuti & Vobis (2002) and Messuti (2003) into *Coccotrema colobinum*, was supposedly collected by the Reverend T. Hill during the Hassler Expedition in Galapagos. Messuti & Vobis (2002) suggest that one of four specimens labelled (a) in the packet is material collected in Galapagos. It is, however, highly doubtful that any one of these four specimens was actually collected in the archipelago. Not a single specimen of that species has ever been found since. Instead, Messuti & Vobis (2002) cite two more specimens collected by Imshaugh & Ohlsson (MSC 43340, MSC 44816) from the Chilean cost. It is therefore much more probable that the type material of this species, like so many other specimens collected by Reverend T. Hill during the Hassler Expedition, was actually mislabeled (previously the same was already suggested by Weber (1086) p. 490); A. Fryday

examined the type from FH and observed: "The "Galapagos" collection has *C. coccophorum* (= *Lepolichen coccophorus*) on the same piece of bark, which makes the possibility that it is really from the Galapagos extremely unlikely. It was even annotated "probably from Str. of Magellan" by Rolf Santesson in 1955.".

34. *Collemopsidium subarenisedum* (G. Salisb.) Coppins & Aptroot
F. Bungartz & R. Miranda: erroneous record based on misidentification of a single specimen (Aptroot 63087 C).
35. *Cryptothecia assimilis* Makhija & Patw.
F. Bungartz: previous records refer to misidentifications of *Cryptothecia darwiniana* Bungartz & Elix
36. *Cryptothecia candida* (Kremp.) R. Sant.
F. Bungartz: all specimens previously identified as *C. candida* are UV+ bright orange and thus belong to *C. darwiniana* Bungartz & Elix
37. *Cryptothecia evergladensis* Seavey
F. Bungartz: previous records based on misidentifications of *Cryptothecia galapagoana* Bungartz & Elix
38. *Cryptothecia punctosorediata* Sparrius
F. Bungartz: all material belongs to *C. striata*
39. *Dictyonema moorei* (Nyl.) A. Henss.
previous reports are based on misidentifications of material that belongs to *D. membranaceum* s.l.
40. *Diploschistes aeneus* (Müll. Arg.) Lumbsch
F. Bungartz: the material previously identified as *D. aeneus* contains only gyrophoric acid and is therefore *D. badius*
41. *Diploschistes hypoleucus* Zahlbr.
F. Bungartz: the material previously identified as *D. hypoleucus* contains lcanoric acid and belongs to *Diploschistes rampoddensis* (Nyl.) Zahlbr.
42. *Diploschistes scruposus* (Schreb.) Norman
Misidentification of *Diploschistes cinereocaesius*, fide F. Bungartz annotations, 2009
43. *Dirina catalinariae f. catalinariae* Hasse
F. Bungartz: according to Tehler et al. (2013) *D. catalinariae* does not occur in Galapagos and the specimens refer to the newly described *Dirina pacifica* Tehler & Ertz; according to Aptroot & Sparrius (2008) specimens reported previously to *Roccellina badia* Tehler belong to *D. catalinariae* Hasse and thus also refer to *D. pacifica*.
44. *Dirina catalinariae f. sorediata* Tehler
F. Bungartz: according to Tehler et al. (2013) *D. catalinariae* does not occur in Galapagos and the specimens refer to the newly described *Dirina pacifica* Tehler & Ertz; according to Aptroot & Sparrius (2008) specimens reported previously to *Roccellina badia* Tehler belong to *D. catalinariae* Hasse and thus also refer to *D. pacifica*.
45. *Dirina herrei* Zahlbr.
Acc. name: *Dirina approximata* Zahlbr. fide Tehler et al. (2013); F. Bungartz: many specimens in COLO are misidentification of *Syncesia psaroleuca*, *S. graphica* and *Dirina approximata* fide F. Bungartz annotations,

2008; Type: Floreana, Post Office Bay, on smooth bark, Herre A.W.C.T. s.n. [LD - lectotype selected by Tehler et al. (2013), UPS - isolectotype and as Zahlbr. Lich. rar. Exs. n 269 in B, W] material not seen by F. Bungartz (no specimens in COLO?)

46. *Dirinaria aspera* (H. Magn.) D.D. Awasthi

Syn.: *Physcia aspera*, fide Elix & McCarthy (1998); Misidentification of *Dirinaria applanata*, fide F. Bungartz annotation, 2008

47. *Everniastrum subplanum* Sipman

F. Bungartz: the only specimen in CDS was collected by Aptroot 63652 on the Teleferico, Quito, mainland Ecuador.

48. *Flavoparmelia baltimorensis* (Gyeln. & Fóriss) Hale

F. Bungartz: misidentifications of *F. leucoxantha*; specimens do not have coarsely pustulate soredia, instead the soredia are mealy.

49. *Flavoparmelia caperata* (L.) Hale

F. Bungartz: misidentifications of *F. leucoxantha*.

50. *Graphina virginea* (Eschw.) Müll. Arg.

Misidentification of *Diorygma poitaei*, fide Bungartz et al. (2009)

51. *Graphis chrysocarpa* (Raddi) Spreng.

Syn.: *Phaeographina chrysocarpa*, fide A. Aptroot (pers. comm.); Misidentification of *Graphis subchrysocarpa*, fide Bungartz et al. (2009); specimens in COLO: Weber (L- 40405, L- 43952)

52. *Graphis desquamescens* Fée

Misidentification of *Graphis adpressa* or *G. anfractuosa*, fide Bungartz et al. (2009)

53. *Graphis striatula* (Ach.) Spreng.

Misidentification of *Graphis rimulosa*, fide Bungartz et al. (2009) and *Opegrapha graphidiza* s.l., fide F. Bungartz annotation, 2008

54. *Graphis subamylacea* Zahlbr.

Misidentification of *Graphis intricata*, fide Bungartz et al. (2009)

55. *Haematomma puniceum* (Sw. ex Ach.) A. Massal.

Syn.: *Lecanora punicea*; Misidentification of *Haematomma persoonii*, fide F. Bungartz annotations, 2008

56. *Herpothallon philippinum* (Vain.) Aptroot & Lücking

F. Bungartz: Galapagos specimens previously reported as *Herpothallon philippinum* (i.e., the *Herpothallon* species cited by Aptroot et al. 2009 to have a partially I+ blue medulla) lack pseudisidia and have ascospores confined to ascigerous cushions; this material thus belongs to *Cryptothecia striata*.

57. *Heterodermia lepidota* Swinscow & Krog

Misidentification of *Heterodermia antillarum*, fide F. Bungartz annot. 2006

58. *Hypotrachyna laevigata* (Sm.) Hale

Specimens reported in previous versions of this online checklist as *H. laevigata* were examined with TLC by A. Yáñez; all Galapagos material contains alectoronic and alpha-colatolic acid and thus belongs to *H. explendens*.

59. *Julella fallaciosa* (Stizenb. ex Arnold) R.C. Harris
60. *Julella lactea* (A. Massal.) M.E. Barr
61. *Lecanographa illecebrosula* (Müll. Arg.) Egea & Torrente
Misidentification of Lecanographa microcarpella, fide Bungartz annotations, 2008
62. *Lecanora conizaea* (Ach.) Nyl.
in Lichenes Exsiccati, Colorado, No. 138, Weber (1981): Incorrect, but no alternative identification available;
name also used in specimen FH 197184 by W.A. Weber, misidentification of Lecanora floridula, fide F.
Bungartz annotation, 2009
63. *Lecanora glaucovirens* Tuck.
Type: Hassler Expedition (FH), coll. Rev. T. Hill; Weber (1986: 489) doubts that this specimen was actually
collected in Galapagos; F. Bungartz: the type specimen (FH-TUCK 197145) does not belong to Lecanora but
to Lecidea s.l. and no similar material was ever collected again in the Galapagos.
64. *Lecanora helva* Stizenb.
Misidentifications based on a confusion with the very similar *L. leprosa*, fide Bungartz annotations 2008;
Guderley (1999) erroneously states that *L. helva* grows in association with the endemic *L. schindleri*, but all
specimens he annotated are identified as *L. leprosa*, not *L. helva*.
65. *Lecanora orosthea* (Ach.) Ach.
F. Bungartz: misidentifications of sterile *L. substrobilina*
66. *Lecanora subcoarctata* (C. Knight) Hertel
F. Bungartz: Identification of the material by Hertel (1989) is incorrect, the record is based on a specimen of
L. avium.
67. *Lecanora thysanophora* R.C. Harris
F. Bungartz: misidentifications of *Herpothallon granulare*
68. *Lecidea flavoareolata* nom. nud.
No author in Index Fungorum; Zahlbrückner's Cat. Lich. Univ. 7: 362; rejected by Weber (1966)
69. *Leprocaulon microscopicum* (Vill.) Gams
Misidentification of *Leprocaulon tenellum*, fide F. Bungartz annot. 2009
70. *Leptogium isidiosellum* (Riddle) Sierk
Misidentification, all Galapagos material belongs to *L. millegratum*, the lobes are fusing and the isidia are
coarsely granular, fide Bungartz (2008)
71. *Leucodection cf. desquamescens* (Vain.) Lücking
F. Bungartz: the only specimen upon which this preliminary identification was based reacts K+ yellow to red;
L. desquamescens does not contain secondary metabolites according to Rivaz-Plata et al. (2010).
72. *Lithothelium obtectum* (Müll. Arg.) Aptroot
F. Bungartz & R. Miranda: misidentifications of *L. microsporum*; the "unpublished" Galapagos specimen
cited by Aptroot (2006) is *L. microsporum*

73. *Lobaria dissecta* (Sw.) Raeusch.
Misidentification of *Lobaria patinifera*, fide F. Bungartz annotations, 2008
74. *Megalospora tuberculosa* (Fée) Sipman
reported in previous versions of this checklist, but later described as the new species: *Megalospora galapagoensis* Bungartz, Ziemmeck & Lücking
75. *Mycoporum pycnocarpoides* Müll. Arg.
Misidentification of *M. compositum*, fide F. Bungartz annotations, 2011
76. *Niebla* sp. nov. ined.
Described by Aptroot & Bungartz (2007) as *Ramalina fragilis*. The specimen mentioned by Weber (1986: 474) Sipman 63573 was not examined.
77. *Opegrapha graphidiza* s.l. Nyl.
Acc. name: *Opegrapha cactacearum* s.str. fide D. Ertz 2010
78. *Parmeliella mariana* (Fr.) P.M. Jørg. & D.J. Galloway
F. Bungartz: cited by Weber (1986) as *Pannaria pannosa*, a synonym of *P. mariana* according to Elix & McCarthy (1998) and Jørgensen & Galloway (1992, p. 275); the only known specimen (COLO (L-63501, Sipman L-26, Isabela, 21-Jun-1976) is, however, densely isidiate and thus refers to *Parmeliella stylophora* (Vain.) P.M. Jørg.
79. *Parmotrema hypotropum* (Nyl.) Hale
A. Spielmann: reports are based on misidentifications of *Parmotrema grayanum* (Hue) Hale.
80. *Parmotrema neotropicum* Kurok.
A. Spielmann: previous records erroneous
81. *Parmotrema peralbidum* (Hale) Hale
Misidentification of *Canoparmelia crozalsiana*, fide F. Bungartz annotations, 2007; Syn.: *Parmelia peralbida*, fide Elix & McCarthy (1998)
82. *Peltigera didactyla* (With.) J.R. Laundon
Syn.: *Peltigera erumpens*, fide Elix & McCarthy (1998); originally reported in Weber (1986) from Isabela (Sipman L-52) and Santa Cruz (L-40302, L-41144), the Santa Cruz specimens are not tomentose and thus misidentifications of *P. ulcerata* Müll. Arg.
83. *Phaeophyscia hispidula* (Ach.) Moberg
Misidentification of *Hyperphyscia adglutinata*, fide A. Aptroot (pers. comm.)
84. *Phaeophyscia rubropulchra* (Degel.) Moberg
F. Bungartz: previously erroneously identified, all specimens at CDS are *Hyperphyscia pandani* (H. Magn.) Moberg
85. *Phyllopsora corallina* (Eschw.) Müll. Arg.
Specimen in COLO: misidentification of *Phyllopsora intermediella*
86. *Phyllopsora furfuracea* Zahlbr.
Misidentifications of *P. intermediella*, fide Timdal 2010

87. *Physcia aipolia* (A. Massal.) Zahlbr.

F. Bungartz: In Weber (1986) and Elix & McCarthy (1998) as *Physcia mexicana*, but this taxon is not included in Moberg (1990). The lichen genus *Physcia* in Central and South America. The Galapagos material has a medulla K+ yellow and therefore does not belong to *P. biziana* as which it previously has been misidentified. Intermittently the specimens were thus referred to *P. aipolia*. However, unlike *P. aipolia*, Galapagos specimens are much finer lobed and do not contain zeorin despite their K+ yellow medulla. The material is thus chemically distinct from *P. aipolia* s.str. and is therefore now treated as *P. aipolia*, unless studies future of the type suggest otherwise.

88. *Physcia biziana* (A. Massal.) Zahlbr.

Acc. Name: *Physcia aipolia*, in Weber (1986) and Elix & McCarthy (1998) as *Physcia mexicana*, fide A. Aptroot (pers. comm.); Galapagos material medulla K+ yellow!!!, F. Bungartz

89. *Physcia convexella* Moberg

F. Bungartz: *P. convexella* was originally published based on a specimen from Cuzco, Peru at 2800 m altitude and it is very unlikely that Galapagos specimens that occur in the dry and coastal zone belong to this species. The only CDS specimen identified by Moberg (UPS) as *P. convexella* (Aptroot 64468) has a K+ yellow medulla and thus does not agree with the description in Moberg (1990), which states that *P. convexella* is K-. The specimen is treated here as *P. mexicana*.

90. *Physcia lacinulata* Müll. Arg.

All specimens have a black lower side and thus belong to *P. lobulata*, fide F. Bungartz

91. *Placopsis cribellans* (Nyl.) Räsänen

Recorded by Lamb (1947) from the Hassler Expedition. Weber (1966: 196) was unable to find the specimen and agrees with P. James, that the material was not collected in Galapagos, since it is on quartzite.

92. *Placynthiella icmalea* (Ach.) Coppins & P. James

F. Bungartz: specimens have been annotated by M. Schultz as *Lecidopyrenopsis corticola* Vain.

93. *Polymeridium sulphurescens* (Müll. Arg.) R.C. Harris

F. Bungartz & R. Miranda: erroneous identification of *Pseudopyrenula diluta*.

94. *Porina chlorotica* (Ach.) Müll. Arg.

F. Bungartz & R. Miranda: previous reports based on misidentifications of *Strigula phaea*.

95. *Porina subinterstes* (Nyl.) Müll. Arg.

F. Bungartz & R. Miranda: The specimen (Bungartz 8758) originally annotated by R. Miranda as *P. subinterstes* has wart-shaped to hemispherical perithecia and thus belongs to *P. melanops*.

96. *Porina tetracerae* (Ach.) Müll. Arg.

In Elix & McCarthy (1998) erroneously cited from Galapagos (Weber 1993: 433), but the citation in Weber (1993) is from the Cocos Islands! Reports of the species in previous versions of this checklist were cited as "rejected"; one single specimen in CDS (Aptroot 64623) annotated by R. Lücking as *P. tetracerae* is extremely poorly developed and lacks perithecia; the specimen is treated here as *P. distans* (= *P. cf. conspersa*).

97. *Pyrenula acutalis* R.C. Harris

F. Bungartz & R. Miranda: all material previously identified as *P. acutalis* does not belong to that species.

98. *Pyrenula aff. septicollaris* (Eschw.) R.C. Harris
F. Bungartz & R. Miranda: misidentifications.
99. *Pyrenula cocoës* Müll. Arg.
F. Bungartz & R. Miranda: all previous reports based on misidentifications.
100. *Pyrenula cruentata* (Müll. Arg.) R.C. Harris
F. Bungartz & R. Miranda: most specimens belong to *P. cruenta*, one specimen is *P. breutelii*
101. *Pyrenula globifera* (Eschw.) Aptroot
F. Bungartz & R. Miranda: previous reports based on misidentifications.
102. *Pyrenula microcarpa* Müll. Arg.
F. Bungartz & R. Miranda: all specimens in CDS misidentifications; specimen in COLO not examined: Santa Cruz, on *Cordia lutea*, Darwin Station, Weber (L-40579), det. Aptroot, 1991, as *P. cinerea* (syn. of *P. microcarpa*)
103. *Pyrenula pyrenuloides* (Mont.) R.C. Harris
COLO 192496 was identified by A. Aptroot as *P. pyrenuloides*, but is fide R. Miranda annot. 2010 refers to *P. thelomorpha*
104. *Pyrrhospora russula* (Ach.) Hafellner
F. Bungartz: All Galapagos material grows on rock, has a creamish white to yellowish thallus that conforms well with *P. sanguinolenta*.
105. *Pyrrhospora sanguinolenta* (Kremp.) Rambold & Hafellner
F. Bungartz: all previous records based on misidentifications of *Ramboldia heterocarpa* (Fée) Kalb, Lumbsch & Elix
106. *Pyxine caesiopruinosa* (Tuck.) Imshaug
Acc. name: *Pyxine subcinerea*, fide Elix & McCarthy (1998)
107. *Ramalina australiensis* Nyl.
Misidentification of *Ramalina sorediosa*, fide A. Aptroot (pers. comm.); Syn.: *Ramalina furcellata*, *Ramalina farinacea* auct. non Ach., *Ramalina dasypoga* auct. non Tuck.
108. *Ramalina denticulata* Nyl.
Misidentification of *Ramalina aspera*, fide A. Aptroot (pers. comm.)
109. *Ramalina farinacea* (L.) Ach.
Acc. name: *Ramalina australiensis*, fide Elix & McCarthy (1998) & misidentification of *Ramalina sorediosa*, fide Aptroot (pers. comm.)
110. *Ramalina furcellata* (Mont.) Zahlbr.
Acc. name: *Ramalina australiensis*, fide Elix & McCarthy (1998) & misidentification of *Ramalina sorediosa*, fide Aptroot & Bungartz (2007)
111. *Ramalina linearis* R.H. Petersen & M. Zang
Misidentification of *Ramalina puiggarii*, fide Aptroot & Bungartz (2007) and of *R. complanata* and *R. sideriza* (FH specimens), fide F. Bungartz annotations, 2009

112. *Roccella lirellina* (Darb.) M. Choisy

Misidentification of *Roccella margaritifera* or *R. nigerrima* fide Tehler et al. (2009); initially presumed to occur in Galapagos (Tehler 2007), but later shown to be restricted to coastal Peru (Tehler et al. 2009). Syn: *Reinkella lirellina* Darb. fide Tehler et al. (2009)

113. *Roccella portentosa* (Mont.) Darb.

Misidentification of various species in the *Roccella galapagoensis* agg., fide Tehler et al. (2009)

114. *Roccellina nigrocincta* Tehler

Wrong reference therefore rejected by Aptroot & Sparrius (2008): Elix & McCarthy (1998: 253) referring to Tehler (1983: 61) who does not mention the species

115. *Sticta filix* (Sw.) Nyl.

Misidentifications of *S. dichotoma* s.l.

116. *Sticta quercizans* Ach.

Misidentification for *Sticta weigelii*, fide Weber (1966); Acc. name: *Sticta weigelii*, fide Elix & McCarthy (1998); Specimens in CAS: Isabela, Iguana Cove, Snodgrass & Heller; Floreana, A. Stewart No. 400; Santa Cruz, NW-side, A. Stewart No. 401

117. *Sticta sylvatica* (Huds.) Ach.

Misidentification of *Sticta weigelii* and *Sticta fuliginosa*, fide F. Bungartz

118. *Syncesia effusa* (Fée) Tehler

Syn.: *Chiodecton effusum*; F. Bungartz: all Galapagos specimens are misidentifications of *Cryptothecia assimilis*

119. *Syncesia flavescens* (Nyl.) Tehler

F. Bungartz: all Galapagos specimens are misidentifications of *Cryptothecia assimilis*

120. *Teloschistes exilis* (Michx.) Vain.

Probably based on misidentifications of *Teloschistes flavicans*, fide Bungartz, 2009

121. *Tephromela atra* (Huds.) Hafellner

Syn.: *Lecanora atra*, fide Elix & McCarthy (1998); F. Bungartz & K. Kalb: all records previously identified as this species are misidentifications of *Tephromela rizophorae*

122. *Trypethelium refertum* Stirt.

F. Bungartz: misidentification of *Anisomeridium tamarindi*

123. *Trypethelium tuberculatum* (Vain.) R.C. Harris

Misidentifications of *T. aff. marcidum* fide annotations F. Bungartz & R. Miranda, 2011

124. *Usnea amabilis* Motyka

Misidentification; Weber (1986: 493) states that *U. longissima* reported by Stewart (1912) is *U. amabilis*, both IDs are misidentifications of *Usnea mexicana*, fide P. Clerc, M. de los Angeles, C. Truong annotations, 2008

125. *Usnea antillarum* (Vain.) Zahlbr.

Misidentification of various different *Usnea* spp., fide Clerc, P. annotations 2008

126. *Usnea arthroclada* Fée

Probably misidentification of *Usnea mexicana*, fide F. Bungartz

127. *Usnea ceratina* Ach.

Misidentification of *Usnea rubicunda*, fide Elix & McCarthy (1998)

128. *Usnea cladocarpa* Fée

Misidentification of *Usnea* spp., fide P. Clerc, M. de los Angeles & C. Truong annotations, 2008

129. *Usnea dasypoga* (Ach.) Röhl.

Misidentification of *Usnea* cf. *bornmuelleri* and *U. baileyi*, fide P. Clerc & C. Truong annotations, 2008

130. *Usnea longissima* Ach.

Misidentification for *U. mexicana*, fide P. Clerc, C. Truong & M. de los Angeles Herrera Campos (pers. comm, 2008)

131. *Usnea mirabilis* Motyka

Probably misidentifications of *Usnea rubicunda* fide C. Truong

132. *Usnea paradoxa* (Zahlbr.) Motyka

Misidentification of *Usnea angulata*, fide P. Clerc & C. Truong annotations, 2008 & F. Bungartz, 2009 (FH 197301)

133. *Usnea plicata* (L.) Weber ex F.H. Wigg.

Syn.: *Usnea dasypoga* var. *plicata*; According to Weber (1986: 493) "reported on the basis of scrappy specimens collected by J.H. Andersson and Charles Darwin" which do not resemble *U. plicata*.

134. *Usnea rubiginea* (Michaux) A. Massal.

C. Truong: Galapagos specimens under the name *U. rubiginea* (Weber 1987) belonged to *U. erinacea*, *U. rubicunda* or *U. poliotrix*; this taxon cited by Weber is listed in Elix & McCarthy (1998) as *Usnea rubicunda*.

135. *Verrucaria cf. pinguicula* A. Massal.

F. Bungartz: records based on misidentifications; all specimens identified by Breuss as *Verrucaria* sp. According to Orange et al. (2009, in Smith et al. Lichens of Great Britain and Ireland p. 954) *Verrucaria pinguicula* grows on hard limestone, a substrate absent from Galapagos.

136. *Verrucaria inficiens* Breuss

F. Bungartz: material initially identified by O. Breuss, later revised to *V. lecideoides*

137. *Verrucaria polysticta* Borrer

F. Bungartz: According to Breuss all with hymenial algae and thus referred to *Staurothele* sp.

138. *Xanthoparmelia microspora* (Müll. Arg.) Hale

F. Bungartz & A. Spielmann: Nash, Gries & Elix (1995) cite a specimen of *Xanthoparmelia microspora* (COLO Weber L-62892) that was originally, in 1976, also identified by Hale as *X. microspora*. However, Hale in 1980 revised his identification to *Pseudoparmelia leucoxantha*, which was recombined into *Flavoparmelia leucoxantha*. The specimen does contain protocetraric acid, not salazinic acid and thus does not belong to *X. microspora*!

139. *Xanthoria candelaria* (L.) Th. Fr.

Misidentification of *Xanthomendoza weberi*, fide F. Bungartz annotations, 2006

140. *Xanthoria cf. tenuiloba* L. Lindblom

Misidentification of *Xanthomendoza weberi*, fide F. Bungartz annotations, 2006

Acknowledgements

We are grateful for the financial report received for this project. Please refer to the website (www.darwinfoundation.org/datazone/checklist/) for a detailed list of all our donors.

This checklist would not be possible without adjunct and collaborating scientists, and volunteers of the Charles Darwin Foundation. The following scientists and volunteers have contributed to the CDF Checklist of Galapagos Lichenized Fungi: Teuvo Ahti, Rosa Batalles, Michel Benatti, Lenyn Betancourt C., Othmar Breuss, Philippe Clerc, Carolina Cornejo, Desiree Cruz, Manuela Dal Forno, Paul Diederich, Valeria Leonor Dután-Patiño, Damien Ertz, Tassilo Feuerer, Anne Guézou, Marusa Herrera, Georg Hillmann, Patricia Jaramillo Díaz, Harald Jonitz, Klaus Kalb, Kerry Knudsen, James Lawrey, Robert Lücking, Danilo Minga, Ricardo Miranda, Roland Moberg, Bibiana Moncado, Catalina Quintana, Eimy Rivaz-Plata, Matthias Schultz, Harrie Sipman, Adriano Spielman, Ulrik Søchting, Anders Tehler, Göran Thor, Leif Tibell, Einar Timdal, Camille Truong, Diego Villagómez, William A. Weber, Martin Westberg.

References

1. Ahti, T. (2000) *Cladoniaceae*. Flora Neotropica Monograph 78, 362 pp., The New York Botanical Garden, Bronx, New York.
2. Andersson, N. J. (1855) *Om Galápagos Öarnes Vegetation*. Kongliga Svenska Vetenskaps Akademien Handl., Stockholm: 61-256.
3. Aptroot, A. (1991) *A monograph of the Pyrenulaceae (excluding Anthracothecium and Pyrenula) and the Requienellaceae, with notes on the Pleomassarinaceae, the Trypetheliaceae and Mycomicrothelia (lichenized and non-lichenized ascomycetes)* Bibliotheca Lichenologica 44: 178 pp.
4. Aptroot, A. (2006) *Three new species of Lithothelium (Pyrenulaceae) from China and Thailand, with a revised world key and annotated list of species*. The Lichenologist 38(6): 541–548.
5. Aptroot, A., Sparrius, L.B., LaGreca, S. & Bungartz, F. (2007) *Angiactis, a new crustose lichen genus in the family Roccellaceae with species from Bermuda, the Galápagos Islands, and Australia*. The Bryologist 111(3): 510-516.
6. Aptroot, A., Bungartz, F. (2007) *The Lichen Genus Ramalina on the Galapagos*. The Lichenologist 39: 519-542.
7. Aptroot, A., Sparrius, L.B. (2008) *Crustose Roccellaceae in the Galapagos Islands, with the new species Schismatomma spierii*. The Bryologist 111(4): 659-666.
8. Aptroot, A., G. Thor, R. Lücking, J.A. Elix & J.L. Chaves (2009) *The lichen genus Herpothallon reinstated*. In: Biodiversity and ecology of lichens - Liber Amicorum Harrie Sipman. Bibliotheca Lichenologica 99: 19-66.
9. Aptroot, A. (2013) *A world key to the species of Anthracothecium and Pyrenula* The Lichenologist 44(1): 5-53
10. Awasthi, D.D. (1975) *A monograph of the lichen genus Dirinaria*. Bibliotheca Lichenologica 2: 1-109.

11. Brodo, I. M., Sharnoff, S. D., Sharnoff, S. (2001) *Lichens of North America* Yale University Press, New Haven.
12. Bungartz, F., Dután-Patiño, V.L., Elix, J.A. () *The lichen genera Cryptothecia, Herpothallon and Helminthocarpon (Arthoniales) in the Galapagos Islands, Ecuador* The Lichenologist 45(6): 1–24 (2013)
13. Bungartz, F. (2008) *Cyanolichens of the Galapagos Islands - The genera Collema and Leptogium*. Sauteria 15: 139-158.
14. Bungartz, F., Lücking, R. & Aptroot, A. (2009) *The lichen family Graphidaceae (Ostropales, Lecanoromycetes) in the Galapagos Islands*. Nova Hedwigia 90: 1-44.
15. Clerc, P. (2008) In: T. Nash III, C. Gries & F. Bungartz (eds.), *Lichen Flora of the Greater Sonoran Desert Region*, vol. 3. Lichen Unlimited Arizona State University, Tempe, pp. 302–335.
16. Clerc, P. (2011) *Notes on the genus Usnea Adans. (lichenized Ascomycota) III*. Bibliotheca Lichenologica 106: 41–51.
17. Darbshire, O.V. (1935) *The Templeton Crocker Expedition of the California Academy of Sciences, 1932. No. 23: The Roccellaceae with Notes on Specimens Collected During The Expedition of 1905-06 to the Galapagos Islands*. Proc. Calif. Acad. Scs., 4th Series, 21(23): 285-294.
18. Diederich, P. (1995) *New or interesting lichenicolous fungi VI. Arthonia follmanniana, a new species from the Galapagos Islands*. In: F.J.A. Daniels, M.Schultz & J. Peine (eds.): Flechten Follmann. Contributions to Lichenology in Honour of Gerhard Follmann, pp. 179-182. Botanical Institute, University of Cologne.
19. Dodge, C.W. (1935) *Lichenes. In: H.K. Svenson: Plants of the Astor Expedition, 1930 (Galápagos and Cocos Islands)*. Amer. J. Bot. 22 (2): 221.
20. Dodge, C.W. (1936) *Lichens of the G. Allan Hancock Expedition of 1934, collected by Wm. R. Taylor*. Hancock Pacific Expeditions 3(3): 33-46.
21. Elix, J.A., McCarthy, P.M. (1998) *Catalogue of the Lichens of the Smaller Pacific Islands*. Bibliotheca Lichenologica 70, J. Cramer, Berlin, Stuttgart, 361pp.
22. Ertz, D. (2009) *Revision of the corticolous Opegrapha species from the Paleotropics*. Bibliotheca Lichenologica 102: 176 pp.
23. Ertz, D., Tehler, A. (2010) *The phylogeny of Arthoniales (Pezizomycotina) inferred from nucLSU and RPB2 sequences* Fungal Diversity DOI 10.1007/s13225-010-0080-y
24. Ertz, D., Bungartz, F., Diederich, P. & Tibell, L. (2011) *Molecular and morphological data place Blarneya in Tylophoron (Arthoniaceae)*. The Lichenologist 43(4): 1–12.
25. Farlow, W.G. (1902) *Lichens*. In: B.L. Robinson: Flora of the Galapagos Islands. Proc. Americ. Acad. Arts Sc. 38(4): 83-89, N.S. 14 (Papers from the Hopkins-Stanford Expedition to the Galapagos Islands)
26. Frisch, A. (2007) *Compsocladium kalbii sp. nova from the Ecuadorian Andes*. Bibliotheca Lichenologica 96: 73–86.

27. Galloway, D.J. (1985) *Nomenclatural notes on Pseudocyphellaria II: Some southern hemisphere taxa*. Lichenologist 17(3): 303-307.
28. Galloway, D.J., Arvidsson, L. (1990) *Studies in Pseudocyphellaria (Lichens) II. Ecuadorean species*. Lichenologist 22(2): 103-135.
29. Grube, M. (2007) *Arthonia*. In: Nash III, T.H., Gries, C. & Bungartz, F.: Lichen Flora of the Greater Sonoran Desert Region. Vol. 3, Lichens Unlimited, Tempe, p. 39-61.
30. Guderley, R. (1999) *Die Lecanora subfuscata-Gruppe in Süd- und Mittelamerica*. J. Hattori Bot. Lab. 87: 131-257.
31. Harris, R.C. (1995) *More Florida Lichens, including the 10 cent tour of the pyrenolichens, a twenty year perspective*. New York Botanical Garden, Bronx (published by the author), 182 pp.
32. Herrera-Campos, M., Clerc, P., Nash III, T.H. (1998) *Pendulous species of Usnea from the temperate forests in Mexico* The Bryologist 101: 303-329.
33. Herrera-Campos, M., T.H. Nash III, Garcia, A. (2001) *Preliminary study of the Usnea fragilescens aggregate in Mexico*. The Bryologist 104: 235-259.
34. Hooker, J.D. (1847) *An Enumeration of the Plants of the Galapagos Archipelago; with descriptions of those which are new*. Transactions of the Linnean Society of London 20: 163-233.
35. Jagadeesh Ram, T.A.M., Aptroot,A., Sinha, G.P. & Singh, K.P. (2007) *A new isidiate megalaria species and new records of lichenized, lichenicolous and non-lichenized ascomycetes from India* Nova Hedwigia 85(1-2): 139-144
36. Jørgensen, P.M. (1998) *Acantholichen pannariooides, a new basidiolichen from South America*. The Bryologist 101(3): 444-447.
37. Kalb, K. (2001) *New or otherwise interesting lichens. I*. In P.M. McCarthy, G. Kantvilas, S.H.J.J. Louwhoff (eds.): Lichenological Contributions in Honour of Jack Elix. Bibliotheca Lichenologica 78, J. Cramer, Berlin, Stuttgart, p. 141-167.
38. Kalb, K. (2008) *New or otherwise interesting lichens. IV. Neue oder anderweitig interessante Flechten. IV*. Sauteria 15: 239-248
39. Kashiwadani, H., Kalb, K. (1993) *The genus Ramalina in Brazil* Lichenologist 25: 1-31.
40. Kirk, P. (ed.) (2010) *Index Fungorum*. CABI Bioscience, CBS and Landcare Research, available online at www.indexfungorum.org.
41. Knudsen, K. (2012) *Some Notes on Acarosporaceae in South America* Opuscula Philolichenum, 11: 31-35.
42. Lamb, I.M. (1977) *A conspectus of the lichen genus Stereocaulon (Schreb.) Hoffm.* Journal of the Hattori Botanical Laboratory 43: 191-355.
43. Landrón C.I. (1972) *The lichen genus Ramalina Ach. in the West Indies with notes on its role in the vegetation of Porto Rico*. PhD Thesis, Michigan State University.

44. Lawrey, J.D., Lücking, R., Sipman, H.J.M., Chaves, J.L., Redhead, S.A., Bungartz, F., Sikaroodi, M. & Gillevet, P.M. (2009) *High concentration of basidiolichens in a single family of agaricoid mushrooms (Basidiomycota: Agaricales: Hygrophoraceae)*. Mycological Research 113: 114-1171.
45. Lendemer, J.C., Hodkinson, B.P. (2013) *A radical shift in the taxonomy of Lepraria s.l.: Molecular and morphological studies shed new light on the evolution of asexuality and lichen growth form diversification* Mycologia doi:10.3852/12-338
46. Linder, D.H. (1934) *The Templeton Crocker Expedition of the California Academy of Sciences, 1932. No. 18: Lichens*. Proc. Calif. Acad. Scs., 4th Series, 21(18): 211-124.
47. Lumbsch, H.T. (1987) *Eine neue Subspecies in der Flechtengattung Diploschistes aus der Südhemisphäre*. Herzogia 7: 601-608.
48. Lumbsch, H.T., Ahti, T., Altermann, S., Amo de Paz, G., Aptroot, A., Arup, U., Bárcenas Peña, A., Bawingan, P.A., Benatti, M.N., Betancourt, L., Björk, C.R., Boonpragob, K., Brand, M., Bungartz, F., Cáceres, M.E.S., Candan, M., Chaves, J.L., Clerc, P., Common, R., Coppins, B.J., Crespo, A., Dal Forno, M., Divakar, P.K., Duya, M.V., Elix, J.A., Elvebakk, A.V., Fankhauser, J., Farkas, E., Ferraro, I.L., Fischer, E., Galloway, D.J., Gaya, E., Giralt, M., Goward, T., Grube, M., Hafellner, J., Hernández M., J.E., De los Ángeles Herrera Campos, M., Kalb, K., Kärnefelt, I., Kantvilas, G., Killmann, D., Kirika, P., Knudsen, K., Komposch, H., Kondratyuk, S., Lawrey, J.D., Mangold, A., Marcelli, M.P., McCune, B., Messuti, M.I., Michlig, A., Miranda González, R., Moncada, B., Naikatini, A., Nelsen, M.P., Øvstedal, D.O., Palice, Z., Papong, K., Parnmen, S., Pérez-Ortega, S., Printzen, C., Rico, V.J., Robayo, J., Rosabal, D., Ruprecht, U., Salazar Allen, N., Sancho, L., Santos de Jesus, L., Santos Vieira, T., Schultz, M., Seaward, M.D.R., Sérusiaux, E., Schmitt, I., Sipman, H.J.M., Sohrabi, M., Søchting, U., Søgaard, M.Z., Sparrius, L.B., Spielmann, A., Spribble, T., Sutjaritturakan, J., Thammathaworn, A., Thor, G., Thüs, H., Timdal, E., Truong, C., Türk, R., Umaña Tenorio, L., Uperti, D.K., Van den Boom, P., Vivas Rebuelta, M., Wedin, M., Will-Wolf, S., Wirth, V., Wirtz, N., Yahr, R., Yeshitela, K., Ziemmeck, F., Lücking, R. (2010) *One hundred new species of lichenized fungi: a signature of undiscovered global diversity*. Phytotaxa 18: 1-127.
49. Lücking, R. (2008) *Foliicolous lichenized fungi*. Flora Neotropica 103: 1-866.
50. Lücking, R., Lawrey, J.D., Sikaroodi, M., Gillevet, P.M., Chaves, J.L., Sipman, H.J.M. & Bungartz, F. (2009) *Do lichens domesticate photobionts like farmers domesticate crops? Evidence from a previously unrecognized lineage of filamentous cyanobacteria*. American Journal of Botany 96(8): 1409-1418.
51. Lücking, R., Tehler, A., Bungartz, F. & Lumbsch, H.T. (2003) *Journey from the West: Did tropical Graphidaceae (lichenized Ascomycota: Ostropales) evolve from a saxicolous ancestor along the American Pacific coast?* American Journal of Botany. DOI 10.3732/ajb.1200548
52. Martin, G.W. (1948) *Additions to Galápagos Fungi* Pacific Science 11(2):71-77
53. Martins, M.F.N. (2007) *O gênero Heterodermia (Physciaceae, Ascomycota liquenizados) no Estado de São Paulo, Brasil* Master Thesis, Universidade Estadual Paulista, São Paulo, 218 pp.
54. McDonald, T., Miadlikowska, J. & Lutzoni, F. (2003) *The lichen genus Sticta in the Great Smoky Mountains: A phylogenetic study of morphological, chemical, and molecular data*. The Bryologist 106(1): 61-79.
55. Moberg, R. (1990) *The lichen genus Physcia in Central and South America*. Nord. J. Bot. 10: 319-342.

56. Motyka, J. (1936–38) *Lichenum Generis Usnea Studium Monographicum Pars Systematica*. Leopoldi (privately printed).
57. Muggia, L., Gueidan, C., Perlmutter, G., Eriksson, O.E. & Grube, M. (2009) *Molecular data confirm the position of Flakea papillata in the Verrucariaceae*. The Bryologist 112(3): 538–543.
58. Nash III, T.H., Ryan, B.D., Gries, C., Bungartz, F. (2002) *Lichen Flora of the Greater Sonoran Desert Region, Volume 1* Lichens Unlimited, Tempe
59. Ohmura, Y. (2001) *Taxonomic study of the genus Usnea (lichenized Ascomycetes) in Japan and Taiwan*. Journal of the Hattori Botanical Laboratory 90: 1–96.
60. Ohmura, Y. (2008) *Molecular phylogeny of Usnea rubicunda and U. rubrotincta (Parmeliaceae, lichenized Ascomycotina) based on ITS rDNA sequences*. Journal of Japanese Botany 83: 347–355.
61. Ohmura, Y., Lin, C.-K. Wang, P.-H. (2010) *Three sorediate species of the genus Usnea (Parmeliaceae, Ascomycota) new to Taiwan*. Memoirs of the National Museum of Natural Science (Tokyo) 46: 69–76.
62. Roth, K., Mathey, A., & Leuckert, C. (1978) *The xanthones of Buellia galapagana Weber (Lichenes, Physciaceae)* Wilidenowia 8 (2): 363–368
63. Santesson, R. (1939) *Amphibious pyrenolichens I*. Arkiv för Botanik 29A(10): 1–67.
64. Schofield, E.K. (1984) *Plants of the Galapagos Islands. Field Guide and Travel Journal*. Universe Books, New York, 159 pp.
65. Schultz, M., Aptroot, A. (2008) *Notes on poorly known, small cyanobacterial lichens from predominantly wet tropical to subtropical regions*. In: Türk, R., John, V. & Hauck, M. (eds.): *Facetten der Flechtenforschung - Festschrift zu Ehren von Volkmar Wirth*. Sauteria 15: 433–458.
66. Stewart, A. (1912) *Expedition of the California Academy of Sciences to the Galapagos Islands 1905–1906. No. 7: Notes on the Lichens of the Galapagos Islands*. Proc. Calif. Acad. Sci., 4th Series, 1(7): 431–446.
67. Svenson, H.K. (1935) *Plants of the Astor Expedition, 1930 (Galapagos and Cocos Islands)*. Am. J. Bot. 22: 208–277.
68. Swinscow, T., Krog, H. (1979) *The fruticose species of Usnea subgenus Usnea in East Africa* Lichenologist 11: 207–252.
69. Tehler, A. (2002) *Roccella* In: T.H. Nash III, B.D. Ryan, C. Gries & F. Bungartz (eds.): *Lichen Flora of the Greater Sonoran Desert Region, vol. 1*. Lichens Unlimited, Arizona State University, Tempe, 451–454 pp.
70. Tehler, A. (2007) *The Roccella lirellina and R. galapagoensis aggregates, taxonomy and nomenclature* In: I. Kärnefelt & A. Thell (eds.): *Lichenological Contributions in Honour of David Galloway*. Bibliotheca Lichenologica 95: 517–530.
71. Tehler, A., Irestedt, M., Bungartz, F. & Wedin, M. (2009) *Evolution and reproduction modes in the Roccella galapagoensis aggregate (Roccellaceae, Arthoniales)* Taxon 58(2): 438–456.
72. Tehler, A., Baloch, E., Tibell, L. & Wedin, M. (2009) *The systematic position of Schistophoron*. Bibliotheca Lichenologica 99: 383–329.

73. Tehler, A., Ertz, D., Irestedt, M. (2013) *The genus Dirina (Roccellaceae, Arthoniales) revisited* The Lichenologist 45(4): 427-476
74. Truong, C., Bungartz, F. & Clerc, P. (2011) *The lichen genus Usnea (Parmeliaceae) in the tropical Andes and the Galapagos: species with a red-orange cortical or subcortical pigmentation.* The Lichenologist, in press.
75. Truong, C., Rodriguez, J.M., Clerc, P. (2013) *Pendulous Usnea species (Parmeliaceae, lichenized Ascomycota) in tropical South America and the Galapagos* The Lichenologist 45(4): 505-543
76. Truong, C., Clerc, P. (2013) *Eumitrioid Usnea species (Parmeliaceae, lichenized Ascomycota) in tropical South America and the Galapagos* The Lichenologist 45(3): 383-395.
77. Vareschi, V. (1973) *Resultados liquenológicos de excusiones efectuadas en Venezuela. No. 3. Catalogo de los líquenes de Venezuela.* Acta Botanica Venezuelica 8: 177–245.
78. Weber, W.A. (1966) *Lichenology and Bryology in the Galápagos Islands, with Check Lists of the Lichens and Bryophytes Thus Far Reported.* In: Bowman, R.I. (ed.): *The Galápagos. Proceedings of the Symposium of the Galápagos International Scientific Project*, University of California Press, Berkeley, p. 197-200.
79. Weber, W.A. (1971) *Four new species of Buellia from western North and South America.* The Bryologist 74: 185-191.
80. Weber, W.A. (1981) *Lichenes Exsiccati, distributed by the University of Colorado Museum, Boulder, Fasc. 1-15, nos. 1-600, 1961-1979.* Mycotaxon 13(1): 85-104.
81. Weber, W.A. (1986) *The Lichen Flora of the Galápagos Islands, Ecuador.* Mycotaxon 27: 451-497.
82. Weber, W.A. (1993) *Additions to the Galápagos and Cocos Islands Lichen and Bryophyte Floras.* The Bryologist 96(3): 431-434.
83. Westberg, M., Arup, U. (2011) *Candelaria pacifica sp. nova (Ascomycota, Candelariales) and the identity of Candelaria vulgaris* Bibliotheca Lichenologica 106, 353-364
84. Yáñez-Ayabaca, A., Dal-Forno, M., Bungartz, F., Lücking, R. (2012) *A first assessment of Galapagos basidiolichens Fungal Diversity* 52:225-244
85. Yáñez-Ayabaca, A., Ahti, T., Bungartz, F. (2013) *The Family Cladoniaceae (Lecanorales) in the Galapagos Islands* Phytotaxa 129 (1): 1–33

Disclaimer

The Charles Darwin Foundation Galapagos Species Checklist is a continuously updated list of all species currently known from the Galapagos Islands and reflects up-to-date knowledge compiled by scientists of the Charles Darwin Foundation (CDF) in collaboration with experts from around the world. CDF shares this data publicly and invites comments, corrections and additions.

Please do not hesitate to contact us; your input is very welcome. However, please understand that additions, changes, and corrections will be posted at periodic intervals after thorough cross-referencing of all new data. As an independent international scientific organization, the Charles Darwin Foundation relies on funding from donors who support our work. Please contact us at datazone@fcdarwin.org.ec if you would like to support the Charles Darwin Foundation Galapagos Species Checklist and help make knowledge of Galapagos biodiversity more widely available.