

**GRANDIDIERELLA HALOPHILUS A NEW SPECIES OF THE FAMILY AORIDAE
(CRUSTACEA: AMPHIPODA) FROM THE SALTPANS OF
THE INNER GULF OF THAILAND**

K. Wongkamhaeng

Excellence Center for Biodiversity of Peninsular Thailand, Faculty of Science, Prince of Songkla University, 90112 Thailand
Prince of Songkla University, 90112 Thailand

P. Pholpunthin

Center of Excellence for Biodiversity of Peninsular Thailand (CBiPT), Department of Biology, Faculty of Science
Prince of Songkla University, 90112 Thailand

B. A. R. Azman

Marine Ecosystem Research Centre (EKOMAR), Faculty of Science and Technology, Universiti Kebangsaan Malaysia
43600 Bangi, Selangor, Malaysia
Email: abrahim@gmail.com

ABSTRACT.—A new species of *Grandidierella* is described from the Inner Gulf of Thailand. This species inhabits high saline waters of abandoned saltpans from the Samut Sakorn district, Thailand. It appears to be closely related to *Grandidierella propodentata* Moore, 1986, also collected at hypersaline environments, in having an accessory flagellum; and arborescent setae on the inner face of the outer lobe of lower lip. The main diagnostic features of *Grandidierella halophilus* new species are a coxal plate 2 with ventral margin posteriorly produced into triangular expansion; uropod 3 peduncle inflated with ramus bearing a small second article; lower lip with arborescent setae on the inner face of the outer lobe; and bifid robust setae on antennae 1–2, gnathopods 1–2, pereopods 5–7, and uropods 1–2. This combination of characters has not been recorded previously in the *Grandidierella*.

KEY WORDS.—*Grandidierella halophilus*, new species, taxonomy, saltpans, Gulf of Thailand

INTRODUCTION

Species of the genus *Grandidierella* Coutière, 1904, are predominant and widespread in brackish, estuarine and coastal waters worldwide. There is extensive literature on the occurrence of *Grandidierella* species from the marine habitats (Myers, 1970, 1981, 1998; Ren, 2006; Bochert & Zettler, 2010; Azman & Othman, 2012), yet there are only two existing records of *Grandidierella* (*G. propodentata* Moore, 1986 and *G. exilis* Myers, 1981) from hypersaline environments. There are many reports of the 40 described species of *Grandidierella* worldwide. However, only three *Grandidierella* species are reported from the waters of Thailand and adjacent archipelagos, namely *G. gilesi* Chilton, 1921, *G. taihuiensis* Morino & Dai, 1990, and *G. gravipes* K. H. Barnard, 1935 (Chilton, 1925; Bussawich, 1985; Angsupanich & Kuwabara, 1995; Angsupanich et al., 2005; Ruensirikul et al., 2007; Ariyama et al., 2010). In the present study we provide a detailed description of both male and female specimens of *G. halophilus* new species collected from abandoned saltpans in the inner Gulf

of Thailand. Interestingly, although *G. halophilus* is fairly abundant in these saltpans, there has been no detailed study on this organism.

MATERIAL AND METHODS

This study is based upon material collected from various abandoned saltpans of the Samut Sakorn Province (Fig. 1), Inner Gulf of Thailand in Feb. 2011. Samples were collected from the host plant (*Ruppia maritima* Linnaeus) and wet sieved through a 0.55-mm mesh sieve. The materials retained on the sieve were then carefully transferred into plastic containers and fixed in 10% buffered formalin. In the laboratory, amphipod specimens were sorted out and stored in 70% alcohol. The animals were then examined under a compound microscope and later selected for dissection. The appendages of the dissected specimens were examined and figures were produced using a Leica DMLB light microscope with a camera lucida. All illustrations were digitally ‘inked’ following Coleman (2003). Figure legend: A, antenna; G,

gnathopod; HD, head; LL, lower lip; MD, mandible; MX, maxilla; MP, maxilliped; P, pereopod; Pl, pleopod; T, telson; U, uropod; UR, urosome; UL, upper lip; R, right; L, left; ♂, male; ♀, female. Type material is deposited at Prince of Songkla University Zoological Collection with the prefix PSUZC for museum numbers and the Universiti Kebangsaan Malaysia Muzium Zoologi with the prefix UKMMZ for museum numbers.

TAXONOMY

Genus *Grandidierella* Coutière, 1904

Diagnosis. — Eyes small to medium. Antenna 1, accessory flagellum minute, 1-segmented. Maxilla 1, inner plate vestigial. Coxae very small, relatively short of various sizes and shapes. Gnathopod 1 (male) complexly subchelate and greatly larger than gnathopod 2. Gnathopod 2 subchelate. Pereopods 6–7, dactylus elongate, falcate. Uropods 1–2 biramous; rami slightly subequal; peduncle with ventrodistal process. Uropod 3 uniramous. Telson entire.

Species composition. — *Grandidierella* contains 41 species: *G. africana* Schellenberg, 1936; *G. bispinosa* Schellenberg, 1938; *G. bonnieroides* Stephensen, 1948; *G. cabindae* (Schellenberg, 1925); *G. chelata* K. H. Barnard, 1951; *G. chaohuensis* Hou & Li, 2002; *G. dentimera* Myers, 1970; *G. elongata* (Chevreux, 1926); *G. exilis* Myers, 1981; *G. fasciata* Ariyama, 1996; *G. gilesi* Chilton, 1921; *G. gravipes* K. H. Barnard, 1935; *G. grossimana* Ledoyer, 1967; ***G. halophilus* new species, 2011**; *G. indentata* Ledoyer, 1979; *G. insulae* Myers, 1981; *G. ischienoplia* Bochert & Zettler, 2010; *G. japonica* Stephensen, 1938; *G. kanakensis* Myers, 1998; *G. koa* J. L. Barnard, 1977; *G. lignorum* K. H. Barnard, 1935; *G. longidactyla* Ledoyer, 1982; *G. lutosa* K.H. Barnard, 1952; *G. macronyx* K. H. Barnard, 1935; *G. mahafalensis* Coutière,

1904 (**Type species**); *G. makena* J. L. Barnard, 1970; *G. malaccaensis* Azman & Othman, 2012; *G. nottoni* Shoemaker, 1935; *G. nyala* Griffiths, 1974; *G. osakaensis* Ariyama, 1996; *G. palama* J. L. Barnard, 1977; *G. perlata* Schellenberg, 1938; *G. propodentata* Moore, 1986; *G. robusta* Ledoyer, 1982; *G. rhizophorae* Myers, 2009; *G. spinicoxa* Myers, 1972; *G. taihuensis* Morino & Dai, 1990; *G. teres* Myers, 1981; *G. trispinosa* Bano & Kazmi, 2010; *G. unidentata* Ren, 2006; *G. vietnamica* Dang, 1968.




Fig. 1. Map showing the sampling area.

Grandidierella halophilus, new species

(Figs. 2–10)

Type Material. — Holotype. Male, THAILAND, Inner Gulf of Thailand, Samut Sakorn (13.5100768°N, 100.3525478°E), abandoned saltpans (associated with *Ruppia maritima* Linnaeus), 24 Feb. 2011, Phenraphai, P., PSUZC-CR-0261.

Allotype. Female, collected with holotype, PSUZC-CR-0262 (adult female, 3.9 mm).

Other material. Same data as holotype, UKMMZ-1431 (10♂; 15♀); PSUZC-CR-0263 (10♂; 20♀).

Description. — **Male (holotype)** (Figs. 3–6). Total body length 6.9 mm. Body slender and subcylindrical.

Head. Head subequal in length to first 2 pereonites; rostrum not developed; inferior antennal sinus moderate and concave, 0.3 times of head length; eye distinct. Antenna 1 (Fig. 3A1) slightly longer than antenna 2, ratio of peduncular articles 1–3 as 1: 1.7: 1.1; article 1 slender, with 2 postero-marginal spines; flagellum with 22 articles, 1.4 times as long as peduncle; accessory flagellum uni-articulate, short. Antenna 2 (Fig. 3A2) peduncle stout; 4 segmented in ratio of 1: 3: 10: 7; inner margin of article 3 with 3 robust setae and 2 posterodistal robust setae; article 5 shorter than 4; flagellum short, subequal in length to peduncular article 5, composed of 7 articles; flagellum articles 2–7 with a pair of curved bifid robust setae on each article.

Upper lip (Fig. 5UL) or labrum round and broad, with small depression in the middle and pubescent on each lobe. Lower lip (Fig. 5LL) inner plates well developed and subtriangular in shape, mandibular process narrow but well developed; outer plate with arborescent setae on the inner face of the outer lobe, covered with thin hairs. Mandible (Fig. 5MD), both similar to each other except for number of accessory blades constituting 4 in right and 5 in left ones; right incisor 4 dentates, left incisor 5 dentates; lacinia mobilis armed with 4 teeth on left side and 5 teeth on right; molar process medium, ridged distally and serrate marginally, armed a single seta; palp triarticulate with ratios of 1: 1.2: 1.1, article 3 with apical setae and marginal setae. Maxilla 1 (Fig. 5MX1), inner plate small; outer plate with 8 apical robust setae; palp extending beyond outer plate, biarticulate with 6 apical spines. Maxilla 2 (Fig. 5MX2), inner plate with mediofacial row of 25 slender setae and one robust seta located in the middle, outer margin naked; outer plate subequal in size with inner plate. Maxilliped (Fig. 5MP), inner plate broad and short, apically provided with several plumose setae and 3 marginal setae; outer plate broad, almost reaching palp

article 2 with 13 marginal spines; palp 4-articulate with ratio of 2: 3: 3.2 : 1.

Pereon. *Gnathopod 1* (Fig. 3G1) carpochelate, larger than gnathopod 2; coxal plate subrectangular, inner side bearing ventral process; length ratio of articles from basis to dactylus 7.8: 1: 2.7: 8: 3.5: 3.5; basis slender, anterior margin straight, 3.9 as long as broad, with one fine seta on posteriodistal margin; ischium short, subrectangular; merus trapezoidal, anterodistally produced; carpus 1.6 times as long as broad, posterodistal corner produced with tooth and a smaller tooth present on inner face, posterior margin setose with a small tooth; propodus subequal to dactylus in length, 0.5 times length of carpus, posterior margin proximally concave, distally expanded; dactylus fairly curved, distal end concave with 2 robust setae. *Gnathopod 2* (Fig. 3G2) subchelate; coxa plate shallow with ventral margins posteriorly produced into triangular expansion; length ratio of articles from basis to dactylus 4.6: 1: 1.7: 3.8: 3.3: 1.8; basis slender, as long as articles 5 and 6 combined, anterior margin straight with 6 fine setae; ischium subrectangular; merus trapezoidal, posterior distal angle produced, posterior margin bearing long setae; carpus 2.5 times as long as wide, posterior margin crenulate, bearing 25 dense setae; propodus palmar margin transverse, with fine setae, bearing a bifid robust seta at posterodistal corner and 2 bifid robust setae on posterior margin; dactylus

slightly longer than palmar margin, inner margin crenulate with several fine setules.

Pereopod 3 (Fig. 4P3) slender; coxa plate small, subquadrate; length ratio of articles from basis to dactylus 4.8: 1: 2.5: 1.7: 1.7: 1.2; basis slender, sparsely setose on anterior and posterior margins and posterodistal corner; ischium short, subrectangular; merus slightly produced anterodistally; carpus suboval, posterior margin setose; propodus narrow, both margin bearing setae; dactylus long and thin, shorter than propodus. *Pereopod 4* (Fig. 4P4) similar to pereopod 3, coxal plate with ventral margin medially produced triangular; length ratio of articles from basis to dactylus 4.1: 1: 2.5: 1.4: 1.6: 1.1; basis slender, with short setae on anterior margin; ischium short, subrectangular; merus slightly produced anterodistally; carpus oval with setae on both anterior and posterior margins; propodus long and narrow, bearing setae on both anterior and posterior margins; dactylus long and thin. *Pereopod 5* (Fig. 4P5) coxa posteroventrally expanded into long, narrowly obtuse angle; length ratio of articles from basis to dactylus 5.8: 1: 2.2: 1.8: 0.8; basis subrectangular with fine setae along anterior margin; ischium short; carpus with two robust setae along posterior margin and on posterodistally; propodus with four robust setae along posterior margin; dactylus short, strongly curved. *Pereopod 6* (Fig. 4P6) elongate, 1.6 times as long as pereopod 5;



Fig. 2. Photography of *Grandidierella halophilus* new species, holotype, male, (PSUZC-CR-0261), 6.9 mm; allotype, female, (PSUZC-CR-0262), 3.9 mm, Samut Sakorn, Inner Gulf of Thailand.




Fig. 3. *Grandidierella halophilus* new species, holotype, male, (PSUZC-CR-0261), 6.9 mm. Samut Sakorn, Inner Gulf of Thailand. All scale bars represent 0.5 mm.




Fig. 4. *Grandidierella halophilus* new species, holotype, male, (PSUZC-CR-0261), 6.9 mm. Samut Sakorn, Inner Gulf of Thailand. All scale bars represent 0.5 mm.




Fig. 5. *Grandidierella halophilus* new species, holotype, male, (PSUZC-CR-0261), 6.9 mm. Samut Sakorn, Inner Gulf of Thailand. All scale bars represent 0.2 mm.




Fig. 6. *Grandidierella halophilus* new species, holotype, male, (PSUZC-CR-0261), 6.9 mm. Samut Sakorn, Inner Gulf of Thailand. Scale bars for U1–U3 represent 0.1 mm; PL1–PL3 represent 0.5 mm; T represents 0.05 mm.

coxa posteriorly produced with rounded lobe; length ratio of articles from basis to dactylus 3.7: 1: 2.7: 1.7: 3.3: 1; basis almost linear, with 7 small bifid robust setae along anterior margin; ischium short with setae on anteroventral corner; merus oblong, with bifid marginal robust setae on both anterior and posterior margins; carpus bearing two rows of bifid robust setae; propodus slender, slightly curved with two rows of marginal robust setae and setose posterodistally; dactylus tapering to a sharply pointed tip, subapex bearing small setae on both sides. *Pereopod 7* (Fig. 4P7) elongate, 1.8 times as long as pereopod 5; coxa bilobed, shallow; length ratio of articles from basis to dactylus 3.3: 1: 2.3: 1.7: 2.7: 1; basis anterior margin with a row of bifid robust setae and posterior margin with long plumose setae bifid robust setae; ischium short and subrectangular; merus elongate with setae on anterior and posterior margins, with one bifid robust seta posterodistally; carpus shorter and narrower than merus with several midlength setae along anterior margin; propodus elongate and slender, slightly curved with bifid robust setae along anterior margin; dactylus tapering to pointed tip, with one thin setae 2/3 from proximal end.

Pleon. *Pleopods 1–3* (Fig. 6PL1–PL3) well developed; peduncles cylindrical, longer than broad and fringed with several plumose setae and a pair of retinaculae on anterior margin; inner ramus slightly longer than peduncle with 9–10 articles, outer ramus shorter than inner ramus, both rami with facial setae.

Uropod 1 (Fig. 6U1) not extending beyond ends of other uropods; peduncle longer than inner and outer rami, fringed with bifid robust setae, peduncular apex bearing a posteroventral process; outer and inner margins of both rami lined with a row of robust setae, distal margins rounded and bearing with several robust setae. *Uropod 2* (Fig. 6U2) slightly longer than uropod 3, peduncle shorter than inner and outer rami, both with two bifid robust setae; outer ramus slightly longer than inner one, both rami lined with a row of bifid robust setae and distal margin bearing short and long robust setae. *Uropod 3* (Fig. 6U3) uniramous, peduncle much shorter than ramus; ramus elongate with short second article, both outer and inner margins with a row of long setae; apex with 4 long stiff setae. *Telson* (Fig. 6T) subtrapezoidal, ending with double pointed apex, each with one long robust seta and one short (thin) seta.

Female. (allotype) (Figs. 7–10). Total body length 3.9 mm (from tip of rostrum to apex of telson).—(sexually dimorphic characters).

Antenna 1 (Fig. 7A1) without accessory flagellum, flagellum with 15 articles. *Antenna 2* (Fig. 7A2) peduncular article 3 with 2 bifid robust setae at ventrodistal corner. *Maxilla 2*, inner plate without robust setae.

Pereon. *Gnathopod 1* (Fig. 7G1) subchelate, smaller than that of male. Coxal plate deeper than those of male; basis more robust, 1.8 times as long as broad; carpus subtriangular, 1.5 times as long as broad, posterior margin setose, with the posterior distal margin produced into a short stout tooth; propodus oval, shorter than carpus, palm oblique and defined

by a large bifid robust seta, palmar margin with short setule, distal end covered sparse plumose setae; dactylus curved with 3 inner marginal short stout robust setae. *Gnathopod 2* (Fig. 7G2) similar to that of male, slightly smaller to gnathopod 1; basis slender; carpus 2.2 times as long as wide; propodus wider than that of male, palm transverse and defined by 2 large bifid robust setae; dactylus curved, grasping margin with fine setule and 5 short stout robust setae.

Etymology. — The specific name “*halophilus*” is a combination of the Greek *halos* = salt and *philus* = friend, loving, referring to the habitat in which this species lives.

Remarks. — The specimens of *Grandidierella halophilus* new species were collected in a hypersaline habitat (~80 ppt) and associated with seagrass (*Ruppia maritima* Linnaeus). To date, only two *Grandidierella* species have been reported from hypersaline environments (*G. propodentata* and *G. exillis*). The former species shows a distinctive similarity with *G. halophilus* in the existence of the unusual arborescent setae on the inner face of the outer lobe of the lower lip (Moore, 1986: Fig. 2e). In addition, several differences also can be seen in comparing with *G. propodentata*, including (1) the spout-like male antenna 2 article 4 (vs. normal) (Moore, 1986: Fig. 2a); (2) the inwardly directed proximal tooth on male gnathopod 1 propodus (vs. posterior margin proximally concave, distally expanded; Moore, 1986: Fig. 1a) and (3) male gnathopod 1 carpus triangular with two large subequal teeth at posterodistal corner (vs. posterodistal corner with produced tooth and posterior margin setose with a small tooth; Moore, 1986: Fig. 1a). The new species differs from *G. exillis* in the presence of an accessory flagellum (Fig. 3A1); presence of the arborescent setae on the inner face of the outer lobe of the lower lip (Fig. 5LL); and by the presence of a second article on uropod 3 ramus (Fig. 6U3).

A sternal process in the male *G. halophilus* new species is seems to be unique among *Grandidierella* members. Only four species, *G. bonnieroides*, *G. exillis*, *G. trispinosa*, and *G. spinicoxa* share this distinct character. The new species, *Grandidierella halophilus*, is similar to *G. bonnieroides* in having the male sternal process; gnathopod 1 carpochelate with posterodistal corner with a produced tooth, inner face with a produced small tooth and posterior margin setose with a small tooth. As reported by Myers (1970), in his investigations on material from the Caribbean and Gulf of Mexico, the sternal process in *G. bonnieroides* shows considerable variation between populations, suggesting that most of the populations are genetically distinct. In agreement with Asari & Myers (1982), the south Indian *G. bonnieroides* also exhibits distinct local variations displaying the possibility of group sibling species.

Grandidierella halophilus can be easily distinguished from all other known species of the genus in the combination of the following characters: (1) coxal plate 2 with ventral margin posteriorly produced into triangular expansion (Fig. 3G2); (2) uropod 3 peduncle inflated with ramus bearing a small second article (Fig. 6U3); (3) lower lip with arborescent setae on the inner face of the outer lobe (Fig. 5LL). Moreover, all




Fig. 7. *Grandidierella halophilus* new species, allotype, female, (PSUZC-CR-0262), 3.9 mm. Samut Sakorn, Inner Gulf of Thailand. All scale bars represent 0.2 mm.




Fig. 8. *Grandidierella halophilus* new species, allotype, female, (PSUZC-CR-0262), 3.9 mm. Samut Sakorn, Inner Gulf of Thailand. All scale bars represent 0.2 mm.




Fig. 9. *Grandidierella halophilus* new species, allotype, female, (PSUZC-CR-0262), 3.9 mm. Samut Sakorn, Inner Gulf of Thailand. All scale bars represent 0.1 mm.




Fig. 10. *Grandidierella halophilus* new species, allotype, female, (PSUZC-CR-0262), 3.9 mm. Samut Sakorn, Inner Gulf of Thailand. Scale bar for U3 represents 0.05 mm; U2, U3 and T represent 0.1 mm, remaining represents 0.2 mm.

Table 1. Comparison of diagnostic character in some closely related *Grandidierella* species.

							Carpus of G1	Basis of G2	Basis of P6 & P7	U3 rami
	Body length (mm)	Accessory flagellum	Lower lip	Sternal spine	Basis of male G1 (Length/Width)	G1 (Carpus: Propodus of male G1)	Propodus of male G1 (posterior margin)	Basis of G2 (length/width)	Coxae 2–4	
<i>G. halophilus</i>	6.9	present	outer plate with two finger-like robust setae on lobes	present	slender (4×)	2.9×	posterior margin distally expanded	slender (2.06×) anterior margin smooth	coxa 2 produced anteriorly	anterior with small bifid robust setae
<i>G. bispinosa</i> Schellenberg, 1938	4.5	present	outer plate well developed with brittle shoulders	absent	stout (2.18×)	2.25×	posterior margin distally expanded	slender (2.53×), crenulate anterior margin	coxae 2–4 not produced	elongate with short second article
<i>G. bonnieirodes</i> Stephensen, 1948	4.5	absent	outer plate well developed with brittle shoulders	present	slender (2.25×)	2×	inflated in the middle	slender (3.2×) anterior margin smooth	coxae 2–4 not produced	elongate with one article
<i>G. exilis</i> Myers, 1981	7.5	absent	outer plate well developed with brittle shoulders	present	slender (2.46×)	2×	inflated in the middle	slender (5×) anterior margin smooth	coxae 2–4 not produced	elongate with one article
<i>G. mahafalensis</i> Coutière, 1904	6.7	present	outer plate well developed with brittle shoulders	absent	slender (2.63×)	2.06×	posterior margin distally expanded	slender (3.04×) anterior margin smooth	coxae 2–4 not produced	anterior margin with long plumose setae
<i>G. propodentata</i> Moore, 1986	8.2	present	outer plate with two finger-like robust setae on lobes	absent	slender (2.76×)	1.46×	with tooth on basal	slender (4.29×) anterior margin smooth	coxae 2–4 not produced	anterior margin with small setae
<i>G. spinicoxa</i> Myers, 1972	8.0	present	outer plate well developed with brittle shoulders	present	slender (2.11×)	2.24×	inflated in the middle	slender (4×) anterior margin smooth	coxa 2–4 produce anteriorly	anterior margin with long setae
<i>G. trispinosa</i> Bano & Kazmi, 2010	6.6	no data	no data	present	slender (2.9×)	3×	posterior margin distally expanded	slender (3.6×) anterior margin smooth	coxa 2–4 produced anteriorly	anterior margin with long setae

of the robust setae occurring on antennae 1–2, gnathopods 1–2, pereopods 5–7 and uropods 1–3 are bifid. This character has not been reported elsewhere in the genus but has been shown in illustrations for *G. mahafalensis* (Ruffo, 1958) and *G. propodentata* (Moore, 1986). A summary of the distinguishing characters of the species discussed here given in Table 1.

ACKNOWLEDGEMENTS

We are grateful to the Department of Biology, Faculty of Science, Prince of Songkla University and Marine Science Programme, School of Environmental and Natural Resource Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia for the laboratory facilities. Thanks are due to the National Research University Project of Thailand's Office of the Higher Education Commission for financial support. Thanks to Phairin Phenphraphai from Marine and Coastal Resources Research Center, the Upper Gulf of Thailand in providing us the specimens for this study. We also thank our colleagues Charles Oliver Coleman from Museum für Naturkunde, Germany for the essential literature.

LITERATURE CITED

- Angsupanich, S. & R. Kuwabara, 1995. Macrobenthic fauna in Thale Sap Songkla, a brackish lake in southern Thailand. *Lakes & Reservoirs: Research and Management*, **1**: 115–125.
- Angsupanich, S., A. Siripech & M. Charoenporntip, 2005. Macrobenthic fauna community in the Middle Songkhla Lake, Southern Thailand. *Songklanakarin Journal of Science and Technology*, **27**: 365–390.
- Ariyama, H., 1996. Four species of the genus *Grandidierella* (Crustacea: Amphipoda: Corophiidae) from Osaka Bay and the northern part of the Kii Channel, central Japan. *Publications of the Seto Marine Biological Laboratory*, **37**: 167–193.
- Ariyama, H., S. Angsupanich & E. Rodcharoen, 2010. Two new species of the genus *Kamaka* (Crustace: Amphipoda: Kamakidae) from the Songkla Lagoon, southern Thailand. *Zootaxa*, **2404**: 55–68.
- Asari, K. P. & A. A. Myers, 1982. Taxonomic studies on the genus *Grandidierella* Coutière (Crustacea, Amphipoda) IV. Indian species. *Bulletin du Museum National d'Histoire Naturelle, Paris*, **4A**: 237–256.
- Azman, B. A. R. & B. H. R. Othman, 2012. Two new species of amphipods of the superfamily Aoroidea (Crustacea: Corophiidae) from the Strait of Malacca, Malaysia, with a description of a new genus. *Zoological Studies*, **51**: 232–247.
- Bano, H. & Q. B. Kazmi, 2010. *Grandidierella trispinosa*, a new species of amphipod from the Karachi coast, Pakistan (Crustacea: Amphipoda: Aoridae). *Turkish Journal of Zoology*, **34**: 151–157.
- Barnard, J. L., 1970. Sublittoral Gammaridea (Amphipoda) of the Hawaiian Islands. *Smithsonian Contributions to Zoology*, **34**: 1–286.
- Barnard, J. L., 1977. The cavernicolous fauna of Hawaiian lava tubes 9. Amphipoda (Crustacea) from brackish lava ponds on Hawaii and Maui. *Pacific Insects*, **17**: 267–299.
- Barnard, K. H., 1935. Report on some Amphipoda, Isopoda, and Tanaidacea in the collections of the Indian Museum. *Records of the Indian Museum*, **37**: 279–319.
- Barnard, K. H., 1951. New records and description of new species of isopods and amphipods from South Africa. *Annals Magazine of Natural History*, **4**: 698–709.
- Barnard, K. H., 1952. Description of a new species of amphipod. *Transactions of the Royal Society of South Africa*, **33**: 279–282.
- Bochert, R. & M. L. Zettler, 2010. *Grandidierella* (Amphipoda, Aoridae) from Angola with description of a new species. *Crustaceana*, **83**: 1209–1219.
- Bussarawich, S., 1985. Gammaridean Amphipoda from mangroves in southern Thailand. *Fifth Seminar on Mangrove Ecosystems*. Phuket, Thailand. Pp. 1–17.
- Chevreux, E., 1926. Amphipodes, 1: Gammariens (Cont.). *Voyage de la Goelette France*, **20**: 365–398.
- Chilton, C., 1921. Fauna of the Chilka Lake. Amphipoda. *Memoirs of the Indian Museum*, **5**: 519–558.
- Chilton, C., 1925. Zoological results of a tour in the Far East. The Amphipoda of Talé Sap. *Memoirs of the Asiatic Society of Bengal*, **6**: 531–539.
- Coleman, C. O., 2003. “Digital inking”: How to make perfect line drawings on computers. *Organism, Diversity and Evolution, Electronic Supplement*, **14**: 1–14. <http://senckenberg.de/odes/03-14.htm>.
- Coutière, H., 1904. Sur un nouveau d'Amphipode *Grandidierella mahafalensis* provenant de Madagascar. *Bulletin de la Societe de Philomath, Paris*, **6**: 166–174.
- Dang, N. T., 1968. Nouveaux amphipods des eaux douces et saumâtres du Nord Viet Nam. *Zooligichesky Zhurnal*, **47**: 212–222.
- Griffiths, C. L., 1974. The Amphipoda of southern Africa. Part 3. The Gammaridea and Caprellidea of Natal. *Annals of the South African Museum*, **62**: 209–264.
- Hou, Z. E. & S. Q. Li, 2002. A new species of the genus *Grandidierella* from Lake Chaohu, China (Crustacea: Amphipoda: Aoridae). *Acta Zootaxonomica Sinica*, **27**: 225–234.
- Ledoyer, M., 1967. Amphipodes gammariens des herbiers de phanerogammes marines de la région de Tulear (République Malgache). *Etude systematique et l'Universite de Madagascar*, **5**: 121–170.
- Ledoyer, M., 1979. Expedition Rumphius II (1975). Crustaces parasites, commensaux etc.VI. Crustaces Amphipodes Gammariens. *Bulletin du Museum National d'Histoire Naturelle, Paris*, **1**: 137–181.
- Ledoyer, M., 1982. Crustaces Amphipodes Gammariens Famille des Acanthonozomatidae a Gammaridae. Faune de Madagascar, **59**: 1–598. *Bulletin du Museum National d'Histoire Naturelle, Paris*, **3**: 213–226.
- Moore, P. G., 1986. A new species in the genus *Grandidierella* Coutière (Crustacea: Amphipoda) from an Australian solar salt-works. *Journal of Natural History*, **20**: 1393–1399.
- Morino, H. & A. Dai, 1990. Three amphipod species (Crustacea) from the east China. *Publications of the Itako Hydrobiological Station*, **4**: 7–27.
- Myers, A. A., 1970. Taxonomic studies on the genus *Grandidierella*, with a description of *G. dentimera* sp. nov. *Bulletin of Marine Science*, **20**: 135–147.

- Myers, A. A., 1972. Taxonomic studies on the genus *Grandidierella* Coutière (Crustacea: Amphipoda) II. The Malagasy species. *Bulletin du Museum National d'Histoire Naturelle, Paris*, **64**: 789–796.
- Myers, A. A., 1981. Taxonomic studies on the genus *Grandidierella* Coutière (Crustacea, Amphipoda). III. Fijian, Australian and Saudi Arabian species. *Bulletin du Museum National d'Histoire Naturelle*, **3A**: 213–226.
- Myers, A. A., 1998. The Amphipoda (Crustacea) of New Caledonia: Aoridae. *Records of the Australian Museum*, **50**: 187–210.
- Myers, A. A., 2009. Aoridae. In: Lowry, J. K. & A. A. Myers (eds.), *Benthic Amphipoda (Crustacea: Peracarida) of the Great Barrier reef, Australia*. *Zootaxa*, **2260**: 220–278.
- Ren, X., 2006. Crustacea Amphipoda Gammaridea (I). *Fauna Sinica Invertebrata*, **41**: 1–588.
- Ruensirikul, J., S. Angsupanich & A. Phongdara, 2007. Abundance and diversity of amphipod crustaceans in the Upper Songkhla Lagoon. *Songklanakarin Journal of Science and Technology*, **29**: 1225–1249.
- Ruffo, S., 1958. Amphipodes Terrestres et des eaux continentales de Madagascar, des Comores et de la Réunion (Etudes sur les Crustacés Amphipodes. - L). *Mémoires de l'institut Scientifique de Madagascar*, **12**: 35–66.
- Schellenberg, A., 1925. Amphipoda, Beiträge zur Kenntnis der Meeresfauna. *Westafrikas*, **3**: 113–204.
- Schellenberg, A., 1936. Zwei neue Amphipoden des Stillen Ozeans und zwei Berichtungen. *Zoologischer Anzeiger*, **116**: 153–156.
- Schellenberg, A., 1938. Littoral Amphipoden des Topischen Pazificks. *Kungliga Svenska Vetenskapsakademiens Handlingar*, **16**: 1–50.
- Shoemaker, C. R., 1935. A new species of amphipod of the genus *Grandidierella* and a new record for *Melita nitida* from Sinaloa, Mexico. *Journal of the Washington Academy of Sciences*, **25**: 65–71.
- Stephensen, K., 1938. *Grandidierella japonica* n.sp. A new amphipod with stridulating organ from brackish water in Japan. *Annotationes Zoologicae Japonenses*, **17**: 179–184.
- Stephensen, K., 1948. Amphipods from Curaçao, Bonaire, Aruba and Margarita. *Studies on the Fauna of Curaçao, Aruba, Bonaire and the Venezuelan Islands*, **3**(11): 1–20.