

## ECOLOGICAL IMPORTANCE OF CHEMOAUTOTROPHIC LUCINID BIVALVES IN A PERI-MANGROVE COMMUNITY IN EASTERN THAILAND

**Erin Meyer**

*Department of Integrative Biology, University of California, Berkeley,  
California 94720-3140, U. S. A.  
Email: meyererini@yahoo.com*

**Bancha Nilkerd**

*Faculty of Marine Technology, Burapha University, Chantaburi 22120, Thailand.*

**Emily A. Glover and John D. Taylor**

*Department of Zoology, The Natural History Museum, London SW7 5BD, United Kingdom.  
Email: emilyglover@mac.com; j.taylor@nhm.ac.uk*

**ABSTRACT.** – A quantitative survey of molluscs inhabiting a disturbed, intertidal muddy-sand flat near mangroves in a sheltered bay in southeastern Thailand recorded a high abundance (to 1,380 m<sup>-2</sup>) of the small, chemosymbiotic lucinid bivalves *Pillucina vietnamica* and *Indoaustriella dalli*. Two other larger lucinids, *Anodontia bullula* and *A. philippiana*, were recorded living near the transects, whereas another, *Ctena delicatula*, inhabited a cobble habitat on a rocky headland at the entrance to the bay. The presence of symbiotic bacteria in the ctenidial filaments was confirmed for each species. Most abundant of the associated molluscan fauna were the epifaunal gastropods *Cerithideopsis cingulata*, *Cerithium coralium* and *Clithon oualaniensis*, and the infaunal bivalves *Gafrarium tumidum*, *Anomalocardia squamosa*, *Pristis capsoides* and *Anadara troscheli*. Biomass estimates for the lucinids ranged from 0.51 to 3.95 gm<sup>-2</sup>. The results highlight the ecological importance of chemoautotrophic bivalves in peri-mangrove communities.

**KEYWORDS.** – chemosymbiosis, Lucinidae, abundance, diversity, inventory.

---

### INTRODUCTION

Although chemosymbiotic communities around deep-sea hydrothermal vents and cold seeps have attracted much attention, by far the most widespread sulphidic habitat is the subsurface anoxic zone of shallow-water marine sediments (Fenchel & Riedl, 1970; Ott et al., 2004; Stewart et al., 2005). This habitat supports a high diversity of chemosymbiotic organisms and prominent amongst these are bivalves of the families Lucinidae and Solemyidae (Fisher, 1990; Reid, 1990; Taylor & Glover, 2000, 2006). Lucinidae have attracted much biological interest because of the likely obligate chemosymbiosis with sulphide-oxidising bacteria housed in the gills and from which they derive much of their nutrition (Reid & Brand, 1986; Distel, 1998; Gros et al., 2003). Typically lucinids live at the boundary between oxic and suboxic zones in the sediment from where sulphide-rich interstitial water is channeled into the mantle cavity by probing and tunneling activities of the foot (Stanley, 1970; Dando et al., 1986, 1994). Lucinids are often associated with organic-rich habitats such as seagrass beds (Jackson, 1972; Barnes & Hickman, 1999; Johnson et al., 2002), mangrove

fringes (Lebata & Primavera, 2001; Glover et al., 2008), and hydrocarbon rich sites such as cold seeps (Oliver & Holmes, 2006), but many others inhabit intertidal and subtidal, unvegetated sand and mud. Although the biology of a few lucinid species, notably from the northern Atlantic, is known in some detail (e.g., Fisher & Hand, 1984; Frenkiel & Mouëza, 1995; Frenkiel et al., 1996; Johnson & Fernandez, 2001; Gros et al., 2003), for the tropical Indo-West Pacific, most species remain unstudied and lack even basic information of distributional range, habitat preferences or estimates of abundance. Johnson et al. (2002) highlighted the importance of chemoautotrophic symbioses in estimates of the primary production of seagrass beds, but for most areas, information is lacking.

Preliminary sampling of intertidal sediments fronting the mangrove fringe of Kungkrabaen Bay, Thailand, indicated that two small (< 10 mm), poorly known lucinids were abundant. In order to make an assessment of the importance of chemosymbiotic bivalves to this community, we decided to make a more detailed quantitative survey from which to estimate species composition, abundance, biomass,

and distribution of the lucinids and associated molluscan fauna.

## MATERIALS AND METHODS

Kungkrabaen Bay is a shallow, semi-enclosed bay on the southeastern coast of the Gulf of Thailand, with an area of approximately 640 hectares (6.4 km<sup>2</sup>) (Fig. 1). The inner bay is fringed by extensive mangrove stands, but the landward margins of these are heavily disturbed by aquacultural development of prawn and fish ponds. Most of the bay is extremely shallow with a large area of sand and mud flats emerged at low spring tides or covered by just a few centimetres of water.

**Faunal sampling.** – Sampling took place over eight days between 22 August and 2 September 2005. Two transects for quantitative sampling were selected in the middle part of the bay (Fig. 1); the shorter transect (2) was used to confirm the general uniformity of the fauna. Transect 1 (started at 12°35.4'N 101°54.39'E) extended for 406 m, with 10 stations sampled for molluscs and sediment analysis. The first station (0) was located 0.5 m from the mangrove prop roots, the next 5 m to seaward, with a further eight stations located 50 m apart. Transect 2 (12°35.89'N 101°54.19'E) comprised three stations between 80 m and 400 m from the mangrove fringe. Extremely soft mud, extending to 80 m from the mangrove margin, precluded sampling in that area. At each station, two quadrat samples (23.5 × 32 cm = 752 cm<sup>2</sup>) were taken; the sediment was excavated with trowels to a depth of 15 cm and sieved through 2 mm-mesh sieves. All molluscs were collected for examination in the laboratory where they were identified, counted, and the shells of sub-samples measured.



Fig. 1. Aerial view of Kungkrabaen Bay showing location of the transects (T1 and T2). The asterisk indicates location of rocky shore inhabited by *Ctena delicatula*.

Other samples were taken from a wide range of intertidal habitats within the bay, including seagrass patches and amongst prop roots at mangrove margins; samples within the bay contained similar species to those encountered along the transects. Additionally, we sampled the fauna of intertidal cobble substrata at the rocky, exposed headland (Fig. 1) at the northern entrance to the bay (Laem Ban Tha Klaeng, 12°35.26'N 101°53.03'E). Away from Kungkrabaen Bay, a similar cobble and boulder habitat was sampled at Koh Nom Sao, Laem Singha District (12°27.9'N 102°0.14'E).

**Biomass.** – Estimates of biomass were made from dry weights of the two lucinid species, and for comparison, four of the most abundant infaunal bivalve species. Shells were measured for 15 individuals of each species; the bodies were then removed and placed on pieces of foil, dried in an 80°C oven for 24 hours, and the dry weights obtained. Regressions of shell length versus dry weight, combined with the size/density data, were used to estimate biomass·m<sup>2</sup> for each species.

**Sediment analysis.** – Particle-size analysis was performed by Gardline Environmental Ltd. using nested sieves for larger grains, with particle size distributions below 1.00 mm determined using a Malvern Mastersizer 2000 particle sizer. To determine total organic matter, 1 g of ground sediment was dried at 50°C to constant weight, and then heated in a muffle furnace at 450°C for 4 hours. The cooled sample was then reweighed with the percentage loss on ignition (LOI) calculated.

**Gill preparations.** – Ctenidial and other tissues were fixed in a cold 2.5% solution of glutaraldehyde in phosphate buffer. Tissue pieces were sliced with a razor blade, dehydrated through an ascending series of acetone solutions, and then critical point dried for examination by scanning electron microscopy (SEM).

**Identification of fauna.** – Species identifications were made using the reference collections and library at The Natural History Museum, London, with comparison with type specimens when appropriate. Voucher material of all species is deposited in the collections of The Natural History Museum, London.

## RESULTS

The two quantitative transects were located in the middle part of the bay on extensive intertidal, gently shelving, muddy sand flats (Fig. 1). Mangroves and their prop roots, largely of *Rhizophora*, form a sharp landward boundary to the mud flat (Fig. 2A), which is emerged at low spring tides to about 450 m from the mangrove fringe. The mud flat is colonized by patchy growths of the seagrass *Halodule pinifolia* (Miki) den Hartog (Fig. 2C), as well as filamentous green algae and *Ulva*. Small patches of the larger seagrass, *Enhalus acoroides* (L.f) Royle, were also present adjacent to seaward parts of the transect. Seagrass growth was generally more luxuriant about 300–400 m from the mangrove fringe.

**Sediment.** – Results of the sediment analysis are summarized in Table 1. Sediments were relatively uniform across the transect, ranging between sandy mud, muddy sand, and fine sand, with a varying gravel component, mainly of shell fragments. The finest sediments occurred around mangrove prop roots (Station 0) or at Stations 6 and 7, approximately 250–300 m from the mangroves, where growth of the seagrass *Halodule* is more abundant with the leaves, roots, and rhizomes likely stabilising the sediment. In general, sediments across the transect were poorly sorted, reflecting significant components of shells and coarser debris. All sediments were blackened just below surface, but in areas where *Halodule* growth was most dense, the sediment was particularly black and smelled strongly sulphurous. Organic content varied between 0.5 and 2.7% LOI with higher values in the seaward part of the transect, where seagrass was more abundant.

**Species composition of the molluscan fauna.** – The molluscan fauna recorded from the transects and other Lucinidae located within the bay are listed in Table 2 and the most abundant species illustrated in Fig. 3. Five species of Lucinidae (Fig. 3A–E) were recorded alive; these are briefly discussed below. The presence of symbiotic bacteria

contained in bacteriocytes within the ctenidial filaments is confirmed in all species. Additionally, a few worn, dead shells of *Cardiolucina macassari* were found in sieved sediment.

*Pillucina vietnamica*, a small lucinid (length to 8.3 mm), is widely distributed along continental shores from the Red Sea to eastern Australia (Glover & Taylor, 2001), usually occurring in sediment close to mangroves. Large pink ctenidia are visible through the shell. The ctenidial filaments are thick, with bacteriocytes packed with coccoid to rod-shaped bacteria to 8  $\mu\text{m}$  long and 3  $\mu\text{m}$  wide (Fig. 4A–B).

*Indoaustraliella dalli*, a small species (length to 9.8 mm) first described from the Gulf of Thailand near Kungkrabaen Bay (Lynge, 1909), has a relatively narrow distribution in Southeast Asia. It belongs to a clade of lucinids closely associated with, or peripheral to, mangrove habitats (Glover et al., 2008). The ctenidia are red-brown and composed of thick filaments with the bacteriocytes containing abundant rod-shaped bacteria to 8–10  $\mu\text{m}$  length and 1–2  $\mu\text{m}$  width (Fig. 4C–D). The ctenidial filaments are notable for the abundant, large (8  $\mu\text{m}$  diameter), spherical, probably sulphur-rich granules (Fig. 4D).



Fig. 2. A, Mangrove fringe and intertidal muddy sand flat at site of Transect 1; B, cobble habit of *Ctena delicatula* at the rocky headland at northern entrance to Kungkrabaen Bay; C, surface of sediment on Transect 1 with *Halophila* seagrass and the gastropods *Clithon oualaniensis*, *Cerithideopsisilla cingulata*, and *Cerithium coralium*; D, local Thai fisherwomen digging for *Lingula* near Transect 1.

Table 1. Summary of sediment analysis across Transect 1.

Station Number	Distance from mangroves (m)	Mean size ( $\mu\text{m}$ )	Mean phi	% Fines (< 63 $\mu\text{m}$ )	% Sand	% Coarse (> 2mm)	Sorting Description	Sediment Description	Total Organic Content (% LOI)
0	0.5	13	6.3	44.9	54.0	1.1	very poor	sandy mud	1.0
1	5.5	29	5.1	25.8	71.7	2.5	very poor	muddy sand	0.5
2	56	25	5.3	14.7	75.2	10.1	very poor	fine sand with gravel	1.2
3	106	16	6.0	5.8	88.7	5.5	poor	fine sand with gravel	0.9
4	156	22	5.5	24.5	72.9	2.7	very poor	muddy sand	0.8
5	206	41	4.6	38.5	57.6	3.9	very poor	sandy mud	0.7
6	256	16	6.0	42.2	55.8	2.0	poor	sandy mud	1.4
7	306	16	5.9	46.5	53.2	0.3	very poor	sandy mud	0.8
8	356	23	5.4	34.3	57.6	8.1	very poor	muddy sand with gravel	1.4
9	406	15	6.1	13.3	82.2	4.5	very poor	muddy sand with gravel	2.7

*Anodontia (Cavatidens) bullula* (length to 25 mm) is known only from a few records from northern Australia and Southeast Asia (Taylor & Glover, 2005). It burrows to depths of approximately 20 cm. The ctenidia are large and purple-black with bacteriocytes containing abundant, elongate bacteria about 5  $\mu\text{m}$  long and 0.5–0.7  $\mu\text{m}$  wide (Fig. 5B).

*Anodontia (Pegophysema) philippiana*, the largest (length to 67 mm) of the lucinids present in the bay, was not recorded in quantitative samples but several individuals were live-collected close to the seaward edge of the mangroves from sediment depths of 25–40 cm. The species is widely distributed across the Indo-West Pacific from East Africa to New Caledonia (Taylor & Glover, 2005), with the usual habitat near the outer fringe of mangroves (Leбата & Primavera, 2001). The ctenidia are large, purple-black in colour, and composed of thick filaments with bacteriocytes containing long, thin bacteria (Taylor & Glover, 2005: Fig. 5D).

*Ctena delicatula* was not recorded within Kungkrabaen Bay but was common at Laem Ban Tha Klaeng, a rocky headland at the northern entrance (12°35.263'N 101°53.034'E) where it occurred along with *Gafrarium dispar* (Dillwyn, 1817) and *Semele carnicolor* (Hanley, 1847) in an intertidal, wave-washed area of cobbles, pebbles, and gravel accumulated between rocky outcrops (Fig. 2B). *Ctena delicatula* was also abundant in a similar boulder and cobble habitat on the island of Koh Nom Sao, Laem Singha District (12°27.9'N 102°0.143'E; site KKB-21) near to the southeastern point of Kungkrabaen Bay. Compared with the other lucinids of Kungkrabaen Bay, *C. delicatula* has thin, flimsy gills with short abfrontal extensions of the filaments and relatively few bacteriocytes containing small rod-shaped bacteria about 2.0  $\mu\text{m}$  length and 0.7  $\mu\text{m}$  width. Cells containing numerous, rounded granules (5  $\mu\text{m}$ ) are concentrated at the proximal ends of the filaments (Fig. 5A).

**Distribution and abundance.** – The distribution of molluscs across the transects is shown in Tables 3 and 4 and the most abundant species are highlighted in Fig. 6.

For Transect 1, 26 molluscan species totaling 2,673 individuals were recorded, numbering 4–13 (mean 8.8) species and 134 animals per quadrat (Table 3). Nine species accounted for 98% of the total number, and 10 species were represented by only one or two individuals. By far the most abundant mollusc was the epifaunal gastropod *Cerithideopsis cingulata* that accounted for 47% of individuals and reached densities of 2,220  $\text{m}^{-2}$ . Next most abundant were the lucinid *Pillucina vietnamica*, the small neritid gastropod *Clithon oualaniensis*, the cerithiid *Cerithium coralium*, the lucinid *Indoaustriella dalli*, and the tellinid bivalve *Pristis capsoides*. The highest densities of molluscs were recorded between 100 and 200 m from the mangroves (Fig. 6), mainly accounted for by the abundance of *Cerithideopsis cingulata* and *Pillucina vietnamica*, and declined both seawards and landwards. The deposit-feeding tellinid *Pristis capsoides* was more abundant in the landward part of the transect. Qualitative sampling near the transect showed that *Anodontia philippiana* and *Psammotaena elongata* (Lamarck, 1818) were also more common near the mangroves.

From the shorter Transect 2, 19 molluscan species and 744 individuals were recorded (Table 4), ranging from 9 to 13 (mean 10) species and 141 individuals per quadrat. Seven of the species accounted for 96% of total number and eight of the species were present as one or two individuals only. Again, *Cerithideopsis cingulata* was the most abundant species with densities to 868  $\text{m}^{-2}$ , followed by *Clithon oualaniensis*, *Cerithium coralium*, *Pillucina vietnamica*, and *Indoaustriella dalli*.

The distribution of the two abundant lucinid species is shown in more detail in Fig. 7. Across Transect 1, *Pillucina vietnamica* reached maximum densities of 800–860  $\text{m}^{-2}$

Table 2. Taxonomic list of molluscan species recorded from transects within Kungkrabaen Bay, with additional lucinids recorded from the bay. Average shell height (H) or length (L) of specimens on transect is given following the name.

## Bivalvia

### Arcidae

*Anadara troscheli* (Dunker, 1882), H to 30 mm

### Ostreidae

*Saccostrea* sp. (juveniles attached to *Cerithideopsis*), H to 12 mm

### Lucinidae

*Anodontia (Cavatidens) bullula* (Reeve, 1850), H to 25 mm

*Anodontia (Pegophysema) philippiana* (Reeve, 1850), H to 67 mm. Not recorded on transect but present near mangroves

*Pillucina vietnamica* Zorina, 1978, H to 8 mm

*Indoaustriella dalli* (Lynge 1909), H to 9.8 mm

*Ctena* cf. *delicatula* (Pilsbry, 1904), H to 11.5 mm. Not recorded from transect but at entrance to bay. (Note: A systematic revision of *Ctena* species from the Indo-Pacific and elsewhere is urgently needed. Although superficially similar to the widespread Pacific species *Ctena bella* (Conrad, 1837), the Kungkrabaen Bay species differs in being smaller, more anteriorly extended and with less prominent ribbing and narrower interspaces. These differences are corroborated by molecular analyses (Taylor & Glover, unpubl. data). The syntypes of *Codakia delicatula* Pilsbry, 1904, described from the Ryuku Islands, have similar morphological features to the KKB specimens and we here use this as the provisional name.)

*Cardiolumacina macassari* (Prashad, 1932), H to 5.7 mm, dead shells on transect

### Solenidae

*Solen* sp., L to 26 mm

### Mactridae

*Meropesta pellucida* (Gmelin, 1791), H = 8 mm

*Mactra* sp. (juvenile), L = 4 mm

### Ungulinidae

*Cycladicama cumingi* (Hanley, 1845), H to 19.5 mm

### Veneridae

*Anomalocardia squamosa* (Linnaeus, 1758), H to 27 mm

*Dosinia cretacea* (Reeve, 1850), H = 20 mm

*Gafrarium tumidum* Röding, 1798, H to 40 mm

*Costellipitar* sp., H = 7 mm

*Marcia hiantina* (Lamarck, 1818), H to 22 mm

*Placamen calophyllum* (Philippi, 1836), H to 22 mm

### Tellinidae

*Pristis capsoides* (Lamarck, 1818), H to 34 mm

*Nitidotellina nitens* (Deshayes, 1855), H to 7 mm

*Pinguitellina pinguis* (Hanley, 1844), H to 10 mm

*Tellinides* sp., H to 7 mm

### Corbulidae

*Corbula* sp., H = 7 mm

### Laternulidae

*Laternula truncata* (Lamarck, 1818), H = 10 mm

## Gastropoda

### Neritidae

*Clithon oualaniensis* (Lesson, 1831), L to 5.5 mm

### Potamididae

*Cerithideopsis cingulata* (Gmelin, 1791), L to 20 mm

*Cerithidea alata* (Philippi, 1849), L to 20 mm

### Cerithiidae

*Cerithium coralium* Kiener, 1841, L to 25 mm

*Clypeomorus pellucida* (Hombron & Jacquinot, 1852), L to 17 mm

*Cerithium* sp., L = 12 mm

### Nassariidae

*Hebra corticata* (A. Adams, 1852), L to 13.7 mm

*Plicarularia leptospira* (A. Adams, 1852), L to 15.5 mm

*Plicarularia pullus* (Linnaeus, 1758), L to 15.5 mm

### Conidae

*Euclithara* sp., L to 12 mm



Fig. 3. Lucinidae and other common molluscs from the transects. A, *Pillucina vietnamica*; B, *Indoaustriella dalli*; C, *Ctena delicatula*; D, *Anodontia (Cavatidens) bullula*; E, *Anodontia (Pegophysema) philippiana*; F, *Pristis capsoides*; G, *Nitidotellina nitens*; H, *Anadara troscheli*; I, *Gafrarium tumidum*; J, *Anomalocardia squamosa*; K, *Solen* sp.; L, *Clithon oualaniensis*; M, *Cerithideopsis cingulata*; N, *Cerithium coralium*; O, *Plicarcularia pullus*.




Fig. 4. Sections through ctenidial filaments of Lucinidae: A, *Pillucina vietnamica* gill filament showing distal ciliated zones and broad bacteriocyte zone; B, *P. vietnamica*, detail of filaments with bacteriocytes packed with symbiotic bacteria; C, *Indoaustriella dalli* gill filaments; D, *I. dalli* detail of gill filament showing bacteria and large spherical granules. Scale bars = A, 20  $\mu$ m; B, D, 10  $\mu$ m; C, 50  $\mu$ m. b, bacteria; bz, bacteriocyte zone; cz, ciliated zone; gr, granule.

between 55–155 m from the mangrove fringe, whereas the less abundant *Indoaustriella dalli* attained maximum densities of 100–112 m<sup>-2</sup> at 200–250 m. Only three individuals of *Anodontia bullula* were recorded in the quadrats. The mean density of all Lucinidae across the transect was 526 m<sup>-2</sup> (range 126–938 m<sup>-2</sup>). On Transect 2, *Pillucina vietnamica* reached a maximum density of 692 m<sup>-2</sup> at 400 m from the mangrove fringe, while *I. dalli* varied between 35–186 m<sup>-2</sup> with a maximum at 300 m. Total Lucinidae had a mean density of 423 m<sup>-2</sup> (range 100–771 m<sup>-2</sup>).

**Size.** – Both *Pillucina vietnamica* and *Indoaustriella dalli* are small species (< 10 mm). The size distribution for *Pillucina vietnamica* across Transect 1 is shown in Fig. 8A; the median size ranged 4.5–5.7 mm, with the individuals from amongst the mangrove prop roots (0.5 m) slightly larger at 6.5 mm. At all 10 stations, 50% of the individuals fell within a narrow range of sizes, variation 0.7–1.3 mm, with 75% of the individuals within 1.7–3.2 mm of each other (Fig. 8A). A one-way Analysis of Variance (ANOVA) followed by a Tukey Test, with  $\alpha = 0.05$ , revealed that the means are significantly different between most of the stations, except those for stations 3 and 4 are equal, as are those for stations 8 and 9. However, no clear relationship between distance from the mangrove and size was indicated in these data.

Median sizes of *Indoaustriella dalli* ranged between 4.5–9.0 mm (Fig. 8B), but the sample sizes were smaller. Because the sample sizes were so small (< 17 individuals) compared to those of *Pillucina vietnamica* (typically >> 17 individuals), the ANOVA and Tukey Test, with  $\alpha = 0.05$ , revealed that the means for eight of the stations were equal. The two exceptions were the station within the mangrove prop roots (0.5 m), which had a sample size of two, and the station at 306 m from the mangrove fringe, which had the largest size range, 4.8 mm (Fig. 8B).

Analysis of size classes for *Pillucina vietnamica* (Fig. 9) shows that the 4.1–6.0 mm size class was dominant at all stations, except within the mangrove prop roots where the 6.1–8.0 mm size class was dominant. However, only 17 individuals were found at station 1 (prop roots), compared to 33–132 individuals collected at the other stations. The 2.1–4.0 mm size class was present between 5.5 and 256 m from the mangroves and completely absent from all other stations. Also, the seaward parts of the transect (206–406 m) had a higher proportion of individuals in the 6.1–8.0 mm size class than shoreward stations, again except for Station 1.

**Biomass.** – Results of the biomass determinations for the more abundant bivalves are shown in Table 5. Dry weights of the two Lucinidae species were estimated at approximately 1.9 gm dry weight m<sup>-2</sup>, with *Pillucina vietnamica* accounting for 1.75 gm m<sup>-2</sup>. By comparison, the larger-bodied, but much less abundant, *Gafrarium tumidum* and *Pristis capsoides* contributed approximately 2.65 gm m<sup>-2</sup> each, with *Anadara troscheli* and *Anomalocardia squamosa* at 0.98 and 0.65 gm·m<sup>-2</sup>, respectively.

**Feeding guilds of molluscan fauna.** – The molluscan

community recorded along the transect included species with a variety of feeding strategies including algal grazing, detritivory, suspension feeding, deposit feeding, carnivory, and chemosymbiosis. We have classified the species into these broad categories (Fig. 10). Grazing and surface browsing gastropods, *Clithon oualaniensis*, *Cerithideopsis cingulata*, and *Cerithium coralium* were the most abundant epifaunal molluscs and responsible for high levels of bioturbation from their surface grazing trails (Fig. 2C). According to Kamimura & Tsuchiya (2004), *Cerithideopsis* is an obligate detritivore, ingesting particulate organic material from the sediment surface layer. Apart from the largely chemoautotrophic lucinids, most other common bivalves are shallow-burrowing suspension feeders, except for the




Fig. 5. Sections through gill filaments of Lucinidae: A, *Ctena delicatula*, showing distal ciliated zone, thin bacteriocyte zone, and proximal granule cells; B, *Anodontia bullula*, section through bacteriocytes with long, thin bacteria. Scale bars = A, 20 µm; B, 5 µm. be, bacteriocyte zone; cil, ciliated zone; gr, granules.



Table 3. Ranked abundance of molluscs recorded from Transect 1. Lucinidae species are highlighted.

Species	Stations																		Total individuals		
	0A	0B	1A	1B	2A	2B	3A	3B	4A	4B	5A	5B	6A	6B	7A	7B	8A	8B		9A	9B
<i>Cerithideopsis cingulata</i>	-	-	-	8	115	22	325	4	255	82	96	88	42	41	12	94	14	12	27	28	1,265
<i>Pillucina vietnamica</i>	12	5	2	34	55	66	16	37	34	96	3	45	63	3	17	16	18	46	27	23	618
<i>Clithon oualaniensis</i>	-	-	3	2	27	58	12	5	16	19	8	6	79	9	4	1	22	16	6	2	295
<i>Cerithium coralium</i>	-	-	-	-	4	7	2	3	-	2	8	8	65	75	3	6	6	3	3	4	199
<i>Indoaustriella dalli</i>	2	-	3	1	6	7	3	2	2	8	4	12	14	3	9	2	4	6	9	1	98
<i>Pristis capsoides</i>	9	8	9	18	5	9	5	-	1	3	1	5	3	3	1	-	-	-	-	1	81
<i>Gafrarium tumidum</i>	-	1	3	7	5	5	4	1	-	1	-	1	-	1	-	-	-	-	-	-	29
<i>Anomalocardia squamosa</i>	1	1	1	3	3	3	3	3	1	1	1	2	1	1	2	-	1	-	-	-	28
<i>Nitidotellina nitens</i>	-	-	-	-	-	-	-	1	-	3	-	2	2	3	-	1	-	3	-	-	15
<i>Solen</i> sp.	-	-	-	-	1	-	-	1	-	-	1	-	1	-	-	-	2	-	1	-	7
<i>Plicarularia pullus</i>	-	-	-	1	-	-	1	-	-	-	1	-	2	-	-	-	-	1	-	-	6
<i>Tellinides</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	1	1	-	5
<i>Anadara troscheli</i>	-	-	-	-	-	-	-	-	-	-	-	-	1	3	-	-	1	-	-	-	5
<i>Anodontia bullata</i>	-	-	-	-	-	1	-	-	1	-	-	-	1	-	-	-	-	-	-	-	3
<i>Dosinia cretacea</i>	-	-	-	-	-	1	1	-	-	-	-	-	-	-	-	-	1	-	-	-	3
<i>Saccostrea</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	-	-	-	-	1	3
<i>Costellipitar</i> sp.	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	-	-	2
<i>Cerithidea alata</i>	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2
<i>Eucithara</i> sp.	-	-	-	-	-	-	-	-	-	1	-	-	-	1	-	-	-	-	-	-	2
<i>Pinguitellina pinguis</i>	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Tellinidae</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	1
<i>Meropesta pellucida</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	1
<i>Maetra</i> sp. (juv.)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	1
<i>Laternula truncata</i>	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1
<i>Cerithium</i> sp.	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Clypeomorus pellucida</i>	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1
Total species	4	6	6	8	10	10	10	9	7	11	9	11	12	13	8	7	9	9	9	7	26
Total individuals	24	18	21	74	222	179	372	57	310	217	123	171	274	147	49	121	69	89	76	60	2,673

Table 4. Ranked abundance of molluscs recorded from Transect 2. Lucinidae species are highlighted.

Species	Station						Total individuals
	1A	1B	2A	2B	3A	3B	
<i>Cerithideopsilla cingulata</i>	122	8	6	1	1	84	222
<i>Clithon oulaniensis</i>	1	28	9	6	4	38	86
<i>Cerithium coralium</i>	1	–	107	33	9	1	151
<i>Pillucina vietnamica</i>	–	2	17	15	74	30	138
<i>Indoaustriella dalli</i>	7	5	11	17	8	4	52
<i>Anomalocardia squamosa</i>	–	1	4	14	4	7	30
<i>Gafrarium tumidum</i>	3	6	12	6	1	1	29
<i>Pristis capsoides</i>	2	3	2	2	2	–	11
<i>Saccostrea</i> sp.	2	–	3	1	2	–	8
<i>Placamen calophyllum</i>	–	–	1	1	1	–	3
<i>Hebra corticata</i>	–	–	3	–	–	–	3
<i>Marcia hiantina</i>	–	–	–	2	–	–	2
<i>Nitidotellina nitens</i>	–	–	–	–	2	–	2
<i>Anadara troscheli</i>	–	–	1	1	–	–	2
<i>Cycladicama cumingi</i>	–	1	–	–	–	–	1
<i>Costellipitar</i> sp.	–	–	–	–	1	–	1
<i>Tellinides</i> sp.	–	–	–	–	–	1	1
<i>Solen</i> sp.	–	–	–	–	–	1	1
<i>Corbula</i> sp.	–	–	–	1	–	–	1
Total species	7	8	12	13	12	9	19
Total individuals	138	54	176	100	109	167	744

deposit-feeding tellinids, *Pristis capsoides* and *Nitidotellina nitens*. The fauna is unusual for the paucity of carnivorous gastropods, including only a few scavenging nassariids and the conoidean *Eucithara*. Although not recorded from the transect quadrats, the molluscivorous muricid gastropod *Chicoreus capucinus* (Lamarck, 1822) is frequent on the inner mudflat and around the mangrove prop roots (Tan, 2008).




Fig. 6. Distribution and abundance of the most common molluscs along Transect 1.




Fig. 7. Distribution of *Pillucina vietnamica* and *Indoaustriella dalli* across: A, Transect 1; B, Transect 2.

Table 5. Biomass estimates for Lucinidae species and other common bivalves.

Species	density (m <sup>-2</sup> ± SD)	range	mean size	dry wt (g·m <sup>-2</sup> )	range
<i>Pillucina vietnamica</i>	408 ± 242	112–858	5.3 ± 0.7	1.75	0.48–3.7
<i>Indoaustraliella dalli</i>	65 ± 32	13–112	5.9 ± 1.2	0.14	0.029–0.246
<i>Gafrarium tumidum</i>	19 ± 26	0–66	25.5 ± 9.3	2.66	–
<i>Pristis capsoides</i>	53 ± 57	0–178	16.9 ± 6.4	2.65	–
<i>Anadara troscheli</i>	7 ± 2	0–3	20.1 ± 10.3	0.98	–
<i>Anomalocardia squamosa</i>	18 ± 14	0–178	17.2 ± 7.4	0.65	–

## DISCUSSION

This study revealed in the intertidal mud and sand flats of Kungkrabaen Bay a high abundance of two small species of Lucinidae, *Pillucina vietnamica* and *Indoaustraliella dalli*. Despite the likely importance of these chemoautotrophic bivalves to the productivity of the mud flat, both of these species are poorly known with only basic taxonomic information available. Indeed *I. dalli* has hardly been mentioned since its original description (Lyngø, 1909). The habitat occupied by the lucinids of sheltered muddy bays backed by mangroves, with nutrient input from terrestrial runoff, is widespread in Southeast Asia and similar

abundances of small lucinids should occur at other localities in the region but are generally unreported.

Although the abundance of lucinids for Kungkrabaen Bay is high, it is not exceptional as shown by comparison with a few other studies (Table 6). There is little published data for the tropical Indo-Pacific; an exception is the high density (ca. 3,000 m<sup>-2</sup>) of *Pillucina* sp. (probably *Pillucina vietnamica*) reported from dugong-grazed seagrass beds in a sheltered bay on the Andaman Sea side of southwestern Thailand (Nakaoka et al., 2002). On Lizard Island, Queensland, *Divaricella irpex* (E. A. Smith, 1885), with densities of to 570 m<sup>-2</sup>, is the most abundant of several lucinids (Glover & Taylor, unpublished data). On an oceanic atoll, Paulay (2000) recorded *Ctena bella* (Conrad, 1834) and *Wallucina fijiensis* (E. A. Smith, 1885) at 40 and 15 m<sup>-2</sup> respectively. Just outside the tropics, at Rottnest Island, Western Australia, Barnes & Hickman (1999) recorded *W. assimilis* (Angas, 1867), which is of similar size (length 5.5–9.0 mm) to our Thailand species, at densities to 1,000 m<sup>-2</sup> in shallow, subtidal seagrass beds. Further afield, the highest densities of any lucinid recorded are for the Mediterranean *Loripes lacteus* (Linnaeus, 1758), to 2,600 m<sup>-2</sup> from *Posidonia* beds of Corsica (Johnson et al., 2002).

Although we recorded four species of lucinids from the muddy sands of Kungkrabaen Bay, *Ctena delicatula* occupied a rather different habitat amongst wave-washed, angular cobbles and gravel on an exposed rocky shore at the entrance to the bay. In our experience, *Ctena* species are often found in sand and gravel pockets on rock platforms. At first sight, this might seem an unsuitable habitat for a chemosymbiotic bivalve, but finer sediments and degrading organic material are often trapped in the quieter environment beneath the cobbles. We observed that *C. delicatula* from Kungkrabaen Bay has thinner gill filaments and smaller, fewer bacteriocytes (Fig. 5A) than the lucinids within the bay, which leads to speculation of perhaps lesser dependence on chemoautotrophy. *Ctena* species are amongst the most abundant and widespread of tropical lucinids but details of biology are known for only one species, *Ctena orbiculata* (Montagu, 1808), from the western Atlantic (Barnes, 1993).

By contrast to the abundant lucinids in sheltered Kungkrabaen Bay, nearby more exposed, intertidal, sandy beaches were sampled for bivalves and no Lucinidae were recorded. The habitat is both more mobile and unvegetated with less accumulation of organic material. The bivalve fauna of these




Fig. 8. Box plots of shell lengths of: A, *Pillucina vietnamica*; B, *Indoaustraliella dalli*, across Transect 1. Box limits are quartiles, black bars are means, grey bars are medians, whiskers extend 3/2 interquartile range, black circles are outliers and open circles extreme outliers.

Table 6. Some population density values for intertidal or shallow subtidal Lucinidae obtained from the literature.

Species	Habitat	Density (/m <sup>2</sup> )	Locality	Reference
<i>Pillucina</i> sp.	seagrass	3,000+	western Thailand	Nakaoka et al., 2002
<i>Ctena bella</i> (Conrad, 1837)	seagrass	40	Tawara Atoll	Paulay, 2000
<i>Wallucina fijiensis</i> (Smith, 1885) (as <i>W. haddoni</i> )	seagrass	15	Tawara Atoll	Paulay, 2000
<i>Divaricella irpex</i> (Smith, 1885)	sand, sparse seagrass	222–570	Lizard Island, Australia	Taylor & Glover, unpublished
<i>Cardiolucina pisiformis</i> (Thiele, 1930)	intertidal sand-mud	41	Dampier, W. Australia	Glover et al., 2003
<i>Wallucina assimilis</i> (Angas, 1867)	seagrass	718–1,048	Rottneest Is., W. Australia	Barnes & Hickman, 1999
<i>Codakia orbicularis</i> (Linnaeus, 1758)	seagrass	19–52	Jamaica Jackson,	1972
<i>Ctena orbiculata</i> (Montagu, 1808)	seagrass	13	Jamaica	Jackson, 1972
“ <i>Parvilucina</i> ” <i>costata</i> (d’Orbigny, 1842)	seagrass	15	Jamaica	Jackson, 1972
<i>Stewartia floridana</i> (Conrad, 1833)	seagrass	84	Florida, USA	Fisher & Hand, 1984
<i>Loripes lacteus</i> (Linnaeus, 1758)	seagrass	242–2,666	Corsica	Johnson et al., 2002
<i>Lucinoma borealis</i> (Linnaeus, 1776)	seagrass	120–1,500	Brittany, France	Monnat, 1970
<i>Lucinoma borealis</i> (Linnaeus, 1776)	sand and seagrass	3.8	Devon, England	Dando et al., 1986
<i>Lucinella divaricata</i> (Linnaeus, 1758)	seagrass	200–300	Brittany, France	Monnat, 1970
<i>Indoaustricella dalli</i> (Lyngé, 1909)	sand/mud, seagrass	13–112	Kungkrabaen Bay, Thailand	this study
<i>Pillucina vietnamica</i> Zorina, 1978	sand mud, seagrass	112–858	Kungkrabaen Bay, Thailand	this study

shores included *Meretrix meretrix* (Linnaeus, 1758), together with *Donax*, *Solen* and *Macra* species.

It should be emphasized that the intertidal habitats of Kungkrabaen Bay are highly disturbed. There is extensive and intensive artisanal shellfish exploitation, notably digging for the burrowing brachiopod *Lingula*, but also the gathering of bivalves including *Anomalocardia*, *Gafrarium*, *Anadara*, and *Psammotaena* (Fig. 2D). This foraging activity has likely taken place for a very long time and has undoubtedly




Fig. 9. Size classes of *Pillucina vietnamica* at stations across Transect 1.




Fig. 10. Feeding guilds of the common molluscs from transects at Kungkrabaen Bay. Origin of the food source is indicated by arrows.

disrupted seagrass growth and influenced the stability of the sediment surface. The lucinids are too small to be of any economic interest but the sediment disturbance and removal of many of the larger bivalves has likely influenced their abundance and distribution. Additionally, the whole bay is influenced by the extensive disturbance and drainage from the massive development of fish and prawn farms (Fig. 1) in the landward parts of the mangrove fringe (Tookwinas, 1998). However, results from isotope studies suggest that effluent nutrients from the prawn farms are largely trapped by the mangroves and in the inner bay sources of particulate organic material are mainly from seagrass, algae, and plankton (Thimdee et al., 2003).

As exemplified by Kungkrabaen Bay, chemoautotrophic lucinid bivalves are often abundant in many tropical shallow-water habitats such as seagrass beds and peri-mangrove habitats. They represent an ecologically important component of these systems in terms of their biomass, production, and cycling of buried organic material via sulphide-oxidising symbiotic bacteria (Johnson et al., 2002). However, details of their role in the productivity of these systems have been little studied or altogether ignored. Past studies have often recognized the abundance of lucinids in seagrass communities (Moore et al., 1968; Taylor & Lewis, 1970; Jackson 1972) but classified the bivalves as particulate suspension feeders and misinterpreted their role in the communities.

#### ACKNOWLEDGMENTS

The International Marine Bivalve Workshop (with contributions on other molluscan groups) in Chantaburi, Thailand, was organized by Kashane Chalermwat (Burapha University), Fred Wells (Western Australian Department of Fisheries), Rüdiger Bieler (Field Museum of Natural History, Chicago) and Paula M. Mikkelsen (American Museum of Natural History), and supported by U.S. National Science Foundation grant PEET DEB-9978119 (to RB and PMM). Field transportation in Thailand and chemicals were provided by the Faculty of Science, Burapha University. Our thanks to them all.

We thank Nopadon Kakhai for local advice, along with Graham Oliver, Paul Valentich-Scott, Rüdiger Bieler, Paula Mikkelsen and Tan Koh Siang for valuable discussions in the field and laboratory. Roger Bamber (NHM) is thanked for arranging the sediment analysis, Paul Callomon (Academy of Natural Sciences, Philadelphia) for images of the syntypes of *Ctena delicatula*, and Alex Ball (NHM Electron Microscope Unit) for much advice.

EAG and JDT are indebted to Phil Rainbow and the Department of Zoology, NHM, for continuing support and facilities.

#### THAI ABSTRACT

จากการสำรวจเชิงปริมาณเกี่ยวกับมอลลัสก์ที่อาศัยอยู่ในพื้นที่ที่ได้รับผลกระทบจากกิจกรรมของมนุษย์ บริเวณพื้นที่ทรายนโคลนในแนวน้ำขึ้นน้ำลงใกล้ป่าชายเลนในอ่าวที่มีคลื่นลมสงบบริเวณภาคตะวันออกเฉียงใต้ของประเทศไทย พบว่ามีความหนาแน่นของหอยสองฝาวงศ์ลูซินิดีขนาดเล็กที่มีแบคทีเรียอาศัยอยู่ในเหงือกแบบ chemosymbiotic ชนิด *Pillucina vietnamica* และ *Indoaustriella dalli* อยู่สูง (1,380 ตัว/ตารางเมตร) ส่วนหอยสองฝาในวงศ์เดียวกันอีกสองชนิดคือ *Anodontia bulla* และ *A. philippiana* ก็มีพบอยู่บ้างในบริเวณใกล้เคียงที่ทำการสำรวจ ในขณะที่หอยลูซินิดีอีกชนิดหนึ่งคือ *Ctena delicatula* จะพบอาศัยอยู่ในพื้นที่ๆ เป็นกรวดบริเวณปากทางเข้าของอ่าวกึ่งกระเบนจากการศึกษาครั้งนี้พบว่าหอยในวงศ์ลูซินิดีทุกชนิดที่สำรวจพบมีแบคทีเรียที่เป็นประโยชน์อาศัยอยู่ในเหงือก ส่วนหอยชนิดอื่นๆ ที่พบว่าอาศัยอยู่ร่วมกับหอยสองฝาคือ *Cerithideopsis cingulata*, *Cerithium coralium* และ *Clithon oualaniensis* ที่เป็นหอยฝาเดียวประเภทที่อาศัยบนพื้นผิว และ พบหอยสองฝาที่ฝังตัวอยู่ในพื้นได้แก่ *Gafrarium tumidum*, *Anomalocardia squamosa*, *Pristis capsoides* และ *Anadara troscheli* จากการประมาณการชีวมวลพบว่าชีวมวลของหอยลูซินิดีมีค่าระหว่าง 0.51 ถึง 3.95 กรัม/ตารางเมตร ผลของการศึกษานี้ชี้ให้เห็นถึงบทบาททางนิเวศวิทยาที่สำคัญของหอยที่มีการดำรงชีวิตแบบ chemoautotrophic ที่อาศัยอยู่ในบริเวณขอบของป่าชายเลนด้านที่ติดกับทะเล

#### LITERATURE CITED

- Adams, A., 1852. Catalogue of the species of *Nassa*, a genus of gasteropodous Mollusca belonging to the family Buccinidae, in the collection of H. Cuming Esq., with the description of some new species. *Proceedings of the Zoological Society of London*, **19**: 94–112.
- Angas, G. F., 1867. Descriptions of thirty-two new species of marine shells from the coast of New South Wales. *Proceedings of the Zoological Society of London*, **1867**: 110–117.
- Barnes, P. A. G., 1993. *Eco-physiology of the Endosymbiont-bearing Lucinid Bivalve, Codakia orbiculata*. Ph. D. Thesis, University of Plymouth, United Kingdom. 359 pp.
- Barnes, P. A. G. & C. S. Hickman, 1999. Lucinid bivalves and marine angiosperms: a search for causal relationships. In: Walker, D. I. & F. E. Wells (eds.), *The Seagrass Flora and Fauna of Rottnest Island, Western Australia*. Western Australian Museum, Perth. Pp. 215–238.
- Conrad, T. A., 1833. On some new fossil and Recent shells of the United States. *American Journal of Science, series 1*, **23**: 339–346.
- Conrad, T. A., 1837. Descriptions of new marine shells, from Upper California. Collected by Thomas Nuttall Esq. *Journal Academy of Natural Sciences, Philadelphia*, **7**(2): 227–268.

- Dando, P. R., A. J. Southward & E. C. Southward, 1986. Chemoautotrophic symbionts in the gills of the bivalve mollusc *Lucinoma borealis* and the sediment chemistry of its habitat. *Proceedings of the Royal Society of London B*, **227**: 227–247.
- Dando, P. R., S. A. Ridgway & B. Spiro, 1994. Sulphide ‘mining’ by lucinid bivalve molluscs: demonstrated by stable sulphur isotope measurements and experimental models. *Marine Ecology Progress Series*, **107**: 169–175.
- Deshayes, G. P., 1855 (“1854”). Descriptions of new shells from the collection of Hugh Cuming, Esq. *Proceedings of the Zoological Society of London*, **1854**(22): 317–320; (280): 321–336; (281): 337–352; (282): 353–368; (283): 369–371.
- Dillwyn, L. W., 1817. *A Descriptive Catalogue of Recent Shells Arranged According to the Linnean Method; with Particular Attention to the Synonymy*, 2 Volumes. J. & A. Arch, London. 1092 pp.
- Distel, D. L., 1998. Evolution of chemoautotrophic endosymbioses in bivalves. *Bioscience*, **48**: 277–286.
- Dunker, W. [G.] B. R. H., 1882. *Novitates Conchologicae. Abbildung und Beschreibung neuer Conchylien. Supplement 7. Index molluscorum maris Japonici*. Fischer, Cassel. vii + 301 pp., 16 pls.
- Fenchel, T. & R. J. Riedl, 1970. The sulfide system: a new biotic community underneath the oxidized layer of marine sand bottoms. *Marine Biology*, **7**: 59–68.
- Fisher, C. R., 1990. Chemoautotrophic and methanotrophic symbioses in marine invertebrates. *Reviews in Aquatic Sciences*, **2**: 399–436.
- Fisher, C. R. & S. C. Hand, 1984. Chemoautotrophic symbionts in the bivalve *Lucina floridana* from seagrass beds. *Biological Bulletin*, **167**: 445–459.
- Frenkiel, L. & M. Mouëza, 1995. Gill ultrastructure and symbiotic bacteria in *Codakia orbicularis* (Bivalvia, Lucinidae). *Zoomorphology*, **115**: 51–61.
- Frenkiel, L., O. Gros & M. Mouëza, 1996. Gill structure in *Lucina pectinata* (Bivalvia: Lucinidae) with reference to hemoglobin in bivalves with symbiotic sulphur-oxidising bacteria. *Marine Biology*, **125**: 511–524.
- Glover, E. A. & J. D. Taylor, 2001. Systematic revision of Australian and Indo-Pacific Lucinidae (Mollusca: Bivalvia): *Pillucina*, *Wallucina* and descriptions of two new genera and four new species. *Records of the Australian Museum*, **53**: 263–292.
- Glover, E. A., J. D. Taylor & J. Whittaker, 2003. Distribution, abundance and foraminiferal diet of an intertidal scaphopod, *Laevidentalium lubricatum*, around the Burrup Peninsula, Dampier, Western Australia. In: Wells, F. E., D. I. Walker & D. S. Jones (eds.) *The Marine Fauna and Flora of Dampier, Western Australia*. Western Australian Museum, Perth. Pp. 225–240.
- Glover, E. A., J. D. Taylor & S. T. Williams, 2008. Mangrove associated lucinid bivalves of the central Indo-West Pacific: review of the “*Austriella*” group with a new genus and species. *Raffles Bulletin of Zoology*, Supplement **18**: 25–40.
- Gmelin, J. F., 1791. *Caroli a Linné ... Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis . . . Editio decima tertia, aucta, reformata. Volume 1, Part 6*. G. E. Beer, Lipsiae [Leipzig]. Pp. 3021–4120.
- Gros, O., M. Liberge & H. Felbeck, 2003. Interspecific infection of aposymbiotic juveniles of *Codakia orbicularis* by various tropical lucinid gill-endosymbionts. *Marine Biology*, **142**: 57–66.
- Hanley, S., 1842–1856. *An Illustrated and Descriptive Catalogue of Recent Bivalve Shells*. Williams & Norgate, London. 392 pp.
- Hanley, S., 1844. Descriptions of new species of *Tellina* collected by H. Cuming Esq. . *Proceedings of the Zoological Society of London*, **12**(134): 59–64.
- Hanley, S., 1845 (“1844”). Descriptions of new species of *Cyrena*, *Venus* and *Amphidesma*. *Proceedings of the Zoological Society of London*, **12**(140): 159–162.
- Hombron, M. M. & H. Jacquinot, 1852. Mollusques. In: Dumont d’Urville (ed.), *Voyage au Pole Sud et dans ‘Océanie sur les Corvettes l’Astrolabe et la Zélée; Exécute par Ordre du Roi Pendant les Années 1837–1838–1839–1840; Atlas*. Gide et J. Baudry, Paris. 27 pls.
- Jackson, J. B. C., 1972. The ecology of molluscs of *Thalassia* communities, Jamaica, West Indies. II. Molluscan population variability along an environmental stress gradient. *Marine Biology*, **14**: 304–337.
- Johnson, M. A. & C. Fernandez, 2001. Bacterial symbiosis in *Loripes lucinalis* (Mollusca: Bivalvia) with comments on reproductive strategy. *Journal of the Marine Biological Association of the United Kingdom*, **81**: 251–257.
- Johnson, M. A. C. Fernandez & G. Pergent, 2002. The ecological importance of an invertebrate chemoautotrophic symbiosis to phanerogam seagrass beds. *Bulletin of Marine Science*, **71**: 1343–1351.
- Kamimura, S. & M. Tsuchiya, 2004. The effect of feeding behaviour of the gastropods *Batillaria zonalis* and *Cerithideopsisilla cingulata* on their ambient environment. *Marine Biology*, **144**: 705–712.
- Kiener, L. C., 1841–1842. *Species Général et Iconographie des Coquilles Vivants, Comprenant la Collection du Muséum d’Histoire Naturelle de Paris, la Collection Lamarck, celle du Prince Masséna (Appartenant Maintenant à M. le Baron Benjamin Delessert), et les Découvertes Récentes des Voyageurs. Genere Cerite*. **5**: 1–102. Rousseau, Paris.
- Lamarck, J. B. P. A. de M. de, 1818. *Histoire Naturelle des Animaux sans Vertèbres, Présentant les Caractères Généraux et Particuliers de ces Animaux, leur Distribution, leurs Classes, leurs Familles, leurs Genres, et la Citation des Principales Espèces qui s’y Rapportent... Volume 5*. Deterville, Verdier, Paris. [iii] + 612 pp.
- Leбата, M. J. H. L. & J. H. Primavera, 2001. Gill structure, anatomy and habitat of *Anodontia edentula*; evidence of endosymbiosis. *Journal of Shellfish Research*, **20**: 1273–1278.
- Lesson, R. P., 1831. *Voyage Autour du Monde sur la Coquille, Pendant 1822–1825 par M. L. J Duperry. Volume 2 Zoophytes*. Paris. 128 pp.
- Linnaeus, C., 1758. *Systema Naturae per Regna Tria Naturae. Tomus I. Editio Decima, Reformata*. Laurentii Salvii, Stockholm. [iii] + 824 pp.
- Lynge, H., 1909. The Danish Expedition to Siam, 1899–1900. IV. Marine Lamellibranchiata. *Kongelige Danske Videnskabernes Selskabs Skrifter, 7. Raekke, Naturvidenskabelig og Mathematisk Afdeling*, Kjøbenhavn, **5**(3): 97–299, pls 1–5.
- Monnat, J. Y., 1970. Introduction à l’étude de la reproduction chez *Lucinoma borealis* (Linné), Bivalvia, Lucinacea. These 3e cycle, Faculté de Science, Brest [not seen; cited by Johnson et al., 2002].

- Moore, H. B., L. T. Davies, T. H. Fraser, R. H. Gore & N. R. Lopez, 1968. Some biomass figures from a tidal flat in Biscayne Bay, Florida. *Bulletin of Marine Science*, **18**: 261–279.
- Montagu, G. 1808. *Testacea Britannica, Supplement*. J. White, London. 183 pp.
- Nakaoka, M., H. Mukai & S. Chunhabundit, 2002. Impacts of dugong foraging on benthic animal communities in a Thailand seagrass bed. *Ecological Research*, **17**: 625–638.
- Oliver, P. G. & A. M. Holmes, 2006. A new species of *Lucinoma* (Bivalvia: Lucinoidea) from the oxygen minimum zone of the Oman Margin, Arabian Sea. *Journal of Conchology*, **39**: 63–77.
- Orbigny, A. d', 1839–1842. Mollusques, Volume 7 and Atlas. In: Sagra, R. de la, *Histoire Physique, Politique et Naturelle de l'Île de Cuba*. Arthus Bertrand, Paris. 380 pp., 28 pls.
- Ott, J., M. Bright & S. Bulgheresi, 2004. Symbioses between marine nematodes and sulfur-oxidising bacteria. *Symbiosis*, **36**: 103–126.
- Paulay, G., 2000. Benthic ecology and biota of Tarawa Atoll lagoon: influence of equatorial upwelling, circulation and human harvest. *Atoll Research Bulletin*, **487**: 1–41.
- Philippi, R. A., 1836. Beschreibung einiger neuen Conchylien-Arten und Bemerkungen über die Gattung *Lacuna* von Turton. *Archiv für Naturgeschichte*, **2**(1): 224–235, Pls. 7–8.
- Philippi, R. A., 1849. *Cerithium*, Tab. I. *Abbildungen und Beschreibungen neuer oder wenig gekannter Conchylien*, **3**(4): 13–20, Pl. 1.
- Pilsbry, H. A., 1904. New Japanese marine Mollusca; Pelecypoda. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **57**: 101–122.
- Prashad, B., 1932. The Lamellibranchia of the Siboga Expedition. Systematic Part. II Pelecypoda (exclusive of the Pectinidae). *Siboga-Expeditie, Uitkomsten op Zoologisch, Botanisch, Oceanographisch en Geologisch Gebeid Verzameld in Nederlandsch Oost-Indië 1899–1900*, **53c**: 1–353, 59 pls.
- Reeve, L. A., 1850a. Monograph of the genus *Artemis*. *Conchologica Iconica; or Illustrations of the Shells of Molluscos Animals*, **6**: 10 pls.
- Reeve, L. A., 1850b. Monograph of the genus *Lucina*. *Conchologica Iconica; or Illustrations of the Shells of Molluscos Animals*, **6**: 11 pls.
- Reid, R. G. B., 1990. Evolutionary implications of sulphide-oxidising symbioses in bivalves. In: Morton, B. (ed.), *The Bivalvia – Proceedings of a Memorial Symposium in Honour of Sir Charles Maurice Yonge, Edinburgh, 1986*. Hong Kong University Press, Hong Kong. Pp. 127–140.
- Reid, R. G. B. & D. G. Brand, 1986. Sulfide-oxidising symbiosis in lucinaceans: implications for bivalve evolution. *The Veliger*, **29**: 3–24.
- Röding, P. F., 1798. *Museum Boltenianum, Pars Secunda Continens Conchlia, sive Testacea Univalvia, Bivalvia et Multivalvia*. Johan Christi Trappii, Hamburg. viii + 199 pp.
- Smith, E. A., 1885. Report on the Lamellibranchiata collected by H. M. S. Challenger during the years 1873–76. *Reports of the Scientific Results of the Exploratory Voyage of H. M. S. Challenger 1873–76, Zoology*, **13**(35): 1–341, 25 pls.
- Stanley, S. M., 1970. Relation of shell form to life habits of the Bivalvia (Mollusca). *The Geological Society of America Memoirs*, **125**: 1–296.
- Stewart, F. J., I. L. G. Newton, & C. M. Cavanaugh, 2005. Chemosymbiotic endosymbioses: adaptations to oxic-anoxic interfaces. *Trends in Microbiology*, **13**: 439–448.
- Tan, K. S., 2008. Mudflat predation on bivalves and gastropods by *Chicoreus capucinus* (Neogastropoda: Muricidae) at Ao Kungkrabaen, Gulf of Thailand. *Raffles Bulletin of Zoology, Supplement* **18**: 235–245.
- Taylor, J. D. & E. A. Glover, 2000. Functional anatomy, chemosymbiosis and evolution of the Lucinidae. In: Harper, E. M., J. D. Taylor & J. A. Crame (eds.), *The Evolutionary Biology of the Bivalvia. Geological Society of London Special Publication*, **177**: 207–225.
- Taylor, J. D. & E. A. Glover, 2005. Cryptic diversity of chemosymbiotic bivalves: a systematic revision of worldwide *Anodontia* (Mollusca: Bivalvia: Lucinidae). *Systematics and Biodiversity*, **3**: 281–338.
- Taylor, J. D. & E. A. Glover, 2006. Lucinidae – the most diverse group of chemosymbiotic molluscs. *Zoological Journal of the Linnean Society*, **148**: 421–438.
- Taylor, J. D. & M. S. Lewis, 1970. The flora, fauna and sediments of the marine grass beds of Mahé, Seychelles. *Journal of Natural History*, **4**: 199–220.
- Thiele, J., 1930. Gastropoda und Bivalvia. In: Michaelson, W. & R. Hartmeyer (eds.), *Die Fauna Südwestaustraliens*. Gustav Fischer, Jena. Pp. 561–596.
- Thimdee, W., G. Deein, C. Sangrungruang, J. Nishuoka & K. Matsunaga, 2003. Sources and fate of organic matter in Khung Krabaen Bay (Thailand) as traced by  $\delta C^{13}$  and C/N ratios. *Wetlands*, **23**: 729–738.
- Tookwinas, S., 1998. The environmental impact of marine shrimp farming effluents and carrying capacity estimation at Kung Krabaen Bay, eastern Thailand. *Asian Fisheries Science*, **11**: 303–316.
- Zorina, I. P., 1978. New species of bivalve molluscs (Bivalvia) of the Gulf of Tonkin (South China Sea). *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR, Leningrad*, **61**: 193–203. [in Russian]