

Grapsoid crabs (Decapoda: Brachyura: Thoracotremata) of the Persian Gulf and the Gulf of Oman

REZA NADERLOO

Marine Zoology, Forschungsinstitut Senckenberg, Senckenberganlage 25, 60325 Frankfurt am Main, Germany and School of Biology, College of Science, University of Tehran, Tehran, Iran. E-mail: rnaderloo@senckenberg.de, naderlo@khayam.ut.ac.ir

Table of contents

Abstract	1
Introduction	2
Taxonomy	2
Family Grapsidae MacLeay, 1838	2
<i>Grapsus albolineatus</i> Latreille in Milbert, 1812	2
<i>Metopograpsus messor</i> (Forskål, 1775)	8
<i>Metopograpsus thukuhar</i> (Owen, 1839)	11
Family Sesarmidae Dana, 1851	15
<i>Chiromantes boulengeri</i> (Calman, 1920)	15
<i>Nanosesarma sarii</i> Naderloo & Türkay, 2009	18
<i>Nanosesarma jousseaumei</i> (Nobili, 1905)	22
<i>Parasesarma persicum</i> Naderloo & Schubart, 2010	26
<i>Perisesarma guttatum</i> (A. Milne-Edwards, 1869)	29
Family Varunidae Alcock, 1900	31
<i>Metaplaax indica</i> H. Milne-Edwards, 1852	31
<i>Thalassograpsus harpax</i> (Hilgendorf, 1892)	34
Family Plagusiidae Dana, 1851	36
<i>Plagusia squamosa</i> (Herbst, 1790)	36
Acknowledgements	40
References	40

Abstract

Grapsoid crabs, together with ocypodoids, are the two most important groups of intertidal brachyurans in the Persian Gulf and the Gulf of Oman. In total, four families, nine genera and 12 species of grapsoids are known from the region. *Perisesarma guttatum* (A. Milne-Edwards, 1869), and *Plagusia squamosa* (Herbst, 1790), are only recorded from the Gulf of Oman; *Grapsus granulosus* H. Milne-Edwards, 1853, and *Thalassograpsus harpax* (Hilgendorf, 1892), are collected for the first time from the Persian Gulf. *Chiromantes boulengeri* (Calman, 1920), *Nanosesarma sarii* Naderloo & Türkay, 2009, and *Parasesarma persicum* Naderloo & Schubart, 2010, are so far known only from the region.

Key words: Crustacea, Brachyura, Grapoidea, Grapsidae, Sesarmidae, Varunidae, Plagusidae, Persian Gulf, Gulf of Oman

Introduction

Several ecological and faunistic studies undertaken along the Arabian coastal region considered grapsoid crabs to be the dominant brachyuran group in the region (e.g., Basson *et al.* 1977; Clayton 1986; Jones 1986a; Al-Khayat & Jones 1996; 1999; Apel 1994a; 1996; Al-Ghais & Cooper 1996; Cooper 1997; Hornby 1997). *Metopograpsus messor* (Forskål, 1775), was first recorded from Bahrain and UAE by Nobili (1906a) and Calman (1920) briefly described a new sesarmid species (*Sesarma boulengeri*) from Basra, Iraq. Stephensen (1945) in his comprehensive work on the brachyuran fauna of the Persian Gulf recorded two more grapsoid species, *M. messor* and *Grapsus maculatus* subsp. *tenuicrustatus* Tesch, 1918, the latter being identified as *Grapsus albolineatus* Latreille in Milbert, 1812, by Apel & Türkay (1999). Apel & Türkay (1999) reviewed the Ocypodoidea and Grapoidea of the Per-