

Article

Diversity, Distribution, and Habitat Occurrence of the Diaptomid Copepods (Crustacea: Copepoda: Diaptomidae) in Freshwater Ecosystems of Thailand

Laorsri Sanoamuang ^{1,2,*}  and Prapatsorn Dabseepai ¹

¹ Applied Taxonomic Research Center, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand; prapda@kku.ac.th

² Department of Biodiversity and Environmental Management, International College, Khon Kaen University, Khon Kaen 40002, Thailand

* Correspondence: la_orsri@kku.ac.th

Abstract: The diversity, distribution, and checklist of diaptomid copepods from various freshwater ecosystems throughout Thailand are presented, based on data from our biodiversity projects during 1993–2019 and literature reviews. Thailand has one of the most diversified diaptomid fauna in the world, with 42 species identified from 2150 localities (4962 samples). *Mongolodiaptomus* has the highest species richness with ten species, followed by *Tropodiaptomus* with seven species. Among these, eight taxa (*Arctodiaptomus* sp., *Mongolodiaptomus pectinidactylus*, *Neodiaptomus meggitti*, *Tropodiaptomus hebereri*, *T. lanaonus*, *T. oryzanus*, *T. ruttneri*, and *Paradiaptomus greeni*) are new to the fauna of Thailand. The rare *P. greeni*, which predominantly occurs in Africa, is also a new record for Southeast Asia. The most frequently encountered species were *Mongolodiaptomus botulifer*, *Phyllodiaptomus praedictus*, *M. calcarus*, *M. dumonti*, *M. malaindosinensis*, *Vietodiaptomus blachei*, *Phyllodiaptomus christineae*, *Eodiaptomus sanoamuangae*, *Neodiaptomus yangtsekiangensis*, *E. draconisignivomi*, *T. vicinus*, and *Heliodiaptomus elegans*. Twelve species appear to be endemic to Thailand, and eight species occur only in the countries belonging to the lower Mekong River Basin (Thailand, Cambodia, Laos, and Vietnam). The uniqueness of the Thai diaptomids is a high degree of co-occurrence of four to seven species in the same localities. Comments on the taxonomic status of the species recorded are provided. In addition, the taxonomic validity of *Mongolodiaptomus malaindosinensis* is discussed.



Citation: Sanoamuang, L.; Dabseepai, P. Diversity, Distribution, and Habitat Occurrence of the Diaptomid Copepods (Crustacea: Copepoda: Diaptomidae) in Freshwater Ecosystems of Thailand. *Water* **2021**, *13*, 2381. <https://doi.org/10.3390/w13172381>

Academic Editors: Kay Van Damme and Alexey A. Kotov

Received: 13 July 2021

Accepted: 23 August 2021

Published: 30 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: *Arctodiaptomus* sp.; biodiversity; Calanoida; lower Mekong River Basin; *Mongolodiaptomus malaindosinensis*; *Paradiaptomus greeni*; Southeast Asia

1. Introduction

Copepods are among the most diverse micro-crustaceans in aquatic habitats [1]. They are abundant in freshwater ecosystems and can be a major component of most planktonic, benthic, and groundwater communities. Members of the family Diaptomidae in the order Calanoida are particularly successful in all kinds of freshwater habitats, comprising over 440 species [2]. Of the 22 genera (92 species) of freshwater diaptomids reported from inland waters of the Oriental biogeographic region, 10 genera (81 species) are endemic to this region [3].

Studies of the freshwater diaptomid copepods in Southeast Asia prior to 1980 were fragmentary. Of the 30 calanoid species, 27 diaptomid species were reported from Southeast Asia [3]. Most of the studies were conducted in Thailand since 1994 [4–22], while a few investigations have been documented in Cambodia [20,21,23], Laos [16,20,24], Malaysia [25–27], Singapore [28], the Philippines [29,30], and Indonesia [31]. A few studies were conducted in Vietnam [20], but most of the publications were published in Vietnamese with an English abstract [32].

In Thailand, Bricker et al. (1978) reported 7 diaptomid species from 11 reservoirs and one natural swampy lake in Thailand [33]. Later, a list of 14 species based on extensive collections sampled throughout the country was documented by Lai and Fernando (1981) [34]. In 1984, Boonsom (1984) examined samples from 39 different kinds of habitats, including rivers, reservoirs, irrigation tanks, ponds, and fish fields, revealing 8 diaptomid species [35]. In 1994–1998, four new species were described based on specimens from Thailand [4–7]. At the same time, Ranga Reddy et al. (1998) [7] published a list of 17 valid species for the country, and 16 of these were reported from the northeast region by Sanoamuang (1999) [9] as well. The misidentifications and taxonomic confusions of two species (*Neodiaptomus mephistopheles* Brehm, 1933 and *Arctodiaptomus bacillifer* (Koelbel, 1885)) [3,33,34,36] by previous researchers and the doubtful records of two taxa, *Tropodiaptomus doriai* (Richard, 1894) and *Sinodiaptomus chaffanjoni* (Richard, 1897), were corrected by Ranga Reddy et al. (1998) [7].

Consequently, there has been a gradual increase in our knowledge of the diversity of diaptomid copepods in Thailand. Since 2001, 13 more new species have been added to the list of Thailand [10–22]. As part of the authors' long-term intensive collections of freshwater copepods from different habitats all over Thailand during 1993–2019 and from the literature review, an updated list of diaptomid copepods known to date for the country is presented in this contribution. Their distributions within the six regions of Thailand, habitat occurrence, and taxonomic status of the diaptomid taxa are presented here.

2. Materials and Methods

2.1. Study Area

Thailand is situated in the heart of the Southeast Asian mainland, covering an area of 513,115 square kilometers. It lies in the humid tropics, having a monsoon climate with three distinct seasons—hot (March to April), rainy or monsoon (May to October), and cool (November to February). The country is officially divided into six geographical regions (Figure 1A) [36]. The northern region (with 9 provinces) is mountainous and was traditionally the most heavily forested area of the country. There are four main river basins (Ping, Wang, Yom, and Nan) in this region (Figure 1B). The northeast region (with 20 provinces) is some 200–300 m above sea level. This region is hemmed in by mountain ridges on the west and the south. This area is drained by the Song Kham, Chi, and Mun rivers, which are tributaries of the Mekong (Figure 1B). The central region (with 22 provinces) comprises the basins of the Chao Phraya and Pa Sak Rivers, which run from north to south and drain into the Gulf of Thailand. The western region (with 5 provinces) is also mountainous, alternating with plains along narrow valleys. The eastern region (with 7 provinces) is characterized by short mountain ranges and fruit plantations are a major component of agriculture in this area. The southern region (with 14 provinces) has the highest amount of rainfall in the country. It is the principal rubber-growing area and contains extensive alluvial deposits of tin.

2.2. Sample Collections and Identifications

Qualitative copepod samples of the pelagic and littoral zones from different freshwater habitats in Thailand were collected during 1993–2019. Information on 18 biodiversity projects in the surveyed water bodies is given in Table 1. A total of 4962 samples were collected from 2150 localities in 6 geographical regions (north, northeast, central, east, west, and south) of Thailand using a plankton net with a mesh size of 60 µm. In each site, sampling was performed either in two (dry and rainy) or three (summer, rainy, and win-ter) seasons, although some were carried out only once. A variety of water bodies, including permanent water habitats—canals, lakes, ponds, swamps, reservoirs, rivers—and temporary water habitats—rice fields, roadside canals, and temporary ponds—were sampled. The animals were preserved on-site in 4% formaldehyde or 70% ethanol. Adult copepods were selected, dissected, and prepared on a glycerin-mounted slide under an

Olympus SZ51 stereomicroscope at 40–100 \times magnification. All un-dissected specimens were stored in 70% ethanol in 1.5 mL microtubes.




Figure 1. The map of Thailand displays six geographical regions (A) and major rivers (B).

Table 1. Information on sampling projects of the surveyed waterbodies in Thailand. The regions in Thailand are given by; N = North, NE = Northeast, C = Central, E = East, W = West, and S = South.

No.	Sampling Habitats	Regions in Thailand	Sampling Years	Number of Localities	Number of Samples
1	Various habitats in Northeast Thailand	NE	1993	93	200
2	Various habitats in Nakhon Ratchasima and Surin Provinces	NE	1996–1999	138	412
3	Man-made reservoirs in the Phuphan National Park	NE	1997–1998	16	336
4	Major rivers in Thailand	N,NE,C,E,W,S	1998–1999	20	60
5	Various habitats throughout Thailand	N,NE,C,E,W,S	1998–2004	294	508
6	Temporary-water habitats in seven provinces of NE Thailand	NE	1999–2003	521	844
7	Various habitats in the floodplain of the Rivers Nan, Chi, Mun and Song Khram	N,NE,C	2000–2004	186	372

Table 1. Cont.

No.	Sampling Habitats	Regions in Thailand	Sampling Years	Number of Localities	Number of Samples
8	Various habitats in Central and Northern Thailand	N,C	2001–2002	120	360
9	Various habitats in Central and Northern Thailand	N,C	2003	55	110
10	Three natural lakes (Buemg Boraphet, Bueng Khong Long, and Lake Kud-Thing)	NE,C	2002–2004	30	60
11	Various habitats in Eastern Thailand	E	2003–2004	55	110
12	Various habitats in Sisaket Province	NE	2006–2007	99	190
13	Various habitats in Suphan Buri, Kanchanaburi, Ratchaburi and Phetchaburi Provinces	W	2007	200	218
14	Various habitats in Nam Nao National Park, Phetchabun Province	C	2007–2009	10	75
15	Lake Kud-Thing and various habitats in Bueng Kan and Udon Thani Provinces	NE	2012–2014	97	408
16	Various habitats in Chiang Rai, Phayao and Nan Provinces	N	2014–2016	67	154
17	Various habitats in Khon Kaen, Kalasin and Buriram Provinces	NE	2014–2018	128	344
18	Various habitats in Saraburi and Lopburi Provinces	C	2019	21	201
Total				2150	4962

All appendages and body ornamentation were examined at 1000- \times magnification under an Olympus CX31 compound microscope. Specimens for scanning electron microscopy (SEM) were dehydrated in an ethanol series (50, 70, 80, 90, 95, 100, and 100%) for 15 min at each concentration. Specimens were dried in a critical-point dryer and were mounted on stubs using adhesive tape under a stereomicroscope. Dried specimens were coated with gold in a sputter coater. The SEM photographs were taken using a scanning electron microscope (LEO, 1450VP).

The abbreviations used in this paper are as follows:

Exp-2 = second exopodal segment;

P5 = leg 5.

2.3. Data Analysis

The relative occurrences of the diaptomid copepods recorded from Thailand are categorized according to the occurrence frequencies of the local populations into five categories as follows: (1) common = species that occurs in more than 45% of the sampled localities; (2) fairly common = species that occurs in 31–45% of the sampled localities; (3) uncommon = species that occurs in 16–30% of the sampled localities; (4) rare = species that occurs in 5–15% of the sampled localities; (5) extremely rare = species that occurs in less than 5% of the sampled localities.

3. Results

3.1. Species Diversity

A list of diaptomid species identified from the samples examined, along with their habitat types, relative occurrences, and distributional regions in Thailand, is provided in

Table 2. Of the 42 species identified, 41 species are members of the subfamily Diaptominae, while only 1 belongs to the subfamily Paradiaptominae. The diversity of species among the 11 genera of the Diaptomidae is dominated by the genus *Mongolodiaptomus*, with 10 known species, followed by *Tropodiaptomus* (7 species), *Neodiaptomus* (6 species) and *Phyllodiaptomus* (6 species). Among these, 17 species (40.5% of the species recorded) have been originally described from specimens from Thailand since 1994, including (1) *Phyllodiaptomus* (*Ctenodiaptomus*) *praedictus* Dumont and Ranga Reddy, 1994 (Figure 2A); (2) *Phyllodiaptomus* (*Phyllodiaptomus*) *christineae* Dumont, Ranga Reddy, and Sanoamuang, 1996 (Figure 2B); (3) *Eodiaptomus sanoamuangae* Ranga Reddy and Dumont, 1998 (Figure 3A); (4) *Mongolodiaptomus rarus* (Ranga Reddy, Sanoamuang, and Dumont, 1998); (5) *Phyllodiaptomus* (*Ctenodiaptomus*) *surinensis* Sanoamuang and Yindee, 2001; (6) *Eodiaptomus phuphanensis* Sanoamuang, 2001 (Figure 3B); (7) *Mongolodiaptomus dumonti* Sanoamuang, 2001 (Figure 4A); (8) *Neodiaptomus songkhramensis* Sanoamuang and Athi-bai, 2002; (9) *Heliodiaptomus phuthaiorum* Sanoamuang, 2004 (Figure 5A); (10) *Eodiaptomus phuvongi* Sanoamuang and Sivongxay, 2005 (Figure 3C); (11) *Phyllodiaptomus* (*Phyllodiaptomus*) *thailandicus* Sanoamuang and Teeramaethee, 2006 (Figure 2C); (12) *Neodiaptomus siamensis* Proogkiat and Sanoamuang, 2008; (13) *Mongolodiaptomus loeiensis* Watiroyram and Sanoamuang, 2017 (Figure 4B); (14) *Mongolodiaptomus mekongensis* Sanoamuang and Watiroyram, 2018 (Figure 4C); (15) *Phyllodiaptomus* (*Phyllodiaptomus*) *roietensis* (Sanoamuang and Watiroyram, 2020) (Figure 2D); (16) *Tropodiaptomus megahyaline* Sae-tang, Sanoamuang, and Maiphae, 2020; (17) *Dentodiaptomus orientalis* Sanoamuang and Watiroyram, 2021. In addition to the above, four more undescribed species (*Arctodiaptomus* sp., *Mongolodiaptomus* sp., *Phyllodiaptomus* sp. and *Tropodiaptomus* sp.) are waiting to be described.

Eight taxa—namely, *Arctodiaptomus* sp. (Figure 5B), *Mongolodiaptomus pectinidactylus* (Shen and Tai, 1964) (Figure 4D), *Neodiaptomus meggitti* Kiefer, 1932, *Tropodiaptomus hebereri* (Kiefer, 1930) (Figure 6A), *T. lanaonus* Kiefer, 1982, *T. oryzanus* Kiefer, 1937 (Figure 6B,C), *T. ruttneri* (Brehm, 1923) (Figure 6D), and *Paradiaptomus greeni* (Gurney, 1906) (Figure 5C,D)—are new to the fauna of Thailand. Moreover, this is the first record of a species complex of *P. greeni* in the subfamily Paradiaptominae in Southeast Asia.

3.2. Relative Occurrences and Distribution

Twelve species (28.6%) are commonly found in different kinds of habitats, and most are widely distributed throughout the six regions of Thailand. The most frequently encountered species in the sampled localities were *Mongolodiaptomus botulifer* (Figure 7A), *Phyllodiaptomus praedictus* (Figure 2A), *M. calcarus* (Figure 7B), *M. dumonti* (Figure 4A), *M. malaindosinensis* (Figure 7C), *Vietodiaptomus blachei* (Figure 8A), *P. christineae* (Figure 2B), *Eodiaptomus sanoamuangae* (Figure 3A), *Neodiaptomus yangtsekiangensis*, *E. draconisignivomi* (Figure 3D), *T. vicinus* (Figure 6E), and *H. elegans* (Figure 8B). The rest, including five fairly common, nine uncommon, seven rare, and nine extremely rare species, are also listed in Table 2.

The species richness of the diaptomids ranged from 1 to 7 species per locality at the same sampling dates. Although 2–3 species usually coexist in the same location, the co-existence of 4–7 diaptomid species has been observed in northeastern Thailand, particularly in temporary water habitats (Table 3). The co-occurrence of congeneric species was about 50% of the localities having co-occurrence records, and it was predominantly found in small water bodies. *M. botulifer* showed the highest co-occurrences with other species, followed by *M. dumonti*, *M. malaindosinensis*, *P. (P.) christineae*, and *V. blachei*.

Table 2. A list of the diaptomid copepods recorded with their habitat types, distributional regions, and relative occurrences in the freshwater ecosystems of Thailand. C = canal, L = lake, Pp/Sw = permanent pond/swamp, Re = reservoir, Ri = river, Rf = rice field, Rc = roadside canal, and Tp = temporary pond/rain pool. The distributional regions of Thailand are given by; N = North, NE = Northeast, C = Central, E = East, W = West, and S = South. Species with bold letters and marked with * are new records for Thailand.

No	Species	Habitat Types								Distributional Regions						Relative Occurrences	
		Permanent Waters				Temporary Waters				N	NE	C	E	W	S		
		C	L	Pp/Sw	Re	Ri	Rf	Rc	Tp								
Subfamily Diaptominae Kiefer, 1932																	
1	<i>Allodiaptomus raoi</i> Kiefer, 1936		✓		✓	✓	✓						✓				uncommon
2	<i>Arctodiaptomus</i> sp.					✓							✓				extremely rare
3	<i>Dentodiaptomus javanus</i> (Grochmalicki, 1915)			✓				✓	✓	✓	✓		✓		✓	✓	fairly common
4	<i>Dentodiaptomus orientalis</i> Sanoamuang & Watiroyram, 2021			✓					✓	✓			✓				uncommon
5	<i>Eodiaptomus draconisignivomi</i> Brehm, 1952	✓	✓	✓	✓	✓							✓	✓	✓	✓	fairly common
6	<i>Eodiaptomus phuphanensis</i> Sanoamuang, 2001				✓	✓			✓	✓			✓			✓	fairly common
7	<i>Eodiaptomus phuvongi</i> Sanoamuang & Sivongxay, 2004	✓		✓						✓			✓				fairly common
8	<i>Eodiaptomus sanoamuangae</i> Ranga Reddy & Dumont, 1998	✓	✓	✓	✓			✓	✓	✓	✓		✓			✓	common
9	<i>Heliodiaptomus elegans</i> Kiefer, 1935		✓	✓	✓	✓				✓	✓					✓	fairly common
10	<i>Heliodiaptomus phuthaiorum</i> Sanoamuang, 2004								✓	✓			✓				rare
11	<i>Heliodiaptomus viduus</i> (Gurney, 1916)	✓	✓	✓	✓						✓		✓	✓	✓		uncommon
12	<i>Mongolodiaptomus botulifer</i> (Kiefer, 1974)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	common
13	<i>Mongolodiaptomus calcarus</i> (Shen & Tai, 1965)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓			✓	✓	common
14	<i>Mongolodiaptomus dumonti</i> Sanoamuang, 2001	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		common
15	<i>Mongolodiaptomus loeiensis</i> Watiroyram & Sanoamuang, 2017									✓			✓				extremely rare
16	<i>Mongolodiaptomus malaindosinensis</i> (Lai & Fernando, 1978)		✓	✓	✓	✓				✓	✓	✓	✓	✓		✓	common
17	<i>Mongolodiaptomus mekongensis</i> Sanoamuang & Watiroyram, 2018			✓					✓	✓	✓		✓				uncommon

Table 2. Cont.

No	Species	Habitat Types								Distributional Regions						Relative Occurrences
		Permanent Waters					Temporary Waters			N	NE	C	E	W	S	
		C	L	Pp/Sw	Re	Ri	Rf	Rc	Tp							
18	<i>Mongolodiptomus pectinidactylus</i> (Shen & Tai, 1964) *		✓						✓		✓					rare
19	<i>Mongolodiptomus rarus</i> (Ranga Reddy, Sanoamuang & Dumont, 1998)							✓	✓	✓	✓					uncommon
20	<i>Mongolodiptomus uenoi</i> (Kikuchi, 1936)	✓		✓	✓	✓		✓	✓		✓					fairly common
21	<i>Mongolodiptomus</i> sp.			✓					✓		✓					extremely rare
22	<i>Neodiptomus laii</i> Kiefer, 1974	✓		✓					✓		✓		✓		✓	uncommon
23	<i>Neodiptomus meggitti</i> Kiefer, 1932 *			✓							✓					extremely rare
24	<i>Neodiptomus schmackeri</i> (Poppe & Richard, 1892)	✓		✓				✓	✓	✓	✓			✓	✓	fairly common
25	<i>Neodiptomus siamensis</i> Proogkiat & Sanoamuang, 2008								✓		✓					extremely rare
26	<i>Neodiptomus songkhramensis</i> Sanoamuang & Athibai, 2002							✓	✓	✓	✓					uncommon
27	<i>Neodiptomus yangtsekiangensis</i> Mashiko, 1951	✓	✓	✓	✓	✓		✓	✓		✓		✓		✓	common
28	<i>Phyllodiptomus (Ctenodiptomus) praedictus</i> Dumont & Ranga Reddy, 1994	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		common
29	<i>Phyllodiptomus (Ctenodiptomus) surinensis</i> Sanoamuang & Yindee, 2001	✓									✓					extremely rare
30	<i>Phyllodiptomus (Phyllodiptomus) christineae</i> Dumont, Ranga Reddy & Sanoamuang, 1996	✓	✓	✓	✓	✓			✓	✓	✓	✓	✓	✓		common
31	<i>Phyllodiptomus (Phyllodiptomus) roietensis</i> Sanoamuang & Watiroyram, 2020							✓	✓	✓	✓					rare
32	<i>Phyllodiptomus (Phyllodiptomus) thailandicus</i> Sanoamuang & Teeramaethee, 2006	✓		✓						✓			✓	✓	✓	rare
33	<i>Phyllodiptomus (Phyllodiptomus)</i> sp.			✓					✓		✓					rare
34	<i>Tropodiptomus hebereri</i> (Kiefer, 1930) *	✓									✓					extremely rare
35	<i>Tropodiptomus lanaonus</i> Kiefer, 1982 *		✓								✓	✓				rare

Table 2. Cont.

No	Species	Habitat Types								Distributional Regions						Relative Occurrences	
		Permanent waters					Temporary waters			N	NE	C	E	W	S		
		C	L	Pp/Sw	Re	Ri	Rf	Rc	Tp								
36	<i>Tropodiptomus megahyaline</i> Saetang, Sanoamuang & Maiphae, 2020		✓	✓			✓					✓					rare
37	<i>Tropodiptomus oryzanus</i> Kiefer, 1937 *		✓				✓	✓	✓			✓					uncommon
38	<i>Tropodiptomus ruttneri</i> (Brehm, 1923) *						✓		✓	✓	✓						uncommon
39	<i>Tropodiptomus vicinus</i> (Kiefer, 1930)	✓	✓	✓		✓				✓	✓	✓			✓	✓	fairly common
40	<i>Tropodiptomus</i> sp.								✓	✓							extremely rare
41	<i>Vietodiptomus blachei</i> (Brehm, 1951)	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓			✓	✓	common
Subfamily Paradiptominae Kiefer, 1932																	
42	<i>Paradiptomus greeni</i> (Gurney, 1906) *								✓		✓						extremely rare
		Total									17	35	14	11	15	8	




Figure 2. SEM photographs of the male leg 5 (P5) of four species of the genus *Phyllodiptomus*: (A) *P. (C.) praedictus*; (B) *P. (P.) christineae*; (C) *P. (P.) thailandicus*; and (D) *P. (P.) roietensis*.




Figure 3. SEM photographs of the male leg 5 (P5) of four species of the genus *Eodiaptomus*: (A) *E. sanoamuangae*; (B) *E. Phuphanensis*; (C) *E. phuvongi*; and (D) *E. draconisignivomi*.




Figure 4. SEM photographs of the male leg 5 (P5) of four species of the genus *Mongolodiptomus*: (A) *M. dumonti*; (B) *M. loeiensis*; (C) *M. mekongensis*; and (D) *M. pectinidactylus*.




Figure 5. Photographs of three species of diaptomid copepods. (A) male P5 of *Heliodiaptomus phuthaiorum*; (B) male P5 of *Arctodiaptomus* sp. (black arrows point to left and right hyaline membranes on left and right bases; (C) habitus of male *Paradiaptomus greenii*; (D) male P5 of *P. greenii*.




Figure 6. Photographs of the male leg 5 (P5) of four species of the genus *Tropodiptomus*: (A) *T. hebereri*; (B) *T. oryzanus*; (C) *T. oryzanus*, showing detail of left P5; and (D) *T. ruttneri*; (E) *T. vicinus*.



Figure 7. SEM photographs of the male leg 5 (P5) of four species of the genus *Mongolodiptomus*: (A) *M. botulifer*; (B) *M. calcarus*; (C) *M. malaindosinensis*; and (D) *M. uenoi*.




Figure 8. SEM photographs of the male leg 5 (P5) of four species of the diaptomid copepods: (A) *Vietodiptomus blachei*; (B) *Heliodiptomus elegans*; (C) *Neodiptomus schmackeri*; and (D) *Dentodiptomus javanus*.

Table 3. Co-occurrences of up to 7 species of diaptomid copepods in freshwater habitats in Northeast Thailand.

Localities	Sampling Dates	Co-occurrence of Diaptomid Species
Canal at Ban Paiklao, Chumphon Buri District, Surin Province	2 April 1999	1. <i>Dentodiaptomus javanus</i> 2. <i>Mongolodiaptomus dumonti</i> 3. <i>Mongolodiaptomus malaindosinensis</i> 4. <i>Mongolodiaptomus uenoi</i> 5. <i>Neodiaptomus yangtsekiangensis</i> 6. <i>Phyllodiaptomus (P.) christineae</i>
Irrigation Canal at Ban Nayom, Sanom District, Surin Province	29 July 1999	1. <i>Dentodiaptomus javanus</i> 2. <i>Mongolodiaptomus dumonti</i> 3. <i>Neodiaptomus laii</i> 4. <i>Phyllodiaptomus (P.) christineae</i> 5. <i>Phyllodiaptomus (C.) surinensis</i> 6. <i>Tropodiaptomus oryzanus</i> 7. <i>Tropodiaptomus vicinus</i>
Temporary pond at Ban Chaiya Buri, Chaiya Buri Sub-district, Tha Uthen District, Nakhon Phanom Province	18 May 2000	1. <i>Dentodiaptomus javanus</i> 2. <i>Eodiaptomus phuphanensis</i> 3. <i>Heliodiaptomus phuthaiorum</i> 4. <i>Mongolodiaptomus malaindosinensis</i> 5. <i>Neodiaptomus yangtsekiangensis</i> 6. <i>Vietodiaptomus blachei</i>
Roadside canal (temporary-water habitat), KM 24, Road #2050, Lao Suekok Sub-district, Ubon Rachathani Province	8 June 2002	1. <i>Eodiaptomus phuphanensis</i> 2. <i>Mongolodiaptomus pectinidactylus</i> 3. <i>Mongolodiaptomus uenoi</i> 4. <i>Neodiaptomus yangtsekiangensis</i> 5. <i>Tropodiaptomus oryzanus</i> , 6. <i>Vietodiaptomus blachei</i>
Lake Bueng Khong Long, Bueng Khong Long District, Bueng Kan Province	13 August 2002	1. <i>Allodiaptomus raoi</i> 2. <i>Heliodiaptomus elegans</i> 3. <i>Mongolodiaptomus pectinidactylus</i> 4. <i>Neodiaptomus yangtsekiangensis</i> 5. <i>Tropodiaptomus megahyaline</i>
Roadside canal (temporary-water habitat), Ban Dirikai, Muangsamsip District, Ubon Rachathani Province	19 October 2002	1. <i>Eodiaptomus phuphanensis</i> 2. <i>Heliodiaptomus elegans</i> 3. <i>Mongolodiaptomus malaindosinensis</i> 4. <i>Neodiaptomus yangtsekiangensis</i> 5. <i>Phyllodiaptomus (P.) christineae</i> 6. <i>Vietodiaptomus blachei</i>
Temporary pond at Ban Thasongcorn, Panna Nikom District, Sakon Nakhon Province	17 May 2004	1. <i>Eodiaptomus phuphanensis</i> 2. <i>Mongolodiaptomus malaindosinensis</i> 3. <i>Mongolodiaptomus uenoi</i> 4. <i>Neodiaptomus songkhramensis</i> 5. <i>Neodiaptomus yangtsekiangensis</i> 6. <i>Tropodiaptomus oryzanus</i> , 7. <i>Vietodiaptomus blachei</i>

It is noted that eight species (*Heliodiaptomus phuthaiorum* (Figure 5A), *Mongolodiaptomus loeiensis* (Figure 4B), *M. rarus*, *Neodiaptomus siamensis*, *N. songkhramensis*, *Phyllodiaptomus roietensis* (Figure 2D), *Tropodiaptomus* sp. and *Paradiaptomus greeni* (Figure 5C,D)) appear to occur only in temporary waters. The northeast is the most diverse region, with 35 recorded species. The other regions have 8–17 species.

The species recorded from Thailand can be classified according to their zoo-geographical distribution into six groups. The list of species endemic to Thailand, the lower Mekong River Basin, and Southeast Asia is provided in Table 4. Overall, 12 (28.6%) of the species recorded appear to be endemic to Thailand. Eight species (19%) have a more re-stricted distribution, occurring in the lower Mekong River Basin (Thailand, Laos, Cam-

bo-dia, and Vietnam), while the other eight species are widely distributed in the countries in the lower Mekong River Basin plus the other Southeast Asian countries, including Malaysia, Singapore, Indonesia, and the Philippines (Table 4). In addition, 10 species (23.8%) are widely distributed in Southeast Asia and East Asia. Three species—*Heliodiaptomus el-egans* (Figure 8B), *Heliodiaptomus viduus*, and *Neodiaptomus schmackeri* (Figure 8C)—are the most widely distributed across Southeast, East, and South Asia. Another species with a wide distribution is *Paradiaptomus greeni* (Figure 5C,D), occurring in Southeast Asia, South Asia, and Africa.

3.3. Taxonomic Checklist

Comments on the relative occurrences, distributional information, and taxonomic status of the species recorded in the present study are provided alphabetically according to the genera. In addition, information on the geographical distribution of the species in other countries, particularly Southeast Asian countries, is documented here. Order Calanoida Sars, 1903; Family Diaptomidae Baird, 1850; Subfamily Diaptominae Kiefer, 1932.

3.3.1. Genus *Allodiaptomus* Kiefer, 1936

Allodiaptomus raoi Kiefer, 1936 is an uncommon species that occurs in lakes, reservoirs, rivers, and rice fields in the northeast of Thailand. It was originally described in south India [37] and has been found in Cambodia [23], Vietnam [32], and China [38].

3.3.2. Genus *Arctodiaptomus* Kiefer, 1932

Arctodiaptomus sp. (Figure 5B) is identified as an undescribed species based on specimens collected from the Mun River, Ubon Ratchathani Province, northeastern Thailand. Since only one male specimen was collected, the description of this species will be provided when more specimens are available. According to the typical pincer-like processes at the end of the exopod of the male left P5 and a distinct hyaline membrane on the inner margin of the basis of both the left and right P5s in the male (Figure 5B), this species is placed in the genus *Arctodiaptomus*. The record of this genus in Thailand is unexpected since members of this genus inhabit water bodies across Eurasia, North Africa, and North Asia, including China [38]. In Southeast Asia, only *Arctodiaptomus dorsalis* (Marsh, 1907) has been recorded in more than 20 lakes in the Philippines [39]. This Neotropical species was originally described from the United States and has been found across Central America to northern South America [40]. *A. dorsalis* has been introduced to the Philippines since 1991 [41]. This species has been documented to prefer eutrophic conditions. Most native calanoids may have been displaced by this invasive species.

3.3.3. Genus *Dentodiaptomus* Shen and Tai, 1964

Dentodiaptomus javanus (Grochmalicki, 1915) (Figure 8D) was originally described from Java (Indonesia) as *Diaptomus javanus* [42]. In 1964, this taxon was raised as a type species of the genus *Dentodiaptomus* by Shen and Tai (1964) [43]. The presence of at least one “denticle” or “tooth-like process” on the male left P5 Exp-2 is a unique character that was used to name the genus *Dentodiaptomus* [43]. Another distinct characteristic of the male right P5 of this genus is the Exp-2, a longitudinal ridge on the posterior surface that terminates near the insertion of the principal spine. This ridge is not observed in other genera of diaptomids. In Thailand, this species commonly occurs in every type of temporary habitat (rice fields, roadside canals, temporary ponds) [9] and permanent ponds or swamps. It is widely distributed in the northeastern region but sporadic in the northern, eastern, and western regions. *D. javanus* has been found in Cambodia [23], Indonesia [31], Vietnam [32], and southern China [38].

Table 4. Diaptomid copepods from Thailand are classified according to their general geographic distribution. The Lower Mekong River Basin region consists of Thailand, Myanmar, Laos, Cambodia, and Vietnam. The Southeast Asian region consists of Thailand, Myanmar, Laos, Cambodia, Vietnam, Malaysia, Singapore, the Philippines, Indonesia, and Brunei. The East Asian region consists of China, Japan, North Korea, South Korea, Mongolia, and Taiwan. The South Asian region consists of India, Sri Lanka, Afghanistan, Bangladesh, Bhutan, Nepal, and Pakistan.

Distributional Regions	Number of Species	% of Species Recorded	List of Species
Thailand only (Endemic species)	12	28.6	<i>Arctodiaptomus</i> sp., <i>Heliodiaptomus phuthaiorum</i> , <i>Mongolodiaptomus dumonti</i> , <i>M. loeiensis</i> , <i>M. rarus</i> , <i>Mongolodiaptomus</i> sp., <i>Neodiaptomus siamensis</i> , <i>N. songkhramensis</i> , <i>Phyllodiaptomus</i> (C.) <i>surinensis</i> , <i>P. (P.) thailandicus</i> , <i>Tropodiaptomus megahyaline</i> , <i>Tropodiaptomus</i> sp.
Lower Mekong River Basin	8	19.0	<i>Dentodiaptomus orientalis</i> , <i>Eodiaptomus draconisignivomi</i> , <i>E. phuphanensis</i> , <i>E. phuvongi</i> , <i>Mongolodiaptomus mekongensis</i> , <i>Phyllodiaptomus</i> (P.) <i>christineae</i> , <i>P. (P.) roietensis</i> , <i>Phyllodiaptomus</i> sp.
Southeast Asia	8	19.0	<i>Mongolodiaptomus botulifer</i> , <i>M. malaindosinensis</i> , <i>Neodiaptomus laii</i> , <i>N. meggitti</i> , <i>Phyllodiaptomus</i> (C.) <i>praedictus</i> , <i>Tropodiaptomus lanaonus</i> , <i>T. vicinus</i> , <i>Vietodiaptomus blachei</i>
Southeast Asia and East Asia	10	23.8	<i>Allodiaptomus raoi</i> , <i>Dentodiaptomus javanus</i> , <i>Eodiaptomus sanoamuangae</i> , <i>Mongolodiaptomus calcarus</i> , <i>M. pectinidactylus</i> , <i>M. uenoi</i> , <i>Neodiaptomus yangtsekiangensis</i> , <i>Tropodiaptomus hebereri</i> , <i>T. oryzanus</i> , <i>T. ruttneri</i>
Southeast Asia, East Asia and South Asia	3	7.2	<i>Heliodiaptomus elegans</i> , <i>Heliodiaptomus viduus</i> , <i>Neodiaptomus schmackeri</i>
Southeast Asia, South Asia and Africa	1	2.4	<i>Paradiaptomus greeni</i>
Total	42	100	

Dentodiaptomus orientalis Sanoamuang and Watirogram, 2021 is a newly described taxon, recorded from five and six localities in the floodplain of the lower Mekong River Basin in Thailand and Cambodia, respectively [44]. *D. orientalis* is an uncommon taxon and has been found only in the northeast of Thailand in both temporary and permanent water habitats throughout the year. The new species differs from its congener in many morphological aspects, especially on their fifth legs. The male left P5 Exp-2 of this species has three inner distal denticles instead of one as in *D. javanus*.

3.3.4. Genus *Eodiaptomus* Kiefer, 1932

Eodiaptomus draconisignivomi Brehm, 1952 (Figure 3D) was described from the neighborhood of the Mekong River, Cambodia, in 1952 [45]. In Thailand, it is a common species occurring in permanent water bodies such as canals, rivers, lakes, reservoirs, and swamps in the northeastern, central, eastern, and western parts of the country [10]. This species appears to be a widely distributed diaptomid in the lower Mekong River Basin, as it is known from Cambodia [46], Laos [47], and Vietnam [32].

Eodiaptomus phuphanensis Sanoamuang, 2001 (Figure 3B) is a fairly common species inhabiting both permanent and temporary habitats but has been so far reported only in the northeast [11] and western Thailand (this study). It was first collected from two res-ervoirs

in the Phu Phan National Park in 2001 [11]. This species has been recorded in Cambodia and central Laos [47].

Eodiaptomus phuvongi Sanoamuang and Sivongxay, 2004 (Figure 3C) was described based on specimens collected from Thailand and Laos [16]. It is a fairly common species, especially in temporary habitats in northeastern Thailand, and is also found in canals, permanent ponds, and swamps. This species is morphologically most similar to *E. phuphanensis* by having a characteristically flat endopod on the male right P5. However, they can be distinguished from each other in the following features of the male P5: (1) the hyaline lamella on the inner margin of the right basis of *E. phuvongi* is semicircular, but it is rectangular in *E. phuphanensis*, and (2) the basis of the left P5 has a distinct spiny knob in *E. phuvongi* but lacks this knob in *E. phuphanensis*.

Eodiaptomus sanoamuangae Ranga Reddy and Dumont, 1998 (Figure 3A) was first collected from a roadside canal in Khon Kaen Province, northeastern Thailand [6]. It is a common species and has been found in every type of both permanent and temporary water bodies, except rivers [10]. This taxon was previously confused with the Chinese *E. sinensis* (Burckhardt, 1913) by Mashiko (1951) [48]. To date, *E. sanoamuangae* is known in the north, northeast, and west of Thailand (this study), and in the Yangtze River, and its tributaries in central China [48].

3.3.5. Genus *Heliodiaptomus* Kiefer, 1932

Heliodiaptomus elegans Kiefer, 1935 (Figure 8B) was briefly described based on specimens from Myanmar [49] and was redescribed from samples collected from Thailand by Ranga Reddy and Dumont (1999) [50]. It is a common species and has been found mostly in permanent water bodies, but sometimes in temporary ponds in the north, northeast, and west of Thailand. It has been reported from Cambodia [23], Bangladesh [50], Vietnam [32], and China [38].

Heliodiaptomus phuthaiorum Sanoamuang, 2004 (Figure 5A) was described from 9 of the 456 sampled localities in northeastern Thailand [13]. This species is rare and has been found only in temporary waters, such as ponds and roadside canals, in the vicinity of the Song Khram River, a tributary of the Mekong River.

Heliodiaptomus viduus (Gurney, 1916) is an uncommon species and has been recorded only in permanent water bodies (canals, lakes, swamps, ponds, and reservoirs) in the north, central, east, and west of Thailand ([9], this study). It was first described as *Diaptomus viduus* from Sri Lanka [51] and has been found in Myanmar and Pakistan. This species is a common diaptomid in southern India and Bangladesh [52].

3.3.6. Genus *Mongolodiaptomus* Kiefer, 1937

A criterion relating to the armature details of the second exopodal segment of the male right P5 has been proposed to differentiate the genus *Mongolodiaptomus* from the close congeners *Neodiaptomus* and *Allodiaptomus* by Ranga Reddy et al. (2000) [8]. Thus, some species of both genera have been transferred to *Mongolodiaptomus*.

Mongolodiaptomus botulifer (Kiefer, 1974) (Figure 7A) was originally described as *Neodiaptomus botulifer* from Malaysia [53] and redescribed using specimens from Thailand by Ranga Reddy et al. (1998) [7]. It is the most common diaptomid copepod and always occurs in great abundance in Thailand. This species inhabits every type of freshwater habitat in every region of the country. *M. botulifer* has also been reported in Singapore [28], Cambodia [3], Vietnam [32], and Laos [47].

Mongolodiaptomus calcarus (Shen and Tai, 1965) (Figure 7B) was originally described as *Allodiaptomus calcarus* from China [54] and redescribed by Ranga Reddy et al. (1998) [7]. Specimens of this species from Malaysia, Singapore, and Thailand have been confused with the Javan *Neodiaptomus mephistopheles* Brehm, 1933 by Lai and Fernando (1981, 1978) [34,55] and Schmoker et al. (2014) [28]. According to Ranga Reddy et al. (1998) [7], it is in fact *M. calcarus*, not *N. mephistopheles*. It is one of the most common diaptomids in Thailand, occurring in every type of freshwater habitat in the north, northeast, central, west, and

south of Thailand. *M. calcarus* has also been found in western Java, Indonesia, Laos [47], and Vietnam [32].

Mongolodiptomus dumonti Sanoamuang, 2001 (Figure 4A) was discovered in a reservoir and several localities in northeastern Thailand and then described by Sanoamuang (2001) [12]. It is a common freshwater calanoid occurring in every type of freshwater habitat in the north, northeast, central, east, and west of Thailand.

Mongolodiptomus loeiensis Watiroyam and Sanoamuang, 2017 (Figure 4B) was collected from a temporary pond in Loei Province of northeastern Thailand and described by Watiroyam and Sanoamuang (2017) [19]. It is an extremely rare taxon and has to date been found only in the type locality. This species was the single calanoid found in the type locality. *M. loeiensis* appears to be an endemic diaptomid of Thailand.

Mongolodiptomus malaindosinensis (Lai and Fernando, 1978) (Figure 7C) was originally described as *Neodiptomus malaindosinensis* based on specimens from Malaysia [56]. However, Ranga Reddy et al. (1998) [7] observed some morphological variabilities in *M. botulifer* from Thailand, leading to serious doubt about the validity of the closely allied *M. malaindosinensis*. Although the morphological features of the two congeners are similar, they can be distinguished from each other by the following characters: (1) male right P5 basis with a large, spherical hyaline lamella on the inner margin in *M. botulifer* (Figure 7A) versus a triangular hyaline lamella in *M. malaindosinensis* (Figure 7C); (2) male right P5 Exp-2 with a blunt chitinous process on the outer proximal third margin in *M. botulifer* versus minute 1–4 hyaline knobs in *M. malaindosinensis*; (3) male right P5 Exp-2 with a narrow chitinous flange on the inner margin in *M. botulifer* versus the more elevated flange in *M. malaindosinensis*; (4) right side of female genital double-somite with an obviously botuliform out-growth (cylindrical form with rounded end or sausage-shaped) in *M. botulifer* versus a small, conical out-growth with a slightly dilated proximal part and gradually tapering to the tip in *M. malaindosinensis*. Thus, the valid status of *M. malaindosinensis* is confirmed. In Thailand, *M. malaindosinensis* is a common species occurring in lakes, permanent ponds, swamps, reservoirs, rivers, and temporary ponds and has been found throughout the country except in the west of Thailand. This species has been reported in Singapore [55] and Cambodia [46]. Recently, *M. malaindosinensis* has been found in two shallow meso-eutrophic and hypereutrophic lakes in Malaysia [26].

Mongolodiptomus mekongensis Sanoamuang and Watiroyam, 2018 (Figure 4C) was discovered in freshwater ecosystems in the floodplain of the lower Mekong River Basin [20]. It is an uncommon taxon in permanent ponds and swamps but is common in temporary habitats such as rice fields, roadside canals, and temporary ponds in the Ubon Ratchathani Province of northeastern Thailand. This species has so far been recorded in Cambodia, Laos, and Vietnam [20].

Mongolodiptomus pectinidactylus (Shen and Tai, 1964) (Figure 4D) was originally described as *Allodiptomus pectinidactylus* from a fish pond on Hainan Island, China [43]. This species was reported for the first time in Thailand. It is a rare taxon and has been found in lakes and temporary ponds in the northeast of Thailand. This species is also found in Vietnam [32] and southern China [38].

Mongolodiptomus rarus (Ranga Reddy, Sanoamuang and Dumont, 1998) was originally described as *Allodiptomus rarus* [7]. It is an uncommon species and has been reported only from temporary water habitats in the northeast of Thailand.

Mongolodiptomus uenoi (Kikuchi, 1936) (Figure 7D) was originally described as *Diptomus uenoi* from Taiwan [57] and redescribed by Ranga Reddy et al. (2000) based on specimens from Thailand [8]. It is a fairly common taxon occurring in canals, swamps, reservoirs, rivers, roadside canals, and temporary ponds in the northeast of Thailand. This species has also been reported in Indonesia [8], Vietnam [32], and southern China [38].

Mongolodiptomus sp. is an extremely rare, undescribed diaptomid found in the northeast of Thailand. It is morphologically similar to *M. mekongensis*. A detailed description of this species will be published separately.

3.3.7. Genus *Neodiaptomus* Kiefer, 1932

Neodiaptomus laii Kiefer, 1974 was described based on specimens from Malaysia [58]. It is an uncommon species and has been found in canals, ponds, swamps, and temporary ponds in the northeast, east, and south of Thailand. This species is also found in Singapore [28] and Laos [47]. The two following diagnostic criteria of the genus *Neodiaptomus* are lacking: (1) the principal lateral spine is inserted generally at the middle of the outer margin of the male right P5 Exp-2 and (2) the right caudal ramus with a tooth-like chitinous structure at the inner ventro-distal corner; thus, it is more appropriate to transfer *N. laii* to the genus *Heliodiaptomus*.

Neodiaptomus meggitti Kiefer, 1932 was first described in Rangoon, Myanmar [59]. It was redescribed based on specimens from the South Andaman Islands, India, by Ranga Reddy (2000) [60]. *N. meggitti* is an extremely rare species. It was recorded for the first time in Thailand, occurring in a permanent pond in Chiang Rai Province, northern Thailand. It is also found in Malaysia [55].

Neodiaptomus schmackeri (Poppe and Richard, 1892) (Figure 8C) was originally described as *Diaptomus schmackeri* [61]. Among the 14 reported species of the genus *Neodiaptomus* [62], *N. schmackeri* is the most widely distributed species, being reported from southern China, the Russian Far East, Nepal, India, Bangladesh, Sri Lanka, Korea, Japan, Taiwan, the Philippines, Malaysia, Singapore [1,63] and Vietnam [32]. Recently, Alfonso et al. (2014) [64] have reported this species in eight lakes in Albania (Europe) for the first time. Since 2003, this species has most likely been introduced to Kazakhstan's Shardara Reservoir (Central Asia) with imported fish [65]. It is a fairly common species, occurring in canals, swamps, rice fields, roadside canals, and temporary ponds in the north, central and west of Thailand.

Neodiaptomus siamensis Proogkiat and Sanoamuang, 2008 was described from specimens recorded in Thailand [18]. It is an extremely rare species and has been found only in three temporary pools in Kamphaeng Phet and Phitsanulok Provinces (these provinces are sometimes designed in the northern region) in the central region of Thailand. This species has not been found in other countries.

Neodiaptomus songkhramensis Sanoamuang and Athibai, 2002 was discovered in several temporary water habitats in the vicinity of the Song Khram River, a tributary of the Mekong River [15]. It is an uncommon species, occurring in rice fields, roadside canals, and temporary ponds in northeastern Thailand. This species has not been reported in other countries.

Neodiaptomus yangtsekiangensis Mashiko, 1951 was described from samples recorded from the Yangtze River, central China [48]. It was redescribed based on specimens from Thailand by Ranga Reddy et al. (1998) [7]. *N. yangtsekiangensis* is a common species occurring in every type of habitat (except in rice fields) in the northeast, east, and south of Thailand. This species has been reported in southern China [38], Laos, Cambodia [46], and Vietnam [32].

3.3.8. Genus *Phyllodiaptomus* Kiefer, 1936

The genus *Phyllodiaptomus* Kiefer, 1936 is among the most common freshwater diaptomid copepods in Southeast Asia [9]. Members of this genus are grouped into two sub-genera, based on the distinctive structures of the male left P5 Exp-2. The subgenus *Phyllo-diaptomus* (*Ctenodiaptomus*) (Dumont, Ranga Reddy and Sanoamuang, 1996) [5] has a ser-rated hyaline fan on the inner margin of the left P5 Exp-2, while in *Phyllodiaptomus* (*Phyl-lodiaptomus*) (Dumont, Ranga Reddy and Sanoamuang, 1996), the corresponding structure bears a field of spinules.

Subgenus *Phyllodiaptomus* (*Ctenodiaptomus*)

Dumont, Ranga Reddy, and Sanoamuang, 1996

Phyllodiaptomus (*Ctenodiaptomus*) *praedictus* Dumont and Ranga Reddy, 1994 (Figure 2A) was described based on specimens from Thailand [4]. It is a common species inhabiting every type of freshwater habitat throughout Thailand, except in the south. This species

has also been recorded in Laos and Cambodia [46]. Another closely related subspecies, *Phyllodiaptomus (Ctenodiaptomus) praedictus sulawesensis* Alekseev and Vaillant, 2013, has been reported in Lake Tondano, North Sulawesi, Indonesia [31].

Phyllodiaptomus (Ctenodiaptomus) surinensis Sanoamuang and Yindee, 2001 was discovered in two irrigation canals in Surin Province, northeastern Thailand [14]. It is an extremely rare species and has never been found again. This species appears to be an endemic species of Thailand.

Subgenus *Phyllodiaptomus (Phyllodiaptomus)*

Dumont, Ranga Reddy and Sanoamuang, 1996

Phyllodiaptomus (Phyllodiaptomus) christineae Dumont, Ranga Reddy, and Sanoamuang, 1996 (Figure 2B) was described based on specimens from Thailand [5]. It is a common species and has been recorded in canals, lakes, permanent ponds, swamps, reservoirs, rivers, and temporary ponds in the north, northeast, central, east, and west of Thailand. This species has not yet been found in the south. This species has also been reported in Laos [46].

Phyllodiaptomus (Phyllodiaptomus) roietensis Sanoamuang and Watiroyram, 2020 (Figure 2D) was described based on specimens from Thailand and Cambodia [21]. It is a rare species, inhabiting rice fields, roadside canals, and temporary ponds only in the northeast of Thailand.

Phyllodiaptomus (Phyllodiaptomus) thailandicus Sanoamuang and Teeramaethee, 2006 (Figure 2C) was described based on specimens from Thailand [17]. It is a rare species occurring in permanent and temporary ponds in the northeast and the south of Thailand. This species appears to be an endemic species of Thailand.

Phyllodiaptomus (Phyllodiaptomus) sp. is a rare, undescribed diaptomid in the north-east of Thailand. It is similar to *P. (P.) christineae*. A description of this species will be published in a separate article.

3.3.9. Genus *Tropodiaptomus* Kiefer, 1932

Tropodiaptomus hebereri (Kiefer, 1930) (Figure 6A) was originally named as *Diaptomus hebereri* [66]. It is an extremely rare taxon that occurs in canals in the north of Thailand. This species has been found in Malaysia [67], Singapore [3], India [68], Indonesia [31], and China [38].

Tropodiaptomus lanaonus Kiefer, 1982 was described from Lake Lanao, Mindanao Island, Philippines [69]. It was recorded as an endemic diaptomid of the Philippines by Lopez et al. (2017) [30]. *T. lanaonus* is a rare species and has been reported from lakes in the northeast and central regions of Thailand.

Tropodiaptomus megahyaline Saetang, Sanoamuang, and Maiphae, 2020 is a newly described species from Thailand [22]. It is a rare species and has been reported from rice fields and permanent ponds in the northeast of Thailand. This species was recorded as *Tropodiaptomus sp.* in Lake Bueng Khong Long in Bueng Kan Province, northeastern Thailand by Teeramaethee (2006) [70].

Tropodiaptomus oryzanus Kiefer, 1937 (Figure 6B,C) was described based on samples from rice fields in Taiwan [71]. It is an uncommon species and has been found in lakes and temporary habitats such as rice fields, roadside canals, and temporary ponds in the northeast of Thailand. This species is also found in Vietnam [32], China, Korea, and Japan [38].

Tropodiaptomus ruttneri (Brehm, 1923) (Figure 6D) was originally named *Diaptomus ruttneri*, based on specimens from Yunnan Province, China. It is an uncommon species and has been reported from rice fields and temporary ponds in the north and northeast of Thailand. This species is also known from Malaysia [67], Japan, and Taiwan [3].

Tropodiaptomus vicinus (Kiefer, 1930) (Figure 6E) was originally described as *Diaptomus vicinus* based on specimens from Indonesia [72]. It is a common species and has been found in irrigation canals, lakes, permanent ponds, swamps, and rivers in the north,

northeast, central, west, and south of Thailand. This species has been found in the Philippines [3], Malaysia [67], Cambodia [46], Indonesia [31], India [68] and Vietnam [32].

Tropodiptomus sp. is an extremely rare, undescribed diaptomid collected from a temporary pond in the north of Thailand. Morphologically, it is similar to *T. ruttneri*. A description of this species will be published separately.

3.3.10. Genus *Vietodiptomus* Dang, 1977

The genus *Vietodiptomus* was created based on specimens from Vietnam, with *Vietodiptomus hatinhensis* Dang, 1977 as the type species [73]. According to Walter and Box-shall (2021), the other two members of this genus are *Vietodiptomus blachei* (Brehm, 1951) and *Vietodiptomus tridentatus* Dang and Ho, 1985 [62]. Since the description of this genus was published in the Vietnamese language, a redescription using English is required.

Vietodiptomus blachei (Brehm, 1951) (Figure 8A) was originally described as *Eodiptomus blachei* based on specimens from Cambodia [23]. It is a common species occurring in canals, lakes, permanent ponds, swamps, reservoirs, rice fields, roadside canals, and temporary ponds, and has been recorded in the north, northeast, central, west, and south of Thailand. This species has not yet been found in the eastern region. This species has been reported from Indonesia [3], Laos [47], Malaysia [67], and Singapore [28].

Subfamily Paradiaptominae Kiefer, 1932

Genus *Paradiaptomus* Sars, 1895

Paradiaptomus greeni (Gurney, 1906) (Figure 5C,D) was originally described as *Diaptomus greeni* based on samples from Sri Lanka (Ceylon) [74]. It is also found in South Africa [75], Tunisia [76], India [77], and China [38]. It is an extremely rare species and has been found so far only in a temporary pond in Khon Kaen Province, northeastern Thailand. This is the first record of a representative of the Paradiaptominae in Southeast Asia.

4. Discussion and Conclusions

The species diversity of freshwater diaptomid copepods (42 species) in Thailand is higher than that reported from the other Southeast Asian countries—8 from the Philippines [30], 12 from Malaysia and Singapore [25], 17 from Indonesia [31], 19 from Laos [24], 24 from Cambodia [46], and 29 from Vietnam [32]. The comparison with other Asian countries revealed that the diaptomid diversity of Thailand is higher than Kazakhstan (26 species) [65] but comparable to that of India (about 50 species) [77] and China (56 species) [38], although the country areas of India and China are larger than Thailand by 6.4 and 18.7 times, respectively. Thus, Thailand is one of the countries with the most diversified diaptomid fauna in the world, and it is a biodiversity hotspot for diaptomid copepods in Southeast Asia.

The new record for Thailand (*Paradiaptomus greeni*) predominantly occurs in Africa but is also known in other Asian countries, including Sri Lanka, India [75], and the Nanning Province of southern China [38]. It is noted that we did not find two species—*Tropodiptomus doriai* and *Sinodiptomus chaffanjoni*—in our extensive collection. Both species were previously recorded in Thailand by Daday (1906) [34] and Boonsom (1984) [35], respectively.

Thailand has a greater record of co-occurrence (up to seven species) than India's Northern Western Ghats (up to four species) [77]. The high degree of diaptomid species co-occurrence in Thailand is remarkable, as it is uncommon to find more than one calan-oid species in a temperate lake [78]. The difference in diversity might also be partly a result of artefacts due to more intensive surveying and a good basis in systematics and taxonomy in Thailand within the region.

The species richness of diaptomid copepods (35 species) in the northeast, which is higher than in the other regions of Thailand (8–17 species), can be explained by the fact that it covers about one-third of the country and contains many temporary water bodies in

the floodplains of major rivers, and the majority of the samples (about 60%) were collected from this region.

In conclusion, a total of 4962 samples were analyzed from 2150 diverse freshwater habitats across Thailand, and 42 species belonging to 11 genera in 2 subfamilies of Diaptomidae were identified. The current number of species listed is the highest among those reported in other Southeast Asian countries, ranging from 8 to 29. *Mongolodiptomus* is the most diverse genus with 10 species, followed by *Tropodiptomus* with 7, and *Neodiptomus* and *Phyllodiptomus* with 6. In total, 17 species (40.5% of the species recorded) have been described originally from Thailand since 1994. Eight taxa (*Arctodiptomus* sp., *Mongolodiptomus pectinidactylus*, *Neodiptomus meggitti*, *Tropodiptomus hebereri*, *T. lanaonus*, *T. oryzanus*, *T. ruttneri*, and *Paradiptomus greeni*) are recorded for the first time in Thailand. This is the first record of a species (*P. greeni*) in the subfamily Paradiptominae in Southeast Asia as well. The relative occurrences of the species recorded are categorized into five groups: 12 common, 5 fairly common, 9 uncommon, 7 rare, and 9 extremely rare species. The most frequently encountered species in the sampled localities, which are widely distributed throughout the six regions of Thailand, were *Mongolodiptomus botulifer*, *Phyllodiptomus praedictus*, *M. calcarus*, *M. dumonti*, *M. malaindosinensis*, *Vietodiptomus blachei*, *P. christineae*, *Eodiptomus sanoamuangae*, *Neodiptomus yangtsekiangensis*, *E. draconisignivomi*, *Tropodiptomus vicinus*, and *Heliodiptomus elegans*. The species richness of the diaptomids ranged from 1 to 7 species per locality at the same sampling dates. The coexistence of 4–7 diaptomid species has been observed in northeastern Thailand, particularly in small-sized temporary-water habitats. The high degree of diaptomid species co-occurrence in Thailand is remarkable, as it is uncommon to find more than one calanoid species in a temperate lake. Eight species appear to occur only in temporary waters. The northeast is the most diverse region, with 35 recorded species. Twelve species appear to be endemic to Thailand, and eight are endemic to the lower Mekong River Basin countries (Thailand, Cambodia, Laos, and Vietnam). This study demonstrated that species diversity is high in temporary water habitats in the floodplains of major rivers, and Thailand is a biodiversity hotspot for diaptomid copepods in Southeast Asia. In addition, the taxonomic status of *Mongolodiptomus malaindosinensis* has been confirmed.

Author Contributions: Conceptualization, L.S. and P.D.; methodology, L.S. and P.D.; formal analysis, L.S.; investigation, P.D.; resources, P.D.; data curation, L.S.; writing—original draft preparation, L.S.; writing—review and editing, L.S.; visualization, P.D.; funding acquisition, L.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by a Research and Graduate Studies Grant, from Khon Kaen University, Grant Number KKU-ATRC-2564-01.

Institutional Review Board Statement: Ethical review and approval were waived for this study due to the fact that the animals that we collected from natural habitats are tiny zooplankton and microscopic in size.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study is available on request from the corresponding author.

Acknowledgments: The authors would like to thank Jittra Teeramaethee, Santi Watiroyram, Sujiphon Athibai, Kamonwan Koomput, Phuttaphanee Boonmak, Rachada Chaichareon, Supatra Tiangnga, Supatra Leckchan, Khunapat Proongkiat, Weera Yindee, Panna Wansuang, Boonthida Boonsit, Sirichai Faitakham, and Parinda Tungpanyaporn, for their assistance in the sample collections and identifications. The authors gratefully acknowledge two anonymous reviewers for their critical review of the manuscript. Nithikarn Sanoamuang and Sirikamon Phlaingam are thanked for linguistic corrections of the manuscript and figure preparation.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Dussart, B.; Defaye, D. *World Directory of Crustacea Copepoda of Inland Waters, I-Calaniformes*; Backhuys Publishers: Leiden, The Netherlands, 2002; pp. 1–276.
- Boxshall, G.A.; Defaye, D. Global diversity of copepods (Crustacea: Copepoda) in freshwater. *Hydrobiologia* **2008**, *595*, 195–207. [[CrossRef](#)]
- Lai, H.C.; Fernando, C.H. Zoogeographical distribution of Southeast Asian freshwater Calanoida. *Hydrobiologia* **1980**, *74*, 53–66. [[CrossRef](#)]
- Dumont, H.J.; Ranga Reddy, Y. *Phyllodiaptomus praedictus* n. sp. (Copepoda, Calanoida) from Thailand. *Hydrobiologia* **1994**, *273*, 101–110. [[CrossRef](#)]
- Dumont, H.J.; Ranga Reddy, Y.; Sanoamuang, L. Description of *Phyllodiaptomus christineae* n. sp. from Thailand, and distinction of two subgenera within *Phyllodiaptomus* Kiefer, 1936 (Copepoda, Calanoida). *Hydrobiologia* **1996**, *323*, 139–148. [[CrossRef](#)]
- Ranga Reddy, Y.; Dumont, H.J. A review of the genus *Eodiaptomus* Kiefer, 1932, with the description of *E. sanuamuangae* n. sp. from Thailand, and a redescription of *E. lumholtzi* (Sars, 1889) from Australia (Copepoda, Calanoida). *Hydrobiologia* **1998**, *361*, 169–189. [[CrossRef](#)]
- Ranga Reddy, Y.; Sanoamuang, L.; Dumont, H.J. A note on the Diaptomidae of Thailand, including redescription of three species and description of a new species (Copepoda, Calanoida). *Hydrobiologia* **1998**, *361*, 201–223. [[CrossRef](#)]
- Ranga Reddy, Y.; Sanoamuang, L.; Dumont, H.J. Amended delimitation of *Mongolodiaptomus* against *Neodiaptomus* and *Allodiaptomus* and redescription of the little known *Mongolodiaptomus uenoi* (Kikuchi, 1936) from Thailand (Copepoda: Calanoida: Diaptomidae). *Hydrobiologia* **2000**, *418*, 99–109. [[CrossRef](#)]
- Sanoamuang, L. Species composition and distribution of freshwater Calanoida and Cyclopoida (Copepoda) of north-east Thailand. In *Crustaceans and Biodiversity Crisis*; Schram, F.R., Klein, J.V.C., Eds.; Brill Academic Publishers: Leiden, The Netherlands, 1999; pp. 217–230.
- Sanoamuang, L. Distributions of three *Eodiaptomus* species (Copepoda: Calanoida) in Thailand, with a redescription of *E. draconisignivomi* Brehm, 1952. *Hydrobiologia* **2001**, *453*, 565–576. [[CrossRef](#)]
- Sanoamuang, L. *Eodiaptomus phuphanensis*, n. sp., a new freshwater copepod (Calanoida: Diaptomidae) from the Phu Phan National Park, Thailand. *Int. Rev. Hydrobiol.* **2001**, *86*, 219–228. [[CrossRef](#)]
- Sanoamuang, L. *Mongolodiaptomus dumonti* n. sp., a new freshwater copepod (Calanoida, Diaptomidae) from Thailand. *Hydrobiologia* **2001**, *448*, 41–52. [[CrossRef](#)]
- Sanoamuang, L. *Heliodiaptomus phuthaiorum* n. sp., a new freshwater copepod (Calanoida, Diaptomidae) from temporary ponds in northeast Thailand. *Int. Rev. Hydrobiol.* **2004**, *89*, 392–406. [[CrossRef](#)]
- Sanoamuang, L.; Yindee, W. A new species of *Phyllodiaptomus* (Copepoda, Diaptomidae) from northeast Thailand. *Crustaceana* **2001**, *74*, 435–448.
- Sanoamuang, L.; Athibai, S. A new species of *Neodiaptomus* (Copepoda, Diaptomidae) from temporary waters in northeast Thailand. *Hydrobiologia* **2002**, *489*, 71–82. [[CrossRef](#)]
- Sanoamuang, L.; Sivongxay, N. Description of *Eodiaptomus phuwoongi*, n. sp. (Copepoda, Calanoida) from Thailand and Laos. *Crustaceana* **2004**, *77*, 1223–1236. [[CrossRef](#)]
- Sanoamuang, L.; Teeramaethee, J. *Phyllodiaptomus thailandicus*, a new freshwater copepod (Copepoda, Calanoida, Diaptomidae) from Thailand. *Crustaceana* **2006**, *79*, 475–487.
- Proongkiat, I.; Sanoamuang, L. Description of *Neodiaptomus siamensis*, a new diaptomid copepod (Copepoda, Calanoida) from temporary pools in northern Thailand. *Crustaceana* **2008**, *81*, 177–189.
- Watiroyram, S.; Sanoamuang, L. A new species of *Mongolodiaptomus* Kiefer, 1938 from northeast Thailand and a key to the species (Crustacea: Copepoda, Calanoida, Diaptomidae). *Zookeys* **2017**, *710*, 15–32. [[CrossRef](#)]
- Sanoamuang, L.; Watiroyram, S. *Mongolodiaptomus mekongensis*, a new species of copepod (Copepoda, Calanoida, Diaptomidae) from temporary waters in the floodplain of the lower Mekong River Basin. *Raffles Bull. Zool.* **2018**, *66*, 782–796.
- Sanoamuang, L.; Watiroyram, S. *Phyllodiaptomus (Phyllodiaptomus) roietensis*, a new diaptomid copepod (Copepoda: Calanoida) from temporary waters in Thailand and Cambodia, with a key to the species. *ZooKeys* **2020**, *911*, 1–20. [[CrossRef](#)]
- Saetang, T.; Sanoamuang, L.; Maiphae, S. A new species of genus *Tropodiaptomus* Kiefer, 1932 (Crustacea: Copepoda: Calanoida: Diaptomidae) from Thailand. *J. Nat. Hist.* **2020**, *54*, 2297–2322. [[CrossRef](#)]
- Brehm, V. Cladocera and Copepoda Calanoida von Cambodja. *Cybiuum* **1951**, *6*, 95–124.
- Sanoamuang, L.; Watiroyram, S. *Allodiaptomus nongensis*, a new diaptomid copepod (Copepoda: Calanoida) from a tributary of the Mekong River, with notes on its consumption by local people in Central Laos. *J. Limnol.* **2019**, *78*, 185–200. [[CrossRef](#)]
- Lim, R.P.; Lai, H.C. Crustacea: Copepoda, Calanoida. In *Freshwater Invertebrates of the Malaysian Region*; Yule, C.M., Sen, Y.H., Eds.; Academy of Sciences Malaysia: Kuala Lumpur, Malaysia, 2014; pp. 254–266.
- Umi, W.A.D.; Yusoff, F.M.; Aris, A.Z.; Sharip, Z.; Sinev, A.Y. Planktonic microcrustacean community structure varies with trophic status and environmental variables in tropical shallow lakes in Malaysia. *Diversity* **2020**, *12*, 322. [[CrossRef](#)]
- Alekseev, V.R.; Yusoff, F.M.; Fefilova, E.B. Continental copepod biodiversity in North-Eastern Borneo, Malaysia. *Arthropoda Sel.* **2016**, *25*, 187–197. [[CrossRef](#)]
- Schmoker, C.; Mahjoub, M.S.; Calbet, A.; Hsiao, S.H.; Russo, F.; Larsen, O.; Trottet, A.; Drillet, G. A review of the zooplankton in Singapore waters. *Raffles Bull. Zool.* **2014**, *62*, 726–749.

29. Mamaril, A.C. Zooplankton diversity in Philippine lakes. In *Conservation and Ecological Management of Philippine Lakes in Relation to Fisheries and Aquaculture*; Santiago, C.B., Cuvin-Aralar, M.L., Basiao, Z.U., Eds.; Southeast Asian Fisheries Development Center, Aquaculture Department: Iloilo, Philippines; Philippine Council for Aquatic and Marine Research and Development: Los Baños, Laguna; Bureau of Fisheries and Aquatic Resources: Quezon City, Philippines, 2001; pp. 81–93.
30. Lopez, M.L.D.; Pascual, J.A.F.; Dela Paz, E.S.P.; Rizo, E.Z.C.; Tordesillas, D.T.; Guinto, S.K.; Han, B.; Dumont, H.J.; Mamaril, A.C.; Papa, R.D.S. Annotated checklist and insular distribution of freshwater microcrustaceans (Copepoda: Calanoida & Cyclopoida; Cladocera: Anomopoda & Ctenopoda) in the Philippines. *Raffles Bull. Zool.* **2017**, *65*, 623–654.
31. Alekseev, V.R.; Haffner, D.G.; Vaillant, J.J.; Yusoff, F.M. Cyclopoid and calanoid copepod biodiversity in Indonesia. *J. Limnol.* **2013**, *72*, 245–274. [[CrossRef](#)]
32. Tran, D.L.; Dang, N.T.; Ho, T.H. An annotated checklist of the family Diaptomidae Sars, 1903 (Copepoda, Calanoida) in Vietnam. *Tap Chi Sinh Hoc.* **2016**, *38*, 384–399. [[CrossRef](#)]
33. Bricker, K.S.; Wongrat, L.; Gannon, J.E. Composition and distribution of Crustacean plankton in twelve inland water bodies of Thailand. *J. Fish. Environ.* **1978**, *10*, 1–14.
34. Lai, H.C.; Fernando, C.H. The freshwater Calanoida (Crustacea: Copepoda) of Thailand. *Hydrobiologia* **1981**, *76*, 161–178. [[CrossRef](#)]
35. Boonsom, J. The freshwater zooplankton of Thailand (Rotifera and Crustacea). *Hydrobiologia* **1984**, *113*, 223–229. [[CrossRef](#)]
36. Wikipedia. Regions of Thailand, Wikimedia Foundation. Available online: https://en.wikipedia.org/wiki/Regions_of_Thailand (accessed on 26 June 2021).
37. Kiefer, F. Indische Ruderfusskrebse (Crustacea, Copepoda). *Zool. Anz.* **1936**, *113*, 136–142.
38. Li, H.; Dumont, H.J.; Han, B.P.; Lin, Q. Updated checklist and distribution of the diaptomid copepods (Copepoda, Calanoida, Diaptomidae) of China. *Crustaceana* **2018**, *91*, 335–352. [[CrossRef](#)]
39. Tordesillas, D.T.; Paredes, P.M.F.; Villaruel, K.P.E.; Queneri, C.A.A.M.; Rico, J.L.; Ban, S.; Papa, R.D.S. Effects of food concentration on the reproductive capacity of the invasive freshwater calanoid copepod *Arctodiaptomus dorsalis* (Marsh, 1907) in the Philippines. *J. Crust. Biol.* **2018**, *38*, 101–106. [[CrossRef](#)]
40. Reid, J.W. *Arctodiaptomus dorsalis* (Marsh): A case history of copepod dispersal. *Banisteria* **2007**, *1860*, 3–18.
41. Papa, R.D.S.; Li, H.; Tordesillas, D.T.; Han, B.; Dumont, H.J. Massive invasion of *Arctodiaptomus dorsalis* (Copepoda, Calanoida, Diaptomidae) in Philippine lakes: A threat to Asian zooplankton biodiversity? *Biol. Invasions* **2012**, *14*, 2471–2478. [[CrossRef](#)]
42. Grochmalicki, J. Beitrag zur Kenntnis der süßwasserefauna Javas: Phyllopora, Copepoda und Ostracod. Bulletin International de l'Académie des Sciences de Cracovie, Classe des Sciences Mathématiques et Naturelles. *Serie B Sci. Nat.* **1915**, *1915*, 217–242.
43. Shen, C.J.; Tai, A.Y. Descriptions of eight new species of freshwater copepods chiefly from the Pearl River delta, south China. *Acta Zootaxo. Sin.* **1964**, *16*, 225–239.
44. Sanoamuang, L.; Watirogram, S. New species of copepod (Copepoda, Calanoida) from the floodplain of the lower Mekong River Basin in Thailand and Cambodia, with amended diagnosis of the genus *Dentodiaptomus* Shen & Tai, 1964. *Raffles Bull. Zool.* **2021**, in press.
45. Brehm, V. Vorläufige Mitteilung über einen neuen *Eodiaptomus* aus Hinterindien. *Anz. Öst. Akad. Wiss. Wien.* **1952**, *89*, 215–217.
46. Chaicharoen, R. Diversity and Distribution of Freshwater Calanoid and Cyclopoid Copepods in 7 Provinces of Cambodia. Ph.D. Thesis, Graduate School, Khon Kaen University, Khon Kaen, Thailand, 2011; pp. 1–216.
47. Sanoamuang, L. *Freshwater Zooplankton: Calanoid Copepods in Thailand*; Klangnatham Publishers: Khon Kaen, Thailand, 2002; pp. 1–159.
48. Mashiko, K. Studies of the freshwater plankton of central China. II. *Sci Rep. Kanazawa Univ.* **1951**, *1*, 137–154.
49. Kiefer, F. Beiträge zur Copepodenkunde (XVII). *Zool. Anz.* **1935**, *109*, 88–93.
50. Ranga Reddy, Y.; Dumont, H.J. Redescription of *Heliodiaptomus elegans* Kiefer, 1935, a rare south-east Asian calanoid copepod. *Hydrobiologia* **1999**, *394*, 145–152. [[CrossRef](#)]
51. Gurney, R. On some Freshwater Entomostraca from Ceylon. *Proc. Zool. Soc. Lond.* **1916**, *86*, 333–343. [[CrossRef](#)]
52. Ranga Reddy, Y. *Copepoda: Calanoida: Diaptomidae. Key to the Genera Heliodiaptomus, Allodiaptomus, Neodiaptomus, Phylloidiaptomus, Eodiaptomus, Arctodiaptomus and Sinodiaptomus*; SPB Academic: The Hague, Philippines, 1994; pp. 1–221.
53. Kiefer, F. Zwei neue Ruderfusskrebse (Crustacea, Copepoda) aus Malaysia. *Beitr. Naturk. Forsch. Südwestdeutschl.* **1974**, *33*, 219–222.
54. Shen, C.J.; Tai, A.Y. Descriptions of six new species of freshwater copepods chiefly from the Pearl River delta, South China. *Acta Zool. Sin.* **1965**, *2*, 126–140.
55. Lai, H.C.; Fernando, C.H. The freshwater Calanoida (Crustacea: Copepoda) of Singapore and peninsular Malaysia. *Hydrobiologia* **1978**, *61*, 113–127. [[CrossRef](#)]
56. Lai, H.C.; Fernando, C.H. Redescription of *Neodiaptomus botulifer* Kiefer and one of its related species. *Hydrobiologia* **1978**, *59*, 229–235. [[CrossRef](#)]
57. Kikuchi, K. A new species of *Diaptomus* from Formosa. *Proc. Imp. Acad. Jpn.* **1936**, *12*, 198–200. [[CrossRef](#)]
58. Kiefer, F. Eine neue Diaptomidenart aus Malaysia (Crustacea, Copepoda, Calanoida). *Zool. Anz.* **1974**, *192*, 420–424.
59. Kiefer, F. Zwei neue Diaptomiden (Copepoda, Calanoida) aus Indien. *Zool. Anz.* **1932**, *100*, 265–270.
60. Ranga Reddy, Y. *Neodiaptomus meggitti* Kiefer, 1932. A rare south-east Asian species from the Andaman Islands, India (Copepoda, Calanoida, Diaptomidae). *Crustaceana* **2000**, *73*, 257–272. [[CrossRef](#)]

61. Poppe, S.A.; Richard, J. Description du *Diaptomus schmackeri* n. sp., recueilli par M. Schmacker dans le lac Tahoo (Chine). *Bull. Soc. Zool. Fr.* **1892**, *17*, 149–151.
62. Walter, T.C.; Boxshall, G. World of Copepods Database. *Neodiaptomus* Kiefer, 1932. Available online: <http://www.marinespecies.org/copepoda/aphia.php?p=taxdetails&id=347568> (accessed on 29 June 2021).
63. Ranga Reddy, Y. *Neodiaptomus prateek* n. sp., a new freshwater copepod from Assam, India, with critical review of the generic assignment of *Neodiaptomus* spp. and a note on diaptomid species richness (Calanoida: Diaptomidae). *J. Crust. Biol.* **2013**, *33*, 849–865. [[CrossRef](#)]
64. Alfonso, G.; Russo, R.; Belmonte, G. First record of the Asian diaptomid *Neodiaptomus schmackeri* (Poppe & Richard, 1892) (Crustacea: Copepoda: Calanoida) in Europe. *J. Limnol.* **2014**, *73*, 584–592.
65. Krupa, E.; Aubakirova, M. Checklist and distribution of Calanoida (Crustacea: Copepoda) in Kazakhstan (Central Asia). *Water* **2021**, *13*, 2015. [[CrossRef](#)]
66. Kiefer, F. Neue ruderfüßkrebse von den Sunda-Inseln. *Zool. Anz.* **1930**, *86*, 185–189.
67. Lim, R.P.; Fernando, C.H. A review of Malaysian freshwater Copepoda with notes on new records and little known species. *Hydrobiologia* **1985**, *128*, 71–89. [[CrossRef](#)]
68. Ranga Reddy, Y. *Tropodiaptomus signatus* Kiefer, 1982, a little-known species from Loktak Lake, Manipur State, India (Copepoda, Calanoida, Diaptomidae). *Crustaceana* **2013**, *86*, 1675–1688. [[CrossRef](#)]
69. Kiefer, F. Vergleichende Untersuchungen über Morphologie, Taxonomie und geographische Verbreitung der Arten der Gattung *Tropodiaptomus* Kiefer (Copepoda: Calanoida) aus asiatischen Binnengewässern. *Hydrobiologia* **1982**, *93*, 223–253. [[CrossRef](#)]
70. Teeramaethee, J. Species Diversity and Abundance of Rotifers, Cladocerans and Copepods in Two Wetlands: Bueng Boraphet, Nakhon Sawan Province and Bueng Khong Long, Nong Khai Province. Ph.D. Thesis, Graduate School, Khon Kaen University, Khon Kaen, Thailand, 2006; pp. 1–161.
71. Kiefer, F. Süßwassercopoden aus Ostasien. II. Neue Diaptomiden und Cyclopiden von der Insel Formosa. *Zool. Anz.* **1937**, *119*, 58–64.
72. Kiefer, F. Beiträge zur Copepodenkunde. 38. Zur Kenntnis des *Diaptomus orientalis* Brady-Sars. *Zool. Anz.* **1930**, *87*, 120–122.
73. Dang, N.T. New diaptomid crustacean from Vietnam. *Tap Chi Sinh Vat-Dia Hoc.* **1977**, *15*, 97–99, (In Vietnamese with English summary).
74. Gurney, R. On two new Entomostraca from Ceylon. *Spolia Zeylan.* **1906**, *4*, 126–134.
75. Rama Devi, C.; Ranga Reddy, Y. The complete postembryonic development of *Paradiaptomus greeni* (Gurney, 1906) (Copepoda, Calanoida) reared in the laboratory. *Crustaceana* **1989**, *56*, 141–161. [[CrossRef](#)]
76. Turki, S.; Turki, B. Copepoda and Branchiopoda from Tunisian temporary waters. *Int. J. Biodivers. Conserv.* **2010**, *2*, 86–97.
77. Kulkarni, M.R.; Pai, K. The freshwater diaptomid copepod fauna (Crustacea: Copepoda: Diaptomidae) of the Western Ghats of Maharashtra with notes on distribution, species richness and ecology. *J. Limnol.* **2016**, *75*, 135–143. [[CrossRef](#)]
78. Burns, C.W. New records of distribution and co-occurrence of freshwater calanoid copepods in New Zealand (Note). *N. Z. J. Mar. Freshw. Res.* **1987**, *21*, 149–150. [[CrossRef](#)]