

Mucopolysaccharidosis type IV

Description

Mucopolysaccharidosis type IV is an inherited disorder characterized by delayed development and vision impairment that worsens over time. The severe form of the disorder is called typical mucopolysaccharidosis type IV, and the mild form is called atypical mucopolysaccharidosis type IV.

Approximately 95 percent of individuals with this condition have the severe form. People with typical mucopolysaccharidosis type IV have delayed development of mental and motor skills (psychomotor delay). Motor skills include sitting, standing, walking, grasping objects, and writing. Psychomotor delay is moderate to severe and usually becomes apparent during the first year of life. Affected individuals have intellectual disability, limited or absent speech, difficulty chewing and swallowing, weak muscle tone (hypotonia) that gradually turns into abnormal muscle stiffness (spasticity), and problems controlling hand movements. Most people with typical mucopolysaccharidosis type IV are unable to walk independently. In about 15 percent of affected individuals, the psychomotor problems worsen over time.

Vision may be normal at birth in people with typical mucopolysaccharidosis type IV, but it becomes increasingly impaired during the first decade of life. Individuals with this condition develop clouding of the clear covering of the eye (cornea) and progressive breakdown of the light-sensitive layer at the back of the eye (retina). By their early teens, affected individuals have severe vision loss or blindness.

People with typical mucopolysaccharidosis type IV also have impaired production of stomach acid (achlorhydria). Achlorhydria does not cause any symptoms in these individuals, but it does result in unusually high levels of gastrin in the blood. Gastrin is a hormone that regulates the production of stomach acid. Individuals with mucopolysaccharidosis type IV may not have enough iron in their blood, which can lead to a shortage of red blood cells (anemia). People with the severe form of this disorder usually survive to adulthood; however, they may have a shortened lifespan.

About 5 percent of affected individuals have atypical mucopolysaccharidosis type IV. These individuals usually have mild psychomotor delay and may develop the ability to walk. People with atypical mucopolysaccharidosis type IV tend to have milder eye abnormalities than those with the severe form of the disorder. Achlorhydria also may be present in mildly affected individuals.

Frequency

Mucopolysaccharidosis type IV is estimated to occur in 1 in 40,000 people. About 70 percent of affected individuals have Ashkenazi Jewish ancestry.

Causes

Mutations in the *MCOLN1* gene cause mucopolysaccharidosis type IV. This gene provides instructions for making a protein called mucopolysaccharin-1. This protein is located in the membranes of lysosomes and endosomes, compartments within the cell that digest and recycle materials. While its function is not completely understood, mucopolysaccharin-1 plays a role in the transport (trafficking) of fats (lipids) and proteins between lysosomes and endosomes. Mucopolysaccharin-1 appears to be important for the development and maintenance of the brain and retina. In addition, this protein is likely critical for normal functioning of the cells in the stomach that produce digestive acids.

Most mutations in the *MCOLN1* gene result in the production of a nonfunctional protein or prevent any protein from being produced. A lack of functional mucopolysaccharin-1 impairs transport of lipids and proteins, causing these substances to build up inside lysosomes. Conditions that cause molecules to accumulate inside the lysosomes, including mucopolysaccharidosis type IV, are called lysosomal storage disorders. Two mutations in the *MCOLN1* gene account for almost all cases of mucopolysaccharidosis type IV in people with Ashkenazi Jewish ancestry. It remains unclear how mutations in this gene lead to the signs and symptoms of mucopolysaccharidosis type IV.

[Learn more about the gene associated with Mucopolysaccharidosis type IV](#)

- MCOLN1

Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- Ganglioside sialidase deficiency
- ML4
- MLIV
- Sialolipidosis

Additional Information & Resources

Genetic Testing Information

- Genetic Testing Registry: Mucopolidosis type IV (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0238286/>)

Genetic and Rare Diseases Information Center

- Mucopolidosis type IV (<https://rarediseases.info.nih.gov/diseases/94/index>)

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

Clinical Trials

- ClinicalTrials.gov ([https://clinicaltrials.gov/search?cond=%22Mucopolidosis type IV%22](https://clinicaltrials.gov/search?cond=%22Mucopolidosis+type+IV%22))

Catalog of Genes and Diseases from OMIM

- MUCOLIPIDOSIS IV; ML4 (<https://omim.org/entry/252650>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28mucopolidosis+type+iv%5BTIAB%5D%29+OR+%28mliv%5BTIAB%5D%29+OR+%28ml4%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>)

References

- Altarescu G, Sun M, Moore DF, Smith JA, Wiggs EA, Solomon BI, Patronas NJ, Frei KP, Gupta S, Kaneshi CR, Quarrell OW, Slaugenhaupt SA, Goldin E, Schiffmann R. The neurogenetics of mucopolidosis type IV. *Neurology*. 2002 Aug 13;59(3):306-13. doi: 10.1212/wnl.59.3.306. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12182165>)
- Dong XP, Cheng X, Mills E, Delling M, Wang F, Kurz T, Xu H. The type IV mucopolidosis-associated protein TRPML1 is an endolysosomal iron release channel. *Nature*. 2008 Oct 16;455(7215):992-6. doi: 10.1038/nature07311. Epub 2008 Sep 14. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18794901>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC430>)

1259/)

- Miedel MT, Rbaibi Y, Guerriero CJ, Colletti G, Weixel KM, Weisz OA, Kiselyov K. Membrane traffic and turnover in TRP-ML1-deficient cells: a revised model formucopolipidosis type IV pathogenesis. *J Exp Med*. 2008 Jun 9;205(6):1477-90. doi: 10.1084/jem.20072194. Epub 2008 May 26. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18504305>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2413042/>)
- Misko A, Grishchuk Y, Goldin E, Schiffmann R. Mucopolipidosis IV. 2005 Jan 28[updated 2021 Feb 11]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews(R)* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1214/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301393>)
- Puertollano R, Kiselyov K. TRPMLs: in sickness and in health. *Am J Physiol Renal Physiol*. 2009 Jun;296(6):F1245-54. doi: 10.1152/ajprenal.90522.2008. Epub 2009 Jan 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19158345>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2692446/>)
- Ruivo R, Anne C, Sagne C, Gasnier B. Molecular and cellular basis of lysosomal transmembrane protein dysfunction. *Biochim Biophys Acta*. 2009 Apr;1793(4):636-49. doi: 10.1016/j.bbamcr.2008.12.008. Epub 2008 Dec 24. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19146888>)
- Venugopal B, Mesires NT, Kennedy JC, Curcio-Morelli C, Laplante JM, Dice JF, Slaugenhaupt SA. Chaperone-mediated autophagy is defective in mucopolipidosis type IV. *J Cell Physiol*. 2009 May;219(2):344-53. doi: 10.1002/jcp.21676. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19117012>)
- Vergarajauregui S, Connelly PS, Daniels MP, Puertollano R. Autophagic dysfunction in mucopolipidosis type IV patients. *Hum Mol Genet*. 2008 Sep 1;17(17):2723-37. doi: 10.1093/hmg/ddn174. Epub 2008 Jun 11. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18550655>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2515373/>)
- Vergarajauregui S, Oberdick R, Kiselyov K, Puertollano R. Mucolipin 1 channel activity is regulated by protein kinase A-mediated phosphorylation. *Biochem J*. 2008 Mar 1;410(2):417-25. doi: 10.1042/BJ20070713. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17988215>)
- Vergarajauregui S, Puertollano R. Mucopolipidosis type IV: the importance of functional lysosomes for efficient autophagy. *Autophagy*. 2008 Aug;4(6):832-4. doi: 10.4161/auto.6567. Epub 2008 Jul 8. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18635948>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2625312/>)
- Wakabayashi K, Gustafson AM, Sidransky E, Goldin E. Mucopolipidosis type IV: an update. *Mol Genet Metab*. 2011 Nov;104(3):206-13. doi: 10.1016/j.ymgme.2011.06.006. Epub 2011 Jun 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21763169>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3205274/>)

Last updated August 1, 2013