

## Myotonia congenita

### Description

Myotonia congenita is a disorder that affects muscles used for movement (skeletal muscles). Beginning in childhood, people with this condition experience bouts of sustained muscle tensing (myotonia) that prevent muscles from relaxing normally. Although myotonia can affect any skeletal muscles, including muscles of the face and tongue, it occurs most often in the legs. Myotonia causes muscle stiffness that can interfere with movement. In some people the stiffness is very mild, while in other cases it may be severe enough to interfere with walking, running, and other activities of daily life. These muscle problems are particularly noticeable during movement following a period of rest. Many affected individuals find that repeated movements can temporarily alleviate their muscle stiffness, a phenomenon known as the warm-up effect.

The two major types of myotonia congenita are known as Thomsen disease and Becker disease. These conditions are distinguished by the severity of their symptoms and their patterns of inheritance. Becker disease usually appears later in childhood than Thomsen disease and causes more severe muscle stiffness, particularly in males. People with Becker disease often experience temporary attacks of muscle weakness, particularly in the arms and hands, brought on by movement after periods of rest. They may also develop mild, permanent muscle weakness over time. This muscle weakness is not seen in people with Thomsen disease.

### Frequency

Myotonia congenita is estimated to affect 1 in 100,000 people worldwide. This condition is more common in northern Scandinavia, where it occurs in approximately 1 in 10,000 people.

### Causes

Mutations in the *CLCN1* gene cause myotonia congenita. The *CLCN1* gene provides instructions for making a protein that is critical for the normal function of skeletal muscle cells. For the body to move normally, skeletal muscles must tense (contract) and relax in a coordinated way. Muscle contraction and relaxation are controlled by the flow of charged atoms (ions) into and out of muscle cells. Specifically, the protein produced from the *CLCN1* gene forms a channel that controls the flow of negatively charged chlorine atoms (chloride ions) into these cells. The main function of this channel is to

stabilize the cells' electrical charge, which prevents muscles from contracting abnormally.

Mutations in the *CLCN1* gene alter the usual structure or function of chloride channels. The altered channels cannot properly regulate ion flow, reducing the movement of chloride ions into skeletal muscle cells. This disruption in chloride ion flow triggers prolonged muscle contractions, which are the hallmark of myotonia.

[Learn more about the gene associated with Myotonia congenita](#)

- CLCN1

## **Inheritance**

The two forms of myotonia congenita have different patterns of inheritance. Thomsen disease is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder. In most cases, an affected person has one parent with the condition.

Becker disease is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. Most often, the parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but do not show signs and symptoms of the condition.

Because several *CLCN1* mutations can cause either Becker disease or Thomsen disease, doctors usually rely on characteristic signs and symptoms to distinguish the two forms of myotonia congenita.

## **Other Names for This Condition**

- Congenital myotonia

## **Additional Information & Resources**

### Genetic Testing Information

- Genetic Testing Registry: Congenital myotonia, autosomal dominant form (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2936781/>)
- Genetic Testing Registry: Congenital myotonia, autosomal recessive form (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0751360/>)
- Genetic Testing Registry: Batten-Turner congenital myopathy (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0027127/>)

### Genetic and Rare Diseases Information Center

- Thomsen and Becker disease (<https://rarediseases.info.nih.gov/diseases/12301/ind>)

ex)

### Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

### Clinical Trials

- ClinicalTrials.gov ([https://clinicaltrials.gov/search?cond=%22Myotonia congenita%22](https://clinicaltrials.gov/search?cond=%22Myotonia%20congenita%22))

### Catalog of Genes and Diseases from OMIM

- MYOTONIA CONGENITA, AUTOSOMAL DOMINANT (<https://omim.org/entry/160800>)
- MYOTONIA CONGENITA, AUTOSOMAL RECESSIVE (<https://omim.org/entry/255700>)

### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28Myotonia+Congenita%5BMAJR%5D%29+AND+%28%28myotonia+congenita%5BTIAB%5D%29+OR+%28Becker&#x27;s+myotonia%5BTIAB%5D%29+OR+%28Thomsen&#x27;s+disease%5BTIAB%5D%29+OR+%28Thomsen+disease%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>)

### **References**

- Chrestian N, Puymirat J, Bouchard JP, Dupre N. Myotonia congenita--a cause of muscle weakness and stiffness. *Nat Clin Pract Neurol.* 2006 Jul;2(7):393-9; quiz following 399. doi: 10.1038/ncpneuro0239. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16932590>)
- Colding-Jorgensen E. Phenotypic variability in myotonia congenita. *Muscle Nerve.* 2005 Jul;32(1):19-34. doi: 10.1002/mus.20295. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15786415>)
- Duno M, Vissing J. Myotonia Congenita. 2005 Aug 3 [updated 2021 Feb 25]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews(R)* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1355/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301529>)
- Pusch M. Myotonia caused by mutations in the muscle chloride channel gene CLCN1. *Hum Mutat.* 2002 Apr;19(4):423-34. doi: 10.1002/humu.10063. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11933197>)

- Sun C, Tranebjaerg L, Torbergesen T, Holmgren G, Van Ghelue M. Spectrum of CLCN1 mutations in patients with myotonia congenita in Northern Scandinavia. *Eur J Hum Genet.* 2001 Dec;9(12):903-9. doi: 10.1038/sj.ejhg.5200736. Erratum In: *Eur J Hum Genet.* 2010 Feb;18(2):264. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11840191>)
- Zhang J, George AL Jr, Griggs RC, Fouad GT, Roberts J, Kwiecinski H, Connolly AM, Ptacek LJ. Mutations in the human skeletal muscle chloride channel gene (CLCN1) associated with dominant and recessive myotonia congenita. *Neurology.* 1996 Oct;47(4):993-8. doi: 10.1212/wnl.47.4.993. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/8857733>)

**Last updated February 1, 2020**