Seeds for Sowing

155.02.05

21 February 2022

TITLE

Import Health Standard: Seeds for Sowing

COMMENCEMENT

This consolidated import health standard comes into force on 21 February 2022

This import health standard amends the import health standard *Seeds for Sowing (155.02.05)*, which came into force on 15 February 2021 and consolidates all amendments made up to commencement of this standard.

The amendment history to this import health standard is set out in Appendix 2.

ISSUING AUTHORITY

This import health standard is issued under section 24A of the Biosecurity Act 1993 and incorporates amendments made in accordance with section 24B(1)(a) of that Act.

Dated at Wellington, 21 February 2022

Director, Animal & Plant Health
Ministry for Primary Industries
(acting under delegated authority of the Director-General)

Contact for further information Ministry for Primary Industries (MPI) Animal and Plant Health Directorate Plant Imports PO Box 2526 Wellington 6140

Email: plantimports@mpi.govt.nz

Ministry for Primary Industries Page 1 of 168

Cont	Contents	
Introdu	ction	5
David 4.	Con and Descriptors and	7
Part 1:	General Requirements	7
1.1 1.2	Application	7
1.2	Incorporation of material by reference Definitions	7
1.3 1.4		7 7
1.4	Requirements for seed for sowing Documentation	9
1.6		11
1.7	Post - entry quarantine Seed for sowing of New Zealand origin	11
1.7		12
1.0	Seed for sowing imported as laboratory specimens Seed imported as pelleted seed	12
1.9	Seed imported as pelieted seed	12
Part 2:	Specific Requirements	16
2.1	Abies	17
2.2	Acer	18
2.3	Acrocomia	19
2.4	Actinidia	20
2.5	Agropyron	22
2.6	Agrostis	23
2.7	Arabidopsis thaliana	24
2.8	Avena	25
2.9	Beta Pression repus	29
2.10 2.11		30 31
	Camellia sinensis Camissonia	32
	Cannabis sativa	33
2.13		35
	Carpinus	38
2.16	Carthamus tinctorius	39
2.17		40
2.18	Castanea	41
2.19	Cicer	42
2.20	Citrus	43
2.21	Cocos	44
2.22	Coffea	45
2.23	Coriandrum	46
2.24	Corylus	47
2.25	•	48
2.26		49
2.27	Cucurbitaceae	52
2.28	Cuminum	56
2.29	Desmodium	57
2.30	Echinochloa	58
2.31	Elaeis	59
2.32	Eriobotrya	60
2.33	Fagus	61

Ministry for Primary Industries Page 2 of 168

2.36 Glycine 66 2.37 Gossypium 68 2.38 Helianthus 69 2.39 Hordeum 71 2.40 Humulus 75 2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 95 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Phoeni	2.34	Fragaria	62
2.37 Gossypium 68 2.38 Helianthus 69 2.39 Hordeum 71 2.40 Humulus 75 2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 10 2.63 Pinus 104 2.65 Populus 113 2.66 Populus 113	2.35	Glebionis	65
2.37 Gossypium 68 2.38 Helianthus 69 2.39 Hordeum 71 2.40 Humulus 75 2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 101 2.65 Populus 111 2.66 Populus 113 2.67 Pseudotsuga menziesii 116<	2.36	Glycine	66
2.38 Helianthus 69 2.39 Hordeum 71 2.40 Humulus 75 2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 104 2.65 Populus 113 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.71 Raphanus sativus 123 </td <td>2.37</td> <td>Gossypium</td> <td>68</td>	2.37	Gossypium	68
2.40 Humulus 75 2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lainum usitatissimum 80 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64	2.38		69
2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densifilorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pisum 110 2.64 Pisum 110 2.65 Psopulus<	2.39	Hordeum	71
2.41 Juglans 76 2.42 Lablab 77 2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus <td>2.40</td> <td>Humulus</td> <td>75</td>	2.40	Humulus	75
2.43 Lavandula 78 2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pisum 110 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus<	2.41	Juglans	76
2.44 Lens 79 2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 111 2.66 Prunus 112 2.67 Pseudotsuga menziesii 114 2.68 Psophocarpus 120 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum Uycopersicum	2.42		77
2.45 Linum usitatissimum 80 2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 111 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 121 2.70 </td <td>2.43</td> <td>Lavandula</td> <td>78</td>	2.43	Lavandula	78
2.46 Lithocarpus densiflorus 81 2.47 Livistona 82 2.48 Lophophora williamsii 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 93 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 104 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72<	2.44	Lens	79
2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 94 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 104 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Rib	2.45	Linum usitatissimum	80
2.47 Livistona 82 2.48 Lophophora williamsii 83 2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 104 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 121 2.71 Raphanus sativus 122 2.72 Ribes 126 2.73 Rubus	2.46	Lithocarpus densiflorus	81
2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 94 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Pyrus 121 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 <td>2.47</td> <td></td> <td>82</td>	2.47		82
2.49 Lotus 84 2.50 Macadamia 85 2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 94 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus	2.48	Lophophora williamsii	83
2.51 Malus 86 2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 126 2.74 Sesamum 130 2.75 Solanum lycopersicum<	2.49		84
2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Propulus 113 2.67 Pseudotsuga menziesii 114 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 121 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 126 2.74 Sesamum 130 2.75 Solanum lycopersicum 131 2.76 Sor	2.50	Macadamia	85
2.52 Mangifera 88 2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 114 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 126 2.74 Sesamum 130 2.75 Solanum lycopersicum 131 2.76 Solan	2.51	Malus	
2.53 Medicago 89 2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Popullus 111 2.66 Prunus 114 2.67 Pseudotsuga menziesii 114 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum lycopersicum 131 2.78 Sorghum 137 2.79 Steno			
2.54 Myrtaceae 90 2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum luberosum 131 2.76 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trig		· ·	
2.55 Nicotiana tabacum 91 2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 111 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum tuberosum 131 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigi			
2.56 Oxyria 93 2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 111 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 T	2.55		
2.57 Panicum 94 2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2	2.56		93
2.58 Papaver somniferum 95 2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 125 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 144			
2.59 Persea 96 2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 131 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.60 Petunia 97 2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.61 Phaseolus 99 2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.62 Phoenix 103 2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144	2.61		
2.63 Pinus 104 2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144	2.62	Phoenix	
2.64 Pisum 110 2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.65 Populus 113 2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.66 Prunus 114 2.67 Pseudotsuga menziesii 116 2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144	2.66		
2.68 Psophocarpus 120 2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144	2.67	Pseudotsuga menziesii	116
2.69 Pyrus 121 2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144		· ·	
2.70 Quercus 123 2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.71 Raphanus sativus 125 2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144		·	
2.72 Ribes 126 2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.73 Rubus 128 2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.74 Sesamum 130 2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.75 Solanum 131 2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.76 Solanum lycopersicum 133 2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.77 Solanum tuberosum 136 2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.78 Sorghum 137 2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.79 Stenotaphrum 138 2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144			
2.80 Trigonella foenum-graecum 139 2.81 Triticum 140 2.82 Ulmus 144		· ·	
2.81 Triticum 140 2.82 Ulmus 144		·	
2.82 <i>Ulmus</i> 144		· · · · · · · · · · · · · · · · · · ·	

Ministry for Primary Industries Page 3 of 168

2.84 2.85 2.86 2.87	Vicia Vigna Vitis Zea	148 150 151 153
Appendi	x 1: Definitions	160
Appendi	x 2: Amendment Record	163
Appendi	x 3: Declaration Form	167
Appendi	x 4: Species on the Plant Biosecurity Index eligible for import into New 2 pelleted seeds for sowing	Zealand as 168



Ministry for Primary Industries Page 4 of 168

Introduction

This introduction is not part of the import health standard (IHS), but is intended to indicate its general effect.

Purpose

This IHS specifies the requirements for the importation of seeds intended for sowing in New Zealand.

Background

The New Zealand Biosecurity Act 1993 provides the legal basis for excluding, eradicating and effectively managing pests and unwanted organisms.

Each IHS issued under the Act specifies requirements to be met for the effective management of risks associated with imported goods that may pose a biosecurity threat to New Zealand. This IHS includes requirements that must be met in the exporting country, during transit and importation, and post clearance if specified before biosecurity clearance is given.

Additional information to the requirements is included in guidance text boxes.

Who should read this import health standard?

This IHS should be read by anyone involved in the process of importing seeds for sowing into New Zealand from all countries and outlines the import requirements that must be met.

Importers of seed for laboratory testing, analysis or research (where biosecurity clearance is not required) should refer to either this IHS or IHS MPI.STD.PLANTMATERIAL: *Dried & Preserved Plant Material, & Fresh Plant Material for Testing, Analysis or Research.*

Products containing viable seed that also contain organic growing media must also meet the requirements of the relevant IHS: MPI.STD.FERTGRO: Fertilisers and Growing Media of Plant Origin.

Why is this important?

It is the importers responsibility to ensure the requirements of this IHS are met. Consignments that do not comply with the requirements of this IHS may not be cleared for entry into New Zealand and /or further information may be sought from importers.

Importers are liable for all associated expenses.

Equivalence

MPI may consider a pre-export application for an equivalent phytosanitary measure to be approved, different from that provided for in this IHS, if in the opinion of the Director-General, it is considered to be equivalent to the current measures taken for managing the risks associated with the importation of those goods.

Equivalence will be considered with reference to the International Standard for Phytosanitary Measures (ISPM), Publication No. 24: *Guidelines for the determination and recognition of equivalence of phytosanitary measures (2011).*

Document history

Refer to Appendix 2 for the amendment record for this IHS.

Ministry for Primary Industries Page 5 of 168

Other Information

This is not an exhaustive list of compliance requirements and it is the importer's responsibility to be familiar with and comply with all New Zealand laws.


Listed below are other New Zealand legislative requirements which may also apply to seeds for sowing.

Importers of *Cannabis sativa*, *Lophophora williamsii and Papaver somniferum* must contact Medsafe before importation for advice on licensing:

Medicines Control Medsafe

Email: medicinescontrol@health.govt.nz

Phone: (04) 816 2444



Part 1: General Requirements

1.1 Application

- (1) This import health standard (IHS) applies to viable seed, and products containing viable seed (including pre-germinated seeds as defined in Appendix 1) from species that are listed in the MPI Plant Biosecurity Index, that are imported for the purpose of planting.
- (2) This IHS does not apply to species listed as "requires assessment" or "prohibited entry" in the MPI Plant Biosecurity Index, which are not eligible to be imported under this standard.
- (3) This IHS applies to seed for sowing from any country, unless otherwise specified in Part 2: Specific Requirements.

Guidance

A guide to importing seeds for sowing can be found on the MPI website.

1.2 Incorporation of material by reference

- (4) The following documents are incorporated by reference under section 142M of the Act;
 - a) <u>ISPM 1 Phytosanitary principles for the protection of plants and the application of phytosanitary</u> measures in international trade. Rome, IPPC, FAO;
 - b) ISPM 4 Requirements for the establishment of pest free areas. Rome, IPPC, FAO;
 - c) ISPM 5 Glossary of phytosanitary terms. Rome, IPPC, FAO;
 - d) <u>ISPM 10 Requirements for the establishment of pest free places of production and pest free production sites.</u> Rome, IPPC, FAO;
 - e) ISPM 12 Guidelines for phytosanitary certificates. Rome, IPPC, FAO;
 - f) ISPM 20 Guidelines for a phytosanitary import regulatory system. Rome, IPPC, FAO;
 - g) ISPM 38 International movement of seeds. Rome, IPPC, FAO;
 - h) MPI Official New Zealand Pest Register (ONZPR);
 - i) MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments;
 - j) MPI Schedule of Regulated (Quarantine) Weed Seeds;
 - k) MPI Plants Biosecurity Index (PBI);
 - I) MPI Protocol for Testing for the Presence of Genetically Modified Plant Material;
- (5) Under section 142O(3) of the Act it is declared that section 142O(1) does not apply, that is, a notice under section 142O(2) of the Act is not required to be published before material that amends or replaces any material incorporated by reference has legal effect as part of those documents.

1.3 Definitions

(1) Definitions that apply to this IHS are listed in Appendix 1.

1.4 Requirements for seed for sowing

- (1) On arrival in New Zealand, seeds must be made available for inspection and examination by MPI inspectors at the importer's expense.
- (2) Seed for sowing must meet the following requirements:
 - all seed for sowing must be clearly identified with the scientific name (i.e. genus and species);
 - b) all hybrid species must be clearly identified as hybrid species, and either:

Ministry for Primary Industries Page 7 of 168

- i) listed in the MPI Plant Biosecurity Index as hybrid species eligible to be imported; or
- ii) identified by the scientific names (genus and species) of both parents;
- all packaging associated with seed for sowing must be clean, free from soil and other contaminants:
- d) all seed for sowing from fleshy fruits (including pods) must have all traces of flesh removed, except:
 - i) Orchidaceae seed (which may be imported in dry/green pods); or
 - ii) when otherwise stated in Part 2: Specific Requirements.
- e) seed for sowing must be free from the following:
 - i) regulated pest(s) including any quarantine pest(s) as listed in Part 2: Specific Requirements;
 - ii) soil particles greater than 0.1% by weight;
 - iii) unidentified seed;
- (3) Quarantine weed seed contamination must not exceed the MPL of 0.01%. To achieve 95% confidence that the MPL (of 0.01% probability) will not be exceeded, no quarantine weeds seeds are permitted (i.e acceptance = No. = 0) in a sample(s) drawn and analysed by a MPI approved method (e.g. ISTA sampling methodology as approved by MPI).
- (4) All importers of seed for sowing must make a <u>declaration</u> to identify whether they are importing pelleted seeds or not.

Guidance

- If undeclared seeds are detected during inspection the seed consignment will be held until an assessment has been made and MPI determine the biosecurity risk of the seeds.
- The Maximum Pest Limit (MPL) for visually detectable regulated pests on seed for sowing is, at a 95% confidence level, not more than 0.5% of the units in the consignment are infested:
 - this equates to an acceptable level of zero units infested by regulated pests in a sample size of a minimum of 5kg.
- Each line of seed must undergo inspection to verify that the seed and associated documentation is compliant with the requirements of this IHS.
- An officially drawn 5 kilogram sample will be inspected for visually regulated pests from each line (or the whole line if less than 5kg).
- An officially drawn sample will be inspected for contaminants from each line as per MPI current procedures.
- A declaration form to be completed by all importers of seeds for sowing can be found on this link: Importer declaration for seed imports.

1.4.1 Requirements for seed for sowing species listed in the MPI <u>Plant Biosecurity Index</u> (PBI) as "Basic"

(1) Seed consignments must meet all the requirements of Part 1: General Requirements.

Guidance

• In order to facilitate processing, the importer may provide a phytosanitary certificate, which must meet the requirements set under Part 1.5.2 of this IHS.

Ministry for Primary Industries Page 8 of 168

1.4.2 Requirements for seed for sowing listed in the PBI with specific import requirements

(1) Seed for Sowing under the import specification as "see 155.02.05 under... " must meet all the requirements of Part 1: General Requirements and any specific requirements in Part 2: Specific Requirements.

1.4.3 Treatment Requirements

- (1) Any approved phytosanitary treatment, as required in Part 2: Specific Requirements, must be completed either:
 - a) offshore prior to export; or
 - b) on arrival in New Zealand, at an MPI approved facility at the importer's expense.
- (2) If performed offshore, the exporting country NPPO must confirm that any treatment(s) as required in Part 2: Specific Requirements is endorsed in the disinfestation and/or disinfection treatment section of the phytosanitary certificate.

1.5 Documentation

1.5.1 Permit to Import

- (1) A permit to import is required if specified in Part 2: Specific Requirements.
- (2) Any permit, as required in Part 2 or Part 1.8, must be presented to MPI at the New Zealand Border together with all other required documentation.

1.5.2 Phytosanitary certificate

- (1) A phytosanitary certificate is required if specified in Part 2: Specific Requirements.
- (2) For consignments arriving with a phytosanitary certificate, the phytosanitary certificate must be issued by the exporting country National Plant Protection Organisation (NPPO) in accordance with **ISPM 12**: *Phytosanitary certificates*.
- (3) The phytosanitary certificate must include any additional declaration(s) required under Part 2: Specific Requirements.
- (4) The phytosanitary certificate must certify that the seed has been inspected in accordance with appropriate official procedures and found to be free of any visually detectable regulated pests and conforms to New Zealand's import requirements.
- (5) If pests are detected that are not listed in this IHS or <u>ONZPR</u>, the exporting country NPPO must establish their regulatory status by contacting MPI prior to issuing the certificate.
- (6) The exporting country NPPO must not issue a phytosanitary certificate if a viable regulated pest is detected, unless the consignment is treated in order to eliminate it.
- (7) If the exporting NPPO is satisfied that the pre-shipment inspection activities have been carried out effectively the following declaration, or a variation that is compliant with **ISPM 12**: *Phytosanitary certificates* and has been approved by MPI prior to shipment must be included on the phytosanitary certificate:
 - "This is to certify that the plants, plant products or other regulated articles described herein have been inspected and/or tested according to appropriate official procedures and are considered to be free from the quarantine pests specified by the importing contracting party and to conform with the current phytosanitary requirements of the importing contracting party, including those for regulated non-quarantine pests."

Ministry for Primary Industries Page 9 of 168

Guidance

Phytosanitary certificate

- Information about the regulated pests for New Zealand is available in <u>ONZPR</u>.
- The phytosanitary certificate should contain sufficient detail to enable identification of the consignment and its component parts. Information should include:
 - lot number(s);
 - number and description of packages;
 - country/place of origin of the seed; and
 - variety name(s).

Seed analysis certificate (SAC)

- In order to facilitate processing, the importer may provide a SAC (original or PDF copy), which must meet the following criteria:
 - be issued by an ISTA or AOSA accredited seed testing station, or an accredited laboratory that follows the ISTA or AOSA methodology;
 - state the actual weight of the sample examined;
 - state the seed lot number and aligns with traceability information on the phytosanitary certificate (for seeds where phytosanitary certificate is presented);
 - be endorsed that the minimum size of the sample examined was as prescribed for the determination of other species by number in ISTA (as published in Seed Science and Technology 24, 1996);
 - state the botanical name of each identified species of seed or nematode gall found in the sample (any unidentified genera or species are to be recorded as such);
 - give the percentage of soil particles present in the sample;
 - certify that no quarantine weed seeds were present in the sample.

1.5.3 Genetically modified testing certificate

- (1) Genetically modified (GM) testing certificates are required for all consignments of *Brassica napus var.* oleifera, Glycine max, Gossypium hirsutum, Medicago sativa, Zea mays var. indentata and Zea mays var. saccharata, unless stated in the MPI *Protocol for Testing for the Presence of Genetically Modified Plant Material*.
- (2) A GM testing certificate is required for *Cucurbita pepo* (see <u>Cucurbitaceae</u>), *Petunia* and <u>Linum usitatissimum</u> unless a non-GM declaration is provided. <u>Cucurbitaceae Linum usitatissimum</u>
- (3) GM testing certificates must:
 - a) state the sampling method used for each seed line (e.g. automatic in-line machine);
 - b) contain the same lot number or unique identifier as stated on all the other import documentation for consignments arriving in New Zealand;
- (4) Testing must be conducted by facilities approved by MPI and a copy of the completed test certificate must accompany the consignment imported into New Zealand.
- (5) Importers must provide all test records when required by an MPI inspector.

Guidance

- A non-GM declaration template is provided in <u>Appendix 3</u>. Refer to specific requirements for each species in Part 2. <u>Appendix 3</u>: <u>Declaration</u>
- Complete guidelines for sampling and testing for the presence of GM seeds are specified in the MPI
 <u>Protocol</u> for Testing for the Presence of Genetically Modified Plant Material. The MPI <u>Protocol</u> for
 Testing for the Presence of Genetically Modified Plant Material and a list of MPI approved facilities for
 testing for genetically modified plant material can be found on MPI website under <u>Genetically modified</u>
 seeds.

Ministry for Primary Industries Page 10 of 168

- MPI will examine the test certificates on arrival to confirm that they reconcile with the actual seed for sowing.
- If consignments arrive at the border without having been tested for the presence of GM seeds, MPI will
 offer the importer the options of re-shipment, destruction, or having the consignment sampled and
 tested according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material
 at the importer's expense.
- Any consignment that is found to contain unapproved GM seeds will not be permitted to enter New
 Zealand and will be re-shipped or destroyed, unless the importer obtains an approval to grow the GM
 variety from the Environmental Protection Authority (EPA).
- All test results must be available to MPI on request.

1.6 Post - entry quarantine

- (1) Seed for sowing must be imported into a post entry quarantine (PEQ) facility if required by Part 2: Specific Requirements.
- (2) The transitional (PEQ) facility must be approved to the MPI Facility Standard: Post Entry Quarantine for Plants (MPI.STD.PEQ).
- (3) Seed for sowing must be actively growing during the quarantine period, and must be tested, treated or inspected for regulated pests at the importer's expense.
- (4) Testing must be undertaken by a diagnostic laboratory approved by MPI to <u>155.04.03</u>: <u>Standard for Transitional Facilities for the Identification of Organisms</u>.

1.6.1 Testing

- (1) The unit for testing is defined as an individual seedling and each seedling must be labelled individually and tested separately, unless one of the following methods has been used:
 - a) Polymerase chain reaction:
 - i) samples taken from up to five seedlings of the same species growing in PEQ can be combined to form a single composite sample for pre-determined testing by polymerase chain reaction (PCR).
 - b) Enzyme-linked immunosorbent assay:
 - for viruses that are not pollen transmitted, samples taken from up to five seedlings can be combined to form a single composite sample for enzyme-linked immunosorbent assay (ELISA) testing:
 - ii) the phytosanitary certificate must be endorsed with an additional declaration (AD) stating that the seeds have been derived from the same parent plant.
 - c) Graft (woody) indexing:
 - i) where prior permission is received from MPI, samples taken from up to five seedlings can be combined to form a single composite sample for testing by graft indexing;
 - ii) the phytosanitary certificate must be endorsed with an AD stating that the seeds have been derived from the same parent plant.

1.7 Seed for sowing of New Zealand origin

- (1) Importers must make the seeds and its documentation available for inspection by an MPI inspector.
- (2) Seed for sowing exported from New Zealand, given clearance into the importing country or rejected prior to clearance, may be returned to New Zealand under one of the following circumstances:

Ministry for Primary Industries Page 11 of 168

1.7.1 Seed for sowing unopened offshore:

(1) Product that remains in its original and unopened packaging must be accompanied with a re-export phytosanitary certificate issued by the NPPO of the overseas country.

1.7.2 Seed for sowing opened offshore:

- (1) Seed that has been grown in New Zealand, exported to another country, packaged in that same country and returned to New Zealand, must be accompanied with:
 - a) the original or a copy of the New Zealand issued phytosanitary certificate,
 - b) an export bill of lading; and
 - c) a declaration from the overseas packaging company manager; stating that:
 - the re-exported seed is the same seed as covered by the attached phytosanitary certificate and bill of lading; and
 - ii) the quality system used by the company ensures that the seed is not contaminated by any other seed lots, residues from grading or packaging machines, or storage pests.

Guidance

 Inspection is required by an MPI inspector to confirm that packaging and labelling is consistent with the documentation provided.

1.8 Seed for sowing imported as laboratory specimens

Guidance

Import requirements for research samples can be found in the <u>Research Samples (excluding animal samples)</u> import health standard.

1.9 Seed imported as pelleted seed

- (1) Only species listed in Appendix 4: Seed species on the Plant Biosecurity Index eligible for import into New Zealand of IHS 155.02.05: Seeds for sowing may be imported into New Zealand as pelleted seeds for sowing from all countries.
- (2) Pelleted seed lots of the following species must have a representative sample officially drawn and sealed (according to ISTA methodology) from each lot and tested for purity at a MPI-approved laboratory for the presence of quarantine weed seeds and other contaminants.
 - a) Beta vulgaris
 - b) Allium cepa
 - c) Allium porrum
 - d) Allium ampeloprasum
 - e) Apium graveolens
 - f) Brassica napus
 - g) Brassica oleracea
 - h) Cichorium intybus
 - i) Cichorium endivia
 - j) Daucus carota
 - k) Foeniculum vulgare
 - Pastinaca sativa
 - m) Spinacia oleracea

Ministry for Primary Industries Page 12 of 168

- (3) The representative seed sample of species listed in Part 1.9(2) must comply with the applicable requirements in Parts 1.9(9) to 1.9(11) of this IHS.
- (4) Pelleted seed lots of *Solanum lycopersicum* species imported for the purpose of rootstock are not required to undergo onshore purity testing.
- (5) If requested by an MPI inspector, pelleted seed lots of all other species listed in Appendix 4 must have a representative sample officially drawn and sealed (according to ISTA methodology) from each lot and tested for purity at a MPI-approved laboratory for the presence of quarantine weed seeds and other contaminants.
- (6) If a pelleted seed lot of *Lactuca sativa* is requested by the MPI inspector to have a representative sample officially drawn and sealed (according to ISTA methodology) and tested for purity at a MPI-approved laboratory for the presence of quarantine weed seeds and other contaminants, the representative sample must comply with the applicable requirements of Part 1.9(12) of this IHS.
- (7) All imported pelleted seed lots that are required to be tested for purity onshore must be held in a MPI-approved transitional facility pending the return of the laboratory results.
- (8) All importers of *Beta vulgaris* (fodder beet) must make a <u>declaration</u> that the *Beta vulgaris* seed in their consignments is not one of the specific seed lots of the following varieties of *Beta vulgaris* var *rapacea* Kyros 16UB128, Bangor 16UB126, Bangor 15UB079, Bangor 16UB114, Feldherr 16UB131 or Troya 16UB112.

Beta vulgaris seed grown in all countries except Italy

- (9) The representative sample for pelleted *Beta vulgaris* seed grown in all countries except Italy must comply with either option 1 or option 2:
 - a) **Option 1:** A 'bare' seed sample of at least 31,540 seeds accompanying the pelleted seed lot, which includes an official label issued by the ISTA-accredited seed laboratory stating the information below will be sent for seed analysis at a MPI approved laboratory at the importers expense:
 - i) that the seeds have been sampled, labelled and sealed according to ISTA rules;
 - ii) the same lot/line number or unique identifier as stated on the pelleted seed lot;
 - iii) the species and variety name;
 - iv) the sample weight, and
 - v) the date, name and signature of the ISTA accredited/approved sampler.
 - b) **Option 2:** For individual lots of pelleted seed not accompanied by a bare seed sample, a representative sample of at least 31,540 pelleted seeds will be drawn according to ISTA methodology and sent for seed analysis at a MPI approved laboratory at the importers expense.

Beta vulgaris seed grown in Italy

- (10) The representative sample of pelleted *Beta vulgaris* seed grown in Italy must comply with either option 1 or option 2.
 - a) Option 1: A 'bare' seed sample of at least 48,480 seeds accompanying the pelleted seed lot, which includes an official label issued by the ISTA-accredited seed laboratory stating the information below will be sent for seed analysis at a MPI approved laboratory at the importers expense:
 - i) that the seeds have been sampled, labelled and sealed according to ISTA rules;
 - ii) the same lot/line number or unique identifier as stated on the pelleted seed lot;
 - iii) the species and variety name;
 - iv) the sample weight, and
 - v) the date, name and signature of the ISTA accredited/approved sampler.

Ministry for Primary Industries Page 13 of 168

b) **Option 2:** For individual lots of pelleted seed not accompanied by a bare seed sample, a representative sample of at least 48,480 pelleted seeds will be drawn according to ISTA methodology and sent for seed analysis at a MPI approved laboratory at the importers expense.

Vegetable species grown in all countries (Allium cepa, Allium porrum, Allium ampeloprasum, Apium graveolens, Brassica napus, Brassica oleracea, Cichorium intybus, Cichorium endivia, Daucus carota, Foeniculum vulgare, Pastinaca sativa and Spinacia oleracea)

- (11) The representative sample of pelleted vegetable species must comply with either option 1 or 2.
 - a) Option 1: A 'bare' seed sample of at least 31,540 seeds accompanying the pelleted seed lot, which includes an official label issued by the ISTA-accredited seed laboratory stating the information below will be sent for seed analysis at a MPI approved laboratory at the importers expense:
 - i) that the seeds have been sampled, labelled and sealed according to ISTA rules;
 - ii) the same lot/line number or unique identifier as stated on the pelleted seed lot;
 - iii) the species and variety name;
 - iv) the sample weight, and
 - v) the date, name and signature of the ISTA accredited/approved sampler.
 - b) Option 2:
 - i) For individual lots of pelleted seed of less than 300,000 seeds and not accompanied by a bare seed sample, a representative sample of at least 10% of the total size of each lot will be drawn according to ISTA methodology and sent for seed analysis at a MPI approved laboratory at the importers expense.
 - ii) For individual lots of pelleted seed of 300,000 pelleted seeds or greater, and not accompanied by a bare seed sample, a representative sample of at least 31,540 pelleted seeds will be drawn according to ISTA methodology and sent for seed analysis at a MPI approved laboratory at the importers expense.

Lettuce (Lactuca sativa)

- The representative sample for pelleted lettuce seeds must comply with either option 1 or option 2. The options for representative samples are:
 - a) Option 1: A 'bare' seed sample of at least 31,540 seeds accompanying the pelleted seed lot, which includes an official label issued by the ISTA-accredited seed laboratory stating the information below will be sent for seed analysis at a MPI approved laboratory at the importers expense:
 - i) that the seeds have been sampled, labelled and sealed according to ISTA rules;
 - ii) the same lot/line number or unique identifier as stated on the pelleted seed lot;
 - iii) the species and variety name;
 - iv) the sample weight, and
 - v) the date, name and signature of the ISTA accredited/approved sampler.
 - b) Option 2: For individual pelleted seed lots of lettuce that are not accompanied by a 'bare' seed sample, a representative sample will be officially drawn and sealed (according to ISTA methodology) and tested for purity at a MPI-approved laboratory for the presence of quarantine weed seeds and other contaminants.
 - i) For individual lots of pelleted seed of less than 300,000 seeds and not accompanied by a bare seed sample, a representative sample of at least 10% of the total size of each lot will be drawn according to ISTA methodology and sent for seed analysis at a MPI approved laboratory at the importers expense.
 - ii) For individual lots of pelleted seed of 300,000 pelleted seeds or greater, and not accompanied by a bare seed sample, a representative sample of at least 31,540 pelleted

Ministry for Primary Industries Page 14 of 168

seeds will be drawn according to ISTA methodology and sent for seed analysis at a MPI approved laboratory at the importers expense.

Guidance

- A declaration form to be completed by all importers of pelleted *Beta vulgaris* seed for sowing can be found on this link: Importer declaration for all importers of pelleted Beta vulgaris seeds for sowing.
- A risk analysis for importation as pelleted seed has been undertaken for the species listed in Appendix 4 of this IHS. Importers who wish to import species that are not listed in Appendix 4 as pelleted seed must contact MPI prior to import as further risk analysis may need to be undertaken for these species.
- The requirements for pelleted seed relate to the country of origin of the seed.
- Identification of a quarantine weed seed or a contaminant in a pelleted seed lot of species listed in Appendix 4 constitutes a non-conformance.
- An MPI inspector may request a sample of a lot of pelleted *Lactuca sativa* seed to be taken for purity testing under Part 1.9(6) of this IHS to validate phytosanitary certification that accompanied the lot.
 - Normally, the sampling regime will be, per importer, per species imported, for sampling and purity testing of one in ten lots (1/10) (randomly).
 - If an importer demonstrates significant compliance in ten consecutive compliance checks, the inspector may only require sampling and purity testing of one in every twenty lots (1/20) (randomly). However, if there is a non-compliance, the frequency of sampling and testing may go back to one in ten (1/10) lots.
 - If there is a non-compliance when the MPI inspector requests sampling and purity testing at the frequency of one every ten lots, then MPI may require that the next ten lots shall be fully verified by purity testing.
- For pelleted seed lots of flowering plants and ornamental species listed in Appendix 4, MPI reserves the
 right to undertake validation audits to confirm that imported consignments are free from quarantine weed
 seeds. Audits may be conducted on a random basis and it will be conducted at an MPI approved facility
 at the expense of the importer.

Ministry for Primary Industries Page 15 of 168

Part 2: Specific Requirements

(1) This part sets out the specific phytosanitary requirements that must be met in addition to Part 1: *General Requirements*, for the following seeds:

Abies Echinochloa Persea Acer Elaeis Petunia Acrocomia Eriobotrya Phaseolus Actinidia Phoenix **Fagus** Pinus Agropyron Fragaria Agrostis Glebionis Pisum Arabidopsis thaliana Glycine **Populus** Avena Gossypium **Prunus**

Beta Helianthus Pseudotsuga menziesii

Brassica napus Hordeum Psophocarpus

Camellia sinensis Pyrus Humulus Camissonia Juglans Quercus Cannabis sativa Lablab Ribes Capsicum Lavandula Rubus Carpinus Lens Sesamum Carthamus tinctorius Linum usitatissimum Solanum

Carya Lithocarpus densiflorus Solanum lycopersicum
Castanea Livistona Solanum tuberosum

Cicer Lophophora williamsii Sorghum

Citrus Lotus Stenotaphrum

Cocos Macadamia Trigonella foenum-graecum

CoffeaMalusTriticumCoriandrumMangiferaUlmusCorylusMedicagoVacciniumCoryphaMyrtaceaeVicia

CucurbitaceaeNicotiana tabacumVignaCuminumOxyriaVitisDesmodiumPanicumZea

Papaver somniferum

Ministry for Primary Industries Page 16 of 168

2.1 Abies

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed for Sowing as "see 155.02.05 under *Abies*"

Approved countries: All

Quarantine pests: Verticillium albo-atrum [strain]

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.1.1 Approved treatment

(1) All Abies seeds must be treated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

2.1.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 17 of 168

2.2 Acer

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed for Sowing as "see 155.02.05 under *Acer*"

Approved countries: All

Quarantine pests: None

Import permit: Required

PEQ: Not required

Phytosanitary certificate: Required

2.2.1 Approved treatment

(1) All Acer seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.2.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 18 of 168

2.3 Acrocomia

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed for Sowing as "see 155.02.05 under *Acrocomia*"

Approved countries: All countries except Guam, the Philippines and the Solomon Islands

Quarantine pests: Coconut cadang-cadang viroid

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.3.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Acrocomia* seeds have been produced in an approved country and have not been produced in Guam, the Philippines or the Solomon Islands".

AND

b) "The Acrocomia seeds have been produced in a 'pest free area' free from Coconut cadang-cadang viroid".

Ministry for Primary Industries Page 19 of 168

2.4 Actinidia

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed for Sowing as "see 155.02.05 under *Actinidia*."

Approved countries: All

Quarantine pests: Apple stem grooving virus [Actinidia infecting strain]

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: six months

Approved treatment: Not required

Phytosanitary certificate: Required

2.4.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.4.2 Testing requirements

Organism	MPI acceptable detection methods
Apple stem grooving virus [Actinida infecting strain]*	ELISA (Bioreba or Loewa) or PCR (Clover et al., 2003) and herbaceous indicators Cq, Nb, Ng, No and Pv.

- (1) Indicator hosts: Chenopodium quinoa (Cq), and Nicotiana benthamiana (Nb), N. occidentalis cv. 37B (No), N. glutinosa (Ng) and Phaseolus vulgaris cv. Prince (Pv). At least two plants of each indicator species must be used in mechanical inoculation tests.
- (2) Indicator plants must be grown under appropriate temperatures and must be shaded for 12-24 hrs prior to inoculation. Maintain post-inoculated indicator species under appropriate glasshouse conditions for at least 4 weeks. Inspect inoculated indicator plants at least twice per week for symptoms of virus infection.
- (3) Testing must be carried out on Actinidia plants while they are in active growth. For bioassay and ELISA, plants shall be sampled from at least two positions on every stem including a young, fully expanded leaflet at the top of the stem and an older leaflet from a midway position.
- (4) PCR and ELISA need to be validated using positive controls/reference material prior to use in quarantine testing.
- (5) Positive and negative controls must be used in ELISA tests.
- (6) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).
- (7) Positive and negative controls (including a blank water control) must be used in PCR.
- (8) Actinidia plants in a PEQ facility must be inspected for signs of pest and disease at least twice per week during periods of active growth and once per week during dormancy.

Ministry for Primary Industries Page 20 of 168

Guidance

- Positive internal controls and a negative plant control should be used to provide confidence in testing results.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at PlantImports@mpi.govt.nz well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit

Reference

Clover, G R G; Pearson, M N; Elliott, D R; Tang, Z; Smales, T E; Alexander, B J R (2003)
 Characterization of a strain of Apple stem grooving virus in *Actinidia chinensis* from China. Plant Pathology 52: 371-378.

Pest list for Actinidia REGULATED PESTS (actionable)

Virus

Capillovirus Apple stem grooving virus [Actinidia infecting strain]

Ministry for Primary Industries Page 21 of 168

2.5 Agropyron

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed for Sowing as "see 155.02.05 under *Agropyron*."

Approved countries: All

Quarantine pests: Tilletia controversa, other Ustilaginales, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.5.1 Approved Treatments

(1) All Agropyron seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.5.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Agropyron seeds have been:
 - i) produced in a 'pest free area' free from Tilletia controversa";

OR

ii) "produced in a crop that has been inspected during the growing season according to appropriate procedures and no *Tilletia controversa* was detected":

OR

iii) "had an representative sample of 600 seeds officially drawn in which no spores of *Tilletia* controversa were found".

2.5.3 Testing requirements

- Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 22 of 168

2.6 Agrostis

The following requirements only apply to species in the Plant Biosecurity listed under Import Specifications for Seed for Sowing as "see 155.02.05 under *Agrostis*."

Approved countries: All

Quarantine pests: Trogoderma spp., Ustilaginales

Import permit: Not Required

PEQ: Not required

Phytosanitary certificate: Required

2.6.1 Approved treatments

(1) All Agrostis seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.6.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 23 of 168

2.7 Arabidopsis thaliana

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Arabidopsis thaliana*."

Approved countries: All

Quarantine pests: None

Import permit: See below

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Not required

2.7.1 GM seed

- (1) A permit to import is required.
- (2) All GM seed must also be imported in accordance with a HSNO approval.

2.7.2 Non-GM seed

- (1) A declaration signed by the exporter and importer must accompany the consignment declaring that the consignment does not contain GM seeds.
- (2) The declaration form is provided in Appendix 3.

Ministry for Primary Industries Page 24 of 168

2.8 Avena

The following requirements only apply to species in the Plant Biosecurity Index listed under import specifications for Seed as "see 155.02.05 under *Avena*".

Approved countries: Australia, Austria, Belgium, Canada, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, The Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom and United States of America.

Quarantine pests: Refer to pest list for Avena

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.8.1 Approved treatments

(1) In lieu of pest free area for *Cephalosporium gramineum* the *Avena* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.8.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Avena seeds have been:
 - i) produced in a 'pest free area' free from Xanthomonas campestris pv. undulosa and High plains virus;

OR

ii) produced in a 'pest free place of production' free from *Xanthomonas campestris* pv. *undulosa* and *High plains virus*".

AND

- b) "The Avena seeds have been:
 - i) produced in a 'pest free area' free from Anguina tritici";

OR

ii) "produced in a 'pest free place of production' free from Anguina tritici';

OR

iii) "inspected microscopically in accordance with official procedures and *Anguina tritici* was not detected".

AND

- c) "The Avena seeds have been:
 - i) produced in a 'pest free area' free from Cephalosporium gramineum";

OR

ii) "treated with a fungicide combination in MPI approved treatments".

Ministry for Primary Industries Page 25 of 168

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments



Ministry for Primary Industries Page 26 of 168

Pest list for Avena REGULATED PESTS (actionable)

secta	
Blattodea	
Blattidae	
Blatta orientalis	oriental cockroach
Coleoptera	
Bostrichidae	
Prostephanus truncatus	larger grain borer
Cryptophagidae	
Cryptophagus schmidti	
Cucujidae	
Cathartus quadricollis	squarenecked grain beetle
Curculionidae	
Caulophilus oryzae	broadnosed grain weevil
Dermestidae	
Trogoderma granarium	khapra beetle
Trogoderma inclusum	trogoderma beetle
Trogoderma ornatum	trogoderma beetle
Trogoderma simplex	dermestid beetle
Trogoderma sternale	dermestid beetle
Trogoderma variabile	warehouse beetle
Mycetophagidae	
Mycetophagus quadriguttatus	spotted hairy fungus beetle
Nitidulidae	
Carpophilus obsoletus	dried fruit beetle
Ptinidae	
Gibbium psylloides	shiny spider beetle
Mezium americanum	american spider beetle
Niptus hololeucus	golden spider beetle
Pseudoeurostus hilleri	spider beetle
Ptinus clavipes	brown spider beetle
Ptinus fur	whitemarked spider beetle
Ptinus villiger	hairy spider beetle
Tipnus unicolor	spider beetle
Trigonogenius globulus	
Tenebrionidae	
Alphitobius laevigatus	black fungus beetle
Alphitophagus bifasciatus	two-banded fungus beetle
Blaps mucronata	cellar beetle
Gnatocerus maxillosus	slenderhorned flour beetle
Latheticus oryzae	longheaded flour beetle
Palorus ratzeburgi	smalleyed flour beetle

Ministry for Primary Industries Page 27 of 168

Palorus subdepressus	depressed flour beetle
Tribolium audax	american black flour beetle
Tribolium destructor	dark flour beetle
Trogossitidae	
Lophocateres pusillus	siamese grain beetle
Hemiptera	
Lygaeidae	
Elasmolomus sordidus	seed bugs
Lepidoptera	
Cosmopterigidae	
Pyroderces rileyi	pink scavenger caterpillar
Oecophoridae	
Anchonoma xeraula	grain moth
Pyralidae	
Corcyra cephalonica	rice moth
Ephestia figulilella	raisin moth
Paralipsa gularis	stored nut moth
Tineidae	
Nemapogon variatella	corn moth
Mite	
Arachnida	
Acarina	
Eriophyidae	
Aceria tosichella	wheat curl mite
Aceria tulipae [vector]	wheat curl mite
Siteroptidae	
Siteroptes cerealium	asparagus spider mite
Tarsonemidae	
Steneotarsonemus spirifex	oat spiral mite
Nematode	
Secernentea	
Tylenchida	
Anguinidae	
Anguina tritici [vector]	seed gall nematode
Fungus	
Hyphomycetales	
Moniliaceae	
Cephalosporium gramineum	
Bacterium	
Pseudomonadaceae	
Xanthomonas campestris pv. undulosa	leaf streak
Virus	
High plains virus	

Ministry for Primary Industries Page 28 of 168

2.9 Beta

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Beta*."

Approved countries: All

Quarantine pests: Clavibacter michiganensis subsp. sepedonicus.

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.9.1 Phytosanitary certificate - Additional Declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Beta seeds have been:
 - i) produced in a 'pest free area' free from Clavibacter michiganensis subsp. sepedonicus;

OR

ii) Clavibacter michiganensis subsp. sepedonicus was not detected in a representative sample of 3200 seeds drawn from this consignment.

2.9.2 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.

Ministry for Primary Industries Page 29 of 168

2.10 Brassica napus

These requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Brassica napus".

Approved countries: All

Quarantine pests: None

Import Permit: Permit not required, unless seeds are to be grown in PEQ.

PEQ: Not required, unless imported under options 2.2.2 or 2.2.3 of the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material.

Approved treatment: Not required

Phytosanitary certificate: Required

2.10.1 Phytosanitary certificate

If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must (1) confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.10.2 GM seed testing

- (1) In addition to the phytosanitary requirements above, all consignments of Brassica napus var. oleifera (oilseed rape) are required to be representatively sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5.3 Genetically modified testing certificate in this IHS).
- (2) The full scientific name of the *Brassica napus* sub-species or variety, plus the appropriate common name, must be specified on the phytosanitary certificate, e.g. Brassica napus var. biennis (forage rape) or Brassica napus var. oleifera (oilseed rape).
- Importers of consignments of Brassica napus that are not identified appropriately will be offered the (3) options of re-shipment, destruction or testing for the presence of unapproved GM seeds.

Guidance

- Validation of Brassica napus varieties MPI reserves the right to undertake validation audits to confirm that the variety matches that which is stated on the phytosanitary certificate. Audits may be conducted on a random basis and if required, grow out testing of samples will be conducted at an MPI accredited facility at the expense of the importer.
- The MPI Protocol for Testing for the Presence of Genetically Modified Plant Material can be found at https://www.mpi.govt.nz/document-vault/10250
- More information on genetically modified seeds can also be found at https://www.mpi.govt.nz/importing/plants/seeds-for-sowing/genetically-modified-seeds/

Ministry for Primary Industries Page 30 of 168

2.11 Camellia sinensis

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Camellia sinensis*".

Approved countries: All

Quarantine pests: Exobasidium vexans, Phloem necrosis

Import permit: Required

PEQ: Level 1

Minimum PEQ period: 1 growing season

Isolation: 50m exclusion area

Phytosanitary certificate: Required

2.11.1 Approved treatments

(1) All Camellia sinensis seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.11.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Camellia sinensis seeds have been produced in a 'pest free area' free from Exobasidium vexans and Phloem necrosis".

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 31 of 168

2.12 Camissonia

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Camissonia*".

Approved countries: All

Quarantine pests: Peronospora arthurii

Import permit: Not Required

PEQ: Not required

Phytosanitary certificate: Required

2.12.1 Approved treatments

(1) All Camissonia seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.12.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 32 of 168

2.13 Cannabis sativa

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Cannabis sativa*".

Approved countries: All

Quarantine pests: Refer to pest list for Cannabis sativa

Import permit: Not Required

PEQ: Not required

Phytosanitary certificate: Required

Guidance

 Importers of Cannabis sativa must contact Medsafe before importation for advice on licensing: Medicines Control

Medsafe

Email: medicinescontrol@health.govt.nz

Phone: (04) 816 2444

2.13.1 Approved treatments

- (1) In lieu of pest free area or pest free place of production for *Pseudomonas syringae* pv. *cannabina* and *Xanthomonas campestris* pv. *cannabis*, the *Cannabis sativa* seeds must be treated using a hot water dip (for bacteria and parasitic weed) prior to shipment or on arrival in New Zealand;
 - a) hot water treatment must be conducted as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.
- (2) In lieu of pest free area for *Leptosphaeria woroninii*, *Septoria cannabis* and *Curvularia cymbopogonis*, the *Cannabis sativa* seeds must be treated with fungicide as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.13.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Cannabis sativa seeds have been:
 - i) produced in a 'pest free area' free from the named regulated bacteria (*Pseudomonas syringae* pv. *cannabina* and *Xanthomonas campestris* pv. *cannabis*)";

OR

ii) "produced in a 'pest free place of production' free from the named regulated bacteria (Pseudomonas syringae pv. cannabina and Xanthomonas campestris pv. cannabis)";

OR

iii) "treated with hot water treatment in MPI approved treatments";

AND

b) "The Cannabis sativa seeds have been:

Ministry for Primary Industries Page 33 of 168

i) produced in a 'pest free area' free from the named regulated fungi (*Leptosphaeria* woroninii, Septoria cannabis and Curvularia cymbopogonis)";

OR

ii) "treated with an approved fungicide combination in MPI approved treatments".

Guidance

- The hot water treatment that would be carried out in New Zealand as an alternative to the same treatment prior to shipment, cannot be permitted as no MPI- approved facility is currently available in New Zealand.
- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

References:

• Hemp Diseases and Pests: Management and Biological Control. J. M. McPartland, R. C. Clarke and D. P. Watson 2000. CAB International.

Pest list for Cannabis sativa REGULATED PESTS (actionable)

Insect		
	Pyrrhocoris apterus	fire bug
	Episyrphus balteatus	
	Ischiodon scutellaris	syrphid fly
	Metasyrphus latifasciatus	syrphid fly
	Sphaerophoria scripta	hover fly
	Syritta pipiens	hover fly
Mite		
	Aculops cannabicola	hemp russett mite
Fungus		
	Curvularia cymbopogonis	
	Leptosphaeria woroninii	
	Septoria cannabis	yellow leaf spot
Bacterium		
	Pseudomonas syringae pv. cannabina	
	Xanthomonas campestris pv. cannabis	
Weed		
	Orobanche ramosa	branched broomrape

Ministry for Primary Industries Page 34 of 168

2.14 Capsicum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Capsicum*".

Approved countries: All countries

Quarantine pests: Pepper chat fruit viroid; Potato spindle tuber viroid, Tomato brown rugose fruit virus,

Tomato mottle mosaic virus

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.14.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - "The [Capsicum annuum; C. baccatum; C. cardenasii; C. chinense; C. eximium; C. frutescens; C. microcarpum; C. pendulum; C. pubescens] seeds for sowing have been
 - a) For Potato spindle tuber viroid (PSTVd):
 - i) produced in (country name) where Potato spindle tuber viroid is not known to occur."

OR

ii) produced in a 'pest free place of production', where parent plants were tested according to a NPPO approved methodology and found free from *Potato spindle tuber viroid*"

OR

officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology using an approved PCR NPPO testing method, and found to be free from *Potato spindle tuber viroid*"

AND

- b) For Pepper chat fruit viroid (PCFVd):
 - i) produced in a 'pest free area' free from Pepper chat fruit viroid"

OR

ii) Pepper chat fruit viroid (PCFVd) is absent/not known to occur in _____(name of country)

OR

iii) produced in a 'pest free place of production' free from Pepper chat fruit viroid"

OR

iv) officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology using an approved PCR NPPO testing method, and found to be free from *Pepper chat fruit viroid*"

AND

Ministry for Primary Industries Page 35 of 168

- c) For Tomato brown rugose fruit virus (TBRFV):
 - i) produced in 'pest free area', free from *Tomato brown rugose fruit virus*".

OR

ii) produced in a 'pest free place of production' free from *Tomato brown rugose fruit virus*".

OR

iii) officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved PCR testing method and found free from *Tomato brown rugose fruit virus*".

AND

- d) For Tomato mottle mosaic virus (ToMMV):
 - i) produced in a 'pest free area' free from Tomato mottle mosaic virus"

OR

ii) produced in a 'pest free place of production' free from *Tomato mottle mosaic virus*"

OR

iii) officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved ELISA or NPPO-approved PCR testing method, and found free from *Tomato mottle mosaic virus*"

2.14.2 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) Testing on-shore will be performed using an MPI-approved testing method.
- (4) For seed lots of 15,000 or more seeds:
 - a) A representative sample of a minimum of 3000 seeds, officially drawn according to ISTA or AOSA methodology is required from each seed lot and tested as specified in the schedule;
- (5) For seed lots with less than 15,000 seeds:
 - a) A composite sample of a minimum of 3000 seeds must be officially drawn across all seed lots of the same seed species in a consignment, which must have been produced at the same place of production or production site.
- (6) In order to achieve a composite sample, proportionate sampling must be carried out across all lots imported. A sample of seeds must be drawn from each imported lot within a consignment, adding up to 3000 seeds. The size of the sample from each lot must be proportionate to the size of the imported lot within the consignment.
- (7) All importers who test a composite sample on arrival in New Zealand must make a <u>declaration</u> to identify the place of production or production site of the lots that form the composite sample.

Guidance

- The sample size from each lot to form the composite sample should be calculated as follows:
 - a) The proportion of each lot in the total consignment (seed number) is calculated using the following equation:


Proportion of total consignment size = $\frac{No.of\ seeds\ in\ each\ lot}{Total\ number\ of\ seeds\ in\ consignment}$

Ministry for Primary Industries Page 36 of 168

b) Calculate the sample size for each lot (number of seeds) using a total composite sample size of 3000 seeds:

Sample size of each line = 3000 seeds x proportion of total consignment size

- c) Take the sum of the sample size for each lot to check that the total composite sample for the consignment is at least 3000 seeds.
- The ISHI-Veg local lesion bioassay for *Tomato brown rugose fruit virus* and *Tomato mottle mosaic virus* is not accepted as a valid test by MPI.
- Additional declarations on phytosanitary certificates to meet the offshore testing requirements for Tomato brown rugose fruit virus in Import Health Standard 155.02.05: Seeds for sowing should be based only on a negative result obtained in an NPPO-approved PCR test and not on results from a bioassay.
- Additional declarations on phytosanitary certificates to meet the offshore testing requirements for Tomato mottle mosaic virus in Import Health Standard 155.02.05: Seeds for sowing should be based only on a negative result obtained in an NPPO-approved ELISA or NPPO-approved PCR test and not on results from a bioassay.



2.15 Carpinus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Carpinus*".

Approved countries: All

Quarantine pests: Cladosporium caryigenum

Import permit: Required

PEQ: Not required

Phytosanitary certificate: Required

2.15.1 Approved treatments

(1) All Carpinus seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.15.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Carpinus seeds have been produced in an area where Cladosporium caryigenum is not known to occur".

Guidance

- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit

Ministry for Primary Industries Page 38 of 168

2.16 Carthamus tinctorius

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Carthamus tinctorius*".

Approved countries: All

Quarantine pests: Alternaria carthami, Cercospora carthami, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.16.1 Approved treatment

(1) All Carthamus tinctorius seeds for sowing must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.16.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 39 of 168

2.17 Carya

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Carya*".

Approved countries: Australia, USA

Quarantine pests: Cladosporium caryigenum, Conotrachelus spp., Curculio caryae, Cydia caryana, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.17.1 Approved Treatments

- (1) In lieu of pest free area for *Conotrachelus* spp., *Curculio caryae* or *Cydia caryana* all *Carya* seeds must be fumigated (in) as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.
- (2) The Carya seeds must be treated with fungicide as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.17.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Carya seed or nuts have been:
 - i) produced in an area where they are not known to be attacked by *Conotrachelus* spp., *Curculio caryae* or *Cydia caryana*;

OR

- ii) fumigated with methyl bromide at _____ pressure for ____ hours at ____ g/m³ at a temperature of ____ C";
- the pressure/time/rate temperature combination used is to be in accordance with the scale in MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

AND

b) "The Carya seeds or nuts have been produced in an area where Cladosporium caryigenum is not known to occur".

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 40 of 168

2.18 Castanea

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Castanea*".

Approved countries: All

Quarantine pests: Ceratocystis fagacearum; Cryphonectria parasitica; Curculio spp.; Cyrtepistomus

castaneus

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 2 years

Approved treatment: Not required

Phytosanitary certificate: Required

2.18.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Castanea seeds have been:
 - i) produced in trees that have been officially inspected and found to be free of diseases caused by *Cryphonectria* spp;

OR

ii) produced in an area where Cryphonectria parasitica is known not to occur".

2.18.2 Inspection and testing requirements

Organism	MPI acceptable detection methods	
Ceratocystis fagacearum	Growing season inspection in PEQ for disease symptom expression	
Cryphonectria parasitica	Growing season inspection in PEQ for disease symptom expression	

Guidance

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at PlantImports@mpi.govt.nz well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 41 of 168

2.19 Cicer

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Cicer*."

Approved countries: All

Quarantine pests: Ascochyta rabiei, Megaselia arietina, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.19.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Cicer seeds have been:
 - i) produced in a 'pest free area' free from Ascochyta rabiei;

OR

ii) produced in a 'pest free place of production' free from Ascochyta rabiei".

Ministry for Primary Industries Page 42 of 168

2.20 Citrus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Citrus*."

Approved countries: Australia, Austria, Belgium, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, United States of America.

Quarantine pests: Xanthomonas campestris pv. citri, 'Candidatus Liberibacter africanus', 'Candidatus Liberibacter asiaticus', 'Candidatus Liberibacter americanus'.

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.20.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Citrus seeds have been produced in an area where Xanthomonas campestris pv. citri is not known to occur":

AND

b) "The Citrus seeds have been produced in an area where 'Candidatus' Liberibacter spp. is not known to occur".

Ministry for Primary Industries Page 43 of 168

2.21 Cocos

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Cocos."

Approved countries: All countries except Guam, the Philippines and the Solomon Islands

Quarantine pests: Coconut cadang-cadang viroid

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.21.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Cocos seeds have been produced in an approved country and have not been produced in Guam, the Philippines or the Solomon Islands"

AND

b) "The Cocos seeds have been produced in a 'pest free area' free from Coconut cadang-cadang viroid"

Ministry for Primary Industries Page 44 of 168

2.22 Coffea

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Coffea*".

Approved countries: Australia, Cook Islands, Hawaii, Samoa, Tonga

Quarantine pests: Stephanoderes hampei

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.22.1 Approved treatments

(1) All Coffea seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.22.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 45 of 168

2.23 Coriandrum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Coriandrum*."

Approved countries: All

Quarantine pests: Ramularia coriandri, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.23.1 Approved treatments

(1) All Coriandrum seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.23.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Coriandrum seeds have been:
 - i) produced in a 'pest free area', free from Ramularia coriandri;

OR

ii) produced in a 'pest free place of production' free from Ramularia coriandri."

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 46 of 168

2.24 Corylus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Corylus."

Approved countries: All

Quarantine pests: Cydia latiferreana, Curculio nucum

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.24.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.24.2 Phytosanitary requirements

(1) All *Corylus* seeds imported into New Zealand must have their shells removed to permit inspection, prior to entry.

Ministry for Primary Industries Page 47 of 168

2.25 Corypha

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Corypha*."

Approved countries: All countries except Guam, the Philippines and the Solomon Islands

Quarantine pests: Coconut cadang-cadang viroid.

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.25.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Corypha* seeds have been produced in an approved country and have not been produced in Guam, the Philippines or the Solomon Islands".

AND

b) "The Corypha seeds have been produced in a'pest free area' free from Coconut cadang-cadang viroid".

Ministry for Primary Industries Page 48 of 168

2.26 Cucumis

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Cucumis*".

Approved countries: All

Quarantine pests: Cucumber green mottle mosaic virus (CGMMV); Kyuri green mottle mosaic virus (KGMMV); Melon necrotic spot virus (MNSV)

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.26.1 Phytosanitary certificate – Additional declarations

- (1) The required additional declarations must be endorsed in full on the phytosanitary certificate. No variations in the wording will be accepted by MPI, with the exception of translation artefacts.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:

The [Citrullus lanatus; Cucumis melo] seeds for sowing in this consignment have been:

- a) For Cucumber green mottle mosaic virus (CGMMV):
 - i) sourced from a "pest free area", free from the named regulated virus Cucumber green mottle mosaic virus;

OR

ii) Cucumber green mottle mosaic virus (CGMMV) is absent/not known to occur in _____(name of country)

OR

iii) sourced from mother plants that were sampled according to an NPPO-approved methodology and tested using an NPPO-approved ELISA or an NPPO-approved PCR method, during the active growing period and found free from *Cucumber green mottle mosaic virus*.

OR

iv) sourced from a seed lot officially sampled according to ISTA or AOSA methodology, and tested using the ISTA-validated ELISA or an NPPO-approved PCR method and found free from *Cucumber green mottle mosaic virus*.

AND

b)) For <i>F</i>	Kyurı green	mottle mosaic	virus	(KGMMV):
----	----------------	-------------	---------------	-------	--------	----

i) sourced from a "pest free area", free from Kyuri green mottle mosaic virus;

OR

ii) Kyuri green mottle mosaic virus (KGMMV) is absent/not known to occur in _____(name of country)

OR

Ministry for Primary Industries Page 49 of 168

iii) sourced from a "pest free place of production", free from Kyuri green mottle mosaic virus;

OR

iv) sourced from a seed lot officially sampled according to ISTA or AOSA methodology, and tested using an NPPO-approved serological (ELISA) or molecular (PCR) method and found free from *Kyuri green mottle mosaic virus*.

AND

- c) For Melon necrotic spot virus (MNSV):
 - i) sourced from a "pest free area", free from Melon necrotic spot virus (MNSV);

OR

ii) Melon necrotic spot virus (MNSV) is absent/not known to occur in _____(name of country)

OR

iii) sourced from mother plants that were sampled according to an NPPO-approved methodology and tested using an NPPO-approved ELISA or an NPPO-approved PCR method, during the active growing period and found free from *Melon necrotic spot virus*

OR

iv) sourced from a seed lot officially sampled according to ISTA or AOSA methodology, and tested using an NPPO- or ISTA-approved ELISA or an NPPO-approved PCR method and found free from *Melon necrotic spot virus*.

2.26.2 Testing Requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) For seed lots of 10,000 or more seeds:
 - a) A representative sample of a minimum of 2000 seeds, officially drawn according to ISTA or AOSA methodology is required from each seed lot and tested as specified in the schedule;
- (4) For seed lots with less than 10,000 seeds:
 - a) A composite sample of a minimum of 2000 seeds must be officially drawn across all seed lots of the same seed species in a consignment, which must have been produced at the same place of production or production site.
- (5) In order to achieve a composite sample, proportionate sampling must be carried out across all lots imported. A sample of seeds must be drawn from each imported lot within a consignment, adding up to 2000 seeds. The size of the sample from each lot must be proportionate to the size of the imported lot within the consignment.
- (6) All importers who test a composite sample on arrival in New Zealand must make a <u>declaration</u> to identify the place of production or production site of the lots that form the composite sample.

Guidance

The sample size from each lot should be calculated as follows:

a) The proportion of each lot in the total consignment (seed number) is calculated using the following equation:

Proportion of total consignment size = $\frac{No.of\ seeds\ in\ each\ lot}{Total\ number\ of\ seeds\ in\ consignment}$

Ministry for Primary Industries Page 50 of 168

b) Calculate the sample size for each lot (number of seeds) using a total composite sample size of 2000 seeds:

Sample size of each line= 2000 seeds x proportion of total consignment size

c) Take the sum of the sample size for each lot to check that the total composite sample for the consignment is at least 2000 seeds.



Ministry for Primary Industries Page 51 of 168

2.27 Cucurbitaceae

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Cucurbitaceae*".

Approved countries: All

Quarantine pests: Cucumber green mottle mosaic virus (CGMMV); Kyuri green mottle mosaic virus (KGMMV)

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.27.1 Phytosanitary certificate – Additional declarations

- (1) The required additional declarations must be endorsed in full on the phytosanitary certificate, no variations in the wording will be accepted by MPI, with exception of translation artifacts.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:

The [Benincasa hispida; Cucumis anguria; Cucumis metulliferus; Cucumis myriocarpus; Cucurbita ficifolia; Cucurbita maxima; Cucurbita mixta Cucurbita moschata; Cucurbita pepo; Cucumis sativus; Lagenaria siceraria; Luffa acutangula; Luffa cylindrical; Luffa aegyptiaca; Momordica charantia; Portulaca oleraceae] seeds for sowing in this consignment have been:

- a) For Cucumber green mottle mosaic virus (CGMMV):
 - i) produced in a Pest free area, free from the named regulated virus Cucumber green mottle mosaic virus:

OR

ii) Cucumber green mottle mosaic virus (CGMMV) is absent/not known to occur in _____(name of country)

OR

iii) produced in mother plants that were sampled according to a NPPO approved methodology and tested using a NPPO approved ELISA or a NPPO approved PCR method, during the active growing period and found free from *Cucumber green mottle mosaic virus*.

OR

iv) produced in a seed lot officially sampled according to ISTA or AOSA methodology, and tested using the ISTA validated ELISA or a NPPO approved PCR method and found free from Cucumber green mottle mosaic virus.

AND

- b) For Kyuri green mottle mosaic virus (KGMMV):
 - i) produced in a "Pest free area", free from Kyuri green mottle mosaic virus;

OR

Ministry for Primary Industries Page 52 of 168

ii) Kyuri green mottle mosaic virus (KGMMV) is absent/not known to occur in _____(name of country)

OR

iii) produced in a "Pest Free Place of Production", free from Kyuri green mottle mosaic virus;

OR

iv) produced in a seed lot officially sampled according to ISTA or AOSA methodology, and tested using a NPPO approved serological (ELISA) or molecular (PCR) method and found free from *Kyuri green mottle mosaic virus*.

2.27.2 Testing Requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) For seed lots of 10,000 or more seeds:
 - A representative sample of a minimum of 2000 seeds, officially drawn according to ISTA or AOSA methodology is required from each seed lot and tested as specified in the schedule;
- (4) For seed lots with less than 10,000 seeds:
 - a) A composite sample of a minimum of 2000 seeds must be officially drawn across all seed lots of the same seed species in a consignment, which must have been produced at the same place of production or production site.
- (5) In order to achieve a composite sample, proportionate sampling must be carried out across all lots imported. A sample of seeds must be drawn from each imported lot within a consignment, adding up to 2000 seeds. The size of the sample from each lot must be proportionate to the size of the imported lot within the consignment.
- (6) All importers who test a composite sample on arrival in New Zealand must make a <u>declaration</u> to identify the place of production or production site of the lots that form the composite sample.

Guidance

The sample size from each lot should be calculated as follows:

a) The proportion of each lot in the total consignment (seed number) is calculated using the following equation:

 $Proportion \ of \ total \ consignment \ size = \frac{\textit{No.of seeds in eac } \ \textit{lot}}{\textit{Total number of seeds in consignment}}$

 Calculate the sample size for each lot (number of seeds) using a total composite sample size of 2000 seeds:

Sample size of each line= 2000 seeds x proportion of total consignment size

c) Take the sum of the sample size for each lot to check the total composite sample for the consignment is at least 2000 seeds.

2.27.3 Cucurbita pepo

- (1) Different varieties of Yellow Straightneck, Yellow Crookneck squash and Green Zucchini seeds have been genetically modified. <u>The following varieties are prohibited entry to New Zealand without HSNO</u> <u>approval by EPA:</u>
 - a) Cucurbita pepo event ZW20;
 - b) Cucurbita pepo event CZW3;

Ministry for Primary Industries Page 53 of 168

- c) Yellow Crookneck squash variety "Revenue"; "Tigress"; "Destiny III"; Prelude II;
- d) Yellow Straightneck squash variety "XPT1832 III"; "Conqueror III"; "Patriot II"; "Liberator III";
- e) Green Zucchini variety "SV6009YG"; "Judgement III"; "Justice III"; "Declaration II"; "Independence II".
- (2) Cucurbita pepo importers are required to comply with one of the two options listed below:

Option 1:

 a) a declaration signed by the exporter and importer must accompany the consignment declaring that the consignment does not contain GM seeds (the declaration form template is provided in Appendix 3).

OR

Option 2:

b) a representative sample from each seed lot of *Cucurbita pepo* must be sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5.3 *Genetically modified testing certificate* in this IHS). More information can also be found at https://www.mpi.govt.nz/importing/plants/seeds-for-sowing/genetically-modified-seeds/

Guidance

- The MPI Protocol for Testing for the Presence of Genetically Modified Plant Material can be found at https://www.mpi.govt.nz/document-vault/10250
- The declaration form template is provided in Appendix 3.

References:

- Ling et al., 2014. First report of *Cucumber green mottle mosaic virus* infecting greenhouse cucumber in Canada. Plant Disease 98 (5): 701-2.
- Reingold et al., 2013. First report of Cucumber green mottle mosaic virus (CGMMV) symptoms in watermelon used for the discrimination of non-marketable fruits in Israeli commercial fields. New Disease Reports 28, 11.
- ISTA https://www.seedtest.org/upload/cms/user/SH-07-026-2014.pdf
- Daryono, B. S., Somowiyarjo, S., Natsuaki, K. T. 2005. Biological and Molecular Characterization of Melon-Infecting Kyuri Green Mottle Mosaic virus in Indonesia. Journal of Phytopathology 153, 588-595.
- Daryono, B.S., Somowiyarjo, S. and Natsuaki, K.T. 2006. Biological characterization and complete nucleotide sequence of coat protein of *Kyuri green mottle mosaic virus* isolated from angled loofah in Indonesia. Jour. Agri. Sci. Tokyo Univ. of Agric. 51 (1), 42-52. (Printed in English)
- Daryono, B. S. and Natsuaki, K.T. 2012. Application of Multiplex RT-PCR for Detection of Cucurbitinfecting Tobamovirus. Jordan Journal of Agricultural Sciences, 8 (1): 46-56.
- Hongyun, C., Wendjun, Z., Qinsheng, G. and Shuifang, Z. 2008. Real time TaqMan RT-PCR assay for the detection of *Cucumber green mottle mosaic virus*. Journal of Virological Methods, 149 (2): 326-9.
- Kwon, J. Y., Hong, J. S., Kim, M. J., Choi, S. H., Byeong, E. M., Song, E. G., Kim, H. H., Ryu, K. H. 2014. Simultaneous multiplex PCR detection of seven cucurbit infecting viruses. Journal of Virological Methods 206, 133-139

Ministry for Primary Industries Page 54 of 168

Pest list for Cucurbitaceae REGULATED PESTS (actionable)

•	•		
١	/ı	rı	10
١	/ I	ı	JO

Cucumber green mottle mosaic virus	CGMMV	
Kyuri green mottle mosaic virus	KGMMV	



Ministry for Primary Industries Page 55 of 168

2.28 Cuminum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Cuminum*."

Approved countries: All

Quarantine pests: Alternaria burnsii

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.28.1 Approved treatments

(1) All *Cuminum* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.28.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Cuminum seeds have been:
 - i) produced in a 'pest free area', free from Alternaria burnsii;

OR

ii) produced in a 'pest free place of production', free from Alternaria burnsii".

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 56 of 168

2.29 Desmodium

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Desmodium*."

Approved countries: All

Quarantine pests: Desmodium mosaic virus, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.29.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Desmodium seeds have been:
 - i) produced in an area where Desmodium mosaic virus is not known to occur;

OR

ii) produced in a crop that has been inspected during the growing season according to appropriate procedures and no *Desmodium mosaic virus* was detected".

Ministry for Primary Industries Page 57 of 168

2.30 Echinochloa

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Echinochloa*."

Approved countries: All

Quarantine pests: Sclerospora graminicola, Trogoderma spp., Ustilaginales

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.30.1 Approved treatments

(1) All *Echinochloa* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.30.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Echinochloa seeds have been:
 - i) produced in a 'pest free area', free from Sclerospora graminicola;

OR

ii) produced in a 'pest free place of production', free from Sclerospora graminicola".

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 58 of 168

2.31 Elaeis

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Elaeis*."

Approved countries: All countries except Guam, the Philippines and the Solomon Islands

Quarantine pests: Coconut cadang-cadang viroid

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.31.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Elaeis* seeds have been produced in an approved country and have not been produced in Guam, the Philippines or the Solomon Islands".

AND

b) "The *Elaeis* seeds have been produced in a 'pest free area' free from *Coconut cadang-cadang viroid*".

Ministry for Primary Industries Page 59 of 168

2.32 Eriobotrya

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Eriobotrya*."

Approved countries: All

Quarantine pests: Pseudomonas syringae pv. eriobotryae

Import permit: Required

Approved treatments: Not required

Phytosanitary certificate: Required

(1) Importers must comply with one of the two options listed below:

2.32.1 Option 1:

PEQ: Not required

Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "the *Eriobotrya* seeds have been produced in an area where *Pseudomonas syringae* pv. eriobotryae is not known to occur".

2.32.2 Option 2:

PEQ: Level 3B

Minimum PEQ Period: 2 growing seasons

Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at PlantImports@mpi.govt.nz well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 60 of 168

2.33 Fagus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Fagus."

Approved countries: All

Quarantine pests: Tortricidae

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.33.1 Approved treatments

(1) All *Fagus* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.33.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 61 of 168

2.34 Fragaria

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Fragaria*."

Approved countries: All

Quarantine pests: Refer to "Pest list for Fragaria.

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 6 months

Approved treatments: Not required

Phytosanitary certificate: Required

2.34.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.34.2 Inspection and testing requirements

Organism	MPI acceptable detection methods
Fragaria chiloensis latent virus	Herbaceous indexing with Cq
Raspberry ringspot virus*	ELISA or PCR and herbaceous indexing with Cq
Strawberry latent ringspot virus*	ELISA or PCR and herbaceous indexing with Cq
Tobacco streak virus*	ELISA or PCR and herbaceous indexing with Cq
Tomato ringspot virus*	ELISA or PCR and herbaceous indexing with Cq

Cq – Chenopodium quinoa


- (1) Tests are to be carried out on plants germinated from the imported seeds.
- (2) Testing must be carried out on plants while they are in active growth.
- (3) Indicator plants must be grown under appropriate temperatures.
- (4) Indicator plants must be shaded for 12-24 hrs prior to inoculation.
- (5) Post-inoculated indicator plants must be maintained under appropriate glasshouse conditions for at least 4 weeks.
- (6) Post-inoculated indicator plants must be inspected at least twice per week for signs of virus infection with observations being recorded on a weekly basis.
- (7) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).
- (8) PCR and ELISA need to be validated using positive controls/reference material prior to use in quarantine testing.
- (9) Positive, negative, and buffer controls must be used in ELISA tests.
- (10) Positive controls must be used in PCR.

Ministry for Primary Industries Page 62 of 168

(11) Fragaria plants in a PEQ facility must be inspected for signs of pest and disease at least once per week

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.



Pest list for Fragaria REGULATED PESTS (actionable)

Virus	
	Fragaria chiloensis latent virus
	Raspberry ringspot virus
	Strawberry latent ringspot virus (strains not in New Zealand)
'	Tobacco streak virus
	Tomato ringspot virus (strains not in New Zealand)



Ministry for Primary Industries Page 64 of 168

2.35 Glebionis

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Glebionis*."

Approved countries: All countries

Quarantine pests: Potato spindle tuber viroid

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.35.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The [insert species name] seeds have been:
 - i) produced in a 'pest free area' free from Potato spindle tuber viroid;

OR

ii) produced in a 'pest free place of production' where parent plants have been tested according to an NPPO approved methodology and found free from *Potato spindle tuber viroid*:

OR

iii) officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology using an NPPO approved PCR testing method, and found to be free from *Potato spindle tuber viroid*".

2.35.2 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) Testing on-shore will be performed using an MPI-approved testing method.

Ministry for Primary Industries Page 65 of 168

2.36 Glycine

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Glycine*."

Approved countries: All

Quarantine pests: Peronospora manshurica, Trogoderma spp.

Import Permit: Permit not required, unless seeds are to be grown in PEQ.

PEQ: Not required, unless imported under options 2.2.2 or 2.2.3 of the MPI <u>Protocol</u> for Testing for the Presence of Genetically Modified Plant Material.

Phytosanitary certificate: Required

2.36.1 Approved treatments

(1) All *Glycine* seeds must be treated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

2.36.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Glycine seeds have been:
 - i) inspected in accordance with appropriate official procedures and found to be free of any visually detectable regulated pests, including *Trogoderma* spp".

AND

- b) "The Glycine seeds have been:
 - i) produced in a 'pest free area' free from *Peronospora manshurica*;

OR

ii) produced in a 'pest free place of production' free from Peronospora manshurica".

AND

c) "The *Glycine* seeds have been treated against *Peronospora manshurica* using one of the approved fungicide combinations".

2.36.3 GM seed testing

(1) In addition to the phytosanitary requirements above, all consignments of *Glycine max* (soybean) are required to be representatively sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5.3: *Genetically Modified Testing Certificate*).

Guidance

• The MPI Protocol for testing for the presence of genetically modified plant material can be found at https://www.mpi.govt.nz/document-vault/10250

Ministry for Primary Industries Page 66 of 168

- More information on genetically modified seeds can also be found at https://www.mpi.govt.nz/importing/plants/seeds-for-sowing/genetically-modified-seeds/
 Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments



Ministry for Primary Industries Page 67 of 168

2.37 Gossypium

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Gossypium*."

Approved countries: Australia

Quarantine pests: Anthonomus grandis, Trogoderma spp.

Import Permit: Permit not required, unless seeds are to be grown in PEQ.

PEQ: Not required, unless imported under options 2.2.2 or 2.2.3 of the MPI <u>Protocol</u> for Testing for the Presence of Genetically Modified Plant Material.

Approved treatment: Not required

Phytosanitary certificate: Required

2.37.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The seed has been cleaned and is completely free of lint".

2.37.2 GM seed testing

- (1) In addition to the phytosanitary requirements above, all consignents of *Gossypium hirsutum* (cotton) are required to be representatively sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5.3: *Genetically Modified Testing Certificate*).
- (2) Importers of consignments of *Gossypium hirsutum* that are not identified appropriately will be offered the options of re-shipment, destruction or testing for the presence of unapproved GM seeds.

Ministry for Primary Industries Page 68 of 168

2.38 Helianthus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Helianthus*."

Approved countries: Australia, Austria, Belgium, Canada, Chile, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, United States of America

Quarantine pests: Alternaria helianthi, Neolasioptera helianthi (syn. Lasioptera murtfeldtiana), Plasmopara halstedii, Septoria helianthi, Sunflower mosaic virus, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.38.1 Approved treatments

(1) All *Helianthus* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.38.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Helianthus seeds have been produced in a crop that has been inspected during the growing season according to appropriate procedures and no Alternaria helianthi, Neolasioptera helianthi, Plasmopara halstedii, Septoria helianthi or Sunflower mosaic virus was detected;

OR

b) "The Helianthus seeds have been produced in an area where Alternaria helianthi, Neolasioptera helianthi, Plasmopara halstedii, Septoria helianthi and Sunflower mosaic virus are not known to occur":

OR

- c) "The Helianthus seeds have:
 - been produced in a crop that has been inspected during the growing season according to appropriate procedures and no Neolasioptera helianthi, Plasmopara halstedii or Sunflower mosaic virus was detected;

AND

ii) had 600 pure seeds drawn and tested in accordance with the general directions for seed health testing in the current International Rules for Seed Testing and no evidence of contamination with Alternaria helianthi or Septoria helianthi was found".

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 69 of 168

Pest list for Helianthus REGULATED PESTS (actionable)

Insect		
Insecta		
Coleoptera		
Dermestidae		
	Trogoderma granarium	khapra beetle
	Trogoderma variabile	warehouse beetle
Diptera		
Asteraceae		
	Neolasioptera helianthi (syn. Lasioptera murtfeldtiana)	midge
Fungus		
Ascomycota		
Pleosporales		
Pleosporaceae	e	
	Alternaria helianthi	
Dothideales		
Leptosphaeria	ceae	
	Leptosphaeria lindquistii	leaf spot
Mitosporic fungi (Coelomycetes)		
Sphaeropsidales		
Sphaerioidaceae		
	Septoria helianthi	septoria leaf spot
Mitosporic fungi (Hyphomycetes)		
Hyphomycetales		
Moniliaceae		
	Aspergillus parasiticus	mould
Oomycota		
Peronosporales		
Peronosporaceae		
	Plasmopara halstedii	downy mildew
Bacterium		
Pseudomonadaceae		
	Pseudomonas syringae pv. aptata	bacterial spot
	Pseudomonas syringae pv. tagetis	bacterial leaf spot
Virus		
Potyviridae		
Potyvirus		
	Sunflower mosaic virus	

Ministry for Primary Industries Page 70 of 168

2.39 Hordeum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Hordeum*."

Approved countries: Australia, Austria, Belgium, Canada, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Japan, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom and United States of America

Quarantine pests: Refer to "Pest list for Hordeum"

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.39.1 Approved treatments

(1) In lieu of pest free area for Cephalosporium gramineum and Fusarium longipes all Hordeum seed for sowing seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.39.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Hordeum seeds have been:
 - i) produced in a 'pest free area' free from the named regulated bacteria (*Pseudomonas* syringae pv. striafaciens, Rathayibacter tritici, Xanthomonas campestris pv. undulosa) and virus (*High plains virus*);

OR

ii) produced in a 'pest free place of production' free from the named regulated bacteria (Pseudomonas syringae pv. striafaciens, Rathayibacter tritici, Xanthomonas campestris pv. undulosa) and virus (High plains virus)";

AND

- b) "The *Hordeum* seeds have been:
 - i) produced in a "pest free area" free from the named regulated fungi (*Cephalosporium gramineum*, *Fusarium longipes*)";

OR

ii) "treated with one of the fungicide combinations in MPI approved treatments";

AND

- c) "The Hordeum seeds have been:
 - i) produced in a 'pest free area' free from *Tilletia controversa*";

OR

Ministry for Primary Industries Page 71 of 168

ii) "produced in a 'pest free place of production' free from *Tilletia controversa*, and treated with one of the fungicide combinations in MPI approved treatments";

OR

"had a representative sample of 600 seeds drawn from this consignment according to the International Seed Testing Association's methodology and have been tested for *Tilletia* controversa, and treated with one of the fungicide combinations in MPI approved treatments."

2.39.3 Testing requirements for *Tilletia controversa*

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 72 of 168

Pest list for Hordeum REGULATED PESTS (actionable)

Insect		
Insecta		
Blattodea		
Blattidae		
	Blatta orientalis	oriental cockroach
Coleoptera		
Curculionidae		
	Caulophilus oryzae	broadnosed grain weevil
Dermestidae		
	Trogoderma granarium	khapra beetle
	Trogoderma grassmani	trogoderma beetle
	Trogoderma inclusum	trogoderma beetle
	Trogoderma irroratum	trogoderma beetle
	Trogoderma ornatum	trogoderma beetle
	Trogoderma simplex	dermestid beetle
	Trogoderma sternale	dermestid beetle
	Trogoderma variabile	warehouse beetle
Languriidae		
	Pharaxonotha kirschii	mexican grain beetle
Tenebrionidae		
	Embaphion muricatum	false wireworm
	Latheticus oryzae	longheaded flour beetle
	Palorus ratzeburgi	smalleyed flour beetle
	Palorus subdepressus	depressed flour beetle
	Tribolium audax	american black flour beetle
	Tribolium destructor	dark flour beetle
Lepidoptera		
Tineidae		
	Haplotinea insectella	casemaking moth
	Tinea fictrix	casemaking moth
Mite		
Arachnida		
Acarina		
Acaridae		
	Acarophenax tribolii [Animals Biosecurity]	grain mite
Eriophyidae		
	Aceria tosichella	wheat curl mite
	Aceria tulipae [vector]	wheat curl mite
Pyemotidae		
i yomodado		

Ministry for Primary Industries Page 73 of 168

Basidiomycot	a: Ustomy	cetes
--------------	-----------	-------

Tilletiaceae		
	Tilletia controversa	dwarf bunt
Mitosporic fungi (Hyphomycete	es)	
Hyphomycetales		
Moniliaceae		
	Cephalosporium gramineum	stripe
Tuberculariales		
Tuberculariaceae		
	Fusarium longipes	fusarium head blight
Bacterium		
Corynebacteriaceae		
	Rathayibacter tritici	yellow ear rot
Pseudomonadaceae		
	Pseudomonas syringae pv. striafaciens	bacterial stripe blight
	Xanthomonas campestris pv. undulosa	leaf streak
Virus		
	High plains virus	
•		

Ministry for Primary Industries Page 74 of 168

2.40 Humulus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Humulus lupulus*."

Approved countries: All

Quarantine pests: Pseudoperonospora humuli, Verticillium albo-atrum

Import permit: Required

PEQ: Level 3B

Minimum PEQ Period: 1 growing season

Approved treatments: Not required

Phytosanitary certificate: Required

2.40.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should
 contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their
 importation plans and apply for an import permit. This is because MPI will need to undertake a risk
 assessment that could potentially lead to changes being made to import requirements and this could
 lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 75 of 168

2.41 Juglans

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Juglans*."

Approved countries: Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, Mexico, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, United States of America

Quarantine pests: Gnomonia leptostyla, Pyralidae; Tortricidae; Trogoderma spp., Cherry leaf roll virus

Import permit: Required

PEQ: Level 1

Minimum PEQ Period: 2 growing seasons

Isolation: 50m exclusion area

Phytosanitary certificate: Required

2.41.1 Approved treatments

(1) All *Juglans* seeds must be fumigated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

2.41.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Juglans seed have been:
 - i) inspected during the growing season according to appropriate procedures, and no Gnomonia leptostyla or Cherry leaf roll virus was detected;

OR

ii) produced in an area where *Gnomonia leptostyla* and *Cherry leaf roll virus* are not known to occur".

AND

b) "The seed was fumigated with methyl bromide at ___ pressure for ___ hours at ___ g/m³ at a temperature of ___ °C ";

Guidance

- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at PlantImports@mpi.govt.nz well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 76 of 168

2.42 Lablab

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Lablab*".

Approved countries: All

Quarantine pests: Earias vitella, Maruca testulali, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

For seed in pods:

2.42.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - "The pods were inspected before export and no caterpillars of *Earias vitella* or *Maruca testulalis* were found in a 600 unit sample".

Ministry for Primary Industries Page 77 of 168

2.43 Lavandula

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under under Lavandula."

Approved countries: All

Quarantine pests: Coniothyrium lavandulae, Phoma lavandulae

Import permit: Not Required

PEQ: Not required

Phytosanitary certificate: Required

2.43.1 Approved treatments

(1) All Lavandula seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.43.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Lavandula seeds have been:
 - i) produced in a 'pest free area', free from *Coniothyrium lavandulae* and *Phoma lavandulae*OR
 - ii) produced in a 'pest free place of production', free from *Coniothyrium lavandulae* or *Phoma lavandulae*".

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 78 of 168

2.44 Lens

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Lens*."

Approved countries: All

Quarantine pests: Trogoderma granarium

Import permit: Not Required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.44.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Lens* seeds have been inspected in accordance with appropriate official procedures and found to be free of *Trogoderma granarium*".

OR

b) "Trogoderma granarium" is absent/not known to occur in _____ (name of country)

Ministry for Primary Industries Page 79 of 168

2.45 Linum usitatissimum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Linum usitatissimum*."

Approved countries: All

Quarantine pests: None

Import permit: Permit not required

PEQ: Not required.

Approved treatments: Not required

Phytosanitary certificate: Required

2.45.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.45.2 GM seed declaration

- (1) There are no specific requirements for *Linum usitatissimum* seeds except for the following GM event which is prohibited entry to New Zealand without HSNO approval by the EPA:
 - a) Linum usitatissimum var. FP967 (CDC Triffid).
- (2) Importers are required to comply with one of the two options listed below:

Option 1:

a) a declaration signed by the exporter and importer must accompany the consignment declaring that the consignment does not contain GM seeds (refer to Appendix 3: Declaration form).

Option 2:

b) samples must be representatively sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5.3 *Genetically modified testing certificate* in this IHS).

Ministry for Primary Industries Page 80 of 168

2.46 Lithocarpus densiflorus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Lithocarpus densiflorus*."

Approved countries: Australia, Canada, Germany, India, Israel, Japan, Mexico, Tunisia, United Kingdom, United States of America

Quarantine pests: Ceratocystis fagacearum, Tortricidae

Import permit: Required

PEQ: Level 1

Isolation: 50 m

Phytosanitary certificate: Required

2.46.1 Approved Treatments

(1) All *Lithocarpus densiflorus* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.46.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Lithocarpus densiflorus seed has been:
 - i) collected from trees that have been officially inspected for disease caused by *Ceratocystis* fagacearum and no disease was detected;

OR

ii) produced in an area where Ceratocystis fagacearum is not known to occur".

Guidance

- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 81 of 168

2.47 Livistona

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Livistona*."

Approved countries: All countries except Guam, the Philippines and the Solomon Islands

Quarantine pests: Coconut cadang-cadang viroid

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.47.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Livistona* seeds have been produced in an approved country and have not been produced in Guam, the Philippines or the Solomon Islands".

AND

b) "The Livistona seeds have been produced in a 'pest free area' free from Coconut cadang-cadang viroid".

Ministry for Primary Industries Page 82 of 168

2.48 Lophophora williamsii

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Lophophora williamsii."

Approved countries: All

Quarantine pests: None

Import permit: Required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.48.1 Phytosanitary certificate

If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

Importers of Lophophora williamsii must contact Medsafe before importation for advice on licensing:
 Medicines Control

Medsafe

Email: medicinescontrol@health.govt.nz

Phone: (04) 816 2444

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at PlantImports@mpi.govt.nz well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 83 of 168

2.49 Lotus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Lotus."

Approved Countries: All

Quarantine pests: Trogoderma spp.

Import permit: Not required


PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.49.1 Phytosanitary certificate - Additional declaration

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.



2.50 Macadamia

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Macadamia*."

Approved countries: All

Quarantine pests: Cryptophlebia ombrodelta, Deudorix epijarbas, Dichocrocis punctiferalis

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.50.1 Approved treatments

(1) All *Macadamia* seeds must be fumigated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

2.50.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

a)	"The Macadamia seed was fumigated with methyl bromide at	pressure for hours at
	g/m³ at a temperature of °C ";	

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 85 of 168

2.51 Malus

These requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Malus*."

Approved countries: All

Quarantine pests: Apple scar skin viroid, Monilinia fructigena, Tomato bushy stunt virus

Import permit: Required

PEQ: Level 2. Herbaceous indexing and PCR testing must be completed, and if seedlings have tested negative, they may be transferred to Level 1 PEQ for woody indexing.

Approved treatments: Not required

Phytosanitary certificate: Required

2.51.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.51.2Inspection and testing requirements

Organism	MPI acceptable detection methods (listed below)	
Apple scar skin viroid	PCR and woody indexing	
Monilinia fructigena	Growing season inspection in PEQ for disease symptom expression	
Tomato bushy stunt virus	PCR and herbaceous Indexing	

- (1) Tests are to be carried out on plants germinated from the imported seeds.
- (2) The quarantine period will begin once the plants have entered a period of active growth and have two fully expanded leaves.
- (3) Virus testing is to be conducted on new spring growth. Viroid testing is to be done during the summer period. For each *Malus* plant, at least two fully-expanded leaves must be sampled from different branches of the main stem, one a younger leaf and one an older leaf.
- (4) Polymerase chain reaction (PCR) tests. All PCR tests must be validated using positive and negative controls prior to use in quarantine testing. Positive and negative controls must be used in all tests. Internal control primers to check the PCR competency of the samples and a negative plant control should also be used in PCR tests.
- (5) Herbaceous indexing will use the indicators Chenopodium quinoa and Nicotiana clevelandii (Nc).
- (6) Woody Indexing will use one of the indicators *Malus* x *domestica* 'Golden Delicious' or 'Red Delicious' Delicious'.
- (7) Inspection of the Malus plants by the Operator of the PEQ facility for signs of pest and disease must be at least twice per week for the first 3 months of active growth, and during spring and autumn. All other times of active growth (summer), plants should be inspected once per week. A record of inspections carried out by the Operator is to be kept and made available to the MPI Inspector on request.

Ministry for Primary Industries Page 86 of 168

Guidance

- Seedlings will be inspected and tested for regulated pests at the expense of the importer. The
 quarantine period may be extended if material is slow growing, pests are detected, or further testing is
 required.
- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.



2.52 Mangifera

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Mangifera*."

Approved countries: All

Quarantine pests: Sternochetus mangiferae, Xanthomonas campestris pv. mangiferae-indicae

Import permit: Required

PEQ: Level 1

Minimum period: 2 growing seasons

Isolation: 50 m exclusion area

Approved treatment: Not required

Phytosanitary certificate: Required

2.52.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Mangifera* seeds have been collected from trees which were inspected during the growing season and *Xanthomonas campestris* pv. *mangiferae-indicae* was not detected".

Guidance

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 88 of 168

2.53 Medicago

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Medicago*."

Approved countries: All

Quarantine pests: Pea early browning virus, Peanut stunt virus, Trogoderma granarium, Xanthomonas campestris pv. alfalfae.

Import Permit: Permit not required, unless seeds are to be grown in PEQ.

PEQ: Not required, unless imported under options 2.2.2 or 2.2.3 of the MPI <u>Protocol</u> for Testing for the Presence of Genetically Modified Plant Material.

Approved treatment: Not required

Phytosanitary certificate: Required

2.53.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Medicago* seeds have been inspected in accordance with appropriate official procedures and found to be free of *Trogoderma granarium*".

OR

b) "Trogoderma granarium" is absent/not known to occur in (name of country)

AND

- c) "The *Medicago* seeds have been:
 - i) produced in a 'pest free area' free from *Pea early browning virus*, *Peanut stunt virus* and *Xanthomonas campestris* pv. *alfalfae*;

OR

ii) produced in a 'pest free place of production' free from *Pea early browning virus, Peanut stunt virus* and *Xanthomonas campestris* pv. *alfalfae*".

2.53.2 GM seed testing

(1) In addition to the phytosanitary requirements above, all consignments of *Medicago sativa* (lucerne/ alfalfa) are required to be representatively sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5. 3: *Genetically Modified Testing Certificate*).

Guidance

- The MPI Protocol for Testing for the Presence of Genetically Modified Plant Material can be found at https://www.mpi.govt.nz/document-vault/10250
- More information on genetically modified seeds can also be found at https://www.mpi.govt.nz/importing/plants/seeds-for-sowing/genetically-modified-seeds/

Ministry for Primary Industries Page 89 of 168

2.54 Myrtaceae

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed for Sowing IHS "see 155.02.05 under *Myrtaceae*".

Approved countries: All

Quarantine pests: Austropuccinia psidii (formerly Puccinia psidii or Uredo rangelii)

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.54.1 Approved treatments

(1) In lieu of country freedom for *Austropuccinia psidii* all *Myrtaceae* seeds must be treated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

2.54.2 Phytosanitary certificate - Additional declarations

- (1) The required additional declarations must be endorsed in full on the phytosanitary certificate, no variations in the wording will be accepted by MPI, with exception of translation artefacts.
- (2) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (3) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this Import Health Standard and also the following additional declaration (s) to the phytosanitary certificate:

a)	"Austropuccinia psidii is	absent in	(name of country)'
ω_j	riada opadonna polan id	- aboont iii	 (marrie or ocurrery)

OR

b) The *Myrtaceae* seeds must be treated with one of the approved fungicide combinations as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 90 of 168

2.55 Nicotiana tabacum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Nicotiana tabacum*."

Approved countries: All

Quarantine pests: Peronospora tabacina

Phytosanitary certificate: Required

(1) Importers must comply with one of the three options listed below:

2.55.1 Option 1: Offshore measure

Import Permit: Not required

PEQ: Not required

Approved treatment

a) All *Nicotiana tabacum* seeds must be treated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

Phytosanitary certificate - Additional declarations:

- a) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- b) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration to the phytosanitary certificate:
 - i) "The *Nicotiana tabacum* seed in this consignment have been inspected during the growing season and no *Peronospora tabacini* was detected";

OR

ii) "produced in an area where Peronospora tabacini is not known to occur".

2.55.2 Option 2: Onshore measure

Import Permit: Not required

PEQ: Not required

Approved treatment

a) All *Nicotiana tabacum* seeds must be treated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

Phytosanitary certificate

a) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Phytosanitary requirements

- a) All seeds must be imported untreated and tested at an MPI-approved testing laboratory for Peronospora tabacini prior to any treatment occurring.
- b) Only seeds tested and found free from *Peronospora tabacini* will be treated with one of the fungicide combinations in the MPI approved treatments prior to biosecurity clearance.

Ministry for Primary Industries Page 91 of 168

2.55.3 Option 3: Onshore measure

Import Permit: Required

PEQ: Level 3B

Minimum PEQ period: 1 growing season


Approved treatment: Not required

Phytosanitary certificate

a) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments



2.56 Oxyria

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Oxyria*."

Approved countries: All

Quarantine pests: Ustilago vinosa

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.56.1 Approved treatments

(1) All Oxyria seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.56.2 Phytosanitary certificate

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 93 of 168

2.57 Panicum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Panicum*."

Approved countries: All

Quarantine pests: Trogoderma spp., Ustilaginales

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.57.1 Approved treatments

(1) All *Panicum* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.57.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 94 of 168

2.58 Papaver somniferum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Papaver somniferum*."

Approved countries: All

Quarantine pests: None

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.58.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

• Importers of *Papaver somniferum* must obtain written approval from Medsafe before importation.

Medicines Control

Medsafe

Email: medicinescontrol@health.govt.nz

Phone: (04) 816 2444

Ministry for Primary Industries Page 95 of 168

2.59 Persea

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Persea*".

Approved countries: USA

Quarantine pests: Avocado sunblotch viroid, Blackstreak

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 1 growing season

Approved treatment: Not required

Phytosanitary certificate: Required

2.59.1 Phytosanitary certificate

(1) If satisfied the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Ministry for Primary Industries Page 96 of 168

2.60 Petunia

These requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Petunia*".

Approved countries: All

Quarantine pests: Potato spindle tuber viroid, Tomato chlorotic dwarf viroid

Import Permit: Permit not required, unless seeds are to be grown in PEQ.

PEQ: Not required, unless imported under options 2.2.2 or 2.2.3 of the MPI <u>Protocol</u> for Testing for the Presence of Genetically Modified Plant Material.

Approved treatment: Not required

Phytosanitary certificate: Required

2.60.1 Phytosanitary requirements

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration(s) to the phytosanitary certificate:
 - a) "The [insert species name] seeds for sowing have been:
 - i) produced in a 'pest free area' free from Potato spindle tuber viroid;

OR

ii) produced in a 'pest free place of production', where parent plants have been tested according to a NPPO approved methodology and found free from *Potato spindle tuber viroid*:

OR

- officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology using a NPPO approved PCR testing method, and found to be free from *Potato spindle tuber viroid*".
- b) "The [insert species name] seeds for sowing have been:
 - i) produced in a 'pest free area' free from Tomato chlorotic dwarf viroid;

OR

ii) produced in a 'pest free place of production', where parent plants have been tested according to a NPPO approved methodology and found free from *Tomato chlorotic dwarf viroid*:

OR

- iii) officially tested, on a representative sample of a minimum of 3000 seeds officially drawn according to the ISTA or AOSA sampling methodology using a NPPO approved PCR testing method, and found to be free from *Tomato chlorotic dwarf viroid*".
- (2) The full scientific name of the *Petunia* species and variety must be specified on the phytosanitary certificate.

2.60.2 GM seed testing

(1) For all lots of *Petunia*, in addition to the phytosanitary requirements above, importers are required to comply with **one of** the two options listed below:

Ministry for Primary Industries Page 97 of 168

Option 1:

a) a declaration signed by the exporter and importer must accompany the consignment declaring that the consignment does not contain GM seeds (refer to Appendix 3: Declaration form).

Option 2:

b) samples from each lot must be representatively sampled, tested, and found to be free of unapproved GM seed according to the MPI Protocol for Testing for the Presence of Genetically Modified Plant Material (refer to Part 1.5.3 Genetically modified testing certificate in this IHS). Every lot tested must be specified on the testing certificate.

2.60.3 Testing requirements

- (2) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (3) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (4) Testing on-shore will be performed using an MPI-approved testing method.

Guidance

- The MPI Protocol for Testing for the Presence of Genetically Modified Plant Material can be found at https://www.mpi.govt.nz/document-vault/10250
- More information on genetically modified seeds can also be found at https://www.mpi.govt.nz/importing/plants/seeds-for-sowing/genetically-modified-seeds/

Ministry for Primary Industries Page 98 of 168

2.61 Phaseolus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Phaseolus*."

Approved countries: Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Luxembourg, the Netherlands, Portugal, Spain, Sweden, United Kingdom and United States of America.

Quarantine pests: Refer to "Pest list for Phaseolus".

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.61.1 Approved treatments

(1) In lieu of pest free area for Cochliobolus miyabeanus, Elsinoe phaseoli, and Phoma exigua var. diversispora all Phaseolus seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.61.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration(s) to the phytosanitary certificate:
 - a) "The Phaseolus seeds have been:
 - i) produced in a 'pest free area' free from the named regulated bacteria (*Curtobacterium flaccumfaciens pv. flaccumfaciens*) and viruses (*Bean common mosaic virus* [blackeye cowpea mosaic strain], *Broad bean mottle virus*, *Cowpea severe mosaic virus*, *Pea early-browning virus*, *Peanut mottle virus*, *Southern bean mosaic virus*);

OR

ii) produced in a 'pest free place of production' free from the named regulated bacteria (*Curtobacterium flaccumfaciens* pv. *flaccumfaciens*) and viruses (*Bean common mosaic virus* [blackeye cowpea mosaic strain], *Broad bean mottle virus*, *Cowpea severe mosaic virus*. *Pea early-browning virus*. *Peanut mottle virus*. *Southern bean mosaic virus*)":

OR

officially tested, on a representative sample of a minimum of 2,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved methodology, and found free from (Bean common mosaic virus [blackeye cowpea mosaic strain], Pea early-browning virus, Peanut mottle virus, Southern bean mosaic virus)";

AND

officially tested, on a representative sample of a minimum of 3,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved methodology, and found free from (*Broad bean mottle virus*, *Cowpea severe mosaic virus*)";

AND

Ministry for Primary Industries Page 99 of 168

officially tested, on a representative sample of a minimum of 4,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved methodology, and found free from *Curtobacterium flaccumfaciens pv. Flaccumfaciens*";

AND

- b) "The *Phaseolus* seeds have been:
 - i) produced in a 'pest free area' free from the named regulated fungi (Cochliobolus miyabeanus, Elsinoe phaseoli, Phoma exigua var. diversispora)";

OR

ii) "treated with one of the fungicide combinations in MPI approved treatments".

2.61.3 Testing requirements

- (1) Testing is required to be completed offshore, prior to export.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 100 of 168

Pest list for *Phaseolus* REGULATED PESTS (actionable)

Insecta		
Coleoptera		
Bostrichidae		
	Prostephanus truncatus	larger grain borer
Bruchidae		
	Acanthoscelides argillaceus	bean weevil
	Acanthoscelides obvelatus	bruchid beetle
	Bruchidius atrolineatus	seed beetle
	Bruchidius incarnatus	seed beetle
	Bruchus pisorum	pea weevil
	Callosobruchus analis	cowpea weevil
	Callosobruchus maculatus	cowpea weevil
	Callosobruchus phaseoli	cowpea weevil
	Zabrotes subfasciatus	mexican bean weevil
Lepidoptera		
Pyralidae		
	Etiella grisea	pod borer
	Etiella grisea drososcia	pod borer
	Etiella zinckenella	limabean pod borer
Tortricidae		
	Cydia fabivora	pod moth
	Matsumuraeses phaseoli	adzuki pod worm
Fungus		·
Ascomycota		
Dothideales		
Elsinoaceae		
	Elsinoe phaseoli	scab
Pleosporaceae		
	Cochliobolus miyabeanus (anamorph Bipolaris oryzae)	
	mitosporic fungi (Coelomycetes)	
Sphaeropsidales		
Sphaerioidaceae		
	Phoma exigua var. diversispora	ascochyta leaf spot
Bacterium	·	
Corynebacteriaceae		
·	Curtobacterium flaccumfaciens pv. flaccumfaciens	bacterium wilt
Virus		
	Bean common mosaic virus [blackeye cowpea mosaic strain]	
	Broad bean mottle virus	

Ministry for Primary Industries Page 101 of 168

Cowpea severe mosaic virus
Pea early-browning virus
Peanut mottle virus
Southern bean mosaic virus



Ministry for Primary Industries Page 102 of 168

2.62 Phoenix

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Phoenix*."

Approved countries: All countries except Guam, the Philippines and the Solomon Islands

Quarantine pests: Coconut cadang-cadang viroid, Fusarium oxysporum f. sp. canariensis

Import permit: Not required

PEQ: Not required

Approved treatments: Not required

Phytosanitary certificate: Required

2.62.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Phoenix* seeds have been produced in an approved country and have not been produced in Guam, the Philippines or the Solomon Islands".
- (2) If the consignment contains *Phoenix canariensis*, *Phoenix dactylifera* or *Phoenix reclinata* seeds:
 - a) "The *Phoenix* seeds have been produced in a 'pest free area' free from *Fusarium oxysporum* f. sp. canariensis".

Ministry for Primary Industries Page 103 of 168

2.63 Pinus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Pinus*."

Approved countries: All

Quarantine pests: Refer to "Pest list for Pinus."

Import permit: Required only for seeds produced in areas not known to be free from *Fusarium circinatum*.

PEQ: Level 3B - Required only for seeds produced in areas not known to be free from *Fusarium circinatum*.

Phytosanitary certificate: Required

2.63.1 Approved Treatments

(1) All *Pinus* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.63.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Pinus seeds have been:
 - i) officially inspected during the growing season according to appropriate procedures and no Dioryctria abietivorella or Conophthorus coniperda was detected.

OR

ii) inspected for evidence of the presence of insect pests and none was found".

AND

b) "The Pinus seeds have been treated with one of the fungicides in MPI approved treatments".

AND

- c) For seeds produced in areas approved by MPI as being free of Fusarium circinatum ONLY:
 - i) "The *Pinus* seeds have been produced in pest free areas that are, as verified by pest surveillance methods, free from *Fusarium circinatum* (syn. *Fusarium subglutinans* f sp. *pini*)".

OR

For seeds produced in areas not recognized by MPI as being free from *Fusarium* circinatum:

ii) Import Permit: Required PEQ: Level 3B

Minimum Period: To be determined at the time of permit issuance.

Guidance

A list of MPI approved pest free areas is provided using this link: <u>Fusarium circinatum</u>

Ministry for Primary Industries Page 104 of 168

2.63.3 Testing requirements

(1) MPI will determine, via the requirements on a permit to import, the testing required for *Pinus* spp. seeds for sowing for quarantine pests. The quarantine period will vary depending on the pests that may be associated with the commodity and the tests required.



Ministry for Primary Industries Page 105 of 168

Pest list for <u>Pinus</u> REGULATED PESTS (actionable)

nsect		
Insecta		
Coleoptera		
Anobiidae		
Ernobius p	unctulatus	borer
Cerambycidae		
Xylotrechu	s schaefferi	longhorn beetle
Curculionidae		
Conotrach	elus neomexicanus	cone borer, curculio
Scolytidae		
	Conophthorus coniperda	white pine cone beetle
	Conophthorus ponderosae	lodgepole cone beetle
	Conophthorus resinosae	red pine cone beetle
Diptera		
Cecidomyi	idae	
	Cecidomyia bisetosa	gall midge
	Resseliella silvana	gall midge
Heteroptera		
Coreidae		
	Lepispilus sulcicollis	seed eater
	Leptoglossus corculus	leaffooted pine seed bug
	Leptoglossus occidentalis	coreid bug
Scutellerid	ae	
	Tetyra bipuctata	shield backed pine seed bug
Hymenoptera		
Torymidae		
	Megastigmus albifrons	seed chalcid
Lepidoptera		
Pyralidae		
	Dioryctria abietivorella	fir coneworm, pine knothorn moth
	Dioryctria amatella	southern pine coneworm
	Dioryctria auranticella	pyralid moth
	Dioryctria clarioralis	coneworm
	Dioryctria disclusa	webbing coneworm
	Dioryctria merkeli	loblolly pine coneworm
	Dioryctria rossi	cone borer, pyralid moth
Tortricidae	;	
	Commophila fuscodorsana	tortricid moth
	Cydia anaranjada	slash pine seedworm
	Cydia ingens	logleaf pine seed worm
	Cydia miscitata	cone borer, tortricid moth

Ministry for Primary Industries Page 106 of 168

	Cydia piperana	cone borer, ponderosa pine seed moth
	Cydia toreuta	cone borer, eastern pine seedworm
Fungus		
Ascomycota		
Diaporthales		
Melancon	iidaceae	
	Melanconis stilbostoma (anamorph Melanconium bicolor)	mould
Dothideales		
Dothioraceae		
	Sydowia polyspora (anamorph Sclerophoma pythiophila)	pine leaf blight, tip dieback
Mycosphaerellac	ceae	
	Mycosphaerella dearnessii (anamorph Lecanosticta acicola)	brown needle spot
Pleosporaceae		
	Setosphaeria rostrata (anamorph Exserohilum rostratum)	leaf blight, black mould
Hypocreales		
Hypocreaceae		
	Nectria inventa (anamorph Verticillium tenerum)	verticillium rot
Pezizales		
Otideaceae		
	Caloscypha fulgens (anamorph Geniculodendron pyriforme)	cold fungus
Pyronemataceae		
	Pyronema omphalodes	mould
Mitosporic fungi		
	Coniosporium aterrimum	mould
	Lacellina graminicola	mould
Mitosporic fungi (Coelomyc	etes)	
Sphaeropsidales		
Sphaerioidaceae)	
	Botryodiplodia acicola	mould
	Coniothyrium quercinum	mould
Unknown (Coelomycetes)		
	Melanconium apiocarpon	mould
	Pestalotia breviseta	mould
	Pestalotia foedans	mould
	Pestalotiopsis glandicola Sirococcus conigenus	mould

Ministry for Primary Industries Page 107 of 168

Hyphomycetales	1	
Hyphomycetales		
71 - 7	Cladosporium cucumerinum	black mould
	Cladosporium naumovi	black mould
	Curvularia inaequalis	black mould
	Stemphylium piriforme	leaf mould
Moniliaceae		
	Acremonium subverticillatum	mould
	Aspergillus funiculosus	mould
	Penicillium arenarium	penicillium mould rot
	Penicillium aurantiogriseum	penicillium mould rot
	Penicillium brevicompactum	penicillium mould rot
	Penicillium canadense	penicillium mould rot
	Penicillium chrysogenum	penicillium mould rot
	Penicillium divergens	penicillium mould rot
	Penicillium fuscum	penicillium mould rot
	Penicillium gladioili	penicillium mould rot
	Penicillium oxalicum	penicillium mould rot
	Penicillium viridicatum	penicillium mould rot
	Torula convoluta	mould
	Verticillium albo-atrum [severe strain]	mould
Tuberculariales	[covere outain]	
Tuberculariaceae		
	Fusarium arthrosporoides	dry rot
	Fusarium chlamydosporum	root and stem rot
	Fusarium circinatum (syn. Fusarium subglutinans f. sp. pini)	pine pitch canker
	Fusarium moniliforme var. intermedium	mould
	Fusarium polyphialidicum	fusarium mould
Unknown (Hyphomycetes)		
	Oidium verticilloides	mould
Oomycota		
Pythiales		
Pythiaceae		
	Pythium aphanidermatum	root and seed rot
Zygomycota: Zygomycetes		
Mucorales		
Mucoraceae		
	Mucor hiemalis	mucor fruit rot
	Mucor mucedo	mucor fruit rot
	Waddi Maddad	- Indoor Halt Fot

Ministry for Primary Industries Page 108 of 168

Muco	r racemosus	storage rot
Muco	r ramanianus	mould
Syncephalastraceae		
Sync	ephalastrum racemosum	mould



Ministry for Primary Industries Page 109 of 168

2.64 Pisum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Pisum*."

Approved countries: Australia, Austria, Belgium, Canada, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, the Netherlands, Portugal, Spain, Sweden, Taiwan, United Kingdom and United States of America.

Quarantine pests: Refer to "Pest list for Pisum".

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.64.1 Approved Fumigation treatment

- (1) All lots of *Pisum* seed imported into New Zealand are required to be fumigated according to the specifications listed in MPI-STD-ABTRT Approved Biosecurity Treatment.
- (2) The treatment is required to be completed offshore prior to export, or on arrival in New Zealand by an MPI approved treatment provider.
- (3) Pre-export treatment for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, where the fumigant used and application rate must be clearly stated, or if done on arrival in New Zealand, must be completed at an MPI-approved facility.

2.64.2 Approved Fungicide Treatments

(1) In lieu of pest free area for *Cladosporium cladosporioides f. sp. pisicola* all pisum seed must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.64.3 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Pisum* seeds have been:
 - i) produced in a 'pest free area' free from Broad bean mottle virus, Broad bean stain virus, Pea early-browning virus, Peanut mottle virus, Peanut stunt virus";

OR

ii) produced in a 'pest free place of production' free from Broad bean mottle virus, Broad bean stain virus, Pea early-browning virus, Peanut mottle virus, Peanut stunt virus";

OR

officially tested, on a representative sample of a minimum of 2,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved methodology, and found free from *Pea early-browning virus* and *Peanut mottle virus*"; AND

Ministry for Primary Industries Page 110 of 168

officially tested, on a representative sample of a minimum of 3,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved methodology, and found free from *Broad bean mottle virus*, *Broad bean stain virus* and *Peanut stunt virus*";

AND

- b) "The Pisum seeds have been:
 - i) produced in a 'pest free area' free from Cladosporium cladosporioides f. sp. Pisicola";

OR

ii) treated with one of the fungicide combinations in MPI approved treatments".

2.64.4 Testing requirements

- (1) Testing is required to be completed offshore, prior to export.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate.

Guidance

- MPI may verify treatment certification provided from both offshore and onshore treatments through an audit sampling regime, as per ISPM 20. Guidelines for a phytosanitary import regulatory system.
- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 111 of 168

Pest list for *Pisum* REGULATED PESTS (actionable)

Insecta	
Coleoptera	
Bruchidae	
Acanthoscelides zeteki	bruchid beetle
Bruchidius atrolineatus	seed beetle
Bruchidius incarnatus	seed beetle
Bruchidius quinqueguttatus	bruchid beetle
Bruchus affinis	bruchid beetle
Bruchus emarginatus	Mediterranean pulse beetle
Bruchus ervi	bruchid beetle
Bruchus lentis	bruchid beetle
Bruchus pisorum	pea weevil
Bruchus rufimanus	broad bean weevil
Bruchus tristis	bruchid beetle
Callosobruchus analis	cowpea weevil
Callosobruchus chinensis	oriental cowpea weevil
Callosobruchus maculatus	cowpea weevil
Dermestidae	
Trogoderma granarium	khapra beetle
Lepidoptera	
Lycaenidae	
Euchrysops cnejus	blue butterfly
Noctuidae	
Spodoptera praefica	western yellowstriped armyworm
Pyralidae	
Etiella zinckenella	limabean pod borer
Tortricidae	
Cydia nigricana	pea moth
Mitosporic fungi (Hyphomycetes)	
Hyphomycetales	
Dematiaceae	
Cladosporium cladosporioides f. sp. pisicola	cladosporium blight
Virus	
Broad bean mottle virus	
Broad bean stain virus	
Pea early-browning virus	
Peanut mottle virus	
Peanut stunt virus	

Ministry for Primary Industries Page 112 of 168

2.65 Populus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Populus*."

Approved countries: Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, and United States of America

Quarantine pests: Marssonina spp.

Import permit: Required

PEQ: Level 2 and Level 1

Minimum PEQ period: 2 growing seasons as follows:

a) in a Level 2 quarantine facility for the first season;

b) in a Level 1 quarantine facility subsequently.

Isolation: 50m exclusion area when planted outside.

Approved treatment: Not required

Phytosanitary certificate: Required

2.65.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard

Guidance

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 113 of 168

2.66 Prunus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Prunus*."

Approved countries: All

Quarantine pests: Eurytoma amygdali, Cherry leaf roll virus [strains not in New Zealand], Cherry rasp leaf virus, Prune dwarf virus [strains not in New Zealand], Prunus necrotic ringspot virus [strains not in New Zealand], Plum pox virus, Tomato bushy stunt virus, Cucumber green mottle mosaic virus (CGMMV), Monilinia fructigena

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 6 months

Approved treatment: Not required

Phytosanitary certificate: Required

2.66.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Prunus* seeds have been inspected in accordance with appropriate official procedures and found to be free of *Eurytoma amygdali*."

2.66.2 Inspection and testing requirements

Organism	MPI acceptable detection methods
Monilinia fructigena	Growing season inspection in PEQ for disease symptom expression.
Cherry leaf roll virus*	ELISA (Agdia) or PCR AND herbaceous indicators Cq, Cs.
Cherry rasp leaf virus	ELISA or PCR using the method of James et al. (1991) AND herbaceous indicators Cq, Cs.
Plum pox virus	Durviz ELISA (Agdia) or PCR using the method of Wetzel et al. (1991) AND herbaceous indicators Nc and Cf.
Prunus necrotic ringspot virus*	ELISA (Agdia) or PCR using the method of Spiegel et al. (1996) AND herbaceous indicators Cs.
Tomato bushy stunt virus*	ELISA (Agdia) or PCR AND herbaceous indicators Cq, Nc.
Cucumber green mottle mosaic virus	ELISA or PCR

Indicator hosts: Chenopodium foetidum (Cf), Chenopodium quinoa (Cq), Cucumis sativus (Cs) and Nicotiana clevelandii (Nc).

- (1) For bioassay and ELISA, plants shall be sampled from at least two positions on every stem including a young, fully expanded leaflet at the top of each stem and an older leaflet from a midway position.
- (2) Indicator plants must be grown under appropriate temperatures.
- (3) Indicator plants must be shaded for 12-24 hrs prior to inoculation.

Ministry for Primary Industries Page 114 of 168

- (4) Maintain post-inoculated indicator species under appropriate glasshouse conditions for at least 4 weeks.
- (5) Inspect plants at least once per week for signs of pest and disease.
- (6) Inspect inoculated herbaceous indicator plants at least twice per week for symptoms of virus infection.
- (7) At least two plants of each indicator species unless otherwise stated must be used in mechanical inoculation tests.
- (8) Positive and negative controls must be used in ELISA tests.
- (9) Testing must be carried out on plants while they are in active growth.
- (10) Positive and negative controls (including a blank water control) must be used in PCR.
- (11) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).
- (12) ELISA or PCR for PPV must test negative before herbaceous indicator tests are conducted.

Guidance:

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.

References:

- James D, Howell WE, Mink GI, 2001. Molecular evidence of the relationship between a virus associated with flat apple disease and Cherry rasp leaf virus as determined by RT-PCR. Plant Disease 85, 47-52.
- Spiegel S, Scott SW, BowmanVance V, Tam Y, Galiakparov NN, Rosner A, 1996. Improved detection
 of prunus necrotic ringspot virus by the polymerase chain reaction. European Journal of Plant
 Pathology 102, 681-685.
- Wetzel T, Candresse T, Ravelonandro M, Dunez J, 1991. A polymerase chain-reaction assay adapted to plum pox potyvirus detection. Journal of Virological Methods 33, 355-365.

Ministry for Primary Industries Page 115 of 168

2.67 Pseudotsuga menziesii

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Pseudotsuga menziesii.*"

Approved countries: All

Quarantine pests: Refer to "Pest list for Pseudotsuga menziesii".

Import permit: Required only for seeds produced in areas not known to be free from *Fusarium circinatum*

PEQ: Level 3B - Required only for seeds produced in areas not known to be free from Fusarium circinatum

Phytosanitary certificate: Required

2.67.1 Approved treatment

(1) All *Pseudotsuga menziesii* seeds must be treated as per MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments.

2.67.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Pseudotsuga menziesii seeds have been:
 - i) collected from trees that have been officially inspected during the growing season according to appropriate procedures and no *Dioryctria abietivorella* was detected.

OR

ii) inspected for evidence of the presence of insect pests and none was found".

AND

b) "The Pseudotsuga menziesii seeds have been treated for regulated pests".

AND

- c) For seeds produced in areas approved by MPI as being free of Fusarium circinatum ONLY:
 - i) The *Pseudotsuga menziesii* seeds for sowing have been produced in pest free areas that are, as verified by pest surveillance methods, free from *Fusarium circinatum* (syn. *Fusarium subglutinans* f sp. *pini*).

OR

For seeds produced in areas not recognized by MPI as being free from *Fusarium* circinatum:

ii) Import Permit: Required

PEQ: Level 3B

Minimum PEQ period: To be determined at the time of permit issuance.

Guidance

- A list of MPI approved pest free areas is provided using this link: Fusarium circinatum.
- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 116 of 168

2.67.3 Testing requirements

(1) MPI will determine, via the requirements on a permit to import, the testing required for *Pseudotsuga menziesii* seeds for sowing for quarantine pests. The quarantine period will vary depending on the pests that may be associated with the commodity and the tests required.



Ministry for Primary Industries Page 117 of 168

Pest list for Pseudotsuga menziesii REGULATED PESTS (actionable)

Insecta		
Coleoptera		
Anobiid		
	Ernobius punctulatus	borer
Curcul	ionidae	
	Lepesoma lecontei	weevil
Scarab	paeidae	
	Melolontha melolontha	cockchafer
Diptera		
Cecido	omyiidae	
	Asynapta keeni	gall midge
	Contarinia constricta	gallmidge
	Contarinia cuniculator	gall midge
	Contarinia oregonensis	douglas fir cone gall midge
	Contarinia pseudotsugae	gall midge
	Contarinia washingtonensis	gall midge
Loncha		
	Earomyia aquilonia	fir seed maggot
	Earomyia barbara	fir seed maggot
Hemiptera		
Coreid	ae	
	Leptoglossus occidentalis	coreid bug
Lepidoptera		
Blastol	pasidae	
	Holcocera augusti	blastobasid moth
Geome	etridae	
	Eupithecia albicapitata	looper
	Eupithecia spermaphaga	looper
Pyralid		
	Dioryctria abietivorella	fir coneworm
Tortrici		
	Barbara colfaxiana	douglas fir cone moth
_	Chionodes periculella	gelechiid moth
	Commophila fuscodorsana	tortricid moth
	Endopiza piceana	tortricid moth
	Laspeyresia bracteatana	leafroller
	Zeiraphera diniana	douglas fir cone moth
ungus		
Ascomycota		
Pezizales		
Otidea	COSO	

Ministry for Primary Industries Page 118 of 168

	Caloscypha fulgens	cold fungus
Mitosporic Fungi (Hyphon	nycetes)	
Hyphomycetales		
Moniliaceae		
	Penicillium chrysogenum	penicillium mould rot
Tuberculariales		
Tuberculariace	eae	
	Fusarium circinatum (syn. Fusarium subglutinans f. sp. pini)	pine pitch canker



Ministry for Primary Industries Page 119 of 168

2.68 Psophocarpus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Psophocarpus*."

Approved countries: All

Quarantine pests: Etiella spp., Maruca testulali, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

For Seed in pods ONLY:

2.68.1 Phytosanitary certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - "The *Psophocarpus* pods have been inspected before export and no caterpillars of *Etiella* spp. or *Maruca testulalis* were found in a 600 unit sample".

Ministry for Primary Industries Page 120 of 168

2.69 Pyrus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Pyrus*."

Approved countries: All

Quarantine pests: Apple scar skin viroid, Monilinia fructigena, Tomato bushy stunt virus, Pear bark measle

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 6 months

Approved treatment: Not required

Phytosanitary certificate: Required

2.69.1 Phytosanitary requirements

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must provide the certifying statement as per Part 1.5.2 of this import health standard.

2.69.2 Inspection and testing requirements

Organism	MPI acceptable detection methods
Monilinia fructigena	Growing season inspection in PEQ for disease symptom expression.
Apple scar skin viroid	PCR using the method of Hadidi et al. (1990).
Tomato bushy stunt virus*	ELISA (Agdia) or PCR AND herbaceous indicators Cq, Nc.
Pear bark measle	Growing season inspection in PEQ for disease expression.

Indicator hosts: Chenopodium quinoa (Cq) and Nicotiana clevelandi (Nc).

- (1) For bioassay and ELISA, plants shall be sampled from at least two positions on every stem including a young, fully expanded leaflet at the top of each stem and an older leaflet from a midway position.
- (2) Indicator plants must be grown under appropriate temperatures.
- (3) Indicator plants must be shaded for 12-24 hrs prior to inoculation.
- (4) Maintain post-inoculated indicator species under appropriate glasshouse conditions for at least 4 weeks.
- (5) Inspect plants at least once per week for signs of pest and disease.
- (6) Inspect inoculated herbaceous indicator plants at least twice per week for symptoms of virus infection.
- (7) PCR and ELISA need to be validated using positive controls/reference material prior to use in quarantine testing.
- (8) At least two plants of each indicator species unless otherwise stated must be used in mechanical inoculation tests.
- (9) Positive and negative controls must be used in ELISA tests.
- (10) Testing must be carried out on plants while they are in active growth.
- (11) Positive and negative controls (including a blank water control) must be used in PCR.

Ministry for Primary Industries Page 121 of 168


(12) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

References:

 Hadidi A, Yang X, 1990. Detection of pome fruit viroids by enzymatic cDNA amplification. Journal of Virological Methods 30, 261-269.



2.70 Quercus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Quercus*."

Approved countries: Australia, Canada, Germany, India, Israel, Japan, Mexico, Spain, Tunisia, United Kingdom and United States of America

Quarantine pests: Ceratocystis fagacearum, Cryphonectria parasitica, Curculionidae

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 2 years

Phytosanitary certificate: Required

2.70.1 Approved treatments

(1) All Quercus seeds must be fumigated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.70.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Quercus seeds have been:
 - i) collected from trees that have been officially inspected during active growth and no diseases caused by *Ceratocystis fagacearum* or *Cryphonectria parasitica* were detected;

OR

ii) produced in an area where *Ceratocystis fagacearum* and *Cryphonectria parasitica* are not known to occur".

AND

b) "The Quercus seeds have been fumigated with methyl bromide at ____ pressure for ___ hours at ____ g/m³ at a temperature of ____ °C";

2.70.3 Inspection and testing requirements

Organism	MPI acceptable detection methods
Ceratocystis fagacearum	Growing season inspection in PEQ for disease symptom expression.
Cryphonectria parasitica	Growing season inspection in PEQ for disease symptom expression.

Guidance

- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at PlantImports@mpi.govt.nz well in advance of their intended import to discuss their

Ministry for Primary Industries Page 123 of 168

importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.



Ministry for Primary Industries Page 124 of 168

2.71 Raphanus sativus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Raphanus sativus*."

Approved countries: All

Quarantine pests: None

Import permit: Not required


PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.71.1 Phytosanitary certificate - Additional declaration

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.



2.72 Ribes

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "See 155.02.05 under *Ribes*."

Approved countries: All

Quarantine pests: Refer to pest list for Ribes

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 6 months

Approved treatment: Not required

Phytosanitary certificate: Required

2.72.1 Phytosanitary requirements

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.72.2 Inspection and testing requirements

Organism	MPI acceptable detection methods	
For both "Currant type" and "Gooseberry types" Ribes		
Raspberry ringspot virus*	ELISA or PCR AND herbaceous indexing with Ca and Cq OR Cq, Cs and Nc	
For "Currant type" Ribes only		
Tobacco rattle virus [strains not in New Zealand]	Herbaceous indexing with Ca and Cq OR Cq, Cs and Nc.	

Indicators: Chenopodium amaranticolor, Ca - Chenopodium quinoa, Cq – Cucumis sativus, Cs – Nicotiana clevelandii, Nc.

- (1) Tests are to be carried out on plants germinated from the imported seeds.
- (2) Testing must be carried out on plants while they are in active growth.
- (3) Indicator plants must be grown under appropriate temperatures.
- (4) Indicator plants must be shaded for 12-24 hrs prior to inoculation.
- (5) For each *Ribes* plant, at least two fully-expanded leaves must be sampled from different branches of the main stem, one a younger leaf and one an older leaf.
- (6) Post-inoculated indicator plants must be maintained under appropriate glasshouse conditions for at least 4 weeks.
- (7) Post-inoculated indicator plants must be inspected at least twice per week for signs of virus infection with observations being recorded on a weekly basis.

Ministry for Primary Industries Page 126 of 168

- (8) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).
- (9) PCR and ELISA need to be validated using positive controls/reference material prior to use in quarantine testing;
- (10) Positive, negative, and buffer controls must be used in ELISA tests.
- (11) Positive controls must be used in PCR.
- (12) Inspection of the *Ribes* plants by the operator of the PEQ facility for signs of pest and disease must be at least once per week.

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Pest list for <u>Ribes</u> REGULATED PESTS (actionable)

Virus	
	Raspberry ringspot virus
	Tobacco rattle virus (strains not in New Zealand)
	*For organisms intercepted that are not listed within this pest list refer to Biosecurity Organisms Register for Imported Commodities (<u>ONZPR</u>) to determine the regulatory status.

Ministry for Primary Industries Page 127 of 168

2.73 Rubus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Rubus*".

Approved countries: All

Quarantine pests: Refer to pest list for Rubus

Import permit: Required

PEQ: Level 2

Minimum PEQ period: 3 months

Approved treatment: Not required

Phytosanitary certificate: Required

2.73.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.73.2 Inspection and testing requirements

Organism	MPI acceptable detection methods
Raspberry ringspot virus	ELISA or PCR and herbaceous indexing with Cq, Cs and Nc
Tomato ringspot virus	ELISA or PCR and herbaceous indexing with Cq, Cs and Nc

Indicators: Chenopodium guinoa, Cg – Cucumis sativus, Cs – Nicotiana clevelandii, Nc.

- (1) Tests are to be carried out on plants germinated from the imported seeds.
- (2) The quarantine period will begin once the plants have entered a period of active growth and have two fully expanded leaves.
- (3) Virus testing needs to be conducted on new spring growth. For each *Rubus* plant, at least two young fully- expanded leaves must be sampled from different branches of the main stem, one a younger leaf and one older leaf.
- (4) Herbaceous Indicator plants must be grown under appropriate temperatures at 18-25 °C.
- (5) Indicator plants must be shaded for 24 hrs prior to inoculation.
- (6) Post-inoculated indicator plants must be maintained under appropriate glasshouse conditions for at least 4 weeks.
- (7) Post-inoculated indicator plants must be inspected at least twice per week for signs of virus infection with observations being recorded on a weekly basis.
- (8) Positive and negative (buffer solution) controls must be used on all herbaceous indexing tests.
- (9) PCR and ELISA tests need to be validated using positive controls/reference material prior to use in quarantine testing;
- (10) Positive, negative, and buffer controls must be used in ELISA tests.
- (11) Positive and negative controls must be used in PCR.

Ministry for Primary Industries Page 128 of 168


(12) Inspection of the *Rubus* plants by the operator of the PEQ facility for signs of pest and disease must be at least twice per week during periods of active growth.

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.
- Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit

Pest list for <u>Rubus</u> REGULATED PESTS (actionable)

Virus		
	Raspberry ringspot virus	
	Tomato ringspot virus	
	*For organisms intercepted that are not listed within this pest list refer to Biosecurity Organisms Register for Imported Commodities (<u>ONZPR</u>) to determine the regulatory status.	



2.74 Sesamum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under Sesamum."

Approved countries: All

Quarantine pests: Alternaria sesami, Cercoseptoria sesami, Xanthomonas campestris pv. sesami, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.74.1 Approved treatment

(1) All Sesamum seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.74.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Sesamum seeds have been:
 - i) produced in a 'pest free area' free from Alternaria sesami, Cercoseptoria sesami and Xanthomonas campestris pv. sesami;

OR

ii) produced in a 'pest free place of production' free from *Alternaria sesami*, *Cercoseptoria sesami* or *Xanthomonas campestris pv. sesami*'.

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 130 of 168

2.75 Solanum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Solanum*." For *Solanum lycopersicum* and *Solanum tuberosum*, please refer to the individual schedules which follow.

Approved countries: All

Quarantine pests: Potato spindle tuber viroid.

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.75.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *[insert species name]* seeds for sowing have been:
 - i) produced in a 'pest free area' free from Potato spindle tuber viroid;

OR

ii) produced in a 'pest free place of production' free from Potato spindle tuber viroid";

OR

b) "The [insert species name] seeds for sowing have been officially tested, on a representative sample and using appropriate methods, and found to be free from Potato spindle tuber viroid".

2.75.2 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) For seed lots of 15,000 or more seeds:
 - A representative sample of a minimum of 3000 seeds, officially drawn according to ISTA or AOSA methodology is required from each seed lot and tested as specified in the schedule;
- (4) For Solanum melongena seed lots with less than 15,000 seeds:
 - a) A composite sample of a minimum of 3000 seeds must be officially drawn across all seed lots of the same seed species in a consignment, which must have been produced at the same place of production or production site.
- (5) In order to achieve a composite sample, proportionate sampling must be carried out across all lots of *Solanum melongena* imported. A sample of seeds must be drawn from each imported lot within a consignment, adding up to 3000 seeds. The size of the sample from each lot must be proportionate to the size of the imported lot within the consignment.
- (6) All importers who test a composite sample on arrival in New Zealand must make a <u>declaration</u> to identify the place of production or production site of the lots that form the composite sample.

Ministry for Primary Industries Page 131 of 168

Guidance


- The sample size from each lot to form the composite sample should be calculated as follows:
 - a) The proportion of each lot in the total consignment (seed number) is calculated using the following equation:

 $Proportion \ of \ total \ consignment \ size = \frac{\textit{No.of seeds in each lot}}{\textit{Total number of seeds in consignmen}}$

b) Calculate the sample size for each lot (number of seeds) using a total composite sample size of 3000 seeds:

Sample size of each line= 3000 seeds x proportion of total consignment size

c) Take the sum of the sample size for each lot to check that the total composite sample for the consignment is at least 3000 seeds.



2.76 Solanum lycopersicum

The following requirements only apply to species in the Plants Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Solanum lycopersicum*."

Approved countries: All countries

Quarantine pests: Columnea latent viroid, Pepino mosaic virus, Potato spindle tuber viroid, Tomato chlorotic dwarf viroid, Tomato brown rugose fruit virus, Tomato apical stunt viroid, Tomato planta macho viroid, Tomato mottle mosaic virus

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.76.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the preshipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declarations to the phytosanitary certificate:
 - a) "The Solanum lycopersicum seeds have been prepared to industry standards with thorough cleaning to remove all traces of flesh from the seeds."

AND

- b) "The Solanum lycopersicum seeds have been:
 - i) produced in a 'pest-free area' free from Pepino mosaic virus.

OR

ii) produced in a 'pest-free place of production' free from *Pepino mosaic virus*.

OR

officially tested, on a representative sample of a minimum of 3,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved ELISA or NPPO-approved PCR testing method, and found to be free from *Pepino mosaic virus*."

AND

- c) "The Solanum lycopersicum seeds have been:
 - produced in a 'pest-free area' free from Columnea latent viroid, Potato spindle tuber viroid, Tomato apical stunt viroid, Tomato chlorotic dwarf viroid, and Tomato planta macho viroid.

OR

ii) produced in a 'pest-free place of production' free from Columnea latent viroid, Potato spindle tuber viroid, Tomato apical stunt viroid, Tomato chlorotic dwarf viroid, and Tomato planta macho viroid.

OR

iii) produced in a 'pest-free place of production' where parent plants have been tested according to an NPPO-approved methodology and found free from *Columnea latent viroid*, *Potato spindle tuber viroid*, *Tomato apical stunt viroid*, *Tomato chlorotic dwarf viroid*, and *Tomato planta macho viroid*.

Ministry for Primary Industries Page 133 of 168

OR

iv) officially tested, on a representative sample of a minimum of 3,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved PCR testing method, and found to be free from *Columnea latent viroid*, *Potato spindle tuber viroid*, *Tomato apical stunt viroid*, *Tomato chlorotic dwarf viroid*, and *Tomato planta macho viroid*."

AND

- d) "The Solanum lycopersicum seeds have been:
 - i) produced in a 'pest-free area', free from Tomato brown rugose fruit virus

OR

ii) produced in a 'pest-free place of production' free from *Tomato brown rugose fruit virus*.

OR

iii) officially tested, on a representative sample of a minimum of 3,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved PCR testing method, and found free from *Tomato brown rugose fruit virus*."

AND

- e) "The Solanum lycopersicum seeds have been:
 - i) produced in a 'pest-free area', free from *Tomato mottle mosaic virus*

OR

ii) produced in a 'pest-free place of production' free from *Tomato mottle mosaic virus*.

OR

iii) officially tested, on a representative sample of a minimum of 3,000 seeds officially drawn according to the ISTA or AOSA sampling methodology, using an NPPO-approved ELISA or NPPO-approved PCR testing method, and found free from *Tomato mottle mosaic virus*."

2.76.2 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate or, if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) Testing onshore will be performed using an MPI-approved testing method.
- (4) For seed lots of 15,000 or more seeds:
 - a) A representative sample of a minimum of 3000 seeds, officially drawn according to ISTA or AOSA methodology is required from each seed lot and tested as specified in the schedule;
- (5) For seed lots with less than 15,000 seeds:
 - a) A composite sample of a minimum of 3000 seeds must be officially drawn across all seed lots of the same seed species in a consignment, which must have been produced at the same place of production or production site.
- (6) In order to achieve a composite sample, proportionate sampling must be carried out across all lots imported. A sample of seeds must be drawn from each imported lot within a consignment, adding up to 3000 seeds. The size of the sample from each lot must be proportionate to the size of the imported lot within the consignment.
- (7) All importers who test a composite sample on arrival in New Zealand must make a <u>declaration</u> to identify the place of production or production site of the lots that form the composite sample.

Ministry for Primary Industries Page 134 of 168

Guidance

- The sample size from each lot to form the composite sample should be calculated as follows:
 - a) The proportion of each lot in the total consignment (seed number) is calculated using the following equation:

 $Proportion \ of \ total \ consignment \ size = \frac{\textit{No.of seeds in each lot}}{\textit{Total number of seeds in consignment}}$

b) Calculate the sample size for each lot (number of seeds) using a total composite sample size of 3000 seeds:

Sample size of each line= 3000 seeds x proportion of total consignment size

- c) Take the sum of the sample size for each lot to check that the total composite sample for the consignment is at least 3000 seeds.
- The ISHI-Veg local lesion bioassay for *Tomato brown rugose fruit virus* and *Tomato mottle mosaic virus* is not accepted as a valid test by MPI.
- The use of a bioassay to detect the presence of *Pepino mosaic virus* on seed samples is not accepted as a valid test by MPI.
- Additional declarations on phytosanitary certificates to meet the offshore testing requirements for Tomato brown rugose fruit virus in Import Health Standard 155.02.05: Seeds for sowing should be based only on a negative result obtained in an NPPO-approved PCR test and not on results from a bioassay.
- Additional declarations on phytosanitary certificates to meet the offshore testing requirements for Pepino
 mosaic virus and Tomato mottle mosaic virus in Import Health Standard 155.02.05: Seeds for sowing
 should be based only on a negative result obtained in an NPPO-approved ELISA or NPPO-approved
 PCR test and not on results from a bioassay.
- For tomato seed lots tested for quarantine pests onshore in New Zealand at an MPI-approved testing laboratory, additional declarations by the exporting NPPO are not required to be endorsed on the phytosanitary certificate.

Ministry for Primary Industries Page 135 of 168

2.77 Solanum tuberosum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Solanum tuberosum*."

Approved countries: All

Quarantine pests: Andean potato latent virus, Andean potato mild mosaic virus, Potato black ring virus, Potato spindle tuber viroid. Potato virus T. Tobacco ringspot virus

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 1 growing season

Approved treatment: Not required

Phytosanitary certificate: Required

2.77.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit

Ministry for Primary Industries Page 136 of 168

2.78 Sorghum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Sorghum*."

Approved countries: Australia, USA

Quarantine pests: Peronosclerospora sorghi, Sclerospora graminicola, Trogoderma spp., Ustilaginales

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.78.1 Approved treatments

(1) All Sorghum seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.78.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Sorghum seeds have been:
 - i) produced in a 'pest free area' free from *Peronosclerospora sorghi* and *Sclerospora graminicola*.

OR

ii) produced in a 'pest free place of production' free from *Peronosclerospora sorghi* and *Sclerospora graminicola*".

Guidance

• Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 137 of 168

2.79 Stenotaphrum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Stenotaphrum*."

Approved countries: All

Quarantine pests: Panicum mosaic virus

Import permit: Required

PEQ: Level 3B

Minimum PEQ period: 1 growing season

Approved treatment: Not required

Phytosanitary certificate: Required

2.79.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

Guidance

 Prospective importers who wish to import eligible species of seed under this IHS schedule should contact MPI at <u>PlantImports@mpi.govt.nz</u> well in advance of their intended import to discuss their importation plans and apply for an import permit. This is because MPI will need to undertake a risk assessment that could potentially lead to changes being made to import requirements and this could lead to longer timeframes for issuance of a permit.

Ministry for Primary Industries Page 138 of 168

2.80 Trigonella foenum-graecum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Trigonella foenum-graecum*."

Approved countries: All

Quarantine pests: Cercosporidium traversiana, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.80.1 Approved treatments

(1) All *Trigonella foenum-graecum* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.80.2 Phytosanitary certificate - Additional declaration

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Trigonella foenum-graecum* seeds have been:
 - i) produced in a 'pest free area' free from Cercosporidium traversiana;

OR

ii) produced in a 'pest free place of production' free from Cercosporidium traversiana.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 139 of 168

2.81 Triticum

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Triticum*."

Approved countries: Australia, Austria, Belgium, Canada, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Luxembourg, The Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom and United States of America.

Quarantine pests: Refer to "Pest List for Triticum".

Import permit: Not required

PEQ: Not required

Phytosanitary certificate: Required

2.81.1 Approved treatments

(1) In lieu of pest free area for *Alternaria triticina*, *Cephalosporium gramineum*, and *Curvularia verruculosa* all *Triticum* seeds must be treatedas per MPI Standard MPI-STD-ABTRT Approved Biosecurity
Treatments

2.81.2 Phytosanitary Certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Triticum* seeds have been:
 - i) produced in a 'pest free area' free from the named regulated bacteria (*Rathayibacter tritici*, *Xanthomonas campestris pv. undulosa*) and viruses (*High plains virus*, *Indian peanut clump virus*)";

OR

ii) "produced in a 'pest free place of production' free from the named regulated bacteria (Rathayibacter tritici, Xanthomonas campestris pv. undulosa) and viruses (High plains virus, Indian peanut clump virus)".

AND

- b) "The *Triticum* seeds have been:
 - i) produced in a 'pest free area' free from Anguina tritici",

OR

ii) "produced in a 'pest free place of production' free from Anguina tritici",

OR

iii) "inspected microscopically for *Anguina tritici* in accordance with appropriate official procedures, and no *Anguina tritici* spores were detected".

AND

Ministry for Primary Industries Page 140 of 168

- c) "The *Triticum* seeds have been:
 - i) produced in a 'pest free area' free from the named regulated fungi (*Alternaria triticina*, Cephalosporium gramineum, Curvularia verruculosa)";

OR

ii) "treated with one of the fungicide combinations in MPI approved treatments";

AND

- d) "The Triticum seeds have been:
 - i) produced in a 'pest free area' free from *Tilletia controversa* and *Tilletia indica*";

OR

ii) "produced in a 'pest free place of production' free from *Tilletia controversa* and *Tilletia indica* and treated with an approved fungicide treatment";

OR

iii) "a representative sample of 600 seeds, drawn from this consignment according to the International Seed Testing Associations methodology, has been tested for *Tilletia controversa* and *Tilletia indica* (and no spores of *Tilletia controversa* or *Tilletia indica* were found in a representative sample of 600 seeds drawn from this consignment) AND treated with an approved fungicide treatment".

2.81.3 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 141 of 168

$\textbf{Pest list for } \underline{\textit{Triticum}} \ \textbf{REGULATED PESTS (actionable)}$

sect	
Insecta	
Blattodea	
Blattidae	
Blatta orientalis	oriental cockroach
Coleoptera	
Bostrichidae	
Dinoderus distinctus	bostrichid beetle
Prostephanus truncatus	larger grain borer
Bruchidae	
Callosobruchus chinensis	oriental cowpea weevil
Curculionidae	
Caulophilus oryzae	broadnosed grain weevil
Dermestidae	
Trogoderma glabrum	khapra beetle
Trogoderma granarium	khapra beetle
Trogoderma grassmani	trogoderma beetle
Trogoderma inclusum	trogoderma beetle
Trogoderma ornatum	trogoderma beetle
Trogoderma simplex	dermestid beetle
Trogoderma sternale	dermestid beetle
Trogoderma variabile	warehouse beetle
Languriidae	
Pharaxonotha kirschii	Mexican grain beetle
Tenebrionidae	
Cynaeus angustus	larger black flour beetle
Latheticus oryzae	longheaded flour beetle
Palorus ratzeburgi	smalleyed flour beetle
Palorus subdepressus	depressed flour beetle
Tribolium audax	american black flour beetle
Tribolium freemani	flour beetle
Ulomoides dermestoides	darkling beetle
Diptera	
Cecidomyiidae	
Contarinia pisi	pea midge
Lepidoptera	
Noctuidae	
Faronta albilinea	wheat head armyworm
Pyralidae	,
Corcyra cephalonica	rice moth
Paralipsa gularis	stored nut moth
Tineidae	_
Cephitinea colonella	grain moth
Haplotinea insectella	casemaking moth
Psocoptera	

Ministry for Primary Industries Page 142 of 168

Liposcelidae	
Troctes minutus	psocid
Mite	
Arachnida	
Acarina	
Acaridae	
Caloglyphus krameri	
Michaelopus macfarlanei	
Eriophyidae	
Aceria tulipae (vector)	wheat curl mite
Aceria tosichella	wheat curl mite
Tarsonemidae	
Tarsonemus granarius	
Tuckerellidae	
Tuckerella ablutus	
unknown Acarina	
Paratriophtydeus coineaurius	
Nematode	
Secernentea	
Tylenchida	
Anguinidae	
Anguina tritici [vector]	seed gall nematode
Fungus	
Basidiomycota: Ustomycetes	
Ustilaginales	
Tilletiaceae	
Tilletia controversa	dwarf bunt
Tilletia indica	karnal bunt
Mitosporic fungi (Hyphomycetes)	
Hyphomycetales	
Dematiaceae	
Alternaria triticina	
Curvularia verruculosa	
Moniliaceae	
Cephalosporium gramineum	stripe
Corynebacteriaceae	
Rathayibacter tritici	yellow ear rot
Pseudomonadaceae	
Xanthomonas campestris pv. undulosa	leaf streak
Virus	
High plains virus	
Indian peanut clump virus	

Ministry for Primary Industries Page 143 of 168

2.82 Ulmus

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Ulmus*."

Approved countries: All

Quarantine pests: Cherry leaf roll virus, Elm mottle virus

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.82.1 Phytosanitary Certificate - Additional declaration

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The *Ulmus* seeds have been:
 - i) produced in trees which were officially inspected during the growing season and no *Cherry leaf roll virus* or *Elm mottle virus* was detected;

OR

ii) produced in an area where Cherry leaf roll virus and Elm mottle virus are not known to occur".

Ministry for Primary Industries Page 144 of 168

2.83 Vaccinium

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Vaccinium*."

Approved countries: All

Quarantine pests: Refer to pest list for Vaccinium

Import permit: Required.

PEQ: Level 3B

Minimum PEQ period: 6 months

Approved treatment: Not required

Phytosanitary certificate: Required

2.83.1 Phytosanitary cerificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.83.2 Inspection and testing requirements

Organism	MPI acceptable detection methods
Diaporthe vaccinii	Growing season inspection in PEQ for disease symptom expression.
Botryosphaeria vaccinii	Growing season inspection in PEQ for disease symptom expression.
Monilinia fructigena	Growing season inspection in PEQ for disease symptom expression.
Monilinia vaccinii-corymbosi	Growing season inspection in PEQ for disease symptom expression.
Blueberry shock virus*	ELISA (Agdia) or PCR AND herbaceous indicators Nb, Nc.
Blueberry leaf mottle virus*	ELISA (Agdia) or PCR AND herbaceous indicators Cq, Nc.
Peach rosette mosaic virus*	ELISA (Agdia) or PCR AND herbaceous indicators Ca, Cq
Tomato ringspot virus*	ELISA (Agdia) or PCR AND herbaceous indicators Cq, Nc.

Indicator hosts: Chenopodium amaranticolor (Ca), C. quinoa (Cq), Nicotiana benthamiana (Nb), and N. clevelandi (Nc).


- (1) For bioassay and ELISA, plants shall be sampled from at least two positions on every stem including a young, fully expanded leaflet at the top of each stem and an older leaflet from a midway position.
- (2) Indicator plants must be grown under appropriate temperatures.
- (3) Indicator plants must be shaded for 12-24 hrs prior to inoculation.
- (4) Maintain post-inoculated indicator species under appropriate glasshouse conditions for at least 4 weeks.
- (5) Inspect plants at least once per week for signs of pest and disease.
- (6) Inspect inoculated herbaceous indicator plants at least twice per week for symptoms of virus infection...
- (7) PCR and ELISA need to be validated using positive controls/reference material prior to use in quarantine testing.

Ministry for Primary Industries Page 145 of 168

- (8) At least two plants of each indicator species unless otherwise stated must be used in mechanical inoculation tests.
- (9) Positive and negative controls must be used in ELISA tests.
- (10) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).
- (11) Testing must be carried out on plants while they are in active growth. Positive and negative controls (including a blank water control) must be used in PCR.

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.



Pest list for $\underline{\textit{Vaccinium}}$ REGULATED PESTS (actionable)

Fungus			
Ascomycota			
Diaporthales			
Valsaceae			
Diaporthe vaccinii (anamorph Phomopsis vaccinii)	twig blight		
Dothideales			
Botryosphaeriaceae			
Botryosphaeria vaccinii (anamorph Phyllosticta elongata)			
Leotiales			
Sclerotiniaceae			
Monilinia fructigena (anamorph Monilia fructigena)	european brown rot		
Monilinia vaccinii-corymbosi	brown rot		
Virus			
Bromoviridae			
llarvirus			
Blueberry shock virus			
Comoviridae			
Nepovirus			
Blueberry leaf mottle virus			
Peach rosette mosaic virus			
Tomato ringspot virus [strains not in New Zealand]			

Ministry for Primary Industries Page 147 of 168

2.84 Vicia

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Vicia*."

Approved countries: Australia, Austria, Belgium, Canada, Czech Republic, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Luxembourg, the Netherlands, Portugal, Spain, Sweden, United Kingdom and United States of America.

Quarantine pests: Refer to pest list for Vicia

Import permit: Not required.

PEQ: Not required

Phytosanitary certificate: Required

2.84.1 Approved treatments

(1) All *Vicia* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.84.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Vicia seeds have been:
 - i) produced in a "pest free area' free from the named regulated viruses (Artichoke yellow ringspot virus, Broad bean mottle virus, Broad bean stain virus, Broad bean true mosaic virus, Pea early-browning virus)".

OR

ii) "produced in a 'pest free place of production' free from the named regulated viruses (Artichoke yellow ringspot virus, Broad bean mottle virus, Broad bean stain virus, Broad bean true mosaic virus, Pea early-browning virus)".

Guidance

Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments

Ministry for Primary Industries Page 148 of 168

Pest list for $\underline{\textit{Vicia}}$ REGULATED PESTS (actionable)

seed beetle
bruchid beetle
bruchid beetle
bruchid beetle
pea weevil
broad bean weevil
oriental cowpea weevil
cowpea weevil
cowpea weevil
khapra beetle
dark flour beetle
pea midge
pomegranate butterfly
_

Ministry for Primary Industries Page 149 of 168

2.85 Vigna

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Vigna*."

Approved countries: All

Quarantine pests: Curtobacterium flaccumfaciens pv. flaccumfaciens, Xanthomonas campestris pv. vignicola, Earias vitella, Maruca testulalis, Trogoderma spp.

Import permit: Not required

PEQ: Not required

Approved treatment: Not required

Phytosanitary certificate: Required

2.85.1 Phytosanitary certificate - Additional declarations

- (1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Vigna seed have been:
 - i) collected from plants which were inspected during the growing season according to appropriate procedures and no *Curtobacterium flaccumfaciens* pv. *flaccumfaciens* or *Xanthomonas campestris* pv. *vignicola* was detected.

OR

ii) produced in an area where *Curtobacterium flaccumfaciens* pv. *flaccumfaciens* and *Xanthomonas campestris* pv. *vignicola* are not known to occur".

AND [For seed in pods]:

b) "The *Vigna* seed pods were inspected before export and no caterpillars of *Earias vitella* or *Maruca testulalis* were found in a 600 unit sample".

Ministry for Primary Industries Page 150 of 168

2.86 Vitis

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Vitis*."

Approved countries: All

Quarantine pests: Grapevine angular mosaic virus, Grapevine Bulgarian latent virus, Grapevine chrome mosaic virus, Grapevine fanleaf virus [strains not in New Zealand], Grapevine line pattern virus, Grapevine yellow speckle viroid-2, Peach rosette mosaic virus, Tomato ringspot virus.

Import permit: Required

PEQ: Level 2

Minimum PEQ period: 3 months

Approved treatment: Not required

Phytosanitary certificate: Required

2.86.1 Phytosanitary certificate

(1) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard.

2.86.2 Inspection and testing requirements

Organism	MPI acceptable detection methods
Grapevine angular mosaic virus	Growing season inspection
Grapevine Bulgarian latent virus	Herbaceous indicators (Ca and Cq)
Grapevine chrome mosaic virus	Herbaceous indicators (Ca, Cq, Cs and Nt)
Grapevine fanleaf virus [strains not in New Zealand]	ELISA or PCR AND herbaceous indicators (Ca, Cq and Cs)
Grapevine line pattern virus	Growing season inspection
Grapevine yellow speckle viroid-2	PCR and Growing season inspection
Peach rosette mosaic virus*	ELISA or PCR AND herbaceous indicators (Ca, Cq, Cs and Nt)
Tomato ringspot virus*	ELISA or PCR AND herbaceous indicators (Ca and Cq)

Herbaceous indexing will use the indicators Ca - Chenopodium amaranticolor, Cq - Chenopodium quinoa, Cs - Cucumis sativus and Nt - Nicotiana tabacum.


- (1) Tests are to be carried out on plants germinated from the imported seeds.
- (2) The quarantine period will begin once the plants have entered a period of active growth and have two fully expanded leaves.
- (3) Virus testing is to be conducted on new spring growth. For each plant, at least two fully-expanded leaves must be sampled from different branches of the main stem, one a younger leaf and one an older leaf.
- (4) For ELISA tests, the unit for testing is an individual seedling because of the presence of pollen transmitted viruses for which pre-determined testing is required (denoted by '*' in the table above).

Ministry for Primary Industries Page 151 of 168

- (5) All PCR and ELISA tests must be validated using positive controls prior to use in quarantine testing. Positive and negative controls (including a blank water control for PCR) must be used in all tests.
- (6) At least two plants of each herbaceous indicator species must be used in each test. Herbaceous indicator plants must be grown under appropriate temperatures and must be shaded for 24 hrs prior to inoculation. Maintain post-inoculated indicator species under appropriate glasshouse conditions for at least 4 weeks. Inspect inoculated indicator plants at least twice per week for symptoms of virus infection.
- (7) Inspection of the *Vitis* plants by the operator of the PEQ facility for signs of pest and disease must be at least twice per week while in active growth. A record of inspections carried out by the Operator is to be kept and made available to the MPI Inspector on request.

Guidance

- Positive internal controls and a negative plant control should be used.
- Internal controls in PCR tests are important to avoid the risk of false negatives.



2.87 Zea

The following requirements only apply to species in the Plant Biosecurity Index listed under Import Specifications for Seed as "see 155.02.05 under *Zea*."

Approved countries: Australia, Austria, Canada, Chile, Finland, France, Germany, Greece, Hungary, Japan, the Netherlands, Norway, South Africa, Sweden, Switzerland, the United Kingdom and United States of America.

Quarantine pests: Acidovorax avenae subsp. avenae, Clavibacter michiganensis subsp. nebraskensis, Pantoea stewartii, High plains virus, Maize dwarf mosaic virus, Maize chlorotic mottle virus, Sugarcane mosaic virus, Botryosphaeria zeae, Cochliobolus pallescens, Cochliobolus tuberculatus, Claviceps gigantea, Gloeocercospora sorghi, Ustilago maydis, Peronosclerospora heteropogoni, P. maydis, P. philippinensis, P. sacchari, P. sorghi, Sclerophthora rayssiae var. zeae, Stenocarpella macrospora and Cephalosporium maydis.

Regulated pests: Refer to pest list for Zea

Import Permit: Permit not required, unless seeds are to be grown in PEQ.

PEQ: Not required, unless imported under options 2.2.2 or 2.2.3 of the MPI <u>Protocol</u> for Testing for the Presence of Genetically Modified Plant Material.

Phytosanitary certificate: Required

2.87.1 Approved treatments

(1) In lieu of pest free area for the regulated fungi specified all *Zea* seeds must be treated as per MPI Standard MPI-STD-ABTRT Approved Biosecurity Treatments.

2.87.2 Phytosanitary certificate - Additional declarations

- (1) The exporting country NPPO must confirm any treatment(s) as required by the IHS in the disinfestation and/or disinfection treatment section.
- (2) If satisfied that the pre-shipment activities have been undertaken, the exporting country NPPO must confirm this by providing the certifying statement as per Part 1.5.2 of this import health standard and also the following additional declaration (s) to the phytosanitary certificate:
 - a) "The Zea seeds have been:
 - i) produced in a 'pest free area' free from the named regulated bacteria *Acidovorax avenae* subsp. *avenae*, *Clavibacter michiganensis* subsp. *nebraskensis*, *Pantoea stewartii* and viruses *High plains virus* and *Maize dwarf mosaic virus*";

OR

ii) "produced in a 'pest free place of production' free from the named regulated bacteria Acidovorax avenae subsp. avenae, Clavibacter michiganensis subsp. nebraskensis, Pantoea stewartii and viruses High plains virus and Maize dwarf mosaic virus";

OR

"a representative sample, officially drawn from this consignment according to ISTA or AOSA methodology, has been tested for the presence of and found free from the named regulated bacteria Acidovorax avenae subsp. avenae, Clavibacter michiganensis subsp. nebraskensis, Pantoea stewartii and viruses High plains virus and Maize dwarf mosaic virus".

AND

Ministry for Primary Industries Page 153 of 168

- b) "The Zea seeds have been:
 - i) produced in a 'pest free area' free from the named regulated viruses *Maize chlorotic mottle virus* and *Sugarcane mosaic virus*";

OR

ii) "a representative sample, officially drawn from this consignment according to ISTA or AOSA methodology, has been tested for and found free from the named regulated viruses Maize chlorotic mottle virus and Sugarcane mosaic virus"

Guidance

- Refer section 1.11 Seeds of MPI Standard MPI-STD-ATBRT Approved Biosecurity Treatments
- Countries that MPI recognise endorsing "Pest free area" as an additional declaration for *Sugarcane mosaic virus* are as follows:
 - Australia, Austria, Canada, Finland, France, Germany, Greece, Hungary, Japan, the Netherlands, Norway, South Africa, Sweden, Switzerland, the United Kingdom and United States of America

AND

- c) "The Zea seeds have been:
 - i) produced in a 'pest free area' free from the named regulated fungi Botryosphaeria zeae, Cochliobolus pallescens, Cochliobolus tuberculatus, Claviceps gigantea, Gloeocercospora sorghi, Ustilago maydis, Peronosclerospora heteropogoni, Peronosclerospora maydis, Peronosclerospora philippinensis, Peronosclerospora sacchari, Peronosclerospora sorghi, Sclerophthora rayssiae var. zeae, Stenocarpella macrospora and Cephalosporium maydis";

OR

ii) "treated with one of the fungicide combinations in MPI approved treatments".

2.87.3 GM seed testing

(1) In addition to the phytosanitary requirements above, all consignments of *Zea mays* (sweet corn, maize) are required to be representatively sampled, tested, and found to be free of unapproved GM seed according to the Protocol (refer to Part 1.5.4: *Genetically Modified Testing Certificate*).

Guidance

- The MPI Protocol for testing for the presence of genetically modified plant material can be found at https://www.mpi.govt.nz/document-vault/10250
- More information on genetically modified seeds can also be found at https://www.mpi.govt.nz/importing/plants/seeds-for-sowing/genetically-modified-seeds/
- Popcorn does not require GM testing. The full scientific name must be specified on the phytosanitary certificate (e.g. Zea mays var. everta) to enable popcorn to be given clearance without a GM testing certificate.

2.87.4 Testing requirements

- (1) Testing is required to be completed offshore prior to export, or on arrival in New Zealand.
- (2) Pre-export testing for each seed lot must be endorsed by the NPPO on the phytosanitary certificate, or if tested on arrival in New Zealand, must be completed by an MPI-approved testing laboratory.
- (3) **Pantoea stewartii**: A negative result from testing a representative sample of a minimum of 400 seeds, using the immunosorbent assay test described by Lamka *et al.* (1991), may be used to show the consignment is free of *Pantoea stewartii* subsp. *stewartii*.;

Ministry for Primary Industries Page 154 of 168

- (4) **Clavibacter michiganensis subsp. nebraskensis**: A negative result from testing a representative sample of a minimum of 400 seeds, using the sCNS Culture Plate Method (Shepherd, 1999; www.seedhealth.org), may be used to show the consignment is *free of Clavibacter michiganensis* subsp. Nebraskensis;
- (5) **Acidovorax avenae subsp. avenae**: A negative result from testing a representative sample of a minimum of 400 seeds, using the methodology of Dange *et al.* (1978), may be used to show the consignment is free of *Acidovorax avenae* subsp. *Avenae*;
- (6) **High plains virus:** A negative result from testing a representative sample of seeds using greenhouse grow-out tests and ELISA testing as described by Forster *et al.* (2001) and Crop Plant Compendium 2003, or a representative sample of a minimum of 3000 seeds, using a PCR NPPO approved method, such as Lebas *et al.* (2005), may be used to show that the consignment is free *of High plains virus*;
- (7) **Maize dwarf mosaic virus:** A negative result from testing a representative sample of a minimum of 2000 seeds, using an NPPO approved method, may be used to show the consignment is free of *Maize dwarf mosaic virus*:
- (8) **Maize chlorotic mottle virus:** A negative result from testing a representative sample of a minimum of 3000 seeds, using ELISA or PCR testing, may be used to show the consignment is free from *Maize chlorotic mottle virus*;
- (9) Sugarcane mosaic virus: A negative result from testing a representative sample of a minimum of 2000 seeds, using an NPPO approved method, may be used to show the consignment is free of Sugarcane mosaic virus.

References:

- Dange SRS, Payak MM, Renfro BL, 1978. Seed transmission of Pseudomonas rubrilineans, the incitant of bacterial leaf stripe of maize. Indian Phytopathology 31(4):523-524.
- Forster RL, Seifers DL, Strausbaugh CA, Jensen SG, Ball EM, Harvey TL, 2001. Seed transmission of the High Plains virus in sweet corn. Plant Disease 85(7):696-699
- Lamka, G L; Hill, J H; McGee, D C; and Braun, E J. 1991: Development of an immunosorbent assay for seedborne Pantoea stewartii subsp. stewartii in corn seeds. Phytopathology 81:839-846
- Lebas, B.S.; Ochoa-Corona, F.M.; Elliot, D.R.; Tang, Z. and Alexander, B.J.R. 2005. Development of an RT-PCR for High Plains virus indexing scheme in New Zealand post entry quarantine. Plant Disease, 89:1103-1108.
- Shepherd, L.M. 1999: Detection and transmission of Clavibacter michiganensis subsp. nebraskensis of corn. Ms Thesis, Iowa State University, Ames, IA.

(10) For seed lots of 15,000 or more seeds:

- a) A representative sample of a minimum of 3400 seeds, officially drawn according to ISTA or AOSA methodology is required from each seed lot and tested as specified in the schedule;
- (11) For seed lots with less than 15,000 seeds:
 - a) Two composite samples, one of a minimum of 3000 seeds and one of a minimum of 400 seeds, must be officially drawn across all seed lots of the same seed species in a consignment, which must have been produced at the same place of production or production site.
- (12) In order to achieve a composite sample, proportionate sampling must be carried out across all lots imported. One sample of seeds must be drawn from each imported lot within a consignment, adding up to 3000 seeds and a second sample of seeds must be drawn from each imported lot within a consignment, adding up to 400 seeds. The size of the sample from each lot must be proportionate to the size of the imported lot within the consignment.
- (13) A minimum of 3000 seeds must be tested for the presence of *High plains virus*, *Maize dwarf mosaic virus*, *Maize chlorotic mottle virus* and *Sugarcane mosaic virus* and a minimum of 400 seeds must be tested for the presence of *Acidovorax avenae subsp. avenae*, *Clavibacter michiganensis subsp. nebraskensis* and *Pantoea stewartii*.

Ministry for Primary Industries Page 155 of 168

(14) All importers who test a composite sample on arrival in New Zealand must make a <u>declaration</u> to identify the place of production or production site of the lots that form the composite sample.

Guidance

- The sample size from each lot to form the composite samples should be calculated as follows:
 - a) The proportion of each lot in the total consignment (seed number) is calculated using the following equation:

 $Proportion \ of \ total \ consignment \ size = \frac{\textit{No.of seeds in each lot}}{\textit{Total number of seeds in consignment}}$

b) Calculate the sample size for each lot (number of seeds) using a total composite sample sizes of 3000 seeds and 400 seeds:

Sample size of each line= 3000 seeds x proportion of total consignment size

Sample size of each line= 400 seeds x proportion of total consignment size

c) For the first composite sample, take the sum of the sample size for each lot to check that the total composite sample for the consignment is at least 3000 seeds. For the second composite sample, take the sum of the sample size for each lot to check that the total composite sample for the consignment is at least 400 seeds.

Ministry for Primary Industries Page 156 of 168

Pest list for $\underline{\textit{Zea}}$ REGULATED PESTS (actionable)

Insect	
Insecta	
Bostrichidae	
Dinoderus distinctus	bostrichid beetle
Dinoderus minutus	bamboo powderpost beetle
Prostephanus truncatus	larger grain borer
Cucujidae	
Cathartus quadricollis	squarenecked grain beetle
Curculionidae	
Caulophilus oryzae	broadnosed grain weevil
Dermestidae	
Attagenus unicolor	black carpet beetle
Trogoderma glabrum	khapra beetle
Trogoderma granarium	khapra beetle
Trogoderma inclusum	trogoderma beetle
Trogoderma variabile	warehouse beetle
Histeridae	
Teretriosoma nigrescens	
Languriidae	
Pharaxonotha kirschil	Mexican grain beetle
Melyridae	
Nitidulidae	
Carpophilus freemani	dried fruit beetle
Carpophilus lugubris	dusky sap beetle
Glischrochilus quadrisignatus	four-spotted sap beetle
Ptinidae	
Gibbium psylloides	shiny spider beetle
Scolytidae	
Pagiocerus frontalis	bark borer
Tenebrionidae	
Alphitobius laevigatus	black fungus beetle
Cynaeus angustus	larger black flour beetle
Gnatocerus maxillosus	slenderhorned flour beetle
Latheticus oryzae	longheaded flour beetle
Palorus ratzeburgi	smalleyed flour beetle
Palorus subdepressus	depressed flour beetle
Tribolium freemani	flour beetle
Diptera	
Otitidae	
Euxesta stigmatias	
Hemiptera	
Coreidae	
Leptoglossus zonatus	coreid bug
Lepidoptera	
Cosmopterigidae	

Ministry for Primary Industries Page 157 of 168

	Pyroderces rileyi	pink scavenger caterpillar	
Noctuidae	,		
	Sesamia calamistis	pink stalk borer	
	Sesamia nonagrioides	pink borer	
Pyralidae			
	Corcyra cephalonica	rice moth	
	Doloessa viridis		
	Mussidia nigrivenella	pyralid moth	
	Paralipsa gularis	stored nut moth	
Tortricidae			
	Cryptophlebia leucotreta	false codling moth	
Psocoptera			
Liposcelidae			
	Liposcelis bostrychophilus	booklouse	
	Liposcelis entomophilus	grain psocid	
	Liposcelis paetus	booklouse	
Trogiidae			
	Lepinotus reticulatus		
Mite			
Arachnida			
Acarina			
Py	vemotidae		
	Acaropsellina sollers		
Fungus			
Ascomycota			
Dothid			
Botryo	sphaeriaceae		
	Botryosphaeria zeae (anamorph macrophoma zeae)	grey ear rot	
Pleosp	oraceae		
	Cochliobolus pallescens (anamorph Curvularia pallescens)	-	
	Cochliobolus tuberculatus (anamorph Curvularia tuberculata)	leaf spot	
	Gloeocercospora sorghi	zonate leaf spot	
Hypocreales		-	
	pitaceae		
	Claviceps gigantea	ergot	
Basidiomyco			
Ustomycetes	5		
Ustilag	inales		
	stilaginaceae		
	Ustilago maydis	boil smut	
Mitosporic Fungi (•		
Sphaerioidales			
Sphaerioidad			
	Stenocarpella macrospora	dry rot of maize	

Ministry for Primary Industries Page 158 of 168

Mitosporic Fungi (Hyphomycetes)	
Hyphomycetales	
Moniliaceae	
Cephalosporium maydis	
Oomycota	
Sclerosporales	
Sclerosporaceae	
Peronosclerospora heteropogoni	
Peronosclerospora maydis	Java downy mildew
Peronosclerospora philippinensis	Philippine downy mildew
Peronosclerospora sacchari	
Peronosclerospora sorghi	sorghum downy mildew
Verrucalvaceae	
Sclerophthora rayssiae var. zeae	
Zygomycota	
Zygomycetes	
Mucorales	
Mucoraceae	
Sclerophthora rayssiae var. zeae	
Stenocarpella macrospora	dry rot
Ustilago maydis	boil smut
Bacterium	
Pseudomonadaceae	
Acidovorax avenae subsp. avenae	bacterial blight
Corynebacteriaceae	
Clavibacter michiganensis subsp. nebraskensis	Goss' bacterial wilt
Enterobacteriaceae	
Pantoea stewartii	Stewart's bacterial wilt
Virus	
High plains virus	
Potyviridae	
Potyvirus	
Maize chlorotic mottle virus	MCMV
Maize dwarf mosaic virus	MDMV
Sugarcane mosaic virus	SCMV
Weed	
Scrophulariales	
Scrophulariaceae	
Striga asiatica	witch-weed
Striga hermonthica	witch-weed

Ministry for Primary Industries Page 159 of 168

Appendix 1: Definitions

Definitions have the same meaning as defined by the Act and ISPM 5: Glossary of Phytosanitary Terms, unless set out below:

a.i.

Active ingredient.

AOSA

The Association of Official Seed Analysts is an organisation comprised of member laboratories which are staffed by certified seed analysts. Such seed testing facilities include official state, federal, and university seed laboratories across the United States of America and Canada.

Basic seed

Refers to seed listed in the Plant Biosecurity Index under "Import Specification for Seed for Sowing".

ONZPR

Biosecurity Organisms Register for Imported Commodities: MPI database which informs on the quarantine status for an organism as either regulated or non- regulated for New Zealand.

Contamination

Presence in a commodity, storage place, conveyance or container, of pests or other regulated articles, not constituting an infestation.

ELISA

Enzyme linked immunosorbent assay.

EPA

Environmental Protection Authority is responsible for administering the Hazardous Substances and New Organisms (HSNO) Act 1996.

Fleshy fruit

Any fruit (matured ovary) that is succulent or semi-succulent e.g. a berry, drupe, pome.

Genetically modified organism (GM)

Any organism in which any of the genes or any of the other genetic material has been modified by in-vitro techniques; or is inherited or otherwise derived, through any number of replications, from any genes or other genetic material which has been modified by *in-vitro* techniques. [as defined by the HSNO Act 1996]

Growing season inspection

Visual inspection by a person authorized by the NPPO during period or periods of the year when plants actively grow in an area, place of production or production site.

Herbaceous Indexing

Virus detection and identification technique where plant viruses are transmitted mechanically or via a vector to a number of herbaceous indicator plants for the observation of characteristic symptoms.

ISTA

International Seed Testing Association.

IPPC

International Plant Protection Convention, as deposited with FAO in Rome in 1951 and as subsequently amended [FAO, 1990].

Ministry for Primary Industries Page 160 of 168

Isolation

Applies to PEQ facilities which must meet the minimum isolation requirements (from plants outside the PEQ facility) listed in this IHS.

ISPM

International Standard for Phytosanitary Measures are the international standards adopted by the Conference of FAO, the Interim Commission on Phytosanitary Measures or the Commission on Phytosanitary Measures, established under the IPPC [CEPM, 1996; revised CEPM, 1999].

Level 1, Level 2 or Level 3 post-entry quarantine

A system of post entry quarantine screening whereby seed is grown under certain specified conditions on a property approved to the Facility Standard: Post Entry Quarantine for Plants (MPI.STD.PEQ).

MPL

Maximum Pest Limit.

NPPO

National Plant Protection Organisation is the official service established by Government to discharge the functions specified by the IPPC. [FAO, 1990; formerly Plant Protection Organisation (National)].

Officially tested

Tested by a laboratory approved by the exporting country NPPO if performed offshore or by the importing country NPPO if performed on-shore

PCR

Polymerase chain reaction

Pelleted seed

Seed encased in a man-made nutritive or protective covering.

PEQ

Post Entry Quarantine. Quarantine applied to a consignment after entry.

Permit

A permit to import issued by MPI that specifies the conditions under which a particular commodity may be imported into New Zealand.

Pest

Any species, strain or biotype of animal or pathogenic agent (fungi, bacteria, viruses, viroids) injurious to plants or plant products.

Note: For the purpose of this import health standard "pest" includes an organism sometimes associated with the pathway, which poses a risk to human or animal or plant life or health (SPS Article 2).

Plant Biosecurity Index

MPI search system for identifying the status of plant species for importing to New Zealand.

Pre-Germinated Seed

Seed with only the radicle (embryonic root) emerged.

Quarantine Pest

A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled [FAO, 1990; revised FAO 1995; IPPC 1997].

Quarantine Weed Seeds

An invasive plant species as set out in the MPI Schedule of Regulated (Quarantine) Weed Seeds.

Ministry for Primary Industries Page 161 of 168

Regulated Pest

A quarantine pest or a regulated non- quarantine pest listed in ONZPR as being regulated for New Zealand. Note: If an intercepted organism is not listed in ONZPR, the NPPO must contact MPI to establish the regulatory status.

Representative sample


A sample that is to be submitted to the approved testing laboratory and may comprise either the whole of the composite seed sample or a subsample thereof.

SAC

Seed Analysis Certificate

Seed

A unit of reproduction used for sowing. This includes spores but excludes vegetative propagules.



Appendix 2: Amendment Record

Amendments to this IHS will be given a consecutive number and dated. The following table provides a summary of the main changes to this IHS for the previous five years.

No:	Details:	Date:
26	Revised schedules of special conditions for <i>Hordeum</i> and <i>Triticum</i> .	7/05/2009
27	Addition of schedule for <i>Linum usitatissimum</i> . Revised schedule of special conditions for <i>Fragaria</i> and <i>Ribes</i> . Removal of <i>Echinacea angustifolia</i> from section 1.5.2	19/03/2010
28	Removal of <i>Xanthomonas translucens</i> pv. <i>translucens</i> from the <i>Hordeum</i> and <i>Triticum</i> schedules. Revised schedule of <i>Zea</i> , including Japan as an approved country with the addition of <i>Gloeocercospora sorghi</i> to the pest list. Addition of a pea seed soak test on arrival in the Pisum schedule.	22/09/2010
29	Revised schedules of special conditions for Acer, Carpinus, Carya ovata, Castanea and Quercus to manage Cryphonectria parasitica.	16/09/2011
30	Addition of section 2.2.7 'Importation of Seed Products', section 2.2.8 "Seed for Sowing of New Zealand Origin' and section 2.4 'Equivalence'.	5/12/2011
31	Revised schedule for <i>Rubus</i> , and removal of [strains not present in New Zealand] from all listings of <i>Tomato ringspot virus</i> in <i>Fragaria, Rubus</i> , and <i>Vaccinium</i> schedules.	20/03/2012
32	Revised schedule of special conditions for Citrus.	3/04/2012
33	Updated fungicide treatment option for Avena, Hordeum and Triticum.	7/05/2012
34	Reformat of complete IHS, including all schedules.	29/06/2012
35	Correction to the <i>Zea</i> schedule: removal of <i>Maize mottle chlorotic stunt virus</i> from the quarantine requirements.	24/07/2012
36	Revision of Section 8 'Equivalence' and Section 9 "Biosecurity clearance'.	27/08/2012
37	New schedule for tomato (Solanum lycopersicum) and minor correction of Macadamia schedule.	19/10/2012
38	New schedule for <i>Brassica</i> (urgent amendment) and minor amendment to <i>Acrocomia</i> schedule.	19/04/2013
39	Revised schedule for Malus (apple) seed for sowing.	24/04/2013
40	Removal of schedule for Brassica, retaining schedule for Brassica napus.	02/08/2013
41	New schedule for grape (Vitis spp.), incorporated as an urgent amendment.	08/08/2013
42	Addition of section 6 (Part A), 'Seed for sowing imported as laboratory specimens'. Revised schedule of special conditions for <i>Arabidopsis thaliana</i> (removal of requirement for a phytosanitary certificate). Revised schedules of special conditions for <i>Fragaria</i> , <i>Phaseolus</i> , <i>Ribes</i> , <i>Rubus idaeus</i> and <i>Vitis</i> (removal of <i>Tomato black ring virus</i> from the quarantine requirements).	12/12/2013
43	Addition of further approved fungicide treatments to the <i>Phaseolus</i> and <i>Pisum</i> schedules	19/6/2014
44	Revised schedule for <i>Zea mays</i> , specifying a seed sample size for <i>Maize dwarf mosaic virus</i> .	18/8/2014
45	New schedule for Capsicum and Solanum	19/8/2014

Ministry for Primary Industries Page 163 of 168

No:	Details:	Date:
46	New schedule for Cucurbitaceae and changes to Zea mays (urgent amendment)	1/12/2014
47	Revised schedule for Zea mays, clarifying the requirements for Sugarcane mosaic virus and Maize chlorotic mottle virus	7/8/2015
48	Publication of the CTO direction for all <i>Zea mays</i> consignment originated from Chile.	11/09/2015
49	New IHS format. Added section 1.6 (pre-determined testing in PEQ), amended phytosanitary certificate requirements and GM testing requirements.	26/11/2015
50	Reinstating section 6 (Part A) now section 1.9 Part 1: "Seed for sowing imported as laboratory specimens". Minor amendment for <i>Beta</i> and <i>Zea</i> schedule. Removal of <i>Barley mosaic virus</i> from the pest list of <i>Hordeum</i> and revised the schedule.	21/12/2015
51	Reinstating and revision of the requirements for species of Rubus and clarification of Section 1.9	21/01/2016
52	Addition to a paragraph related to importation of GMO seeds for reseach purposes and also the addition of the pathogen Andean potato mild mosaic virus (APMMV) to the Solanum tuberosum schedule as a regulated pest.	02/03/2016
53	Revised the Capsicum schedule: addition of PCFVd as a quarantine pest	09/10/2016
54	Orthographic corrections under Zea, Triticum and Lavandula schedules and amendment to the Zea schedule to allow for testing onshore for all quanrantine pests listed in the Zea mays pest list, reformatting of Appendix 3: Declaration form to facilitate its use and the addition of a hyperlink to the protocol for GMO testing under the Zea schedule.	25/11/2016
55	Review of the Cucurbitaceae schedule: new measures for CGMMV and addition of KGMMV as a quarantine pest. Update of the Capsicum schedule.	26/01/2017
56	Removal of <i>Clover yellow mosaic virus</i> and <i>Red clover vein mosaic virus</i> . Addition of onshore testing for the Agropyron schedule. Update name of PEQ standard, update references to Level 3 PEQ to reflect requirements of the reissued PEQ standard.	09/03/2017
57	Addition of Apiaceae and Petunia schedules and other minor changes.	09/06/2017
58	Change to the treatment requirements for the Apiaceae schedule and other minor changes.	14/12/2017
59	Removal of pea soak test, addition of compulsory fumigation for pea seeds and migration of all approved treatments to MPI-STD-ABTRT.	22/02/2018
60	Addition of Myrtaceae Specific Requirements	11/07/2018
61	Addition of onshore testing for Grapevine yellow speckle viroid-2 to the specific requirements for Vitis.	11/01/2019
62	Addition of requirements for Tomato brown rugose fruit virus (TBRFV) to specific requirements for Capsicum and Solanum lycopersicum.	19/03/2019
63	Review of the standard to remove errors and clarify ambiguities and current requirements as part of the tidying the room project.	02/09/2019
64	Apiaceae and Cannabis sativa specific requirements: Addition of onshore hot water treatment as a phytosanitary option; Capsicum and Solanum lycopersicum specific requirements: addition of molecular testing as an offshore phytosanitary measure for Tomato brown rugose fruit virus and inclusion of guidance	28/04/2020

Ministry for Primary Industries Page 164 of 168

No:	Details:	Date:
	information that the ISHI-Veg local lesion bioassay is not accepted by MPI as a valid test for <i>Tomato brown rugose fruit virus</i> ; <i>Petunia</i> specific requirements: addition of option for importers to provide a non-GMO declaration to meet the genetically modified (GM) requirements for <i>Petunia</i> seeds for sowing and removal of requirement for 'appropriate common name' to be specified on phytosanitary certificates for <i>Petunia</i> seeds for sowing.	
65	Addition of requirements for pelleted seeds for sowing in Section 1.9 of the IHS, and inclusion of a requirement for all importers of seed for sowing to make a declaration whether their consignment contains pelleted seed or not in Section 1.4 of the IHS.	16/06/2020
66.	Removal of specific requirements for the Apiaceae schedule.	16/06/2020
67	Addition of additional declaration requirements for <i>Tomato mottle mosaic virus</i> in the specific requierments for <i>Capsicum</i> spp. and <i>Solanum lycopersicum</i> .	22/07/2020
68	Removal of requirement for phytosanitary certificates for 'basic' pelleted seeds within part 1.9.	22/07/2020
69	Amendment to the <i>Petunia</i> specific requirements with addition of measures for <i>Tomato chlorotic dwarf viroid</i> and to the <i>Solanum lycopersicum</i> specific requirements with addition of measures for <i>Columnea latent viroid</i> , <i>Tomato apical stunt viroid</i> and <i>Tomato planta macho viroid</i> . Harmonization of measures for <i>Potato spindle tuber viroid</i> and <i>Tomato chlorotic dwarf viroid</i> in the <i>Solanum lycopersicum</i> specific requirements.	22/07/2020
70	Amendment to the specific requirements for <i>Capsicum</i> and <i>Solanum Iycopersicum</i> to include a suspension notice for import pathways from Israel.	17/12/2020
71	Amendment to the specific requirements for <i>Capsicum</i> and <i>Solanum Iycopersicum</i> to remove notices of suspension for import pathways from Israel.	15/02/2021
72	Amendment to section 1.8 Seeds for sowing imported as laboratory specimens to remove contents of this section and include guidance directing to import requirements under the Research Samples (excluding animal samples) import health standard.	21/06/2021
73	Amendment of 2.72.1(1)(a)(i), 2.82.1(1)(a)(ii) Correction of formatting, terminology and grammar in Appendix 3 & 4, paragraph 2 under "Who should read this IHS?", Parts 2.42.1(1)(a), 2.73, 2.80, 2.70, 2.14, 2.74 pest list, 2.44.2, 2.59.2, 2.27.3, 2.1.2(1), 2.42.1 (1)a, 2.68 Isolation, 2.17 pest list, 2.80 (2), 2.64, 1.9 (2)(10), 2.61 pest list and in terms used throughout the document. Addition of guidance to 2.4 Actinidia, 2.15 Carpinus, 2.18 Castanea, 2.31 Eriobotrya, 2.33 Fragaria, 2.38 Humulus, 2.39 Juglans, 2.44 Lithocarpus densiflorus, 2.46 Lophophora williamsii, 2.49 Malus, 2.50 Mangifera, 2.50 Persea, 2.63 Populus, 2.64 Prunus, 2.67 Pyrus, 2.68 Quercus, 2.69 Ribes, 2.70 Rubus, 2.74 Solanum tuberosum and 2.76 Stenotaphrum. Correction of references to the 'seeds imported as laboratory specimens' section to direct to the research samples IHS in Part 1, Section 1.5.1(1)(b) and 1.5.1(2)	12/08/2021
74	Removal of the following pests from pest lists and measures for these pests: Hemp streak virus and Hemp mosaic virus in the specific requirements of Cannabis sativa, Artichoke yellow ringspot virus in the specific requirements of Phaseolus, Pea enation mosaic virus in the specific requirements of Pisum and Vicia, Peanut stunt virus in the specific requirements of Phaseolus and Vicia; Removal of ELISA as a testing option for Tomato brown rugose fruit virus in the	20/09/2021

Ministry for Primary Industries Page 165 of 168

No:	Details:	Date:
	specific requirements of Capsicum and Solanum lycopersicum; Addition of measures for Potato spindle tuber viroid in the specific requirements for Glebionis (new schedule) and Petunia.	
75	Addition of Cucumis schedule for measures for Cucumis melo and Citrullus lanatus for Melon necrotic spot virus and removal of Cucumis melo and Citrullus lanatus from Cucurbitaceae schedule.	5/11/2021
76	Amendment to the testing requirements for <i>Pepino mosaic virus</i> in the specific requirements for <i>Solanum lycopersicum</i>	21/02/2022



Ministry for Primary Industries Page 166 of 168

Appendix 3: Declaration Form

(Exporter's name and address)...

To be completed and signed by the exporter and importer.

As defined by the New Zealand HSNO Act 1996, Genetically modified organism means, unless expressly provided otherwise by regulations, any organism in which any of the genes or any other genetic material (a) have been modified by in vitro techniques; or (b) are inherited or otherwise derived, through any number of replications, from any genes or other genetic material which has been modified by in vitro techniques.

Note that under the Hazardous Substances and New Organisms (HSNO) Act 1996, the import and release of any genetically modified crop without approval from the Environmental Protection Authority (EPA) is unlawful.

declare that according to the requirements set out in the Standard: 155.02.05: Seeds for Sowing - https://www.mpi.g	Seed for Sowing Import Health Standard (MPI Import Health ovt.nz/document-vault/1151),		
Insert species name and lot/line number or unique ic	lentifier as stated on all the other import documentation		
was produced neither "from" nor "by" genetically modified co	rops.		
I undertake to inform immediately the importer and the Ministry for Primary Industries, MPI, New Zealand of any information that can undermine the accuracy of this declaration.			
Note that MPI may request evidence as to how production, handling and transport of these seeds is performed in the field, or require and audit as a way to provide quality to the production system.			
I (Importer's name and address)			
declare to the best of my knowledge that according to the re Standard (MPI Import Health Standard: 155.02.05: Seeds for			
Insert species name and lot/line number or unique identifier as stated on all the other import documentation			
was produced neither "from" nor "by" genetically modified on	rops.		
Signed by Exporter and Company Name (details) and date	Signed by Importer and Company Name (details) and date		

Warning: Any person who knowingly makes a statement of information or a declaration that is false or misleading in a material particular may on summary conviction, be sentenced to a term of imprisonment and/or fined not exceeding \$500,000.00.

Ministry for Primary Industries Page 167 of 168

Appendix 4: Species on the Plant Biosecurity Index eligible for import into New Zealand as pelleted seeds for sowing

Ageratum houstonianum	Solanum lycopersicum
Allium cepa	Spinacia oleracea
Allium porrum	Lisianthus russellianus (= Eustoma grandiflorum)
Anethum graveolens	Lobelia sp.
Angelonia salicariifolia	Lobularia maritima
Antirrhinum sp.	Matthiola incana
Apium graveolens	Mimulus sp.
Begonia sp.	Nemesia sp.
Bellis perennis	Nicotiana sp.
Beta vulgaris	Ocimum basilicum
Brassica napus	Origanum vulgare
Brassica oleracea	Papaver sp.
Calceolaria sp.	Pentas sp.
Calibrachoa hybrida	Pericallis hybrida (= Pericallis x hybrida)
Campanula sp.	Petroselinum crispum
Celosia sp.	Petunia sp.
Chaenorhinum sp.	Portulaca sp.
Chrysanthemum sp	Primula sp.
Cichorium endivia	Pyrethrum sp.
Cichorium intybus	Ranunculus sp.
Cineraria maritima (= Senecio cineraria)	Rosmarinus officinalis
Cyperus papyrus	Rudbeckia sp.
Daucus carota	Salpiglossis sinuata
Dianthus sp.	Salvia officinalis
Diascia barberae	Saxifraga sp.
Dichondra sp.	Senecio cruentus (= Pericallis cruenta)
Digitalis sp.	Silene sp.
Eruca sativa	Solenostemon scutellarioides
Exacum affine	Streptocarpus sp.
Foeniculum vulgare	Sutera sp.
Gazania sp.	Tagetes sp.
Geranium sp.	Tanacetum parthenium
Gerbera jamesoni	Thymus vulgaris
Gloxinia speciosa (= Sinningia speciosa)	Torenia fournieri
Gypsophila sp.	Trachelium caeruleum
Helichrysum sp.	Verbascum sp.
Heuchera sp.	Verbena sp.
Isolepis sp.	Veronica sp.
Juncus sp.	Viola sp.
Lactuca sativa	Zinnia sp.
Laurentia axillaris (= Isotoma axillaris)	· ·
Linaria sp.	
Pastinaca sativa	

Ministry for Primary Industries Page 168 of 168