

# CHECKLIST OF INSECT AND MITE SPECIES ATTACKING CONES AND SEEDS OF WORLD CONIFERS

Harry O. Yates III  
Research Entomologist  
Southeastern Forest Experiment Station  
Athens, GA 30602  
USDA Forest Service  
(Accepted for publication May 5, 1986)

## ABSTRACT

Listed alphabetically by order, family, genus, and species are 360 insect and mite species that attack flowers, cones, and seeds of world conifers. Synonyms, misspellings, invalid combinations, erroneous author names, and abbreviations of author names are listed exactly as published following the currently accepted species binomen. All species names listed are referenced to the cone and seed insect literature where that terminology appears.

**Key Words:** Cone and seed insects and mites, world checklist, Coleoptera, Diptera, Heteroptera, Homoptera, Hymenoptera, Lepidoptera, Thysanoptera, Acari.

---

J. Entomol. Sci. 21(2): 142-168 (April 1986)

## INTRODUCTION

Insects and mites that attack and destroy the cones and seeds of conifers throughout the world have received increased attention in recent years. This interest parallels the growing emphasis on production of large quantities of viable seed for use in tree regeneration programs. The emphasis on production and protection of large quantities of genetically improved seed in seed orchards also has heightened this interest.

In 1983, the Cone and Seed Insects Working Party (S2.07-01) of the International Union of Forestry Research Organisations (IUFRO) solicited workers throughout the world to develop a list of all insect and mite species that attack, damage, or kill the flowers, cones, or seeds of conifers. Simultaneously, the literature was reviewed and a master list of all species was produced. These lists were to form a basis for developing a publication on the world cone and seed insects and mites of conifers similar to those produced for North America [23] and France [48].

It became apparent that these lists were in need of considerable taxonomic editing in order to develop a master list that reflects the current state of taxonomic nomenclature. With the assistance of taxonomists from the Insect Identification and Beneficial Insect Introduction Institute of the USDA in Washington, D.C., the University of Minnesota, and North Carolina State University, a revised list reflecting the current taxonomy of the groups was developed and all species from the original lists were incorporated under this revised scheme. This included synonyms, invalid combinations, misspellings, erroneous authorships, and various abbreviations for authors' names. In some cases these author abbreviations created major problems in determining the species being referenced. To assist the nontaxonomist I include, exactly as they appear in print, all such abbreviations in the list below each species. With the exception of "L." for Linnaeus and "F." for

Fabricius, all author names are spelled out in the main entries. Species within the class Insecta are listed first, followed by species within the class Arachnida. Within each class the orders, families, and species names are listed alphabetically. The sources of inclusions in this checklist are in brackets [ ] by reference number; those not referenced are from personal correspondence.

No attempt was made to include all the world cone and seed insect references, but rather I tried to include the important monographs on the subject or other significant references that represent major listings or additions to insect and mite species attacking conifer cones and seeds.

## CLASS INSECTA

### COLEOPTERA

#### Anobiidae

*Ernobius abietinus* (Gyllenhal)

E. abietinus Gyll. [48,83]

*Ernobius abietis* (F.)

E. abietis F. [10,11,48,73]

E. abietis L. [74,77]

E. abietis Fabr. [83]

*Anobium abietes* Fb. [6]

*Ernobius alutaceus* (LeConte) [27]

*Ernobius anatolicus* Johnson

E. anatolicus Johns. [48]

*Ernobius angusticollis* (Ratzeburg)

E. angusticollis Ratzbg. [10,11]

E. angusticollis Ratz. [48]

E. parvicollis Muls. [48]

E. tabidus Kiesw. [48]

E. tabidus Kzw. [65,74]

E. tabidus Ksw. [72,73,77]

*Ernobius bicolor* White [82]

*Ernobius conicola* Fisher [23]

E. cupressi VanDyke [26]

E. cupressi Van Dyke [23]

*Ernobius cupressi* Chobaut

E. cupressi Chob. [48]

*Ernobius fructuum* Peyerimhoff

E. fructuum Peyer [48]

*Ernobius granulatus* LeConte [23]

E. granulatus Lec. [39]

*Ernobius kailidis* Johnson [31]

*Ernobius longicornis* (Sturm)

E. longicornis Sturm. [48]

E. densicornis Muls. and Rey [48]

*Ernobius melanoventris* Ruckes [23]

*Ernobius mollis* (L.)*E. mollis* Linné [88]*E. mollis* L. [38,48]*E. mollis* LeConte [23]*E. mollis* F. [48]*Ernobius montanus* Fall [23,26]*Ernobius nigrans* Fall [23,54]*Ernobius oertzeni* Schilsky*E. oertzenii* Schils. [48]*Ernobius oxycedri* Reitter*E. oxycedri* Reitt. [48]*Ernobius pallitarsis* Fall [23,26]*Ernobius parens* (Mulsant and Rey)*E. parens* Muls. [48]*Ernobius pini* (Sturm)*E. pini* Sturm. [48,50]*Ernobius pini* var. *crassiusculus* Mulsant and Rey*E. pini* var. *crassiusculus* Muls. [11,48]*E. pini* Strm. var. *crassiusculus* Muls. [10]*Ernobius pinicola* Ruckes [23]*Ernobius punctulatus* (LeConte) [21,23]*E. punctulatus* Leconte [48,53]*E. punctulatus* (Lec.) [26]

## Apionidae

*Nanodiscus transversus* (Aubé)*N. transversus* Aubé [48,51]

## Buprestidae

*Chrysobothris cupressicona* Barr and Wescott [23]

## Curculionidae

*Anthonomus varians* (Paykull)*A. varians* Payk. [76]*Conotrachelus neomexicanus* Fall [23]*Pissodes validirostris* Gyllenhal*P. validirostris* Gyll. [5,10,11,12,20,46,47,48,50,65,73,83,88]

## Meloidae

*Lytta polita* Say [23]*Pomphopea polita* (Say) [39]

## Nemonychidae

*Neocimberis elongatus* (LeConte)*Cimberis elongatus* (LeConte) [23]*Neocimberis pilosus* (LeConte)*Cimberis pilosus* (LeConte) [23]

## Ptinidae

*Ptinus fur* (L.)

*P. fur.* L. [38,48]

*P. fur* [37]

## Scarabaeidae

*Dichelonyx backi* (Kirby)

*D. backi* Kirby [23]

*Phyllophaga micans* (Knoch) [15,16,23]

## Scolytidae

*Conophthorus apachecae* Hopkins [23,86]

*C. apachecae* Hopk. [26]

*Conophthorus cembroides* Wood [23,86]

*Conophthorus conicolens* Wood [23,86]

*Conophthorus coniperda* (Schwarz) [15,16,23,84,86]

*C. clunicus* Hopkins [23,86]

*C. taedae* Hopkins [23,86]

*Conophthorus echinatae* Wood [23,86]

*Conophthorus edulis* Hopkins [23,86]

*C. edulis* Hopk. [26]

*Conophthorus mexicanus* Wood [23,86]

*Conophthorus michoacanae* Wood [86]

*Conophthorus michoacanae* Wood [85]

*Conophthorus monophyllae* Hopkins [23,86]

*C. monophyllae* Hopk. [26]

*Conophthorus ponderosae* Hopkins [23,27,86]

*C. ponderosae* Hopk. [26]

*C. contortae* Hopkins [23,86]

*C. contortae* Hopk. [26]

*C. flexilis* Hopkins [23,86]

*C. flexilis* Hopk. [26]

*C. lambertiana* Hopkins [23,86]

*C. lambertiana* Hopk. [26]

*C. monticolae* Hopkins [21,23,86]

*C. monticolae* Hopk. [26]

*C. scopulorum* Hopkins [23,86]

*C. scopulorum* Hopk. [26]

*Conophthorus radiatae* Hopkins [23,86]

*C. radiatae* Hopk. [26]

*Conophthorus resinosae* Hopkins [23,86]

*C. resinosae* Hopk. [43,84]

*C. resinosa* Hopk. [44]

*C. virginianae* Hopkins [23,86]

*Conophthorus teocotum* Wood [85,86]

*Pityophthorus aztecus* Bright [23]

*Pityophthorus orarius* Bright [21,53]

*Pityophthorus pubescens* (Marsham)

*Pityophthorus pubescens* Marsh. [50]

*Pityophthorus pulicarius* (Zimmermann) [23]

*P. pulicarius* (Zimm.) [14,69]

*P. pulicarius* Zimm. [39]

*P. pulicarius* Zinn. [39]

*Pityophthorus schwerdtfergeri* (Schedl)

## DIPTERA

### Anthomyiidae

*Lasiomma abietis* (Huckett)<sup>1</sup> [23]

*L. abietis* Huckett [78]

*L. abietis* Huck. [48,60,73]

*L. abietum* Huck. [48]

*Hylemya abietis* Huckett [23]

*H. (Lasiomma) abietis* Huckett [23]

*H. (Pegohylemyia) abietis* Huckett [26]

*Hylemiaia abietis* Huck. [42]

*Lasiomma anthracinum* (Czerny) [18]

*L. anthracina* (Czerny) [23,78]

*L. anthracina* Czerny [48,72]

*L. anthracina* Cerny [73]

*L. anthracinum* Czerny [42]

*L. anthracium* (Gerny) [88]

*Hylemya anthracina* (Czerny) [23,54]

*H. anthracina* Czerny [21]

*H. (Lasiomma) anthracina* (Czerny) [23]

*H. (Pegohylemyia) anthracina* (Czerny) [26]

*Pegohylemyia anthracina* Czerny [42,65,74,83]

*P. anthrazina* Czerny [77]

*P. anthracini* (Czerny) [7]

*Pegohylemia anthracina* Czerny [84]

*Hylemyia anthracina* Czerny [61]

*Lasiomma baicalense* Elberg [18,42,75,88]

*L. baicalensis* Elberg [48,49]

*Lasiomma carbonarium* (Ringdahl)

*Hylemya (Lasiomma) carbonarium* (Ringdahl) [23]

*Lasiomma infrequens* Ackland [18,42,75]

*L. infrequens* Ackl. [48,49]

*L. infreguens* Ackland [88]

*Lasiomma jurtschenkoi* Elberg [48,49,73]

*L. jurtschenkovi* Elberg [75]

*Lasiomma laricicola* (Karl)<sup>2</sup> [18,88]

*L. laricicola* Karl [42,48,49,65,73,75]

*Hylemyia laricicola* Karl [42,74,77,83]

<sup>1</sup> Records of *L. abietis* from Europe probably refer to a different species. Personal communication, F. C. Thompson, ARS Systematic Entomology Laboratory, USDA, Washington, DC.

<sup>2</sup> North American *Larix* feeding species now called *L. laricicola* (Karl) is a new species. Personal communication, F. C. Thompson, ARS Systematic Entomology Laboratory, USDA, Washington, DC.

- H. laricicola* (Karl) [30,57,59]  
*Hylemya laricicola* (Karl) [28]  
*H. (Lasiomma) laricicola* (Karl) [23]  
*Chortophila laricicola* Karl [42,49,59,90]  
*Phorbia laricicola* (Karl) [63]  
*Lasiomma luteoforceps* Fan et Fang [18]  
*Lasiomma melania* Ackland [42,71,72,73]  
  *L. melania* Ackl. [48,49]  
  *L. melania* Acland [75]  
*Lasiomma melania melaniola* Fan [18,88]  
*Lasiomma oriens* Suwa<sup>3</sup>  
*Lasiomma todocola* Suwa<sup>3</sup>

#### Cecidomyiidae

- Asynapta hopkinsi* Felt [23,78]  
  *A. hopkinsi* (Felt) [43,44]  
  *A. keeni* (Foote) [17,23,27]  
  *A. keeni* Foote [39]  
*Rubaamenia keeni* Foote [23,26,78]  
*Ruebsaamenia keeni* Foote [78]  
*Asynapta laricis* Skrzypczynska [64]  
  *A. laricis* Skrzypcz. [59]  
  *A. laricis* Skrz. [48]  
*Asynapta strobilophila* (Foote) [23]  
  *Holoneurus strobilophilus* Foote [26,78]  
*Camptomyia pinicola* Mamaev  
  *C. pinicolana* Mam. [48,50]  
*Camptomyia pseudotsugae* Hedlin and Johnson [21,23]  
*Cecidomyia bisetosa* Gagné [16,23]  
*Cecidomyia pini* DeGeer  
  *C. pini* Deg. [48]  
  *C. pini* (DeGeer) [63]  
*Contarinia oregonensis* Foote [21,23,26,48,53,78]  
*Contarinia washingtonensis* Johnson [21,23,53,78]  
*Dasineura abiesemias* Foote [21,23,78]  
  *Dasyneura abiesemias* Foote [26]  
*Dasineura canadensis* Felt [21,23,54,78]  
  *Dasyneura canadensis* Felt [26]  
*Dasineura laricis* (F. Loew) [64]  
  *Dasyneura laricis* [48]  
*Dasineura rachiphaga* Trip [21,23,54,78]  
  *Dasyneura rachiphaga* Tripp [26]  
*Janetiella siskiyou* Felt [13,26,48,78]  
  *Craneiobia lawsonianae* DeMeijere [13]  
*Kaltenbachiola strobi* (Winnertz) [64,83]  
  *K. strobi* (Winn.) [7,61]  
  *K. strobi* Winn. [46,48,65,72,73,77]

---

<sup>3</sup> Personal communication, Verner Michelson, Zoologisk Museum, Kobenhavn, Denmark.

- Cecidomyia strobi* Winnertz [7]  
*Kaltenbachiella strobi* (Winnertz) [6]  
*K. strobi* (Winn.) [6]  
*Kalt. strobi* (Winn.) [6]  
*Kaltenbrachiola strobi* Winn. [74]  
*Kaitenbachiola strobi* Winn. [77]  
*Kaltembakiola strobi* Winn. [38]  
*Mayetiola carpophaga* (Tripp) [21,23,54,78]  
*Phytophaga carpophaga* Tripp [26]  
*Mayetiola thujae* (Hedlin) [23,78]  
*Phytophaga thujae* Hedlin [21,23]  
*Mycodiplosis pucciniae* Kieffer [77]  
*Mycodiplosis thoracica* (Fitch) [17,23]  
*Plemeliella abietina* Seitner [64,73,74,77]  
*P. abietina* Seitn. [6,7,48,61,72,83]  
*P. abietina* Stein [48]  
*Resseliella ingrlica* (Mamaev)  
*Thomasiniana ingrlica* Mam. [74]  
*T. ingrlica* Mamajev [61,72,73]  
*Thomasinianna ingrlica* Mam. [48]  
*Resseliella piceae* Seitner [38,64,73]  
*R. piceae* [37]  
*R. piceae* Seit. [60]  
*R. piceae* Seitn. [62]  
*Reseliella piceae* Seitn. [48]  
*Resseliella sibirica* (Mamaev)  
*Thomasiniana sibirica* Mamaev [75]  
*T. sibirica* Mamajev [71,73]  
*Thomasinianna sibirica* Mam. [48]  
*Camptomyia laricis* Mam. [48]  
*C. laricis* M. [73]  
*C. laricis* Mar. [65]  
*Resseliella silvana* (Felt) [17,23,43,44]  
*Mycodiplosis silvana* Felt [78]  
*Resseliella skuhravyorum* Skrzypczynska [64]  
*R. skuhravyorum* Skrzyp. [75]  
*R. skuhravyorum* Skrzypcz. [30,57,59,63]  
*Reseliella skuhravyorum* Skrz. [48]  
*Sequoioomyia taxodii* (Felt) [23]  
*Retinodiplosis taxodii* Felt [23,78]  
*Taxodiomyia cupressi* (Schweinitz) [23]<sup>4</sup>

#### Chloropidae

- Hapleginella conicola* (Greene) [23,27,78]  
*Oscinella conicola* Greene [23]  
*O. conicola* (Greene) [26,39]  
*Madiza conicola* Greene [26]

<sup>4</sup> This is a gall forming species that was incorrectly cited. Personal communication, R. J. Gagné, ARS Systematic Entomology Laboratory, USDA, Washington, DC.

*Hapleginella laevifrons* (Loew)<sup>5</sup> [63]

*H. laevifrons* Lw. [20,48,50,75]

*H. laevifrons* Loew [10,11,30]

*H. loevifrons* Lw. [48]

*Haploginella laevifrons* Lw. [77]

*Oscinus laevifrons* Loew [78]

### Lonchaeidae

*Earomyia abietum* McAlpine [21,23,26,32,78]

*Earomyia aquilonia* McAlpine [21,23,26,32,78]

*Earomyia barbara* McAlpine [21,23,26,32,53,78]

*Earomyia bazini* (Seguy) [30,57,59,63]

*Earomyia brevistylata* McAlpine [23,26,32,78]

*Earomyia grusia* Morge [60,73]

*E. grusia* [48]

*E. gruzia* Morge [38]

*E. gruzia* [37]

*Loncheae fugax* Beck [73]

*Earomyia impossibile* Morge [38,60,62]

*E. impossibile* [37]

*E. impossibilis* Morge [48,73]

*Earomyia lonchaeoides* Zetterstedt

*E. lonchaeoides* Zett. [76]

*Earomyia longistylata* McAlpine [23,26,32,78]

*Earomyia schistopyga* Collin [61,73,75]

*E. schistopyga* Coelin [71]

*Earomyia viridana* (Meigen)<sup>6</sup> [60,63,78]

*E. viridana* (Meig.) [57]

*Spermatolonchea viridana* Meig. [48]

*Lonchaea polita* Say [23,78]

### Trypetidae

*Rhagoletis flavigenualis* Hering [10,11,48]

## HETEROPTERA

### Acanthosomatidae

*Cyphostethus tristriatus* (F.)

*C. tristriatus* F. [48]

### Coreidae

*Gonocerus juniperi* (Herrich-Schäffer)

*G. juniperi* H. Sch. [48]

*Leptoglossus corculus* (Say) [15,16,23,44]

*L. corculus* Say [48]

---

<sup>5</sup> Two references indicate direct cone feeding [10,11] while remainder list it as a secondary insect found in dead or dying cones.

<sup>6</sup> At least one reference lists this as an insect predator [48].

*Leptoglossus occidentalis* Heidemann [21,23,53,54]

*L. occidentalis* Heid. [48]

*Palomena prasina* (L.)

*P. prasina* L. [65]

#### Lygaeidae

*Eremocoris ferus* (Say) [79]

*Eremocoris borealis* (Dallas) [79]

*Gastrodes abietum* Bergroth [48]

*G. abietum* Bergr. [48]

*G. abietis* Bergr. [48,77]

*G. abietis* Linn. [4]

*G. abietis* Deg. [65]

*G. abietis* L. [83]

*Gastrodes grossipes* DeGeer [48,77]

*G. grossipes* deGeer [48]

*G. ferrugineus* L. [48,83]

*G. ferrugineus* Linn. [4]

*Orsillus depressus* Dallas

*O. depressus* Dl. [48]

*Orsillus maculatus* (Fieber)

*C. maculatus* Fb. [48]

#### Miridae

*Platyllygus luridus* (Reuter) [43,45]

*P. tinctus* Uhler [44]

#### Pentatomidae

*Chlorochroa juniperina* (L.)

*Pitedia juniperina* L. [48]

*P. (Chlorochroa) juniperina* L. [25]

*Chlorochroa pinicola* (Mulsant and Rey)

*Pitedia pinicola* Muls. and Rey. [48]

*P. pinicola* Mls. [65]

*Dolycoris baccarum* (L.)

*Dolycorus baccarum* L. [65]

*Halyomorpha mista* Uhler [29]

*Plautia stali* Scott [29,88]

*P. stali* [28]

#### Scutellaridae

*Poecilocoris lewisi* (Distant) [28]

*P. lewisi* Distant [88]

*Tetyra bipunctata* (Herrich-Schäffer) [23,44]

*T. bipunctata* (H.-S.) [15,16]

*T. bipunctata* H.S. [48]

*T. bipunctata* (Herrich-Schaeffer) [43]

## HOMOPTERA

## Phylloxeridae

- Adelges cooleyi* (Gillette) [23,53,54]
- Adelges lariciatus* (Patch) [23]
- Adelges laricis* Vallot [23]
  - A. strobilobius* (Kaltenbach) [23]
- Adelges piceae* (Ratzeburg) [23]

## HYMENOPTERA

## Diprionidae

- Augomonoctenus libocedrii* Rohwer [23,26]
- Augomonoctenus smithi* (Xiao and Wu) [87]
- Neodiprion pratti pratti* (Dyar) [15,16,23]

## Eurytomidae

- Eurytoma bouceki* Skrzypczynska
  - E. bouceki* Skrz. [48]
  - E. bouceki* Skrzyp. [75]
  - E. bouceki* Skrzypcz. [30,57,63]
- Eurytoma conica* Provancher [23]
- Eurytoma juniperina* Marcovitch [23]
  - E. juniperiana* Marcovitch [21]
  - E. juniperina* Marc. [26]
- Eurytoma laricis* Yano [28,48,72,75]
  - E. laricis* Jano [73]
  - Euritoma laricis* Yano [65]

## Torymidae

- Megastigmus albifrons* Walker [23,26,27,70]
  - M. grandiosus* Yoshimoto [23]
- Megastigmus amicorum* Bouček [51]
- Megastigmus atedius* Walker [9,23,54,70]
  - M. atedius* Walk. [48]
  - M. piceae* Rohwer [9,21,23,26,54]
  - M. zweelferi* Schefer-Immell [70]
  - M. zwolferi* Scheffer-Immell [9,73]
  - M. atedius* (Walker) [15,16]
- Megastigmus atedius atedius* Walker
- Megastigmus atedius montana* Milliron
  - M. piceae* var. *montana* Milliron [26]
- Megastigmus bipunctatus* (Swederus)
  - M. bipunctatus* Swed. [48,51,73]
  - M. bipunctatus* (Swed.) [25]
  - M. kuntzei* Kapuściński [25,40]
  - M. kuntzei* Kap. [24,25,48]
- Megastigmus borriesi* Crosby [28]
  - M. borriesi* [48]

- Megastigmus caperatus* Milliron  
*Megastigmus certus* Nikolskaya  
   *M. certus* Nikolskaja [40]  
   *M. certus* Nik. [48]  
   *M. cerfus* Nic. [73]  
*Megastigmus chamaecyparidis* Kamijo  
*Megastigmus cryptomeriae* Yano [25]  
   *M. cryptomeriae* [28]  
*Megastigmus ezomatsuanus* Hussey and Kamijo [28]  
*Megastigmus fidus* Nikolskaya  
   *M. fidus* Nik. [48]  
   *M. fidus* Nic. [73]  
*Megastigmus firmae* Kamijo  
*Megastigmus gravis* Nikolskaya  
   *M. gravis* Nik. [48]  
*Megastigmus hoffmeyeri* Walley  
*Megastigmus inamurae* Yano [28]  
*Megastigmus juniperi* Nikolskaya  
   *M. juniperi* Nik. [25,48]  
   *M. juniperi* Nic. [73]  
   *M. juniperi* Nikolskaja [40]  
*Megastigmus laricis* Marcovitch [23]  
*Megastigmus lasiocarpae* Crosby [21,23,26]  
   *M. lasiocarpae* [48]  
*Megastigmus milleri* Milliron [23,26]  
   *M. milleri* [48]  
*Megastigmus pictus* (Förster) [9,30]  
   *M. pictus* Hoffm. [48,73]  
   *M. pictus* (Först.) [56,57,59,63]  
   *M. pictus* (Fötrst.) [63]  
   *M. pictus* Först. [48]  
   *M. seitneri* Hoffmeyer [9]  
   *M. seitneri* Hoffmr. [30,56]  
   *M. seitneri* Hoffm. [57,65,71,72,75]  
   *M. seitneri* Ratz. [75]  
   *M. seitneri* Hffm. [74]  
   *M. seitneri* Hefm. [83]  
   *Magastigmus pictus* (Först.) [63]  
   *Torymus pictus* Förster [9]  
*Megastigmus pinus* Parfitt [9,21,23,26,48,70]  
   *M. pinus* Parf. [48]  
   *M. pinus* [24]  
*Megastigmus pinus crosbyi* Hoffmeyer  
*Megastigmus pinus marginatus* Hoffmeyer  
*Megastigmus pinus pinus* Parfitt  
*Megastigmus rafni* Hoffmeyer [21,23,26]  
   *M. rafni* [48]

- Megastigmus schimitscheki* Novitzky [10]  
*M. schimitscheki* Nozitzky [11]  
*M. schimitschekii* Novitz. [48]
- Megastigmus somaliensis* Hussey [24,25]
- Megastigmus specularis* Walley [9,23,48,70]  
*M. groenblomi* Kangas [9,48]  
*M. specularis* [48]
- Megastigmus spermotrophus* Wachtl [21,23,24,26,38,48,53,70]  
*M. spermotrophus* Ratz. [73]
- Megastigmus spermotrophus nigrodorsatus* Milliron  
*M. spermotrophus* var. *nigrodorsatus* Milliron [26]
- Megastigmus spermotrophus spermotrophus* Wachtl
- Megastigmus strobilobius* Ratzeburg [9,60,70]  
*M. strobilobius* Ratz. [38,48,61,73,74,83]  
*M. strobilobius* Seitn. [61]  
*M. abietis* Seitner [9,73]  
*M. abietis* Seitn. [72,83]
- Megastigmus suspectus* Borries [9,38,48,56,60]  
*M. suspectus* [37]  
*M. piceae* Seitn. [48,83]  
*M. piceae* Seitner [9]  
*M. bornmulleriana* Hussey [9,10,11,48]  
*M. suspectus* Borr. [48,62]  
*Mefastigmus suspectus* Borries [83]
- Megastigmus suspectus pinsapinis* Hoffmeyer  
*M. suspectus* var. *pinsapinis* Hoff. [48]  
*M. suspectus* Borries var. *pinsapinis* Hoff. [48]  
*M. suspectus* var. *pinsapinis* Hoffm. [48]
- Megastigmus suspectus suspectus* Borries
- Megastigmus thuyopsis* Yano [25]
- Megastigmus tsugae* Crosby [23,26]
- Megastigmus tsugae heterophyllae* Milliron  
*M. tsugae* var. *heterophyllae* Milliron [21,26]
- Megastigmus tsugae tsugae* Crosby
- Megastigmus tsugaphilus* Kamijo
- Megastigmus validus* Nikolskaya  
*M. validus* Nik. [48]  
*M. validus* Nic. [73]
- Megastigmus variegatus* Strand
- Megastigmus wachtli* Seitner [11]  
*M. wachtli* Seitn. [25]  
*M. waachtli* Seitner [48]  
*M. Wachtli* Seitner [10]

## Xyelidae

- Xyela alberta* (Curran) [23,66]
- Xyela apligena* (Strobl) [23,66]  
*X. brunneiceps* Rohwer [23,66]  
*X. middlekauffi* Burdick [23,66]

- Xyela bakeri* Konow [14,15,16,23,66]  
*Xyela californica* Rohwer [23,66]  
*Xyela cheloma* Burdick [23,66]  
*Xyela concava* Burdick [23,66]  
*Xyela curva* Benson [66]  
*Xyela deserti* Burdick [23,66]  
*Xyela dodgei* Greenbaum [15,16,23,66]  
*Xyela exilicornis* Maa [66]  
*Xyela graeca* Stein [66]  
*Xyela helvetica* (Benson) [66]  
*Xyela julii* (Brebisson) [66]  
  *X. julii* Breb. [76]  
*Xyela kamtshatica* Gussakovskij [66]  
*Xyela linsleyi* Burdick [23,66]  
*Xyela longula* Dalman [66]  
*Xyela lugdunensis* (Berland) [66]  
*Xyela lunata* Burdick [23,66]  
*Xyela menelaus* Benson [66]  
*Xyela minor* Norton [14,15,16,23,66]  
*Xyela obscura* (Strobl) [15,16,23,66]  
  *X. pini* Rohwer [14,23,66]  
*Xyela piliserra* Thomson [66]  
*Xyela priceae* Burdick [23,66]  
*Xyela radiatae* Burdick [23,66]  
*Xyela serrata* Burdick [23,66]  
*Xyela sinicola* Maa [66]  
*Xyela styrax* Burdick [23,66]  
*Xyela ussuriensis* Rasnitsyn [66]

## LEPIDOPTERA

### Argyresthiidae

- Argyresthia anthocephala* Meyrick  
  *A. anthocephala* [28]  
*Argyresthia chrysidella* (Peyerimhoff)  
  *Blastotere* (= *Argyresthia*) *chrysidella* Peyer. [48]  
  *B. chrysidella* Peyer [48,51]  
*Argyresthia libocedrella* Busck [23,26]  
*Argyresthia pseudotsuga* Freeman [53]  
*Blastotere praecocella* Zeller  
  *B. (= Argyresthia) praecocella* Zell. [48]  
  *B. praecocella* Zell. [48,51]  
*Argyresthia praecocella* Zell. [10,25]  
  *A. praecocella* Z. [25]  
*Argyresthia praecocella* Zell. [11]

### Blastobasidae

- Holcocera augusti* Heinrich [23,26]  
*Holcocera lepidophaga* Clarke [14,23]

*Holcocerina immaculella* (McDunnough)  
*Holcocera immaculella* McDunnough [23]  
*H. immaculella* McDermott [53]

#### Cochylidae

*Henricus fuscodorsanus* (Kearfott)  
*H. fuscodorsana* (Kearfott) [23]  
*H. fuscodorsana* (Kearf.) [26]  
*Henricus macrocarpanus* (Walsingham)  
*H. macrocarpana* (Walsingham) [23]  
*H. macrocarpana* (Wlshm.) [26]  
*Henricus melanoleucus* (Clarke)  
*Irazona melanoleuca* Dyar

#### Cosmopterigidae

*Periploca atrata* Hodges [23]

#### Gelechiidae

*Aristotelia latipennella* (Rebel)  
*Acanthophila latipennella* Rbl. [76]  
*A. latepiella* Rebl. [73]  
*A. letepellia* Ratz. [48]  
*A. piceana* Balz. [48]  
*A. piceana* Salz. [74]  
*A. piceana* Sulcz. [73]  
*Acantophila piceana* [77]  
*Battaristis vittella* (Busck)  
*Duvita vittella* Busck [23]  
*Brachyacma oxycedrella* (Millière)  
*B. oxycedrella* Mill. [48,51]  
*Chionodes periculella* (Busck) [23,26]  
*Chionodes sabinianus* Powell  
*C. sabinianae* Powell [23]  
*Exoteleia dodecella* (L.)  
*E. dodecella* Linnaeus [88]  
*E. (= Heringia) dodecella* L. [76]  
*Exoteleia burkei* Keifer [23]  
*Exoteleia nepheos* Freeman [23]

#### Geometridae

*Eupithecia abietaria* Goeze [48,72,73]  
*E. abietaria* Goetze [77]  
*E. abietaria* Goeze [74,77]  
*E. abietaria* [6]  
*E. pini* Ratz. [48]  
*E. pini* (Retz.) [6,83]  
*E. strobilata* [6]  
*E. togata* [6]  
*E. bilunulata* [6]

- Eupithecia abietaria debrunneata* Staudinger  
*E. abietaria* var. *debrunneata* St. [73]  
*E. abietaria* var. *debrumeata* St. [60]  
*E. strobilata debrunneata* Staudinger [28]  
*Eupithecia abietaria gigantea* Staudinger [88]  
*Eupithecia albicapitata* Packard [23,26]  
*Eupithecia bilunulata* (Zetterstedt)  
*E. bilunulata* (Zett.) [6,83]  
*E. bilunulata* Zett. [48]  
*E. strobilata* Hb. [48,72,73,74,77]  
*Eupithecia columbrata* McDunnough [23,26]  
*Eupithecia gigantea* Staudinger [28]  
*E. gigantea* St. [60,73]  
*Eupithecia mutata* Pearsall [23]  
*Eupithecia rufescens* Butler  
*E. rufescens* [28]  
*Eupithecia spermaphaga* (Dyar) [23,26]  
*Nepytia semiclusaria* (Walker) [15,16,23,43]  
*N. semiclusaria* (Wlk.) [14]  
*Sabulodes niveostriata* (Cockerell)  
*Phengommataea niveostriata* (Cockerell) [27]

#### Pyralidae

- Assara cedrella* (Hampson)  
*Cateremna cedrella* Hamp. [1]  
*Assara conicolella* (Constant)  
*Cateremna conicolella* Const. [48]  
*C. conicolella* Constant [48]  
*Assara terebrella* (Zincken)  
*A. terebrella* Znek. [65]  
*A. terebrella* Zinck [48,61]  
*Cateremna terebrella* Zinck [48]  
*Hyphantidium terebrella* Zinck [48]  
*H. terebrellum* Zinck. [60,72,73]  
*H. terebrellum* Zink. [74,77]  
*Cryptoblabes angustipennella* Ragonot  
*C. laricana* Matsuura [28]  
*C. laricana* Mat. [88]  
*Dichocrocis punctiferalis* (Guenée)  
*D. punctiferalis* [28]  
*D. punctiferalis* Guenée [81,88]  
*Dicocrocis punctiferalis* [28]  
*Dioryctria abietella* (Denis and Schiffermüller) [27,34,35]  
*D. abietella* Denis & Schiffermüller [28]  
*D. abietella* [28,37]  
*D. abietella* Denis and Schiff [60]  
*D. abietella* (Den. et Schiff.) [30,57,58,59,62,63]  
*D. abietella* (D. and S.) [5,14,26,39,84]  
*D. abietella* D. and S. [52]

- D. abietella* Den. and Schiff. [36,38,61]  
*D. abietella* (Schiff.) [1,3,7,12]  
*D. abietella* Schiff. [6,9,20,65,71,72,73,74,75,77,83,88]  
*D. abietella* F. [41,48,89]  
*D. abietella* Schiff. [11]  
*D. ebietella* Schiff. [11]  
*D. abietivorella* [88]
- Dioryctria abietivorella* (Grote) [21,23,34,53,54,55]  
*D. abietivorella* Grote [89]  
*D. reniculella* Grt. [52]  
*D. abietella* (D. and S.) [23]  
*D. abietinella* Grote [48]  
*Pinipestis reniculella* Grote [34]
- Dioryctria albovittella* (Hulst) [23,26]  
*D. albovittella* Hulst [89]
- Dioryctria amatella* (Hulst) [14,15,16,23,39,55]  
*D. amatella* Hulst [89]
- Dioryctria auranticella* (Grote) [21,23,26,27,33,52]  
*Dioryctria caesirufella* Blanchard and Knudson [8]  
*Dioryctria clarioralis* (Walker) [15,16,23,39,55]  
*D. clarioralis* Walk. [80]  
*D. clarioralis* (Wlk.) [14]
- Dioryctria disclusa* Heinrich [16,23,25,33,39,43,44,55,84]  
*D. disclusa* (Heinrich) [15]
- Dioryctria ebeli* Mutuura and Munroe [16,23,35]  
*D. abietella* D. and S. [15]  
*D. abietella* (D. and S.) [16,23,39]  
*D. abietivorella* (Grote) [15,16,23]  
*D. new species (abietella group)* [15]
- Dioryctria erythropasa* (Dyar) [23,26]  
*Dioryctria horneana* (Dyar) [23]  
*D. horneana* Dyar [80]
- Dioryctria majorella* Dyar [23]
- Dioryctria mendacella* Staudinger [81]  
*D. mendacella* (Stgr.) [12]  
*D. mendacella* Stgr. [10,11,48,88,89]
- Dioryctria merkeli* Mutuura and Munroe [16,23,35]  
*D. zimmermani* (Grote) [15,16,35,39]  
*D. new species (zimmermani group)* [15]
- Dioryctria mutatella* Fuchs [5,12,36,38,46,47,48,50,73]  
*D. mutatella* Fchs. [89]  
*D. mutatella* [37]
- Dioryctria peltieri* Joannis  
*D. peltieri* Joannis [48]
- Dioryctria pinicolella* Amsel
- Dioryctria pineae* Staudinger  
*D. pineae* Stgr. [10,11,48,89]

- Doryctria pryeri* Ragonot [33,88]  
*Salebria laruata* Heinrich [28,33]  
*Phycita pryeri* Leech [88]  
*Dioryctria pseudotsugella* Munroe [23,34]  
*Dioryctria pygmaeella* Ragonot [8,23]  
*Dioryctria reniculelloides* Mutuura and Munroe [23,34,54]  
*D. reniculelloides* [21]  
*D. reniculella* Grote [23]  
*D. reniculella* (Grote) [26,34,84]  
*D. reniculella* Grt. [52]  
*Pinipestis reniculella* Grote [34]  
*Dioryctria resinosella* Mutuura  
*Dioryctria rossi* Munroe [23,33]  
*D. rossi* [21]  
*Dioryctria rubella* Hampson<sup>7</sup>  
*Dioryctria schutzeella* Fuchs [5,34,81]  
*D. schutzeella* (Fuchs) [12]  
*D. schutzeella* Fuchs [36,48,74,77]  
*D. schutzella* Fuchs [73]  
*D. schutzeela* Fuchs [83]  
*Dioryctria simplicella* Heinemann  
*D. simplicella* Heinem. [36]  
*Dioryctria sylvestrella* (Ratzeburg)<sup>8</sup>  
*D. sylvestrella* Rtz. [89]  
*D. sylwestrella* (Ratz.) [5,12]  
*D. splendidella* (H.S.) [12]  
*D. splendidella* Herrich-Schaeffer [81]  
*D. splendidella* Herrich-Schäffer [28,88]  
*D. splendidella* H.S. [5,36]  
*Dioryctria sysstratiotes* Dyar [23]  
*Dioryctria taedae* Schaber and Wood [15,23,55]  
*Dioryctria yatesi* Mutuura and Munroe [16,23,35]  
*D. taedae* Schaber and Wood [23,35]  
*Herculia phoezalis* Dyar  
*Moodna biviella* L. [76]  
*Vitula lugubrella* (Ragonot)  
*V. sp. near lugubrella* (Ragonot) [23]  
*Vitula pinei* Heinrich [23,27]

#### Tortricidae

- Archips oporana* (L.)  
*A. piceana* L. [28]  
*A. piceana* L. [75]  
*Cacoecia piceana* L. [76,77]  
*Argyrotaenia urbana* (Busck)  
*A. urbana* Busck [23]

<sup>7</sup> Personal communication, Ping-Yuan Wang, Institute of Zoology, Academia Sinica, Beijing, People's Republic of China.

<sup>8</sup> Listed as damaging second-year cones, shoots, and trunk of pine in Japan [28]; other citations [5,12,36] indicate pine shoot and trunk attacks only.

- Barbara colfaxiana* (Kearfott) [21,23,26,53]
- B. colfaxiana* Kearfott [48]
  - B. colfaxiana colfaxiana* (Kearf.) [26]
  - Cydia pseudotsugana* Kearf. [26]
- Barbara colfaxiana coloradensis* (Heinrich) [26]
- Barbara colfaxiana taxifoliella* (Busck) [26]
- Barbara fulgens* V. I. Kuznetsov
- B. fulgens* VI. Kuzn. [73]
- Barbara herrichiana* Obraztsov [38,58,60]
- B. herrichiana* [37,48]
  - B. herrichiana* Obr. [48,58,62]
  - B. margarotana* Obr. [48]
  - B. margarotana* H.S. [46,47]
  - Evetria margarotana* H.S. [11,58]
  - Retinia margarotana* Wocke [58]
- Barbara mappana* Freeman [23,26]
- Barbara osmana* Obraztsov
- B. osmana* Obr. [10,11,48]
- Barbara ulteriorana* (Heinrich) [26]
- Barbara* sp.<sup>9</sup> [21,23,26]
- B. colfaxiana siskiyouana* (Kearfott) [23]
  - B. colfaxiana siskiyouana* (Kft.) [21,22]
  - B. colfaxiana siskiyouana* (Kearf.) [26]
- Choristoneura diversana* (Hübner) [88]
- C. diversana* Hb. [48,65,73,75]
- Choristoneura fumiferana* (Clemens) [23,43]
- Choristoneura lambertiana* (Busck) [23]
- Choristoneura lambertiana lambertiana* (Busck) [23]
- Choristoneura lambertiana subretiniana* Obraztsov [23]
- Choristoneura murinana* (Hübner)
- C. murinana* Hb. [73]
- Choristoneura occidentalis* Freeman [21,23,53]
- Choristoneura pinus* Freeman [23]
- Choristoneura pinus maritima* Freeman [23]
- Choristoneura pinus pinus* Freeman [23,43,44]
- Cydia anaranjada* (Miller)
- Laspeyresia anaranjada* Miller [14,15,16,23,39]
- Cydia araucariae* (Pastrana)
- C. (Laspeyresia) araucariae* Pastrana
- Cydia bracteatana* (Fernald)
- Laspeyresia bracteatana* (Fernald) [23]
  - L. bracteatana* (Fern.) [26]
  - L. pallidibasalis* Hein. [26]
- Cydia colorana* Kearfott
- Laspeyresia colorana* (Kearfott) [23]

<sup>9</sup> Formerly known as *Barbara colfaxiana siskiyouana* (Kearfott); however, this is an unfortunate misapplication of the specific name of *siskiyouana* that pertains to another moth, *Eucosma siskiyouana* (Kearfott). This true fir cone attacking *Barbara* species differs enough from *Barbara colfaxiana* (Kearfott) to merit a specific name.

- Cydia conicolana* (Heylaerts)  
*C. conicolana* Heyl. [46,48,50]  
*Laspeyresia conicolana* Heyl. [10,11,48]  
*Enarmonia conicolana* Heyl. [47,48]
- Cydia cryptomeriae* (Issiki)  
*Grapholitha cryptomeriae* [28]
- Cydia cupressana* Kearfott  
*Laspeyresia cupressana* (Kearfott) [23]  
*L. cupressana* (Kearf.) [26]
- Cydia ethelinda* (Meyrick)  
*C. ethelinda* Meyr. [3]
- Cydia illutana* (Herrich-Shäffer)  
*Laspeyresia illutana* H.-S. [71,72,73,74,75,77,83]  
*L. illutana* H.S. [48]
- Cydia illutana dahuricolana* (V. I. Kuznetsov)  
*Laspeyresia illutana* var *dahuricolana* Vl. Kuzn. [73]  
*L. illutana dahuricolana* Kuzn. [75]  
*L. illutana* Vl. Kurz. [65]
- Cydia ingens* (Heinrich)  
*Laspeyresia ingens* Heinrich [14,15,16,23,39]
- Cydia injectiva* (Heinrich)  
*Hedulia injectiva* Heinrich [23,26]  
*Laspeyresia injectiva* (Heinrich) [23]
- Cydia kamijoi* (Oku)
- Cydia miscitata* (Heinrich)  
*Laspeyresia miscitata* Heinrich [23,26]  
*L. miscitata* [21]
- Cydia nigra* (Miller)  
*Laspeyresia nigra* Miller [23]
- Cydia piperana* Kearfott  
*Laspeyresia piperana* (Kearfott) [21,23,27]  
*L. piperana* (Kearf.) [26]
- Cydia strobilella* (L.)  
*C. (Laspeyresia) youngana* (Kearfott) [54]  
*Laspeyresia strobilella* L. [10,11,28,61,65,72,73,74,77]  
*L. strobilella* (L.) [6,7]  
*L. strobiliella* Ratz. [83]  
*L. youngana* (Kearfott) [21,23,27]  
*L. youngana* (Kearf.) [26]  
*L. youngana* (Kft.) [84]  
*Lasp. strobilella* (L.) [6]  
*Pseudotomoides strobilellus* L. [38]  
*P. strobilella* L. [48]  
*P. strobilellus* (Linnaeus) [88]
- Cydia toreuta* (Grote)  
*Laspeyresia toreuta* (Grote) [15,16,23,27,43,44,70]  
*L. toreuta* Grote [14,39]
- Endopiza piceana* (Freeman)  
*Polychrosis piceana* Freeman [23,26]

- Epinotia hopkinsana* (Kearfott) [23]  
*E. hopkinsana* (Kearf.) [26]
- Epinotia hopkinsana cupressi* Heinrich [23,26]
- Epinotia nigricana* (Herrich-Shäffer)  
*E. nigricana* H.S. [38,48]  
*E. nigricana* [37]
- Eucosma bobana* Kearfott [19,23]  
*E. bobana* Kearf. [26]
- Eucosma cocana* Kearfott [15,16,23]
- Eucosma monitorana* Heinrich [23,43,84]  
*E. monitorana* Kearf. [44]
- Eucosma ponderosa* Powell [23]
- Eucosma pylonitis* Meyrick  
*E. pylonitis* Meyr. [1,2]
- Eucosma rescissoriana* Heinrich [23,26]  
*E. rescissoriana* Heinrich [21]
- Eucosma siskiyouana* (Kearfott) [23]
- Eucosma tocullionana* Heinrich [15,16,23]
- Exapate duratella* Heyden  
*E. duratella* Heyd. [48]
- Gravitarmata margarotana* (Heinemann) [88]  
*G. retiferana* Wocke [28]  
*G. margarotana* (H.S.) [12]  
*G. margarotana* H.S. [48,50]  
*G. margarotana* H.-S. [73]
- Evetria (Gravitarmata) retiferana* Weck. [11]  
*E. (Gravitarmata) retiferana* Wck. [10]  
*E. retiferana* Wck. [20]  
*E. margarotana* HS. [10,11]  
*E. mararotana* [88]  
*E. retiferana* Wocke [48]  
*Retinia retiferana* [88]
- Lobesia aeolopa* Meyrick [28]
- Lobesia reliquana* Hübner  
*L. reliquana* HB. [71,72,73]
- Pammene juniperana* (Millière)  
*P. juniperana* Mill. [48,51]  
*P. thuriferana* Cleu [48]
- Pammene oxycedrana* (Millière)  
*P. oxycedrana* Mill. [48,51]
- Pammene pontica* Obraztsov [10]  
*P. pontica* Obr. [11,48]
- Pseudococcyx tessulatana* (Staudinger)  
*P. tessulatana* Stgr. [48]  
*P. tessulanata* Stgr. [48]  
*Evetria tessulatana* Stgr. [10,11]

- Ptycholoma lecheanum* (L.)  
*P. lecheana* (Linnaeus) [88]  
*Cacoecia lecheana* L. [48,73,74,77]  
*C. lecheana* [88]
- Retinia cristata* (Walsingham)  
*Petrova cristata* (Walsingham) [88]  
*Evetria cristata* Walshingham [28]
- Retinia edemoidana* (Dyar)  
*Petrova* sp. poss. *edemoidana* (Dyar) [23]
- Retinia perangustana* Snellen  
*Petrova perangustana* Snellen [30,48,57,58,59]  
*P. perangustana* Snell. [48,63,73,75,88]  
*P. perangustana* Shell. [75]  
*Semasia perangustana* Snell. [83]  
*Laspeyresia zonovae* Flor. [73]
- Retinia immanitana* (V. I. Kuznetsov)  
*Petrova immanitana* Vl. Kuzn. [73]
- Retinia pini* (V. I. Kuznetsov)  
*Petrova pini* Vl. Kuzn. [73]
- Rhyacionia buoliana* (Denis and Schiffermüller)  
*R. buoliana* Schiff. [48,50]  
*R. buoliana* (Schiffermuller) [23]  
*R. buoliniana* Schiff. [46,47]  
*Rhuacionia buoliana* Den. et Schiff. [76]
- Rhyacionia duplana* (Hübner) [88]  
*Rhyacionia frustrana* (Comstock) [15,16,23]  
*Rhyacionia pinicolana* Doubleday  
*R. pinicolana* (Doubleday) [88]  
*R. pinicolana* Dbld. [47,48,50]
- Satronia tantilla* Heinrich [14,23]
- Spilonota laricana* Heinemann  
*S. laricana* Heineman [28]  
*S. laricana* Hein. [48,71,72,75]  
*S. laricana* Z. [73]  
*S. lariciana* (Heinemann) [88]  
*Spilonata laricana* Hein. [48]
- Zeiraphera canadensis* Mutuura and Freeman  
*Z. ratzeburgiana* Saxen [48]  
*Z. ratzeburgiana* Sax. [73,74,76,77]  
*Semasia ratzeburgiana* Rtz. [72]
- Zeiraphera diniana* Guenée [48]  
*Z. diniana* Guen. [48]  
*Zeirafera diniana* Gn. [75]
- Zeiraphera improbana* (Walker)  
*Z. diniana* (Guenee) [27]  
*Z. diniana* Guenée [88]
- Zeiraphera rufimitrana* (Herrich-Schäffer) [88]  
*Z. rufimitrana* H.S. [38,48,73]  
*Z. rufimitrana* [37]

## THYSANOPTERA

### Phlaeothripidae

- Gnophothrips fuscus* (Morgan) [15,16,23]
- G. piniphilus* Cwfd. [14]
- Leptothrips pini* (Watson) [23]
- Cryptothrips pini* (Watson) [23]

### Thripidae

- Chilothonips pini* Hood [23]
- Frankliniella bispinosa* (Morgan) [23]
- Frankliniella occidentalis* (Pergande) [23]
- Frankliniella tritici* (Fitch) [14,23]
- Oxythrips ajugae* Uzel [65,76]
- Oxythrips bicolor* (Reuter)
  - O. brevistylis* Tryb. [76]
- Oxythrips pallidiventris* Hood [23]
  - O. pallidiventris* Hd. [14]
- Thrips pini* (Uzel)
  - Taeniothrips pini* Uzel. [76]

## CLASS ARACHNIDA

### ACARI

#### Eriophyidae

- Eriophyes* (? = *Trisetacus*) *pini floricolus* Trotter [68]
- Eriophyes* (? = *Trisetacus*) *ramosus* Hodgkiss [68]
  - E. ramosus* Hodgkiss [24,67,68]
- Phytoptus* (? = *Trisetacus*) *quadrisetus* Thomas [68]
  - Trisetacus quadrisetus* (Thomas) [67,68]
  - T. quadrisetus* Thomas [21]
  - T. quadrisetus* Thom. [48,51]
- Trisetacus abietivagrans* Kadono [68]
- Trisetacus batonrougei* Smith [67,68]
- Trisetacus chamaecypari* Smith [68]
- Trisetacus kirghisorum* Schevchenko [67,68]
- Trisetacus macrocarpae* Smith [68]
- Trisetacus neoquadrisetus* (Smith) [67,68]

### ACKNOWLEDGMENT

I wish to acknowledge the invaluable assistance of taxonomists with the Systematic Entomology Laboratory, United States Department of Agriculture, Washington, D.C.: specifically D. M. Anderson; E. B. Baker; D. C. Ferguson; R. J. Gagné; R. D. Gordon; E. E. Grissell; T. J. Henry; R. W. Hodges; J. W. Kingsolver; P. M. Marsh; S. Nakahara; D. R. Smith; T. J. Spilman; M. B. Stoetzel; F. C. Thompson; R. E. White; D. R. Whitehead; and W. L. Murphy, technical information specialist; W. D. Duckworth with the Department of Entomology, Smithsonian Institution; H. H. Neunzig, North Carolina State University, Raleigh, N.C.; W. E. Miller, University of Minnesota, St. Paul, Minn.; and Ping Yuan Wang,

Institute of Zoology, Academia Sinica, Beijing, People's Republic of China. They are acknowledged for their careful checking of species names and clarification of currently accepted nomenclature.

I am also indebted to the following colleagues for their review of the manuscript: E. Annila, The Finnish Forest Research Institute, Helsinki, Finland; W. J. Mattson, North Central Forest Experiment Station, East Lansing, Mich.; G. Miller, Pacific Forest Research Centre, Victoria, Canada; A. Roques, Institut National de la Recherche Agronomique, Orleans, France; M. Skrzypczynska, Agricultural Academy of Krakow, Krakow, Poland; and G. Stadnitskii, Leningrad Forestry Research Institute, Leningrad, USSR.

#### LITERATURE CITED

1. Ahmad, S., R. Ahmad, and M. A. Ghani. 1977. Notes on the incidence and biology of some parasites of the pine cone-borers in Pakistan. *Entomophaga* 22: 107-10.
2. Ahmad, R., S. Ahmad, M. Salim, and M. A. Ghani. 1976. Incidence of *Eucosma pylonitis* Meyr. (Lepidoptera: Tortricidae) and its natural enemies in Pakistan. *Proc. Pakistan Acad. Sci.* 12: 85-94.
3. Ahmad, R., M. Salim, S. Ahmad, and M. A. Ghani. 1977. Studies on the seasonal incidence and biological mortality factors of the pine cone borer *Dioryctria abietella* (Schiff.) (Lep., Pyralidae) in Pakistan. *Tech. Bull.* 18, Common Inst. Biol. Control, pp. 113-27.
4. Aitkins, A. E. 1936. The 'spruce cone bug,' *Gastrodes abietis* Linn. (Hemiptera Heteroptera: Lygaeidae). *Entomol. Mon. Mag.* 72: 139-49.
5. Annila, E. 1979. The life cycles of the cone-infesting *Dioryctria* species (Lepidoptera, Pyralidae) in Finland. *Not. Entomol.* 59: 69-74.
6. Bakke, A. 1955. Insects reared from spruce cones in northern Norway. 1951. *Norsk. Entomol. Tidsskr.* 9(3-4): 152-212.
7. Bakke, A. 1963. Studies on the spruce-cone insects *Laspeyresia strobilella* (L.) (Lepidoptera: Tortricidae), *Kaltenbachiola strobi* (Winn.) (Diptera: Itonidae) and their parasites (Hymenoptera) in Norway. *Rep. Norw. For. Res. Inst.* 19(67). 151 p.
8. Blanchard, A., E. C. Knudson. 1963. A new species of *Dioryctria* Zeller (Lepidoptera: Pyralidae) from Texas. *Proc. Entomol. Soc. Wash.* 85: 116-20.
9. Bouček, Z. 1970. On some British *Megastigmus* (Hym. Torymidae), with a revised key to the west European species. *Entomol. Gaz.* 21: 265-75.
10. Canakcioglu, H. 1959. Studies on insects which are injurious to the Turkish forest tree seeds and control of some of the important species. Faculty of Forestry. Rev. Fac. Sci. For. Univ. Istanbul 155-167.
11. Canakcioglu, H. 1969. Insect damage on cones and seeds of forest trees in Turkey. *Istanbul. Univ. Orman Fakultesi* 19A: 83-6.
12. Charles, P. J., A. Roques. 1977. Observations sur la biologie de *Dioryctria mutatella* Fuchs (Lepidoptera: Phycitidae), ravageur des pousses et des cônes de pin sylvestre en forêt de Fontainebleau. *Ann. Zool. Ecol. Anim.* 9:117-31.
13. Coutin, R. 1976. Une cécidomyie nouvelle en France, ravageur des graines du cyprès de Lawson: *Janetiella siskiyou* Felt, 1917 (= *Craneobia lawsoniae* DeMeijere, 1935) (Dipt. Cecidomyiidae). *Bull. Soc. Entomol. Fr.* 81(1/2): 2-8.
14. Ebel, B. H. 1963. Insects affecting seed production of slash and longleaf pines: their identification and biological annotation. U. S. Dept. Agric. For. Ser. Res. Pap. SE-6. 24 p.
15. Ebel, B. H., T. H. Flavell, L. E. Drake, H. O. Yates III, and G. L. DeBarr. 1975. Seed and cone insects of southern pines. U. S. Dep. Agric. For. Ser. Gen. Tech. Rep. SE-8. 40 p.
16. Ebel, B. H., T. H. Flavell, L. E. Drake, H. O. Yates III, and G. L. DeBarr. 1980. Seed and cone insects of southern pines. U. S. Dept. Agric. For. Serv. Gen. Tech. Rep. SE-8 (Rev.). 43 p.

17. Ebel, B. H., R. J. Gagné, and E. P. Merkel. 1975. Cecidomyiidae from pine cones in Florida, with a generic key to larvae. Fla. Entomol. 58: 193-7.
18. Fan, Z. D., F. X. Ge, R. S. Zhang, and S. Y. Fang. 1982. [On Chinese species of genus *Lasiomma* especially those injurious to the coniferous cones (Diptera: Anthomyiidae).] J. North-Eastern Forestry Inst., China 1; 1-12. [in Chinese, with English summary].
19. Forcella, F. 1978. Irregularity of pinyon cone production and its relation to pinyon cone moth predation. Madrona 25(3): 170-2.
20. Gaidene, E. K., E. P. Nartshuk. 1963. On the biology of *Hapleginella laevifrons* Lw. (Diptera: Chloropidae) the cone pest. Entomol. Rev. 42: 411-3.
21. Hedlin, A. F. 1974. Cone and seed insects of British Columbia. Info. Rep. BC-X-90: Victoria, BC: Canadian Forestry Service, Environment Canada, Pacific Forest Research Centre. 63 p.
22. Hedlin, A. F., and D. S. Ruth. 1974. *Barbara califaxiana siskiyouana* (Kft), a pest in cones of *Abies grandis*. J. Entomol. Soc. BC: 71:13.
23. Hedlin, A. F., H. O. Yates III, D. Cibrian-Tovar, B. H. Ebel, T. W. Koerber, and E. P. Merkel. 1981. Cone and seed insects of North America conifers. 122 p. Can. For. Serv., U. S. Dep. Agric. For. Serv., and Secr. Agric. Recur. Hidrául. México. 122 p. (coop. publ., unnumbered, 2nd printing).
24. Hussey, N. W. 1960. Two new species of *Mesopolobus* Westwood (Hym., Pteromalidae) with notes on other parasites of *Megastigmus* Dalman (Hym., Torymidae). Entomol. Mon. Mag. 95: 237-9.
25. Kapuściński, S. 1964. *Megastigmus kuntzei* Kap. spotted juniper wasp (Hymenoptera, Chalcidoidea, Torymidae), a destructive insect feeding in seeds of common juniper (*Juniperus communis* L.), and other insects of genus *Megastigmus* Dalm. injurious to the seeds of Cupressaceae. Translated from Polish for USDA and NSF. 105 p. [Institut Polanais des Recherches Forestieres, Ser. A, No. 47; 1946. 129 p.]
26. Keen, F. P. 1958. Cone and seed insects of western forest trees. U. S. Dep. Agric. For. Serv. Tech. Bull. 1169. 168 p.
27. Kinzer, H. G., B. J. Ridgill, and J. G. Watts. 1972. Seed and cone insects of ponderosa pine. New Mexico St. Univ. Agric. Exp. Stn. Bull. 594. 36 p.
28. Kobayashi, K. 1981. Cone and seed insects of Japanese conifers. In Proc. Forest Plants and Forest Protection, XVII IUFRO World Cong. Div. 2, Kyoto, Japan. IUFRO Secretariat Vienna, Austria. pp. 455-61.
29. Kobayashi, K. T. Yokoyama. 1984. [Seed reduction in *Cryptomeria japonica* by infestation of stinkbugs.] For. Tree Breed. 133: 16-9. [in Japanese, with English abstract].
30. Kříštek, J., M. Skrzypczynska, J. Vrana. 1976. The insect pests attacking the ripening cones and seeds of European larch in southern Moravia. Acta Univ. Agric. 45(3-4): 149-61 [includes Czech summary].
31. Markalas, S., and D. S. Kailidis. 1982. *Ernobius kailidis* Johnson (1975) (Coleoptera, Anobiidae), ein neuer Tannenzapfenschadling in Griechenland. Anz. Schadlings. Pflanzl. 55: 23-7 [includes English abstract].
32. McAlpine, J. R. 1956. Cone-infesting lonchaeids of the genus *Earomyia* Zett., with descriptions of five new species from western North America (Diptera: Lonchaeidae). Can. Entomol. 88: 178-96.
33. Mutuura, A., and E. Munroe. 1972. American species of *Dioryctria* (Lepidoptera: Pyralidae) III. Grouping of species: species of the *auranticella* group, including the Asian species, with the description of a new species. Can. Entomol. 104: 609-25.
34. Mutuura, A., and E. Munroe. 1973. American species of *Dioryctria* (Lepidoptera: Pyralidae) IV. The *schuetzeella* group and the taxonomic status of the spruce cone moth. Can. Entomol. 105: 653-68.
35. Mutuura, A. and E. Munroe. 1979. American species of *Dioryctria* (Lepidoptera: Pyralidae) V. Three new cone-feeding species from the southeastern United States. J. Ga. Entomol. Soc. 14: 290-304.

36. Nanu, N. 1975. [Contribution to the knowledge of the pest *Dioryctria mutatella* Fuchs] Rev. Padurilor Ind. Lemnului Ser. Silvic. Exploataarea Padurilor 90(3): 150-2 [in Rumanian].
37. Nanu, N. 1978. The destroying insects of fir-tree (*Abies alba* Mill.) cone and seed: biology and ecology. In IUFRO Symposium on flowering and seed development in trees. 1978 May 15-18; Miss. State Univ., Starkville, Miss. p. 363.
38. Nanu, N. 1980. Biologia si combaterea principalelor insecte daunatoare fructificatiei rasinoaselor (duglas, molid, brad). Inst. Cercetari Amenajair Silvice II-a; 46 p. [in Rumanian, with English and German summaries].
39. Neel, W. W., and C. F. Sartor. 1969. Notes on insects infesting pine cones in Mississippi. Entomol. News 80: 159-67.
40. Nikol'skaya, M. N. 1966. Species of *Megastigmus* Dalm. (Hymenoptera, Torymidae) from seeds of *Juniperus*. Entomol. Rev. 45: 481-2.
41. Popova, A. A. 1973. Infestation of the seeds and cones of Siberian larch and Scots pine in the Baikal region by larvae of injurious insects. Entomol. Rev. 4: 495-9.
42. Popova, A. A., and K. J. Elberg. 1970. Species of the genus *Lasiomma* (Diptera, Anthomyiidae) which damage larch seeds and cones in the Baikal area. Entomol. Rev. 49: 340-5.
43. Rauf, A., D. M. Benjamin, R. A. Cecich. 1985. Insects affecting seed production of jack pine, and life tables of conelet and cone mortality in Wisconsin. For. Sci. 31: 271-81.
44. Rauf, A., R. A. Cecich, and D. M. Benjamin. 1981. Life table evaluation of conelet and cone mortality for jack pine, *Pinus banksiana* Lamb., in Wisconsin. R. P. Guries, ed. Proc. 2d North Central For. Tree Improv. Assoc. Conf. 1981 August 4-5, Lincoln, Neb. pp. 166-75.
45. Rauf, A., R. A. Cecich, and D. M. Benjamin. 1984. Conelet abortion in jack pine caused by *Platylygus luridus* (Hemiptera: Miridae) Can. Entomol. 116: 1213-8.
46. Roques, A. 1976. Observations sur la biologie des Diptères présents dans les cônes de pin sylvestre à Fontainebleau (Seine-et-Marne). Bull. Soc. Entomol. Fr. 81: 65-74.
47. Roques, A. 1977. Étude du complexe entomologique lié aux cônes de pin sylvestre en forêt de Fontainebleau. Bull. Ecol. 8: 415-34.
48. Roques, A. 1983. Les insectes ravageurs des cônes et graines de conifères en France. Paris: Institut National De Recherche Agronomique; 135 p.
49. Roques, A., M. Martinez, and A. Delplanque. 1983. Les diptères Anthomyiidae du genre *Lasiomma* Stein. ravageurs des cônes et graines de mélèze d'Europe (*Larix decidua* Mill.) en France. I: Taxonomie et répartition. Z. Angew. Entomol. 95: 429-39 [includes English abstract].
50. Roques, A., J. P. Raimbault, and F. Goussard. 1983. Impact des insectes ravageurs des cônes et graines sur la régénération naturelle du pin à crochets dans quelques forêts de montagne des Alpes du sud et des Pyrénées orientales. Acta. Biol. Mont. 2: 331-47 [includes English summary].
51. Roques, A., J. P. Raimbault, and F. Goussard. 1984. La colonisation des cônes et galbules des genévrier méditerranéens par les insectes et acariens et son influence sur les possibilités de régénération naturelle de ces essences. Ecol. Mediterr. 10 (Fac. 1-2): 147-69 [includes English summary].
52. Ross, D. A., and D. Evans. 1957. Annotated list of forest insects of British Columbia Part V — *Dioryctria* spp. (Pyralidae). Entomol. Soc. BC Proc. 53: 10-11.
53. Ruth, D. S. 1980. A guide to insect pests of Douglas-fir seed orchards. Info. Rep. BC-X-204. Victoria, BC: Can. For. Serv., Environ. Can., Pac. For. Res. Centre. 19 p.
54. Ruth, D. S., G. E. Miller, and J. R. Sutherland. 1982. A guide to common insect pests and diseases in spruce seed orchards in British Columbia. Info. Rep. BC-X-231. Victoria, BC: Environment Canada, Canadian Forestry Service, Pacific Forest Research Centre. 28 p.

55. Schaber, B. D., and F. E. Wood. 1971. A new species of *Dioryctria* infesting loblolly pine (Lepidoptera: Pyralidae). Proc. Entomol. Soc. Wash. 73: 215-23.
56. Skrzypczynska, M. 1974. [On the occurrence of male *Megastigmus pictus* (Forst.) (Hym., Torymidae).] Pol. Pismo Entomol. 44: 571-6 [in Polish].
57. Skrzypczynska, M. 1977A. Entomofauna of the cones of the European larch (*Larix decidua* Mill.) and Polish larch (*Larix polonica* Rac.) in Poland. Z. Angew. Entomol. 83: 238-47.
58. Skrzypczynska, M. 1977b. [*Petrova perangustana* Snellen (Lepidoptera, Tortricidae) injurious to seeds and cones of larches in Poland.] Pol. Pismo Entomol. 47: 117-21 [in Polish, with English abstract].
59. Skrzypczynska, M. 1978. Parasites of insects damaging seeds and cones of the European (*Larix decidua*) and Polish larches (*L. polonica*) in Poland. Entomophaga 23: 303-7.
60. Skrzypczynska, M. 1981. The entomofauna of the cones of fir (*Abies alba*) in Poland. Bull. Soc. Entomol. Suisse 54: 291-95.
61. Skrzypczynska, M. 1982a. The entomofauna of the cones of spruce, *Picea abies* (L.) Karst. in Poland. Z. Angew. Entomol. 94: 21-32.
62. Skrzypczynska, M. 1982b. [Pest of the cone seeds of *Abies alba* Mill. in the selected localities of the Beskid Sadecki region in the years 1976-1980.] Acta Agrar. Silvestria 21: 79-97 [in Polish, with English and Russian summaries].
63. Skrzypczynska, M. 1983. [Species composition and significance of insects found in the cones of European larch *Larix decidua* Mill. in the regions of the Forest Inspectorate at Piwniczna and Forest Experimental Station at Krynica.] Acta Agrar. Silvestria 22: 55-69 [in Polish, with English summary].
64. Skuhrava, M., and M. Skrzypczynska. 1983. [A review of gall-midges (Cecidomyiidae, Diptera) of Poland.] Acta Zool. Cracov. 26: 387-420 [in Polish, with English summary].
65. Smetanin, A. N. 1981. [Protection of seeds of coniferous trees under conditions of protected plantations.] Translated from the Russia by Amerind Publ. Co. Pvt. Ltd., New Delhi. 79 p. [Zashchita semyan khvoinykh porod v usloviyakh zapovednykh nasazhdennii. Kiev: Naukova Dumka Publ.; 1977. 59 p.]
66. Smith, D. R. 1978. Suborder Symphyta. Part 14. In van der Vecht, J., and R. D. Shenefelt, eds. Hymenopterorum catalogus. Holland, The Hague: Dr. W. Junk B. V. Publ. pp. 7-17 [Xyelidae].
67. Smith, I. M. 1978. Two new species of *Trisetacus* (Prostigmata: Eriophyoidea) from berries of junipers in North America. Can. Entomol. 110: 1157-60.
68. Smith, I. M. 1984. Review of species of *Trisetacus* (Acar: Eriophyoidea) from North America, with comments on all nominate taxa in the genus. Can. Entomol. 116: 1157-1211.
69. Smith, R. H., F. Mergen. 1954. *Pityophthorus pulicarius* (Zimm.), a bark beetle attacking scions of grafted slash pines. U. S. Dep. Agric. For. Ser. Res. Note 64. 2 p.
70. Speers, C. F. 1975. Seed chalcids: new insect pests of eastern white pine and Fraser fir seeds. In Southeastern Nurserymen's Conference Eastern Region, 1975. Gainesville, FL. p. 129-34.
71. Stadnitskii, G. V., L. S. Golutvina, and V. P. Grebenshchikova. 1976. [Pest of larch seed in seed-production areas.] Lesn. Zh. 2:151-5 [in Russian, with English available from Environ. Can. Transl. OOENV-1341].
72. Stadnitskii, G. V., V. P. Grebenshchikova, F. V. Naumov, and A. M. Bortnik. 1974. [Management and monitoring of pests of the reproductive organs of coniferous species and control of such pests in the seed sectors and plantations of the European part of the USSR.] Leningrad: Leningrad Science Research Institute Forestry, 73 p. [translated from Russian].
73. Stadnitskii, G. V., G. I. Iurchenko, A. N. Smetanin, V. P. Grebenshchikova, and M. V. Pribylova. 1978. [Conifer cone and seed pests.] Moscow: Lesn. Prom-st. 168 p. [translated from Russian].

74. Stadnitskiy, G. V. 1971. Characteristics of cones of Norway spruce (*Picea abies* (L.) Karst.) as a habitat and the complex of insects infesting them. Entomol. Rev. 50: 50-5.
75. Stadnitskiy, G. V., and V. P. Grebenshchikova, 1977. [The formation of an insect fauna in the reproductive organs of Siberian larch in artificial plantations.] Entomol. Rev. 56: 28-32.
76. Stadnitskiy, G. V., and V. P. Grebenshchikova, 1979. Insects developing in the male reproductive organs of conifers. Entomol. Rev. 58: 46-51.
77. Stadnitzky, G. W. 1969. Die entomofauna der Fichtenzapfen (*Picea abies* (L.) Karst). des europaischen teiles der UdSSR. Anz. Schadlingskd. Pflanzen. Unweltsch. 10 Schadlingsbekaempf. 42: 145-50.
78. Stone, A., C. W. Sabrosky, W. W. Wirth, R. H. Foote, and J. R. Coulson. 1965. A catalog of the Diptera of America north of Mexico. U. S. Dep. of Agric. Ag. Handb. 276. 1696 p.
79. Sweet, M. H. 1977. Elevation of the seedbug *Eremocoris borealis* (Dallas) from synonymy with *Eremocoris ferus* (Say) (Hemiptera: Lygaeidae). Entomol. News 88: 169-76.
80. Tsankov, G. 1977. [A study of insect pests of coniferous tree species in Cuba.] Gorskostop. Nauk. 14(2): 54-63 [in Russian, with English summary].
81. Wang, P.-y. 1980. [Economic insect fauna of China, Lepidoptera: Pyralidae.] Fasc. 21. China: Institute Zoology Academia Sinica Beijing. 229 p., 32 plates [in Chinese].
82. White, R. E. 1983. A new species of *Ernobius* (Coleoptera: Anobiidae) injurious to spruce. Proc. Entomol. Soc. Wash. 85: 557-9.
83. Wiersma, N. 1978. Cone and seed insects in Europe. In IUFRO Symposium on flowering and seed development in trees. 1978 May 15-18. Miss. State Univ.; Starkville, Miss., pp. 291-304.
84. Wilson, L. F. 1977. A guide to insect injury of conifers in the Lake States. U. S. Dep. Agric. For. Serv. Agric. Handb. 501. 218 p.
85. Wood, S. L. 1980. New American bark beetles (Coleoptera: Scolytidae), with two recently introduced species. Great Basin Nat. 40: 353-8.
86. Wood, S. L. 1982. The bark and ambrosia beetles of north and central America (Coleoptera: Scolytidae), a taxonomic monograph. Great Basin Nat. Mem. 6. 1359 p.
87. Xiao, G-r; Jian Wu. 1983. A new species of the genus *Augmonocetus* (Hymenoptera, Diprionidae) from China [Abstract]. Sci. Silvae Sinicae 19(2): 141-144 [in Chinese and English].
88. Xiu-Yaun, Y., and Jian Wu. 1981. [A checklist of the forest insects of China.] Chinese Acad. For., For. Res. Inst., Beijing, China. 444 p. [in Chinese].
89. Zocchi, R. 1961. Contributi alla conoscenza degli insetti delle piante forestali. V. II Gen. *Dioryctria* Zell. (Lepidoptera Pyralidae) in Italia. Redia 46: 9-143 + III, IV [in Italian, with English summary].
90. Zonova, E. V. 1935. [The larch fly (*Chortophila laricicola* Karl) as a larch seed pest.] Entomol. Obozr. 25: 206-20 [translated from Russian].