

Key to genera of Cactus Moths and their Relatives (Pyralidae: Phycitinae)

Thomas Simonsen

Department of Entomology, The Natural History Museum
Cromwell Road, London, SW7 5BD, United Kingdom

This key was modified from Neunzig 1997, Simonsen 2008, and Heinrich 1956.

- | | | |
|-----|---|-----------------------|
| 1. | Male..... | 2 |
| - | Female..... | 21 |
| 2. | Antenna bipectinate..... | 3 |
| - | Antenna not bipectinate..... | 8 |
| 3. | Flagellum of antenna with dorso-basal patch of scale-like sensilla | <i>Cactobrosis</i> |
| - | Flagellum of antenna without such patch..... | 4 |
| 4. | Abdomen 8 with two pair of ventro-lateral scale tufts..... | <i>Amalafrida</i> |
| - | Abdomen 8 without two such tufts..... | 5 |
| 5. | Forewing with M ₂ and M ₃ divided for less than half their length..... | <i>Melitara</i> |
| - | Forewing with M ₂ and M ₃ divided for more than half their length..... | 6 |
| 6. | Sharp ridge between eye and labial palpus; ocellus placed in an anterior incision of chaetosomata | 7 |
| - | No ridge between eye and labial palpus; ocellus placed adjacent to chaetosomata..... | <i>Alberada</i> |
| 7. | Antecosta of male T8 laterally lyre shaped..... | <i>Sigelgaita</i> |
| - | Antecosta of male T8 laterally curved but not lyre shaped..... | <i>Nanaia</i> |
| 8. | Flagellum of antenna with dorso-basal patch of scale-like sensilla..... | 9 |
| - | Flagellum of antenna without such patch..... | 11 |
| 9. | Hair-like sensilla on antenna almost as long as segments are wide..... | 10 |
| - | Hair-like sensilla on antenna clearly shorter than width of segment..... | <i>Ozamia</i> |
| 10. | Forewing with Rs ₂₊₃ and Rs ₄ diverging after the end of Rs ₁ ; forewing with M ₂ and M ₃ divided for less than half their length..... | <i>Baphala</i> |
| - | Forewing with Rs ₂₊₃ and Rs ₄ diverging before the end of Rs ₁ ; forewing with M ₂ and M ₃ diverged for more than half their length..... | <i>Zophodia</i> |
| 11. | Haustellum absent..... | <i>Rostrolaetilia</i> |
| - | Haustellum present..... | 12 |
| 12. | Antenna serrate..... | 13 |
| - | Antenna not serrate..... | 18 |

13.	Maxillary palpus fan-shaped.....	14
-	Maxillary palpus not fan-shaped.....	<i>Echinocereta</i>
14.	Hair-like sensilla on antenna almost as long as segments are wide.....	15
	Hair-like sensilla on antenna clearly shorter than width of segment..	<i>Salambola</i>
15.	Sharp ridge between eye and labial palpus.....	16
-	No ridge between eye and labial palpus.....	<i>Tucumania</i>
16.	Terminal segment of maxillary palpus pointed outwards.....	<i>Cactoblastis</i>
-	Terminal segment of maxillary palpus not pointed outwards.....	17
17.	Head of gnathos fused except most terminally.....	<i>Cahela</i>
-	Head of gnathos not fused except dorso-basally.....	<i>Eremberga</i>
18.	Dorsal base of uncus with a pair of rounded bumps.....	<i>Rumatha</i>
-	Dorsal base of uncus without rounded bumps.....	19
19.	Sharp ridge between eye and labial palpus.....	20
-	No ridge between eye and labial palpus.....	<i>Yosemitia</i>
20.	Chaetosomata placed posterior to dorso-basal corner of eye; tergal antecosta of A3 M-shaped.....	<i>Laetilia</i>
-	Chaetosomata placed anterior to dorso-basal corner of eye; tergal antecosta of A3 ring-shaped.....	<i>Rhagea</i>
21.	Dorsal sclerotization of abdomen 8 in lateral view triangular and saddle-shaped.....	22
-	Dorsal sclerotization of abdomen 8 different.....	25
22.	Antenna bipectinate.....	<i>Melitara</i>
-	Antenna not bipectinate.....	23
23.	Lamella antevaginalis prominent, entire and sclerotized.....	24
-	Lamella antevaginalis not prominent.....	<i>Alberada</i>
24.	Lamella antevaginalis with latero-anterior horns.....	<i>Olyca</i>
-	Lamella antevaginalis anteriorly smooth and rounded.....	<i>Cactoblastis</i>
25.	Ductus seminalis attached anteriorly on corpus bursae.....	26
-	Ductus seminalis attached more posterior on corpus bursae.....	34
26.	Maxillary palpus fan-shaped.....	27
-	Maxillary palpus not fan-shaped.....	29
27.	Dorsal sclerotization of abdomen 8 with large, unsclerotized, V-shape basal incision.....	28
-	Unsclerotized incision smaller and rounded.....	<i>Amalafrida</i>

28. Hind wing cell with vestigial anterior cross veins.....*Salambola*
 - Hind wing cell fully open without vestigial anterior cross veins.....*Nanaia*
29. Forewing Rs_{2+3} diverge well after the end of Rs_130
 - Forewing Rs_{2+3} diverge before the end of Rs_1*Zophodia*
30. Forewing M_2 and M_3 diverged for more than half their length.....31
 - Forewing M_2 and M_3 diverged for less than half their length.....*Baphala*
31. Forewing CuA_1 and CuA_2 close together at base, CuA_2 then sigmoid
 swung away from CuA_132
 - Forewing CuA_1 and CuA_2 not particularly close together, CuA_2 not
 sigmoid.....33
32. Hind wing cell with vestigial anterior cross veins.....*Cactobrosis*
 - Hind wing cell fully open without vestigial anterior cross veins...*Echinocereta*
33. Base of anterior apophysis narrow.....*Laetilia*
 - Base of anterior apophysis broad and diamond-shaped.....*Rhagea*
34. Dorsal sclerotization of abdomen 8 entire anteriorly; haustellum
 present.....35
 - Dorsal sclerotization of abdomen 8 incomplete anteriorly; haustellum
 absent.....*Rostrolaetilia*
35. Sharp ridge between eye and labial palpus.....36
 - No ridge between eye and labial palpus.....39
36. Dorsal sclerotization of abdomen 8 cylindrical.....37
 - Dorsal sclerotization of abdomen 8 flattened and clearly longer than high....
*Sigelgaita*
37. Base of posterior apophysis broad and diamond-shaped.....38
 - Base of posterior apophysis narrow.....*Rumatha*
38. Base of anterior apophysis broad with a ventral point; bursa ostium
 membranous ventrally.....*Cahela*
 - Base of anterior apophysis broad and diamond-shaped; bursa ostium
 sclerotized ventrally.....*Eremberga*
39. Ocellus placed in an anterior incision of chaetosomata.....*Tucumania*
 - Ocellus not placed in incision of chaetosomata.....40
40. Dorsal sclerotization of abdomen 8 with anterior centro-anterior incision.....
*Yosemitia*
 Dorsal sclerotization of abdomen 8 anteriorly entire.....*Ozamia*

References

- Heinrich, C. 1956. American moths of the subfamily Phycitinae. United States National Museum Bulletin 207, Washington, D.C., USA.
- Neunzig, H.H. 1997. Pyraloidea, Pyralidae, Phycitinae (Part). In R.B. Dominick, et al. [ed.] The Moths of America North of Mexico, Fascicle 15.4. The Wedge Entomological Research Foundation, Washington, D.C., USA.
- Simonsen, T.J. 2008. Phylogeny of the cactus-feeding phycitines and their relatives (Lepidoptera, Pyralidae) based on adult morphology: Evaluation of adult character-systems in phycitine systematics and evidence for a single origin of Cactaceae-feeding larvae. *Insect Systematics and Evolution* 39: 303-325.