

Acta Scientifica Naturalis

Former Annual of Konstantin Preslavsky University – Chemistry, Physics, Biology, Geography

Journal homepage: asn.shu.bg

Received: 03.2019

Accepted: 04.2019

Assessment of Antioxidant effect of the essential oil and methanol extract of *Centaurea dimorpha* Viv. aerial parts from Algeria.

 Abdelwahab Belkassam^{1,2}, Amar Zellaoui², Noureddine Gherraf³, Guido Flamini⁴, Pier Luigi Cioni⁴,
 Khellaf Rebbas¹ and Tahar Smaili^{1,2}
¹ *Département de Sciences de la Nature et de la Vie, Faculté de Sciences, Université de M'sila, 28000 M'sila, Algeria*
² *Laboratory of Biomolecules and Plant Breeding, Life Science and Nature Department, Faculty of Exact Science and Life Science and Nature, University of Larbi Ben M'hidi Oum El Bouaghi, Algeria*
³ *Laboratory of Natural Resources and Management of Sensitive Environments, Larbi ben M'hidi University, Oum El Bouaghi, Algeria,*
⁴ *Dipartimento di Scienze Farmaceutiche, Sede di Chimica Bioorganica e Biofarmacia, University of Pisa, Via Bonanno 33, 56126 Pisa, Italy*

Abstract: *The essential oils obtained by hydrodistillation of aerial parts of *Centaurea dimorpha* Viv. were analyzed by means of gas chromatography-mass spectrometry (GC-MS). The main constituents were Caryophyllene oxide (09.88%), limonene (5.73%), tetradecanoic acid (5.68%), spathulenol (5.44%), methyl hexadecanoate (4.45%), and α -pinene (03.08%). To our best knowledge this is the first study of essential oils from the aerial parts of *Centaurea dimorpha* growing in Algeria. Moreover the essential oil and methanolic extract were screened for their possible in vitro antioxidant activity by DPPH free radical-scavenging test. The findings showed that the percentage inhibition is 10.67 and 77.01 respectively at a concentration of 1mole/L.*

Keywords: *Centaurea dimorpha*; Essential Oil; GC-MS; aerial parts; Antioxidant activity

Introduction

Centaurea is the largest genus in Asteraceae family, comprising almost 400-700 species of annual, biennial and perennial grassy plants, predominately distributed in Mediterranean region and Asia [1-3].

Centaurea species have been used for their anti-dandruff, antidiarrhoic, antirheumatic, anti-inflammatory, choleric, diuretic, digestive, stomachic, astringent, antipyretic, cytotoxic, and antibacterial properties [4-6].

In Chinese folk medicine, *Centaurea uniflora* was used against fever and for detoxification. Ethyl acetate extract has an anti-atherosclerotic effect and was shown to inhibit membrane lipid peroxidation and [7,8]. Aqueous extract of *Centaurea chilensis* and *Centaurea ornata* was used to reduce fever and rheumatic pain [9, 10]. Tea prepared from the aerial parts of several *Centaurea* species are also used as

hypoglycaemic, such as *Centaurea ornato*, *Centaurea aspera*, *Centaurea seridis* and *Centaurea melitensis*, due to the bitter taste [11]. *Centaurea* species are also used as digestive tonic or stomachic, namely *Centaurea melitensis*, *Centaurea pallaszens* [12]. *Centaurea melitensis* and *Centaurea pallaszens* as well as *Centaurea sinaica* are also used as diuretic [13]. In Spain, aerial part of *Centaurea ornato* were used as depurative and cholagogue, while the underground part was used as antispasmodic. For *Centaurea sinaica* cytostatic, diuretic, antimalarial, astringent, antineoplastic, allergenic, stomachic, tonic and emmenagogue properties are attributed [8, 10].

Significant biological activities of *Centaurea* species led to deeper search on identifying responsible compounds. *Centaurea* species contain secondary metabolites, mainly, acetylenic compounds, flavonoids, terpenoids and, the most important ones are sesquiterpene lactones, having guaiane, germacrane, elemene and eudesmane skeletons [14-16].

To our best knowledge this paper is the first contribution in studying the composition and the antioxidant activity of the essential oil obtained from the aerial parts of *Centaurea dimorpha* (endemic in north Africa). Hence the main objectives of this study were:

- (i) Determine the chemical composition of hydrodistilled oils of the aerial parts of *Centaurea dimorpha* Viv. growing in Algeria by gas chromatography/mass spectrometry (GC/MS).
- (ii) Evaluate the antioxidant capacity of the plant essential oils by DPPH.

Materials and Methods

Plant material

The Aerial parts of *Centaurea dimorpha* Viv. were collected during February 2011 (flowering stage) near Bousaada M'sila (pre-saharan area) on a sandy soil, approximately 250 km south-east of Algiers, Algeria. Fresh aerial parts were dried to constant weight at room temperature. The plants were identified by Dr. Rebbas khellaf and a voucher sample was deposited in the Laboratory of Biomolecules and Plant Breeding, University of Larbi Ben M'hidi Oum El Bouaghi, under number ZARk 201). The dried plant material (146 g for aerial parts,) were hydrodistilled in a Clevenger-type apparatus for 2 h.

Gas chromatography-mass spectrometry:

GC-MS analyses were performed with a Varian CP-3800 gas chromatograph equipped with a DB-5 capillary column (30 m × 0.25 mm; coating thickness 0.25 μm) and a Varian Saturn 2000 ion trap mass detector. Analytical conditions: injector and transfer line temperatures 220 and 240°C, respectively; oven temperature programmed from 60°C to 240°C at 3°C/min; carrier gas helium at 1 mL/min; injection 0.2 μL (10% n-hexane solution); split ratio 1:30. Identification of the constituents was based on comparison of the retention times with those of authentic samples, comparing their linear retention indices relative to the series of n-hydrocarbons, and by computer matching against commercial (NIST 98 and ADAMS) and homemade library mass spectra built up from pure substances and components of known oils and MS literature data [17].

DPPH radical-scavenging activity

The capacity of essential oil and methanolic extract isolated from *Centaurea dimorpha* Viv. to reduce the radical 2,2-diphenyl-1-picrylhydrazyl (DPPH) was assessed using the method of Masuda *et al.* [18]. 15 μl of the essential oil and methanolic extract at different concentrations was added to 15 μl of a 55

Corresponding author: zellaguia@yahoo.com

DPPH ethanolic solution. The mixture was shaken vigorously and left standing at room temperature for 30 min in the dark. The absorbance of the resulting solution was then measured at 517 nm. The normal purple color of DPPH will turn into yellow when its singlet electron is paired with a hydrogen atom coming from a potential antioxidant. The scavenging activity of essential oil and methanolic extract was evaluated according to the formula:

DPPH· scavenging effect (%) = $(A_0 - A_1)/A_0 \times 100$, where A_0 is the absorbance of the control at 30 min, and A_1 is the absorbance of the sample at 30 min. All samples were analyzed in three replications.

Results and Discussion:

The essential oil composition of the aerial parts is reported in Table 1. In total, 99 Compounds were identified, representing about 79.21% of the whole oil. The essential oil (yield 0.02 , w/w) was mainly composed of oxygenated sesquiterpenes (27.49%), with caryophyllene oxide (09.88%) as The major constituent.

Other important classes of chemicals were monoterpene derivatives (17.29%), monoterpene hydrocarbons (10.65%), oxygenated monoterpenes (08.42 %), apocarotenoids (6.98%), sesquiterpene hydrocarbons (6.08 %) and phenylpropanoids compounds (2.30%). Other studies showed that the important classes for island *Centaurea* species (*C. veneris* *C. gymnocarpa* *C. aetaliae* *C. ilvensis* *C. aeolica* *C. busambarensis* *C. panormitana* *C. panormitana* *C. panormitana* *C. panormitana*) dominated by sesquiterpenes (22.35- 61.67%) and the main constituents is germacrene D in all species[19].

The main constituent of the essential oil under study was caryophyllene oxide (09.88%), followed by limonene (05.73%), tetradecanoic acid (5.68%), spathulenol (5.44%), methyl hexadecanoate (4.45%) , and α -pinene (3.08%). Similarly, the essential oil of the aerial parts of *Centaurea pichleri* ssp. *pichleri* growing in turkey consists of caryophyllene oxide (6.4%), spathulenol (6.2%) and dodecanoic acid (4.5%) [20].

Concerning the chemical composition of the essential oil of other *Centaurea* species it is worth noting the abundance of β -eudesmol (26.5%), spathulenol (06.3%) and caryophyllene oxide (02.9%) in *C. cuneifolai*, while *C. euxina* reveals mainly the presence of spathulenol (10.8%), caryophyllene oxide (06.2%) and β -eudesmol (03.9%) [21]. β -eudesmol (12.4%), caryophyllene oxide (10%), phytol (06.4) and spathulenol (04.9%) are present in *C. sessilis* [3]; Germacrene D (29.3%), β -eudesmol (17.4%) and β -caryophyllene (07.3%) in *C. mucronifera*; germacrene D (27.4%) , caryophyllene oxide (09.5%) and bicyclgermacrene (05.4%) in *C. chrysantha* [22]; hexadecanoic acid (39.3%), caryophyllene oxide (06.6%) and hexahydro farnesyl acetone (04.3%) in *C. aladagensis* , endemic in Turkey [6]. These differences, in most of the species, could be due to the local, ecological, climatic conditions of their habitat [23].

DPPH radical-scavenging activity

The highest DPPH radical scavenging activity (%) was shown by methanol extract which demonstrated a strong ability to act as a donor for hydrogen atoms or electrons. the reduction of a stable radical DPPH to yellow colored was obtained with 77.01% at 1M. Moreover essential oil exerted low antioxidant activity (10.67 %) at 1M compared to standard vitamin C (71%), fig.1. This is due to the influence of their different constituents especially the presence of phenolic compounds. The findings showed that the methanolic extract of *Centaurea pulchella* has the strongest antioxidant capacity

56
Corresponding author: zellaguia@yahoo.com

compared to other two *Centaurea* species and the highest level of phenolics was found in *C. pulchella*. The order of the antioxidant properties is *C. pulchella* > *C. patula* > *C. tchihatcheffii* [24]. It has been indicated in the literature that phenolic compounds have strong antioxidant properties [25]. Concerning Algerian *Centaurea*, it was reported that 3,5,7,4'-tetrahydroxy-6-methoxyflavone (6-methoxy kaempferol) isolated from *C. microcarpa* presented a more active antiradical scavenger than standard vitamin C [26].

In comparison with other *Centaurea* species, Marian et al (2017) reported that the DPPH Inhibition scavenging of *C. cyanus* growing in Romania exhibited a percentage of 83.42%, which is approximately in accordance with that found by Chougule et al. (2012) dealing with ethanol extract of the roots of *Centaurea behen* growing in Mumbai, India [27-28]. In the case of *C. calcitrapa* subsp. *calcitrapa*, *C. spicata*, *C. ptosimopappa* The results show that all of the three *Centaurea* species have antioxidant potential. DPPH scavenging activity seems to be higher in aqueous extracts of in *C. calcitrapa* and *C. spicata*, but it is higher in methanolic extract of *C. ptosimopappa*. The scavenging capacity is correlated to increasing concentration of extracts in all species [29].




Fig. 1: Scavenging activity of essential oils against the DPPH at different concentrations




Fig. 2: Scavenging activity of methanol extract against the DPPH at different concentrations

Table 1: Percentage Composition of the Essential Oil of the Aerial Parts of *Centaurea Dimorpha* from Algeria

Constituents	I.r.i. ^a	%
Furfural	830	0.53
(E)-2-hexanol	858	tr
1-hexanol	871	tr

2-heptanone	889	tr
Heptanal	899	0.20
(E,E)-2,4-hexadienal	913	tr
α – thujene	931	tr
α – pinene	939	3.08
camphene	954	tr
thuja- 2,4(10)-diene	957	0.19
benzaldehyde	963	0.23
1-heptanol	973	tr
β –pinene	980	0.19
6-methyl-5-hepten-2-one	987	tr
2-carene	1001	0.22
α –terpinene	1018	tr
ρ –cymene	1027	0.31
Limonene	1031	5.73
(Z)- β – ocimene	1040	tr
Phenylacetaldehyde	1045	1.01
(E) – β –ocimene	1051	0.45
γ - terpinene	1062	0.31
Acetophenone	1068	tr
p-mentha-2,4(8) diene	1086	0.17
6-camphenone	1095	0.30
Linalool	1100	0.56
Lonanal	1103	0.99
α –thujone	1105	tr
<i>cis</i> -p-menth-2-en-1-ol	1123	0.52
α –campholenol	1127	0.27
<i>cis</i> –limonene oxide	1134	tr
<i>trans</i> - pinocarveol	1140	0.29
<i>cis</i> - verbenol	1148	0.29
<i>trans</i> - verbenol	1144	1.69
<i>cis</i> –chrysanthenol	1164	tr
1-nonanal	1172	0.63
4-terpineol	1178	0.41
Nephthalene	1181	0.19
ρ –cymen-8-ol	1185	0.27
dihydrocarveol	1194	0.72
decanal	1205	0.50
verbenone	1209	0.40
<i>trans</i> –carveol	1219	0.58
nerol	1228	tr
<i>cis</i> - carveol	1231	tr
cuminaldehyde	1239	tr
carvone	1245	0.54
geraniol	1257	0.18
<i>cis</i> -chrysanthenyl acetate	-	tr
nonanoic acid	1278	0.86
isobornyl acetate	1286	0.28
thymol	1291	0.82
perilla alcohol	1295	tr
carvacrol	1300	tr
(E,E)-2,4-decadienal	1314	0.19
Myrtenyl acetate	1327	tr
α –terpinyl acetate	1351	0.30
daucene	1380	0.23
(E) – β -damascenone	1383	1.17
1-tetradecene	1392	tr

α -cedrene	1409	tr
β -cedrene	1418	0.90
2-methylbutyl benzoate	1436	tr
(E)-geranylacetone	1454	1.04
9- <i>epi</i> -(E)-caryophyllene	1467	tr
β -chamigrene	1475	0.47
germacrene D	1480	0.45
(E)- β -ionone	1485	2.13
α -bulnesene	1505	2.64
cubebol	1516	tr
myristicin	1523	2.30
β -thujaplicinol	1534	0.62
β -calacorene	1563	1.39
longicamphenylone	1559	1.14
spathulenol	1576	5.44
caryophyllene oxide	1581	9.88
globulol	1583	0.21
cedrol	1597	2.88
humulene oxide II	1606	1.88
1,10-di- <i>epi</i> -cubenol	1614	0.75
α -acorenol	1630	0.33
T-muurolol	1645	0.34
α -endesmol	1653	0.53
selin-11-en-4- α -ol	1652	1.36
14-hydroxy-9- <i>epi</i> -(E)-caryophyllene	1664	0.61
(Z)- α -santalool	1678	0.66
<i>cis</i> -14-nor-muurool-5-en-4-one	1682	0.71
acorenone	1685	0.40
n-heptadecane	1700	0.46
tetradecanoic acid	-	5.68
n-octadecane	1800	0.20
(E,E)- α -farnesyl acetate	1843	0.37
6,10,14-trimethylpentadecanone	1845	2.64
n-nonadecane	1900	0.16
methyl hexadecanoate	1927	4.45
n-docosane	2200	0.23
n-tricosane	2300	tr
n-pentacosane	2500	0.16

I.r.i.^a = Linear retention indices (HP-5 column).

tr = trace (< 0.1 %).

- = not detected.

Table 2: Chemical classes of the essential oil composition

Monoterpene hydrocarbons	10.65
Oxygenated monoterpenes	08.42
Sesquiterpene hydrocarbons	6.08
Oxygenated sesquiterpenes	27.49
Phenylpropanoids	2.30
Non-terpene derivatives	17.29
Apocarotenoids	6.98
Total identified (%)	79.21

Number of compounds	99
Essential oil yield (% , w/w)	0.02

Conflict of Interest: The authors declare no conflict of interest

Conclusion

Centaurea Centaurea Dimorpha from Algeria showed high diversity of essential oil components which may have a direct effect on antioxidant capacity. Moreover, methanol extract has been determined to have high Scavenging inhibition percentage against DPPH due to total phenolic content. As a result, this species can be used as a source of natural antioxidant in pharmaceutical and food processing fields. These results will contribute to further investigations for sources of natural antioxidant substances from other *Centaurea*.

References

- [1] Quezel, P. ; Santa, S., Nouvelle Flore de L'Algérie et des Régions Désertiques Méridionales. Editions Du Centre National de la Recherche Scientifique (C.N.R.S), France, Paris, **1963**, Tome II. 663 p.
- [2] Ozenda, P., Flore et végétation du Sahara. Edition, Centre National de la Recherche Scientifique (C.N.R.S) , Paris, **1991**. 453 p .
- [3] Yayli, N.; Yasar, A.; Güleç, C.; Usta, A.; Kolayli, S.; Coskunçelebi.; Karaoglu, S., Composition and antimicrobial activity of essential oils from *Centaurea sessilis* and *Centaurea armena*, *Phytochemistry*, **2005**, *66*, 1741-1745 .
- [4] Csupor, D.; Blazso, G.; Balogh, A.; Hohmann, J., The traditional Hungarian medicinal plant *Centaurea sadleriana* Janka accelerates wound healing in rats . *Journal of Ethnopharmacology*, **2010**, *127*, 193-195.
- [5] Zengin, G.; Cakmak, Y. S.; Guler, G.O.; Aktumsek, A., In vitro antioxidant capacities and fatty acid compositions of three *Centaurea* species collected from Central Anatolia region of Turkey, *Food and Chemical Toxicology*, **2010**, *48*, 2638-2641.
- [6] Köse, Y.B.; Iscan, G.; Demirci, B.; Baser, K.H.C. Celik, S., Antimicrobial activity of the essential oil of *Centaurea aladagensis*, *Fitoterapia*, **2007**, *78*, 253-254 .
- [7] Wei, H.X.; Gao, W.Y.; Tian, Y.K.; Guan, Y.K.; Huang, M.H.; Cheng, D.L., New sesquiterpene and thiophene derivatives from the roots of *Rhaponticum uniflorum*, *Die Pharmazie*, **1997**, *52*, 245–247.
- [8] Akkol, E.K.; Arif, R.; Ergun, F.; Yesilada, E., Sesquiterpene lactones with antinociceptive and antipyretic activity from two *Centaurea* species, *Journal of Ethnopharmacology*, **2009**, *122*, 210–215.
- [9] Sepulveda, S.; Delhvi, S.; Koch, B.; Zilliken, F., Constituents of *Centaurea chilensis*, *Fitoterapia*, **1994**, *65*, 88–89.
- [10] Al-Easa, H.S.; Kamel, A.; Rızık, A.F., Flavonoids from *Centaurea sinaica*, *Fitoterapia*, **1992**, *63*, 468–469.
- [11] Vazquez, F.M.; Suarez, M.A.; Perez, A., Medicinal plants used in the Barros Area, Badajoz Province (Spain), *Journal of Ethnopharmacology*, **1997**, *55*, 81-85.
- [12] Ali, Y.E.; Omar, A.A.; Sarg, T.M.; Slatkin, D.J., Chemical constituents of *Centaurea pallescens*, *Planta Medica*, **1987**, *53*, 503–504.

- [13] Kamanzi, K.; Raynaud, J.; Voirin, B., The C-glycosyl flavonoids from flowers of *Centaurea malitensis*, *Plantes Medicinal et Phytotherapie*, **1983**, *17*, 47–51.
- [14] Gonzalez, A.G.; Darias, V.; Alonso, G.; Boada, J.N.; Feria, M., Cytostatic activity of sesquiterpene lactones from compositae of the Canary Island, *Planta Medica*, **1978**, *33*, 356–359..
- [15] Gurbuz, I.; Yesilada, E., Evaluation of the anti-ulcerogenic effect of sesquiterpene lactones from *Centaurea solstitialis* L. ssp. *solstitialis* by using various in vivo and biochemical techniques, *Journal of Ethnopharmacology*, **2007**, *112*, 284– 291.
- [16] Koca, U.; Nahar, L.; Süntar, I. P.; Keles, H.; Yesilada, E.; Akkol, E. K., In vivo anti-inflammatory and wound healing activities of *Centaurea iberica* Trev . ex Spreng, *Journal of Ethnopharmacology*, **2009**, *126*, 551-556.
- [17] Rice-Evans, C.A.; Miller, N.J.; Bolwell, P.G.; Bramley, P.M.; Pridham, J.B., The relative antioxidant activities of plant derived polyphenolic flavonoids, *Free Radical Research*, **1995**, *22*, 375–383
- [18] Masuda, T.; Yonemori, S.; Oyama, Y.; Takeda, Y.; Tanaka, T.; Andoh, T.; Shinohara, A.; Nakata, M., Evaluation of the antioxidant activity of environmental plants: Activity of the leaf extracts from seashore plants, *Journal of Agriculture and Food Chemistry*, **1999**, *47*(4), 1749–1754.
- [19] Viegi, L.; Vangelisti, R.; Cecotti, R.; Tava, A., Essential oil composition of some *Centaurea* Sp. (Asteraceae) from different Italian islands, *Atti Soc. Tosc. Sci. Nat., Mem., Serie B*, **2014**, 121
- [20] Tastan, P.; Fafal, T.; Tuzun, B. S.; Gonenc, T.; Demirci, B.; Kivcak, B., Composition of essential oil and fatty acids of *Centaurea pichleri* ssp. *Pichleri*, *Int. J. Sec. Metabolite*, **2017**, *4*(3), 37-42
- [21] Rosselli, S.; Bruno, M.; Maggio, A.; Raccuglia, R.A.; Bancheva, S.; Senatore, F.; Formisano, C., Essential oils from the aerial parts of *Centaurea cuneifolia* Sibth. & Sm . and *C. euxina* Velen ., tow species growing wild in Bulgaria, *Biochemical Systematics and Ecology*, **2009**, *37*, 426-431 .
- [22] Rosselli, S.; Bruno, M.; Maggio, A.; Raccuglia, R.A.; Bancheva, S.; Senatore, F.; Formisano, C., Essential oil composition of two endemic *Centaurea* species from Turkey, *Centaurea mucronifera* and *Centaurea chrysantha* , collected in the same habitat, *Biochemical Systematics and Ecology*, **2009**, *31*, 1417-1425 .
- [23].Kilic, O., Essential oil compounds of three *Centaurea* L. taxa from Turkey and their chemotaxonomy. *J. Medic. Plants Res.*, **2013**, *7*(19), 1344-1350.
- [24] Davies., N.W., Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and Carbowax 20M phases, *Journal of Chromatography*, **1990**, *50*, 1-24
- [25] Zengin, Y.S.; Cakmak, G.O.; Guler, A.; Aktumsek, A., *In vitro* antioxidant capacities and fatty acid compositions of three *Centaurea* species collected from Central Anatolia region of Turkey, *Food and Chemical Toxicology*, **2010**, *48*, 2638-2641.
- [26] Louaar, S.; Zellagui, A.; Gherraf, N.; Medjroubi, K.; Derbre, S.; Seguin, E.; Laouer, H.; Akkal, S., Antiradical Activity of Flavonoids from the Algerian Native Plant: *Centaurea microcarpa* Coss. et Dur. *Journal of Biologically Active Products from Nature TBAP*, **2014**, *4* (3), 249 – 253.
- [27] Marian, E.; Vicas, L. G.; Tunde, J.; Muresan, M.; Stan, R. L.; Sevastere, B. Z.; Diaconeasa, C.; Ionescu, A.; Hangan, C., Comparative Study on the biologic activity of *Centaurea cyanus* versus, *Calendula officinalis*, *Farmacica*, **2017**, *65*, 6.

- [28]. Chougule, P.; Pawar, R.; Limaye, D.; Joshi, Y. M.; Kadam, V., *In-Vitro* Antioxidant Activity of Ethanolic Extract of *Centaurea behen*, *Journal of Applied Pharmaceutical Science*, **2012**, 02 (04), 106-110.
- [29] Öa, E.-D.; Pekmez, M.B.; Bona, M.C.; Aras-Perk, A.C.; Arda, N., Total Phenolic Contents, Antioxidant Activities and Cytotoxicity of Three *Centaurea* Species: *C. calcitrapa* subsp. *calcitrapa*, *C. ptosimopappa* and *C. spicata*, *Free Radicals and Antioxidants*, **2011**, 1, (2), 31-36