

Copyright © 2019 by Academic Publishing House Researcher s.r.o.



Published in the Slovak Republic
Central European Journal of Botany
Has been issued since 2015.
E-ISSN 2413-757X
2019, 5(1): 3-21

DOI: [10.13187/cejb.2019.1.3](https://doi.org/10.13187/cejb.2019.1.3)
www.ejournal34.com



Articles and Statements

Floristic Diversity and Ecological Characteristics of Historical Kalash Valley (Pakistan)

Fazal Hadi ^{a,*}, Muhammad Ibrar ^a, Ömer Kılıç ^b

^a University of Peshawar, Peshawar, Pakistan

^b Technical Science Vocational College, Bingol University, Turkey

Abstract

Kalash valley is famous all over the world for its inhabitants having indigenous culture, religion, language and considered to be the descendants of Alexander the Great living in this rugged mountainous series of Hindu Kush range. Frequent field visits were carried out during 2013–2016 years for the collection of plant samples and documentation of related data. Then the collected plants samples were properly pressed and identified with the help of different volumes of Flora of Pakistan and other available literature. The study showed 389 species belong to 237 genera and 88 families including 62 dicots, 11 monocots, 7 pteridophytes, 4 bryophytes, 3 gymnosperms and 1 fungi family. Species distribution indicated 307 dicots, 55 monocots, 12 pteridophytes, 10 gymnosperms, 4 bryophytes and 1 fungi species. Asteraceae was detected dominant (49 taxa) family followed by Poaceae (36 spp.), Brassicaceae (25 taxa), Lamiaceae (22 spp.), Rosaceae (22 spp.), Fabaceae (16 spp.) and Apiaceae (13 spp.). The remaining families had 12 or less species. The life form classes indicated 205 therophytes, 56 geophytes and 46 hemicryptophytes species while nanophylls (178 spp.) mesophylls (72 spp.) and leptophylls (53 spp.) were dominant leaf spectra classes. The present study is first ever record of floristic diversity of Kalash valley and includes some new reports. The findings will be helpful in future detail studies on plant resources of the area.

Keywords: floristic diversity, plant ecology, Kalash valley, district Chitral, Pakistan.

1. Introduction

Chitral is the largest district of Khyber Pakhtunkhwa and lies between $35^{\circ} 15' 06''$ to $36^{\circ} 55' 32''$ north latitudes and $71^{\circ} 11' 32''$ to $73^{\circ} 51' 34''$ east longitudes (Anonymous, 1998). The district has boundaries with Ghizer valley (Gilgit-Baltistan) in the east, with districts of Dir and Swat in the south and with Nooristan province and Wakhan corridor of Afghanistan in the west and north-west respectively (Figure 1). The present study area “Kalash valley” lies to the extreme south-west of district Chitral and is bounded with Nooristan (Afghanistan) in the west and to other parts of Chitral by the three sides. The valley has three sub localities viz: Bumburet, Birir, Rumbur and inhabits people that practice an old, unique and indigenous religion called Kalasha (Figure 2).

* Corresponding author
E-mail addresses: hadi@uop.edu.pk (F. Hadi)



Fig. 1. Map of Pakistan on the globe showing Chitral Valley in Black Square box




Fig. 2. Map of district Chitral showing the study area in black square box

The people are mostly dependant on different products of plants for food, fodder, medicines and shelter while the natural forests are comprise of *Cedrus deodora*, *Juniperus communis*, *Juniperus excelsa*, *Pinus gerardiana*, *Pinus wallichiana*, *Quercus baloot* and *Quercus incana*. Cattle rearing are major income source for the locals and the valley has rangelands where the Kalash people graze their animals and also collect plants for domestic and commercial use. The extensive exploitations of these rangelands for different purposes have caused serious drastic changes in annual climatic conditions that lead to severe floods which affected infrastructures, cultivated land, service roads and irrigation channels of the valley.

Flora is the sum of all plant species in a particular habitat and ecosystem. Floristic diversity and its ecological characteristics depend upon climate, altitudinal and related habitat conditions (Ali, 2008). Many researcher have studied on floristic checklists of local flora in different regions

(Durrani et al., 2005; Sher, Khan 2007; Qureshi, Bhatti, 2008; Abdullahi et al., 2009; Jabeen et al., 2009; Saima et al., 2009; Durrani et al., 2010; Udayakumar et al., 2011; Hussain et al., 2012; Qin et al., 2012; Youcef et al., 2012; Badshah et al., 2013; Sher et al., 2014; Hussain et al., 2015; Badshah et al., 2016; Baykal, Atamov 2016; Shaheen et al., 2016).

The present study was initiated to enlist the unexplored flora of Kalash valley, district Chitral which will help for future studies and planning in the area regarding plant resources and their utilization for different purposes.

2. Material and Methods

Plant specimens were collected from entire Kalash valley during 2014-2016. They were dried and identified with the help of available literature (Nasir & Ali, 1970-1989; Ali & Nasir, 1989-1991; Ali & Qaiser, 1993-2015). A complete floristic list was prepared and plants were placed into genera and family. The identified plant specimens were deposited at Herbarium of Department of Botany, University of Peshawar. The ecological characteristics of species viz: leaf persistence, spiny nature, habitat, light requirement, cultivation/wild and leaf shape were observed in the field. Life-form and leaf size spectra were determined by following Raunkiaer (1934) and Hussain (1989).

3. Results and Discussion

a. Flora and its ecological characteristics:

The present study revealed that the Kalash valley contained 389 species of 273 genera and 88 families including 62 (70.45 %) dicots, 11 (12.50 %) monocots, 7 (7.95 %) pteridophytes, 4 (4.54 %) bryophytes, 3 (3.41 %) gymnosperms and 1 (1.13 %) fungi families. Dicots had maximum 307 (78.92 %) species followed by monocot with 55 (14.13 %) species, pteridophytes 12 (3.08 %), gymnosperms 10 (2.57 %), bryophytes 4 (1.03 %) and 1 (0.26 %) was fungi species. Asteraceae had 49 (12.60%) taxa, followed by Poaceae having 36 (9.25 %) species, Brassicaceae 25 (6.42 %) taxa, Lamiaceae and Rosaceae 22 (5.65 %) each, Fabaceae 16 (4.11%), Apiaceae 13 (3.34 %), Boraginaceae and Polygonaceae 12 (3.08 %) each and Schrophulariaceae and Caryophyllaceae had 10 (2.57 %) species each. The remaining families have less than 10 species (Table 1, Figures 3-5). Our findings are supported by different workers viz: Shah and Hussain (2007) that reported 42 species belonging to 31 families from Chamkani area, Peshawar with similar dominant families. Sher and Khan (2007) reported 222 species belonging to 89 families with maximum representation (21 taxa) in Asteraceae. Durrani et al., (2010) reported 123 species of 36 families from Aghberg rangelands Quetta, whereas Asteraceae, Boraginaceae, Brassicaceae, Fabaceae, Lamiaceae and Poaceae were dominant families of the area. 571 species belonging to 82 families were reported from Mastuj valley, Chitral and Asteraceae was leading family (Hussain et al., 2015). Badshah et al., (2016) reported 283 species belonging to 85 families from Parachinar, Kurram Agency where Asteraceae was dominant family with 29 species. These findings are in support of our results as Asteraceae, Poaceae, Fabaceae, Chenopodiaceae and Moraceae were also leading families in their research areas.

The ecological characteristics indicates that the valley included 328 (84.32%) deciduous, 48 (12.34 %) evergreen and 13 (3.34 %) leafless species. This high percentage of deciduous plants indicates the xeric and dry conditions of the area. The leaf appearance showed that 266 (68.38 %) species had simple, 66 (16.97 %) species had compound, 44 (11.31 %) species had dissected and 13 (3.34 %) species were aphyllous. 369 (94.85 %) species were spineless (non-spiny) and 20 (5.15 %) were spiny species (Figures 8-10). The results showed that 291 (74.81%) species were in dry and 98 (25.19 %) were of moist habitats like *Fumaria hygrometrica*, *Marchantia polymorpha*, *Adiantum venustum*, *Equisetum arvense*, *Selaginella indica*, *Phragmites karka*, *Mentha longifolia*, *Nasturtium officinale* and *Ranunculus muricatus* etc. which agrees with Hussain et al., (2015) and Badshah et al., (2016) with similar species in such dry climatic conditions of the valley where only few localized points having aquatic/moist species. 363 (93.31 %) species were heliophytes (light loving) and only 26 (6.69 %) species were sciophytes (shade loving) as the area consists of scattered vegetation with open sunny environment and favours sun loving plants. Similarly, 340 (87.40 %) species were wild and 49 (12.60 %) were cultivated (Figures 11-13). The cultivated species included crops, fruit plants and vegetables.

b. Life form spectrum

The life form spectrum shows all the life processes of flora and vegetation. It usually represents climatic, microclimatic and bioclimatic habitat conditions of an area (Cain, Castro, 1959). It plays key role in the detection of environmental amplification of plant life. The life form spectrum may be changed due to introduction of therophytes like annual weed, due to anthropogenic activities like agricultural practices, deforestation, grazing and trampling etc. (Malik et al., 2007). In the present study therophyte was dominant class with 205 (52.70 %) species followed by geophytes with 56 (14.40 %) species, hemicryptophytes 46 (11.82 %), nanophanerophytes 29 (7.45 %), chamaephytes 21 (5.40 %), megaphanerophytes 19 (4.90 %), climbers 6 (1.54 %), mesophanerophytes and macrophanerophytes 5 (1.28 %) species each and parasites had 3 (0.077 %) species (Figure 6).

The dominance of therophytes and hemicryptophytes in the valley indicates that the area is under great anthropogenic pressure like deforestation and overgrazing that is destroying the natural habitat for other classes and the vegetation is dominating by annual plants. Our findings are supported by Nazir, Malik 2006; Guo et al., 2009; Kar et al., 2010; Manhas et al., 2010; Fatna et al., 2014; Khan et al., 2014; Osman et al., 2014; Hussain et al., 2015; Badshah et al., 2016; Baykal, Atamov, 2016 and Shaheen et al., 2016 that enlisted therophyte plants dominating in such degraded dry and arid environments.

c. Leaf size spectrum

The leaf size classes show that nanophylls dominated the valleys with 178 (45.76 %) species followed by mesophylls with 72 (18.51 %), leptophylls 53 (13.62 %), microphylls 44 (11.31 %), macrophylls 29 (7.45 %) species and 13 (3.34 %) species were without leaves (Figure 7). Leaf size classes are considered to be useful in determination of plant associations. The Kalash valley is dominated by reduced leaf sized plants (nanophylls) that indicates its extreme condition. The findings of Durrani et al., 2005; Alelign et al., 2007; Sher, Khan, 2007; Parswan et al., 2010; Hussain et al., 2015 and Badshah, 2016 support our results that found nannophylls as dominant class in their studies from such severe climatic areas.

Table 1. Floristic list and ecological characteristics of plants of Kalash Valley, District Chitral, Hindukush range, Pakistan

S#	Plant name	Kalash valley			Plant characteristics								
		B	R	BR	1	2	3	4	5	6	7	8	
FUNGI													
1. Family Halveliaceae													
1.	<i>Morchella esculenta</i> (L.) Pers ex. Fr	+	+	+	Th	Ap	-	-	M	Sd	W	Ap	
BRYOPHYTES													
2. Family Funariaceae													
2.	<i>Funaria hygrometrica</i> Hedw.	+	+	+	Th	Ap	-	-	M	Sd	W	Ap	
3. Family Marchantiaceae													
3.	<i>Marchantia polymorpha</i> L.	+	-	+	Th	Ap	-	-	M	Sd	W	Ap	
4. Family Atyoniaceae													
4.	<i>Reboulia hemisphaerica</i> (L.) Raddi	+	-	-	Th	Ap	-	-	M	Sd	W	Ap	
5. Family Polytrichaceae													
5.	<i>Polytrichum</i> sp.	+	+	+	Th	Ap	-	-	M	Sd	W	Ap	
PTERIDOPHYTES													
6. Family Adiantaceae													
6.	<i>Adiantum capillus-veneris</i> L.	+	+	+	G	N	E	-	M	Sd	W	Com	
7.	<i>Adiantum venustum</i> D. Don.	+	-	+	G	N	E	-	M	Sd	W	Com	
8.	<i>Cheilanthes pteridioides</i> (Reichb.) C. Chr	-	-	+	G	N	E	-	M	Sd	W	Com	
7. Family Aspleniaceae													
9.	<i>Asplenium trichomanes</i> L.	-	-	+	G	N	E	-	M	Sd	W	Com	

8. Family Cystopteridaceae												
10.	<i>Cystopteris fragilis</i> (L.) Bernh.	-	+	+	G	N	E	-	M	Sd	W	Com
9. Family Dryopteridaceae												
11.	<i>Dryopteris ramosa</i> (Hope) C. Chr.	-	-	+	G	L	E	-	M	Sd	W	Com
10. Family Equisetaceae												
12.	<i>Equisetum arvense</i> L.	-	+	-	G	Ap	E	-	M	Sd	W	Ap
13.	<i>Equisetum ramosissimum</i> Desf.	+	+	-	G	Ap	E	-	M	Sd	W	Ap
11. Family Pteridaceae												
14.	<i>Cystopteris fragilis</i> (L.) Bernh.	-	-	+	G	N	E	-	M	Sd	W	Com
15.	<i>Polystichum acanthophyllum</i> (Franch) Christ	-	-	+	G	N	E	-	M	Sd	W	Com
16.	<i>Pteris cretica</i> L.	-	-	+	G	N	E	-	M	Sd	W	Com
12. Family Selaginellaceae												
17.	<i>Selaginella indica</i> (Milde) Alston.	+	+	+	G	N	E	-	M	Sd	W	Com
GYMNOSPERMS												
13. Family Cupressaceae												
18.	<i>Juniperus communis</i> L.	-	+	+	NP	L	E	-	D	L	W	Dis.
19.	<i>Juniperus excelsa</i> M. Bieb.	-	+	+	Meg P	L	E	-	D	L	W	Dis.
20.	<i>Juniperus squamata</i> Buch.	-	-	+	NP	L	E	-	D	L	W	Dis.
21.	<i>Thuja orientalis</i> L.	+	-	-	NP	L	E	-	D	L	C	Dis.
14. Family Ephedraceae												
22.	<i>Ephedra gerardiana</i> Wall ex. Stapf.	-	+	+	Ch	Ap	E	-	D	L	W	Ap
23.	<i>Ephedra intermedia</i> Schrenk & Meyer	-	+	+	Ch	Ap	E	-	D	L	W	Ap
15. Family Pinaceae												
24.	<i>Cedrus deodora</i> (Roxb. Ex Lamb.) G.Don	+	+	+	Meg P	L	E	-	D	L	W	Com
25.	<i>Picea smithiana</i> (Wall.) Boiss.	+	+	+	MegP	L	E	-	D	L	W	Com
26.	<i>Pinus gerardiana</i> Wall. ex Lamb.	+	+	+	MegP	L	E	-	D	L	W	Com
27.	<i>Pinus wallichiana</i> A.B. Jackson	+	+	+	MegP	L	E	-	D	L	W	Com
MONOCOTYLEDANAE												
16. Family Alliaceae												
28.	<i>Allium carolinianum</i> DC.	+	-	+	G	Mes	Dec	-	D	L	W	S
29.	<i>Allium cepa</i> L.	+	+	+	G	Mes	Dec	-	D	L	C	S
30.	<i>Allium chitralicum</i> Wang & Tang	+	+	-	G	Mes	Dec	-	D	L	W	S
31.	<i>Allium sativum</i> L.	+	+	+	G	Mes	Dec	-	D	L	C	S
17. Family Amaryllidaceae												
32.	<i>Ixiolirion tataricum</i> (Pall.) Herb.	-	-	+	G	Mes	Dec	-	M	L	W	S
18. Family Araceae												
33.	<i>Arum jacquemontii</i> Blume	+	+	+	G	Mic	Dec	-	M	L	W	S
19. Family Colchicaceae												
34.	<i>Colchicum luteum</i> Baker	-	-	+	G	N	Dec	-	M	Sd	W	S
20. Family Cyperaceae												
35.	<i>Carex orbicularis</i> Boott.	-	+	-	G	L	E	-	M	L	W	S
36.	<i>Cyperus diluta</i> M. Bieb.	+	+	-	G	L	E	-	M	L	W	S
37.	<i>Scirpus setaceus</i> L.	+	+	-	G	Mes	E	-	M	L	W	S

21. Family Iridaceae												
38.	<i>Iris germinea</i> L.	-	+	+	G	Mes	Dec	-	D	L	C	S
39.	<i>Iris lactea</i> Pallas	+	-	+	G	Mes	Dec	-	D	L	W	S
22. Family Juncaceae												
40.	<i>Juncus articulata</i> L.	+	+	+	H	Mic	E	-	M	L	W	S
23. Family Liliaceae												
41.	<i>Fritillaria imperialis</i> var. <i>chitralensis</i> Hort.	-	-	+	G	Mes	E	-	M	L	W	Com
42.	<i>Tulipa stellata</i> Hk. f.	+	+	+	G	L	Dec	-	M	L	W	S
24. Family Orchidaceae												
43.	<i>Epipactis wallichii</i> Schlech.	+	-	-	G	N	Dec	-	M	Sd	W	S
44.	<i>Spiranthes lancea</i> (Thunb.) Backer	+	-	-	G	N	Dec	-	M	Sd	W	S
25. Family Poaceae												
45.	<i>Agropyron semicostatum</i> Nees ex Steud	-	+	+	Th	N	E	-	D	L	W	S
46.	<i>Agrostis viridis</i> Gouan.	-	+	-	Th	N	E	-	D	L	W	S
47.	<i>Aristida cynantha</i> Nees ex Steud	-	+	+	H	N	E	-	D	L	W	S
48.	<i>Arundo donax</i> L.	+	-	-	Ch	Mac	Dec	-	M	L	W	S
49.	<i>Avena fatua</i> L.	+	-	+	Th	N	E	-	D	L	W	S
50.	<i>Avena sativa</i> L.	+	+	+	Th	N	E	-	D	L	W	S
51.	<i>Bromus danthoniae</i> Trin.	+	+	-	Th	Mes	Dec	-	D	L	W	S
52.	<i>Bromus tectorum</i> L.	+	+	-	Th	Mes	Dec	-	D	L	W	S
53.	<i>Calamagrostis emodensis</i> Griseb.	+	-	-	G	Mes	E	-	M	L	W	S
54.	<i>Chrysopogon echinulatus</i> (Nees ex Steud.) W. Wats	-	+	-	Th	Mes	E	-	D	L	W	S
55.	<i>Cynodon dactylon</i> (L.) Pers.	+	+	+	H	N	Dec	-	D	L	W	S
56.	<i>Dactylis glomerata</i> L.	+	-	+	Th	N	Dec	-	D	L	W	S
57.	<i>Dichanthium annulatum</i> (Forssk.) Stapf.	-	-	+	H	N	Dec	-	M	L	W	S
58.	<i>Echinochloa crusgalli</i> (L.) P. Beauv.	+	-	+	Th	N	Dec	-	M	L	W	S
59.	<i>Eragrostis poaeoides</i> P. Beave.	+	+	+	H	N	Dec	-	M	L	W	S
60.	<i>Hordeum murinum</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
61.	<i>Hordeum vulgare</i> L.	+	+	+	Th	N	Dec	-	D	L	C	S
62.	<i>Koeleria gracilis</i> Pers.	-	-	+	H	N	Dec	-	D	L	W	S
63.	<i>Lolium rigidum</i> Gaud.	+	+	-	Th	N	Dec	-	D	L	W	S
64.	<i>Lolium temulentum</i> L.	+	-	-	Th	N	Dec	-	D	L	W	S
65.	<i>Melica inaequiglumis</i> (Boiss.) Bor.	+	-	-	H	N	Dec	-	D	L	W	S
66.	<i>Oryza sativa</i> L.	-	+	+	G	Mic	Dec	-	M	L	C	S
67.	<i>Panicum antidotale</i> Retz.	-	-	+	Th	N	Dec	-	D	L	W	S
68.	<i>Phleum pretense</i> L.	+	-	-	Th	N	Dec	-	D	L	W	S
69.	<i>Phragmites karka</i> (Retz.) Trin. ex Steud	+	+	-	G	Mac	Dec	-	M	L	W	S
70.	<i>Piptatherum wendelboi</i> Bor	-	+	-	Th	N	Dec	-	D	L	W	S
71.	<i>Poa annua</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
72.	<i>Poa bulbosa</i> L.	-	-	+	Th	N	Dec	-	D	L	W	S
73.	<i>Poa pratensis</i> L.	-	-	+	H	N	Dec	-	D	L	W	S
74.	<i>Poa supina</i> Schrad.	-	+	-	Th	N	Dec	-	D	L	W	S
75.	<i>Polypogon fugax</i> Nees ex Steud	-	-	+	Th	Mic	Dec	-	M	L	W	S
76.	<i>Saccharum spontaneum</i> L.	+	-	-	Ch	Mac	Dec	-	D	L	W	S
77.	<i>Setaria viridis</i> (L.) P. Beauv.	+	-	-	Th	N	Dec	-	M	L	W	S
78.	<i>Triticum aestivum</i> L.	+	+	+	Th	Mes	Dec	-	D	L	C	S
79.	<i>Vulpia myuros</i> (L.) Gmel.	-	+	+	Th	N	Dec	-	D	L	W	S
80.	<i>Zea mays</i> L.	+	+	+	Th	Mac	Dec	-	D	L	C	S
26. Family Typhaceae												
81.	<i>Typha angustata</i> Bory & Chaub	-	+	+	G	Mac	E	-	M	L	W	S
DICOTYLEDONAE												
27. Family Aceraceae												
82.	<i>Acer pentapomicum</i> J.L. Stewart	-	+	+	Meg P	Mic	E	-	D	L	W	Com

28. Family Amaranthaceae												
83.	<i>Amaranthus viridis</i> L.	+	+	+	Th	Mes	Dec	-	D	L	W	S
29. Family Anacardiaceae												
84.	<i>Pistacia integerrima</i> (L.) Steward ex Brandis	+	+	+	NP	Mes	Dec	-	D	L	W	S
85.	<i>Pistacia khinjuk</i> Stocks	+	-	+	NP	N	Dec	-	D	L	W	S
86.	<i>Rhus punjabensis</i> Steward ex Brandis	+	-	+	H	N	Dec	-	D	L	W	S
30. Family Apiaceae												
87.	<i>Ammi visnaga</i> (L.) Lamk.	+	-	-	Th	N	Dec	-	D	L	W	S
88.	<i>Bunium persicum</i> (Boiss.) B. Fedtsch.	+	+	+	G	N	Dec	-	D	L	W	S
89.	<i>Bupleurum exaltatum</i> Clarke	+	+	-	G	N	Dec	-	D	L	W	S
90.	<i>Bupleurum gilesii</i> Wolff	-	+	-	G	N	Dec	-	D	L	W	S
91.	<i>Carum capticum</i> L.	+	+	+	Th	N	Dec	-	D	L	W	Dis.
92.	<i>Carum carvi</i> L.	+	+	+	Th	N	Dec	-	D	L	W	Dis.
93.	<i>Coriandrum sativum</i> L.	-	-	+	Th	N	Dec	-	D	L	C	Dis.
94.	<i>Daucus carota</i> L.	+	+	+	G	Mes	Dec	-	D	L	C	Dis.
95.	<i>Foeniculum vulgare</i> Mill.	+	+	+	Th	N	Dec	-	D	L	C	Dis.
96.	<i>Pimpinella stewartii</i> (Dunn) Wolff.	-	-	+	Th	N	Dec	-	D	L	W	Com
97.	<i>Pimpinella diversifolia</i> DC.	+	+	-	Th	N	Dec	-	D	L	W	Com
98.	<i>Prangos pabularia</i> Lindl.	+	-	-	H	N	Dec	-	D	L	W	Com
99.	<i>Trachyspermum ammi</i> (L.) Sprague	-	+	+	G	N	Dec	-	D	L	W	Com
31. Family Apocynaceae												
100.	<i>Vinca major</i> L.	+	-	-	G	Mic	E	-	D	L	C	S
32. Family Asclepiadaceae												
101.	<i>Cynanchum acutum</i> L.	+	+	+	Ch	Mes	Dec	-	D	L	W	S
102.	<i>Periploca aphylla</i> Dcne.	-	-	+	Ch	Ap	E	-	D	L	W	Ap
33. Family Asteraceae												
103.	<i>Anaphalis contorta</i> L.	+	+	+	H	Mes	Dec	-	D	L	W	S
104.	<i>Anaphalis nepalensis</i> (Spreng.) Hand.	+	+	+	H	Mes	Dec	-	D	L	W	S
105.	<i>Artemisia brevifolia</i> Wall. Ex DC.	+	+	+	H	L	Dec	-	D	L	W	Dis.
106.	<i>Artemisia maritima</i> L.	+	+	-	H	L	Dec	-	D	L	W	Dis.
107.	<i>Artemisia parviflora</i> Roxb.	+	+	+	Th	Mic	Dec	-	D	L	W	Dis.
108.	<i>Artemisia sacrorum</i> Ledeb.	+	+	-	Th	Mic	Dec	-	D	L	W	Dis.
109.	<i>Artemisia scoparia</i> Waldst. & Kit.	+	+	+	H	N	Dec	-	D	L	W	Dis.
110.	<i>Aster flaccidus</i> var. <i>flaccidus</i> Bunge	-	-	+	Th	N	Dec	-	D	L	W	S
111.	<i>Bellis perennis</i> L.	+	-	-	Th	N	Dec	-	D	L	W	S
112.	<i>Bidens tripartita</i> L.	-	-	+	Th	Mes	Dec	-	D	L	W	Dis.
113.	<i>Calendula officinalis</i> L.	+	-	-	Th	Mes	Dec	-	D	L	C	S
114.	<i>Carduus edelbergii</i> Rech.	+	-	+	Th	Mes	Dec	+	D	L	W	Dis.
115.	<i>Centaurea iberica</i> Trev. ex Spreng.	+	+	+	Th	N	Dec	+	D	L	W	Dis.
116.	<i>Cichorium intybus</i> L.	+	+	+	Th	Mes	Dec	-	D	L	W	S
117.	<i>Cirsium arvense</i> (L.) Scop.	+	+	+	Th	Mac	Dec	+	D	L	W	S
118.	<i>Cirsium falconeri</i> (Hk. f.) Petrak	-	-	+	Th	Mac	Dec	+	D	L	W	S
119.	<i>Conyza bonariensis</i> (L.) Cronquist	+	+	-	Th	N	Dec	-	D	L	W	S
120.	<i>Conyza canadensis</i> (L.) Cronquist	+	+	+	Th	N	Dec	-	D	L	W	S
121.	<i>Conyza japonica</i> Less.	+	+	+	Th	N	Dec	-	D	L	W	S
122.	<i>Cousinia buphtalmoides</i> Regel & Schm.	+	-	-	Th	L	Dec	-	D	L	W	S
123.	<i>Cousinia mattfeldii</i> Bornm.	+	-	-	Th	Mic	Dec	+	D	L	W	S
124.	<i>Cousinia racemosa</i> Boiss.	+	-	-	Th	L	Dec	+	D	L	W	S
125.	<i>Echinops cornigerus</i> DC.	+	+	+	Th	Mes	Dec	+	D	L	W	S
126.	<i>Erigeron canadensis</i> L.	+	-	+	Th	Mic	Dec	-	D	L	W	S

127.	<i>Filago arvensis</i> L.	-	-	+	Th	N	Dec	-	D	L	W	S
128.	<i>Filago desertorum</i> Pomel	-	-	+	Th	N	Dec	-	D	L	W	S
129.	<i>Galinsoga parviflora</i> Cav.	+	-	+	Th	N	Dec	-	D	L	W	S
130.	<i>Inula rhizocephala</i> Schrenk Enum.	-	+	+	Th	N	Dec	-	D	L	W	S
131.	<i>Lachnophyllum gossypinum</i> Bunge.	-	-	+	Th	Mic	Dec	+	D	L	W	S
132.	<i>Lactuca decipiens</i> (H. & T.) Clarke	-	-	+	Th	N	Dec	-	D	L	W	S
133.	<i>Lactuca sativa</i> L.	+	+	+	Th	Mac	Dec	-	D	L	C	S
134.	<i>Lactuca persica</i> Boiss.	+	+	-	Th	N	Dec	-	D	L	W	S
135.	<i>Lactuca serriola</i> L.	+	+	+	Th	Mes	Dec	-	D	L	W	S
136.	<i>Luctuca dolichophylla</i> Kitam.	-	-	+	Th	N	Dec	-	D	L	W	S
137.	<i>Matricaria aurea</i> L.	+	+	+	Th	N	Dec	-	D	L	W	Dis.
138.	<i>Matricaria disciformis</i> (C.A.M.) DC. Prodr.	+	+	+	Th	N	Dec	-	D	L	W	Dis.
139.	<i>Pentanema indicum</i> (L.) Y. Ling	-	+	-	Th	N	Dec	-	D	L	W	S
140.	<i>Phagnalon acuminatum</i> Boiss.	+	+	-	Th	L	Dec	-	D	L	W	S
141.	<i>Pulicaria salviaefolia</i> Bunge	+	-	+	H	N	Dec	-	D	L	W	S
142.	<i>Saussurea heteromalla</i> (D. Don.) Hannd.	-	+	-	Th	Mic	Dec	-	D	L	W	S
143.	<i>Scorzonera virgata</i> DC	+	-	-	H	L	Dec	-	D	L	W	Dis.
144.	<i>Sonchus arvensis</i> Boiss.	+	+	+	Th	Mes	Dec	-	M	L	W	Dis.
145.	<i>Sonchus asper</i> (L.) Hill	+	+	+	Th	Mes	Dec	-	M	L	W	Dis.
146.	<i>Sonchus oleraceus</i> L.	+	+	+	Th	N	Dec	-	M	L	W	Dis.
147.	<i>Senecio dubitabilis</i> C. Jeffery & Y.L. Chen	+	-	-	Th	N	Dec	-	M	L	W	S
148.	<i>Tagetes erecta</i> L.	+	+	-	Th	N	Dec	-	D	L	C	Dis.
149.	<i>Taraxacum officinale</i> Weber	+	+	+	G	Mes	Dec	-	M	L	W	S
150.	<i>Tragopogon gracilis</i> D. Don.	-	+	-	H	N	Dec	-	D	L	W	S
151.	<i>Xanthium strumarium</i> L.	+	+	+	Th	Mes	Dec	+	D	L	W	S

34. Family Berberidaceae

152.	<i>Berberis lycium</i> Royle	-	-	+	NP	N	Dec	+	D	L	W	S
------	------------------------------	---	---	---	----	---	-----	---	---	---	---	---

35. Family Balsaminaceae

153.	<i>Impatiens bicolor</i> Royle	-	+	+	Th	Mic	Dec	-	M	Sd	W	S
154.	<i>Impatiens edgeworthii</i> Hk. f. FBI	-	+	+	Th	Mic	Dec	-	M	Sd	W	S

36. Family Bignoniaceae

155.	<i>Campsis grandiflora</i> L.	+	-	-	Ch (Cl)	Mes	E	-	D	L	C	S
------	-------------------------------	---	---	---	------------	-----	---	---	---	---	---	---

37. Family Boraginaceae

156.	<i>Arnebia decumbens</i> (Vent) Coss & Kral.	-	-	+	H	N	Dec	-	D	L	W	S
157.	<i>Asperugo procumbens</i> L.	+	-	-	Th	Mic	Dec	-	D	L	W	S
158.	<i>Cynoglossum glochidiatum</i> Wall. ex Benth.	+	+	-	H	N	Dec	-	D	L	W	S
159.	<i>Cynoglossum lanceolatum</i> Forssk.	+	+	-	H	N	Dec	-	D	L	W	S
160.	<i>Lithospermum arvense</i> L.	+	-	-	Th	N	Dec	-	D	L	W	S
161.	<i>Lappula microcarpa</i> (Ledeb.) Gurke	+	+	-	Th	L	Dec	-	D	L	W	S
162.	<i>Lepechinella inconspicua</i> (Brand) Reidl.	+	-	-	H	Mes	Dec	-	D	L	W	S
163.	<i>Lindelofia longiflora</i> (Benth.) Baill.	-	+	+	N	N	Dec	-	D	L	W	S
164.	<i>Onosma dichroanthum</i> Boiss.	+	-	-	H	N	Dec	-	D	L	W	S
165.	<i>Onosma hispidum</i> Wall. ex. G. Don.	+	-	-	H	N	Dec	-	D	L	W	S
166.	<i>Pseudomertensia chitalensis</i> (Riedl) Reidl in Rechinger	-	+	-	H	N	Dec	-	D	L	W	S
167.	<i>Solenanthus circinnatus</i> Ledeb.	-	+	-	H	N	Dec	-	D	L	W	S

38. Family Brassicaceae

168.	<i>Alliaria petiolata</i> (M. B.) Cav. & Grande Boll.	+	-	+	Th	N	Dec	-	D	L	W	S
169.	<i>Alyssum desertorum</i> Stapf.	+	-	+	Th	N	Dec	-	D	L	W	S
170.	<i>Arabidopsis wallichii</i> (H. & T.) N. Busch.	+	-	+	H	N	Dec	-	D	L	W	S

171.	<i>Arabis auriculata</i> Lam.	+	+	+	Th	N	Dec	-	D	L	W	S
172.	<i>Brassica campestris</i> L.	+	+	+	Th	Mac	Dec	-	D	L	C	Dis.
173.	<i>Brassica napus</i> L.	+	+	+	Th	Mac	Dec	-	D	L	C	Dis.
174.	<i>Capsella bursa-pastoris</i> (L.) Medic.	+	+	+	Th	Mes	Dec	-	M	L	W	Dis.
175.	<i>Cardamine hirsuta</i> L.	+	-	+	Th	N	Dec	-	M	L	W	Com
176.	<i>Coronopus didymus</i> (L.) Sm.	+	+	+	Th	L	Dec	-	M	L	W	Dis.
177.	<i>Descurainia sophia</i> (L.) Webb. & Berth.	-	-	+	Th	N	Dec	-	M	L	W	Dis.
178.	<i>Draba lanceolata</i> Royle	+	+	-	H	N	Dec	-	D	L	W	S
179.	<i>Draba altaica</i> (C.A.M.) Bunge	-	+	-	Th	N	Dec	-	D	L	W	S
180.	<i>Draba olgae</i> var. <i>chitralensis</i> O.E. Schulz	-	+	-	Th	N	Dec	-	D	L	W	S
181.	<i>Draba trinervis</i> O.E.S.	-	+	-	Th	N	Dec	-	D	L	W	S
182.	<i>Eruca sativa</i> Mill.	+	-	-	Th	Mic	Dec	-	D	L	W	S
183.	<i>Lepidium sativum</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
184.	<i>Malcolmia africana</i> (L.) R. Br.	+	+	-	Th	L	Dec	-	D	L	W	S
185.	<i>Malcolmia cabulica</i> (Boiss.) H. & T.	-	+	-	Th	N	Dec	-	D	L	W	S
186.	<i>Matthiola tenera</i> Rech.f.,	+	-	+	Ch	Mes	Dec	-	M	L	W	S
187.	<i>Nasturtium officinale</i> R. Br.	+	+	-	G	N	E	-	M	L	W	Dis.
188.	<i>Neslia apiculata</i> Fisch.	+	+	+	Th	L	Dec	-	M	L	W	S
189.	<i>Raphanus sativus</i> L.	+	+	+	Th	Mac	Dec	-	M	L	C	Dis.
190.	<i>Sisymbrium brassiciforme</i> C. A. Mey.	+	-	-	Th	N	Dec	-	D	L	W	Dis.
191.	<i>Sisymbrium irio</i> L.	+	+	+	Th	Mic	Dec	-	M	L	W	Dis.
192.	<i>Thlaspi andersonii</i> (H. & T.) O.E.S.	-	+	-	Th	Mic	Dec	-	D	L	W	S

39. Family Campanulaceae

193.	<i>Campanula staintonii</i> Rech.	-	-	+	Th	N	Dec	-	M	L	W	S
194.	<i>Codonopsis ovata</i> Bth.	+	+	-	H	Mes	Dec	-	M	L	W	S

40. Family Cannabinaceae

195.	<i>Cannabis sativa</i> L.	+	+	+	Th	N	Dec	-	D	L	W	Com
------	---------------------------	---	---	---	----	---	-----	---	---	---	---	-----

41. Family Capparidaceae

196.	<i>Capparis spinosa</i> L.	+	+	-	H	Mes	Dec	+	D	L	W	S
197.	<i>Cleome ariana</i> Hedge & Lamond	+	+	-	Th	N	Dec	-	D	L	W	S

42. Family Caprifoliaceae

198.	<i>Viburnum cotinifolium</i> D. Don.	+	+	-	H	Mac	Dec	-	M	L	W	Com
------	--------------------------------------	---	---	---	---	-----	-----	---	---	---	---	-----

43. Family Caryophyllaceae

199.	<i>Arenaria griffithii</i> Boiss.	+	+	-	Ch	L	E	-	D	L	W	S
200.	<i>Arenaria serpyllifolia</i> L.	+	-	-	Th	N	Dec	-	D	L	W	S
201.	<i>Herniaria hirsuta</i> L.	+	-	+	Th	L	Dec	-	D	L	W	S
202.	<i>Lepydodiscus holosteoides</i> (C. A. Mey.) Fenzl.	+	+	-	Th	N	Dec	-	D	L	W	S
203.	<i>Silene conoidea</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
204.	<i>Silene pseudo-verticillata</i> Nasir	-	-	+	Th	Mic	Dec	-	D	L	W	S
205.	<i>Silene viscosa</i> (L.) Pers.	+	+	+	Th	N	Dec	-	D	L	W	S
206.	<i>Silene vulgaris</i> (Moench) Garcke.	-	+	-	H	N	Dec	-	D	L	W	S
207.	<i>Spergula arvensis</i> L.	-	+	-	Th	L	Dec	-	D	L	W	S
208.	<i>Stellaria media</i> (L.) Cyr.	+	+	+	Th	N	Dec	-	D	L	W	S

44. Family Chenopodiaceae

209.	<i>Atriplex crassifolia</i> C. A. Mey.	+	+	+	Th	N	E	-	D	L	W	S
210.	<i>Chenopodium album</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
211.	<i>Chenopodium ambrosioides</i> L.	+	-	-	Th	L	Dec	-	M	L	W	S
212.	<i>Chenopodium botrys</i> L.	-	-	+	Th	L	Dec	-	D	L	W	S
213.	<i>Chenopodium foliosum</i> (Merrich.) Aschers.	-	-	+	Th	N	Dec	-	D	L	W	S
214.	<i>Chenopodium murale</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
215.	<i>Chenopodium novopokrovskyanum</i> (Aellen) Uotila	-	+	-	Th	N	Dec	-	D	L	W	S
216.	<i>Kochia indica</i> Wight.	+	+	+	Th	L	Dec	-	D	L	W	S

45. Family Convolvulaceae													
217.	<i>Convolvulus arvensis</i> L.	+	+	+	Th (Cl)	Mes	Dec	-	D	L	W	S	
46. Family Crassulaceae													
218.	<i>Sedum ewersii</i> Ledeb.	+	+	-	G	N	Dec	-	M	L	W	S	
219.	<i>Sedum quadrifidum</i> Pall.	-	+	-	G	N	Dec	-	M	L	W	S	
47. Family Cucurbitaceae													
220.	<i>Citrullus vulgaris</i> L.	+	+	+	Th	Mes	Dec	-	D	L	C	Com	
221.	<i>Cucumis sativa</i> L.	+	+	+	Th	Mes	Dec	-	D	L	C	Com	
222.	<i>Cucurbita maxima</i> Duch ex Lam	+	+	+	Th	Mac	Dec	-	D	L	C	Com	
223.	<i>Cucurbita pepo</i> L.	+	+	+	Th	Mac	Dec	-	D	L	C	Com	
48. Family Cuscutaceae													
224.	<i>Cuscuta brevistyla</i> A. Braun. ex A. Rich.	-	-	+	P	Ap	Ap	-	D	L	W	Ap	
225.	<i>Cuscuta lehmanniana</i> Bunge	-	-	+	P	Ap	Ap	-	D	L	W	Ap	
226.	<i>Cuscuta reflexa</i> Roxb.	+	+	+	P	Ap	Ap	-	D	L	W	Ap	
49. Family Dipsacaceae													
227.	<i>Dipsacus inermis</i> var. <i>mitis</i> (D. Don) Y. Nasir	-	+	-	Th	L	Dec	-	D	L	W	S	
50. Family Elaeagnaceae													
228.	<i>Elaeagnus angustifolia</i> L.	+	-	-	Mac P	Mes	Dec	-	D	L	W	S	
51. Family Euphorbiaceae													
229.	<i>Euphorbia granulata</i> L.	-	+	-	Th	N	Dec	-	D	L	W	S	
230.	<i>Euphorbia falcata</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S	
231.	<i>Euphorbia helioscopia</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S	
232.	<i>Euphorbia peplus</i> L.	+	+	+	Th	L	Dec	-	D	L	W	S	
233.	<i>Euphorbia prostrata</i> L.	-	-	+	Th	L	Dec	-	D	L	W	S	
52. Family Fabaceae													
234.	<i>Astragalus grahamianus</i> Royle. ex Bth.	+	-	+	Ch	L	Dec	+	D	L	W	Com	
235.	<i>Astragalus subumbellatus</i> Klotzsch.	+	+	-	H	N	Dec	-	D	L	W	Com	
236.	<i>Astragalus nivales</i> Kar & Kir	+	+	-	H	L	Dec	-	D	L	W	Com	
237.	<i>Indigofera heterantha</i> var. <i>heterantha</i> L.	-	+	-	Ch	Mic	Dec.	-	D	L	W	Com	
238.	<i>Lathyrus aphaca</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S	
239.	<i>Lotus corniculatus</i> L.	+	+	+	H	N	Dec	-	D	L	W	S	
240.	<i>Medicago lupulina</i> L.	+	-	-	Th	N	Dec	-	D	L	W	Com	
241.	<i>Medicago polymorpha</i> L.	+	+	+	Th	N	Dec	-	D	L	W	Com	
242.	<i>Melilotus indica</i> (L.) All.	+	-	+	Th	N	Dec	-	D	L	W	Com	
243.	<i>Robinia pseudoacacia</i> L.	+	-	+	NP	N	E	+	D	L	C	Com	
244.	<i>Sophora mollis</i> (Royle) Baker	+	-	+	NP	N	Dec	+	D	L	W	S	
245.	<i>Trifolium pratense</i> L.	+	+	+	G	N	Dec	-	M	L	W	S	
246.	<i>Trifolium repens</i> L.	+	+	+	G	N	Dec	-	M	L	W	S	
247.	<i>Trigonella incisa</i> Bth.	+	+	+	Th	N	Dec	-	M	L	W	S	
248.	<i>Vicia monentha</i> L.	+	+	+	Th (Cl)	N	Dec	-	M	L	W	S	
249.	<i>Vicia sativa</i> L.	+	+	+	Th (Cl)	N	Dec	-	M	L	W	S	
53. Family Fumariaceae													
250.	<i>Fumaria indica</i> (Hausskn.) H. N. Pugsley	+	+	+	Th	N	Dec	-	D	L	W	Dis.	
54. Family Fagaceae													
251.	<i>Quercus baloot</i> Griffith	+	+	+	NP	Mic	E	+	D	L	W	Com	
252.	<i>Quercus dilatata</i> Lindl.	+	+	+	Meg P	Mic	E	-	D	L	W	Com	

55. Family Geraniaceae													
253.	<i>Erodium cicutarium</i> (L.) L' Herit.	+	-	-	Th	Mic	Dec	-	D	L	W	Com	
254.	<i>Geranium rotundifolium</i> L.	+	+	+	Th	N	Dec	-	M	L	W	Com	
255.	<i>Geranium wallichianum</i> D. Don. Ex Sweet	+	-	-	Th	Mes	Dec	-	M	L	W	Com	
56. Family Juglandaceae													
256.	<i>Juglans regia</i> L.	+	+	+	Meg P	Mac	Dec	-	D	L	W	S	
57. Family Lamiaceae													
257.	<i>Calamintha umbrosa</i> (M. Bieb.) Fisch. & Mey.	+	+	-	Th	N	Dec	-	D	L	W	S	
258.	<i>Calamintha vulgaris</i> (L.) Druce	+	+	-	Th	N	Dec	-	D	L	W	S	
259.	<i>Lagocheirus cabulicus</i> Bth. in DC. Prodr.	+	+	-	Th	N	Dec	-	D	L	W	S	
260.	<i>Lallemantia royleana</i> (Bth.) Bth.	-	+	-	Th	L	Dec	-	D	L	W	S	
261.	<i>Lamium amplexicaule</i> L.	+	+	+	Th	N	Dec	-	M	L	W	S	
262.	<i>Marrubium vulgare</i> L.	+	-	-	Ch	Mes	Dec	-	D	L	W	S	
263.	<i>Mentha longifolia</i> (L.) Huds	+	+	+	G	Mes	Dec	-	M	L	W	S	
264.	<i>Mentha piperata</i> Benth.	+	+	+	G	N	Dec	-	D	L	W	S	
265.	<i>Nepeta brachyantha</i> Rech.	+	-	+	Ch	Mes	Dec	-	D	L	W	S	
266.	<i>Nepeta clerkei</i> Hk. f.	-	-	+	Th	Mic	Dec	-	M	L	W	S	
267.	<i>Nepeta raphanorhiza</i> Bth.	-	+	+	Th	N	Dec	-	D	L	W	S	
268.	<i>Nepeta sewerzowii</i> Regel	-	+	+	Th	N	Dec	-	D	L	W	S	
269.	<i>Ocimum basilicum</i> L.	+	+	+	Ch	N	Dec	-	D	L	C	S	
270.	<i>Phlomis cashmeriana</i> Royle	+	+	+	Th	N	Dec	-	D	L	W	S	
271.	<i>Prunella vulgaris</i> L.	+	+	-	Th	N	Dec	-	D	L	W	S	
272.	<i>Salvia aegyptiaca</i> L.	-	+	+	Ch	N	Dec	-	D	L	W	S	
273.	<i>Salvia moorcroftiana</i> Wall. ex Bth.	+	-	-	Th	Mac	Dec	-	D	L	W	S	
274.	<i>Salvia nubicola</i> wall. ex Sweet	+	+	-	Ch	Mes	Dec	-	D	L	W	S	
275.	<i>Salvia plebeia</i> R. Br.	+	+	+	Th	N	Dec	-	D	L	W	S	
276.	<i>Scutellaria nuristanica</i> Rech.	-	-	+	H	N	Dec	-	D	L	W	S	
277.	<i>Thymus linearis</i> L.	+	+	+	H	N	Dec	-	D	L	W	S	
278.	<i>Ziziphora tenuior</i> L.	-	-	+	Th	L	Dec	-	D	L	W	S	
58. Family Malvaceae													
279.	<i>Abelmoschus esculentus</i> (L.) Moench Meth.	+	+	+	Th	Mes	Dec	-	D	L	C	Dis.	
280.	<i>Althaea rosea</i> (L.) Cav. Diss.	+	-	-	H	Mac	Dec	-	D	L	C	Com	
281.	<i>Malva neglecta</i> Wallr.	+	-	-	Th	Mes	Dec	-	D	L	W	Com	
282.	<i>Malva peruviana</i> L.	+	-	-	Th	N	Dec	-	D	L	W	S	
59. Family Moraceae													
283.	<i>Ficus carica</i> L.	+	+	+	Mac P	Mac	Dec	-	D	L	W	S	
284.	<i>Morus alba</i> L.	+	+	+	Meg P	Mac	Dec	-	D	L	C	S	
285.	<i>Morus nigra</i> L.	+	+	+	Meg P	Mac	Dec	-	D	L	C	S	
60. Family Oleaceae													
286.	<i>Fraxinus xanthoxyloides</i> (Wall. ex G. Don) DC.	+	+	+	NP	Mic	Dec	-	D	L	W	Com	
61. Family Onagraceae													
287.	<i>Epilobium hirsutum</i> L.	+	+	+	Th	N	Dec	-	M	L	W	S	
288.	<i>Epilobium latifolium</i> L.	-	-	+	Th	Mes	Dec	-	M	L	W	S	
289.	<i>Epilobium royleanum</i> Hausskn.	-	+	+	Th	N	Dec	-	M	L	W	S	
62. Family Oxalidaceae													
290.	<i>Oxalis corniculata</i> L.	+	+	+	Th	Mes	Dec	-	M	L	W	S	
63. Family Paeoniaceae													
291.	<i>Paeonia emodi</i> Wall ex G. Don	-	+	-	G	N	Dec	-	M	Sd	W	Com	
64. Family Papaveraceae													
292.	<i>Papaver dubium</i> L.	-	-	+	Th	L	Dec	-	D	L	W	Dis	

65. Family Plantaginaceae												
293.	<i>Plantago lanceolata</i> L.	+	+	+	Th	Mes	Dec	-	D	L	W	S
294.	<i>Plantago major</i> Aitch.	+	+	+	G	Mac	Dec	-	M	L	W	S
295.	<i>Plantago ovata</i> Forssk.	-	-	+	Th	Mic	Dec	-	M	L	W	S
66. Family Platanaceae												
296.	<i>Platanus orientalis</i> L.	+	+	+	Meg P	Mac	Dec	-	D	L	C	Com
67. Family Plumbaginaceae												
297.	<i>Acantholimon lycopodioides</i> (Girard) Boiss.	-	-	+	Ch	L	Dec	-	D	L	W	S
298.	<i>Limonium griffithii</i> (Aitch. & Hemsl.) O. Ktze.	-	+	+	H	Mic	Dec	-	D	L	W	S
68. Family Polygonaceae												
299.	<i>Bistorta affinis</i> (D. Don) Green	-	-	+	Ch	N	Dec	-	D	L	W	S
300.	<i>Bistorta vivipara</i> (L.) S.F. Gray	-	-	+	H	N	Dec	-	D	L	W	S
301.	<i>Oxyria digyna</i> (L.) Hill.	+	+	+	Th	N	Dec	-	M	L	W	S
302.	<i>Polygonum amplexicaule</i> D Don.	-	+	+	Th	L	Dec	-	D	L	W	S
303.	<i>Polygonum aviculare</i> L.	-	-	+	Th	L	Dec	-	D	L	W	S
304.	<i>Polygonum convolvulus</i> L.	-	+	+	Th	L	Dec	-	D	L	W	S
305.	<i>Polygonum paronychioides</i> C,A, Mey. ex Hohen.	-	+	+	H	L	Dec	-	D	L	W	S
306.	<i>Polygonum nepalense</i> Meissn.	+	+	+	G	N	Dec	-	D	L	W	S
307.	<i>Rheum webbianum</i> Royle	+	-	-	G	Mac	Dec	-	D	L	W	Com
308.	<i>Rumex hastatus</i> D. Don	+	+	+	Ch	Mes	Dec	-	D	L	W	S
309.	<i>Rumex longifolius</i> DC	+	+	+	Th	Mes	Dec	-	D	L	W	S
310.	<i>Rumex nepalensis</i> Spreng	+	+	+	H	Mes	Dec	-	D	L	W	S
69. Family Portulacaceae												
311.	<i>Portulaca oleracea</i> L.	+	+	+	Th	N	Dec	-	D	L	W	S
70. Family Primulaceae												
312.	<i>Anagallis arvensis</i> L.	+	+	+	Th	L	Dec	-	D	L	W	S
313.	<i>Androsace harrissii</i> Du.	-	-	+	H	N	Dec	-	M	L	W	S
314.	<i>Primula denticulata</i> Sm.	-	-	+	H	N	Dec	-	M	L	W	S
315.	<i>Primula elliptica</i> Royle	-	+	+	H	N	Dec	-	M	L	W	S
71. Family Punicaceae												
316.	<i>Punica granatum</i> L.	+	+	+	NP	Mes	Dec	-	D	L	C	S
72. Family Ranunculaceae												
317.	<i>Adonis aestivalis</i> L.	-	-	+	Th	L	Dec	-	D	L	W	S
318.	<i>Ceratocephalus falcatus</i> (L.) Pers.	-	-	+	Th	Mic	Dec	-	D	L	W	S
319.	<i>Clematis orientalis</i> L.	+	-	+	NP (Cl)	N	Dec	-	D	L	W	Com
320.	<i>Delphinium denudatum</i> Wall. ex H. & T	+	+	-	Ch	Mic	Dec	-	D	L	W	Dis
321.	<i>Thalictrum elegans</i> Wall. ex Royle	+	-	-	Th	N	Dec	-	M	L	W	Dis.
322.	<i>Ranunculus aucheri</i> L.	+	+	+	Th	N	Dec	-	M	L	W	Dis.
323.	<i>Ranunculus lanata</i> L.	+	+	+	Th	N	Dec	-	M	L	W	Dis.
324.	<i>Ranunculus muricatus</i> L.	+	+	-	Th	N	Dec	-	M	L	W	Dis.
325.	<i>Ranunculus sceleratus</i> L.	-	+	-	Th	N	Dec	-	M	L	W	Dis.
73. Family Rosaceae												
326.	<i>Cotoneaster microphylla</i> Wall. ex Lindl.	+	+	-	NP	L	Dec	+	D	L	W	S
327.	<i>Cotoneaster nummularia</i> Fisch. & Mey	+	+	-	NP	N	Dec	+	D	L	W	S
328.	<i>Crataegus songarica</i> C. Koch.	+	-	-	Meg P	Mes	Dec	-	D	L	C	Com
329.	<i>Cydonia oblonga</i> Mill.	+	-	-	NP	Mac	Dec	-	D	L	C	S
330.	<i>Filipendula vestita</i> (Wall. ex G. Don) Maxim.	-	+	+	NP	Mic	Dec	-	D	L	W	S
331.	<i>Potentilla pamirica</i> Wolf.	-	-	+	Th	N	Dec	-	D	L	W	S
332.	<i>Potentilla sericea</i> L.	+	+	-	Th	L	Dec	-	D	L	W	S

333.	<i>Prunus armeniaca</i> L.	+	+	+	Meg P	Mes	Dec	-	D	L	C	S
334.	<i>Prunus domestica</i> L.	+	-	-	Meg P	Mes	Dec	-	D	L	C	S
335.	<i>Prunus griffithii</i> (Boiss.) C. K. Sch.	-	-	+	NP	Mes	Dec	-	D	L	W	Dis.
336.	<i>Prunus kuramica</i> (Korsh.) Kitam.	+	+	-	Meg P	Mic	Dec	-	D	L	W	S
337.	<i>Prunus persica</i> L.	+	+	+	Mes P	Mes	Dec	-	D	L	C	S
338.	<i>Pyrus communis</i> L.	+	+	+	Meg P	Mac	Dec	-	D	L	C	S
339.	<i>Pyrus pashia</i> Ham. ex D. Don	-	-	+	Mes P	Mic	Dec	-	D	L	W	S
340.	<i>Pyrus malus</i> L.	+	+	+	Meg P	Mac	Dec	-	D	L	C	S
341.	<i>Rosa alba</i> L.	+	+	+	NP	Mic	Dec	+	D	L	C	Com
342.	<i>Rosa ecae</i> Aitch.	+	-	-	NP	N	Dec	+	D	L	W	Com
343.	<i>Rosa webbiana</i> Wall. ex Royle	+	+	+	NP	N	Dec	+	D	L	W	Com
344.	<i>Rubus fruticosus</i> L.	+	-	+	NP	Mic	Dec	-	D	L	W	Com
345.	<i>Rubus irritans</i> Focke	+	-	+	NP	Mic	Dec	-	D	L	W	Com
346.	<i>Spiraea canescens</i> Lour.	+	+	-	NP	N	Dec	-	D	L	W	S
347.	<i>Sorbaria tomentosa</i> (Lindl.) Rehder	+	+	-	Ch	Mic	Dec	-	D	L	W	Com

74. Family Rubiaceae

348.	<i>Galium aparine</i> L.	+	+	+	Th (Cl)	N	Dec	-	D	L	W	S
349.	<i>Galium tricornutum</i> L.	+	+	+	Th (Cl)	N	Dec	-	D	L	W	S
350.	<i>Rubia cordifolia</i> L.	+	+	-	Th	N	Dec	-	D	L	W	S

75. Family Salicaceae

351.	<i>Populus alba</i> L.	-	+	+	Mac P	Mes	Dec	-	M	L	C	S
352.	<i>Populus nigra</i> L.	+	+	+	Mac P	Mac	Dec	-	M	L	C	S
353.	<i>Salix acmophylla</i> Boiss.	+	+	+	Mes P	Mes	Dec	-	D	L	C	S
354.	<i>Salix linearifolia</i> E. Wolf.	+	-	-	Mes P	Mes	Dec	-	D	L	C	S
355.	<i>Salix tetrasperma</i> Roxb.	+	+	+	Meg P	Mes	Dec	-	D	L	C	S
356.	<i>Salix denticulata</i> Anderson subsp. <i>denticulata</i>	+	+	-	Mes P	Mes	Dec	-	D	L	C	S

76. Family Saxifragaceae

357.	<i>Bergenia ciliata</i> (Haw.) Sternb.	+	-	+	G	Mic	E	-	M	Sd	W	S
358.	<i>Bergenia himalaica</i> Boriss	+	-	+	G	Mic	E	-	M	Sd	W	S

77. Family Scrophulariaceae

359.	<i>Euphrasia aristulata</i> Penn.	+	-	+	Th	L	Dec	-	D	L	W	S
360.	<i>Kickxia ramossissima</i> (Wall.) Janchen	-	-	+	Th	L	Dec	-	M	Sd	W	S
361.	<i>Linaria bamianica</i> Patzak	+	+	+	Th	L	Dec	-	D	L	W	S
362.	<i>Pedicularis chitralensis</i> Penn.	-	+	-	H (P)	L	Dec	-	D	L	W	S
363.	<i>Scrophularia robusta</i> Penn.	-	-	+	Th	Mic	Dec	-	D	L	W	S
364.	<i>Scrophularia scoraria</i> Penn.	-	-	+	Th	Mic	Dec	-	D	L	W	S
365.	<i>Verbascum thapsus</i> L.	+	+	+	G	Mes	Dec	-	D	L	W	S
366.	<i>Veronica anagallis aquatica</i> L.	+	+	+	G	Mes	Dec	-	M	L	W	Com
367.	<i>Veronica persica</i> L.	+	-	-	Th	L	Dec	-	M	L	W	S
368.	<i>Veronica verna</i> L.	+	-	-	Th	Mic	Dec	-	M	L	W	Com

78. Family Simaroubaceae

369.	<i>Ailanthus altissima</i> (Mill.) Swingle	+	+	+	NP	Mic	Dec	-	M	L	W	Com
------	--	---	---	---	----	-----	-----	---	---	---	---	-----

79. Family Solanaceae												
370.	<i>Datura stramonium</i> L.	+	+	+	Th	Mic	Dec	-	D	L	W	S
371.	<i>Lycopersicon esculentum</i> Mill.	+	+	+	Th	Mic	Dec	-	D	L	C	Com
372.	<i>Solanum melongena</i> L.	+	+	+	Th	Mac	Dec	-	D	L	C	S
373.	<i>Solanum nigrum</i> L.	+	+	+	Th	Mes	Dec	-	M	L	W	S
374.	<i>Solanum tuberosum</i> L.	+	+	+	G	Mes	Dec	-	D	L	C	Com
80. Family Tamaricaceae												
375.	<i>Myricaria squamosa</i> Desv.	+	-	-	NP	L	Dec	-	M	L	W	S
376.	<i>Tamarix dioica</i> Roxb. ex Roth	+	+	+	NP	L	Dec	-	M	L	W	S
81. Family Thymelaeaceae												
377.	<i>Daphne oleoides</i> Schreb.	-	+	-	NP	L	E	-	M	L	W	S
378.	<i>Thymelaea passerina</i> (L.) Coss. & Germ.	+	-	-	Th	N	Dec	-	D	L	W	S
82. Family Ulmaceae												
379.	<i>Celtis australis</i> L.	+	-	-	Mac P	Mes	Dec	-	D	L	W	Com
83. Family Urticaceae												
380.	<i>Parietaria debilis</i> Forst.	-	+	-	Th	N	Dec	-	D	L	W	Com
381.	<i>Parietaria judaica</i> Starand.	-	+	-	Th	N	Dec	-	D	L	W	Com
382.	<i>Urtica dioica</i> L.	+	+	+	Th	Mac	Dec	-	D	L	W	Com
84. Family Valerianaceae												
383.	<i>Valeriana ficariifolia</i> Boiss.	+	-	-	Th	N	Dec	-	D	L	W	S
85. Family Verbenaceae												
384.	<i>Duranta repens</i> L.	+	-	-	NP	Mes	Dec	-	D	L	C	Com
385.	<i>Verbena officinalis</i> L.	+	+	-	Th	Mes	Dec	-	D	L	W	S
86. Family Violaceae												
386.	<i>Viola canescens</i> Wall. ex Roxb.	+	+	+	G	Mic	Dec	-	M	L	W	S
87. Family Vitaceae												
387.	<i>Vitis vinifera</i> L.	+	+	+	NP	Mac	Dec	-	D	L	C	Com
88. Family Zygophyllaceae												
388.	<i>Peganum harmala</i> L.	-	-	+	H	N	Dec	-	D	L	W	S
389.	<i>Tribulus terrestris</i> L.	+	-	+	Th	N	Dec	-	D	L	W	S

Keys: B = Bumburet, R = Rombor, BR = Birir

(1) Life form: Th. Therophytes, G. Geophytes, H. Hemicryptophytes, Ch. Chamaephytes, NP. Nanophanerophytes, MacP. Macrophanerophytes, MesP. Mesophanerophytes, MegP. Megaphanerophytes, Cl. Climber, P. Parasite.

(2) Leaf Size: AP. Aphyllous, L. Leptophyll, N. Nanophyll, Mic. Microphyll, Mes. Mesophyll, Mac. Macrophyll, Meg. Megaphyll.

(3) Leaf Persistence: E. Evergreen, Dec. Deciduous

(4) Spininess: + Spiny, – Non spiny

(5) Habitat: M. Moist, D. Dry

(6) Light requirement: L. light, Sd. Shade

(7) Habit: C. Cultivated, W. Wild

(8) Leaf appearance: S. Simple, Com. Compound, Disec. Dissected

Appendix




Fig. 3. Graphical presentation of total taxa belong to specific plant group




Fig. 4. Graphical presentations of total genera belong to specific plant group




Fig. 5. Graphical presentation of total no. of families belongs to specific plant group




Fig. 6. Graphical presentation of no. of species in different life form




Fig. 7. Graphical presentation of no. of species in different leaf size classes




Fig. 8. Graphical presentation of no. of species on the basis of leaf persistence




Fig. 9: Graphical presentation of spininess of plants




Fig. 10. Graphical presentation of habitat of plants




Fig. 11. Graphical presentation of light requirement of plants




Fig. 12. Graphical presentation of nature of plants




Fig. 13. Graphical presentation of leaf shapes of plants

Pakistan is naturally gifted with diverse flora containing more than 6000 identified vascular plants. The country has mangrove forests at zero feet altitude in Arabian Sea to highest peaks and forests in Hindukush-Karakoram-Himalayan ranges. District Chitral is located to extreme Northwest of Pakistan with diverse plant resources. Kalash valley has international fame due to its inhabitants considered as the descendants of Alexander the Great. The Kalash people have their own way of life and mostly depend on plants and their products for daily life needs. In the present study the unexplored floristic wealth of the Kalash valley was documented along with their ecological characteristics. The study showed that the conservation status of plants is alarming and most of them are vulnerable as the area is under heavy anthropogenic pressure due to deforestation

for timber and fuel wood utilization, unscientific medicinal plants collection and overgrazing by livestock. These factors have severely damaged the habitats and causing soil erosion. The drastic changes in the climatic conditions during the last few years have resulted in continuous floods and caused tremendous loses to plant habitats, to cultivated farms, infrastructures and to fruit gardens. Perhaps the locals have realized that it is their moral and ethnic duty to save the plants and other natural resources and have felt that they are destroying themselves by destroying natural resources. Therefore, from last few years the young generation is providing volunteer services in controlling of wood smuggling and deforestation in the area.

4. Conclusion

The present findings suggest further extensive and intensive exploration and attention on plant resources of the area specially the members of families Asteraceae, Poaceae, Rosaceae and some gymnosperm families. The present study might also be helpful for the same type of studies of much closed and unexplored Nooristan Province of war affected Afghanistan.

5. Acknowledgement

The present article is a part of Ph.D. dissertation of the first author.

References

- [Abdullahi et al., 2009](#) – *Abdullahi, M.B., S.S. Sanusi, S.D. Abdul., Sawa, F.B.J.* (2009). An assessment of the herbaceous species vegetation of Yankari Game Reserve, Bauchi, Nigeria, *Am-Eur. J. Agric. & Environ Sci.*, 6(1): 20-25.
- [Aelign et al., 2007](#) – *Aelign, A., Teketay D., Yemshaw Y., Edwards, S.* (2007). Diversity and status of regeneration of woody plants on the peninsula of Zegie, northwestern Ethiopia. *Tropical Ecology*, 48(1): 37-49.
- [Ali, 2008](#) – *Ali, S.I.* (2008). The significance of flora with special reference to Pakistan. *Pak. J. Bot.*, 40(30): 967-971.
- [Ali, Nasir, 1989-1991](#) – *Ali, S.I., Nasir, Y.J.* (1989-1991). Flora of Pakistan. Department of Botany, University of Karachi.
- [Ali, Qaiser, 1993-2013](#) – *Ali, S.I., Qaiser, M.* (1993-2013). Flora of Pakistan. Department of Botany, University of Karachi. Pakistan.
- [Anonymous, 1998](#) – Anonymous (1998). District Census Report of Chitral. PCO, Govt. of Pakistan.
- [Badshah et al., 2013](#) – *Badshah, L., Hussain F., Sher, Z.* (2013). Floristic inventory, ecological characteristics and biological spectrum of rangeland, District Tank, Pakistan. *Pak. J. Bot.*, 45(4): 1159-1168.
- [Badshah et al., 2016](#) – Badshah, L., Hussain F., Sher Z. (2016). Floristic inventory, ecological characteristics and biological spectrum of plants of Parachinar, Kurram Agency, Pakistan. *Pak. J. Bot.* 48(4): 1547-1558.
- [Baykal, Atamov, 2016](#) – *Baykal, H., Atamov V.* (2016). Floristic diversity in Bashemsin valley of Kackar mountains national park of Rize. *Pak. J. Bot.*, 48(5): 1871-1876.
- [Cain, Castro, 1959](#) – *Cain, S.A., Castro G.M.* (1959). Manual of Vegetation analysis. Harper, NY. 325 p.
- [Durrani et al., 2005](#) – Durrani, M.J., Hussain F., Rehman S. (2005). Ecological characteristics of plants of Harboi rangelands, Kalat, Pakistan. *J. Trop. & Subtrop. Bot.*, 13(2): 130-138.
- [Durrani et al., 2010](#) – *Durrani, M.J., Razaq A., Muhammad S.G., Hussain F.* (2010). Floristic diversity, ecological, characteristics and ethnobotonical plants of Aghberg rangelands, Balochistan. *Pak. J. Pl. Sci.*, 16 (1): 29-36.
- [Fatna et al., 2014](#) – *Fatna, B., Noury B., Souhila B., Djamel B.* (2014). Research of the eco-floristic data Into the contribution of groups to Halophyte, hammam boughrara area West Algeria. *European Scientific Journal*. 10 (29): 296-307.
- [Guo et al., 2009](#) – *Guo, Q.S., Wang X.F., Bar G., Kang Y., Hong M., Pei S.X., Zhang F.J.* (2009). Life form spectra, leaf character, and hierarchical-synusia structure of vascular plants in *Thuja sutchuehensis*. *PubMed.*, 20 (9): 2057-2062.
- [Hussain, 1989](#) – *Hussain, F.* (1989). Field and laboratory manual of Plant Ecology. UGC, Islamabad.

Hussain et al., 2012 – Hussain, F., Shah S.M., Badshah Lal (2012). Floristic and vegetation diversity of some aquatic habitats of Mastuj Valley, Hindukush region, District Chitral, Pakistan. *Pak. J. Pl. Sci.*, 18(1): 55-68.

Hussain et al., 2015 – Hussain, F., Shah S.M., Badshah L., Durrani M.J. (2015). Diversity and ecological characteristics of flora of Mastuj valley, district Chitral, Hindukush range. *Pakistan. Pak. J. Bot.*, 47(2): 495-510.

Jabeen et al., 2009 – Jabeen, A., Khan M.A., Ahmad M., Zafar M., Ahmad F. (2009). Indigenous uses of economically important Flora of Margallah Hills National Park, Islamabad, Pakistan. *Afric. J. Biotech.*, 8(5): 763-784.

Kar et al., 2010 – Kar, P.K., Biswal A.K., Barik K.L. (2010). Floristic composition and biological spectrum of a grassland community of Rangamatia in the district of Mayurbhanj, Odisha. *J. Curr. Sci.*, 15(2): 465-469.

Khan et al., 2014 – Khan, M.A., Ullah A., Rashid A., Shah S.M., Fida S. (2014). Floristic leaf-size and life form spectra of Asshab Baba Graveyard Chaghar Matti, district Peshawar, Khyber Pakhtunkhwa, Pakistan. *Int. J. Biol. Biotech.*, 11 (1): 167-171.

Malik et al., 2007 – Malik, Z.H., Hussain F., Malik N.Z. (2007). Life form and Leaf Size Spectra of Plant Communities Harboring Ganga Chotti and Bedori Hills during 1999-2000. *International J. Agriculture & Biology*, 9(6): 833-838.

Manhas et al., 2010 – Manhas, R.K., Singh L., Vasistha H.B., Negi M. (2010). Floristic diversity of protected ecosystems of Kandi Region of Punjab, India. *New York Science Journal*, 3(4): 96-103.

Nasir, Ali, 1970–1989 – Nasir, Y.J., Ali S.I. (1970-1989). Flora of Karachi and Islamabad.

Nazir, Malik, 2006 – Nazir, A., Malik Z.H. (2006). Life form and index of similarity of plant communities recorded at Sarsawa Hill District Kotri. *J. Res. Sci.*, 17(1): 27-33.

Osman et al., 2014 - Osman, A.K., Al-Ghamdi F., Bawadekji A. (2014). Floristic diversity and vegetation analysis of Wadi Arar: A typical desert Wadi of the Northern Border region of Saudi Arabia. *Saudi Journal of Biological Sciences*, 21(6): 554-565.

Parswan et al., 2010 – Parswan, K., Mehta J.P., Sobodh (2010). Floristic composition and biological spectrum of vegetation of alpine meadows of Kedarnath: Garhwal Himalaya. *Nature and Sci.*, 8(7): 109-115.

Qin et al., 2012 – Qin, X., Zhang R., Xing F. (2012). A Study on the Flora and vegetation of Cat Dua Island, Northeastern Vietnam. *Pak. J. Bot.*, 44(4): 1229-1232.

Qureshi, Bhatti, 2008 – Qureshi, R., Bhatti G.R. (2008). Diversity of micro-habitats and their plant resources in Nara Desert, Pakistan. *Pak. J. Bot.* 40(3): 979-992.

Raunkiaer, 1934 – Raunkiaer, C. (1934). The life forms of plants and Statistical Plant Geography. Clarendon Press, Oxford.

Saima et al., 2009 – Saima, S., Dasti A.A., Abbas Q., Hussain F. (2009). Floristic diversity during monsoon in Ayubia National Park, District Abbottabad, Pakistan. *Pak. J. Pl. Sci.*, 16(1): 43-50.

Shah, Hussain, 2007 – Shah, S.M., Hussain F. (2007). Ecology of waterlogged habitats around Chamkani, district Peshawar. *Putaj*. 14: 55-64.

Shaheen et al., 2016 – Shaheen, S., Iqbal Z., Ijaz F., Alam J., Rahman I. (2016). Floristic composition, biological spectrum and phenology of Tehsil Havelian, District Abbottabad, Pakistan. *Pak. J. Bot.*, 48(5): 1849-1859.

Shaheen, Qureshi, 2011 – Shaheen, H., Qureshi R.A. (2011). Vegetation types of Sheosar Lake and surrounding landscape in Deosai Plains of North Pakistan, Western Himalayas. *J. Med. Plant Res.*, 5(4): 599-603.

Sher et al., 2014 – Sher, Z., Hussain F., Badshah L. (2014). Biodiversity and ecological characterization of the flora of Gadoon Rangeland, District Swabi, Khyber Pakhtunkhwa, Pakistan. *Iran. J. Bot.*, 20(1): 96-108.

Sher, Khan, 2007 – Sher, Z., Khan Z. (2007). Floristic composition, life form and leaf size spectra of the vegetation of Chagharzai valley, district Buner. *Pak. J. Pl. Sci.*, 13(1): 55-64.

Udayakumar et al., 2011 – Udayakumar, M., Ayyanar M., Sekar T. (2011). Angiosperms, Pachaiyappa's College, Chennai, Tamil Nadu, India, Check List, 7(1): 37-48.

Youcef et al., 2012 – Youcef, B., Lamine M., Hocine B., Rabah M., Ali L., Belhamra M.B., (2012). Diversity of halophyte desert vegetation of the different saline habitats in the valley of Oued Righ, Low Sahara Basin, Algeria. *Res. J. Environ. Earth Sci.*, 4(3): 308-315.