

With the Author's compliments

G B Sowerby

MARINE SHELLS  
OF  
SOUTH AFRICA.

---

(APPENDIX.)

1897



QL  
427  
S4248  
1892  
Acc.  
Moll.

APPENDIX TO  
MARINE SHELLS  
OF  
SOUTH AFRICA

A CATALOGUE OF ALL THE KNOWN SPECIES

WITH

REFERENCES TO FIGURES IN VARIOUS WORKS, DESCRIPTIONS  
OF NEW SPECIES, AND FIGURES OF SUCH AS ARE NEW,  
LITTLE KNOWN, OR HITHERTO UNFIGURED.

Division of Mollusks  
Sectional Library

BY  
G. B. SOWERBY, F.L.S., F.Z.S.

LONDON:  
SOWERBY, 121, FULHAM ROAD, S.W.

1897.



PRINTED BY  
HUTCHINGS AND CROWLEY,  
123, FULHAM ROAD, SOUTH KENSINGTON, S.W.

## PREFACE.

SINCE the publication of "Marine Shells of South Africa" in 1892, a considerable number of additional species has been discovered, some well known as inhabiting other seas, and others new to science. Descriptions without figures of some of these were published in the "Journal of Conchology," 1894. The time has now come for the publication, in the form of an appendix to the original work, of descriptions of the new species, with figures of those hitherto unfigured, together with a list (as complete as possible) of all the species found up to date in South African Waters, and not included in my work of 1892, giving references to figures in well-known works. I include a list of emendations referring to the pages of the 1st Volume, altering some of the generic names in accordance with the nomenclature now in use. These changes are somewhat inconvenient, but they are rendered necessary by the law of priority. The number of additional South African Marine Mollusca enumerated in the present work is 311, bringing up the grand total to 1051. The greater number of these have come under my observation, but some are quoted on the authority of Krauss, Gould, and other authors, and are unknown to me. The three plates in the present work following the five originally published, and numbered accordingly, vi., vii., viii., give 74 figures of 56 species.

Again I have pleasure in tendering my grateful acknowledgments to friends resident in South Africa, after whom I have named some of the species, and particularly to John H. Ponsonby, Esq., for the material which has enabled me to make this contribution to South African Conchology.

G. B. SOWERBY.



APPENDIX TO  
MARINE SHELLS  
OF  
SOUTH AFRICA.

ADDITIONAL SPECIES.

GASTROPODA.

*MUREX RAMOSUS*, Linn., Sowerby, Thes. Conch., vol. iv., pl. 387, fig. 69.—Natal. Common in the Indian Ocean.

*MUREX TRIVIALIS*, A. Adams, Sow. Thes. Conch., vol. iv., pl. 388, fig. 8o.—Natal.

*FULGUR AFRICANUS*. *Sowerby*, nov. sp. (pl. 6, fig. 19.)  
Testa elongato-pyriformis, albida, fusco suffusa, flammis conspicuis fuscis longitudinaliter picta; spira breviter conica, gradata; apice? (fracto); anfractus spiraliter confertim sulcati, angulati, ad angulum nodulis parum elevatis muniti; anfractus ultimus supra angulum planato dechvis, infra convexiusculus, in cauda longiuscula desinens, ubique densissime longitudinaliter striatus, spiraliter liratus, liris numerosis, laevigatis, parum elevatis; apertura oblongo-ovata; canalis longus, mediocriter latus, rectus; columella leviter incurvata; labrum tenue.

Long. 32, diam. 16 millim.

Hab. Port Elizabeth.

Although this may be a young shell, it is certainly distinct from the dextral form of *F. berversus*.

The longitudinal flames are rather broad, distant and irregularly waved on the back of the shell, not (as in *F. carica*) marking the periods of growth.

I have only seen two specimens of this species, both collected at Port Elizabeth. They are about equal in size.

*FUSUS CRENULATUS*, Sowerby, Thes. Conch., vol iv., p. 85, pl. 14, fig. 170.—Cape of Good Hope.

Tryon opines that this is a *Siphonalia*, but in the absence of evidence, I think it best for the present to place it with *Fusus*.

*EUTHRIA MAGELLANI*, Vélin, Arch. Zool. Exper. et Gen. 1877, vol. vi., p. 104, pl. 2, figs. 8-11.—Natal. The type is from St. Paul's Island.

*PISANIA TRITONOIDES*, Reeve. (*Buccinum*), Conch. Icon. sp. 77.—Cape of Good Hope (Gould). Durban.

*METULA CLATHRATA*, Adams and Reeve, Voy. Samarang, p. 32, pl. 11, fig. 12.—Cape of Good Hope, 136 fathoms.

✓ *PLEUROTOMA TIGRINA*, Lamarck, Reeve, Conch. Icon. (*Pleurotom.t*) fig. 3.—Durban. Common in many localities from Mauritius to the Philippines and South Sea Islands.

✓ *PLEUROTOMA CINGULIFERA*, Lamarck, Reeve, Conch. Icon. fig. 1.—Durban. Distribution about the same as the last.

✓ *PLEUROTOMA MONILIFERA*, Pease, Proc. Zool. Soc. 1860, p. 398. Tryon, Man. of Conch. vol. vi., p. 173, pl. 4, fig. 52 (as var. of *P. gemmata*, Hinds).

✓ *PLEUROTOMA (DRILLIA) LAYARDI*, Sowerby, n. sp. pl. 8, fig. 3, 4. Testa turrita, rufo-tusca, laevigata, longitudinaliter costata; anfractus 7 obtusissime angulati, supra angulum leviter concavi, infra convexiusculi, costis circ. 11 rotundatis levibus, obliquiusculis instructi; ultimus spiram fere equans ad basin leviter alternatus haud rostratus, spiraliter tenuisulcatus; apertura ob'ongo-ovata; columella rectiuscula superne valde callosa; labrum arcuatum, sinu latiusculo, haud profundo emarginatum.

Long 11, diam 4 millim.

Hab. Pondoland. Port Elizabeth.

This species has long been mistaken for *P. castanea* Reeve, but upon comparison with the type of that species it is found to be quite distinct.

**PLEUROTOMA (DRILLIA) BURNUPI, Sowerby, n. sp. (Plate 8,  
fig. 1, 2.)**

Testa turrita, rufo-fusca, fascia albida interrupta et linea fusca ornata; spira acuta; anfractus 8, obtuse angulati, superne declives, costis circ. 10 lœvibus parum elevatis leviter obliquis instructi; ultimus spiram vix æquans, leviter intortus, inferne lœvigatus, ad basim liratus, haud rostratus; apertura latiuscula, intus dilute fusca; columella suboblique arcuata, superne tuberculata parviuscula instructa; labrum tenue, sinu postico latiusculo.

Long. 10, diam. 4 millim.

Hab. Durban.

Allied to *P. Layardi*, with a sharper spire; the somewhat oblique and distant ribs run to the top of the whorls, and become obsolete on the lower half of the body whorl. The reddish brown colour of the shell is relieved by an interrupted whitish band at the angle, surmounted by a thin dark brown line.

**CLATHURELLA VERRUCOSA, Sowerby, nov. sp. (pl. 6,  
fig. 14).** Testa parva, breviter ovata, rugosa, fusca, spira conica, gradata; anfractus 5, primi 2 lœves, sequentes biangulati, liris spiralibus 2, crassis confertis valde nodulosis instructi; sutura late canaliculata; anfractus ultimus spiram superans, superne angulatus, basin versus leviter attenuatus, haud rostratus, undique nodulosus; apertura latiuscula; labrum vix incrassatum; sinu postico latiusculo.

Long. 2.75., diam. 1.50 millim.

Hab Port Elizabeth.

This species was erroneously referred by me on page 7 of "Marine Shells of South Africa," as *Mangelia clathrata*, M. de Serres, from which upon further examination I find it to be quite distinct.

**CLATHURELLA COMMODA, Smith, Ann. and Mag. of Nat. Hist., 1882.—South Africa (Smith).**

**MANGILIA CEREA, Carpenter, Ann. and Mag. of Nat. Hist., 1865, p. 400.—Cape Town.**

**MANGILIA CASTA, Reeve, Conch. Icon. (Mang.), pl. 7, f. 55.—Kalk Bay.** Tryon (vol. vi., p. 305) proposed the name of *Daphnella Reeveana* for this species, on account of the pre-occupation of the name by Hinds in the genus *Pleurotomæ*, and considering both to be *Daphnella*. It seems to me, however, that Reeve's species is a true *Mangilia*, while that of Hinds is, though somewhat uncertain, probably a *Daphnella*, certainly not a *Mangilia*.

- ✓ *LOTORIUM GEMMATEUM*, Reeve, Conch. Icon. (*Triton*), sp. 60.—Natal. This species is common in many localities from Mauritius to the South Pacific Islands.
- ✓ *LOTORIUM (PRIENE) MURRAYI*, Smith (*Lampusia*), Proc. Zool. Soc. 1891, p. 436, pl. 34, fig. 1.—Off Cape of Good Hope in 150 fathoms (Voy. of Challenger).
- ✓ *LOTORIUM GALLINAGO*, Reeve, Conch. Icon. (*Triton*), pl. 1, fig. 3.—Durban.
- LOTORIUM CHLOROSTOMA*, Lamarck, Reeve, Conch. Icon. (*Triton*), fig. 3.—Durban. Widely distributed in the Indian Ocean, etc.
- LOTORIUM MONILIFERUM*, Adams and Reeve, Voy. of Samarang Moll., pl. 10, fig. 18.—Durban.
- LOTORIUM TUBEROSUM*, Lamarck, Reeve, Conch. Icon. (*Triton*), fig. 1.—Durban. Widely distributed in the Indian and Pacific Oceans.
- LOTORIUM CYNOCEPHALUM*, Lamarck, Reeve, Conch. Icon. (*Triton*), fig. 26.—Durban; also Mauritius, Ceylon, Philippines, etc.
- LOTORIUM GRACILE*, Reeve, Conch. Icon. (*Triton*), fig. 58.—Durban.
- LOTORIUM ELONGATUM*, Reeve, Conch. Icon. (*Triton*), sp. 59.—Durban.
- LOTORIUM PYRUM*, Reeve, Conch. Icon. (*Triton*), sp. 33.—Durban.
- LOTORIUM ENCAUSTICUM*, Reeve, Conch. Icon. (*Triton*), sp. 43.—Durban.
- LOTORIUM (LAGENA) CINGULATUM*, Pfeiffer, Reeve, Conch. Icon. (*Triton*), fig. 35.—Durban.
- LOTORIUM (PERSONA) ANUS*, Lamarck, Reeve, Conch. Icon. (*Triton*), fig. 44.—Natal. This species is pretty common at Mauritius.
- RANELLA AFFINIS*, Broderip, Reeve, Conch. Icon. (*Ranella*), fig. 19.—Durban. I have some doubt whether this is more than a variety of *R. granifera*, Lam.
- COMINELLA UNIFASCIATA*, VAR. *CONCOLOR*, Sowerby, var. nov. In form the same as type, but of a uniform reddish brown colour, without band.—Natal.

*BULLIA PUSTULOSA*, *Sowerby* (plate 6, fig. 1). *Journal of Conchology*, April 1894, p. 5.—Durban.

*BULLIA CAPENSIS*, *Euthyme*, Bull. de la Soc. Malac. de France, vol. ii. p. 237.—Port Elizabeth. Unfigured, and unknown to me. May be a form of *B. digitalis*.

v *BULLIA SIMILIS*, *Sowerby*, nov. sp. (pl. 7, fig. 1). Testa acuminata, solidiuscula, albida, lineis numerosis stramineis undulatis longitudinis picta; spira acuta; anfractus 8. convexiusculi, undique spiraliter sulcati, superne liris granulosis instructi; sutura anguste canaliculata, lira pustulata marginata; anfractus ultimus spiram fere æquans, ad basin profunde canaliculatus; columella leviter contorta, tenuiter callosa, extus acute lirata; apertura ovata, fauce dilute fusca.

Long. 31, diam. maj. 15 millim.

Hab. Natal.

This species is very like *B. Belangeri*, Kiener, in form and colouring, but distinguished by the distinctly granular ridges on the upper part of the whorls. It may, possibly, be only a variety, but I have examined more than a hundred specimens of *B. Belangeri* without finding the slightest trace of these characteristic bead-like granules on any.

*NASSA ALGIDA*, *Reeve*, *Conch. Icon. (Nassa)*, fig. 145.—Natal.

*NASSA PICTA*, *Dunker*, *Reeve*, *Conch. Icon. (Nassa)*, pl. 2, fig. 9.—Natal (Gibbons.)

*NASSA COCCINELLA*, *Lamarek*, *An. Sans. Vert.*, vii., p. 274.—Algoa Bay (Gibbons). This is probably the *N. coccinea*, A. Adams. I doubt its identity with *N. incrassata*.

*NASSA RUFULA*, *Kiener (Buccinum)*, *Reeve*, *Conch. Icon. (Nassa)*, pl. 2, fig. 14.—Natal. Known as an Australian species.

v *NASSA HORRIDA*, *Dunker (Buccinum)*, *Reeve*, *Conch. Icon. pl. 11*, fig. 69.—Natal.

*NASSA FENESTRATA*, *Marrat*, "New Forms of *Nassa*" 1887, p. 10; Tr. on Man. of Conch., vol. iv., p. 57, pl. 16, fig. 280, as var. of *N. albescens*.—Natal (Gibbons).

v *NASSA SIGNATA*, *Dunker*, *Tryon*, Man. of Conch., vol. iv., p. 57, pl. 17, fig. 327—South Africa (Tryon).

*NASSA KOCHIANA*, *Dunker*, *Tryon*, Man. of Conch., vol. iv., pl. 17, fig. 334.—Cape (Krauss), Table Bay (Tryon).

*NASSA PRODUCTA*, Sowerby, n. sp. (plate 8, fig. 4, 5.)  
 Testa elongato-turrita, pallide lutescens, fusco unifasciata, spira perelata, leviter convexa, ad apicem acuta; anfractus 9, rotunde convexi, sutura impressa sejuncti, costis longitudinis numerosis crassis instructi, spiraliter tenuiter lirati; ultimus brevis, rotundatus, ad basin excavatus; apertura rotunde ovata, intus lirata; labrum tenuie, extus varicosum; columella rugosa.

Long. 13.50, diam. 5 millim.

Hab. Durban.

A shell allied to *N. incrassata*, but of a very elongated form.

- v NASSARIA ACUMINATA, Reeve, Conch. Icon. (*Trilon*), pl. 14, fig. 5½.—Durban.

PURPURA DUBIA, Krauss, is COMINELLA LAGENARIA.

PURPURA CASTANEA, Krauss, Küster Monog. p. 170, pl. 28, figs. 8, 9.—Cape of Good Hope (Krauss).

PURPURA (IOPAS) SITULA, Reeve, Conch. Icon. (*Buccinum*) pl. 6, fig. 40.—Durban (Turton).

SISTRUM UNDATUM, Chemnitz, Tryon, Man. of Conch. vol. ii., p. 189, pl. 59, fig. 259 and 261. *Murex margariticola*, Brod., Reeve, Conch. Icon (*Murex*), fig. 178.—Durban. De-lagoa Bay (Gibbons).

SISTRUM ELONGATUM, Blainville, Reeve, Conch. Icon. (*Ricinula*), pl. 4, fig. 25.—Durban.

SISTRUM MUTICUM, Lamk, Reeve, Conch. Icon. (*Ricinula*), pl. 2, fig. 6.—Durban.

SISTRUM AFFINE, Pease, Proc. Zool. Soc. 1862. Tryon vol. ii., pl. 57, fig. 228, as var. of *S. Marginatum*.

SISTRUM PARVULUM, Gould, Proc. Bost. Soc. vii., p. 328. Simon's Bay (Stimpson). Probably a var. of *S. Marginatum*.

SISTRUM CONCATENATUM, Lamarck, Reeve, Conch. Icon. (*Ricinula*), sp. 18.—Durban.

SISTRUM CORONATUM, H. Adams, Proc. Zool. Soc. 1869, p. 272, pl. 19, fig. 4.—Natal. A specimen much larger than those from Mauritius.

CORALLIOPHILA FRITSCHI, Martens, Jahrb. Mal. Gesell. i., p. 135, pl. 6, fig. 3. Tryon vol. ii., p. 208, pl. 65, fig. 352.—False Bay (Martens).

*OLIVA TRUNCATA*, *Marrat*, Sowerby, Thes. Conch., vol. iv., pl. 331, fig. 41.—Cape of Good Hope (*Marrat*).

*OLIVA MICANS*, *Solander*, Sowerby, Thes. Conch., vol. iv., pl. 345, figs. 294-296. *O. nama*, *Lam.*—S Africa (Tryon).

✓ *OLIVA ISPIDULA*, *Linn*, Sow., Thes. Conch., vol. iv., p. 21, pl. 343, fig. 248—Knysna.

✓ *OLIVA SCITULA*, *Marrat*, Sow., Thes. Conch., vol. vi., pl. 333, fig. 76.—Durban.

✓ *ANCILLA DECIPIENS*, Sowerby, n. sp. (pl. 6, fig. 23). Testa oblongo-ovata, crassa, pallide purpurascens, antice et postice fusco fasciata; spira mediocriter elata, obtusiuscula; anfractus 6, planato convexi; callo tenui obtecti; callositate polita, fusco fulvoque zonata; anfractus ultimus convexiusculus, haud angulatus; columella arcuata, callosa, purpurea, superne callo fusco latemarginato effuso; apertura oblongovalis, intus purpurea.

Long. 28, diam. 15 millim.

Hab., Port Elizabeth.

Allied to *A. obtusa* and *australis*, but distinct from both.

✓ *ANCILLA CONTUSA*, *Reeve*, Conch. Icon. (*Ancillaria*), pl. ix, fig. 31.—Durban.

*ANCILLA OPTIMA*, *Sowerby*, n. sp. (Plate vi., fig. 31). Testa subfusiformi-ovata, solidiuscula, pallide fulvo-carnea, longitudinaliter sparsim rufescente strigata, antice et postice vivide fusco fasciata; spira mediocriter elata, acutiuscula; anfractus 6, planato-convexi, callo tenui obtecti; callositate polita, fusco fulvoque balteata; anfractus ultimus convexiusculus, superne obtusissime angulatus, liris obliquis numerosis angustis instructus, infra angulum callo fusco sulca alba marginato, infra medium zona angusta alba, sulca vix profunda marginata, cinctus; ad basin varice calloso fusco unisulcato terminante; columella alba, polita, levissime contorta, extus biplicata, superne callo albo effuso; apertura latiuscula, alba.

Long. 38, maj. diam. 21 millim.

Hab., Durban.

Of this fine species, I have at present only seen two specimens. It is allied to *A. australis* and *A. obtusa*, from both of which, however, it differs very materially. It may be readily distinguished from its congeners, by its acutely conical spire.

FASCIOLARIA TRAPEZIUM, *Linn*, Sowerby, Thes. Conch., vol. v., pl. 426, fig. 22.; var. AUDOUINI, *Jonas* (fig. 23.)—Durban.

FASCIOLARIA RUTILA, *Watson*, Voy Challenger, vol. xv., p. 242, pl. 13, fig. 6.—Off Cape of Good Hope 150 fath. This species bears a remarkable resemblance to *Sipho Islandicus*.

FASCIOLARIA FILAMENTOSA, *Chemnitz*, Sowerby, Thes. Conch., vol. v., pl. 424., fig. 9.—Natal.

LATIRUS ABNORMIS, *Sowerby* (Plate 6, fig. 7), Journal of Conchology, vol. viii., p. 6., 1894.—Natal.

LATIRUS TURRITUS, *Gmelin*, Tryon, Man. of Conch., vol. iii., p. 93, pl. 69, figs. 160, 161.—Natal.

PERISTERIA FENESTRATA, *Gould*, Proc. Bost. Soc., vol. vii., p. 327.—Simon's Bay (Gould). A doubtful species, unfigured.

PERISTERIA CROCEA, *Gray*, var. sulcata, *Gray* (*Turbinella*), Zool. Beechy's Voy. 116.—Natal.

VOLUTA FESTIVA, *Lamarck*, Sowerby, Thes. Conch., vol. i., pl. 52, figs. 79, 80.—A beautiful shell of great rarity. A young specimen in fair condition was found on the Natal coast by Mrs. Trotter.

MITRA LITERATA, *Lamarck*, Sowerby, Thes. Conch., vol. iv., pl. 372, fig. 436.—Natal.

MITRA CHINENSIS, *Gray*, Sowerby, Thes. Conch., vol iv., pl. 357, fig. 81.—Natal. A small specimen.

MITRA INTERLIRATA, *Reeve*, Conch. Icon. (*Mitra*), fig. 70.

MITRA CRENIFERA, *Lamarck*, Sowerby, Thes. Conch., vol. iv., pl. 354, fig. 30.—Durban.

MITRA TEXTURATA, *Lamarck*, Sowerby, Thes. Conch., vol. iv., pl. 358, fig. 84.—Durban.

MITRA CADAVEROSEA, *Reeve*, Proc. Zool. Soc. 1844, Sowerby, Thes. Conch., vol. iv., pl. 370, fig. 386.—Durban.

MITRA EXASPERATA, *Gmelin*, Sowerby, Thes. Conch., vol. iv. pl. 371, fig. 419.—Durban.

MITRA CIRCULATA, *Kiener*, Sowerby, Thes. Conch., vol. iv., pl. 358, fig. 86.—Durban.

*MITRA FLAMMEA*, *Quoy*, Sowerby, Thes. Conch., vol. iv., pl. 262, fig. 173.—Durban.

*MITRA RUFESCENS*, *A. Adams*, Proc. Zool. Soc. 1851, p. 137.—Durban. I doubt whether this is specifically distinct from *M. circulata*. The body whorl is more cylindrical.

*MITRA PRETIOSA*, *Reeve*, Sowerby, Thes. Conch., vol. iv., pl. 14, fig. 236.—Natal.

*MITRA BOVEI*, *Kiener*, Sowerby, Thes. Conch., vol. iv., pl. 5, fig. 60.—Natal (Gibbons).

*MITRA LUCTUOSA*, *A. Adams*, Sowerby, Thes. Conch., vol. iv., pl. 14, fig. 229.—Durban.

*MITRA CINNAMOMEA*, *A. Adams* (*Volutomitra*), Proc. Zool. Soc. 1854, p. 134—Natal (Adams).

*MITRA ZEPHYRINA*, *Duclos*, Sowerby, Thes. Conch., vol. iv., pl. 18, fig. 306.—Durban.

*MITRA FUSCESCENS*, *Pease*, Sow., Thes. Conch., vol. v., pl. 367, fig. 303.—Durban.

*MARGINELLA BENSONI*, *Reeve*, Conch. Icon., fig. 151.—Algoa Bay (Gibbons).

*MARGINELLA LUCIDA*, *Marrat*, Journ. of Conch., vol. i., p. 205. Tryon, vol. v., pl. 12, fig. 25.—Algoa Bay (Gibbons). This appears scarcely to differ from *M. pellicula* of the same author.

*MARGINELLA CINEREA*, *Jousseaume*, Reeve, Conch. Icon. (*Marginella*), sp. 145.—(*M. semen*, Reeve, non Lea).

*MARGINELLA PUELLA*, *Gould*, Proc. Bost. Soc., vol. 7, p. 385.—Simon's Bay (Gould). Unfigured, and unknown to me.

*MARGINELLA QUADRIFASCIATA*, *Marrat*, Ann. and Mag. Nat. Hist. 4th Series, vol. xii., p. 426 (unfigured).—S. Africa.

*MARGINELLA ALBANYANA*, *Gaskoin*, Ann. and Mag. Nat. Hist. 2 Series, vol. xi., p. 104.—Albany (Gaskoin). Unknown to me.

*MARGINELLA RUFULA*, *Gaskoin*, Reeve, Conch. Icon. (*Marginella*), fig. 149.—Green Point.

*MARGINELLA PERMINIMA*, *Sowerby* (pl. 6, fig. 36), Journal of Conchology, vol. viii., p. 7.—S. Africa (Turton).

**MARGINELLA BURNUPI**, *Sowerby*, n. sp. (pl. 6, fig. 35).  
Testa cylindrica, alba, solidiuscula; spira fere plana; apertura angusta; columella quinqueplicata; plicis acutis; labrum crassiusculum.

Long. 4, diam. 2 millim.

Hab. Port Elizabeth.

A small cylindrical white shell with the spire almost flat.

**MARGINELLA PONSONBYI**, *Sowerby*, n. sp. (pl. 6, fig. 2).  
Testa subcylindrica, nitens, sub-pellucida, pallide cornea, strigis fulvis transversis irregulariter ornata; antice leviter attenuata, postice vix inflata; spira obtuse conica, parum elevata; apertura angustiuscula, antice paulo latior; columella triplicata; labrum incrassatum.

Hab. S. Africa.

Long. 9, diam. 5 millim.

A sub-pellucid shell of somewhat cylindrical form, pale in colour with transverse fulvous streaks, some of which run obliquely.

**COLUMBELLA FULMINATA**, *Gould*, "Otia," p. 131.—Simon's Bay (Stimpson).

**COLUMBELLA OBTUSA**, *Sowerby*, Thes. Conch., vol. i., p. 120, pl. 37, fig. 63.—Durban.

**COLUMBELLA PYRAMIDALIS**, *Sowerby* (pl. 6, fig. 4), Journal of Conchology, vol. vii. p. 7.—Port Elizabeth.

**COLUMBELLA KITCHINGI**, *Sowerby*, pl. 6, fig. 3, Journal of Conchology, vol. vii., p. 7.—Green Point, Cape of Good Hope.

**COLUMBELLA MERCATORIA**, *Lamarck*, Sow., Thes. Conch., vol. i., p. 115, pl. 36, figs. 28-32.—Natal. Very common in the West Indies.

**COLUMBELLA LIGULA**, *Duclos*, Monog., pl. 2, figs. 11-16, Sow., Thes. Conch. vol. i. p. 123, pl. 38, figs. 83-85.—Natal (Gibbons). Is. Ticao (Cuming), Solomon Islands, and Australia.

**COLUMBELLA AVENA**, *Reeve*, Conch. Icon. (*Columbella*), fig. 158.—Buffalo, Cape Colony (Reeve).

**COLUMBELLA LANGLEYI**, n. sp. (pl. 8, f. 8, 9).—Testa elongata, pellucida, succinacea, polita; spira acuminata, acutiuscula, ad apicem obtusa; anfractus 5, convexi, longitudinaliter obscurissime plicati, superne obtusissime angulati; ultimus spiram fere æquans, leviter inflata, basim versus leviter attenuatus; apertura oblonga; columella levissime contorta; labrum tenue.

Long. 4.50, diam. 1.50 millim.

Hab. Port Elizabeth (Crawford).

A smooth, transparent, amber-coloured little shell.

**COLUMBELLA CONSANGUINEA**, n. sp. (pl. viii. f. 6, 7). Testa elongata, fulvo fusca, crassiuscula, undique sculpta; spira acuminata, ad apicem obtusa; anfractus 5, convexi, rotundati, spiraliter striati, longitudinaliter plicati, plicis pernumerosis, interstitiis puncturatis; anfractus ultimus spiram paulo superans, basim versus leviter attenuatus; apertura oblonga; columella rectiuscula, labrum leviter incrassatum.

Long. 4.25, diam. 1.50 millim.

Hab. Port Elizabeth.

Similar in form to *C. Langleyi*, but of a thicker substance, and sculptured with fine, close, longitudinal riblets, and minute spiral striae.

✓ *ENGINA NATALENSIS*, *Melvill*, Proc. Malac. Soc., vol. i. p. 226.—Durban.

✓ *HARPA VENTRICOSA*, *Lamarck*, Sow., Thes. Conch., vol. iii., p. 169, pl. 232, figs. 18, 22.—Durban.

✓ *CASSIS GLAUCA*, *Lamarck*, *Reeve*, Conch. Icon. (*Cassis*), fig. 33.—Natal. Common in the Indian Ocean.

✓ *CASSIS AREOLA*, *Lamarck*, *Reeve*, Conch. Icon. (*Cassis*), fig. 24.—Durban. Widely distributed in the Indian and Pacific Oceans.

✓ *CASSIS BISULCATA*, *Schub & Wagn.*, *Reeve*, Conch. Icon. (*Cassis*), fig. 6.—Natal.

*CASSIS PILA*, *Reeve*, Conch. Icon. (*Cassis*), fig. 31.—A variety without blotches, specimen in perfect condition, found at Durban.

✓ *CASSIS CRATICULATUS*, *Euthyme*, Bull. Soc. Mal. de France, vol. ii., p. 250 (unfigured). I think from the description that this is *C. pila*.—S. Africa.

*DOLIUM FIMBRIATUM*, *Sowerby*, *Reeve*, Conch. Icon. (*Dolium*) fig. 3.—Durban.

*DOLIUM PROCELLARUM*, *Euthyme*, Bull. de la Soc. Mal. de France, vol. ii., p. 247.—Pt. Elizabeth (Euthyme). Having read the description (without figure) I feel almost sure that this is the same as *D. luteostoma*.

✓ *PYRULA FICUS*, *Linn.*, Sow., Thes. Conch., vol. iv. pl. 423, fig. 4.—Natal.

- ✓ PYRULA FICOIDES, Lamarck, Tryon, Man. of Conch., vol. vii., pl. v., fig. 28. P. reticulata, auctorum, sed non Lamk.—Durban.
- ✓ NATICA QUEKETTI, Sowerby (pl. 6, fig. 6), Journal of Conch., vol. vii., p. 8.—Natal.
- ✓ NATICA SEBÆ<sup>o</sup>, Souleyet, Voy. de la Bonite, pl. 35, figs. 6, 7.—Thes. Conch., vol. v., pl. 457, fig. 79.—Natal.
- ✓ NATICA TÆNIATA, Menhe (*N. ala-papilionis*, Ch.), Sow., Thes. Conch., vol. v., pl. 457, fig. 4.—Natal.
- ✓ NATICA ZANZIBARICA, Recluz, Sow., Thes. Conch., vol. v., pl. 457, fig. 73. (A young specimen).—Durban.
- ✓ NATICA MELANOSTOMA, Gmelin, Sow., Thes. Conch., vol. iv., pl. 457, fig. 72.—Durban.
- ✓ NATICA SIMPLEX, Sow., n. sp., pl. 6, fig. 18.—Testa parva, gibbosa, alba, lævis, aut obscure longitudinaliter sculpta; spira breviter conica; anfractus 4, convexi, sutura vix impressa; anfractus ultimus rotundatus; columella rectiuscula, callosa; umbilicus parvus, angustus; apertura medio-criter lata, fauce alba, lœvis.  
Long. 6, diam. 5.50 millim.  
Hab. South Africa (Turton).  
A small white shell of very simple and regular form.
- ✓ NATICINA PAPILLA, Gmelin, Tryon, Man. of Conch., vol. 8, pl. 25, fig. 87.—Durban.
- ✓ SIGARETUS PLANULATUS, Recluz, Thes. Conch., vol. v., pl. 442, fig. 29.—Durban.
- ✓ SIGARETUS DELESSERTI, Recluz, Sow., Thes. Conch., vol. v., pl. 442, figs. 20-22.—Durban.
- ✓ VANIKORO DESHAYESIANA, Recl., Sow., Thes. Conch., vol. v., pl. 482, fig. 17.—Durban.
- ✓ SCALARIA JUKESIANA, Forbes, Voy. Rattlesnake, Append., p. 383, pl. 3, fig. 7.—Port Elizabeth.
- ✓ SCALARIA SIMPLEX, Sow. (pl. 6, fig. 5), Journal of Conchology, vol. vii., p. 8, 1894.—Natal.
- TEREBRA RUFOPUNCTATA, Smith, Ann. and Mag. of Nat. Hist., 1877, vol. xix., p. 29.—Durban.
- TEREBRA DIMIDIATA, Lamk., Sow., Thes. Conch., vol. i., pl. 41, fig. 7, 8.—Durban.

TEREBRA CASTA, *Hinds*, Sow., Thes. Conch., vol. i., pl. 44, fig. 84.—Natal.

TEREBRA STRAMINEA, *Gray*, Sow., Thes. Conch. vol. i., pl. 42, fig. 22, 23.—Natal.

TEREBRA LÆVIGATA, *Gray*, Sow., Thes. Conch., vol. i., pl. 44, fig. 93.—Natal.

TEREBRA LONGISCATA, *Deshayes*, Reeve, Conch. Icon., fig. 103.—Durban.

TEREBRA FICTILIS, *Hinds*, Sow., Thes. Conch. vol. i., pl. 45, f. 109, 110.—Durban.

TEREBRA GEMINATA, *Deshayes*, Proc. Zool. Soc. 1859, p. 296 (unfigured).—Natal (Deshayes). Probably var. of *T. spectabilis*, Hinds.

TEREBRA NEBULOSA, *Sowerby*, Thes. Conch., vol. i., pl. 43, fig. 51. (?—*T. nubeculata*, Sow.)—Durban.

TEREBRA BABYLONIA, *Lamarche*, Sow., Thes. Conch., vol. i., pl. 43, fig. 67.—Durban.

TEREBRA SUBULATA, *Lamarck*, Sow., Thes. Conch., vol. i., pl. 41, fig. 16.—Durban.

✓ PYRAMIDELLA MITRALIS, A. Adams, Sow., Thes. Conch., vol. ii., p. 814, pl. 172, fig. 9.—Durban.

✓ OBELISCUS TERES, A. Adams, Sow., Thes. Conch., vol. ii., p. 807, pl. 171, f. 31, 32.—Durban.

✓ OBELISCUS DOLABRATUS, *Linn.*, Sow., Thes. Conch., vol. ii., pl. 171, fig. 1.—Durban.

✓ CIONISCUS UNILINEATUS, *Sowerby* (Plate 6., fig. 8), Journal of Conchology, vol. vii., p. 8 (as *Aclis*).—Port Elizabeth.

✓ CIONISCUS PELLUCIDUS, *Sowerby* n. sp. (Plate 6., fig. 10). Testa parva, turrata, alba, pellucida; anfractus  $6\frac{1}{2}$ , convexi, spiraliter subtilissime striati, liris tenuissimis 2-3, superne instructi; sutura impressa, anguste lirata; nucleus subretrorsus; anfractus ultimus  $\frac{1}{4}$  longitudinis æquans, leviter rotundatus; apertura ovata; labrum tenué; columella tenuissima, leviter incurvata, haud plicata.

Long. 3.50. diam. 1.50 millim.

Hab. Port Elizabeth.

A little white pellucid shell, very lightly spirally striated; whorls slightly shouldered, with two or three fine ridges.

TURBONILLA TROCHLEARIS, *Gould* (*Chemnitzia*), *Otia Conchologica*, p. 151 (unfigured).—Simon's Bay (*Gould*).

TURBONILLA OBELISCUS, *Gould* (*Chemnitzia*), *Otia Conchologica*, p. 150 (unfigured).—Simon's Bay.

EULIMA SIMPLEX, *Sowerby*, n. sp. (Plate 6, fig. 9.) Testa subulata, polita, semipellucida, alba; spira acutiuscula, ad apicem obtusa; anfractus 7, vix convexi; anfr. ultimus  $\frac{1}{2}$  longitudinis æquans; apertura ovata, leviter obliqua; columella oblique rectiuscula.

Long. 3.50, diam. 1.50. millim.

Hab. South Africa.

A little white species of very simple character.

ODOSTOMIA LUCIDA, *Sowerby*, n. sp. (Plate 6, fig. 11). Testa oblonga, alba, subpellucida, lævis, antice inflata, postice acuminata; spira mediocriter elongata, convexiuscula; anfractus 5, convexiusculi; sutura impressa, lira angusta inserta; anfractus ultimus spiram æquans, basin versus inflatus, rotundatus; apertura ovata; columella leviter sinuata.

Long. 3, diam. 1 $\frac{1}{2}$  millim.

Hab. Port Elizabeth.

CERITHIOPSIS LIRATA, *Sowerby*, n. sp. (Plate 6, fig. 12). Testa elongato pyramidalis, fulva, antice angulata; spira acuta, ad apicem obtusiuscula; anfractus 11, primi 3 læves, rotundati, sequentes planati, spiraliter tri-lirati; sutura vix conspicua; anfractus ultimus brevis, quadriliratus; basis concava, lira unica marginata; columella brevis, rectiuscula, apertura quadrata.

Long. 4.25, diam. 1.75.

Hab. Port Elizabeth.

A little shell of very distinct pyramidal form, with a flattened, slightly concave base. The spiral ridges are like those prevailing in the genus *Turritella*.

CERITHIOPSIS EXQUISITA, *Sowerby*, n. sp. (Plate 6, fig. 13). Testa turricula, fulva; spira convexiuscula, ad apicem acuta; anfractus 10, primi minut, læves, sequentes convexi, biangulati, liris spiralibus et longitudinalibus decussatim exigue sculpta; sutura impressa, lira angusta munita; anfractus ultimus breviusculus superne leviter concavus, ad basin convexiusculus, anguste umbilicatus; columella brevis, rectiuscula, apertura subquadrata.

Long. 4.50, diam. 1.50 millim.

Hab. Natal.

A fulvous crisply sculptured little shell.

- / SOLARIUM MAXIMUM, *Philippi*, Sow., Thes. Conch. vol iii., pl. 250, figs. 5-6.—Durban. Usually found in the China Sea.
- / SOLARIUM LEVIGATUM, *Lamarch*, Sow., Thes. Conch., vol. iii., pl. 251, figs. 21-22.—Durban, Persian Gulf, Indian Ocean, &c.
- / SOLARIUM PERSPECTIVUM, *Linn.*, Sow., Thes. Conch. vol. iii., pl. 253, figs. 36-38.—Natal, and Indian and Pacific Oceans.
- / SOLARIUM (PHILIPPIA) HYBRIDUM, *Linn.*, Sow., Thes. Conch., vol. iii., pl. 253, fig. 39—St. John's (Turton).
- / SOLARIUM (TORINIA) DORSUOSUM, *Hinds*, Sow., Thes. Conch., vol. iii., pl. 254, fig. 73.—St. John's (Turton), Natal. Also Persian Gulf, and Japan.
- / SOLARIUM (TORINIA) CYLINDRACEUM, *Chemnitz*, Sow., Thes. Conch., vol. iii., pl. 254, figs. 98, 99.—Natal.
- / IANTHINA PALLIDA, *Harvey*, Sow., Thes. Conch., vol. v., pl. 444, figs. 19, 20.—Port Elizabeth.
- CONUS CEYLANENSIS, *Hwass*, Sow., Thes. Conch., vol. iii., pl. 192, figs. 139-41.—Durban.
- CONUS QUERCINUS, *Bruguiere*, Sow., Thes. Conch., vol. iii., pl. 197, fig. 239.—Natal. Common in the Indian Ocean, Persian Gulf, East Africa, &c.
- CONUS MILES, *Linn.*, Sow., Thes. Conch., vol. iii., pl. 193, fig. 157.—Durban. Widely distributed, principally in the Indian Ocean.
- CONUS VERMICULATUS, *Lamarck*, Sow., Thes. Conch. vol. iii., pl. 189 fig. 52.—Durban. Distr., N.E. Africa, Red Sea, Indian Ocean, &c.
- CONUS LEGATUS, *Lamarck*, Sow., Thes. Conch., vol. iii., pl. 209, fig. 566 (*C. musivus*, Brod).—Natal (Gibbons).
- CONUS LAMARCKI, *Kiener*, Coq. Viv., p. 240, pl. 83, fig. 4.—S. Africa (Tryon).
- CONUS PRIMULA, *Reeve*, Conch. Icon. Supp., pl. 6, fig. 259.—Natal (Tryon).
- CONUS PLUMBEUS, *Reeve*, Conch. Icon. (*Conus*), fig. 253.—S. Africa (Tryon).
- CONUS RATTUS, *Lamarck*, Sow., Thes. Conch., vol. iii., pl. 193, fig. 161.—Durban.

**CONUS ARACHNOIDEUS**, *Gmelin*, Sow., Thes. Conch., vol. iii., pl. i, fig. 14.—Natal.

**CONUS ARENATUS**, *Hwass*, Sow., Thes. Conch., vol. iii., pl. 188, fig. 7.—Durban.

**CONUS OBSCURUS**, *Humph.* Sow., Thes. Conch., vol. iii., pl. 208, fig. 526.—Durban.

**STROMBUS FUSIFORMIS**, *Sowerby*, Thes. Conch., vol. i., pl. 9, figs. 91, 92.—Durban.

**CYPRÆA CLANDESTINA**, *Linn.*, Sow., Thes. Conch., vol. iv., pl. 310, f. 139.—Durban.

**CYPRÆA STERCUS-MUSCARUM**, *Lamarck*, Sow., Thes. Conch., vol. iv., pl. 323, f. 363.—Natal (Gibbons).

**CYPRÆA CASTANEA**, *Higgins*, Sow., Thes. Conch., vol. iv., pl. 320, figs. 302, 303.—Cape of Good Hope (Higgins).

**CYPRÆA CRIBRARIA**, *Linn*, Sow., Thes. Conch., vol. iv., pl. 311, fig. 164.—Uenzinto, Natal.

**CYPRÆA STAPHYLÆA**, *Linn*, Sow., Thes. Conch., vol. iv., pl. 316, fig. 225. (var. *Limacina*, Lam.)

**TRIVIA PRODUCTA**, *Gaskoin*, Sow., Thes. Conch., vol. iv., pl. 327, f. 495.—Aguilhas Bank.

**TRIVIA SULCATA**, *Gaskoin*, Sow., Thes. Conch., vol. iv., pl. 326, f. 454.—Pondoland (Layard).

**TRIVIA QUADRIPUNCTATA**, *Gray*, Sow., Thes. Conch., vol. iv., pl. 326, f. 460.—Pondoland (Layard). Common in West Indies, also reported from Philippines.

**TRIVIA INSECTA**, *Mighels*, Sow., Thes. Conch., vol. iv., pl. 326, fig. 478.—S. Africa.

**TRIVIA ORYZA**, *Lamarck*, Sow., Thes. Conch., vol. iv., pl. 326, fig. 474.—Durban.

**TRIVIA VITREA**, *Gaskoin*, Sow., Thes. Conch., vol. iv., pl. 326, fig. 456.—Durban.

**CANCELLARIA IMBRICATA**, *Watson*, Linn. Soc. Journ. Zool. xvi., p. 325. Moll. Voy. Challenger vol. xv., pl. 18, fig. 10.—Off Cape of Good Hope (Watson).

**CERITHIUM CRASSILABRUM**, *Krauss*, Sow., Thes. Conch., vol. ii., pl. 182, f. 203.—Durban.

✓ PYRAZUS PALUSTRIS, *Linn.*, Sow., Thes. Conch., voi. ii., pl. 185, f. 261.—Natal.

✓ TRIFORIS CORRUGATUS, *Hinds*. Tryon, Man. of Conch., vol. ix., pl. 39, f. 59.—Durban.

✓ RISSOA (CINGULA) CAFFRA, n. sp. (Plate 6, fig. 15.) Testa ovato-conoidea, anguste umbilicata, tenuis, polita, fulvo-cornea, fusco obscure trifasciata; spira conica; anfractus 5, convexi; sutura profunda; anfractus ultimus spiram superans, inflata, rotundata; apertura ovata, peristoma simplex.

Long. 2.50, diam. 1.50 millim.

Hab. Port Elizabeth.

A thin, polished, horny little shell with three faint bands.

✓ TURRITELLA DECLIVIS, *Adams & Reeve*, Voy. of Samarang, Moll. p. 48, pl. 2, fig. 10.—Hout's Bay.

✓ XENOPHORA CORRUGATA, *Reeve*, Conch. Icon., sp. 6.—Durban.

✓ CALYPTERA CICATRICOSA, *Reeve*, Conch. Icon. (Calyptrea), fig. 3.—Durban.

✓ CALYPTERA POROSA, *Reeve*, Conch. Icon. (Calyptrea), fig. 20.—Durban.

✓ TURBO INTERCOSTALIS, *Menke*, Sow., Thes. Conch., vol. v., pl. 494, f. 14.—Durban.

✓ TURBO (LEPTOTHYRA) PILLULA, *Dunker*, Sow., Thes Conch., vol. 5, pl. 505, f. 160.—St. John's (Turton).

✓ TURBO TRICARINULATUS, *Euthyme*, Bull. de la Soc. Malac de France, vol. ii., p. 252. (unfigured). This is surely a small form of *T. cidaris*, with the keels of var. *nataleensis* (*Reeve*).

✓ TURBO PONSONBYI, n. sp. (Plate 6, fig. 20.) Testa subglobosa, crassa, profunde umbilicata, grisea, fusco maculata, spiraliter dense lirata; liris rotundatis, maculis parvis fuscis articulatis, lira minima interveniente; spira breviter conica; anfractus 4, convexi; sutura canaliculata; anfr. ultimus rotundatus; basis convexa, costa crassa nodosa in regione umbilicali, sulca profunda et latiuscula marginata, instructa; apertura fere circularis, fauce alba; peristoma crassum. Operculum intus planum, extus convexum, profunde umbilicatum, corrugatum, carina angustiuscula marginatum.

Long 8, diam. 8 millim.

Hab. Durban.

A small shell resembling one of the larger species of *Leptothyra* (*Collonia*), from which section it differs principally in the form of its operculum.

PACHYPOMA TAYLORIANUM, *Smith*, Proc. Zool. Soc. 1880, pl. 48, fig. 9, p. 483.—Port Elizabeth. This species was described from a single specimen without locality in the collection of the late Thos. Lombe Taylor.

GIBBULA TRYONI, *Pilsbry* (Plate 6, fig. 22.) Manual of Conchology, vol. xi., p. 239, pl. 69, figs. 20, 21. *G. incincta*, Sow. Journ. of Conch., April 1894, is a synonym.

GIBBULA FUCATA, *Gould*, Proc. Boston Soc. Nat. Hist. p. 8, p. 20.—Simon's Bay (Turton).

GIBBULA LOCULOSA, *Gould*, "Otia," p. 159.—False Bay.

MARGARITA PINTADO, *Gould*, "Otia," p. 154.—Simon's Bay.

MARGARITA ARTICULATA, *Gould*, "Otia," p. 154.—Simon's Bay.

MONILEA SPURIA, *Gould*, "Otia," p. 155.—Simon's Bay.

SOLARIELLA SPLENDENS, *Sowerby*, n. sp. (pl. 6, fig. 21).—Testa depresso orbicularis, late et profunde umbilicata, aureo-nitens, fusco maculata ad basim fusco oblique strigata; spira breviter conica, gradata, ad apicem acuta; anfractus  $6\frac{1}{2}$  tabulati, spirali ter grano-lirati, sutura canaliculata sejuncti; anfractus ultimus depresso rotundatus, supra angulatus, infra rotundatus, liris 7, supremis granulatis, coeteris planis; basis convexa, laevis; umbilicus perspectivus, intus grano-liratus; apertura sub-quadrata, vix obliqua; columella oblique rectiuscula, vix callosa.

Alt. 6, diam. 9 millim.

Hab. Natal.

A pretty little shining shell, smooth below and with about 7 ridges above the periphery, the upper ridges being beaded.

CALLIOSTOMA LAYARDI, *Sowerby*, n. sp. (pl. 8, f. 10, 11).—Testa conica, elatiuscula, angustissime umbilicata, argenteo-nitens, ad angulum vivide rubro bicingulata, ad basim rubra; spira acuta, ad apicem vivide rubra; anfractus  $6\frac{1}{2}$ , primi (2) laeves rotundati, sequentes plano declives, liris 4 leviter nodulosis instructi, ad angulum rubro bicarinati; sutura

canaliculata; basis vix convexa, valde lirata, liris 7-8 confertiusculis, rotundatis; columella leviter obliqua, callosa; apertura oblique sub-quadrata, intus argentea.

Alt. 13, diam. 14 millim.

Hab. Pondoland (Layard).

A regularly conical typical *Calliostoma*, distinguished by the bright ruby double keel at the angle of the whorls. In the type specimen the narrow umbilicus is only partially covered, but probably in older shells it would be entirely closed.

CLANCULUS KRAUSSI, *Philippi* (*Monodontia*), Conch. Cab., p. 82, pl. 14, f. 14.—Durban.

EUCHELUS QUADRICARINATUS, *Chemnitz* (*Trochus*), Conch. Cab. xi., p. 167, pl. 196, f. 1892.—Durban.

UMBONIUM VESTARIUM, *Linn.*, Sow., Thes. Conch., vol. v., pl. 472, figs. 1-5.—Durban.

FISSURELLA SIMILIS, *Sowerby*, Thes. Conch., vol. iii., p. 194, pl. 241, f. 143.—Durban

FISSURELLA GRECA, *Linn.*, Sow., Illust. Index Brit. Shells, pl. 11, figs. 1, 2 (*F. reticulata*, Donov.).—Durban.

MACROCHISMA COMPRESSA, *A. Adams*, Sow., Thes. Conch., vol. iii., p. 205, pl. 244, f. 218.—S Africa (Turton).

CHITON LYRATUS, *Sowerby*, Conch. Illust., fig. 126.—Port Elizabeth.

CHITON (ISCHNOCHITON) ELIZABETHENSIS, *Pilsbry*. “The Nautilus,” vol. viii., p. 9.—Port Elizabeth. This is the species quoted by me as *Chiton marginatus*, which it very closely resembles. Mr. E. R. Sykes (Proc. Malac. Soc., vol. i., p. 133) expressed the opinion that it was identical with *C. oniscus*, Krauss.

CHITON (ACANTHOBLEURA) AFRA, *Rochebrune*, Bull. Soc. Philom. 1882, p. 192.—Cape of Good Hope (Verreaux). Unknown to me.

CHITON (ACANTHOBLEURA) QUATREFAGESI, *Rochebrune*, Journ. de Conch., 1881, p. 44.—Cape of Good Hope (Verreaux). Unknown to me.

CHITON (ACANTHOCHITES) CARPENTERI, *Pilsbry*, Man. of Conch., vol. xv., p. 35, pl. 1, f. 14-22.—Pt. Elizabeth.

- ✓ CHITON (ONITHOCHECITON) LITERATUS, Krauss, sudafric, Moll., p. 36.—Natal (Krauss).
- ✓ CHITON (ONITHOCHECITON) ALVEOLATUS, Rochebrune, Bull. Soc. Phil. de Paris, 1883, p. 32.—Cape of Good Hope (Paris Museum).
- ✓ CHITON (GYMNOPLAX) ANAGLYPTUS, Rochebrune, Bull. Soc. Phil. de Paris, 1883, p. 33.—Cape of Good Hope (Paris Museum.).
- ✓ CHITON (GYMNOPLAX) MELANOTREPHEUS, Rochebrune, Bull. Soc. Phil. de Paris, 1883, p. 34.—Cape of Good Hope (Paris Museum.).
- ✓ ACTÆON FLAMMEUS, Gmelin, Reeve, Conch. Icon. (Tornatella) fig 2.—Durban.
- ✓ ACTÆON (SOLIDULA) AFFINIS, A. Adams, Proc. Zool. Soc., 1854, p. 61.—Pilsbry, Man. of Conch., vol. xv., pl. 20 A, fig. 52.—Durban.
- ✓ HYDATINA UNDATA, Bruguière, Pilsbry, Man. of Conch., vol. xv., pl. 59., f. 20, 21.—Natal.
- BULLINA OBLONGA, Sowerby, n. sp. (plate 8, fig. 12, 13). Testa oblongo-ovalis, postice acuminata, antice leviter attenuata, undique confertim sulcata, grisea, fusco tri-zonata, strigis numerosis fuscis irregularibus ornata; spira elatiuscula, convexa, ad apicem acuta; anfractus  $4\frac{1}{2}$ , rotunde convexit; sutura anguste canaliculata; anfr. ultimus elongatus, superne leviter rotundatus; apertura oblonga, mediocriter lata, utrinque angustior; labrum tenui, columella rectiuscula, obscurissime plicata.  
 Long. 13, diam. 7 millim  
 Hab. Pondoland (Layard)  
 Compared with *B. Zizzac* the spire is much higher.
- HAMINEA GRACILIS, Sowerby, n. sp. (Pl. 6, fig. 16.) Testa anguste rimata, oblongo sub-cylindracea, tenuis, pallide straminea, antice mediocriter lata, postice angustior, undique spiraliter minutissime striata, longitudinaliter leviter et irregulariter plicata; vertice anguste sed profunde umbilicata; apertura antice lata, postice angusta.  
 Long. 11 diam. 6.50 millim.  
 Hab. Durban.
- ✓ APLUSATRUM AMPLUSTRE, Linn., Pilsbry, Man. of Conch. vol. xv., pl. 44, figs. 1-6. Synonyms, *Bulla aplustre*, Lamk. *Aplustrum fasciatum*, Schum., *Aplustra pulchella*, Swains.

- ✓ *Amplustre thalassiarchi*, Martini, &c.—Durban. A species of wide geographical distribution.
- ✓ *ATYS ELONGATA*, A. Adams (Bulla), Sow., Thes. Conch., vol. ii., pl. 125, f. 121.—Durban. Widely distributed in the Indian and Pacific Oceans.
- ✓ *CYLINDROBULLA SCULPTA*, G. & H. Nevill, Journ. Asiatic Soc., p. 2, No. 2, 1869, p. 68, pl. 13, fig. 3.—Hab., Port Elizabeth.
- ✓ *VOLVATELLA LAGUNCULA*, Sowerby (pl. 6, fig. 17), Journal of Conchology, vol. vii., p. 10, 1894.—Port Elizabeth.

## PELECYPODA.

- ✓ *SOLEN CORNEUS*, Lamarck, Sow., Conch. Icon. (*Solen*), fig. 19.—Durban.
- ✓ *SOLEN SLOANEI*, Gray, Sow., Conch. Icon., f. 10.—Durban.
- *PANDORA DISSIMILIS*, Sowerby (pl. 6, fig. 33), Journal of Conchology, vol. vii., p. 11, April, 1894.—Sea Point, Cape Town.
- ✓ *BASTEROTIA OBTUSA*, Sowerby (pl. 6, fig. 27), Journal of Conchology, vol. vi., p. 11, April, 1894.—Durban.
- ✓ *BASTEROTIA TRICOSTALIS*, Sowerby, n. sp. (pl. 8, figs. 14, 15).—Testa aequivalvis, inaequilateralis, pallide straminea, elongato-subtrigona, crassiscula, antice breviuscula, rotundata, postice longior, oblique, declivis, angulata, ad angulum acute carinata, irregulariter rugosa; area postica leviter concava, valde tricostalis; umbones acuti, approximati, lunula parva; cardo in utraque valva dentem unicum leviter excurvatum munitus.  
Antero-post 12, umbono marg. 7 millim.  
Hab. Durban (Burnup).
- This second species of *Basterotia*, which I have described from South Africa, is more like the type of the genus than the first (*B. obtusa*). It has a prominent keel ridge at the posterior angle, behind which the somewhat concave area is furnished with two additional keels or ribs.
- ✓ *MACTRA ACHATINA*, Chemnitz, Conch. Cab. vol. ii., p. 218, fig. 1957.—Reeve, Conch. Icon. (*Mactra*), fig. 51.
- ✓ *MACTRA AEQUISULCATA*, Sowerby (pl. 8, fig. 26), Journal of Conchology, vol. vii., p. 12.—Durban.

- ✓ *RAETA PELLICULA*, Deshayes, Reeve, Conch. Icon. (*Mactra*), pl. 21, fig. 124.—Durban.
- ✓ *MACTRINULA OVALINA*, Lamk, Reeve, Conch. Icon. (*Mactra*), fig. 66 (= *depressa*, Speng).—Durban.
- ✓ *SENELE CORDIFORMIS*, Chemnitz, Reeve, Conch. Icon. (*Amphidesma*), pl. 5, fig. 30.—Natal.
- ✓ *IACRA SEYCHELLARUM*, H. & A. Adams, Rec. Moll. vol. ii., p. 409.—Natal.
- ✓ *PSAMMOBIA BURNUPI*, Sowerby (plate 6, fig. 30), Journal of Conchology, vol. vii., p. 1894.
- ✓ *PSAMMOBIA ORNATA*, Deshayes, Reeve, Conch. Icon. (*Psam-mobia*), fig. 36.—Durban.
- ✓ *TELLINA DISPAR*, Conrad, Sow., Thes. Conch. vol. i., p. 306, pl. 59, fig. 108.—Natal.
- ✓ *TELLINA VIRGATA*, Linn., Sow., Thes. Conch., vol. i., p. 228, pl. 65, fig. 212.—Natal.
- ✓ *TELLINA RASTELLUM*, Hanley, Sow., Thes. Conch., vol. i., p. 223, pl. 64, fig. 231.—Natal.
- ✓ *TELLINA PLANISSIMA*, Anton, Sow., Thes. Conch., vol. i., p. 295, pl. 59, fig. 124.—Durban.
- ✓ *TELLINA PHARAONIS*, Hanley, Sow., Thes. Conch., vol. i., p. 235, pl. 63, f. 215.—Durban.
- ✓ *TELLINA PERNA*, Spengler, Sow., Thes. Conch. vol. i., p. 236, pl. 63, f. 202.—Natal.
- ✓ *TELLINA QUEKETTI*, n. sp. (plate 8, fig. 16.) Testa trigono sub-ovalis, solidiuscula, inaequilateralis, alba, nitens, concentrica levissime rugata; umbones leviter prominentes, acutiusculi, post medium locati; margo dorsalis utrinque declivis; latus anticum rotundatum, posticum leviter truncatum, margo ventralis arcuatus.  
Antero-post 15.50, umbono-marg. 11.50.  
Hab. Durban.  
Var. *radiata*. (fig. 17) Extus intusque purpureo radiata.
- ✓ *TELLINA VIRGULATA*, Hanley, Sow., Thes. Conch., vol. i., pl. 56, fig. 5.—Durban.
- ✓ *TELLINA PRISMATICA*, Sowerby, n. sp. (pl. vi., fig. 29). Testa oblonga, tenuissima, compressa, iridescent, valde

inæquilateralis, concentrica striata; margo dorsalis anticus leviter arcuatus, posticus valde declivis.

Antero-post 16, umbono-marg. 8 millim.

Hab. Durban.

A thin iridescent species, somewhat resembling *T. Valtonis*.

- ✓ TELLINA VULSELLA, *Chemnitz*, Thes. Conch., vol. i., pl. 61, fig. 162.—Natal.
- ✓ TELLINA (MACOMA) CANDIDATA, *Sowerby* (pl. vi., fig. 25), Journal of Conchology, vol. vii., p. 12, April 1894.—Durban.
- ✓ STRIGILLIA TROTTERIANA, *Sowerby* (pl. 6, fig. 32), Journal of Conchology, vol. viii., p. 12, April, 1896.—Durban.
- ✓ DONAX NITIDUS, *Deshayes*, Sow., Thes. Conch., vol. 3, p. 314, pl. 282, fig. 75.—Natal. Also East Australia.
- ✓ DONAX LUBRICUS, *Hanley*, Reeve, "Conch. Icon." (*Donax*), pl. 7, fig. 46.—Natal.
- ✓ DONAX ÆMULUS, *Smith*, Proc. Zool. Soc., 1877, p. 271, pl. 75, figs. 23-25.—Durban.
- ✓ DONAX BURNUPI, *Sowerby* (pl. 6, fig. 26), Journal of Conchology, vol. vii., p. 14.—Natal.
- ✓ DONAX SIMPLEX, *Sowerby*, n. sp. (pl. 8, fig. 18, 19). Testa cuneiformis solida, postice tumida, antice attenuata, luteo albida, radiatim striata; striis confertis, puncturatis, antice evanidis; umbones obtusi; margo dorsalis anticus oblique declivis, posticus sub rotunde truncatus, ventralis rectiusculus.  
Antero-post 7, umbono-marg. 5, crass. 3 millim.  
Hab. Umzinto, Natal.  
This shell is somewhat similar to *D. æmulus* Smith, but the posterior side is more tumid and rounded.
- ✓ DONAX FABA, *Chemnitz*, Sow., Thes. Conch., vol. iii., p. 312, plate 283, figs. 108, 109.—S. Africa.
- ✓ CYTHEREA (TIVELA) TRANSVERSA, n. sp. (plate 7, figs. 2 and 3). Testa transverse oblonga, crassiuscula, leviter compressa, fere æquilateralis, utrinque leviter acuminate, laeviuscula, nitens, albida, dilute fusco radiata et fasciata; umbones acutiusculi; margo dorsalis utrinque declivis, ventralis arcuatus; cardo normalis; pagina interna roseo tincta; impressiones musculares fuscia, sinu pallii producto.

Antero-post 59, umbono-marg. 42, crass. 24 millim.  
Hab. Natal.

This species may be readily distinguished from its congeners by its transversely oblong form.

CY THEREA (TIVELA) ALUCINANS, Sowerby, n. sp. (plate 7, figs. 5, 6.) Testa subtrigona, leviter inaequilateralis, crassa, laevis, nitens, pallida, lineis fuscis angulatum undulatis parum picta, epidermide tenui cornea induta; umbones acutiusculi, prominentes, paulo ante medium locati; margo dorsalis utrinque declivis, anticus rectiusculus, posticus leviter arcuatus, ventralis arcuatus; area postica planulata, antica levissime concavo depressa; pagina interna alba, postice liraea violacea picta; impressiones musculares parviusculi, sinu pallii lato paulo producto.

Antero-post 63, umbono-marg. 55 millim.  
Hab. Natal.

A worn valve of this species was mistaken by me for *Meretrix zonaria* (Journal of Conchology, vol. 7, p. 14.) I have since seen fine fresh specimens, and find it to be a *Tivela*.

CY THEREA (TIVELA) POLITA, Sowerby, Thes. Conch., vol. ii., p. 618, pl. 127, fig. 14.—Natal. *C. dolabella*, Sowerby, is only a form of this species.

✓ CY THEREA (CARYATIS) MANILLÆ, Sowerby, Thes. Conch., vol. ii., p. 634, pl. 136, figs. 180, 181.—Natal.

✓ CIRCE (CRISTA) DIVARICATA, Chemnitz, Sow.. Thes. Conch., vol. ii., p. 250, pl. 137, figs. 8, 9.—Natal.

✓ LIOCONCHA PICTA, Lamarck, Sow., Thes. Conch. (Cytherea), vol. ii., p. 642, pl. 134, f. 149.—Durban.

✓ SUNETTA CONTEMPTA, Smith, Proc. Zool. Soc., 1891, p. 422.—*Meretrix menstrualis*, Reeve (non. Menke) Conch. Icon., vol. xiv., fig. 9.—Natal.

✓ TAPES SULCARIA, Lamarck, Sow., Thes. Conch., vol. ii., p. 685, pl. 146, f. 30-32.—Durban.

✓ VENUS (CHIONE) LISTERI, Gray, Sow., Thes. Conch., vol. ii., p. 705, pl. 152, f. 7-9.—Durban.

✓ VENUS (CHIONE) ARAKANA, Nevill (*Cryptogramma*), Journ. Asiatic Soc., Bengal, vol. xxxix., part 2, p. 10, pl. 1, f. 16.—Port Elizabeth and Durban.

✓ VENUS (ANAITIS) LATILIRATA, Sowerby, n. sp. (pl. vi. fig. 24). Testa triangularis, crassa, liris concentricis latis

rotundatis et obsolete angulatis instructa, fulvescente alba, maculis numerosis parvis strigisque fuscis et carneo purpurascensibus radiata et aspersa; umbones prominentes, complanati, ante medium locati; latus anticum subtruncatum, posticum longius; lunula parviuscula, area ligamenti late excavata.

Antero-post 25, umbono-marg, 20 millim.

Hab. Durban.

This shell has somewhat the appearance of the West-Indian *V. paphia*, but the ridges are not produced posteriorly into short laminæ as in that species.

- ✓ VENERUPIS LAJONKARI, Payrandeau, Sowerby, Thes. Conch. (*Tapes*), vol. ii., p. 695, pl. 150, f. 120.—Durban. The usual habitat of this species is the Mediterranean, but it does not appear to be common anywhere.
- ✓ CYPRICARDIA ANGULATA, Lamarck, Reeve, Conch. Icon., fig. 2.—Natal.
- ✓ CARDIUM TENUICOSTATUM, Lamarck, Reeve, Conch. Icon. *Cardium*, pl. 10, fig. 50.—Natal. Common in several localities on the Australian coast.
- ✓ CARDIUM RUGOSUM, Lamarck, Reeve, Conch. Icon. *Cardium*, pl. 14, fig. 68.—Natal. Widely distributed in the Indian Ocean, Australia, and South Sea.
- ✓ CARDIUM RUBICUNDUM, Reeve, Conch. Icon. *Cardium*, pl. 9, fig. 44.—Natal.
- ✓ CARDIUM PAPYRACEUM, Chemnitz, Reeve, Conch. Icon., pl. 2, fig. 9.—Natal. China Sea, Philippines, &c.
- ✓ CARDIUM TURTONI, Sowerby, pl. vii., fig. 4, Journal of Conchology, vol. 7, p. 14.—Port Elizabeth.
- ✓ CARDIUM BURNUPI, Sowerby, n. sp., pl. viii., f. 25. Testa subcircularis inflata, crassiscula, straminea, carneo-fusco tincta radiatim costata; costis 29 angustis, sub-remotis anticis fornicato squamatis, medianis serratis, posticis aculeatis.  
Diam antero-post 44, umbono-marg, 43 millim.  
Hab. Durban.  
This species (of which I have only seen odd valves) is intermediate in character between *C lima*, Gmelin, and *C. aculeatum*, Linn.
- ✓ CARDIUM ADAMSI, Reeve, Moll. Voy. Samarang, pl. 22, fig. 2

✓ CHAMA IOSTOMA, Conrad, Reeve, Conch. Icon. (*Chama*), pl. 2, fig. 7.—Natal.

LUCINA EXASPERATA, Reeve, Conch. Icon. (*Lucina*), pl. 1, fig. 4.—Durban.

LUCINA (CODAKIA) PECTEN, Lamarck, Reeve, Conch. Icon., (*Lucina*) sp. 38.—St. John's, Durban.

LUCINA (DIVARICELLA) QUADRISULCATA, D'Orbigny.—*Lucina Eburnea*, Reeve, "Conch. Icon." (*Lucina*), pl. 8, sp. 49,—Natal. A species of wide distribution, Hong Kong, St. Helena, &c.

✓ LORIPES CLAUSUS, Philippi, Abbild. und Beschr. Conch. vol. iii., Lucina pl. 2, fig. 2.—Natal. This is the shell quoted by me, p. 61, as *Loripes lacteus*. There are specimens in the British Museum from Mozambique.

SCINTILLA COMPTA, n. sp. (pl. 6, fig. 28). Testa oblongo-ovalis, compressiuscula, tenuis, æquivalvis, fere æquilateralis, alba, nitens; margo dorsalis utrinque arcuatim declivis; valvis utrinque paulo hiantibus, concentrica leviter et irregulariter rugatis; umbones leviter prominentes, parviusculi.

Antero-post. 12.50, umbono-marg. 8 millim.  
Hab. Durban.

SCINTILLA QUEKETTI, n. sp. (pl. 8, figs. 20, 21). Testa sub-quadrato oblonga, æquilateralis, leviter inflata, alba, nitens, irregulariter concentrica rugata, clausa; umbones vix prominentes, obtusiusculi; margo dorsalis rectiusculus, lateribus leviter rotundatis, margo ventralis levissime arcuatus, vel rectiusculus.

Antero-post 14, umbono-marg. 9.50.  
Hab. Durban (Quekett).

SCINTILLA ELONGATA, n. sp. (pl. 8, fig. 24). Testa auguste oblonga, cylindracea, alba, tenuissima, leviter inæquilateralis, lateribus regulariter rotundatis; umbones vix prominentes, paulo ante medium locati; valvis vix hiantibus.

Antero-post 12, umbono-marg. 5.50 millim.  
Hab. Durban (Burnup).

A very thin white shell, with valves scarcely gaping, distinguished by its narrow elongated form.

SCINTILLA DURBANENSIS, n. sp. (pl. 8, figs. 22, 23). Testa oblongo-ovalis, tenuis, alba, fere æquilateralis; umbones acuti, vix prominentes; margo dorsalis utrinque arcuatim paulo declivis; hiatus ventralis angustus.

Antero-post 9, umbono—marg. 5 millim.  
Hab., Durban (Burnup).

This shell is narrower than *S. Compta*, and has a rather narrow but decided ventral hiatus.

- ✓ SOLENOMYA TOGATA, Poli (= *Mediterranea*, Lamk), Reeve, Conch. Icon. (*Solemya*), fig. 2.—Durban.
- ✓ SEPTIFER NICOBARICUS, Chemnitz, Reeve, Conch. Icon. (*Mytilus*), pl. 9, fig. 42.—Durban.
- ✓ PINNA MADIDA, Reeve, Conch. Icon., pl. 17, fig. 31.—Durban.
- ✓ PINNA SACCATA, Linn., Reeve, Conch. Icon., pl. 4, fig. 6.—Natal.
- ✓ PINNA SERRA, Reeve, Conch. Icon., pl. 23, f. 43.—Durban.
- ✓ PINNA VEXILLUM, Born, Reeve, Conch. Icon., pl. 19, fig. 36.—Durban.
- ✓ PINNA SQUAMOSISSIMA, Philippi, Reeve, Conch. Icon., fig. 2 (as *P. seminuda*, Lam.)
- ✓ AVICULA (MELEAGRINA) FLABELLUM, Reeve, Conch. Icon. (*Avicula*), figs. 7, 8.—Durban. With this species I unite *A. imbricata*, *lacunata* and *muricata*, Reeve.
- ✓ AVICULA ZEBRA, Reeve, Conch. Icon., figs. 3, 6.—Port Elizabeth.
- ✓ MALLEUS TIGRINUS, Reeve, Conch. Icon. (*Malleus*), pl. 3, fig. 7.—Durban.
- ✓ MALLEUS LEGUMEN, Reeve, Conch. Icon., pl. 1, fig. 2.—Durban.
- ✓ PERNA ANOMIOIDES, Reeve, Conch. Icon. (*Perna*), pl. 3, fig. 11.—Durban.
- ✓ ARCA DIVARICATA, Reeve, Conch. Icon. (*Arca*), pl. 16, fig. 108.—Durban.
- ✓ ARCA (BARBATIA) CÆLATA, Reeve, Conch. Icon. (*Arcæ*), fig. 110.—Durban.
- ✓ PECTUNCULUS QUEKETTI, n. sp. (plate 7, figs. 8, 9).—Testa subquadrato rotundata, crassa, convexa, fusca, lineis numerosis albidis radiata, undique subtilissime radiatim striata; umbones centrales prominentes, acutiusculi, fere approximati; area elongata, angusta, sub-profundâ; ligam-

entum crassiusculum; cardo arcuatus, planulatus, latus; dentes 18, plus minusve elongati, rectiusculi; pagina interna alba; impressiones musculares vivide rufo-fusco radiatim tincti; marginibus valde crenulatis.

Antero-post. 65, umbono-marg. 65 m.m.

Hab. Durban.

This shell is quite distinct from the Mediterranean species, and the vivid colouring of the muscular impressions appears to be characteristic, although having only seen one perfect specimen and two or three odd valves, I cannot with certainty pronounce the character invariable.

✓ *PECTEN SQUAMOSUS*, Gmelin, Sow., Thes. Conch., vol. i., pl. 13, figs. 48, 50.—Natal.

✓ *SPONDYLUS DUCALIS*, Chemnitz, Sow., Thes. Conch., vol. i., pl. 85, fig. 16.—Natal.

✓ *SPONDYLUS NICOBARICUS*, Chemnitz, Sow., Thes. Conch., vol. i., pl. 88, fig. 48.—Natal.

✓ *LIMA SQUAMOSA*, Lamarck, Sow., Thes. Conch., vol. i., pl. 21, fig. 1.—Natal. A species of very wide distribution from the Mediterranean to Polynesia.

✓ *LIMA HIANS*, Gmelin, var. *tenuer*, Turton (= *L. fragilis*, Ch.), Reeve, Conch. Icon. (*Lima*), fig. 18.

ANOMIA EPHIPPUM, Linn., Sow., Ill. Index of Brit. Shells, pl. 8., fig. 18.—Durban.

OSTREA LACERATA, Hanley, Proc. Zool. Soc. 1845, p. 106, Conch. Icon., figs. 51 and 78, (as *O. lacerans* and *pes-tigris*)—Limpopo R.

#### BRACHIOPODA.

*Agulhasia Davidoni*, King, Ann. and Mag. Nat. Hist., 1871, vol. vii., p. 111, pl. xi., fig. 1.—Aguilhas Bank.

#### ADDENDA.

(Received too late for figuring).

✓ *APICALIA BIFORMIS*, Sowerby, n. sp.—Testa ovato-pyramidata, solidiuscula, polita, alba, anguste umbilicata, antice globosa, postice acuminata; spira elatiuscula, ad apicem mucronata; sutura angustissime canaliculata; anfractus 10, convexi, primi 5 contracti; ultimus rotundatus; apertura latiuscula; labrum tenue, utrinque leviter sinuatum.

Hab. Durban.

An interesting species of which I have only seen one specimen. The first 5 whorls are narrowly contracted, from thence the spire suddenly widens, becoming regularly pyramidal, the body whorl being shortly rounded at the base. In form this shell resembles certain species of *Stylifer*, but its more opaque substance and smooth polished surface induce me to place it in the genus *Apicalia*, although I doubt whether the generic distinction is of much value. Unfortunately, I omitted to take the dimensions of this shell before returning it to S. Africa, but I think its length is about 12 or 13 m.m., or half an inch.

*PANDORA SIMILIS*, n. sp.—*Testa oblonga, latiuscula, compressa, leviter inaequivalvis, valde inaequilateralis, concentrica irregulariter plicata, valva dextra subconcavo planulata, sinistra leviter convexa; umbones leviter prominentes, conjuncti; margo dorsalis anticus brevis declivis, posticus elongatus, incurvatus, duplicatus, vix rostratus; margo ventralis valde arcuatus.*

Antero-post 16.50, umbono-marg. 10 millim.

Hab. Durban.

This species is broader than *P. inaequivalvis*, it closely resembles *P. rostrata*, Sow., but wants the posterior rostrum characteristic of that species.

*SOLARIELLA SCULPTA*, *Sowerby*, n. sp.—*Testa depresso orbicularis, profunde umbilicata, supra fulvo-viridula, obscure fusco maculata, infra albida, fusco maculata; spira vix elevata; sutura planato depressa, plicata; anfractus 5, valde clathrati, convexi, superne angulati, spiraliter conspicue carinati, liris angustis numerosis obliquis clathrati; anfractus ultimus depresso rotundatus, infra convexus, costis spiralibus numerosis confertis instructus; umbilicus circularis, perspectivus, latiusculus, intus clathratus, extus crenulatus; apertura latiuscula, subcircularis, leviter obliqua, intus margaritacea; labrum tenue.*

Diam. 4.50, alt. 2.25 millim.

Hab. Durban.

A depressed little shell, conspicuously latticed above the periphery, and closely lirate beneath. It has much the appearance of a small species of *Torinia*.

✓ *HAMINEA SUBCYLINDRICA*, *Sowerby*, n. sp.—*Testa oblonga, subcylindrica, tenuissima, alba, postice vix acuminata, leviter umbilicata, undique densissime transversim striata, striis*

utrinque fortioribus; lateribus convexiusculis; apertura mediocriter lata, postice producta, columella rectiuscula.

Hab. Durban.

Long. 8, lat. 4 millim.

A very fragile shell, very finely striated. In form it somewhat resembles *H. brevis*, (*Quoy and Gaim.*), but it is rather more attenuated at the extremities, and consequently less cylindrical.

---

#### NOTE.

✓ *ZIZIPHINUS MULTILIRATUS*, Sow. (P.Z.S., 1875, p. 127, pl. 24, f. 10), is, I think, not S. African. The locality given "Cape of Good Hope," being a mistake.

---

---

## MARINE SHELLS OF SOUTH AFRICA, 1892.

---

### EMENDATIONS, 1897.

- Page 4. For PYRULA read MELONGENA. The genus *Pyrula*, Lam. belongs to the family Dolidae, and is restricted to the group of which *Bulla ficus*, Linn. is the type.
- Page 4. For METULA read TRITONIDEA, and add locality Durban.
- Page 6. For DEFRACTIA AMPLEXA read MANGILIA AMPLEXA.
- Page 6. For DEFRACTIA CAPENSIS read CLATHURELLA CAPENSIS.
- Page 7. For DEFRACTIA read CLATHURELLA.
- Page 7. Omit MANGILIA CLATHRATA. This was wrongly identified, see CLATHURELLA VERRUCOSA, Sow., n. sp.
- Page 7. MANGILIA COSTATA, add var. *coarctata*, Forbes.
- Pages 7-9. For TRITON read LOTORIUM.<sup>1</sup> The well-known generic name of *Triton* (Montfort) having been previously used in another branch of Zoology, it is unfortunately illegal to continue its use in Malacology.
- Page 9. RANELLA ARGUS, add locality Kalk Bay.
- Page 9. RANELLA SEMIGRANOSA, R. CRUMENA, R. PUSILLA.—Durban.

---

<sup>1</sup>LOTORIUM, Montfort, 1810.

- Page 20. Omit *MARGINELLA EPIGRUS*. See *M. BURNUPI*, *Sow.*, n. sp. .
- Page 31. *CYPRÆA FIMBRIATA*, add locality Durban.
- Page 32. *CYPRÆA CITRINA*, add locality Port Elizabeth.
- Page 32. *CYPRÆA HELVOLA*, *C. CARNEOLA*, *C. CAURICA*, *C. MANRITIANA*, and *C Lamarcki*.—Durban.
- Page 33. *CYPRÆA UNDATA*.—Durban.
- Page 35. *CERITHUIM KOCHI*, and *C. MONILIFERUM*.—Durban.
- Page 40. *HIPPONYX AUSTRALIS*.—Durban.
- Page 44. *TROCHUS NIGROPUNCTATUS*.—Durban.
- Page 45. *CLANCULUS PUNICEUS*.—Durban.
- Page 50. Omit *CHITON MARGINATUS*. The specimens referred to are *C. ELIZABETHENSIS*, Pilsbry.
- Pages 50 and 51. The *POLYPLACOPHORA* or *Chitons* are divided into the following sub-genera :—  
*DINOPLAX* for *Chiton gigas*.  
*ACANTHOCHITES* for *Chiton Garnoti*.  
*ISCHNOCHITON* for *Chitons textilis*, *pruniosus*, *pertusus*, *tigrinus*, and *oviscus*.  
*CHÄTOPLEURA* for *Chitons Watsoni*, and *pustulatus*.  
*CALLOCHITON* for *Chiton castaneus*.  
*PLAXIPHORA* for *Chiton Wahlbergi*.
- Page 51. Omit *CHITON CAPENSIS* (is *C. nigrovirens*).
- Page 51. „ *CHITON MAGGILLIVRAVI*.
- Page 51. „ *CHITON CARMICHAELIS*.
- Page 51. For *CHITON NIGROVIRESCENS* read *C. NIGROVIRENS*.
- Page 55. For *MACTRA ADANSONI* read *M. GLABRATA*, *Linn*, and place *M. Adansonii* in the synonymy. Add locality Durban.
- Page 56. *LUTRARIA OBLONGA*.—Natal.
- Page 57. *TELLINA LITTORALIS*.—Sub-genus *MACOMA*.
- Page 58. *CY THEREA (TIVELA) COMPRESSA*.—Remove *C. polita* from synonymy.


- Page 58. Omit *CY THEREA (TIVELA) DOLABELLA* as a species,  
and place it as synonym of *C. POLITA*, *Sowerby*.
- Page 59. *MEROE OVALIS*.—add loc., Durban.
- Page 60. Omit *VENUS LAYARDI*. The shells mistaken for  
this species are *V. ARAKANA*, *Nevill*.
- Page 60. *PETRICOLA TYPICA*.—Add loc., Durban.
- Page 61. *LORIPES GLOBOSUS*.—Add loc., Durban.
- Page 65. *ARCA NIVEA*.—Add loc. Durban.





SOUTH AFRICAN SHELLS.

Plate VI



## DESCRIPTION OF PLATE VI.


- Fig. 1. *Bullia pustulosa*.  
,, 2. *Marginella Ponsonbyi*.  
,, 3. *Columbella Kitchingi*.  
,, 4. " *pyramidalis*.  
,, 5. *Scalaria simplex*.  
,, 6. *Natica Queketti*.  
,, 7. *Latirus abnormis*.  
,, 8. *Cioniscus unilineatus*.  
,, 9. *Eulima simplex*.  
,, 10. *Cioniscus pellucidus*.  
,, 11. *Odostomia lucida*.  
,, 12. *Cerithiopsis lirata*.  
,, 13. " *exquisita*.  
,, 14. *Clathurella verrucosa*.  
,, 15. *Rissoa (Cingula) Caffra*.  
,, 16. *Haminea gracilis*.  
,, 17. *Volvatella laguncula*.  
,, 18. *Natica simplex*.  
,, 19. *Fulgur africanus*.  
,, 20. *Turbo Ponsonbyi*.  
,, 21. *Solariella splendens*.  
,, 22. *Gibbula Tryoni*.  
,, 23. *Ancilla decipiens*.  
,, 24. *Venus latilirata*.  
,, 25. *Tellina (Macoma) candidata*.  
,, 26. *Donax Burnupi*.  
,, 27. *Basterotia obtusa*.  
,, 28. *Scintilla compta*.  
,, 29. *Tellina prismatica*.  
,, 30. *Psammobia Burnupi*.  
,, 31. *Ancilla optima*.  
,, 32. *Strigillia Trotteriana*.  
,, 33. *Pandora dissimilis*.  
,, 34. *Tellina dispar*.  
,, 35. *Marginella Burnupi*.  
,, 36. *Marginella perminima*.





SOUTH AFRICAN SHELLS.


Plate VII



2




1



3




4




5



6



7



8

DESCRIPTION OF PLATE VII.


- Fig. 1. *Bullia similis*.  
,, 2, 3. *Cytherea (Tivela) transversa*.  
,, 4. *Cardium Turtoni*.  
,, 5, 6. *Cytherea (Tivela) alucinans*  
,, 7, 8. *Pectunculus Queketti*.





SOUTH AFRICAN SHELLS

Plate VIII.



### DESCRIPTION OF PLATE VIII.

- Figs. 1, 2. *Pleurotoma (Drillia) Burnupi.*  
" 3. " " Layardi.  
" 4, 5. *Nassa producta.*  
" 6, 7. *Columbella consanguinea.*  
" 8, 9. " Langleyi.  
" 10, 11. *Calliostoma Layardi.*  
" 12, 13. *Bullina oblonga.*  
" 14. *Basterotia tricostalis.*  
" 15. " " (posterior side).  
" 16. *Tellina Queketti.*  
" 17. " " var.  
" 18. *Donax simplex.*  
" 19. " " (posterior).  
" 20. *Scintilla Queketti.*  
" 21. " " (posterior).  
" 22. " *durbanensis.*  
" 23. " " (ventral hiatus).  
" 24. " *elongata.*  
" 25. *Cardium Burnupi.*  
" 26. *Mactra æquisulcata.*



## INDEX TO GENERA.

—o—

|                      | PAGE |        | PAGE                 |     |        |
|----------------------|------|--------|----------------------|-----|--------|
| <i>Acanthochites</i> | ...  | 19     | <i>Cryptogamma</i>   | ... | 24     |
| <i>Acanthopleura</i> | ...  | 19     | <i>Cylindrobulla</i> | ... | 21     |
| <i>Action</i>        | ...  | 22     | <i>Cypraea</i>       | ... | 16     |
| <i>Agulhasia</i>     | ...  | 23     | <i>Cypocardia</i>    | ... | 25     |
| <i>Amphidesma</i>    | ...  | 22     | <i>Cytherea</i>      | ... | 23, 24 |
| <i>Anaitis</i>       | ...  | 24     |                      |     |        |
| <i>Ancilla</i>       | ...  | 7      | <i>Divaricella</i>   | ... | 26     |
| <i>Anomia</i>        | ...  | 28     | <i>Doliium</i>       | ... | 11     |
| <i>Apicalia</i>      | ...  | 28     | <i>Donax</i>         | ... | 23     |
| <i>Aplustrum</i>     | ...  | 20     | <i>Drillia</i>       | ... | 2      |
| <i>Arca</i>          | ...  | 7      |                      |     |        |
| <i>Atys</i>          | ...  | 21     | <i>Engina</i>        | ... | 11     |
| <i>Avicula</i>       | ...  | 27     | <i>Euchelus</i>      | ... | 19     |
|                      |      |        | <i>Eulima</i>        | ... | 14     |
| <i>Basterotia</i>    | ...  | 21     | <i>Euthria</i>       | ... | 2      |
| <i>Bullia</i>        | ...  | 5      |                      |     |        |
| <i>Bullina</i>       | ...  | 20     | <i>Fasciolaria</i>   | ... | 8      |
| <i>Callistoma</i>    | ...  | 18     | <i>Fissurella</i>    | ... | 19     |
| <i>Calyptrea</i>     | ...  | 17     | <i>Fulgor</i>        | ... | 1      |
| <i>Cancellaria</i>   | ...  | 16     | <i>Fusus</i>         | ... | 2      |
| <i>Cardium</i>       | ...  | 25     | <i>Gibbula</i>       | ... | 18     |
| <i>Cassis</i>        | ...  | 11     |                      |     |        |
| <i>Caryatis</i>      | ...  | 24     | <i>Haninea</i>       | ... | 20, 29 |
| <i>Cerithiopsis</i>  | ...  | 14     | <i>Harpa</i>         | ... | 11     |
| <i>Cerithium</i>     | ...  | 16     | <i>Hydatina</i>      | ... | 20     |
| <i>Chama</i>         | ...  | 26     |                      |     |        |
| <i>Chione</i>        | ...  | 24     | <i>Iacra</i>         | ... | 22     |
| <i>Chiton</i>        | ...  | 19, 20 | <i>Ianthina</i>      | ... | 15     |
| <i>Cioniscus</i>     | ...  | 13     | <i>Iopas</i>         | ... | 6      |
| <i>Clanculus</i>     | ...  | 19     | <i>Ischnochiton</i>  | ... | 19     |
| <i>Cluthurella</i>   | ...  | 3      |                      |     |        |
| <i>Codakia</i>       | ...  | 26     | <i>Lagena</i>        | ... | 4      |
| <i>Columbella</i>    | ...  | 10, 11 | <i>Latus</i>         | ... | 8      |
| <i>Cominella</i>     | ...  | 4      | <i>Leptothyra</i>    | ... | 17     |
| <i>Conus</i>         | ...  | 15, 16 | <i>Lima</i>          | ... | 28     |
| <i>Coralliphila</i>  | ...  | 6      | <i>Lioconcha</i>     | ... | 24     |
| <i>Crista</i>        | ...  | 24     | <i>Lotorium</i>      | ... | 4      |

## INDEX TO GENERA.

|                  | PAGE   |                 | PAGE       |
|------------------|--------|-----------------|------------|
| Loripes ...      | 26     | Pyrazus ...     | ... 17     |
| Lucina ...       | 26     | Pyrula ...      | ... 11, 12 |
|                  |        | Raeta ...       | 22         |
| Macrochisma ...  | 19     | Ranella ...     | 4          |
| Macoima ...      | 23     | Ricinula ...    | 6          |
| Mactra ...       | 21     | Rissoa ...      | 17         |
| Mactrinula ...   | 22     |                 |            |
| Malleus ...      | 27     | Scalaria ...    | 12         |
| Magilia ...      | 3      | Scintilla ...   | 26         |
| Marginella ...   | 9, 10  | Semele ...      | 22         |
| Margarita ...    | 18     | Septifer ...    | 27         |
| Meroe ...        | 24     | Sigaretus ...   | 12         |
| Metula ...       | 2      | Sistrium ...    | 6          |
| Mitra ...        | 8, 9   | Solariella ...  | 18, 29     |
| Monilea ...      | 18     | Solarium ...    | 15         |
| Murex ...        | 1      | Solen ...       | 21         |
| Nassa ...        | 5      | Solenomya ...   | 27         |
| Nassaria ...     | 6      | Spondylus ...   | 28         |
| Natica ...       | 12     | Strigililia ... | 23         |
| Naticina ...     | 12     | Strombus ...    | 16         |
|                  |        | Sunetta ...     | 24         |
| Obeliscus ...    | 13     | Tapes ...       | 24         |
| Odostomia ...    | 14     | Tellina ...     | ... 22, 23 |
| Oliva ...        | 7      | Terebra ...     | ... 12, 13 |
| Onithochiton ... | 20     | Tivela ...      | ... 23, 24 |
| Ostrea ...       | 28     | Torinia ...     | 15         |
| Pachypoma ...    | 18     | Triforis ...    | 17         |
| Pecten ...       | 28     | Trivia ...      | 16         |
| Pectunculus ...  | 27     | Triton ...      | 4          |
| Pandora ...      | 21, 29 | Turbanilla ...  | 14         |
| Peristernia ...  | 8      | Turbo ...       | 17         |
| Perna ...        | 27     | Turritella ...  | 17         |
| Persona ...      | 4      | Umbonium ...    | 19         |
| Philippia ...    | 15     | Vanikoro ...    | 12         |
| Pinna ...        | 27     | Venerupis ...   | ... 25     |
| Pisania ...      | 2      | Venus ...       | 24         |
| Pleurotoma ...   | 2, 3   | Voluta ...      | 8          |
| Priere ...       | 6      | Volvatella ...  | 21         |
| Psammobia ...    | 22     | Xenophora ...   | 17         |
| Purpura ...      | 6      |                 |            |
| Pyramidella ...  | 13     |                 |            |



