



**Marine Species Survey of Johnston Atoll,
Central Pacific Ocean, June 2000**

**Marine Species Survey of Johnston Atoll,
Central Pacific Ocean, June 2000**

A report to

U. S. Fish and Wildlife Service
Pacific Islands Area Office
Honolulu, Hawaii

S. L. Coles¹
R. C. DeFelice^{1,2}
D. Minton²

¹Hawaii Biological Survey
Bishop Museum
Honolulu, Hawaii

²Department of Zoology
University of Hawaii
Honolulu, Hawaii

Bishop Museum Technical Report No. 19

January, 2001

Published by
Bishop Museum Press
1525 Bernice Street
Honolulu, Hawai'i



Copyright © 2001 Bishop Museum
All Rights Reserved
Printed in the United States of America

ISSN 1085-455X

Contribution No. 2001-001 to the Hawaii Biological Survey

EXECUTIVE SUMMARY

The marine biota of Johnston Atoll was surveyed for nonindigenous species in June, 2000 with observations and collections made by investigators using Scuba. Eleven stations were surveyed, including sites at the Johnston Island pier, the north and south lagoon and the outer reef slope. A total of 668 taxa were determined, with 462 of these identified to species. This compares with 865 taxa reported by the 21 previous studies that have been made at Johnston Atoll. Three hundred twenty five, or 49%, of the taxa of the present study were new reports for the Atoll. Most of the previous reports were for macroalgae, corals and fishes, and the present study has the first reports that have been made for organisms in the major groups Porifera, Hydroida, Sipunculida, Bryozoa and Ascidiacea, and of the reef coral *Montipora hoffmeisteri*. Most of the species found in the present study (91%) are known from Hawaii, similar to findings of previous studies.

Little difference was found among the 11 stations in terms of numbers of total taxa, taxa in major taxonomic groups or Shannon's H'_{10} diversity indices. Similarity analysis using the Sorensen's Index of the total taxa among stations showed no clusters grouping above 55% and indicated no unique clusters of stations. The general result indicated by these analyses was of a well mixed system with little variation among sites. However, on-site observations clearly showed that the station on the south side of Johnston Island in the vicinity of the island's sewer outfall was highly eutrophicated, with a heavy bloom of the alga *Byopsis hypnoides* over most of the reef surface and sand channels. However, this infestation appeared to have only a limited impact on other reef organisms, since the site had an intermediate total number of other taxa compared to the other ten stations, including seven species of reef corals, although in relatively low abundance.

Only ten nonindigenous or cryptogenic species were found at Johnston Atoll, and these usually occurred as single specimens at one or two stations. Compared with total reports for this study, nonindigenous or cryptogenic species comprised only 1.5% of the total taxa, or 2% of those identified to species. These introduced or potentially introduced organisms were hydroids, polychaete, bryozoans and ascidians, all of which are likely to have come as fouling on vessels arriving in the Atoll over the last 60 years. This proportion of introduced species to total biota is in good agreement with values that have been determined for Midway Atoll and Kaho'olawe in the Hawaiian Islands, and for studies that have been made in Guam and Australia. The proportion of introductions is much less than the approximate 20% value that has been determined for Oahu harbors.

TABLE OF CONTENTS

	Page
EXECUTIVE SUMMARY	iv
INTRODUCTION	1
HISTORY OF USE	2
PHYSICAL ENVIRONMENT AND SITE DESCRIPTION	3
METHODS	4
RESULTS	
<u>Station Descriptions</u>	7
<u>Previous Species Reports</u>	10
<u>Current Study</u>	11
DISCUSSION	15
CONCLUSIONS	17
ACKNOWLEDGMENTS	18
REFERENCES	18
PLATES	23
APPENDICES	
A. Previous Reports of Marine Organisms Observed or Collected at Johnston Atoll.	25
B. Previous Reports of Marine Organisms Observed or Collected at Johnston Atoll with 1-2 Listings.	46
C. Marine Organisms Observed or Collected in Present Study Listed by Station.	48
TABLES	
1. Sampling station information	5
2. Total numbers of taxa of marine organisms reported by 21 previous studies at Johnston Atoll and by present study, numbers of new reports, identified species and proportion of Hawaiian species component.	11
3. Cryptogenic and nonindigenous species found on June 2000 Johnston Atoll Surveys.	12
4. Data for Figure 2 and Shannon's H'_{10} diversity indices.	13
FIGURES	
1. Map of Johnston Atoll showing locations of sampling Stations 1 to 11.	5
2. Numbers of taxa in major taxonomic groups at Johnston Atoll stations.	13
3. Dendrograph of Sorenson similarities among station for all taxa using UPGMA method for calculating intercluster distances	14

INTRODUCTION

Johnston Atoll lies approximately 1325 km southwest of Honolulu Hawaii in the vicinity of 16° 45' N, 169° 31' W in the north central Pacific Ocean (Figure 1). The nearest other landfalls are French Frigate Shoals, over 800 km to the north-northeast, and the Line Islands, over 1500 km to the southeast. Johnston Atoll is thus one of the most isolated landmasses in the world. It consists of two islands that have been greatly enlarged by dredging and filling and two islands that were totally created by this process. The total land area of the four islands at present is approximately 2.63 km², more than 10 times the area of the original natural islands. Most of this was added in dredge and fill operations completed in 1963-64, which created Akau (North) Island and Hikina (East) Island and enlarged Sand Island from about 0.04 to 0.16 km² and Johnston Island from about 0.85 to 2.30 km². Johnston had previously been enlarged in 1949-50 from its original size of 0.19 km² (Amerson 1973). No original shorelines therefore exist at Johnston Atoll and the topography of the lagoon has been highly modified by dredging.

Because of its remote location and restriction from most public access for the last 50 years, relatively little information has been available for most marine taxonomic groups at Johnston Atoll. The most comprehensive sampling of invertebrates was done by the members of the Tanager Expedition in 1923 which concentrated on crustaceans and echinoderms, reported in Edmondson et al. (1925) and Clark (1949). More is known about the reef fish and nearshore pelagic fishes of Johnston Island from expeditions dating back as far as the 1880s (Smith and Swain 1882; Fowler and Ball 1925; Schultz and Collaborators 1953; Halstead and Bunker 1954; Gosline 1955; Brock et al. 1965; Brock et al. 1966; Randall et al. 1985; Kosaki et al. 1991; Chave and Mundy 1994), and reef corals are well known from studies conducted since the late 1940s (Wells 1954; Grigg 1981; Cairns 1984; Maragos and Jokiel 1986; Jokiel and Tyler 1992). Shallow reef macroalgae were comprehensively surveyed in 1964 (Buggeln and Tsuda 1966) and deep algae surveyed using a submersible in 1983 (Agegian 1985). All information for the environment and biota of Johnston Atoll prior to 1973 was assembled and synthesized in two reports (Amerson 1973; Amerson and Shelton 1976) and information for deep-water invertebrates and fishes is summarized in (Chave and Malahoff 1998). Other pertinent marine biota related studies or surveys conducted at Johnston Atoll are (Moul 1964; Baker et al. 1997) for benthic algae; (Bailey-Brock 1976; Ward 1981) for polychaetes; (Kay 1961; Brock 1973; Brock 1979) for mollusks and the spiny lobster; (Jones 1967; Randall 1972; Randall 1977; Lobel 1984; Randall and Ralston 1984; Ralston, et al. 1985; Anderson 1986; Irons 1989; Kosaki 1989; Irons 1990; Winterbottom and Burrige 1993; Dee and Parrish 1994; Lobel and Mann 1995; Mann and Lobel 1995; Kerr and Lobel 1997; Lobel 1997; McCosker and Smith 1997; Sancho et al. 1997; Economakis and Lobel 1998; Randall 1999) for fishes and (Balazs 1986; Balazs 1986; Ackman et al. 1992; Balazs 1994) for sea turtles.

As the agency responsible for stewardship of the atoll, the U.S. Fish and Wildlife Service (USFWS) is concerned with evaluation of the present status of atoll organism populations, including those which inhabit the marine coral reef ecosystem. With the high degree of marine traffic that has frequented the atoll in the last 60 years, there has been ample opportunity for introductions of nonindigenous marine species to the area that may have developed resident populations that could compete or displace native biota. Staff of the Bishop Museum have been conducting studies of nonindigenous marine species over the past five years in harbors on Oahu (Coles et al. 1997; Coles et al. 1999a, 1999b), the island of

Kaho'olawe (Coles et al. 1998) and Midway Atoll (DeFelice et al. 1998) which have shown various levels of nonindigenous species occurrence at these locations in the Hawaiian chain. In order to determine whether such introductions have occurred at Johnston Atoll, and to develop a current baseline of knowledge of the marine biotic community at Johnston, the present study was conducted in June 2000.

HISTORY OF USE

There are no reports of discovery or habitation of Johnston Atoll by Polynesians and, although the atoll may have been sighted by Spanish sailors (Amerson and Shelton 1976), its first recorded discovery was by the American brig *Sally* on September 2, 1796. First landing was by Captain J. Johnston of the HMS *Cornwallis*, for whom the atoll was later named, on December 14, 1807 (Amerson 1973; Amerson and Shelton 1976). The first habitation and use of the atoll was for mining of bird guano in 1858-60. Following the end of this enterprise, visits were few until the first scientific expedition, that of the *Tanager-Whippoorwill* in 1923. Much of the focus of this expedition was on birds, which led to the atoll being made a federal bird refuge by executive order on July 29, 1926. Responsibility for stewardship of the atoll was originally placed with U. S. Department of Agriculture and passed to the Department of Interior in 1940. However, "ownership" from 1934 to 1973 was designated by executive orders to be with the Department of Navy, while operational control was at various times with the U. S. Navy, the U. S. Air Force, Joint Task Forces (JTC) 7 and 8, the Atomic Energy Commission (AEC) and the Defense Nuclear Agency (DNA) (Amerson and Shelton 1976).

Development at Johnston Atoll began shortly after a contract was awarded in 1939 for construction of a small navy base, initially composed of a lagoon seaplane landing area and headquarters on Sand Island. The naval air station was commissioned in August 1941 and was shelled briefly by the Japanese in December 1941, causing extensive damage to the facilities. These were reconstructed and substantial dredging was conducted during WW II of channel approaches and a seaplane runway, and the atoll was also used as refueling base for Pacific submarines. By the end of the war Johnston was one of the busiest air terminals in the Pacific (Amerson 1973; Amerson and Shelton 1976).

Following WW II, activity decreased and Sand Island was abandoned in 1946. Operational control was transferred to the U. S. Air Force in 1948, and activity resumed in 1951-52 during the Korean War when the Johnston airstrip was enlarged and new buildings were built on Johnston Island. A LORAN transmitting station and a weather station were established on Johnston Island in 1957. The LORAN station and Coast Guard facility were transferred to Sand Island in 1959 and 1961, respectively. A U. S. Air Force tracking station was established on Sand Island in 1964.

In 1958 a new era of operations began with assumption of operational control by the commander of Joint Task Force 7 to conduct stratospheric testing of thermonuclear devices. Two >1 megaton bombs were exploded in 1958, the first on August 1 at an altitude of 76 km and the second on August 12 at approximately 33 km. Two more devices were planned for testing in 1962, and one 1.5 bomb was exploded at 200 km altitude on July 9, while the other planned for July 25 was aborted when the test missile was destroyed. Subsequently Johnston Atoll came under the joint operational control of JTF 8 and the AEC as a headquarters and base of operations in the Pacific in case the Nuclear Test Ban Treaty

were nullified, and the island supported personnel of the AIR Defense Command and Pacific Missile Range. In 1973 operational control passed to the Defense Nuclear Agency (DNA). In 1976 the USFWS and the DNA completed a Memorandum of Agreement granting USFWS jurisdiction and responsibility over the atoll's natural resources, including all waters and coral reefs within three nautical miles of the atoll's land areas.

In 1985 a permit was issued under the Resource Conservation and Recovery Act (RCRA) to build and operate the Johnston Atoll Chemical Disposal System (JACADS) facility to incinerate the U. S. Army's stocks of chemical agents and munitions that were stored on Johnston Island. The JACADS facility is located at the west end of the island, downwind from the prevailing trade winds. It has entered the final stages of agent destruction and is preparing to undergo closure activities in the next two years, and the population on the atoll will decrease dramatically from its present number of over 1000 personnel. The atoll will thereupon revert to its former status of a remote natural wildlife preserve with usage directed toward maintaining undisturbed populations of resident organisms, with a yet to be determined level of human visitors.

This brief history indicates that although remote, Johnston Atoll has been highly utilized for various military and civilian activities over much of the past century, and this usage has provided ample opportunity for introduction of nonindigenous species by supply and transport vessels. At present a supply barge visits Johnston Atoll monthly that moves between Honolulu, Johnston Atoll and Kwajalein Atoll. The two small islands that originally were at the atoll have been greatly enlarged and altered, and the highly modified shorelines of the four islands now present consist primarily of concrete structures that are highly suitable for settlement of fouling organisms that could be transported by ship traffic.

PHYSICAL ENVIRONMENT AND SITE DESCRIPTION

Johnston Atoll consists of a shallow coral reef platform approximately 130 km² in area (Figure 1) composed of alternating sand, loose coral and large formations of live coral, especially of the genus *Acropora*. Unlike most coral atolls, the lagoon at Johnston is not enclosed by a ring of reef and islands but only has a protective reef along its northwest margin. This is on a broad shallow ridge that extends from the west end of the marginal reef eastward about 14 km and protects a shallow lagoon about 11 km X 3 km from the predominant waves from the northwest. Depths in the lagoon vary about from about 3 to 10 m. The atoll is not enclosed because the platform on which it sits has subsided and is tilting to the southeast. Most of the reef platform lies outside of the lagoon, extending about 19 km east-southeast and 8 km south of Johnston Island. The platform slopes gently to about 7 to 18 m depth then much more steeply to 180 m (Emery 1956; Amerson and Shelton 1976; Anderson 1986; Jokiel and Tyler 1992).

The oceanographic environment at Johnston is typical of coral reef conditions, with a small and moderate annual temperature range of about 25-27°C, clear oceanic water and surface salinity of 34.6-34.8 ‰ (Wennekens 1969; Amerson 1973). Johnston Atoll lies near the southern edge of the North Pacific gyre and is in the zone of the North Pacific Equatorial Current which moves water past the southern Hawaiian Islands to the atoll (Amerson and Shelton 1976; Maragos and Jokiel 1986). The atoll is also probably

affected by the eastwardly flowing North Equatorial Countercurrent, which brings water from more tropical regions (Grigg 1981).

Although the Johnston Atoll reef faces northwest and would therefore normally be on the leeward side of waves generated by the prevailing northeast tradewinds, these waves are influenced by shoaling on shallow areas east of Johnston Island, which refracts them around the platform to break on the northwest side of the reef. The prevailing currents, influenced by both tradewinds and North Pacific equatorial circulation have net flow to the west both inside and outside of the lagoon, stronger during winter months than in summer. Breaking of waves over the reef and entering the lagoon creates a “pumping action” (Jokiel and Tyler 1992) that augments prevailing oceanic currents to generate a westward flowing current. Tides are mixed and semi-diurnal, with strong daily inequalities and maximum amplitudes for highest tides of about 1m. The effect of tides on circulation is to deflect currents to the left on rising tides and to the right on falling tides, and falling tides can reverse the usual westerly flow during weaker circulation in summer months. Ship channels to the southeast and southwest also influence current flow by providing for movement of water out of the lagoon (Wennekens 1969; Amerson and Shelton 1976; Anderson 1986; Jokiel and Tyler 1992).

METHODS

Samples were collected from 11 sites at Johnston Atoll (Figure 1) using methods previously employed on nonindigenous species surveys in Hawaii and Midway (Coles et al. 1997; Coles et al. 1998; Coles et al. 1999a; Coles et al. 1999b; DeFelice et al. 1998). Some of the sampling stations were located to be near sites previously sampled by (Brock et al. 1965; Brock et al. 1966) and (Jokiel and Tyler 1992), others were located in important sites such as in the harbor, along dredge channels or near the Johnston Island sewage outfall. Sampling station locations, dates coordinates and depths are summarized in Table 1.

Collections and observations were made by three experienced investigators sampling as large a variety of habitats as possible at each station while using Scuba. Two divers sampled organisms growing on hard surfaces and in sediments from the intertidal zone to the base of the reef. The third diver recorded the identities of fishes swimming in the area, noted the presence of abundant invertebrate macrofauna and macroalgae, and photographed and collected corals for identification of both the corals and their commensal organisms. Macro-organisms were collected by hand, hard surfaces were scraped with a chisel, and several liters of coral rubble were placed in an 80 µm mesh bag and transported back to the laboratory for later inspection and removal of cryptic organisms. Where sediments were present at a station, 500 cc of sand was collected in plastic bags and later air-dried prior to hand sorting for micromollusks under a dissecting microscope. A subsample of 20 cc was sorted from each sample. When present at a survey site, whole macroalgae plants were collected and preserved in 70% ethanol, and epiphytic organisms were later rinsed from the algae and preserved in ethanol for future processing. Collected organisms, which range 4-8 liters in total volume for each station were inspected on site, and selected




Figure 1. Map of Johnston Atoll showing locations of sampling Stations 1 to 11.

Table 1 Sampling station information

Station	Location	Date	Latitude N	Longitude W	Depth (m)
1	Pier	16-Jun-00	16°44'7.0"	169°31'42.3"	0-12
2	Lagoon (Nr. Brock Sta. 3)	16-Jun-00	16°44'15.5"	169°31'32.5"	3-5.5
3	Lagoon (Nr. Brock Sta. 1)	17-Jun-00	16°45'53.3"	169°31'2.4"	2-3.5
4	Slope, Donovans Reef	17-Jun-00	16°47'10.0"	169°27'58"	8.5-31.5
5	Lagoon (Range Finder)	18-Jun-00	16°44'52.3"	169°31'3.5"	2-9.5
6	NE of Sand Island	18-Jun-00	16°44'51.6"	169°30'28.9"	1.5
7	E of Johnston island	18-Jun-00	16°44'18.0"	169°31'0"	0.5-9
8	Reef Slope	19-Jun-00	16°46'47.2"	169°29'28.7"	3-20
9	Lagoon (Nr.Brock Sta. 2)	19-Jun-00	16°45'29.5"	169°31'29.3"	1-5
10	South Channel	20-Jun-00	16°42'43.7"	169°31'2.4"	2-9.5
11	Sewage Outfall	20-Jun-00	16°43'18.0"	169°31'58"	3-10.0

hydroids, anemones and tunicates were removed and relaxed in a solution of Epsom salts and seawater before preserving in 5% formalin. The remaining organisms were preserved on site in 70% alcohol before returning the samples to the laboratory for sorting and identification of organisms.

Specimens collected were sorted and identified to species or the lowest practicable taxa, using dissecting or compound microscope magnification when necessary. Identifications were made using descriptions available in Reef and Shore Fauna of Hawai'i Sections 1 to 4 (published), 5 and 6 (unpublished), various taxonomic references, and voucher specimens in the Bishop Museum collections. Specimens from various groups were sent to taxonomic experts for final identifications (see Acknowledgments).

All available sources of information for the marine environment of Johnston Atoll were investigated for previous reports of marine organisms collected or sighted at the atoll. Literature consulted included published papers in the open scientific literature, taxonomy-based monographs and reports and unpublished reports for environmental studies. Resources that were consulted in this search were the libraries of Bishop Museum and the University of Hawaii, and literature search engines such as the Zoological Record, Biological and Oceanic Abstracts and the Aquatic Sciences and Fisheries Abstracts (ASFA) available on the University of Hawaii Library web site. The Bishop Museum invertebrate collection database was reviewed for all marine or estuarine organisms indicated to have been collected at Johnston Atoll. The retrieved data were assembled into a combined database containing taxa identity, taxonomic authority, and collection date, when available. Species identified from the present study were compared with listings of known introductions for Hawaii (Carlton and Eldredge in prep.) and the Pacific (Eldredge, 1987) and results of other field studies of nonindigenous species [Coles et al. 1997; Coles et al. 1998; 1999a, 1999b; DeFelice et al. 1998].

The Sorenson's Index of percent similarity, based on presence-absence of species at station pairs, was used to measure the degree of association between stations. By this index, the more species two stations share relative to their total species complements, the greater their ecological similarity. Based on a matrix of Sorensen Index values, cluster analysis was used to arrange stations into groups or clusters. Intercluster distances were calculated using an unweighted pair group average (UPGMA) method. In this analysis, similar stations will form clusters distinct from other stations. These clusters are arranged in a hierarchical, treelike structure called a dendrogram based upon the station pairs similarities. Calculation of the similarity measures and cluster analysis were performed using the Multi Variate Statistical Package (MVSP), Version 3.0 (Kovach 1998).

RESULTS

Station Descriptions

Collections and observations were made at 11 stations, with one of these at the Johnston Island pier, nine on lagoon and outer slope reefs, and one near the sewer outfall on the south side of Johnston Island (Figure 1). Dates of sampling and geographic coordinates for these stations are shown in Table 1. Descriptions of the environment at each station are as follows:

Station 1 (Johnston Island Pier). Near the east end of the north side of Johnston Island, the pier is the docking site for all ships and barges that supply the atoll. The pier is made of interlocked vertical metal sheet piling rising out a depth of about 12 m depth, and the pier surfaces is quite clean of fouling organisms, with few of the barnacles, sponges, bryozoans and tunicates that usually cover pier surfaces in Hawaiian harbors. The substratum outside of the pier is coral rubble and coarse sand, with abundant coral interspersed among the sand patches mostly composed of *Montipora capitata*. Mollusks found at this site include 36 common taxa which inhabit rubble/algae areas. High numbers of cerithids, limpets such as *Diodora granifera*, chitons, and nudibranchs were found, all characteristic of this habitat type. This was one of two sites (the other being site 7) with the vermetid *Dendropoma platypus*, which seems to favor the flat surfaces associated with man-made structures. This site also had two species of littorinid, *Littoraria pintado* and *Littoraria scabra*, due to the site's close proximity to the shore. Fishes were quite abundant, especially in the coral offshore of the pier, with 25 species recorded.

Station 2 (Near Brock et al. 1965 Sta. 3). In the lagoon seaward of Johnston Island near inner reef edge. The substratum is primarily live linear *Montipora capitata* reefs and *Acropora cythera* tables divided by sand channels 3 to 6 m deep. A single *A cythera* table was observed to have an number of aberrant skeletal growths on its surface that may indicate hyperplasia and possible tumor formation (Plate 1). This site contained 69 mollusk species, the highest of any area surveyed, which can probably be attributed to the diversity of habitat types at this location. This diverse community contained common Indo-Pacific reef flat species like *Vasum turbinelles* and *V. ceramicum*, which are among the most common reef flat mollusks in the Pacific, numerous micromollusks and a high diversity of common Pacific reef flat bivalves. These coincided with the extensive patches of coral sand found at this site, a feature uncommon at most of the other survey sites. By contrast with the diverse mollusk community, 32 fish species occurred, one of the lowest values found at the reef stations.

Station 3 (Near Brock et al. 1965 Sta. 1). Another lagoon station, located north of Akau (North) Island in 2-3 m depth. The area had virtually 100% coral cover, almost entirely composed of tabular *Acropora cythera* showing annular growth bands (Plate 2) and growing in a variety of beautiful formations (Plate 3). Almost no sand channels were

present. Probably as a result, mollusk diversity was intermediate among the reef stations, with only 35 species found. Of these species, several unusual micro-bivalves were noted, particularly *Crenella* sp. and *Cratis* sp. (*kaneko?*), two species that have been found in submarine caves off Okinawa. These cave bivalves, among others, are often found in deepwater sediments off Oahu (Swartz et al. 2000, but are not reported from shallow water sediments. These bivalves were collected from a single, small sand patch that was located underneath extensive *Acropora* cover (almost 100% cover). This “cave-like” environment may explain their presence at his site and no other. Fish diversity was also intermediate among the reef stations, with 36 species sighted.

Station 4 (Donovan’s Reef). This area, approximately 8 km northeast of Johnston Island, is a popular dive site with residents and the most remote from usual activity on the atoll. The station was located seaward of the isolated reef which is about 1 km east of the main Johnston Atoll reef. The relief of the upper part of the outer reef slope is very gradual from 8.5 to 21 m depth, at which the slope becomes almost vertical to the reef abase at about 32 m. Observations were made throughout the entire depth range. The substratum is consolidated limestone with moderate coral cover and little relief, with coarse sand at the reef base. Coral diversity was among the highest found in the study, with 11 species found. Diversity of mollusks was moderate with 39 species. Cone shells, particularly *Conus lividus*, *C. rattus*, and *C. miles*, were common on the current-swept pavement regions. All three of these species are common cones on Indo-Pacific reef flats, and they tend to favor areas with exposed/bare substrate. Fish diversity was the highest of any site, with 52 species found. These included the only sighting for the study of the Johnston Atoll endemic rainbow angelfish *Centropyge nahackyi* (Plate 4), which was observed and photographed under a ledge near the bottom of the outer reef slope, and of the whaleshark *Rhincodon typus*, which was observed near the surface above the reef edge.

Station 5 (Near Brock et al. 1965 Sta. 7). The station was near a range finder marking the main channel thorough the lagoon. Substratum over 2 to 9.5 m depth consisted of the concrete surface of range finder pilings, high coverage of coral dominated by *Acropora cythera*, *Montipora capitata* and *M. patula*, and coarse sand in the channel and among the coral formations. This site had the second highest diversity of mollusks, composed of 52 species with a large number of bivalves and several sand dwelling species including two species of tellinid and two species of mesodesmatids. Also present at this site was a cave species, *Barbatia decorata*. This species was described from submarine caves off Okinawa, but may also occur in deep water sediments off Hawaii. Fish diversity was relatively high, with 39 species.

Station 6 (Shallow area NE of Sand island). An area of sparse coral cover on a flat limestone bottom only about 1.5 m deep, with a few concrete structures and interspersed areas of sand. Despite the low coral coverage, diversity was among the highest of any station, with 11 species found. Diversity of mollusks was relatively low (25 species) and

composed of common Indo-Pacific species. Sand dwellers such as nassarids were common. Cerithids, especially *Cerithium nesioticum*, were common in the rubble and the on the pavement. Fish diversity was the lowest of any station, with only 20 species observed.

Station 7_(Near Brock et al. 1965 Sta. 8). Along the eastern side of the dredged channel east of Johnston Island and adjacent to a shallow reef area with numerous, repeating linear reefs mostly composed of *Montipora* and *Acropora* species divided by narrow sand channels. Depth ranged 0.5 m on top of the linear reefs to around 10 m at the edge of the dredged channel, with a maximum depth of 12 m. Coral diversity was high with 11 species, but mollusk diversity low with only 19 species, the second lowest station. Thirty eight species of fish were observed, the fourth highest of all the stations.

Station 8 (Outer Reef Slope). The station was located in a break between the main reef and a smaller reef to the northeast, on the outer reef slope. Like Station 4 at Donovan's Reef, the outer slope is gradual from down to about 20 m depth then breaks to a nearly vertical face extending to the reef base. Observations were made from about 3 m depth to the slope break. This area has low relief and mostly dead coral rubble and consolidated limestone until near the slope break where coral becomes more abundant. An exception to this condition occurred in the shallow area near the start of the transect made down the slope, where a prolific growth of *Acropora cythera* provides extensive relief that supports an abundant population of the yellow tang *Zebrasoma flavescens* (Plate 5). This was the area of highest diversity for corals, with 12 species found, but with only 27 species of mollusks. This modest mollusk diversity probably does not typify the area because collection time was restricted and nearly every species collected was a small sand dweller. Large common reef mollusks were present however, including *Cypraea tigris*, and several common thaidids (*Drupella ochrostoma* and *Morula uva*).

Station 9 (Cucumber Flats). Another popular dive site for Johnston Island residents, this site gets its name from the abundant holothurians that occur in the coarse sand among the reefs. Coral is also abundant and diverse (Plate 6), with 11 species found, including *Montipora hoffmeisteri* (Wells, 1956), the first report of this species from Johnston Atoll. Thirty-two species of mollusks were found, including the holothurian parasite *Balcis*, numerous sand dwellers including *Conus pulicaris*, various nassarids, and tellinid bivalves. The large number of bivalve species (7 species) is attributable to the extensive sand patches at this site. With the exception of *Isognomon perna*, all of the bivalves were infaunal species. Fish diversity was second lowest of any reef station, with only 25 species observed.

Station 10 (South Channel). The most southerly located station of the study, this site is at the east edge of the channel dredged in 1963-64 for fill material used in increasing the size of Johnston Island. The substratum is mostly rubble cobbles next to a reef with high coverage of *Acropora cythera* (Plates 7 and 8) and coarse sand in the sand channel. Depths range from 2 m on top of the reef to 9.5 m in the sand channel at the edge of the

reef. The *A. cythera* tables can be very large, e.g. one the upper surface of one colony was approximately 6 m in longest diameter. This station had the lowest diversity of both corals (6 species) and mollusks (16 species), but fish diversity was intermediate for the reef stations, with 37 species observed.

Station 11 (Sewage Outfall). The site is along the sewage discharge pipe that extends from the treatment facility on the south jetty of Johnston Island to the point of effluent discharge about 500 m south of the jetty shore. The pipe is approximate 0.5 m in diameter and passes through and over numerous sand patches and reef areas along its route. Unlike all the other stations that had high water clarity and virtually no large macroalgae, algal blooms dominated the benthos at this site. By far the dominant species was the chlorophyte *Bryopsis hypnoides*, which overgrew most hard surfaces on the reefs and collected in mass deposits in sand channels. Another common and obvious species was *Caulerpa racemosa*, large growths of which occurred near the shoreline. The area is therefore under obvious stress from nutrients released from the sewage outfall, which is causing eutrophication in the lagoon south of Johnston Island. The impact of this eutrophication extends as far east and south as Station 10, where *Bryopsis hypnoides* was also observed, although in low abundance.

Despite this eutrophication impact, 7 species of corals occurred at Station 11 in moderate abundance. This site had 36 mollusk species, but these had an unusual composition. Few macro-mollusks were observed at this site; the fauna was almost exclusive micromollusks. The site also had a high number of pyramidellid species, a feature common to high nutrient locations around Hawaii (Swartz et al. 2000), supporting a conclusion of an impact related to the outfall. This conclusion is also sustained by the number of mollusk individuals collected at this site, which had at least four times as many individuals as any other station. Only 30 species of fishes were observed at this station, suggesting that the outfall's impact was reducing the diversity of the fish community, but this effect may also be in part due to the reduced visibility at this station compared the other sites.

Previous Species Reports

A total of 865 taxa were reported by 21 previous surveys and studies at Johnston Atoll or are in the Bishop Museum collections (Appendix Table A), with an additional 24 papers or unpublished reports listing one or two species. The species listed in Appendix A are summarized in Table 2 as follows: 91 macroalgae, 57 cnidarian, 1 aschelminth, 20 polychaetes, 112 crustaceans, 221 molluscs, 56 echinoderms, 306 fishes and 1 reptile. The greatest number of species for each of the previous 21 studies or surveys were 148 for Brock et al. (1965); 149 each for Amerson (1973) and Amerson and Shelton (1976), and 195 by Kay (Unpublished). One hundred sixty-five voucher specimens for these and other studies are in the Bishop Museum collections. As indicated above, the previous reports are highly dominated by fishes, which were the main focus of many of the investigations (Smith and Swain 1982; Fowler and Ball 1925; Schultz and

collaborators 1953; Halsted and Bunker; Brock et al. 1965, 1966; Randall et al. 1985; Kosaki et al. 1991). The second most frequently reported group was the Cnidaria, primarily corals (Wells, 1954; Brock et al. 1965, 1966; Amerson 1973; Amerson and Shelton 1976; Grigg 1981; Cairns 1984; Agegian 1985; Maragos and Jokiel 1986; Jokiel 1992; Cohen 1997).

Table 2. Total numbers of taxa of marine organisms reported by 21 previous studies at Johnston Atoll and by present study, numbers of new reports, identified species and proportion of Hawaiian species component.

Taxa	Previous Reports	Present Study	1 st J. A. Report	% New Reports	Identified Species	Hawaiian Species	% Hawaiian
Macroalgae	91	100	69	69	73	73	100
Porifera	0	16	16	100	1	1	100
Cnidaria	57	28	5	18	23	16	70
Aschelminthes	1	0	-	-	-	-	-
Polychaeta	20	59	48	81	20	18	90
Sipunculida	0	4	4	100	4	4	100
Crustacea	112	135	96	71	79	69	87
Mollusca	221	178	55	31	126	115	91
Bryozoa	0	12	12	100	11	7	63
Echinodermata	56	30	7	23	26	24	92
Ascidacea	0	13	13	100	7	4	57
Fish	306	93	0	0	92	90	97
Reptilia	1	0	-	-	-	-	-
Total Taxa	865	668	325	49	462	421	91

Current Study

A total of 668 taxa of macroalgae, invertebrates and fishes were observed or collected at the 11 stations in this survey, (Appendix C), or equal to about 77% of the total number of taxa that had been reported by all previous surveys and studies at Johnston Atoll since 1882 (Table 2). For many taxonomic groups i.e. sponges (16 taxa), sipunculids (4 taxa) bryozoans (30 taxa) and ascidians (13 taxa) the present study provides the first reports and records for the atoll. No new fish taxa were found in the present study, and only five new cnidarians were found. These were the reef coral *Montipora hoffmeisteri*, previously not reported north of Fanning Island (Coles, in press) and four hydroids, one of them the nonindigenous *Pennaria disticha*. For the remaining groups of macroalgae, polychaetes, crustaceans, molluscs, and echinoderms, the number of taxa newly reported taxa range from 13 (echinoderms) to 96 (crustaceans). Alternatively, new reports expressed as a percentage of total taxa range from 23% (echinoderms) to 81% (polychaetes), with nearly half (48%) of the total taxa being previously unreported for the atoll. Within the crustacea, in which new reports were 71 % of the total, some groups such as amphipods, cumaceans and tanaids had no previously reported specimens (Appendix A).

Of the 668 total taxa found in the present study 462 (69%) were identified to species (Appendix B), and 421 of these identified species (91%) are known to occur in Hawaii (Table 2), indicating that the majority of the Johnston Atoll marine flora and fauna is composed of Hawaiian species.

This was particularly the case for macroalgae, Porifera, Polychaeta, Sipunculida Mollusca, Echinodermata and fish, where Hawaiian species made up 90% or more of the identified organisms (Table 2). Only bryozoans (Ectoprocta) and ascidian had a substantial proportion of their identified species which were not previous known from Hawaii, suggesting that these two groups, which are susceptible to transport as fouling organisms, include organisms that may have been humanly introduced from areas outside of Hawaii.

Introduced organisms made up only a small fraction of the total Johnston Atoll biota. Only ten of the 668 taxa (1.5%) were species designated as introduced or cryptogenic in Carlton and Eldredge (in prep.) or by taxonomists familiar with the respective groups (Table 3). Alternatively, nonindigenous or cryptogenic species composed only 2% of the 462 taxa identified to species. These introduced or potentially introduced species included the a nonindigenous hydrozoan and polychaete found in Hawaii, a one nonindigenous bryozoan found in Hawaii, three previously unreported cryptogenic bryozoans, and three nonindigenous ascidians. Only one of these, the bryozoan *Didymozoum triseriale* (Philipps, 1899), has a previously known distribution restricted to the Indo-Pacific, while the others were tropical worldwide or undetermined (Table 3).

Table 3. Cryptogenic and nonindigenous species found on June 2000 Johnston Atoll Surveys

Taxa	Species	Status	Origin or Range
Hydrozoa	<i>Pennaria disticha</i>	Nonindigenous	Tropical Worldwide
Polychaeta	<i>Branchiomma nigromaculata</i>	Cryptogenic	Tropical Worldwide
	<i>Armandia intermedia</i>	Cryptogenic	Undetermined
Bryozoa	<i>Bugula vectifera?</i>	Cryptogenic	Undetermined
	<i>Caulibugula dendrograpta</i>	Nonindigenous	Undetermined
	<i>Didymozoum triseriale</i>	Cryptogenic	Indo-Pacific
	<i>Halysis diaphana</i>	Cryptogenic	Tropical Worldwide
Ascidacea	<i>Ascidia sydneyensis</i>	Nonindigenous	Undetermined
	<i>Diplosoma listerianum</i>	Nonindigenous	Tropical Worldwide
	<i>Microcosmus exasperatus</i>	Nonindigenous	Tropical Worldwide

Comparison of numbers of taxa among stations for various taxonomic groups and for all taxa combined (Figure 2, Table 4) showed little variation among stations for any major group nor for total taxa. Maximum numbers of taxa occurred at Station 4 (Donovan's Reef) for combined taxa, crustaceans and fish, but the differences from other stations were not substantial. For example, the 215 total taxa occurring at this site were only 71 more than the 144 taxa occurring at Station 9 (Cucumber Flats), the site of fewest reports. Even lower variation among stations was indicated by Shannon H'_{10} diversity indices, which ranged only 2.16 at Stations 6 and 9 to 2.33 at Station 4. The most species rich taxonomic groups by stations were the macroalgae (9-28 taxa), polychaetes (14-27 taxa), crustaceans (26-49 taxa), mollusks (19-71 taxa) and fishes (20-52 taxa). The only spatial pattern suggested by these data are increased numbers of taxa and diversity at Station 2 on the or at Stations 4 and 8 on the reef slope, which ranked 1 to 3 in total taxa, respectively. However, Station 9 outer back reef flat ranked last of the 11 stations in total taxa, 10th in macroalgae and 9th in fishes.




Figure 2 Numbers of taxa in major taxonomic groups at Johnston Atoll stations

Table 4. Data for Figure 2 and Shannon's H'_{10} diversity indices.

Taxa	Station										
	1	2	3	4	5	6	7	8	9	10	11
Algae	9	28	24	25	19	16	11	17	10	21	13
Porifera	3	0	2	4	0	1	3	1	1	5	0
Cnidaria	7	9	7	12	11	11	13	13	11	7	7
Polychaeta	25	17	18	14	24	27	20	12	17	15	17
Sipunculida	3	2	1	2	1	1	0	0	0	0	1
Crustacea	32	37	40	49	26	37	39	45	43	42	40
Mollusca	37	71	36	37	53	29	19	27	32	19	40
Ectoprocta	5	1	1	3	1	0	1	0	0	1	0
Echinodermata	4	10	10	13	2	2	2	15	5	1	1
Ascidia	3	1	0	3	1	1	6	3	0	4	0
Fish	25	32	36	52	39	20	38	47	25	37	30
All Taxa	153	208	175	214	177	145	152	180	144	152	149
H'_{10} Diversity	2.18	2.32	2.24	2.33	2.25	2.16	2.17	2.26	2.16	2.17	2.18




Figure 3. Dendrogram of Sorensen similarities among station for all taxa using UPGMA method for calculating intercluster distances

Calculation of Sorensen similarity indices species presence-absence (Figure 3) for all taxa indicates that similarities within clusters were not high in any case, with the highest similarity about 55% occurring between stations 4 and 8. Nonetheless, the dendrogram suggests patterns among station clusters that correspond to reef location and environment, despite the relatively small differences among stations for species numbers and diversity. Stations 1 and 10 did not group with any other, and these were in unique locations, Station 1 being on and near the Johnston Island pier, and Station 10 along the channel in the south lagoon. The Station 4 and 8 cluster contains the two stations which were on the outer reef slope where overall taxa numbers and diversity were high (Table 3), especially for crustaceans and fishes.

The cluster containing Stations 2, 3 and 9 corresponds to locations in the north lagoon that were on the outer back reef flat. Running similar analysis for the individual taxonomic groups indicates that this cluster results primarily from taxa within the polychaetes, crustaceans, echinoderms and fishes. Stations 2 and 3 grouped at about 60% similarity for the crustaceans and echinoderms.

Stations 2 and 9 grouped at about 77% similarity for the fishes, and Stations 3 and 9 at about 70% similarity for the polychaetes.

The largest cluster of four stations for total taxa is formed by Stations 5, 6 and 7, all along dredged areas in the north lagoon, with Station 11, near the sewage outfall south of Johnston Island. This cluster appears to result from a number of associations within the individual taxonomic groups. Station 5 groups at about 75% similarity with Station 6 for the polychaetes, at about 65% similarity with Station 7 for the fishes and at about 55% similarity with Station 11 for the crustaceans. Station 6 clusters with Station 11 at about 72% similarity for the fishes, at 92% similarity with Station 7 for the cnidarians; and with Station 7 at about 40% similarity for mollusks. Station clusters at about 63% similarity with both Stations 5 and 6 for the polychaetes.

Station 11 was the most unusual of the reef sites surveyed, with the alga *Bryopsis hypnoides* growing over the reef in high densities, apparently dominating the benthos in competition with reef corals and other sessile forms. Despite this apparent dominance by a species indicating eutrophication by the sewage outfall and discharge, the similarity analysis based on species presence-absence did not separate this station from any other for total taxa or any major taxonomic group. Nor were species diversity or numbers of taxa particularly low at this station except for echinoderms, where it tied with Station 7 for least number of species. This indicates, that despite the apparent dominance of *Bryopsis hypnoides* in this area, the site still supports a large variety of biota typical for the Johnston Atoll lagoon.

Stations 1 and 10 did not group with any sites for total taxa (Figure 3) or in similarity dendrographs for all invertebrates excluding algae and fishes, and for mollusks alone. Station 1 was unique in its location on and near the concrete Johnston Island Pier where most benthic organisms had little three dimensional relief for settlement and growth, and many molluscs which are usually found on flat manmade structures occurred only at this site. Also the fewest numbers of algal found on the study were at this site, and fish species tied with second from last. Station 10, at the end of the south lagoon channel, had next to the lowest number of total taxa and the fewest taxa of molluscs and coral species found on the study.

DISCUSSION

Compared with harbors which have been surveyed on Oahu, Hawaii (Coles et al. 1997, 1999a, 1999b) and found to have a introduced component of around 100 species or about 20% of their total biota, cryptogenic or nonindigenous are only a minor portion of the Johnston Atoll marine community. With only ten potentially introduced species comprising only 1-2% of the total species or taxa, the nonindigenous component approximates that which has been found for previous surveys of non harbor areas in the Hawaiian Islands such as Midway Atoll and Kaho'olawe (Coles et al. 1998, DeFelice et al. 1998). Similarly, studies in Guam (Paulay et al. in prep.) and Australia (Hewitt et al, 1998; Hoedt et al. 2000) which have found a nonindigenous component of around 1-2% of the total biota identified (Table 5). Nonindigenous species conspicuously absent from Johnston Atoll are the Caribbean intertidal barnacle *Cthamalus*

proteus, which is extremely abundant in the intertidal zone in harbors and bays in Hawaii and has spread as far westward as Guam (Southward et al. 1998), and invasive nonindigenous red algae (Russell 1992) which cover reef flats in many areas of the main Hawaiian Islands. With monthly visits by a supply barge from Honolulu plus previous military vessel movement during the atoll's occupation in the last 50 years, there has been ample opportunity for species introductions. The relative lack of introduced species on the atoll is probably in large part due to the inability for nonindigenous forms adapted to harbor conditions to establish themselves in the oligotrophic reef environment where they must also compete with a diverse biota adapted to tropical conditions. Higher diversity of the Australian native fauna has been proposed to account for the relatively low nonindigenous component found in North Queensland ports and harbors (Hutchings et al. ms submitted) although such areas are still susceptible to invasion by an foreign organism that is adapted to tropical reef conditions (Pyne 1999; Willan et al. 2000).

Table 5. Numbers of marine nonindigenous, cryptogenic and total species determined on Hawaii, Guam and Australian surveys. (Modified from Coles and Eldredge, ms submitted).

<i>Location</i>	Nonindigenous (N)	Cryptogenic (C)	Total N + C	Total Species	% N + C
<u>Hawaii</u>					
Oahu, Pearl Harbor	69	26	95	419	23.0
Oahu, S. & W. Shores	73	27	100	585	17.0
Midway	4	0	4	444	1.5
Kaho'olawe	3	0	3	298	1.0
<u>Guam</u>					
Apra Harbor	27*	29*	46*	682	6.7
Island-wide			104	4635	2.2
<u>Australia</u>					
Hay Point Port	8	2	10	506	2.0
Mourilyan Harbour	2	2	4	401	1.0
Abbot Point Port	0	5	5	593	0.8

As previously reported for the algal fish and reef coral assemblages at Johnston Atoll, the invertebrates found in this study are highly dominated by organisms that occur in the Hawaiian Islands. This would be expected given Johnston Atoll's location in the North Pacific gyre where it receives water that has passed the Hawaiian Islands. However, under the influence of the North Equatorial Countercurrent, the Johnston Atoll biota also has a number of tropical species not found in the main Hawaiian Islands which probably originate from the central equatorial Pacific. Some taxonomic groups such as reef corals, are highly dominated by tropical species which are not found in most of the main Hawaiian Islands, and Johnston Atoll may be responsible for the reintroduction of these organisms into the Northwest Hawaiian Islands (Grigg 1981).

The data and similarity analyses indicate that there are no large differences among the sites in terms of their species presence-absence, although there is an overall pattern in the similarities that corresponds to location and environmental conditions. Small differences among the biota of

the eleven sites would be expected in a well-mixed atoll lagoon where no outstanding differences occur among sites in depth, water quality or sedimentation. The only area that was dramatically different from the other areas observed on the survey was at Station 11 in the vicinity of the Johnston Island sewer outfall. The bloom of *Byopsis hypnoides* and, to lesser extent, *Caulerpa racemosa* in this area was conspicuous and unique for the atoll, and clearly indicated eutrophication of the waters of the south lagoon by the sewage discharge. This is somewhat surprising, given the openness of the receiving water to unrestricted flow of open ocean water, depth of the discharge which should encourage mixing as effluent rises to the surface, and assumed low volume of discharge.

The occurrence of a colony of *Acropora cythera* with aberrant growth on its surface at Station 2 (Plate 1) is interesting and may warrant further surveys for corals with such growths in the Johnston Atoll lagoon, given the use of the atoll for aerial nuclear bomb testing 40-50 years ago. These growths are reminiscent, but not the same as spherical calciblastic tumors that have been observed on *Acropora* corals in Oman (Coles and Seapy 1998) and elsewhere (Peters et al. 1986). The factor or factors that causes such aberrant coral growth is unknown, and the growth form found on the Johnston Atoll coral has not previously been observed.

CONCLUSIONS

The Johnston Atoll marine community is a representative of a typical coral reef environment that is composed primarily of organisms found in the Hawaiian Islands. However, its megafauna is highly dominated by live reef corals of the genus *Acropora*, which is not found in any of the main Hawaiian Islands except Kauai, where it occurs only rarely. The present study has substantially increased the knowledge of taxonomic groups other than macroalgae, reef corals and fishes with 325 taxa newly reported, with about 90% of these composed of Hawaiian species or higher taxa.

A very small component of the Johnston Atoll marine community is composed of introduced species, with only ten cryptogenic or recognized nonindigenous species found among the 668 taxa or 452 identified species. None of these introduced organisms were abundant, and they usually occurred as single specimens at one or two stations. Johnston Atoll is therefore consistent with other coral reef areas in Hawaii, Guam and Australia that have been found to have few nonindigenous species occurring in low abundance.

ACKNOWLEDGMENTS

We are grateful to the U. S. Fish and Wildlife Service for funding and logistic support that made this study possible. We greatly appreciate the kind assistance of the following individuals that provided identification and verification of species identifications for the respective taxonomic groups:

Macroalgae: Mr. Jack Fisher, Bishop Museum

Hydroids: Dr Dale Calder, Royal Ontario Museum

Zoantharians: Dr. Daphne Fautin, University of Kansas

Isopods: Dr. Brian Kensley, U.S. National Museum of Natural History

Bryozoans: Ms. Chela Zabin, University of Hawaii

Ophiuroids: Dr. Gordon Hendler, Los Angeles County Museum

Ascidians: Dr. Gretchen Lambert, California State University at Fullerton

REFERENCES

- Ackman, R. G., T. Takeuchi and G. H. Balazs. 1992. Fatty acids in depot fats of green turtles *Chelonia mydas* from the Hawaiian Islands and Johnston Atoll. *Comp. Biochem. Physiol. B Comp. Biochem* 1992: 813-819.
- Agegian, C. R. 1985. Deep water macroalgal communities: a comparison between Penguin Bank, Hawaii and Johnstons Atoll. *Proc. 5th Int Coral Reef Congress, Tahiti* 5: 47-50.
- Amerson, A. B. 1973. Ecological baseline survey of Johnston Atoll, Central Pacific Ocean. Smithsonian Institution, Washington, D. C.
- Amerson, A. B. and P. C. Shelton. 1976. The natural history of Johnston Atoll, Central Pacific Ocean. *Atoll Res. Bull.* 192: 1-477.
- Anderson, J. D. 1986. Age, growth, reproduction and diet of *Myripristis aeneana* at Johnston Atoll. Masters thesis, Depart. of Zoology, University of Hawaii, Honolulu. 73 pp.
- Bailey-Brock, J. 1976. Habitats of tubicolous polychaetes from the Hawaiian Islands and Johnston Atoll. *Pac. Sci.* 30: 69-81.
- Baker, H. K., G. M. Patterson and M. Ikagawa. 1997. Preliminary description of *Borzia elongata* sp. nov., a representative of a valid genus in the Oscillatoriales. *Arch. Hydrobiologie Suppl.* 118: 1-12.
- Balazs, G. H. 1994. Howeward bound: satellite tracking of green turtles from nesting beaches to foraging pastures. NMFS, NOAA Tech. Mem. NMFS-SEFSC 341,
- Balazs, G. W. 1986. Status and ecology of marine turtles at Johnston Atoll. *Atoll Res. Bull.* 285: 1-46.
- Balazs, G. W. 1986. Status and ecology of marine turtles at Johnston Atoll: 1985 assessment. Southwest Fisheries Center, Honolulu Laboratory, Admin. Rep. H-86-9, Honolulu.
- Brock, R. E. 1973. A new distributional record for *Panulirus marginatus* (Quoy and Gaimard 1925). *Crustaceana* 25: 111-112.
- Brock, R. E. 1979. A statistical study of *Cypraea tigris* in the central Pacific. *Veliger* 22: 166-170.

- Brock, V. E., R. S. Jones and P. Helfrich. 1965. An ecological reconnaissance of Johnston Island and the effects of dredging. Annual report. Univ. of Hawaii, Haw. Inst. Mar. Biol., HIMB Tech. Rep. No. 5, Honolulu.
- Brock, V. E., W. van Heukelem and P. Helfrich. 1966. An ecological reconnaissance of Johnston Island and the effects of dredging. Second annual report. Univ. of Hawaii, Hawaii Inst. Mar. Biol., HIMB Tech. Rep. No. 11, Honolulu.
- Buggeln, R. G. and R. Tsuda. 1966. A preliminary marine algal flora from selected habitats on Johnston Atoll. Univ. of Hawaii, Hawaii Inst. Mar. Biol., HIMB Tech. Rep. No. 9, Honolulu.
- Cairns, S. D. 1984. New records of ahermatypic corals (Scleractinia) from the Hawaiian and Line Islands. Occas. Pap. Bishop Mus. 25: 1-30.
- Chave, E. H. and A. Malahoff 1998. In Deeper Waters. Photographic studies of Hawaiian Deep-sea Habitats and Life-forms. Univ. of Hawaii Press, Honolulu. 125 pp.
- Chave, E. H. and B. C. Mundy. 1994. Deep-sea benthic fish of the Hawaiian Archipelago, Cross Seamount and Johnston Atoll. Pac. Sci. 48: 36-404.
- Clark, A. H. 1949. Ophiuroidea of the Hawaiian Islands. Bishop Museum Bull. 195: 1-133.
- Cohen, A. L., P. S. Lobel and G. L. Tomasky. 1997. Coral bleaching on Johnston Atoll, Central Pacific Ocean. Biol. Bull. 193: 276-279.
- Coles, S. L. In press. First record of the reef coral; *Montipora hoffmeisteri* Wells, 1954 at Johnston Atoll (Cnidaria: Anthozoa: Scleractinia). Occ Pap. Bernice Bishop Mus.
- Coles, S. L., R. C. DeFelice and L. G. Eldredge. 1999a. Historical and recent introductions to non-indigenous marine species into Pearl Harbor, Oahu, Hawaiian Islands. Mar. Biol. 135: 1247-158.
- Coles, S. L., R. C. DeFelice and L. G. Eldredge. 1999b. Nonindigenous marine species introductions in the harbors of the south and west shores of Oahu, Hawaii. Bishop Museum, Tech. Rep. No. 15, Honolulu.
- Coles, S. L., R. C. DeFelice, L. G. Eldredge and J. T. Carlton. 1997. Biodiversity of marine communities in Pearl Harbor, Oahu, Hawaii with observations on introduced species. Bishop Museum, Tech. Rep. No. 10, Honolulu.
- Coles, S. L., R. C. DeFelice, J. E. Smith, D. Muir and L. G. Eldredge. 1998. Determination of baseline conditions for introduced marine species in nearshore waters of the island of Kaho'olawe, Hawaii. Bishop Museum, Tech. Rep. No. 14, Honolulu.
- Coles, S. L. and L. G. Eldredge. Ms. submitted. Nonindigenous species introductions on coral reefs: a need for information. Pac. Sci.
- Coles, S. L. and D. G. Seapy. 1998. Ultra-violet absorbing compounds and tumorous growths on acroporid corals from Bandar Khayran, Gulf of Oman, Indian Ocean. Coral Reefs 17: 195-198.
- Dee, A. J. and J. D. Parrish. 1994. Reproductive and trophic ecology of the soldierfish *Myripristis amaena* in tropical fisheries. USNMFS Fish. Bull. 92: 516-530.
- DeFelice, R. C., S. L. Coles, D. Muir and L. G. Eldredge. 1998. Investigation of the marine communities of Midway Harbor and adjacent lagoon, Midway Atoll, Northwestern Hawaiian Islands. Bishop Museum, Hawaiian Biological Survey Contr. No. 1998-014, Honolulu.

- Economakis, A. E. and P. S. Lobel. 1998. Aggregation behavior of the grey reef shark, *Carcharinus amblyrhinchos* at Johnston Atoll, central Pacific Ocean. *Environ. Biol. Fish.* 51: 129-139.
- Edmondson, C. H., W. K. Fisher, H. L. Clark, A. L. Treadwell and J. A. Cushman. 1925. Marine zoology of the tropical central Pacific. Tanager Expedition. Bishop Mus. Bull. 27: 1-148.
- Emery, K. O. 1956. Marine geology of Johnston Island and its surrounding shallows, central Pacific Ocean. *Geol. Soc. Amer. Bull.* 67: 105-1520.
- Fowler, H. W. and S. C. Ball. 1925. Fishes of Hawaii, Johnston Island and Wake Island. Bishop Mus. Bull. 26: 1-31.
- Gosline, W. A. 1955. The inshore fishes of Johnston Island, a central Pacific atoll. *Pac. Sci.* 9: 442-480.
- Grigg, R. W. 1981. *Acropora* in Hawaii. Part 2. Zoogeography. *Pac. Sci.* 35: 14-24.
- Halstead, B. W. and N. C. Bunker. 1954. A survey of the poisonous fishes of Johnston Island. *Zoologica* 39: 67-81.
- Hewitt, C. L., M. L. Campbell, K. M. Moore, N. B. Murfet and B. Robertson. 1998. Introduced species survey. Port of Hay Point, Queensland. CSIRO Centre for Research on Introduced Marine Pests to Ports Corporation of Queensland, Unpublished Report, Brisbane.
- Hoedt, F. E., J. H. Choat, J. Collins and J. J. Cruz. 2000. Mourilyan Harbour and Abbot Point surveys: port marine baseline surveys and surveys for introduced marine pests. School of Marine Sciences and Aquaculture, James Cook University to Ports Corporation of Queensland, Brisbane.
- Hutchings, P., R. Hilliard and S. L. Coles. Ms submitted. Assessment of non-indigenous species introductions and potential for marine pest invasions into tropical marine communities, with special references to the Indo-Pacific. *Pac. Sci.*
- Irons, D. K. 1989. Temporal and feeding behavior of the butterflyfish, *Chaetodon trifascialis*, at Johnston Atoll. *Environ. Biol. Fish.* 25: 187-193.
- Irons, D. K. 1990. Natural history and behavioral ecology of the butterflyfish, *Chaetodon trifascialis*, at Johnston Atoll. Masters thesis, Depart. of Zoology, University of Hawaii, Honolulu. 89 pp.
- Jokiel, P. L. and W. A. Tyler, III. 1992. Distribution of stony corals in Johnston Atoll lagoon. *Proc. 7th Intern. Coral Reef Symp.* 2: 683-692.
- Jones, R. S. 1967. Ecological relationships in Hawaiian and Johnston Island Acanthuridae (surgeonfishes) with emphasis on food and feeding habits. Ph. D. thesis, Dept. of Zoology, University of Hawaii, Honolulu. 245 pp.
- Kay, E. A. 1961. On *Cypraea tigris schilderiana* Cate. *Veliger* 4: 36-40.
- Kay, E. A. Unpublished. List of marine molluscs reported from Johnston Atoll.
- Kerr, L. M. and P. S. Lobel. 1997. PCB contamination relative to age for a Pacific damselfish *Abudefduf sordidus* (Pomacentridae). *Biol. Bull.* 193: 279-281.
- Kosaki, R. K. 1989. *Centropyge nahackyi*, a new species of angelfish from Johnston Atoll (Teleostei: Pomacanthidae). *Copeia* 1989: 880-886.
- Kosaki, R. K., R. L. Pyle, J. R. Randall and D. K. Irons. 1991. New records of fishes from Johnston Atoll, with notes on biogeography. *Pac. Sci.* 45: 186-203.

- Lobel, P. S. 1984. Ecological investigations to assess the impact of proposed deep ocean disposal of brine waste off Johnston Atoll. Unpublished report,
- Lobel, P. S. 1997. Comparative settlement age of damselfish larvae (*Plectroglydodon imparipennis*, Pomacentridae) from Hawaii and Johnston Atoll. Biol. Bull. 193: 281-283.
- Lobel, P. S. and D. A. Mann. 1995. Spawning sounds of the damselfish *Dascyllus albisella* (Pomacentridae), and relationship to male size. Bioacoustics 6: 187-198.
- Mann, D. A. and P. S. Lobel. 1995. Passive acoustic detection of sounds produced by the damselfish *Dascyllus albisella* (Pomacentridae). Bioacoustics 6: 199-213.
- Mann, D. A. and P. S. Lobel. 1998. Acoustic behavior of the damsel fish *Dascyllus albisella*: behavior and geographic variation. Environ. Biol. Fish. 51: 421-428.
- Maragos, J. E. and P. L. Jokiel. 1986. Reef corals of Johnston Atoll: one of the world's most isolated reefs. Coral Reefs 4: 141-150.
- McCosker, J. E. and D. G. Smith. 1997. Two new Indo-Pacific morays of the genus *Uropterygius* (Anguilliformes:Muraenidae). Bull. Mar. Sci. 60: 1005-1014.
- Moul, E. T. 1964. New records of *Halimeda* and *Udotea* for the Pacific area. Atoll Res. Bull. 106: 1-10.
- Paulay, G., L. Kirkendale, G. Lambert and C. Meyer. In prep. Anthropogenic biotic interchange in a coral reef ecosystem: a case study from Guam. Pac. Sci.
- Paulay, G., L. Kirkendale, G. Lambert and J. Starmer. In prep. The marine invertebrate biodiversity of Apra Harbor: significant areas and introduced species, with focus on sponges echinoderms and ascidians. Naval Activities Guam, Cooperative agreement N68711-97-LT-70001, Agana, Guam.
- Peters, E. C., J. C. Halas and H. B. McCarty. 1986. Calicoblastic neoplasms in *Acropora palmata*, with a review of reports on anomalies of growth and form in corals. Journal of the National Cancer Institute 76: 895-911.
- Pyne, R. 1999. The black striped mussel (*Mytilopsis sallei*) infestation in Darwin: a clean-up strategy. The Ballast Water Problem: Where to from here? Ecoports Monograph Series, Brisbane. 77-83.
- Ralston, S., R. M. Gooding and G. M. Ludwig. 1985. An ecological survey and comparison of bottom fish resource assessment (submersible versus handline fishing) at Johnston Atoll. Fish. Bull.
- Randall, J. E. 1972. A revision of the labrid fish genus *Anampses*. Micronesica 8: 151-195.
- Randall, J. E. 1977. Contribution to the biology of the whitetip reef shark (*Triaenodon obesus*). Pac. Sci. 31: 143-164.
- Randall, J. E. 1999. Review of the dragonets (Pisces: Callionymidae) of the Hawaiian Islands, with descriptions of two new species. Pac. Sci. 53: 185-207.
- Randall, J. E., P. S. Lobel and E. H. Chave. 1985. Annotated checklist of the fishes of Johnston Island. Pac. Sci. 39: 24-80.
- Randall, J. E. and S. Ralston. 1984. A new species of serranid fish of the genus *Anthias* from the Hawaiian Islands and Johnston Island. Pac. Sci. 38: 220-227.
- Russell, D. J. 1992. The ecological invasion of Hawaiian reefs by two marine red algae, *Acanthophora spicifera* (Vahl) Boerg. and *Hypnea musciformis* (Wulfen) J. Ag., and their

- association with two native species, *Laurencia nidifica* J. Ag. and *Hypnea cervicornis* J. Ag. ICES Mar. Sci. Symp. 194: 110-125.
- Sancho, G., D. Ma and P. S. Lobel. 1997. Behavioral observations of an upcurrent reef colonization event by larval surgeonfish *Ctenochaetus strigosus* (Acanthuridae). 311-315
- Schultz, L. P. and Collaborators. 1953. Fishes of the Marshall and Marianas Islands. Bull. U. S. National Mus. 202:
- Smith, R. M. and J. Swain. 1882. Notes on a collection of fishes from Johnston Island, including descriptions of five new species. Proc. U. S. Nat. Mus. 5: 119-143.
- Ward, L. A. 1981. A taxonomic study of the Spionidae (Annelida: Polychaeta) from the Hawaiian Islands and Johnston Atoll with notes on their ecology and biogeographical distribution. Ph. D. thesis, Dept. of Zoology, University of Hawaii, Honolulu. 224 pp.
- Wells, J. W. 1954. Recent corals of the Marshall Islands. Prof. Pap. U.S. Geolog. Surv. 260-I: 385-486.
- Wennekens, M. P. 1969. Johnston Island regional oceanography. Forecasting currents, eddies, island wake. Office of Naval Research, San Francisco.
- Willan, R. C., B. C. Russell, N. B. Muref, K. L. Moore, F. R. Mcennulty, S. K. Horner, C. L. Hewitt, G. M. Dally, M. L. Campbell and S. T. Bourke. 2000. Outbreak of *Mytilopsis sallei* (Recluz, 1849) (Bivalvia: Dreissenidae) in Australia. Molluscan Res. 20: 25-30.
- Winterbottom, R. and M. Burrige. 1993. Revision of Indo-Pacific *Priolepsis* species possessing a reduced transverse pattern of cheek papillae, and predorsal scales (Teleostei: Gobiidae). Can. Rev. Zool. 71: 2056-2076.

PLATES

Plate 1. Colony of the table coral *Acropora cythera* at Station 2 showing aberrant growths on upper surface.

Plate 2. *Acropora cythera* at Station 3 showing annular growth bands.

Plate 3. Mixed growth of *Acropora* and *Montipora* coral species at Station 3.


Plate 4. The damselfish *Centropyge nahackyi* Kosaki, 1989 at the base of the reef slope in 20 depth at Station 4.

Plate 5. School of surgeonfish *Zebrasoma flavescens* (Bennett, 1828) near top of reef slope in about 5 m depth at Station 8.

Plate 6. Mixed growth of *Acropora* and *Montipora* coral species at Station 9.

Plate 7. Close-up of *Acropora cythera* at Station 10 showing annular growth bands.

Plate 8. Medium distance view of *Acropora cythera* at Station 10 showing annular growth bands.



APPENDIX A

Previous Reports of Marine Organisms Observed or Collected at Johnston Atoll

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Goniotrichum alsidii</i>	(Zanardini) M. Howe																						x
		<i>Griffithsia metcalfi</i>	Tseng																						x
		<i>Griffithsia ovalis</i>	Harv.																						x
		<i>Griffithsia sp.</i>																							x
		<i>Griffithsia tenuis</i>	C. Agardh																						x
		<i>Hypnea esperi</i>	Bory																						x
		<i>Jania capillacea</i>	Harv.																						x
		<i>Jania decussato-dichotoma</i>	(Yendo) Yendo																						x
		<i>Lomentaria hakodatensis</i>	Yendo																						x
		<i>Taenioma macrourum</i>																							x
		<i>Wurdemanina sp.</i>																							x
Hydrozoa	Milleporidae	<i>Millepora tenera</i>	Boschma, 1949						x																x
		<i>Millepora tenera?</i>	Boschma, 1949																						x
	Stylasterinidae	<i>Distichopora violacea</i>	(Pallas, 1766)																						x
		<i>Stylaster sp.</i>																							x
														x		x									x
Anthozoa	Acroporidae	<i>Acropora cerialis</i>	(Dana, 1846)																						x
		<i>Acropora cythera</i>	(Dana, 1846)						x																x
		<i>Acropora elseyi</i>	(Brook, 1892)																						x
		<i>Acropora humilis</i>	(Dana, 1846)						x																x
		<i>Acropora nasuta</i>	(Dana, 1846)																						x
		<i>Acropora paniculata</i>	(Verrill, 1902)																						x
		<i>Acropora selago</i>	(Studer, 1878)																						x
		<i>Acropora valida</i>	(Dana, 1846)																						x
		<i>Acropora yongei</i>	Veron & Wallace 1984																						x
		<i>Montipora capitata?</i>	(Lamarck, 1816)						x																x
		<i>Montipora studeri</i>	(Vaughan, 1907)																						x
		<i>Montipora tuberculosa</i>	(Lamarck, 1816)																						x
		<i>Montipora verrilli</i>	Vaughan, 1907																						x
		<i>Montipora verrucosa</i>	(Lamarck, 1816)																						x
	Agariciidae	<i>Leptastrea purpurea</i>	(Dana, 1846)																						x
		<i>Leptastrea sp.</i>																							x
		<i>Leptoseris hawiiensis</i>	Vaughan, 1907																						x
		<i>Leptoseris incrustans</i>	(Quelch 1886)																						x
		<i>Leptoseris papyracea</i>	(Dana, 1846)																						x
		<i>Leptoseris scabra</i>	(Vaughan, 1907)																						x
		<i>Pavona duerdeni</i>	(Vaughan, 1907)																						2
		<i>Pavona maldivensis</i>	(Gardiner, 1905)																						2
		<i>Pavona varians</i>	Verrill, 1864																						x
	Alcyonacea	<i>Bebryce brunnea</i>	(Nutting)																						x
		<i>Corallium tortuosum</i>	Bayer																						2
		<i>Sinularia abrupta</i>	Tixier-Durivault, 1970																						x
	Antipatharia	<i>Antipathes dichotoma</i>	Pallas																						x
		<i>Antipathes intermedia</i>	(Brook)																						x
		<i>Antipathes punctata</i>	(Roule)																						x

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Antipathes subpinnata</i>	Ellis & Solander																					x	
		<i>Antipathes ulex</i>	Ellis & Solander																					x	
		<i>Bathypathes conferta</i>	(Brook)																				x		
		<i>Cirripathes spiralis</i>	Linn., 1758																				x		
		<i>Leopathes glaberrima</i>	(Esper)																				x		
	Balanophyllidae	<i>Balanophyllia hawaiiensis</i>	Vaughan, 1907																				x		
	Dendrophyllidae	<i>Cladopsammia echinata</i>	Cairns, 1984																				2		
		<i>Dendrophyllia oahuensis</i>	Vaughan, 1907																			x	x		
		<i>Enallopsammia rostrata</i>	Pourtales, 1878																				2		
	Faviidae	<i>Cyphastrea ocellina</i>	(Dana, 1846)																				x		
		<i>Plesiastrea versipora</i>	(Lamarck, 1816)																				x		
	Flabellidae	<i>Javania lamprotichum</i>	Mosley, 1880																				x		
	Fungiidae	<i>Cycloseris vaughani</i>	Lamarck, 1801																				x		
		<i>Fungia scutaria</i>	Lamarck, 1801																				x		
	Isopheliidae	<i>Telmatactis decora</i>	(Hemprich and Ehrenberg, 1834)																				x		
	Pocilloporidae	<i>Madracis kauaiensis</i>	Vaughan, 1907																				x		
		<i>Pocillopora damicornis</i>	(Linnaeus, 1758)																				x		
		<i>Pocillopora eydouxi</i>	(Milne Edwards and Haime 1860)																				x		
		<i>Pocillopora meandrina</i>	(Dana, 1846)																				x		
	Poritidae	<i>Portites lobata</i>	(Dana, 1846)																				x		
		<i>Portites lutea</i>	(Milne Edwards and Haime 1860)																				x		
	Siderastreidae	<i>Psammocora nierstrazi</i>	Van der Horst, 1922																				x		
		<i>Psammocora stellata</i>	Verrill, 1864																				x		
		<i>Oulangia bradleyi</i>	(Boschma, 1923)																				x		
Aschelminthes	Nemertea	<i>Baseodiscus cingulatum</i>	(Coe, 1906)																				x		
Polychaeta	Amphinomidae	<i>Eurythoe complanata</i>	(Pallas, 1766)																				x		
		<i>Eurythoe pacifica</i>	Kinberg																				x		
		<i>Hermodice pinnata</i>	Treadwell																				x		
		<i>Notopygos albiseta</i>	Holly, 1939																				x		
		<i>Pherecardia striata</i>	(Kinberg, 1857)																				x		
	Cirratulidae	<i>Cirratulus sp.</i>																					x		
	Eunicidae	<i>Eunice sp.</i>																					x		
	Leodiciidae	<i>Lysidice fusca</i>	Treadwell																				x		
		<i>Lysidice sp.</i>																					x		
	Leodocidae	<i>Leodice sp.</i>																					x		
	Nereidae	<i>Perinereis helleri</i>	(Grube, 1878)																				x		
		<i>Perinereis sp.</i>																					x		
		<i>Platynereis pulchella</i>	Gravier, 1901																				x		
	Phyllodocidae	<i>Phyllodoce stigmata</i>	Treadwell																				2		
	Polynoidae	<i>Hololepidella nigropunctata</i>	(Horst, 1915)																				x		
		<i>Neodexiospira foraminosa</i>	(Morre & Bush, 1904)																				x		
		<i>Neodexiospira pseudocorrugata</i>	(Bush, 1904)																				x		
		<i>Pileolaria pseudomilitaris</i>	(Thirot-Quievreux, 1965)																				x		
		<i>Vinaria koehlerii</i>	(Caullery & Mesnil, 1897)																				x		

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Oenone fulgida</i>																						x	
Crustacea	Alpheidae	<i>Alpheus brevipes</i>	Stimpson, 1860																						
		<i>Alpheus bucephalus</i>	Coutiere, 1905		x																				x
		<i>Alpheus clypeatus</i>	Coutiere, 1905		x																				x
		<i>Alpheus collumianus</i>	Stimpson, 1860		x																				
		<i>Alpheus crassimanus</i>	Heller		x																				
		<i>Alpheus diadema</i>	Dana, 1852		x																				
		<i>Alpheus gracilis simplex</i>	(Banner, 1953)		x																				
		<i>Alpheus paracrinitus</i>	Miers, 1881		x																				
		<i>Alpheus paragracilis</i>	Coutiere, 1905		x																				
		<i>Alpheus edmondsoni</i>	(Banner, 1953)																						x
		<i>Alpheus leviusculus</i>	Dana, 1852		x																				
		<i>Alpheus lottini</i>	Guerin, 1830																						x
		<i>Synalpheus paraneomeris</i>	Coutiere, 1905		x																				x
	Axiidae	<i>Axiopsis johnstoni</i>	Edmondson		x																				x
	Calappidae	<i>Calappa hepatica</i>	(Linnaeus, 1758)		x																				x
	Chirostylidae	<i>Eumunida smithii</i>	Henderson, 1885																						x
	Cirripedia	<i>Lepas anatifera</i>	L., 1758																						x
	Cryptochiridae	<i>Hapalocarcinus marsupialis</i>	Stimpson, 1859		x																				x
		<i>Pseudocryptochirus crescentus</i>	(Edmondson, 1925)		x																				2
	Diogenidae	<i>Aniculus aniculus</i>	(Fabricius, 1787)		x																				x
		<i>Calcinus elegans</i>	(N. Milne-Edwards, 1836)		x																				x
		<i>Calcinus herbstii</i>	deMan		x																				x
		<i>Calcinus latens</i>	(Randall, 1840))		x																				x
		<i>Dardanus haanii</i>	Rathbun																						
		<i>Dardanus megistos</i>	(Herbst)																						
		<i>Dardanus punctulatus</i>			x																				
		<i>Trizopagurus tenebrarum</i>	(Alcock, 1905)																						x
	Dynomenidae	<i>Dynomene devaneyi</i>	Takeda																						2
		<i>Dynomene hispida</i>	Desmarest, 1825																						x
		<i>Dynomene hispida</i>	Desmarest, 1825		x																				
	Galatheidae	<i>Galathea spinosorostris</i>	Dana																						
		<i>Munida brucei</i>	Baba																						x
		<i>Munida heteracantha</i>	Ortmann, 1892																						
	Gnathophyllidae	<i>Gnathophyllum americanum</i>	Guerin, 1856		x																				
		<i>Gnathophyllum fasciolatum</i>																							x
	Grapsidae	<i>Grapsus strigosus</i>	(Herbst)		x																				x
		<i>Grapsus tenuicrustatus</i>	(Herbst, 1783)		x																				2
		<i>Pachygrapsus minutus</i>	A. Milne-Edwards		x																				x
		<i>Pachygrapsus plicatus</i>	A. Milne-Edwards																						
		<i>Pachygrapsus plicatus</i>	A. Milne-Edwards																						x
	Hippolytidae	<i>Lysmata paucidens</i>	(Rathbun)																						x
		<i>Saron marmoratus</i>	(Olivier, 1811)		x																				x
	Isopoda	<i>Limnoria tripunctata</i>	Menzies, 1951																						x

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
	Latrellidae	<i>Latrellia metanese</i>	(Williams)																				x		
	Leucosidae	<i>Nucia speciosa</i>	Dana										x		x										
		<i>Randallia distincta</i>	Rathbun, 1906																						
	Leucosiidae	<i>Nucia speciosa</i>	Dana		x																				
		<i>Randallia distincta</i>	Rathbun																			x			
	Majidae	<i>Perinea tumida</i>	Dana, 1852																						
		<i>Schizophrys hilensis</i>	Rathbun, 1906																						
	Ocypodidae	<i>Ocypode pallidula</i>	Jacquinet																						
			Jaguinot, 1852																						
	Paguridae	<i>Pagurus haani</i>																							
		<i>Pagurus megistos</i>																							
	Palaemonidae	<i>Coralliocaris graminea</i>	(Dana, 1852)		x									x		x									
		<i>Harpiliopsis depressus</i>	(Stimpson, 1860)		x								x		x										
		<i>Jocaste lucina</i>	(Nobili, 1901)		x									x		x									
		<i>Palaemonella tenuipes</i>	Dana, 1852		x																				
		<i>Periclimenaeus tridentatus</i>	(Miers, 1884)		x									x		x									
	Panuliridae	<i>Panulirus marginatus</i>	(Quoy & Gaimard)											x		x						x			
		<i>Panulirus pencillatus</i>	(Olivier, 1888)		x									x		x									
	Parapaguridae	<i>Parapagurus dofleini</i>	Balss																			x			
	Parthenopidae	<i>Parthenope stellata</i>	Rathbun, 1906																			x			
	Plandonidae	<i>Plesionika alcocki</i>	(Anderson)																						
		<i>Plesionika pacificus</i>	Edmondson, 1952																						
	Portunidae	<i>Catoptrus inaequalis</i>	(Rathbun, 1906)		x										x		x								
		<i>Libystes edwardsi</i>	Alcock, 1900																						
		<i>Portunus longispinosus</i>	(Dana)		x										x		x								
		<i>Thalamita admete</i>	(Herbst)		x										x		x								
		<i>Thalamitoides quadridens</i>	(A. Milne-Edwards, 1869)																						
		<i>Thalamitoides quidridens</i>	(A. Milne-Edwards, 1869)		x												x		x						
	Processidae	<i>Heterocarpus ensifer</i>	A. Milne Edwards, 1881																						
	Scyllaridae	<i>Parribacus antarcticus</i>	(Lund)			x									x		x								
	Squillidae	<i>Pseudosquilla oculata</i>	(Brulle)			x									x		x								
	Stenopididae	<i>Stenopus hispidus</i>	Rathbun, 1907																						
		<i>Stenopus pyrrsonotus</i>	Goy & Devaney																				x		
	Xanthidae	<i>Actaea speciosa</i>	(Dana, 1852)		x										x		x								
		<i>Carpilius convexus</i>	(Forsskal, 1775)		x																				
		<i>Carpilodes bellus</i>	(Dana, 1852)																						
		<i>Chlorodiella asper</i>	Edmondson																						
		<i>Chlorodiella laevisima</i>	A. Milne-Edwards, 1873																						
		<i>Chlorodiella nigra</i>	(Forskal, 1775)																						
		<i>Chlorodopsis aberrans</i>	Rathbun, 1906																						
		<i>Chlorodopsis areolata</i>	A. Milne-Edwards, 1873																						
		<i>Domecia hispida</i>	Eydoux & Souleyet, 1842																						
		<i>Etisus electra</i>	(Herbst, 1801)		x										x		x								
		<i>Leptodius sanguineus</i>	(H. Milne-Edwards, 1834)		x												x		x						

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Leptodius waialuanus</i>	(Rathbun, 1906)		x																				
		<i>Liocarpilodes biunguis</i>	(Rathbun, 1906)																					x	
		<i>Liocarpilodes integerrimus</i>	Dana, 1852																					x	
		<i>Liomera belia</i>	(Dana, 1852)		x								x												
		<i>Lophozozymus dodone</i>	(Herbst, 1801)										x		x									x	
		<i>Phymodius laysani</i>	(Herbst, 1801)		x								x		x									x	
		<i>Phymodius nitidus</i>	(Dana, 1852)		x								x		x									x	
		<i>Pilodius aberrans</i>	(Rathbun, 1906)		x								x		x										
		<i>Pilodius areolata</i>	(H. Milne-Edwards)										x		x										
		<i>Platypodia eydouxi</i>	(A. Milne-Edwards, 1865)		x									x	x									x	
		<i>Portunus longispinosus</i>	(Dana, 1852)										x											x	
		<i>Pseudoliomera speciosa</i>	(Dana, 1852)		x									x		x									
		<i>Tetralia glaberrima</i>	(Herbst)		x																			x	
		<i>Tetralia spp.</i>	spp.										x												
		<i>Trapezia cymodoce</i>	(Herbst, 1801)										x												
		<i>Trapezia digitalis</i>	(Dana, 1852)		x								x											x	
		<i>Trapezia ferruginea</i>	Latreille, 1823		x									x		x								x	
		<i>Trapezia intermedia</i>	Miers, 1886		x									x		x								x	
		<i>Trapezia rufopunctata</i>	(Herbst)		x																				
		<i>Trapezia tigrina</i>	Eydoux & Souleyet, 1842										x		x		x							x	
Gastropoda	Actaeonidae	<i>Pupa tessellata</i>	Reeve, 1842																					x	
	Aplustridae	<i>Hydratina amplustre</i>	Linn., 1758																					x	
	Architectonidae	<i>Heliacus implexus</i>	Mighels, 1845																					x	
	Assumineidae	<i>Assumiea nitida</i>	Pease, 1865																					x	
	Buccinidae	<i>Cantharis farinosus</i>	Gould, 1850																					x	
		<i>Pisania ignea</i>	Gmelin, 1791											x		x									
		<i>Prodotia iostomus</i>	Gray, 1834																					x	
	Bullidae	<i>Bulla vernicosa</i>	Gould, 1859																					x	
	Bursidae	<i>Bursa cruentata</i>	Sowerby, 1841																					x	
		<i>Bursa rosa</i>	Perry, 1811																					x	
	Caecidae	<i>Caecum arcuatum</i>	de Folin, 1867																					x	
		<i>Caecum sepimentum</i>	de Folin, 1867																					x	
	Cassidae	<i>Casmaria erinaceus kalosmodix</i>	Melville, 1883																					x	
		<i>Cassus cornuta</i>	Linnaeus, 1758																					x	
	Cerithiidae	<i>Bittium impendens</i>	Hedley, 1899																					x	
		<i>Bittium zebrum</i>	(Kiener, 1841)																					x	
		<i>Cerithium atomarginatum</i>	Dautzenberg and Bouge, 1933																					x	
		<i>Cerithium columna</i>	Sowerby, 1834																					x	
		<i>Cerithium interstatum</i>	Sowerby, 1841																					x	
		<i>Cerithium mutatum</i>	Sowerby, 1834											x		x								x	

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																				
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
		<i>Cerithium nesioticum</i>	Pilsbry & Vanatta, 1905										x		x									x
		<i>Cerithium placidum</i>	Gould, 1861																					x
		<i>Plesiotrochus luteus</i>	Gould, 1861																					x
		<i>Rhinoclavis articulatus</i>	Adams & Reeve										x		x									
		<i>Rhinoclavis sinensis</i>	(Gmelin)										x		x									
		<i>Rhinoclavus articulata</i>	Adams and Reeve, 1850																					x
		<i>Rhinoclavus fasciata</i>	Bruguière, 1792																					x
		<i>Rhinoclavus sinensis</i>	Gmelin, 1791																					x
	Columbellidae	<i>Mitolumna metula</i>	Hinds, 1843																					x
		<i>Seminella virginea</i>	Gould, 1860																					x
	Conidae	<i>Conus abbreviatus</i>	Reeve, 1843																					x
		<i>Conus bandanus</i>																						x
		<i>Conus ebraeus</i>	Linn., 1758																					x
		<i>Conus flavidus</i>	Lamarck											x		x								
		<i>Conus imperialis</i>	Linn., 1758																					x
		<i>Conus lividus</i>	Hwass, 1792																					x
		<i>Conus miles</i>	Linn., 1758											x		x								x
		<i>Conus millepunctatus</i>																						x
		<i>Conus nanus</i>	Sowerby											x		x								
		<i>Conus obscurus</i>	Sowerby, 1833																					x
		<i>Conus pulicarius</i>	Hwass, 1792											x		x								x
		<i>Conus rattus</i>	Hwass, 1792											x		x								x
		<i>Conus retifer</i>	Menke, 1829																					x
		<i>Conus sponsalis</i>	Hwass, 1792																					x
		<i>Conus textile</i>	Linn., 1758																					x
		<i>Conus vitulinus</i>	Hwass, 1792											x		x								x
	Coralliophilidae	<i>Coralliophila erosa</i>	(Roding, 1798)																					x
		<i>Coralliophila violacea</i>	(Kiener, 1836)																					x
		<i>Coralliphilia erosa</i>	Röding, 1798																					x
		<i>Coralliphilia vilacea</i>	(Kiener, 1836)											x		x								
		<i>Quoyula madroporarum</i>	Sowerby, 1834											x		x								
	Cyclostrematidae	<i>Cyclostremicus emeryi</i>	(Ladd, 1966)																					x
	Cymatidae	<i>Cymatium aquatile</i>	Reeve, 1844											x		x								
		<i>Cymatium gemmatum</i>	Reeve, 1844											x		x								
		<i>Cymatium muricinum</i>	Röding, 1798											x		x								
		<i>Cymatium nicobaricum</i>	Röding, 1798											x		x								
		<i>Distorsio anus</i>	Linn., 1758											x		x								
		<i>Charonia tritonis</i>	Linnaeus, 1767																					x
		<i>Cymatium aquatile</i>	Reeve, 1844																					x
		<i>Cymatium gemmatum</i>	Reeve, 1844																					x
		<i>Cymatium intermedium</i>	Pease, 1869																					x
		<i>Cymatium muricinum</i>	(Roding, 1798)																					x
		<i>Cymatium nicobaricum</i>	(Roding, 1798)																					x
		<i>Cymatium rubeculum</i>	Linnaeus, 1758																					x

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																				
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
		<i>Distorsio anus</i>	Linn., 1758																					X
	Cypraeidae	<i>Cymatium aquatile</i>	Reeve, 1844																					X
		<i>Cypraea caputserpentis</i>	Linn., 1758										X		X									X
		<i>Cypraea carneola</i>	Linn., 1758										X		X									X
		<i>Cypraea chinensis</i>	Gmelin, 1791																					X
		<i>Cypraea granulata</i>	Pease, 1863											X		X								X
		<i>Cypraea helvola</i>	Linn., 1758										X		X									X
		<i>Cypraea isabella</i>	Linn., 1758										X		X									X
		<i>Cypraea lynx</i>	Linn., 1758																					X
		<i>Cypraea maculifera</i>	Schilder, 1932											X		X								X
		<i>Cypraea mauritiana</i>	Linn., 1758																					X
		<i>Cypraea moneta</i>	Linn., 1758											X		X								X
		<i>Cypraea nucleus</i>	Linn., 1758																					X
		<i>Cypraea poraria</i>	Linn., 1758											X		X								X
		<i>Cypraea schildererorum</i>	Iredale, 1939											X		X								X
		<i>Cypraea scurra</i>	Gmelin, 1791																					X
		<i>Cypraea talpa</i>	Linnaeus, 1758																					X
		<i>Cypraea teres</i>	Gmelin, 1791																					X
		<i>Cypraea tigris</i>	Linn., 1758																					X
		<i>Cypraea ventriculus</i>																						X
	Dialidae	<i>Cerithidium diplax</i>	Watson, 1886																					X
		<i>Dendrodoris nigra</i>	Stimpson, 1856																					X
	Fasciolaridae	<i>Peristernia crocea</i>	(Gray)													X		X						
		<i>Peristernia chlorostoma</i>	Sowerby, 1825																					X
		<i>Peristernia crocea</i>																						X
	Fissurellidae	<i>Diodora foveolata</i>																						X
		<i>Diodora granifera</i>	Pease, 1861																					X
		<i>Emarginula cf. dilecta</i>	A. Adams, 1852																					X
	Harpidae	<i>Harpa amouretta</i>	Röding, 1798																					X
	Hipponicidae	<i>Antisabia foliaceus</i>	Quoy and Gaimard, 1835																					X
		<i>Sabia conica</i>	(Schumacher, 1817)											X		X								X
	Juliidae	<i>Julia exquisita</i>	Gould, 1862																					X
	Littorinidae	<i>Littoraria coccinea</i>	Gmelin, 1791																					X
		<i>Littoraria pintado</i>	(Wood, 1828)											X		X								X
		<i>Littoraria scabra</i>	(Linnaeus, 1758)																					X
		<i>Littoraria undulata</i>	Gray, 1839											X		X								X
		<i>Littorina coccinea</i>	(Gmelin)											X		X								X
		<i>Peasiella tantilla</i>	Gould, 1849																					X
	Marginellidae	<i>Volvarina fusiformis</i>	Hinds, 1844																					X
	Mitridae	<i>Imbricaria olivaeformis</i>	Swainson, 1821																					X
		<i>Mitra acuminata</i>	Swainson, 1824																					X
		<i>Mitra columbelliformis</i>	Kiener													X		X						
		<i>Mitra cucumerina</i>	Lamarck, 1811																					X

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Mitra papalis</i>	Linn., 1758																						X
		<i>Scabricola casta</i>																							X
	Modulidae	<i>Modulus tectum</i>	Gmelin, 1791											X		X									X
	Muricidae	<i>Aspella cf. producta</i>	Pease, 1861																						X
	Nassariidae	<i>Nassarius crematus</i>	Hinds, 1844																						X
		<i>Nassarius dermstina</i>	(Gould)												X		X								X
		<i>Nassarius papliiosus</i>	Linnaeus, 1758																						X
		<i>Nassarius pauperus</i>	Gould, 1850																						X
	Naticidae	<i>Natica bougei</i>	Sowerby, 1908																						X
		<i>Natica gaulteriana</i>	Récluz, 1844																						X
		<i>Polinices melanostomus</i>	Gmelin, 1791												X		X								X
		<i>Polynices melanostomus</i>	Gmelin, 1791																						X
	Neritidae	<i>Nerita albicilla</i>	Linn., 1758												X		X								X
		<i>Nerita picea</i>	(Recluz, 1841)												X		X								X
		<i>Nerita plicata</i>	Linn., 1758												X		X								X
		<i>Nerita polita</i>	Linn., 1758												X		X								X
	Olividae	<i>Oliva paxillus</i>	Pease, 1860																						X
	Omalogyridae	<i>Omalogyra japonica</i>	Habe, 1972																						X
	Orbistellidae	<i>Orbistella regina</i>	Kay, 1979																						X
	Phasianellidae	<i>Tricolia variabilis</i>	Pease, 1861																						X
	Planaxidae	<i>Planaxis zonatus</i>	A. Adams												X		X								X
	Pyramidellidae	<i>Odostomia gulicki</i>	Pilsbry, 1918																						X
		<i>Odostomia stearnsiella</i>	Pilsbry, 1918																						X
		<i>Pyramidella sulcata</i>	A. Adams, 1854																						X
	Rissoidae	<i>Merelina wanawana</i>	Kay, 1979																						X
		<i>Powellsetia fallax</i>	Kay, 1979																						X
		<i>Pyramidelloides miranda</i>	A. Adams, 1861																						X
		<i>Rissoina abigua</i>	Gould, 1849																						X
		<i>Rissoina costata</i>	A. Adams, 1851																						X
		<i>Rissoina ephamilla</i>	Watson, 1886																						X
		<i>Schwartziella horrida</i>																							X
	Scaphandridae	<i>Acteocina hawaiiensis</i>	Pilsbry, 1921																						X
	Scissurellidae	<i>Scissurella coronata</i>																							X
	Siphonariidae	<i>Siphonaria normalis</i>	Gould, 1846																						X
		<i>Williamia radiata</i>	Pease, 1861																						X
	Strombidae	<i>Strombus dentatus</i>	Linn., 1759																						X
		<i>Strombus maculatus</i>	Sowerby, 1842																				X		X
	Terebridae	<i>Hastula lanceata</i>	Linn., 1758																						X
		<i>Terebra achatas</i>	Weaver, 1960																						X
		<i>Terebra affinis</i>	Gray, 1834																						X
		<i>Terebra babylonia</i>	Lamarck, 1822																						X
		<i>Terebra cf. argus</i>	Pilsbry, 1921																						X
		<i>Terebra crenulata</i>	(Linn., 1758)												X		X								X
		<i>Terebra felina</i>	Dillwyn, 1817																						X

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																						
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Osteichthyes	Acanthuridae	<i>Acanthurus achilles</i>	Shaw, 1803			x					x	x	x													
		<i>Acanthurus blochii</i>	Valenciennes, 1835										x					x								
		<i>Acanthurus dussimeri</i>	Valenciennes, 1835															x			x	x				
		<i>Acanthurus glaucopareius</i>	Cuvier, 1829									x		x				x								
		<i>Acanthurus guttatus</i>	Schneider, 1801											x												
		<i>Acanthurus nigricauda</i>	Duncker & Mohr, 1929																							
		<i>Acanthurus nigroris</i>	Valenciennes, 1835										x	x	x											
		<i>Acanthurus olivaceus</i>	Forster & Schneider, 1801										x	x	x											
		<i>Acanthurus thompsoni</i>	(Fowler, 1923)																x			x	x			
		<i>Acanthurus triostegus</i>	(Linnaeus, 1758)										x	x	x											
		<i>Ctenochaetus hawaiiensis</i>	Randall 1955												x	x										
		<i>Ctenochaetus marginatus</i>	(Valenciennes, 1835)													x										
		<i>Ctenochaetus strigosus</i>	(Bennett, 1828)											x	x	x										
		<i>Naso brevirostris</i>	(Valenciennes, 1839)																			x	x	x		
		<i>Naso hexacanthus</i>	(Bleeker, 1855)																							
		<i>Naso lituratus</i>	Forster & Schneider, 1801											x	x	x										
		<i>Naso unicornis</i>	(Forsskal, 1775)											x	x											
		<i>Zebrasoma flavescens</i>	(Bennett, 1828)											x	x	x										
		<i>Zebrasoma veliferum</i>	(Bloch, 1797)													x										
		Albulidae		<i>Albula glossodonta</i>	(Forsskal, 1775)																					
		Antennariidae		<i>Antennarius coccineus</i>	(Cuvier, 1831)																					
		Apogonidae		<i>Apogon coccineus</i>	Ruppell, 1838																					
				<i>Apogon erythrinus</i>	Snyder, 1904										x											
	<i>Apogon kallopterus</i>		Bleeker, 1856											x	x											
	<i>Apogon perdix</i>		Bleeker, 1854											x												
	<i>Apogon taeniopterus</i>		Bennett, 1835											x	x	x										
	<i>Epigonus atherinoides</i>		(Gilbert)																					x	x	
	<i>Pseudamiops gracilicauda</i>		Lachner, 1953																							
Aulostomidae		<i>Aulostomus chinensis</i>	(Linnaeus, 1766)										x	x	x								x	x		
	Balistidae	<i>Melichthys niger</i>	(Bloch, 1786)											x	x	x										
<i>Melichthys vidua</i>		(Solander, 1844)												x	x	x										
<i>Rhinecanthus aculeatus</i>		(L., 1758)												x	x	x										
<i>Sufflamen bursa</i>		(Bloch & Schneider, 1801)														x										
<i>Sufflamen fraenatus</i>		(Bloch & Schneider, 1801)																						x	x	
<i>Xanthichthys auromarginatus</i>		(Bennett, 1831)																					x	x	x	
		<i>Xanthichthys strigatus</i>	(Bennett, 1831)																							
Belonidae		<i>Platybelone argalus</i>	(Lesueur, 1821)																							
	Blenniidae	<i>Cirripectes vanderbilti</i>	(Fowler, 19380)																							
<i>Cirripectes variolosus</i>		(Valenciennes, 1836)																						x		
<i>Exallias brevis</i>		(Kner, 1868)																								
<i>Istiblennius gibbifrons</i>		(Quoy & Gaimard, 1824)																								
<i>Bothus mancus</i>		(Broussonet, 1782)																								
Bothidae		<i>Bothus mancus</i>	(Broussonet, 1782)																							
Bromidae		<i>Eumegistus illusris</i>	Jordan & Jordan, 1922																					x		
Brotulidae		<i>Brotula multibarbata</i>	Temminck & Schlegel, 1846																							
		<i>Brotula townsendi</i>	Fowler, 1900																							

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																						
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
	Callanthiidae	<i>Grammatonatus laysanus</i>	Gilbert, 1905															x					x	x		
	Callionymidae	<i>Synchiropus rosulentus</i>	Randall, 1999																							
	Carangidae	<i>Carangoides ferdau</i>	Forsskal, 1775				x				x	x						x								
		<i>Carangoides orthogrammus</i>	Jordan & Gilbert, 1881		x														x				x	x		
		<i>Caranx equula</i>	Schlegel, 1844					x																		
		<i>Caranx ignobilis</i>	(Forsskal, 1775)																	x						
		<i>Caranx lugubris</i>	Poey, 1860					x												x				x	x	
		<i>Caranx melampygus</i>	Cuvier and Valenciennes, 1833							x										x						
		<i>Caranx sexfasciatus</i>	Quoy & Gaimard, 1824																			x				
		<i>Decapterus marcarellus</i>	Cuvier, 1833																							
		<i>Elegatis bipinnulatus</i>	(Quoy & Gaimard, 1825)																							
		<i>Gnathanodon speciosus</i>	Forsskal, 1775																							
		<i>Scomberoides lysan</i>	(Forsskal, 1775)																							
		<i>Selar crumenophthalmus</i>	(Bloch, 1793)																							
		<i>Seriola dumerili</i>	(Risso, 1810)																							
		Carapodidae	<i>Carapus mourlani</i>	(Petit, 1934)																						
		Chaetodontidae	<i>Chaetodon auriga</i>	Forsskal, 1775		x						x	x	x										x	x	
			<i>Chaetodon citrinellus</i>	Cuvier 1831									x	x	x											
			<i>Chaetodon ephippium</i>	Cuvier 1831										x	x	x										
	<i>Chaetodon lineolatus</i>		Cuvier 1831																							
	<i>Chaetodon lunula</i>		(Lacepede, 1803)																							
	<i>Chaetodon lunulatus</i>		Quoy & Gaimard, 1825																							
	<i>Chaetodon miliaris</i>		(Quoy & Gaimard, 1835)																							
	<i>Chaetodon modestus</i>		Temminck & Schlegel, 1842																							
	<i>Chaetodon multicinctus</i>		Garrett, 1863										x	x	x											
	<i>Chaetodon ornatissimus</i>		Cuvier 1831										x	x	x											
	<i>Chaetodon quadrimaculatus</i>		Gray, 1831										x	x	x									x	x	
	<i>Chaetodon reticulatus</i>		Cuvier 1831																							
	<i>Chaetodon tinkeri</i>		Schultz, 1951																							
	<i>Chaetodon trifascialis</i>		(Quoy & Gaimard, 1825)																							
	<i>Chaetodon unimaculatus</i>		Bloch, 1788																							
	<i>Forciper flavissimus</i>		Jorfán & McGregor, 1898																							
	<i>Hemitaurichthys thompsoni</i>		Fowler, 1923																							
	<i>Heniochus diphreutes</i>	Jordan, 1903																								
	Cirrhitidae	<i>Amblycirrhites bimacula</i>	(Jenkins, 1903)																							
		<i>Cirrhites alternatus</i>	Gill, 1862																							
		<i>Cirrhites pinnulatus</i>	Bloch & Schneider, 1801																							
		<i>Paracirrhites arcatus</i>	(Cuvier 1831)																							
	Congridae	<i>Paracirrhites forsteri</i>	(Bloch & Schneider, 1801)																							
		<i>Ariosoma marginatum</i>	(Vaillant & Sauvage)																							
	Coryphaenidae	<i>Conger cinereus</i>	Ruppell, 1830																							
		<i>Conger oligoporus</i>	Kanazawa																							
	Diodontidae	<i>Coryphaena hippurus</i>	L., 1758																							
		<i>Diodon hystrix</i>	Linnaeus, 1758																							

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Parupeneus bifasciatus</i>	(Lacepede, 1801)	x							x	x													
		<i>Parupeneus chrysonemus</i>	Jordan & Evermann																			x	x		
		<i>Parupeneus cyclostomus</i>	(Lecepede, 1801)					x				x					x					x	x		
		<i>Parupeneus multifasciatus</i>	Quoy & Gaimard, 1824	x							x	x					x					x	x		
		<i>Parupeneus pleurostigma</i>	(Bennett, 1830)														x								
	Muraenidae	<i>Anarchias allardicei</i>	Jordan & Starks, 1906								x	x													
		<i>Anarchias cantonensis</i>	(Schultz, 1943)								x														
		<i>Anarchias leucurus</i>	Jordan and Starcks, 1906								x	x													
		<i>Echidna leucotaenia</i>	Schultz, 1943								x														
		<i>Echidna polyzona</i>	(Richardson, 1844)								x														
		<i>Echidna unicolor</i>	Schultz, 1953										x												
		<i>Enchelycore pardalis</i>	(Temminck & Schlegel, 1846)																			x			
		<i>Gymnomuraena zebra</i>	(Shaw, 1797)								x														
		<i>Gymnothorax berndti</i>	Snyder, 1904															x				2	2		
		<i>Gymnothorax buroensis</i>	(Bleeker, 1857)															x							
		<i>Gymnothorax eurostus</i>	(Abbott, 1860)								x	x													
		<i>Gymnothorax flavimarginatus</i>	(Ruppell, 1828)															x							
		<i>Gymnothorax fuscomaculatus</i>	(Schultz, 1953)								x	x													
		<i>Gymnothorax gracilicaudus</i>	Jenkins, 1903								x	x													
		<i>Gymnothorax javanicus</i>	(Bleeker, 1859)								x	x													
		<i>Gymnothorax margaritophorus</i>	(Bleeker, 1865)																			x			
		<i>Gymnothorax meleagris</i>	(Shaw and Nodder, 1795)								x	x													
		<i>Gymnothorax pindae</i>	(Smith, 1962)								x	x													
		<i>Gymnothorax undulatus</i>	(Lacepede, 1803)								x	x													
		<i>Gymnothorax nudivomer</i>	(Bleeker, 1864)																			x			
		<i>Gymnothorax nuttingi</i>	Snyder, 1904																			x			
		<i>Gymnothorax zonipectis</i>	Seale, 1917																			x			
		<i>Siderea picta</i>	(Ahl, 1789)																						
		<i>Uropterygius fuscoguttatus</i>	McCosker & Smith, 1997																						
		<i>Uropterygius inornatus</i>	Schultz, 1953								x	x													
		<i>Uropterygius inornatus</i>	Gosline, 1958																			x			
		<i>Uropterygius polyspilus</i>	Regan, 1905									x													
		<i>Uropterygius supraforatus</i>	(Regan, 1909)									x	x												
		<i>Uropterygius macrocephalus</i>	(Bleeker, 1865)										x												
		<i>Uropterygius tigrinus</i>	(Lesson, 1829)									x													
	Ophichthidae	<i>Brachysomophis sauropsis</i>	Schultz, 1943									x	x												
		<i>Leiuranus semicinctus</i>	(Lay & Bennett, 1839)									x	x												
		<i>Muraenichthys cookei</i>	Fowler, 1928									x	x												
		<i>Muraenichthys gymnotus</i>	Bleeker, 1857									x													
		<i>Muraenichthys schultzei</i>	Bleeker, 1864									x	x												
		<i>Myrichthys bleekeri</i>										x													
		<i>Myrichthys maculosus</i>	(Cuvier, 1917)									x	x									x	x		
		<i>Phyllophichthus xenodontus</i>	Gosline, 1951										x												
		<i>Schizomorhynchus labialis</i>	Seale, 1917									x													

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																					
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
		<i>Schultzia johnstonensis</i>	(Schultz & Woods, 1917)							x	x														
	Oplegnathidae	<i>Oplegnathus punctatus</i>	(Temminck & Schlegel, 1844)														x								
	Ostraciontidae	<i>Ostracion cubicus</i>	L., 1758					x																	
		<i>Ostracion meleagris</i>	Shaw, 1796	x		x				x	x	x													
		<i>Ostracion whitleyi</i>	Fowler, 1931			x						x	x												
	Percophididae	<i>Chrionema chyseres</i>	Gilbert, 1905														x								
		<i>Chrionema squamiceps</i>	Gilbert, 1905														x								
	Pleuronectidae	<i>Samariscus triocellatus</i>	Woods, 1966																		x				
	Polynemidae	<i>Polydactylus sexfilis</i>	(Cuvier and Valenciennes, 1931)	x																					
	Pomacanthidae	<i>Centropyge fisheri</i>	(Snyder, 1904)														x								
		<i>Centropyge loriculus</i>	(Gunther, 1873)			x	x			x		x													
		<i>Centropyge multicolor</i>	Randall & Wass, 1974																		x				
		<i>Centropyge nahackyi</i>	Kosaki, 1989																		x				
		<i>Centropyge nigriocellus</i>	Woods & Schultz, 1953					x				x													
		<i>Centropyge potteri</i>	(Jordan & Metz, 1912)														x								
	Pomacentridae	<i>Desmoholocanthus arcuatus</i>	(Gray, 1831)														x				x	x			
		<i>Abudefduf abdominalis</i>	(Quoy & Gaimard, 1824)																		x				
		<i>Abudefduf sordidus</i>	(Forsskal, 1775)			x							x												
		<i>Chromis acares</i>	Randall & Swerdloff, 1973								x						x								
		<i>Chromis agilis</i>	Smith, 1950			x				x	x	x													
		<i>Chromis verater</i>	Jordan & Metz																		x	x			
		<i>Dascyllus albisella</i>	Gill, 1863								x	x	x								x	x			
		<i>Dascyllus marginatus</i>				x																			
		<i>Plectroglyphidodon imparipennis</i>	(Vaillant & Sauvage, 1875)					x			x		x												
		<i>Plectroglyphidodon johnstonianus</i>	Fowler and Ball, 1925					x			x	x	x												
		<i>Plectroglyphidodon phoenixensis</i>	Schultz, 1843					x			x														
	Priacanthidae	<i>Cookeolus boops</i>	(Schneider, 1801)																		x				
		<i>Heteropriacanthus cruentatus</i>	(Lacepede, 1801)					x			x	x													
		<i>Priacanthus alalaua</i>	Jordan & Evermann																			x	x		
		<i>Priacanthus meeki</i>	Jenkins, 1903																		x				
	Scaridae	<i>Calotomus carolinus</i>	(Valenciennes, 1839)								x	x	x												
		<i>Chlorurus sordidus</i>	(Forsskal, 1775)					x			x	x	x												
		<i>Scarus cyanogrammus</i>		x																					
		<i>Scarus dubius</i>	Bennett, 1828	x							x	x	x												
		<i>Scarus duperreyi</i>		x																					
		<i>Scarus perspicillatus</i>	Steindacher, 1897	x							x	x	x												
		<i>Scarus psittacus</i>	Forsskal, 1775											x											
		<i>Scarus rubroviolaceus</i>	Bleeker, 1849																			x			
	Scombridae	<i>Acanthocybium solanderi</i>	(Cuvier, 1831)																			x			
		<i>Euthynnus affinis</i>	(Cantor, 1849)																				x		
		<i>Katsuwonis pelamis</i>	(L., 1758)																				x		

Taxa1	Taxa 2	Genus/Species	Authority	Reference Number																						
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
		<i>Thunnus albacarea</i>	(Bonnaterre, 1788)														x									
	Scorpaenidae	<i>Dendrochirus barberi</i>	(Steindachner, 1900)														x									
		<i>Neomerinthe rufescens</i>	(Gilbert)																		x	x				
		<i>Pontinus macrocephalus</i>	(Sauvage, 1882)														x				x	x				
		<i>Scorpaena colorata</i>	(Gilbert, 1905)														x									
		<i>Scorpaena kelloggi</i>	(Jenkins, 1903)														x									
		<i>Scorpaenodes hirsutus</i>	(Smith, 1957)																	x						
		<i>Scorpaenodes parvipinnis</i>	(Garrett, 1864)									x														
		<i>Scorpaenopsis diabolus</i>	(Cuvier and Valenciennes, 1829)														x									
		<i>Scorpaenopsis fowleri</i>	(Pietschmann, 1934)																	x						
		<i>Sebastapistes ballieui</i>	(Sauvage, 1875)								x															
		<i>Sebastapistes conioarta</i>	(Jenkins, 1903)								x															
	Serranidae	<i>Anthias fuscinus</i>	Randall & Ralston, 1984														x									
		<i>Anthias ventalis</i>	Randall 1979														x									
		<i>Aporops bilinearis</i>	Schultz, 1943																	x						
		<i>Epinephalus quernus</i>	Seale, 1901														x				x	x				
		<i>Holanthias elizabethae</i>	(Fowler, 1923)														x				x	x				
		<i>Holanthias fuscipinnis</i>	(Jenkins, 1901)														x				x	x				
		<i>Liopropoma collettei</i>	Randall & Taylor, 1988																	x						
		<i>Plectranthias helenae</i>	Randall, 1980														x				x	x				
		<i>Plectranthias winniensis</i>	(Tyler, 1966)																	x						
		<i>Promicrops lanceolatus</i>	(Bloch, 1790)														x									
		<i>Pseudanthias bicolor</i>	(Randall, 1979)																	x						
		<i>Pseudanthias fucinus</i>	(Randall & Ralston)																		x	x				
		<i>Pseudanthias randalli</i>	(Lubbock & Allen, 1978)																	x						
	Sphyraenidae	<i>Sphyraena barracuda</i>	(Walbaum, 1792)														x									
		<i>Sphyraena helleri</i>	Jenkins, 1901				x																			
	Syngnathidae	<i>Doryrhamphus excisis</i>	Kaup, 1856																			x				
	Synodontidae	<i>Saurida flamma</i>	Waples, 1982																			x				
		<i>Saurida gracillis</i>	Quoy & Gaimard, 1824								x	x														
		<i>Synodus binotatus</i>	Schultz, 1953								x	x														
		<i>Synodus engelmani</i>	Schultz, 1953																			x				
		<i>Synodus variegatus</i>	Lacepede, 1803																			x				
	Tetraodontidae	<i>Arothron meleagris</i>	(Bloch & Schneider, 1801)			x					x	x	x													
		<i>Canthigaster coronata</i>	(Vaillant & Sauvage, 1875)																			x				
		<i>Canthigaster inframacula</i>	Allan & Randall, 1977																			x				
		<i>Canthigaster jactator</i>	(Jenkins, 1901)																							
	Triacanthodidae	<i>Hollardia goslinei</i>	Tyler, 1968									x	x	x								x				
	Triglidae	<i>Satyricthys engyceros</i>	(Gunther, 1871)																			x	x			
	Xenocongridae	<i>Kaupichthys hyproroides</i>	(Stromamm, 1896)																							
	Zanclidae	<i>Zanclus cornutus</i>	(Linnaeus, 1758)				x					x		x								x	x			
Reptilia	Cheloniidae	<i>Chelonia mydas</i>	Bocourt, 1835																							
Total Reports					25	92	68	4	12	11	113	148	77	85	149	4	149	6	102	32	26	12	61	102	195	165

APPENDIX B

Previous Reports of Marine Organisms Observed or Collected at Johnston Atoll with 1-2 Listings

Reference	Taxa 1	Taxa2	Genus, Species	Authority, Date	Cited as	Year
Baker et al. 1997	Planta	Chlorophyta	<i>Borzia elongata</i>			1995
Moul 1964	Planta	Chlorophyta	<i>Halimeda tuna</i>	Moul		1953
Cohen 1997	Anthozoa	Pocilloporidae	<i>Pocillopora meandrina</i>	(Dana, 1846)		1996
Cohen 1997	Anthozoa	Acroporidae	<i>Acropora cythera</i>	(Dana, 1846)		1996
Cairns 1984	Anthozoa	Flabellidae	<i>Javania lamprotichum</i>	Mosley, 1880		
Kay 1961	Gastropoda	Cypraeidae	<i>Cypraea tigris</i>	Linn., 1758	<i>Cypraea tigris schilderiana</i>	
Brock 1979	Gastropoda	Cypraeidae	<i>Cypraea tigris</i>	Linn., 1758		1970
Economakis and Lobel 1998	Elasmobranchii	Carcharhinidae	<i>Carcharhinus amblyrhynchos</i>	(Bleeker, 1856)		1992-95
Randall et al. 1977	Elasmobranchii	Carcharhinidae	<i>Triaenodon obesus</i>	(Ruppel, 1837)		1968-71
McCosker and Smith 1997	Osteichthyes	Muraenidae	<i>Uropterygius fuscoguttatus</i>	McCosker & Smith, 1997		1968
Dee and Parrish 1994	Osteichthyes	Holocentridae	<i>Myripristis amaena</i>	(Castelnau, 1873)		1993
Randall 1999	Osteichthyes	Callionymidae	<i>Synchiropus rosulentus</i>	Randall, 1999		
Irons 1989	Osteichthyes	Chaetodontidae	<i>Chaetodon trifascialis</i>	(Quoy & Gaimard, 1825)		1988
Irons 1990	Osteichthyes	Chaetodontidae	<i>Chaetodon trifascialis</i>	(Quoy & Gaimard, 1825)		1988
Kosaki 1989	Osteichthyes	Pomacanthidae	<i>Centropyge nahackyi</i>	Kosaki, 1989		1987-88
Kerr and Lobel 1997	Osteichthyes	Pomacentridae	<i>Abudefduf sordidus</i>	(Forsskal, 1775)		1995
Mann and Lobel 1995	Osteichthyes	Pomacentridae	<i>Dascyllus albisella</i>	Gill, 1863		1995
Mann and Lobel 1998	Osteichthyes	Pomacentridae	<i>Dascyllus albisella</i>	Gill, 1863		1995
Lobel 1997	Osteichthyes	Pomacentridae	<i>Plectroglyphidodon imparipennis</i>	(Vaillant & Sauvage, 1875)		1995
Randall 1972	Osteichthyes	Labridae	<i>Anampses cuvier</i>	Quoy & Gaimard, 1824		1970
Randall 1972	Osteichthyes	Labridae	<i>Anampses cuvier</i>	Quoy & Gaimard, 1824		1970
Gorka et al. 1997	Osteichthyes	Acanthuridae	<i>Ctenochaetus strigosus</i>	(Bennett, 1828)		1995
Ackman et al. 1992	Reptilia	Cheloniidae	<i>Chelonia mydas</i>	Bocourt, 1835		1992
Balazs 1994	Reptilia	Cheloniidae	<i>Chelonia mydas</i>	Bocourt, 1835		1992

APPENDIX C

Marine Organisms Observed or Collected in Present Study Listed by Station
Nonindigenous or Cryptogenic Species in Bold
New Johnston Atoll Reports Marked by Asterisks

Taxa	Genus_Species	Author, Date	Station											
			1	2	3	4	5	6	7	8	9	10	11	
Macroalgae														
Cyanophyta	<i>Ulothrix pseudoflacca*</i>	Wille		X		X								
	Unidentified spp.			X					X					
Chlorophyta	<i>Acetabularia</i> sp.								X					
	<i>Bryopsis hypnoides*</i>	Lamouroux		X					X				X	X
	<i>Bryopsis pennata</i>	Lamouroux										X	X	
	<i>Bryopsis</i> sp.						X							
	<i>Caulerpa ambigua</i>	(Okamura) Prudhomme van Reine & Lockhorst						X						
	<i>Caulerpa lentillifera*</i>	J. Agardh		X										
	<i>Caulerpa racemosa</i>	(Forsskål) J. Agardh			X									X
	<i>Caulerpa serrulata*</i>	(Forsskål) J. Agardh	X	X		X	X							
	<i>Caulerpa webbiana*</i>	Montagne			X								X	
	<i>Cladophora</i> sp.			X						X			X	
	<i>Dictyosphaeria cavernosa*</i>	(Forsskål) Børgesen			X	X						X		
	<i>Dictyosphaeria versluysii</i>	Weber-van Bosse		X					X	X				
	<i>Enteromorpha clathrata*</i>	(Roth) Greville		X									X	
	<i>Halimeda discoidea</i>	Decaisne			X	X								
	<i>Halimeda opuntia*</i>	(Linnaeus) Lamouroux								X				
	<i>Trichosolen oahuensis*</i>	Egerod						X						
	<i>Trichosolen</i> sp.		X											
	Unidentified sp.				X									
Phaeophyta	<i>Dictyopteris repens*</i>	(Okamura) Børgesen	X					X						
	<i>Dictyopteris</i> sp.			X										
	<i>Dictyota acutiloba*</i>	J. Agardh											X	
	<i>Dictyota divaricata*</i>	Lamouroux		X				X						
	<i>Dictyota</i> sp.				X	X		X	X				X	
	<i>Lobophora variegata</i>	(Lamouroux) Womersley	X	X	X	X	X	X	X	X	X	X		X
	<i>Rosenvingea</i> sp.*							X	X					
	<i>Sphacelaria novae-hollandiae</i>	Sonder		X						X				
	<i>Sphacelaria tribuloides</i>	Meneghini								X				
Rhodophyta	<i>Aglaothamnion boergesenii*</i>	(Aponte & Ballantine) L'Hardy-Halos												X
	<i>Anotrichium secundum</i>	(C. Agardh) Nageli				X								
	<i>Anotrichium tenue</i>	(C. Agardh) Nageli	X		X				X		X			
	<i>Antithamnion antillanum</i>	Børgesen			X		X			X			X	
	<i>Antithamnion</i> sp.					X								
	<i>Antithamnionella breviramosa*</i>	Dawson		X										
	<i>Botryocladia</i> sp.*									X				
	<i>Caulacanthus ustulatus*</i>	(Mertens) Kützing	X					X	X					
	<i>Centroceras clavulatum</i>	(C. Agardh) Montagne						X						
	<i>Centroceras minutum*</i>	Yamada							X					
	<i>Ceramium aduncum*</i>	Nakamura		X										
	<i>Ceramium borneense*</i>	Weber Bosse											X	
	<i>Ceramium codii</i>	(Richards) G. Mazoyer	X							X				
	<i>Ceramium flaccidum</i>	(Harvey) Mazoyer		X	X	X	X							
	<i>Ceramium macilentum*</i>	J. Agardh				X		X		X				X
	<i>Ceramium serpens*</i>	Setchell & Gardner									X			
	<i>Ceramium vagans</i>	Silva	X		X									
	<i>Champia parvula</i>	(C. Agardh) Harvey				X							X	
	<i>Chondracanthus</i> sp*.					X								
	<i>Chondria polyrhiza</i>	Collins & Hervey												X
	<i>Chondria</i> sp.				X	X								
	<i>Corallophila apiculata*</i>	(Yamada) R.E. Norris				X								
	<i>Corallophila huysmansii</i>	(Weber Bosse) R.E. Norris			X	X				X				
	<i>Crouania</i>	Itono								X				

<i>mageshimensis*</i>												
<i>Dasya iridescens*</i>	(Schlech) Milar & Abbott	X										
<i>Dasya kristeniae*</i>	Abbott			X								
<i>Dasya murrayana*</i>	Abbott & Millar		X									
<i>Dasya sp.*</i>								X		X		
<i>Diplothamnion jolyi</i>	van den Hoek									X		
<i>Gelidiella sp.*</i>				X	X						X	
<i>Gelidiopsis variabilis*</i>	(J. Agardh) Schmitz					X						
<i>Griffithsia heteromorpha*</i>	Kützing	X		X	X		X	X	X		X	
<i>Griffithsia schousboei*</i>	Montagne										X	
<i>Griffithsia subcylindrica*</i>	Okamura		X						X			
<i>Herposiphonia arcuata*</i>	Hollenberg					X						
<i>Herposiphonia crassa*</i>	Hollenberg					X						
<i>Herposiphonia delicatula*</i>	Hollenberg	X	X									
<i>Herposiphonia parca*</i>	Setchell										X	
<i>Herposiphonia secunda*</i>	(C. Agardh) Ambronn							X				
<i>Herposiphonia sp.*</i>						X				X	X	
<i>Herposiphonia variabilis*</i>	Hollenberg			X								
<i>Heterosiphonia crispella*</i>	(C. Agardh) Wynne			X							X	
<i>Hypnea pannosa*</i>	J. Agardh	X										
<i>Hypnea sp.</i>		X									X	
<i>Hypnea spinella*</i>	(C. Agardh) Kützing	X	X	X		X		X				
<i>Hypnea valentiae*</i>	(Turner) Montagne					X					X	
<i>Jania adhaerens*</i>	Lamouroux										X	
<i>Jania pumila*</i>	Lamouroux				X	X						
<i>Jania sp.</i>		X										
<i>Kallymenia sp.*</i>		X										
<i>Laurencia majuscula*</i>	(Harvey) Lucas			X								
<i>Laurencia sp.*</i>				X	X			X				
<i>Lomentaria hakodatensis</i>	Yendo			X		X	X		X	X		
<i>Lomentaria sp.*</i>					X						X	
<i>Lophosiphonia prostrata*</i>	(Harvey) Falkenberg							X				
<i>Monosporus indicus*</i>	Børgesen							X			X	
Neosiphonia						X						
sphaerocarpa*												
<i>Peyssonnelia conchicola*</i>	Piccone & Grunow			X								
<i>Peyssonnelia inamoena*</i>	Pilger						X				X	
<i>Peyssonnelia sp.*</i>		X			X				X	X		
<i>Polysiphonia delicatula*</i>	Hollenberg	X										
<i>Polysiphonia exilis*</i>	Harvey								X			
<i>Polysiphonia howei*</i>	Hollenberg	X										
<i>Polysiphonia scopulorum*</i>	Harvey					X		X				
<i>Polysiphonia sp.*</i>		X	X	X			X					
<i>Pterocladella caloglossoides*</i>	(Howe) Santelices					X						
<i>Ptilothamnion cladophorae*</i>	(Yamada & Tanaka) Feldmann	X										
<i>Spermathamnion sp.*</i>				X		X						
<i>Stylonema alsidii*</i>	(Zanardini) Howe										X	
<i>Taenioma perpusillum*</i>	(J. Agardh) J. Agardh							X				
<i>Tiffaniella sp.*</i>						X						
Total Algae		9	28	24	25	19	16	11	17	10	21	13
Porifera												
Callyspongiidae	<i>Callyspongia sp.*</i>					X						
Chondrillidae	<i>Chondrosia sp.*</i>			X	X							
Clionidae	<i>Cliona sp.*</i>										X	
Dysideidae	<i>Dysidea sp.1*</i>	X										
	<i>Dysidea sp.2*</i>	X						X				

Olividae	<i>Peristernia sp. (juv)</i>						X			X
	<i>Oliva sp.</i>						X			X
Marginellidae	<i>Dentimargo sp.</i>			X			X		X	X
	<i>Gramelina sp.</i>			X			X			X
	<i>Granula sandwichensis*</i>	Pease, 1860					X			X
	<i>Granulina</i>		X	X			X	X	X	X
	<i>Gyrincum sp.</i>								X	
	<i>Marginella sp.1</i>		X				X			X
	<i>Marginella sp.2</i>			X					X	
	<i>Volvarina sp.</i>			X						
Mitridae	<i>Imbricaria olivaeformis</i>	Swainson, 1821					X		X	X
	<i>Mitridae sp.</i>								X	
Costellariidae	<i>Vexillum unifasciatum*</i>	Wood, 1828					X			X
Pyramidellidae	<i>Evalea sp. (waikikiensis?)*</i>	Pilsbry, 1918							X	
	<i>Herviera gliriella*</i>	Melville and Standen, 1896				X				
	<i>Kolonella sp. (hawaiiensis?)*</i>	Kay, 1979						X	X	
	<i>Miralda sp.</i>		X	X					X	X
	<i>Odostomia gulicki</i>	Pilsbry, 1918								X
	<i>Odostomia oxia*</i>	Watson, 1886								X
	<i>Otopleura mitralis*</i>	Adams, 1854			X					
	<i>Turbonilla cornelliana*</i>	Newcomb, 1870						X		
Actaeonidae	<i>Pupa tessellata</i>	Reeve, 1842				X				X
Siphonariidae	<i>Williamia radiata</i>	Pease, 1861	X							X
	<i>Siphonaria sp.</i>							X		
Scaphandridae	<i>Acteocina sp.</i>			X						
Conidae	<i>Conus lividus</i>	Hwass in Bruguière, 1792						X		
	<i>Conus lividus (juv)</i>	Hwass in Bruguière, 1792						X	X	X
	<i>Conus miles</i>	Linnaeus, 1758						X		
	<i>Conus pulicarius</i>	Hwass in Bruguière, 1792			X				X	
	<i>Conus rattus</i>	Hwass in Bruguière, 1792						X	X	
	<i>Conus sp. (juv)</i>							X		
Costellariidae	c.f. <i>Pusia microzonias*</i>	Lamarck, 1811				X				
Mollusca-Opisthobranchia										
Atyidae	<i>Atys curta*</i>	Adams, 1850						X		
	<i>Atys debilis</i>	Pease, 1860			X	X		X	X	
	<i>Atys semistriata</i>	Pease, 1860			X					
	<i>Diniatys dentifer</i>	Adams, 1850				X		X		
Aglajidae	<i>Philinopsis sp.</i>				X					
Aplysiidae	<i>Aplysia sp</i>						X			X
	<i>Aplysia sp. (juiliana?)*</i>	Quoy and Gaimard, 1832								X
	<i>Dolabella auricularia*</i>		X							X
Pleurobranchidae	<i>Berthellina citrina*</i>	Rupell and Leuckart, 1831			X	X				
	<i>Pleurobranchia sp.</i>				X					
Juliidae	<i>Julia exquisita*</i>	Gould, 1862	X	X				X		X
Dorididae	<i>Chromodoris vibrata</i>	Pease, 1860								X
	<i>Chromodoris sp.</i>				X					
	<i>Risbecia imperialis*</i>	Pease, 1860								X
Dendrodorididae	<i>Dendrodoris nigra*</i>	Stimpson, 1856								X
Mollusca-Bivalvia										
Arcidae	<i>Arca ventricosa</i>	Lamarck, 1819			X	X			X	
	<i>Barbatia decussata*</i>	Sowerby, 1823			X					
	<i>Barbatia divaricata*</i>	Sowerby, 1833						X		
	<i>Benthacara decorata?</i>	Hayami and Kase, 1993						X		
Mytilidae	<i>Crenella sp. (c.f. Hayami and Kase, 1993)</i>					X				
	<i>Lithophaga sp.</i>							X		
	<i>Septifer bryanae</i>	Pilsbry, 1921					X		X	X
Pinnidae	<i>Pinna muricata</i>	Linnaeus, 1758			X				X	X
Pteriidae	<i>Pinctada margaritifera</i>	Linnaeus, 1758			X	X				
	<i>Pinctada radiata*</i>	Leach, 1814			X					X
	<i>Pteria sp.</i>					X				
Isognomonidae	<i>Isognomon perna</i>	Linnaeus, 1767			X	X	X		X	
Malleidae	<i>Malleus regula</i>	Forskål, 1775				X				
Pectinidae	<i>Chlamys coruscans hawaiiensis*</i>	Dall, Bartsch, and Rehder, 1938			X					
	<i>Mirapecten mirificus</i>	Reeve, 1853					X			
Spondylidae	<i>Spondylus sp.1.</i>		X						X	
	<i>Spondylus sp.2.</i>								X	
	<i>Spondylus sp.3.</i>								X	

	<i>Labroides phthirophagus</i>	Randall 1958							X						
	<i>Novaculichthys taeniourus</i>	(Lacepede, 1801)					X								X
	<i>Oxycheilinus unifasciatus</i>	Streets, 1877			X	X			X		X	X		X	X
	<i>Pseudocheilinus octotaenia</i>	Jenkins, 1900				X								X	
	<i>Stethojulis balteata</i>	(Quoy & Gaimard, 1824)				X	X				X			X	
	<i>Thalassoma ballieui</i>	(Vaillant and Sauvage, 1875)				X	X	X			X				
	<i>Thalassoma duperreyi</i>	(Quoy and Gaimard, 1824)	X	X	X	X	X	X	X	X	X	X	X	X	X
	<i>Thalassoma lutescens</i>	(Lay and Bennett, 1839)			X		X	X			X	X			
Scaridae	<i>Chlorurus perspicillatus</i>	Steindacher, 1897									X				X
	<i>Chlorurus sordidus</i>	(Forskål, 1775)	X	X	X	X	X	X	X	X	X	X	X	X	X
Gobiidae	<i>Nemateleotris magnifica</i>	Fowler, 1938					X				X				
Scombridae	<i>Thunnus albacarea</i>	(Bonnaterre, 1788)	X												
Acanthuridae	<i>Acanthurus achilles</i>	Shaw, 1803			X	X	X	X	X		X	X			X
	<i>Acanthurus blochii</i>	Valenciennes, 1835	X	X	X	X	X	X	X	X	X	X	X	X	X
	<i>Acanthurus dussimeri</i>	Valenciennes, 1835									X				
	<i>Acanthurus nigroris</i>	Valenciennes, 1835	X	X	X	X	X			X	X	X	X	X	X
	<i>Acanthurus olivaceus</i>	Forster & Schneider, 1801	X			X	X			X	X			X	
	<i>Acanthurus thompsoni</i>	(Fowler, 1923)					X								
	<i>Acanthurus triostegus</i>	(Linnaeus, 1758)	X	X	X			X	X		X	X	X	X	X
	<i>Ctenochaetus hawaiiensis</i>	Randall 1955				X	X				X				
	<i>Ctenochaetus strigosus</i>	(Bennett, 1828)	X	X	X	X	X	X	X	X	X	X	X	X	X
	<i>Naso lituratus</i>	Forster & Schneider, 1801	X	X	X	X	X	X	X	X	X	X	X	X	X
	<i>Zebrasoma flavescens</i>	(Bennett, 1828)			X	X	X	X		X	X			X	X
	<i>Zebrasoma veliferum</i>	(Bloch, 1797)												X	
Zanclidae	<i>Zanclus cornutus</i>	(Linnaeus, 1758)	X	X	X	X	X			X	X	X			
Bothidae	<i>Bothus mancus</i>	(Broussonet, 1782)	X												
Balistidae	<i>Melichthys niger</i>	(Bloch, 1786)			X	X	X	X			X	X			
	<i>Melichthys vidua</i>	(Solander, 1844)	X	X		X	X				X			X	
	<i>Rhinecanthus aculeatus</i>	(L., 1758)	X			X	X			X					X
	<i>Sufflamen bursa</i>	(Bloch & Schneider, 1801)					X				X				
	<i>Xanthichthys auromarginatus</i>	(Bennett, 1831)					X								
Monocanthidae	<i>Cantherhines dumerilii</i>	(Hollard, 1854)				X									
	<i>Cantherhines sandwichiensis</i>	(Quoy and Gaimard, 1824)					X	X							
	<i>Pervagor spilosoma</i>	(Lay & Bennett, 1839)													X
Ostraciontidae	<i>Ostracion meleagris</i>	Shaw, 1796					X	X		X		X			
	<i>Ostracion whitleyi</i>	Fowler, 1931						X							X
Tetraodontidae	<i>Arothron meleagris</i>	(Bloch & Schneider, 1801)			X	X		X	X	X	X	X	X		X
	<i>Canthigaster jactator</i>	(Jenkins, 1901)	X								X				
	Total Fish		25	32	36	52	39	20	38	47	25	37	30		
	Total Taxa		153	208	175	214	177	145	152	180	144	152	149		