


DT: 76565

NIOZ-RAPPORT 1994 - 1
NIOO-CEMO RAPPORTEN EN VERSLAGEN 1994 - 1

DISTRIBUTION OF THE ZOOBENTHOS ON THE DUTCH CONTINENTAL SHELF: THE WESTERN FRISIAN FRONT, BROWN BANK AND BROAD FOURTEENS (1992/1993)

S. E. Holtmann, A. Groenewold



Nederlands Instituut voor Onderzoek der Zee

MILZON - BENTHOS II



© 1994

This report is not to be cited without the
acknowledgement of the source.

Netherlands Institute for Sea Research (NIOZ)
P.O. Box 59, 1790 AB Den Burg,
Texel, The Netherlands

Netherlands Institute of Ecology-
Centre for Estuarine and Coastal Ecology (NIOO-CEMO)
Vlierstraat 28, 4401 EA Yerseke
The Netherlands

North Sea Directorate
Ministry of Transport and Public Works
P.O. Box 5807, 2280 HV Rijswijk
The Netherlands

ISSN 0923 - 3210

Cover design: H. Hobbelink

Rijkswaterstaat

Bibliotheek

Directie Noord-Nederland

**DISTRIBUTION OF THE ZOOBENTHOS ON THE DUTCH
CONTINENTAL SHELF: THE WESTERN FRISIAN FRONT,
BROWN BANK AND BROAD FOURTEENS
(1992/1993).**

S.E. Holtmann¹, A. Groenewold²

This study was commissioned by the North Sea Directorate of the Ministry of Transport, Public Works and Water Management.

**¹ NETHERLANDS INSTITUTE FOR SEA RESEARCH
Department of Benthic Systems**

**² NETHERLANDS INSTITUTE OF ECOLOGY-CENTRE
FOR ESTUARINE AND COASTAL ECOLOGY**

MILZON - BENTHOS II
NIOZ-RAPPORT 1994 - 1
NIOO-CEMO RAPPORTEN EN VERSLAGEN 1994 - 1

SUMMARY

This report presents the results of the 1992 and 1993 surveys of the MILZON-BENTHOS II project. The aim of the research program was to study the spatial distribution of the zoobenthos in the area of the Broad Fourteens, Brown Bank (Southern North Sea) and the western Frisian Front (transition Southern-Central North Sea).

The fieldwork was carried out in spring 1992 and 1993 and included sediment sampling at 82 stations. From these samples the meiobenthos, the macrobenthos and selected sediment parameters, such as median grain size, the mud content, chlorophyll-a and POC, were analyzed. Additional data from 3 stations located in the area, which were sampled during the EXP*BMN program 1992 (Duineveld & Belgers, 1993; Huys & De Smet, 1993), have been incorporated in the results.

The investigated area can be described as follows. The southern part of the area is situated in medium to coarse sand, with depths around 30 m at the Broad Fourteens and between 20 and 40 m around the Brown Bank area. Further north, around 53°30' N, between the 30 and 40 m isobaths, a rather sharp boundary separates the sandy Southern Bight from the muddy Frisian Front area and Central North Sea. The north-west corner of the research area covered a sandy part of the Cleaver Bank, whereas the north-eastern area was located in the muddy Oyster Ground (depth > 40 m). The north-western part of the area, investigated in 1993, included the Western Mud Hole and partly the Botney Ground and one station was situated in the Botney Cut.

The results of the zoobenthos survey can be summarized as follows:

- * - In total 14 permanent (= real) meiobenthic taxa were found in the whole area. The density of the meiobenthos varied between 412 and 6815 ind./10 cm² with an average value of 1783 ind./10 cm².
- The Nematoda formed the most dominant taxon of the meiobenthos, accounting for 82 % of the total density. Highest densities were found in the muddy areas (western Frisian Front and Oyster Ground), whereas they were lowest in the southern area. The sandy southern part (Brown Bank and Broad Fourteens) and the Cleaver Bank contained relatively high densities of the meiobenthic taxa Copepoda (interstitial), Gastrotricha, Turbellaria, Annelida, Hydrozoa and Tardigrada. The Frisian Front area was marked by high Nematoda densities and relatively high abundances of Halacarida and Kinorhyncha, whereas the Oyster Ground contained high densities of Nematoda, burrowing Copepoda and the presence of Kinorhyncha, Rotifera and Priapulida.
- The meiobenthic diversities ($Hill_{0..\infty}$) were highest in the southern sandy area and the Cleaver Bank, whereas they were lowest in the western Frisian Front and in the Oyster Ground.
- * - From 85 sorted macrobenthos samples in total 160 species were identified. The taxa of the Polychaeta and Crustacea represent the main part of the identified macrobenthic species.
- The macrobenthic density varied between 257 and 4126 ind./m² (mean: 1308 ind./m²). The biomass of the macrobenthos showed values between 1 and 136 g AFDW/m² (mean: 18 g AFDW/m²). In the northern part the density was higher than in the area of the Southern Bight. Highest biomass values were found in the north and in the south-western part of the investigated area. About 40 % of the total density was made up by the Polychaeta. The biomass of the Echinodermata formed 52 % of the total biomass.
- The number of macrobenthic species per boxcore sample ($Hill_0$) varied between 11 and 33, with the lowest values occurring in the south-eastern part of the studied area. Stations with relatively low values of the diversity numbers $Hill_1$ to $Hill_\infty$ were situated in the north, where a few species numerically dominate the macrobenthic assemblage.

- * The observed zoobenthos composition showed high correlations with the sediment composition, especially with median grain size, mud content and, to a less extent, with the waterdepth. In areas where the sediment has a high mud content and a low median grain size, the macrobenthic assemblages were characterized by a high total biomass, total density and species richness. On the contrary, these areas revealed low meiobenthic diversity values, as a result of the absence of many meiobenthic taxa.

In order to get more information about the spatial distribution of the meio- and macrobenthic assemblages, the stations were classified using the TWINSPAN-cluster analysis based on taxon-level for the meiobenthic density, respectively on species-level for the macrobenthic density. Meio- and macrobenthos clusters showed in general the same distribution; two clusters in the northern muddy area and two clusters in the medium to coarser sandy area in the south. The borders of the two southern meio- and macrobenthos clusters did not coincide perfectly, probably due to different critical grain size ranges for both groups. The meiobenthos seems to be more sensitive to the range of median grain sizes between 250 and 300 μm , whereas the macrobenthos reacts to changes between 200 and 250 μm .

The clusters of the meio- and macrobenthos can be described as follows:

- * In the north cluster 1 of the macrobenthos showed the highest values of the total density and biomass, whereas in the south (cluster 4) the lowest values of the density and biomass were found. In the central area between these two clusters the macrobenthos was intermediate in terms of density and biomass (clusters 2 & 3).
- * The meiobenthos was represented in relatively high densities but with low diversities in the north in the western Frisian Front and Oyster Ground area (cluster 1). This area is surrounded by an area, around the 30 and 40 m isobaths, with lower densities but still impoverished in taxa (cluster 2). In the south total densities were generally lower, but a more diverse assemblage could be found (cluster 3), especially in the coarsest sandy parts (cluster 4).

The distribution of the meio- and macrobenthic assemblages in the Southern Bight as well as in the southern part of the Oyster Ground and the Frisian Front area seemed to be relatively stable in the period between 1988 and 1993 (MILZON-BENTHOS I+II).

The clusters of the meio- and macrobenthos of the total MILZON-BENTHOS area 1988-1993 were situated as follows:

- * The macrobenthos showed two assemblages in the muddy sand area of the Oyster Ground and Frisian Front (clusters I & II), a coastal assemblage (cluster III) and an assemblage of coarse sand area (cluster IV).
- * The meiobenthos showed an assemblage (cluster I) of stations in the muddy fine grained sand (Oyster Ground, Frisian Front and near river outflows). Cluster II is concentrated in the coastal area, in the fine-medium sandy area between 53° N and the 30 m isobath and in the less muddy parts of the northern area. Cluster III is mainly found in the Southern Bight, whereas cluster IV is found in the coarsest sediments.

SAMENVATTING

Dit rapport geeft een verslag van de resultaten van de inventarisaties van 1992 en 1993 van het MILZON-BENTHOS II project. Het doel van het onderzoeksprogramma was om de ruimtelijke verspreiding van het zoöbenthos van de Breeveertien, Bruine Bank (Zuidelijke Noordzee) en het westelijke Friese Front (overgang Zuidelijke-Centrale Noordzee) te bestuderen.

Het veldwerk vond in het voorjaar van 1992 en 1993 plaats, waarbij op 82 stations bodemonsters werden genomen. Van deze monsters werden meiobenthos, macrobenthos en enkele geselecteerde sediment parameters (medianen korrelgrootte, slibgehalte, chlorofyl-a en POC) geanalyseerd. Aanvullende data van 3 in hetzelfde gebied gelegen stations, bemonsterd gedurende het EXP*BMN programma 1992 (Duineveld & Belgers, 1993; Huys & De Smet, 1993), zijn in de resultaten geïntegreerd.

Het betrokken gebied kan als volgt worden omschreven. Het zuidelijke deel bestaat uit medium tot grofzandige gebieden, met dieptes van rond de 30 m in de Breeveertien en tussen de 20 en 40 m rond de Bruine Bank. Verder naar het noorden, ter hoogte van 53°30' N, tussen de 30 en 40 m dieptelijnen, scheidt een tamelijk scherpe grens de zandige Zuidelijke Noordzee van het slibrijke Friese Front gebied en de Centrale Noordzee. De noordwest hoek van het studiegebied bedekte het zandige deel van de Klaverbank, terwijl het noordoostelijke deel in de uit fijn slijf bestaande Oestergronden lag (> 40 m diep). Het noordwestelijke deel van het gebied, dat in 1993 werd bemonsterd, omvatte tevens de Western Mud Hole en gedeeltelijk de Botney Ground, met tevens één station in de Botney Cut.

De resultaten van de studie aan het zoöbenthos kunnen als volgt worden samengevat:

- * - In totaal zijn 14 permanente (= echte) meiobenthische taxa in het gebied van 1992/93 gevonden. De dichtheid van het meiobenthos varieerde tussen 412 en 6815 ind./10 cm², met een gemiddelde van 1783 ind./10 cm².
- De Nematoda vormden het meest dominante taxon in het meiobenthos, waarbij ze verantwoordelijk waren voor 82 % van de totale dichtheid. De hoogste dichtheden werden in slibrijke gebieden aangetroffen (westelijke Friese Front en Oestergronden), terwijl de laagste dichtheden gevonden werden in het zuidelijke gebied. Het zandige zuidelijke deel (Bruine Bank en Breeveertien) en de Klaver Bank bevatten hogere dichtheden van de meiobenthische taxa Copepoda (interstitieel), Gastrotricha, Turbellaria, Annelida, Hydrozoa en Tardigrada. Het Friese Front werd gekenmerkt door hoge Nematoda dichtheden, hogere aantallen aan Halacarida en Kinorhyncha, terwijl de Oestergronden gekenmerkt werden door hoge dichtheden aan Nematoda en gravende Copepoda alsmede door de aanwezigheid van Kinorhyncha, Rotifera en Priapulida.
- De meiobenthische diversiteit ($H_{\text{ill}, \infty}$) was het hoogst in het zandige zuidelijke gebied en in de Klaverbank, terwijl ze het laagst was in westelijke Friese Front en in de Oestergronden.
- * - In de 85 onderzochte macrobenthos monsters werden 160 soorten gedetermineerd. De taxa Polychaeta en Crustacea vertegenwoordigden een meerderheid van het aantal gevonden macrobenthos soorten.
- De macrobenthos dichtheid varieerde tussen 257 en 4126 ind./m² (gemiddeld: 1308 ind./m²). De biomassa van het macrobenthos vertoonde waarden tussen 1 en 136 g AFDW/m² (gemiddeld: 18 g AFDW/m²). In het noordelijke gebied was de dichtheid hoger dan in het gebied in de Zuidelijke Bocht. De hoogste biomassa waarden werden in het noorden en in het zuidwestelijke deel aangetroffen. Circa 40 % van de totale dichtheid werd gevormd door de Polychaeta. De Echinodermata vormden 52 % van de totale biomassa.

- Het aantal macrobenthos soorten per boxcore ($Hill_0$) varieerde tussen 11 en 33, waarbij de laagste waarden gevonden zijn in het zuidoostelijke deel van het onderzoeksgebied. Stations met een lage diversiteit ($Hill_1$ tot $Hill_{\infty}$) zijn in het noorden gevonden, waar een paar soorten dominant zijn.
- * De waargenomen samenstelling van het zoöbenthos vertoonde hoge correlaties met de sediment samenstelling vooral met de mediane korrelgrootte, het slibgehalte en, in mindere mate, met de waterdiepte. Gebieden met een hoog slibgehalte en een lage mediane korrelgrootte, werden bevolkt door macrobenthische gemeenschappen die gekarakteriseerd worden door een hoge totale biomassa, totale dichtheid en soortenrijkdom. Daarentegen vertoonde de meiobenthische diversiteit lage waarden in deze gebieden, als gevolg van de afwezigheid van veel meiobenthische taxa.

Om meer informatie te verkrijgen over de ruimtelijke verspreiding van de meio- en macrobenthos gemeenschappen, zijn de stations geklassificeerd met behulp van de TWINSPAN-clustering methode gebaseerd op taxon-niveau voor de meiobenthos dichtheid, respectievelijk op soortsniveau voor de macrobenthos dichtheid. Meio- en macrobenthos clusters vertoonden in het algemeen dezelfde ruimtelijke verdeling; twee clusters in het noordelijke slibrijke gebied en twee clusters in het medium tot grofzandige gebied in het zuiden. De grenzen van de twee zuidelijke meio- en macrobenthos clusters kwamen niet geheel overeen, waarschijnlijk als gevolg van verschillen in de kritische korrelgrootte grenzen voor beide groepen. Het meiobenthos lijkt gevoeliger te zijn in het korrelgrootte bereik tussen 250 en 300 μm , terwijl het macrobenthos reageert op verschillen tussen 200 en 250 μm .

De clusters van het meio- en macrobenthos kunnen als volgt worden omschreven:

- * In het noorden vertoonde cluster 1 van het macrobenthos de hoogste waarden van de totale dichtheid en biomassa, terwijl in het zuiden (cluster 4) de laagste waarden van de dichtheid en biomassa gevonden zijn. In het centrale gebied tussen deze twee clusters was het macrobenthos gemiddeld wat betreft de dichtheid en biomassa (clusters 2 & 3).
- * Het meiobenthos was in het noorden in het westelijke Friese Front en de Oestergronden in relatief hoge dichtheden aanwezig, maar de diversiteit was laag (cluster 1). Rond dit gebied, ter hoogte van de 30 en 40 m dieptelijnen, ligt een gebied met lagere dichtheden maar nog steeds arm aan taxa (cluster 2). In het zuiden waren de dichtheden in het algemeen lager, maar werd er een gemeenschap met een hogere diversiteit aangetroffen (cluster 3), vooral in de gebieden met het grofste zand (cluster 4).

De verspreiding van de meio- en macrobenthische gemeenschappen in de Zuidelijk Bocht alsook in het zuidelijke deel van de Oestergronden en het Friese Front leken relatief stabiel te zijn gebleven in de periode tussen 1988 en 1993 (MILZON-BENTHOS I+II).

De clusters van het meio- en macrobenthos van het totale MILZON-BENTHOS 1988-1993 gebied werden in de volgende gebieden gevonden:

- * Het macrobenthos vertoonde twee gemeenschappen in het slibrijke gebied van de Oestergronden en het Friese Front (clusters I & II), een kustgemeenschap (cluster III) en een gemeenschap van in grofzandige gebieden gelegen stations (cluster IV).
- * Het meiobenthos gaf een gemeenschap (cluster I) te zien in het slibrijke fijnzandige gebied (Oester Gronden, Friese Front en in de buurt van riviermondingen). Cluster II is geconcentreerd in het kustgebied, in het fijn-medium zandige gebied tussen 53° N en de 30 m dieptelijn en in de slibarmere gebieden in het noorden. Cluster III wordt voornamelijk aangetroffen in de Zuidelijk Bocht, terwijl cluster IV in de grofste sedimenten voorkomt.

1 PREFACE

The MILZON-BENTHOS II (MILieuZONering/Environmental Zonation) project (1991-1993) was carried out under commission of the North Sea Directorate of the Ministry of Transport, Public Works and Water Management (RWS-DNZ) as a continuation of the MILZON-BENTHOS I project (1988-1989) (Zevenboom & Leewis, 1987; Zevenboom, 1993). The aim of the project is to map in great detail the spatial distribution of the zoobenthos in the Dutch part of the North Sea and, on basis of this information, to classify homogenous areas. Within the framework of these projects the Dutch coastal area has been surveyed in the years 1988-89 (Groenewold & van Scheppingen, 1989; 1990; van Scheppingen & Groenewold, 1990), the area north of the Dutch Wadden islands, the Oyster Ground and Frisian Front area in 1991 (Holtmann & Groenewold, 1992). Together with the investigation of 1992 and 1993, which covered the area west of the Frisian Front, the Broad Fourteens and the Brown Bank (this report), almost the total Dutch Continental Shelf (DCS) of the North Sea was studied in the course of the MILZON-BENTHOS I+II projects.

The achievement of a deeper understanding of the functioning of the North Sea ecosystem is necessary in order to carry out an appropriate and efficient management of the North Sea. This knowledge is of importance not only in the assessment of the effect of human activities (e.g. discharges into the sea, offshore mining, fishery etc.), but also in relation to the drafting/framing of international conventions/treaties. Benthic ecosystem information has already played a dominant role in determining the Environmental Zone on the Dutch Continental Shelf, which is part of the Water System Management Plan North Sea 1991-1995. Furthermore a "stable and diverse benthic ecosystem" is one of the policy aims in this plan.

The objectives of the project in 1992/93 are to:

1. Make an inventory of the macrozoobenthos (species composition, density and biomass) and meiozoobenthos (taxon composition and density) in the area west of the Frisian Front, in the Broad Fourteens and the Brown Bank.
2. Determine the heterogeneity in the spatial distribution of the meio- and macrozoobenthos, by quantifying the natural variability in meio- and macrobenthos assemblages.
3. Distinguish several assemblages by statistical analysis.
4. Compare and integrate the present data with overlapping stations of the long term monitoring program of Rijkswaterstaat (EXP*BMN project) as well as with the survey area that has already been mapped (MILZON-BENTHOS I project).

2 INTRODUCTION

2.1 General abiotic features

The Dutch Continental Shelf (DCS) of the North Sea is a rather diverse area in terms of water depth and sediment composition. In the southern part (the Southern Bight), the depth ranges from 0 m (coastline) to 30 m. Beyond the 30 m isobath at approximately 53° N, a narrow zone with increasing depths separates this shallow southern part from the deeper northern area (Oyster Ground, Botney Ground)(Figs. 1 and 2). Further north, the depth gradually increases to about 50 m, with a deeper spot (50-60 m) in the Oyster Ground (Puzzle Hole) as well as in the Botney Ground (Botney Cut). Between these two Grounds a relatively shallow (30-40 m) area is found, running north into the sand and gravel area of the Cleaver Bank. The northern border of the Oyster Ground is formed by the shallow and sandy Dogger Bank (10-30 m).

Strong tidal currents occur in offshore parts in the Southern Bight, where the bottom sediment can generally be characterized as medium to coarse sands with small mud contents. Near to the coast this is sometimes mixed with mud, while in some areas the sediment consists of gravel. In the area north of the 30 m isobath, called the Frisian Front area (de Gee et al., 1991), tidal current velocities decrease sharply, resulting in the deposition of mud and detritus on the bottom. This muddy area is rich in organic material and in macrobenthic fauna (Creutzberg et al., 1984). North of the 40 m isobath, in the fine-grained silty area of the Oyster Ground, tidal currents are weak. In summertime, under favourable conditions, stratification of the water column can occur in these areas with low current velocities. The upper warmer layer is separated from the colder bottom layer by a thermocline, restricting exchange between the two layers. In the Southern Bight the strong tidal currents prevent such a stratification.

In the Dutch part of the Southern and Central North Sea, several water masses are formed as a result of the tidal movement and the influence of predominantly westerly winds. Depending on its origin, these water masses are: continental coastal water (mixture of Atlantic Ocean- and continental river water), English Channel water (unmixed Atlantic Ocean water) and Central North Sea water (unmixed Atlantic Ocean water) (Lee, 1970) (Fig. 3). Fronts mark the boundaries between the water masses, which can differ in biological as well as chemical characteristics (Pingree & Griffiths, 1978).

2.2 General biotic features

Since the introduction of the concept of large-scale marine assemblages at the beginning of this century, interest in the composition of the zoobenthos in the North Sea and its relationship with the abiotic environment (depth, sediment composition, salinity etc.) has increased. In 1986 the Benthos Ecology Working Group of ICES studied the distribution of marine bottom assemblages in the North Sea. This synoptic mapping of the infauna of the North Sea therefore should act as a basis for all further comparisons on natural and anthropogenic changes in the benthos since it is the first synoptic overview.

More detailed investigations have been carried out during the subsequent MILZON-BENTHOS projects I+II, which showed results that were generally in line with those of the synoptic mapping of ICES. Due to the finer sampling grid, the MILZON-BENTHOS I project (1988-1989) revealed the presence of three benthic zones parallel to the Dutch coast, both of the meio- as well as of the macrobenthos (Groenewold & van Scheppingen, 1989; 1990; van Scheppingen & Groenewold, 1990). The results of the MILZON-BENTHOS II project (1991) in the area north of the Dutch Wadden islands, around the Frisian Front and the Oyster Ground, were comparable with those of

the ICES survey (Holtmann & Groenewold, 1992). In the sandy area of 1991 stations were found to contain a relatively low macrobenthos biomass and density and low meiobenthos density, whereas the muddy northern area was characterized by a rich zoobenthos assemblage in terms of macrobenthos density and biomass and also of meiobenthos density (Holtmann & Groenewold, 1992).

2.3 Acknowledgements

The project was carried out under the direction of the Ministry of Transport, Public Works and Water Management (RWS, North Sea Directorate). The authors would like to thank the captain and the crew of the R/V Holland for their ambitious help during the fieldwork.

Thanks are also due to:

- J. Asjes, W. Zevenboom (RWS-North Sea Directorate), S.A. de Jong (RWS-Division Zeeland), M. van Arkel, G. Duineveld (NIOZ-Texel), C. Heip, J. Craeymeersch, L. Goud (CEMO-Yerseke) and A. Smaal (RWS-RIKZ-Middelburg) for their efforts having the project running as smoothly as possible.
- R. Huys (Natural History Museum-London) and G. Duineveld (NIOZ-Texel) for the use of their data.
- G. Duineveld (NIOZ-Texel), J. Asjes, W. Zevenboom, S. van Laar, H. van Bostelen (RWS-North Sea Directorate), J. Craeymeersch (CEMO-Yerseke), K. Essink (RWS-RIKZ-Haren) and A. Smaal (RWS-RIKZ-Middelburg) for critically reading the manuscript.
- H. Hobbelink (NIOZ-Texel) for adaptation of the cover picture.
- W. Schreurs, F. Geyp, M. Steendijk and G. den Hartog (RIKZ-Middelburg) for analysing sediment samples.
- J. Buijs, J. Polderman, R. Braat (CEMO-Yerseke), A. Kok (NIOZ-Texel), W. Bil, and J. van Dienst (RIKZ-Middelburg) for assistance with and improvement of computer programs.
- J. Kamphuis, C. Bijleveld, M. de Back, J. van Bree, D. Heijkoop, B. de Jong, M. Kints and T. van Schie (RWS-North Sea Directorate) for assistance during the fieldwork and deliverance of the samples and parameters measured during the field survey.

3 MATERIAL AND METHODS

3.1 Sampling

In the period 27 April to 22 May 1992 and 24 May to 4 June 1993, sediment samples were collected at a total of 82 stations with the R/V Holland (Groenewold & Holtmann, 1992; 1993). Additional data from another 3 stations (N 50, N 70 and META 1) located within the present study area, which were sampled on 31 March 1992 during the EXP*BMN program (Duineveld & Belgers, 1993; Huys & De Smet, 1993), have been incorporated in the dataset (Fig. 2). These samples have been treated in the same way as the ones collected for this survey. All stations had at least a distance of 2 nautical miles from operational or abandoned drilling locations (drilled since 1985). The existence of peat-banks in the Brown Bank area of 1992 (compartments 6, 7 and 8), caused the extra limited condition that 3 shallow (stations 1-3) and 3 deeper (stations 4-6) stations had to be selected.

Prior to sampling, the investigated area was divided into 19 homogeneous compartments in terms of sediment composition and water depth, under the surmise that there exists a relationship between the zoobenthos and the environment (Duineveld et al., 1990; Künitzer et al., 1992)(Figs. 1 and 2). Depending on the surface area of the compartment, 3-6 stations were selected according to the "stratified random sampling"-method.

The following parameters were collected for investigation:

- * macrozoobenthos (species, density, biomass)
- * meiocoobenthos (taxa, density)
- * median grain size and mud content
- * POC (Particulate Organic Carbon) content of the sediment
- * chl.-a (chlorophyll-a) content of the sediment
- * visual oxidized layer of the sediment

At each station two Reineck boxcore samples were taken with a surface of 0.068 m² (at some stations it was 0.078 m²). The required minimal depth of the boxcore samples was set at 15 cm. One boxcore was subsampled for meiobenthos (1 core of minimal 15 cm depth and ϕ of 3.4 cm), sediment grain size (1 core of 10 cm depth and ϕ of 3.4 cm), and for POC and chlorophyll-a content (2 cores each of 3 cm depth and ϕ of 11 mm). These subsample cores were also used for visual observations of the "oxidized layer". The other boxcore sample was used for macrobenthos analysis. For this purpose, the sample was washed on board through a sieve with round holes of 1 mm diameter. The macrobenthic residue as well as the entire meiobenthos subsample were preserved in a borax-buffered solution of 4-6 % formaldehyde in seawater. The subsamples collected for sediment grain size analysis, mud content (fraction < 63 μ m), chlorophyll-a and POC analyses were immediately frozen at -20°C and analyzed within 6 months after sampling.

3.2 Treatment of samples

The subsamples collected for sediment grain size, mud content (fraction < 63 μ m), chlorophyll-a and POC contents were analyzed at the analytical department of RWS-National Institute for Coastal and Marine Management (RIKZ) in Middelburg. Procedures were according to the standard regulations for meio- and macrobenthos in the Southern North Sea (Holtmann, 1990a & b).

In the laboratory, rose-bengal was added to the meio- and macrobenthic samples to facilitate sorting. The meiofauna was separated by means of decantation through a 38 μm sieve and sorted at taxon level (densities/10 cm²).

Prior to sorting, the macrobenthos samples were fractionated over a set of 4 nested stainless steel sieves with 0.5 mm as the smallest mesh size. The roughly equal particle size in the four residues improved the efficiency of sorting organisms from the debris. Density (individuals/m²) and biomass (g Ash-Free Dry Weight/m²) were, if possible, recorded at the species level and otherwise at higher taxonomic level. For some groups of macrofauna (Bivalvia, Echinoidea) the sizes of individuals were noted and, if clearly discernible, the juvenile species were recorded, but not further used for data analysis. The biomass (g AFDW/m²) of the macrobenthos was determined by drying the organisms at 60°C for at least 60 hours and subsequently incinerating them at 520°C for 4 hours. Some representatives of the Crustacea (amphipods, isopods and cumaceans), were assigned an average individual AFDW (0.3-0.5 mg) which was derived from data obtained during the MILZON-BENTHOS I+II projects (Duineveld and Holtmann, pers. comm.).

More information about sampling and sample treatment on board the ship can be found in the cruise reports of MILZON-BENTHOS II 1992 and 1993 (Groenewold & Holtmann, 1992; 1993).

3.3 Sediment classification

The grain size distribution of the sand fraction in the sediment was analyzed by means of a Malvern (RIKZ-Middelburg). The contribution of the mud fraction (% particles < 63 μm) was determined separately. The sediment types were classified on basis of the median grain size as follows:

Characterization of the sedimenttype according to
the median grain size (after Gullentops et al., 1977).

< 175 μm	Very fine sand
175 - 250 μm	Fine sand
250 - 300 μm	Fine-medium sand
300 - 350 μm	Medium-coarse sand
> 350 μm	Coarse sand

3.4 Statistical analysis

3.4.1 Diversity

The zoobenthic diversity in the investigated area is expressed by Hill's diversity numbers (Hill, 1973). The relationship between the commonly used measures of diversity and Hill's numbers is as follows:

- Hill₀ = N₀ = the number of species/taxa
Hill₁ = N₁ = exp(H') (H' = Shannon-Wiener index; Shannon & Weaver, 1949)
Hill₂ = N₂ = 1/SI (SI = Simpson index of dominance; Simpson, 1949).
Hill _{∞} = N _{∞} = the reciprocal of the proportional abundance of the most common species/taxa


3.4.2 Classification of assemblages

3.4.2.1 TWINSPAN

All stations were classified into groups by means of the FORTRAN program TWINSPAN (Hill, 1979) using the taxon, respectively species abundance of the meio- and macrozoobenthos. Within the TWINSPAN method, the stations are ordered along an ordination axis according to their faunal characteristics. This axis is split in two parts and each set of stations is separately classified again and so on. This leads to a dichotomous division of the total set of stations up to a preset level.

The resulting clusters of the classification of the 1992/93 area were numbered from 1 to 4 and of the 1988-1993 area from I to IV. For meio- and macrobenthos identical symbols were used for comparable TWINSPAN clusters.

To illustrate the differences between the clusters with respect to selected biotic and abiotic parameters, notched box and whisker plots were drawn using SYSTAT 5.0 (SYSTAT inc. IL, USA). Because these plots show the medians and their simultaneous confidence intervals (the notches) of the different groups, they allow a direct assessment of significant differences (at a 95 % confidence level) between median values of the groups (McGill et al., 1978).



3.4.2.2 Multi-dimensional scaling (MDS)

Another method to classify several stations, is non-metric multi-dimensional scaling (MDS) ordination (Kruskal & Wish, 1978). Prior to MDS ordination all data were $\log(x+1)$ transformed, in order to prevent numerically dominant species/taxa to swamp the other data, which is usually desirable for density and biomass data (Clarke & Green, 1988). The faunal (dis)similarity between the stations was calculated with the Bray-Curtis similarity index, whereas dissimilarity in terms of abiotic data was based on Euclidean distances. Separate data of meio- and macrobenthos as well as (a combination) of environmental parameters were used for similarity analyses. The resulting (dis)similarity matrices between the stations were utilized as input for the MDS ordination. In the MDS plot, the dissimilarity ranking is preserved as Euclidean distances. For simplicity only the first two axes of the multidimensional ordination were plotted.

The stress factor expresses the accuracy with which the dissimilarity of the stations is presented by the two dimensional MDS plot; increasing stress denotes less accuracy.

By replacing the stations symbols in the MDS plot by symbols of variable size representing values of abiotic parameters, the relation between the stations ordination and the selected parameter is visualized.

3.4.3 Correlations

In order to gain more insight into the factors influencing the zoobenthos distribution, linear correlations (Pearson's r) between abiotic and biotic parameters were tested for significance ($P < 0.05$). The Spearman rank correlation was used to calculate the correlation between the benthic assemblages and the abiotic parameters (Clarke & Ainsworth, 1993).

Prior to the calculations of Pearson or Spearman rank correlation, all data were $\log(x+1)$ -transformed.

4 RESULTS

4.1 The investigated area

4.1.1 Environmental parameters

Depths in the investigated area of the MILZON-BENTHOS II 1992/93 project were recorded, but not corrected for differences due to the tidal cycle (mean tidal amplitude ± 1.5 m). Fig. 4 shows these results in combination with the official hydrographic soundings (isobaths). In the southern part depths ranged generally between 20 and 30 m, with the exception of the Brown Bank area (compartments 6-8, Table 1 and Fig. 4). Further north, in the compartments 11-13, the depth increased from circa 30 m to 40 m in the vicinity of the western Frisian Front. In the northern part of compartments 18 and 19 (Oyster Ground) depth increased from 40 m to 50 m. In the north-west corner (compartment 17) the part of the Botney Cut with depths over 50 m lies just within the study area.

The median grain size of the sediment decreased from south to north (Fig. 5). South of the 30 m isobath, the sediment consists of locally coarse to predominantly medium/fine sand with a low mud content (0-10 %), except for some deeper stations in the Brown Bank area. Between the 30 and 40 m isobath the mud percentage ($< 63 \mu\text{m}$) of the sediment increases rapidly with maximum levels surpassing 20 % at Botney Cut (Fig. 6). The Oyster Ground sediment consists of fine sand with a mud percentage of 5-10 %. In the extreme north-eastern part of the investigated area the Cleaver Bank is just visible, with coarse sand and low mud contents.

High concentrations of chlorophyll-a and POC in the upper 3 cm of the sediment were found near the Frisian Front area, Botney Cut and in some parts of the Brown Bank (Figs. 7 and 8, Table 2). However, measurements took place at various times over a 2 month period (Groenewold & Holtmann, 1993). Lowest POC contents were invariably found in the sandy sediment south of the 30 m isobath and in the Cleaver Bank area (Fig. 8).

Fig. 9. shows the depth of the "oxidized layer" based on visual observations of the subsample cores. In the western Frisian Front area and in the Oyster Ground area, only a few centimetres of aerated sediment were observed. In the sandy sediment of the Southern Bight a deeper visual "oxygen layer" was found. Values of the oxygen penetration in the Frisian Front area, measured with a microelectrode, are much lower (0.45 cm in August and 1.85 cm in February) (Lohse et al., 1993), than the visual "oxidized layer" observed during the MILZON-BENTHOS II project 1992/93.

4.1.2 Correlations among the abiotic parameters

The matrix of the correlation coefficients (Pearson's r) between the abiotic parameters is given in Table 3. A very high correlation was found between the sediment parameters mud and median grain size. In deep areas the sediment consisted of a low median grain size with a relatively high mud content. A significant correlation was also found between the sediment composition and the amount of Particulate Organic Carbon (POC) in the sediment. The chlorophyll-a content of the sediment only correlated with the mud percentage and the content of POC.

4.2 Meiobenthos

4.2.1 Taxon composition and diversity

The present survey revealed the presence of 14 permanent meiobenthic taxa over the whole study area (Table 4). Species belonging to the permanent taxa are considered to spent their juvenile and adult life stage in the meiobenthic size range ($38 \mu - \pm 1 \text{ mm}$). An additional number of 13 temporary (e.g. juvenile macrobenthos) taxa were found. Appendix 1 presents the densities of both types. Meiofaunal densities of the EXP*BMN stations 2-6 (N 50), 7-7 (N 70) and 3-6 (META 1) were given by Huys & de Smet (1993)(Appendix 3.1). Only the permanent taxa have been used for further analyses.

The Nematoda turned out to be the dominant group, accounting on average for 82 % of the total meiobenthos density (min: 44.6 %, max: 99.6 %)(Table 5). Copepoda ranked second in contribution to the total abundance (mean: 9 %). The soft-bodied taxa Gastrotricha and Turbellaria together formed 5 % of the total meiobenthic abundance. All other taxa represented about 2.5 % of the total.

Means and ranges of Hill's diversity numbers for taxa composition of the meiofauna were as follows:

$$\begin{aligned} \text{Hill}_0 &= N_0 = 3.0 - 11.0 \text{ (mean} = 7.0\text{)} \\ \text{Hill}_1 &= N_1 = 1.0 - 3.9 \text{ (mean} = 1.8\text{)} \\ \text{Hill}_2 &= N_2 = 1.0 - 3.0 \text{ (mean} = 1.4\text{)} \\ \text{Hill}_{\infty} &= N_{\infty} = 1.0 - 2.2 \text{ (mean} = 1.2\text{)} \end{aligned}$$

The distributions of Hill's numbers ($\text{Hill}_{0-\infty}$)(Figs. 10-13) show that taxon diversity was highest in the coarse and medium/fine sandy sediments in the southern part of the study area and in the Cleaver Bank. Diversity was low(est) in the western Frisian Front and in the Oyster Ground. Considering the whole area, Hill's diversity numbers decreased rapidly with increasing order due to the high Nematoda numbers. This is especially valid in the western Frisian Front and Oyster Ground where most individuals belonged to the Nematoda.

4.2.2 Density

Total and mean density and respective taxon densities are given in Table 6. High meiobenthic densities were found in the western Frisian Front (east side $> 3500 \text{ ind./10 cm}^2$) and Oyster Ground area with a maximum of 6815 ind./ 10 cm^2 (Fig. 14). Densities below 1000 ind./ 10 cm^2 were only recorded in the southern sandy area. On average, 1783 ind./ 10 cm^2 were present in the study area (median: 1472 ind./ 10 cm^2).

The distribution of the Nematoda (Fig. 15) showed a pattern similar to the total meiobenthos density in the northern area (western Frisian Front and Oyster Ground), but the higher total densities in the southern area (Brown Bank and Broad Fourteens) were only partly due to the Nematoda densities.

Copepoda comprised bigger mud dwelling and burrowing forms (well suited for muddy substrates) and small interstitial specimen (adapted to the interstices in coarser sand). Representatives of the latter category were most abundant in the coarser sandy southern area and the Cleaver Bank area (Fig. 16). 85 % of all Copepoda in the 1992/93 area belonged to the interstitial Copepoda and were mainly found in the sandy sediment, whereas the bigger burrowing Copepoda predominated in the muddy western Frisian Front and Oyster Ground (Fig. 17 a/b).

The ratio between Nematoda and Copepoda (N/C ratio) was relatively high in the western Frisian Front and Oyster Ground and also in the area directly south of the 30 m isobath. The ratio was rather low in the medium-coarse sandy parts (range: 1 to 832, mean 67) (Fig. 18).

Of the other taxa, Gastrotricha (generally preferring a fine-medium ($\sim 250 \mu\text{m}$) sand type) hardly occurred in the western Frisian Front and Oyster Ground, but they were abundant in the sandy Southern Bight (Fig. 19). Turbellaria were present in relatively high densities in the sandy southern area, but also occurred in the muddy parts (Fig. 20).

The taxa of the Annelida, Archiannelida and Oligochaeta, were scarce and primarily present in sandy substrates, whereas Polychaeta were present everywhere with no clear pattern in area distribution (Figs. 21, 22 and 23).

Hydrozoa and Tardigrada were mainly found in the sandy areas (Figs. 24 and 25). Ostracoda were found both in the sandy part as well as in the muddy part (Fig. 26). Halacarida were scarce and found mainly in the sand (Fig. 27).

Kinorhyncha (Fig. 28) and Priapulida (1 specimen at 2 stations in the western Frisian Front area) were restricted to the muddy parts. Rotifera were found twice: 14 ind./10 cm² on station 13-2 and 1 ind./10 cm² on station 16-3.

4.2.3 Correlations among the meiobenthos

Within the meiobenthos several taxa were positively correlated to each other (Table 7). Nematoda, Ostracoda and Kinorhyncha appeared in higher densities in areas with high total meiobenthic densities. On the contrary, interstitial Copepoda and Gastrotricha were present in low densities in these areas. Interstitial Copepoda, Gastrotricha, Turbellaria, Archiannelida, Hydrozoa and Tardigrada were positively correlated to each other, especially in areas with coarser sand and a low mud content. The diversity was generally high in these latter areas, while most taxa were found in relatively high densities, except Nematoda and Kinorhyncha. Therefore the total density was often rather low. Halacarida and Rotifera did not correlate with any parameter and were thus excluded from the table.

4.2.4 Meiobenthic assemblages

4.2.4.1 TWINSPLAN ordination

TWINSPLAN-classification, based on taxon abundance (Copepoda split into burrowing and interstitial groups), of the combined stations of the 1992/93 area resulted in the formation of 4 clusters (Table 8; Figs. 29 and 30). Table 8 represents the mean values of the meiobenthic and abiotic parameters per meiobenthos TWINSPLAN cluster. Clusters 1 (western Frisian Front and Oyster Ground) & 2 (around the 30 and 40 m isobaths) were situated in the deeper northern muddy area, whereas the clusters 3 & 4 were found in the sandy Southern Bight, with cluster 4 in the coarsest parts. Characteristic for the 'north-south division' were burrowing Copepoda (>20 ind./10 cm²) (clusters 1 & 2), whereas Gastrotricha (>10 ind./10 cm²), interstitial Copepoda and Tardigrada were characteristic for the clusters 3 & 4. The box and whisker plots of Fig. 31 illustrate the differences between the meiobenthic clusters (Table 8).

The EXP*BMN stations of 1992, N 70 and META 1 were grouped in the same clusters as the stations surrounding them (cluster 3), whereas N 50 was grouped in the nearby cluster 1.

- cluster 1 = The western Frisian Front and part the Oyster Ground, consisting of muddy very fine sand with relatively high contents of mud and POC, and intermediate chlorophyll-a contents. Total meiobenthic as well as Nematoda densities were very high, resulting in low diversities. Copepoda mainly consisted of burrowing forms. Kinorhyncha were characteristic for this area, whereas Archiannelida, Hydrozoa, Tardigrada and Halacarida were absent. Gastrotricha were scarce.
 - cluster 2 = Mainly the northern part around the 30 and 40 m isobaths, with sediments built up of fine sand with still relatively high mud-, chlorophyll-a and POC content. Densities were intermediate. All taxa, except the Nematoda, were generally low in density or even absent, consequently resulting in low diversities.
 - cluster 3 = Most stations belong to the fine-medium grained sandy southern part with low mud- and POC contents. One station is found in the western Frisian Front. Chlorophyll-a contents were intermediate. Diversities were higher than those of the first two clusters. The interstitial taxa appeared in higher densities than in the north. Half of the investigated stations were grouped in this cluster.
 - cluster 4 = A group of stations with low mud-, chlorophyll-a- and POC contents situated in the medium-coarse sandy areas in the south, in the eastern corner of compartments 5 and 13 (coarse relict Pleistocene sand), and in the Cleaver Bank. Diversities were relatively high. All taxa, except Nematoda, Kinorhyncha and Priapulida, were present in relatively high densities. Copepoda were found to be mainly interstitial.

4.2.4.2 MDS ordination

Fig. 32 shows the MDS plot of the dissimilarity between the stations in the investigated area. Stations are marked with the symbols of the TWINSPAN clusters. The stress value was 0.10, giving a plot which is unlikely to mislead (Warwick & Clarke, 1993). The four station groups distinguished by TWINSPAN can also be distinguished in the MDS plot, although they do not form 4 highly segregated groups. The first division in TWINSPAN (clusters 1/2 <-> 3/4) occurs at the first dimension axis of the MDS plot.

4.3 Macrobenthos

4.3.1 Species composition and diversity

From 85 boxcore samples a total of 160 species were identified (Table 9 and 10). The majority of the species belonged to the Polychaeta (59 species) and Crustacea (57 species). From the taxon of the Mollusca 26 species and of the Echinodermata 8 species were found. The remaining 10 taxonomic groups were not identified to species level. The most common species, i.e. those being present at more than 25 % of the stations, are listed in Table 11.

The number of species per boxcore sample ($Hill_0$) varied between 11 and 33 (Table 12, Fig. 33), with the lowest values occurring in the south-eastern part of the studied area (compartments 1, 2 and 3), at stations with medium to coarse sand. The diversity numbers $Hill_1$ to $Hill_\infty$ are given in Figs. 34-36. Stations with relatively low values for these diversity numbers are situated in the north, where a few species numerically dominate the macrobenthic assemblage. Examples of these species are the polychaete Magelona papillicornis, the bivalve Mysella bidentata, the decapod Callianassa subterranea and the ophiuroid Amphiura filiformis.

4.3.2 Density and biomass

The data on the macrobenthos abundance and biomass are given in Appendix 2. All data of the EXP*BMN project 1992 can be found in Appendix 3.2. The total density of macrofauna ranged from 257 to 4126 ind./m² (mean: 1308 ind./m²) (Fig. 37). In the northern part the total density was consistently higher than in the area of the Southern Bight, where the macrobenthic abundance seemed to be more heterogeneous. Total macrofaunal biomass ranged from 1 to 136 g AFDW/m² with a mean value of 18 g AFDW/m². Fig. 38 shows that higher biomass values were generally found in the north and in the south-western part of the area.

The distribution of the density and biomass of the four major taxa Polychaeta, Crustacea, Mollusca and Echinodermata are given in Figs. 39-42 and in Figs. 43-46 respectively. The spatial distribution of the most common species are shown in Figs. 47-74 (alphabetized consecutively per taxon).

The Polychaeta were found at all stations, with higher densities south to the 30 m isobath. From the total macrobenthic density, 40 % belongs to the Polychaeta (Table 12). The taxon of the Crustacea showed higher abundances in the centre of the study area caused by high densities of the amphipods Bathyporeia elegans and Bathyporeia guilliamsoniana (Figs. 63 and 64). The distribution of the Crustacea biomass showed higher values in the north because in this area many specimens of Callianassa subterranea (Fig. 65) were found. In the northern part of the area high densities of the taxa Mollusca and Echinodermata were found, corresponding with high abundances of the bivalve Mysella bidentata (Fig. 60) and the brittle-star Amphiura filiformis (Fig. 70). Frequently high values of the Mollusca and Echinodermata biomass were caused by a single adult specimen of a large-sized species such as Echinocardium cordatum (Fig. 71) and the bivalves Ensis siliqua or Lutraria lutraria, or by the dominance of a single species, as Amphiura filiformis (Fig. 70). About 52 % of the total biomass in the whole area was formed by the taxon Echinodermata (Table 12).

4.3.3 Correlations among the macrobenthos

Between the macrobenthic parameters many positive correlations were found (Table 13). The total macrofaunal density showed close relationships with almost all of the measured

macrobenthic factors. In areas with a high macrobenthic abundance, stations with a high species number ($Hill_0$) and high biomass were found (cf. TWINSPAN clustering in section 4.3.4.1). The Polychaeta (density and biomass), the Crustacea density as well as the biomass of the Mollusca and Echinodermata showed no correlation with the other macrobenthic parameters. This could be expected considering that these parameters of the taxa showed no clear distribution pattern (cf. Figs. 39-42 and 43-46). Negative correlations were found between the diversity numbers ($Hill_{1-\infty}$) and total macrobenthos density.

4.3.4 Macrobenthic assemblages

4.3.4.1 TWINSPAN ordination

Using TWINSPAN classification based on species abundance, the stations of the 1992/93 area were grouped into four clusters (Fig. 75). The characteristic 'indicator species' of the TWINSPAN clustering are given in Fig. 76. The geographical position of the clusters in Fig. 75 shows a high degree of spatial segregation between clusters. First the whole area can be split into a northern and a southern part along the 30 m depth line. Two northern clusters (1 & 2) with 23 stations around the area of the Frisian Front and the Oyster Ground in the north and clusters 3 & 4 with 59 stations in the area of the Broad Fourteens and the Brown Bank in the south. At the second division level of TWINSPAN, in the north the deeper stations with high mud content of cluster 1 were separated from cluster 2 and in the south the sandy area of cluster 4 from cluster 3. In Fig. 77, notched box and whisker plots illustrate the differences between the four macrobenthic clusters in terms of abiotic and biotic parameters as given in Table 14.

The two stations META 1 and N 50 of the EXP*BMN project 1992, incorporated in the dataset of MILZON-BENTHOS II 1992/93, belong to the same cluster as the area around the stations (cluster 4), but in compartment 7 station N 70 was grouped into a northern cluster (cluster 3).

Since the distribution of the clusters formed four well-defined adjacent areas with different sediment types (Table 14 and Fig. 77), the four clusters can be described as follows:

- ☒ cluster 1 = The deep area of the Oyster Ground and the western part of the Frisian Front area, with sediments composed of muddy very fine sand with a relatively high content of mud and POC and low chlorophyll-a content. Cluster 1 has the highest average density and biomass and the lowest diversity numbers ($Hill_{1..}$), which are presented in Table 14.
 - ☐ cluster 2 = The muddy very fine sand area, around the 30 m isobath, with an intermediate mud, chlorophyll-a and POC content. Cluster 2 has the highest diversity numbers ($Hill_{1..}$).
 - ◻ cluster 3 = The shallow, fine sand area south of the Frisian Front, with the highest content of chlorophyll-a and an intermediate POC content. Cluster 3 has the highest average species richness.
 - ◊ cluster 4 = The shallow fine-medium sand area in the south covers almost the whole area of the Broad Fourteens and the Brown Bank. The sandy sediment has low contents of mud, chlorophyll-a and POC. Cluster 4 has the lowest average values for macrobenthic density, biomass and for species richness.

The most abundant species of each cluster, i.e. those found at more than 60 % of the stations, are listed in Table 15 (Figs. 47-74). These macrobenthic species can be used to describe the main assemblages of the MILZON-BENTHOS II area 1992/93. In the northern part with muddy fine sand, an Amphiura filiformis-, Callianassa subterranea- and Mysella bidentata- assemblage (cluster 1) and a Callianassa subterranea-, Cultellus pellucidus-, Lumbrineris latreilli-, Magelona papillicornis- and Nephtys hombergii- assemblage was found (cluster 2). In the southern part, the area with fine sand showed a Chaetozone setosa-, Magelona papillicornis- and Spiophanes bombyx- assemblage (cluster 3) and the medium to coarse sand area was characterized by the macrobenthic species Nephtys cirrosa and Scoloplos armiger (cluster 4).

4.3.4.2 MDS ordination

The macrobenthos was also classified by MDS. Fig. 78 shows the first two axes of the MDS plot, based on the dissimilarity between the stations, using the macrobenthos abundance (stress value: 0.16). The locations are marked with the same symbols as chosen for the clusters of the TWINSPAN classification. The MDS plot produces in general the same grouping as found by TWINSPAN. On the first axis of the MDS plot clusters 1 & 2 were separated from clusters 3 & 4 and the second axis of MDS corresponded with the second division of the TWINSPAN classification. This indicates a high degree of dissimilarity between the four TWINSPAN assemblages.

4.4 Comparison with ICES and MILZON-BENTHOS I+II

4.4.1 Meiobenthos

The most extensive study into the spatial distribution of the meiobenthos in the North Sea, the synoptic mapping of the zoobenthos of the Benthos Ecology Working Group of ICES, was carried out in 1986. 14 stations of this program were situated in the MILZON-BENTHOS II 1992/93 area. Therefore these ICES stations were incorporated into the present dataset, in order to assess possible changes in the meiobenthic assemblages after TWINSPAN analysis based on taxon densities (Fig. 79). Comparison with Fig. 29 (without ICES stations) shows that these classifications match fairly well in the Southern Bight. The ICES stations belong to the same clusters as adjacent stations without changing the original pattern. The only exception is formed by ICES station 31, with a totally different meiobenthic composition (high Nematoda density, low diversity) than the area around this station. In the northern muddy part, the ICES stations effected cluster 1, in the area of the Oyster Ground and western Frisian Front, to extend southward (30 m isobath), westward (Botney Ground) and northward (40 m isobath). The ICES stations in the north caused the enlargement of the group of deeper, muddy stations with fine grained sediment (cluster 1) characterized by relatively high Nematoda densities and, except Copepoda and Kinorhyncha, low densities or absences of the remaining taxa.

In order to compare the present results on meiobenthos with those from the previous MILZON surveys (MILZON-BENTHOS I+II) in other parts of the Dutch Continental Shelf (Groenewold & van Scheppingen, 1989; 1990; Holtmann & Groenewold, 1992), all data were combined for a TWINSPAN analysis (Figs. 80 and 81). The classification of the present study area fits well into the pattern found on basis of the combined results from MILZON-BENTHOS I+II (Holtmann & Groenewold, 1992). (These new TWINSPAN cluster groupings are marked with Roman numerals). Cluster I stations were all situated in muddy fine grained sediment, with high total and Nematoda densities (Fig. 82 a/b) and low densities of most other taxa (Fig. 82 c-e) resulting in low diversities ($Hill_{1..}$). They were found in the near coastal area in the vicinity of the river outflows (North Sea Channel and Rhine), in the Oyster Ground and in the Frisian Front, which extends westward into the western Frisian Front area. The cluster II assemblage is concentrated in the coastal area, in the fine-medium coarse sand area between 53° N and the 30 m isobath, and also in the less muddy Oyster Ground and western Frisian Front parts. The main part of the investigated Southern Bight west of the Dutch coastal area (including the Brown Bank and Broad Fourteens) and south of the 30 m isobath belongs to the relatively diverse ($Hill_{0..}$) cluster III (medium-coarse sand, low mud contents, lowest Nematoda densities and highest Gastrotricha densities). The most diverse assemblage (cluster IV), is found in the coarsest sediments, in the south and in the relict Pleistocene sands around Texel/Vlieland and Borkum. In overlapping compartments most of the stations from two different sampling years belonged to the same clusters as the adjacent stations, indicating that in the period between 1988 and 1992 (Southern Bight) as well as between 1991 and 1993 (Frisian Front/Oyster Ground), the composition and densities of the meiobenthic taxa have remained relatively stable.

4.4.2 Macrofauna

13 stations of the ICES investigation (1986), located in the area of MILZON-BENTHOS II (1992/93), have been included in the data of the present investigation (Fig. 2), looking for possible changes of the macrobenthic assemblages. The results of ICES station 51 (only two samples) were not used for comparison. Fig. 83 shows the TWINSPAN classification of the combined data. The 5 replicate samples, taken at one ICES location, are found in nearly all cases in the same cluster, therefore only one symbol for the ICES stations was printed. The classification shows in general the same pattern as the clustering without the ICES locations (Fig. 75). Almost all stations of the ICES survey were grouped into the same cluster as the adjacent MILZON stations. Only some differences are found. In the north, the station ICES 42 belongs to the southern cluster (cluster 4; cf. Holtmann & Groenewold, 1992) and in the south the station of the EXP*BMN 1992 (N 70) was found in a cluster of the northern area (MILZON cluster 2).

In order to compare the MILZON-BENTHOS II (1992/93) project with the investigations of MILZON-BENTHOS from 1988 to 1991, the TWINSPAN classification was used for the whole area. The compartments in the east of the present study area (compartments 1 to 5, 13, 16 and 19) overlapped with the area of earlier MILZON-BENTHOS investigations. The 513 stations of MILZON-BENTHOS I+II were also split into 4 clusters (I-IV) (Figs. 84 and 85). Taking the characteristic species into account, the following assemblages can be distinguished:

In the muddy sand area of the Oyster Ground and the Frisian Front two clusters can be described as Callianassa subterranea, Amphiura filiformis, Pholoe minuta and Harpinia antennaria assemblages (clusters I & II). A coastal assemblage, characterized by the species Magelona papillicornis, Tellina fabula, Nephtys cirrosa, Lanice conchilega and Urothoe poseidonis (cluster III), an assemblage of coarse sand area in the Southern Bight (cluster IV).

The stations in the overlapping compartments were found in the same clusters (cluster I in the north and clusters III & IV in the south). This shows that the macrobenthic assemblages remained unchanged in the period between 1988 and 1992 in the Southern Bight (compartments 1 to 5 and 13) and between 1991 and 1993 in the area around the Oyster Ground and the Frisian Front (compartments 16 and 19).

4.5 Correlations between the zoobenthos and the environment

4.5.1 Separate zoobenthic components and abiotic parameters

The sediment characteristics and selected benthic parameters were correlated, using Pearson's correlation coefficient (Tables 7 and 13). This gives an idea about the relations between the separate meio- and macrofauna components and the environment. The chlorophyll-a content showed no correlations with any of the zoobenthos factors.

The total meiobenthic density correlated with most of the measured abiotic parameters (Table 7). The median grain size showed a significant positive correlation with several taxa (especially with interstitial Copepoda, Gastrotricha, Turbellaria, Hydrozoa and Tardigrada), whereas the same taxa were negatively correlated with areas enriched with mud and POC. The meiobenthic diversity was highest in shallow coarse sand areas and lowest in deeper muddy areas. Nematoda and Kinorhyncha on the contrary, were concentrated in deeper areas with high mud- and POC contents.

Table 13 presents the correlations between the macrobenthos and the abiotic parameters. Stations with high total density, high density of Mollusca and Echinodermata, high total biomass, high Crustacea biomass and high species richness ($Hill_0$) were concentrated in deeper areas with a high mud content and low median grain size. The amount of organic material (POC) correlated positively with some macrobenthic parameters (total density, density of Mollusca, total biomass and biomasses of Polychaeta and Crustacea). No correlation was found between any of the sediment parameters and the diversity numbers $Hill_{1-\infty}$ of the macrobenthos.

4.5.2 Total benthic fauna and abiotic parameters

The relation between abiotic parameters and differences in the total fauna was determined in two ways.

First, by correlating the TWINSPLAN scores on the first axis, with selected abiotic parameters of the stations. The scores or positions of the stations on the ordination axis, as shown in Fig. 86 and 87, were compared with the environmental parameters of those stations, using the Spearman rank correlation method. The median grain size showed the highest correlation with the benthic fauna, whereas the mud content also seemed to influence the benthic assemblages (Table 16).

Second, by correlating the dissimilarities between the stations on basis of the fauna with the dissimilarities of their abiotic properties (Clarke & Ainsworth, 1993). Table 17 shows the Spearman rank correlation between biotic and abiotic station similarities (cf. section 3.3.3). Both meio- and macrobenthos seemed to correlate best with the median grain size of the sediment. A combination of median grain size and mud content, with an important role of the water depth, seem to provide the best environment to explain the (dis)similarity between the stations in the MILZON-BENTHOS II 1992/93 area. These correlations are also illustrated in the 2-dimensional plots of Fig. 88 a/b.

4.6 Relations between the meio- and macrobenthos

Table 18 shows the correlations between the separate meio- and macrobenthos components. The taxon of the Nematoda, which presented 80 % of the meiobenthos, was positively correlated with the total macrobenthic density and biomass. The total meiobenthos density and the N/C ratio also showed a positive relationship with the macrobenthos, as was also observed earlier during the MILZON-BENTHOS I project (van Scheppingen & Groenewold, 1990). Besides the Nematoda and Kinorhyncha (only found in the muddy northern part) all the other meiobenthic taxa appear to be negatively correlated with the macrobenthos. The diversity of the macro- and meiobenthos showed a negative correlation; the areas with high meiobenthic diversity ($Hill_{1-\infty}$), showed low densities, biomasses and numbers of species ($Hill_0$) of the macrobenthos. This was also found in the MILZON-BENTHOS II project 1991 (Holtmann & Groenewold, 1992).

The distribution of the TWINSPLAN clusters for the meio- and macrobenthos showed in general the same pattern (Figs. 29 and 75): two northern assemblages in the muddy sand area and two southern assemblages in the medium-coarser sand area. However in the south of the investigated area the meiobenthos clusters seemed to be more heterogenous (Fig. 29). This can partly be explained by the fact that the zoobenthos are strongly tied to specific sediment types (see high correlations) which are heterogeneously distributed in the southern part of the study area (cf. Figs. 5 and 6). The meiobenthos taxa responsible for the TWINSPLAN division in clusters 3 & 4 (cf. Fig. 30), for example interstitial Copepoda, Archiannelida, Oligochaeta and Hydrozoa, were mainly found in medium-coarser sand (median grain size $> 300 \mu\text{m}$). The macrobenthos seemed to be more sensitive to differences in the median grain size between 200 and 250 μm . Some dominant macrobenthic species like *Tellina fabula* and *Magelona papillicornis* were only found in an area with a median grain size between 200 and 250 μm (macrobenthos cluster 3) and found to be 'indicator species' of TWINSPLAN classification. The MDS-ordination (Figs. 32 and 78) also emphasizes the general similarity in cluster pattern.

5 DISCUSSION

5.1 Meiobenthos

Due to their small size and the use of wide meshed sieves the study of the meiobenthos in the North Sea was started much later (Smidt, 1951) than the study of the macrobenthos (Petersen, 1914). Although considerable effort has been put in meiobenthos studies in the North Sea (Heip et al., 1990), few studies have been carried out on the spatial distribution of the meiobenthos on the Dutch Continental Shelf. Smol et al. (1989) studied the meiobenthos in TiO_2 waste dumping-area just south of the MILZON-BENTHOS 1992/93 study area. The sediment composition coincided with those of the stations of compartments 1 and 6. The composition of the meiobenthos was quite comparable among these two areas: relatively high densities of the main taxa Nematoda, Copepoda, Gastrotricha and Turbellaria and also higher densities of the remaining taxa.

In the synoptic mapping of the infauna of the ICES stations in 1986 Huys et al. (1992) determined the meiobenthos on taxon level and additionally determined the Copepoda on species level. TWINSPLAN analysis of the ICES data, based on copepod species composition, showed five main assemblages in the North Sea. Two of them were situated in the present study area. A clearly distinctive Southern Bight assemblage was found south of the 30 m isobath, containing relatively high numbers of harpacticoid Copepoda with a majority of interstitial species. In this area nematode numbers were relatively low. In the western Frisian Front/Botney Ground/Oyster Ground area an assemblage was found that formed a transition between the (Dutch) coastal and deepwater (north of the Dogger Bank) copepod associations. The other meiobenthic taxa were present especially in the medium to coarse sand of the Southern Bight assemblage. These patterns coincide with the results found in the present study. The Southern Bight, south of the 30 m isobath, together with station 17-1 on the coarse sandy Cleaver Bank showed the highest diversity (Hill...), highest densities of (interstitial) Copepoda, Gastrotricha, Turbellaria, Hydrozoa and Tardigrada in comparison with the deeper stations in the silty western Frisian Front, Oyster Ground and Botney Ground. In general the meiobenthic community seems to remain relatively stable in composition, although larger year to year fluctuations in densities and even in copepod composition occasionally occur at some stations, as was found in the EXP*BMN studies of 1991 and 1992 (Huys & De Smet, 1993).

In the Southern Bight (present study) the meiobenthic assemblages seem to be more patchy distributed than the macrobenthos, probably due to the differences in sensitivity to interstitial space between the sand particles. The range between 250 and 300 μm is therefore critical for the meiobenthos and acts a selection criterium for the presence or absence of the interstitial meiobenthic species/taxa.

The N/C ratio, a still discussible tool for detecting the presence of certain forms of pollution (Raffaelli, 1987; Coull et al., 1981; Lambshead, 1984; Huys et al., 1992), can only be used to give an indication in areas with the same sediment composition. They will therefore be only valuable for comparison with future spatial distribution investigations.

The different sampling method used in shallower/deeper peat-banks in the Brown Bank in order to get more detailed information about the distribution of the meiofauna in the area did not show any clear differences in zoobenthos composition or density.

5.2 Macrobenthos

5.2.1 Relations between the macrobenthos and the environment

Many research programs were carried out to study the distribution of macrobenthic assemblages in the North Sea and found high correlations between the species distribution and the sediment composition. Salzwedel et al. (1985) compared for the German Bight the types of sediment with the distribution of the macrofauna. In 1984, Creutzberg et al. published the results of the macrobenthic fauna in the Frisian Front area and found a rich benthic macrofauna, corresponding with the zonation of the sediment. Künitzer et al. (1992) reports that most of the macrobenthic species are restricted to a special sediment and therefore these species are limited in their distribution.

The bottom of the Southern North Sea shows a major gradient in sediment composition: the Southern Bight contains medium-coarse grained sediment, which in northern direction is replaced by finer-grained sand. All these sandy bottoms contain little mud, except for some local patches. The macrobenthos composition found in this study follows a similar gradient. Besides the abiotic factors, such as hydrographic circumstances, depth and sediment structure, interactions with other species and food availability play an important role in the stability of the species distribution in the North Sea (Künitzer et al., 1992; Heip & Craeymeersch, 1994).

The area studied during the MILZON-BENTHOS II project 1992/93, presented in this report, covered a part of the North Sea with different types of sediment. Also in this particular area we found a clear relation between the distribution of macrobenthic species and the sediment type. The muddy area north of the 30 m depth line contained other assemblages than south of the 30 m isobath, corresponding with the different sediment composition. The sandy station on the Cleaver Bank (17-1) in the north-western part of the research area belonged to the cluster of the southern part (MILZON cluster 4). The macrobenthic composition found at this station was typical for assemblages in sandy sediments, a striking example of the close relationship between sediment grain size and macrobenthos. In the same compartment one station was situated in the Botney Cut (17-3), with a depth of 53 m. This station had a macrobenthic assemblage with high abundance and biomass values (MILZON cluster 1). The Oyster Ground, in the north-eastern part of the study area (compartments 18 and 19), belonged to a macrobenthic cluster (MILZON cluster 1) being rich in terms of species richness, biomass and density, as also found earlier in the MILZON-BENTHOS project 1991 (Holtmann & Groenewold, 1992). In the south only one deeper station (7-4) of the Brown Bank was found in another cluster (MILZON cluster 3) than the area around this station, because of the high mud content of this station. The area of the Brown Bank in the compartments 6, 7 and 8 contained an assemblage typical for a coarse sand assemblage (MILZON cluster 4). In order to get more detailed information about the distribution of the macrobenthic fauna in the area of the peat-banks at the Brown Bank other sampling methods have to be used.

5.2.2 Stability of the macrobenthos

Comparing the present data with those of earlier investigations we can discern some light stability in time of the assemblages. In April 1986 the Benthos Ecology Working Group of ICES recorded a synoptic mapping of the infauna of the North Sea (de Wilde & Duineveld, 1988; Künitzer et al., 1992). In the area of MILZON-BENTHOS II project 1992/93 they found in the south a shallow coarse-sand assemblage (MILZON cluster 4) with Scoloplos armiger and Nephthys cirrosa as two of the most abundant species and a shallow fine-sand assemblage characterized by

Bathyporeia elegans, Spiophanes bombyx, Magelona papillicornis, Tellina fabula and Urothoe poseidonis (MILZON cluster 3). In the north they observed shallow muddy sand assemblages described by some of the species of the MILZON clusters 1 & 2, namely Amphiura filiformis, Mysella bidentata, Magelona papillicornis and Pholoe minuta (Duineveld et al., 1990). All macrobenthic species mentioned here have been found to be dominant in the respective MILZON clusters 1 to 4 (Table 13). The stability of the macrobenthos assemblages is also demonstrated by the results of the TWINSPLAN classification of the MILZON-BENTHOS survey 1992/93 including the ICES data of 1986, where ICES locations grouped into the same assemblages as the adjacent MILZON stations (cf. 4.4.2).

Several investigations of the macrobenthos published the biomass distribution of the Southern North Sea (Creutzberg, 1984; Cadée, 1984; de Wilde et al., 1984), prior to the ICES survey. The biomass in the western area of the Frisian Front showed in the present study average values between 36 g AFDW/m² of cluster 1 and 15 g AFDW/m² of cluster 2 (Table 12). Similar high values of biomass were also found by de Wilde et al. (1984) (21.4 g AFDW/m²). Creutzberg et al. (1984) reported average biomass values of 13.2 g AFDW/m² at the Frisian Front. In the same year Cadée (1984) found at the Front values of 10.6 g AFDW/m² and south to the 30 m depth line an average biomass of 5.7 g AFDW/m², much lower than found during the MILZON project 1992/93 (MILZON cluster 4 in the south = 10.5 g AFDW/m²). The low biomass values in the latter two studies may have been due to the use of Van Veen grabs instead of boxcorers. By using the Van Veen grab deeper burrowing animals, which are common in the Oyster Ground, are easily missed.

6 CONCLUSIONS

Meiobenthos:

- * High total densities, mainly Nematoda, were found in the muddy areas, especially the western Frisian Front area.
- * Nematoda dominated the meiofauna (82 %).
- * The medium-coarse sandy Southern Bight (south of the 53°30' latitude) showed the most diverse (Hill_{1-∞}) meiobenthic assemblage.
- * The results of the MILZON-BENTHOS II project 1992/93 in general showed a similar distribution of taxon density and composition as found during the ICES investigation in 1986.
- * Four assemblages could be distinguished in the meiobenthos:
In the deeper muddy northern parts two assemblages were found: in the western Frisian Front and Oyster Ground the meiobenthos was represented in relatively high densities but with low diversities (cluster 1). This area is surrounded by an area around the 30 and 40 m isobaths with lower densities but still impoverished in taxa (cluster 2).
In the sandy Southern Bight total densities were generally lower, but a more diverse assemblage could be found (cluster 3), especially in the coarsest sandy parts (cluster 4).
- * Between 1988 and 1993 (MILZON-BENTHOS I+II) the meiobenthic density and composition in the Southern Bight as well as the Central North Sea (southern part of the Oyster Ground and Frisian Front) seemed to be relatively stable on taxon level.
- * In the period between 1988 and 1993 (MILZON-BENTHOS I+II) the meiobenthic assemblages of the Southern Bight and the southern part of the Oyster Ground seemed to be relatively stable. The following assemblages were found:
 - an assemblage (cluster I) of stations in the muddy fine grained sand (Oyster Ground, Frisian Front and near river outflows).
 - cluster II is concentrated in the coastal area, in the fine-medium coarse sand area between 53° N and the 30 m isobath and in the less muddy parts of the northern area.
 - cluster III is mainly found in the Southern Bight,
 - cluster IV is found in the coarsest sediments.

Macrobenthos:

- * The total density (ind./m²), biomass (g AFDW/m²) and species richness of the macrobenthos showed higher values in the northern part of the area.
- * The majority of the total density belongs to the taxon of the Polychaeta (40 %) and 52 % of the total biomass is represented by the Echinodermata.
- * The macrobenthos can be divided into four assemblages:
North of the 30 m depth line
 - an Amphiura filiformis-, Callianassa subterranea and Mysella bidentata- assemblage (cluster 1) and
 - a Callianassa subterranea-, Cultellus pellucidus-, Lumbrineris latreilli-, Magelona papillicornis- and Nephtys hombergii- assemblage (cluster 2).
- South of the 30 m depth line
 - a Chaetozone setosa-, Magelona papillicornis- and Spiophanes bombyx- assemblage (cluster 3) and
 - a Nephtys cirrosa- and Scoloplos armiger- assemblage (cluster 4).

- * The results of the MILZON-BENTHOS II project 1992/93 shows the same distribution of species composition as found during the ICES investigation in 1986.
- * In the period between 1988 and 1993 (MILZON-BENTHOS I+II) the macrobenthic assemblages of the Southern Bight and the southern part of the Oyster Ground seemed to be stable. The following assemblages were found:
 - two assemblages of the muddy sand area from the Oyster Ground and the Frisian Front, characterized by Callianassa subterranea, Amphiura filiformis, Pholoe minuta and Harpinia antennaria (clusters I & II),
 - a coastal assemblage characterized by Magelona papillicornis, Tellina fabula, Nephtys cirrosa, Lanice conchilega and Urothoe poseidonis (cluster III) and
 - an assemblage of the coarse sand area in the Southern Bight (cluster IV).

Meio-/macrobenthos:

- * The distribution of the TWINSPAN clusters showed for the meio- and macrobenthos in general the same pattern, although in the southern part of the investigated area the macrobenthos clusters seemed to be more confined.
- * The distribution of the meio- and macrobenthic assemblages showed a high relationship with the sediment composition, especially with median grain size, mud content and, to a less extent, with the water depth.
- * The meiobenthos seems to be more sensitive to the range of median grain sizes between 250 and 300 µm, whereas the macrobenthos reacts to changes between 200 and 250 µm. This explained the different cluster distribution of meio- and macrobenthos in the southern part of the studied area.
- * The total density of the meiobenthos and the Nematoda density showed positive correlations with the macrobenthos.
- * The sampling method used in shallower/deeper peat-banks in the Brown Bank did not show any clear differences in zoobenthos composition or density. Therefore other sampling methods have to be used to get more detailed information about the distribution of the meio- and macrobenthic fauna in the area of the peat-banks at the Brown Bank.

7 RECOMMENDATIONS

The MILZON-BENTHOS I and II projects have resulted in a large dataset of the meio- and macrobenthos distribution on the Dutch Continental Shelf. In order to present all this information the results should be published in a benthic atlas. Such an atlas could serve two goals: provide information necessary for management policy in the North Sea as well as act as an information source for further investigation.

The aim of the MILZON-BENTHOS I+II research was to map the spatial distribution of the meio- and macrobenthos. More information about temporal changes of the zoobenthos in the Southern North Sea is to be expected from the long term monitoring program EXP*BMN, which started in spring 1990 (Holtmann et al., 1990c) and which has developed into a cooperative effort of the National Institute for Coastal and Marine Management (RIKZ), the North Sea Directorate of Rijkswaterstaat (RWS-DNZ), the Netherlands Institute for Sea Research (NIOZ) and the Netherlands Institute of Ecology/Centre for Estuarine and Coastal Ecology (NIOO/CEMO) (Duineveld, 1992; Duineveld & Belgers, 1993; Huys & De Smet, 1992; 1993). To check if the chosen stations of the monitoring program are still representative for the specific area, the mapping of the zoobenthos, as was carried out in the MILZON-BENTHOS I+II projects, needs to be repeated every 10-15 years. Because of the useful results presented in this report, it would be better to alter the present strategy of sampling fixed stations in the EXP*BMN project to a strategy of stratified random sampling.

The results from the MILZON-BENTHOS surveys were used in the decision of the establishment of the Environmental Zone in the Dutch area of the North Sea. It should be meaningful if the surrounding North Sea countries be brought together to join into a zoobenthos mapping program in the context of the closer cooperation and concern of the common North Sea environment.

8 REFERENCES

- CADÉE, G.C. 1984. Macrofauna and macrobenthic remains on the Oyster Ground, North Sea. -Neth. J. Sea Res. 18(1/2): 160-178.
- CLARKE, K.R. & R.H. Green, 1988. Statistical design and analysis for a 'biological effects' study. -Mar. Ecol. Prog. Ser. 46: 213-226.
- CLARKE, K.R. & M. Ainsworth, 1993. A method of linking multivariate community structure to environmental variables. -Mar. Ecol. Prog. Ser. 92: 205-219.
- COULL, B.C., G.R.F. Hicks & J.B.J. Wells, 1981. Nematode/copepod ratios for monitoring pollution: a rebuttal. -Mar. Poll. Bull. 12: 378-381.
- CREUTZBERG, F., P. Wapenaar, G. Duineveld & N. Lopez Lopez, 1984. Distribution and density of benthic fauna in the southern North Sea in relation to bottom characteristics and hydrographic conditions. -Rapp. P.-v. Réun. Cons. perm. int. Explor. Mer 183: 101-110.
- DUINEVELD, G.C.A., P.A.W.J. de Wilde & A. Kok, 1990. A synopsis of the macrobenthic assemblages and benthic ETS activity in the Dutch sector of the North Sea. -Neth. J. Sea Res. 26(1): 125-138.
- DUINEVELD, G.C.A., (1992). The macrobenthic fauna in the Dutch sector of the North Sea in 1991. -NIOZ-RAPPORT 1992-6: 17pp.
- DUINEVELD, G.C.A. & J. Belgers, (1993). The macrobenthic fauna in the Dutch sector of the North Sea in 1992. -NIOZ-RAPPORT 1993-11: 38pp.
- GEE, A. de, M.A. Baars & H.W. van der Veer, 1991. De ecologie van het Friese Front. Waarnemingen aan een biologisch-rijke zone in de Noordzee, gelegen tussen de Zuidelijke Bocht en de Oestergronden. -NIOZ-RAPPORT 1991-2: 96 pp.
- GROENEWOLD, A. & Y.C.M. van Scheppingen, 1989. De ruimtelijke verspreiding van het benthos in de zuidelijke Noordzee, voorjaar 1988. -MILZON-BENTHOS rapport nr. 89-05 (89-010). Rijkswaterstaat.
- GROENEWOLD, A. & Y.C.M. van Scheppingen, 1990. De ruimtelijke verspreiding van het benthos in de zuidelijke Noordzee, 1989. -MILZON-BENTHOS rapport nr. 90-01 (90-001). Rijkswaterstaat.
- GROENEWOLD, A. & S.E. Holtmann, 1992. Cruise report of the MILZON-BENTHOS survey in the North Sea: the Broad Fourteens and Brown Bank. Spring 1992. -MILZON-BENTHOS report. Rijkswaterstaat, North Sea Directorate.
- GROENEWOLD, A. & S.E. Holtmann, 1993. Cruise report of the MILZON-BENTHOS survey in the North Sea: the Western Frisian Front. Spring 1993. -MILZON-BENTHOS report. Rijkswaterstaat, North Sea Directorate.
- GULLENTOPS, F., M. Moens, A. Ringel & R. Sengier, 1977. Geologische kenmerken van de suspensies en de sedimenten. -In: J. Nihoul & F. Gullentops (eds): Mathematisch Model Noordzee. Vol4. Sedimentologie.
- HEIP, C.H.R., R. Huys, M. Vincx, A. Vanreusel, N. Smol, R. Herman & P.M.J. Herman, 1990. Composition, distribution, biomass and production of North Sea meiofauna. -Neth. J. Sea Res. 26: 333-342.
- HEIP, C. & J.A. Craeymeersch, 1994. Benthic community structures in the North Sea. -Helgoländer wiss. Meeresunters. (in press).
- HILL, M.O. 1973. Diversity and evenness: a unifying notation and its consequences. -Ecology, 54: 427-432.

- HILL, M.O., 1979. TWINSPLAN. A FORTRAN program for arranging Multivariate data in an Ordered Two-Way Table by Classification of the Individuals and Attributes. -Ecology and Systematics, Cornell University. Ithaca, New York 14850.
- HOLTMANN, S.E., 1990a. Biomonitoring van macro- en meiofauna in de zuidelijke Noordzee I. Macrofauna (april 1990). -MILZON-BENTHOS rapport nr. 90-02.1. Rijkswaterstaat.
- HOLTMANN, S.E., 1990b. Biomonitoring van macro- en meiofauna in de zuidelijke Noordzee. II. Meiofauna (april 1990). -MILZON-BENTHOS rapport nr. 90-02.2. Rijkswaterstaat.
- HOLTMANN, S.E., Y.C.M. van Scheppingen & A. Groenewold, 1990c. Biomonitoring van het zoobenthos in de zuidelijke Noordzee, voorjaar 1990. Deelrapport monitoring MILZON-BENTHOS. -MILZON-BENTHOS rapport 90. Rijkswaterstaat.
- HOLTMANN, S.E. & A. Groenewold, 1992. Distribution of the zoobenthos on the Dutch Continental Shelf: the Oyster Ground, Frisian Front, Vlieland Ground and Terschelling Bank (1991) -NIOZ-RAPPORT 1992-8/NIOO-CEMO rapporten en Verslagen 1992-6: 129pp.
- HUYS, R., C.H.R. Heip, P.M.J. Herman & K. Soetaert, 1992. The meiobenthos of the North Sea: density, biomass trends and distribution of copepod communities. - ICES J. mar. Sci., 49: 23-44.
- HUYS, R. & G. De Smet, 1992. Het meiobenthos van het Nederlands Continentaal Plat. Voorjaar 1991. -Rapportage in het kader van het Biologisch Monitoring Project: 57 pp.
- HUYS, R. & G. De Smet, 1993. Het meiobenthos van het Nederlands Continentaal Plat. Voorjaar 1992. -Rapportage in het kader van het Biologisch Monitoring Project: ??pp
- KRUSKAL, J.B. & M. Wish, 1978. Multidimensional scaling. -Sage Publications, Beverley Hills, California.
- KÜNTZER, A., G.C.A Duineveld, D. Basford, J.M. Dewarumez, J. Dörjes, A. Elefteriou, C. Heip, P.M.J. Herman, P. Kingston, U. Niermann, H. Rumohr & P.A.J.W. de Wilde 1992. The benthic infauna of the North Sea: Species distribution and assemblages. -ICES J. mar. Sci. 49: 127-143.
- LAMBSHEAD, P.J.D., 1984. The nematode/copepod ratio. Some anomalous results from the Firth of Clyde. -Mar. Poll. Bull. 15: 256-259.
- LEE, A., 1970. The currents and watermasses of the North Sea. -Oceanogr. mar. Biol. ann. Rev. 8: 33-71.
- LOHSE, L., J.F.P. Malschaert, C.P. Slomp, W. Helder & W.v. Raaphorst, 1993. Nitrogen cycling in North Sea sediments: interactions of denitrification and nitrification in offshore and coastal areas. -Mar. Ecol. Prog. Ser. 101: 283-296.
- MCGILL, R., J.W. Tukey & W.A. Larsen 1978. Variations of box plots. -The American Statistics 32: 12-16.
- PINGREE, R.D. & D.H. Griffiths, 1978. Tidal fronts on the shelf sea around the British Isles. -J. geophys. Res. 83 (C9): 4615-4622.
- RAFFAELLI, D.G., 1987. The behaviour of the nematode/copepod ratio in organic pollution studies. -Mar. Environm. Res. 23: 135-153.
- SALZWEDEL, H., E. Rachor & D. Gerdes, 1985. Benthic macrofauna communities in the German Bight. -Veröff. Inst. Meeresforsch. Bremerh. 20: 199-267.

- SCHEPPINGEN, Y.C.M. van & A. Groenewold, 1990. De ruimtelijke verspreiding van het benthos in de zuidelijke Noordzee. De Nederlandse kustzone 1988-1989. - MILZON-BENTHOS rapport nr. 90-03 (90-003). Rijkswaterstaat.
- SHANNON, C.E. & W. Weaver, 1949. The mathematical theory of communication. -Univ. of Illinois Press, Urbana.
- SIMPSON, E.H., 1949. Measurements of diversity. -Nature, 163: 688-688.
- SMIDT, E.L.B., 1951. Animal production in the Danish Waddensea. -Medd. Danm. Fisk. Havundersøg., Bd II (6): 151 pp.
- SMOL, N., R. Huys & M. Vincx, 1989. Studie van het meiobenthos van een dumpingsgebied van titaandioxyde-afval in de Nederlandse kustwateren, periode 1986-1987. -Rapport RWS Directie Noordzee.
- WARWICK, R.M. & K.R. Clarke, 1993. Comparing the severity of disturbance: a meta-analysis of marine macrobenthic community data. -Mar. Ecol. Prog. Ser. 92: 221-231.
- WILDE, P.A.W.J. de, E.M. Berghuis & A. Kok, 1984. Structure and energy demand of the benthic community of the Oyster Ground, Central North Sea. -Neth. J. Sea Res. 18(1/2): 143-159.
- WILDE, P.A.W.J. de & G.C.A. Duineveld, 1988. Macrofauna van het Nederlands Continentale Plat, verzameld tijdens de ICES North Sea benthos survey april 1986. - NIOZ RAPPORT, APRIL 1988.
- ZEVENBOOM, W. & R. Leewis, 1987. Projectplan MILIEUZONERING. -MILZON notitie 87-05. Directie Noordzee, en Dienst Getijdewateren.
- ZEVENBOOM, W., 1993. Projectplan MILZON-2: 2^e fase project MILZON. Eindversie, d.d. 22 juni 1993. -MILZON notitie 93-02-c. Directie Noordzee, en Dienst Getijdewateren.

Tables

Tables :

I Abiotic parameters

1. (a/b) Position, depth and sediment composition of the MILZON-BENTHOS II area 1992/93.
2. (a/b) Chlorophyll-a ($\mu\text{g/g}$) and POC (% C) content of the upper 3 cm of the sediment and depth of the oxidized layer (cm) of the MILZON-BENTHOS II area 1992/93.
3. Correlations between the abiotic parameters found in the MILZON-BENTHOS II area 1992/93.

II Meiobenthos

4. Systematics of the meiobenthos.
5. Mean, median, minimum and maximum values of densities with the standard deviation of the meiobenthic taxa, expressed in percentages, in the MILZON-BENTHOS II 1992/93 area.
6. Mean, median, minimum and maximum values of the meiobenthos densities, together with the corresponding standard deviation and presence (in percentages) of the meiobenthic taxa in the MILZON-BENTHOS II area 1992/93.
7. Correlations between the meiobenthos and the abiotic parameters found in the MILZON-BENTHOS II area 1992/93.
8. Mean values of abiotic and meiobenthic parameters of each meiobenthic TWINSPAN cluster, as shown in Figs. 28 and 29, with their standard deviation.

III Macrobenthos

9. Systematics of the macrobenthos.
10. (a-c) List of macrobenthic species found in the MILZON-BENTHOS II area 1992/93. The mean values with standard deviation of density (ind./ m^2) and biomass (g AFDW/ m^2).
11. Presence (n, %) of the most abundant macrobenthos species, found at more than 25 % of the stations in the MILZON-BENTHOS II area 1992/93.
12. Presence (n, %), minimum, maximum, mean values with standard deviation and percentages of the mean of some taxonomic macrobenthic groups in the MILZON-BENTHOS II area 1992/93.
13. Correlations between the macrobenthos and the abiotic parameters found in the MILZON-BENTHOS II area 1992/93.
14. Mean values of abiotic and macrobenthic parameters of each macrobenthic TWINSPAN cluster, as shown in Figs. 45 and 46, with their standard deviation.
15. List of the most abundant macrobenthos species of each macrobenthic TWINSPAN cluster, as shown in Figs. 45 and 46. The mean values with standard deviation of density (ind./ m^2) and biomass (g AFDW/ m^2).

IV Meio- and macrobenthos

16. Spearman rank correlations between benthic (meio- and macrobenthos) TWINSPAN scores and abiotic parameters.
17. Spearman rank correlations between benthic (meio- and macrobenthos) and abiotic dissimilarity matrices.
18. Correlations between the meio- and macrobenthos found in the MILZON-BENTHOS II area 1992/93.

Table 1a. Position, depth and sediment composition of the MILZON-BENTHOS II area 1992/93.
 (*) = stations sampled during EXP*BMN 1992)

STATION NUMBER	NR	GEOGRAPHICAL POSITION	DEPTH (m)	MEDIAN (μm)	MUD (%)
1-1	1	52-15-10 N 003-24-42 E	32.8	325.0	1.83
1-2	2	52-19-36 N 003-23-48 E	33.5	387.7	0.71
1-3	3	52-21-32 N 003-46-06 E	26.7	350.6	0.61
1-4	4	52-25-19 N 003-33-12 E	31.7	317.8	3.12
1-5	5	52-21-27 N 003-33-25 E	33.6	327.4	1.84
2-1	6	52-31-14 N 003-36-25 E	32.5	287.7	1.43
2-2	7	52-29-24 N 003-43-00 E	30.9	284.0	2.25
2-3	8	52-32-18 N 003-52-00 E	29.7	266.8	1.64
2-4	9	52-35-50 N 003-58-00 E	26.1	283.2	1.42
2-5	10	52-40-00 N 003-40-30 E	31.2	273.5	1.54
* 2-6 N 50	11	52-28-51 N 003-47-12 E	30.4	/	/
3-1	12	52-41-42 N 003-55-09 E	25.5	264.1	1.71
3-2	13	52-46-58 N 003-58-20 E	29.7	247.3	1.77
3-3	14	52-49-12 N 003-51-08 E	27.4	264.7	1.54
3-4	15	52-53-33 N 003-44-17 E	28.0	247.3	3.04
3-5	16	52-57-48 N 003-48-34 E	35.1	388.4	1.88
* 3-6 META 1	17	52-57-53 N 003-55-00 E	27.3	247.4	1.68
4-1	18	53-02-36 N 004-20-00 E	30.4	228.7	2.07
4-2	19	53-02-28 N 003-50-25 E	31.4	246.5	3.04
4-3	20	53-07-38 N 003-52-25 E	28.4	221.7	2.47
4-4	21	53-07-38 N 004-02-00 E	28.2	213.0	2.48
4-5	22	53-05-48 N 004-21-30 E	31.2	245.3	2.57
5-1	23	53-12-42 N 003-50-03 E	26.1	208.0	2.08
5-2	24	53-19-20 N 003-47-48 E	27.6	201.9	4.23
5-3	25	53-21-38 N 004-06-20 E	28.3	198.5	8.08
5-4	26	53-14-40 N 004-23-40 E	29.8	338.4	2.14
5-5	27	53-21-28 N 004-33-20 E	27.4	345.7	1.61
6-1	28	52-20-25 N 003-11-25 E	32.4	293.2	4.09
6-2	29	52-27-43 N 003-12-12 E	34.0	283.6	1.44
6-3	30	52-26-20 N 003-06-04 E	34.5	290.1	2.37
6-4	31	52-24-08 N 003-14-30 E	41.2	343.6	2.87
6-5	32	52-27-19 N 003-23-21 E	34.7	292.7	2.75
6-6	33	52-28-36 N 003-09-13 E	40.5	299.3	2.68
7-1	34	52-35-59 N 003-19-04 E	19.3	284.3	1.71
7-2	35	52-36-11 N 003-13-15 E	32.6	295.2	2.94
7-3	36	52-41-02 N 003-13-10 E	33.7	269.1	3.25
7-4	37	52-39-04 N 003-17-00 E	41.2	245.9	12.61
7-5	38	52-32-40 N 003-21-35 E	37.4	327.1	1.71
7-6	39	52-38-42 N 003-37-42 E	33.3	318.3	2.47
* 7-7 N 70	40	52-34-11 N 003-31-53 E	31.3	294.1	1.12
8-1	41	52-46-06 N 003-18-26 E	25.4	271.7	2.95
8-2	42	52-53-53 N 003-17-20 E	33.4	277.6	2.03
8-3	43	52-57-21 N 003-34-47 E	28.1	248.9	2.93
8-4	44	52-47-30 N 003-22-13 E	41.3	278.1	6.94
8-5	45	52-50-12 N 003-17-18 E	34.6	271.8	2.54
8-6	46	52-56-31 N 003-22-06 E	34.2	290.9	2.35
9-1	47	53-02-58 N 003-11-36 E	29.5	254.4	3.77
9-2	48	53-03-55 N 003-23-30 E	27.9	258.2	2.63
9-3	49	53-08-43 N 003-23-00 E	26.9	245.1	3.74
9-4	50	53-07-48 N 003-31-42 E	29.5	238.3	2.35
9-5	51	53-05-30 N 003-43-00 E	28.1	227.7	3.37

Table 1b. Position, depth and sediment composition of the MILZON-BENTHOS II area 1992/93.
(* = stations sampled during EXP*BMN 1992)

STATION NUMBER	NR	GEOGRAPHICAL POSITION	DEPTH (m)	MEDIAN (μm)	MUD (%)
10-1	52	53-12-43 N 003-26-36 E	34.4	208.1	9.34
10-2	53	53-14-55 N 003-20-37 E	26.7	211.3	2.92
10-3	54	53-21-00 N 003-22-18 E	28.2	187.2	6.49
10-4	55	53-17-00 N 003-31-18 E	26.6	198.9	3.93
10-5	56	53-18-20 N 003-42-58 E	27.4	343.2	3.33
11-1	57	53-31-30 N 003-18-21 E	34.3	124.1	11.70
11-2	58	53-26-02 N 003-12-03 E	29.2	167.0	6.02
11-3	59	53-30-00 N 003-29-46 E	32.7	127.2	12.19
12-1	60	53-25-31 N 003-44-02 E	30.4	167.8	6.05
12-2	61	53-30-24 N 003-53-45 E	30.8	152.7	11.98
12-3	62	53-33-58 N 003-37-40 E	37.2	115.2	16.82
13-1	63	53-32-14 N 004-24-31 E	28.0	205.1	10.92
13-2	64	53-24-47 N 004-32-33 E	30.0	360.6	0.30
13-3	65	53-23-36 N 004-06-48 E	28.0	201.3	11.32
14-1	66	53-41-01 N 003-00-11 E	42.3	194.9	7.96
14-2	67	53-37-40 N 003-09-38 E	39.2	122.4	11.78
14-3	68	53-42-20 N 003-13-28 E	43.4	127.7	13.56
14-4	69	53-38-09 N 003-29-52 E	40.0	139.1	13.80
15-1	70	53-45-20 N 003-37-50 E	37.6	181.4	5.26
15-2	71	53-37-11 N 003-57-57 E	37.9	120.1	15.55
15-3	72	53-38-54 N 003-38-52 E	39.9	134.5	11.38
16-1	73	53-40-19 N 004-06-07 E	40.5	154.7	15.46
16-2	74	53-43-27 N 004-13-38 E	41.0	149.2	15.72
16-3	75	53-37-40 N 004-16-37 E	37.0	118.3	19.48
16-4	76	53-46-36 N 004-29-31 E	41.0	125.8	16.73
17-1	77	54-00-39 N 003-02-41 E	40.9	290.7	2.16
17-2	78	54-00-21 N 003-25-08 E	43.2	188.0	4.38
17-3	79	53-51-49 N 003-11-30 E	53.2	106.0	32.79
18-1	80	53-59-47 N 003-55-17 E	46.5	132.7	9.59
18-2	81	53-52-24 N 003-39-04 E	34.1	208.3	2.06
18-3	82	53-51-31 N 003-52-24 E	39.1	157.3	5.59
19-1	83	53-58-40 N 004-14-05 E	44.5	115.7	8.40
19-2	84	53-50-42 N 004-09-06 E	42.0	139.7	7.39
19-3	85	53-59-46 N 004-23-24 E	44.5	116.4	9.45

Table 2a. Chlorophyll-a content ($\mu\text{g/g}$) and POC content (% C) of the upper 3 cm of the sediment and depth of the oxidized layer (cm) of the MILZON-BENTHOS II area 1992/93.
 (/=values not available)
 {*} = stations sampled during EXP*BMN 1992)

STATION NUMBER	NUMBER	CHL.A ($\mu\text{g/g}$)	POC (% C)	O2 (cm)
1-1	1	0.83	0.0	10
1-2	2	0.19	0.0	22
1-3	3	0.00	0.2	10
1-4	4	0.37	0.2	22
1-5	5	0.42	0.2	23
2-1	6	0.12	0.1	21
2-2	7	0.22	0.1	23
2-3	8	0.00	0.2	19
2-4	9	0.00	0.1	23
2-5	10	0.13	0.1	23
* 2-6 N 50	11	0.11	0.0	/
3-1	12	0.12	0.2	20
3-2	13	0.16	0.2	20
3-3	14	0.15	0.1	17
3-4	15	1.79	0.2	20
3-5	16	2.09	0.3	3
* 3-6 META 1	17	0.51	0.0	/
4-1	18	0.00	0.2	20
4-2	19	1.15	0.1	21
4-3	20	1.08	0.3	20
4-4	21	1.44	0.2	5
4-5	22	0.00	0.1	17
5-1	23	1.84	0.0	5
5-2	24	1.54	0.0	3
5-3	25	14.15	0.6	2
5-4	26	0.00	0.0	30
5-5	27	0.00	0.0	17
6-1	28	0.27	0.0	20
6-2	29	0.56	0.0	17
6-3	30	5.93	0.0	21
6-4	31	0.68	0.0	20
6-5	32	0.26	0.0	/
6-6	33	4.17	0.0	20
7-1	34	0.30	0.0	25
7-2	35	0.59	0.0	10
7-3	36	1.01	0.0	10
7-4	37	0.81	0.1	13
7-5	38	1.00	0.0	20
7-6	39	0.36	0.0	22
* 7-7 N 70	40	0.52	0.0	/
8-1	41	0.51	0.0	22
8-2	42	0.22	0.0	22
8-3	43	2.79	0.0	/
8-4	44	4.95	0.0	4
8-5	45	0.23	0.0	20
8-6	46	0.21	0.0	21
9-1	47	0.11	0.0	21
9-2	48	0.27	0.0	19
9-3	49	10.20	0.0	21
9-4	50	0.38	0.0	18
9-5	51	1.77	0.5	20

Table 2b. Chlorophyll-a content ($\mu\text{g/g}$) and POC content (% C) of the upper 3 cm of the sediment and depth of the oxidized layer (cm) of the MILZON-BENTHOS II area 1992/93.
 {/ = values not available)
 {*} = stations sampled during EXP*BMN 1992)

STATION NUMBER	NUMBER	CHL.A ($\mu\text{g/g}$)	POC (% C)	O2 (cm)
10-1	52	0.78	0.2	3
10-2	53	0.24	0.0	/
10-3	54	0.25	0.0	21
10-4	55	0.53	0.0	8
10-5	56	0.75	0.0	5
11-1	57	1.58	0.3	6
11-2	58	0.95	0.1	4
11-3	59	1.48	0.4	5
12-1	57	0.25	0.1	9
12-2	58	2.58	0.4	7
12-3	59	1.57	0.5	4
13-1	60	33.33	1.5	14
13-2	61	0.28	0.2	20
13-3	62	20.89	0.8	10
14-1	63	1.58	0.3	3
14-2	64	0.86	0.4	7
14-3	65	0.83	0.5	1
14-4	66	1.34	0.5	5
15-1	67	0.17	0.2	6
15-2	68	1.32	0.5	4
15-3	69	0.97	0.4	6
16-1	70	1.01	0.7	3
16-2	71	1.29	0.5	3
16-3	72	1.31	0.6	3
16-4	73	0.45	0.4	10
17-1	74	0.13	0.1	30
17-2	75	0.39	0.2	5
17-3	76	3.31	1.0	1
18-1	77	0.57	0.3	7
18-2	78	0.00	0.2	9
18-3	79	0.26	0.3	6
19-1	80	0.28	0.3	6
19-2	81	0.35	0.4	7
19-3	82	0.32	0.3	5

Table 3. Correlations between the abiotic parameters, $\log(x+1)$ transformed, found in the MILZON-BENTHOS II area 1992/93.
 $(n=85, \alpha=0.001, r=0.347, 95\% \text{ reliability})$
*=significant

	abiotic parameters					
	Depth	Median	Mud	Chl.a	POC	Ox.l.
Depth	1.00					
Median gr.size	-0.47*	1.00				
Mud content	0.55*	-0.86*	1.00			
Chl.a content	0.01	-0.18	0.40*	1.00		
POC content	0.35*	-0.64*	0.70*	0.50*	1.00	
Oxidized layer	-0.46*	0.67*	-0.70*	-0.30	-0.53*	1.00

Table 4. Systematics of the meiobenthos found in the MILZON-BENTHOS II area 1992/93 (relevant taxa are underlined).

Phylum Cnidaria
 Classis Hydrozoa
 Phylum Plathyhelminthes
 Classis Turbellaria
 Phylum Gastrotrichia
 Phylum Kinorhyncha
 Phylum Priapulida
 Phylum Rotifera
 Phylum Nematoda
 Phylum Annelida
 Classis Polychaeta
 Ordo Archiannelida
 Classis Oligochaeta
 Phylum Arthropoda
 Classis Arachnida
 Fam. Halacarida
 Subphylum Crustacea
 Classis Ostracoda
 Classis Copepoda
 Ordo Harpacticoida
 Phylum Tardigrada




Table 5. Mean, median, minimum and maximum values of densities expressed in percentages, together with the corresponding standard deviation of the meiobenthic taxa in the MILZON-BENTHOS II area 1992/93.

	Mean	Median	Min.	Max.	St.dev.
Nematoda	82.38	85.56	44.64	99.64	15.25
Copepoda total	9.10	4.73	0.11	40.15	10.06
burrowing	1.02	0.73	0.00	4.81	1.10
interstitial	8.08	3.16	0.00	40.03	10.38
Gastrotricha	4.41	2.62	0.00	23.81	5.21
Turbellaria	1.74	1.21	0.00	6.95	1.73
Archiannelida	0.09	0.00	0.00	0.90	0.20
Oligochaeta	0.04	0.00	0.00	0.73	0.13
Polychaeta	0.23	0.12	0.00	1.42	0.30
Hydrozoa	0.32	0.00	0.00	3.46	0.61
Tardigrada	1.41	0.51	0.00	12.27	2.14
Ostracoda	0.21	0.08	0.00	2.44	0.41
Halacarida	0.02	0.00	0.00	0.49	0.06
Rotifera	0.01	0.00	0.00	0.66	0.07
Kinorhyncha	0.05	0.00	0.00	0.73	0.14
Priapulida	0.00	0.00	0.00	0.04	0.00

Table 6. Mean, median, minimum, and maximum values of densities (ind./10 cm²) with the standard deviation and presence (in percentages) of the meiobenthic taxa in the MILZON-BENTHOS II area 1992/93. The Nematoda/Copepoda (N/C) ratio is also given.

	Mean	Median	Min.	Max.	St.dev.	%
Density	1783.0	1472.0	412	6815	1213.8	
Taxa	7.2	7.0	4	12	1.9	
Nematoda	1540.0	1062.0	327	6711	1234.7	100
Copepoda total	131.0	65.0	1	1067	163.4	100
burrowing	20.1	8.0	0	238	32.3	92
interstitial	111.0	25.0	0	1067	169.1	78
Gastrotricha	54.1	33.0	0	293	66.3	84
Turbellaria	21.1	17.0	0	80	18.9	98
Archiannelida	1.3	0.0	0	15	3.1	28
Oligochaeta	0.8	0.0	0	12	2.4	22
Polychaeta	3.2	2.0	0	27	4.2	76
Hydrozoa	4.2	0.0	0	58	8.3	49
Tardigrada	21.2	5.0	0	296	41.4	69
Ostracoda	3.9	1.0	0	63	8.8	55
Halacarida	0.3	0.0	0	10	1.3	11
Rotifera	0.2	0.0	0	14	1.5	2
Kinorhyncha	1.8	0.0	0	36	5.7	17
Priapulida	0.0	0.0	0	1	0.2	2
N/C ratio	67.2	19.2	1	832	138.3	

Table 7. Correlations between the meiobenthos and the abiotic parameters, $\log(x+1)$ transformed, found in the MILZON-BENTHOS II area 1992/93. ($n=85$, $\alpha = 0.001$, $r=0.347$, 95 % reliability) **=significant

	Total	Nem.	Cop.	Copbur.	Copint.	Gastr.	Turb.	Arch.	Oligo.	Poly.	Hydr.	Tard.	Ostr.	Kin.
Total	1.00													
Nematoda	0.96*	1.00												
Copepoda	0.13	-0.10	1.00											
,, burrowing	-0.30	-0.50*	0.78*	1.00										
,, interstitial	-0.30	-0.50*	0.78*	-0.52*	1.00									
Gastropoda	-0.41*	-0.56*	0.39*	-0.62*	0.74*	1.00								
Turbellaria	-0.29	-0.41*	0.34	-0.45*	0.58*	0.76*	1.00							
Archiannelida	-0.05	-0.16	0.45*	-0.31	0.52*	0.41*	0.30	1.00						
Oligochaeta	0.16	0.10	0.34	-0.12	0.34	0.27	0.23	0.59*	1.00					
Polychaeta	0.07	-0.05	0.51*	-0.15	0.47*	0.24	0.22	0.50*	0.42*	1.00				
Hydrozoa	-0.15	-0.32	0.51*	-0.54*	0.67*	0.59*	0.42*	0.57*	0.33	0.46*	1.00			
Tardigrada	-0.18	-0.35*	0.63*	-0.47*	0.84*	0.67*	0.48*	0.47*	0.39*	0.40*	0.64*	1.00		
Ostracoda	0.42*	0.28	0.51*	0.16	0.25	0.02	-0.01	0.36*	0.38*	0.42*	0.30	0.33*	1.00	
Kinorhyncha	0.63*	0.63*	0.07	0.20	-0.49*	-0.49*	-0.42*	-0.07	0.01	-0.03	-0.27	-0.32	0.32	1.00
Depth	0.39*	0.45*	0.00	0.65	-0.32	-0.58*	-0.46*	-0.16	-0.10	-0.04	-0.41*	-0.31	0.20	0.47*
Median gr.size	-0.54*	-0.66*	0.40*	-0.70*	0.80*	0.80*	0.60*	0.46*	0.32	0.34	0.57*	0.75*	0.12	-0.57*
Mud content	0.52*	0.63*	-0.34	0.72*	-0.73*	-0.79*	-0.61*	-0.40*	-0.21	-0.23	-0.57*	-0.65*	-0.03	0.54*
Chl.a content	0.02	0.10	-0.34	0.19	-0.36*	-0.19	-0.07	-0.23	-0.20	-0.07	-0.36*	-0.25	-0.11	-0.02
POC content	0.39*	0.47*	-0.33	0.48*	-0.62*	-0.54*	-0.38*	-0.27	-0.16	-0.03	-0.38*	-0.62*	-0.04	0.41*
Oxidized layer	-0.41*	-0.54*	0.49*	-0.53*	0.77*	0.68*	0.58*	0.49*	0.22	0.40*	0.60*	0.62*	0.05	-0.35*
Taxa	0.04	-0.14	0.65*	-0.30	0.69*	0.51*	0.36*	0.66*	0.56*	0.67*	0.69*	0.68*	0.61*	-0.04
Hill ₁	-0.43*	-0.64*	0.70*	-0.54*	0.89*	0.80*	0.66*	0.52*	0.26	0.43*	0.70*	0.77*	0.25	-0.36*
Hill ₂	-0.39*	-0.62*	0.70*	-0.53*	0.85*	0.74*	0.60*	0.45*	0.18	0.40*	0.66*	0.71*	0.26	-0.34
Hill _m	-0.36*	-0.60*	0.69*	-0.53*	0.80*	0.68*	0.53*	0.40*	0.13	0.37*	0.64*	0.65*	0.28	-0.31
N/C ratio	0.30	0.50*	-0.91*	0.25	-0.88*	-0.57*	-0.46*	-0.45*	-0.26	-0.46*	-0.57*	-0.69*	-0.30	0.20

	Taxa	Hill ₁	Hill ₂	Hill ₃	N/C
Depth	-0.17	-0.43*	-0.42*	-0.40*	0.17
Median gr.size	0.53*	0.77*	0.72*	0.67*	-0.60*
Mud content	-0.44*	-0.72*	-0.67*	-0.64*	0.54*
Chl.a content	-0.23	-0.29	-0.29	-0.29	0.33
Feo.a content	-0.19	-0.29	-0.27	-0.25	0.28
POC content	-0.34	-0.54*	-0.50*	-0.46*	0.47*
Oxidized layer	0.49*	0.74*	0.68*	0.61*	-0.65*

Table 8. Mean values and (/)standard deviation of the meiobenthic and abiotic parameters per meiobenthic TWINSPAN cluster of the MILZON-BENTHOS II 1992/93 area.

Symbol Nr of locations	Cluster 1	Cluster 2	Cluster 3	Cluster 4
	12 mean/st.d.	19 mean/st.d.	43 mean/st.d.	11 mean/st.d.
<u>Abiotic parameter</u>				
Depth (m)	39.9 / 3.1	35.6 / 7.6	30.9 / 4.7	32.0 / 3.7
Median gr. size (μm)	126.1 / 11.6	181.7 / 58.2	268.3 / 37.8	320.2 / 40.7
Mud content (%)	13.67 / 3.28	8.94 / 6.70	2.89 / 2.28	2.02 / 0.95
Chl-a content ($\mu\text{g/g}$)	0.99 / 0.46	3.47 / 7.88	1.56 / 3.58	0.37 / 0.50
POC content (% C)	0.45 / 0.12	0.36 / 0.37	0.09 / 0.16	0.08 / 0.10
Oxidized layer (cm)	5.00 / 2.30	6.16 / 4.69	17.18 / 5.93	21.50 / 5.82
<u>Meiobenthos</u>				
Total (ind./10 cm ²)	3480.3 / 1540.2	2055.6 / 1254.9	1101.3 / 395.3	2125.5 / 741.1
Nematoda	3383.8 / 1513.9	2009.4 / 1242.9	841.7 / 361.1	1447.0 / 592.5
Copepoda total	73.7 / 57.5	24.8 / 17.2	130.7 / 121.9	378.6 / 254.8
burrowing	72.0 / 58.2	23.4 / 16.1	6.8 / 6.2	9.3 / 16.5
interstitial	1.7 / 4.9	1.4 / 1.6	123.8 / 122.8	369.3 / 256.9
Gastrotricha	0.2 / 0.4	8.0 / 9.7	72.6 / 66.9	120.1 / 64.6
Turbellaria	6.2 / 5.8	9.0 / 8.0	26.3 / 16.9	38.4 / 25.6
Archannelida	0.0 / 0.0	0.0 / 0.0	0.9 / 2.0	6.4 / 5.2
Oligochaeta	0.1 / 0.3	0.2 / 0.4	0.4 / 1.6	4.3 / 4.8
Polychaeta	2.2 / 2.3	1.3 / 1.8	2.7 / 2.6	9.4 / 7.5
Hydrozoa	0.0 / 0.0	0.2 / 0.4	4.2 / 6.1	15.4 / 15.2
Tardigrada	0.0 / 0.0	0.9 / 2.1	19.7 / 23.0	84.9 / 79.7
Ostracoda	4.0 / 2.6	1.2 / 2.3	2.0 / 3.4	16.1 / 19.7
Halacarida	0.0 / 0.0	0.1 / 0.2	0.1 / 0.3	2.0 / 3.2
Rotifera	0.1 / 0.3	0.0 / 0.0	0.0 / 0.0	1.3 / 4.2
Kinorhyncha	10.0 / 11.1	0.6 / 2.1	0.0 / 0.0	1.7 / 5.7
Priapulida	0.1 / 0.3	0.1 / 0.2	0.0 / 0.0	0.0 / 0.0
Nr of taxa (Hill ₀)	6.1 / 0.7	5.6 / 1.4	7.4 / 1.4	10.1 / 1.4
Hill ₁	1.16 / 0.08	1.16 / 0.09	2.27 / 0.61	2.83 / 0.50
Hill ₂	1.06 / 0.04	1.06 / 0.04	1.72 / 0.46	2.03 / 0.44
Hill _∞	1.03 / 0.02	1.03 / 0.02	1.37 / 0.28	1.52 / 0.27
N/C ratio	65.0 / 54.3	166.2 / 186.4	40.1 / 127.9	4.8 / 2.7

Table 9. Systematics of the macrobenthos found in the MILZON-BENTHOS II area 1992/92.
(dominant taxa are underlined)

Phylum Cnidaria

Class Hydrozoa
Class Anthozoa

Phylum Plathyhelminthes

Phylum Nemertinae

Symbols:

Phylum Mollusca

Class Bivalvia
Class Gastropoda




Phylum Sipunculida

Phylum Priapulida

Phylum Annelida

Class Polychaeta
Order Errantia
Order Sedentaria


Class Oligochaeta



Phylum Arthropoda

Subphylum Crustacea

Class Malacostraca
Order Decapoda
Order Mysidacea
Order Cumacea
Order Tanaidacea
Order Isopoda
Order Amphipoda




Phylum Tentaculata

Class Phoronida

Phylum Enteropneusta

Phylum Echinodermata

Class Echinoidea
Class Asteroidea
Class Ophiuroidea
Class Holothurioidea



Phylum Acrania

Family Branchiostoma

Table 10a. List of macrobenthic species (with shortening) found in the MILZON-BENTHOS II area 1992/93 and the number of location (n, %) were the species have been found. The mean values of density (ind./m²) and biomass (g/AFDW/m²) are given with their standard deviation.

I. POLYCHAETA : (59 species) (36.9 % of the total species)

		n	%	density main	st.dev.	biomass main	st.dev.
Anaitides groenlandica	(ANAI GROE)	4	4	0.8	3.8	0.0117	0.1045
Anaitides rosea	(ANAI ROSE)	11	12	4.6	20.5	0.0020	0.0079
Anaitides subulifera	(ANAI SUBU)	1	1	0.2	1.6	0.0001	0.0006
Aonides paucibranchiata	(AONI PAUC)	2	2	0.3	2.1	0.0001	0.0004
Aphrodite aculeta	(APHRACUL)	1	1	0.2	1.6	0.0001	0.0003
Aricidea jeffreysii	(ARIC JEFF)	1	1	0.3	3.2	0.0003	0.0024
Aricidea minuta	(ARIC MINU)	38	44	20.5	42.5	0.0050	0.0088
Chaetopterus variopedatus	(CHAEVARI)	6	7	3.3	19.6	0.7231	3.7254
Diplocirrus glaucus	(DIPL GLAU)	7	8	1.9	7.1	0.0013	0.0054
Eteone flava	(ETEO FLAV)	1	1	0.2	1.4	0.0008	0.0078
Eteone foliosa	(ETEO FOLI)	3	3	0.5	2.7	0.0009	0.0051
Eteone longa	(ETEO LONG)	6	7	1.1	4.2	0.0012	0.0048
Eumida sanguinea	(EUMISANG)	5	5	2.9	15.4	0.0017	0.0099
Exogone hebes	(EXOGHEBE)	4	4	0.8	3.9	0.0002	0.0010
Exogone naidina	(EXOGNAID)	9	10	28.0	216.0	0.0009	0.0049
Gattyana cirrosa	(GATT CIRR)	3	3	1.2	8.2	0.0243	0.1572
Glycinde nordmanni	(GLYCNORD)	1	1	0.2	1.4	0.0005	0.0047
Glycera rouxii	(GLYC ROUX)	2	2	1.0	8.1	0.0027	0.0244
Goniada maculata	(GONIMACU)	36	42	8.6	12.3	0.0358	0.0881
Gyptis capensis	(GYPT CAPE)	14	16	3.3	8.3	0.0023	0.0057
Harmothoe longisetis	(HARMLONG)	6	7	1.1	4.5	0.0010	0.0043
Harmothoe lunulata	(HARMLUNU)	17	20	8.7	22.8	0.0583	0.2576
Hesionura augeneri	(HESIAUGE)	7	8	2.4	8.7	0.0002	0.0009
Lanice conchilega	(LANICONC)	18	21	14.5	40.3	0.7707	2.9428
Lumbrineris latreilli	(LUMBLATR)	14	16	10.3	28.5	0.0305	0.0868
Magelona allenii	(MAGE ALLE)	2	2	1.0	8.1	0.0019	0.0168
Magelona papillicornis	(MAGEPAPI)	43	50	67.2	198.6	0.1762	0.3391
Mediomastus fragilis	(MEDIFRAG)	4	4	1.5	8.7	0.0003	0.0017
Nephtys cirrosa	(NEPH CIRR)	51	60	53.9	62.0	0.1104	0.1335
Nephtys hombergii	(NEPH HOMB)	32	37	12.2	21.3	0.3498	0.8674
Nephtys incisa	(NEPH INCI)	2	2	0.7	5.0	0.0129	0.0838
Nephtys longosetosa	(NEPH LONG)	11	12	3.7	11.9	0.0533	0.2208
Nereis longissima	(NERE LONG)	5	5	1.5	7.5	0.0635	0.5030
Notomastus latericeus	(NOTOLATE)	14	16	6.1	18.3	0.1412	0.4484
Ophelina acuminata	(OPHEACUM)	3	3	0.7	3.8	0.0003	0.0015
Ophelia limacina	(OPHELIMA)	8	9	2.1	6.8	0.0031	0.0123
Ophidromus flexuosus	(OPHIFLEX)	11	12	3.1	9.6	0.0101	0.0331
Orbinia sertulata	(ORBISERT)	2	2	0.5	3.5	0.0431	0.3818
Owenia fusiformis	(OWENFUSI)	2	2	0.7	5.0	0.0006	0.0041
Paraonis gracilis	(PARAGRAC)	2	2	0.5	3.5	0.0001	0.0009
Pectinaria auricoma	(PECTAURI)	1	1	0.2	1.6	0.0001	0.0002
Pectinaria koreni	(PECT KORE)	2	2	1.5	12.8	0.0053	0.0455
Pholoe minuta	(PHOLMINU)	18	21	9.8	26.9	0.0052	0.0157
Pista cristata	(PIST CRIS)	1	1	0.3	3.2	0.0009	0.0079
Poecilochaetus serpens	(POEC SERP)	10	11	7.9	39.3	0.0072	0.0265
Polydora guillei	(POLY GUIL)	8	9	8.6	30.7	0.0067	0.0292
Prionospio cirrfera	(PRIOCIRR)	5	5	7.4	33.4	0.0054	0.0257
Scalibregma inflatum	(SCALINFL)	3	3	0.5	2.6	0.0002	0.0011
Scoloplos armiger	(SCOLARMI)	51	60	44.7	85.7	0.2889	0.4567
Scolelepis bonnieri	(SCOLBONN)	6	7	1.3	5.3	0.0178	0.1196
Scolelepis squamata	(SCOL SQUA)	14	16	4.5	12.9	0.0769	0.3097
Sigalion mathildae	(SIGAMATH)	12	14	2.8	8.6	0.0439	0.1558
Spiophanes bombyx	(SPIOBOMB)	60	70	111.4	265.7	0.1083	0.4088
Spio filicornis	(SPIO FILI)	30	35	20.4	47.1	0.0060	0.0153
Sthenelais limicola	(STHE LIMI)	21	24	4.6	9.0	0.0408	0.1275
Streptosyllis websteri	(STREWEBS)	6	7	0.9	3.5	0.0001	0.0005
Synelmis klatti	(SYNEKLAT)	10	11	3.3	10.2	0.0008	0.0028
Travisia forbesi	(TRAVFORB)	1	1	0.2	1.6	0.0001	0.0003

Table 10b. List of macrobenthic species (with shortening) found in the MILZON-BENTHOS II area 1992/93 and the number of location (n, %) were the species have been found. The mean values of density (ind./m²) and biomass (g/AFDW/m²) are given with their standard deviation.

II. CRUSTACEA : (57 species) (35.6 % of the total species)

		n	%	density main	st.dev.	biomass main	st.dev.
Ampelisca tenuicornis	(AMPETENU)	7	8	1.5	5.5	0.0008	0.0028
Amphilochus neapolitanus	(AMPHNEAP)	1	1	0.2	1.6	0.0001	0.0006
Aora typica	(AORATYPI)	1	1	0.3	3.2	0.0003	0.0032
Apherusa ovalipes	(APHEOVAL)	2	2	0.3	2.0	0.0002	0.0010
Argissa hamatipes	(ARGIHAMA)	2	2	1.2	9.6	0.0004	0.0029
Atylus falcatus	(ATYLFALC)	2	2	0.3	2.1	0.0002	0.0010
Atylus swammerdami	(ATYLSWAM)	6	7	2.6	12.8	0.0008	0.0035
Bathyporeia elegans	(BATHELEG)	31	36	45.0	115.5	0.0133	0.0306
Bathyporeia guilliamsoniana	(BATHGUIL)	37	43	43.7	108.4	0.0262	0.0567
Bathyporeia tenuipes	(BATHTENU)	6	7	2.9	12.9	0.0011	0.0048
Bodotria arenosa	(BODOAREN)	1	1	0.2	1.6	0.0001	0.0008
Bodotria scorpioides	(BODOSCOR)	1	1	0.2	1.6	0.0001	0.0003
Callianassa subterranea	(CALLSUBT)	26	30	39.7	77.9	1.0351	2.1963
Caprellidae spec.	(CAPRELLI)	5	5	0.9	4.0	0.0002	0.0010
Cirolana borealis	(CIROBORE)	1	1	0.2	1.6	0.0001	0.0008
Coryistes cassivelaunus	(CORYCASS)	5	5	0.8	3.4	0.1408	1.1126
Crangon crangon	(CRANCRAN)	1	1	0.2	1.4	0.0002	0.0021
Diastylis bradyi	(DIASBRAD)	6	7	1.4	5.3	0.0008	0.0034
Diastylis lucifera	(DIASLUCI)	1	1	0.5	4.8	0.0002	0.0016
Diastylis rathkei	(DIASRATH)	3	3	0.5	2.4	0.0005	0.0027
Diastylis rugosa	(DIASRUGO)	1	1	0.2	1.6	0.0003	0.0024
Ebalia chanchii	(EBALCRAN)	3	3	0.5	2.6	0.0005	0.0030
Eudorellopsis deformis	(EUDODEFO)	2	2	0.9	6.5	0.0002	0.0016
Eudorella truncatula	(EUDOTRUN)	9	10	1.7	5.3	0.0005	0.0015
Eurydice spinigera	(EURYSPIN)	1	1	0.3	2.8	0.0006	0.0059
Gastrosaccus spinifer	(GASTSPIN)	18	21	5.6	13.1	0.0091	0.0255
Harpinia antennaria	(HARPANTE)	16	18	12.7	37.1	0.0034	0.0097
Harpinia crenulata	(HARPCREN)	2	2	0.7	4.5	0.0002	0.0013
Hippomedon denticutaus	(HIPPDENT)	17	20	3.8	9.5	0.0015	0.0036
Iphinoe trispinosa	(IPHITRIS)	1	1	0.2	1.6	0.0001	0.0008
Leucothoe incisa	(LEUCINCI)	5	5	1.6	8.8	0.0011	0.0053
Leucothoe lilljeborgii	(LEUCLILL)	11	12	2.6	7.8	0.0012	0.0037
Leucothoe richiardii	(LEUCRICH)	3	3	1.0	5.9	0.0005	0.0029
Liocarcinus pusillus	(LIOPCUSI)	1	1	0.2	1.6	0.0001	0.0008
Megaluropus agilis	(MEGAAGIL)	44	51	18.2	26.3	0.0068	0.0108
Melita obtusata	(MELIOBTU)	1	1	0.2	1.4	0.0002	0.0014
Metopa borealis	(METOBORE)	1	1	0.2	1.6	0.0001	0.0008
Mysidacea spec.	(MYSIDACE)	1	1	0.2	1.6	0.0001	0.0008
Orchomene humilis	(ORCHHUMI)	1	1	0.3	2.8	0.0002	0.0014
Periocolodes longimanus	(PERILONG)	15	17	3.8	10.5	0.0014	0.0041
Phoxichilidium femoratum	(PHOXFEMO)	1	1	0.2	1.4	0.0001	0.0003
Pinnotheres pisum	(PINNPISU)	2	2	0.3	2.0	0.0005	0.0029
Pontocrates altamarinus	(PONTALTA)	20	23	8.0	23.6	0.0026	0.0079
Pontocrates arenarius	(PONTAREN)	7	8	3.6	14.3	0.0012	0.0049
Pontophylus trispinosus	(PONTTRIS)	5	5	0.9	3.5	0.0004	0.0017
Processa nouveli	(PROCNOUV)	1	1	0.2	1.4	0.0006	0.0055
Pseudocuma longicornis	(PSEULONG)	44	51	19.7	31.1	0.0041	0.0058
Pseudocuma similis	(PSEUSIMI)	1	1	0.5	4.8	0.0003	0.0011
Siphonoecetes kroyeranus	(SIPHKROY)	1	1	0.2	1.4	0.0001	0.0004
Siriella clausii	(SIRICLAU)	1	1	0.2	1.4	0.0002	0.0014
Synchelidium maculatum	(SYNCMACU)	1	1	0.2	1.6	0.0001	0.0008
Tanaidacea spec.	(TANAIDAC)	1	1	0.3	2.8	0.0001	0.0001
Thia scutellata	(THIASCUT)	8	9	1.7	5.8	0.0494	0.2284
Upogebia deltaura	(UPOGDELT)	10	11	4.3	13.7	0.9676	3.6008
Urothoe brevicornis	(UROTBREV)	30	35	24.9	47.4	0.0152	0.0308
Urothoe elegans	(UROTELEG)	6	7	1.9	7.8	0.0006	0.0022
Urothoe poseidonis	(UROPOSE)	21	24	38.7	118.2	0.0120	0.0336

Table 10c. List of macrobenthic species (with shortening) found in the MILZON-BENTHOS II area 1992/93 and the number of location (n, %) were the species have been found. The mean values of density (ind./m²) and biomass (g/AFDW/m²) are given with their standard deviation.

III. MOLLUSCA : (26 species) (16.2 % of the total species)

		n	%	density main	st.dev.	biomass main	st.dev.
Abra alba	(ABRAALBA)	7	8	3.3	16.9	0.0009	0.0042
Abra prismatica	(ABRAPRIS)	1	1	0.3	3.2	0.0015	0.0136
Acanthocardia echinata	(ACANECHI)	1	1	0.2	1.6	0.0001	0.0006
Cochlodesma pratenue	(COCHPRAE)	1	1	1.7	15.9	0.0167	0.1539
Corbula gibba	(CORBGIBB)	2	2	0.3	2.2	0.0009	0.0083
Cultellus pellucidus	(CULTPELL)	13	15	3.6	9.8	0.0012	0.0037
Cylinchna cylindracea	(CYLICLYI)	4	4	0.7	3.0	0.0002	0.0010
Donax vittatus	(DONAVITT)	6	7	1.8	8.5	0.1740	1.0884
Dosinia exolata	(DOSIEXOL)	1	1	1.9	17.5	0.0017	0.0157
Dosinia lupinus	(DOSILUPI)	2	2	0.3	2.1	0.0217	0.1996
Ensis ensis	(ENSIENSI)	2	2	0.3	2.1	0.0180	0.1164
Ensis siliqua	(ENSISILI)	3	3	0.6	3.5	1.0332	7.7873
Gari fervensis	(GARIFERV)	1	1	0.2	1.6	0.0038	0.0348
Lepton squamosum	(LEPTSQUA)	1	1	0.2	1.6	0.0001	0.0002
Lutraria lutraria	(LUTRLUTR)	1	1	0.2	1.6	1.2271	11.3134
Mactra corallina	(MACTCORA)	1	1	1.5	14.3	0.0002	0.0014
Montacuta ferruginosa	(MONTFERR)	23	27	11.5	28.0	0.0069	0.0218
Mysella bidentata	(MYSEBIDE)	24	28	129.7	306.4	0.0263	0.0644
Natica alderi	(NATIALDE)	30	35	10.0	18.0	0.0137	0.0489
Nucula turgida	(NUCUTURG)	7	8	2.7	10.7	0.0119	0.0736
Spisula subtruncata	(SPISSUBT)	1	1	0.2	1.4	0.0751	0.6929
Tellina fabula	(TELLFABU)	24	28	16.7	48.2	0.0795	0.3196
Tellina pygmæus	(TELLPYGM)	1	1	0.2	1.6	0.0001	0.0002
Thrasia convexa	(THRACONV)	1	1	0.2	1.6	0.0123	0.1130
Turbonilla lactea	(TURBLACT)	1	1	0.2	1.6	0.0001	0.0003
Venus striatula	(VENUSTRI)	4	4	0.6	2.9	0.1011	0.6641

IV. ECHINODERMATA : (8 species) (6.3 % of the total species)

		n	%	density main	st.dev.	biomass main	st.dev.
Amphiura filiformis	(AMPHFILI)	22	25	163.2	374.4	2.3258	5.8756
Acrocnida brachiate	(ACROBRAC)	1	1	0.2	1.6	0.0001	0.0002
Brissopsis lyrifera	(BRISLYRI)	1	1	0.2	1.6	0.1274	1.1743
Echinocardium cordatum	(ECHICORD)	44	51	18.3	29.9	6.8456	13.3072
Echinocyamus pusillus	(ECHIPUSI)	25	29	17.6	63.0	0.0100	0.0228
Leptosynapta inhaerens	(LEPTINHA)	4	4	1.0	4.9	0.0308	0.1732
Ophiura albida	(OPHIALBI)	14	16	10.1	46.0	0.0206	0.0872
Ophiura texturata	(OPHITEXT)	2	2	0.3	2.0	0.0058	0.0530

V. REST : (10 species) (5.0 % of the total species)

		n	%	density main	st.dev.	biomass main	st.dev.
Anthozoa spec.	(ANTHOZOA)	7	8	7.4	53.1	0.0052	0.0419
Branchiostoma lanceolatum	(BRANLANC)	1	1	1.5	13.9	0.0348	0.3212
Enteropneusta	(ENTEROPN)	1	1	0.2	1.4	0.0001	0.0001
Hydrozoa spec.	(HYDROZOA)	4	4	0.8	4.0	0.0001	0.0006
Nemertinae spec.	(NEMERTIN)	68	80	21.6	20.1	0.1167	0.2614
Oligochaeta spec.	(OLIGOCHA)	6	7	4.1	22.0	0.0003	0.0015
Phoronida spec.	(PHORONID)	26	30	36.3	124.2	0.0158	0.0560
Plathyhelminthes spec.	(PLATHYHE)	5	5	1.5	7.1	0.0016	0.0121
Priapulida spec.	(PRIAPULI)	1	1	0.2	1.6	0.0001	0.0008
Sipunculida spec.	(SIPUNCUL)	6	7	7.7	39.0	0.0819	0.4529

VI. TOTAL NUMBER OF SPECIES : 160

Table 11. Presence (n, %) of the most abundant macrobenthos species, found at more than 25 % of the stations in the MILZON-BENTHOS II area 1992/93. The mean values of density (ind./m²) and biomass (g AFDW/m²) are given with their standard deviation. Abundance patterns are shown in indicated maps.

species	Fig.	Number of presence			density (ind./m ²)		biomass (g AFDW/m ²)	
		n	%	mean	st.dev.	mean	st.dev.	
Nemertinae spec.	73.	68	80	21.6	20.2	0.1167	0.2614	
Spiophanes bombyx	56.	60	70	111.4	265.7	0.1083	0.4088	
Nephtys cirrosa	52.	51	60	53.9	62.0	0.1104	0.1335	
Scoloplos armiger	55.	51	60	44.7	85.7	0.2889	0.4567	
Echinocardium cordatum	71.	44	51	18.3	29.9	6.8456	13.3072	
Megaluropus agilis	67.	44	51	18.2	26.3	0.0068	0.0108	
Pseudocuma longicornis	68.	44	51	19.7	31.1	0.0041	0.0058	
Magelona papillicornis	51.	43	50	67.2	198.6	0.1762	0.3391	
Aricidea minuta	47.	38	44	20.5	42.5	0.0050	0.0088	
Bathyporeia guilliamsoniana	64.	37	43	43.7	108.4	0.0262	0.0567	
Goniada maculata	49.	36	42	8.6	12.3	0.0358	0.0881	
Chaetozone setosa	48.	35	41	21.5	66.0	0.0084	0.0242	
Nephtys hombergii	53.	32	37	12.2	21.3	0.3498	0.8674	
Bathyporeia elegans	63.	31	36	45.0	115.5	0.0133	0.0306	
Spio filicornis	57.	30	35	20.4	47.1	0.0060	0.0153	
Natica alderi	61.	30	35	10.0	18.0	0.0137	0.0489	
Urothoe brevicornis	69.	30	35	24.9	47.4	0.0152	0.0308	
Callianassa subterranea	65.	26	30	39.7	77.9	1.0351	2.1963	
Phoronide spec.	74.	26	30	36.3	124.2	0.0158	0.0560	
Echinocyamus pusillus	72.	25	29	17.6	63.0	0.0100	0.0228	
Mysella bidentata	60.	24	28	129.7	306.4	0.0263	0.0644	
Tellina fabula	62.	24	28	16.7	48.2	0.0795	0.3196	
Montacuta ferruginosa	59.	23	27	11.5	28.0	0.0069	0.0218	
Amphiura filiformis	70.	22	25	163.2	374.4	2.3258	5.8756	

Table 12. Presence (number and percentage of stations), minimum, maximum, mean value with standard deviation and percentage of the mean of some taxonomic macrobenthic groups in the MILZON-BENTHOS II area 1992/93.

	presence		min.	max.	mean	st.dev.	% of mean
	n	%					
Density:							
Polychaeta	85	100	87.8	2311.5	528.1	441.5	40.36
Crustacea	85	100	29.3	1385.6	302.7	245.8	23.14
Mollusca	67	79	0.0	1477.6	189.2	313.9	14.46
Echinodermata	69	81	0.0	1741.0	210.9	377.2	16.12
Branchiostoma	1	1	0.0	128.3	1.5	13.9	0.12
Cnidaria	11	13	0.0	487.5	8.3	53.2	0.63
Enteropneusta	1	1	0.0	12.8	0.2	1.4	0.02
Nemertinae	68	80	0.0	117.0	21.6	20.1	1.65
Phoronida	26	31	0.0	1024.1	36.3	124.2	2.77
Plathyhelminthes	5	6	0.0	43.9	1.5	7.1	0.12
Priapulida	1	1	0.0	14.6	0.2	1.6	0.02
Sipunculida	6	7	0.0	278.0	7.7	39.0	0.59
Sum density (ind./m ²)			256.6	4125.7	1308.4	835.5	100.00
Biomass :							
Polychaeta	85	100	0.2852	31.5919	3.2657	5.3357	18.12
Crustacea	85	100	0.0103	25.1841	2.3039	4.8537	12.79
Mollusca	67	79	0.0000	104.5139	2.8280	13.6605	15.69
Echinodermata	69	81	0.0000	93.3789	9.3660	13.9716	51.98
Branchiostoma	1	1	0.0000	2.9612	0.0348	0.3212	0.19
Cnidaria	11	13	0.0000	0.3962	0.0053	0.0419	0.03
Enteropneusta	1	1	0.0000	0.0012	0.0001	0.0001	0.00
Nemertinae	68	80	0.0000	1.7644	0.1167	0.2614	0.65
Phoronida	26	31	0.0000	0.4550	0.0158	0.0560	0.09
Plathyhelminthes	5	6	0.0000	0.1112	0.0016	0.0012	0.01
Priapulida	1	1	0.0000	0.0073	0.0001	0.0008	0.00
Sipunculida	6	7	0.0000	3.6165	0.0819	0.4529	0.45
Sum biomass (g AFDW/m ²)			0.79	135.58	18.02	21.81	100.00
Hill ₀ (number of species)	11.0		33.0	18.4	4.7		
Hill ₁		2.8	23.4	9.8	3.4		
Hill ₂		1.6	17.5	6.8	3.0		
Hill _∞		1.3	7.9	3.7	1.5		

Table 13. Correlation between the macrobenthos and the abiotic parameters, log (x+1) transformed, found in the MILZON-BENTHOS II area 1992/93. (n=85, $\alpha=0.001$, $r=0.347$, 95 % reliability) * = significant

	Density (ind./m ²):				Biomass (g AFDW/m ²):				Diversity :			
	total	Poly.	Crus.	Moll.	total	Poly.	Crus.	Moll.	Spec.	Hill ₁	Hill ₂	Hill ₃
total density	1.00											
Polychaeta "	0.52*	1.00										
Crustacea "	0.34	-0.05	1.00									
Mollusca "	0.67*	0.15	0.16	1.00								
Echinoderm."	0.63*	-0.06	0.27	0.55*	1.00							
total biomass	0.48*	-0.00	0.24	0.38*	0.62*	1.00						
Polychaeta "	0.36*	0.45*	0.07	0.22	0.14	0.33	1.00					
Crustacea "	0.52*	-0.16	0.20	0.45*	0.57*	0.51*	0.16	1.00				
Mollusca "	0.25	0.04	0.25	0.32	0.11	0.36*	-0.02	0.07	1.00			
Echinoderm."	0.31	-0.15	0.19	0.25	0.62*	0.81*	0.03	0.33	0.03	1.00		
Species number	0.65*	0.33	0.30	0.64*	0.62*	0.48*	0.42*	0.38*	0.33	0.26	1.00	
Hill :	-0.38*	-0.16	0.02	-0.02	-0.03	0.02	0.13	-0.14	0.01	-0.04	0.35*	1.00
Hill :	-0.53*	-0.24	-0.07	-0.16	-0.18	-0.07	0.06	-0.22	-0.05	-0.11	0.11	0.95*
Hill -	-0.52*	-0.24	-0.09	-0.18	-0.19	-0.07	0.06	-0.22	-0.03	-0.12	0.01	0.85*
Depth	0.43*	-0.05	0.03	0.37*	0.63*	0.45*	0.17	0.55*	-0.08	0.41*	0.35*	-0.12
Median gr.size	-0.50*	0.15	-0.12	-0.60*	-0.52*	-0.47*	-0.30	-0.67*	-0.08	-0.33	-0.39*	0.12
Mud content	0.45*	-0.08	0.03	0.50*	0.49*	0.51*	0.32	0.70*	0.05	0.34	0.41*	-0.04
Chl.a content	0.15	0.11	0.01	0.14	-0.01	0.21	0.11	0.07	0.02	0.20	0.14	-0.02
POC content	0.49*	0.17	-0.08	0.48*	0.28	0.35*	0.35*	0.59*	0.05	0.20	0.28	-0.21
Oxidized layer	-0.41*	0.02	-0.09	-0.54*	-0.50*	-0.55*	-0.37*	-0.55*	-0.19	-0.39*	-0.54*	-0.14
												0.02

Table 14. Mean values of abiotic and macrobenthic parameters of each macrobenthic TWINSPAN cluster, as shown in Figs. 75 and 76, with their standard deviation.

Symbols	Cluster 1	Cluster 2	Cluster 3	Cluster 4
	⊗	◐	◀	◇
Number of location	18	5	19	43
Abiotic parameters :	mean/st.d.	mean/st.d.	mean/st.d.	mean/st.d.
Depth (m)	41.8 / 3.9	33.2 / 2.9	29.8 / 3.8	31.5 / 4.5
Median grain size (μm)	136.5 / 24.5	150.6 / 25.0	229.9 / 55.0	288.7 / 36.6
Mud content (%)	13.10 / 6.5	9.44 / 3.5	5.11 / 3.6	2.34 / 1.1
Chl.a content ($\mu\text{g/g}$)	1.00 / 0.7	1.21 / 0.5	4.39 / 8.8	0.96 / 1.9
POC content (% C)	0.45 / 0.2	0.28 / 0.1	0.25 / 0.4	0.06 / 0.1
Oxidized layer (cm)	0.7 / 0.2	0.9 / 0.1	0.9 / 0.3	1.3 / 0.1
Macrobenthos :				
Total number of species	83	50	98	87
Density (ind./ m^2)	2177.4 / 862.9	921.0 / 594.7	1588.5 / 686.7	865.4 / 530.5
Biomass (g AFDW/ m^2)	36.01 / 28.34	15.33 / 12.5	18.71 / 16.58	10.50 / 17.20
Polychaeta (ind./ m^2)	372.3 / 162.7	298.5 / 135.9	838.3 / 538.2	483.0 / 437.0
Crustacea (")	290.2 / 138.5	155.1 / 39.5	413.2 / 368.0	276.3 / 214.5
Mollusca (")	588.5 / 416.2	280.9 / 482.8	142.8 / 100.0	31.9 / 69.8
Echinodermata (")	778.6 / 481.3	84.9 / 93.3	103.2 / 161.2	35.4 / 45.8
Polychaeta (g AFDW/ m^2)	5.64 / 8.44	3.90 / 5.10	4.76 / 6.50	1.49 / 1.02
Crustacea (")	8.20 / 7.34	2.49 / 1.83	1.33 / 3.49	0.24 / 0.32
Mollusca (")	6.33 / 24.54	0.19 / 0.28	5.77 / 16.10	0.42 / 1.54
Echinodermata(")	15.11 / 9.56	8.57 / 12.44	6.76 / 9.00	8.21 / 16.89
Hill ₀ (species richness)	21.3 / 3.9	19.6 / 3.0	21.8 / 4.7	15.5 / 3.2
Hill ₁ diversity	8.5 / 3.1	12.6 / 3.9	11.4 / 4.4	9.3 / 2.4
Hill ₂ "	5.3 / 2.5	9.6 / 4.3	7.6 / 3.8	6.8 / 2.3
Hill _∞ "	2.9 / 1.0	5.3 / 2.5	3.8 / 1.7	3.7 / 1.3

Table 15. List of abundant macrobenthos species in each macrobenthic TWINSPLAN cluster of the MILZON-BENTHOS II area 1992/93. The presence (number and percentage of stations) and mean values with the standard deviation of density (ind./m²) and biomass (g AFDW/m²) (cf. Fig. 75). **Bold-printed** species were 'indicator species' of the 4 clusters.

	species	presence n %	density mean	density st.dev.	biomass mean	biomass st.dev.
Cluster 1						
	Amphiura filiformis	18 100	752.6	471.8	10.7055	8.6680
	Callianassa subterranea	18 100	154.4	94.0	3.5557	2.2278
	Mysella bidentata	17 94	552.7	407.9	0.1163	0.0931
	Pholoe minuta	14 77	40.6	46.1	0.0216	0.0285
	Harpinia antennaria	13 72	54.5	64.4	0.0144	0.0165
	Nephtys hombergii	13 72	28.4	29.0	0.4357	0.5950
	Echinocardium cordatum	12 66	20.3	22.5	3.6612	5.4112
Cluster 2						
	Callianassa subterranea	5 100	96.6	48.1	2.2422	2.0803
	Cultellus pellucidus	4 80	23.4	13.1	0.0111	0.0063
	Lumbrineris latreilli	4 80	43.9	38.7	0.1235	0.0983
	Magelona papillicornis	4 80	20.5	13.1	0.0108	0.0088
	Nephtys hombergii	4 80	20.5	22.2	1.2471	2.2435
Cluster 3						
	Chaetozone setosa	18 94	75.2	125.9	0.0248	0.0451
	Magelona papillicornis	18 94	256.5	365.7	0.4785	0.5040
	Spiophanes bombyx	16 84	209.5	317.9	0.1642	0.4414
	Tellina fabula	15 78	63.7	85.8	0.3073	0.6200
	Natica alderi	14 73	24.7	24.0	0.0409	0.0883
	Nephtys hombergii	14 73	21.3	22.1	0.7461	1.1416
	Goniada maculata	12 63	16.6	16.4	0.1082	0.1593
Cluster 4						
	Nephtys cirrosa	40 93	92.4	58.6	0.1899	0.1339
	Scoloplos armiger	39 90	76.0	104.8	0.5108	0.5235
	Aricidea minuta	35 81	37.6	53.0	0.0091	0.0105
	Megaluropus agilis	35 81	20.5	30.2	0.0115	0.0130
	Spiophanes bombyx	34 79	116.3	297.4	0.1350	0.4929
	Pseudocuma longicornis	31 72	30.4	37.5	0.0062	0.0066
	Urothoe brevicornis	29 67	48.6	57.5	0.0298	0.0381
	Bathyporeia guilliamsoniana	27 62	46.9	94.3	0.0324	0.0526

Table 16. Spearman rank correlations between the meio- and macrobenthic TWINSPLAN scores and abiotic parameters. Prior to the calculations, all data were $(x+1)$ -transformed. Number of observations = 3570 ($n*(n-1)/2$; $n=85$).

Abiotic parameters TWINSPLAN Scores	Depth	Median	Mud	Chl.a	POC	Oxidized layer
Meiobenthos	0.568	-0.823	0.780	0.451	0.643	-0.739
Macrobenthos	0.535	-0.740	0.708	0.348	0.647	-0.721

Table 17. Spearman rank correlations between benthic (meio- and macrobenthos) and abiotic dissimilarity matrices. k gives the number of combinations of environmental parameters, yielding the 'best matches' of the respective dissimilarity matrices. Number of observations = 3570 ($n*(n-1)/2$; $n=85$).

A. Meiobenthos

k Best environmental parameter combinations

1	Depth	Median	Mud	Chl-a	POC	Oxygen layer
	.261	.762	.657	.091	.441	.564
2	Depth/Median	Depth/Mud		Median/Mud		
	.743		.665		.724	
3	Depth/Median/Mud					.727

B. Macrobenthos

k Best environmental parameter combinations

1	Depth	Median	Mud	Chl-a	POC	Oxygen layer
	.407	.729	.666	.061	.456	.504
2	Depth/Median	Depth/Mud		Median/Mud		
	.767		.694		.723	
3	Depth/Median/Mud					.743

Table 18. Correlations between the meio- and macrobenthos, $\log(x+1)$ transformed, found in the MILZON-BENTHOS II area 1992/1993.
(n=85, $\alpha=0.001$, $r=0.347$, 95 % reliability) *=significant

	Macrofauna:											
	Density (ind./m²)				Biomass (g AFDW/m²)				Diversity			
Meiobenthos:												
Total density	0.42*	0.01	-0.02	0.49*	0.54*	0.39*	0.12	0.47*	0.08	0.31	0.34	-0.03
Taxa	-0.26	-0.01	-0.15	-0.45*	-0.21	-0.34	-0.24	-0.22	-0.19	-0.30	-0.28	-0.05
Hill ₁	-0.55*	-0.06	-0.20	-0.71*	-0.50*	-0.57*	-0.32	-0.50*	-0.27	-0.40*	-0.55*	-0.04
Hill ₂	-0.55*	-0.09	-0.22	-0.70*	-0.51*	-0.55*	-0.30	-0.46*	-0.28	-0.38*	-0.57*	-0.06
Hill _m	-0.52*	-0.08	-0.22	-0.68*	-0.50*	-0.53*	-0.27	-0.43*	-0.27	-0.36*	-0.56*	-0.09
Nematoda	0.52*	0.03	0.05	0.62*	0.61*	0.49*	0.19	0.53*	0.15	0.37*	0.46*	-0.00
Copepoda	-0.25	-0.11	-0.22	-0.45*	-0.13	-0.24	-0.28	-0.12	-0.20	-0.16	-0.38*	-0.15
burrowing	0.49*	-0.09	0.01	0.52*	0.66*	0.52*	0.11	0.61*	0.11	0.40*	0.38*	-0.13
interstitial	-0.49*	-0.00	-0.10	-0.68*	-0.42*	-0.50*	-0.33	-0.50*	-0.22	-0.36*	-0.47*	0.02
Gastrotricha	-0.45*	0.11	-0.08	-0.53*	-0.46*	-0.56*	-0.31	-0.60*	-0.18	-0.40*	-0.45*	-0.01
Turbellaria	-0.38*	0.04	-0.09	-0.44*	-0.36*	-0.41*	-0.28	-0.47*	-0.16	-0.25	-0.32	0.01
Archiannelida	-0.23	0.09	-0.26	-0.26	-0.25	-0.29	-0.10	-0.28	-0.07	-0.28	-0.19	0.01
Oligochaeta	-0.04	0.16	-0.24	-0.12	-0.05	-0.19	-0.04	-0.10	0.09	-0.33	0.10	0.13
Polychaeta	-0.13	0.11	-0.32	-0.38*	-0.21	-0.21	-0.06	-0.13	-0.22	-0.22	-0.26	-0.11
Hydrozoa	-0.43*	-0.02	-0.12	-0.46*	-0.39*	-0.53*	-0.23	-0.37*	-0.20	-0.44*	-0.43*	-0.01
Tardigrada	-0.39*	0.06	-0.03	-0.51*	-0.36*	-0.46*	-0.33	-0.49*	-0.09	-0.38*	-0.30	0.10
Ostracoda	0.01	-0.08	-0.17	-0.00	0.18	0.04	-0.06	0.12	-0.13	-0.00	0.05	0.04
Halacarida	0.00	0.21	-0.03	-0.04	-0.12	-0.27	-0.11	-0.10	-0.04	-0.28	0.03	0.01
Rotifera	0.14	0.10	-0.09	0.11	0.03	0.01	-0.00	0.11	-0.05	-0.07	0.09	0.04
Kinorhyncha	0.51*	0.03	0.04	0.46*	0.53*	0.39*	0.15	0.60*	0.09	0.30	0.27	-0.23
N/C ratio	0.44*	0.13	0.22	0.64*	0.36*	0.43*	0.33	0.32	0.24	0.30	0.52*	0.13
											0.01	-0.01

Figures

Figures :

I Abiotic parameters

1. Survey area of the MILZON-BENTHOS I (1988-1989) and II (1991-1993) projects.
2. Locations of ICES (1986) and the EXP*BMN project 1992 in the MILZON-BENTHOS II area 1992/93.
3. Water types and their general boundaries in the North Sea in the summer period.
4. Depth (m) recordings, uncorrected for differences due to the tidal cycle.
5. Median grain size (for the fraction 63 - 2000 µm) of the upper 10 cm of the sediment.
6. Mud content (fraction < 63 µm, %) of the upper 10 cm of the sediment.
7. Chlorophyll-a content (µg/g) of the upper 3 cm of the sediment.
8. POC content (% C) of the upper 3 cm of the sediment.
9. Depth of the (visible) oxygen layer (cm) of the sediment.

II Meiobenthos

10. Total number of meiobenthic taxa ($Hill_0$) .
11. Diversity ($Hill_1$) of the meiobenthos.
12. Diversity ($Hill_2$) of the meiobenthos.
13. Diversity ($Hill_\infty$) of the meiobenthos.
14. Total density of the meiobenthos (ind./10 cm²) .
15. Spatial distribution of the Nematoda (ind./10 cm²) .
16. Spatial distribution of the Copepoda (ind./10 cm²) .
- 17a/b. Spatial distribution of the interstitial Copepoda (ind./10 cm²) .
18. Spatial distribution of the Nematoda/Copepoda (N/C) ratio.
19. Spatial distribution of the Gastrotricha (ind./10 cm²) .
20. Spatial distribution of the Turbellaria (ind./10 cm²) .
21. Spatial distribution of the Archiannelida (ind./10 cm²) .
22. Spatial distribution of the Oligochaeta (ind./10 cm²) .
23. Spatial distribution of the Polychaeta (< 1mm) (ind./10 cm²) .
24. Spatial distribution of the Hydrozoa (ind./10 cm²) .
25. Spatial distribution of the Tardigrada (ind./10 cm²) .
26. Spatial distribution of the Ostracoda (ind./10 cm²) .
27. Spatial distribution of the Halacarida (ind./10 cm²) .
28. Spatial distribution of the Kinorhyncha (ind./10 cm²) .
29. TWINSPLAN clustering of the meiobenthos density (ind./10 cm²) .
30. TWINSPLAN dichotomy of the meiobenthos density (ind./10 cm²) .
31. (a-c) Box and whisker plots of some average biotic and abiotic variables after TWINSPLAN analysis using the data of the meiobenthos density (ind./10 cm²) .
32. Multidimensional scaling (MDS) ordination of the meiobenthos density (ind./10 cm²) in relation to the TWINSPLAN clusters of the MILZON-BENTHOS area 1992/93.

III Macrobenthos

33. Total number of macrobenthic species ($Hill_0$) .
34. Diversity ($Hill_1$) of the macrobenthos.
35. Diversity ($Hill_2$) of the macrobenthos.
36. Diversity ($Hill_\infty$) of the macrobenthos.
37. Total density of the macrobenthos (ind./m²) .
38. Total biomass of the macrobenthos (g AFDW/m²) .
39. Spatial distribution of the Polychaeta density (ind./m²) .
40. Spatial distribution of the Mollusca density (ind./m²) .
41. Spatial distribution of the Crustacea density (ind./m²) .
42. Spatial distribution of the Echinodermata density (ind./m²) .
43. Spatial distribution of the Polychaeta biomass (g AFDW/m²) .
44. Spatial distribution of the Mollusca biomass (g AFDW/m²) .
45. Spatial distribution of the Crustacea biomass (g AFDW/m²) .
46. Spatial distribution of the Echinodermata biomass (g AFDW/m²) .
47. Spatial distribution (ind./m²) of Aricidea minuta (Polychaeta) .
48. Spatial distribution (ind./m²) of Chaetozone setosa (Polychaeta) .

49. Spatial distribution (ind./m²) of Goniada maculata (Polychaeta).
 50. Spatial distribution (ind./m²) of Lumbrineris latreilli (Polychaeta).
 51. Spatial distribution (ind./m²) of Magelona papillicornis (Polychaeta).
 52. Spatial distribution (ind./m²) of Nephtys cirrosa (Polychaeta).
 53. Spatial distribution (ind./m²) of Nephtys hombergii (Polychaeta).
 54. Spatial distribution (ind./m²) of Pholoe minuta (Polychaeta).
 55. Spatial distribution (ind./m²) of Scoloplos armiger (Polychaeta).
 56. Spatial distribution (ind./m²) of Spiophanes bombyx (Polychaeta).
 57. Spatial distribution (ind./m²) of Spio filicornis (Polychaeta).
 58. Spatial distribution (ind./m²) of Cultellus pellucidus (Mollusca).
 59. Spatial distribution (ind./m²) of Montacuta ferruginosa (Mollusca).
 60. Spatial distribution (ind./m²) of Mysella bidentata (Mollusca).
 61. Spatial distribution (ind./m²) of Natica alderi (Mollusca).
 62. Spatial distribution (ind./m²) of Tellina fabula (Mollusca).
 63. Spatial distribution (ind./m²) of Bathyporeia elegans (Crustacea).
 64. Spatial distribution (ind./m²) of Bathyporeia gulliamsoniana (Crustacea).
 65. Spatial distribution (ind./m²) of Callianassa subterranea (Crustacea).
 66. Spatial distribution (ind./m²) of Harpinia antennaria (Crustacea).
 67. Spatial distribution (ind./m²) of Megaluporus agilis (Crustacea).
 68. Spatial distribution (ind./m²) of Pseudocuma longicornis (Crustacea).
 69. Spatial distribution (ind./m²) of Urothoe brevicornis (Crustacea).
 70. Spatial distribution (ind./m²) of Amphiura filiformis (Echinodermata).
 71. Spatial distribution (ind./m²) of Echinocardium cordatum (Echinodermata).
 72. Spatial distribution (ind./m²) of Echinocyamus pusillus (Echinodermata).
 73. Spatial distribution (ind./m²) of the Nemertinae.
 74. Spatial distribution (ind./m²) of the Phoronida.
 75. TWINSPLAN clustering of the macrobenthos density (ind./m²).
 76. TWINSPLAN dichotomy of the macrobenthos density (ind./m²).
 77. (a-d) Box and whisker plots of some average biotic and abiotic variables
after TWINSPLAN analysis using the data of the macrobenthos density
(ind./m²).
 78. Multidimensional scaling (MDS) ordination of the macrobenthos density
(ind./m²) in relation to the TWINSPLAN clusters of the MILZON-BENTHOS area
1992/93.

IV Meio- and macrobenthos

79. TWINSPLAN clustering of the meiobenthos density (ind./10 cm²) by including
the ICES data of 1986.
 80. TWINSPLAN clustering of the meiobenthos density (ind./10 cm²) including the
MILZON-BENTHOS data of 1988, 1989 and 1991.
 81. TWINSPLAN dichotomy of the meiobenthos density (ind./10 cm²) including the
MILZON-BENTHOS data of 1988, 1989 and 1991.
 82. (a-e) Distribution of the total density (ind./10 cm²), Nematoda,
Copepoda, Gastrotricha and Turbellaria in the total study area of 1988,
1989 and 1991.
 83. TWINSPLAN clustering of the macrobenthos density (ind./m²) including the
ICES data of 1986.
 84. TWINSPLAN clustering of the macrobenthos density (ind./m²) including the
MILZON-BENTHOS I data of 1988, 1989 and 1991.
 85. TWINSPLAN dichotomy of the macrobenthos density (ind./m²) including the
MILZON-BENTHOS I data of 1988, 1989 and 1991.
 86. Axis 1 contours of the meiobenthos TWINSPLAN stations scores.
 87. Axis 1 contours of the macrobenthos TWINSPLAN stations scores.
 88. (a-b) Super-imposition of environmental parameters on the 2-dimensional MDS
ordination of the A. meiobenthic taxa densities (ind./10 cm²)
B. macrobenthic species densities (ind./m²), in relation to the TWINSPLAN
clusters of the MILZON-BENTHOS area 1992/93.




Fig. 1. Survey area of the MILZON-BENTHOS I (1988-1989) and II (1991-1993) projects.




Fig. 2. Location of ICES (1986) and the EXP*BMN project 1992 in the MILZON-BENTHOS area 1992/93.




Fig. 3. Water types and their general boundaries in the North Sea in the summer period.




Fig. 4. Depth (m) recordings, uncorrected for differences due to the tidal cycle. Official 30, 40, 50 and 60 m isobaths are indicated.




Fig. 5. Median grain size (for the fraction 63 - 2000 μm) of the upper 10 cm of the sediment.




Fig. 6. Mud content (%, particles $< 63 \mu\text{m}$) of the upper 10 cm of the sediment.




Fig. 7. Chlorophyll-a content ($\mu\text{g/g}$ dry weight) of the upper 3 cm of the sediment.




Fig. 8. POC content (% C) of the upper 3 cm of the sediment.




Fig. 9. Depth of the (visible) oxidized layer (cm) of the sediment.




Fig. 10. Total number of meiobenthic taxa (Hill (0)).




Fig. 11. Diversity (Hill (1)) of the meiobenthos.




Fig. 12. Diversity (Hill (2)) of the meiobenthos.




Fig. 13. Diversity (Hill (∞))) of the meiobenthos.




Fig. 14. Total density of the meiobenthos (ind./10 cm²).




Fig. 15. Spatial distribution of the Nematoda (ind./10 cm²).




Fig. 16. Spatial distribution of the (total) Copepoda (ind./10 cm²).




Fig. 17a. Spatial distribution of the burrowing Copepoda (ind./10 cm²).




Fig. 17b. Spatial distribution of the interstitial Copepoda (ind./10 cm²).




Fig. 18. Spatial distribution of the Nematoda/Copepoda ratio.




Fig. 19. Spatial distribution of the Gastrotricha (ind./10 cm²).




Fig. 20. Spatial distribution of the Turbellaria (ind./10 cm²).




Fig. 21. Spatial distribution of the Archiannelida (ind./10 cm^2).




Fig. 22. Spatial distribution of the Oligochaeta (ind./10 cm^2).




Fig. 23. Spatial distribution of the Polychaeta (ind./10 cm^2).




Fig. 24. Spatial distribution of the Hydrozoa (ind./10 cm^2).




Fig. 25. Spatial distribution of the Tardigrada (ind./10 cm²).




Fig. 26. Spatial distribution of the Ostracoda (ind./10 cm²).




Fig. 27. Spatial distribution of the Halacarida (ind./10 cm²).




Fig. 28. Spatial distribution of the Kinorhyncha (ind./10 cm²).




Fig. 29. TWINSPLAN clustering of the meiobenthos density (ind./10 cm²).




Fig. 30. Dichotomy of the TWINSPAN cluster found in the MILZON-BENTHOS II area 1992/93 by using the meiobenthos density (ind./10 cm²).
Classes : 0, 5, 10, 20, 50, 100, >1000 ind./10 cm²




Fig. 31a. Box and whisker plots of the average abiotic parameters ($\log(x+1)$ -transformed) after TWINSPLAN analysis using the data of the meiobenthos density (ind./ 10 cm^2) (cf. Figs. 28, 29 and Table 6).




Fig. 31b. Box and whisker plots of some average meiobenthic parameters ($\log(x+1)$ -transformed) after TWIN-SPAN analysis using the data of the meiobenthos density (Ind./10 cm²) (cf. Figs . 28, 29 and Table 6).




Fig. 31c. Box and whisker plots of the average density (log_e(x+1)-transformed) after TWINSPAN analysis using the data of the meiobenthos density (Ind./10 cm²) (cf. Figs . 28, 29 and Table 6).





Fig. 32. Multidimensional scaling (MDS) ordination of the meiobenthos density (ind./10 cm²) in relation to the TWINSPAN clusters of the MILZON-BENTHOS area 1992/93.
Stress = 0.10




Fig. 33. Total number of macrobenthic species (Hill (0)).




Fig. 34. Diversity (Hill (1)) of the macrobenthos.




Fig. 35. Diversity (Hill (2)) of the macrobenthos.




Fig. 36. Diversity (Hill (∞)) of the macrobenthos.




Fig. 37. Total density (ind./m²) of the macrobenthos.








Fig. 38. Total biomass (g AFDW/m²) of the macrobenthos.






Fig. 43. Spatial distribution of the Polychaeta biomass (g AFDW/m^2).




Fig. 44. Spatial distribution of the Mollusca biomass (g AFDW/m^2).




Fig. 45. Spatial distribution of the Crustacea biomass (g AFDW/m^2).




Fig. 46. Spatial distribution of the Echinodermata biomass (g AFDW/m^2).




Fig. 47. Spatial distribution (ind./m²) of *Aricidea minuta* (Polychaeta).




Fig. 48. Spatial distribution (ind./m²) of *Chaetozone setosa* (Polychaeta).




Fig. 49. Spatial distribution (ind./m²) of *Goniada maculata* (Polychaeta).




Fig. 50. Spatial distribution (ind./m²) of *Lumbrineris latreilli* (Polychaeta).




Fig. 51. Spatial distribution (ind./m^2) of *Magelona papillicornis* (Polychaeta).




Fig. 52. Spatial distribution (ind./m^2) of *Nephtys cirrosa* (Polychaeta).




Fig. 53. Spatial distribution (ind./m^2) of *Nephtys hombergii* (Polychaeta).




Fig. 54. Spatial distribution (ind./m^2) of *Pholoe minuta* (Polychaeta).




Fig. 55. Spatial distribution (ind./m^2) of *Scoloplos armiger* (Polychaeta).




Fig. 56. Spatial distribution (ind./m^2) of *Spiophanes bombyx* (Polychaeta).




Fig. 57. Spatial distribution (ind./m^2) of *Spio filicornis* (Polychaeta).




Fig. 58. Spatial distribution (ind./m^2) of *Cultellus pellucidus* (Mollusca).




Fig. 59. Spatial distribution (ind./m²) of *Montacuta ferruginea* (Mollusca).




Fig. 60. Spatial distribution (ind./m²) of *Mysella bidentata* (Mollusca).




Fig. 61. Spatial distribution (ind./m²) of *Natica alderi* (Mollusca).




Fig. 62. Spatial distribution (ind./m²) of *Tellina fabula* (Mollusca).




Fig. 63. Spatial distribution (ind./m²) of *Bathyporeia elegans* (Crustacea).




Fig. 64. Spatial distribution (ind./m²) of *Bathyporeia guilliemoniana* (Crustacea).




Fig. 65. Spatial distribution (ind./m²) of *Callianassa subterranea* (Crustacea).




Fig. 66. Spatial distribution (ind./m²) of *Harpinia antennaria* (Crustacea).




Fig. 67. Spatial distribution (ind./m²) of *Megalopus agilis* (Crustacea).




Fig. 68. Spatial distribution (ind./m²) of *Pseudocuma longicornis* (Crustacea).




Fig. 69. Spatial distribution (ind./m²) of *Urothoe brevicornis* (Crustacea).




Fig. 70. Spatial distribution (ind./m²) of *Amphiura filiformis* (Echinodermata).




Fig. 71. Spatial distribution (ind./m²) of *Echinocardium cordatum* (Echinodermata).




Fig. 72. Spatial distribution (ind./m²) of *Echinocystus pusillus* (Echinodermata).




Fig. 73. Spatial distribution (ind./m²) of the Nemertinae.




Fig. 74. Spatial distribution (ind./m²) of the Phoronida.




Fig. 75. TWINSPAN clustering of the macrobenthos density (ind./m²).




Fig. 76. Dichotomy of the TWINSPAN cluster found in the MILZON-BENTHOS II area 1992/1993 by using the macrobenthos density (ind./m²).
Classes : 0, 15, 30, 75, 150, 350, 1000 ind./m²




Fig. 77a. Box and whisker plots of the average abiotic parameters ($\log(x+1)$ -transformed) after TWINSPAN analysis using the data of the macrobenthos density (ind./m^2) (cf. Figs. 45, 46 and Table 11).




Fig. 77b. Box and whisker plots of some average macrobenthic parameters ($\log(x+1)$ -transformed) after TWINSPLAN analysis using the data of the macrobenthos density (ind./m²) (cf. Figs. 45, 46 and Table 11).




Fig. 77c. Box and whisker plots of the average density ($\log(x+1)$ -transformed) after TWINSPAN analysis using the data of the macrobenthos density (ind./m²) (cf. Figs. 45, 46 and Table 11).




Fig. 77d. Box and whisker plots of the average biomass (log(x+1)-transformed) after TWINSPAN analysis using the data of the macrobenthos density (ind./m²) (cf. Figs. 45, 46 and Table 11).




Fig. 78. Multidimensional scaling (MDS) ordination of the macrobenthos density (ind./m²) in relation to the TWINSPAN clusters of the MILZON-BENTHOS area 1992/93.
Stress = 0.16




Fig. 79. TWINSPAN clustering of the meiobenthos density (ind./10 cm²) including the ICES data of 1986.




Fig. 80. TWINSPAN clustering of the meiobenthos density (ind./10 cm²) including the MILZON-BENTHOS I + II data of 1988/89 and 1991.




Fig. 81. Dichotomy of the TWINSPAN cluster found in the MILZON-BENTHOS I + II area 1988-1993 by using the meiobenthos density (ind./10 cm²).
 Classes : 0, 10, 150, 350, 700, 1000, 2000, 3000, >3000
 ind./10 cm²




Fig. 82a. Spatial distribution of the total meiobenthic density (ind./10 cm²) including the MILZON-BENTHOS I + II data of 1988/89 and 1991.




Fig. 82b. Spatial distribution of the Nematoda (Ind./10 cm²) including the MILZON-BENTHOS I + II data of 1988/89 and 1991.




Fig. 82c. Spatial distribution of the Copepoda (ind./10 cm²) including the MILZON-BENTHOS I + II data of 1988/89 and 1991.




Fig. 82d. Spatial distribution of the Gastrotricha (ind./10 cm²) including the MILZON-BENTHOS I + II data of 1988/89 and 1991.




Fig. 82e. Spatial distribution of the Turbellaria (ind./10 cm²) including the MILZON-BENTHOS I + II data of 1988/89 and 1991.




Fig. 83. TWINSPAN clustering of the macrobenthos density (ind./m²) including the ICES data 1986.




Fig. 84. TWINSPAN clustering of the macrobenthos density (ind./m²) including the MILZON-BENTHOS I + II data of 1988-1990.




Fig. 85. Dichotomy of the TWINSPAN cluster found in the MILZON-BENTHOS I + II area 1988-1993 by using the macrobenthos density (ind./m²).

Classes : 0, 15, 30, 75, 150, 350 ind./m²




Fig. 86. Axis 1 contours of the meiobenthos.




Fig. 87. Axis 1 contours of the macrobenthos.




Fig. 88a. Super-imposition of scaled sedimental parameters on the 2-dimensional MDS ordination of the meiobenthic taxa (ind./10 cm²).




Fig. 88b. Super-imposition of scaled sedimental parameters on the 2-dimension MDS ordination of the macrobenthic species (ind./m²).

Appendices

Appendix 1-1. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on all MILZON-BENTHOS II 1992 locations.

Station	1-1	1-2	1-3	1-4	1-5	2-1	2-2	2-3	2-4	2-5	3-1	3-2	3-3	3-4	3-5
Permanent															
Nematoda	1047	1505	358	1124	1105	531	660	327	378	513	599	787	937	1730	623
Copepoda total	155	1067	322	378	213	412	92	17	23	160	106	38	194	226	1
burrowing	14	.	1	5	11	.	7	4	.	10	4	5	5	2	1
interstitial	141	1067	321	373	202	412	85	13	23	150	102	33	189	224	.
Gastrotricha	288	157	70	99	109	68	81	50	145	74	112	150	125	167	7
Turbellaria	22	51	11	17	28	19	29	16	37	37	27	63	30	72	5
Archiannelida	.	2	.	2	4	1	8	.	2	1	1	.	.	5	.
Oligochaeta	.	.	.	1	1	2	.	3	.
Polychaeta	2	18	3	10	9	3	12	.	1	4	4	4	1	2	.
Hydrozoa	2	58	9	12	12	.	6	2	18	.	3	6	10	18	.
Tardigrada	95	296	24	39	68	4	2	.	5	4	13	3	48	60	.
Ostracoda	.	63	5	7	4	.	1
Halacarida	.	2
Temporary															
Nauplii	222	1511	96	160	106	24	10	8	9	35	37	10	55	54	15
Bivalvia	1	2	.	.	1	.	2	1	.
Foraminifera	27	91	1	10	21	5	1	.	1	10	5	.	3	9	.
Isopoda
Cumacea
Nemertina
Tanaidacea
Amphipoda
Echinoidea
Ophiuroidea
Cnidaria other	.	.	1
Bryozoa
Taxa permanent	7	10	8	10	10	7	9	5	8	7	8	8	7	9	4
Taxa temporary	3	3	3	2	3	2	3	1	2	2	2	1	2	3	1
Taxa total	10	13	11	12	13	9	12	6	10	9	10	9	9	12	5

Appendix 1-2. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on all MILZON-BENTHOS II 1992 locations.

Station	4-1	4-2	4-3	4-4	4-5	5-1	5-2	5-3	5-4	5-5	6-1	6-2	6-3	6-4	6-5	6-6
<u>Permanent</u>																
Nematoda	1443	576	1400	1359	978	832	1702	1820	2807	794	888	547	1619	812	1047	424
Copepoda	161	100	10	10	72	1	5	23	443	276	253	65	193	56	373	117
burrowing	6	7	3	6	5	1	5	23	5	.	6	4	14	2	3	13
interstitial	155	93	7	4	67	.	.	.	438	276	247	61	179	54	370	104
Gastropoda	18	43	38	39	7	11	26	4	51	66	35	24	31	34	43	21
Turbellaria	16	11	18	24	18	36	10	6	31	22	20	48	20	11	16	20
Archannelida	.	5	10	10	.	.	.	2	10	.
Oligochaeta	12	9	1
Polychaeta	2	3	.	1	7	.	1	1	8	9	2	3	7	2	3	3
Hydrozoa	4	9	2	.	8	.	.	.	24	8	.	.	1	2	6	1
Tardigrada	1	18	4	14	7	1	.	1	136	30	14	15	34	12	20	13
Ostracoda	.	3	2	.	1	.	.	.	15	3	1	.	6	1	5	.
Halacarida	1
<u>Temporary</u>																
Nauplii	29	26	47	38	26	2	16	95	130	141	82	8	142	16	40	80
Bivalvia	.	9	.	.	.	3	6	1	2	1	3	1	1	.	3	.
Foraminifera	4	39	2	5	14	.	4	.	18	18	10	31	14	14	60	9
Isopoda
Cumacea	2
Nemertina	2
Tanaidacea
Amphipoda	.	.	.	1	.	.	1
Echinoidea
Ophiuroidea	.	.	.	1	1
Cnidaria other	.	1	.	1
Bryozoa
Taxa permanent	7	9	7	6	8	5	5	7	10	11	8	6	8	9	9	7
Taxa temporary	3	4	2	5	2	2	4	2	3	3	4	3	3	2	3	3
Taxa total	10	13	9	11	10	7	9	9	13	14	12	9	11	11	12	10

Appendix 1-3. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on all MILZON-BENTHOS II 1992 locations.

Station	7-1	7-2	7-3	7-4	7-5	7-6	8-1	8-2	8-3	8-4	8-5	8-6
Permanent												
Nematoda	372	1032	493	487	913	487	393	705	551	1258	1020	758
Copepoda	169	538	30	66	452	90	297	16	144	33	187	242
burrowing	.	7	9	30	8	3	6	2	7	8	18	10
interstitial	169	531	21	36	444	87	291	14	137	25	169	232
Gastrotricha	135	43	59	15	73	5	76	72	32	4	87	226
Turbellaria	23	11	18	12	47	5	23	26	13	5	18	72
Archannelida	1	1	1	.	1	.	.	12
Oligochaeta	.	1	.	1	.	1	1
Polychaeta	4	9	.	9	1	4	1	2	1	1	.	4
Hydrozoa	26	5	.	2	3	4	18	.	.	2	.	6
Tardigrada	15	77	1	41	17	85	7	3	20	9	14	24
Ostracoda	7	43	.	2	16	10	1	.	.	4	.	11
Halacarida	.	2	.	.	1	1
Temporary												
Nauplii	153	188	18	58	115	65	50	3	29	44	56	44
Bivalvia	.	.	.	1	.	1	1	.	.	7	1	1
Foraminifera	14	5	17	3	31	8	5	1	2	4	10	16
Isopoda	1
Cumacea	1	2	1	.	.	.
Nemertina
Tanaidacea
Amphipoda
Echinidea
Ophiuroidea	2	.	.	.	1	1	.	.
Cnidaria other	1	.	.	1	1	.
Bryozoa	1
Taxa permanent	9	10	5	9	9	11	9	6	7	8	5	10
Taxa temporary	5	3	2	4	3	4	3	2	3	4	4	4
Taxa total	14	13	7	13	12	15	12	8	10	12	9	14

Appendix 1-4. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on all MILZON-BENTHOS II 1992 locations.

Station	9-1	9-2	9-3	9-4	9-5	10-1	10-2	10-3	10-4	10-5
<u>Permanent</u>										
Nematoda	833	1026	567	1311	956	1548	823	1257	1062	1587
Copepoda	271	381	37	32	9	4	16	19	14	107
burrowing	11	1	6	6	1	4	.	18	13	6
interstitial	260	380	31	26	8	.	16	1	1	101
Gastrotricha	23	77	55	44	58	.	33	5	3	79
Turbellaria	23	32	50	8	34	3	14	3	3	30
Archannelida	.	.	1	.	.	.	1	.	.	1
Oligochaeta	.	1	10
Polychaeta	2	3	6	.	.	4
Hydrozoa	.	7	.	2	2	1	4	.	.	.
Tardigrada	34	38	8	75	2	1	9	.	8	39
Ostracoda	.	.	1	1	5
Halacarida
<u>Temporary</u>										
Nauplii	40	82	35	46	9	17	82	33	45	42
Bivalvia	.	.	1	.	.	1	2	6	5	4
Foraminifera	2	34	8	5	1	9	22	15	2	7
Isopoda	.	1
Cumacea	.	2	.	1
Nemertina	.	3	.	2	1
Tanaidacea	1	.	.	1	.	1
Amphipoda
Echinoidea	.	.	.	1
Ophiuroidea
Cnidaria other	1	5
Bryozoa
Taxa permanent	6	8	7	6	6	5	8	4	6	9
Taxa temporary	3	5	3	6	3	4	3	3	4	4
Taxa total	9	13	10	12	9	9	11	7	10	13

Appendix 1-5. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on all MILZON-BENTHOS II 1993 locations.

Station	11-1	11-2	11-3	12-1	12-2	12-3	13-1	13-2	13-3	14-1	14-2	14-3	14-4	15-1	15-2	15-3
Permanent																
Nematoda	3921	1967	3039	2111	1284	2833	1058	1599	1297	1415	4644	2827	2740	929	2738	2179
Copepoda	18	25	5	11	51	41	8	278	21	59	238	85	58	32	80	109
burrowing	17	25	5	11	48	41	8	2	19	56	238	85	58	27	80	109
interstitial	1	.	.	.	3	.	.	276	2	3	.	.	.	5	.	.
Gastrotricha	.	26	1	9	.	.	5	115	293	7	.	1	.	17	1	.
Turbellaria	5	11	4	18	9	8	13	23	77	5	22	4	6	35	1	4
Archiamelida	10
Oligochaeta	.	1	.	.	.	1	.	12	1	.	.
Polychaeta	2	.	1	1	2	1	7	27	7	.	3	.	1	.	3	7
Hydrozoa	.	1	.	.	1	.	.	8
Tardigrada	14	1
Ostracoda	1	6	.	.	.	5	.	12	1	.	7	4	.	.	5	4
Halacarida	1
Rotifera	14
Kinorhyncha	3	1	36	13	6	.	7	3
Friapulida
Temporary																
Nauplii	40	76	15	16	26	50	20	126	59	34	779	30	167	40	119	343
Bivalvia	5	9	.	3	4	.	1	8	1	5	1	4	.	8	1	9
Foraminifera	13	40	7	18	15	28	.	27	17	.	104	64	56	43	131	69
Isopoda	.	.	2	1
Cumacea	.	1	.	.	1
Tanaidacea	1
Amphipoda	1
Echinidea	1	2
Ophiuroidea	1	1	.	.	1
Gastropoda	20
Cnidaria other	6	.	6	.	.	1	2	.	18	.	3	.
Bryozoa	1	10	.	.	2	4	2	.
Taxa permanent	6	7	5	5	5	7	5	12	7	4	6	6	5	5	7	6
Taxa temporary	5	5	4	3	5	4	3	5	6	2	5	5	3	3	5	3
Taxa total	11	12	9	8	10	11	8	17	13	6	11	11	8	8	12	9

Appendix 1-6. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on all MILZON-BENTHOS II 1993 locations.

Station	16-1	16-2	16-3	16-4	17-1	17-2	17-3	18-1	18-2	18-3	19-1	19-2	19-3
<u>Permanent</u>													
Nematoda	4950	5225	6711	3738	1953	2318	1007	4411	867	1729	1005	3673	2320
Copepoda	56	48	68	60	361	36	27	41	44	33	28	29	43
burrowing	54	46	68	43	58	32	25	40	19	31	28	26	43
interstitial	2	2	.	17	303	4	2	1	25	2	.	3	.
Gastroparia	.	31	.	.	68	1	.	1	58	2	.	7	.
Turbellaria	12	12	3	2	15	15	3	13	22	3	4	.	3
Archannelida	15
Oligochaeta	.	1	.	.	7
Polychaeta	2	1	.	6	7	5	2	1	3	1	.	1	1
Hydrozoa	8	.	.	.	3
Tardigrada	.	1	.	.	120	2	.	.	1	.	.	5	.
Ostracoda	9	8	6	3	19	2	.	2	2	.	2	3	2
Halacarida	6	.	1
Rotifera	.	.	1	1	3	5
Kinorhyncha	2	9	25	18	19
Priapulida	.	.	1	.	.	1
<u>Temporary</u>													
Nauplii	33	238	160	53	524	51	15	57	18	22	59	56	53
Bivalvia	1	6	5	4	8	6	4	4	2	7	9	8	8
Foraminifera	69	162	32	36	88	149	.	76	5	18	60	228	178
Isopoda
Cumacea	1	.	.
Tanaidacea
Amphipoda
Echinoidea	1	.	.	.	3	2	.	.	.
Ophiuroidea	1	1	2	1	.	1	.	1	.	.	1	.	1
Gastropoda
Cnidaria other	3	15	.	9	.	.	.	7	2
Bryozoa	.	3	.	6	.	.	3	2
Taxa permanent	6	9	7	6	12	8	5	6	8	5	5	7	6
Taxa temporary	5	6	4	6	4	4	3	6	4	4	5	3	5
Taxa total	11	15	11	12	16	12	8	12	12	9	10	10	11

Appendix 2-1. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 1).

Density (ind./m ²) :						Biomass (g AFDW/m ²) :					
Station	11	12	13	14	15	11	12	13	14	15	
Compartment	1	1	1	1	1	1	1	1	1	1	
Number	1	2	3	4	5	1	2	3	4	5	
Date	280492	280492	280492	280492	280492	280492	280492	280492	280492	280492	
Species	15	13	11	11	14	15	13	11	11	14	
Density	746.13	570.57	497.42	468.16	570.57	746.13	570.57	497.42	468.16	570.57	
Biomass	1.79	1.64	1.40	2.48	26.82	1.79	1.64	1.40	2.48	26.82	
Depth	32.8	33.5	26.7	31.7	33.6	32.8	33.5	26.7	31.7	33.6	
<u>Polychaeta (12 species) :</u>											
ANAIROSE	14.630219	
AONIPAU	.	14.630015	.	.	.	
ARICMINU	.	29.26	29.26	102.41	14.63	.	.0029	.0029	.0205	.0029	
CHAESETO	29.260117	
GONIMACU	.	14.63	14.630044	.0293	.	.	
MAGEPAPI	.	29.26	29.26	58.52	14.63	.	.7359	.3555	.7783	.0293	
NEPHCIRR	146.30	73.15	29.26	102.41	87.78	.4184	.4843	.2619	.1053	.1565	
NEPHLONG	29.26	.	29.26	29.26	.	.1024	.	.	.2824	.	
SCOLARMI	29.26	.	29.26	43.89	29.26	.0614	.	.6306	1.2330	.1009	
SCOLSQUA	14.63	14.63	.	.	43.89	.4155	.1434	.	.	1.5800	
SPIOBOMB	117.04	131.67	14.63	.	160.93	.3628	.1258	.0146	.	.2502	
SPIOFILI	73.15	131.67	204.82	.	.	.0146	.0688	.0410	.	.	
<u>Mollusca (2 species) :</u>											
MONTFERR	.	.	.	14.630073	.	
NATIALDE	14.63	14.630073	.0059	.	.	.	
<u>Crustacea (8 species) :</u>											
BATHGUIL	.	43.89	14.63	14.63	43.89	.	.0263	.0044	.0073	.0132	
GASTSPIN	.	43.89	14.63	29.26	.	.	.0263	.0088	.0176	.	
MEGAAGIL	58.52	14.63	43.89	14.63	29.26	.0176	.0044	.0132	.0044	.0088	
PONTALTA	14.630044	
PSEULONG	.	.	.	43.89	14.630088	.0029	
THIASCUT	29.262824	
UROTREV	160.93	.	.	.	43.89	.08050219	
UROTPOSE	.	.	73.15	.	14.63	.	.	.0366	.	.0044	
<u>Echinodermata (2 species) :</u>											
ECHICORD	14.63	.	.	.	29.26	.2165	.	.	.	24.1219	
OPHIALBI	14.630483	
<u>Rest (1 species) :</u>											
NEMERTIN	14.63	14.63	.	14.63	29.26	.0073	.0073	.	.0117	.2472	

Appendix 2-2. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 2).

Density (ind./m²) :						Biomass (g AFDW/m²) :					
Station	21	22	23	24	25	21	22	23	24	25	
Compartment	2	2	2	2	2	2	2	2	2	2	
Number	6	7	8	9	10	6	7	8	9	10	
Date	280492	280492	060592	060592	080592	280492	280492	060592	060592	080592	
Species	14	14	11	11	11	14	14	11	11	11	
Density	2399.32	819.28	461.88	256.60	577.35	2399.32	819.28	461.88	256.60	577.35	
Biomass	5.71	47.71	1.48	1.09	1.30	5.71	47.71	1.48	1.09	1.30	
Depth	32.5	30.9	29.7	26.1	31.2	32.5	30.9	29.7	26.1	31.2	
<u>Polychaeta (13 species) :</u>											
ANAIROSE	29.260585	
ARICMINU	117.04	14.63	12.83	51.32	115.47	.0293	.0029	.0026	.0103	.0231	
CHAESETO	14.630029	
EXOGHEBE	.	.	12.830077	.	.	
GONIMACU	.	.	.	12.83	12.830398	.0898	
MAGEPAPI	14.63	73.15	.	64.15	25.66	.0102	.2414	.	.7647	.3028	
NEPHCIRR	87.78	117.04	205.28	12.83	128.30	.1448	.1419	.2925	.0128	.1514	
NEPHLONG	29.26	.	.	.	12.83	1.61374478	
SCOLARMI	29.26	73.15	38.49	12.83	.	.0146	.9027	.5389	.1347	.	
SCOLSQUA	.	87.78	.	.	25.66	.	2.2091	.	.	.0629	
SPIOBOMB	1887.27	117.04	.	12.83	89.81	3.2230	.3321	.	.0731	.0180	
SPIOFILI	.	.	.	12.83	64.150026	.0128	
STHELIMI	14.635208	
<u>Mollusca (1 species) :</u>											
TELLFABU	.	.	12.83	12.830077	.0026	.	
<u>Crustacea (8 species) :</u>											
BATHELEG	14.63	14.630044	.0029	.	.	.	
BATHGUIL	.	14.63	89.81	.	25.66	.	.0059	.1103	.	.0128	
CALLSUBT	.	.	12.834980	.	.	
GASTSPIN	29.26	14.63	12.83	25.66	.	.0176	.0088	.0077	.0192	.	
LEUCLILL	.	.	12.830044	.	.	
MEGAAGIL	58.52	29.26	25.66	25.66	.	.0176	.0088	.0077	.0077	.	
PSEULONG	.	14.63	25.660029	.0051	.	.	
UROTBREV	29.26	146.30	.	.	51.32	.0146	.0995	.	.	.0205	
<u>Echinodermata (1 species) :</u>											
ECHICORD	.	73.15	43.7481	.	.	.	
<u>Rest (2 species) :</u>											
HYDROZOA	.	29.260044	.	.	.	
NEMERTIN	43.89	.	.	12.83	25.66	.0366	.	.	.0231	.1552	

Appendix 2-3. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 3).

Density (ind./m ²) :						Biomass (g AFDW/m ²) :					
Station	31	32	33	34	35		31	32	33	34	35
Compartment	3	3	3	3	3		3	3	3	3	3
Number	12	13	14	15	16		12	13	14	15	16
Date	060592	060592	060592	060592	060592		060592	060592	060592	060592	060592
Species	14	12	15	17	28		14	12	15	17	28
Density	667.16	731.31	910.93	1154.70	3399.95		667.16	731.31	910.93	1154.70	3399.95
Biomass	3.98	1.34	12.46	3.79	39.43		3.98	1.34	12.46	3.79	39.43
Depth	25.5	29.7	27.4	28.0	35.1		25.5	29.7	27.4	28.0	35.1
<u>Polychaeta (16 species) :</u>											
ANAIROE	25.66	9635
ARICMINU	269.43	25.66	38.49	25.66	.	.0539	.0051	.0077	.0051	.	.
CHAESETO	.	.	.	25.66	89.810154	.0269
EXOGHEBE	.	12.83	12.830026	.0026	.	.	.
GONIMACU	.	.	12.83	12.83	38.49	.	.	.0308	.0462	.	.4144
HARMLUNU	76.981450
LANICONC	166.7914.6467
MAGEPAPI	12.83	12.83	25.66	12.83	12.83	.0885	.0321	.1514	.1180	.	.0128
NEPHCIRR	141.13	205.28	230.94	89.81	38.49	.1950	.2925	.2476	.1925	.	.0385
NEPHHOMB	12.832.8842
NEPHLONG	12.83	.	.	12.83	.	1.09570244	.
POECSERP	38.490924
SCOLARMI	.	64.15	115.47	551.69	256.60	.	.5530	1.0161	2.4480	.	.6146
SCOLSQUA	12.83	.	.	38.49	12.83	.01927480	.0321
SPIOBOMB	12.83	89.81	218.11	12.83	1308.66	.1296	.0860	.4349	.0359	.	.1.9656
SPIOFILI	12.83	.	12.83	38.49	.	.0192	.	.0026	.0077	.	.
<u>Mollusca (4 species) :</u>											
DONAVITT	38.49	2.3286
MONTFERR	51.320192
NATIALDE	.	.	.	25.66	76.980128	.	.3759
TELLFABU	51.322.2940
<u>Crustacea (11 species) :</u>											
ATYLSWAM	12.830051
BATHELEG	38.490115
BATHGUIL	25.66	38.49	51.32	230.94	295.09	.0154	.0192	.0205	.1091	.	.0885
LEUCLILL	.	.	12.83	.	51.32	.	.	.0051	.	.	.0205
MEGAAGIL	25.66	12.83	.	12.83	38.49	.0077	.0038	.	.0038	.	.0115
ORCHHUMI	25.660128
PERILONG	12.830038
PONTALTA	.	12.830038
PSEULONG	38.49	.	.	25.66	25.66	.0115	.	.	.0026	.	.0051
UROTBREV	12.83	192.45	76.98	.	.	.0038	.1552	.0308	.	.	.
UROTPOSE	.	.	25.66	.	449.05	.	.	.0077	.	.	.1347
<u>Echinodermata (3 species) :</u>											
ECHICORD	.	.	25.66	.	25.66	.	.	10.4937	.	.	.13.9975
ECHIPUSI	.	25.66	.	12.83	76.98	.	.0398	.	.0128	.	.0411
OPHIALBI	.	.	.	12.83	38.490013	.	.4940
<u>Rest (2 species) :</u>											
HYDROZOA	.	.	12.83	.	12.83	.	.	.0013	.	.	.0026
NEMERTIN	25.66	38.49	38.49	12.83	64.15	.0026	.1488	.0026	.0038	.	.0834

Appendix 2-4. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 4).

Density (ind./m²) :					Biomass (g AFDW/m²) :					
Station	41	42	43	44	45	41	42	43	44	45
Compartment	4	4	4	4	4	4	4	4	4	4
Number	18	19	20	21	22	18	19	20	21	22
Date	070592	060592	070592	070592	070592	070592	060592	070592	070592	070592
Species	17	19	19	26	19	17	19	19	26	19
Density	1667.90	1154.70	2129.78	1809.03	1295.83	1667.90	1154.70	2129.78	1809.03	1295.83
Biomass	45.49	3.01	5.16	18.10	1.47	45.49	3.01	5.16	18.10	1.47
Depth	30.4	31.4	28.4	28.2	31.2	30.4	31.4	28.4	28.2	31.2
<u>Polychaeta (19 species) :</u>										
ANAIROSE	.	.	12.83	12.830192	.0103	
ARICMINU	12.83	51.32	.	.	51.32	.0013	.0154	.	.0321	
CHAESETO	.	.	25.66	128.300321	.0205	
ETEOLONG	12.830128	.	.	.	
EUMISANG	.	12.830128	.	.	
GONIMACU	.	12.830885	.	.	
HARMLUNU	.	12.83	25.66	38.49	.	.	.0192	.0257	.0500	
LANICONC	51.32	12.83	51.32	89.81	.	.5145	.7826	1.0572	3.0587	
MAGEFAPI	51.32	12.83	141.13	102.64	.	.2014	.0962	.6915	.5312	
NEPHCIRR	.	25.66	64.15	89.81	102.64	.	.0218	.1694	.1168	
NEPHHOMB	.	.	.	12.83	1.7936	
NOTOLATE	.	.	.	12.832053	
POECSERP	.	.	.	38.490321	
SCOLARMI	51.32	205.28	.	12.83	192.45	.0334	1.5858	.	.7287	
SCOLSQUA	.	12.830718	.	.	
SIGAMATH	.	12.832925	.	
SPIOBOMB	166.79	.	192.45	603.01	615.84	.0654	.	.1617	.0282	
SPIOFILI	38.490128	
STREWEBS	.	.	.	12.830013	.	
<u>Mollusca (3 species) :</u>										
DONAVITT	64.15	12.83	12.83	.	.	9.5891	.0064	1.9925	.	
NATIALDE	12.83	12.83	25.66	12.83	12.83	.0064	.0064	.0128	.0064	
TELLFABU	.	25.66	115.47	141.13	.	.	.0064	.1694	1.0867	
<u>Crustacea (14 species) :</u>										
APHEOVAL	.	.	12.830051	
ATYLEG	.	.	.	12.830051	
BATHELEG	.	115.47	538.86	51.32	25.66	.	.0462	.1078	.0257	
BATHGUIL	166.79	513.20	487.54	76.98	76.98	.1001	.2053	.2951	.0552	
DIASRATH	.	.	12.830128	
HIPPEND	.	12.83	12.83	12.830051	.0051	
MEGAAGIL	12.83	38.49	12.83	25.66	38.49	.0038	.0115	.0038	.0077	
PERILONG	12.83	.	.	25.66	.	.0051	.	.	.	
PONTALTA	12.83	.	102.64	.	12.83	.0038	.	.0308	.	
PROCNOUN	12.380505	
PSEULONG	12.83	12.83	.	25.66	12.83	.0013	.0026	.	.0103	
THIASCUT	25.664952	.	.	.	
UROTBREV	.	25.660154	.	.	
UROPOSE	795.46	.	230.94	166.79	.	.0282	.	.0398	.1334	
<u>Echinodermata (3 species) :</u>										
ECHICORD	141.13	.	.	12.83	.	34.3523	.	.	10.0651	
ECHIPUSI	.	12.83	51.32	25.66	12.83	.	.0026	.0449	.0321	
OPHIALBI	.	12.830026	.	.	.0026	
<u>Rest (3 species) :</u>										
ENTEROPN	.	.	.	12.830013	
NEMERTIN	64.15	25.66	12.83	51.32	25.66	.0757	.0231	.0128	.0654	
PHORONID	12.830192	

Appendix 2-5. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 5).

Density (ind.m ⁻²) :						Biomass (g AFDW/m ⁻²) :					
Station	51	52	53	54	55	51	52	53	54	55	
Compartment	5	5	5	5	5	5	5	5	5	5	
Number	23	24	25	26	27	23	24	25	26	27	
Date	070592	070592	070592	070592	070592	070592	070592	070592	070592	070592	
Species	23	24	23	21	16	23	24	23	21	16	
Density	1398.47	1141.87	962.25	718.48	603.01	1398.47	1141.87	962.25	718.48	603.01	
Biomass	14.82	26.96	14.50	3.23	1.94	14.82	26.96	14.50	3.23	1.94	
Depth	26.1	27.6	28.3	29.8	27.4	26.1	27.6	28.3	29.8	27.4	
<u>Polychaeta (25 species) :</u>											
ANAIROSE	.	12.83	.	12.83	.	.	.0064	.	.0051	.	
ACNIPAU	.	.	.	12.830038	.	
ARICMINU	.	.	.	12.83	25.660064	.0128	
CHAESETO	12.83	102.64	12.83	.	.	.0064	.0026	.0038	.	.	
ETEOLONG	.	.	.	12.830128	.	
EUMISANG	.	25.660192	.	.	.	
EXOGNAID	.	.	.	76.98	76.980077	.0077	
GONIMACU	12.83	25.66	12.83	25.66	12.83	.0744	.1706	.0192	.0269	.0128	
HARMLONG	.	.	.	12.83	12.830128	.0257	
HARMLUNU	.	25.660488	.	.	.	
HESIAUGE	.	.	.	25.660026	.	
LANICONC	.	51.32	1.2650	.	.	.	
MAGEPAPI	218.11	166.79	128.30	.	38.49	.8352	.2758	.3233	.	.5145	
NEPHCIRR	38.49	.	.	38.49	64.15	.0488	.	.	.1565	.0872	
NEPHHOMB	12.83	.	12.83	.	.	.1963	.	.1232	.	.	
NOTOLATE	.	25.66	25.668237	.6274	.	.	
OPHELIMA	.	.	.	25.66	25.660385	.0192	
POECSERP	.	.	230.940872	.	
SCOLARMI	12.83	.	51.32	12.83	12.83	.0231	.	.0488	.0115	.0180	
SCOLEBONN	38.49	1.0880	
SCOLSQUA	.	25.660192	.	.	.	
SIGAMATH	12.83	38.49	12.83	.	.	.2361	.5453	.0462	.	.	
SPIOBOMB	166.79	.	102.64	25.66	76.98	.0629	.	.0642	.0064	.0064	
SPIOFILI	.	.	.	102.64	128.300180	.0128	
STREWEBS	12.830013	
<u>Mollusca (8 species) :</u>											
DONAVITT	.	.	.	12.838622	.	
DOSILUPI	.	.	.	12.83	1.8398	.	
MONTFERR	38.49	179.62	51.32	.	.	.0629	.0847	.0103	.	.	
MYSEBIDE	.	12.830026	.	.	.	
NATIALDE	.	12.83	38.490026	.0192	.	.	
SPISSUBT	12.83	6.3881	
TELLFABU	38.49	64.15	25.66	.	.	.1129	.0282	.0334	.	.	
VENUSTRI	.	.	12.83	2.9188	.	.	
<u>Crustacea (16 species) :</u>											
APHEOVAL	.	12.830077	.	.	.	
BATHELEG	51.32	38.49	51.32	.	.	.0205	.0154	.0205	.	.	
BATHGUIL	577.35	25.66	12.83	.	.	.2502	.0128	.0064	.	.	
CALLSUBT	.	.	12.833759	.	.	
CAPRELLI	.	25.660077	.	.	.	
DIASRATH	12.83	.	.	12.83	.	.0077	.	.	.0205	.	
EBALCRAN	.	.	12.830154	.	.	
HIPPEND	12.830051	
LEUCLILL	.	12.83	25.660051	.0205	.	.	
MEGAAGIL	.	.	.	153.96	25.660616	.0103	
PERILONG	12.830038	
PINNFISU	12.83	.	12.83	.	.	.0192	.	.0192	.	.	
PONTALTA	64.15	38.49	12.83	.	12.83	.0192	.0115	.0038	.	.0038	
PSEULONG	12.83	.	.	12.83	12.83	.0013	.	.	.0051	.0038	
SIPHKROY	.	.	12.830038	.	.	
UROTPOSE	25.66	141.130128	.0847	.	.	.	
<u>Echinodermata (3 species) :</u>											
ECHICORD	12.83	51.32	12.83	.	.	5.9416	23.5046	9.6187	.	.	
ECHIPUSI	12.83	.	.	76.98	.	.0026	.	.	.0372	.	
OPHITEXT	12.83	12.834888	.0064	.	.	.	
<u>Rest (2 species) :</u>											
NEMERTIN	.	12.83	76.98	12.83	25.66	.	.0051	.0949	.0834	.1193	
OLIGOCHA	.	.	38.490090	.	

Appendix 2-6. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 6).

Density (ind./m²) :						Biomass (g AFDW/m²) :						
Station	61	62	63	64	65	66	61	62	63	64	65	66
Compartment	6	6	6	6	6	6	6	6	6	6	6	6
Number	28	29	30	31	32	33	28	29	30	31	32	33
Date	290492	180592	290492	180592	210592	290492	290492	180592	290492	180592	210592	290492
Species	16	11	14	16	11	15	16	11	14	16	11	15
Density	804.65	603.01	907.06	538.86	500.37	760.76	804.65	603.01	907.06	538.86	500.37	760.76
Biomass	20.75	9.68	94.14	11.35	13.36	14.16	20.75	9.68	94.14	11.35	13.36	14.16
Depth	32.4	34.0	34.5	41.2	34.7	40.5	32.4	34.0	34.5	41.2	34.7	40.5
<u>Polychaeta (13 species) :</u>												
ARICMINU	14.63	.	14.63	12.83	51.32	14.63	.0044	.	.0059	.0064	.0205	.0073
ETEOFFLAV	.	12.830718
HARMLONG	.	29.26	.	12.83	14.63	.	.	.0219	.	.0077	.0146	.
MAGEPAPI	58.52	.	58.529627	.	.2546	.	.	.
NEPHCIRR	.	76.98	73.15	89.81	128.30	43.89	.	.1771	.2063	.3785	.1103	.0644
NEPHOMB	14.63	1.4806
NEPHLONG	73.15	.	.	12.83	.	.	.1873	.	.1373	.	.	.
SCOLARMI	29.26	38.49	.	64.15	25.66	58.52	.4155	.6004	.	1.5358	.7108	.5413
SCOLSQUA	14.63	25.664155	.2155
SPIOBOMB	307.23	179.62	29.26	76.98	115.47	87.78	.0146	.0359	.0146	.0154	.0423	.0059
SPIOFILI	14.63	12.83	.	12.83	89.81	.	.0029	.0051	.	.0026	.0180	.
<u>Mollusca (2 species) :</u>												
MONTFERR	.	.	.	12.830051	.	.	.
NATIALDE	.	.	.	12.830038	.	.	.
<u>Crustacea (12 species) :</u>												
ATYLSWAM	.	12.830051
BATHELEG	.	14.63	12.830059	.0064	.	.	.
BATHGUIL	14.63	.	14.63	.	12.83	.	.0278	.	.0366	.	.0205	.
GASTSPIN	.	.	38.49	12.831578	.0038	.	.
HIPPENT	14.63	14.63	.00440044
LEUCLILL	.	14.630088
MEGAAGIL	58.52	25.66	14.63	64.15	.	14.63	.0219	.0103	.0059	.0257	.	.0044
PERILONG	58.520176
PONTALTA	14.630044
PSEULONG	29.26	141.13	43.89	12.83	.	14.63	.0088	.0192	.0132	.0038	.	.0029
UROTBREV	102.41	64.15	29.26	64.15	12.83	102.41	.0863	.0667	.0146	.0385	.0077	.0688
UROTOPSE	.	453.53	.	.	.	263.34	.	.	.1653	.	.	.1551
<u>Echinodermata (2 species) :</u>												
ECHICORD	14.63	12.83	87.78	12.83	12.83	29.26	18.1895	8.4768	93.3789	9.0092	12.1654	11.7918
ECHIPUSI	.	.	25.660128	.	.	.
<u>Rest (3 species) :</u>												
ANTHOZOA	.	29.26	.	.	14.63	.	.	.00880029
NEMERTIN	14.63	.	12.83	25.66	14.63	.	.3892	.	.0064	.2540	.	.0029
OLIGOCHA	29.260117

Appendix 2-7. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 7).

Density (ind./m ²) :							Biomass (g AFDW/m ²) :						
Station	71	72	73	74	75	76	71	72	73	74	75	76	
Compartment	7	7	7	7	7	7	7	7	7	7	7	7	
Number	34	35	36	37	38	39	34	35	36	37	38	39	
Date	210592	290492	290492	290492	210592	080592	210592	290492	290492	290492	210592	080592	
Species	13	15	15	24	22	18	13	15	15	24	22	18	
Density	397.73	555.94	570.57	2413.95	910.93	667.16	397.73	555.94	570.57	2413.95	910.93	667.16	
Biomass	0.93	1.40	3.89	14.16	28.69	1.30	0.93	1.40	3.89	14.16	28.69	1.30	
Depth	19.3	32.6	33.7	41.2	37.4	33.3	19.3	32.6	33.7	41.2	37.4	33.3	
<u>Polychaeta (22 species) :</u>													
ANAGROE	.	.	.	14.630146	.	.	
ARICMINU	.	14.63	14.63	.	.	12.83	.	.0073	.0073	.	.	.0038	
CHAESETO	.	.	14.63	102.41	38.490073	.0205	.0642	.	
ETEOLONG	.	.	.	14.630219	.	.	
EXOGHEBE	.	.	.	29.260029	.	.	
GONIMACU	.	.	14.63	14.63	12.830527	.0980	.0218	.	
GYPTCAFE	.	.	.	43.890263	.	.	
HARMLUNU	.	.	.	43.89	12.830512	.0128	.	
HESIAUGE	.	43.89	.	.	.	12.83	.	.00440013	
NEPHCIRR	.	58.52	131.67	190.19	141.13	38.49	.	.0571	.2414	.2911	.2104	.2104	
NEPHHOMB	.	.	.	29.262546	.	.	
NEPHLONG	.	14.633892	
PECTKORE	.	.	.	14.634184	.	.	
POECSERP	.	.	.	14.63	.	12.830746	.	.0654	
OPHELIMA	12.83	.	.	14.63	.	25.66	.0192	.	.	.0219	.	.0257	
SCOLARMI	38.49	58.52	14.63	.	12.83	102.64	.2322	.6203	.0980	.	.4824	.2874	
SCOLBONN	12.831257	
SCOLSQLA	25.663990	
SPIOBOMB	12.83	.	14.63	131.67	.	153.96	.0038	.	.0073	.0644	.	.0513	
SPIOFILI	25.66	58.52	.	.	282.26	25.66	.0077	.0439	.	.	.0449	.0077	
STHELIMI	12.833580	
STREWEBS	.	14.63	.	.	.	12.83	.	.00150013	
<u>Mollusca (3 species) :</u>													
ENSIENSI	.	.	14.637330	.	.	.	
MYSEBIDE	12.83	12.830038	.0038	
NATIALDE	12.830064	.	
<u>Crustacea (18 species) :</u>													
AMPHNEAP	.	.	.	14.630059	.	.	
ATYLSWAM	.	.	14.630059	.	.	
BATHELEG	.	14.63	43.89	.	12.83	.	.	.0059	.0176	.	.0064	.	
BATHGUIL	12.83	58.52	14.63	.	25.66	38.49	.0282	.0629	.0322	.	.0192	.0449	
CALLSUBT	.	.	.	87.78	11.8942	.	.	
CRANCRAN	12.830192	
EURYSPIN	25.660539	
GASTSPIN	51.32	73.15	14.63	.	25.66	12.83	.0436	.0819	.0658	.	.0372	.1052	
HIPPIDENT	.	.	.	14.630059	.	.	
LEUCLILL	.	.	14.630059	.	.	
MEGAAGIL	.	43.89	14.63	14.63	12.83	64.15	.	.0176	.0059	.0059	.0051	.0257	
METOBORE	12.830073	
PERILONG	12.830051	.	
PHOXFEMO	.	.	.	12.830026	.	
PSEULONG	141.13	58.52	29.26	.	38.49	25.66	.0282	.0176	.0059	.	.0077	.0051	
THIASCUT	.	.	.	12.836274	.	
UROTBREV	.	14.63	190.19	29.26	64.15	51.32	.	.0088	.0775	.0117	.0539	.0308	
UROTPOSE	.	.	.	25.660128	.	.	
<u>Echinodermata (3 species) :</u>													
ECHICORD	.	.	14.63	.	89.81	.	.	.	2.5295	.	.26.8968	.	
ECHIPUSI	.	14.63	.	131.67	12.83	.	.	.0585	.	.1302	.0051	.	
OPHIALBI	25.66	.	.	395.01	25.66	.	.0064	.	.	.2677	.0077	.	
<u>Rest (2 species) :</u>													
NEMERTIN	12.83	14.63	29.26	.	12.83	25.66	.0026	.0234	.0044	.	.1565	.0282	
PHORONID	.	.	.	1024.104550	.	.	

Appendix 2-8. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 8).

Density (ind./m²) :							Biomass (g AFDW/m²) :						
Station	81	82	83	84	85	86	81	82	83	84	85	86	
Compartment	8	8	8	8	8	8	8	8	8	8	8	8	
Number	41	42	43	44	45	46	41	42	43	44	45	46	
Date	290492	300492	070592	290492	290492	300492	290492	300492	070592	290492	290492	300492	
Species	14	19	19	15	17	18	14	19	19	15	17	18	
Density	482.79	877.80	1372.81	585.20	950.95	672.98	482.79	877.80	1372.81	585.20	950.95	672.98	
Biomass	8.64	21.63	8.28	19.29	1.16	1.99	8.64	21.63	8.28	19.29	1.16	1.99	
Depth	25.4	33.4	28.1	41.3	34.6	34.2	25.4	33.4	28.1	41.3	34.6	34.2	
<u>Polychaeta (15 species) :</u>													
ARICMINU	.	14.63	12.83	.	190.19	58.52	.	.0073	.0051	.	.0161	.0117	
CHAESETO	.	14.63	.	.	14.63	.	.	.0044	.	.	.0219	.	
ETEOFOLI	.	14.63	.	.	14.63	.	.	.0293	.	.	.0366	.	
GONIMACU	.	.	12.830411	.	.	.	
HARMLONG	.	.	.	14.630059	.	.	.	
HESIAUGA	14.630015	
LANICONC	.	.	12.83	.	.	14.63	.	.	.8378	.	.	.4857	
MAGEPAPI	.	25.661065	.	.	.	
NEPHCIRR	102.41	190.19	153.96	146.30	43.89	29.26	.2911	.1595	.3015	.2414	.1214	.0454	
SCOLARMI	43.89	87.78	102.64	29.26	87.78	117.04	.0775	.5194	.6056	.1346	.4940	1.0665	
SCOLBONN	14.630117	
SPIOBOMB	29.26	29.26	25.66	58.52	43.89	.	.0468	.0146	.0154	.0146	.0088	.	
SPIOFILI	.	12.830026	.	.	.	
STHELIMI	14.636042	
STREWEBS	14.630015	
<u>Mollusca (5 species) :</u>													
ENSIENSI	.	.	12.837929	.	.	.	
MONTFERR	.	14.63	.	.	14.63	.	.	.0044	.	.	.0029	.	
MYSEBIDE	.	.	.	14.630029	.	.	
NATIALDE	.	14.63	.	29.26	73.15	29.26	.	.0117	.	.0117	.0629	.0146	
TELLFABU	.	14.63	.	.	87.78	.	.	.0073	.	.	.1551	.	
<u>Crustacea (15 species) :</u>													
BATHELEG	43.89	29.26	436.22	.	58.52	131.67	.0219	.0088	.0975	.	.0234	.0336	
BATHGUIL	.	43.89	282.260219	.2207	
DIASRUGO	14.630219	
EBALCRAN	14.630073	
GASTSPIN	14.63	14.63	.06580658	
HIPPIDENT	.	.	.	14.63	14.630044	.0044	.	.	
LEUCINCII	.	.	.	14.630146	.	.	.	
LEUCILILL	14.630073	
MEGAAGIL	43.89	29.26	12.83	29.26	.	29.26	.0176	.0117	.0051	.0117	.	.0117	
PERILONG	.	.	.	14.630059	.	.	
PONTALTA	.	29.26	76.98	14.63	14.63	14.63	.	.0088	.0205	.0059	.0059	.0059	
PSEULONG	87.78	43.89	51.32	.	14.63	58.52	.0176	.0088	.0103	.	.0044	.0176	
THIASCUT	14.63	.	25.668690	.	.0975	.	.	.	
UROTBREV	29.26	190.19	38.49	14.63	73.15	58.52	.0146	.0995	.0257	.0088	.0336	.0293	
UROTPOSE	.	58.52	.	.	.	14.63	.	.03070059	
<u>Echinodermata (3 species) :</u>													
ECHICORD	14.63	29.26	25.66	73.15	.	.	7.1936	20.0592	4.9575	18.5596	.	.	
ECHIPUSI	.	14.63	25.66	.	73.15	14.63	.	.0146	.0192	.	.0658	.0439	
OPHALIBI	.	.	.	87.78	102.41	14.632575	.0293	.0234	
<u>Rest (1 species) :</u>													
NEMERTIN	14.63	.	25.66	29.26	29.26	29.26	.0073	.	.1168	.0059	.0688	.0966	

Appendix 2-9. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 9).

Density (ind./m²) :						Biomass (g AFDW/m²) :					
Station	91	92	93	94	95	91	92	93	94	95	
Compartment	9	9	9	9	9	9	9	9	9	9	
Number	47	48	49	50	51	47	48	49	50	51	
Date	300492	300492	300492	300492	070592	300492	300492	300492	300492	070592	
Species	17	13	17	19	21	17	13	17	19	21	
Density	672.98	643.72	629.09	2165.24	1526.77	672.98	643.72	629.09	2165.24	1526.77	
Biomass	6.72	1.15	0.79	4.27	23.90	6.72	1.15	0.79	4.27	23.90	
Depth	29.5	27.9	26.9	29.5	28.1	29.5	27.9	26.9	29.5	28.1	
<u>Polychaeta (17 species) :</u>											
ARICMINU	14.63	29.26	29.26	14.63	.	.0073	.0146	.0117	.0044	.	
CHAESETO	.	14.63	.	14.63	25.66	.	.0102	.	.0015	.0192	
ETEOFOLI	.	.	.	14.630073	.	
GONIMACU	29.26	29.26	14.63	43.89	.	.1097	.1697	.1112	.2341	.	
HARMLUNU	12.830103	
LANICONC	12.830783	
MAGEPAPI	.	14.63	.	131.67	320.75	.	.0585	.	.4330	1.4857	
NEPHCIRR	102.41	102.41	43.89	29.26	38.49	.2165	.1024	.1478	.1565	.0398	
NEPHHOMB	.	.	.	14.635238	.	
NEPHLONG	29.26	.	58.52	.	.	.1756	.	.0702	.	.	
OPHELIMA	29.260293	
SCOLARMI	14.63	102.41	14.63	29.26	12.83	.1200	.6832	.0088	.0293	.0128	
SCOLBONN	14.63	.	14.63	.	.	.0702	.	.1156	.	.	
SIGAMATH	.	.	.	12.835902	
SPIOBOMB	.	14.63	.	614.46	153.96	.	.0146	.	.2750	.0475	
SPIOFILI	.	.	14.630044	.	.	
STHELIMI	.	.	29.26	14.631287	.7871	.	
<u>Mollusca (4 species) :</u>											
DONAVITT	.	.	.	14.630073	.	
MONTFERR	64.151398	
NATIALDE	.	.	14.63	29.26	38.490102	.0293	.0385
TELLFABU	14.63	.	.	365.75	115.47	.0073	.	.	1.4601	.1232	.
<u>Crustacea (12 species) :</u>											
ATYLFALC	.	.	14.630073	.	.	
BATHELEG	117.04	43.89	73.15	629.09	205.28	.0468	.0176	.0336	.1624	.0500	
BATHGUIL	14.63	.	.	.	102.64	.00590718	
CORYCASS	12.83	10.1986	
DIASBRAD	.	.	29.260219	.	.	
GASTSPIN	.	14.63	29.260146	.0205	.	.	
MEGAAGIL	14.63	43.89	.	43.89	51.32	.0059	.0176	.	.0176	.0205	
PONTALTA	14.63	29.26	160.93	.	.	.0059	.0088	.0585	.	.	
PONTAREN	.	.	.	58.52	38.490234	.0154	
PSEULONG	43.89	146.30	58.52	29.26	89.81	.0132	.0117	.0117	.0059	.0180	
UROTBREV	131.67	58.52	14.63	.	.	.1185	.0293	.0073	.	.	
UROTPOSE	58.52	.	.	.	166.79	.01760667	
<u>Echinodermata (2 species) :</u>											
ECHICORD	14.63	.	.	.	12.83	5.7598	.	.	.	10.8593	
ECHIPUSI	.	.	.	43.89	25.660219	.0154	
<u>Rest (1 species) :</u>											
NEMERTIN	14.63	.	14.63	29.26	12.83	.0059	.	.0176	.0863	.0013	

Appendix 2-10. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 10).

Density (ind./m²) :						Biomass (g AFDW/m²) :					
Station	101	102	103	104	105	101	102	103	104	105	
Compartment	10	10	10	10	10	10	10	10	10	10	
Number	52	53	54	55	56	52	53	54	55	56	
Date	300492	200592	200592	200592	070592	300492	200592	200592	200592	070592	
Species	24	16	18	17	33	24	16	18	17	33	
Density	1492.26	1257.34	1372.81	808.29	987.91	1492.26	1257.34	1372.81	808.29	987.91	
Biomass	11.29	1.53	7.27	70.71	25.52	11.29	1.53	7.27	70.71	25.52	
Depth	34.4	26.7	28.2	26.6	27.4	34.4	26.7	28.2	26.6	27.4	
<u>Polychaeta (23 species) :</u>											
ANAIROSE	12.830103	
ARICMINU	.	89.81	.	.	25.66	.	.0180	.	.	.0103	
CHAESETO	29.26	38.49	102.64	.	12.83	.0146	.0115	.0128	.	.0192	
ETEOLONG	.	12.830192	.	.	
EUMISANG	117.04	.	.	12.83	76.98	.0878	.	.	.0128	.0077	
GLYCNORD	.	12.830436	.	.	
GONIMACU	.	38.49	.	12.83	12.83	.	.1771	.	.0462	.0744	
GYPTCAPE	14.630219	
HARMLUNU	117.04	.	.	.	76.98	.85731591	
HESIAUGE	38.490038	
LANICONC	219.45	12.83	25.66	.	128.30	.7856	.8237	.0192	.	6.6716	
MAGEPAFI	73.15	38.49	603.01	89.81	38.49	.0512	.0565	.9379	.1681	.1142	
NEPHCIRR	.	.	.	51.32	38.490821	.2540	
NEPHHOMB	73.15	25.66	64.15	.	.	3.6692	.1039	.6223	.	.	
NERELONG	12.830975	
NOTOLATE	12.830475	
OPELIMA	12.830128	
POECSERP	277.97	.	12.83	.	.	.1858	.	.0115	.	.	
SCOLARMI	25.660192	
SPIOBOMB	58.52	153.96	141.13	102.64	89.81	.0322	.1001	.1142	.0718	.0860	
SPIOFILI	12.830090	
STHELMIMI	29.26	12.83	12.83	.	.	.0263	.0603	.0269	.	.	
STREWEBS	12.830013	
<u>Mollusca (8 species) :</u>											
ABRALBALA	14.630059	
CYCLICYLI	14.63	.	12.83	.	.	.0073	.	.0051	.	.	
ENSISIL	.	.	.	12.83	25.66	.	.	.	69.8812	17.3487	
MONTFERR	102.41	.	.	12.83	.	.0293	.	.	.0038	.	
NATIALDE	29.26	.	76.98	12.83	12.83	.1258	.	.0269	.0038	.0064	
NUCUTURG	.	.	38.494516	.	.	
TELLFABU	.	64.15	115.47	25.66	.	.	.0103	.0385	.1681	.	
VENUSTRI	12.832540	
<u>Crustacea (20 species) :</u>											
AORATYPI	29.260293	
ARGIHAMA	14.630059	
ATYLSWAM	102.41	.	.	.	25.66	.02490128	
BATHELEG	.	410.56	.	128.30	.	.	.0565	.	.0513	.	
BATHGUIL	.	38.49	.	64.15	.	.	.0282	.	.0257	.	
BATHTENU	.	.	89.81	.	12.83	.	.	.0359	.	.0051	
CAPRELLI	.	.	12.83	12.830038	.0026	.	
EBALCRAN	14.630219	
HIPPIDENT	.	.	.	12.83	12.830038	.0038	
MEGAAGIL	14.630059	
MELIOTBTU	12.830128	
LEUCINCII	.	12.83	.	76.98	.	.	.0257	.	.0385	.	
PERILONG	14.63	.	.	.	12.83	.00590038	
PONTALTA	.	.	12.830051	.	.	
PONTAREN	.	64.15	.	89.81	.	.	.0231	.	.0269	.	
PSEULONG	.	25.66	12.830064	.0026	.	.	
SIRICLAU	12.830128	
TANAIDAC	25.660013	
THIASCUT	12.830090	
UROTOPSE	14.63	205.28	.	76.98	.	.0029	.0282	.	.0385	.	
<u>Echinodermata (3 species) :</u>											
ECHICORD	14.63	.	12.83	.	12.83	5.2478	.	4.8959	.	.0680	
ECHIPUSI	89.810847	
OPHIALBI	87.78	.	.	.	12.83	.01460103	
<u>Rest (3 species) :</u>											
NEMERTIN	.	25.66	.	12.83	12.83	.	.0013	.	.0872	.0872	
OLIGOCHA	38.490013	
PHORONID	14.630293	

Appendix 2-11. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 11).

Density (ind./m ²) :			Biomass (g AFDW/m ²) :		
Station	111	112	113	111	112
Compartment	11	11	11	11	11
Number	57	58	59	57	58
Date	030693	030693	030693	030693	030693
Species	20	17	23	20	17
Density	713.50	892.43	687.61	713.50	892.43
Biomass	13.97	23.12	35.59	13.97	23.12
Depth	34.3	29.2	32.7	34.3	29.2
<u>Polychaeta (16 species) :</u>					
ANAGROE	14.63	.	.	.0073	.
CHAESETO	.	29.26	.	.	.0102
DIPLGLAU	29.26	.	.	.0146	.
GONIMACU	14.63	.	14.63	.0146	.
GYPTCAPE	14.63	29.26	14.63	.0088	.0117
HARMLUNU	.	102.41	14.63	.	.1551
LANICONG	.	175.56	43.89	.	20.9882
LUMBLATR	29.26	.	29.26	.0878	.0863
MAGEPAPI	.	263.34	29.26	.	.5179
NEPHHOMB	14.63	14.63	14.63	.3219	.5940
NEREOLONG	.	.	14.63	.	.6188
OPHIFLEX	43.89	.	.	.0717	.
PHOLMINU	14.63	.	29.26	.0073	.
SCALINFL	14.63	.	.	.0044	.
SCOLARMI	.	.	14.63	.	.0073
SIGAMATH	.	58.52	14.63	.	.2326
<u>Mollusca (5 species) :</u>					
ABRAALBA	.	.	14.63	.	.0029
CULTELL	.	.	29.26	.	.0146
MONTFERR	14.63	14.63	58.52	.0073	.0029
TELLFABU	.	14.63	.	.	.0176
NATIALDE	.	.	29.26	.	.1990
<u>Crustacea (11 species) :</u>					
BODOSCOR	.	.	14.63	.	.0029
CALLSUBT	146.30	.	87.78	4.7182	.0907
CAPRELLI	.	14.63	.	.	.0015
CORYCASS	.	14.63	.	.	.0146
EUDOTRUN	14.63	.	14.63	.0029	.
HARPANTE	14.63	.	.	.0044	.
HIPPDENT	.	43.89	.	.	.0176
LEUCLILL	.	29.26	.	.	.0117
PONTALTA	14.63	.	.	.0044	.
PONTRIS	.	.	14.63	.	.0073
PSEULONG	14.63	43.89	.	.0044	.0073
<u>Echinodermata (3 species) :</u>					
AMPHFILI	29.26	.	.	.3540	.
ECHICORD	43.89	.	87.78	7.9748	.29.9052
OPHALALBI	.	14.63	14.63	.	.5340
<u>Rest (4 species) :</u>					
ANTHOZOA	.	14.63	.	.	.0073
HYDROZOA	14.63	.	.	.0029	.
NEMERTIN	29.26	14.63	14.63	.3233	.0015
PHORONID	204.82	.	73.15	.0351	.0059
					.0044

Appendix 2-12. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 12).

Density (ind./m ²) :			Biomass (g AFDW/m ²) :		
Station	121	122	123	121	122
Compartment	12	12	12	12	12
Number	60	61	62	60	61
Date	010693	010693	020693	010693	010693
Species	16	17	15	16	17
Density	409.64	848.54	936.32	409.64	848.54
Biomass	1.97	16.00	16.97	1.97	16.00
Depth	30.4	30.8	37.2	30.4	30.8
<u>Polychaeta (14 species) :</u>					
APHRACUL	14.63	.	.	.0029	.
CHAESETO	43.89	.	.	.0102	.
GATTIRR	.	14.63	.	.	.0073
GONIMACU	14.63	.	.	.0044	.
HARMLUNU	.	73.15	.	.	.2326
LANICONC	.	87.78	14.63	.	6.3875
LUMBLATR	58.52	102.41	.	.2487	.1946
MAGEPAPI	29.26	29.26	.	.0234	.0146
NEPHHOMB	14.63	58.52	14.63	.2209	5.2536
OPHIFLEX	14.63	.	14.63	.0410	.
SCALINFL	.	14.63	.	.	.0073
SPIOBOMB	.	117.04	.	.	.1814
SPIOFILI	.	.	14.63	.	.
STHELIMI	.	.	14.63	.	.
<u>Mollusca (4 species) :</u>					
ABRAALBA	.	29.26	.	.	.0029
CULTPELL	29.26	29.26	.	.0146	.0146
MYSEBIDE	14.63	.	58.52	.0029	.
TELLFABU	.	14.63	.	.	.0044
<u>Crustacea (9 species) :</u>					
BATHTENU	14.63	.	.	.0073	.
CALLSUBT	58.52	146.30	102.41	.0497	3.1659
CORYCASS	14.63	.	14.63	1.1294	.
DIASBRAD	.	.	14.63	.	.0102
EUDOTRUN	.	14.63	.	.	.0044
HIPPIDENT	.	14.63	.	.	.0132
LEUCINCI	.	14.63	.	.	.0088
PONTAREN	29.26	.	.	.0088	.
PONTTRIS	.	.	14.63	.	.
<u>Echinodermata (2 species) :</u>					
AMPHFILI	14.63	.	190.19	.1682	.
ECHICORD	.	.	29.26	.	3.3356
<u>Rest (4 species) :</u>					
ANTHOZOA	14.63	.	.	.0044	.
NEMERTIN	29.26	29.26	43.89	.0336	.4784
OLIGOCHA	.	.	14.63	.	.0366
PHORONID	.	58.52	380.38	.	.0029
				.0263	.2019

Appendix 2-13. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 13).

Density (ind./m²) :			Biomass (g AFDW/m²) :		
Station	131	132	133	131	132
Compartment	13	13	13	13	13
Number	63	64	65	63	64
Date	250593	240593	250593	250593	240593
Species	16	21	17	16	21
Density	1653.19	1700.53	2501.73	1653.19	1700.53
Biomass	2.74	7.04	14.72	2.74	7.04
Depth	28.0	30.0	28.0	28.0	30.0
<u>Polychaeta (15 species) :</u>					
ANAIROSE	175.56	38.49	.	.0307	.0064
CHAESETO	29.26	12.83	570.57	.0117	.0051
ETEOLONG	.	25.66	.	.	.0038
EXOGNAID	.	102.64	.	.	.0103
GONIMACU	43.89	51.32	29.26	.5925	.0141
HESIAUGE	.	38.49	.	.	.0051
MAGEPAPI	775.39	.	1521.52	.6671	.
NEPHCIRR	.	51.32	.	.	.6210
NEPHHOMB	29.26	.	58.52	.5486	.
NOTOLATE	.	.	29.26	.	.0922
SCALINFL	.	12.83	.	.	.1.9399
SCOLARMI	.	192.45	.	.	.0064
SCOLBONN	.	.	14.63	.	.
SIGAMATH	14.63	.	14.63	.0351	.
SPIOBOMB	131.67	.	14.63	.0263	.
<u>Mollusca (9 species) :</u>					
ABRAALBA	146.30	14.63	.	.0293	.0015
CORBGIBB	14.63	.	.	.0029	.
CULTPELL	.	.	14.63	.	.
MACTCORA	.	131.67	.	.	.0015
MONTFERR	.	.	29.26	.	.
NATIALDE	43.89	.	14.63	.0146	.
TELLFABU	29.26	.	29.26	.1902	.
TELLPYGM	.	14.63	.	.	.1039
THIASCUT	.	12.83	.	.	.1156
<u>Crustacea (7 species) :</u>					
ATYLSWAM	.	51.32	.	.	.0154
BATHTENU	.	.	43.89	.	.
DIASBRAD	.	12.83	.	.	.0132
EUDOTRUN	14.63	.	.	.0059	.
PONTAREN	.	.	14.63	.	.
PSEULONG	.	25.66	29.26	.	.0044
UROPOSE	14.63	.	.	.0038	.0029
<u>Echinodermata (2 species) :</u>					
ECHICORD	.	.	14.63	.	9.2813
ECHIPUSI	.	25.66	.	.	.0064
<u>Rest (6 species) :</u>					
ANTHOZOA	.	487.54	.	.	.3862
BRANLANC	.	128.30	.	.	2.9612
NEMERTIN	117.04	76.98	58.52	.5647	.1617
OLIGOCHA	.	192.45	.	.	.0366
PHORONID	29.26	.	.	.0117	.
PLATHYHE	43.89	.	.	.0015	.

Appendix 2-14. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 14).

Density (ind./m²) :				Biomass (g AFDW/m²) :				
Station	141	142	143	144	141	142	143	144
Compartment	14	14	14	14	14	14	14	14
Number	66	67	68	69	66	67	68	69
Date	030693	030693	030693	030693	030693	030693	030693	030693
Species	19	17	26	25	19	17	26	25
Density	892.43	1726.34	2443.21	2443.21	892.43	1726.34	2443.21	2443.21
Biomass	21.72	29.00	52.68	135.58	21.72	29.00	52.68	135.58
Depth	42.3	39.2	43.4	40.0	42.3	39.2	43.4	40.0
<u>Polychaeta (19 species) :</u>								
CHAESETO	43.89	.	14.63	.	.0146	.	.0073	.
DIPGLAU	.	43.89	.	14.63	.	.0132	.	.0410
EXOGNAID	.	14.630029	.	.
GONIMACU	14.630205	.	.	.
GYPTCAPE	29.260146	.	.	.
HARMLUNU	.	.	14.630790
LUMBLATR	.	.	14.630117	.
MEDIFRAG	.	.	14.63	14.63	.	.	.0044	.0073
NEPHHOMB	43.89	117.04	43.89	14.63	.1068	.4610	.5516	.0380
NOTOLATE	.	.	87.78	29.26	.	.	2.3057	1.7673
OPHEACUM	.	.	14.630073	.
OPHIFLEX	.	14.63	29.26	.	.	.0541	.1068	.
PHOLMINU	.	73.15	29.26	29.26	.	.0366	.0146	.0146
POLYGUIL	.	.	146.30	29.26	.	.	.2209	.0600
PRIOCRIRR	.	.	.	43.890146
SPIOBOMB	58.52	14.63	43.89	29.26	.0146	.0059	.0088	.0073
SPIOFILI	.	.	87.780146	.
STHELIMI	14.631024	.	.	.
SYNEKLAT	29.260117	.	.	.
<u>Mollusca (6 species) :</u>								
CYLCYLI	.	14.630015	.	.
LUTRLULTR	.	.	.	14.63104.3046
MONTFERR	43.89	102.41	14.63	14.63	.0980	.0380	.0044	.0059
MYSEBIDE	87.78	438.90	482.79	950.95	.0176	.0878	.0966	.1975
NUCUTURG	.	.	.	14.630059
THRACONV	.	.	14.63	.	.	.	1.0417	.
<u>Crustacea (14 species) :</u>								
AMPETENU	.	.	.	29.260146
CALLSUBT	87.78	87.78	146.30	248.71	2.4491	1.1192	3.1484	2.5515
DIASBRAD	14.630059	.	.	.
EUDOTRUN	14.63	14.63	29.26	.	.0059	.0029	.0088	.
HARPANTE	.	.	29.26	43.89	.	.	.0088	.0130
HIPPIDENT	29.260117	.	.	.
LEUCILL	14.630059	.	.	.
LEUCRICH	.	.	43.89	14.63	.	.	.0219	.0073
LIOCUPUSI	.	.	.	14.630073
MEGAAGIL	14.630044	.	.	.
MYSIDACE	.	.	.	14.630073
PERILONG	.	.	14.630044	.
PONTTRIS	.	14.63	.	.	.0073	.	.	.
UPOGDELT	.	.	58.52	14.63	.	.	21.9918	6.7532
<u>Echinodermata (4 species) :</u>								
AMPHFILI	204.82	585.20	702.24	746.13	1.9882	7.3369	10.7355	15.9760
BRISLYRI	.	.	14.63	.	.	.	10.8262	.
ECHICORD	43.89	87.78	.	.	16.8172	18.0593	.	.
LEPTINHA	.	.	14.636525	.
<u>Rest (5 species) :</u>								
NEMERTIN	14.63	29.26	14.63	14.63	.0029	1.7644	.2253	.0527
OLIGOCHA	.	43.890015	.	.
PHORONID	87.78	29.26	292.60	58.52	.0234	.0117	.1302	.0290
FRIAPULI	.	.	.	14.630073
SIPUNCUL	.	.	43.89	14.63	.	.	.5325	3.6165

Appendix 2-15. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 15).

Density (ind./m²) :				Biomass (g AFDW/m²) :		
Station	151	152	153	151	152	153
Compartment	15	15	15	15	15	15
Number	70	71	72	70	71	72
Date	020693	020693	020693	020693	020693	020693
Species	22	18	18	22	18	18
Density	1945.79	3364.90	3203.97	1945.79	3364.90	3203.97
Biomass	9.14	38.71	43.46	9.14	38.71	43.46
Depth	37.6	37.9	39.9	37.6	37.9	39.9
<u>Polychaeta (20 species) :</u>						
CHAESETO	29.26	.	.	.0219	.	.
ETEOLONG	14.63	.	.	.0293	.	.
GONIMACU	.	14.63	.	.	.0146	.
GYPTCAPE	14.63	.	.	.0146	.	.
LUMBLATR	.	58.52	.	.	.1770	.
MAGEPAPI	14.63	14.63	.	.0073	.0117	.
NEPHHOMB	.	29.26	29.26	.	1.9473	.1156
NERELONG	.	14.63	.	.	.0424	.
NOTOLATE	.	.	43.89	.	.	.9232
OPHEACUM	14.63	.	29.26	.0102	.	.0059
OPHIFLEX	14.63	.	.	.0102	.	.
OWENFUSI	.	14.63	.	.	.0146	.
PECTKORE	117.04	.	.	.0351	.	.
PHOLMINU	14.63	87.78	29.26	.0117	.0395	.0146
POLYGUIL	.	58.52	160.93	.	.0293	.1229
SCOLARMI	14.63	.	.	.0512	.	.
SIGAMATH	14.63	.	.	.6203	.	.
SPIOBOMB	73.15	.	29.26	.0146	.	.0059
STHELIMI	14.63	.	.	.0146	.	.
SYNEKLAT	29.26	43.89	14.63	.0117	.0176	.0029
<u>Mollusca (4 species) :</u>						
ABRALBA	43.89	.	.	.0058	.	.
CULTEPELL	29.26	.	14.63	.0117	.	.0044
MYSEBIDE	994.84	833.91	760.76	.1258	.2414	.1902
NUCUTURG	73.15	.	.	.5121	.	.
<u>Crustacea (5 species) :</u>						
CALLSUBT	43.89	336.49	117.04	3.1864	5.7686	2.7475
HARPANTE	73.15	117.04	29.26	.0219	.0351	.0073
HIPPDENT	14.63	.	.	.0044	.	.
LEUCINCI	.	.	14.63	.	.	.0073
UPOGDELT	.	29.26	43.89	.	.0673	14.1106
<u>Echinodermata (3 species) :</u>						
AMPHFILI	234.08	1536.15	1682.45	4.3875	26.3384	18.4879
ECHICORD	.	14.63	29.26	.	1.8258	4.8659
LEPTINHA	.	29.26	29.26	.	1.3708	.5398
<u>Rest (2 species) :</u>						
NEMERTIN	.	14.63	29.26	.	.7330	1.2509
PHORONID	58.52	117.04	117.04	.0366	.0351	.0541

Appendix 2-16. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 16).

Density (ind./m ²) :				Biomass (g AFDW/m ²) :				
Station	161	162	163	164	161	162	163	164
Compartment	16	16	16	16	16	16	16	16
Number	73	74	75	76	73	74	75	76
Date	250593	250593	250593	250593	250593	250593	250593	250593
Species	29	16	21	24	29	16	21	24
Density	2443.21	1872.64	4125.66	1784.86	2443.21	1872.64	4125.66	1784.86
Biomass	44.24	26.69	57.31	39.75	44.24	26.69	57.31	39.75
Depth	40.5	41.0	37.0	41.0	40.5	41.0	37.0	41.0
<u>Polychaeta (21 species) :</u>								
CHAESETO	14.630073	.	.	.
CHAEVARI	14.63	.	14.63	175.56	2.8924	.	2.1272	27.0787
DIPLGLAU	29.26	.	14.63	.	.0146	.	.0146	.
EXOGNAID	.	14.63	.	.	.0015	.	.	.
GATTCIRR	14.63	.	.	73.15	1.0065	.	.	1.0548
GONIMACU	14.63	14.63	.	.	.0073	.0117	.	.
GYPTCAPE	14.63	.	.	14.63	.0073	.	.	.0146
LUMBLATR	117.04	43.89	87.78	146.30	.2633	.3775	.3072	.1843
MEDIFRAG	29.26	.	.	73.15	.0059	.	.	.0117
NEPHHOMB	.	.	43.896291	.
NEPHINCI	.	.	.	14.635837
NERELONG	58.52	.	.	29.26	4.6041	.	.	.0351
NOTOLATE	14.63	.	14.63	117.04	.1507	.	.0073	1.5230
OPHIFLEX	.	.	14.630410	.
PARAGRAC	.	.	.	29.260073
PHOLMINU	29.26	14.63	190.19	29.26	.0146	.0073	.1229	.0146
POLYGUIL	73.15	73.15	146.30	43.89	.0132	.0366	.0746	.0146
SCOLARMI	.	.	.	14.639802
SPIOFILI	14.63	29.26	14.63	.	.0029	.0029	.0073	.
STHELIAMI	14.63	.	14.63	14.63	.0336	.	.0775	.0849
SYNEKLAT	58.52	.	.	14.63	.0029	.	.	.0044
<u>Mollusca (5 species) :</u>								
CORBGBIB	.	.	14.630761	.
CULTPELL	14.63	.	.	14.63	.0015	.	.	.0015
MONTFERR	.	43.89	14.63	.	.	.0161	.0015	.
MYSEBIDE	512.05	234.08	1448.37	117.04	.1024	.0702	.3804	.0293
VENUSTRI	.	14.630044	.	.
<u>Crustacea (8 species) :</u>								
AMPETENU	14.63	14.63	14.63	.	.0117	.0117	.0059	.
CALLSUBT	307.23	234.08	234.08	160.93	3.9603	5.2756	6.6991	3.7102
CIROBORE	14.630073	.	.	.
HARPANTE	43.89	.	190.19	29.26	.0146	.	.0190	.0088
LEUCRICH	.	.	29.26	.	.	.0146	.	.
PERILONG	29.26	14.63	.	.	.0088	.0044	.	.
UPOGDELT	73.15	58.52	29.26	29.26	15.5034	9.4466	10.0654	1.6927
UROTELEG	43.890117	.	.	.
<u>Echinodermata (2 species) :</u>								
AMPHFILI	775.39	1024.10	1448.37	292.60	13.4494	7.4277	34.0411	1.7541
ECHICORD	14.63	29.26	43.89	14.63	.0746	3.9911	2.5661	.0468
<u>Rest (3 species) :</u>								
NEMERTIN	14.63	.	.	43.89	.0073	.	.	.4813
PHORONID	14.63	14.63	102.41	73.15	.0059	.0059	.0322	.0310
SIPUNCUL	58.52	.	.	219.45	2.0497	.	.	.4050

Appendix 2-17. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 17).

Density (ind./m²) :				Biomass (g AFDW/m²) :		
Station	171	172	173	171	172	173
Compartment	17	17	17	17	17	17
Number	77	78	79	77	78	79
Date	020693	020693	030693	020693	020693	030693
Species	22	23	21	22	23	21
Density	3072.30	1916.53	819.28	3072.30	1916.53	819.28
Biomass	2.94	12.87	19.41	2.94	12.87	19.41
Depth	40.9	43.2	53.2	40.9	43.2	53.2
<u>Polychaeta (24 species) :</u>						
ANAIROSE	29.26	.	.	.0015	.	.
ARICJEFF	29.26	.	.	.0219	.	.
CHAESETO	.	.	14.63	.	.	.0102
CHAEVARI	.	.	14.63	.	.	4.1169
EXOGNAID	1989.68	.	.	.0424	.	.
GLYCRoux	.	.	14.63	.	.	.0073
GYPTCAPE	.	14.63	.	.	.0073	.
HESIAUGE	29.26	.	.	.0015	.	.
LUMBLATR	.	.	14.63	.	.	.0629
MAGEALLE	.	73.15	.	.	.1551	.
NEPHCIRR	43.89	.	.	.1609	.	.
NEPHHOMB	.	14.63	29.26	.	.1361	.1024
NEPHINCI	.	.	43.89	.	.	.5135
NOTOLATE	.	.	43.89	.	.	1.4162
OPHIFLEX	.	.	58.52	.	.	.1522
ORBISERT	29.26	.	14.63	.1405	.	3.5185
PARAGRAC	.	.	14.63	.	.	.0044
PHOLMINU	.	43.89	.	.	.0219	.
PISTCRIS	29.26	.	.	.0731	.	.
PRIOCIRR	.	.	87.78	.	.	.0366
SCOLARMI	58.52	14.63	.	.1726	.0146	.
SPIOBOMB	14.63	87.78	14.63	.0658	.0322	.0073
SPIOFILI	43.89	131.67	.	.0146	.0995	.
STHELIMI	14.63	29.26	.	.0585	.1814	.
<u>Mollusca (9 species) :</u>						
ABRAPRIS	29.26	.	.	.1258	.	.
ACANECHI	.	.	14.63	.	.	.0058
CULTPELL	58.52	14.63	.	.0146	.0029	.
COCHPRAE	146.30	.	.	1.4191	.	.
DOSIEXOL	160.93	.	.	.1448	.	.
ENSISILI	.	14.63	.	.	.5954	.
GARIFERV	14.63	.	.	.3204	.	.
MYSEBIDE	.	336.49	.	.	.0673	.
NUCUTURG	.	14.63	.	.	.0146	.
<u>Crustacea (13 species) :</u>						
AMPETENU	.	14.63	14.63	.	.0059	.0088
ARGIHAMA	87.78	.	.	.0263	.	.
BATHELEG	.	43.89	.	.	.0176	.
CALLSUBT	.	58.52	248.71	.	1.9092	8.8848
DIASLUCI	.	.	43.89	.	.	.0146
EUDODEFO	.	14.63	.	.	.0029	.
HARPANTE	.	29.26	.	.	.0117	.
HARPCREN	.	.	29.26	.	.	.0088
HIPPDENT	58.52	.	.	.0176	.	.
PERILONG	.	14.63	.	.	.0044	.
PONTTRIS	.	.	14.63	.	.	.0073
PSEULONG	14.63	.	.	.0059	.	.
UROTELEG	.	29.26	.	.	.0088	.
<u>Echinodermata (2 species) :</u>						
AMPHFILI	.	848.54	14.63	.	9.4700	.4594
ECHIPUSI	117.04	.	.	.0805	.	.
<u>Rest (4 species) :</u>						
ANTHOZOA	43.89	29.26	.	.0161	.0117	.
NEMERTIN	29.26	14.63	.	.0205	.0761	.
PHORONID	.	29.26	58.52	.	.0190	.0688
PLATHYHE	.	.	14.63	.	.	.0073

Appendix 2-18. Macrofauna density and biomass of MILZON-BENTHOS II 1992 (compartment 18).

Density (ind./m²) :			Biomass (g AFDW/m²) :		
Station	181	182	183	181	182
Compartment	18	18	18	18	18
Number	80	81	82	80	81
Date	250593	020693	020693	250593	020693
Species	23	28	18	23	28
Density	2370.06	1550.78	2106.72	2370.06	1550.78
Biomass	20.08	3.53	21.47	20.08	3.53
Depth	46.5	34.1	39.1	46.5	34.1
<u>Polychaeta (22 species) :</u>					
ANAIROE	.	.	14.63	.	.
ANAIROSE	.	43.89	.	.	.0015
CHAESETO	.	58.52	.	.	.0380
DIPLGLAU	.	14.63	.	.	.0073
EXOGNAID	14.63	58.52	.	.0015	.0029
GONIMACU	.	29.26	.	.	.1331
GYPTCAPE	14.63	.	14.63	.0073	.
MAGEALLE	.	14.63	.	.	.0073
MAGEPAPI	.	219.45	87.78	.	.1902
NEPHCIRR	.	14.63	.	.	.0059
NEPHHOMB	43.89	14.63	58.52	1.6166	2.7534
OPHIFLEX	29.26	.	.	.2063	.
OWENFUSI	.	43.89	.	.	.0351
PHOLMINU	29.26	.	.	.0146	.
POECSERP	14.63	14.63	.	.0176	.0336
PRIOCIRR	160.93	.	.	.1463	.
SCOLARMI	.	73.15	.	.	.0585
SIGAMATH	.	14.63	.	.	.0307
SPIOBOMB	14.63	.	.	.0073	.
SPIOFILI	.	14.63	.	.	.0029
STHELIAMI	.	14.63	43.89	.	.1214
SYNEKLAT	29.26	.	.	.0073	.
<u>Mollusca (7 species) :</u>					
CULTPELL	.	29.26	14.63	.	.0146
DOSILUPI	.	14.63	.	.	.0029
MONTFERR	.	.	43.89	.	.
MYSEBIDE	570.57	14.63	1302.07	.1024	.0029
TELLFABU	.	.	14.63	.	.
TURBLACT	.	14.63	.	.	.0029
VENUSTRI	.	.	14.63	.	5.4160
<u>Crustacea (14 species) :</u>					
AMPETENU	29.26	.	.	.0088	.
BATHELEG	.	102.41	.	.	.0307
BATHGUIL	.	14.63	.	.	.0044
BATHTENU	58.52	.	29.26	.0176	.
CALLSUBT	87.78	.	14.63	.5720	.
CAPRELLI	.	14.63	.	.	.0015
EUDODEFO	58.52	.	.	.0146	.
EUDOTRUN	14.63	.	.	.0029	.
HARPANTE	204.82	.	73.15	.0614	.
HARPCREN	29.26	.	.	.0088	.
MEGAAGIL	.	14.63	.	.	.0044
PSEULONG	.	29.26	14.63	.	.0059
UPOGDELT	14.63	.	.	.0483	.
UROTELEG	.	43.89	14.63	.	.0132
<u>Echinodermata (3 species) :</u>					
AMPHFILI	833.91	.	307.23	12.3155	.
ECHICORD	14.63	.	14.63	4.4709	.
ECHIPUSI	.	541.31	.	.	.0117
<u>Rest (3 species) :</u>					
NEMERTIN	29.26	29.26	.	.3789	.0088
PHORONID	58.52	43.89	29.26	.0410	.0088
PLATHYHE	14.63	.	.	.0073	.

Appendix 2-19. Macrobenthos density and biomass of MILZON-BENTHOS II 1992 (compartment 19).

Density (ind./m ²) :			Biomass (g AFDW/m ²) :		
Station	191	192	193	191	192
Compartment	19	19	19	19	19
Number	83	84	85	83	84
Date	250593	250593	250593	250593	250593
Species	25	21	24	25	21
Density	1828.75	2062.83	2852.85	1828.75	2062.83
Biomass	17.02	13.20	38.03	17.02	13.20
Depth	44.5	42.0	44.5	44.5	42.0
<u>Polychaeta (18 species) :</u>					
CHAESETO	.	43.89	.	.0292	.
CHAEVARI	29.26	.	29.26	4.6070	.20.6371
DIPLGLAU	14.63	.	.	.0044	.
EXOGNAID	.	29.26	.	.0015	.
GLYCROUX	.	.	73.15	.	.2253
GONIMACU	.	.	14.63	.	.0146
GYPTCAPE	14.63	.	29.26	.0146	.0219
HARMLUNU	29.26	.	43.89	1.9780	.1.0343
LUMBLATR	14.63	87.78	73.15	.0234	.4565
NEPHCIRR	.	.	14.63	.	.0161
NEPHHOMB	29.26	.	.	.7096	.
NOTOLATE	.	14.63	43.89	.	.0146
OPHIFLEX	14.63	14.63	.	.0761	.0775
POECSERP	.	.	14.63	.	.0132
PHOLMINU	29.26	29.26	87.78	.0146	.0146
PRIOCIRR	146.30	190.19	.	.1170	.1463
STHELIMI	14.63	14.63	.	.0044	.0702
SYNEKLAT	14.63	14.63	29.26	.0029	.0029
<u>Mollusca (7 species) :</u>					
ABRAALBA	14.63	.	.	.0249	.
CULTFELL	.	.	14.63	.	.0029
CYLCYLI	.	14.63	.	.0015	.
LEPTSQUA	.	.	14.63	.	.0015
MONTFERR	.	.	29.26	.	.0102
MYSEBIDE	526.68	497.42	790.02	.1053	.0878
NUCUTURG	29.26	.	14.63	.0190	.
<u>Crustacea (10 species) :</u>					
CALLSUBT	43.89	175.56	87.78	.5881	2.5062
CORYCASS	14.63	.	.	.6145	.
DIASBRAD	.	14.63	.	.	.0117
EUDOTRUN	14.63	.	.	.0029	.
HARPANTE	131.67	14.63	43.89	.0395	.0044
LEUCLILL	.	14.63	.	.	.0044
PERILONG	.	14.63	.	.	.0044
PONTRIS	14.63	.	.	.0073	.
UPOGDELT	.	14.63	.	.	2.5705
UROTELEG	14.63	14.63	.	.0044	.
<u>Echinodermata (3 species) :</u>					
AMPHFILI	555.94	746.13	1053.36	7.8314	7.1511
ECHICORD	.	.	29.26	.	.3.1937
LEPTINHA	.	.	14.63	.	.0585
<u>Rest (4 species) :</u>					
NEMERTIN	14.63	14.63	14.63	.0029	.0278
PHORONID	14.63	87.78	.	.0073	.0132
PLATHYHE	43.89	.	14.63	.1112	.
SIPUNCUL	43.89	.	277.97	.1053	.2546

Appendix 3-1. Densities (ind./10 cm²) of meiobenthic taxa (permanent and temporary) on the EXP*BMN 1992 locations.

Station	N 50	META 1	N 70
<hr/>			
<u>Permanent</u>			
Nematoda	1510	1347	840
Copepoda	143	93	194
burrowing	3	3	11
interstitial	140	90	183
Gastropoda	220	237	100
Turbellaria	80	21	80
Archiannelida	.	.	.
Oligochaeta	.	.	.
Polychaeta	.	3	.
Hydrozoa	10	20	.
Tardigrada	70	19	60
Ostracoda	.	.	10
Halacarida	10	1	.
<hr/>			
<u>Temporary</u>			
Nauplii	.	.	.
Bivalvia	.	.	.
Foraminifera	.	.	.
Isopoda	.	.	.
Cumacea	.	.	.
Nemertina	.	.	.
Tanaidacea	.	.	.
Amphipoda	.	.	.
Echinoidea	.	.	.
Ophiuroidea	.	.	.
Cnidaria other	.	.	.
Bryozoa	.	.	.
Taxa true	7	8	6
Taxa temporary	.	.	.
Taxa total	7	8	6

Appendix 3-2. Macrobenthos density and biomass of EXP*BMN 1992 (compartment 2, 3 and 7).

Density (ind./m ²) :				Biomass (g AFDW/m ²) :		
Station	N 50	META 1	N 70	N 50	META 1	N 70
Compartment	2	3	7	2	3	7
Number	11	17	40	11	17	40
Date	310392	310392	200592	310392	310392	200592
Species	20	14	21	20	14	21
Density	1316.70	746.13	716.87	1316.70	746.13	716.87
Biomass	2.12	1.90	33.73	2.12	1.90	33.73
Depth	30	27	32	30	27	32
<u>Polychaeta :</u>						
ANAIISUBU	14.63	.	.	.0059	.	.
ARICMINU	29.26	87.78	.	.0044	.0132	.
CHAESETO	14.63	.	43.89	.	.	.0132
HARMLUNU	.	.	14.63	.	.	.0702
LANICONC	.	.	58.52	.	.	.6935
MAGEPAPI	.	.	29.26	.	.	.0073
NEPHCIRR	131.67	160.93	.	.0600	.3701	.
NEPHHOMB	.	.	29.26	.	.	.0161
OPHELIMA	29.26	.	.	.0951	.	.
PECTAURI	.	.	14.63	.	.	.0015
PHOLMINU	.	.	43.89	.	.	.0219
SCOLARMI	409.64	73.15	.	.9056	.7739	.
SCOLSQUA	.	29.26	.	.	.2063	.
SIGAMATH	.	.	14.63	.	.	.0922
SPIOBOMB	43.89	29.26	14.63	.0102	.0775	.0044
STHELIMI	.	.	29.26	.	.	.0073
TRAVFORB	.	14.63	.	.	.0029	.
<u>Mollusca :</u>						
NATIALDE	14.63	58.52	43.89	.0015	.0234	.0102
NUCUTURG	.	.	43.89	.	.	.0073
TELLFABU	14.63	.	14.63	.6013	.	.
<u>Crustacea :</u>						
BATHELEG	321.86	14.63	.	.1609	.0073	.
BATHGUIL	29.26	102.41	.	.0146	.1302	.
BODOAREN	.	.	14.63	.	.	.0073
DIASBRAD	29.26	.	.	.0146	.	.
HARPARANTE	.	.	14.63	.	.	.0073
HIPPDDENT	14.63	.	.	.0073	.	.
IPHITRIS	.	.	14.63	.	.	.0073
MEGAAGIL	102.41	73.15	.	.0512	.0366	.
PERILONG	.	.	58.52	.	.	.0293
PSEULONG	.	.	14.63	.	.	.0073
PSEUSIMI	43.89	.	.	.0219	.	.
SYNCMACU	14.63	.	.	.0073	.	.
THIASCUT	14.63	.	.	.0600	.	.
UROPOSE	.	14.63	.	.	.0073	.
UROTBREV	.	43.89	.	.	.0205	.
<u>Echinodermata :</u>						
ACROBRAC	.	.	14.63	.	.	.0015
AMPHFILI	.	.	43.89	.	.	.0805
ECHICORD	14.63	.	146.30	.0761	.	32.6395
ECHIPUSI	14.63	14.63	.	.0146	.0468	.
<u>Rest :</u>						
NEMERTIN	14.63	29.26	.	.0029	.1829	.

NIOZ-RAPPORT 1994 - 1
NIOO-CEMO RAPPORTEN EN VERSLAGEN 1994 - 1

CONTENTS

Summary	i
Samenvatting	iii
1 Preface	1
2 Introduction	2
2.1 General abiotic features	2
2.2 General biotic features	2
2.3 Acknowledgements	3
3 Material and methods	4
3.1 Sampling	4
3.2 Treatment of samples	4
3.3 Sediment classification	5
3.4 Statistical analysis	5
4 Results	7
4.1 Environmental parameters	7
4.2 Meiobenthos	8
4.3 Macrobenthos	11
4.4 Comparison ICES/MILZON I+II	14
4.5 Correlations benthos/environment	16
4.6 Relationship meio-/macrobenthos	17
5 Discussion	18
5.1 Meiobenthos	18
5.2 Macrobenthos	19
6 Conclusions	21
7 Recommendations	23
8 References	24
9 Tables	27
10 Figures	51
11 Appendices	109